



HU000033509T2

(19) **HU**(11) Lajstromszám: **E 033 509**(13) **T2****MAGYARORSZÁG****Szellemi Tulajdon Nemzeti Hivatala**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 12 746134**(22) A bejelentés napja: **2012. 07. 20.**(51) Int. Cl.: **A01P 7/02** (2006.01)**A01N 43/54** (2006.01)**A01N 43/76** (2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20120746134

(97) Az európai bejelentés közzétételi adatai:

EP 2734045 A1 **2013. 01. 24.**

(86) A nemzetközi (PCT) bejelentési szám:

PCT/JP 12/069071

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2734045 B1 **2016. 12. 28.**

(87) A nemzetközi közzétételi szám:

WO 13012099

(30) Elsőbbségi adatok:

2011159711 **2011. 07. 21.** **JP**

(72) Feltaláló(k):

NATSUHARA, Katsuya, Tokyo 104-8260 (JP)**TANAKA, Azusa, Takarazuka-shiHyogo 665-8555 (JP)**

(73) Jogosult(ak):

Sumitomo Chemical Company, Limited, Tokyo 104-8260 (JP)

(74) Képviselő:

Danubia Szabadalmi és Jogi Iroda Kft., Budapest

(54)

Kártevőirtó készítmény

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.

(19)



(11)

EP 2 734 045 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
28.12.2016 Bulletin 2016/52

(51) Int Cl.:
A01P 7/02 (2006.01) **A01N 43/76** (2006.01)
A01N 43/54 (2006.01)

(21) Application number: **12746134.1**

(86) International application number:
PCT/JP2012/069071

(22) Date of filing: **20.07.2012**

(87) International publication number:
WO 2013/012099 (24.01.2013 Gazette 2013/04)

(54) PEST CONTROL COMPOSITION

SCHÄDLINGSBEKÄMPFUNGZUSAMMENSETZUNG

COMPOSITION ANTIPARASITAIRE

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **21.07.2011 JP 2011159711**

(43) Date of publication of application:
28.05.2014 Bulletin 2014/22

(73) Proprietor: **Sumitomo Chemical Company,
Limited
Tokyo 104-8260 (JP)**

(72) Inventors:
• **NATSUHARA, Katsuya**
Tokyo 104-8260 (JP)
• **TANAKA, Azusa**
Takarazuka-shi
Hyogo 665-8555 (JP)

(74) Representative: **Vossius & Partner**
Patentanwälte Rechtsanwälte mbB
Siebertstrasse 3
81675 München (DE)

(56) References cited:
CN-A- 102 228 055 US-A1- 2010 216 738

EP 2 734 045 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

Technical Field


[0001] The present application is filed claiming the priority of the Japanese Patent Application No. 2011-159711.
[0002] The present invention relates to a pest control composition and a pest control method.

Background Art

[0003] Etoxazole, 2-(2,6-difluorophenyl)-4-[4-(1,1-dimethyl ethyl)-2-ethoxyphenyl]-4,5-dihydrooxazole) is known as an active ingredient of a pest control agent (see, e.g., Patent Literature 1).

[0004] Also, fenazaquin, 4-tert-butylphenethylquinazolin-4-yl ether, is known as an active ingredient of a pest control agent (see, e.g., Patent Literature 2).

[0005] Patent Literature 3 relates to the use of active substance combinations comprising the compound of the following formula (I)



(I)

and active substances from the IRAC classes of the sodium channel modulators/blockers and/or site I electron transport inhibitors and/or chloride channel activators and/or inhibitors of the magnesium-stimulated ATPase and/or bifenazate for controlling animal pests. These active substance combinations may also comprise the compound of formula (I) and at least one of the compounds bifenazate, fenpyroximate, pyridaben, fenazaquin, abamectin, emamectin, fenpropathrin, and propargite.

Citation List

Patent Literature

[0006]

Patent Literature 1: WO93/22297

Patent Literature 2: EP0326329A

Patent Literature 3: US2010/0216738A1

Summary of Invention

Technical Problem

[0007] An object of the present invention is to provide a pest control composition having an excellent control effect on pests and a pest control method.

Solution to Problem

[0008] The present inventors have intensively studied and finally found that a combination of etoxazole and fenazaquin has an excellent control effect on pests. Thus, the present invention has been completed.

[0009] Namely, the present invention includes the followings:

[1] A pest control composition comprising etoxazole and fenazaquin.

[2] The pest control composition according to the above [1], wherein the weight ratio of etoxazole to fenazaquin is from 5000:1 to 1:5000.

[3] The pest control composition according to the above [1], wherein the weight ratio of etoxazole to fenazaquin is from 100:1 to 1:100.

[4] A pest control method, which comprises applying effective amounts of etoxazole and fenazaquin to a pest or an area where a pest lives, with the proviso that a method for the treatment of the human or animal body by therapy is excluded.

[5] The pest control method according to the above [4], wherein the weight ratio of etoxazole to fenazaquin is from 5000:1 to 1:5000.

[6] The pest control method according to the above [4], wherein the weight ratio of etoxazole to fenazaquin is from 100:1 to 1:100.

[7] Use of etoxazole and fenazaquin as a pest control agent, with the proviso that a use for the treatment of the human or animal body by therapy is excluded.

Effects of Invention

[0010] The pest control composition of the present invention has an excellent control effect on pests.

Description of Embodiments

[0011] The pest control composition of the present invention contains etoxazole and fenazaquin.

[0012] Etoxazole is a known compound and can be produced by a process described in, for example, WO93/22297.

[0013] Fenazaquin is described in, for example, EP0326329A and can be produced by a process described therein.

[0014] In the pest control composition of the present disclosure, the weight ratio of etoxazole to Fenazaquin is, for example, from 5000:1 to 1:5000, from 500:1 to 1:5000, from 300:1 to 1:4100, from 250:1 to 1:4097, from 100:1 to 1:100, from 16:1 to 1:16 and from 1:3 to 1:16.

[0015] The pest control composition of the present invention may be a simple mixture of etoxazole and fenazaquin. However, the pest control composition of the present acetone, and water.

[0016] Examples of the gaseous carrier include fluorocarbon, butane gas, liquefied petroleum gas (LPG), dimethyl ether, and carbon dioxide.

[0017] Examples of the surfactant include anionic surfactants (e.g. alkyl sulfate ester salts, alkylaryl sulfonates, dialkyl sulfosuccinates, polyoxyethylene alkylaryl ether phosphate ester salts, ligninsulfonates, naphthalene sulfonate formaldehyde polycondensates, styrene-acrylate copolymers, and methyl oleyl taurate sodium salts), nonionic surfactants (e.g. polyoxyethylene alkylaryl ethers, polyoxyethylene alkylpolyoxypropylene block copolymers, and sorbitan fatty acid esters), and cationic surfactants (e.g. alkyl trimethyl ammonium salts).

[0018] Examples of the other formulation additives include water-soluble polymers (e.g., polyvinyl alcohol, and polyvinyl pyrrolidone), polysaccharides [e.g., gum arabic, alginic acid and a salt thereof, CMC (carboxymethyl cellulose), and xanthane gum], inorganic substances (e.g. aluminum magnesium silicate, smectite, and alumina-sol), preservatives (e.g. 5-chloro-2-methyl-4-isothiazolin-3-one, 1,2-benzothiazolin-3-one, and 2-bromo-2-nitropropane-1,3-diol), colorants, and stabilizers [e.g. PAP (isopropyl acid phosphate), and BHT (2,6-di-tert-butyl-4-methylphenol)].

[0019] Examples of the pest on which the pest control composition of the present invention has a controlling effect include arthropods such as insects and mites, and Nematelminthes such as nematodes, as listed below.

Hemiptera:

[0020] Delphacidae such as *Laodelphax striatellus*, *Nilaparvata lugens*, and *Sogatella furcifera*; Deltocephalidae such as *Nephotettix cincticeps*, and *Nephotettix virescens*; Aphididae such as *Aphis gossypii*, *Myzus persicae*, *Brevicoryne brassicae*, *Macrosiphum euphorbiae*, *Aulacorthum solani*, *Rhopalosiphum padi*, and *Toxoptera citricidus*; Pentatomidae such as *Nezara antennata*, *Riptortus clavatus*, *Leptocoris chinensis*, *Eysarcoris parvus*, *Halyomorpha mista*, and *Lygus lineolaris*; Aleyrodidae such as *Trialeurodes vaporariorum*, *Bemisia tabaci*, *Bemisia argentifolii*, and *Aleurocanthus spiniferus*; Coccidae such as *Aonidiella aurantii*, *Comstockaspis perniciosus*, *Unaspis citri*, *Ceroplastes rubens*, *Icerya purchasi*, and *Pseudaulacaspis pentagona*; Tingidae; Psyllidae; etc.

Lepidoptera:

[0021] Pyralidae such as *Chilo suppressalis*, *Tryporyza incertulas*, *Cnaphalocrocis medinalis*, *Notarcha derogata*, *Plodia interpunctella*, *Ostrinia furnacalis*, *Ostrinia nubilalis*, *Hellula undalis*, and *Pediasia teterrellus*; Noctuidae such as

Spodoptera litura, *Spodoptera exigua*, *Pseudaletia separata*, *Mamestra brassicae*, *Agrotis ipsilon*, *Plusia nigrisigna*, *Thoricoplusia* spp., *Heliothis* spp., and *Helicoverpa* spp.; Pieridae such as *Pieris rapae*; Tortricidae such as *Adoxophyes* spp., *Grapholita molesta*, *Leguminivora glycinivorella*, *Matsumuraeses azukivora*, *Adoxophyes orana fasciata*, *Adoxophyes* sp., *Homona magnanima*, *Archips fuscocupreanus*, and *Cydia pomonella*; Gracillariidae such as *Caloptilia theivora*, and *Phyllonorycter ringoneella*; Carposinidae such as *Carposina niponensis*; Lyonetiidae such as *Lyonetia* spp.; Lymantriidae such as *Lymantria* spp., and *Euproctis* spp.; Yponomeutidae such as *Plutella xylostella*; Gelechiidae such as *Pectinophora gossypiella*, and *Phthorimaea operculella*; Arctiidae such as *Hyphantria cunea*; Tineidae such as *Tinea translucens*, and *Tineola bisselliella*; etc.

Thysanoptera:

[0022] Thripidae such as *Frankliniella occidentalis*, *Thrips palmi*, *Scirtothrips dorsalis*, *Thrips tabaci*, *Frankliniella intonsa*, and *Frankliniella fusca*; etc.

Diptera:

[0023] *Musca domestica*, *Culex pipiens pallens*, *Tabanus trigonus*, *Hylemya antiqua*, *Hylemya platura*, *Anopheles sinensis*, *Agromyza oryzae*, *Hydrellia griseola*, *Chlorops oryzae*; Agromyzidae such as *Liriomyza trifolii*; *Dacus cucurbitae*, *Ceratitis capitata*; etc.

Coleoptera:

[0024] *Epilachna vigintioctopunctata*, *Aulacophora femoralis*, *Phyllotreta striolata*, *Oulema oryzae*, *Echinocnemus squameus*, *Lissorhoptrus oryzophilus*, *Anthonomus grandis*, *Callosobruchus chinensis*, *Sphenophorus venatus*, *Popillia japonica*, *Anomala cuprea*, *Diabrotica* spp., *Leptinotarsa decemlineata*, *Agriotes* spp., *Lasioderma serricorne*, *Anthrenus verbasci*, *Tribolium castaneum*, *Lyctus brunneus*, *Anoplophora malasiaca*, *Tomicus piniperda*, etc.

Orthoptera:

[0025] *Locusta migratoria*, *Grylotalpa africana*, *Oxya yezoensis*, *Oxya japonica*, etc.

Hymenoptera:

[0026] *Athalia rosae*, *Acromyrmex* spp., *Solenopsis* spp., etc.

Blattodea:

[0027] *Blattella germanica*, *Periplaneta fuliginosa*, *Periplaneta americana*, *Periplaneta brunnea*, *Blatta orientalis*, etc.

Acarina:

[0028] Tetranychidae such as *Tetranychus urticae*, *Panonychus citri*, and *Oligonychus* spp.; Eriophyidae such as *Aculops pelekassi*; Tarsonemidae such as *Polyphagotarsonemus latus*; Tenuipalpidae; Tuckerellidae; Acaridae such as *Tyrophagus putrescentiae*; Pyroglyphidae such as *Dermatophagoides farinae*, and *Dermatophagoides pteronyssus*; Cheyletidae such as *Cheyletus eruditus*, *Cheyletus malaccensis*, and *Chelacaropsis moorei*; etc.

Nematoda:

[0029] *Aphelenchoides besseyi*, *Nothotylenchus acris*, etc.

[0030] The pest control method of the present invention comprises applying effective amounts of etoxazole and fenazaquin to a pest or an area where a pest lives.

[0031] Herein, the "effective amounts" mean the total amount of etoxazole and fenazaquin in which amount the application of both compounds can make a pest controlled.

[0032] Examples of the area where a pest lives include crops and soil where crops are grown.

[0033] The pest control method of the present invention can be carried out by applying the pest control composition of the present invention to a pest or an area where a pest lives. The pest control method of the present invention can be also carried out by applying etoxazole and fenazaquin separately to a pest or an area where a pest lives.

[0034] In the pest control method of the present disclosure, the weight ratio of etoxazole to fenazaquin is, for example,

from 5000:1 to 1:5000, from 500:1 to 1:5000, from 300:1 to 1:4100, from 250:1 to 1:4097, from 100:1 to 1:100, from 16:1 to 1:16 and from 1:3 to 1:16.

[0035] In the pest control method of the present invention, application of etoxazole and fenazaquin can be carried out by, for example, spraying the foliage of crops with etoxazole and fenazaquin, irrigating soil where crops are grown with etoxazole and fenazaquin, or treating the seeds of crops with etoxazole and fenazaquin.

[0036] When etoxazole and fenazaquin are applied to the foliage of crops or the soil where crops are grown, the application rate is generally from 0.1 to 1000 g per 10000 m², preferably from 1 to 200 g per 10000 m², in terms of the total amount of etoxazole and fenazaquin, although it may be varied depending on the kinds of crops to be protected from pests, the kinds of target pests, the population size of target pests, the type of a formulation, the application period, and climate conditions.

[0037] When the seeds of crops are treated with etoxazole and fenazaquin, the treatment rate is generally from 0.001 to 20 g, preferably from 0.01 to 10 g per 1 kg of seeds, in terms of the total amount of etoxazole and fenazaquin.

[0038] When etoxazole and fenazaquin are formulated into emulsifiable concentrate, wettable powder or suspension concentrate, the formulation is generally diluted with water and then sprayed. In this case, the formulation is diluted so that the total concentration of etoxazole and fenazaquin becomes generally from 1 to 10000 ppm, preferably from 10 to 500 ppm.

[0039] When etoxazole and fenazaquin are formulated into dusts or granules, the formulation is generally applied as it is without diluting it.

[0040] The pest control composition of the present invention can be used in pest control for plants including, but not limited to, "crops" listed below.

[0041] "Crops":

Agricultural crops: corn, wheat, barley, rye, oat, sorghum, cotton, soybean, kidney bean, azuki bean, rice, peanut, sarrazin, sugar beet, rapeseed, sunflower, sugar cane, tobacco etc.;

Vegetables: Solanaceae vegetables (eggplant, tomato, green pepper, hot pepper, potato, etc.), Cucurbitaceae vegetables (cucumber, pumpkin, zucchini, watermelon, melon, etc.), Cruciferae vegetables (Japanese radish, turnip, horseradish, kohlrabi, Chinese cabbage, cabbage, brown mustard, broccoli, cauliflower, etc.), Compositae vegetables (burdock, garland chrysanthemum, artichoke, lettuce, etc.), Liliaceae vegetables (Welsh onion, onion, garlic, asparagus, etc.), Umbelliferae vegetables (carrot, parsley, celery, parsnip, etc.), Chenopodiaceae vegetables (spinach, chard, etc.), Labiatae vegetables (Japanese basil, mint, basil, etc.), strawberry, sweet potato, yam, aroid etc.; Flowers and ornamental plants: acanthus, morning glory, azalea, hydrangea, anemone raddeana, rhodohypoxis baurii; anemone, polygonatum odoratum, amaryllis, iris, alyssum, armeria, arctotis, China aster, edible flower, Bauera rubioides, Cuban lily, Hosta montana, Mexican aster, four o'clock, Hypericum, oriental poppy, gentiana makinoi, Hosta aureomarginata, Japanese iris, clematis patens, gazania, Casa Blanca, carnation, showy lily, gerbera, kalanchoe, calceolaria, curry plant, Carolina jasmine, canna, chrysanthemum, Brugmansia, yellow cosmos, plantain lily, KimJongilia, tea tree (Manuka), pot marigold, myrtle, nasturtium, gladiolus, Siam tulip, clematis, cockscomb, shrimp plant, midday flower, cosmos, Hosta sieboldii, Convolvulus arvensis, Hosta sagae, primrose, saffron crocus, salvia, cyclamen, moss phlox, Paeonia lactiflora, Anemone hupehensis, Bletilla striata, sweet pea, lily of the valley, snowflake, portulaca, violet, rose of Sharon, yarrow, Chinese pink, zephyranthes, pelargonium, geum, zepher lily, dahlia, tithonia, tulip, chocolate cosmos, Vinca major, scilla, downy myrtle, German iris, passionflower, dianthus, rape blossom, Madagascar periwinkle, soft windflower, nemophila, Nerine, swamp chrysanthemum (North pole), Japanese water iris (Iris ensata var. spontanea), verbena, hibiscus, Joseph's coat, coral flower, Japanese water iris (Iris ensata), eastern redbud, spring starflower, wavyleaf sea-lavender, California poppy, pansy, Virginia stock, daisy, corn poppy, Himalayan creeping saxifrage, sunflower, hyacinth, crape-myrtle, Geranium, fuchsia, freesia, primula, garden balsam, ground-cherry, peony, Tricyrtis, marguerite, marigold, Gymnaster savatieri, strawflower, muscari, Japanese kerria, lily, ranunculus, lantana, gentian, Lupinus, lobelia, etc.;

Ornamental foliage plants: ivy, cat tail, aglaonema, adiantum, asparagus, asplenium, ananas, aphelandra, alocasia, anthurium, Indian rubber tree, nepenthes, aechmea, aeschynanthus, episcia, strelitzia augusta, spiders plant, Chinese banyan, kapok, caladium, calathea, velvet plant (Gynura), Guzumania, Ctenanthe, gum tree, crassula, croton, Alocasia odora, orange jessamine, coffee tree, massangeana, conifers, coleus, cordyline, columnnea, sansevieria, sansevieria, Chinese ixora, schefflera, cissus, cyperus, reed rhaps, silk jessamine, syngonium, strelitzia, spathiphyllum, senecio, zebrina, Japanese sago palm, tillandsia, tupidanthus, coral tree, dizygotheca, dieffenbachia, duranta, bottle palm, dracaena, tradescantia, neoregelia, nephrolepis, hearts vine, hibiscus, pachypodium, Guiana chestnut (Pachira), ponytail, staghorn fern, pilea, fatshedera, ficus pumila, philodendron, bougainvillea, phoenix, fittonia, pteris, bridal veil, vriesia, plectranthus, begonia, peperomia, heliconia, benjamina, poinsettia, pothos, hoyo, maranta, Belgian evergreen, milkbush, oyster plant, monstera, palm, yucca, lantana, etc.;

Fruit trees: pomaceous fruits (apple, common pear, Japanese pear, Chinese quince, quince, etc.), stone fruits (peach, plum, nectarine, Japanese plum, cherry, apricot, prune, etc.), citrus plants (Satsuma mandarin, orange,

lemon, lime, grapefruit, etc.), nuts (chestnut, walnut, hazel nut, almond, pistachio, cashew nut, macadamia nut, etc.), berry fruits (blueberry, cranberry, blackberry, raspberry, etc.), grape, persimmon, olive, loquat, banana, coffee, date, coconut, etc.;

Trees other than fruit trees: tea, mulberry, flowering trees and shrubs, street trees (ash tree, birch, dogwood, eucalyptus, ginkgo, lilac, maple tree, oak, poplar, cercis, Chinese sweet gum, plane tree, zelkova, Japanese arborvitae, fir tree, Japanese hemlock, needle juniper, pine, spruce, yew), etc.

[0042] The above-described "crops" include plants having the resistance to herbicides which is imparted by a classic breeding method or a genetic engineering technique.

[0043] In the present invention, etoxazole and fenazaquin may be used in admixture with or in combination with other active ingredients such as other insecticides, acaricides, nematocides, fungicides, herbicides, plant hormones, and plant growth regulators; synergists; safeners; pigments, fertilizers; soil conditioners; and/or animal feed.

Examples

[0044] Hereinafter, the present invention is described specifically by way of Formulation Examples and Test Examples to which the present invention is not limited.

[0045] First, Formulation Examples are described. In Examples, the term "part(s)" means part(s) by weight.

Formulation Example 1

[0046] Five parts of etoxazole, 5 parts of fenazaquin, 8 parts of polyoxyethylene styrylphenyl ether, 2 parts of calcium dodecylbenzenesulfonate, and 80 parts of xylene are mixed to obtain an emulsifiable concentrate.

Formulation Example 2

[0047] A mixture of 20 parts of etoxazole, 4 parts of fenazaquin, 3 parts of sodium dodecylbenzenesulfonate, 3 parts of sodium ligninsulfonate, and 70 parts of diatomaceous earth is pulverized in a jet air mill to obtain a wettable powder.

Formulation Example 3

[0048] One part of etoxazole, 0.5 parts of fenazaquin, 48.5 parts of talc, and 50 parts of clay are mixed to obtain dusts.

Formulation Example 4

[0049] A mixture of 1 part of etoxazole, 4 parts of fenazaquin, 5 parts of sodium dodecylbenzenesulfonate, 30 parts of bentonite, and 60 parts of clay is stirred with an appropriate amount of water, granulated in a granulator, and then dried under ventilation to obtain granules.

Formulation Example 5

[0050] To a mixture of 5 parts of polyoxyethylene styrylphenyl ether sulfate, 20 parts of a 1% aqueous xanthan gum solution, 3 parts of a smectite mineral, and 60 parts of water are added 5 parts of etoxazole and 5 parts of fenazaquin. The mixture is stirred and then wet ground in a sand mill to obtain a suspension concentrate.

Formulation Example 6

[0051] Firstly, 0.1 part of etoxazole and 0.02 parts of fenazaquin are solved in 10 parts of acetone. Then, the solution is uniformly mixed with 99.88 parts of animal solid feed powder (CE-2: a solid powdery diet for growing and breeding manufactured by CLEA Japan, Inc.), and then the acetone is removed by air drying to obtain a poison bait.

Formulation Example 7

[0052] Firstly, 0.1 part of etoxazole and 0.1 part of fenazaquin are solved in 5 parts of xylene and 5 parts of trichloroethane. Then, the solution is mixed with 89.8 parts of deodorized kerosene to obtain an oil solution.

[0053] Next, Test Examples for pest control by the present invention are described.

Test Example 1

[0054] Kidney beans (Nagauzura saitou) were planted (one plant per container) in plastic containers (volume: 90 ml) and grown until the primary leaves had fully expanded. Only one leaf of each kidney bean plant was left on the plant and all the other leaves were removed from the plant. Then, 5 adults of *Tetranychus urticae* were released on each leaf.

[0055] A suspension concentrate containing 10.0% by weight of etoxazole [product name: Baroque (registered trademark) flowable, manufactured by Kyoyu Agri Co., Ltd.] was diluted with water containing 0.02% by volume of a spreading agent [product name: Sindain (registered trademark), manufactured by Sumitomo Chemical Co., Ltd.] so that the concentration of etoxazole became 12.5 ppm.

[0056] A reference standard containing 99.9% by weight of fenazaquin was diluted with water containing 0.02% by volume of a spreading agent [product name: Sindain (registered trademark), manufactured by Sumitomo Chemical Co., Ltd.] so that the concentration of fenazaquin became 50 ppm.

[0057] The water dilution of etoxazole, the water dilution of fenazaquin, and water containing 0.02% by volume of a spreading agent [product name: Sindain (registered trademark), manufactured by Sumitomo Chemical Co., Ltd.] were mixed to prepare a test solution with a predetermined concentration of etoxazole and fenazaquin.

[0058] Three days after the release of the insects, each test solution was sprayed onto the leaves of kidney bean plants and the leaves were air-dried. Two days after the treatment, the tested insects were observed for life or death. An insect death rate was calculated according to the following equation. For each treatment there are duplicates. Results are shown in Table 1.

$$\text{Insect death rate (\%)} = 100 \times (\text{number of dead insects} / \text{number of tested insects})$$

Table 1

Etoxazole	Fenazaquin	Insect death rate (%)
Concentration (ppm)	Concentration (ppm)	
0.000781	-	10
-	3.2	40
0.000781	3.2	90

Test Example 2

[0059] Kidney beans (Nagauzura saitou) were planted (one plant per container) in plastic containers (volume: 90 ml) and grown until the primary leaves had fully expanded. Only one leaf of each kidney bean plant was left on the plant and all the other leaves were removed from the plant. Then, 10 adults of *Tetranychus urticae* were released on each leaf.

[0060] A suspension concentrate containing 10.0% by weight of etoxazole [product name: Baroque (registered trademark) flowable, manufactured by Kyoyu Agri Co., Ltd.] was diluted with water containing 0.02% by volume of a spreading agent [product name: Sindain (registered trademark), manufactured by Sumitomo Chemical Co., Ltd.] so that the concentration of etoxazole became 50 ppm.

[0061] A reference standard containing 99.9% by weight of fenazaquin was diluted with water containing 0.02% by volume of a spreading agent [product name: Sindain (registered trademark), manufactured by Sumitomo Chemical Co., Ltd.] so that the concentration of fenazaquin became 50 ppm.

[0062] The water dilution of etoxazole, the water dilution of fenazaquin, and water containing 0.02% by volume of a spreading agent [product name: Sindain (registered trademark), manufactured by Sumitomo Chemical Co., Ltd.] were mixed to prepare a test solution with a predetermined concentration of etoxazole and fenazaquin.

[0063] Four days after the release of the insects, each test solution was sprayed onto the leaves of kidney bean plants and the leaves were air-dried. Three days after the treatment, the tested insects were observed for life or death. An insect death rate was calculated according to the following equation. For each treatment there are duplicates. Results are shown in Table 2.

Insect death rate (%) = $100 \times (\text{number of dead insects} / \text{number of tested insects})$

Table 2

Etoxazole	Fenazaquin	Insect death rate (%)
Concentration (ppm)	Concentration (ppm)	
0.8	-	15
-	3.2	40
-	12.5	50
0.8	3.2	60
0.8	12.5	95

Test Example 3

[0064] Cabbages (Green ball) were planted (one plant per container) in plastic containers (volume: 90 ml) and grown until the third to fourth leaf stage. Only one leaf of each cabbage plant was left on the plant and all the other leaves were removed from the plant. The plastic containers were put into a net cage containing a lot of living *Bemisia tabaci* and allowed to stand for 24 hours. After that, the plastic containers were taken out from the cage and the number of living insects (*Bemisia tabaci*) on the cabbage seedlings was determined (This is called the "number of insects before treatment").

[0065] A suspension concentrate containing 10.0% by weight of etoxazole [product name: Baroque (registered trademark) flowable, manufactured by Kyoyu Agri Co., Ltd.] was diluted with water containing 0.02% by volume of a spreading agent [product name: Sindain (registered trademark), manufactured by Sumitomo Chemical Co., Ltd.] so that the concentration of etoxazole became 800 ppm.

[0066] A reference standard containing 99.9% by weight of fenazaquin was diluted with water containing 0.02% by volume of a spreading agent [product name: Sindain (registered trademark), manufactured by Sumitomo Chemical Co., Ltd.] so that the concentration of fenazaquin became 50 ppm.

[0067] The water dilution of etoxazole, the water dilution of fenazaquin, and water containing 0.02% by volume of a spreading agent [product name: Sindain (registered trademark), manufactured by Sumitomo Chemical Co., Ltd.] were mixed to prepare a test solution with a predetermined concentration of etoxazole and/or fenazaquin.

[0068] Then, each test solution was sprayed onto the plastic containers to sufficiently soak cabbage seedlings, and allowed to stand for room temperature (about 25°C). Two days after the treatment, the number of living insects (*Bemisia tabaci*) on the cabbage seedlings was determined (This is called the "number of insects after treatment").

[0069] An insect death rate was calculated according to the following equation. For each treatment there are duplicates. Results are shown in Table 3.

Insect death rate (%) = $100 - (\text{number of insects after treatment} / \text{number of insects before treatment}) \times 100$

Table 3

Etoxazole	Fenazaquin	Insect death rate (%)
Concentration (ppm)	Concentration (ppm)	
50	-	15
200	-	17
800	-	32

(continued)

Etoxazole	Fenazaquin	Insect death rate (%)
Concentration (ppm)	Concentration (ppm)	
-	3.2	11
-	50	64
50	3.2	65
200	3.2	82
800	3.2	63
50	50	100
200	50	100
800	50	100

Claims

1. A pest control composition comprising etoxazole and fenazaquin.
2. The pest control composition according to claim 1, wherein the weight ratio of etoxazole to fenazaquin is from 5000:1 to 1:5000.
3. The pest control composition according to claim 1, wherein the weight ratio of etoxazole to fenazaquin is from 100:1 to 1:100.
4. A pest control method, which comprises applying effective amounts of etoxazole and fenazaquin to a pest or an area where a pest lives, with the proviso that a method for the treatment of the human or animal body by therapy is excluded.
5. The pest control method according to claim 4, wherein the weight ratio of etoxazole to fenazaquin is from 5000:1 to 1:5000.
6. The pest control method according to claim 4, wherein the weight ratio of etoxazole to fenazaquin is from 100:1 to 1:100.
7. Use of etoxazole and fenazaquin as a pest control agent, with the proviso that a use for the treatment of the human or animal body by therapy is excluded.

Patentansprüche

1. Eine Schädlingsbekämpfungszusammensetzung, umfassend Etoxazol und Fenazaquin.
2. Die Schädlingsbekämpfungszusammensetzung gemäß Anspruch 1, wobei das Gewichtsverhältnis von Etoxazol zu Fenazaquin 5000:1 bis 1:5000 beträgt.
3. Die Schädlingsbekämpfungszusammensetzung gemäß Anspruch 1, wobei das Gewichtsverhältnis von Etoxazol zu Fenazaquin 100:1 bis 1:100 beträgt.
4. Ein Schädlingsbekämpfungsverfahren, welches das Anwenden von wirksamen Mengen an Etoxazol und Fenazaquin auf einen Schädling oder einen Ort, den ein Schädling befallen hat, umfasst, mit der Maßgabe, dass ein Verfahren zur therapeutischen Behandlung des menschlichen oder tierischen Körpers ausgeschlossen ist.
5. Das Schädlingsbekämpfungsverfahren gemäß Anspruch 4, wobei das Gewichtsverhältnis von Etoxazol zu Fenazaquin 5000:1 bis 1:5000 beträgt.

EP 2 734 045 B1

6. Das Schädlingsbekämpfungsverfahren gemäß Anspruch 4, wobei das Gewichtsverhältnis von Etoxazol zu Fenzaquin 100:1 bis 1:100 beträgt.
7. Verwendung von Etoxazol und Fenzaquin als ein Schädlingsbekämpfungsmittel, mit der Maßgabe, dass eine Verwendung zur therapeutischen Behandlung des menschlichen oder tierischen Körpers ausgeschlossen ist.

Revendications


1. Composition antiparasitaire comprenant de l'étoxazole et de la fénazaquine.
2. Composition antiparasitaire selon la revendication 1, dans laquelle le rapport pondéral de l'étoxazole à la fénazaquine est de 5 000/1 à 1/5 000.
3. Composition antiparasitaire selon la revendication 1, dans laquelle le rapport pondéral de l'étoxazole à la fénazaquine est de 100/1 à 1/100.
4. Procédé de lutte contre les parasites, qui comprend l'application de quantités efficaces d'étoxazole et de fénazaquine sur un parasite ou une zone où vit un parasite, à condition qu'un procédé de traitement du corps humain ou animal par le traitement soit exclu.
5. Procédé de lutte contre les parasites selon la revendication 4, dans lequel le rapport pondéral de l'étoxazole à la fénazaquine est de 5 000/1 à 1/5 000.
6. Procédé de lutte contre les parasites selon la revendication 4, dans lequel le rapport pondéral de l'étoxazole à la fénazaquine est de 100/1 à 1/100.
7. Utilisation d'étoxazole et de fénazaquine en tant qu'agent antiparasitaire, à condition qu'une utilisation pour le traitement du corps humain ou animal par le traitement soit exclue.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- JP 2011159711 A [0001]
- WO 9322297 A [0006] [0012]
- EP 0326329 A [0006] [0013]
- US 20100216738 A1 [0006]



Szabadalmi igénypontok

1. Kártevőirtő készítmény, amely tartalmaz etoxazolt és fenazakint.
2. Az 1. igénypont szerinti kártevőirtő készítmény, ahol az etoxazolnak fenazakinre vetített tömegaránya 5000:1 - 1:5000.
3. Az 1. igénypont szerinti kártevőirtő készítmény, ahol az etoxazolnak fenazakinre vetített tömegaránya 100:1 - 1:100.
4. Kártevőirtási eljárás, amely tartalmazza az etoxazol és a fenazakin hatásos mennyiségének az alkalmazását a kártevőre vagy a területre, ahol a kártevő él, azzal a kikötéssel, hogy az emberi vagy állati test terápiás kezelési eljárása ki van zárva.
5. A 4. igénypont szerinti kártevőirtási eljárás, ahol az etoxazolnak fenazakinre vetített tömegaránya 5000:1 - 1:5000.
6. A 4. igénypont szerinti kártevőirtási eljárás, ahol az etoxazolnak fenazakinre vetített tömegaránya 100:1 - 1:100.
7. Etoxazol és fenazakin alkalmazása kártevőirtó szerként, azzal a kikötéssel, hogy az emberi vagy állati test terápiás kezelésére történő alkalmazás kizárt.