

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 April 2001 (12.04.2001)


PCT

(10) International Publication Number
WO 01/25191 A1

- (51) International Patent Classification⁷: C07C 281/02 (CA). DEKEYSER, Mark, Achiel; 333 Boxbury Drive, Waterloo, Ontario N2K 1W3 (CA).
- (21) International Application Number: PCT/US00/26465
- (22) International Filing Date: 26 September 2000 (26.09.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 09/413,072 6 October 1999 (06.10.1999) US
- (71) Applicants: UNIROYAL CHEMICAL COMPANY, INC. [US/US]; 199 Benson Road, Middlebury, CT 06749 (US). UNIROYAL CHEMICAL CO./UNIROYAL CHEMICAL CIE. [CA/CA]; 25 Erb Street, Elmira, Ontario N3B 3A3 (CA).
- (72) Inventors: CHEE, Gaik-Lean; 3 Pondview Court, Guelph, Ontario N1E 3K2 (CA). PARK, Sheldon, Bernard; 286 Pastern Trail, Waterloo, Ontario N2K 3W7
- (74) Agent: REITENBACH, Daniel; Uniroyal Chemical Company, Inc., 199 Benson Road, Middlebury, CT 06749 (US).
- (81) Designated States (*national*): AU, BR, CA, CN, CZ, HU, IL, JP, KR, KZ, PL, RO, RU, SD, SK, TR, UA, UZ, ZA.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 4-HYDROXYBIPHENYL HYDRAZIDE DERIVATIVES

WO 01/25191 A1



(57) Abstract: Compounds having formula (I) wherein R is hydrogen or CO₂CH(CH₃)₂, useful as intermediate in the preparation of the miticide bifenazate, methods for their preparation, and methods for the preparation of bifenazate.

- 1 -

4-HYDROXYBIPHENYL HYDRAZIDE DERIVATIVES

5 Field of Invention

 This invention relates to certain 4-hydroxybiphenyl hydrazide derivatives useful as intermediates in the preparation of isopropyl-2-(4-methoxy-[1,1'-biphenyl]-3-yl)hydrazine carboxylate (bifenazate).

10 Background of The Invention

 U. S. Patent No. 5,367,093 describes a method for the preparation of the miticidal phenylhydrazine derivative, isopropyl-2-(4-methoxy-[1,1'-biphenyl]-3-yl)hydrazine carboxylate (bifenazate), using a six-step procedure which comprises the undesirable steps of preparation and reduction of a diazonium salt.


 Certain phenylhydrazine derivatives can be prepared using the methods described in U. S. Patent No. 4,864,032 (amination of Grignard); in Mitchell, J. Org. Chem. 59: 682 (1994) (amination of electron-rich arenes); and in Lenarsic, J. Org. Chem. 64: 2558 (1999) (by electrophilic azodicarboxylates).

 It is the purpose of this invention to provide new intermediates useful in the preparation of bifenazate. It is also a purpose of this invention to provide a new method for the preparation of bifenazate.

30 Summary of the Invention


 The present invention relates to compounds of the formula:

- 2 -




(I)

wherein R is hydrogen (IA) or isopropyl ester ($\text{CO}_2\text{CH}(\text{CH}_3)_2$) (IB).



(IA)



(IB)

The compounds of formulas IA and IB are useful as intermediates in the preparation of bifenazate.

The present invention also relates to a method for preparing the compound of IA which comprises hydrolyzing

- 3 -


the compound of IB in the presence of an effective amount of a base in a suitable organic solvent.

The present invention further relates to a method for preparing bifenazate comprising methylating Compound IA in the presence of an effective amount of a methylating agent and a base, in a suitable solvent.

Detailed Description of the Invention

The compounds of this invention can be prepared as described below in SCHEME 1.

1. Amination




4-Hydroxybiphenyl is reacted with di-isopropyl azodicarboxylate in the presence of an effective amount of Lewis acid in a suitable solvent, to produce Compound IB. Useful Lewis acids include, e.g., boron trifluoride etherate and aluminum chloride. The concentration of the Lewis acid to the 4-hydroxybiphenyl in the reaction mixture can be between about 1:0.2 to about 1:1.1 (mol/mol), preferably about 1:1.1 (mol/mol). Suitable organic solvents are those organic solvents which are not

- 4 -

deleterious to the amination reaction and include, e.g., ethyl acetate, dichloromethane, toluene, and glyme. The temperature of the reaction mixture should be from about 0°C to about 60°C, preferably at room temperature.

5

2. Selective Hydrolysis



Compound IB is treated with an effective amount of a base in a suitable organic solvent to produce Compound IA. Useful base compounds are those base compounds which can hydrolyze the ester functionality and include, e.g., sodium hydroxide and potassium hydroxide. The concentration of the base to Compound IB can be between about 7:1 to about 10:1 (mol/mol), preferably about 9:1 (mol/mol). Suitable organic solvents are those organic solvents which are not deleterious to the hydrolysis reaction and include, e.g., toluene, dimethyl sulfoxide, and glyme. Preferably, the temperature of the organic solvent should be above room temperature and below 110°C.


20

Compound IA is then methylated in the presence of an effective amount of a methylating agent and a base, in a suitable organic solvent. For the purpose of this

- 5 -

invention, a "methylating agent" is any compound which is capable of substituting a methyl group for the hydrogen atom in the 4-hydroxy group in Compound IA. Useful methylating agents include, e.g., dimethylsulfate and methyl iodide. The concentration of methylating agent to Compound IA can be between about 1:1 to about 1:1.2 (mol/mol), preferably about 1:1 (mol/mol). Useful base compounds are those base compounds which can deprotonate phenols and include, e.g., sodium carbonate and potassium carbonate. The concentration of base to Compound IA can be between about 1:1 to about 3:1 (mol/mol), preferably about 2:1 (mol/mol). Suitable organic solvents are those organic solvents which are not deleterious to the methylation reaction and include, e.g., toluene and acetone. The methylation reaction can be conducted at about room temperature. The methylation process is exemplified below in Scheme 2.

SCHEME 2



The following examples are provided to illustrate the present invention.

- 6 -

Example 1

Preparation of 1,2-Hydrazinedicarboxylic acid, 1-(4-hydroxy-[1,1'-biphenyl]-3-yl)-, bis(1-methylethyl) ester (Compound IB)

5

To a solution of 4-hydroxybiphenyl (5.50 g) in ethyl acetate (60 mL) at room temperature was added boron trifluoride etherate (4.1 mL). The resultant mixture was cooled to -5°C, added dropwise di-isopropyl azodicarboxylate (6.3 mL), and stirred at that temperature for 30 min. and then stirred at room temperature for two hours. The mixture was then quenched with water (100 mL) and extracted with ethyl acetate (50 mL). The organic phase was separated, dried over magnesium sulfate, and concentrated under vacuum to leave an oil which was chromatographed on silica gel using 20-30% ethyl acetate/hexane to produce Compound IB as beige-colored solid (10.65 g, 88% yield). ¹H-NMR (ppm, in CDCl₃): m(12)1.30; m(2)5.04; m(2)7.10; m(1)7.32; dd(2)7.43; m(3)7.51-7.55; br s(1)8.53.

10

15

20

Example 2

Preparation of hydrazinecarboxylic acid, 2-(4-hydroxy-[1,1'-biphenyl]-3-yl)-, 1-methylethyl ester (Compound IA)

25

Potassium hydroxide (5.0 g) was added to a stirred suspension of Compound IB produced above in Example 1 (5.15 g) in toluene (50 mL). The resultant purple mixture was bubbled with nitrogen for 20 min, and then heated at 45°C for 4 days. The mixture was then cooled to 0°C and then 6M HCl was added to the mixture until the pH of the mixture was about 1. The mixture was then extracted with ethyl acetate. The organic phase was separated, washed with brine, dried over magnesium sulfate, and concentrated to a brown solid. Recrystallization from toluene produced Compound IA as

30

35

- 7 -

beige-colored powder (3.35 g, 85% yield). ¹H-NMR (ppm, in CDCl₃): d(6)1.81; septet(1)5.04; br d(1)5.91; br s(1)6.61; d(1)6.74; dd(1)7.01; d(1)7.14; dd(1)7.32; dd(2)7.42; dd(2)7.61.

5

Example 3

Preparation of hydrazinecarboxylic acid, 2-(4-methoxy-[1,1'-biphenyl]-3-yl)-, 1-methylethyl ester (bifenazate)

10

A suspension of Compound IA prepared above in Example 2 (2.63 g) and potassium carbonate (2.50 g) in acetone (40 mL) was bubbled with nitrogen for 20 minutes. Dimethyl sulfate (0.96 mL) at room temperature was then added to the suspension. After 2 hours, the resultant reaction mixture was cooled in ice bath. 2M HCl was then cautiously added (ca. 30 mL) to the reaction mixture until the pH of the reaction mixture was about 1. The reaction mixture was then concentrated to remove most acetone. The solid formed from the concentrated reaction mixture was filtered, washed with water, hexane, and dried under air with suction to produce bifenazate (2.60 g). ¹H-NMR spectral data is consistent with the ¹H-NMR reported for bifenazate in U. S. Patent No. 5,367,093.

25


- 8 -

What is claimed is:

1. A compound having the formula:

5

10



(I)


15

wherein R is hydrogen or CO₂CH(CH₃)₂.

20

2. A compound as recited in claim 1 having the formula:


25




(IA)

30

3. A compound as recited in claim 1 having the formula:



4. A method for preparing a compound of the formula



which method comprises reacting 4-hydroxybiphenyl with di-isopropyl azodicarboxylate in the presence of an effective amount of a Lewis acid, in a suitable solvent.

- 10 -


5 5. A method as recited in claim 4 wherein the concentration of the Lewis acid to the 4-hydroxybiphenyl in the reaction mixture is between about 1:0.2 to about 1:1.1 (mol/mol).

10 6. A method as recited in claim 5 wherein the concentration of the Lewis acid to the 4-hydroxybiphenyl in the reaction mixture is about 1:1.1 (mol/mol).

7. A method as recited in claim 5 wherein the Lewis acid is selected from the group consisting of boron trifluoride etherate, and aluminum chloride.

15 8. A method as recited in claim 4 wherein the organic solvent is selected from the group consisting of ethyl acetate, dichloromethane, toluene, glyme, and diethyl ether.


20 9. A method for preparing a compound of the formula:



30 (IA)

which comprises hydrolyzing a compound of the formula:

- 11 -



(IB)

in the presence of an effective amount of base and a suitable organic solvent.

10. A method as recited in claim 9 wherein the base is selected from the group consisting of sodium hydroxide and potassium hydroxide.

11. A method as recited in claim 9 wherein the organic solvent is selected from the group consisting of toluene, dimethyl sulfoxide, and glyme.

12. A method as recited in claim 9 conducted at a temperature between room temperature and 110°C.


13. A method as recited in claim 9 wherein the concentration of base to Compound IB is about 7:1 to about 10:1 (mol/mol).

14. A method as recited in claim 13 wherein the concentration of base to Compound IB is about 7:1 (mol/mol).


15. A process for the preparation of a compound of

- 12 -

the formula



which process comprises methylating a compound of the formula



(IA)

in the presence of an effective amount of a methylating agent and a base, and a suitable organic solvent.

16. A method as recited in claim 15 wherein the methylating agent is selected from the group consisting of dimethylsulfate and methyl iodide.

17. A method as recited in claim 15 wherein the base is selected from the group consisting of sodium carbonate and potassium carbonate.

18. A method as recited in claim 15 wherein the

- 13 -

organic solvent is selected from the group consisting of toluene and acetone.

19. A method as recited in claim 15 conducted at about room temperature.


20. A method as recited in claim 15 wherein the concentration of methylating agent to Compound IA is about 1:1 to about 1.2:1 (mol/mol).

21. A method as recited in claim 20 wherein the concentration of methylating agent to Compound IA is between about 1:1 (mol/mol).

22. A method as recited in claim 15 wherein the concentration of base to Compound IA is about 1:1 to about 3:1 (mol/mol).

23. A method as recited in claim 22 wherein the concentration of base to Compound IA is about 2:1 (w/w).

24. A process for the preparation of a compound of the formula




(I)

which process comprises:


- 14 -

(a) reacting 4-hydroxybiphenyl with di-isopropyl azodicarboxylate in the presence of an effective amount of Lewis acid, in a suitable solvent, to produce a compound of the formula:



(IB)

(b) hydrolyzing the compound of the formula IB in the presence of an effective amount of base and a suitable organic solvent, to produce a compound of the formula



(IA)

and


(c) methylating the compound of formula IA in the presence of an effective amount of a methylating agent and

- 15 -

a base, in a suitable organic solvent,
to produce the compound of the formula

5

10



(I).

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/26465

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C281/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 367 093 A (MARK A. DEKEYSER ET AL.) 22 November 1994 (1994-11-22) cited in the application column 2, line 32 -column 6, line 68; example 135; table 4 ---	1,15,24
A	YVES LEBLANC ET AL.: "Para-Directed Amination of Electron-Rich Arenes with Bis(2,2,2-trichloroethyl)Azodicarboxylate" JOURNAL OF ORGANIC CHEMISTRY., vol. 60, 1995, pages 4268-4271, XP002155948 AMERICAN CHEMICAL SOCIETY. EASTON., US ISSN: 0022-3263 page 4268, column 1, line 32 -page 4270, column 2, line 4 --- -/--	1,15,24



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

19 December 2000

Date of mailing of the international search report

11/01/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Zervas, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/26465

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 864 032 A (JAMES P. DEMERS) 5 September 1989 (1989-09-05) cited in the application column 1 -column 2; claims; examples -----	1,15,24

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/26465

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5367093 A	22-11-1994	BG 98785 A NO 941876 A RU 2109730 C SK 60094 A US 5438123 A US 5536746 A AT 148104 T AU 670927 B AU 3137493 A BR 9206803 A CA 2123885 A CN 1075952 A, B CZ 9401248 A DE 69217085 D DE 69217085 T DK 641316 T EP 0641316 A ES 2097372 T FI 942355 A GR 3023097 T HU 68639 A IL 103828 A JP 2552811 B JP 7502267 T KR 235246 B PL 171968 B RO 112860 B WO 9310083 A ZA 9208915 A	31-05-1995 29-06-1994 27-04-1998 10-05-1995 01-08-1995 16-07-1996 15-02-1997 08-08-1996 15-06-1993 02-05-1995 27-05-1993 08-09-1993 12-07-1995 06-03-1997 28-05-1997 11-08-1997 08-03-1995 01-04-1997 20-07-1994 30-07-1997 28-07-1995 10-01-1997 13-11-1996 09-03-1995 15-12-1999 31-07-1997 30-01-1998 27-05-1993 06-08-1993
US 4864032 A	05-09-1989	NONE	