



HU000024912T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 024 912**(13) **T2****EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 06 737611**(22) A bejelentés napja: **2006. 03. 06.**(51) Int. Cl.: **C07D 233/61** (2006.01)**C07D 233/64** (2006.01)**C07D 401/04** (2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20060737611

(97) Az európai bejelentés közzétételi adatai:

EP 1858850 A2 **2006. 09. 21.**

(86) A nemzetközi (PCT) bejelentési szám:

PCT/US 06/008450

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 1858850 B1 **2015. 01. 21.**

(87) A nemzetközi közzétételi szám:

WO 06099060

(30) Elsőbbségi adatok:

661784 P**2005. 03. 14.****US**

(73) Jogosult(ak):

Janssen Pharmaceutica NV, 2340 Beerse (BE)

(72) Feltaláló(k):

CHAOZHONG, Cai, North Wales, PA 19403 (US)**HE, Wei, Audubon, PA 19403 (US)**

(74) Képviselő:

**dr. Fehérvári Flóra, DANUBIA Szabadalmi és
Jogi Iroda Kft., Budapest**

(54)

Eljárás opioid modulátorok előállítására

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.

(19)



(11)

EP 1 858 850 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
21.01.2015 Bulletin 2015/04

(51) Int Cl.:
C07D 233/61 *(2006.01)* **C07D 233/64** *(2006.01)*
C07D 401/04 *(2006.01)*

(21) Application number: **06737611.1**

(86) International application number:
PCT/US2006/008450

(22) Date of filing: **06.03.2006**

(87) International publication number:
WO 2006/099060 (21.09.2006 Gazette 2006/38)

(54) **PROCESS FOR THE PREPARATION OF OPIOID MODULATORS**

HERSTELLUNGSVERFAHREN FÜR OPIOIDMODULATOREN

PROCEDE POUR PREPARER DES MODULATEURS OPIOIDES

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI
SK TR**
Designated Extension States:
AL BA HR MK YU

(30) Priority: **14.03.2005 US 661784 P**

(43) Date of publication of application:
28.11.2007 Bulletin 2007/48

(73) Proprietor: **Janssen Pharmaceutica NV**
2340 Beerse (BE)

(72) Inventors:
• **CHAOZHONG, Cai**
North Wales, PA 19403 (US)
• **HE, Wei**
Audubon, PA 19403 (US)

(74) Representative: **Warner, James Alexander et al**
Carpmaels & Ransford LLP
One Southampton Row
London WC1B 5HA (GB)

(56) References cited:
WO-A-03/092688 WO-A-2005/090315

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 858 850 B1


Description

FIELD OF THE INVENTION

[0001] The present invention is directed to a novel process for the preparation of opioid modulators (agonists and antagonists), and intermediates in their synthesis. The opioid modulators are useful in the treatment and prevention of such disorders as pain, visceral pain including post-operative pain, gastrointestinal disorders including diarrheic syndromes, motility disorders including post-operative ileus, constipation, irritable bowel syndrome and inflammatory bowel disorders.


BACKGROUND OF THE INVENTION

[0002] The present invention is directed to the preparation of novel opioid receptor modulators and intermediates in their synthesis. More specifically, the present invention is directed to novel processes for the preparation of compounds of formula (II)



wherein all variables are as hereinafter defined, disclosed in United States Patent Application No. 11/079,647, filed March 15, 2004, and published as US Patent Publication US-2005-0203143-A1, September 15, 2005, which is hereby incorporated by reference in its entirety.


[0003] WO 03/092688 discloses opioid receptor modulators represented by the general formula:



[0004] Known methods for the preparation of the compounds of formula (II) and compounds of formula (I), as herein defined, require the use of dimethyl-tyrosine, which is expensive and thus not suitable for large scale synthesis. Thus there remains a need for a process for the preparation of compounds of formula (I) and compounds of formula (II) which is suitable for large scale production.

SUMMARY OF THE INVENTION

[0005] The present invention is directed to a process for the preparation of compounds of formula (I)



wherein




is C₆₋₁₀aryl or a heteroaryl selected from the group consisting of furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, indolinyl, benzofuryl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinoliziny, quinoliny, isoquinoliny and quinazoliny;

each R^{41P} is independently selected from C₁₋₆alkyl, C₁₋₆alkoxy or fluoro;



R^J and R^K are each independently selected from hydrogen or C₁₋₄alkyl; alternatively, R^J and R^K are taken together with the nitrogen atom to which they are bound to form a five to seven membered heterocyclyl;

Pg¹ is a nitrogen protecting group;

comprising




reacting a compound of formula (X), wherein X^P is selected from CN, -CO₂H, -C(O)-Cl or -C(O)-OC₁₋₄alkyl and wherein Y^P is selected from Br, Cl or I, to yield the corresponding compound of formula (XII);




25

reacting the compound of formula (XII) with a suitably substituted compound of formula (XVIII); in the presence of palladium catalyst; in the presence of an organic or inorganic base; in an organic solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XIX);



35

reacting the compound of formula (XIX) with hydrogen or a source of hydrogen; in the presence of a catalyst; in a solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XX);




50

reacting the compound of formula (XX) with an aqueous base; in an organic solvent; to yield the corresponding compound of formula (I).

55

[0006] The present invention is further directed to a process for the preparation of a compound of formula (I)



wherein




is C₆₋₁₀aryl or a heteroaryl selected from the group consisting of furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, indolyl, benzofuryl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinoliziny, quinoliny, isoquinoliny and quinazoliny;


each R^{41P} is independently selected from C₁₋₆alkyl, C₁₋₆alkoxy or fluoro;

R^J and R^K are each independently selected from hydrogen or C₁₋₄alkyl; alternatively, R^J and R^K are taken together with the nitrogen atom to which they are bound to form a five to seven membered heterocyclyl;

Pg¹ is a nitrogen protecting group; comprising




reacting the compound of formula (XIX) with hydrogen or a source of hydrogen; in the presence of a catalyst; in a solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XX);




reacting the compound of formula (XX) with an aqueous base; in an organic solvent; to yield the corresponding compound of formula (I).


[0007] The present invention is further directed to a process for the preparation of a compound of formula (Ia) (also known as, 4-(aminocarbonyl)-N-[(1,1-dimethylethoxy)carbonyl]-2,6-dimethyl-L-phenylalanine)




comprising




reacting a compound of formula (Xa), wherein X^P is selected from CN, -CO₂H, -C(O)-Cl or -C(O)-OC₁₋₄alkyl and wherein Y^P is selected from Br, Cl or I, to yield the corresponding compound of formula (XIIa);



reacting the compound of formula (XIIa) with a suitably substituted compound of formula (XVIIIa); in the presence of palladium catalyst; in the presence of an organic or inorganic base; in an organic solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XIXa);




reacting compound of formula (XIXa) with hydrogen gas, at a pressure sufficient to hydrogenate; in the presence of a suitable chiral catalyst; at a temperature greater than about room temperature; in an organic solvent; to yield the corresponding compound of formula (XXa);




reacting the compound of formula (XXa) with an aqueous base; in an organic solvent; to yield the corresponding compound of formula (Ia).


[0008] The present invention is further directed to a process for the preparation of the compound of formula (Ia)



comprising




reacting compound of formula (XIXa) with hydrogen gas, at a pressure sufficient to hydrogenate; in the presence of a suitable chiral catalyst; at a temperature greater than about room temperature; in an organic solvent; to yield the corresponding compound of formula (XXa);




reacting the compound of formula (XXa) with an aqueous base; in an organic solvent; to yield the corresponding compound of formula (Ia).


[0009] The present invention is further directed to processes for the preparation of compounds of formula (XIX)



[0010] The present invention is further directed to processes for the preparation of the compound of formula (XIXb)



[0011] The present invention is further directed to a process for the preparation of compounds of formula (II)



wherein



is C₆₋₁₀aryl or a heteroaryl selected from the group consisting of furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, indolinyl, benzofuryl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinoliziny, quinoliny, isoquinoliny and quinazoliny;

each R^{41P} is independently selected from C₁₋₆alkyl, C₁₋₆alkoxy or fluoro;

R^J and R^K are each independently selected from hydrogen or C₁₋₄alkyl; alternatively, R^J and R^K are taken together with the nitrogen atom to which they are bound to form a five to seven membered heterocycl;

R¹ is selected from the group consisting of hydrogen, C₁₋₆alkyl, cycloalkyl, heterocycl, aryl(C₁₋₆)alkyl, and heteroaryl(C₁₋₆)alkyl; wherein when R¹ is phenyl(C₁₋₆)alkyl, phenyl is optionally fused to a heterocycl or cycloalkyl; wherein when R¹ is C₁₋₂alkyl, said C₁₋₂alkyl is optionally substituted with one to two substituents independently

selected from the group consisting of C₁₋₆alkoxy, aryl, cycloalkyl, heterocyclyl, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluoromethyl, and carboxy;

and further, wherein when R¹ is C₃₋₆alkyl, said C₃₋₆alkyl is optionally substituted with one to three substituents independently selected from the group consisting of C₁₋₆alkoxy, aryl, cycloalkyl, heterocyclyl, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluoromethyl, and carboxy;

wherein the cycloalkyl and heterocyclyl of C₁₋₂alkyl and C₃₋₆alkyl are optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₆alkyl, hydroxy(C₁₋₆)alkyl, C₁₋₆alkoxy, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluoromethyl, carboxy, aryl(C₁₋₆)alkoxycarbonyl, C₁₋₆alkoxycarbonyl, aminocarbonyl, C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl, and aminosulfonyl;

furthermore, wherein the cycloalkyl and heterocyclyl of R¹ are optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₆alkyl, hydroxy(C₁₋₆)alkyl, C₁₋₆alkoxy, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluoromethyl, carboxy, aryl(C₁₋₆)alkoxycarbonyl, C₁₋₆alkoxycarbonyl, aminocarbonyl, C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl, and aminosulfonyl;


furthermore, wherein the aryl and heteroaryl portion of the R¹ substituents aryl(C₁₋₆)alkyl and heteroaryl(C₁₋₆)alkyl, are optionally substituted with one to three R¹¹ substituents independently selected from the group consisting of C₁₋₆alkyl; hydroxy(C₁₋₆)alkyl; C₁₋₆alkoxy; C₆₋₁₀aryl(C₁₋₆)alkyl; C₆₋₁₀aryl(C₁₋₆)alkoxy; C₆₋₁₀aryl; heteroaryl optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₄alkyl, C₁₋₄alkoxy, and carboxy; cycloalkyl; heterocyclyl; C₆₋₁₀aryloxy; heteroaryloxy; cycloalkyloxy; heterocyclyloxy; amino; C₁₋₆alkylamino; (C₁₋₆alkyl)₂amino; C₃₋₆cycloalkylaminocarbonyl; hydroxy(C₁₋₆)alkylaminocarbonyl; C₆₋₁₀arylaminocarbonyl wherein C₆₋₁₀aryl is optionally substituted with carboxy or C₁₋₄alkoxycarbonyl; heterocyclyl-carbonyl; carboxy; C₁₋₆alkylcarbonyloxy; C₁₋₆alkoxycarbonyl; C₁₋₆alkylcarbonyl; C₁₋₆alkylcarbonylamino; aminocarbonyl; C₁₋₆alkylaminocarbonyl; (C₁₋₆alkyl)₂aminocarbonyl; cyano; halogen; trifluoromethyl; trifluoromethoxy; and hydroxy;

provided that no more than one R¹¹ substituent is selected from the group consisting of C₆₋₁₀aryl(C₁₋₆)alkyl; C₆₋₁₀aryl(C₁₋₆)alkoxy; C₆₋₁₀aryl; heteroaryl optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₄alkyl, C₁₋₄alkoxy, and carboxy; cycloalkyl; heterocyclyl; C₆₋₁₀aryloxy; heteroaryloxy; cycloalkyloxy; C₆₋₁₀arylaminocarbonyl, heterocyclylcarbonyl; and heterocyclyloxy;

R² is hydrogen, C₁₋₈alkyl, hydroxy(C₁₋₈)alkyl, C₆₋₁₀aryl(C₁₋₆)alkoxy(C₁₋₆)alkyl, or C₆₋₁₀aryl(C₁₋₈)alkyl;

wherein the C₆₋₁₀aryl group in the C₆₋₁₀aryl-containing substituents of R² is optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₆alkyl, C₁₋₆alkoxy, hydroxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, aminocarbonyl, C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl, cyano, fluoro, chloro, bromo, trifluoromethyl, and trifluoromethoxy; and, wherein the C₁₋₆alkyl and C₁₋₆alkoxy substituents of aryl are optionally substituted with hydroxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, or C₆₋₁₀aryl;

A is selected from the group consisting of aryl, ring system **a-1**, **a-2**, **a-3**, and **a-4**, optionally substituted with R³ and R⁵;



wherein A-B is selected from the group consisting of N-C, C-N, N-N and C-C; wherein D-E is selected from the group consisting of O-C, S-C, and O-N; and wherein F-G is selected from the group consisting of N-O and C-O;


R³ is one to two substituents independently selected from the group consisting of C₁₋₆alkyl, aryl, aryl(C₁₋₆)alkyl, aryl(C₂₋₆)alkenyl, aryl(C₂₋₆)alkynyl, heteroaryl, heteroaryl(C₁₋₆)alkyl, heteroaryl(C₂₋₆)alkenyl, heteroaryl(C₂₋₆)alkynyl, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, arylamino, heteroarylamino, aryloxy, heteroaryloxy, trifluoromethyl, and halogen;

wherein the aryl, heteroaryl, and the aryl and heteroaryl of aryl(C₁₋₆)alkyl, aryl(C₂₋₆)alkenyl, aryl(C₂₋₆)alkynyl, heteroaryl(C₁₋₆)alkyl, heteroaryl(C₂₋₆)alkenyl, heteroaryl(C₂₋₆)alkynyl, arylamino, heteroarylamino, aryloxy, and heteroaryloxy, are optionally substituted with one to five fluoro substituents or one to three substituents independently selected from the group consisting of C₁₋₆alkyl, hydroxy(C₁₋₆)alkyl, C₁₋₆alkoxy, C₆₋₁₀aryl(C₁₋₆)alkyl,


C₆₋₁₀aryl(C₁₋₆)alkoxy, C₆₋₁₀aryl, C₆₋₁₀aryloxy, heteroaryl(C₁₋₆)alkyl, heteroaryl(C₁₋₆)alkoxy, heteroaryl, heteroaryloxy, C₆₋₁₀arylamino, heteroarylamino, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carboxy(C₁₋₆)alkylamino, carboxy, C₁₋₆alkylcarbonyl, C₁₋₆alkoxycarbonyl, C₁₋₆alkylcarbonylamino, aminocarbonyl, C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl, carboxy(C₁₋₆)alkylaminocarbonyl, cyano, halogen, trifluoromethyl, trifluoromethoxy, hydroxy, C₁₋₆alkylsulfonyl, and C₁₋₆alkylsulfonylamino; provided that not more than one such substituent on aryl and heteroaryl portion of R³ is selected from the group consisting of C₆₋₁₀aryl, heteroaryl, C₆₋₁₀aryl(C₁₋₆)alkyl, C₆₋₁₀aryl(C₆₋₁₀)alkoxy, aryl(C₂₋₆)alkenyl, aryl(C₂₋₆)alkynyl, heteroaryl, heteroaryl(C₁₋₆)alkyl, heteroaryl(C₂₋₆)alkoxy, C₆₋₁₀arylamino, heteroarylamino, C₆₋₁₀aryloxy, and heteroaryloxy; and wherein C₁₋₆alkyl, and C₁₋₆alkyl of aryl(C₁₋₆)alkyl and heteroaryl(C₁₋₆)alkyl, are optionally substituted with a substituent selected from the group consisting of hydroxy, carboxy, C₁₋₄alkoxycarbonyl, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, aminocarbonyl, (C₁₋₄)alkylaminocarbonyl, di(C₁₋₄)alkylaminocarbonyl, aryl, heteroaryl, arylamino, heteroarylamino, aryloxy, heteroaryloxy, aryl(C₁₋₄)alkoxy, and heteroaryl(C₁₋₄)alkoxy;

R⁵ is a substituent on a nitrogen atom of ring A selected from the group consisting of hydrogen and C₁₋₄alkyl;

R^a and R^b are independently selected from the group consisting of hydrogen, C₁₋₆alkyl, and C₁₋₆alkoxycarbonyl; alternatively, when R^a and R^b are each other than hydrogen, R^a and R^b are optionally taken together with the nitrogen atom to which they are both attached to form a five to eight membered monocyclic ring; and pharmaceutically acceptable enantiomers, diastereomers, racemates, and salts thereof; comprising




reacting a compound of formula (X), wherein X^P is selected from CN, -CO₂H, -C(O)-Cl or -C(O)-OC₁₋₄alkyl and wherein Y^P is selected from Br, Cl or I, to yield the corresponding compound of formula (XII);




reacting the compound of formula (XII) with a suitably substituted compound of formula (XVIII) wherein Pg¹ is a nitrogen protecting group; in the presence of palladium catalyst; in the presence of an organic or inorganic base; in an organic solvent; at a temperature greater than about room temperature; to yield the corresponding compound


of formula (XIX);



reacting the compound of formula (XIX) with hydrogen or a source of hydrogen; in the presence of a catalyst; in a solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XX);




reacting the compound of formula (XX) with an aqueous base; in an organic solvent; to yield the corresponding compound of formula (I);




reacting the compound of formula (I), to yield the corresponding compound of formula (II).

[0012] In an embodiment, the present invention is directed to processes for the preparation of the compound of formula (IV)



also known as, 5-({[2-amino-3-(4-carbamoyl-2,5-dimethyl-phenyl)-propionyl]-[1-(4-phenyl-1H-imidazol-2-yl)-ethyl]-amino}-methyl)-2-methoxy-benzoic acid.

[0013] In an embodiment, the present invention is directed to processes for the preparation of the compound of formula (V)




also known as, 5-[[[2-Amino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionyl]-{1-[4-(4-iodo-phenyl)-1H-imidazol-2-yl]-ethyl}-amino]-methyl]-2-methoxy-benzoic acid.

[0014] The present invention is further directed to a product prepared according to any of the processes described herein.

DETAILED DESCRIPTION OF THE INVENTION

[0015] The present invention is directed to processes for the preparation of compounds of formula (I)



wherein Pg^1 ,



R^J , R^K and R^{41P} are as herein defined. The compounds of formula (I) are useful in the preparation of opioid receptor modulators - compounds of formula (II) as defined herein. The present invention is further directed to processes for the preparation of the compound of formula (Ia) as herein defined, useful as intermediates in the synthesis of opioid receptor modulators.

[0016] In an embodiment, the present invention is directed to processes for the preparation of compounds wherein the



ring is unsubstituted. In an embodiment of the present invention, the




ring is substituted with one R^{41P} group, which is bound at the 2- or 6-position. In another embodiment, the present invention is directed to processes for the preparation of compounds wherein the




ring is substituted with two R^{41P} groups, which are bound at the 2- and 6-positions. For example, processes wherein



is phenyl, the compound of formula (I) is of the following structure:




[0017] In an embodiment, the present invention is directed to processes for the preparation of compounds of formula (lc)



wherein R^{41Q} is selected from methyl, ethyl, methoxy, ethoxy or fluoro and wherein R^J , R^K and Pg^1 are as herein defined.

[0018] In another embodiment, the present invention is directed to processes for the preparation of the compound of formula (lb)




(i.e. a compound of formula (I) wherein




is phenyl; R^J and R^K are each hydrogen; the phenyl ring is further substituted with two R^{41P} groups, which are each methyl and Pg^1 is Boc), also known as 2-tert-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionic acid.


[0019] In another embodiment, the present invention is directed to processes for the preparation of the compound of formula (la)




[0020] The present invention is further directed to processes for the preparation of compounds of formula (XIX)



[0021] In an embodiment, the present invention is directed to processes for the preparation of the compound of formula (XIXb),




[0022] Preferably, the present invention is directed to processes for the preparation of the compound of formula (XIXa)



[0023] The compounds of formula (XIX) are useful as intermediates in the synthesis of compounds of formula (II).


[0024] The present invention is further directed to processes for the preparation of compound of formula (II)



wherein



R^J , R^K , R^{41P} , R^a , R^b , R^1 , R^2 , R^3 , R^5 , and



are as herein defined. The compounds are opioid receptor modulators, useful in the treatment of disorders mediated by at least one opioid receptor (preferably δ or μ opioid receptor), including, but not limited to pain and gastrointestinal disorders.

[0025] Embodiments of the present invention include processes for the preparation of compounds wherein R^1 is selected from the group consisting of hydrogen, C_{1-6} alkyl, aryl(C_{1-4})alkyl, and heteroaryl(C_{1-4})alkyl; wherein the aryl and heteroaryl portion of aryl(C_{1-4})alkyl and heteroaryl(C_{1-4})alkyl are optionally substituted with one to three R^{11} substituents independently selected from the group consisting of C_{1-6} alkoxy; heteroaryl optionally substituted with one to two substituents independently selected from the group consisting of C_{1-4} alkyl, C_{1-4} alkoxy, and carboxy; carboxy; C_{1-4} alkoxycarbonyl; C_{1-4} alkoxycarbonyloxy; aminocarbonyl; C_{1-4} alkylaminocarbonyl; C_{3-6} cycloalkylaminocarbonyl; hydroxy(C_{1-6})alkylaminocarbonyl; C_{6-10} arylaminocarbonyl wherein C_{6-10} aryl is optionally substituted with carboxy or C_{1-4} alkoxycarbonyl; heterocyclylcarbonyl; cyano; halogen; trifluoromethoxy; and hydroxy; provided that no more than one R^{11} is heteroaryl (optionally substituted with one to two C_{1-4} alkyl substituents); C_{6-10} arylaminocarbonyl wherein C_{6-10} aryl is optionally substituted with carboxy or C_{1-4} alkoxycarbonyl; or heterocyclylcarbonyl.

[0026] Embodiments of the present invention further include processes for the preparation of compounds wherein R^1 is selected from the group consisting of C_{6-10} aryl(C_{1-4})alkyl, pyridinyl(C_{1-4})alkyl, and furanyl(C_{1-4})alkyl; wherein C_{6-10} aryl, pyridinyl, and furanyl are optionally substituted with one to three R^{11} substituents independently selected from the group consisting of C_{1-3} alkoxy; tetrazolyl; carboxy; C_{1-4} alkoxycarbonyl; aminocarbonyl; C_{1-4} alkylaminocarbonyl; C_{3-6} cycloalkylaminocarbonyl; hydroxy(C_{1-4})alkylaminocarbonyl; C_{6-10} arylaminocarbonyl wherein C_{6-10} aryl is optionally substituted with carboxy or C_{1-4} alkoxycarbonyl; morpholin-4-ylcarbonyl; cyano; halogen; and trifluoromethoxy; provided that that no more than one R^{11} is C_{6-10} arylaminocarbonyl.

[0027] Embodiments of the present invention further include processes for the preparation of compounds wherein R^1 is selected from the group consisting of phenyl(C_{1-3})alkyl, pyridinyl(C_{1-3})alkyl, and furanyl(C_{1-3})alkyl; wherein phenyl, pyridinyl, and furanyl are optionally substituted with one to three R^{11} substituents independently selected from the group consisting of C_{1-3} alkoxy; tetrazolyl, C_{3-6} cycloalkylaminocarbonyl; hydroxy(C_{1-4})alkylaminocarbonyl; C_{6-10} arylaminocarbonyl wherein C_{6-10} aryl is optionally substituted with carboxy or C_{1-4} alkoxycarbonyl; morpholin-4-ylcarbonyl; chloro; fluoro; trifluoromethoxy; C_{1-4} alkoxycarbonyl; and carboxy; provided that that no more than one R^{11} is C_{6-10} arylaminocarbonyl.

[0028] Embodiments of the present invention further include processes for the preparation of compounds wherein R^1 is phenylmethyl, pyridinylmethyl, or furanylmethyl; wherein phenyl, pyridinyl, and furanyl are optionally substituted with one to three R^{11} substituents independently selected from the group consisting of methoxy; tetrazolyl; cyclopropylaminocarbonyl; (2-hydroxyethyl)aminocarbonyl; methoxycarbonyl; phenylaminocarbonyl wherein phenyl is optionally substituted with carboxy; morpholin-4-ylcarbonyl; and carboxy; provided that that no more than one R^{11} is phenylaminocarbonyl.

[0029] Embodiments of the present invention include processes for the preparation of compounds wherein R^2 is a substituent selected from the group consisting of hydrogen, C_{1-4} alkyl, hydroxy(C_{1-4})alkyl, and phenyl(C_{1-6})alkoxy(C_{1-4})alkyl; wherein said phenyl is optionally substituted with one to two substituents independently selected from the group consisting of C_{1-3} alkyl, C_{1-3} alkoxy, hydroxy, cyano, fluoro, chloro, bromo, trifluoromethyl, and trifluoromethoxy.

[0030] Embodiments of the present invention further include processes for the preparation of compounds wherein R^2 is selected from the group consisting of hydrogen and C_{1-4} alkyl. Embodiments of the present invention further include processes for the preparation of those compounds wherein R^2 is hydrogen or methyl.

[0031] Embodiments of the present invention include processes for the preparation of compounds wherein ring A is a-1. Embodiments of the present invention further include processes for the preparation of compounds wherein A-B of ring a-1 is N-C.

[0032] Embodiments of the present invention include processes for the preparation of compounds wherein R^3 is one to two substituents independently selected from the group consisting of C_{1-6} alkyl, halogen, and aryl; wherein aryl is

optionally substituted with one to three substituents independently selected from the group consisting of halogen, carboxy, aminocarbonyl, C₁₋₃alkylsulfonylamino, cyano, hydroxy, amino, C₁₋₃alkylamino, and (C₁₋₃alkyl)₂amino.

[0033] Embodiments of the present invention further include processes for the preparation of compounds wherein R³ is one to two substituents independently selected from the group consisting of C₁₋₃alkyl, bromo, and phenyl; wherein phenyl is optionally substituted with one to three substituents independently selected from the group consisting of chloro, fluoro, carboxy, aminocarbonyl, and cyano.

[0034] Embodiments of the present invention further include processes for the preparation of compounds wherein R³ is one to two substituents independently selected from the group consisting of methyl and phenyl; wherein phenyl is optionally substituted with one to three substituents independently selected from the group consisting of chloro and carboxy.

[0035] Embodiments of the present invention further include processes for the preparation of compounds wherein at least one R³ substituent is phenyl.

[0036] Embodiments of the present invention further include processes for the preparation of compounds wherein R³ is a substituent selected from the group consisting of methyl and phenyl; wherein phenyl is optionally substituted with one to two substituents independently selected from the group consisting of chloro and carboxy.

[0037] Embodiments of the present invention include processes for the preparation of compounds wherein



is C₆₋₁₀aryl. Embodiments of the present invention further include processes for the preparation of compounds wherein




is phenyl.

[0038] Embodiments of the present invention include processes for the preparation of compounds wherein R^{41P} is selected from C₁₋₃alkyl, C₁₋₆alkoxy or fluoro. Embodiments of the present invention further include processes for the preparation of compounds wherein R^{41P} is selected from C₁₋₃alkyl or C₁₋₃alkoxy. Embodiments of the present invention further include processes for the preparation of compounds wherein R^{41P} is selected from methyl, ethyl, methoxy, ethoxy or fluoro. Embodiments of the present invention further include processes for the preparation of compounds wherein R^{41P} is selected from methyl or methoxy.

[0039] Embodiments of the present invention include processes for the preparation of compounds wherein R⁵ is hydrogen or methyl. Embodiments of the present invention further include processes for the preparation of compounds wherein R⁵ is hydrogen.

[0040] Embodiments of the present invention include processes for the preparation of compounds wherein R^a and R^b are independently selected from the group consisting of hydrogen and C₁₋₃alkyl; or, when R^a and R^b are each other than hydrogen, R^a and R^b are optionally taken together with the nitrogen atom to which they are both attached to form a five to seven membered monocyclic ring. Embodiments of the present invention further include processes for the preparation of compounds wherein R^a and R^b are independently hydrogen or methyl. Embodiments of the present invention further include processes for the preparation of compounds wherein R^a and R^b are each hydrogen.


[0041] Embodiments of the present invention include processes for the preparation of compounds of formula (I) wherein the stereo-center denoted with a "*" as shown below,



is in the S-configuration. In another embodiment are processes for the preparation of compounds of formula (I) wherein the stereo-center denoted with a "*" on the compound of formula (I) is in the R-configuration.

[0042] Embodiments of the present invention include processes for the preparation of compounds of formula (II) that are present in their RR, SS, RS, or SR configuration. Embodiments of the present invention further include processes for the preparation of compounds of formula (II) that are present in their S,S configuration.

[0043] Embodiments of the present invention include processes for the preparation of compounds of formula (IIe)



wherein:

R¹ is selected from the group consisting of hydrogen, C₁₋₆alkyl, aryl(C₁₋₄)alkyl, and heteroaryl(C₁₋₄)alkyl; wherein the aryl and heteroaryl portion of aryl(C₁₋₄)alkyl and heteroaryl(C₁₋₄)alkyl are optionally substituted with one to three R¹¹ substituents independently selected from the group consisting of C₁₋₆alkoxy; heteroaryl optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₄alkyl, C₁₋₄alkoxy, and carboxy; carboxy; C₁₋₄alkoxycarbonyloxy; C₁₋₄alkoxycarbonyl; aminocarbonyl; C₁₋₄alkylaminocarbonyl; C₃₋₆cycloalkylaminocarbonyl; hydroxy(C₁₋₆)alkylaminocarbonyl; C₆₋₁₀arylaminocarbonyl wherein C₆₋₁₀aryl is optionally substituted with carboxy or C₁₋₄alkoxycarbonyl; heterocyclylcarbonyl; cyano; halogen; trifluoromethoxy; and hydroxy; provided that no more than one R¹¹ is heteroaryl (optionally substituted with one to two C₁₋₄alkyl substituents); C₆₋₁₀arylaminocarbonyl wherein C₆₋₁₀aryl is optionally substituted with carboxy or C₁₋₄alkoxycarbonyl; or heterocyclylcarbonyl;

R² is selected from the group consisting of hydrogen, C₁₋₄alkyl, hydroxy(C₁₋₄)alkyl, and phenyl(C₁₋₆)alkoxy(C₁₋₄)alkyl; wherein said phenyl is optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₃alkyl, C₁₋₃alkoxy, hydroxy, cyano, fluorine, chlorine, bromine, trifluoromethyl, and trifluoromethoxy; R³ is one to two substituents independently selected from the group consisting of C₁₋₆alkyl, halogen, and aryl; wherein aryl is optionally substituted with one to three substituents independently selected from the group consisting of halogen, carboxy, aminocarbonyl, C₁₋₃alkylsulfonylamino, cyano, hydroxy, amino, C₁₋₃alkylamino, and (C₁₋₃alkyl)₂amino;

R⁵ is hydrogen or methyl;

R^a and R^b are independently hydrogen or C₁₋₃alkyl; or, when R^a and R^b are each other than hydrogen, R^a and R^b are optionally taken together with the nitrogen atom to which they are both attached to form a five to seven membered monocyclic ring;



is C₆₋₁₀aryl;

R^{41P} is selected from C₁₋₃alkyl, C₁₋₆alkoxy or fluoro;

R^J and R^K are each independently selected from hydrogen or C₁₋₄alkyl; alternatively, R^J and R^K are taken together with the nitrogen atom to which they are bound to form a five to seven membered heterocyclyl; and pharmaceutically acceptable enantiomers, diastereomers, racemates, and salts thereof.

[0044] Embodiments of the present invention further include processes for the preparation of compounds of formula (IIe) wherein

R¹ is selected from the group consisting of C₆₋₁₀aryl(C₁₋₄)alkyl, pyridinyl(C₁₋₄)alkyl, and furanyl(C₁₋₄)alkyl; wherein C₆₋₁₀aryl, pyridinyl, and furanyl are optionally substituted with one to three R¹¹ substituents independently selected from

the group consisting of C₁₋₃alkoxy; tetrazolyl; carboxy; C₁₋₃alkoxycarbonyl; aminocarbonyl; C₁₋₄alkylaminocarbonyl; C₁₋₃alkylaminocarbonyl; C₃₋₆cycloalkylaminocarbonyl; hydroxy(C₁₋₄)alkylaminocarbonyl; C₆₋₁₀arylaminocarbonyl wherein C₆₋₁₀aryl is optionally substituted with carboxy or C₁₋₄alkoxycarbonyl; morpholin-4-ylcarbonyl; cyano; halogen; and trifluoromethoxy; provided that no more than one R¹¹ is C₆₋₁₀arylaminocarbonyl;

R² is hydrogen or C₁₋₄alkyl;

R³ is one to two substituents independently selected from the group consisting of C₁₋₃alkyl, bromo, and phenyl; wherein phenyl is optionally substituted with one to three substituents independently selected from the group consisting of chloro, fluoro, carboxy, aminocarbonyl, and cyano;

R⁵ is hydrogen;

R^a and R^b are independently hydrogen or methyl;



is C₆₋₁₀aryl;

R^{41P} is selected from C₁₋₃alkyl or C₁₋₆alkoxy;

R^J and R^K are each independently selected from hydrogen or C₁₋₄alkyl;

and pharmaceutically acceptable enantiomers, diastereomers, racemates, and salts thereof.

[0045] Embodiments of the present invention further include processes for the preparation of compounds of formula (Ile) wherein

R¹ is selected from the group consisting of phenyl(C₁₋₃)alkyl, pyridinyl(C₁₋₃)alkyl, and furanyl(C₁₋₃)alkyl; wherein phenyl, pyridinyl, and furanyl are optionally substituted with one to three R¹¹ substituents independently selected from the group consisting of C₁₋₃alkoxy; tetrazolyl, C₃₋₆cycloalkylaminocarbonyl; hydroxy(C₁₋₄)alkylaminocarbonyl; C₆₋₁₀arylaminocarbonyl wherein C₆₋₁₀aryl is optionally substituted with carboxy or C₁₋₄alkoxycarbonyl; morpholin-4-ylcarbonyl; chloro; fluoro; trifluoromethoxy; and carboxy;

R² is hydrogen or methyl;

R³ is one to two substituents independently selected from the group consisting of methyl and phenyl; wherein phenyl is optionally substituted with one to three substituents independently selected from the group consisting of chloro and carboxy;

R⁵ is hydrogen;

R^a and R^b are each hydrogen;



is phenyl;

R^{41P} is selected from methyl, ethyl, methoxy, ethoxy or fluoro.

R^J and R^K are each independently selected from hydrogen or C₁₋₃alkyl;

and pharmaceutically acceptable enantiomers, diastereomers, racemates, and salts thereof.

[0046] Additional embodiments of the present invention, include processes for the preparation of compounds wherein the substituents for one or more of the variables defined herein (i.e.



R^J, R^K, R^{41P}, Pg¹, etc.) are independently selected to be any individual substituent or any subset of substituents selected from the complete list as defined herein.

[0047] As used herein, unless otherwise noted, "alkyl" whether used alone or as part of a substituent group refers to straight and branched carbon chains having 1 to 8 carbon atoms or any number within this range. The term "alkoxy" refers to an -Oalkyl substituent group, wherein alkyl is as defined supra. Similarly, the terms "alkenyl" and "alkynyl" refer to straight and branched carbon chains having 2 to 8 carbon atoms or any number within this range, wherein an alkenyl chain has at least one double bond in the chain and an alkynyl chain has at least one triple bond in the chain.

An alkyl and alkoxy chain may be substituted on a carbon atom. In substituent groups with multiple alkyl groups such as (C₁₋₆alkyl)₂amino- the C₁₋₆alkyl groups of the dialkylamino may be the same or different.

[0048] The term "**cycloalkyl**" refers to saturated or partially unsaturated, monocyclic or polycyclic hydrocarbon rings of from 3 to 14 carbon atom members. Examples of such rings include, and are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and adamantyl. Alternatively, the cycloalkyl ring may be fused to a benzene ring (benzo fused cycloalkyl), a 5 or 6 membered heteroaryl ring (containing one of O, S or N and, optionally, one additional nitrogen) to form a heteroaryl fused cycloalkyl.

[0049] The term "**heterocyclyl**" refers to a nonaromatic cyclic ring of 5 to 7 members in which 1 to 2 members are nitrogen, or a nonaromatic cyclic ring of 5 to 7 members in which zero, one or two members are nitrogen and up to two members are oxygen or sulfur, wherein, optionally, the ring contains zero to one unsaturated bonds, and, optionally, when the ring is of 6 or 7 members, it contains up to two unsaturated bonds. The term heterocyclyl includes 5 to 7 membered monocycle wherein the heterocyclyl may be fused to a benzene ring (benzo fused heterocyclyl), a 5 or 6 membered heteroaryl ring (containing one of O, S or N and, optionally, one additional nitrogen), a 5 to 7 membered cycloalkyl or cycloalkenyl ring, a 5 to 7 membered heterocyclyl ring (of the same definition as above but absent the option of a further fused ring) or fused with the carbon of attachment of a cycloalkyl, cycloalkenyl or heterocyclyl ring to form a spiro moiety. For some compounds, the carbon atom ring members that form the heterocyclyl ring are fully saturated. Other compounds may have a partially saturated heterocyclyl ring. The term heterocyclyl include a 5 to 7 membered monocyclic ring bridged to form bicyclic rings. Such compounds are not considered to be fully aromatic and are not referred to as heteroaryl compounds. Examples of heterocyclyl groups include, and are not limited to, pyrrolinyl (including 2*H*-pyrrole, 2-pyrrolinyl or 3-pyrrolinyl), pyrrolidinyl, 2-imidazolinyl, imidazolidinyl, 2-pyrazolinyl, pyrazolidinyl, piperidinyl, morpholinyl, thiomorpholinyl and piperazinyl.

[0050] The term "**aryl**" refers to an unsaturated, aromatic monocyclic ring of 6 carbon members or to an unsaturated, aromatic polycyclic ring of from 10 to 14 carbon members. Examples of such aryl rings include, and are not limited to, phenyl, naphthalenyl or anthracenyl. Preferred aryl groups for the practice of this invention are phenyl and naphthalenyl.

[0051] The term "**heteroaryl**" refers to an aromatic ring of 5 or 6 members wherein the ring consists of carbon atoms and has at least one heteroatom member. Suitable heteroatoms include nitrogen, oxygen or sulfur. In the case of 5 membered rings, the heteroaryl ring contains one member of nitrogen, oxygen or sulfur and, in addition, may contain up to three additional nitrogens. In the case of 6 membered rings, the heteroaryl ring may contain from one to three nitrogen atoms. For the case wherein the 6 membered ring has three nitrogens, at most two nitrogen atoms are adjacent. Optionally, the heteroaryl ring is fused to a benzene ring (benzo fused heteroaryl), a 5 or 6 membered heteroaryl ring (containing one of O, S or N and, optionally, one additional nitrogen), a 5 to 7 membered cycloalkyl ring or a 5 to 7 membered heterocyclo ring (as defined supra but absent the option of a further fused ring). Examples of heteroaryl groups include, and are not limited to, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl or pyrazinyl; fused heteroaryl groups include indolyl, isoindolyl, indolinyl, benzofuryl, benzothienyl, indazolyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, benzothiadiazolyl, benzotriazolyl, quinoliziny, quinolinyl, isoquinolinyl or quinazolinyl.

[0052] The term "**arylalkyl**" means an alkyl group substituted with an aryl group (e.g., benzyl, phenethyl). Similarly, the term "**arylalkoxy**" indicates an alkoxy group substituted with an aryl group (e.g., benzyloxy).


[0053] The term "**halogen**" refers to fluorine, chlorine, bromine and iodine. Substituents that are substituted with multiple halogens are substituted in a manner that provides compounds, which are stable.

[0054] Whenever the term "alkyl" or "aryl" or either of their prefix roots appear in a name of a substituent (e.g., arylalkyl, alkylamino) it shall be interpreted as including those limitations given above for "alkyl" and "aryl." Designated numbers of carbon atoms (e.g., C₁₋₆) shall refer independently to the number of carbon atoms in an alkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root. For alkyl, and alkoxy substituents the designated number of carbon atoms includes all of the independent members included in the range specified individually and all the combination of ranges within in the range specified. For example C₁₋₆ alkyl would include methyl, ethyl, propyl, butyl, pentyl and hexyl individually as well as sub-combinations thereof (e.g. C₁₋₂, C₁₋₃, C₁₋₄, C₁₋₅, C₂₋₆, C₃₋₆, C₄₋₆, C₅₋₆, C₂₋₅, etc.).

[0055] Where the compounds have at least one **chiral center**, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. Furthermore, some of the crystalline forms for the compounds may exist as polymorphs. In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents.

[0056] An "**independently**" selected substituent refers to a group of substituents, wherein the substituents may be different. Therefore, designated numbers of carbon atoms (e.g. C₁₋₆) shall refer independently to the number of carbon atoms in an alkyl or cycloalkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.

[0057] Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. Thus, for example, a "phenyl-(C₁₋₆alkyl)amino-carbonyl-(C₁₋₆alkyl)" substituent refers to a group of the formula



[0058] Abbreviations used in the specification, particularly the Schemes and Examples, are as follows:

10	Ac	=	Acetyl group (-C(O)-CH ₃)
	Ac ₂ O	=	Acetic anhydride
	Cbz or CBZ	=	Benzyloxy-carbonyl-
15	Cpd	=	Compound
	DBU	=	1,8-Diazabicyclo[5.4.0]undec-7-ene
	DCM	=	Dichloromethane
	DIPEA or DIEA	=	Diisopropylethylamine
20	DMF	=	N,N-Dimethylformamide
	DPPE	=	1,2-Bis(diphenylphosphino)ethane
	DPPF	=	1,1'-Bis(diphenylphosphino)ferrocene
25	DPPP	=	1,3-Bis(diphenylphosphino)propane
	EDCI or EDC	=	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
	Et ₂ O	=	Diethyl ether
	EtOAc	=	Ethyl acetate
30	Fmoc	=	9-fluorenyl-methoxy-carbonyl-
	HOAc	=	Acetic acid
	HOBT	=	1-Hydroxybenzotriazole
35	Me		Methyl
	MeOH	=	Methanol
	MeO	=	Methoxy
	MTBE	=	Methyl- <i>tert</i> -butyl ether
40	NaBH(OAc) ₃	=	Sodium triacetoxymethylborohydride
	Pd-C	=	Palladium on Carbon Catalyst
	Pd ₂ (OAc) ₂	=	Palladium(II)acetate
45	Pd ₂ (dba) ₃	=	Tris(dibenzylidene acetone)dipalladium(0)
	Pd(PPh ₃) ₄	=	Tetrakis(triphenylphosphine)palladium (0)
	Ph	=	Phenyl
	P(Ph) ₃	=	Triphenylphosphine
50	PyBop	=	Benzotriazol-1-yloxy-tris(pyrrolidino)phosphonium hexafluorophosphate
	PyBrop	=	Bromotri(pyrrolidino)phosphonium hexafluorophosphate
	[Rh(cod)(R,R-DIPAMP)] ⁺ BF ₄ ⁻		(R,R)-(-)-Bis[(o-methoxyphenyl)(phenyl)phosphino]ethane(1,5-cyclo-octadiene)rhodium (I) tetrafluoroborate
55	rt or RT	=	Room temperature
	t-BOC or Boc	=	<i>tert</i> -Butoxycarbonyl

(continued)

TEA	=	Triethylamine
Tf	=	Trifluoromethyl-sulfonyl- (-SO ₂ -CF ₃)
TFA	=	Trifluoroacetic acid
THF	=	Tetrahydrofuran
Tyr	=	Tyrosine

[0059] As used herein, the term "**pain**" shall include centrally mediated pain, peripherally mediated pain, structural or soft tissue injury related pain, pain related to inflammation, progressive disease related pain, neuropathic pain, acute pain and chronic pain. Further, the term "**chronic pain**" shall include neuropathic pain conditions, diabetic peripheral neuropathy, post-herpetic neuralgia, trigeminal neuralgia, post-stroke pain syndromes and cluster or migraine headaches.

[0060] As used herein, the term "**gastrointestinal disorder**" shall include diarrheic syndromes, motility disorders such as diarrhea-predominant, constipation-predominant, alternating irritable bowel syndrome, post-operative ileus and constipation, and inflammatory bowel disease. Further, the term "**inflammatory bowel disease**" shall include ulcerative colitis and Crohn's disease.

[0061] The term "**subject**" as used herein, refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

[0062] The term "**therapeutically effective amount**" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

[0063] As used herein, the term "**composition**" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

[0064] To provide a more concise description, some of the quantitative expressions given herein are not qualified with the term "**about**". It is understood that whether the term "about" is used explicitly or not, every quantity given herein is meant to refer to the actual given value, and it is also meant to refer to the approximation to such given value that would reasonably be inferred based on the ordinary skill in the art, including approximations due to the experimental and/or measurement conditions for such given value.

[0065] As used herein, unless otherwise noted, the term "**aprotic solvent**" shall mean any solvent that does not yield a proton. Suitable examples include, but are not limited to DMF, dioxane, THF, acetonitrile, pyridine, dichloroethane, dichloromethane, MTBE, toluene, and the like.

[0066] One skilled in the art will recognize that wherein a reaction step of the present invention may be carried out in a variety of solvents or solvent systems, said reaction step may also be carried out in a mixture of the suitable solvents or solvent systems.

[0067] Where the processes for the preparation of the compounds give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-D-tartaric acid and/or (+)-di-p-toluoyl-L-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

[0068] During any of the processes for preparation of the compounds, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis. John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

[0069] As used herein, unless otherwise noted, the term "**nitrogen protecting group**" shall mean a group which may be attached to a nitrogen atom to protect said nitrogen atom from participating in a reaction and which may be readily removed following the reaction. Suitable nitrogen protecting groups include, but are not limited to carbamates - groups of the formula -C(O)O-R wherein R is for example methyl, ethyl, t-butyl, benzyl, phenylethyl, CH₂=CH-CH₂-, and the like; amides - groups of the formula -C(O)-R' wherein R' is for example methyl, phenyl, trifluoromethyl, and the like; N-sulfonyl derivatives - groups of the formula -SO₂-R" wherein R" is for example tolyl, phenyl, trifluoromethyl, 2,2,5,7,8-


pentamethylchroman-6-yl-, 2,3,6-trimethyl-4-methoxybenzene, and the like. Other suitable nitrogen protecting groups may be found in texts such as T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991.

[0070] For use in medicine, the salts of the compounds refer to nontoxic pharmaceutically acceptable salts.

[0071] The present invention is directed to processes for the preparation of compounds of formula (I) as outlined in Scheme 1 below.

STEP 1: Preparation of Compounds of Formula (XII), wherein X is -OC(O)-C₁₋₄alkyl and Y is selected from Br, Cl or I


[0072]




or

STEP 1: Preparation of Compounds of Formula (XII), wherein X is -CN and Y is selected from Br, Cl or I


[0073]




STEP 2: Preparation of Compound of Formula (XIX)



STEP 3: Preparation of Compounds of Formula (XX)



STEP 4: Preparation of Compounds of Formula (I)



Scheme 1

STEP 1: Wherein X is -C(O)-OC₁₋₄alkyl and wherein Y^P is Br, Cl or I

[0074] A suitably substituted compound of formula (X), wherein X^P is -C(O)-OC₁₋₄alkyl and wherein Y^P is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with a suitably substituted compound of formula (XIV), a known compound or compound prepared by known methods, or when R^J and R^K are each hydrogen, with a suitable source of ammonia such as NH₄OH, HMDS, ammonia gas, and the like; at a temperature greater than room temperature, preferably at about reflux temperature; to yield the corresponding compound of formula (XII).

[0075] Alternatively, a suitably substituted compound of formula (X), wherein X^P is -C(O)-OC₁₋₄alkyl and wherein Y^P is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with a suitably substituted compound of formula (XIV), a known compound or compound prepared by known methods, or when R^J and R^K are each hydrogen, with a suitable source of ammonia such as NH₄OH, HMDS, ammonia gas, and the like; in the presence of an activating agent such as trimethylaluminum, triisopropylaluminum, and the like; in an aprotic organic solvent such as THF, dioxane, toluene, DCM, and the like; preferably at a temperature in the range of about 0°C to about reflux temperature; to yield the corresponding compound of formula (XII).

[0076] Alternatively, a suitably substituted compound of formula (X), wherein X^P is -C(O)-OC₁₋₄alkyl and wherein Y^P is selected from Br, Cl or I, a known compound or compound prepared by known methods, is hydrolyzed according to known methods, for example by reacting with a base such as NaOH, LiOH, KOH, and the like, or by reacting with an acid such as HCl, H₂SO₄, and the like; preferably, the compound of formula (X) is reacted with an acid at a temperature greater than about room temperature, preferably at a temperature in the range of from about 60° to about 120°C, more preferably at a temperature of about 100°C; to yield the corresponding compound of formula (XIII).

[0077] The compound of formula (XIII) is reacted with a suitably substituted compound of formula (XIV), a known compound or compound prepared by known methods, or when R^J and R^K are each hydrogen, with a suitable source of ammonia such as HMDS, ammonia gas, and the like, preferably HMDS; in the presence of a coupling agent such as EDCI, HOBT, PyBop, PyBrop, and the like; preferably in the presence of an organic base such as TEA, DIPEA, pyridine, the like, or in the presence of an amount of the compound of formula (XIV) or source of ammonia sufficient to act as the base, preferably greater than about 2 equivalents; in an organic solvent such as THF, dioxane, DMF, and the like, to yield the corresponding compound of formula (XII).

[0078] Alternatively, the compound of formula (X), wherein X^P is -C(O)-OC₁₋₄alkyl and wherein Y^P is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with carbon monoxide or a source of carbon monoxide such as Ac₂O in combination with HCOONa (see for example, S. Cacchi, G. Fabrizi, A. Goggiamani, Org. Lett. (2003), 5(23), pp4269-4272); in the presence of an inorganic base such as K₂CO₃, Na₂CO₃, and the like, in an organic solvent such as DMF, dioxane, THF, and the like; at a temperature in the range of from about 50°C to about 160°C; preferably at a temperature in the range of from about 60°C to about 80°C; to yield the corresponding compound of formula (XIII).

[0079] The compound of formula (XIII) is reacted with a suitable source of chlorine such as thionyl chloride, PCl₃, PCl₅, oxalyl chloride, oxalyl chloride in DMF, and the like; in an organic solvent such as DCM, chloroform, and the like, preferably at a temperature greater than about room temperature, more preferably at a temperature in the range of about 35°C to about 60°C, to yield the corresponding compound of formula (XV).

[0080] The compound of formula (XV) is reacted with a suitably substituted compound of formula (XIV), a known compound or compound prepared by known methods, or when R^J and R^K are each hydrogen, with a suitable source of ammonia such as ammonium chloride, NH₄OH, HMDS, ammonia gas, and the like, preferably ammonium chloride; preferably in the presence of an organic base such as TEA, DIPEA, pyridine, the like, or in the presence of an amount of the compound of formula (XIV) or source of ammonia sufficient to act as the base, preferably greater than about 2 equivalents; in an organic solvent such as THF, dioxane, DMF, and the like; to yield the corresponding compound of formula (XII).

[0081] Alternatively, the compound of formula (X), wherein X^P is -C(O)-OC₁₋₄alkyl and wherein Y^P is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with carbon monoxide or a source of carbon monoxide such as Ac₂O in combination with HCOONa (see for example, S. Cacchi, G. Fabrizi, A. Goggiamani, Org. Lett. (2003), 5(23), pp4269-4272); in the presence of an inorganic base such as K₂CO₃, Na₂CO₃, and the like, in an organic solvent such as DMF, dioxane, THF, and the like; at a temperature in the range of from about 50°C to about 160°C, preferably at a temperature in the range of from about 60°C to about 80°C; to yield the corresponding compound of formula (XIII).

[0082] Alternatively, the compound of formula (XIII) is reacted with C₁₋₄alkyl-chloroformate, preferably, methylchloroformate; in the presence of an organic base such as TEA, DIPEA, pyridine and the like; preferably at a temperature less than about room temperature, more preferably at a temperature of about 0°C; in an organic solvent such as DMF, DCM, chloroform, THF, and the like; to yield the corresponding compound of formula (XVI), wherein A¹ is the corresponding C₁₋₄alkyl, preferably methyl.

[0083] The compound of formula (XVI) is reacted with a suitably substituted compound of formula (XIV), a known compound or compound prepared by known methods, or when R^J and R^K are each hydrogen, with a suitable source of ammonia such as NH₄OH, HMDS, ammonia gas, and the like, preferably NH₄OH; in the presence of a palladium catalyst such PdCl₂, Pd₂(OAc)₂, and the like, in combination with a ligand, such DPPP, DPPF, P(Ph)₃, and the like; or in the presence of a palladium:ligand complex such as Pd(PPh₃)₄, and the like; at a temperature in the range of from about 50°C to about 160°C, preferably at a temperature in the range of from about 60°C to about 80°C; to yield the corresponding compound of formula (XII).

STEP 1: Wherein X is -CN and wherein Y^P is Br, Cl or I

[0084] Alternatively, a suitably substituted compound of formula (X), wherein X^P is CN and wherein Y^P is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with a suitably substituted compound of formula (XIV), a known compound or compound prepared by known methods, according to known methods (for example as described in Parris, C.L., Org. Syn. Coll., (1973), 5, p73; Lin, S., Synthesis, (April 1978), p.330; Murahashi, S., Takeshi Naota, T., and Eiichiro Saito, E., JACS, (1986), 108(24), p7846), to yield the corresponding compound of formula (XII).

[0085] Alternatively, a suitably substituted compound of formula (X), wherein X^P is CN and wherein Y^P is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with an acid such as concentrated sulfuric acid, and the like; at a temperature greater than about room temperature, preferably at reflux temperature; to yield the corresponding compound of formula (XVI).

[0086] Alternatively, a suitably substituted compound of formula (X), wherein X^P is CN and wherein Y^P is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with an inorganic base such as NaOH, KOH, and the like; at a temperature greater than about room temperature, preferably at about reflux temperature; to yield the corresponding compound of formula (XVI).

[0087] The compound of formula (XVI) is reacted according to known methods, for example, by alkylating in the presence of a base, to yield the corresponding compound of formula (XII).

STEP 2:

[0088] The compound of formula (XII) is reacted with a suitably substituted compound of formula (XVII), wherein Pg¹ is a suitable nitrogen protecting group such as Boc, Cbz, Fmoc, acetyl, and the like, preferably Pg¹ is Boc, a known compound or compound prepared by known methods; in the presence of palladium catalyst such as Pd₂(dba)₃, Pd(OAc)₂, PdCl₂, and the like, preferably Pd₂(dba)₃; and preferably in the presence of a phosphorous ligand such as P(o-toluene)₃, P(Ph)₃, P(t-butyl)₃, DPPE, and the like, preferably P(t-butyl)₃ or P(o-toluene)₃; or in the presence of a palladium:ligand complex such as Pd(PPh₃)₄, and the like; in the presence of an organic or inorganic base such as dicyclohexylmethylamine, Na₂CO₃, K₂CO₃, TEA, DIPEA, pyridine, and the like, preferably TEA; in an organic solvent such as DMF, dioxane, and the like; at a temperature greater than about room temperature, preferably at a temperature in the range of about 60°C to about 120°C; to yield the corresponding compound of formula (XIX).

STEP 3:

[0089] The compound of formula (XIX) is hydrogenated according to known methods; for example by reacting with hydrogen or a source of hydrogen (such as cyclohexadiene, and the like); in the presence of a catalyst such as platinum oxide, palladium on carbon, nickel, ClRh(PPh₃)₃, RuCl₂, and the like, preferably palladium on carbon; in an organic solvent such as methanol, ethanol, THF, ethyl acetate, and the like, preferably methanol; at a temperature greater than room temperature, preferably at a temperature in the range of about 60°C to about 120°C, to yield the corresponding compound of formula (XX).

[0090] One skilled in the art will recognize that the compound of formula (XIX) may be optionally reacted in the presence of a chiral catalyst, to yield the corresponding compound of formula (XX), wherein one stereo-isomer is present in an enantiomeric excess.

STEP 4:

[0091] The compound of formula (XX) is reacted with an aqueous base such as NaOH, LiOH, KOH, and the like; in an organic solvent such as methanol, THF, ethanol, and the like; to yield the corresponding compound of formula (I).

[0092] In an embodiment, the present invention is directed to processes for the preparation of compounds of formula (Ic).

[0093] In an embodiment, the present invention is directed to processes for the preparation of a compound of formula (Ib), a compound of formula (I) wherein




is phenyl; R^J and R^K are each hydrogen; the phenyl ring is further substituted with two R^{41P} groups, which are each methyl and Pg^1 is Boc, also known as 2-tert-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionic acid, as described in Scheme 1 above.

[0094] The present invention is further directed to processes for the preparation of the compound of formula (Ia) as outlined in Scheme 2 below.

STEP 1a: Preparation of the Compound of Formula (XIIa), wherein X is -OC(O)-C₁₋₄alkyl and Y is selected from Br, Cl or I


[0095]




or

STEP 1a: Preparation of the Compound of Formula (XIIa), wherein X is -CN and Y is selected from Br, Cl or I


[0096]




STEP 2a: Preparation of the Compound of Formula (XIXa)



STEP 3a: Preparation of the Compound of Formula (XXa)



STEP 4a: Preparation of the Compound of Formula (Ia)



Scheme 2

STEP 1a: Wherein X is $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkyl}$ and wherein Y^P is Br, Cl or I

[0097] A suitably substituted compound of formula (Xa), wherein X^P is $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkyl}$ and wherein Y^P is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with a suitable source of ammonia such as NH_4OH , HMDS, ammonia gas, and the like; at a temperature greater than room temperature, preferably at about reflux temperature; to yield the corresponding compound of formula (XIIa).

[0098] Alternatively, a suitably substituted compound of formula (Xa), wherein X^P is $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkyl}$ and wherein Y^P

is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with a suitable source of ammonia such as NH_4OH , HMDS, ammonia gas, and the like; in the presence of a activating agent such as trimethylaluminum, triisopropylaluminum, and the like; in an aprotic organic solvent such as THF, dioxane, toluene, DCM, and the like; preferably, at a temperature in the range of about 0°C to reflux temperature; to yield the corresponding compound of formula (XIIa).

[0099] Alternatively, a suitably substituted compound of formula (Xa), wherein X^{P} is $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkyl}$ and wherein Y^{P} is selected from Br, Cl or I, a known compound or compound prepared by known methods, is hydrolyzed according to known methods; for example by reacting with a base such as NaOH, LiOH, KOH, and the like, or by reacting with an acid such as HCl, H_2SO_4 , and the like; preferably, the compound of formula (Xa) is reacted with an acid at a temperature greater than about room temperature, preferably at a temperature in the range of from about 60° to about 120°C , preferably at a temperature of about 100°C ; to yield the corresponding compound of formula (XIIIa).

[0100] The compound of formula (XIII₂) is reacted with a suitable source of ammonia such as HMDS, ammonia gas, and the like, preferably HMDS; in the presence of a coupling agent such as EDCI, HOBt, PyBop, PyBrop, and the like; preferably in the presence of an organic base such as TEA, DIPEA, pyridine, the like, or in the presence of an amount of the source of ammonia sufficient to act as the base, preferably greater than about 2 equivalents; in an organic solvent such as THF, dioxane, DMF, and the like, to yield the corresponding compound of formula (XIIa).

[0101] Alternatively, the compound of formula (Xa), wherein X^{P} is $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkyl}$ and wherein Y^{P} is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with carbon monoxide or a source of carbon monoxide such as Ac_2O in combination with HCOONa (see for example, S. Cacchi, G. Fabrizi, A. Goggiamani, Org. Lett. (2003), 5(23), pp4269-4272); in the presence of an inorganic base such as K_2CO_3 , Na_2CO_3 , and the like; in an organic solvent such as DMF, dioxane, THF, and the like; at a temperature in the range of from about 50°C to about 160°C , preferably at a temperature in the range of from about 60°C to about 80°C ; to yield the corresponding compound of formula (XIIIa).

[0102] The compound of formula (XIIIa) is reacted with a suitable source of chlorine such as thionyl chloride, PCl_3 , PCl_5 , oxalyl chloride, oxalyl chloride in DMF, and the like; in an organic solvent such as DCM, chloroform, and the like; preferably at a temperature greater than about room temperature, more preferably at a temperature in the range of about 35°C to about 60°C , to yield the corresponding compound of formula (XVa).

[0103] The compound of formula (XVa) is reacted with a suitable source of ammonia such as ammonium chloride, NH_4OH , HMDS, ammonia gas, and the like, preferably ammonium chloride; preferably in the presence of an organic base such as TEA, DIPEA, pyridine, the like; or in the presence of an amount of the source of ammonia sufficient to act as the base, preferably greater than about 2 equivalents; in an organic solvent such as THF, dioxane, DMF, and the like; to yield the corresponding compound of formula (XIIa).

[0104] Alternatively, the compound of formula (Xa), wherein X^{P} is $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkyl}$ and wherein Y^{P} is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with carbon monoxide or a source of carbon monoxide such as Ac_2O in combination with HCOONa (see for example, S. Cacchi, G. Fabrizi, A. Goggiamani, Org. Lett. (2003), 5(23), pp4269-4272); in the presence of an inorganic base such as K_2CO_3 , Na_2CO_3 , and the like; in an organic solvent such as DMF, dioxane, THF, and the like; at a temperature in the range of from about 50°C to about 160°C , preferably at a temperature in the range of from about 60°C to about 80°C ; to yield the corresponding compound of formula (XIIIa).

[0105] Alternatively, the compound of formula (XIIIa) is reacted with $\text{C}_{1-4}\text{alkylchloroformate}$, preferably, methylchloroformate; in the presence of an organic base such as TEA, DIPEA, pyridine and the like; preferably at a temperature less than about room temperature, more preferably at a temperature of about 0°C ; in an organic solvent such as DMF, DCM, chloroform, THF, and the like; to yield the corresponding compound of formula (XVIa), wherein A^1 is the corresponding $\text{C}_{1-4}\text{alkyl}$, preferably methyl.

[0106] The compound of formula (XVIa) is reacted with a suitable source of ammonia such as NH_4OH , HMDS, ammonia gas, and the like, preferably NH_4OH ; in the presence of a palladium catalyst such as PdCl_2 , $\text{Pd}_2(\text{OAc})_2$, and the like, in combination with a ligand, such as DPPP, DPPF, $\text{P}(\text{Ph})_3$, and the like, or in the presence of a palladium:ligand complex such as $\text{Pd}(\text{PPh}_3)_4$, and the like; at a temperature in the range of from about 50°C to about 160°C , preferably at a temperature in the range of from about 60°C to about 80°C ; to yield the corresponding compound of formula (XIIa).

STEP 1a: Wherein X is -CN and wherein Y^{P} is Br, Cl or I

[0107] Alternatively, a suitably substituted compound of formula (Xa), wherein X^{P} is CN and wherein Y^{P} is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with an acid such as concentrated sulfuric acid, and the like; at a temperature greater than about room temperature, preferably at about reflux temperature; to yield the corresponding compound of formula (XVIa).

[0108] Alternatively, a suitably substituted compound of formula (Xa), wherein X^{P} is CN and wherein Y^{P} is selected from Br, Cl or I, a known compound or compound prepared by known methods, is reacted with an inorganic base such

as NaOH, KOH, and the like; at a temperature greater than about room temperature, preferably at about reflux temperature; to yield the corresponding compound of formula (XVIa).

[0109] Preferably, a suitably substituted compound of formula (Xa), a known compound or compound prepared by known methods, is reacted with a triflating reagent such as triflic anhydride, N-phenyltrifluoromethanesulfonimide, and the like, preferably triflic anhydride; in the presence of an organic or inorganic base such as pyridine, TEA, DIPEA, K_3PO_4 , K_2CO_3 , and the like, preferably pyridine; optionally in an organic solvent such as DCM, chloroform, THF, and the like; to yield the corresponding compound of formula (XIa).

[0110] The compound of formula (XIa) is reacted with carbon monoxide or a source of carbon monoxide such as Ac_2O in combination with $HCOONa$ (see for example, S. Cacchi, G. Fabrizi, A. Goggiani, *Org. Lett.* (2003), 5(23), pp4269-4272) and a suitable source of ammonia such as HMDS, ammonia gas, and the like; preferably the compound of formula (XIa) is reacted with carbon monoxide and HMDS; in the presence of a palladium catalyst such as $PdCl_2$, $Pd_2(OAc)_2$, and the like, in combination with a suitable ligand, such as DPPP, DPPF, $P(Ph)_3$, and the like; or in the presence of a palladium:ligand complex such as $Pd(PPh_3)_4$, and the like; preferably, in the presence of $PdCl_2$ in combination with DPPP; at a temperature in the range of from about 50°C to about 160°C, preferably at a temperature in the range of from about 60°C to about 120°C, more preferably, at a temperature of about 100°C; in an organic solvent such as DMF, THF, dioxane, and the like, preferably, in DMF; to yield the corresponding compound of formula (XII).

STEP 2a:


[0111] The compound of formula (XIIa) is reacted with a suitably substituted compound of formula (XVIIIa), a known compound or compound prepared by known methods, in the presence of palladium catalyst such as $Pd_2(dba)_3$, $Pd(OAc)_2$, $PdCl_2$, and the like, preferably $Pd_2(dba)_3$; and preferably in the presence of a phosphorous ligand such as $P(o-toluen)_3$, $P(Ph)_3$, $P(t-butyl)_3$, DPPE, and the like, preferably $P(o-toluen)_3$; or in the presence of a palladium:ligand complex such as $Pd(PPh_3)_4$, and the like; in the presence of an organic or inorganic base such as dicyclohexylmethylamine, Na_2CO_3 , K_2CO_3 , TEA, DIPEA, pyridine, and the like, preferably TEA; in an organic solvent such as DMF, dioxane, and the like, preferably DMF; at a temperature greater than about room temperature, preferably at a temperature in the range of about 60°C to about 120°C, preferably at about 120°C; to yield the corresponding compound of formula (XIXa).

STEP 3a:

[0112] The compound of formula (XIXa) is reacted with hydrogen gas, at a pressure sufficient to hydrogenate, preferably at a pressure greater than about 500 psi, more preferably, at a pressure greater than about 800 psi, more preferably still, at a pressure about 1000 psi; in the presence of a suitable chiral catalyst such as $[Rh(cod)(R,R-DIPAMP)]^+BF_4^-$, $[Rh(cod)(R,R-DIPAMP)]^+SO_2CF_3^-$, and the like; wherein the chiral catalyst is preferably present in an amount greater than about 0.01 equivalents, more preferably, in an amount of about 0.04 equivalents; at a temperature greater than about room temperature, preferably at a temperature in the range of about 60°C to about 100°C, more preferably, at a temperature of about 60°C; in an organic solvent such as methanol, ethanol, THF, ethyl acetate, and the like, preferably methanol; preferably not under vacuum; to yield the corresponding compound of formula (XXa), wherein the S-enantiomer is present in an enantiomeric excess of greater than about 80%, preferably, in an enantiomeric excess of greater than about 90%, more preferably, in an enantiomeric excess of greater than about 95%, more preferably, in an enantiomeric excess of greater than about 98%, most preferably, in an enantiomeric excess of greater than about 99%.

[0113] One skilled in the art will recognize that if the chiral catalyst is oxygen sensitive, then the hydrogenation reaction vessel is purged with an inert gas such as argon, nitrogen, and the like, prior to charging the vessel with the oxygen sensitive catalyst reagents and hydrogen gas.

[0114] One skilled in the art will recognize that the compound of formula (XIXa) may be optionally reacted to yield the corresponding racemic compound of formula (XXb), as outlined in the scheme below,



by hydrogenating the compound of formula (XIXa) according to known methods, for example, by reacting with hydrogen

or a source of hydrogen (such as cyclohexadiene, and the like); in the presence of a catalyst such as platinum oxide, palladium on carbon, nickel, $\text{ClRh(PPh}_3)_3$, RuCl_2 , and the like, preferably palladium on carbon; in a solvent such as methanol, ethanol, THF, ethyl acetate, and the like; in an organic solvent such as methanol, ethanol, THF, ethyl acetate, and the like, preferably methanol; at a temperature greater than room temperature, preferably at a temperature in the

[0115] Preferably, for the preparation of the compound of formula (Ib), the compound of formula (XIXa) is reacted with hydrogen gas; at a pressure sufficient to hydrogenate, preferably at a pressure greater than about 40 psi, more preferably at a pressure of about 51 psi; in a solvent such as methanol, ethanol, THF, and the like, preferably methanol; preferably, at about room temperature; to yield the corresponding compound of formula (XXb).


[0116] The compound of formula (XXb) is then reacted according to the process described in Step 4a below, to yield the corresponding compound of formula (Ib).


STEP 4a:

[0117] The compound of formula (XXa) is reacted with an aqueous base such as NaOH, LiOH, KOH, and the like, preferably LiOH; in an organic solvent such as methanol, THF, ethanol, and the like, preferably THF; to yield the corresponding compound of formula (Ia).

[0118] The present invention is further directed to processes for the preparation of compounds of formula (II).

[0119] The compounds of formula (I) may be further reacted according to known processes, for example as disclosed in United States Patent Application No. 11/079,647, filed March 14, 2005, and published as US Patent Publication US-2005-0203143-A1, September 15, 2005, to yield the corresponding compounds of formula (II). More specifically, the compounds of formula (II) may be prepared according to the process outlined in Scheme 3 below.






Scheme 3

[0120] Accordingly, a suitably substituted compound of formula (I) is reacted with a suitably substituted compound of formula (L), a known compound or compound prepared by known methods, under standard peptide coupling conditions (for example, with a coupling agent such as EDCI and an additive such as HOBT), to yield the corresponding compound of formula (LI).

[0121] The compound of formula (LI) is then de-protected according to known methods, and then further, optionally reacted according to known methods, to yield the corresponding compound of formula (II) wherein R^a and R^b are each other than hydrogen. For example, the compound of formula (LI) is de-protected and the alkylated, according to known methods, to yield the corresponding compound of formula (II) wherein one or both of R^a and R^b is alkyl. Alternatively, for compounds of formula (II) wherein R^a and R^b are taken together to form a ring, the compound of formula (LI) is de-protected and then converted to the corresponding ring by reductive cyclization with a suitably selected di-aldehyde.

[0122] A process for the preparation of compounds of formula (XIX) is outlined in Scheme 4.




Scheme 4

[0123] Accordingly, a suitably substituted compound of formula (XIII), wherein Y^P is Br or Cl, is reacted with a formylating reagent such as a DMF, $HC(O)-N(CH_3)(OCH_3)$, and the like; in the presence of a base such as n-butyl lithium, NaH, and the like; in an organic solvent such as THF, dioxane, and the like; at a temperature less than about room temperature, preferably at a temperature in the range of about $-130^\circ C$ to about $0^\circ C$, more preferably, at about $-100^\circ C$; to yield the corresponding compound of formula (XXI).

[0124] The compound of formula (XXI) is reacted with a suitably substituted compound of formula (XIV), a known compound or compound prepared by known methods, or when R^J and R^K are each hydrogen, with a suitable source of ammonia such as HMDS, ammonia gas, and the like, preferably HMDS; in the presence of a coupling agent such as EDCI, HOBt, PyBop, PyBrop, and the like; preferably in the presence of an organic base such as TEA, DIPEA, pyridine, the like, or an amount of the compound of formula (XIV) or source of ammonia sufficient to act as the base, preferably greater than about 2 equivalents; in an organic solvent such as THF, dioxane, DMF, and the like; to yield the corresponding compound of formula (XXII).

[0125] The compound of formula (XXII) is reacted with a suitably selected compound of formula (XXIII), a known compound or compound prepared by known methods; in the presence of a base such as DBU, potassium t-butoxide, NaH, and the like; in an organic solvent such as THF, dioxane, and the like; preferably at about room temperature, to yield the corresponding compound of formula (XIX).

[0126] A further process for the preparation of compounds of formula (XIX) is outlined in Scheme 5.



Scheme 5.


[0127] Accordingly, a suitably substituted compound of formula (XXIV), a known compound or compound prepared by known methods, is reacted with a suitably substituted compound of formula (XIV), a known compound or compound prepared by known methods, or when R^J and R^K are each hydrogen, with a suitable source of ammonia such as HMDS, ammonia gas, and the like, preferably HMDS; in the presence of a coupling agent such as EDCI, HOBT, PyBop, PyBrop, and the like; preferably in the presence of an organic base such as TEA, DIPEA, pyridine, the like, or an amount of the compound of formula (XIV) or source of ammonia sufficient to act as the base, preferably greater than about 2 equivalents; in an organic solvent such as THF, dioxane, DMF, and the like; to yield the corresponding compound of formula (XXV).

[0128] The compound of formula (XXV) is reacted with triflating reagent such as triflic anhydride, N-phenyltrifluoromethanesulfonimide, and the like; in the presence of an organic or inorganic base such as pyridine, TEA, DIPEA, K_3PO_4 , K_2CO_3 , and the like; optionally in an organic solvent such as DCM, chloroform, THF, and the like; to yield the corresponding compound of formula (XXVI).

[0129] The compound of formula (XXVI) is reacted with carbon monoxide; in the presence of a palladium catalyst such as $PdCl_2$, $Pd_2(OAc)_2$, and the like, in combination with a suitable ligand, such as DPPP, DPPF, $P(Ph)_3$, and the like; or in the presence of a palladium:ligand complex such as $Pd(PPh_3)_4$, and the like; in the presence of an organic base such as TEA, DIPEA, pyridine, and the like; in the presence of $(alkyl)_3SiH$; in an organic solvent such as DMF, THF, dioxane, and the like; to yield the corresponding compound of formula (XXVII).

[0130] The compound of formula (XXVII) is reacted with a suitably selected compound of formula (XXIII), a known compound or compound prepared by known methods; in the presence of a base such as DBU, potassium t-butoxide, NaH, and the like; in an organic solvent such as THF, dioxane, and the like; preferably at about room temperature, to yield the corresponding compound of formula (XIX).

[0131] A process for the preparation of compounds of formula (XIXb) is outlined in Scheme 6.




Scheme 6

[0132] Accordingly, a suitably substituted compound of formula (XIIIa), wherein Y^P is Br or Cl, is reacted with a formylating reagent such as a DMF, $HC(O)-N(CH_3)(OCH_3)$, and the like; in the presence of a base such as n-butyl lithium, NaH, and the like; in an organic solvent such as THF, dioxane, and the like; at a temperature less than about room temperature, preferably at a temperature in the range of about -130°C to about 0°C , more preferably, at about -100°C ; to yield the corresponding compound of formula (XXIa).

[0133] The compound of formula (XXIa) is reacted with a suitable source of ammonia such as HMDS, ammonia gas, and the like, preferably HMDS; in the presence of a coupling agent such as EDCI, HOBT, PyBop, PyBrop, and the like; preferably in the presence of an organic base such as TEA, DIPEA, pyridine, the like, or an amount of the source of ammonia sufficient to act as the base, preferably greater than about 2 equivalents; in an organic solvent such as THF, dioxane, DMF, and the like; to yield the corresponding compound of formula (XXIIa).

[0134] The compound of formula (XXIIa) is reacted with a suitably selected compound of formula (XXIIIa), wherein Pg^1 is a suitable nitrogen protecting group such as Boc, Cbz, and the like, a known compound or compound prepared by known methods; in the presence of a base such as DBU, potassium t-butoxide, NaH, and the like; in an organic solvent such as THF, dioxane, and the like; preferably at about room temperature, to yield the corresponding compound of formula (XIXb).

[0135] A process for the preparation of compounds of formula (XIXb) as outlined in Scheme 7.



Scheme 7

[0136] Accordingly, a suitably substituted compound of formula (XXIVa), a known compound or compound prepared by known methods, is reacted with a suitable source of ammonia such as HMDS, ammonia gas, and the like, preferably HMDS; in the presence of a coupling agent such as EDCI, HOBt, PyBop, PyBrop, and the like; preferably in the presence of an organic base such as TEA, DIPEA, pyridine, the like, or in the presence of an amount of the source of ammonia sufficient to act as the base, preferably greater than about 2 equivalents; in an organic solvent such as THF, dioxane, DMF, and the like; to yield the corresponding compound of formula (XXVa).

[0137] The compound of formula (XXVa) is reacted with triflating reagent such as triflic anhydride, N-phenyltrifluoromethanesulfonimide, and the like; in the presence of an organic or inorganic base such as pyridine, TEA, DIPEA, K_3PO_4 , K_2CO_3 , and the like; optionally in an organic solvent such as DCM, chloroform, THF, and the like; to yield the corresponding compound of formula (XXVIa).

[0138] The compound of formula (XXVIa) is reacted with carbon monoxide; in the presence of a palladium catalyst such as $PdCl_2$, $Pd_2(OAc)_2$, and the like, in combination with a suitable ligand, such as DPPP, DPPF, $P(Ph)_3$, and the like; or in the presence of a palladium:ligand complex such as $Pd(PPh_3)_4$, and the like; in the presence of an organic base such as TEA, DIPEA, pyridine, and the like; in the presence of $(alkyl)_3SiH$; in an organic solvent such as DMF, THF, dioxane, and the like; to yield the corresponding compound of formula (XXVIIa).

[0139] The compound of formula (XXVIIa) is reacted with a suitably selected compound of formula (XXIIIa), wherein Pg^1 is a suitable nitrogen protecting group such as Boc, Cbz, and the like, a known compound or compound prepared by known methods; in the presence of a base such as DBU, potassium t-butoxide, NaH, and the like; in an organic solvent such as THF, dioxane, and the like; preferably at about room temperature, to yield the corresponding compound of formula (XIXb).

[0140] Also disclosed are pharmaceutical compositions containing one or more compounds prepared according to any of the processes described herein with a pharmaceutically acceptable carrier. Pharmaceutical compositions containing one or more of the compounds described herein as the active ingredient can be prepared by intimately mixing the compound or compounds with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending upon the desired route of administration (e.g., oral, parenteral). Thus for liquid oral preparations such as suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, stabilizers, coloring agents and the like; for solid

oral preparations, such as powders, capsules and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Solid oral preparations may also be coated with substances such as sugars or be enteric-coated so as to modulate major site of absorption. For parenteral administration, the carrier will usually consist of sterile water and other ingredients may be added to increase solubility or preservation. Injectable suspensions or solutions may also be prepared utilizing aqueous carriers along with appropriate additives.

[0141] Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of the disease condition. In addition, factors associated with the particular patient being treated, including patient age, weight, diet and time of administration, will result in the need to adjust dosages.


[0142] The following Examples are set forth to aid in the understanding of the invention, and are not intended and should not be construed to limit in any way the invention set forth in the claims which follow thereafter.

[0143] In the Examples which follow, some synthesis products are listed as having been isolated as a residue. It will be understood by one of ordinary skill in the art that the term "residue" does not limit the physical state in which the product was isolated and may include, for example, a solid, an oil, a foam, a gum, a syrup, and the like.

Reference Example 1

(S)-2-tert-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionic acid

[0144]



STEP A: Trifluoromethanesulfonic acid 4-bromo-3,5-dimethyl-phenyl ester

[0145] To a cooled (0°C) solution of 4-bromo-3,5-dimethylphenol (3.05 g, 15.2 mmol) in pyridine (8 mL) was added trifluoromethanesulfonic anhydride (5.0 g, 17.7 mmol) dropwise. After completion of addition, the resulting mixture was stirred at 0°C for 15 min, and then at room temperature overnight. The reaction was quenched by addition of water, and then extracted with EtOAc. The organic extracts were washed sequentially with water, 2N HCl (2x), brine, and then dried over MgSO₄. Filtration and evaporation to dryness yielded compound **1b** as a colorless oil.

[0146] ¹H NMR (300 MHz, CDCl₃): δ 2.45 (6H, s), 7.00 (2H, s).

Step B: 4-Bromo-3,5-dimethylbenzoic acid

[0147] Into a solution of compound **1b** (6.57 g, 19.7 mmol) in DMF (65 mL) were added K₂CO₃ (13.1 g, 94.7 mmol), Pd(OAc)₂ (0.44 g, 1.97 mmol) and 1,1'-bis(diphenylphosphino)ferrocene (2.29 g, 4.14 mmol). The resulting mixture was bubbled in gaseous CO for 10 min and was heated to 60°C for 7.5h with a CO_(g) balloon. The cooled mixture was partitioned between aqueous NaHCO₃ and EtOAc, and filtered. The aqueous phase was separated, acidified with aqueous 6N HCl, extracted with EtOAc, and then dried over Na₂SO₄. Filtration and concentration of the filtrate yielded crude compound **1c** as a brown residue, which was used in the next step without further purification.

STEP C: Method A: 4-Bromo-3,5-dimethyl-benzamide

[0148] Into a suspension of compound **1c** in DCM (40 mL) was added SOCl₂ (3.1 mL, 42 mmol) and the mixture was heated at reflux for 2 h. Upon removal of the solvent by evaporation, the residue was dissolved in DCM (40 mL) and then ammonium hydroxide (28% NH₃ in water, 2.8 mL) was added. The reaction mixture was heated at 50°C for 2 h and concentrated. The residue was diluted with H₂O, extracted with EtOAc, and the organic portion was dried over Na₂SO₄. After filtration and evaporation, the residue was purified by flash column chromatography (eluent: EtOAc) to yield compound **1d** as an off-white solid.

[0149] ¹H NMR (300 MHz, CD₃CN): δ 2.45 (6H, s), 5.94 (1 H, br s), 6.71 (1 H, br s), 7.57 (2H, s)

[0150] MS(ES⁺)(relative intensity): 228.0 (100%) (M+1).

Step C: Method B: 4-Bromo-3,5-dimethyl-benzamide

[0151] A mixture of compound **1b** (3.33 g, 10 mmol), Pd-Cl₂(0.053 g, 0.3 mmol), hexamethyldisilazane (HMDS, 8.4 mL, 40 mmol), and DPPP (0.12 g, 0.3 mmol) was bubbled with a gaseous CO for 5 min and then stirred in a CO balloon at 80°C for 4 h. To the reaction mixture was added MeOH (5 mL). The reaction mixture was stirred for 10 min, diluted with 2N H₂SO₄ (200 mL), and then extracted with EtOAc. The EtOAc extract was washed with saturated aqueous NaHCO₃, brine, and then dried over Na₂SO₄. Filtration and evaporation of the resultant filtrate yielded a residue, which was purified by flash column chromatography (eluent: EtOAc) to yield compound **1d** as a white solid.

Step D: 2-tert-Butoxycarbonylaminoacrylic acid methyl ester

[0152] To a suspension of *N*-Boc-serine methyl ester (Compound **1e**, 2.19 g, 10 mmol) and EDCI (2.01 g, 10.5 mmol) in DCM (70 mL) was added CuCl (1.04 g, 10.5 mmol). The reaction mixture was stirred at room temperature for 72 h. Upon removal of the solvent, the residue was diluted with EtOAc, washed sequentially with water and brine and then dried over MgSO₄. The crude product was purified by flash column chromatography (eluent: EtOAc:hexane ~1:4) to yield compound **1f** as a colorless oil.

[0153] ¹H NMR (300 MHz, CDCl₃): δ 1.49 (9H, s), 3.83 (3H, s), 5.73 (1 H, d, *J* = 1.5 Hz), 6.16 (1 H, s), 7.02 (1 H, s).

STEP E: (Z)-2-tert-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)acrylic acid methyl ester

[0154] A flask charged with compound **1d** (0.46 g, 2.0 mmol), compound **1f** (0.80 g, 4.0 mmol), tri-*o*-tolylphosphine (0.098 g, 0.32 mmol) and DMF (8 mL) was purged with N_{2(g)} 3 times. After the addition of tris(dibenzylideneacetone)dipalladium (0) (0.074 g, 0.08 mmol) and TEA (0.31 mL, 2.2 mol), the reaction mixture was heated at 110°C for 24 h. At that time, the reaction was quenched by addition of water, and then extracted with EtOAc. The organic phase was washed with 1 N HCl, saturated aqueous NaHCO₃, brine, and dried over MgSO₄. The mixture was concentrated to a residue, which was purified by flash column chromatography (eluent: EtOAc:hexane~1:1 to EtOAc only) to yield compound **1g** as a white solid.

[0155] ¹H NMR (300 MHz, CD₃OD): δ 1.36 (9H, s), 2.26 (6H, s), 3.83 (3H, s), 7.10 (1 H, s), 7.56 (2H, s);

[0156] ¹³C NMR (75 MHz, DMSO-*d*₆): δ 17.6, 25.7, 50.2, 78.7, 124.9, 126.4, 128.3, 131.2, 135.2, 135.5, 152.8, 164.3,

169.6;

[0157] MS (ES⁺) (relative intensity): 349.1 (38%)(M+1).**STEP F: (S)-2-tert-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)propionic acid methyl ester**

[0158] Into a reactor charged with a solution of compound **1g** (0.56 g, 1.6 mmol) in degassed MeOH (80 mL) was added [Rh(cod)(R,R-DIPAMP)]⁺BF₄⁻ under a stream of argon. The reactor was sealed and flushed with H₂, stirred at 60°C under 1000 psi of H₂ for 14 days. The crude product was purified by flash column chromatography (eluent: EtOAc:hexane ~1:1) to yield compound **1h** as a white solid.

[0159] ee: >99%;

[0160] ¹H NMR (300 MHz, CDCl₃): δ 1.36 (9H, s), 2.39 (6H, s), 3.11 (2H, J = 7.2 Hz), 3.65 (3H, s), 4.53-4.56 (1H, m), 5.12 (1H, d, J = 8.7 Hz), 5.65 (1H, br s), 6.09 (1H, br s), 7.46 (2H, s);


[0161] MS(ES⁺) (relative intensity): 250.9 (100) (M-Boc)⁺.**STEP G: (S)-2-tert-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)propionic acid**

[0162] Into an ice-cooled solution of compound **1h** (0.22 g, 0.63 mmol) in THF (3.5 mL) was added an aqueous LiOH solution (1 N, 3.5 mL) and the reaction mixture stirred at 0°C. Upon completion of the reaction, the reaction mixture was concentrated and the aqueous phase was neutralized with cooled aqueous 1 N HCl at 0°C, and then extracted with EtOAc. The combined extracts were dried over Na₂SO₄ overnight. Filtration and evaporation of the filtrate to dryness yielded compound **1j** as a white solid.

[0163] ¹H NMR (300 MHz, DMSO-*d*₆): δ 1.30 (9H, s), 2.32 (6H, s), 2.95 (1H, dd, J = 8.8, 13.9 Hz), 3.10 (1H, dd, J = 6.2, 14.0 Hz), 4.02-4.12 (1H, m), 7.18-7.23 (2H, m), 7.48 (2H, s), 7.80 (1H, s);

[0164] MS(ES⁺) (relative intensity): 236.9 (6) (M-Boc)⁺.**Example 2****Racemic 2-tert-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)-propionic acid**

[0165]

**STEP A: Racemic 2-tert-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)propionic acid methyl ester**

[0166] To a reactor charged with a solution of compound **1g** (0.68 g, 1.95 mmol) in MeOH (80 mL) was added 10% Pd-C (0.5 g). The reactor was connected to a hydrogenator and shaken under 51 psi of H₂ overnight. The mixture was filtered through a pad of Celite and the filtrate was concentrated to dryness to yield compound **2a** as a white solid.


[0167] The ¹H NMR spectrum was identical to that of (S)-2-tert-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)propionic acid methyl ester, compound **1h**.

STEP B: Racemic 2-*tert*-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)propionic acid

[0168] Following the procedure described for Example 1, STEP G (preparation of (*S*)-2-*tert*-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)propionic acid), compound **2b** - racemic 2-*tert*-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)propionic acid - was prepared.

Reference Example 3**2-Amino-3-(4-hydroxy-2,6-dimethyl-phenyl)-N-isopropyl-N-[1-(4-phenyl-1H-imidazol-2-yl)-ethyl]-propionamide**

[0169]

**STEP A. [1-(2-Oxo-2-phenyl-ethylcarbamoyl)-ethyl]-carbamic acid benzyl ester.**

[0170] To a solution of commercially available N- α -CBZ-L-alanine (2.11 g, 9.5 mmol) in dichloromethane (50 mL) was added 2-aminoacetophenone hydrochloride (1.62g, 9.5 mmol). The resulting solution was cooled to 0°C and N-methylmorpholine (1.15 g, 11 mmol), 1-hydroxybenzotriazole (2.55 g, 18.9 mmol) and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (2.35 g, 12.3 mmol), in that order, were added under an Argon atmosphere. The reaction mixture was warmed to room temperature and stirred overnight. The reaction was quenched by addition of saturated aqueous NaHCO₃ solution; the separated organic phase was washed with 2N citric acid, saturated NaHCO₃ solution and brine, then dried over MgSO₄ overnight. After filtration and concentration, the residue was purified by column chromatography on silica gel (eluent, EtOAc:hexane-1:1) to yield the title compound, [1-(2-oxo-2-phenylethylcarbamoyl)-ethyl]-carbamic acid benzyl ester.

[0171] ¹H NMR (300 MHz, CDCl₃): δ 1.46 (3H, d), 4.39 (1H, m), 4.75 (2H, d), 5.13 (2H, d), 5.40 (1H, m), 7.03 (1H, m), 7.36 (5H, m), 7.50 (2H, m), 7.63 (1H, m), 7.97 (2H, m)

[0172] MS(ES⁺): 341.1 (100%).

step B. [1-(4-Phenyl-1H-imidazol-2-yl)-ethyl]-carbamic acid benzyl ester.

[0173] To a suspension of [1-(2-oxo-2-phenyl-ethylcarbamoyl)-ethyl]-carbamic acid benzyl ester (2.60 g, 7.64 mmol) in xylene (60 mL) was added NH₄OAc (10.3 g, 134 mmol) and HOAc (5 mL). The resulting mixture was heated at reflux for 7 h. After being cooled to room temperature, brine was added and the mixture was separated. The aqueous phase

was extracted with EtOAc, and the combined organic phases were dried over Na₂SO₄ overnight. After filtration and concentration, the residue was purified by column chromatography on silica gel (eluent, EtOAc:hexane-1:1) to yield the title compound.

[0174] ¹H NMR (300 MHz, CDCl₃): δ 1.65 (3H, d), 5.06 (1H, m), 5.14 (2H, q), 5.94 (1H, d), 7.32 (10H, m), 7.59 (2H, d)

[0175] MS(ES⁺): 322.2 (100%).

Step C. 1-(4-Phenyl-1*H*-imidazol-2-yl)-ethylamine

[0176] To a solution of [1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-carbamic acid benzyl ester (1.5 g, 4.67 mmol) in methanol (25 mL) was added 10% palladium on carbon (0.16 g). The mixture was shaken in a hydrogenation apparatus at rt under a hydrogen atmosphere (10 psi) for 8 h. Filtration followed by evaporation to dryness under reduced pressure yielded the crude product 1-(4-phenyl-1*H*-imidazol-2-yl)-ethylamine.

[0177] ¹H NMR (300 MHz, CDCl₃): δ 1.53 (3H, d), 4.33 (1H, q), 7.23 (3H, m), 7.37 (2H, m), 7.67 (2H, m)

[0178] MS(ES⁺): 188.1 (38%).

STEP D. Isopropyl-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-amine

[0179] 1-(4-Phenyl-1*H*-imidazol-2-yl)-ethylamine (0.20 g, 1.07 mmol) and acetone (0.062 g, 1.07 mmol) were mixed in 1,2-dichloroethane (4 mL), followed by the addition of NaBH(OAc)₃ (0.34 g, 1.61 mmol). The resulting mixture was stirred at rt for 3 h. The reaction was quenched with saturated NaHCO₃ solution. The mixture was extracted with EtOAc and the combined extracts were dried over Na₂SO₄. Filtration followed by evaporation to dryness under reduced pressure yielded crude isopropyl-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-amine, which was used for the next reaction without further purification.

[0180] ¹H NMR (300 MHz, CDCl₃): δ 1.10 (3H, d), 1.18 (3H, d), 1.57 (3H, d), 2.86 (1H, m), 4.32 (1H, m), 7.24 (2H, m), 7.36 (2H, m), 7.69 (2H, m)

[0181] MS(ES⁺): 230.2 (100%).

STEP E. (2-(4-Hydroxy-2,6-dimethyl-phenyl)-1-{isopropyl-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-carbamoyl}-ethyl)-carbamic acid *tert*-butyl ester

[0182] Into a solution of 2-*tert*-Butoxycarbonylamino-3-(4-hydroxy-2,6-dimethylphenyl)-propionic acid (0.18 g, 0.6 mmol) in DMF (7 mL) was added isopropyl-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-amine (0.11 g, 0.5 mmol), 1-hydroxy-benzotriazole (0.22 g, 1.6 mmol) and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.12 g, 0.6 mmol). The resulting mixture was stirred under an Argon atmosphere at room temperature overnight. The reaction mixture was extracted with EtOAc and the combined organic extracts were washed sequentially with saturated aqueous NaHCO₃ solution, 1N HCl, saturated aqueous NaHCO₃ solution, and brine. The organic phase was then dried over MgSO₄, filtered, and the filtrate was concentrated under reduced pressure. The resulting residue was purified by flash column chromatography (eluent: EtOAc) to yield the product (2-(4-hydroxy-2,6-dimethyl-phenyl)-1-{isopropyl-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-carbamoyl}-ethyl)-carbamic acid *tert*-butyl ester.

[0183] MS(ES⁺): 521.5 (100%).

Step F. 2-Amino-3-(4-hydroxy-2,6-dimethyl-phenyl)-*N*-isopropyl-*N*-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-propionamide

[0184] A solution of (2-(4-hydroxy-2,6-dimethyl-phenyl)-1-{isopropyl-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-carbamoyl}-ethyl)-carbamic acid *tert*-butyl ester (0.13 g, 0.25 mmol) in trifluoroacetic acid (5 mL) was stirred at room temperature for 2 h. Upon removal of the solvents, the residue was purified by preparative LC and lyophilized to yield the TFA salt of the title compound as a white powder.


[0185] ¹H NMR (300 MHz, CDCl₃): δ 0.48 (3H, d), 1.17 (3H, d), 1.76 (3H, d), 2.28 (6H, s), 3.19 (2H, m), 3.74 (1H, m), 4.70 (1H, m), 4.82 (1H, q), 6.56 (2H, s), 7.45 (4H, m), 7.74 (2H, m)

[0186] MS(ES⁺): 421.2 (100%).

Example 4

(3,4-Dimethoxy-benzyl)-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-amine

[0187]



[0188] A solution of 1-(4-phenyl-1*H*-imidazol-2-yl)-ethylamine (0.061 g, 0.33 mmol) of Example 3, and 0.55 g (0.33 mmol) of 3,4-dimethoxybenzaldehyde in 5 mL of anhydrous methanol was stirred at room temperature for 1 h and then cooled to about 0-10°C in an ice bath for 1 h. The reaction was treated carefully with 0.019 g (0.49 mmol) of sodium borohydride in one portion and maintained at about 0-10°C for 21 h. Cold 2M aqueous HCl was added dropwise (30 drops), the mixture was stirred for 5 min, and then partially concentrated *in vacuo* unheated. The residual material was taken up in EtOAc to yield a suspension that was treated with 5 mL of cold 3M aqueous NaOH and stirred vigorously until clear. The phases were separated and the aqueous layer was extracted three times additional with EtOAc. The combined extracts were dried over MgSO₄, filtered, and concentrated to yield (3,4-dimethoxybenzyl)-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-amine as a light yellow oil (HPLC: 87% @ 254nm and 66% @ 214 nm).


[0189] MS (ES⁺) (relative intensity): 338.1 (100) (M+1)

[0190] This sample was of sufficient quality to use in the next reaction without further purification.

Example 5

5-{{[2-Amino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionyl]-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-amino}-methyl)-2-methoxy-benzoic acid

[0191]



STEP A. 2-Methoxy-5-([1-(4-phenyl-1*H*-imidazol-2-yl)-ethylamino]-methyl)-benzoic acid methyl ester

[0192] Using the procedures described for Example 4, substituting 5-formyl-2-methoxy-benzoic acid methyl ester (WO 02/22612) for 3,4-dimethoxybenzaldehyde, 2-methoxy-5-([1-(4-phenyl-1 *H*-imidazol-2-yl)-ethylamino]-methyl)-benzoic acid methyl ester was prepared.

STEP B. 5-([2-*tert*-Butoxycarbonylmethyl-3-(4-carbamoyl-2,6-dimethylphenyl)-propionyl]-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-amino)-methyl)-2-methoxy-benzoic acid methyl ester

[0193] Using the procedure of Example 3 for the conversion of Cpd **3d** to Cpd **3e**, substituting 2-methoxy-5-([1-(4-phenyl-1 *H*-imidazol-2-yl)-ethylamino]-methyl)-benzoic acid methyl ester for Cpd **3d** and substituting 2-*tert*-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionic acid for 2-*tert*-Butoxycarbonylamino-3-(4-hydroxy-2,6-dimethyl-phenyl)-propionic acid, Cpd **5a** was prepared.

STEP C. 5-([2-*tert*-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)-propionyl]-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-amino)-methyl)-2-methoxy-benzoic acid


[0194] 5-([2-*tert*-Butoxycarbonylmethyl-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionyl]-[1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-amino)-methyl)-2-methoxybenzoic acid methyl ester was dissolved in an ice-chilled (0-10°C), mixed solvent system of THF (10 mL) and MeOH (5 mL). A LiOH·H₂O/water suspension (2.48 M; 3.77 mL) was added dropwise, then the reaction was allowed to warm to room temperature and stirred overnight. The resulting mixture was cooled in an ice bath and the basic solution was neutralized with 2N citric acid until slightly acidic. The mixture was concentrated under reduced pressure to remove the volatile materials, after which time the remaining aqueous phase was extracted with EtOAc (3 x 26 mL). These combined organic phases were dried over MgSO₄, filtered, and concentrated under reduced pressure to yield a pale yellowish white solid. This crude material was dissolved in a 10% MeOH/CH₂Cl₂ solution and adsorbed onto 30 g of silica. The adsorbed material was divided and chromatographed on an ISCO normal phase column over two runs, using a 40 g Redi-Sep column for both runs. The solvent system was a gradient MeOH/CH₂Cl₂ system as follows: Initial 100% CH₂Cl₂, 98%-92% over 40 min; 90% over 12 min, and then 88% over 13 min. The desired product eluted cleanly between 44-61 min. The desired fractions were combined and concentrated under reduced pressure to yield 5-([2-*tert*-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionyl]-[1-(4-phenyl-1 *H* imidazol-2-yl)-ethyl]-amino)-methyl)-2-methoxy-benzoic acid, Cpd **5b**, as a white solid.

STEP D. 5-([2-Amino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionyl]-[1-(4-phenyl-1 *H*-imidazol-2-yl)-ethyl]-amino)-methyl)-2-methoxy-benzoic acid

[0195] A portion of Cpd **5b** (0.27g, 0.41 mmol) was dissolved in EtOAc (39 mL)/ THF (5 mL), filtered, and subsequently treated with gaseous HCl for 15 min. After completion of the HCl addition, the reaction was slowly warmed to room temperature and a solid precipitate formed. After 5 h the reaction appeared >97% complete by LC (@214nm; 2.56 min.). The stirring was continued over 3 d, then the solid was collected and rinsed with a small amount of EtOAc. The resulting solid was dried under high vacuum under refluxing toluene for 2.5 h to yield Cpd **5c** as a white solid di-HCl salt.

Example 6**4-{2-Amino-2-([1-[4-(2-cyano-phenyl)-1*H*-imidazol-2-yl]-ethyl]-(3,4-dimethoxy-benzyl)-carbamoyl]-ethyl}-3,5-dimethyl-benzamide**

[0196]



STEP A: {1-[2-(2-Bromo-phenyl)-2-oxo-ethylcarbamoyl]-ethyl}-carbamic acid *tert*-butyl ester

[0197] Compound **6a** was prepared according to Example 3 using the appropriate reagents, starting materials and methods known to those skilled in the art.

STEP B: {1-[4-(2-Bromo-phenyl)-1*H*-imidazol-2-yl]-ethyl}-carbamic acid *tert*-butyl ester

[0198] Following the procedure described in Example 3 for the conversion of Compound **3a** to Compound **3b**, and using the appropriate reagents and methods known to those skilled in the art, Cpd **6b**, was prepared.

STEP C. 1-[4-(4-Bromo-phenyl)-1H-imidazol-2-yl]-ethylamine

[0199] Using the procedure described for the conversion of Cpd **3e** to **3f**, Compound **6c** was prepared.

STEP D. [1-[[1-[4-(2-Bromo-phenyl)-1H-imidazol-2-yl]-ethyl]-(3,4-dimethoxy-benzyl)-carbamoyl]-2-(4-carbamoyl-2,6-dimethyl-phenyl)-ethyl]-carbamic acid *tert*-butyl ester

[0200] Using the procedure described in Example 5, STEP B, and substituting 1-[4-(4-bromo-phenyl)-1H-imidazol-2-yl]-ethylamine for 1-(4-phenyl-1H-imidazol-2-yl)-ethylamine, the product was prepared.

STEP E. {2-(4-Carbamoyl-2,6-dimethyl-phenyl)-1-[[1-[4-(2-cyano-phenyl)-1H-imidazol-2-yl]-ethyl]-(3,4-dimethoxy-benzyl)-carbamoyl]-ethyl}-carbamic acid *tert*-butyl ester

[0201] To a solution of [1-[[1-[4-(2-bromo-phenyl)-1H-imidazol-2-yl]-ethyl]-(3,4-dimethoxy-benzyl)-carbamoyl]-2-(4-carbamoyl-2,6-dimethyl-phenyl)-ethyl]-carbamic acid *tert*-butyl ester (294 mg; 0.4 mmol) in DMF (2 mL) was added Zn(CN)₂ (28 mg; 0.24 mmol). The resulting mixture was degassed with Argon for 5 min, then Pd(PPh₃)₄ (92 mg; 0.08 mmol) was added neat, and the system was immediately warmed to 100°C. After heating for 6 h, the reaction was cooled to room temperature and partitioned between EtOAc and water. The organic phase was dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The crude material was subjected to reverse phase HPLC (water/ acetonitrile/ 0.1% TFA). The fractions of interest were combined, basified with saturated aqueous NaHCO₃ and extracted twice with EtOAc. The EtOAc extracts were combined, dried over Na₂SO₄, filtered, and concentrated to yield {2-(4-carbamoyl-2,6-dimethyl-phenyl)-1-[[1-[4-(2-cyano-phenyl)-1H-imidazol-2-yl]-ethyl]-(3,4-dimethoxy-benzyl)-carbamoyl]-ethyl}-carbamic acid *tert*-butyl ester (HPLC: 96% @ 254 nm and 97% @ 214 nm). This sample was of sufficient quality to use in the next reaction without further purification.


STEP F. 4-{2-Amino-2-[[1-[4-(2-cyano-phenyl)-1H-imidazol-2-yl]-ethyl]-(3,4-dimethoxy-benzyl)-carbamoyl]-ethyl}-3,5-dimethyl-benzamide

[0202] {2-(4-carbamoyl-2,6-dimethyl-phenyl)-1-[[1-[4-(2-cyano-phenyl)-1H-imidazol-2-yl]-ethyl]-(3,4-dimethoxy-benzyl)-carbamoyl]-ethyl}-carbamic acid *tert*-butyl ester may be BOC-deprotected using the procedure described in Example 3 for the conversion of Cpd **3e** to Cpd **3f** to yield the title compound.

Example 7

3-(2-{1-[[2-Amino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionyl]-(3,4-dimethoxy-benzyl)-amino]-ethyl}-1H-imidazol-4-yl)-benzoic acid

[0203]



STEP A. 1-[4-(3-Bromo-phenyl)-1H-imidazol-2-yl]-ethylamine

[0204] Using the procedure described in Example 6, and the appropriately substituted starting materials and reagents, 1-[4-(3-bromo-phenyl)-1H-imidazol-2-yl]-ethylamine was prepared.

STEP B. {1-[4-(3-Bromo-phenyl)-1H-imidazol-2-yl]-ethyl}-(3,4-dimethoxybenzyl)-amine

[0205] Using the procedure described in Example 4, and substituting 1-[4-(3-bromo-phenyl)-1H-imidazol-2-yl]-ethylamine for 1-(4-phenyl-1H-imidazol-2-yl)-ethylamine, the product was prepared.

STEP C. [1-[[1-[4-(3-Bromo-phenyl)-1H-imidazol-2-yl]-ethyl}-(3,4-dimethoxy-benzyl)-carbamoyl]-2-(4-carbamoyl-2,6-dimethyl-phenyl)-ethyl]-carbamic acid *tert*-butyl ester

[0206] Using the procedure of Example 3 for the conversion of Cpd 3d to Cpd 3e, substituting {1-[4-(3-Bromo-phenyl)-1H-imidazol-2-yl]-ethyl}-(3,4-dimethoxybenzyl)-amine for Cpd 3d and substituting 2-*tert*-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionic acid for 2-*tert*-Butoxycarbonylamino-3-(4-hydroxy-2,6-dimethyl-phenyl)-propionic acid, the product was prepared.

Step D. 3-(2-{1-[2-*tert*-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)-propionyl]-(3,4-dimethoxybenzyl)-amino}-ethyl)-1H-imidazol-4-yl)-benzoic acid

[0207] To a solution of [1-[[1-[4-(3-bromo-phenyl)-1H-imidazol-2-yl]-ethyl}-(3,4-dimethoxybenzyl)-carbamoyl]-2-(4-

carbamoyl-2,6-dimethyl-phenyl)-ethyl]-carbamic acid *tert*-butyl ester (290 mg; 0.40 mmol) in DMF (5mL) was added K_2CO_3 (262 mg; 1.9 mmol) and the resulting mixture was degassed with Argon for 5 min. At this time, $Pd(OAc)_2$ (8.9 mg; 0.04 mmol) and 1,1-bis(diphenylphosphino) ferrocene (46 mg; 0.083 mmol) were added. Carbon monoxide was then bubbled through the resulting mixture for 10 min at room temperature, the reaction was capped, and warmed to 100°C for 6 h. After cooling to room temperature the mixture was partitioned between EtOAc and water, filtered through Celite, and then separated. The aqueous phase was then washed with a second portion of EtOAc. The aqueous phase was then acidified to pH 5 with 2N citric acid and the resulting aqueous solution extracted with EtOAc (4x). These latter EtOAc extracts were combined, dried over Na_2SO_4 , filtered, and concentrated under reduced pressure to yield the crude product (HPLC: 87% at 254 nm).


Step E. 3-(2-{1-[[2-Amino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionyl]-(3,4-dimethoxy-benzyl)-amino]-ethyl}-1H-imidazol-4-yl)-benzoic acid

[0208] 3-(2-{1-[[2-*tert*-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)-propionyl]-(3,4-dimethoxy-benzyl)-amino]-ethyl}-1H-imidazol-4-yl)-benzoic acid may be BOC-de-protected using the procedure described in Example 3 for the conversion of Cpd **3e** to Cpd **3f** to yield the title compound.

Example 8

4-(2-Amino-2-{[2-hydroxy-1-(4-phenyl-1H-imidazol-2-yl)-ethyl]-isopropylcarbamoyl}-ethyl)-3,5-dimethyl-benzamide

[0209]



STEP A. [2-Benzyloxy-1-(2-oxo-2-phenyl-ethylcarbamoyl-ethyl)]-carbamic acid *tert* butyl ester

[0210] The product was prepared using the procedure described in Example 3 and substituting N- α -BOC-*L*-serine benzyl ester for N- α -CBZ-*L*-alanine.

STEP B. [2-Benzyloxy-1-(4-phenyl-1*H*-imidazol-2-yl-ethyl)]-carbamic acid *tert* butyl ester

[0211] By the procedure described in Example 3 for the conversion of Cpd 3a to Cpd 3b, [2-benzyloxy-1-(2-oxo-2-phenyl-ethylcarbamoyl-ethyl)]-carbamic acid *tert* butyl ester was converted to the product.

STEP C. [2-Benzyloxy-1-(4-phenyl-1*H*-imidazol-2-yl-ethyl)amine

[0212] [2-benzyloxy-1-(4-phenyl-1*H*-imidazol-2-yl-ethyl)]-carbamic acid *tert* butyl ester may be BOC-deprotected using the procedure described in Example 3 for the conversion of Cpd 3e to Cpd 3f to give the product.

STEP D. [2-Benzyloxy-1-(4-phenyl-1*H*-imidazol-2-yl-ethyl)]-isopropyl-amine

[0213] By the procedure described in Example 3 for the conversion of Cpd 3c to Cpd 3d, [2-benzyloxy-1-(4-phenyl-1*H*-imidazol-2-yl-ethyl)amine was converted to the product.

STEP E. [1-([2-Benzyloxy-1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-isopropylcarbamoyl)-2-(4-carbamoyl-2,6-dimethyl-phenyl)-ethyl]-carbamic acid *tert*-butyl ester


[0214] Using the procedure of Example 3 for the conversion of Cpd 3d to Cpd 3e, substituting [2-benzyloxy-1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-isopropylamine for Cpd 3d and substituting 2-*tert*-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)-propionic acid for 2-*tert*-butoxycarbonylamino-3-(4-hydroxy-2,6-dimethyl-phenyl)-propionic acid, the product was prepared.


[0215] **STEP F. 4-(2-Amino-2-([2-hydroxy-1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-isopropyl-carbamoyl)-ethyl)-3,5-dimethyl-benzamide (TFA salt).** A solution of [1-([2-benzyloxy-1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-isopropyl-carbamoyl)-2-(4-carbamoyl-2,6-dimethyl-phenyl)-ethyl]-carbamic acid *tert*-butyl ester, (0.287 g, 0.439 mmol), in chloroform (10 mL) was cooled in an ice bath and treated with 0.62 mL (4.4 mmol) of iodotrimethylsilane. The reaction, which immediately clouded, was warmed slowly to room temperature while stirring. After 16 h, the reaction was cooled in an ice bath to 5-10°C and treated with 100 mL of MeOH. The quenched mixture was stirred at 5-10°C for 30 min, removed from the ice bath and stirred for an additional 30 min, and concentrated *in vacuo* to yield an orange residue that was subjected to reverse phase HPLC (water/ acetonitrile / 0.1 % TFA). The fractions of interest were combined and the sample was lyophilized to yield 4-(2-amino-2-([2-hydroxy-1-(4-phenyl-1*H*-imidazol-2-yl)-ethyl]-isopropyl-carbamoyl)-ethyl)-3,5-dimethylbenzamide (TFA salt) as a white powder (HPLC: 99% @ 254 nm and 100% @ 214 nm)

[0216] MS (ES⁺) (relative intensity): 464.1 (100) (M+1).

Reference Example 9**(*S*)-2-*tert*-Butoxycarbonylamino-3-(2,6-dimethyl-4-trifluoromethanesulfonylphenyl)-propionic acid methyl ester**

[0217]





STEP A. (S)-2-tert-Butoxycarbonylamino-3-(2,6-dimethyl-4-trifluoromethanesulfonylphenyl)-propionic acid methyl ester

[0218] Into a cool solution of Boc-L-(2,6-diMe)Tyr-OMe (7.0 g, 21.6 mmol; Sources: Chiramer or RSP AminoAcidAnalogues) and *N*-phenyltrifluoromethanesulfonimide (7.9 g, 22.0 mmol) in dichloromethane (60 mL) was added triethylamine (3.25 mL, 23.3 mmol). The resulting solution was stirred at 0°C for 1 h and slowly warmed to room temperature. Upon completion, the reaction was quenched by addition of water. The separated organic phase was washed with 1N NaOH aqueous solution, water and dried over Na₂SO₄ overnight. After filtration and evaporation, the residue was purified by flash column chromatography (eluent: EtOAc-hexane: 3:7) to yield the desired product as a clear oil.

[0219] ¹H NMR (300 MHz, CDCl₃): δ 1.36 (9H, s), 2.39 (6H, s), 3.06 (2H, d, *J* = 7.7 Hz), 3.64 (3H, s), 4.51-4.59 (1H, m), 5.12 (1H, d, *J* = 8.5 Hz), 6.92 (2H, s)

[0220] MS (ES⁺) (relative intensity): 355.8 (100) (M-Boc)⁺.

STEP B. (S)-4-(2-tert-Butoxycarbonylamino-2-methoxycarbonylethyl)-3,5-dimethylbenzoic acid

[0221] To a suspension of (S)-2-tert-butoxycarbonylamino-3-(2,6-dimethyl-4-trifluoromethanesulfonylphenyl)-propionic acid methyl ester (9.68 g, 21.3 mmol), K₂CO₃ (14.1 g, 0.102 mol), Pd(OAc)₂ (0.48 g, 2.13 mmol) and 1,1'-bis(diphenylphosphino)ferrocene (2.56 g, 4.47 mmol) in DMF (48 mL) was bubbled in gaseous CO for 15 min. The mixture was heated to 60°C for 8 h with a CO balloon. The cool mixture was partitioned between NaHCO₃ and EtOAc, and filtered. The aqueous layer was separated, acidified with 10% citric acid aqueous solution, extracted with EtOAc, and finally dried over Na₂SO₄. Filtration and concentration of the filtrate resulted in a residue. The residue was recrystallized from EtOAc-hexanes to yield the desired product.

[0222] ¹H NMR (300 MHz, CDCl₃): δ 1.36 (9H, s), 2.42 (6H, s), 3.14 (2H, *J* = 7.4 Hz), 3.65 (3H, s), 4.57-4.59 (1H, m), 5.14 (1H, d, *J* = 8.6 Hz), 7.75 (2H, s)

[0223] MS(ES⁺) (relative intensity): 251.9 (100) (M-Boc)⁺.

STEP C. (S)-2-tert-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)propionic acid methyl ester

[0224] Into a stirring solution of (S)-4-(2-tert-butoxycarbonylamino-2-methoxycarbonylethyl)-3,5-dimethylbenzoic acid (3.00 g, 8.54 mmol), PyBOP (6.68 g, 12.8 mmol) and HOBT (1.74 g, 12.8 mmol) in DMF (36 mL) was added DIPEA (5.96 mL, 34.2 mmol) and NH₄Cl (0.92 g, 17.1 mmol). The resulting mixture was stirred at rt for 40 min before being partitioned between aqueous NH₄Cl solution and EtOAc. The separated organic phase was washed sequentially with 2N citric acid aqueous solution, saturated aqueous NaHCO₃ solution, and brine, then dried over Na₂SO₄ overnight. After filtration and concentration, the residue was purified by flash column chromatography (eluent: EtOAc) to yield the product.

[0225] ¹H NMR (300 MHz, CDCl₃): δ 1.36 (9H, s), 2.39 (6H, s), 3.11 (2H, *J* = 7.2 Hz), 3.65 (3H, s), 4.53-4.56 (1H, m), 5.12 (1H, d, *J* = 8.7 Hz), 5.65 (1H, br s), 6.09 (1H, br s), 7.46 (2H, s)

[0226] MS(ES⁺) (relative intensity): 250.9 (100) (M-Boc)⁺.

STEP D. (S)-2-tert-Butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)propionic acid

[0227] Into an ice-cooled solution of methyl ester from Step C (2.99 g, 8.54 mmol) in THF (50 mL) was added an aqueous LiOH solution (1N, 50 mL) and stirred at 0°C. Upon consumption of the starting materials, the organic solvents were removed and the aqueous phase was neutralized with cooled 1 N HCl at 0°C, and extracted with EtOAc, and dried over Na₂SO₄ overnight. Filtration and evaporation to dryness yielded the title acid (S)-2-tert-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)propionic acid.


[0228] ¹H NMR (300 MHz, DMSO-*d*₆): δ 1.30 (9H, s), 2.32 (6H, s), 2.95 (1H, dd, *J* = 8.8, 13.9 Hz), 3.10 (1H, dd, *J* = 6.2, 14.0 Hz), 4.02-4.12 (1H, m), 7.18-7.23 (2H, m), 7.48 (2H, s), 7.80 (1H, s)

[0229] MS(ES⁺) (relative intensity): 236.9 (6) (M-Boc)⁺.

Example 10

(Z)-2-Benzoyloxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)acrylic acid methyl ester

[0230]



STEP A. Trifluoromethanesulfonic acid 4-bromo-3,5-dimethyl-phenyl ester

[0231] To a cooled (0°C) solution of 4-bromo-3,5-dimethylphenol (3.05 g, 15.2 mmol) in pyridine (8 mL) was added trifluoromethanesulfonic anhydride (5.0 g, 17.7 mmol) dropwise. After completion of addition, the resulting mixture was stirred at 0°C for 15 min and at room temperature overnight. The reaction was then quenched by addition of water, then extracted with EtOAc. The EtOAc extracts were washed with water, 2N HCl (2 x), brine and dried over MgSO₄. Filtration and evaporation to dryness yield the product (10a) as a colorless oil.

[0232] ¹H NMR (300 MHz, CDCl₃): δ 2.45 (6H, s), 7.00 (2H, s).

STEP B. 4-Bromo-3,5-dimethylbenzoic acid

[0233] Into a solution of trifluoro-methanesulfonic acid 4-bromo-3,5-dimethylphenyl ester (6.57 g, 19.7 mmol) in DMF (65 mL) were added K₂CO₃ (13.1 g, 94.7 mmol), Pd(OAc)₂ (0.44 g, 1.97 mmol) and 1,1'-bis(diphenylphosphino)ferrocene (2.29 g, 4.14 mmol). The resulting mixture was bubbled in gaseous CO for 10 min and was then heated to 60°C for 7.5 h with a CO balloon. The cooled mixture was partitioned between aqueous NaHCO₃ and EtOAc, and filtered. The aqueous phase layer was separated, acidified with aqueous 6N HCl, extracted with EtOAc, and then dried over Na₂SO₄. Filtration and concentration of the filtrate resulted in the crude product (10b) as a brown residue, which was used in the next step without further purification.

STEP C. 4-Formyl-3,5-dimethyl-benzoic acid

[0234] A solution of 4-bromo-3,5-dimethylbenzoic acid (0.92 g, 4 mmol) in THF (10 mL) was cooled down to -100°C with N₂(l)-Et₂O bath and added n-butyllithium (1.6 M in hexanes, 5 mL, 8 mmol) slowly. After completion of addition, the reaction mixture was warmed to -78°C and DMF (0.74 mL, 8 mmol) was added dropwise. The resulting mixture was stirred at -78°C for 1.5 h and allowed to warm to -20°C, followed by the addition of 2N aqueous HCl (30 mL). The organic phase was separated and the aqueous phase was extracted with EtOAc, the combined organic phases were dried over MgSO₄. The solvent was removed and the resulting residue was purified by flash column chromatography (eluent: EtOAc-hexanes~1:1) to yield 4-formyl-3,5-dimethyl-benzoic acid (10c).

[0235] ¹H NMR (300 MHz, CDCl₃): δ 2.65 (6H, s), 7.82 (2H, s), 10.67 (1 H, s).

STEP D. 4-Formyl-3,5-dimethyl-benzamide

[0236] To a solution of 4-formyl-3,5-dimethyl-benzoic acid (0.15 g, 0.85 mmol) in DMF (6 mL) were added PyBOP (1.0 g, 1.92 mmol), HOBT (0.26 g, 1.92 mmol), DIPEA (0.89 mL, 5.12 mmol) and NH₄Cl (0.14 g, 2.56 mmol). The resulting mixture was stirred at room temperature for 1 h, and quenched by addition of brine, then extracted with EtOAc. The organic phase was washed with 2N aqueous HCl, saturated NaHCO₃, brine and then dried over MgSO₄. The solvent was removed to yield the crude product (**10d**), which was used in the next step without further purification.


STEP E. (Z)-2-Benzoyloxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)acrylic acid methyl ester

[0237] Into a solution of N-(benzyloxycarbonyl)-α-phosphinoglycine trimethyl ester (0.46 g, 1.4 mmol) in DCM (5 mL) was added DBU (0.21 mL, 1.4 mmol). After stirring for 10 min, a solution of the above made 4-formyl-3,5-dimethylbenzamide in DCM (5 mL) was added dropwise. The resulting mixture was stirred at room temperature for 5.5 h and the solvent was removed by rotary evaporation. The residue was dissolved in EtOAc and washed with 1 N aqueous HCl, brine and then dried over MgSO₄. The solvent was removed and the residue purified by flash column chromatography (eluent: EtOAc-hexanes~1:1) to yield (Z)-2-*tert*-butoxycarbonylamino-3-(4-carbamoyl-2,6-dimethyl-phenyl)acrylic acid methyl ester (**10e**) as a white solid.

[0238] MS(ES⁺) (relative intensity): 383.4 (10%)(M+1).

Example 11**(Z)-2-Benzoyloxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)acrylic acid methyl ester**

[0239]

**STEP A. 4-Hydroxy-3,5-dimethyl-benzamide**

[0240] Using the procedure described in Example 10, Step D, 4-hydroxy-3,5-dimethyl-benzamide (**11a**) was prepared as a yellowish solid.

[0241] ¹H NMR (300 MHz, CDCl₃): δ 2.82 (6H, s), 5.51 (1H, br s), 5.90 (1H, br s), 7.48 (2H, s);

[0242] MS(ES⁺) (relative intensity): 166.2 (8%)(M+1).

STEP B. Trifluoromethanesulfonic acid 4-carbamoyl-2,6-dimethyl-phenyl ester

[0243] Into a solution of 4-hydroxy-3,5-dimethyl-benzamide (3.72 g, 22.5 mmol) and N-phenyltrifluoromethanesulfoniunimide (9.4 g, 25 mmol) in DCM (80 mL) was added TEA (3.48 mL, 25 mmol) at room temperature, then the resulting mixture was stirred at room temperature overnight. After the reaction was quenched by addition of water, the separated

organic phase was washed with 1N NaOH, water and then dried over MgSO_4 . The solvent was removed and the residue purified by flash column chromatography (eluent: EtOAc-hexanes~1:1) to yield trifluoromethanesulfonic acid 4-carbamoyl-2,6-dimethylphenyl ester (**11b**) as a white solid.

[0244] ^1H NMR (300 MHz, CDCl_3): δ 2.42 (6H, s), 6.28 (2H, br s), 7.57 (2H, s) MS(ES^+) (relative intensity): 298.1 (63%)(M+1).

STEP C. 4-Formyl-3,5-dimethyl-benzamide

[0245] Into a solution of trifluoro-methanesulfonic acid 4-carbamoyl-2,6-dimethyl-phenyl ester (1.49 g, 5 mmol), $\text{Pd}(\text{OAc})_2$ (0.037 g, 0.15 mmol), DPPP (0.062 g, 0.15 mmol) and TEA (1.74 mL, 12.5 mmol) in DMF (25 mL) was bubbled CO (gas) for 10 min, then triethylsilane (1.6 mL, 10 mmol) was added. The resulting mixture was stirred at 75°C under a CO gas balloon for 6.5 hr. After cooling to room temperature, the reaction was quenched by addition of water, then extracted with EtOAc. The EtOAc extracts were washed with water, brine and then dried over MgSO_4 . After filtration and evaporation, the residue was purified by column chromatography (eluent, EtOAc-hexanes~1:1) to yield 4-formyl-3,5-dimethyl-benzamide (**11c**) as a yellowish solid.

[0246] ^1H NMR (300 MHz, CDCl_3): δ 2.65 (6H, s), 5.75 (1H, br s), 6.13 (1H, br s), 7.52 (2H, s), 10.64 (1H, s).

STEP D. (Z)-2-Benzyloxycarbonylamino-3-(4-carbamoyl-2,6-dimethylphenyl)acrylic acid methyl ester

[0247] The title compound was prepared as described in Example 10, Step E.


Example 12 Optical Rotation Measurements

[0248] The optical rotation of a representative sample of the compound of formula (Ia), prepared as in Example 1, was measured as $[\alpha]_D = -12$ (c 1.5, MeOH).

[0249] The optical rotation of a representative sample of the compound of formula (Ia), prepared as in Example 9, from commercially purchased (S)-N-BOC-Tyr-OMe was measured as $[\alpha]_D = -10.8$ (c 1.7, MeOH).

Claims

1. A process for the preparation of a compound of formula (I)




wherein




is C_{6-10} aryl or a heteroaryl selected from the group consisting of furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, indolinyl, benzofuryl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinoliziny, quinoliny, isoquinoliny and quinazolinyl;

each $\text{R}^{41\text{P}}$ is independently selected from C_{1-6} alkyl, C_{1-6} alkoxy or fluoro;


R^J and R^K are each independently selected from hydrogen or C_{1-4} alkyl;
 alternatively, R^J and R^K are taken together with the nitrogen atom to which they are bound to form a five to seven membered heterocyclyl;
 Pg^1 is a nitrogen protecting group;
 comprising




reacting a compound of formula (X), wherein X^P is selected from CN, $-CO_2H$, $-C(O)-Cl$ or $-C(O)-OC_{1-4}alkyl$ and wherein Y^P is selected from Br, Cl or I, to yield the corresponding compound of formula (XII);



reacting the compound of formula (XII) with a suitably substituted compound of formula (XVIII); in the presence of palladium catalyst; in the presence of an organic or inorganic base; in an organic solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XIX);



reacting the compound of formula (XIX) with hydrogen or a source of hydrogen; in the presence of a catalyst; in a solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XX);



reacting the compound of formula (XX) with an aqueous base; in an organic solvent; to yield the corresponding compound of formula (I).

2. The process as in Claim 1, wherein



is phenyl;

the phenyl is substituted with a R^{41P} group at the 2-position and a second R^{41P} group at the 4- position; each R^{41P} is independently selected from C_{1-2} alkyl, C_{1-2} alkoxy or fluoro; R^J and R^K are each independently selected from hydrogen or C_{1-4} alkyl;

alternatively, R^J and R^K are taken together with the nitrogen atom to which they are bound to form a five to seven membered heterocyclyl; and


Pg^1 is a nitrogen protecting group.

3. The process as in Claim 1, wherein



is phenyl; the phenyl is substituted with a R^{41P} group at the 2-position and a second R^{41P} group at the 4- position; each R^{41P} is methyl; R^J and R^K are each hydrogen; and Pg^1 is a t-butoxycarbonyl.

4. A process for the preparation of a compound of formula (I)



wherein




is C₆₋₁₀aryl or a heteroaryl selected from the group consisting of furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, indolyl, benzofuryl, benzothieryl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinoliziny, quinoliny, isoquinoliny and quinazoliny;


each R^{41P} is independently selected from C₁₋₆alkyl, C₁₋₆alkoxy or fluoro; R^J and R^K are each independently selected from hydrogen or C₁₋₄alkyl;

alternatively, R^J and R^K are taken together with the nitrogen atom to which they are bound to form a five to seven membered heterocyclyl;

Pg¹ is a nitrogen protecting group; comprising




reacting the compound of formula (XIX) with hydrogen or a source of hydrogen; in the presence of a catalyst; in a solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XX);




reacting the compound of formula (XX) with an aqueous base; in an organic solvent; to yield the corresponding compound of formula (I).

5. A process for the preparation of a compound of formula (Ia)




comprising




15

reacting a compound of formula (Xa), wherein X^P is selected from CN, $-\text{CO}_2\text{H}$, $-\text{C}(\text{O})-\text{Cl}$ or $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkyl}$ and wherein Y^P is selected from Br, Cl or I, to yield the corresponding compound of formula (XIIa);




30

reacting the compound of formula (XIIa) with a suitably substituted compound of formula (XVIIIa); in the presence of palladium catalyst; in the presence of an organic or inorganic base; in an organic solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XIXa);



45

reacting compound of formula (XIXa) with hydrogen gas, at a pressure sufficient to hydrogenate; in the presence of a suitable chiral catalyst; at a temperature greater than about room temperature; in an organic solvent; to yield the corresponding compound of formula (XXa);




reacting the compound of formula (XXa) with an aqueous base; in an organic solvent; to yield the corresponding compound of formula (Ia).


6. The process as in Claim 5, wherein the compound of formula (XIIa) is reacted with the compound of formula (XVIIIa) in the presence of $\text{Pd}_2(\text{dba})_3$ and $\text{P}(\text{o-toluene})_3$.

7. The process as in Claim 5, wherein the chiral catalyst is $[\text{Rh}(\text{cod})(\text{R,R-DIPAMP})]^+\text{BF}_4^-$.


8. A process for the preparation of the compound of formula (Ia)



comprising




reacting compound of formula (XIXa) with hydrogen gas, at a pressure sufficient to hydrogenate; in the presence of a suitable chiral catalyst; at a temperature greater than about room temperature; in an organic solvent; to yield the corresponding compound of formula (XXa);



reacting the compound of formula (XXa) with an aqueous base; in an organic solvent; to yield the corresponding compound of formula (Ia).

9. A process for the preparation of compounds of formula (II)



wherein



is C₆₋₁₀aryl or a heteroaryl selected from the group consisting of furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, indolinyl, benzofuryl, benzothienyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, quinoliziny, quinoliny, isoquinoliny and quinazoliny;

each R^{41P} is independently selected from C₁₋₆alkyl, C₁₋₆alkoxy or fluoro;

R^J and R^K are each independently selected from hydrogen or C₁₋₄alkyl; alternatively, R^J and R^K are taken together with the nitrogen atom to which they are bound to form a five to seven membered heterocyclyl;

R¹ is selected from the group consisting of hydrogen, C₁₋₆alkyl,

cycloalkyl, heterocyclyl, aryl(C₁₋₆)alkyl, and heteroaryl(C₁₋₆)alkyl;

wherein when R¹ is phenyl(C₁₋₆)alkyl, phenyl is optionally fused to a heterocyclyl or cycloalkyl;

wherein when R¹ is C₁₋₂alkyl, said C₁₋₂alkyl is optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₆alkoxy, aryl, cycloalkyl, heterocyclyl, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluoromethyl, and carboxy;

and further, wherein when R¹ is C₃₋₆alkyl, said C₃₋₆alkyl is optionally substituted with one to three substituents independently selected from the group consisting of C₁₋₆alkoxy, aryl, cycloalkyl, heterocyclyl, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluoromethyl, and carboxy;


wherein the cycloalkyl and heterocyclyl of C₁₋₂alkyl and C₃₋₆alkyl are optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₆alkyl, hydroxy(C₁₋₆)alkyl, C₁₋₆alkoxy, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluoromethyl, carboxy, aryl(C₁₋₆)alkoxycarbonyl, C₁₋₆alkoxycarbonyl, aminocarbonyl, C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl, and aminosulfonyl;

furthermore, wherein the cycloalkyl and heterocyclyl of R¹ are optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₆alkyl, hydroxy(C₁₋₆)alkyl, C₁₋₆alkoxy, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluoromethyl, carboxy, aryl(C₁₋₆)alkoxycarbonyl, C₁₋₆alkoxycarbonyl, aminocarbonyl, C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl, and aminosulfonyl;

furthermore, wherein the aryl and heteroaryl portion of the R¹ substituents aryl(C₁₋₆)alkyl and heteroaryl(C₁₋₆)alkyl, are optionally substituted with one to three R¹¹ substituents independently selected from the group consisting of C₁₋₆alkyl; hydroxy(C₁₋₆)alkyl; C₁₋₆alkoxy; C₆₋₁₀aryl(C₁₋₆)alkyl; C₆₋₁₀aryl(C₁₋₆)alkoxy; C₆₋₁₀aryl; heteroaryl optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₄alkyl, C₁₋₄alkoxy, and carboxy; cycloalkyl; heterocyclyl; C₆₋₁₀aryloxy; heteroaryloxy; cycloalkyloxy; heterocyclyloxy; amino; C₁₋₆alkylamino; (C₁₋₆alkyl)₂amino; C₃₋₆cycloalkylaminocarbonyl; hydroxy(C₁₋₆)alkylaminocarbonyl; C₆₋₁₀arylaminocarbonyl wherein C₆₋₁₀aryl is optionally substituted with carboxy or C₁₋₄alkoxycarbonyl; heterocyclylcarbonyl; carboxy; C₁₋₆alkylcarbonyloxy; C₁₋₆alkoxycarbonyl; C₁₋₆alkylcarbonyl; C₁₋₆alkylcarbonylamino; aminocarbonyl; C₁₋₆alkylaminocarbonyl; (C₁₋₆alkyl)₂aminocarbonyl; cyano; halogen; trifluoromethyl; trifluoromethoxy; and hydroxy;

provided that no more than one R¹¹ substituent is selected from the group consisting of C₆₋₁₀aryl(C₁₋₆)alkyl; C₆₋₁₀aryl(C₁₋₆)alkoxy; C₆₋₁₀aryl; heteroaryl optionally substituted with one to two substituents independently selected from the group consisting of C₁₋₄alkyl, C₁₋₄alkoxy, and carboxy; cycloalkyl; heterocyclyl; C₆₋₁₀aryloxy;

heteroaryloxy; cycloalkyloxy; C₆₋₁₀arylamino, heterocyclylcarbonyl; and heterocyclyloxy;
 R² is hydrogen, C₁₋₈alkyl, hydroxy(C₁₋₈)alkyl, C₆₋₁₀aryl(C₁₋₆)alkoxy(C₁₋₆)alkyl, or C₆₋₁₀aryl(C₁₋₈)alkyl;
 wherein the C₆₋₁₀aryl group in the C₆₋₁₀aryl-containing substituents of R² is optionally substituted with one to
 two substituents independently selected from the group consisting of C₁₋₆alkyl, C₁₋₆alkoxy, hydroxy, amino,
 C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, aminocarbonyl, C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl, cyano,
 fluoro, chloro, bromo, trifluoromethyl, and trifluoromethoxy; and, wherein the C₁₋₆alkyl and C₁₋₆alkoxy substituents
 of aryl are optionally substituted with hydroxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, or aryl;
 A is selected from the group consisting of aryl, ring system a-1, a-2, a-3, and a-4, optionally substituted with R⁵
 and R⁵;




wherein A-B is selected from the group consisting of N-C, C-N, N-N and C-C; wherein D-E is selected from the
 group consisting of O-C, S-C, and ON; and wherein F-G is selected from the group consisting of N-O and C-O;
 R³ is one to two substituents independently selected from the group consisting of C₁₋₆alkyl, aryl, aryl(C₁₋₆)alkyl,
 aryl(C₂₋₆)alkenyl, aryl(C₂₋₆)alkynyl, heteroaryl, heteroaryl(C₁₋₆)alkyl, heteroaryl(C₂₋₆)alkenyl, heteroar-
 yl(C₂₋₆)alkynyl, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, arylamino, heteroarylamino, aryloxy, heteroaryloxy,
 trifluoromethyl, and halogen;


wherein the aryl and heteroaryl, as well as the aryl and heteroaryl of aryl(C₁₋₆)alkyl, aryl(C₂₋₆)alkenyl, ar-
 yl(C₂₋₆)alkynyl, heteroaryl(C₁₋₆)alkyl, heteroaryl(C₂₋₆)alkenyl, heteroaryl(C₂₋₆)alkynyl, arylamino, heteroarylami-
 no, aryloxy, and heteroaryloxy, are optionally substituted with one to five fluoro substituents or one to three
 substituents independently selected from the group consisting of C₁₋₆alkyl, hydroxy(C₁₋₆)alkyl, C₁₋₆alkoxy,
 C₆₋₁₀aryl(C₁₋₆)alkyl, C₆₋₁₀aryl(C₁₋₆)alkoxy, C₆₋₁₀aryl, C₆₋₁₀aryloxy, heteroaryl(C₁₋₆)alkyl, heteroaryl(C₁₋₆)alkoxy,
 heteroaryl, heteroaryloxy, C₆₋₁₀arylamino, heteroarylamino, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, car-
 boxy(C₁₋₆)alkylamino, carboxy, C₁₋₆alkylearbonyl, C₁₋₆alkoxycarbonyl, C₁₋₆alkylcarbonylamino, aminocarbonyl,
 C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl, carboxy(C₁₋₆)alkylaminocarbonyl, cyano, halogen, trifluor-
 omethyl, trifluoromethoxy, hydroxy, C₁₋₆alkylsulfonyl, and C₁₋₆alkylsulfonylamino; provided that not more than
 one R³ is selected from the group consisting of aryl, heteroaryl, aryl(C₁₋₆)alkyl, aryl(C₂₋₆)alkenyl, aryl(C₂₋₆)alky-
 nyl, heteroaryl, heteroaryl(C₁₋₆)alkyl, heteroaryl(C₂₋₆)alkenyl, heteroaryl(C₂₋₆)alkynyl, arylamino, heteroarylami-
 no, aryloxy, and heteroaryloxy;

and wherein C₁₋₆alkyl, and C₁₋₆alkyl of aryl(C₁₋₆)alkyl and heteroaryl(C₁₋₆)alkyl, are optionally substituted with
 a substituent selected from the group consisting of hydroxy, carboxy, C₁₋₄alkoxycarbonyl, amino, C₁₋₆alkylamino,
 (C₁₋₆alkyl)₂amino, aminocarbonyl, (C₁₋₄)alkylaminocarbonyl, di(C₁₋₄)alkylaminocarbonyl, aryl, heteroaryl, ar-
 ylamino, heteroarylamino, aryloxy, heteroaryloxy, aryl(C₁₋₄)alkoxy, and heteroaryl(C₁₋₄)alkoxy;

R⁵ is a substituent on a nitrogen atom of ring A selected from the group consisting of hydrogen and C₁₋₄alkyl;
 R^a and R^b are independently selected from the group consisting of hydrogen, C₁₋₆alkyl, and C₁₋₆alkoxycarbonyl;
 alternatively, when R^a and R^b are each other than hydrogen, R^a and R^b are optionally taken together with the
 nitrogen atom to which they are both attached to form a five to eight membered monocyclic ring;
 and pharmaceutically acceptable enantiomers, diastereomers, racemates, and salts thereof;
 comprising




reacting a compound of formula (X), wherein X^P is selected from CN, $-\text{CO}_2\text{H}$, $-\text{C}(\text{O})-\text{Cl}$ or $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkyl}$ and wherein Y^P is selected from Br, Cl or I, to yield the corresponding compound of formula (XII);



25

30


reacting the compound of formula (XII) with a suitably substituted compound of formula (XVIII), wherein Pg^1 is a nitrogen protecting group; in the presence of palladium catalyst; in the presence of an organic or inorganic base; in an organic solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XIX);




45

50

reacting the compound of formula (XIX) with hydrogen or a source of hydrogen; in the presence of a catalyst; in a solvent; at a temperature greater than about room temperature; to yield the corresponding compound of formula (XX);



reacting the compound of formula (XX) with an aqueous base; in an organic solvent; to yield the corresponding compound of formula (I);



reacting the compound of formula (I), to yield the corresponding compound of formula (II).

10. The process as in Claim 9, wherein



is phenyl;

the phenyl is substituted with a R^{41P} group at the 2-position and a second R^{41P} group at the 4- position;

each R^{41P} is independently selected from C_{1-2} alkyl, C_{1-2} alkoxy or fluoro;

R^J and R^K are each independently selected from hydrogen or C_{1-4} alkyl;

alternatively, R^J and R^K are taken together with the nitrogen atom to which they are bound to form a five to seven membered heterocyclyl; and

Pg^1 is a nitrogen protecting group.

11. The process as in Claim 9, wherein



is phenyl, the phenyl is substituted with a R^{41P} group at the 2-position and a second R^{41P} group at the 4- position;

each R^{41P} is methyl; R^J and R^K are each hydrogen; and Pg^1 is a t-butoxycarbonyl.


12. The process as in any one of Claims 1-3, 5-7 and 9-11, wherein X^P is selected from CN, $-C(O)-Cl$ or $-C(O)-OC_{1-4}$ alkyl.

13. The process as in any one of Claims 1-3, 5-7 and 9-11, wherein Y^P is selected from Cl or I.

Patentansprüche

1. Verfahren zur Herstellung einer Verbindung der

Formel (I)




worin




für C₆₋₁₀-Aryl oder ein Heteroaryl aus der Gruppe bestehend aus Furyl, Thienyl, Pyrrolyl, Oxazolyl, Thiazolyl, Imidazolyl, Pyrazolyl, Pyridinyl, Pyrimidinyl, Pyrazinyl, Indolyl, Isoindolyl, Indolinyl, Benzofuryl, Benzothienyl, Benzimidazolyl, Benzothiazolyl, Benzoxazolyl, Chinolizinyll, Chinolinyll, Isochinolinyll und Chinazolinyll steht;

R^{41P} jeweils unabhängig aus C₁₋₆-Alkyl, C₁₋₆-Alkoxy oder Fluor ausgewählt ist;


R^J und R^K jeweils unabhängig aus Wasserstoff oder C₁₋₄-Alkyl ausgewählt sind; alternativ dazu R^J und R^K zusammen mit dem Stickstoffatom, an das sie gebunden sind, ein fünf- bis siebengliedriges Heterocyclyl bilden; Pg¹ für eine Stickstoff-Schutzgruppe steht; bei dem man




eine Verbindung der Formel (X), worin X^P aus CN, -CO₂H, -C(O)-Cl oder -C(O)-O-C₁₋₄-Alkyl ausgewählt ist und worin Y^P aus Br, Cl oder I ausgewählt ist, zu der entsprechenden Verbindung der Formel (XII) umsetzt;



die Verbindung der Formel (XII) in Gegenwart von Palladiumkatalysator in Gegenwart einer organischen oder anorganischen Base in einem organischen Lösungsmittel bei einer Temperatur, die über etwa Raumtemperatur liegt, mit einer geeignet substituierten Verbindung der Formel (XVIII) zu der entsprechenden Verbindung der Formel (XIX) umsetzt;



die Verbindung der Formel (XIX) in Gegenwart eines Katalysators in einem Lösungsmittel bei einer Temperatur, die über etwa Raumtemperatur liegt, mit Wasserstoff oder einer Wasserstoffquelle zu der entsprechenden Verbindung der Formel (XX) umsetzt;



die Verbindung der Formel (XX) in einem organischen Lösungsmittel mit einer wässrigen Base zu der entsprechenden Verbindung der Formel (I) umsetzt.

2. Verfahren nach Anspruch 1, bei dem



für Phenyl steht;

das Phenyl in der 2-Position mit einer R^{41P} -Gruppe und in der 4-Position mit einer zweiten R^{41P} -Gruppe substituiert ist;

R^{41P} jeweils unabhängig aus C_{1-2} -Alkyl, C_{1-2} -Alkoxy oder Fluor ausgewählt ist;


R^J und R^K jeweils unabhängig aus Wasserstoff oder C_{1-4} -Alkyl ausgewählt sind; alternativ dazu R^J und R^K zusammen mit dem Stickstoffatom, an das sie gebunden sind, ein fünf- bis siebengliedriges Heterocycl cycl bilden; und Pg^1 für eine Stickstoff-Schutzgruppe steht.

3. Verfahren nach Anspruch 1, bei dem



für Phenyl steht; das Phenyl in der 2-Position mit einer R^{41P} -Gruppe und in der 4-Position mit einer zweiten R^{41P} -Gruppe substituiert ist; R^{41P} jeweils für Methyl steht; R^J und R^K jeweils für Wasserstoff stehen und Pg^1 für ein t-Butoxycarbonyl steht.

4. Verfahren zur Herstellung einer Verbindung der Formel (I)



worin




für C_{6-10} -Aryl oder ein Heteroaryl aus der Gruppe bestehend aus Furyl, Thienyl, Pyrrolyl, Oxazolyl, Thiazolyl, Imidazolyl, Pyrazolyl, Pyridinyl, Pyrimidinyl, Pyrazinyl, Indolyl, Isoindolyl, Indoliny, Benzofuryl, Benzothienyl, Benzimidazolyl, Benzothiazolyl, Benzoxazolyl, Chinoliziny, Chinoliny, Isochinoliny und Chinazoliny steht;


R^{41P} jeweils unabhängig aus C_{1-6} -Alkyl, C_{1-6} -Alkoxy oder Fluor ausgewählt ist;

R^J und R^K jeweils unabhängig aus Wasserstoff oder C_{1-4} -Alkyl ausgewählt sind; alternativ dazu R^J und R^K zusammen mit dem Stickstoffatom, an das sie gebunden sind, ein fünf- bis siebengliedriges Heterocycl cycl bilden; Pg^1 für eine Stickstoff-Schutzgruppe steht;

bei dem man




die Verbindung der Formel (XIX) in Gegenwart eines Katalysators in einem Lösungsmittel bei einer Temperatur, die über etwa Raumtemperatur liegt, mit Wasserstoff oder einer Wasserstoffquelle zu der entsprechenden Verbindung der Formel (XX) umsetzt;




die Verbindung der Formel (XX) in einem organischen Lösungsmittel mit einer wässrigen Base zu der entsprechenden Verbindung der Formel (I) umsetzt.


5. Verfahren zur Herstellung einer Verbindung der Formel (Ia)




bei dem man




eine Verbindung der Formel (Xa), worin X^P aus CN , $-\text{CO}_2\text{H}$, $-\text{C}(\text{O})-\text{Cl}$ oder $-\text{C}(\text{O})-\text{O}-\text{C}_{1-4}\text{-Alkyl}$ ausgewählt ist und worin Y^P aus Br , Cl oder I ausgewählt ist, zu der entsprechenden Verbindung der Formel (XIIa) umsetzt;



die Verbindung der Formel (XIIa) in Gegenwart von Palladiumkatalysator in Gegenwart einer organischen oder anorganischen Base in einem organischen Lösungsmittel bei einer Temperatur, die über etwa Raumtemperatur liegt, mit einer geeignet substituierten Verbindung der Formel (XVIIIa) zu der entsprechenden Verbindung der Formel (XIXa) umsetzt;




die Verbindung der Formel (XIXa) in Gegenwart eines geeigneten chiralen Katalysators bei einer Temperatur, die über etwa Raumtemperatur liegt, in einem organischen Lösungsmittel mit Wasserstoffgas bei einem zur Hydrierung ausreichenden Druck zu der entsprechenden Verbindung der Formel (XXa) umsetzt;




die Verbindung der Formel (XXa) in einem organischen Lösungsmittel mit einer wässrigen Base zu der entsprechenden Verbindung der Formel (Ia) umsetzt.


6. Verfahren nach Anspruch 5, bei dem man die Verbindung der Formel (XIIa) in Gegenwart von $\text{Pd}_2(\text{dba})_3$ und $\text{P}(\text{o-Toluol})_3$ mit der Verbindung der Formel (XVIIa) umsetzt.
7. Verfahren nach Anspruch 5, bei dem es sich bei dem chiralen Katalysator um $[\text{Rh}(\text{cod})(\text{R}, \text{R-DIPAMP})]^+\text{BF}_4^-$ handelt.
8. Verfahren zur Herstellung der Verbindung der Formel (Ia)



bei dem man




eine Verbindung der Formel (XIXa) in Gegenwart eines geeigneten chiralen Katalysators bei einer Temperatur, die über etwa Raumtemperatur liegt, in einem organischen Lösungsmittel mit Wasserstoffgas bei einem zur Hydrierung ausreichenden Druck zu der entsprechenden Verbindung der Formel (XXa) umsetzt;



die Verbindung der Formel (XXa) in einem organischen Lösungsmittel mit einer wässrigen Base zu der entsprechenden Verbindung der Formel (Ia) umsetzt.

9. Verfahren zur Herstellung von Verbindungen der Formel (II) worin



für C₆₋₁₀-Aryl oder ein Heteroaryl aus der Gruppe bestehend aus Furyl, Thienyl, Pyrrolyl, Oxazolyl, Thiazolyl, Imidazolyl, Pyrazolyl, Pyridinyl, Pyrimidinyl, Pyrazinyl, Indolyl, Isoindolyl, Indolinyl, Benzofuryl, Benzothienyl, Benzimidazolyl, Benzothiazolyl, Benzoxazolyl, Chinolizinyl, Chinoliny, Isochinoliny und Chinazoliny steht;

R^{41P} jeweils unabhängig aus C₁₋₆-Alkyl, C₁₋₆-Alkoxy oder Fluor ausgewählt ist;

R^J und R^K jeweils unabhängig aus Wasserstoff oder C₁₋₄-Alkyl ausgewählt sind; alternativ dazu R^J und R^K zusammen mit dem Stickstoffatom, an das sie gebunden sind, ein fünf- bis siebengliedriges Heterocyclyl bilden; R¹ aus der Gruppe bestehend aus Wasserstoff, C₁₋₆-Alkyl, Cycloalkyl, Heterocyclyl, Aryl-(C₁₋₆)-alkyl und Heteroaryl-(C₁₋₆)-alkyl ausgewählt ist;

wobei dann, wenn R¹ für Phenyl-(C₁₋₆)-alkyl steht, Phenyl gegebenenfalls mit einem Heterocyclyl oder Cycloalkyl kondensiert ist;

wobei dann, wenn R¹ für C₁₋₂-Alkyl steht, das C₁₋₂-Alkyl gegebenenfalls durch einen oder zwei Substituenten, die unabhängig voneinander aus der Gruppe bestehend aus C₁₋₆-Alkoxy, Aryl, Cycloalkyl, Heterocyclyl, Hydroxy, Cyano, Amino, C₁₋₆-Alkyl-amino, (C₁₋₆-Alkyl)₂amino, Trifluormethyl und Carboxy ausgewählt sind, substituiert ist;

und wobei weiterhin dann, wenn R¹ für C₃₋₆-Alkyl steht, das C₃₋₆-Alkyl gegebenenfalls durch einen bis drei Substituenten, unabhängig voneinander aus der Gruppe bestehend aus C₁₋₆-Alkoxy, Aryl, Cycloalkyl, Heterocyclyl, Hydroxy, Cyano, Amino, C₁₋₆-Alkylamino, (C₁₋₆-Alkyl)₂amino, Trifluormethyl und Carboxy ausgewählt sind, substituiert ist; wobei das Cycloalkyl und Heterocyclyl von C₁₋₂-Alkyl und C₃₋₆-Alkyl gegebenenfalls durch einen oder zwei Substituenten, unabhängig voneinander aus der Gruppe bestehend aus C₁₋₆-Alkyl, Hydroxy-(C₁₋₆)-alkyl, C₁₋₆-Alkoxy, Hydroxy, Cyano, Amino, C₁₋₆-Alkylamino, (C₁₋₆-Alkyl)₂amino, Trifluormethyl, Carboxy, Aryl-(C₁₋₆)-alkoxycarbonyl, C₁₋₆-Alkoxycarbonyl, Aminocarbonyl, C₁₋₆-Alkylaminocarbonyl, (C₁₋₆-Al-

kyl)₂aminocarbonyl und Aminosulfonyl ausgewählt sind, substituiert sind;

wobei weiterhin das Cycloalkyl und Heterocyclyl von R¹ gegebenenfalls durch einen oder zwei Substituenten, die unabhängig voneinander aus der Gruppe bestehend aus C₁₋₆-Alkyl, Hydroxy-(C₁₋₆)-alkyl, C₁₋₆-Alkoxy, Hydroxy, Cyano, Amino, C₁₋₆-Alkylamino, (C₁₋₆-Alkyl)₂amino, Trifluormethyl, Carboxy, Aryl-(C₁₋₆)-alkoxycarbonyl, C₁₋₆-Alkoxycarbonyl, Aminocarbonyl, C₁₋₆-Alkylaminocarbonyl, (C₁₋₆-Alkyl)₂aminocarbonyl und Aminosulfonyl ausgewählt sind, substituiert sind;


wobei weiterhin der Aryl- und Heteroarylteil der R¹-Substituenten Aryl-(C₁₋₆)-alkyl und Heteroaryl-(C₁₋₆)-alkyl gegebenenfalls durch einen bis drei R¹¹-Substituenten, die unabhängig voneinander aus der Gruppe bestehend aus C₁₋₆-Alkyl; Hydroxy-(C₁₋₆)-alkyl; C₁₋₆-Alkoxy; C₆₋₁₀-Aryl-(C₁₋₆)-alkyl; C₆₋₁₀-Aryl-(C₁₋₆)-alkoxy; C₆₋₁₀-Aryl; Heteroaryl, das gegebenenfalls durch einen oder zwei Substituenten, die unabhängig voneinander aus der Gruppe bestehend aus C₁₋₄-Alkyl, C₁₋₄-Alkoxy und Carboxy ausgewählt sind, substituiert ist; Cycloalkyl; Heterocyclyl; C₆₋₁₀-Aryloxy; Heteroaryloxy; Cycloalkyloxy; Heterocyclyloxy; Amino; C₁₋₆-Alkylamino; (C₁₋₆-Alkyl)₂amino; C₃₋₆-Cycloalkylaminocarbonyl; Hydroxy-(C₁₋₆)-alkylaminocarbonyl; C₆₋₁₀-Arylamino-carbonyl, wobei C₆₋₁₀-Aryl gegebenenfalls durch Carboxy oder C₁₋₄-Alkoxycarbonyl substituiert ist; Heterocyclylcarbonyl; Carboxy; C₁₋₆-Alkylcarbonyloxy; C₁₋₆-Alkoxycarbonyl; C₁₋₆-Alkylcarbonyl; C₁₋₆-Alkylcarbonylamino; Aminocarbonyl; C₁₋₆-Alkylaminocarbonyl; (C₁₋₆-Alkyl)₂aminocarbonyl; Cyano; Halogen; Trifluormethyl; Trifluormethoxy und Hydroxy ausgewählt sind, substituiert ist;

mit der Maßgabe, dass nicht mehr als ein R¹¹-Substituent aus der Gruppe bestehend aus C₆₋₁₀-Aryl-(C₁₋₆)-alkyl; C₆₋₁₀-Aryl-(C₁₋₆)-alkoxy; C₆₋₁₀-Aryl; Heteroaryl, das gegebenenfalls durch einen oder zwei Substituenten, die unabhängig voneinander aus der Gruppe bestehend aus C₁₋₄-Alkyl, C₁₋₄-Alkoxy und Carboxy ausgewählt sind, substituiert ist; Cycloalkyl; Heterocyclyl; C₆₋₁₀-Aryloxy; Heteroaryloxy; Cycloalkyloxy; C₆₋₁₀-Arylamino-carbonyl; Heterocyclylcarbonyl und Heterocyclyloxy ausgewählt ist;

R² für Wasserstoff, C₁₋₈-Alkyl, Hydroxy-(C₁₋₈)-alkyl, C₆₋₁₀-Aryl-(C₁₋₆)-alkoxy-(C₁₋₆)-alkyl oder C₆₋₁₀-Aryl-(C₁₋₈)-alkyl steht;

wobei die C₆₋₁₀-Arylgruppe in den C₆₋₁₀-Aryl enthaltenden Substituenten von R² gegebenenfalls durch einen oder zwei Substituenten, unabhängig voneinander aus der Gruppe bestehend aus C₁₋₆-Alkyl, C₁₋₆-Alkoxy, Hydroxy, Amino, C₁₋₆-Alkylamino, (C₁₋₆-Alkyl)₂amino, Aminocarbonyl, C₁₋₆-Alkylamino-carbonyl, (C₁₋₆-Alkyl)₂aminocarbonyl, Cyano, Fluor, Chlor, Brom, Trifluormethyl und Trifluormethoxy ausgewählt sind, substituiert ist; und wobei die C₁₋₆-Alkyl- und C₁₋₆-Alkoxysubstituenten von Aryl gegebenenfalls durch Hydroxy, Amino, C₁₋₆-Alkyl-amino, (C₁₋₆-Alkyl)₂amino oder Aryl substituiert sind;

A aus der Gruppe bestehend aus Aryl, Ringsystem a-1, a-2, a-3 und a-4, das gegebenenfalls durch R³ und R⁵ substituiert ist, ausgewählt ist;



wobei A-B aus der Gruppe bestehend aus N-C, C-N, N-N und C-C ausgewählt ist; wobei D-E aus der Gruppe bestehend aus O-C, S-C und O-N ausgewählt ist und wobei F-G aus der Gruppe bestehend aus N-O und C-O ausgewählt ist;

R³ für einen oder zwei Substituenten, die unabhängig voneinander aus der Gruppe bestehend aus C₁₋₆-Alkyl, Aryl, Aryl-(C₁₋₆)-alkyl, Aryl-(C₂₋₆)-alkenyl, Aryl-(C₂₋₆)-alkinyl, Heteroaryl, Heteroaryl-(C₁₋₆)-alkyl, Heteroaryl-(C₂₋₆)-alkenyl, Heteroaryl-(C₂₋₆)-alkinyl, Amino, C₁₋₆-Alkylamino, (C₁₋₆-Alkyl)₂amino, Arylamino, Heteroarylamino, Aryloxy, Heteroaryloxy, Trifluormethyl und Halogen ausgewählt sind, steht;

wobei das Aryl und Heteroaryl sowie das Aryl und Heteroaryl von Aryl-(C₁₋₆)-alkyl, Aryl-(C₂₋₆)-alkenyl, Aryl-(C₂₋₆)-alkinyl, Heteroaryl-(C₁₋₆)-alkyl, Heteroaryl-(C₂₋₆)-alkenyl, Heteroaryl-(C₂₋₆)-alkinyl, Arylamino, Heteroarylamino, Aryloxy und Heteroaryloxy gegebenenfalls durch einen bis fünf Fluorsubstituenten oder einen bis drei Substituenten, unabhängig voneinander aus der Gruppe bestehend aus C₁₋₆-Alkyl, Hydroxy-(C₁₋₆)-alkyl, C₁₋₆-Alkoxy, C₆₋₁₀-Aryl-(C₁₋₆)-alkyl, C₆₋₁₀-Aryl-(C₁₋₆)-alkoxy, C₆₋₁₀-Aryl, C₆₋₁₀-Aryloxy, Heteroaryl-(C₁₋₆)-alkyl, Heteroaryl-(C₁₋₆)-alkoxy, Heteroaryl, Heteroaryloxy, C₆₋₁₀-Arylamino, Heteroarylamino, Amino, C₁₋₆-Alkylamino, (C₁₋₆-Alkyl)₂amino, Carboxy-(C₁₋₆)-alkylamino, Carboxy, C₁₋₆-Alkylcarbonyl, C₁₋₆-Alkoxycarbonyl, C₁₋₆-Alkylcarbonylamino, Aminocarbonyl, C₁₋₆-Alkylaminocarbonyl, (C₁₋₆-Alkyl)₂aminocarbonyl, Carboxy-(C₁₋₆)-alkylamino-carbonyl, Cyano, Halogen, Trifluormethyl, Trifluormethoxy, Hydroxy, C₁₋₆-Alkylsulfonyl und C₁₋₆-Alkyl-

sulfonylamino ausgewählt sind, substituiert sind;


mit der Maßgabe, dass nicht mehr als ein R^3 aus der Gruppe bestehend aus Aryl, Heteroaryl, Aryl-(C₁₋₆)-alkyl, Aryl-(C₂₋₆)-alkenyl, Aryl-(C₂₋₆)-alkinyl, Heteroaryl, Heteroaryl-(C₁₋₆)-alkyl, Heteroaryl-(C₂₋₆)-alkenyl, Heteroaryl-(C₂₋₆)-alkinyl, Arylamino, Heteroarylamino, Aryloxy und Heteroaryloxy ausgewählt ist;

und wobei C₁₋₆-Alkyl und C₁₋₆-Alkyl von Aryl-(C₁₋₆)-alkyl und Heteroaryl-(C₁₋₆)-alkyl gegebenenfalls durch einen Substituenten, der aus der Gruppe bestehend aus Hydroxy, Carboxy, C₁₋₄-Alkoxycarbonyl, Amino, C₁₋₆-Alkylamino, (C₁₋₆-Alkyl)₂amino, Aminocarbonyl, (C₁₋₄)-Alkylaminocarbonyl, Di-(C₁₋₄)alkylaminocarbonyl, Aryl, Heteroaryl, Arylamino, Heteroarylamino, Aryloxy, Heteroaryloxy, Aryl-(C₁₋₄)-alkoxy und Heteroaryl-(C₁₋₄)-alkoxy ausgewählt ist, substituiert sind;


R^5 für einen Substituenten an einem Stickstoffatom des Rings A, der aus der Gruppe bestehend aus Wasserstoff und C₁₋₄-Alkyl ausgewählt ist, steht;

R^a und R^b unabhängig voneinander aus der Gruppe bestehend aus Wasserstoff, C₁₋₆-Alkyl und C₁₋₆-Alkoxycarbonyl ausgewählt sind; alternativ dazu dann, wenn R^a und R^b jeweils von Wasserstoff verschieden sind, R^a und R^b gegebenenfalls zusammen mit dem Stickstoffatom, an das sie beide gebunden sind, einen fünf- bis achtgliedrigen monocyclischen Ring bilden;


und pharmazeutisch unbedenklichen Enantiomeren, Diastereomeren, Racematen und Salzen davon; bei dem man




eine Verbindung der Formel (X), worin X^P aus CN, -CO₂H, -C(O)-Cl oder -C(0) -O-C₁₋₄-Alkyl ausgewählt ist und worin Y^P aus Br, Cl oder I ausgewählt ist, zu der entsprechenden Verbindung der Formel (XII) umsetzt;




die Verbindung der Formel (XII) in Gegenwart von Palladiumkatalysator in Gegenwart einer organischen oder anorganischen Base in einem organischen Lösungsmittel bei einer Temperatur, die über etwa Raumtemperatur liegt, mit einer geeignet substituierten Verbindung der Formel (XVIII), wobei Pg^1 für eine Stickstoff-Schutzgruppe steht, zu der entsprechenden Verbindung der Formel (XIX) umsetzt;



die Verbindung der Formel (XIX) in Gegenwart eines Katalysators in einem Lösungsmittel bei einer Temperatur, die über etwa Raumtemperatur liegt, mit Wasserstoff oder einer Wasserstoffquelle zu der entsprechenden Verbindung der Formel (XX) umsetzt;



die Verbindung der Formel (XX) in einem organischen Lösungsmittel mit einer wässrigen Base zu der entsprechenden Verbindung der Formel (I) umsetzt;



die Verbindung der Formel (I) zu der entsprechenden Verbindung der Formel (II) umsetzt.

10. Verfahren nach Anspruch 9, bei dem



für Phenyl steht;

das Phenyl in der 2-Position mit einer R^{41P} -Gruppe und in der 4-Position mit einer zweiten R^{41P} -Gruppe substituiert ist;

R^{41P} jeweils unabhängig aus C_{1-2} -Alkyl, C_{1-2} -Alkoxy oder Fluor ausgewählt ist;

R^J und R^K jeweils unabhängig aus Wasserstoff oder C_{1-4} -Alkyl ausgewählt sind; alternativ dazu R^J und R^K zusammen mit dem Stickstoffatom, an das sie gebunden sind, ein fünf- bis siebengliedriges Heterocyclyl bilden; und
 Pg^1 für eine Stickstoff-Schutzgruppe steht.

11. Verfahren nach Anspruch 9, bei dem




für Phenyl steht; das Phenyl in der 2-Position mit einer R^{41P} -Gruppe und in der 4-Position mit einer zweiten R^{41P} -Gruppe substituiert ist; R^{41P} jeweils für Methyl steht; R^J und R^K jeweils für Wasserstoff stehen und Pg^1 für ein t-Butoxycarbonyl steht.

12. Verfahren nach einem der Ansprüche 1-3, 5-7 und 9-11, bei dem X^P aus CN, $-C(O)-Cl$ oder $-C(O)-O-C_{1-4}$ -Alkyl ausgewählt ist.

13. Verfahren nach einem der Ansprüche 1-3, 5-7 und 9-11, bei dem Y^P aus Cl oder I ausgewählt ist.

Revendications

1. Procédé de préparation d'un composé de formule (I)



dans laquelle




est C_{6-10} aryle ou un groupement hétéroaryle choisi dans le groupe constitué par furyle, thiényle, pyrrolyle, oxazolyle, thiazolyle, imidazolyle, pyrazolyle, pyridinyle, pyrimidinyle, pyrazinyle, indolyle, isoindolyle, indolinyle, benzofuryle, benzothiényle, benzimidazolyle, benzothiazolyle, benzoxazolyle, quinolizinyle, quinoléinyle, isoquinoléinyle et quinazolinyle ;


chaque R^{41P} est choisi indépendamment parmi C_{1-6} -alkyle, C_{1-6} alcoxy ou fluoro ;

R^J et R^K sont choisis chacun indépendamment parmi hydrogène ou C_{1-4} alkyle ; de manière alternative, R^J et R^K sont pris ensemble avec l'atome d'azote auquel ils sont liés pour former hétérocyclyle de cinq à sept chaînons ;


Pg^1 est un groupement protecteur d'azote ; comprenant




la réaction d'un composé de formule (X), où X^P est choisi parmi CN, $-\text{CO}_2\text{H}$, $-\text{C}(\text{O})-\text{Cl}$ ou $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkyle}$ et où Y^P est choisi parmi Br, Cl ou I, pour conduire au composé correspondant de formule (XII) ;



la réaction du composé de formule (XII) avec un composé de formule (XVIII) convenablement substitué, en présence d'un catalyseur à base de palladium ; en présence d'une base organique ou inorganique ; dans un solvant organique ; à une température supérieure à environ la température ambiante ; pour conduire au composé correspondant de formule (XIX) ;



la réaction du composé de formule (XIX) avec de l'hydrogène ou une source d'hydrogène ; en présence d'un catalyseur ; dans un solvant ; à une température supérieure à environ la température ambiante ; pour conduire au composé correspondant de formule (XX) ;



la réaction du composé de formule (XX) avec une base aqueuse ; dans un solvant organique ; pour conduire au composé correspondant de formule (I).

2. Procédé selon la revendication 1, dans lequel



est phényle ;

le groupement phényle est substitué par un groupement R^{41P} au niveau de la position 2 et un deuxième groupement R^{41P} au niveau de la position 4 ;

chaque R^{41P} est choisi indépendamment parmi C_{1-2} -alkyle, C_{1-2} alcoxy ou fluoro ;


R^J et R^K sont choisis chacun indépendamment parmi hydrogène ou C_{1-4} alkyle ; de manière alternative, R^J et R^K sont pris ensemble avec l'atome d'azote auquel ils sont liés pour former hétérocyclyle de cinq à sept chaînons ; et Pg^1 est un groupement protecteur d'azote.

3. Procédé selon la revendication 1, dans lequel



est phényle ; le groupement phényle est substitué par un groupement R^{41P} au niveau de la position 2 et un deuxième groupement R^{41P} au niveau de la position 4 ; chaque R^{41P} est méthyle ; R^J et R^K sont chacun hydrogène ; et Pg^1 est t-butoxycarbonyl.

4. Procédé de préparation d'un composé de formule (I)




dans laquelle




est C_{6-10} aryle ou un groupement hétéroaryle choisi dans le groupe constitué par furyle, thiényle, pyrrolyle, oxazolyle, thiazolyle, imidazolyle, pyrazolyle, pyridinyle, pyrimidinyle, pyrazinyle, indolyle, isoindolyle, indolinyle, benzofuryle, benzothiényle, benzimidazolyle, benzothiazolyle, benzoxazolyle, quinolizinyne, quinoléinyle, isoquinoléinyle et quinazolinyle ;

chaque R^{41P} est choisi indépendamment parmi C_{1-6} -alkyle, C_{1-6} alcoxy ou fluoro ;

R^J et R^K sont choisis chacun indépendamment parmi hydrogène ou C_{1-4} alkyle ; de manière alternative, R^J et R^K sont pris ensemble avec l'atome d'azote auquel ils sont liés pour former hétérocyclyle de cinq à sept chaînons ; Pg^1 est un groupement protecteur d'azote ; comprenant




la réaction du composé de formule (XIX) avec de l'hydrogène ou une source d'hydrogène ; en présence d'un catalyseur ; dans un solvant ; à une température supérieure à environ la température ambiante ; pour conduire au composé correspondant de formule (XX) ;




la réaction du composé de formule (XX) avec une base aqueuse ; dans un solvant organique ; pour conduire au composé correspondant de formule (I).


5. Procédé de préparation d'un composé de formule (Ia)




comprenant




la réaction d'un composé de formule (Xa), où X^P est choisi parmi CN, $-\text{CO}_2\text{H}$, $-\text{C}(\text{O})-\text{Cl}$ ou $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkyle}$ et où Y^P est choisi parmi Br, Cl ou I, pour conduire au composé correspondant de formule (XIIa) ;



la réaction du composé de formule (XIIa) avec un composé de formule (XVIIIa) convenablement substitué, en présence d'un catalyseur à base de palladium ; en présence d'une base organique ou inorganique ; dans un solvant organique ; à une température supérieure à environ la température ambiante ; pour conduire au composé correspondant de formule (XIXa) ;




la réaction du composé de formule (XIXa) avec de l'hydrogène gazeux, à une pression suffisante pour une hydrogénation ; en présence d'un catalyseur chiral convenable ; à une température supérieure à environ la température ambiante ; dans un solvant organique ; pour conduire au composé correspondant de formule (XXa) ;




la réaction du composé de formule (XXa) avec une base aqueuse ; dans un solvant organique ; pour conduire au composé correspondant de formule (Ia).


6. Procédé selon la revendication 5, dans lequel le composé de formule (XIIa) est réagi avec le composé de formule (XVIIIa) en présence de $\text{Pd}_2(\text{dba})_3$ et de $\text{P}(\text{o-toluène})_3$.
7. Procédé selon la revendication 5, dans lequel le catalyseur chiral est $[\text{Rh}(\text{cod})(\text{R,R-DIPAMP})]^+\text{BF}_4^-$.
8. Procédé de préparation du composé de formule (Ia)



comprenant




la réaction du composé de formule (XIXa) avec de l'hydrogène gazeux, à une pression suffisante pour une hydrogénation ; en présence d'un catalyseur chiral convenable ; à une température supérieure à environ la température ambiante ; dans un solvant organique ; pour conduire au composé correspondant de formule (XXa) ;



la réaction du composé de formule (XXa) avec une base aqueuse ; dans un solvant organique ; pour conduire au composé correspondant de formule (Ia).

9. Procédé de préparation de composés de formule (II)



dans laquelle



est C₆₋₁₀aryle ou un groupement hétéroaryle choisi dans le groupe constitué par furyle, thiényle, pyrrolyle, oxazolyle, thiazolyle, imidazolyle, pyrazolyle, pyridinyle, pyrimidinyle, pyrazinyle, indolyle, isoindolyle, indolinyle, benzofuryle, benzothiényle, benzimidazolyle, benzothiazolyle, benzoxazolyle, quinolizinyne, quinoléinyle, isoquinoléinyle et quinazolinyle ;

chaque R^{41P} est choisi indépendamment parmi C₁₋₆-alkyle, C₁₋₆alcoxy ou fluoro ;

R^J et R^K sont choisis chacun indépendamment parmi hydrogène ou C₁₋₄alkyle ; de manière alternative, R^J et R^K sont pris ensemble avec l'atome d'azote auquel ils sont liés pour former hétérocyclyle de cinq à sept chaînons ; R¹ est choisi dans le groupe constitué par hydrogène, C₁₋₆alkyle, cycloalkyle, hétérocyclyle, aryl(C₁₋₆)alkyle et hétéroaryl(C₁₋₆)alkyle ;

où lorsque R¹ est phényl(C₁₋₆)alkyle, phényle est éventuellement condensé sur hétérocyclyle ou cycloalkyle ;
où lorsque R¹ est C₁₋₂alkyle, ledit C₁₋₂alkyle est éventuellement substitué par de un à deux substituants choisis indépendamment dans le groupe constitué par C₁₋₆alcoxy, aryle, cycloalkyle, hétérocyclyle, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluorométhyle et carboxy ;

et en outre, où lorsque R¹ est C₃₋₆alkyle, ledit C₃₋₆alkyle est éventuellement substitué par de un à trois substituants choisis indépendamment dans le groupe constitué par C₁₋₆alcoxy, aryle, cycloalkyle, hétérocyclyle, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluorométhyle et carboxy ;

où les groupements cycloalkyle et hétérocyclyle de C₁₋₂alkyle et C₃₋₆alkyle sont éventuellement substitués par de un à deux substituants choisis indépendamment dans le groupe constitué par C₁₋₆alkyle, hydroxy(C₁₋₆)alkyle, C₁₋₆alcoxy, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluorométhyle, carboxy, aryl(C₁₋₆)-alcoxycarbonyl, C₁₋₆alcoxycarbonyl, aminocarbonyl, C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl et aminosulfonyl ;


en outre, où les groupements cycloalkyle et hétérocyclyle de R¹ sont éventuellement substitués par de un à deux substituants choisis indépendamment dans le groupe constitué par C₁₋₆alkyle, hydroxy(C₁₋₆)alkyle, C₁₋₆alcoxy, hydroxy, cyano, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, trifluorométhyle, carboxy, aryl(C₁₋₆)-alcoxycarbonyl, C₁₋₆alcoxycarbonyl, aminocarbonyl, C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl et aminosulfonyl ;

en outre, où les portions aryle et hétéroaryle des substituants de R¹ aryl (C₁₋₆) alkyle et hétéro (C₁₋₆) alkyle sont éventuellement substituées par de un à trois substituants R¹¹ choisis indépendamment dans le groupe constitué par C₁₋₆alkyle, hydroxy(C₁₋₆)alkyle, C₁₋₆alcoxy, C₆₋₁₀aryl (C₁₋₆) alkyle, C₆₋₁₀aryl (C₁₋₆) alcoxy, C₆₋₁₀aryle, hétéroaryle éventuellement substitué par de un à deux substituants choisis indépendamment dans le groupe constitué par C₁₋₄alkyle, C₁₋₄alcoxy et carboxy ; cycloalkyle, hétérocyclyle, C₆₋₁₀aryloxy, hétéroaryloxy, cycloalkyloxy, hétérocyclyloxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, C₃₋₆cycloalkylaminocarbonyl, hydroxy-(C₁₋₆) alkylaminocarbonyl, C₆₋₁₀arylaminocarbonyl où C₆₋₁₀aryle est éventuellement substitué par carboxy ou C₁₋₄-alcoxycarbonyl, hétérocyclylcarbonyl, carboxy, C₁₋₆-alkylcarbonyloxy, C₁₋₆alcoxycarbonyl, C₁₋₆alkyl-carbonyl, C₁₋₆alkylcarbonylamino, aminocarbonyl, C₁₋₆-alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl, cyano, halogène, trifluorométhyle, trifluorométhoxy, et hydroxy ;

à condition que pas plus d'un substituant R¹¹ ne soit choisi dans le groupe constitué par C₆₋₁₀aryl (C₁₋₆) alkyle, C₆₋₁₀aryl (C₁₋₆) alcoxy, C₆₋₁₀aryle, hétéroaryle éventuellement substitué par de un à deux substituants choisis indépendamment dans le groupe constitué par C₁₋₄alkyle, C₁₋₄alcoxy et carboxy ; cycloalkyle, hétérocyclyle, C₆₋₁₀aryloxy, hétéroaryloxy, cycloalkyloxy, C₆₋₁₀arylaminocarbonyl, hétérocyclylcarbonyl et hétérocyclyloxy ;

R² est hydrogène, C₁₋₈alkyle, hydroxy(C₁₋₈)alkyle, C₆₋₁₀-aryl (C₁₋₆) alcoxy(C₁₋₆) alkyle ou C₆₋₁₀aryl (C₁₋₈) alkyle ;
où le groupement C₆₋₁₀aryle dans les substituants de R² contenant C₆₋₁₀aryle est éventuellement substitué par de un à deux substituants choisis indépendamment dans le groupe constitué par C₁₋₆alkyle, C₁₋₆alcoxy, hydroxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, aminocarbonyl, C₁₋₆alkylaminocarbonyl, (C₁₋₆alkyl)₂aminocarbonyl, cyano, fluoro, chloro, bromo, trifluorométhyle et trifluorométhoxy ; et où les substituants C₁₋₆alkyle et C₁₋₆alcoxy d'aryle sont éventuellement substitués par hydroxy, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino ou aryle ;


A est choisi dans le groupe constitué par aryle, un noyau a-1, a-2, a-3 et a-4, éventuellement substitué par R³ et R⁵ ;




où A-B est choisi dans le groupe constitué par N-C, C-N, N-N et C-C ; où D-E est choisi dans le groupe constitué par O-C, S-C et O-N ; et où F-G est choisi dans le groupe constitué par N-O et C-O ;

R³ est un à deux substituants choisis indépendamment dans le groupe constitué par C₁₋₆alkyle, aryle, aryl(C₁₋₆)alkyle, aryl (C₂₋₆) alcényle, aryl (C₂₋₆) alcynyle, hétéroaryle, hétéroaryl (C₁₋₆)alkyle, hétéroaryl(C₂₋₆)-alcényle, hétéroaryl (C₂₋₆) alcynyle, amino, C₁₋₆-alkylamino, (C₁₋₆alkyl)₂amino, arylamino, hétéro-arylamino, aryloxy, hétéroaryloxy, trifluorométhyle et halogène ;

où les groupements aryle et hétéroaryle, ainsi que les groupements aryle et hétéroaryle d'aryl(C₁₋₆)alkyle, aryl(C₂₋₆)alcényle, aryl(C₂₋₆)alcynyle, hétéroaryl(C₁₋₆)alkyle, hétéroaryl(C₂₋₆)alcényle, hétéroaryl(C₂₋₆)alcynyle, arylamino, hétéroarylamino, aryloxy et hétéroaryloxy, sont éventuellement substitués par de un à cinq substituants fluoro ou de un à trois substituants choisis indépendamment dans le groupe constitué par C₁₋₆alkyle, hydroxy(C₁₋₆)alkyle, C₁₋₆alcoxy, C₆₋₁₀aryl(C₁₋₈)alkyle, C₆₋₁₀aryl(C₁₋₈)alcoxy, C₆₋₁₀aryle, C₆₋₁₀aryloxy, hétéroaryl(C₁₋₆)alkyle, hétéroaryl(C₁₋₆)alcoxy, hétéroaryle, hétéroaryloxy, C₆₋₁₀arylamino, hétéro-arylamino, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, carboxy(C₁₋₆)alkylamino, carboxy, C₁₋₆alkylcarbonyle, C₁₋₆alcoxycarbonyle, C₁₋₆alkylcarbonylamino, aminocarbonyle, C₁₋₆alkylaminocarbonyle, (C₁₋₆alkyl)₂aminocarbonyle, carboxy(C₁₋₆)alkylaminocarbonyle, cyano, halogène, trifluorométhyle, trifluorométhoxy, hydroxy, C₁₋₆alkylsulfonyle et C₁₋₆alkylsulfonylamino ; à condition que pas plus d'un R³ ne soit choisi dans le groupe constitué par aryle, hétéroaryle, aryl(C₁₋₆)alkyle, aryl(C₂₋₆)alcényle, aryl(C₂₋₆)alcynyle, hétéroaryle, hétéroaryl(C₁₋₆)alkyle, hétéroaryl(C₂₋₆)alcényle, hétéroaryl(C₂₋₆)alcynyle, arylamino, hétéroarylamino, aryloxy et hétéroaryloxy ; et où C₁₋₆alkyle et C₁₋₆alkyle d'aryl(C₁₋₆)alkyle et d'hétéroaryl(C₁₋₆)alkyle sont éventuellement substitués par un substituant choisi dans le groupe constitué par hydroxy, carboxy, C₁₋₄alcoxycarbonyle, amino, C₁₋₆alkylamino, (C₁₋₆alkyl)₂amino, aminocarbonyle, (C₁₋₄)-alkylaminocarbonyle, di(C₁₋₄)alkylaminocarbonyle, aryle, hétéroaryle, arylamino, hétéroarylamino, aryloxy, hétéroaryloxy, aryl(C₁₋₄)alcoxy et hétéroaryl(C₁₋₄)-alcoxy ; R⁵ est un substituant sur un atome d'azote du cycle A choisi dans le groupe constitué par hydrogène et C₁₋₄-alkyle ; R^a et R^b sont choisis indépendamment dans le groupe constitué par hydrogène, C₁₋₆alkyle et C₁₋₆-alcoxycarbonyle ; de manière alternative, lorsque R^a et R^b sont chacun autres qu'hydrogène, R^a et R^b sont éventuellement pris conjointement avec l'atome d'azote auquel ils sont tous deux fixés pour former un cycle monocyclique de cinq à huit chaînons ; et les énantiomères, diastéréoisomères, racémates et sels pharmaceutiquement acceptables de ceux-ci ;




la réaction d'un composé de formule (X), où X^P est choisi parmi CN, -CO₂H, -C(O)-Cl ou -C(O)-OC₁₋₄alkyle et où Y^P est choisi parmi Br, Cl ou I, pour conduire au composé correspondant de formule (XII) ;




la réaction du composé de formule (XII) avec un composé de formule (XVIII) convenablement substitué, où Pg¹ est un groupement protecteur d'azote, en présence d'un catalyseur à base de palladium ; en présence d'une base organique ou inorganique ; dans un solvant organique ; à une température supérieure à environ la


température ambiante ; pour conduire au composé correspondant de formule (XIX) ;



la réaction du composé de formule (XIX) avec de l'hydrogène ou une source d'hydrogène ; en présence d'un catalyseur ; dans un solvant ; à une température supérieure à environ la température ambiante ; pour conduire au composé correspondant de formule (XX) ;



la réaction du composé de formule (XX) avec une base aqueuse ; dans un solvant organique ; pour conduire au composé correspondant de formule (I) ;



la réaction du composé de formule (I) pour conduire au composé correspondant de formule (II).

10. Procédé selon la revendication 9, dans lequel



est phényle ;

le groupement phényle est substitué par un groupement R^{41P} au niveau de la position 2 et un deuxième groupement

R^{41P} au niveau de la position 4 ;
 chaque R^{41P} est choisi indépendamment parmi C₁₋₂-alkyle, C₁₋₂alcoxy ou fluoro ;
 R^J et R^K sont choisis chacun indépendamment parmi hydrogène ou C₁₋₄alkyle ; de manière alternative, R^J et R^K
 sont pris ensemble avec l'atome d'azote auquel ils sont liés pour former hétérocyclyle de cinq à sept chaînons ; et
 Pg¹ est un groupement protecteur d'azote.

11. Procédé selon la revendication 9, dans lequel



est phényle ; le groupement phényle est substitué par un groupement R^{41P} au niveau de la position 2 et un deuxième
 groupement R^{41P} au niveau de la position 4 ; chaque R^{41P} est méthyle ; R^J et R^K sont chacun hydrogène ; et Pg¹
 est t-butoxycarbonyl.

12. Procédé selon l'une quelconque des revendications 1-3, 5-7 et 9-11, dans lequel X^P est choisi parmi CN, -C(O)-Cl
 ou -C(O)-OC₁₋₄alkyle.

13. Procédé selon l'une quelconque des revendications 1-3, 5-7 et 9-11, dans lequel Y^P est choisi parmi Cl ou I.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- US 11079647 B [0002] [0119]
- US 20050203143 A1 [0002] [0119]
- WO 03092688 A [0003]
- WO 0222612 A [0192]

Non-patent literature cited in the description

- Protective Groups in Organic Chemistry. Plenum Press, 1973 [0068]
- T.W. GREENE ; P.G.M. WUTS. Protective Groups in Organic Synthesis. John Wiley & Sons, 1991 [0068] [0069]
- S. CACCHI ; G. FABRIZI ; A. GOGGIAMANI. *Org. Lett.*, 2003, vol. 5 (23), 4269-4272 [0078] [0081] [0101] [0104] [0110]
- PARRIS, C.L. *Org. Syn. Coll.*, 1973, vol. 5, 73 [0084]
- LIN, S. *Synthesis*, April 1978, 330 [0084]
- MURAHASHI, S. ; TAKESHI NAOTA, T. ; EIICHIRO SAITO, E. *JACS*, 1986, vol. 108 (24), 7846 [0084]

Szabadalmi igénypontok

1. Eljárás az (I) képletű vegyület előállítására



ahol



jelentése C_{6-10} arilcsoport vagy egy heteroarilcsoport furil-, tienil-, pirrolil-, oxazolil-, flazolil-, imidazolil-, pirazolil-, piridinil-, pirimidinil-, pirazinil-, indolil-, izoindolil-, indolinil-, benzofuril-, benzotienil-, benzimidazolil-, benzisazolil-, benzoxazolil-, kinolizinil-, kinolinil-, izokinolinil- és kinazolinilcsoport közül kiválasztva;


R^{41P} mindegyik jelentése egymástól függetlenül C_{1-6} alkil-, C_{1-6} alkoxycsoport vagy fluoratom közül van kiválasztva;

R^J és R^K jelentése egymástól függetlenül hidrogénatom vagy C_{1-6} alkilcsoport közül van kiválasztva;


alternatív módon R^J és R^K azzal a nitrogénatommal együtt, amelyhez kapcsolódnak, öt-hét tagú heterocikluscsoportot képeznek;

PG^1 jelentése nitrogén-védőcsoport;


amelynek során




reagáltatunk egy (X) képletű vegyületet, ahol X^P jelentése CN-, $-CO_2H$ -, $-C(O)-Cl$ vagy $-C(O)-OC_{1-6}$ alkil közül van kiválasztva és ahol Y^P jelentése Br, Cl vagy I közül van kiválasztva, így kapjuk a megfelelő (XII) képletű vegyületet;



a (XII) képletű vegyületet egy megfelelően szubsztituált (XVIII) képletű vegyülettel reagáltatjuk; palládium katalizátor jelenlétében; egy szerves vagy szervetlen bázis jelenlétében; egy szerves oldószerben; körülbelül szobahőmérsékletnél magasabb hőmérsékleten; így kapjuk a megfelelő (XIX) képletű vegyületet;



a (XIX) képletű vegyületet hidrogénnel vagy egy hidrogénforrással reagáltatjuk; egy katalizátor jelenlétében; oldószerben; körülbelül szobahőmérsékletnél magasabb hőmérsékleten; így kapjuk a megfelelő (XX) képletű vegyületet;



a (XX) képletű vegyületet egy vizes bázissal reagáltatjuk; egy szerves oldószerben; így kapjuk a megfelelő (I) képletű vegyületet.

2. Az 1. igénypont szerinti eljárás, ahol



jelentése fenilcsoport;

a fenilcsoport 2-helyzetben egy R^{41P} csoporttal és 4-helyzetben egy második R^{41P} csoporttal van szubsztituálva;

R^{41P} mindegyik jelentése egymástól függetlenül C_{1-2} alkil-, C_{1-2} alkoxycsoport vagy fluoratom;

R^J és R^K mindegyikének jelentése egymástól függetlenül hidrogénatom vagy C_{1-4} alkilcsoport közül van kiválasztva;

alternatív módon R^J és R^K azzal a nitrogénatommal együtt, amelyhez kapcsolódnak, öt-hét tagú heterocikluscsoportot képeznek; és


Pg^1 jelentése nitrogén-védőcsoport.

3. Az 1. igénypont szerinti eljárás, ahol



jelentése fenilcsoport; a fenilcsoport 2-helyzetben szubsztituálva van egy R^{41P} csoporttal és 4-helyzetben egy második R^{41P} csoporttal; mindegyik R^{41P} jelentése metilcsoport; R^J és R^K mindegyikének jelentése hidrogénatom; és Pg^1 jelentése terc-butoxikarbonil-csoport.

4. Eljárás (I) képletű vegyület előállítására



ahol



jelentése C_{6-10} arilcsoport vagy egy heteroarilcsoport furil-, tienil-, pirrolil-, oxazolil-, tiazolil-, imidazolil-, pirazolil-, piridinil-, pirimidinil-, pirazinil-, indolil-, izoindolil-, indolinil-, benzofuril-, benzo tienil-, benzimidazolil-, benziazolil-, benzoxazolil-, kinolizinil-, kinolinil-, izokinolinil- és kinazolinilcsoport közül kiválasztva;


R^{41P} mindegyikének jelentése egymástól függetlenül C_{1-4} alkil-, C_{1-4} alkoxycsoport vagy fluoratom;

R^J és R^K mindegyikének jelentése egymástól függetlenül hidrogénatom vagy C_{1-4} alkilcsoport közül van kiválasztva;


alternatív módon R^J és R^K azzal a nitrogénatommal együtt, amelyhez kapcsolódnak, öt-hét tagú heterocikluscsoportot képeznek;

Pg^1 jelentése nitrogén-védőcsoport;

amelynek során




a (XIX) képletű vegyületet hidrogénnel vagy egy hidrogénforrással reagáltatjuk; katalizátor jelenlétében; oldószerben; körülbelül szobahőmérsékletnél magasabb hőmérsékleten; így kapjuk a megfelelő (XX) képletű vegyületet;




a (XX) képletű vegyületet egy vizes bázissal reagáltatjuk; szerves oldószerben; így kapjuk a megfelelő (I) képletű vegyületet.


5. Eljárás (Ia) képletű vegyület előállítására




amelynek során




reagáltatunk egy (Xa) képletű vegyületet, ahol X^p jelentése CN, $-\text{CO}_2\text{H}$, $-\text{C}(\text{O})-\text{Cl}$ vagy $-\text{C}(\text{O})-\text{OC}_{1-4}\text{alkil}$ közül van kiválasztva és ahol Y^p jelentése Br, Cl vagy I közül van kiválasztva, így kapjuk a megfelelő (XIa) képletű vegyületet;



a (XIIa) képletű vegyületet egy megfelelően szubsztituált (XVIIIa) képletű vegyülettel reagáltatjuk; palládium katalizátor jelenlétében; szerves vagy szervetlen bázis jelenlétében; szerves oldószerben; körülbelül szobahőmérsékletnél magasabb hőmérsékleten; így kapjuk a megfelelő (XIXa) képletű vegyületet;



a (XIXa) képletű vegyületet hidrogéngázzal reagáltatjuk, hidrogénezéshez szükséges nyomáson; megfelelő királis katalizátor jelenlétében; körülbelül szobahőmérsékletnél magasabb hőmérsékleten; egy szerves oldószerben; így kapjuk a megfelelő (XXa) képletű vegyületet;




a (XXa) képletű vegyületet egy vízes bázissal reagáltatjuk; egy szerves oldószerben; így kapjuk a megfelelő (Ia) képletű vegyületet.

6. Az 5. igénypont szerinti eljárás, ahol a (XIIa) képletű vegyületet (XVIIIa) képletű vegyülettel reagáltatjuk $\text{Pd}_2(\text{dba})_3$ és $\text{P}(o\text{-toluol})_3$ jelenlétében.

7. Az 5. igénypont szerinti eljárás, ahol a királis katalizátor $[\text{Rh}(\text{cod})(\text{R,R-DIPAMP})]^+\text{BF}_4^-$.

8. Eljárás (Ia) képletű vegyület előállítására



amelynek során

ahol amikor R^1 jelentése fenil(C_{1-6})alkilcsoport, a fenilcsoport adott esetben heterociklil- vagy cikloalkilcsoporttal van kondenzálva;

ahol amikor R^1 jelentése C_{1-2} alkilcsoport, az említett C_{1-2} alkilcsoport adott esetben szubsztituálva van egy-kettő szubsztituenssel, amelyek egymástól függetlenül C_{1-6} alkoxi-, aril-, cikloalkil-, heterociklil-, hidroxil-, ciano-, amino-, C_{1-6} alkilamino-, $(C_{1-6}$ alkil)₂amino-, trifluormetil- és karboxilcsoport közül vannak kiválasztva;

továbbá, ahol amikor R^1 jelentése C_{1-6} alkilcsoport, az említett C_{1-6} alkilcsoport adott esetben szubsztituálva van egy-három szubsztituenssel, amelyek egymástól függetlenül C_{1-6} alkoxi-, aril-, cikloalkil-, heterociklil-, hidroxil-, ciano-, amino-, C_{1-6} alkilamino-, $(C_{1-6}$ alkil)₂amino-, trifluormetil- és karboxilcsoport közül vannak kiválasztva;

ahol a C_{1-6} alkil- és C_{1-6} alkilcsoport cikloalkil- és heterociklilcsoportja adott esetben szubsztituálva van egy-kettő szubsztituenssel, amelyek egymástól függetlenül C_{1-6} alkil-, hidroxil(C_{1-6})alkil-, C_{1-6} alkoxi-, hidroxil-, ciano-, amino-, C_{1-6} alkilamino-, $(C_{1-6}$ alkil)₂amino-, trifluormetil-, karboxil-, aril(C_{1-6})alkoxikarbonil-, C_{1-6} alkoxikarbonil-, aminokarbonil-, C_{1-6} alkilaminokarbonil-, $(C_{1-6}$ alkil)₂aminokarbonil- és aminoszulfonil-csoport közül vannak kiválasztva;

továbbá, ahol R^1 jelentésében a cikloalkil- és heterociklilcsoport adott esetben szubsztituálva van egy-kettő szubsztituenssel, amelyek egymástól függetlenül C_{1-6} alkil-, hidroxil(C_{1-6})alkil-, C_{1-6} alkoxi-, hidroxil-, ciano-, amino-, C_{1-6} alkilamino-, $(C_{1-6}$ alkil)₂amino-, trifluormetil-, karboxil-, aril(C_{1-6})alkoxikarbonil-, C_{1-6} alkoxikarbonil-, aminokarbonil-, C_{1-6} alkilaminokarbonil-, $(C_{1-6}$ alkil)₂aminokarbonil- és aminoszulfonil-csoport közül vannak kiválasztva;


továbbá, ahol az R^1 szubsztituensek jelentésében az aril(C_{1-6})alkil- és heteroaril(C_{1-6})alkilcsoport aril- és heteroaril-része adott esetben szubsztituálva van egy-három R^{11} szubsztituenssel, amelyek egymástól függetlenül a következő csoportból vannak kiválasztva: C_{1-6} alkilcsoport; hidroxil(C_{1-6})alkil-csoport; C_{1-6} alkoxycsoport; C_{6-10} aril(C_{1-6})alkil-csoport; C_{6-10} aril(C_{1-6})alkoxi-csoport; C_{6-10} arilcsoport; heteroarilcsoport, amely adott esetben C_{1-6} alkil-, C_{1-6} alkoxi- és karboxilcsoport közül egymástól függetlenül kiválasztott egy-kettő szubsztituenssel van szubsztituálva; cikloalkilcsoport; heterociklilcsoport; C_{6-10} ariloxycsoport; heteroariloxi-csoport; cikloalkiloxi-csoport; heterocikliloxi-csoport; aminocsoport; C_{1-6} alkilamino-csoport; $(C_{1-6}$ alkil)₂amino-csoport; C_{1-6} cikloalkilaminokarbonil-csoport; hidroxil(C_{1-6})alkilaminokarbonil-csoport; C_{6-10} arilaminokarbonil-csoport, ahol a C_{6-10} arilcsoport adott esetben karboxil- vagy C_{1-6} alkoxikarbonil-csoporttal van szubsztituálva; heterociklilkarbonil-csoport; karboxilcsoport; C_{1-6} alkilkarboniloxi-csoport; C_{1-6} alkoxikarbonil-csoport; C_{1-6} alkilkarbonil-csoport; C_{1-6} alkilkarbonilamino-csoport; aminokarbonil-csoport; C_{1-6} alkilaminokarbonil-csoport; $(C_{1-6}$ alkil)₂aminokarbonil-csoport; cianocsoport; halogénatom; trifluormetilcsoport; trifluor-metoxycsoport; és hidroxilcsoport;

azzal a megkötéssel, hogy egy-nél nem több R^{11} szubsztituens a következő csoportból van kiválasztva: C_{6-10} aril(C_{1-6})alkil-csoport; C_{6-10} aril(C_{1-6})alkoxi-csoport; C_{6-10} arilcsoport; heteroarilcsoport, amely adott esetben szubsztituálva van egy-kettő szubsztituenssel, amelyek egymástól függetlenül C_{1-6} alkil-, C_{1-6} alkoxi- és karboxilcsoport közül vannak kiválasztva; cikloalkilcsoport; heterociklilcsoport; C_{6-10} ariloxycsoport; heteroariloxi-csoport; cikloalkiloxi-csoport; C_{6-10} arilaminokarbonil-, heterociklilkarbonil-csoport; és heterocikliloxi-csoport;

R^2 jelentése hidrogénatom, C_{1-6} alkil-, hidroxil(C_{1-6})alkil-, C_{6-10} aril(C_{1-6})alkoxi(C_{1-6})alkil-, vagy C_{6-10} aril(C_{1-6})alkilcsoport;

ahol R^2 jelentésében az C_{6-10} aril-tartalmú szubsztituensnek C_{6-10} aril-csoportja adott esetben szubsztituálva van egy-kettő szubsztituenssel, amelyek egymástól függetlenül C_{1-6} alkil-, C_{1-6} alkoxi-, hidroxil-, amino-, C_{1-6} alkilamino-, $(C_{1-6}$ alkil) $_2$ amino-, aminokarbonil-, C_{1-6} alkilaminokarbonil-, $(C_{1-6}$ alkil) $_2$ aminokarbonil-, cianocsoport, fluor-, klór-, brómatom, trifluometil- és trifluormetoxicsoport közül vannak kiválasztva; és ahol az arilcsoport C_{1-6} alkil- és C_{1-6} alkoxi-szubsztituensei adott esetben szubsztituálva vannak hidroxil-, amino-, C_{1-6} alkilamino-, $(C_{1-6}$ alkil) $_2$ amino- vagy arilcsoporttal;

A jelentése arilcsoport és adott esetben R^3 és R^5 csoporttal szubsztituált a-1, a-2, a-3 és a-4 gyűrűrendszerek csoportjából van kiválasztva;



ahol A-B jelentése N-C, C-N, N-N és C-C csoportjából van kiválasztva; ahol D-E jelentése O-C, S-C, és ON csoportjából van kiválasztva; és ahol F-G jelentése N-O és C-O csoportjából van kiválasztva;

R^2 jelentése egy-kettő szubsztituens, amelyek egymástól függetlenül C_{1-6} alkil-, aril-, aril(C_{1-6}) alkil-, aril(C_{2-6}) alkenil-, aril(C_{2-6}) alkínil-, heteroaril-, heteroaril(C_{1-6}) alkil-, heteroaril(C_{2-6}) alkenil-, heteroaril(C_{2-6}) alkínil-, amino-, C_{1-6} alkilamino-, $(C_{1-6}$ alkil) $_2$ amino-, arilamino-, heteroarilamino-, ariloxi-, heteroariloxi-, trifluometilcsoport és halogénatom közül vannak kiválasztva;


ahol az aril- és a heteroarilcsoport, valamint az aril(C_{1-6}) alkil-, aril(C_{2-6}) alkenil-, aril(C_{2-6}) alkínil-, heteroaril(C_{1-6}) alkil-, heteroaril(C_{2-6}) alkenil-, heteroaril(C_{2-6}) alkínil-, arilamino-, heteroarilamino-, ariloxi- és heteroariloxi-csoport aril- és heteroarilcsoportja adott esetben szubsztituálva van egy-öt fluor-szubsztituenssel vagy egy-három C_{1-6} alkil-, hidroxil(C_{1-6}) alkil-, C_{1-6} alkoxi-, C_{6-10} aril(C_{1-6}) alkil-, C_{6-10} aril(C_{1-6}) alkoxi-, C_{6-10} aril-, C_{6-10} ariloxi-, heteroaril(C_{1-6}) alkil-, heteroaril(C_{1-6}) alkoxi-, heteroaril-, heteroariloxi-, C_{6-10} arilamino-, heteroarilamino-, amino-, C_{1-6} alkilamino-, $(C_{1-6}$ alkil) $_2$ amino-, karboxi(C_{1-6}) alkilamino-, karboxil-, C_{1-6} alkilkarbonil-, C_{1-6} alkoxikarbonil-, C_{1-6} alkilkarbonilamino-, aminokarbonil-, C_{1-6} alkilaminokarbonil-, $(C_{1-6}$ alkil) $_2$ aminokarbonil-, karboxi(C_{1-6}) alkilaminokarbonil-, cianocsoport, halogénatom, trifluometil-, trifluormetoxi-, hidroxil-, C_{1-6} alkilszulfonil- és C_{1-6} alkilszulfonilamino-csoport közül egymástól függetlenül kiválasztott szubsztituenssel; azzal a megkötéssel, hogy nem több mint egy R^3 csoport aril-, heteroaril-, aril(C_{1-6}) alkil-, aril(C_{2-6}) alkenil-, aril(C_{2-6}) alkínil-, heteroaril-, heteroaril(C_{1-6}) alkil-, heteroaril(C_{2-6}) alkenil-, heteroaril(C_{2-6}) alkínil-, arilamino-, heteroarilamino-, ariloxi- és heteroariloxi-csoport közül van kiválasztva;

és ahol a C_{1-6} alkilcsoport, valamint az aril(C_{1-6}) alkil- és heteroaril(C_{1-6}) alkilcsoport C_{1-6} alkilcsoportja adott esetben szubsztituálva van egy, hidroxil-, karboxil-, C_{1-6} alkoxikarbonil-, amino-, C_{1-6} alkilamino-, $(C_{1-6}$ alkil) $_2$ amino-, aminokarbonil-, $(C_{1-6}$) alkilaminokarbonil-, di(C_{1-6}) alkilaminokarbonil-, aril-, heteroaril-, arilamino-, heteroarilamino-, ariloxi-, heteroariloxi-, aril(C_{1-6}) alkoxi- és heteroaril(C_{1-6}) alkoxi-csoport közül kiválasztott szubsztituenssel;


R^5 jelentése egy, az A gyűrű nitrogénatomján lévő szubsztituens, amely hidrogénatom és C_{1-6} alkilcsoport közül van kiválasztva;

R^a és R^b jelentése egymástól függetlenül hidrogénatom, C_{1-4} alkil- és C_{1-4} alkoxikarbonil-csoport közül van kiválasztva; alternatív módon amikor R^a és R^b mindegyikének jelentése hidrogénatomtól eltérő, R^a és R^b adott esetben azzal a nitrogénatommal egyült, amelyhez kapcsolódnak, öt-nyolc tagú monociklusos gyűrűt képeznek;


és ennek gyógyszerészetileg elfogadható enantiomerei, diasztereomerei, racémái és sói, amelynek során




reagáltatunk egy (X) képletű vegyületet, ahol X^P jelentése CN , $-CO_2H$, $-C(O)-Cl$ vagy $-C(O)-OC_{1-4}$ alkil közül van kiválasztva és ahol Y^P jelentése Br , Cl vagy I közül van kiválasztva, így kapjuk a megfelelő (XII) képletű vegyületet;




a (XII) képletű vegyületet reagáltatjuk egy megfelelően szubsztituált (XVIII) képletű vegyülettel, ahol Pg^1 jelentése nitrogén-védőcsoport; palládium katalizátor jelenlétében; egy szerves vagy szervetlen bázis jelenlétében; egy szerves oldószerben; körülbelül szobahőmérsékletnél magasabb hőmérsékleten; így kapjuk a megfelelő (XIX) képletű vegyületet;



a (XIX) képletű vegyületet hidrogénnel vagy egy hidrogénforrással reagáltatjuk; katalizátor jelenlétében; oldószerben; körülbelül szobahőmérsékletnél magasabb hőmérsékleten; így kapjuk a megfelelő (XX) képletű vegyületet;



a (XX) képletű vegyületet egy vizes bázissal reagáltatjuk; egy szerves oldószerben; így kapjuk a megfelelő (I) képletű vegyületet;



az (I) képletű vegyületet reagáltatva kapjuk a megfelelő (II) képletű vegyületet.

10. A 9. igénypont szerinti eljárás, ahol



jelentése fenilcsoport;

amely fenilcsoport 2-helyzetben szubsztituálva van egy R^{41P} csoporttal és 4-helyzetben egy második R^{41P} csoporttal;

R^{41P} mindegyik jelentése egymástól függetlenül C_{1-2} alkil-, C_{1-2} alkoxycsoport vagy fluoratom közül van kiválasztva;

R^j és R^k mindegyikének jelentése egymástól függetlenül hidrogénatom vagy C_{1-4} alkilcsoport közül van kiválasztva;

alternatív módon R^j és R^k azzal a nitrogénatommal együtt, amelyhez kapcsolódnak, öt-hét tagú heterocikluscsoportot képeznek; és

Pg^1 jelentése nitrogén-védőcsoport.

11. A 9. igénypont szerinti eljárás, ahol



jelentése fenilcsoport; amely fenilcsoport 2-helyzetben szubsztituálva van egy $R^{4'p}$ csoporttal és 4-helyzetben egy második $R^{4'p}$ csoporttal; mindegyik $R^{4'p}$ jelentése metilcsoport; R^1 és R^2 mindegyikének jelentése hidrogénatom; és Pg^1 jelentése terc-butoxikarbonil-csoport.

12. Az 1-3., 5-7. és 9-11. igénypontok bármelyike szerinti eljárás, ahol X^p jelentése CN, $-C(O)-Cl$ vagy $-C(O)-OC_{1-4}alkil$ közül van kiválasztva.

13. Az 1-3., 5-7. és 9-11. igénypontok bármelyike szerinti eljárás, ahol Y^p jelentése Cl vagy I közül van kiválasztva.