



(11) (21) (C) **2,058,822**
(22) 1992/01/06
(43) 1992/07/19
(45) 2000/06/27

(72) Ternansky, Robert J., US



(73) ELI LILLY AND COMPANY, US

(51) Int.Cl.⁵ C07D 519/00, C07D 501/59, A61K 31/545, C07F 7/10

(30) 1991/01/18 (07/642,959) US

(54) **AGENTS ANTIBACTERIENS**


(54) **ANTIBACTERIAL AGENTS**



(57) The present invention provides cephalosporins of the following general formula: (see formula I) wherein R is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₆ cycloalkyl, or C₁-C₆ haloalkyl; A and A' are taken together to form a group of the formula (see formula II) wherein X is hydrogen, halo, C₁-C₆ alkyl, -C₁-C₆ alkoxy, C₁-C₆ alkoxy-carbonyl, amino, nitro, or carboxy; and Y is nitrogen; and pharmaceutically acceptable salts thereof. These compounds are useful as antibacterial agents. Also provided by the invention are pharmaceutical formulations incorporating a compound of the above formula as the active ingredient, and the use of such compounds as antibiotics.


ABSTRACT OF THE DISCLOSURE

The present invention provides cephalosporins of the following general formula:



5 wherein R is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₆ cycloalkyl, or C₁-C₆ haloalkyl;

A and A' are taken together to form a group of the formula




10 wherein X is hydrogen, halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxy carbonyl, amino, nitro, or carboxy; and Y is nitrogen; and pharmaceutically acceptable salts thereof. These compounds are useful as antibacterial agents. Also provided by the invention are pharmaceutical formulations incorporating a compound of the above formula as the active ingredient, and the use
15 of such compounds as antibiotics.

ANTIBACTERIAL AGENTS

This invention relates to cephalosporin anti-
biotics, to pharmaceutical formulations comprising the
5 antibiotics, and to a method for the treatment of infec-
tious diseases in man and other animals.

Cephalosporin antibiotics have the bicyclic
ring system represented by the following formula wherein
the numbering system is that commonly employed in the
10 arbitrary cepham nomenclature system.



15

In the field of antibacterial therapy, the
need for new chemotherapeutic agents is one that will
never extinguish. Mutant strains resistant to existing
20 antibacterial agents are encountered frequently. In
particular, many strains of Staph. aureus and Staph. epi
(so-called methicillin resistant Staph. (MRS)) are becom-
ing increasingly resistant to available antibacterial
agents. (see, for example, Phillips, I., and Cookson,
25 B., J. Appl. Bacteriology 67(6)1989). To meet this need,
considerable research effort continues to focus on such
new agents. The present invention provides antibacterial


agents useful against a wide variety of gram-positive and gram-negative bacteria. The compounds of the present invention are especially useful against these methicillin-resistant Staph. organisms.

5


The present invention provides various 3-thiazolothio cephalosporins useful as antibacterial agents. In particular, the present invention provides 7 β -(2-aminothiazol-4-yl)oximino-(or alkoximino)acetylamino-3-
10 optionally-substituted-thiazolothio-3-cephem-4-carboxylic acids useful as antibacterial agents. The invention also provides pharmaceutical formulations and a therapeutic method useful in the treatment of antibacterial infections in man and other animals.

15

The present invention provides compounds of Formula (1):



wherein R is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₆ cycloalkyl, or C₁-C₆ haloalkyl; A and A' are independently hydrogen, C₁-C₆ alkyl, nitro, amino, C₁-C₆ alkoxy, a 5 or 6 membered heterocycle containing
 5 nitrogen or sulfur, or phenyl; or A and A' taken together form a group of the formula



wherein X is hydrogen, halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxycarbonyl, amino, nitro, or carboxy; and Y is
 15 nitrogen or carbon; or a pharmaceutically acceptable salt thereof.

The term "pharmaceutically-acceptable salt" encompasses those salts that form with the carboxylate anions and includes salts formed with the organic and inorganic cations such as counterions chosen from the
 20 alkali and alkaline earth metals, (such as lithium, sodium, potassium, barium and calcium); ammonium; and the organic cations (such as dibenzylammonium, benzylammonium, 2-hydroxyethylammonium, bis(2-hydroxyethyl)ammonium, phenylethylbenzylammonium, dibenzylethylene-
 25 diammonium, and like cations). Other cations encompassed by the above term include the protonated form of procaine, quinine and N-methylglucosamine, and the protonated forms of basic amino acids such as glycine, ornithine, histidine, phenylglycine, lysine and arginine. Further-
 30 more, any zwitterionic form of the compounds represented

by formula (1) formed by a carboxylic acid and an amino group is referred to by this term. A preferred cation for the carboxylate anion is the sodium cation. Furthermore, the term includes salts that form by standard acid-base reactions with basic groups (such as amino groups) and organic or inorganic acids. Such acids include hydrochloric, sulfuric, phosphoric, acetic, succinic, citric, lactic, maleic, fumaric, palmitic, cholic, pamoic, mucic, D-glutamic, d-camphoric, glutaric, phthalic, tartaric, lauric, stearic, salicylic, methanesulfonic, benzenesulfonic, sorbic, picric, benzoic, cinnamic, and like acids.

In the above Formula (1), the term " C_1-C_6 alkyl" denotes such radicals as methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, amyl, tert-amyl, hexyl and the like. The preferred " C_1-C_6 alkyl" group is methyl.

The term " C_2-C_6 alkenyl" is a straight chain or branched lower alkenyl and is exemplified by vinyl, allyl, 1-propenyl, isopropenyl, 1-butenyl, 2-butenyl, 3-butenyl, methallyl, or 1,1-dimethylallyl.

The term " C_2-C_6 alkynyl" is a straight chain or branched lower alkynyl group and is exemplified by ethynyl, 1-propynyl, or propargyl.

The term " C_3-C_{10} cycloalkyl" is exemplified by cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, norbornyl or adamantyl.

The term "C₁-C₆ haloalkyl" denotes the above C₁-C₆ alkyl groups that are substituted by one halogen, wherein "halo" or "halogen" denotes the chloro, bromo, iodo, and fluoro groups. Fluoro C₁-C₆ alkyl is preferred.

5 Fluoroethyl is a further preferred "C₁-C₆ haloalkyl" group.

The term "C₁-C₆ alkoxy" refers to such groups as methoxy, ethoxy, 3-propoxy, butyloxy, and the like.

The term "halo" includes fluoro, bromo,


10 chloro and iodo.

The term "C₁-C₆ alkoxy-carbonyl" refers to such groups as methoxycarbonyl, ethoxycarbonyl, 3-propoxycarbonyl, 3-ethoxycarbonyl, 4-t-butyloxycarbonyl, 3-methoxycarbonyl, 6-methoxycarbonyl, and the like.

15 The term "5 or 6 membered heterocycle containing nitrogen or sulfur" includes pyridine and thiophene, and may include more than a nitrogen or sulfur, and combinations thereof. Other examples include those described in Fletcher, Dermer, & Otis, Nomenclature of


20 Organic Compounds, pp. 49-64 (1974).

Compounds of Formula (1) may be prepared according to Scheme 1:



The starting material (A), (wherein R is methyl), 2-(trityl)amino- α -(methoxy-imino)-4-thiazole-acetic acid may be prepared from the corresponding free amine (available from Aldrich Chemical Co., Inc., 940 West Saint Paul Avenue, Milwaukee, Wisconsin 53233) utilizing methodology well-known in the β -lactam art. Starting material (B), or benzhydryl 7-amino-3-trifluoromethanesulfonyl-3-cephem-4-carboxylate may be prepared using known methodology, such as from the corresponding 3-enol-3-cephem and trifluoromethanesulfonyl-acetic anhydride. (Syn. Commun. 20(14), 2185-2189 (1990)).

In Scheme 1, the acid chloride of (A) can be prepared by known methodology, for example, by reaction with phosphoryl chloride, and reacted with the free amine (B) to form the 7-acyl-3-triflate (C). The thiazolothio group can then be introduced by reacting the triflate (C) with a compound of formula




in the presence of a base such as NaH. The final product (1) can then be prepared by removal of amino and carboxy protecting groups. In the above scheme, $\text{CF}_3\text{CO}_2\text{H}/(\text{CH}_3\text{CH}_2)_3\text{SiH}$ is utilized to remove the trityl and benzhydryl groups. One of ordinary skill in the

25

art of β -lactam chemistry will appreciate that other protecting groups would be efficacious. Further, one may also introduce the thiazolothio function into the 3-position of the cephem nucleus (B) prior to the insertion of the 7-acyl functions to provide useful intermediates set forth in formula (2) below.


Alternatively, the compounds may be prepared according to Scheme (2) below:

Scheme 2



In Scheme (2), the acetic acid (A), dissolved in DMF, is treated with N-methylmorpholine and oxalyl chloride. A mixture of 7 β -amino-3-chloro-cephem dissolved in DMF and treated with bis(dimethylsilyl)urea (BSU) and pyridine is combined with the acetic acid, to form (C). Compound (C) is then treated with diphenyldiazomethane, and the thiazolothio group is introduced in the presence of a base such as NaH, to form compound (D). The removal of the benzhydryl and trityl groups may be removed as in Scheme (1).


Compounds of the formula



15

where A and A' are taken together to form a group of the formula


20



and Y is nitrogen may be prepared according to the Scheme (3):


25

30

Scheme (3)


In Scheme (3), 3-aminopyridine is acylated with di-t-butyldicarbonate to introduce the t-butoxycarbonyl (t-BOC) protecting group. (It will be appreciated that two other pyridinothiazolothio mercaptans may be prepared by known methodology using other amino pyridine isomers.) The t-BOC protected 3-aminopyridine is then treated with n-butyllithium in tetrahydrofuran followed by elemental sulfur (S_8), followed by treatment with saturated ammonium chloride. The resulting 3-t-butoxycarbonylamino-4-thia-pyridine is treated with a mixture of acetic acid and HCl to provide 3-amino-4-mercaptopyridine hydrochloride. The desired 5-pyridinothiazolo thiomercaptan can then be prepared by treating this compound with carbon disulfide under basic conditions.

When A and A' are taken together to form a group of the formula



20


the desired thiol of the formula



25

may be made as shown in scheme (4) below:


Scheme (4)



20 In the above scheme, 2-chloro-3-nitropyridine is treated with potassium isothiocyanate to provide 2-isocyanato-3-nitropyridine, which is in turn hydrolyzed to provide 2-mercapto-3-nitropyridine. The 3-nitro intermediate is then reduced by treatment with

25 SnCl_2/HCl to provide 2-mercapto-3-amino pyridine. The desired pyridinothiazolothio mercaptan is then prepared by base catalyzed condensation with CS_2 ($\text{KOH}/\text{CH}_3\text{OH}/\text{CS}_2/\text{H}_2\text{O}$).

Examples of compounds falling within the scope of formula 1 are set forth in the table below:



10

TABLE 1

	<u>R</u>	<u>A</u>	<u>A'</u> (independently)
	methyl	H	H
15	ethyl	H	H
	propyl	H	H
	butyl	H	H
	pentyl	H	H
	hexyl	H	H
20	isopropyl	H	H
	isobutyl	H	H
	t-butyl	H	H
	isopentyl	H	H
	isohexyl	H	H
25	fluoromethyl	H	H
	1-fluoroethyl-2-yl	H	H
	1-fluoroprop-3-yl	H	H
	1-fluoro-but-4-yl	H	H
	1-fluoro-hex-5-yl	H	H
30	chloromethyl	H	H

TABLE 1 (CONTINUED)

	1-chloreth-2-yl	H	H
	1-chloroprop-3-yl	H	H
5	1-chlorobut-4-yl	H	H
	bromomethyl	H	H
	1-bromoeth-2-yl	H	H
	1-bromoprop-3-yl	H	H
	1-bromobut-4-yl	H	H
10	vinyl	H	H
	1-propene-2-yl	H	H
	1-butene-4-yl	H	H
	1-pentene-5-yl	H	H
	1-hexene-6-yl	H	H
15	cyclopropyl	H	H
	cyclobutyl	H	H
	cyclopentyl	H	H
	cyclohexyl	H	H
20	<u> R</u>	<u> A</u>	<u> A'</u> (independently)
	methyl	NO ₂	H
	ethyl	NO ₂	H
	propyl	NO ₂	H
25	butyl	NO ₂	H
	pentyl	NO ₂	H
	hexyl	NO ₂	H
	isopropyl	NO ₂	H
	isobutyl	NO ₂	H
30	t-butyl	NO ₂	H
	isopentyl	NO ₂	H

TABLE 1 (CONTINUED)

	isohexyl	NO ₂	H
	fluoromethyl	NO ₂	H
5	1-fluoroethyl-2-yl	NO ₂	H
	1-fluoroprop-3-yl	NO ₂	H
	1-fluoro-but-4-yl	NO ₂	H
	1-fluoro-hex-5-yl	NO ₂	H
	chloromethyl	NO ₂	H
10	1-chloroeth-2-yl	NO ₂	H
	1-chloroprop-3-yl	NO ₂	H
	1-chlorobut-4-yl	NO ₂	H
	bromomethyl	NO ₂	H
	1-bromoeth-2-yl	NO ₂	H
15	1-bromoprop-3-yl	NO ₂	H
	1-bromobut-4-yl	NO ₂	H
	vinyl	NO ₂	H
	1-propene-2-yl	NO ₂	H
	1-butene-4-yl	NO ₂	H
20	1-pentene-5-yl	NO ₂	H
	1-hexene-6-yl	NO ₂	H
	cyclopropyl	NO ₂	H
	cyclobutyl	NO ₂	H
	cyclopentyl	NO ₂	H
25	cyclohexyl	NO ₂	H

TABLE 1 (CONTINUED)

	<u>R</u>	<u>A</u>	<u>A'</u> (independently)
	methyl	NH ₂	H
5	ethyl	NH ₂	H
	propyl	NH ₂	H
	butyl	NH ₂	H
	pentyl	NH ₂	H
	hexyl	NH ₂	H
10	isopropyl	NH ₂	H
	isobutyl	NH ₂	H
	t-butyl	NH ₂	H
	isopentyl	NH ₂	H
	isohexyl	NH ₂	H
15	fluoromethyl	NH ₂	H
	1-fluoroethyl-2-yl	NH ₂	H
	1-fluoroprop-3-yl	NH ₂	H
	1-fluoro-but-4-yl	NH ₂	H
	1-fluoro-hex-5-yl	NH ₂	H
20	chloromethyl	NH ₂	H
	1-chloroeth-2-yl	NH ₂	H
	1-chloroprop-3-yl	NH ₂	H
	1-chlorobut-4-yl	NH ₂	H
	bromomethyl	NH ₂	H
25	1-bromoeth-2-yl	NH ₂	H
	1-bromoprop-3-yl	NH ₂	H
	1-bromobut-4-yl	NH ₂	H
	vinyl	NH ₂	H
	1-propene-2-yl	NH ₂	H
30	1-butene-4-yl	NH ₂	H

TABLE 1 (CONTINUED)

	1-pentene-5-yl	NH ₂	H
	1-hexene-6-yl	NH ₂	H
5	cyclopropyl	NH ₂	H
	cyclobutyl	NH ₂	H
	cyclopentyl	NH ₂	H
	cyclohexyl	NH ₂	H
10	<u>R</u>	<u>A</u>	<u>A'</u> (independently)
	methyl	CH ₃	H
	ethyl	CH ₃	H
	propyl	CH ₃	H
15	butyl	CH ₃	H
	pentyl	CH ₃	H
	hexyl	CH ₃	H
	isopropyl	CH ₃	H
	isobutyl	CH ₃	H
20	t-butyl	CH ₃	H
	isopentyl	CH ₃	H
	isohexyl	CH ₃	H
	fluoromethyl	CH ₃	H
	1-fluoroethyl-2-yl	CH ₃	H
25	1-fluoroprop-3-yl	CH ₃	H
	1-fluoro-but-4-yl	CH ₃	H
	1-fluoro-hex-5-yl	CH ₃	H
	chloromethyl	CH ₃	H
	1-chloroeth-2-yl	CH ₃	H

TABLE 1 (CONTINUED)

	1-chloroprop-3-yl	CH ₃	H
	1-chlorobut-4-yl	CH ₃	H
5	bromomethyl	CH ₃	H
	1-bromoeth-2-yl	CH ₃	H
	1-bromoprop-3-yl	CH ₃	H
	1-bromobut-4-yl	CH ₃	H
	vinyl	CH ₃	H
10	1-propene-2-yl	CH ₃	H
	1-butene-4-yl	CH ₃	H
	1-pentene-5-yl	CH ₃	H
	1-hexene-6-yl	CH ₃	H
	cyclopropyl	CH ₃	H
15	cyclobutyl	CH ₃	H
	cyclopentyl	CH ₃	H
	cyclohexyl	CH ₃	H
	<u> R</u>		<u>A and A'</u> together forming
20	methyl		
	ethyl		
	propyl		
	butyl		
25	pentyl		
	hexyl		
	isopropyl		
	isobutyl		
	t-butyl		




TABLE 1 (CONTINUED)

	isopentyl
	isohexyl
5	fluoromethyl
	1-fluoroethyl-2-yl
	1-fluoroprop-3-yl
	1-fluoro-but-4-yl
	1-fluoro-hex-5-yl
10	chloromethyl
	1-chloroeth-2-yl
	1-chloroprop-3-yl
	1-chlorobut-4-yl
	bromomethyl
15	1-bromoeth-2-yl
	1-bromoprop-3-yl
	1-bromobut-4-yl
	vinyl
	1-propene-2-yl
20	1-butene-4-yl
	1-pentene-5-yl
	1-hexene-6-yl
	cyclopropyl
	cyclobutyl
25	cyclopentyl
	cyclohexyl

TABLE 1 (CONTINUED)


	<u>R</u>	<u>A and A'</u> together forming
5	methyl ethyl propyl butyl pentyl hexyl	
10	isopropyl isobutyl t-butyl isopentyl isohexyl	
15	fluoromethyl 1-fluoroethyl-2-yl 1-fluoroprop-3-yl 1-fluoro-but-4-yl 1-fluoro-hex-5-yl	
20	chloromethyl 1-chloroprop-3-yl 1-chlorobut-4-yl bromomethyl 1-bromoeth-2-yl	
25	1-bromoprop-3-yl 1-bromobut-4-yl vinyl 1-propene-2-yl 1-butene-4-yl	

TABLE 1 (CONTINUED)


	1-pentene-5-yl	
	1-hexene-6-yl	
	cyclopropyl	
5	cyclobutyl	
	cyclopentyl	
	cyclohexyl	
	<u>R</u>	<u>A and A' together forming</u>
10	methyl	
	ethyl	
	propyl	
	butyl	
15	pentyl	
	hexyl	
	isopropyl	
	isobutyl	
	t-butyl	
20	isopentyl	
	isohexyl	
	fluoromethyl	
	1-fluoroethyl-2-yl	
	1-fluoroprop-3-yl	
25	1-fluoro-but-4-yl	
	1-fluoro-hex-5-yl	
	chloromethyl	
	1-chloroeth-2-yl	
	1-chloroprop-3-yl	
30	1-chlorobut-4-yl	

Table 1, (CONTINUED)


	bromomethyl	
	1-bromoeth-2-yl	
	1-bromoprop-3-yl	
5	1-bromobut-4-yl	
	vinyl	
	1-propene-2-yl	
	1-butene-4-yl	
	1-pentene-5-yl	
10	1-hexene-6-yl	
	cyclopropyl	
	cyclobutyl	
	cyclopentyl	
	cyclohexyl	
15		
	<u>R</u>	<u>A and A'</u> taken together forming
	methyl	
	ethyl	
20	propyl	
	butyl	
	pentyl	
	hexyl	
	isopropyl	
25	isobutyl	
	t-butyl	
	isopentyl	
	isoheptyl	
	fluoromethyl	
30	1-fluoroethyl-2-yl	


TABLE 1 (CONTINUED)

	1-fluoroprop-3-yl
	1-fluoro-but-4-yl
5	1-fluoro-hex-5-yl
	chloromethyl
	1-chloroeth-2-yl
	1-chloroprop-3-yl
	1-chlorobut-4-yl
10	bromomethyl
	1-bromoeth-2-yl
	1-bromoprop-3-yl
	1-bromobut-4-yl
	vinyl
15	1-propene-2-yl
	1-butene-4-yl
	1-pentene-5-yl
	1-hexene-6-yl
	cyclopropyl
20	cyclobutyl
	cyclopentyl
	cyclohexyl

In the above Formula (1), R is preferably C₁-C₆ alkyl or C₁-C₆ haloalkyl. A preferred C₁-C₆ alkyl group is methyl. A preferred C₁-C₆ haloalkyl group is methyl. A preferred C₁-C₆ haloalkyl group is fluoro-C₁-C₆ alkyl. A further preferred
 5 fluoro-C₁-C₆ alkyl group is the 2-fluoroeth-1-yl group.

In the above Formula (1), it is preferred that A and A' are taken together to form a group of the formula


10



15


It is further preferred that Y is nitrogen and A and A' are taken together to form a group of the formula

20



25

for example, providing a compound of the formula



or a pharmaceutically acceptable salt thereof. Two further preferred compounds of the above formula are where R is methyl or 2-fluoroethyl.

This invention also provides a method for
5 treating infectious diseases in man and other animals and pharmaceutical formulations suitable for administration in the treatment method. The therapeutic method of this invention comprises administering to man or other animals an antibiologically effective non-toxic
10 dose of a compound represented by Formula (1) or a pharmaceutically acceptable salt thereof.

An antibiologically effective amount is an amount between about 25 mg and about 2 grams. The compound, salt or ester may be administered in a single
15 dose or in multiple doses throughout the day. Treatment may continue for a week to ten days or longer depending upon the duration of the infection. The particular dose and regimen can depend on such factors as the weight and age of the patient, the particular
20 causative organism, the severity of the infection, the general health of the patient, and the tolerance of the individual to the antibiotic.

The cephalosporin may be administered parenterally, subcutaneously or rectally. As with other β -lactam
25 antibiotics, the method of this invention may be used prophylactically to prevent infections after exposure or before possible exposure, e.g., preoperatively. The antibiotic may be administered by conventional methods, e.g., by syringe or by intravenous drip.

The pharmaceutically-acceptable salts as noted above can be useful forms of the antibiotics for preparing antibiotic formulations.


5 The pharmaceutical formulations of the invention comprise an antibiotically effective non-toxic amount of a compound represented by Formula (1) or a pharmaceutically acceptable non-toxic salt thereof, and a pharmaceutically acceptable carrier.

10 Parenteral formulations of the antibacterial agent for injection are formulated with Water-for-Injection, Ringer's solution, physiological saline or glucose solution. The antibiotic also may be administered in an intravenous fluid by the drip method.


15 For parenteral use the antibacterial agent of Formula (1) or a pharmaceutically acceptable salt thereof, can be made up preferably in dry crystalline powder form or as a lyophilized powder and filled into vials. Such vials may contain between about 100 mg and about 2 grams of antibiotic per vial.

20 As a further aspect of the present invention, there are provided novel intermediates of Formula (2):

25



wherein R^o is amino or a a protected amino group; R' is hydrogen or a carboxy-protecting group; and A and A' are independently hydrogen, C₁-C₆ alkyl, phenyl, nitro, amino, a 5-6 membered heterocycle containing nitrogen or sulfur, or C₁-C₆ alkoxy; or A and A' taken together form a group of the formulae



wherein X is hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxy-carbonyl, amino, nitro, or carboxy, and Y is nitrogen or carbon.

In Formula (2), the term "carboxy-protecting group" refers to one of the ester derivatives of the carboxylic acid group commonly employed to block or protect the carboxylic acid group while reactions are carried out on other functional groups on the compound. Examples of such carboxylic acid protecting groups include 4-nitrobenzyl, 4-methoxybenzyl, 3,4-di-methoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl, 2,4,6-trimethylbenzyl, pentamethylbenzyl, 3,4-methylene-dioxybenzyl, benzhydryl, 4,4'-dimethoxybenzhydryl, 2,2',4,4'-tetramethoxybenzhydryl, t-butyl, t-amyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, 4,4',4''-trimethoxytrityl, 2-phenylprop-2-yl, trimethylsilyl, t-butyl-dimethylsilyl, phenacyl, 2,2,2-trichloroethyl,

β -(trimethylsilyl)ethyl, β -(di(n-butyl)methyl)dilyl)-ethyl, p-toluenesulfonylethyl, 4-nitrobenzylsulfonylethyl, allyl, cinnamyl, 1-(trimethylsilylmethyl)prop-1-en-3-yl, and like moieties. The species of carboxy-protecting group employed is not critical so long as the derivatized carboxylic acid is stable to the condition of subsequent reaction(s) on other positions of the molecule and can be removed at the appropriate point without disrupting the remainder of the molecule. In particular, it is important not to subject the carboxy-protected molecule to strong nucleophilic bases or reductive conditions employing highly activated metal catalysts such as Raney nickel. (Such harsh removal conditions are also to be avoided when removing amino-protecting groups discussed herein.) Preferred carboxylic acid protecting groups are the allyl, the benzhydryl, and the p-nitro benzyl groups. Similar carboxy-protecting groups used in the cephalosporin, penicillin and peptide arts can also be used to protect a carboxy group. Further examples of these groups are found in E. Haslam, "Protective Groups in Organic Chemistry", J.G.W. McOmie, Ed., Plenum Press, New York, N.Y., 1973, Chapter 5, and T.W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1981, Chapter 5.

The term "protected amino group" as used in Formula (2) refers to an amino group substituted by a group commonly employed to block or protect the amino functionality while reacting other functional groups on the compound. Examples of such amino-protecting


groups include the formyl group, the trityl group, the t-butoxycarbonyl group, the phthalimido group, the phenoxyacetyl, trichloroacetyl group, the chloroacetyl, bromoacetyl and iodoacetyl groups, urethane-type blocking
5 groups such as benzyloxycarbonyl, 4-phenylbenzyloxy-carbonyl, 2-methylbenzyloxycarbonyl, 4-methoxybenzyloxy-carbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxy-carbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyl-oxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenxyloxy-
10 carbonyl, 4-cyanobenzyloxycarbonyl, 2-(4-xenyl)iso-propoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenyl-prop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(p-toluyl)prop-2-yloxycarbonyl, cyclo-pentanyloxy-carbonyl, 1-methylcyclopentanyloxycarbonyl,
15 cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxy-carbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluyl-sulfonyl)ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl, 9-fluorenyl-methoxycarbonyl ("Fmoc"), 2-(trimethylsilyl)ethoxy-
20 carbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)-prop-1-enyloxycarbonyl, 5-benzisoxazolylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxy-carbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropyl-methoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl, iso-bornyloxycarbonyl, 1-piperidyloxycarbonyl and the like;
25 the benzoylmethylsulfonyl group, the 2-(nitro)phenyl-sulfenyl group, the diphenylphosphine oxide group and like amino-protecting groups. The species of amino-protecting group employed is not critical so long as the




2058822

derivatized amino group is stable to the condition of subsequent reaction(s) on other positions of the molecule and can be removed at the appropriate point without disrupting the remainder of the molecule. Preferred amino-protecting groups are the allyloxycarbonyl, the phenoxyacetyl, the t-butoxycarbonyl, and the trityl groups. Similar amino-protecting groups used in the cephalosporin, penicillin and peptide art are also embraced by the above terms. Further examples of groups referred to by the above terms are described by J.W. Barton, "Protective Groups in Organic Chemistry", J.G. W. McOmie, Ed., Plenum Press, New York, N.Y., 1973, Chapter 2, and T.W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1981, Chapter 7.

In Formula (2), it is preferred that A and A' are taken together to form a group of the formula




It is especially preferred that A and A' are taken together to form a group of the formula




thus providing a compound of the formula

5



The compounds of formula (2) are useful as
 10 intermediates in the preparation of the antibacterial
 agents of Formula(1) above. The compounds of formula
 (2) may be prepared by the methodology as taught in
 scheme (1) above displacing the 3-triflate moiety with
 the desired thiol of the formula

15



20 utilizing a 7-protected amino nucleus.

The final products (1) may then be prepared
 from intermediates of formula (2) by deprotection of the
 7-amino function if necessary, followed by acylation
 with a desired acyl group, and subsequent amino/carboxy
 25 protecting group removal.

The following Experimental Section provides
 further examples of the various aspects of the present
 invention but is not to be construed as limiting the
 scope therefore.

30

Experimental SectionPreparation 17 β -amino-3-chloro-3-cephem-4-carboxylic acid

5

The title compound may be prepared by the method of Chauvette, U.S. Patent No. 4,064,343, dated December 20, 1977.

10

Preparation 23-(t-butyloxycarbonyl)amino pyridine

A 76.13 g (0.81 mol) sample of 3-aminopyridine was dissolved in 500 ml of water, along with 150 ml of t-butanol and 34 g (0.85 mol) of NaOH, cooled in an ice bath, and treated with 200 g (0.92 mol) of di-t-butyldi-carbonate. After about 2.5 days, another 100 g of di-t-butyl dicarbonate was added. The reaction mixture was then poured into an ethyl acetate/water mixture. The organic phase was separated and the remaining aqueous phase was extracted with ethyl acetate. The combined organic portions were dried over anhydrous sodium sulfate, concentrated in vacuo, and purified via flash chromatography to provide 97 g (80%) of the title compound.

25

¹H NMR: (300 MHz, CDCl₃) δ 8.43(d, J=1.5Hz, 1H), 8.26 (d, J=3Hz, 1H), 7.97 (br d, J=6Hz, 1H), 7.24-7.20 (m, 1H), 6.81 (br s, 1H), 1.51 (s, 9H).

30



IR: (KBr, cm^{-1}) 3167, 2986, 1716, 1598, 1545, 1407,
1566, 1288, 1233, 1154, 1017

MS: FDMS m/e 195 (M⁺)

5

UV: (ethanol) $\lambda=281$ nm ($\epsilon=3350$)
 $\lambda=235$ nm ($\epsilon=15200$)

Preparation 3

10

3-(t-Butyloxycarbonyl)amino-4-mercaptopyridine

A 10 g (51.5 mmol) sample of 3-(t-butyloxy-
carbonyl)amino pyridine was dissolved in 110 ml of
15 tetrahydrofuran and cooled to -78°C under nitrogen.
An 80 ml (128 mmol, 1.6 M in hexanes) sample of n-
butyllithium was then added in two portions. The
reaction mixture was then placed in an acetone/ice bath
to allow the resulting solid to dissolve. After about
20 2 hours, the reaction mixture was then cooled to -78°C
and treated with 2 g (7.8 mmol) of elemental sulfur.
After about $\frac{1}{2}$ hour, the reaction mixture was allowed
to warm to room temperature and was quenched with a
saturated NH_4Cl solution. Work-up and flash chroma-
25 tography (50% Hexane/ethyl acetate) provided 5.24 g
(45%) of the title compound.

m.p. = 170° - 171°C (dec.)

¹H NMR: (300 MHz, DMSO-d₆) δ 12.88 (br s, 1H), 8.95 (s, 1H), 8.45 (br s, 1H), 7.62 (br d, J=3Hz, 1H), 7.44 (d, J=3Hz, 1H), 1.49 (s, 9H).

5 IR: (KBr, cm⁻¹) 3239, 2978, 2885, 2741, 1721, 1608, 1530, 1492, 1436, 1384, 1213, 1161, 1085

MS: FDMS m/e 227 (M⁺)

10 UV: (ethanol) λ=345nm (ε=19600)
λ=259nm (ε=10200)
λ=224nm (ε=17200)

Preparation 4

15

3-Amino-4-mercapto-pyridine hydrochloride

A 13.78 g (0.06 mol) sample of 3-(t-butyloxy-carbonyl)amino-4-mercapto pyridine was dissolved with
20 acetic acid (250 mL) and added to an ice cold solution of ~3N HCl in acetic acid which had been made by bubbling HCl_(g) through glacial acetic acid (100 mL). After about four hours the resulting solid was filtered, washed with diethylether and dried in vacuo to yield 10.4 g (~100%)
25 of the title compound.

m.p.: >200°C

¹H NMR: (300 MHz, DMSO-d₆) δ 8.17 (s, 1H), 7.99 (d, J=3 Hz, 1H), 7.81 (d, J=3 Hz, 1H), 5.60-4.00 (br, 4H).

5 IR: (KBr, cm⁻¹) 3184, 3054, 2848, 1639, 1586, 1482, 1442, 1134, 1123

MS: FDMS m/e 126 (M-36)

10 UV: (ethanol) λ=355nm (ε=13900)
λ=264nm (ε= 6830)
λ=223nm (ε=13100)

Preparation 5

15

2-Mercapto-5-pyridinothiazole

A 13 g (0.198 mol) sample of potassium hydroxide was dissolved in 32 ml of water and 154 ml of methanol. This solution was then treated with 3.8 ml (0.063 mol) of CS₂, followed by a 10.4 g (0.06 mol) sample of 3-amino-4-mercaptopyridine hydrochloride. After stirring at reflux overnight, the reaction mixture was treated with decolorizing carbon and filtered through Hyflo Super Cel[™]. The filtrate was acidified with acetic acid causing a solid to form. The resulting solid was dried in vacuo at 50°C for about 3 hours and at room temperature for about 2.5 days to provide 8.19 g (81%) of the title compound.

30

m.p. >310 dec.

¹H NMR: (300 MHz, DMSO-d₆) δ 14.03 (br s, 1H), 8.46
5 (s, 1H), 8.33 (d, J=6Hz, 1H) 7.75 (d, J=6Hz,
1H)

IR: (KBr cm⁻¹) 3440(br), 2650(br), 2510(br), 1528,
1457, 1305, 1294, 1265, 1256, 1039, 1024, 815

10 MS: EI MS m/e 168 (M+)

Preparation 6

2-isothiocyanato-3-nitro pyridine

15

A 10 g sample of 2-chloro-3-nitropyridine, an
8 g sample of potassium isothiocyanate, and 75 ml of
acetic acid were combined and refluxed for 2 h. The
reaction mixture was then cooled and poured into 400 ml
20 of ice/H₂O. The resulting solid was washed with water,
redissolved in ethyl acetate and washed (4x) with
water. The ethyl acetate solution was then treated
with activated carbon, dried over anhydrous Na₂SO₄,
filtered and evaporated to dryness to provide 3.72 g of
25 the title compound. m.p. = 115°-118°C.

¹H NMR: (300 MHz, CDCl₃) δ 8.62 (m, 1H), 8.22 (d,
J = 6Hz, 1H), 7.46 (m, 1H).

Preparation 72-Mercapto-3-nitropyridine

5 A 50 ml sample of ethanol was treated with
612 mg of sodium at reduced temperature (ice bath)
under substantially anhydrous conditions. The reaction
mixture was then treated with a 3.6 g (0.02 mol) sample
(in portions) of the title compound of preparation 6.
10 The reaction mixture was stirred for 2 h, diluted with
250 ml of H₂O and evaporated in vacuo. The resulting
solid was filtered off and discarded. The solution
was then acidified with acetic acid to pH=4.5 and
yellowish-red crystals formed. The title compound was
15 filtered off, washed with water and dried under vacuum
over a desiccant to provide 1.1 g (m.p.=185°-7°C
(dec.))

¹H NMR: (300 MHz, CDCl₃) δ 8.09 (d, J = 7Hz, 1H), 7.89
(d, J = 7Hz, 1H), 6.84 (dd, J = 6, 3Hz, 1H).

20

IR: (KBr cm⁻¹) 3119, 2872, 1611, 1577, 1527, 1349,
1330, 1240, 1141

MS: EI MS m/e 126 (M⁺)

25

Preparation 82-Mercapto-3-aminopyridine

5 A 100 ml sample of concentrated HCL(aq) was cooled in an ice bath and treated with 100 g (0.53 mol) of SnCl₂. The reaction mixture was then treated with a 14 g (0.11 mol) sample of the title compound from preparation 7, in portions, and stirred for 3 hours.

10 The reaction mixture was then evaporated to a solid, dissolved in 1 L H₂O, and treated with H₂S(g) for 30 min., while heating over a steam bath. The resulting solid was filtered off, washed with hot H₂O and discarded. The combined aqueous portions were
15 evaporated to afford a solid. The resulting solid was digested (2x) with hot concentrated NH₄OH. The resulting solid was filtered and discarded and the NH₄OH solution was evaporated to afford a wet solid, which was, in turn, mobilized in H₂O. The resulting
20 yellow/green title compound was filtered, washed with H₂O, and dried in vacuo at 40° over desiccant.

Yield=4.20 g (30%)

m.p.=127°-128°C

25 ¹H NMR: 300 MHz, CDCl₃/DMSO-d₆) δ 6.91 (m, 1H), 6.65 (d, J = 5Hz, 1H), 6.46 (m, 1H), 5.03 (s, 2H).



Preparation 92-Mercapto-7-pyridinothiazole

5 A 2.8 g (85%) sample of KOH was dissolved in
16 ml of H₂O and 50 ml of methanol. A 2.6 g sample of
CS₂ was then added and washed in with 30 ml of methanol.
A 4 g (23.8 mmol) sample of 2-mercapto-3-aminopyridine
was added and the reaction mixture refluxed overnight.
10 After cooling, the reaction mixture was treated with
activated carbon and filtered through "Super Cel"[™], while
washing the "Super Cel"[™] pad with a small amount of
methanol. The solution was then acidified to pH=5.5
with acetic acid. The title compound precipitated from
15 this solution as a yellowish solid and was dried at 60°C
over a desiccant. Yield=3.29 g

m.p. = 285-287°C (dec)

20 ¹H NMR: (300 MHz, DMSO-d₆) δ 8.38 (dd, J = 3, 1.5 Hz,
1H), 7.61 (dd, J = 4, 1.5 Hz, 1H), 7.43 (dd,
J = 5, 3Hz, 1H), 3.33 (br s, 1H) .

IR: (KBr cm⁻¹) 3040, 2700, 2540, 1597, 1523, 1399,
25 1311, 1302, 1274, 1132, 876.

MS: EI MS m/e 169 (m+1)

Preparation 10Ethyl(2-(triphenylmethyl)-aminothiazol-4-yl)-
2-bromoeth-1-yl-oximinoacetate

5

A 9.88 g (0.02 mol) sample of ethyl-(2-(triphenylmethyl)aminothiazol-4-yl)oximinoacetate was dissolved in 20 ml of N,N'-dimethylformamide and treated with 8.28 g (0.06 mol) of powdered potassium carbonate. After $\frac{1}{2}$ h of stirring, 17.3 ml of 1,2-dibromoethane was added and the reaction mixture was stirred overnight under argon.

The reaction mixture was then poured into 100 ml of CH₂Cl₂/200 ml H₂O. The aqueous layer was again extracted with CH₂Cl₂. The combined CH₂Cl₂ phase was washed with H₂O and brine, dried over anhydrous MgSO₄, filtered, and evaporated in vacuo to provide an oil. Liquid chromatography (25% hexane/CH₂Cl₃) provided 7.16 g (63.4%) of the title compound. m.p. = 55°C.

15
20

¹H NMR: (300 MHz, CDCl₃) δ 7.32 (s, 15H), 6.52 (s, 1H), 4.55-4.46 (m, 2H), 4.38 (q, J = 4 Hz, 2H), 3.63-3.53 (m, 2H), 1.37 (t, J = 4 Hz, 3H)

25

Elem. Anal:

calc'd: C: 59.58; H: 4.64; N: 7.44

obs'd: C: 59.36; H: 4.61; N: 7.18

Preparation 11Ethyl(2-(triphenylmethyl)aminothiazol-4-yl)-
2-fluoroeth-1-yl-oximino acetate

5

The title compound was prepared in a manner analogous to that of Preparation 10, substituting 1-bromo-2-fluoroethane as the alkylating agent.

Yield = 3.3 g

10

¹H NMR: (300 MHz, DMSO-d₆) δ 8.77 (s, 1H), 7.39-7.12 (m, 15H), 6.92 (s, 1H), 4.60 (t, J = 3Hz, 1H), 4.44 (t, J = 3Hz, 1H), 4.26 (t, J = 3Hz, 1H), 4.16 (t, J = 3Hz, 1H), 4.16 (t, J = 3Hz, 1H), 3.90 (q, J = 4Hz, 2H), 1.06 (t, J = 4Hz, 3H).

15

Preparation 12(2-(Triphenylmethyl)aminothiazol-4-yl)-2-
20 fluoroeth-1-yl-oximinoacetic acid

A 2.5 g (5 mmol) sample of the title compound of preparation 11 was dissolved in 20 ml of ethanol and 5 ml (10 mmol) of 2N NaOH. After stirring for 2 h at 25 50°C, the sodium salt of the acid crystallized. This solid was slurried in H₂O/CHCl₃ and acidified with 1N HCl. The aqueous layer was extracted again with CHCl₃ and the combined CHCl₃ phase was washed with water,

A

brine, and dried over anhydrous Na_2SO_4 . The CHCl_3 phase was then evaporated in vacuo to provide 1.52 g (63.9%) of the title compound as a foam.

5 m.p.=125.33°C (dec)

^1H NMR: (300 MHz, CDCl_3) δ 9.70 (br s, 1H), 7.30-7.22 (m, 15H), 6.52 (s, 1H), 4.65 (t, J = 3Hz, 1H), 4.49 (t, J = 3Hz, 1H), 4.37 (t, J = 3Hz, 1H),
10 4.27 (t, J = 3Hz, 1H)

IR: (CDCl_3 , cm^{-1}) 3000, 1735, 1592, 1529, 1449, 1186, 1070, 1035

15

Example 1

7 β -[(2-aminothiazol-4-yl)-(Z)-methoximinoacetyl]amino-3-[2-(5-pyridinothiazolothio)]-3-cephem-4-carboxylic acid

20

A. 7 β -[(2-(triphenylmethyl)aminothiazol-4-yl)-(Z)-methoximinoacetyl]amino-3-chloro-3-cephem-4-carboxylic acid

25 A 39.8 g (0.17 mol) sample of 7 β -amino-3-chloro-3-cephem-4-carboxylic acid was suspended in 800 ml of N,N'-dimethylformamide and treated with 100 g (0.49 mol) of bis(dimethylsilyl)urea and heated to about 50°-65°C for about 1 hour.

In another reaction vessel, a 100 g (0.21 mol) sample of 2-(triphenylmethyl)aminothiazol-4-yl-(Z)-methoximinoacetic acid was dissolved in 800 ml of N,N'-dimethylformamide and cooled in an ice/acetone bath.

5 The reaction mixture was then treated with 23 ml (0.21 mol) of N-methylmorpholine followed by 25 g (0.20 mol) of oxalyl chloride.

10 In the first reaction vessel above, the reaction mixture was treated with 32 ml (0.40 mol) of pyridine and transferred via cannula to the second reaction vessel over 50 minutes.

The reaction mixture was then poured into about 2.5 L of ice/H₂O and the resulting solid air dried to provide 116 g of the title compound (3:1 Δ^3 : Δ^2 mixture).

¹H NMR: (300 MHz, DMSO-d₆) δ 9.61 (d, J = 9Hz, 1H), 8.83 (s, 1H x $\frac{1}{4}$), 8.80 (s, 1H x $\frac{3}{4}$), 7.46-7.10 (br m, 15H), 6.83 (s, 1H), 6.68 (s, 1H x $\frac{1}{4}$), 5.72-5.66 (m, 1H x $\frac{3}{4}$), 5.60-5.54 (m, 1H x $\frac{1}{4}$), 5.23-5.17 (m, 1H x $\frac{1}{4}$), 5.20 (d, J = 5Hz, 1H x $\frac{3}{4}$), 4.83 (s, 1H x $\frac{1}{4}$), 3.80 (s, 3H), 3.79 (ABq, J = 20Hz, 2H x $\frac{3}{4}$).

25 B. Benzhydryl 7 β -[2-(triphenylmethyl)aminothiazol-4-yl-(Z)-methoximinoacetyl]amino-3-chloro-3-cephem-4-carboxylate

30 The material from part A, above, was dissolved in 500 ml of CH₃CN and treated with 10 g (XS) of diphenyldiazomethane and stirred at room temperature for about

2.5 days. The reaction mixture was then quenched with acetic acid and concentrated in vacuo, utilizing toluene to azeotrope excess acetic acid. Purification via flash chromatography (25% and 50% ethyl acetate/hexane) provided 15.46 g of a 2:1 (Δ^2/Δ^3) mixture.

^1H NMR: (300 MHz, DMSO- d_6) δ 9.60 (d, 6Hz, 1H), 8.80 (s, 1H), 7.46-7.02 (br. m, 25H), 6.92 (s, 1H x 1/3), 6.88 (s, 1H x 2/3), 6.84 (s, 1H x 2/3), 6.78 (s, 1H x 2/3), 6.67 (s, 1H x 1/3), 5.76-5.70 (m, 1H x 1/3), 5.51-5.45 (m, 1H x 2/3), 5.28-5.22 (m, 1H x 1/3), 5.26 (s, 1H x 2/3), 5.12 (d, J = 4Hz, 1H x 2/3), 3.79 (ABq, J = 19Hz, 2H x 1/3), 3.77 (s, 3H).

15

C. Benzhydryl 7 β -[2-(triphenylmethyl)amino-thiazol-4-yl-(Z)-methoximino]acetyl-3-[2-(5-pyridinothiazolothio)]-3-cephem-4-carboxylate

A 92 mg (2.3 mmol; 60% in oil) sample of NaH was washed with hexanes and suspended in 50 ml of tetrahydrofuran and treated with a 390.9 mg (2.3 mmol) sample of 2-mercapto-5-pyridinothiazole and heated. This solution was transferred via cannula to a 5.7 g (2.3 mmol) sample of the compound prepared in part B, above, dissolved in 50 ml of tetrahydrofuran. The reaction mixture was then treated with 15 ml of 1N HCl and poured into an ethyl acetate/water mixture. The organic phase was washed

25

with brine, dried over anhydrous Na_2SO_4 , filtered and concentrated in vacuo. After column chromatography (75%-90% ethyl acetate/hexane), the pure Δ^3 isomer crystallized out. (0.31 g, 34%).

5

^1H NMR: (300 MHz, DMSO-d_6) δ 9.70 (d, $J = 9\text{Hz}$, 1H), 9.10 (s, 1H), 8.79 (s, 1H), 8.47 (d, $J = 7\text{Hz}$, 1H), 8.08 (d, $J = 7\text{Hz}$, 1H), 7.38-7.03 (br m, 25H), 6.91 (s, 1H), 6.68 (s, 1H), 5.90-5.82 (m, 1H), 5.37 (d, $J = 8\text{Hz}$, 1H), 3.83 (ABq, $J = 20\text{Hz}$, 2H), 3.78 (s, 3H).

10

IR: (KBr, cm^{-1}) 3402 (br), 3030, 2938, 1786, 1738, 1695, 1522, 1496, 1371, 1278, 1223, 1044, 700.

15

MS: FABMS m/e 958 (m^+)

OR: $[\alpha]_D = -133.33$ degrees @ 589 nm, 5 mg/DMSO

20

Elem. Anal:

calc'd: C: 63.93; H: 4.10; N: 10.23

obs'd: C: 64.19; H: 4.06; N: 10.43.

D. Deprotection to provide title compound

25

A 0.42 g (438 μmol) sample of the product from part C, above, was suspended with 7 ml of triethylsilane and 10 ml of CH_2Cl_2 and treated with 5 ml of trifluoroacetic acid and stirred at room temperature.



The reaction mixture was then concentrated, in vacuo, utilizing toluene to azeotrope excess trifluoroacetic acid. The resulting residue was purified by reverse phase chromatography (10%-20% CH₃CN/H₂O).

5

¹H NMR: (300 MHz, DMSO-d₆) δ 9.75 (d, J = 9Hz, 1H), 9.18 (s, 1H), 8.49 (d, J = 6Hz, 1H), 8.19 (d, J = 6Hz, 1H), 7.21 (br s, 2H), 6.71 (s, 1H), 5.94 (dd, J = 5Hz, 10Hz, 1H), 5.35 (d, J = 6Hz, 1H), 3.88 (ABq, J = 15Hz, 2H), 3.85 (s, 3H).

10

IR: (KBr. cm⁻¹) 3395, 1782, 1621, 1532, 1381, 1037.

MS: FABMS m/e 550 (m⁺)

15

UV: (EtOH) λ = 286 nm (E 22700)
λ = 231 nm (E 34200)


OR: 5 mg/DMSO [α]_D = -123.26 degrees @ 589 nm

20

Examples 2-5

Examples 2 through 5, which follow, were prepared in a manner essentially as described in Example 1, by utilizing different mercaptans of the formula

25



30

Example 2

5 7 β -[(2-aminothiazol-4-yl)-(Z)-(2-fluoroeth-1-yl)-oximinoacetyl]amino-3-[2-(5-pyridinothiazolo)]thio-3-cephem-4-carboxylic acid

10 ¹H NMR: (300 MHz, DMSO-d₆): δ 9.70 (d, 1H, J=10Hz); 9.03 (s, 1H); 8.39 (d, 1H, J=5Hz); 8.03 (d, 1H, J=5Hz); 7.20 (s, 2H); 6.72 (s, 1H); 5.73 (m, 1H); 5.19 (d, 1H, J=7Hz); 4.67 (t, 1H, J=5Hz); 4.55 (t, 1H, J=5Hz); 4.32 (t, 1H, J=5Hz); 4.22 (t, 1H, J=5Hz); 3.63 (ABq, 2H, J=18Hz)

15 IR: (KBr) 3420, 1774, 1668, 1663, 1653, 1617, 1534, 1388 cm⁻¹

MS: (FAB) m/e = 604 (m+1)

20 UV: (EtOH) λ = 288 nm (ϵ = 21700); 232 nm (ϵ = 31400)

OR: $[\alpha]_{\text{DMSO}}^{\circ}$ = -89.22°

Example 3

25

7 β -[(2-aminothiazol-4-yl)-(Z)-(2-fluoroeth-1-yl)-oximinoacetyl]amino-3-[2-(7-pyridinothiazolo)]-3-cephem-4-carboxylic acid

30 Yield = 13% overall (22.8 mg)

¹H NMR: (300 MHz, DMSO-d₆) δ 9.68 (d, 1H, J=10Hz);
8.25 (d, 1H, J=5Hz); 8.14 (d, 1H, J=10Hz);
7.45 (m, 1H); 7.20 (s, 2H); 6.72 (s, 1H);
5.70 (m, 1H); 5.20 (d, 1H, J=5Hz); 4.70 (t,
5 1H, J=5Hz); 4.53 (t, 1H, J=5Hz); 4.30 (t, 1H,
J=5Hz); 4.20 (t, 1H, J=5Hz); 3.63 (ABq, 2H,
J=15Hz)

MS: (FAB) m/e = 604 (m+1)
10

Example 4

7β-[(2-aminothiazol-4-yl)-(Z)-(2-fluoroeth-1-yl)-
oximinoacetyl]amino-3-(thiazol-2-yl)thio-3-cephem-
15 4-carboxylic acid

Yield = 63 mg (71%)

¹H NMR: (300 MHz, DMSO-d₆) δ 9.67 (d, 1H, J=10Hz);
20 7.76 (s, 2H); 7.20 (s, 2H); 6.72 (s, 1H);
5.80-5.70 (m, 1H); 5.20 (d, 1H, J=5Hz); 4.66
(t, 1H, J=5Hz); 4.50 (t, 1H, J=5Hz); 4.28 (t,
1H, J=5Hz); 4.19 (t, 1H, J=5Hz); 3.50 (ABq,
2H, J=15Hz)

25 IR: (KBr) 3400, 1768, 1653, 1614, 1535, 1389, 1350,
1035 cm⁻¹

MS: (FAB) m/e=(m-1) 553
30

2058822

X-7505

-49-

UV: (ethanol) $\lambda=284$ nm ($\epsilon=14900$); 231 nm ($\epsilon=18100$)

Example 5


5 7 β -[2-aminothiazol-4-yl-(Z)-(2-fluoroeth-1-yl)-
oximinoacetyl]amino-3-[(benzothiazol-2-yl)thio]-3-
 cephem-4-carboxylic acid, sodium salt

10 ¹H NMR: (300 MHz, DMSO-d₆): δ 9.67 (d, 1H, J=10Hz);
 7.92 (d, 1H, J=10Hz); 7.78 (d, 1H, J=10Hz);
 7.43-7.26 (m, 2H); 7.20 (s, 2H); 6.73 (s, 1H);
 5.66 (m, 1H); 5.15 (d, 1H, J=5Hz); 4.70 (t, 1H,
 J=5Hz); 4.53 (t, 1H, J=5Hz); 4.30 (t, 1H,
 J=5Hz); 4.20 (t, 1H, J=5Hz); 3.64 (ABq, 2H,
15 J=15Hz)

MS: (FAB) m/e=603 (m+1)


THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A compound of the formula:



wherein R is hydrogen, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₆ cycloalkyl, or C₁-C₆ haloalkyl;

A and A' are taken together to form a group of the formula



wherein X is hydrogen, halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkoxy carbonyl, amino, nitro, or carboxy; and Y is nitrogen; or a pharmaceutically acceptable salt thereof.


2. The compound as recited in claim 1 wherein R is C₁-C₆ alkyl or C₁-C₆ haloalkyl.

3. The compound as recited in claim 2 wherein R is methyl or fluoro-C₁-C₆ haloalkyl.

4. The compound as recited in claim 3 wherein R is 2-fluoroethyl.

5. The compound as recited in claim 1 wherein A and A' form

2058822



6. The compound as recited in claim 5 wherein R is methyl or 2-fluoroethyl.

7. A pharmaceutical composition which comprises a compound of any one of claims 1 to 6 combined with one or more pharmaceutically acceptable carriers, excipients or diluents.

8. A compound of any one of claims 1 to 6 for use as an antibiotic.