



(11) **EP 2 203 430 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:  
**15.06.2011 Bulletin 2011/24**

(21) Application number: **08831558.5**

(22) Date of filing: **17.09.2008**

(51) Int Cl.:  
**C07D 239/22** <sup>(2006.01)</sup> **C07D 239/54** <sup>(2006.01)</sup>  
**C07D 401/10** <sup>(2006.01)</sup> **C07D 403/10** <sup>(2006.01)</sup>  
**C07D 405/10** <sup>(2006.01)</sup> **C07D 409/10** <sup>(2006.01)</sup>  
**C07D 413/10** <sup>(2006.01)</sup> **C07D 239/553** <sup>(2006.01)</sup>  
**A61K 31/513** <sup>(2006.01)</sup> **A61K 31/506** <sup>(2006.01)</sup>  
**A61P 31/14** <sup>(2006.01)</sup>

(86) International application number:  
**PCT/US2008/076594**

(87) International publication number:  
**WO 2009/039135 (26.03.2009 Gazette 2009/13)**

(54) **N-PHENYL-DIOXO-HYDROPYRIMIDINES USEFUL AS HEPATITIS C VIRUS (HCV) INHIBITORS**

ALS INHIBITOREN DES HEPATITIS-C-VIRUS (HCV) GEEIGNETE N-PHENYLDIOXOHYDROPYRIMIDINE

N-PHENYL-DIOXO-HYDROPYRIMIDINES UTILES EN TANT QU'INHIBITEURS DU VIRUS DE L'HEPATITE C (VHC)

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR**

(30) Priority: **17.09.2007 US 972887 P**  
**13.09.2008 US 96794 P**

(43) Date of publication of application:  
**07.07.2010 Bulletin 2010/27**

(60) Divisional application:  
**11163527.2**

(73) Proprietor: **Abbott Laboratories**  
**Abbott Park, Illinois 60064 (US)**

(72) Inventors:  
• **DONNER, Pamela L.**  
**Mundelein**  
**IL 60060 (US)**  
• **RANDOLPH, John T.**  
**Libertyville**  
**IL 60048 (US)**  
• **KRUEGER, Allan C.**  
**Gurnee**  
**IL 60031 (US)**  
• **BETEBENNER, David A.**  
**Libertyville**  
**IL 60048 (US)**

- **HUTCHINSON, Douglas K.**  
**Antioch**  
**IL 60002-1863 (US)**
- **LIU, Dachun**  
**Waukegan, IL 60085 (US)**
- **LIU, Yaya**  
**Buffalo Grove**  
**IL 60089 (US)**
- **LONGENECKER, Kenton L.**  
**Grayslake**  
**IL 60030 (US)**
- **MARING, Clarence J.**  
**Palatine**  
**IL 60067 (US)**
- **PRATT, John K.**  
**Kenosha**  
**WI 53142 (US)**
- **ROCKWAY, Todd W.**  
**Grayslake**  
**IL 60030 (US)**
- **STEWART, Kent D.**  
**Gurnee**  
**IL 60031 (US)**
- **WAGNER, Rolf**  
**Antioch**  
**IL 60002 (US)**
- **BARNES, David M.**  
**Bristol**  
**WI 53104 (US)**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 2 203 430 B1**

- CHEN, Shuang  
Gurnee  
IL 60031 (US)
- FRANCZYK II, Thaddeus S.  
Lake Villa  
IL 60046 (US)
- GAO, Yi  
Vernon Hills  
IL 60061 (US)
- HAIGHT, Anthony R.  
Wadsworth  
IL 60083 (US)
- HENGEVELD, John E.  
Kenosha  
WI 53143 (US)
- HENRY, Rodger F.  
Wildwood  
IL 60030 (US)
- KOTECKI, Brian J.  
Oak Creek  
IL 53154 (US)
- LOU, Xiaochun  
Long Grove  
IL 60047 (US)
- ZHANG, Geoff G. Z.  
Libertyville  
IL 60048 (US)

(74) Representative: **Wright, Robert Gordon McRae**  
**Elkington and Fife LLP**  
**Prospect House**  
**8 Pembroke Road**  
**Sevenoaks**  
**Kent TN13 1XR (GB)**

(56) References cited:

- **RAFFAELE DE FRANCESCO ET AL:**  
**"Approaching a new era for hepatitis C virus**  
**therapy: inhibitors of the NS3-4A serine protease**  
**and the NS5B RNA-dependent RNA polymerase"**  
**ANTIVIRAL RESEARCH, ELSEVIER SCIENCE**  
**BV., AMSTERDAM, NL, vol. 58, 1 January 2003**  
**(2003-01-01), pages 1-16, XP002414639 ISSN:**  
**0166-3542**
- **DE FRANCESCO RAFFAELE ET AL: "Challenges**  
**and successes in developing new therapies for**  
**hepatitis C" NATURE 18 AUG 2005,, vol. 436, no.**  
**7053, 18 August 2005 (2005-08-18), pages**  
**953-960, XP002504755**
- **KOCH UWE ET AL: "2-(2-thienyl)-5,6-dihydroxy-**  
**4-carboxypyridines as inhibitors of the**  
**hepatitis C virus NS5B polymerase: Discovery,**  
**SAR, modeling, and mutagenesis" JOURNAL OF**  
**MEDICINAL CHEMISTRY, vol. 49, no. 5, March**  
**2006 (2006-03), pages 1693-1705, XP002508123**  
**ISSN: 0022-2623**

**Description**

## CROSS-REFERENCE TO RELATED PATENT APPLICATIONS

5 **[0001]** This patent application claims priority to U.S. Provisional Patent Application No. 60/972,887 (filed September 17, 2007) and U.S. Provisional Patent Application No. 61/096,794 (filed September 13, 2008). The entire text of those applications is incorporated by reference into this application.

## FIELD OF THE INVENTION

10 **[0002]** This invention is directed to: (a) compounds and salts thereof that, *inter alia*, are useful as hepatitis C virus (HCV) inhibitors; (b) intermediates useful for the preparation of such compounds and salts; (c) compositions comprising such compounds and salts; (d) methods for preparing such intermediates, compounds, salts, and compositions; (e) methods of use of such compounds, salts, and compositions; and (f) kits comprising such compounds, salts, and compositions.

## BACKGROUND OF THE INVENTION

20 **[0003]** Hepatitis C is a blood-borne, infectious, viral disease that is caused by a hepatotropic virus called HCV. At least six different HCV genotypes (with several subtypes within each genotype) are known to date. In North America, HCV genotype 1a predominates, followed by HCV genotypes 1b, 2a, 2b, and 3a. In the United States, HCV genotypes 1, 2, and 3 are the most common, with about 80% of the hepatitis C patients having HCV genotype 1. In Europe, HCV genotype 1b is predominant, followed by HCV genotypes 2a, 2b, 2c, and 3a. HCV genotypes 4 and 5 are found almost exclusively in Africa. As discussed below, the patient's HCV genotype is clinically important in determining the patient's potential response to therapy and the required duration of such therapy.

25 **[0004]** An HCV infection can cause liver inflammation (hepatitis) that is often asymptomatic, but ensuing chronic hepatitis can result in cirrhosis of the liver (fibrotic scarring of the liver), liver cancer, and/or liver failure. The World Health Organization estimates that about 170 million persons worldwide are chronically infected with HCV, and from about three to about four million persons are newly infected globally each year. According to the Centers for Disease Control and Prevention, about four million people in the United States are infected with HCV. Co-infection with the human immunodeficiency virus (HIV) is common, and rates of HCV infection among HIV positive populations are higher.

30 **[0005]** There is a small chance of clearing the virus spontaneously, but the majority of patients with chronic hepatitis C will not clear it without treatment. Indications for treatment typically include proven HCV infection and persistent abnormal liver function tests. There are two treatment regimens that are primarily used to treat hepatitis C: monotherapy (using an interferon agent - either a "conventional" or longer-acting pegylated interferon) and combination therapy (using an interferon agent and ribavirin). Interferon, which is injected into the bloodstream, works by bolstering the immune response to HCV; and ribavirin, which is taken orally, is believed to work by preventing HCV replication. Taken alone, ribavirin does not effectively suppress HCV levels, but an interferon/ribavirin combination is more effective than interferon alone. Typically, hepatitis C is treated with a combination of pegylated interferon alpha and ribavirin for a period of 24 or 48 weeks, depending on the HCV genotype.

35 **[0006]** The goal of treatment is sustained viral response - meaning that HCV is not measurable in the blood after therapy is completed. Following treatment with a combination of pegylated interferon alpha and ribavirin, sustained cure rates (sustained viral response) of about 75% or better occur in people with HCV genotypes 2 and 3 in 24 weeks of treatment, about 50% in those with HCV genotype 1 with 48 weeks of treatment, and about 65% in those with HCV genotype 4 in 48 weeks of treatment.


40 **[0007]** Treatment may be physically demanding, particularly for those with prior history of drug or alcohol abuse, because both interferon and ribavirin have numerous side effects. Common interferon-associated side effects include flu-like symptoms, extreme fatigue, nausea, loss of appetite, thyroid problems, high blood sugar, hair loss, and skin reactions at the injection site. Possible serious interferon-associated side effects include psychoses (e.g., suicidal behavior), heart problems (e.g., heart attack, low blood pressure), other internal organ damage, blood problems (e.g., blood counts falling dangerously low), and new or worsening autoimmune disease (e.g., rheumatoid arthritis). Ribavirin-associated side effects include anemia, fatigue, irritability, skin rash, nasal stuffiness, sinusitis, and cough. Ribavirin can also cause birth defects, so pregnancy in female patients and female partners of male patients must be avoided during treatment and for six months afterward.

45 **[0008]** Some patients do not complete treatment because of the serious side effects discussed above; other patients (non-responders) continue to have measurable HCV levels despite treatment; and yet other patients (relapsers) "clear" the virus during therapy, but the virus returns sometime after completion of the treatment regimen. Thus, there continues to be a need for alternative compounds, compositions, and methods of treatment (used either in combination with or in

lieu of an interferon agent and/or ribavirin) to alleviate the symptoms of hepatitis C, thereby providing partial or complete relief. This invention provides compounds (including salts thereof), compositions, and methods of treatment that generally address such a need.

5 SUMMARY OF THE INVENTION

[0009] This invention is directed to compounds that correspond in structure to formula I:



20 [0010] In formula I:

$\text{---}^*\text{---}$  is selected from the group consisting of single carbon-carbon bond and double carbon-carbon bond;

$R^1$  is selected from the group consisting of hydrogen and methyl,

25  $R^2$  is selected from the group consisting of hydrogen, halo, hydroxy, methyl, cyclopropyl, and cyclobutyl;

$R^3$  is selected from the group consisting of hydrogen, halo, oxo, and methyl;

$R^4$  is selected from the group consisting of halo, alkyl, alkenyl, alkynyl, nitro, cyano, azido, alkyloxy, alkenyloxy, alkynyloxy, amino, aminocarbonyl, aminosulfonyl, alkylsulfonyl, carbocyclyl, and heterocyclyl, wherein:

30 (a) the amino, aminocarbonyl, and aminosulfonyl optionally are substituted with:

(1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, and alkylsulfonyl, or

(2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl, and

35 (b) the alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, alkynyloxy, and alkylsulfonyl, optionally are substituted with one or more substituents independently selected from the group consisting of halo, oxo, nitro, cyano, azido, hydroxy, amino, alkyloxy, trimethylsilyl, carbocyclyl, and heterocyclyl, wherein:

the amino optionally is substituted with:

40 (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylcarbonyl, alkylsulfonyl, alkyloxy, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl, or

(2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl, and

45 (c) the carbocyclyl and heterocyclyl optionally are substituted with up to three substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, halo, oxo, nitro, cyano, azido, hydroxy, amino, alkyloxy, trimethylsilyl, carbocyclyl, and heterocyclyl, wherein:

the amino optionally is substituted with:

50 (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylcarbonyl, alkylsulfonyl, alkyloxy, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl, or

(2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl;

55  $R^5$  is selected from the group consisting of hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, alkynyloxy, alkylsulfonyloxy, carbocyclylsulfonyloxy, haloalkylsulfonyloxy, and halo;

L is selected from the group consisting of  $C(R^A)=C(R^B)$ , ethylene, and cyclopropyl-1,2-ene;

## EP 2 203 430 B1

**R<sup>A</sup>** and **R<sup>B</sup>** are independently selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkyloxy, C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, and halo, wherein:

5 the C<sub>1</sub>-C<sub>6</sub>-alkyl optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, halo, hydroxy, nitro, oxo, amino, cyano, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy, carbocyclyl, and heterocyclyl;

**R<sup>G</sup>** is selected from the group consisting of C<sub>5</sub>-C<sub>6</sub>-carbocyclyl, 5-6-membered heterocyclyl, fused 2-ring carbocyclyl, and fused 2-ring heterocyclyl, wherein each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of **R<sup>E</sup>**, **R<sup>F</sup>**, **R<sup>G</sup>**, **R<sup>H</sup>**, **R<sup>I</sup>**, **R<sup>J</sup>**, and **R<sup>K</sup>**;  
10 each **R<sup>E</sup>** is independently selected from the group consisting of halo, nitro, hydroxy, oxo, carboxy, cyano, amino, imino, azido, and aldehydo, wherein:

15 the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl;

each **R<sup>F</sup>** is independently selected from the group consisting of alkyl, alkenyl, and alkynyl, wherein:

20 each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, imino, nitro, azido, oxo, aminosulfonyl, alkylsulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:  
the amino, imino, aminosulfonyl, aminocarbonyl, carbocyclyl, and heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylsulfonyl,  
25 alkenylsulfonyl, alkynylsulfonyl, alkylsulfonylamino, hydroxy, and alkyloxy,  
wherein:

30 amino portion of the alkylsulfonylamino optionally is substituted with a substituent selected from the group consisting of alkyl, alkenyl, and alkynyl;

each **R<sup>G</sup>** is independently selected from the group consisting of carbocyclyl and heterocyclyl, wherein:

35 each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

40 the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl;

each **R<sup>H</sup>** is independently selected from the group consisting of alkyloxy, alkenyloxy, alkynyloxy, alkylsulfonyloxy, alkenylsulfonyloxy, and alkynylsulfonyloxy, wherein:

45 each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

50 the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl;

55 each **R<sup>I</sup>** is independently selected from the group consisting of alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl, aminocarbonyl, alkyloxycarbonyl, carbocyclylcarbonyl, and heterocyclylcarbonyl, wherein:

(a) the alkylcarbonyl, alkenylcarbonyl, and alkynylcarbonyl optionally are substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo,

aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, and

(b) the aminocarbonyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkyloxyalkyl, carbocyclyl, heterocyclyl, alkylsulfonyl, and alkylsulfonylamino, wherein:

the carbocyclyl and heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of halo, alkyl, and oxo;

each  $R^J$  is independently selected from the group consisting of carbocyclylsulfonylamino, heterocyclylsulfonylamino, alkylcarbonylamino, alkenylcarbonylamino, alkynylcarbonylamino, alkyloxycarbonylamino, alkenyloxycarbonylamino, alkynyloxycarbonylamino, alkylsulfonylamino, alkenylsulfonylamino, alkynylsulfonylamino, aminocarbonylamino, alkyloxycarbonylaminoimino, alkylsulfonylaminoimino, alkenylsulfonylaminoimino, and alkynylsulfonylaminoimino, wherein:

(a) the amino portion of such substituents optionally is substituted with a substituent independently selected from the group consisting of carbocyclylalkyl, heterocyclylalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkenyl, alkynyl, alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl, alkyloxycarbonyl, alkyloxyalkyloxycarbonyl, alkylcarbonyloxyalkyl, and alkylsulfonyl, wherein:

(1) the carbocyclyl portion of the carbocyclylalkyl and the heterocyclyl portion of the heterocyclylalkyl optionally are substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, alkyloxy, alkenyloxy, alkynyloxy, halo, nitro, cyano, azido, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl,

(b) the alkyl, alkenyl, and alkynyl portion of such substituents optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxycarbonyl, alkylcarbonyloxy, hydroxy, alkyloxy, carbocyclyl, heterocyclyl, and cyano, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, and alkynyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy;

(c) the carbocyclyl and heterocyclyl portions of such substituents optionally are substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, alkyloxy, alkenyloxy, alkynyloxy, halo, nitro, cyano, azido, and amino, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl; and

each  $R^K$  is independently selected from the group consisting of aminosulfonyl, alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl, wherein:

(a) the alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl optionally are substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl; and

(b) the aminosulfonyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl.

[0011] This invention also is directed to the salts (including pharmaceutically acceptable salts) of the compounds of the invention.

[0012] This invention also is directed to compositions (including pharmaceutical compositions) that comprise one or more compounds and/or salts of the invention, and, optionally, one or more additional therapeutic agents.

[0013] This invention also is directed to kits that comprise one or more compounds and/or salts of the invention, and, optionally, one or more additional therapeutic agents.

[0014] This invention also is directed to the compounds, salts, compositions and/or kits of the invention for use in, for example, inhibiting replication of an RNA virus (including HCV), treating a disease treatable by inhibiting HCV ribonucleic acid (RNA) polymerase (including hepatitis C).

[0015] This invention also is directed to a use of one or more compounds and/or salts of the invention to prepare a medicament. The medicament optionally can comprise one or more additional therapeutic agents. In some embodiments, the medicament is useful for treating hepatitis C.

[0016] Further benefits of Applicants' invention will be apparent to one skilled in the art from reading this patent application.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0017] Figure 1 shows an illustrative PXRD pattern for the disodium salt nonahydrate of compound **IB-L1-1.1**.

[0018] Figure 2 shows an illustrative PXRD pattern for the disodium salt tetrahydrate of compound **IB-L1-1.1**.

[0019] Figure 3 shows an illustrative TGA profile of the disodium salt tetrahydrate of compound **IB-L1-1.1**.

[0020] Figure 4 shows an illustrative PXRD pattern for the dipotassium salt tetrahydrate of compound **IB-L1-1.1**.

[0021] Figure 5 shows an illustrative PXRD pattern for the monopotassium salt trihydrate of compound **IB-L1-1.1**.

[0022] Figure 6 shows an illustrative PXRD pattern for the monopotassium salt dihydrate of compound **IB-L1-1.1**.

[0023] Figure 7 shows an illustrative TGA profile of the monopotassium salt dihydrate of compound **IB-L1-1.1**.

[0024] Figure 8 shows an illustrative PXRD pattern for the 1/7 potassium salt of compound **IB-L1-1.1**.

[0025] Figure 9 shows an illustrative PXRD pattern for the monodiethylamine salt tetrahydrate of compound **IB-L1-1.1**.

[0026] Figure 10 shows an illustrative TGA profile of the monodiethylamine salt tetrahydrate of compound **IB-L1-1.1**.

[0027] Figure 11 shows an illustrative PXRD pattern for the pattern A polymorph of compound **IB-L1-1.1**.

[0028] Figure 12 shows an illustrative DSC profile of the pattern A polymorph of compound **IB-L1-1.1**.

[0029] Figure 13 shows an illustrative PXRD pattern for the pattern B polymorph of compound **IB-L1-1.1**.

[0030] Figure 14 shows an illustrative PXRD pattern for the pattern C polymorph of compound **IB-L1-1.1**.

[0031] Figure 15 shows an illustrative PXRD pattern for the pattern D polymorph of compound **IB-L1-1.1**.

[0032] Figure 16 shows an illustrative PXRD pattern for the pattern A hydrate of compound **IB-L1-1.1**.

[0033] Figure 17 shows an illustrative TGA profile of the pattern A hydrate of compound **IB-L1-1.1**.

[0034] Figure 18 shows an illustrative PXRD pattern for the pattern B hydrate of compound **IB-L1-1.1**.

[0035] Figure 19 shows an illustrative TGA profile of the pattern B hydrate of compound **IB-L1-1.1**.

[0036] Figure 20 shows an illustrative PXRD pattern for the pattern C hydrate of compound **IB-L1-1.1**.

[0037] Figure 21 shows an illustrative TGA profile of the pattern C hydrate of compound **IB-L1-1.1**.

[0038] Figure 22 shows an illustrative PXRD pattern for the pattern D hydrate of compound **IB-L1-1.1**.

[0039] Figure 23 shows an illustrative PXRD pattern for the pattern E hydrate of compound **IB-L1-1.1**.

#### DETAILED DESCRIPTION OF THE INVENTION

[0040] This detailed description is intended only to acquaint others skilled in the art with Applicants' invention, its principles, and its practical application so that others skilled in the art may adapt and apply the invention in its numerous forms, as they may be best suited to the requirements of a particular use. This description and its specific examples are intended for purposes of illustration only. This invention, therefore, is not limited to the embodiments described in this patent application, and may be variously modified.

##### A. Definitions.

[0041] The term "alkyl" (alone or in combination with another term(s)) means a straight-or branched-chain saturated hydrocarbyl substituent typically containing from 1 to about 20 carbon atoms, more typically from 1 to about 8 carbon atoms, and even more typically from 1 to about 6 carbon atoms. Examples of such substituents include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, and hexyl. As in this definition, throughout this detailed description Applicants have provided illustrative examples. The provision of such illustrative examples should not be interpreted as if the provided illustrative examples are the only options available to one skilled in the art.

[0042] The term "alkenyl" (alone or in combination with another term(s)) means a straight- or branched-chain hydro-

carbonyl substituent containing one or more double bonds and typically from 2 to about 20 carbon atoms, more typically from about 2 to about 8 carbon atoms, and even more typically from about 2 to about 6 carbon atoms. Examples of such substituents include ethenyl (vinyl), 2-propenyl, 3-propenyl, 1,4-pentadienyl, 1,4-butadienyl, 1-butenyl, 2-butenyl, and 3-butenyl.

**[0043]** The term "alkynyl" (alone or in combination with another term(s)) means a straight- or branched-chain hydrocarbonyl substituent containing one or more triple bonds and typically from 2 to about 20 carbon atoms, more typically from about 2 to about 8 carbon atoms, and even more typically from about 2 to about 6 carbon atoms. Examples of such substituents include ethynyl, 2-propynyl, 3-propynyl, 2-butylnyl, and 3-butylnyl.

**[0044]** The term "carbocyclyl" (alone or in combination with another term(s)) means a saturated cyclic (*i.e.*, "cycloalkyl"), partially saturated cyclic (*i.e.*, "cycloalkenyl"), or completely unsaturated (*i.e.*, "aryl") hydrocarbonyl substituent containing from 3 to 14 carbon ring atoms ("ring atoms" are the atoms bound together to form the ring or rings of a cyclic substituent). A carbocyclyl may be a single ring, which typically contains from 3 to 6 ring atoms. Examples of such single-ring carbocyclyls include cyclopropyl (cyclopropanyl), cyclobutyl (cyclobutanyl), cyclopentyl (cyclopentanyl), cyclopentenyl, cyclopentadienyl, cyclohexyl (cyclohexanyl), cyclohexenyl, cyclohexadienyl, and phenyl. A carbocyclyl alternatively may be 2 or 3 rings fused together, such as naphthalenyl, tetrahydronaphthalenyl (tetralinyl), indenyl, indanyl (dihydroindenyl), anthracenyl, phenanthrenyl, and decalinyl.

**[0045]** The term "cycloalkyl" (alone or in combination with another term(s)) means a saturated cyclic hydrocarbonyl substituent containing from 3 to 14 carbon ring atoms. A cycloalkyl may be a single carbon ring, which typically contains from 3 to 6 carbon ring atoms. Examples of single-ring cycloalkyls include cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. A cycloalkyl alternatively may be 2 or 3 carbon rings fused together, such as, decalinyl.

**[0046]** The term "aryl" (alone or in combination with another term(s)) means an aromatic carbocyclyl containing from 6 to 14 carbon ring atoms. Examples of aryls include phenyl, naphthalenyl, and indenyl.

**[0047]** In some instances, the number of carbon atoms in a hydrocarbonyl substituent (*e.g.*, alkyl, alkenyl, alkynyl, or cycloalkyl) is indicated by the prefix "C<sub>x</sub>-C<sub>y</sub>", wherein x is the minimum and y is the maximum number of carbon atoms in the substituent. Thus, for example, "C<sub>1</sub>-C<sub>6</sub>-alkyl" refers to an alkyl substituent containing from 1 to 6 carbon atoms. Illustrating further, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl means a saturated hydrocarbonyl ring containing from 3 to 6 carbon ring atoms.

**[0048]** The term "hydrogen" (alone or in combination with another term(s)) means a hydrogen radical, and may be depicted as -H.

**[0049]** The term "hydroxy" (alone or in combination with another term(s)) means -OH.

**[0050]** The term "nitro" (alone or in combination with another term(s)) means -NO<sub>2</sub>.

**[0051]** The term "cyano" (alone or in combination with another term(s)) means -CN, which also may be depicted as -C≡N.

**[0052]** The term "keto" (alone or in combination with another term(s)) means an oxo radical, and may be depicted as =O.

**[0053]** The term "carboxy" (alone or in combination with another term(s)) means -C(O)-OH.

**[0054]** The term "amino" (alone or in combination with another term(s)) means -NH<sub>2</sub>.

**[0055]** The term "imino" (alone or in combination with another term(s)) means =NH.

**[0056]** The term "aminoimino" (alone or in combination with another term(s)) means =NNH<sub>2</sub>.

**[0057]** The term "halogen" or "halo" (alone or in combination with another term(s)) means a fluorine radical (which may be depicted as -F), chlorine radical (which may be depicted as -Cl), bromine radical (which may be depicted as -Br), or iodine radical (which may be depicted as -I).

**[0058]** A substituent is "substitutable" if it comprises at least one carbon or nitrogen atom that is bonded to one or more hydrogen atoms. Thus, for example, hydrogen, halogen, and cyano do not fall within this definition. In addition, a sulfur atom in a heterocyclyl containing such atom is substitutable with one or two oxo substituents.

**[0059]** If a substituent is described as being "substituted", a non-hydrogen radical is in the place of hydrogen radical on a carbon or nitrogen of the substituent. Thus, for example, a substituted alkyl substituent is an alkyl substituent in which at least one non-hydrogen radical is in the place of a hydrogen radical on the alkyl substituent. To illustrate, monofluoroalkyl is alkyl substituted with a fluoro radical, and difluoroalkyl is alkyl substituted with two fluoro radicals. It should be recognized that if there are more than one substitution on a substituent, each non-hydrogen radical may be identical or different (unless otherwise stated).

**[0060]** If a substituent is described as being "optionally substituted", the substituent may be either (1) not substituted or (2) substituted. If a substituent is described as being optionally substituted with up to a particular number of non-hydrogen radicals, that substituent may be either (1) not substituted; or (2) substituted by up to that particular number of non-hydrogen radicals or by up to the maximum number of substitutable positions on the substituent, whichever is less. Thus, for example, if a substituent is described as a heteroaryl optionally substituted with up to 3 non-hydrogen radicals, then any heteroaryl with less than 3 substitutable positions would be optionally substituted by up to only as many non-hydrogen radicals as the heteroaryl has substitutable positions. To illustrate, tetrazolyl (which has only one substitutable position) would be optionally substituted with up to one non-hydrogen radical. To illustrate further, if an amino nitrogen is described as being optionally substituted with up to 2 non-hydrogen radicals, then a primary amino

nitrogen will be optionally substituted with up to 2 non-hydrogen radicals, whereas a secondary amino nitrogen will be optionally substituted with up to only 1 non-hydrogen radical.

**[0061]** This patent application uses the terms "substituent" and "radical" interchangeably.

**[0062]** The prefix "halo" indicates that the substituent to which the prefix is attached is substituted with one or more independently selected halogen radicals. For example, haloalkyl means an alkyl substituent in which at least one hydrogen radical is replaced with a halogen radical. Examples of haloalkyls include chloromethyl, 1-bromoethyl, fluoromethyl, difluoromethyl, trifluoromethyl, and 1,1,1-trifluoroethyl. It should be recognized that if a substituent is substituted by more than one halogen radical, those halogen radicals may be identical or different (unless otherwise stated).

**[0063]** The prefix "perhalo" indicates that every hydrogen radical on the substituent to which the prefix is attached is replaced with independently selected halogen radicals, i.e., each hydrogen radical on the substituent is replaced with a halogen radical. If all the halogen radicals are identical, the prefix typically will identify the halogen radical. Thus, for example, the term "perfluoro" means that every hydrogen radical on the substituent to which the prefix is attached is substituted with a fluorine radical. To illustrate, the term "perfluoroalkyl" means an alkyl substituent wherein a fluorine radical is in the place of each hydrogen radical.

**[0064]** The term "carbonyl" (alone or in combination with another term(s)) means -C(O)-.

**[0065]** The term "aminocarbonyl" (alone or in combination with another term(s)) means -C(O)-NH<sub>2</sub>.

**[0066]** The term "oxy" (alone or in combination with another term(s)) means an ether substituent, and may be depicted as -O-.

**[0067]** The term "alkoxy" (alone or in combination with another term(s)) means an alkylether substituent, i.e., -O-alkyl. Examples of such a substituent include methoxy (-O-CH<sub>3</sub>), ethoxy, n-propoxy, isopropoxy, n-butoxy, iso-butoxy, sec-butoxy, and tert-butoxy.

**[0068]** The term "alkylcarbonyl" (alone or in combination with another term(s)) means -C(O)-alkyl.

**[0069]** The term "aminoalkylcarbonyl" (alone or in combination with another term(s)) means -C(O)-alkyl-NH<sub>2</sub>.

**[0070]** The term "alkoxycarbonyl" (alone or in combination with another term(s)) means -C(O)-O-alkyl.

**[0071]** The term "carbocyclylcarbonyl" (alone or in combination with another term(s)) means -C(O)-carbocyclyl.

**[0072]** Similarly, the term "heterocyclylcarbonyl" (alone or in combination with another term(s)) means -C(O)-heterocyclyl.

**[0073]** The term "carbocyclylalkylcarbonyl" (alone or in combination with another term(s)) means -C(O)-alkyl-carbocyclyl.

**[0074]** Similarly, the term "heterocyclylalkylcarbonyl" (alone or in combination with another term(s)) means -C(O)-alkyl-heterocyclyl.

**[0075]** The term "carbocyclyloxycarbonyl" (alone or in combination with another term(s)) means -C(O)-O-carbocyclyl.

**[0076]** The term "carbocyclylalkoxycarbonyl" (alone or in combination with another term(s)) means -C(O)-O-alkyl-carbocyclyl.

**[0077]** The term "thio" or "thia" (alone or in combination with another term(s)) means a thiaether substituent, i.e., an ether substituent wherein a divalent sulfur atom is in the place of the ether oxygen atom. Such a substituent may be depicted as -S-. This, for example, "alkyl-thio-alkyl" means alkyl-S-alkyl (alkyl-sulfanyl-alkyl).

**[0078]** The term "thiol" or "sulfhydryl" (alone or in combination with another term(s)) means a sulfhydryl substituent, and may be depicted as -SH.

**[0079]** The term "(thiocarbonyl)" (alone or in combination with another term(s)) means a carbonyl wherein the oxygen atom has been replaced with a sulfur. Such a substituent may be depicted as -C(S)-.

**[0080]** The term "sulfonyl" (alone or in combination with another term(s)) means -S(O)<sub>2</sub>-.

**[0081]** The term "aminosulfonyl" (alone or in combination with another term(s)) means -S(O)<sub>2</sub>-NH<sub>2</sub>-.

**[0082]** The term "sulfinyl" or "sulfoxido" (alone or in combination with another term(s)) means -S(O)-.

**[0083]** The term "heterocyclyl" (alone or in combination with another term(s)) means a saturated (i.e., "heterocycloalkyl"), partially saturated (i.e., "heterocycloalkenyl"), or completely unsaturated (i.e., "heteroaryl") ring structure containing a total of 3 to 14 ring atoms. At least one of the ring atoms is a heteroatom (i.e., oxygen, nitrogen, or sulfur), with the remaining ring atoms being independently selected from the group consisting of carbon, oxygen, nitrogen, and sulfur.

**[0084]** A heterocyclyl may be a single ring, which typically contains from 3 to 7 ring atoms, more typically from 3 to 6 ring atoms, and even more typically 5 to 6 ring atoms. Examples of single-ring heterocyclyls include furanyl, dihydrofuranyl, tetrahydrofuranyl, thiophenyl (thiofuranyl), dihydrothiophenyl, tetrahydrothiophenyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolyl, imidazoliny, imidazolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, triazolyl, tetrazolyl, oxazolyl, oxazolidinyl, isoxazolidinyl, isoxazolyl, thiazolyl, isothiazolyl, thiazolinyl, isothiazolinyl, thiazolidinyl, isothiazolidinyl, thiodiazolyl, oxadiazolyl (including 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl (furazanyl), or 1,3,4-oxadiazolyl), oxatriazolyl (including 1,2,3,4-oxatriazolyl or 1,2,3,5-oxatriazolyl), dioxazolyl (including 1,2,3-dioxazolyl, 1,2,4-dioxazolyl, 1,3,2-dioxazolyl, or 1,3,4-dioxazolyl), oxathiazolyl, oxathioly, oxathiolanyl, pyranyl, dihydropyranyl, thiopyranyl, tetrahydrothiopyranyl, pyridinyl (azinyl), piperidinyl, diazinyl (including pyridazinyl (1,2-diazinyl), pyrimidinyl (1,3-diazinyl), or pyrazinyl (1,4-diazinyl)), piperazinyl, triazinyl (including 1,3,5-triazinyl, 1,2,4-triazinyl, and 1,2,3-triazinyl), oxazinyl (including 1,2-

oxazinyl, 1,3-oxazinyl, or 1,4-oxazinyl)), oxathiazinyl (including 1,2,3-oxathiazinyl, 1,2,4-oxathiazinyl, 1,2,5-oxathiazinyl, or 1,2,6-oxathiazinyl)), oxadiazinyl (including 1,2,3-oxadiazinyl, 1,2,4-oxadiazinyl, 1,4,2-oxadiazinyl, or 1,3,5-oxadiazinyl)), morpholinyl, azepinyl, oxepinyl, thiepinyl, and diazepinyl.

**[0085]** A heterocyclyl alternatively may be 2 or 3 rings fused together, such as, for example, indolizinyl, pyranopyrrolyl, 4H-quinolizinyl, purinyl, naphthyridinyl, pyridopyridinyl (including pyrido[3,4-b]-pyridinyl, pyrido[3,2-b]-pyridinyl, or pyrido[4,3-b]-pyridinyl), and pteridinyl. Other examples of fused-ring heterocyclyls include benzo-fused heterocyclyls, such as indolyl, isoindolyl (isobenzazolyl, pseudoisoindolyl), indoleninyl (pseudoindolyl), isoindazolyl (benzpyrazolyl), benzazinyl (including quinolinyl (1-benzazinyl) or isoquinolinyl (2-benzazinyl)), phthalazinyl, quinoxalinyl, quinazoliny, benzodiazinyl (including cinnolinyl (1,2-benzodiazinyl) or quinazoliny (1,3-benzodiazinyl)), benzopyranyl (including chromanyl or isochromanyl), benzoxazinyl (including 1,3,2-benzoxazinyl, 1,4,2-benzoxazinyl, 2,3,1-benzoxazinyl, or 3,1,4-benzoxazinyl), and benzisoxazinyl (including 1,2-benzisoxazinyl or 1,4-benzisoxazinyl).

**[0086]** The term "2-fused ring" heterocyclyl (alone or in combination with another term(s)) means a saturated, partially saturated, or aryl heterocyclyl containing 2 fused rings. Examples of 2-fused-ring heterocyclyls include indolizinyl, quinolizinyl, purinyl, naphthyridinyl, pteridinyl, indolyl, isoindolyl, indoleninyl, isoindazolyl, phthalazinyl, quinoxalinyl, quinazoliny, benzodiazinyl, benzopyranyl, benzothiopyranyl, benzoxazolyl, anthranilyl, benzodioxolyl, benzodioxanyl, benzoxadiazolyl, benzofuranyl, isobenzofuranyl, benzothiazolyl, benzothiadiazolyl, benzimidazolyl, benzotriazolyl, benzoxazinyl, and tetrahydroisoquinolinyl.

**[0087]** The term "heteroaryl" (alone or in combination with another term(s)) means an aromatic heterocyclyl containing from 5 to 14 ring atoms. A heteroaryl may be a single ring or 2 or 3 fused rings. Examples of heteroaryl substituents include 6-membered ring substituents such as pyridyl, pyrazyl, pyrimidinyl, pyridazinyl, and 1,3,5-, 1,2,4- or 1,2,3-triazinyl; 5-membered ring substituents such as imidazolyl, furanyl, thiophenyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, 1,2,3-, 1,2,4-, 1,2,5-, or 1,3,4-oxadiazolyl and isothiazolyl; 6/5-membered fused ring substituents such as benzothiofuranyl, benzisoxazolyl, benzoxazolyl, purinyl, and anthranilyl; and 6/6-membered fused rings such as benzopyranyl, quinolinyl, isoquinolinyl, cinnolinyl, quinazoliny, and benzoxazinyl.

**[0088]** A prefix attached to a multi-component substituent only applies to the first component. To illustrate, the term "alkylcycloalkyl" contains two components: alkyl and cycloalkyl. Thus, the C<sub>1</sub>-C<sub>6</sub>-prefix on C<sub>1</sub>-C<sub>6</sub>-alkylcycloalkyl means that the alkyl component of the alkylcycloalkyl contains from 1 to 6 carbon atoms; the C<sub>1</sub>-C<sub>6</sub>-prefix does not describe the cycloalkyl component. To illustrate further, the prefix "halo" on haloalkoxyalkyl indicates that only the alkoxy component of the alkoxyalkyl substituent is substituted with one or more halogen radicals. If halogen substitution may alternatively or additionally occur on the alkyl component, the substituent would instead be described as "halogen-substituted alkoxyalkyl" rather than "haloalkoxyalkyl." And finally, if the halogen substitution may only occur on the alkyl component, the substituent would instead be described as "alkoxyhaloalkyl."

**[0089]** If substituents are described as being "independently selected" from a group, each substituent is selected independent of the other. Each substituent therefore may be identical to or different from the other substituent(s).

**[0090]** When words are used to describe a substituent, the rightmost-described component of the substituent is the component that has the free valence.

**[0091]** When a chemical formula is used to describe a substituent, the dash on the left side of the formula indicates the portion of the substituent that has the free valence.

**[0092]** When a chemical formula is used to describe a linking element between two other elements of a depicted chemical structure, the leftmost dash of the substituent indicates the portion of the substituent that is bound to the left element in the depicted structure. The rightmost dash, on the other hand, indicates the portion of the substituent that is bound to the right element in the depicted structure. To illustrate, if the depicted chemical structure is X-L-Y and L is described as -C(O)-N(H)-, then the chemical would be X-C(O)-N(H)-Y.

**[0093]** With reference to the use of the words "comprise" or "comprises" or "comprising" in this patent application (including the claims), Applicants note that unless the context requires otherwise, those words are used on the basis and clear understanding that they are to be interpreted inclusively, rather than exclusively, and that Applicants intend each of those words to be so interpreted in construing this patent application, including the claims below.

**[0094]** ChemDraw software has been used to generate the compound names in this patent application.

**[0095]** The term "amorphous" as applied to a compound refers to a solid-state in which the compound molecules are present in a disordered arrangement and do not form a distinguishable crystal lattice or unit cell. When subjected to X-ray powder diffraction, an amorphous compound does not produce any characteristic crystalline peaks.

**[0096]** The term "crystalline form" as applied to a compound refers to a solid-state in which the compound molecules are arranged to form a distinguishable crystal lattice (i) comprising distinguishable unit cells, and (ii) yielding diffraction pattern peaks when subjected to X-ray radiation.

**[0097]** The term "purity", unless otherwise qualified, means the chemical purity of a compound according to conventional HPLC assay.

**[0098]** The term "phase purity" means the solid-state purity of a compound with regard to a particular crystalline or amorphous form of the compound as determined by X-ray powder diffraction analytical methods.

[0099] The term "phase pure" refers to purity with respect to other solid-state forms of the compound, and does not necessarily imply a high degree of chemical purity with respect to other compounds.


[0100] The term "PXRD" means X-ray powder diffraction.

[0101] The term "TGA" means thermogravimetric analysis.

[0102] The term "DSC" means differential scanning calorimetry.


**B. Compounds.**

[0103] This invention is directed, in part, to compounds that are phenyl-uracil derivatives that correspond in structure to formula I:




[0104] In these compounds,  $\overset{*}{\text{---}}$  is selected from the group consisting of single carbon-carbon bond and double carbon-carbon bond.

[0105] In some embodiments,  $\overset{*}{\text{---}}$  is a single carbon-carbon bond. In these embodiments, the compounds of formula I correspond in structure to the following formula (i.e., formula IA):



[0106] In other embodiments,  $\overset{*}{\text{---}}$  is a double carbon-carbon bond. In these embodiments, the compounds of formula I correspond in structure to the following formula (i.e., formula IB):



**B1. Substituent R<sup>1</sup>.**

- [0107] R<sup>1</sup> is selected from the group consisting of hydrogen, and methyl,  
 [0108] In some embodiments, R<sup>1</sup> is hydrogen.  
 5 [0109] In some embodiments, R<sup>1</sup> is methyl.  
 [0110] In some embodiments, R<sup>1</sup> is selected from the group consisting of hydrogen and methyl.

**B2. Substituent R<sup>2</sup>.**

- 10 [0111] R<sup>2</sup> is selected from the group consisting of hydrogen, halo, hydroxy, methyl, cyclopropyl, and cyclobutyl.  
 [0112] In some embodiments, R<sup>2</sup> is hydrogen.  
 [0113] In some embodiments, R<sup>1</sup> is halo. In some such embodiments, R<sup>2</sup> is selected from the group consisting of  
 fluoro and chloro. In other such embodiments, R<sup>2</sup> is fluoro. In yet other such embodiments, R<sup>2</sup> is chloro. In yet other  
 such embodiments, R<sup>2</sup> is bromo. In further such embodiments, R<sup>2</sup> is iodo.  
 15 [0114] In some embodiments, R<sup>2</sup> is hydroxy.  
 [0115] In some embodiments, R<sup>2</sup> is methyl.  
 [0116] In some embodiments, R<sup>2</sup> is cyclopropyl.  
 [0117] In some embodiments, R<sup>2</sup> is cyclobutyl.  
 [0118] In some embodiments, R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, hydroxy, and halo. In  
 20 some such embodiments, R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, hydroxy, fluoro, and chloro. In  
 other such embodiments, R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, hydroxy, and fluoro. In yet other  
 such embodiments, R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, hydroxy, and chloro. In yet other such  
 embodiments, R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, hydroxy, and bromo. In further such em-  
 bodiments, R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, hydroxy, and iodo.  
 25 [0119] In some embodiments, R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, and halo. In some such  
 embodiments, R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, fluoro, and chloro. In other such embodi-  
 ments, R<sup>2</sup> is selected from the group consisting of hydrogen, methyl, and fluoro. In yet other such embodiments, R<sup>2</sup>  
 is selected from the group consisting of hydrogen, methyl, and chloro. In yet other such embodiments, R<sup>2</sup> is selected from  
 the group consisting of hydrogen, methyl, and bromo. In further such embodiments, R<sup>2</sup> is selected from the group  
 30 consisting of hydrogen, methyl, and iodo.  
 [0120] In some embodiments, R<sup>2</sup> is selected from the group consisting of hydrogen and halo. In some such embod-  
 iments, R<sup>2</sup> is selected from the group consisting of hydrogen, fluoro, and chloro. In other such embodiments, R<sup>2</sup>  
 is selected from the group consisting of hydrogen and fluoro. In yet other such embodiments, R<sup>2</sup> is selected from the group  
 consisting of hydrogen and chloro. In yet other such embodiments, R<sup>2</sup> is selected from the group consisting of hydrogen  
 35 and bromo. In further such embodiments, R<sup>2</sup> is selected from the group consisting of hydrogen and iodo.

**B3. Substituent R<sup>3</sup>.**

- 40 [0121] R<sup>3</sup> is selected from the group consisting of hydrogen, halo, oxo, and methyl. In some such embodiments, R<sup>3</sup>  
 is selected from the group consisting of hydrogen, fluoro, oxo, and methyl. In other such embodiments, R<sup>3</sup> is selected  
 from the group consisting of hydrogen, chloro, oxo, and methyl. In yet other such embodiments, R<sup>3</sup> is selected from the  
 group consisting of hydrogen, bromo, oxo, and methyl. In yet other such embodiments, R<sup>3</sup> is selected from the group  
 consisting of hydrogen, iodo, oxo, and methyl.  
 [0122] In some embodiments, R<sup>3</sup> is selected from the group consisting of hydrogen, halo, and oxo. In some such  
 45 embodiments, R<sup>3</sup> is selected from the group consisting of hydrogen, fluoro, and oxo. In other such embodiments, R<sup>3</sup>  
 is selected from the group consisting of hydrogen, chloro, and oxo. In yet other such embodiments, R<sup>3</sup> is selected from  
 the group consisting of hydrogen, bromo, and oxo. In yet other such embodiments, R<sup>3</sup> is selected from the group  
 consisting of hydrogen, iodo, and oxo.  
 [0123] In some embodiments, R<sup>3</sup> is selected from the group consisting of hydrogen and methyl.  
 50 [0124] In some embodiments, R<sup>3</sup> is hydrogen.  
 [0125] In some embodiments, R<sup>3</sup> is methyl.  
 [0126] In some embodiments, R<sup>3</sup> is oxo.  
 [0127] In some embodiments, R<sup>3</sup> is halo. In some such embodiments, R<sup>3</sup> is fluoro. In other such embodiments, R<sup>3</sup> is  
 chloro. In yet other such embodiments, R<sup>3</sup> is bromo. In further such embodiments, R<sup>3</sup> is iodo.  
 55

**B4. Substituent R<sup>4</sup>.**

- [0128] R<sup>4</sup> is selected from the group consisting of halo, alkyl, alkenyl, alkynyl, nitro, cyano, azido, alkyloxy, alkenyloxy,

alkynyloxy, amino, aminocarbonyl, aminosulfonyl, alkylsulfonyl, carbocyclyl, and heterocyclyl, wherein:

(a) the amino, aminocarbonyl, and aminosulfonyl optionally are substituted with:

- 5 (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, and alkylsulfonyl, or  
 (2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl,

10 (b) the alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, alkynyloxy, and alkylsulfonyl, optionally are substituted with one or more substituents independently selected from the group consisting of halo, oxo, nitro, cyano, azido, hydroxy, amino, alkyloxy, trimethylsilyl, carbocyclyl, and heterocyclyl, wherein:  
 the amino optionally is substituted with:

- 15 (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylcarbonyl, alkylsulfonyl, alkyloxycarbonyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl, or  
 (2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl, and

20 (c) the carbocyclyl and heterocyclyl optionally are substituted with up to three substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, halo, oxo, nitro, cyano, azido, hydroxy, amino, alkyloxy, trimethylsilyl, carbocyclyl, and heterocyclyl, wherein:  
 the amino optionally is substituted with:

- 25 (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylcarbonyl, alkylsulfonyl, alkyloxycarbonyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl, or  
 (2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl.

30 **[0129]** In some embodiments,  $R^4$  is selected from the group consisting of halo, alkyl, alkenyl, alkynyl, nitro, cyano, azido, alkyloxy, alkenyloxy, alkynyloxy, amino, aminocarbonyl, aminosulfonyl, alkylsulfonyl, carbocyclyl, and heterocyclyl, wherein:

the amino, aminocarbonyl, and aminosulfonyl optionally are substituted with:

- 35 (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, and alkylsulfonyl, or  
 (2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl.

40 **[0130]** In some embodiments,  $R^4$  is selected from the group consisting of halo, alkyl, alkenyl, alkynyl, nitro, cyano, azido, alkyloxy, alkenyloxy, alkynyloxy, amino, aminocarbonyl, aminosulfonyl, alkylsulfonyl, carbocyclyl, and heterocyclyl, wherein:

the alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, alkynyloxy, and alkylsulfonyl, optionally are substituted with one or more substituents independently selected from the group consisting of halo, oxo, nitro, cyano, azido, hydroxy, amino, alkyloxy, trimethylsilyl, carbocyclyl, and heterocyclyl, wherein:

45 the amino optionally is substituted with:

- 50 (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylcarbonyl, alkylsulfonyl, alkyloxycarbonyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl, or  
 (2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl.

55 **[0131]** In some embodiments,  $R^4$  is selected from the group consisting of halo, alkyl, alkenyl, alkynyl, nitro, cyano, azido, alkyloxy, alkenyloxy, alkynyloxy, amino, aminocarbonyl, aminosulfonyl, alkylsulfonyl, carbocyclyl, and heterocyclyl, wherein:

the carbocyclyl and heterocyclyl optionally are substituted with up to three substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, halo, oxo, nitro, cyano, azido, hydroxy, amino, alkyloxy, trimethylsilyl, carbocyclyl, and heterocyclyl, wherein:

the amino optionally is substituted with:

- (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylcarbonyl, alkylsulfonyl, alkyloxycarbonyl, carbocyclyl, heterocyclyl, carbocyclalkyl, and heterocyclalkyl, or  
 (2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl.

**[0132]** In some embodiments,  $R^4$  is selected from the group consisting of halo, alkyl, alkenyl, alkynyl, nitro, cyano, azido, alkyloxy, alkenyloxy, alkynyloxy, amino, aminocarbonyl, aminosulfonyl, alkylsulfonyl, carbocyclyl, and heterocyclyl, wherein:

(a) the amino, aminocarbonyl, and aminosulfonyl optionally are substituted with:

- (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl, or,  
 (2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl; and

(b) the alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, alkynyloxy, alkylsulfonyl, carbocyclyl, and heterocyclyl optionally are substituted with up to three substituents independently selected from the group consisting of halo, oxo, nitro, cyano, azido, hydroxy, amino, alkyloxy, carbocyclyl, and heterocyclyl, wherein the amino optionally is substituted with:

- (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylcarbonyl, alkylsulfonyl, alkyloxycarbonyl, carbocyclyl, heterocyclyl, carbocyclalkyl, and heterocyclalkyl, or,  
 (2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl.

**[0133]** In some embodiments,  $R^4$  is selected from the group consisting of halo, alkyl, alkenyl, alkynyl, nitro, cyano, azido, alkyloxy, alkenyloxy, alkynyloxy, amino, aminocarbonyl, aminosulfonyl, alkylsulfonyl, carbocyclyl, and heterocyclyl, wherein:

the amino, aminocarbonyl, and aminosulfonyl optionally are substituted with:

- (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl, or,  
 (2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl.

**[0134]** In some embodiments,  $R^4$  is selected from the group consisting of halo, alkyl, alkenyl, alkynyl, nitro, cyano, azido, alkyloxy, alkenyloxy, alkynyloxy, amino, aminocarbonyl, aminosulfonyl, alkylsulfonyl, carbocyclyl, and heterocyclyl, wherein:

the alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, alkynyloxy, alkylsulfonyl, carbocyclyl, and heterocyclyl optionally are substituted with up to three substituents independently selected from the group consisting of halo, oxo, nitro, cyano, azido, hydroxy, amino, alkyloxy, carbocyclyl, and heterocyclyl, wherein the amino optionally is substituted with:

- (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylcarbonyl, alkylsulfonyl, alkyloxycarbonyl, carbocyclyl, heterocyclyl, carbocyclalkyl, and heterocyclalkyl, or,  
 (2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl.

**[0135]** In some embodiments,  $R^4$  is selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl,  $C_2$ - $C_4$ -alkenyl,  $C_2$ - $C_4$ -alkynyl, amino,  $C_1$ - $C_4$ -alkylsulfonyl,  $C_3$ - $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl, wherein:

(a) the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, and alkylsulfonyl,

(b) the  $C_1$ - $C_4$ -alkyl,  $C_2$ - $C_4$ -alkenyl, and  $C_2$ - $C_4$ -alkynyl optionally are substituted with one or more substituents independently selected from the group consisting of halo, oxo, hydroxy, alkyloxy, and trimethylsilyl, and

(c) the  $C_3$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl optionally are substituted with up to three substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, halo, and amino, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, and alkylsulfonyl.

**[0136]** In some embodiments,  $R^4$  is selected from the group consisting of  $C_1$ - $C_4$ -alkyl,  $C_2$ - $C_4$ -alkenyl,  $C_2$ - $C_4$ -alkynyl, amino,  $C_1$ - $C_4$ -alkylsulfonyl,  $C_3$ - $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl, wherein:

(a) the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, and alkylsulfonyl,

(b) the  $C_1$ - $C_4$ -alkyl,  $C_2$ - $C_4$ -alkenyl, and  $C_2$ - $C_4$ -alkynyl optionally are substituted with one or more substituents independently selected from the group consisting of halo, oxo, hydroxy, alkyloxy, and trimethylsilyl, and

(c) the  $C_3$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl optionally are substituted with up to three substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, halo, and amino, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, and alkylsulfonyl.

**[0137]** In some embodiments,  $R^4$  is selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl, wherein:

(a) the  $C_1$ - $C_4$ -alkyl optionally is substituted with up to three substituents independently selected from the group consisting of halo, oxo, hydroxy, alkyloxy, and trimethylsilyl, and

(b) the  $C_3$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, halo, and alkylsulfonylamino.

**[0138]** In some embodiments,  $R^4$  is selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl, wherein:

(a) the  $C_1$ - $C_4$ -alkyl optionally is substituted with one or two substituents independently selected from the group consisting of halo, oxo, hydroxy, alkyloxy, and trimethylsilyl, and

(b) the  $C_3$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl optionally are substituted with a substituent selected from the group consisting of alkyl, halo, and alkylsulfonylamino.

**[0139]** In some embodiments,  $R^4$  is selected from the group consisting of  $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl, wherein:

(a) the  $C_1$ - $C_4$ -alkyl optionally is substituted with up to three substituents independently selected from the group consisting of halo, oxo, hydroxy, alkyloxy, and trimethylsilyl, and

(b) the  $C_3$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, halo, and alkylsulfonylamino.

**[0140]** In some embodiments,  $R^4$  is selected from the group consisting of halo, tert-butyl,  $C_3$ - $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl, wherein:

the  $C_3$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl optionally are substituted with a substituent selected from the group consisting of alkyl, halo, and alkylsulfonylamino.

**[0141]** In some embodiments,  $R^4$  is selected from the group consisting of tert-butyl,  $C_3$ - $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl, wherein:

the  $C_3$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl optionally are substituted with a substituent selected from the group consisting of alkyl, halo, and alkylsulfonylamino.

**[0142]** In some embodiments,  $R^4$  is selected from the group consisting of halo, alkyl, haloalkyl, carboxyalkyl, hydroxyalkyl, alkyloxyalkyl, trimethylsilylalkynyl, alkylcarbocyclyl, carbocyclyl, alkylheterocyclyl, heterocyclyl, halocarbocyclyl, alkylsulfonylamino, and alkylsulfonyl.

**[0143]** In some embodiments,  $R^4$  is selected from the group consisting of halo, alkyl, alkenyl, alkynyl, nitro, cyano, azido, alkyloxy, alkenyloxy, alkynyloxy, amino, aminocarbonyl, aminosulfonyl, alkylsulfonyl, carbocyclyl, and heterocyclyl.

**[0144]** In some embodiments,  $R^4$  is selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl,  $C_2$ - $C_4$ -alkenyl,  $C_2$ - $C_4$ -alkynyl, amino,  $C_1$ - $C_4$ -alkylsulfonyl,  $C_3$ - $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl. In some such embodiment,  $R^4$  is selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl,  $C_2$ - $C_4$ -alkenyl,  $C_2$ - $C_4$ -alkynyl, amino,  $C_1$ - $C_4$ -alkylsulfonyl,  $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl. In other such embodiment,  $R^4$  is selected from the group consisting of halo,

C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, amino, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, phenyl, and 5-6-membered heteroaryl.

[0145] In some embodiments, R<sup>4</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, amino, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>3</sub>-C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In some such embodiment, R<sup>4</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, amino, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In other such embodiment, R<sup>4</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, amino, C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl, phenyl, and 5-6-membered heteroaryl.

[0146] In some embodiments, R<sup>4</sup> is selected from the group consisting of halo, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In some such embodiments, R<sup>4</sup> is selected from the group consisting of halo, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In other such embodiments, R<sup>4</sup> is selected from the group consisting of halo, C<sub>1</sub>-C<sub>4</sub>-alkyl, phenyl, and 5-6-membered heteroaryl.

[0147] In some embodiments, R<sup>4</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In some such embodiments, R<sup>4</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In other such embodiments, R<sup>4</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, phenyl, and 5-6-membered heteroaryl.

[0148] In some embodiments, R<sup>4</sup> is selected from the group consisting of halo, tert-butyl, C<sub>3</sub>-C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In some such embodiments, R<sup>4</sup> is selected from the group consisting of halo, tert-butyl, C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In other such embodiments, R<sup>4</sup> is selected from the group consisting of halo, tert-butyl, phenyl, and 5-6-membered heteroaryl.

[0149] In some embodiments, R<sup>4</sup> is selected from the group consisting of tert-butyl, C<sub>3</sub>-C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In some such embodiments, R<sup>4</sup> is selected from the group consisting of tert-butyl, C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In other such embodiments, R<sup>4</sup> is selected from the group consisting of tert-butyl, phenyl, and 5-6-membered heteroaryl.

[0150] In some embodiments, R<sup>4</sup> is selected from the group consisting of C<sub>3</sub>-C<sub>6</sub>-carbocyclyl and 5-6-membered heterocyclyl. In some such embodiments, R<sup>4</sup> is selected from the group consisting of C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl. In other such embodiments, R<sup>4</sup> is selected from the group consisting of phenyl and 5-6-membered heteroaryl.

[0151] Suitable carbocyclyls for the above embodiments include, for example, cyclopropyl and phenyl.

[0152] Suitable heterocyclyls for the above embodiments include, for example, furanyl, thienyl, and pyridinyl.

[0153] In some embodiments, R<sup>4</sup> is selected from the group consisting of halo, alkyl, and alkyloxy.

[0154] In some embodiments, R<sup>4</sup> is alkyl.

[0155] In some embodiments, R<sup>4</sup> is tert-butyl.

#### B5. Substituent R<sup>5</sup>.

[0156] R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, alkynyloxy, alkylsulfonyloxy, carbocyclylsulfonyloxy, haloalkylsulfonyloxy, and halo.

[0157] In some embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, alkyloxy, and halo. In some such embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, alkyloxy, and fluoro. In other such embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, alkyloxy, and fluoro. In yet other such embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, alkyloxy, and chloro. In yet other such embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, alkyloxy, and bromo. In further such embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, alkyloxy, and iodo.

[0158] In some embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, methoxy, and halo. In some such embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, methoxy, and fluoro. In other such embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, methoxy, and chloro. In yet other such embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, methoxy, and bromo. In further such embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, methoxy, and iodo.

[0159] In some embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, and alkyloxy. In some such embodiments, R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy, methoxy, and ethoxy.

[0160] In some embodiments, R<sup>5</sup> is hydrogen.

[0161] In some embodiments, R<sup>5</sup> is hydroxy.

[0162] In some embodiments, R<sup>5</sup> is alkyloxy.

[0163] In some embodiments, R<sup>5</sup> is methoxy.

[0164] In some embodiments, R<sup>5</sup> is ethoxy.


#### B6. Substituent L.

[0165] L is selected from the group consisting of C(R<sup>A</sup>)=C(R<sup>B</sup>), ethylene, and cyclopropyl-1,2-ene, wherein R<sup>A</sup> and R<sup>B</sup> are as discussed below.

[0166] In some embodiments, L is  $C(R^A)=C(R^B)$ , wherein  $R^A$  and  $R^B$  are as discussed below. In these embodiments, the compounds of formula I correspond in structure to formula I-L1:

5

10




15

[0167] In some such embodiments, the compounds correspond in structure to formula IA-L1:

20

25




30

[0168] In other such embodiments, the compounds correspond in structure to formula IB-L1:

35

40




[0169] Typically, the compounds of formula I-L1 are more potent if  $R^6$  and the phenyl-uracil are on opposite sides of the double bond (i.e., in trans configuration in relation to the double bond).

45


[0170] In some embodiments, L is ethylene. In these embodiments, the compounds of formula I correspond in structure to I-L5-2:

50


55




[0171] In some such embodiments, the compounds correspond in structure to formula IA-L5-2:




15 [0172] In other such embodiments, the compounds correspond in structure to formula IB-L5-2:




30 [0173] In some embodiments, L is cyclopropyl-1,2-ene. In these embodiments, the compounds of formula I correspond in structure to formula I-L8:



45 [0174] In some such embodiments, the compounds correspond in structure to formula IA-L8:



[0175] In other such embodiments, the compounds correspond in structure to formula IB-L8:



15 **B7. Substituents  $R^A$  and  $R^B$ .**

**[0176]**  $R^A$  and  $R^B$  are independently selected from the group consisting of hydrogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkyloxy,  $C_3$ - $C_8$ -cycloalkyl, and halo, wherein:

20 the  $C_1$ - $C_6$ -alkyl optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, halo, hydroxy, nitro, oxo, amino, cyano, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy, carbocyclyl, and heterocyclyl.

**[0177]** In some embodiments, one of  $R^A$  and  $R^B$  is hydrogen, and the other is selected from the group consisting of  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkyloxy,  $C_3$ - $C_8$ -cycloalkyl, and halo, wherein:

the  $C_1$ - $C_6$ -alkyl optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, halo, hydroxy, nitro, oxo, amino, cyano, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy, carbocyclyl, and heterocyclyl.

**[0178]** In some embodiments,  $R^A$  and  $R^B$  are independently selected from the group consisting of hydrogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkyloxy,  $C_3$ - $C_8$ -cycloalkyl, and halo.

**[0179]** In some of the above embodiments,  $R^A$  is hydrogen. In other of the above embodiments,  $R^B$  is hydrogen.

**[0180]** In some embodiment, one of  $R^A$  and  $R^B$  is hydrogen, and the other is selected from the group consisting of hydrogen, methyl, methoxy, and halo.

**[0181]** In some embodiments,  $R^A$  is hydrogen, and  $R^B$  is selected from the group consisting of methyl, methoxy, and halo. In some such embodiments,  $R^B$  is selected from the group consisting of methyl, methoxy, and fluoro. In other such embodiments,  $R^B$  is selected from the group consisting of methyl, methoxy, and chloro. In yet other such embodiments,  $R^B$  is selected from the group consisting of methyl, methoxy, and bromo. In further such embodiments,  $R^B$  is selected from the group consisting of methyl, methoxy, and iodo. In yet further such embodiments,  $R^B$  is selected from the group consisting of methyl, methoxy, chloro, and fluoro.

**[0182]** In some embodiments,  $R^B$  is hydrogen, and  $R^A$  is selected from the group consisting of methyl, methoxy, and halo. In some such embodiments,  $R^A$  is selected from the group consisting of methyl, methoxy, and fluoro. In other such embodiments,  $R^A$  is selected from the group consisting of methyl, methoxy, and chloro. In yet other such embodiments,  $R^A$  is selected from the group consisting of methyl, methoxy, and bromo. In further such embodiments,  $R^A$  is selected from the group consisting of methyl, methoxy, and iodo. In yet further such embodiments,  $R^A$  is selected from the group consisting of methyl, methoxy, chloro, and fluoro.

**[0183]** In some embodiments,  $R^A$  is hydrogen, and  $R^B$  is hydrogen.

50 **B8. Substituent  $R^6$ .**

**[0184]**  $R^6$  is selected from the group consisting of  $C_5$ - $C_6$ -carbocyclyl, 5-6-membered heterocyclyl, fused 2-ring carbocyclyl, and fused 2-ring heterocyclyl, wherein each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ , wherein  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$  are as described below. In some such embodiments, the  $C_5$ - $C_6$ -carbocyclyl, 5-6-membered heterocyclyl, fused 2-ring carbocyclyl, and fused 2-ring heterocyclyl are not substituted. In other such embodiments, the  $C_5$ - $C_6$ -carbocyclyl, 5-6-membered heterocyclyl, fused 2-ring carbocyclyl, and fused 2-ring heterocyclyl are substituted with a substituent selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In other such embodiments, the  $C_5$ - $C_6$ -car-



substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In further such embodiments, the 5-6-membered heterocyclyl is substituted with one, two, or three substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ .

**[0188]** In some embodiments,  $R^6$  is selected from the group consisting of fused 2-ring carbocyclyl and fused 2-ring heterocyclyl, wherein each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In some such embodiments, the fused 2-ring carbocyclyl and fused 2-ring heterocyclyl are not substituted. In other such embodiments, the fused 2-ring carbocyclyl and fused 2-ring heterocyclyl are substituted with a substituent selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In yet other such embodiments, the fused 2-ring carbocyclyl and fused 2-ring heterocyclyl are substituted with two substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In further such embodiments, the fused 2-ring carbocyclyl and fused 2-ring heterocyclyl are substituted with three substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ .

**[0189]** In some embodiments,  $R^6$  is fused 2-ring carbocyclyl optionally substituted with one or more substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In some such embodiments, the fused 2-ring carbocyclyl is not substituted. In other such embodiments, the fused 2-ring carbocyclyl is substituted with a substituent selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In yet other such embodiments, the fused 2-ring carbocyclyl is substituted with two substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In further such embodiments, the fused 2-ring carbocyclyl is substituted with three substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In further such embodiments, the fused 2-ring carbocyclyl is substituted with one, two, or three substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ .

**[0190]** In some embodiments,  $R^6$  is fused 2-ring heterocyclyl optionally substituted with one or more substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In some such embodiments, the fused 2-ring heterocyclyl is not substituted. In other such embodiments, the fused 2-ring heterocyclyl is substituted with a substituent selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In yet other such embodiments, the fused 2-ring heterocyclyl is substituted with two substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In further such embodiments, the fused 2-ring heterocyclyl is substituted with three substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ . In further such embodiments, the fused 2-ring heterocyclyl is substituted with one, two, or three substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$ .

**[0191]** In some of the above embodiments, the optionally substituted  $C_5$ - $C_6$ -carbocyclyl is selected from the group consisting of cyclopentyl, cyclopentenyl, cyclopentadienyl, cyclohexyl, cyclohexenyl, cyclohexadienyl, and phenyl. In some such embodiments, the optionally substituted  $C_5$ - $C_6$ -carbocyclyl is phenyl.

**[0192]** In some of the above embodiments, the optionally substituted  $C_5$ - $C_6$ -carbocyclyl is  $C_5$ -carbocyclyl. Examples of  $C_5$ -carbocyclyls include cyclopentyl, cyclopentenyl, and cyclopentadienyl.

**[0193]** In other of the above embodiments, the optionally substituted  $C_5$ - $C_6$ -carbocyclyl is  $C_6$ -carbocyclyl. Examples of  $C_6$ -carbocyclyls include cyclohexyl, cyclohexenyl, cyclohexadienyl, and phenyl.

**[0194]** In some of the above embodiments, the optionally substituted 5-6-membered-heterocyclyl is selected from the group consisting of furanyl, dihydrofuranyl, tetrahydrofuranyl, thiophenyl (thiofuranyl), dihydrothiophenyl, tetrahydrothiophenyl, pyrrolyl, pynoliny, pyrrolidinyl, oxazolyl, dihydrooxazolyl, isoxazolyl, dihydroisoxazolyl, oxazolidinyl, isoxazolidinyl, thiazolyl, isothiazolyl, thiazoliny, isothiazoliny, thiazolidiny, isothiazolidiny, imidazolyl, imidazolidinyl, pyrazolyl, pyrazoliny, pyrazolidiny, oxathioly, oxathiolany, triazolyl, oxadiazolyl, furazanyl, tetrazolyl, oxatriazolyl, dioxazolyl, oxathiazolyl, oxathiazolidiny, dihydrooxadiazolyl, dioxazolidinyl, pyranyl, dihydropyranyl, tetrahydropyranyl, pyridiny, dihydropyridiny, tetrahydropyridiny, piperidiny, diaziny, pyraziny, pyridaziny, pyrimidiny, dihydropyraziny, tetrahydropyraziny, piperaziny, triaziny, dihydrotriaziny, tetrahydrotriaziny, triazinany, oxaziny, dihydrooxaziny, morpholiny, oxathiaziny, dihydrooxathiaziny, oxathiazinany, oxadiaziny, dihydrooxadiaziny, oxadiazinany, thiopyranyl, dihydrothiopyranyl, and tetrahydrothiopyranyl.
















**[0195]** In some of the above embodiments, the optionally substituted 5-6-membered-heterocyclyl is 5-membered heterocyclyl. Examples of such 5-membered heterocyclyl include furanyl, dihydrofuranyl, tetrahydrofuranyl, thiophenyl (thiofuranyl), dihydrothiophenyl, tetrahydrothiophenyl, pyrrolyl, pyrroliny, pyrrolidinyl, oxazolyl, dihydrooxazolyl, isoxazolyl, dihydroisoxazolyl, oxazolidinyl, isoxazolidinyl, thiazolyl, isothiazolyl, thiazoliny, isothiazoliny, thiazolidiny, isothiazolidiny, imidazolyl, imidazolidinyl, pyrazolyl, pyrazoliny, pyrazolidiny, oxathioly, oxathiolany, triazolyl, oxadiazolyl, furazanyl, tetrazolyl, oxatriazolyl, dioxazolyl, oxathiazolyl, oxathiazolidiny, dihydrooxadiazolyl, and dioxazolidinyl.


**[0196]** In other of the above embodiments, the optionally substituted 5-6-membered-heterocyclyl is 6-membered heterocyclyl. Examples of 6-membered heterocyclyls include pyranyl, dihydropyranyl, tetrahydropyranyl, pyridiny, dihydropyridiny, tetrahydropyridiny, piperidiny, diaziny, pyraziny, pyridaziny, pyrimidiny, dihydropyraziny, tetrahydro-

pyrazinyl, piperazinyl, triazinyl, dihydrotriazinyl, tetrahydrotriazinyl, triazinanyl, oxazinyl, dihydrooxazinyl, morpholinyl, oxathiazinyl, dihydrooxathiazinyl, oxathiazinanyl, oxadiazinyl, dihydrooxadiazinyl, oxadiazinanyl, thiopyranyl, dihydrothiopyranyl, and tetrahydrothiopyranyl.

[0197] In some of the above embodiments, the optionally substituted fused 2-ring carbocyclyl is selected from the group consisting of naphthalenyl, dihydronaphthalenyl, tetrahydronaphthalenyl, hexahydronaphthalenyl, octahydronaphthalenyl, decahydronaphthalenyl, indenyl, dihydroindenyl, hexahydroindenyl, octahydroindenyl, pentalenyl, octahydropentalenyl, and hexahydropentalenyl. In some such embodiments, the optionally substituted fused 2-ring carbocyclyl is selected from the group consisting of naphthalenyl and dihydroindenyl. In some such embodiments, the optionally substituted fused 2-ring carbocyclyl is naphthalenyl. In other such embodiments, the optionally substituted fused 2-ring carbocyclyl is dihydroindenyl. In further such embodiments, the optionally substituted fused 2-ring carbocyclyl is indenyl.

[0198] In some of the above embodiments, the optionally substituted fused 2-ring heterocyclyl is selected from the group consisting of







- $X^1$ ,  $X^2$ , and  $X^3$  are independently selected from the group consisting of N and C(H);
- $X^4$  is selected from the group consisting of N(H), O, and S;
- $X^5$ ,  $X^6$ , and  $X^7$  are independently selected from the group consisting of N and C(H);
- $X^8$  is selected from the group consisting of N(H), O, and S;
- $X^9$  is selected from the group consisting of N(H), O, and S;
- $X^{10}$ ,  $X^{11}$ ,  $X^{12}$ , and  $X^{13}$  are independently selected from the group consisting of N and C(H);
- $X^{14}$  is selected from the group consisting of N(H), O, and S;
- $X^{15}$ ,  $X^{16}$ ,  $X^{17}$ , and  $X^{18}$  are independently selected from the group consisting of N and C(H);
- one or more of  $X^{19}$ ,  $X^{20}$ , and  $X^{21}$  is N, and the remaining one(s) is/are C(H);
- one or more of  $X^{22}$ ,  $X^{23}$ ,  $X^{24}$ , and  $X^{25}$  is N, and the remaining one(s) is/are C(H);
- one or more of  $X^{26}$ ,  $X^{27}$ , and  $X^{28}$  is N, and the remaining one(s) is/are C(H);
- one or more of  $X^{29}$ ,  $X^{30}$ ,  $X^{31}$ , and  $X^{32}$  is N, and the remaining one(s) is/are C(H);
- one or more of  $X^{33}$ ,  $X^{34}$ , and  $X^{35}$  is N, and the remaining one(s) is/are C(H);
- one or more of  $X^{36}$ ,  $X^{37}$ ,  $X^{38}$ , and  $X^{39}$  is N, and the remaining one(s) is/are C(H);
- $X^{40}$ ,  $X^{41}$ , and  $X^{42}$  are independently selected from the group consisting of N and C(H);
- one of  $X^{43}$ ,  $X^{44}$ , and  $X^{45}$  is selected from the group consisting of N(H), O, and S, and the remaining two are C(H)<sub>2</sub>;
- one of  $X^{46}$  and  $X^{47}$  is selected from the group consisting of N(H), O, and S, and the other one is C(H)<sub>2</sub>;
- $X^{48}$ ,  $X^{49}$ ,  $X^{50}$ , and  $X^{51}$  are independently selected from the group consisting of N and C(H);
- $X^{52}$ ,  $X^{53}$ , and  $X^{54}$  are independently selected from the group consisting of N and C(H);
- $X^{55}$  is selected from the group consisting of N(H), O, and S;
- $X^{56}$ ,  $X^{57}$ , and  $X^{58}$  are independently selected from the group consisting of N and C(H);
- $X^{59}$  is selected from the group consisting of N(H), O, and S;
- $X^{60}$  is selected from the group consisting of N(H), O, and S;
- $X^{61}$ ,  $X^{62}$ ,  $X^{63}$ , and  $X^{64}$  are independently selected from the group consisting of N and C(H);
- $X^{65}$  is selected from the group consisting of N(H), O, and S;
- $X^{66}$ ,  $X^{67}$ ,  $X^{68}$ , and  $X^{69}$  are independently selected from the group consisting of N and C(H);
- one or more of  $X^{70}$ ,  $X^{71}$ , and  $X^{72}$  is N, and the remaining one(s) is/are C(H);
- one or more of  $X^{73}$ ,  $X^{74}$ ,  $X^{75}$ , and  $X^{76}$  is N, and the remaining one(s) is/are C(H); and
- one of  $X^{77}$  and  $X^{78}$  is N(H), and the remaining one is C(H)<sub>2</sub>.

[0199] In some of the above embodiments, the optionally substituted fused 2-ring heterocyclyl is selected from the

group consisting of



[0200] In some of the above embodiments, the optionally substituted fused 2-ring heterocyclyl is selected from the group consisting of:



[0201] In some of the above embodiments, X<sup>1</sup>, X<sup>2</sup>, and X<sup>3</sup> are C(H).

[0202] In some of the above embodiments, X<sup>5</sup>, X<sup>6</sup>, and X<sup>7</sup> are C(H).

[0203] In some of the above embodiments, X<sup>10</sup>, X<sup>11</sup>, X<sup>12</sup>, and X<sup>13</sup> are C(H).

[0204] In some of the above embodiments, X<sup>15</sup>, X<sup>16</sup>, X<sup>17</sup>, and X<sup>18</sup> are C(H).

[0205] In some of the above embodiments, one of X<sup>19</sup>, X<sup>20</sup>, and X<sup>21</sup> is N.

[0206] In some of the above embodiments, one of X<sup>22</sup>, X<sup>23</sup>, X<sup>24</sup>, and X<sup>25</sup> is N.

[0207] In some of the above embodiments, one of X<sup>26</sup>, X<sup>27</sup>, and X<sup>28</sup> is N, and one of X<sup>29</sup>, X<sup>30</sup>, X<sup>31</sup>, and X<sup>32</sup> is N.

[0208] In some of the above embodiments, X<sup>40</sup>, X<sup>41</sup>, and X<sup>42</sup> are C(H).

[0209] In some of the above embodiments, X<sup>48</sup>, X<sup>49</sup>, X<sup>50</sup>, and X<sup>51</sup> are C(H).

[0210] In some of the above embodiments, X<sup>52</sup>, X<sup>53</sup>, and X<sup>54</sup> are C(H).

[0211] In some of the above embodiments, X<sup>56</sup>, X<sup>57</sup>, and X<sup>58</sup> are C(H).

[0212] In some of the above embodiments, X<sup>61</sup>, X<sup>62</sup>, X<sup>63</sup>, and X<sup>64</sup> are C(H).

[0213] In some of the above embodiments, X<sup>66</sup>, X<sup>67</sup>, X<sup>68</sup>, and X<sup>69</sup> are C(H).

[0214] In some of the above embodiments, one or more of X<sup>70</sup>, X<sup>71</sup>, and X<sup>72</sup> is N, and the remaining one(s) is/are C(H).

[0215] In some of the above embodiments, one or more of X<sup>73</sup>, X<sup>74</sup>, X<sup>75</sup>, and X<sup>76</sup> is N, and the remaining one(s) is/are C(H).

### B9. Substituent R<sup>E</sup>.

[0216] Each R<sup>E</sup> is independently selected from the group consisting of halo, nitro, hydroxy, oxo, carboxy, cyano, amino, imino, azido, and aldehydo, wherein the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl.

[0217] In some embodiment, each R<sup>E</sup> is independently selected from the group consisting of halo, nitro, hydroxy, oxo,

carboxy, amino, imino, and aldehydo, wherein the amino optionally is substituted with one or two independently selected alkyl.

[0218] In some embodiment, each  $R^E$  is independently selected from the group consisting of halo, nitro, hydroxy, oxo, carboxy, amino, imino, aldehydo, and alkylamino.

[0219] In some embodiment, each  $R^E$  is independently selected from the group consisting of chloro, fluoro, nitro, hydroxy, oxo, carboxy, amino, imino, aldehydo, and alkylamino.

[0220] In some embodiment, each  $R^E$  is independently selected from the group consisting of halo, nitro, hydroxy, oxo, carboxy, cyano, amino, imino, and azido. In some such embodiments, each  $R^E$  is halo. In other such embodiments, each  $R^E$  is nitro. In yet other such embodiments, each  $R^E$  is hydroxy. In yet other such embodiments, each  $R^E$  is oxo. In yet other such embodiments, each  $R^E$  is carboxy. In yet other such embodiments, each  $R^E$  is cyano. In yet other such embodiments, each  $R^E$  is amino. In further such embodiments, each  $R^E$  is imino. In yet further such embodiments, each  $R^E$  is and azido.

[0221] In some embodiments, each  $R^E$  is independently selected from the group consisting of halo, nitro, hydroxy, oxo, carboxy, cyano, amino, and imino.

#### **B10. Substituent $R^F$ .**

[0222] Each  $R^F$  is independently selected from the group consisting of alkyl, alkenyl, and alkynyl, wherein:

each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, imino, nitro, azido, oxo, aminosulfonyl, alkylsulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, imino, aminosulfonyl, aminocarbonyl, carbocyclyl, and heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylsulfonyl, alkenylsulfonyl, alkynylsulfonyl, alkylsulfonylamino, hydroxy, and alkyloxy,

wherein:

amino portion of the alkylsulfonylamino optionally is substituted with a substituent selected from the group consisting of alkyl, alkenyl, and alkynyl.

[0223] In some embodiment, each  $R^F$  is independently selected from the group consisting of alkyl, alkenyl, and alkynyl, wherein:

each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, imino, nitro, azido, oxo, aminosulfonyl, alkylsulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, imino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylsulfonyl, alkenylsulfonyl, alkynylsulfonyl, and alkylsulfonylamino, wherein:

amino portion of the alkylsulfonylamino optionally is substituted with a substituent selected from the group consisting of alkyl, alkenyl, and alkynyl.

[0224] In some of the above embodiments, each  $R^F$  is independently selected from the group consisting of the alkyl, alkynyl, and alkenyl, wherein such substituents are not substituted.

[0225] In some embodiments, each  $R^F$  is independently selected from the group consisting of alkyl, alkenyl, and alkynyl, wherein:

each such substituent optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, imino, nitro, oxo, aminosulfonyl, alkylsulfonyl, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, imino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents

independently selected from the group consisting of alkyl, alkylsulfonyl, and alkylsulfonylamino,

wherein:

5 amino portion of the alkylsulfonylamino optionally is substituted with alkyl.

[0226] In some embodiments, each  $R^F$  is an independently selected alkyl optionally substituted with a substituent selected from the group consisting of carboxy, hydroxy, halo, amino, imino, nitro, oxo, aminosulfonyl, alkylsulfonyl, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

10 the amino, imino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkylsulfonyl, and alkylsulfonylamino, wherein:

15 amino portion of the alkylsulfonylamino optionally is substituted with alkyl.

[0227] In some embodiments, each  $R^F$  is an independently selected alkyl optionally substituted with a substituent selected from the group consisting of carboxy, halo, amino, imino, and aminosulfonyl, wherein:

20 the amino, imino, and aminosulfonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkylsulfonyl, and alkylsulfonylamino.

[0228] In some embodiments, each  $R^F$  is an independently selected alkyl optionally substituted with amino, wherein the amino optionally is substituted with alkylsulfonyl.

[0229] In some embodiments, each  $R^F$  is an independently selected alkyl substituted with amino, wherein the amino is substituted with alkylsulfonyl. In some such embodiments, each  $R^F$  is methylsulfonylaminoethyl.

[0230] In some embodiments, each  $R^F$  is independently selected from the group consisting of alkyl, alkenyl, and alkynyl, wherein:

30 each such substituent optionally is substituted with one, two, or three substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, imino, nitro, azido, oxo, aminosulfonyl, alkylsulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl.

[0231] In some embodiments, each  $R^F$  is independently selected alkyl substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, imino, nitro, azido, oxo, aminosulfonyl, alkylsulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl.

#### 40 **B11. Substituent $R^G$ .**

[0232] Each  $R^G$  is independently selected from the group consisting of carbocyclyl and heterocyclyl, wherein:

45 each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

50 the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl.

[0233] In some of the above embodiments, each  $R^G$  is independently selected from the group consisting of carbocyclyl and heterocyclyl, wherein such substituents are not substituted.

[0234] In some embodiments, each  $R^G$  is independently selected from the group consisting of carbocyclyl and heterocyclyl, wherein:

55 each such substituent optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, halo, amino, nitro, oxo, aminosulfonyl, alkyloxycarbonyl, alkylcarbonyloxy,

alkyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl and alkylsulfonyl.

**[0235]** In some of the above embodiments, the carbocyclyl is C<sub>3</sub>-C<sub>6</sub>-carbocyclyl.

**[0236]** In some of the above embodiments, the heterocyclyl is 5-6-membered heterocyclyl.

**B12. Substituent R<sup>H</sup>.**

**[0237]** Each R<sup>H</sup> is independently selected from the group consisting of alkyloxy, alkenyloxy, alkynyloxy, alkylsulfonyloxy, alkenylsulfonyloxy, and alkynylsulfonyloxy, wherein:

each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxy, alkenyloxy, alkynyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl.

**[0238]** In some of the above embodiments, each R<sup>H</sup> is independently selected from the group consisting of alkyloxy, alkenyloxy, alkynyloxy, alkylsulfonyloxy, alkenylsulfonyloxy, and alkynylsulfonyloxy, wherein such substituents are not substituted.

**[0239]** In some embodiments, each R<sup>H</sup> is independently selected from the group consisting of alkyloxy and alkylsulfonyloxy, wherein:

each such substituent optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, oxo, aminosulfonyl, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl and alkylsulfonyl.

**[0240]** In some embodiments, each R<sup>H</sup> is independently selected from the group consisting of alkyloxy and alkylsulfonyloxy, wherein:

each such substituent optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, oxo, aminosulfonyl, alkyloxy, alkenyloxy, alkynyloxy, cyano, and aminocarbonyl, wherein:

the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl and alkylsulfonyl.

**[0241]** In some embodiments, each R<sup>H</sup> is independently selected from the group consisting of alkyloxy and alkylsulfonyloxy, wherein:

each such substituent optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, oxo, aminosulfonyl, alkyloxy, alkenyloxy, alkynyloxy, cyano, and aminocarbonyl.

**[0242]** In some embodiments, each R<sup>H</sup> is independently selected alkyloxy.

**[0243]** In some embodiments, each R<sup>H</sup> is independently selected alkylsulfonyloxy.

**B13. Substituent R<sup>I</sup>.**

**[0244]** Each R<sup>I</sup> is independently selected from the group consisting of alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl,

aminocarbonyl, alkyloxycarbonyl, carbocyclylcarbonyl, and heterocyclylcarbonyl, wherein:

(a) the alkylcarbonyl, alkenylcarbonyl, and alkynylcarbonyl optionally are substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, and

(b) the aminocarbonyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkyloxyalkyl, carbocyclyl, heterocyclyl, alkylsulfonyl, and alkylsulfonylamino, wherein:

the carbocyclyl and heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of halo, alkyl, and oxo.

**[0245]** In some embodiments, each  $R^I$  is independently selected from the group consisting of alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl, aminocarbonyl, alkyloxycarbonyl, carbocyclylcarbonyl, and heterocyclylcarbonyl, wherein such substituents are not substituted.

**[0246]** In some embodiments, each  $R^I$  is independently selected from the group consisting of alkylcarbonyl, aminocarbonyl, alkyloxycarbonyl, carbocyclylcarbonyl, and heterocyclylcarbonyl, wherein:

(a) the alkylcarbonyl optionally is substituted with a substituent selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, oxo, aminosulfonyl, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy, and aminocarbonyl, and

(b) the aminocarbonyl optionally is substituted with a substituent selected from the group consisting of alkyl, alkyloxyalkyl, alkylsulfonyl, and alkylsulfonylamino.

**[0247]** In some embodiments, each  $R^I$  is independently selected from the group consisting of alkylcarbonyl and aminocarbonyl, wherein:

the aminocarbonyl optionally is substituted with a substituent selected from the group consisting of alkyl, alkyloxyalkyl, alkylsulfonyl, and alkylsulfonylamino.

**[0248]** In some embodiment, each  $R^I$  is independently selected from the group consisting of alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl, and aminocarbonyl, wherein:

(a) the alkylcarbonyl, alkenylcarbonyl, and alkynylcarbonyl optionally are substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, and

(b) the aminocarbonyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, and alkylsulfonylamino.

**[0249]** In some of the above embodiments, each  $R^I$  is independently selected from the group consisting of alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl, and aminocarbonyl, wherein such substituents are not substituted.

**[0250]** In some embodiments, each  $R^I$  is independently selected from the group consisting of alkylcarbonyl and aminocarbonyl, wherein:

(a) the alkylcarbonyl optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, and

(b) the aminocarbonyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl and alkylsulfonylamino.

**[0251]** In some embodiments, each  $R^I$  is independently selected from the group consisting of alkylcarbonyl and aminocarbonyl, wherein:

(a) the alkylcarbonyl optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, oxo, aminosulfonyl, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy, cyano, and aminocarbonyl, and

(b) the aminocarbonyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl and alkylsulfonylamino.

[0252] In some embodiments, each  $R^I$  is independently selected from the group consisting of alkylcarbonyl and aminocarbonyl, wherein:

the alkylcarbonyl optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl.

[0253] In some embodiments, each  $R^I$  is independently selected alkylcarbonyl.

[0254] In some embodiments, each  $R^I$  is independently selected aminocarbonyl.

#### **B14. Substituent $R^J$ .**

[0255] Each  $R^J$  is independently selected from the group consisting of carbocyclylsulfonylamino, heterocyclylsulfonylamino, alkylcarbonylamino, alkenylcarbonylamino, alkynylcarbonylamino, alkyloxycarbonylamino, alkenyloxycarbonylamino, alkynyloxycarbonylamino, alkylsulfonylamino, alkenylsulfonylamino, alkynylsulfonylamino, aminocarbonylamino, alkyloxycarbonylaminoimino, alkylsulfonylaminoimino, alkenylsulfonylaminoimino, and alkynylsulfonylaminoimino, wherein:

(a) the amino portion of such substituents optionally is substituted with a substituent independently selected from the group consisting of carbocyclylalkyl, heterocyclylalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkenyl, alkynyl, alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl, alkyloxycarbonyl, alkyloxyalkyloxycarbonyl, alkylcarbonyloxyalkyl, and alkylsulfonyl, wherein:

(1) the carbocyclyl portion of the carbocyclylalkyl and the heterocyclyl portion of the heterocyclylalkyl optionally are substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, alkyloxy, alkenyloxy, alkynyloxy, halo, nitro, cyano, azido, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl,

(b) the alkyl, alkenyl, and alkynyl portion of such substituents optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxycarbonyl, alkylcarbonyloxy, hydroxy, alkyloxy, carbocyclyl, heterocyclyl, and cyano, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, and alkynyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy;

(c) the carbocyclyl and heterocyclyl portions of such substituents optionally are substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, alkyloxy, alkenyloxy, alkynyloxy, halo, nitro, cyano, azido, and amino, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl.

[0256] In some embodiment, each  $R^J$  is independently selected from the group consisting of carbocyclylsulfonylamino, heterocyclylsulfonylamino, alkylcarbonylamino, alkenylcarbonylamino, alkynylcarbonylamino, alkyloxycarbonylamino, alkenyloxycarbonylamino, alkynyloxycarbonylamino, alkylsulfonylamino, alkenylsulfonylamino, alkynylsulfonylamino, aminocarbonylamino, alkylsulfonylaminoimino, alkenylsulfonylaminoimino, and alkynylsulfonylaminoimino, wherein:

(a) the amino portion of such substituents optionally is substituted with a substituent independently selected from the group consisting of carbocyclylalkyl, heterocyclylalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkenyl, alkynyl, alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl, alkyloxycarbonyl, alkyloxyalkyloxycarbonyl, alkylcarbonyloxyalkyl, and alkylsulfonyl, wherein:

(1) the carbocyclyl portion of the carbocyclylalkyl and the heterocyclyl portion of the heterocyclylalkyl optionally

are substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, alkyloxy, alkenyloxy, alkynyloxy, halo, nitro, cyano, azido, oxo, and amino, and (2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl,

5

(b) the alkyl, alkenyl, and alkynyl portion of such substituents optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxycarbonyl, alkylcarbonyloxy, hydroxy, alkyloxy, carbocyclyl, heterocyclyl, and cyano, wherein:

10 the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, and alkynyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy;

15 (c) the carbocyclyl and heterocyclyl portions of such substituents optionally are substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, alkyloxy, alkenyloxy, alkynyloxy, halo, nitro, cyano, azido, and amino, wherein:

20 the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl; and

[0257] In some of the above embodiments, each  $R^J$  is independently selected from the group consisting of carbocyclysulfonylamino, heterocyclysulfonylamino, alkylcarbonylamino, alkenylcarbonylamino, alkynylcarbonylamino, alkyloxycarbonylamino, alkenyloxycarbonylamino, alkynyloxycarbonylamino, alkylsulfonylamino, alkenylsulfonylamino, alkynylsulfonylamino, aminocarbonylamino, alkylsulfonylaminoimino, alkenylsulfonylaminoimino, and alkynylsulfonylaminoimino, wherein such substituents are not substituted.

25

[0258] In some embodiments, each  $R^J$  is independently selected from the group consisting of carbocyclysulfonylamino, heterocyclysulfonylamino, alkylcarbonylamino, alkyloxycarbonylamino, alkylsulfonylamino, aminocarbonylamino, and alkylsulfonylaminoimino, wherein:

30

(a) the amino portion of such substituents optionally is substituted with a substituent independently selected from the group consisting of carbocyclylalkyl, heterocyclylalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkylcarbonyl, alkyloxycarbonyl, alkyloxyalkyloxycarbonyl, alkylcarbonyloxyalkyl, and alkylsulfonyl, wherein:

35 (1) the carbocyclyl portion of the carbocyclylalkyl and the heterocyclyl portion of the heterocyclylalkyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl,

40

(b) the alkyl portion of such substituents optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxycarbonyl, alkylcarbonyloxy, hydroxy, alkyloxy, carbocyclyl, heterocyclyl, and cyano, wherein:

45 the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl and alkyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy;

50 (c) the carbocyclyl and heterocyclyl portions of such substituents optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, and amino, wherein:

the amino optionally is substituted with one or two substituents independently selected alkyl.

55

[0259] In some embodiments, each  $R^J$  is independently selected from the group consisting of carbocyclysulfonylamino, heterocyclysulfonylamino, alkylsulfonylamino, and alkylsulfonylaminoimino, wherein:

(a) the amino portion of such substituents optionally is substituted with a substituent independently selected from the group consisting of carbocyclalkyl, heterocyclalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkylcarbonyl, alkyloxy, alkyloxyalkyloxy, alkyloxyalkyloxy, alkyloxyalkyloxy, and alkylsulfonyl, wherein:

(1) the carbocycl portion of the carbocyclalkyl and the heterocycl portion of the heterocyclalkyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl,

(b) the alkyl portion of such substituents optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxy, alkyloxy, alkyloxy, alkyloxy, carbocycl, heterocycl, and cyano, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl and alkyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy;

(c) the carbocycl and heterocycl portions of such substituents optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, and amino, wherein:

the amino optionally is substituted with one or two substituents independently selected alkyl.

**[0260]** In some embodiments, each  $R^J$  is independently selected from the group consisting of carbocyclsulfonylamino, heterocyclsulfonylamino, alkylsulfonylamino, and alkylsulfonylaminoimino, wherein:

the amino portion of such substituents optionally is substituted with a substituent independently selected from the group consisting of carbocyclalkyl, heterocyclalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkylcarbonyl, alkyloxy, alkyloxyalkyloxy, alkyloxyalkyloxy, alkyloxyalkyloxy, and alkylsulfonyl, wherein:

(1) the carbocycl portion of the carbocyclalkyl and the heterocycl portion of the heterocyclalkyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl.

**[0261]** In some embodiments, each  $R^J$  is independently selected from the group consisting of carbocyclsulfonylamino, heterocyclsulfonylamino, alkylsulfonylamino, and alkylsulfonylaminoimino, wherein:

the alkyl portion of the alkylsulfonylamino and alkylsulfonylaminoimino optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxy, alkyloxy, alkyloxy, alkyloxy, carbocycl, heterocycl, and cyano, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl and alkyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy.

**[0262]** In some embodiments, each  $R^J$  is independently selected from the group consisting of carbocyclsulfonylamino, heterocyclsulfonylamino, alkylsulfonylamino, and alkylsulfonylaminoimino, wherein:

the carbocycl and heterocycl portions of such substituents optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, and amino.

**[0263]** In some embodiments, each  $R^J$  is independently selected from the group consisting of carbocyclsulfonylamino and heterocyclsulfonylamino, wherein:

the carbocyclyl and heterocyclyl portions of such substituents optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, and amino.

**[0264]** In some embodiments, each  $R^J$  is independently selected from the group consisting of alkylsulfonylamino, alkenylsulfonylamino, alkynylsulfonylamino, and alkylsulfonylaminoimino, wherein:

(a) the amino portion of such substituents optionally is substituted with a substituent independently selected from the group consisting of carbocyclylalkyl, heterocyclylalkyl, alkylcarboxyloxy, aminocarbonylalkyl, alkyl, alkylcarbonyl, alkyloxycarbonyl, alkyloxyalkyloxycarbonyl, alkylcarboxyloxyalkyl, and alkylsulfonyl, wherein:

(1) the carbocyclyl portion of the carbocyclylalkyl and the heterocyclyl portion of the heterocyclylalkyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl,

(b) the alkyl, alkenyl, and alkynyl portion of such substituents optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxycarbonyl, alkylcarboxyloxy, hydroxy, alkyloxy, carbocyclyl, heterocyclyl, and cyano, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl and alkyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy.

**[0265]** In some embodiments, each  $R^J$  is an independently selected alkylsulfonylamino, wherein:

(a) the amino portion of the alkylsulfonylamino optionally is substituted with a substituent independently selected from the group consisting of carbocyclylalkyl, heterocyclylalkyl, alkylcarboxyloxy, aminocarbonylalkyl, alkyl, alkylcarbonyl, alkyloxycarbonyl, alkyloxyalkyloxycarbonyl, alkylcarboxyloxyalkyl, and alkylsulfonyl, wherein:

(1) the carbocyclyl portion of the carbocyclylalkyl and the heterocyclyl portion of the heterocyclylalkyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl,

(b) the alkyl portion of the alkylsulfonylamino optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxycarbonyl, alkylcarboxyloxy, hydroxy, alkyloxy, carbocyclyl, heterocyclyl, and cyano, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl and alkyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy.

**[0266]** In some embodiments, each  $R^J$  is an independently selected alkylsulfonylamino, wherein:

the amino portion of the alkylsulfonylamino optionally is substituted with a substituent independently selected from the group consisting of carbocyclylalkyl, heterocyclylalkyl, alkylcarboxyloxy, aminocarbonylalkyl, alkyl, alkylcarbonyl, alkyloxycarbonyl, alkyloxyalkyloxycarbonyl, alkylcarboxyloxyalkyl, and alkylsulfonyl, wherein:

(1) the carbocyclyl portion of the carbocyclylalkyl and the heterocyclyl portion of the heterocyclylalkyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl.

[0267] In some embodiments, each  $R^J$  is an independently selected alkylsulfonylamino, wherein:

the amino portion of the alkylsulfonylamino optionally is substituted with a substituent independently selected from the group consisting of carbocyclalkyl, heterocyclalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkylcarbonyl, alkyloxy, alkyloxyalkyloxy, alkyloxyalkyloxy, alkylcarbonyloxyalkyl, and alkylsulfonyl.

[0268] In some embodiments, each  $R^J$  is an independently selected alkylsulfonylamino, wherein:

the alkyl portion of the alkylsulfonylamino optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxy, alkyloxyalkyloxy, hydroxy, alkyloxy, carbocycl, heterocycl, and cyano, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl and alkyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy.

[0269] In some embodiments, each  $R^J$  is an independently selected alkylsulfonylamino, wherein:

the alkyl portion of the alkylsulfonylamino optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxy, alkyloxyalkyloxy, hydroxy, alkyloxy, carbocycl, heterocycl, and cyano.

[0270] In some embodiments, each  $R^J$  is an independently selected alkylsulfonylamino. In some such embodiments, each  $R^J$  is methylsulfonylamino.

[0271] In some embodiments, each  $R^J$  is an independently selected alkylsulfonylaminoimino, wherein:

(a) the amino portion of the alkylsulfonylaminoimino optionally is substituted with a substituent independently selected from the group consisting of carbocyclalkyl, heterocyclalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkylcarbonyl, alkyloxy, alkyloxyalkyloxy, alkyloxyalkyloxy, alkylcarbonyloxyalkyl, and alkylsulfonyl, wherein:

(1) the carbocycl portion of the carbocyclalkyl and the heterocycl portion of the heterocyclalkyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl,

(b) the alkyl portion of the alkylsulfonylaminoimino optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxy, alkyloxyalkyloxy, hydroxy, alkyloxy, carbocycl, heterocycl, and cyano, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl and alkyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy.

[0272] In some embodiments, each  $R^J$  is an independently selected alkylsulfonylaminoimino, wherein:

the amino portion of the alkylsulfonylaminoimino optionally is substituted with a substituent independently selected from the group consisting of carbocyclalkyl, heterocyclalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkylcarbonyl, alkyloxy, alkyloxyalkyloxy, alkyloxyalkyloxy, alkylcarbonyloxyalkyl, and alkylsulfonyl, wherein:

(1) the carbocycl portion of the carbocyclalkyl and the heterocycl portion of the heterocyclalkyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl.

**[0273]** In some embodiments, each  $R^J$  is an independently selected alkylsulfonylaminoimino, wherein:

the amino portion of the alkylsulfonylaminoimino optionally is substituted with a substituent independently selected from the group consisting of carbocyclalkyl, heterocyclalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkyl-  
 5 carbonyl, alkyloxycarbonyl, alkyloxyalkyloxycarbonyl, alkylcarbonyloxyalkyl, and alkylsulfonyl.

**[0274]** In some embodiments, each  $R^J$  is an independently selected alkylsulfonylaminoimino, wherein:

the alkyl portion of the alkylsulfonylaminoimino optionally is substituted with one or two substituents independently  
 10 selected from the group consisting of carboxy, halo, oxo, amino, alkyloxycarbonyl, alkylcarbonyloxy, hydroxy, alkyl-  
 loxy, carbocyclyl, heterocyclyl, and cyano, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting  
 15 of alkyl and alkyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy.

**[0275]** In some embodiments, each  $R^J$  is an independently selected alkylsulfonylaminoimino, wherein:

the alkyl portion of the alkylsulfonylaminoimino optionally is substituted with one or two substituents independently  
 20 selected from the group consisting of carboxy, halo, oxo, amino, alkyloxycarbonyl, alkylcarbonyloxy, hydroxy, alkyl-  
 loxy, carbocyclyl, heterocyclyl, and cyano.

**[0276]** In some embodiments, each  $R^J$  is an independently selected alkylsulfonylaminoimino. In some such embodi-  
 25 ments, each  $R^J$  is methylsulfonylaminoimino.

**[0277]** In some embodiments, each  $R^J$  is independently selected from the group consisting of alkylcarbonylamino and  
 alkyloxycarbonylamino, wherein:

the alkyl portion of such substituents optionally is substituted with one or two substituents independently selected  
 30 from the group consisting of carboxy, halo, oxo, amino, alkyloxycarbonyl, alkylcarbonyloxy, hydroxy, alkyloxy, carb-  
 ocyclyl, heterocyclyl, and cyano.

#### **B15. Substituent $R^K$ .**

**[0278]** Each  $R^K$  is independently selected from the group consisting of aminosulfonyl, alkylsulfonyl, alkenylsulfonyl,  
 35 and alkynylsulfonyl, wherein:

(a) the alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl optionally are substituted with one or more substituents  
 40 independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl,  
 alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbony-  
 loxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independ-  
 45 ently selected from the group consisting of alkyl, alkenyl, and alkynyl; and

(b) the aminosulfonyl optionally is substituted with one or two substituents independently selected from the group  
 consisting of alkyl, alkenyl, and alkynyl.

**[0279]** In some of the above embodiments, each  $R^K$  is independently selected from the group consisting of amino-  
 50 sulfonyl, alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl, wherein such substituents are not substituted.

**[0280]** In some embodiments, each  $R^K$  is independently selected from the group consisting of aminosulfonyl and  
 alkylsulfonyl, wherein:

(a) the alkylsulfonyl optionally is substituted with one or two substituents independently selected from the group  
 55 consisting of carboxy, hydroxy, halo, amino, nitro, oxo, aminosulfonyl, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy,  
 carbocyclyl, heterocyclyl, cyano, and aminocarbonyl; and

(b) the aminosulfonyl optionally is substituted with one or two substituents independently selected alkyl.

[0281] In some embodiments, each  $R^K$  is independently selected from the group consisting of aminosulfonyl and alkylsulfonyl.

C. Embodiments of Compounds of Formula I.

[0282] Various embodiments of substituents  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , L,  $R^A$ ,  $R^B$ ,  $R^C$ ,  $R^D$ ,  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ , and  $R^K$  have been discussed above. These substituent embodiments can be combined to form various embodiments of compounds of formula I. All embodiments of compounds of formula I formed by combining the substituent embodiments discussed above are within the scope of Applicants' invention, and some illustrative embodiments of the compounds of formula I are provided below.

[0283] In some embodiments, in the compounds of formula I:

$\text{---}\overset{\star}{\text{C}}\text{---}$  is selected from the group consisting of single carbon-carbon bond and double carbon-carbon bond;

$R^1$  is selected from the group consisting of hydrogen and methyl;

$R^1$  is selected from the group consisting of hydrogen and halo;

$R^3$  is selected from the group consisting of hydrogen and halo;

$R^4$  is selected from the group consisting of  $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl, wherein:

(a) the  $C_1$ - $C_4$ -alkyl optionally is substituted with up to three substituents independently selected from the group consisting of halo, oxo, hydroxy, alkyloxy, and trimethylsilyl, and

(b) the  $C_3$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, halo, and alkylsulfonylamino;

$R^5$  is selected from the group consisting of hydrogen, hydroxy, alkyloxy, and halo;

L is selected from the group consisting of  $C(R^A)=C(R^B)$ , ethylene, and cyclopropyl-1,2-ene; one of  $R^A$  and  $R^B$  is hydrogen, and the other is selected from the group consisting of hydrogen, methyl, methoxy, and halo;

$R^6$  is selected from the group consisting of  $C_5$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl, wherein each such substituent is substituted with one, two, or three substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ , and  $R^J$ ;

each  $R^E$  is independently selected from the group consisting of chloro, fluoro, nitro, hydroxy, oxo, carboxy, amino, imino, aldehyde, and alkylamino;

each  $R^F$  is an independently selected alkyl optionally substituted with a substituent selected from the group consisting of carboxy, halo, amino, imino, and aminosulfonyl, wherein:

the amino, imino, and aminosulfonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkylsulfonyl, and alkylsulfonylamino;

each  $R^I$  is independently selected from the group consisting of alkylcarbonyl and aminocarbonyl, wherein:

the aminocarbonyl optionally is substituted with a substituent selected from the group consisting of alkyl, alkyloxyalkyl, alkylsulfonyl, and alkylsulfonylamino; and

each  $R^J$  is independently selected from the group consisting of alkylsulfonylamino, alkenylsulfonylamino, alkynylsulfonylamino, and alkylsulfonylaminoimino, wherein:

(a) the amino portion of such substituents optionally is substituted with a substituent independently selected from the group consisting of carbocyclylalkyl, heterocyclylalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkylcarbonyl, alkyloxyalkyl, alkyloxyalkylalkyl, alkylcarbonyloxyalkyl, and alkylsulfonyl, wherein:

(1) the carbocyclyl portion of the carbocyclylalkyl and the heterocyclyl portion of the heterocyclylalkyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, carboxy, hydroxy, alkyloxy, halo, nitro, cyano, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl,


(b) the alkyl, alkenyl, and alkynyl portion of such substituents optionally is substituted with one or two substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxyalkyl, alkylcarbo-

nyloxy, hydroxy, alkyloxy, carbocyclyl, heterocyclyl, and cyano, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl and alkyloxy, wherein the alkyl optionally is substituted with one or more hydroxy.




[0284] Examples of compounds of formula I (and salts thereof) are shown in Tables 1-7 below. The synthesis examples below provide step-by-step preparation instructions for some of these compounds. The remaining compounds were prepared utilizing the general method-of-preparation discussion, specific synthesis examples below, and/or the discussion throughout this application.

TABLE 1



compound	R <sup>5</sup>	R <sup>B</sup>	substituent(s)
IA-L1-1.3	-OCH <sub>3</sub>	-Cl	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [Z]
IA-L1-1.4	-OCH <sub>3</sub>	-F	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [Z]
IA-L1-1.5	-OCH <sub>3</sub>	-F	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IA-L1-1.6	-OCH <sub>3</sub>	-CH <sub>3</sub>	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IA-L1-1.9	-OCH <sub>3</sub>	-H	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IA-L1-1.10	-OCH <sub>3</sub>	-H	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [Z]
IA-L1-1.11	-OCH <sub>3</sub>	-H	-4-N[C(O)CH <sub>3</sub> ]S(O) <sub>2</sub> CH <sub>3</sub> [E]
IA-L1-1.12	-OCH <sub>3</sub>	-H	-4-F [E]
IA-L1-1.13	-OCH <sub>3</sub>	-H	-4-NH <sub>2</sub> [E]
IA-L1-1.14	-OCH <sub>3</sub>	-H	-4-OCH <sub>3</sub> [E]
IA-L1-1.16	-H	-H	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IA-L1-1.17	-OCH <sub>3</sub>	-OCH <sub>3</sub>	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [Z]
IA-L1-1.18	-OCH <sub>3</sub>	-H	-- [E]
IA-L1-1.20	-OCH <sub>3</sub>	-H	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [Z]
IA-L1-1.21	-OCH <sub>3</sub>	-F	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [Z]:[E] (1:1)
IA-L1-1.22	-OCH <sub>3</sub>	-H	-4-NO <sub>2</sub> [E]
IA-L1-1.23	-OCH <sub>3</sub>	-Cl	-4-NO <sub>2</sub> [Z]
IA-L1-1.24	-OCH <sub>3</sub>	-CH <sub>3</sub>	-4-NO <sub>2</sub> [E]
IA-L1-1.25	-H	-H	-4-NO <sub>2</sub> [E]
IA-L1-1.26	-OCH <sub>3</sub>	-H	-3-F and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IA-L1-1.27	-OCH <sub>3</sub>	-H	-2-OCH <sub>3</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]

TABLE 2

compound	substituent(s)
IB-L1-1.1	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.4	-2-C(O)OH and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.5	-3-F and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.6	-2-C(O)H and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.7	-2-C(O)OCH <sub>3</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.8	-2-C(H)=N(OH) and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.9	-2-C(O)N(H)CH <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.10	-2-CH <sub>2</sub> OH and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.11	-2-C(O)OC(H) <sub>2</sub> CH <sub>3</sub> and 4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.13	-2-C(H) <sub>2</sub> OCH <sub>3</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.14	-2-C(O)N(CH <sub>3</sub> ) <sub>2</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.15	-2-CH <sub>3</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> and -5-F [E]
IB-L1-1.16	imidazol-2-yl and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.17	-2-C(O)N(H)CH <sub>3</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.18	 -2 and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.19	-2-C(H)=NOCH <sub>3</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.21	-2-C(O)NH <sub>2</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.22	 -2 and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.23	 -2 and 4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.14	-2-C(O)N(CH <sub>3</sub> )C(H) <sub>2</sub> C(H) <sub>2</sub> OCH <sub>3</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.25	-2-C(H) <sub>2</sub> OC(H)(CH <sub>3</sub> ) <sub>2</sub> and 4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]

EP 2 203 430 B1

(continued)





compound	substituent(s)
IB-L1-1.26	 -2 and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.27	 -2 and 4-N(MS(O) <sub>2</sub> CH <sub>3</sub> ) [E]
IB-L1-1.28	-2-NH <sub>2</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.29	 -2 and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.31	-2-C(H) <sub>2</sub> N(H)C(H) <sub>2</sub> C(H)C(H)(CH <sub>3</sub> ) <sub>2</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.32	-2-N(H)C(O)OC(CH <sub>3</sub> ) <sub>3</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.33	 -2 and 4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.34	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub> [Z]

TABLE 3

compound	R <sup>4</sup>
IB-L1-1.45	-C(CH <sub>3</sub> ) <sub>2</sub> C(H) <sub>2</sub> OH [E]
IB-L1-1.46	furan-2-yl [E]
IB-L1-1.47	 [E]
IB-L1-1.48	 [E]
IB-L1-1.49	-S(O) <sub>2</sub> CH <sub>3</sub> [E]
IB-L1-1.50	furan-3-yl [E]
IB-L1-1.51	-I [E]

(continued)

compound	R <sup>4</sup>
IB-L1-1.52	-Br [E]
IB-L1-1.53	pyridin-3-yl [E]
IB-L1-1.55	pyridin-4-yl [E]


TABLE 4

compound	R <sup>2</sup>	R <sup>5</sup>
IB-L1-1.2	-F	-OCH <sub>3</sub> [E]
IB-L1-1.12	-H	-Cl [E]
IB-L1-1.20	-Cl	-OCH <sub>3</sub> [E]
IB-L1-1.30	-H	-OCH <sub>2</sub> CH <sub>3</sub> [E]

TABLE 5


compound	R <sup>5</sup>
IA-L5-2-1.1	-OCH <sub>3</sub>
IA-L5-2-1.2	-H

TABLE 6

	substituent(s) as described in the table below
--	--

(continued)

compound	substituent(s)
IB-L5-2-1.1	-2-C(O)OCH <sub>3</sub> and -4-N(H)S(O) <sub>2</sub> CH <sub>3</sub>
IB-L5-2-1.2	-4-N(H)S(O) <sub>2</sub> CH <sub>3</sub>

**TABLE 7****IA-L8-1.1****D. Isomert.**

**[0285]** This invention also is directed, in part, to all stereoisomers of the compounds of formula I (and their salts). Stereoisomers include E/Z isomers (*i.e.*, isomers with regard to one or more double bonds), enantiomers (*i.e.*, stereoisomers that have opposite configurations at all stereogenic centers), and diastereoisomers (*i.e.*, stereoisomers that have the same configuration at one or more stereogenic centers, but differ at other stereogenic centers).

**E. Salts.**

**[0286]** This invention also is directed, in part, to all salts of the compounds of formula I. A salt of a compound may be advantageous due to one or more of the salt's properties, such as, for example, enhanced pharmaceutical stability in differing temperatures and humidities, or a desirable solubility in water or other solvents. Where a salt is intended to be administered to a patient (as opposed to, for example, being in use in an *in vitro* context), the salt preferably is pharmaceutically acceptable and/or physiologically compatible. The term "pharmaceutically acceptable" is used adjectivally in this patent application to mean that the modified noun is appropriate for use as a pharmaceutical product or as a part of a pharmaceutical product. Pharmaceutically acceptable salts include salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. In general, these salts typically may be prepared by conventional means by reacting, for example, the appropriate acid or base with a compound of the invention.

**[0287]** Pharmaceutically acceptable acid addition salts of the compounds of formula I can be prepared from an inorganic or organic acid. Examples of often suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric, and phosphoric acid. Suitable organic acids generally include, for example, aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic, and sulfonic classes of organic acids. Specific examples of often suitable organic acids include acetate, trifluoroacetate, formate, propionate, succinate, glycolate, gluconate, digluconate, lactate, malate, tartaric acid, citrate, ascorbate, glucuronate, maleate, fumarate, pyruvate, aspartate, glutamate, benzoate, anthranilic acid, mesylate, stearate, salicylate, p-hydroxybenzoate, phenylacetate, mandelate, embonate (pamoate), ethanesulfonate, benzenesulfonate, pantothenate, 2-hydroxyethanesulfonate, sulfanilate, cyclohexylaminosulfonate, algenic acid, beta-hydroxybutyric acid, galactarate, galacturonate, adipate, alginate, bisulfate, butyrate, camphorate, camphorsulfonate, cyclopentanepropionate, dodecylsulfate, glycoheptanoate, glycerophosphate, heptanoate, hexanoate, nicotinate, oxalate, palmoate, pectinate, 2-naphthalesulfonate, 3-phenylpropionate, picrate, pivalate, thiocyanate, tosylate, and undecanoate.

**[0288]** Pharmaceutically acceptable base addition salts of the compounds of formula I include, for example, metallic salts and organic salts. Preferred metallic salts include alkali metal (group Ia) salts, alkaline earth metal (group IIa) salts, and other physiologically acceptable metal salts. Such salts may be made from aluminum, calcium, lithium, magnesium, potassium, sodium, and zinc. Preferred organic salts can be made from amines, such as tromethamine, diethylamine, N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine), and procaine. Basic nitrogen-containing groups can be quaternized with agents such as lower alkyl (C<sub>1</sub>-C<sub>6</sub>) halides (*e.g.*, methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides), dialkyl sulfates (*e.g.*, dimethyl, diethyl,

dibutyl, and diamyl sulfates), long chain halides (e.g., decyl, lauryl, myristyl, and stearyl chlorides, bromides, and iodides), arylalkyl halides (e.g., benzyl and phenethyl bromides), and others.

[0289] In some embodiments, the salt is sodium salt of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

[0290] In some embodiments, the salt is disodium salt of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

[0291] In some embodiments, the salt is potassium salt of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

[0292] In some embodiments, the salt is monopotassium salt of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

#### F. Purity.

[0293] Compounds of formula I (and salts thereof) with any level of purity (including pure and substantially pure) are within the scope of Applicants' invention. The term "substantially pure" in reference to a compound/salt/isomer, means that the preparation/composition containing the compound/salt/isomer contains more than about 85% by weight of the compound/salt/isomer, preferably more than about 90% by weight of the compound/salt/isomer, preferably more than about 95% by weight of the compound/salt/isomer, preferably more than about 97% by weight of the compound/salt/isomer, and preferably more than about 99% by weight of the compound/salt/isomer.

#### G. Crystalline Forms of Some Specific Compounds and Salts of The Invention.

##### G1. Crystalline Forms of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide Disodium Salt.

[0294] This invention also relates, in part, to crystalline forms of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide disodium salt, namely the nonahydrate and tetrahydrate crystalline forms discussed below.

[0295] This invention relates, in part, to a nonahydrate crystalline disodium salt. The crystallographic unit cell parameters of the nonahydrate crystalline disodium salt have been determined to be as follows: a is 8.9Å, b is 9.4Å, and c is 20.7Å (more precisely, a is 8.926(2)Å, b is 9.415(2)Å, and c is 20.674(5)Å); the cell angles are:  $\alpha$  - 94.8°,  $\beta$  - 93.3°, and  $\gamma$  - 107.0° (more precisely,  $\alpha$  is 94.796(4)°,  $\beta$  is 93.345(4)°, and  $\gamma$  is 107.013(4)°); and the cell volume is 1649Å<sup>3</sup> (more precisely, 1649.3(7)Å<sup>3</sup>). The salt crystallizes in the P-1 space group.

[0296] In some embodiments, the disodium salt nonahydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 4.3±0.2, 10.4±0.2, 10.9±0.2, 11.6±0.2, 12.9±0.2, 14.7±0.2, 16.4±0.2, 17.8±0.2, 19.4±0.2, 19.8±0.2, 20.8±0.2, 21.9±0.2, and 23.5±0.2 degrees 2 $\theta$ . In some such embodiments, the disodium salt nonahydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 4.3±0.2, 10.4±0.2, 10.9±0.2, 11.6±0.2, 12.9±0.2, 14.7±0.2, 16.4±0.2, 17.8±0.2, 19.4±0.2, 19.9±0.2, 20.8±0.2, 21.9±0.2, and 23.5±0.2 degrees 2 $\theta$ . In other such embodiments, the disodium salt nonahydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 4.3±0.2, 10.4±0.2, 10.9±0.2, 11.6±0.2, 12.9±0.2, 14.7±0.2, 16.4±0.2, 17.9±0.2, 19.4±0.2, 19.8±0.2, 20.8±0.2, 21.9±0.2, and 23.5±0.2 degrees 2 $\theta$ .

[0297] In some embodiments, the disodium salt nonahydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 4.3±0.2, 10.4±0.2, 10.9±0.2, 11.6±0.2, 12.9±0.2, 14.7±0.2, 14.9±0.2, 16.4±0.2, 17.8±0.2, 19.4±0.2, 19.7±0.2, 19.8±0.2, 20.8±0.2, 20.9±0.2, 21.9±0.2, 22.1±0.2, and 23.5±0.2 degrees 2 $\theta$ . In some such embodiments, the disodium salt nonahydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 4.3±0.2, 10.4±0.2, 10.9±0.2, 11.6±0.2, 12.9±0.2, 14.7±0.2, 14.9±0.2, 16.4±0.2, 17.8±0.2, 19.4±0.2, 19.7±0.2, 19.8±0.2, 20.8±0.2, 20.9±0.2, 21.9±0.2, 22.1±0.2, and 23.5±0.2 degrees 2 $\theta$ . In other such embodiments, the disodium salt nonahydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 4.3±0.2, 10.4±0.2, 10.9±0.2, 11.6±0.2, 12.9±0.2, 14.7±0.2, 14.9±0.2, 16.4±0.2, 17.8±0.2, 19.4±0.2, 19.7±0.2, 19.8±0.2, 20.8±0.2, 20.9±0.2, 21.9±0.2, 22.1±0.2, and 23.5±0.2 degrees 2 $\theta$ .

[0298] In some embodiments, the disodium salt nonahydrate has an X-ray powder diffraction pattern substantially as shown in Figure 1. The 2 $\theta$  values for the peaks in Figure 1 (and their intensities) are as follows: 4.31 (100), 10.36 (12), 10.91 (23), 11.61 (52), 12.93 (24), 14.73 (65), 14.89 (20), 16.44 (41), 17.80 (38), 19.44 (26), 19.67 (37), 19.83 (59), 20.75 (69), 20.89 (21), 21.92 (43), 22.13 (40), and 22.42 (24).

[0299] This invention also relates, in part, to a process for preparing the disodium salt nonahydrate. It was prepared in aqueous medium. Aqueous NaOH (1M, 1.18ml) was added to (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-

1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.1**) (27.82mg) (molar ratio 1:20 acid:base). The resulting suspension was equilibrated at ambient conditions. The disodium salt nonahydrate formed seven days later through a solution-mediated process. Alternatively, the disodium salt nonahydrate was prepared by suspending 278.8mg of compound **IB-L1-1.1** in 1.25ml THF while heated to about 50°C. Aqueous NaOH (1N, 1.5ml, 2.2molar equivalents) was added. The solid dissolved completely to yield a clear solution, which was naturally cooled to ambient temperatures. The salt crystallized spontaneously. The molecular structure was determined by single crystal diffraction.

**[0300]** This invention relates, in part, to a tetrahydrate crystalline disodium salt.

**[0301]** In some embodiments, the disodium salt tetrahydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $4.8 \pm 0.2$ ,  $12.1 \pm 0.2$ ,  $14.0 \pm 0.2$ ,  $17.0 \pm 0.2$ ,  $17.5 \pm 0.2$ ,  $20.9 \pm 0.2$ ,  $21.6 \pm 0.2$ ,  $25.0 \pm 0.2$ , and  $29.5 \pm 0.2$  degrees  $2\theta$ . In some such embodiments, the disodium salt tetrahydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $4.8 \pm 0.2$ ,  $12.1 \pm 0.2$ ,  $14.0 \pm 0.2$ ,  $17.0 \pm 0.2$ ,  $17.5 \pm 0.2$ ,  $20.9 \pm 0.2$ ,  $21.6 \pm 0.2$ ,  $25.0 \pm 0.2$ , and  $29.5 \pm 0.2$  degrees  $2\theta$ . In other such embodiments, the disodium salt tetrahydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $4.8 \pm 0.2$ ,  $12.1 \pm 0.2$ ,  $14.0 \pm 0.2$ ,  $17.0 \pm 0.2$ ,  $17.5 \pm 0.2$ ,  $20.9 \pm 0.2$ ,  $21.6 \pm 0.2$ ,  $25.0 \pm 0.2$ , and  $29.5 \pm 0.2$  degrees  $2\theta$ .

**[0302]** In some embodiments, the disodium salt tetrahydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $4.8 \pm 0.2$ ,  $12.1 \pm 0.2$ ,  $14.0 \pm 0.2$ ,  $14.4 \pm 0.2$ ,  $17.0 \pm 0.2$ ,  $17.5 \pm 0.2$ ,  $20.9 \pm 0.2$ ,  $21.6 \pm 0.2$ ,  $25.0 \pm 0.2$ ,  $29.5 \pm 0.2$ , and  $34.2 \pm 0.2$  degrees  $2\theta$ . In some such embodiments, the disodium salt tetrahydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $4.8 \pm 0.2$ ,  $12.1 \pm 0.2$ ,  $14.0 \pm 0.2$ ,  $14.4 \pm 0.2$ ,  $17.0 \pm 0.2$ ,  $17.5 \pm 0.2$ ,  $20.9 \pm 0.2$ ,  $21.6 \pm 0.2$ ,  $25.0 \pm 0.2$ ,  $29.5 \pm 0.2$ , and  $34.2 \pm 0.2$  degrees  $2\theta$ . In other such embodiments, the disodium salt tetrahydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $4.8 \pm 0.2$ ,  $12.1 \pm 0.2$ ,  $14.0 \pm 0.2$ ,  $14.4 \pm 0.2$ ,  $17.0 \pm 0.2$ ,  $17.5 \pm 0.2$ ,  $20.9 \pm 0.2$ ,  $21.6 \pm 0.2$ ,  $25.0 \pm 0.2$ ,  $29.5 \pm 0.2$ , and  $34.2 \pm 0.2$  degrees  $2\theta$ .

**[0303]** In some embodiments, the disodium salt tetrahydrate has an X-ray powder diffraction pattern substantially as shown in Figure 2. The  $2\theta$  values for the peaks in Figure 2 (and their intensities) are as follows: 4.81 (100), 12.07 (7), 14.01 (27), 14.41 (8), 16.96 (18), 17.53 (11), 20.87 (18), 21.58 (22), 24.99 (11), 29.47 (9), and 34.20 (9).

**[0304]** This invention also relates, in part, to a process for preparing the disodium salt tetrahydrate by suspending the nonahydrate disodium salt in an organic solvent (e.g., ethanol, 1-propanol, or 2-propanol).

**G2. Crystalline Form of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide Dipotassium Salt.**

**[0305]** This invention also relates, in part, to a crystalline (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide dipotassium salt tetrahydrate.

**[0306]** The crystallographic unit cell parameters of the dipotassium salt tetrahydrate have been determined to be as follows: a is 14.5Å, b is 10.8Å, and c is 35.8Å (more precisely, a is 14.454(14)Å, b is 10.763(14)Å, and c is 35.75(4)Å); the cell angle iso  $\beta$  98.8° (more precisely,  $\beta$  is 98.82(3)°); and the cell volume is 5499Å<sup>3</sup> (more precisely, 5499(11)Å<sup>3</sup>). The salt crystallizes in the C2/c space group.

**[0307]** In some embodiments, the dipotassium salt tetrahydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $5.0 \pm 0.2$ ,  $11.9 \pm 0.2$ ,  $12.4 \pm 0.2$ ,  $13.7 \pm 0.2$ ,  $15.0 \pm 0.2$ ,  $16.5 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $20.8 \pm 0.2$ ,  $21.3 \pm 0.2$ ,  $22.2 \pm 0.2$ ,  $24.0 \pm 0.2$ ,  $26.4 \pm 0.2$ , and  $29.3 \pm 0.2$  degrees  $2\theta$ . In some such embodiments, the dipotassium salt tetrahydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $5.0 \pm 0.2$ ,  $11.9 \pm 0.2$ ,  $12.4 \pm 0.2$ ,  $13.7 \pm 0.2$ ,  $15.0 \pm 0.2$ ,  $16.5 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $20.8 \pm 0.2$ ,  $21.3 \pm 0.2$ ,  $22.2 \pm 0.2$ ,  $24.0 \pm 0.2$ ,  $26.4 \pm 0.2$ , and  $29.3 \pm 0.2$  degrees  $2\theta$ . In other such embodiments, the dipotassium salt tetrahydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $5.0 \pm 0.2$ ,  $11.9 \pm 0.2$ ,  $12.4 \pm 0.2$ ,  $13.7 \pm 0.2$ ,  $15.0 \pm 0.2$ ,  $16.5 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $20.8 \pm 0.2$ ,  $21.3 \pm 0.2$ ,  $22.2 \pm 0.2$ ,  $24.0 \pm 0.2$ ,  $26.4 \pm 0.2$ , and  $29.3 \pm 0.2$  degrees  $2\theta$ .

**[0308]** In some embodiments, the dipotassium salt tetrahydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $5.0 \pm 0.2$ ,  $11.9 \pm 0.2$ ,  $12.4 \pm 0.2$ ,  $12.6 \pm 0.2$ ,  $13.7 \pm 0.2$ ,  $15.0 \pm 0.2$ ,  $16.5 \pm 0.2$ ,  $16.7 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $20.7 \pm 0.2$ ,  $20.8 \pm 0.2$ ,  $21.3 \pm 0.2$ ,  $22.2 \pm 0.2$ ,  $22.4 \pm 0.2$ ,  $24.0 \pm 0.2$ ,  $26.4 \pm 0.2$ , and  $29.3 \pm 0.2$  degrees  $2\theta$ . In some such embodiments, the dipotassium salt tetrahydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $5.0 \pm 0.2$ ,  $11.9 \pm 0.2$ ,  $12.4 \pm 0.2$ ,  $12.6 \pm 0.2$ ,  $13.7 \pm 0.2$ ,  $15.0 \pm 0.2$ ,  $16.5 \pm 0.2$ ,  $16.7 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $20.7 \pm 0.2$ ,  $20.8 \pm 0.2$ ,  $21.3 \pm 0.2$ ,  $22.2 \pm 0.2$ ,  $22.4 \pm 0.2$ ,  $24.0 \pm 0.2$ ,  $26.4 \pm 0.2$ , and  $29.3 \pm 0.2$  degrees  $2\theta$ . In other such embodiments, the dipotassium salt tetrahydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $5.0 \pm 0.2$ ,  $11.9 \pm 0.2$ ,  $12.4 \pm 0.2$ ,  $12.6 \pm 0.2$ ,  $13.7 \pm 0.2$ ,  $15.0 \pm 0.2$ ,  $16.5 \pm 0.2$ ,  $16.7 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $20.7 \pm 0.2$ ,  $20.8 \pm 0.2$ ,  $21.3 \pm 0.2$ ,  $22.2 \pm 0.2$ ,  $22.4 \pm 0.2$ ,  $24.0 \pm 0.2$ ,  $26.4 \pm 0.2$ , and  $29.3 \pm 0.2$  degrees  $2\theta$ .

**[0309]** In some embodiments, the dipotassium salt tetrahydrate has an X-ray powder diffraction pattern substantially as shown in Figure 4. The  $2\theta$  values for the peaks in Figure 4 (and their intensities) are as follows: 5.00 (100), 11.86 (34), 12.39 (32), 12.64 (19), 13.70 (23), 15.03 (21), 16.47 (24), 16.66 (24), 17.12 (28), 20.75 (29), 20.81 (33), 21.34 (22), 22.15 (46), 22.38 (31), 24.02 (24), 26.44 (24), and 29.32 (21).

**[0310]** This invention also relates, in part, to a process for preparing the dipotassium salt tetrahydrate by suspending compound **D3-L1-1.1** (261.13mg) in 1.25min THF while heated to about 50°C. Aqueous KOH (1N, 1.3ml, 2.2 molar equivalent) was added. The solid dissolved completely to yield a clear solution, which was naturally cooled to ambient temperatures. Crystallization occurred during the slow evaporation process.

**G3. Crystalline Forms of (E)-N-(4-(3-tert-butyl-5-(2,4-diaro-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl) methanesulfonamide Monopotassium Salt.**

**[0311]** This invention also relates, in part, to crystalline forms of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide monopotassium salt, namely the trihydrate and dihydrate crystalline forms discussed below.

**[0312]** This invention relates, in part, to a monopotassium salt trihydrate. The crystallographic unit cell parameters of the trihydrate crystalline monopotassium salt have been determined to be as follows: a is 9.0Å, b is 8.3Å, and c is 18.6Å (more precisely, a is 9.0393(16)Å, b is 8.3332(15)Å, and c is 18.582(3)Å); the cell angles are:  $\alpha$  - 80.5°,  $\beta$  - 85.1°, and  $\gamma$  - 80.5° (more precisely,  $\alpha$  is 80.511 (2)°,  $\beta$  is 85.134(3)°, and  $\gamma$  is 80.531(2)°); and the cell volume is 1359Å<sup>3</sup> (more precisely, 1359.3(4)Å<sup>3</sup>). The salt crystallizes in the P-1 space group.

**[0313]** In some embodiments, the monopotassium salt trihydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 4.8±0.2, 10.8±0.2, 11.3±0.2, 13.4±0.2, 15.3±0.2, 16.9±0.2, 21.2±0.2, 21.7±0.2, 22.1±0.2, 22.5±0.2, and 23.0±0.2 degrees 2 $\theta$ . In some such embodiments, the monopotassium salt trihydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 4.8±0.2, 10.8±0.2, 11.3±0.2, 13.4±0.2, 15.3±0.2, 16.9±0.2, 21.2±0.2, 21.7±0.2, 22.1±0.2, 22.5±0.2, and 23.0±0.2 degrees 2 $\theta$ . In other such embodiments, the monopotassium salt trihydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 4.8±0.2, 10.8±0.2, 11.3±0.2, 13.4±0.2, 15.3±0.2, 16.9±0.2, 21.2±0.2, 21.7±0.2, 22.1±0.2, 22.5±0.2, and 23.0±0.2 degrees 2 $\theta$ .

**[0314]** In some embodiments, the monopotassium salt trihydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 4.8±0.2, 10.8±0.2, 11.3±0.2, 13.4±0.2, 13.6±0.2, 15.3±0.2, 16.9±0.2, 21.2±0.2, 21.7±0.2, 21.7±0.2, 22.1±0.2, 22.5±0.2, 22.6±0.2, and 23.0±0.2 degrees 2 $\theta$ . In some such embodiments, the monopotassium salt trihydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 4.8±0.2, 10.8±0.2, 11.3±0.2, 13.4±0.2, 13.6±0.2, 15.3±0.2, 16.9±0.2, 21.2±0.2, 21.7±0.2, 21.7±0.2, 22.1±0.2, 22.5±0.2, 22.6±0.2, and 23.0±0.2 degrees 2 $\theta$ . In other such embodiments, the monopotassium salt trihydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 4.9±0.2, 10.8±0.2, 11.3±0.2, 13.4±0.2, 13.6±0.2, 15.3±0.2, 16.9±0.2, 21.2±0.2, 21.7±0.2, 21.7±0.2, 22.1±0.2, 22.5±0.2, 22.6±0.2, and 23.0±0.2 degrees 2 $\theta$ .

**[0315]** In some embodiments, the monopotassium salt trihydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 4.8±0.2, 10.8±0.2, 11.3±0.2, 13.4±0.2, 13.6±0.2, 15.3±0.2, 16.9±0.2, 21.2±0.2, 21.7±0.2, 21.7±0.2, 22.1±0.2, 22.5±0.2, 22.6±0.2, and 23.0±0.2 degrees 2 $\theta$ . In some such embodiments, the monopotassium salt trihydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 4.8±0.2, 10.8±0.2, 11.3±0.2, 13.4±0.2, 15.3±0.2, 16.9±0.2, 21.2±0.2, 21.7±0.2, 22.1±0.2, 22.5±0.2, and 23.0±0.2 degrees 2 $\theta$ . In other such embodiments, the monopotassium salt trihydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 4.8±0.2, 10.8±0.2, 11.3±0.2, 13.4±0.2, 15.3±0.2, 16.9±0.2, 21.2±0.2, 21.7±0.2, 22.1±0.2, 22.5±0.2, and 23.0±0.2 degrees 2 $\theta$ .

**[0316]** In some embodiments, the monopotassium salt trihydrate has an X-ray powder diffraction pattern substantially as shown in Figure 5. The  $2\theta$  values for the peaks in Figure 5 (and their intensities) are as follows: 4.83 (60), 10.79 (100), 11.31 (22), 13.42 (41), 13.59 (18), 15.32 (21), 16.90 (38), 21.24 (22), 21.68 (20), 21.68 (21), 22.15 (22), 22.55 (29), 22.63 (23), and 23.02 (27).

**[0317]** This invention also relates, in part, to a process for preparing the monopotassium salt trihydrate. It was prepared by suspending compound **IB-L11.1** (108.81mg) in 0.4ml THF while heated to about 50°C. Aqueous KOH solution (1N, 0.278ml, 1.2 molar equivalent) was added. The solid dissolved completely to yield a clear solution. Additional 1.6ml THF was added to the solution, which was then naturally cooled to ambient temperatures and crystallization was observed. Alternatively, the monopotassium salt trihydrate was prepared by suspending compound **IB-L11.1** (343.89mg) in 1.0ml THF while heated to 50°C. Aqueous KOH (1 N, 0.878ml, 1.2 molar equivalent) was added. The solid dissolved completely to yield a clear solution. Ethanol was added to the solution dropwise to a total volume of 4.0ml. The solution was then naturally cooled to ambient temperature and crystallization was observed.

**[0318]** This invention relates, in part, to a monopotassium salt dihydrate.

**[0319]** In some embodiments, the monopotassium salt dihydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $7.7\pm 0.2$ ,  $8.8\pm 0.2$ ,  $16.1\pm 0.2$ , and  $19.7\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the monopotassium salt dihydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of degrees  $2\theta$ .

**[0320]** In some embodiments, the monopotassium salt dihydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $7.7\pm 0.2$ ,  $8.8\pm 0.2$ ,  $12.4\pm 0.2$ ,  $14.0\pm 0.2$ ,  $16.1\pm 0.2$ ,  $17.7\pm 0.2$ ,  $19.2\pm 0.2$ ,  $19.7\pm 0.2$ ,  $23.1\pm 0.2$ , and  $29.2\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the monopotassium salt dihydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $7.7\pm 0.2$ ,  $8.8\pm 0.2$ ,  $12.4\pm 0.2$ ,  $14.0\pm 0.2$ ,  $16.1\pm 0.2$ ,  $17.7\pm 0.2$ ,  $19.2\pm 0.2$ ,  $19.7\pm 0.2$ ,  $23.1\pm 0.2$ , and  $29.2\pm 0.2$  degrees  $2\theta$ . In other such embodiments, the monopotassium salt dihydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $7.7\pm 0.2$ ,  $8.8\pm 0.2$ ,  $12.4\pm 0.2$ ,  $14.0\pm 0.2$ ,  $16.1\pm 0.2$ ,  $17.7\pm 0.2$ ,  $19.2\pm 0.2$ ,  $19.7\pm 0.2$ ,  $23.1\pm 0.2$ , and  $29.2\pm 0.2$  degrees  $2\theta$ .

**[0321]** In some embodiments, the monopotassium salt dihydrate has an X-ray powder diffraction pattern substantially as shown in Figure 6. The  $2\theta$  values for the peaks in Figure 6 (and their intensities) are as follows: 7.68 (19), 8.83 (100), 12.40 (7), 13.97 (10), 16.12 (25), 17.75 (9), 19.22 (12), 19.73 (40), 23.05 (9), and 29.21 (7).

**[0322]** This invention also relates, in part, to a process for preparing the monopotassium salt dihydrate. It was prepared by suspending the monopotassium salt trihydrate in media of low water activity, such as an ethanol/H<sub>2</sub>O mixture (50/1 v/v). Alternatively, the monopotassium salt dihydrate was prepared by dissolving potassium trihydrate solid (1.8g) in 36mL of IPA and 4ml water at 80°C. The resulting solution was cooled to 55°C over 1h. The solution was then seeded with 7.5mg of dihydrate crystals at 55°C and maintained at 55°C for 1h. Heptane (36ml) was then added over 3h. The reaction mixture was cooled to 0°C, and filtration yielded a material containing both di- and trihydrate crystals. The solid was then reslurried in 20mL of 10:1 v/v EtOH/H<sub>2</sub>O at 50°C for 3h and cooled to 25°C over 5h. The slurry was then mixed at 25°C for additional 3 days and cooled to 0°C over 3h and held at this temperature for 2h. The resulting crystals were filtered and air-dried on filter funnel for 1h to give dihydrate. The dihydrate monopotassium salt was also prepared by slurrying a mixture of dihydrate and trihydrate crystals in 10:1 v/v EtOH/H<sub>2</sub>O at 80°C for 2 days. The potassium content was confirmed by ion chromatography.

**G4. Crystalline Form of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3, 4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl) methanesulfonamide 1/7 Potassium Salt.**

**[0323]** This invention also relates, in part, to a crystalline form of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide 1/7 potassium salt.

**[0324]** In some embodiments, the 1/7 potassium salt has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $7.7\pm 0.2$ ,  $8.3\pm 0.2$ ,  $10.1\pm 0.2$ ,  $10.6\pm 0.2$ ,  $11.4\pm 0.2$ ,  $12.0\pm 0.2$ ,  $13.4\pm 0.2$ ,  $15.6\pm 0.2$ ,  $16.3\pm 0.2$ ,  $16.7\pm 0.2$ ,  $17.2\pm 0.2$ ,  $18.3\pm 0.2$ ,  $18.8\pm 0.2$ ,  $19.4\pm 0.2$ ,  $19.9\pm 0.2$ ,  $20.2\pm 0.2$ ,  $20.5\pm 0.2$ ,  $21.2\pm 0.2$ ,  $22.1\pm 0.2$ , and  $22.9\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the 1/7 potassium salt has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $7.7\pm 0.2$ ,  $8.3\pm 0.2$ ,  $10.1\pm 0.2$ ,  $10.6\pm 0.2$ ,  $11.4\pm 0.2$ ,  $12.0\pm 0.2$ ,  $13.4\pm 0.2$ ,  $15.6\pm 0.2$ ,  $16.3\pm 0.2$ ,  $16.7\pm 0.2$ ,  $17.2\pm 0.2$ ,  $18.3\pm 0.2$ ,  $18.8\pm 0.2$ ,  $19.4\pm 0.2$ ,  $19.9\pm 0.2$ ,  $20.2\pm 0.2$ ,  $20.5\pm 0.2$ ,  $21.2\pm 0.2$ ,  $22.1\pm 0.2$ , and  $22.9\pm 0.2$  degrees  $2\theta$ . In other such embodiments, the 1/7 potassium salt has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $7.7\pm 0.2$ ,  $8.3\pm 0.2$ ,  $10.1\pm 0.2$ ,  $10.6\pm 0.2$ ,  $11.4\pm 0.2$ ,  $12.0\pm 0.2$ ,  $13.4\pm 0.2$ ,  $15.6\pm 0.2$ ,  $16.3\pm 0.2$ ,  $16.7\pm 0.2$ ,  $17.2\pm 0.2$ ,  $18.3\pm 0.2$ ,  $18.8\pm 0.2$ ,  $19.4\pm 0.2$ ,  $19.9\pm 0.2$ ,  $20.2\pm 0.2$ ,  $20.5\pm 0.2$ ,  $21.2\pm 0.2$ ,  $22.1\pm 0.2$ , and  $22.9\pm 0.2$  degrees  $2\theta$ .

**[0325]** In some embodiments, the 1/7 potassium salt has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $7.7\pm 0.2$ ,  $8.3\pm 0.2$ ,  $10.1\pm 0.2$ ,  $10.6\pm 0.2$ ,  $11.4\pm 0.2$ ,  $12.0\pm 0.2$ ,  $13.4\pm 0.2$ ,  $15.6\pm 0.2$ ,  $16.3\pm 0.2$ ,  $16.7\pm 0.2$ ,  $17.2\pm 0.2$ ,  $18.3\pm 0.2$ ,  $18.8\pm 0.2$ ,  $19.4\pm 0.2$ ,  $19.9\pm 0.2$ ,  $20.2\pm 0.2$ ,  $20.5\pm 0.2$ ,  $20.8\pm 0.2$ ,  $21.2\pm 0.2$ ,  $22.1\pm 0.2$ ,  $22.9\pm 0.2$ ,  $24.3\pm 0.2$ ,  $24.9\pm 0.2$ , and  $25.1\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the 1/7 potassium salt has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $7.7\pm 0.2$ ,  $8.3\pm 0.2$ ,  $10.1\pm 0.2$ ,  $10.6\pm 0.2$ ,  $11.4\pm 0.2$ ,  $12.0\pm 0.2$ ,  $13.4\pm 0.2$ ,  $15.6\pm 0.2$ ,  $16.3\pm 0.2$ ,  $16.7\pm 0.2$ ,  $17.2\pm 0.2$ ,  $18.3\pm 0.2$ ,  $18.8\pm 0.2$ ,  $19.4\pm 0.2$ ,  $19.9\pm 0.2$ ,  $20.2\pm 0.2$ ,  $20.5\pm 0.2$ ,  $20.8\pm 0.2$ ,  $21.2\pm 0.2$ ,  $22.1\pm 0.2$ ,  $22.9\pm 0.2$ ,  $24.3\pm 0.2$ ,  $24.9\pm 0.2$ , and  $25.1\pm 0.2$  degrees  $2\theta$ . In other such embodiments, the 1/7 potassium salt has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $7.7\pm 0.2$ ,  $8.3\pm 0.2$ ,  $10.1\pm 0.2$ ,  $10.6\pm 0.2$ ,  $11.4\pm 0.2$ ,  $12.0\pm 0.2$ ,  $13.4\pm 0.2$ ,  $15.6\pm 0.2$ ,  $16.3\pm 0.2$ ,  $16.7\pm 0.2$ ,  $17.2\pm 0.2$ ,  $18.3\pm 0.2$ ,  $18.8\pm 0.2$ ,  $19.4\pm 0.2$ ,  $19.9\pm 0.2$ ,  $20.2\pm 0.2$ ,  $20.5\pm 0.2$ ,  $20.8\pm 0.2$ ,  $21.2\pm 0.2$ ,  $22.1\pm 0.2$ ,  $22.9\pm 0.2$ ,  $24.3\pm 0.2$ ,  $24.9\pm 0.2$ , and  $25.1\pm 0.2$  degrees  $2\theta$ .

**[0326]** In some embodiments, the 1/7 potassium salt has an X-ray powder diffraction pattern substantially as shown in Figure 8. The  $2\theta$  values for the peaks in Figure 8 (and their intensities) are as follows: 7.71 (19), 8.33 (34), 10.10 (100), 10.66 (29), 11.39 (27), 12.04 (22), 13.39 (39), 15.56 (41), 16.27 (62), 16.69 (70), 17.22 (59), 18.31 (18), 18.78 (47), 19.44 (36), 19.89 (28), 20.19 (33), 20.54 (87), 20.80 (33), 21.15 (47), 22.05 (24), 22.82 (67), 24.32 (22), 24.87 (22), and 25.07 (33).

[0327] This invention also relates, in part, to a process for preparing the 1/7 potassium salt. It was prepared by suspending compound **IB-L1-1.1** (2g) 6ml THF at 50°C. One molar equivalent of KOH dissolved in 4.3ml water was added, and the reaction mixture was heated to 65°C to dissolve all solids. The solution was then cooled to ambient temperatures over 2h and spontaneous crystallization took place. The slurry was then cooled to 5°C and held at that temperature for 2h. The pale yellow crystals were filtered and air-dried for 24h at ambient conditions. The potassium content was determined by ion chromatography.

**G5. Crystalline Form of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide Monodiethylamine Salt Tetrahydrate.**

[0328] This invention also relates, in part, to crystalline (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide monodiethylamine salt tetrahydrate.

[0329] In some embodiments, the monodiethylamine salt tetrahydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $9.5\pm 0.2$ ,  $10.0\pm 0.2$ ,  $11.8\pm 0.2$ ,  $12.1\pm 0.2$ ,  $14.4\pm 0.2$ ,  $16.8\pm 0.2$ ,  $17.6\pm 0.2$ ,  $19.8\pm 0.2$ ,  $20.8\pm 0.2$ ,  $21.4\pm 0.2$ ,  $21.8\pm 0.2$ , and  $29.8\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the monodiethylamine salt tetrahydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $9.5\pm 0.2$ ,  $10.0\pm 0.2$ ,  $11.8\pm 0.2$ ,  $12.1\pm 0.2$ ,  $14.4\pm 0.2$ ,  $16.8\pm 0.2$ ,  $17.6\pm 0.2$ ,  $19.8\pm 0.2$ ,  $20.8\pm 0.2$ ,  $21.4\pm 0.2$ ,  $21.8\pm 0.2$ , and  $29.8\pm 0.2$  degrees  $2\theta$ . In other such embodiments, the monodiethylamine salt tetrahydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $9.5\pm 0.2$ ,  $10.0\pm 0.2$ ,  $11.8\pm 0.2$ ,  $12.1\pm 0.2$ ,  $14.4\pm 0.2$ ,  $16.8\pm 0.2$ ,  $17.6\pm 0.2$ ,  $19.8\pm 0.2$ ,  $20.8\pm 0.2$ ,  $21.4\pm 0.2$ ,  $21.8\pm 0.2$ , and  $29.8\pm 0.2$  degrees  $2\theta$ .

[0330] In some embodiments, the monodiethylamine salt tetrahydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $9.5\pm 0.2$ ,  $10.0\pm 0.2$ ,  $11.8\pm 0.2$ ,  $12.1\pm 0.2$ ,  $14.4\pm 0.2$ ,  $16.8\pm 0.2$ ,  $17.6\pm 0.2$ ,  $19.4\pm 0.2$ ,  $19.8\pm 0.2$ ,  $20.8\pm 0.2$ ,  $21.4\pm 0.2$ ,  $21.8\pm 0.2$ ,  $21.9\pm 0.2$ , and  $29.8\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the monodiethylamine salt tetrahydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $9.5\pm 0.2$ ,  $10.0\pm 0.2$ ,  $11.8\pm 0.2$ ,  $12.1\pm 0.2$ ,  $14.4\pm 0.2$ ,  $16.8\pm 0.2$ ,  $17.6\pm 0.2$ ,  $19.4\pm 0.2$ ,  $19.8\pm 0.2$ ,  $20.8\pm 0.2$ ,  $21.4\pm 0.2$ ,  $21.8\pm 0.2$ ,  $21.9\pm 0.2$ , and  $29.8\pm 0.2$  degrees  $2\theta$ . In other such embodiments, the monodiethylamine salt tetrahydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $9.5\pm 0.2$ ,  $10.0\pm 0.2$ ,  $11.8\pm 0.2$ ,  $12.1\pm 0.2$ ,  $14.4\pm 0.2$ ,  $16.8\pm 0.2$ ,  $17.6\pm 0.2$ ,  $19.4\pm 0.2$ ,  $19.8\pm 0.2$ ,  $20.8\pm 0.2$ ,  $21.4\pm 0.2$ ,  $21.8\pm 0.2$ ,  $21.9\pm 0.2$ , and  $29.8\pm 0.2$  degrees  $2\theta$ .

[0331] In some embodiments, the monodiethylamine salt tetrahydrate has an X-ray powder diffraction pattern substantially as shown in Figure 9. The  $2\theta$  values for the peaks in Figure 9 (and their intensities) are as follows: 9.45 (100), 9.97 (31), 11.85 (67), 12.09 (16), 14.38 (22), 16.80 (9), 17.59 (10), 19.39 (8), 19.83 (21), 20.85 (25), 21.37 (12), 21.75 (34), 21.87 (8), and 29.78 (7).

[0332] This invention also relates, in part, to a process for preparing the monodiethylamine salt tetrahydrate. It was prepared in aqueous medium. Compound **IB-L1-1.1** was slowly added to 500ul of 1M diethylamine until no more solid can be dissolved into the solution. The solution was then evaporated slowly at ambient temperatures and the salt crystallized 2 days later. Alternatively, the monodiethylamine salt tetrahydrate was prepared by suspending 64.15mg of compound **IB-L1-1.1** in 400ul 1M diethylamine while heated to 50°C. About 5 drops of THF (- 20ul) was added. The solid dissolved completely upon addition to yield a clear solution. The solution was then evaporated at ambient temperature, and the salt crystallized 4 days later. The stoichiometry of the salt was confirmed by solution  $^1\text{H}$  NMR.

**G6. Crystalline Forms of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.1**).**

[0333] This invention also relates, in part, to crystalline forms of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.1**), namely the true polymorphs (pattern A, pattern B, pattern C, and pattern D) and hydrate (pattern AH, pattern BH, pattern CH, and pattern DH) crystalline forms discussed below.

**G6A. **IB-L1-1.1** True Polymorphs.**

[0334] This invention relates, in part, to pattern A crystalline (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

[0335] In some embodiments, the pattern A polymorph has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $5.8\pm 0.2$ ,  $9.9\pm 0.2$ ,  $11.8\pm 0.2$ ,  $12.4\pm 0.2$ ,  $14.5\pm 0.2$ ,  $18.8\pm 0.2$ ,  $22.7\pm 0.2$ , and  $29.2\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the pattern A polymorph has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $5.8\pm 0.2$ ,  $9.9\pm 0.2$ ,  $11.8\pm 0.2$ ,  $12.4\pm 0.2$ ,  $14.5\pm 0.2$ ,

18.8±0.2, 22.7±0.2, and 29.2±0.2 degrees 2θ. In other such embodiments, the pattern A polymorph has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 5.8±0.2, 9.9±0.2, 11.8±0.2, 12.4±0.2, 14.5±0.2, 18.8±0.2, 22.7±0.2, and 29.2±0.2 degrees 2θ.

5 [0336] In some embodiments, the pattern A polymorph has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 5.8±0.2, 9.9±0.2, 11.8±0.2, 12.4±0.2, 14.0±0.2, 14.5±0.2, 15.3±0.2, 18.5±0.2, 18.8±0.2, 22.2±0.2, 22.7±0.2, 23.8±0.2, 26.0±0.2, and 29.2±0.2 degrees 2θ. In some such embodiments, the pattern A polymorph has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 5.8±0.2, 9.9±0.2, 11.8±0.2, 12.4±0.2, 14.0±0.2, 14.5±0.2, 15.3±0.2, 18.5±0.2, 18.8±0.2, 22.2±0.2, 22.7±0.2, 23.8±0.2, 26.0±0.2, and 29.2±0.2 degrees 2θ. In other such embodiments, the pattern A polymorph has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 5.8±0.2, 9.9±0.2, 11.8±0.2, 12.4±0.2, 14.0±0.2, 14.5±0.2, 15.3±0.2, 18.5±0.2, 18.8±0.2, 22.2±0.2, 22.7±0.2, 23.8±0.2, 26.0±0.2, and 29.2±0.2 degrees 2θ.

10 [0337] In some embodiments, the pattern A polymorph has an X-ray powder diffraction pattern substantially as shown in Figure 11. The 2θ values for the peaks in Figure 11 (and their intensities) are as follows: 5.85 (28), 9.88 (51), 11.79 (73), 12.38 (56), 14.03 (38), 14.45 (100), 15.27 (29), 18.52 (39), 18.80 (47), 22.24 (40), 22.72 (77), 23.76 (39), 25.98 (22), and 29.21 (64).

15 [0338] This invention also relates, in part, to a process for preparing pattern A polymorph. Pattern A polymorph was prepared as discussed in Example E below.

20 [0339] This invention relates, in part, to pattern B crystalline (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

25 [0340] In some embodiments, the pattern B polymorph has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 11.5±0.2, 13.3±0.2, 15.4±0.2, 16.4±0.2, 17.1±0.2, 18.6±0.2, 19.4±0.2, 20.4±0.2, 21.6±0.2, 22.4±0.2, 24.0±0.2, 26.8±0.2, and 29.0±0.2 degrees 2θ. In some such embodiments, the pattern B polymorph has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 11.5±0.2, 13.3±0.2, 15.4±0.2, 16.4±0.2, 17.1±0.2, 18.6±0.2, 19.4±0.2, 20.4±0.2, 21.6±0.2, 22.4±0.2, 24.0±0.2, 26.8±0.2, and 29.0±0.2 degrees 2θ. In other such embodiments, the pattern B polymorph has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 11.5±0.2, 13.3±0.2, 15.4±0.2, 16.4±0.2, 17.1±0.2, 18.6±0.2, 19.4±0.2, 20.4±0.2, 21.6±0.2, 22.4±0.2, 24.0±0.2, 26.8±0.2, and 29.0±0.2 degrees 2θ.

30 [0341] In some embodiments, the pattern B polymorph has an X-ray powder diffraction pattern substantially as shown in Figure 13. The 2θ values for the peaks in Figure 13 (and their intensities) are as follows: 11.52(71), 13.30(87), 15.37 (100), 16.42 (60), 17.13 (69), 18.60 (97), 19.37 (56), 20.40 (62), 21.55 (55), 22.41 (39), 23.99 (33), 26.81 (31), and 28.98 (50).

35 [0342] This invention relates, in part, to pattern C crystalline (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

40 [0343] In some embodiments, the pattern C polymorph has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 7.7±0.2, 10.1±0.2, 10.0±0.2, 12.0 ±0.2, 13.4±0.2, 16.2±0.2, 19.4±0.2, 20.5±0.2, 21.4±0.2, 22.0±0.2, 22.6±0.2, 24.3±0.2, and 27.6±0.2 degrees 2θ. In some such embodiments, the pattern C polymorph has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 7.7±0.2, 10.1±0.2, 10.6±0.2, 12.0 ±0.2, 13.4±0.2, 16.2±0.2, 19.4±0.2, 20.5±0.2, 21.4±0.2, 22.0±0.2, 22.6±0.2, 24.3±0.2, and 27.6±0.2 degrees 2θ. In other such embodiments, the pattern C polymorph has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 7.7±0.2, 10.1±0.2, 10.6±0.2, 12.0 ±0.2, 13.4±0.2, 16.2±0.2, 19.4±0.2, 20.5±0.2, 21.4±0.2, 22.0±0.2, 22.6±0.2, 24.3±0.2, and 27.6±0.2 degrees 2θ.

45 [0344] In some embodiments, the pattern C polymorph has an X-ray powder diffraction pattern substantially as shown in Figure 14. The 2θ values for the peaks in Figure 14 (and their intensities) are as follows: 7.69 (27), 10.13 (27), 10.64 (49), 12.01 (31), 13.39 (33), 16.25 (91), 19.44 (46), 20.49 (100), 21.40 (35), 22.03 (37), 22.60 (30), 24.32 (23), and 27.55 (27).

50 [0345] This invention relates, in part, to pattern D crystalline (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

55 [0346] In some embodiments, the pattern D polymorph has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 5.8±0.2, 10.7±0.2, 11.2±0.2, 15.2±0.2, 16.1±0.2, 16.9±0.2, 19.9±0.2, 22.1±0.2, 24.7±0.2, and 26.0±0.2 degrees 2θ. In some such embodiments, the pattern D polymorph has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 5.8±0.2, 10.7±0.2, 11.2±0.2, 15.2±0.2, 16.1±0.2, 16.9±0.2, 19.9±0.2, 22.1±0.2, 24.7±0.2, and 26.0±0.2 degrees 2θ. In other such embodiments, the pattern D polymorph has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 5.8±0.2, 10.7±0.2, 11.2±0.2, 15.2±0.2, 16.1±0.2, 16.9±0.2, 19.9±0.2, 22.1±0.2, 24.7±0.2, and 26.0±0.2 degrees 2θ.

[0347] In some embodiments, the pattern D polymorph has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $5.8\pm 0.2$ ,  $10.7\pm 0.2$ ,  $11.2\pm 0.2$ ,  $15.2\pm 0.2$ ,  $16.1\pm 0.2$ ,  $16.9\pm 0.2$ ,  $17.1\pm 0.2$ ,  $19.9\pm 0.2$ ,  $20.1\pm 0.2$ ,  $22.1\pm 0.2$ ,  $24.7\pm 0.2$ , and  $26.0\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the pattern D polymorph has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $5.9\pm 0.2$ ,  $10.7\pm 0.2$ ,  $11.2\pm 0.2$ ,  $15.2\pm 0.2$ ,  $16.1\pm 0.2$ ,  $16.9\pm 0.2$ ,  $17.1\pm 0.2$ ,  $19.9\pm 0.2$ ,  $20.1\pm 0.2$ ,  $22.1\pm 0.2$ ,  $24.7\pm 0.2$ , and  $26.0\pm 0.2$  degrees  $2\theta$ . In other such embodiments, the pattern D polymorph has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $5.8\pm 0.2$ ,  $10.7\pm 0.2$ ,  $11.2\pm 0.2$ ,  $15.2\pm 0.2$ ,  $16.1\pm 0.2$ ,  $16.9\pm 0.2$ ,  $17.1\pm 0.2$ ,  $19.9\pm 0.2$ ,  $20.1\pm 0.2$ ,  $22.1\pm 0.2$ ,  $24.7\pm 0.2$ , and  $26.0\pm 0.2$  degrees  $2\theta$ .

[0348] In some embodiments, the pattern D polymorph has an X-ray powder diffraction pattern substantially as shown in Figure 15. The  $2\theta$  values for the peaks in Figure 15 (and their intensities) are as follows: 5.81 (24), 10.70 (91), 11.23 (60), 15.17 (28), 16.10 (48), 16.89 (100), 17.10 (42), 19.88 (81), 20.12 (100), 22.12 (59), 24.72 (37), and 25.91 (24).

[0349] This invention also relates, in part, to a process for preparing pattern B, C, and D polymorphs by heating pattern A polymorph to about 160, about 225, and about 268°C, respectively using DSC.

#### 15 **G6B. IB-L1-1.1 Hydrates.**

[0350] This invention also relates, in part, to hydrates of compound **IB-L1-1.1**, namely to hydrates A, B, C, D, and E discussed below.

[0351] This invention relates, in part, to a pattern A (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide hydrate.

[0352] In some embodiments, the pattern A hydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $5.1\pm 0.2$ ,  $7.9\pm 0.2$ ,  $9.5\pm 0.2$ ,  $10.3\pm 0.2$ ,  $13.7\pm 0.2$ ,  $16.5\pm 0.2$ ,  $17.1\pm 0.2$ ,  $17.5\pm 0.2$ ,  $18.8\pm 0.2$ ,  $19.2\pm 0.2$ ,  $20.7\pm 0.2$ ,  $21.3\pm 0.2$ ,  $21.6\pm 0.2$ ,  $25.8\pm 0.2$ ,  $26.8\pm 0.2$ , and  $28.4\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the pattern A hydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $5.1\pm 0.2$ ,  $7.9\pm 0.2$ ,  $9.5\pm 0.2$ ,  $10.3\pm 0.2$ ,  $13.7\pm 0.2$ ,  $16.5\pm 0.2$ ,  $17.1\pm 0.2$ ,  $17.5\pm 0.2$ ,  $18.8\pm 0.2$ ,  $19.2\pm 0.2$ ,  $20.7\pm 0.2$ ,  $21.3\pm 0.2$ ,  $21.6\pm 0.2$ ,  $25.8\pm 0.2$ ,  $26.8\pm 0.2$ , and  $28.4\pm 0.2$  degrees  $2\theta$ . In other such embodiments, the pattern A hydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $5.1\pm 0.2$ ,  $7.9\pm 0.2$ ,  $9.5\pm 0.2$ ,  $10.3\pm 0.2$ ,  $13.7\pm 0.2$ ,  $16.5\pm 0.2$ ,  $17.1\pm 0.2$ ,  $17.5\pm 0.2$ ,  $18.8\pm 0.2$ ,  $19.2\pm 0.2$ ,  $20.7\pm 0.2$ ,  $21.3\pm 0.2$ ,  $21.6\pm 0.2$ ,  $25.8\pm 0.2$ ,  $26.8\pm 0.2$ , and  $28.4\pm 0.2$  degrees  $2\theta$ .

[0353] In some embodiments, the pattern A hydrate has an X-ray powder diffraction pattern substantially as shown in Figure 16. The  $2\theta$  values for the peaks in Figure 16 (and their intensities) are as follows: 5.13 (13), 7.87 (80), 9.45 (100), 10.29 (60), 13.7 (28), 16.54 (30), 17.07 (17), 17.51 (40), 18.80 (99), 19.18 (74), 20.69 (21), 21.25 (21), 21.63 (23), 25.85 (32), 26.81 (20), and 28.35 (27).

[0354] This invention also relates, in part, to a process for preparing the pattern A hydrate by suspending pattern A polymorph (discussed above) in ethyl acetate. The recovered pattern A hydrate contain ~1 water molecules per molecule of compound **IB-L1-1.1**.

[0355] This invention also relates, in part, to a pattern B (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide hydrate.

[0356] In some embodiments, the pattern B hydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $6.3\pm 0.2$ ,  $7.7\pm 0.2$ ,  $10.4\pm 0.2$ ,  $12.7\pm 0.2$ ,  $13.3\pm 0.2$ ,  $14.9\pm 0.2$ ,  $15.4\pm 0.2$ ,  $16.4\pm 0.2$ ,  $18.6\pm 0.2$ ,  $18.9\pm 0.2$ ,  $19.4\pm 0.2$ ,  $22.5\pm 0.2$ ,  $23.5\pm 0.2$ ,  $24.0\pm 0.2$ ,  $26.8\pm 0.2$ , and  $29.0\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the pattern B hydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $6.3\pm 0.2$ ,  $7.7\pm 0.2$ ,  $10.4\pm 0.2$ ,  $12.7\pm 0.2$ ,  $13.3\pm 0.2$ ,  $14.9\pm 0.2$ ,  $15.4\pm 0.2$ ,  $16.4\pm 0.2$ ,  $18.6\pm 0.2$ ,  $18.9\pm 0.2$ ,  $19.4\pm 0.2$ ,  $22.5\pm 0.2$ ,  $23.5\pm 0.2$ ,  $24.0\pm 0.2$ ,  $26.8\pm 0.2$ , and  $29.0\pm 0.2$  degrees  $2\theta$ . In other such embodiments, the pattern B hydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $6.3\pm 0.2$ ,  $7.7\pm 0.2$ ,  $10.4\pm 0.2$ ,  $12.7\pm 0.2$ ,  $13.3\pm 0.2$ ,  $14.9\pm 0.2$ ,  $15.4\pm 0.2$ ,  $16.4\pm 0.2$ ,  $18.6\pm 0.2$ ,  $18.9\pm 0.2$ ,  $19.4\pm 0.2$ ,  $22.5\pm 0.2$ ,  $23.5\pm 0.2$ ,  $24.0\pm 0.2$ ,  $26.8\pm 0.2$ , and  $29.0\pm 0.2$  degrees  $2\theta$ .

[0357] In some embodiments, the pattern B hydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $6.3\pm 0.2$ ,  $7.7\pm 0.2$ ,  $10.4\pm 0.2$ ,  $12.7\pm 0.2$ ,  $13.3\pm 0.2$ ,  $13.5\pm 0.2$ ,  $14.9\pm 0.2$ ,  $15.4\pm 0.2$ ,  $16.4\pm 0.2$ ,  $18.5\pm 0.2$ ,  $18.6\pm 0.2$ ,  $18.9\pm 0.2$ ,  $19.4\pm 0.2$ ,  $22.5\pm 0.2$ ,  $23.5\pm 0.2$ ,  $24.0\pm 0.2$ ,  $26.8\pm 0.2$ , and  $29.0\pm 0.2$  degrees  $2\theta$ . In some such embodiments, the pattern B hydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $6.3\pm 0.2$ ,  $7.7\pm 0.2$ ,  $10.4\pm 0.2$ ,  $12.7\pm 0.2$ ,  $13.3\pm 0.2$ ,  $13.5\pm 0.2$ ,  $14.9\pm 0.2$ ,  $15.4\pm 0.2$ ,  $16.4\pm 0.2$ ,  $18.5\pm 0.2$ ,  $18.6\pm 0.2$ ,  $8.9\pm 0.2$ ,  $19.4\pm 0.2$ ,  $22.5\pm 0.2$ ,  $23.5\pm 0.2$ ,  $24.0\pm 0.2$ ,  $26.8\pm 0.2$ , and  $29.0\pm 0.2$  degrees  $2\theta$ . In other such embodiments, the pattern B hydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $6.3\pm 0.2$ ,  $7.7\pm 0.2$ ,  $10.4\pm 0.2$ ,  $12.7\pm 0.2$ ,  $13.3\pm 0.2$ ,  $13.5\pm 0.2$ ,  $14.9\pm 0.2$ ,  $15.4\pm 0.2$ ,  $16.4\pm 0.2$ ,  $18.5\pm 0.2$ ,  $18.6\pm 0.2$ ,  $18.9\pm 0.2$ ,  $19.4\pm 0.2$ ,  $22.5\pm 0.2$ ,  $23.5\pm 0.2$ ,  $24.0\pm 0.2$ ,  $26.8\pm 0.2$ , and  $29.0\pm 0.2$  degrees  $2\theta$ .

[0358] In some embodiments, the pattern B hydrate has an X-ray powder diffraction pattern substantially as shown

in Figure 18. The  $2\theta$  values for the peaks in Figure 18 (and their intensities) are as follows: 6.31 (7), 7.72 (14), 10.45 (24), 12.67 (26), 13.30 (88), 13.50 (44), 14.89 (70), 15.40 (100), 16.43 (43), 18.46 (47), 18.63 (86), 18.91 (26), 19.42 (33), 22.52 (47), 23.52 (44), 24.02 (20), 26.82 (40), and 28.97 (49).

[0359] This invention also relates, in part, to a process for preparing the pattern B hydrate by suspending pattern A polymorph (discussed above) in acetonitrile/water (9/1 v/v). The recovered pattern B hydrate contains  $\sim 0.7$  water molecules per molecule of compound **IB-L1-1.1**.

[0360] This invention also relates, in part, to a pattern C (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide hydrate.

[0361] In some embodiments, the pattern C hydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $10.5 \pm 0.2$ ,  $13.3 \pm 0.2$ ,  $14.9 \pm 0.2$ ,  $15.4 \pm 0.2$ ,  $16.4 \pm 0.2$ ,  $18.6 \pm 0.2$ ,  $19.0 \pm 0.2$ ,  $19.4 \pm 0.2$ ,  $22.5 \pm 0.2$ ,  $23.5 \pm 0.2$ ,  $26.9 \pm 0.2$ , and  $29.0 \pm 0.2$  degrees  $2\theta$ . In some such embodiments, the pattern C hydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $10.5 \pm 0.2$ ,  $13.3 \pm 0.2$ ,  $14.9 \pm 0.2$ ,  $15.4 \pm 0.2$ ,  $16.4 \pm 0.2$ ,  $18.6 \pm 0.2$ ,  $19.0 \pm 0.2$ ,  $19.4 \pm 0.2$ ,  $22.5 \pm 0.2$ ,  $23.5 \pm 0.2$ ,  $26.9 \pm 0.2$ , and  $29.0 \pm 0.2$  degrees  $2\theta$ . In other such embodiments, the pattern C hydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $10.5 \pm 0.2$ ,  $13.3 \pm 0.2$ ,  $14.9 \pm 0.2$ ,  $15.4 \pm 0.2$ ,  $16.4 \pm 0.2$ ,  $18.6 \pm 0.2$ ,  $19.0 \pm 0.2$ ,  $19.4 \pm 0.2$ ,  $22.5 \pm 0.2$ ,  $23.5 \pm 0.2$ ,  $26.9 \pm 0.2$ , and  $29.0 \pm 0.2$  degrees  $2\theta$ .

[0362] In some embodiments, the pattern C hydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $10.5 \pm 0.2$ ,  $13.3 \pm 0.2$ ,  $13.5 \pm 0.2$ ,  $14.9 \pm 0.2$ ,  $15.4 \pm 0.2$ ,  $16.4 \pm 0.2$ ,  $18.6 \pm 0.2$ ,  $19.0 \pm 0.2$ ,  $19.4 \pm 0.2$ ,  $22.5 \pm 0.2$ ,  $23.5 \pm 0.2$ ,  $26.9 \pm 0.2$ , and  $29.0 \pm 0.2$  degrees  $2\theta$ . In some such embodiments, the pattern C hydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $10.5 \pm 0.2$ ,  $13.3 \pm 0.2$ ,  $13.5 \pm 0.2$ ,  $14.9 \pm 0.2$ ,  $15.4 \pm 0.2$ ,  $16.4 \pm 0.2$ ,  $18.6 \pm 0.2$ ,  $19.0 \pm 0.2$ ,  $19.4 \pm 0.2$ ,  $22.5 \pm 0.2$ ,  $23.5 \pm 0.2$ ,  $26.9 \pm 0.2$ , and  $29.0 \pm 0.2$  degrees  $2\theta$ . In other such embodiments, the pattern C hydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $10.5 \pm 0.2$ ,  $13.3 \pm 0.2$ ,  $13.5 \pm 0.2$ ,  $14.9 \pm 0.2$ ,  $15.4 \pm 0.2$ ,  $16.4 \pm 0.2$ ,  $18.6 \pm 0.2$ ,  $19.0 \pm 0.2$ ,  $19.4 \pm 0.2$ ,  $22.5 \pm 0.2$ ,  $23.5 \pm 0.2$ ,  $26.9 \pm 0.2$ , and  $29.0 \pm 0.2$  degrees  $2\theta$ .

[0363] In some embodiments, the pattern C hydrate has an X-ray powder diffraction pattern substantially as shown in Figure 20. The  $2\theta$  values for the peaks in Figure 20 (and their intensities) are as follows: 10.47 (21), 13.31 (56), 13.49 (31), 14.91 (28), 15.40 (86), 16.43 (48), 18.61 (100), 18.96 (20), 19.44 (19), 22.55 (26), 23.54 (39), 26.84 (29), and 28.99 (54).

[0364] This invention also relates, in part, to a process for preparing the pattern C hydrate by suspending pattern A polymorph (discussed above) in water. The recovered pattern C hydrate contains  $\sim 1$  water molecules per molecule of compound **IB-L1-1.1**.

[0365] This invention also relates, in part, to a pattern D (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide hydrate.

[0366] The crystallographic unit cell parameters of the pattern D hydrate salt have been determined to be as follows: a is  $17.8 \text{ \AA}$ , b is  $9.6 \text{ \AA}$ , and c is  $27.0 \text{ \AA}$  (more precisely, a is  $17.783(2) \text{ \AA}$ , b is  $9.5651(12) \text{ \AA}$ , and c is  $27.014(4) \text{ \AA}$ ); the cell angle is:  $\beta = 93.3^\circ$  (more precisely,  $\beta$  is  $93.256(2)^\circ$ ); and the cell volume is  $4588 \text{ \AA}^3$  (more precisely,  $4587.5(10) \text{ \AA}^3$ ). The salt crystallizes in the C2/c space group.

[0367] In some embodiments, the pattern D hydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $6.6 \pm 0.2$ ,  $10.0 \pm 0.2$ ,  $10.5 \pm 0.2$ ,  $11.1 \pm 0.2$ ,  $11.6 \pm 0.2$ ,  $12.2 \pm 0.2$ ,  $14.2 \pm 0.2$ ,  $16.6 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $17.7 \pm 0.2$ ,  $18.5 \pm 0.2$ ,  $18.8 \pm 0.2$ ,  $19.3 \pm 0.2$ ,  $21.4 \pm 0.2$ ,  $22.7 \pm 0.2$ ,  $23.1 \pm 0.2$ ,  $23.6 \pm 0.2$ ,  $24.6 \pm 0.2$ ,  $25.2 \pm 0.2$ ,  $27.2 \pm 0.2$ ,  $29.1 \pm 0.2$ , and  $31.0 \pm 0.2$  degrees  $2\theta$ . In some such embodiments, the pattern D hydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $6.6 \pm 0.2$ ,  $10.0 \pm 0.2$ ,  $10.5 \pm 0.2$ ,  $11.1 \pm 0.2$ ,  $11.6 \pm 0.2$ ,  $12.2 \pm 0.2$ ,  $14.2 \pm 0.2$ ,  $16.6 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $17.7 \pm 0.2$ ,  $18.5 \pm 0.2$ ,  $18.8 \pm 0.2$ ,  $19.3 \pm 0.2$ ,  $21.4 \pm 0.2$ ,  $22.7 \pm 0.2$ ,  $23.1 \pm 0.2$ ,  $23.6 \pm 0.2$ ,  $24.6 \pm 0.2$ ,  $25.2 \pm 0.2$ ,  $27.2 \pm 0.2$ ,  $29.1 \pm 0.2$ , and  $31.0 \pm 0.2$  degrees  $2\theta$ . In other such embodiments, the pattern D hydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $6.6 \pm 0.2$ ,  $10.0 \pm 0.2$ ,  $10.5 \pm 0.2$ ,  $11.1 \pm 0.2$ ,  $11.6 \pm 0.2$ ,  $12.2 \pm 0.2$ ,  $14.2 \pm 0.2$ ,  $16.6 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $17.7 \pm 0.2$ ,  $18.5 \pm 0.2$ ,  $18.8 \pm 0.2$ ,  $19.3 \pm 0.2$ ,  $21.4 \pm 0.2$ ,  $22.7 \pm 0.2$ ,  $23.1 \pm 0.2$ ,  $23.6 \pm 0.2$ ,  $24.6 \pm 0.2$ ,  $25.2 \pm 0.2$ ,  $27.2 \pm 0.2$ ,  $29.1 \pm 0.2$ , and  $31.0 \pm 0.2$  degrees  $2\theta$ .

[0368] In some embodiments, the pattern D hydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of  $6.6 \pm 0.2$ ,  $10.0 \pm 0.2$ ,  $10.5 \pm 0.2$ ,  $11.1 \pm 0.2$ ,  $11.6 \pm 0.2$ ,  $12.2 \pm 0.2$ ,  $12.5 \pm 0.2$ ,  $14.2 \pm 0.2$ ,  $16.6 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $17.7 \pm 0.2$ ,  $18.5 \pm 0.2$ ,  $18.8 \pm 0.2$ ,  $19.3 \pm 0.2$ ,  $21.4 \pm 0.2$ ,  $22.7 \pm 0.2$ ,  $22.8 \pm 0.2$ ,  $23.1 \pm 0.2$ ,  $23.6 \pm 0.2$ ,  $24.6 \pm 0.2$ ,  $24.9 \pm 0.2$ ,  $25.2 \pm 0.2$ ,  $27.2 \pm 0.2$ ,  $29.1 \pm 0.2$ , and  $31.0 \pm 0.2$  degrees  $2\theta$ . In some such embodiments, the pattern D hydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of  $6.6 \pm 0.2$ ,  $10.0 \pm 0.2$ ,  $10.5 \pm 0.2$ ,  $11.1 \pm 0.2$ ,  $11.6 \pm 0.2$ ,  $12.2 \pm 0.2$ ,  $12.5 \pm 0.2$ ,  $14.2 \pm 0.2$ ,  $16.6 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $17.7 \pm 0.2$ ,  $18.5 \pm 0.2$ ,  $18.8 \pm 0.2$ ,  $19.3 \pm 0.2$ ,  $21.4 \pm 0.2$ ,  $22.7 \pm 0.2$ ,  $22.8 \pm 0.2$ ,  $23.1 \pm 0.2$ ,  $23.6 \pm 0.2$ ,  $24.6 \pm 0.2$ ,  $24.9 \pm 0.2$ ,  $25.2 \pm 0.2$ ,  $27.2 \pm 0.2$ ,  $29.1 \pm 0.2$ , and  $31.0 \pm 0.2$  degrees  $2\theta$ . In other such embodiments, the pattern D hydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of  $6.6 \pm 0.2$ ,  $10.0 \pm 0.2$ ,  $10.5 \pm 0.2$ ,  $11.1 \pm 0.2$ ,  $11.6 \pm 0.2$ ,  $12.2 \pm 0.2$ ,  $12.5 \pm 0.2$ ,  $14.2 \pm 0.2$ ,  $16.6 \pm 0.2$ ,  $17.1 \pm 0.2$ ,  $17.7 \pm 0.2$ ,  $18.5 \pm 0.2$ ,  $18.8 \pm 0.2$ ,  $19.3 \pm 0.2$ ,  $21.4 \pm 0.2$ ,  $22.7 \pm 0.2$ ,  $22.8 \pm 0.2$ ,  $23.1 \pm 0.2$ ,  $23.6 \pm 0.2$ ,  $24.6 \pm 0.2$ ,  $24.9 \pm 0.2$ ,  $25.2 \pm 0.2$ ,  $27.2 \pm 0.2$ ,  $29.1 \pm 0.2$ , and  $31.0 \pm 0.2$  degrees  $2\theta$ .

19.3±0.2, 21.4±0.2, 22.7±0.2, 22.8±0.2, 23.1±0.2, 23.6±0.2, 24.6±0.2, 24.9±0.2, 25.2±0.2, 27.2±0.2, 29.1±0.2, and 31.0±0.2 degrees 2θ.

[0369] In some embodiments, the pattern D hydrate has an X-ray powder diffraction pattern substantially as shown in Figure 22. The 2θ values for the peaks in Figure 22 (and their intensities) are as follows: 6.55 (10), 9.96 (12), 10.51 (37), 11.09 (31), 11.62 (100), 12.24 (44), 12.54 (40), 14.22 (15), 16.62 (68), 17.07 (22), 17.77 (21), 18.52 (82), 18.84 (47), 19.30 (63), 21.45 (34), 22.67 (30), 22.80 (34), 23.08 (20), 23.57 (58), 24.63 (73), 24.88 (26), 25.24 (21), 27.23 (36), 29.06 (41), and 31.04 (21).

[0370] This invention also relates, in part, to a process for preparing the pattern D hydrate. It was prepared by suspending pattern A polymorph (discussed above) in ethanol. Alternatively, it was prepared by suspending compound **IB-L1-1.1** (103.03mg) in 400ul THF while heated to about 55°C. Aqueous NaOH (1M, 264ul, 1.2 molar equivalent) was added. The solid dissolved completely to yield a clear solution. Ethanol (1.6ml) was added to the solution. The solution was allowed to cool naturally to ambient temperatures. Crystals were formed during the slow evaporation process. Although it appears that the lattice can accommodate as much as 0.5 water molecules per molecule of compound **IB-L1-1.1**, the recovered pattern D hydrate contained ~0.2 water molecules per molecule of compound **IB-L1-1.1**.

[0371] This invention also relates, in part, to a pattern E (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide hydrate.

[0372] The crystallographic unit cell parameters of the pattern E hydrate crystalline disodium salt have been determined to be as follows: a is 9.5Å, b is 14.5Å, and c is 17.3Å (more precisely, a is 9.462(2)Å, b is 14.462(3)Å, and c is 17.281(4)Å); the cell angles are: α - 84.9°, β - 80-8°, and γ - 81.8° (more precisely, α is 84.863(4)°, β is 80.760(4)°, and γ is 81.751(4)°; and the cell volume is 2304Å<sup>3</sup> (more precisely, 2304.4(9)Å<sup>3</sup>). The salt crystallizes in the P-1 space group.

[0373] In some embodiments, the pattern E hydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 6.2±0.2, 7.8±0.2, 10.2±0.2, 10.7±0.2, 12.1±0.2, 16.3±0.2, 19.7±0.2, 20.9±0.2, 21.8±0.2, 24.5±0.2, and 28.0±0.2 degrees 2θ. In some such embodiments, the pattern E hydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 6.2±0.2, 7.8±0.2, 10.2±0.2, 10.7±0.2, 12.1±0.2, 16.3±0.2, 19.7±0.2, 20.9±0.2, 21.8±0.2, 24.5±0.2, and 28.0±0.2 degrees 2θ. In other such embodiments, the pattern E hydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 6.2±0.2, 7.8±0.2, 10.2±0.2, 10.7±0.2, 12.1±0.2, 16.3±0.2, 19.7±0.2, 20.9±0.2, 21.8±0.2, 24.5±0.2, and 28.0±0.2 degrees 2θ.

[0374] In some embodiments, the pattern E hydrate has an X-ray powder diffraction pattern comprising one or more peaks selected from the group consisting of 6.2±0.2, 7.8±0.2, 10.2±0.2, 10.4±0.2, 10.7±0.2, 12.1±0.2, 16.3±0.2, 19.7±0.2, 20.9±0.2, 21.8±0.2, 24.5±0.2, and 28.0±0.2 degrees 2θ. In some such embodiments, the pattern E hydrate has an X-ray powder diffraction pattern comprising three or more peaks selected from the group consisting of 6.2±0.2, 7.8±0.2, 10.2±0.2, 10.4±0.2, 10.7±0.2, 12.1±0.2, 16.3±0.2, 19.7±0.2, 20.9±0.2, 21.8±0.2, 24.5±0.2, and 28.0±0.2 degrees 2θ. In other such embodiments, the pattern E hydrate has an X-ray powder diffraction pattern comprising five or more peaks selected from the group consisting of 6.2±0.2, 7.8±0.2, 10.2±0.2, 10.4±0.2, 10.7±0.2, 12.1±0.2, 16.3±0.2, 19.7±0.2, 20.9±0.2, 21.8±0.2, 24.5±0.2, and 28.0±0.2 degrees 2θ.

[0375] In some embodiments, the pattern E hydrate has an X-ray powder diffraction pattern substantially as shown in Figure 23. The 2θ values for the peaks in Figure 23 (and their intensities) are as follows: 6.19 (6), 7.81 (18), 10.17 (13), 10.40 (14), 10.68 (39), 12.06 (20), 16.29 (78), 19.72 (32), 20.88 (100), 21.77 (27), 24.52 (25), and 28.01 (27).

[0376] This invention also relates, in part, to a process for preparing the pattern E hydrate. It was prepared by suspending compound **IB-L1-1.1** (56.76mg) in 200ul THF while heated. Aqueous NaOH (1M, 146ul, 1.2 molar equivalent) was added, which yielded a clear solution. Ethanol (800ul) was added to the solution. The solution was allowed to cool naturally to ambient temperatures. Crystals were formed during the slow evaporation process. Although it appears that the lattice can accommodate as much as one water molecule per molecule of compound **IB-L1-1.1**, the recovered pattern D hydrate contained ~0.25 water molecules per molecule of compound **IB-L1-1.1**.

#### H. Compositions.

[0377] This invention also is directed, in part, to compositions comprising one or more compounds and/or salts of the invention (including the crystalline compounds and salts discussed in section G above). In some embodiments, the compositions comprise one or more substantially phase pure crystalline forms (compounds/salts/solvates/hydrates) discussed in section G above. The compositions can be pharmaceutical compositions.

[0378] In some embodiments, the compositions further comprise one or more additional therapeutic agents. Such therapeutic agents can, but need not be, additional HCV inhibitors.

[0379] The preferred composition depends on the method of administration, and typically comprises one or more conventional pharmaceutically acceptable carriers, adjuvants, and/or vehicles (together referred to as "excipients"). Formulation of drugs is generally discussed in, for example, Hoover, J., Remington's Pharmaceutical Sciences (Mack Publishing Co., 1975) and Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems (Lippincott Williams &

Wilkins, 2005).

**[0380]** Solid dosage forms for oral administration include, for example, capsules, tablets, pills, powders, and granules. In such solid dosage forms, the compounds or salts are ordinarily combined with one or more excipients. If administered per os, the compounds or salts can be mixed with, for example, lactose, sucrose, starch powder, cellulose esters of

alkanoic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets can contain a controlled-release formulation, as can be provided in, for example, a dispersion of the compound or salt in hydroxypropylmethyl cellulose. In the case of capsules, tablets, and pills, the dosage forms also can comprise buffering agents, such as sodium citrate, or magnesium or calcium carbonate or bicarbonate.

Tablets and pills additionally can be prepared with enteric coatings.

**[0381]** Liquid dosage forms for oral administration include, for example, pharmaceutically acceptable emulsions (including both oil-in-water and water-in-oil emulsions), solutions (including both aqueous and non-aqueous solutions), suspensions (including both aqueous and non-aqueous suspensions), syrups, and elixirs containing inert diluents commonly used in the art (e.g., water). Such compositions also can comprise, for example, wetting, emulsifying, suspending, flavoring (e.g., sweetening), and/or perfuming agents.

**[0382]** Parenteral administration includes subcutaneous injections, intravenous injections, intramuscular injections, intrasternal injections, and infusion. Injectable preparations (e.g., sterile injectable aqueous or oleaginous suspensions) can be formulated according to the known art using suitable dispersing, wetting agents, and/or suspending agents. Acceptable vehicles and solvents include, for example, water, 1,3-butanediol, Ringer's solution, isotonic sodium chloride solution, bland fixed oils (e.g., synthetic mono- or diglycerides), fatty acids (e.g., oleic acid), dimethyl acetamide, surfactants (e.g., ionic and non-ionic detergents), and/or polyethylene glycols.

**[0383]** Formulations for parenteral administration may, for example, be prepared from sterile powders or granules having one or more of the excipients mentioned for use in the formulations for oral administration. A compound or salt of the invention can be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. The pH may be adjusted, if necessary, with a suitable acid, base, or buffer.

**[0384]** Suppositories for rectal administration can be prepared by, for example, mixing a compound or salt of the invention with a suitable nonirritating excipient that is solid at ordinary temperatures, but liquid at the rectal temperature, and will therefore melt in the rectum to release the drug. Suitable excipients include, for example, cocoa butter; synthetic mono-, di-, or triglycerides, fatty acids, and/or polyethylene glycols.

**[0385]** Topical administration includes the use of transdermal administration, such as transdermal patches or iontophoresis devices.

**[0386]** Other excipients and modes of administration known in the pharmaceutical art also may be used.

**[0387]** Applicants have discovered that some **I-L1** compounds in which **R<sup>6</sup>** and the phenyluracil are in trans-position relative to the double bond, when in solution, tend to convert into the corresponding cis-isomer upon exposure to light; thus, it may be desirable to store such solutions under conditions that reduce exposure to light (e.g., in an amber bottle or in a dark place).

**[0388]** The preferred total daily dose of the compound or salt (administered in single or divided doses) is typically from about 0.001 to about 100mg/kg, more preferably from about 0.001 to about 30mg/kg, and even more preferably from about 0.01 to about 10mg/kg (i.e., mg of the compound or salt per kg body weight). Dosage unit compositions can contain such amounts or submultiples thereof to make up the daily dose. In many instances, the administration of the compound or salt will be repeated a plurality of times. Multiple doses per day typically may be used to increase the total daily dose, if desired.

**[0389]** Factors affecting the preferred dosage regimen include the type, age, weight, sex, diet, and condition of the patient; the severity of the pathological condition; the severity of the pathological condition; the route of administration; pharmacological considerations, such as the activity, efficacy, pharmacokinetic, and toxicology profiles of the particular compound or salt used; whether a drug delivery system is utilized; and whether the compound or salt is administered as part of a drug combination. Thus, the dosage regimen actually employed can vary widely, and therefore, can derive from the preferred dosage regimen set forth above.

#### *I. Kits.*

**[0390]** This invention also is directed, in part, to a kit comprising one or more compounds and/or salts of the invention. The kit can optionally contain one or more additional therapeutic agents and/or instructions for, for example, using the kit.

### J. Methods of Use.

5 [0391] This invention also is directed, in part, to a method for inhibiting replication of an RNA virus comprising exposing the virus to one or more compounds and/or salts of this invention *in vitro*. Replication of the RNA virus is inhibited *in vitro*. This invention is also directed, in part, to the compounds and/or salts of the invention for use in inhibiting replication of the RNA virus *in vivo*. In some embodiments, the RNA virus whose replication is being inhibited is a single-stranded, positive sense RNA virus. In some such embodiments, the RNA virus whose replication is being inhibited is a virus from the *Flaviviridae* family. In some such embodiments, the RNA virus whose replication is being inhibited is HCV.

10 [0392] This invention also is directed, in part, to a method for inhibiting HCV RNA polymerase *in vitro*. The method comprises exposing the polymerase with one or more compounds and/or salts of this invention. In some embodiments, HCV RNA polymerase activity is inhibited *in vitro*. This invention is also directed, in part, to the compounds and/or salts of the invention for use in inhibiting HCV RNA polymerase activity *in vivo*.

15 [0393] The term "inhibiting" means reducing the level of RNA virus replication/HCV polymerase activity either *in vitro* or *in vivo*. For example, if a compound/salt of the invention reduces the level of RNA virus replication by at least about 10% compared to the level of RNA virus replication before the virus was exposed to the compound/salt, then the compound/salt inhibits RNA virus replication. In some embodiments, the compound/salt can inhibit RNA virus replication by at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, or at least about 95%.

20 [0394] This invention also is directed, in part, to the compounds and/or salts of the invention for use in method for treating a disease that can be treated by inhibiting HCV RNA polymerase. Thus, this invention also is directed, in part, to the compounds and/or salts of the invention for use in a method for treating hepatitis C in an animal in need of such treatment. These methods comprise administering to the animal one or more compounds and/or salts of the invention, and, optionally, one or more additional therapeutic agents. In some embodiments, a therapeutically effective amount of the compound(s) and/or salt(s) is administered to the animal. "Treating" means ameliorating, suppressing, eradicating, preventing, reducing the risk of, and/or delaying the onset of the disease being treated. Applicants specifically intend that the term "treating" encompass administration of the compounds and/or salts of the invention to an HCV-negative patient that is a candidate for an organ transplant. The methods of treatment are particularly suitable for use with humans, but may be used with other animals, particularly mammals. A "therapeutically-effective amount" or "effective amount" is an amount that will achieve the goal of treating the targeted condition.

25 [0395] In some embodiments, the methods comprise combination therapy, wherein the compound(s) and/or salt(s) of the invention is/are co-administered with a second (or even a third, fourth, etc.) compound, such as, for example, another therapeutic agent used to treat hepatitis C (e.g., interferon or interferon/ribavirin combination, or an HCV inhibitor such as, for example, an HCV polymerase inhibitor or an HCV protease inhibitor). The compound(s) and/or salt(s) of this invention can also be co-administered with therapeutic agents other than therapeutic agents used to treat hepatitis C (e.g., anti-HIV agents). In these co-administration embodiments, the compound(s) and/or salt(s) of the invention and the second, etc. therapeutic agent(s) may be administered in a substantially simultaneous manner (e.g., or within about 5 minutes of each other), in a sequential manner, or both. It is contemplated that such combination therapies may include administering one therapeutic agent multiple times between the administrations of the other. The time period between the administration of each agent may range from a few seconds (or less) to several hours or days, and will depend on, for example, the properties of each composition and active ingredient (e.g., potency, solubility, bioavailability, half-life, and kinetic profile), as well as the condition of the patient. The compound(s) and/or salt(s) of this invention and the second, etc. therapeutic agent may also be administered in a single formulation.

30 [0396] This invention also is directed, in part, to a use of one or more compounds and/or salts of the invention, and, optionally one or more additional therapeutic agents to prepare a medicament. In some embodiments, the medicament is for co-administration with one or more additional therapeutic agents.


35 [0397] In some embodiments, the medicament is for inhibiting replication of an RNA virus.

[0398] In some embodiments, the medicament is for treating hepatitis C.

40 [0399] This invention also is directed, in part, to one or more compounds and/or salts of the invention, and, optionally one or more additional therapeutic agents, for use as a medicament. In some embodiments, the medicament is for inhibiting replication of an RNA virus. In other embodiments, the medicament is for treating hepatitis C.

### K. Intermediate Compounds.

45 [0400] Disclosed are also intermediates that correspond in structure to formula II that can be used to prepare the compounds of formula I (and their salts)(although some intermediates can also be used, just like the compounds of formula I, as HCV inhibitors, and one skilled in the art can determine such ability of the compounds of formula II by utilizing, for example, the methods discussed below):



15 [0401] In formula II:



20  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ , and  $R^5$  are as discussed above for the compounds of formula I; and  $X^2$  is halo.

25 [0402] The various embodiments for




30  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ , and  $R^5$  (as well as their combinations) discussed above apply to the compounds of formula II. As to  $X^2$ , in some embodiments,  $X^2$  is selected from the group consisting of chloro, bromo, and iodo. In other embodiments,  $X^2$  is selected from the group consisting of chloro and bromo. In yet other embodiments,  $X^2$  is selected from the group consisting of chloro and iodo. In yet other embodiments,  $X^2$  is selected from the group consisting of iodo and bromo. In further embodiments,  $X^2$  is fluoro. In yet further embodiments,  $X^2$  is chloro. In yet further embodiments,  $X^2$  is bromo. And in yet further embodiments,  $X^2$  is iodo.

35 [0403] The various embodiments for




40  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , and  $X^2$  discussed above can be combined to form various embodiments of compounds of formula II, and all embodiments of compounds of formula II so formed are within the scope of Applicants' invention. Some exemplary embodiments of the compounds (and salts thereof) of formula II are discussed below.

45 [0404] In some embodiments, the compounds of formula II correspond in structure to formula IIA:



[0405] In other embodiments, the compounds of formula II correspond in structure to formula IIB:



15 [0406] In some embodiments of the compounds of formula II:

**R<sup>1</sup>** is selected from the group consisting of hydrogen, methyl, and nitrogen-protecting group;

**R<sup>2</sup>** is selected from the group consisting of hydrogen and halo;

**R<sup>3</sup>** is selected from the group consisting of hydrogen and halo;

20 **R<sup>4</sup>** is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-carbocyclyl, and 5-6-membered heterocyclyl, wherein:

(a) the C<sub>1</sub>-C<sub>4</sub>-alkyl optionally is substituted with up to three substituents independently selected from the group consisting of halo, oxo, hydroxy, alkyloxy, and trimethylsilyl, and

25 (b) the C<sub>3</sub>-C<sub>6</sub>-carbocyclyl and 5-6-membered heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, halo, and alkylsulfonylamino;

**R<sup>5</sup>** is selected from the group consisting of hydrogen, hydroxy, alkyloxy, and halo; and

**X<sup>2</sup>** is selected from the group consisting of chloro, bromo, and iodo.

30 [0407] In some embodiments of the compounds of formula II:



is a double carbon-carbon bond;

**R<sup>1</sup>** is hydrogen;

**R<sup>2</sup>** is selected from the group consisting of hydrogen and halo;

**R<sup>3</sup>** is hydrogen;

40 **R<sup>4</sup>** is tert-butyl;

**R<sup>5</sup>** is selected from the group consisting of hydrogen, hydroxy, and methoxy; and

**X<sup>2</sup>** is selected from the group consisting of bromo and iodo.

45 [0408] In some embodiments of the compounds of formula II:

**R<sup>1</sup>** is selected from the group consisting of hydrogen and methyl;

**R<sup>2</sup>** is selected from the group consisting of hydrogen and methyl;

**R<sup>3</sup>** is selected from the group consisting of hydrogen and methyl;

**R<sup>4</sup>** is tert-butyl;

50 **R<sup>5</sup>** is selected from the group consisting of hydroxy and methoxy; and

**X<sup>2</sup>** is selected from the group consisting of chloro, bromo, and iodo.

[0409] In some embodiments of the compounds of formula II:


55



is a double carbon-carbon bond;

$R^1$  is hydrogen;  
 $R^2$  is hydrogen;  
 $R^3$  is hydrogen;  
 $R^4$  is tert-butyl;  
 $R^5$  is selected from the group consisting of hydroxy and methoxy; and  
 $X^2$  is selected from the group consisting of chloro, bromo, and iodo.


[0410] In some embodiments, the compound of formula II is selected from the group consisting of



[0411] The discussion below provides instructions for the preparation of intermediate compounds of formula II (and salts thereof).

#### L. Starting Compounds.

[0412] Disclosed are also starting compounds that correspond in structure to formula III that can be used to prepare the compounds of formulas II and I (and their salts):



[0413] In formula III,  $\text{---}^*$ ,  $R^1$ ,  $R^2$ , and  $R^3$  are as discussed above for the compounds of formula I and II. The various embodiments for  $\text{---}^*$ ,  $R^1$ ,  $R^2$ , and  $R^3$  (as well as their combinations) discussed above apply to the compounds of formula III. The various embodiments for  $\text{---}^*$ ,  $R^1$ ,  $R^2$ , and  $R^3$  discussed above can be combined to form various embodiments of compounds of formula III, and all embodiments of compounds of formula III so formed are within the scope of Applicants' invention. Some exemplary embodiments of the compounds (and salts thereof) of formula III are discussed below.

[0414] In some embodiments of the compounds of formula III:

$R^1$  is selected from the group consisting of hydrogen, methyl, and nitrogen-protecting group;  
 $R^2$  is selected from the group consisting of hydrogen and halo; and  
 $R^3$  is selected from the group consisting of hydrogen and halo.

[0415] In some embodiments of the compounds of formula III:

$\text{---}^*$  is a double carbon-carbon bond;  
 $R^1$  is selected from the group consisting of hydrogen;  
 $R^2$  is selected from the group consisting of hydrogen and halo; and


$R^3$  is selected from the group consisting of hydrogen.

[0416] In some embodiments of the compounds of formula III:

$R^1$  is selected from the group consisting of hydrogen and methyl;  
 $R^2$  is selected from the group consisting of hydrogen and methyl; and  
 $R^3$  is selected from the group consisting of hydrogen and methyl.

[0417] In some embodiments, the compound of formula III is uracil.

[0418] Disclosed are also starting compounds that correspond in structure to formula IV that can be used to prepare the compounds of formulas II and I (and their salts):



[0419] In formula IV:

$R^4$ ,  $R^5$ , and  $X^2$  are as discussed above for the compounds of formula I and II; and  
 $X^1$  is halo.

[0420] The various embodiments for  $R^4$ ,  $R^5$ , and  $X^2$  (as well as their combinations) discussed above apply to the compounds of formula IV. As to  $X^1$ , in some embodiments,  $X^1$  is selected from the group consisting of chloro, bromo, and iodo. In other embodiments,  $X^1$  is selected from the group consisting of chloro and bromo. In yet other embodiments,  $X^1$  is selected from the group consisting of chloro and iodo. In yet other embodiments,  $X^1$  is selected from the group consisting of iodo and bromo. In further embodiments,  $X^1$  is fluoro. In yet further embodiments,  $X^1$  is chloro. In yet further embodiments,  $X^1$  is bromo. And in yet further embodiments,  $X^1$  is iodo. As to  $X^1$  and  $X^2$ , in some embodiments,  $X^1$  and  $X^2$  are identical.

[0421] The various embodiments for  $R^4$ ,  $R^5$ ,  $X^1$ , and  $X^2$  discussed above can be combined to form various embodiments of compounds of formula IV, and all embodiments of compounds of formula III so formed are within the scope of Applicants' invention. Some exemplary embodiments of the compounds (and salts thereof) of formula IV are discussed below.

[0422] In some embodiments of the compounds of formula IV:

$R^4$  is selected from the group consisting of  $C_1$ - $C_4$ -alkyl,  $C_3$ - $C_6$ -carbocyclyl, and 5-6-membered heterocyclyl, wherein:

(a) the  $C_1$ - $C_4$ -alkyl optionally is substituted with up to three substituents independently selected from the group consisting of halo, oxo, hydroxy, alkyloxy, and trimethylsilyl, and

(b) the  $C_3$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, halo, and alkylsulfonylamino;

$R^5$  is selected from the group consisting of hydrogen, hydroxy, and alkyloxy;  
 $X^1$  is selected from the group consisting of chloro, bromo, and iodo; and  
 $X^2$  is selected from the group consisting of chloro, bromo, and iodo.

[0423] In some embodiments of the compounds of formula IV:

$R^4$  is selected from the group consisting of tert-butyl;  
 $R^2$  is selected from the group consisting of hydrogen, hydroxy, and methoxy;  
 $X^1$  is selected from the group consisting of bromo and iodo; and  
 $X^2$  is selected from the group consisting of bromo and iodo.


[0424] In some embodiments of the compounds of formula IV:

$R^4$  is selected from the group consisting of tert-butyl;  
 $R^5$  is selected from the group consisting of hydroxy and methoxy;  
 $X^1$  is selected from the group consisting of chloro, bromo, and iodo; and  
 $X^2$  is selected from the group consisting of chloro, bromo, and iodo.

[0425] In some embodiments of the compounds of formula IV:

$R^4$  is tert-butyl;  
 $R^5$  is selected from the group consisting of hydroxy and methoxy;  
 $X^1$  is selected from the group consisting of chloro, bromo, and iodo; and  
 $X^2$  is selected from the group consisting of chloro, bromo, and iodo.


[0426] In some embodiments, the compound of formula IV is selected from the group consisting of



[0427] The discussion below provides instructions for the preparation of starting compounds of formula IV (and salts thereof).

*L. Methods for Preparation.*

[0428] Disclosed is also a process for preparing compounds of formula II. The process comprises reacting a compound of formula III with a compound of formula IV in the presence of (i) copper (I) salt catalyst and (ii) nitrogenous heteroaryl ligand:



[0429] In the above process,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $X^1$ , and  $X^2$  are as discussed above.

[0430] Applicants have discovered that the process generally results in the substitution of the N1 hydrogen of uracil derivative compound III thus resulting in intermediate compound II. When  $X^2$  in intermediate compound II is chloro, bromo, or iodo, then compound II is suitable for subsequent reaction (e.g., Suzuki coupling with an appropriate boronic acid or boronate ester) to provide compound of formula I. In other words, when  $X^2$  in intermediate compound II is chloro, bromo, or iodo, the above process is suitable for preparing compounds of formula I as well.

[0431] In some embodiments, compound III is uracil, and compound IV corresponds in structure to a compound selected from the group consisting of compound IV-I, IV-Br, and IV-Cl, with compounds IV-I and IV-Br typically resulting in better yield than compound IV-Cl.


[0432] Suitable Cu(I) catalysts include, for example, CuI, CuBr, CuCl, Cu<sub>2</sub>O, and CH<sub>3</sub>C(O)OCu. In some embodiments, the catalyst is selected from the group consisting of CuI and CuBr. In some such embodiments, the catalyst is CuI. In other such embodiments, the catalyst is CuBr.

**[0433]** In some embodiments, the process is conducted in the presence of a base. In some such embodiments, the base is an inorganic base. Suitable inorganic bases include, for example, potassium, sodium, and cesium salts (e.g.,  $K_2CO_3$ ,  $K_3PO_4$ ,  $Cs_2CO_3$ ,  $Na_2CO_3$ ). In some embodiments, the base is selected from the group consisting of potassium salt and cesium salt. In some such embodiments, the salt is selected from the group consisting of  $K_3PO_4$  and  $Cs_2CO_3$ . In some embodiments, the base comprises a potassium salt. In some such embodiments, the potassium salt is  $K_2CO_3$ . In other such embodiments, the potassium salt is  $K_3PO_4$ . In some embodiments, the base comprises a cesium salt. In some such embodiments, the potassium salt is  $Cs_2CO_3$ .

**[0434]** Typically, the process is conducted in the presence of a solvent. Suitable solvents include, for example, dimethylsulfoxide (DMSO), dimethylformamide (DMF), and acetonitrile (MeCN). In some embodiments, the solvent is DMSO.

**[0435]** Typically, the process is conducted at a temperature of from about 40 to about 130°C.


**[0436]** In some embodiments, the nitrogenous heteroaryl ligand comprises 8-hydroxyquinoline. In other embodiments, the ligand comprises 2-(2-pyridyl)-benzimidazole. In yet other embodiments, the ligand comprises a picolinamide compound corresponding in structure to formula **V**:



**[0437]** In formula **V**,  $R^{11}$ ,  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ , and  $R^{17}$  are independently selected from the group consisting of hydrogen,  $C_{1-4}$ -perfluoroalkyl,  $C_{1-4}$ -alkyloxy,  $C_{1-4}$ -haloalkyl, chloro, or cyano. In some embodiments,  $R^{11}$ ,  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ , and  $R^{17}$  are independently selected from the group consisting of hydrogen, methyl, methoxy, trifluoromethyl, chloro, and cyano. In some embodiments, the ligand of formula **V** comprises N-(4-cyanophenyl)picolinamide. In other embodiments, the ligand of formula **V** comprises N-(2-cyanophenyl)picolinamide.

**[0438]** In some embodiments, the process comprises (a) preparing a compound of formula **IV**; and (b) reacting a compound of formula **III** with a compound of formula **IV** in the presence of (i) copper (I) salt catalyst and (ii) nitrogenous heteroaryl ligand, optionally in the presence of inorganic base.


**[0439]** Compound of formula **IV-I** can be prepared by, for example, converting 2-tert-butylphenol into 2-tert-butyl-4,6-diiodophenol (by, for example, reacting it with NaI and NaOCl), and then converting the 2-tert-butyl-4,6-diiodophenol into 1-tert-butyl-3,5-diiodo-2-methoxybenzene (by, for example, treating it with  $CH_3I$  in the presence of a base, such as, for example, NaOH).



**[0440]** Compound of formula **IV-Br** can be prepared by, for example, converting 2-tert-butylphenol into 2,4-dibromo-6-tert-butylphenol (by, for example, reacting it with 1,3-dibromo-5,5-dimethylimidazolidine-2,4-dione), and then converting the 2,4-dibromo-6-tert-butylphenol into 1,5-dibromo-3-tert-butyl-2-methoxybenzene (by, for example, treating it with  $CH_3I$  in the presence of  $KOtBu$ ).

**[0441]** Additional information about the preparation of compounds of formulas **I** and **II** (and their salts) is provided in the general discussion and/or specific synthesis examples below. In the discussion below,  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $L$ ,  $R^A$ ,  $R^B$ ,  $R^C$ ,  $R^D$ ,  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$ ,  $R^K$ ,  $X^1$ , and  $X^2$  have the meaning discussed above unless otherwise stated.

## SCHEME 1



[0442] Compound (1-1), wherein R<sup>7</sup> is, for example, hydrogen or -CO<sub>2</sub>Me, and R<sup>8</sup> is, for example, hydrogen or t-butyl, may be treated with nitric acid in solvents such as, for example, acetic acid or water in a temperature range of about 0 to about 35°C over about 1 to about 5h to provide compound (1-2). Compound (1-2) may then be reduced using conditions known to those skilled in the art to furnish the corresponding aniline (1-3). Typical conditions for this reduction include using hydrogen at a pressure of about 1 to about 5 atmospheres in the presence of a catalyst such as, for example, palladium or platinum on charcoal in a solvent such as, for example, tetrahydrofuran, ethyl acetate, ethanol, or hexane at or near ambient temperature over a period of about 1 to about 12h. Dependent on the functional groups present, an alternative reduction procedure may be more appropriate such as, for example, using iron powder in the presence of a mild acid such as, for example, ammonium chloride or dilute hydrochloric acid at reflux temperatures in a mixture of solvents containing, for example, methanol, water, and/or tetrahydrofuran over about 1 to about 12h. Another set of reduction conditions includes the use of sodium borohydride in a solvent mixture such as, for example, water and tetrahydrofuran. Yet another set of reduction conditions includes the use of tin(II) chloride in the presence of hydrochloric acid in such solvents as, for example, water and methanol or mixtures thereof.


[0443] Compound (1-2) may be modified prior to reduction. For example, treatment of compound (1-2), wherein R<sup>7</sup> is hydrogen, with iodine monochloride in a mixture of methanol and water at or near ambient temperature over a period of about 8 to about 24h supplies compound (1-4), wherein X<sup>1</sup> is iodine. Alternatively, compound (1-2) can be treated with pyridinium hydrobromide perbromide in a solvent such as, for example, acetic acid at or near ambient temperature over a period of about 2 to about 16h to provide compound (1-4), wherein X<sup>1</sup> is bromine. Modifications may be introduced at the phenol moiety in compound (1-4). For example, the phenol may be alkylated with alkyl halides (e.g., methyl iodide), alkyl sulfates (e.g., methyl sulfate), alkenyl halides (e.g., allyl bromide), alkynyl halides (e.g., propargyl bromide) in the presence of a base such as, for example, potassium carbonate in acetone, sodium hydride in dimethylformamide, or potassium t-butoxide in tetrahydrofuran, at temperatures from about 0 to about 35°C over a period of about 1 to about 24h to provide compound (1-5), wherein R<sup>9</sup> is, for example, alkyl, alkenyl, or alkynyl. Alternatively, alkylation may be achieved by using a reagent such as (trimethylsilyl) diazomethane in solvents such as, for example, methanol or t-butyl methyl ether, or mixtures thereof in a sealed tube at or near room temperature over about 8 to about 24h. Compound (1-5) may subsequently be reduced to compound (1-6) using the iron powder or tin(II) chloride conditions described above. An alternative reduction procedure employs hydrogenation at approximately 1 atmosphere pressure with a catalyst such as 5% platinum on sulfided carbon in a solvent such as methanol. Protection of the resultant aniline of

compound (1-6) with, for example, a t-butyl carbamate can be achieved by treatment with di-tert-butyl dicarbonate in a solvent such as, for example, tetrahydrofuran or dioxane at a temperature of about 50 to about 65 °C for about 1 to about 8h provides compound (1-7).


[0444] Modifications may also occur at the phenol moiety in compound (1-2). One skilled in the art may alkylate the phenol of compound (1-2) using, for example, the conditions described above to obtain compound (1-8). Compound (1-8) is transformed into compound (1-9) using, for example, one or more of the appropriate reduction conditions described above.

[0445] Another modification of the phenol group in compound (1-2) is sulfonylation to furnish compound (1-8), wherein R<sup>9</sup> is alkylsulfonyl, carbocyclisulfonyl, or haloalkylsulfonyl. Such a compound may be prepared by exposing compound (1-2) to sulfonyl chlorides such as, for example, methanesulfonyl chloride, cyclohexanesulfonyl chloride, benzenesulfonyl chloride, or 3-chloropropane sulfonyl chloride in the presence of a base such as, for example, triethylamine, diisopropylethylamine, or pyridine in a solvent such as, for example, dichloromethane at or near ambient temperature for a period of about 1 to about 24h. One skilled in the art can then transform compound (1-8) into compound (1-9) with an appropriate set of reduction conditions.


### SCHEME 2



[0446] Aniline (2-4) can be prepared through use of the Curtius rearrangement. To this end, compound (2-1), wherein R<sup>4</sup> is not amino, can be treated in refluxing thionyl chloride with a catalytic amount of dimethylformamide for about 1 to about 4h to obtain acid chloride (2-2). Treatment with thionyl chloride at the reflux temperature in solvents such as, for example, chloroform or toluene also furnishes compound (2-2). Compound (2-2) can be reacted with an aqueous solution of sodium azide in a solvent such as, for example, acetone over about 1 to about 8h to provide acyl azide (2-3). Compound (2-3) can then undergo a Curtius rearrangement in refluxing solvents such as dioxane or toluene. The intermediate isocyanate is hydrolyzed with an aqueous acid such as dilute hydrochloric acid in a solvent such as dimethoxyethane to provide compound (2-4).


**SCHEME 3**

[0447] Compound (3-1), wherein R<sup>10</sup> is, for example, hydrogen, bromine, iodine, or -CO<sub>2</sub>Me, can be treated with an acrylic acid either neat at or near ambient temperature in a solvent such as, for example, toluene and heated to reflux over a period of about 15 to about 48h to supply compound (3-2). When excess of an acrylic acid is used, compound (3-3) is produced. Compound (3-2) or (3-3) can be treated with urea in a solvent such as, for example, acetic acid at about 100 to about 120°C over about 2 to about 48h to supply compound (3-4).

**SCHEME 4**


[0448] Compound (4-2) can be prepared from compound (3-1) dissolved in solvents such as, for example, dimethylformamide or dimethylacetamide by the addition of a benzene solution of (E)-3-methoxyacryloyl isocyanate (prepared as described by Santana, L.; et al. J. Heterocyclic Chem. 1999, 36, 293-295.) at a temperature of about -40 to about -15°C under an inert atmosphere and then warming to ambient temperature for from about 30 min to about 4h. Compound (4-2) can be treated with an acid such as, for example, sulfuric acid in mixtures of water and ethanol in a temperature range of from about 90 to about 110°C for about 1 to about 8h to supply compound (4-3). Alternatively, compound (4-2) can be cyclized to uracil (4-3) under the basic conditions described by Ueno, Y.; et al. J. Org. Chem. 70:7925-7935 (2005).

## SCHEME 5




[0449] Compound (9-1) can be treated in refluxing thionyl chloride for about 1 to about 4h to obtain acid chloride (9-2). Treatment with thionyl chloride at the reflux temperature in solvents such as, for example, chloroform or toluene also furnishes compound (9-2). Compound (2) is converted to the corresponding aldehyde (9-3) by reduction with lithium *t*-butoxyaluminum hydride in a solvent such as, for example, tetrahydrofuran at about -78 °C over from about 1 to about 8h. The reduction can also be achieved by treatment with indium chloride and tributyltin hydride in the presence of triphenylphosphine in a solvent such as tetrahydrofuran or toluene at temperatures from about -40 to about 0°C. Compound (9-3) can be treated with compound (9-4) in the presence of a base such as potassium *t*-butoxide in a solvent such as dichloromethane at or near room temperature over a period of about 1 to about 8h to provide compound (9-5).

## SCHEME 6




[0450] Compound (10-1), wherein X<sup>1</sup> is halo (e.g., bromine, iodine) can undergo a Suzuki reaction with vinyl boronic acid (10-2) to provide compound (10-3). The reaction typically requires the use of a base and a catalyst. Examples of bases include, for example, potassium carbonate, potassium phosphate, potassium *t*-butoxide, sodium carbonate, cesium carbonate, and cesium fluoride. Examples of catalysts include, for example, tris(dibenzylideneacetone)dipalladium (0), palladium acetate, bis(triphenyl phosphine)palladium (II) chloride, tetrakis(triphenylphosphine)palladium, dichloro [1,1'-bis(di-*tert*-butylphosphino)ferrocene] palladium (II), or dichloro[1,1'-bis(diphenylphosphino)ferrocene] palladium (II) dichloromethane adduct. The reaction may be conducted in a solvent such as, for example, water, dioxane, dimethoxyethane, dimethylformamide, toluene, ethanol, tetrahydrofuran and the like or mixtures thereof. The reaction may be conducted at ambient or elevated temperatures.

## SCHEME 7




15 **[0451]** Compound (11-1) can be converted to compound (11-2) by treatment with diazomethane in a solvent such as, for example, tetrahydrofuran in the presence of palladium acetate at or near room temperature over a period of about 30 min to about 4h.

## SCHEME 8




35 **[0452]** Compound (11-1) is reduced to supply compound (14-2). Typical conditions for this reduction include using hydrogen at a pressure of about 1 to about 5 atmospheres in the presence of a catalyst such as, for example, palladium or platinum on charcoal in a solvent such as, for example, tetrahydrofuran, ethyl acetate, ethanol, or hexane at or near ambient temperature over a period of about 1 to about 12h.

## SCHEME 9




55 **[0453]** Compound (15-1) can be converted in a two-step sequence to compound (15-2). The initial step involves reduction of the aromatic nitro moiety with iron powder in the presence of a mild acid such as, for example, ammonium chloride or dilute hydrochloric acid at temperatures from about 60 to about 80°C in a mixture of solvents containing, for example, methanol, water, and tetrahydrofuran over about 1 to about 12h. The second step consists of exposure of the aniline, prepared in the first step, to methanesulfonyl chloride in the presence of a base such as pyridine in a solvent such as dichloromethane at or near ambient temperature.

## SCHEME 10




[0454] Compound (17-1) can be mesylated to provide compound (17-2) by treatment with methanesulfonyl chloride in the presence of a base such as, for example, pyridine in a solvent such as, for example, dichloromethane. Compound (17-3) can be exposed to borane dimethyl sulfide complex in a solvent such as, for example, tetrahydrofuran at approximately about 0 to about 10°C to supply compound (17-4). Compounds (17-2) and (17-4) can be combined with acetaldehyde in refluxing tetrahydrofuran. Subsequent treatment with water at room temperature yields compound (17-5).

## SCHEME 11



[0455] Carboxylic acid (18-1) can be reduced with boron tetrahydrofuran complex with heating to provide alcohol (18-2). Compound (18-2) is converted to the corresponding bromide (18-3) with N-bromosuccinimide and triphenylphosphine in solvents such as, for example, dichloromethane at room temperature in several hours. Treatment of compound (18-3) with triethyl phosphite at about 120°C for about 1 to about 3h supplies compound (18-4). Compound (18-4) can be used for example to make compound (9-5) as described in Scheme 5.

## SCHEME 12



[0456] Benzaldehyde (19-1) can be treated with diethyl phosphonate in the presence of a base such as, for example, sodium methoxide in a solvent such as, for example, methanol at room temperature to provide compound (19-2). Compound (19-2) can be treated with N-chlorosuccinimide and triphenylphosphine in dichloromethane at room temperature to yield compound (19-3). Compound (19-2) can also be reacted with (diethylamino)sulfur trifluoride (DAST) to supply compound (19-4).

[0457] Compound (19-1) can also be treated with p-toluenesulfonic acid and trimethyl orthoformate in methanol at

about 50°C to provide acetal (19-5). Compound (19-5) can be converted to compound (19-6) by exposure to triethyl phosphite and boron trifluoride diethyl etherate at about -20°C to about ambient temperature.

[0458] Compounds (19-3), (19-4), and (19-6) can be used for example to make compound (9-5) as described in Scheme 5.

5


### SCHEME 13

10

15

20

25



30

35

[0459] Phenol (20-1), wherein R<sup>4</sup> is other than amino, is treated with a source of electrophilic halide, such as, for example, iodine monochloride to provide dihalogenated compound (20-2), wherein X<sup>1</sup> and X<sup>2</sup> are independently bromine or iodine. Compound (20-2) is transformed to compound (20-3) by reaction of an alkylating agent such as, for example, methyl sulfate with a base such as, for example, potassium carbonate in refluxing acetone. Alternatively, methyl iodide in the presence of a base such as, for example, potassium *t*-butoxide in a solvent such as, for example, tetrahydrofuran, or dimethylformamide also furnish compound (20-3). In yet another alternative, compound (20-2) can be methylated with (trimethylsilyl)diazomethane in a solvent such as, for example, *t*-butyl methyl ether. Compound (20-3) can be reacted with uracil, ligand (20-4), copper (I) iodide, and potassium phosphate in dimethyl sulfoxide at about 40°C to about 100°C to supply compound (20-5).

40


[0460] For example, when in compound (20-3), R<sup>4</sup> is *tert*-butyl, X<sup>1</sup> is iodo, and X<sup>2</sup> is iodo or bromo, compound (20-3) can be stirred with uracil and compound (20-4) in the presence of CuI and K<sub>2</sub>PO<sub>4</sub> in DMSO for about 15 to about 24h at about 60°C to supply compound (20-5). Alternatives to ligand (20-4) for making (20-5) are 8-hydroxyquinoline and 2-(2-pyridyl)-benzimidazole.

45

50

55

SCHEME 14




[0461] Compound (21-1) can be nitrated with nitric acid in acetic acid in a temperature range of about 10 to about 15°C to give compound (21-2). The phenol moiety of compound (21-2) can be protected as a silyl ether, e.g. t-butyl dimethylsilyl ether, by treatment with a silyl chloride such as, for example, t-butyl dimethylsilyl chloride and imidazole in a solvent such as, for example, dimethyl formamide at ambient temperature to furnish compound (21-3). Compound (21-3) may then be reduced using conditions known to those skilled in the art to furnish the corresponding aniline (21-4).

[0462] Typical conditions for this reduction include using hydrogen at a pressure of about 1 to about 5 atmospheres in the presence of a catalyst such as, for example, palladium or platinum on charcoal in a solvent such as, for example, tetrahydrofuran, ethyl acetate, ethanol, methanol, or hexane at or near ambient temperature over a period of about 1 to about 12h. Dependent on the functional groups present, an alternative reduction procedure may be more appropriate such as, for example, using iron powder in the presence of a mild acid such as, for example, ammonium chloride or dilute hydrochloric acid at reflux temperatures in a mixture of solvents containing, for example, methanol, water, and tetrahydrofuran over about 1 to about 12h.

[0463] Aniline (21-4) can then be sulfonated with methanesulfonyl chloride in the presence of pyridine in a solvent such as, for example, dichloromethane. The starting material and reagents are combined at about 0°C and then allowed to gradually warm to ambient temperature over the course of the reaction to supply compound (21-5). The silyl ether protecting group is removed under conditions familiar to one skilled in the art. For example, tetrabutylammonium fluoride in tetrahydrofuran at room temperature transforms compound (21-5) to compound (21-6). The phenol group of compound (21-6) may be sulfonated with trifluoromethanesulfonyl anhydride in the presence of a base such as, for example, pyridine in a solvent such as, for example, dichloromethane at room temperature to provide compound (21-7). Compound (21-7) can be used as described in Scheme 12 to make compound (12-3).

## SCHEME 15




[0464] Compound (22-1) is converted to compound (22-2) in a two-step sequence. First, compound (22-1) can be hydrolyzed with a base such as, for example, sodium hydroxide, lithium hydroxide, or potassium hydroxide in a solvent such as, for example, methanol, ethanol, or tetrahydrofuran, or mixtures thereof. The resultant reaction mixture can be stirred for a period of about 6 to about 48h at ambient temperature. Second, the intermediate carboxylic acid is treated in refluxing thionyl chloride with or without a catalytic amount of dimethylformamide for about 1 to about 4h to deliver acid chloride (22-2). Treatment with thionyl chloride at reflux temperature in solvents such as, for example, chloroform or toluene also furnishes compound (22-2). Treatment of the carboxylic acid with oxalyl chloride in dichloromethane with a catalytic amount of dimethylformamide also furnishes compound (22-2).

[0465] Compound (22-2) can be treated with an amine or the corresponding salt in a solvent such as, for example, dioxane, dimethylformamide, dimethylacetamide, or dichloromethane optionally in the presence of a base such as, for example, pyridine, triethylamine or diisopropylethylamine at temperatures ranging from at or near ambient to about 100°C for between about 1 and about 24h to provide compound (22-4) wherein R<sup>11</sup> and R<sup>12</sup> are independently hydrogen or R<sup>F</sup>, or taken together with the nitrogen to which they are attached form a 5-6-membered heterocyclyl or a fused 2-ring heterocyclyl.

[0466] Compound (22-2) is converted to the corresponding aldehyde (22-3) by reduction with lithium tri-*t*-butoxyaluminum hydride in a solvent such as, for example, tetrahydrofuran at about -60°C to about -78°C.

## SCHEME 16



[0467] Compound (22-3) can be converted to compound (23-2) wherein  $R^{11}$  and  $R^{12}$  are independently hydrogen or  $R^F$ , or taken together with the nitrogen to which they are attached form a 5-6-membered heterocyclyl or a fused 2-ring heterocyclyl by treatment with an amine,  $N(R^{11})(R^{12})$ , in the presence of a reductant such as, for example, sodium triacetoxyborohydride or sodium cyanoborohydride in a solvent such as, for example, methanol, ethanol, dichloromethane, dimethylacetamide, or dimethylformamide over a period of about 1 to about 24h. The reaction often proceeds best at an acidic pH that can be maintained by the addition of acetic acid or hydrochloric acid.


[0468] Compound (22-3) can also be converted to compound (23-3) by reduction with lithium tri-*t*-butoxyaluminum hydride in a solvent such as tetrahydrofuran at room temperature.

40

45


50

55

**SCHEME 17**

30

[0469] Compound (23-3) can be converted to compound of formula (24-2) by treatment with thionyl chloride in dichloromethane at room temperature. Compound (24-2) can be treated with a sodium alkoxide,  $R^{13}ONa$ , in a heated solution of the corresponding alcohol to provide compound (24-3), wherein  $R^{13}$  is hydrogen or  $R^F$ .


**SCHEME 18**

[0470] Compound (25-1) can be brominated by treatment with, for example, pyridinium hydrobromide perbromide in a solvent such as, for example, acetic acid at or near ambient temperature over a period of about 1 to about 8h to give compound (25-2). The amino group of compound (25-2) can be removed by exposure to t-butyl nitrite in a solvent such as, for example, dimethylformamide at a temperature initially at ambient temperature and then increased to the range of about 50 to about 65 °C to give compound (25-3). Additional aliquots of t-butyl nitrite can be added at ambient temperature followed by heating until the transformation is complete. Compound (25-3) can be reduced to compound (25-4) by, for example, treatment with iron and ammonium chloride.

**EXAMPLES**

[0471] The following examples are merely illustrative, and not limiting to this disclosure in any way.

[0472] **Example A.** Preparation of (E)-N-(3-tert-butyl-5-iodo-4-methoxyphenylcarbamoyl)-3-methoxy acrylamide.



[0473] **Part A.** Preparation of 2-tert-butyl-4-nitrophenol.

[0474] To a vigorously stirred solution of 2-tert-butylphenol (10g, 66.6mmol) in heptane (67ml) was added at a fast drip a solution of 70% nitric acid (4.25ml, 66.6mmol) diluted with water (4.25ml). The resulting dark red/brown mixture was stirred vigorously for 2h. The suspended solid was collected by filtration washed with hexane (300mL), water (200mL) and once again with hexane (200mL) to give a cocoa colored powder that was dried to constant mass (4.65g, 35.6%).

[0475] **Part B.** Preparation of 2-tert-butyl-6-iodo-4-nitrophenol.

[0476] To the product from **Part A** (4.5g, 23.05mmol) dissolved in MeOH (120ml) and water (30mL) was added iodine monochloride (1.155ml, 23.05mmol) drop wise over a period of 10min. The mixture was stirred for 2h and diluted into 1L of water and allowed to stand overnight. The solid material was collected by filtration and washed 3x50mL with water and dried under vacuum overnight to give a tan solid (7.14g, 96%).

[0477] **Part C.** Preparation of 1-tert-butyl-3-iodo-2-methoxy-5-nitrobenzene.

To an ice bath cooled solution of the product from **Part B** (5.5g, 17.13mmol) in MTBE (15ml) in a 50mL pressure vessel was added 2.0M TMS diazomethane (12.85ml, 25.7mmol) followed by drop-wise addition of methanol (1.0mL) resulting in calm bubbling. The vessel was sealed and stirred at room temperature for 16h, cooled and the pressure was released. The solution was partitioned between EtOAc and water. The organic layer was washed with 1.0M HCl, saturated potassium carbonate solution, and saturated NaCl. The organic layer was dried over sodium sulfate, filtered and concentrated to give a red oil that was used without purification (5.4g, 84%).


[0478] **Part D.** Preparation of 3-tert-butyl-5-iodo-4-methoxyaniline.

[0479] A mixture of the product from **Part C** (5.80g, 17.31mmol), ammonium chloride (1.389g, 26.0mmol), and iron (4.83g, 87mmol) in THF/MeOH/water (200mL total, 2/2/1) was refluxed for 2h, cooled and filtered through Celite. The filtrate was evaporated and the residue was partitioned between water and EtOAc. The organic layer was washed with saturated brine, dried with sodium sulfate, filtered and evaporated to give a brown oil (5.28g, 100% yield).

[0480] **Part E.** Preparation of (E)-N-(3-tert-butyl-5-iodo-4-methoxyphenylcarbamoyl)-3-methoxy acrylamide.


[0481] To a solution of the product from **Part E** (3.05g, 10mmol) in DMF (50ml) at -20°C under N<sub>2</sub> was added at a fast drip a 0.4M solution in benzene of (E)-3-methoxyacryloyl isocyanate (50.0ml, 20.00mmol, prepared by the method of Santana et al., J. Heterocyclic Chem. 36:293 (1999)). The solution was stirred for 15min at -20°C, warmed to room temperature for 45min and diluted into EtOAc. The EtOAc layer was washed 4 x 300mL with water, 2 x 100mL with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to a brown solid. The residue was triturated in Et<sub>2</sub>O/hexane to give a fine powder that was collected by filtration and dried to give a tan powder (2.46g, 57%).

[0482] **Example B.** Preparation of 1-(3-tert-butyl-5-iodo-4-methoxyphenyl)dihydropyrimidine-2,4(1H,3H)-dione.



[0483] To a suspension of the product from **Example A** (2.46g, 5.69mmol) in ethanol (50ml) was added a solution of 5.5mL of H<sub>2</sub>SO<sub>4</sub> in 50mL water and the mixture was heated at 110°C for 2.5h to give a clear solution. The solution was cooled and diluted with 50mL of water while stirring to give an off-white solid that was collected by filtration, washed with water and dried (2.06g, 90%).

[0484] **Example C.** Preparation of 1-(3-tert-butyl-5-iodo-4-methoxyphenyl)pyrimidine-2,4(1H,3H)-dione.



5  
10  
[0485] **Part A.** Preparation of 2-tert-butyl-4,6-diiodophenol.

[0486] A solution of 2-tert-butylphenol (20.0g, 133mmol) in methanol (266mL) was treated with sodium hydroxide pellets (6.39g, 160mmol). The mixture was stirred until all the sodium hydroxide had dissolved and was then cooled in an ice-salt bath to -2°C. Sodium iodide (15.0g, 100mmol) was added and then 10 % sodium hypochlorite solution (45mL, 73.3mmol) was added drop wise at a rate such that the solution temperature rose no higher than 1.3°C. This sequence of events was repeated (3x) until a total of 60g (400mmol) of sodium iodide had been added and the sodium hypochlorite solution was added until the solution color changed from a light green-yellow color to the color of weak iced tea. This required all but 16mL of the 180mL total sodium hypochlorite solution measured out. With continued cooling at ca. 2°C, a solution of sodium thiosulfate pentahydrate (20g) in water (100mL) was added drop wise over 20min. After addition, the solution was acidified to pH 3 by drop wise addition of concentrated hydrochloric acid (ca. 35mL required of 40mL placed in the addition funnel). The precipitate was collected by filtration and washed with >1 liter of water. The salmon-colored solid was sucked as dry as possible, and dried in a vacuum oven at 50°C for 18h. These procedures afforded the product (49.61 g, 93%) as a tan solid.


25 [0487] **Part B.** Preparation of 1-tert-butyl-3,5-diiodo-2-methoxybenzene.

[0488] A solution of the product from **Part A** (20.0g, 49.7mmol) in acetone (140mL) was treated with methyl iodide (3.9mL, 8.83g, 62.2mmol) and 50 % (w/w) sodium hydroxide solution (3.02mL, 4.58g, 57.2mmol) followed by stirring at ambient temperature for 48h. The mixture was concentrated in vacuo to a volume of ca. 50-60mL, followed by dilution with heptane (80mL) and water (50mL). The layers were separated and the organic layer was extracted with saturated sodium chloride solution. Drying (Na<sub>2</sub>SO<sub>4</sub>) and concentration in vacuo afforded the product (20.59g, 99%) as a light yellow oil.

[0489] **Part C.** Preparation of 1-(3-tert-butyl-5-iodo-4-methoxyphenyl)pyrimidine-2,4(1H,3H)-dione.

[0490] A suspension of the product from **Part B** (12.04g, 28.9mmol), uracil (3.89g, 34.7mmol), N-(2-cyanophenyl) picolinamide (1.29g, 5.79mmol) and tribasic potassium phosphate (12.9g, 60.8mmol) in DMSO (181mL) was degassed by nitrogen sparge for 1 h. The mixture was then treated with copper (I) iodide (551mg, 2.89mmol) and degassing was continued for another 10min. The mixture was then warmed at 60°C for 18h. The mixture was then poured into water (600mL) and acidified to pH 3 by addition of 4N hydrochloric acid solution. The mixture was diluted with ethyl acetate, and the organic layer was extracted with water (3x), saturated ammonium chloride solution (1x) and saturated sodium chloride solution. The solution was dried and treated with (3-mercaptopropyl) silica gel, followed by stirring for 2h. The mixture was filtered and concentrated in vacuo. The solid obtained was triturated with ether-ethyl acetate (>10:1) and collected by filtration and washed with ether. After drying in a vacuum oven at 50°C for 2h, these procedures afforded the product (2.75 g) as a white solid. The mother liquors were concentrated in vacuo to afford an amber solid. This material was chromatographed over a Flash 65 silica gel cartridge, eluting with 20-100 % ethyl acetate in hexanes. These procedures afforded a nearly white solid, which was triturated with ether-hexanes and collected by filtration. After drying in a vacuum oven for 3h, these procedures afforded another 4.31g of the product as a white solid. Total yield: 7.06g(61 %).

[0491] **Example D.** Preparation of 1-(3-tert-Butyl-5-iodo-4-methoxyphenyl)pyrimidine-2,4(1H,3H)-dione.



[0492] **Part A.** Preparation of 2-*tert*-butyl-4,6-diiodophenol.

[0493] 2-*tert*-Butylphenol (99.95g, 665.36mmol) was dissolved in 1250mL methanol and converted to the corresponding phenoxide with 31.96g (799.0mmol, 1.2equiv.) of sodium hydroxide by stirring the sodium hydroxide pellets at room temperature, and then cooling the reaction mixture in an ice/salt bath. Sodium iodide (299.34g, 1997.07mmol, 3.0equiv.) and 8.3% bleach (1265.83g, 1411.39mmol, 2.1equiv.) were added to the cold reaction solution in four equal portions, the bleach being added while keeping the reaction mixture at <math>0^{\circ}\text{C}</math>. 500mL of 20% (w/w) sodium thiosulfate solution was added over an 18-minute period, with the temperature rising from


[0494] **Part B.** Preparation of 1-*tert*-butyl-3,5-diiodo-2-methoxybenzene.

[0495] The product from **Part A** (93% assay, 21.6g, 50mmol) was dissolved in 140mL of acetone. Methyl iodide (4.2mL, 67.5mmol, 1.35equiv.) was added, followed by 50% aqueous sodium hydroxide (5.0g, 62.5mmol, 1.25equiv.). The reaction was stirred overnight, then concentrated to approximately 50-60mL. 80mL of heptanes was added followed by 50mL of water, and the layers were shaken and separated, and the aqueous layer was back extracted with 20mL of heptanes. The organic layers were combined and washed twice with 50mL each of 10% aqueous NaCl to afford 91.1 grams of a heptane solution, which assayed to 19.1 g of the title compound.

[0496] **Part C.** Preparation of 1-(3-*tert*-Butyl-5-iodo-4-methoxyphenyl)pyrimidine-2,4(1H,3H)-dione.

[0497] Uracil (33.3g, 297mmol, 1.2equiv.),  $\text{K}_3\text{PO}_4$  (106g, 500mmol, 2.1equiv.), CuI (4.6g, 24.2mmol, 0.1equiv.), and *N*-(2-cyanophenyl)picolinamide (6.4g, 28.7mmol, 0.12equiv.) were charged to a flask and inerted with argon. The 1-*tert*-butyl-3,5-diiodo-2-methoxybenzene was solvent switched into MeCN, dissolved in 1L DMSO and sparged with argon and added to the solids. The reaction was heated to (\text{CuOAc})\_2 then 2 x 830mL 15%  $\text{NH}_4\text{Cl}$  then 800mL brine. The organic layer was then concentrated and chased with 1L heptane, then triturated with refluxing 85:15 (v/v) heptane:iPrOAc for 4h. After cooling, the product was collected by filtration and washed with an additional 330mL of 85:15 v/v heptanes:EtOAc to yield after drying 66.9g (70% yield) of the product as a white solid.

[0498] **Example E.** Preparation of (E)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.




[0499] The boronic acid (96% potency) (3.75g, 15.6mmol, 1.2eq), product from **Example D** (5.0g, 12.5mmol), Cytec

## EP 2 203 430 B1

ligand (175mg, 5mol%), Pd<sub>2</sub>(dba)<sub>3</sub> (46mg, 0.4mol%) and potassium phosphate (5.25g, 25.0mmol, 2eq.) were charged to a 3 neck RB flask. The solids were purged with nitrogen for 10min. 75mL 4:1 THF:water was sparged 10min and charged to the flask. The mixture was stirred to dissolve the solids followed by heating the mixture at 50 °C in darkness overnight. HPLC showed the reaction was not complete after stirring overnight (~ 2% iodouracil remained). The reaction mixture was diluted with 375mL DCM and 250ml 10% citric acid. The mixture was shaken in a sep funnel and the layers were separated. The DCM layer was washed with a solution of 0.6g L-cysteine in 250ml 5% NaHCO<sub>3</sub> for 30min which changed the DCM layer color from orange to yellow. Repeated the 0.6g L-cysteine in 250ml 5% NaHCO<sub>3</sub> for 30min treatment followed by a 250ml 5% NaHCO<sub>3</sub> wash, and a 250ml 10% NaCl wash. The DCM layer was treated with 2gm thiourea silica for 30min. Added 1gm carbon to decolorize mixed 5min and filtered through hy-flo. The wet cake was washed with DCM. The DCM solution was then stripped to give 6.74g of a light yellow solid. The solids were ~92% pure. The solids were heated in a mixture of 192ml DCM and 9mL MeOH. They never completely dissolved. Cooled to room temp with mixing. 80ml heptane was added and more product began to crystallize. The slurry stirred over the weekend. Added 50ml heptane in portions until a total of 230ml heptane was added. The product was filtered. Filtrate was measured at 1.21 mg/mL at 210nm and 1.35 at 220nm, which equals a 522-582mg loss in the liquors or 9-10% loss vs. theoretical. The wet cake was washed with 50ml of a 27ml Heptane:22ml DCM: 1 ml MeOH mixture. The wash contained 0.5mg/mL product or 25mg (0.4% vs. theoretical). Product yield 5.22gm (88.9%), purity 99.2%PA. Iodouracil was removed in the crystallization. Samples were submitted to solid state for analysis and analytical for Pd determination. NMR did not show any residual solvent.

**[0500] Example 1.** Preparation of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IA-L1-1.9**).



**[0501] Part A.** Preparation of methyl 3-tert-butyl-2-hydroxy-5-nitrobenzoate.

**[0502]** Methyl 3,5-di-tert-butyl-2-hydroxybenzoate (28.66g, 108.4mmol) was dissolved with stirring in 430mL glacial acetic acid and the resulting mixture was treated drop wise with fuming nitric acid (90%, 179.26mL). When the addition was complete, the resulting mixture was stirred for 2.5h. The reaction mixture was poured into a 2.0L of crushed ice and allowed to stand 30min. Afterwards, 1.0L of water was added and the ice water mixture was allowed to melt. The mixture was then filtered, washed with water and dried to provide the title compound (24.57g, 89%).

**[0503] Part B.** Preparation of methyl 3-tert-butyl-2-methoxy-5-nitrobenzoate.

**[0504]** Methyl 3-tert-butyl-2-hydroxy-5-nitrobenzoate (11.41g, 45.0mmol), potassium carbonate(9.34g, 67.6mmol), acetone (200mL), and dimethyl sulfate (6.46g, 67.6mmol) were added together. The resultant mixture was then heated to reflux for 16h. The mixture was then filtered and the solid was washed with ethyl acetate. The resulting organic liquid was then concentrated under vacuum to an oil and redissolved in ethyl acetate (600mL). The organic solution was then washed with water, dried, filtered and concentrated under vacuum to an oil that was then subjected to purification via column chromatography (gradient of 5% to 40% EtOAc/Hexanes) to yield the title compound as an oil (10.42, 87%).

**[0505] Part C.** Preparation of methyl 5-amino-3-tert-butyl-2-methoxybenzoate.

**[0506]** Methyl 3-tert-butyl-2-methoxy-5-nitrobenzoate(10.42g, 39.0mmol), iron powder (325mesh, 10.89g, 195mmol), ammonium chloride (3.13g, 58.5mmol), water (30mL), and methanol (150mL) were added together. The resultant mixture was then refluxed for 1h. The mixture was then cooled to room temperature, filtered through celite, and the celite washed with methanol. The filtrate was then concentrated under vacuum and dissolved in ethyl acetate (600mL). The resultant solution was then washed with water and brine. The organic extract was then dried, filtered and concentrated under vacuum to yield the title compound as an oil (9.25g, 100%).

**[0507] Part D.** Preparation of 3-(3-tert-butyl-4-methoxy-5-(methoxycarbonyl)phenylamino) propanoic acid.

**[0508]** The product from **Part C** (16.44g, 69.3mmol) was dissolved in toluene (200mL). This mixture was heated to reflux and acrylic acid added over time (1 mL of acrylic acid added every 3h, 5.23mL total, 76.2mmol). The mixture was then refluxed for 24h. The mixture was then cooled and concentrated under vacuum to dryness to yield an oil as the

crude title compound that was used directly in the next reaction.

**[0509] Part E.** Preparation of methyl 3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxybenzoate.

**[0510]** The product from **Part D** (21.43g, 69.3mmol), urea (10.4g, 173mmol) and acetic acid (glacial, 200mL) were added together. The mixture was then heated to 120°C for 18.5h followed by concentration under vacuum to give an oil. To this oil was added methanol (13mL), and ethyl acetate (350mL). The resultant mixture was allowed to stand for 24-48h whereby a precipitate formed. The resulting solid was filtered off and washed with a small amount of methanol (10mL) and then air dried to yield the title compound as a solid (15.26g, 66%).

**[0511] Part F.** Preparation of 3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxy benzoic acid.

**[0512]** The product from **Part D** (4.52g, 13.52mmol), methanol (70mL), and tetrahydrofuran (70mL) were added together. The mixture was then stirred vigorously until a homogenous solution resulted. Once homogenous, a solution of aqueous sodium hydroxide (1.0M, 68mL) was added. The mixture was then stirred for 12h, the mixture was then concentrated under vacuum to remove the organic solvent, followed by the addition of aqueous hydrochloric acid (1.0M, 80mL) that resulted in solid formation. The mixture was then concentrated under vacuum. To this material was added hydrochloric acid (12M, 100mL) and the resultant material heated to 100°C for 1.5h. The reaction was then cooled and water added. The resulting solid was filtered, washed with water, and dried to yield the title compound as a solid (3.55g, 82%).

**[0513] Part G.** Preparation of 3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxybenzaldehyde.

**[0514]** The product obtained in **Part F** (4.07g, 12.71mmol) and thionyl chloride (40.82mL, 559mmol) were combined and the mixture was refluxed for 2h, followed by concentration under vacuum to provide a light yellow colored solid product. The solid was dissolved in tetrahydrofuran (125mL), the solution cooled to -78°C and LiAlH(O<sup>t</sup>Bu)<sub>3</sub> (1M, 14mL) was added slowly over 10min while maintaining the temperature at -78°C. The mixture was stirred at -78°C for 2h, and the reaction was quenched with hydrochloric acid (aq., 1M, 25mL) at -78°C. The mixture was warmed to room temperature and ethyl acetate was added. The layers were separated and the aqueous layer was washed with ethyl acetate. The organic extracts were combined and washed with half saturated sodium bicarbonate solution. The organic layer was dried, filtered and concentrated under vacuum to yield the title compound as a solid (3.73g, 96%).

**[0515] Part H.** Preparation of 1-(3-tert-butyl-4-methoxy-5-(4-nitrostyryl)phenyl)dihydro- pyrimidine-2,4(1H,3H)-dione.


**[0516]** The product prepared in **Part G** (1.00g, 3.29mmol) and diethyl 4-nitrobenzyl- phosphonate (0.853g, 3.12mmol) were dissolved in dichloromethane (50mL). Solid potassium *tert*-butoxide (0.737g, 6.57mmol) was added portion wise at room temperature. The resultant dark red solution was stirred for 1.5h at room temperature. 1N aqueous HCl (50mL) solution was added and the mixture was stirred 30min, and then diluted with dichloromethane (50mL). The resultant organic layer was separated and dried. The material was purified by column chromatography on silica gel using 99/1 dichloromethane/methanol as eluent to obtain the title compound as a solid (1.12g, 80%).

**[0517] Part I.** Preparation of Preparation of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

**[0518]** The product obtained in **Part H** (1.1g, 2.60mmol), iron (0.725g, 12.99mmol), and ammonium chloride (0.208g, 3.90mmol) was added to a mixture of tetrahydrofuran (40mL), ethanol (40mL) and water (12mL). The slurry was heated to 90°C for 45min, and then cooled to ambient temperature. The solution was filtered through a pad of celite (10g), washed with ethanol (20mL), and the filtrate concentrated under vacuum to a solid. The resulting solid was dissolved in ethyl acetate (100mL), and the solution was washed with water (50mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. The drying agent was filtered off and the solvent removed under vacuum to give the aniline adduct as a yellow solid (830mg).

**[0519]** The solid (830mg, 2.109mmol) was dissolved in dichloromethane (50mL), and pyridine (0.512mL, 6.33mmol) and methanesulfonyl chloride (0.181mL, 2.32mmol) were added and the resulting solution was stirred at room temperature 16h. Dichloromethane (100mL) was added followed by extraction with a 1N aq. HCl solution (2 x 50mL). The organic layer was dried, concentrated under vacuum and purified by column chromatography on silica gel using 98/2 CH<sub>2</sub>Cl<sub>2</sub>/MeOH to provide the title compound as a solid (480mg, 39%, two steps). m.p. = 260-261°C (trans-isomer) <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>): δ ppm 1.37 (s, 9H), 2.71 (t, J=6.7Hz, 2H), 3.01 (s, 3H), 3.75 (s, 3H), 3.79 (t, J=6.6Hz, 2H), 7.13 (d, J=16.5Hz, 1H), 7.15 (d, J=2.4Hz, 2H), 7.23 (d, J=8.5Hz, 2H), 7.25 (d, J=16.5 Hz, 1H), 7.51 (d, J=2.4Hz, 1H), 7.61 (d, J=8.6Hz, 2H), 9.80(bs, 1H), 10.30 (s, 1H). (trans-isomer).

**[0520] Example 2.** Preparation of (Z)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-chlorovinyl)phenyl)methanesulfonamide (compound IA-L1-1.3).



[0521] **Part A.** Preparation of diethyl hydroxy(4-nitrophenyl)methylphosphonate.

[0522] The title compound was prepared as described in Taylor, WP, et. Al, Bioorg. Med. Chem. 4:1515-1520 (1996). 4-Nitrobenzaldehyde (3.0g, 19.85mmol) and diethyl phosphonate (2.74g, 19.85mmol) were combined and treated with a 0.5N solution of sodium methoxide in methanol (0.993mL, 0.496mmol). The resulting red-orange solution was stirred 12h at room temperature. The reaction mixture was extracted with dichloromethane (20mL) followed by half saturated ammonium chloride (20mL). The organic layer was separated, dried and concentrated under vacuum to provide the title compound as a semi-solid (5.1g, 89%).


[0523] **Part B.** Preparation of diethyl chloro(4-nitrophenyl)methylphosphonate.

[0524] The product prepared in **Part A** (500mg, 1.729mmol) was dissolved in dichloromethane (10mL) and treated with triphenylphosphine (998mg, 3.80mmol), followed by N-chlorosuccinimide (462mg, 3.46mmol). The mixture was stirred at room temperature for 18h. The solution was concentrated under vacuum and the residue was purified by column chromatography using silica gel eluting with a 1/1 mixture of hexanes/ethyl acetate to provide the title compound as an oil (262mg, 49%).

[0525] **Part C.** Preparation of (Z)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-chlorovinyl)phenyl)methanesulfonamide.


[0526] The product prepared in **Example 1, Part G** (100mg, 0.329mmole) was treated with the product obtained from **Part B** using the procedures described in **Example 1, Part H and Example 1, Part I** to provide 39mg of the title compound. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.36 (s, 9H), 2.71 (t, J=6.8Hz, 2H), 3.06 (s, 3H), 3.71 (s, 3H), 3.78 (t, J=6.8Hz, 2H), 7.23 (d, J=2.6 Hz, 1H), 7.27 (s, 1H), 7.28 (d, J=8.6Hz, 2H), 7.48 (d, J=2.6 Hz, 1H), 7.78 d, J=8.8 Hz, 1H), 10.05(s, 1H), 10.34 (s, 1H).

[0527] **Example 3.** Preparation of (E)-1-(3-tert-butyl-5-(4-fluorostyryl)-4-methoxyphenyl) dihydropyrimidine-2,4(1H, 3H)-dione (compound **IA-L1-1.12**).



[0528] The title compound was prepared according the procedures described in **Example 1, Part H and Example 1, Part I** using the product obtained in **Example 1, Part G** (50mg, 0.164mmol) and diethyl 4-fluorobenzylphosphonate (40.5mg, 0.164mmol). The title compound was obtained as a solid (30mg, 46%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.37 (s, 9H), 2.72 (t, J=6.6Hz, 2H), 3.76 (s, 3H), 3.79 (t, =6.6Hz, 2H), 7.21 (m, 4H), 7.30 (d, J=16.3Hz, 1H), 7.53 (d, J=2.6Hz, 1H), 7.73 (m, 2H), 10.35 (s, 1H).

[0529] **Example 4.** Preparation of (Z)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-fluorovinyl)phenyl)methanesulfonamide (compound **IA-L1-1.4**).



**[0530] Part A.** Preparation of diethyl fluoro(4-nitrophenyl)methylphosphonate.


**[0531]** The title compound was prepared as described in Taylor, WP, et. Al, Bioorg. Med. Chem. 4:1515-1520 (1996).

The product from **Example 2, Part A** (500mg, 1.729mmol) was dissolved in dichloromethane (10mL) and treated by drop wise addition of (diethylamino)sulfur trifluoride (DAST) (2.5mL, 18.9mmol). The mixture was stirred at room temperature for 18h. A solution of half saturated sodium phosphate monobasic (20mL) was added followed by dichloromethane (20mL) addition and separation of the resulting organic phase. The organic solution was dried and concentrated under vacuum, and then subjected to column chromatography using silica gel eluting with a 1/1 mixture of hexanes/ethyl acetate to provide the title compound as an oil (215mg, 43%).

**[0532] Part B.** Preparation of (Z)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-fluorovinyl)phenyl)methanesulfonamide.


The product prepared as described in **Part A** (100mg, 0.329mmole) was treated with the product prepared in **Example 1, Part G** (96mg, 0.329mmole) according to the procedures described in **Example 1, Part H** and **Example 1, Part I** to provide 53mg of the title compound as a 1/1 mixture of cis/trans isomers. Reverse phase HPLC chromatographic separation using a 40-100% gradient of acetonitrile in 0.1% aqueous trifluoroacetic acid provided the title compound as a solid (20mg). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.37 (s, 9H), 2.71 (t, J=6.8Hz, 2H), 3.06 (s, 3H), 3.77 (s, 3H), 3.78 (m, 2H), 6.62 (d, J=40.4Hz, 1H), 7.18 (d, J=2.6Hz, 1H), 7.30 (d, J=8.4Hz, 2H), 7.55 (d, J=2.6Hz, 1H), 7.75 (d, J=8.8Hz, 2H), 10.08 (s, 1H), 10.33 (s, 1H).

**[0534] Example 5.** Preparation of (E)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-fluorovinyl)phenyl)methanesulfonamide (compound **IA-L1-1.5**).



Reverse phase HPLC chromatographic separation of the 1/1 mixture of cis/trans isomeric material (53mg) from **Example 4, Part A** using a 40-100% gradient of acetonitrile in 0.1% aqueous trifluoroacetic acid provided the title compound as a solid (16.5mg). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.33 (s, 9H), 2.60 (t, J=6.6Hz, 2H), 3.01 (s, 3H), 3.57 (t, J=6.6Hz, 2H), 3.79 (s, 3H), 6.46 (d, J=21.3Hz, 1H), 6.87 (d, J=2.2Hz, 1H), 7.14 (m, 3H), 7.36 (d, J=8.8Hz, 2H), 10.02 (s, 1H), 10.24 (s, 1H).

**[0536] Example 6.** Preparation of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)-2-fluorophenyl)methanesulfonamide (compound **IA-L1-1.26**)



**[0537] Part A.** Preparation of 4-(bromomethyl)-2-fluoro-1-nitrobenzene.

**[0538]** (3-Fluoro-4-nitrophenol)methanol (1.24g, 7.25mmol) was dissolved in dichloromethane (25mL) and treated with triphenylphosphine (2.28 g, 8.70mmol) followed by N-bromosuccinimide (1.548g, 8.70mmol). The mixture was stirred at room temperature for 2h. Water (50mL) and dichloromethane (40mL) were added, and the organic layer was separated and dried. The solution was concentrated under vacuum and purified by column chromatography using silica gel eluting with a 5/1 mixture of hexanes/ethyl acetate to provide the title compound as a solid (1.27g, 75%).


**[0539] Part B.** Preparation of diethyl 3-fluoro-4-nitrobenzylphosphonate.

**[0540]** The product prepared in **Part A** (1.27g, 5.43mmol) was added to triethyl phosphite (8mL, 54.3mmol) and the solution heated to 120°C for 1hr. After cooling, the excess triethyl phosphite was removed by heating under vacuum and the residue subjected to column chromatography on silica gel using 99/1 dichloromethane/methanol as eluent to obtain the crude title compound as an oil (800mg).

**[0541] Part C.** Preparation of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)-2-fluorophenyl)methanesulfonamide.


**[0542]** The product described in **Example 1, Part G** (533mg, 1.751mmole) was treated with the product described in **Part B** (510mg, 1.751 mmole) according to the procedures described in **Example 1, Part H** and **Example 1, Part I** to provide 80mg of the title compound. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.37 (s, 9H), 2.71 (t, J=6.5Hz, 2H), 3.05 (s, 3H), 3.76 (s, 3H), 3.79 (t, J=6.6Hz, 2H), 7.18 (m, 2H), 7.36 (d, J=16.5Hz, 1H), 7.39 (m, 1H), 7.44 (m, 1H), 7.52 (d, J=2.6Hz, 1H), 7.63 (m, 1H), 9.65 (s, 1H), 10.35 (s, 1H).

**[0543] Example 7.** Preparation of N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)cyclopropyl)phenyl)methanesulfonamide (compound **IA-L8-1.1**).




**[0544]** The product obtained as described in **Example 1, Part I** (30mg, 0.064mmol) was dissolved in tetrahydrofuran (2mL) and treated with 0.95mL of a 0.67M ether solution of diazomethane (0.636mmol) followed by palladium acetate (0.7mg, 0.0031mmol). The mixture was stirred for 30 min at room temperature followed by removal of the solid by filtration and concentration of the filtrate. The filtrate was purified by column chromatography on silica gel using 98/2 dichloromethane/methanol as eluent to obtain the title compound as a solid (21.6mg, 70%). m.p. 265-266°C. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.33 (s, 9H) 1.50 (m, 2H), 2.13 (m, 1H), 2.27 (m, 1H), 2.69 (t, J=6.6Hz, 2H), 2.94 (s, 3H), 3.63 (s, 3H), 3.74 (t, J=6.6Hz, 2H), 6.84 (d, J=2.6Hz, 1H), 7.04 (d, J=2.6Hz, 1H), 7.14 (d, J=8.8Hz, 2H), 7.20 (d, J=8.8Hz, 2H), 9.60 (s, 1H), 10.29 (s, 1H).

**[0545] Example 8.** Preparation of N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenethyl)phenyl)methanesulfonamide (compound **IA-L5-2-1.1**).



15 **[0546]** The product obtained as described in **Example 1, Part I** (415mg, 0.88mmol) was dissolved in methanol (30mL) and treated with 50mg of 10% palladium on carbon. The slurry was stirred for 48h at room temperature under latm of hydrogen. The reaction mixture was filtered through celite and concentrated in vacuo to provide the title compound as a solid (230mg, 55%). m.p. 233-234°C. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.34 (s, 9H), 2.68 (t, J=6.8Hz, 2H), 2.86 (s, 4H), 2.93 (s, 3H), 3.70 (m, 2H), 3.74 (s, 3H), 7.11 (m, 4H), 7.23 (m, 2H), 9.59 (s, 1H), 10.29 (s,).

20 **[0547] Example 9.** Preparation of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)styryl)phenyl)methanesulfonamide (compound **IA-L1-1.16**)



35 **[0548] Part A.** Preparation of methyl 3-tert-butyl-5-(chlorocarbonyl)benzoate.

**[0549]** A mixture of 3-tert-butyl-5-(methoxycarbonyl)benzoic acid (9.18g, 38.9mmol, prepared by the method of Carter et. al., WO2005021500A1), thionyl chloride (75mL) and 1 drop of DMF in toluene (200mL) was heated at reflux for 2h, cooled and concentrated. The residue was azeotroped with toluene (3 x 50mL) and dried under high vacuum to give the title compound as an off-white waxy solid (9.9g, quantitative yield).

40 **[0550] Part B.** Preparation of methyl 3-(azidocarbonyl)-5-tert-butylbenzoate.

**[0551]** To the product of **Part A** (9.9g, 38.9mmol) in acetone (200ml) was added at a fast drip a solution of sodium azide (10.12g, 156mmol) dissolved in water (20mL). The mixture was stirred for 2h and diluted with EtOAc. The organic layer was washed with H<sub>2</sub>O, saturated brine, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to give the title compound as a white solid (9.9g, 97%).

45 **[0552] Part C.** Preparation of methyl 3-amino-5-tert-butylbenzoate.

**[0553]** The product from **Part B** (9.9g, 37.9mmol) in toluene (100mL) was heated at reflux for 1 h and concentrated to give the intermediate isocyanate which was dissolved in DME (60mL) treated with 8% HCl (150mL) and stirred for 16h. The mixture was concentrated and the residue was dissolved in water, neutralized with solid sodium bicarbonate and extracted 3 x 100mL with EtOAc. The organics were combined, washed with saturated NaCl, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The crude product was chromatographed on silica eluting with 2:1 hexane/EtOAc to give the title compound as an oil (2.7g, 35%).

**[0554] Part D.** Preparation of methyl 3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxybenzoate.

50 **[0555]** A mixture of the product of **Part C** (2.34g, 11.29mmol) and acrylic acid (2.32ml, 33.9mmol) in toluene (60ml) was heated at reflux under nitrogen for 24h, cooled and concentrated. The resulting residue was then treated with urea (2.03g, 33.9mmol) in acetic acid (35ml) and heated at 120°C for 24h, cooled and concentrated. The residue was azeotroped 3 x 50mL with toluene and dissolved in 100mL of EtOAc. The organic layer was washed with dilute aqueous NaHCO<sub>3</sub>, H<sub>2</sub>O, saturated brine, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to give the title compound as a white solid (2.1g, 61 %).


**[0556]** Part E. Preparation of 3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)benzoic acid.

**[0557]** A mixture of the product from **Part D** (1.8g, 5.91mmol) and 1M NaOH (29.6ml, 29.6mmol) in MeOH (15ml) and THF (15mL) was stirred for 24h and concentrated. The residue was treated with 50mL of 1M HCl and extracted into EtOAc. The EtOAc layer was washed with H<sub>2</sub>O, saturated brine, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to give a white solid. This intermediate urea was combined with 20mL of concentrated HCl and heated at 100°C for 1h, cooled and diluted with 75mL of ice water to give a white powder which was collected by filtration and dried to constant mass to give the title compound (1.6g, 93%).

**[0558]** **Part F.** Preparation of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl) styryl)phenyl)methanesulfonamide.

**[0559]** The product described in **Part E** was treated with thionyl chloride and lithium tri-tert-butoxyaluminum hydride according to procedures described in **Example 1, Part G** to produce 3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)benzaldehyde. The aldehyde was treated with diethyl 4-nitrobenzylphosphonate according to the procedures described in **Example 1, Part H** and **Example 1, Part I** to provide the title compound (85mg). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.32 (s, 9 H) 2.72 (t, J=6.43 Hz, 2 H) 3.01 (s, 3 H) 3.82 (t, J=6.62 Hz, 2 H) 7.18 - 7.25 (m, 5 H) 7.39 (s, 1H) 7.46 (s, 1H) 7.58 (d, J=8.46 Hz, 2 H) 9.84 (s, 1H) 10.37 (s, 1H).

**[0560]** **Example 10.** Preparation of (Z)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-methoxyvinyl)phenyl)methanesulfonamide (compound **IA-L1-1.17**).



**[0561]** **Part A.** Preparation of 1-(dimethoxymethyl)-4-nitrobenzene.

**[0562]** A flask equipped with a magnetic stir bar and vigreux column was charged with 4-nitrobenzaldehyde (5.0g, 33.1mmol), pyridinium *p*-toluenesulfonate (1.66g, 6.62mmol), trimethoxymethane (3.51g, 33.1mmol) and methanol (100mL). The mixture was heated at 50°C for 12h and was concentrated in vacuo. The residue was redissolved in EtOAc and washed with aq. NaOH (1M), H<sub>2</sub>O and brine. The mixture was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated in vacuo to yield the title compound as a clear, light yellow oily product (6.36g, 97%).


**[0563]** **Part B.** Preparation of diethyl methoxy(4-nitrophenyl)methylphosphonate.

**[0564]** The product from **Part A** (3.0g, 15.2mmol) and triethyl phosphite (2.53g, 15.2mmol) were dissolved in dichloromethane (30mL) under a nitrogen atmosphere, cooled to -20°C and treated with drop wise addition of boron trifluoride etherate (2.27g, 16mmol). The mixture was allowed to slowly warm to room temperature overnight with stirring. Water was added and the resulting mixture was stirred 5min, separated and the organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated in vacuo to a solid residue. The residue was purified on silica gel (100% EtOAc to 3% CH<sub>3</sub>OH/EtOAc) to yield the title compound as a light yellow oily product (3.78 g, 82%).

**[0565]** **Part C.** Preparation of (Z)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-methoxyvinyl)phenyl)methanesulfonamide.

**[0566]** The product obtained according to the procedure described in **Example 1, Part G** (400mg, 1.314mmole) was treated with the product obtained in **Part B** (399mg, 1.314mmole) according to the procedures described in **Example 1, Part H** and **Example 1, Part I** to provide the title compound (17mg, 6%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.36 (s, 9 H) 2.71 (t, J=6.62 Hz, 2 H) 3.05 (s, 3 H) 3.58 (s, 3 H) 3.75 (s, 3 H) 3.76 - 3.81 (m, 2 H) 6.25 (s, 1H) 7.11 (d, J=2.57 Hz, 1H) 7.27 (d, J=8.46 Hz, 2 H) 7.60 (d, J=8.82 Hz, 2 H) 7.67 (d, J=2.57 Hz, 1H) 9.96 (s, 1H) 10.32 (s, 1H).


**[0567]** **Example 11.** Preparation of (E)-1-(3-tert-butyl-4-methoxy-5-styrylphenyl)dihydro-pyrimidine-2,4(1H,3H)-dione (compound **IA-L1-1.18**).



[0568] The product obtained according to procedure described in **Example 1, Part G** (50mg, 0.164mmole) was treated with diethyl benzylphosphonate (0.034ml, 0.164mmole) according to the procedure described in **Example 1, Part H** to provide the title compound (13mg, 19%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.37 (s, 9 H) 2.72 (t, J=6.62 Hz, 2 H) 3.76 (s, 3 H) 3.80 (t, J=6.80 Hz, 2 H) 7.16 -7.18 (m, 1H) 7.21 - 7.23 (m, 1H) 7.29 - 7.33 (m, 2 H) 7.36 - 7.43 (m, 2 H) 7.54 (d, J=2.57 Hz, 1H) 7.64 (d, J=7.35 Hz, 2 H) 10.35 (s, 1H).

[0569] **Example 12.** Preparation of (E)-1-(3-tert-butyl-4-methoxy-5-(4-methoxystyryl)phenyl)dihydro pyrimidine-2,4 (1H,3H)-dione (compound **IA-L1-1.14**).


20



[0570] The product obtained according to procedure described in **Example 1, Part G** (50mg, 0.164mmole) was treated with diethyl 4-methoxybenzylphosphonate (0.028ml, 0.164mmole) according to the procedure described in **Example 1, Part H** to provide the title compound (4mg, 4%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ ppm 1.37 (s, 9 H) 2.71 (t, J=6.62 Hz, 2 H) 3.70 - 3.81 (m, 8H) 6.96 (d, J=8.82 Hz, 2 H) 7.13 (d, J=2.21 Hz, 1H) 7.15 (d, J=2.57 Hz, 2 H) 7.50 (d, J=2.57 Hz, 1H) 7.58 (d, J=8.46 Hz, 2 H) 10.34 (s, 1H).

[0571] **Example 13A.** Preparation of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.1**).

40



[0572] **Part A.** Preparation of (E)-methyl 3-tert-butyl-2-methoxy-5-(3-(3-methoxyacryloyl)ureido) benzoate.

[0573] The product obtained as described in **Example 1, Part C** (2.0g, 8.43mmol) was dissolved in 30mL of N,N-dimethylacetamide and cooled to -25°C. A 0.5Molar solution of E-3-methoxyacryloyl isocyanate in benzene (21.9mL, 10.96mmol) was added drop wise and the resulting solution was stirred at ambient temperature for 4h, and then poured into water. The product was extracted into dichloromethane, washed with brine, dried over sodium sulfate, filtered and evaporated under vacuum to give the title compound.

55

[0574] **Part B.** Preparation of methyl 3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxybenzoate.

[0575] The product from **Part A** (3.1g, 8.51mmol) was dissolved in ethanol (60mL). To this solution was added a mixture of concentrated sulfuric acid (6mL) and water (60mL). The heterogeneous mixture was heated at 100°C for 3h. The ethanol was removed under vacuum, and then the aqueous solution was extracted with dichloromethane and evaporated to dryness. This residue was purified by column chromatography on silica gel, eluting with 1% methanol/dichloromethane to yield the title compound (1.23g, 44%).

[0576] **Part C.** Preparation of 3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy benzoic acid.

[0577] The product from **Part B** (1.23g, 3.7mmol) was taken up in ethanol (5mL) and 1M sodium hydroxide solution (10mL) and stirred at ambient temperature for 18h. The solution was acidified with 1M HCl and the resulting solid was filtered and dried to give the title compound (0.945 g, 80%).

[0578] **Part D.** Preparation of 3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy benzaldehyde.

[0579] The product from **Part C** (0.945g, 2.97mmol) was taken up in thionyl chloride (4.5mL) and the mixture was heated at 80°C for 40min. After evaporation to dryness, the acid chloride was dissolved in dry THF (8mL) and cooled to -78°C. A 1 M solution of lithium tri-tert-butoxyaluminum hydride in THF (3.0mL, 3.0mmol) was added drop wise. After 45min the cold reaction was quenched with 1M HCl (5mL), extracted into ethyl acetate, and purified by column chromatography on silica gel, eluting with dichloromethane followed by 1% methanol/dichloromethane to give the title compound (0.635 g, 71%).

[0580] **Part E.** Preparation of (E)-1-(3-tert-butyl-4-methoxy-5-(4-nitrostyryl)phenyl)pyrimidine-2,4(1H,3H)-dione.

[0581] The product of **Part D** (0.634g, 2.1 mmol) and diethyl 4-nitrobenzylphosphonate (0.573g, 2.1 mmol) were combined in dichloromethane (25mL) at ambient temperature. Potassium tert-butoxide (0.494g, 4.4mmol) was added portion wise and the resulting red/brown heterogeneous mixture was stirred for 1.5h. This mixture was quenched with 1M HCl (15mL), poured into water and extracted into ethyl acetate, and the crude product was purified by column chromatography on silica gel, eluting with 1% methanol/dichloromethane to give the title compound (0.735g, 83%).


[0582] **Part F.** Preparation of (E)-1-(3-(4-aminostyryl)-5-tert-butyl-4-methoxyphenyl)pyrimidine-2,4(1H,3H)-dione.

[0583] The product from **Part E** (0.735g, 1.74mmol), ammonium chloride (0.14g, 2.62mmol), and iron (0.487g, 8.72mmol) were combined in a solution of ethanol (10mL) water (5mL), and THF (10mL) and heated at 75°C for 1h. The mixture was filtered through diatomaceous earth, rinsing well with THF and concentrated to give the title compound.

[0584] **Part G.** Preparation of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

[0585] The product from **Part F** (0.683g, 1.75mmol) and pyridine (0.564mL, 6.98mmol) were combined in dichloromethane (15mL) at ambient temperature. Methane sulfonylchloride (0.163mL, 2.1 mmol) was added drop wise and the solution was stirred for 18h. The mixture was poured into 1 M HCl and extracted into dichloromethane, concentrated, and purified by column chromatography on silica gel, eluting with 1%, 2% methanol/dichloromethane. Trituration from dichloromethane provided a solid that was filtered and dried to give the title compound as a colorless powder (0.465g, 57%). <sup>1</sup>H NMR (300 MHz, DMSO-D<sub>6</sub>) δ ppm 1.38 (s, 9 H), 3.01 (s, 3H), 3.79 (s, 3 H) 5.65 (d, J=7.72 Hz, 1H), 7.17 - 7.28 (m, 5 H), 7.58 - 7.70 (m, 3 H), 7.75 (d, J=7.72 Hz, 1H), 9.86 (s, 1H), 11.42 (s, 1H).

[0586] **Example 13B.** Preparation of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.1**).



[0587] **Part A.** Preparation of N-(4-ethynylphenyl)methanesulfonamide.

[0588] In a 2L, 3-neck round-bottom flask equipped with an overhead stirrer was added 4-ethynylaniline (30g, 256mmol) and pyridine (42.5ml, 525mmol) in dichloromethane (512ml) to give an orange solution. The mixture was cooled to 5°C and methanesulfonyl chloride (19.96ml, 256mmol) was added drop wise over 15min. The reaction solution was stirred at 5°C for 2h and washed with 1M aqueous HCl (3x250mL). The dichloromethane layer was then washed sequentially with saturated aqueous NaHCO<sub>3</sub>, water, and saturated aqueous NaCl. The dichloromethane layer was dried over sodium sulfate and treated simultaneously with decolorizing charcoal for 30min, the solution then filtered through Celite and the

filtrate was concentrated. The pink/orange solid was dissolved in a minimal amount of hot ethyl acetate (50-75mL) and slowly diluted with hexanes (500-600ml) to give orange crystals that were collected by filtration and dried to provide the title compound (40.0g, 80%).

**[0589] Part B.** Preparation of (E)-4-(methylsulfonamido)styrylboronic acid.

**[0590]** (Reference: Org. Prep. Proc. Int., 2004, 36, 573-579) To a flask was added borane-methyl sulfide complex (8.03mL, 85mmol) followed by tetrahydrofuran (16mL) and the mixture then cooled to 0°C. (1R)-(+)-alpha-pinene (26.2mL, 169mmol) was then added drop wise (over 10min) to the ice-cooled solution. The mixture was then stirred at 0°C for 1h followed by stirring 2h at room temperature. The resulting thick white slurry was cooled to -40°C in a dry ice/acetone bath, followed by the addition of the product from **Part A** (15.0g, 77mmol) dissolved in 60mL of THF, drop wise over 30min. After the addition was complete, the mixture was stirred for an additional hour at -35°C, then 1h at room temperature. The light yellow solution was then cooled to 0°C and acetaldehyde (61.4mL, 1088mmol) added, then the mixture refluxed at 50°C for 18h. The solvent was then removed under vacuum to provide an orange syrup, to which water (115mL) was added and the heterogeneous mixture stirred for 3h at room temperature. The light yellow solid generated was collected and washed with water (250mL) then dried in a vacuum oven overnight. The resultant material was then dissolved in boiling acetone (190mL), which provided a homogenous yellow solution, followed by removal of the solution from heating and the addition of hexanes (365ml) over 5min time. A white solid formed in the solution and the mixture was stirred until the solution cooled to room temperature, then the white solid was collected and dried in a vacuum oven for 1hr to provide the title compound (12.1g, 85%).

**[0591] Part C.** Preparation of 2-tert-butyl-4-nitrophenol.

**[0592]** To a vigorously stirred solution of 2-tert-butylphenol (10g, 66.6mmol) in heptane (67ml) was added at a fast drip a solution of 70% nitric acid (4.25ml, 66.6mmol) diluted with water (4.25ml). The resulting dark red/brown mixture was stirred vigorously for 2h. The suspended solid was collected by filtration washed with hexane (300mL), water (200mL) and once again with hexane (200mL) to give a cocoa colored powder that was dried to constant mass (4.65g, 35.6%).

**[0593] Part D.** Preparation of 2-bromo-6-tert-butyl-4-nitrophenol.

**[0594]** A solution of the product from **Part C** (1.0g, 5.12mmol) in glacial acetic acid (10.25mL) was treated portion wise with pyridine hydrobromide perbromide (1.80g, 5.63mmol) followed by stirring at room temperature for 2h. Additional pyridinium hydrobromide perbromide (3.6g) was added in two portions and after another 3h of stirring, the reaction was complete. The mixture was poured into ice water, and the mixture treated with a small amount of sodium sulfite. The resulting solid was filtered and dried under vacuum to give the title compound as a brown solid (1.40g, 100 %).

**[0595] Part E.** Preparation of 1-bromo-3-tert-butyl-2-methoxy-5-nitrobenzene.

**[0596]** A solution of the product from **Part D** (1.40g, 5.11mmol) in 10:1 t-butylmethylether-methanol (25.5mL) was treated with 2.0M trimethylsilyldiazomethane in ether (5.1mL, 10.21mmol), followed by stirring at room temperature for 18h. The mixture was concentrated under vacuum to afford a yellow oil, which was purified by silica gel column chromatography eluting with EtOAc/hexanes to give the title compound as a yellow oil (1.36g, 92 %).

**[0597] Part F.** Preparation of tert-butyl 3-bromo-5-tert-butyl-4-methoxyphenylcarbamate.

**[0598]** A solution of the product from **Part E** (960mg, 3.33mmol) in methanol (17mL) was treated with 5 % platinum on sulfided carbon (100mg), followed by hydrogenation under balloon pressure for 3h, and then filtered through celite and concentrated under vacuum to afford the 3-bromo-5-tert-butyl-4-methoxyaniline as a yellow oil (860mg, 3.33mmol, 100%). A solution of this material in THF (17mL) was treated with di-tert-butyl dicarbonate (800mg, 3.66mmol) followed by warming at reflux for 2h. Concentration under vacuum afforded a beige solid, which was purified by silica gel column chromatography eluting with EtOAc/hexanes. Solid was triturated with hexanes, collected by filtration, and dried under vacuum to give the title compound as a nearly white solid (890mg, 75 %).

**[0599] Part G.** Preparation of (E)-N-(3-bromo-5-tert-butyl-4-methoxyphenylcarbamoyl)-3-methoxy acrylamide.

**[0600]** The product from **Part F** (2.0g, 5.58mmol) was dissolved in dichloromethane (10mL) and trifluoroacetic acid (5mL) added. The solution was stirred at room temperature for 1h followed by concentration under vacuum and the addition of 10% aqueous sodium bicarbonate (50mL), followed by extraction with ethyl acetate (3 x 50mL). The combined organic extracts were dried and concentrated to provide a residue that was dissolved in 10mL of N,N-dimethylacetamide and cooled to -25°C. A 0.5 molar solution of E-3-methoxyacryloyl isocyanate in benzene (20.3mL, 11.16mmol) was added drop wise and the resulting solution was stirred at ambient temperature for 4h, and then poured into water. The product was extracted into dichloromethane, washed with brine, dried over sodium sulfate, filtered and evaporated under vacuum to give the title compound.

**[0601] Part H.** Preparation of 1-(3-bromo-5-tert-butyl-4-methoxyphenyl)pyrimidine-2,4(1H,3H)-dione.


**[0602]** The product from **Part G** (2.15g, 5.58mmol) was dissolved in ethanol (10mL). To this solution was added a mixture of concentrated sulfuric acid (1mL) and water (10mL). The heterogeneous mixture was heated at 100°C for 2h. The ethanol was removed under vacuum, and then the aqueous solution was extracted with dichloromethane and evaporated to dryness. This residue was purified by column chromatography on silica gel, eluting with 1% methanol/dichloromethane to yield the title compound (1.35g, 69%).

**[0603] Part I.** Preparation of (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phe-

nyl)methanesulfonamide.

[0604] The product from **Part H** (8.0g, 22.65mmol), the product from **Part B** (5.90g, 24.46mmol), 1,1'-bis(di-tert-butylphosphino)ferrocene palladium dichloride (0.738g, 1.132mmol), and potassium phosphate (9.62g, 45.3mmol) were dissolved in a mixture of tetrahydrofuran (128mL) and water (32mL). Nitrogen gas was bubbled through the resultant mixture for 10 min followed by heating the solution at 50 °C for 5h in darkness. The reaction was allowed to cool to room temperature followed by the addition of saturated aqueous ammonium chloride (50mL), water (200mL), and the solution extracted with dichloromethane (600mL). To the organic extract was added magnesium sulfate, and 3-mercaptopropyl-functionalized silica gel (20 g) and the resultant solution stirred in darkness for 18h. The solids were then removed by filtration and the filtrate concentrated under vacuum and subjected to silica gel column chromatography using a 99/1 to 99/2 dichloromethane/methanol gradient to provide the title compound (7.4 g, 70%). <sup>1</sup>H NMR (300 MHz, DMSO-D<sub>6</sub>) δ ppm 1.38 (s, 9 H), 3.01 (s, 3 H), 3.79 (s, 3 H) 5.65 (d, J=7.72 Hz, 1H), 7.17 - 7.28 (m, 5 H), 7.58 - 7.70 (m, 3 H), 7.75 (d, J=7.72 Hz, 1H), 9.86 (s, 1H), 11.42 (s, 1H).

[0605] **Example 14.** Preparation of (E)-N-(4-(3-tert-butyl-5-(5-fluoro-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1.1.2**).



[0606] **Part A.** Preparation of methyl 3-tert-butyl-5-(5-fluoro-6-methoxy-2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxybenzoate.

[0607] The fluorination procedure was performed as described in Lal, GS, et al. J. Org Chem., 60:7340-7342 (1995). The product from **Example 13A, Part B** (0.42g, 1.26mmol) and Selectfluor™ (0.672g, 1.9mmol) were combined in a mixture of acetonitrile (8mL) and methanol (1mL) and heated at 90 °C under N<sub>2</sub> for 5h. The solution was diluted with water, extracted into ethyl acetate, washed with sodium bicarbonate solution, concentrated and purified by column chromatography on silica gel to give the title compound (0.138g, 29%).

[0608] **Part B.** Preparation of methyl 3-tert-butyl-5-(5-fluoro-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxybenzoate.

[0609] The product from **Part A** (0.134g, 0.35mmol) and triethylamine (1mL) were combined in methanol (4mL) and stirred at ambient temperature for 18h. The solution was quenched with 1M HCl, extracted into dichloromethane and concentrated to give the title compound (0.113g, 92%).

[0610] **Part C.** Preparation of 3-tert-butyl-5-(5-fluoro-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxybenzoic acid.

[0611] The product from **Part B** (0.113g, 0.32mmol) was treated as described in **Example 13A, Part C** to give the title compound (0.088g, 81%).

[0612] **Part D.** Preparation of 3-tert-butyl-5-(5-fluoro-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxybenzaldehyde.

[0613] The product from **Part C** (0.088g, 0.26mmol) was treated as described in **Example 13A, Part D** to give the title compound (0.075g, 90%).

[0614] **Part E.** Preparation of (E)-1-(3-tert-butyl-4-methoxy-5-(4-nitrostyryl)phenyl)-5-fluoropyrimidine-2,4(1H,3H)-dione.

[0615] The product of **Part D** (0.075g, 0.23mmol) was treated as described in **Example 13A, Part E** to give 0.077g (75%).

[0616] **Part F.** Preparation of (E)-1-(3-(4-aminostyryl)-5-tert-butyl-4-methoxyphenyl)-5-fluoro pyrimidine-2,4(1H,3H)-dione.


[0617] The product of **Part E** (0.077g, 0.18mmol) was treated as described in **Example 13A, Part F** to give the title compound (0.071g, 94%).

[0618] **Part G.** Preparation of (E)-N-(4-(3-tert-butyl-5-(5-fluoro-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-

styryl)phenyl)methanesulfonamide.

[0619] The product of **Part F** (0.071g, 0.17mmol) was treated as described in **Example 13A, Part G** to give the title compound (0.048g, 57%). <sup>1</sup>H NMR (300 MHz, DMSO-D<sub>6</sub>): δ ppm 1.38(s, 9 H), 3.01(s, 3 H), 3.79 (s, 3 H) 7.19 - 7.27 (m, 5 H), 7.62 (d, J=8.82 Hz, 2 H), 7.66 (d, J=2.57 Hz, 1H), 8.25 (d, J=6.99 Hz, 1 H).

[0620] **Example 15.** Preparation of (E)-N-(4-(3-bromo-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.52**).



[0621] **part A.** Preparation of 2-bromo-4,6-diiodophenol.

[0622] A 1L round-bottom flask was charged with 2-bromophenol (8.65g, 50mmol) and methanol (100ml) to give a colorless solution. Sodium hydroxide (2.40g, 60.0mmol) was added and stirred until the hydroxide pellets had dissolved. The solution was cooled in an ice water bath and sodium iodide (5.6g, 37.4mmol) was added followed by drop-wise addition of sodium hypochlorite (17mL, 27. mmol) to give a transparent brown/red solution and gradual precipitation of a thick, white solid. The addition of sodium iodide and bleach was repeated 3 times to give an orange mixture that was stirred for 2h, treated with a solution of sodium thiosulfate in water (20g in 100mL), stirred for 15min and treated drop-wise with concentrated HCl to a constant pH of 1. The mixture was stirred for 15min and filtered to collect a white solid that was washed repeatedly with water and dried to constant mass (14.7g, 69%).

[0623] **Part B.** Preparation of 1-bromo-3,5-diiodo-2-methoxybenzene.

[0624] A 500mL round-bottom flask was charged with the product from **Part A** (14.7g, 34.6mmol), iodomethane (2.70ml, 43.3mmol), and sodium hydroxide (2.101ml, 39.8mmol) in acetone (96ml) to give a tan solution. The mixture was stirred for 24h and concentrated. The residue was dissolved in ethyl acetate, washed with water and saturated sodium chloride, dried over sodium sulfate, filtered and concentrated to give a white solid. The solid was recrystallized from hot hexane to give a white solid that was collected by filtration (12.3g, 81%).


[0625] **Part C.** Preparation of 1-(3-bromo-5-iodo-4-methoxyphenyl)pyrimidine-2,4(1H,3H)-dione.

[0626] A 250mL round-bottom flask was charged with the product from **Part B** (8.09g, 18.44mmol), pyrimidine-2,4(1H,3H)-dione(2.273g, 20.28mmol), N-(2-cyanophenyl)picolinamide (0.823g, 3.69mmol), copper (I) iodide (0.351g, 1.844mmol) and potassium phosphate (8.22g, 38.7mmol) in DMSO (70ml). The mixture was sealed, sparged with nitrogen for 15min and heated at 60°C for 16h. The mixture was partitioned with ethyl acetate and water. The organic layer was washed with 1M HCl, water, brine, dried with sodium sulfate, and filtered. The filtrate was treated with 3-mercaptopropyl functionalized silica gel (Aldrich catalog #538086), filtered through celite and evaporated to give an off-white solid (3.92g, 50%).

[0627] **Part D.** Preparation of (E)-N-(4-(3-bromo-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.


[0628] To a 100ml round-bottom flask was added the product from **Part C** (846mg, 2.0mmol), the product from **Example 13B, Part B** (482mg, 2.000mmol), potassium phosphate (892mg, 4.20mmol), 1,3,5,7-tetramethyl-6-phenyl-2,4,8-trioxa-6-phosphaadamante (PA-Ph) (CAS 97739-46-3) (17.54mg, 0.060mmol) and tris(dibenzylideneacetone) dipalladium(0)(18.31mg, 0.020mmol) in THF (12.0ml) and water (4.0ml). The flask was sealed and the mixture was sparged with nitrogen for 5min and stirred at ambient temperature for 72h. The mixture was partitioned with ethyl acetate and 1M HCl. The organic layer was washed with saturated sodium bicarbonate, brine, dried with sodium sulfate and filtered. The filtrate was treated with 3-mercaptopropyl functionalized silica gel, filtered and evaporated. The residue was triturated with a minimal amount of methanol/CH<sub>2</sub>Cl<sub>2</sub> to give the title compound as a white solid (595mg, 60%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ ppm 3.03 (s, 3 H) 3.82 (s, 3 H) 5.69(dd, J=7.72, 1.50 Hz, 1H) 7.24 (d, J=8.46 Hz, 2 H) 7.35 (m, 2 H) 7.61(d, J=8.46 Hz, 2 H) 7.69 (d, J=2.21 Hz, 1H) 7.78 (d, J=8.09 Hz, 1H) 7.87 (d, J=2.21 Hz, 1H) 9.90 (s, 1H) 11.50 (s, 1H). MS (ESI-) m/z 490,492 (M-H)+.

[0629] **Example 16.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(thiophen-2-yl)styryl)phenyl)methanesulfonamide (compound **IB-L1-1.48**).




**[0630]** To a 5ml microwave tube was added the product from **Example 15, Part D** (40mg, 0.081mmol), thiophen-2-ylboronic acid (10.40mg, 0.081mmol), 1,1'-bis(di-tert-butylphosphino)ferrocene palladium dichloride (2.65mg, 4.06 $\mu$ mol) and potassium phosphate (34.5mg, 0.162mmol) in THF (3.0ml) and water (1.0ml). The vessel was sealed and the mixture was sparged by nitrogen for 5min and heated at 50°C for 3h. The mixture was partitioned with ethyl acetate and 1M HCl. The organic layer was washed with saturated sodium bicarbonate, brine, dried with sodium sulfate and filtered. The filtrate was treated with 3-mercaptopropyl functionalized silica gel, filtered through celite and evaporated. The residue was purified by reverse phase chromatography to give the title compound as a white solid (20mg, 50%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  ppm 3.03 (s, 3 H) 3.70 (s, 3 H) 5.70 (dd, J=7.72, 2.21 Hz, 1 H) 7.18 (dd, J=5.43, 4.05 Hz, 1 H) 7.25 (d, J=8.82 Hz, 2 H) 7.35 (s, 2 H) 7.63 (d, J=8.82 Hz, 2 H) 7.68 (m, 2 H) 7.77 (m, 2 H) 7.83 (d, J=7.72 Hz, 1 H) 9.89 (s, 1 H) 11.49 (d, J=2.21 Hz, 1 H). MS (ESI+) m/z 496 (M+H)<sup>+</sup>.

**[0631] Example 17.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-(furan-2-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.46**).




**[0632]** The title compound was prepared according to the procedure of **Example 16** substituting furan-2-ylboronic acid for thiophen-2-ylboronic acid to give a white solid (22mg, 56%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  ppm 3.03 (s, 3 H) 3.76 (s, 3 H) 5.69 (d, J=7.72 Hz, 1 H) 6.69 (dd, J=3.31, 1.84 Hz, 1 H) 7.08 (d, J=2.57 Hz, 1 H) 7.25 (d, J=8.46 Hz, 2 H) 7.36 (m, 2 H) 7.63 (d, J=8.82 Hz, 2 H) 7.67 (d, J=2.57 Hz, 1 H) 7.77 (d, J=2.57 Hz, 1 H) 7.82 (m, J=7.72 Hz, 2 H) 9.88 (s, 1 H) 11.48 (s, 1 H). MS (ESI+) m/z 497 (M+NH<sub>4</sub>)<sup>+</sup>.

**[0633] Example 18.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(pyridin-4-yl)styryl)phenyl)methanesulfonamide (compound **IB-L1-1.55**).




[0634] The title compound was prepared according to the procedure of **Example 16** substituting 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine for thiophen-2-ylboronic acid to give a white solid (15mg, 38%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ ppm 3.03 (s, 3 H) 3.49 (s, 3 H) 5.72 (dd, J=7.72, 2.21 Hz, 1 H) 7.25 (d, J=8.46 Hz, 2 H) 7.38 (d, J=4.41 Hz, 2 H) 7.51 (d, J=2.57 Hz, 1 H) 7.63 (d, J=8.82 Hz, 2 H) 7.80 (d, J=5.88 Hz, 2 H) 7.85 (d, J=7.72 Hz, 1 H) 7.97 (d, J=2.57 Hz, 1 H) 8.77 (d, J=6.25 Hz, 2 H) 9.90 (s, 1 H) 11.51 (d, J=2.21 Hz, 1 H). MS (ESI+) m/z 491 (M+H)+.

[0635] **Example 19.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(pyridin-3-yl)styryl)phenyl)methanesulfonamide (compound **IB-L1-1.53**).




[0636] The title compound was prepared according to the procedure of **Example 16** substituting 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine for thiophen-2-ylboronic acid to give a white solid (19mg, 48%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ ppm 3.02 (s, 3 H) 3.45 (s, 3 H) 5.71 (dd, J=8.09, 2.21 Hz, 1 H) 7.24 (d, J=8.46 Hz, 2 H) 7.37 (d, J=2.94 Hz, 2 H) 7.47 (d, J=2.57 Hz, 1 H) 7.63 (m, 3 H) 7.85 (d, J=7.72 Hz, 1 H) 7.93 (d, J=2.57 Hz, 1 H) 8.15 (m, 1 H) 8.68 (dd, J=4.80 Hz, 1.47 Hz, 1 H) 8.86 (d, J=1.84 Hz, 1 H) 9.89 (s, 1 H) 11.50 (d, J=2.21 Hz, 1 H). MS (ESI+) m/z 491 (M+H)+.

[0637] **Example 20.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(thiophen-3-yl)styryl)phenyl)methanesulfonamide (compound **IB-L1-1.47**).




[0638] The title compound was prepared according to the procedure of **Example 16** substituting thiophen-3-ylboronic acid for thiophen-2-ylboronic acid to give a white solid (19mg, 38%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ ppm 3.02 (s, 3 H) 3.55 (s, 3 H) 5.69 (d, J=8.09 Hz, 1 H) 7.24 (d, J=8.46 Hz, 2 H) 7.36 (s, 2 H) 7.55 (m, 2 H) 7.61 (d, J=8.46 Hz, 2 H) 7.67 (dd, J=5.15, 2.94 Hz, 1 H) 7.78 (d, J=2.57 Hz, 1 H) 7.83 (d, J=7.72 Hz, 1 H) 7.93 (dd, J=2.57, 0.96 Hz, 1 H) 9.88 (s, 1 H) 11.48 (s, 1 H). MS (ESI-) m/z 494 (M-H)+.

[0639] **Example 21.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-(furan-3-yl)-2-methoxy-styryl)phenyl)methanesulfonamide (compound **IB-L1-1.50**).



**[0640]** The title compound was prepared according to the procedure of **Example 16** substituting furan-3-ylboronic acid for thiophen-2-ylboronic acid to give a white solid (14mg, 29%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ ppm 3.02 (s, 3 H) 3.69 (s, 3 H) 5.69 (d, J=8.09 Hz, 1 H) 7.05 (dd, J=2.57, 0.90 Hz, 1 H) 7.24 (d, J=8.82 Hz, 2 H) 7.34 (s, 2 H) 7.61 (m, 3 H) 7.74 (d, J=2.57 Hz, 1 H) 7.80 (m, 2 H) 8.25 (s, 1 H) 9.88 (s, 1 H) 11.49 (s, 1 H). MS (ESI-) m/z 478 (M-H)+.

**[0641]** **Example 22.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-(1-hydroxy-2-methylpropan-2-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.45**).



**[0642]** **Part A.** Preparation of 2-(2-hydroxy-3,5-diiodophenyl)acetic acid.

**[0643]** To a 250mL round-bottom flask was added 2-(2-hydroxyphenyl)acetic acid (Aldrich, 3.04g, 20mmol) in acetonitrile (50ml) to give a colorless solution. N-iodosuccinimide (9.00g, 40.0mmol) was added portionwise over 15min to give a red/brown transparent solution that was stirred for 16h. The mixture was concentrated and the resulting solid was triturated in 75mL of water and filtered to collect an orange solid that was dried under vacuum. The crude solid was recrystallized from toluene to give a light orange powder (6.0g, 74%).

**[0644]** **Part B.** Preparation of methyl 2-(3,5-diiodo-2-methoxyphenyl)acetate.

**[0645]** To a 250mL round-bottom flask was added the product from **Part A** (6g, 14.85mmol), potassium carbonate (6.16g, 44.6mmol), and dimethyl sulfate (4.12g, 32.7mmol) in acetone (49.5ml) to give a brown suspension. The suspension was heated at reflux for 16h, cooled, concentrated and the residue was partitioned between EtOAc and water. The EtOAc layer was washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to a brown oil that was chromatographed on a 40g silica cartridge eluting with 3:1 hexane/EtOAc to give a yellow oil (6.0g, 94%).

**[0646]** **Part C.** Preparation of methyl 2-(3,5-diiodo-2-methoxyphenyl)-2-methylpropanoate.

**[0647]** To a 100mL round-bottom flask under nitrogen was added the product from **Part B** (1.728g, 4mmol) in anhydrous THF (20ml) and HMPA (2ml) to give a colorless solution. Methyl iodide (1.251ml, 20.00mmol) was added and the solution was cooled to -40°C. Potassium t-butoxide (12.00ml, 12.00mmol) was added drop-wise and the mixture was stirred at -40 to -20°C for 30min and quenched with 1M HCl to a pH of 1. The mixture was extracted 3 X 40ml with EtOAc. The extracts were combined, washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The crude product was flash chromatographed on a 40g ISCO silica cartridge eluting with 9:1 hexane/EtOAc to give the bis-methylated product as a yellow oil (1.63g, 89%).

**[0648]** **Part D.** Preparation of 2-(3,5-diiodo-2-methoxyphenyl)-2-methylpropanoic acid.

**[0649]** A suspension of the product from **Part C** (2.63g, 5.72mmol) in MeOH (40ml) and THF (40ml) was treated with 4.0M sodium hydroxide (28ml, 112mmol) and heated at 80°C for 48h. The organic solvent was evaporated and the remaining aqueous solution was acidified with 1 M HCl producing a solid that was collected by filtration, washed with water and dried to give the desired carboxylic acid (2.46g, 96%).

**[0650]** **Part E.** Preparation of 2-(3,5-diiodo-2-methoxyphenyl)-2-methylpropan-1-ol.

[0651] A solution of the product from **Part D** (1.00g, 2.242mmol) in THF (40ml) was treated drop-wise with borane THF complex 1.0M (20ml, 20mmol) and then heated at 50°C for 24h. The mixture was treated with methanol (20mL), refluxed for 30min and concentrated. The resulting residue was washed with water, brine, dried with sodium sulfate, filtered and evaporated. The residue was chromatographed on silica gel eluting with hexane/EtOAc (4:1) to give the desired product (810mg, 84%).

[0652] **Part F.** Preparation of tert-butyl(2-(3,5-diiodo-2-methoxyphenyl)-2-methylpropoxy)-dimethylsilane.

[0653] A solution of the product from **Part E** (432mg, 1.000mmol) in DMF (5ml) was treated with tert-butyldimethylchlorosilane (301mg, 2.000mmol), and imidazole (204mg, 3.00mmol) and stirred for 2h. The mixture was partitioned between 1 M HCl and ethyl acetate. The organic layer was washed with saturated sodium bicarbonate, brine, dried with sodium sulfate, filtered and evaporated. The residue was chromatographed on silica gel eluting with hexane/EtOAc (9:1) to give the desired product (522mg, 96%).

[0654] **Part G.** Preparation of 1-(3-(1-(tert-butyldimethylsilyloxy)-2-methylpropan-2-yl)-5-iodo-4-methoxyphenyl)pyrimidine-2,4(1H,3H)-dione.

[0655] To a 50mL round-bottom flask was added the product from **Part F** (520mg, 0.952mmol), pyrimidine-2,4(1H,3H)-dione (117mg, 1.047mmol), N-(2-cyanophenyl)picolinamide (42.5mg, 0.190mmol), copper(I) iodide (18.13mg, 0.095mmol) and potassium phosphate (424mg, 1.999mmol) in DMSO (5ml). The vessel was sealed, sparged with nitrogen and then heated at 60°C for 24h. The mixture was partitioned between 1M HCl and ethyl acetate. The organic layer was washed with saturated sodium bicarbonate, brine, dried with sodium sulfate, and filtered. The filtrate was treated with 3-mercaptopropyl functionalized silica gel, filtered and evaporated. The residue was chromatographed on silica gel eluting with hexane/EtOAc (3:2) to give the product as a solid (285mg, 65%).


[0656] **Part H.** Preparation of (E)-N-(4-(3-(1-(tert-butyldimethylsilyloxy)-2-methylpropan-2-yl)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

[0657] To a 5ml microwave tube was added the product from **Part G** (53mg, 0.1mmol), the product from **Example 13B, Part B** (24mg, 0.1mmol), potassium phosphate (44.0mg, 0.2mmol), PA-Ph (CAS 97739-46-3) (0.87mg, 3.0μmol) and tris(dibenzylideneacetone)palladium(0) (0.9mg, 1μmol) in THF (3.0ml) and water (1.0ml). The vessel was sealed and the mixture was sparged with nitrogen for 5min and then heated at 50°C for 2h. The mixture was partitioned between 1M HCl and ethyl acetate. The organic layer was washed with saturated sodium bicarbonate, brine, dried with sodium sulfate and filtered. The filtrate was treated with 3-mercaptopropyl functionalized silica gel, filtered and evaporated. The residue was chromatographed on silica gel eluting with hexane/EtOAc(1:1) to give a solid (50mg, 83 %).

[0658] **Part I.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-(1-hydroxy-2-methylpropan-2-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

[0659] A solution of the product from **Part H** (120mg, 0.20mmol) in THF (5.0ml) was treated with 1 M TBAF (0.800ml, 0.800mmol) in THF and stirred for 16h. The mixture was partitioned with water and ethyl acetate. The organic layer was washed (3 X brine), dried with sodium sulfate, filtered and evaporated. The residue was chromatographed on silica gel eluting with 4% methanol in CH<sub>2</sub>Cl<sub>2</sub> to give a solid (85mg, 88 %). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ ppm 1.30 (s, 6 H) 3.01 (s, 3 H) 3.62 (d, J=5.52 Hz, 2 H) 3.77 (s, 3 H) 4.67 (t, J=5.33 Hz, 1 H) 5.66 (d, J=8.09 Hz, 1 H) 7.21 (m, 5 H) 7.62 (m, 3 H) 7.72 (d, J=8.09 Hz, 1 H) 9.85 (s, 1 H) 11.42 (s, 1 H). MS (ESI+) m/z 503 (M+NH<sub>4</sub>)<sup>+</sup>.

[0660] **Example 23.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-iodo-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.51**).



[0661] **Part A.** Preparation of 1,3,5-triiodo-2-methoxybenzene.

[0662] In a 250mL pressure vessel was added 2,4,6-triiodophenol (5g, 10.60mmol) in MTBE (60ml) to give a yellow solution. The solution was cooled in an ice bath and 2.0M trimethylsilyldiazomethane (7.95ml, 15.90mmol) was added at a fast drip followed by dropwise addition of methanol (6mL) resulting in calm bubbling. The vessel was sealed and stirred at room temperature for 4h. The reaction solution was partitioned between EtOAc and water and the organic

layer was washed with 1M HCl, saturated NaHCO<sub>3</sub>, and saturated NaCl. The EtOAc was dried (MgSO<sub>4</sub>), filtered and concentrated to give a tan solid that was used without purification (4.8g, 94 %).


**[06663] Part B.** Preparation of 1-(3,5-diiodo-4-methoxyphenyl)pyrimidine-2,4(1H,3H)-dione.

**[06664]** To a 100mL round-bottom flask under N<sub>2</sub> was added the product from **Part A** (3.5g, 7.2mmol), 1H-pyrimidine-2,4-dione (0.97g, 8.64mmol), and potassium phosphate tribasic (3.2g, 15.0mmol) in DMSO (50ml) to give a colorless suspension. N-(2-cyanophenyl)picolinamide (320mg, 1.44mmol) was added and the mix was sparged with N<sub>2</sub> for 5min. Copper(I) iodide (137mg, 0.72mmol) was added and the mix was sparged once again for 10min, placed under N<sub>2</sub> and heated at 60 °C for 18h. The mixture was cooled and partitioned between EtOAc and water adjusting the pH to 1 with HCl. The aqueous layer was extracted 2X with EtOAc. The organics were combined, washed with water, saturated NaHCO<sub>3</sub>, and saturated NaCl, dried (Na<sub>2</sub>SO<sub>4</sub>), treated with 3-mercaptopropyl functionalized silica, filtered and concentrated. The resulting solid was triturated in 2:1 hexane/EtOAc to give an off white powder (2.2g, 62 %).

**[06665] Part C.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-iodo-2-methoxystyryl)phenyl)methanesulfonamide.

**[06666]** In a 5ml microwave tube was mixed the product from **Part B** (141mg, 0.30mmol), the product from **Example 13B, Part B** (72.3mg, 0.300mmol), 1,1'-bis(diphenylphosphino)ferrocene-palladium(II)dichloride CH<sub>2</sub>Cl<sub>2</sub> complex (12.25mg, 0.015mmol) and potassium phosphate (70.0mg, 0.330mmol) in THF (3.0ml) and water (1.0ml). The mixture was sparged with nitrogen for 5min and heated at 50 °C for 2h. The mixture was partitioned with ethyl acetate and 1M HCl. The organic layer was washed with saturated sodium bicarbonate, brine, dried with sodium sulfate and filtered. The filtrate was treated with 3-mercaptopropyl functionalized silica gel, filtered and evaporated. The residue was chromatographed on silica eluting with 5% methanol in CH<sub>2</sub>Cl<sub>2</sub> to give a solid (47mg, 29%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ ppm 3.02 (s, 3 H) 3.77 (s, 3 H) 5.67 (d, J=7.72 Hz, 1 H) 7.28 (m, 4 H) 7.60 (d, J=8.82 Hz, 2 H) 7.76 (d, J=8.09 Hz, 1 H) 7.81 (d, J=2.57 Hz, 1 H) 7.86 (d, J=2.21 Hz, 1 H) 9.90 (s, 1 H) 11.48 (s, 1 H). MS (ESI-) m/z 538 (M-H)+.

**[06667] Example 24.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(methylsulfonyl)styryl)phenyl)methanesulfonamide (compound **IB-L1-1.49**).



**[06668] Part A.** Preparation of 4-nitrobenzene-2-diazo-1-oxide.

**[06669]** To a 250mL round-bottom flask was added 2-amino-4-nitrophenol (6.165g, 40.0mmol) in 48% tetrafluoroboric acid (15ml). Sodium nitrite (2.76g, 40.0mmol) in water (6ml) was added dropwise at 0 °C and the mixture was stirred at room temperature for 30min. The solid was collected by filtration, washed with tetrafluoroboric acid and water. The solid was suspended in acetone (500ml), filtered and dried to give a solid (3.31g, 50%).

**[0670] Part B.** Preparation of 2-(methylthio)-4-nitrophenol.

**[0671]** To a 1L beaker was added the product from **Part A** (2.70g, 16.35mmol) in ice water (250g) to give a brown suspension. Copper (0.520g, 8.18mmol) was added, followed by addition of sodium thiomethoxide (2.292g, 32.7mmol) in water (50ml) slowly. The mixture was stirred at room temperature for 24h. The mixture was filtered and the filtrate was acidified with 1M HCl producing a solid that was collected by filtration and dried (2.53g, 84%).

**[0672] Part C.** Preparation of 2-(methylsulfonyl)-4-nitrophenol.

**[0673]** To a 250mL round-bottom flask was added the product from **Part B** (1.11g, 6.00mmol) in MeOH (20ml) to give a brown suspension. Oxone (7.746g, 12.60mmol) in water (20ml) was added slowly at 0 °C. The mixture was warmed to room temperature, stirred for 1h and partitioned with ethyl acetate and 1M HCl. The organic layer was washed with brine, dried with sodium sulfate, filtered and evaporated. The residue was chromatographed on silica gel eluting with 1% to 5% methanol in CH<sub>2</sub>Cl<sub>2</sub> to give a solid (0.472g, 36%).

**[0674] Part D.** Preparation of 2-iodo-6-(methylsulfonyl)-4-nitrophenol.

**[0675]** To a 50mL round-bottom flask was added the product from **Part C** (470mg, 2.164mmol) in MeOH (10ml) and water (2.5ml). Iodine monochloride (0.130ml, 2.60mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2.0mL) was added drop-wise and the mixture was stirred at room temperature, poured into water (200mL) and stirred for 10min. The resulting solid was collected by

filtration and dried (636mg, 86%).

**[0676] Part E.** Preparation of 1-iodo-2-methoxy-3-(methylsulfonyl)-5-nitrobenzene.

**[0677]** To a 50mL pressure vessel was added the product from **Part D** (630mg, 1.836mmol) in MTBE (6ml) to give a yellow solution. The mixture was cooled in an ice bath and 2M trimethylsilyldiazomethane (1.377ml, 2.75mmol) was added at a fast drip followed by drop-wise addition of MeOH (0.4ml) resulting in calm bubbling. The vessel was sealed and stirred at room temperature for 1h. The mixture was partitioned with ethyl acetate and 1M HCl. The organic layer was washed with saturated sodium bicarbonate, brine, dried with sodium sulfate, filtered and evaporated to give an off-white solid (655mg, 100%).

**[0678] Part F.** Preparation of 3-iodo-4-methoxy-5-(methylsulfonyl)aniline.

**[0679]** To a 250mL round-bottom flask was added the product from **Part E** (0.650g, 1.820mmol), ammonium chloride (0.146g, 2.73mmol), and iron (0.508g, 9.10mmol) in THF/MeOH/water (50ml, 2/2/1). The mixture was refluxed for 2h, cooled and filtered. The filtrate was evaporated and the residue was partitioned with ethyl acetate and water. The organic layer was washed with brine, dried with sodium sulfate, filtered and evaporated to give a solid (590mg, 99%).

**[0680] Part G.** Preparation of (E)-N-(3-iodo-4-methoxy-5-(methylsulfonyl)phenyl)carbamoyl)-3-methoxyacrylamide.

**[0681]** To a 100mL round-bottom flask was added the product from **Part F** (500mg, 1.528mmol) in DMF (15.0ml). The solution was cooled under nitrogen to -20°C and (E)-3-methoxyacryloyl isocyanate (15.28ml, 6.11mmol; prepared as described by Santana, L.; et al. J. Heterocyclic Chem. 1999, 36, 293-295) was added dropwise. The mixture was stirred at this temperature for 15min, then warmed to room temperature and stirred for 45min. The mixture was diluted with ethyl acetate and washed by water (3 x 50ml), brine (3 x 50ml), dried with sodium sulfate, filtered and evaporated. The residue was triturated with ethyl acetate/hexane to give a solid (425mg, 61%).


**[0682] Part H.** Preparation of 1-(3-iodo-4-methoxy-5-(methylsulfonyl)phenyl)pyrimidine-2,4(1H,3H)-dione.

**[0683]** To a 100mL round-bottom flask was added the product from **Part G** (420mg, 0.925mmol) in ethanol (10ml) to give a suspension. Concentrated sulfuric acid (1mL, 18.76mmol) in water (10ml) was added and the mixture was heated at 110°C for 2h. The reaction mix was cooled, diluted with water (50ml) and stirred for 10min. The solid material was collected by filtration, washed with water and dried to give a white solid (325mg, 83%).

**[0684] Part I.** Preparation of (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(methylsulfonyl)styryl)phenyl)methanesulfonamide.

**[0685]** In a 5ml microwave tube was added the product from **Part H** (63.3mg, 0.15mmol), the product from **Example 13B, Part B** (36.2mg, 0.150mmol), potassium phosphate (66.9mg, 0.315mmol), PA-Ph (CAS 97739-46-3) (1.315mg, 4.50μmol) and tris(dibenzylideneacetone)dipalladium(0) (1.374mg, 1.500μmol) in THF (3.0ml) and water (1.0ml). The vessel was sealed and the mixture was sparged with nitrogen for 5min and heated at 50°C for 2h. The mixture was partitioned with ethyl acetate and 1M HCl. The organic layer was washed with saturated sodium bicarbonate, brine, dried with sodium sulfate and filtered. The filtrate was treated with 3-mercaptopropyl functionalized silica gel, filtered and evaporated. The residue was triturated with methanol CH<sub>2</sub>Cl<sub>2</sub> to give a solid (62mg, 84%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ ppm 3.03 (s, 3 H) 3.37 (s, 3 H) 3.94 (s, 3 H) 5.72 (d, J=7.72 Hz, 1 H) 7.26 (m, 3 H) 7.45 (m, 1 H) 7.65 (d, J=8.46 Hz, 2 H) 7.77 (d, J=2.57 Hz, 1 H) 7.81 (d, J=8.09 Hz, 1 H) 8.21 (d, J=2.57 Hz, 1 H) 9.93 (s, 1 H) 11.52 (s, 1 H). MS (ESI+) m/z 509 (M+NH<sub>4</sub>)<sup>+</sup>.

**[0686] Example 25.** Preparation of (E)-methyl 2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-5-(methylsulfonamido)benzoate (compound **IB-L1-1.7**).



**[0687] Part A.** Preparation of methyl 2-((diethoxyphosphoryl)methyl)-5-nitrobenzoate.

**[0688]** To a solution of methyl 2-methyl-5-nitrobenzoate (0.40 g, 2.05mmol) in CCl<sub>4</sub> (20ml) was added *N*-bromosuccinimide (365mg, 2.05mmol) and 2,2'-azobisisobutyronitrile (34mg, 0.21mmol). The resulting mixture was stirred at reflux for 18h, cooled to room temperature and partitioned between EtOAc (50ml) and H<sub>2</sub>O (50ml). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The crude product was purified by column chromatography

on silica gel using 1:3 EtOAc:hexanes as the eluent to give the bromide as an oil (345mg, 61%). The oil was placed in triethylphosphite (5ml) and heated with stirring at 120°C for 3h. The mixture was allowed to cool to room temperature, and the crude product was purified by column chromatography on silica gel using 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as the eluent. The title compound was obtained as an oil (313mg, 75%).


**[0689] Part B.** Preparation of (*E*)-methyl 2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-5-nitrobenzoate.

**[0690]** To a solution of the product from **Part A** (360mg, 1.09mmol) and the product from **Example 13A, Part D** (329mg, 1.09mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (10ml) was added potassium *tert*-butoxide (305mg, 2.72mmol). The resulting dark red solution was stirred at room temperature for 1h, and then poured into 1 N aq. HCl (10ml). The resulting mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (10ml), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo to give a solid. A solution of the solid in thionyl chloride (2.3ml) was heated at 85°C for 30min, and the thionyl chloride was removed in vacuo. The residue was stirred in a 2:1 mixture of CH<sub>2</sub>Cl<sub>2</sub> and MeOH (3ml) for 30min, and evaporated to dryness in vacuo. The crude product was purified by column chromatography on silica gel using 3% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as the eluent to give the title compound (350mg, 69%).

**[0691] Part C.** Preparation of (*E*)-methyl 2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-5-(methylsulfonamido)benzoate.


**[0692]** To a solution of the product from **Part B** (465mg, 0.97mmol) in a 2:2:1 mixture of TBF:MeOH:H<sub>2</sub>O (10ml) was added iron powder (271mg, 4.85mmol), and ammonium chloride (78mg, 1.46mmol). The mixture was heated at 80°C for 45min, filtered through celite, and concentrated to dryness in vacuo. The residue was combined with methanesulfonyl chloride (0.16ml, 2.0mmol) and triethylamine (0.392ml, 4.85mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (10ml) and the resulting mixture was stirred at room temperature for 3h. The mixture was partitioned between 1 N HCl (20ml) and CH<sub>2</sub>Cl<sub>2</sub> (20ml), and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 3% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as the eluent to give the title compound (270mg, 53%). <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 11.42 (s, 1 H) 10.07 (s, 1 H) 7.90 (d, *J*=8.82 Hz, 1 H) 7.66 - 7.79 (m, 3 H) 7.52 (d, *J*=2.57 Hz, 1 H) 7.44 (dd, *J*=8.64, 2.39 Hz, 1 H) 7.14 - 7.26 (m, 2 H) 5.65 (dd, *J*=7.72, 1.84 Hz, 1 H) 3.86 (s, 3 H) 3.79 (s, 3 H) 3.04 (s, 3 H) 1.38 (s, 9 H).

**[0693] Example 26.** Preparation of (*E*)-2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-5-(methylsulfonamido)benzoic acid (compound **IB-L1-1.4**).



**[0694]** A solution of the product from **Example 25** (55mg, 0.104mmol) in THF (1ml) and 1N aq. NaOH (1ml) was stirred in the dark at room temperature for 1.5h. 1N aqueous HCl was added until pH 3, and the resulting mixture was extracted with EtOAc (2 x 2ml). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give the title compound (53mg, 99%). <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 13.22 (br s, 1 H) 11.40 (d, *J*=2.21 Hz, 1 H) 10.02 (s, 1 H) 7.72-7.91 (m, 3 H) 7.68 (d, *J*=2.57 Hz, 1 H) 7.49 (d, *J*=2.57 Hz, 1 H) 7.42 (dd, *J*=8.64, 2.39 Hz, 1 H) 7.21 (d, *J*=2.57 Hz, 1 H) 7.16 (d, *J*=16.18 Hz, 1 H) 5.64 (dd, *J*=7.72, 2.2 Hz, 1 H) 3.79 (s, 3 H) 3.04 (s, 3 H) 1.38 (s, 9 H).

**[0695] Example 27.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-3-(morpholine-4-carbonyl)phenyl)methanesulfonamide (compound **IB-L1-1.23**).




**[0696] Part A.** Preparation of (*E*)-2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-5-(methylsulfonamido)benzoyl chloride.

**[0697]** A solution of the product from **Example 26** (257mg, 0.50mmol) in thionyl chloride (1.5ml) was heated at 85°C for 40min and then concentrated and dried in vacuo to give the title compound as a solid (0.27 g).

**[0698] Part B.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-3-(morpholine-4-carbonyl)phenyl)methanesulfonamide.


**[0699]** To a solution of the product from **Part A** (24mg, 0.045mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (1ml) was added morpholine (0.02ml, 0.226mmol). The mixture was stirred at room temperature for 2h, and then partitioned between 1 N aq. HCl (5ml) and EtOAc (2 x 5ml). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 4% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as the eluent to give the title compound (19mg, 71%). <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ ppm 11.41 (d, *J*=1.84 Hz, 1 H) 10.04 (s, 1 H) 7.85 (d, *J*=8.46 Hz, 1 H) 7.75 (d, *J*=8.09 Hz, 1 H) 7.52 (d, *J*=2.57 Hz, 1 H) 6.99 - 7.34 (m, 5 H) 5.65 (dd, *J*=7.72, 1.84 Hz, 1 H) 3.76 (s, 3 H) 3.56 - 3.71 (m, 4 H) 3.40 - 3.51 (m, 2 H) 3.11 - 3.22 (m, 2 H) 3.06 (s, 3H) 1.38 (s, 9 H).

**[0700] Example 28.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-3-(hydroxymethyl)phenyl)methanesulfonamide (compound **IB-L1-1.10**).




**[0701]** To a solution of the product from **Example 27, Part A** (375mg, 0.705mmol) in anhydrous THF (5ml) at 0°C under N<sub>2</sub> gas was added a 1.0 M solution of lithium *tert*-butoxyaluminumhydride (1.8ml, 1.8mmol) dropwise. The resulting mixture was stirred at 0°C for 30min, and then allowed to warm to room temperature and was stirred for 1h. The mixture was partitioned between 1 N aq. HCl (10ml) and EtOAc (2 x 10ml). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 3% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as the eluent to give the title compound (220mg, 63%). <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ ppm 11.41 (s, 1 H) 9.82 (s, 1 H) 7.73 (t, *J*=8.27 Hz, 2 H) 7.66 (d, *J*=2.57 Hz, 1 H) 7.31 - 7.39 (m, 2 H) 7.20 (d, *J*=2.57 Hz, 1 H) 7.12 - 7.19 (m, 2 H) 5.65 (d, *J*=8.09 Hz, 1 H) 5.28 (t, *J*=5.52 Hz, 1 H) 4.65 (d, *J*=5.52 Hz, 2 H) 3.79 (s, 3 H) 3.00 (s, 3 H) 1.38 (s, 9 H).

**[0702] Example 29.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-3-(methoxymethyl)phenyl)methanesulfonamide (compound **IB-L1-1.13**).



15 [0703] To a solution of the product from **Example 28** (32mg, 0.064mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (1ml) was added thionyl chloride (23 $\mu\text{L}$ , 0.32mmol), and the resulting mixture was stirred at room temperature for 30min. The mixture was partitioned between saturated aq.  $\text{NaHCO}_3$  (5ml) and  $\text{CH}_2\text{Cl}_2$  (5ml) and the organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated. The residue was dissolved in MeOH (1ml), and a solution of 25% NaOMe in MeOH (58 $\mu\text{L}$ , 0.254mmol) was added. The resulting mixture was stirred at 50°C for 2h. The mixture was partitioned between 1 N aq. HCl (10ml) and EtOAc (2 x 10ml). The combined organic layers were dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 3% MeOH in  $\text{CH}_2\text{Cl}_2$  as the eluent to give the title compound (15mg, 46%).  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO-d}_6$ )  $\delta$  11.43 (s, 1 H) 9.86 (s, 1 H) 7.62 - 7.87 (m, 3 H) 7.12 - 7.39 (m, 5 H) 5.66 (d,  $J=7.72$  Hz, 1 H) 4.58 (s, 2 H) 3.78 (s, 3 H) 3.35 (s, 3 H) 3.00 (s, 3 H) 1.38 (s, 9 H).

20 [0704] **Example 30.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-3-((isopentylamino)methyl)phenyl)methanesulfonamide (compound **IB-L1-1.31**).



[0705] **Part A.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-3-formylphenyl)methanesulfonamide.


40 [0706] To a solution of the product from **Example 28** (0.60 g, 1.20mmol) in anhydrous DMA (15ml) was added 2-iodoxybenzoic acid (336mg, 1.20mmol). The mixture was stirred at room temperature for 1h, and then partitioned between EtOAc (20ml) and  $\text{H}_2\text{O}$  (2x 20ml). The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 2% MeOH in  $\text{CH}_2\text{Cl}_2$  as the eluent to give the title compound as a colorless solid (395mg, 66%).  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO-d}_6$ )  $\delta$  ppm 11.43 (d,  $J=2.21$  Hz, 1 H) 10.45 (s, 1 H) 10.15 (s, 1 H) 8.06 (d,  $J=16.18$  Hz, 1 H) 7.97 (d,  $J=8.82$  Hz, 1 H) 7.73 - 7.78 (m, 2 H) 7.69 (d,  $J=2.57$  Hz, 1 H) 7.51 (dd,  $J=8.64, 2.39$  Hz, 1 H) 7.30 (d,  $J=16.18$  Hz, 1 H) 7.26 (d,  $J=2.57$  Hz, 1 H) 5.66 (dd,  $J=7.72, 2.21$  Hz, 1 H) 3.81 (s, 3H) 3.07 (s, 3 H) 1.39 (s, 9 H).

45 [0707] **Part B.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-3-((isopentylamino)methyl)phenyl)methanesulfonamide.

50 [0708] To a solution of the product from **Part A** (50mg, 0.10mmol) and 3-methylbutan-1-amine (12 $\mu\text{L}$ , 0.10mmol) in anhydrous THF (3ml) was added sodium triacetoxyborohydride (32mg, 0.15mmol) and AcOH (9 $\mu\text{L}$ , 0.15mmol). The resulting mixture was stirred at room temperature for 4 h, and then partitioned between  $\text{H}_2\text{O}$  (10ml) and EtOAc (2 x 10ml). The combined organic layers were dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 3% MeOH in  $\text{CH}_2\text{Cl}_2$  as the eluent to give the title compound (37mg, 65%).  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO-d}_6$ )  $\delta$  11.45 (d,  $J=1.84$  Hz, 1 H) 10.04 (s, 1 H) 8.80 - 8.87 (m, 1 H) 7.88 (d,  $J=8.46$  Hz, 1 H) 7.71 - 7.77 (m, 2 H) 7.41 - 7.48 (m, 1 H) 7.37 (d,  $J=2.21$  Hz, 1 H) 7.21 - 7.29 (m, 3 H) 5.67 (dd,  $J=7.91, 2.02$  Hz, 1 H) 4.30 - 4.38 (m, 2 H) 3.80 (s, 3 H) 3.10 (s, 3 H) 2.95 - 3.04 (m, 2 H) 1.49 - 1.67 (m, 3 H) 1.38 (s, 9 H) 0.86 (d,  $J=6.25$  Hz, 6 H).


55 [0709] **Example 31.** Preparation of *N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-5-methoxystyryl)-

3-((*E*)-(methoxyimino)methyl)phenyl)methanesulfonamide (compound **IB-L1-1.19**).




**[0710]** To a solution of the product from **Example 30, Part A** (35mg, 0.070mmol) in EtOH (2ml) was added *O*-methoxyamine hydrochloride (29mg, 0.35mmol) and sodium bicarbonate (30mg, 0.35mmol). The resulting mixture was stirred at 70°C for 2h. To the mixture was added 1 N aq. HCl (1ml) to give a colorless precipitate that was filtered and dried to give the title compound as a colorless solid (24mg, 64%). <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ ppm 11.43 (d, *J*=2.21 Hz, 1 H) 9.94 (s, 1 H) 8.74 (s, 1 H) 7.79 - 7.85 (m, 2 H) 7.76 (d, *J*=7.72 Hz, 1 H) 7.57 - 7.65 (m, 2 H) 7.32 (dd, *J*=8.64, 2.39 Hz, 1 H) 7.23 (d, *J*=2.57 Hz, 1 H) 7.18 (d, *J*=16.18 Hz, 1 H) 5.66 (dd, *J*=7.72, 2.21 Hz, 1 H) 3.93 (s, 3 H) 3.79 (s, 3 H) 3.03 (s, 3 H) 1.38 (s, 9 H).

**[0711] Example 32.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-d)oxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-3-(oxazol-2-yl)phenyl)methanesulfonamide (compound **IB-L1-1.26**).



**[0712]** To a solution of the product from **Example 27, Part A** (80mg, 0.15mmol) in tetramethylene sulfone (1.5ml) was added 1*H*-1,2,3-triazole (10μL, 0.17mmol) and potassium carbonate (73mg, 0.53mmol). The mixture was heated for 35min at 130°C in a microwave reactor. After cooling to room temperature, the mixture was partitioned between 1 N aqueous HCl (10ml) and EtOAc (2 x 10ml). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 3% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as the eluent to give the title compound (37mg, 46%). <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 11.41 (d, *J*=1.84 Hz, 1 H) 10.10 (s, 1 H) 8.29 (d, *J*=1.10 Hz, 1 H) 8.05 (d, *J*=16.18 Hz, 1 H) 7.95 (d, *J*=8.82 Hz, 1 H) 7.82 (d, *J*=2.21 Hz, 1 H) 7.74 (d, *J*=8.09 Hz, 1 H) 7.51 (d, *J*=2.57 Hz, 1 H) 7.46 (d, *J*=0.74 Hz, 1 H) 7.39 (dd, *J*=8.64, 2.39 Hz, 1 H) 7.20 - 7.30 (m, 2 H) 5.65 (dd, *J*=7.91, 2.02 Hz, 1 H) 3.80 (s, 3 H) 3.07 (s, 3 H) 1.38 (s, 9 H).

**[0713] Example 33.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-3-(1*H* imidazol-2-yl)phenyl)methanesulfonamide (compound **IB-L1-1.16**).




15 [0714] To a solution of the product from **Example 30, Part A** (50mg, 0.10mmol) in EtOH (2ml) was added glyoxal (57 $\mu$ L, 0.50mmol) and concentrated aqueous NH<sub>4</sub>OH (70 $\mu$ L, 0.50mmol). The resulting mixture was stirred at room temperature for 16h. To the mixture was added 1 N aq. HCl until pH = 7, and the mixture was partitioned between H<sub>2</sub>O (10ml) and EtOAc (2 x 10ml). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as the eluent to give the title compound (27mg, 50%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.39 (s, 1 H) 11.40 (d, *J*=1.84 Hz, 1 H) 9.98 (s, 1 H) 7.89 (d, *J*=8.82 Hz, 1 H) 7.66 - 7.76 (m, 2 H) 7.38 (t, *J*=2.21 Hz, 2 H) 7.23 - 7.31 (m, 2 H) 7.06 - 7.21 (m, 3 H) 5.63 (dd, *J*=8.09, 1.84 Hz, 1 H) 3.78 (s, 3 H) 3.07 (s, 3 H) 1.37 (s, 9 H).

20

[0715] **Example 34.** Preparation of (*E*)-*tert*-butyl 2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-5-(methylsulfonamido)phenylcarbamate (compound **IB-L1-1.32**).

25




[0716] To a solution of the product from **Example 26** (75mg, 0.146mmol) in *tert*-butanol (4ml) was added diphenylphosphoryl azide (47 $\mu$ L, 0.219mmol) and triethylamine (31 $\mu$ L, 0.219mmol). The resulting mixture was stirred at 80°C for 18h. The cooled mixture was partitioned between H<sub>2</sub>O (10ml) and EtOAc (2 x 10ml). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 3% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as the eluent to give the title compound (16mg, 19%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  11.45 (d, *J*=1.84 Hz, 1 H) 9.86 (s, 1 H) 9.03 (s, 1 H) 7.75 (d, *J*=7.72 Hz, 2 H) 7.55 (d, *J*=2.57 Hz, 1 H) 7.10 - 7.33 (m, 4 H) 7.04 (dd, *J*=8.64, 2.39 Hz, 1 H) 5.66 (dd, *J*=7.91, 2.02 Hz, 1 H) 3.78 (s, 3 H) 3.02 (s, 3 H) 1.45 (s, 9 H) 1.38 (s, 9 H).

40


[0717] **Example 35.** Preparation of (*E*)-*N*-(3-amino-4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.28**).

45



[0718] The procedure described for the preparation of **Example 34** provided the title compound, which was purified by column chromatography on silica gel using 5% methanol in CH<sub>2</sub>Cl<sub>2</sub> as the eluent (6mg, 9%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.44 (d, *J*=2.21 Hz, 1 H) 9.55 (s, 1 H) 7.77 (d, *J*=2.57 Hz, 1 H) 7.75 (d, *J*=8.09 Hz, 1 H) 7.45 (d, *J*=8.46 Hz, 1 H) 7.33 (d, *J*=15.81 Hz, 1 H) 7.15 (d, *J*=2.57 Hz, 1 H) 7.00 (d, *J*=16.18 Hz, 1 H) 6.56 (d, *J*=2.21 Hz, 1 H) 6.44 (dd, *J*=8.46, 2.21 Hz, 1 H) 5.66 (dd, *J*=7.91, 2.02 Hz, 1 H) 5.56 (s, 2 H) 3.78 (s, 3 H) 2.97 (s, 3 H) 1.37 (s, 9 H).

[0719] **Example 36.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-2-fluorophenyl)methanesulfonamide (compound **IB-L1-1.5**).



[0720] **Part A.** Preparation of (3-fluoro-4-nitrophenyl)methanol.

[0721] To a solution of 3-fluoro-4-nitrobenzoic acid (2.0 g, 10.8mmol) in THF (50ml) at 0°C was added BH<sub>3</sub>-Me<sub>2</sub>S complex (2.215ml, 22.15mmol) drop-wise. The mixture was stirred at 0°C for 3h, and was then stirred at 65°C for 18h. To the cooled mixture was added ice (50g), followed by 1N aq. HCl (100ml), and the resulting mixture was extracted with EtOAc (200ml). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo to provide the title compound as a white solid (1.79g, 97%).

[0722] **Part B.** Preparation of 4-(bromomethyl)-2-fluoro-1-nitrobenzene.

[0723] A solution of the product from **Part A** (1.79 g, 10.46mmol), *N*-bromosuccinimide (2.234 g, 12.55mmol) and triphenylphosphine (3.29 g, 12.55mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100ml) and THF (50ml) was stirred at room temperature for 3h. The mixture was partitioned between H<sub>2</sub>O (200ml) and EtOAc (400ml), and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using 1:1 EtOAc:hexanes as the eluent to give the title compound (1.14 g, 47%).


[0724] **Part C.** Preparation of diethyl 3-fluoro-4-nitrobenzylphosphonate.

[0725] The product from **Part B** (1.25 g, 5.34mmol) was subjected to the conditions described for **Example 6, Part B** to provide the title product (0.75 g, 48%).

[0726] **Part D.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-2-fluorophenyl)methanesulfonamide.

[0727] The product from **Part C** (0.193 g, 0.662mmol) was subjected to the conditions described for **Example 13A, Part E, Part F, and Part G** to provide the title product as a colorless solid (15mg, 5%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.43(s, 1H), 9.67(s, 1H), 7.76(d, *J*=8.1Hz, 1H), 7.62(m, 2H), 7.41(m, 2H), 7.38(m, 1H), 7.23(m, 2H), 5.66(dd, *J*=8.0, 2.0Hz, 1H), 3.80(s, 3H), 3.05(s, 3H), 1.38(s, 9H).

[0728] **Example 37.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-2-fluoro-5-methylphenyl)methanesulfonamide (compound **IB-L1-1.15**).



[0729] **Part A.** Preparation of *N*-(4-bromo-2-fluoro-5-methylphenyl)methanesulfonamide.

[0730] To a solution of 4-bromo-2-fluoro-5-methylaniline (2.04 g, 10.0mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (20ml) and pyridine (3.23ml, 40.0mmol) was added methanesulfonyl chloride (0.86ml, 11.0mmol) and the resulting mixture was stirred at room temperature for 2h. Solvent was removed in vacuo, and the residue was partitioned between EtOAc and 1M aq. HCl. The organic layer was washed with saturated aqueous NaHCO<sub>3</sub>, brine and then dried over Na<sub>2</sub>SO<sub>4</sub>. The drying

agent was filtered off, and the filtrate was concentrated to give the title compound as a solid (2.80 g, 99%).

**[0731] Part B.** Preparation of *N*-(4-ethynyl-2-fluoro-5-methylphenyl)methanesulfonamide.

**[0732]** A mixture of the product from Part A (3.0 g, 10.63mmol), triphenylphosphine (0.279 g, 1.06mmol), trimethylsilylacetate (6.0ml, 42.5mmol) and palladium(II) acetate (0.12 g, 0.53mmol) in triethylamine (30ml) and toluene (15ml) under N<sub>2</sub> was heated at 80 °C for 5h. The mixture was allowed to cool to room temperature, and was partitioned between EtOAc and 1M aq. HCl. The organic layer was washed with saturated NaHCO<sub>3</sub> and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The crude product was purified by column chromatography on silica gel using a solvent gradient of 10% to 35% EtOAc in hexanes to give an oil (3.0 g, 94%). To a solution of the oil (3.0 g, 10.0mmol) in MeOH (50ml) was added 1M aq. NaOH (21ml, 21.0mmol), and the resulting mixture was stirred at room temperature for 45min. The mixture was partitioned between EtOAc and 1M aq. HCl, and the organic layer was washed with brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The drying agent was filtered off, and the filtrate was concentrated in vacuo to give the title compound as a solid (2.3 g, quant.).


**[0733] Part C.** Preparation of (*E*)-5-fluoro-2-methyl-4-(methylsulfonamido)styrylboronic acid.

**[0734] The product from Part B (0.20 g, 0.88mmol) was subjected to the conditions described for the preparation of Example 13B, Part B to give the title compound (42mg, 17%).**

**[0735] Part D.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxystyryl)-2-fluoro-5-methylphenyl)methanesulfonamide.


**[0736]** The product from **Part C** (40mg, 0.15mmol) was subjected to the conditions described for the preparation of **Example 13B, Part I** to give the title compound (51mg, 83%). <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 11.42 (d, *J*=2.21 Hz, 1 H) 9.59 (s, 1 H) 7.70 - 7.78 (m, 2 H) 7.66 (d, *J*=11.77 Hz, 1 H) 7.20 - 7.32 (m, 3 H) 5.65 (dd, *J*=7.72, 2.21 Hz, 1 H) 3.79 (s, 3 H) 3.05 (s, 3 H) 2.38 (s, 3 H) 1.38 (s, 9 H).

**[0737] Example 38.** Preparation of methyl 2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxyphenethyl)-5-(methylsulfonamido)benzoate (compound **IB-L5-2-1.1**).



**[0738]** To a solution of the product from **Example 25** (40mg, 0.076mmol) in MeOH (2ml) and THF (2ml) was added 10% Pd/C (20mg) and the resulting mixture was stirred at room temperature under 1 atm H<sub>2</sub> for 16h. The mixture was filtered through celite and concentrated in vacuo to give a solid (27.5mg, 68%). <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 11.39 (s, 1 H) 9.88 (s, 1 H) 7.61 - 7.71 (m, 2 H) 7.28 - 7.36 (m, 2 H) 7.20 (d, *J*=2.57 Hz, 1 H) 7.13 (d, *J*=2.94 Hz, 1 H) 5.64 (d, *J*=7.72 Hz, 1 H) 3.83 (s, 3 H) 3.75 (s, 3 H) 3.14 (dd, *J*=10.30, 5.88 Hz, 2 H) 2.96 (s, 3 H) 2.83 - 2.92 (m, 2 H) 1.34 (s, 9 H).


**[0739] Example 39.** Preparation of *N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-methoxyphenethyl)phenyl)methanesulfonamide (compound **IB-L5-2-1.2**).



**[0740]** The product from **Example 13B, Part M** (200mg, 0.426mmol) was dissolved in MeOH (10ml) followed by the addition of 10% Palladium on activated Carbon (50mg). The resultant mixture was evacuated and a hydrogen balloon

attached then stirred at room temperature for 48h. The mixture was then filtered through celite and the filtrate concentrated under vacuum to an oil which was dissolved in ethanol (4ml) then a 1N solution of aqueous sodium hydroxide (3.8ml, 3.8mmol) was added and the solution stirred at room temperature for 18h. The ethanol was then removed under vacuum and a 1N solution of aqueous hydrochloric acid (4ml) was added to acidify the mixture followed by extraction with EtOAc (2 x 10mL). The organic extracts were combined, dried and purified by column chromatography on silica gel using 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as the eluent to provide the title compound as a colorless solid (82mg, 41%). <sup>1</sup>H NMR (300MHz, DMSO-d<sub>6</sub>) δ 11.39 (s, 1H), 9.60 (s, 1H), 7.65 (d, J=8.1Hz, 1H), 7.23 (m, 3H), 7.17 (m, 3H), 5.64 (d, J=7.7Hz, 1 H), 3.77 (s, 3H), 2.93 (s, 3H), 2.88 (br s, 4H), 1.35 (s, 9H).

**[0741] Example 40.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-ethoxystyryl)phenyl)methanesulfonamide (compound **IB-L1-1.30**).



**[0742] Part A.** Preparation of 2-*tert*-butyl-4-iodophenol.

**[0743]** To a 250mL round-bottom flask was added 2-*tert*-butylphenol (3.76g, 25mmol) in MeOH (50.0ml) to give a colorless solution. Sodium hydroxide (1.200g, 30.0mmol) was added and the mix was stirred until the hydroxide was completely dissolved. The solution was cooled to 0°C and treated with sodium iodide (1.75g, 11.6mmol) followed by drop-wise addition of 10% sodium hypochlorite solution (7.2ml, 11.6mmol). The addition of sodium iodide followed by sodium hypochlorite was repeated twice and the mixture was stirred at 0°C for 30min. The mixture was treated with 10% w/w solution of sodium thiosulfate, stirred for 30min and treated with concentrated HCl dropwise to a constant pH of 1. The mixture was extracted 3X with EtOAc. The extracts were combined, washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated. The crude oil was flash chromatographed on an Isco 80g silica cartridge eluting with hexane to > 4:1 hexane/EtOAc to give a yellow oil (5.2g, 75%).

**[0744] Part B.** Preparation of 2-bromo-6-*tert*-butyl-4-iodophenol.

**[0745]** To a 250mL round-bottom flask was added the product from **Part A** (4.8g, 17.38mmol) and 1,3-dibromo-5,5-dimethylhydantoin (2.61 g, 9.13mmol) in chloroform (87ml) to give an orange solution. The reaction mixture was stirred for 2h resulting in a black solution that was washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. The black oil was flash chromatographed on a 120g Isco silica cartridge eluting with hexane to give a pinkish solid (4.84g, 78%).

**[0746] Part C.** Preparation of 1-bromo-3-*tert*-butyl-2-ethoxy-5-iodobenzene.

**[0747]** To a 50mL round-bottom flask was added the product from **Part B** (888mg, 2.5mmol), ethyl iodide (409mg, 2.63mmol), and potassium carbonate (415mg, 3.00mmol) in acetone (12ml) to give a green suspension. The mixture was heated at reflux for 16h, cooled and concentrated. The residue was partitioned between water and EtOAc. The organic layer was washed twice with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to a red oil. The oil was flash chromatographed on an Isco 40g silica cartridge eluting with hexane to give a clear oil (820mg, 86%).

**[0748] Part D.** Preparation of 1-(3-bromo-5-*tert*-butyl-4-ethoxyphenyl)pyrimidine-2,4(1*H*,3*H*)-dione.

**[0749]** In a 20mL microwave tube under nitrogen flush was added the product from **Part C** (0.4g, 1.044mmol), 1*H*-Pyrimidine-2,4-dione (0.140g, 1.253mmol), and potassium phosphate tribasic (0.465g, 2.193mmol) in DMSO (5ml) to give a colorless suspension. *N*-(2-cyanophenyl)picolinamide (0.047g, 0.209mmol) was added and the mix was sparged with nitrogen for 10min. Copper(I) iodide (0.020g, 0.104mmol) was added and the mix was sparged once again for 10min, placed under nitrogen and heated at 60°C for 18h. The mixture was cooled and partitioned between EtOAc and water adjusting the pH to 1 with HCl. The aqueous layer was extracted 2X with EtOAc. The organics were combined, washed with water, saturated NaHCO<sub>3</sub>, and saturated NaCl. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), stirred with 3-mercaptopropyl functionalized silica for 1h, filtered and concentrated. The crude product was purified by chromatography on an Isco 12g silica cartridge eluting with 2% MeOH in CH<sub>2</sub>Cl<sub>2</sub> to give a white powder (266mg, 69%).

**[0750] Part E.** Preparation of (*E*)-*N*-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2*H*)-yl)-2-ethoxystyryl)phenyl)methanesulfonamide.

**[0751]** A mixture of the product from **Part D** (55.1 mg, 0.15mmol), the product from **Example 13B, Part B** (36.2mg,

0.150mmol), potassium phosphate tribasic (63.7mg, 0.300mmol) and 1,1'-bis(di-tert-butylphosphino)ferrocene palladium dichloride (4.89mg, 7.50 $\mu$ mol) in THF (3ml) water (1ml) was sparged for 10min with nitrogen, and then sealed and heated at 50 °C for 4h. The mixture was cooled to room temperature and diluted into EtOAc. The EtOAc layer was washed with 1M HCl, saturated NaHCO<sub>3</sub>, saturated NaCl, dried (Na<sub>2</sub>SO<sub>4</sub>) and treated simultaneously with mercaptopropyl silica gel, filtered and concentrated. The crude product was purified by column chromatography on silica gel using 2% MeOH in CH<sub>2</sub>Cl<sub>2</sub> as the eluent to give the title compound as a solid (40mg, 55%) m.p. 265-266 °C. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  11.42 (s, 1 H) 9.87 (s, 1 H) 7.76 (d, J=8.09 Hz, 1 H) 7.55 - 7.66 (m, 3 H) 7.17 - 7.27 (m, 5 H) 5.65 (dd, J=7.72, 1.47 Hz, 1 H) 3.89 (q, J=6.74 Hz, 2 H) 3.02 (s, 3 H) 1.45 (t, J=6.99 Hz, 3 H) 1.39 (s, 9 H).

**[0752]** The following compounds were prepared utilizing the above discussion:

**[0753]** (E)-N-(4-(1-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl) prop-1-en-2-yl)phenyl) methanesulfonamide (compound **IA-L1-1.6**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  2.14 (s, 3 H) 2.70 (t, J=6.62 Hz, 2 H) 3.01 (s, 3 H) 3.68 (s, 3 H) 3.78 (t, J=6.62 Hz, 2 H) 6.82 (s, 1 H) 7.10 - 7.17 (m, 2 H) 7.23 (d, J=8.46 Hz, 2 H) 7.59 (d, J=8.46 Hz, 2 H) 9.78 (s, 1 H) 10.32 (s, 1 H).

**[0754]** (Z)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl) methanesulfonamide (compound **IA-L1-1.10**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  10.23 (s, 1 H) 9.74 (s, 1 H) 7.23 (d, J=8.46 Hz, 2H) 7.13 (d, J=2.57 Hz, 1 H) 7.06 (d, J=8.82 Hz, 2 H) 6.92 (d, J=2.57 Hz, 1 H) 6.54 - 6.67 (m, 2 H) 3.78 (s, 3 H) 3.57 (t, J=6.62 Hz, 2 H) 2.96 (s, 3 H) 2.60 (t, J=6.80 Hz, 2 H) 1.34 (s, 9 H).

**[0755]** (E)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl) phenyl)-N-(methylsulfonyl) acetamide (compound **IA-L1-1.11**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  10.36 (s, 1 H) 7.77 (d, J=8.46 Hz, 2 H) 7.56 (d, J=2.21 Hz, 1 H) 7.39 - 7.50 (m, 3 H) 7.25 (d, J=16.55 Hz, 1 H) 7.19 (d, J=2.57 Hz, 1 H) 3.74 - 3.85 (m, 5 H) 3.54 (s, 3 H) 2.72 (t, J=6.62 Hz, 2 H) 1.94 (s, 3 H) 1.38 (s, 9 H).

**[0756]** (E)-1-(3-(4-aminostyryl)-5-tert-butyl-4-methoxyphenyl)dihydropyrimidine-2,4(1H,3H)-dione (compound **IA-L1-1.13**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  1.36 (s, 9 H) 2.70 (t, J=6.62 Hz, 2 H) 3.74 (s, 3 H) 3.77 (t, J=6.62 Hz, 2 H) 5.34 (s, 1 H) 6.57 (d, J=8.46 Hz, 2 H) 6.98 (s, 1 H) 7.07 (d, J=2.21 Hz, 1 H) 7.17 (s, 2 H) 7.30 (d, J=8.09 Hz, 2 H) 7.45 (d, J=2.21 Hz, 1 H) 10.32 (s, 1 H).

**[0757]** (Z)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1 (2H)-yl)-2-methoxystyryl)phenyl) methanesulfonamide (compound **IA-L1-1.20**). <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>):  $\delta$  ppm 1.37 (s, 9H), 2.71 (t, J=6.7Hz, 2H), 3.01 (s, 3H), 3.75 (s, 3H), 3.79 (t, J=6.6Hz, 2H), 7.13 (d, J=16.5Hz, 1H), 7.15 (d, J=2.4Hz, 2H), 7.23 (d, J=8.5Hz, 2H), 7.25 (d, J=16.5 Hz, 1H), 7.51 (d, J=2.4Hz, 1H), 7.61 (d, J=8.6Hz, 2H), 9.80(bs, 1H), 10.30 (s, 1H).

**[0758]** N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin- 1(2H)-yl)-2-methoxyphenyl)-1-fluorovinyl)phenyl)methanesulfonamide (compound **IA-L1-1.21**). (racemic mixture (1:1) of compounds **IA-L1-1.4** and **IA-L1-1.5**).

**[0759]** (E)-1-(3-tert-butyl-4-methoxy-5-(4-nitrostyryl)phenyl)dihydropyrimidine-2,4(1H,3H)-dione (compound **IA-L1-1.22**).

**[0760]** 1-{3-tert-butyl-5-[(Z)-2-chloro-2-(4-nitro-phenyl)-vinyl]-4-methoxy-phenyl}-dihydropyrimidine-2,4-dione (compound **IA-L1-1.23**).

**[0761]** 1-{3-tert-butyl-4-methoxy-5-[(E)-2-(4-nitro-phenyl)-propenyl]-phenyl}-dihydro-pyrimidine-2,4-dione (compound **IA-L1-1.24**).

**[0762]** 1-{3-tert-Butyl-5-[(E)-2-(4-nitro-phenyl)-vinyl]-phenyl}-dihydro-pyrimidine-2,4-dione (compound **IA-L1-1.25**). <sup>1</sup>H NMR (300 MHz, DMSO-D6)  $\delta$  ppm 1.33 (s, 9 H) 2.70 - 2.77 (m, 2 H) 3.84 (t, J=6.80 Hz, 2 H) 7.33 (s, 1 H) 7.49 (d, J=4.04 Hz, 2 H) 7.56 (d, J=5.88 Hz, 2 H) 7.89 (d, J=8.82 Hz, 2 H) 8.25 (d, J=8.82 Hz, 2 H) 10.40 (s, 1 H)

**[0763]** N-(4-{(E)-2-[3-tert-Butyl-5-(dioxo-tetrahydro-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl}-3-methoxy-phenyl)-methanesulfonamide (compound **IA-L1-127**). <sup>1</sup>H NMR (300 MHz, DMSO-D6)  $\delta$  ppm 10.33 (s, 1 H) 9.86 (s, 1 H) 7.64 (d, J=8.46 Hz, 1 H) 7.45 (d, J=2.21 Hz, 1 H) 7.26 (s, 2 H) 7.12 (d, J=2.21 Hz, 1 H) 6.89 (s, 1 H) 6.85 (dd, J=8.46, 1.84 Hz, 1 H) 3.84 (s, 3 H) 3.78 (t, J=6.80 Hz, 2H) 3.74 (s, 3H) 3.04 (s, 3 H) 2.71 (t, J=6.62 Hz, 2 H) 1.37 (s, 9 H)

**[0764]** N-(4-{(E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl}-3-formyl-phenyl)-methanesulfonamide (compound **IB-L1-1.6**). <sup>1</sup>H NMR (300 MHz, DMSO-D6)  $\delta$  ppm 1.39 (s, 9 H) 3.07 (s, 3 H) 3.81 (s, 3H) 5.66 (dd, J=7.72, 2.21 Hz, 1 H) 7.26 (d, J=2.57 Hz, 1 H) 7.30 (d, J=16.18 Hz, 1 H) 7.51 (dd, J=8.64, 2.39 Hz, 1 H) 7.69 (d, J=2.57 Hz, 1 H) 7.73 - 7.78 (m, 2 H) 7.97 (d, J=8.82 Hz, 1H) 8.06 (d, J=16.18 Hz, 1 H) 10.15 (s, 1H) 10.45 (s, 1H) 11.43 (d, J=2.21 Hz, 1 H)

**[0765]** N-[4-{(E)-2-[3-tert-Butyl-5(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl}-3-(hydroxyimino-methyl)-phenyl]-methanesulfonamide (compound **IB-L1-1.8**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  1.38 (s, 9 H) 3.03 (s, 3 H) 3.79 (s, 3 H) 5.66 (dd, J=7.91, 2.02 Hz, 1 H) 7.16 (d, J=15.81 Hz, 1 H) 7.22 (d, J=2.57 Hz, 1 H) 7.26 (dd, J=8.64, 2.39 Hz, 1 H) 7.59 (d, J=16.18 Hz, 1 H) 7.63 (d, J=2.21 Hz, 1 H) 7.73 - 7.83 (m, 3 H) 8.64 (s, 1 H) 9.96 (s, 1 H) 11.42 (d, J=2.21 Hz, 1 H) 11.50 (s, 1 H).

**[0766]** 2-[(E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl]-5-methanesulfonylamino-N-(2-methoxy-ethyl)-benzamide (compound **IB-L1-1.9**). <sup>1</sup>H NMR (300 MHz, DMSO-D6)  $\delta$  1.38 (s, 9 H) 3.05 (s, 3 H) 3.20 (s, 3 H) 3.37 - 3.49 (m, 4 H) 3.78 (s, 3 H) 5.64 (d, J=7.72 Hz, 1 H) 7.15 (d, J=2.57 Hz, 1 H) 7.20 (d, J=2.57 Hz, 1 H) 7.24 (s, 2 H) 7.28 (dd, J=8.46, 2.21 Hz, 1 H) 7.42 (d, J=2.57 Hz, 1 H) 7.73 (d, J=7.72 Hz, 1 H) 7.87 (d, J=8.82

Hz, 1 H) 8.49 (t, J=5.15 Hz, 1 H) 9.99 (s, 1 H) 11.42 (s, 1 H).

**[0767]** 2-((E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-benzoic acid ethyl ester (compound **IB-L1-1.11**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 1.31 (t, J=7.17 Hz, 3 H) 1.38 (s, 9 H) 3.05 (s, 3 H) 3.79 (s, 3 H) 4.33 (q, J=7.23 Hz, 2 H) 5.65 (dd, J=7.72, 2.21 Hz, 1 H) 7.15 - 7.25 (m, 2 H) 7.46 (dd, J=8.64, 2.39 Hz, 1 H) 7.52 (d, J=2.57 Hz, 1 H) 7.68 (d, J=2.57 Hz, 1 H) 7.71 - 7.81 (m, 2 H) 7.90 (d, J=8.46 Hz, 1 H) 10.06 (s, 1 H) 11.42 (d, J=1.84 Hz, 1 H).

**[0768]** N-(4-((E)-2-[3-tert-Butyl-2-chloro-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-phenyl]-vinyl)-phenyl)-methanesulfonamide (compound **IB-L1-1.12**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ ppm 1.49 (s, 9 H) 3.02 (s, 3 H) 5.69 (d, J=7.72 Hz, 1 H) 7.22 (m, 3 H) 7.41 (d, J=2.21 Hz, 1 H) 7.51 (d, J=16.18 Hz, 1 H) 7.59 (d, J=8.82 Hz, 2 H) 7.78 (d, J=2.21 Hz, 1 H) 7.80 (d, J=8.09 Hz, 1 H) 9.90 (s, 1 H) 11.47 (s, 1 H).

**[0769]** 2-((E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-N,N-dimethyl-benzamide (compound **IB-L1-1.14**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 1.37 (s, 9 H) 2.76 (s, 3 H) 3.03 (s, 3 H) 3.05 (s, 3 H) 3.76 (s, 3 H) 5.64 (dd, J=7.91, 1.65 Hz, 1 H) 6.95 (d, J=16.55 Hz, 1 H) 7.02 (d, J=2.21 Hz, 1 H) 7.17 - 7.25 (m, 2 H) 7.27 (dd, J=8.64, 2.39 Hz, 1 H) 7.48 (d, J=2.57 Hz, 1 H) 7.74 (d, J=8.09 Hz, 1 H) 7.82 (d, J=8.82 Hz, 1 H) 10.03 (s, 1 H) 11.39 - 11.43 (m, 1 H).

**[0770]** 2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-N-methyl-benzamide (compound **IB-L1-1.17**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 1.38 (s, 9 H) 2.77 (d, J=4.41 Hz, 3 H) 3.06 (s, 3 H) 3.77 (s, 3 H) 5.64 (dd, J=7.72, 1.84 Hz, 1 H) 7.16-7.33 (m, 5 H) 7.43 (d, J=2.21 Hz, 1 H) 7.73 (d, J=7.72 Hz, 1 H) 7.84 (d, J=8.46 Hz, 1 H) 8.37 (q, J=4.41 Hz, 1 H) 10.00 (s, 1 H) 11.40 (d, J=1.84 Hz, 1 H).

**[0771]** 2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-N-(1,1-dioxo-tetrahydro-1λ<sup>6</sup>-thiophen-3-yl)-5-methanesulfonylamino-N-methyl-benzamide (compound **IB-L1-1.18**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 1.37 (s, 9 H) 2.17 - 2.47 (m, 2 H) 2.70 (s, 3 H) 3.06 (s, 3 H) 3.15 - 3.31 (m, 2 H) 3.36 - 3.51 (m, 2 H) 3.77 (s, 3 H) 5.37 (dt, J=17.74, 8.96 Hz, 1 H) 5.65 (dd, J=7.91, 2.02 Hz, 1 H) 6.93 (d, J=16.18 Hz, 1 H) 7.05 (d, J=2.2 Hz, 1 H) 7.19 - 7.35 (m, 3 H) 7.50 (d, J=2.57 Hz, 1 H) 7.76 (d, J=8.09 Hz, 1 H) 7.87 (d, J=8.82 Hz, 1 H) 10.04 (s, 1 H) 11.38 (d, J=2.21 Hz, 1 H).

**[0772]** N-(4-((E)-2-[3-tert-butyl-5-(5-chloro-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]-vinyl)-phenyl)-methanesulfonamide (Compound **IB-L1-1.20**). <sup>1</sup>H NMR (300 MHz, DMSO-D<sub>6</sub>) δ ppm 11.31 (s, 1H) 9.77 (s, 1H) 7.53 (d, J=8.09 Hz, 1H) 7.23 (d, J=8.46 Hz, 2H) 7.17 (d, J=2.57 Hz, 1 H) 7.06 (d, J=8.82 Hz, 2 H) 7.01 (d, J=2.57 Hz, 1 H) 6.53 - 6.71 (m, 2 H) 5.56 (d, J=7.72 Hz, 1 H) 3.81 (s, 3 H) 2.96 (s, 3 H) 1.35 (s, 9 H)

**[0773]** 2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-benzamide (compound **IB-L1-1.21**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 1.38 (s, 9 H) 3.07 (s, 3 H) 3.78 (s, 3 H) 5.64 (d, J=7.72 Hz, 1 H) 7.18 - 7.34 (m, 5 H) 7.43 (d, J=2.21 Hz, 1 H) 7.54 (s, 1 H) 7.73 (d, J=7.72 Hz, 1 H) 7.84 (d, J=8.46 Hz, 1 H) 7.93 (s, 1 H).

**[0774]** N-(3-(azetidine-1-carbonyl)-4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-phenyl)-methanesulfonamide (compound (compound **IB-L1-1.22**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 1.38 (s, 9 H) 3.07 (s, 3 H) 3.78 (s, 3 H) 5.64 (d, J=7.72 Hz, 1 H) 7.18 - 7.34 (m, 5 H) 7.43 (d, J=2.21 Hz, 1 H) 7.54 (s, 1 H) 7.73 (d, J=7.72 Hz, 1 H) 7.84 (d, J=8.46 Hz, 1 H) 7.93 (s, 1 H).

**[0775]** 2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-N-(2-methoxy-ethyl)-N-methyl-benzamide (compound **IB-L1-1.24**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 1.40 (s, 9 H) 2.81 (s, 3 H) 3.07 (s, 3 H) 3.23 (s, 3 H) 3.29 (t, J=5.33 Hz, 1 H) 3.39 (t, J=4.96 Hz, 1 H) 3.62 (t, J=4.78 Hz, 2 H) 3.82 (s, 3 H) 5.68 (d, J=8.09 Hz, 1 H) 6.96 - 7.07 (m, 1 H) 7.09 - 7.17 (m, 1 H) 7.23 - 7.38 (m, 3 H) 7.49 (dd, J=16.55, 2.57 Hz, 1 H) 7.71 - 7.76 (m, 1 H) 7.83 - 7.94 (m, 1 H).

**[0776]** N-(4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-isopropoxymethyl-phenyl)-methanesulfonamide (compound **IB-L1-1.25**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 1.16 (d, J=5.88 Hz, 6 H) 1.38 (s, 9 H) 3.01 (s, 3H) 3.69 (dt, J=12.13, 6.07 Hz, 1H) 3.79 (s, 3 H) 4.59 (s, 2 H) 5.65 (dd, J=7.91, 2.02 Hz, 1 H) 7.13 - 7.29 (m, 4 H) 7.32 - 7.40 (m, 1 H) 7.59 (d, J=2.57 Hz, 1 H) 7.75 (d, J=8.09 Hz, 2 H) 9.86 (s, 1 H) 11.43 (d, J=1.84 Hz, 1 H).

**[0777]** N-(4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-(pyrrolidine-1-carbonyl)-phenyl)-methanesulfonamide (compound **IB-L1-1.27**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 1.37 (s, 9 H) 1.73 - 1.89 (m, 4 H) 3.03 - 3.12 (m, 5 H) 3.51 (t, J=6.80 Hz, 2 H) 3.76 (s, 3 H) 5.64 (dd, J=7.91, 2.02 Hz, 1 H) 6.99 - 7.06 (m, 1 H) 7.08 (d, J=2.21 Hz, 1 H) 7.19 - 7.31 (m, 3 H) 7.46 (d, J=2.57 Hz, 1 H) 7.75 (d, J=8.09 Hz, 1 H) 7.82 (d, J=8.82 Hz, 1 H) 10.01 (s, 1 H) 11.41 (d, J=2.21 Hz, 1 H).

**[0778]** N-(4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-(3-hydroxyazetidin-1-ylmethyl)-phenyl)-methanesulfonamide (compound **IB-L1-1.29**). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 1.38 (s, 9 H) 2.78 - 2.85 (m, 2 H) 2.99 (s, 3 H) 3.50 - 3.58 (m, 2 H) 3.71 (s, 2 H) 3.79 (s, 3 H) 4.19 (td, J=12.41, 6.07 Hz, 1 H) 5.29 (d, J=6.25 Hz, 1 H) 5.66 (d, J=8.09 Hz, 1 H) 7.10 - 7.18 (m, 2 H) 7.20 (t, J=2.21 Hz, 2 H) 7.35 - 7.42 (m, 1 H) 7.63 (d, J=2.57 Hz, 1 H) 7.69 (d, J=8.46 Hz, 1 H) 7.76 (d, J=7.72 Hz, 1 H) 9.78 (s, 1 H) 11.42 (s, 1 H).

**[0779]** N-(4-((E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-pyrrolidin-1-

ylmethyl-phenyl)-methanesulfonamide (compound **IB-L1-133**).  $^1\text{H NMR}$  (500 MHz,  $\text{DMSO-d}_6$ )  $\delta$  1.40 (s, 9 H) 1.72 - 1.95 (m, 4 H) 2.84 (s, 2 H) 2.88 - 2.98 (m, 2 H) 3.01 (s, 3H) 3.81 (s, 3H) 3.86 - 4.23 (m, 2 H) 5.63 (d,  $J=7.81$  Hz, 1H) 7.17 (d,  $J=15.63$  Hz, 1 H) 7.21 - 7.28 (m, 2 H) 7.32 - 7.38 (m, 1 H) 7.47 (d,  $J=16.11$  Hz, 1 H) 7.53 - 7.59 (m, 1 H) 7.61 (d,  $J=7.81$  Hz, 1 H) 7.70 (d,  $J=6.35$  Hz, 1 H) 9.42 (s, 1 H) 10.88 (s, 1 H).

**[0780]** N-(4-((Z)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-phenyl)-methanesulfonamide (compound **IB-L1-1.34**)  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-D}_6$ )  $\delta$  ppm 11.31(s, 1H) 9.77 (s, 1 H) 7.53 (d,  $J=8.09$  Hz, 1H) 7.23 (d,  $J=8.46$  Hz, 2 H) 7.17 (d,  $J=2.57$  Hz, 1H) 7.06 (d,  $J=8.82$  Hz, 2 H) 7.01 (d,  $J=2.57$  Hz, 1H) 6.53 - 6.71 (m, 2 H) 5.56 (d,  $J=7.72$  Hz, 1 H) 3.81 (s, 3 H) 2.96 (s, 3 H) 1.35 (s, 9 H)


**[0781]** N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)phenethyl)phenyl)methane sulfonamide (compound **IA-L5-2-1.2**).  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-d}_6$ )  $\delta$  1.25 (s, 9 H) 2.69 (t,  $J=6.62$  Hz, 2 H) 2.83 (s, 4 H) 2.91 (s, 3 H) 3.75 (t,  $J=6.62$  Hz, 2 H) 6.99 - 7.21 (m, 7 H) 9.60 (s, 1 H) 10.31 (s, 1 H).

**[0782]** methyl 2-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxyphen ethyl)-5-(methylsulfonamido)benzoate (compound **IB-L5-2-1.1**).  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-d}_6$ )  $\delta$  1.34 (s, 9 H) 2.83 - 2.92 (m, 2 H) 2.96 (s, 3 H) 3.14 (dd,  $J=10.30, 5.88$  Hz, 2 H) 3.75 (s, 3 H) 3.83 (s, 3H) 5.64 (d,  $J=7.72$  Hz, 1 H) 7.13 (d,  $J=2.94$  Hz, 1H) 7.20 (d,  $J=2.57$  Hz, 1 H) 7.28 - 7.36 (m, 2 H) 7.61 - 7.71 (m, 2 H) 9.88 (s, 1 H) 11.39 (s, 1 H)


**[0783]** N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxyphenethyl)phenyl) methanesulfonamide (compound **IB-L5-2-1.2**).  $^1\text{H NMR}$  (300 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  11.39 (s,1H), 9.60 (s,1H), 7.65 (d,  $J=8.1$ Hz,1H), 7.23 (m,3H), 7.17(m,3H), 5.64(d, $J=7.7$ Hz,1 H), 3.77(s,3H), 2.93(s,3H), 2.88(bs,4H), 1.35(s,9H)

**[0784]** The following compounds can be prepared utilizing the above discussion:


**TABLE A**



**IA-L1-1.2**



**IA-L1-1.7**



**IA-L1-1.15**

**[0785] HCV Polymerase Inhibition Assay**

**[0786]** Either two-fold serial dilutions (fractional inhibition assay) or a narrower range of dilutions spanning the  $\text{IC}_{50}$  of the inhibitor (tight binding assay) of the inhibitors were incubated with 20mM Tris-Cl pH 7.4, 2mM  $\text{MnCl}_2$ , 1mM dithiothreitol, 1mM ethylene diamine tetraacetic acid (EDTA), 60 to 125 $\mu\text{M}$  GTP and 20 to 50nM  $\Delta 21$  NS5B (HCV Strain 1B (BK, Genbank accession number M58335, or H77, Genbank accession number AF011751)) for 15min at room temperature. The reaction was initiated by the addition of 20 $\mu\text{M}$  CTP, 20 $\mu\text{M}$  ATP, 1 $\mu\text{M}$   $^3\text{H}$ -UTP (10mCi/ $\mu\text{mol}$ ), 5nM template RNA and 0.1 U/ $\mu\text{l}$  RNase inhibitor (RNasin, Promega), and allowed to proceed for 2 to 4h at room temperature. Reaction volume was 50 $\mu\text{l}$ . The reaction was terminated by the addition of 1 volume of 4mM spermine in 10mM Tris-

EP 2 203 430 B1

Cl pH 8.0, 1mM EDTA. After incubation for at least 15 min at room temperature, the precipitated RNA was captured by filtering through a GF/B filter (Millipore) in a 96 well format. The filter plate was washed three times with 200µl each of 2mM spermine, 10mM Tris-Cl pH 8.0, 1mM EDTA, and 2 times with ethanol. After air-drying, 30µl of Microscint 20 scintillation cocktail (Packard) was added to each well, and the retained cpm were determined by scintillation counting. IC<sub>50</sub> values were calculated by a two-variable nonlinear regression equation using an uninhibited control and a fully inhibited control sample to determine the minimum and maximum for the curve. Tight-binding assays were performed on those compounds exhibiting IC<sub>50</sub> values less than 0.005µM in the fractional inhibition assay in order to more precisely measure the IC<sub>50</sub> values. Retained cpm were plotted vs. inhibitor concentration and fit to equation 1 using non-linear regression (ref. 1) to obtain the IC<sub>50</sub> values:

$$\text{Retained cpm} = A \left[ \sqrt{(IC_{50} + I_t - E_t)^2 + 4 * IC_{50} * E_t} - (IC_{50} + I_t - E_t) \right] \quad (\text{eqn 1})$$

where A=Vmax[S]/2(Km+[S]); I<sub>t</sub>=total inhibitor concentration and E<sub>t</sub>=total active concentration of enzyme.

[0787] Ref. Morrison, J. F. and S. R. Stone. 1985. Approaches to the study and analysis of the inhibition of enzymes by slow- and tight-binding inhibitors. Comments Mol. Cell. Biophys. 2: 347-368.

[0788] The sequence of the template RNA used was: 5'-GGGCGAAUUG GGCCCUUAG AUGCAUGCUC GAGCG-GCCGC CAGUGUGAUG GAUAUCUGCA GAUUUCGCC UUGGUGGUC CAUCUUAGCC CUAGUCACGG CUAGCUGUGA AAGGUCCGUG AGCCGCUUGA CUGCAGAGAG UGCUGAUACU GGCCUCUCUG CAGAU-CAAGUC-3'

[0789] When tested by the above method, the compounds of this invention inhibit HCV polymerase 1A and/or 1B. The legend in the table below is as follows: A - IC<sub>50</sub> ≤ 0.01µM; B -- 0.1µM ≥ IC<sub>50</sub> > 0.01µM; C -- 1 µM ≥ IC<sub>50</sub> > 0.1 µM; and D -- IC<sub>50</sub> > 1 µM; ND - not determined.

Table IC<sub>50</sub>

compound	1a	1b	compound	1a	1b
IA-L1-1.3	A	A	IA-L1-1.4	A	A
IA-L1-1.5	A	B	IA-L1-1.6	A	B
IA-L1-1.9	A	B	IA-L1-1.10	B	B
IA-L1-1.11	B	B	IA-L1-1.12	C	C
IA-L1-1.13	C	C	IA-L1-1.14	D	D
IA-L1-1.16	A	A	IA-L1-1.17	B	B
IA-L1-1.18	C	C	IA-L1-1.20	A	B
IA-L1-1.21	B	B	IA-L1-1.22	C	C
IA-L1-1.23	C	C	IA-L1-1.24	D	D
IA-L1-1.25	D	D	IA-L1-1.26	B	B
IA-L1-1.27	A	B	IB-L1-1.1	A	A
IB-L1-1.2	B	B	IB-L1-1.4	A	A
IB-L1-1.5	A	A	IB-L1-1.6	A	B
IB-L1-1.7	A	B	IB-L1-1.8	A	B
IB-L1-1.9	A	B	IB-L1-1.10	A	B
IB-L1-1.11	A	B	IB-L1-1.12	A	B
IB-L1-1.13	A	B	IB-L1-1.14	A	B
IB-L1-1.15	A	B	IB-L1-1.16	A	B
IB-L1-1.17	A	B	IB-L1-1.18	A	B
IB-L1-1.19	A	B	IB-L1-1.20	A	B
IB-L1-1.21	A	B	IB-L1-1.22	B	B

(continued)

compound	1a	1b	compound	1a	1b
IB-L1-1.23	B	B	IB-L1-1.24	B	B
IB-L1-1.25	B	B	IB-L1-1.26	B	B
IB-L1-1.27	B	B	IB-L1-1.28	B	B
IB-L1-1.29	B	B	IB-L1-1.30	B	B
IB-L1-1.31	B	C	IB-L1-1.32	C	C
IB-L1-1.33	C	C	IB-L1-1.34	D	D
IB-L1-1.45	A	B	IB-L1-1.46	B	B
IB-L1-1.47	B	B	IB-L1-1.48	B	B
IB-L1-1.49	B	C	IB-L1-1.50	B	B
IB-L1-1.51	B	B	IB-L1-1.52	C	C
IB-L1-1.53	D	D	IB-L1-1.55	D	D
IA-L5-2-1.1	B	B	IA-L5-2-1.2	B	B
IB-L5-2-1.1	A	B	IB-L5-2-1.2	B	B
IA-L8-1.1	C	C			

**[0790] HCV Polymerase Replicon Assay**

**[0791]** Two stable subgenomic replicon cell lines were used for compound characterization in cell culture: one derived from genotype 1a-H77 and one derived from genotype 1b-Con1 (obtained from Apath, LLC, St. Louis, MO). All replicon constructs were bicistronic subgenomic replicons similar to those described by Bartenschiager and coworkers (Lohmann et al., Replication of Subgenomic Hepatitis C Virus RNAs in a Hepatoma Cell Line, SCIENCE 285:110-3(1999)). The genotype 1a replicon construct contains NS3-NS5B coding region derived from the H77 strain of HCV (1a-H77) (Blight et al., Efficient Replication of Hepatitis C Virus Genotype 1a RNAs in Cell Culture, J. VIROL. 77:3181-90 (2003)). The replicon also has a firefly luciferase reporter and a neomycin phosphotransferase (Neo) selectable marker. These two coding regions, separated by the FMDV 2a protease, comprise the first cistron of the bicistronic replicon construct, with the second cistron containing the NS3-NS5B coding region with addition of adaptive mutations E1202G, K1691R, K2040R and S22041. The 1b-Con1 replicon construct is identical to the 1a-H77 replicon, except that the NS3-NS5B coding region was derived from the 1b-Con1 strain, and the adaptive mutations are E1202G, T12801 and S22041. Replicon cell lines were maintained in Dulbecco's modified Eagles medium (DMEM) containing 10% (v/v) fetal bovine serum (FBS), 100 IU/ml penicillin, 100mg/ml streptomycin (Invitrogen), and 200mg/ml G418 (Invitrogen).

**[0792]** The inhibitory effects of compounds on HCV replication were determined by measuring activity of the luciferase reporter gene. Briefly, replicon-containing cells were seeded into 96 well plates at a density of 5000 cells per well in 100ul DMEM containing 5% FBS. 16-24h later, the compounds were diluted in dimethyl sulfoxide (DMSO) to generate a 200x stock in a series of eight half-log dilutions. The dilution series was then further diluted 100-fold in the medium containing 5% FBS. Medium with the inhibitor was added to the overnight cell culture plates already containing 100ul of DMEM with 5% FBS. In assays measuring inhibitory activity in the presence of human plasma, the medium from the overnight cell culture plates was replaced with DMEM containing 40% human plasma and 5% FBS. The cells were incubated for three days in the tissue culture incubators and were then lysed for RNA extraction. For the luciferase assay, 30ul of Passive Lysis buffer (Promega) was added to each well, and then the plates were incubated for 15min with rocking to lyse the cells. Luciferin solution (50 to 100ul, Promega) was added to each well, and luciferase activity was measured with a Victor II luminometer (Perkin-Elmer). The percent inhibition of HCV RNA replication was calculated for each compound concentration and the EC<sub>50</sub> value was calculated using nonlinear regression curve fitting to the 4-parameter logistic equation and GraphPad Prism 4 software.

**[0793]** When tested by the above method, the compounds of this invention inhibit HCV polymerase 1A and/or 1B. The legend in the table below is as follows: A -- EC<sub>50</sub> ≤ 0.01uM; B -- 0.1uM ≥ EC<sub>50</sub> > 0.01uM; C - 1uM ≥ EC<sub>50</sub> > 0.1uM; and D - EC<sub>50</sub> > 1uM; ND - not determined.

EP 2 203 430 B1


Table EC<sub>50</sub>

compound	1a	1b	compound	1a	1b
IA-L1-13	B	A	IA-L1-1.4	A	A
IA-L1-1.5	B	A	IA-L1-1.6	B	B
IA-L1-1.9	B	A	IA-L1-1.10	B	B
IA-L1-1.11	A	A	IA-L1-1.12	C	C
IA-L1-1.13	D	C	IA-L1-1.14	D	D
IA-L1-1.16	B	B	IA-L1-1.17	B	B
IA-L1-1.18	C	C	IA-L1-1.20	B	B
IA-L1-1.21	A	A	IA-L1-1.22	D	C
IA-L1-1.23	D	D	IA-L1-1.24	D	D
IA-L1-1.25	ND	ND	IA-L1-1.26	B	B
IA-L1-1.27	B	A	IB-L1-1.1	A	A
IB-L1-1.2	ND	B	IB-L1-1.4	B	A
IB-L1-1.5	B	A	IB-L1-1.6	A	A
IB-L1-1.7	A	A	IB-L1-1.8	B	A
IB-L1-1.9	B	A	IB-L1-1.10	A	A
IB-L1-1.11	B	A	IB-L1-1.12	B	B
IB-L1-1.13	B	A	IB-L1-1.14	B	A
IB-L1-1.15	A	A	IB-L1-1.16	C	B
IB-L1-1.17	B	A	IB-L1-1.18	B	B
IB-L1-1.19	B	A	IB-L1-1.20	B	A
IB-L1-1.21	B	A	IB-L1-1.22	B	A
IB-L1-1.23	C	A	IB-L1-1.24	B	A
IB-L1-1.25	B	A	IB-L1-1.26	B	A
IB-L1-1.27	B	A	IB-L1-1.28	A	A
IB-L1-1.29	C	C	IB-L1-1.30	C	B
IB-L1-1.31	D	D	IB-L1-1.32	C	B
IB-L1-1.33	C	B	IB-L1-1.34	B	A
IB-L1-1.45	B	A	IB-L1-1.46	C	A
IB-L1-1.47	C	B	IB-L1-1.48	C	A
IB-L1-1.49	D	D	IB-L1-1.50	C	B
IB-L1-1.51	D	B	IB-L1-1.52	D	C
IB-L1-1.53	ND	ND	IB-L1-1.55	ND	ND
IA-L5-2-1.1	C	B	IA-L5-2-1.2	C	C
IB-L5-2-1.1	B	A	IB-L5-2-1.2	C	B
IA-L6-1.1	C	B	IA-L8-1.1	C	C

Claims

1. A compound or salt thereof, wherein:

the compound corresponds in structure to formula I:



20 is selected from the group consisting of single carbon-carbon bond and double carbon-carbon bond;

**R<sup>1</sup>** is selected from the group consisting of hydrogen and methyl;

**R<sup>2</sup>** is selected from the group consisting of hydrogen, halo, hydroxy, methyl, cyclopropyl, and cyclobutyl;

**R<sup>3</sup>** is selected from the group consisting of hydrogen, halo, oxo, and methyl;

25 **R<sup>4</sup>** is selected from the group consisting of halo, alkyl, alkenyl, alkynyl, nitro, cyano, azido, alkyloxy, alkenyloxy, alkynyloxy, amino, aminocarbonyl, aminosulfonyl, alkylsulfonyl, carbocyclyl, and heterocyclyl, wherein:

(a) the amino, aminocarbonyl, and aminosulfonyl optionally are substituted with:

30 (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, and alkylsulfonyl, or

(2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl, and

(b) the alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, alkynyloxy, and alkylsulfonyl, optionally are substituted with one or more substituents independently selected from the group consisting of halo, oxo, nitro, cyano, azido, hydroxy, amino, alkyloxy, trimethylsilyl, carbocyclyl, and heterocyclyl, wherein:

the amino optionally is substituted with:

40 (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylcarbonyl, alkylsulfonyl, alkyloxycarbonyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl, or

(2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl, and

(c) the carbocyclyl and heterocyclyl optionally are substituted with up to three substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, halo, oxo, nitro, cyano, azido, hydroxy, amino, alkyloxy, trimethylsilyl, carbocyclyl, and heterocyclyl, wherein:

the amino optionally is substituted with:

50 (1) one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylcarbonyl, alkylsulfonyl, alkyloxycarbonyl, carbocyclyl, heterocyclyl, carbocyclylalkyl, and heterocyclylalkyl, or

(2) two substituents that, together with the amino nitrogen, form a single-ring heterocyclyl;

55 **R<sup>5</sup>** is selected from the group consisting of hydrogen, hydroxy, alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, alkynyloxy, alkylsulfonyloxy, carbocyclylsulfonyloxy, haloalkylsulfonyloxy, and halo;

**L** is selected from the group consisting of C(**R<sup>A</sup>**)=C(**R<sup>B</sup>**), ethylene, and cyclopropyl-1,2-ene;

**R<sup>A</sup>** and **R<sup>B</sup>** are independently selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkyloxy, C<sub>3</sub>-C<sub>5</sub>-cycloalkyl, and halo, wherein:

the C<sub>1</sub>-C<sub>6</sub>-alkyl optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, halo, hydroxy, nitro, oxo, amino, cyano, alkyloxycarbonyl, alkylcarbonyloxy, alkyloxy, carbocyclyl, and heterocyclyl;

**R<sup>6</sup>** is selected from the group consisting of C<sub>5</sub>-C<sub>6</sub>-carbocyclyl and 5-6-membered heterocyclyl, wherein each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of **R<sup>E</sup>**, **R<sup>F</sup>**, **R<sup>G</sup>**, **R<sup>H</sup>**, **R<sup>I</sup>**, **R<sup>J</sup>**, and **R<sup>K</sup>**;

each **R<sup>E</sup>** is independently selected from the group consisting of halo, nitro, hydroxy, oxo, carboxy, cyano, amino, imino, azido, and aldehydo, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl;

each **R<sup>F</sup>** is independently selected from the group consisting of alkyl, alkenyl, and alkynyl, wherein:

each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, imino, nitro, azido, oxo, aminosulfonyl, alkylsulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, imino, aminosulfonyl, aminocarbonyl, carbocyclyl, and heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylsulfonyl, alkenylsulfonyl, alkynylsulfonyl, alkylsulfonylamino, hydroxy, and alkyloxy,

wherein:

amino portion of the alkylsulfonylamino optionally is substituted with a substituent selected from the group consisting of alkyl, alkenyl, and alkynyl;

each **R<sup>G</sup>** is independently selected from the group consisting of carbocyclyl and heterocyclyl, wherein:

each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl;

each **R<sup>H</sup>** is independently selected from the group consisting of alkyloxy, alkenyloxy, alkynyloxy, alkylsulfonyloxy, alkenylsulfonyloxy, and alkynylsulfonyloxy, wherein:

each such substituent optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl;

each **R<sup>I</sup>** is independently selected from the group consisting of alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl, aminocarbonyl, alkyloxycarbonyl, carbocyclylcarbonyl, and heterocyclylcarbonyl, wherein:

(a) the alkylcarbonyl, alkenylcarbonyl, and alkynylcarbonyl optionally are substituted with one or more

substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, and

(b) the aminocarbonyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkyloxyalkyl, carbocyclyl, heterocyclyl, alkylsulfonyl, and alkylsulfonylamino, wherein:

the carbocyclyl and heterocyclyl optionally are substituted with one or two substituents independently selected from the group consisting of halo, alkyl, and oxo;

each  $R^J$  is independently selected from the group consisting of carbocyclylsulfonylamino, heterocyclylsulfonylamino, alkylcarbonylamino, alkenylcarbonylamino, alkynylcarbonylamino, alkyloxycarbonylamino, alkenyloxycarbonylamino, alkynyloxycarbonylamino, alkylsulfonylamino, alkenylsulfonylamino, alkynylsulfonylamino, aminocarbonylamino, alkyloxycarbonylaminoimino, alkylsulfonylaminoimino, alkenylsulfonylaminoimino, and alkynylsulfonylaminoimino, wherein:

(a) the amino portion of such substituents optionally is substituted with a substituent independently selected from the group consisting of carbocyclylalkyl, heterocyclylalkyl, alkylcarbonyloxy, aminocarbonylalkyl, alkyl, alkenyl, alkynyl, alkylcarbonyl, alkenylcarbonyl, alkynylcarbonyl, alkyloxycarbonyl, alkyloxyalkyloxycarbonyl, alkylcarbonyloxyalkyl, and alkylsulfonyl, wherein:

(1) the carbocyclyl portion of the carbocyclylalkyl and the heterocyclyl portion of the heterocyclylalkyl optionally are substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, alkyloxy, alkenyloxy, alkynyloxy, halo, nitro, cyano, azido, oxo, and amino, and

(2) the amino portion of the aminocarbonylalkyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl,

(b) the alkyl, alkenyl, and alkynyl portion of such substituents optionally is substituted with one or more substituents independently selected from the group consisting of carboxy, halo, oxo, amino, alkyloxycarbonyl, alkylcarbonyloxy, hydroxy, alkyloxy, carbocyclyl, heterocyclyl, and cyano, wherein: the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkyloxy, alkenyloxy, and alkynyloxy, wherein:

the alkyl optionally is substituted with one or more hydroxy;

(c) the carbocyclyl and heterocyclyl portions of such substituents optionally are substituted with one or more substituents independently selected from the group consisting of alkyl, alkenyl, alkynyl, carboxy, hydroxy, alkyloxy, alkenyloxy, alkynyloxy, halo, nitro, cyano, azido, and amino, wherein:

the amino optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl; and

each  $R^K$  is independently selected from the group consisting of aminosulfonyl, alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl, wherein:

(a) the alkylsulfonyl, alkenylsulfonyl, and alkynylsulfonyl optionally are substituted with one or more substituents independently selected from the group consisting of carboxy, hydroxy, halo, amino, nitro, azido, oxo, aminosulfonyl, alkyloxycarbonyl, alkenyloxycarbonyl, alkynyloxycarbonyl, alkylcarbonyloxy, alkenylcarbonyloxy, alkynylcarbonyloxy, alkyloxy, alkenyloxy, alkynyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl, wherein:

the amino, aminosulfonyl, and aminocarbonyl optionally are substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl; and

(b) the aminosulfonyl optionally is substituted with one or two substituents independently selected from the group consisting of alkyl, alkenyl, and alkynyl.



nyloxy, alkyloxy, carbocyclyl, heterocyclyl, cyano, and aminocarbonyl; and

(b) the aminosulfonyl optionally is substituted with one or two substituents independently selected alkyl.

3. The compound or salt of any one of claims 1 and 2, wherein  $R^6$  is selected from the group consisting of  $C_5$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl, wherein each such substituent is substituted with one, two, or three substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ ,  $R^I$ ,  $R^J$ , and  $R^K$ .

4. The compound or salt of any one of claims 1-3, wherein:

$R^1$  is hydrogen;

$R^2$  is selected from the group consisting of hydrogen and halo;

$R^3$  is hydrogen;

$R^4$  is tert-butyl;

$R^5$  is selected from the group consisting of hydroxy and methoxy;

$R^A$  is hydrogen; and

$R^B$  is hydrogen.

5. The compound or salt of any one of claims 1-4, wherein:

$R^6$  is selected from the group consisting of  $C_5$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl, wherein each such substituent is substituted with one, two, or three substituents independently selected from the group consisting of  $R^E$ ,  $R^F$ , and  $R^J$ .

6. The compound or salt of any one of claims 1-5, wherein:

$R^6$  is selected from the group consisting of  $C_5$ - $C_6$ -carbocyclyl and 5-6-membered heterocyclyl, wherein each such substituent is substituted with a substituent selected from the group consisting of  $R^F$  and  $R^J$ ;

$R^F$  is alkylsulfonylaminoalkyl; and

$R^J$  is alkylsulfonylamino.

7. The compound or salt of any one of claims 1-6, wherein:

$R^6$  is phenyl substituted with a substituent selected from the group consisting of  $R^F$  and  $R^J$ ;

$R^F$  is alkylsulfonylaminoalkyl; and

$R^J$  is alkylsulfonylamino.

8. The compound or salt of any one of claims 1-7, wherein  $L$  is  $C(R^A)=C(R^B)$ .

9. The compound or salt of any one of claims 1-8, wherein  $\overset{\ast}{\text{---}}\text{---}$  is a double carbon-carbon bond.

10. The compound or salt of claim 1, wherein:

$R^A$  is hydrogen; and

$R^B$  is hydrogen.

11. The compound or salt of claim 1, wherein the compound is selected from the group consisting of:

(E)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl) methanesulfonamide;  
(Z)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-chlorovinyl)phenyl) methanesulfonamide;

(E)-1-(3-tert-butyl-5-(4-fluorostyryl)-4-methoxyphenyl)dihydropyrimidine-2,4(1H,3H)-dione;

(Z)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-fluorovinyl)phenyl) methanesulfonamide;

(E)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-fluorovinyl)phenyl) methanesulfonamide;

(E)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)-2-fluorophenyl) methanesulfonamide;

N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)cyclopropyl)phenyl) meth-

anesulfonamide;  
N-(4-(3-*tert*-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenethyl)phenyl) methanesulfonamide;  
5 (E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)styryl)phenyl)methane sulfonamide;  
(Z)-N-(4-(2-(3-*tert*-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-methoxyvinyl)phenyl) methanesulfonamide;  
(E)-1-(3-*tert*-butyl-4-methoxy-5-styrylphenyl)dihydropyrimidine-2,4(1H,3H)-dione;  
(E)-1-(3-*tert*-butyl-4-methoxy-5-(4-methoxystyryl)phenyl)dihydropyrimidine-2,4(1H,3H)-dione;  
10 (E)-N-(4-(3-*tert*-butyl-5-(5-fluoro-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl) phenyl)methanesulfonamide;  
(E)-N-(4-(3-bromo-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl) methanesulfonamide;  
(E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(thiophen-2-yl)styryl) phenyl)methanesulfonamide;  
15 (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-(furan-2-yl)-2-methoxystyryl)phenyl) methanesulfonamide;  
(E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(pyridin-4-yl)styryl) phenyl)methanesulfonamide;  
(E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(pyridin-3-yl)styryl) phenyl)methanesulfonamide;  
20 (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(thiophen-3-yl)styryl) phenyl)methanesulfonamide;  
(E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-(furan-3-yl)-2-methoxystyryl)phenyl) methanesulfonamide;  
(E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-(1-hydroxy-2-methylpropan-2-yl)-2-methoxystyryl) phenyl)methanesulfonamide;  
25 (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-iodo-2-methoxystyryl)phenyl) methanesulfonamide;  
(E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(methylsulfonyl)styryl) phenyl)methanesulfonamide;  
(E)-methyl 2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-5-(methylsulfonamido) benzoate;  
30 (E)-2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-5-(methylsulfonamido)benzoic acid;  
(E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-(morpholine-4-carbonyl) phenyl)methanesulfonamide;  
35 (E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-(hydroxymethyl)phenyl) methanesulfonamide;  
(E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-(methoxymethyl)phenyl) methanesulfonamide;  
(E)-N-(4(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-((isopentylamino)methyl) phenyl)methanesulfonamide;  
40 N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-((E)-(methoxyimino)methyl) phenyl)methanesulfonamide;  
(E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-(oxazol-2-yl) phenyl) methanesulfonamide;  
45 (E)-N-(4(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-(1H-imidazol-2-yl)phenyl) methanesulfonamide;  
(E)-*tert*-butyl 2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-5-(methylsulfonamido)phenylcarbamate;  
50 (E)-N-(3-amino-4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl) phenyl)methanesulfonamide;  
(E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-2-fluorophenyl) methanesulfonamide;  
(E)-N-(4(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-2-fluoro-5-methylphenyl) methanesulfonamide;  
55 methyl 2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxyphenethyl)-5-(methylsulfonamido)benzoate;  
N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxyphenethyl)phenyl) methanesulfonamide;

(E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-ethoxystyryl)phenyl) methanesulfonamide;  
(E)-N-(4-(1-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl) prop-1-en-2-yl)phenyl) methanesulfonamide;  
5 (Z)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl) methanesulfonamide;  
(E)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)-N-(methylsulfonyl) acetamide;  
(E)-1-(3-(4-aminostyryl)-5-tert-butyl-4-methoxyphenyl)dihydropyrimidine-2,4(1H,3H)-dione;  
(Z)-N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl) methanesulfonamide;  
10 N-(4-(2-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-fluorovinyl)phenyl) methanesulfonamide;  
(E)-1-(3-tert-butyl-4-methoxy-5-(4-nitrostyryl)phenyl)dihydropyrimidine-2,4(1H,3H)-dione;  
1-(3-tert-butyl-5-[(Z)-2-chloro-2-(4-nitro-phenyl)-vinyl]-4-methoxy-phenyl)-dihydropyrimidine-2,4-dione;  
1-(3-tert-butyl-4-methoxy-5-[(E)-2-(4-nitro-phenyl)-propenyl]-phenyl)-dihydropyrimidine-2,4-dione;  
15 1-(3-tert-butyl-5-[(E)-2-(4-nitro-phenyl)-vinyl]-phenyl)-dihydro-pyrimidine-2,4-dione;  
N-(4-((E)-2-[3-tert-butyl-5-(dioxo-tetrahydro-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-methoxy-phenyl)-methanesulfonamide;  
N-(4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-formyl-phenyl)-methanesulfonamide;  
N-[4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-(hydroxyimino-methyl)-phenyl]-methanesulfonamide;  
20 2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-N-(2-methoxy-ethyl)-benzamide;  
2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-benzoic acid ethyl ester;  
25 N-(4-((E)-2-[3-tert-butyl-2-chloro-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-phenyl]-vinyl)-phenyl)-methanesulfonamide;  
2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-N,N-dimethyl-benzamide;  
2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-N-methyl-benzamide ;  
30 2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-N-(1,1-dioxo-tetrahydro-1λ<sup>6</sup>-thiophen-3-yl)-5-methanesulfonylamino-N-methyl-benzamide;  
N-(4-((E)-2-[3-tert-butyl-5-(5-chloro-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]-vinyl)-phenyl)-methanesulfonamide;  
35 2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-benzamide;  
N-(3-(azetidine-1-carbonyl)-4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-phenyl)-methanesulfonamide;  
2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-5-methanesulfonylamino-N-(2-methoxy-ethyl)-N-methyl-benzamide;  
40 N-(4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-isopropoxymethyl-phenyl)-methanesulfonamide;  
N-[4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-(pyrrolidine-1-carbonyl)-phenyl]-methanesulfonamide;  
45 N-[4-1((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-(3-hydroxyazetid-1-ylmethyl)-phenyl]-methanesulfonamide;  
N-(4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-3-pyrrolidin-1-ylmethyl-phenyl)-methanesulfonamide;  
N-(4-((Z)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxy-phenyl]-vinyl)-phenyl)-methanesulfonamide; and  
50 N-(4-(3-tert-butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)phenethyl)phenyl)methanesulfonamide.

12. The compound or salt of claim 1, wherein the compound is (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide.

13. The salt of claim 1, wherein the salt is (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide, sodium salt.

14. The salt of claim 1, wherein the salt is (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide, potassium salt.

15. The salt of claim 1, wherein the salt is (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methanesulfonamide, monopotassium salt.


16. A composition comprising (a) one or more compounds and/or salts recited in any one of claims 1-15; (b) one or more excipients; and, optionally, (c) one or more additional therapeutic agents.

17. One or more compounds and/or salts recited in any one of claims 1-15, and, optionally, one or more additional therapeutic agents, for use in treating hepatitis C in a mammal in need of such treatment.

### Patentansprüche

1. Verbindung oder Salz davon, worin:

die Verbindung in der Struktur der Formel I entspricht:



aus der Gruppe ausgewählt ist, die aus Kohlenstoff-Kohlenstoff-Einfachbindung und Kohlenstoff-Kohlenstoff-Doppelbindung besteht;

R<sup>1</sup> aus der Gruppe ausgewählt ist, die aus Wasserstoff und Methyl besteht;

R<sup>2</sup> aus der Gruppe ausgewählt ist, die aus Wasserstoff, Halogen, Hydroxy, Methyl, Cyclopropyl und Cyclobutyl besteht;

R<sup>3</sup> aus der Gruppe ausgewählt ist, die aus Wasserstoff, Halogen, Oxo und Methyl besteht;

R<sup>4</sup> aus der Gruppe ausgewählt ist, die aus Halogen, Alkyl, Alkenyl, Alkynyl, Nitro, Cyano, Azido, Alkyloxy, Alkenyloxy, Alkinyloxy, Amino, Aminocarbonyl, Aminosulfonyl, Alkylsulfonyl, Carbocyclyl und Heterocyclyl besteht, worin:

(a) das Amino, Aminocarbonyl und Aminosulfonyl optional mit Folgenden substituiert sind:

(1) einem oder zwei Substituenten, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl, Alkynyl und Alkylsulfonyl besteht, oder

(2) zwei Substituenten, die zusammen mit dem Aminostickstoff einen einzelnen Heterocyclylring bilden, und

(b) das Alkyl, Alkenyl, Alkynyl, Alkyloxy, Alkenyloxy, Alkinyloxy und Alkylsulfonyl optional mit einem oder mehreren Substituenten substituiert sind, die unabhängig aus der Gruppe ausgewählt sind, die aus Halogen, Oxo, Nitro, Cyano, Azido, Hydroxy, Amino, Alkyloxy, Trimethylsilyl, Carbocyclyl und Heterocyclyl besteht, worin:

das Amino optional mit Folgenden substituiert ist:

- (1) einem oder zwei Substituenten, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl, Alkynyl, Alkylcarbonyl, Alkylsulfonyl, Alkyloxycarbonyl, Carbocyclyl, Heterocyclyl, Carbocyclalkyl und Heterocyclalkyl besteht, oder  
 (2) zwei Substituenten, die zusammen mit dem Aminostickstoff einen einzelnen Heterocyclring bilden, und

(c) das Carbocyclyl und Heterocyclyl optional mit bis zu drei Substituenten substituiert sind, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl, Alkynyl, Halogen, Oxo, Nitro, Cyano, Azido, Hydroxy, Amino, Alkyloxy, Trimethylsilyl, Carbocyclyl und Heterocyclyl besteht, worin:  
 das Amino optional mit Folgenden substituiert ist:

- (1) einem oder zwei Substituenten, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl, Alkynyl, Alkylcarbonyl, Alkylsulfonyl, Alkyloxycarbonyl, Carbocyclyl, Heterocyclyl, Carbocyclalkyl und Heterocyclalkyl besteht, oder  
 (2) zwei Substituenten, die zusammen mit dem Aminostickstoff einen einzelnen Heterocyclring bilden;

**R<sup>5</sup>** aus der Gruppe ausgewählt ist, die aus Wasserstoff, Hydroxy, Alkyl, Alkenyl, Alkynyl, Alkyloxy, Alkenyloxy, Alkinyloxy, Alkylsulfonyloxy, Carbocyclylsulfonyloxy, Halogenalkylsulfonyloxy und Halogen besteht;

**L** aus der Gruppe ausgewählt ist, die aus C(**R<sup>A</sup>**)=C(**R<sup>B</sup>**), Ethylen und Cyclopropyl-1,2-en besteht;

**R<sup>A</sup>** und **R<sup>B</sup>** unabhängig aus der Gruppe ausgewählt sind, die aus Wasserstoff, C<sub>1</sub>-C<sub>6</sub>-Alkyl, C<sub>1</sub>-C<sub>6</sub>-Alkyloxy, C<sub>3</sub>-C<sub>8</sub>-Cycloalkyl und Halogen besteht, worin:

das C<sub>1</sub>-C<sub>6</sub>-Alkyl optional mit einem oder mehreren Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Carboxy, Halogen, Hydroxy, Nitro, Oxo, Amino, Cyano, Alkyloxycarbonyl, Alkylcarbonyloxy, Alkyloxy, Carbocyclyl und Heterocyclyl besteht;

**R<sup>6</sup>** aus der Gruppe ausgewählt ist, die aus C<sub>5</sub>-C<sub>6</sub>-Carbocyclyl und 5-6-gliedrigem Heterocyclyl besteht, worin jeder dieser Substituenten optional mit einem oder mehreren Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus **R<sup>E</sup>**, **R<sup>F</sup>**, **R<sup>G</sup>**, **R<sup>H</sup>**, **R<sup>I</sup>**, **R<sup>J</sup>** und **R<sup>K</sup>** besteht;

jedes **R<sup>E</sup>** unabhängig aus der Gruppe ausgewählt ist, die aus Halogen, Nitro, Hydroxy, Oxo, Carboxy, Cyano, Amino, Imino, Azido und Aldehydo besteht, worin:

das Amino optional mit einem oder zwei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl und Alkynyl besteht;

jedes **R<sup>F</sup>** unabhängig aus der Gruppe ausgewählt ist, die aus Alkyl, Alkenyl und Alkynyl besteht, worin:

jeder dieser Substituenten optional mit einem oder mehreren Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Carboxy, Hydroxy, Halogen, Amino, Imino, Nitro, Azido, Oxo, Aminosulfonyl, Alkylsulfonyl, Alkyloxycarbonyl, Alkenyloxycarbonyl, Alkinyloxycarbonyl, Alkylcarbonyloxy, Alkenylcarbonyloxy, Alkynylcarbonyloxy, Alkyloxy, Alkenyloxy, Alkinyloxy, Carbocyclyl, Heterocyclyl, Cyano und Aminocarbonyl besteht, worin:

das Amino, Imino, Aminosulfonyl, Aminocarbonyl, Carbocyclyl und Heterocyclyl optional mit einem oder zwei Substituenten substituiert sind, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl, Alkynyl, Alkylsulfonyl, Alkenylsulfonyl, Alkynylsulfonyl, Alkylsulfonylamino, Hydroxy und Alkyloxy besteht, worin:

der Aminoteil des Alkylsulfonylamino optional mit einem Substituenten substituiert ist, der aus der Gruppe ausgewählt ist, die aus Alkyl, Alkenyl und Alkynyl besteht;

jedes **R<sup>G</sup>** unabhängig aus der Gruppe ausgewählt ist, die aus Carbocyclyl und Heterocyclyl besteht, worin:

jeder dieser Substituenten optional mit einem oder mehreren Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl, Alkynyl, Carboxy, Hydroxy, Halogen, Amino, Nitro, Azido, Oxo, Aminosulfonyl, Alkyloxycarbonyl, Alkenyloxycarbonyl, Alkinyloxycarbonyl, Alkylcarbonyloxy,



## EP 2 203 430 B1

das Amino optional mit einem oder zwei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl, Alkynyl, Alkyloxy, Alkenyloxy und Alkinyloxy besteht, worin:

das Alkyl optional mit einem oder mehreren Hydroxy substituiert ist;

(c) das Carbocyclyl- und Heterocyclylteil derartiger Substituenten optional mit einem oder mehreren Substituenten substituiert sind, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl, Alkynyl, Carboxy, Hydroxy, Alkyloxy, Alkenyloxy, Alkinyloxy, Halogen, Nitro, Cyano, Azido und Amino besteht, worin:

das Amino optional mit einem oder zwei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl und Alkynyl besteht; und

jedes  $R^K$  unabhängig aus der Gruppe ausgewählt ist, die aus Aminosulfonyl, Alkylsulfonyl, Alkenylsulfonyl und Alkynylsulfonyl besteht, worin:

(a) das Alkylsulfonyl, Alkenylsulfonyl und Alkynylsulfonyl optional mit einem oder mehreren Substituenten substituiert sind, die unabhängig aus der Gruppe ausgewählt sind, die aus Carboxy, Hydroxy, Halogen, Amino, Nitro, Azido, Oxo, Aminosulfonyl, Alkyloxycarbonyl, Alkenyloxycarbonyl, Alkinyloxycarbonyl, Alkyl-carbonyloxy, Alkenylcarbonyloxy, Alkynylcarbonyloxy, Alkyloxy, Alkenyloxy, Alkinyloxy, Carbocyclyl, Heterocyclyl, Cyano und Aminocarbonyl besteht, worin:

das Amino, Aminosulfonyl und Aminocarbonyl optional mit einem oder zwei Substituenten substituiert sind, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl und Alkynyl besteht; und

(b) das Aminosulfonyl optional mit einem oder zwei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl und Alkynyl besteht.

### 2. Verbindung oder Salz nach Anspruch 1, worin:

$R^2$  aus der Gruppe ausgewählt ist, die aus Wasserstoff und Halogen besteht;  
 $R^3$  aus der Gruppe ausgewählt ist, die aus Wasserstoff und Halogen besteht;  
 $R^4$  aus der Gruppe ausgewählt ist, die aus  $C_1$ - $C_4$ -Alkyl,  $C_3$ - $C_6$ -Carbocyclyl und 5-6-gliedrigem Heterocyclyl besteht, worin:

(a) das  $C_1$ - $C_4$ -Alkyl optional mit bis zu drei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Halogen, Oxo, Hydroxy, Alkyloxy und Trimethylsilyl besteht, und  
(b) das  $C_3$ - $C_6$ -Carbocyclyl und 5-6-gliedrige Heterocyclyl optional mit einem oder zwei Substituenten substituiert sind, die unabhängig aus der Gruppe ausgewählt sind, die Alkyl, Halogen und Alkylsulfonylamino besteht;

$R^5$  aus der Gruppe ausgewählt ist, die aus Wasserstoff, Hydroxy, Alkyloxy und Halogen besteht;  
eines von  $R^A$  und  $R^B$  Wasserstoff ist und das andere aus der Gruppe ausgewählt ist, die aus Wasserstoff, Methyl, Methoxy und Halogen besteht;

$R^6$  aus der Gruppe ausgewählt ist, die aus  $C_5$ - $C_6$ -Carbocyclyl und 5-6-gliedrigem Heterocyclyl besteht, worin jeder dieser Substituenten mit einem, zwei, drei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus  $R^E$ ,  $R^F$ ,  $R^G$ ,  $R^H$ ,  $R^I$ ,  $R^J$  und  $R^K$  besteht;

jedes  $R^E$  unabhängig aus der Gruppe ausgewählt ist, die aus Chlor, Fluor, Nitro, Hydroxy, Oxo, Carboxy, Amino, Imino, Aldehydo und Alkylamino besteht;

jedes  $R^F$  ein unabhängig ausgewähltes Alkyl ist, das optional mit einem Substituenten substituiert ist, der aus der Gruppe ausgewählt ist, die aus Carboxy, Halogen, Amino, Imino und Aminosulfonyl besteht, worin:

das Amino, Imino und Aminosulfonyl optional mit einem oder zwei Substituenten substituiert sind, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkylsulfonyl und Alkylsulfonylamino besteht;

jedes  $R^I$  unabhängig aus der Gruppe ausgewählt ist, die aus Alkylcarbonyl und Aminocarbonyl besteht, worin:

das Aminocarbonyl optional mit einem Substituenten substituiert ist, der aus der Gruppe ausgewählt ist, die aus Alkyl, Alkyloxyalkyl, Alkylsulfonyl und Alkylsulfonylamino besteht;

## EP 2 203 430 B1

jedes  $R^J$  unabhängig aus der Gruppe ausgewählt ist, die aus Alkylsulfonylamino, Alkenylsulfonylamino, Alkylsulfonylamino und Alkylsulfonylaminoimino besteht, worin:

(a) der Aminoteil derartiger Substituenten optional mit einem Substituenten substituiert ist, der unabhängig aus der Gruppe ausgewählt ist, die aus Carbocyclalkyl, Heterocyclalkyl, Alkylcarbonyloxy, Aminocarbonylalkyl, Alkyl, Alkylcarbonyl, Alkyloxy, Alkyloxyalkyl, Alkylcarbonyloxyalkyl und Alkylsulfonyl besteht, worin:

(1) der Carbocyclteil des Carbocyclalkyls und der Heterocyclteil des Heterocyclalkyls optional mit einem oder zwei Substituenten substituiert sind, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Carboxy, Hydroxy, Alkyloxy, Halogen, Nitro, Cyano, Oxo und Amino besteht, und  
(2) der Aminoteil des Aminocarbonylalkyls optional mit einem oder zwei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl, Alkenyl und Alkynyl besteht,

(b) der Alkyl-, Alkenyl- und Alkynylteil derartiger Substituenten optional mit einem oder zwei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Carboxy, Halogen, Oxo, Amino, Alkyloxy, Alkylcarbonyloxy, Hydroxy, Alkyloxy, Carbocycl, Heterocycl und Cyano besteht, worin:

das Amino optional mit einem oder zwei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Alkyl und Alkyloxy besteht, worin:

das Alkyl optional mit einem oder mehreren Hydroxy substituiert ist; und

jedes  $R^K$  unabhängig aus der Gruppe ausgewählt ist, die aus Aminosulfonyl und Alkylsulfonyl besteht, worin:

(a) das Alkylsulfonyl optional mit einem oder zwei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus Carboxy, Hydroxy, Halogen, Amino, Nitro, Oxo, Aminosulfonyl, Alkyloxy, Alkylcarbonyloxy, Alkyloxy, Carbocycl, Heterocycl, Cyano und Aminocarbonyl besteht; und  
(b) das Aminosulfonyl optional mit einem oder zwei Substituenten, unabhängig ausgewähltem Alkyl, substituiert ist.

3. Verbindung oder Salz nach einem der Ansprüche 1 und 2, worin  $R^6$  aus der Gruppe ausgewählt ist, die aus  $C_5$ - $C_6$ -Carbocycl und 5-6-gliedrigem Heterocycl besteht, worin jeder dieser Substituenten mit einem, zwei oder drei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus  $R^E$ ,  $R^F$ ,  $R^I$ ,  $R^J$  und  $R^K$  besteht.

4. Verbindung oder Salz nach einem der Ansprüche 1-3, worin:

$R^1$  Wasserstoff ist;

$R^2$  aus der Gruppe ausgewählt ist, die aus Wasserstoff und Halogen besteht;

$R^3$  Wasserstoff ist;

$R^4$  tert-Butyl ist;

$R^5$  aus der Gruppe ausgewählt ist, die aus Hydroxy und Methoxy besteht;

$R^A$  Wasserstoff ist; und

$R^B$  Wasserstoff ist.

5. Verbindung oder Salz nach einem der Ansprüche 1-4, worin:

$R^6$  aus der Gruppe ausgewählt ist, die aus  $C_5$ - $C_6$ -Carbocycl und 5-6-gliedrigem Heterocycl besteht, worin jeder dieser Substituenten mit einem, zwei oder drei Substituenten substituiert ist, die unabhängig aus der Gruppe ausgewählt sind, die aus  $R^E$ ,  $R^F$  und  $R^J$  besteht.

6. Verbindung oder Salz nach einem der Ansprüche 1-5, worin:

$R^6$  aus der Gruppe ausgewählt ist, die aus  $C_5$ - $C_6$ -Carbocycl und 5-6-gliedrigem Heterocycl besteht, worin jeder dieser Substituenten mit einem Substituenten substituiert ist, der aus der Gruppe ausgewählt ist, die aus  $R^F$  und  $R^J$  besteht;

$R^F$  Alkylsulfonylaminoalkyl ist; und

$R^J$  Alkylsulfonylamino ist.

7. Verbindung oder Salz nach einem der Ansprüche 1-6, worin:

$R^6$  phenyl ist, das mit einem Substituenten substituiert ist, der aus der Gruppe ausgewählt ist, die aus  $R^F$  und  $R^J$  besteht;  
 $R^F$  Alkylsulfonylaminoalkyl ist; und  
 $R^J$  Alkylsulfonylamino ist.

8. Verbindung oder Salz nach einem der Ansprüche 1-7, worin  $L C(R^A)=C(R^B)$  ist.

9. Verbindung oder Salz nach einem der Ansprüche 1-8, worin  $\text{---}^*\text{---}$  eine Kohlenstoff-Kohlenstoff-Doppelbindung ist.

10. Verbindung oder Salz nach Anspruch 1, worin:

$R^A$  Wasserstoff ist; und  
 $R^B$  Wasserstoff ist.

11. Verbindung oder Salz nach Anspruch 1, worin die Verbindung aus der Gruppe ausgewählt ist, die aus Folgenden besteht:

(E)-N-(4-(3-tert-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methansulfonamid ;  
 (2)-N-(4-(2-(3-tert-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-chlorvinyl)phenyl)methansulfonamid ;  
 (E)-1-(3-tert-Butyl-5-(4-fluorstyryl)-4-methoxyphenyl)dihydropyrimidin-2,4(1H,3H)-dion ;  
 (2)-N-(4-(2-(3-tert-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-fluorvinyl)phenyl)methansulfonamid ;  
 (E)-N-(4-(2-(3-tert-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-fluorvinyl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-tert-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)-2-fluorphenyl)methansulfonamid ;  
 N-(4-(2-(3-tert-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)cyclopropyl)phenyl)methansulfonamid ;  
 N-(4-(3-tert-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenethyl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-tert-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)styryl)phenyl)methansulfonamid ;  
 (2)-N-(4-(2-(3-tert-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-methoxyvinyl)phenyl)methansulfonamid ;  
 (E)-1-(3-tert-Butyl-4-methoxy-5-styrylphenyl)dihydropyrimidin-2,4(1H,3H)-dion ;  
 (E)-1-(3-tert-Butyl-4-methoxy-5-(4-methoxystyryl)phenyl)dihydropyrimidin-2,4(1H,3H)-dion ;  
 (E)-N-(4-(3-tert-Butyl-5-(5-fluor-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-Brom-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(5-(2,4-Dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(thiophen-2-yl)styryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(5-(2,4-Dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-(furan-2-yl)-2-methoxystyryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(5-(2,4-Dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(pyridin-4-yl)styryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(5-(2,4-Dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(pyridin-3-yl)styryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(5-(2,4-Dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(thiophen-3-yl)styryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(5-(2,4-Dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-(furan-3-yl)-2-methoxystyryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(5-(2,4-Dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-(1-hydroxy-2-methylpropan-2-yl)-2-methoxystyryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(5-(2,4-Dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-iod-2-methoxystyryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(5-(2,4-Dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxy-3-(methylsulfonyl)styryl)phenyl)methansulfonamid  
 (E)-Methyl-2-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-5-(methylsulfonamido)

benzoat ;  
 (E)-2-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-5-(methylsulfonamido)benzoesäure ;  
 (E)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-(morpholin-4-carbonyl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-(hydroxymethyl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-(methoxymethyl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-((isopentylamino)methyl)phenyl)methansulfonamid ;  
 N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-((E)-(methoxyimino)methyl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-(oxazol-2-yl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-3-(1H-imidazol-2-yl)phenyl)methansulfonamid ;  
 (E)-*tert*-Butyl-2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-5-(methylsulfonamido)phenylcarbamat ;  
 (E)-N-(3-Amino-4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-2-fluorphenyl)methansulfonamid ;  
 (E)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxystyryl)-2-fluor-5-methylphenyl)methansulfonamid ;  
 Methyl-2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxyphenethyl)-5-(methylsulfonamido)benzoat ;  
 N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-methoxyphenethyl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-ethoxystyryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(1-(3-*tert*-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)prop-1-en-2-yl)phenyl)methansulfonamid ;  
 (Z)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methansulfonamid ;  
 (E)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)-N-(methylsulfonyl)acetamid ;  
 (E)-1-(3-(4-Aminostyryl)-5-*tert*-butyl-4-methoxyphenyl)dihydropyrimidin-2,4(1H,3H)-dion ;  
 (Z)-N-(4-(3-*tert*-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methansulfonamid ;  
 N-(4-(2-(3-*tert*-Butyl-5-(2,4-dioxotetrahydropyrimidin-1(2H)-yl)-2-methoxyphenyl)-1-fluorvinyl)phenyl)methansulfonamid ;  
 (E)-1-(3-*tert*-Butyl-4-methoxy-5-(4-nitrostyryl)phenyl)dihydropyrimidin-2,4(1H,3H)-dion ;  
 1-(3-*tert*-Butyl-5-[(2)-2-chlor-2-(4-nitrophenyl)vinyl]-4-methoxyphenyl)dihydropyrimidin-2,4-dion ;  
 1-(3-*tert*-Butyl-4-methoxy-5-[(E)-2-(4-nitrophenyl)propenyl]phenyl)dihydropyrimidin-2,4-dion ;  
 1-(3-*tert*-Butyl-5-[(E)-2-(4-nitrophenyl)vinyl]phenyl)dihydropyrimidin-2,4-dion ;  
 N-(4-{(E)-2-[3-*tert*-Butyl-5-(dioxotetrahydropyrimidin-1-yl)-2-methoxyphenyl]vinyl}-3-methoxyphenyl)methansulfonamid ;  
 N-(4-{(E)-2-[3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl}-3-formylphenyl)methansulfonamid ;  
 N-[4-{(E)-2-[3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl}-3-(hydroxyimino-methyl)phenyl]methansulfonamid ;  
 2-[(E)-2-[3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl]-5-methansulfonylamino-N-(2-methoxyethyl)benzamid ;  
 2-{(E)-2-[3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl}-5-methansulfonylamino-N,N-dimethylbenzamid ;  
 2-[(E)-2-[3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl]-5-methansulfonyl-

amino-N-methylbenzamid ;  
 2-((E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl)-N-(1,1-dioxotetrahydro-1λ<sup>6</sup>-thiophen-3-yl)-5-methansulfonylamino-N-methylbenzamid;  
 N-(4-((E)-2-[3-tert-Butyl-5-(5-chlor-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl)phenyl) methansulfonamid ;  
 2- { (E)- 2-[3- tert- Butyl- 5-(2,4- dioxo- 3,4- dihydro- 2H- pyrimidin- 1- yl)- 2- methoxyphenyl] vinyl}- 5- methansulfonylamino-N-methylbenzamid ;  
 N-(3-(Azetidin-1-carbonyl)-4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl] vinyl)phenyl)methansulfonamid ;  
 2-((E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl)-5-methansulfonylamino-N-(2-methoxyethyl)-N-methylbenzamid ;  
 N-(4-((E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl)-3-isopropoxymethylphenyl)methansulfonamid ;  
 N-[4-((E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl)-3-(pyrrolidin-1-carbonyl)phenyl]methansulfonamid ;  
 N-[4-((E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl)-3-(3-hydroxyazetidin-1-ylmethyl)phenyl]methansulfonamid ;  
 N-(4-((E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-methoxyphenyl]vinyl)-3-pyrrolidin-1-yl-methylphenyl)methansulfonamid ;  
 N-(4- { (Z)- 2-[3- tert- Butyl- 5-(2,4- dioxo- 3,4- dihydro- 2H- pyrimidin- 1- yl)- 2- methoxyphenyl] vinyl} phenyl) methansulfonamid ; und  
 N-(4-(3-tert-Butyl-5-(2,4-dioxotetrahydropyrimidin-1 (2H)-yl) phenethyl) phenyl) methansulfonamid .

12. Verbindung oder Salz nach Anspruch 1, worin die Verbindung (E)-N-(4-(3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methansulfonamid ist.

13. Salz nach Anspruch 1, worin das Salz (E)-N-(4-(3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methansulfonamid-Natriumsalz ist.

14. Salz nach Anspruch 1, worin das Salz (E)-N-(4-(3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methansulfonamid-Kaliumsalz ist.

15. Salz nach Anspruch 1, worin das Salz (E)-N-(4-(3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1(2H)-yl)-2-methoxystyryl)phenyl)methansulfonamid-Monokaliumsalz ist.


16. Zusammensetzung, die Folgendes umfasst: (a) eine oder mehrere Verbindungen und/oder Salze, die in einem der Ansprüche 1-15 vorgetragen wurden, (b) einen oder mehrere Hilfsstoffe und optional (c) ein oder mehrere zusätzliche Therapeutika.

17. eine oder mehrere Verbindungen und/oder Salze, die in einem der Ansprüche 1-15 vorgetragen wurden, und optional ein oder mehrere zusätzliche Therapeutika zur Verwendung bei der Behandlung von Hepatitis C in einem Säuger, der eine derartige Behandlung benötigt.

## Revendications

1. Composé ou sel de celui-ci, dans lequel :

le composé correspond au niveau de sa structure à la formule I :



20 est sélectionné parmi le groupe constitué d'une liaison simple carbone-carbone et d'une double liaison carbone-carbone ;

**R<sup>1</sup>** est sélectionné parmi le groupe constitué d'hydrogène et de méthyle ;

**R<sup>2</sup>** est sélectionné parmi le groupe constitué d'hydrogène, d'halogéno, d'hydroxy, de méthyle, de cyclopropyle et de cyclobutyle ;

**R<sup>3</sup>** est sélectionné parmi le groupe constitué d'hydrogène, d'halogéno, d'oxo et de méthyle ;

25 **R<sup>4</sup>** est sélectionné parmi le groupe constitué d'halogéno, d'alkyle, d'alkényle, d'alkynyle, de nitro, de cyano, d'azido, d'alkyloxy, d'alkényloxy, d'alkynyloxy, d'amino, d'aminocarbonyle, d'aminosulfonyle, d'alkylsulfonyle, de carbocyclyle et d'hétérocyclyle, dans lequel :

30 (a) les amino, aminocarbonyle et aminosulfonyle sont substitués en option par :

(1) un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle et d'alkylsulfonyle, ou

(2) deux substituants qui forment avec l'azote aminé un hétérocyclyle à cycle unique, et

35 (b) les alkyle, alkényle, alkynyle, alkyloxy, alkényloxy, alkynyloxy et alkylsulfonyle sont substitués en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué d'halogéno, d'oxo, de nitro, de cyano, d'azido, d'hydroxy, d'amino, d'alkyloxy, de triméthylsilyle, de carbocyclyle et d'hétérocyclyle, dans lequel :

l'amino est substitué en option par :

40 (1) un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, d'alkylcarbonyle, d'alkylsulfonyle, d'alkyloxycarbonyle, de carbocyclyle, d'hétérocyclyle, de carbocyclylalkyle et d'hétérocyclylalkyle, ou

(2) deux substituants qui forment avec l'azote aminé un hétérocyclyle à cycle unique, et

45 (c) les carbocyclyle et hétérocyclyle sont substitués en option par jusqu'à trois substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, d'halogéno, d'oxo, de nitro, de cyano, d'azido, d'hydroxy, d'amino, d'alkyloxy, de triméthylsilyle, de carbocyclyle et d'hétérocyclyle, dans lequel :

50 l'amino est substitué en option par :

55 (1) un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, d'alkylcarbonyle, d'alkylsulfonyle, d'alkyloxycarbonyle, de carbocyclyle, d'hétérocyclyle, de carbocyclylalkyle et d'hétérocyclylalkyle, ou

(2) deux substituants qui forment avec l'azote aminé un hétérocyclyle à cycle unique ;

**R<sup>5</sup>** est sélectionné parmi le groupe constitué d'hydrogène, d'hydroxy, d'alkyle, d'alkényle, d'alkynyle, d'alkyloxy, d'alkényloxy, d'alkynyloxy, d'alkylsulfonyloxy, de carbocyclylsulfonyloxy, d'halogénoalkylsulfonyloxy et

d'halogéno ;

**L** est sélectionné parmi le groupe constitué de  $C(R^A)=C(R^B)$ , d'éthylène et de cyclopropyl-1,2-ène ;

**RA** et **RB** sont sélectionnés indépendamment parmi le groupe constitué d'hydrogène, de  $C_1-C_6$ -alkyle, de  $C_1-C_6$ -alkyloxy, de  $C_3-C_8$ -cycloalkyle et d'halogéno, dans lequel :

5

le  $C_1-C_6$ -alkyle est substitué en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué de carboxy, d'halogéno, d'hydroxy, de nitro, d'oxo, d'amino, de cyano, d'alkyloxycarbonyle, d'alkylcarbonyloxy, d'alkyloxy, de carbocyclyle et d'hétérocyclyle ;

10

**R<sup>6</sup>** est sélectionné parmi le groupe constitué de  $C_5-C_6$ -carbocyclyle et d'hétérocyclyle à 5 à 6 éléments, dans lequel chaque substituant de ce type est substitué en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué de **RE**, **RF**, **RG**, **RH**, **RI**, **RJ** et **RK** ;

chaque **RE** est sélectionné indépendamment parmi le groupe constitué d'halogéno, de nitro, d'hydroxy, d'oxo, de carboxy, de cyano, d'amino, d'imino, d'azido et d'aldéhydo, dans lequel :

15

l'amino est substitué en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle et d'alkynyle ;

20

chaque **RF** est sélectionné indépendamment parmi le groupe constitué d'alkyle, d'alkényle et d'alkynyle, dans lequel :

25

chaque substituant de ce type est substitué en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué de carboxy, d'hydroxy, d'halogéno, d'amino, d'imino, de nitro, d'azido, d'oxo, d'aminosulfonyle, d'alkylsulfonyle, d'alkyloxycarbonyle, d'alkényloxycarbonyle, d'alkynylloxycarbonyle, d'alkylcarbonyloxy, d'alkénylcarbonyloxy, d'alkynylcarbonyloxy, d'alkyloxy, d'alkényloxy, d'alkynyloxy, de carbocyclyle, d'hétérocyclyle, de cyano et d'aminocarbonyle, dans lequel :

30

les amino, imino, aminosulfonyle, aminocarbonyle, carbocyclyle et hétérocyclyle sont substitués en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, d'alkylsulfonyle, d'alkénylsulfonyle, d'alkynylsulfonyle, d'alkylsulfonylamino, d'hydroxy et d'alkyloxy,

dans lequel :

35

la portion amino de l'alkylsulfonylamino est substituée en option par un substituant sélectionné parmi le groupe constitué d'alkyle, d'alkényle et d'alkynyle ;

40

chaque **RG** est sélectionné indépendamment parmi le groupe constitué de carbocyclyle et d'hétérocyclyle, dans lequel :

45

chaque substituant de ce type est substitué en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, de carboxy, d'hydroxy, d'halogéno, d'amino, de nitro, d'azido, d'oxo, d'aminosulfonyle, d'alkyloxycarbonyle, d'alkényloxycarbonyle, d'alkynylloxycarbonyle, d'alkylcarbonyloxy, d'alkénylcarbonyloxy, d'alkynylcarbonyloxy, d'alkyloxy, d'alkényloxy, d'alkynyloxy, de carbocyclyle, hétérocyclyle, de cyano et d'aminocarbonyle, dans lequel :

50

les amino, aminosulfonyle et aminocarbonyle sont substitués en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, d'alkylsulfonyle, d'alkénylsulfonyle et d'alkynylsulfonyle ;

55

chaque **RH** est sélectionné indépendamment parmi le groupe constitué d'alkyloxy, d'alkényloxy, d'alkynyloxy, d'alkylsulfonyloxy, d'alkénylsulfonyloxy et d'alkynylsulfonyloxy, dans lequel :

chaque substituant de ce type est substitué en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué de carboxy, d'hydroxy, d'halogéno, d'amino, de nitro, d'azido, d'oxo, d'aminosulfonyle, d'alkyloxycarbonyle, d'alkényloxycarbonyle, d'alkynylloxycarbonyle, d'alkylcarbonyloxy, d'alkénylcarbonyloxy, d'alkynylcarbonyloxy, d'alkyloxy, d'alkényloxy, d'alkynyloxy, de carbocyclyle, d'hétérocyclyle, de cyano et d'aminocarbonyle, dans lequel :

les amino, aminosulfonyle et aminocarbonyle sont substitués en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, d'alkylsulfonyle, d'alkénylsulfonyle et d'alkynylsulfonyle ;

5 chaque **R<sup>I</sup>** est sélectionné indépendamment dans le groupe constitué d'alkylcarbonyle, d'alkénylcarbonyle, d'alkynylcarbonyle, d'aminocarbonyle, d'alkyloxycarbonyle, de carbocyclcarbonyle et d'hétérocyclcarbonyle, dans lequel :

10 (a) les alkylcarbonyle, alkénylcarbonyle et alkynylcarbonyle sont substitués en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué de carboxy, d'hydroxy, d'halogéno, d'amino, de nitro, d'azido, d'oxo, d'aminosulfonyle, d'alkyloxycarbonyle, d'alkényloxycarbonyle, d'alkynylcarbonyloxy, d'alkylcarbonyloxy, d'alkénylcarbonyloxy, d'alkynylcarbonyloxy, d'alkyloxy, d'alkényloxy, d'alkynyloxy, de carbocyclyle, d'hétérocyclyle, de cyano et d'aminocarbonyle, et

15 (b) l'aminocarbonyle est substitué en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, d'alkyloxyalkyle, de carbocyclyle, d'hétérocyclyle, d'alkylsulfonyle et d'alkylsulfonylamino, dans lequel :

les carbocyclyle et hétérocyclyle sont substitués en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'halogéno, d'alkyle et d'oxo ;

20 chaque **R<sup>J</sup>** est sélectionné indépendamment parmi le groupe constitué de carbocyclsulfonylamino, d'hétérocyclsulfonylamino, d'alkylcarbonylamino, d'alkénylcarbonylamino, d'alkynylcarbonylamino, d'alkyloxycarbonylamino, d'alkényloxycarbonylamino, d'alkynylsulfonylamino, d'alkénylsulfonylamino, d'alkynylsulfonylamino, d'aminocarbonylamino, d'alkyloxycarbonylaminoimino, d'alkylsulfonylaminoimino, d'alkénylsulfonylaminoimino et d'alkynylsulfonylaminoimino, dans lequel :

30 (a) la portion amino de tels substituants est substituée en option par un substituant sélectionné indépendamment parmi le groupe constitué de carbocyclalkyle, d'hétérocyclalkyle, d'alkylcarbonyloxy, d'aminocarbonylalkyle, d'alkyle, d'alkényle, d'alkynyle, d'alkylcarbonyle, d'alkénylcarbonyle, d'alkynylcarbonyle, d'alkyloxycarbonyle, d'alkyloxyalkyloxycarbonyle, d'alkylcarbonyloxyalkyle et d'alkylsulfonyle, dans lequel :

35 (1) la portion carbocyclyle du carbocyclalkyle et la portion hétérocyclyle de l'hétérocyclalkyle sont substituées en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, de carboxy, d'hydroxy, d'alkyloxy, d'alkényloxy, d'alkynyloxy, d'halogéno, de nitro, de cyano, d'azido, d'oxo et d'amino, et

(2) la portion amino de l'aminocarbonylalkyle est substituée en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle et d'alkynyle,

40 (b) la portion alkyle, alkényle et alkynyle de tels substituants est substituée en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué de carboxy, d'halogéno, d'oxo, d'amino, d'alkyloxycarbonyle, d'alkylcarbonyloxy, d'hydroxy, d'alkyloxy, de carbocyclyle, d'hétérocyclyle et de cyano, dans lequel :

45 l' amino est substitué en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, d'alkyloxy, d'alkényloxy et d'alkynyloxy, dans lequel :

l'alkyle est substitué en option par un ou plusieurs hydroxy ;

50 (c) les portions carbocyclyle et hétérocyclyle de tels substituants sont substituées en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle, d'alkynyle, de carboxy, d'hydroxy, d'alkyloxy, d'alkényloxy, d'alkynyloxy, d'halogéno, de nitro, de cyano, d'azido et d'amino, dans lequel :

55 l' amino est substitué en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle et d'alkynyle ; et

chaque **R<sup>K</sup>** est sélectionné indépendamment parmi le groupe constitué d'aminosulfonyle, d'alkylsulfonyle, d'alkénylsulfonyle et d'alkynylsulfonyle, dans lequel :

(a) les alkylsulfonyle, alkénylsulfonyle et alkynylsulfonyle sont substitués en option par un ou plusieurs substituants sélectionnés indépendamment parmi le groupe constitué de carboxy, d'hydroxy, d'halogéno, d'amino, de nitro, d'azido, d'oxo, d'aminosulfonyle, d'alkyloxycarbonyle, d'alkényloxycarbonyle, d'alkynylloxycarbonyle, d'alkylcarbonyloxy, d'alkénylcarbonyloxy, d'alkynylcarbonyloxy, d'alkyloxy, d'alkényloxy, d'alkynyloxy, de carbocyclyle, d'hétérocyclyle, de cyano et d'aminocarbonyle, dans lequel :

les amino, aminosulfonyle et aminocarbonyle sont substitués en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle et d'alkynyle ; et

(b) l'aminosulfonyle est substitué en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle et d'alkynyle.

2. Composé ou sel selon la revendication 1, dans lequel :

**R<sup>2</sup>** est sélectionné parmi le groupe constitué d'hydrogène et d'halogéno ;

**R<sup>3</sup>** est sélectionné parmi le groupe constitué d'hydrogène et d'halogéno ;

**R<sup>4</sup>** est sélectionné parmi le groupe constitué de C<sub>1</sub>-C<sub>4</sub>-alkyle, de C<sub>3</sub>-C<sub>6</sub>-carbocyclyle et d'hétérocyclyle à 5 à 6 éléments, dans lequel :

(a) le C<sub>1</sub>-C<sub>4</sub>-alkyle est substitué en option par jusqu'à trois substituants sélectionnés indépendamment parmi le groupe constitué d'halogéno, d'oxo, d'hydroxy, d'alkyloxy et de triméthylsilyle, et

(b) les C<sub>3</sub>-C<sub>6</sub>-carbocyclyle et hétérocyclyle à 5 à 6 éléments sont substitués en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'halogéno et d'alkylsulfonylamino ;

**R<sup>5</sup>** est sélectionné parmi le groupe constitué d'hydrogène, d'hydroxy, d'alkyloxy et d'halogéno ;

un de **R<sup>A</sup>** et **R<sup>B</sup>** est hydrogène et l'autre est sélectionné parmi le groupe constitué d'hydrogène, de méthyle, de méthoxy et d'halogéno ;

**R<sup>6</sup>** est sélectionné parmi le groupe constitué de C<sub>5</sub>-C<sub>6</sub>-carbocyclyle et d'hétérocyclyle à 5 à 6 éléments, dans lequel chaque substituant de ce type est substitué par un, deux, trois substituants sélectionnés indépendamment parmi le groupe constitué de **R<sup>E</sup>**, **R<sup>F</sup>**, **R<sup>G</sup>**, **R<sup>H</sup>**, **R<sup>I</sup>**, **R<sup>J</sup>** et **R<sup>K</sup>** ;

chaque **R<sup>E</sup>** est sélectionné indépendamment parmi le groupe constitué de chloro, de fluoro, de nitro, d'hydroxy, d'oxo, de carboxy, d'amino, d'imino, d'aldéhyde et d'alkylamino ;

chaque **R<sup>F</sup>** est un alkyle indépendamment sélectionné substitué en option par un substituant sélectionné parmi le groupe constitué de carboxy, d'halogéno, d'amino, d'imino et d'aminosulfonyle, dans lequel :

les amino, imino et aminosulfonyle sont substitués en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkylsulfonyle et d'alkylsulfonylamino ;

chaque **R<sup>I</sup>** est sélectionné indépendamment parmi le groupe constitué d'alkylcarbonyle et d'aminocarbonyle, dans lequel :

l'aminocarbonyle est substitué en option par un substituant sélectionné parmi le groupe constitué d'alkyle, d'alkyloxyalkyle, d'alkylsulfonyle et d'alkylsulfonylamino ;

chaque **R<sup>J</sup>** est sélectionné indépendamment parmi le groupe constitué d'alkylsulfonylamino, d'alkénylsulfonylamino, d'alkynylsulfonylamino et d'alkylsulfonylaminoimino, dans lequel :

(a) la portion amino de tels substituants est substituée en option par un substituant sélectionné indépendamment parmi le groupe constitué de carbocyclylalkyle, d'hétérocyclylalkyle, d'alkylcarbonyloxy, d'aminocarbonylalkyle, d'alkyle, d'alkylcarbonyle, d'alkyloxycarbonyle, d'alkyloxyalkyloxycarbonyle, d'alkylcarbonyloxyalkyle et d'alkylsulfonyle, dans lequel :

(1) la portion carbocyclyle du carbocyclylalkyle et la portion hétérocyclyle de l'hétérocyclylalkyle sont substituées en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, de carboxy, d'hydroxy, d'alkyloxy, d'halogéno, de nitro, de cyano, d'oxo et d'amino, et  
(2) la portion amino de l'aminocarbonylalkyle est substituée en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle, d'alkényle et d'alkynyle,

(b) la portion alkyle, alkényle et alkynyle de tels substituants est substituée en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué de carboxy, d'halogéno, d'oxo, d'amino, d'alkyloxy, d'alkylcarbonyloxy, d'hydroxy, d'alkyloxy, de carbocyclyle, d'hétérocyclyle et de cyano, dans lequel :

5

l'amino est substitué en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué d'alkyle et d'alkyloxy, dans lequel :

10

l'alkyle est substitué en option par un ou plusieurs hydroxy ; et

chaque  $R^K$  est sélectionné indépendamment parmi le groupe constitué d'aminosulfonyle et d'alkylsulfonyle, dans lequel :

15

(a) l'alkylsulfonyle est substitué en option par un ou deux substituants sélectionnés indépendamment parmi le groupe constitué de carboxy, d'hydroxy, d'halogéno, d'amino, de nitro, d'oxo, d'aminosulfonyle, d'alkyloxy, d'alkylcarbonyloxy, d'alkyloxy, de carbocyclyle, d'hétérocyclyle, de cyano et d'aminocarbonyle ; et

(b) l'aminosulfonyle est substitué en option par un ou deux substituants sélectionnés indépendamment alkyle.

20

3. Composé ou sel selon l'une quelconque des revendications 1 et 2, dans lequel  $R^6$  est sélectionné parmi le groupe constitué de  $C_5$ - $C_6$ -carbocyclyle et d'hétérocyclyle à 5 à 6 éléments, dans lequel chaque substituant de ce type est substitué par un, deux ou trois substituants sélectionnés indépendamment parmi le groupe constitué de  $R^E$ ,  $R^F$ ,  $R^I$ ,  $R^J$  et  $R^K$ .

25

4. Composé ou sel selon l'une quelconque des revendications 1 à 3, dans lequel :

$R^1$  est hydrogène ;

$R^2$  est sélectionné parmi le groupe constitué d'hydrogène et d'halogéno ;

$R^3$  est hydrogène ;

$R^4$  est tert-butyle ;

$R^5$  est sélectionné parmi le groupe constitué d'hydroxy et de méthoxy ;

$R^A$  est hydrogène ; et

$R^B$  est hydrogène.

35

5. Composé ou sel selon l'une quelconque des revendications 1 à 4, dans lequel :

$R^6$  est sélectionné parmi le groupe constitué de  $C_5$ - $C_6$ -carbocyclyle et d'hétérocyclyle à 5 à 6 éléments, dans lequel chaque substituant de ce type est substitué par un, deux ou trois substituants sélectionnés indépendamment parmi le groupe constitué de  $R^E$ ,  $R^F$  et  $R^J$ .

40

6. Composé ou sel selon l'une quelconque des revendications 1 à 5, dans lequel :

$R^6$  est sélectionné parmi le groupe constitué de  $C_5$ - $C_6$ -carbocyclyle et d'hétérocyclyle à 5 à 6 éléments, dans lequel chaque substituant de ce type est substitué par un substituant sélectionné parmi le groupe constitué de  $R^F$  et  $R^J$ ;

$R^F$  est alkylsulfonylaminoalkyle ; et

$R^J$  est alkylsulfonylamino.

45

7. Composé ou sel selon l'une quelconque des revendications 1 à 6, dans lequel :

$R^6$  est phényle substitué par un substituant sélectionné parmi le groupe constitué de  $R^F$  et  $R^J$ ;

$R^F$  est alkylsulfonylaminoalkyle ; et

$R^J$  est alkylsulfonylamino.

50

8. Composé ou sel selon l'une quelconque des revendications 1 à 7, dans lequel  $L$  est  $C(R^A)=C(R^B)$ .

9. Composé ou sel selon l'une quelconque des revendications 1 à 8, dans lequel  $\text{--}\overset{\star}{\text{C}}\text{--}\text{C}\text{--}$  est une double liaison carbone-

carbone.

10. Composé ou sel selon la revendication 1, dans lequel :

5            **R<sup>A</sup>** est hydrogène ; et  
               **R<sup>B</sup>** est hydrogène.

11. Composé ou sel selon la revendication 1, dans lequel le composé est sélectionné parmi le groupe constitué de :

10            (E)-N-(4-(3- tert- butyl- 5-(2,4- dioxotétrahydropyrimidin- 1 (2H)-yl)- 2- méthoxystyryl) phényl) méthanesulfonamide ;  
               (Z)-N-(4-(2-(3- tert- butyl- 5-(2,4- dioxotétrahydropyrimidin- 1 (2H)-yl)- 2- méthoxyphényl)- 1-chlorovinyl) phényl) méthanesulfonamide ;  
               (E)-1-(3-tert-butyl-5-(4-fluorostyryl)-4-méthoxyphényl)dihydropyrimidine-2,4(1H,3H)-dione  
 15            (Z)-N-(4-(2-(3- tert- butyl- 5-(2,4- dioxotétrahydropyrimidin- 1 (2H)-yl)- 2- méthoxyphényl)- 1-fluoro- vinyl) phényl) méthanesulfonamide ;  
               (E)-N-(4-(2-(3- tert- butyl- 5-(2,4- dioxotétrahydropyrimidin- 1 (2H)-yl)- 2- méthoxyphényl)- 1-fluoro- vinyl) phényl) méthanesulfonamide ;  
               (E)-N-(4-(3- tert- butyl- 5-(2,4- dioxotétrahydropyrimidin- 1 (2H)-yl)- 2- méthoxystyryl)- 2- fluoro- phényl) méthanesulfonamide ;  
 20            N-(4-(2-(3- tert- butyl- 5-(2,4- dioxotétrahydropyrimidin- 1 (2H)-yl)- 2- méthoxyphényl) cyclopropyl) phényl) méthanesulfonamide ;  
               N-(4-(3- tert- butyl- 5-(2,4- dioxotétrahydropyrimidin- 1 (2H)-yl)- 2- méthoxyphénéthyl) phényl) méthanesulfonamide ;  
 25            (E)-N-(4-(3-tert-butyl-5-(2,4-dioxotétrahydropyrimidin-1(2H)-yl)styryl)phényl)méthanesulfonamide ;  
               (Z)-N-(4-(2-(3-tert-butyl-5-(2,4-dioxotétrahydropyrimidin-1(2H)-yl)-2-méthoxyphényl)-1-méthoxyvinyl)phényl) méthanesulfonamide ;  
               (E)-1-(3-tert-butyl-4-méthoxy-5-styrylphényl)dihydropyrimidine-2,4(1H,3H)-dione ;  
               (E)-1-(3-tert-butyl-4-méthoxy-5-(4-méthoxystyryl)phényl)dihydropyrimidine-2,4(1H,3H)-dione ;  
 30            (E)-N-(4-(3- tert- butyl- 5-(5- fluoro- 2,4- dioxo- 3,4- dihydropyrimidin- 1 (2H)-yl)- 2- méthoxystyryl) phényl) méthanesulfonamide ;  
               (E)-N-(4-(3-bromo-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)phényl)méthanesulfonamide ;  
               (E)-N-(4-(5-(2,4- dioxo- 3,4- dihydropyrimidin- 1 (2H)-yl)- 2- méthoxy- 3-(thiophén- 2- yl) styryl) phényl) méthanesulfonamide ;  
 35            (E)-N-(4-(5-(2,4- dioxo- 3,4- dihydropyrimidin- 1 (2H)-yl)- 3-(furan- 2- yl)- 2- méthoxystyryl) phényl) méthanesulfonamide ;  
               (E)-N-(4-(5-(2,4- dioxo- 3,4- dihydropyrimidin- 1 (2H)-yl)- 2- méthoxy- 3-(pyridin- 4- yl) styryl) phényl) méthanesulfonamide ;  
               (E)-N-(4-(5-(2,4- dioxo- 3,4- dihydropyrimidin- 1 (2H)-yl)- 2- méthoxy- 3-(pyridin- 3- yl) styryl) phényl) méthanesulfonamide ;  
 40            (E)-N-(4-(5-(2,4- dioxo- 3,4- dihydropyrimidin- 1 (2H)-yl)- 2- méthoxy- 3-(thiophén- 3- yl) styryl) phényl) méthanesulfonamide ;  
               (E)-N-(4-(5-(2,4- dioxo- 3,4- dihydropyrimidin- 1 (2H)-yl)- 3-(furan- 3- yl)- 2- méthoxystyryl) phényl) méthanesulfonamide ;  
 45            (E)-N-(4-(5-(2,4- dioxo- 3,4- dihydropyrimidin- 1 (2H)-yl)- 3-(1- hydroxy- 2- méthylpropan- 2- yl)- 2- méthoxystyryl) phényl)méthanesulfonamide ;  
               (E)-N-(4-(5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3-iodo-2-méthoxystyryl)phényl)méthanesulfonamide ;  
               (E)-N-(4-(5-(2,4- dioxo- 3,4- dihydropyrimidin- 1 (2H)-yl)- 2- méthoxy- 3-(méthylsulfonyl) styryl) phényl) méthanesulfonamide ;  
 50            2-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-5-(méthylsulfonamido)benzoate de (E)-méthyle ;  
               acide (E)-2-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-5-(méthylsulfonamido) benzoïque ;  
               (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-3-(morpholine-4-carbonyl) phényl)méthanesulfonamide ;  
 55            (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-3-(hydroxyméthyl)phényl) méthanesulfonamide ;  
               (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-3-(méthoxyméthyl)phényl)

méthanesulfonamide ;  
 (E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-3-((isopentylamino)méthyl)phényl)méthanesulfonamide ;  
 N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-3-((E)-(méthoxyimino)méthyl)phényl)méthanesulfonamide ;  
 5 (E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-3-(oxazol-2-yl)phényl)méthanesulfonamide ;  
 (E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-3-(1H-imidazol-2-yl)phényl)méthanesulfonamide ;  
 10 2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-5-(méthylsulfonamido)phénylcarbamate de (E)-*tert*-butyle ;  
 (E)-N-(3-amino-4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)phényl)méthanesulfonamide ;  
 (E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-2-fluorophényl)méthanesulfonamide ;  
 15 (E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)-2-fluoro-5-méthylphényl)méthanesulfonamide ;  
 2-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxyphénéthyl)-5-(méthylsulfonamido)benzoate de méthyle  
 20 N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxyphénéthyl)phényl)méthanesulfonamide ;  
 (E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-éthoxystyryl)phényl)méthanesulfonamide ;  
 (E)-N-(4-(1-(3-*tert*-butyl-5-(2,4-dioxotétrahydropyrimidin-1(2H)-yl)-2-méthoxyphényl)prop-1-én-2-yl)phényl)méthanesulfonamide ;  
 25 (Z)-N-(4-(3-*tert*-butyl-5-(2,4-dioxotétrahydropyrimidin-1(2H)-yl)-2-méthoxystyryl)phényl)méthanesulfonamide ;  
 (E)-N-(4-(3-*tert*-butyl-5-(2,4-dioxotétrahydropyrimidin-1(2H)-yl)-2-méthoxystyryl)phényl)-N-(méthylsulfonyl)acétamide ;  
 30 (E)-1-(3-(4-aminostyryl)-5-*tert*-butyl-4-méthoxyphényl)dihydropyrimidine-2,4(1H,3H)-dione ;  
 (Z)-N-(4-(3-*tert*-butyl-5-(2,4-dioxotétrahydropyrimidin-1(2H)-yl)-2-méthoxystyryl)phényl)méthanesulfonamide ;  
 N-(4-(2-(3-*tert*-butyl-5-(2,4-dioxotétrahydropyrimidin-1(2H)-yl)-2-méthoxyphényl)-1-fluorovinyl)phényl)méthanesulfonamide ;  
 35 (E)-1-(3-*tert*-butyl-4-méthoxy-5-(4-nitrostyryl)phényl)dihydropyrimidine-2,4(1H,3H)-dione ;  
 1-(3-*tert*-butyl-5-[(2)-2-chloro-2-(4-nitro-phényl)-vinyl]-4-méthoxy-phényl)-dihydropyrimidine-2,4-dione ;  
 1-(3-*tert*-butyl-4-méthoxy-5-[(E)-2-(4-nitro-phényl)-propényl]-phényl)-dihydropyrimidine-2,4-dione ;  
 1-(3-*tert*-Butyl-5-[(E)-2-(4-nitro-phényl)-vinyl]-phényl)-dihydro-pyrimidine-2,4-dione ;  
 40 N-(4-{(E)-2-[3-*tert*-Butyl-5-(dioxo-tétrahydropyrimidin-1-yl)-2-méthoxy-phényl]-vinyl}-3-méthoxy-phényl)-méthanesulfonamide ;  
 N-(4-{(E)-2-[3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl}-3-formyl-phényl)-méthanesulfonamide ;  
 N-[4-{(E)-2-[3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl}-3-(hydroxyiminométhyl)-phényl]-méthanesulfonamide ;  
 45 2-[(E)-2-[3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl]-5-méthanesulfonylamino-N-(2-méthoxy-éthyl)-benzamide ;  
 ester éthylique d'acide 2-[(E)-2-[3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl]-5-méthanesulfonylamino-benzoïque ;  
 N-(4-{(E)-2-[3-*tert*-Butyl-2-chloro-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-phényl]-vinyl}-phényl)-méthanesulfonamide ;  
 50 2-[(E)-2-[3-*tert*-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl]-5-méthanesulfonylamino-N,N-diméthyl-benzamide ;  
 2-[(E)-2-[3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl]-5-méthanesulfonylamino-N-méthyl-benzamide ;  
 55 2-[(E)-2-[3-*tert*-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl]-N-(1,1-dioxotétrahydro-1λ<sup>6</sup>-thiophén-3-yl)-5-méthanesulfonylamino-N-méthyl-benzamide ;  
 N-(4-{(E)-2-[3-*tert*-butyl-5-(5-chloro-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl}-phényl)-méthanesulfonamide ;

2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl)-5-méthanesulfonylamino-benzamide ;

N-(3-(azétidine-1-carbonyl)-4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxyphényl]-vinyl)-phényl)-méthanesulfonamide ;

5 2-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl)-5-méthanesulfonylamino-N-(2-méthoxy-éthyl)-N-méthyl-benzamide ;

N-(4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl)-3-isopropoxy-méthyl-phényl)-méthanesulfonamide ;

10 N-[4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl)-3-(pyrrolidine-1-carbonyl)-phényl]-méthanesulfonamide ;

N-[4-((E)-2-[3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl)-3-(3-hydroxyazétidin-1-ylméthyl)-phényl]-méthanesulfonamide ;

N-(4-((E)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl)-3-pyrrolidin-1-ylméthyl-phényl)-méthanesulfonamide ;

15 N-(4-((2)-2-[3-tert-Butyl-5-(2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-yl)-2-méthoxy-phényl]-vinyl)-phényl)-méthanesulfonamide ; et

N-(4-(3-tert-butyl-5-(2,4-dioxotétrahydropyrimidin-1(2H)-yl)phénéthyl)phényl)méthane-sulfonamide.

20 **12.** Composé ou sel selon la revendication 1, dans lequel le composé est le (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)phényl)méthanesulfonamide.

**13.** Sel selon la revendication 1, dans lequel le sel est le sel sodique de (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-2-méthoxystyryl)phényl)méthanesulfonamide.

25 **14.** Sel selon la revendication 1, dans lequel le sel est le sel potassique de (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-pyrimidin-1(2H)-yl)-2-méthoxystyryl)phényl)méthanesulfonamide.

**15.** Sel selon la revendication 1, dans lequel le sel est le sel monopotassique de (E)-N-(4-(3-tert-butyl-5-(2,4-dioxo-3,4-dihydro-pyrimidin-1(2H)-yl)-2-méthoxystyryl)phényl)méthanesulfonamide.

30 **16.** Composition comprenant (a) un ou plusieurs composés et/ou sels selon l'une quelconque des revendications 1 à 15 ; (b) un ou plusieurs excipients ; et, en option, (c) un ou plusieurs agents thérapeutiques supplémentaires.

35 **17.** Un ou plusieurs composés et/ou sels selon l'une quelconque des revendications 1 à 15, et, en option, (c) un ou plusieurs agents thérapeutiques supplémentaires destinés à être utilisés dans le traitement de l'hépatite C chez un mammifère ayant besoin d'un tel traitement.

40

45

50

55

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21






Figure 22





## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- US 97288707 P [0001]
- US 61096794 A [0001]
- WO 2005021500 A1 [0549]

## Non-patent literature cited in the description

- **Hoover, J.** Remington's Pharmaceutical Sciences. Mack Publishing Co, 1975 [0379]
- Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems. Lippincott Williams & Wilkins, 2005 [0379]
- **Santana, L. et al.** *J. Heterocyclic Chem.*, 1999, vol. 36, 293-295 [0448] [0681]
- **Ueno, Y. et al.** *J. Org. Chem.*, 2005, vol. 70, 7925-7935 [0448]
- **Santana et al.** *J. Heterocyclic Chem.*, 1999, vol. 36, 293 [0481]
- **Taylor, WP.** *Bioorg. Med. Chem.*, 1996, vol. 4, 1515-1520 [0522] [0531]
- *Org. Prep. Proc. Int.*, 2004, vol. 36, 573-579 [0590]
- **Lal, GS et al.** *J. Org Chem.*, 1995, vol. 60, 7340-7342 [0607]
- **Morrison, J. F. ; S. R. Stone.** Approaches to the study and analysis of the inhibition of enzymes by slow- and tight-binding inhibitors. *Comments Mol. Cell. Biophys.*, 1985, vol. 2, 347-368 [0787]
- **Lohmann et al.** Replication of Subgenomic Hepatitis C Virus RNAs in a Hepatoma Cell Line. *SCIENCE*, 1999, vol. 285, 110-3 [0791]
- **Blight et al.** Efficient Replication of Hepatitis C Virus Genotype 1a RNAs in Cell Culture. *J. VIROL*, 2003, vol. 77, 3181-90 [0791]