

A&A Ref: 155742

PUBLICATION PARTICULARS AND ABSTRACT
(Section 32(3)(a) - Regulations 22(1)(g) and 31)

21	01	PATENT APPLICATION NO	22	LODGING DATE	43	ACCEPTANCE DATE
----	----	-----------------------	----	--------------	----	-----------------

07465

6 September 2006

2006-09-06

51	INTERNATIONAL CLASSIFICATION	NOT FOR PUBLICATION
----	------------------------------	---------------------

C07C

CLASSIFIED BY: ISA

71	FULL NAME(S) OF APPLICANT(S)
----	------------------------------

Syngenta Participations AG

72	FULL NAME(S) OF INVENTOR(S)
----	-----------------------------

BENKE, Alan Henry

WICHERT, Julie Marie

EARLIEST PRIORITY CLAIMED	COUNTRY	NUMBER	DATE
	33 GB	31 0406894.6	32 26 March 2004

NOTE: The country must be indicated by its International Abbreviation - see schedule 4 of the Regulations

54	TITLE OF INVENTION
----	--------------------

Process for purifying mesotrione

57	ABSTRACT (NOT MORE THAN 150 WORDS)
----	------------------------------------

NUMBER OF SHEETS	
------------------	--

The sheet(s) containing the abstract is/are attached.

If no classification is furnished, Form P.9 should accompany this form.

~~The figure of the drawing to which the abstract refers is attached.~~

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
6 October 2005 (06.10.2005)

PCT

(10) International Publication Number
WO 2005/092846 A1

(51) International Patent Classification⁷: C07C 315/06, 317/24

(21) International Application Number:
PCT/EP2005/002230

(22) International Filing Date: 3 March 2005 (03.03.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0406894.6 26 March 2004 (26.03.2004) GB

(71) Applicant (for all designated States except US): SYNGENTA PARTICIPATIONS AG [CH/CH]; Intellectual Property Department, Schwarzwaldallee 215, CH-4058 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BENKE, Alan, Henry [US/US]; Cold Creek Plant, Bucks, AL (US). WICHERT, Julie, Marie [US/US]; Cold Creek Plant, Bucks, AL (US).

(74) Agent: SWIFT, Jane, Elizabeth; Intellectual Property Department, Syngenta Limited, P.O. Box 3538, Jealott's Hill International Research Centre, Bracknell, Berks RG42 6YA (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SI, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR PURIFYING MESOTRIONE


(57) Abstract: A method for reducing the cyanide levels in a mesotronic sample, said method comprising: (i) taking an aqueous solution of the mesotronic sample in an aqueous solvent, (ii) adjusting the pH of said aqueous solution to a value of 9.5 or higher, and (iii) crystallising the mesotrone out of solution is disclosed.

WO 2005/092846 A1


PROCESS FOR PURIFYING MESOTRIONE

The present invention relates to a novel method for reducing the cyanide content of a mesotrione sample.

Mesotrione (2-(2'-nitro-4'-methylsulphonyl benzoyl)-1,3-cyclohexanedione) is a selective corn herbicide and has the structure of formula (I)



Mesotrione is prepared by reacting 2-nitro-4-methylsulphonyl benzoyl chloride with cyclohexanedione to give the enol ester, followed by a rearrangement reaction to give mesotrione, as shown in the following reaction scheme:



2-Nitro-4-methylsulphonyl benzoyl chloride is prepared from the corresponding benzoic acid, which in turn is prepared by oxidation of 2-nitro-4-methylsulphonyl toluene. More details on the preparative route may be found in US4695673.

During the rearrangement process, the mesotrione sample is contaminated with cyanide residues from the acetone cyanohydrin catalyst. It is therefore an object of the present invention to provide a simple, but effective method for reducing the level of cyanide residues in the mesotrione sample to an acceptable level.

It has surprisingly been found that adjusting the pH of a mesotrione sample in aqueous solution has a significant impact on the resulting cyanide levels.

Accordingly the present invention provides a method for reducing the cyanide levels in a mesotrione sample, said method comprising:

- (i) taking an aqueous solution of the mesotrione sample in an aqueous solvent,
- 5 (ii) adjusting the pH of said aqueous solution to a value of 9.5 or higher, and
- (iii) crystallising the mesotrione out of solution.

In one embodiment of the invention, the mesotrione sample has previously been isolated, and the aqueous solution is formed by dissolving the isolated sample in an aqueous solvent.

- 10 In a second embodiment of the invention, the mesotrione sample has not previously been isolated and remains dissolved in the aqueous solvent used in the condensation/rearrangement reaction described above.

The aqueous solvent may be selected from the group consisting of water and a water soluble solvent, such as acetonitrile, triethylamine, methanol, ethanol, acetone.

- 15 Preferably, the aqueous solvent is water. The mesotrione sample is suitably dissolved in the aqueous solvent to give a solution concentration of from 1% to 30%, suitably from 5% to 15%, and preferably from 8% to 11%.

- Suitably, the pH of the aqueous solution is raised to a pH of at least 11, and preferably at least 11.5. Suitably, the aqueous mesotrione sample is held at a pH of at least 9.5 for at least 5 minutes, suitably at least 15 minutes and preferably at least 30 minutes.
- 20

Suitably, the temperature of the aqueous solution should not be greater than 30°C.

- The crystallisation is carried out according to standard laboratory procedures. For example, for a batch crystallisation, the final pH is adjusted from its starting value of 9.5 or higher to pH 2.5 by charging hydrochloric acid to the crystalliser. The hydrochloric acid should be charged in a manner to ensure adequate mixing. The crystallisation process may also be carried out as a semi-batch or continuous process. The crystallisation step may also include a nitrogen-sparging step, wherein nitrogen is bubbled through the crystallisation vessel in a continuous fashion and sent to a scrubber.
- 25

- 30 The method of the invention may further include a distillation step to remove solvents when the mesotrione sample has not previously been isolated (i.e. the second embodiment of the invention). The distillation step may be carried out either before or after adjusting the pH to 9.5 or higher. The distillation step is suitably carried out using a sufficient amount of steam to remove the solvents from the aqueous solution.

Suitably, the method of the invention reduces the cyanide levels in the mesotrione sample to 150ppm or less, more suitably 100ppm or less and preferably 50ppm or less.

The invention will now be described further with reference to the following examples, which are illustrative but not limiting of the invention.

5

Example 1

Wet paste mesotrione that was high in total CN was subjected to different treatments in an effort to reduce the total CN content. The results are shown in Table 1.

Table 1				
Example No.	Treatment	Original Cyanide Content (PPM)	Cyanide Content after Treatment (PPM)	% reduction in Cyanide content
1A	Mesotrione was placed in water to make an aqueous solution at a concentration of ~10%, pH was adjusted to >13, ACN was charged, and the mixture was then batch crystallised following standard lab procedures.	546	15	97%
1B	Mesotrione was placed in water to make an aqueous solution at a concentration of ~10%, pH was adjusted to 11.3, ACN was charged, and the mixture was batch crystallised following standard lab procedures.	1114	557	50%
1C	Mesotrione was placed in water to make an aqueous solution at a concentration of ~10%, pH was adjusted to >13, ACN was charged, and the mixture was batch crystallised following standard lab procedures.	1114	50	96%
1D	Mesotrione was placed in water to make an aqueous solution at a concentration of ~10%, pH was adjusted to 11.3, ACN was charged, and the mixture was batch crystallised following standard lab procedures.	690	150	78%
1E	Mesotrione was placed in water to make an aqueous solution at a concentration of ~10%, pH was adjusted to 12-13, ACN was charged, and the mixture was batch crystallised following standard lab procedures.	690	170	75%

Example 2

Mesotrione was crystallised from samples taken from the plant during the solvent distillation. Samples were taken from the same batch after both 4500 lbs steam (distillation not finished) and 5000 lbs steam (distillation finished) had been used during the distillation process. The pH of the sample was adjusted and the samples were crystallised via standard lab procedures. Total CN content was measured by titration of the wet paste. The results are given in Table 2.

Table 2			
Example No.	Distillation Complete?	Starting pH of crystallisation	Total CN of wet paste (PPM)
2A	No	9.5	278
2B	Yes	9.5	651
2C	No	11.1	120
2D	Yes	11.1	26
2E	No	12.8	121
2F	Yes	12.8	20

10

Example 3

Mesotrione was crystallised from samples taken from the plant during the solvent distillation. The effects of varying the starting pH of the crystallisation and purging the headspace of the crystallisation vessel with nitrogen were looked at. Samples were taken from the same batch after both 4500 lbs steam (distillation not finished) and 5045 lbs steam (distillation finished) has been used during the distillation. The pH of the sample was adjusted and the samples were crystallised via standard lab procedures. Total CN was measured by titration of the wet paste or filtrate. The results are shown in table 3.

15

Table 3					
Example No.	Distillation Complete	Starting pH of crystallisation	N₂ purge?	Total CN content of wet paste (PPM)	% reduction in cyanide content from control
3A	No	5.2	No	311	Control
3B	Yes	5.2	No	236	Control
3C	No	9.5	No	91	70
3D	Yes	9.5	No	63	74
3E	No	11.0	No	20	94
3F	Yes	11.0	No	50	79
3G	No	13	No	15	95
3H	Yes	13	No	56	76
3I	No	5.2	Yes	294	Control
3J	No	11.0	Yes	46	85
3K	No	11.3	Yes	15	95

Example 4

- 5 This example looks at the cyanide content of mesotrione crystallised from different feed pH in a continuous crystallisation. The results are shown in Table 4.

Table 4			
Sample	pH of feed to crystalliser	Total CN (ppm)	% Reduction in Cyanide Content from Control
7 th (final) sample from crystalliser	5.0	217	Control
7 th (final) sample from crystalliser	5.0	181	Control
7 th (final) sample from crystalliser	11	15	92-93
Final sample from crystalliser	11	15	92-93
Final sample from crystalliser	11	15	92-93
Final sample from crystalliser	9.5	15	92-93

Example 5

- 10 Mesotrione was produced from the acid chloride by a standard condensation/rearrangement reaction. After the condensation/rearrangement reaction, water was added and the pH was adjusted to >11 and held for ½ hour. The pH was then adjusted to ~5, the mixture distilled and then batch crystallised from either pH 5 or 9.5. The results are shown in Table 5.

Table 5			
Example No.	pH held at after condensation/rearrangement reaction	Starting pH of crystallisation	Total CN in mesotrione (ppm)
5A	11.9	5.0	15
5B	11.3	9.5	40

Example 6

- 5 A large sample of mesotrione was obtained at the end of the distillation. This sample was divided into aliquots which were adjusted to a pH >11. A series of batch samples were made up and held agitated at the given pH for the amount of time specified in the table before being quickly adjusted to pH 2.4, filtered, washed and submitted for total cyanide analysis. The results are shown in Table 6.

Table 6			
Example No.	Starting pH	Time kept at starting pH (min)	Total CN of mesotrione (ppm)
6A	11.6	0	111
6B	11.3	5	76
6C	11.5	10	72
6D	11.6	15	73
6E	11.5	20	55
6F	11.5	25	72
6G	11.4	30	76
6H	11.7	60	15
6I	12.3	90	15

Claims

1. A method for reducing the cyanide levels in a mesotrione sample, said method comprising:
 - 5 (i) taking an aqueous solution of the mesotrione sample in an aqueous solvent,
 - (ii) adjusting the pH of said aqueous solution to a value of 9.5 or higher, and
 - (iii) crystallising the mesotrione out of solution.
- 10 2. The method of claim 1, wherein the mesotrione sample has previously been isolated and is redissolved in an aqueous solvent.
3. The method of claim 1, wherein the mesotrione sample has not previously been isolated and is already present as an aqueous solution in an aqueous solvent.
- 15 4. The method of claim 3, which further includes a distillation step.
5. The method of claim 4, wherein the distillation step is carried out prior to adjusting the pH of the aqueous solution to a value of 9.5 or higher.
- 20 6. The method of claim 4, wherein the distillation step is carried out after adjusting the pH of the aqueous solution to a value of 9.5 or higher.
7. The method according to any one of claim 1 to 6, wherein the crystallisation step also includes a nitrogen sparging step.
- 25 8. A method according to any one of claims 1 to 7, wherein the cyanide levels in the mesotrione sample are reduced to 150ppm or less.