



(86) Date de dépôt PCT/PCT Filing Date: 2003/07/02  
(87) Date publication PCT/PCT Publication Date: 2004/01/15  
(85) Entrée phase nationale/National Entry: 2004/12/15  
(86) N° demande PCT/PCT Application No.: US 2003/020779  
(87) N° publication PCT/PCT Publication No.: 2004/004689  
(30) Priorité/Priority: 2002/07/02 (10/188,554) US

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> A61K 9/06, A61K 47/18, A61K 9/00

(71) Demandeur/Applicant:  
NEXMED (HOLDINGS), INC., US

(72) Inventeurs/Inventors:  
YEAGER, JAMES L., US;  
BUEYUEKTIMKIN, NADIR, US;  
BUEYUEKTIMKIN, SERVET, US;  
LU, MINGQI, US;  
MO, Y. JOSEPH, US

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : COMPOSITIONS ET METHODES DE CORRECTION DE DYSFONCTIONNEMENTS SEXUELS CHEZ LA FEMME

(54) Title: COMPOSITIONS AND METHODS FOR AMELIORATION OF HUMAN FEMALE SEXUAL DYSFUNCTION

(57) **Abrégé/Abstract:**

The invention provides compositions and methods suitable for ameliorating female sexual dysfunction, and in particular, female sexual arousal disorder. In one embodiment, the invention provides a semisolid composition suitable for topical application comprising: an effective amount of a vasoactive prostaglandin, a penetration enhancer, a polymer thickener, a lipophilic component, and an acidic buffer system.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number  
**WO 2004/004689 A1**

- (51) International Patent Classification<sup>7</sup>: **A61K 9/06**, 47/18, 9/00
- (21) International Application Number: PCT/US2003/020779
- (22) International Filing Date: 2 July 2003 (02.07.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 10/188,554 2 July 2002 (02.07.2002) US
- (71) Applicant: **NEXMED (HOLDINGS), INC.** [US/US]; 350 Corporate Drive, Robbinsville, NJ 08691 (US).
- (72) Inventors: **YEAGER, James, L.**; 476 Oakwood Avenue, Lake Forest, IL 60045 (US). **BÜYÜKTIMKIN, Nadir**; 299 Meadowbrook Road, Robbinsville, NJ 08691 (US). **BÜYÜKTIMKIN, Servet**; 299 Meadowbrook Road, Robbinsville, NJ 08691 (US). **LU, Mingqi**; 12 Fountayne Lane, Lawrenceville, NJ 08648 (US). **MO, Y., Joseph**; 1 Bellevue Terrace, Princeton, NJ 08540 (US).
- (74) Agents: **ZIMMERMAN, Roger, P.** et al.; Bowditch & Dewey, LLP, 161 Worcester Road, P.O. Box 9320, Framingham, MA 01701-9320 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT (utility model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (utility model), DE, DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK (utility model), SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— with international search report
- (48) Date of publication of this corrected version: 15 April 2004
- (15) Information about Correction: see PCT Gazette No. 16/2004 of 15 April 2004, Section II
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS AND METHODS FOR AMELIORATION OF HUMAN FEMALE SEXUAL DYSFUNCTION

(57) Abstract: The invention provides compositions and methods suitable for ameliorating female sexual dysfunction, and in particular, female sexual arousal disorder. In one embodiment, the invention provides a semisolid composition suitable for topical application comprising: an effective amount of a vasoactive prostaglandin, a penetration enhancer, a polymer thickener, a lipophilic component, and an acidic buffer system.

WO 2004/004689 A1

## COMPOSITIONS AND METHODS FOR AMELIORATION OF HUMAN FEMALE SEXUAL DYSFUNCTION

### BACKGROUND OF THE INVENTION

Sexual dysfunction has been a persistent problem, more frequent in an aging  
5 population, that has only recently been addressed with frank evaluation, scientific  
investigation and effective treatment. Male impotence, especially male erectile  
dysfunction, has received the most attention. Female sexual dysfunction has been  
considered in the context of male erectile dysfunction, in part because of the  
anatomical and physiological parallels between the male and female genitalia, and in  
10 part, with the hope that effective treatments for male erectile dysfunction could  
provide some relief for female sexual dysfunction.

Both male and female sexual behavior is viewed from the standpoint of a  
four-phase sexual response cycle consisting of the stages of desire, excitement,  
orgasm and resolution. Studies have shown that while there are many similarities  
15 between male and female sexual response, significant differences exist. Specific  
dysfunctions have been correlated with the phases of the model. The female sexual  
response and its dysfunctions remain poorly understood.

Female sexual arousal disorder (FSAD) is the persistent or recurrent inability  
to attain, or to maintain, sufficient sexual excitement, which causes personal distress.  
20 It may be expressed as lack of subjective excitement, lack of genital response, such as  
lubrication and swelling, or lack of other somatic responses. Female sexual arousal  
disorder is one form of female sexual dysfunction, and is associated with the  
excitement phase. See Basson, R., et al., Report of the international consensus  
development conference on female sexual dysfunction: definitions and  
25 classifications, *J. Urol.* 2000;163(3):888-93.

While increased understanding of the pathophysiology of male erectile  
dysfunction has progressed rapidly in the past decade and led to new therapeutic  
modalities, little has been done to address similar issues in women. Cardiovascular  
risk factors have been shown to correlate with complaints of vaginal and clitoral  
30 dysfunction. Goldstein, M. K., et al., Gynecological factors in sexual dysfunction of  
the older woman. *Clin Geriatr Med* 7: 41-61, (1991); Sadeghi-Nejad, H., et al.:

Impotence is a couple's disease: studies in female sexual dysfunction. J Urol 155: 677A, (1996); Slob, A. K., et al.: Sexuality and psychophysiological functioning in women with diabetes mellitus. J Sex Marital Ther: 59-69, (1990).

The correlation of cardiovascular risk factors and complaints of vaginal and clitoral dysfunction have led to suggestions that a significant degree of female sexual dysfunction is due to vascular insufficiency and therefore amenable to treatment with vasoactive agents. The underlying foundations of the normal and dysfunctional female sexual response must be considered in the context of the anatomy and physiology, summarized below. See, generally, Goldstein, I., and Berman, J.R., Vasculogenic female sexual dysfunction: vaginal engorgement and clitoral erectile insufficiency syndromes, Int. J. Impotence Research 10: Suppl. 2, S84-S90 (1998).

#### Anatomy of the Vagina

The vagina is the canal that connects the uterus with the external genital organs. Its design easily accommodates penetration of a rigid penile erection. At the posterior end the rounded neck of the uterus, the cervix, projects into the space known as the fornix or vaginal vault. Anteriorly, two pleats of sensitive tissue, the labia minora, surround the opening of the vagina and are further protected by larger folds known as the labia majora.

The walls of the vagina consist of three layers - an inner mucosa, an aglandular mucous membrane epithelium, an intermediate, highly vascularized muscularis layer, and an outer supportive fibrous mesh. The vaginal mucosa is a mucous type stratified squamous cell epithelium that undergoes hormone-related cyclical changes, such as a slight keratinization of the superficial cells during the menstrual cycle. The muscularis portion comprises smooth muscle and an extensive arborization of blood vessels that may swell during intercourse. The surrounding fibrous layer provides structural support to the vagina; this layer consists of elastin and collagen fibers that allow for expansion of the vaginal vault during sexual arousal or childbirth. Large blood vessels run within the mucosa, and nerve plexuses are present within muscular and adventitial layers. The vagina has many rugae or folds that are necessary for the distensibility of the organ during intercourse and childbirth. Smaller ridges lend to the frictional tension that exists during intercourse.

The arterial supply to the vagina is derived from an extensive network of

branching vessels surrounding it from all sides. The anterior branch of the internal iliac artery continually bifurcates as it descends through the pelvis with a series of the newly generated vessels, each supplying the vagina to some degree. After giving off an obturator artery branch, the umbilical, and the middle rectal arteries diverge off to supply a superior and inferior vesical artery, respectively. Between the umbilical and the mid-rectal branches there is a generation of a uterine artery, which further bifurcates to give the vaginal artery. The internal pudendal and accessory pudendal artery also send a branch to the vaginal artery. Finally, the common clitoral artery sends a branch to the vaginal muscularis.

10 The neurologic innervation of the vagina originates from two separate plexuses, the superior hypogastric plexus and the sacral plexus. The hypogastric nerve plexus descends on the great vessels spreading into an inferior hypogastric plexus, which systematically branches further into a uterovaginal nerve. The somatic pudendal nerve originates off the pelvic splanchnic branches from the secret plexus. 15 Pudendal branching innervates the vagina towards the opening of the introitus as the perineal and posterior labial nerves.

Immunohistochemistry studies have been utilized to better understand the innervation of the human vaginal mucosa. In a study by Hilliges *et al.* using protein gene product 9.5, more distal areas of the vagina had significantly more nerve fibers compared to the more proximal parts, and the anterior wall showed a denser innervation than the posterior wall (Hilliges, M. et al., Innervation of the human vaginal mucosa as revealed by PGP 9.5 immunohistochemistry, Acta Anatomica 153: 119 (1995)). Graf *et al* studied the distribution patterns and the occurrence of helospectin and pituitary adenylate cyclase activating polypeptide (PACAP) immunoreactivity (Graf, A.H., et al. Helospectin and pituitary adenylate cyclase activating polypeptide in the human vagina, Regul. Pept. 55: 277 (1995)). They confirmed a dense network of vasoactive intestinal peptide (VIP) immunoreactive nerve fibers showing sub-populations of helospectin and LI-type PACAP. Nerve fibers of the vagina had previously been shown to be active in association with specific peptides that include VIP, peptide histidine methionine (PHM), calcitonin gene related peptide (CGPP), and galanin. Genital vasodilation and subsequent increase in vaginal blood flow and lubrication have been observed upon exposure of vessels to VIP. VIP has been implicated as the neurotransmitter for mediating

vaginal vasodilation and the formation of lubricating fluid during sexual arousal. Helospectin and PACAP, a potent vasodilator, belong to the same peptide family as VIP and PHM, and recent observations have been made to the effect that distributions and co-localizations of helospectin and VEP as well as PACAP and VIP have been reported in the mammalian gastrointestinal tract.

The vaginal canal is lubricated primarily from a transudate originating from the subepithelial vascular bed passively transported through the interepithelial spaces, sometimes referred to as intercellular channels. Additional moistening during intercourse comes from secretion of the paired greater vestibular or Bartholin's glands.

Estrogen effects on the maintenance and function of female genitalia have been well documented in studies. Estrogen receptors have been shown to exist throughout the vaginal epithelium, in stromal cells, and in the smooth muscle fibers in the muscularis. Weaker conformations of estrogen such as estriol appear more effective in stimulating the vagina as opposed to the uterus. Thickness and rugae of the vaginal wall, as well as vaginal lubrication, have been shown to be estrogen dependent. Although this fluid production has been shown to be hormone-dependent both in the resting state and during sexual excitement, quantitative changes apparently do not occur during the menstrual cycle. An insufficient amount of estrogen will result in thin vaginal walls more easily susceptible to trauma with a decreased ability to heal, as well as a drier and less acidic vaginal environment more vulnerable to infection. Vaginal dryness is associated with ovarian failure and is effectively controlled by estrogen replacement therapy. Some women who are not sexually active may not notice the extent of vaginal atrophy but when coitus does resume, pain and discomfort from intercourse can be considerable.

#### Anatomy of the Clitoris

The clitoris is the homologue of the penis arising from the embryological genital tubercle. The clitoris consists of a cylindrical, erectile organ composed of three parts: the outermost glans or head, the middle corpus or body, and the innermost crura. The glans of the clitoris is visualized as it emerges from the labia minora, which bifurcate to form the upper prepuce anteriorly and the lower fronulum posteriorly. The body of the clitoris consists of two paired corpora cavernosa of

about 2.5 cm in length and lacks a corpus spongiosum. The body extends under the skin at the corona to the crura. The two crura of the clitoris, formed from the separation of the most proximal portions of the corpora in the perineum, attach bilaterally to the undersurface of the symphysis pubis at the ischiopubic rami.

5 A fibrous tunica albuginea ensheathes each corporal body made up of lacunar space sinusoids surrounded by trabecula of vascular smooth muscle and collagen connective tissue. No retractor clitoridis muscle exists in humans as it does in other animals such as cattle and sheep, however a supporting suspensory ligament does hold the clitoris in the introital region.

10 The main arterial supply to the clitoris is from the ilio-hypogastric-pudendal arterial bed. The internal pudendal artery is the last anterior branch off the internal iliac artery. Distally, the internal pudendal artery traverses Alcock's canal, a position of the obturator fascia and lies on the inner side in apposition to the ischio-pubic ramus. In this latter location, the artery is susceptible to blunt perineal trauma. The  
15 internal pudendal artery terminates as it supplies the inferior rectal and perineal artery, which supplies the labia. The common clitoral artery continues to the clitoris. This artery bifurcates into a dorsal clitoral artery and a cavernosal clitoral artery.

Autonomic efferent innervation of the clitoris passes from the pelvic and hypogastric nerves to the clitoris through the urogenital diaphragm. Pelvic nerve  
20 stimulation results in clitoral smooth muscle relaxation and arterial smooth muscle dilation. There is a rise in clitoral cavernosal artery inflow, an increase in clitoral intracavernous pressure which lead to tumescence and extrusion of the glans clitoris.

Anatomical studies using female rats have indicated that the major neuronal input to the clitoris was seen in spinal segments from L5-S1, and to a lesser extent in  
25 T12-L4 as well as S2-S4. When a label that is taken up by nerve terminals and transported retrogradely to the nerve cell bodies (pseudorabies virus) was injected into the clitoris, labeled nerve cell bodies were found in the brain in multiple locations, including the nucleus paragigantocellularis, raphe pallidus, raphe magnus, Barrington's nucleus, ventrolateral central gray, hypothalamus, and the medial pre-  
30 optic region. This implies a multisynaptic circuit of neurons may be involved in clitoral neurological control rather than just a simple somatic reflex connection.

Morphological studies have been performed using wheat germ agglutinin conjugated with horseradish peroxidase (WGA / HRP) injected into the clitoris of the

female cat to compare afferent pathways to the entire population of pudendal nerve afferents. Central projections of the clitoral afferents were identified in the L7-S3 segments with the most prominent labeling in S1-S2. In the same study, electrophysiological analysis of the clitoris performed under constant mechanical pressure stimulation indicated both phasic and tonic discharges in L7-S2, but most prominently in S1. In contrast electrical stimulation of the clitoris evoked discharges at S1 only. The neurotransmitters mediating clitoral and arterial smooth muscle dilation remain undetermined, however preliminary studies suggest that nitric oxide is involved. Histochemical studies have revealed VIP and neuropeptide Y (NPY) immunoreactive nerves in the clitoral erectile tissues. Somatic sensory pathways originate from the clitoral skin. There exists a dense collection of Pacinian corpuscles innervated by rapidly adapting myelinated afferents, as well as Meissner's corpuscles, Merckel tactile disks, and free nerve endings. These sensory afferents pass from the dorsal clitoral nerve to the pudendal nerve.

15

#### The Grafenberg Spot

The Grafenberg spot (or G-spot) can also play a role in female sexual arousal. The current information regarding the Grafenberg zone (also known as Grafenberg spot, or G-spot) was recently summarized (Goldstein, I., et al., "Female Sexual Dysfunction" pp. 507-557, at 523 in Jardin, A, et al., editors, *Erectile Dysfunction*, (First International Consultation on Erectile Dysfunction, co-sponsored by the World Health Organization (WHO), International Consultation on Urological Diseases (ICUD) and Societe Internationale d'Urologie (SIU), held July 1-3, 1999, Paris. 2000). Grafenberg reported that the digital stroking of the anterior vagina along the urethra, especially in the region of the base of the bladder, sexually aroused female subjects greatly (Grafenberg E. (1950): The role of the urethra in the female orgasm. *Int. J. Sexology*. 3: 145-148). In a number of women this area swelled up to the size of a kidney bean and projected into the vaginal lumen. Few took any notice of this finding. The area was rediscovered and renamed the G-spot in honor of Grafenberg (Ladad, A.K., et al., (1982): *The G spot and other recent discoveries about Human Sexuality*. Holt, Rinehart & Winston, New York). Other investigators could not locate a "spot" but found, rather than a punctate locus, a general excitable area along the whole length of the urethra running along the anterior vaginal wall (Hoch Z.

30



(1986): Vaginal erotic sensitivity by sexological examination. *Acta Obstet. et Gynecol. Scand.* 65: 768-773). When this was stimulated manually, the sexual arousal induced was almost immediate. Alzate & Londono located the erotic sensitive area in closer relation to the bladder base than the urethra (Alzate H. & Londono M.L. 5 (1984): Vaginal erotic sensitivity. *J. Sex & Marital Therapy.* 10: 49-56). Lenck, et al. localized by ultrasound in the living subjects the underlying structure in the anterior vaginal wall that gave the erotic sensations on stimulation as the urethral sphincter confirming it by dissection in the cadaver (Lenck L. Ch., et al., (1992): Sphincter uretral (point G) correlations anatomo-cliniques. *Revue Française de Gynecologie et* 10 *Obstrique.* 87: 65-69.). Other investigators have implied that the G spot/area represents that part of the urethra that contains the periglandular or paraurethral tissue, corresponding to the female equivalent of the prostate (See Zaviacic M. & Whipple B. (1993): Update on the female prostate and the phenomenon of female ejaculation. *J. Sex Research.* 30: 148-151, for references). These glands are present to 15 a greater or lesser degree in about 90% of women.

#### Physiology of Female Sexual Arousal

The female sexual response phase of arousal is not easily distinguished from the phase of desire until physiological changes begin to take place in the vagina and clitoris as well as other sexual organs. Sexual excitement and pleasure are 20 accompanied by pelvic vasocongestion and swelling of the external genitalia including vaginal engorgement and clitoral erection.

Vaginal engorgement enables a process of plasma transudation to occur, allowing a flow through the epithelium and onto the vaginal surface. Plasma transudation results from the rising pressure in the vaginal capillary bed during the 25 arousal state. In addition there is an increase in vaginal length and luminal diameter, especially in the distal 2/3 of the vaginal canal.

#### Dissociation Of Genital Reflexes From Subjective Arousal

Central nervous system areas primarily implicated in sexual arousal, based on animal research, include the medial preoptic, anterior hypothalamic region and related 30 limbic-hippocampal structures. Cognitive effects have been investigated, and in one study the results suggest that the greatest contribution to sexual arousal in the female

results from cognitive processing of stimulus content and meaning, and not from peripheral vasocongestive feedback (Laan, E., et al., Determinants of subjective experience of sexual arousal in women. Feedback from genital arousal and erotic stimulus content, Psychophysiol. 32: 44- (1995)).

5           The distinction between local physiological aspects of sexual response, such as genital vasocongestion measured by vaginal photoplethysmography, and subjective sexual arousal, measured by self-reporting rating scales and inventories has been clearly demonstrated in both normal and sexually dysfunctional women (Palace, E.M. and Goralka, B.B., Differential patterns of arousal in sexually functional and  
10           dysfunctional women: Physiological and subjective components of sexual response, Arch. Sexual Behav. 21: 135-159 (1992)). Several reliable and validated self-report inventories are recognized for measurement of female sexual function (Derogatis, L.R. and Conklin-Powers, B., Psychological assessment measures of female sexual functioning in clinical trials, Int. J. Impot. Res. 10 Suppl. 2: S111-S116 (1998)).

15           There does not appear to be a relation between menstrual phases and physiologic arousability. Meuwissen and Over (Habituation and Dishabituation of Female Sexual Arousal, Behav. Res. Ther. 28: 217- (1990)) have found that neither film-induced nor fantasy-induced levels of sexual arousal varied significantly throughout the menstrual cycle. There are conflicting reports as well as to the  
20           habituation of the female sexual response. Some claim that levels of subjective and physiologic sexual arousal decrease over repeated exposure to sexual stimuli. Others could not elucidate similar results even after 21 trials, yet both concur that the subsequent presentation of a novel stimulus will increase the female sexual response. The desire for increased sexual performance on sexual arousal in functional women  
25           have been found to facilitate genital responses, most prominently with the stimulus of erotic fantasy as opposed to erotic film. Interestingly, masturbation frequency had no affect on genital responses despite its significance on subjective reports of arousal. (Laan et al, 1995; Meuwissen and Over, 1990).

30           Clinicians and researchers have assumed that sexual arousal, is inhibited by the sympathetic nervous system, while facilitation and maintenance are through the parasympathetic nervous system. However, studies have challenged these notions in the woman. Intense exercise, consisting of twenty-minute bike riding sessions, increased physiological sexual arousal measured by vaginal photoplethysmography.

This challenged the notion that sympathetic nervous system stimulation inhibited sexual arousal in women and further provided evidence that sexual arousal was actually facilitated by the sympathetic nervous system. Another study examined the temporal effect of sympathetic activation through acute exercise on immediate  
5 delayed, and residual sexual arousal. Sexual arousal was objectively assessed by vaginal plethysmography. A relationship between sympathetic nervous system activation and sexual arousal was found, such that sexual arousability was inhibited five minutes post-exercise and was facilitated fifteen minutes post-exercise and only marginally increased thirty minutes post-exercise. The two studies suggest that  
10 sympathetic nerve stimulation activation plays an important facilitatory role in the early stages of sexual arousal.

The clitoris may play a major role during sexual activity in that it is not only part of what makes the sexual act enjoyable for the woman but also enhances her response to coitus upon clitoral stimulation. Clitoral stimulation may induce local  
15 autonomic and somatic reflexes causing vaginal vasocongestion, engorgement, and subsequent transudation, lubricating the introital canal making the sexual act easier, more comfortable, and more pleasurable. The more stimulation, the higher the level of arousal and the easier it is to further increase stimulations.

## 20 Vasculogenic Female Sexual Dysfunction

Female sexual dysfunction has traditionally included disorders of desire/libido, disorders of arousal, pelvic pain disorders, and inhibited orgasm. Patient surveys estimate that 18-76% of adult women have such complaints during sexual activity. Female sexual dysfunction which may have its origin in abnormal  
25 arterial circulation into the vagina or clitoris during sexual stimulation, usually from atherosclerotic vascular disease may be considered a disorder of arousal. This vasculogenic female sexual dysfunction may include such clinical symptoms as delayed vaginal engorgement, diminished vaginal lubrication, pain or discomfort with intercourse, diminished vaginal sensation, diminished vaginal orgasm, diminished  
30 clitoral sensation or diminished clitoral orgasm. Traumatic injury to the ilio-hypogastric-pudendal arterial bed from pelvic fractures or blunt perineal trauma may also result in diminished vaginal/clitoral blood flow following sexual stimulation and fall into this vasculogenic category.

## Prostaglandins

The prostaglandins are a series of cyclic derivatives of certain unsaturated fatty acids. They are found in a variety of tissues, including the prostate gland, the seminal vesicles, the lungs and the brain. These naturally occurring prostaglandins are derived by cyclization of 20-carbon unsaturated fatty acids such as arachidonic acid. See Lehninger, Albert L., *Biochemistry*, 2d ed. (1975), p. 300 (hereinafter "Lehninger").

Prostaglandins as a class of compounds have diverse pharmacologic activity, including stimulation of gastrointestinal and reproductive smooth muscle, relaxation and contraction of respiratory smooth muscle, hypotensive activity, inhibition of fatty acid lipolysis, inhibition of blood platelet aggregation, and inhibition of gastric acid secretion. Therapeutic utility of prostaglandins in general is correspondingly broad. As for prostaglandin E<sub>1</sub> ("PGE<sub>1</sub>") in particular, this compound, salts thereof, and lower alkyl esters thereof are well known and disclosed, e.g., in U.S. Pat. Nos. 3,069,322 (Bergstrom et al.), 5,219,885 (Froelich et al.) and in *J. Org. Chem.* 1974, 37, 2921. PGE<sub>1</sub> has found utility in the treatment of peripheral occlusive diseases, acute myocardial infarction, angina pectoris, acute ischemic stroke, asthma, gastrointestinal ulcers, ulcers of the skin, and organ rejection. Various routes of administration have been described, including oral, intravenous, buccal, rectal, intra-arterial, subcutaneous, and sublingual. The preferred route of administration of PGE<sub>1</sub> will of course be dependent on the particular intended therapeutic use.

Prostaglandins are well known to those skilled in the art. This class of drugs includes those derivatives of prostanic acid (5-octylcyclopentaneheptanoic acid) referred to as A-I series prostaglandins. Prostaglandin nomenclature is well known and disclosed, e.g., in page 409, Remington's Pharmaceutical Sciences, 18th Edition, 1990, A. R. Gennaro, Ed., Mack Publishing Company, Easton, Pa. The term "prostaglandin" as used generically herein refers to the prostaglandin free acid and pharmaceutically acceptable derivatives thereof, including PGE<sub>1</sub>, PGA<sub>1</sub>, PGB<sub>1</sub>, PGF<sub>1α</sub>, 19-hydroxy-PGA<sub>1</sub>, 19-hydroxy-PGB<sub>1</sub>, PGE<sub>2</sub>, PGA<sub>2</sub>, PGB<sub>2</sub>, 19-hydroxy-PGA<sub>2</sub>, 19-hydroxy-PGB<sub>2</sub>, PGE<sub>3</sub>, PGF<sub>3α</sub>, carboprost tromethamine, dinoprost tromethamine, dinoprostone, lipoprost, gemeprost, metenoprost, sulprostone and tiaprostone as well as salts and esters thereof. Preferred prostaglandins for use in the

formulations of this invention include those prostaglandins comprising a  $\beta$  - hydroxyketone moiety, including D-series and E-series prostaglandins, preferably E-series prostaglandins such as prostaglandin E<sub>1</sub>, including pharmaceutically acceptable salts and lower alkyl esters thereof (the term "lower alkyl" as used herein means  
5 straight chain or branched chain alkyl containing one to four carbon atoms). Of the lower alkyl esters, the ethyl ester of prostaglandin E<sub>1</sub> (commercially available from Sigma Chemical Company, St. Louis, Mo., and preparable as disclosed, e.g., in U.S. Pat. No. 5,219,885, incorporated herein by reference) is preferred.

The biosynthesis of prostaglandins has been well characterized. See, e.g.,  
10 Lehninger at p. 687. In a typical biosynthetic pathway, exemplified by production of PGE<sub>2</sub>, the essential fatty acid linoleic acid is converted into the 20-carbon arachidonic acid, which is then acted upon by prostaglandin synthase, a dioxygenase enzyme. Oxygen atoms are added at carbon atoms 9 and 15, and the product is cyclized by formation of a bond between carbon atoms 8 and 12. In the presence of reduced  
15 glutathione, this cyclized product undergoes conversion into prostaglandin PGE<sub>2</sub>. Other types of naturally occurring prostaglandins are derived from different polyunsaturated fatty acids.

In about the 1960s, prostaglandins were isolated from a particular species of Caribbean coral, which made them more widely available for research. Catanzarite,  
20 Valerian A. and Gary Aisenbrey, *Contemporary OB/GYN* (October 1987), p. 22. A large number of natural and synthetic analogues of the prostaglandins are now known. Lehninger at 687.

The prostaglandins are known to produce often unpredictable effects over a very wide range of biological activities of a hormonal or regulatory nature.  
25 Prostaglandins have been reported to both lower and raise blood pressure, to inhibit gastric secretion, dilate bronchi, inhibit lipolysis, antagonize vasopressin-induced anti-diarrhesis, constrict the pupil, increase and decrease the intraocular pressure and produce contraction of the uterus. See, e.g., Ganong, William F., *Review of Medical Physiology*, 7th ed. (1975), p. 226 (hereinafter "Ganong"). The naturally occurring  
30 prostaglandins all appear to be capable of affecting the control of vascular and other smooth muscle contractions. In the central nervous system, prostaglandins are known to modify responses to certain synaptic transmitters. They have been reported to mimic the actions of some hormones and to inhibit the actions of certain others. See

Ganong at 226.

Two of the most extensively studied of the prostaglandins are PGE<sub>2</sub> and PGF<sub>2α</sub>. Both of these molecules are synthesized within the pregnant and non-pregnant uterus. While PGE<sub>2</sub> and PGF<sub>2α</sub> are similar in mediating some effects, they are different with respect to certain others. Both cause uterine contractions, but they predominate at different sites within the uterus—PGE<sub>2</sub> in the lower uterine segment, PGF<sub>2α</sub> in the fundal region. Both play important roles during labor, but PGE<sub>2</sub> has its major effect in cervical ripening, whereas PGF<sub>2α</sub> is more important in generating uterine contractions. PGE<sub>2</sub> elevates body temperature, whereas PGF<sub>2α</sub> has no apparent effect on body temperature. PGE<sub>2</sub> is vasodilator and bronchodilator, while PGF<sub>2α</sub> is a bronchoconstrictor and vasoconstrictor. See Catanzarite at 21–22.

Prostaglandins have been used in gynecology for pregnancy termination. Preparing the cervix with a prostaglandin suppository has been found to reduce the incidence of cervical laceration and significant bleeding. See Catanzarite at page 22. Synthetic analogues of prostaglandin PGE<sub>2</sub>, such as 16-16-dimethyl PGE<sub>2</sub> and 9-methylene PGE<sub>2</sub>, have proven useful for the induction of first trimester abortions. Such procedures typically use vaginal suppositories containing 20 milligrams PGE<sub>2</sub> or 3 milligrams of 15-methyl PGF<sub>2α</sub>, or by repeated intramyometrial injections of 15-methyl PGF<sub>2α</sub>, or by infusing a PGF<sub>2α</sub>-urea mixture (20 milligrams of PGF<sub>2α</sub> and 40 milligrams of urea in 100 mL of 5% dextrose in water) into the amniotic sac.

In obstetrics, prostaglandins have been used for cervical ripening, labor induction and control of post-partum hemorrhage. Catanzarite at 29. For cervical ripening, PGE<sub>2</sub> has been given intravenously, orally and vaginally, but the preferred route is intracervically. A PGE<sub>2</sub> gel is now commercially available in Scandinavia, and another PGE<sub>2</sub> gel is being investigated in the United States. The PGE<sub>2</sub> gel can also be used for labor induction (3–5 mg of PGE<sub>2</sub>, prepared by blending a 20 mg suppository with 60 mL of lubricating jelly and using 9–15 mL of the mixture, is placed in the vagina). Catanzarite at 32. Prostaglandins have also been utilized to control post-partum hemorrhage.

Topical and transdermal drug formulations are designed to deliver a therapeutically effective amount of drug to or across the skin of a patient. Devices known to the art include reservoir type devices involving membranes that control the rate of drug release to the skin, gels and creams, and devices involving a dispersion of

the drug in a matrix such as a pressure sensitive adhesive. As the skin presents a barrier to the drug it is often desirable or necessary to incorporate certain materials that enhance the rate at which the drug passes through the skin. For any particular drug, however, the type of device, the transdermal flux rate that is suitable, and suitable formulation components, are dependent upon the particular drug to be delivered.

Topical and transdermal administration of PGE<sub>1</sub> and PGE<sub>1</sub> derivatives have also been described, e.g., in U.S. Pat. Nos. 4,889,845 (Ritter et al.), 4,515,810 (Chow et al.), and 5,219,885 (Froelich et al.) and in Japanese Kokai 2-264725 (Morimoto et al.) and 63-135333 (Nakano et al.). In order for a transdermal formulation of PGE<sub>1</sub> or a derivative thereof to be effective and suitable it is desirable that the formulation have a high transdermal flux rate, allowing a therapeutically effective blood level of the drug to be achieved or maintained when the formulation is applied to a relatively small area of the skin. Furthermore PGE<sub>1</sub> readily undergoes certain reactions and rearrangements (see. e.g., J. Chromatography, 1991, 555, 73 (Lee et al.). This instability of the prostaglandin can be problematic in providing a suitable transdermal formulation.

#### SUMMARY OF THE INVENTION

The present invention provides topical compositions and methods of treatment for female sexual dysfunction. In preferred embodiments the present invention provides topical compositions and methods of treatment for female sexual arousal disorder. The invention also provides methods for ameliorating female sexual dysfunction by modulating arousal and the excitation and plateau phases of the female sexual response on demand by topical application of an effective amount of prostaglandin E<sub>1</sub>. In preferred embodiments, the topical composition of the present invention is applied about five to about twenty minutes before sexual intercourse. In preferred embodiments, the topical composition of the present invention is applied to two structures involved in female sexual arousal, the clitoris and the Graffenberg spot.

The composition of the invention is suitable for topical application, and comprises a vasoactive prostaglandin, more preferably prostaglandin E<sub>1</sub>, a penetration enhancer, a polymer thickener, a lipophilic component, and an acidic buffer system.

In some embodiments, the polymer thickener is a polyacrylic acid polymer. In other preferred embodiments, the polymer thickener is a polysaccharide gum or a modified polysaccharide gum. The lipophilic component is selected from the group consisting of the C<sub>1</sub> to C<sub>8</sub>, aliphatic alcohols, the C<sub>2</sub> to C<sub>30</sub> aliphatic esters and mixtures thereof.

5 The acidic buffer system is chosen to provide a suitable pH to minimize irritation of skin and mucous membranes. The composition is typically in the form of a cream, lotion, gel or other form suitable for topical application to skin and mucous membranes.

In a preferred embodiment, the present invention provides a composition

10 suitable for topical application comprising an effective amount of a vasoactive prostaglandin; a penetration enhancer selected from the group consisting of an alkyl-(N-substituted amino) alkanoate, an alkyl-2-(N,N-disubstituted amino) alkanoate, an (N-substituted amino) alkanol alkanoate, an (N,N-disubstituted amino) alkanol alkanoate, pharmaceutically acceptable salts thereof and mixtures

15 thereof; a polymer thickener selected from the group consisting of a polyacrylic acid polymer, a polysaccharide gum, a modified polysaccharide gum and mixtures thereof; a lipophilic component; and a buffer system wherein the pH of the composition is 3.0 to 7.4. Preferably the vasoactive prostaglandin is selected from the group consisting of prostaglandin E<sub>1</sub>, prostaglandin E<sub>1</sub> alkyl esters,

20 pharmaceutically acceptable salts thereof and mixtures thereof. In preferred embodiments, the composition further comprises polyethylene glycol.

In preferred embodiments, the invention provides a composition suitable for topical application comprising 0.001 weight percent to 1 weight percent of a vasoactive prostaglandin selected from the group consisting of PGE<sub>1</sub>,

25 pharmaceutically acceptable salts thereof, lower alkyl esters thereof and mixtures thereof; 0.01 weight percent to 5 weight percent modified polysaccharide gum; 0.5 weight percent to 10 percent weight dodecyl N,N-dimethylamino isopropionate or pharmaceutically acceptable salts thereof; 0.5 weight percent to 10 weight percent of a lower alcohol selected from the group consisting of ethanol, propanol, isopropanol

30 and mixtures thereof; 0.5 weight percent to 10 weight percent of an ester selected from the group consisting of ethyl laurate, isopropyl myristate, isopropyl laurate and mixtures thereof, based on the total weight of the composition, and a buffer system wherein the pH of the composition is 3.0 to 7.4. Preferably, the composition further



comprises 1 weight percent to 25 weight percent polyethylene glycol 400, based on the total weight of the composition.

The prostaglandin is present in a formulation of the invention in a therapeutically effective amount. Therapeutic effectiveness can be assessed, in part, with increase in vaginal secretion, increase in vaginal engorgement, increase in sexual responsiveness and increase in arousal.

The present invention provides the use of compositions comprising prostaglandin E<sub>1</sub> for the manufacture of a medicament for topical or transdermal administration to modulate sexual response in a human female. While not being tied to a specific mechanism, it is believed that prostaglandin E<sub>1</sub> acts directly on local tissues to produce effects such as increases in vaginal secretion, increases in vaginal engorgement, and acts indirectly on the central nervous system to increase sexual responsiveness and arousal.

The methods of the present invention may be used to improve or enhance arousal and sexual response in women whose sexual response is impaired as evidenced by diminished capacity to produce sufficient vaginal lubrication to facilitate comfortable penile penetration and by other symptoms of impaired sexual responsiveness. The invention thus provides a method of ameliorating female sexual arousal disorder, comprising the step of administering to a human female a composition suitable for topical application comprising an effective amount of a vasoactive prostaglandin, a penetration enhancer, a polymer carrier, a lipophilic component, and a buffer system, typically in a cream, lotion, gel or other suitable form. In some embodiments, the method comprises the step of applying a semi-solid prostaglandin composition to the genitalia of a human female in need of treatment, the composition comprising: a modified polysaccharide gum; an effective amount of a vasoactive prostaglandin selected from the group consisting of PGE<sub>1</sub>, pharmaceutically acceptable salts thereof, lower alkyl esters thereof and mixtures thereof; 0.5 percent to 10 percent DDAIP or a pharmaceutically acceptable salt thereof, based on the total weight of the composition; 0.5 percent to 10 percent, based on the total weight of the composition, of a lower alcohol selected from the group consisting of ethanol, propanol, isopropanol and mixtures thereof; 0.5 percent to 10 percent of an ester selected from the group consisting of ethyl laurate, isopropyl myristate, isopropyl laurate and mixtures thereof, based on the total weight of the

composition; and an acidic buffer system.

In a preferred embodiment, the topical composition of the present invention is applied approximately 5 to 20 minutes prior to sexual intercourse. In one embodiment, the composition is applied to the vaginal Grafenberg spot (G-spot) as well as the clitoris.

The present invention also provides a method for ameliorating female sexual arousal disorder, comprising the step of administering a composition to the genitalia of a human female in need of treatment, the composition comprising: an amount of a vasoactive prostaglandin effective to produce an increase in sexual arousal; a penetration enhancer selected from the group consisting of an alkyl-(N-substituted amino) alkanoate, an alkyl-2-(N,N-disubstituted amino) alkanoate, an (N-substituted amino) alkanol alkanoate, an (N,N-disubstituted amino) alkanol alkanoate, pharmaceutically acceptable salts thereof and mixtures thereof; a shear-thinning polymer; a lipophilic component selected from the group consisting of an aliphatic C<sub>1</sub> to C<sub>8</sub> alcohol, an aliphatic C<sub>8</sub> to C<sub>30</sub> ester, and mixtures thereof; and an acidic buffer system.

The effective amount of prostaglandin to be administered is selected to provide increased blood flow to the genitalia, which may be assessed by visual inspection, vaginal photoplethysmography, vaginal lubrication or engorgement. Alternatively, the effective amount to be administered is selected to provide increased sexual response, which may be assessed by visual inspection, vaginal photoplethysmography, vaginal lubrication and engorgement of the genitalia.

The effective amount of prostaglandin to be administered to increase arousal is selected to provide increased arousal as measured by self-report by a suitable questionnaire.

In the absence of any clinically diagnosed dysfunction in the female sexual response, the methods of the present invention may also be used to enhance the sexual response in a human female not suffering from a sexual dysfunction. The present invention will allow a more rapid response to sexual stimulation along with heightened sensation associated with excitement and plateau stages of the female sexual response by virtue of the increased blood flow to the tissues, as well as enhance subjective aspects, thereby leading to relatively increased arousal. The invention thus provides a method of enhancing female sexual arousal, comprising the

step of administering to a human female a composition suitable for topical application comprising an effective amount of a vasoactive prostaglandin, a penetration enhancer, a polymer carrier, a lipophilic component, and a buffer system, typically in a cream, lotion, gel or other suitable form.

5           The invention further provides a method of enhancing female sexual response, comprising the step of administering to a human female a composition suitable for topical application comprising an effective amount of a vasoactive prostaglandin, a penetration enhancer, a polymer carrier, a lipophilic component, and a buffer system, typically in a cream, lotion, gel or other suitable form.

## 10   BRIEF DESCRIPTION OF THE DRAWINGS

In the drawings,

Figure 1 is a graphical representation of the results of experiments in which the permeation of prostaglandin E<sub>1</sub> after application of different compositions to a model system, shed snake skin;

15           Figure 2 is a graphical representation of the results of experiments in which the permeation of prostaglandin E<sub>1</sub> after application of compositions comprising different concentrations of prostaglandin E<sub>1</sub> to a model system, stripped shed snake skin;

Figure 3 is a graphical representation of the results of experiments in which  
20 the permeation of prostaglandin E<sub>1</sub> is compared in two model systems, stripped shed snake skin and sheep vaginal membrane;

Figure 4 is a graphical representation of the baseline responses and the maximum treatment responses of vaginal blood flow;

25           Figure 5 is a graphical representation of the baseline responses and the maximum treatment responses to question 2 of the video assessment questionnaire;

Figure 6 is a graphical representation of the baseline responses and the maximum treatment responses to question 3 of the video assessment questionnaire;

Figure 7 is a graphical representation of the baseline responses and the maximum treatment responses to question 4 of the video assessment questionnaire;

30           Figure 8 is a graphical representation of the baseline responses and the maximum treatment responses to question 5 of the video assessment questionnaire;

Figure 9 is a graphical representation of the baseline responses and the maximum treatment responses to question 6 of the video assessment questionnaire; and

Figure 10 is a graphical representation of the baseline responses and the maximum treatment responses to question 9 of the video assessment questionnaire.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Before describing the present invention in detail, it is to be understood that this invention is not limited to particular drugs or drug delivery systems, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

It must be noted that, as used in this specification and the appended claims, the singular forms “a”, “an” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a vasoactive agent” includes a mixture of two or more such drugs, reference to “a penetration enhancer” includes mixtures of two or more enhancers, and the like.

In describing and claiming the present invention, the following terminology will be used in accordance with the definitions set out below.

The term “drug” or “pharmacologically active agent” as used herein is intended to mean a compound or composition of matter which, when administered to an organism (human or animal) induces a desired pharmacologic and/or physiologic effect by local and/or systemic action. As noted above, the pharmacologically active agents used in conjunction with the present invention are vasoactive agents.

By “transdermal” drug delivery, applicant is using the term in its conventional sense, i.e., to indicate delivery of a drug by passage into and through the skin and the underlying tissues and into the blood stream. By “transmucosal” drug delivery, applicant intends delivery of a drug by passage of a drug through the mucosal and underlying tissue into the blood stream. The compositions, systems, and methods of the invention, unless explicitly stated otherwise, should be presumed to be equally applicable to either transdermal or transmucosal modes of drug delivery.

“Penetration enhancement” or “permeation enhancement” as used herein relates to an increase in the permeability of the skin or mucosal tissue to a selected pharmacologically active agent, i.e., so that the rate at which the drug permeates

through the skin or mucosal tissue is increased. "Carriers" or "vehicles" as used herein refer to carrier materials suitable for transdermal or transmucosal drug administration, and include any such materials known in the art, e.g., any liquid, gel, solvent, liquid diluent, solubilizer, or the like, which is nontoxic and which does not interact with other components of the composition in a deleterious manner.

By an "effective" amount of a drug or pharmacologically active agent is meant a nontoxic but sufficient amount of the drug or agent to provide the desired effect.

In order to carry out the method of the invention, a composition suitable for topical application comprising a selected vasoactive agent is administered about fifteen minutes to about one hour prior to the time of desired effect. Preferably, the topical composition is applied once, twice or three times within a twenty-four hour period.

Suitable vasoactive agents include, but are not limited to: nitrates such as nitroglycerin, isosorbide dinitrate, erythrityl tetranitrate, amyl nitrate, sodium nitroprusside, molsidomine, linsidomine chlorhydrate and S-nitroso-N-acetyl-d,l-penicillamine ("SNAP"); long and short acting  $\alpha$ -blockers such as phenoxybenzamine, dibenamine, doxazosin, terazosin, phentolamine, tolazoline, prazosin, trimazosin, alfuzosin, tamsulosin and indoramin; ergot alkaloids such as ergotamine and ergotamine analogs, e.g., acetergamine, brazergoline, bromerguride, cianergoline, delorgotrile, disulergine, ergonovine maleate, ergotamine tartrate, etisulergine, lergotrile, lysergide, mesulergine, metergoline, metergotamine, nicergoline, pergolide, propisergide, proterguride and terguride; antihypertensive agents such as diazoxide, hydralazine and minoxidil; vasodilators such as nimodipine, pinacidil, cyclandelate, dipyridamole and isoxsuprine; chlorpromazine; haloperidol; yohimbine; trazodone and vasoactive intestinal peptides. Prostaglandin  $E_1$  and phentolamine are particularly preferred vasoactive agents for use in conjunction with the present method.

A dose of a prostaglandin  $E_1$  in an amount sufficient to enhance engorgement or vaginal secretion is topically administered to a woman. The appropriate doses of the particular vasodilating agent may be readily determined using methods described in Examples 3 and 4, below. The female response may be measured using methods described in Masters, W. H. and Johnson, V. E., *Human Sexual Response*, Little, Brown, and Co., Boston (1966) which is incorporated herein by reference.

Engorgement and redness of the external genitalia can be assessed by visual inspection. Methods for measuring blood flow, including Doppler ultrasonic velocimetry, thermography using for example an isothermal blood flow transducer, radioscintigraphic methods, vaginal photoplethysmography may be used as well as  
5 other methods well known in the art. In addition, measuring the contraction of the distal 1/3 as is characteristic of the plateau phase of female sexual response of the vagina may be measured using methods and equipment well known in the art including but not limited to strain gauges or other devices for measuring muscular contraction or muscle tension.

10 In addition, enhanced sexual response and heightened arousal can be measured using a questionnaire that requests that the female subject to describe any change in sensation brought about by administration of the prostaglandin composition by the methods of the present invention. In determining as suitable effective dose, appropriate placebo controls can be used to determine whether or not the observed  
15 effect is directly attributable to the administration of the prostaglandin composition. A suitable questionnaire for the measurement of enhanced sexual response and heightened arousal is provided below in Example 4.

A preferred embodiment of the present invention involves the topical administration of from at least 0.1 mg to about 6 mg of prostaglandin E<sub>1</sub> to a female  
20 from about one minute to about one hour before sexual intercourse, preferably about five minutes to about twenty minutes before sexual intercourse. In a preferred embodiment of the present invention, about 0.1 mg to about 2 mg of prostaglandin E<sub>1</sub> is administered topically. In another preferred embodiment of the present invention, about 1.4 mg to about 6 mg of prostaglandin E<sub>1</sub> is administered topically to a female.  
25 In another preferred embodiment of the present invention, about 1 mg to about 3 mg of prostaglandin E<sub>1</sub> is administered topically to a female.

A preferred composition comprises about 0.07 weight percent to about 0.4 weight percent of prostaglandin E<sub>1</sub> and a pharmaceutically acceptable excipient to form a composition suitable for topical application.

30 More particularly, in a preferred embodiment, the composition suitable for topical application of the present invention comprises:

- a) about 0.07 percent by weight of the total composition to about 0.4 percent by weight of the total composition of prostaglandin E<sub>1</sub>;

b) about 0.5 to about 5 percent by weight of the total composition of a suitable polymer;

c) about 70 to about 90 percent by weight of the total composition of a buffer;

5 d) about 0.5 to about 15 percent by weight of the total composition of a lipophilic component;

e) about 0.4 to about 5 percent by weight of the total composition of an emulsifier; and

f) about 50 to about 90 percent by weight of the total composition of water.

10 In one embodiment, the topical composition comprises 0.01 weight percent to 5 weight percent modified polysaccharide gum; 0.001 weight percent to 1 weight percent of a vasoactive prostaglandin selected from the group consisting of PGE1, pharmaceutically acceptable salts thereof, lower alkyl esters thereof and mixtures thereof; 0.5 weight percent to 10 percent weight dodecyl N,N-dimethylamino  
15 isopropionate or pharmaceutically acceptable salts thereof; 0.5 weight percent to 10 weight percent of a lower alcohol selected from the group consisting of ethanol, propanol, isopropanol and mixtures thereof; 0.5 weight percent to 10 weight percent of an ester selected from the group consisting of ethyl laurate, isopropyl myristate, isopropyl laurate and mixtures thereof, based on the total weight of the composition,  
20 and an acid buffer. In a preferred embodiment, the topical composition further comprises 1 weight percent to 25 weight percent, more preferably 3 weight percent to 20 weight percent polyethylene glycol 400, based on the total weight of the composition.

In addition, the present invention is concerned with a method for treating an  
25 human female suffering from sexual dysfunction. The method comprises administering an effective amount of the above disclosed topical composition by applying the topical composition to the genital area of a human female. In one embodiment of the present invention, the topical composition is applied to the labia, clitoris and the vulvar region of the vagina. In one embodiment of the present  
30 invention, the topical composition is applied to the vaginal Grafenberg spot (G-spot) and to the clitoris.

The stable, uniform, composition suitable for topical application of the present invention preferably contains prostaglandin as a vasoactive agent. Vasoactive

prostaglandins are those that act as peripheral vasodilators, including naturally occurring prostaglandins such as PGE<sub>1</sub>, PGA<sub>1</sub>, PGB<sub>1</sub>, PGF<sub>1 $\alpha$</sub> , 19-hydroxy-PGA<sub>1</sub>, 19-hydroxy-PGB<sub>1</sub>, PGE<sub>2</sub>, PGA<sub>2</sub>, PGB<sub>2</sub>, 19-hydroxy-PGA<sub>2</sub>, 19-hydroxy-PGB<sub>2</sub>, PGE<sub>3</sub>, PGF<sub>2 $\alpha$</sub> , PGF<sub>3 $\alpha$</sub> ; semisynthetic or synthetic derivatives of natural prostaglandins, including carboprost tromethamine, dinoprost tromethamine, dinoprostone, lipoprost, gemeprost, metenoprost, sulprostone and tiaprost. Prostaglandin E<sub>1</sub> and prostaglandin E<sub>2</sub> are particularly preferred vasoactive prostaglandins for use in conjunction with the present method.

The quantity of vasoactive prostaglandin, such as prostaglandin E<sub>1</sub>, in the pharmaceutical composition is a therapeutically effective amount and necessarily varies according to the desired dose, the dosage form (e.g., suppository or topical), and the particular form of vasoactive prostaglandin used. The term "prostaglandin" as used generically herein refers to the prostaglandin free acid and pharmaceutically acceptable derivatives thereof, including, for example PGE<sub>1</sub>, pharmaceutically acceptable salts and lower alkyl esters thereof (the term "lower alkyl" as used herein means straight chain or branched chain alkyl containing one to four carbon atoms). The composition generally contains between 0.001 weight percent to 1 weight percent of vasoactive prostaglandin, e.g., prostaglandin E<sub>1</sub>, typically contains between 0.05 weight percent to 1 weight percent, preferably from 0.1 weight percent to 0.5 weight percent, based on the total weight of the composition.

The amount that constitutes a therapeutically effective amount varies according to the particular prostaglandin to be delivered, the indication to be treated, the surface area of the skin and mucous membrane over which the formulation is to be placed, and on the other components of the composition. Accordingly it is not practical to enumerate particular preferred amounts but such can be readily determined by those skilled in the art with due consideration of these factors. Generally, however, the prostaglandin is present in an amount of about 0.07 to about 1 percent, preferably about 0.1 to about 1 percent by weight based on the total weight of the composition. In one preferred embodiment, prostaglandin E<sub>1</sub> is present in an amount of about 0.07 to about 0.4 percent by weight based on the total weight of the composition. The prostaglandin can be dissolved or substantially uniformly dispersed in the topical composition. It is preferably soluble (and dissolved) in the topical composition.




The topical composition can contain one or more penetration enhancers. Among the preferred penetration enhancers for the present invention are ethanol, propylene glycol, glycerol, ethyl laurate, isopropyl palmitate, isopropyl myristate, laurocapram (Azone™), dioxolanes (described in U.S. Patent No. 4,861,764),  
 5 macrocyclic ketones, HP-101, oxazolidones and biodegradable penetration enhancers (described in U.S. Patents Nos. 4,980,378 and 5,082,866 to Wong et al. and U.S. Patent Number 6,118,020 to Büyüktimkin et al. such as alkyl-2-(N-substituted amino) alkanooates (e.g., dodecyl N,N-dimethylamino isopropionate (DDAIP)), N-substituted amino alkanol alkanooates), acid addition salts and mixtures thereof.

10 The penetration enhancer is present in an amount sufficient to enhance the penetration of the prostaglandin E<sub>1</sub>. The specific amount varies necessarily according to the desired release rate and the specific form of prostaglandin E<sub>1</sub> used. Generally, the penetration enhancer is present in an amount ranging from about 0.5 weight percent to about 20 weight percent, based on the total weight of the composition.  
 15 Preferably, the penetration enhancer is present in an amount ranging from about 1 weight percent to about 10 weight percent of the composition. More preferably, the penetration enhancer is present in an amount ranging from about 1 weight percent to about 5 weight percent of the composition.

Preferred penetration enhancers include an alkyl-2-(N-substituted amino)-  
 20 alkanooate, a (N-substituted amino)-alkanol alkanooate, or a mixture of these. For convenient reference, alkyl-2-(N-substituted amino)-alkanoates and (N-substituted amino)-alkanol alkanooates can be grouped together under the term alkyl (N-substituted amino) esters.

25 Alkyl-2-(N-substituted amino)-alkanoates suitable for the present invention can be represented as follows:




wherein n is an integer having a value in the range of about 4 to about 18; R is

a member of the group consisting of hydrogen, C<sub>1</sub> to C<sub>7</sub> alkyl, benzyl and phenyl; R<sub>1</sub> and R<sub>2</sub> are members of the group consisting of hydrogen and C<sub>1</sub> to C<sub>7</sub> alkyl; and R<sub>3</sub> and R<sub>4</sub> are members of the group consisting of hydrogen, methyl and ethyl.

Preferred are alkyl (N,N-disubstituted amino)-alkanoates such as C<sub>4</sub> to C<sub>18</sub> alkyl (N,N-disubstituted amino)-acetates and C<sub>4</sub> to C<sub>18</sub> alkyl (N,N-disubstituted amino)-propionates and pharmaceutically acceptable salts and derivatives thereof. Exemplary specific alkyl-2-(N,N-disubstituted amino)-alkanoates include dodecyl 2-(N,N dimethylamino)-propionate (DDAIP). For example, the preparation of crystalline acid addition salts of DDAIP by cooled mixing of DDAIP with one of a select group of acids in the presence of a water-immiscible solvent such as hexane, is disclosed in U.S. Patent Number 6,118,020, the contents of which is incorporated herein by reference in its entirety. Acid addition salts of dodecyl 2-(N,N-dimethylamino)-propionate (DDAIP) can be inorganic as well as organic. Representative inorganic acid addition salts include the hydrochloric, hydrobromic, sulfuric, phosphoric, nitric acid addition salts of DDAIP, and their solvates. Exemplary organic acid addition salts include acetic, benzoic, salicylic, glycolic, succinic, nicotinic, tartaric, maleic, malic, palmoic, methanesulfonic, cyclohexanesulfamic, picric, and lactic acid addition salts, as well as their respective solvates. Preferred among the inorganic acid addition salts are DDAIP hydrogen chloride, and DDAIP dihydrogen sulfate.

Suitable (N-substituted amino)-alkanol alkanoates can be represented by the formula:



25

wherein n is an integer having a value in the range of about 5 to about 18; y is an integer having a value in the range of 0 to about 5; and R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and R<sub>7</sub> are members of the group consisting of hydrogen, C<sub>1</sub> to C<sub>8</sub> alkyl, and C<sub>3</sub> to C<sub>8</sub> aryl; and R<sub>8</sub> is a member of the group consisting of hydrogen, hydroxyl, C<sub>1</sub> to C<sub>8</sub>

alkyl, and C<sub>3</sub> to C<sub>8</sub> aryl. Preferred are (N-substituted amino)-alkanol alkanoates such as C<sub>5</sub> to C<sub>18</sub> carboxylic acid esters and pharmaceutically acceptable salts thereof. Preferred (N-substituted amino)-alkanol alkanoates include (N,N-disubstituted amino)-alkanol alkanoates.

5

In general, suitable penetration enhancers can be chosen from those listed above as well as sulfoxides, alcohols, fatty acids, fatty acid esters, polyols, amides, surfactants, terpenes, alkanones, organic acids and mixtures thereof. See generally Chattaraj, S.C. and Walker, R.B., Penetration Enhancer Classification, pp.5-20 in  
10 Maibach, H.I., and Smith, H.E., (eds.), Percutaneous Penetration Enhancers, CRC Press, Inc., Boca Raton, FL (1995) and Büyüktimkin, N., et al., Chemical Means of Transdermal Drug Permeation Enhancement, in Gosh, T.K., et al., (eds.) Transdermal and Topical Drug Delivery Systems, Interpharm Press, Inc., Buffalo Grove, IL  
(1997). Suitable sulfoxides include dimethylsulfoxide, decylmethylsulfoxide and  
15 mixtures thereof. Suitable alcohols include ethanol, propanol, butanol, pentanol, hexanol, octanol, nonanol, decanol, 2-butanol, 2-pentanol, benzyl alcohol, caprylic alcohol, decyl alcohol, lauryl alcohol, 2-lauryl alcohol, myristyl alcohol, cetyl alcohol, stearyl alcohol, olcyl alcohol, linolyl alcohol, linolenyl alcohol and mixtures thereof. Suitable fatty acids include valeric, heptanoic, pelargonic, caproic, capric,  
20 lauric, myristic, stearic, oleic, linoleic, linolenic, caprylic, isovaleric, neopentanoic, neoheptanoic, neononanoic, trimethyl hexanoic, neodecanoic and isostearic acids and mixtures thereof.

Suitable fatty acid esters include isopropyl n-butyrate, isopropyl n-hexanoate, isopropyl n-decanoate, isopropyl myristate, isopropyl palmitate, octyldodecyl  
25 myristate, ethyl acetate, butyl acetate, methyl acetate, methylvalerate, methylpropionate, diethyl sebacate, ethyl oleate, ethyl laurate and mixtures thereof. Suitable polyols include propylene glycol, polyethylene glycol, ethylene glycol, diethylene glycol, triethylene glycol, dipropylene glycol, glycerol, propanediol, sorbitol, dextrans, butanediol, pentanediol, hexanetriol and mixtures thereof.

30 Suitable amides include urea, dimethylacetamide, diethyltoluamide, dimethylformamide, dimethyloctamide, dimethyldecamide, 1-alkyl-4-imidazolin-2-one, pyrrolidone derivatives, cyclic amides, hexamethylenelauramide and its derivatives, diethanolamine, triethanolamine and mixtures thereof. Suitable

pyrrolidone derivatives include 1-methyl-2-pyrrolidone, 2-pyrrolidone, 1-lauryl-2-pyrrolidone, 1-methyl-4-carboxy-2-pyrrolidone, 1-hexyl-4-carboxy-2-pyrrolidone, 1-lauryl-4-carboxy-2-pyrrolidone, 1-decyl-thioethyl-2-pyrrolidone (HP-101), 1-methyl-4-methoxycarbonyl-2-pyrrolidone, 1-hexyl-4-methoxycarbonyl-2-pyrrolidone, 1-lauryl-4-methoxycarbonyl-2-pyrrolidone, N-cyclohexylpyrrolidone, N-dimethylaminopropylpyrrolidone, N-cocoalkylpyrrolidone, N-tallowalkylpyrrolidone, fatty acid esters of N-(2-hydroxymethyl)-2-pyrrolidone and mixtures thereof. Suitable cyclic amides include 1-dodecylazacycloheptane-2-one (laurocapram, Azone®), 1-geranylazacycloheptan-2-one, 1-farnesylazacycloheptan-2-one, 1-geranylgeranylazacycloheptan-2-one, 1-(3,7-dimethyloctyl)azacycloheptan-2-one, 1-(3,7,11-trimethyloctyl)azacycloheptan-2-one, 1-geranylazacyclohexane-2-one, 1-geranylazacyclopentan-2,5-dione, 1-farnesylazacyclopentan-2-one and mixtures thereof.

Suitable surfactants include anionic surfactants, cationic surfactants, nonionic surfactants, bile salts and lecithin. Suitable anionic surfactants include sodium laurate, sodium lauryl sulfate and mixtures thereof. Suitable cationic surfactants include cetyltrimethylammonium bromide, tetradecyltrimethylammonium bromide, benzalkonium chloride, octadecyltrimethylammonium chloride, cetylpyridinium chloride, dodecyltrimethylammonium chloride, hexadecyltrimethylammonium chloride, and mixtures thereof. Suitable nonionic surfactants include  $\alpha$ -hydro- $\omega$ -hydroxy-poly(oxyethylene)-poly(oxypropyl) poly(oxyethylene)block copolymers, polyoxyethylene ethers, polyoxyethylene sorbitan esters, polyethylene glycol esters of fatty alcohols and mixtures thereof. Suitable  $\alpha$ -hydro- $\omega$ -hydroxy-poly(oxyethylene)-poly(oxypropyl) poly(oxyethylene)block copolymers include Poloxamers 231, 182, and 184 and mixtures thereof. Suitable polyoxyethylene ethers include 4-lauryl ether (BRIJ 30™), (BRIJ 93™), (BRIJ 96™), 20-oleyl ether (BRIJ 99™) and mixtures thereof. Suitable polyoxyethylene sorbitan esters include the monolaurate (TWEEN 20™, SPAN 20™) the monopalmitate (TWEEN 40™), the monostearate (TWEEN 60™), and the monooleate (TWEEN 80™) and mixtures thereof. Suitable polyethylene glycol esters of fatty acids include the 8-oxyethylene stearate ester (MYRJ 45™), (MYRJ 51™), the 40-oxyethylene stearate ester (MYRJ 52™) and mixtures thereof. Suitable bile salts include sodium cholate, sodium salts of

laurocholic, glycolic and desoxycholic acids and mixtures thereof.

Suitable terpenes include D-limonene,  $\alpha$ -pinene,  $\beta$ -enrene,  $\alpha$ -terpineol, terpinen-4-ol, carvol, carvone, pulegone, piperitone, menthone, menthol, geraniol, cyclohexene oxide, limonene oxide,  $\alpha$ -pinene oxide, cyclopentene oxide, 1,8-cineole, ylang ylang oil, anise oil, chenopodium oil, eucalyptus oil and mixtures thereof.

Suitable alkanones include N-heptane, N-octane, N-nonane, N-decane, N-undecane, N-dodecane, N-tridecane, N-tetradecane, N-hexadecane and mixtures thereof.

Suitable organic acids include citric acid, succinic acid, salicylic acid, salicylates (including the methyl, ethyl and propyl glycol derivatives), tartaric acid and mixtures thereof.

Natural and modified polysaccharide gums are also an important ingredient of the composition. Suitable representative gums are those in the natural and modified galactomannan gum category. A galactomannan gum is a carbohydrate polymer containing D-galactose and D-mannose units, or other derivatives of such a polymer. There is a relatively large number of galactomannans, which vary in composition depending on their origin. The galactomannan gum is characterized by a linear structure of  $\beta$ -D-mannopyranosyl units linked (1 $\rightarrow$ 4). Single membered  $\alpha$ -D-manopyranosyl units, linked (1 $\rightarrow$ 6) with the main chain, are present as side branches. Galactomannan gums include guar gum, which is the pulverized endosperm of the seed of either of two leguminous plants (*Cyamopsis tetragonalobus* and *psoraloids*) and locust bean gum, which is found in the endosperm of the seeds of the carobtree (*ceratonia siliqua*). Suitable modified polysaccharide gums include ethers of natural or substituted polysaccharide gums, such as carboxymethyl ethers, ethylene glycol ethers and propylene glycol ethers. An exemplary substituted polysaccharide gum is methylcellulose.

Other suitable representative gums include agar gum, carrageenan gum, ghatti gum, karaya gum, rhamsan gum and xanthan gum. The composition of the present invention may contain a mixture of various gums, or mixture of gums and acidic polymers.

Gums, and galactomannan gums in particular, are well-known materials. See for instance, *Industrial Gums: Polysaccharides & Their Derivatives*, Whistler R. L. and BeMiller J.N. (eds.), 3rd Ed. Academic Press (1992) and Davidson R. L., *Handbook of Water-Soluble Gums & Resins*, McGraw-Hill, Inc., N.Y. (1980). Most

gums are commercially available in various forms, commonly a powder, and ready for use in foods and topical compositions. For example, locust bean gum in powdered form is available from Tic Gums Inc. (Belcam, MD).

When present, the polysaccharide gums are present in the range from about  
5 0.1 percent to about 5 percent, based on the total weight of the composition, with the preferred range being from 0.5 percent to 3 percent. In one preferred embodiment, 2.5 percent by weight of a polysaccharide gum is present. Illustrative compositions are given in the examples, below.

An optional alternative to the polysaccharide gum is a polyacrylic acid  
10 polymer. A common variety of polyacrylic acid polymer is known generically as "carbomer." Carbomer is polyacrylic acid polymers lightly cross-linked with polyalkenyl polyether. It is commercially available from the B. F. Goodrich Company (Akron, Ohio) under the designation "CARBOPOL™." A particularly preferred variety of carbomer is that designated as "CARBOPOL 940."

15 Other polyacrylic acid polymers suitable for use are those commercially available under the designations "Pemulen™" (B. F. Goodrich Company) and "POLYCARBOPHIL™" (A.H. Robbins, Richmond, VA). The Pemulen™ polymers are copolymers of C<sub>10</sub> to C<sub>30</sub> alkyl acrylates and one or more monomers of acrylic acid, methacrylic acid or one of their simple esters crosslinked with an allyl ether of  
20 sucrose or an allyl ether of pentaerythritol. The POLYCARBOPHIL™ enhancer is a polyacrylic acid cross-linked with divinyl glycol.

Where polyacrylic acid polymers are present, they represent about 0.5 percent to about 5 percent of the composition, based on its total weight.

Another important component of the present invention is a lipophilic  
25 component. The term lipophilic component as used herein refers to an agent, preferably a mixture of agents, that is both lipophilic and hydrophilic. The C<sub>1</sub> to C<sub>8</sub>, aliphatic alcohols, the C<sub>2</sub> to C<sub>30</sub> aliphatic esters, and their mixtures can serve as lipophilic component. Illustrative suitable alcohols are ethanol, n-propanol and isopropanol, while suitable esters are ethyl acetate, butyl acetate, ethyl laurate, methyl  
30 propionate, isopropyl palmitate and isopropyl myristate. As used herein, the term "aliphatic alcohol" includes polyols such as glycerol, propylene glycol and polyethylene glycols. A mixture of alcohol and ester is preferred, and in particular, a mixture of ethanol and ethyl laurate is most preferred. The concentration of

lipophilic component required necessarily varies according to other factors such as the desired semi-solid consistency and the desired skin penetration promoting effects. The preferred topical composition contains lipophilic compound in the range of 7 percent to 40 percent by weight based on the total weight of the composition. Where  
5 a lipophilic component that is a mixture of aliphatic alcohol and aliphatic ester is used, the preferred amount of alcohol is in the range of 5 percent to 15 percent, while that of aliphatic ester is in the range from 2 percent to 15 percent (again based on the total weight of the composition). Where polyethylene glycol is used, polyethylene glycol is present in the amount of about 1 weight percent to about 25 weight percent,  
10 based on the total weight of the composition. A preferred polyethylene glycol is polyethylene glycol 400 (PEG 400). When present, polyethylene glycol 400 is about about 1 weight percent to about 25 weight percent, preferably about 3 weight percent to about 20 weight percent, based on the total weight of the composition.

In one embodiment, the C<sub>2</sub> to C<sub>30</sub> aliphatic esters, and their mixtures  
15 comprising the lipophilic compound include C<sub>8</sub> to C<sub>30</sub> aliphatic esters of glycerol selected from the group consisting monoglycerides, diglycerides, triglycerides, and mixtures thereof. Suitable aliphatic esters include glyceryl esters of saturated fatty acids, unsaturated fatty acids and mixtures thereof. Suitable saturated fatty acids include caproic acid, caprylic acid, capric acid, lauric acid, myristic acid, palmitic  
20 acid, stearic acid, arachidic acid, behenic acid and lignoceric acid. Suitable unsaturated fatty acids include oleic acid, linoleic acid and linolenic acid. Suitable glyceryl esters include glyceryl monooleate, triolein, trimyristin and tristearin, preferably trimyristin.

The concentration of lipophilic compound required necessarily varies  
25 according to other factors such as the desired semi-solid consistency and the desired skin penetration promoting effects. Suitably the concentration of lipophilic compound is in the range of 0.5 percent to 40 percent by weight based on the total weight of the composition. The preferred topical composition contains lipophilic compound in the range of 7 percent to 40 percent by weight based on the total weight of the  
30 composition.

Where a mixture of aliphatic alcohol and aliphatic ester are employed, the suitable amount of alcohol is in the range of about 0.5 percent to 10 percent. In one preferred embodiment, the amount of alcohol is in the range of about 5 percent to 15

percent, while that of aliphatic ester is in the range of about 2 percent to 15 percent (again based on the total weight of the composition). In another preferred embodiment, the amount of alcohol is in the range of about 0.5 percent to 10 percent, while that of aliphatic ester is in the range of 0 percent to 10 percent (again based on  
5 the total weight of the composition).

An optional, but preferred, component of the present invention is an emulsifier. Although not a critical factor, preferable emulsifiers generally exhibit a hydrophilic-lipophilic balance (HLB) number of at least 9. Sucrose esters, and specifically sucrose stearate, can serve as emulsifiers for the topical composition of  
10 the present invention. Sucrose stearate is a well known emulsifier available from various commercial sources.

Typical non-ionic surfactants include the polysorbates, which are mixtures of partial esters of sorbitol and its mono- and dianhydrides, typically condensed with approximately 20 mol of ethylene oxide; polyethoxylated alkyl ethers and esters, in  
15 which the alkyl chain can be either saturated, unsaturated, branched or linear; polyethoxylated alkyl phenols, in which the hydrophobic group normally octyl or nonylphenol; and poloxamers, polyoxyethylene-polyoxypropylene block copolymers, in which the polyoxypropylene chain acts as the hydrophobic moiety. Some commercially available non-ionic surfactants are BRIJ 99™, BRIJ 78™, polyoxyl 40  
20 stearate and polysorbate 80. BRIJ 99™ and BRIJ 78™ are polyethylene glycol fatty alcohol ethers. Polyoxyl 40 stearate is a mixture of mono- and distearate esters of polyoxyethylene and of free polyoxyethylene. Polysorbate 80 is polyoxyethylene (20) sorbitan monooleate.

When an emulsifier is used, sucrose stearate present up to about 2 percent, based on the total weight of the composition, is preferred. The preferred amount of sucrose stearate emulsifier can also be expressed as a weight ratio of emulsifier to polysaccharide gum. A ratio of 1 to 6 emulsifier to gum is preferred generate the  
25 desired semi-solid consistency and separation resistance.

The present invention includes a buffer system. Buffer systems serve to  
30 maintain or buffer the pH of compositions within a desired range. The term "buffer system" or "buffer" as used herein has reference to a solute agent or agents which, when in a water solution, stabilize such solution against a major change in pH (or hydrogen ion concentration or activity) when acids or bases are added thereto. Solute



agent or agents which are thus responsible for a resistance to change in pH from a starting buffered pH value in the range indicated above are well known. While there are numerous other suitable buffers, such as acetate buffers, potassium phosphate monohydrate has proven effective for compositions of the present invention.

5           The final pH value of the pharmaceutical composition of the present invention may vary within the physiologically compatible range. Necessarily, the final pH value is not irritating to human skin. Without violating this constraint, the pH may be selected to improve prostaglandin E<sub>1</sub>, stability and to adjust consistency when required. With these factors accounted for, the preferred pH value is about 3.0 to 7.4.  
10   The most preferred pH range is from about 3.5 to about 6.0.

          The remaining component of the composition is water, which is preferably purified. The composition contains water in the range of about 50 to about 90 percent, optionally including at least some of the water in the buffer, based on the total weight of the composition. The specific amount of water present is not critical,  
15   however, being adjustable to obtain the desired consistency and/or concentration of the other components.

          Additionally, known transdermal penetration enhancers can also be added, if desired. Illustrative are dimethyl sulfoxide (DMSO), dimethyl acetamide (DMA), 2-pyrrolidone, N,N-diethyl-m-toluamide (DEET), 1-dodecylazacycloheptane-2-one  
20   (laurocapram, Azone®, a registered trademark of Nelson Research), N,N-dimethylformamide, N-methyl-2-pyrrolidone, calcium thioglycolate, oxazolidinones, alkyl-2-(N-substituted amino) alkanooates and their acid addition salts (e.g., dodecyl N,N-dimethylamino isopropionate (DDAIP)), N-substituted amino alkanol  
25   alkanoates and their acid addition salts, dioxolane derivatives, laurocapram derivatives, macrocyclic enhancers such as macrocyclic ketones and mixtures thereof.

          Stabilizers, coloring agents, rheological agents, fragrances and preservatives can be added to the extent that they do not overly limit prostaglandin E<sub>1</sub> skin penetration or prevent the desired semi-solid consistency. When present, such are usually added in amounts of about 0.05 to about 0.30%. Suitable preservatives  
30   include methylparabens (methyl PABA), propylparabens (propyl PABA) and butylhydroxy toluene (BHT).

          The compositions of the present invention can also include a small amount of a topical anesthetic, if desired. When present, the topical anesthetic comprises about

0.01 to about 20 percent by weight, preferably about 0.01 to about 10 percent by weight based on the weight of the composition. As can be recognized, the suitable concentration of topical anesthetic will vary, depending the specific anesthetic and the presence of other components. For example, suitable concentrations include  
5 about 1 to about 20 percent by weight of benzocaine, about 0.25 to about 2.5 percent by weight of dibucaine, about 0.01 to about 10 percent by weight of lidocaine, or about 0.25 to about 1 percent by weight of tetracaine.

In preferred embodiments, the topical composition comprises at least one local anesthetic. Suitable local anesthetics include those approved for topical  
10 application, including, but not limited to ambucaine, amolanone, amylocaine hydrochloride, articaine, benoxinate, benzocaine, betoxycaine, biphenamine, bupivacaine, butacaine, butamben, butanilicaine, butethamine, butoxycaine, carticaine, chlorprocaine hydrochloride, cocaethylene, cocaine, cyclomethycaine, dibucaine hydrochloride, dimethocaine, diperodon hydrochloride, dyclonine,  
15 ecgonidine, ecgonine, ethyl chloride, etidocaine, beta-eucaine, euprocin, fenalcomine, fomocaine, hexylcaine hydrochloride, hydroxytetracaine, isobutyl p-aminobenzoate, leucinocaine mesylate, levoadrol, lidocaine, mepivacaine, meprylcaine, metabutoxycaine, methyl chloride, myrtecaine, naepaine, octacaine, orthocaine, oxethazaine, parethoxycaine, phenacaine hydrochloride, phenol, piperocaine,  
20 piridocaine, polidocanol, pramoxine, prilocaine, procaine, propanocaine, proparacaine, propipocaine, propoxycaine hydrochloride, pseudococaine, pyrrocaine, ropivacaine, salicyl alcohol, tetracaine hydrochloride, tolycaine, trimecaine, zolamine and mixtures thereof.

In general, with few exceptions, useful local anesthetics contain a lipophilic  
25 radical (mostly of aromatic structure), an intermediate chain and a hydrophilic radical (often an amino group). Local anesthetics can be further classified chemically as alcohols and alkyl ethers (such as chlorbutanol, benzyl alcohol, saligenine and pistocaine), amines, amino alcohols, amino alkyl ethers (such as pramocaine and dimethisoquine), amino ketones (such as falicaine), carboxylic acid esters (such as  
30 benzocaine, procaine and parophoxycaine), carboxylic acid amides (such as lidocaine and dibucaine), carbamic acid esters (such as diperodone) and amidines and guanidines (such as phenacaine and guanicaine). See Büchi, J., and Perlia, X., "Structure - Activity Relations and Physico-Chemical Properties of Local

Anesthetics. Part I. Relations between Chemical Structure and Local Anesthetic Activity," pp. 39-130 in Int. Encycl. Pharm. Therapeut., Local Anesthetics, Vol. I, Pergamon Press, New York, 1971.

In preferred embodiments, the local anesthetic molecular structure consists of a tertiary amine linked to a substituted aromatic ring by an intermediate chain. In some embodiments, the intermediate chain includes both a carbonyl group and one or more alkyl groups. The intermediate chain may further contain an ester linkage or an amide linkage. Suitable aminoamide local anesthetics include lidocaine, bupivacaine, mepivacaine, dibucaine, propivacaine, etidocaine and tlocainide. Suitable aminoester local anesthetics include procaine, chlorprocaine, tetracaine, isocaine, benzocaine, and monocaine. In embodiments in which the intermediate chain includes both a carbonyl group and one or more alkyl groups, a preferred local anesthetic is dyclonine, 1-(4-butoxyphenyl)-3-(1-piperidynyl)-1-propanone.

Preferred local anesthetics are those producing a moderate duration of anesthesia, more preferably those having a long duration of anesthetic action. For example, procaine and chlorprocaine have a short duration of action. Lidocaine, mepivacaine and prilocaine produce a moderate duration of anesthesia. Suitable long-acting local anesthetics include ropivacaine, tetracaine, bupivacaine and etidocaine.

Contemplated dosage forms of the semi-solid pharmaceutical composition of the present invention are creams, gels, and the like, also including but not limited to compositions suitable for use with transdermal patches and like devices.

The semi-solid composition of the present invention has a suitably chosen viscosity such that the composition is naturally retained where applied. The semi-solid composition can exhibit Newtonian or non-Newtonian rheological characteristics. In some preferred embodiments, the semi-solid composition of the present invention exhibits non-Newtonian rheological characteristics, i.e. in which the apparent viscosity is dependent on the shear rate applied to the composition. Preferably the composition has "shear-thinning" rheological properties. As used herein, "shear-thinning" refers to a reduction in apparent viscosity (the ratio of shear stress to the shear rate) with increasing shear rate, whether the reduction in apparent viscosity is time independent (pseudoplastic), time dependent (thixotropic) or

associated with a yield stress, defined as a stress that must be exceeded before flow starts, (Bingham plastics and generalized Bingham plastics). See, generally, Harris, J., & Wilkinson, W.L., "Non-newtonian Fluid," pp.856-858 in Parker, S.P., ed., McGraw-Hill Encyclopedia of Physics, Second Edition, McGraw-Hill, New York, 1993. Suitable viscosity ranges from about 5,000 centipoise (cps) to about 20,000 cps, preferably from about 7,000 cps to about 13,000 cps.

The topical composition is applied about five to about twenty minutes before sexual intercourse to the labia, clitoris and vagina and massaged until absorption is complete. In a preferred embodiment of the present invention, the topical composition is applied to the clitoris and the vaginal Grafenberg spot (also known as Grafenberg zone or G-spot, a region on the inner anterior wall of the vagina about 5 cm from the vaginal opening about five to about twenty minutes before sexual intercourse. Amounts of the topical composition ranging between about 0.1 and about 10 grams, preferably about 0.1 to about 3 grams are sufficient for vasodilation and the beneficial effects to occur. The present invention can be used with or without benefit of erotic stimuli. The determination of an ideal dose of the composition should be determined with each individual by one skilled in the art, such as a physician or sex therapist. The effective amount to be administered is selected to provide increased blood flow to the genitalia, which may be assessed by visual inspection, vaginal photoplethysmography, vaginal lubrication or engorgement. The preferred active component is prostaglandin, most preferably prostaglandin E<sub>1</sub>. Suitable doses of these selected drugs and other suitable drugs, such as phentolamine will be apparent to those skilled in the art, or may be deduced from the literature in combination with the teaching of the present disclosure.

While this invention has been described by way of preferred embodiments, the examples set out herein are not intended to limit the scope of the invention which contemplates the use of any pharmacologic vasodilating drug capable of absorption into the local and systemic circulation upon administration of the drug via the transmucosal, transdermal, intranasal, buccal or rectal routes of administration.

Numerous other advantages of the present invention will be apparent from the following detailed description of the invention including the accompanying examples and the appended claims.

## EXAMPLE 1

## Formulation of Suitable Compositions

Composition I was prepared as follows according to Formulation I (Table 1, below). Part A was formed by dissolving about 0.4 parts prostaglandin E<sub>1</sub> (Alprostadi USP) in about 5 parts ethyl alcohol. Next, about 5 parts ethyl laurate were mixed into the alcohol-prostaglandin E<sub>1</sub> solution. Part B was prepared starting from a pH 5.5 water/buffer solution. The water/buffer solution was prepared by adding sufficient potassium phosphate monobasic to purified water to create a 0.1 M solution. The water/buffer solution diluted to a final concentration of about 0.05M and about pH 5.5, adjusted with a strong base solution (1 N sodium hydroxide) and a strong acid (1 N phosphoric acid). Suitable buffer concentrations range from about 0.005M to about 1.0M. Preferred buffer concentrations range from about 0.05M to about 0.2M. In several preferred embodiments the buffer concentration is 0.1M. Propylene glycol (about 5 parts) was added to the water / buffer solution, and then the polyacrylic polymer (about 1 part ) was dispersed in the propylene glycol / water / buffer solution. All parts specified herein are parts by weight.

Parts A and B were mixed and homogenized using a homogenizer. Table 1, below, contains a list of ingredients and proportions. The resulting composition was a spreadable, semi-solid suitable for application to the skin and mucous membranes without the need for supporting devices such as patches and adhesive strips. The composition was both homogenous in appearance and resistant to separation. Compositions based on formulations II-VII were prepared following the same procedure.

Table 1							
Component	Formulation I (weight %)	II	III	IV	V	VI	VII
Noveon AA-1	1	1	1	1			
Ethanol	5	5		5	5	5	5
Propylene glycol	5	5	5				
Ethyl laurate	5	5		5	5	3	3
70% Sorbitol		5					
Glycerol			5				
DDAIP				5		2.5	
DDAIP HCl							2.5
Sesame oil			5				
Squalene			5				
Prehydrated Locust bean gum					3		
Modified Guar Gum						3	2.5
Sucrose stearate					0.5		

0.05M pH 5.5 buffer	78.85	73.85	73.85	78.85	86.1	81	
0.1M pH 5.5 buffer							87
1 M NaOH	4.75	4.75	4.75	4.75			
Prostaglandin E <sub>1</sub>	0.4	0.4	0.4	0.4	0.4	0.3	0.3

As noted above, in other embodiments, such as Compositions VI and VII, the composition may include a modified polysaccharide gum, suitably a modified galactomannan gum, such as a guar gum. Alternatively, a polyacrylic polymer may be used instead of the polysaccharide gum.

5

## EXAMPLE 2

### *In Vitro* Penetration Of Different Formulations

The relative ability of compositions prepared according to the formulations of Table 1 to provide prostaglandin E<sub>1</sub> was studied in two *in vitro* model systems corresponding to skin and mucosal membranes: shed snake skin and sheep vaginal  
10 membrane. The results are presented in Figures 1 – 3.

Compositions were evaluated for skin penetration using shed snake skin as a model barrier. Shed snake skin was obtained from the Animal Care Unit of the University of Kansas. With head and tail sections removed, the skin was randomly divided into test sections and then hydrated by soaking.

15

Samples of the compositions listed in Table 1 were evaluated using modified Franz-type diffusion cells (surface area 1.8 cm<sup>2</sup>). Specifically, skin pieces were mounted on top of a receptor cell of a vertical diffusion cell assembly in which a small magnetic bar was inserted and filled with an isotonic buffer. A seal was placed on top of the skin section followed by a donor cell. The two cells were then clamped  
20 together. Known amounts of the formulations were applied on the bottom of a small capped vial (weight about 5 grams) which fits exactly to the donor cell to ensure uniform distribution. The vials were placed on the skin in the donor cell. To reduce the evaporation of the ingredients, the donor cell and vial were gently taped together with a water-resistant adhesive band. The cells were transferred to a stirred water  
25 bath that was maintained at 37 degrees Celsius. Samples were withdrawn from the cells each hour for four hours and analyzed for the concentration of prostaglandin E<sub>1</sub>, with changes in concentration indicating the amount penetrating. Tests with multiple skin samples from the same snake yielded data that were averaged.

For a discussion of the use of shed snake skin in the evaluation of drug

penetration, see U.S. Patent No. 4,771,004 to Higuchi, which is incorporated here by reference to the extent that it is not inconsistent.

The results of the penetration study are presented in Figure 1 and in Table 2, below. Prostaglandin E<sub>1</sub> penetrated quickly at a relatively sustained rate for four hours from compositions prepared based on Formulations I, II and III. In contrast, relatively little penetration was observed using compositions based on Formulations IV and V.

Table 2  
Prostaglandin E<sub>1</sub> Average Cumulative Amount ( $\mu\text{g}/\text{cm}^2$ )

Hour	Formulation I	Formulation II	Formulation III	Formulation IV	Formulation V
1	5.00	2.89	1.58	3.55	0.39
2	8.42	6.32	2.11	8.42	0.92
3	12.37	11.58	2.11	16.58	2.11
4	18.68	17.11	1.58	23.82	4.21

10

### EXAMPLE 3

#### Concentration Effects On *In Vitro* Penetration

The effect of the prostaglandin E<sub>1</sub> concentration on permeation was studied using stripped shed snake skin. Stripped shed snake skin was prepared by removing the outer scale layer of the shed snake skin by 3-5 cycles of application and removal of adhesive tape (Minnesota Mining and Manufacturing Co., St. Paul, MN). The compositions tested were prepared as described in Example 1, and had final proportions (parts) of prostaglandin E<sub>1</sub>, (either 0.05%, 0.1%, or 0.2%); ethanol, about 5 parts; propylene glycol, about 5 parts; ethyl laurate, about 5 parts; polyacrylic polymer, about 1 part; 1M NaOH about 4.75 parts; 0.005M phosphate buffer, about pH 5.5, q.s. 100.

20

Penetration studies were performed as described in Example 2. The results are shown in Figure 2 and Table 3, below. Higher prostaglandin E<sub>1</sub> concentrations produce both more rapid permeation and a higher amount delivered.

Table 3  
Prostaglandin E<sub>1</sub>  
Cumulative Amount (μg/cm<sup>2</sup>)

Hour	0.05% Prostaglandin E <sub>1</sub> (Open Squares)	0.1% Prostaglandin E <sub>1</sub> (Filled Triangles)	0.2% Prostaglandin E <sub>1</sub> (Filled Squares)
1	25	33.75	41
2	40	65	81
3	50	85	118
4	58.75	102.5	143

#### EXAMPLE 4

##### 5 Comparison of Permeation in Two Model Membrane Systems

The permeation of prostaglandin E<sub>1</sub> in a topical composition of the present invention was compared using the stripped shed snake skin of Example 3 and the sheep vaginal membrane *in vitro* system. The topical composition used was the 0.2% prostaglandin E<sub>1</sub> composition of Example 3.

10 Sheep vaginas were obtained from a local slaughterhouse. The freshly excised organ was refrigerated and used immediately. After excision, the outer wall of the vagina was carefully separated from any adhering tissue, taking care to avoid damage. The vagina was cut open longitudinally (vertically) (Kabadi, M.B., and Chien, Y.W., Intravaginal controlled administration of Flurogestone acetate. II: 15 Development of an *in vitro* system for studying the intravaginal release and permeation of Flurogestone acetate, J. Pharm. Sci. 73: 1464-1468 (1984)). The vaginal mucosa were separated from the interior of the vaginal wall, soaked in nanopure water, cut into appropriately sized pieces, and mounted in using modified Franz-type diffusion cells as described in Example 2.

20 Penetration studies were performed as described in Example 2. The results are shown in Figure 3 and Table 4, below. The penetration measured in stripped shed snake is comparable to that measured in sheep vaginal membrane over the first two hours, diverging slightly at three hours.

Table 4  
Prostaglandin E<sub>1</sub>  
Average Cumulative Amount (μg/cm<sup>2</sup>)

Hour	Stripped Shed Snake Skin (Membrane I)	Sheep Vaginal Membrane (Membrane II)
1	11.67	8.33
2	30.00	32.92
3	50.00	60.00
4	55.83	85.42

25



## EXAMPLE 5

## Clinical Study in Women Suffering from Sexual Dysfunction

This study was conducted to evaluate the efficacy and safety of placebo and 3 doses of topical prostaglandin E<sub>1</sub> cream (compositions based on Formulation I of Example 1 containing either 0.05%, 0.1% or 0.2% prostaglandin E<sub>1</sub>) in female subjects with FSAD in a controlled laboratory setting. Efficacy was assessed by vaginal photoplethysmography (Geer Gauge) during visual sexual stimulation (VSS), and by the use of quantitative patient questionnaires and diaries. Premenopausal subjects were enrolled under the assumption that their inherent magnitude of physiologic response is greater than that of postmenopausal subjects and thus will improve the likelihood of measuring pharmacologic effects. The study assessed the safety of 3 doses of topical prostaglandin E<sub>1</sub> cream in women with sexual dysfunction. The study also assessed the efficacy of 3 doses of topical prostaglandin E<sub>1</sub> cream in affecting vaginal blood flow and exudates and refinement of a quality of life instrument.

The study was a single center, single-blind, escalating dose, placebo-controlled pilot study to investigate the physiologic action, and the efficacy and safety of 3 doses of topical prostaglandin E<sub>1</sub> cream in women with FSAD. A total of 8 subjects were enrolled in this study. The study measured the dose-response characteristics of the safety and efficacy of the prostaglandin E<sub>1</sub> cream in terms of physiological response as well as the subjects' signs and symptoms and perceptions of the physiologic responses.

After signing the informed consent, at screening (Visit 1), the subjects underwent an adaptation session in the sexual response assessment laboratory. This session was intended to allow the patient to become familiar and comfortable with the procedures that will be carried out at subsequent visits. A complete medical history (including the Sexual Activity Questionnaire and the Brief Index of Sexual Function for Women (BISF-W; Taylor, J. F., et al., Self-report assessment of female sexual function: psychometric evaluation of the Brief Index of Sexual Functioning for Women. Arch Sexual Behavior: 23: 627-643, 1994.)) were collected. All medications taken by the patient were recorded. Baseline safety assessments included ECG, physical exam (including pelvic exam), clinical laboratory tests, and

vital signs. Subjects who met all inclusion and no exclusion criteria continued in the study.

The inclusion criteria for the study were that the female subjects aged 21 and provided written, informed consent; had a history of female sexual dysfunction (defined as impairment of the woman's ability to experience vaginal lubrication or engorgement sufficient for intercourse on at least 50% of attempts) of at least 6 months duration; were premenopausal; used adequate contraception (oral hormonal contraceptives, hormonal implants, or tubal ligation); had regular menses (cycles consistent in duration  $\pm$  2 days and between 25 and 31 days in length; and had a normal Pap smear within the past year. Any patient with an ASCUS Pap, except, "ASCUS, favor dysplasia" was admitted. Any Pap smear with inflammation or inflammatory changes in the absence of clinically significant vaginitis was admitted.

Subjects with any of the following conditions or meeting any of the following criteria were excluded from the study: female sexual dysfunction caused by untreated endocrine disease, e.g., hypopituitarism, hypothyroidism, diabetes mellitus; positive serum beta HCG or UPT result; a history of chronic or complicated urinary tract or vaginal infections within previous 12 months; a history of pelvic inflammatory disease within previous 12 months; history of dyspareunia not attributable to vaginal dryness within previous 12 months; significant (moderate to severe) vaginal atrophy; presence of moderate to severe vaginitis on pelvic examination; cervical dysplasia; significant cervicitis as manifested by mucopurulent discharge from the cervix; evidence of clinically significant hepatic disease as evidenced by SGOT or SGPT  $>3$  times the upper limit of normal within the last 6 months; evidence of clinically significant renal disease as evidenced by a serum creatinine  $> 2.5$  mg% within the last 6 months; a history of myocardial infarction within previous 12 months; symptomatic coronary artery disease, i.e., angina pectoris; symptomatic hypotension requiring medical consultation within the last 6 months; psychoses, uncontrolled bipolar disorder, uncontrolled depression; acute or chronic disease requiring frequent changes (changes within previous two months or anticipated in following two months) in medications or doses of chronic therapy; significant central nervous system diseases within the last 6 months i.e., stroke, spinal cord injury, etc.; participation in another study with an investigational drug or device during the 30 days prior to study entry, or planned during the study; any condition which would interfere with the patient's

ability to provide informed consent, to comply with study instructions, or which might confound the interpretation of the study results; or any condition which would endanger the participant if she participated in this trial.

At Visit 2, subjects received a single-blinded intravaginal dose of placebo.  
5 The patient underwent visual sexual stimulation in the sexual response assessment laboratory. The Geer Gauge (vaginal photoplethysmograph) was applied according to the manufacturer's instructions and vaginal photoplethysmography was recorded continuously from 15 minutes prior to dosing until the end of the visual sexual stimulation, approximately 60 minutes postdose. Safety was assessed from vital sign  
10 measurements and by monitoring the occurrence of adverse events. The patient's external genitalia, the vagina, and the cervix were inspected. Questionnaires were administered.

At Visit 3, subjects received a single-blinded dose of prostaglandin E<sub>1</sub> cream (containing 0.5 mg PGE<sub>1</sub>) applied to the labia, clitoris and the vulvar region of the  
15 vagina. All procedures from Visit 2 were repeated. If the patient tolerated this dose, she continued to the next visit. If the patient did not tolerate the dose in the clinic, she would be dismissed from the study but receive appropriate follow-up medical care.

At Visit 4, subjects returned to the clinic and received a single-blinded dose of prostaglandin E<sub>1</sub> cream (containing 1.0 mg PGE<sub>1</sub>) applied to the labia, clitoris and the  
20 vulvar region of the vagina. All procedures from Visit 2 were repeated. As before, if the patient tolerated this dose, she continued to the next visit. If the patient did not tolerate the dose in the clinic, she would be dismissed from the study but receive appropriate follow-up medical care.

At Visit 5, subjects returned to the clinic and received a single-blinded dose of prostaglandin E<sub>1</sub> cream (containing 2.0 mg PGE<sub>1</sub>) applied to the labia, clitoris and the  
25 vulvar region of the vagina. All procedures from Visit 2 were repeated. As before, if the patient tolerated this dose, she continued to the next visit. If the patient did not tolerate the dose in the clinic, she would be dismissed from the study but receive appropriate follow-up medical care.

30 The eight subjects had an average age of  $40.4 \pm 7.7$  years, an average weight of  $150.3 \pm 42.3$  pounds, and an average height of  $63.9 \pm 2.5$  inches. There were six Caucasian, one black and one Asian. All 8 subjects completed the study. The results are presented in Tables 4-14 and Figures 4-10.

Vaginal blood flow measurement, as maximum amplitude change in photoplethysmography measurements, did not show a statistically significant increase (Figure 4, Table 5). However, there were no decreases seen on treatment. The lack of statistical significance may be related to the high baseline levels.

5 TABLE 5: VAGINAL BLOOD FLOW MEASUREMENT  
MAXIMUM AMPLITUDE CHANGE IN PHOTOPLETHYSMOGRAPHY MEASUREMENT

Means for Placebo, Treatment, and Within-Subject Change from Placebo

	PLACEBO (Visit 2) N=6	PGE <sub>1</sub> 0.05% (Visit 3) N=8	PGE <sub>1</sub> 0.1% (Visit 4) N=8	PGE <sub>1</sub> 0.2% (Visit 5) N=7
Mean at Visit	10.00 ± 4.6*	8.50 ± 4.1	10.88 ± 4.6	5.57 ± 3.5
Change from Placebo		-1.50 ± 2.4** (p = 0.56)	0.50 ± 1.6 (p = 0.77)	-1.50 ± 2.4 (p = 0.56)

\*Standard deviation

\*\*Standard error

## Video Assessment Questionnaire

Please answer the following questions in regard to the erotic video that you have just seen. For Questions #2-4, circle the number that best describes your response on a scale from 0 to 10, with 0 meaning "not at all" and 10 meaning "very much."

- 5 1. Which of the videos did you find most arousing?  
A B C D
2. How much subjective arousal did you have during this video?  
Not at All Very Much  
0 1 2 3 4 5 6 7 8 9 10
- 10 3. How much lubrication (wetness) did you feel during this video?  
Not at All Very Much  
0 1 2 3 4 5 6 7 8 9 10
4. How much engorgement (fullness) did you feel during this video?  
Not at All Very Much  
15 0 1 2 3 4 5 6 7 8 9 10
5. How much tingling did you feel in your vagina during this video?  
Not at All Very Much  
0 1 2 3 4 5 6 7 8 9 10
- 20 6. How pleasurable were the feelings you had during the presentation of the video?  
Not at All Very Much  
0 1 2 3 4 5 6 7 8 9 10
7. Did you notice any other physical sensations during presentation of the video?  
No   
Yes  Please describe those sensations:
- 
- 25 8. Please rate the sensations described above in Question 7.  
I did not describe any sensations in Question 7.
- The sensations described in Question 7 were:  
Very Uncomfortable Very  
Pleasurable
- 30 0 1 2 3 4 5 6 7 8 9 10
9. How relaxed did you feel during the video presentation?  
Not at All Very Much  
0 1 2 3 4 5 6 7 8 9 10
10. Did you have any problems/difficulties in watching the videos? Please specify:
-

The visual inspection by the investigator revealed several significant increases in objective measures related to the treatment with the topical composition with some analyses (Table 6), in particular an increase in erythema at all dosage levels and increases in exudates at the two higher dosages at each visit. In other analyses (Tables 5 7, 8), the changes seen did not reach the  $p=0.05$  criterion of significance, with the exception of the exudate observations using 0.05% PGE<sub>1</sub> as reference.

TABLE 6: VISUAL INSPECTION

Pre-Treatment and Post-Treatment Means at Each Visit

	PLACEBO (Visit 2) N=8	PGE <sub>1</sub> 0.05% (Visit 3) N=8	PGE <sub>1</sub> 0.1% (Visit 4) N=8	PGE <sub>1</sub> 0.2% (Visit 5) N=7
<b>ERYTHEMA</b>				
Pre	1.00 ± 0**	1.00 ± 0	1.13 ± .35	1.00 ± 0
Post	1.25 ± .46	1.63 ± .52	2.13 ± .83	2.29 ± .76
Pre-Post*	p = 0.17	p = 0.01	p = 0.02	p = 0.004
<b>SWELLING</b>				
Pre	1.00 ± 0	1.00 ± 0	1.00 ± 0	1.00 ± 0
Post	1.13 ± .35	1.50 ± .53	1.25 ± .46	1.29 ± .49
Pre-Post	p = 0.35	p = 0.03	p = 0.17	p = 0.17
<b>EXUDATES</b>				
Pre	1.38 ± 0.52	1.88 ± .64	1.63 ± .52	1.50 ± .53
Post	2.00 ± 0.76	2.25 ± .46	2.63 ± .52	2.57 ± .98
Pre-Post	p = 0.05	p = 0.08	p = 0.001	p = 0.005

\*Change from pre-treatment to post-treatment, p-values based on paired t-tests.

\*\*Standard deviation

TABLE 7: VISUAL INSPECTION  
Comparisons of Mean Change From Pre- to Post-Treatment Within Subjects, Using Placebo as the Reference

	PLACEBO (Visit 2) N=8	PGE <sub>1</sub> 0.05% (Visit 3) N=8	PGE <sub>1</sub> 0.1% (Visit 4) N=8	PGE <sub>1</sub> 0.2% (Visit 5) N=7
ERYTHEMA*	0.25 ± .16***	0.63 ± .18	1.00 ± .33	1.29 ± .29
Change**		0.38 ± .26 (p = 0.20)	0.75 ± .37 (p = 0.08)	1.00 ± .31 (p = 0.018)
SWELLING	0.13 ± .13	0.50 ± .19	0.25 ± .16	0.29 ± .18
Change		0.38 ± .26 (p = 0.20)	0.13 ± .23 (0.60)	0.14 ± .26 (0.60)
EXUDATES	0.63 ± .26	0.38 ± .18	1.00 ± .19	1.14 ± .26
Change		-0.25 ± .31 (p = 0.45)	0.38 ± .26 (p = 0.20)	0.57 ± .43 (0.23)

\* Mean of the difference between pre-treatment and post-treatment, using paired t-tests.

\*\*Mean of the pre- to post-treatment change at each active drug visit, correcting for the change during treatment with placebo.

\*\*\*Standard error

5 TABLE 8: VISUAL INSPECTION  
Comparisons of Mean Change From Pre- to Post-Treatment Within Subjects, Using PGE<sub>1</sub> 0.05% as the Reference

	PGE <sub>1</sub> 0.05% (Visit 3) N=8	PGE <sub>1</sub> 0.1% (Visit 4) N=8	PGE <sub>1</sub> 0.2% (Visit 5) N=7
ERYTHEMA*	0.63 ± .18***	1.00 ± .33	1.29 ± .29
Change**		0.38 ± .32 (p = 0.28)	0.71 ± .29 (p = 0.047)
SWELLING	0.50 ± .19	0.25 ± .16	0.29 ± .18
Change		-0.25 ± .25 (p = 0.35)	-0.14 ± .14 (p = 0.36)
EXUDATES	0.38 ± .18	1.00 ± .19	1.14 ± .26
Change		0.63 ± .18 (p = 0.011)	0.71 ± .29 (p = 0.047)

\* Mean of the difference between pre-treatment and post-treatment, using paired t-tests.

\*\*Mean of the pre- to post-treatment change at each active drug visit, correcting for the change during treatment with PGE 0.05%.

\*\*\*Standard error

As with the vaginal blood flow measurements, the responses to the video assessment questionnaire showed high baseline responses (e.g., Table 9). The differences in responses between the lowest dose of PGE and the two higher doses are significant for question 3 (related to lubrication (Table 11) and engorgement at the highest dose level (Q4, Table 11). The graphs of the maximum responses show increased responses relative to baseline in all but one subject (Figures 6, 7). The

responses to questions 2 (subjective arousal) and 6 (pleasurable feelings) are near the  $p=0.05$  level if the 0.05% and 0.2% dose levels are compared (Table 11, compare to Figures 5, 9).

5

TABLE 9: VIDEO ASSESSMENT QUESTIONNAIRE RESULTS  
Visit Means

	Q2	Q3	Q4	Q5	Q6	Q9
Placebo (N = 8)	4.13 ± 1.6*	4.25 ± 2.3	2.13 ± 2.0	2.00 ± 2.2	4.25 ± 1.9	6.50 ± 2.7
PGE <sub>1</sub> 0.05% (N = 8)	2.25 ± 1.9	2.25 ± 1.7	1.38 ± 2.2	2.00 ± 2.3	2.75 ± 2.1	7.50 ± 1.5
PGE <sub>1</sub> 0.1% (N = 8)	4.13 ± 1.7	3.88 ± 1.9	2.88 ± 2.7	2.38 ± 2.3	3.88 ± 2.1	7.75 ± 1.8
PGE <sub>1</sub> 0.2% (N = 7)	4.57 ± 1.7	4.43 ± 2.5	3.71 ± 2.4	3.29 ± 1.8	5.00 ± 1.9	8.00 ± 2.2

\*Standard deviation

TABLE 10: VIDEO ASSESSMENT QUESTIONNAIRE RESULTS  
Means of Within-Subject Differences Between Placebo and Treatment

	Q2	Q3	Q4	Q5	Q6	Q9
Placebo vs. PGE <sub>1</sub> 0.05% (N = 8)	-1.88 ± 1.1* (p = 0.13)	-2.00 ± 1.2 (p = 0.13)	-0.75 ± 1.1 (p = 0.52)	0 ± 1.2 (p = 1.0)	-1.50 ± 1.3 (p = 0.27)	1.00 ± 0.62 (p = 0.15)
Placebo vs. PGE <sub>1</sub> 0.1% (N = 8)	0 ± 0.8 (p = 1.0)	-0.38 ± 1.0 (p = 0.73)	0.75 ± 0.5 (p = 0.20)	0.38 ± 0.8 (p = 0.64)	-0.38 ± 1.0 (p = 0.73)	1.25 ± 1.0 (p = 0.27)
Placebo vs. PGE <sub>1</sub> 0.2% (N = 7)	0.29 ± 0.4 (p = 0.46)	0 ± 0.5 (p = 1.0)	1.57 ± 1.0 (p = 0.17)	1.14 ± 0.5 (p = 0.07)	0.57 ± 0.5 (p = 0.32)	1.00 ± 1.0 (p = 0.38)

\*Standard error

10

TABLE 11: VIDEO ASSESSMENT QUESTIONNAIRE  
Means of Within-Subject Differences Between PGE<sub>1</sub> 0.05% and Treatment

	Q2	Q3	Q4	Q5	Q6	Q9
PGE <sub>1</sub> 0.05% vs. PGE <sub>1</sub> 0.1% (N = 8)	1.88 ± 0.8* (p = 0.054)	1.63 ± 0.7 (p = .04)	1.50 ± 1.0 (p = 0.18)	0.38 ± 0.7 (p = 0.62)	1.13 ± 0.8 (p = 0.22)	0.25 ± 0.6 (p = 0.68)
PGE <sub>1</sub> 0.05% vs. PGE <sub>1</sub> 0.2% (N = 7)	2.57 ± 1.1 (p = 0.06)	2.57 ± 1.0 (p = .04)	3.00 ± 1.2 (p = .04)	1.71 ± 1.0 (p = 0.14)	2.57 ± 1.2 (p = 0.08)	0.29 ± 0.6 (p = 0.65)

\*Standard error



TABLE 12: VITAL SIGNS  
Means at Admission and Discharge at Each Visit

	PLACEBO (Visit 2)	PGE <sub>1</sub> 0.05% (Visit 3)	PGE <sub>1</sub> 0.1% (Visit 4)	PGE <sub>1</sub> 0.2% (Visit 5)
	N=8	N=8	N=8	N=7
<u>Standing</u>				
Systolic				
Admission	108.8 ± 11.5*	107.5 ± 12.6	103.0 ± 5.0	106.5 ± 7.8
Discharge	110.0 ± 11.3	103.5 ± 11.5	103.5 ± 11.4	104.0 ± 10.6
Diastolic				
Admission	70.8 ± 5.8	70.8 ± 6.9	69.5 ± 7.2	71.0 ± 5.1
Discharge	72.3 ± 6.5	66.0 ± 7.3	68.8 ± 4.4	68.6 ± 4.1
Pulse				
Admission	74.0 ± 5.1	75.0 ± 7.3	70.5 ± 5.7	77.3 ± 8.6
Discharge	72.3 ± 4.3	69.8 ± 5.5	69.8 ± 4.2	73.4 ± 8.0
<u>Supine</u>				
Systolic				
Admission	110.5 ± 8.5	108.8 ± 6.0	110.5 ± 11.4	107.8 ± 10.8
Discharge	105.5 ± 6.7	102.5 ± 11.2	106.8 ± 9.9	108.6 ± 10.8
Diastolic				
Admission	69.3 ± 6.0	69.8 ± 3.1	67.8 ± 6.5	67.3 ± 6.9
Discharge	67.0 ± 4.5	65.3 ± 7.1	68.3 ± 5.2	68.9 ± 5.0
Pulse				
Admission	70.3 ± 7.5	71.5 ± 6.7	72.8 ± 4.7	75.5 ± 10.9
Discharge	68.8 ± 4.7	66.8 ± 5.4	68.3 ± 5.9	72.6 ± 7.4

\*\* Standard deviation

TABLE 13: VITAL SIGNS  
Means of Within-Subject Differences Between Pre-Treatment  
and Post-Treatment at Each Visit

	PLACEBO (Visit 2)	PGE <sub>1</sub> 0.05% (Visit 3)	PGE <sub>1</sub> 0.1% (Visit 4)	PGE <sub>1</sub> 0.2% (Visit 5)
	N=8	N=8	N=8	N=7
<u>Standing</u>				
Systolic	1.25 ± 3.0* (p = 0.68)	-4.00 ± 4.3 (p = 0.39)	0.50 ± 4.7 (p = 0.92)	-4.57 ± 4.2 (p = 0.32)
Diastolic	1.50 ± 1.8 (p = 0.42)	-4.75 ± 3.5 (p = 0.22)	-0.75 ± 3.0 (p = 0.81)	-3.71 ± 2.3 (p = 0.16)
Pulse	-1.75 ± 2.7 (p = 0.54)	-5.25 ± 2.7 (p = 0.09)	-0.75 ± 2.6 (p = 0.78)	-5.71 ± 2.7 (p = 0.08)
<u>Supine</u>				
Systolic	-5.00 ± 2.2 (p = 0.055)	-6.25 ± 2.9 (p = 0.07)	-3.75 ± 3.9 (p = 0.37)	-0.57 ± 4.0 (p = 0.89)
Diastolic	-2.25 ± 1.2 (p = 0.11)	-4.50 ± 2.5 (p = 0.11)	0.50 ± 1.9 (p = 0.80)	0.57 ± 3.5 (p = 0.88)
Pulse	-1.50 ± 3.7 (p = 0.70)	-4.75 ± 2.7 (p = 0.12)	-4.50 ± 2.2 (p = 0.08)	-4.86 ± 3.4 (p = 0.20)

\* Standard error

5

The minimal adverse events and the comparison of vital signs (Tables 13 and 14) indicated that the medication was well tolerated.

The raw data presented graphically in Figures 4-10 are presented in Table 14, below. Baseline response levels were recorded following the administration of placebo (visit 2). Maximum response levels were recorded following the administration of PGE<sub>1</sub> at 0.05%, 0.1% or 0.2% (visits 3-5). For all questions, the maximum responses showed increased means compared to baseline means (Table 14, bottom row).

The results of analysis of the raw data of Table 14 are presented in Table 15, below, where statistical significance at the  $p < 0.05$  level is designated by “\*” and at the  $p < 0.01$  level by “\*\*”. All statistical analysis of the difference between the mean baseline value and the mean maximum response were performed by paired t test with two-tailed P value as described in Bancroft, H.; 1957, Introduction to Biostatistics, 172-182.

The increases in maximum response means compared to baseline means were statistically significant for Questions No. 2, No. 4, No. 5, and No. 6 as shown in Table 15. These results indicate that PGE<sub>1</sub> treatment significantly increased female sexual response including subjective arousal, engorgement, vaginal tingling, and pleasurable feelings. The significant ( $p < 0.05$ ) response to question No. 2, “How much subjective arousal did you have during this video?”, shows that the method of the invention is effective in increasing subjective arousal in even this small sample of treated patients. The significant ( $p < 0.01$ ) response to question No. 4, “How much engorgement (fullness) did you feel during this video?”, shows that the method of the invention is effective in subjective awareness of increased engorgement in even this small sample of treated patients. The significant ( $p < 0.01$ ) response to question No. 5, “How much vaginal tingling did you feel during this video?”, shows that the method of the invention is effective in increasing subjective awareness of increase vaginal tingling in even this small sample of treated patients. The significant ( $p < 0.05$ ) response to question No. 6, “How pleasurable were the feelings you had during the presentation of this video?”, shows that the method of the invention is effective in increasing subjective pleasurable feelings in even this small sample of treated patients. These data thus show significant increases in subjective aspects that are specific to female sexual arousal disorder.

Patient No.	Question #2		Question #3		Question #4		Question #5		Question #6		Question #9	
	Base-line	Max. Resp.	Base-Line	Max. Resp.	Base-line	Max. Resp.	Base-line	Max. Resp.	Base-line	Max. Resp.	Base-line	Max. Resp.
1	3	7	3	7	2	6	1	5	3	7	3	7
2	3	4	3	4	1	1	0	0	4	4	9	10
3	6	7	8	6	2	7	4	5	7	8	9	10
4	7	6	5	6	7	8	6	6	6	6	2	8
5	3	5	2	5	2	5	3	5	4	6	6	8
6	4	5	7	8	1	6	0	3	6	6	9	7
7	4	5	4	5	1	3	2	4	2	4	7	10
8	3	5	2	3	1	2	0	4	2	6	7	7
Mean	4.1	5.5	4.3	5.5	2.1	4.8	2.0	4.0	4.3	5.9	6.5	8.4

Question #2	Question #3	Question #4	Question #5	Question #6	Question #9
P=0.028*	P=0.0835	P=0.0062**	P=0.0096**	P=0.0284*	P=0.0693
Statistical significance at the p<0.05 level is designated by "*" and at the p<0.01 level by "**".					

### EXAMPLE 6

#### Further Clinical Study in Women Suffering from Female Sexual Arousal Disorder

A randomized, placebo-controlled double blind crossover clinical efficacy and safety study was performed on eleven female patients 21-45 years of age. The patients who were selected for the study had been diagnosed with Female Sexual Arousal Disorder (FSAD) of more than 6 months duration.

The ingredients of the study medications are summarized in Table 16, below. Composition A was a topical prostaglandin E<sub>1</sub> composition comprising 0.2 weight percent prostaglandin E<sub>1</sub>. The placebo used, Composition B, was the composition lacking the active ingredient prostaglandin E<sub>1</sub>. A dose of 500 mg was packaged in a single-use applicator.

**Table 16: Composition of Study Medications**

<b>Ingredient</b>	<b>(%w/w)</b>
Ethyl Alcohol, dehydrated, USP	5.0
DDAIP HCl	1.6
Ethyl Laurate, FCC	3.0
Guar Gum, modified	2.5
Sodium Hydroxide Solution, 1M	Qs pH 5.5 (adjust pH of cream)
Sterile Water for Injection, USP, buffered to pH 5.5 with 0.1M Potassium Phosphate Monobasic, NF	qs 100

To 450 milligrams of the above cream base was added 50 milligrams of polyethylene glycol (PEG 400) that contained 1 mg prostaglandin E<sub>1</sub> to make Composition A. 50 milligrams of polyethylene glycol (PEG 400) alone was added to 450 milligrams of the above cream base to make Composition B, which contained no prostaglandin E<sub>1</sub>.

The patients were instructed regarding administration of the topical composition. The recommended dosage was stated as no more than one applicator per day. The patients were instructed to apply the cream as directed by the physician approximately 5-20 minutes prior to attempting intercourse. The patients were told to micturate and wash their hands and vulva by using soapsuds before applying the topical composition. After removing the cap from the tip of the applicator, the patient was to expel the cream onto a fingertip; and equally spread the cream over the clitoris. The patients were directed to spread the remaining cream into vaginal anterior G-spot at about 5 cm deep, 1/3 of the middle passage of the vagina. Each patient was asked to apply one dose of Composition A and one dose of Composition B during the course of the study. Either Composition A or Composition B was applied first, with a waiting period of at least five days before the application of the second study medication. The patients were asked to keep a diary, and note whether the use of the medication produced an improvement. Contact was lost with three of the eleven patients. No adverse effects were reported in the remaining population of eight patients. No patient diaries were returned by three patients; of these, two patients reported that neither composition was effective, while the other patient reported that Composition A but not B was effective. Overall, five of the eight patients reporting indicated that Composition A, containing 0.2 weight percent prostaglandin E<sub>1</sub>, was effective. Only one of the eight patients reported that the placebo was effective; she also reported that composition A was effective. Three patients reported that neither composition was effective. No patients reported that composition B was effective, but not composition A. While the size of the study population is small, the difference

between the reported efficacy of composition A (5/8 or 62.5%) and the reported efficacy of composition B (1/8 or 12.5%) is significantly different at the  $P \leq 0.01$  level. Thus, in this limited preliminary study, the composition containing the active ingredient, prostaglandin E<sub>1</sub>, was found to be more effective than the placebo when  
5 applied to the clitoris and intravaginally to the Grafenberg spot (G-spot) about 5-20 minutes prior to intercourse.

We claim:

1. A composition suitable for topical application comprising:  
an effective amount of a vasoactive prostaglandin;  
a penetration enhancer selected from the group consisting of an alkyl-(N-  
5 substituted amino) alkananoate, an alkyl-2-(N,N-disubstituted amino) alkananoate, an  
(N-substituted amino) alkanol alkananoate, an (N,N-disubstituted amino) alkanol  
alkanoate, pharmaceutically acceptable salts thereof and mixtures thereof;  
a polymer thickener selected from the group consisting of a polyacrylic acid  
polymer, a polysaccharide gum, a modified polysaccharide gum and mixtures thereof;  
10 a lipophilic component; and  
a buffer system wherein the pH of the composition is 3.0 to 7.4.
2. The composition of claim 1 wherein the vasoactive prostaglandin is  
chosen from the group consisting of prostaglandin E<sub>1</sub>, prostaglandin E<sub>1</sub> alkyl esters,  
pharmaceutically acceptable salts thereof and mixtures thereof.
- 15 3. The composition of claim 1 wherein the vasoactive agent is present in  
an amount of 0.07 weight percent to 1 weight percent, based on the total weight of the  
composition.
4. The composition of claim 1 wherein the vasoactive agent is  
prostaglandin E<sub>1</sub> present in an amount of 0.07 weight percent to 0.4 weight percent,  
20 based on the total weight of the composition.
5. The composition of claim 1 wherein the penetration enhancer is an  
alkyl-2-(N,N-disubstituted amino) alkananoate.
6. The composition of claim 1 wherein the penetration enhancer is  
selected from the group consisting of dodecyl N,N-dimethylamino isopropionate  
25 (DDAIP) and dodecyl N,N-dimethylamino isopropionate hydrochloride (DDAIP  
HCl).
7. The composition of claim 1 further comprising polyethylene glycol  
400.
8. The composition of claim 1 wherein the lipophilic component is  
30 chosen from the group consisting of the C<sub>1</sub> to C<sub>8</sub>, aliphatic alcohols, the C<sub>2</sub> to C<sub>30</sub>  
aliphatic esters and mixtures thereof.
9. The composition of claim 1 wherein the lipophilic component  
comprises ethanol.

10. The composition of claim 1 wherein the lipophilic component comprises propylene glycol.
11. The composition of claim 1 wherein the pH of the composition is about 3.5 to about 6.0.
- 5 12. The composition of claim 1 further comprising an emulsifier.
13. The composition of claim 1 further comprising a fragrance.
14. The composition of claim 1 further comprising a topical anesthetic.
15. A composition suitable for topical application comprising:  
0.001 weight percent to 1 weight percent of a vasoactive prostaglandin  
10 selected from the group consisting of PGE<sub>1</sub>, pharmaceutically acceptable salts thereof, lower alkyl esters thereof and mixtures thereof;  
0.01 weight percent to 5 weight percent modified polysaccharide gum;  
0.5 weight percent to 10 percent weight dodecyl N,N-dimethylamino isopropionate or pharmaceutically acceptable salts thereof;  
15 0.5 weight percent to 10 weight percent of a lower alcohol selected from the group consisting of ethanol, propanol, isopropanol and mixtures thereof;  
0.5 weight percent to 10 weight percent of an ester selected from the group consisting of ethyl laurate, isopropyl myristate, isopropyl laurate and mixtures thereof, based on the total weight of the composition, and  
20 a buffer system wherein the pH of the composition is 3.0 to 7.4.
16. The composition of claim 15 further comprising 1 weight percent to 25 weight percent polyethylene glycol 400, based on the total weight of the composition.
17. The composition of claim 15 further comprising 3 weight percent to 25 20 weight percent polyethylene glycol 400, based on the total weight of the composition.
18. The composition of claim 15 further comprising an emulsifier.
19. The composition of claim 15 wherein the pH of the composition is 3.5 to 6.0.
- 30 20. The use of a composition in accordance with any one of claims 1-19 for the preparation of a pharmaceutical composition for transdermal or transmucosal administration.
21. The use of a topical composition in accordance with any one of claims

1-19 for the preparation of a pharmaceutical composition for the treatment of female sexual arousal disorder.

22. The use of a topical composition in accordance with any one of claims 1-19 for the preparation of a pharmaceutical composition for increasing female sexual  
5 arousal.

23. A method for ameliorating female sexual arousal disorder, comprising the step of administering a composition in accordance with any one of claims 1-19 to the genitalia of a human female in need of treatment.

24. A method for ameliorating female sexual arousal disorder, comprising  
10 the step of administering a composition in accordance with any one of claims 1-19 to the clitoris and Grafenberg spot of a human female in need of treatment.

25. A method for ameliorating female sexual arousal disorder, comprising the step of administering a composition to the genitalia of a human female in need of treatment, the composition comprising:

15 an amount of a vasoactive prostaglandin effective to produce an increase in sexual arousal;

a penetration enhancer selected from the group consisting of an alkyl-(N-substituted amino) alkanoate, an alkyl-2-(N,N-disubstituted amino) alkanoate, an (N-substituted amino) alkanol alkanoate, an (N,N-disubstituted amino) alkanol  
20 alkanoate, pharmaceutically acceptable salts thereof and mixtures thereof;

a shear-thinning polymer;

a lipophilic component selected from the group consisting of an aliphatic C<sub>1</sub> to C<sub>8</sub> alcohol, an aliphatic C<sub>8</sub> to C<sub>30</sub> ester, and mixtures thereof; and an acidic buffer system.

25 26. The method in accordance with claim 25, wherein the vasoactive prostaglandin is selected from the group consisting of PGE<sub>1</sub>, PGA<sub>1</sub>, PGB<sub>1</sub>, PGF<sub>1α</sub>, 19-hydroxy-PGA<sub>1</sub>, 19-hydroxy-PGB<sub>1</sub>, PGE<sub>2</sub>, PGA<sub>2</sub>, PGB<sub>2</sub>, 19-hydroxy-PGA<sub>2</sub>, 19-hydroxy-PGB<sub>2</sub>, PGE<sub>3</sub>, PGF<sub>2α</sub>, and mixtures thereof.

30 27. The method in accordance with claim 25, wherein the shear-thinning polymer is selected from the group consisting of a polyacrylic acid polymer, a polysaccharide gum, a modified polysaccharide gum and mixtures thereof.



28. The method in accordance with claim 27, wherein the modified polysaccharide gum is a modified guar gum.
29. The method in accordance with claim 25, wherein the lipophilic component includes at least one aliphatic C<sub>8</sub> to C<sub>30</sub> ester.
- 5 30. The method in accordance with claim 25, wherein the lipophilic component includes at least one glyceryl ester selected from the group consisting monoglycerides, diglycerides, triglycerides, and mixtures thereof.
- 10 31. The method in accordance with claim 25, wherein the lipophilic component includes at least one glyceryl ester selected from the group consisting of glyceryl monooleate, triolein, trimyristin, tristearin, and mixtures thereof.
- 15 32. The method in accordance with claim 25, wherein the composition further comprises an emulsifier selected from the group consisting of sucrose esters, polyoxyethylene sorbitan esters, long chain alcohols, and glyceryl esters.
33. The method in accordance with claim 25, wherein the acidic buffer system provides a buffered pH value for said composition in the range of about 3 to about 6.5.
- 20 34. The method in accordance with claim 25 wherein the composition is applied to the clitoris and the Grafenberg spot.
35. The method in accordance with claim 25 wherein the composition is applied about five minutes to about twenty minutes before sexual intercourse.
36. The method in accordance with claim 25, wherein the composition further comprises a fragrance.
- 25 37. The method in accordance with claim 25, wherein the composition further comprises a preservative.
38. The method in accordance with claim 25, wherein the composition further comprises a topical anesthetic.
- 30 39. A method for ameliorating female sexual arousal disorder, comprising the step of applying a semi-solid prostaglandin composition to the genitalia of a human female in need of treatment, the composition comprising:  
a modified polysaccharide gum;

an effect amount of a vasoactive prostaglandin selected from the group consisting of PGE<sub>1</sub>, pharmaceutically acceptable salts thereof, lower alkyl esters thereof and mixtures thereof;

0.5 percent to 10 percent DDAIP or a pharmaceutically acceptable salt thereof, based on the total weight of the composition;

0.5 percent to 10 percent, based on the total weight of the composition, of a lower alcohol selected from the group consisting of ethanol, propanol, isopropanol and mixtures thereof;


0.5 percent to 10 percent of an ester selected from the group consisting of ethyl laurate, isopropyl myristate, isopropyl laurate and mixtures thereof, based on the total weight of the composition; and

an acidic buffer system.


40. The method in accordance with claim 39 wherein the composition is applied to the clitoris and the Grafenberg spot.

41. The method in accordance with claim 39 wherein the composition is applied about five minutes to about twenty minutes before sexual intercourse.


1/10

**Fig. 1**

2/10

**Fig. 2**

3/10

**Fig. 3**

4/10




Fig. 4




Fig. 5

6/10




Fig. 6



7/10




Fig. 7

8/10






Fig. 8

9/10



**Fig. 9**



**Fig. 10**