



PCT

特許協力条約に基づいて公開された国際出願

(51) 国際特許分類6 C12N 15/11, C12Q 1/68	A1	(11) 国際公開番号 WO00/09683 (43) 国際公開日 2000年2月24日(24.02.00)
(21) 国際出願番号 PCT/JP99/04341 (22) 国際出願日 1999年8月11日(11.08.99) (30) 優先権データ 特願平10/227177 1998年8月11日(11.08.98) JP (71) 出願人 (米国を除くすべての指定国について) アサヒビール株式会社(ASAHI BREWERIES, LTD.)(JP/JP) 〒104-0031 東京都中央区京橋3丁目7番1号 Tokyo, (JP) (72) 発明者 ; および (75) 発明者 / 出願人 (米国についてのみ) 本山靖朗(MOTOYAMA, Yasuo)(JP/JP) 尾形智夫(OGATA, Tomoo)(JP/JP) 坂井和久(SAKAI, Kazuhisa)(JP/JP) 〒302-0106 茨城県北相馬郡守谷町緑1-1-21 アサヒビール株式会社 酒類研究所内 Ibaraki, (JP) (74) 代理人 弁理士 舟橋榮子(FUNAHASHI, Eiko) 〒105-0003 東京都港区西新橋1丁目16番4号 ノアックスビル3階 Tokyo, (JP)		(81) 指定国 AU, CA, CN, CZ, JP, KR, US, 欧州特許 (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE) 添付公開書類 国際調査報告書
(54) Title: GENES FOR DETECTING BACTERIA AND DETECTION METHOD BY USING THE SAME (54) 発明の名称 細菌検出のための遺伝子及びそれを用いた検出法 (57) Abstract Genes for detecting respectively beer-clouding bacteria belonging to the genus <i>Pectinatus</i> , i.e., <i>P. frisingensis</i> and <i>P. cerevisiiphillus</i> and a method for detecting these bacteria by using these genes. Namely, the gene sequence of the spacer region constructed between 16SrRNA gene and 23SrRNA gene, which are specific to the bacterium belonging to the genus <i>Pectinatus</i> causing beer clouding; and a method for quickly detecting these bacteria at a high sensitivity by using this sequence.		

Genes for detecting bacteria and
a method for detecting bacteria by using the genes

Field of the Invention

The present invention relates to genes for detecting *Pectinatus*
5 *frisingensis* or *Pectinatus cerevisiiphilus* of the genus *Pectinatus*, which is
known as beer-spoilage bacteria, and a method for detecting the bacteria by
using the genes.

Description of the Prior Art

Bacteria of the genus *Pectinatus* have been known as beer-spoilage
10 bacteria. In the genus, two kinds of *Pectinatus frisingensis* and *Pectinatus*
cerevisiiphilus have been known. For detecting the bacteria of the genus
Pectinatus, the bacteria must be isolated after multiplication culture and
separation culture. It takes at least seven days. Then, isolated bacteria are
multiplied and tested by many qualitative tests such as morphological
15 observation, gram stainability, a catalase test, utilization of various carbon
sources and the like to identify the bacteria.

These tests are very troublesome, and it takes much time and it costs
much. In addition to these common identification tests, there is a method that
DNA is extracted from isolated bacteria, fixed on a membrane, and conducted a
20 hybridization test by using standard bacteria DNA as a probe to identify the
class. However, it takes some days, and it is difficult to obtain necessary
detective sensitivity and selectivity.

Lately, a method for detection of bacteria of the genus *Pectinatus* is
disclosed by using a monoclonal antibody that specifically reacts with *Pectinatus*
25 *cerevisiiphilus* (ASBC Journal: 51(4)158-163, 1993). However, the method is
insufficient to the detective sensitivity. The method has a problem that
Pectinatus frisingensis can not be detected.

The other detection method has been reported. It can detect *Pectinatus*
frisingensis and *Pectinatus cerevisiiphilus* by a Ribotyping method that

polymorphism of a ribosomal RNA gene is detected (J. Am. Soc. Chem.: 56 (1) 19-23, 1998). However, since the method needs operation for isolating the bacteria, it has problems of detective sensitivity and speed.

Considering these problems, further quick detection methods have been
5 studied. WO97/20071 discloses a method for detecting *Pectinatus* comprising extracting DNA of the test microorganism, and using a PCR method that a complementary oligonucleotide of the DNA functionates as a primer. However, the base sequences of 16S rRNA gene used in the technique are sometimes similar to those of microorganisms of the other genera, so that there are
10 problems that the other microorganisms are detected in addition to particular microorganisms to be detected.

The gene in a spacer between a 16S rRNA gene and a 23S rRNA gene has a specific gene sequence. Though methods for detecting microorganisms using the gene sequence are disclosed in Japanese Jozo Ronbunshu 50, 22-31
15 (1995), APPL. ENVIRON. MICROBIOL. VOL.62, NO.5, 1683-1688(1996), FEMS MICROBIOL LETT. VOL. 84, NO.3, 307-312(1991), Japanese Patent Kokai Publication No. 6-98800 and the like, gene sequences of the spacers of the genus *Pectinatus* have not been found.

Summary of the Invention

20 The present invention aims to provide gene sequences of a spacer region that is constituted between a 16S rRNA gene and a 23S rRNA gene specific for the genus *Pectinatus* relating to beer-spoilage, and to provide a method for sensitively and quickly detecting the genus by using the sequences.

(1) The first invention is a gene sequence of a spacer region between a gene
25 coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus frisingensis* containing a part of the base sequence or the whole base sequence represented by SEQ ID NO: 1.

(2) The second invention is a gene sequence of a spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus frisingensis*

containing a part of the base sequence or the whole base sequence represented by SEQ ID NO: 2.

(3) The third invention is a gene sequence of a spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus cerevisiiphilus*
 5 containing a part of the base sequence or the whole base sequence represented by SEQ ID NO: 3.

(4) The fourth invention is a gene sequence of a spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus cerevisiiphilus*
 containing a part of the base sequence or the whole base sequence represented
 10 by SEQ ID NO: 4.

(5) The fifth invention is an oligonucleotide characterized in that the gene sequence of a spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus frisingensis* has at least one of the following sequence group or the corresponding complementary sequence:

15 5'-CCATCCTCTTGAAAATCTC-3' ①

5'-TCTCRTCTCACAAGTTTGGC-3' ②.

(6) The sixth invention is an oligonucleotide characterized in that the gene sequence of a spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus cerevisiiphilus* has at least one of the following sequence
 20 group or the corresponding complementary sequence:

5'-CACTCTTACAAGTATCTAC-3' ③

5'-CCACAATATTTCCGACCAGC-3' ④

5'-AGTCTTCTCTACTGCCATGC-3' ⑤.

(7) The seventh invention is a method for detecting *Pectinatus frisingensis*,
 25 wherein the oligonucleotide made from the gene sequence described in (1) or (2) uses as a primer for synthesis of nucleic acids, and the nucleic acid is treated by gene amplification to detect the bacteria.

(8) The eighth invention is a method for detecting *Pectinatus cerevisiiphilus*, wherein the oligonucleotide made from the gene sequence described in (3) or (4)

uses as a primer for synthesis of nucleic acids, and the nucleic acid is treated by gene amplification to detect the bacteria.

(9) The ninth invention is a method for detecting *Pectinatus frisingensis*, wherein the oligonucleotide made from the gene sequence described in (1) or (2),
 5 or the oligonucleotide made from the gene sequence described in (5), and a nucleotide sequence coding 16S rRNA gene of *Pectinatus frisingensis* use as primers for synthesis of nucleic acids, and the nucleic acid is treated by gene amplification to detect the bacteria.

(10) The tenth invention is a method for detecting *Pectinatus cerevisiophilus*,
 10 wherein the oligonucleotide made from the gene sequence described in (3) or (4) or the oligonucleotide made from the gene sequence described in (6), and a nucleotide sequence coding 16S rRNA gene of *Pectinatus cerevisiophilus* use as primers for synthesis of nucleic acids, and the nucleic acid is treated by gene amplification to detect the bacteria.

15 (11) The eleventh invention is a method as in (9), wherein the nucleotide sequence coding the 16S rRNA gene of *Pectinatus frisingensis* has the following sequence:

5'-CGTATCCAGAGATGGATATT-3' ⑥

(12) The twelfth invention is a method as in (10), wherein the nucleotide
 20 sequence coding the 16S rRNA gene of *Pectinatus cerevisiophilus* has the following sequence:

5'-CGTATGCAGAGATGCATATT-3' ⑦

Brief Description of Drawings

- 25 Figure 1. It shows Electrophoretogram in Example 3.
 Figure 2. It shows Electrophoretogram in Example 5.

Detailed Description of the Invention

Since the technique of gene amplification is well known, it is conducted

under the polymerase chain reaction method which has been developed by Saiki et al. (abbreviated as PCR method hereinafter; Science 230, 1350, 1985).

This method is conducted by amplification reaction of particular gene sequences. Since the method shows quick reaction, high sensitivity and
5 specificity and convenience, applications has been tried to quickly judge viruses in medical fields or quickly detect noxious bacteria in food fields. By the PCR method, even if only a few nucleotide sequences are present in test samples, the target nucleotide sequence between two primers is amplified several hundred times, and the copies are produced in large quantities to be detectable. For
10 conducting the PCR method, the nucleic acid ingredient should be liberated from the bacteria in the test samples. However, in the PCR method, when several or more molecules exist in the target sequence, the amplification reaction proceeds. Accordingly, samples of the PCR method can be provided by a simple pretreatment of the bacteria with a lytic enzyme or a surfactant. For this
15 reason, the method for detecting bacteria has merits higher than conventional methods.

The present invention provides gene sequences of a spacer region between a gene coding 16SrRNA and a gene coding 23SrRNA in each *Pectinatus frisingensis* or *Pectinatus cerevisiiphilus*. By using a nucleotide sequence
20 coding a 16SrRNA gene or oligonucleotide which is selected from the sequence as a primer for nucleic acid synthesis in the PCR method, and by gene amplification treatment, the present inventors have developed a quick and high sensitive method for judging the existence of *Pectinatus frisingensis* or *Pectinatus cerevisiiphilus* in samples.

25 The test samples may be beer or semi-products of beer, or a sample extracted from sewage and the like. The oligonucleotide for a primer may be a chemical synthetic or natural product.

Description of the Preferred Embodiments

As shown hereinafter, in the method of the present invention,

Pectinatus frisingensis or *Pectinatus cerevisiiphilus* is detected by the PCR method. The base sequences used in the PCR method are, not by way of limitation, for example, above-mentioned (5), (6), (11) and (12). The primer length used in the PCR method is, not by way of limitation, 19-20 base length in
 5 above-mentioned (5), (6), (11) and (12), preferably, 10-50 base length.

When *Pectinatus frisingensis* is detected by the PCR method, the existence of the bacteria is judged by that the DNA fragments amplified in case of the combination of ① and ⑥ as the primer are about 700 base pairs and about 900 base pairs, and the DNA fragments amplified in case of the
 10 combination of ② and ⑥ as the primer are about 700 base pairs and about 900 base pairs. When these bands are detected by electrophoresis, it is judged that *Pectinatus frisingensis* is present. Since the combination of the primers, in any cases, is specific for *Pectinatus frisingensis* bacteria, the genus can be detected. By parallel using two of the combination, further precise determination becomes
 15 possible. By changing the base sequences of the primers used in the PCR method, the length of the nucleotide sequences amplified can be changed.

On the other hand, when *Pectinatus cerevisiiphilus* is detected by the PCR method, the existence of the bacteria is judged by that the DNA fragments amplified in case of the combination of ③ and ⑦ are about 600 base pairs, the
 20 DNA fragments amplified in case of the combination of ④ and ⑦ are about 650 base pairs, and the DNA fragments amplified in case of the combination of ⑤ and ⑦ are about 700 base pairs. When these bands are detected by electrophoresis, it is judged that *Pectinatus cerevisiiphilus* is present. Since the combination of the primers, in any cases, is specific for *Pectinatus cerevisiiphilus*
 25 bacteria, the genus can be detected. By parallel using two or more of the combination, further precise determination becomes possible. By changing the base sequences of the primers used in the PCR method, the length of the nucleotide sequences amplified can be changed.

The temperature conditions of one cycle in the PCR method are 90-98°C

in a thermal denaturation reaction in which double-stranded DNA is changed to single-stranded DNA, 37-65°C in an annealing reaction in which DNA is hybridized into primer template DNA, and 50-75°C in a chain elongation reaction in which DNA polymerase is reacted. The target sequences can be amplified by several ten cycles. After PCR reaction, the reactant is separated by electrophoresis, and the nucleic acid is stained with ethidium bromide or the like. When the base length of the amplified nucleotide sequence is equal to the base length of the above target sequence, it can be judged that the bacteria to be detected are in the test sample. To detect the amplified nucleotide sequence, chromatography is usable.

The sequences of the present invention are described in the following:

- SEQ ID NO: 1 The sequence length is 624, the sequence type is nucleic acid, the strandness is double, the topology is linear, the molecule type is genomic DNA, and the original source is *Pectinatus frisingensis* DSM6306.
- SEQ ID NO: 2 The sequence length is 442, the sequence type is nucleic acid, the strandness is double, the topology is linear, the molecule type is genomic DNA, and the original source is *Pectinatus frisingensis* DSM6306.
- SEQ ID NO: 3 The sequence length is 724, the sequence type is nucleic acid, the strandness is double, the topology is linear, the molecule type is genomic DNA, and the original source is *Pectinatus cerevisiiphilus* DSM20467.
- SEQ ID NO: 4 The sequence length is 399, the sequence type is nucleic acid, the strandness is double, the topology is linear, the molecule type is genomic DNA, and the original source is *Pectinatus cerevisiiphilus* DSM20467.
- SEQ ID NO: 5 The sequence length is 19, the sequence type is nucleic acid, the strandness is single, the topology is linear, the molecule type is genomic DNA, and the original source is *Pectinatus frisingensis* DSM6306.
- SEQ ID NO: 6 The sequence length is 20, the sequence type is nucleic acid, the strandness is single, the topology is linear, the molecule type is genomic DNA, and the original source is *Pectinatus frisingensis* DSM6306.

SEQ ID NO: 7 The sequence length is 19, the sequence type is nucleic acid, the strandness is single, the topology is linear, the molecule type is genomic DNA, and the original source is *Pectinatus cerevisiiphilus* DSM20467.

5 SEQ ID NO: 8 The sequence length is 20, the sequence type is nucleic acid, the strandness is single, the topology is linear, the molecule type is genomic DNA, and the original source is *Pectinatus cerevisiiphilus* DSM20467.

SEQ ID NO: 9 The sequence length is 20, the sequence type is nucleic acid, the strandness is single, the topology is linear, the molecule type is genomic DNA, and the original source is *Pectinatus cerevisiiphilus* DSM20467.

10 SEQ ID NO: 10 The sequence length is 20, the sequence type is nucleic acid, the strandness is single, the topology is linear, the molecule type is genomic DNA, and the original source is *Pectinatus frisingensis* DSM6306.

SEQ ID NO: 11 The sequence length is 20, the sequence type is nucleic acid, the strandness is single, the topology is linear, the molecule type is genomic DNA,
15 and the original source is *Pectinatus cerevisiiphilus* DSM20467.

The present invention is described by working examples in the following. The present invention is not limited by these examples.

Example 1

Preparation of test samples

20 *Pectinatus frisingensis* DSM6306 and *Pectinatus cerevisiiphilus* DSM20467 were used as bacterial strains belonging to *Pectinatus*. To confirm the specificity of *Pectinatus frisingensis* and *Pectinatus cerevisiiphilus* primers shown in SEQ ID NO:5, 6, 7, 8, 9, 10 and 11 in the present invention, the other bacteria shown in Table 1 were used. These bacteria were cultivated on
25 suitable culture mediums, and the strains were collected by centrifugation. The DNA from the strains were extracted in accordance with the description of SHIN-SEIKAGAKU-JIKKEN-KOZA 2, Nucleic acid I, Separation and Purification, p.p. 20-21 (edited by Japan Biochemical Learned Society, Tokyo-Kagaku-Dojin), and a DNA solution was obtained.

[Table 1]

Bacteria No.	Bacteria type	Strain name	Remarks
1	<i>Pectinatus frisingensis</i>	DSM6306	type strain
2	<i>Pectinatus cerevisiiphilus</i>	DSM20467	type strain
3	<i>Selenomonas lacticifex</i>	DSM20757	type strain
4	<i>Zymophilus raffinovorans</i>	DSM20765	type strain
5	<i>Zymophilus paucivorans</i>	DSM20756	type strain
6	<i>Escherichia coli</i>	IF03301	K-12
7	<i>Megasphaera cerevisiae</i>	DSM20462	type strain
8	<i>Lactobacillus acidophilus</i>	IF013951	type strain
9	<i>Lactobacillus plantarum</i>	JCM1149	type strain
1 0	<i>Lactobacillus brevis</i>	JCM1059	type strain
1 1	<i>Lactococcus lactis</i>	JCM5805	type strain
1 2	<i>Leuconostoc mesenteroides</i>	JCM6124	type strain
1 3	<i>Pediococcus damnosus</i>	JCM5886	type strain

Example 2

Cloning of spacer regions between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus frisingensis*, and determination of the base sequences

- 5 (1) Selection and synthesis of oligonucleotide primers for amplification of 16S/23S rRNA spacer region by the PCR method

Since the base sequences of the 16S ribosomal RNA gene of *Pectinatus frisingensis* were apparent (International Journal of Systematic Bacteriology, Vol. 40, p.p. 19-27 (1990)), the primers were selected on the basis of the 557-576th
10 base sequences.

Since the base sequences of the 23S ribosomal RNA gene of *Pectinatus frisingensis* were apparent (Systematic Applied Microbiology, Vol. 15, p.p. 487-501 (1990), EMBL Accession Number X48423), the primers were selected on the basis of the 1-20th base sequences to obtain corresponding comprehensive
15 sequences. The synthesis was entrusted to Sawady Technology Co., Ltd.

- (2) Amplification of 16S/23S rRNA spacer regions by the PCR method

The *Pectinatus frisingensis* DNA solution 0.1 μ g, which was prepared in Example 1, was placed in a 0.2 ml tube (manufactured by Perkin-Elmer), 5 μ l of 10X buffer in a rTaq DNA Polymerase Kit (Toyobo Co., Ltd.), 3 μ l of 25mM
20 $MgCl_2$, 5 μ l of a 2mM dNTP mixture solution (dATP, dGTP, dCTP and dTTP), 0.5 μ l of 5 units / μ l of rTaq polymerase, and each 0.5 μ l of 100 mM primers prepared in Example 2-(1) were added to the solution, and then sterilized distilled water was added to obtain final volume of 50 μ l. The tube was set on a thermal cycler of an automatic gene amplification device (Perkin Elmer) and
25 the amplification method was conducted. The reaction was repeated by 30 cycles, and one cycle had the following conditions:

Denaturation at 94°C for 2.5 minutes; Denaturation at 94°C for 30 seconds; Annealing of primers at 55°C for 30 seconds; and synthetic reaction at 72°C for 30seconds. After the reaction, using 5 μ l of the solution,

electrophoresis was conducted by agarose gel. DNA was dyed with ethidium bromide, and amplified DNA was observed. The result shows that about 1600 bp (abbreviated as "long" hereinafter) DNA and about 1400 bp (abbreviated as "short" hereinafter) DNA were amplified.

5 (3) Cloning and sequencing of the spacer region "long"

Using a high pure PCR product purification kit (Baringer Mannheim), unreactive dNTPs was removed from the solution after the PCR reaction. To the resulting amplified DNA 100 ng, 2 μ l of plasmid pCR 2.1 contained in a TA cloning kit (INVITROGEN), 1 μ l of ligase and 1 μ l of buffer were added, and
10 then sterilized water was added to obtain the total volume of 10 μ l. After the solution was reacted at 14°C for 4 hours, 2 μ l of the solution and 2 μ l of 0.5 M β -mercaptoethanol were added to *Escherichia coli* INV α 'F competent cells, and placed in ice for 30 minutes. Then, the solution was heated at 42°C for 30 seconds, and plasmid transformation to the bacteria was conducted. To the
15 transformed bacteria, 250 μ l of a SOC culture (2.0% Tryptone, 0.5% yeast extract, 10.0 mM NaCl, 2.5 mM KCl, 10.0 mM MgCl₂·6H₂O, and 20.0 mM glucose) was added, and the mixture was shaken at 37°C for 60 minutes, then transferred to a LB plate culture medium containing 50 μ g / ml of ampicillin and 40 μ g / ml X-Gal, and cultured at 37°C overnight. The expressed white
20 colony was transferred to 3 ml of a LB liquid culture medium containing 50 μ g / ml of ampicillin, and cultured at 37°C overnight.

After the cultivation, plasmids were extracted from the bacteria with a plasmid mini kit (QIAGEN). A part of the resulting plasmids was taken out and reacted with a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) at 37
25 °C for 60 minutes, and separated by agarose electrophoresis. The DNA was dyed with ethidium bromide, and insertion of "long" was confirmed. 500 ng of the residual plasmid was reacted with restriction enzyme *Sma*I (manufactured by TOYOBO Co., Ltd.) at 30°C for 60 minutes. To the reactant, 2 μ l of 3 M sodium acetate and 500 μ l of 100% ethanol were added, and the mixture was

placed in ice for 15 minutes and centrifuged at 15000 rpm for 15 minutes, and the supernatant was removed. To the precipitate, 500 μ l of 70% ethanol was added, the mixture was centrifuged at 15000 rpm for 15 minutes, and the supernatant was removed, and dried for 10 minutes under reduced pressure.

5 Sterilized water was added to dissolve the precipitate, and the mixture was reacted with restriction enzyme *Xba*I (Baringer Mannheim) at 37°C for 60 minutes. To the reactant, equivalent phenol / chloroform (equivalent mixture liquid) was added and gently mixed, the mixture was centrifuged at 15000 rpm for 15 minutes, and the water layer (upper layer) was recovered.

10 To the recovery liquid, equivalent water-saturated ether was added and gently mixed, and the mixture was centrifuged at 15000 rpm for 15 minutes to remove the ether layer (upper layer). To the remaining water layer, 2 μ l of 3M sodium acetate and 500 μ l of 100% ethanol were added, and the mixture was placed in ice for 15 minutes and centrifuged at 15000rpm for 15 minutes to
 15 remove the supernatant. To the precipitate, 500 μ l of 70% ethanol was added, and the mixture was centrifuged at 15000 rpm for 15 minutes to remove the supernatant, and the residue was dried under reduced pressure for 10 minutes, and 20 μ l of sterilized distillation water was added. To 5 μ l of the solution, 1 μ l of 10X buffer contained in a blunting kit (Takara Shuzo Co., Ltd.) and 3 μ l of
 20 sterilized distillation water were added, and the mixture was maintained at 70 °C for 5 minutes, 1 μ l of T4 DNA polymerase was added, and the mixture was maintained at 37°C for 5 minutes to obtain blunt ends. After T4 DNA polymerase was inactivated by stirring, 40 μ l of ligation solution A and 10 μ l of ligation solution B were added, and the mixture was maintained at 16°C for
 25 30 minutes to conduct internal ligation.

The reactant 2 μ l and 2 μ l of 0.5M β -mercaptoethanol were added to a *Escherichia coli* INV α 'F competent cell, and the mixture was placed in ice for 30 minutes and heated at 42°C for 30 seconds, and the plasmid was transformed to the *Escherichia coli*. To the transformed *Escherichia coli*, a SOC culture

medium (2.0% Tryptone, 0.5% Yeast extract, 10.0 mM NaCl, 2.5 mM KCl, 10.0 mM MgCl₂·6H₂O, 20.0 mM glucose) 250 μ l was added, and the mixture was shaken at 37°C for 60 minutes and spread on a LB plate culture medium containing 50 μ g / ml ampicillin to culture at 37°C overnight. Appeared white colonies were inoculated into 3 ml of a LB liquid culture medium containing 50 μ g / ml of ampicillin and cultured at 37°C overnight. After the culture, the plasmid was extracted from the *Escherichia coli* with a plasmid mini kit (QIAGEN Company).

Using such obtained plasmid as a template, a sequence reaction was conducted. As the sequencing primers, an IRD41 Infrared Dye Labeled M13 Forward primer and an IRD41 Infrared Dye Labeled M13 Reverse primer (manufactured by Nisshinbo, sold by Aroka Co., Ltd.) were used. As the reaction liquid, SequiTherm (trademark) Long-Read (trademark) Cycle Sequencing Kit-LC (manufactured by EPICENTRE TECHNOLOGIES) was used. 4000L Long ReadIR (trademark) DNA Sequencing System (manufactured by LI-COR) was used for the determination of the base sequences.

The gene sequence of spacer region "long" between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus frisingensis* DSM6306 bacteria is shown in SEQ ID NO: 1.

(4) Cloning and sequencing of spacer region "short"

Using a high pure PCR product purification kit (Baringer Mannheim), unreactive dNTPs was removed from the solution after the PCR reaction in Example 2-(2). To the resulting amplified DNA 100 ng, 2 μ l of plasmid pCR 2.1 contained in a TA cloning kit (INVITROGEN), 1 μ l of ligase and 1 μ l of buffer were added, and then sterilized water was added to obtain the total volume of 10 μ l. After the solution was reacted at 14°C for 4 hours, 2 μ l of the solution and 2 μ l of 0.5 M β -mercaptoethanol were added to *Escherichia coli* INV α 'F competent cells, and placed in ice for 30 minutes. Then, the

solution was heated at 42°C for 30 seconds, and plasmid transformation to the bacteria was conducted. To the transformed bacteria, 250 μ l of a SOC culture (2.0% Tryptone, 0.5% yeast extract, 10.0 mM NaCl, 2.5 mM KCl, 10.0 mM $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, and 20.0 mM glucose) was added, and the mixture was shaken at
5 37°C for 60 minutes, then transferred to a LB plate culture medium containing 50 μ g / ml of ampicillin and 40 μ g / ml X-Gal, and cultured at 37°C overnight. The appeared white colony was transferred to 3 ml of a LB liquid culture medium containing 50 μ g / ml of ampicillin, and cultured at 37°C overnight. After the cultivation, plasmid was extracted from the bacteria with a plasmid
10 mini kit (QIAGEN).

A part of the resulting plasmid was taken out and reacted with a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) at 37°C for 60 minutes, and separated by agarose electrophoresis. The DNA was dyed with ethidium bromide, and insertion of "short" was confirmed. 500 ng of the
15 residual plasmid was reacted with restriction enzyme *Sma*I (manufactured by TOYOBO Co., Ltd.) at 30°C for 60 minutes. To the reactant, 2 μ l of 3 M sodium acetate and 500 μ l of 100% ethanol were added, and the mixture was placed in ice for 15 minutes and centrifuged at 15000 rpm for 15 minutes, and the supernatant was removed. To the precipitate, 500 μ l of 70% ethanol was
20 added, the mixture was centrifuged at 15000 rpm for 15 minutes, and the supernatant was removed, and dried for 10 minutes under reduced pressure. Sterilized water was added to dissolve the precipitate, and the mixture was reacted with restriction enzyme *Xba*I (Baringer Mannheim) at 37°C for 60 minutes. To the reactant, equivalent phenol / chloroform (equivalent mixture
25 liquid) was added and gently mixed, the mixture was centrifuged at 15000 rpm for 15 minutes, and the water layer (upper layer) was recovered. To the recovery liquid, equivalent water-saturated ether was added and gently mixed, and the mixture was centrifuged at 15000 rpm for 15 minutes to remove the ether layer (upper layer).

To the remaining water layer, 2 μ l of 3M sodium acetate and 500 μ l of 100% ethanol were added, and the mixture was placed in ice for 15 minutes and centrifuged at 15000rpm for 15 minutes to remove the supernatant. To the precipitate, 500 μ l of 70% ethanol was added, and the mixture was centrifuged
 5 at 15000 rpm for 15 minutes to remove the supernatant, and the residue was dried under reduced pressure for 10 minutes, and 20 μ l of sterilized distilled water was added. To 5 μ l of the solution, 1 μ l of 10X buffer contained in a blunting kit (Takara Shuzo Co., Ltd.) and 3 μ l of sterilized distilled water were added, and the mixture was maintained at 70°C for 5 minutes, 1 μ l of T4 DNA
 10 polymerase was added, and the mixture was maintained at 37°C for 5 minutes to obtain blunt ends. After T4 DNA polymerase was inactivated by stirring, 40 μ l of ligation solution A and 10 μ l of ligation solution B were added, and the mixture was maintained at 16°C for 30 minutes to conduct internal ligation. 2
 15 μ l of the reactant and 2 μ l of 0.5M β -mercaptoethanol were added to a *Escherichia coli* INV α 'F competent cell, and the mixture was placed in ice for 30 minutes and heated at 42°C for 30 seconds, and the plasmid was transformed to the *Escherichia coli*.

To the transformed *Escherichia coli*, 250 μ l of SOC culture medium (2.0% Tryptone, 0.5% Yeast extract, 10.0 mM NaCl, 2.5 mM KCl, 10.0 mM
 20 MgCl₂·6H₂O, 20.0 mM glucose) was added, and the mixture was shaken at 37°C for 60 minutes and spread on a LB plate culture medium containing 50 μ g / ml ampicillin to culture at 37°C overnight. Appeared white colonies were inoculated into 3 ml of a LB liquid culture medium containing 50 μ g / ml of ampicillin and cultured at 37°C overnight. After the culture, the Plasmid was
 25 extracted from the *Escherichia coli* with a plasmid mini kit (QIAGEN Company).

Using such obtained plasmid as a template, a sequence reaction was conducted. As the sequencing primers, an IRD41 Infrared Dye Labeled M13 Forward primer and an IRD41 Infrared Dye Labeled M13 Reverse primer (manufactured by Nisshinbo, sold by Arok co., Ltd.) were used. As the reaction

liquid, SequiTherma (trademark) Long-Read (trademark) Cycle Sequencing Kit-LC (manufactured by EPICENTRE TECHNOLOGIES) was used. 4000L Long ReadIR (trademark) DNA Sequencing System (manufactured by LI-COR) was used for the determination of the base sequences.

- 5 The gene sequence of spacer region "short" between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus frisingensis* is shown in SEQ ID NO: 2.

Example 3

Detection of *Pectinatus frisingensis* by the PCR method

- 10 (1) Selection and synthesis of a primer for *Pectinatus frisingensis*

The sequences specific for *Pectinatus frisingensis* by using DNASIS (tradename of Hitachi Soft Engineering Ltd., Co.) on the basis of SEQ ID NO: 1 and SEQ ID NO: 2 were analyzed. The result selected a sequence of 377th to 395th on the gene sequence of the spacer region between the gene coding 16S rRNA and the
15 gene coding 23S rRNA of *Pectinatus frisingensis* of SEQ ID NO: 1, and a sequence of 195th to 213th on the gene sequence of the spacer region between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus frisingensis* of SEQ ID NO: 2. (SEQ ID NO: 5.)

In addition, the similar analysis selected a sequence of 361st to 380th on
20 the gene sequence of the spacer region between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus frisingensis* of SEQ ID NO: 1, and a sequence of 179th to 198th on the gene sequence of the spacer region between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus frisingensis* of SEQ ID NO: 2. (SEQ ID NO: 6.)

- 25 Further, specific primer showing in SEQ ID NO: 10 was selected by a gene sequence coding 16S rRNA of *Pectinatus frisingensis*. The oligonucleotides were chemically synthesized by the same method as in Example 2-(1).

(2) Detection and identification of *Pectinatus frisingensis* by the primers

having the sequences of SEQ ID NO: 6 and SEQ ID NO: 10.

The DNA solutions of bacteria prepared in Example 1 were treated with the primers synthesized in Example 3 (SEQ ID NO:6 and SEQ ID NO:10) by PCR. The temperature conditions of the PCR were as follows:

- 5 Thermal denaturation; 94°C, 30 seconds
- Annealing; 55°C, 30 seconds
- Chain elongation reaction; 72°C, 30 seconds

One cycle of the conditions was repeated 35 times. After the PCR, the reactant was electrophoresed with agarose gel at constant 100 V for 30 minutes. A pHY marker was also electrophoresed at the same time as a molecular weight marker. After the electrophoresis, the agarose gel was stained in about 0.5 μ g/ml of an ethidium bromide solution for 20 minutes, and ultraviolet was applied to observe the gel and take a photograph of the gel. By the observation or the photography of the gel, the base length of the amplified products was determined from the relative migration distance of the molecular marker.

As shown in Fig. 1, bands of about 700 bps and about 900 bps were detected only in case of *Pectinatus frisingensis*.

From the results, when the oligonucleotides of SEQ ID NO: 6 and SEQ ID NO: 10 were used as PCR primers, the bands having objective length were detected only in case of *Pectinatus frisingensis*. Accordingly, it was shown that each oligonucleotide of the present invention correctly recognized the gene sequences of the spacer region between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus frisingensis*, and the base sequence targeted on the gene coding 16S rRNA. Moreover, the bands having the aimed length were not observed in the same genus *Pectinatus cerevisiiphilus*, and relative strictly anaerobic bacteria and Gram-positive bacteria. Accordingly, *Pectinatus frisingensis* can be specifically detected, and at the same time also determined by the present invention.

Example 4

Cloning and determination of the base sequence of the spacer regions between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus cerevisiiphilus*

(1) Selection and synthesis of oligonucleotide primers for amplifying

5 16S/23S rRNA spacer regions by PCR

As the base sequence of 16S ribosome RNA gene of *Pectinatus cerevisiiphilus* is disclosed in International Journal of Systematic Bacteriology, Vol. 40, pages 19-27 (1990), the primers were selected on the basis of the base sequence of 557th -576th.

10 The base sequence of 23 ribosome RNA gene of *Pectinatus cerevisiiphilus* had not been disclosed, but the base sequence of 23 ribosome RNA gene of *Pectinatus frisingensis* had been disclosed in Systematic Applied Microbiology, Vol. 15, pages 487-501 (1990), EMBL Accession Number X48423. The primer was selected to obtain the complementary sequence corresponding to
15 the base sequence of 1st-20th of 23 ribosome RNA gene of *Pectinatus frisingensis*. Sawaday Technology was entrusted with the synthesis.

(2) Amplification of 16S/23S rRNA by PCR

The DNA solution 0.1 μ g of *Pectinatus cerevisiiphilus* prepared in Example 1 was charged in a 0.2 ml tube (Perkin-Elmer Co.), 5 μ l of 10x buffer
20 in rTaq DNA Polymerase Kit (TOYOBO Co., Ltd.), 3 μ l of 25mM MgCl₂, 5 μ l of 2mM dNTP mixture solution (dATP, dGTP, dCTP and dTTP), 0.5 μ l of 5 unit/ μ l rTaq-polymerase, each 0.5 μ l of the 100mM primers prepared in Example 2-(1) were added, and sterilized water was added to obtain final volume of 50 μ l. The tube was set in a thermal cycler of an automatic gene amplification device
25 (Perkin-Elmer Co.) and amplification reaction was conducted. 30 cycles were carried under the reaction conditions of one cycle of denaturation at 94°C for 2.5 minutes, denaturation at 94°C for 30 seconds, primer annealing at 55°C for 30 seconds and synthetic reaction at 72°C for 30 seconds. After the reaction, 5 μ l of the reactant was used in agarose gel electrophoresis, the DNA was stained

with ethidium bromide, and the amplified DNA was observed. As a result, DNA of about 1700 bp (abbreviated as "long") and DNA of about 1400 bp (abbreviated as "short") were amplified.

(3) Cloning and sequencing of the spacer region "long"

5 Using a high pure PCR product purification kit (Baringer Mannheim), unreactive dNTPs was removed from the solution after the PCR reaction. To 10 ng of the resulting amplified DNA, 2 μ l of plasmid pCR 2.1 contained in a TA cloning kit (INVITROGEN), 1 μ l of ligase and 1 μ l of buffer were added, and then sterilized water was added to obtain the total volume of 10 μ l. After the
10 solution was reacted at 14°C for 4 hours, 2 μ l of the solution and 2 μ l of 0.5 M β -mercaptoethanol were added to *Escherichia coli* INV α 'F competent cells, and placed in ice for 30 minutes. Then, the solution was heated at 42°C for 30 seconds, and plasmid transformation to the bacteria was conducted. To the transformed bacteria, 250 μ l of a SOC culture (2.0% Tryptone, 0.5% yeast
15 extract, 10.0 mM NaCl, 2.5 mM KCl, 10.0 mM MgCl₂·6H₂O, and 20.0 mM glucose) was added, and the mixture was shaken at 37°C for 60 minutes, then transferred to a LB plate culture medium containing 50 μ g / ml of ampicillin and 40 μ g / ml X-Gal, and cultured at 37°C overnight. The expressed white colony was transferred to 3 ml of a LB liquid culture medium containing 50 μ g /
20 ml of ampicillin, and cultured at 37°C overnight.

After the cultivation, plasmids were extracted from the bacteria with a plasmid mini kit (QIAGEN). A part of the resulting plasmids was taken out and reacted with a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) at 37 °C for 60 minutes, and separated by agarose electrophoresis. The DNA was
25 dyed with ethidium bromide, and insertion of "long" was confirmed. 500 ng of the residual plasmid was reacted with restriction enzyme *Sma*I (manufactured by TOYOBO Co., Ltd.) at 30°C for 60 minutes. To the reactant, 2 μ l of 3 M sodium acetate and 500 μ l of 100% ethanol were added, and the mixture was placed in ice for 15 minutes and centrifuged at 15000 rpm for 15 minutes, and

the supernatant was removed. To the precipitate, 500 μ l of 70% ethanol was added, the mixture was centrifuged at 15000 rpm for 15 minutes, and the supernatant was removed, and the residual was dried for 10 minutes under reduced pressure. Sterilized water was added to dissolve the precipitate, and
5 the mixture was reacted with restriction enzyme *Xba*I (Baringer Manhaim) at 37 °C for 60 minutes. To the reactant, equivalent phenol / chloroform (equivalent mixture liquid) was added and gently mixed, the mixture was centrifuged at 15000 rpm for 15 minutes, and the water layer (upper layer) was recovered. To the recovery liquid, equivalent water-saturated ether was added and gently
10 mixed, and the mixture was centrifuged at 15000 rpm for 15 minutes to remove the ether layer (upper layer). To the remaining water layer, 2 μ l of 3M sodium acetate and 500 μ l of 100% ethanol were added, and the mixture was placed in ice for 15 minutes and centrifuged at 15000rpm for 15 minutes to remove the supernatant.

15 To the precipitate, 500 μ l of 70% ethanol was added, and the mixture was centrifuged at 15000 rpm for 15 minutes to remove the supernatant, and the residue was dried under reduced pressure for 10 minutes, and 20 μ l of sterilized distillation water was added. To 5 μ l of the solution, 1 μ l of 10X buffer contained in a blunting kit (Takara Shuzo Co., Ltd.) and 3 μ l of sterilized
20 distillation water were added, and the mixture was maintained at 70°C for 5 minutes, 1 μ l of T4 DNA polymerase was added, and the mixture was maintained at 37°C for 5 minutes to obtain blunt ends. After T4 DNA polymerase was inactivated by stirring, 40 μ l of ligation solution A and 10 μ l of ligation solution B were added, and the mixture was maintained at 16°C for
25 30 minutes to conduct internal ligation. 2 μ l of the reactant and 2 μ l of 0.5M β -mercaptoethanol were added to *Escherichia coli* INV α 'F competent cells, and the mixture was placed in ice for 30 minutes and heated at 42°C for 30 seconds, and the plasmid was transformed to the *Escherichia coli*.

To the transformed *Escherichia coli*, 250 μ l of a SOC culture medium

(2.0% Tryptone, 0.5% Yeast extract, 10.0 mM NaCl, 2.5 mM KCl, 10.0 mM $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 20.0 mM glucose) was added, and the mixture was shaken at 37°C for 60 minutes and spread on a LB plate culture medium containing 50 $\mu\text{g} / \text{ml}$ of ampicillin to culture at 37°C overnight. Appeared white colonies were
 5 inoculated into 3 ml of a LB liquid culture medium containing 50 $\mu\text{g} / \text{ml}$ of ampicillin and cultured at 37°C overnight. After the culture, the plasmid was extracted from the *Escherichia coli* with a plasmid mini kit (QIAGEN Company).

Using such obtained plasmid as a template, a sequence reaction was conducted. As the sequencing primers, an IDRD41 Infrared Dye Labeled M13
 10 Forward primer and an IRD41 Infrared Dye Labeled M13 Reverse primer (manufactured by Nisshinbo, sold by Aroka Co., Ltd.) were used. As the reaction liquid, SequiTherm (trademark) Long-Read (trademark) Cycle Sequencing Kit-LC (manufactured by EPICENTRE TECHNOLOGIES) was used. 4000L Long ReadIR (trademark) DNA Sequencing System
 15 (manufactured by LI-COR) was used for the determination of the base sequences.

The gene sequence of spacer region "long" between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus cerevisiiphilus* is shown in SEQ ID NO: 3.

20 (4) Cloning and sequencing of spacer region "short"

Using a high pure PCR product purification kit (Baringer Mannheim), unreactive dNTPs was removed from the solution after the PCR reaction in Example 4-(2). To 100 ng of the resulting amplified DNA, 2 μl of plasmid pCR 2.1 contained in a TA cloning kit (INVITROGEN), 1 μl of ligase and 1 μl of
 25 buffer were added, and then sterilized water was added to obtain the total volume of 10 μl . After the solution was reacted at 14°C for 4 hours, 2 μl of the solution and 2 μl of 0.5 M β -mercaptoethanol were added to *Escherichia coli* INV α 'F competent cells, and placed in ice for 30 minutes. Then, the solution was heated at 42°C for 30 seconds, and plasmid transformation to the

bacteria was conducted. To the transformed bacteria, 250 μ l of a SOC culture (2.0% Tryptone, 0.5% yeast extract, 10.0 mM NaCl, 2.5 mM KCl, 10.0 mM $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, and 20.0 mM glucose) was added, and the mixture was shaken at 37°C for 60 minutes, then transferred to a LB plate culture medium containing
 5 50 μ g / ml of ampicillin and 40 μ g / ml X-Gal, and cultured at 37°C overnight. The appeared white colony was transferred to 3 ml of a LB liquid culture medium containing 50 μ g / ml of ampicillin, and cultured at 37°C overnight. After the cultivation, plasmid was extracted from the bacteria with a plasmid mini kit (QIAGEN).

10 A part of the resulting plasmid was taken out and reacted with a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) at 37°C for 60 minutes, and the reactant was separated by agarose electrophoresis. The DNA was dyed with ethidium bromide, and insertion of "short" was confirmed. 500 ng of the residual plasmid was reacted with restriction enzyme *Sma*I
 15 (manufactured by TOYOBOKU Co., Ltd.) at 30°C for 60 minutes. To the reactant, 2 μ l of 3 M sodium acetate and 500 μ l of 100% ethanol were added, and the mixture was placed in ice for 15 minutes and centrifuged at 15000 rpm for 15 minutes, and the supernatant was removed. To the precipitate, 500 μ l of 70% ethanol was added, the mixture was centrifuged at 15000 rpm for 15 minutes,
 20 and the supernatant was removed, and the residue was dried for 10 minutes under reduced pressure. Sterilized water was added to dissolve the precipitate, and the mixture was reacted with restriction enzyme *Bam*HI (Takara Shuzo Co.) at 37°C for 60 minutes. To the reactant, equivalent phenol / chloroform (equivalent mixture liquid) was added and gently mixed, the mixture was
 25 centrifuged at 15000 rpm for 15 minutes, and the water layer (upper layer) was recovered. To the recovery liquid, equivalent water-saturated ether was added and gently mixed, and the mixture was centrifuged at 15000 rpm for 15 minutes to remove the ether layer (upper layer). To the remaining water layer, 2 μ l of 3M sodium acetate and 500 μ l of 100% ethanol were added, and the mixture

was placed in ice for 15 minutes and centrifuged at 15000rpm for 15 minutes to remove the supernatant.

To the precipitate, 500 μ l of 70% ethanol was added, and the mixture was centrifuged at 15000 rpm for 15 minutes to remove the supernatant, and the
5 residue was dried under reduced pressure for 10 minutes, and 20 μ l of sterilized distilled water was added. To 5 μ l of the solution, 1 μ l of 10X buffer contained in a blunting kit (Takara Shuzo Co., Ltd.) and 3 μ l of sterilized distilled water were added, and the mixture was maintained at 70°C for 5 minutes, 1 μ l of T4 DNA polymerase was added, and the mixture was maintained at 37°C for 5
10 minutes to obtain blunt ends. After T4 DNA polymerase was inactivated by stirring, 40 μ l of ligation solution A and 10 μ l of ligation solution B were added, and the mixture was maintained at 16°C for 30 minutes to conduct internal ligation. 2 μ l of the reactant and 2 μ l of 0.5M β -mercaptoethanol were added to a *Escherichia coli* INV α 'F competent cell, and the mixture was
15 placed in ice for 30 minutes and heated at 42°C for 30 seconds, and the plasmid was transformed to the *Escherichia coli*. To the transformed *Escherichia coli*, 250 μ l of a SOC culture medium (2.0% Tryptone, 0.5% Yeast extract, 10.0 mM NaCl, 2.5 mM KCl, 10.0 mM MgCl₂·6H₂O, 20.0 mM glucose) was added, and the mixture was shaken at 37°C for 60 minutes and spread on a LB plate culture
20 medium containing 50 μ g / ml ampicillin to culture at 37°C overnight. Appeared white colonies were inoculated into 3 ml of a LB liquid culture medium containing 50 μ g / ml of ampicillin and cultured at 37°C overnight. After the culture, the plasmid was extracted from the *Escherichia coli* with a plasmid kit (QIAGEN Company).

25 Using such obtained plasmid as a template, a sequence reaction was conducted. As the sequencing primers, an IRD41 Infrared Dye Labeled M13 Forward primer and an IRD41 Infrared Dye Labeled M13 Reverse primer (manufactured by Nisshinbo, sold by Aroka Co., Ltd.) were used. As the reaction liquid, SequiTherm (trademark) Long-Read (trademark) Cycle

Sequencing Kit-LC (manufactured by EPICENTRE TECHNOLOGIES) was used. 4000L Long ReadIR (trademark) DNA Sequencing System (manufactured by LI-COR) was used for the determination of the base sequences.

- 5 The gene sequence of spacer region "short" between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus cerevisiiphilus* is shown in SEQ ID NO: 4.

Example 5

Detection of *Pectinatus cerevisiiphilus* by the PCR method

- 10 (1) Selection and synthesis of a primer for *Pectinatus cerevisiiphilus*

The sequences specific for *Pectinatus cerevisiiphilus* using DNASIS (tradename of Hitachi Soft Engineering Ltd., Co.) on the basis of SEQ ID NO: 3 were analyzed. The result selected a sequence of 135th to 153rd on the gene sequence of the spacer region between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus cerevisiiphilus* of SEQ ID NO: 3. (SEQ ID NO: 7.)

In addition, the similar analysis selected a sequence of 172nd to 191st on the gene sequence of the spacer region between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus cerevisiiphilus* of SEQ ID NO: 3. (SEQ ID NO: 8.)

- 20 The similar analysis also selected a sequence of 203rd to 222nd on the gene sequence of the spacer region between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus cerevisiiphilus* of SEQ ID NO: 3. (SEQ ID NO: 9.)

Further, specific primer showing in SEQ ID NO: 11 was selected by a gene sequence coding 16S rRNA of *Pectinatus cerevisiiphilus*. The oligonucleotides were chemically synthesized by the same method as in Example 2-(1).

(2) Detection and identification of *Pectinatus cerevisiiphilus* by the primers having the sequences of SEQ ID NO: 7 and SEQ ID NO: 11.

The DNA solutions of bacteria prepared in Example 1 were treated with the primers synthesized in Example 5-(1) (SEQ ID NO: 7 and SEQ ID NO: 11) by PCR. The temperature conditions of the PCR were as follows:

Thermal denaturation; 94°C, 30 seconds

5 Annealing; 55°C, 30 seconds

Chain elongation reaction; 72°C, 30 seconds

One cycle of the conditions was repeated 35 times. After the PCR, the reactant was electrophoresed with agarose gel at constant 100 V for 30 minutes. A pHY marker was also electrophoresed at the same time as a molecular weight
10 marker. After the electrophoresis, the gel was stained with 5 μ g/ml of an ethidium bromide solution for 20 minutes, and ultraviolet was applied to observe the gel and take a photograph of the gel. By the observation or the photography of the gel, the base length of the amplified products was determined from the relative migration distance with a molecular weight marker.

15 As shown in Fig. 2, a band of about 600 bps was detected only in case of *Pectinatus cerevisiiphilus*.

From the results, when the oligonucleotides of SEQ ID NO: 7 and SEQ ID NO: 11 were used as PCR primers, the band having objective length was detected only in case of *Pectinatus cerevisiiphilus*. Accordingly, it was shown
20 that each oligonucleotide of the present invention correctly recognized the gene sequences of the spacer region between the gene coding 16S rRNA and the gene coding 23S rRNA of *Pectinatus cerevisiiphilus*, and the base sequence targeted on the gene coding 16S rRNA. Moreover, the bands having the aimed length were not observed in the same genus *Pectinatus frisingensis*, and relative
25 strictly anaerobic bacteria and Gram-positive bacteria. Accordingly, *Pectinatus cerevisiiphilus* can be specifically detected, and at the same time also determined by the present invention.

By the present invention, the genes of the spacer region constituted between the 16S rRNA genes and the 23S rRNA genes of *Pectinatus frisingensis*

and *Pectinatus cerevisiiphilus* have been proved, and a method for quickly and reliably detecting *Pectinatus frisingensis* and *Pectinatus cerevisiiphilus* can be provided by using a part or all of the gene sequences.

Claims

1. A gene sequence of a spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus frisingensis* containing a part of the base sequence or the whole base sequence represented by SEQ ID NO: 1.
- 5 2. A gene sequence of a spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus frisingensis* containing a part of the base sequence or the whole base sequence represented by SEQ ID NO: 2.
3. A gene sequence of a spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus cerevisiiphilus* containing a part of
10 the base sequence or the whole base sequence represented by SEQ ID NO: 3.
4. A gene sequence of a spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus cerevisiiphilus* containing a part of the base sequence or the whole base sequence represented by SEQ ID NO: 4.
5. An oligonucleotide characterized in that the gene sequence of a
15 spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus frisingensis* has at least one of the following sequence group or the corresponding complementary sequence:
 - 5'-CCATCCTCTTGAAAATCTC-3' ①
 - 5'-TCTCRTCTCACAAGTTTGGC-3' ②.
- 20 6. An oligonucleotide that the gene sequence of a spacer region between a gene coding 16S rRNA and a gene coding 23S rRNA of *Pectinatus cerevisiiphilus* has at least one of the following sequence group or the corresponding complementary sequence:
 - 5'-CACTCTTACAAGTATCTAC-3' ③
 - 25 5'-CCACAATATTTCCGACCAGC-3' ④
 - 5'-AGTCTTCTCTACTGCCATGC-3' ⑤.
7. A method for detecting *Pectinatus frisingensis*, comprising employing the oligonucleotide made from the gene sequence described in claim 1 or 2 as a primer for synthesis of nucleic acids, and treating the nucleic acid by

gene amplification to detect the bacteria.

8. A method for detecting *Pectinatus cerevisiiphilus*, comprising
employing the oligonucleotide made from the gene sequence described in claim 3
or 4 as a primer for synthesis of nucleic acids, and treating the nucleic acid by
5 gene amplification to detect the bacteria.

9. A method for detecting *Pectinatus frisingensis*, comprising
employing the oligonucleotide made from the gene sequence described in claim 1
or 2, or the oligonucleotide made from the gene sequence described in claim 5,
and a nucleotide sequence coding 16S rRNA gene of *Pectinatus frisingensis* as
10 primers for synthesis of nucleic acids, and treating the nucleic acid by gene
amplification to detect the bacteria.

10. A method for detecting *Pectinatus cerevisiiphilus*, comprising
employing the oligonucleotide made from the gene sequence described in claim 3
or 4 or the oligonucleotide made from the gene sequence described in claim 6,
15 and a nucleotide sequence coding 16S rRNA gene of *Pectinatus cerevisiiphilus* as
primers for synthesis of nucleic acids, and treating the nucleic acid by gene
amplification to detect the bacteria.

11. A method as claimed in claim 9, wherein the nucleotide sequence
coding the 16S rRNA gene of *Pectinatus frisingensis* has the following sequence:
20 5'-CGTATCCAGAGATGGATATT-3' ⑥

12. A method as claimed in claim 10, wherein the nucleotide sequence
coding the 16S rRNA gene of *Pectinatus cerevisiiphilus* has the following
sequence:

5'-CGTATGCAGAGATGCATATT-3' ⑦

Abstract

The present invention relates to genes for detecting the genus *Pectinatus* *frisingensis* or *Pectinatus cerevisiiphilus* of the genus *Pectinatus*, which is known as beer-spoilage bacteria, and a method for detecting the bacteria by
5 using the genes.

The present invention provides gene sequences of spacer regions between 16S rRNA genes and 23S rRNA genes specific for the genus *Pectinatus* relating to beer-spoilage and a method for quickly and sensitively detecting the bacteria by using the sequences.

Fig. 1





Fig. 2



Sequence Listing

<110> Asahi Breweries, Ltd.
 <120> Genes for detecting bacteria and a detection method using them
 <130> 98T3-10
 5 <160> 11
 <210> 1
 <211> 624
 <212> DNA
 <213> *Pectinatus frisingensis*
 10 <400> 1
 gaagtcgtaa caaggtagcc gtatcggaag gtgcggctgg atcacctcct ttctaaggat 60
 taaaacaatc cgtcgagcac atccggaaca tgtattgttt ggttttgagg gtttctccct 120
 caaaaaaata gatagaacta atgggggcgt agctcagctg ggagagcacc tgccttgcaa 180
 gcagggggtc aggagttcaa atctcctcgt ctccaccaga agagaaatgg gcctatagct 240
 15 cagctggtta gagegcacgc ctgataagcg tgaggtcagt agttcaagtc tacttaggcc 300
 caccataatt gcacattgaa aactacacag aagaaaagca aagaacaatt aatcaccaat 360
 gccaaacttg tgagaggaga ttttcaagag gatggcgggg aatagttgga ccaagcacia 420
 ttaggaaact aaaaacaagc taagacaaaa catataaact taagctaaag gtgatattct 480
 ggaggagact cgagaatata ataaacttac cagaagcggt cagatgcaag gaagcatgaa 540
 20 agcgaatgaa gaaggcgtat tagtatacgc cgatgagtga gctgaaatga tgacgaagca 600
 gatgagcggg tatggaaagt ttaa 624

 <210> 2
 <211> 442
 25 <212> DNA
 <213> *Pectinatus frisingensis*
 <400> 2
 gaagtcgtaa caaggtagcc gtatcggaag gtgcggctgg atcacctcct ttctaaggat 60
 taaaacaatc cgtcgagcac atccggaaca tgtattgttt ggttttgagg gtttctccct 120
 30 caaatattgc acattgaaaa ctacacagaa gaaaagcaaa gaacaattaa tcaccaatgc 180
 caaacttgat agaagagatt ttcaagagga tggcggggaa tagttggacc aagcacaatt 240
 aggaaactaa aaacaagcta agacaaaaca tataaactta agctaaaggt gatattcttg 300

aggagactcg agaataataa aaacttacca gaagcggttca gatgcaagga agcatgaaag 360
 cgaatgaaga aggcgtatta gtatacgccg atgagtgagc tgaaatgatg acgaagcaga 420
 tgagcggtta tggaaagttt aa 442

5 <210> 3
 <211> 724
 <212> DNA
 <213> *Pectinatus cerevisiiphilus*
 <400> 3

10 gaagtcgtaa caaggtagcc gtatcggaag gtgcggctgg atcacctcct ttctaaggat 60
 ttgacaaaaa tctgtcgagt acatccggaa tatgtattgt ttggttttga gggtttctcc 120
 ctcataaata tatagtagat acttgtaaga gtgtttatgg tatgtttaaa agctggctcg 180
 aaatattgtg gtgcaaaaaa atgcatggca gtagagaaga ctggtaaaaa aagaatgaac 240
 taatgggggc gtagctcaga tgggagagca cctgccttgc aagcaggggg tcaggagttc 300
 15 aactctctctc gtctccacca gaagagaaag ggcctatagc tcagctgggt agagcgcacg 360
 cctgataagc gtgaggtcag tagttcaagt ctacttaggc ccaccaatat tgcacattga 420
 aaactacaca gaagaaagca aagaacaatt atcaccaatg ccaaacttgt aagagaaatc 480
 gaggagagaa tggcggggaa tagttggacc aagcacaat taggaaaaga aacaaacgct 540
 aagaacaaaa catataaact taagcgaaaa ggtgatattc tggaggaaac ttcagagtat 660
 20 ataaacttac cagaagcgtt cagatgcgag gaagggcaaa gctgagagaa gaaagcgtat 660
 taatatacgc tgatgaacga agcaaagcac tgacaaagca gatggatggt tatgggaagt 720
 taca
 724

25 <210> 4
 <211> 399
 <212> DNA
 <213> *Pectinatus cerevisiiphilus*
 <400> 4

30 gaagtcgtaa caaggtagcc gtatcggaag gtgcggctgg atcacctcct ttctaaggat 60
 ttgacaaaaa tctgtcgagt acatccggaa tatgtattgt ttggttttga gggtttctcc 120

ctcataaata ttgcacattg aaaactacac agaagaaagc aaagaacaat taccaccaat 180
 gccaaacttg taagagaaat cgagaagaga atggcgggga atagttggac caagcacaaa 240
 ttaggaaaag aaacaaacgc taagaaacaa acatataaac ttaagcgaag aggtgatatt 300
 ctggaggaaa cttcagagta tataaactta ccagaagcgt tcagatgcga ggaagggcaa 360
 5 agcactgaca aagtagatgg atggttatgg gaagttaca 399

<210> 5
 <211> 19
 <212> DNA
 10 <213> *Pectinatus frisingensis*
 <400> 5
 ccacctctt gaaaatctc 19

<210> 6
 15 <211> 19
 <212> DNA
 <213> *Pectinatus frisingensis*
 <400> 6
 tctcrtctca caagtttgcc 20

20 <210> 7
 <211> 19
 <212> DNA
 <213> *Pectinatus cerevisiiphilus*
 25 <400> 7
 cactcttaca agtatctac 19

<210> 8
 <211> 20
 30 <212> DNA
 <213> *Pectinatus cerevisiiphilus*
 <400> 8
 ccacaatatt tccgaccagc 20

35 <210> 9
 <211> 20

<212> DNA
<213> *Pectinatus cerevisiiphilus*
<400> 9
5 agtcttctct actgccatgc 20

<210> 10
<211> 20
<212> DNA
<213> *Pectinatus frisingensis*
10 <400> 10
cgtatccaga gatggatatt 20

<210> 11
<211> 20
15 <212> DNA
<213> *Pectinatus cerevisiiphilus*
<400> 11
cgtatccaga gatggatatt 20