

(19)



(11)

EP 4 494 710 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
22.01.2025 Bulletin 2025/04

(51) International Patent Classification (IPC):
A61P 35/00 ^(2006.01)

(21) Application number: **24203877.6**

(52) Cooperative Patent Classification (CPC):
(C-Sets available)
**A61K 39/39558; A61K 47/68037; A61K 47/6851;
A61K 47/6855; A61K 47/6857; A61K 47/6861;
A61K 47/6863; A61K 47/6869; A61K 47/6889;
A61P 1/02; A61P 1/04; A61P 1/16; A61P 1/18;
A61P 11/00; A61P 13/02;**

(Cont.)

(22) Date of filing: **10.04.2015**

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

- **ABRAHAM, Reimar**
82152 Martinsried (DE)
- **BLUM, Sabine**
82152 Martinsried (DE)
- **UENO, Suguru**
Tokyo, 140-8710 (JP)

(30) Priority: **10.04.2014 JP 2014081454**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
**20200710.0 / 3 789 042
15719011.7 / 3 129 063**

(74) Representative: **Marks & Clerk LLP**
15 Fetter Lane
London EC4A 1BW (GB)

(71) Applicants:

- **Daiichi Sankyo Company, Limited**
Tokyo 103-8426 (JP)
- **Daiichi Sankyo Europe GmbH**
81379 München (DE)

Remarks:


- The complete document including Reference Table(s) and the Sequence Listing(s) can be downloaded from the EPO website
- This application was filed on 01-10-2024 as a divisional application to the application mentioned under INID code 62.

(72) Inventors:

- **HETTMANN, Thore**
82152 Martinsried (DE)

(54) **PHARMACEUTICAL COMPOSITION COMPRISING AN ANTI-HER3 ANTIBODY-DRUG CONJUGATE FOR USE IN TREATING BREAST CANCERS**

(57) Provided is a pharmaceutical composition comprising an anti-HER3 antibody-drug conjugate, for use in treating breast cancers. In the antibody-drug conjugate, an antitumor compound represented by the following formula is conjugated to an anti-HER3 antibody via a linker having a structure represented by the formula: $-L^1-L^2-L^P-NH-(CH_2)_n-L^A-(CH_2)_n-C(=O)-$ or $-L^1-L^2-L^P-$ (the anti-HER3 antibody is connected to the terminal of L^1 , the antitumor compound is connected to the carbonyl group of $-(CH_2)_n-C(=O)-$ moiety or the C terminal of L^P , with the nitrogen atom of the amino group at position 1 as a connecting position):

**EP 4 494 710 A2**

(52) Cooperative Patent Classification (CPC): (Cont.)

A61P 13/08; A61P 13/10; A61P 13/12;

A61P 15/00; A61P 17/00; A61P 25/00;

A61P 35/00; A61P 43/00; C07K 16/2863;

C07K 16/3015; C07K 16/3023; C07K 16/32;

A61K 2039/505; C07K 2317/21; C07K 2317/24;

C07K 2317/76; C07K 2317/77

C-Sets

A61K 39/39558, A61K 2300/00

Description

Technical Field


[0001] The present invention relates to an antibody-drug conjugate having an anti-HER3 antibody and an antitumor drug conjugated to each other via a linker structure moiety, the conjugate being useful as an antitumor drug.

Background Art

[0002] An antibody-drug conjugate (ADC) having a drug with cytotoxicity conjugated to an antibody which binds to an antigen expressed on a surface of cancer cells and capable of cellular internalization (the antibody which binds to the antigen is also capable of cellular internalization), can deliver the drug selectively to the cancer cells and is thus expected to cause accumulation of the drug in the cancer cells and to kill the cancer cells (see, Non Patent Literatures 1 to 3). As an ADC, Mylotarg (registered trademark; Gemtuzumab ozogamicin) in which calicheamicin is conjugated to an anti-CD33 antibody is approved as a therapeutic agent for acute myeloid leukemia. Further, Adcetris (registered trademark; Brentuximab vedotin), in which auristatin E is conjugated to an anti-CD30 antibody, has recently been approved as a therapeutic agent for Hodgkin's lymphoma and anaplastic large cell lymphoma (see, Non Patent Literature 4). The drugs contained in ADCs which have been approved until now target DNA or tubulin.

[0003] As an antitumor, low-molecular-weight compounds, camptothecin derivatives, which inhibit topoisomerase I to exhibit an antitumor effect, are known. Among them, an antitumor compound represented by the formula below (exatecan, chemical name: (1S,9S)-1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-10,13(9H,15H)-dione) is a water soluble derivative of camptothecin (Patent Literature 1 and 2).

[Chem. 1]



[0004] Unlike irinotecan currently used in clinical settings, this compound does not require activation by an enzyme for exerting its antitumor effect. Further, compared to SN-38 as a main pharmaceutically active ingredient of irinotecan and topotecan also used in clinical settings, it has higher inhibitory activity on topoisomerase I and has higher cytotoxic activity in vitro against various cancer cells. In particular, it exhibits the effect against cancer cells which have resistance to SN-38 or the like due to expression of P-glycoprotein. Further, in a mouse model with a human tumor subcutaneously transplanted, it exhibited a potent antitumor effect, and thus has undergone the clinical studies, but has not been put on the market yet (see, Non Patent Literatures 5 to 10). It remains unclear whether or not exatecan functions effectively as an ADC.

[0005] DE-310 is a complex in which exatecan is conjugated to a biodegradable carboxymethyldextran polyalcohol polymer via a GGFG peptide spacer (Patent Literature 3). By making exatecan into a form of a polymer prodrug, a high blood retention property can be maintained and also a high penetration property to a tumor area is passively increased by utilizing the increased permeability of newly formed tumor vessels and retention property in tumor tissues. With DE-310, the peptide spacer is cleaved by an enzyme to continuously release exatecan as a main active ingredient and exatecan with glycine bonded to an amino group, and as a result, the pharmacokinetics are improved. According to various tumor evaluation models in non-clinical studies, it was found that higher effectiveness was obtained by DE-310 than exatecan administered alone even though the total amount of exatecan contained therein is lower than the case of administration of exatecan alone. A clinical study was conducted for DE-310, and effective cases were confirmed. There is also a report suggesting that the main active ingredient accumulates in a tumor than in normal tissues. However, there is also a report

indicating that the accumulation of DE-310 and the main active ingredient in a tumor is not much different from the accumulation in normal tissues, and thus no passive targeting is observed in humans (see, Non Patent Literatures 11 to 14). As a result, DE-310 was not also commercialized, and it remains unclear whether or not exatecan effectively functions as a drug oriented for such targeting.

[0006] As a compound relating to DE-310, a complex in which a structure moiety represented by $\text{-NH-(CH}_2\text{)}_4\text{-C(=O)-}$ is inserted between -GGFG -spacer and exatecan to form $\text{-GGFG-NH-(CH}_2\text{)}_4\text{-C(=O)-}$ used as a spacer structure is also known (Patent Literature 4). However, the antitumor effect of the complex is not known at all.

[0007] The human epidermal growth factor receptor 3 (also known as HER3 and ErbB3) is a receptor protein tyrosine kinase and belongs to the epidermal growth factor receptor (EGFR) subfamily of receptor protein tyrosine kinases, which also includes HER1 (also known EGFR), HER2, and HER4 (see Non Patent Literatures 15 to 17). As with the prototypical epidermal growth factor receptor, the transmembrane receptor HER3 consists of an extracellular ligand-binding domain (ECD), a dimerization domain within the ECD, a transmembrane domain, and a carboxyl-terminal phosphorylation domain. HER1, HER2, and HER4 carry an intracellular protein tyrosine kinase domain (TKD) in addition to these domains, while HER3 lacks this domain and is thus unable to be autophosphorylated.

[0008] The ligand Heregulin (HRG) binds to the extracellular domain of HER3 and activates the receptor-mediated signaling pathway by promoting dimerization with other human epidermal growth factor receptor (HER) family members and transphosphorylation of its intracellular domain. The dimer formation of HER3 with other HER family members expands the signaling potential of HER3 and serves as means not only for signal diversification but also for signal amplification. For example, the HER2/ HER3 heterodimer induces one of the most important mitogenic signals among HER family members. HER3 is overexpressed in several types of cancers such as breast, gastrointestinal, and pancreatic cancers. Interestingly, a correlation between the expression of HER2/HER3 and the progression from a non-invasive stage to an invasive stage has been shown (see Non Patent Literatures 18 to 20). Accordingly, agents that interfere with HER3-mediated signaling are desirable. Anti-HER3 antibodies and immunoconjugates thereof have been reported in, for example, Patent Literatures 5 to 10, respectively.

Citation List

Patent Literature

[0009]

- [PTL 1] Japanese Patent Laid-Open No. 5-59061
- [PTL 2] Japanese Patent Laid-Open No. 8-337584
- [PTL 3] International Publication No. WO 1997/46260
- [PTL 4] International Publication No. WO 2000/25825
- [PTL 5] U.S. Patent No. 5968511
- [PTL 6] U.S. Patent No. 5480968
- [PTL 7] International Publication No. WO 2003/013602
- [PTL 8] International Publication No. WO 2007/077028
- [PTL 9] International Publication No. WO 2008/100624
- [PTL 10] International Publication No. WO 2012/019024

Non Patent Literature

[0010]

- [NPL 1] Ducry, L., et al., Bioconjugate Chem. (2010) 21, 5-13.
- [NPL 2] Alley, S. C., et al., Current Opinion in Chemical Biology (2010) 14, 529-537.
- [NPL 3] Damle N.K. Expert Opin. Biol. Ther. (2004) 4, 1445-1452.
- [NPL 4] Senter P. D., et al., Nature Biotechnology (2012) 30, 631-637.
- [NPL 5] Kumazawa, E., Tohgo, A., Exp. Opin. Invest. Drugs (1998) 7, 625-632.
- [NPL 6] Mitsui, I., et al., Jpn J. Cancer Res. (1995) 86, 776-786.
- [NPL 7] Takiguchi, S., et al., Jpn J. Cancer Res. (1997) 88, 760-769.
- [NPL 8] Joto, N. et al., Int J Cancer (1997) 72, 680-686.
- [NPL 9] Kumazawa, E. et al., Cancer Chemother. Pharmacol. (1998) 42, 210-220.
- [NPL 10] De Jager, R., et al., Ann N Y Acad Sci (2000) 922, 260-273.
- [NPL 11] Inoue, K. et al. Polymer Drugs in the Clinical Stage, Edited by Maeda et al., (2003) 145-153.
- [NPL 12] Kumazawa, E. et al., Cancer Sci (2004) 95, 168-175.

- [NPL 13] Soepenbergh, O. et al., Clinical Cancer Research, (2005) 11, 703-711.
 [NPL 14] Wente M. N. et al., Investigational New Drugs (2005) 23, 339-347.
 [NPL 15] Plowman, et al., Proc. Natl. Acad. Sci. U.S.A. (1990) 87, 4905-4909.
 [NPL 16] Kraus et al., Proc. Natl. Acad. Sci. U.S.A. (1989) 86, 9193-9197.
 [NPL 17] Kraus et al., Proc. Natl. Acad. Sci. U.S.A. (1993) 90, 2900-2094.
 [NPL 18] Alimandi et al., Oncogene (1995) 10, 1813-1821.
 [NPL 19] DeFazio et al., Int. J. Cancer (2000) 87, 487-498.
 [NPL 20] Nadiu et al., Br. J. Cancer (1998) 78, 1385-1390.

Summary of Invention

Technical Problem

[0011] With regard to the treatment of tumor using an antibody, an insufficient antitumor effect may be observed even when the antibody recognizes an antigen and binds to tumor cells, and thus a more effective antitumor antibody is sometimes needed. Further, many antitumor low-molecular-weight compounds have a problem in safety like side effect and toxicity even the compounds have an excellent antitumor effect. As such, it remains as a subject to achieve a superior therapeutic effect by further enhancing the safety. Thus, an object of the present invention is to provide an antitumor drug having an excellent therapeutic effect, which is excellent in terms of antitumor effect and safety.

Solution to Problem


[0012] The inventors thought that, since the anti-HER3 antibody is an antibody capable of targeting tumor cells, that is, it is an antibody having a property of recognizing tumor cells, a property of binding to tumor cells, a property of internalizing in tumor cells, a cytotoxic activity against tumor cells, or the like, when exatecan as an antitumor compound is converted into an antibody-drug conjugate by conjugation to the antibody via a linker structure moiety, the antitumor compound can be more surely delivered to tumor cells to specifically exhibit the antitumor effect of the compound in tumor cells, and thus the antitumor effect can be surely exhibited and also an enhanced cytotoxic effect of the anti-HER3 antibody is expected, and a dose of the antitumor compound can be reduced compared to a case of administering the compound alone, and thus an influence of the antitumor compound on normal cells can be alleviated so that higher safety can be achieved.

[0013] In this connection, the inventors created a linker with a specific structure and succeeded in obtaining an antibody-drug conjugate in which the anti-HER3 antibody and exatecan are conjugated to each other via the linker, and confirmed an excellent antitumor effect exhibited by the conjugate to thereby complete the present invention.


[0014] Specifically, the present invention relates to the followings.

[1] An antibody-drug conjugate wherein an antitumor compound represented by the following formula

[Chem. 2]



is conjugated to an anti-HER3 antibody by a thioether bond which is formed at a disulfide bond moiety present in a hinge part of the anti-HER3 antibody via a linker having a structure represented by the following formula:



[0015] Here, the anti-HER3 antibody is connected to the terminal of L¹, the antitumor compound is connected to the carbonyl group of -(CH₂)^{n²}-C(=O)- moiety or the C terminal of L^P, with the nitrogen atom of the amino group at position 1 as a connecting position.

In the formula, n¹ represents an integer of 0 to 6,

n² represents an integer of 0 to 5,

L¹ represents -(Succinimid-3-yl-N)-(CH₂)^{n³}-C(=O)-,

wherein n³ represents an integer of 2 to 8,

L² represents -NH-(CH₂CH₂-O)^{n⁴}-CH₂CH₂-C(=O)- or a single bond,


wherein n⁴ represents an integer of 1 to 6,

L^P represents a peptide residue consisting of 2 to 7 amino acids,

L^a represents -O- or a single bond,

-(Succinimid-3-yl-N)- has a structure represented by the following formula:

[Chem. 3]



which is connected to the anti-HER3 antibody at position 3 thereof and is connected on the nitrogen atom at position 1 to a methylene group in the linker structure containing this structure.

[0016] The present invention further relates to each of the followings.

[2] The antibody-drug conjugate according to [1], wherein the peptide residue of L^P is a peptide residue comprising an amino acid selected from phenylalanine, glycine, valine, lysine, citrulline, serine, glutamic acid, and aspartic acid.

[3] The antibody-drug conjugate according to [1] or [2], wherein L^P is a peptide residue selected from the following group

-GGF-,

-DGGF-,

-(D-)D-GGF-,

-EGGF-,

-GGFG-,

-SGGF-,

-KGGF-,

-DGGFG-,

-GGFGG-,

-DDGGFG-,

-KDGGFG-,

and

-GGFGGGF-;

(wherein, "(D-)D" represents D-aspartic acid).

[4] The antibody-drug conjugate according to [1] or [2], wherein L^P is a peptide residue comprising 4 or 5 amino acids.

[5] The antibody-drug conjugate according to any one of [1] to [4], wherein L^P is -GGFG- or -DGGFG-.

[6] The antibody-drug conjugate according to any one of [1] to [4], wherein L^P is -GGFG-.

[7] The antibody-drug conjugate according to any one of [1] to [6], wherein n^3 is an integer of 2 to 5 and L^2 is a single bond.

[8] The antibody-drug conjugate according to any one of [1] to [7], wherein the linker is $-L^1-L^2-L^P-NH-(CH_2)^{n^1}-L^a-(CH_2)^{n^2}-C(=O)-$.

[9] The antibody-drug conjugate according to [8], wherein n^3 is an integer of 2 to 5, L^2 is $-NH-(CH_2CH_2-O)^{n^4}-CH_2CH_2-C(=O)-$, and n^4 is 2 or 4.

[10] The antibody-drug conjugate according to [8] or [9], wherein $-NH-(CH_2)^{n^1}-L^a-(CH_2)^{n^2}-C(=O)-$ is a partial structure having chain length of 4 to 7 atoms.

[11] The antibody-drug conjugate according to [8] or [9], wherein $-NH-(CH_2)^{n^1}-L^a-(CH_2)^{n^2}-C(=O)-$ is a partial structure having chain length of 5 or 6 atoms.

[12] The antibody-drug conjugate described in [10] or [11], wherein $-NH-(CH_2)^{n^1}-L^a-(CH_2)^{n^2}-C(=O)-$ is

$-NH-CH_2CH_2-C(=O)-$,

$-NH-CH_2CH_2CH_2-C(=O)-$,

$-NH-CH_2CH_2CH_2CH_2-C(=O)-$,

$-NH-CH_2CH_2CH_2CH_2CH_2-C(=O)-$,

$-NH-CH_2-O-CH_2-C(=O)-$,

or

$-NH-CH_2CH_2-O-CH_2-C(=O)-$.

[13] The antibody-drug conjugate according to [12], wherein $-NH-(CH_2)^{n^1}-L^a-(CH_2)^{n^2}-C(=O)-$ is any one of the followings:

$-NH-CH_2CH_2CH_2-C(=O)-$,

$-NH-CH_2-O-CH_2-C(=O)-$,

or

$-NH-CH_2CH_2-O-CH_2-C(=O)-$.

[14] The antibody-drug conjugate according to any one of [1] to [5], wherein the linker is $-L^1-L^2-L^P-$.

[15] The antibody-drug conjugate according to [14], wherein L^P is -DGGFG-.

[16] The antibody-drug conjugate according to [15], wherein n^3 is an integer of 2 to 5 and L^2 is a single bond.


[17] The antibody-drug conjugate according to [1], wherein the drug-linker structure moiety in which a drug is bound to $-L^1-L^2-L^P-NH-(CH_2)^{n^1}-L^a-(CH_2)^{n^2}-C(=O)-$ or $-L^1-L^2-L^P-$ is one drug-linker structure selected from the following group:

- (Succinimid-3-yl-N)- $CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2CH_2CH_2-C(=O)-(NH-DX)$,

- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-NH-CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-NH-CH₂CH₂CH₂CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-GGFG-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-DGGFG-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX).

[0017] In the above, -(Succinimid-3-yl-N)- has a structure represented by the following formula:


[Chem. 4]



which is connected to the anti-HER3 antibody at position 3 and is connected to a methylene group in the linker structure containing it on the nitrogen atom at position 1,

- (NH-DX) represents a group represented by the following formula, wherein the nitrogen atom of the amino group at position 1 is the connecting position,

[Chem. 5]



- GGFG- represents a tetrapeptide residue of -Gly-Gly-Phe-Gly- and -DGGFG-represents a pentapeptide residue of -Asp-Gly-Gly-Phe-Gly-.

[0018] [18] The antibody-drug conjugate described in [1], wherein the drug-linker structure moiety having a drug bonded


to $-L^1-L^2-L^P-NH-(CH_2)_{n^1}-L^a-(CH_2)_{n^2}-C(=O)-$ is one drug-linker structure selected from the following group:

- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-DGGFG-NH-CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2-O-CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2-C(=O)-NH-CH_2CH_2-O-CH_2CH_2-O-CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-DGGFG-(NH-DX)$.


Here, $-(Succinimid-3-yl-N)-$, $-(NH-DX)$, $-GGFG-$, and $-DGGFG-$ are as described above.

[19] An antibody-drug conjugate comprising an antitumor compound represented by the following formula:

[Chem. 6]



conjugated to an anti-HER3 antibody by a thioether bond which is formed at a disulfide bond moiety present in a hinge part of the anti-HER3 antibody via a linker having a structure represented by the following formula:



Here, the anti-HER3 antibody is connected to the terminal of L^1 and the antitumor compound is connected to the carbonyl group of $-(CH_2)_{n^2}-C(=O)-$ moiety. In the formula, n^1 represents an integer of 0 to 6,

n^2 represents an integer of 0 to 5,

L^1 represents $-(Succinimid-3-yl-N)-(CH_2)_{n^3}-C(=O)-$,

wherein n^3 represents an integer of 2 to 8,

L^2 represents $-NH-(CH_2CH_2-O)_{n^4}-CH_2CH_2-C(=O)-$ or a single bond,


wherein n^4 represents an integer of 1 to 6,

L^P represents a tetrapeptide residue of $-GGFG-$,

L^a represents $-O-$ or a single bond,

- (Succinimid-3-yl-N)- has a structure represented by the following formula:

[Chem. 7]



which is connected to the anti-HER3 antibody at position 3 thereof and binds on the nitrogen atom at position 1 to a methylene group in a linker structure containing this structure.

[20] The antibody-drug conjugate according to [19], wherein n^1 is 3, n^2 is 0, n^3 is 2, L^2 is $-\text{NH}-(\text{CH}_2\text{CH}_2-\text{O})$ n^4 - $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-$, n^4 is 2, and L^a is a single bond, or

n^1 is 1, n^2 is 1, n^3 is 5, L^2 is a single bond, and L^a is $-\text{O}-$, or
 n^1 is 2, n^2 is 1, n^3 is 5, L^2 is a single bond, and L^a is $-\text{O}-$.

[21] The antibody-drug conjugate according to [19] or [20], wherein n^3 is 2 or 5 and L^2 is a single bond.

[22] The antibody-drug conjugate according to [19] or [20], wherein n^3 is 2 or 5, L^2 is $-\text{NH}-(\text{CH}_2\text{CH}_2-\text{O})$ n^4 - $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-$, and n^4 is 2 or 4.

[23] The antibody-drug conjugate described in any one of [19] to [22], wherein $-\text{NH}-(\text{CH}_2)n^1-L^a-(\text{CH}_2)n^2-\text{C}(=\text{O})-$ is

$-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-$,

$-\text{NH}-\text{CH}_2-\text{O}-\text{CH}_2-\text{C}(=\text{O})-$,

or


$-\text{NH}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2-\text{C}(=\text{O})-$.

[24] The antibody-drug conjugate described in any one of [19] to [23], wherein the drug-linker structure moiety having a drug bonded to $-\text{L}^1-\text{L}^2-\text{L}^p-\text{NH}-(\text{CH}_2)n^1-L^a-(\text{CH}_2)n^2-\text{C}(=\text{O})-$ is one drug-linker structure selected from the following group:

- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX),
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX),
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX),
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX),
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX),
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2-\text{O}-\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX),
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX),
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -NH- $\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX),
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -NH- $\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX),
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -NH- $\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX);
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -NH- $\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -GGFG-NH- $\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})$ -(NH-DX),

[0020] In the above, $-(\text{Succinimid-3-yl-N})-$ has a structure represented by the following formula:


[Chem. 8]



which is connected to the anti-HER3 antibody at position 3 thereof and is connected on the nitrogen atom at position 1 to a methylene group in a linker structure containing this structure.

- (NH-DX) represents a group represented by the following formula, wherein the nitrogen atom of the amino group at position 1 is the connectig position:

[Chem. 9]



- GGFG- represents a tetrapeptide residue of -Gly-Gly-Phe-Gly-.

[0021] [25] The antibody-drug conjugate described in any one of [19] to [23], wherein the drug-linker structure moiety having a drug connected to $-L^1-L^2-L^P-NH-(CH_2)^{n1}-L^a-(CH_2)^{n2}-C(=O)-$ is one drug-linker structure selected from the following group:

- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2-O-CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2-O-CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2-C(=O)-NH-CH_2CH_2-O-CH_2CH_2-O-CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2CH_2-C(=O)-(NH-DX)$.

In the above, -(Succinimid-3-yl-N)-, -(NH-DX), and -GGFG- are as defined above.

[0022] [26] The antibody-drug conjugate according to any one of [1] to [25], wherein the average number of units of the selected one drug-linker structure conjugated per antibody is in a range of from 1 to 10.

[0023] [27] The antibody-drug conjugate according to any one of [1] to [25], wherein the average number of units of the selected one drug-linker structure conjugated per antibody is in a range of from 2 to 8.

[0024] [28] The antibody-drug conjugate according to any one of [1] to [25], wherein the average number of units of the selected one drug-linker structure conjugated per antibody is in a range of from 3 to 8.

[0025] [29] A medicine comprising the antibody-drug conjugate according to any one of [1] to [28], a salt thereof or a hydrate thereof.

[0026] [30] An antitumor medicine and/or anticancer medicine comprising the antibody-drug conjugate according to any one of [1] to [28], a salt thereof or a hydrate thereof.

[0027] [31] The antitumor medicine and/or anticancer medicine according to [30], which is applied to lung cancer, kidney cancer, urothelial cancer, colorectal cancer, prostate cancer, glioblastoma multiforme, ovarian cancer, pancreatic cancer, breast cancer, melanoma, liver cancer, bladder cancer, stomach cancer, gastrointestinal stromal tumor, cervical cancer, head and neck cancer, esophageal cancer, epidermoid cancer, peritoneal cancer, adult glioblastoma multiforme, hepatic cancer, hepatocellular carcinoma, colon cancer, rectal cancer, colon and rectal cancer, endometrial cancer, uterus cancer, salivary cancer, renal cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anus carcinoma, or penis cancer.

[0028] [32] A pharmaceutical composition comprising the antibody-drug conjugate according to any one of [1] to [28], a salt thereof or a hydrate thereof as an active component, and a pharmaceutically acceptable formulation component.

[0029] [33] The pharmaceutical composition according to [32], which is applied to lung cancer, kidney cancer, urothelial cancer, colorectal cancer, prostate cancer, glioblastoma multiforme, ovarian cancer, pancreatic cancer, breast cancer, melanoma, liver cancer, bladder cancer, stomach cancer, gastrointestinal stromal tumor, cervical cancer, head and neck cancer, esophageal cancer, epidermoid cancer, peritoneal cancer, adult glioblastoma multiforme, hepatic cancer, hepatocellular carcinoma, colon cancer, rectal cancer, colon and rectal cancer, endometrial cancer, uterus cancer, salivary cancer, renal cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anus carcinoma, or penis cancer.


[0030] [34] A method for treating a tumor and/or cancer comprising administering the antibody-drug conjugate according to any one of [1] to [28], a salt thereof or a hydrate thereof.

[0031] [35] The medicine according to [29], the antitumor medicine and/or anticancer medicine according to [30] or [31],


the pharmaceutical composition according to [32] or [33], or the treatment method according to [34], which is used in administration in combination with an additional medicine.

[0032] [36] The pharmaceutical composition according to [32] or [33], further comprising even an additional medicine as an active ingredient.

[0033] [35] A method for producing an antibody-drug conjugate comprising reacting a compound represented by the following formula:



or



with an anti-HER3 antibody or a reactive derivative thereof and conjugating a drug-linker moiety to the antibody by a method for forming a thioether bond on a disulfide bond moiety present at a hinge part of the antibody.

[0034] In the formula, n^3 represents an integer of 2 to 8,

L^2 represents $-\text{NH}-(\text{CH}_2\text{CH}_2-\text{O})_{n^4}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-$ or a single bond wherein n^4 represents an integer of 1 to 6, L^{P} represents a peptide residue consisting of 2 to 7 amino acids selected from phenylalanine, glycine, valine, lysine, citrulline, serine, glutamic acid, and aspartic acid,


n^1 represents an integer of 0 to 6,

n^2 represents an integer of 0 to 5,

L^{a} represents $-\text{O}-$ or a single bond,


(maleimid-N-yl)- is a group represented by the following formula and has a nitrogen atom as a connecting position.

[Chem. 10]



-(NH-DX) represents a group represented by the following formula, wherein the nitrogen atom of the amino group at position 1 is the connectig position:


[Chem. 11]



CH₂CH₂-O-CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂-C(=O)-GGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂-C(=O)-GGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂-C(=O)-GGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂-C(=O)-DGGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX), or
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX).

[0041] In the above, (maleimid-N-yl)- is a group represented by the following formula:


[Chem. 12]



which has a nitrogen atom as a connecting position.

- (NH-DX) represents a group represented by the following formula, the nitrogen atom of the amino group at position 1 being a connecting position.

[Chem. 13]



- GGFG- represents a tetrapeptide residue of -Gly-Gly-Phe-Gly- and -DGGFG- represents pentapeptide residue of -Asp-Gly-Gly-Phe-Gly-.

[0042] [42] An antibody-drug conjugate obtained by forming a thioether bond at a disulfide bond site present in a hinge part of an anti-HER3 antibody, and characterized by treating the anti-HER3 antibody with a reducing condition and thereafter reacting with a compound selected from the compound group shown below:

(maleimid-N-yl)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-(NH-DX), or
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-(NH-DX).

In the above, (maleimid-N-yl)-, -(NH-DX), and -GGFG- are as defined above.

[0043] [43] The antibody-drug conjugate according to [41] or [42], wherein an average conjugated number of the selected one drug-linker structure per antibody is in a range of from 1 to 10.

[0044] [44] The antibody-drug conjugate according to [41] or [42], wherein an average conjugated number of the selected one drug-linker structure per antibody is in a range of from 2 to 8.

[0045] [45] The antibody-drug conjugate according to [41] or [42], wherein an average conjugated number of the selected one drug-linker structure per antibody is in a range of from 3 to 8.

[0046] [46] A medicine comprising the antibody-drug conjugate according to any one of [40] to [45], a salt thereof or a hydrate thereof.

[0047] [47] An antitumor medicine and/or anticancer medicine comprising the antibody-drug conjugate according to any one of [40] to [45], a salt thereof or a hydrate thereof.

[0048] [48] The antitumor medicine and/or anticancer medicine according to [47], which is applied to lung cancer, kidney cancer, urothelial cancer, colorectal cancer, prostate cancer, glioblastoma multiforme, ovarian cancer, pancreatic cancer, breast cancer, melanoma, liver cancer, bladder cancer, stomach cancer, gastrointestinal stromal tumor, cervical cancer, head and neck cancer, esophageal cancer, epidermoid cancer, peritoneal cancer, adult glioblastoma multiforme, hepatic cancer, hepatocellular carcinoma, colon cancer, rectal cancer, colon and rectal cancer, endometrial cancer, uterus cancer, salivary cancer, renal cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anus carcinoma, or penis cancer.

[0049] [49] A pharmaceutical composition comprising the antibody-drug conjugate according to any one of [40] to [45], a salt thereof or a hydrate thereof as an active component, and a pharmaceutically acceptable formulation component.

[0050] [50] The pharmaceutical composition according to [49], which is applied to lung cancer, kidney cancer, urothelial cancer, colorectal cancer, prostate cancer, glioblastoma multiforme, ovarian cancer, pancreatic cancer, breast cancer, melanoma, liver cancer, bladder cancer, stomach cancer, gastrointestinal stromal tumor, cervical cancer, head and neck cancer, esophageal cancer, epidermoid cancer, peritoneal cancer, adult glioblastoma multiforme, hepatic cancer, hepatocellular carcinoma, colon cancer, rectal cancer, colon and rectal cancer, endometrial cancer, uterus cancer, salivary cancer, renal cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anus carcinoma, or penis cancer.

[0051] [51] A method for treating a tumor and/or cancer comprising administering the antibody-drug conjugate according to any one of [40] to [45], a salt thereof or a hydrate thereof.

[0052] [52] The medicine according to [46], the antitumor medicine and/or anticancer medicine according to [47] or [48], the pharmaceutical composition according to [49] or [50], or the treatment method according to [51], which is used in administration in combination with an additional medicine.

[0053] [53] The pharmaceutical composition according to [49] or [50], further comprising even an additional medicine as an active ingredient.

Advantageous Effects of Invention

[0054] With an anti-HER3 antibody-drug conjugate having an antitumor compound exatecan conjugated via a linker with a specific structure, an excellent antitumor effect and safety can be achieved.

Brief Description of Drawings

[0055]

[Fig. 1]

Figure 1 shows the full-length amino acid sequence of a heavy chain of anti-HER3 human antibody U1-59 (SEQ ID NO: 583).

[Fig. 2]

Figure 2 shows the full-length amino acid sequence of a light chain of anti-HER3 human antibody U1-59 (SEQ ID NO: 584).

[Fig. 3]

Figure 3 shows the mean fluorescence intensity of HCC1569 treated with serial dilutions of U1-59 or each antibody-drug conjugate. KD and Bmax values were calculated using GraphPad Prism Software.

[Fig. 4]

A549 cells were cultured for 2 days with U1-59 or varied antibody-drug conjugates. HER3 or phosphorylated HER3 was evaluated by Western blotting. pan-Actin was detected as an electrophoresis control.

[Fig. 5]

Figure 5 shows an average value of reduction in HER3 expression on the surface of HCC1569 cells treated with U1-59 or each antibody-drug conjugate (37C ("C" represents "degrees Celsius"), 1 hr).

[Fig. 6]

Figure 6 shows results of a test on the inhibition of mitogenic or survival signals by each HER3 antibody-drug

conjugate in a human breast cancer line (HCC1569). Figure 6A shows cell growth or survival derived from the antibody-drug conjugate in the presence of 10% FBS. The data is indicated by mean $\pm$ standard deviation of triplicates. The ordinate depicts a luminescence value indicating the ATP activity of each sample. The abscissa depicts the concentration of each antibody-drug conjugate. Figure 6B shows the rate of reduction in luminescence caused by antibody-drug conjugate treatment when the luminescence of an untreated group was defined as 100%. [Fig. 7]

Figure 7 shows results of a test on the inhibition of mitogenic or survival signals by each HER3 antibody-drug conjugate in a human breast cancer line (MDA-MB 453). Figure 7A shows cell growth or survival derived from the antibody-drug conjugate in the presence of 10% FBS. The ordinate depicts a luminescence value indicating the ATP activity of each sample. The abscissa depicts the concentration of each antibody-drug conjugate. The data is indicated by mean $\pm$ standard deviation of triplicates. Figure 7B shows the rate of reduction in luminescence caused by antibody-drug conjugate treatment when the luminescence of an untreated group was defined as 100%. [Fig. 8]

Figure 8 shows results of a test on the inhibition of mitogenic or survival signals by each HER3 antibody-drug conjugate in a human melanoma line (A375). Figure 8A shows cell growth or survival derived from the antibody-drug conjugate in the presence of 10% FBS. The ordinate depicts a luminescence value indicating the ATP activity of each sample. The abscissa depicts the concentration of each antibody-drug conjugate. The data is indicated by mean $\pm$ standard deviation of triplicates. Figure 8B shows the rate of reduction in luminescence caused by antibody-drug conjugate treatment when the luminescence of an untreated group was defined as 100%. [Fig. 9]

Figure 9 shows results of a test on the inhibition of mitogenic or survival signals by each HER3 antibody-drug conjugate in a human colorectal cancer line (HT29). Figure 9A shows cell growth or survival derived from the antibody-drug conjugate in the presence of 10% FBS. The ordinate depicts a luminescence value indicating the ATP activity of each sample. The abscissa depicts the concentration of each antibody-drug conjugate. The data is indicated by mean $\pm$ standard deviation of triplicates. Figure 9B shows the rate of reduction in luminescence caused by antibody-drug conjugate treatment when the luminescence of an untreated group was defined as 100%. [Fig. 10]

Figure 10 shows results of a test on the inhibition of mitogenic or survival signals by each HER3 antibody-drug conjugate in a human lung cancer line (A549). Figure 10A shows cell growth or survival derived from the antibody-drug conjugate in the presence of 10% FBS. The ordinate depicts a luminescence value indicating the ATP activity of each sample. The abscissa depicts the concentration of each antibody-drug conjugate. The data is indicated by mean $\pm$ standard deviation of triplicates. Figure 10B shows the rate of reduction in luminescence caused by antibody-drug conjugate treatment when the luminescence of an untreated group was defined as 100%. [Fig. 11]

Figure 11 shows results of comparing the rate of inhibition of cell growth or survival between the antibody-drug conjugate (3) and the antibody-drug conjugate (4). The left diagram shows the rate of inhibition of cell growth or survival derived from the antibody-drug conjugate in the presence of 10% FBS. The ordinate depicts luminescence indicating the ATP activity of each sample. The abscissa depicts the concentration of each antibody-drug conjugate. The data is indicated by mean $\pm$ standard deviation of triplicates. The right diagram shows the comparison of the rate of reduction in luminescence caused by antibody-drug conjugate treatment between high drug loading (HDL) and middle drug loading (MDL) when the luminescence of an untreated group was defined as 100%. [Fig. 12]

Figure 12 shows results of comparing the rate of inhibition of cell growth or survival between the antibody-drug conjugate (10) and the antibody-drug conjugate (11). The left diagram shows the rate of inhibition of cell growth or survival derived from the antibody-drug conjugate in the presence of 10% FBS. The ordinate depicts luminescence indicating the ATP activity of each sample. The abscissa depicts the concentration of each antibody-drug conjugate. The data is indicated by mean $\pm$ standard deviation of triplicates. The right diagram shows the comparison of the rate of reduction in luminescence caused by antibody-drug conjugate treatment between high drug loading (HDL) and middle drug loading (MDL) when the luminescence of an untreated group was defined as 100%. [Fig. 13]

Figure 13 shows results of comparing the rate of inhibition of cell growth or survival between the antibody-drug conjugate (13) and the antibody-drug conjugate (14). The left diagram shows the rate of inhibition of cell growth or survival derived from the antibody-drug conjugate in the presence of 10% FBS. The ordinate depicts luminescence indicating the ATP activity of each sample. The abscissa depicts the concentration of each antibody-drug conjugate. The data is indicated by mean $\pm$ standard deviation of triplicates. The right diagram shows the comparison of the rate of reduction in luminescence caused by antibody-drug conjugate treatment between high drug loading (HDL) and middle drug loading (MDL) when the luminescence of an untreated group was defined as 100%. [Fig. 14]

Figure 14 shows results of a human breast cancer (HCC1569) antitumor test using the antibody-drug conjugate (3), (10), or (13). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 15]

Figure 15 shows results of a human melanoma (HT-144) antitumor test using the antibody-drug conjugate (3), (10), or (13). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 16]

Figure 16 shows results of a human breast cancer (MDA-MB-453) antitumor test using the antibody-drug conjugate (3), (10), or (13). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from administration. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 17]

Figure 17 shows results of a human colorectal cancer line (HT-29) antitumor test using the antibody-drug conjugate (3), (10), or (13). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from administration. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 18]

Figure 18 shows results of a human lung cancer line (A549) antitumor test using the antibody-drug conjugate (3), (10), or (13). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 19]

Figure 19 shows results of a human triple-negative breast cancer line (MDA-MB-468) antitumor test using the antibody-drug conjugate (13). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 20]

Figure 20 shows results of a human luminal breast cancer line (MCF-7) antitumor test using the antibody-drug conjugate (16a). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 21]

Figure 21 shows results of a human melanoma line (WM-266-4) antitumor test using the antibody-drug conjugate (16a). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 22]

Figure 22 shows results of a human ovarian cancer line (OVCAR-8) antitumor test using the antibody-drug conjugate (16a). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 23]

Figure 23 shows results of a human bladder cancer line (SW-780) antitumor test using the antibody-drug conjugate (16a). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 24]

Figure 24 shows results of a human breast cancer line (MDA-MB-453) antitumor test using the antibody-drug conjugate (16a). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

[Fig. 25]

Figure 25 shows results of a human breast cancer line (MDA-MB-453) antitumor test using the antibody-drug

conjugate (16a). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009. [Fig. 26]

Figure 26 shows results of a human breast cancer line (JIMT-1) antitumor test using the antibody-drug conjugate (15). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009. [Fig. 27]

Figure 27 shows results of a human lung cancer line (PC9) antitumor test using the antibody-drug conjugate (16a). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009. [Fig. 28]

Figure 28 shows results of a human triple-negative breast cancer line (MDA-MB-468) antitumor test using the antibody-drug conjugate (16a). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009. [Fig. 29]

Figure 29 shows results of a human head and neck cancer line (Fadu) antitumor test using the antibody-drug conjugate (16a). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009. [Fig. 30]

Figure 30 shows results of an antitumor test using a human stomach cancer patient-derived tumor section (NIBIO-G016) and the antibody-drug conjugate (16a). The ordinate depicts an average tumor volume. The abscissa depicts the number of days from cell transplantation. All values are indicated by mean $\pm$ standard deviation. The initial tumor volume and the initial mouse weight were analyzed on the basis of descriptive data (mean and standard deviation) using Microsoft Excel 2009.

Description of Embodiments

[0056] Hereinbelow, the preferred embodiments for carrying out the present invention are explained in view of the drawings. Meanwhile, the embodiments explained below are the examples of the representative embodiments of the present invention and the scope of the present invention shall not be narrowly interpreted based on them.

[0057] The present invention provides HER3 binding protein-drug conjugate. Preferably, the HER3 binding protein of the invention is a scaffold protein having an antibody like binding activity or an antibody, i.e. an anti-HER3 antibody.

[0058] The anti-HER3 antibody-drug conjugate of the present invention is an antitumor medicine in which an anti-HER3 antibody is conjugated to an antitumor compound via a linker structure moiety and explained in detail hereinbelow.

[0059] Within the context of the present invention, the term "scaffold protein", as used herein, means a polypeptide or protein with exposed surface areas in which amino acid insertions, substitutions or deletions are highly tolerable. Examples of scaffold proteins that can be used in accordance with the present invention are protein A from *Staphylococcus aureus*, the bilin binding protein from *Pieris brassicae* or other lipocalins, ankyrin repeat proteins, and human fibronectin (reviewed in Binz and Pluckthun, *Curr Opin Biotechnol*, 16, 459-69). Engineering of a scaffold protein can be regarded as grafting or integrating an affinity function onto or into the structural framework of a stably folded protein. Affinity function means a protein binding affinity according to the present invention. A scaffold can be structurally separable from the amino acid sequences conferring binding specificity. In general, proteins appearing suitable for the development of such artificial affinity reagents may be obtained by rational, or most commonly, combinatorial protein engineering techniques such as panning against HER3, either purified protein or protein displayed on the cell surface, for binding agents in an artificial scaffold library displayed in vitro, skills which are known in the art (Skerra, *J. Mol. Recog.*, 2000; Binz and Pluckthun, 2005). In addition, a scaffold protein having an antibody like binding activity can be derived from an acceptor polypeptide containing the scaffold domain, which can be grafted with binding domains of a donor polypeptide to confer the binding specificity of the donor polypeptide onto the scaffold domain containing the acceptor polypeptide. Said inserted binding domains may be, for example, the complementarity determining region (CDR) of an antibody, in particular an anti-HER3 antibody. Insertion can be accomplished by various methods known to those skilled in the art including, for example, polypeptide synthesis, nucleic acid synthesis of an encoding amino acid as well by various forms of recombinant methods well known to those skilled in the art.

{Antibody}

[0060] Moreover, the term "antibody" or "anti-HER3 antibody", as used herein, means a monoclonal antibody, a polyclonal antibody, a recombinant antibody, a humanized antibody (Jones et al., Nature 321 (1986), 522-525; Riechmann et al., Nature 332 (1988), 323-329; and Presta, Curr. Op. Struct. Biol. 2 (1992), 593-596), a chimeric antibody (Morrison et al., Proc. Natl. Acad. Sci. U.S.A. 81 (1984), 6851-6855), a human antibody and fully human antibody, (Tomizuka, K. et al., Nature Genetics (1997) 16, p.133-143.; Kuroiwa, Y. et al., Nucl. Acids Res. (1998) 26, p.3447-3448; Yoshida, H. et al., Animal Cell Technology: Basic and Applied Aspects vol. 10, p.69-73 (Kitagawa, Y., Matsuda, T. and Iijima, S. eds.), Kluwer Academic Publishers, 1999.; Tomizuka, K. et al., Proc. Natl. Acad. Sci. USA (2000) 97, p.722-727, International Publication No. WO 2007/077028, and so on), a multispecific antibody (e.g. a bispecific antibody) formed from at least two antibodies, or an antibody fragment thereof. The term "antibody fragment" comprises any portion of the aforementioned antibodies, preferably their antigen binding region or variable regions. Examples of antibody fragments include Fab fragments, Fab' fragments, F(ab')₂ fragments, Fv fragments, diabodies (Hollinger et al., Proc. Natl. Acad. Sci. U.S.A. 90 (1993), 6444-6448), single chain antibody molecules (Pluckthun in: The Pharmacology of Monoclonal Antibodies 113, Rosenberg and Moore, EDS, Springer Verlag, N.Y. (1994), 269-315) and other fragments as long as they exhibit the desired capability of binding to HER3.

[0061] In addition, the term "antibody" or "anti-HER3 antibody", as used herein, may include antibody-like molecules that contain engineered sub-domains of antibodies or naturally occurring antibody variants. These antibody-like molecules may be single-domain antibodies such as VH-only or VL-only domains derived either from natural sources such as camelids (Muyldermans et al., Reviews in Molecular Biotechnology 74, 277-302) or through in vitro display of libraries from humans, camelids or other species (Holt et al., Trends Biotechnol., 21, 484-90).

[0062] In accordance with the present invention, the "Fv fragment" is the minimum antibody fragment that contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDR's of each variable domain interact to define an antigen-binding site on the surface of the V_H-V_L dimer. Collectively, the six CDR's confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDR's specific for an antigen) has the ability to recognize and bind the antigen, although usually at a lower affinity than the entire binding site.

[0063] The "Fab fragment" also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. The "Fab fragment" differs from the "Fab' fragment" by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. The "F(ab')₂ fragment" originally is produced as a pair of "Fab' fragments" which have hinge cysteines between them. Methods of preparing such antibody fragments, such as papain or pepsin digestion, are known to those skilled in the art.

[0064] In another preferred embodiment of the present invention, the anti-HER3 antibody of the invention is an anti-HER3 antibody directed against the extracellular domain (ECD) of HER3.

[0065] The anti-HER3 antibody used in an anti-HER3 antibody-drug conjugate of the present invention may be derived from any species. Preferred examples of the species can include humans, rats, mice, and rabbits. The anti-HER3 antibody derived from other than human species is preferably chimerized or humanized using a well known technique. The antibody of the present invention may be a polyclonal antibody or a monoclonal antibody and is preferably a monoclonal antibody.

[0066] The anti-HER3 antibody is may be those which are capable of targeting tumor cells and thus possesses the property of being capable of recognizing tumor cells, the property of being capable of binding to tumor cells, the property of being internalized into tumor cells, and cytotoxic activity against tumor cells, etc. The anti-HER3 antibody can be conjugated with a compound having antitumor activity via a linker to form an antibody-drug conjugate.

[0067] The binding activity of the antibody against tumor cells can be confirmed using flow cytometry. The internalization of the antibody into tumor cells can be confirmed using (1) an assay of visualizing an antibody incorporated in cells under a fluorescence microscope using a secondary antibody (fluorescently labeled) binding to the therapeutic antibody (Cell Death and Differentiation (2008) 15, 751-761), (2) an assay of measuring the amount of fluorescence incorporated in cells using a secondary antibody (fluorescently labeled) binding to the therapeutic antibody (Molecular Biology of the Cell, Vol. 15, 5268-5282, December 2004), or (3) a Mab-ZAP assay using an immunotoxin binding to the therapeutic antibody wherein the toxin is released upon incorporation into cells to inhibit cell growth (Bio Techniques 28: 162-165, January 2000). A recombinant complex protein of a diphtheria toxin catalytic domain and protein G may be used as the immunotoxin.

[0068] The antitumor activity of the antibody can be confirmed *in vitro* by determining inhibitory activity against cell growth. For example, a cancer cell line overexpressing a target protein for the antibody is cultured, and the antibody is added at varying concentrations into the culture system to determine inhibitory activity against focus formation, colony formation, and spheroid growth. The antitumor activity can be confirmed *in vivo*, for example, by administering the antibody to a nude mouse with a transplanted tumor cell line highly expressing the target protein, and determining change in the cancer (tumor) cells.

[0069] Since the compound conjugated in the antibody-drug conjugate exerts an antitumor effect, it is preferred but not

essential that the antibody itself should have an antitumor effect. For the purpose of exerting the cytotoxicity of the antitumor compound specifically and selectively for tumor cells, it is important and also preferred that the antibody should have the property of being internalized to migrate into tumor cells.

[0070] The anti-HER3 antibody can be obtained using a method usually carried out in the art, which involves immunizing animals with an antigenic polypeptide and collecting and purifying antibodies produced *in vivo*. The origin of the antigen is not limited to humans, and the animals may be immunized with an antigen derived from a non-human animal such as a mouse or a rat and the like. In this case, the cross-reactivity of antibodies binding to the obtained heterologous antigen with human antigens can be tested to screen for an antibody applicable to a human disease.

[0071] Alternatively, antibody-producing cells which produce antibodies against the antigen are fused with myeloma cells according to a method known in the art (e.g., Kohler and Milstein, *Nature* (1975) 256, p. 495-497; and Kennet, R. ed., *Monoclonal Antibodies*, p. 365-367, Plenum Press, N.Y. (1980)) to establish hybridomas, from which monoclonal antibodies can in turn be obtained.

[0072] The antigen can be obtained by genetically engineering host cells to produce a gene encoding the antigenic protein. Specifically, vectors that permit expression of the antigen gene are prepared and transferred to host cells so that the gene is expressed. The antigen thus expressed can be purified. The antibody can also be obtained by use of a method which involves immunizing animals with the genetically engineered antigen-expressing cells or a cell line with an expressed antigen.

[0073] The anti-HER3 antibody can be obtained by means known in the art.

[0074] The anti-HER3 antibody that can be used in the present invention is not particularly limited and is desirably, for example, any of antibodies having properties as described below.

(1) An anti-HER3 antibody having the following properties:

(a) specifically binding to HER3, and/or

(b) having the activity of being internalized into HER3-expressing cells through binding to HER3.

(2) The antibody according to (1), wherein the antibody binds to the extracellular domain of HER3.

(3) The antibody according to (1) or (2), wherein the antibody is a monoclonal antibody.

(4) The antibody according to any of (1) to (3), wherein the antibody has antibody-dependent cell-mediated cytotoxicity (ADCC) activity and/or complement-dependent cytotoxicity (CDC) activity.

(5) The antibody according to any of (1) to (4), wherein the antibody is a mouse monoclonal antibody, a chimeric monoclonal antibody, a humanized monoclonal antibody, or a human or fully human (monoclonal) antibody.

(6) The antibody according to any of (1) to (5), wherein the antibody is a humanized monoclonal antibody comprising a heavy chain comprising the amino acid sequence represented by SEQ ID NO: 1 and a light chain comprising the amino acid sequence represented by SEQ ID NO: 2.

(7) The antibody according to any of (1) to (6), wherein the antibody lacks a lysine residue at the carboxy terminus of the heavy chain.

(8) The antibody according to (7), wherein the antibody comprises a heavy chain variable region represented by the amino acid sequence represented by SEQ ID NO: 70 and a light chain variable region represented by the amino acid sequence represented by SEQ ID NO: 72.

(9) An antibody obtained by a method for producing the antibody according to any of (1) to (8), the method comprising the steps of: culturing a host cell transformed with an expression vector comprising a polynucleotide encoding the antibody; and collecting the antibody of interest from the cultures obtained in the preceding step.

[0075] Hereinafter, the anti-HER3 antibody used in the present invention will be described.

[0076] In the present specification, the terms "cancer" and "tumor" are used interchangeably.

[0077] In the present specification, the term "gene" includes not only DNA but its mRNA, cDNA, and cRNA thereof.

[0078] In the present specification, the term "polynucleotide" is used interchangeably with a nucleic acid and also includes DNA, RNA, probes, oligonucleotides, and primers.

[0079] In the present specification, the terms "polypeptide" and "protein" are used interchangeably.

[0080] In the present specification, the term "cell" also includes cells in animal individuals and cultured cells.

[0081] In the present specification, the term "HER3" is used interchangeably with HER3 protein.

[0082] In the present specification, the term "CDR" means a complementarity determining region (CDR). An antibody molecule is known to have three CDRs in each of heavy and light chains. CDRs, also called hypervariable domains, are located in the variable regions of the antibody heavy and light chains. These sites have a particularly highly variable primary structure and are separated at three positions on the respective primary structures of heavy and light chain polypeptide strands. In the present specification, the antibody CDRs are referred to as CDRH1, CDRH2, and CDRH3 from the amino terminus of the heavy chain amino acid sequence as to heavy chain CDRs and as CDRL1, CDRL2, and CDRL3

from the amino terminus of the light chain amino acid sequence as to light chain CDRs. These sites are proximal to each other on the three-dimensional structure and determine specificity for the antigen to be bound.

[0083] In the present invention, the phrase "hybridizing under stringent conditions" refers to hybridization at 68C in a commercially available hybridization solution ExpressHyb Hybridization Solution (manufactured by Clontech Laboratories, Inc.), or identifiable hybridization under conditions involving hybridization at 68C in the presence of 0.7 to 1.0 M NaCl using a DNA-immobilized filter, followed by washing at 68C using 0.1 to 2 × SSC solution (1 × SSC is composed of 150 mM NaCl and 15 mM sodium citrate), or hybridization under conditions equivalent thereto.

1. HER3

[0084] The human epidermal growth factor receptor 3 (HER3, also known as ErbB3) is a receptor protein tyrosine kinase and belongs to the epidermal growth factor receptor (EGFR) subfamily of receptor protein tyrosine kinases, which also includes HER1 (also known as EGFR), HER2, and HER4. HER3 is a transmembrane receptor and consists of an extracellular ligand-binding domain (ECD), a dimerization domain within the ECD, a transmembrane domain, an intracellular protein tyrosine kinase domain (TKD) and a C-terminal phosphorylation domain. HER3 has been found to be overexpressed in several types of cancer such as breast, gastrointestinal and pancreatic cancers. A correlation between the expression of HER2/HER3 and the progression from a non-invasive to an invasive stage has been shown.

[0085] The HER3 protein used in the present invention can be used after direct purification from HER3-expressing human or non-human mammalian (rat, mouse, etc.) cells or can be used by preparing cell membrane fractions of the cells. Alternatively, HER3 may be synthesized *in vitro* or may be produced from host cells by genetic engineering. In the genetic engineering, specifically, HER3 cDNA is integrated into vectors that permit expression thereof, and HER3 can then be expressed by synthesis in a solution containing enzymes necessary for transcription and translation, substrates, and energy substances or by transformation of other host prokaryotic cells or host eukaryotic cells to yield the protein. Alternatively, the genetically engineered HER3-expressing cells described above or a cell line with expressed HER3 may be used as the HER3 protein.

[0086] An RNA sequence, a cDNA sequence, and an amino acid sequence of HER3 are available in public database, and can be referred to by an accession number such as AAA35979 (precursor including a signal sequence consisting of amino terminus 19 amino acid residue), M34309 (NCBI), for example.

[0087] The above amino acid sequence of HER3 consists of an amino acid sequence which is subjected to replacements, deletions, additions and/or insertions of at least one amino acid, and proteins having a biological activity equivalent to that of the protein are also included in HER3.

2. Production of anti HER3 antibody

[0088] The antibody against HER3 of the present invention can be obtained by immunizing an animal with HER3 or an arbitrary polypeptide selected from the amino acid sequence of HER3, and collecting and purifying the antibody produced in vivo according to a method usually carried out in the art. The biological species of HER3 to be used as an antigen is not limited to being human, and an animal can be immunized with HER3 derived from an animal other than humans such as a mouse or a rat. In this case, by examining the cross-reactivity between an antibody binding to the obtained heterologous HER3 and human HER3, an antibody applicable to a human disease can be selected.

[0089] Further, a monoclonal antibody can be obtained from a hybridoma established by fusing antibody-producing cells which produce an antibody against HER3 with myeloma cells according to a known method (for example, Kohler and Milstein, Nature, (1975) 256, pp. 495-497; Kennet, R. ed., Monoclonal Antibodies, pp. 365-367, Plenum Press, N.Y. (1980)).

[0090] HER3 to be used as an antigen can be obtained by expressing HER3 gene in a host cell using genetic engineering.

[0091] Specifically, a vector capable of expressing HER3 gene is produced, and the resulting vector is transfected into a host cell to express the gene, and then, the expressed HER3 is purified.

[0092] It is also possible to use HER3 expressing cells obtained by the genetic engineering or a cell line expressing HER3 as HER3 protein. Hereinbelow, a method for obtaining an antibody against HER3 is explained specifically.

(1) Preparation of antigen

[0093] Examples of the antigen to be used for producing the anti HER3 antibody include HER3, a polypeptide consisting of a partial amino acid sequence comprising at least 6 consecutive amino acids of HER3, and a derivative obtained by adding a given amino acid sequence or carrier thereto.

[0094] HER3 can be purified directly from human tumor tissues or tumor cells and used. Further, HER3 can be obtained by synthesizing it in vitro or by producing it in a host cell by genetic engineering.

[0095] With respect to the genetic engineering, specifically, after HER3 cDNA is integrated into a vector capable of expressing HER3 cDNA, HER3 can be obtained by synthesizing it in a solution containing an enzyme, a substrate and an energy substance required for transcription and translation, or by expressing HER3 in another prokaryotic or eucaryotic transformed host cell.

[0096] Further, the antigen can also be obtained as a secretory protein by expressing a fusion protein obtained by ligating the extracellular domain of HER3, which is a membrane protein, to the constant region of an antibody in an appropriate host-vector system.

[0097] HER3 cDNA can be obtained by, for example, a so-called PCR method in which a polymerase chain reaction (referred to as "PCR"; see Saiki, R. K., et al., Science, (1988) 239, pp. 487-489) is performed using a cDNA library expressing HER3 cDNA as a template and primers which specifically amplify HER3 cDNA.

[0098] As the in vitro synthesis of the polypeptide, for example, Rapid Translation System (RTS) manufactured by Roche Diagnostics, Inc. can be exemplified, but it is not limited thereto.

[0099] Examples of the prokaryotic host cells include Escherichia coli and Bacillus subtilis. In order to transform the host cells with a target gene, the host cells are transformed by a plasmid vector comprising a replicon, i.e., a replication origin derived from a species compatible with the host, and a regulatory sequence. Further, the vector preferably has a sequence capable of imposing phenotypic selectivity on the transformed cell.

[0100] Examples of the eucaryotic host cells include vertebrate cells, insect cells, and yeast cells. As the vertebrate cells, for example, simian COS cells (Gluzman, Y., Cell, (1981) 23, pp. 175-182, ATCC CRL-1650; ATCC: American Type Culture Collection), murine fibroblasts NIH3T3 (ATCC No. CRL-1658), and dihydrofolate reductase-deficient strains (Urlaub, G. and Chasin, L. A., Proc. Natl. Acad. Sci. USA (1980) 77, pp. 4126-4220) of Chinese hamster ovarian cells (CHO cells; ATCC: CCL-61); and the like are often used, however, the cells are not limited thereto.

[0101] The thus obtained transformant can be cultured according to a method usually carried out in the art, and by the culturing of the transformant, a target polypeptide is produced intracellularly or extracellularly.

[0102] A suitable medium to be used for the culturing can be selected from various commonly used culture media depending on the employed host cells. If Escherichia coli is employed, for example, an LB medium supplemented with an antibiotic such as ampicillin or IPMG as needed can be used.

[0103] A recombinant protein produced intracellularly or extracellularly by the transformant through such culturing can be separated and purified by any of various known separation methods utilizing the physical or chemical property of the protein.

[0104] Specific examples of the methods include treatment with a common protein precipitant, ultrafiltration, various types of liquid chromatography such as molecular sieve chromatography (gel filtration), adsorption chromatography, ion exchange chromatography, and affinity chromatography, dialysis, and a combination thereof.

[0105] Further, by attaching a tag of six histidine residues to a recombinant protein to be expressed, the protein can be efficiently purified with a nickel affinity column. Alternatively, by attaching the IgG Fc region to a recombinant protein to be expressed, the protein can be efficiently purified with a protein A column.

[0106] By combining the above-described methods, a large amount of a target polypeptide can be easily produced in high yield and high purity.

[0107] It is also possible to use the aforementioned transformant itself as an antigen. It is also possible to use a cell line expressing HER3 as an antigen. Examples of the cell line include. However, as long as HER3 is expressed, it is not limited to those cell lines.

(2) Production of anti HER3 monoclonal antibody

[0108] Examples of the antibody specific binding to HER3 include a monoclonal antibody specific binding to HER3, and a method of obtaining the antibody is as described below.

[0109] The production of a monoclonal antibody generally requires the following operational steps of:

- (a) Purification of a biopolymer used as an antigen or preparation of cells expressing antigen;
- (b) preparing antibody-producing cells by immunizing an animal by injection of the antigen, collecting the blood, assaying its antibody titer to determine when the spleen is excised;
- (c) preparing myeloma cells (hereinafter referred to as "myeloma");
- (d) fusing the antibody-producing cells with the myeloma;
- (e) screening a group of hybridomas producing a desired antibody;
- (f) dividing the hybridomas into single cell clones (cloning);
- (g) optionally, culturing the hybridoma or rearing an animal implanted with the hybridoma for producing a large amount of a monoclonal antibody;
- (h) examining the thus produced monoclonal antibody for biological activity and binding specificity, or assaying the same for properties as a labeled reagent; and the like.

[0110] Hereinafter, the method of producing a monoclonal antibody will be described in detail following the above steps, however, the method is not limited thereto, and, for example, antibody-producing cells other than spleen cells and myeloma can be used.

5 (a) Purification of antigen

[0111] As the antigen, HER3 prepared by the method as described above or a partial peptide thereof can be used.

[0112] Further, a membrane fraction prepared from recombinant cells expressing HER3 or the recombinant cells expressing HER3 themselves, and also a partial peptide of the protein of the invention chemically synthesized by a method
10 known to those skilled in the art can also be used as the antigen.

[0113] Further, a cell line expressing HER3 can be also used as an antigen.

(b) Preparation of antibody-producing cells

[0114] The antigen obtained in the step (a) is mixed with an adjuvant such as Freund's complete or incomplete adjuvant or aluminum potassium sulfate and the resulting mixture is used as an immunogen to immunize an experimental animal. In an alternative method, a test animal is immunized with cells expressing antigen as an immunogen. As the experimental animal, any animal used in a known hybridoma production method can be used without any trouble. Specifically, for example, a mouse, a rat, a goat, sheep, cattle, a horse, or the like can be used. However, from the viewpoint of ease of
20 availability of myeloma cells to be fused with the extracted antibody-producing cells, a mouse or a rat is preferably used as the animal to be immunized.

[0115] Further, the strain of a mouse or a rat to be used is not particularly limited, and in the case of a mouse, for example, various strains such as A, AKR, BALB/c, BDP, BA, CE, C3H, 57BL, C57BL, C57L, DBA, FL, HTH, HT1, LP, NZB, NZW, RF, R III, SJL, SWR, WB, and 129 and the like can be used, and in the case of a rat, for example, Wistar, Low, Lewis, Sprague,
25 Dawley, ACI, BN, Fischer and the like can be used.

[0116] These mice and rats are commercially available from breeders/distributors of experimental animals, for example, CLEA Japan, Inc. and Charles River Laboratories Japan, Inc.

[0117] As the animal to be immunized, in consideration of compatibility of fusing with myeloma cells described below, in the case of a mouse, BALB/c strain, and in the case of a rat, Wistar and Low strains are particularly preferred.

[0118] Further, in consideration of antigenic homology between humans and mice, it is also preferred to use a mouse having decreased biological function to remove auto-antibodies, that is, a mouse with an autoimmune disease.

[0119] The age of such mouse or rat at the time of immunization is preferably 5 to 12 weeks of age, more preferably 6 to 8 weeks of age.

[0120] In order to immunize an animal with HER3 or a recombinant thereof, for example, a known method described in detail in, for example, Weir, D. M., Handbook of Experimental Immunology Vol. I. II. III., Blackwell Scientific Publications, Oxford (1987); Kabat, E. A. and Mayer, M. M., Experimental Immunochemistry, Charles C Thomas Publisher Springfield, Illinois (1964) or the like can be used.

[0121] Among these immunization methods, a preferred specific method in the invention is, for example, as follows.

[0122] That is, first, a membrane protein fraction serving as the antigen or cells caused to express the antigen is/are intradermally or intraperitoneally administrated to an animal. However, the combination of both routes of administration is preferred for increasing the immunization efficiency, and when intradermal administration is performed in the first half and intraperitoneal administration is performed in the latter half or only at the last dosing, the immunization efficiency can be particularly increased.

[0123] The administration schedule of the antigen varies depending on the type of animal to be immunized, individual difference or the like. However, in general, an administration schedule in which the frequency of administration of the antigen is 3 to 6 times and the dosing interval is 2 to 6 weeks is preferred, and an administration schedule in which the frequency of administration of the antigen is 3 to 4 times and the dosing interval is 2 to 4 weeks is more preferred.

[0124] Further, the dose of the antigen varies depending on the type of animal, individual differences or the like, however, the dose is generally set to 0.05 to 5 mg, preferably about 0.1 to 0.5 mg.

[0125] A booster immunization is performed 1 to 6 weeks, preferably 1 to 4 weeks, more preferably 1 to 3 weeks after the administration of the antigen as described above. When the immunogen is a cell, 1×10^6 to 1×10^7 cells are used.

[0126] The dose of the antigen at the time of performing the booster immunization varies depending on the type or size of animal or the like, however, in the case of, for example, a mouse, the dose is generally set to 0.05 to 5 mg, preferably 0.1 to 0.5 mg, more preferably about 0.1 to 0.2 mg. When the immunogen is a cell, 1×10^6 to 1×10^7 cells are used.

[0127] Spleen cells or lymphocytes including antibody-producing cells are aseptically removed from the immunized animal 1 to 10 days, preferably 2 to 5 days, more preferably 2 to 3 days after the booster immunization. At this time, the antibody titer is measured, and if an animal having a sufficiently increased antibody titer is used as a supply source of the antibody-producing cells, the subsequent procedure can be carried out more efficiently.

[0128] Examples of the method of measuring the antibody titer to be used here include an RIA method and an ELISA method, but the method is not limited thereto. For example, if an ELISA method is employed, the measurement of the antibody titer in the invention can be carried out according to the procedures as described below.

[0129] First, a purified or partially purified antigen is adsorbed to the surface of a solid phase such as a 96-well plate for ELISA, and the surface of the solid phase having no antigen adsorbed thereto is covered with a protein unrelated to the antigen such as bovine serum albumin (hereinafter referred to as "BSA"). After washing the surface, the surface is brought into contact with a serially-diluted sample (for example, mouse serum) as a primary antibody to allow the antibody in the sample to bind to the antigen.

[0130] Further, as a secondary antibody, an antibody labeled with an enzyme against a mouse antibody is added and is allowed to bind to the mouse antibody. After washing, a substrate for the enzyme is added and a change in absorbance which occurs due to color development induced by degradation of the substrate or the like is measured and the antibody titer is calculated based on the measurement.

[0131] The separation of the antibody-producing cells from the spleen cells or lymphocytes of the immunized animal can be carried out according to a known method (for example, Kohler et al., *Nature* (1975), 256, p. 495; Kohler et al., *Eur. J. Immunol.* (1977), 6, p. 511; Milstein et al., *Nature* (1977), 266, p. 550; Walsh, *Nature* (1977), 266, p. 495). For example, in the case of spleen cells, a general method in which the antibody-producing cells are separated by homogenizing the spleen to yield the cells through filtration with a stainless steel mesh and suspending the cells in Eagle's Minimum Essential Medium (MEM) can be employed.

(c) Preparation of myeloma cells (hereinafter referred to as "myeloma")

[0132] The myeloma cells to be used for cell fusion are not particularly limited and suitable cells can be selected from known cell lines. However, in consideration of convenience when a hybridoma is selected from fused cells, it is preferred to use an HGPRT (hypoxanthine-guanine phosphoribosyl transferase) deficient strain whose selection procedure has been established.

[0133] More specifically, examples of the HGPRT-deficient strain include X63-Ag8(X63), NS1-ANS/1(NS1), P3X63-Ag8.U1(P3U1), X63-Ag8.653(X63.653), SP2/0-Ag14(SP2/0), MPC11-45.6TG1.7(45.6TG), FO, S149/5XXO, and BU.1 derived from mice; 210.RSY3.Ag.1.2.3(Y3) derived from rats; and U266AR(SKO-007), GM1500 GTG-A12(GM1500), UC729-6, LICR-LOW-HMy2(HMy2) and 8226AR/NIP4-1(NP41) derived from humans. These HGPRT-deficient strains are available from, for example, ATCC or the like.

[0134] These cell strains are subcultured in an appropriate medium such as an 8-azaguanine medium [a medium obtained by adding 8-azaguanine to an RPMI 1640 medium supplemented with glutamine, 2-mercaptoethanol, gentamicin, and fetal calf serum (hereinafter referred to as "FBS")], Iscove's Modified Dulbecco's Medium; IMDM), or Dulbecco's Modified Eagle Medium (hereinafter referred to as "DMEM"). In this case, 3 to 4 days before performing cell fusion, the cells are subcultured in a normal medium [for example, an ASF104 medium (manufactured by Ajinomoto Co., Ltd.) containing 10% FCS] to ensure not less than 2×10^7 cells on the day of cell fusion.

(d) Cell fusion

[0135] Fusion between the antibody-producing cells and the myeloma cells can be appropriately performed according to a known method (Weir, D. M. *Handbook of Experimental Immunology* Vol. I. II. III., Blackwell Scientific Publications, Oxford (1987); Kabat, E. A. and Mayer, M. M., *Experimental Immunochemistry*, Charles C Thomas Publisher, Springfield, Illinois (1964), etc.), under conditions such that the survival rate of cells is not excessively reduced.

[0136] As such a method, for example, a chemical method in which the antibody-producing cells and the myeloma cells are mixed in a solution containing a polymer such as polyethylene glycol at a high concentration, a physical method using electric stimulation, or the like can be used. Among these methods, a specific example of the chemical method is as described below.

[0137] That is, in the case where polyethylene glycol is used in the solution containing a polymer at a high concentration, the antibody-producing cells and the myeloma cells are mixed in a solution of polyethylene glycol having a molecular weight of 1500 to 6000, more preferably 2000 to 4000 at a temperature of from 30 to 40°C, preferably from 35 to 38°C for 1 to 10 minutes, preferably 5 to 8 minutes.

(e) Selection of a group of hybridomas

[0138] The method of selecting hybridomas obtained by the above-described cell fusion is not particularly limited. Usually, an HAT (hypoxanthine, aminopterin, thymidine) selection method (Kohler et al., *Nature* (1975), 256, p. 495; Milstein et al., *Nature* (1977), 266, p. 550) is used.

[0139] This method is effective when hybridomas are obtained using the myeloma cells of an HGPRT-deficient strain

which cannot survive in the presence of aminopterin. That is, by culturing unfused cells and hybridomas in an HAT medium, only hybridomas resistant to aminopterin are selectively allowed to survive and proliferate.

(f) Division into single cell clone (cloning)

[0140] As a cloning method for hybridomas, a known method such as a methylcellulose method, a soft agarose method, or a limiting dilution method can be used (see, for example, Barbara, B. M. and Stanley, M. S.: Selected Methods in Cellular Immunology, W. H. Freeman and Company, San Francisco (1980)). Among these methods, particularly, a three-dimensional culture method such as a methylcellulose method is preferred. For example, the group of hybridomas produced by cell fusion are suspended in a methylcellulose medium such as ClonaCell-HY Selection Medium D (manufactured by StemCell Technologies, inc., #03804) and cultured. Then, the formed hybridoma colonies are collected, whereby monoclonal hybridomas can be obtained. The collected respective hybridoma colonies are cultured, and a hybridoma which has been confirmed to have a stable antibody titer in an obtained hybridoma culture supernatant is selected as an anti-HER3 monoclonal antibody-producing hybridoma strain.

(g) Preparation of monoclonal antibody by culturing hybridoma

[0141] By culturing the thus selected hybridoma, a monoclonal antibody can be efficiently obtained. However, prior to culturing, it is preferred to perform screening of a hybridoma which produces a target monoclonal antibody.

[0142] In such screening, a known method can be employed.

[0143] The measurement of the antibody titer in the invention can be carried out by, for example, an ELISA method explained in item (b) described above.

[0144] The hybridoma obtained by the method described above can be stored in a frozen state in liquid nitrogen or in a freezer at -80°C or below.

[0145] After completion of cloning, the medium is changed from an HT medium to a normal medium, and the hybridoma is cultured.

[0146] Large-scale culture is performed by rotation culture using a large culture bottle or by spinner culture. From the supernatant obtained by the large-scale culture, a monoclonal antibody which specifically binds to the protein of the invention can be obtained by purification using a method known to those skilled in the art such as gel filtration.

[0147] Further, the hybridoma is injected into the abdominal cavity of a mouse of the same strain as the hybridoma (for example, the above-described BALB/c) or a Nu/Nu mouse to proliferate the hybridoma, whereby the ascites containing a large amount of the monoclonal antibody of the invention can be obtained.

[0148] In the case where the hybridoma is administrated in the abdominal cavity, if a mineral oil such as 2,6,10,14-tetramethyl pentadecane (pristane) is administrated 3 to 7 days prior thereto, a larger amount of the ascites can be obtained.

[0149] For example, an immunosuppressant is previously injected into the abdominal cavity of a mouse of the same strain as the hybridoma to inactivate T cells. 20 days thereafter, 10^6 to 10^7 hybridoma clone cells are suspended in a serum-free medium (0.5 mL), and the suspension is administrated in the abdominal cavity of the mouse. In general, when the abdomen is expanded and filled with the ascites, the ascites is collected from the mouse. By this method, the monoclonal antibody can be obtained at a concentration which is about 100 times or much higher than that in the culture solution.

[0150] The monoclonal antibody obtained by the above-described method can be purified by a method described in, for example, Weir, D. M.: Handbook of Experimental Immunology Vol. I, II, III, Blackwell Scientific Publications, Oxford (1978).

[0151] The thus obtained monoclonal antibody has high antigen specificity for HER3.

(h) Assay of monoclonal antibody

[0152] The isotype and subclass of the thus obtained monoclonal antibody can be determined as follows.

[0153] First, examples of the identification method include an Ouchterlony method, an ELISA method, and an RIA method.

[0154] An Ouchterlony method is simple, but when the concentration of the monoclonal antibody is low, a condensation operation is required.

[0155] On the other hand, when an ELISA method or an RIA method is used, by directly reacting the culture supernatant with an antigen-adsorbed solid phase and using antibodies corresponding to various types of immunoglobulin isotypes and subclasses as secondary antibodies, the isotype and subclass of the monoclonal antibody can be identified.

[0156] In addition, as a simpler method, a commercially available identification kit (for example, Mouse Typer Kit manufactured by Bio-Rad Laboratories, Inc.) or the like can also be used.

[0157] Further, the quantitative determination of a protein can be performed by the Folin Lowry method and a method of calculation based on the absorbance at 280 nm [$1.4 \text{ (OD 280)} = \text{Immunoglobulin } 1 \text{ mg/mL}$].

[0158] Further, even when the monoclonal antibody is separately and independently obtained by performing again the steps of (a) to (h) in (2), it is possible to yield an antibody having a cytotoxic activity equivalent to that of the anti-HER3 antibody. As one example of such an antibody, an antibody which binds to the same epitope as the anti-HER3 antibody can be exemplified. If a newly produced monoclonal antibody binds to a partial peptide or a partial tertiary structure to which the anti-HER3 antibody binds, it can be determined that the monoclonal antibody binds to the same epitope as the anti-HER3 antibody. Further, by confirming the competition by the monoclonal antibody for binding of the anti-HER3 antibody to HER3 (binding between the anti-HER3 antibody and HER3 is interfered by the monoclonal antibody), it can be determined that the monoclonal antibody binds to the same epitope as the anti-HER3 antibody even though a specific sequence or structure of the epitope has not been identified. Once the epitope is confirmed to be the same, it is strongly expected that the monoclonal antibody has the same antigen binding capacity or biological activity as the anti-HER3 antibody.

(3) Other antibodies

[0159] The antibody of the invention includes not only the above-described monoclonal antibody against HER3 but also a recombinant antibody obtained by artificial modification for the purpose of decreasing heterologous antigenicity to humans such as a chimeric antibody, a humanized antibody and a human antibody. These antibodies can be produced using a known method.

[0160] As the chimeric antibody, an antibody in which antibody variable and constant regions are derived from different species, for example, a chimeric antibody in which a mouse- or rat-derived antibody variable region is connected to a human-derived constant region can be exemplified (see Proc. Natl. Acad. Sci. USA, 81, 6851-6855, (1984)).

[0161] As the humanized antibody, an antibody obtained by integrating only a complementarity determining region (CDR) into a human-derived antibody (see Nature (1986) 321, pp. 522-525), and an antibody obtained by grafting a part of the amino acid residues of the framework as well as the CDR sequence to a human antibody by a CDR-grafting method (WO 90/07861) can be exemplified.

[0162] The term "several" as used herein refers to 1 to 10, 1 to 9, 1 to 8, 1 to 7, 1 to 6, 1 to 5, 1 to 4, 1 to 3, or 1 or 2.

[0163] In accordance with the present invention, it is to be understood, that the amino acid sequence of the binding protein of the invention is not limited to the twenty conventional amino acids (See Immunology - A Synthesis (2nd Edition, E. S. Golub and D. R. Gren, Eds., Sinauer Associates, Sunderland, Mass. (1991)), which is incorporated herein by reference). For example, the amino acids may include stereoisomers (e.g. D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as alpha-,alpha-disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids. Examples of unconventional amino acids, which may also be suitable components for the binding protein of the invention, include: 4-hydroxyproline, gamma-carboxyglutamate, epsilon-N,N,N-trimethyllysine, epsilon-N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, sigma-N-methylarginine, and other similar amino acids and imino acids, e.g. 4-hydroxyproline.

[0164] As the amino acid substitution in this specification, a conservative amino acid substitution is preferred. The conservative amino acid substitution refers to a substitution occurring within a group of amino acids related to amino acid side chains. Preferred amino acid groups are as follows: an acidic group (aspartic acid and glutamic acid); a basic group (lysine, arginine, and histidine); a non-polar group (alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, and tryptophan); and an uncharged polar family (glycine, asparagine, glutamine, cysteine, serine, threonine, and tyrosine). More preferred amino acid groups are as follows: an aliphatic hydroxy group (serine and threonine); an amide-containing group (asparagine and glutamine); an aliphatic group (alanine, valine, leucine, and isoleucine); and an aromatic group (phenylalanine, tryptophan, and tyrosine). Such an amino acid substitution is preferably performed within a range which does not impair the properties of a substance having the original amino acid sequence. When the heavy and light chains of the antibody of the present invention have glutamate as the N-terminal amino acid, it may be cyclized (in the form of pyroglutamate). In the present invention, such pyroglutamate is not differentiated from normal glutamine on amino acid sequences. In the heavy and light chains of the antibody of the present invention, cysteine may be in the form of cysteinyl. In the present invention, such a cysteinyl form is not differentiated from normal cysteine on amino acid sequences.

[0165] Further, the antibody of the invention includes a human antibody which binds to the HER3. An anti HER3 human antibody refers to a human antibody having only a gene sequence of an antibody derived from a human chromosome. The anti HER3 human antibody can be obtained by a method using a human antibody-producing mouse having a human chromosome fragment comprising heavy and light chain genes of a human antibody (see Tomizuka, K. et al., Nature Genetics (1997) 16, pp. 133-143; Kuroiwa, Y. et al., Nucl. Acids Res. (1998) 26, pp. 3447-3448; Yoshida, H. et al., Animal Cell Technology: Basic and Applied Aspects vol. 10, pp. 69-73 (Kitagawa, Y., Matuda, T. and Iijima, S. eds.), Kluwer Academic Publishers, 1999; Tomizuka, K. et al., Proc. Natl. Acad. Sci. USA (2000) 97, pp. 722-727, etc.).

[0166] Such a human antibody-producing mouse can be created specifically as follows. A genetically modified animal in which endogenous immunoglobulin heavy and light chain gene loci have been disrupted, and instead, human immunoglobulin heavy and light chain gene loci have been introduced via a yeast artificial chromosome (YAC) vector or the like

is created by producing a knockout animal and a transgenic animal and mating these animals.

[0167] Further, according to a recombinant DNA technique, by using cDNAs encoding each of such a heavy chain and a light chain of a human antibody, and preferably a vector comprising such cDNAs, eukaryotic cells are transformed, and a transformant cell which produces a recombinant human monoclonal antibody is cultured, whereby the antibody can also be obtained from the culture supernatant.

[0168] Here, as the host, for example, eukaryotic cells, preferably mammalian cells such as CHO cells, lymphocytes, or myeloma cells can be used.

[0169] With regard to preparation of a human antibody, detailed descriptions are given in International Publication No. WO 2007/077028. The contents of International Publication No. WO 2007/077028 are incorporated herein by reference.

[0170] Further, a method of obtaining a phage display-derived human antibody selected from a human antibody library (see Wormstone, I. M. et al., *Investigative Ophthalmology & Visual Science*. (2002) 43 (7), pp. 2301-2308; Carmen, S. et al., *Briefings in Functional Genomics and Proteomics* (2002), 1 (2), pp. 189-203; Siriwardena, D. et al., *Ophthalmology* (2002) 109 (3), pp. 427-431, etc.) is also known.

[0171] For example, a phage display method in which a variable region of a human antibody is expressed on the surface of a phage as a single-chain antibody (scFv), and a phage which binds to an antigen is selected (*Nature Biotechnology* (2005), 23, (9), pp. 1105-1116) can be used.

[0172] By analyzing the gene of the phage selected based on the binding to an antigen, a DNA sequence encoding the variable region of a human antibody which binds to an antigen can be determined.

[0173] If the DNA sequence of scFv which binds to an antigen is determined, a human antibody can be obtained by preparing an expression vector comprising the sequence and introducing the vector into an appropriate host to express it (WO 92/01047, WO 92/20791, WO 93/06213, WO 93/11236, WO 93/19172, WO 95/01438, WO 95/15388; *Annu. Rev. Immunol.* (1994) 12, pp. 433-455; *Nature Biotechnology* (2005) 23 (9), pp. 1105-1116).

[0174] One aspect of the present invention relates to an isolated protein that binds to HER3. In one embodiment of the present invention, an isolated HER3-binding protein of the invention comprises a heavy chain variable region amino acid sequence comprising: (a) CDRH1 comprised in the amino acid sequence represented by SEQ ID NO: 2, 6, 10, 14, 18, 22, 26, 30, 34, 36, 40, 42, 46, 50, 54, 60, 62, 66, 70, 74, 78, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 122, 126, 130, 134, 138, 142, 146, 150, 154, 158, 162, 166, 170, 174, 178, 182, 186, 190, 194, 198, 202, 206, 210, 214, 218, 222, 226 or 230, (b) CDRH2 comprised in the amino acid sequence represented by SEQ ID NO: 2, 6, 10, 14, 18, 22, 26, 30, 34, 36, 40, 42, 46, 50, 54, 60, 62, 66, 70, 74, 78, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 122, 126, 130, 134, 138, 142, 146, 150, 154, 158, 162, 166, 170, 174, 178, 182, 186, 190, 194, 198, 202, 206, 210, 214, 218, 222, 226 or 230, and (c) CDRH3 comprised in the amino acid sequence represented by SEQ ID NO: 2, 6, 10, 14, 18, 22, 26, 30, 34, 36, 40, 42, 46, 50, 54, 60, 62, 66, 70, 74, 78, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 122, 126, 130, 134, 138, 142, 146, 150, 154, 158, 162, 166, 170, 174, 178, 182, 186, 190, 194, 198, 202, 206, 210, 214, 218, 222, 226 or 230, and a light chain variable region amino acid sequence comprising: (d) CDRL1 comprised in the amino acid sequence represented by SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 38, 44, 48, 52, 56, 58, 64, 68, 72, 76, 82, 86, 90, 94, 98, 102, 106, 110, 114, 118, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228 or 232, (e) CDRL2 comprised in the amino acid sequence represented by SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 38, 44, 48, 52, 56, 58, 64, 68, 72, 76, 82, 86, 90, 94, 98, 102, 106, 110, 114, 118, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228 or 232, and (f) CDRL3 comprised in the amino acid sequence represented by SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 38, 44, 48, 52, 56, 58, 64, 68, 72, 76, 82, 86, 90, 94, 98, 102, 106, 110, 114, 118, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228 or 232.

[0175] The isolated HER3-binding protein of the present invention preferably comprises a heavy chain amino acid sequence comprising (a) CDRH1 comprising the amino acid sequence represented by one selected from the group consisting of SEQ ID NOs: 236, 251, 252, and 256; (b) CDRH2 comprising the amino acid sequence represented by one selected from the group consisting of SEQ ID NOs: 258, 278, 280, and 282; and (c) CDRH3 comprising the amino acid sequence represented by one selected from the group consisting of SEQ ID NOs: 283, 285, 309, 313, and 315, and a light chain amino acid sequence comprising (d) CDRL1 comprising the amino acid sequence represented by one selected from the group consisting of SEQ ID NOs: 320, 334, 337, and 340; (e) CDRL2 comprising the amino acid sequence represented by one selected from the group consisting of SEQ ID NOs: 343, 356, 351, and 344; and (f) CDRL3 comprising the amino acid sequence represented by one selected from the group consisting of SEQ ID NOs: 360, 381, 385, and 387.

[0176] In another embodiment of the present invention, an isolated binding protein of the invention comprises a heavy chain variable region amino acid sequence selected from the group consisting of SEQ ID Nos: 2, 6, 10, 14, 18, 22, 26, 30, 34, 36, 40, 42, 46, 50, 54, 60, 62, 66, 70, 74, 78, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 122, 126, 130, 134, 138, 142, 146, 150, 154, 158, 162, 166, 170, 174, 178, 182, 186, 190, 194, 198, 202, 206, 210, 214, 218, 222, 226 and 230, and/or a light chain variable region amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 38, 44, 48, 52, 56, 58, 64, 68, 72, 76, 82, 86, 90, 94, 98, 102, 106, 110, 114, 118, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228 and 232.

[0177] In yet another embodiment of the present invention, an isolated binding protein of the invention comprises a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 2 and 4, 6 and 8, 10 and 12, 14 and 16, 18 and 20, 22 and 24, 26 and 28, 30 and 32, 36 and 38, 42 and 44, 46 and 48, 50 and 52, 54 and 56, 60 and 58, 62 and 64, 66 and 68, 70 and 72, 74 and 76, 78 and 82, 80 and 82, 84 and 86, 88 and 90, 92 and 94, 96 and 98, 100 and 102, 104 and 106, 108 and 110, 112 and 114, 116 and 118, 122 and 124, 126 and 128, 130 and 132, 134 and 136, 138 and 140, 142 and 144, 146 and 148, 150 and 152, 154 and 156, 158 and 160, 162 and 164, 166 and 168, 170 and 172, 174 and 176, 178 and 180, 182 and 184, 186 and 188, 190 and 192, 194 and 196, 198 and 200, 202 and 204, 206 and 208, 210 and 212, 214 and 216, 218 and 220, 222 and 224, 226 and 228 or 230 and 232, or, a heavy chain variable region amino acid sequence represented by SEQ ID NO: 34, 40, 60, 62 or 120 and a light chain variable region amino acid sequence represented by SEQ ID NO: 58 or 64, respectively.

[0178] The isolated HER3-binding protein of the present invention more preferably comprises a heavy chain variable region amino acid sequence represented by SEQ ID NO: 42, 54, 70, 92, or 96 and a light chain variable region amino acid sequence represented by SEQ ID NO: 44, 56, 72, 94, or 98.

[0179] An antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 2 and 4 is referred to as "U1-39", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 6 and 8 is referred to as "U1-40", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 10 and 12 is referred to as "U1-38", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 14 and 16 is referred to as "U1-41", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 18 and 20 is referred to as "U1 -42", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 22 and 24 is referred to as "U1-43", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 26 and 28 is referred to as "U1 -44", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 30 and 32 is referred to as "U1-45", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 36 and 38 is referred to as "U1-47", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 42 and 44 is referred to as "U1 -49", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 46 and 48 is referred to as "U1-50", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 50 and 52 is referred to as "U1-51", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 54 and 56 is referred to as "U1-53", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 60 and 58 is referred to as "U1-55", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 62 and 64 is referred to as "U1 -57", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 66 and 68 is referred to as "U1-58", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 70 and 72 is referred to as "U1 -59", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 74 and 76 is referred to as "U1-52", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 78 and 82 is referred to as "U1-61", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 80 and 82 is referred to as "U1 -61.1", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 84 and 86 is referred to as "U1-62", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 88 and 90 is referred to as "U1-2", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 92 and 94 is referred to as "U1-7", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 96 and 98 is referred to as "U1-9", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 100 and 102 is referred to as "U1-10", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 104 and 106 is referred to as "U1-12", an antibody comprising a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 108 and 110 is referred to as "U1-13", an antibody comprising a heavy

[illegible]

[0180] The isolated HER3-binding protein of the present invention even more preferably comprises a heavy chain

variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 42 and 44, respectively, a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 54 and 56, respectively, a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 70 and 72, respectively, a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 92 and 94, respectively, or a heavy chain variable region amino acid sequence and a light chain variable region amino acid sequence represented by SEQ ID NOs: 96 and 98, respectively, and still even more preferably, the HER3-binding proteins is U1-49, U1-53, U1-59, U1-7, or U1-9, which are an anti-HER3 antibody.

[Chem. 14]

Sequence Listing

Antibody U1-39

1 Heavy Chain DNA:

GAGGTGCAGCTGGTGGAGTCTGGAGGAGCCTTGATCCAGCCTGGGGGGTCCCTGAGACTC
 TCCTGTGCAGCCTCTGGGTTACCGTCAGTAGCAACTACATGAGCTGGGTCCGCCAGGCT
 CCAGGGAAGGGGCTGGATTGGGTCTCAGTTATTTATAGCGGTGGTAGCACATACTACGCA
 GACTCCGTGAAGGGCCGATTACCATCTCCAGAGACAATCCAAGAACACGCTGTATCTT
 CAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGTATTACTGTGCGAGAGGGCAGTGG
 CTGGACGTCTGGGGCCAAGGGACCACGGTCAACCGTCTCCTCA

2 Heavy Chain Protein:

EVQLVESGGGLIQPGGSLRLSCAASGFTVSSNYMSWVRQAPGKGLDWVSVIYSGGSTYYA
 DSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGQWLDVWGQGT TTVTVSS

3 Light Chain DNA:

GATATTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTGGAGAGCCGGCCTCC
 ATCTCCTGCAGGTCAAGTCAGAGCCTCCTGCATAGTAATGGATACAAC TATTGGATTGG
 TACCTGCAGAGGCCAGGGCAGTCTCCACAACTCCTGTTCTATTTGGGTTTTCATCGGGCC
 TCCGGGGTCCCTGACAGGTTCACTGGCAGTGGATCAGGCACAGATTTTACACTGAAAATC
 AGCAGAGTGGAGGCTGAGGATGTTGGGGTTTATTACTGCAGGCAAGCTCTACAACTCCG
 CTCAC TTTCCGGCGAGGGACCAAGGTGGAGATCAAA

4 Light Chain Protein:

DIVMTQSPLSLPVTPGEPASISCRSSQSLLSNGYNYLDWYLQRPQSPQLLFYLG FHRA
 SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCRQALQTF LTFGGGT KVEIK

Antibody U1-40

5 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGGCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGTACTGTCTCTGGTGGCTCCATCAGCAGTGGTGGTTACTACTGGAGCTGGATCCGC
 CAGCACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTCCAGTGGGAGCACCTAC
 TACAACCCGTCCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGAT
 AGGGA ACTGGA ACTTTACTACTACTACTACGGTATGGACGTCTGGGGCCAAGGGACCACG
 GTCACCGTCTCCTC

6 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQHPGKGLEWIGYIYSSGSTY
 YNP SLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDRELELYYYYYYGM DVWGQGT
 TVTS

7 Light Chain DNA:

GATATTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTGGAGAGCCGGCCTCC
 ATCTCCTGCAGGTCTAGTCAGAGCCTCCTGTATAGTAATGGATACAAC TATTGGATTGG
 TACCTGCAGAAGCCAGGGCAGTCTCCACAGCTCCTGATCTATTTGGGTTCTAATCGGGCC
 TCCGGGGTCCCTGACAGGTTCACTGGCAGTGGATCAGGCACAGATTTTACACTGAAAATC
 AGCAGAGTGGAGGCTGAGGATGTTGGGATTTATTACTGCATGCAAGCTCTACAACTCCG
 CTCAC TTTCCGGCGAGGGACCAAGGTGGAGATCAAA

8 Light Chain Protein:

DIVMTQSPLSLPVTPGEPASISCRSSQSLLSNGYNYLDWYLQKPGQSPQLLIYLG SNRA
 SGVPDRFSGSGSGTDFTLKISRVEAEDVGIYYCMQALQTF LTFGGGT KVEIK

Antibody U1-38

9 Heavy Chain DNA:

CAGATCACCTTGAAGGAGTCTGGTCC TACGCTGGTGA AACCACACAGACCCTCAGCTG
 ACCTGCACCTTCTCTGGGTTCTCACTCAGCACTAGTGGAGTGGGTGTGGCTGGATCCGT
 CAGCCCCCAGGAAAGGCCCTGGACTGGCTTGCACTCATTTATTGGAATGATGATAAGCGC

TACAGCCCATCTCTGAAGAGCAGGCTCACCATCACCAGGACACCTCCAAAAACCAGGTG
 GTCCTTACAATGACCAACATGGATCTTGTGGACACAGCCACATATTACTGTGTACACAGA
 GACCAAGTTCGAGGGTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCCTCA

10 Heavy Chain Protein:

QITLKESGPTLVKPTQTLTCTFSGFSLSTSGVGVGWIRQPPGKALDWLALIYWNDDKR
 YSPSLKSRLTITKDTSKNQVVLMTNMDLVDATYYCVHRDEVRFDYWGQGLVTVSS

11 Light Chain DNA:

GATGTTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTTGGACAGCCGGCCTCC
 ATCTCCTGCAGGTCTAGTCAAAGCCTCGTATACAGTGATGGATACACCTACTTGCATTGG
 TTTCAGCAGAGGCCAGGCCAATCTCCAAGGCGCCTTATTTATAAGGTTTCTAACTGGGAC
 TCTGGGGTCCCAGACAGATTTCAGCGGCAGTGGGTGAGGCACTGATTTCACACTGAAAATC
 AGCAGGGTGGAGGCTGAGGATGTTGGGGTTTATTACTGCATGCAAGGTGCACACTGGCCG
 ATCACCTTCGGCCAAGGGACACGACTGGAGATTAAA

12 Light Chain Protein:

DVVMTQSPSLPVTGLQBPASISCRSSQSLVYSDGYTYLHWFOQRPGQSPRRLIYKVSND
 SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQGAHWPIITFCQGRLEIK

Antibody U1-41

13 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGGCCAGGACTGGTGAAGCCTTCACAGACCCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGGGTACTACTGGAGCTGGATCCGC
 CAGCACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
 TACAACCCGTCCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTTCTGTGCGAGAGAT
 CGGGAACCTTGAGGGTTACTCCAATACTACGGTGTGGACGTCTGGGGCCAAGGGACACG
 GTCACCGTCTCCTC

14 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQHPGKGLEWIGYIYYSGSTY
 YNPSLKSRLTISVDTSKNQFSLKLSSVTAADTAVYFCARDRELEGYSNYYGVVDVWGQGT
 VTVS

15 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGGCCATTAGCAACTATTTAAATTGGTATCAGCAGAAACCA
 GGGAAAGCCCCCTAAGCTCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCAGCCT
 GAAGATTTTGCAACTTATTACTGTCAACAGAATAATAGTCTCCCGATCACCTTCGGCCAA
 GGGACACGACTGGAGATTAAA

16 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQAISNYLNWYQQKPGKAPKLLIYAASSLQSGVPS
 RFGSGSGTDFTLTISLQPEDFATYYCQQNNSLPITFCQGRLEIK

Antibody U1-42

17 Heavy Chain DNA:

GAGGTGCAGCTGGTGCAGTCTGGAGCAGAGGTGAAAAAGCCCGGGGAGTCTCTGAAGATC
 TCCTGTAAGGGTTCTGGATACAGCTTTACCAGCTACTGGATCGGCTGGGTGCGCCAGATG
 CCCGGGAAAGGCCTGGAGTGGATGGGGATCATCTATCCTGGTGAATCTGATACCAGATAC
 AGCCCGTCCCTCCAAGGCCAGGTCAACATCTCAGCCGACAAGTCCATCAGCACCAGCCTAC
 CTGCACTGGAGCAGCCTGAAGGCCTCGGACACCGCCATGTATTACTGTGCGAGACATGAA
 AACTACGGTGAATACAATACTGGGGCCAGGGAACCCTGGTCACCGTCTCCTCA

18 Heavy Chain Protein:

EVQLVQSGAEVKKPGESLKISCKSGSYSTSYWIGWVRQMPGKGLEWMGLIYPGDSDFRY
 SPSFQGGVTTISADKSIATAYLQWSSLKASDTAMYICARHENYGDYNYWGQGLVTVSS

19 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTGGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGAGCATTGCGAGCTATTTAAATTGGTATCAGCAGAAACCA
 GGGAAAGCCCCCTAAGCTCCTGATCTATGCTGCTTCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCT
 GAAGATTTTGCACCTTACTGCTGTCAACAGAGTAACGGTCCCCGCTCACTTTGCGCGGA
 GGGACCAAGGTGGAGATCAAA

20 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQSIRSYLNWYQQKPKAPKLLIYAASSLQSGVPS
 RFSGSGSGTDFTLTISLQPEDFALYCCQSQNSGSPFTFGGGTKVEIK

Antibody U1-43

21 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCCGGGCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGGTTACTACTGGAGCTGGATCCGC
 CAGCACCACAGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
 TACAACCCGTCCCTCAGGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACTGCCGCGACACGGCCGTGTATTACTGTGCGAGAGAT
 AGAGAGAGAGAGTGGGATGATTACGGTGACCCCCAAGGTATGGACGTCTGGGGCCAAGGG
 ACCACGGTCACCGTCTCCTC

22 Heavy Chain Protein:

QVQLQESGPEGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQHPGKGLEWIGYIYYSGSTY
 YNPSLRSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDREWDYDGPQGMVWGQG
 TTVTVS

23 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGAGCATTAGCAGCTATTTACATTGGTATCAGCAGAAACCA
 GGGAAAGCCCCCTAAGCTCCTGATCCATGCTGCATCCAGTTTACAAAGTGGGGTCCCATCA
 AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGTAGTCTGCAACCT
 GAAGATTTTGCAACTTACTACTGTCAACAGAGTTACAGTAACCCGCTCACTTTGCGCGGA
 GGGACCAAGGTGGAGATCCAA

24 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQSISSYLHWYQQKPKAPKLLIHAASSLQSGVPS
 RFSGSGSGTDFTLTISLQPEDFATYYCQSYSNPLTFGGGTKVEIQ

Antibody U1-44

25 Heavy Chain DNA:

GAGGTGCAGCTGGTGCAGTCTGGAGCAGAGGTGAAAAAGCCCGGGAGTCTCTGAAGATC
 TCCTGTAAGGGTTCTGGATACAGCTTTACCAGCTACTGGATCGGCTGGGTGCGCCAGATG
 CCCGGGAAAGGCCTGGAGTGGATGGGGATCATCTGGCCTGGTGACTCTGATACCATATAC
 AGCCCGTCCCTCCAAGGCCAGGTCAACATCTCAGCCGACAAGTCCATCAGCACCGCCTAC
 CTGCAGTGGAGCAGCCTGAAGGCCTCGGACACCGCCATGTATTACTGTGCGAGACATGAA
 AACTACGGTGACTACAACTACTGGGGCCAGGGAACCCTGGTCACCGTCTCCTCA

26 Heavy Chain Protein:

EVQLVQSGAEVKKPGESLKISKSGSYFTSYWIGWVRQMPGKGLEWMGIWPGDSDTIY
 SPSFQGGVTTISADKSISTAYLQWSSSLKASDTAMYICARHENYGDYNYWGQGLTVTVSS

27 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTGGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGAGCATTGGAAGTTATTTAAATTGGTATCAGCAGAAACCG
 GGGAAATGCCCTAAACTCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCT
 GAAGATTTTGCACCTTACTACTGTCAACAGAGTATCAGTTCCCCGCTCACTTTGCGCGGA
 GGGACCAAGGTGGAGATCAAA

28 Light Chain Protein:
 DIQMTQSPSSLSASVGDRTITCRASQSIRSYLNWYQQKPGNAPKLLIYAASSLQSGVPS
 RFGSGSGTDFTLTISLQPEDFALYYCQSSISPLTFGGGTKEIK

Antibody (U1-45)

29 Heavy Chain DNA:
 CAGGTGCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGAAGCCTGGGGCCTCAGTGAAGGTC
 TCCTGCAAGGCTTCTGGATACACCTTCACCAGTTATGATATCAACTGGGTGCGACAGGCC
 ACTGGACAAGGGCTTGAGTGGATGGGATGGATGAACCTAACAGTGGTGACACTGGCTAT
 GCACAGGTGTTCCAGGGCAGAGTCACCATGACCTGGAACACCTCCATAAGCACAGCCTAC
 ATGGAAGTGAAGCAGCCTGAGATCTGAGGACACGGCCGTCTATTACTGTGCGAGATTTGGG
 GATCTCCCGTATGACTACAGTTACTACGAATGGTTCGACCCCTGGGGCCAGGGAACCCCTG
 GTCACCGTCTCCTC

30 Heavy Chain Protein:
 QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYDINWVRQATGQGLEWMGWMNPNSGDTGY
 AQVFQGRVTMTWNTSISTAYMELSSLRSEDYAVYYCARFGDLPYDYSYYEWFDPWGQGT
 LTVTS

31 Light Chain DNA:
 GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGCCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAGACCA
 GGGAAAGCCCCCTAAGCTCCTGATCTATGCAGCATCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCT
 GAAGATTTTGCAACTTACTACTGTCAACAGAGTTACAGTACCCCGCTCACTTTCCGGCGGA
 GGGACCAAGGTGGAGATCAAA

32 Light Chain Protein:
 DIQMTQSPSSLSASVGDRTITCRASQSISSYLNWYQQRPGKAPKLLIYAASSLQSGVPS
 RFGSGSGTDFTLTISLQPEDFATYYCQSYSTPLTFGGGTKEIK

Antibody (U1-46)

33 Heavy Chain DNA:
 CAGGTACAGCTGCAGCAGTCAGGTCCAGGACTGGTGAAGCCCTCGCAGACCCTCTCACTC
 ACCTGTGCCATCTCCGGGGACAGTGTCTCTAGCAACAGTGCTGCTTGGAAGTGGATCAGG
 CAGTCCCCATCGAGAGGCCTTGAGTGGCTGGGAAGGACATACTACAGGTCCAAGTGGTAT
 AATGATTATGCAGTATCTGTGAAAAGTGAATAACCATCAACCCAGACACATCCAAGAAC
 CAGTTCTCCCTGCAGCTGAACTCTGTGACTCCCGAGGACACGGCTGTGTATTACTGTGCA
 AGAGATCTCTACGATTTTGGAGTGGTTATCCCTACTACTACGGTATGGACGTCTGGGGC
 CAAGGGACCACGGTCACCGTCTCCTC

34 Heavy Chain Protein:
 QVQLQQSGPGLVKPSQTLSTCAISGDSVSSNSAAWNWIQSPSRGLEWLGRTYYRSKWY
 NDYAVSVKSRITINPDTSKNQFSLQLNSVTPEDTAVYYCARDLYDFWSGYPPYYGMDVWG
 QGTTVTVTS

Antibody U1-47

35 Heavy Chain DNA:
 CAGGTACAGCTGCAGCAGTCAGGTCCAGGACTGGTGAAGCCCTCGCAGACCCTCTCACTC
 ACCTGTGCCATCTCCGGGGACAGTGTCTCTAGCAACAGTGCTGCTTGGAAGTGGATCAGG
 CAGTCCCCATCGAGAGGCCTTGAGTGGCTGGGAAGGACATACTACAGGTCCAAGTGGTAT
 AATGATTATGCAGTATCTGTGAAAAGTGAATAACCATCAACCCAGACACATCCAAGAAC
 CAGTTCTCCCTGCAGCTGAACTCTGTGACTCCCGAGGACACGGCTGTGTATTACTGTGCA
 AGAGATTACTATGGTTCGGGGAGTTTCTACTACTACTACGGTATGGACGTCTGGGGCCAA
 GGGACCACGGTCACCGTCTCCTC

36 Heavy Chain Protein:
 QVQLQQSGPGLVKPSQTLSTCAISGDSVSSNSAAWNWIQSPSRGLEWLGRTYYRSKWY

NDYAVSVKSRITINPDTSKNQFSLQLNSVTPEDTAVYYCARDYYGSGSFYYYYGMDVWGQ
GTTVTVS

37 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCGGGCAAGTCAGAGCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCA
GGGAAAGCCCCCTAAGGTCCCTGATCTATGCTGCATCCAATTTGCAAAGTGGGGTCCCATCA
AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCT
GAAGATTTTGCAACTTACTACTGTCAACAGAGTTACAGTACCCCTCGGACGTTCCGGCCAA
GGGACCAAGGTGGAAATCAAA

38 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQSISSYLNWYQQKPGKAPKVLIIAASNLQSGVPS
RFSGSGSGTDFTLTISLQPEDFATYYCQQSYSTPRTFGQGTKVEIK

Antibody U1-48

39 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGAGACCCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGTAGTTACTACTGGAGCTGGATCCGGCAGCCC
GCCGGAAGGGACTGGAGTGGATGGGCATATCTATACCAGTGGGAGCACCAACTACAAC
CCCTCCCTCAAGAGTCGAGTCACCATGTCAGTAGACACGTCCAAGAACCAGTTCTCCCTG
AAGCTGAGCTCTGTGACCGCCGCGACACGGCCGTGTATTACTGTGCGAGAGAAGCGATT
TTTGGAGTGGGCCCCCTACTACTACTACGGTATGGACGTCTGGGGCCAAGGGACCACGGTC
ACCGTCTCCTC

40 Heavy Chain Protein:

QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWSWIRQAGKGLEWIGHIYTSGSTNMYN
PSLKSRVTMSVDTSKNQFSLKLSSVTAADTAVYYCAREAI FGVGPYYYYGMDVWGQGT
TVS

Antibody U1-49

41 Heavy Chain DNA:

CAGGTGCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGAAGCCTGGGGCCTCAGTGAAGGTC
TCCTGCAAGGCTTCTGGATACACCTTCACCGCTACTATATGCACTGGGTGCGACAGGCC
CCTGGACAAGGGCTTGAGTGGATGGGATGGATCAACCCCTAATATTGGTGGCACAACTGT
GCACAGAAGTTTCAGGGCAGGGTCACCATGACCAGGGACACGTCCATCAGCACAGCCTAC
ATGGAGCTGAGCAGGCTGAGATCTGACGACACGGCCGTGTATTACTGTGCGAGAGGGGGA
CGGTATAGCAGCAGCTGGTCCCTACTACTACTACGGTATGGACGTCTGGGGCCAAGGGACC
ACGGTCACCGTCTCCTC

42 Heavy Chain Protein:

QVQLVQSGAEVKKPGASVKVSCKASGYTFTGYMHVVRQAPGQGLEWMGWINPNIGGTNC
AQKFQGRVTMTTRDTSISTAYMELSRRLRSDDTAVYYCARGGRYSSWSYYYYGMDVWGQGT
TVTVS

43 Light Chain DNA:

GATATTCTGATGACCCAGACTCCACTCTCTCTGTCCGTCACCCCTGGACAGCCGGCCTCC
ATCTCCTGCAAGTCTAGTCAGAGCCTCCTGCTTAGTGATGGAGGGACCTATTTGTATTGG
TACCTGCAGAAGCCAGGCCAGCCTCCACAGCTCCTGATCTATGAAGTTTCAACCGGTTT
TCTGGAGTGCCAGATAGGTTTCAGTGGCAGCGGTCAGGGACAGATTTCACTGAAAATC
AGCCGGGTGGAGGCTGAGGATGTTGGGGTTTATTACTGCATGCAAAGTATGCAGCTTCCG
ATCACCTTCGGCCAAGGGACACGACTGGAAATTAA

44 Light Chain Protein:

DILMTQTPLSLSVTPGQPASISCKSSQSLLSLSDGGTYLYWYLQKPGQPPQLLIYEVSNRF
SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQSMQLPITFGQGTRLKIK

Antibody U1-50

45 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCGTCAGCAGTGGTGGTTACTACTGGAGCTGGATCCGG
 CAGCCCCCAGGGAAGGGACTGGAGTGGATTGGGTATATCTATTACAGTGGGAGCACC AAC
 TACAACCCCTCCCTCAAGAGTCGAGTCACCATATCAGTAGACACGTCCAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACCGCTGCGGACACGGCCGTGTATTACTGTGCGAGAGGG
 GGGGACAGTAAC TACGAGGATTACTACTACTACTACGGTATGGACGTCTGGGGCCAAGGG
 ACCACGGTCACCGTCTCCTC

46 Heavy Chain Protein:

QVQLQESGPGLVKPSETLSLTCTVSGGSVSSGGYYWSWIRQPPGKGLEWIGYIYYSGSTN
 YNP SLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARGGDSNYEDYYYYYGMVWVGQG
 TTVTVS

47 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGAGCATTAGCATCTATTACATTGGTATCAGCAGAAACCA
 GGGAAAGCCCCCTAAGCTCTTGATCTCTGCTGCATCCAGTTTGCAAAGTGGGGTCCCGTCA
 AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGAAGTCTGCAACCT
 GAAGATTTTGCAACTTACTACTGTCAACAGAGTTACACTTCCCGATCACCTTCCGCCAA
 GGGACACGACTGGAGATTAA

48 Light Chain Protein:

DIQMTQSPSSLSASVGRVTITCRASQSSISYIYLHWYQKPGKAPKLLISAASSLQSGVPS
 RFGSGSGTDFTLTIRSLQPEDFATYYCQQSYTSPITFGQGRLEIK

49 Antibody U1-51

Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGTAGTTACTACTGGAGCTGGATCCGGCAGCCC
 CCAGGGAAGGGACTGGAGTGGATTGGGTATATCTATTACAGTGGGAGCACC AACTACAAC
 CCCTCCCTCAAGAGTCGAGTCACCATATCAGTAGACACGTCCAAGCACCAGTTCTCCCTG
 AAGCTGAGCTCTGTGACCGCTGCGGACACGGCCGTGTATTACTGTGCGAGAGATTGAGT
 TACTATGATAGTAGTGGTTATTACTTATACTACTACGCTATGGACGTCTGGGGCCAAGGG
 ACCACGGTCACCGTCTCCTC

50 Heavy Chain Protein:

QVQLQESGPGLVKPSETLSLTCTVSGGSISYIYWSWIRQPPGKGLEWIGYIYYSGSTNYN
 PSLKSRVTISVDTSKHQFSLKLSSVTAADTAVYYCARDSSYYDSSGYLYYYAMVWVGQG
 TTVTVS

51 Light Chain DNA:

GACATCGTGATGACCCAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGAGGGCCACC
 ATCAACTGCAAGTCCAGCCAGAGTGTTTTATACAGCTCCAACAATAAGAACTACTTAGCT
 TGGTACCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTCCCTGGGCATCTACCCGG
 GAATCCGGGGTCCCTGACCGATTTCAGTGGCAGCGGTCTGGGACAGATTTCACTCTCACC
 ATCAGCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTGACCAATATTATACTACT
 CCTCTCACTTTCCGGCCCTGGGACCAAAGTGCATATCAA

52 Light Chain Protein:

DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQKPGQP KLLISWASTR
 ESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCQQYYTTPLTFGPGTKVDIX

Antibody U1-53

53 Heavy Chain DNA:

GAGGTGCAACTGGTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGAGACTC
 TCCTGTGCAGCCTCTGGATTACCTTCAGTATCTATAGCATGAACTGGGTCCGCCAGGCT
 CCAGGGAAGGGGCTGGAGTGGGTTTCATACATTAGTAGTAGTAGTAGTACCATATACTAC
 GCAGACTCTGTGAAGGGCCGATTACCATCTCCAGAGACAATGCCAAGAACTCACTGTAT
 CTGCAAATGAACAGCCTGAGAGACGAGGACACGGCTGTGTATTACTGTGCGAGAGATAGG

CGTGACTTCGATGCTTTTGATATCTGGGGCCAAGGGACAATGGTCACCGTCTCTTCA
 54 Heavy Chain Protein:
 EVQLVESGGGLVQPGGSLRLSCAASGFTFSIYSMNWVRQAPGKGLEWVSYISSSSSSTIYY
 ADSVKGRFTISRDNAKNSLYLQMNSLRDEDTAVYYCARDRGDFDAFDIWGQGTMTVTVSS
 55 Light Chain DNA:
 GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCAGGCGAGTCAGGACATTACCAACTATTTGAATTGGTATCAGCAGAAACCA
 GGGAAAGCCCTAAGCTCCTGATCTACGATGCATCCAATTTGGAAACAGGGGTCCCATCA
 AGGTTCAGTGGAAGTGGATCTGGGACAGATTTTACTTTCACCATCAGCAGCCTGCAGCCT
 10 GAAGATATTGCAACATATAACTGTCAACAGTGTGAAAATTTCCCGATCACCTTCGGCCAA
 GGGACACGACTGGAGATTAA
 56 Light Chain Protein:
 DIQMTQSPSSLSASVGRVTITCQASQDITNYLWYQQKPKGKAPKLLIYDASNLETGVPS
 15 RFGSGSGTDFTFITISLQPEDATYNCQQCENFFITFGQGTREIK

Antibody U1-55

57 Light Chain DNA:
 GATATTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTGGAGAGCCGGCCCTCC
 20 ATCTCCTGCAGGTCTAGTCAGAGCCTCCTGTATAGTAATGGATACAAGTATTTGGATTGG
 TACCTGCAGAAGCCAGGGCAGTCTCCACAGCTCCTGATCTATTTGGGTCTAAATCGGGCC
 TCCGGGGTCCCTGACAGGTTCACTGGCAGTGGATCAGGCACAGATTTTACACTGAAAATC
 AGCAGAGTGGAGGCTGAGGATGTTGGGGTTTATTATTGCATGCAGGCTCTACAACTCCG
 25 ATCACCTTCGGCCAAGGGACACGACTGGAGATTAA
 58 Light Chain Protein:
 DIVMTQSPPLSLPVTPEPASISCRSSQSLLYSNGYKYLDWYLQKPGQSPQLLIYLGSNRA
 SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQALQTPITFGQGTREIK

Antibody (U1-55.1)

59 Heavy Chain DNA:
 CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTC
 30 ACCTGCACTGTCTCTGGTGGCTCCGTCAGCAGTGGTGGTTACTACTGGAACTGGATCCGG
 CAGCCCCCAGGGAAGGGACTGGAGTGGATTGGGTATATCAATTACAGTGGGAGCACCAAC
 35 TACAACCCCTCCCTCAAGAGTCGAGTCACCATATCAGTAGACACGTCCAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACCGCTGCGGACACGGCCGTGTATTACTGTGCGAGAGAT
 CGAGAACTGGAACCTTACTACTACTACTACGGTATGGACGTCTGGGGCCAAGGGACCACG
 40 GTCACCGTCTCCTC
 60 Heavy Chain Protein:
 QVQLQESGPGLVKPSSETLSLTCTVSGGSVSSGGYYWNWIRQPPGKLEWIGYINYSGSTN
 YNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDRELELYYYYYGMDVWGQGT
 TVTS

Antibody (U1-57)

61 Heavy Chain DNA:
 CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCTGAGACCCTGTCCCTC
 45 ACCTGCACTGTCTCTGGTGGCTCCGTCAGCAGTGGTGGTTACTACTGGAACTGGATCCGG
 CAGCCCCCAGGGAAGGGACTGGAGTGGATTGGGTATATCAATTACAGTGGGAGCACCAAC
 50 TACAACCCCTCCCTCAAGAGTCGAGTCACCATATCAGTAGACACGTCCAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACCGCTGCGGACACGGCCGTGTATTACTGTGCGAGAGAT
 CGAGAACTGGAACCTTACTACTACTACTACGGTATGGACGTCTGGGGCCAAGGGACCACG
 55 GTCACCGTCTCCTC
 62 Heavy Chain Protein:
 QVQLQESGPGLVKPSSETLSLTCTVSGGSVSSGGYYWNWIRQPPGKLEWIGYINYSGSTN
 YNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDRELELYYYYYGMDVWGQGT

Antibody U1-57.1

63 Light Chain DNA:

GATATTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTGGAGAGCCGGCCTCC
 ATCTCCTGCAGGTCTAGTCAGAGCCTCCTGTATAGTAATGGATACAAGTATTTGGATTGG
 TACCTGCAGAAGCCAGGGCAGTCTCCACAGCTCATGATCTATTTGGGTTCTAATCGGGCC
 TCCGGGGTCCCTGACAGGTTTCAGTGGCAGTGGATCAGGCACAGATTTTACACTGAAAATC
 AGCAGAGTGGAGGCTGAGGATGTTGGGGTTTATTATTGCATGCAGGCTCTACAACTCCG
 ATCACCTTCGGCCAAGGGACACGACTGGAGATTAAA

64 Light Chain Protein:

DIVMTQSPLSLPVTGPGEPAISCRSSQSLLYSNGYKYLDWYLQKPGQSPQLMIYLGSNRA
 SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQALQTPITFGQTRLEIK

Antibody U1-58

65 Heavy Chain DNA:

CAGGTGCAGCTGGTGGAGTCTGGGGGAGGCGTGGTCCAGCCTGGGAGGTCCCTGAGACTC
 TCCTGTGCAGCGTCTGGATTCACCTTCAGTAGCTATGGCATGCACTGGGTCCGCCAGGCT
 CCAGGCAAGGGGCTGGAGTGGGTGGCAGTTATATGGTATGATGGAAGTAATAAATACTAT
 GCAGACTCCGTGAAGGGCCGATTACCATCTCCAGAGACAATTCCAAGAACACGCTGTAT
 CTGCAAAATGAACAGCCTGAGAGCCGAGGACACGGCTGTGTATTACTGTGCGAGAGCAGCT
 CGCCTTGACTACTACTACGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCC
 TCA

66 Heavy Chain Protein:

QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYGMHWVRQAPGKGLEWVAVIWDGSGNKYY
 ADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARAARLDYVYGMDEVWGQGTITVTVS
 S

67 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCTCC
 ATCACTTGCCCGGCAAGTCAGAGCATTAACAGCTATTTAAATTGGTTTCAGCAGAAGCCA
 GGGAAAGCCCCCTCAGCTCCTGATCTTTGGTGCATCCGGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTCACTCTCAACATCAACAGTCTGCAACCT
 GAAGATTTTGCAACTTACTACTGTCAACAGAGTTACAGTTCCCCGCTCACCTTCGGCCAA
 GGGACACGACTGGAGATTAAA

68 Light Chain Protein:

DIQMTQSPSSLSASVGRVSIICRASQSINSYLNWFQKPKAPQLLIIFGASGLQSGVPS
 RFGSGSGSGTDFTLTINSIQPEDFATYYCQSYSSPLTFGQGTTRLEIK

Antibody U1-59

69 Heavy Chain DNA:

CAGGTGCAGCTACAGCAGTGGGGCGCAGGACTGTTGAAGCCTTCGGAGACCCTGTCCCTC
 ACCTGCGCTGTCTATGGTGGGTCTTCAGTGGTTACTACTGGAGCTGGATCCGCCAGCCC
 CCAGGGAAGGGGCTGGAGTGGATTGGGGAAATCAATCATAGTGGGAAGCACCACACTACAAC
 CCGTCCCTCAAGAGTCGAGTCACCATATCAGTAGAAACGTCCAAGAACCAGTTCTCCCTG
 AAGCTGAGCTCTGTGACCGCCGCGGACACGGCTGTGTATTACTGTGCGAGAGATAAGTGG
 ACCTGGTACTTCGATCTCTGGGGCCGTGGCACCCCTGGTCACTGTCTCCTCA

70 Heavy Chain Protein:

QVQLQQWGAGLLKPSETLSLTCAVYGGSFSGYYWSWIRQPPGKGLEWIGEINHSGSTNYN
 PSLKSRVTISVETSKNQFSLKLSSVTAADTAVYYCARDKWTWYFDLWGRGTLTVTVSS

71 Light Chain DNA:

GACATCGAGATGACCCAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGAGGGCCACC
 ATCAACTGCAGGTCCAGCCAGAGTGTTTTATACAGCTCCAGCAATAGGAACACTTAGCT
 TGGTACCAGCAGAACCAGGACAGCCTCCTAAGCTGCTCATTTACTGGGCTTCTACCCGG

GAATCCGGGGTCCCTGACCGATTTCAGTGGCAGCGGGTCTGGGACAGATTTCACTCTCACC
ATCAGCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTCAGCAATATTATAGTACT
CCTCGGACGTTCCGCCAAGGGACCAAGGTGGAAATCAAA

72 Light Chain Protein:

DIEMTQSPDSLAVSLGERATINCRSSQSVLYSSSNRNYLAWYQQNPGQPPKLLIYWASTR
ESGVPDRFSGSGSGTDFTLTISLSLOAEDVAVYYCQQYYSTPRTFGQGTKVEIK

Antibody U1-52

73 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGGTTACTACTGGAGCTGGATCCGC
CAGCAGCCAGGGAAGGGCCTGGAGTGGATGGGGAACATCTATTACAGTGGGAGCACCTAC
TACAACCCGTCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTGAGAACCAGTTC
TCCCTGAAGCTGAACTCTGTGACTGCCGCGGACACGGCCGTATATTACTGTGCGAGAGGG
GGAAGTGAACCAATTACTACTACTACTACGGTATGGACGTCTGGGGCCAAGGGACCACG
GTCACCGTCTCCTC

74 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQHPGKGLEWMENIYYSGSTY
YNPSLKSRTISVDTSNQFSLKLSVTAADTAVYYCARGGTGTNYYYYYGMDEVWGQGT
TVTS

75 Light Chain DNA:

GAAATTGTGTTGACGCAGTCTCCAGGCACCCTGTCTTTGTCTCCAGGGGAAAGAGCCACC
CTCTCCTGCAGGGCCAGTCAGAGTGTAGCAGCAGCTACTTAGCCTGGTACCAGCAGAAA
CCTGGCCAGGCTCCCAGGCTCCTCATCTATGGTGCATCCAGCTGGGCCACTGGCATCCCA
AACAGGTTTCAGTGGCAGTGGTCTGGGACAGACTTCACTCTCACCATCAGCAGACTGGAG
CCTGAAGATTTTGCAGTGTATTACTGTTCAGCAGTATGGTAGCTCACCCTCACTTTCCGC
GGAGGGACCAAGGTGGAGATCAAA

76 Light Chain Protein:

EIVLTQSPGTLSTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGASSWATGIP
NRFSGSGSGTDFTLTISRLEPEDFAVYYCQYGSSPLTFGGGTKVEIK

Antibody U1-61

77 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGTCTCCATCAGCAGTGGTGGTTACTACTGGAGCTGGATCCGC
CAGCAGCCAGGGATGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
TACAACCCGTCCTCAAGAGTCGAGTCACCATATCAGAAGACACGTCTAAGAACCAGTTC
TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGAT
TCCGAGTCCGAGTATAGCAGCTCGTCGAACCTACGGTATGGACGTCTGGGGCCAAGGGACC
ACGGTCACCGTCTCCTC

78 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGVSISSGGYYWSWIRQHPGMGLEWIGYIYYSGSTY
YNPSLKSRTISEDTSKNQFSLKLSSVTAADTAVYYCARDSESEYSSSSNYGMDEVWGQGT
TVTTS

Antibody U1-61.1

79 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGTCTCCATCAGCAGTGGTGGTTACTACTGGAGCTGGATCCGC
CAGCAGCCAGGGATGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
TACAACCCGTCCTCAAGAGTCGAGTCACCATATCAGAAGACACGTCTAAGAACCAGTTC
TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGAT
TCCGAGTCCGAGTATAGCAGCTCGTCGAACCTACGGTATGGACGTCTGGGGCCAAGGGACC

ACGGTCACCGTCTCCTC

80 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGVSISSGGYYWSWIRQHFGMGLEWIGYIYYSGSTY
YNPSLKSRVTISEDTSKNQFSLKLSSVTAADTAVYYCARDSESEYSSSSNYGMDVWGQGT
TVTVS

81 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAATCACC
ATCACTTGCCGGGCAAGTCAGACCATTAGCAGCTATTTAAATTGGTATCAGCAGAAACCA
GGGAAAGCCCCAAGCTCCTGATCTATGCTGCATCCAGTTTGCAAGGTGGGGTCCCATCA
AGGTTTCAGTGGCAGTGTATCTGGGACAGATTTACCCCTCACCGTCAGCAGTCTGCAACCT
GAAGATTTTGCAACTTACTACTGTCAACAGAGTTACAGTAACCCGCTCACTTTCCGGCGGA
GGGACCAAGGTGGAGATCAAA

82 Light Chain Protein:

DIQMTQSPSSLSASVGDRIITICRASQTISSYLNWYQQKPGKAPKLLIYAASSLQGGVPS
RFSGSVSGTDFTLTVSSLPEDFATYYCQQSYSNPLTFGGGTKEIK

Antibody U1-62 (2.9.1)

83 Heavy Chain DNA:

GAGGTGCAGCTGGTGCAGTCTGGAGCAGAGGTGAAAAAGCCCGGGGAGTCTCTGAAGATC
TCCTGTAAGGGTTCTGGATACAGTTTACCAGCTACTGGATCGGCTGGGTGCGCCAGATG
CCCGGGAAAGGCCTGGAGTGGATGGGGATCATCTATCCTGGTGACTCTGATACCAGATAC
AGCCCGTCCTTCCAAGGCCAGGTCACCATGTCAGCCGACAAGTCCATCAGTACCGCCTAC
CTGCAGCTGAGCAGCCATGAAGGCCTCGGACACCGCCATGTATTACTGTGCGAGACAGAT
GGCTGGAAACTACGTACATCACGGGTGATCGAGACGTCCTGGGGCCAAGGGACCACGGTC
ACCGTCTCCTC

84 Heavy Chain Protein:

EVQLVQSGAEVKKPGESLKIISCKGSGYSFTSYWIGWVRQMPGKGLEWMGIIPGDSDFTRY
SPSFQGVMTSADKSIISTAYLQLSSHEGLGHRVLLCETDQWKLRTSRVIETSWGQGTTV
TVS

85 Light Chain DNA:

GAAATTGTGTTGACGCAGTCTCCAGGCACCCTGTCTTTGTCTCCAGGGGAAAGAGCCACC
CTCTCCTGCAGGGCCAGTCAGAGTGTATCAGCATCTACTTAGCCTGGTACCAGCAGAAA
CCTGGCCAGGCTCCCAGGCTCCTCATCTATGGTGCATCCAGCAGGGCCACTGGCATCCCA
GACAGGTTTCAGTGGCAGTGGGTCTGGGACAGACTTCACTCTCACCATCAGCAGACTGGAG
CCTGAAGATTTTGCAGTGTATTACTGTCTCAGCAGTATGGTAGCTCACCCTGCAGTTTGGC
CAGGGGACCAAACTGGAGATCAAA

86 Light Chain Protein:

EIVLTQSPGTLSTLSPGERATLSCRASQSVISYLAWYQQKPGQAPRLLIYGASSRATGIP
DRFSGSGSGTDFTLTISRLEPEDFAVYYCQYGSFPCSFQGTKEIK

Antibody U1-2

87 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
CAGCACCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
TACAACCCGTCCCTCAGGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
TCCCTGAAGCTCAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCG
GATTACGATTTTGGAGTGGTTATTTTGACTACTGGGGCCAGGGAACCCCTGGTCACCGTC
TCCTCA

88 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTY
YNPSLRSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGQGTLVTV
SS

89 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGATACCT
 GGGAAAGCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTCAGTCTCACAATCAACAGCCTGCAGCCT
 GAAGATTTTGCAACTTATTACTGTCTACAGCATAATGGTTACCCGTGGACGTTGCGCCAA
 GGGACCAAGGTGGAAATCAAAC

90 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQGIRNDLGWYQQIPGKAPKRLIYAASSLQSGVPS
 RFSGSGSGTEFTLTINSIQPEDFATYYCLQHNGYPWTFGQGTKVEIK

Antibody U1-7

91 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
 CAGCACCCAGGGAAGGGCCTGGAGTGGATTGGATACATCTATTACAGTGGGAGCACCTAC
 TACAACCCGTCCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCG
 GATTACGATTTTGGAGTGGTTATTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTC
 TCCTCA

92 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTY
 YNPSLKSRTVISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGQGLTVTV
 SS

93 Light Chain DNA:

GACTTCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGGACATTGAAATGATTTAGGCTGGTATCGGCAGAAACCT
 GGGAAAGCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTCAGTCTCACAATCAGCAGCCTGCAGCCT
 GAAGATTTTGCAACTTATTACTGTCTACAGCATAATAGTTACCCGTGGACGTTGCGCCAA
 GGGACCAAGGTGGAAATCAAAC

94 Light Chain Protein:

DFQMTQSPSSLSASVGDRTITCRASQDIRNDLGWYRQKPGKAPKRLIYAASSLQSGVPS
 RFSGSGSGTEFTLTISLQPEDFATYYCLQHSYPWTFGQGTKVEIK

Antibody U1-9

95 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
 CAGCACCCAGGGAAGGGCCTGGAGTGGATTGGATACATCTATTACAGTGGGAGCACCTAC
 TACAACCCGTCCCTCAAGAGTCGAGTTACCATATCAATAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCG
 GATTACGATTTTGGAAATGGTTATTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTC
 TCCTCA

96 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTY
 YNPSLKSRTVISIDTSKNQFSLKLSSVTAADTAVYYCARADYDFWNGYFDYWGQGLTVTV
 SS

97 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGGACATTAGAAATGATTTAGGCTGGTATCGGCAGAAACCT
 GGGAAAGCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTCAGTCTCACAATCAGCAGCCTGCAGCCT

GAAGATTTTGCAACTTATTACTGTCTACAGCATAATAGTTACCCGTGGACGTTTCGGCCAA
GGGACCAAGGTGGAAATCAAA

98 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQDIRNDLGWYRQKPGKAPKRLIYAASSLQSGVPS
RFSGSGSGTEFTLTITSSSLQPEDFATYYCLQHNSYPWTFGQGTKVEIK

Antibody U1-10

99 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGGCCAGGACTGGTGAAGCCTACACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
CAGCACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
TACAACCCGTCCCTCAAGAGTCGACTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCA
GATTACGATTTTGGAGTGGTTACTTTGACTACTGGGGCCAGGGAACCCCTGGTCACCGTC
TCCTCA

100 Heavy Chain Protein:

QVQLQESGPGLVKPTQTLSTCTVSGGSISGDIYWSWIRQHPGKGLEWIGYIYYSGSTY
YNPSLKSRLTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGQGLVTV
SS

101 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
GGGAAAGCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
AGGTTCAGCGGCAGTGGATCTGGGACAGAATTCACTCTCACAATCAGCAGCCTGCAGCCT
GAAGATTTTGCAACTTATTACTGTCTACAGCATAATAATTACCCGTGGACGTTTCGGCCAA
GGGACCAAGGTGGAAATCAAA

102 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQGI RNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
RFSGSGSGTEFTLTITSSSLQPEDFATYYCLQHNNYPWTFGQGTKVEIK

Antibody U1-12

103 Heavy Chain DNA

CAGGTGCAGCTGCAGGAGTCGGGGCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGTAGTGGTGATTACTACTGGAGCTGGATCCGC
CAGCACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
TACAACCCGTCCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
TCCCTGAAGTTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCC
GATTACGATTTTGGAGTGGTTATTTGACTACTGGGGCCAGGGAACCCCTGGTCACCGTC
TCCTCA

104 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISGDIYWSWIRQHPGKGLEWIGYIYYSGSTY
YNPSLKSRLTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGQGLVTV
SS

105 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
GGGAAAGCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
AGGTTCAGCGGCAGTGGATCTGGGACAGAATTCACTCTCACAATCAGCAGCCTGCAGCCT
GAAGATTTTGCAACTTATTACTGTCTACAGCATAATAATTACCCGTGGACGTTTCGGCCAA
GGGACCAAGGTGGAAATCAAA

106 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQGI RNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
RFSGSGSGTEFTLTITSSSLQPEDFATYYCLQHNNYPWTFGQGTKVEIK

Antibody U1- 13

107 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGGTTACTACTGGAGCTGGATCCGC
 CAGCAGCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
 TACAACCCGTCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGAG
 GACGACGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCA

108 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYVWSWIRQHPGKLEWIGYIYYSGSTY
 YNPSLKSRTVISVDTSKNQFSLKLSSVTAADTAVYYCAREDDGMDVWGQGTITVTVSS

109 Light Chain DNA:

GATATTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTGGAGAGCCGGCCTCC
 ATTTCTGCAGGTCTAGTCAGAGCCTCCTGCATAGTAATGGATACAACATTTTGGAAATGG
 TACCTGCAGAAGCCAGGGCAGTCCCCACAGTTCATGATTTATTTGGGGTCTAATCGGGCC
 TCCGGGGTCCCTGACAGGTTCAGTGGCAGTGGATCAGGCACAGATTTTACACTGAAAATC
 AGCAGAGTGGAGGCTGAGGATGTTGGGGTTTATTACTGCATGCAAGCTCTACAACTCCG
 ATCACCTTCGGCCAAGGGACACGACTGGAGATTAAA

110 Light Chain Protein:

DIVMTQSPSLPVTTPGEPASISCRSSQSLHNSNGYNYLEWYLOKPGQSPQFMIYLGSNRA
 SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQALQTPITFGQGTREIK

Antibody U1-14

111 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGGTTACTACTGGAGCTGGATCCGC
 CAGTACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
 TACAACCCGTCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGAGGTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCG
 GATTACCATTTTTGGAGTGGTTATTTGACTACTGGGGCCAGGGAACCCCTGGTCACCGTC
 TCCTCA

112 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQYPGKLEWIGYIYYSGSTY
 YNPSLKSRTVISVDTSKNQFSLKLRSVTAADTAVYYCARADYDFWSGYFDYWGQGTITVTV
 SS

113 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
 GGGAAAGCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGCGGCAGTGGATCTGGGACAGAATCACTCTCACAATCAGCAGCCTGCAGCCT
 GAAGATTTTGCAACTTATTACTGTCTACAGCATAATACTTACCCGTGGACGTTCCGGCCAA
 GGGACCAAGGTGGAAATCAAAC

114 Light Chain Protein:

DIQMTQSPSSLSASVGRVTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
 RFGSGSGSTEFTLTISLQPEDFATYYCLQHNTYPTWTFGQGTKVEIK

Antibody U1-15

115 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCGGAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCGTCAGCAGTGGTGGTTACTACTGGAGCTGGATCCGG
 CAGCCCCAGGGAAGGGACTGGAGTGGATTGGGTATATCTATTACAGTGGGAGCACCAAC
 TACAACCCCTCCCTCAAGAGTCGAGTCACCATATCAGTAGACACGTCCAAGAACCAGTTC

TCCCTGAAGCTGAGCTCTGTGACCGCTGCGGACACGGCCGTGTATTACTGTGCGAGAGAT
GGGGACGTGGATACAGCTATGGTCGATGCTTTTGATATCTGGGGCCAAGGGACAATGGTC
ACCGTCTCCTCA

116 Heavy Chain Protein:

QVQLQESGPGLVKPSSETLSLTCTVSGGSVSSGGYYWSWIRQPPGKGLEWIGYIYYSGSTN
YNPSLKSRTVISVDTSKNQFSLKLSSVTAADTAVYYCARDGDVDAMVDAFDIWGQGTMTV
TVSS

117 Light Chain DNA:

GAAATTGTATTGACGCAGTCTCCAGGCACCCCTGTCTTTGTCTCCAGGGGAAAAGAGCCACC
CTCTCCTGCAGGGCCAGTCAGAGTTTAAGCGGCAACTACTTAGCCTGGTACCAGCAGAAG
CCTGGCCAGGCTCCCAGGCTCATCATCTGTGGTGCATCCAGCAGGGCCACTGGCATCCCA
GACAGGTTCACTGGCAGTGGGTCTGGGACAGACTTCCTCTCACCATCACAAGACTGGAG
CCTGAAGATTTTGCAGTGTATTACTGTGACAGTATGATAGGTACCCGCTCACTTTCGGC
GGAGGGACCAAGGTGGAGATCAAA

118 Light Chain Protein:

EIVLTQSPGTLSPGERATLSCRASQSLSGNYLAWYQQKPGQAPRLIICGASSRATGIP
DRFSGSGSGTDFTLTITRLEPEDFAVYYCQQYDRSPLTFGGGTKEIK

Antibody U1-19

119 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
CAGCACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
TACAACCCGTCCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGGA
GATTACGATTTTGGAGTGGAGAGTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTC
TCCTCA

120 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTY
YNPSLKSRTVISVDTSKNQFSLKLSSVTAADTAVYYCARGDYDFWSGEFDYWQGTLLTV
SS

Antibody U1-20

121 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGGTTACTACTGGAGCTGGATCCGC
CAGCACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATGACAGTGGGAGCACCTAC
TACAACCCGTCCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGAT
CAGGGGCAGGACGATACAGCTATGGTTACGGCTACTACTACGGTATGGACGTCTGGGGC
CAAGGGACCACGGTCACCGTCTCCTC

122 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYWSWIRQHPGKGLEWIGYIYDSGSTY
YNPSLKSRTVISVDTSKNQFSLKLRSVTAADTAVYYCARDQGDGYSYGYGYGMDVWG
QGTTVTVS

123 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCAGGCGAGTCAGGACATTAGCAATTATTTAAATTGGTATCAGCAGAAACCA
GGGAAAGCCCCCTAACTCCTGATCTACGTTGCATCCAATTTGGAAACAGGGGTCCCATCA
AGGTTCACTGGAAGTGGATCTGGGACAGATTTTACTTTACCATCAGCAGCCTGCAGCCT
GAAGATATTGCAACATATTACTGTCAACAGTGTGATAATCTCCCTCTCACTTTCCGGCGGA
GGGACCAAGGTGGAGATCAAA

124 Light Chain Protein:

124 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCQASQDISNYLNWYQQKPGKAPKLLIYVASNLETGVPS
RFSGSGSGTDFTTITSSLPEDIATYYCQQCDNLPITFGGGTKVEIK

Antibody U1-21

125 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGGCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
CAGCACCAGGGAAGGGCCTGGAGTGGATTGGATACATCTATTACAGTGGGAGCACCTAC
TACAACCCGTCCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCG
GATTACGATTTTGGAGTGGTTATTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTC
TCCTC

126 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTY
YNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGQGLVTV
S

127 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCGGGCAAGTCAGGACATTAGAAATGATTTAGGCTGGTATCGGCAGAAACCT
GGGAAAGCCCCTAAGCGCCTGATCTATGCTGCATCCCGTTTGCAAAGTGGGTCCCATCA
AGGTTTCAGCGGCAGTGGATCTGGGACAGAATCACTCTCACAATCAGCAGCCTGCAGCCT
GAAGATTTTGCAACTTATTACTGTCTACAGCATAATAGTTACCCGTGGACGTTCCGGCCAA
GGGACCAAGGTGGAAATCAAAC

128 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQDIRNDLGWYRQKPGKAPKRLIYAASRLQSGVPS
RFSGSGSGTEFTLTITSSLPEDFATYYCLQHNSYPWTFGQGTKVEIK

Antibody U1-22

129 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGGCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
CAGCACCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
TACAACCCGTCCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCC
GATTACGATTTTGGAGTGGTTATTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTC
TCCTCA

130 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTY
YNPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGQGLVTV
SS

131 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
GGGAAAGCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAATGGGGTCCCATCA
AGGTTTCAGCGGCAGTGGATCTGGGACAGAATCACTCTCACAATCAGCAGCCTGCAGCCT
GAAGATTTTGCAACTTATTACTGTCTACAGCATAATAGTTACCCGTGGACGTTCCGGCCAA
GGGACCAAGGTGGAAATCAAAC

132 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQIRNDLGWYQQKPGKAPKRLIYAASSLQNGVPS
RFSGSGSGTEFTLTITSSLPEDFATYYCLQHNSYPWTFGQGTKVEIK

Antibody U1-23

Antibody U1-23

133 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
 CAGCAGCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
 TACAACCCGTCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCG
 GATTACGATTTTTGGAGTGGTTATTTGACTACTGGGGCCAGGGAATCCTGGTCACCGTC
 TCCTC

134 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTY
 YNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGQGILVTV
 S

135 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
 GGGAAAGCCCCCTAAGCGCCTGATTTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTCACCTCTCACAATCAGCAGCCTGCAGCCT
 GAAGATTTTCAACTTATTACTGTCTACAGCATAATAGTTACCCGTGGACGTTCCGGCCAA
 GGGACCAAGGTGGAAATCAAAC

136 Light Chain Protein:

DIQMTQSPSSLSASVGDRVTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
 RFGSGSGTEFTLTISLQPEDFATYYCLQHNSYPWTFGQGTKVEIK

Antibody U1-24

137 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGTAGTGGTGATTACTACTGGAGCTGGATCCGC
 CAGCAGCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
 TACAACCCGTCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGTTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCC
 GATTACGATTTTTGGAATGGTTATTTTGAATACTACTGGGGCCAGGGAACCCTGGTCACCGTC
 TCCTCA

138 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTY
 YNPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWNGYFDYWGQGLTVTV
 SS

139 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
 GGGAAAGCCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTCACCTCTCACAATCAGCAGCCTGCAGCCT
 GAAGATTTTCAACTTATTACTGTCTACAGCATAATAATTACCCGTGGACGTTCCGGCCAA
 GGGACCAAGGTGGAAATCAAA

140 Light Chain Protein:

DIQMTQSPSSLSASVGDRVTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
 RFGSGSGTEFTLTISLQPEDFATYYCLQHNNYPWTFGQGTKVEIK

Antibody U1-25

141 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
 CAGCAGCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC

TACAACCCGTCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCC
 GATTACGATTTTGGAGTGGTTATTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTC
 TCCTCA

142 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYWSWIRQHPGKGLEWIGYIYYSGSTY
 YNPSLKSRTVISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGQSTLTV
 SS

143 Light Chain DNA:

GACATCCAGCTGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
 GGGAAAGCCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAATGGGGTCCCATCA
 AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTCCTCTCACAATCAGCAGCCTGCAGCCT
 GAAGATTTTGCAACTTATTACTGTCTACAGCATAATAGTTACCCGTGGACGTTTCGGCCAA
 GGGACCAAGGTGGAAATCAAAC

144 Light Chain Protein:

DIQLTQSPSSLSASVGDRTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQNGVPS
 RFSGSGSGTEFTLTISLQPEDFATYYCLQHNSYPWTFGQGTKVEIK

Antibody U1-26

145 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
 CAGTACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
 TACAACCCGTCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGGGCTCTGTGACTGCCGCGGACACGGCCGTGTATTTCTGTGCGAGAGCC
 GATTACGATTTTGGAGTGGTTATTTTGACTTCTGGGGCCAGGGAACCCTGGTCACCGTC
 TCCTC

146 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYWSWIRQYPGKGLEWIGYIYYSGSTY
 YNPSLKSRTVISVDTSKNQFSLKLGSVTAADTAVYFCARADYDFWSGYFDYWGQSTLTV
 S

147 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
 ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
 GGGAAAGCCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
 AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTCCTCTCACAATCAGCAGCCTGCAGCCT
 GAAGATTTTGCAACTTATTACTGTCTACAGCATAATGGTTACCCGTGGACGTTTCGGCCAA
 GGGACCAAGGTGGAAATCAAAC

148 Light Chain Protein:

DIQMTQSPSSLSASVCDRTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
 RFSGSGSGTEFTLTISLQPEDFATYYCLQHNGYPWTFGQGTKVEIK

Antibody U1-27

149 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
 ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
 CAGTACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
 TACAACCCGTCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
 TCCCTGAAGCTGGGCTCTGTGACTGCCGCGGACACGGCCGTGTATTTCTGTGCGAGAGCC
 GATTACGATTTTGGAGTGGTTATTTTGACTTCTGGGGCCAGGGAACCCTGGTCACCGTC
 TCCTC

150 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQYPGKGLEWIGYIYYSGSTY
YNPSLKSRVTISVDTSKNQFSLKLSVTAADTAVYFCARADYDFWSGYFDWFGQGLVTV
S

151 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
GGGAAAGCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
AGGTTCAAGCGGCAGTGGATCTGGGACAGAATTCAGTCTCACAATCAGCAGCCTGCAGCCT
GAAGATTTTGCAACTTATTACTGTCTACAGCATAATGGTTACCCGTGGACGTTCCGGCCAA
GGGACCAAGGTGGAAATCAAAC

152 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQGIKNDLGWYQQKPKAPKRLIYAASSLQSGVPS
RFGSGSGTEFTLTISLQPEDFATYYCLQHNGYPWTFGQGTKVEIK

Antibody U1-28

153 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGTAGTGGTGATTACTACTGGAGCTGGATCCGC
CAGCACCCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGAGCACCTAC
TACAACCCGTCCTCAAGAGTCGAGTTACCATATCAGTAGACACGTCCTAAGAACCAGTTC
TCCCTGAAGCTGAGCTCTGTGACTGCCGCGACACGGCCGTGTATTACTGTGCGAGAGCG
GATTACGATTTTGGAGTGGTTATTTTACTCTCGGGCCAGGGAACCCTGGTCACCGTC
TCCTCA

154 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHHPKGLEWIGYIYYSGSTY
YNPSLKSRVTISVDTSKNQFSLKLSVTAADTAVYYCARADYDFWSGYFDSWGQGLVTV
SS

155 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGATACCT
GGGAAAGCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
AGGTTCAAGCGGCAGTGGATCTGGGACAGAATTCAGTCTCACAATCAGCAGCCTGCAGCCT
GAAGATTTTGCAACTTATTACTGTCTACAGCATAATGGTTACCCGTGGACGTTCCGGCCAA
GGGACCAAGGTGGAAATCAAAC

156 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQGIKNDLGWYQQIPGKAPKRLIYAASSLQSGVPS
RFGSGSGTEFTLTISLQPEDFATYYCLQHNGYPWTFGQGTKVEIK

Antibody U1-31

157 Heavy Chain DNA:

CAGGTTCAAGCTGGTGCAGTCTGGAGCTGAGGTGAAGAAGCCTGGGGCCTCAGTGAAGGTC
TCCTGCAAGGCTTCTGGTTACACCTTTACCACTATGGTATCAGCTGGGTGCGGCAGGCC
CCTGGACAAGGGCTTGAGTGGATGGGATGGATCAGCGCTTACGATGGTTACAGAACTAT
GCACAGAAGCTCCAGGGCAGAGTCACCATGACCACAGACACATCCACGACCACTGCCTAC
ATGGAGCTGAGGAGCCTGAGATCTGACGACACGGCCGTGTATTACTGTGCGAGAGATGTT
CAAGACTACGGTGACTACGACTACTTTGACTACTGGGGCCAGGGAACCCCTGGTCACCGTC
TCCTCA

158 Heavy Chain Protein:

QVQLVQSGAEVKKPGASVKVSCKASGYFTFTNYGISWVRQAPGQGLEWMGWISAYDGYRNY
AQKLQGRVTMTTDTSTTTAYMELRSLRSDDTAVYYCARDVQDYGDYDFDYWGQGLVTV
SS

159 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC

ATCACTTGCCGGGCAAGTCAGAGCATTAGCAGTTATTTAAATTGCTATCAGCAGAAACCA
GGGAAAGCCCCTAACCTCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
AGATTTCAGGGGCAGTGGATCTGGGACAGATTTCACTCTCACCATCAGCAGTCTGCAACCT
GAAGATTTTCCAACCTTACTACTGTCAACAGAGTTACAGTACCCCCATCACCTTCGGCCAA
GGGACACGACTGGAGATTAAA

160 Light Chain Protein:

DIQMTQSPSSLSASVGDRVITITCRASQSISSYLNWYQQKPGKAPNLLIYAASSLQSGVPS
RFRSGSGTDFTLTISLQPEDFATYYCQQSYSTPITFGQGRLEIK

Antibody U1-32

161 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGCCCAGGACTGGTGAAGCCTTTACAGACCCTGTCCCTC
ACCTGCACTGTCTCTGGTGGCTCCATCAGCAGTGGTGATTACTACTGGAGCTGGATCCGC
CAGCACCAGGGAAGGGCCTGGAGTGGATTGGGTACATCTATTACAGTGGGACCACCTAC
TACAACCCGTCCTCAAGAGTCGAGTTACCATATCAGTAGACACGCTAAGAACCAGTTC
GCCCTGAAGCTGAACCTCTGTGACTGCCGCGACACGGCCGTGTATTACTGTGCGAGAGCC
GATTACGATTTTGGAGTGGTTATTTGACTACTGGGGCCAGGGAACCTGGTCACCGTC
TCCTCA

162 Heavy Chain Protein:

QVQLQESGPGLVKPLQLTSLTCTVSGGSISSGDYYSWIRQHPGKGLEWIGYIYYSGTTY
YNPSLKSRTISVDTSKNQFALKLNSVTAADTAVYYCARADYDFWSGYFDYWGQGLTVT
SS

163 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCGGGCAGGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
GGGAAAGCCCCTCAGCCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTCTCTCTCACAATCTCCAGCCTGCAGCCT
GAAGATTTTGCAACTTATTACTGTCTACAGCATAATAGTTACCCGTGGACGTTCCGGCCAA
GGGACCAAGGTGGAAATCAAAC

164 Light Chain Protein:

DIQMTQSPSSLSASVGDRVITITCRAGQGIRNDLGWYQQKPGKAPQRLIYAASSLQSGVPS
RFSGSGSGTEFSLTISLQPEDFATYYCLQHNSYPWTFGQGTKVEIK

Antibody U1-35

165 Heavy Chain DNA:

CAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTCAAGCCTGGAGGGTCCCTGAGACTC
TCCTGTGCAGCCTCTGGATTACCTTCAGTGACTACTACATGAGCTGGATCCGCCAGGCT
CCAGGGAAGGGGCTGGAGTGGGTTTCATATATTAGTAGTAGTGGTAATAACATATACCAC
GCAGACTCTGTGAAGGGCCGATTCACCATCTCCAGGGACAACGCCAAGAACTCACTGTAT
CTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCCGTGTATTACTGTGCGAGAGAGAGA
TATAGTGGCTACGACGACCCTGATGGTTTTGATATCTGGGGCCAAGGGACAATGGTCACC
GTCTCTTCA

166 Heavy Chain Protein:

QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMSWIRQAPGKGLEWVSYISSSGNNIYH
ADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARERYSGYDDPDGFDIWGQGTMTV
VSS

167 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCAGGCGAGTCAGGACATTAGCAACTATTTAAGTTGGTTTCAGCAGAAACCA
GGGAAAGCCCCTAAGCTCCTGATCCACGATGCATCCAATTTGGAAACAGGGGTCCCTTCA
AGGTTTCAGTGAAGTGGATCTGGGACAGATTTTACTTTACCATCAGCAGCCTGCAGCCT
GAAGATATTGCAACATATTACTGTCAACAGTATGATAATCCCCCGTCAGTTTGGCCAG
GGGACCAAGCTGGAGATCAAA

168 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCQASQDISNYLSWFQQKPGKAPKLLIHDAENLETGVPS
RFGSGSGTDFTTITSSLPEDVATYYCQYDNPPCSFGQGTKLEIK

Antibody U1-36

169 Heavy Chain DNA:

CAGGTGCAGCTGCAGGAGTCGGGGCCAGGACTGGTGAAGCCTTCACAGACCCTGTCCCTC
ACCTGCAGCTGTCTCTCGGTGGCTCCATCAGCAGTGGTTATTACTACTGGAGCTGGATCCGC
CAGCAGCCAGGGAAGGGCCCTGGAGTGGATTGGGTACATCTATTACAGTGGGACCACCTAC
TACAATCCGTCCTTCAAGAGTCGAGTTACCATATCAGTAGACACGTCTAAGAACCAGTTC
TCCCTGAAACTGAGCTCTGTGACTGCCGCGGACACGGCCGTGTATTACTGTGCGAGAGCC
GATTACGATTTTTGGAGTGGTCACCTTTGACTACTGGGGCCAGGGAACCCTGGTCACCGTC
TCCTCA

170 Heavy Chain Protein:

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGYYWSWIRQHPGKGLEWIGYIYYSGTTY
YNPSFKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGHFDYWGQGLVTV
SS

171 Light Chain DNA:

GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCATCTGTAGGAGACAGAGTCACC
ATCACTTGCCGGGCAAGTCAGGGCATTAGAAATGATTTAGGCTGGTATCAGCAGAAACCA
GGGAAAGCCCCCTAAGCGCCTGATCTATGCTGCATCCAGTTTGCAAAGTGGGGTCCCATCA
AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTCACCTCTCACAATCAGCAGCCTGCAGCCT
GAAGATTTTGCAACTTATTACTGTCTACAGCATAATAGTTACCCGTGGACGTTCCGCCAA
GGGACCAAGGTGGAAATCAAA

172 Light Chain Protein:

DIQMTQSPSSLSASVGDRTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
RFGSGSGTEFTLTITSSLPEDFATYYCLQHNSYPWTFGQGTKVEIK

Antibody U1-37

173 Heavy Chain DNA:

CAGGTTTCAGCTGGTGCAGTCTGGAGCTGAGGTGAAGAAGCCTGGGGCCTCAGTGAAGGTC
TCCTGCAAGGCTTCTGGTTACACCTTTACCAGCTATGGTATCAGCTGGGTGCGACAGGCC
CCTGGACAAGGACTTGAGTGGATGGGATGGATCAGCGCTTACGATGGTCACACAACTAT
GCACAGAAGCTCCAGGGCAGAGTCACCATGACCACAGACACATCCACGAACACAGCCTAC
ATGGAGCTGAGGAGCCTGAGATCTGACGACACGGCCGTTTATTACTGTGCGAGAGACCCC
CATGACTACAGTAACTACGAGGCTTTTGACTTCTGGGGCCAGGGAACCCTGGTCACCGTC
TCCTC

174 Heavy Chain Protein:

QVQLVQSGAEVKKPGASVKVSCKASGYTFSTSYGISWVRQAPGQGLEWMGWISAYDGHNTY
AQLQLGRVTMTTDTSTNTAYMELRSLRSDDTAVYYCARDPHDYSNIEAFDFWGQGLVTV
S

175 Light Chain DNA

atgaggtccctgtctcagctcctggggctcctgtctactctggctccgaggtgccagatgtg
acatccagatgacccagctctccatcctccctgtctgcatctgtaggagacagagtcacat
cacttgccgggcaagtcagagcatttagcagttatthaaattggtatcagcagaaaccaggg
aaagcccctaaccctcctgatctatgtctgcatccagtttgcaaagtggggtcccatcaagat
tcagtgccagtggtctctgggacagatttcactctcaccatcagcagctctgcaacctgaaga
ttttgcaacttactactgtcaacagagttacagtagcccccacaccttcggccaagggaaca
cgactggagattaaacgaactgtggctgcaccatctgtcttcatcttcccgccatctgatg
agcagttgaaatctggaactgcctctgtgtgtgctgctgaataacttctatcccagaga
ggccaaagtacagtggaaggtggataacgcc

176 Light Chain Protein:

DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPNLLIYAASSLQSGVPSR
FSGSGSGTDFTLTISLQPEDFATYYCQQSYSTPITFGQGRLEIK

5

Antibody U1-34

177 Heavy Chain DNA:

10 accatggactggacctggagggtccttttcttggtggcagcagcaacaggtgcccactccca
gggttcagctggtgcagctctggagctgaggtgaagaagcctggggcctcagtgaaaggtctcct
gcaaggcttctggttacacctttaccaactatggtatcagctgggtgcggcaggccctgga
caagggcttgagtggatgggatggatcagcgcttacgatgggttacagaaactatgcacagaa
15 gctccagggcagagtcacatgaccacagacacatccacgaccactgcctacatggagctga
ggagcctgagatctgacgacacggccgtgtattactgtgcgagagatggttcaagactacggt
gactacgactactttgactactggggccagggaaccctggtcaccgtctcctcagcttccac
caagggcccatccgtcttccccctgggtgcccgtgtccagagacacctccgagagcacagccg
ccttggtgctgctggtcaaggactacttccccgaaccg

20

178 Heavy Chain Protein

QVQLVQSGAEVKKPGASVKVSKASGYTFTNYGISWVRQAPGQGLEWMGWISAYDGYRNYA
QKLQGRVTMTTDTSTTTAYMELRSLRSDDTAVYYCARDVQDYGDDYFDYWGQGLTVTVSS

25

179 Light Chain DNA:

25 cagctcctggggctcctgctactctggctccgaggtgccagatgtgacatccagatgaccc
agctccatcctcctgtctgcactctgtaggagacagagtcacatcacttgccgggcaag
tcagagcattagcagttatttaaattggtatcagcagaaaccagggaagccctaacctc
ctgatctatgctgcacccagtttgcaaagtgggttcccatcaagattcagtggcagtggt
30 ctgggacagatttcaactctcaacatcagcagctctgcaacctgaagattttgcaacttacta
ctgtcaacagagttacagtacccccatcaccttcggccaagggaacagactggagattaaa
cgaaactgtggctgcaccatctgtcttcatcttcccgccatctgatgagcagttgaaatctg
gaactgcctctgttgtgtgctgctgaataacttctatccagagaggccaaagtacagtg
35 gaaggtggataacgcc

35

180 Light Chain Protein:

DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPNLLIYAASSLQSGVPSR
FSGSGSGTDFTLTISLQPEDFATYYCQQSYSTPITFGQGRLEIK

40

Antibody U1-1

181 Heavy Chain DNA:

45 catctgtggttcttctcctctgctggtggcagctcccagatgggtcctgtcccaggtgcagc
tgcaggagtcgggcccaggactgggtgaagccttcacagaccctgtccctcacctgcactgt
ctctggtggctccatcaacagtgggtgattactactggagctggatccgccagcaccaggg
aagggcctggagtggattgggtacatctattacagtgggagcacctactacaaccogtccc
tcaagagtcaggttaccatatacgtagacacgtctaagaaccagttctccctgaagctgag
50 ctctgtgactgcccggacacggccgtgtattactgtgcgagagcagattacgatttttg
agtgggttactttgactactggggccagggaaccctggtcaccgtctcctcagcctccacca
agggcccatcggtcttccccctggcaccctcctccaagagcacctctgggggcacaacggc
cctgg

55

182 Heavy Chain Protein

QVQLQESGPGLVKPSQTLSTCTVSGGSINSGDYYSWIRQHHPKGLEWIGYIYYSGSTYY
NPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWQGQGLVTVSS

183 Light Chain DNA:

atgaggggtccctgctcagctcctggggctcctgctgctctgggttcccagggtgccagggtgtga
catccagatgacccagctctccatccctcctgtctgcatctgtaggagacagagtcacccatca
cttgccggggcaagtcagggtcattagaaatgatttaggctgggtatcagcagaaaccaggga
gccccctaagcgcctgatctatgctgcacccagtttgcaaagtgggggtcccatcaagggttcag
cggcagtggtatctgggacagaattcactctcacaatcagcagcctgcagcctgaagattttg
caacttattactgtctacagcataatagttaccctggacgttcggccaagggaaccaagggtg
gaaatcaaacgaactgtggctgcaccatctgtcttcacatctcccgccatctgatgagcagtt
gaaatctggaactgcctctgttggtgctgctgaataacttctatcccagagaggccaaag
tacagtggaagggtggataacgc

184 Light Chain Protein

DIQMTQSPSSLSASVGDRTITCRASQGIKNDLWYQQKPKAPKRLIYAASSLQSGVPSR
FSGSGSGTEFTLTISSLQPEDFATYYCLQHNSYPWTFGQGTKVEIK

Antibody U1-3

185 Heavy Chain DNA:

tgggtcttctctctgctgggtggcagctcccagatgggtcctgtcccagggtgcagctgcagga
gtcggggcccaggactgggtgaagccttcacagaccctgtccctcacctgcactgtctctgggtg
gtcccatcagcagtggtgggttactactggagctggatccgccagcaccagggaagggtcctg
gagtggtattgggtacatctattacagtgggagcaccctactacaaccctccctcaagagtcg
agttaccatatcagtagacacgtctaagaaccagttctccctgaagctgagctctgtgactg
ccggcgacacggccgtgtattactgtgagagagatggctatgatagtagtggttattaccac
ggctactttgactactggggccagggaaccctgggtcaccgtctcctcagcctccaccaaggg
cc

186 Heavy Chain Protein

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGGYYSWIRQHHPKGLEWIGYIYYSGSTYY
NPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARDGYDSSGGYHGYFDYWQGQGLVTVSS

187 Light Chain DNA:

H3_130_1N1K

cagggtcttctctctgctgggtggcagctcccagatgggtcctgtcccagggtgcagctgcagga
gtcggggcccaggactgggtgaagccttcacagaccctgtccctcacctgcactgtctctgggtg
gtcccatcagcagtggtgggttactactggagctggatccgccagcaccagggaagggtcctg
gagtggtattgggtacatctattacagtgggagcaccctactacaaccctccctcaagagtcg
agttaccatatcagtagacacgtctaagaaccagttctccctgaagctgagctctgtgactg
ccggcgacacggccgtgtattactgtgagagagatggctatgatagtagtggttattaccac
ggctactttgactactggggccagggaaccctgggtcaccgtctcctcagcctccaccaaggg
cc

188 Light Chain Protein:

DIVMTQSPDSLAVSLGERATINCKSSQSVLYSSNNKNYLAWYQQKPGQPPKLLIYWASTRE
SGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCQYYSTPLTFGGGTKVEIK

Antibody U1-4

H3_133_1N1G1

189 Heavy Chain DNA

ctgtggttcttctctctgtgtggtggcagctcccagatgggtcctgtcccaggtgcagctgca
 ggagtcgggcccaggactggtgaagccttcacagaccctgtccctcacctgcactgtctctg
 gtggctccatcagtagtggtgattactactggagctggatccgccagcaccaggggaagggc
 ctggagtggattgggtacatctattacagtgggagcacctactacaaccgctccctcaagag
 tccagttaccatatcagtagacacgtctaaagaaccagttctccctgaagttgagctctgtga
 ctgcgcggacacggcgtgtattactgtgcgagagccgattacgatttttggagtggttat
 ttgactactggggccagggaaccctggtcaccgtctcctcagcctccaccaagggcccatc
 ggtcttccccctggcaccctc

190 Heavy Chain Protein

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTYY
 NPSLKSRTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGQGLTVTVSS

191 Light Chain DNA

H3_133_1N1K

gtgcccgtcagcgcctggggctcctgtgtctctggttcccaggtgccaggtgtgacatcc
 agatgacccagtcctccatcctcctgtctgtcatctgtaggagacagagtcaccatcacttg
 ccgggcaagtcagggcattagaaatgatttaggctgggtatcagcagaaaccagggaagcc
 cctaagcgcctgatctatgtctgtcatccagtttgcaaagtggggctccatcaaggttcagcg
 gcagtggtatctgggacagaattcactctcacaatcagcagcctgcagcctgaagattttgc
 aacttattactgtctacagcataataattaccgctggacgttcggccaagggaaccaaggtg
 gaaatcaaacgaactgtggctgcaccatctgtcttcatcttcccgcctctgatgagcagt
 tgaaatctggaactg

192 Light Chain Protein

DIQMTQSPSSLSASVGDRTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPSRF
 SGSGSGTEFTLTITSSLPEDFATYYCLQHNYPWTFGQGTKVEIK

Antibody U1-5

193 Heavy Chain DNA:

H3_138_1N1G1

tggttcttctctctgtgtggtggcagctcccagatgggtcctgtcccaggtgcagctgcagga
 gtcgggcccaggactggtgaagccttcacagaccctgtccctcacctgcactgtctctggtg
 gctccatcagcagtggtgattactactggagctggatccgccagcaccaggggaagggcctg
 gagtggattgggtacatctattacagtgggagcacctactacaaccgctccctcaagagtcg
 agttaccatatcagtagacacgtctaaagaaccagttctccctgaagctgagctctgtgactg
 ccgcgacacggcgtgtatttctgtgcgagagccgattacgatttttggagtggttatttt
 gactactggggccagggaaccctggtcaccgtctcctcagcctccaccaagggcc

194 Heavy Chain Protein

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTYYN
 PSLKSRTISVDTSKNQFSLKLSSVTAADTAVYFCARADYDFWSGYFDYWGQGLTVTVSS

195 Light Chain DNA:

H3_138_1N1K

atgagggtccccgtcagctcctggggctcctgtgtctctggttcccaggtgccaggtgtga
 catccagatgacccagtcctccatcctcctgtctgtcatctgtaggagacagagtcaccatca
 cttgccgggcaagtcagggcattagaaatgatttaggctgggtatcagcagaaaccaggga

gccccctaagcgccctgatctatgctgcatccagtttgcaaagtggggtcccatcaagggtcag
 cggcagtggtatctgggacagaattcactctcacaatcagcagcctgcagcctgaagattttg
 caacttattactgtctacagcataatacttaccctggacgttcggccaagggaccaagggtg
 gaaatcaaacgaactgtggctgcaccatctgtcttcattctcccgccatctgatgagcagtt
 gaaatctggaactgcctctgttgtgtgcctgctgaataacttctatcccagagaggccaaag
 tacagtgggaaggtggataacgc

196 Light Chain Protein

DIQMTQSPSSLSASVGDRVTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
 RFSGSGSGTEFTLTISLQPEDFATYYCLQHNTYPWTFGQGTKVEIK

Antibody U1-6

197 Heavy Chain DNA:

H3_162_1N1G1

tgggtcttccctctgctgggtggcagctcccagatgggtcctgtcccaggtgcagctgcagga
 gtccggcccaggactgggtgaagccttcacagaccctgtccctcactgcactgtctctgggtg
 gctccatcagcagtggtgattactactggagctggatccgccagcaccagggaagggcctg
 gagtggattgggtacatctattacagtgggagcacctactacaaccgctccctcaagagtcg
 agttaccatatcagtagacacgtctaagaaccagttctccctgaagctgagctctgtgactg
 ccgcggacacggccgtgtatttctgtgagagagccgattacgatttttgggaatgggtatttt
 gactactggggccagggaaccctgggtcacccgtctcctcagcctccaccaagggccc

198 Heavy Chain Protein

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTY
 YNPSLKRVTISVDTSKNQFSLKLSSVTAADTAVYFCARADYDFWNGYFDYWGQGLVTV
 SS

199 Light Chain DNA:

H3_162_1N1K

atgagggtccccgctcagctcctggggctcctgctgctctgggtcccaggtgccaggtgtga
 catccagatgacccagctctccatcctccctgtctgcattctgtaggagacagagtcaccatca
 cttgcggggcaagtcagggcattagaaatgatttaggctgggtatcagcagaaaccaggga
 gccccctaagcgccctgatctatgctgcttccagtttgcaaagtggggtcccatcaagggttcag
 cggcagtggtatctgggacagaattcactctcacaatcagcagcctgcagcctgaagattttg
 caacttattactgtctacagcataatacttaccctggacgttcggccaagggaccaagggtg
 gaaatcaaacgaactgtggctgcaccatctgtcttcattctcccgccatctgatgagcagtt
 gaaatctggaactgcctctgttgtgtgcctgctgaataacttctatcccagagaggccaaag
 tacagtgggaaggtggataacgc

200 Light Chain Protein

DIQMTQSPSSLSASVGDRVTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
 RFSGSGSGTEFTLTISLQPEDFATYYCLQHNTYPWTFGQGTKVEIK

Antibody U1-8

201 Heavy Chain DNA:

H3_174_1N1G1

ttgggtggcagcagctacaggcacccacgcccagggtccagctggtacagtctggggctgaggt
 gaagaagcctggggcctcagtggaaggtctcctgcaaggtttccggatacaccctcactgaat
 tatccatgtactgggtgcgacaggctcctggaaaagggttgagtggatgggaggttttgat

cctgaagatgggtgaaacaatctacgcacagaagtccagggcagagtcaccatgaccgagga
 cacatctacagacacagcctacatggagctgagcagcctgagatctgaggacacggccgtgt
 attactgtgcaactgggtggaactacgtctttgactactggggccagggaaccctggtcacc
 gtctctcagcctccaccaagggcc

5

202 Heavy Chain Protein

QVQLVQSGAEVKKPGASVKVSKVSGYTLTELSMYWVRQAPGKGLEWMGGFDPEDGETIYA
 QKFQGRVTMTEDTSTDYAMELSSLRSEDYAVYYCATGWNVYFDYWGQGLVTVSS

10

203 Light Chain DNA:

H3_174_1N1K

ggatccagtgaggatattgtgatgactcagtcctccactctccctgcccgtcaccctggaga
 gcggcctccatctccctgcaggtccagtcagagcctcctgcataagtaatggatacaactatt
 tggatttggtacctgcagaagccagggcagtcctccacagctcctgatctatttggattctcat
 cgggcctccggggtccctgcaggttcagtgaggatcaggcacagattttacactgaa
 aatcagcagagtgagggtgaggatgttgggggtttattactgcatgcaagctctacaaactc
 cgctcacttttcggcggagggaaccaaggtggagatcaaacgaactgtggctgcaccatctgtc
 ttcactctcccgccat

15

20

204 Light Chain Protein

DIVMTQSPLSLPVTPGEPASISCRSSQSLLSHNGYNYLDWYLQKPGQSPQLLIYLD SHRA
 SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQALQTPITFGGGTKVEIK

25

Antibody U1-11

205 Heavy Chain DNA:

H3_178_1N1G1

tggttcttccctctgtctgggtggcagctcccagatgggtcctgtcccaggtgcagctgcagga
 gtccggcccaggactgggtgaagccttcacagaccctgtccctcactgcactgtctctgggtg
 gctccatcagcagtggtgattactactggagctggatccgccagcaccaggggaagggcctg
 gagtgattgggtacatctattacagtgggagcactactacaaccctgcccctcaagagtcg
 agttaccatatcagtagacacgtctaagaaccagttctccctgaagctgagctctgtgactg
 ccggggacacggccgtgtatttctgtgagagccgattacgatttttggagtgggtattttt
 gactactggggccagggaaccctgggtcaccgtctcctcagcctccaccaaggggcccatcgag
 tcttccccctgg

30

35

206 Heavy Chain Protein

QVQLQESGPGIIVKPSQTLSTCTVSGGSISGDIYWSWIRQHPGKGLEWIGYIYSGSTY
 YNPSLKSRTISVDTSKNQFSLKLSSVTAADYAVYFCARADYDFWSGYFDYWGQGLVTV
 SS

40

207 Light Chain DNA:

H3_178_1N1K

atgaggggtcccgcctcagctcctggggctcctgctgctctgggtcccaggtgccaggtgtg
 acatccagatgaccagctctccatcctccctgtctgcactctgtaggagacagagtcacat
 cacttgccgggcaagtcagggcattagaaatgatttaggctgggtatcagcagaaaccaggg
 aaagcccctaagcgccctgatctatgctgcacccagtttgcaaagtgggggtcccatcaaggt
 tcagcggcagtggtctgtgggacaaaattcactctcactatcagcagcctgcagcctgaaga
 ttttgcaacttattactgtctacagcataatacttaccgtggagcttcggccaagggaacc
 aaggtggaaatcagacgaactgtggctgcaccatctgtcttcatcttcccgccatctgatg
 agcagttgaaatctggaactgcctctgttgtgtgacctgtgaataacttctatcccagaga
 ggccaaagtacagtgggaaggtggataacgcc

50

55

208 Light Chain Protein

DIQMTQSPSSLSASVGDRTTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVP
SRFSGSGSGTKFTLTISLQPEDFATYYCLOHNTYPWTFGQGTKVEIR

5

Antibody U1-16

209 Heavy Chain DNA:

H3_221_1N1G1

accatgaaacatctgtggttcttctctctgctggtggcagctcccagatgggtcctgtccc
aggtgcagctgcaggagtcgggcccaggactggtgaagccttcacagaccctgtccctcac
ctgcactgtctctggtggctccatcagcagtggtgattactactggagctggatccgccag
caaccagggaaagggcctggagtggttgggtacatctattacagtgggagcacctactaca
acccgtccctcaagagtcgagttaccatatacagtagacacgtctaagaaccagttctccct
gaagctgagctctgtgactgccgcggacacggccgtgtattactgtgcgagagcggattac
gatttttggagtggttattttgactactggggccagggaatccctgggtcaccgtctctcag
cctccaccaagggcccatcgggtcttccccctggcaccctcctccaagaacacctctggggg
cacagcggccctgggctgctggtcaaggactacttccccgaaccggtgacggtgtcctgg
aactcaggcgccctg

10

15

20

210 Heavy Chain Protein

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTYY
NPSLKSRTVISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWQGILVTVSS

25

211 Light Chain DNA:

H3_221_1N1K

atgaggggtccccgctcagctcctggggctcctgctgctctggttcccagggtgccagggtgt
gacatccagatgacccagctctccatcctccctgtctgcatctgttaggagacagagtcacc
atcaacttgccgggcaagtccagggcattagaaatgatttaggctgggtatcagcagaaacca
gggaaagccccctaagcgctgatctatgctgcatccagtttgcaaagtgggggtcccatca
aggttcagcggcagtggtctgtggacagaattcactctcacaatcagcagcctgcagcct
gaagattttgcaacttattactgtctacagcataatagttaccogtggaagcttcggccaa
gggaccaaggtggaaatcaaacgaactgtggctgcaccatctgtcttcatcttcccgcca
tctgatgagcagttgaaatctggaactgcctctgttgtgtgctgctgaataactctat
cccagagaggccaaagtacagtggaaggtggataacgcc

30

35

212 Light Chain Protein

DIQMTQSPSSLSASVGDRTTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPSR
FSGSGSGTEFTLTISLQPEDFATYYCLOHNSYPWTFGQGTKVEIK

40

45

Antibody U1-17

213 Heavy Chain DNA:

H3_224_1N1G1

tggttcttctctctgctggtggcagctcccagatgggtcctgtcccagggtgcagctgcagg
agtcgggcccaggactggtgaagccttcacagaccctgtccctcacctgcactgtctctgg
tggctccatcagcagtggtgattactactggagctggatccgccagcaccagggaaagggc
ctggagtggttggatacatctattacagtgggagcacctactacaattcgctccctcaaga
gtcgagttaccatatacagtagacacgtctaagaaccagttctccctgaagctgagctctgt
gactgccgcggacacggccgtgtattactgtgcgagagcggattacgatttttggagtggt
tattttgactactggggccagggaaccctgggtcaccgtctcctcagcctccaccaagggcc
catcg

50

55

214 Heavy Chain Protein

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTYY
NSSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGGQTLTVSS

215 Light Chain DNA:

H3_224_1N1K

ggtgccaggtgtgacatccagatgacccagtcctccatccctccctgtctgcatctgtaggag
acagagtcaccatcacttgccgggcaagtcagggcattagaaatgatttaggctggtatca
gcagaaacctgggaaagccctaaagcgctgatctatgctgcatccagtttgcaaagtggg
gtccatcaagggttcagcggcagtggtatctgggacagaattcactctcacaatcagcagcc
tgcagcctgaagatcttgcaacttattactgtctacagcacaatagttaccctgggacgtt
cggccaaagggaaccaaggtggaaatcaaacgaactgtggctgcaccatctgtcttcattctc
ccgcca

216 Light Chain Protein

DIQMTQSPSSLSASVGDRTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
RFGSGSGTEFTLTISSLQPEDFATYYCLQHNSYPWTFGQGTKVEIK

Antibody U1-18

217 Heavy Chain DNA:

H3_227_1N1G1

aggttcttctctctgtctggtggcagctcccagatgggtccctgtcccaggtgcagctgcagg
agtcgggcccaggactggtgaagccttcacagaccctgtccctcactgcactgtctctgg
tggctccatcagcagtggtgattactactggagctggatccgccagcaccaggggaagggc
ctggagtggattggatacatctattacagtgggagcacctactacaacccgtccctcaaga
gtcgagttaccatatcagtagacacgtctaaagaaccagttctccctgaagctgagctctgt
gactgccgcggacacggccgtgtattactgtgcgagagccgattacgatttttggagtggg
tattttgactactggggccagggaaccctggtcacctgtccctcagcctccaccaagggcc
catcggtcttccccctggcaccctcctccaagagcacctctgggggacagcggccctggg
ctgcctgggcaaggactacttccccgaaccggtgacggtgtcgtggaactcaggcgccct

218 Heavy Chain Protein

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTYY
NPSLKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGYFDYWGGQTLTVSS

219 Light Chain DNA:

H3_227_1N1K

atgaggggtccccgctcagctcctggggctcctgtctgtctgtggttcccaggtgccaggtgtga
catccagatgacccagtcctccatccctccctgtctgcatctgtaggagacagagtcaccatca
cttgccggggcaagtcagggcattagaaatgatttaggctggtatcagcagaaaccaggga
gcccctaaagcgctgatctatgctgcatccagtttgcaaagtgggtcccatcaagggttcag
cggcagtggtatctgggacagaattcactctcacaatcagcagcctgcagcctgaagatcttg
caacttattactgtctacagcataatagttaccctgggacgttcggccaagggaaccaaggtg
gaaatcaaacgaactgtggctgcaccatctgtcttcattctcccgccatctgatgagcagtt
gaaatctggaactgcctctgtgtgtgcctgctgaataacttctatcccagagagggccaaag
tacagtggaaggtggataacg

220 Light Chain Protein

DIQMTQSPSSLSASVGDRTITCRASQGIRNDLGWYQQKPGKAPKRLIYAASSLQSGVPS
RFGSGSGTEFTLTISSLQPEDFATYYCLQHNSYPWTFGQGTKVEIK

Antibody U1-33

221 Heavy Chain DNA:

H4_14_1N1G4

ctgtgggttcttctctctgctgggtggcagctcccagatgggtcctgtcccaggtgcagctgc
 aggagtcggggcccaggactgggtgaagccttcacagaccctgtccctcacctgcactgtctc
 tgggtggctccatcagcagtggtgattactactggagctggatccgccagcaccagggaag
 ggcctggagtggttgggtacatctattacagtgaggagcactactacaaccctgcccctca
 agagtcgagttaccatgtcagtagacacgtctaagaaccagttctccctgaagctgagctc
 tgtgactgcgcggacacggccgtgtattactgtgcgagagccgattacgatttttgagtg
 ggtcacttttgactgctggggccagggaaccctgggtcacctgtctctcagcttccaccaagg
 gcccctccgtcttccccc

222 Heavy Chain Protein

QVQLQESGPGLVKPSQTLSTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGSTYY
 NPSLKSRVTMSVDTSKNQFSLKLSSVTAADTAVYYCARADYDFWSGHFDCWGQGTILVTVSS

223 Light Chain DNA:

H4_14_1N1K

atgaggggtcccgcctcagctcctggggctcctgtctgtctctgggtcccaggtgccaggtgtga
 catccagatgacccagtcctccatcctccctgtctgtcatctgttaggagacagagtcaccatca
 cttgccggggcaagtcagggcatttagagatgatttaggctgggtatcagcagaaaccaggga
 gcccttaagcgccctgatctatgtgaatccagtttgcaaagtgggggtcccatacaaggttcag
 cggcagtggtatctgggacagaattcaactctcacaatcagcagcctgcagcctgaagattttg
 caacttattactgtctacagcatcatagttaccctggacgttcggccaagggaaccaaggtg
 gaaatcaaacgaactgtggctgcaccatctgtcttcatcttcccgcc

224 Light Chain Protein

DIQMTQSPSSLSASVGDRTITCRASQGIRDDLGWYQQKPGKAPKRLIYAESSLQSGVPSR
 FSGSGSGTEFTLTISSLQPEDFATYYCLQHHSYPWTFGQGTKVEIK

Antibody U1-29

225 Heavy Chain DNA:

H4_107_1N1G4

tggctgagctgggttttccctcgcttctcttttaagaggtgtccagtggtcaggtgcagctgg
 tggagtcctgggggaggcgtgggtccagcctgggaggtccctgagactctcctgtgcagcgtc
 tggattcaccttcaatagctatgacatgcactgggtccgccaggctccaggcaaggggtg
 gagtgggtggcagttatatgggtatgatggaagtaataaatactatgcagactccgtgaagg
 gccgattcaccatctctagagacaattccaagaacacgctgtatctgcaaatgaacagcct
 gagagccgaggacacggctgtgtattactgtgcgagagaccgcttgtgtactaatggtgta
 tgctatgaagactacggtatggacgtctggggccaagggaaccaggtcacctgtctctcag
 cttccaccaagggcccatccgtcttccccctgggcgcctgtctccaggagcacctccgagag
 cacagccgcctgggc

226 Heavy Chain Protein

QVQLVESGGGVVQPGRSLRLSCAASGFTFNSYDMHWVRQAPGKGLEWVAVIWDGSKNYA
 DSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARDRLCTNGVCYEDYGMDVWGQGT
 TVSS

227 Light Chain DNA:

H4_107_1N1K

atgaggggtccctgtctcagctcctggggctcctgtctgtctctggctctcaggtgccagatgtga
 catccagatgacccagtcctccatcctccctgtctgtcatctgttaggagacagagtcaccatca

cltgcagggcgagtcaggacattagcaactatTTTaaattgggtatcagcagaaaccagggaaa
 gcccttaaggtcctgatctacgatgcattccaatttggaaacaggggtcccatcaaggttcag
 tggagtggtatctgggacagattttactttaccatcagcagcctgcagcctgaagatgttg
 caacataattactgtcaacactatgatactctcccgtcactttcggcggaggggaccaaggtg
 gagatcaaacgaactgtgggtgcaccatctgtcttcactctcccggccatctgatgagcagtt
 gaaatctggaactgcctctgttgtgtgcctgctgaataacttctatcccagagaggccaaag
 tacagtgg

228 Light Chain Protein

DIQMTQSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPGKAPKVLIIYDASNLETGVPSR
 FSGSGSGTDFTFTISSLPEDVATYYCQHYDTLPITFGGGTKVEIK

Antibody U1-30

229 Heavy Chain DNA:

H4_116_1_1N1G4

ggactgtgcaagaacatgaaacacctgtggttcttccctcctgctggtggcagctcccagatg
 ggtcctgtcccaggtgcagctgcaggagtcggggccagggaactggtgaagcctttacagaccc
 tgtccctcacctgcactgtctctggtggctccatcagcagtggtgattactactggagctgg
 atccgccagcaccaggggaagggcctggagtggtattgggtacatctattacagtgggaccac
 ctactacaaccgtccctcaagagtcgagttaccatatacagtagacacgtctaagaaccagt
 tcgccctgaagctgaactctgtgactgccggcgacacggccgtgtattactgtgcgagagcc
 gattacgatttttggagtggttattttgactactggggccagggaaccctggtcacgctctc
 ctacagctccaccaagggcccatcctcttccccctgg

230 Heavy Chain Protein

QVQLQESGPGLVKPLQLTSLTCTVSGGSISSGDYYWSWIRQHPGKGLEWIGYIYYSGT'TYY
 NPSLKSRTISVDTSKNQFALKLNSVTAADTAVYYCARADYDFWSGYFDYWQGTLVTVSS

231 Light Chain DNA:

H4_116_1_1N1K

atgaggggtccctgctcagctcctggggctcctgctgctctggttcccaggtgccaggtgtg
 acatccagatgacccagtcctccatcctccctgtctgcattctgtaggagacagagtcacccat
 cacttgccgggaggtcagggcattagaaatgatttaggctggtatcagcagaaaccaggg
 aaagccctcagcgcctgatctatgctgcattccagtttgcaaagtgggtcccataaggt
 tcagcggcagtggtatctgggacagaattctctctcacaatctccagcctgcagcctgaaga
 ttttgcaacttattactgtctacagcataatagttaccctggagcttcggccaagggacc
 aaggtggaaatcaaacgaactgtgggtgcaccatctgtcttcattctcccggccatctgatg
 agcagttgaaatctggaactgcctctgttgtgtgcctgctgaataacttctatcccagaga
 ggccaaagtacagtgggaaggtggataacgccttccaatcggg

232 Light Chain Protein

DIQMTQSPSSLSASVGDRVTITCRAGQGIRNDLGWYQQKPGKAPQRLIYAASSLQSGVPSR
 FSGSGSGTEFSLTISLQPEDFATYYCLQHNSTPWFQGGTKVEIK

[Chem. 15]

[0181]

Sequence List of ODR

Chain antibodies	Pat. number:	CDR1	CDR2	CDR3
Heavy	U1-1	GGINSISGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY

EP 4 494 710 A2

(continued)

	Sequence List of ODR				
5	Chain antibodies	Pat. number:	CDR1	CDR2	CDR3
	Light		RASQGIRNDLG	AASSLQS	LQHNSYPWT
	Heavy	U1-2	GGSISSGDYYWS	YIYYSGSTYYNPSLRS	ADYDFWSGYFDY
10	Light		RASQGIRNDLG	AASSLQS	LQHNGYPWT
	Heavy	U1-3	GGSISSGGYYWS	YIYYSGSTYYNPSLKS	DGYDSSGGYYHGYFDY
	Light		KSSQSVLYSSNNKNYLA	WASTRES	QQYYSTPLT
	Heavy	U1-4	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
15	Light		RASQGIRNDLG	AASSLQS	LQHNNYPWT
	Heavy	U1-5	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQGIRNDLG	AASSLQS	LQHNTYPWT
	Heavy	U1-6	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWNGYFDY
20	Light		RASQGIRNDLG	AASSLQS	LQHNTYPWT
	Heavy	U1-7	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQDIRNDLG	AASSLQS	LQHNSYPWT
25	Heavy	U1-8	GYTLTELSMY	GFPEDGETIYAQKFQG	GWNYVFDY
	Light		RSSQSLLHSNGYNYLD	LDSHRAS	MQALQTPLT
	Heavy	U1-9	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWNGYFDY
	Light		RASQDIRNDLG	AASSLQS	LQHNSYPWT
30	Heavy	U1-10	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQGIRNDLG	AASSLQS	LQHNNYPWT
	Heavy	U1-11	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
35	Light		RASQGIRNDLG	AASSLQS	LQHNTYPWT
	Heavy	U1-12	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQGIRNDLG	AASSLQS	LQHNNYPWT
	Heavy	U1-13	GGSISSGGYYWS	YIYYSGSTYYNPSLKS	EDDGMDV
40	Light		RSSQSLLHSNGYNYLE	LGSNRAS	MQALQTPIT
	Heavy	U1-14	GGSISSGDYYWS Y	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQGIRNDLG	AASSLQS	LQHNTYPWT
45	Heavy	U1-15	GGSVSSGGYYWS	YIYYSGSTNYNPSLKS	DGDVDTAMVDAFDI
	Light		RASQSLSGNYLA	GASSRAT	QQYDRSPLT
	Heavy	U1-16	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQGIRNDLG	AASSLQS	LQHNSYPWT
50	Heavy	U1-17	GGSISSGDYYWS	YIYYSGSTYYNSSLKS	ADYDFWSGYFDY
	Light		RASQGTRNDLG	AASSLQS	LQHNSYPWT
	Heavy	U1-18	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQGIRNDLG	AASSLQS	LQHNSYPWT
55	Heavy	U1-19	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	GDYDFWSGEFDY
	Light		sequence not available		

EP 4 494 710 A2

(continued)

	Sequence List of ODR				
5	Chain antibodies	Pat. number:	CDR1	CDR2	CDR3
	Heavy	U1-20	GGSISSGGYYWS	YIYDSGSTYYNPSLKS	DQGGDGYSGYGYYYGM DV
	Light		QASQDISNYLN	VASNLET	QQCDNLPLT
10	Heavy	U1-21	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQDIRNDLG	AASRLQS	LQHNSYPWT
	Heavy	U1-22	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQGIRNDLG	AASSLQN	LQHNSYPWT
15	Heavy	U1-23	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQGIRNDLG	AASSLQS	LQHNSYPWT
	Heavy	U1-24	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWNGYFDY
20	Light		RASQGIRNDLG	AASSLQS	LQHNNYPWT
	Heavy	U1-25	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDY
	Light		RASQGIRNDLG	AASSLQN	LQHNSYPWT
25	Heavy	U1-26	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDF
	Light		RASQGIRNULG	AASSLQS	LQHNGYPWT
	Heavy	U1-27	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDF
	Light		RASQGIRNDLG	AASSLQS	LQHNGYPWT
30	Heavy	U1-28	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGYFDS
	Light		RASQGIRNDLG	AASSLQS	LQHNGYPWT
	Heavy	U1-29	GFTFNSYDMH	VIWYDGSNKYYADSVKG	DRLCTNGVCYEDYGMDV
35	Light		QASQDISNYLN	DASNLET	QHYDTLPLT
	Heavy	U1-30	GGSISSGDYYWS	YIYYSGTTYNPSLKS	ADYDFWSGYFDY
	Light		RAGQGIRNDLG	AASSLQS	LQHNSYPWT
	Heavy	U1-31	GYTFTNYGIS	WISAYDGYRNYAQLQG	DVQDYGDYDYFDY
40	Light		RASQSISSYLN	AASSLQS	QQSYSTPIT
	Heavy	U1-32	GGSISSGDYYWS	YIYYSGTTYNPSLKS	ADYDFWSGYFDY
	Light		RAGQGIRNDLG	AASSLQS	LQHNSYPWT
	Heavy	U1-33	GGSISSGDYYWS	YIYYSGSTYYNPSLKS	ADYDFWSGHFDC
45	Light		RASQGIRDDL	AESSLQS	LQHNSYPWT
	Heavy	U1-34	GYTFTNYGIS	WISAYDGYRNYAQLQG	DVQDYGDYDYFDY
	Light		RASQSISSYLN	AASSLQS	QQSYSTPIT
50	Heavy	U1-35	GFTFSDYYMS	YISSSGNNIYHADSVKG	ERYSGYDDPDGFDI
	Light		QASQDISNYLS	DASNLET	QQYDNPPCS
	Heavy	U1-36	GGSISSGYYYWS	YIYYSGTTYNPSFKS	ADYDFWSGHFDY
	Light		RASQGIRNDLG	AASSLQS	LOHNSYPWT
55	Heavy	U1-37	GYTFTSYGIS	WISAYDGHTNYAQLQG	DPHDYSNIEAFDF
	Light		RASQSISSYLN	AASSLQS	QQSYSTPIT
	Heavy	U1-38	GFSLSGSGVGVG	LIYWNDDKRYSPSLKS	RDEVRGFDY

EP 4 494 710 A2

(continued)

	Sequence List of ODR				
5	Chain antibodies	Pat. number:	CDR1	CDR2	CDR3
	Light		RSSQSLVYSDGYTYLH	KVSNWDS	MQGAHWPIT
	Heavy	U1-39	GFTVSSNYMS	VIYSGGSTYYADSVKG	GQWLDV
10	Light		RSSQSLLHSNGYNYLD	LGFHRAS	RQALQTPLT
	Heavy	U1-40	GGSISSGGYYWS	YIYSSGSTYYNPSLKS	DRELELYYYYYGMDV
	Light		RSSQSLLYSNGYNYLD	LGSNRAS	MQALQTPLT
	Heavy	U1-41	GGSISSGGYYWS	YIYSGSTYYNPSLKS	DRELEGYSNYYGVDV
15	Light		RASQAISNYLN	AASSLQS	QQNNSLPIT
	Heavy	U1-42	GYSFTSYWIG	IIYPGDS DTRYSPSFQG	HENYGDYNY
	Light		RASQSIRSYLN	AASSLQS	QQSHGSPLT
	Heavy	U1-43	GGSISSGGYYWS	YIYYSGSTYYNPSLRS	DREREWDYDGPQGMDV
20	Light		RASQSISSYLH	AASSLQS	QQSYSNPLT
	Heavy	U1-44	GYSFTSYWIG	IIWPGDS DTYSPSFQG	HENYGDYNY
	Light		RASQSIRSYLN	AASSLQS	QQSISSPLT
25	Heavy	U1-45	GYTFTSYDIN	WMNPNSGDTGYAQVFQG	FGDLPYDYSYEWFD P
	Light		RASQSISSYLN	AASSLQS	QQSYSTPLT
	Heavy	U1-46	GDSVSSNSAAWN	RTYYRSKWYNDYAVSVK S	DLYDFWSGYPYYYYGMDV
30	Light		sequence not available		
	Heavy	U1-47	GDSVSSNSAAWN	RTYYRSKWYNDYAVSVK S	DYYGSGSFYYYYGMDV
35	Light		RASQSISSYLN	AASN LQS	QQSYSTPRT
	Heavy	U1-48	GGSISSYYWS	HIYTS GSTNYNPSLKS	EAIFGVGPYYYYGMDV
	Light		sequence not available		
	Heavy	U1-49	GYTFTGYMH	WINPNIGGTNCAQKFQG	GGRYSSSWSYYYYGMDV
40	Light		KSSQSLLLSDGGTYLY	EVS NRFS	MQSMQLPIT
	Heavy	U1-50	GGSVSSGGYYWS	YIYSGSTNYNPSLKS	GGDSNYEDYYYYY GMDV
	Light		RASQSISIY LH	AASSLQS	QQSYTSPIT
45	Heavy	U1-51	GGSISSYYWS Y	YIYSGSTNYNPSLKS D D	JSSYYDSSGYLYYYAM DV
	Light		KSSQSVLYSSNNKNYLA	WASTRES	QQYYTTPLT
	Heavy	U1-52	GGSISSGGYYWS N	NYYS GSTYYNPSLKS	GGTGTNYYYYY GMDV
	Light		RASQSVSSSYLA	GASSWAT	QQYGSSPLT
50	Heavy	U1-53	GFTFSIYSMN	YISSSSSTIYYADSVKG	DRGDFDAFDI
	Light		QASQDITNYLN	DASNLET	QQCENFPIT
	Heavy	U1-55.1	GGSVSSGGYYWN	YINYS GSTNYNPSLKS	DRELELYYYYYGMDV
55	Light		Identical with U1-55		
	Heavy	U1-55	Identical with U1-55.1		
	Light		RSSQSLLYSNGYKYLD	LGSNRAS	MQALQTPIT

(continued)

Sequence List of ODR				
Chain antibodies	Pat. number:	CDR1	CDR2	CDR3
Heavy	U1-57.1	Identical with U1-57		
Light		RSSQSLLYSNGYKYLD	LGSNRAS	MQALQTPIT
Heavy	U1-57	GGSVSSGGYYWH	YINYSGSTNYNPSLKS	DRELELYYYYYGMDV
Light		Identical with U1-57.1		
Heavy	U1-58	GFTFSSYGMII	VIWYDGSNKYYADSVKG	AARLDYYYYGMDV
Light		RASQSINSYLN	GASGLQS	QQSYSSPLT
Heavy	U1-59	GGSFSGYYWS	EINHSGSTNYNPSLKS	DKWTWYFDL
Light		RSSQSVLYSSSNRNYLA	WASTRES	QQYYSTPRT
Heavy	U1-61.1	GVSISSGGYYWS	YIYSGSTYYNPSLKS	DSESEYSSSSNYGMDV
Light		RASQTISSYLN	AASSLQG	QQSYSNPLT
Heavy	U1-61	GVSISSGGYYWS	YIYSGSTYYNPSLKS	DSESEYSSSSNYGMDV
Light		Identical with U1-61.1		
Heavy	U1-62	GYSFTSYWIG	IIYPGDS DTRYSPSFQG	QMAGNYYYYGMDV
Light		RASQSVISIYLA	GASSRAT	QQYGSSPCS

[0182] When a newly produced monoclonal antibody binds to a partial peptide or a partial tertiary structure to which the U1-49, U1-53, U1-59, U1-7, or U1-9 antibody binds, it can be determined that the antibody binds to the same epitope as the U1-49, U1-53, U1-59, U1-7, or U1-9 antibody. Further, by confirming that the antibody competes with the U1-49, U1-53, U1-59, U1-7, or U1-9 antibody for binding to HER3 (that is, the antibody inhibits the binding between the U1-49, U1-53, U1-59, U1-7, or U1-9 antibody and HER3), it can be determined that the antibody binds to the same epitope as the U1-49, U1-53, U1-59, U1-7, or U1-9 antibody even when the specific sequence or structure of an epitope is not defined. Once the epitope is confirmed to be the same, it is strongly expected that the antibody has a biological activity equivalent to that of the U1-49, U1-53, U1-59, U1-7, or U1-9 antibody.

[0183] According to the present invention, the binding protein of the invention interacts with at least one epitope in the extracellular part of HER3. The epitopes are preferably located in domain L1 (aa 19-184), which is the amino terminal domain, in domain S1 (aa 185-327) and S2 (aa 500-632), which are the two Cysteine-rich domains, in domain L2 (328-499), which is flanked by the two Cysteine-rich domains or in a combination of HER3 domains. The epitopes may also be located in combinations of domains such as but not limited to an epitope comprised by parts of L1 and S1. Moreover, the binding protein of the invention is further characterized in that its binding to HER3 reduces HER3-mediated signal transduction. In accordance with the present invention, a reduction of HER3-mediated signal transduction may, e.g. be caused by a downregulation of HER3 resulting in an at least partial disappearance of HER3 molecules from the cell surface or by a stabilization of HER3 on the cell surface in a substantially inactive form, i.e. a form which exhibits a lower signal transduction compared to the non-stabilized form. Alternatively, a reduction of HER3-mediated signal transduction may also be caused by influencing, e.g. decreasing or inhibiting, the binding of a ligand or another member of the HER family to HER3, of GRB2 to HER-2 or of GRB2 to SHC, by inhibiting receptor tyrosine phosphorylation, AKT phosphorylation, PYK2 tyrosine phosphorylation or ERK2 phosphorylation, or by decreasing tumor invasiveness. Alternatively, a reduction of HER3 mediated signal transduction may also be caused by influencing, e.g., decreasing or inhibiting, the formation of HER3 containing dimers with other HER family members. One example among others may be the decreasing or inhibiting of the HER3-EGFR protein complex formation.

[0184] Furthermore, in accordance with the present invention, minor variations in the amino acid sequences shown in SEQ ID NOs: 1-232 are contemplated as being encompassed by the present invention, providing that even the variations in the amino acid sequence still maintain at least 75 %, more preferably at least 80 %, 90 %, 95 %, and most preferably 99 % of the sequences shown in SEQ ID NOs: 1-232. The variations may occur within the framework regions (i.e. outside the CDRs), within the CDRs, or within the framework regions and the CDRs. Preferred variations in the amino acid sequences shown in SEQ ID NOs: 1-232, i.e. deletions, insertions and/or replacements of at least one amino acid, occur near boundaries of functional domains. Structural and functional domains can be identified by comparison of the nucleotide and/or amino acid sequence data to public or proprietary sequence databases. Computerized comparison methods can

be used to identify sequence motifs or predicted protein conformation domains that occur in other binding proteins of known structure and/or function. Methods to identify protein sequences that fold into a known three-dimensional structure are known. See e.g. Bowie et al, Science 253, 164 (1991); Proteins, Structures and Molecular Principles (Creighton, Ed., W. H. Freeman and Company, New York (1984)); Introduction to Protein Structure (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton et al., Nature 354, 105 (1991), which are all incorporated herein by reference. Thus, those of skill in the art can recognize sequence motifs and structural conformations that may be used to define structural and functional domains in accordance with the invention. Among antibodies obtained by combining heavy and light chains having variations in such an amino acid sequences, an antibody equivalent to the original antibody (parent antibody) or more excellent than a parent antibody may be selected. As mentioned above, the HER3-binding protein, the anti-HER3 antibody, and the like of the present invention maintain the HER3-binding activity even if having variations in their amino acid sequences.

[0185] In the present invention, the term "homology" has the same meaning as the "identity. The homology between two amino acid sequences can be determined using default parameters of Blast algorithm version 2.2.2 (Altschul, Stephen F., Thomas L. Madden, Alejandro A. Schaeffer, Jinghui Zhang, Zheng Zhang, Webb Miller, and David J. Lipman (1997), "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs", Nucleic Acids Res. 25: 3389-3402). The Blast algorithm can be used also through the Internet by accessing the site www.ncbi.nlm.nih.gov/blast.

[0186] The chimeric antibody, humanized antibody, or human antibody obtained by the aforementioned method can be subjected to a known method for evaluating the binding property to an antigen for selecting preferable antibodies.

[0187] In the anti-HER3 antibody of the present invention, MEHD-7945A (or duligotuzumab), RG-7116, MM-111, MM-121 (or seribantumab, MM-141, LJM-716, huHER3-8, tri-specific anti-EGFR/ErbB3 zyboby, GSK-2849330, REGN-1400, KTN-3379, AV-203, monospecific surrobody (ErbB3), lumretuzumab, MP-EV-20, ZW-9, Dimercept™, anti-Erb3 surrobody (SL-175 or SL-176), SYM-013, variants, active fragments, modified products thereof, and the like are also included.

[0188] As one example of another index for use in the comparison of the properties of antibodies, the stability of antibodies can be exemplified. The differential scanning calorimetry (DSC) is a device capable of quickly and accurately measuring a thermal denaturation midpoint temperature (T_m) to be used as a favorable index of the relative conformational stability of proteins. By measuring the T_m values using DSC and comparing the values, a difference in thermal stability can be compared. It is known that the storage stability of antibodies shows some correlation with the thermal stability of antibodies (Lori Burton, et. al., Pharmaceutical Development and Technology (2007) 12, pp. 265-273), and a preferred antibody can be selected by using thermal stability as an index. Examples of other indices for selecting antibodies include the following features: the yield in an appropriate host cell is high; and the aggregability in an aqueous solution is low. For example, an antibody which shows the highest yield does not always show the highest thermal stability, and therefore, it is necessary to select an antibody most suitable for the administration to humans by making comprehensive evaluation based on the above-described indices.

[0189] The antibody of the present invention encompasses a modified product of the antibody. The modified variant refers to a variant obtained by subjecting the antibody of the invention to chemical or biological modification. Examples of the chemically modified variant include variants chemically modified by linking a chemical moiety to an amino acid skeleton, variants chemically modified with an N-linked or O-linked carbohydrate chain, etc. Examples of the biologically modified variant include variants obtained by modification after translation (such as N-linked or O-linked glycosylation, N- or C-terminal processing, deamidation, isomerization of aspartic acid, or oxidation of methionine), and variants in which a methionine residue has been added to the N terminus by being expressed in a prokaryotic host cell. Further, an antibody labeled so as to enable the detection or isolation of the antibody or an antigen of the invention, for example, an enzyme-labeled antibody, a fluorescence-labeled antibody, and an affinity-labeled antibody are also included in the meaning of the modified variant. Such a modified variant of the antibody of the invention is useful for improving the stability and blood retention of the antibody, reducing the antigenicity thereof, detecting or isolating the antibody or the antigen, and so on.

[0190] Further, by regulating the modification of a glycan which is linked to the antibody of the invention (glycosylation, defucosylation, etc.), it is possible to enhance an antibody-dependent cellular cytotoxic activity. As the technique for regulating the modification of a glycan of antibodies, International Publication WO 1999/54342, WO 2000/61739, WO 2002/31140, etc. are known. However, the technique is not limited thereto. In the antibody of the invention, an antibody in which the modification of a glycan is regulated is also included.

[0191] In the case where an antibody is produced by first isolating an antibody gene and then introducing the gene into an appropriate host, a combination of an appropriate host and an appropriate expression vector can be used. Specific examples of the antibody gene include a combination of a gene encoding a heavy chain sequence of an antibody and a gene encoding a light chain sequence thereof described in this specification. When a host cell is transformed, it is possible to insert the heavy chain sequence gene and the light chain sequence gene into the same expression vector, and also into different expression vectors separately.

[0192] In the case where eukaryotic cells are used as the host, animal cells, plant cells, and eukaryotic microorganisms can be used. As the animal cells, mammalian cells, for example, simian COS cells (Gluzman, Y., Cell, (1981) 23, pp.

175-182, ATCC CRL-1650), murine fibroblasts NIH3T3 (ATCC No. CRL-1658), and dihydrofolate reductase-deficient strains (Urlaub, G. and Chasin, L. A., Proc. Natl. Acad. Sci. USA (1980) 77, pp. 4126-4220) of Chinese hamster ovarian cells (CHO cells; ATCC: CCL-61) can be exemplified.

[0193] In the case where prokaryotic cells are used, for example, *Escherichia coli* and *Bacillus subtilis* can be exemplified.

[0194] By introducing a desired antibody gene into these cells through transformation, and culturing the thus transformed cells in vitro, the antibody can be obtained. In the above-described culture method, the yield may sometimes vary depending on the sequence of the antibody, and therefore, it is possible to select an antibody which is easily produced as a pharmaceutical by using the yield as an index among the antibodies having an equivalent binding activity. Therefore, in the antibody of the invention, an antibody obtained by a method of producing an antibody, characterized by including a step of culturing the transformed host cell and a step of collecting a desired antibody or a functional fragment of the antibody from a cultured product obtained in the culturing step is also included.

[0195] It is known that a lysine residue at the carboxyl terminus of the heavy chain of an antibody produced in a cultured mammalian cell could be deleted/eliminated (Journal of Chromatography A, 705: 129-134 (1995)), and it is also known that two amino acid residues (glycine and lysine) at the carboxyl terminus of the heavy chain of an antibody produced in a cultured mammalian cell could be deleted/eliminated and a proline residue newly located at the carboxyl terminus could be amidated (Analytical Biochemistry, 360: 75-83 (2007)). However, such deletion/elimination and modification of the heavy chain sequence do not affect the antigen-binding affinity and the effector function (the activation of a complement, the antibody-dependent cellular cytotoxicity, etc.) of the antibody. Therefore, in the antibody of the invention, an antibody and a functional fragment of the antibody subjected to such modification are also included, and a deletion variant in which one or two amino acids have been deleted at the carboxyl terminus of the heavy chain, a variant obtained by amidation of the deletion variant (for example, a heavy chain in which the carboxyl terminal proline residue has been amidated), and the like are also included. The type of deletion variant having a deletion at the carboxyl terminus of the heavy chain of the antibody according to the invention is not limited to the above variants as long as the antigen-binding affinity and the effector function are conserved. The two heavy chains constituting the antibody according to the invention may be of one type selected from the group consisting of a full-length heavy chain and the above-described deletion variant, or may be of two types in combination selected therefrom. The ratio of the amount of each deletion variant can be affected by the type of cultured mammalian cells which produce the antibody according to the invention and the culture conditions, however, a case where one amino acid residue at the carboxyl terminus has been deleted in both of the two heavy chains contained as main components in the antibody according to the invention can be exemplified. The scope of the whole antibody (in the present invention, also simply referred to as an "antibody") of the present invention also includes deletion variants thereof, mixtures containing one or two or more deletion variants thereof, etc. The "antibody" of the present invention includes an antibody comprising a heavy or light chain in which N-terminal glutamate is in the form of pyroglutamate by cyclization and/or a heavy or light chain in which a portion of cysteine residues are in the form of cysteinyl.

[0196] In a preferred embodiment of the present invention, the anti-HER3 antibody of the invention is of the IgA, IgD-, IgE₁ IgG- or IgM-type, preferably of the IgG- or IgM-type including, but not limited to, the IgG1-, IgG2-, IgG3-, IgG4-, IgM1- and IgM2-type. In most preferred embodiments, the antibody is of the IgG1-, IgG2- or IgG4- type.

[0197] As the biological activity of the antibody, generally, an antigen-binding activity, activity of internalizing an antigen in cells expressing the antigen by binding with the antigen, an activity of neutralizing the activity of an antigen, an activity of enhancing the activity of an antigen, an antibody-dependent cellular cytotoxicity (ADCC) activity, a complement-dependent cytotoxicity (CDC) activity, and an antibody-dependent cell-mediated phagocytosis (ADCP) can be exemplified. The function of the antibody according to the invention is a binding activity to HER3, preferably, activity of internalizing HER3 in HER3 expressing cells by binding with HER3. Further, the antibody of the invention may have an ADCC activity, a CDC activity and/or an ADCP activity in addition to the cell internalization activity.

[0198] In certain respects, e.g. in connection with the generation of antibodies as therapeutic candidates against HER3, it may be desirable that the anti-HER3 antibody of the invention is capable of fixing complement and participating in complement-dependent cytotoxicity (CDC). There are a number of isotypes of antibodies that are capable of the same including without limitations the following: murine IgM, murine IgG2a, murine IgG2b, murine IgG3, human IgM, human IgG1, human IgG3, and human IgA. It will be appreciated that antibodies that are generated need not initially possess such an isotype but, rather the antibody as generated can possess any isotype and the antibody can be isotype switched by appending the molecularly cloned V region genes or cDNA to molecularly cloned constant region genes or cDNAs in appropriate expression vectors using conventional molecular biological techniques that are well known in the art and then expressing the antibodies in host cells using techniques known in the art. The isotype-switched antibody may also possess an Fc region that has been molecularly engineered to possess superior CDC over naturally occurring variants (Idusogie et al., J Immunol., 166, 2571-2575) and expressed recombinantly in host cells using techniques known in the art. Such techniques include the use of direct recombinant techniques (see e.g. U.S. Patent No. 4,816,397), cell-cell fusion techniques (see e.g. U.S. Patent Nos. 5,916,771 and 6,207,418), among others. In the cell-cell fusion technique, a myeloma or other cell line such as CHO is prepared that possesses a heavy chain with any desired isotype and another

myeloma or other cell line such as CHO is prepared that possesses the light chain. Such cells can, thereafter, be fused and a cell line expressing an intact antibody can be isolated. By way of example, a human anti-HER3 IgG4 antibody, that possesses the desired binding to the HER3 antigen, could be readily isotype switched to generate a human IgM, human IgG1 or human IgG3 isotype, while still possessing the same variable region (which defines the antibody's specificity and some of its affinity). Such molecule might then be capable of fixing complement and participating in CDC.

[0199] Moreover, it may also be desirable for the anti-HER3 antibody of the invention to be capable of binding to Fc receptors on effector cells, such as monocytes and natural killer (NK) cells, and participate in antibody-dependent cellular cytotoxicity (ADCC). There are a number of isotypes of antibodies that are capable of the same, including without limitations the following: murine IgG2a, murine IgG2b, murine IgG3, human IgG1 and human IgG3. It will be appreciated that antibodies that are generated need not initially possess such an isotype but, rather the antibody as generated can possess any isotype and the antibody can be isotype switched by appending the molecularly cloned V region genes or cDNA to molecularly cloned constant region genes or cDNAs in appropriate expression vectors using conventional molecular biological techniques that are well known in the art and then expressing the antibodies in host cells using techniques known in the art. The isotype-switched antibody may also possess an Fc region that has been molecularly engineered to possess superior ADCC over naturally occurring variants (Shields et al. J Biol Chem., 276, 6591-6604) and expressed recombinantly in host cells using techniques known in the art. Such techniques include the use of direct recombinant techniques (see e.g. U.S. Patent No. 4,816,397), cell-cell fusion techniques (see e.g. U.S. Patent Nos. 5,916,771 and 6,207,418), among others. In the cell-cell fusion technique, a myeloma or other cell line such as CHO is prepared that possesses a heavy chain with any desired isotype and another myeloma or other cell line such as CHO is prepared that possesses the light chain. Such cells can, thereafter, be fused and a cell line expressing an intact antibody can be isolated. By way of example, a human anti-HER3 IgG4 antibody, that possesses the desired binding to the HER3 antigen, could be readily isotype switched to generate a human IgG1 or human IgG3 isotype, while still possessing the same variable region (which defines the antibody's specificity and some of its affinity). Such molecule might then be capable of binding to FcγR on effectors cells and participating in ADCC.

[0200] The obtained antibody can be purified to be homogeneous. The separation and purification of the antibody may be performed employing a conventional protein separation and purification method. For example, the antibody can be separated and purified by appropriately selecting and combining column chromatography, filter filtration, ultrafiltration, salt precipitation, dialysis, preparative polyacrylamide gel electrophoresis, isoelectric focusing electrophoresis, and the like (Strategies for Protein Purification and Characterization: A Laboratory Course Manual, Daniel R. Marshak et al. eds., Cold Spring Harbor Laboratory Press (1996); Antibodies: A Laboratory Manual. Ed Harlow and David Lane, Cold Spring Harbor Laboratory (1988)), but the method is not limited thereto.

[0201] Examples of such chromatography include affinity chromatography, ion exchange chromatography, hydrophobic chromatography, gel filtration chromatography, reverse phase chromatography, and adsorption chromatography.

[0202] Such chromatography can be performed employing liquid chromatography such as HPLC or FPLC.

[0203] As a column to be used in affinity chromatography, a Protein A column and a Protein G column can be exemplified. For example, as a column using a Protein A column, Hyper D, POROS, Sepharose FF (Pharmacia Corp.) and the like can be exemplified.


[0204] Further, by using a carrier having an antigen immobilized thereon, the antibody can also be purified utilizing the binding property of the antibody to the antigen.

{Antitumor compound}

[0205] The antitumor compound to be conjugated to the anti-HER3 antibody-drug conjugate of the present invention is explained. The antitumor compound used in the present invention is not particularly limited if it is a compound having an antitumor effect and a substituent or a partial structure allowing connecting to a linker structure. When a part or whole linker of the antitumor compound is cleaved in tumor cells, the antitumor compound moiety is released to exhibit the antitumor effect. As the linker is cleaved at a connecting position with a drug, the antitumor compound is released in its unmodified structure to exhibit its intrinsic antitumor effect.

[0206] As an antitumor compound used in the present invention, exatecan, a camptothecin derivative ((1S,9S)-1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1H,12H-benzo[de]pyrano[3',4'-6,7]indolizino[1,2-b]quino-
lin-10,13(9H,15H)-dione shown in the following formula) can be preferably used.

[Chem. 16]



[0207] Although having an excellent antitumor effect, exatecan has not been commercialized as an antitumor drug. The compound can be easily obtained by a known method and the amino group at position 1 can be preferably used as a connecting position to the linker structure. Further, exatecan can be also released in tumor cells while part of the linker is still attached thereto. However, it is an excellent compound exhibiting an excellent antitumor effect even in such structure.

[0208] Because exatecan has a camptothecin structure, it is known that the equilibrium shifts to a structure with a closed lactone ring (closed ring) in an acidic aqueous medium (for example, pH 3 or so) but it shifts to a structure with an open lactone ring (open ring) in a basic aqueous medium (for example, pH 10 or so). A drug conjugate being introduced with an exatecan residue corresponding to the closed ring structure and the open ring structure is also expected to have the same antitumor effect and it is needless to say that any of them is within the scope of the present invention.

[0209] Examples of other antitumor compounds include doxorubicin, daunorubicin, mitomycin C, bleomycin, cyclocytidine, vincristine, vinblastine, methotrexate, platinum-based antitumor agent (cisplatin or derivatives thereof), taxol or derivatives thereof, and other camptothecin or derivatives thereof (antitumor agent described in Japanese Patent Laid-Open No. 6-87746).


[0210] With regard to the antibody-drug conjugate, the number of conjugated drug molecules per antibody molecule is a key factor having an influence on the efficacy and safety. Production of the antibody-drug conjugate is performed by defining the reaction condition including the amounts of use of raw materials and reagents for reaction so as to have a constant number of conjugated drug molecules, a mixture containing different numbers of conjugated drug molecules is generally obtained unlike the chemical reaction of a low-molecular-weight compound. The number of drugs conjugated in an antibody molecule is expressed or specified by the average value, that is, the average number of conjugated drug molecules. Unless specifically described otherwise as a principle, the number of conjugated drug molecules means an average value except in a case in which it represents an antibody-drug conjugate having a specific number of conjugated drug molecules that is included in an antibody-drug conjugate mixture having different number of conjugated drug molecules.

[0211] The number of exatecan molecules conjugated to an antibody molecule is controllable, and as an average number of conjugated drug molecules per antibody, about 1 to 10 exatecans can be bound. Preferably, it is 2 to 8, and more preferably 3 to 8. Meanwhile, a person skilled in the art can design a reaction for conjugating a required number of drug molecules to an antibody molecule based on the description of the Examples of the present application and can obtain an antibody-drug conjugate with a controlled number of conjugated exatecan molecules.

[0212] The antibody-drug conjugate of the present invention is unlikely to have an occurrence of aggregation, insolubility, fragmentation, or the like, even when the number of conjugated drug molecules per antibody molecule is increased.

{Linker structure}

[0213] With regard to the anti-HER3 antibody-drug conjugate of the present invention, the linker structure for conjugating an antitumor compound to the anti-HER3 antibody is explained. The linker has the following structure:




the antibody is connected to the terminal of L^1 (opposite terminal to which L^2 is connected), and the antitumor compound is connected to the carbonyl group of $-L^a-(CH_2)_{n^2}-C(=O)-$ moiety or the C terminal of L^P .

n^1 represents an integer of 0 to 6, preferably, an integer of 1 to 5, and more preferably 1 to 3.


1. L¹

[0214] L¹ is represented by a structure shown below:




[0215] In the above, n³ is an integer of 2 to 8, and "-(Succinimid-3-yl-N)-" has a structure represented by the following formula:

[Chem. 17]







[0216] Position 3 of the above partial structure is the connecting position to the anti-HER3 antibody. The connection to the antibody at position 3 is characterized by forming a thioether bond. The nitrogen atom at position 1 of the structure moiety is connected to the carbon atom of methylene which is present within the linker including the structure. Specifically, $-(\text{Succinimid-3-yl-N})-(\text{CH}_2)_{n^3}-\text{C}(=\text{O})-\text{L}^2-$ is a structure represented by the following formula (herein, "antibody -S-" is derived from an antibody).

[Chem. 18]




[0217] In the formula, n³ is an integer of 2 to 8, and preferably 2 to 5.

[0218] Specific examples of L¹ include the followings.



2. L²






[0219] L² has a structure represented by the following formula:



[0220] L² may not be present, and in such a case, L² is a single bond. In the drug-linker structure of the present invention,

in particular, L^P may be directly connected to a drug, and in such a case, L^2 is particularly preferably a single bond. n^4 is an integer of 1 to 6, and preferably 2 to 4. L^2 is connected to L^1 at its terminal amino group and is connected to L^P at the carbonyl group of the opposite terminal.

[0221] Specific examples of L^2 include the followings.



3. L^P

[0222] L^P is a peptide residue consisting of 2 to 7 amino acids. Specifically, it consists of an oligopeptide residue in which 2 to 7 amino acids are linked by a peptide bond. L^P is connected to L^2 at N terminal and it is connected to the amino group of -NH-(CH₂)_n¹-L^a-(CH₂)_n²-C(=O)- moiety of the linker at C terminal.

[0223] The amino acid constituting L^P is not particularly limited, and the examples thereof include an L- or a D-amino acid, preferably an L-amino acid. Further, it can be an amino acid having a structure such as beta-alanine, epsilon-aminocaproic acid, or gamma-aminobutyric acid in addition to an alpha-amino acid, further, it can be a non-natural type amino acid such as N-methylated amino acid.

[0224] Sequence of the amino acid of L^P is not particularly limited, but examples of the constituting amino acid include phenylalanine (Phe; F), tyrosine (Tyr; Y), leucine (Leu; L), glycine (Gly; G), alanine (Ala; A), valine (Val; V), lysine (Lys; K), citrulline (Cit), serine (Ser; S), glutamic acid (Glu; E), and aspartic acid (Asp; D). Among them, preferred examples include phenylalanine, glycine, valine, lysine, citrulline, serine, glutamic acid, and aspartic acid. Depending on the type of the amino acid, drug release pattern can be controlled. The number of the amino acid can be between 2 to 7.

[0225] Specific examples of L^P include the followings.



[0226] The "(D-)D" described above means D-aspartic acid. Examples of the particularly preferred L^P of the antibody-

drug conjugate of the present invention include -GGFG- and -DGGFG- peptide residue. Further, in the drug-linker structure of the present invention, L^P may be directly connected to the drug, and for such a case, preferred examples of L^P include a pentapeptide residue of -DGGFG-.

4. $L^a-(CH_2)n^2-C(=O)-$

[0227] L^a in $L^a-(CH_2)n^2-C(=O)-$ is a structure of -O- or a single bond. n^2 is an integer of 0 to 5, preferably, 0 to 3, and more preferably 0 or 1.

[0228] Examples of $L^a-(CH_2)n^2-C(=O)-$ include the followings.

-O-CH₂-C(=O)-,

-O-CH₂CH₂-C(=O)-,

-O-CH₂CH₂CH₂-C(=O)-,

-O-CH₂CH₂CH₂CH₂-C(=O)-,

-O-CH₂CH₂CH₂CH₂CH₂-C(=O)-,

-CH₂-C(=O)-,

-CH₂CH₂-C(=O)-,

-CH₂CH₂CH₂-C(=O)-,

-CH₂CH₂CH₂CH₂-C(=O)-,

-CH₂CH₂CH₂CH₂CH₂-C(=O)-.

[0229] Among them, those with

-O-CH₂-C(=O)-,

-O-CH₂CH₂-C(=O)-

or those in which L^a is a single bond and n^2 is 0 are preferable.

[0230] Specific examples of the linker structure represented by -NH-(CH₂) n^1 - L^a -(CH₂) n^2 -C(=O)- include the followings.

-NH-CH₂-C(=O)-,

-NH-CH₂CH₂-C(=O)-,

-NH-CH₂-O-CH₂-C(=O)-,

-NH-CH₂CH₂-O-C(=O)-,

-NH-CH₂CH₂-O-CH₂-C(=O)-,



-NH-CH₂CH₂CH₂-C(=O)-,

-NH-CH₂CH₂CH₂CH₂-C(=O)-,

-NH-CH₂CH₂CH₂CH₂CH₂-C(=O)-





[0231] Among them, the examples are more preferably the followings.

-NH-CH₂CH₂CH₂-C(=O)-,



[0232] As for the linker $-\text{NH}-(\text{CH}_2)_n-\text{L}^a-(\text{CH}_2)_n-\text{C}(=\text{O})-$, the chain length of 4 to 7 atoms is preferable, and more preferably, are those having the chain length of 5 or 6 atoms.


[0233] With regard to the anti-HER3 antibody-drug conjugate of the present invention, when it is transferred to the inside of tumor cells, it is thought that the linker moiety is cleaved and the drug derivative having a structure represented by $\text{NH}_2-(\text{CH}_2)_n-\text{L}^a-(\text{CH}_2)_n-\text{C}(=\text{O})-(\text{NH}-\text{DX})$ is released to express an antitumor action. Examples of the antitumor derivative exhibiting an antitumor effect by releasing from the antibody-drug conjugate of the present invention include an antitumor derivative having a structure moiety in which the terminal of the structure represented by $-\text{NH}-(\text{CH}_2)_n-\text{L}^a-(\text{CH}_2)_n-\text{C}(=\text{O})-$ of the linker is an amino group, and the particularly preferred include the followings.




[0234] Meanwhile, in case of $\text{NH}_2-\text{CH}_2-\text{O}-\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$, it was confirmed that, as the amination structure in the molecule is unstable, it again undergoes a self-degradation to release the following $\text{HO}-\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$. Those compounds can be also preferably used as a production intermediate of the antibody-drug conjugate of the present invention.

[0235] Further, in the drug-linker structure of the present invention, there arises a case in which L^P may be directly connected to the drug. In such a case, when the C terminal of L^P is glycine, the antitumor drug to be released is exatecan itself or a compound having glycine bonded to the amino group of exatecan.

[0236] For the antibody-drug conjugate of the present invention in which exatecan is used as a drug, the drug-linker structure moiety having the following structure



or



to which the antibody is connected is preferable. The conjugated number of these drug-linker structure moiety may be from 1 to 10 as the average conjugated number per antibody, preferably, 2 to 8, and more preferably 3 to 8.

- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{GGFG}-\text{NH}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{GGFG}-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{GGFG}-\text{NH}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{GGFG}-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{GGFG}-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{DGGFG}-\text{NH}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{DGGFG}-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{DGGFG}-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{GGFG}-\text{NH}-\text{CH}_2-\text{O}-\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{GGFG}-\text{NH}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{NH}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{GGFG}-\text{NH}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{NH}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{GGFG}-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,
- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{NH}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-\text{GGFG}-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-(\text{NH}-\text{DX})$,

NH-CH₂CH₂CH₂-C(=O)-(NH-DX),

- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-GGFG-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-DGGFG-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX).

[0237] Among them, the more preferred are the followings.

- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-DGGFG-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX).

[0238] The still more preferred are the followings.

- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX).

[0239] The particularly preferred are the followings.

- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-(NH-DX),
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX).

[0240] With regard to the linker structure for conjugating the anti-HER3 antibody and a drug in the antibody-drug conjugate of the present application, the preferred linker can be constructed by connecting preferred structures shown for each part of the linker explained above. As for the linker structure, those with the following structure can be preferably used. Meanwhile, the left terminal of the structure is a connecting position to the antibody and the right terminal is a connecting position to the drug.

- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-NH-CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-NH-CH₂CH₂CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-GGFG-,

- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-DGGFG-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-.

5 [0241] Among them, more preferred are the followings.

- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-NH-CH₂CH₂CH₂-C(=O)-,
- 10 - (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- 15 - (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-DGGFG-
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-.

[0242] Still more preferred are the followings.

- 20 - (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-.

25 [0243] Particularly preferred are the followings.

- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-,
- (Succinimid-3-yl-N)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-.


{Production method}

[0244] Next, explanations are given for the representative method for producing the antibody-drug conjugate of the present invention or a production intermediate thereof. Meanwhile, the compounds are hereinbelow described with the number shown in each reaction formula. Specifically, they are referred to as a "compound of the formula (1)", a "compound (1)", or the like. The compounds with numbers other than those are also described similarly.

1. Production method 1


40 [0245] The antibody-drug conjugate represented by the formula (1) in which the antibody is conjugated to the drug-linker structure via thioether can be produced by the following method, for example.

[Chem. 19]



[in the formula, AB represents an antibody with a sulfhydryl group and L¹ corresponds to L¹ having a structure in which the linker terminal is converted to a maleimidyl group (formula shown below).


[Chem. 20]



(in the formula, the nitrogen atom is the connecting position)

[0246] Specifically, it represents a linker having a structure which, within the structure of L¹ represented as -(Succinimid-3-yl-N)-(CH₂)_n-C(=O)-, said -(Succinimid-3-yl-N)-moiety is converted into a maleimidyl group. Further, the -(NH-DX) represents a structure represented by the following formula:

[Chem. 21]



and it represents a group that is derived by removing one hydrogen atom of the amino group at position 1 of exatecan.]

[0247] Further, the compound of the formula (1) in the above reaction formula can be interpreted as a structure in which one structure moiety from drug to the linker terminal is connected to one antibody. However, it is only the description given for the sake of convenience, and there are actually many cases in which a plurality of said structure moieties is connected to one antibody molecule. The same applies to the explanation of the production method described below.

[0248] Specifically, the antibody-drug conjugate (1) can be produced by reacting the compound (2), which is obtainable by the method described below, with the antibody (3a) having a sulfhydryl group.

[0249] The antibody (3a) having a sulfhydryl group can be obtained by a method well known in the art (Hermanson, G.T., Bioconjugate Techniques, pp. 56-136, pp. 456-493, Academic Press (1996)). Examples include: Traut's reagent is reacted with the amino group of the antibody; N-succinimidyl S-acetylthioalkanoates are reacted with the amino group of the antibody followed by reaction with hydroxylamine; after reacting with N-succinimidyl 3-(pyridyldithio)propionate, it is reacted with a reducing agent; the antibody is reacted with a reducing agent such as dithiothreitol, 2-mercaptoethanol, and tris(2-carboxyethyl)phosphine hydrochloride (TCEP) to reduce the disulfide bond at a hinge part in the antibody to form a sulfhydryl group, but it is not limited thereto.

[0250] Specifically, using 0.3 to 3 molar equivalents of TCEP as a reducing agent per disulfide bonds at hinge part in the antibody and reacting with the antibody in a buffer solution containing a chelating agent, the antibody which the disulfide bonds at hinge part in the antibody is partially or completely reduced can be obtained. Examples of the chelating agent include ethylenediamine tetraacetic acid (EDTA) and diethylenetriamine pentaacetic acid (DTPA). It can be used at concentration of 1 mM to 20 mM. Examples of the buffer solution which may be used include a solution of sodium phosphate, sodium borate, or sodium acetate. Specifically, by reacting the antibody with TCEP at 4°C to 37°C for 1 to 4 hours, the antibody (3a) having partially or completely reduced sulfhydryl groups can be obtained.

[0251] Meanwhile, by performing an addition reaction of a sulfhydryl group to a drug-linker moiety, the drug-linker moiety can be conjugated by a thioether bond.

[0252] Using 2 to 20 molar equivalents of the compound (2) per the antibody (3a) having a sulfhydryl group, the antibody-drug conjugate (1) in which 2 to 8 drug molecules are conjugated per antibody can be produced. Specifically, it is sufficient that the solution containing the compound (2) dissolved therein is added to a buffer solution containing the antibody (3a)

having a sulfhydryl group for the reaction. Herein, examples of the buffer solution which may be used include sodium acetate solution, sodium phosphate, and sodium borate. pH for the reaction is 5 to 9, and more preferably the reaction is performed near pH 7. Examples of the solvent for dissolving the compound (2) include an organic solvent such as dimethyl sulfoxide (DMSO), dimethylformamide (DMF), dimethylacetamide (DMA), and N-methyl-2-pyrrolidone (NMP).

[0253] The reaction may be carried out by adding the organic solvent solution containing the compound (2) dissolved therein at 1 to 20% v/v to a buffer solution containing the antibody (3a) having a sulfhydryl group. The reaction temperature is 0 to 37°C, more preferably 10 to 25°C, and the reaction time is 0.5 to 2 hours. The reaction can be terminated by deactivating the reactivity of unreacted compound (2) with a thiol-containing reagent. Examples of the thiol-containing reagent include cysteine and N-acetyl-L-cysteine (NAC). More specifically, by adding 1 to 2 molar equivalents of NAC to the compound (2) used and, by incubating at room temperature for 10 to 30 minutes, the reaction can be terminated.

[0254] The produced antibody-drug conjugate (1) can be subjected to, after concentration, buffer exchange, purification, and measurement of antibody concentration and average number of conjugated drug molecules per antibody molecule according to common procedures described below, to make an identification of the antibody-drug conjugate (1).

Common procedure A: Concentration of aqueous solution of antibody or antibody-drug conjugate

[0255] To a Amicon Ultra (50,000 MWCO, Millipore Corporation) container, a solution of antibody or antibody-drug conjugate was added and the solution of the antibody or antibody-drug conjugate was concentrated by centrifugation (centrifuge for 5 to 20 minutes at 2000 G to 3800 G) using a centrifuge (Allegra X-15R, Beckman Coulter, Inc.)

Common procedure B: Measurement of antibody concentration

[0256] Using a UV detector (Nanodrop 1000, Thermo Fisher Scientific Inc.), measurement of the antibody concentration was performed according to the method defined by the manufacturer. Here, 280 nm absorption coefficient can be estimated from the amino acid sequence of an antibody using a known calculation method (Protein Science, 1995, vol. 4, 2411-2423), and 280 nm absorption coefficient different for each antibody was used ($1.3 \text{ mLmg}^{-1}\text{cm}^{-1}$ to $1.8 \text{ mLmg}^{-1}\text{cm}^{-1}$). In the case of U1-59, 280 nm absorption coefficient of $1.768 \text{ mLmg}^{-1}\text{cm}^{-1}$ was used as an estimated value according to its amino acid sequence.

Common procedure C: Buffer Exchange for antibody

[0257] NAP-25 column (Cat. No. 17-0852-02, GE Healthcare Japan Corporation) using Sephadex G-25 carrier was equilibrated with phosphate buffer (10 mM, pH 6.0; it is referred to as PBS6.0/EDTA in the specification) containing sodium chloride (137 mM) and ethylene diamine tetraacetic acid (EDTA, 5 mM) according to the method defined by the manufacturer. Aqueous solution of the antibody was applied in an amount of 2.5 mL to single NAP-25 column, and then the fraction (3.5 mL) eluted with 3.5 mL of PBS6.0/EDTA was collected. The resulting fraction was concentrated by the Common procedure A. After measuring the concentration of the antibody using the Common procedure B, the antibody concentration was adjusted to 10 mg/mL using PBS6.0/EDTA.

Common procedure D: Purification of antibody-drug conjugate

[0258] NAP-25 column was equilibrated with acetate buffer containing sorbitol (5%) (10 mM, pH 5.5; it is referred to as ABS in the specification). Aqueous reaction solution of the antibody-drug conjugate (about 2.5 mL) was applied to the NAP-25 column, and then eluted with the buffer in an amount as defined by the manufacturer to collect the antibody fraction. By conducting a gel filtration purification process, in which said collected fraction was again applied to the NAP-25 column and eluted with buffer, was repeated 2 to 3 times in total, the antibody-drug conjugate excluding non-conjugated drug linker and a low-molecular-weight compound (tris(2-carboxyethyl)phosphine hydrochloride (TCEP), N-acetyl-L-cysteine (NAC), and dimethyl sulfoxide) was obtained.

Common procedure E: Measurement of antibody concentration in antibody-drug conjugate and average number of conjugated drug molecules per antibody molecule (1)

[0259] The conjugated drug concentration in the antibody-drug conjugate can be calculated by measuring UV absorbance of an aqueous solution of the antibody-drug conjugate at two wavelengths of 280 nm and 370 nm, followed by performing the calculation shown below.

[0260] Because the total absorbance at any wavelength is equal to the sum of the absorbance of every light-absorbing chemical species that are present in a system [additivity of absorbance], when the molar absorption coefficients of the antibody and the drug remain the same before and after conjugation between the antibody and the drug, the antibody

concentration and the drug concentration in the antibody-drug conjugate are expressed with the following equations.

$$A_{280} = A_{D,280} + A_{A,280} = E_{D,280}C_D + E_{A,280}C_A \quad \text{Equation (1)}$$

$$A_{370} = A_{D,370} + A_{A,370} = E_{D,370}C_D + E_{A,370}C_A \quad \text{Equation (2)}$$

[0261] In the above, A_{280} represents the absorbance of an aqueous solution of the antibody-drug conjugate at 280 nm, A_{370} represents the absorbance of an aqueous solution of the antibody-drug conjugate at 370 nm, $A_{A,280}$ represents the absorbance of an antibody at 280 nm, $A_{A,370}$ represents the absorbance of an antibody at 370 nm, $A_{D,280}$ represents the absorbance of a conjugate precursor at 280 nm, $A_{D,370}$ represents the absorbance of a conjugate precursor at 370 nm, $E_{A,280}$ represents the molar absorption coefficient of an antibody at 280 nm, $E_{A,370}$ represents the molar absorption coefficient of an antibody at 370 nm, $E_{D,280}$ represents the molar absorption coefficient of a conjugate precursor at 280 nm, $E_{D,370}$ represents the molar absorption coefficient of a conjugate precursor at 370 nm, C_A represents the antibody concentration in an antibody-drug conjugate, and C_D represent the drug concentration in an antibody-drug conjugate.

[0262] As for $E_{A,280}$, $E_{A,370}$, $E_{D,280}$, and $E_{D,370}$ in the above, previously prepared values (estimated value based on calculation or measurement value obtained by UV measurement of the compound) are used. For example, $E_{A,280}$ can be estimated from the amino acid sequence of an antibody using a known calculation method (Protein Science, 1995, vol. 4, 2411-2423). $E_{A,370}$ is generally zero. In the case of U1-59, $E_{A,280}$ of 259400 was used as an estimated value according to its amino acid sequence. $E_{D,280}$ and $E_{D,370}$ can be obtained based on Lambert-Beer's law (Absorbance = molar concentration $\times$ molar absorption coefficient $\times$ cell path length) by measuring the absorbance of a solution in which the conjugate precursor to be used is dissolved at a certain molar concentration. By measuring A_{280} and A_{370} of an aqueous solution of the antibody-drug conjugate and solving the simultaneous equations (1) and (2) using the values, C_A and C_D can be obtained. Further, by dividing C_D by C_A , the average drug binding number per antibody can be obtained.

[0263] In the present invention, the method for determining the average number of conjugated drug molecules per antibody as described above is referred to as a "UV method".

Common procedure F: Measurement of average number of conjugated drug molecules per antibody molecule in antibody-drug conjugate - (2)

[0264] The average number of conjugated drug molecules per antibody molecule in the antibody-drug conjugate can also be determined by high-performance liquid chromatography (HPLC) analysis using the following method, in addition to the aforementioned Common procedure E.

{F-1. Preparation of sample for HPLC analysis (Reduction of antibody-drug conjugate)}

[0265] An antibody-drug conjugate solution (about 1 mg/mL, 60 μ L ("u" represents "micro")L) is mixed with an aqueous solution of dithiothreitol (DTT) (100 mM, 15 μ L). By incubating the mixture at 37°C for 30 minutes, the disulfide bond between the L and H chains of the antibody-drug conjugate is cleaved. The resulting sample is used in HPLC analysis.

{F-2. HPLC analysis}

[0266] The HPLC analysis is carried out under the following measurement conditions.

HPLC system: Agilent 1290 HPLC system (Agilent Technologies)

Detector: Ultraviolet absorption spectrometer (measurement wavelength: 280 nm)

Column: PLRP-S (2.1 $\times$ 50 mm, 8 μ m, 1000 angstroms; Agilent Technologies, PIN PL1912-1802)

Column temperature: 80°C

Mobile phase A: 0.04% aqueous trifluoroacetic acid (TFA) solution

Mobile phase B: Acetonitrile solution containing 0.04% TFA

Gradient program: 29%-36% (0 min.-12.5 min.), 36%-42% (12.5-15 min.), 42%-29% (15 min.-15.1 min.), 29%-29% (15.1 min.-25 min.)

Sample injection: 15 μ L

{F-3. Data analysis}

[0267] [F-3-1] Compared with non-conjugated antibody L (L_0) and H (H_a) chains, drugconjugated L (L chain bound to one drug molecule: L_1) and H (H chain bound to one drug molecule: H_1 , H chain bound to two drug molecule: H_2 , H chain

bound to three drug molecules: H₃) chains exhibit higher hydrophobicity in proportion to the number of conjugated drug molecules and thus have a larger retention time. These chains are therefore eluted in the order of L₀ and L₁ or H₀, H₁, H₂, and H₃. Detection peaks can be assigned to any of L₀, L₁, H₀, H₁, H₂, and H₃ by the comparison of retention times with L₀ and H₀.

[0268] [F-3-2] Since the drug linker has UV absorption, peak area values are corrected in response to the number of conjugated drug linker molecules according to the following expression using the molar absorption coefficients of the L or H chain and the drug linker.

[Math. 1]

Corrected value of the peak area of the L chain (L_i)

= Peak area

$$\times \frac{\text{Molar absorption coefficient of the L chain}}{\text{Molar absorption coefficient of the L chain} + \text{The number of conjugated drug molecules} \times \text{Molar absorption coefficient of the drug linker}}$$

[Math. 2]

Corrected value of the peak area of the H chain (H_i)

= Peak area

$$\times \frac{\text{Molar absorption coefficient of the H chain}}{\text{Molar absorption coefficient of the H chain} + \text{The number of conjugated drug molecules} \times \text{Molar absorption coefficient of the drug linker}}$$

[0269] Here, a value estimated from the amino acid sequence of the L or H chain of each antibody using a known calculation method (Protein Science, 1995, vol. 4, 2411-2423) can be used as the molar absorption coefficient (280 nm) of the L or H chain of each antibody. In the case of U1-59, a molar absorption coefficient of 34690 and a molar absorption coefficient of 95000 were used as estimated values for the L and H chains, respectively, according to its amino acid sequence. The actually measured molar absorption coefficient (280 nm) of a compound in which the maleimide group has been converted to succinimide thioether by the reaction of each drug linker with mercaptoethanol or N-acetylcysteine was used as the molar absorption coefficient (280 nm) of the drug linker.

[F-3-3] The peak area ratio (%) of each chain is calculated for the total of the corrected values of peak areas.

[Math. 3]

$$\text{Peak area ratio of the L chain} = \frac{A_{Li}}{A_{L0} + A_{L1}} \times 100$$

$$\text{Peak area ratio of the H chain} = \frac{A_{Hi}}{A_{H0} + A_{H1} + A_{H2} + A_{H3}} \times 100$$


Corrected values of respective peak areas of Au, A_{Hi}:L_i, H_i

[0270] [F-3-4] The average number of conjugated drug molecules per antibody molecule in the antibody-drug conjugate is calculated according to the following expression.


Average number of conjugated drug molecules = (L₀ peak area ratio x 0 + L₀ peak area ratio x 1 + H₀ peak area ratio x 0 + H₁ peak area ratio x 1 + H₂ peak area ratio x 2 + H₃ peak area ratio x 3) / 100 x 2

[0271] Hereinbelow, production intermediate compounds used in Production method 1 are described. The compound

represented by the formula (2) in the production method 1 is a compound represented by the following formula:



or



In the formula,

n^3 represents an integer of 2 to 8,

L^2 represents $-\text{NH}-(\text{CH}_2\text{CH}_2-\text{O})_{n^4}-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-$ or a single bond,

wherein n^4 represents an integer of 1 to 6,

L^{P} represents a peptide residue consisting of 2 to 7 amino acids selected from phenylalanine, glycine, valine, lysine, citrulline, serine, glutamic acid, and aspartic acid,


n^1 represents an integer of 0 to 6,

n^2 represents an integer of 0 to 5,

L^{a} represents $-\text{O}-$ or a single bond,


(maleimid-N-yl)- is a maleimidyl group (2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl group) represented by the following formula:

[Chem. 22]



wherein the nitrogen atom is the connecting position, and $-(\text{NH-DX})$ is a group represented by the following formula:

[Chem. 23]



wherein the nitrogen atom of the amino group at position 1 is the connecting position.

[0272] As for the peptide residue L^{P} , those consisting of an amino acid selected from phenylalanine, glycine, valine, lysine, citrulline, serine, glutamic acid, and aspartic acid is preferred as a production intermediate. Among the peptide residue L^{P} , those consisting of 4 or 5 amino acids is preferred as a production intermediate. More specifically, those in

which LP is a tetrapeptide residue of -GGFG- or a pentapeptide of -DGGFG- is preferred as a production intermediate, more preferably, -GGFG-.

[0273] Further, as for the $-\text{NH}-(\text{CH}_2)^{n1}-\text{L}^a-(\text{CH}_2)^{n2}-$, those having $-\text{NH}-\text{CH}_2\text{CH}_2-$, $-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{NH}-\text{CH}_2-\text{O}-\text{CH}_2-$, or $-\text{NH}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2-$ is preferred as a production intermediate. A compound of $-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{NH}-\text{CH}_2-\text{O}-\text{CH}_2-$, or $-\text{NH}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2-$ is more preferred.

[0274] As for n^3 , those in which it is an integer of 2 to 8 is preferred as a production intermediate.

[0275] As for L², those in which it is a single bond or -NH-(CH₂CH₂-O)ⁿ-CH₂CH₂-C(=O)- and ⁿ is an integer of 2 to 4 is preferred as a production intermediate.

[0276] Further, those in which n^3 is an integer of 2 to 5, L^2 is a single bond, and $-NH-(CH_2)n^1-L^a-(CH_2)n^2-$ is $-NH-CH_2CH_2-$, $-NH-CH_2CH_2CH_2-$, $-NH-CH_2CH_2CH_2CH_2-$, $-NH-CH_2CH_2CH_2CH_2CH_2-$, $-NH-CH_2-O-CH_2-$, or $-NH-CH_2CH_2-O-CH_2-$ is preferred as a production intermediate. Further, more preferred among them is those in which $-NH-(CH_2)n^1-L^a-(CH_2)n^2-$ is $-NH-CH_2CH_2-$, $-NH-CH_2CH_2CH_2-$, $-NH-CH_2-O-CH_2-$, or $-NH-CH_2CH_2-O-CH_2-$. Further, those in which n^3 is an integer of 2 or 5 is preferred.

[0277] Further, those in which n^3 is an integer of 2 to 5, L^2 is $-\text{NH}-(\text{CH}_2\text{CH}_2-\text{O})n^4-\text{CH}_2\text{CH}_2-\text{C}(=\text{O})-$, n^4 is an integer of 2 to 4, and $-\text{NH}-(\text{CH}_2)n^1-\text{L}^a-(\text{CH}_2)n^2-$ is $-\text{NH}-\text{CH}_2\text{CH}_2-$, $-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{NH}-\text{CH}_2-\text{O}-\text{CH}_2-$, or $-\text{NH}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2-$ is preferred as a production intermediate. More preferred among them is those in which n^4 is an integer of 2 or 4. Further, those in which $-\text{NH}-(\text{CH}_2)n^1-\text{L}^a-$ is $-\text{NH}-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{NH}-\text{CH}_2-\text{O}-\text{CH}_2-$, or $-\text{NH}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2-$ is preferred.

[0278] Preferred examples of the intermediate that are useful for production of the compound of the present invention include those exemplified below:

[illegible]

(maleimid-N-yl)-CH₂CH₂-C(=O)-DGGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX), or
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX).

[0279] By the reaction of the drug-linker compound selected from the aforementioned group of intermediate compounds with an anti-Her3 antibody or a reactive derivative thereof, a thioether bond can be formed at a disulfide bond moiety present in a hinge part of the anti-Her3 antibody, and as a result, the anti-Her3 antibody-drug conjugate of the present invention can be produced. In this case, it is preferable to use a reactive derivative of an anti-Her3 antibody. A reactive derivative obtained by reducing an anti-Her3 antibody is particularly preferred.

[0280] The followings are a compound which is more preferred as a production intermediate.

(maleimid-N-yl)-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂-C(=O)-DGGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX).

[0281] Further, among the aforementioned intermediate compound group, the intermediates represented by the following formula are a more preferred compound:

(maleimid-N-yl)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX), or
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-DGGFG-(NH-DX).


[0282] Particularly preferred are the compounds that are represented by the following formula:

(maleimid-N-yl)-CH₂CH₂-C(=O)-NH-CH₂CH₂-O-CH₂CH₂-O-CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂CH₂-C(=O)-(NH-DX),
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂-O-CH₂-C(=O)-(NH-DX), or
 (maleimid-N-yl)-CH₂CH₂CH₂CH₂CH₂-C(=O)-GGFG-NH-CH₂CH₂-O-CH₂-C(=O)-(NH-DX).

2. Production method 2

[0283] The compound represented by the formula (2) or a pharmacologically acceptable salt thereof used as an intermediate in the previous production method can be produced by the following method, for example.

[Chem. 24]



[in the formula, L¹ corresponds to L¹ having a structure in which the terminal is converted to a maleimidyl group and P¹, P², and P³ represent a protecting group].

[0284] The compound (6) can be produced by derivatizing the carboxylic acid (5) into an active ester, mixed acid anhydride, acid halide, or the like and, in the presence of base, reacting it with NH₂-DX [indicating exatecan; chemical name: (1S,9S)-1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino [1,2-b]quinolin-10,13(9H,15H)-dione] (4) or a pharmacologically acceptable salt thereof.

[0285] Reaction reagents and conditions that are commonly used for peptide synthesis can be employed for the reaction. There are various kinds of active ester, for example, it can be produced by reacting phenols such as p-nitrophenol, N-hydroxy benzotriazole, N-hydroxy succinimide, or the like, with the carboxylic acid (5) using a condensing agent such as N,N'-dicyclohexylcarbodiimide or 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride; further, the active ester can be also produced by a reaction of the carboxylic acid (5) with pentafluorophenyl trifluoroacetate or the like; a reaction of the carboxylic acid (5) with 1-benzotriazolyl oxytripyrrolidinophosphonium hexafluorophosphate; a reaction of the carboxylic acid (5) with diethyl cyanophosphonate (Shioiri method); a reaction of the carboxylic acid (5) with triphenylphosphine and 2,2'-dipyridyl disulfide (Mukaiyama method); a reaction of the carboxylic acid (5) with a triazine derivative such as 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM); or the like. Further, the reaction can be also performed by, e.g., an acid halide method by which the carboxylic acid (5) is treated with acid halide such as thionyl chloride and oxalyl chloride in the presence of a base.

[0286] By reacting the active ester, mixed acid anhydride, or acid halide of the carboxylic acid (5) obtained as above with the compound (4) in the presence of a suitable base in an inert solvent at a reaction temperature of -78°C to 150°C, the compound (6) can be produced. Meanwhile, "inert solvent" indicates a solvent which does not inhibit a desired reaction for which the solvent is used.

[0287] Specific examples of the base used for each step described above include a carbonate, an alkoxide, a hydroxide or a hydride of an alkali metal or an alkali earth metal such as sodium carbonate, potassium carbonate, sodium ethoxide, potassium butoxide, sodium hydroxide, potassium hydroxide, sodium hydride, or potassium hydride; organometallic base represented by an alkyl lithium such as n-butyl lithium, or dialkylamino lithium such as lithium diisopropylamide; organometallic base such as bisilylamine including lithium bis(trimethylsilyl)amide; and organic base such as pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, triethylamine, N-methyl morpholine, diisopropylethylamine, and diaza-bicyclo[5.4.0]undec-7-ene (DBU).

[0288] Examples of the inert solvent which is used for the reaction of the present invention include a halogenated hydrocarbon solvent such as dichloromethane, chloroform, and carbon tetrachloride; an ether solvent such as tetrahydrofuran, 1,2-dimethoxyethane, and dioxane; an aromatic hydrocarbon solvent such as benzene and toluene; and an amide solvent such as N,N-dimethylformamide, N,N-dimethylacetamide, and N-methylpyrrolidin-2-one. In addition to them, a sulfoxide solvent such as dimethyl sulfoxide and sulfolane; a ketone solvent such as acetone and methyl ethyl ketone; and an alcohol solvent such as methanol and ethanol may be used in some case. Alternatively, these solvents may be used as a mixed solvent.

[0289] As for the protecting group P¹ for the terminal amino group of the compound (6), a protecting group for an amino group which is generally used for peptide synthesis, for example, tert-butyloxy carbonyl group, 9-fluorenylmethyloxy carbonyl group, and benzyloxy carbonyl group, can be used. Examples of the other protecting group for an amino group

include an alkanoyl group such as acetyl group; an alkoxycarbonyl group such as methoxycarbonyl group and ethoxycarbonyl group; an arylmethoxy carbonyl group such as paramethoxybenzyloxy carbonyl group, and para (or ortho)nitrobenzyloxy carbonyl group; an arylmethyl group such as benzyl group and triphenyl methyl group; an aroyl group such as benzoyl group; and an aryl sulfonyl group such as 2,4-dinitrobenzene sulfonyl group and orthonitrobenzene sulfonyl group. The protecting group P¹ can be selected depending on, e.g., properties of a compound having an amino group to be protected.

[0290] By deprotecting the protecting group P¹ for the terminal amino group of the compound (6) obtained, the compound (7) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0291] The compound (9) can be produced by derivatizing the peptide carboxylic acid (8) having the N terminal protected with P² into an active ester, mixed acid anhydride, or the like and reacting it with the compound (7) obtained. The reaction conditions, reagents, base, and inert solvent used for a peptide bond formation between the peptide carboxylic acid (8) and the compound (7) can be suitably selected from those described for the synthesis of the compound (6). The protecting group P² can be suitably selected from those described for the protecting group of the compound (6), and the selection can be made based on, e.g., the properties of the compound having an amino group to be protected. As it is generally used for peptide synthesis, by repeating sequentially the reaction and deprotection of the amino acid or peptide constituting the peptide carboxylic acid (8) for elongation, the compound (9) can be also produced.

[0292] By deprotecting P² as the protecting group for the amino group of the compound (9) obtained, the compound (10) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0293] It is possible to produce the compound (2) by derivatizing the carboxylic acid (11) into an active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (10) obtained. The reaction conditions, reagents, base, and inert solvent used for forming a peptide bond between the carboxylic acid (11) and the compound (10) can be suitably selected from those described for the synthesis of the compound (6).

[0294] The compound (9) can be also produced by the following method, for example.

[0295] The compound (13) can be produced by derivatizing the peptide carboxylic acid (8) having the N terminal protected with P² into active ester, mixed acid anhydride, or the like and reacting it with the amine compound (12) having the carboxy group protected with P³ in the presence of a base. The reaction conditions, reagents, base, and inert solvent used for forming a peptide bond between the peptide carboxylic acid (8) and the compound (12) can be suitably selected from those described for the synthesis of the compound (6).

[0296] The protecting group P² for the amino group of the compound (13) is not particularly limited if it is a protecting group which is commonly used. Specifically, examples of the protecting group for a hydroxyl group include an alkoxymethyl group such as methoxymethyl group; an arylmethyl group such as benzyl group, 4-methoxybenzyl group, and triphenylmethyl group; an alkanoyl group such as acetyl group; an aroyl group such as benzoyl group; and a silyl group such as tert-butyl diphenylsilyl group. Carboxy group can be protected by an ester with an alkyl group such as methyl group, ethyl group, and tert-butyl group, an allyl group, or an arylmethyl group such as benzyl group. As for the amino group, an alkyloxy carbonyl group such as tert-butyloxy carbonyl group, methoxycarbonyl group, and ethoxycarbonyl group; an arylmethoxy carbonyl group such as allyloxycarbonyl group, 9-fluorenylmethyloxy carbonyl group, benzyloxy carbonyl group, paramethoxybenzyloxy carbonyl group, and para (or ortho)nitrobenzyloxy carbonyl group; an alkanoyl group such as acetyl group; an arylmethyl group such as benzyl group and triphenyl methyl group; an aroyl group such as benzoyl group; and an aryl sulfonyl group such as 2,4-dinitrobenzene sulfonyl group or orthonitrobenzene sulfonyl group can be mentioned.

[0297] As for the protecting group P³ for a carboxy group, a protecting group commonly used as a protecting group for a carboxy group in organic synthetic chemistry, in particular, peptide synthesis can be used. A carboxyl group can be protected as an ester with an alkyl group such as a methyl group, an ethyl group, or a tert-butyl, an allyl group, and an arylmethyl group such as a benzyl group.

[0298] In such case, it is preferable that the protecting group for an amino group and the protecting group for a carboxy group can be removed by a different method or different conditions. For example, a representative example includes a combination in which P² is a tert-butyloxy carbonyl group and P³ is a benzyl group. The protecting groups can be selected from the aforementioned ones depending on, e.g., the properties of a compound having an amino group and a carboxy group to be protected. For removal of the protecting groups, reagents and conditions can be selected depending on the protecting group.

[0299] By deprotecting the protecting group P³ for the carboxy group of the compound (13) obtained, the compound (14) can be produced. In this deprotection, reagents and conditions are selected depending on the protecting group.

[0300] The compound (9) can be produced by derivatizing the compound (14) obtained into active ester, mixed acid anhydride, acid halide, or the like and reacting with the compound (4) in the presence of a base. For the reaction, reaction reagents and conditions that are generally used for peptide synthesis can be also used, and the reaction conditions, reagents, base, and inert solvent used for the reaction can be suitably selected from those described for the synthesis of the compound (6).

[0301] The compound (2) can be also produced by the following method, for example.

[0302] By deprotecting the protecting group P² for the amino group of the compound (13), the compound (15) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0303] The compound (16) can be produced by derivatizing the carboxylic acid derivative (11) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (15) obtained in the presence of a base. The reaction conditions, reagents, base, and inert solvent used for forming an amide bond between the peptide carboxylic acid (11) and the compound (15) can be suitably selected from those described for the synthesis of the compound (6).


[0304] By deprotecting the protecting group for the carboxy group of the compound (16) obtained, the compound (17) can be produced. In this deprotection, it can be carried out similar to deprotecting carboxy group for producing the compound (14).

[0305] The compound (2) can be produced by derivatizing the compound (17) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (4) in the presence of a base. For the reaction, reaction reagents and conditions that are generally used for peptide synthesis can be also used, and the reaction conditions, reagents, base, and inert solvent used for the reaction can be suitably selected from those described for the synthesis of the compound (6).

3. Production method 3

[0306] The compound represented by the formula (2) used as an intermediate can be also produced by the following method.

[Chem. 25]



[in the formula, L^{1'} corresponds to L¹ having a structure in which the terminal is converted to a maleimidyl group and P⁴ represents a protecting group].

[0307] The compound (19) can be produced by derivatizing the compound (11) into active ester, mixed acid anhydride, or the like and reacting it with the peptide carboxylic acid (18) having the C terminal protected with P⁴ in the presence of a base. The reaction conditions, reagents, base, and inert solvent used for forming a peptide bond between the peptide carboxylic acid (18) and the compound (11) can be suitably selected from those described for the synthesis of the compound (6). The protecting group P⁴ for the carboxy group of the compound (18) can be suitably selected from the aforementioned protective groups.


[0308] By deprotecting the protecting group for the carboxy group of the compound (19) obtained, the compound (20) can be produced. In this deprotection, it can be performed similar to the deprotection of the carboxy group for producing the compound (14).

[0309] The compound (2) can be produced by derivatizing the compound (20) obtained into active ester, mixed acid anhydride, or the like and reacting it with the compound (7). For the reaction, reaction reagents and conditions that are generally used for peptide synthesis can be also used, and the reaction conditions, reagents, base, and inert solvent used for the reaction can be suitably selected from those described for the synthesis of the compound (6).

4. Production method 4

[0310] Hereinbelow, within the production intermediate (10) described in production method 2, the method for producing the compound (10b) having $n^1 = 1$ and $L^a = 0$ is described in detail. The compound represented by the formula (10b), a salt or a solvate thereof can be produced according to the following method, for example.

[Chem. 26]



[in the formula, L^P is as defined above, L represents an acyl group including an alkanoyl group such as acetyl group or an aroyl group such as benzoyl group, or represents a hydrogen atom or the like, X and Y represent an oligopeptide consisting of 1 to 3 amino acids, P^5 and P^7 represent a protecting group for an amino group, and P^6 represents a protecting group for a carboxy group].

[0311] A compound represented by the formula (21) can be produced by using or applying the method described in Japanese Patent Laid-Open No. 2002-60351 or the literature (J. Org. Chem., Vol. 51, page 3196, 1986), and if necessary, by removing the protecting groups or modifying the functional groups. Alternatively, it can be also obtained by treating an amino acid with a protected terminal amino group or acid amide of oligopeptide with protected amino group with aldehyde or ketone.

[0312] By reacting the compound (21) with the compound (22) having a hydroxyl group under temperature conditions ranging from under cooling to room temperature in an inert solvent in the presence of an acid or a base, the compound (23) can be produced.

[0313] Here, examples of the acid which may be used can include inorganic acid such as hydrofluoric acid, hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid, and boric acid; an organic acid such as acetic acid, citric acid, paratoluene sulfonic acid, and methane sulfonic acid; and a Lewis acid such as tetrafluoroborate, zinc chloride, tin chloride, aluminum chloride, and iron chloride. Among them, sulfonic acids are preferable, and paratoluene sulfonic acid is particularly preferable. As for the base, any one of the aforementioned base can be suitably selected and used. Preferred examples thereof include an alkali metal alkoxide such as potassium tertbutoxide, an alkali metal or alkaline earth metal hydroxide such as sodium hydroxide and potassium hydroxide; alkali metal hydride such as sodium hydride and potassium hydride; organometallic base represented by dialkylamino lithium such as lithium diisopropylamide; and organometallic base of bissilylamine such as lithium bis(trimethylsilyl)amide.

[0314] Examples of the solvent to be used for the reaction include an ether solvent such as tetrahydrofuran and 1,4-dioxane; and an aromatic hydrocarbon solvent such as benzene and toluene. Those solvents can be prepared as a mixture with water.

[0315] Further, the protecting group for an amino group as represented as P^5 is not particularly limited if it is a group commonly used for protection of an amino group. Representative examples can include the protecting groups for an amino group that are described in Production method 2. However, the protecting group for an amino group as represented as P^5 may be cleaved off within the course of the reaction. In such case, a protecting group can be re-introduced by appropriately performing a reaction with a suitable reagent for protecting an amino group as required.

[0316] The compound (24) can be derived by removing the protecting group P^6 of the compound (23). Herein, although the representative examples of the protecting group for a carboxy group as represented as P^6 are described in Production method 2, it can be appropriately selected from these examples. In the compound (23), it is desirable that the protecting group P^5 for an amino group and the protecting group P^6 for a carboxy group are the protecting groups that can be removed

by a different method or different conditions. For example, a representative example can include a combination in which P⁵ is a 9-fluorenylmethyloxy carbonyl group and P⁶ is a benzyl group. The protecting groups can be selected depending on, e.g., the properties of a compound having an amino group and a carboxy group to be protected. For removal of the protecting groups, reagents and conditions are selected depending on the protecting group.


[0317] The compound (26) can be produced by derivatizing the compound (24) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (4) or a pharmacologically acceptable salt thereof in the presence of a base to produce the compound (25) followed by removing the protecting group P⁵ of the compound (25) obtained. For the reaction between the compound (4) and the carboxylic acid (24) and the reaction for removing the protecting group P⁶, the same reagents and reaction conditions as those described for Production method 2 can be used.

[0318] The compound (10b) can be produced by reacting the compound (26) with an amino acid with protected terminal amino group or the oligopeptide (27) with protected amino group to produce the compound (9b) and removing the protective group P⁷ of the compound (9b) obtained. The protective group for an amino group as represented as P⁷ is not particularly limited if it is generally used for protection of an amino group. Representative examples thereof include the protecting groups for an amino group that are described in Production method 2. For removing the protective group, reagents and conditions are selected depending on the protecting group. For the reaction between the compound (26) and the compound (27), reaction reagents and conditions that are commonly used for peptide synthesis can be employed. The compound (10b) produced by the aforementioned method can be derivatized into the compound (1) of the present invention according to the method described above.

5. Production method 5

[0319] The compound represented by the formula (2) as an intermediate can be also produced by the method shown below.

[Chem. 27]



[in the formula, $L^{1'}$ corresponds to L^1 having a structure in which the terminal is converted to a maleimidyl group, L^P represents a structure consisting of $-L^{P1}-L^{P2}-$, and P^3 , P^8 , P^9 , P^{10} , P^{11} , and P^{12} represent a protecting group].

[0320] Because L^P is formed by connecting L^{P1} to L^{P2} , the hydrophilic amino acid at N terminal of L^P is derived from L^{P1} , and thus, those that having a hydrophilic amino acid at the N terminal are suitably employed as L^{P1} . Meanwhile, plural hydrophilic amino acids may be present therein. Further, when L^{P2} with hydrophilic amino acid is employed, L^P having plural hydrophilic amino acids at the N terminal of L^P or at the N terminal and at other positions can be produced depending on the location of the hydrophilic amino acid.

[0321] The compound (29) can be produced by derivatizing the peptide or amino acid (28) having the N terminal protected with P^2 into active ester, mixed acid anhydride, or the like and reacting it with the compound (7) obtained. The reaction conditions, reagents, base, and solvent used for forming an amide bond between the peptide or amino acid (28) and the compound (7) can be suitably selected from those described for the synthesis of the compound (6). The protecting group P^8 for an amino group can be suitably selected from those described for the protecting group of the compound (6), and the selection can be made based on the properties of the compound or the like. As it is generally used for peptide synthesis, by repeating sequentially the reaction and deprotection of the amino acid or peptide constituting the peptide or amino acid (28) for elongation, the compound (29) can be also produced.

[0322] By deprotection of P^8 as a protecting group of the amino group of the compound (29) obtained, the compound (23) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0323] The compound (32) can be produced by derivatizing the amino acid or peptide (31) having the N terminal protected with P^8 and the protected carboxy group, hydroxy group, or amino group in side chain protected into active ester, mixed acid anhydride, or the like and reacting it with the compound (30) obtained. The reaction conditions, reagents, base, and inert solvent used for forming a peptide bond between the amino acid or peptide (31) and the compound (30) can be

suitably selected from those described for the synthesis of the compound (6). As for the protecting groups P⁸ and P⁹, the protecting groups can be suitably selected from those described as protecting group for an amino group, carboxy group, or hydroxy group of the compound (6). However, in such case, it is necessary that the protecting group P⁹ for an amino group and the protecting group P¹⁰ for a functional group in side chain can be removed by a different method or different conditions. For example, a representative example includes a combination in case P⁹ is a 9-fluorenylmethyloxy carbonyl group and P¹⁰ is a tert-butyl group or the like as a protecting group for a carboxy group, a methoxymethyl group or the like as a protecting group for a hydroxy group, or a tert-butyloxycarbonyl group or the like as a protecting group for an amino group. The protective group P¹⁰ for a functional group in a side chain is preferably a protecting group which can be deprotected by a treatment under acidic conditions. However, it is not limited thereto, and it can be selected from the aforementioned ones depending on, e.g., the properties of amino group, carboxy group, or a hydroxy group of a compound to be protected. For removal of the protecting groups, reagents and conditions are selected depending on the protecting group. As it is generally used for peptide synthesis, by repeating sequentially the reaction and deprotection of the constituting amino acid or peptide for elongation, the compound (32) can be also produced.

[0324] By deprotection of P⁹ as a protecting group of the terminal amino group of the compound (32) obtained, the compound (33) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0325] It is possible to produce the compound (34) by derivatizing the carboxylic acid derivative (11) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (33) obtained. Herein, the carboxylic acid derivative (11) is a compound with a structure in which the linker terminal of L^{1'} has a maleimidyl group.

[0326] The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the carboxylic acid derivative (11) and the compound (33) can be suitably selected from those described for the synthesis of the compound (6).

[0327] By deprotecting the protecting group P¹⁰ for the carboxy group, hydroxy group, or amino group in the amino acid side chain of the peptide moiety of the compound (34) obtained, the compound (2) can be produced. Reagents and conditions can be selected depending on the protecting group.

[0328] The compound (29) can be also produced by the following method, for example.

[0329] The compound (35) can be produced by derivatizing the peptide or amino acid (28) having the N terminal protected with P⁸ into active ester, mixed acid anhydride, or the like and reacting it with the amine compound (12) having the terminal carboxy group protected with P³ in the presence of a base. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the peptide or amino acid (28) and the compound (12) can be suitably selected from those described for the synthesis of the compound (6). The protecting group P⁸ for an amino group of the compound (35) can be suitably selected and used from those described as a protecting group for the compound (6). As for the protecting group P³ for a carboxy group, a protecting group commonly used as a protecting group for a carboxy group in organic synthetic chemistry, in particular, peptide synthesis can be used. Specific examples include alkyl ester such as methyl group, ethyl group, and tert-butyl, allyl ester, and benzyl ester, and it can be suitably selected and used from the protecting groups that are described for the compound (6). In such case, it is necessary that the protecting group P⁸ for an amino group and the protecting group P³ for a carboxy group can be removed by a different method or different conditions. For example, a representative example includes a combination in which P⁸ is a tert-butyloxy carbonyl group and P³ is a benzyl group. The protecting groups can be selected from the aforementioned ones depending on, e.g., the properties of a compound having an amino group and a carboxy group to be protected. For removal of the protecting groups, reagents and conditions are selected depending on the protecting group.

[0330] By deprotecting the protecting group P³ for the carboxy group of the compound (35) obtained, the compound (36) can be produced. In this deprotection, reagents and conditions are selected depending on the protecting group.

[0331] The compound (29) can be produced by derivatizing the compound (36) obtained into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (4) in the presence of a base. For the reaction, reaction reagents and conditions that are generally used for peptide synthesis can be also used, and the reaction conditions, reagents, base, and solvent used for the reaction can be suitably selected from those described for the synthesis of the compound (6).

[0332] The compound (32) can be also produced by the following method, for example.

[0333] By deprotecting the protecting group P⁸ for the amino group of the compound (35), the compound (37) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0334] The compound (38) can be produced by derivatizing the amino acid or peptide (31) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (37) obtained in the presence of a base. The reaction conditions, reagents, base, and solvent used for forming an amide bond between the amino acid or peptide (31) and the compound (37) can be suitably selected from those described for the synthesis of the compound (6). In such case, it is necessary that the protecting group P⁹ and P¹⁰ for the amino acid or peptide (31) and the protecting group P³ for the compound (37) can be removed by a different method or different conditions. For example, a representative example includes a combination in which P⁹ is a 9-fluorenylmethyloxy carbonyl group, P¹⁰ is a tert-butyloxy carbonyl group, tert-butyl group, or a methoxymethyl group, and P³ is a benzyl group. Further, the protective group P¹⁰ for a functional group in

a side chain is preferably a protecting group which can be deprotected by a treatment under acidic conditions as described above. However, it is not limited thereto, and it can be selected from the aforementioned ones depending on, e.g., the properties of amino group, carboxy group, or a hydroxy group of a compound to be protected. For removal of the protecting groups, reagents and conditions are selected depending on the protecting group.

[0335] By deprotecting the protecting group P³ for the carboxy group of the compound (38) obtained, the compound (39) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0336] The compound (32) can be produced by derivatizing the compound (39) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (4) in the presence of a base. For the reaction, reaction reagents and conditions that are generally used for peptide synthesis can be also used, and the reaction conditions, reagents, base, and solvent used for the reaction can be suitably selected from those described for the synthesis of the compound (6).

[0337] The compound (34) can be also produced by the following method, for example.

[0338] By deprotecting the protecting group P⁹ for the amino group of the compound (38), the compound (40) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0339] The compound (41) can be produced by derivatizing the carboxylic acid derivative (11) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (40) obtained in the presence of a base. The reaction conditions, reagents, base, and solvent used for forming an amide bond between the carboxylic acid derivative (11) and the compound (40) can be suitably selected from those described for the synthesis of the compound (6).

[0340] By deprotecting the protecting group P³ for the carboxy group of the compound (41) obtained, the compound (42) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0341] The compound (34) can be produced by derivatizing the compound (42) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (4) in the presence of a base. For the reaction, reaction reagents and conditions that are generally used for peptide synthesis can be also used, and the reaction conditions, reagents, base, and solvent used for the reaction can be suitably selected from those described for the synthesis of the compound (6).

[0342] The compound (34) can be also produced by the following method, for example.

[0343] The compound (44) can be produced by derivatizing the carboxylic acid derivative (11) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the amino acid or peptide (43) having the carboxy group protected with P¹¹ and the carboxy group, hydroxy group, or amino group in side chain protected with P¹⁰ in the presence of a base. The reaction conditions, reagents, base, and solvent used for forming an amide bond between the carboxylic acid derivative (11) and the compound (43) can be suitably selected from those described for the synthesis of the compound (6).

As for the protecting groups P¹⁰ and P¹¹ of the compound (44), the protecting groups can be suitably selected from those described as protecting group for a carboxy group, hydroxy group, or amino group of the compound (6). Meanwhile, in such case, it is necessary that the protecting group P¹¹ for a carboxy group and the protecting group P¹⁰ for a functional group in side chain can be removed by a different method or different conditions. For example, a representative example includes a combination in which P¹¹ is a benzyl group and P¹⁰ is a tert-butyl group or the like as a protecting group for a carboxy group, a methoxymethyl group or the like as a protecting group for a hydroxy group, or a tert-butyloxycarbonyl group or the like as a protecting group for an amino group. The protective group P¹⁰ for a functional group in a side chain is preferably a protecting group which can be deprotected by a treatment under acidic conditions. However, it is not limited thereto, and it can be selected from the aforementioned ones depending on, e.g., the properties of amino group, carboxy group, or a hydroxy group of a compound to be protected. For removing the protecting group, the reagents and conditions can be selected depending on the protecting group.

[0344] By deprotecting the protecting group P¹¹ for the carboxy group of the compound (44) obtained, the compound (45) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0345] The compound (34) can be produced by derivatizing the compound (45) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (30) in the presence of a base. For the reaction, reaction reagents and conditions that are generally used for peptide synthesis can be also used, and the reaction conditions, reagents, base, and solvent used for the reaction can be suitably selected from those described for the synthesis of the compound (6).

[0346] The compound (47) can be produced by derivatizing the compound (45) into active ester, mixed acid anhydride, acid halide or the like and reacting it with the amino acid or peptide (46) having the carboxy group protected with P¹² in the presence of a base. For the reaction, the reaction reagents and conditions commonly used for peptide synthesis can be used and the reaction conditions, reagents, base, and solvent can be suitably selected from those described for the synthesis of the compound (6). As for the protecting groups P¹⁰ and P¹² of the compound (47), the protecting groups can be suitably selected and used from those described as protecting group for a carboxy group, hydroxy group, or amino group of the compound (6). Meanwhile, in such case, it is necessary that the protecting group P¹² for a carboxy group and the protecting group P¹⁰ for a functional group in side chain can be removed by a different method or different conditions. For example, a representative example includes a combination in which P¹² is a benzyl group and P¹⁰ is a tert-butyl group or the like as a protecting group for a carboxy group, a methoxymethyl group or the like as a protecting group for a hydroxy group, or a tert-butyloxycarbonyl group or the like as a protecting group for an amino group. The protective group P¹⁰ for a functional group in a side chain is preferably a protecting group which can be deprotected by a treatment under acidic

conditions. However, it is not limited thereto, and it can be selected from the aforementioned ones depending on, e.g., the properties of amino group, carboxy group, or a hydroxy group of a compound to be protected. For removing the protecting group, the reagents and conditions can be selected depending on the protecting group. Further, the compound (47) can be also produced by repeating sequentially the reaction and deprotection of constituting amino acid or peptide for elongation.

[0347] By deprotecting the protecting group P^{12} for the carboxy group of the compound (47) obtained, the compound (48) can be produced. Reagents and conditions can be selected depending on the protecting group.

[0348] The compound (34) can be produced by derivatizing the compound (48) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (7) in the presence of a base. For the reaction, reaction reagents and conditions that are generally used for peptide synthesis can be also used, and the reaction conditions, reagents, base, and solvent used for the reaction can be suitably selected from those described for the synthesis of the compound (6).

[0349] The compound (47) can be also produced by the following method, for example.

[0350] The peptide (49) can be produced by derivatizing the amino acid or peptide (46) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the amino acid or peptide (31) having the N terminal protected with P^9 and the carboxy group, hydroxy group, or amino group in side chain protected with P^{10} in the presence of a base. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the amino acid or peptide (46) and the amino acid or peptide (31) can be suitably selected from those described for the synthesis of the compound (6). Meanwhile, in this case, it is necessary that the protecting group P^{12} for a carboxy group of the amino acid or peptide (46) and the protecting group P^9 and P^{10} for the amino acid or peptide (31) can be removed in the same manner as described above but by a different method or different conditions. For example, a representative example includes a combination in which P^9 is a 9-fluorenylmethyloxy carbonyl group, P^{10} is a tert-butyl group or the like as a protecting group for a carboxy group, a methoxymethyl group or the like as a protecting group for a hydroxy group, or a tert-butyloxycarbonyl group as a protecting group or the like for an amino group, and P^{12} is a benzyl group. The protective group P^{10} for a functional group in a side chain is preferably a protecting group which can be deprotected by a treatment under acidic conditions. However, it is not limited thereto, and it can be selected from the aforementioned ones depending on, e.g., the properties of amino group, carboxy group, or a hydroxy group of a compound to be protected. For removing the protecting group, the reagents and conditions can be selected depending on the protecting group.


[0351] By deprotecting the protecting group P^9 for the N terminal of the peptide (49) obtained, the compound (50) can be produced. Reagents and conditions can be selected depending on the protecting group.

[0352] The compound (47) can be produced by derivatizing the carboxylic acid derivative (11) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the peptide (50) obtained in the presence of a base. The reaction conditions, reagents, base, and solvent used for forming an amide bond between the carboxylic acid derivative (11) and the peptide (50) can be suitably selected from those described for the synthesis of the compound (6).

6. Production method 6

[0353] Within the production intermediate (2), those in which the linker has a structure represented by $-L^1-L^2-L^P-$, and said L^P is the peptide residue containing a hydrophilic amino acid at the N terminal and said hydrophilic amino acid located at the N terminal is other than glycine, can be also produced by the following method.

[Chem. 28]



[in the formula, L^{1'} corresponds to L¹ having a structure in which the terminal is modified to maleimidyl group, L^P represents a structure consisting of -L^{P1}-L^{P2}-, and P⁸, P⁹, P¹⁰, and P¹² represent a protecting group].

[0354] Because L^P is formed by connecting L^{P1} to L^{P2}, the hydrophilic amino acid at N terminal of L^P is derived from L^{P1}, and thus, those that having a hydrophilic amino acid at the N terminal are suitably employed as L^{P1}. Meanwhile, plural hydrophilic amino acids may present therein. Further, when L^{P2} with hydrophilic amino acid is employed, L^P having plural hydrophilic amino acids at the N terminal of L^P or at the N terminal and at other positions can be produced depending on its location of hydrophilic amino acid.

[0355] The compound (51) can be produced by derivatizing the peptide or amino acid (28) described in Production method 5, which has the N terminal protected with P⁸, into active ester, mixed acid anhydride, or the like, and reacting with the compound (4) and a salt thereof. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the peptide or amino acid (28) and the compound (4) can be suitably selected from those described for the synthesis of the compound (6). The protective group P⁸ can be suitably selected and used from those described as the protective group for the compound (6), and it can be selected depending on, e.g., a property of the compound having an amino group to be protected. Further, as it is generally used for peptide synthesis, by repeating sequentially the reaction and deprotection of the amino acid or peptide constituting the peptide or amino acid (28) for elongation, the compound (51) can be also produced.

[0356] By deprotecting the protecting group P^8 for the amino group of the compound (51) obtained, the compound (52) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0357] The compound (53) can be produced by derivatizing the amino acid or peptide (31) having the N terminal protected with P^9 and the carboxy group, hydroxy group, or amino group in side chain protected with P^{10} as described in Production method 4 into active ester, mixed acid anhydride, or the like and reacting it with the compound (52) obtained. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the amino acid or peptide (31) and the compound (52) can be suitably selected from those described for the synthesis of the compound (6). The protecting group P^9 and P^{10} are the same as those described in Production method 5. Further, as it is generally used for peptide synthesis, by repeating sequentially the reaction and deprotection of the constituting amino acid or peptide for elongation, the compound (53) can be also produced.

[0358] By deprotection of P^9 as the protecting group of the amino group of the compound (53) obtained, the compound (54) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0359] It is possible to produce the compound (55) by derivatizing the carboxylic acid derivative (11) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (54) obtained. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the carboxylic acid derivative (11) and the compound (54) can be suitably selected from those described for the synthesis of the compound (6).

[0360] By deprotecting the protecting group P^{10} for the carboxy group, hydroxy group, or amino group of the compound (55) obtained, the compound (2) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0361] The compound (53) can be also produced by the following method, for example.

[0362] By deprotecting the protecting group P^{12} for the carboxy group of the compound (49) described in Production method 5, the peptide (56) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0363] The compound (53) can be produced by derivatizing the peptide (56) obtained into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (4) or a salt thereof. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the compound (56) and the compound (4) can be suitably selected from those described for the synthesis of the compound (6).


[0364] The compound (55) can be also produced by the following method, for example.

[0365] The compound (55) can be produced by derivatizing the compound (48) described in Production method 5 into active ester, mixed acid anhydride, or the like, and reacting it with the compound (4) in the presence of a base, or derivatizing the amino acid or peptide (45) described in Production method 5 into active ester, mixed acid anhydride, or the like, and reacting it with the compound (52) in the presence of a base. The reaction conditions, reagents, base, and solvent used for forming each peptide bond can be suitably selected from those described for the synthesis of the compound (6).

7. Production method 7

[0366] Within the production intermediate represented by the formula (2), those having the linker structure of $-L^1-L^2-L^P-NH-(CH_2)n^1-L^a-NH-(CH_2)n^2-C(=O)-$, and said L^P is the peptide residue having a hydrophilic amino acid at the N terminal, and said hydrophilic amino acid located at N terminal is other than glycine can be also produced by the following method.

[Chem. 29]



[in the formula, $L^{1'}$ corresponds to L^1 having a structure in which the terminal is modified to maleimidyl group, L^P represents a structure consisting of $-L^{P1}-L^{P2}-$, and P^9 and P^{13} represent a protecting group].

[0367] The production intermediate represented by the formula (2) includes the following two modes, that is, a structure in which the linker is represented by $-L^1-L^2-L^P-NH-(CH_2)_n-L^a-NH-(CH_2)_n-C(=O)-$ and a structure in which the linker is represented by $-L^1-L^2-L^P-$.

[0368] The compound (2) with a structure in which the linker is represented by $-L^1-L^2-L^P-NH-(CH_2)_n-L^a-NH-(CH_2)_n-C(=O)-$ can be produced as follows.

[0369] The compound (57) can be synthesized in the same manner as the compound (32) described in Production method 5. However, unlike the compound (32), it is not necessary that the protecting group P^9 for the amino group and the protecting group P^{13} for the functional group in side chain can be removed by a different method or different conditions. The functional group in side chain is a carboxy group or a hydroxy group, and the protecting group P^9 for the amino group and the protecting group P^{13} for the carboxy group or hydroxy group in side chain can be simultaneously deprotected. For example, a representative example includes a combination in which P^9 is a tert-butyloxy carbonyl group and P^{13} is a tert-butyl group or a trityl group, or P^3 is a benzyloxy carbonyl group and P^{13} is a benzyl group. The protecting groups can be suitably selected from the aforementioned ones with regard to the protecting groups for the compound (6) depending on, e.g., the properties of an amino group, a carboxy group, or a hydroxy group of the compound to be protected. For removal of the protecting groups, reagents and conditions are selected depending on the protecting group. By using the protected amino acid or peptide satisfying above properties, the compound (57) can be synthesized in the same manner as Production method 5.

[0370] By sequential or simultaneous deprotection of the protecting group P^9 and P^{13} of the compound (57), the compound (51) can be produced. Reagents and conditions can be selected depending on the protecting group.

[0371] A functional group in hydrophilic side chain of L^P in the compound (58) is not particularly protected, however, by reaction with the compound (11) derivatized into active ester, mixed acid anhydride, or the like in the presence of a base, the compound (2) can be produced. The reaction conditions, reagents, base, and solvent used for forming each peptide bond can be suitably selected from those described for the synthesis of the compound (6).

[0372] The compound (2) with a structure in which the linker is represented by $-L^1-L^2-L^P-$ can be produced as follows.

[0373] The compound (59) can be also synthesized in the same manner as the compound (53) described in Production method 6. However, unlike the compound (53), it may not be necessary that the protecting group P^3 for the amino group and the protecting group P^8 for the functional group in side chain can be removed by a different method or different conditions. The functional group in side chain is a carboxy group or a hydroxy group, and the protecting group P^9 for the amino group and the protecting group P^{13} for the carboxy group or hydroxy group in side chain can be simultaneously deprotected. For example, a representative example includes a combination in which P^9 is a tert-butyloxy carbonyl group and P^{13} is a tert-butyl group or a trityl group, or P^3 is a benzyloxy carbonyl group and P^{13} is a benzyl group. The protecting groups can be suitably selected from the aforementioned ones with regard to the protecting groups for the compound (6) depending on, e.g., the properties of an amino group, a carboxy group, or a hydroxy group of the compound to be protected. For removal of the protecting groups, reagents and conditions are selected depending on the protecting group. By using the protected amino acid or peptide satisfying above properties, the compound (59) can be synthesized in the same manner as Production method 6.

[0374] By sequential or simultaneous deprotection of the protecting group P^9 and P^{13} of the compound (59), the compound (53) can be produced. Reagents and conditions can be selected depending on the protecting group.

[0375] A functional group in hydrophilic side chain of L^P in the compound (60) is not particularly protected. However, by reaction with the compound (11) derivatized into active ester, mixed acid anhydride, or the like in the presence of a base, the compound (2) can be produced. The reaction conditions, reagents, base, and solvent used for forming each peptide bond can be suitably selected from those described for the synthesis of the compound (6).


8. Production method 8

[0376] The compound (43) shown in Production method 5 in which the linker $-L^P-$ has a structure of $-L^{P1}-Gly-Gly-Phe-Gly-$ can be also produced by the following method.

[Chem. 30]



61



31



62


[in the formula, P^9 and P^{10} represent a protecting group].

[0377] The compound (62) can be produced by derivatizing the amino acid or peptide (31) described in Production method 5 into active ester, mixed acid anhydride, acid halide, or the like and reacting it with glycyglycyl-L-phenylalanyl-N-[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano [3',4':6,7]indolizino[1,2-b]quinolin-1-yl]glycinamide (that is, free form of the pharmaceutical compound disclosed in International Publication No. WO 1997/46260) (61) or a salt thereof in the presence of a base. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the amino acid or peptide (31) and the compound (61) can be suitably selected from those described for the synthesis of the compound (6). The protecting group P^3 for N terminal and the protecting group P^{10} for the functional group in side chain are the same as those described in Production method 5. Meanwhile, the protecting group P^{10} for the functional group in side chain may not be present, and by performing the reaction using the amino acid or peptide (31) with N terminal protected only, the compound (62) can be obtained.

9. Production method 9

[0378] Among the compounds represented by the formula (2), a compound, in which the linker has a structure represented by $\text{-L}^1\text{-L}^2\text{-L}^{\text{P}}\text{-}$, and said L^{P} is the oligopeptide in which the C terminal is composed of 2 or 3 or more glycines and is connected to a drug, and the N terminal of said peptide residue is glycine in case a hydrophilic amino acid is present at N terminal, can be also produced according to the following method.

[Chem. 31]



[in the formula, $\text{L}^1\text{'}$ corresponds to L^1 having a structure in which the terminal is converted to maleimidyl group, L^{P} represents a structure consisting of $\text{L}^{\text{P}^1}\text{-L}^{\text{P}^2}$, and P^{12} and P^{14} represent a protecting group].

[0379] Because L^{P} is formed by connecting L^{P^1} to L^{P^2} , the number of glycines for constituting the C terminal of L^{P} contained therein can be designed in consideration of the number of glycines at C terminal in L^{P} and the number of repeated use thereof during the reaction.

[0380] The peptide (63) is an oligopeptide in which the C terminal is composed of 2 or 3 or more glycines, and the N

terminal is glycine in case the N terminal of said peptide residue is a hydrophilic amino acid, and further, said N terminal is protected with P¹⁴. As commonly employed for peptide synthesis, the peptide (63) can be synthesized by repeating sequentially the condensation reaction of the constituting amino acid or peptide and deprotection.

[0381] The compound (64) can be produced by derivatizing the peptide (63) into active ester, mixed acid anhydride, or the like and reacting it with the compound (4) or a salt thereof. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between peptide (63) and the compound (4) can be suitably selected from those described for the synthesis of the compound (6). The protecting group P¹⁴ can be suitably selected and used from those described for synthesis of the compound (6).

[0382] The compound (64) can be also produced by derivatizing the amino acid or peptide (65) with the N terminal protected with P¹⁴ into active ester, mixed acid anhydride, or the like and reacting it with the compound (52) described in Production method 6. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between amino acid or peptide (65) and the compound (52) can be suitably selected from those described for the synthesis of the compound (6). The protecting group P¹⁴ can be suitably selected and used from those described for synthesis of the compound (6).

[0383] By deprotecting the protecting group P¹⁴ for the amino group of the compound (64) obtained, the compound (66) can be produced. Reagents and conditions can be selected depending on the protecting group.

[0384] The compound (2) can be produced by derivatizing the carboxylic acid derivative (11) into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (66) obtained. The reaction conditions, reagents, base, and solvent used for forming an amide bond between the carboxylic acid derivative (11) and the compound (66) can be suitably selected from those described for the synthesis of the compound (6).

[0385] The compound (2) can be also produced by the following method.

[0386] In the compound (67), of which glycine at N terminal of L^{P1} is connected to L², and it can be produced in the same manner as the compound (45) described in Production method 5. The compound (68) can be produced by derivatizing the amino acid or peptide (46) described in Production method 5 into active ester, mixed acid anhydride, acid halide, or the like and reacting it with the compound (67). Herein, the amino acid or peptide (46) is an oligopeptide consisting of glycine or having C terminal consisting of 2 or 3 or more glycines, in which the C terminal is protected with P¹². The reaction conditions, reagents, base, and solvent used for forming an amide bond between amino acid or peptide (46) and the compound (67) can be suitably selected from those described for the synthesis of the compound (6).

[0387] The compound (68) can be also produced by derivatizing the compound (11) into active ester, mixed acid anhydride, or the like and reacting it with the peptide (69) having the C terminal protected with P¹². Herein, the peptide (69) is an oligopeptide in which the C terminal is composed of 2 or 3 or more glycines and the N terminal is glycine in case the N terminal of said peptide residue is a hydrophilic amino acid. As commonly employed for peptide synthesis, the peptide (69) can be produced by repeating sequentially the condensation reaction of the constituting amino acid or peptide and deprotection. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the peptide (69) and the compound (11) can be suitably selected from those described for the synthesis of the compound (6). The protecting group P¹² is preferably a protecting group which can be deprotected under acidic conditions but it is not limited thereto, and can be suitably selected and used from those described for synthesis of the compound (6).

[0388] By deprotecting the protecting group P¹² for the carboxy group of the compound (68) obtained, the compound (70) can be produced. In this deprotection, reagents and conditions can be selected depending on the protecting group.

[0389] The compound (2) can be produced by derivatizing the compound (70) into active ester, mixed acid anhydride, or the like and reacting it with the compound (4) or a salt thereof. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the compound (70) and the compound (4) can be suitably selected from those described for the synthesis of the compound (6).

[0390] In addition to above, the compound (2) can be also produced according to the following method.

[0391] The compound (2) can be produced by derivatizing the compound (52) described in Production method 6 into active ester, mixed acid anhydride, or the like and reacting it with the compound (67) in the presence of a base. The reaction conditions, reagents, base, and solvent used for forming a peptide bond between the compound (67) and the compound (52) can be suitably selected from those described for the synthesis of the compound (6).

[0392] Meanwhile, it is also possible that all of the intermediate compounds of Production method 1 to Production method 9 may be present as in form of salt and/or hydrate.

[0393] The antibody-drug conjugate of the present invention may absorb moisture to have adsorption water, for example, or turn into a hydrate when it is left in air or subjected to purification procedures such as recrystallization. Such a compound or a salt containing water are also included in the present invention.

[0394] A compound labeled with various radioactive or non-radioactive isotopes is also included in the present invention. One or more atoms constituting the antibody-drug conjugate of the present invention may contain an atomic isotope at non-natural ratio. Examples of the atomic isotope can include deuterium (²H), tritium (³H), iodine-125 (¹²⁵I), and carbon-14 (¹⁴C). Further, the compound of the present invention may be radioactive-labeled with a radioactive isotope such as tritium (³H), iodine-125 (¹²⁵I), carbon-14 (¹⁴C), copper-64 (⁶⁴Cu), zirconium-89 (⁸⁹Zr), iodine-124 (¹²⁴I),

fluorine-18 (^{18}F), indium-111 (^{111}I), carbon-11 (^{11}C), or iodine-131 (^{131}I). The compound labeled with a radioactive isotope is useful as a therapeutic or prophylactic agent, a reagent for research such as an assay reagent and an agent for diagnosis such as an *in vivo* diagnostic imaging agent. Without being related to radioactivity, any isotope variant type of the antibody-drug conjugate of the present invention is within the scope of the present invention.

{Pharmaceuticals/Medicines}

[0395] The anti-HER3 antibody-drug conjugate of the present invention exhibits a cytotoxic activity against cancer cells, and thus, as a medicine, it can be particularly used as a therapeutic agent and/or prophylactic agent for cancer.

[0396] Specifically, the anti-HER3 antibody-drug conjugate of the present invention can be selectively used as a medicine for chemotherapy, which is a major method for treating cancer, and as a result, can delay development of cancer cells, inhibit growth thereof, and further destroy the cancer cells. This can allow cancer patients to be free from symptoms caused by cancer or achieve improvement in QOL of cancer patients and attains a therapeutic effect by sustaining the lives of the cancer patients. Even if the anti-HER3 antibody-drug conjugate of the present invention does not reach to destroying cancer cells, it can achieve higher QOL of cancer patients while achieving long-term survival, by inhibiting or controlling the growth of cancer cells.

[0397] The anti-HER3 antibody-drug conjugate of the present invention can be used as a medicine alone in such medicinal therapy. In addition, the anti-HER3 antibody-drug conjugate of the present invention can be used as a medicine in combination with an additional therapy in adjuvant therapy and can be combined with surgical operation, radiotherapy, hormone therapy, or the like. Furthermore, the anti-HER3 antibody-drug conjugate of the present invention can also be used as a medicine for drug therapy in neoadjuvant therapy.

[0398] In addition to the therapeutic use as described above, the effect of suppressing the growth of tiny metastatic cancer cells and further destroying them can also be expected. Particularly, when the expression of HER3 is confirmed in primary cancer cells, inhibition of cancer metastasis or a prophylactic effect can be expected by administering the anti-HER3 antibody-drug conjugate of the present invention. For example, the effect of inhibiting and destroying cancer cells in a body fluid in the course of metastasis or the effect of, for example, inhibiting and destroying tiny cancer cells immediately after implantation in any tissue can be expected. Accordingly, inhibition of cancer metastasis or a prophylactic effect can be expected, particularly, after surgical removal of cancer.

[0399] The anti-HER3 antibody-drug conjugate of the present invention can be expected to produce a therapeutic effect by administration as systemic therapy to patients as well as by local administration to cancer tissues.

[0400] The antibody-drug conjugate (1) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, and anti-melanoma effects *in vitro*.

[0401] The antibody-drug conjugate (2) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, anti-colorectal cancer, and anti-melanoma effects *in vitro* and stronger anti-breast cancer and anti-melanoma effects *in vivo* than those of U1-59.

[0402] The antibody-drug conjugate (3) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, anti-ovarian, anti-colorectal cancer, and anti-melanoma effects *in vitro* and stronger anti-breast cancer, anti-lung cancer, anti-stomach cancer, and anti-melanoma effects *in vivo* than those of U1-59.

[0403] The antibody-drug conjugate (4) had excellent antitumor activity, safety, and physical properties and exhibited an anti-breast cancer effect *in vitro*.

[0404] The antibody-drug conjugate (5) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, and anti-melanoma effects *in vitro*.

[0405] The antibody-drug conjugate (6) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, and anti-melanoma effects *in vitro* and a stronger anti-breast cancer effect *in vivo* than that of U1-59.

[0406] The antibody-drug conjugate (7) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, and anti-melanoma effects *in vitro*.

[0407] The antibody-drug conjugate (8) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, and anti-melanoma effects *in vitro* and a stronger anti-breast cancer effect *in vivo* than that of U1-59.

[0408] The antibody-drug conjugate (9) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, anti-ovarian cancer, anti-colorectal cancer, and anti-melanoma effects *in vitro*.

[0409] The antibody-drug conjugate (10) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, anti-colorectal cancer, and anti-melanoma effects *in vitro* and stronger anti-breast cancer, anti-lung cancer, anti-colorectal cancer, anti-stomach cancer, and anti-melanoma effects *in vivo* than those of U1-59.

[0410] The antibody-drug conjugate (11) had excellent antitumor activity, safety, and physical properties and exhibited an anti-breast cancer effect *in vitro*.

[0411] The antibody-drug conjugate (12) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, anti-ovarian cancer, anti-colorectal cancer, and anti-melanoma effects *in vitro*.

[0412] The antibody-drug conjugate (13) had excellent antitumor activity, safety, and physical properties and exhibited anti-breast cancer, anti-lung cancer, anti-colorectal cancer, and anti-melanoma effects *in vitro* and stronger anti-breast cancer (including triple-negative breast cancer), anti-lung cancer, anti-stomach cancer, anti-pancreatic cancer, and anti-melanoma effects *in vivo* than those of U1-59.

[0413] The antibody-drug conjugate (14) had excellent antitumor activity, safety, and physical properties and exhibited an anti-breast cancer effect *in vitro*.

[0414] The antibody-drug conjugate (16a) had excellent antitumor activity, safety, and physical properties and exhibited an anti-breast cancer (including luminal and triple negative), anti-melanoma, anti-ovarian cancer, anti-bladder cancer, anti-lung cancer, anti-head and neck cancer, and anti-gastric cancer effects *in vivo* when it was administered alone or in combination with trastuzumab, gefinitib, cetuximab, panitumumab or pertuzumab.

[0415] Examples of the cancer type to which the anti-HER3 antibody-drug conjugate of the present invention is applied include lung cancer, kidney cancer, urothelial cancer, colorectal cancer, prostate cancer, glioblastoma multiforme, ovarian cancer, pancreatic cancer, breast cancer, metastatic breast cancer, luminal breast cancer, melanoma, liver cancer, bladder cancer, gastric (stomach) cancer, gastrointestinal stromal tumor, cervical cancer, head and neck cancer, esophageal cancer, epidermoid cancer, peritoneal cancer, adult glioblastoma multiforme, hepatic cancer, hepatocellular carcinoma, colon cancer, rectal cancer, colon and rectal cancer, endometrial cancer, uterus cancer, salivary cancer, renal cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anus carcinoma, penis cancer. Chemotherapy is the only current treatment indicated for particularly triple negative breast cancer (that lacks the expression of HER2, estrogen, and progesteron receptors) among breast cancers, which is said to have a poor prognosis. There have been almost no reports of HER3 expression in triple negative breast cancer. However if HER3 expression is observed in patients with triple negative breast cancer, then the anti-HER3 antibody-drug conjugate of the present invention can be used as a therapeutic agent and/or a preventive agent. However, it is not limited to them as long as it is a cancer cell expressing, in a cancer cell as a treatment subject, a protein which the antibody of the antibody-drug conjugate can recognize.

[0416] The treatment using the anti-HER3 antibody-drug conjugate of the present invention can target a cancer cell expressing, in a cancer cell as a treatment subject, HER3 protein which the antibody of the antibody-drug conjugate can recognize. In the present specification, the "cancer expressing HER3 protein" is cancer comprising cells having HER3 protein on their surface or cancer secreting HER3 protein into blood. The HER3 protein is overexpressed in various human tumors and can be evaluated using a method usually carried out, such as immunohistochemical staining method (IHC) for evaluating the overexpression of the HER3 protein in tumor (primary, metastatic) specimens, fluorescence in situ hybridization method (FISH) for evaluating the amplification of the HER3 gene, or enzyme-linked immunosorbent assay (ELISA) for evaluating the overexpression of the HER3 protein in blood specimens.

[0417] The anti-HER3 antibody-drug conjugate of the present invention exhibits an antitumor effect by recognizing and further internalizing HER3 protein expressed on cancer cell surface. Thus, the treatment subject of the anti-HER3 antibody-drug conjugate of the present invention is not limited to the "cancer expressing HER3 protein" and can also be, for example, leukemia, malignant lymphoma, plasmacytoma, myeloma, or sarcoma.

[0418] The anti-HER3 antibody-drug conjugate of the present invention can be preferably administered to a mammal, but it is more preferably administered to a human.

[0419] Substances used in a pharmaceutical composition comprising the anti-HER3 antibody-drug conjugate of the present invention can be suitably selected and applied from formulation additives or the like that are generally used in the art, in view of the dosage or administration concentration.

[0420] The anti-HER3 antibody-drug conjugate of the present invention can be administered as a pharmaceutical composition comprising at least one pharmaceutically compatible ingredient. For example, the pharmaceutical composition typically comprises at least one pharmaceutical carrier (for example, sterilized liquid). As described herein, examples of the liquid include water and oil (petroleum oil and oil of animal origin, plant origin, or synthetic origin). The oil may be, for example, peanut oil, soybean oil, mineral oil, sesame oil or the like. Water is a more typical carrier when the pharmaceutical composition is intravenously administered. Saline solution, an aqueous dextrose solution, and an aqueous glycerol solution can be also used as a liquid carrier, in particular, for an injection solution. A suitable pharmaceutical vehicle can be appropriately selected from those known in the art. If desired, the composition may also comprise a trace amount of a moisturizing agent, an emulsifying agent, or a pH buffering agent. Examples of suitable pharmaceutical carrier are disclosed in "Remington's Pharmaceutical Sciences" by E. W. Martin. The formulations correspond to an administration mode.

[0421] Various delivery systems are known and they can be used for administering the anti-HER3 antibody-drug conjugate of the present invention. Examples of the administration route can include intradermal, intramuscular, intraperitoneal, intravenous, and subcutaneous routes, but not limited thereto. The administration can be made by injection or bolus injection, for example. According to a specific preferred embodiment, the administration of the ligand-drug conjugate is performed by injection. Parenteral administration is a preferred administration route.

[0422] According to a representative embodiment, the pharmaceutical composition is prescribed, as a pharmaceutical composition suitable for intravenous administration to human, according to the conventional procedures. The composition for intravenous administration is typically a solution in a sterile and isotonic aqueous buffer solution. If necessary, the drug may contain a solubilizing agent and local anesthetics to alleviate pain at injection area (for example, lignocaine). Generally, the ingredient is provided individually as any one of lyophilized powder or an anhydrous concentrate contained in a container which is obtained by sealing in an ampoule or a sachet having an amount of the active agent or as a mixture in a unit dosage form. When the pharmaceutical is the form to be administered by injection, it may be administered from an injection bottle containing water or saline of sterile pharmaceutical grade. When the pharmaceutical is administered by injection, an ampoule of sterile water or saline for injection may be provided such that the aforementioned ingredients are admixed with each other before administration.

[0423] The pharmaceutical composition of the present invention may comprise only the anti-HER3 antibody-drug conjugate of the present application as an active ingredient or may comprise the anti-HER3 antibody-drug conjugate and at least one medicine (e.g., cancer-treating agent) other than the conjugate. The anti-HER3 antibody-drug conjugate of the present invention can be administered with another cancer-treating agent, and the anti-cancer effect may be enhanced accordingly. For example, another medicine such as an anti-cancer agent used for such purpose may be administered before administration of the pharmaceutical composition comprising the anti-HER3 antibody-drug conjugate of the present invention as an active ingredient or after administration of the pharmaceutical composition comprising the anti-HER3 antibody-drug conjugate as an active ingredient, or may be administered simultaneously with, separately (individually) from, or subsequently to the antibody-drug conjugate, and it may be administered while varying the administration interval for each. In the present invention, the case where the anti-HER3 antibody-drug conjugate of the present invention is administered simultaneously with another medicine as a single formulation containing the antibody-drug conjugate and the medicine and the case where the anti-HER3 antibody-drug conjugate of the present invention and another medicine are administered simultaneously or subsequently as separate formulations or administered while varying the administration interval for each are both included in the scope of the "pharmaceutical composition comprising the antibody-drug conjugate and another medicine". Examples of the cancer-treating agent include 5-FU, trastuzumab, trastuzumab emtansine (T-DM1), cetuximab, gefitinib, panitumumab, pertuzumab, abraxane, erlotinib, carboplatin, cisplatin, gemcitabine, capecitabine, irinotecan (CPT-11), paclitaxel, docetaxel, pemetrexed, sorafenib, vinblastine, vinorelbine, vemurafenib, medicines described in International Publication No. WO 2003/038043, LH-RH analogues (leuporelin, goserelin, or the like), estramustine phosphate, estrogen antagonist (tamoxifen, raloxifene, or the like), and an aromatase inhibitor (anastrozole, letrozole, exemestane, or the like), but it is not limited as long as it is a medicine having an antitumor activity. These cancer-treating agents can be classified, according to their targets, into: anti-FGFR agents such as cetuximab, gefitinib, and panitumumab; anti-HER2 agents such as trastuzumab, T-DM1, and pertuzumab; anti-HER3 agents such as patritumab, MM-121, and MM-111; anti-VEGF agents such as infliximab and adalimumab; etc. Further, they can be classified into: anti-EGFR antibodies such as cetuximab and panitumumab; anti-HER2 antibodies such as trastuzumab and pertuzumab; anti-HER3 antibodies such as patritumab, MM-121, and MM-111; anti-VEGF antibodies such as infliximab and adalimumab; etc. The anti-HER3 antibody-drug conjugate of the present invention exerts an excellent therapeutic effect when administered in combination with an anti-HER2 agent or an anti-HER2 antibody in i) the treatment of stomach cancer, breast cancer, triple-negative breast cancer, or the like or when administered in combination with an anti-EGFR agent or an anti-EGFR antibody in ii) the treatment of lung cancer, head and neck cancer, stomach cancer, breast cancer, triple-negative breast cancer, or the like. One or two or more medicines other than the conjugate can be used, and these medicines may be anti-cancer agents or may be medicines for alleviating side effect caused by companion medicines.

[0424] In the present invention, the "pharmaceutical composition comprising the anti-HER3 antibody-drug conjugate and another medicine" has the same meaning as a "pharmaceutical composition in which the anti-HER3 antibody-drug conjugate is to be administered in combination with another medicine". In the present invention, the phrase "administered in combination" used for the anti-HER3 antibody-drug conjugate and another medicine means that the anti-HER3 antibody-drug conjugate and another medicine are incorporated in the body of a recipient within a certain period. A single formulation containing the anti-HER3 antibody-drug conjugate and another medicine may be administered, or the anti-HER3 antibody-drug conjugate and another medicine may be separately formulated and administered as separate formulations. In the case of the separate formulations, the timings of administration thereof are not particularly limited, and the formulations may be administered at the same time or may be administered at different times or different days in a staggered manner. In the case where the anti-HER3 antibody-drug conjugate and another medicine are separately administered at different times or different days, the order of administration thereof is not particularly limited. Since separate formulations are usually administered according to their respective administration methods, the frequency of administration thereof may be the same or may be different. Further, such separate formulations may be administered by the same administration method (administration route) or different administration methods (administration routes). It is not necessary that the anti-HER3 antibody-drug conjugate and another medicine exist in the body at the same time, and it is sufficient that the anti-HER3 antibody-drug conjugate and another medicine are incorporated in the body within a certain

period (e.g., 1 month, preferably 1 week, more preferably several days, even more preferably 1 day). Alternatively, when one of the active ingredients is administered, the other active ingredient may have already disappeared from the body.

[0425] Examples of the dosage form of the "pharmaceutical composition in which the anti-HER3 antibody-drug conjugate is to be administered in combination with another medicine" can include: 1) the administration of a single formulation comprising the anti-HER3 antibody-drug conjugate and another medicine, 2) the simultaneous administration through the same administration route of two formulations obtained by separately formulating the anti-HER3 antibody-drug conjugate and another medicine, 3) the administration in a staggered manner through the same administration route of two formulations obtained by separately formulating the anti-HER3 antibody-drug conjugate and another medicine, 4) the simultaneous administration through different administration routes of two formulations obtained by separately formulating the anti-HER3 antibody-drug conjugate and another medicine, and 5) the administration in a staggered manner through different administration routes of two formulations obtained by separately formulating the anti-HER3 antibody-drug conjugate and another medicine. The dose, dosing interval, dosage form, formulation, etc., of the "pharmaceutical composition in which the anti-HER3 antibody-drug conjugate is to be administered in combination with another medicine" abide by those of the pharmaceutical composition containing the anti-HER3 antibody-drug conjugate of the present invention, but are not limited thereto.

[0426] Such a pharmaceutical composition formulated in two different formulations may be in the form of a kit containing them.

[0427] In the present invention, the "combination" of the anti-HER3 antibody-drug conjugate and another medicine means that the anti-HER3 antibody-drug conjugate and the medicine are "administered in combination".

[0428] The pharmaceutical composition can be formulated into a lyophilization formulation or a liquid formulation as a formulation having desired composition and required purity. When formulated as a lyophilization formulation, it may be a formulation containing suitable formulation additives that are used in the art. Also for a liquid formulation, it can be formulated as a liquid formulation containing various formulation additives that are used in the art.

[0429] Constituents and concentration of the pharmaceutical composition may vary depending on administration method. However, the anti-HER3 antibody-drug conjugate comprised in the pharmaceutical composition of the present invention can exhibit the pharmaceutical effect even at a small dosage when the antibody-drug conjugate has higher affinity for an antigen, that is, higher affinity (= lower K_d value) in terms of the dissociation constant (that is, K_d value) for the antigen. Thus, for determining dosage of the antibody-drug conjugate, the dosage can be determined in view of a situation relating to the affinity between the antibody-drug conjugate and antigen. When the antibody-drug conjugate of the present invention is administered to a human, for example, about 0.001 to 100 mg/kg can be administered once or administered several times with an interval of one time for 1 to 180 days.

Examples

[0430] The present invention is specifically described in view of the examples shown below. However, the present invention is not limited to them. Further, it is by no means interpreted in a limited sense. Further, unless specifically described otherwise, the reagent, solvent, and starting material described in the specification can be easily obtained from a commercial supplier.


Reference Example 1 Production of U1-59

[0431] U1-59 was produced on the basis of the method described in International Publication No. WO 2007/077028.

Example 1 Antibody-drug conjugate (1)

[0432]

[Chem. 32]



Process 1: tert-Butyl 4-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-4-oxobutyl]carbamate

[0433] 4-(tert-Butoxycarbonylamino)butanoic acid (0.237 g, 1.13 mmol) was dissolved in dichloromethane (10 mL), charged with N-hydroxysuccinimide (0.130 g, 1.13 mmol) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.216 g, 1.13 mmol), and stirred for 1 hour. The reaction solution was added dropwise to an N,N-dimethylformamide solution (10 mL) charged with exatecan mesylate (0.500 g, 0.94 mmol) and triethylamine (0.157 mL, 1.13 mmol), and stirred at room temperature for 1 day. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - chloroform : methanol = 8 : 2 (v/v)] to yield the titled compound (0.595 g, quantitative).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.87 (3H, t, J = 7.2 Hz), 1.31 (9H, s), 1.58 (1H, t, J = 7.2 Hz), 1.66 (2H, t, J = 7.2 Hz), 1.89-1.82 (2H, m), 2.12-2.21 (3H, m), 2.39 (3H, s), 2.92 (2H, t, J = 6.5 Hz), 3.17 (2H, s), 5.16 (1H, d, J = 19.2 Hz), 5.24 (1H, d, J = 18.8 Hz), 5.42 (2H, s), 5.59-5.55 (1H, m), 6.53 (1H, s), 6.78 (1H, t, J = 6.3 Hz), 7.30 (1H, s), 7.79 (1H, d, J = 11.0 Hz), 8.40 (1H, d, J = 8.6 Hz).

MS (APCI) m/z: 621 (M+H)⁺.

Process 2: 4-Amino-N-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]]butanamide trifluoroacetate

[0434] The compound (0.388 g, 0.61 mmol) obtained in above Process 1 was dissolved in dichloromethane (9 mL). After adding trifluoroacetic acid (9 mL), it was stirred for 4 hours. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - partitioned organic layer of chloroform :

methanol : water = 7 : 3 : 1 (v/v/v)] to yield the titled compound (0.343 g, quantitative).

¹H-NMR (400 MHz, DMSO-d₆) 0.87 (3H, t, J=7.2 Hz), 1.79-1.92 (4H, m), 2.10-2.17 (2H, m), 2.27 (2H, t, J=7.0 Hz), 2.40 (3H, s), 2.80-2.86 (2H, m), 3.15-3.20 (2H, m), 5.15 (1H, d, J=18.8 Hz), 5.26 (1H, d, J=18.8 Hz), 5.42 (2H, s), 5.54-5.61 (1H, m), 6.55 (1H, s), 7.32 (1H, s), 7.72 (3H, brs), 7.82 (1H, d, J=11.0 Hz), 8.54 (1H, d, J=8.6 Hz).
MS (APCI) m/z: 521 (M+H)⁺.

Process 3: N-(tert-butoxycarbonyl)glycylglycyl-L-phenylalanyl-N-(4-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-4-oxobutyl)glycinamide

[0435] N-(tert-butoxycarbonyl)glycylglycyl-L-phenylalanylglycine (0.081 g, 0.19 mmol) was dissolved in dichloromethane (3 mL), charged with N-hydroxysuccinimide (0.021 g, 0.19 mmol) and 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (0.036 g, 0.19 mmol), and stirred for 3.5 hours. The reaction solution was added dropwise to an N,N-dimethylformamide solution (1.5 mL) charged with the compound (0.080 g, 0.15 mmol) which has been obtained in above Process 2 and stirred at room temperature for 4 hours. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - chloroform : methanol = 8 : 2 (v/v)] to yield the titled compound (0.106 g, 73%).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.87 (3H, t, J = 7.4 Hz), 1.36 (9H, s), 1.71 (2H, m), 1.86 (2H, t, J = 7.8 Hz), 2.15-2.19 (4H, m), 2.40 (3H, s), 2.77 (1H, dd, J = 12.7, 8.8 Hz), 3.02 (1H, dd, J = 14.1, 4.7 Hz), 3.08-3.11 (2H, m), 3.16-3.19 (2H, m), 3.54 (2H, d, J=5.9 Hz), 3.57-3.77 (4H, m), 4.46-4.48 (1H, m), 5.16 (1H, d, J=19.2 Hz), 5.25 (1H, d, J = 18.8 Hz), 5.42 (2H, s), 5.55-5.60 (1H, m), 6.53 (1H, s), 7.00 (1H, t, J = 6.3 Hz), 7.17-7.26 (5H, m), 7.31 (1H, s), 7.71 (1H, t, J = 5.7 Hz), 7.80 (1H, d, J = 11.0 Hz), 7.92 (1H, t, J = 5.7 Hz), 8.15 (1H, d, J = 8.2 Hz), 8.27 (1H, t, J = 5.5 Hz), 8.46 (1H, d, J = 8.2 Hz).
MS (APCI) m/z: 939 (M+H)⁺.

Process 4: Glycylglycyl-L-phenylalanyl-N-(4-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-4-oxobutyl)glycinamide trifluoroacetate

[0436] The compound (1.97 g, 2.10 mmol) obtained in above Process 3 was dissolved in dichloromethane (7 mL). After adding trifluoroacetic acid (7 mL), it was stirred for 1 hour. The solvent was removed under reduced pressure, and it was charged with toluene for azeotropic distillation. The obtained residues were purified by silica gel column chromatography [chloroform - partitioned organic layer of chloroform : methanol : water = 7 : 3 : 1 (v/v/v)] to yield the titled compound (1.97 g, 99%).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.87 (3H, t, J = 7.4 Hz), 1.71-1.73 (2H, m), 1.82-1.90 (2H, m), 2.12-2.20 (4H, m), 2.40 (3H, s), 2.75 (1H, dd, J = 13.7, 9.4 Hz), 3.03-3.09 (3H, m), 3.18-3.19 (2H, m), 3.58-3.60 (2H, m), 3.64 (1H, d, J = 5.9 Hz), 3.69 (1H, d, J = 5.9 Hz), 3.72 (1H, d, J = 5.5 Hz), 3.87 (1H, dd, J = 16.8, 5.9 Hz), 4.50-4.56 (1H, m), 5.16 (1H, d, J = 19.2 Hz), 5.25 (1H, d, J = 18.8 Hz), 5.42 (2H, s), 5.55-5.60 (1H, m), 7.17-7.27 (5H, m), 7.32 (1H, s), 7.78-7.81 (2H, m), 7.95-7.97 (3H, m), 8.33-8.35 (2H, m), 8.48-8.51 (2H, m).
MS (APCI) m/z: 839 (M+H)⁺.

Process 5: N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]glycylglycyl-L-phenylalanyl-N-(4-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-4-oxobutyl)glycinamide

[0437] To an N,N-dimethylformamide (1.2 mL) solution of the compound (337 mg, 0.353 mmol) obtained in above Process 4, triethylamine (44.3 mL, 0.318 mmol) and N-succinimidyl 6-maleimidehexanoate (119.7 mg, 0.388 mmol) were added and stirred at room temperature for 1 hour. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - chloroform : methanol = 5:1 (v/v)] to yield the titled compound as a pale yellow solid (278.0 mg, 76%).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.87 (3H, t, J=7.3 Hz), 1.12-1.22 (2H, m), 1.40-1.51 (4H, m), 1.66-1.76 (2H, m), 1.80-1.91 (2H, m), 2.05-2.21 (6H, m), 2.39 (3H, s), 2.79 (1H, dd, J=14.0, 9.8 Hz), 2.98-3.21 (5H, m), 3.55-3.77 (8H, m), 4.41-4.48 (1H, m), 5.15 (1H, d, J=18.9 Hz), 5.24 (1H, d, J=18.9 Hz), 5.40 (1H, d, J=17.1 Hz), 5.44 (1H, d, J=17.1 Hz), 5.54-5.60 (1H, m), 6.53 (1H, s), 6.99 (2H, s), 7.20-7.27 (5H, m), 7.30 (1H, s), 7.70 (1H, t, J=5.5 Hz), 7.80 (1H, d, J=11.0

Hz), 8.03 (1H, t, J=5.8 Hz), 8.08 (1H, t, J=5.5 Hz), 8.14 (1H, d, J=7.9 Hz), 8.25 (1H, t, J=6.1 Hz), 8.46 (1H, d, J=8.5 Hz). MS (APCI) m/z: 1032 (M+H)⁺.

Process 6: Antibody-drug conjugate (1)

[0438] Reducing the antibody: U1-59 produced in Reference Example 1 was prepared to have antibody concentration of 10 mg/mL by replacing the medium with PBS6.0/EDTA by using the Common procedure B and Common procedure C described in Production method 1. The solution (1.00 mL) was added to a 2.0 mL polypropylene tube and charged with an aqueous solution of 10 mM TCEP (Tokyo Chemical Industry Co., Ltd.) (0.0307 mL; 4.6 equivalents per antibody molecule) and a 1 M aqueous solution of dipotassium hydrogen phosphate (Nacalai Tesque, Inc.; 0.050 mL). After confirming that the solution has pH of 7.4 +/- 0.1, the disulfide bond at hinge part in the antibody was reduced by incubating at 37C for 1 hour.

[0439] Conjugation between antibody and drug linker: After incubating the solution in a water bath at 22C for 10 minutes, dimethyl sulfoxide (Sigma-Aldrich Co. LLC; 0.0586 mL) and a dimethyl sulfoxide solution (0.0615 mL; 9.2 equivalents per antibody molecule) containing 10 mM of the compound obtained in above Process 5 were added thereto and incubated in a water bath at 22C for 40 minutes for conjugating the drug linker to the antibody. Next, an aqueous solution (0.0123 mL) of 100 mM NAC (Sigma-Aldrich Co. LLC) was added thereto and stirred by using a tube rotator (MTR-103, manufactured by AS ONE Corporation) at room temperature for 20 minutes to terminate reaction of the drug linker.

[0440] Purification: The above solution was subjected to purification using the Common procedure D (ABS is used as buffer solution) described in Production method 1 to yield 6 mL of a solution containing the titled antibody-drug conjugate.


[0441] Physicochemical characterization: By using the Common procedure E described in Production method 1 (as molar absorption coefficient of the drug linker, $E_{D,280} = 7280$ and $E_{D,370} = 23400$ were used), the following characteristic values were obtained.

[0442] Antibody concentration: 1.29 mg/mL, antibody yield: 7.74 mg (77%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E: 4.9.

Example 2 Antibody-drug conjugate (2)

[0443]

[Chem. 33]



Process 1: Antibody-drug conjugate (2)

[0444] By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 5 of Example 1, the titled antibody-drug conjugate was obtained in the same manner as Process 6 of Example 1.


[0445] Antibody concentration: 12.0 mg/mL, antibody yield: 226.8 mg (91%), and average number of conjugated drug

molecules (n) per antibody molecule measured by Common procedure E: 4.9.

Example 3 Antibody-drug conjugate (3)

[0446]

[Chem. 34]



Process 1: Antibody-drug conjugate (3)


[0447] By using U1-59 produced in Reference Example 1 and the compound obtained in Process 5 of Example 1, the titled antibody-drug conjugate was obtained in the same manner as Process 6 of Example 1.

[0448] Antibody concentration: 16.9 mg/mL, antibody yield: 219.7 mg (88%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E: 4.9.

Example 4 Antibody-drug conjugate (4)

[0449]

[Chem. 35]



Process 1: Antibody-drug conjugate (4)

[0450] Reducing the antibody: U1-59 produced in Reference Example 1 was prepared to have antibody concentration of 10 mg/mL by replacing the medium with PBS6.0/EDTA by using the Common procedure B and Common procedure C described in Production method 1. The solution (1.00 mL) was added to a 1.5 mL polypropylene tube and charged with an aqueous solution of 10 mM TCEP (Tokyo Chemical Industry Co., Ltd.) (0.0187 mL; 2.8 equivalents per antibody molecule) and a 1 M aqueous solution of dipotassium hydrogen phosphate (Nacalai Tesque, Inc.; 0.0170 mL). After confirming that the solution has pH of 7.0 +/- 0.1, the disulfide bond at hinge part in the antibody was reduced by incubating at 37°C for 1 hour.

[0451] Conjugation between antibody and drug linker: After adding a dimethyl sulfoxide solution (0.0314 mL; 4.7 equivalents per antibody molecule) containing 10 mM of the compound obtained in above Process 5 to the solution at room temperature, it was incubated at 15°C for 1 hour for conjugating the drug linker to the antibody. Next, an aqueous solution (0.0123 mL; 18.4 equivalents per antibody molecule) of 100 mM NAC (Sigma-Aldrich Co. LLC) was added thereto and incubated at room temperature for another 20 minutes to terminate reaction of the drug linker.

[0452] Purification: The above solution was subjected to purification using the Common procedure D (ABS is used as buffer solution) described in Production method 1 to yield 6 mL of a solution containing the titled antibody-drug conjugate. After that, the solution was concentrated by the Common procedure A.


[0453] Physicochemical characterization: By using the Common procedure E described in Production method 1 (as molar absorption coefficient of the drug linker, $E_{D,280} = 5000$, and $E_{D,370} = 19000$ were used), the following characteristic values were obtained.

[0454] Antibody concentration: 1.02 mg/mL, antibody yield: 6.1 mg (61%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E: 2.9; and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure F (as molar absorption coefficient of the drug linker, $E_{D,280} = 5000$ were used): 3.2.

Example 5 Antibody-drug conjugate (5)

[0455]

[Chem. 36]



Process 1: tert-Butyl (5S,14S)-5-benzyl-1-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-14-[[[(9H-fluoren-9-ylmethoxy)carbonyl]amino]-1,4,7,10,13-pentaoxo-3,6,9,12-tetraazahexadecan-16-oate

[0456] Under ice cooling, to an N,N-dimethylformamide (10.0 mL) solution of glycylglycyl-L-phenylalanyl-N-[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]glycinamide (free form of the pharmaceutical compound described in International Publication No. WO 1997/46260; 0.250 g, 0.332 mmol), N-hydroxysuccinimide (57.2 mg, 0.497 mmol), and 4-tert-butyl N-[(9H-fluoren-9-ylmethoxy)carbonyl]-L-aspartic acid (0.205 g, 0.497 mmol), N,N'-dicyclohexylcarbodiimide (0.123 g, 0.497 mmol) was added and stirred at room temperature for 2 days. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - chloroform : methanol = 9 : 1 (v/v)] to yield the titled compound as a pale yellow solid (0.278 g, 73%).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.86 (3H, t, J = 7.1 Hz), 1.35 (9H, s), 1.79-1.90 (2H, m), 2.03-2.25 (2H, m), 2.40 (3H, s), 2.40-2.51 (2H, m), 2.64-2.82 (2H, m), 2.98 (1H, dd, J = 13.7, 4.6 Hz), 3.16 (2H, brs), 3.55 (1H, dd, J = 16.7, 5.7 Hz), 3.63-3.80 (4H, m), 4.16-4.34 (3H, m), 4.36-4.50 (2H, m), 5.23 (2H, s), 5.37 (1H, d, J = 16.5 Hz), 5.43 (1H, d, J = 16.5 Hz), 5.51-5.62 (1H, m), 6.52 (1H, s), 7.10-7.25 (5H, m), 7.26-7.33 (3H, m), 7.39 (2H, t, J = 7.3 Hz), 7.65-7.72 (3H, m), 7.80 (1H, d, J = 11.0 Hz), 7.86 (2H, d, J = 7.3 Hz), 7.98 (1H, t, J = 5.5 Hz), 8.07 (1H, d, J = 7.8 Hz), 8.15 (1H, t, J = 5.5 Hz), 8.31 (1H, t, J = 5.5 Hz), 8.41 (1H, d, J = 8.7 Hz).

MS (ESI) m/z: 1147 (M+H)⁺.

EP 4 494 710 A2

Process 2: tert-Butyl (5S,14S)-14-amino-5-benzyl-1-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-1,4,7,10,13-pentaoxo-3,6,9,12-tetraazahexadecan-16-oate

- 5 **[0457]** To an N,N-dimethylformamide (2.00 mL) solution of the compound (0.279 g, 0.242 mmol) obtained in above Process 1, piperidine (0.240 mL, 2.42 mmol) was added and stirred at room temperature for 1 hour. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - chloroform : methanol = 2:1 (v/v)] to yield the titled compound as a pale yellow solid (0.265 g, quantitative).
¹H-NMR (400 MHz, DMSO-d₆) delta: 0.88 (3H, t, J = 7.2 Hz), 1.39 (9H, s), 1.81-1.94 (1H, m), 2.07-2.28 (2H, m), 2.37 (1H, dd, J = 15.8, 8.0 Hz), 2.43 (3H, s), 2.60 (1H, dd, J = 15.8, 4.9 Hz), 2.75-2.82 (1H, m), 3.00 (1H, dd, J = 13.9, 4.5 Hz), 3.16-3.25 (2H, m), 3.50-3.61 (2H, m), 3.65-3.81 (5H, m), 4.40-4.51 (1H, m), 5.27 (2H, dd, J = 24.1, 19.0 Hz), 5.43 (2H, dd, J = 21.3, 16.2 Hz), 5.56-5.65 (1H, m), 6.55 (1H, s), 7.15-7.28 (5H, m), 7.33 (1H, s), 7.83 (1H, d, J = 11.0 Hz), 8.04 (1H, t, J = 5.7 Hz), 8.09 (1H, d, J = 8.2 Hz), 8.26-8.39 (2H, m), 8.44 (1H, d, J = 8.2 Hz).

- 15 Process 3: tert-Butyl (5S,14S)-5-benzyl-14-[[[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]amino]-1-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-1,4,7,10,13-pentaoxo-3,6,9,12-tetraazahexadecan-16-oate

- 20 **[0458]** To an N,N-dimethylformamide (2.00 mL) solution of the compound (0.100 g, 0.108 mmol) obtained in above Process 2, N-succinimidyl 6-maleimide hexanoate (40.0 mg, 0.130 mmol) was added and stirred at room temperature for 2 days. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - chloroform : methanol = 9 : 1 (v/v)] to yield the titled compound as a pale yellow solid (80.0 mg, 66%).

- 25 ¹H-NMR (400 MHz, DMSO-d₆) delta: 0.88 (3H, t, J = 7.2 Hz), 1.13-1.23 (2H, m), 1.37 (9H, s), 1.42-1.54 (4H, m), 1.80-1.96 (2H, m), 2.08-2.25 (4H, m), 2.35-3.76 (15H, m), 2.43 (3H, s), 4.39-4.49 (1H, m), 4.55-4.67 (1H, m), 5.21-5.34 (2H, m), 5.43 (2H, dd, J = 21.1, 16.4 Hz), 5.56-5.64 (1H, m), 6.55 (1H, s), 7.01 (2H, d, J = 0.8 Hz), 7.16-7.26 (5H, m), 7.33 (1H, s), 7.83 (1H, d, J = 11.3 Hz), 8.04-8.18 (3H, m), 8.30-8.37 (1H, m), 8.43 (1H, d, J = 8.6 Hz).
 MS (ESI) m/z: 1118 (M+H)⁺.

- 30 Process 4: N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]-L-alpha-aspartylglycylglycyl-L-phenylalanyl-N-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]glycinamide

- 35 **[0459]** Under ice cooling, trifluoroacetic acid (4.00 mL) was added to the compound (70.0 mg, 62.6 μmol) obtained in above Process 3 and stirred at room temperature for 1 hour. The solvent was removed under reduced pressure to yield the titled compound as a pale yellow solid (55.0 mg, 83%).

- 40 ¹H-NMR (400 MHz, DMSO-d₆) delta: 0.88 (3H, t, J = 7.4 Hz), 1.14-1.24 (2H, m), 1.41-1.53 (4H, m), 1.79-1.95 (2H, m), 2.08-2.28 (4H, m), 2.37-2.60 (2H, m), 2.42 (3H, s), 2.63-2.82 (2H, m), 2.99 (1H, dd, J = 14.1, 5.1 Hz), 3.12-3.25 (2H, m), 3.29-3.44 (1H, m), 3.52-3.80 (6H, m), 4.38-4.48 (1H, m), 4.56 (1H, dd, J = 13.7, 7.4 Hz), 5.27 (2H, dd, J = 24.3, 18.8 Hz), 5.43 (2H, dd, J = 21.5, 16.4 Hz), 5.57-5.62 (1H, m), 6.55 (1H, s), 7.01 (2H, s), 7.15-7.26 (5H, m), 7.33 (1H, s), 7.82 (1H, d, J = 11.0 Hz), 7.98 (1H, brs), 8.08 (1H, d, J = 6.7 Hz), 8.15 (1H, d, J = 7.8 Hz), 8.34 (1H, brs), 8.44 (1H, d, J = 8.6 Hz), 12.26 (1H, brs).
 45 MS (ESI) m/z: 1062 (M+H)⁺.

Process 5: Antibody-drug conjugate (5)


- 50 **[0460]** By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 4, the titled antibody-drug conjugate was obtained in the same manner as Process 6 of Example 1.

[0461] Antibody concentration: 1.36 mg/mL, antibody yield: 8.16 mg (82%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, E_{D,280} = 7620, E_{D,370} = 23700 were used): 5.0.

- 55 Example 6 Antibody-drug conjugate (6)

[0462]

[Chem. 37]



Process 1: Antibody-drug conjugate (6)


[0463] By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 4 of Example 5, the titled antibody-drug conjugate was obtained in the same manner as Process 6 of Example 1.

[0464] Antibody concentration: 11.5 mg/mL, antibody yield: 224.2 mg (90%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 7620$, $E_{D,370} = 23700$ were used): 4.6.

Example 7 Antibody-drug conjugate (7)

[0465]

[Chem. 38]



Process 1: tert-Butyl (3S,12S)-12-benzyl-21-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-3-[(9H-fluoren-9-ylmethoxy)carbonyl]amino]-4,7,10,13,16,21-hexaoxo-5, 8,11,14,17-pentaazahenicosan-1-oate

[0466] (2S)-4-tert-Butoxy-2-[[[(9H-fluoren-9-ylmethoxy)carbonyl]amino]-4-oxobutanoic acid (0.625 g, 1.52 mmol) was dissolved in dichloromethane (10.0 mL), charged with N-hydroxysuccinimide (0.175 g, 1.52 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.291 g, 1.52 mmol), and stirred for 1 hour. The reaction solution was added dropwise to an N,N-dimethylformamide solution (10.0 mL) charged with the compound (1.00 g, 1.01 mmol) obtained in above Process 4 of Example 1 and stirred at room temperature for 20 hours. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - chloroform : methanol = 8 : 2 (v/v)] to yield the titled compound as a pale yellow solid (0.873 g, 70%).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.88 (3H, t, J=7.4 Hz), 1.37 (9H, s), 1.68-1.78 (2H, m), 1.81-1.93 (2H, m), 2.10-2.23 (4H, m), 2.41 (3H, s), 2.68-2.85 (3H, m), 2.99-3.22 (5H, m), 3.58-3.81 (6H, m), 4.19-4.36 (3H, m), 4.38-4.52 (2H, m), 5.17 (1H, d, J=19.2 Hz), 5.25 (1H, d, J=19.2 Hz), 5.43 (2H, s), 5.54-5.62 (1H, m), 6.55 (1H, s), 7.15-7.34 (8H, m), 7.41 (2H, t, J=7.2 Hz), 7.66-7.75 (4H, m), 7.81 (1H, d, J=11.0 Hz), 7.88 (2H, d, J=7.4 Hz), 8.01-8.06 (1H, m), 8.14 (1H, d, J=8.2 Hz), 8.17-8.22 (1H, m), 8.25-8.30 (1H, m), 8.47 (1H, d, J=8.6 Hz).

MS (APCI) m/z: 1232 (M+H)⁺.

EP 4 494 710 A2

Process 2: tert-Butyl (3S,12S)-12-benzyl-3-[[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]amino]-21-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-4,7,10,13,16,21-hexaoxo-5,8,11,14,17-pentaazahenicosan-1-oate

[0467] The compound (0.800 g, 0.649 mmol) obtained in above Process 1 was dissolved in N,N-dimethylformamide (3.00 mL), charged with piperidine (0.643 mL, 6.49 mmol), and stirred for 1 hour. The solvent was removed to dryness under reduced pressure and the obtained residues were dissolved in N,N-dimethylformamide (10 mL). After adding N-succinimidyl 6-maleimide hexanoate (0.300 g, 0.974 mmol), it was stirred for 20 hours. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - chloroform : methanol = 8 : 2 (v/v)] to yield the titled compound as a pale yellow solid (0.224 g, 29%).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.87 (3H, t, J = 7.6 Hz), 1.15-1.22 (2H, m), 1.35 (9H, s), 1.44-1.47 (4H, m), 1.71-1.73 (2H, m), 1.80-1.91 (2H, m), 2.08 (2H, t, J = 7.6 Hz), 2.13-2.20 (4H, m), 2.40 (3H, s), 2.67 (1H, dt, J = 11.1, 4.8 Hz), 2.78 (1H, dd, J = 13.6, 9.4 Hz), 2.99-3.17 (6H, m), 3.31-3.36 (2H, m), 3.57-3.76 (6H, m), 4.45-4.47 (1H, m), 4.57-4.60 (1H, m), 5.16 (1H, d, J = 18.7 Hz), 5.25 (1H, d, J = 18.7 Hz), 5.42 (2H, s), 5.55-5.60 (1H, m), 6.53 (1H, s), 6.99 (2H, s), 7.15-7.27 (5H, m), 7.31 (1H, s), 7.70 (1H, t, J = 5.4 Hz), 7.80 (1H, d, J = 10.9 Hz), 7.99 (1H, t, J = 5.7 Hz), 8.09-8.12 (3H, m), 8.25 (1H, t, J = 6.0 Hz), 8.45 (1H, d, J = 9.1 Hz).
MS (APCI) m/z: 1203 (M+H)⁺.

Process 3: N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]-L-alpha-aspartylglycylglycyl-L-phenylalanyl-N-(4-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-4-oxobutyl)glycinamide

[0468] The compound (0.224 g, 0.186 mmol) obtained in above Process 2 was reacted in the same manner as Process 2 of Example 1 to yield the titled compound as a pale yellow solid (21.2 mg, 10%).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.87 (3H, t, J = 7.2 Hz), 1.13-1.21 (2H, m), 1.42-1.45 (6H, m), 1.70-1.72 (2H, m), 1.85-1.88 (2H, m), 2.06-2.20 (6H, m), 2.39 (3H, s), 2.63-2.67 (1H, m), 2.78-2.81 (1H, m), 3.04-3.12 (6H, m), 3.63-3.70 (6H, m), 4.46-4.52 (2H, m), 5.16 (1H, d, J = 19.2 Hz), 5.25 (1H, d, J = 18.8 Hz), 5.42 (2H, s), 5.55-5.58 (1H, m), 6.53 (1H, s), 6.99 (2H, s), 7.18-7.23 (6H, m), 7.30 (1H, s), 7.71 (1H, t, J = 5.5 Hz), 7.79 (1H, d, J = 10.9 Hz), 7.99-8.02 (1H, m), 8.10-8.11 (3H, m), 8.27-8.30 (1H, m), 8.47-8.50 (1H, m).
MS (APCI) m/z: 1147 (M+H)⁺.

Process 4: Antibody-drug conjugate (7)


[0469] By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 3, the titled antibody-drug conjugate was obtained in the same manner as Process 6 of Example 1.

[0470] Antibody concentration: 1.39 mg/mL, antibody yield: 8.34 mg (83%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, E_{D,280} = 7670, E_{D,370} = 24800 were used): 4.7.

Example 8 Antibody-drug conjugate (8)

[0471]

[Chem. 39]



Process 1: Antibody-drug conjugate (8)

30 **[0472]** By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 3 of Example 7, the titled antibody-drug conjugate was obtained in the same manner as Process 6 of Example 1.

[0473] Antibody concentration: 11.2 mg/mL, antibody yield: 228.5 mg (91%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 7670$, $E_{D,370} = 24800$ were used): 4.7.

35 Example 9 Antibody-drug conjugate (9)

[0474]


40

45

50

55

[Chem. 40]



Process 1: N-{3-[2-(2-[[3-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)propanoyl]amino}ethoxy)ethoxy]propanoyl}glycylglycyl-L-phenylalanyl-N-(4-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-4-oxobutyl]glycinamide

[0475] The compound (100 mg, 0.119 mmol) obtained in above Process 4 of Example 1 was reacted in the same manner as Process 5 of Example 1 by using N-succinimidyl 3-(2-(2-(3-maleimidepropanamide)ethoxy)ethoxy)propanoate (50.7 mg, 0.119 mmol) instead of N-succinimidyl 6-maleimide hexanoate to yield the titled compound as a pale yellow solid (66.5 mg, 48%).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.85 (3H, t, J = 7.4 Hz), 1.65-1.74 (2H, m), 1.77-1.90 (2H, m), 2.07-2.19 (4H, m), 2.30 (2H, t, J = 7.2 Hz), 2.33-2.36 (2H, m), 2.38 (3H, s), 2.76 (1H, dd, J = 13.7, 9.8 Hz), 2.96-3.18 (9H, m), 3.42-3.44 (4H, m), 3.53-3.76 (10H, m), 4.43 (1H, td, J = 8.6, 4.7 Hz), 5.14 (1H, d, J = 18.8 Hz), 5.23 (1H, d, J = 18.8 Hz), 5.38 (1H, d, J = 17.2 Hz), 5.42 (1H, d, J = 17.2 Hz), 5.52-5.58 (1H, m), 6.52 (1H, s), 6.98 (2H, s), 7.12-7.17 (1H, m), 7.18-7.25 (4H, m), 7.29 (1H, s), 7.69 (1H, t, J = 5.5 Hz), 7.78 (1H, d, J = 11.3 Hz), 7.98-8.03 (2H, m), 8.11 (1H, d, J = 7.8 Hz), 8.16 (1H, t, J = 5.7 Hz), 8.23 (1H, t, J = 5.9 Hz), 8.44 (1H, d, J = 9.0 Hz). MS (APCI) m/z: 1149 (M+H)⁺.

Process 2: Antibody-drug conjugate (9)


[0476] By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 1, the titled antibody-drug conjugate was obtained in the same manner as Process 6 of Example 1.

[0477] Antibody concentration: 2.08 mg/mL, antibody yield: 18.7 mg (94%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, E_{D,280} = 4964, E_{D,370} = 18982 were used): 5.6.

Example 10 Antibody-drug conjugate (10)

[0478]

[Chem. 41]



Process 1: Antibody-drug conjugate (10)


[0479] By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 1 of Example 9, the titled antibody-drug conjugate was obtained in the same manner as Process 6 of Example 1.

[0480] Antibody concentration: 19.7 mg/mL, antibody yield: 236.4 mg (95%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 4964$, $E_{D,370} = 18982$ were used): 6.2; and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure F (as molar absorption coefficient of the drug linker, $E_{D,280} = 4964$ were used): 6.4.

Example 11 Antibody-drug conjugate (11)

[0481]

[Chem. 42]



Process 1: Antibody-drug conjugate (11)


[0482] By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 1 of Example 9, the titled antibody-drug conjugate was obtained in the same manner as Process 1 of Example 4.

[0483] Antibody concentration: 0.88 mg/mL, antibody yield: 5.28 mg (53%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 4964$, $E_{D,370} = 18982$ were used): 3.0; and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure F (as molar absorption coefficient of the drug linker, $E_{D,280} = 4964$ were used): 3.3.

Example 12 Antibody-drug conjugate (12)

[0484]

[Chem. 43]



Process 1: ({N-}[(9H-fluoren-9-ylmethoxy)carbonyl]glycyl)amino)methyl acetate

[0485] To a mixture containing N-9-fluorenylmethoxycarbonyl-L-glycyl-L-glycine (4.33 g, 12.2 mmol), tetrahydrofuran (120

EP 4 494 710 A2

ml), and toluene (40.0 ml), pyridine (1.16 mL, 14.7 mmol) and lead tetraacetate (6.84 g, 14.7 mmol) were added and refluxed under heating for 5 hours. After the reaction solution was cooled to room temperature, the insoluble material was removed by filtration through Celite, and the filtrate was concentrated under reduced pressure. The obtained residues were dissolved in ethyl acetate, washed with water and saturated brine, and then the organic layer was dried over anhydrous magnesium sulfate. After the solvent was removed under reduced pressure, the obtained residues were purified by silica gel column chromatography [hexane : ethyl acetate = 9 : 1 (v/v) - ethyl acetate] to yield the titled compound as colorless solid (3.00 g, 67%).

¹H-NMR (400 MHz, CDCl₃) delta: 2.07 (3H, s), 3.90 (2H, d, J = 5.1 Hz), 4.23 (1H, t, J = 7.0 Hz), 4.46 (2H, d, J = 6.6 Hz), 5.26 (2H, d, J = 7.0 Hz), 5.32 (1H, brs), 6.96 (1H, brs), 7.32 (2H, t, J = 7.3 Hz), 7.41 (2H, t, J = 7.3 Hz), 7.59 (2H, d, J = 7.3 Hz), 7.77 (2H, d, J = 7.3 Hz).

Process 2: Benzyl [(N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycyl)amino]methoxy]acetate

[0486] To a tetrahydrofuran (40.0 mL) solution of the compound (3.68 g, 10.0 mmol) obtained in above Process 1 and benzyl glycolate (4.99 g, 30.0 mmol), potassium tert-butoxide (2.24 g, 20.0 mmol) was added at 0°C and stirred at room temperature for 15 minutes. The reaction solution was charged with ethyl acetate and water at 0°C and extracted with ethyl acetate and chloroform. The obtained organic layer was dried over sodium sulfate and filtered. The solvent was removed under reduced pressure. The obtained residues were dissolved in dioxane (40.0 mL) and water (10.0 mL), charged with sodium hydrogen carbonate (1.01 g, 12.0 mmol) and 9-fluorenylmethyl chloroformate (2.59 g, 10.0 mmol), and stirred at room temperature for 2 hours. The reaction solution was charged with water and extracted with ethyl acetate. The obtained organic layer was dried over sodium sulfate and filtered. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [hexane : ethyl acetate = 100 : 0 (v/v) - 0 : 100] to yield the titled compound in colorless oily substance (1.88 g, 40%).

¹H-NMR (400 MHz, CDCl₃) delta: 3.84 (2H, d, J = 5.5 Hz), 4.24 (3H, t, J = 6.5 Hz), 4.49 (2H, d, J = 6.7 Hz), 4.88 (2H, d, J = 6.7 Hz), 5.15-5.27 (1H, m), 5.19 (2H, s), 6.74 (1H, brs), 7.31-7.39 (7H, m), 7.43 (2H, t, J = 7.4 Hz), 7.61 (2H, d, J = 7.4 Hz), 7.79 (2H, d, J = 7.4 Hz).

Process 3: [(N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycyl)amino]methoxy]acetic acid

[0487] The compound (1.88 g, 3.96 mmol) obtained in above Process 2 was dissolved in ethanol (40.0 mL) and ethyl acetate (20.0 mL). After adding palladium carbon catalyst (376 mg), it was stirred at room temperature under hydrogen atmosphere for 2 hours. The insoluble material was removed by filtration through Celite, and the solvent of the filtrate was removed under reduced pressure to yield the titled compound as colorless solid (1.52 g, quantitative).

¹H-NMR (400 MHz, DMSO-d₆) delta: 3.62 (2H, d, J = 6.3 Hz), 3.97 (2H, s), 4.18-4.32 (3H, m), 4.60 (2H, d, J = 6.7 Hz), 7.29-7.46 (4H, m), 7.58 (1H, t, J = 5.9 Hz), 7.72 (2H, d, J = 7.4 Hz), 7.90 (2H, d, J = 7.4 Hz), 8.71 (1H, t, J = 6.5 Hz).

Process 4: 9H-Fluoren-9-ylmethyl(2-[(2-[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl)amino]-2-oxoethoxy)methyl]amino)-2-oxoethyl)carbamate

[0488] Under ice cooling, to an N,N-dimethylformamide (10.0 mL) solution of exatecan mesylate (0.283 g, 0.533 mmol), N-hydroxysuccinimide (61.4 mg, 0.533 mmol), and the compound (0.205 g, 0.533 mmol) obtained in above Process 3, N,N-diisopropylethylamine (92.9 µL, 0.533 mmol) and N,N'-dicyclohexylcarbodiimide (0.143 g, 0.693 mmol) were added and stirred at room temperature for 3 days. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - partitioned organic layer of chloroform : methanol : water = 7 : 3 : 1 (v/v/v)] to yield the titled compound as a pale brown solid (0.352 g, 82%).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.81 (3H, t, J = 7.4 Hz), 1.73-1.87 (2H, m), 2.06-2.20 (2H, m), 2.34 (3H, s), 3.01-3.23 (2H, m), 3.58 (2H, d, J = 6.7 Hz), 3.98 (2H, s), 4.13-4.25 (3H, m), 4.60 (2H, d, J = 6.7 Hz), 5.09-5.22 (2H, m), 5.32-5.42 (2H, m), 5.50-5.59 (1H, m), 6.49 (1H, s), 7.24-7.30 (3H, m), 7.36 (2H, t, J = 7.4 Hz), 7.53 (1H, t, J = 6.3 Hz), 7.66 (2H, d, J = 7.4 Hz), 7.75 (1H, d, J = 11.0 Hz), 7.84 (2H, d, J = 7.4 Hz), 8.47 (1H, d, J = 8.6 Hz), 8.77 (1H, t, J = 6.7 Hz). MS (ESI) m/z: 802 (M+H)⁺.

Process 5: N-[(2-[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl)amino]-2-oxoethoxy)methyl]glycinamide

[0489] To an N,N-dimethylformamide (11.0 mL) solution of the compound (0.881 g, 1.10 mmol) obtained in above Process 4, piperidine (1.1 mL) was added and stirred at room temperature for 2 hours. The solvent was removed under

reduced pressure to yield a mixture containing the titled compound. The mixture was used for the next reaction without further purification.

Process 6: N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl-L-phenylalanyl-N-[(2-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-2-oxoethoxy)methyl]glycinamide

[0490] Under ice cooling, to an N,N-dimethylformamide (50.0 mL) solution of the mixture (0.439 mmol) obtained in above Process 5, N-hydroxysuccinimide (0.101 g, 0.878 mmol), and N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl-L-phenylalanine (the compound described in Japanese Patent Laid-Open No. 2002-60351; 0.440 g, 0.878 mmol), N,N'-dicyclohexylcarbodiimide (0.181 g, 0.878 mmol) was added and stirred at room temperature for 4 days. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - chloroform : methanol = 9 : 1 (v/v)] to yield the titled compound as a pale orange solid (0.269 g, 58%). MS (ESI) m/z: 1063 (M+H)⁺.

Process 7: Glycylglycyl-L-phenylalanyl-N-[(2-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-2-oxoethoxy)methyl]glycinamide

[0491] To an N,N-dimethylformamide (4.00 mL) solution of the compound (0.269 g, 0.253 mmol) obtained in above Process 6, piperidine (0.251 mL, 2.53 mmol) was added and stirred at room temperature for 2 hours. The solvent was removed under reduced pressure to yield a mixture containing the titled compound. The mixture was used for the next reaction without further purification.

Process 8: N-[6-(2,5-dioxo-2,5-dihydro-1H-pyrrol-1-yl)hexanoyl]glycylglycyl-L-phenylalanyl-N-[(2-[[[(1S,9S)-9-ethyl-5-fluoro-9-hydroxy-4-methyl-10,13-dioxo-2,3,9,10,13,15-hexahydro-1H,12H-benzo[de]pyrano[3',4':6,7]indolizino[1,2-b]quinolin-1-yl]amino]-2-oxoethoxy)methyl]glycinamide

[0492] To an N,N-dimethylformamide (10.0 mL) solution of the compound (0.253 mmol) obtained in above Process 7, N-succinimidyl 6-maleimide hexanoate (0.156 g, 0.506 mmol) was added and stirred at room temperature for 3 days. The solvent was removed under reduced pressure and the obtained residues were purified by silica gel column chromatography [chloroform - chloroform : methanol = 9 : 1 (v/v)] to yield the titled compound as a pale yellow solid (0.100 g, 38%).

¹H-NMR (400 MHz, DMSO-d₆) delta: 0.83 (3H, t, J = 7.2 Hz), 1.09-1.21 (2H, m), 1.33-1.47 (4H, m), 1.75-1.90 (2H, m), 2.00-2.23 (4H, m), 2.36 (3H, s), 2.69-2.81 (1H, m), 2.94-3.03 (1H, m), 3.06-3.22 (2H, m), 3.23-3.74 (6H, m), 3.98 (2H, s), 4.39-4.50 (1H, m), 4.60 (2H, d, J = 6.7 Hz), 5.17 (2H, s), 5.39 (2H, s), 5.53-5.61 (1H, m), 6.50 (1H, s), 6.96 (2H, s), 7.11-7.24 (5H, m), 7.28 (1H, s), 7.75 (1H, d, J = 11.0 Hz), 7.97 (1H, t, J = 5.7 Hz), 8.03 (1H, t, J = 5.9 Hz), 8.09 (1H, d, J = 7.8 Hz), 8.27 (1H, t, J = 6.5 Hz), 8.48 (1H, d, J = 9.0 Hz), 8.60 (1H, t, J = 6.5 Hz).

MS (ESI) m/z: 1034 (M+H)⁺.

Process 9: Antibody-drug conjugate (12)


[0493] By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 8, the titled antibody-drug conjugate was obtained in the same manner as Process 6 of Example 1.

[0494] Antibody concentration: 2.11 mg/mL, antibody yield: 19.0 mg (95%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, E_{D,280} = 5178, E_{D,370} = 20217 were used): 4.9.

Example 13 Antibody-drug conjugate (13)

[0495]

[Chem. 44]



Process 1: Antibody-drug conjugate (13)


[0496] By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 8 of Example 12, the titled antibody-drug conjugate was obtained in the same manner as Process 6 of Example 1.

[0497] Antibody concentration: 22.2 mg/mL, antibody yield: 244.2 mg (98%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$, $E_{D,370} = 20217$ were used): 6.2; and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure F (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$ were used): 7.0.

Example 14 Antibody-drug conjugate (14)

[0498]

[Chem. 45]



Process 1: Antibody-drug conjugate (14)

[0499] Reduction of the antibody: U1-59 produced in Reference Example 1 was prepared to have antibody concentration of 10 mg/mL by replacing the medium with PBS6.0/EDTA by using the Common procedure B and Common procedure C described in Production method 1. The solution (1.00 mL) was added to a 2.0 mL polypropylene tube and charged with an aqueous solution of 10 mM TCEP (Tokyo Chemical Industry Co., Ltd.) (0.0160 mL; 2.4 equivalents per antibody molecule) and a 1 M aqueous solution of dipotassium hydrogen phosphate (Nacalai Tesque, Inc.; 0.0150 mL). After confirming that the solution has pH of 7.0 +/- 0.1, the disulfide bond at hinge part in the antibody was reduced by incubating at 37C for 1 hour.

[0500] Conjugation between antibody and drug linker: After incubating the solution in a water bath at 15C for 10 minutes, dimethyl sulfoxide (Sigma-Aldrich Co. LLC; 0.0209 mL) and a dimethyl sulfoxide solution (0.0315 mL; 5.0 equivalents per antibody molecule) containing 10 mM of the compound obtained in above Process 8 of Example 12 was added thereto and incubated in a water bath at 15C for 60 minutes for conjugating the drug linker to the antibody. Next, an aqueous solution (0.0050 mL) of 100 mM NAC (Sigma-Aldrich Co. LLC) was added thereto and stirred by using a tube rotator (MTR-103, manufactured by AS ONE Corporation) at room temperature for another 20 minutes to terminate reaction of the drug linker.


[0501] According to the same purification processes and physicochemical characterizations as Process 6 of Example 1, the following characteristics values were obtained.

[0502] Antibody concentration: 1.46 mg/mL, antibody yield: 8.76 mg (88%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$, $E_{D,370} = 20217$ were used): 2.5; and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure F (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$ were used): 2.9.


Example 15 Antibody-drug conjugate (15)

[0503]

[Chem. 46]



Process 1



Process 1: Antibody-drug conjugate (15)

[0504] Reduction of the antibody: U1-59 produced in Reference Example 1 was prepared to have antibody concentration of 10 mg/mL by replacing the medium with PBS6.0/EDTA by using the Common procedure B and Common procedure C described in Production method 1. The solution (100 mL) was added to a 250 mL polycarbonate Erlenmeyer flask and charged with a 1 M aqueous solution of dipotassium hydrogen phosphate (1.70 mL) and then an aqueous solution of 10 mM TCEP (4.010 mL; 6.0 equivalents per antibody molecule) at room temperature with stirring using a magnetic stirrer.

After confirming that the solution has pH of 7.0 +/- 0.1, the stirring was stopped, and the disulfide bond at hinge part in the antibody was reduced by incubating at 37C for 1 hour.

[0505] Conjugation between antibody and drug linker: After cooling the above solution to 15C, a DMSO solution (6.684 mL; 10.0 equivalents per antibody molecule) containing 10 mM of the compound obtained in above Process 8 of Example 12 was gradually added thereto with stirring. The mixture was stirred at 15C for the first 30 minutes and, after stopping the stirring, incubated for another 1 hour for conjugating the drug linker to the antibody. Next, an aqueous solution (0.862 mL; 12.9 equivalents per antibody molecule) of 100 mM NAC was added thereto with stirring and incubated at room temperature for 20 minutes to terminate the reaction of unreacted drug linker.


[0506] Purification: A 20% aqueous acetic acid solution (about 0.6 mL) and ABS (100 mL) were gradually added to the solution with stirring to adjust pH of the solution to 5.5 +/- 0.1. This solution was subjected to microfiltration (Millipore Corp. Millex-HV filter, 0.45 μ m, PVDF membrane) to remove whitish matter. This solution was subjected to ultrafiltration purification using a ultrafiltration apparatus constituted by a ultrafiltration membrane (Merck Japan, Ltd., Pellicon XL Cassette, Biomax 50 KDa), a tube pump (Cole-Parmer International, USA, MasterFlex pump model 77521-40, pump head model 7518-00), and a tube (Cole-Parmer International, USA, MasterFlex tube L/S16). Specifically, by adding ABS dropwise (a total of 1600 mL) as a buffer solution for purification to the reaction solution while carrying out ultrafiltration purification, non-conjugated drug linkers and other low-molecular-weight reagents were removed while the buffer solution was replaced with ABS and further the solution was concentrated. The obtained purified solution was subjected to microfiltration (0.22 μ m (Millipore Corp. Millex-GV filter, PVDF membrane) to yield 37.5 mL of a solution containing the titled antibody-drug conjugate.

[0507] Antibody concentration: 26.5 mg/mL, antibody yield: 993.0 mg (90%), average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$ and $E_{D,370} = 20217$ were used): 6.3, and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure F (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$ was used): 7.3.

Example 16a Antibody-drug conjugate (16a)

[0508]

[Chem. 47]



Process 1: Antibody-drug conjugate (16a)

[0509] Reduction of the antibody: U1-59 produced in Reference Example 1 was prepared to have antibody concentration of 10 mg/mL by replacing the medium with PBS6.0/EDTA by using the Common procedure B and Common procedure C described in Production method 1. The solution (15 mL) was added to a 50 mL polyethylene terephthalate container and charged with a 1 M aqueous solution of dipotassium hydrogen phosphate (0.255 mL) and then an aqueous solution of 10

mM TCEP (0.601 mL; 6.0 equivalents per antibody molecule) at room temperature with stirring using a magnetic stirrer. After confirming that the solution has pH of 7.0 +/- 0.1, the disulfide bond at hinge part in the antibody was reduced by incubating at 37C for 2 hours.


[0510] Conjugation between antibody and drug linker: After cooling the above solution to 15C, a DMSO solution (1.002 mL; 10.0 equivalents per antibody molecule) containing 10 mM of the compound obtained in above Process 8 of Example 12 was gradually added thereto with stirring. The mixture was stirred at 15C for 30 minutes for conjugating the drug linker to the antibody. Next, an aqueous solution (0.129 mL; 12.9 equivalents per antibody molecule) of 100 mM NAC was added thereto with stirring and incubated at room temperature for 20 minutes to terminate the reaction of unreacted drug linker. According to the same purification processes and physicochemical characterizations as Process 6 of Example 1, the following characteristic values were obtained.

[0511] Antibody concentration: 2.36 mg/mL, antibody yield: 140 mg (59.5 mL) (94%), average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$ and $E_{D,370} = 20217$ were used): 6.4, and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure F (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$ was used): 7.7.


Example 16b Antibody-drug conjugate (16b)

[0512]

[Chem. 48]



Process 1



[0513] Reduction of the antibody: U1-59 produced in Reference Example 1 was prepared to have antibody concentration of 10 mg/mL by replacing the medium with PBS6.0/EDTA by using the Common procedure B and Common procedure C described in Production method 1. The solution (900 mL) was added to a 2000 mL polycarbonate Erlenmeyer flask and charged with a 1 M aqueous solution of dipotassium hydrogen phosphate (15.3 mL) and then an aqueous solution of 10 mM TCEP (36.1 mL; 6.0 equivalents per antibody molecule) at room temperature with stirring using a magnetic stirrer. After confirming that the solution has pH of 7.0 +/- 0.1, the stirring was stopped, and the disulfide bond at hinge part in the antibody was reduced by incubating at 37C for 2 hours.

[0514] Conjugation between antibody and drug linker: After cooling the above solution to 15C, a DMSO solution (60.16 mL; 10.0 equivalents per antibody molecule) containing 10 mM of the compound obtained from Process 8 of Example 12 was gradually added thereto with stirring. The mixture was stirred at 15C for 30 minutes for conjugating the drug linker to the antibody. Next, an aqueous solution (7.76 mL; 12.9 equivalents per antibody molecule) of 100 mM NAC was added thereto with stirring and incubated at room temperature for 20 minutes to terminate the reaction of unreacted drug linker.

[0515] Purification: A 20% aqueous acetic acid solution (about 5 mL) and ABS (1000 mL) were gradually added to the solution with stirring to adjust pH of the solution to 5.5 +/- 0.1. This solution was subjected to microfiltration (Millipore Corp. Stericup, 0.45 um, PVDF membrane) to remove whitish matter. This solution was subjected to ultrafiltration purification


using a ultrafiltration apparatus constituted by a ultrafiltration membrane (Merck Japan, Ltd., Pellicon 2 mini cassette, Ultracel 30 KDa, 0.1 m²), a tube pump (Cole-Parmer International, USA, MasterFlex pump model 7528-20, pump head model 77800-62), and a tube (Cole-Parmer International, USA, MasterFlex tubes L/S24 and 25). Specifically, by adding ABS dropwise (a total of 16 L) as a buffer solution for purification to the reaction solution while carrying out ultrafiltration purification, non-conjugated drug linkers and other low-molecular-weight reagents were removed while the buffer solution was replaced with ABS and further the solution was concentrated to yield about 500 mL of a solution containing the titled antibody-drug conjugate.

[0516] Antibody concentration: 19.66 mg/mL, antibody yield: 9830 mg (109%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$ and $E_{D,370} = 20217$ were used): 6.5.

Example 16c Antibody-drug conjugate (16c)

[0517]

[Chem. 49]




[0518] By using U1-59 produced in Reference Example 1 and the compound obtained in above Process 8 of Example 12, the titled antibody-drug conjugate was obtained in the same manner as Example 16b.

[0519] Antibody concentration: 16.21 mg/mL, antibody yield: 9726 mg (600 mL, 108%), and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$ and $E_{D,370} = 20217$ were used): 6.5.


Example 16d Antibody-drug conjugate (16d)

[0520]

[Chem. 50]



Process 1



[0521] The antibody-drug conjugates (16a), (16b), and (16c) produced in Examples 16a, 16b, and 16c, respectively, were mixed (a total of about 18 g) and further subjected to ultrafiltration in the same manner as Example 16b (11 L of ABS was used). The obtained purified solution was subjected to microfiltration (Millipore Corp. Stericup, 0.45 μ m and 0.22 μ m, PVDF membrane) to yield 745 mL of a solution containing the titled antibody-drug conjugate. By further adding 110 mL of ABS, 855 mL of a solution containing the titled antibody-drug conjugate was obtained.

[0522] Antibody concentration: 20.0 mg/mL, antibody yield: 17.1 g (94%), average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure E (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$ and $E_{D,370} = 20217$ were used): 6.5, and average number of conjugated drug molecules (n) per antibody molecule measured by Common procedure F (as molar absorption coefficient of the drug linker, $E_{D,280} = 5178$ was used): 7.8.

Test Example 1 HER3 binding affinity of antibody-drug conjugate compared with U1-59

Method:

[0523] A human breast cancer cell line HCC1569 (CRL-2330) from ATCC was cultured in an RPMI1640 medium (purchased from Invitrogen Corp., containing 10% bovine serum albumin (manufactured by Invitrogen Corp.) and 2 mM L-glutamine (manufactured by Invitrogen Corp.)). The cells were dissociated from the culture plate using ACCUTASE(R) SOLUTION (Millipore Corp., SCR005) or EDTA (5 mM, phosphate buffered saline (PBS, 137 mM sodium chloride, 2.7 mM potassium chloride, 1.47 mM potassium dihydrogen phosphate, and 10.5 mM disodium hydrogen phosphate)), and the number of living cells was measured by trypan blue treatment. The same numbers of cells suspended in a fluorescence-activated cell sorting (FACS) buffer (PBS containing 3% FBS and 0.004% sodium azide) were inoculated to 96-well U-bottom plates, and the cells were precipitated by centrifugation and suspended in 100 μ L of an ice-cooled antibody or antibody-drug conjugate dilution or FACS buffer.

[0524] The antibody or each antibody-drug conjugate was serially diluted at a ratio of 1/3 with a FACS buffer and adjusted to 30 μ g/mL to 5 ng/mL (200 nM to 0.03 nM). Cells treated with a FACS buffer without the addition of a primary antibody were used as a control group.

[0525] U1-59, the antibody-drug conjugate (3), the antibody-drug conjugate (10), or the antibody-drug conjugate (13) was evaluated as the antibody or the antibody-drug conjugates.

[0526] The cells of each group were reacted with a primary antibody dilution for 45 minutes on ice and then washed with a FACS buffer. Further, 100 μ L of a FACS buffer or a reaction solution of a secondary antibody diluted 1/100 (phycoerythrin (PE)-coupled anti human antibody, Dianova GmbH #709-116-149) was added thereto. The cells were treated for 45 minutes on ice in the dark and then washed with a FACS buffer, and dead cells were excluded using a FACS buffer or a FACS buffer supplemented with 7-aminoactinomycin D (7AAD, Sigma-Aldrich Co. LLC, #A9400, 1.1 μ g/mL).

[0527] Fluorescence signals from living cells were evaluated using Accuri C6 Flow cytometer (BD Biosciences/Accuri(R) Cytometers Inc., serial number 1424) and CFlow software (CFlow sampler Version 1.0.264.13).

[0528] For the correction of PE- and 7-AAD-derived signals, the fluorescence signals of U1-59 (30 μ g/mL) and cells stained with the PE-labeled secondary antibody or 7-AAD were evaluated.

[0529] In order to quantify U1-59-specific fluorescence signals in the cells, values obtained by the subtraction of FL-2 signals of cells treated with only the secondary antibody or 7-AAD were used. The equilibrium binding affinity (KD) and the maximum binding strength (Bmax) were calculated using GraphPad Prism software (version 5.04 for Windows(R) (one-site-specific binding)).

[0530] The results are shown in Figure 3 and Table 1. Figure 3 and Table 1 show the mean fluorescence intensity of HCC1569 treated with serial dilutions of U1-59 or each antibody-drug conjugate. The equilibrium binding affinity KD and the maximum binding strength Bmax were calculated using GraphPad Prism software.

[Table 1]

	U1-59	Antibody-drug conjugate (3)	Antibody-drug conjugate (10)	Antibody-drug conjugate (13)
Bmax	31830	28981	28841	28415
Kd (nM)	1.672	1.653	1.554	2.757

[0531] The antibody-drug conjugate (3) or the antibody-drug conjugate (10) exhibited average binding affinity KD for HCC-1569 equivalent to KD of the non-conjugated anti-HER3 antibody U1-59. The antibody-drug conjugate (13) also exhibited average binding affinity KD equivalent to KD of the non-conjugated anti-HER3 antibody U1-59 (2.7 nM vs. 1.6 nM). The KD values of the different antibody-drug conjugates suggested that the antibody-drug conjugation processes do not significantly impair the binding affinity of U1-59.

Test Example 2 Inhibition of HER3 signal by anti-HER3 antibody-drug conjugate

Method:

[0532] A human lung cancer cell line A549 (CRS-300114) from Cell Lines Service was dissociated by trypsin treatment, and 50,000 living cells were inoculated to 3 mL of DMEM/F12 (Invitrogen Corp., #21331-020) + 10% FBS (Invitrogen Corp., #10270-106) in each of 6 wells. After culturing the cells for 3 days, the medium was replaced with 2 mL of a fresh medium.

[0533] The antibody or each antibody-drug conjugate was added directly to 2 mL of a medium in each of the 6 wells such that the final concentration was 10 ug/mL (20 uL of a stock solution of 1 ug/uL antibody or antibody-drug conjugate was added).

[0534] U1-59, the antibody-drug conjugate (3), the antibody-drug conjugate (10), or the antibody-drug conjugate (13) was used as the antibody or the antibody-drug conjugate. An untreated group was used as a control.

[0535] The cells were cultured for 2 days, washed once with PBS, and treated with 100 uL of an ice-cooled buffer (50 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), pH 7.5, 150 mM sodium chloride, 1 mM ethylenediaminetetraacetic acid (EDTA), 12.5% glycerin, 1% Triton X-100, and 10 mM sodium pyrophosphate tetrabasic supplemented with proteinase inhibitors (Roche Diagnostics, Inc., #11697 498 001), 10 mM sodium fluoride, 1 mM sodium vanadate, 1 mM phenyl-methanesulfonyl-fluoride (PMSF), and 10 ug/mL aprotinin (Sigma-Aldrich Co. LLC, A1153)) for 30 minutes at 4C for lysis. The lysate was washed for 20 minutes at 13000 rpm at 4C, and the supernatant was used in protein concentration measurement by Bradford assay (Sigma-Aldrich Co. LLC, #B6916, BSA standard was from Thermo Fisher Scientific Inc., #23209). To each sample (amount of proteins: 120 ug), a 4-fold concentration of an LDS buffer (Invitrogen Corp., containing DTT (final concentration: 166.67 mM)) was added, and its volume was finally adjusted to 40 uL with water. The sample was boiled for 10 minutes at 70C and added to wells of NuPage Mini Bis-Tris gel (4%-12%, 1.5 mm thick, 10 slots/gel, Invitrogen Corp.). As protein standards, 7.5 uL of Novex(R) sharp ladder (Invitrogen Corp., PIN 57318) was added. The sample was electrophoresed for 70 minutes at 175 V with 1 × MOPS Running buffer (Invitrogen Corp.) containing NuPage antioxidant (Invitrogen Corp., NP0005, Lot 1356629 added to the internal chamber. Proteins separated by the gel electrophoresis were transferred to a nitrocellulose membrane (GE Healthcare Life Sciences) having a pore size of 0.45 um using NuPage transfer buffer (Invitrogen Corp.) containing 10% methanol and NuPage antioxidant (Invitrogen Corp., NP0005, Lot 1356629, 1:1000 dilution). The proteins were transferred for 80 minutes at a constant volume of 30 V.

[0536] The transfer membrane was cut, separated into fractions of 100 kDa or larger and 30 to 100 kDa, washed twice with PBS containing 0.1% Tween-20, and blocked by shaking for 1 hour at room temperature using Odyssey Blocking solution (LI-COR, Inc., #927-40000). The transfer membrane thus blocked was treated overnight at 4C with a solution of a diluted primary antibody (mixture of Odyssey blocking solution and PBS in equal amounts).

[0537] An anti-HER3 antibody (Santa Cruz Biotechnology, Inc., SC-81455, dilution 1:500) and an anti-phosphorylated HER antibody (Cell Signaling Technology, Inc., #4791, 1:1000) were used as the primary antibody, and an anti-actin

antibody (Neomarkers, #MS1295, dilution 1:3333) was used as an electrophoresis control.

[0538] The transfer membrane was washed three times (5 minutes for each) with PBS containing 0.1% Tween-20 and reacted with a dilution containing a secondary antibody (mixture of Odyssey blocking solution and PBS in equal amounts) for 1 hour at room temperature in a dark room.

[0539] Goat anti mouse IRDye 680RD (LI-COR, Inc., #926-68070, dilution 1:25000) or goat anti rabbit IR Dye 800CW (LI-COR, Inc., #926-32211, dilution 1:10000) was used as the secondary antibody. The transfer membrane was washed three times (6 minutes for each) with PBS containing 0.1% Tween-20, followed by signal detection using Odyssey infrared imager (LI-COR, Inc.).

[0540] The results are shown in Figure 4. A549 cells were cultured for 2 days with U1-59 or different antibody-drug conjugates. HER3 or phosphorylated HER3 was evaluated by Western blotting. pan-Actin was detected as an electrophoresis control.

[0541] As a result of culturing A549 for 2 days with 10 ug/mL U1-59 or antibody-drug conjugate, HER3 phosphorylation was reduced as compared with untreated cells. This reduction in HER3 phosphorylation was equivalent between U1-59 and the antibody-drug conjugate, suggesting that a plurality of drug conjugation processes did not impair a HER3 signal-inhibiting function derived from U1-59.

[0542] When A549 was treated with U1-59 or each antibody-drug conjugate for 2 days, reduction in HER3 expression was also observed as compared with untreated cells. The degree of this reduction in expression was equivalent between U1-59 and the antibody-drug conjugate. This suggests that a plurality of drug conjugation processes did not impair U1-59-mediated internalization (see Test Example regarding internalization) and U1-59-induced HER downregulation (see Test Example regarding signal inhibition).

Test Example 3 Reduction in expression of HER3 on cell surface by U1-59 and antibody-drug conjugate

Method:

[0543] HER3 internalization by U1-59 and each antibody-drug conjugate was evaluated by flow cytometry. 70,000 living cells of HCC1569 (from ATCC) were suspended in 0.5 mL of RPMI1640 (Invitrogen Corp., #31870-025) (containing 10% FBS (Invitrogen Corp., #10270-106 or PAN Biotech GmbH, #1505-P131304) and 2 mM glutamine (Invitrogen Corp., #25030-024)) and inoculated to each well of a 24-well plate. The cells were cultured for 4 days, and the medium was replaced with 0.5 mL of a fresh medium before the start of the internalization test. 5 ug of the antibody or each antibody-drug conjugate was added to 0.5 mL in each well of the 24-well plate such that the final concentration was 10 ug/mL. The cells were cultured for 1 hour at 37C in the presence of the antibody or the antibody-drug conjugate. U1-59, the antibody-drug conjugate (3), the antibody-drug conjugate (10), or the antibody-drug conjugate (13) was used as the antibody or the antibody-drug conjugate. Cells as a positive control or a negative control were untreated in some procedures.

[0544] For flow cytometry analysis, the cells were washed once with PBS and dissociated from the plate using 5 mM EDTA (100 uL/well) dissolved in ACCUTASE(R) SOLUTION (Millipore Corp., SCR005) or PBS. The cells were suspended in 200 uL of an ice-cooled FACS buffer (PBS containing 3% FBS and 0.004% sodium azide), then added to each well of a 96-well U-bottom plate, and left on ice. The cells were washed once with a FACS buffer. To each sample, 100 uL of U1-59 (10 ug/mL) diluted with a FACS buffer or only a FACS buffer was added. The cells were treated for 45 minutes with shaking on ice and then washed with a FACS buffer, and 100 uL of a PE-labeled secondary antibody anti-human antibody (Dianova GmbH, 709-116-149) dissolved at a ratio of 1:100 in a FACS buffer, or only a FACS buffer was added to each well. The cells were treated for 45 minutes in a dark room with shaking on ice. The cells were washed with a FACS buffer and treated with a FACS buffer or a FACS buffer containing 7AAD (Sigma-Aldrich Co. LLC, A9400, 1.1 ug to 1.25 ug/mL) for staining dead cells. The fluorescence signals of living cells were measured using Accuri C6 Flow cytometer. For the correction of PE and 7-AAD signals, U1-59 (10 ug/mL) and cells stained with only the PE-labeled secondary antibody or 7-AAD were used.

[0545] In order to quantify HER3-specific signals, values of cells stained with only the secondary antibody or 7-AAD were subtracted from FL-2 values of cells stained with the primary antibody and the secondary antibody, and further 7-AAD.

[0546] When the FL-2 signals of cells untreated with U1-59 or the antibody-drug conjugate (without internalization) were defined as the maximum value, reduction in HER3 (internalization) on the surface of the cells treated with U1-59 or the antibody-drug conjugate at 37C was calculated.

[0547] An average value calculated from 2 to 3 wells was used for the positive control (without treatment at 37C) and the negative control (without the addition of the primary antibody), and internalization in the wells of each treatment group was quantified.

[0548] The results are shown in Figure 5. This diagram shows an average value of reduction in HER3 expression on the surface of HCC1569 cells treated with U1-59 or each antibody-drug conjugate (37C, 1 hr). The HER3 expression of a group without the addition of U1-59 or the antibody-drug conjugate was defined as the maximum value of HER3 expression in cells. The values of groups treated with only the secondary antibody or 7-AAD were used as backgrounds. Groups treated with U1-59 or each antibody-drug conjugate for 1 hour were used as treatment groups. The FL-2 values were almost the

same between U1-59 and the antibody-drug conjugate.

[0549] Reduction in generated fluorescence caused by the treatment of HCC-1569 with U1-59 or each antibody-drug conjugate indicates reduction in HER3 expression. As compared with about 50% reduction in HER3 expression by U1-59, reduction in HER3 expression by each antibody-drug conjugate also exhibited a value equivalent to or higher than it, suggesting that the drug conjugation processes of the antibody did not impair the HER3-internalizing function of the antibody.

Test Example 4 Inhibition of *in vitro* mitogenic or survival signal by HER3 antibody-drug conjugate in human cancer cell line

Method:

[0550] The inhibitory activity of each HER3 antibody-drug conjugate against mitogenic or survival signals was measured in the presence of 10% FBS. The growth and development of cells were evaluated by measuring adenosine triphosphate (ATP) activity in untreated and antibody-drug conjugate-treated groups. Adherent cancer cell lines (human breast cancer cell line HCC1569 (CRL-2330) from ATCC, human breast cancer cell line MDA-MB-453 (CLB-22) from ATCC, and human colorectal cancer cell line HT-29 (CPQ-57) from ProQinase GmbH) were cultured in 2D culture systems, and floating (non-adherent) cancer cell lines (human melanoma cell line A375 (CRL-1619) from ATCC and human lung cancer cell line A549 (CRS-300114) from Cell Lines Service) were cultured in 3D culture systems.

Treatment of adherent cell

[0551] Each cancer cell line was suspended in 100 μ L of each medium at a low density (500 cells/well for HT-29, 800 cells/well for MDA-MB-453, and 1000 cells/well for HCC-1569) and inoculated to 96-MicroWell Optical Bottom plate (Thermo Fisher Scientific Inc./Nunc, #165306, white wall and clear bottom). As for HCC-1569 and MDA-MB-453, the cells were cultured in an RPMI1640 medium (Invitrogen Corp., 31870-025) containing 10% FBS (Invitrogen Corp., 10270-106) and 2 mM glutamine (Invitrogen Corp., 25030-024). As for HT-29, the cells were cultured in a McCoy's 5A medium (Invitrogen Corp., 26600-023) containing 10% FBS (Invitrogen Corp., 10270-106) and 2 mM glutamine (Invitrogen Corp., 25030-024). The edge wells of each plate were filled with 100 μ L of a medium.

[0552] The cells were cultured for 3 days, and the medium was replaced with 95 μ L of a fresh medium before antibody-drug conjugate treatment.

[0553] The antibody-drug conjugate (3), the antibody-drug conjugate (10), and the antibody-drug conjugate (13) were used. An untreated group was set as a control for measuring normal cell growth.

[0554] By adding 5 μ L of each antibody-drug conjugate concentrated into a 20-fold concentration to 95 μ L of a culture medium (10% FBS) contained in each well of the 96-well plate, the final concentration was established. Only 5 μ L of a medium was added to each control well. The test was conducted in triplicate per sample.

[0555] In order to calculate the concentration of the antibody-drug conjugate at which the growth or survival of cells was reduced by 50%, concentrations of the antibody-drug conjugate were prepared by 4-fold dilutions (10 μ g/mL to 0.15 ng/mL or 40 μ g/mL to 0.15 ng/mL), and the cells were treated with these concentrations of the antibody-drug conjugate and compared with the untreated group in terms of ATP activity. Evaluation was carried out by 5-day culture of HT-29 and 7-day culture of HCC-1569 and MDA-MB-453 thus supplemented with the antibody-drug conjugate. CellTiter-Glo(R) Luminescent Cell Viability Assay was used for evaluating the activity of the antibody-drug conjugate. This method involves measuring living cells having metabolic activity on the basis of ATP activity and finally estimating the number of living cells and employed CellTiter-Glo(R) Luminescent Cell (Promega K.K., G7573) as a kit.

[0556] 100 μ L of CellTiter-Glo(R) reagent was added to each well of the 96-well plate and stored for 25 minutes to 65 minutes at room temperature in a dark room before measurement using Wallac Victor2 1420 Multilabel Counter (program luminescence, measurement time: 0.5 s). Wells containing only a culture solution without the inoculation of the cells were assayed as blanks. In order to measure reduction in ATP activity, an average luminescence value of 3 wells was calculated under each condition (Microsoft Excel 2010). In order to remove cell-independent signals, the average luminescence value of the blanks was subtracted from the average luminescence value of the cells treated with the antibody-drug conjugate (Microsoft Excel 2010). The rate of reduction (%) in luminescence was calculated by comparison with the cells of the untreated group (Microsoft Excel 2010). This value was interpreted as the rate of inhibition (%) of cell growth or survival.

Treatment of floating cells

[0557] Since A375 and A549 have a faster growth rate than that of other cell lines, growth measurement was carried out in non-adherent 3D culture systems.

[0558] Each cancer cell line was suspended in 75 μ L of each medium at a low density (500 cells/well for A375 and 1500

cells/well for A549) and inoculated to a 96-well round-bottom non-adherent 3D culture plate (Prime Surface 96U; Sumitomo Bakelite Co, Ltd.; order no. MS-9096U). As for A375, the cells were cultured in a DMEM medium (Invitrogen Corp., 41965-039) containing 10% FBS (Invitrogen Corp., 10270-106) and 2 mM glutamine (Invitrogen Corp., 25030-024). As for A549, the cells were cultured in a DMEM/F12 medium (Invitrogen Corp., 21331-020) containing 10% FBS (Invitrogen Corp., 10270-106) and 2 mM glutamine (Invitrogen Corp., 25030-024). The edge wells of each plate were filled with 150 μ L of a medium. The cells were cultured for 3 days, and the final dose was adjusted to 142.5 μ L or 150 μ L by adding 67.5 or 75 μ L of a fresh medium before antibody-drug conjugate addition. The antibody-drug conjugate (3), the antibody-drug conjugate (10), and the antibody-drug conjugate (13) were used. An untreated group was set as a control for measuring normal cell growth.

[0559] By adding 7.5 or 8 μ L of each antibody-drug conjugate concentrated into a 20-fold concentration to 142.5 or 150 μ L of a culture medium (10% FBS) contained in each well of the 96-well plate, the final concentration was established. The final dose was set to 150 μ L or 158 μ L. Only 7.5 or 8 μ L of a medium was added to each control well. The test was conducted in triplicate per sample.

[0560] In order to calculate the concentration of the antibody-drug conjugate at which the growth or survival of cells was reduced by 50%, concentrations of the antibody-drug conjugate were prepared by 4-fold dilutions (10 μ g/mL to 0.15 ng/mL or 40 μ g/mL to 0.15 ng/mL), and the cells were treated with these concentrations of the antibody-drug conjugate and compared with the untreated group in terms of ATP activity. Evaluation was carried out by 7-day culture of the cells thus supplemented with the antibody-drug conjugate. CellTiter-Glo(R) Luminescent Cell Viability Assay was used for evaluating the activity of the antibody-drug conjugate. This method involves measuring living cells having metabolic activity on the basis of ATP activity and finally estimating the number of living cells and employed CellTiter-Glo(R) Luminescent Cell (Promega K.K., G7573) as a kit.

[0561] Before measurement, 50 μ L of the medium was removed from each well, and 100 μ L of CellTiter-Glo(R) reagent was added to each well of the 96-well plate and stored for 30 minutes to 55 minutes at room temperature in a dark room before measurement using Wallac Victor2 1420 Multilabel Counter (program luminescence, measurement time: 0.5 s). Before measurement, 180 μ L was collected from each well and transferred to measurable 96-MicroWell Optical Bottom white plate. Wells containing only a culture solution without the inoculation of the cells were assayed as blanks. The method for calculating the concentration of the antibody-drug conjugate at which the growth or survival of cells was inhibited by 50% was described in the evaluation method as to the adherent cells.

[0562] The results about the human breast cancer lines HCC1569 and MDA-MB453 are shown in Figures 6 and 7, respectively. The results about the human melanoma line A375 is shown in Figure 8. The results about the human colorectal cancer line HT29 are shown in Figure 9. The results about the human lung cancer line A549 are shown in Figure 10. A of each figure shows cell growth or survival derived from the antibody-drug conjugate in the presence of 10% FBS. The ordinate depicts a luminescence value indicating the ATP activity of each sample. The abscissa depicts the concentration of each antibody-drug conjugate. The data is indicated by mean $\pm$ standard deviation of triplicates. B of each figure shows the rate of reduction in luminescence caused by antibody-drug conjugate treatment when the luminescence of an untreated group was defined as 100%.

[0563] Three types of antibody-drug conjugates were added to various human cancer cell lines in the presence of 10% FBS and evaluated for *in vitro* growth in 2D or 3D systems. The rate of inhibition of cell growth or development in the untreated group or by each antibody-drug conjugate was calculated from the CellTiter-Glo(R) assay of ATP activity. In the evaluation of the ATP activity, these antibody-drug conjugates strongly inhibited the cell growth or survival of two types of breast cancer cell lines (HCC-1569 and MDA-MB-453) and one type of human melanoma line (A375).

[0564] The addition of each antibody-drug conjugate and culture (in the presence of 10% FBS) for 7 days reduced the ATP activity by 55 to 75% in HCC1569, by 60 to 83% in MDA-MB-453, and by 60 to 70% in A375. The inhibitory activity of the antibody-drug conjugate against cell growth or survival was not strong in the human colorectal cancer line HT-29 and the human lung cancer line A549 compared with the human breast cancer and human melanoma lines. In HCC-1569 and MDA-MB-453, the antibody-drug conjugate (10) exhibited strong inhibitory activity, which did not largely differ from the inhibitory activity of the antibody-drug conjugate (3) *in vitro*. By contrast, in both of the human breast cancer lines, the antibody-drug conjugate (13) exhibited low activity and required a concentration of 15 nM for achieving 50% inhibition of cell growth or survival, though the antibody-drug conjugate (3) or the antibody-drug conjugate (10) achieved this inhibition at 1 nM or lower. In the human melanoma line compared with the human breast cancer lines, the activity of the antibody-drug conjugate (13) was equivalent to that of the antibody-drug conjugate (3) or the antibody-drug conjugate (10).

[0565] All of the antibody-drug conjugates supported a maximum rate of inhibition on the order of 61 to 68%. The antibody-drug conjugates achieved 50% inhibition of ATP activity at a concentration of 1 to 4 nM.

[0566] In addition to the aforementioned test, the inhibitory activity of the antibody-drug conjugate (13) against the cell growth or survival of a human ovarian cancer cell line OVCAR-8 was also confirmed *in vitro* (data not shown).

Test Example 5 Comparison of rate of inhibition of *in vitro* cell growth or survival of human cancer cell line depending on the number of drug molecules (high or medium) loaded on antibody-drug conjugate

Method:

[0567] Antibody-drug conjugates differing in the number of loaded drug molecules were evaluated for *in vitro* inhibitory activity against cell growth or survival. The high drug loading represents the state where 5 to 7 drug molecules are conjugated with an antibody, and the middle drug loading represents the state where about 3 drug molecules are conjugated with an antibody. The average number of drug molecules conjugated with one antibody was measured by the UV method (described in other parts of the present invention).

[0568] The average number of drug molecules conjugated with one antibody:

High drug loading <HDL>

Antibody-drug conjugate (3): 4.9
Antibody-drug conjugate (10): 6.2
Antibody-drug conjugate (13): 6.2

Middle drug loading <MDL>

Antibody-drug conjugate (4): 2.9
Antibody-drug conjugate (11): 3.0
Antibody-drug conjugate (14): 2.5

[0569] The inhibitory activity of each HER3 antibody-drug conjugate against mitogenic or survival signals was measured in the presence of 10% FBS. The growth and development of cells were evaluated by measuring adenosine triphosphate (ATP) activity in untreated and antibody-drug conjugate-treated groups. The cancer cell line was suspended in 100 μ L of each medium at a low density (750 cells/well for a human breast cancer cell line MDA-MB-453 (CLB-22)) from ATCC) and inoculated to 96-MicroWell Optical Bottom white plate (Thermo Fisher Scientific Inc./Nunc, #165306). The cells were cultured in an RPMI1640 medium (Invitrogen Corp., 31870-025) containing 10% FBS (Invitrogen Corp., 10270-106) and 2 mM glutamine (Invitrogen Corp., 25030-024). The edge wells of each plate were filled with 100 μ L of a medium.

[0570] The cells were cultured for 3 days, and the medium was replaced with 95 μ L of a fresh medium before antibody-drug conjugate addition. The antibody-drug conjugate (3), the antibody-drug conjugate (10), the antibody-drug conjugate (13), the antibody-drug conjugate (4), the antibody-drug conjugate (11), and the antibody-drug conjugate (14) were used. An untreated group was set as a control for measuring normal cell growth.

[0571] By adding 5 μ L of each antibody-drug conjugate concentrated into a 20-fold concentration to 95 μ L of a culture medium (10% FBS) contained in each well of the 96-well plate, the final concentration was established. Only 5 μ L of a medium was added to each control well. The test was conducted in triplicate per sample. In order to calculate the concentration of the antibody-drug conjugate at which the growth or survival of cells was reduced by 50%, concentrations of the antibody-drug conjugate were prepared by 4-fold dilutions (10 μ g/mL to 0.15 ng/mL), and the cells were treated with these concentrations of the antibody-drug conjugate and compared with the untreated group in terms of ATP activity. Evaluation was carried out by 7-day culture of the cells thus supplemented with the antibody-drug conjugate. CellTiter-Glo(R) Luminescent Cell Viability Assay was used for evaluating the activity of the antibody-drug conjugate. This method involves measuring living cells having metabolic activity on the basis of ATP activity and finally estimating the number of living cells and employed CellTiter-Glo(R) Luminescent Cell (Promega K.K., G7573) as a kit. 100 μ L of CellTiter-Glo(R) reagent was added to each well of the 96-well plate and stored for 25 minutes to 55 minutes at room temperature in a dark room before measurement using Wallac Victor2 1420 Multilabel Counter (program luminescence, measurement time: 0.5 s). Wells containing only a culture solution without the inoculation of the cells were assayed as blanks. In order to measure reduction in ATP activity, an average luminescence value of 3 wells was calculated under each condition (Microsoft Excel 2010). In order to remove cell-independent signals, the average luminescence value of the blanks was subtracted from the average luminescence value of the cells treated with the antibody-drug conjugate (Microsoft Excel 2010). The rate of reduction (%) in luminescence was calculated by comparison with the cells of the untreated group (Microsoft Excel 2010). This value was interpreted as the rate of inhibition (%) of cell growth or survival.

[0572] The results are shown in Figures 11 to 13. Figure 11 shows the results of comparing the antibody-drug conjugate (3) with the antibody-drug conjugate (4). Figure 12 shows the results of comparing the antibody-drug conjugate (10) with the antibody-drug conjugate (11). Figure 13 shows the results of comparing the antibody-drug conjugate (13) with the antibody-drug conjugate (14). In each figure, the left diagram shows the rate of inhibition of cell growth or survival derived

from the antibody-drug conjugate in the presence of 10% FBS in one of the triplicate tests. The ordinate depicts luminescence indicating the ATP activity of each sample. The abscissa depicts the concentration of each antibody-drug conjugate. The right diagram shows the comparison of the rate of reduction in luminescence caused by antibody-drug conjugate treatment between high drug loading (HDL) and middle drug loading (MDL) when the luminescence of an untreated group was defined as 100%.

[0573] The high drug loading and middle drug loading antibody-drug conjugates inhibited cell growth or survival through the treatment and 7-day culture of MDA-MB-453. As already shown in the results about the high drug loading, the middle drug loading antibody-drug conjugate (11) exhibited high activity at the same level as that of the antibody-drug conjugate (4). In the comparison between the numbers of loaded drug molecules, the antibody-drug conjugates having a high number of loaded drug molecules exhibited higher reduction in ATP than that of the middle drug loading ones. The antibody-drug conjugate (3), the antibody-drug conjugate (10), and the antibody-drug conjugate (13) having a high number of loaded drug molecules exhibited rates of inhibition of 68%, 76%, and 56%, respectively, at a concentration of 10 ug/mL, whereas the antibody-drug conjugate (4), the antibody-drug conjugate (11), and the antibody-drug conjugate (14) having a middle number of loaded drug molecules merely exhibited rates of inhibition of 44%, 47%, and 27%, respectively, at this concentration. The antibody-drug conjugates having a high number of loaded drug molecules were superior in 50% inhibition concentration of cancer cell growth or survival to the antibody-drug conjugates having a middle number of loaded drug molecules. The antibody-drug conjugate (3) or the antibody-drug conjugate (10) having a high number of loaded drug molecules required 15 ng/mL (1 nM) of the antibody-drug conjugate for reducing the ATP activity value by 50%. The antibody-drug conjugate (13) reduced the ATP activity by 50% at least at the highest concentration tested. By contrast, reduction in ATP activity corresponding to this was not observed at 1000 ng/mL (67 nM) or lower within the range of concentrations of the evaluated antibody-drug conjugates having a middle number of loaded drug molecules. The *in vitro* comparison between the high number of loaded drug molecules and the middle number of loaded drug molecules suggested that an antibody-drug conjugate having a high number of loaded drug molecules is also superior in *in vivo* inhibitory activity against the growth of cancer cells.

Test Example 6 Antibody-drug conjugates (3), (10), and (13) exhibited antitumor effect in *in vivo* antitumor test using human breast cancer

[0574] Five-week-old female BALB/C nude mice having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day. 5×10^6 cells of a human breast cancer cell line HCC1569 (CRL-2330) from ATCC were suspended in a solution prepared from 50 uL of PBS and Matrigel (PBS: PAA #H21-002, Matrigel: BD #354230) mixed at a ratio of 1:1, and subcutaneously transplanted to the right side area of the body of each BALB/C nude mouse using a 29 G needle.

[0575] The body weights were measured using a weight scale (Mettler Toledo PB602-L). The major axis and minor axis of the tumor were measured twice a week using an electronic digital caliper (manual caliper, OMC Fontana), and the tumor volume (mm³) was calculated. The calculation was carried out according to the following expression (the same holds true for Test Examples described below).

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0576] At Day 19 when the tumor size reached about 150 mm³, 70 animals were randomly divided into 7 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (3), (10), or (13) or PBS for a control group were administered into the tail vein of each animal at the following doses.

[0577] Administration group: PBS was intravenously injected once a week at the same dose as the antibody-drug conjugate.

[0578] Administration group: The antibody-drug conjugate (3) was intravenously injected once a week at 3 or 10 mg/kg.

[0579] Administration group: The antibody-drug conjugate (10) was intravenously injected once a week at 3 or 10 mg/kg.

[0580] Administration group: The antibody-drug conjugate (13) was intravenously injected once a week at 3 or 10 mg/kg.

[0581] All data were indicated by mean $\pm$ SEM. The tumor sizes and the body weights were evaluated by mean $\pm$ SEM. All data were analyzed using Microsoft Excel 2009 (the same holds true for Test Examples described below).

[0582] The results are shown in Figure 14. The PBS administration group was euthanized at Day 53 after the transplantation, because the tumor sizes exceeded the acceptable maximum level. The inhibition of tumor growth of the human breast cancer cell line was observed in all of the antibody-drug conjugate administration groups compared with the control group. No weight loss was observed in the mice of the treated groups.

Test Example 7 Antibody-drug conjugates (3), (10), and (13) exhibited antitumor effect in antitumor test using human melanoma

[0583] Five- to 6-week-old female NMRI nude mice having a body weight of 22 to 26 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0584] 5×10^6 cells of a human melanoma cell line HT-144 (HTB-63) from ATCC were suspended in a solution prepared from 50 μ L of PBS and Matrigel (PBS: PAA #H21-002, Matrigel: BD #354230) mixed at a ratio of 1:1, and subcutaneously transplanted to the right side area of the body of each NMRI nude mouse using a 29 G needle.

[0585] The measurement of the body weights and the tumor sizes and the measurement and calculation of the tumor volumes were carried out in the same manner as Test Example 6.

[0586] At Day 22 when the tumor size reached about 150 mm³, 80 animals were randomly divided into 8 groups on the basis of their tumor sizes. At the same day, U1-59 or the antibody-drug conjugate (3), (10), or (13) or PBS for a control group were administered into the tail vein of each animal at the following doses.

[0587] Administration group: PBS was intravenously injected once a week at the same dose as the antibody-drug conjugate.

[0588] Administration group: U1-59 was subcutaneously injected twice a week at 25 mg/kg.

[0589] Administration group: The antibody-drug conjugate (3) was intravenously injected once a week at 3 or 10 mg/kg.

[0590] Administration group: The antibody-drug conjugate (10) was intravenously injected once a week at 3 or 10 mg/kg.

[0591] Administration group: The antibody-drug conjugate (13) was intravenously injected once a week at 3 or 10 mg/kg.

[0592] The results are shown in Figure 15. The PBS and U1-59 administration groups were euthanized at Days 48 and 52, respectively, after the transplantation, because the tumor sizes exceeded the acceptable maximum level. The inhibition of tumor growth of the human melanoma cell line was observed in all of the antibody-drug conjugate administration groups compared with the control group and the U1-59 administration group.

Test Example 8 Antibody-drug conjugates (3), (10), and (13) exhibited antitumor effect in antitumor test using human breast cancer line

[0593] Sixteen-week-old female SCID nude mice having a body weight of 17 to 25.5 g after acclimation were used. The mice were placed in individually ventilated cages which were kept at room temperature and a constant humidity.

[0594] For solid tumors derived from a human breast cancer cell line MDA-MB-453 (CLB-22) from ATCC, MDA-MB-453 in the first passage (Batch 1089) was transplanted to 3 mice (two areas per mouse: right and left side areas of the body). After 13 to 17 weeks, tumor sections were recovered and cryopreserved. For the second passage, the tumor section ($2 \times 2 \times 2$ mm) of the first passage was further subcutaneously transplanted (10 mice, two areas per mouse: right and left side areas of the body) and allowed to grow for tumor formation for 7 weeks. The tumor thus formed was prepared into a tumor section ($2 \times 2 \times 2$ mm, second passage) and transplanted to the right side area of the body of each SCID nude mouse.

[0595] The body weights were measured using a weight scale. The major axis (length) and diameter of the tumor were measured using an electronic digital caliper (Pro-Max 150 mm hand-held calipers, Fred V. Fowler Co., Inc.), and the tumor volume (mm³) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = \pi/6 \times \text{Major axis (mm)} \times [\text{Diameter (mm)}]^2$$

[0596] At Day 40 when the tumor size reached about 143 mm³, 72 animals were randomly divided into 8 groups on the basis of their tumor sizes. At the same day, U1-59 or the antibody-drug conjugate (3), (10), or (13) or PBS for a control group were administered into the tail vein of each animal at the following doses.

[0597] Administration group: PBS was intravenously injected once a week at the same dose as the antibody-drug conjugate.

[0598] Administration group: U1-59 was subcutaneously injected twice a week at 25 mg/kg.

[0599] Administration group: The antibody-drug conjugate (3) was intravenously injected once a week at 3 or 10 mg/kg.

[0600] Administration group: The antibody-drug conjugate (10) was intravenously injected once a week at 3 or 10 mg/kg.

[0601] Administration group: The antibody-drug conjugate (13) was intravenously injected once a week at 3 or 10 mg/kg.

[0602] The results are shown in Figure 16. The inhibition of tumor growth of the human breast cancer cell line was observed in all of the antibody-drug conjugate administration groups compared with the control group and the U1-59 administration group. No weight loss was observed in the mice of the treated groups. Further, in other antitumor test using human breast cancer cell line HCC1954 or JIMT1-PR10 (trastuzumab-, pertuzumab- and T-DM1- resistant), the inhibition of tumor growth was also observed in the antibody-drug conjugate (16a) administration group compared with the control

group.

Test Example 9 Antibody-drug conjugates (3), (10), and (13) exhibited antitumor effect in antitumor test using human colorectal cancer line

[0603] Five- to 6-week-old female NMRI nude mice having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0604] 4×10^6 cells of a human colorectal cancer cell line HT-29 (CPQ-57) from ProQinase GmbH were suspended in a solution prepared from PBS and Matrigel (PBS: PAA #H21-002, Matrigel: BD #354230) mixed at a ratio of 1:1, and subcutaneously transplanted to the right side area of the body of each NMRI nude mouse using a 29 G needle.

[0605] The measurement of the body weights and the tumor sizes and the measurement and calculation of the tumor volumes were carried out in the same manner as Test Example 6.

[0606] At Day 8 when the tumor size reached about 150 mm^3 , 70 animals were randomly divided into 7 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (3), (10), or (13) or PBS for a control group were administered into the tail vein of each animal at the following doses.

[0607] Administration group: PBS was intravenously injected once a week at the same dose as the antibody-drug conjugate.

[0608] Administration group: The antibody-drug conjugate (3) was intravenously injected once a week at 3 or 10 mg/kg.

[0609] Administration group: The antibody-drug conjugate (10) was intravenously injected once a week at 3 or 10 mg/kg.

[0610] Administration group: The antibody-drug conjugate (13) was intravenously injected once a week at 3 or 10 mg/kg.

[0611] The results are shown in Figure 17. The PBS administration group (6 out of the 10 mice), the groups given the antibody-drug conjugate (3) at 3 mg/kg (3 out of the 10 mice) and at 10 mg/kg (4 out of the 10 mice), the groups given the antibody-drug conjugate (10) at 3 mg/kg (2 out of the 10 mice) and at 10 mg/kg (2 out of the 10 mice), and the group given the antibody-drug conjugate (13) at 3 mg/kg (2 out of the 10 mice) were euthanized at Day 50 after the transplantation, because the tumor sizes exceeded the acceptable maximum level or ulcer was formed. The inhibition of tumor growth of the human colorectal cancer cell line was observed in all of the antibody-drug conjugate administration groups compared with the control group. The antibody-drug conjugate (13) exhibited stronger antitumor activity than that of the antibody-drug conjugate (3) or the antibody-drug conjugate (10). No weight loss was observed in the mice of the treated groups.

Test Example 10 Antibody-drug conjugates (3), (10), and (13) exhibited antitumor effect in antitumor test using human lung cancer line

[0612] Five- to 6-week-old female CD1 nude mice having a body weight of 24 to 28 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0613] 5×10^6 cells of a human lung cancer cell line A549 (CRS-300114) from Cell Lines Service were suspended in a solution prepared from 200 μL of PBS and Matrigel (PBS: PAA #H21-002, Matrigel: BD #354230) mixed at a ratio of 1:1, and subcutaneously transplanted to the right side area of the body of each CD1 nude mouse using a 29 G needle.

[0614] The measurement of the body weights and the tumor sizes and the measurement and calculation of the tumor volumes were carried out in the same manner as Test Example 6.

[0615] At Day 38 when the tumor size reached about 200 mm^3 , 70 animals were randomly divided into 7 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (3), (10), or (13) or PBS for a control group were administered into the tail vein of each animal at the following doses.

[0616] Administration group: PBS was intravenously injected once a week at the same dose as the antibody-drug conjugate.

[0617] Administration group: The antibody-drug conjugate (3) was intravenously injected once a week at 3 or 10 mg/kg.

[0618] Administration group: The antibody-drug conjugate (10) was intravenously injected once a week at 3 or 10 mg/kg.

[0619] Administration group: The antibody-drug conjugate (13) was intravenously injected once a week at 3 or 10 mg/kg.

[0620] The results are shown in Figure 18. The inhibition of tumor growth of the human lung cancer cell line was observed in all of the antibody-drug conjugate administration groups compared with the control group. No weight loss was observed in the mice of the treated groups.

Test Example 11 Antibody-drug conjugate (13) exhibited antitumor effect in *in vivo* antitumor test using human triple-negative breast cancer line

[0621] The triple-negative breast cancer refers to a breast cancer that neither expresses hormone receptors (estrogen receptor and progesterone receptor) nor expresses HER2. Since these receptors are not expressed, hormone treatment (tamoxifen, etc.) or anti-HER2 treatment (trastuzumab, trastuzumab emtansine, or pertuzumab) cannot be applied to the cancer. This breast cancer therefore leads to low survival rates, and many therapeutic agents are still under clinical trial. As the expression of HER3 was confirmed in a human triple-negative breast cancer line MDA-MB-468 (data not shown), the antitumor activity of the antibody-drug conjugate was evaluated. Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrJrlj [Foxn1nu/Foxn1nu] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0622] 5×10^6 cells of a human triple-negative breast cancer cell line MDA-MB-468 (CRL-2322) from ATCC were suspended in a solution prepared from 200 μ L of PBS and Matrigel (PBS: PAA #10010-023, Matrigel: BD #354234) mixed at a ratio of 1:1, and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0623] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm^3) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0624] At Day 20 when the tumor size reached about 170 mm^3 , 18 animals were randomly divided into 3 groups on the basis of their tumor sizes. At the same day, U1-59 or the antibody-drug conjugate (13) or PBS for a control group were administered into the tail vein of each animal at the following doses.

[0625] Administration group: PBS was intravenously injected once a week at the same dose as the antibody-drug conjugate.

[0626] Administration group: U1-59 was intravenously injected once a week at 10 mg/kg.

[0627] Administration group: The antibody-drug conjugate (13) was intravenously injected once a week at 10 mg/kg.

[0628] The results are shown in Figure 19. The inhibition of tumor growth of the human triple-negative breast cancer line was observed in the antibody-drug conjugate administration group compared with the control group and the U1-59 administration group. No weight loss was observed in the mice of the treated groups.

Test Example 12 Antibody-drug conjugate (16a) exhibited antitumor effect in *in vivo* antitumor test using human luminal breast cancer line

[0629] The human luminal breast cancer refers to a breast cancer that expresses hormone receptors (estrogen receptor), but expresses no HER2. Since these receptors are not expressed, anti-HER2 treatment (trastuzumab, trastuzumab emtansine, pertuzumab) cannot be applied to the cancer. This breast cancer therefore leads to low survival rates, and many therapeutic agents are still under clinical trial. As the expression of HER3 was confirmed in a human luminal breast cancer line MCF-7 (data not shown), the antitumor activity of the antibody-drug conjugate was evaluated.

[0630] Five- to 6-week-old female athymic nude mice Nude-Foxn1^{nu} (ProQinase GmbH) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0631] 5×10^6 cells of a human luminal breast cancer cell line MCF-7 (CRQ-#327) were suspended in a solution prepared from 200 μ L of PBS and Matrigel (PBS: PAA #10010-023, Matrigel: BD #354234) mixed at a ratio of 1:1, and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0632] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm^3) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0633] At Day 11 when the tumor size reached about 250 mm³, 20 animals were randomly divided into 2 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (16a) or PBS for a control group were administered at the following doses.

[0634] Administration group: PBS was intravenously injected at the same single dose as the antibody-drug conjugate.

[0635] Administration group: The antibody-drug conjugate (16a) was intravenously injected at a single dose of 10 mg/kg.

[0636] The results are shown in Figure 20. The inhibition of tumor growth of the human luminal breast cancer line was observed in the antibody-drug conjugate administration group compared with the control group. No weight loss was observed in the mice of the treated groups.

Test Example 13 Antibody-drug conjugate (16a) exhibited antitumor effect in *in vivo* antitumor test using human melanoma line

[0637] Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrIcrlj [Foxn1nu/Foxn1nu] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0638] 3×10⁶ cells of a human melanoma cell line WM-266-4 (CRL-1676) from ATCC were mixed and suspended in Matrigel (BD #354234) and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0639] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm³) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0640] At Day 19 when the tumor size reached about 220 mm³, 8 animals were randomly divided into 2 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (16a) or PBS for a control group were administered at the following doses.

[0641] Administration group: PBS was intravenously injected at the same single dose as the antibody-drug conjugate.

[0642] Administration group: The antibody-drug conjugate (16a) was intravenously injected at a single dose of 10 mg/kg.

[0643] The results are shown in Figure 21. The inhibition of tumor growth of the human melanoma line was observed in the antibody-drug conjugate administration group compared with the control group. No weight loss was observed in the mice of the treated groups. In other human melanoma model C32, the inhibition of tumor growth was also observed in the antibody-drug conjugate (16a) administration group compared with the control group.

Test Example 14 Antibody-drug conjugate (16a) exhibited antitumor effect in *in vivo* antitumor test using human ovarian cancer line

[0644] Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrIcrlj [(Foxn1nu/Foxn1nu)] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0645] 5×10⁶ cells of a human ovarian cancer cell line OVCAR-8 (HTB-161) from ATCC were mixed and suspended in Matrigel (BD #354234) and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0646] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm³) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0647] At Day 21 when the tumor size reached about 140 mm³, 8 animals were randomly divided into 2 groups on the

basis of their tumor sizes. At the same day, the antibody-drug conjugate (16a) or PBS for a control group were administered at the following doses.

[0648] Administration group: PBS was intravenously injected at the same single dose as the antibody-drug conjugate.

[0649] Administration group: The antibody-drug conjugate (16a) was intravenously injected at a single dose of 10 mg/kg.

[0650] The results are shown in Figure 22. The inhibition of tumor growth of the human ovarian cancer line was observed in the antibody-drug conjugate administration group compared with the control group. No weight loss was observed in the mice of the treated groups.

Test Example 15 Antibody-drug conjugate (16a) exhibited antitumor effect in *in vivo* antitumor test using human bladder cancer line

[0651] Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrIcrlj [Foxn1nu/Foxn1nu] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0652] 8×10^6 cells of a human bladder cancer cell line SW-780 (CRL-2169) from ATCC were mixed and suspended in Matrigel (BD #354234) and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0653] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm^3) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0654] At Day 7 when the tumor size reached about 190 mm^3 , 10 animals were randomly divided into 2 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (16a) or PBS for a control group were administered at the following doses.

[0655] Administration group: PBS was intravenously injected at the same single dose as the antibody-drug conjugate.

[0656] Administration group: The antibody-drug conjugate (16a) was intravenously injected at a single dose of 10 mg/kg.

[0657] The results are shown in Figure 23. The inhibition of tumor growth of the human bladder cancer line was observed in the antibody-drug conjugate administration group compared with the control group. No weight loss was observed in the mice of the treated groups.

Test Example 16 Antibody-drug conjugate (16a) exhibited HER3-dependent antitumor effect in *in vivo* antitumor test using human breast cancer line

[0658] The human breast cancer line MDA-MB-453 expresses HER3 and responds to the antibody-drug conjugate (13) as described in Test Example 8. However, as it had not been demonstrated that this pharmaceutical effect was mediated by HER3 yet, HER3 was veiled by administering U1-59 beforehand, and whether or not to reduce the pharmaceutical effect was evaluated.

[0659] Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrIcrlj [Foxn1nu/Foxn1nu] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0660] 1×10^7 cells of a human breast cancer cell line MDA-MB-453 (CLB-22) from ATCC were mixed and suspended in Matrigel (BD #354234) and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0661] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm^3) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0662] At Day 11 when the tumor size reached about 130 mm³, 16 animals were randomly divided into 4 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (16a) and/or U1-59 or PBS for a control group were administered at the following doses.

[0663] Administration group: PBS was intravenously injected at the same single dose as the antibody-drug conjugate.

[0664] Administration group: U1-59 was intravenously injected at a single dose of 30 mg/kg.

[0665] Administration group: The antibody-drug conjugate (16a) was intravenously injected at a single dose of 3 mg/kg.

[0666] Administration group: 30 minutes after administration (intravenous injection at a single dose) of U1-59, the antibody-drug conjugate (16a) was intravenously injected at a single dose of 3 mg/kg.

[0667] The results are shown in Figure 24. The inhibition of tumor growth of the human breast cancer line was observed in the antibody-drug conjugate administration group compared with the control group, whereas this tumor inhibitory effect was attenuated by administering U1-59 beforehand. These results demonstrated that the tumor inhibitory effect of the antibody-drug conjugate is a pharmaceutical effect mediated by HER3. No weight loss was observed in the mice of the treated groups.

Test Example 17 Antibody-drug conjugate (16a) exhibited antitumor effect in combined use with trastuzumab in *in vivo* antitumor test using human breast cancer line

[0668] Trastuzumab has been approved as a therapeutic agent for human HER2-positive breast cancer. However, trastuzumab resistance is known, and a mutation in PIK3CA <H1047R or H420R> has been reported to participate in one of the mechanisms underlying this resistance. In this test, whether or not the combined use of trastuzumab and the antibody-drug conjugate was effective for a trastuzumab-resistant breast cancer line was evaluated.

[0669] Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrIcrIj [Foxn1nu/Foxn1nu] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0670] 1×10^7 cells of a human breast cancer cell line MDA-MB-453 (CLB-22, H1047R mutation in PIK3CA) from ATCC were mixed and suspended in Matrigel (PBS: PAA #10010-023, Matrigel: BD #354234) and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0671] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm³) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0672] At Day 11 when the tumor size reached about 130 mm³, 16 animals were randomly divided into 4 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (16a), trastuzumab, combined use of the conjugate and trastuzumab, or PBS for a control group were administered at the following doses.

[0673] Administration group: PBS was intravenously injected at the same single dose as the antibody-drug conjugate.

[0674] Administration group: Trastuzumab (Roche Diagnostics, Inc.) was intravenously injected at a single dose of 1 mg/kg.

[0675] Administration group: The antibody-drug conjugate (16a) was intravenously injected at a single dose of 3 mg/kg.

[0676] Administration group: 30 minutes after administration (intravenous injection at a single dose) of trastuzumab (Roche Diagnostics, Inc.), the antibody-drug conjugate (16a) was intravenously injected at a single dose of 3 mg/kg.

[0677] The results are shown in Figure 25. The antitumor effect brought about by combined use on the human breast cancer line (PIK3CA H1047R) was observed in the administration of trastuzumab and the antibody-drug conjugate compared with the administration of each medicine alone. These results demonstrated that the pharmaceutical effect of the antibody-drug conjugate is potentiated by the combined use thereof with trastuzumab. No weight loss was observed in the mice of the treated groups. In other antitumor test using a human breast cancer cell line HCC1954(PIK3CA H1047R), the combined antitumor effect was also observed in the administration of trastuzumab and the antibody-drug conjugate (16a) compared with the administration of each medicine alone.

Test Example 18 Antibody-drug conjugate (15) exhibited antitumor effect in combined use with trastuzumab in *in vivo* antitumor test using human breast cancer line

[0678] Trastuzumab has been approved as a therapeutic agent for human HER2-positive breast cancer. However,

trastuzumab resistance is known, and a mutation in PIK3CA <H1047R or H420R> has been reported to participate in one of the mechanisms underlying this resistance. In this test, whether or not the combined use of trastuzumab and the antibody-drug conjugate was effective for a trastuzumab-resistant breast cancer line was evaluated.

[0679] Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrIcrlj [Foxn1nu/Foxn1nu] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0680] 5×10^6 cells of a human breast cancer cell line JIMT-1 (ACC-589, H420R mutation in PIK3CA) from ATCC were suspended in PBS (PAA #10010-023) and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0681] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm³) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0682] At Day 10 when the tumor size reached about 200 mm³, 24 animals were randomly divided into 4 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (15), trastuzumab, combined use of the conjugate and trastuzumab, or PBS for a control group were administered at the following doses.

[0683] Administration group: PBS was intravenously injected once a week at the same dose as the antibody-drug conjugate.

[0684] Administration group: Trastuzumab (Roche Diagnostics, Inc.) was intravenously injected once a week at 10 mg/kg.

[0685] Administration group: The antibody-drug conjugate (15) was intravenously injected once a week at 10 mg/kg.

[0686] Administration group: 30 minutes after administration (intravenous injection once a week) of trastuzumab (Roche Diagnostics, Inc.), the antibody-drug conjugate (15) was intravenously injected once a week at 10 mg/kg.

[0687] The results are shown in Figure 26. The antitumor effect brought about by combined use on the human breast cancer line (PIK3CA H420R) was observed in the administration of trastuzumab and the antibody-drug conjugate compared with the administration of each medicine alone. These results demonstrated that the pharmaceutical effect of the antibody-drug conjugate is potentiated by the combined use thereof with trastuzumab. No weight loss was observed in the mice of the treated groups.

Test Example 19 Antibody-drug conjugate (16a) exhibited antitumor effect in combined use with gefitinib in *in vivo* antitumor test using human lung cancer line

[0688] Gefitinib has been approved as a therapeutic agent for human lung cancer. In this test, whether or not the combined use of gefitinib and the antibody-drug conjugate was effective was evaluated.

[0689] Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrIcrlj [Foxn1nu/Foxn1nu] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0690] 3×10^6 cells of a human lung cancer cell line PC-9 (RCB0446) from ATCC were suspended in PBS (PAA #10010-023) and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0691] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm³) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0692] At Day 14 when the tumor size reached about 270 mm³, 16 animals were randomly divided into 4 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (16a), gefitinib, combined use of the conjugate and gefitinib, or PBS for a control group were administered at the following doses.

[0693] Administration group: PBS was intravenously injected once a week at the same dose as the antibody-drug

conjugate.

[0694] Administration group: Gefitinib (AstraZeneca) was orally administered once a day at 6 mg/kg.

[0695] Administration group: The antibody-drug conjugate (16a) was intravenously injected once a week at 10 mg/kg.

[0696] Administration group: 30 minutes after administration (oral administration once a day) of gefitinib (AstraZeneca), the antibody-drug conjugate (16a) was intravenously injected once a week at 10 mg/kg.

[0697] The results are shown in Figure 27. The antitumor effect brought about by combined use on the human lung cancer line was observed in the administration of gefitinib and the antibody-drug conjugate compared with the administration of each medicine alone. These results demonstrated that the pharmaceutical effect of the antibody-drug conjugate is potentiated by the combined use thereof with gefitinib. No weight loss was observed in the mice of the treated groups.

Test Example 20 Antibody-drug conjugate (16a) exhibited antitumor effect in combined use with cetuximab or panitumumab in *in vivo* antitumor test using human triple-negative breast cancer line

[0698] An anti-EGFR antibody cetuximab or panitumumab is under clinical trial against human triple-negative breast cancer. In this test, whether or not the combined use of cetuximab or panitumumab and the antibody-drug conjugate was effective was evaluated.

[0699] Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrlCrlj [Foxn1nu/Foxn1nu] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0700] 5×10^6 cells of a human triple-negative breast cancer cell line MDA-MB-468 (CRL-2322) from ATCC were suspended in a solution prepared from 200 μ L of PBS and Matrigel (PBS: PAA #10010-023, Matrigel: BD #354234) mixed at a ratio of 1:1, and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0701] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm^3) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0702] At Day 21 when the tumor size reached about 160 mm^3 , 30 animals were randomly divided into 6 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (16a), cetuximab or panitumumab, combined use of the conjugate and cetuximab or panitumumab, or PBS for a control group were administered at the following doses.

[0703] Administration group: PBS was administered at the same single dose as the antibody-drug conjugate.

[0704] Administration group: Cetuximab (Bristol-Myers Squibb Company) was administered at a single dose of 10 mg/kg.

[0705] Administration group: Panitumumab (Amgen Inc.) was administered at a single dose of 10 mg/kg.

[0706] Administration group: The antibody-drug conjugate (16a) was administered at a single dose of 10 mg/kg.

[0707] Administration group: 30 minutes after administration (administration at a single dose) of cetuximab (Bristol-Myers Squibb Company), the antibody-drug conjugate (16a) was administered at a single dose of 10 mg/kg.

[0708] Administration group: 30 minutes after administration (administration at a single dose) of panitumumab (Amgen Inc.), the antibody-drug conjugate (16a) was administered at a single dose of 10 mg/kg.

[0709] The results are shown in Figure 28. The antitumor effect brought about by combined use on the human triple-negative breast cancer line was observed in the administration of cetuximab (Figure 28A) or panitumumab (Figure 28B) and the antibody-drug conjugate compared with the administration of each medicine alone. These results demonstrated that the pharmaceutical effect of the antibody-drug conjugate is potentiated by the combined use thereof with cetuximab or panitumumab. No weight loss was observed in the mice of the treated groups. The antitumor effect brought about by combined use on the other human triple-negative breast cancer line in MDA-MB-231 was also observed in the administration of cetuximab or panitumumab and the antibody-drug conjugate (16a) compared with the administration of each medicine alone.

Test Example 21 Antibody-drug conjugate (16a) exhibited antitumor effect in combined use with cetuximab or panitumumab in *in vivo* antitumor test using human head and neck cancer line

[0710] An anti-EGFR antibody cetuximab has been approved against human head and neck cancer, while panitumu-

mab is under clinical trial against this cancer. In this test, whether or not the combined use of cetuximab or panitumumab and the antibody-drug conjugate was effective was evaluated.

[0711] Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrIcrlj [Foxn1nu/Foxn1nu] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0712] 4×10^6 cells of a human head and neck cancer cell line Fadu (HTB-43) from ATCC were suspended in 200 μ L of PBS (PAA #10010-023) and subcutaneously transplanted to the right side area of the body of each nude mouse using a 29 G needle. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0713] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm^3) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0714] At Day 6 when the tumor size reached about 330 mm^3 , 30 animals were randomly divided into 6 groups on the basis of their tumor sizes. At the same day, the antibody-drug conjugate (16a), cetuximab or panitumumab, combined use of the conjugate and cetuximab or panitumumab, or PBS for a control group were administered at the following doses.

[0715] Administration group: PBS was administered at the same single dose as the antibody-drug conjugate.

[0716] Administration group: Cetuximab (Bristol-Myers Squibb Company) was administered at a single dose of 5 mg/kg.

[0717] Administration group: Panitumumab (Amgen Inc.) was administered at a single dose of 5 mg/kg.

[0718] Administration group: The antibody-drug conjugate (16a) was administered at a single dose of 10 mg/kg.

[0719] Administration group: 30 minutes after administration (administration at a single dose) of cetuximab (Bristol-Myers Squibb Company), the antibody-drug conjugate (16a) was administered at a single dose of 10 mg/kg.

[0720] Administration group: 30 minutes after administration (administration at a single dose) of panitumumab (Amgen Inc.), the antibody-drug conjugate (16a) was administered at a single dose of 10 mg/kg.

[0721] The results are shown in Figure 29. The antitumor effect brought about by combined use on the human head and neck cancer line was observed in the administration of cetuximab (Figure 29A) or panitumumab (Figure 29B) and the antibody-drug conjugate compared with the administration of each medicine alone. These results demonstrated that the pharmaceutical effect of the antibody-drug conjugate is potentiated by the combined use thereof with cetuximab or panitumumab. No weight loss was observed in the mice of the treated groups.

Test Example 22 Antibody-drug conjugate (16a) exhibited antitumor effect in combined use with cetuximab or pertuzumab in *in vivo* antitumor test by transplantation of tumor from stomach cancer patient

[0722] An anti-EGFR antibody cetuximab and an anti-HER2 antibody pertuzumab are under clinical trial against human stomach cancer. In this test, whether or not the combined use of cetuximab or pertuzumab and the antibody-drug conjugate was effective was evaluated. The evaluation was carried out by conducting an antitumor test close to clinical condition involving the stroma of a patient using the patient-derived tumors transplanted in mice instead of the general evaluation system using a human cancer line.

[0723] Five- to 6-week-old female CAnN.Cg-Foxn1[nu]/CrIcrlj [Foxn1nu/Foxn1nu] nude mice (Charles River Laboratories Japan, Inc.) having a body weight of 15 to 20 g after acclimation were used. The mice were placed in individually ventilated cages (IVC, 4 mice at maximum per cage) which were kept at room temperature and a constant humidity. After randomization, the body weights of the mice were measured every other day, and the behaviors of the animals were recorded every day.

[0724] A stomach cancer patient-derived tumor NIBIO-G016 from National Institute of Biomedical Innovation was subcutaneously transplanted to the right side area of the body of each nude mouse. The body weights were measured using a weight scale (Mettler Toledo PB602-L).

[0725] The major axis and minor axis of the tumor were measured using an electronic digital caliper (CD-15CX, Mitsutoyo Corp), and the tumor volume (mm^3) was calculated. The calculation was carried out according to the following expression.

$$\text{Tumor volume (mm}^3\text{)} = 1/2 \times \text{Major axis (mm)} \times [\text{Minor axis (mm)}]^2$$

[0726] At Day 56 when the tumor size reached about 220 mm^3 , 30 animals were randomly divided into 6 groups on the

basis of their tumor sizes. At the same day, the antibody-drug conjugate (16a), cetuximab or pertuzumab, combined use of the conjugate and cetuximab or pertuzumab, or PBS for a control group were administered at the following doses.

[0727] Administration group: PBS was administered at the same single dose as the antibody-drug conjugate.

[0728] Administration group: Cetuximab (Bristol-Myers Squibb Company) was administered at a single dose of 10 mg/kg.

[0729] Administration group: Pertuzumab (Roche Diagnostics, Inc.) was administered at a single dose of 10 mg/kg.

[0730] Administration group: The antibody-drug conjugate (16a) was administered at a single dose of 10 mg/kg.

[0731] Administration group: 30 minutes after administration (administration at a single dose) of cetuximab (Bristol-Myers Squibb Company), the antibody-drug conjugate (16a) was administered at a single dose of 10 mg/kg.

[0732] Administration group: 30 minutes after administration (administration at a single dose) of pertuzumab (Roche Diagnostics, Inc.), the antibody-drug conjugate (16a) was administered at a single dose of 10 mg/kg.

[0733] The results are shown in Figure 30. The antitumor effect brought about by combined use on the stomach cancer patient-derived tumor model was observed in the administration of cetuximab (Figure 30A) or pertuzumab (Figure 30B) and the antibody-drug conjugate compared with the administration of each medicine alone. These results demonstrated that the pharmaceutical effect of the antibody-drug conjugate is potentiated by the combined use thereof with cetuximab or panitumumab in the cancer patient-derived tumor model. No weight loss was observed in the mice of the treated groups. Further, in an antitumor test using a human pancreatic cancer cell line BxPC3, the antibody-drug conjugate (13) exhibited a stronger antitumor effect as compared with the PBS administration group or the U1-59 administration group. The antibody-drug conjugate also exhibited a strong antitumor effect on *in vivo* antitumor model using a HER2-positive breast cancer cell line JIMT1 which had acquired resistance to combined use of trastuzumab and pertuzumab or to trastuzumab emtansine.

Test Example 23 Safety of antibody-drug conjugate for non-human animal

[0734] The anti-HER3 antibody-drug conjugate of the present invention and the pharmaceutical composition containing this anti-HER3 antibody-drug conjugate have excellent safety as a therapeutic or prophylactic agent for a disease. For example, when the antibody-drug conjugate (5), (10), or (13) was administered up to 30 mg/kg to a cross-breed rat, two times in total with an interval of once per week, no serious toxicity findings were observed from any of the antibody-drug conjugates as a result of observation until Day 7 after the final administration. Further, when the antibody-drug conjugate (5), (10), or (13) was administered up to 30 mg/kg to a cross-breed monkey, no remarkable toxicity findings were observed from any of the antibody-drug conjugates as a result of observation for 7 days.

[0735] Further, when antibody-drug conjugate (5), (10), or (13) was administered at a plurality of doses to a monkey (3-week intervals), no remarkable toxicity findings were observed from any of the antibody-drug conjugates as a result of observation. Accordingly, the antibody-drug conjugate of the present invention has excellent safety as a pharmaceutical composition for treatment or prevention of a disease.

[Sequence Listing Free Text]

[0736]

SEQ ID NO: 1 - Nucleotide sequence encoding a heavy chain variable region amino acid sequence of the anti-HER3 human antibody U1-39

SEQ ID NO: 2 - Heavy chain variable region amino acid sequence of the anti-HER3 human antibody U1-39

SEQ ID NO: 3 - Nucleotide sequence encoding a light chain variable region amino acid sequence of the anti-HER3 human antibody U1-39

SEQ ID NO: 4 - Light chain variable region amino acid sequence of the anti-HER3 human antibody U1-39

SEQ ID NO: 5 - Nucleotide sequence encoding a heavy chain variable region amino acid sequence of the anti-HER3 human antibody U1-40

SEQ ID NO: 6 - Heavy chain variable region amino acid sequence of the anti-HER3 human antibody U1-40

SEQ ID NO: 7 - Nucleotide sequence encoding a light chain variable region amino acid sequence of the anti-HER3 human antibody U1-40

SEQ ID NO: 8 - Light chain variable region amino acid sequence of the anti-HER3 human antibody U1-40

SEQ ID NO: 9 - Nucleotide sequence encoding a heavy chain variable region amino acid sequence of the anti-HER3 human antibody U1-38

SEQ ID NO: 10 - Heavy chain variable region amino acid sequence of the anti-HER3 human antibody U1-38

SEQ ID NO: 11 - Nucleotide sequence encoding a light chain variable region amino acid sequence of the anti-HER3 human antibody U1-38

SEQ ID NO: 12 - Light chain variable region amino acid sequence of the anti-HER3 human antibody U1-38

SEQ ID NO: 13 - Nucleotide sequence encoding a heavy chain variable region amino acid sequence of the anti-HER3

EP 4 494 710 A2

[illegible]

[illegible]

[illegible]

[illegible]

5

10

15

20

25

30

35

40

45

50

55

[illegible]


55

SEQ ID NO: 542 - Light chain CDRL2 amino acid sequence of the anti-HER3 human antibody U1-53
 SEQ ID NO: 543 - Light chain CDRL3 amino acid sequence of the anti-HER3 human antibody U1-53
 SEQ ID NO: 544 - Heavy chain CDRH1 amino acid sequence of the anti-HER3 human antibody U1-55.1
 SEQ ID NO: 545 - Heavy chain CDRH2 amino acid sequence of the anti-HER3 human antibody U1-55.1
 5 SEQ ID NO: 546 - Heavy chain CDRH3 amino acid sequence of the anti-HER3 human antibody U1-55.1
 SEQ ID NO: 547 - Light chain CDRL1 amino acid sequence of the anti-HER3 human antibody U1-55
 SEQ ID NO: 548 - Light chain CDRL2 amino acid sequence of the anti-HER3 human antibody U1-55
 SEQ ID NO: 549 - Light chain CDRL3 amino acid sequence of the anti-HER3 human antibody U1-55
 SEQ ID NO: 550 - Light chain CDRL1 amino acid sequence of the anti-HER3 human antibody U1-57.1
 10 SEQ ID NO: 551 - Light chain CDRL2 amino acid sequence of the anti-HER3 human antibody U1-57.1
 SEQ ID NO: 552 - Light chain CDRL3 amino acid sequence of the anti-HER3 human antibody U1-57.1
 SEQ ID NO: 553 - Heavy chain CDRH1 amino acid sequence of the anti-HER3 human antibody U1-57
 SEQ ID NO: 554 - Heavy chain CDRH2 amino acid sequence of the anti-HER3 human antibody U1-57
 SEQ ID NO: 555 - Heavy chain CDRH3 amino acid sequence of the anti-HER3 human antibody U1-57
 15 SEQ ID NO: 556 - Heavy chain CDRH1 amino acid sequence of the anti-HER3 human antibody U1-58
 SEQ ID NO: 557 - Heavy chain CDRH2 amino acid sequence of the anti-HER3 human antibody U1-58
 SEQ ID NO: 558 - Heavy chain CDRH3 amino acid sequence of the anti-HER3 human antibody U1-58
 SEQ ID NO: 559 - Light chain CDRL1 amino acid sequence of the anti-HER3 human antibody U1-58
 SEQ ID NO: 560 - Light chain CDRL2 amino acid sequence of the anti-HER3 human antibody U1-58
 20 SEQ ID NO: 561 - Light chain CDRL3 amino acid sequence of the anti-HER3 human antibody U1-58
 SEQ ID NO: 562 - Heavy chain CDRH1 amino acid sequence of the anti-HER3 human antibody U1-59
 SEQ ID NO: 563 - Heavy chain CDRH2 amino acid sequence of the anti-HER3 human antibody U1-59
 SEQ ID NO: 564 - Heavy chain CDRH3 amino acid sequence of the anti-HER3 human antibody U1-59
 SEQ ID NO: 565 - Light chain CDRL1 amino acid sequence of the anti-HER3 human antibody U1-59
 25 SEQ ID NO: 566 - Light chain CDRL2 amino acid sequence of the anti-HER3 human antibody U1-59
 SEQ ID NO: 567 - Light chain CDRL3 amino acid sequence of the anti-HER3 human antibody U1-59
 SEQ ID NO: 568 - Heavy chain CDRH1 amino acid sequence of the anti-HER3 human antibody U1-61.1
 SEQ ID NO: 569 - Heavy chain CDRH2 amino acid sequence of the anti-HER3 human antibody U1-61.1
 SEQ ID NO: 570 - Heavy chain CDRH3 amino acid sequence of the anti-HER3 human antibody U1-61.1
 30 SEQ ID NO: 571 - Light chain CDRL1 amino acid sequence of the anti-HER3 human antibody U1-61.1
 SEQ ID NO: 572 - Light chain CDRL2 amino acid sequence of the anti-HER3 human antibody U1-61.1
 SEQ ID NO: 573 - Light chain CDRL3 amino acid sequence of the anti-HER3 human antibody U1-61.1
 SEQ ID NO: 574 - Heavy chain CDRH1 amino acid sequence of the anti-HER3 human antibody U1-61
 SEQ ID NO: 575 - Heavy chain CDRH2 amino acid sequence of the anti-HER3 human antibody U1-61
 35 SEQ ID NO: 576 - Heavy chain CDRH3 amino acid sequence of the anti-HER3 human antibody U1-61
 SEQ ID NO: 577 - Heavy chain CDRH1 amino acid sequence of the anti-HER3 human antibody U1-62
 SEQ ID NO: 578 - Heavy chain CDRH2 amino acid sequence of the anti-HER3 human antibody U1-62
 SEQ ID NO: 579 - Heavy chain CDRH3 amino acid sequence of the anti-HER3 human antibody U1-62
 SEQ ID NO: 580 - Light chain CDRL1 amino acid sequence of the anti-HER3 human antibody U1-62
 40 SEQ ID NO: 581 - Light chain CDRL2 amino acid sequence of the anti-HER3 human antibody U1-62
 SEQ ID NO: 582 - Light chain CDRL3 amino acid sequence of the anti-HER3 human antibody U1-62
 SEQ ID NO: 583 - Full-length amino acid sequence of a heavy chain of anti-HER3 human antibody U1-59
 SEQ ID NO: 584 - Full-length amino acid sequence of a light chain of anti-HER3 human antibody U1-59


Claims

1. A pharmaceutical composition for use in the treatment of triple-negative breast cancer and/or luminal breast cancer comprising an antibody-drug conjugate, a salt thereof or a hydrate thereof as an active ingredient, and a pharmaceutically acceptable formulation ingredient,


wherein the antibody-drug conjugate is an antibody-drug conjugate wherein an antitumor compound represented by the following formula:



is conjugated to an anti-HER3 antibody by a thioether bond which is formed at a disulfide bond moiety present in a hinge part of the anti-HER3 antibody via a linker having a structure represented by the following formula:



wherein, the anti-HER3 antibody is connected to the terminal of L^1 ,
the antitumor compound is connected to the carbonyl group of $-(CH_2)_{n^2}-C(=O)-$ moiety or the C terminal of L^P , with
the nitrogen atom of the amino group at position 1 as a connecting position,
wherein, n^1 represents an integer of 0 to 6,
 n^2 represents an integer of 0 to 5,
 L^1 represents $-(\text{Succinimid-3-yl-N})-(CH_2)_{n^3}-C(=O)-$,
wherein n^3 represents an integer of 2 to 8,
 L^2 represents $-NH-(CH_2CH_2-O)_{n^4}-CH_2CH_2-C(=O)-$ or a single bond,
wherein n^4 represents an integer of 1 to 6,
 L^P represents a tetrapeptide residue $-GGFG-$ or pentapeptide residue $-DGGFG-$,
 L^a represents $-O-$ or a single bond,
 $-(\text{Succinimid-3-yl-N})-$ has a structure represented by the following formula:




which is connected to the anti-HER3 antibody at position 3 thereof and is connected to a methylene group in the linker structure containing this structure on the nitrogen atom at position 1.

2. The pharmaceutical composition according to claim 1, wherein the drug-linker structure moiety in which a drug is connected to $-L^1-L^2-L^P-NH-(CH_2)_{n^1}-L^a-(CH_2)_{n^2}-C(=O)-$ or $-L^1-L^2-L^P-$ is one drug-linker structure selected from the following group:

- (Succinimid-3-yl-N)- $CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-DGGFG-NH-CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-DGGFG-NH-CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-DGGFG-NH-CH_2CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2-O-CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2-O-CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-NH-CH_2CH_2-O-CH_2CH_2-O-CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-NH-CH_2CH_2-O-CH_2CH_2-O-CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-NH-CH_2CH_2-O-CH_2CH_2-O-CH_2CH_2-O-CH_2CH_2-O-CH_2CH_2-C(=O)-GGFG-NH-CH_2CH_2-C(=O)-(NH-DX)$,
- (Succinimid-3-yl-N)- $CH_2CH_2CH_2CH_2CH_2-C(=O)-NH-CH_2CH_2-O-CH_2CH_2-O-CH_2CH_2-O-CH_2CH_2-O-$


$\text{CH}_2\text{CH}_2\text{-C(=O)-GGFG-NH-CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-(NH-DX)}$,
 - (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{-C(=O)-GGFG-(NH-DX)}$,
 - (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{-C(=O)-DGGFG-(NH-DX)}$,
 - (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-GGFG-(NH-DX)}$,
 - (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-DGGFG-(NH-DX)}$;

wherein, -(Succinimid-3-yl-N)- has a structure represented by the following formula:



which is connected to the anti-HER3 antibody at position 3 thereof and is connected to a methylene group in the linker structure containing this structure on the nitrogen atom at position 1,

-(NH-DX) represents a group represented by the following formula, with the nitrogen atom of the amino group at position 1 being a connecting position:



-GGFG- represents a tetrapeptide residue of -Gly-Gly-Phe-Gly- and -DGGFG- represents a pentapeptide residue of -Asp-Gly-Gly-Phe-Gly-.

3. The pharmaceutical composition according to claim 2, wherein a drug-linker structure moiety is one drug-linker structure selected from the following group:

- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-GGFG-NH-CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-(NH-DX)}$,
 - (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-DGGFG-NH-CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-(NH-DX)}$,
 - (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-GGFG-NH-CH}_2\text{-O-CH}_2\text{-C(=O)-(NH-DX)}$,
 - (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{-C(=O)-NH-CH}_2\text{CH}_2\text{O-CH}_2\text{CH}_2\text{O-CH}_2\text{CH}_2\text{-C(=O)-GGFG-NH-CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-(NH-DX)}$,
 - (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-DGGFG-(NH-DX)}$.

4. The pharmaceutical composition according to claim 3, wherein the drug-linker structure is:

-(Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-GGFG-NH-CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-(NH-DX)}$.

5. The pharmaceutical composition according to claim 3, wherein the drug-linker structure is:

-(Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-DGGFG-NH-CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-(NH-DX)}$.


6. The pharmaceutical composition according to claim 3, wherein the drug-linker structure is:

-(Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-GGFG-NH-CH}_2\text{-O-CH}_2\text{-C(=O)-(NH-DX)}$.

7. The pharmaceutical composition according to claim 3, wherein the drug-linker structure is:

- (Succinimid-3-yl-N)- $\text{CH}_2\text{CH}_2\text{-C(=O)-NH-CH}_2\text{CH}_2\text{O-CH}_2\text{CH}_2\text{O-CH}_2\text{CH}_2\text{-C(=O)-GGFG-NH-CH}_2\text{CH}_2\text{CH}_2\text{-C(=O)-(NH-DX)}$.

8. The pharmaceutical composition according to claim 3, wherein the drug-linker structure is:



9. The pharmaceutical composition according to any one of claims 1 to 8, wherein the anti-HER3 antibody comprises the CDRH1 to CDRH3 and CDRL1 to CDRL3 of U1-49, U1-53, U1-59, U1-7 or U1-9 in the heavy and light chains, respectively.

10. The pharmaceutical composition according to any one of claims 1 to 8, wherein the anti-HER3 antibody comprises the heavy chain variable region and the light chain variable chain of U1-49, U1-53, U1-59, U1-7 or U1-9 on the heavy and light chains, respectively.

11. The pharmaceutical composition according to any one of claims 1 to 8, wherein the anti-HER3 antibody comprises the amino acid sequences represented by or comprising SEQ ID No: 42 and 44, SEQ ID No: 54 and 56, SEQ ID No: 70 and 72, SEQ ID No: 92 and 94, or, SEQ ID No: 96 and 98, in the heavy and light chains, respectively.

12. The pharmaceutical composition according to any one of claims 1 to 8, wherein the anti-HER3 antibody comprises the amino acid sequences represented by or comprising SEQ ID No: 583 and 584 in the heavy and light chains, respectively.

13. The pharmaceutical composition according to claim 12, wherein the anti-HER3 antibody lacks a lysine residue at the carboxyl terminus of the heavy chain.

14. The pharmaceutical composition according to any one of claims 1 to 13, wherein the average number of units of the selected one drug-linker structure conjugated per antibody is in a range of from 2 to 8.

15. The pharmaceutical composition according to any one of claims 1 to 13, wherein the average number of units of the selected one drug-linker structure conjugated per antibody is in a range of from 3 to 8.

FIG.1

Full-length amino acid sequence of heavy chain of anti-HER3
human antibody U1-59 (SEQ ID NO: 583)

Q V Q L Q Q W G A G L L K P S E T L S L T C A V Y G G S F S G Y Y W S W
I R Q P P G K G L E W I G E I N H S G S T N Y N P S L K S R V T I S V E
T S K N Q F S L K L S S V T A A D T A V Y Y C A R D K W T W Y F D L W G
R G T L V T V S S A S T K G P S V F P L A P S S K S T S G G T A A L G C
L V K D Y F P E P V T V S W N S G A L T S G V H T F P A V L Q S S G L Y
S L S S V V T V P S S S L G T Q T Y I C N V N H K P S N T K V D K R V E
P K S C D K T H T C P P C P A P E L L G G P S V F L F P P K P K D T L M
I S R T P E V T C V V V D V S H E D P E V K F N W Y V D G V E V H N A K
T K P R E E Q Y N S T Y R V V S V L T V L H Q D W L N G K E Y K C K V S
N K A L P A P I E K T I S K A K G Q P R E P Q V Y T L P P S R E E M T K
N Q V S L T C L V K G F Y P S D I A V E W E S N G Q P E N N Y K T T P P
V L D S D G S F F L Y S K L T V D K S R W Q Q G N V F S C S V M H E A L
H N H Y T Q K S L S L S P G K

FIG.2

Full-length amino acid sequence of light chain of anti-HER3 human antibody U1-59 (SEQ ID NO: 584)

```

D I E M T Q S P D S L A V S L G E R A T I N C R S S Q S V L Y S S S N R
N Y L A W Y Q Q N P G Q P P K L L I Y W A S T R E S G V P D R F S G S G
S G T D F T L T I S S L Q A E D V A V Y Y C Q Q Y Y S T P R T F G Q G T
K V E I K R T V A A P S V F I F P P S D E Q L K S G T A S V V C L L N N
F Y P R E A K V Q W K V D N A L Q S G N S Q E S V T E Q D S K D S T Y S
L S S T L T L S K A D Y E K H K V Y A C E V T H Q G L S S P V T K S F N
R G E C

```

FIG.3




FIG.4




FIG.5





FIG.6

A.



B.




FIG. 7




FIG. 8





FIG.9

A.



B.





FIG.10

A.



B.




FIG.11




FIG.12




FIG.13




FIG.14




FIG.15




FIG.16




FIG.17




FIG.18




FIG.19




FIG.20




FIG.21




FIG.22




FIG.23




FIG.24




FIG.25




FIG.26




FIG.27




FIG.28





FIG.29

(A)



(B)






FIG.30

(A)



(B)



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- JP 5059061 A [0009]
- JP 8337584 A [0009]
- WO 199746260 A [0009] [0377] [0456]
- WO 200025825 A [0009]
- US 5968511 A [0009]
- US 5480968 A [0009]
- WO 2003013602 A [0009]
- WO 2007077028 A [0009] [0060] [0169] [0431]
- WO 2008100624 A [0009]
- WO 2012019024 A [0009]
- WO 9007861 A [0161]
- WO 9201047 A [0173]
- WO 9220791 A [0173]
- WO 9306213 A [0173]
- WO 9311236 A [0173]
- WO 9319172 A [0173]
- WO 9501438 A [0173]
- WO 9515388 A [0173]
- WO 199954342 A [0190]
- WO 200061739 A [0190]
- WO 200231140 A [0190]
- US 4816397 A [0198] [0199]
- US 5916771 A [0198] [0199]
- US 6207418 B [0198] [0199]
- JP 6087746 A [0209]
- JP 2002060351 A [0311] [0490]
- WO 2003038043 A [0423]

Non-patent literature cited in the description

- DUCRY, L. et al. *Bioconjugate Chem.*, 2010, vol. 21, 5-13 [0010]
- ALLEY, S. C. et al. *Current Opinion in Chemical Biology*, 2010, vol. 14, 529-537 [0010]
- DAMLE N.K. *Expert Opin. Biol. Ther.*, 2004, vol. 4, 1445-1452 [0010]
- SENTER P. D. et al. *Nature Biotechnology*, 2012, vol. 30, 631-637 [0010]
- KUMAZAWA, E. ; TOHGO, A. *Exp. Opin. Invest. Drugs*, 1998, vol. 7, 625-632 [0010]
- MITSUI, I. et al. *Jpn J. Cancer Res.*, 1995, vol. 86, 776-786 [0010]
- TAKIGUCHI, S. et al. *Jpn J. Cancer Res.*, 1997, vol. 88, 760-769 [0010]
- JOTO, N. et al. *Int J Cancer*, 1997, vol. 72, 680-686 [0010]
- KUMAZAWA, E. et al. *Cancer Chemother. Pharmacol.*, 1998, vol. 42, 210-220 [0010]
- DE JAGER, R. et al. *Ann N Y Acad Sci*, 2000, vol. 922, 260-273 [0010]
- INOUE, K. et al. *Polymer Drugs in the Clinical Stage*, 2003, 145-153 [0010]
- KUMAZAWA, E. et al. *Cancer Sci*, 2004, vol. 95, 168-175 [0010]
- SOEPENBERG, O. et al. *Clinical Cancer Research*, 2005, vol. 11, 703-711 [0010]
- WENTE M. N. et al. *Investigational New Drugs*, 2005, vol. 23, 339-347 [0010]
- PLOWMAN et al. *Proc. Natl. Acad. Sci. U.S.A.*, 1990, vol. 87, 4905-4909 [0010]
- KRAUS et al. *Proc. Natl. Acad. Sci. U.S.A.*, 1989, vol. 86, 9193-9197 [0010]
- KRAUS et al. *Proc. Natl. Acad. Sci. U.S.A.*, 1993, vol. 90, 2900-2904 [0010]
- ALIMANDI et al. *Oncogene*, 1995, vol. 10, 1813-1821 [0010]
- DEFAZIO et al. *Int. J. Cancer*, 2000, vol. 87, 487-498 [0010]
- NADIU et al. *Br. J. Cancer*, 1998, vol. 78, 1385-1390 [0010]
- BINZ ; PLUCKTHUN. *Curr Opin Biotechnol*, vol. 16, 459-69 [0059]
- SKERRA. *J. Mol. Recog.*, 2000 [0059]
- JONES et al. *Nature*, 1986, vol. 321, 522-525 [0060]
- RIECHMANN et al. *Nature*, 1988, vol. 332, 323-329 [0060]
- PRESTA. *Curr. Op. Struct. Biol.*, 1992, vol. 2, 593-596 [0060]
- MORRISON et al. *Proc. Natl. Acad. Sci. U.S.A.*, 1984, vol. 81, 6851-6855 [0060]
- TOMIZUKA, K. et al. *Nature Genetics*, 1997, vol. 16, 133-143 [0060] [0165]
- KUROIWA, Y. et al. *Nucl. Acids Res.*, 1998, vol. 26, 3447-3448 [0060] [0165]
- YOSHIDA, H. et al. *Animal Cell Technology: Basic and Applied Aspects*. Kluwer Academic Publishers, 1999, vol. 10, 69-73 [0060] [0165]
- TOMIZUKA, K. et al. *Proc. Natl. Acad. Sci. USA*, 2000, vol. 97, 722-727 [0060] [0165]
- HOLLINGER et al. *Proc. Natl. Acad. Sci. U.S.A.*, 1993, vol. 90, 6444-6448 [0060]

- **PLUCKTHUN.** The Pharmacology of Monoclonal Antibodies. Springer Verlag, 1994, vol. 113, 269-315 [0060]
- **MUYLDERMANS et al.** *Reviews in Molecular Biotechnology*, vol. 74, 277-302 [0061]
- **HOLT et al.** *Trends Biotechnol.*, vol. 21, 484-90 [0061]
- *Cell Death and Differentiation*, 2008, vol. 15, 751-761 [0067]
- *Molecular Biology of the Cell*, vol. 15, 5268-5282 [0067]
- *Bio Techniques*, January 2000, vol. 28, 162-165 [0067]
- **KOHLER ; MILSTEIN.** *Nature*, 1975, vol. 256, 495-497 [0071] [0089]
- Monoclonal Antibodies. Plenum Press, 1980, 365-367 [0071] [0089]
- **SAIKI, R. K. et al.** *Science*, 1988, vol. 239, 487-489 [0097]
- **GLUZMAN, Y.** *Cell*, 1981, vol. 23, 175-182 [0100] [0192]
- **URLAUB, G. ; CHASIN, L. A.** *Proc. Natl. Acad. Sci. USA*, 1980, vol. 77, 4126-4220 [0100] [0192]
- **WEIR, D. M.** Handbook of Experimental Immunology. Blackwell Scientific Publications, 1987, vol. I. II. III [0120] [0135]
- **KABAT, E. A. ; MAYER, M. M.** Experimental Immunochemistry. Charles C Thomas Publisher, 1964 [0120] [0135]
- **KOHLER et al.** *Nature*, 1975, vol. 256, 495 [0131] [0138]
- **KOHLER et al.** *Eur. J. Immunol.*, 1977, vol. 6, 511 [0131]
- **MILSTEIN et al.** *Nature*, 1977, vol. 266, 550 [0131] [0138]
- **WALSH.** *Nature*, 1977, vol. 266, 495 [0131]
- **BARBARA, B. M. ; STANLEY, M. S.** Selected Methods in Cellular Immunology. W. H. Freeman and Company, 1980 [0140]
- **WEIR, D. M.** Handbook of Experimental Immunology. Blackwell Scientific Publications, 1978, vol. I, II, III [0150]
- *Proc. Natl. Acad. Sci. USA*, 1984, vol. 81, 6851-6855 [0160]
- *Nature*, 1986, vol. 321, 522-525 [0161]
- Immunology - A Synthesis. Sinauer Associates, 1991 [0163]
- **WORMSTONE, I. M. et al.** *Investigative Ophthalmology & Visual Science*, 2002, vol. 43 (7), 2301-2308 [0170]
- **CARMEN, S. et al.** *Briefings in Functional Genomics and Proteomics*, 2002, vol. 1 (2), 189-203 [0170]
- **SIRIWARDENA, D. et al.** *Ophthalmology*, 2002, vol. 109 (3), 427-431 [0170]
- *Nature Biotechnology*, 2005, vol. 23 (9), 1105-1116 [0171] [0173]
- *Annu. Rev. Immunol.*, 1994, vol. 12, 433-455 [0173]
- **BOWIE et al.** *Science*, 1991, vol. 253, 164 [0184]
- Proteins, Structures and Molecular Principles. W. H. Freeman and Company, 1984 [0184]
- Introduction to Protein Structure. Garland Publishing, 1991 [0184]
- **THORNTON.** *Nature*, 1991, vol. 354, 105 [0184]
- **ALTSCHUL, STEPHEN F. ; THOMAS L. MADDEN ; ALEJANDRO A. SCHAEFFER ; JINGHUI ZHANG ; ZHENG ZHANG ; WEBB MILLER ; DAVID J. LIPMAN.** Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.*, 1997, vol. 25, 3389-3402 [0185]
- **LORI BURTON.** *Pharmaceutical Development and Technology*, 2007, vol. 12, 265-273 [0188]
- *Journal of Chromatography A*, 1995, vol. 705, 129-134 [0195]
- *Analytical Biochemistry*, 2007, vol. 360, 75-83 [0195]
- **IDUSOGIE et al.** *J Immunol.*, vol. 166, 2571-2575 [0198]
- **SHIELDS et al.** *J Biol Chem.*, vol. 276, 6591-6604 [0199]
- Strategies for Protein Purification and Characterization: A Laboratory Course Manual. Cold Spring Harbor Laboratory Press, 1996 [0200]
- Antibodies: A Laboratory Manual. Cold Spring Harbor Laboratory, 1988 [0200]
- **HERMANSON, G.T.** Bioconjugate Techniques. Academic Press, 1996, 56-136 [0249]
- *Protein Science*, 1995, vol. 4, 2411-2423 [0256] [0262] [0269]
- *J. Org. Chem.*, 1986, vol. 51, 3196 [0311]