



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/04/24
(87) Date publication PCT/PCT Publication Date: 2018/11/01
(85) Entrée phase nationale/National Entry: 2019/10/17
(86) N° demande PCT/PCT Application No.: EP 2018/060477
(87) N° publication PCT/PCT Publication No.: 2018/197492
(30) Priorité/Priority: 2017/04/24 (US62/489,226)

(51) Cl.Int./Int.Cl. *C07K 14/725* (2006.01),
C07K 14/47 (2006.01)
(71) Demandeurs/Applicants:
OSPEDALE SAN RAFFAELE S.R.L., IT;
FONDAZIONE CENTRO SAN RAFFAELE, IT
(72) Inventeurs/Inventors:
BONINI, MARIA CHIARA, IT;
RUGGIERO, ELIANA, IT;
MAGNANI, ZULMA IRENE, IT;
VAGO, LUCA ALDO EDOARDO, IT;
BONDANZA, ATTILIO, IT;
CICERI, FABIO, IT
(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : TCR ET PEPTIDES
(54) Title: TCR AND PEPTIDES

(57) Abrégé/Abstract:
TCR AND PEPTIDES A T-cell receptor (TCR), which binds to a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
01 November 2018 (01.11.2018)



(10) International Publication Number
WO 2018/197492 A1

- (51) International Patent Classification:
C07K 14/725 (2006.01) C07K 14/47 (2006.01)
- (21) International Application Number:
PCT/EP2018/060477
- (22) International Filing Date:
24 April 2018 (24.04.2018)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
62/489,226 24 April 2017 (24.04.2017) US
- (71) Applicants: OSPEDALE SAN RAFFAELE S.R.L. [IT/IT]; Via Olgettina 60, 20132, Milan (IT). FON-DAZIONE CENTRO SAN RAFFAELE [IT/IT]; Via Olgettina 60, 20132, Milan (IT).
- (72) Inventors: BONINI, Maria Chiara; Via Olgettina 58, 20132 Milan (IT). RUGGIERO, Eliana; Via Olgettina 58, 20132 Milan (IT). MAGNANI, Zulma Irene; Via Olgettina 58, 20132 Milan (IT). VAGO, Luca Aldo Edoardo; Via Olgettina 58, 20132 Milan (IT). BONDANZA, Attilio; Via Olgettina 58, 20132 Milan (IT). CICERI, Fabio; Via Olgettina 58, 20132 Milan (IT).
- (74) Agent: D YOUNG & CO LLP; 120 Holborn, London EC1N 2DY (GB).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: TCR AND PEPTIDES

(57) Abstract: TCR AND PEPTIDES A T-cell receptor (TCR), which binds to a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC).

WO 2018/197492 A1

TCR AND PEPTIDES

FIELD OF THE INVENTION

The present invention relates to T-cell receptors (TCRs) which bind to peptides derived from Wilms tumour 1 protein (WT1) when presented by a major histocompatibility complex. In this regard, the present invention relates to complementarity determining regions (CDRs) which specifically recognise WT1 peptides. The present invention further relates to immunogenic peptides derived from WT1.

BACKGROUND TO THE INVENTION

T cell receptor (TCR) gene therapy is based on the genetic transfer of high-avidity tumour-specific TCR genes into T lymphocytes, thus enabling the specific targeting of the desired tumour-associated antigens and leading to a less toxic and more specific and effective therapy. This approach has shown promise in clinical trials. One of the main barriers limiting the exploitation of TCR gene therapy for clinical treatment of cancers is the lack of tumour-specific T-cells and corresponding TCRs. Thus, the low availability of tumour-specific TCRs still remains an open issue limiting the broad exploitation of TCR-based immunotherapeutic approaches.

The majority of tumour-associated antigens (TAAs) are self antigens, thus T-cells specific for such molecules are either destroyed or anergized due to central and peripheral tolerance. Despite this, naturally occurring tumour-specific T-cells have been observed in healthy donors and patients, particularly in patients affected by hematological malignancies, after allogeneic hematopoietic stem cell transplantation (allo-HSCT) where frequencies of tumor-specific lymphocytes have been correlated with disease regression (Kapp, M. et al. Bone Marrow Transplantation 43,399–410 (2009); and Tyler, E.M. et al. Blood 121,308–317 (2013)).

The choice of a tumor antigen to be targeted by immunotherapeutic approaches is still a matter of debate. Ideal TAAs are highly expressed on tumor cells while being minimally expressed in healthy tissue.

Wilms tumor 1 (WT1) is an intracellular protein encoding a zinc finger transcription factor that plays an important role in cell growth and differentiation (Yang, L. et al. Leukemia 21, 868-876 (2007)). WT1 is widely expressed on a variety of hematological and solid tumors, while showing limited expression on various healthy tissues (e.g. gonads, uterus, kidney, mesothelium, progenitor cells in different tissues). Recent evidence suggests a role for WT1 in leukemogenesis and tumorigenesis.

Several ongoing clinical trials rely on the generation of cytotoxic T lymphocyte (CTL) responses upon vaccination with WT1 peptides. However, despite the recognition that WT1 is useful for immunotherapy, a small number of WT1 epitopes, which are restricted to a limited number of HLA alleles, are presently used for vaccination purposes (Di Stasi, A. et al. Front. Immunol. (2015)). One such epitope is the WT1 126-134 epitope (RMFPNAPYL; SEQ ID NO: 255), which is presented by MHC encoded by the HLA-A*0201 allele (i.e. the epitope is HLA-A*0201 restricted).

HLA-A*0201 restricted epitopes and corresponding TCRs are of interest since major histocompatibility complex (MHC) having the HLA-A*0201 haplotype are expressed in the vast majority (60%) of the Caucasian population. Accordingly, TCRs that target HLA-A*0201-restricted WT1 epitopes are particularly advantageous since an immunotherapy making use of such TCRs may be widely applied.

The WT1 126-134 epitope has been widely studied in several trials, alone or in combination with additional tumor antigens. However, recent reports have highlighted a major concern regarding the processing of this particular epitope, which may impair its use for immunotherapy purposes. Notably, the WT1 126-134 epitope is more efficiently processed by the immunoproteasome compared with standard proteasomes (Jaigirdar, A. et al. J Immunother. 39(3):105-16 (2016)), which leads to poor recognition of many HLA-A*0201 tumour cell lines or primary leukemia cells that endogenously express WT1.

Thus, there remains a need for new WT1 epitopes, particularly those presented by MHC with prevalent HLA haplotypes (e.g. HLA-A*0201).

One naturally processed HLA-A*0201 restricted epitope that has been identified is WT1 37-45, which has the amino acid sequence VLDFAPPGA (SEQ ID NO: 157, see e.g. Smithgall et al 2001; Blood 98 (11 Part 1): 121a). However, few TCR amino acid sequences, particularly CDR sequences, specific for this peptide sequence have been reported (Schmitt, T.M. et al. (2017) Nat Biotechnol 35: 1188-1195).

Accordingly, there remains a need for new WT1 epitopes, particularly those restricted to common HLA alleles and a need for new TCRs capable of binding to WT1 epitopes.

SUMMARY OF THE INVENTION

We have identified novel TCRs that bind to WT1 peptides when presented by an MHC. Further, we have determined the amino acid sequences of the TCRs, including the amino acid sequences of their CDR regions, which are responsible for binding specificity for WT1. Moreover, we have demonstrated that T-cells expressing TCRs according to the present

invention specifically target and kill cells that overexpress the WT1 protein. In addition, it has been shown that the TCRs of the present invention are restricted to MHC encoded by HLA class 1 and 2 alleles common in the Caucasian population, such as HLA-A*0201 and HLA-B*3501 or HLA-B*3502.

5 Accordingly, in a first aspect, the present invention provides a T-cell receptor (TCR), which binds to a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR:

- 10 (i) comprises a CDR3 α comprising the amino acid sequence of CGTAWINDYKLSF (SEQ ID NO: 3) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRKTGGYSNQPQHF (SEQ ID NO: 8) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- 15 (ii) comprises a CDR3 α comprising the amino acid sequence of CVVNLLSNQGGKLIF (SEQ ID NO: 36) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSQDYLVSNKLFF (SEQ ID NO: 41) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- 20 (iii) comprises a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- 25 (iv) comprises a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- 30 (v) comprises a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions;

- (vi) comprises a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (vii) comprises a CDR3 α comprising the amino acid sequence of CATDAYSGNTPLVF (SEQ ID NO: 47) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRAAGLDTEAFF (SEQ ID NO: 57) or a variant thereof having up to three amino acid substitutions, additions or deletions; .
- (viii) comprises a CDR3 α comprising the amino acid sequence of CATDAYSGNTPLVF (SEQ ID NO: 47) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASTQTPYEQYF (SEQ ID NO: 63) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (ix) comprises a CDR3 α comprising the amino acid sequence of CATDAYSGNTPLVF (SEQ ID NO: 47) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSTVGGEDYGYTF (SEQ ID NO: 69) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (x) comprises a CDR3 α comprising the amino acid sequence of CAVRAEIYNQGGKLIF (SEQ ID NO: 52) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRAAGLDTEAFF (SEQ ID NO: 57) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xi) comprises a CDR3 α comprising the amino acid sequence of CAVRAEIYNQGGKLIF (SEQ ID NO: 52) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASTQTPYEQYF (SEQ ID NO: 63) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xii) comprises a CDR3 α comprising the amino acid sequence of CAVRAEIYNQGGKLIF (SEQ ID NO: 52) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of

CASSTVGGEDYGYTF (SEQ ID NO: 69) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xiii) comprises a CDR3 α comprising the amino acid sequence of CAASMAGAGSYQLTF (SEQ ID NO: 75) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CAISVGQGALYEQYF (SEQ ID NO: 80) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xiv) comprises a CDR3 α comprising the amino acid sequence of CAASMAGAGSYQLTF (SEQ ID NO: 75) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSVARDRRNYGYTF (SEQ ID NO: 86) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xv) comprises a CDR3 α comprising the amino acid sequence of CAANNARLMF (SEQ ID NO: 92) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSDTRAREQFF (SEQ ID NO: 97) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xvi) comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xvii) comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSVGGSGSYNEQFF (SEQ ID NO: 169) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xviii) comprises a CDR3 α comprising the amino acid sequence of CAVEATDSWGKLQF (SEQ ID NO: 108) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions;

- (xix) comprises a CDR3 α comprising the amino acid sequence of CAVEATDSWGKLQF (SEQ ID NO: 108) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSVGGSGSYNEQFF (SEQ ID NO: 169) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xx) comprises a CDR3 α comprising the amino acid sequence of CAVRTSYDKVIF (SEQ ID NO: 113) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxi) comprises a CDR3 α comprising the amino acid sequence of CAVRTSYDKVIF (SEQ ID NO: 113) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSVGGSGSYNEQFF (SEQ ID NO: 169) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxii) comprises a CDR3 α comprising the amino acid sequence of CAVTVGNKLVF (SEQ ID NO: 175) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRGWREQFF (SEQ ID NO: 180) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxiii) comprises a CDR3 α comprising the amino acid sequence of CAARSYNTDKLIF (SEQ ID NO: 186) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSWGYQETQYF (SEQ ID NO: 196) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxiv) comprises a CDR3 α comprising the amino acid sequence of CAARSYNTDKLIF (SEQ ID NO: 186) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPTGGEYYGYTF (SEQ ID NO: 202) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxv) comprises a CDR3 α comprising the amino acid sequence of CAARSYNTDKLIF (SEQ ID NO: 186) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of

CASSSYPLRTGRYNSYNSPLHF (SEQ ID NO: 208) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxvi) comprises a CDR3 α comprising the amino acid sequence of CAASYNNARLMF (SEQ ID NO: 191) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSWGYQETQYF (SEQ ID NO: 196) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxvii) comprises a CDR3 α comprising the amino acid sequence of CAASYNNARLMF (SEQ ID NO: 191) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPTGGEYYGYTF (SEQ ID NO: 202) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxviii) comprises a CDR3 α comprising the amino acid sequence of CAASYNNARLMF (SEQ ID NO: 191) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSSYPLRTGRYNSYNSPLHF (SEQ ID NO: 208) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxix) comprises a CDR3 α comprising the amino acid sequence of CAASGGRDDKIIF (SEQ ID NO: 214) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSYSRTESTDTQYF (SEQ ID NO: 219) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxx) comprises a CDR3 α comprising the amino acid sequence of CAANNARLMF (SEQ ID NO: 92) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPGQHGELEFF (SEQ ID NO: 271) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxi) comprises a CDR3 α comprising the amino acid sequence of CAASATGNQFYF (SEQ ID NO: 266) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSDTRAREQFF (SEQ ID NO: 97) or a variant thereof having up to three amino acid substitutions, additions or deletions;

- (xxxii) comprises a CDR3 α comprising the amino acid sequence of CAASATGNQFYF (SEQ ID NO: 266) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPGQHGELEFF (SEQ ID NO: 271) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxxiii) comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxxiv) comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSVGGSGSYNEQFF (SEQ ID NO: 169) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxxv) comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxxvi) comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLGLSISQETQYF (SEQ ID NO: 288) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxxvii) comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLGLSISQETQYF (SEQ ID NO: 288) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxxviii) comprises a CDR3 α comprising the amino acid sequence of CAVEATDSWGKLQF (SEQ ID NO: 108) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino

acid sequence of CASSLGLSISQETQYF (SEQ ID NO: 288) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxix) comprises a CDR3 α comprising the amino acid sequence of CAVRTSYDKVIF (SEQ ID NO: 113) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLGLSISQETQYF (SEQ ID NO: 288) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxx) comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxxi) comprises a CDR3 α comprising the amino acid sequence of CAVEATDSWGKLQF (SEQ ID NO: 108) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions; or

(xxxxii) comprises a CDR3 α comprising the amino acid sequence of CAVRTSYDKVIF (SEQ ID NO: 113) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions.

In one embodiment, the present invention provides a TCR of the present invention comprising the following CDR sequences:

(i) CDR1 α - KALYS (SEQ ID NO: 1),
 CDR2 α - LLKGGEQ (SEQ ID NO: 2),
 CDR3 α - CGTAWINDYKLSF (SEQ ID NO: 3),
 CDR1 β - SGHDY (SEQ ID NO: 6),
 CDR2 β - FNNNVP (SEQ ID NO: 7), and
 CDR3 β - CASRKTGGYSNQPQHF (SEQ ID NO: 8),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (ii) CDR1 α - NSASQS (SEQ ID NO: 34),
 CDR2 α - VYSSGN (SEQ ID NO: 35),
 CDR3 α - CVVNLLSNQGGKLIF (SEQ ID NO: 36),
 CDR1 β - LGHNA (SEQ ID NO: 39),
 5 CDR2 β - YSLEER (SEQ ID NO: 40), and
 CDR3 β - CASSQDYLVSNKELFF (SEQ ID NO: 41),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (iii) CDR1 α - SSVPPY (SEQ ID NO: 12),
 10 CDR2 α - YTSAATLV (SEQ ID NO: 13),
 CDR3 α - CAVRLSGSARQLTF (SEQ ID NO: 14),
 CDR1 β - SGHAT (SEQ ID NO: 22),
 CDR2 β - FQNNGV (SEQ ID NO: 23), and
 CDR3 β - CASSLLGDEQYF (SEQ ID NO: 24),
 15 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (iv) CDR1 α - SSVPPY (SEQ ID NO: 12),
 CDR2 α - YTSAATLV (SEQ ID NO: 13),
 CDR3 α - CAVRLSGSARQLTF (SEQ ID NO: 14),
 20 CDR1 β - SGHTA (SEQ ID NO: 28),
 CDR2 β - FQGNSA (SEQ ID NO: 29), and
 CDR3 β - CASSLVALQGAGEQYF (SEQ ID NO: 30),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- 25 (v) CDR1 α - TSESDYY (SEQ ID NO: 17),
 CDR2 α - QEAYKQQN (SEQ ID NO: 18),
 CDR3 α - CAYRSLKYGNKLVF (SEQ ID NO: 19),
 CDR1 β - SGHAT (SEQ ID NO: 22),
 CDR2 β - FQNNGV (SEQ ID NO: 23), and
 30 CDR3 β - CASSLLGDEQYF (SEQ ID NO: 24),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (vi) CDR1 α - TSESDYY (SEQ ID NO: 17),
 CDR2 α - QEAYKQQN (SEQ ID NO: 18),
 35 CDR3 α - CAYRSLKYGNKLVF (SEQ ID NO: 19),

CDR1 β - SGHTA (SEQ ID NO: 28),
 CDR2 β - FQGNSA (SEQ ID NO: 29), and
 CDR3 β - CASSLVALQGAGEQYF (SEQ ID NO: 30),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(vii) CDR1 α – TSINN (SEQ ID NO: 45),
 CDR2 α - IRSNERE (SEQ ID NO: 46),
 CDR3 α – CATDAYSGNTPLVF (SEQ ID NO: 47),
 CDR1 β - MNHNS (SEQ ID NO: 55),
 CDR2 β - SASEGT (SEQ ID NO: 56), and
 CDR3 β - CASRAAGLDTEAFF (SEQ ID NO: 57),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(viii) CDR1 α – TSINN (SEQ ID NO: 45),
 CDR2 α - IRSNERE (SEQ ID NO: 46),
 CDR3 α – CATDAYSGNTPLVF (SEQ ID NO: 47),
 CDR1 β – MNHNY (SEQ ID NO: 61),
 CDR2 β - SVGAGI (SEQ ID NO: 62), and
 CDR3 β - CASTQTPYEQYF (SEQ ID NO: 63),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(ix) CDR1 α – TSINN (SEQ ID NO: 45),
 CDR2 α - IRSNERE (SEQ ID NO: 46),
 CDR3 α – CATDAYSGNTPLVF (SEQ ID NO: 47),
 CDR1 β – SGHNS (SEQ ID NO: 67),
 CDR2 β - FNNNVP (SEQ ID NO: 68), and
 CDR3 β – CASSTVGGEDYGYTF (SEQ ID NO: 69),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(x) CDR1 α - DSAIYN (SEQ ID NO: 50),
 CDR2 α – IQSSQRE (SEQ ID NO: 51),
 CDR3 α - CAVRAEIYNQGGKLIF (SEQ ID NO: 52),
 CDR1 β - MNHNS (SEQ ID NO: 55),
 CDR2 β - SASEGT (SEQ ID NO: 56), and
 CDR3 β - CASRAAGLDTEAFF (SEQ ID NO: 57),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xi) CDR1 α - DSAIYN (SEQ ID NO: 50),

CDR2 α - IQSSQRE (SEQ ID NO: 51),

5 CDR3 α - CAVRAEIYNQGGKLIF (SEQ ID NO: 52),

CDR1 β - MNHNY (SEQ ID NO: 61),

CDR2 β - SVGAGI (SEQ ID NO: 62), and

CDR3 β - CASTQTPYEYF (SEQ ID NO: 63),

10 or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xii) CDR1 α - DSAIYN (SEQ ID NO: 50),

CDR2 α - IQSSQRE (SEQ ID NO: 51),

CDR3 α - CAVRAEIYNQGGKLIF (SEQ ID NO: 52),

CDR1 β - SGHNS (SEQ ID NO: 67),

15 CDR2 β - FNNNVP (SEQ ID NO: 68), and

CDR3 β - CASSTVGGEDYGYTF (SEQ ID NO: 69),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xiii) CDR1 α - DSASNY (SEQ ID NO: 73),

20 CDR2 α - IRSNVGE (SEQ ID NO: 74),

CDR3 α - CAASMAGAGSYQLTF (SEQ ID NO: 75),

CDR1 β - ENHRY (SEQ ID NO: 78),

CDR2 β - SYGVKD (SEQ ID NO: 79), and

CDR3 β - CAISVGQGALYEQYF (SEQ ID NO: 80),

25 or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xiv) CDR1 α - DSASNY (SEQ ID NO: 73),

CDR2 α - IRSNVGE (SEQ ID NO: 74),

CDR3 α - CAASMAGAGSYQLTF (SEQ ID NO: 75),

30 CDR1 β - SGDLS (SEQ ID NO: 84),

CDR2 β - YYNGEE (SEQ ID NO: 85), and

CDR3 β - CASSVARDRRNYGYTF (SEQ ID NO: 86),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (xv) CDR1 α - NSMFDY (SEQ ID NO: 90),
 CDR2 α – ISSIKDK (SEQ ID NO: 91),
 CDR3 α - CAANNARLMF (SEQ ID NO: 92),
 CDR1 β – SGHNS (SEQ ID NO: 95),
 5 CDR2 β – FNNNVP (SEQ ID NO: 96), and
 CDR3 β – CASSDTRAREQFF (SEQ ID NO: 97),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (xvi) CDR1 α - DSSSTY (SEQ ID NO: 101),
 10 CDR2 α – IFSNMDM (SEQ ID NO: 102),
 CDR3 α - CAERLNTDKLIF (SEQ ID NO: 103),
 CDR1 β – DFQATT (SEQ ID NO: 161),
 CDR2 β – SNEGSKA (SEQ ID NO: 162), and
 CDR3 β – CSARDSVSGNTIYF (SEQ ID NO: 163),
 15 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (xvii) CDR1 α - DSSSTY (SEQ ID NO: 101),
 CDR2 α – IFSNMDM (SEQ ID NO: 102),
 CDR3 α - CAERLNTDKLIF (SEQ ID NO: 103),
 20 CDR1 β – SQVTM (SEQ ID NO: 167),
 CDR2 β – ANQGSEA (SEQ ID NO: 168), and
 CDR3 β – CSVGGSGSYNEQFF (SEQ ID NO: 169),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- 25 (xviii) CDR1 α - DSVNN (SEQ ID NO:106),
 CDR2 α – IPSGT (SEQ ID NO: 107),
 CDR3 α - CAVEATDSWGKLQF (SEQ ID NO: 108),
 CDR1 β – DFQATT (SEQ ID NO: 161),
 CDR2 β – SNEGSKA (SEQ ID NO: 162), and
 30 CDR3 β – CSARDSVSGNTIYF (SEQ ID NO: 163),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (xix) CDR1 α - DSVNN (SEQ ID NO:106),
 CDR2 α – IPSGT (SEQ ID NO: 107),
 35 CDR3 α - CAVEATDSWGKLQF (SEQ ID NO: 108),

CDR1 β – SQVTM (SEQ ID NO: 167),
 CDR2 β – ANQGSEA (SEQ ID NO: 168), and
 CDR3 β – CSVGGSGSYNEQFF (SEQ ID NO: 169),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xx) CDR1 α - DSASNY (SEQ ID NO: 111),
 CDR2 α – IRSNVGE (SEQ ID NO: 112),
 CDR3 α - CAVRTSYDKVIF (SEQ ID NO: 113),
 CDR1 β – DFQATT (SEQ ID NO: 161),
 CDR2 β – SNEGSKA (SEQ ID NO: 162), and
 CDR3 β – CSARDSVSGNTIYF (SEQ ID NO: 163),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxi) CDR1 α - DSASNY (SEQ ID NO: 111),
 CDR2 α – IRSNVGE (SEQ ID NO: 112),
 CDR3 α - CAVRTSYDKVIF (SEQ ID NO: 113),
 CDR1 β – SQVTM (SEQ ID NO: 167),
 CDR2 β – ANQGSEA (SEQ ID NO: 168), and
 CDR3 β – CSVGGSGSYNEQFF (SEQ ID NO: 169),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxii) CDR1 α - VGISA (SEQ ID NO: 173),
 CDR2 α – LSSGK (SEQ ID NO: 174),
 CDR3 α - CAVTVGNKLVF (SEQ ID NO: 175),
 CDR1 β – MNHNS (SEQ ID NO: 178),
 CDR2 β – SASEGT (SEQ ID NO: 179), and
 CDR3 β – CASRGWREQFF (SEQ ID NO: 180),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxiii) CDR1 α - VGISA (SEQ ID NO: 184),
 CDR2 α – LSSGK (SEQ ID NO: 185),
 CDR3 α - CAARSYNTDKLIF (SEQ ID NO: 186),
 CDR1 β – SGHTS (SEQ ID NO: 194),
 CDR2 β – YDEGEE (SEQ ID NO: 195), and
 CDR3 β – CASSWGYQETQYF (SEQ ID NO: 196),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxiv) CDR1 α - VGISA (SEQ ID NO: 184),
 CDR2 α - LSSGK (SEQ ID NO: 185),
 5 CDR3 α - CAARSYNTDKLIF (SEQ ID NO: 186),
 CDR1 β - KGHSH (SEQ ID NO: 200),
 CDR2 β - LQKENI (SEQ ID NO: 201), and
 CDR3 β - CASSPTGGEYYGYTF (SEQ ID NO: 202),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxv) CDR1 α - VGISA (SEQ ID NO: 184),
 CDR2 α - LSSGK (SEQ ID NO: 185),
 CDR3 α - CAARSYNTDKLIF (SEQ ID NO: 186),
 CDR1 β - MNHEY (SEQ ID NO: 206),
 15 CDR2 β - SVGAGI (SEQ ID NO: 207), and
 CDR3 β - CASSYPLRTGRYNSYNSPLHF (SEQ ID NO: 208),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxvi) CDR1 α - NSMFDY (SEQ ID NO: 189),
 20 CDR2 α - ISSIKDK (SEQ ID NO: 190),
 CDR3 α - CAASYNNARLMF (SEQ ID NO: 191),
 CDR1 β - SGHTS (SEQ ID NO: 194),
 CDR2 β - YDEGEE (SEQ ID NO: 195), and
 CDR3 β - CASSWGYQETQYF (SEQ ID NO: 196),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxvii) CDR1 α - NSMFDY (SEQ ID NO: 189),
 CDR2 α - ISSIKDK (SEQ ID NO: 190),
 CDR3 α - CAASYNNARLMF (SEQ ID NO: 191),
 30 CDR1 β - KGHSH (SEQ ID NO: 200),
 CDR2 β - LQKENI (SEQ ID NO: 201), and
 CDR3 β - CASSPTGGEYYGYTF (SEQ ID NO: 202),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (xxviii) CDR1 α - NSMFDY (SEQ ID NO: 189),
 CDR2 α – ISSIKDK (SEQ ID NO: 190),
 CDR3 α - CAASYNNARLMF (SEQ ID NO: 191),
 CDR1 β – MNHEY (SEQ ID NO: 206),
 5 CDR2 β – SVGAGI (SEQ ID NO: 207), and
 CDR3 β – CASSYPLRTGRYNSYNSPLHF (SEQ ID NO: 208),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (xxix) CDR1 α - NSMFDY (SEQ ID NO: 212),
 10 CDR2 α – ISSIKDK (SEQ ID NO: 213),
 CDR3 α - CAASGGRDDKIIF (SEQ ID NO: 214),
 CDR1 β – MNHEY (SEQ ID NO: 217),
 CDR2 β – SVGAGI (SEQ ID NO: 218), and
 CDR3 β – CASSYSRTESTDTQYF (SEQ ID NO: 219),
 15 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (xxx) CDR1 α - NSMFDY (SEQ ID NO: 90),
 CDR2 α – ISSIKDK (SEQ ID NO: 91),
 CDR3 α - CAANNARLMF (SEQ ID NO: 92),
 20 CDR1 β – SGHRS (SEQ ID NO: 269),
 CDR2 β – YFSETQ (SEQ ID NO: 270), and
 CDR3 β – CASSPGQHGELEFF (SEQ ID NO: 271),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- 25 (xxxi) CDR1 α - NSMFDY (SEQ ID NO: 264),
 CDR2 α – ISSIKDK (SEQ ID NO: 265),
 CDR3 α - CAASATGNQFYF (SEQ ID NO: 266),
 CDR1 β – SGHNS (SEQ ID NO: 95),
 CDR2 β – FNNNVP (SEQ ID NO: 96), and
 30 CDR3 β – CASSDTRAREQFF (SEQ ID NO: 97),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (xxxii) CDR1 α - NSMFDY (SEQ ID NO: 264),
 CDR2 α – ISSIKDK (SEQ ID NO: 265),

CDR3 α - CAASATGNQFYF (SEQ ID NO: 266),
 CDR1 β - SGHRS (SEQ ID NO: 269),
 CDR2 β - YFSETQ (SEQ ID NO: 270), and
 CDR3 β - CASSPGQHGELEFF (SEQ ID NO: 271),

5 or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxxiii) CDR1 α - TSINN (SEQ ID NO: 275),
 CDR2 α - IRSNERE (SEQ ID NO: 276),
 CDR3 α - CATDGDSSYKLIF (SEQ ID NO: 277),

10 CDR1 β - DFQATT (SEQ ID NO: 161),
 CDR2 β - SNEGSKA (SEQ ID NO: 162), and
 CDR3 β - CSARDSVSGNTIYF (SEQ ID NO: 163),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;

15 (xxxiv) CDR1 α - TSINN (SEQ ID NO: 275),
 CDR2 α - IRSNERE (SEQ ID NO: 276),
 CDR3 α - CATDGDSSYKLIF (SEQ ID NO: 277),
 CDR1 β - SQVTM (SEQ ID NO: 167),
 CDR2 β - ANQGSEA (SEQ ID NO: 168), and
 20 CDR3 β - CSVGGSGSYNEQFF (SEQ ID NO: 169),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxxv) CDR1 α - TSINN (SEQ ID NO: 275),
 CDR2 α - IRSNERE (SEQ ID NO: 276),
 25 CDR3 α - CATDGDSSYKLIF (SEQ ID NO: 277),
 CDR1 β - DFQATT (SEQ ID NO: 280),
 CDR2 β - SNEGSKA (SEQ ID NO: 281), and
 CDR3 β - CSARDVLTGDYGYTF (SEQ ID NO: 282),

30 or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxxvi) CDR1 α - TSINN (SEQ ID NO: 275),
 CDR2 α - IRSNERE (SEQ ID NO: 276),
 CDR3 α - CATDGDSSYKLIF (SEQ ID NO: 277),
 CDR1 β - SGHDY (SEQ ID NO: 286),
 35 CDR2 β - FNNNVP (SEQ ID NO: 287), and

CDR3 β – CASSLGLSISQETQYF (SEQ ID NO: 288),
or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxxvii) CDR1 α - DSSSTY (SEQ ID NO: 101),

5 CDR2 α – IFSNMDM (SEQ ID NO: 102),

CDR3 α - CAERLNTDKLIF (SEQ ID NO: 103),

CDR1 β – SGHDY (SEQ ID NO: 286),

CDR2 β – FNNNVP (SEQ ID NO: 287), and

CDR3 β – CASSLGLSISQETQYF (SEQ ID NO: 288),

10 or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxxviii) CDR1 α - DSVNN (SEQ ID NO: 106),

CDR2 α – IPSGT (SEQ ID NO: 107),

CDR3 α - CAVEATDSWGKLQF (SEQ ID NO: 108),

15 CDR1 β – SGHDY (SEQ ID NO: 286),

CDR2 β – FNNNVP (SEQ ID NO: 287), and

CDR3 β – CASSLGLSISQETQYF (SEQ ID NO: 288),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

20 (xxxix) CDR1 α - DSASNY (SEQ ID NO: 111),

CDR2 α – IRSNVGE (SEQ ID NO: 112),

CDR3 α - CAVRTSYDKVIF (SEQ ID NO: 113),

CDR1 β – SGHDY (SEQ ID NO: 286),

CDR2 β – FNNNVP (SEQ ID NO: 287), and

25 CDR3 β – CASSLGLSISQETQYF (SEQ ID NO: 288),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxxx) CDR1 α - DSSSTY (SEQ ID NO: 101),

CDR2 α – IFSNMDM (SEQ ID NO: 102),

30 CDR3 α - CAERLNTDKLIF (SEQ ID NO: 103),

CDR1 β – DFQATT (SEQ ID NO: 280),

CDR2 β – SNEGSKA (SEQ ID NO: 281), and

CDR3 β – CSARDVLTGDYGYTF (SEQ ID NO: 282),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

35

(xxxxi) CDR1 α - DSVNN (SEQ ID NO: 106),
 CDR2 α – IPSGT (SEQ ID NO: 107),
 CDR3 α - CAVEATDSWGKLQF (SEQ ID NO: 108),
 CDR1 β – DFQATT (SEQ ID NO: 280),
 5 CDR2 β – SNEGSKA (SEQ ID NO: 281), and
 CDR3 β – CSARDVLTGDYGYTF (SEQ ID NO: 282),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions; or

(xxxxii) CDR1 α - DSASNY (SEQ ID NO: 111),
 10 CDR2 α – IRSNVGE (SEQ ID NO: 112),
 CDR3 α - CAVRTSYDKVIF (SEQ ID NO: 113),
 CDR1 β – DFQATT (SEQ ID NO: 280),
 CDR2 β – SNEGSKA (SEQ ID NO: 281), and
 CDR3 β – CSARDVLTGDYGYTF (SEQ ID NO: 282),
 15 or variants thereof each having up to three amino acid substitutions, additions or
 deletions.

In one embodiment, the present invention provides a TCR of the present invention comprising:

- (i) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 4 or
 20 a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at
 least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at
 least 75%, sequence identity thereto; and a β chain variable domain comprising the
 amino acid sequence of SEQ ID NO: 9 or a variant thereof having at least 75%, at
 least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at
 25 least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (ii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 37
 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at
 least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at
 least 75%, sequence identity thereto; and a β chain variable domain comprising the
 30 amino acid sequence of SEQ ID NO: 42 or a variant thereof having at least 75%, at
 least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at
 least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (iii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 15
 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at

- least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 25 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- 5 (iv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 15 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 31 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- 10 (v) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 20 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 25 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- 15 (vi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 20 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 31 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- 20 (vii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 48 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 58 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- 25
- 30

- (viii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 48 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 64 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (ix) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 48 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 70 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (x) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 53 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 58 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 53 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 64 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 53 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 70 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

- (xiii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 76 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 81 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xiv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 76 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 87 or a variant thereof at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 93 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 98 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xvi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 104 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 164 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xvii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 104 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 170 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

- (xviii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 109 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 164 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xix) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 109 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 170 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xx) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 114 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 164 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 114 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 170 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 176 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 181 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

- (xxiii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 187 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 197 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxiv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 187 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 203 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 187 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 209 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxvi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 192 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 197 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxvii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 192 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 203 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

- (xxviii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 192 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 209 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; or
- (xxix) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 215 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 220 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxx) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 93 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 272 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 267 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 98 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 267 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 272 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

- (xxxiii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 278 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 164 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxiv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 278 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 170 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 278 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 283 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxvi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 278 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 289 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxvii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 104 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 289 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

- (xxxviii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 109 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 289 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxix) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 114 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 289 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxx) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 104 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 283 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxxi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 109 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 283 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; or
- (xxxxii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 114 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 283 or a variant thereof having at least 75%, at

least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto.

In one embodiment, the present invention provides a TCR of the present invention comprising:

- 5 (i) an α chain comprising the amino acid sequence of SEQ ID NO: 5 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 10, SEQ ID NO: 11 and variants of SEQ ID NOs: 10
10 and 11 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (ii) an α chain comprising the amino acid sequence of SEQ ID NO: 38 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at
15 least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 43, SEQ ID NO: 44 and variants of SEQ ID NOs: 43 and 44 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%,
20 preferably at least 75%, sequence identity thereto;
- (iii) an α chain comprising the amino acid sequence of SEQ ID NO: 16 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at
25 least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 26, SEQ ID NO: 27 and variants of SEQ ID NOs: 26 and 27 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (iv) an α chain comprising the amino acid sequence of SEQ ID NO: 16 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at
30 least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 32, SEQ ID NO: 33 and variants of SEQ ID NOs: 32 and 33 having at least 75%, at least 80%, at least 85%, at least

90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

- 5 (v) an α chain comprising the amino acid sequence of SEQ ID NO: 21 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 26, SEQ ID NO: 27 and variants of SEQ ID NOs: 26 and 27 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- 10 (vi) an α chain comprising the amino acid sequence of SEQ ID NO: 21 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 32, SEQ ID NO: 33 and variants of SEQ ID NOs: 32 and 33 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- 15 (vii) an α chain comprising the amino acid sequence of SEQ ID NO: 49 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 59, SEQ ID NO: 60 and variants of SEQ ID NOs: 59 and 60 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- 20 (viii) an α chain comprising the amino acid sequence of SEQ ID NO: 49 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 65, SEQ ID NO: 66 and variants of SEQ ID NOs: 65 and 66 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- 25
- 30

- (ix) an α chain comprising the amino acid sequence of SEQ ID NO: 49 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 71, SEQ ID NO: 72 and variants of SEQ ID NOs: 71 and 72 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (x) an α chain comprising the amino acid sequence of SEQ ID NO: 54 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 59, SEQ ID NO: 60 and variants of SEQ ID NOs: 59 and 60 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xi) an α chain comprising the amino acid sequence of SEQ ID NO: 54 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 65, SEQ ID NO: 66 and variants of SEQ ID NOs: 65 and 66 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xii) an α chain comprising the amino acid sequence of SEQ ID NO: 54 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 71, SEQ ID NO: 72 and variants of SEQ ID NOs: 71 and 72 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xiii) an α chain comprising the amino acid sequence of SEQ ID NO: 77 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%,

sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 82, SEQ ID NO: 83 and variants of SEQ ID NOs: 82 and 83 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xiv) an α chain comprising the amino acid sequence of SEQ ID NO: 77 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 88, SEQ ID NO: 89 and variants of SEQ ID NOs: 88 and 89 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xv) an α chain comprising the amino acid sequence of SEQ ID NO: 94 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 99, SEQ ID NO: 100 and variants of SEQ ID NO: 99, SEQ ID NO: 100 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xvi) an α chain comprising the amino acid sequence of SEQ ID NO: 105 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 165, SEQ ID NO: 166 and variants of SEQ ID NOs: 165 and 166 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xvii) an α chain comprising the amino acid sequence of SEQ ID NO: 105 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 171, SEQ ID NO: 172 and variants of SEQ ID NOs: 171 and 172 having at least 75%, at least 80%, at least 85%, at least

90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xviii) an α chain comprising the amino acid sequence of SEQ ID NO: 110 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 165, SEQ ID NO: 166 and variants of SEQ ID NOs: 165 and 166 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xix) an α chain comprising the amino acid sequence of SEQ ID NO: 110 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 171, SEQ ID NO: 172 and variants of SEQ ID NOs: 171 and 172 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xx) an α chain comprising the amino acid sequence of SEQ ID NO: 160 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 165, SEQ ID NO: 166 and variants of SEQ ID NOs: 165 and 166 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxi) an α chain comprising the amino acid sequence of SEQ ID NO: 160 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 171, SEQ ID NO: 172 and variants of SEQ ID NOs: 171 and 172 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

- (xxii) an α chain comprising the amino acid sequence of SEQ ID NO: 177 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 182, SEQ ID NO: 183 and variants of SEQ ID NOs: 182 and 183 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxiii) an α chain comprising the amino acid sequence of SEQ ID NO: 188 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 198, SEQ ID NO: 199 and variants of SEQ ID NOs: 198 and 199 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxiv) an α chain comprising the amino acid sequence of SEQ ID NO: 188 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 204, SEQ ID NO: 205 and variants of SEQ ID NOs: 204 and 205 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxv) an α chain comprising the amino acid sequence of SEQ ID NO: 188 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 210, SEQ ID NO: 211 and variants of SEQ ID NOs: 210 and 211 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxvi) an α chain comprising the amino acid sequence of SEQ ID NO: 193 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%,

sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 198, SEQ ID NO: 199 and variants of SEQ ID NOs: 198 and 199 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxvii) an α chain comprising the amino acid sequence of SEQ ID NO: 193 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 204, SEQ ID NO: 205 and variants of SEQ ID NOs: 204 and 205 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxviii) an α chain comprising the amino acid sequence of SEQ ID NO: 193 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 210, SEQ ID NO: 211 and variants of SEQ ID NOs: 210 and 211 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxix) an α chain comprising the amino acid sequence of SEQ ID NO: 216 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 221, SEQ ID NO: 222 and variants of SEQ ID NOs: 221 and 222 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxx) an α chain comprising the amino acid sequence of SEQ ID NO: 94 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 273, SEQ ID NO: 274 and variants of SEQ ID NOs: 273 and 274 having at least 75%, at least 80%, at least 85%, at

least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxxi) an α chain comprising the amino acid sequence of SEQ ID NO: 268 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 99, SEQ ID NO: 100 and variants of SEQ ID NOs: 99 and 100 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxxii) an α chain comprising the amino acid sequence of SEQ ID NO: 268 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 273, SEQ ID NO: 274 and variants of SEQ ID NOs: 273 and 274 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxxiii) an α chain comprising the amino acid sequence of SEQ ID NO: 279 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 165, SEQ ID NO: 166 and variants of SEQ ID NOs: 165 and 166 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxxiv) an α chain comprising the amino acid sequence of SEQ ID NO: 279 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 171, SEQ ID NO: 172 and variants of SEQ ID NOs: 171 and 172 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

- (xxxv) an α chain comprising the amino acid sequence of SEQ ID NO: 279 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 284, SEQ ID NO: 285 and variants of SEQ ID NOs: 284 and 285 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxvi) an α chain comprising the amino acid sequence of SEQ ID NO: 279 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 290, SEQ ID NO: 291 and variants of SEQ ID NOs: 290 and 291 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxvii) an α chain comprising the amino acid sequence of SEQ ID NO: 105 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 290, SEQ ID NO: 291 and variants of SEQ ID NOs: 290 and 291 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxviii) an α chain comprising the amino acid sequence of SEQ ID NO: 110 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 290, SEQ ID NO: 291 and variants of SEQ ID NOs: 290 and 291 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;
- (xxxix) an α chain comprising the amino acid sequence of SEQ ID NO: 160 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%,

sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 290, SEQ ID NO: 291 and variants of SEQ ID NOs: 290 and 291 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxxx) an α chain comprising the amino acid sequence of SEQ ID NO: 105 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 284, SEQ ID NO: 285 and variants of SEQ ID NOs: 284 and 285 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto;

(xxxxi) an α chain comprising the amino acid sequence of SEQ ID NO: 110 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 284, SEQ ID NO: 285 and variants of SEQ ID NOs: 284 and 285 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; or

(xxxxii) an α chain comprising the amino acid sequence of SEQ ID NO: 160 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 284, SEQ ID NO: 285 and variants of SEQ ID NOs: 284 and 285 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto.

In one embodiment, the present invention provides a TCR of the present invention comprising an α chain comprising the amino acid sequence of SEQ ID NO: 257 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence of SEQ ID NO: 259 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least

96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto.

In one embodiment, the present invention provides a TCR of the present invention comprising an α chain comprising the amino acid sequence of SEQ ID NO: 261 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto; and a β chain comprising an amino acid sequence of SEQ ID NO: 263 or a variant thereof having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, preferably at least 75%, sequence identity thereto.

A TCR of the present invention may bind to a WT1 peptide comprising an amino acid sequence selected from the group consisting of EPASQHTLRSG (SEQ ID NO: 123), YESDNHTTPIL (SEQ ID NO: 126), NHTTPILCGAQYRIH (SEQ ID NO: 127), QCLSAFTVHFSGQFT (SEQ ID NO: 118), EDPMGQQGSLGEQQY (SEQ ID NO: 119), SQLECMTWNQMNLGA (SEQ ID NO: 120), APVLDFAPPGA (SEQ ID NO: 117), NQMNLGATLKG (SEQ ID NO: 250), DPGGIWAKLGAAEAS (SEQ ID NO: 251), NHTTPILCGAQYRIH (SEQ ID NO: 252), KRHQRRHTGVKPFQC (SEQ ID NO: 253), PSCQKKFARSDELVR (SEQ ID NO: 254) and variants thereof each having up to three amino acid substitutions, additions or deletions.

In another aspect, the present invention provides a T-cell receptor (TCR), which binds to a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the WT1 peptide comprises an amino acid sequence selected from the group consisting of EPASQHTLRSG (SEQ ID NO: 123), YESDNHTTPIL (SEQ ID NO: 126), NHTTPILCGAQYRIH (SEQ ID NO: 127), QCLSAFTVHFSGQFT (SEQ ID NO: 118), EDPMGQQGSLGEQQY (SEQ ID NO: 119), SQLECMTWNQMNLGA (SEQ ID NO: 120), APVLDFAPPGA (SEQ ID NO: 117), NQMNLGATLKG (SEQ ID NO: 250), DPGGIWAKLGAAEAS (SEQ ID NO: 251), NHTTPILCGAQYRIH (SEQ ID NO: 252), KRHQRRHTGVKPFQC (SEQ ID NO: 253), PSCQKKFARSDELVR (SEQ ID NO: 254) and variants thereof each having up to three amino acid substitutions, additions or deletions.

In one embodiment, a TCR of the present invention binds to an MHC I and/or MHC II peptide complex.

In one embodiment, a TCR of the present invention is restricted to a human leukocyte antigen (HLA) allele. In one embodiment, a TCR of the present invention is restricted to a HLA-A or a HLA-B allele. In one embodiment, a TCR of the present invention is restricted to

a HLA-A allele selected from the group consisting of HLA-A*0201, HLA-A*0101, HLA-A*2402 and HLA-A*0301 or a HLA-B allele selected from the group consisting of HLA-B*0702, HLA-B*3501 and HLA-B*3502.

In one embodiment, a TCR of the present invention is restricted to HLA-A*0201.

5 In one embodiment, a TCR of the present invention is restricted to HLA-B*3502.

In one embodiment, a TCR of the present invention is restricted to HLA-B*3501.

In one embodiment, a TCR of the present invention is restricted to a HLA-C allele. In one embodiment, a TCR of the present invention is restricted to a HLA-C allele selected from the group consisting of HLA-C*07:01, HLA-C*03:04, HLA-C*04:01, HLA-C*05:01, HLA-C*06:02
10 and HLA-C*07:02.

In one embodiment, a TCR of the present invention comprises one or more mutations at the α chain/ β chain interface, such that when the α chain and the β chain are expressed in a T-cell, the frequency of mispairing between said chains and endogenous TCR α and β chains is reduced.

15 In one embodiment, a TCR of the present invention comprises one or more mutations at the α chain/ β chain interface, such that when the α chain and the β chain are expressed in a T-cell, the level of expression of the TCR α and β chains is increased.

In one embodiment, the one or more mutations introduce a cysteine residue into the constant region domain of each of the α chain and the β chain, wherein the cysteine
20 residues are capable of forming a disulphide bond between the α chain and the β chain.

A TCR of the present invention may comprise a murinized constant region.

In one embodiment, the TCR of the invention is a soluble TCR.

In another aspect, the present invention provides an isolated polynucleotide encoding the α chain of a T-cell receptor (TCR) of the present invention, and/or the β chain of a TCR of the
25 present invention.

In another aspect, the present invention provides an isolated polynucleotide comprising a nucleotide sequence of SEQ ID NO: 256 and/or a nucleotide sequence of SEQ ID NO: 258, or variants thereof having at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at
30 least 98%, or at least 99% sequence identity thereto.

In another aspect, the present invention provides an isolated polynucleotide comprising a nucleotide sequence of SEQ ID NO: 260 and/or a nucleotide sequence of SEQ ID NO: 262, or variants thereof having at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at
 5 least 98%, or at least 99% sequence identity thereto.

In one embodiment, the isolated polynucleotide encodes the α chain linked to the β chain. In one embodiment, the isolated polynucleotide encodes one or more short interfering RNA (siRNA) sequences and/or one or more other agents capable of reducing or preventing expression of one or more endogenous TCR genes.

10 In another aspect, the present invention provides a vector comprising a polynucleotide of the present invention. In one embodiment, the vector comprises a polynucleotide which encodes one or more CD3 chains, CD8, a suicide gene, and/or a selectable marker.

In another aspect, the present invention provides a cell comprising a TCR of the present invention, a polynucleotide of the present invention, or a vector of the present invention.

15 In one embodiment, the cell further comprises a vector which encodes one or more CD3 chains, CD8, a suicide gene and/or a selectable marker.

In one embodiment, the cell is a T-cell, a lymphocyte or a stem cell, such as hematopoietic stem cells or induced pluripotent stem cells (iPS). The T-cell, the lymphocyte, or the stem cell may be selected from the group consisting of CD4 cells, CD8 cells, Th0 cells, Tc0 cells,
 20 Th1 cells, Tc1 cells, Th2 cells, Tc2 cells, Th17 cells, Th22 cells, gamma/delta T-cells, natural killer (NK) cells, natural killer T (NKT) cells, double negative T-cells, naive T-cells, memory stem T-cells, central memory T-cells, effector memory T-cells, effector T cells, hematopoietic stem cells and pluripotent stem cells.

In one embodiment, the cell is a T-cell which has been isolated from a subject.

25 In one embodiment, an endogenous gene encoding a TCR α chain and/or an endogenous gene encoding a TCR β chain in the cell is disrupted, preferably such that the endogenous gene encoding a TCR α chain and/or the endogenous gene encoding a TCR β chain is not expressed. In one embodiment, the endogenous gene encoding a TCR α chain and/or the
 30 endogenous gene encoding a TCR β chain is disrupted by insertion of an expression cassette comprising a polynucleotide sequence encoding a TCR of the present invention. In one embodiment, one or more endogenous genes encoding an MHC in the cell is disrupted, preferably wherein the cell is a non-alloreactive universal T-cell. In one embodiment, an endogenous gene involved in persistence, expansion, activity, resistance to

exhaustion/senescence/inhibitory signals, homing capacity, or other T-cell functions in the cell is disrupted, preferably wherein the endogenous gene involved in persistence, expansion, activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other T-cell functions is selected from the group consisting of *PD1*, *TIM3*, *LAG3*, *2B4*,
 5 *KLRG1*, *TGFbR*, *CD160* and *CTLA4*. In one embodiment, the endogenous gene involved in persistence, expansion, activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other T-cell functions is disrupted by integration of an expression cassette, wherein the expression cassette comprises a polynucleotide sequence encoding a TCR of the present invention.

10 In another aspect, the present invention provides a method of preparing a cell, which comprises the step of introducing a vector of the invention into a cell *in vitro*, *ex vivo* or *in vivo*, for example by transfection or transduction.

In another aspect, the present invention provides a method of preparing a cell, which comprises the step of transducing a cell *in vitro*, *ex vivo* or *in vivo* with one or more vectors
 15 of the present invention.

In one embodiment, the cell to be transduced with the one or more vectors is selected from the group consisting of T-cells, lymphocytes or stem cells, such as hematopoietic stem cells or induced pluripotent stem cells (iPS), optionally the T-cell, the lymphocyte or the stem cell may be selected from the group consisting of CD4 cells, CD8 cells, Th0 cells, Tc0 cells, Th1
 20 cells, Tc1 cells, Th2 cells, Tc2 cells, Th17 cells, Th22 cells, gamma/delta T-cells, natural killer (NK) cells, natural killer T (NKT) cells, double negative T-cells, naive T-cells, memory stem T-cells, central memory T-cells, effector memory T-cells, effector T cells, hematopoietic stem cells and pluripotent stem cells.

In one embodiment, the method comprises the step of T-cell editing, which comprises
 25 disrupting an endogenous gene, for example an endogenous gene encoding a TCR α chain and/or an endogenous gene encoding a TCR β chain with an artificial nuclease, preferably wherein the artificial nuclease is selected from the group consisting of zinc finger nucleases (ZFN), transcription activator-like effector nucleases (TALEN) and CRISPR/Cas system.

In one embodiment, the method comprises the step of T-cell editing, which comprises
 30 disrupting an endogenous gene encoding a TCR α chain and/or an endogenous gene encoding a TCR β chain with an artificial nuclease, preferably wherein the artificial nuclease is selected from the group consisting of zinc finger nucleases (ZFN), transcription activator-like effector nucleases (TALEN) and CRISPR/Cas system.

In one embodiment, the method comprises the step of targeted integration of an expression cassette into the endogenous gene encoding the TCR α chain gene and/or the endogenous gene encoding the TCR β chain disrupted by the artificial nuclease, wherein the expression cassette comprises a polynucleotide encoding a TCR of the present invention or a
 5 polynucleotide sequence of the present invention.

In one embodiment, the method comprises the step of disrupting one or more endogenous genes encoding an MHC, preferably wherein the cell prepared by the method is a non-alloreactive universal T-cell.

In one embodiment, the method comprises the step of disrupting one or more endogenous
 10 MHC genes, preferably wherein the cell prepared by the method is a non-alloreactive universal T-cell.

In one embodiment, the method comprises the step of disrupting one or more endogenous genes to modify the persistence, expansion, activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other T-cell functions,
 15 preferably wherein the method comprises the step of targeted integration of an expression cassette into an endogenous gene involved in persistence, expansion, activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other T-cell functions disrupted by an artificial nuclease, wherein the expression cassette comprises a polynucleotide sequence encoding a TCR of the present invention, preferably wherein the
 20 endogenous gene is selected from the group consisting of *PD1*, *TIM3*, *LAG3*, *2B4*, *KLRG1*, *TGFbR*, *CD160* and *CTLA4*.

In another aspect, the present invention provides a cell of the present invention or a cell prepared by a method of the present invention for use in adoptive cell transfer, preferably adoptive T-cell transfer, optionally the adoptive T-cell transfer may be allogenic adoptive T-
 25 cell transfer, universal non-alloreactive T-cell transfer, or autologous adoptive T-cell transfer.

In another aspect, the present invention provides a TCR of the present invention, an isolated polynucleotide of the present invention, a vector of the present invention, a cell of the present invention, a cell prepared by a method of the present invention, or a chimeric molecule of the present invention for use in therapy.

30 In another aspect, the present invention provides a TCR of the present invention, an isolated polynucleotide of the present invention, a vector of the present invention, a cell of the present invention, a cell prepared by a method the present invention for use in treating and/or preventing a disease associated with expression of WT1.

In another aspect, the present invention provides a T-cell genetically engineered (genetically edited) to modify the persistence, expansion, activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other T cell functions, wherein the T-cell expresses a TCR α chain of the present invention and/or a TCR β chain of the present invention.

In another aspect, the present invention provides a T cell genetically engineered (genetically edited) by a protocol which comprises the step of targeted integration of an expression cassette into an endogenous gene involved in persistence, expansion, activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other T-cell functions disrupted by an artificial nuclease, wherein the expression cassette comprises a polynucleotide sequence encoding TCR α chain of the present invention and/or a TCR β chain of the present invention.

In another aspect, the present invention provides a method for treating and/or preventing a disease associated with expression of WT1, which comprises the step of administering a TCR of the present invention, an isolated polynucleotide of the present invention, a vector of the present invention, a cell of the present invention, a cell prepared by a method of the present invention, or a chimeric molecule of the present invention to a subject in need thereof.

The disease associated with expression of WT1 may be a proliferative disorder. Preferably the proliferative disorder may be selected from the group consisting of hematological malignancies, such as acute myeloid leukemia (AML), chronic myeloid leukemia (CML), lymphoblastic leukemia, myelodysplastic syndromes, multiple myeloma, non Hodgkin lymphoma, Hodgkin lymphoma. The proliferative disorder may be selected from the group of solid tumors, such as lung cancer, breast cancer, oesophageal cancer, gastric cancer, colon cancer, cholangiocarcinoma, pancreatic cancer, ovarian cancer, head and neck cancers, synovial sarcoma, angiosarcoma, osteosarcoma, thyroid cancer, endometrial cancer, neuroblastoma, rhabdomyosarcoma, liver cancer, melanoma, prostate cancer, renal cancer, soft tissue sarcoma, urothelial cancer, biliary cancer, glioblastoma, mesothelioma, cervical cancer, and colorectal cancer.

In a preferred embodiment, the disease associated with expression of WT1 is acute myeloid leukemia (AML).

In another preferred embodiment, the disease associated with expression of WT1 is chronic myeloid leukemia (CML).

In another aspect, the present invention provides an isolated immunogenic WT1 peptide comprising an amino acid sequence selected from the group consisting of EPASQHTLRSG (SEQ ID NO: 123), YESDNHTTPIL (SEQ ID NO: 126), NHTTPILCGAQYRIH (SEQ ID NO: 127), QCLSAFTVHFSGQFT (SEQ ID NO: 118), EDPMGQQGSLGEQQY (SEQ ID NO: 119),
 5 SQLECMTWNQMNLGA (SEQ ID NO: 120), APVLDFAPPGA (SEQ ID NO: 117), NQMNLGATLKG (SEQ ID NO: 250), DPGGIWAKLGAAEAS (SEQ ID NO: 251), NHTTPILCGAQYRIH (SEQ ID NO: 252), KRHQRRHTGVKPFQC (SEQ ID NO: 253), PSCQKKFARSDELVR (SEQ ID NO: 254) and variants thereof each having up to three amino acid substitutions, additions or deletions.

10 DESCRIPTION OF THE DRAWINGS

Figure 1. Plots showing the results of *in vitro* expansion of functional WT-1 specific T-cells from peripheral blood of ten healthy donors

Peripheral blood mononuclear cells of ten healthy donors (HD) were stimulated with pooled, overlapping WT1 15-mer peptides for 26-30 hours, enriched for CD137⁺ cells, and expanded
 15 for 9-19 days. Expanded T-cells were re-stimulated for 6 hours with autologous antigen presenting cells (APCs) loaded with an unrelated peptide pool or WT1 peptide pool. Additionally, negative (T-cells unstimulated) and positive (T-cells cultured in the presence of PMA and Ionomycin) controls were included in the experimental setting (not shown). Dot plots indicate the results of the intracellular staining for IFN γ production and CD107a
 20 exposure on cell surface. After several re-stimulations with autologous APCs loaded with WT1 peptide pool, T-cells specificity was tested by intracellular staining as previously described. Results showed an enrichment of WT1-specific T-cells in the CD8 T cell compartment for HD1 (a), HD3 (c), HD4 (d), HD5 (e), HD6 (f), HD7 (g), and HD10 (j) and in the CD4 T cell compartment for HD2 (b), HD8 (h) and HD9 (i). WT1, Wilms Tumor 1; PMA,
 25 Phorbol 12-myristate 13-acetate; IFN γ , interferon- γ ; S, stimulation.

Figure 2. Grid and plots showing the identification of WT1-immunogenic peptides by a mapping grid strategy

Epitopes recognized by T-cells sensitized *in vitro* by repeated stimulations with the pool of overlapping WT1 peptides were identified by intracellular staining. In particular, the
 30 percentage of specific T-cells responding to the mapping grid of subpools of WT1 pentadecapeptides loaded on APCs was assessed. Additionally, negative (T-cells unstimulated and T-cells co-cultured with APCs loaded with an unrelated peptide pool) and positive (T-cells cultured in the presence of PMA and Ionomycin) controls were included in the experimental setting (T-cells unstimulated and PMA/Iono conditions are not shown).

(a) Deconvolution grid indicating the percentage of T-cells expressing IFN γ and CD107a after co-culture with APCs loaded with the different subpools (denoted SP1-24). IFN γ and CD107a values in bold text denote subpools that contain the WT1 epitope recognized by the T-cells. Representative dot plots relative to the co-culture of the T-cells with APCs loaded with the responsive subpools and indicating the expression of IFN γ and CD107a are reported. Dominant responses were observed for: subpools 4, 5, 16 in HD1 (**b**), HD3 (**d**), HD6 (**g**), HD7 (**h**), HD10 (**k**); subpools 6, 16, 17, 20, 23 in HD2 (**c**); subpools 4, 5, 6, 14, 18, 21 in HD4 (**e**); subpools 5, 11, 12, 21, 22 in HD5 (**f**); subpools 12, 14 for HD8 (**i**); subpools 5, 13, 21 for HD9 (**j**). For HD7, we also observed an increased IFN γ secretion and CD107a expression in response to subpools 7, 8, 20, even though at lower percentages compared to the response observed with subpools 4, 5, 16. SP, subpools; WT1, Wilms Tumor 1; APC, antigen-presenting cells; PMA, Phorbol 12-myristate 13-acetate; IFN γ , interferon- γ .

Figure 3. Epitope specificity of the WT1-specific T cells generated by sensitization with the pooled peptides.

In order to validate the WT1 immunogenic peptides, T-cells expanded from each HD were co-cultured for 6 hours in the presence of APCs loaded with the peptides identified after deconvolution of the mapping grid and with at least one unrelated peptide as negative control. Additionally, negative (T cells unstimulated) and positive (T cells cultured in the presence of PMA and Ionomycin) controls were included in the experimental setting (not shown). Dot plots show for each HD the results of the intracellular staining for IFN γ and/or surface CD107a. Enrichment of CD107a and/or IFN γ positive cells was respectively observed for T-cells co-cultured with peptides 40 and 41 for HD1 (**a**) and not for peptide 42 and 43 (unrelated peptides); peptides 54, 77, 90 for HD2 (**b**) and not for peptide 42 and 138 (unrelated peptides); peptide VLDFAPPGA (SEQ ID NO: 157, VLD, which is a nonamer of the peptide represented by SEQ ID NO: 117 (referred to as “11 mer” in Figure 3c)) for HD3 (**c**) and low response with peptides PVLDFAPPG (SEQ ID NO: 158, PVL, which is another nonamer of the peptide represented by SEQ ID NO: 117) and LDFAPPGAS (SEQ ID NO: 159, LDF, which is a nonamer of the peptide represented by SEQ ID NO: 116, previously described as an immunogenic peptide (Doubrovina, E. et al. (2012) Blood 120: 1633–1646)); peptides 17, 18, 99, 100 for HD4 (**d**, **e**) and not for the unrelated peptides (15, 16, 63-66, 101, 102 and 132); peptide 101 for HD5 (**f**) and not for the unrelated peptides (63, 107, 108, 113, 119 and 120); peptide VLDFAPPGA (SEQ ID NO: 157, VLD, which is a nonamer of the peptide represented by SEQ ID NO: 117 (referred to as “11 mer” in Figure 3c)) for HD6 (**g**) and peptide PVLDFAPPG (SEQ ID NO: 158, PVL, which is another nonamer of the peptide represented by SEQ ID NO: 117) and not for peptide LDFAPPGAS (SEQ ID NO: 159, LDF, which is a nonamer of the peptide represented by SEQ ID NO: 116, previously described as

an immunogenic peptide (Doubrovina, E. et al. (2012) Blood 120: 1633–1646)); peptides 101, 125, 137 for HD9 (h); peptide VLDFAPPGA (SEQ ID NO: 157, VLD, which is a nonamer of the peptide represented by SEQ ID NO: 117 (referred to as “11 mer” in Figure 3c)) for HD10 (i) and not for the unrelated peptide. For HD7 and HD8, due to a reduced fitness of T cells, it was not possible to perform functional tests to verify the peptide predicted by the deconvolution of the mapping grid, i.e. peptides 40, 41, 91, 92 for HD7 and peptide 24 for HD8.

In order to determine the HLA restriction of the WT1 epitopes identified for HD4, HD5 and HD10 T-cells, donor DNA was sequenced to determine the HLA typing. Afterwards, WT1-specific T-cells were co-cultured with different antigen presenting EBV-BLCL cell lines, each one harboring a specific HLA allele of interest that was identified by sequencing of the HD4, HD5 or HD10 DNA. The EBV-BLCL cells were pulsed with peptide 17 for HD4, peptide 101 for HD5 and peptide VLDFAPPGA (SEQ ID NO: 157) or with an unrelated control peptide. After co-culture for 6 hours, we observed a substantial response to WT1 by the WT1-specific T-cells that had been co-cultured with EBV-BLCL cells expressing the HLA-B*3502 allele and pulsed with peptide 17 for HD4 (j), EBV-BLCL cells expressing the HLA-B*3501 allele and pulsed with peptide 101 for HD5 (k) and EBV-BLCL cells expressing the HLA-A*0201 allele and pulsed with peptide VLDFAPPGA (SEQ ID NO: 157) for HD10 (l). (m) Table showing the peptides recognized by T-cells expanded from HD1-HD10. For HD3, HD6 and HD10, the specific nonamer overlapping peptides 40 and 41 and eliciting an immune response is shown. Wilms' Tumor 1; APC, antigen-presenting cells; PMA, 2; Phorbol 12-myristate 13-acetate; IFN γ , interferon- γ ; S, stimulation.

Figure 4. Graphs and plots showing that expanded T-cells of HD1, HD3 and HD4 recognize a naturally processed WT1 epitope

(a) Graph depicting CD107 expression by CD8⁺ T-cells expanded from HD1 following co-culture with T2 cells pulsed with WT1 pool, K562 cells genetically modified to express the HLA-A*0201 allele and to overexpress the WT1 protein, or T2 cells pulsed with the non-specific control MelanA/MART1 pool as a negative control.

(b) Graph depicting the results of experiments to determine the ability of HD3 expanded T-cells to target WT1-expressing cells. The results are represented as an elimination index, which is calculated as the total number of target cells still present after co-culture with the WT1-specific T-cells divided by the total number of target cells alone. HD3 T-cells were co-cultured with T2 cells pulsed with the subpool 16 (SP16) containing the immunogenic peptide eliciting the immune response; T2 cells pulsed with the MelanA/MART1 pool (Melan

A) as negative control; K562 cells either wild type (K562) or genetically modified in order to express both the HLA-A*0201 allele and to overexpress the WT1 protein (K562 A2+WT1+).

(c) Plots depicting the results of experiments to determine the ability of WT1-specific T-cells from HD4 to eliminate target cells. HD4 T-cells were co-cultured with primary CD33+ blasts harvested from a HLA-B*3502 patient at a ratio of 10:1 or, as control, with leukemic cells from a patient not harboring the HLA-B*3502 allele. After 3 days of co-culture, results indicate a nearly complete clearance of the CD33+ HLA-B*3502 blasts when seeded with WT1-specific T-cells (CD3⁺ cells). E, effector; T, target.

Figure 5. Graph showing results of V β profiling of WT1-specific T-cells

WT1-specific T-cells generated from the different HDs after several stimulations with the WT1 pool were stained with the V β Immunoprofiling kit in order to determine the clonality of the population. In particular, the expression of the variable (V) genes of the β -chain was determined by FACS analysis. Results indicate the expression of a highly dominant V β gene in HD1 (TRBV12-3;12-4), HD2 (TRBV11-2), HD3 (TRBV4-3), HD5 (TRBV20-1) whereas for HD4, HD6, HD10 a clear enrichment of a defined V β was not detected. HD4 SP14 indicates T cells stimulated with subpool 14 which contains peptides 17-18 eliciting the highest immune response; HD4 SP18+21 indicates T cells stimulated with subpools 18 and 21 which contain peptides 63-64-65-66 and 99-100-101-102, respectively, eliciting a minimal immune response as shown in Figure 3. For HD7, HD8 and HD9, it was not possible to perform the V β Immunoprofiling analysis due to a reduced cell fitness.

Figure 6. Graphs showing results of TCR sequencing of enriched WT1-specific T-cells over time

T-cells generated from each healthy donor included in the experimental setting were characterized by TCR $\alpha\beta$ sequencing after several stimulations with the WT1 pool. Sequencing results indicated the presence of predominant clonotypes for HD1 (a), HD2 (b) and HD3 (c), HD4 (d), HD5 (e), HD6 (f), HD7 (g), HD8 (h), HD9 (i), HD10 (j). Bar charts depict the ten most predominant CDR3 amino acid sequences identified at each time point (e.g. S9 corresponds to the sequencing results obtained following the 9th round of stimulation). For each bar, starting from the x-axis, the bottom segment represents the most predominant CDR sequence. The next nine most predominant sequences are stacked above the bottom segment and are ordered by decreasing frequency going upwards. The remaining sequences are grouped together in top segment. WT1, Wilms Tumor 1; CDR3, complementarity determining region 3; S, stimulation.

Figure 7. Functional activity of genetically-modified T lymphocytes.

T cells isolated from PBMCs of healthy individuals were transduced with a bidirectional lentiviral vector encoding for the α and the β chain of TCRs isolated from HD1 and HD3. As control we transduced T cells with a previously published TCR recognizing the WT1 126-134 (RMFPNAPYL; SEQ ID NO: 255) peptide when presented by the HLA-A*0201 allele.

Transfer (TR) T lymphocytes were co-cultured for 3 days with (a) T2 cells either pulsed or not with the WT1 126-134 peptide or with the VLDFAPPGA (SEQ ID NO: 157) peptide (effector:target ratio=1:1); (b) K562 cells either wild type (K562) or genetically modified in order to express the HLA-A*0201 allele (effector:target ratio=1:1); (c) 3 different primary AML blasts selected according to the expression of the HLA-A*0201 allele and of the WT1 antigen (effector:target ratio=5:1). For the co-culture with T2 and K562 cell lines, we included untransduced T cells as control. Results indicated the ability of each TCR in recognizing the target peptide when presented by the HLA-A*0201 allele (Figure 7a) and the greater potential of HD1 TCR-transduced T cells in mediating a specific and near complete elimination of K562 cells harbouring the HLA*A0201 allele compared to HD3-TR T cells. Of note, no substantial killing of target cells was observed in the co-culture of K562 HLA*A0201 cells with WT1 126-134 TR T cells (Figure 7b). These results were further confirmed by the outcome of the co-culture experiment performed using as target cells primary AML blasts derived from 3 different AML patients (pAML1 blasts: WT1-/HLA-A*0201+; pAML2 and pAML3 blasts: WT1+/HLA-A*0201+). In this experimental setting, each individual T cell population was sorted with specific dextramers, before co-culture with targets, to enrich the purity of effector cells. We observed a greater elimination of both pAML blasts harbouring the HLA-A*0201 allele upon co-culture with HD1 TR T cells, whereas only blasts from pAML3 were recognized by HD3 T and WT1 126-134 T cells. UT, untransduced, pAML, primary acute myeloid leukemia; TR, transfer; Dx, dextramer.

DETAILED DESCRIPTION

The terms “comprising”, “comprises” and “comprised of” as used herein are synonymous with “including” or “includes”; or “containing” or “contains”, and are inclusive or open-ended and do not exclude additional, non-recited members, elements or steps. The terms “comprising”, “comprises” and “comprised of” also include the term “consisting of”.

T-cell receptor

During antigen processing, antigens are degraded inside cells and then carried to the cell surface by major histocompatibility complex (MHC) molecules. T-cells are able to recognise this peptide:MHC complex at the surface of the antigen presenting cell. There are two

different classes of MHC molecules: MHC I and MHC II, each class delivers peptides from different cellular compartments to the cell surface.

A T cell receptor (TCR) is a molecule which can be found on the surface of T-cells that is responsible for recognizing antigens bound to MHC molecules. The naturally-occurring TCR
 5 heterodimer consists of an alpha (α) and beta (β) chain in around 95% of T-cells, whereas around 5% of T-cells have TCRs consisting of gamma (γ) and delta (δ) chains.

Engagement of a TCR with antigen and MHC results in activation of the T lymphocyte on which the TCR is expressed through a series of biochemical events mediated by associated enzymes, co-receptors, and specialized accessory molecules.

10 Each chain of a natural TCR is a member of the immunoglobulin superfamily and possesses one N-terminal immunoglobulin (Ig)-variable (V) domain, one Ig-constant (C) domain, a transmembrane/cell membrane-spanning region, and a short cytoplasmic tail at the C-terminal end.

The variable domain of both the TCR α chain and β chain have three hypervariable or
 15 complementarity determining regions (CDRs). A TCR α chain or β chain, for example, comprises a CDR1, a CDR2, and a CDR3 in amino to carboxy terminal order. In general, CDR3 is the main CDR responsible for recognizing processed antigen, although CDR1 of the alpha chain has also been shown to interact with the N-terminal part of the antigenic peptide, whereas CDR1 of the beta chain interacts with the C-terminal part of the peptide.
 20 CDR2 is thought to recognize the MHC molecule.

A constant domain of a TCR may consist of short connecting sequences in which a cysteine residue forms a disulfide bond, making a link between the two chains.

An α chain of a TCR of the present invention may have a constant domain encoded by a TRAC gene. An example amino acid sequence of an α chain constant domain encoded by a
 25 TRAC gene is shown below:

```
IQNPDPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDSDVYITDKTVLDMRSMDFKSNSAV
AWSNKSDFACANAFNNSIIPEDTFFPSPESSCDVKLVEKSFETDTNLFQNLSVIGFRILL
KVAGFNLLMTLRLWSS
(SEQ ID NO: 128)
```

30 A TCR of the present invention may comprise an α chain comprising the amino acid sequence of SEQ ID NO: 128 or a variant thereof having at least 75%, at least 80%, at least

85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% sequence identity thereto, preferably at least 75% sequence identity thereto.

A β chain of a TCR of the present invention may have a constant domain encoded by a TRBC1 or a TRBC2 gene. An example amino acid sequence of a β chain constant domain
5 encoded by a TRBC1 gene is shown below:

```
DLNKVFPPEVAVFEPSEAEISHTQKATLVCLATGFFPDHVELSWVNGKEVHSGVSTDPQPL
KEQPALNDSRYCLSSRLRVSATFWQNPRNHFRCQVQFYGLSENDEWTQDRAKPVTQIVSAEA
WGRADCGFTSVSYQQGVLSATILYEILLGKATLYAVLVSAVLMLMAMVKRKDF
(SEQ ID NO: 129)
```

10 An example amino acid sequence of a β chain constant domain encoded by a TRBC2 gene is shown below:

```
DLKNVFPPEVAVFEPSEAEISHTQKATLVCLATGFYPDHVELSWVNGKEVHSGVSTDPQPL
KEQPALNDSRYCLSSRLRVSATFWQNPRNHFRCQVQFYGLSENDEWTQDRAKPVTQIVSAEA
WGRADCGFTSESYQQGVLSATILYEILLGKATLYAVLVSAVLMLMAMVKRKDSRG
15 (SEQ ID NO: 130)
```

A TCR of the present invention may comprise a β chain comprising the amino acid sequence of SEQ ID NO: 129, SEQ ID NO: 130, or variants of SEQ ID NOs: 129 and 130 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% sequence identity thereto, preferably at least 75%
20 sequence identity thereto.

The TCR of the present invention may have one or more additional cysteine residues in each of the α and β chains such that the TCR may comprise two or more disulphide bonds in the constant domains.

The structure allows the TCR to associate with other molecules like CD3 which possess
25 three distinct chains (γ , δ , and ϵ) in mammals and the ζ -chain. These accessory molecules have negatively charged transmembrane regions and are vital to propagating the signal from the TCR into the cell. The CD3- and ζ -chains, together with the TCR, form what is known as the T cell receptor complex.

The signal from the T cell complex is enhanced by simultaneous binding of the MHC
30 molecules by a specific co-receptor. For helper T-cells, this co-receptor is CD4 (specific for class II MHC); whereas for cytotoxic T-cells, this co-receptor is CD8 (specific for class I MHC). The co-receptor allows prolonged engagement between the antigen presenting cell

and the T cell and recruits essential molecules (e.g., LCK) inside the cell involved in the signalling of the activated T lymphocyte.

Accordingly, as used herein the term "T-cell receptor" (TCR) refers to molecule capable of recognising a peptide when presented by an MHC molecule. The molecule may be a heterodimer of two chains α and β (or optionally γ and δ) or it may be a single chain TCR construct. A TCR of the present invention may be a soluble TCR, e.g. omitting or altering one or more constant domains. A TCR of the present invention may comprise a constant domain.

The present invention also provides an α chain or a β chain from such a T cell receptor.

The TCR of the present invention may be a hybrid TCR comprising sequences derived from more than one species. For example, it has surprisingly been found that murine TCRs are more efficiently expressed in human T-cells than human TCRs. The TCR may therefore comprise a human variable region and murine sequences within a constant region.

A disadvantage of this approach is that the murine constant sequences may trigger an immune response, leading to rejection of the transferred T-cells. However, the conditioning regimens used to prepare patients for adoptive T-cell therapy may result in sufficient immunosuppression to allow the engraftment of T-cells expressing murine sequences.

Complementarity determining (CDR) regions

The portion of the TCR that establishes the majority of the contacts with the antigenic peptide bound to the major histocompatibility complex (MHC) is the complementarity determining region 3 (CDR3), which is unique for each T cell clone. The CDR3 region is generated upon somatic rearrangement events occurring in the thymus and involving non-contiguous genes belonging to the variable (V), diversity (D, for β and δ chains) and joining (J) genes. Furthermore, random nucleotides inserted/deleted at the rearranging loci of each TCR chain gene greatly increase diversity of the highly variable CDR3 sequence. Thus, the frequency of a specific CDR3 sequence in a biological sample indicates the abundance of a specific T cell population. The great diversity of the TCR repertoire in healthy human beings provides a wide range protection towards a variety of foreign antigens presented by MHC molecules on the surface of antigen presenting cells. In this regard, it is of note that theoretically up to 10^{15} different TCRs can be generated in the thymus.

T-cell receptor diversity is focused on CDR3 and this region is primarily responsible for antigen recognition.

The sequences of the CDR3 regions of the TCR of the present invention may be selected from those set out in Table 1 below. A TCR may comprise CDRs that comprise or consist of a CDR3 α and a CDR3 β pair described below.

5 The CDRs may, for example, comprise one, two, or three substitutions, additions or deletions from the given sequence, provided that the TCR retains the capacity to bind a WT1 peptide when presented by an MHC molecule.

As used herein, the term “protein” includes single-chain polypeptide molecules as well as multiple-polypeptide complexes where individual constituent polypeptides are linked by covalent or non-covalent means. As used herein, the term “polypeptide” refers to a polymer
10 in which the monomers are amino acids and are joined together through peptide or disulphide bonds.

Variants, derivatives, analogues, homologues and fragments

In addition to the specific proteins and polynucleotides mentioned herein, the present invention also encompasses the use of variants, derivatives, analogues, homologues and
15 fragments thereof.

In the context of the present invention, a variant of any given sequence is a sequence in which the specific sequence of residues (whether amino acid or nucleic acid residues) has been modified in such a manner that the polypeptide or polynucleotide in question substantially retains at least one of its endogenous functions. A variant sequence can be
20 obtained by addition, deletion, substitution, modification, replacement and/or variation of at least one residue present in the naturally-occurring protein.

A variant amino acid sequence of the present invention referred to as having up to three amino acid substitutions, additions or deletions may have, for example, one, two or three amino acid substitutions, additions or deletions.

25 The term “derivative” as used herein, in relation to proteins or polypeptides of the present invention includes any substitution of, variation of, modification of, replacement of, deletion of and/or addition of one (or more) amino acid residues from or to the sequence providing that the resultant protein or polypeptide substantially retains at least one of its endogenous functions.

30 The term “analogue” as used herein, in relation to polypeptides or polynucleotides includes any mimetic, that is, a chemical compound that possesses at least one of the endogenous functions of the polypeptides or polynucleotides which it mimics.

Proteins used in the present invention may also have deletions, insertions or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent protein. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues as long as the endogenous function is retained. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include asparagine, glutamine, serine, threonine and tyrosine.

A substitution may involve replacement of an amino acid for a similar amino acid (a conservative substitution). A similar amino acid is one which has a side chain moiety with related properties as grouped together, for example as shown below:

(i) basic side chains: lysine (K), arginine (R), histidine (H);

(ii) acidic side chains: aspartic acid (D) and glutamic acid (E);

(iii) uncharged polar side chains: asparagine (N), glutamine (Q), serine (S), threonine (T) and tyrosine (Y); or

(iv) non-polar side chains: glycine (G), alanine (A), valine (V), leucine (L), isoleucine (I), proline (P), phenylalanine (F), methionine (M), tryptophan (W) and cysteine (C).

Any amino acid changes should maintain the capacity of the TCR to bind WT1 peptide presented by MHC molecules.

Variant sequences may comprise amino acid substitutions, additions, deletions and/or insertions. The variation may be concentrated in one or more regions, such as the constant regions, the linker, or the framework regions of the α or β chains, or they may be spread throughout the TCR molecule.

Conservative substitutions, additions or deletions may be made, for example according to the Table below. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	G A P
		I L V
	Polar – uncharged	C S T M

		N Q
	Polar - charged	D E
		K R
AROMATIC		H F W Y

The present invention also encompasses homologous substitution (substitution and replacement are both used herein to mean the interchange of an existing amino acid residue, with an alternative residue), e.g. like-for-like substitution such as basic for basic, acidic for acidic, polar for polar etc. Non-homologous substitution may also occur e.g. from one class of residue to another or alternatively involving the inclusion of unnatural amino acids, such as ornithine.

The term “variant” as used herein may mean an entity having a certain homology with the wild type amino acid sequence or the wild type nucleotide sequence. The term “homology” can be equated with “identity”.

A variant sequence may include an amino acid sequence which may be at least 50%, 55%, 65%, 75%, 85% or 90% identical, preferably at least 95%, at least 97%, or at least 99% identical to the subject sequence. Typically, the variants will comprise the same active sites etc. as the subject amino acid sequence. Although homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity.

A variant sequence may include a nucleotide sequence which may be at least 40%, 45%, 50%, 55%, 65%, 75%, 85% or 90% identical, preferably at least 95%, at least 97%, or at least 99% identical to the subject sequence. Although homology can also be considered in terms of similarity, in the context of the present invention it is preferred to express homology in terms of sequence identity.

Preferably, reference to a sequence which has a percent identity to any one of the SEQ ID NOs detailed herein refers to a sequence which has the stated percent identity over the entire length of the SEQ ID NO referred to.

Identity comparisons can be conducted by eye or, more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate percentage homology or identity between two or more sequences.

5 Percentage homology may be calculated over contiguous sequences, i.e. one sequence is aligned with the other sequence and each amino acid in one sequence is directly compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an “ungapped” alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues.

10 Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion in the nucleotide sequence may cause the following codons to be put out of alignment, thus potentially resulting in a large reduction in percent homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without
15 penalising unduly the overall homology score. This is achieved by inserting “gaps” in the sequence alignment to try to maximise local homology.

However, these more complex methods assign “gap penalties” to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible, reflecting higher relatedness between the two compared
20 sequences, will achieve a higher score than one with many gaps. “Affine gap costs” are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is
25 preferred to use the default values when using such software for sequence comparisons. For example when using the GCG Wisconsin Bestfit package the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension.

Calculation of maximum percentage homology therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for
30 carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A.; Devereux *et al.* (1984) *Nucleic Acids Res.* 12: 387). Examples of other software that can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel *et al.* (1999) *ibid* – Ch. 18), FASTA (Atschul *et al.* (1990) *J. Mol. Biol.* 403-410) and the GENWORKS suite of comparison tools. Both BLAST and FASTA are

available for offline and online searching (see Ausubel *et al.* (1999) *ibid*, pages 7-58 to 7-60). However, for some applications, it is preferred to use the GCG Bestfit program. Another tool, called BLAST 2 Sequences is also available for comparing protein and nucleotide sequences (see *FEMS Microbiol. Lett.* (1999) 174: 247-50; *FEMS Microbiol. Lett.* (1999) 177: 187-8).

Although the final percentage homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix – the default matrix for the BLAST suite of programs. GCG Wisconsin programs generally use either the public default values or a custom symbol comparison table if supplied (see the user manual for further details). For some applications, it is preferred to use the public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62.

Once the software has produced an optimal alignment, it is possible to calculate percentage homology, preferably percentage sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

“Fragments” are also variants and the term typically refers to a selected region of the polypeptide or polynucleotide that is of interest either functionally or, for example, in an assay. “Fragment” thus refers to an amino acid or nucleic acid sequence that is a portion of a full-length polypeptide or polynucleotide.

Such variants may be prepared using standard recombinant DNA techniques such as site-directed mutagenesis. Where insertions are to be made, synthetic DNA encoding the insertion together with 5' and 3' flanking regions corresponding to the naturally-occurring sequence either side of the insertion site may be made. The flanking regions will contain convenient restriction sites corresponding to sites in the naturally-occurring sequence so that the sequence may be cut with the appropriate enzyme(s) and the synthetic DNA ligated into the cut. The DNA is then expressed in accordance with the invention to make the encoded protein. These methods are only illustrative of the numerous standard techniques known in the art for manipulation of DNA sequences and other known techniques may also be used.

Major histocompatibility complex (MHC) molecules

Typically, TCRs bind to peptides as part of peptide:MHC complex.

The MHC molecule may be an MHC class I or II molecule. The complex may be on the surface of an antigen presenting cell, such as a dendritic cell or a B cell, or any other cell, including cancer cells, or it may be immobilised by, for example, coating on to a bead or plate.

- 5 The human leukocyte antigen system (HLA) is the name of the gene complex which encodes major histocompatibility complex (MHC) in humans and includes HLA class I antigens (A, B & C) and HLA class II antigens (DP, DQ, & DR). HLA alleles A, B and C present peptides derived mainly from intracellular proteins, e.g. proteins expressed within the cell. This is of particular relevance since WT1 is an intracellular protein.
- 10 During T-cell development *in vivo*, T-cells undergo a positive selection step to ensure recognition of self MHCs followed by a negative step to remove T-cells that bind too strongly to MHC which present self-antigens. As a consequence, certain T-cells and the TCRs they express will only recognise peptides presented by certain types of MHC molecules - i.e. those encoded by particular HLA alleles. This is known as HLA restriction.
- 15 One HLA allele of interest is HLA-A*0201, which is expressed in the vast majority (>50%) of the Caucasian population. Accordingly, TCRs which bind WT1 peptides presented by MHC encoded by HLA-A*0201 (i.e. are HLA-A*0201 restricted) are advantageous since an immunotherapy making use of such TCRs will be suitable for treating a large proportion of the Caucasian population.
- 20 Other HLA-A alleles of interest are HLA-A*0101, HLA-A*2402, and HLA-A*0301.

Widely expressed HLA-B alleles of interest are HLA-B*3501, HLA-B*0702 and HLA-B*3502.

A TCR of the present invention may be HLA-A*0201-restricted.

- 25 In one aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CGTAWINDYKLSF (SEQ ID NO: 3) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRKTGGYSNQPQHF (SEQ ID NO: 8) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

- 30 In another aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CVVNLLSNQGGKLIF (SEQ ID NO: 36) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSQDYLVSNKLF (SEQ ID NO: 41) or a variant thereof

having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In another aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CAANNARLMF (SEQ ID NO: 92) or a variant thereof having up to
5 three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSDTRAREQFF (SEQ ID NO: 97) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In another aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up
10 to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In another aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CAASGGRDDKIIF (SEQ ID NO: 214) or a variant thereof having up
15 to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSYSRTESTDTQYF (SEQ ID NO: 219) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In another aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CAANNARLMF (SEQ ID NO: 92) or a variant thereof having up to
20 three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPGQHGELEFF (SEQ ID NO: 271) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In another aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CAASATGNQFYF (SEQ ID NO: 266) or a variant thereof having up
25 to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSDTRAREQFF (SEQ ID NO: 97) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In another aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CAASATGNQFYF (SEQ ID NO: 266) or a variant thereof having up
30 to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPGQHGELEFF (SEQ ID NO: 271) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In another aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In another aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In another aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In one embodiment, a TCR of the present invention that is HLA-A*0201 restricted binds to a WT1 peptide comprising amino acid sequence APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions.

In one aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of CGTAWINDYKLSF (SEQ ID NO: 3) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRKTGGYSNQPQHF (SEQ ID NO: 8) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein the WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions.

In another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of CVVNLLSNQGGKLIF (SEQ ID NO: 36) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSQDYLVSNKELFF (SEQ ID NO: 41) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein

thaw WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions.

In another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC),
5 wherein the TCR comprises a CDR3 α comprising the amino acid sequence of CAANNARLMF (SEQ ID NO: 92) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSDTRAREQFF (SEQ ID NO: 97) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein
10 the WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions.

In another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of
15 CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein the WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117)
20 or a variant thereof having up to three amino acid substitutions, additions or deletions.

In another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of
25 CAASGGRDDKIIF (SEQ ID NO: 214) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSYSRTESTDTQYF (SEQ ID NO: 219) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein the WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or
30 deletions.

In another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of
35 CAANNARLMF (SEQ ID NO: 92) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of

CASSPGQHGELEFF (SEQ ID NO: 271) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein the WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions.

- 5 In another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of CAASATGNQFYF (SEQ ID NO: 266) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of
- 10 CASSDTRAREQFF (SEQ ID NO: 97) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein the WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions.

- In another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of CAASATGNQFYF (SEQ ID NO: 266) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of
- 15 CASSPGQHGELEFF (SEQ ID NO: 271) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein the WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions.
- 20

- In another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein
- 25 the WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions.
- 30

- In another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of
- 35 CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid

substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein the WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions.

In another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-A*0201 restricted, and wherein the WT1 peptide comprises the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions.

Another widely expressed HLA allele of interest is HLA-B*3501. A TCR of the present invention may be HLA-B*3501 restricted.

In one aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CAASMAGAGSYQLTF (SEQ ID NO: 75) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CAISVGQGALYEQYF (SEQ ID NO: 80) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-B*3501 restricted.

Thus, in another aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of CAASMAGAGSYQLTF (SEQ ID NO: 75) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CAISVGQGALYEQYF (SEQ ID NO: 80) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-B*3501 restricted, and wherein the WT1 peptide comprises the amino acid sequence of NHTTPILCGAQYRIH (SEQ ID NO: 127) or a variant thereof having up to three amino acid substitutions, additions or deletions.

In one aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CAASMAGAGSYQLTF (SEQ ID NO: 75) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the

amino acid sequence of CASSVARDRRNYGYTF (SEQ ID NO: 86) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-B*3501 restricted.

Thus, in another aspect, the present invention provides a TCR which binds a Wilms tumour
 5 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of CAASMAGAGSYQLTF (SEQ ID NO: 75) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSVARDRRNYGYTF (SEQ ID NO: 86) or a variant thereof having up to three amino acid
 10 substitutions, additions or deletions, wherein the TCR is HLA-B*3501 restricted, and wherein the WT1 peptide comprises the amino acid sequence of NHTTPILCGAQYRIH (SEQ ID NO: 127) or a variant thereof having up to three amino acid substituions, additions or deletions.

Another widely expressed HLA allele of interest is HLA-B*3502. A TCR of the present invention may be HLA-B*3502 restricted.

15 In one aspect, where a TCR of the present invention comprises a CDR3 α comprising the amino acid sequence of CATDAYSGNTPLVF (SEQ ID NO: 47) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRAAGLDTEAFF (SEQ ID NO: 57) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-B*3502
 20 restricted.

Thus, in one aspect, the present invention provides a TCR which binds a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR comprises a CDR3 α comprising the amino acid sequence of CATDAYSGNTPLVF (SEQ ID NO: 47) or a variant thereof having up to three amino acid
 25 substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRAAGLDTEAFF (SEQ ID NO: 57) or a variant thereof having up to three amino acid substitutions, additions or deletions, wherein the TCR is HLA-B*3502 restricted, and wherein the WT1 peptide comprises the amino acid sequence of EPASQHTLRSG (SEQ ID NO: 123) or a variant thereof having up to three amino acid substituions, additions or deletions.

30 We have demonstrated that T-cells expressing TCRs of the present invention which bind to WT1 peptides comprising an amino acid sequence of EPASQHTLRSG (SEQ ID NO: 123) are able to selectively eliminate cancer (AML) cells expressing the HLA-B*3502 allele – see Example 4 and Figure 4c.

In one aspect, where a TCR of the present invention binds to a WT1 peptide comprising an amino acid sequence of EPASQHTLRSG (SEQ ID NO: 123), or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-B*3502 restricted.

In one embodiment, where a TCR of the present invention binds to a WT1 peptide comprising an amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-A*0201 restricted.

In one embodiment, where a TCR of the present invention binds to a WT1 peptide comprising an amino acid sequence of NHTTPILCGAQYRIH (SEQ ID NO: 127) or a variant thereof having up to three amino acid substitutions, additions or deletions, the TCR is HLA-B*3501 restricted.

Wilms tumor 1 (WT1) protein

Wilms tumor 1 (WT1) is an intracellular protein encoding a zinc finger transcription factor that plays an important role in cell growth and differentiation (Yang, L. *et al. Leukemia* 21, 868-876 (2007)). It is widely expressed on a variety of hematological and solid tumors, while showing limited expression on other tissues (gonads, uterus, kidney, mesothelium, progenitor cells in different tissues). Recent evidence suggests that WT1 plays a role in leukemogenesis and tumorigenesis.

WT1 has several isoforms, some of which result from alternative splicing of mRNA transcripts encoding WT1. The complete amino acid sequence of a WT1 isoform was previously published (Gessler, M. *et al. Nature*; 343(6260):774-778; (1990)). This particular isoform consists of 575 amino acids and includes a first 126 amino acids at the N terminus which are lacking in the exon 5+ and the KTS+ isoforms of WT1.

An example WT1 protein has the amino acid sequence set out in UniProt entry J3KNN9. Another example WT1 protein has the amino acid sequence set out below:

SRQRPHPGALRNPTACPLPHFPPSLPPTHSPTHPPRAGTAAQAPGPRRLLAAILDFLLLQDPASTCVP
EPASQHTLRSGPGCLQQPEQQGVDRDPGGIWAKLGAAEASAERLQGRRSRGASGSEPQQMGSDVRDLNA
LLPAVPSLGGGGGCALPVSGAAQWAPVLDFAPPGASAYGSLGGPAPPPAPPPPPPPPHSFIKQEPSW
GGAEPHEEQCLSAFTVHFSGQFTGTAGACRYGPFPPPSQASSGQARMFPNAPYLPSCLESQPAIRN
QGYSTVTFDGTTPSYGHTPSHHAAQFPNHSFKHEDPMGQQGSLGEQQYSVPPPVYGCHTPTDSC TGSQA
LLLRTPYSSDNLYQMTSQLECM TWNQMNLGATLKGVAAGSSSVKWTEGQSNHSTGYESDNHTTPILC
GAQYRIHTHG VFRGIQDVRRVPGVAPTLVRSASETSEKRPFMCAYPGCNKRYFKLSHLQMHSRKHTGE
KPYQCDFKDCERRFSRSDQLKRHQRRHTGVKPFQCKTCQRKFSRSDHLKTHTRHTGTSEKPFSCRW
PSCQKKFARSDELVRHHNMHQRMNTKLQLAL
(SEQ ID NO: 131)

WT1 peptides

As used herein the term peptide refers to a plurality of amino acid residues linked by peptide bonds. As defined herein a peptide may consist of less than about 30, less than about 25, less than about 20, less than 19, less than 18, less than 17, less than 16, less than 15, less than 14, less than 13, less than 12, less than 11, less than 10, less than 9, less than 8, less than 7, less than 6, or less than 5 amino acid residues in length. Preferably, a peptide is about 5 to 20 amino acids in length, more preferably, a peptide is about 8 to 15 amino acid residues in length.

The TCRs of the present invention bind to a WT1 peptide when presented by an MHC. As used herein, the term WT1 peptide is understood to mean a peptide comprising an amino acid sequence derived from a WT1 protein.

For example, a WT1 peptide may comprise at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, or at least 25 contiguous amino acid residues of a WT1 protein amino acid sequence.

The WT1 peptide may comprise or consist of the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions. Examples of WT1 peptides comprising the amino acid sequence are AAQWAPVLDFAPPGA (SEQ ID NO: 115) and APVLDFAPPGASAYG (SEQ ID NO: 116).

The WT1 peptide may comprise or consist of an amino acid sequence selected from the group consisting of QCLSAFTVHFSGQFT (SEQ ID NO: 118), EDPMGQQGSLGEQQY (SEQ ID NO: 119), SQLECMTNQMNLGA (SEQ ID NO: 120), and variants of SEQ ID NOs: 118-120 each having up to three amino acid substitutions, additions or deletions.

The WT1 peptide may comprise or consist of an amino acid sequence selected from the group consisting of EPASQHTLRSG (SEQ ID NO: 123), YESDNHTTPIL (SEQ ID NO: 126), and variants of SEQ ID NOs: 123 and 126 each having up to three amino acid substitutions, additions or deletions. Example WT1 peptides may have an amino acid sequence selected from the group consisting of TCVPEPASQHTLRSG (SEQ ID NO: 121), EPASQHTLRSGPGCL (SEQ ID NO: 122), HSTGYESDNHTTPIL (SEQ ID NO: 124) and YESDNHTTPILCGAQ (SEQ ID NO: 125).

The WT1 peptide may comprise or consist of the amino acid sequence of NHTTPILCGAQYRIH (SEQ ID NO: 127) or a variant thereof having up to three amino acid substitutions, additions or deletions.

The WT1 peptide may comprise or consist of the amino acid sequence of NQMNLGATLKG (SEQ ID NO: 250) or a variant thereof having up to three amino acid substitutions, additions or deletions. Example WT1 peptides may have an amino acid sequence selected from the group consisting of CMTWNQMNLGATLKG (SEQ ID NO: 248) and NQMNLGATLKGVAAG (SEQ ID NO: 249).

The WT1 peptide may comprise or consist of the amino acid sequence of DPGGIWAKLGAAEAS (SEQ ID NO: 251) or a variant thereof having up to three amino acid substitutions, additions or deletions.

The WT1 peptide may comprise or consist of an amino acid sequence selected from the group consisting of NHTTPILCGAQYRIH (SEQ ID NO: 252), KRHQRRTGTVKPFQC (SEQ ID NO: 253), PSCQKKFARSDELVR (SEQ ID NO: 254), and variants of SEQ ID NOs: 252, 253 and 254 each having up to three amino acid substitutions, additions or deletions.

In some embodiments, for WT1 peptides which bind to MHC molecules encoded by HLA-A*0201 allele it may be preferred that the amino acids at position 2 of the peptide (i.e. the second amino acid from the N-terminus) are leucine or methionine, although isoleucine, valine, alanine and threonine may also be preferable. It may also be preferred that the amino acid at position 9 or 10 is valine, leucine or isoleucine, although alanine, methionine and threonine may also be preferable. The preferred MHC binding motifs of other HLA alleles are disclosed in Celis *et al* (Molecular Immunology, Vol. 31, 8, December 1994, pages 1423 to 1430).

Various uses of the WT1 peptides described herein are contemplated by the present invention. For example, the WT1 peptides described herein may be administered to a subject, e.g. a human subject. Administration of the WT1 peptides of the present invention may elicit an immune response against cells expressing or overexpressing WT1 protein, i.e. the WT1 peptides are immunogenic WT1 peptides.

Thus in another aspect, the present invention provides an isolated immunogenic WT1 peptide comprising an amino acid sequence selected from the group consisting of EPASQHTLRSG (SEQ ID NO: 123), YESDNHTTPIL (SEQ ID NO: 126), NHTTPILCGAQYRIH (SEQ ID NO: 127), QCLSAFTVHFSGQFT (SEQ ID NO: 118), EDPMGQQGSLGEQQY (SEQ ID NO: 119), SQLECMTWNQMNLGA (SEQ ID NO: 120), APVLDFAPPGA (SEQ ID NO: 117), NQMNLGATLKG (SEQ ID NO: 250), DPGGIWAKLGAAEAS (SEQ ID NO: 251), NHTTPILCGAQYRIH (SEQ ID NO: 252), KRHQRRTGTVKPFQC (SEQ ID NO: 253), PSCQKKFARSDELVR (SEQ ID NO: 254), and variants thereof each having up to three amino acid substitutions, additions or deletions.

The WT1 peptides described herein, e.g. WT1 peptides comprising an amino acid sequence selected from the group consisting of EPASQHTLRSG (SEQ ID NO: 123) and YESDNHTTPIL (SEQ ID NO: 126), NHTTPILCGAQYRIH (SEQ ID NO: 127), QCLSAFTVHFSGQFT (SEQ ID NO: 118), EDPMGQQGSLGEQQY (SEQ ID NO: 119),
 5 SQLECMTWNQMNLGA (SEQ ID NO: 120), APVLDFAPPGA (SEQ ID NO: 117), NQMNLGATLKG (SEQ ID NO: 250), DPGGIWAKLGAAEAS (SEQ ID NO: 251), NHTTPILCGAQYRIH (SEQ ID NO: 252), KRHQRRHTGVKPFQC (SEQ ID NO: 253) PSCQKKFARSDELVR (SEQ ID NO: 254), and variants thereof each having up to three amino acid substitutions, additions or deletions, may be used to screen for and/or identify
 10 new TCR sequences which bind to WT1 cells. For example, T2 cells may be pulsed with a WT1 peptide mentioned in the present invention and incubated with a T-cell population isolated from a donor. In this approach, expression of cytokines, e.g. CD107a and IFN γ , may be indicative of T-cells which recognise WT1 peptides.

Accordingly, in one aspect, the present invention provides a T-cell receptor (TCR), which
 15 binds to a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the WT1 peptide comprises an amino acid sequence selected from the group consisting of EPASQHTLRSG (SEQ ID NO: 123), YESDNHTTPIL (SEQ ID NO: 126), NHTTPILCGAQYRIH (SEQ ID NO: 127), QCLSAFTVHFSGQFT (SEQ ID NO: 118), EDPMGQQGSLGEQQY (SEQ ID NO:
 20 119), SQLECMTWNQMNLGA (SEQ ID NO: 120), APVLDFAPPGA (SEQ ID NO: 117), NQMNLGATLKG (SEQ ID NO: 250), DPGGIWAKLGAAEAS (SEQ ID NO: 251), NHTTPILCGAQYRIH (SEQ ID NO: 252), KRHQRRHTGVKPFQC (SEQ ID NO: 253), PSCQKKFARSDELVR (SEQ ID NO: 254), and variants thereof each having up to three amino acid substitutions, additions or deletions.

25 **TCR sequences**

We have determined the amino acid sequences for TCRs that bind to WT1 peptides described herein. In particular, we have determined the amino acid sequences of the TCR CDRs, which are important for WT1 peptide recognition and binding.

Thus, in one embodiment, the present invention provides a TCR comprising a CDR3 α
 30 comprising the amino acid sequence of CGTAWINDYKLSF (SEQ ID NO: 3) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRKTGGYSNQPQHF (SEQ ID NO: 8) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID

NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Thus, in one embodiment, the present invention provides, a TCR comprising a CDR3 α comprising the amino acid sequence of CVVNLLSNQGGKLIF (SEQ ID NO: 36) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSQDYLVSNKLFF (SEQ ID NO: 41) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Thus, in one embodiment, the present invention provides, a TCR comprising a CDR3 α comprising the amino acid sequence of CAANNARLMF (SEQ ID NO: 92) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSDTRAREQFF (SEQ ID NO: 97) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Thus, in one embodiment, the present invention provides, a TCR comprising a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, the present invention provides, a TCR comprising a CDR3 α comprising the amino acid sequence of CAVEATDSWGKLQF (SEQ ID NO: 108) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSVGGSGSYNEQFF (SEQ ID NO: 169) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of NQMNLGATLKG (SEQ ID NO: 250) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, the present invention provides, a TCR comprising a CDR3 α comprising the amino acid sequence of CAVRTSYDKVIF (SEQ ID NO: 113) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSVGGSGSYNEQFF (SEQ ID NO: 169) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of NQMNLGATLKG (SEQ ID NO: 250) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Thus, in one embodiment, the present invention provides, a TCR comprising a CDR3 α comprising the amino acid sequence of CAASGGRDDKIIF (SEQ ID NO: 214) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSYSRTESTDTQYF (SEQ ID NO: 219) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

The present invention also provides a TCR comprising:

a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions;

a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions;

a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions; or

a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions;

wherein the TCR binds to a WT1 peptide comprising or consisting of an amino acid sequence selected from the group consisting of QCLSAFTVHFSGQFT (SEQ ID NO: 118), EDPMGQQGSLGEQQY (SEQ ID NO: 119), and SQLECMTWNQMNLGA (SEQ ID NO: 120) or variants thereof each having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Thus, in one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of the amino acid sequence of QCLSAFTVHFSGQFT (SEQ ID NO: 118) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of the amino acid sequence of EDPMGQQGSLGEQQY (SEQ ID NO: 119) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of the amino acid sequence of SQLECMTWNQMNLGA (SEQ ID NO: 120) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of the amino acid sequence of QCLSAFTVHFSGQFT (SEQ ID NO:

118) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of the amino acid sequence of EDPMGQQGSLGEQQY (SEQ ID NO: 119) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of the amino acid sequence of SQLECMTWNQMNLGA (SEQ ID NO: 120) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of the amino acid sequence of QCLSAFTVHFSGQFT (SEQ ID NO: 118) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of the amino acid sequence of EDPMGQQGSLGEQQY (SEQ ID NO: 119) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of the amino acid sequence of SQLECMTWNQMNLGA (SEQ ID NO: 120) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of the amino acid sequence of QCLSAFTVHFSGQFT (SEQ ID NO: 118) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of an amino acid sequence of EDPMGQQGSLGEQQY (SEQ ID NO: 119) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

In one embodiment, there is provided a TCR comprising a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising or consisting of an amino acid sequence of SQLECMTWNQMNLGA (SEQ ID NO: 120) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CATDAYSGNTPLVF (SEQ ID NO: 47) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the

amino acid sequence of CASRAAGLDTEAFF (SEQ ID NO: 57) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising comprising or consisting of an amino acid sequence of EPASQHTLRSG (SEQ ID NO: 123) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAVRAEIYNQGGLIF (SEQ ID NO: 52) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASTQTPYEQYF (SEQ ID NO: 63) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising comprising or consisting of an amino acid sequence of YESDNHTTPIL (SEQ ID NO: 126) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAVRAEIYNQGGLIF (SEQ ID NO: 52) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSTVGGEDYGYTF (SEQ ID NO: 69) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising comprising or consisting of an amino acid sequence of YESDNHTTPIL (SEQ ID NO: 126) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAASMAGAGSYQLTF (SEQ ID NO: 75) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CAISVGQGALYEQYF (SEQ ID NO: 80) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of NHTTPILCGAQYRIH (SEQ ID NO: 127) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAASMAGAGSYQLTF (SEQ ID NO: 75) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSVARDRRNYGYTF (SEQ ID NO: 86) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of NHTTPILCGAQYRIH (SEQ ID NO: 127) or a

variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAVTVGNKLVF (SEQ ID NO: 175) or a variant thereof having up to
5 three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRGWREQFF (SEQ ID NO: 180) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of DPGGIWAKLGAAEAS (SEQ ID NO: 251) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an
10 MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAARSYNTDKLIF (SEQ ID NO: 186) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSWGYQETQYF (SEQ ID NO: 196) or a variant thereof having up to
15 three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of NHTTPILCGAQYRIH (SEQ ID NO: 252) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAASYNNARLMF (SEQ ID NO: 191) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPTGGEYYGYTF (SEQ ID NO: 202) or a variant thereof having up to
20 three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of KRHQRRTGKPFQC (SEQ ID NO: 253) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.
25

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAASYNNARLMF (SEQ ID NO: 191) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSYPLRTGRYNSYNSPLHF (SEQ ID NO: 208) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of PSCQKKFARSDELVR (SEQ ID NO: 254) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.
30

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAANNARLMF (SEQ ID NO: 92) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPGQHGELEFF (SEQ ID NO: 271) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAASATGNQFYF (SEQ ID NO: 266) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSDTRAREQFF (SEQ ID NO: 97) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAASATGNQFYF (SEQ ID NO: 266) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPGQHGELEFF (SEQ ID NO: 271) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β

comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAVEATDSWGKLQF (SEQ ID NO: 108) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLGLSISQETQYF (SEQ ID NO: 288) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of NQMNLGATLKG (SEQ ID NO: 250) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAVRTSYDKVIF (SEQ ID NO: 113) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLGLSISQETQYF (SEQ ID NO: 288) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of NQMNLGATLKG (SEQ ID NO: 250) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant

thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Further provided by the present invention is a TCR comprising a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions, which binds to a WT1 peptide comprising the amino acid sequence of APVLDFAPPGA (SEQ ID NO: 117) or a variant thereof having up to three amino acid substitutions, additions or deletions when presented by an MHC.

Example TCR amino acid sequences of the present invention are provided in Table 1.

Table 1

Donor: HD1

Chain	Region	Amino acid sequence	SEQ ID NO
Alpha (α)	CDR1 α	KALYS	SEQ ID NO: 1
	CDR2 α	LLKGGEQ	SEQ ID NO: 2
	CDR3 α	CGTAWINDYKLSF	SEQ ID NO: 3
	Variable	METLLKVLSGTLLWQLTWVRSQQPVQSPQAVILREGEDAVIN CSSSKALYSVHWYRQKHGEAPVFLMILLKGGEQKGHEKISAS FNEKKQQSSLYLTASQLSYSGTYFCGTAWINDYKLSFGAGTT VTVRAN	SEQ ID NO: 4
	Full – with TRAC constant domain	METLLKVLSGTLLWQLTWVRSQQPVQSPQAVILREGEDAVIN CSSSKALYSVHWYRQKHGEAPVFLMILLKGGEQKGHEKISAS FNEKKQQSSLYLTASQLSYSGTYFCGTAWINDYKLSFGAGTT VTVRANIQNPDPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQS KDSDVYITDKTVLDMRSMDFKSNSAVAWSNKSDFACANAFNN SIIPEDTFFPSPRESSCDVKLVEKSFETDTNLFQNLVIGFR ILLKLVAGFNLLMTLRLWSS	SEQ ID NO: 5
Beta (β)	CDR1 β	SGHDY	SEQ ID NO: 6
	CDR2 β	FNNNV	SEQ ID NO: 7
	CDR3 β	CASRKTGGYSNQPQHF	SEQ ID NO: 8
	Variable	MGSWTLCCVSLCILVAKHTDAGVIQSPRHEVTEMGQEVTLRC KPI SGHDYLFWYRQTM MRGLELLIYFNNNV IDDSGMPEDRF SAKMPNASFSTLKI QPSEPRDSAVYFCASRKTGGYSNQPQHF GDGTRLSILE	SEQ ID NO: 9
	Full – with TRBC1 constant domain	MGSWTLCCVSLCILVAKHTDAGVIQSPRHEVTEMGQEVTLRC KPI SGHDYLFWYRQTM MRGLELLIYFNNNV IDDSGMPEDRF SAKMPNASFSTLKI QPSEPRDSAVYFCASRKTGGYSNQPQHF GDGTRLSILEDLNKVFPEVAVFEPSEAEISHTQKATLVCLA TGFFPDHVELS W W V N G K E V H S G V S T D P Q P L K E Q P A L N D S R Y C LSSRLRVSATFWQNPRNHFRQVQFYGLSENDEWTQDRAKPV TQIVSAEAWGRADCGFTSVSYQQGVLSATILYEILLGKATLY AVLVSAVLMLAMVVRKDF	SEQ ID NO: 10
	Full – with TRBC2 constant domain	MGSWTLCCVSLCILVAKHTDAGVIQSPRHEVTEMGQEVTLRC KPI SGHDYLFWYRQTM MRGLELLIYFNNNV IDDSGMPEDRF SAKMPNASFSTLKI QPSEPRDSAVYFCASRKTGGYSNQPQHF GDGTRLSILEDLNKVFPEVAVFEPSEAEISHTQKATLVCLA TGFFPDHVELS W W V N G K E V H S G V S T D P Q P L K E Q P A L N D S R Y C LSSRLRVSATFWQNPRNHFRQVQFYGLSENDEWTQDRAKPV TQIVSAEAWGRADCGFTSESYQQGVLSATILYEILLGKATLY	SEQ ID NO: 11

		AVLVSAVLMLAMVKRKDSRG	
--	--	----------------------	--

Donor: HD2

Clonotype	Chain	Region	Sequence	SEQ ID NO
HD2-1	Alpha (α)	CDR1α	SSVPPY	SEQ ID NO: 12
		CDR2α	Y TSAATLV	SEQ ID NO: 13
		CDR3α	CAVRLSGSARQLTF	SEQ ID NO: 14
		Variable domain	MLLLLVPVLEVI FT LGGTRAQSVTQLGSHVSV SEGALVLLRCNYSSSVPPYLFWYVQYPNQGLQ LLLKYTSAATLVK GINGFEAEFKKSETSFHLT KPSAHMSDAAEYFCAVRLSGSARQLTFGSGTQ LTVLPD	SEQ ID NO: 15
		Full – with TRAC constant domain	MLLLLVPVLEVI FT LGGTRAQSVTQLGSHVSV SEGALVLLRCNYSSSVPPYLFWYVQYPNQGLQ LLLKYTSAATLVK GINGFEAEFKKSETSFHLT KPSAHMSDAAEYFCAVRLSGSARQLTFGSGTQ LTVLPDIQNPDPAVYQLRDSKSSDKSVCLFTD FDSQTNVSQSKSDVYITDKTVLDMRSMDFKS NSAVAWSNKSDFACANAFNNSIIPEDTFFPSP ESSCDVKLVEKSFETDTNLFQNL SVIGFRIL LLKVAGFNLLMTLRLWSS	SEQ ID NO: 16
HD2-2	Alpha (α)	CDR1α	TSESDYY	SEQ ID NO: 17
		CDR2α	QEAYKQQN	SEQ ID NO: 18
		CDR3α	CAYRSLKYGNKLVF	SEQ ID NO: 19
		Variable domain	MACPGFLWALVISTCLEFSMAQTVTQSQPEMS VQEAEVTLSCTYDTSESDYYLFWYKQPPSRQ MILVIRQEAYKQQNATENRFSVNFQKA AKSFS LKISDSQLGDAAMYFCAYRSLKYGNKLVFGAG TILRVKSY	SEQ ID NO: 20
		Full – with TRAC constant domain	MACPGFLWALVISTCLEFSMAQTVTQSQPEMS VQEAEVTLSCTYDTSESDYYLFWYKQPPSRQ MILVIRQEAYKQQNATENRFSVNFQKA AKSFS LKISDSQLGDAAMYFCAYRSLKYGNKLVFGAG TILRVKSYIQNPDP AVYQLRDSKSSDKSVCLF TDFDSQTNVSQSKSDVYITDKTVLDMRSMDF KSNSAVAWSNKSDFACANAFNNSIIPEDTFFP SPESSCDVKLVEKSFETDTNLFQNL SVIGFR ILLKVAGFNLLMTLRLWSS	SEQ ID NO: 21
HD2-1β	Beta (β)	CDR1β	SGHAT	SEQ ID NO: 22
		CDR2β	FQNGV	SEQ ID NO: 23
		CDR3β	CASSLLGDEQYF	SEQ ID NO: 24
		Variable domain	MGTRLLCWAALCLLGAELTEAGVAQSPRYKII EK RQSVAFWCNPISGHATLYWYQQILGQGPKL LIQFQNGVVDDSQLPKDRFSAERLKGVDSTL KIQPAKLEDSAVYLCASSLLGDEQYFGPGTRL TVTE	SEQ ID NO: 25
		Full – with TRBC1 constant domain	MGTRLLCWAALCLLGAELTEAGVAQSPRYKII EK RQSVAFWCNPISGHATLYWYQQILGQGPKL LIQFQNGVVDDSQLPKDRFSAERLKGVDSTL KIQPAKLEDSAVYLCASSLLGDEQYFGPGTRL TVTEDLNKVPPEVAVFEPSEAEISHTQKATL VCLATGFFPDHVELSWWVNGKEVHSGVSTDPQ PLKEQPALNDSRYCLSSRLRVSATFWQNPRNH FRCQVQFYGLSENDEWTQDRAK PVTQIVSAEA WGRADCGFTSVSYQQGVLSATILYEILLGKAT LYAVLVSAVLMLAMVKRKDF	SEQ ID NO: 26
		Full – with TRBC2 constant domain	MGTRLLCWAALCLLGAELTEAGVAQSPRYKII EK RQSVAFWCNPISGHATLYWYQQILGQGPKL LIQFQNGVVDDSQLPKDRFSAERLKGVDSTL KIQPAKLEDSAVYLCASSLLGDEQYFGPGTRL TVTEDLNKVPPEVAVFEPSEAEISHTQKATL VCLATGFYPDHVELSWWVNGKEVHSGVSTDPQ PLKEQPALNDSRYCLSSRLRVSATFWQNPRNH	SEQ ID NO: 27

			FRCQVQFYGLSENDEWTQDRAKPVTQIVSAEA WGRADCGFTSESYQQGVLSATILYEILLGKAT LYAVLV SALVLMAMVKRKDSRG	
HD2-2β	Beta (β)	CDR1β	SGHTA	SEQ ID NO: 28
		CDR2β	FQNSA	SEQ ID NO: 29
		CDR3β	CASSLVALQGAGEQYF	SEQ ID NO: 30
		Variable domain	MGTRLLFWVAFCLLGADHTGAGVSQSPSNKVT EKGKDVELRCDPISGHTALYWYRQSLGQGLEF LIYFQGNSAPDKSGLPSDRFSAERTGGSVSTL TIQRTQQEDSAVYLCASSLVALQGAGEQYFGP GTRLTVTE	SEQ ID NO: 31
		Full – with TRBC1 constant domain	MGTRLLFWVAFCLLGADHTGAGVSQSPSNKVT EKGKDVELRCDPISGHTALYWYRQSLGQGLEF LIYFQGNSAPDKSGLPSDRFSAERTGGSVSTL TIQRTQQEDSAVYLCASSLVALQGAGEQYFGP GTRLTVTEDLNKVFPEVAVFEPSEAEISHTQ KATLVCLATGFFPDHVELSWWVNGKEVHSGVS TDPQPLKEQPALNDSRYCLSSRLRVSATFWQN PRNHFRQVQFYGLSENDEWTQDRAKPVTQIV SAEAWGRADCGFTSVSYQQGVLSATILYEILL GKATLYAVLV SALVLMAMVKRKDF	SEQ ID NO: 32
		Full – with TRBC2 constant domain	MGTRLLFWVAFCLLGADHTGAGVSQSPSNKVT EKGKDVELRCDPISGHTALYWYRQSLGQGLEF LIYFQGNSAPDKSGLPSDRFSAERTGGSVSTL TIQRTQQEDSAVYLCASSLVALQGAGEQYFGP GTRLTVTEDLNKVFPEVAVFEPSEAEISHTQ KATLVCLATGFYPDHVELSWWVNGKEVHSGVS TDPQPLKEQPALNDSRYCLSSRLRVSATFWQN PRNHFRQVQFYGLSENDEWTQDRAKPVTQIV SAEAWGRADCGFTSESYQQGVLSATILYEILL GKATLYAVLV SALVLMAMVKRKDSRG	SEQ ID NO: 33

Donor: HD3

Chain	Type	Sequence	SEQ ID NO
Alpha (α)	CDR1α	NSASQS	SEQ ID NO: 34
	CDR2α	VYSSGN	SEQ ID NO: 35
	CDR3α	CVVNLLSNQGGKLIF	SEQ ID NO: 36
	Variable domain	MISLRVLLVILWLQLSWVWSQRKEVEQDPGPFNVPEGATVAFNCTYS NSASQSFFWYRQDCRKEPKLLMSVYSSGNEDGRFTAQLNRASQYISL LIRDSKLSDSATYLCVVNLLSNQGGKLIFGQGTELS VKPN	SEQ ID NO: 37
	Full – with TRAC constant domain	MISLRVLLVILWLQLSWVWSQRKEVEQDPGPFNVPEGATVAFNCTYS NSASQSFFWYRQDCRKEPKLLMSVYSSGNEDGRFTAQLNRASQYISL LIRDSKLSDSATYLCVVNLLSNQGGKLIFGQGTELS VKPNIQNPDA VYQLRDSKSSDKSVCLFTDFDSQTNVSQSKSDVYITDKTVLDMRSM DFKSNSAWAWSNKSDFACANAFNNSIIPEDTFFPSPESSCDVKLVEK SFETDTNLFQNL SVIGFRILLK VAGFNLLMTLRLWSS	SEQ ID NO: 38
Beta (β)	CDR1β	LGHNA	SEQ ID NO: 39
	CDR2β	YSLEER	SEQ ID NO: 40
	CDR3β	CASSQDYLV SNEKLFF	SEQ ID NO: 41
	Variable domain	MGCRLLC A VLCLLGAGELVPMETGVTQT PRHLVMGMTNKKSLKCEQ HLGHNAMYWKQSAKKPLELMFVYSLEERVENNSVPSRFSPEC PNSS HLFLHLHTLQPEDSALYLCASSQDYLV SNEKLFFGSGTQLSVLE	SEQ ID NO: 42
	Full – with TRBC1 constant domain	MGCRLLC A VLCLLGAGELVPMETGVTQT PRHLVMGMTNKKSLKCEQ HLGHNAMYWKQSAKKPLELMFVYSLEERVENNSVPSRFSPEC PNSS HLFLHLHTLQPEDSALYLCASSQDYLV SNEKLFFGSGTQLSVLEDLN KVFPPEVAVFEPSEAEISHTQKATLVCLATGFFPDHVELSWWVNGKE VHSGVSTD P QPLKEQPALNDSRYCLSSRLRVSATFWQNPRNHFRQV QFYGLSENDEWTQDRAKPVTQIVSAEAWGRADCGFTSVSYQQGVLSA TILYEILLGKATLYAVLV SALVLMAMVKRKDF	SEQ ID NO: 43
	Full – with TRBC2 constant domain	MGCRLLC A VLCLLGAGELVPMETGVTQT PRHLVMGMTNKKSLKCEQ HLGHNAMYWKQSAKKPLELMFVYSLEERVENNSVPSRFSPEC PNSS HLFLHLHTLQPEDSALYLCASSQDYLV SNEKLFFGSGTQLSVLEDLK NVFPPEVAVFEPSEAEISHTQKATLVCLATGFYPDHVELSWWVNGKE	SEQ ID NO: 44

		VHSGVSTDPQPLKEQPALNDSRYCLSSRLRVSATFWQNPRNHFRCQV QFYGLSENDEWTQDRAKPVTQIVSAEAWGRADCGFTSESYQQGVLSA TILYEILLGKATLYAVLVLSALVLMAMVKRKDSRG	
--	--	---	--

Donor: HD4

Clonotype	Chain	Region	Sequence	SEQ ID NO
HD4-1	Alpha (α)	CDR1α	TSINN	SEQ ID NO: 45
		CDR2α	IRSNERE	SEQ ID NO: 46
		CDR3α	CATDAYSGNTPLVF	SEQ ID NO: 47
		Variable domain	METLLGVSLVILWLQLARVNSQQGEEDPQALS IQEGENATMNC SYKTSINN LQWYRQNSGRGLV HLILIRSNEREKHSGRRLRVTLDTSKKSSSLLI TASRAADTASYFCATDAYSGNTPLVFGKGTRL SVIAN	SEQ ID NO: 48
		Full – with TRAC constant domain	METLLGVSLVILWLQLARVNSQQGEEDPQALS IQEGENATMNC SYKTSINN LQWYRQNSGRGLV HLILIRSNEREKHSGRRLRVTLDTSKKSSSLLI TASRAADTASYFCATDAYSGNTPLVFGKGTRL SVIANIQNPDP AVYQLRDSKSSDKSVCLFTDF DSQTNVSQSKSDVYITDKTVLDMRSMDFKSN SAVAWSNKSD FACANAFNNSIIPEDTFFPSPE SSCDVKLVEKSFETDTNLFQNL SVIGFRILL LKVAGFNLLMTLRLWSS	SEQ ID NO: 49
HD4-2	Alpha (α)	CDR1α	DSAIYN	SEQ ID NO: 50
		CDR2α	IQSSQRE	SEQ ID NO: 51
		CDR3α	CAVRAEIYNQGGKLIF	SEQ ID NO: 52
		Variable domain	METLLGLLILWLQLQWVSSKQEV TQIPAALSV PEGENLVNCSFTDSAIYNLQWFRQDPGKGLT SLLLIQSSQREQ TSGRLNASLDKSSGRSTLYI AASQPGDSATYLC AVRAEIYNQGGKLIFGQGT ELSVKPN	SEQ ID NO: 53
		Full – with TRAC constant domain	METLLGLLILWLQLQWVSSKQEV TQIPAALSV PEGENLVNCSFTDSAIYNLQWFRQDPGKGLT SLLLIQSSQREQ TSGRLNASLDKSSGRSTLYI AASQPGDSATYLC AVRAEIYNQGGKLIFGQGT ELSVKPNIQNPDP AVYQLRDSKSSDKSVCLFT DFDSQTNVSQSKSDVYITDKTVLDMRSMDFK SNSAVAWSNKSD FACANAFNNSIIPEDTFFPS PESSCDVKLVEKSFETDTNLFQNL SVIGFRI LLLKVAGFNLLMTLRLWSS	SEQ ID NO: 54
HD4-1	Beta (β)	CDR1β	MNHNS	SEQ ID NO: 55
		CDR2β	SASEGT	SEQ ID NO: 56
		CDR3β	CASRAAGLDTEAFF	SEQ ID NO: 57
		Variable domain	MSIGLLCCVAFSLLWASPVNAGVTQTPKFQVL KTGQSM TLQCAQDMNHNSMYWYRQDPGMGLRL IYY SASEGTTDKGEVPNGYNVSRLNKREFSLR LESAAPSQTSVYFCASRAAGLDTEAFFGQGTR LTVVE	SEQ ID NO: 58
		Full – with TRBC1 constant domain	MSIGLLCCVAFSLLWASPVNAGVTQTPKFQVL KTGQSM TLQCAQDMNHNSMYWYRQDPGMGLRL IYY SASEGTTDKGEVPNGYNVSRLNKREFSLR LESAAPSQTSVYFCASRAAGLDTEAFFGQGTR LTVVEDLNKVFPP EVAVFEPSEAEISHTQKAT LVCLATGF FPDHVELSWVNGKEVHSGVSTDP QPLKEQPALNDSRYCLSSRLRV SATFWQNPRN HFRCQVQFYGLSENDEWTQDRAKPVTQIVSAE AWGRADCGFTSVSYQQGVLSATILYEILLGKA TLYAVLV SALVLMAMVKRKDF	SEQ ID NO: 59
		Full – with TRBC2 constant domain	MSIGLLCCVAFSLLWASPVNAGVTQTPKFQVL KTGQSM TLQCAQDMNHNSMYWYRQDPGMGLRL IYY SASEGTTDKGEVPNGYNVSRLNKREFSLR LESAAPSQTSVYFCASRAAGLDTEAFFGQGTR	SEQ ID NO: 60

			LTVVEDLKNVFPPEVAVFEPSEAEISHTQKAT LVCLATGFYPDHVELSWWVNGKEVHSGVSTDP QPLKEQPALNDSRYCLSSRLRVSATFWQNPRN HFRQCQVQFYGLSENDEWTQDRAKPVTQIVSAE AWGRADCGFTSESYQQGVLSATILYEILLGKA TLYAVLVSALVLMAMVKRKDSRG	
HD4-2	Beta (β)	CDR1β	MNHNY	SEQ ID NO: 61
		CDR2β	SVGAGI	SEQ ID NO: 62
		CDR3β	CASTQTPYEQYF	SEQ ID NO: 63
		Variable domain	MSISLLCCAAFP LLWAGPVNAGVTQTPKFRIL KIGQSMTLQCTQDMNHNYMYWYRQDPGMGLKL IYYSVGAGITDKGEVPNGYNVSRSTTEDFPLR LELAAPSQTSVYFCASTQTPYEQYFGPGTRLT VTE	SEQ ID NO: 64
		Full – with TRBC1 constant domain	MSISLLCCAAFP LLWAGPVNAGVTQTPKFRIL KIGQSMTLQCTQDMNHNYMYWYRQDPGMGLKL IYYSVGAGITDKGEVPNGYNVSRSTTEDFPLR LELAAPSQTSVYFCASTQTPYEQYFGPGTRLT VTEDLNKVFPPEVAVFEPSEAEISHTQKATLV CLATGFFPDHVELSWWVNGKEVHSGVSTDPQP LKEQPALNDSRYCLSSRLRVSATFWQNPRNHF RCQVQFYGLSENDEWTQDRAKPVTQIVSAEAW GRADCGFTSVSYQQGVLSATILYEILLGKATL YAVLVSALVLMAMVKRKDF	SEQ ID NO: 65
		Full – with TRBC2 constant domain	MSISLLCCAAFP LLWAGPVNAGVTQTPKFRIL KIGQSMTLQCTQDMNHNYMYWYRQDPGMGLKL IYYSVGAGITDKGEVPNGYNVSRSTTEDFPLR LELAAPSQTSVYFCASTQTPYEQYFGPGTRLT VTEDLKNVFPPEVAVFEPSEAEISHTQKATLV CLATGFYPDHVELSWWVNGKEVHSGVSTDPQP LKEQPALNDSRYCLSSRLRVSATFWQNPRNHF RCQVQFYGLSENDEWTQDRAKPVTQIVSAEAW GRADCGFTSESYQQGVLSATILYEILLGKATL YAVLVSALVLMAMVKRKDSRG	SEQ ID NO: 66
HD4-3	Beta (β)	CDR1β	SGHNS	SEQ ID NO: 67
		CDR2β	FNNNVP	SEQ ID NO: 68
		CDR3β	CASSTVGGEDYGYTF	SEQ ID NO: 69
		Variable domain	MDSWTFCCVSLCILVAKHTDAGVIQSPRHEVT EMGQEVTLRCKPISGHNSLFWYRQTMMRGLEL LIYFNNNVPIDDSGMPEDRFSAKMPNASFSTL KIQPSEPRDSAVYFCASSTVGGEDYGYTFGSG TRLTVVE	SEQ ID NO: 70
		Full – with TRBC1 constant domain	MDSWTFCCVSLCILVAKHTDAGVIQSPRHEVT EMGQEVTLRCKPISGHNSLFWYRQTMMRGLEL LIYFNNNVPIDDSGMPEDRFSAKMPNASFSTL KIQPSEPRDSAVYFCASSTVGGEDYGYTFGSG TRLTVVEDLNKVFPPEVAVFEPSEAEISHTQK ATLVCLATGFFPDHVELSWWVNGKEVHSGVST DPQPLKEQPALNDSRYCLSSRLRVSATFWQNP RNHFRQCQVQFYGLSENDEWTQDRAKPVTQIVS AEAWGRADCGFTSVSYQQGVLSATILYEILLG KATLYAVLVSALVLMAMVKRKDF	SEQ ID NO: 71
		Full – with TRBC2 constant domain	MDSWTFCCVSLCILVAKHTDAGVIQSPRHEVT EMGQEVTLRCKPISGHNSLFWYRQTMMRGLEL LIYFNNNVPIDDSGMPEDRFSAKMPNASFSTL KIQPSEPRDSAVYFCASSTVGGEDYGYTFGSG TRLTVVEDLKNVFPPEVAVFEPSEAEISHTQK ATLVCLATGFYPDHVELSWWVNGKEVHSGVST DPQPLKEQPALNDSRYCLSSRLRVSATFWQNP RNHFRQCQVQFYGLSENDEWTQDRAKPVTQIVS AEAWGRADCGFTSESYQQGVLSATILYEILLG KATLYAVLVSALVLMAMVKRKDSRG	SEQ ID NO: 72

Donor: HD5

Clonotype	Chain	Region	Sequence	SEQ ID NO
HD5-1	Alpha (α)	CDR1α	DSASNY	SEQ ID NO: 73
		CDR2α	IRSNVGE	SEQ ID NO: 74
		CDR3α	CAASMAGAGSYQLTF	SEQ ID NO: 75
		Variable domain	MTSIRAVFIFLWLQLDLVNGENVEQHPSTLSV QEGDSAVIKCTYSDSASNYFPWYKQELGKRPO LIIDIRSNVGEKKDQRIAVTLNKTAKHFSLHI TETQPEDSAVYFCAASMAGAGSYQLTFGKGTK LSVIPN	SEQ ID NO: 76
		Full – with TRAC constant domain	MTSIRAVFIFLWLQLDLVNGENVEQHPSTLSV QEGDSAVIKCTYSDSASNYFPWYKQELGKRPO LIIDIRSNVGEKKDQRIAVTLNKTAKHFSLHI TETQPEDSAVYFCAASMAGAGSYQLTFGKGTK LSVIPNIQNPDPAVYQLRDSKSSDKSVCLFTD FDSQTNVSQSKSDVYITDKTVLDMRSMDFKS NSAVAWSNKSDFACANAFNNSIIPEDTFFPSP ESSCDVKLVEKSFETDTNLFQNLVIGFRIL LLKVAGFNLLMTLRLWSS	SEQ ID NO: 77
HD5-1	Beta (β)	CDR1β	ENHRY	SEQ ID NO: 78
		CDR2β	SYGVKD	SEQ ID NO: 79
		CDR3β	CAISVGQGALYEQYF	SEQ ID NO: 80
		Variable domain	MGTRLFFFYVALCLLWTGHMDAGITQSPRHKVT ETGTPVTLRCHQ TENHRYMYWYRQDPGHGLRL IHYSYGVKDTDKGEVSDGYSVSRSKTEDFLLT LESATSSQTSVYFCAISVGQGALYEQYFGPGT RLTVTE	SEQ ID NO: 81
		Full – with TRBC1 constant domain	MGTRLFFFYVALCLLWTGHMDAGITQSPRHKVT ETGTPVTLRCHQ TENHRYMYWYRQDPGHGLRL IHYSYGVKDTDKGEVSDGYSVSRSKTEDFLLT LESATSSQTSVYFCAISVGQGALYEQYFGPGT RLTVTEDLNKVFPEVAVFEPSEAEISHTQKA TLVCLATGFFPDHVELSWWVNGKEVHSGVSTD PQPLKEQPALNDSRYCLSSRLRVSATFWQNPR NHFRQCQVQFYGLSENDEWTQDRAKPVTQIVSA EAWGRADCGFTSVSYQQGVLSATILYEILLGK ATLYAVLVSAVLMLMAMVKRKDF	SEQ ID NO: 82
		Full – with TRBC2 constant domain	MGTRLFFFYVALCLLWTGHMDAGITQSPRHKVT ETGTPVTLRCHQ TENHRYMYWYRQDPGHGLRL IHYSYGVKDTDKGEVSDGYSVSRSKTEDFLLT LESATSSQTSVYFCAISVGQGALYEQYFGPGT RLTVTEDLNKVFPEVAVFEPSEAEISHTQKA TLVCLATGFYPDHVELSWWVNGKEVHSGVSTD PQPLKEQPALNDSRYCLSSRLRVSATFWQNPR NHFRQCQVQFYGLSENDEWTQDRAKPVTQIVSA EAWGRADCGFTSESYQQGVLSATILYEILLGK ATLYAVLVSAVLMLMAMVKRKDSRG	SEQ ID NO: 83
HD5-2	Beta (β)	CDR1β	SGDLS	SEQ ID NO: 84
		CDR2β	YYNGEE	SEQ ID NO: 85
		CDR3β	CASSVARDRRNYGYTF	SEQ ID NO: 86
		Variable domain	MGFRLCCVAFCLLGAGPVD SGVTQTPKHLIT ATGQRVTLRCS PRSGDLSVYWYQQSLDQGLQF LIQYYNGEERAKGNILERFSAQQFPDLHSELN LSSLELGDSALYFCASSVARDRRNYGYTFGSG TRLTVVE	SEQ ID NO: 87
		Full – with TRBC1 constant domain	MGFRLCCVAFCLLGAGPVD SGVTQTPKHLIT ATGQRVTLRCS PRSGDLSVYWYQQSLDQGLQF LIQYYNGEERAKGNILERFSAQQFPDLHSELN LSSLELGDSALYFCASSVARDRRNYGYTFGSG TRLTVVEDLNKVFPEVAVFEPSEAEISHTQK ATLVCLATGFFPDHVELSWWVNGKEVHSGVST DPQPLKEQPALNDSRYCLSSRLRVSATFWQNP RNHFRQCQVQFYGLSENDEWTQDRAKPVTQIVS AEAWGRADCGFTSVSYQQGVLSATILYEILLG KATLYAVLVSAVLMLMAMVKRKDF	SEQ ID NO: 88
		Full – with TRBC2	MGFRLCCVAFCLLGAGPVD SGVTQTPKHLIT	SEQ ID NO: 89

		constant domain	ATGQRVTLRCSPRSGDLSVYWYQQSLDQGLQF LIQYYNGEERAKGNILERFSAQQFPDLHSELN LSSLELGDSALYFCASSVARDRRNYGYTFGSG TRLTVVEDLKNVFPPEVAVFEPSEAEISHTQK ATLVCLATGFYPDHVELSWWVNGKEVHSGVST DPQPLKEQPALNDSRYCLSSRLRVSATFWQNP RNHFRQCQVQFYGLSENDEWTQDRAKPVTQIVS AEAWGRADCGFTSESYQQGVLSATILYEILLG KATLYAVLVSALVLMAMVKRKDSRG	
--	--	------------------------	--	--

Donor: HD6

Clonotype	Chain	Region	Amino acid sequence	SEQ ID NO
HD6-1	Alpha (α)	CDR1α	NSMFDY	SEQ ID NO: 90
		CDR2α	ISSIKDK	SEQ ID NO: 91
		CDR3α	CAANNARLMF	SEQ ID NO: 92
		Variable domain	MAMLLGASVLILWLQPDWVNSQQKNDDQQVKQ NSPSLSVQEGRISILNCDYTNSMFDYFLWYKK YPAEGPTFLISISSIKDKNEDGRFTVFLNKSA KHLSLHIVPSQPGDSAVYFCAANNARLMFGDG TQLVVKPN	SEQ ID NO: 93
		Full – with TRAC constant domain	MAMLLGASVLILWLQPDWVNSQQKNDDQQVKQ NSPSLSVQEGRISILNCDYTNSMFDYFLWYKK YPAEGPTFLISISSIKDKNEDGRFTVFLNKSA KHLSLHIVPSQPGDSAVYFCAANNARLMFGDG TQLVVKPNIQNPDAVYQLRDSKSSDKSVCLF TDFDSQTNVSQSKSDVYITDKTVLDMRSMDF KSNSAVAWSNKSDFACANAFNNSIIPEDTFFP SPESSCDVKLVEKSFETDTNLFQNLVIGFR ILLKLVAGFNLLMTLRLWSS	SEQ ID NO: 94
HD6-2	Alpha (α)	CDR1α	NSMFDY	SEQ ID NO: 264
		CDR2α	ISSIKDK	SEQ ID NO: 265
		CDR3α	CAASATGNQFYF	SEQ ID NO: 266
		Variable domain	MAMLLGASVLILWLQPDWVNSQQKNDDQQVKQ NSPSLSVQEGRISILNCDYTNSMFDYFLWYKK YPAEGPTFLISISSIKDKNEDGRFTVFLNKSA KHLSLHIVPSQPGDSAVYFCAASATGNQFYFG TGTSLTVIPN	SEQ ID NO: 267
		Full – with TRAC constant domain	MAMLLGASVLILWLQPDWVNSQQKNDDQQVKQ NSPSLSVQEGRISILNCDYTNSMFDYFLWYKK YPAEGPTFLISISSIKDKNEDGRFTVFLNKSA KHLSLHIVPSQPGDSAVYFCAASATGNQFYFG TGTSLTVIPNIQNPDAVYQLRDSKSSDKSVC LFTDFDSQTNVSQSKSDVYITDKTVLDMRSM DFKSNSAVAWSNKSDFACANAFNNSIIPEDTF FPSPSSCDVKLVEKSFETDTNLFQNLVIG FRILLKLVAGFNLLMTLRLWSS	SEQ ID NO: 268
HD6-1	Beta (β)	CDR1β	SGHNS	SEQ ID NO: 95
		CDR2β	FNNNVP	SEQ ID NO: 96
		CDR3β	CASSDTRAREQFF	SEQ ID NO: 97
		Variable domain	MDSWTFCCVSLCILVAKHTDAGVIQSPRHEVT EMGQEVTLRCKPISGHNSLFWYRQTMMRGLEL LIYFNNNVPIDDSGMPEDRFSKMPNASFSTL KIQPSEPRDSAVYFCASSDTRAREQFFGPGTR LTVLE	SEQ ID NO: 98
		Full – with TRBC1 constant domain	MDSWTFCCVSLCILVAKHTDAGVIQSPRHEVT EMGQEVTLRCKPISGHNSLFWYRQTMMRGLEL LIYFNNNVPIDDSGMPEDRFSKMPNASFSTL KIQPSEPRDSAVYFCASSDTRAREQFFGPGTR LTVLEDLNKVFPEVAVFEPSEAEISHTQKAT LVCLATGFFPDHVELSWWVNGKEVHSGVSTDP QPLKEQPALNDSRYCLSSRLRVSATFWQNPRN HFRQCQVQFYGLSENDEWTQDRAKPVTQIVSAE AWGRADCGFTSVSYQQGVLSATILYEILLGKA	SEQ ID NO: 99

			TLYAVLVSAVLMLMAMVKRKDF	
		Full – with TRBC2 constant domain	MDSWTFCCVSLCILVAKHTDAGVIQSPRHEVT EMGQEVTLRCKPISGHNSLFWYRQTMMRGLEL LIYFNNNVPIDDSGMPEDRFSAKMPNASFSTL KIQPSEPRDSAVYFCASSDTRAREQFFGPGTR LTVLEDLKNVFPPEVAVFEPSEAEISHTQKAT LVCLATGFYPDHVELSWWVNGKEVHSGVSTDP QPLKEQPALNDSRYCLSSRLRVSATFWQNPRN HFRCQVQFYGLSENDEWTQDRAKPVQTQIVSAE AWGRADCGFTSESYQQGVLSATILYEILLGKA TLYAVLVSAVLMLMAMVKRKDSRG	SEQ ID NO: 100
HD6-2	Beta (β)	CDR1β	SGHRS	SEQ ID NO: 269
		CDR2β	YFSETQ	SEQ ID NO: 270
		CDR3β	CASSPGQHGELEFF	SEQ ID NO: 271
		Variable domain	MGSRLLCWVLLCLLGAGPVKAGVTQTTPRYLIK TRGQQVTLSCSPI SGHRSSVSWYQQTPGQGLQF LFEYFSETQRNKGNFPGRFSGRQFSNSRSEM NSTLELGDSALYLCASSPGQHGELEFFGEGSRL TVLE	SEQ ID NO: 272
		Full – with TRBC1 constant domain	MGSRLLCWVLLCLLGAGPVKAGVTQTTPRYLIK TRGQQVTLSCSPI SGHRSSVSWYQQTPGQGLQF LFEYFSETQRNKGNFPGRFSGRQFSNSRSEM NSTLELGDSALYLCASSPGQHGELEFFGEGSRL TVLEDLNKVFPEVAVFEPSEAEISHTQKATL VCLATGFFPDHVELSWWVNGKEVHSGVSTDPQ PLKEQPALNDSRYCLSSRLRVSATFWQNPRNH FRCQVQFYGLSENDEWTQDRAKPVQTQIVSAEA WGRADCGFTSVSYQQGVLSATILYEILLGKAT LYAVLVSAVLMLMAMVKRKDF	SEQ ID NO: 273
		Full – with TRBC2 constant domain	MGSRLLCWVLLCLLGAGPVKAGVTQTTPRYLIK TRGQQVTLSCSPI SGHRSSVSWYQQTPGQGLQF LFEYFSETQRNKGNFPGRFSGRQFSNSRSEM NSTLELGDSALYLCASSPGQHGELEFFGEGSRL TVLEDLKNVFPPEVAVFEPSEAEISHTQKATL VCLATGFYPDHVELSWWVNGKEVHSGVSTDPQ PLKEQPALNDSRYCLSSRLRVSATFWQNPRNH FRCQVQFYGLSENDEWTQDRAKPVQTQIVSAEA WGRADCGFTSESYQQGVLSATILYEILLGKAT LYAVLVSAVLMLMAMVKRKDSRG	SEQ ID NO: 274

Donor: HD7

Clonotype	Chain	Region	Sequence	SEQ ID NO
HD7-1	Alpha (α)	CDR1α	DSSSTY	SEQ ID NO: 101
		CDR2α	IFSNMDM	SEQ ID NO: 102
		CDR3α	CAERLNTDKLIF	SEQ ID NO: 103
		Variable domain	MKTFAGFSFLFLWLQLDCMSRGEDVEQSLFLS VREGDSSVINCTYTDSSSTYLYWKQEPGAGL QLLTYIFSNMDMKQDQRLTVLLNKKDKHLSLR IADTQTGDSAIYFCAERLNTDKLIFGTGTRLQ VFPN	SEQ ID NO: 104
		Full – with TRAC constant domain	MKTFAGFSFLFLWLQLDCMSRGEDVEQSLFLS VREGDSSVINCTYTDSSSTYLYWKQEPGAGL QLLTYIFSNMDMKQDQRLTVLLNKKDKHLSLR IADTQTGDSAIYFCAERLNTDKLIFGTGTRLQ VFPNIQNPDPAVYQLRDSKSSDKSVCLFTDFD SQTNVSQSKSDVYITDKTVLDMRSMDFKSNS AWAWSNKSDFACANAFNNSIIPEDTFFPSPES SCDVKLVEKSFETDTNLFQNLVIGFRILL KVAGFNLLMTLRLWSS	SEQ ID NO: 105
HD7-2	Alpha (α)	CDR1α	DSVNN	SEQ ID NO: 106
		CDR2α	IPSGT	SEQ ID NO: 107
		CDR3α	CAVEATDSWGKLQF	SEQ ID NO: 108
		Variable domain	MKRILGALLGLLSAQVCCVRGIQVEQSPPDLI	SEQ ID NO: 109

			LQEGANSTLRCNFSDSVNNLQWFHQNPWGQLI NLFYIPSGTKQNGRLSATTVATERYSLLYISS SQTTDSGVYFCAVEATDSWGKLQFGAGTQVVV TPD	
		Full – with TRAC constant domain	MKRILGALLGLLSAQVCCVRGIQVEQSPPDLI LQEGANSTLRCNFSDSVNNLQWFHQNPWGQLI NLFYIPSGTKQNGRLSATTVATERYSLLYISS SQTTDSGVYFCAVEATDSWGKLQFGAGTQVVV TPDIQNPDPVYQLRDSKSSDKSVCLFTDFDS QTNVSQSKSDVYITDKTVLDMRSMDFKSNSA VAWSNKSDFACANAFNNSIIPEDTFFPSPESS CDVKLVEKSFETDTNLFQNLQNLVIGFRILLK VAGFNLLMTLRLWSS	SEQ ID NO: 110
HD7-3	Alpha (α)	CDR1α	DSASNY	SEQ ID NO: 111
		CDR2α	IRSNVGE	SEQ ID NO: 112
		CDR3α	CAVRTSYDKVIF	SEQ ID NO: 113
		Variable domain	MTSIRAVFIFLWLQLDLVNGENVEQHPSTLSV QEGDSAVIKCTYSDSASNYFPWYKQELGKRPO LIIDIRSNVGEKKDQRIAVTLNKTAKHFSLHI TETQPEDSAVYFCAVRTSYDKVIFGPGTSLSV IPN	SEQ ID NO: 114
		Full – with TRAC constant domain	MTSIRAVFIFLWLQLDLVNGENVEQHPSTLSV QEGDSAVIKCTYSDSASNYFPWYKQELGKRPO LIIDIRSNVGEKKDQRIAVTLNKTAKHFSLHI TETQPEDSAVYFCAVRTSYDKVIFGPGTSLSV IPNIQNPDPVYQLRDSKSSDKSVCLFTDFDS QTNVSQSKSDVYITDKTVLDMRSMDFKSNSA VAWSNKSDFACANAFNNSIIPEDTFFPSPESS CDVKLVEKSFETDTNLFQNLQNLVIGFRILLK VAGFNLLMTLRLWSS	SEQ ID NO: 160
HD7-4	Alpha (α)	CDR1α	TSINN	SEQ ID NO: 275
		CDR2α	IRSNERE	SEQ ID NO: 276
		CDR3α	CATDGDSSYKLIF	SEQ ID NO: 277
		Variable domain	METLLGVSLVILWLQLARVNSQQGEEDPQALS IQEGENATMNCSYKTSINNLOWYRQNSGRGLV HLILIRSNEREKHSGRRLRVTLDTSKKSSSLI TASRAADTASYFCATDGDSSYKLIFGSGTRLL VRPD	SEQ ID NO: 278
		Full – with TRAC constant domain	METLLGVSLVILWLQLARVNSQQGEEDPQALS IQEGENATMNCSYKTSINNLOWYRQNSGRGLV HLILIRSNEREKHSGRRLRVTLDTSKKSSSLI TASRAADTASYFCATDGDSSYKLIFGSGTRLL VRPDIQNPDPVYQLRDSKSSDKSVCLFTDFD SQTNVSQSKSDVYITDKTVLDMRSMDFKSNS AVAWSNKSDFACANAFNNSIIPEDTFFPSPESS SCDVKLVEKSFETDTNLFQNLQNLVIGFRILLK KVAGFNLLMTLRLWSS	SEQ ID NO: 279
HD7-1	Beta (β)	CDR1β	DFQATT	SEQ ID NO: 161
		CDR2β	SNEGSKA	SEQ ID NO: 162
		CDR3β	CSARDSVSGNTIYF	SEQ ID NO: 163
		Variable domain	MLLLLLLLGPGISLLPGSLAGSGLGAVVSQH PSWVICKSGTSVKIECRSLDFQATTMFWYRQF PKQSLMLMATSNEGSKATYEQGVEKDKFLINH ASLTTLSTLTVTSAHPEDSSFYICARDSVSGN TIYFGECSWLTVE	SEQ ID NO: 164
		Full – with TRBC1 constant domain	MLLLLLLLGPGISLLPGSLAGSGLGAVVSQH PSWVICKSGTSVKIECRSLDFQATTMFWYRQF PKQSLMLMATSNEGSKATYEQGVEKDKFLINH ASLTTLSTLTVTSAHPEDSSFYICARDSVSGN TIYFGECSWLTVEEDLNKVPPEVAVFEPSEA EISHTQKATLVCLATGFFPDHVELSWWVNGKE VHSGVSTDPOPLKEQPALNDSRYCLSSRLRVS ATFWQNPRNHFRQCQVQFYGLSENDEWTQDRAK PVTQIVSAEAWGRADCGFTSVSYQQGVLSATI	SEQ ID NO: 165

			LYEILLGKATLYAVLVSAVLMLAMVKKRDF	
		Full – with TRBC2 constant domain	MLLLLLLLGPGISLLLPGSLAGSGLGAVVSQH PSWVICKSGT SVKIECRSLDFQATTMFWYRQF PKQSLMLMATSNEGSKATYEQGVEKDKFLINH ASLTTLSTLTVTSAHPEDSSFYICSARDSVSGN TIYFGECSWLT VVEDLKNVFPPEVAVFEPSEA EISHTQKATLVCLATGFYDPDHVELSWWVNGKE VHSGVSTDPQPLKEQPALNDSRYCLSSRLRVS ATFWQNPRNHFRCQVQFYGLSENDEWTQDRAK PVTQIVSAEAWGRADCGFTSESYQQGVLSATI LYEILLGKATLYAVLVSAVLMLAMVKKRDSRG	SEQ ID NO: 166
HD7-2	Beta (β)	CDR1β	SQVTM	SEQ ID NO: 167
		CDR2β	ANQGSEA	SEQ ID NO: 168
		CDR3β	CSVGGSGSYNEQFF	SEQ ID NO: 169
		Variable domain	MLLLLLLLGGLGVSFSAVISQKPSRDICQRT SLTIQCQVDSQVTMMFWYRQQPGQSLTLIATA NQGSEATYESGFVIDKFPISRPNLTFSTLTVS NMSPEDSSIYLC SVGGSGSYNEQFFGPGTRLT VLE	SEQ ID NO: 170
		Full – with TRBC1 constant domain	MLLLLLLLGGLGVSFSAVISQKPSRDICQRT SLTIQCQVDSQVTMMFWYRQQPGQSLTLIATA NQGSEATYESGFVIDKFPISRPNLTFSTLTVS NMSPEDSSIYLC SVGGSGSYNEQFFGPGTRLT VLEDLNKVFPEVAVFEPSEAEISHTQKATLV CLATGFFPDHVELSWWVNGKEVHSGVSTDPQP LKEQPALNDSRYCLSSRLRVSATFWQNPRNHF RCQVQFYGLSENDEWTQDRAK PVTQIVSAEAW GRADCGFTSVSYQQGVLSATILYEILLGKATL YAVLVSAVLMLAMVKKRDF	SEQ ID NO: 171
		Full – with TRBC2 constant domain	MLLLLLLLGGLGVSFSAVISQKPSRDICQRT SLTIQCQVDSQVTMMFWYRQQPGQSLTLIATA NQGSEATYESGFVIDKFPISRPNLTFSTLTVS NMSPEDSSIYLC SVGGSGSYNEQFFGPGTRLT VLEDLNKVFPEVAVFEPSEAEISHTQKATLV CLATGFYDPDHVELSWWVNGKEVHSGVSTDPQP LKEQPALNDSRYCLSSRLRVSATFWQNPRNHF RCQVQFYGLSENDEWTQDRAK PVTQIVSAEAW GRADCGFTSESYQQGVLSATILYEILLGKATL YAVLVSAVLMLAMVKKRDSRG	SEQ ID NO: 172
HD7-3	Beta (β)	CDR1β	DFQATT	SEQ ID NO: 280
		CDR2β	SNEGSKA	SEQ ID NO: 281
		CDR3β	CSARDVLTGDYGYTF	SEQ ID NO: 282
		Variable domain	MLLLLLLLGPGISLLLPGSLAGSGLGAVVSQH PSWVICKSGT SVKIECRSLDFQATTMFWYRQF PKQSLMLMATSNEGSKATYEQGVEKDKFLINH ASLTTLSTLTVTSAHPEDSSFYICSARDVLTGD YGYTFGSGTRLT VV	SEQ ID NO: 283
		Full – with TRBC1 constant domain	MLLLLLLLGPGISLLLPGSLAGSGLGAVVSQH PSWVICKSGT SVKIECRSLDFQATTMFWYRQF PKQSLMLMATSNEGSKATYEQGVEKDKFLINH ASLTTLSTLTVTSAHPEDSSFYICSARDVLTGD YGYTFGSGTRLT VVEDLNKVFPEVAVFEPSE AEISHTQKATLVCLATGFFPDHVELSWWVNGK EVHSGVSTDPQPLKEQPALNDSRYCLSSRLRV SATFWQNPRNHFRCQVQFYGLSENDEWTQDRA KPVTQIVSAEAWGRADCGFTSVSYQQGVLSAT ILYEILLGKATLYAVLVSAVLMLAMVKKRDF	SEQ ID NO: 284
		Full – with TRBC2 constant domain	MLLLLLLLGPGISLLLPGSLAGSGLGAVVSQH PSWVICKSGT SVKIECRSLDFQATTMFWYRQF PKQSLMLMATSNEGSKATYEQGVEKDKFLINH ASLTTLSTLTVTSAHPEDSSFYICSARDVLTGD YGYTFGSGTRLT VVEDLNKVFPEVAVFEPSE AEISHTQKATLVCLATGFYDPDHVELSWWVNGK EVHSGVSTDPQPLKEQPALNDSRYCLSSRLRV SATFWQNPRNHFRCQVQFYGLSENDEWTQDRA	SEQ ID NO: 285

			KPVTQIVSAEAWGRADCGFTSESYQQGVLSAT ILYEILLGKATLYAVLVSALVLMAMVKRKDSR G	
HD7-4	Beta (β)	CDR1β	SGHDY	SEQ ID NO: 286
		CDR2β	FNNNVP	SEQ ID NO: 287
		CDR3β	CASSLGLSISQETQYF	SEQ ID NO: 288
		Variable domain	MGSWTLCCVSLCILVAKHTDAGVIQSPRHEVT EMGQEVTLRCKPISGHDYLFWYRQTMMRGLEL LIYFNNNVPIDDSGMPEDRFSAKMPNASFSTL KIQPSEPRDSAVYFCASSLGLSISQETQYFGP GTRLLVLE	SEQ ID NO: 289
		Full – with TRBC1 constant domain	MGSWTLCCVSLCILVAKHTDAGVIQSPRHEVT EMGQEVTLRCKPISGHDYLFWYRQTMMRGLEL LIYFNNNVPIDDSGMPEDRFSAKMPNASFSTL KIQPSEPRDSAVYFCASSLGLSISQETQYFGP GTRLLVLEDLNKVFPEVAVFEPSEAEISHTQ KATLVCLATGFFPDHVELSWWVNGKEVHSGVS TDPQPLKEQPALNDSRYCLSSRLRVSATFWQN PRNHFRQCQVQFYGLSENDEWTQDRAKPV TQIVSAEAWGRADCGFTSVSYQQGVLSATILYEILL GKATLYAVLVSALVLMAMVKRKDF	SEQ ID NO: 290
		Full – with TRBC2 constant domain	MGSWTLCCVSLCILVAKHTDAGVIQSPRHEVT EMGQEVTLRCKPISGHDYLFWYRQTMMRGLEL LIYFNNNVPIDDSGMPEDRFSAKMPNASFSTL KIQPSEPRDSAVYFCASSLGLSISQETQYFGP GTRLLVLEDLNKVFPEVAVFEPSEAEISHTQ KATLVCLATGFYPDHVELSWWVNGKEVHSGVS TDPQPLKEQPALNDSRYCLSSRLRVSATFWQN PRNHFRQCQVQFYGLSENDEWTQDRAKPV TQIVSAEAWGRADCGFTSESYQQGVLSATILYEILL GKATLYAVLVSALVLMAMVKRKDSRG	SEQ ID NO: 291

Donor: HD8

Chain	Region	Amino acid sequence	SEQ ID NO
Alpha (α)	CDR1α	VGISA	SEQ ID NO: 173
	CDR2α	LSSGK	SEQ ID NO: 174
	CDR3α	CAVTVGNKLVF	SEQ ID NO: 175
	Variable domain	MVKIRQFLLAILWLQ LSCVSAAKNEVEQSPQNLTAQEGEFIT INCSYSVGISALHWLQ QHPGGGIVSLFMLSSGKKKHGRLIAT INIQEKHSSLHITASHPRDSAVYICAVTVGNKLVFGAGTILR VKSY	SEQ ID NO: 176
	Full – with TRAC constant domain	MVKIRQFLLAILWLQ LSCVSAAKNEVEQSPQNLTAQEGEFIT INCSYSVGISALHWLQ QHPGGGIVSLFMLSSGKKKHGRLIAT INIQEKHSSLHITASHPRDSAVYICAVTVGNKLVFGAGTILR VKS YIQNPDPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKD SDVYITDKTVLDMRSMDFKSN SAVAWSNKSDFACANAFNNSI IPEDTFFPSPESSCDVKLVEKSFETDTNLFQNL SVIGFRIL LLKVAGFNLLMTLRLWSS	SEQ ID NO: 177
Beta (β)	CDR1β	MNHNS	SEQ ID NO: 178
	CDR2β	SASEGT	SEQ ID NO: 179
	CDR3β	CASRGWREQFF	SEQ ID NO: 180
	Variable domain	MSIGLLCCVAFSLLWASPVNAGVTQTPKFQVLKTGQSMTLQC AQDMNHNSMYWYRQDPGMGLRLIYY SASEGTTDKGEVPNGYN VSRLNKREFSLRLESAAPSQTSVYFCASRGWREQFFGPGTRL TVLE	SEQ ID NO: 181
	Full – with TRBC1 constant domain	MSIGLLCCVAFSLLWASPVNAGVTQTPKFQVLKTGQSMTLQC AQDMNHNSMYWYRQDPGMGLRLIYY SASEGTTDKGEVPNGYN VSRLNKREFSLRLESAAPSQTSVYFCASRGWREQFFGPGTRL TVLEDLNKVFPEVAVFEPSEAEISHTQKATLVCLATGFFPD HVELSWWVNGKEVHSGVSTDPQPLKEQPALNDSRYCLSSRLR VSATFWQNPRNHFRQCQVQFYGLSENDEWTQDRAKPV TQIVSAEAWGRADCGFTSVSYQQGVLSATILYEILLGKATLYAVLVSA LVLAMVKRKDF	SEQ ID NO: 182

	Full – with TRBC2 constant domain	MSIGLLCCVAFSLLWASPVNAGVTQTPKFQVLKTGQSMTLQC AQDMNHNSMYWYRQDPGMGLRLIYYSASEGTTDKGEVPNGYN VSRLNKREFSLRLESAAPSQTSVYFCASRGWREQFFGPGTRL TVLEDLKNVFPPEVAVFEPSEAEISHTQKATLVCLATGFYPD HVELSWWVNGKEVHSGVSTDPQPLKEQPALNDSRYCLSSRLR VSATFWQNPRNHFRQVQFYGLSENDEWTQDRAKPVQTIVSA EAWGRADCGFTSESYQQGVLSATILYEILLGKATLYAVLVSA LVLAMVKKRDSRG	SEQ ID NO: 183
--	--	--	----------------

Donor: HD9

Clonotype	Chain	Region	Sequence	SEQ ID NO
HD9-1	Alpha (α)	CDR1α	VGISA	SEQ ID NO: 184
		CDR2α	LSSGK	SEQ ID NO: 185
		CDR3α	CAARSYNTDKLIF	SEQ ID NO: 186
		Variable domain	MVKIRQFLLAILWLQLSCVSAAKNEVEQSPQN LTAQEGEFITINCSYSGISALHWLQQHPGGG IVSLFMLSSGKKKHGRLIATINIQEKHSSLHI TASHPRDSAVYICAARSYNTDKLIFGTGTRLQ VFPN	SEQ ID NO: 187
		Full – with TRAC constant domain	MVKIRQFLLAILWLQLSCVSAAKNEVEQSPQN LTAQEGEFITINCSYSGISALHWLQQHPGGG IVSLFMLSSGKKKHGRLIATINIQEKHSSLHI TASHPRDSAVYICAARSYNTDKLIFGTGTRLQ VFPNIQNPDPAVYQLRDSKSSDKSVCLFTDFD SQTNVSQSKSDVYITDKTVLDMRSMDFKSNS AWAWSNKSDFACANAFNNSIIPEDTFFPSPES SCDVKLVEKSFETDTNLFQNLVIGFRILL KVAGFNLLMTLRLWSS	SEQ ID NO: 188
HD9-2	Alpha (α)	CDR1α	NSMFDY	SEQ ID NO: 189
		CDR2α	ISSIKDK	SEQ ID NO: 190
		CDR3α	CAASYNARLMF	SEQ ID NO: 191
		Variable domain	MAMLLGASVLILWLQPDWVNSQQKNDDQQVKQ NSPSLSVQEGRISILNCDYTNSMFDYFLWYKK YPAEGPTFLISISSIKDKNEDGRFTVFLNKSA KHLHLHIVPSQPGDSAVYFCAASYNARLMFG DGTQLVVKPN	SEQ ID NO: 192
		Full – with TRAC constant domain	MAMLLGASVLILWLQPDWVNSQQKNDDQQVKQ NSPSLSVQEGRISILNCDYTNSMFDYFLWYKK YPAEGPTFLISISSIKDKNEDGRFTVFLNKSA KHLHLHIVPSQPGDSAVYFCAASYNARLMFG DGTQLVVKPNIQNPDPAVYQLRDSKSSDKSVC LFTDFDSQTNVSQSKSDVYITDKTVLDMRSM DFKSNSAWAWSNKSDFACANAFNNSIIPEDTF FPSPSSCDVKLVEKSFETDTNLFQNLVIG FRILLKLVAGFNLLMTLRLWSS	SEQ ID NO: 193
HD9-1	Beta (β)	CDR1β	SGHTS	SEQ ID NO: 194
		CDR2β	YDEGEE	SEQ ID NO: 195
		CDR3β	CASSWGYQETQYF	SEQ ID NO: 196
		Variable domain	MGPRLFWALLCCLGTGPVEAGVTQSPTHLIK TRGQQATLRCSPISGHTSVYWYQQALGLGLQF LLWYDEGEERNRGNFPPRFSGRQFPNYSSELN VNALELEDSALYLCASSWGYQETQYFGPGTRL LVLE	SEQ ID NO: 197
		Full – with TRBC1 constant domain	MGPRLFWALLCCLGTGPVEAGVTQSPTHLIK TRGQQATLRCSPISGHTSVYWYQQALGLGLQF LLWYDEGEERNRGNFPPRFSGRQFPNYSSELN VNALELEDSALYLCASSWGYQETQYFGPGTRL LVLEDLKNVFPPEVAVFEPSEAEISHTQKATL VCLATGFFPDHVELSWWVNGKEVHSGVSTDPQ PLKEQPALNDSRYCLSSRLRVSATFWQNPRNH FRQVQFYGLSENDEWTQDRAKPVQTIVSAEA WGRADCGFTSVSYQQGVLSATILYEILLGKAT LYAVLVSAVLAMVKKRDF	SEQ ID NO: 198

		Full – with TRBC2 constant domain	MGPRLFWALLCLLGTGPVEAGVTQSPHLLIK TRGQQATLRCSPISGHTSVYWYQQALGLGLQF LLWYDEGEERNRGNFPFRFSGRQFPNYSSELN VNALELEDSELYLCASSWGYQETQYFGPGTRL LVLEDLKNVFPPEVAVFEPSEAEISHTQKATL VCLATGFYDPDHVELSWWVNGKEVHSGVSTDPQ PLKEQPALNDSRYCLSSRLRVSATFWQNP RNHFRQCQVQFYGLSENDEWTQDRAKPV TQIVSAEAWGRADCGFTSESYQQGVLSATILYEILLGKAT LYAVLVSAVLVLMAMVKRKDSRG	SEQ ID NO: 199
HD9-2	Beta (β)	CDR1β	KGHSH	SEQ ID NO: 200
		CDR2β	LQKENI	SEQ ID NO: 201
		CDR3β	CASSPTGGEYYGYTF	SEQ ID NO: 202
		Variable domain	MDTRVLCCAIVICLLGAGLSNAGVMQNPRHLVR RRGQEARLRCSPMKGHSHVYWRQLPEEGLKF MVYLQKENIIDESGMPKERFSAEFPKEGPSIL RIQQVVRGDSAAAYFCASSPTGGEYYGYTFGSG TRLTVVE	SEQ ID NO: 203
		Full – with TRBC1 constant domain	MDTRVLCCAIVICLLGAGLSNAGVMQNPRHLVR RRGQEARLRCSPMKGHSHVYWRQLPEEGLKF MVYLQKENIIDESGMPKERFSAEFPKEGPSIL RIQQVVRGDSAAAYFCASSPTGGEYYGYTFGSG TRLTVVEDLNKVFPPEVAVFEPSEAEISHTQK ATLVCLATGFFPDHVELSWWVNGKEVHSGVST DPQPLKEQPALNDSRYCLSSRLRVSATFWQNP RNHFRQCQVQFYGLSENDEWTQDRAKPV TQIVSAEAWGRADCGFTSVSYQQGVLSATILYEILLG KATLYAVLVSAVLVLMAMVKRKDF	SEQ ID NO: 204
		Full – with TRBC2 constant domain	MDTRVLCCAIVICLLGAGLSNAGVMQNPRHLVR RRGQEARLRCSPMKGHSHVYWRQLPEEGLKF MVYLQKENIIDESGMPKERFSAEFPKEGPSIL RIQQVVRGDSAAAYFCASSPTGGEYYGYTFGSG TRLTVVEDLKNVFPPEVAVFEPSEAEISHTQK ATLVCLATGFYDPDHVELSWWVNGKEVHSGVST DPQPLKEQPALNDSRYCLSSRLRVSATFWQNP RNHFRQCQVQFYGLSENDEWTQDRAKPV TQIVSAEAWGRADCGFTSESYQQGVLSATILYEILLG KATLYAVLVSAVLVLMAMVKRKDSRG	SEQ ID NO: 205
HD9-3	Beta (β)	CDR1β	MNHEY	SEQ ID NO: 206
		CDR2β	SVGAGI	SEQ ID NO: 207
		CDR3β	CASSSYPLRTGRYNSYNSPLHF	SEQ ID NO: 208
		Variable domain	MSIGLLCCAALSLLWAGPVNAGVTQTPKFQVL KTGQSMTLQCAQDMNHEYMSWYRQDPGMGLRL IHYSVGAGITDQGEVPNGYNVSRSTTEDFPLR LLSAAPSQTSVYFCASSSYPLRTGRYNSYNSP LHFGNGTRLTVTE	SEQ ID NO: 209
		Full – with TRBC1 constant domain	MSIGLLCCAALSLLWAGPVNAGVTQTPKFQVL KTGQSMTLQCAQDMNHEYMSWYRQDPGMGLRL IHYSVGAGITDQGEVPNGYNVSRSTTEDFPLR LLSAAPSQTSVYFCASSSYPLRTGRYNSYNSP LHFGNGTRLTVTEDLNKVFPPEVAVFEPSEAE ISHTQKATLVCLATGFFPDHVELSWWVNGKEV HSGVSTDPQPLKEQPALNDSRYCLSSRLRVSATFWQNP RNHFRQCQVQFYGLSENDEWTQDRAKPV TQIVSAEAWGRADCGFTSVSYQQGVLSATILYEILLG KATLYAVLVSAVLVLMAMVKRKDF	SEQ ID NO: 210
		Full – with TRBC2 constant domain	MSIGLLCCAALSLLWAGPVNAGVTQTPKFQVL KTGQSMTLQCAQDMNHEYMSWYRQDPGMGLRL IHYSVGAGITDQGEVPNGYNVSRSTTEDFPLR LLSAAPSQTSVYFCASSSYPLRTGRYNSYNSP LHFGNGTRLTVTEDLKNVFPPEVAVFEPSEAE ISHTQKATLVCLATGFYDPDHVELSWWVNGKEV HSGVSTDPQPLKEQPALNDSRYCLSSRLRVSATFWQNP RNHFRQCQVQFYGLSENDEWTQDRAKPV TQIVSAEAWGRADCGFTSESYQQGVLSATIL	SEQ ID NO: 211

		YEILLGKATLYAVLVVSALVLMAMVKRKDSRG	
--	--	----------------------------------	--

Donor: HD10

Chain	Region	Amino acid sequence	SEQ ID NO
Alpha (α)	CDR1 α	NSMFDY	SEQ ID NO: 212
	CDR2 α	ISSIKDK	SEQ ID NO: 213
	CDR3 α	CAASGGRDDKIIF	SEQ ID NO: 214
	Variable domain	MAMLLGASVLILWLQPDWVNSQQKNDDQQVKQNSP SLSVQEG RISILNCDYTNSMFDYFLWYKKYP AEGPTFLISSISSIKDKNE DGRFTVFLNKS AKHLSLHIVPSQPGDSAVYFCAASGGRDDKI IFGKGTRLHILPN	SEQ ID NO: 215
	Full – with TRAC constant domain	MAMLLGASVLILWLQPDWVNSQQKNDDQQVKQNSP SLSVQEG RISILNCDYTNSMFDYFLWYKKYP AEGPTFLISSISSIKDKNE DGRFTVFLNKS AKHLSLHIVPSQPGDSAVYFCAASGGRDDKI IFGKGTRLHILPNIQNPDPAVYQLRDSKSSDKSVCLFTDFDS QTNVSQSKSDSDVYITDKTVLDMRSMDFKSNSAVAWSNKSDFA CANAFNNSIIPEDTFFPSPESSCDVKLVEKSFETDTNLFQN LSVIGFRILLKLVAGFNLLMTLRLWSS	SEQ ID NO: 216
Beta (β)	CDR1 β	MNHEY	SEQ ID NO: 217
	CDR2 β	SVGAGI	SEQ ID NO: 218
	CDR3 β	CASSYSRTESTDTQYF	SEQ ID NO: 219
	Variable domain	MSIGLLCCAALSLLWAGPVNAGVTQTPKFQVLKTGQSMTLQC AQDMNHEYMSWYRQDPGMGLRLIHYSVGAGITDQGEVPNGYN VSRSTTEDFPLRLLSAAPSQTSVYFCASSYSRTESTDTQYFG PGTRLTVLE	SEQ ID NO: 220
	Full – with TRBC1 constant domain	MSIGLLCCAALSLLWAGPVNAGVTQTPKFQVLKTGQSMTLQC AQDMNHEYMSWYRQDPGMGLRLIHYSVGAGITDQGEVPNGYN VSRSTTEDFPLRLLSAAPSQTSVYFCASSYSRTESTDTQYFG PGTRLTVLEDLNKVFPPEVAVFEPSEAEISHTQKATLVCLAT GFFPDHVELSWWWNGKEVHSGVSTDPQPLKEQPALNDSRYCL SSRLRVSATFWQNPRNHFRCQVQFYGLSENDEWTQDRAKPVT QIVSAEAWGRADCGFTSVSYQQGVLSATILYEILLGKATLYA VLVSALVLMAMVKRKDF	SEQ ID NO: 221
	Full – with TRBC2 constant domain	MSIGLLCCAALSLLWAGPVNAGVTQTPKFQVLKTGQSMTLQC AQDMNHEYMSWYRQDPGMGLRLIHYSVGAGITDQGEVPNGYN VSRSTTEDFPLRLLSAAPSQTSVYFCASSYSRTESTDTQYFG PGTRLTVLEDLNKVFPPEVAVFEPSEAEISHTQKATLVCLAT GFYPDHVELSWWWNGKEVHSGVSTDPQPLKEQPALNDSRYCL SSRLRVSATFWQNPRNHFRCQVQFYGLSENDEWTQDRAKPVT QIVSAEAWGRADCGFTSESYQQGVLSATILYEILLGKATLYA VLVSALVLMAMVKRKDSRG	SEQ ID NO: 222

Accordingly, the present invention provides isolated polypeptides comprising one or more amino acid sequences selected from the group consisting of SEQ ID NOs: 1-114 and 160-222, fragments, variants and homologues thereof.

In one aspect, the invention provides a TCR comprising a TCR alpha chain sequence selected from the group consisting of the HD1-HD10 alpha chain sequences of Table 1, and a TCR beta chain sequence independently selected from the group consisting of the HD1-HD10 beta chain sequences of Table 1.

10 Reduced mispairing and improved TCR expression

The TCR of the invention may be expressed in a T-cell to alter the antigen specificity of the T-cell. TCR-transduced T-cells may express at least two TCR alpha and two TCR beta

chains. While the endogenous TCR alpha/beta chains form a receptor that is self-tolerant, the introduced TCR alpha/beta chains form a receptor with defined specificity for the given target antigen.

However, TCR gene therapy requires sufficient expression of transferred TCRs. Transferred
 5 TCR might be diluted by the presence of the endogeneous TCR, resulting in suboptimal expression of the tumor specific TCR. Furthermore, mispairing between endogenous and introduced chains may occur to form novel receptors, which might display unexpected specificities for self-antigens and cause autoimmune damage when transferred into patients.

Hence, several strategies have been explored to reduce the risk of mispairing between
 10 endogenous and introduced TCR chains. Mutations of the TCR alpha/beta interface is one strategy currently employed to reduce unwanted mispairing. For example, the introduction of a cysteine in the constant domains of the alpha and beta chain allows the formation of a disulfide bond and enhances the pairing of the introduced chains while reducing mispairing with wild type chains.

Accordingly, the TCRs of the present invention may comprise one or more mutations at the
 15 α chain/ β chain interface, such that when the α chain and the β chain are expressed in a T-cell, the frequency of mispairing between said chains and endogenous TCR α and β chains is reduced. In one embodiment, the one or more mutations introduce a cysteine residue into the constant region domain of each of the α chain and the β chain, wherein the cysteine
 20 residues are capable of forming a disulphide bond between the α chain and the β chain.

Another strategy to reduce mispairing relies on the introduction of polynucleotide sequences encoding siRNA, added to the genes encoding for the tumor specific TCR α and or β chains, and designed to limit the expression of the endogenous TCR genes (Okamoto S. *Cancer research* 69, 9003-9011, 2009).

Accordingly, the vector or polynucleotide encoding the TCRs of the present invention may
 25 comprise one or more siRNA or other agents aimed at limiting or abrogating the expression of the endogenous TCR genes.

It is also possible to combine artificial nucleases, such as zinc finger nucleases (ZFN), transcription activator-like effector nucleases (TALEN) or CRISPR/Cas systems, designed to
 30 target the constant regions of the endogenous genes, e.g. TCR genes (TRAC and, or TRBC), to obtain the permanent disruption of the endogenous TCR alpha and/or beta chain genes, thus allowing full expression of the tumor specific TCR and thus reducing or abrogating the risk of TCR mispairing. This process, known as the TCR gene editing proved

superior to TCR gene transfer *in vitro* and *in vivo* (Provasi E., Genovese P., *Nature Medicine* May; 18(5):807-15; 2012).

Accordingly, the TCRs of the present invention may be used to edit T cell specificity by TCR disruption and genetic addition of the tumor specific TCR.

- 5 In addition, the genome editing technology allows targeted integration of a expression cassette, comprising a polynucleotide encoding a TCR of the present invention, and optionally one or more promoter regions and/or other expression control sequences, into an endogenous gene disrupted by the artificial nucleases (Lombardo A., *Nature biotechnology* 25, 1298-1306; 2007).
- 10 Accordingly, the TCRs of the present invention may be used to edit T-cell specificity by targeted integration of a polynucleotide encoding a TCR of the present invention at a genomic region. The integration may be targeted by an artificial nuclease.

- Another strategy developed to increase expression of the transferred TCR and to reduce TCR mispairing consists in “murinization,” which replaces the human TCR α and TCR β constant regions (e.g. the TRAC, TRBC1 and TRBC2 regions) by their murine counterparts.
- 15 Murization of TCR constant regions is described in, for example, Sommermeyer and Uckert J Immunol; 2010 (184:6223-6231). Accordingly, the TCRs of the present invention may be murinized.

Isolated polynucleotide

- 20 The present invention relates to an isolated polynucleotide encoding a TCR receptor of the invention or a part thereof, such as the α chain and/or the β chain, a variable domain or a portion thereof.

The isolated polynucleotide may be double or single stranded, and may be RNA or DNA.

- It will be understood by a skilled person that numerous different polynucleotides can encode the same polypeptide as a result of the degeneracy of the genetic code. In addition, it is to be understood that the skilled person may, using routine techniques, make nucleotide substitutions, additions or deletions that do not affect the polypeptide sequence encoded by the polynucleotides of the invention to reflect the codon usage of any particular host organism in which the polypeptides of the invention are to be expressed.
- 25

- 30 The polynucleotides described herein may be modified by any method available in the art. Such modifications may be carried out in order to enhance the *in vivo* activity or lifespan of the polynucleotides of the invention.

Polynucleotides such as DNA polynucleotides may be produced recombinantly, synthetically or by any means available to those of skill in the art. They may also be cloned by standard techniques.

Longer polynucleotides will generally be produced using recombinant means, for example using polymerase chain reaction (PCR) cloning techniques. This will involve making a pair of primers (e.g. of about 15 to 30 nucleotides) flanking the target sequence which it is desired to clone, bringing the primers into contact with Mrna or Cdna obtained from an animal or human cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region, isolating the amplified fragment (e.g. by purifying the reaction mixture with an agarose gel) and recovering the amplified DNA. The primers may be designed to contain suitable restriction enzyme recognition sites so that the amplified DNA can be cloned into a suitable vector.

Examples of nucleotide sequences encoding TCRs according to the present invention are provided in the Table 2.

Table 2

Donor	Chain	Nucleotide sequence	SEQ ID NO
HD1	α (with TRAC)	ATGGAGACTCTCCTGAAAGTGCTTTCAGGCACCTTGTTGTGGCAGTTGACCTGGG TGAGAAGCCAACAACCAGTGCAGAGTCCTCAAGCCGTGATCCTCCGAGAAGGGGA AGATGCTGTCATCAACTGCAGTTCCTCCAAGGCTTTATATTCTGTACACTGGTAC AGGCAGAAGCATGGTGAAGCACCCGCTTCCTGATGATATTACTGAAGGGTGGAG AACAGAAGGGTCATGAAAAAATATCTGCTTCATTTAATGAAAAAAGCAGCAAAG CTCCCTGTACCTTACGGCCTCCCAGCTCAGTTACTCAGGAACCTACTTCTGCGGC ACAGCTTGGATTAACTGACTACAAGCTCAGCTTTGGAGCCGGAACACAGTAACTG TAAGAGCAAATATCCAGAACCCTGACCCTGCCGTGTACCAGCTGAGAGACTCTAA ATCCAGTGACAAGTCTGTCTGCCTATTACCGATTTTGATTCTCAAACAAATGTG TCACAAAGTAAGGATTCTGATGTGTATATCACAGACAAAAGTGTGCTAGACATGA GGTCTATGGACTTCAAGAGCAACAGTGCTGTGGCCTGGAGCAACAAATCTGACTT TGCATGTGCAAACGCCTTCAACAACAGCATTATTCCAGAAGACACCTTCTTCCCC AGCCCAGAAAAGTTCTGTGATGTCAAGCTGGTGCAGAAAAGCTTTGAAACAGATA CGAACCTAACTTTCAAAACCTGTGAGTATTGGGTTCGGAATCCTCCTCCTGAA AGTGGCCGGGTTTAATCTGCTCATGACGCTGCGGCTGTGGTCCAGC	SEQ ID NO: 132
	β (with TRBC1)	ATGGGCTCCTGGACCCTCTGCTGTGTGTCCCTTTGCATCCTGGTAGCAAAGCACA CAGATGCTGGAGTTATCCAGTCACCCCGGCACGAGGTGACAGAGATGGGACAAGA AGTGACTCTGAGATGTAAACCAATTTCAAGACACGACTACCTTTTCTGGTACAGA CAGACCATGATGCGGGGACTGGAGTTGCTCATTTACTTTAACAACAACGTTCCGA TAGATGATTCAAGGATGCCCCGAGGATCGATTCTCAGCTAAGATGCCTAATGCATC ATTCTCCACTCTGAAGATCCAGCCCTCAGAACCCAGGGACTCAGCTGTGTACTTC TGTGCCAGCAGAAAAACCGGGGGATATAGCAATCAGCCCCAGCATTTTGGTGATG GGACTCGACTCTCCATCCTAGAGGACCTGAACAAGGTGTTCCACCCGAGGTCGC TGTGTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTG TGCTTGCCACAGGCTTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATG GGAAGGAGGTGCACAGTGGGGTCAGCACGGACCCGAGCCCCCTCAAGGAGCAGCC CGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCTGAGGGTCTCGGCCACC TTCTGGCAGAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCT CGGAGAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTCACCCAGATCGTCAG CGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCTACCAGCAA GGGGTCTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCCTGT ATGCTGTGCTGGTCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAGGATTT C	SEQ ID NO: 133
	β (with TRBC2)	ATGGGCTCCTGGACCCTCTGCTGTGTGTCCCTTTGCATCCTGGTAGCAAAGCACA CAGATGCTGGAGTTATCCAGTCACCCCGGCACGAGGTGACAGAGATGGGACAAGA AGTGACTCTGAGATGTAAACCAATTTCAAGACACGACTACCTTTTCTGGTACAGA CAGACCATGATGCGGGGACTGGAGTTGCTCATTTACTTTAACAACAACGTTCCGA TAGATGATTCAAGGATGCCCCGAGGATCGATTCTCAGCTAAGATGCCTAATGCATC ATTCTCCACTCTGAAGATCCAGCCCTCAGAACCCAGGGACTCAGCTGTGTACTTC	SEQ ID NO: 134

		TGTGCCAGCAGAAAAACCGGGGGATATAGCAATCAGCCCCAGCATTTTGGTGATG GGACTCGACTCTCCATCCTAGAGGACCTGAAAAACGTGTTCCCACCCGAGGTCGC TGTGTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTG TGCCTGGCCACAGGCTTCTACCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATG GGAAGGAGGTGCACAGTGGGGTCAGCACAGACCCGCAGCCCCTCAAGGAGCAGCC CGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACC TTCTGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCT CGGAGAATGACGAGTGGACCCAGGATAGGGCCAAACCTGTCACCCAGATCGTCAG CGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTCACCTCCGAGTCTTACCAGCAA GGGGTCCTGTCTGCCACCATCCTCTATGAGATCTTGCTAGGGAAGGCCACCTTGT ATGCCGTGCTGGTCAGTGCCCTCGTGCTGATGGCCATGGTCAAGAGAAAGGATTC CAGAGGC	
HD2-1	α (with TRAC)	ATGCTCCTGCTGCTCGTCCCAGTGCTCGAGGTGATTTTTACCCTGGGAGGAACCA GAGCCCAGTCGGTGACCCAGCTTGGCAGCCACGTCTCTGTCTCTGAAGGAGCCCT GGTTCTGCTGAGGTGCAACTACTCATCGTCTGTTCCACCATATCTCTTCTGGTAT GTGCAATACCCCAACCAAGGACTCCAGCTTCTCCTGAAGTACACATCAGCGGCCA CCCTGGTTAAAGGCATCAACGTTTTTGAGGCTGAATTTAAGAAGAGTGAAACCTC CTTCCACCTGACGAAACCTCAGCCCATATGAGCGACGCGGCTGAGTACTTCTGT GCTGTGAGATTATCTGGTTCTGCAAGGCAACTGACCTTTGGATCTGGGACACAAT TGACTGTTTTACCTGATATCCAGAACCCTGACCCTGCCGTGTACCAGCTGAGAGA CTCTAAATCCAGTGACAAGTCTGTCTGCCTATTACCGATTTTGATTCTCAAACA AATGTGTCACAAAGTAAGGATTCTGATGTGTATATCACAGACAAAACCTGTGCTAG ACATGAGGTCTATGGACTTCAAGAGCAACAGTGCTGTGGCCTGGAGCAACAAATC TGACTTTGCATGTGCAAACGCCTTCAACAACAGCATTATTCCAGAAGACACCTTC TTCCCCAGCCCAGAAAGTTTCTGTGATGTCAAGCTGGTCGAGAAAAGCTTTGAAA CAGATACGAACCTAAACTTTCAAAACCTGTCAGTGATTGGGTTCCGAATCCTCCT CCTGAAAGTGGCCGGGTTTAATCTGCTCATGACGCTGCGGCTGTGGTCCAGC	SEQ ID NO: 135
	β (with TRBC1)	ATGGGCACCAGGCTCCTCTGCTGGGCGGCCCTCTGTCTCCTGGGAGCAGAACTCA CAGAAGCTGGAGTTGCCAGTCTCCCAGATATAAGATTATAGAGAAAAGGCAGAG TGTGGCTTTTTGGTGCAATCCTATATCTGGCCATGCTACCCTTTACTGGTACCAG CAGATCCTGGGACAGGGCCCCAAAGCTTCTGATTTCAGTTTCAGAATAACGGTGTAG TGGATGATTACAGTTGCCTAAGGATCGATTTTCTGCAGAGAGGCTCAAAGGAGT AGACTCCACTCTCAAGATCCAGCCTGCAAAGCTTGAGGACTCGGCCGTGTATCTC TGTGCCAGCAGCTTACTGGGAGACGAGCAGTACTTCGGGCCGGGCACCAGGCTCA CGGTCACAGAGGACCTGAACAAGGTGTTCCCACCCGAGGTCGCTGTGTTTGAGCC ATCAGAAGCAGAGATCTCCCACACCCAAAAGGCCACACTGGTGTGCCTGGCCACA GGCTTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGAAGGAGGTGC ACAGTGGGGTCAGCACGGACCCGCAGCCCCCTCAAGGAGCAGCCCGCCCTCAATGA CTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTCTGGCAGAAC CCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGAATGACG AGTGGACCCAGGATAGGGCCAAACCCGTACCCAGATCGTCAGCGCCGAGGCCTG GGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCTTACCAGCAAGGGGTCTCTGTCT GCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCCTGTATGCTGTGCTGG TCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAGGATTTC	SEQ ID NO: 136
	β (with TRBC2)	ATGGGCACCAGGCTCCTCTGCTGGGCGGCCCTCTGTCTCCTGGGAGCAGAACTCA CAGAAGCTGGAGTTGCCAGTCTCCCAGATATAAGATTATAGAGAAAAGGCAGAG TGTGGCTTTTTGGTGCAATCCTATATCTGGCCATGCTACCCTTTACTGGTACCAG CAGATCCTGGGACAGGGCCCCAAAGCTTCTGATTTCAGTTTCAGAATAACGGTGTAG TGGATGATTACAGTTGCCTAAGGATCGATTTTCTGCAGAGAGGCTCAAAGGAGT AGACTCCACTCTCAAGATCCAGCCTGCAAAGCTTGAGGACTCGGCCGTGTATCTC TGTGCCAGCAGCTTACTGGGAGACGAGCAGTACTTCGGGCCGGGCACCAGGCTCA CGGTCACAGAGGACCTGAAAAACGTGTTCCCACCCGAGGTCGCTGTGTTTGAGCC ATCAGAAGCAGAGATCTCCCACACCCAAAAGGCCACACTGGTGTGCCTGGCCACA GGCTTCTACCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGAAGGAGGTGC ACAGTGGGGTCAGCACAGACCCGCAGCCCCCTCAAGGAGCAGCCCGCCCTCAATGA CTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTCTGGCAGAAC CCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGAATGACG AGTGGACCCAGGATAGGGCCAAACCTGTCACCCAGATCGTCAGCGCCGAGGCCTG GGGTAGAGCAGACTGTGGCTTCACCTCCGAGTCTTACCAGCAAGGGGTCTCTGTCT GCCACCATCCTCTATGAGATCTTGCTAGGGAAGGCCACCCTGTATGCCGTGCTGG TCAGTGCCCTCGTGCTGATGGCCATGGTCAAGAGAAAGGATTCCAGAGGC	SEQ ID NO: 137
HD2-2	α (with TRAC)	ATGGCATGCCCTGGCTTCCTGTGGGCACTTGTGATCTCCACCTGTCTTGAATTTA GCATGGCTCAGACAGTCACTCAGTCTCAACCAGAGATGTCTGTGCAGGAGGCAGA GACCGTGACCCTGAGCTGCACATATGACACCAGTGAGAGTGATTATTATTTATTC TGGTACAAGCAGCCTCCCAGCAGGCAGATGATTCTCGTTATTCGCCAAGAAGCTT ATAAGCAAACAGAATGCAACAGAGAATCGTTTTCTCTGTGAACTTCCAGAAAGCAGC CAAATCCTTCAGTCTCAAGATCTCAGACTCACAGCTGGGGGATGCCGCGATGTAT TTCTGTGCTTATAGGAGTCTAAAATATGGAACAAAACCTGGTCTTTGGCGCAGGAA CCATTCTGAGAGTCAAGTCCATATATCCAGAACCCTGACCCTGCCGTGTACCAGCT GAGAGACTCTAAATCCAGTGACAAGTCTGTCTGCCTATTACCGATTTTGATTCT CAAAACAAATGTGTACAAAAGTAAGGATTCTGATGTGTATATCACAGACAAAACCTG TGCTAGACATGAGGTCTATGGACTTCAAGAGCAACAGTGCTGTGGCCTGGAGCAA CAAATCTGACTTTGCATGTGCAAACGCCTTCAACAACAGCATTATTCCAGAAGAC ACCTTCTTCCCCAGCCCAGAAAGTTCTGTGATGTCAAGCTGGTCGAGAAAAGCT	SEQ ID NO: 138

		TTGAAACAGATACGAACCTAAACTTTCAAACCTGTCAGTGATTGGGTTCCGAATCCTCCTCCTGAAAGTGGCCGGGTTTAACTCTGCTCATGACGCTGCGGCTGTGGTCCAGC	
	β (with TRBC1)	ATGGGCACCAGGCTCCTCTTCTGGGTGGCCTTCTGTCTCCTGGGGGCAGATCACA CAGGAGCTGGAGTCTCCCAGTCCCCCAGTAACAAGGTCACAGAGAAGGGAAAGGA TGTAGAGCTCAGGTGTGATCCAATTTTCAGGTCATACTGCCCTTTACTGGTACCGA CAGAGCCTGGGGCAGGGCCTGGAGTTTTTAATTTACTTCCAAGGCAACAGTGCAC CAGACAAATCAGGGCTGCCCAGTGATCGCTTCTCTGCAGAGAGGACTGGGGGATC CGTCTCCACTCTGACGATCCAGCGCACACAGCAGGAGGACTCGGCCGTGTATCTC TGTGCCAGCAGCTTGGTAGCTTTACAGGGTGCGGGCGAGCAGTACTTCGGGCCGG GCACCAGGCTCACGGTCACAGAGGACCTGAACAAGGTGTTCCACCCGAGGTCGC TGTGTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTG TGCCTGGCCACAGGCTTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATG GGAAGGAGGTGCACAGTGGGGTCAGCACGGACCCGCAGCCCCTCAAGGAGCAGCC CGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACC TTCTGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCT CGGAGAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTACCCAGATCGTCAG CGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCCTACCAGCAA GGGGTCTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCCTGT ATGCTGTGCTGGTCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAGGATTT C	SEQ ID NO: 139
	β (with TRBC2)	ATGGGCACCAGGCTCCTCTTCTGGGTGGCCTTCTGTCTCCTGGGGGCAGATCACA CAGGAGCTGGAGTCTCCCAGTCCCCCAGTAACAAGGTCACAGAGAAGGGAAAGGA TGTAGAGCTCAGGTGTGATCCAATTTTCAGGTCATACTGCCCTTTACTGGTACCGA CAGAGCCTGGGGCAGGGCCTGGAGTTTTTAATTTACTTCCAAGGCAACAGTGCAC CAGACAAATCAGGGCTGCCCAGTGATCGCTTCTCTGCAGAGAGGACTGGGGGATC CGTCTCCACTCTGACGATCCAGCGCACACAGCAGGAGGACTCGGCCGTGTATCTC TGTGCCAGCAGCTTGGTAGCTTTACAGGGTGCGGGCGAGCAGTACTTCGGGCCGG GCACCAGGCTCACGGTCACAGAGGACCTGAAAAACGTGTTCCACCCGAGGTCGC TGTGTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTG TGCCTGGCCACAGGCTTCTACCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATG GGAAGGAGGTGCACAGTGGGGTCAGCACAGACCCGCAGCCCCTCAAGGAGCAGCC CGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACC TTCTGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCT CGGAGAATGACGAGTGGACCCAGGATAGGGCCAAACCTGTACCCAGATCGTCAG CGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTCACCTCCGAGTCTTACCAGCAA GGGGTCTGTCTGCCACCATCCTCTATGAGATCTTGCTAGGGAAGGCCACCTTGT ATGCCGTGCTGGTCAGTGCCCTCGTGCTGATGGCCATGGTCAAGAGAAAGGATTC CAGAGGC	SEQ ID NO: 140
HD3	α (with TRAC)	ATGATATCCTTGAGAGTTTTACTGGTGATCCTGTGGCTTCAGTTAAGCTGGGTTT GGAGCCAACGGAAGGAGGTGGAGCAGGATCCTGGACCCTTCAATGTTCCAGAGGG AGCCACTGTCGCTTTCAACTGTACTTTACAGCAACAGTGCTTCTCAGTCTTTCTTC TGGTACAGACAGGATTGCAGGAAAGAACCTAAGTTGCTGATGTCCGTATACTCCA GTGGTAATGAAGATGGAAGGTTTACAGCACAGCTCAATAGAGCCAGCCAGTATAT TTCCCTGCTCATCAGAGACTCCAAGCTCAGTGATTCAGCCACCTACCTCTGTGTG GTGAACCTCCTGTCTAACCAGGGAGGAAAGCTTATCTTCGGACAGGGAACGGAGT TATCTGTGAAACCCAATATCCAGAACCCTGACCCTGCCGTGTACCAGCTGAGAGA CTCTAAATCCAGTGACAAGTCTGTCTGCCTATTACCCGATTTTGATTCTCAAACA AATGTGTCACAAAGTAAGGATTCTGATGTGTATATCACAGACAAAACCTGTGCTAG ACATGAGGTCTATGGACTTCAAGAGCAACAGTGCTGTGGCCTGGAGCAACAAATC TGACTTTGCATGTGCAAACGCCTTCAACAACAGCATTATTCCAGAAGACACCTTC TTCCCCAGCCCAGAAAGTTTCTGTGATGTCAAGCTGGTTCGAGAAAAGCTTTGAAA CAGATACGAACCTAAACTTTCAAACCTGTCAAGTGATTGGGTTCCGAATCCTCCT CCTGAAAGTGGCCGGGTTTAACTCTGCTCATGACGCTGCGGCTGTGGTCCAGC	SEQ ID NO: 141
	β (with TRBC1)	ATGGGCTGCAGGCTGCTCTGCTGTGCGGTTCTCTGTCTCCTGGGAGCGGGTGAGT TGGTCCCCATGGAACGGGAGTTACGCAGACACCAAGACACCTGGTCATGGGAAT GACAAATAAGAAGTCTTTGAAATGTGAACAACATCTGGGTCATAACGCTATGTAT TGGTACAAGCAAAGTGCTAAGAAGCCACTGGAGCTCATGTTTGTCTACAGTCTTG AAGAACGGGTTGAAAACAACAGTGTGCCAAGTCGCTTCTCACCTGAATGCCCCAA CAGCTCTCACTTATTCCTTCACCTACACACCCTGCAGCCAGAAGACTCGGCCCTG TATCTCTGCGCCAGCAGCCAAGATTACTTGGTTTCTAATGAAAACTGTTTTTTG GCAGTGGAACCCAGCTCTCTGTCTTGGAGGACCTGAACAAGGTGTTCCACCCGA GGTCGCTGTGTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACA CTGGTGTGCCCTGGCCACAGGCTTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGG TGAATGGGAAGGAGGTGCACAGTGGGGTCAGCACGGACCCGCAGCCCCTCAAGGA GCAGCCCCGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCG GCCACCTTCTGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACG GGCTCTCGGAGAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTACCCAGAT CGTCAGCGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCCTAC CAGCAAGGGGTCCTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCA CCCTGTATGCTGTGCTGGTCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAA GGATTC	SEQ ID NO: 142
	β (with TRBC2)	ATGGGCTGCAGGCTGCTCTGCTGTGCGGTTCTCTGTCTCCTGGGAGCGGGTGAGT TGGTCCCCATGGAACGGGAGTTACGCAGACACCAAGACACCTGGTCATGGGAAT GACAAATAAGAAGTCTTTGAAATGTGAACAACATCTGGGTCATAACGCTATGTAT	SEQ ID NO: 143

		TGGTACAAGCAAAGTGCTAAGAAGCCACTGGAGCTCATGTTTGTCTACAGTCTTG AAGAACGGGTTGAAAACAACAGTGTGCCAAGTCGCTTCTCACCTGAATGCCCCAA CAGCTCTCACTTATTCCTTCACCTACACACCCTGCAGCCAGAAGACTCGGCCCTG TATCTCTGCGCCAGCAGCCAAGATTACTTGGTTTCTAATGAAAACTGTTTTTTG GCAGTGGAACCCAGCTCTCTGTCTTGGAGGACCTGAAAAACGTGTTCCCACCCGA GGTCGCTGTGTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACA CTGGTGTGCCTGGCCACAGGCTTCTACCCCGACCACGTGGAGCTGAGCTGGTGGG TGAATGGGAAGGAGGTGCACAGTGGGGTCAGCACAGACCCGCAGCCCCCTCAAGGA GCAGCCCCGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCG GCCACCTTCTGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACG GGCTCTCGGAGAATGACGAGTGGACCCAGGATAGGGCCAAACCTGTCACCCAGAT CGTCAGCGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTCACCTCCGAGTCTTAC CAGCAAGGGGTCTGTCTGCCACCATCCTCTATGAGATCTTGCTAGGGAAGGCCA CCTTGTATGCCGTGCTGGTCAGTGCCCTCGTGCTGATGGCCATGGTCAAGAGAAA GGATTCCAGAGGC	
HD4-1	α1 (with TRAC)	ATGGAAACTCTCCTGGGAGTGTCTTTGGTGATTCTATGGCTTCAACTGGCTAGGG TGAACAGTCAACAGGGAGAAGAGGATCCTCAGGCCTTGAGCATCCAGGAGGGTGA AAATGCCACCATGAACTGCAGTTACAAAAGTAGTATAAAACAATTTACAGTGGTAT AGACAAAATTCAGGTAGAGGCCTTGTCCACCTAATTTTAATACGTTCAAATGAAA GAGAGAAACACAGTGGAAGATTAAAGAGTCACGCTTGACACTTCCAAGAAAAGCAG TTCCTTGTTGATCACGGCTTCCCGGGCAGCAGACACTGCTTCTTACTTCTGTGCT ACGGACGCGTATTCAGGAAACACACCTCTTGTCTTTGGAAAGGGCACAAGACTTT CTGTGATTGCAAATATCCAGAACCCTGACCCTGCCGTGTACCAGCTGAGAGACTC TAAATCCAGTGACAAGTCTGTCTGCCTATTACCGATTTTGATTCTCAAACAAAT GTGTCACAAAGTAAGGATTCTGATGTGTATATCACAGACAAAAGTGTGCTAGACA TGAGGTCTATGGACTTCAAGAGCAACAGTGCTGTGGCCTGGAGCAACAAATCTGA CTTTGCATGTGCAAACGCCTTCAACAACAGCATTATTCCAGAAGACACCTTCTTC CCCAGCCCAGAAAGTTCTGTGATGTCAAGCTGGTCGAGAAAAGCTTTGAAACAG ATACGAACCTAAACTTTCAAACCTGTGAGTGATTGGGTTCGGAATCCTCCTCCT GAAAGTGGCCGGGTTTAATCTGCTCATGACGCTGCGGCTGTGGTCCAGC	SEQ ID NO: 144
	β1 (with TRBC1)	ATGAGCATCGGGCTCCTGTGCTGTGTGGCCTTTTCTCTCCTGTGGGCAAGTCCAG TGAATGCTGGTGTCACTCAGACCCCAAAATTCAGGTCCCTGAAGACAGGACAGAG CATGACACTGCAGTGTGCCCAGGATATGAACCATAACTCCATGTACTGGTATCGA CAAGACCCAGGCATGGGACTGAGGCTGATTTATTACTCAGCTTCTGAGGGTACCA CTGACAAAGGAGAAGTCCCCAATGGCTACAATGTCTCCAGATTAAACAAACGGGA GTTCTCGCTCAGGCTGGAGTCGGCTGCTCCCTCCCAGACATCTGTGTACTTCTGT GCCAGCAGGGCAGCAGGGTTGGACACTGAAGCTTTCTTTGGACAAGGCACCAGAC TCACAGTTGTAGAGGACCTGAACAAGGTGTTCCACCCGAGGTCGCTGTGTTTGA GCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGCCTGGCC ACAGGCTTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGAAGGAGG TGCACAGTGGGGTCAGCACGACCCGCAGCCCCCTCAAGGAGCAGCCCGCCCTCAA TGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTCTGGCAG AACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGAATG ACGAGTGGACCCAGGATAGGGCCAAACCCGTCACCCAGATCGTCAGCGCCGAGGC CTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCTACCAGCAAGGGGTCCTG TCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCCTGTATGCTGTGC TGGTCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAGGATTTC	SEQ ID NO: 145
	β1 (with TRBC2)	ATGAGCATCGGGCTCCTGTGCTGTGTGGCCTTTTCTCTCCTGTGGGCAAGTCCAG TGAATGCTGGTGTCACTCAGACCCCAAAATTCAGGTCCCTGAAGACAGGACAGAG CATGACACTGCAGTGTGCCCAGGATATGAACCATAACTCCATGTACTGGTATCGA CAAGACCCAGGCATGGGACTGAGGCTGATTTATTACTCAGCTTCTGAGGGTACCA CTGACAAAGGAGAAGTCCCCAATGGCTACAATGTCTCCAGATTAAACAAACGGGA GTTCTCGCTCAGGCTGGAGTCGGCTGCTCCCTCCCAGACATCTGTGTACTTCTGT GCCAGCAGGGCAGCAGGGTTGGACACTGAAGCTTTCTTTGGACAAGGCACCAGAC TCACAGTTGTAGAGGACCTGAAAAACGTGTTCCACCCGAGGTCGCTGTGTTTGA GCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGCCTGGCC ACAGGCTTCTACCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGAAGGAGG TGCACAGTGGGGTCAGCACAGACCCGCAGCCCCCTCAAGGAGCAGCCCGCCCTCAA TGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTCTGGCAG AACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGAATG ACGAGTGGACCCAGGATAGGGCCAAACCTGTACCCAGATCGTCAGCGCCGAGGC CTGGGGTAGAGCAGACTGTGGCTTCACCTCCGAGTCTTACCAGCAAGGGGTCCTG TCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCCTGTATGCCGTGC TGGTCAGTGCCCTCGTGCTGATGGCCATGGTCAAGAGAAAGGATTCCAGAGGC	SEQ ID NO: 146
HD4-2	α2 (with TRAC)	ATGGAGACCCTCTTGGGCCTGCTTATCCTTTGGCTGCAGCTGCAATGGGTGAGCA GCAAAACAGGAGGTGACGCAGATTCCCTGCAGCTCTGAGTGTCCAGAAGGAGAAAA CTTGGTTCTCAACTGCAGTTTCACTGATAGCGCTATTTACAACCTCCAGTGGTTT AGGCAGGACCCTGGGAAAGGTCTCACATCTCTGTTGCTTATTCAAGTCAAGTCAGA GAGAGCAAACAAGTGGAAGACTTAAATGCCTCGCTGGATAAATCATCAGGACGTAG TACTTTATACATTGCAGCTTCTCAGCCTGGTGACTCAGCCACCTACCTCTGTGCT GTCCGGGCAGAGATTTATAACCAGGGAGGAAAGCTTATCTTCGGACAGGGAACGG AGTTATCTGTGAAACCCAATATCCAGAACCCTGACCCTGCCGTGTACCAGCTGAG AGACTCTAAATCCAGTGACAAGTCTGTCTGCCTATTACCGATTTTGATTCTCAA ACAAATGTGTACAAAGTAAGGATTCTGATGTGTATATCACAGACAAAAGTGTGC TAGACATGAGGTCTATGGACTTCAAGAGCAACAGTGCTGTGGCCTGGAGCAACAA	SEQ ID NO: 147

		ATCTGACTTTTGCATGTGCAAACGCCTTCAACAACAGCATTATTCCAGAAGACACC TTCTTCCCCAGCCCAGAAAAGTTCCTGTGATGTCAAGCTGGTCGAGAAAAGCTTTG AAACAGATACGAACCTAAACTTTTCAAACCTGTCAGTGATTGGGTTCGAATCCT CCTCCTGAAAGTGGCCGGGTTTAATCTGCTCATGACGCTGCGGCTGTGGTCCAGC	
	β2 (with TRBC1)	ATGAGCATCAGCCTCCTGTGCTGTGCAGCCTTTTCTCTCCTGTGGGCAGGTCCAG TGAATGCTGGTGTCACTCAGACCCCCAAAATTCCGCATCCTGAAGATAGGACAGAG CATGACACTGCAGTGTACCCAGGATATGAACCATAACTACATGTACTGGTATCGA CAAGACCCAGGCATGGGGCTGAAGCTGATTTATTATTTCAGTTGGTGCTGGTATCA CTGATAAAGGAGAAGTCCCGAATGGCTACAACGTCTCCAGATCAACCACAGAGGA TTTCCCGCTCAGGCTGGAGTTGGCTGCTCCCTCCCAGACATCTGTGTACTTCTGT GCCAGTACCCAAACTCCCTACGAGCAGTACTTCGGGCCGGGCACCAGGCTCACGG TCACAGAGGACCTGAACAAGGTGTTCCACCCGAGGTCGCTGTGTTTGAGCCATC AGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGCCTGGCCACAGGC TTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGAAGGAGGTGCACA GTGGGGTCAGCACGGACCCGCAGCCCCTCAAGGAGCAGCCCGCCCTCAATGACTC CAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTCTGGCAGAACCCC CGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGAATGACGAGT GGACCCAGGATAGGGCCAAACCCGTCACCCAGATCGTCAGCGCCGAGGCCTGGGG TAGAGCAGACTGTGGCTTTACCTCGGTGTCTACCAGCAAGGGGTCTGTCTGCC ACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCTGTATGCTGTGCTGGTCA GCGCCCTTGTTGATGGCCATGGTCAAGAGAAAGGATTTTC	SEQ ID NO: 148
	β2 (with TRBC2)	ATGAGCATCAGCCTCCTGTGCTGTGCAGCCTTTTCTCTCCTGTGGGCAGGTCCAG TGAATGCTGGTGTCACTCAGACCCCCAAAATTCCGCATCCTGAAGATAGGACAGAG CATGACACTGCAGTGTACCCAGGATATGAACCATAACTACATGTACTGGTATCGA CAAGACCCAGGCATGGGGCTGAAGCTGATTTATTATTTCAGTTGGTGCTGGTATCA CTGATAAAGGAGAAGTCCCGAATGGCTACAACGTCTCCAGATCAACCACAGAGGA TTTCCCGCTCAGGCTGGAGTTGGCTGCTCCCTCCCAGACATCTGTGTACTTCTGT GCCAGTACCCAAACTCCCTACGAGCAGTACTTCGGGCCGGGCACCAGGCTCACGG TCACAGAGGACCTGAAAAACGTGTTCCACCCGAGGTCGCTGTGTTTGAGCCATC AGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGCCTGGCCACAGGC TTCTACCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGAAGGAGGTGCACA GTGGGGTCAGCACAGACCCGCAGCCCCTCAAGGAGCAGCCCGCCCTCAATGACTC CAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTCTGGCAGAACCCC CGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGAATGACGAGT GGACCCAGGATAGGGCCAAACCTGTACCCAGATCGTCAGCGCCGAGGCCTGGGG TAGAGCAGACTGTGGCTTACCTCCGAGTCTTACCAGCAAGGGGTCTGTCTGCC ACCATCCTCTATGAGATCTTGCTAGGGAAGGCCACCTTGTATGCCGTGCTGGTCA GTGCCCTCGTGCTGATGGCCATGGTCAAGAGAAAGGATTCCAGAGGC	SEQ ID NO: 149
	β3 (with TRBC1)	ATGGACTCCTGGACCTTCTGCTGTGTGTCCCTTTGCATCCTGGTAGCGAAGCATA CAGATGCTGGAGTTATCCAGTCACCCCGCCATGAGGTGACAGAGATGGGACAAGA AGTGACTCTGAGATGTAAACCAATTTCAGGCCACAACCTCCCTTTTCTGGTACAGA CAGACCATGATGCGGGGACTGGAGTTGCTCATTTACTTTAACAACAACGTTCCGA TAGATGATTTCAGGGATGCCCCGAGGATCGATTCTCAGCTAAGATGCCTAATGCATC ATTCTCCACTCTGAAGATCCAGCCCTCAGAACCCAGGGACTCAGCTGTGTACTTC TGTGCCAGCAGCACAGTGGGAGGGGAGGATTATGGCTACACCTTCGGTTCGGGGA CCAGGTTAACCGTTGTAGAGGACCTGAACAAGGTGTTCCACCCGAGGTTCGCTGT GTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGC CTGGCCACAGGCTTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGA AGGAGGTGCACAGTGGGGTCAGCACGGACCCGCAGCCCCTCAAGGAGCAGCCCGC CCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTC TGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGG AGAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTCACCCAGATCGTCAGCGC CGAGGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCTACCAGCAAGGG GTCTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCTGTATG CTGTGCTGGTTCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAGGATTTTC	SEQ ID NO: 150
	β3 (with TRBC2)	ATGGACTCCTGGACCTTCTGCTGTGTGTCCCTTTGCATCCTGGTAGCGAAGCATA CAGATGCTGGAGTTATCCAGTCACCCCGCCATGAGGTGACAGAGATGGGACAAGA AGTGACTCTGAGATGTAAACCAATTTCAGGCCACAACCTCCCTTTTCTGGTACAGA CAGACCATGATGCGGGGACTGGAGTTGCTCATTTACTTTAACAACAACGTTCCGA TAGATGATTTCAGGGATGCCCCGAGGATCGATTCTCAGCTAAGATGCCTAATGCATC ATTCTCCACTCTGAAGATCCAGCCCTCAGAACCCAGGGACTCAGCTGTGTACTTC TGTGCCAGCAGCACAGTGGGAGGGGAGGATTATGGCTACACCTTCGGTTCGGGGA CCAGGTTAACCGTTGTAGAGGACCTGAAAAACGTGTTCCACCCGAGGTTCGCTGT GTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGC CTGGCCACAGGCTTCTACCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGA AGGAGGTGCACAGTGGGGTCAGCACGGACCCGCAGCCCCTCAAGGAGCAGCCCGC CCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTC TGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGG AGAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTCACCCAGATCGTCAGCGC CGAGGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCTACCAGCAAGGG GTCTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCTGTATG CCGTGCTGGTTCAGTGCCTCGTGCTGATGGCCATGGTCAAGAGAAAGGATTCCAG AGGC	SEQ ID NO: 151
HD5	α (with TRAC)	ATGACATCCATTTCGAGCTGTATTTATATTCTGTGGCTGCAGCTGGACTTGGTGA ATGGAGAGAATGTGGAGCAGCATCCTTCAACCCTGAGTGTCCAGGAGGGAGACAG	SEQ ID NO: 152

		CGCTGTTATCAAGTGTACTTATTTCAGACAGTGCCTCAAACACTACTTCCCTTGGTAT AAGCAAGAACTTGGA AAAAGACCTCAGCTTATTATAGACATTTCGTTCAAATGTGG GCGAAAAGAAAGACCAACGAATTGCTGTTACATTGAACAAGACAGCCAAACATTT CTCCCTGCACATCACAGAGACCCAACCTGAAGACTCGGCTGTCTACTTCTGTGCA GCAAGTATGGCTGGGGCTGGGAGTTACCAACTCACTTTCGGGAAGGGGACCAAAC TCTCGGTCATACCAAATATCCAGAACCCTGACCCTGCCGTGTACCAGCTGAGAGA CTCTAAATCCAGTGACAAGTCTGTCTGCCTATTACCGATTTTGATTCTCAAACA AATGTGTCACAAAGTAAGGATTCTGATGTGTATATCACAGACAAAACCTGTGCTAG ACATGAGGTCTATGGACTTCAAGAGCAACAGTGCTGTGGCCTGGAGCAACAAATC TGACTTTGCATGTGCAAACGCCTTCAACAACAGCATTATTCCAGAAGACACCTTC TTCCCCAGCCCAGAAAGTTCTGTGATGTCAAGCTGGTCGAGAAAAGCTTTGAAA CAGATACGAACCTAAACTTTCAAAACCTGTCAAGTGATTGGGTTCCGAATCCTCCT CCTGAAAGTGGCCGGGTTTAATCTGCTCATGACGCTGCGGCTGTGGTCCAGC	
	β1 (with TRBC1)	ATGGGCACAAGGTTGTTCTTCTATGTGGCCCTTTGTCTCCTGTGGACAGGACACA TGGATGCTGGAATCACCCAGAGCCCAAGACACAAGGTCACAGAGACAGGAACACC AGTGACTCTGAGATGTCAACAGACTGAGAACCACCGCTATATGTACTGGTATCGA CAAGACCCGGGGCATGGGCTGAGGCTGATCCATTACTCATATGGTGT TAAAGATA CTGACAAAGGAGAAGTCTCAGATGGCTATAGTGTCTCTAGATCAAAGACAGAGGA TTTCCTCCTCACTCTGGAGTCCGCTACCAGCTCCCAGACATCTGTGTACTTCTGT GCCATCTCGGTGGGACAGGGGGCCCTCTACGAGCAGTACTTCGGGCCGGGCACCA GGCTCACGGTCACAGAGGACCTGAACAAGGTGTTCCCAACCCGAGGTCGCTGTGTT TGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGCCTG GCCACAGGCTTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGAAGG AGGTGCACAGTGGGGTCAGCACGGACCCGCAGCCCCCTCAAGGAGCAGCCCGCCCT CAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTCTGG CAGAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGA ATGACGAGTGGACCCAGGATAGGGCCAAACCCGTCACCCAGATCGTCAGCGCCGA GGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCCTACCAGCAAGGGGTC CTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCCTGTATGCTG TGCTGGTCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAGGATTTTC	SEQ ID NO: 153
	β1 (with TRBC2)	ATGGGCACAAGGTTGTTCTTCTATGTGGCCCTTTGTCTCCTGTGGACAGGACACA TGGATGCTGGAATCACCCAGAGCCCAAGACACAAGGTCACAGAGACAGGAACACC AGTGACTCTGAGATGTCAACAGACTGAGAACCACCGCTATATGTACTGGTATCGA CAAGACCCGGGGCATGGGCTGAGGCTGATCCATTACTCATATGGTGT TAAAGATA CTGACAAAGGAGAAGTCTCAGATGGCTATAGTGTCTCTAGATCAAAGACAGAGGA TTTCCTCCTCACTCTGGAGTCCGCTACCAGCTCCCAGACATCTGTGTACTTCTGT GCCATCTCGGTGGGACAGGGGGCCCTCTACGAGCAGTACTTCGGGCCGGGCACCA GGCTCACGGTCACAGAGGACCTGAAAAACGTGTTCCCAACCCGAGGTCGCTGTGTT TGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGCCTG GCCACAGGCTTCTACCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGAAGG AGGTGCACAGTGGGGTCAGCACAGACCCGCAGCCCCCTCAAGGAGCAGCCCGCCCT CAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTCTGG CAGAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGA ATGACGAGTGGACCCAGGATAGGGCCAAACCTGTCACCCAGATCGTCAGCGCCGA GGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCCGAGTCTTACCAGCAAGGGGTC CTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCCTGTATGCCG TGCTGGTCAGTGCCCTCGTGCTGATGGCCATGGTCAAGAGAAAGGATTCCAGAGG C	SEQ ID NO: 154
	β2 (with TRBC1)	ATGGGCTTCAGGCTCCTCTGCTGTGTGGCCTTTTGTCTCCTGGGAGCAGGCCCCAG TGGATTCTGGAGTCACACAAACCCCAAAGCACCTGATCACAGCAACTGGACAGCG AGTGACGCTGAGATGCTCCCCTAGGTCTGGAGACCTCTCTGTGTACTGGTACCAA CAGAGCCTGGACCAGGGCCTCCAGTTCCCTCATT CAGTATTATAATGGAGAAGAGA GAGCAAAAGGAAACATTCTTGAACGATTCTCCGCACAACAGTTCCCTGACTTGCA CTCTGAACTAAACCTGAGCTCTCTGGAGCTGGGGGACTCAGCTTTGTATTTCTGT GCCAGCAGCGTAGCTCGGGACAGGCGGAACCTATGGCTACACCTTCGGTTCGGGGA CCAGGTTAACCGTTGTAGAGGACCTGAACAAGGTGTTCCCAACCCGAGGTCGCTGT GTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGC CTGGCCACAGGCTTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGA AGGAGGTGCACAGTGGGGTCAGCACGGACCCGCAGCCCCCTCAAGGAGCAGCCCGC CCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTC TGGCAGAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGG AGAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTCACCCAGATCGTCAGCGC CGAGGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCCTACCAGCAAGGG GTCTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCCTGTATG CTGTGCTGGTCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAGGATTTTC	SEQ ID NO: 155
	β2 (with TRBC2)	ATGGGCTTCAGGCTCCTCTGCTGTGTGGCCTTTTGTCTCCTGGGAGCAGGCCCCAG TGGATTCTGGAGTCACACAAACCCCAAAGCACCTGATCACAGCAACTGGACAGCG AGTGACGCTGAGATGCTCCCCTAGGTCTGGAGACCTCTCTGTGTACTGGTACCAA CAGAGCCTGGACCAGGGCCTCCAGTTCCCTCATT CAGTATTATAATGGAGAAGAGA GAGCAAAAGGAAACATTCTTGAACGATTCTCCGCACAACAGTTCCCTGACTTGCA CTCTGAACTAAACCTGAGCTCTCTGGAGCTGGGGGACTCAGCTTTGTATTTCTGT GCCAGCAGCGTAGCTCGGGACAGGCGGAACCTATGGCTACACCTTCGGTTCGGGGA CCAGGTTAACCGTTGTAGAGGACCTGAACAAGGTGTTCCCAACCCGAGGTCGCTGT GTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTGTGC CTGGCCACAGGCTTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGA AGGAGGTGCACAGTGGGGTCAGCACGGACCCGCAGCCCCCTCAAGGAGCAGCCCGC CCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTC TGGCAGAACCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGG AGAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTCACCCAGATCGTCAGCGC CGAGGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCCTACCAGCAAGGG GTCTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCCTGTATG CTGTGCTGGTCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAGGATTTTC	SEQ ID NO: 156

		CCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCCTGAGGGTCTCGGCCACCTTC TGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGG AGAATGACGAGTGGACCCAGGATAGGGCCAAACCTGTCACCCAGATCGTCAGCGC CGAGGCCTGGGGTAGAGCAGACTGTGGCTTACCTCCGAGTCTTACCAGCAAGGG GTCCTGTCTGCCACCATCCTCTATGAGATCTTGCTAGGGAAGGCCACCTTGTATG CCGTGCTGGTCAGTGCCCTCGTGCTGATGGCCATGGTCAAGAGAAAGGATTCCAG AGGC	
HD6	α1 (with TRAC)	atggccatgctcctgggggcatcagtgctgattctgtggcttcagccagactggg taaacagtcaacagaagaatgatgaccagcaagttaagcaaaattcaccatccct gagcgtccaggaaggaagaatttctattctgaactgtgactataactaacagcatg tttgattatttcctatggtacaaaaaataccctgctgaaggctcctacattcctga tatctataagttccattaaaggataaaaaatgaagatggaagattcactgtcttctt aaacaaaagtgccaaagcacctctctctgcacattgtgccctcccagcctggagac tctgcagtgtacttctgtgcagcaaacaatgccagactcatgtttggagatggaa ctcagctggtggtgaagcccaatatccagaaccctgaccctgccgtgtaccagct gagagactctaaatccagtgacaagtctgtctgcctattcaccgattttgattct caaacaaatgtgtcacaaagtaaggattctgatgtgtatatcacagacaaaactg tgctagacatgaggtctatggacttcaagagcaacagtgctgtggcctggagcaa caaatctgactttgcatgtgcaaacgccttcaacaacagcattattccagaagac accttcttccccagcccagaaaagttcctgtgatgtcaagctggtcgagaaaagct ttgaaacagatacgaacctaaactttcaaaacctgtcagtgattggggtccgaat cctcctcctgaaagtggccgggtttaatctgctcatgacgctgcggctgtggtcc agc	SEQ ID NO: 223
	α2 (with TRAC)	ATGGCCATGCTCCTGGGGGCATCAGTGCTGATTCTGTGGCTTCAGCCAGACTGGG TAAACAGTCAACAGAAGAATGATGACCAGCAAGTTAAGCAAAATTCACCATCCCT GAGCGTCCAGGAAGGAAGAATTTCTATTCTGAACTGTGACTATACTAACAGCATG TTTGATTATTTCCCTATGGTACAAAAAATACCCTGCTGAAGGTCCTACATTCTGA TATCTATAAGTTCCATTAAGGATAAAAAATGAAGATGGAAGATTCAGTGTCTTCTT AAACAAAAGTGCCAAGCACCTCTCTCTGCACATTGTGCCCTCCCAGCCTGGAGAC TCTGCAGTGTACTTCTGTGCAGCAAGCGCTACCGGTAACCAGTTCTATTTTGGA CAGGGACAAGTTTGACGGTCATTCCAAATATCCAGAACCCTGACCCTGCCGTGTA CCAGCTGAGAGACTCTAAATCCAGTGACAAGTCTGTCTGCCTATTCACCGATTTT GATTCTCAAACAAATGTGTACAAAGTAAGGATTCTGATGTGTATATCACAGACA AACTGTGCTAGACATGAGGTCTATGGACTTCAAGAGCAACAGTGCTGTGGCCTG GAGCAACAAATCTGACTTTGCATGTGCAAACGCCTTCAACAACAGCATTATTCCA GAAGACACCTTCTTCCCCAGCCCAGAAAAGTTCTGTGATGTCAAGCTGGTCGAGA AAAGCTTTGAAACAGATACGAACCTAAACTTTCAAAACCTGTCAGTGATTGGGTT CCGAATCCTCCTCCTGAAAGTGCCGGGTTTAATCTGCTCATGACGCTGCGGCTG TGGTCCAGCTGA	SEQ ID NO: 292
	β1 (with TRBC1)	atggactcctggaccttctgctgtgtgtccctttgcatcctggtagcgaagcata cagatgctggagttatccagtcaccccgccatgaggtgacagagatgggacaaga agtgactctgagatgtaaaccaatttcaggccacaaactcccttttctggtacaga cagaccatgatgcggggactggagttgctcatttactttaacaacaacgttccga tagatgattcagggatgcccgaggatcgattctcagctaagatgcctaataatgcac attctccactctgaagatccagccctcagaacccagggaactcagctgtgtacttc tgtgccagcagtgataaccaggggcccgggagcagttcttcggggccagggaacacggc tcaccgtgctagaggacctgaacaaggtgttcccacccgaggtcgctgtgtttga gccatcagaagcagagatctcccacacccaaaaggccacactggtgtgcctggcc acaggcttcttccccgaccacgtggagctgagctggtgggtgaatgggaaggagg tgcacagtggggtcagcacaggacccgcagccctcaaggagcagcccgcctcaa tgactccagatactgcctgagcagccgcctgagggctctcgccaccttctggcag aacccccgcaaccacttccgctgtcaagtccagttctacgggctctcggagaatg acgagtggaccaggatagggccaaacctgtcaccagatcgtcagcgccgaggc ctggggtagagcagactgtggctttacctcgggtgtcctaccagcaaggggtcctg tctgccaccatcctctatgagatcctgctagggaaggccacctgtatgctgtgc tggtcagcgcccttggtgttgatggccatggtcaagagaaaggatttc	SEQ ID NO: 224
	β1 (with TRBC2)	atggactcctggaccttctgctgtgtgtccctttgcatcctggtagcgaagcata cagatgctggagttatccagtcaccccgccatgaggtgacagagatgggacaaga agtgactctgagatgtaaaccaatttcaggccacaaactcccttttctggtacaga cagaccatgatgcggggactggagttgctcatttactttaacaacaacgttccga tagatgattcagggatgcccgaggatcgattctcagctaagatgcctaataatgcac attctccactctgaagatccagccctcagaacccagggaactcagctgtgtacttc tgtgccagcagtgataaccaggggcccgggagcagttcttcggggccagggaacacggc tcaccgtgctagaggacctgaaaaacgtgttcccacccgaggtcgctgtgtttga gccatcagaagcagagatctcccacacccaaaaggccacactggtgtgcctggcc acaggcttctaccccgaccacgtggagctgagctggtgggtgaatgggaaggagg tgcacagtggggtcagcacagacccgcagccctcaaggagcagcccgcctcaa tgactccagatactgcctgagcagccgcctgagggctctcgccaccttctggcag aacccccgcaaccacttccgctgtcaagtccagttctacgggctctcggagaatg acgagtggaccaggatagggccaaacctgtcaccagatcgtcagcgccgaggc ctggggtagagcagactgtggctttacctcgggtgtcctaccagcaaggggtcctg tctgccaccatcctctatgagatcctgctagggaaggccacctgtatgccgtgc tggtcagtgccctcgtgctgatggccatggtcaagagaaaggattccagaggc	SEQ ID NO: 225
	β2 (with TRBC1)	ATGGGCTCCAGGCTGCTCTGTTGGGTGCTGCTTTGTCTCCTGGGAGCAGGCCAG TAAAGGCTGGAGTCACTCAAACCTCAAGATATCTGATCAAAACGAGAGGACAGCA	SEQ ID NO: 293

		AGTGACACTGAGCTGCTCCCCTATCTCTGGGCATAGGAGTGTATCCTGGTACCAA CAGACCCCAGGACAGGGCCTTCAGTTCCTCTTTGAATACTTCAGTGAGACACAGA GAAACAAAGGAAACTTCCCTGGTCGATTCTCAGGGCGCCAGTTCTCTAACTCTCG CTCTGAGATGAATGTGAGCACCTTGGAGCTGGGGGACTCGGCCCTTTATCTTTGC GCCAGCAGCCCTGGACAGCACGGGGAGCTGTTTTTTGGAGAAGGCTCTAGGCTGA CCGTACTGGAGGACCTGAACAAGGTGTTCCCACCCGAGGTCGCTGTGTTTGAGCC ATCAGAAGCAGAGATCTCCCACACCCAAAAGGCCACACTGGTGTGCCTGGCCACA GGCTTCTTCCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGAAGGAGGTGC ACAGTGGGGTCAGCACGGACCCGCAGCCCCTCAAGGAGCAGCCCGCCCTCAATGA CTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTCTGGCAGAAC CCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGAATGACG AGTGGACCCAGGATAGGGCCAAACCCGTCACCCAGATCGTCAGCGCCGAGGCCTG GGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCTTACCAGCAAGGGGTCTGTCT GCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCCTGTATGCTGTGCTGG TCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAAGGATTTCTC	
	β2 (with TRBC2)	ATGGGCTCCAGGCTGCTCTGTTGGGTGCTGCTTTGTCTCCTGGGAGCAGGCCCAG TAAAGGCTGGAGTCACTCAAACCTCCAAGATATCTGATCAAAACGAGAGGACAGCA AGTGACACTGAGCTGCTCCCCTATCTCTGGGCATAGGAGTGTATCCTGGTACCAA CAGACCCCAGGACAGGGCCTTCAGTTCCTCTTTGAATACTTCAGTGAGACACAGA GAAACAAAGGAAACTTCCCTGGTCGATTCTCAGGGCGCCAGTTCTCTAACTCTCG CTCTGAGATGAATGTGAGCACCTTGGAGCTGGGGGACTCGGCCCTTTATCTTTGC GCCAGCAGCCCTGGACAGCACGGGGAGCTGTTTTTTGGAGAAGGCTCTAGGCTGA CCGTACTGGAGGACCTGAAAAACGTGTTCCCACCCGAGGTCGCTGTGTTTGAGCC ATCAGAAGCAGAGATCTCCCACACCCAAAAGGCCACACTGGTGTGCCTGGCCACA GGCTTCTACCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATGGGAAGGAGGTGC ACAGTGGGGTCAGCACAGACCCGCAGCCCCTCAAGGAGCAGCCCGCCCTCAATGA CTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACCTTCTGGCAGAAC CCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCTCGGAGAATGACG AGTGGACCCAGGATAGGGCCAAACCTGTCACCCAGATCGTCAGCGCCGAGGCCTG GGGTAGAGCAGACTGTGGCTTCACCTCCGAGTCTTACCAGCAAGGGGTCTGTCT GCCACCATCCTCTATGAGATCTTGCTAGGGAAGGCCACCCTGTATGCCGTGCTGG TCAGTGGCCCTCGTGCTGATGGCCATGGTCAAGAGAAAAGGATTCCAGAGGC	SEQ ID NO: 294
HD7	α1 (with TRAC)	atgaagacatttgctggattttcggttcctggttttgtggctgcagctggactgta tgagtagaggagaggatgtggagcagagtcttttcctgagtgtccgagagggaga cagctccgttataaaactgcacttacacagacagctcctccacctacttatactgg tataagcaagaacctggagcaggtctccagttgctgacgtatattttttcaaata tggacatgaaacaagaccaaaagactcactgttctattgaataaaaaggataaaca tctgtctctgcgcattgcagacacccagactggggactcagctatctacttctgt gcagagaggcttaacaccgcacaagctcatctttgggactgggaccagattacaag tctttccaaatatccagaaccttgacctgcccgtgtaccagctgagagactctaa atccagtgacaagtctgtctgcctattcaccgattttgattctcaaacaaatgtg tcacaaagtaaggattctgatgtgtatatcacagacaaaactgtgctagacatga ggtctatggacttcaagagcaacagtgctgtggcctggagcaacaaatctgactt tgcatgtgcaaacgccttcaacaacagcattattccagaagacaccttcttcccc agcccagaaaagttcctgtgatgtcaagctggtcgagaaaagctttgaaacagata cgaacctaaactttcaaaacctgtcagtgattgggttccgaatcctcctcctgaa agtggccgggtttaatctgctcatgacgctgcggctgtggtccagc	SEQ ID NO: 226
	α2 (with TRAC)	atgaagaggatattgggagctctgctggggctcttgagtgcccagggttgctgtg tgagaggaatacaagtggagcagagtctccagacctgattctccaggagggagc caattccacgctgccgtgcaatttttctgactctgtgaacaatttgcagtggttt catcaaaaccttggggacagctcatcaacctgttttacattccctcagggacaa aacagaatggaagattaagcgccacgactgtcgctacggaacgctacagcttatt gtacatttctcttcccagaccacagactcaggcggtttatttctgtgctgtggag gcaactgacagctgggggaaattgcagtttgagcagggacccagggttggtgtca ccccagatatccagaaccttgacctgcccgtgtaccagctgagagactctaaatc cagtgacaagtctgtctgcctattcaccgattttgattctcaaacaaatgtgtca caaagtaaggattctgatgtgtatatcacagacaaaactgtgctagacatgaggt ctatggacttcaagagcaacagtgctgtggcctggagcaacaaatctgactttgc atgtgcaaacgccttcaacaacagcattattccagaagacaccttcttccccagc ccagaaaagttcctgtgatgtcaagctggtcgagaaaagctttgaaacagatacga acctaaaactttcaaaacctgtcagtgattgggttccgaatcctcctcctgaaagt ggccgggtttaatctgctcatgacgctgcccgtgtggtccagc	SEQ ID NO: 227
	α3 (with TRAC)	atgacatccattcgagctgtattttatattcctgtggctgcagctggacttggtga atggagagaatgtggagcagcatccttcaacctgagtgtccaggaggggagacag cgctgttatcaagtgtacttattcagacagtgccctcaaactacttcccttggtat aagcaagaacttggaaaaagacctcagcttattatagacattcgttcaaagtgtgg gcgaaaagaaagaccaacgaattgctgttacattgaacaagacagccaaacattt ctccctgcacatcacagagacccaacctgaagactcggctgtctacttctgtgca gtacgaacctcctacgacaaggatgatatttggggccagggaagcttatcagtc ttccaaatatccagaaccttgacctgcccgtgtaccagctgagagactctaaatc cagtgacaagtctgtctgcctattcaccgattttgattctcaaacaaatgtgtca caaagtaaggattctgatgtgtatatcacagacaaaactgtgctagacatgaggt ctatggacttcaagagcaacagtgctgtggcctggagcaacaaatctgactttgc atgtgcaaacgccttcaacaacagcattattccagaagacaccttcttccccagc ccagaaaagttcctgtgatgtcaagctggtcgagaaaagctttgaaacagatacga ccctaaaactttcaaaacctgtcagtgattgggttccgaatcctcctcctgaaagt ggccgggtttaatctgctcatgacgctgcccgtgtggtccagc	SEQ ID NO: 228

		acctaaactttcaaaacctgtcagtgattgggttccgaatcctcctcctgaaagt ggccgggttttaatctgctcatgacgctgcggtgtggtccagc	
	α-4 (with TRAC)	ATGGAAACTCTCCTGGGAGTGTCTTTGGTGATTCTATGGCTTCAACTGGCTAGGG TGAACAGTCAACAGGGAGAAGAGGATCCTCAGGCCTTGAGCATCCAGGAGGGTGA AAATGCCACCATGAACTGCAGTTACAAAAGTAGTATAAAACAATTTACAGTGGTAT AGACAAAATTCAGGTAGAGGCCTTGTCACCTAATTTTAATACGTTCAAATGAAA GAGAGAAACACAGTGGAAGATTAAGAGTCACGCTTGACACTTCCAAGAAAAGCAG TTCCTTGTTGATCACGGCTTCCCGGGCAGCAGACACTGCTTCTTACTTCTGTGCT ACGGACGGGGATAGCAGCTATAAATTGATCTTCGGGAGTGGGACCAGACTGCTGG TCAGGCCTGATATCCAGAACCCTGACCCTGCCGTGTACCAGCTGAGAGACTCTAA ATCCAGTGACAAGTCTGTCTGCCTATTCACCGATTTTGATTCTCAAACAAATGTG TCACAAAGTAAGGATTCTGATGTGTATATCACAGACAAAAGTGTGCTAGACATGA GGTCTATGGACTTCAAGAGCAACAGTGCTGTGGCCTGGAGCAACAAATCTGACTT TGCATGTGCAAACGCCTTCAACAACAGCATTATTCCAGAAGACACCTTCTTCCCC AGCCCAGAAAGTTCCTGTGATGTCAAGCTGGTCGAGAAAAGCTTTGAAACAGATA CGAACCTAAACTTTCAAACCTGTCAGTGATTGGGTTCGAATCCTCCTCCTGAA AGTGGCCGGGTTTAATCTGCTCATGACGCTGCGGCTGTGGTCCAGC	SEQ ID NO: 295
	β1 (with TRBC1)	atgctgctgcttctgctgcttctggggccaggtataagcctccttctacctggga gcttggcaggctccgggcttggtgctgtcgtctctcaacatccgagctgggttat ctgtaagagtggaacctctgtgaagatcgagtgccgttccctggactttcaggcc acaactatgttttggtatcgtcagttcccgaacagagtctcatgctgatggcaa cttccaatgagggctccaaggccacatacgagcaaggcgtcgagaaggacaagtt tctcatcaaccatgcaagcctgaccttgtccactctgacagtgaccagtgcccat cctgaagacagcagcttctacatctgcagtgctagggacagtgtgtctggaaaca ccatatattttgagaggggaagttggctcactgttgtagaggacctgaacaaggt gttcccacccgaggtegctgtgtttgagccatcagaagcagagatctcccacacc caaaaggccacactggtgtgcctggccacaggcttcttcccgaccacgtggagc tgagctggtgggtgaatgggaaggaggtgcacagtggggtcagcacggacccgca gcccctcaaggagcagcccgccctcaatgactccagatactgcctgagcagccgc ctgaggggtctcggccaccttctggcagaacccccgcaaccacttccgctgtcaag tccagttctacgggctctcggagaatgacgagtggaccaggatagggccaaacc cgtcaccagatcgtcagcgccgaggcctggggtagagcagactgtggctttacc tcggtgtcctaccagcaaggggtcctgtctgccaccatcctctatgagatcctgc tagggaaggccacctgtatgctgtgctggtcagcgcccttgtgttgatggccat ggtcaagagaaaggatttc	SEQ ID NO: 229
	β1 (with TRBC2)	atgctgctgcttctgctgcttctggggccaggtataagcctccttctacctggga gcttggcaggctccgggcttggtgctgtcgtctctcaacatccgagctgggttat ctgtaagagtggaacctctgtgaagatcgagtgccgttccctggactttcaggcc acaactatgttttggtatcgtcagttcccgaacagagtctcatgctgatggcaa cttccaatgagggctccaaggccacatacgagcaaggcgtcgagaaggacaagtt tctcatcaaccatgcaagcctgaccttgtccactctgacagtgaccagtgcccat cctgaagacagcagcttctacatctgcagtgctagggacagtgtgtctggaaaca ccatatattttgagaggggaagttggctcactgttgtagaggacctgaaaaacgt gttcccacccgaggtegctgtgtttgagccatcagaagcagagatctcccacacc caaaaggccacactggtgtgcctggccacaggcttctaccccgaccacgtggagc tgagctggtgggtgaatgggaaggaggtgcacagtggggtcagcacagacccgca gcccctcaaggagcagcccgccctcaatgactccagatactgcctgagcagccgc ctgaggggtctcggccaccttctggcagaacccccgcaaccacttccgctgtcaag tccagttctacgggctctcggagaatgacgagtggaccaggatagggccaaacc tgtcaccagatcgtcagcgccgaggcctggggtagagcagactgtggcttcacc tccgagtcttaccagcaaggggtcctgtctgccaccatcctctatgagatcttgc tagggaaggccacctgtatgccgtgctggtcagtgccctcgtgctgatggccat ggtcaagagaaaggattccagaggc	SEQ ID NO: 230
	β2 (with TRBC1)	atgctgagtccttctgctccttctcctgggactaggctctgtgttcagtgctgtca tctctcaaaagccaagcaggggatatctgtcaacgtggaacctcctgacgatcca gtgtcaagtcgatagccaagtaccatgatgttctggtaccgtcagcaacctgga cagagcctgacactgatcgcaactgcaaatacagggtctgaggccacatatgaga gtggatttgtcattgacaagtttcccatcagccgccccaaacctaacattctcaac tctgactgtgagcaacatgagccctgaagacagcagcatatatctctgcagcgtt gggggtagcgggagttacaatgagcagttcttcgggccagggacacggctcaccg tgctagaggacctgaacaagggtgttcccacccgaggtcgctgtgtttgagccatc agaagcagagatctcccacacccaaaaggccacactggtgtgcctggccacaggc ttcttcccgcaccacgtggagctgagctggtgggtgaatgggaaggaggtgcaca gtggggtcagcacggacccgcagccctcaaggagcagcccgccctcaatgactc cagatactgcctgagcagccgcctgagggtctcggccaccttctggcagaacccc cgcaaccacttccgctgtcaagtccagttctacgggctctcggagaatgacgagt ggaccaggatagggccaaacccgtcaccagatcgtcagcgccgaggcctgggg tagagcagactgtggctttacctcggtgtcctaccagcaaggggtcctgtctgcc accatcctctatgagatcctgctagggaaggccacctgtatgctgtgctggtca gcgcccttgtgttgatggccatggtcaagagaaaggatttc	SEQ ID NO: 231
	β2 (with TRBC2)	atgctgagtccttctgctccttctcctgggactaggctctgtgttcagtgctgtca tctctcaaaagccaagcaggggatatctgtcaacgtggaacctcctgacgatcca gtgtcaagtcgatagccaagtaccatgatgttctggtaccgtcagcaacctgga cagagcctgacactgatcgcaactgcaaatacagggtctgaggccacatatgaga gtggatttgtcattgacaagtttcccatcagccgccccaaacctaacattctcaac tctgactgtgagcaacatgagccctgaagacagcagcatatatctctgcagcgtt gggggtagcgggagttacaatgagcagttcttcgggccagggacacggctcaccg tgctagaggacctgaacaagggtgttcccacccgaggtcgctgtgtttgagccatc agaagcagagatctcccacacccaaaaggccacactggtgtgcctggccacaggc ttcttcccgcaccacgtggagctgagctggtgggtgaatgggaaggaggtgcaca gtggggtcagcacggacccgcagccctcaaggagcagcccgccctcaatgactc cagatactgcctgagcagccgcctgagggtctcggccaccttctggcagaacccc cgcaaccacttccgctgtcaagtccagttctacgggctctcggagaatgacgagt ggaccaggatagggccaaacccgtcaccagatcgtcagcgccgaggcctgggg tagagcagactgtggctttacctcggtgtcctaccagcaaggggtcctgtctgcc accatcctctatgagatcctgctagggaaggccacctgtatgctgtgctggtca gcgcccttgtgttgatggccatggtcaagagaaaggatttc	SEQ ID NO: 232

		gggggtagcgggagttacaatgagcagttcttcggggccagggacacgggtcaccg tgctagaggacctgaaaaacgtgttcccacccgaggtcgctgtgtttgagccatc agaagcagagatctcccacacccaaaaggccacactggtgtgcctggccacaggc ttctaccccgaccacgtggagctgagctggtgggtgaatgggaaggaggtgcaca gtggggtcagcacagacccgcagcccctcaaggagcagcccgccctcaatgactc cagatactgcctgagcagccgcctgaggggtctcgggcaccttctggcagaacccc cgcaaccacttccgctgtcaagtccagttctacgggctctcggagaatgacgagt ggaccacaggatagggccaaacctgtcacccagatcgtcagcgccgaggcctgggg tagagcagactgtggcttcacctccgagtcttaccagcaaggggtcctgtctgcc accatcctctatgagatcttgctaggggaaggccaccttgatgccgtgctggtca gtgccctcgtgctgatggccatggtcaagagaaaggattccagaggc	
	β-3 (with TRBC1)	ATGCTGCTGCTTCTGCTGCTTCTGGGGCCAGGTATAAGCCTCCTTCTACCTGGGA GCTTGGCAGGCTCCGGGCTTGGTGCTGTGCTCTCTCAACATCCGAGCTGGGTTAT CTGTAAGAGTGGAACCTCTGTGAAGATCGAGTGCCGTTCCCTGGACTTTCAGGCC ACAACATATGTTTTGGTATCGTCAGTTCCCGAAACAGAGTCTCATGCTGATGGCAA CTTCCAATGAGGGCTCCAAGGCCACATACGAGCAAGGCGTCGAGAAGGACAAGTT TCTCATCAACCATGCAAGCCTGACCTTGTCCACTCTGACAGTGACCAGTGCCCAT CCTGAAGACAGCAGCTTCTACATCTGCAGTGCTAGAGACGTACTGACAGGGGACT ATGGCTACACCTTCGGTTCGGGGACCAGGTAAACCGTTGTAGAGGACCTGAACAA GGTGTTCACCCAGGTCGCTGTGTTTGAGCCATCAGAAGCAGAGATCTCCAC ACCCAAAAGGCCACACTGGTGTGCCTGGCCACAGGCTTCTTCCCGACCACGTGG AGCTGAGCTGGTGGGTGAATGGGAAGGAGGTGCACAGTGGGGTCAGCACGGACCC GCAGCCCCCTCAAGGAGCAGCCCCGCCCTCAATGACTCCAGATACTGCCTGAGCAGC CGCCTGAGGGTCTCGGCCACCTTCTGGCAGAACCCCCGCAACCACTTCCGCTGTC AAGTCCAGTTCTACGGGCTCTCGGAGAATGACGAGTGGACCCAGGATAGGGCCAA ACCCGTCACCCAGATCGTCAGCGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTT ACCTCGGTGTCCTACCAGCAAGGGTCTGTCTGCCACCATCCTCTATGAGATCC TGCTAGGGAAGGCCACCTGTATGCTGTGCTGGTCAGCGCCCTTGTGTTGATGGC CATGGTCAAGAGAAAGGATTTTC	SEQ ID NO: 296
	β-3 (with TRBC2)	ATGCTGCTGCTTCTGCTGCTTCTGGGGCCAGGTATAAGCCTCCTTCTACCTGGGA GCTTGGCAGGCTCCGGGCTTGGTGCTGTGCTCTCTCAACATCCGAGCTGGGTTAT CTGTAAGAGTGGAACCTCTGTGAAGATCGAGTGCCGTTCCCTGGACTTTCAGGCC ACAACATATGTTTTGGTATCGTCAGTTCCCGAAACAGAGTCTCATGCTGATGGCAA CTTCCAATGAGGGCTCCAAGGCCACATACGAGCAAGGCGTCGAGAAGGACAAGTT TCTCATCAACCATGCAAGCCTGACCTTGTCCACTCTGACAGTGACCAGTGCCCAT CCTGAAGACAGCAGCTTCTACATCTGCAGTGCTAGAGACGTACTGACAGGGGACT ATGGCTACACCTTCGGTTCGGGGACCAGGTAAACCGTTGTAGAGGACCTGAAAAA CGTGTTCACCCAGGTCGCTGTGTTTGAGCCATCAGAAGCAGAGATCTCCAC ACCCAAAAGGCCACACTGGTGTGCCTGGCCACAGGCTTCTACCCCGACCACGTGG AGCTGAGCTGGTGGGTGAATGGGAAGGAGGTGCACAGTGGGGTCAGCACAGACCC GCAGCCCCCTCAAGGAGCAGCCCCGCCCTCAATGACTCCAGATACTGCCTGAGCAGC CGCCTGAGGGTCTCGGCCACCTTCTGGCAGAACCCCCGCAACCACTTCCGCTGTC AAGTCCAGTTCTACGGGCTCTCGGAGAATGACGAGTGGACCCAGGATAGGGCCAA ACCTGTCACCCAGATCGTCAGCGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTC ACCTCCGAGTCTTACCAGCAAGGGTCTGTCTGCCACCATCCTCTATGAGATCT TGCTAGGGAAGGCCACCTTGTATGCCGTGCTGGTCAGTGCCCTCGTGCTGATGGC CATGGTCAAGAGAAAGGATTCCAGAGGC	SEQ ID NO: 297
	β-4 (with TRBC1)	ATGGGCTCCTGGACCCTCTGCTGTGTGTCCCTTTGCATCCTGGTAGCAAAGCACA CAGATGCTGGAGTTATCCAGTCACCCCGGCACGAGGTGACAGAGATGGGACAAGA AGTGACTCTGAGATGTAAACCAATTTCAGGACACGACTACCTTTTCTGGTACAGA CAGACCATGATGCGGGGACTGGAGTTGCTCATTTACTTTAACAACAACGTTCCGA TAGATGATTACAGGGATGCCCCGAGGATCGATTCTCAGCTAAGATGCCTAATGCATC ATTCTCCACTCTGAAGATCCAGCCCTCAGAACCCAGGGACTCAGCTGTGTACTTC TGTGCCAGCAGTTTAGGACTGAGCATTTCCCAAGAGACCCAGTACTTCGGGCCAG GCACGCGGCTCCTGGTGCTCGAGGACCTGAACAAGGTGTTCCACCCGAGGTGCG TGTGTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTG TGCCTGGCCACAGGCTTCTTCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATG GGAAGGAGGTGCACAGTGGGGTCAGCACGGACCCGCAGCCCCTCAAGGAGCAGCC CGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACC TTCTGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCT CGGAGAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTACCCAGATCGTCAG CGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCCTACCAGCAA GGGGTCCTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCTGT ATGCTGTGCTGGTCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAGGATTT C	SEQ ID NO: 298
	β-4 (with TRBC2)	ATGGGCTCCTGGACCCTCTGCTGTGTGTCCCTTTGCATCCTGGTAGCAAAGCACA CAGATGCTGGAGTTATCCAGTCACCCCGGCACGAGGTGACAGAGATGGGACAAGA AGTGACTCTGAGATGTAAACCAATTTCAGGACACGACTACCTTTTCTGGTACAGA CAGACCATGATGCGGGGACTGGAGTTGCTCATTTACTTTAACAACAACGTTCCGA TAGATGATTACAGGGATGCCCCGAGGATCGATTCTCAGCTAAGATGCCTAATGCATC ATTCTCCACTCTGAAGATCCAGCCCTCAGAACCCAGGGACTCAGCTGTGTACTTC TGTGCCAGCAGTTTAGGACTGAGCATTTCCCAAGAGACCCAGTACTTCGGGCCAG GCACGCGGCTCCTGGTGCTCGAGGACCTGAACAAGGTGTTCCACCCGAGGTGCG TGTGTTTGAGCCATCAGAAGCAGAGATCTCCACACCCAAAAGGCCACACTGGTG TGCCTGGCCACAGGCTTCTTCCCGACCACGTGGAGCTGAGCTGGTGGGTGAATG GGAAGGAGGTGCACAGTGGGGTCAGCACGGACCCGCAGCCCCTCAAGGAGCAGCC CGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACC TTCTGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCT CGGAGAATGACGAGTGGACCCAGGATAGGGCCAAACCCGTACCCAGATCGTCAG CGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTTACCTCGGTGTCCTACCAGCAA GGGGTCCTGTCTGCCACCATCCTCTATGAGATCCTGCTAGGGAAGGCCACCTGT ATGCTGTGCTGGTCAGCGCCCTTGTGTTGATGGCCATGGTCAAGAGAAAGGATTT C	SEQ ID NO: 299

		CGCCCTCAATGACTCCAGATACTGCCTGAGCAGCCGCCTGAGGGTCTCGGCCACC TTCTGGCAGAACCCCCGCAACCACTTCCGCTGTCAAGTCCAGTTCTACGGGCTCT CGGAGAATGACGAGTGGACCCAGGATAGGGCCAAACCTGTCACCCAGATCGTCAG CGCCGAGGCCTGGGGTAGAGCAGACTGTGGCTTCACCTCCGAGTCTTACCAGCAA GGGGTCCTGTCTGCCACCATCCTCTATGAGATCTTGCTAGGGAAGGCCACCTTGT ATGCCGTGCTGGTCAGTGCCCTCGTGCTGATGGCCATGGTCAAGAGAAAGGATTC CAGAGGC	
HD8	α (with TRAC)	atgggtgaagatccggcaatttttgttggctattttgtggcttcagctaagctgtg taagtgccgccaaaaatgaagtggagcagagtcctcagaacctgactgcccagga aggagaatttatcacaatcaactgcagttactcggtaggaataagtgccttacac tggctgcaacagcatccaggaggaggcattgtttccttgtttatgctgagctcag ggaagaagaagcatggaagattaattgccacaataaacatacaggaaaagcacag ctccctgcacatcacagcctccatcccagagactctgccgtctacatctgtgct gtcacagtcggaacaaaactggctcttggcgcgaggaaccattctgagagtcaagt cctatatccagaacctgacctgccgtgtaccagctgagagactctaaatccag tgacaagtctgtctgcctattcaccgattttgattctcaaacaaatgtgtcacia agtaaggattctgatgtgtatatcacagacaaaactgtgctagacatgaggtcta tggacttcaagagcaacagtgctgtggcctggagcaacaaatctgactttgcatg tgcaaacgccttcaacaacagcattattccagaagacaccttcttcccagccca gaaagttcctgtgatgtcaagctggtcgagaaaagctttgaaacagatacgaacc taaactttcaaaacctgtcagtgattgggttccgaatcctcctcctgaaagtggc cgggtttaatctgctcatgacgctgcggctgtggtccagc	SEQ ID NO: 233
	β (with TRBC1)	atgagcatcgggctcctgtgctgtgtggccttttctctcctgtgggcaagtccag tgaatgctggtgtcactcagacccccaaaattccagggtcctgaagacaggacagag catgacactgcagtggtgccaggatatgaaccataactccatgtactggtatcga caagacccaggcatgggactgaggctgatttattactcagcttctgagggtagca ctgacaaaaggagaagtccccaatggctacaatgtctccagattaaacaaacggga gttctcgctcaggctggagtcggctgctccctcccagacatctgtgtacttctgt gccagcaggggggtggcgtgagcagttcttcgggccagggaacaggctcacctgtc tagaggacctgaacaagggtgttcccaccgaggtcgctgtgtttgagccatcaga agcagagatctcccacacccaaaaggccacactggtgtgcctggccacaggcttc ttcccgcaccacgtggagctgagctggtgggtgaatgggaaggaggtgcacagtg gggtcagcacggacccgcagccctcaaggagcagccgcctcaatgactccag atactgcctgagcagccgcctgagggctctcgccaccttctggcagaacccccgc aaccacttccgctgtcaagtccagttctacgggctctcggagaatgacgagtga cccaggatagggccaaacctgtcacccagatcgtcagcgccgaggcctggggtag agcagactgtggctttacctcgggtgtctaccagcaaggggtcctgtctgccacc atcctctatgagatcctgctagggaaggccacctgtatgctgtgctggtcagcg cccttgtgttgatggccatggtcaagagaaaggatttcc	SEQ ID NO: 234
	β (with TRBC2)	atgagcatcgggctcctgtgctgtgtggccttttctctcctgtgggcaagtccag tgaatgctggtgtcactcagacccccaaaattccagggtcctgaagacaggacagag catgacactgcagtggtgccaggatatgaaccataactccatgtactggtatcga caagacccaggcatgggactgaggctgatttattactcagcttctgagggtagca ctgacaaaaggagaagtccccaatggctacaatgtctccagattaaacaaacggga gttctcgctcaggctggagtcggctgctccctcccagacatctgtgtacttctgt gccagcaggggggtggcgtgagcagttcttcgggccagggaacaggctcacctgtc tagaggacctgaaaaacgtgttcccaccgaggtcgctgtgtttgagccatcaga agcagagatctcccacacccaaaaggccacactggtgtgcctggccacaggcttc taccgcgaccacgtggagctgagctggtgggtgaatgggaaggaggtgcacagtg gggtcagcacagacccgcagccctcaaggagcagccgcctcaatgactccag atactgcctgagcagccgcctgagggctctcgccaccttctggcagaacccccgc aaccacttccgctgtcaagtccagttctacgggctctcggagaatgacgagtga cccaggatagggccaaacctgtcacccagatcgtcagcgccgaggcctggggtag agcagactgtggctttacctccgagtcctaccagcaaggggtcctgtctgccacc atcctctatgagatcctgctagggaaggccacctgtatgccgtgctggtcagtg ccctcgtgctgatggccatggtcaagagaaaggattccagagggc	SEQ ID NO: 235
HD9	α1 (with TRAC)	atgggtgaagatccggcaatttttgttggctattttgtggcttcagctaagctgtg taagtgccgccaaaaatgaagtggagcagagtcctcagaacctgactgcccagga aggagaatttatcacaatcaactgcagttactcggtaggaataagtgccttacac tggctgcaacagcatccaggaggaggcattgtttccttgtttatgctgagctcag ggaagaagaagcatggaagattaattgccacaataaacatacaggaaaagcacag ctccctgcacatcacagcctccatcccagagactctgccgtctacatctgtgct gcccgatcttataacacccgacaaagctcatcttgggactgggaccagattacaag tctttccaaatatccagaacctgacctgccgtgtaccagctgagagactctaa atccagtgacaagtctgtctgcctattcacccgattttgattctcaaacaaatgtg tcacaaagtaaggattctgatgtgtatatcacagacaaaactgtgctagacatga gggtctatggacttcaagagcaacagtgctgtggcctggagcaacaaatctgactt tgcatgtgcaaacgccttcaacaacagcattattccagaagacaccttcttcccc agcccagaaaagttcctgtgatgtcaagctggtcgagaaaagctttgaaacagata cgaacctaaactttcaaaacctgtcagtgattgggttccgaatcctcctcctgaa agtggccgggtttaatctgctcatgacgctgcggctgtggtccagc	SEQ ID NO: 236
	α2 (with TRAC)	atggccatgctcctgggggcatcagtgctgattctgtggcttcagccagactggg taaacagtcaacagaagaatgatgaccagcaagttaagcaaaattcaccatccct gagcgtccaggaaggaagaatttctattctgaactgtgactatactaacagcatg	SEQ ID NO: 237

		tttgattatttcctatggtacaaaaaataccctgctgaaggctcctacattcctga tatctataagttccattaaggataaaaaatgaagatggaagattcactgtcttctt aaacaaaagtgccaaagcacctctctctgcacattgtgccctcccagcctggagac tctgcagtgtacttctgtgcagcaagttacaacaatgccagactcatgtttggag atggaactcagctggtggtgaagcccaatatccagaaccctgaccctgccgtgta ccagctgagagactctaaatccagtgacaagtctgtctgcctattcaccgatttt gattctcaaacaaatgtgtcacaagtaaggattctgatgtgtatatcacagaca aaactgtgctagacatgaggtctatggacttcaagagcaacagtgctgtggcctg gagcaacaaatctgactttgcatgtgcaaacgccttcaacaacagcattattcca gaagacaccttcttccccagcccagaaagttcctgtgatgtcaagctggtcgaga aaagctttgaaacagatacgaacctaaactttcaaaacctgtcagtgattgggtt ccgaatcctcctcctgaaagtggccgggtttaatctgctcatgacgctgcggtg tgggccagc	
	α3 (with TRAC)	atggccatgctcctgggggcatcagtgtgattctgtggcttcagccagactggg taaacagtcaacagaagaatgatgaccagcaagttaagcaaaattcaccatccct gagcgtccaggaaggaagaatttctattctgaactgtgactatactaacagcatg tttgattatttcctatggtacaaaaaataccctgctgaaggctcctacattcctga tatctataagttccattaaggataaaaaatgaagatggaagattcactgtcttctt aaacaaaagtgccaaagcacctctctctgcacattgtgccctcccagcctggagac tctgcagtgtacttctgtgcagcaagttacaacaatgccagactcatgtttggag atggaactcagctggtggtgaagcccaatatccagaaccctgaccctgccgtgta ccagctgagagactctaaatccagtgacaagtctgtctgcctattcaccgatttt gattctcaaacaaatgtgtcacaagtaaggattctgatgtgtatatcacagaca aaactgtgctagacatgaggtctatggacttcaagagcaacagtgctgtggcctg gagcaacaaatctgactttgcatgtgcaaacgccttcaacaacagcattattcca gaagacaccttcttccccagcccagaaagttcctgtgatgtcaagctggtcgaga aaagctttgaaacagatacgaacctaaactttcaaaacctgtcagtgattgggtt ccgaatcctcctcctgaaagtggccgggtttaatctgctcatgacgctgcggtg tgggccagc	SEQ ID NO: 238
	β1 (with TRBC1)	atgggacccaggctcctcttctgggcaactgctttgtctcctcggaacaggcccag tggaggctggagtcacacaaagtcccacacacctgatcaaaacgagaggacagca agcgactctgagatgctctcctatctctgggcacaccagtgtgtactggtacca caggccctgggtctgggcctccagttcctcctttggtatgacgaggggtgaagaga gaaacagaggaaacttccctcctagattttcagggtcgccagttccctaattatag ctctgagctgaatgtgaacgccttggagctggaggactcgccctgtatctctgt gccagcagctgggggtaccaagagacccagtaacttcgggccaggcacgcggtcc tggtgctcgaggacctgaacaaggtgttcccacccgaggtcgctgtgtttgagcc atcagaagcagagatctcccacacccaaaaggccacactggtgtgcctggccaca ggcttcttccccgaccacgtggagctgagctggtgggtgaatgggaaggaggtgc acagtggggtcagcacggacccgcagcccctcaaggagcagcccgcctcaatga ctccagatactgcctgagcagccgcctgaggggtctcggccaccttctggcagaac ccccgcaaccacttccgctgtcaagtccagttctacgggctctcggagaatgacg agtggacccaggatagggccaaacctgtaccccagatcgtcagcgccgaggcctg gggtagagcagactgtggctttacctcggtgtcctaccagcaaggggtcctgtct gccaccatcctctatgagatcctgctaggggaaggccacctgtatgctgtgctgg tcagcgcccttgtgttgatggccatggtcaagagaaaggatttcc	SEQ ID NO: 239
	β1 (with TRBC2)	atgggacccaggctcctcttctgggcaactgctttgtctcctcggaacaggcccag tggaggctggagtcacacaaagtcccacacacctgatcaaaacgagaggacagca agcgactctgagatgctctcctatctctgggcacaccagtgtgtactggtacca caggccctgggtctgggcctccagttcctcctttggtatgacgaggggtgaagaga gaaacagaggaaacttccctcctagattttcagggtcgccagttccctaattatag ctctgagctgaatgtgaacgccttggagctggaggactcgccctgtatctctgt gccagcagctgggggtaccaagagacccagtaacttcgggccaggcacgcggtcc tggtgctcgaggacctgaaaaacgtgttcccacccgaggtcgctgtgtttgagcc atcagaagcagagatctcccacacccaaaaggccacactggtgtgcctggccaca ggcttctaccccgaccacgtggagctgagctggtgggtgaatgggaaggaggtgc acagtggggtcagcacagacccgcagcccctcaaggagcagcccgcctcaatga ctccagatactgcctgagcagccgcctgaggggtctcggccaccttctggcagaac ccccgcaaccacttccgctgtcaagtccagttctacgggctctcggagaatgacg agtggacccaggatagggccaaacctgtaccccagatcgtcagcgccgaggcctg gggtagagcagactgtggctttacctcggtgtcctaccagcaaggggtcctgtct gccaccatcctctatgagatcctgctaggggaaggccacctgtatgctgtgctgg tcagtgccctcgtgctgatggccatggtcaagagaaaggattccagaggc	SEQ ID NO: 240
	β2 (with TRBC1)	atggacaccagagtactctgctgtgcggtcatctgtcttctgggggcaggctctct caaatgccggcgtcatgcagaacccaagacacctggtcaggaggaggggacagga ggcaagactgagatgcagcccaatgaaaggacacagtcatgtttactggtatcgg cagctcccagagggaaggtctgaaattcatggtttatctccagaaagaaaatatca tagatgagtcaggaatgccaaaggaacgattttctgctgaatttcccaaagaggg ccccagcatcctgaggatccagcaggtagtgcgaggagattcggcagcttatttc tgtgccagctcaccgacaggtggcgagtactatggctacaccttcgggttcgggga ccaggttaaccgttgtagaggacctgaacaaggtgttcccacccgaggtcgctgt gtttgagccatcagaagcagagatctcccacacccaaaaggccacactggtgtgc ctggccacaggcttcttccccgaccacgtggagctgagctggtgggtgaatggga aggaggtgcacagtggggtcagcacggacccgcagcccctcaaggagcagcccgc cctcaatgactccagatactgcctgagcagccgcctgaggggtctcggccaccttc tggcagaacccccgcaaccacttccgctgtcaagtccagttctacgggctctcgg	SEQ ID NO: 241

		agaatgacgagtggaacccaggatagggccaaacccgtcaccagatcgtcagcgc cgaggcctggggtagagcagactgtggctttacctcgggtgtcctaccagcaaggg gtcctgtctgccaccatcctctatgagatcctgctaggggaagggccacctgtatg ctgtgctgggtcagcgcccttgtgttgatggccatgggtcaagagaaaggatttc	
	β2 (with TRBC2)	atggacaccagagtactctgctgtgcggteatctgtcttctgggggcaggctctct caaatgccggcgtcatgcagaacccaagacacctggtcaggaggaggggacagga ggcaagactgagatgcagcccaatgaaaggacacagtcatgtttactggatatcgg cagctcccagaggaaggctctgaaattcatggtttatctccagaaagaaaatatca tagatgagtcaggaatgccaaaggaacgattttctgctgaatttcccaaagaggg ccccagcatcctgaggatccagcaggtagtgcgaggagattcggcagcttatttc tgtgccagctcacgcacaggtggcgagtaactatggctacaccttcgggttcgggga ccaggttaaccgttgtagaggacctgaaaaacgtgttcccaccgcagggtcgctgt gtttgagccatcagaagcagagatctcccacacccaaaaggccacactggtgtgc ctggccacaggcttctaccccgaccacgtggagctgagctggtgggtgaatggga aggaggtgcacagtggggtcagcacagaccgcagccctcaaggagcagccgc cctcaatgactccagatactgcctgagcagccgcctgaggggtctcgccaccttc tggcagaacccccgcaaccacttcgctgtcaagtccagttctacgggctctcgg agaatgacgagtggaacccaggatagggccaaacctgtcaccagatcgtcagcgc cgaggcctggggtagagcagactgtggcttcacctccgagctttaaccagcaaggg gtcctgtctgccaccatcctctatgagatcttgctaggggaagggccaccttgtatg ccgtgctgggtcagtgccctcgtgctgatggccatgggtcaagagaaaggattccag aggc	SEQ ID NO: 242
	β3 (with TRBC1)	atgagcatcggcctcctgtgctgtgcagccttgtctctcctgtgggcagggtccag tgaatgctggtgtcactcagacccccaaaattccagggtcctgaagacaggacagag catgacactgcagtggtgccaggatatgaacctgaatacatgtcctgggtatcga caagaccaggcatggggctgaggctgattcattactcagttgggtgctgggtatca ctgaccaaggagaagtcaccaatggctacaatgtctccagatcaaccacagagga tttcccgtcaggctgctgtcggtgctccctcccagacatctgtgtacttctgt gccagcagttcatacccccttcggacagggcgatacaactcctataattcacccc tccactttgggaacgggaccagggtcactgtgacagaggacctgaacaagggtgtt cccaccgcagggtcgctgtgtttgagccatcagaagcagagatctcccacacccaa aaggccacactggtgtgcctggccacaggcttcttcccgcaccacgtggagctga gctggtgggtgaatgggaaggaggtgcacagtggggtcagcacggacccgcagcc cctcaaggagcagcccgccctcaatgactccagatactgcctgagcagccgcctg aggggtctcgccaccttctggcagaacccccgcaaccacttcgctgtcaagtcc agttctacgggctctcggagaatgacgagtggaaccaggatagggccaaaacccgt caccagatcgtcagcgccgaggcctggggtagagcagactgtggctttacctcg gtgtcctaccagcaaggggtcctgtctgccaccatcctctatgagatcctgctag ggaaggccaccttgtatgctgtgctggtcagcgcccttgtgttgatggccatgggt caagagaaaggatttctga	SEQ ID NO: 243
	β3 (with TRBC2)	atgagcatcggcctcctgtgctgtgcagccttgtctctcctgtgggcagggtccag tgaatgctggtgtcactcagacccccaaaattccagggtcctgaagacaggacagag catgacactgcagtggtgccaggatatgaacctgaatacatgtcctgggtatcga caagaccaggcatggggctgaggctgattcattactcagttgggtgctgggtatca ctgaccaaggagaagtcaccaatggctacaatgtctccagatcaaccacagagga tttcccgtcaggctgctgtcggtgctccctcccagacatctgtgtacttctgt gccagcagttcatacccccttcggacagggcgatacaactcctataattcacccc tccactttgggaacgggaccagggtcactgtgacagaggacctgaaaaacgtgtt cccaccgcagggtcgctgtgtttgagccatcagaagcagagatctcccacacccaa aaggccacactggtgtgcctggccacaggcttctaccccgaccacgtggagctga gctggtgggtgaatgggaaggaggtgcacagtggggtcagcacagacccgcagcc cctcaaggagcagcccgccctcaatgactccagatactgcctgagcagccgcctg aggggtctcgccaccttctggcagaacccccgcaaccacttcgctgtcaagtcc agttctacgggctctcggagaatgacgagtggaaccaggatagggccaaaacctgt caccagatcgtcagcgccgaggcctggggtagagcagactgtggcttcacctcc gagttaccagcaaggggtcctgtctgccaccatcctctatgagatcttgctag ggaaggccaccttgtatgctgtgctggtcagtgccctcgtgctgatggccatgggt caagagaaaggattccagaggc	SEQ ID NO: 244
HD10	α (with TRAC)	atggccatgctcctgggggcatcagtgctgattctgtggcttcagccagactggg taaacagtcaacagaagaatgatgaccagcaagttaagcaaaattcaccatccct gagcgtccaggaaggaagaatttctattctgaactgtgactatactaacagcatg tttgattatttcctatggtacaaaaaataccctgctgaaggctctacattcctga tatctataagttccattaaaggataaaaaatgaagatggaagattcactgtcttctt aaacaaaagtgccaaagcacctctctctgcacattgtgcctcccagcctggagac tctgcagtgtaacttctgtgcagcaagcggaggaagagatgacaagatcatctttg gaaaagggacacgacttcataattctcccgaatatccagaacctgacctgccgt gtaccagctgagagactctaaatccagtgaacagtctgtctgcctattcaccgat tttgattctcaaaacaaatgtgtcacaaagtaaggattctgatgtgtatatcacag acaaaactgtgctagacatgaggtctatggacttcaagagcaacagtgctgtggc ctggagcaacaaatctgactttgcatgtgcaaacgccttcaacaacagcattatt ccagaagacaccttcttccccagcccagaaagttcctgtgatgtcaagctggtcg agaaaagctttgaaacagatacgaacctaaactttcaaaacctgtcagtgattgg gttccgaatcctcctcctgaaagtggccgggtttaatctgctcatgacgctgcgg ctgtggtccagc	SEQ ID NO: 245
	β (with	atgagcatcggcctcctgtgctgtgcagccttgtctctcctgtgggcagggtccag	SEQ ID NO: 246

	TRBC1)	tgaatgctggtgtcactcagacccccaaaattccaggtcctgaagacaggacagag catgacactgcagtggtgccaggatatgaaccatgaatacatgtcctgggtatcga caagacccaggcatggggctgaggctgattcattactcagttggtgctgggtatca ctgaccaaggagaagtccccaatggctacaatgtctccagatcaaccacagagga tttcccgctcaggctgctgtcggtgctccctcccagacatctgtgtacttctgt gccagcagctactcccggacagagagcacagatacgcagtattttggcccaggca cccggctgacagtgtcgcaggacctgaacaagggtgtcccacccgaggtcgctgt gtttgagccatcagaagcagagatctcccacacccaaaaggccacactggtgtgc ctggccacaggcttcttccccgaccacgtggagctgagctggtgggtgaatggga aggaggtgcacagtgggtcagcacggacccgcagccctcaaggagcagccgc cctcaatgactccagatactgctgagcagccgcctgaggggtctcgccaccttc tggcagaacccccgcaaccacttccgctgtcaagtccagttctacgggctctcgg agaatgacgagtggacccaggatagggccaaacccgtcaccagatcgtcagcgc cgaggcctgggttagagcagactgtggctttacctcggtgtcctaccagcaagg gtcctgtctgccaccatcctctatgagatcctgctaggggaaggccaccttgatg ctgtgctggtcagcgcccttgtgttgatggccatggtcaagagaaaggatttc	
	β (with TRBC2)	atgagcatcggcctcctgtgctgtgcagccttgtctctcctgtgggcagggtccag tgaatgctggtgtcactcagacccccaaaattccaggtcctgaagacaggacagag catgacactgcagtggtgccaggatatgaaccatgaatacatgtcctgggtatcga caagacccaggcatggggctgaggctgattcattactcagttggtgctgggtatca ctgaccaaggagaagtccccaatggctacaatgtctccagatcaaccacagagga tttcccgctcaggctgctgtcggtgctccctcccagacatctgtgtacttctgt gccagcagctactcccggacagagagcacagatacgcagtattttggcccaggca cccggctgacagtgtcgcaggacctgaaaaacgtgtcccacccgaggtcgctgt gtttgagccatcagaagcagagatctcccacacccaaaaggccacactggtgtgc ctggccacaggcttctaccccgaccacgtggagctgagctggtgggtgaatggga aggaggtgcacagtgggtcagcacagacccgcagccctcaaggagcagccgc cctcaatgactccagatactgctgagcagccgcctgaggggtctcgccaccttc tggcagaacccccgcaaccacttccgctgtcaagtccagttctacgggctctcgg agaatgacgagtggacccaggatagggccaaacctgtcaccagatcgtcagcgc cgaggcctgggttagagcagactgtggcttcacctccgagtcttaccagcaagg gtcctgtctgccaccatcctctatgagatcctgctaggggaaggccaccttgatg ccgtgctggtcagtgccctcgtgctgatggccatggtcaagagaaaggattccag aggc	SEQ ID NO: 247

Accordingly, the present invention provides an isolated polynucleotide comprising one or more nucleotide sequences selected from the group consisting of SEQ ID NOs: 132-156, 223-247 and 292-299, or variants thereof having at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity thereto.

The present invention also provides a TCR comprising an α chain encoded by a nucleotide sequence selected from the group consisting of SEQ ID NOs: 132, 135, 138, 141, 144, 147, 152, 223, 226, 227, 228, 233, 236, 237, 238, 245, 292, 295, and variants thereof having at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity thereto.

The present invention also provides a TCR comprising a β chain encoded by a nucleotide sequence selected from the group consisting of SEQ ID Nos: 133, 134, 136, 137, 139, 140, 142, 143, 145, 146, 148, 149, 150, 151, 153, 154, 155, 156, 224, 225, 229, 230, 231, 232, 234, 235, 239, 240, 241, 242, 243, 244, 246, 247, 293, 294, 296-299, and variants thereof having at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity thereto.

Further provided by the present invention are isolated polynucleotide sequences derived from the sequences present in Table 2. For example, the present invention provides an isolated polynucleotide encoding a variable region of a TCR according to the present invention, wherein the isolated polynucleotide comprises a stretch of nucleotides of any one
5 of SEQ ID Nos: 132-156, 223-247 and 292-299.

The variant sequences may have additions, deletions or substitutions, of one or more bases. If the variation involves addition(s) or deletion(s) they may either occur in threes or be balanced (i.e. an addition for each deletion) so that the variation does not cause a frame-shift for translation of the remainder of the sequence.

10 Some or all of the variations may be "silent" in the sense that they do not affect the sequence of the encoded protein due to the degeneracy of the genetic code.

Some or all of the variations may produce conservative amino acid substitutions, additions or deletions as explained above. The variation may be concentrated in one or more regions, such as the regions encoding the constant regions, the linker, or the framework regions of
15 the α or β chains, or they may be spread throughout the molecule.

The variant sequence should retain the capacity to encode all or part of a TCR amino acid sequence which binds to a WT1 peptide.

Codon optimisation

The polynucleotides used in the present invention may be codon-optimised. Codon
20 optimisation has previously been described in WO 1999/41397 and WO 2001/79518. Different cells differ in their usage of particular codons. This codon bias corresponds to a bias in the relative abundance of particular tRNAs in the cell type. By altering the codons in the sequence so that they are tailored to match with the relative abundance of corresponding tRNAs, it is possible to increase expression. By the same token, it is possible to decrease
25 expression by deliberately choosing codons for which the corresponding tRNAs are known to be rare in the particular cell type. Thus, an additional degree of translational control is available.

Many viruses, including HIV and other lentiviruses, use a large number of rare codons and by changing these to correspond to commonly used mammalian codons, increased
30 expression of the packaging components in mammalian producer cells can be achieved. Codon usage tables are known in the art for mammalian cells, as well as for a variety of other organisms.

Codon optimisation may also involve the removal of mRNA instability motifs and cryptic splice sites.

Vector

The present invention provides a vector comprising a polynucleotide described herein.

5 A vector is a tool that allows or facilitates the transfer of an entity from one environment to another. In accordance with the present invention, and by way of example, some vectors used in recombinant nucleic acid techniques allow entities, such as a segment of nucleic acid (e.g. a heterologous DNA segment, such as a heterologous Cdna segment), to be transferred into a target cell. The vector may serve the purpose of maintaining the
10 heterologous nucleic acid (DNA or RNA) within the cell, facilitating the replication of the vector comprising a segment of nucleic acid, or facilitating the expression of the protein encoded by a segment of nucleic acid. Vectors may be non-viral or viral. Examples of vectors used in recombinant nucleic acid techniques include, but are not limited to, plasmids, chromosomes, artificial chromosomes and viruses. The vector may be single stranded or
15 double stranded. It may be linear and optionally the vector comprises one or more homology arms. The vector may also be, for example, a naked nucleic acid (e.g. DNA). In its simplest form, the vector may itself be a nucleotide of interest.

The vectors used in the invention may be, for example, plasmid or virus vectors and may include a promoter for the expression of a polynucleotide and optionally a regulator of the
20 promoter.

Vectors comprising polynucleotides used in the invention may be introduced into cells using a variety of techniques known in the art, such as transformation, transfection and transduction. Several techniques are known in the art, for example transduction with recombinant viral vectors, such as retroviral, lentiviral, adenoviral, adeno-associated viral, baculoviral and herpes simplex viral vectors, Sleeping Beauty vectors; direct injection of
25 nucleic acids and biolistic transformation.

Non-viral delivery systems include but are not limited to DNA transfection methods. Here, transfection includes a process using a non-viral vector to deliver a gene to a target cell. Typical transfection methods include electroporation, DNA biolistics, lipid-mediated
30 transfection, compacted DNA-mediated transfection, liposomes, immunoliposomes, lipofectin, cationic agent-mediated transfection, cationic facial amphiphiles (CFAs) (Nature Biotechnology 1996 14; 556) and combinations thereof.

The term “transfection” is to be understood as encompassing the delivery of polynucleotides to cells by both viral and non-viral delivery.

In addition, the invention may employ gene targeting protocols, for example the delivery of DNA-modifying agents.

- 5 The term “vector” includes an expression vector i.e. a construct capable of *in vivo* or *in vitro/ex vivo* expression. Expression may be controlled by a vector sequence, or, for example in the case of insertion at a target site, expression may be controlled by a target sequence. A vector may be integrated or tethered to the cell’s DNA.

10 Viral delivery systems include but are not limited to adenovirus vector, an adeno-associated viral (AAV) vector, a herpes viral vector, a retroviral vector, a lentiviral vector, and a baculoviral vector.

15 Retroviruses are RNA viruses with a life cycle different to that of lytic viruses. In this regard, a retrovirus is an infectious entity that replicates through a DNA intermediate. When a retrovirus infects a cell, its genome is converted to a DNA form by a reverse transcriptase enzyme. The DNA copy serves as a template for the production of new RNA genomes and virally encoded proteins necessary for the assembly of infectious viral particles.

20 There are many retroviruses, for example murine leukemia virus (MLV), human immunodeficiency virus (HIV), equine infectious anaemia virus (EIAV), mouse mammary tumour virus (MMTV), Rous sarcoma virus (RSV), Fujinami sarcoma virus (FuSV), Moloney murine leukemia virus (Mo-MLV), FBR murine osteosarcoma virus (FBR MSV), Moloney murine sarcoma virus (Mo-MSV), Abelson murine leukemia virus (A-MLV), Avian myelocytomatosis virus-29 (MC29), and Avian erythroblastosis virus (AEV) and all other retroviridae including lentiviruses.

25 A detailed list of retroviruses may be found in Coffin *et al* (“Retroviruses” 1997 Cold Spring Harbour Laboratory Press Eds: JM Coffin, SM Hughes, HE Varmus pp 758-763).

Lentiviruses also belong to the retrovirus family, but they can infect both dividing and non-dividing cells (Lewis et al (1992) EMBO J. 3053-3058).

30 The vector may be capable of transferring a nucleotide sequence encoding a WT1-specific TCR described herein to a cell, such as a T-cell, such that the cell expresses the WT1-specific TCR. Preferably the vector will be capable of sustained high-level expression in T-cells, so that the introduced TCR may compete successfully with the endogenous TCR for a limited pool of CD3 molecules.

Increasing the supply of CD3 molecules may increase TCR expression, for example, in a cell that has been modified to express the TCRs of the present invention. Accordingly, the vector of the present invention may further comprise one or more genes encoding CD3-gamma, CD3-delta, CD3-epsilon and/or CD3-zeta. In one embodiment, the vector of the present invention comprises a gene encoding CD3-zeta. The vector may comprise a gene encoding CD8. The vector may encode a selectable marker or a suicide gene, to increase the safety profile of the genetically engineered cell, e.g. a cell of the present invention, or a cell that has been modified to express the TCRs of the present invention (Bonini, Science 1997, Ciceri, Bonini Lancet Oncol. 2009, Oliveira et al., STM 2015). The genes comprised in the vector of the present invention may be linked by self-cleaving sequences, such as the 2A self-cleaving sequence.

Alternatively one or more separate vectors encoding a CD3 gene may be provided for co-transfer to a cell simultaneously, sequentially or separately with one or more vectors of the present invention, e.g. one or more vectors encoding TCRs of the present invention.

Cell

The present invention relates to a cell comprising a polynucleotide or a vector according to the present invention.

The cell may be a T-cell, a lymphocyte, or a stem cell. The T-cell, the lymphocyte, or the stem cell may be selected from the group consisting of CD4 cells, CD8 cells, naive T-cells, memory stem T-cells, central memory T-cells, double negative T-cells, effector memory T-cells, effector T-cells, Th0 cells, Tc0 cells, Th1 cells, Tc1 cells, Th2 cells, Tc2 cells, Th17 cells, Th22 cells, gamma/delta T-cells, natural killer (NK) cells, natural killer T (NKT) cells, hematopoietic stem cells and pluripotent stem cells.

The type of cell may be selected in order to provide desirable and advantageous *in vivo* persistence and to provide desirable and advantageous functions and characteristics to the cells of present invention.

The cell may have been isolated from a subject.

The cell of the present invention may be provided for use in adoptive cell transfer. As used herein the term “adoptive cell transfer” refers to the administration of a cell population to a patient. Typically, the cells are T-cells isolated from a subject and then genetically modified and cultured *in vitro* in order to express a TCR of the present invention before being administered to the patient.

Adoptive cell transfer may be allogenic or autologous.

By “autologous cell transfer” it is to be understood that the starting population of cells (which are then transduced according to a method of the invention, or are transduced with a vector according to the present invention) is obtained from the same subject as that to which the transduced T-cell population is administered. Autologous transfer is advantageous as it avoids problems associated with immunological incompatibility and are available to subjects irrespective of the availability of a genetically matched donor.

By “allogeneic cell transfer” is to be understood that the starting population of cells (which are then transduced according to a method of the invention, or are transduced with a vector according to the present invention) is obtained from a different subject as that to which the transduced cell population is administered. Preferably, the donor will be genetically matched to the subject to which the cells are administered to minimise the risk of immunological incompatibility. Alternatively, the donor may be mismatched and unrelated to the patient.

Suitable doses of transduced cell populations are such as to be therapeutically and/or prophylactically effective. The dose to be administered may depend on the subject and condition to be treated, and may be readily determined by a skilled person.

The cell may be derived from a T-cell isolated from a subject. The T-cell may be part of a mixed cell population isolated from the subject, such as a population of peripheral blood lymphocytes (PBL). T-cells within the PBL population may be activated by methods known in the art, such as using anti-CD3 and/or anti-CD28 antibodies or cell sized beads conjugated with anti-CD3 and/or anti-CD28 antibodies.

The T-cell may be a CD4⁺ helper T cell or a CD8⁺ cytotoxic T cell. The cell may be in a mixed population of CD4⁺ helper T cell/CD8⁺ cytotoxic T-cells. Polyclonal activation, for example using anti-CD3 antibodies optionally in combination with anti-CD28 antibodies will trigger the proliferation of CD4⁺ and CD8⁺ T-cells.

The cell may be isolated from the subject to which the genetically modified cell is to be adoptively transferred. In this respect, the cell may be made by isolating a T-cell from a subject, optionally activating the T-cell, transferring the TCR gene to the cell *ex vivo*. Subsequent immunotherapy of the subject may then be carried out by adoptive transfer of the TCR-transduced cells. As used herein this process refers to autologous T-cell transfer - i.e. the TCR-transduced cells are administered to the same subject from which the T-cells were originally derived.

Alternatively the T-cell may be isolated from a different subject, such that it is allogeneic. The T-cell may be isolated from a donor subject. For example, if the subject is undergoing

allogeneic haematopoietic stem cell transplantation (Allo-HSCT) or solid organ transplantation or cell transplantation or stem cell therapy, the cell may be derived from the donor, from which the organs, tissues or cells are derived. The donor and the subject undergoing treatment may be siblings.

- 5 Alternatively the cell may be, or may be derived from, a stem cell, such as a haemopoietic stem cell (HSC). Gene transfer into HSCs does not lead to TCR expression at the cell surface as stem cells do not express CD3 molecules. However, when stem cells differentiate into lymphoid precursors that migrate to the thymus, the initiation of CD3 expression leads to the surface expression of the introduced TCR in thymocytes.
- 10 An advantage of this approach is that the mature T-cells, once produced, express only the introduced TCR and little or no endogenous TCR chains, because the expression of the introduced TCR chains suppresses rearrangement of endogenous TCR gene segments to form functional TCR alpha and beta genes. A further benefit is that the gene-modified stem cells are a continuous source of mature T-cells with the desired antigen specificity. The cell
- 15 may therefore be a gene-modified stem cell, preferably a gene-modified hematopoietic stem cell, which, upon differentiation, produces a T-cell expressing a TCR of the invention.

Other approaches known in the art may be used to reduce, limit, prevent, silence, or abrogate expression of endogenous genes in the cells of the present invention or cells prepared by the methods of the present invention.

- 20 As used herein the term “disrupting” refers to reducing, limiting, preventing, silencing, or abrogating expression of a gene. The person skilled in the art is able to use any method known in the art to disrupt an endogenous gene, e.g., any suitable method for genome editing, gene silencing, gene knock-down or gene knock-out.

- For example, an endogenous gene may be disrupted with an artificial nuclease. An artificial
- 25 nuclease is, e.g., an artificial restriction enzyme engineered to selectively target a specific polynucleotide sequence (e.g. encoding a gene of interest) and induce a double strand break in said polynucleotide sequence. Typically, the double strand break (DSB) will be repaired by error-prone non-homologous end joining (NHEJ) thereby resulting in the formation of a non-functional polynucleotide sequence, which may be unable to express an
- 30 endogenous gene.

In some embodiments, the artificial nuclease is selected from the group consisting of zinc finger nucleases (ZFN), transcription activator-like effector nucleases (TALEN) and CRISPR/Cas (e.g. CRISPR/Cas9).

The methods of preparing a cell (e.g. a T-cell) of the present invention may comprise the step of targeted integration of a expression cassette into an endogenous gene (e.g. an endogenous TCR α chain gene and/or an endogenous TCR β chain gene). As used herein the term expression cassette refers to a polynucleotide sequence (e.g. a DNA polynucleotide sequence) comprising one or more polynucleotide sequences encoding one or more genes of interest such that said genes of interest are capable of expression. Endogenous sequences may facilitate expression from the expression cassette, and/or transcription control sequences within the expression cassette may facilitate expression. For example, the expression cassette may comprise a polynucleotide sequence of the present invention, or a polynucleotide sequence encoding a TCR of the present invention, operably linked to an expression control sequence, e.g. a promoter or an enhancer sequence. The one or more genes of interest may be located between one or more sets of restriction sites. Suitably, the restriction sites may facilitate the integration of the expression cassette into, e.g., a vector, a plasmid, or genomic DNA (e.g. host cell genomic DNA).

For example, an expression cassette of the present invention may be transferred from a first polynucleotide sequence, e.g. on a vector, to another by 'cutting', e.g. excising, the expression cassette using one or more suitable restriction enzymes and 'pasting', e.g. integrating, the expression cassette into a second polynucleotide sequence.

The expression cassette may comprise a polynucleotide of the present invention. The expression cassette may comprise a polynucleotide encoding one or more TCRs of the present invention. The expression cassette may further comprise an antibiotic resistance gene or other selectable marker gene that allows cells that have successfully integrated the expression cassette into their DNA to be identified. The polynucleotide sequences comprised in the expression cassette may be operably linked to expression control sequences, e.g. a suitable promoter or enhancer sequence. The person skilled in the art will be able to select suitable expression control sequences.

The present invention also contemplates a cell expressing a TCR of the present invention, which has been engineered to disrupt one or more endogenous MHC genes. Disruption of an endogenous MHC gene can reduce or prevent expression of MHC on the engineered cell surface. Accordingly, such an engineered cell with reduced or no MHC expression will have limited or no capacity to present antigens on its cell surface. Such a cell is particularly advantageous for adoptive cell transfer since the cell will be non-alloreactive, e.g., the cell will not present antigens which could be recognized by the immune system of a subject receiving the adoptively transferred cell. As a result, the transferred cell will not be recognized as 'non-self' and an adverse immune reaction to the cell can be avoided. Such a

cell is termed a 'universal cell' since it is suitable for adoptive transfer to a variety of different hosts regardless of HLA type.

Accordingly, the present invention provides a method of preparing a non-alloreactive universal T-cell, which expresses a TCR of the present invention. Further provided by the present invention is a non-alloreactive universal T-cell, which expresses a TCR of the present invention.

The present invention further contemplates cells which have been engineered to disrupt one or more endogenous genes to modify the cell to enhance advantageous properties, characteristics or functions of the cell and/or reduce undesirable properties, characteristics or functions. For example, by disrupting an endogenous cell the persistence, expansion, activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other cell functions may be modified. As used in this context, the term 'modify' refers to a change in one or more characteristics relative to an equivalent unmodified cell, e.g. a cell in which an endogenous gene has not been disrupted. For example, the change may be an increase, an enhancement or an introduction of a characteristic or function of the cell relative to an equivalent unmodified cell. Alternatively, the change may be a decrease, suppression or abrogation of a characteristic or function of the cell relative to an equivalent unmodified cell.

The polynucleotides and vectors of the present invention may be transferred into specific T-cell subsets, including CD4 and or CD8, naive, memory stem T cells, central memory, effector memory or effector cells, or in other cellular subsets such as to promote different *in vivo* length of persistence and function in the cells of the present invention.

The polynucleotides and vectors of the present invention may also be transferred into T-cell subsets such as naïve, memory stem T cells, central memory cells, effector memory cells, effectors.

The polynucleotides and vectors of the present invention may also be transferred into T-cell subsets with different polarizations, such as Th0/Tc0, Th1/Tc1, Th2/Tc2, Th17, Th22 or others, depending on the cytokine background most appropriate to target a particular tumor type.

Furthermore, the polynucleotides and vectors of the present invention encoding the antigen-specific regions of the TCRs of the present invention may be transferred in other cellular subsets, including gamma/delta T-cells, NK cells, NKT cells, hematopoietic stem cells or other cells, in order to obtain the therapeutic effect.

Further provided by the present invention is a method of preparing a cell, which comprises the step of transducing a cell *in vitro* or *ex vivo* with a vector of the present invention. Various methods for transduction of a cell with a vector are known in the art (see e.g. Sambrook et al).

- 5 The present invention also provides a method of producing a T-cell expressing a TCR of the invention by inducing the differentiation of a stem cell which comprises a polynucleotide or a vector of the present invention.

A population of cells may be purified selectively for cells that exhibit a specific phenotype or characteristic, and from other cells which do not exhibit that phenotype or characteristic, or
 10 exhibit it to a lesser degree. For example, a population of cells that expresses a specific marker (e.g. CD3, CD4, CD8, CD25, CD127, CD152, CXCR3, or CCR4) may be purified from a starting population of cells. Alternatively, or in addition, a population of cells that does not express another marker may be purified.

By “enriching” a population of cells for a certain type of cells it is to be understood that the
 15 concentration of that type of cells is increased within the population. The concentration of other types of cells may be concomitantly reduced.

Purification or enrichment may result in the population of cells being substantially pure of other types of cell.

Purifying or enriching for a population of cells expressing a specific marker (e.g. CD3, CD4,
 20 CD8, CD25, CD127, CD152, CXCR3, or CCR4) may be achieved by using an agent that binds to that marker, preferably substantially specifically to that marker. An agent that binds to a cellular marker may be an antibody, for example antibody which binds to CD3, CD4, CD8, CD25, CD127, CD152, CXCR3, or CCR4.

The term “antibody” refers to complete antibodies or antibody fragments capable of binding
 25 to a selected target, and including Fv, ScFv, F(ab') and F(ab')₂, monoclonal and polyclonal antibodies, engineered antibodies including chimeric, CDR-grafted and humanised antibodies, and artificially selected antibodies produced using phage display or alternative techniques.

In addition, alternatives to classical antibodies may also be used in the invention, for
 30 example “avibodies”, “avimers”, “anticalins”, “nanobodies” and “DARPs”.

The agents that bind to specific markers may be labelled so as to be identifiable using any of a number of techniques known in the art. The agent may be inherently labelled, or may be

modified by conjugating a label thereto. By “conjugating” it is to be understood that the agent and label are operably linked. This means that the agent and label are linked together in a manner which enables both to carry out their function (e.g. binding to a marker, allowing fluorescent identification, or allowing separation when placed in a magnetic field) substantially unhindered. Suitable methods of conjugation are well known in the art and would be readily identifiable by the skilled person.

A label may allow, for example, the labelled agent and any cell to which it is bound to be purified from its environment (e.g. the agent may be labelled with a magnetic bead or an affinity tag, such as avidin), detected or both. Detectable markers suitable for use as a label include fluorophores (e.g. green, cherry, cyan and orange fluorescent proteins) and peptide tags (e.g. His tags, Myc tags, FLAG tags and HA tags).

A number of techniques for separating a population of cells expressing a specific marker are known in the art. These include magnetic bead-based separation technologies (e.g. closed-circuit magnetic bead-based separation), flow cytometry, fluorescence-activated cell sorting (FACS), affinity tag purification (e.g. using affinity columns or beads, such as biotin columns to separate avidin-labelled agents) and microscopy-based techniques.

It may also be possible to perform the separation using a combination of different techniques, such as a magnetic bead-based separation step followed by sorting of the resulting population of cells for one or more additional (positive or negative) markers by flow cytometry.

Clinical grade separation may be performed, for example, using the CliniMACS[®] system (Miltenyi). This is an example of a closed-circuit magnetic bead-based separation technology.

It is also envisaged that dye exclusion properties (e.g. side population or rhodamine labelling) or enzymatic activity (e.g. ALDH activity) may be used to enrich for HSCs.

Chimeric molecules

In another aspect, the present invention provides a chimeric molecule comprising a TCR of the present invention, a TCR encoded by a polynucleotide of the present invention, or a portion thereof, conjugated to a non-cellular substrate. The conjugation may be covalent or non-covalent.

The non-cellular substrate may be a nanoparticle, an exosome, or any non-cellular substrate known in the art.

The chimeric molecule of the present invention may be soluble.

In another aspect the present invention provides a chimeric molecule comprising a TCR of the present invention, a TCR encoded by a polynucleotide of the present invention, or a portion thereof, conjugated to a toxin or an antibody.

- 5 The toxin or antibody may be cytotoxic. The toxin may be a cytotoxic molecule or compound, e.g. a radioactive molecule or compound. The TCR portion of the chimeric molecule may confer the ability to recognize cells expressing WT1 protein or peptides. Thus, the chimeric molecule may specifically recognize and/or bind to WT1-expressing tumor cells. Accordingly, the chimeric molecules of the present invention may provide WT1-
10 targeted delivery of cytotoxic toxins, antibodies and/or compounds.

WT1-related diseases

- WT1 is widely expressed on a variety of hematological and solid tumors, while showing limited expression on various healthy tissues (e.g. gonads, uterus, kidney, mesothelium, progenitor cells in different tissues). The present inventors have identified and determined
15 the amino acid sequences of TCRs that recognise WT1 peptides. Furthermore, they have demonstrated that T-cells expressing TCRs according to the present invention target and kill cells which present WT1 peptide or overexpress WT1 protein.

- Accordingly, the present invention provides a method for treating and/or preventing a disease associated with expression of WT1, which comprises the step of administering a
20 TCR, an isolated polynucleotide, a vector, or a cell of the present invention to a subject in need thereof. The present invention also provides a method for treating and/or preventing a disease associated with expression of WT1, comprises the step of administering a cell prepared by the method of the present invention to a subject in need thereof.

- Further provided by the present invention is a TCR of the present invention, an isolated
25 polynucleotide of the present invention, a vector of the present invention, a cell according of the present invention, or a cell prepared by the method of the present invention for use in treating and/or preventing a disease associated with expression of WT1.

- The term 'preventing' is intended to refer to averting, delaying, impeding or hindering the contraction of the disease. The treatment may, for example, prevent or reduce the likelihood
30 of developing or contracting a disease associated with expression of WT1.

'Treating' as used herein refers to caring for a diseased subject, in order to ameliorate, cure or reduce the symptoms of the disease, or in order to reduce, halt or delay the progression of the disease.

5 The subject may be a human subject. The human subject may be a child. For example, the child may be less than 10 years in age, less than 9 years in age, less than 8 years in age, less than 7 years in age, less than 6 years in age, less than 5 years in age, less than 4 years in age, less than 3 years in age, or less than 2 years in age. The human subject may be an infant.

10 The subject may have been previously determined to be in need of a TCR, an isolated polynucleotide, a vector, or a cell of the present invention, or a cell prepared by the method of the present invention on the basis of expression of WT1. For example, the subject may have a cell population that exhibits increased expression of WT1 relative to a healthy control cell population. A variety of techniques known in the art may be used to determine WT1 expression - e.g. quantitative RT-PCR can be used to determine the amount of WT1 RNA
15 transcript, which is indicative of WT1 protein expression. The person skilled in the art will also appreciate that WT1 protein expression may be determined by performing western blots using commercially available antibodies specific for WT1.

The subject may also have been previously identified as having an alteration (e.g. mutation or deletion) in a WT1 gene. Such an alteration may be hereditary. Thus, the disease
20 associated with expression of WT1 may be a hereditary disease. Examples of hereditary diseases associated with expression of WT1 include but are not limited to WAGR (Wilms tumor-Aniridia-Genitourinary malformation-Retardation) syndrome, Denys-Drash syndrome (DDS), Frasier syndrome (FS), genitourinary anomalies (abnormalities of the reproductive and urinary systems) syndrome.

25 Subjects with hereditary diseases associated with expression of WT1 may be at higher risk of developing a proliferative disorder (e.g. a cancer).

The disease associated with expression of WT1 may be a proliferative disorder.

The proliferative disorder may be a hematological malignancy or a solid tumor. The hematological malignancy may be selected from the group consisting of acute myeloid
30 leukemia (AML), chronic myeloid leukemia (CML), lymphoblastic leukemia, myelodysplastic syndromes, lymphoma, multiple myeloma, non Hodgkin lymphoma, and Hodgkin lymphoma.

The solid tumor may be selected from the group consisting of lung cancer, breast cancer, oesophageal cancer, gastric cancer, colon cancer, cholangiocarcinoma, pancreatic cancer,

ovarian cancer, head and neck cancers, synovial sarcoma, angiosarcoma, osteosarcoma, thyroid cancer, endometrial cancer, neuroblastoma, rhabdomyosarcoma, liver cancer, melanoma, prostate cancer, renal cancer, soft tissue sarcoma, urothelial cancer, biliary cancer, glioblastoma, mesothelioma, cervical cancer, and colorectal cancer.

- 5 The disease associated with expression of WT1 may be selected from a group consisting of acute myeloid leukemia (AML), chronic myeloid leukemia (CML), lymphoblastic leukemia, myelodysplastic syndromes, lymphoma, multiple myeloma, non Hodgkin lymphoma, and Hodgkin lymphoma, lung cancer, breast cancer, oesophageal cancer, gastric cancer, colon cancer, cholangiocarcinoma, pancreatic cancer, ovarian cancer, head and neck cancers, synovial sarcoma, angiosarcoma, osteosarcoma, thyroid cancer, endometrial cancer, neuroblastoma, rhabdomyosarcoma, liver cancer, melanoma, prostate cancer, renal cancer, soft tissue sarcoma, urothelial cancer, biliary cancer, glioblastoma, mesothelioma, cervical cancer, and colorectal cancer.

Pharmaceutical composition

- 15 The TCRs of the present invention, the polynucleotides of the present invention, the vectors of the present invention, the cells of the present invention, the cells prepared by the methods of the present invention, the chimeric molecules of the present invention, and the mixed cell population of the present invention may be formulated for administration to subjects with a pharmaceutically acceptable carrier, diluent or excipient. Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline, and potentially contain human serum albumin.

Handling of the cell therapy products is preferably performed in compliance with FACT-JACIE International Standards for cellular therapy.

Method of treatment

- 25 In another aspect, the present invention provides a method for treating and/or preventing a disease associated with expression of WT1, which comprises the step of administering a TCR of the present invention, an isolated polynucleotide of the present invention, a vector of the present invention, a cell of the present invention, a cell prepared by a method of the present invention, a chimeric molecule of the present invention, or a mixed cell population of the present invention to a subject in need thereof.

The subject may be a human subject. The subject may be a non-human animal subject.

The subject may have a disease associated with expression of WT1. The subject may be at risk of developing a diseases associated with expression of WT1. The subject may have been

previously determined to be at risk of developing a disease associated with expression of WT1. The subject may have an increased risk of developing a disease associated with WT1.

5 The increased risk may have been determined by genetic screening and/or by reviewing the subject's family history. The subject may express genetic markers indicative of increased risk of developing a disease associated with expression of WT1.

Suitably, a person skilled in the art will be aware of genetic risk factors (e.g. genetic markers) associated with increased risk of developing a disease associated with WT1. The skilled person may be able to use any suitable method or technique known in the art to determine
10 whether the subject has an increased risk of developing a disease associated with expression of WT1.

The subject may have previously received treatment for a disease associated with expression of WT1. The subject may be in remission. The subject may be resistant to chemotherapy. The subject may be resistant to an anti-WT1 therapy.

15 In one embodiment, the method for treating and/or preventing a disease associated with expression of WT1 comprises the step of administering a chemotherapy to the subject. The chemotherapy may be administered to the subject simultaneously, sequentially or separately with the TCR of the present invention, the isolated polynucleotide of the present invention, the vector of the present invention, the cell according of the present invention, the cell
20 prepared by the method of the present invention, or the chimeric molecule of the present invention.

In another aspect, the present invention provides a method of treating and/or preventing a disease associated with expression of WT1, which comprises the step of administering a mixed cell population, wherein the mixed cell population comprises a plurality of cell
25 populations each expressing a different TCR of the present invention.

In another aspect, the present invention provides a mixed cell population comprising a plurality of cell populations each expressing a different TCR of the present invention.

In another aspect, the present invention provides a method for preparing a mixed cell population comprising a plurality of cell populations each expressing a different TCR of the
30 present invention, wherein the method comprises the step of transducing a cell *in vitro* or *ex vivo* with a vector of the present invention.

In another aspect, the present invention provides a mixed cell population for use in treating and/or preventing a disease associated with expression of WT1, wherein the mixed cell population comprises a plurality of cell populations each expressing a different TCR of the present invention.

- 5 For example, the mixed cell population may comprise a first cell population expressing a first TCR of the present invention and a second cell population expressing a second TCR of the present invention. For example, the mixed cell population may comprise a first cell population expressing a first TCR of the present invention, a second cell population expressing a second TCR of the present invention, and a third cell population expressing a
10 third TCR of the present invention, and so on.

Each cell population of the mixed cell population may, for example, express a single TCR of the present invention only. The endogenous TCR genes of the cell populations in the mixed cell population may be disrupted or deleted. Expression of endogenous TCR genes of the cells in the mixed cell population may be disrupted, e.g. by gene editing with an artificial
15 nuclease.

In another aspect, the present invention provides use of TCR of the present invention, an isolated polynucleotide of the present invention, a vector of the present invention, a cell of the present invention, a cell prepared by a method of the present invention, a chimeric molecule of the present invention, or a mixed cell population of the present invention, for the
20 manufacture of a medicament for the treatment of a disease associated with expression of WT1.

Both human and veterinary treatments are within the scope of the present invention.

The practice of the present invention will employ, unless otherwise indicated, conventional
25 techniques of cell biology, molecular biology, histology, immunology, oncology, which are within the capabilities of a person of ordinary skill in the art. Such techniques are explained in the literature.

See, for example, Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) Molecular Cloning: A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory Press; Ausubel, F.M. et al.
30 (1995 and periodic supplements) Current Protocols in Molecular Biology, Ch. 9, 13 and 16, John Wiley & Sons; Roe, B., Crabtree, J. and Kahn, A. (1996) DNA Isolation and Sequencing: Essential Techniques, John Wiley & Sons; Polak, J.M. and McGee, J.O'D. (1990) In Situ Hybridization: Principles and Practice, Oxford University Press; Gait, M.J. (1984) Oligonucleotide Synthesis: A Practical Approach, IRL Press; and Lilley, D.M. and

Dahlberg, J.E. (1992) Methods in Enzymology: DNA Structures Part A: Synthesis and Physical Analysis of DNA, Academic Press. Each of these general texts is herein incorporated by reference.

5 Various preferred features and embodiments of the present invention will now be described by way of non-limiting examples.

EXAMPLES

Example 1 - Generation of functional WT1-specific cytotoxic T-lymphocytes (CTLs) from healthy donors (HDs)

10 In order to identify novel TCRs specific for WT1 epitopes restricted to different HLA alleles, we stimulated peripheral blood mononuclear cells (PBMCs) from ten different HDs with a pool of pentadecapeptides (15mer) with an 11 amino acid overlap spanning the complete sequence of the WT1 protein (see materials and methods). This peptide design ensures the optimal stimulation of both CD4⁺ and CD8⁺ T-cells.

15 After 26-30 hours of stimulation, we enriched T-cells expressing CD137. CD137 is molecule upregulated upon T cell receptor engagement and has been previously shown to be a reliable marker for the rapid identification, isolation and expansion *in vitro* of antigen-specific memory and naive CD4⁺ and CD8⁺ T-cells. The CD137-negative fraction was further depleted of the CD3 fraction, then irradiated at 30 Gy and used as antigen presenting cells (APCs) for the CD137⁺ fraction. Sorted CD137⁺ cells were expanded *in vitro* for ~9 days and
20 restimulated with autologous APCs represented by CD3-depleted cells or by immortalized autologous B cells loaded with the peptide pool every 7-14 days. This procedure led to the enrichment of the WT1-specific T lymphocytes as shown by the cytofluorimetric results presented in Figure 1 (a-j). Functional characterization of T cells was performed at different time points. More in detail, T cells were co-cultured with autologous APCs loaded with the
25 peptide pool and, after 6 hours of co-culture, expression of CD107a and IFN γ in the T-cell population was identified by intracellular staining. The expression of CD107a after antigen encounter indicates antigen-induced degranulation and lytic potential.

30 As a negative control, cells were stimulated with an unrelated peptide pool. Control stimulation resulted in minimal secretion of IFN γ and CD107a by T-cells derived from each healthy donor.

Example 2 - Mapping of WT1 epitopes eliciting a T cell response

To identify the WT1 epitope recognized by T-cells, IFN γ secretion was quantified after 6 hours of *in vitro* co-culture of the WT1-stimulated/enriched T-cells with autologous APCs loaded with peptide subpools each containing up to 12 peptides according to a mapping grid. The mapping grid consists of 24 subpools with each peptide being uniquely contained within two intersecting subpools (Doubrovina, E. et al. Blood 120, 1633–1646 (2012)). Results are summarized in Figure 2a.

FACS analysis showed substantial expression of IFN γ and CD107a by the HD1-, HD3-, HD6-, HD7-, HD10-derived T-cells after stimulation with subpools 4, 5 and 16 (Figure 2b, d, g, h, k). Substantial expression of IFN γ was observed for HD2-derived T-cells stimulated by subpools 6, 16, 17, and 20 (Figure 2c), whereas subpools 4, 5, 6 14, 18, 21 stimulated expression of IFN γ and CD107a by HD4-derived T-cells (Figure 2e) and subpools 5, 11, 12, 21, 22 stimulated expression of IFN γ in HD5 (Figure 2f). For HD7, we additionally observed an increased expression of IFN γ and CD107a, even though at a lower level compared to the one observed with subpools 4, 5 and 16, after stimulation with subpools 7, 8, 20 (Figure 2h). Furthermore, we observed an increased expression of IFN γ and CD107a after stimulation of HD8-derived T cells cells with subpools 12 and 14 (Figure 2i) and of HD9-derived T cells with subpools 5,13, 21 (Figure 2j).

Afterwards, the HD-derived T-cells were stimulated for 6 hours with APCs pulsed with the single pentadecapeptides shared by the subpools eliciting the highest immune response and with at least one unrelated 15mer. FACS analysis indicated an increased expression of CD107a and/or IFN γ for peptides 40 and 41 in HD1 T-cells (Figure 3a), peptides 54, 77, 90 for HD2 T-cells (Figure 3b), peptide VLDFAPPGA (SEQ ID NO: 157; which is a nonamer of the peptide represented by SEQ ID NO: 117) for HD3 T-cells (Figure 3c), peptides 17, 18, 99, 100 for HD4 (Figure 3d, e), peptide 101 for HD5 (Figure 3f), peptide VLDFAPPGA (SEQ ID NO: 157; which is a nonamer of the peptide represented by SEQ ID NO: 117) for HD6 T-cells (Figure 3g), peptides 101, 125 and 137 for HD9 T-cells (Figure 3h), peptide VLDFAPPGA (SEQ ID NO: 157; which is a nonamer of the peptide represented by SEQ ID NO: 117) for HD10 T-cells (Figure 3i). In this way, the dominant immunogenic sequences were identified. No relevant immune responses (i.e. increased expression of CD107a and/or IFN γ) were observed after co-culture with unrelated control peptides. For HD7 and HD8, due to the reduced cell fitness, it was not possible to perform culture experiments to identify the immunogenic peptides. Still, we could predict the recognized peptide by the deconvolution of the mapping grid. We identified the overlapping sequence of peptides 41 and 42 (originated

from SP4, 5, 16) and the overlapping sequence of peptides 91 and 92 (originated from SP7, 8, 20) for HD7 and peptide 24 for HD8.

To determine the HLA-restriction of the WT1 immunogenic peptide recognized by T-cells expanded from HD4, HD5 and HD10, the T-cells from these were co-cultured for 6 hours with a panel of different target EBV-BLCL cells each expressing a different HLA-A or HLA-B allele that had been pulsed with a relevant peptide (peptide 17 for HD4; peptide 101 for HD5; peptide VLDFAPPGA (SEQ ID NO: 157) for HD10) or an unrelated control peptide. Results of this experiment showed that peptide 17 is presented by the HLA-B*3502 allele and is recognized by T-cells derived from HD4 (Figure 3j), peptide 101 is presented by the HLA-B*3501 and is recognized by T-cells derived from HD5 (Figure 3k); peptide VLDFAPPGA (SEQ ID NO: 157) is presented by the HLA-A*0201 and is recognized by T-cells derived from HD10 (Figure 3l).

Sequences of the WT1 peptides recognized by WT1-specific T-cells expanded from HD1-HD10 are shown in Table 3 below.

Table 3

Binding TCR donor	Peptide	Sequence	SEQ ID NO
HD1/HD3/HD6/HD7/HD10	40	AAQWAPVLDFAPPGA	SEQ ID NO: 115
	41	APVLDFAPPGASAYG	SEQ ID NO: 116
	Overlapping sequence ("11 mer" in Figure 3c)	APVLDFAPPGA	SEQ ID NO: 117
HD2	54	QCLSAFTVHFSGQFT	SEQ ID NO: 118
	77	EDPMGQQGSLGEQQY	SEQ ID NO: 119
	90	SQLECMTWNQMNLGA	SEQ ID NO: 120
HD4	17	TCVPEPASQHTLRSG	SEQ ID NO: 121
	18	EPASQHTLRSGPGCL	SEQ ID NO: 122
	Overlapping sequence	EPASQHTLRSG	SEQ ID NO: 123
	99	HSTGYESDNHTTPIL	SEQ ID NO: 124
	100	YESDNHTTPILCGAQ	SEQ ID NO: 125

	Overlapping sequence	YESDNHHTTPIL	SEQ ID NO: 126
HD5	101	NHTTPILCGAQYRIH	SEQ ID NO: 127
HD7	91	CMTWNQMNLGATLKG	SEQ ID NO: 248
	92	NQMNLGATLKGVAAG	SEQ ID NO: 249
	Overlapping sequence	NQMNLGATLKG	SEQ ID NO: 250
HD8	24	DPGGIWAKLGAAEAS	SEQ ID NO: 251
HD9	101	NHTTPILCGAQYRIH	SEQ ID NO: 252
	125	KRHQRRHTGVKPFQC	SEQ ID NO: 253
	137	PSCQKKFARSDDELVR	SEQ ID NO: 254

Example 3 - WT1-specific T-cells selectively eliminate WT1 expressing cells

To determine the HLA restriction of the WT1 specific T-cells and their ability to eliminate WT1-expressing cells, T cells were co-cultured with different target cells.

HD1-derived T-cells

- 5 Being aware that HD1 harbors the HLA-A*0201 allele, we co-cultured enriched WT1-specific T-cells with different target cells: T2 cells pulsed with the overlapping peptide pool comprising peptides 40 and 41 (see Table 3); T2 cells pulsed with the MelanA/MART1 pool as a negative control (T2 MelanA/MART1 pool); or K562 cells genetically modified in order to express both the HLA-A*0201 allele and to overexpress the WT1 protein (K562 HLA-A*0201 WT1).
- 10

After 6 hours of co-culture, expression of CD107a was established by FACS. Results for HD1 indicate the expression of CD107a in >60% of CD8⁺ T-cells following co-culture with T2 cells pulsed with the WT1 pool (Figure 4a). Similarly, co-culture of WT1 specific T-cells with genetically modified K562 cells (expressing HLA-A*0201 allele and the WT1 protein) resulted in CD107a expression by >60% of CD8⁺ T-cells (Figure 4a).

15

In contrast, CD107a expression by CD8⁺ T-cells co-cultured with T2 cells pulsed with the negative control MelanA/MART1 pool was minimal (Figure 4a).

These results demonstrate that isolated HD1-derived T-cells specifically recognize WT1 peptide comprising the sequence APVLDFAPPGA (SEQ ID NO: 117) when presented by MHC molecules encoded by the HLA-A*0201 allele. Moreover, the results show that HD1-derived T-cells are able to specifically target cells overexpressing the WT1 protein.

- 5 Accordingly, these experimental data demonstrate that TCRs expressed by HD1-derived T-cells specifically bind to the peptides comprising the APVLDFAPPGA (SEQ ID NO: 117) amino acid sequence and that such TCRs are HLA-A*0201 restricted.

HD3-derived T-cells

- 10 Being aware that HD3 harbors the HLA-A*0201 allele, we co-cultured enriched WT1-specific T-cells with different target cells: T2 cells pulsed with subpool 16 which was previously determined to contain the immunogenic peptide eliciting immune response (T2-SP16); T2 cells pulsed with the MelanA/MART1 pool as a negative control (T2-Melan A); wild type K562 cells (K562) as negative control; or K562 cells genetically modified to express both the HLA-A*0201 allele and to overexpress the WT1 protein (K562 A2+WT1+).

- 15 After 4 days of co-culture, the ability of the HD3 derived T-cells to kill target cells was expressed as elimination index - calculated as the total number of target cells still present after co-culture with the WT1-specific T-cells divided by the total number of target cells alone.

- 20 The results demonstrate the ability of WT1-specific T-cells to eliminate target cells expressing the identified specific WT1 epitope (Figure 4b). In particular, the HD3-derived T-cells eliminated about 95% of the T2 cells pulsed with WT1 peptides comprising APVLDFAPPGA (SEQ ID NO: 117) amino acid sequence (subpool 16; SP16). Furthermore, the HD3-derived T-cells eliminated about 78% of the K562 cells expressing MHC molecules encoded by the HLA-A*0201 allele and overexpressing WT1 protein. In contrast, none of
25 the negative control MelanA/MART1 pool-pulsed T2 cells were eliminated by the HD3-derived T-cells. Similarly, there was minimal elimination of control wild-type K562 cells (Figure 4b).

- 30 These results demonstrate that isolated HD3-derived T-cells specifically recognize WT1 peptide comprising the amino acid sequence APVLDFAPPGA (SEQ ID NO: 117). Moreover, the results show that HD3-derived T-cells are able to specifically target and kill cells overexpressing the WT1 protein *via* peptide presentation by HLA-A*0201 encoded MHC. Accordingly, these experimental data demonstrate WT1 peptide specificity for TCRs expressed by HD3-derived T-cells and that HD3-derived TCRs are HLA-A*0201 restricted.

HD4-derived T-cells

The ability of HD4-derived T-cells to eliminate target cells was assessed by co-culturing the T-cells with primary leukemic blasts (CD33⁺ cells) isolated from an acute myeloid leukemia (AML) patient who was selected on the basis of high expression of the WT1 antigen and
 5 HLA typing (HLA-B*3502). As a negative control, HD4-derived T-cells were co-cultured with leukemic blasts from an AML patient who did not express the HLA-B*3502 allele.

After three days of co-culture at an effector to target ratio of 10 to 1, FACS analysis showed the nearly complete clearance of the leukemic blasts (CD33⁺) harvested from the AML patient expressing the HLA-B*3502 allele following the co-culture with HD4 WT1-specific T-
 10 cells (CD3⁺ cells) (Figure 4c, upper panel). Indeed, only 0.54% of the remaining total cell population was positive for CD33 expression.

In contrast, no clearance of CD33⁺ cells was seen following co-culture of WT1-specific T-cells with the unrelated control AML blasts (Figure 4c, lower panel). Indeed, in the control sample 7.9% of the total cell population was positive for CD33 expression following co-
 15 culture with the HD4-derived WT1-specific T-cells.

Importantly, these results demonstrate the ability of the WT1-specific T-cells derived from HD4 to specifically target and kill leukemic blasts (AML cancer cells) overexpressing WT1 and MHC encoded by the HLA-B*3502. Thus, TCRs derived from HD4 are able to specifically target and kill cancer cells in HLA-B*3502 restricted manner.

20 **Example 4 - Immunoprofiling of V β sequences for WT1 specific T-cells**

To better identify the TCRs involved in antigenic recognition by the WT1 specific T-cells from HD1-6 and 10 (for HD7, HD8 and HD9, it was not possible to perform the V β Immunoprofiling analysis due to a reduced cell fitness), we first performed a multi-parametric FACS analysis to quantitatively determine the TCR V β repertoire. Thus, in order to
 25 determine the clonality of the expanded WT1-specific T-cells, we used the IO Test Beta Mark TCR V beta repertoire kit according to manufacturer's recommendations. This kit allows the detection of the expression of 24 different V beta genes in eight individual tubes. In particular, coverage of 75% of the complete repertoire of V beta is guaranteed by using this approach. Results of FACS staining indicated the great prevalence of a specific V β for
 30 HD1, HD2, HD3, HD5 - see Figure 5. For HD4, HD6 and HD10, an exhaustive determination of the predominant V β was not possible, likely due to the intrinsic limitation of the kit which includes antibodies covering 75% of the existing V β proteins.

Example 5 – High throughput sequencing of TCR α and β chains isolated from WT1-specific T cells

WT1-specific T-cells were collected at different time points over the co-culture time frame and their RNA was extracted by using the Arcturus Pico Pure RNA extraction kit. CDR3
 5 sequences of the WT1-specific T-cells were amplified by using a modified RACE approach in which a magnetic capture was included after the cDNA synthesis in order to increase the specificity of the reaction and eliminate unwanted templates (Ruggiero et al. Nat. Commun. 6, 8081 (2015)). Samples were sequenced using an Illumina MiSeq sequencer and the CDR3 clonotypes were identified using the MiXTCR software (Bolotin, DA et al. Nature
 10 Methods 12, 380–381 (2015)) Additionally, CDR1, CDR2 and CDR3 were further determined using the IMGT V-quest tool (Brochet, X. et al., Nucl. Acids Res. 36, W503-508 (2008). PMID: 18503082; Giudicelli, V., Brochet, X., Lefranc, M.-P., Cold Spring Harb Protoc. 2011 Jun 1;2011(6). pii: pdb.prot5633. doi: 10.1101/pdb.prot5633. PMID: 21632778 Abstract also in IMGT booklet with generous provision from Cold Spring Harbor (CSH)
 15 Protocol).

Sequencing results demonstrated the increasing predominance of a specific CDR3 clonotype in the WT1-specific T cell population over time for both TCR chains in HD1-10. Predominant α and β chain genotypes, and CDR3 sequences are provided in Figure 6.

In addition, the full length α and β chain amino acid sequences for TCRs derived from HD1-
 20 10 were determined - see Table 1. The corresponding nucleotide sequences were also determined - see Table 2.

Example 6 – Functional validation of the newly identified TCRs

TCRs α and β sequences isolated from HD1 and HD3 and recognizing the WT1 VLDFAPPGA (SEQ ID NO: 157) peptide when presented by the HLA-A*0201 allele were
 25 cloned into a lentiviral vector under the control of a bidirectional promoter to promote robust and coordinate expression of both TCR chains in transduced lymphocytes. T cells from a healthy individual were transduced with the viral vector encoding either the HD1 TCR or the HD3 TCR. As control, we also transduced cells with the WT1 126-134 TCR. Transduced T cells were functionally validated by co-culture with different target cells represented by the
 30 T2 cells pulsed with one of the 2 recognized peptides (VLDFAPPGA (SEQ ID NO: 157) for HD1 and HD3 TCR; RMFPNAPYL (SEQ ID NO: 255) for WT1 126-134 TCR) (Figure 7a), K562 cells either wild type or engineered in order to express the HLA-A*0201 allele (Figure 7b), primary AML blasts derived from 3 AML patients and selected according to the expression of the HLA-A*0201 allele and the WT1 expression (Figure 7c). Upon 3 days of

co-culture we observed the ability of each transduced T cell population in specifically recognizing the target peptide when presented by the HLA-A*0201 allele (Figure 7a) and the greater potential of HD1 T cells in mediating a near complete elimination of the engineered K562 cells. The higher potency of HD1 TCR in recognizing the target antigen was further confirmed by the results of the co-culture with pAML blasts. Here, we observed a greater elimination of both pAML blasts harbouring the HLA-A*0201 allele upon co-culture with HD1 TR T cells compared to the conditions in which HD3 TR and WT1 126-134 TR T lymphocytes were used as effector cells.

MATERIALS AND METHODS

The WT1 protein sequence previously published by Gessler et al. (Gessler, M. et al. (1990) Nature 343: 774-778) was used to design the peptides used for the stimulation and isolation of WT1-specific T cells. This sequence contains 575 amino acids and includes the first 126 amino acids in the N-terminus missing in the (exon 5+, KTS+) isoform of WT1. We designed 141 pentadecapeptides spanning the whole sequence of the WT1 protein, each overlapping the next one by 11 amino acids.

Peptides were synthesized by PRIMM to specifications of validated sequence, 70% purity, sterility and absence of endotoxin. These peptides were mixed in equal amounts in the WT1 pool at a concentration of 1 µg/ml per peptide. Additionally, 24 subpools were generated, each containing up to 12 peptides (4.17 µg/ml/ per peptide) according to a specific mapping matrix in order to have each peptide included in only two overlapping subpools as shown in Table 4 (see mapping grid strategy in Doubrovina, E. et al. Blood 120, 1633–1646 (2012)).

Table 4.

	SP1	SP2	SP3	SP4	SP5	SP6	SP7	SP8	SP9	SP10	SP11	SP12
SP13	1	2	3	4	5	6	7	8	9	10	11	12
SP14	13	14	15	16	17	18	19	20	21	22	23	24
SP15	25	26	27	28	29	30	31	32	33	34	35	36
SP16	37	38	39	40	41	42	43	44	45	46	47	48
SP17	49	50	51	52	53	54	55	56	57	58	59	60
SP18	61	62	63	64	65	66	67	68	69	70	71	72
SP19	73	74	75	76	77	78	79	80	81	82	83	84
SP20	85	86	87	88	89	90	91	92	93	94	95	96
SP21	97	98	99	100	101	102	103	104	105	106	107	108
SP22	109	110	111	112	113	114	115	116	117	118	119	120
SP23	121	122	123	124	125	126	127	128	129	130	131	132
SP24	133	134	135	136	137	138	139	140	141			

Isolation of peripheral blood mononuclear cells

Peripheral blood was obtained from ten healthy donors at San Raffaele Hospital upon informed consent. Peripheral blood mononuclear cells were isolated using Ficoll-Hypaque density gradient centrifugation.

Immortalized B cells

5 Autologous B cells were isolated from PBMCs of healthy donors using the CD19 Microbeads (Miltenyi Biotech). Cells were transduced with a lentiviral vector harbouring the BCL-6/BCL-XL transgene (Kwakkenbos, M. J. et al. Nat. Med.16, 123 (2009)) and the H/F pseudotype (Lévy, C. et al. Molecular Therapy20 9, 1699–1712, (2012)) and cultured in IMDM supplemented with 10% fetal bovine serum (FBS), penicillin-streptomycin, and 10 ng/ml of
10 IL21 (Miltenyi Biotech). B-cells were re-stimulated every 5 days by co-culture with irradiated (50 Gy) mouse L-cell fibroblasts expressing CD40L (3T3-CD40L) at a B-cell:3T3-CD40L ratio of 10:1.

Cell lines

We cultured the T2 and K562 cell lines in RPMI 1640 (GIBCO-BRL) supplemented with
15 penicillin, streptomycin, glutamine and 10% FBS (BioWhittaker).

Leukemic cells

Primary AML cells were obtained from San Raffaele Hospital (OSR) Leukemia biobank and selected according to the expression of WT1 by quantitative PCR and to the HLA typing. All EBV-BLCLs and primary leukemia cells were typed for HLA-A, HLA-B, HLA-C, HLA-DR and
20 HLA-DQ alleles at high resolution at the HLA laboratory of the OSR.

Flow cytometry

We used FITC-, PE-, PerCP-, APC-, PE-Cychrome 7–, APC Cychrome 7–, Pacific Blue and Brilliant Violet–conjugated antibodies directed to human CD3, CD4, CD8, CD107a, IFN γ , TNF α , CD33, CD117, CD34, CD14, V β 21.3, V β 8, V β 7.2 and HLA-A2. APC fluorescently-
25 labelled WT1 VLDFAPPGA (SEQ ID NO: 157) and PE fluorescently-labelled WT1 RMFPNAPYL (SEQ ID NO: 255) dextramers were used following the manufacturer's instructions. Cells were incubated with antibodies for 15 minutes at 4°C and washed with phosphate-buffered saline containing 1% FBS. Samples were run through a fluorescence-activated cell sorter (FACS) Canto II flow cytometer (BD Biosciences), and data were
30 analyzed by Flow Jo software (Tree star Inc). For intracellular evaluation of cytokine secretion and expression of degranulation markers, the Fix/Perm buffer set (Biolegend) was used according to manufacturer instructions.

Stimulation, isolation and expansion of WT1-specific T-cells

Freshly isolated PBMCs were resuspended in X-VIVO supplemented with 5% human AB serum, 2 mM glutamine and 1 µg/ml CD28 monoclonal antibody, seeded at a density of 10^7 cells/ml and stimulated with the WT1 overlapping peptide pool, each peptide present at a concentration of 1 µg/ml.

Antigen-specific T-cells were isolated after 26-30 hours by CD137 expression. More specifically, cells were stained with the PE-conjugated CD137 antibody and sorted using anti-PE microbeads (Miltenyi Biotech). The CD137⁻ fraction was depleted of the CD3 cells using CD3-Microbeads (Miltenyi Biotech), irradiated 30 Gy and used as peptide-loaded APCs in a co-culture with the CD137⁺ fraction at a ratio of 100:1 when possible or at least 20:1 and a final density of 5×10^6 cells/ml. X-VIVO supplemented with 5% human AB serum, 5 ng/ml IL7, 5 ng/ml IL15 and 10 ng/ml IL21 was used as the medium. Media, including cytokines, was replaced every 2-3 days.

Re-stimulation of expanded antigen-specific T-cells

Cells were re-stimulated every 7-14 days with WT1-pulsed autologous APCs (PBMC CD3-depleted cells; immortalized B cells). In the initial re-stimulations, cells were washed 2 days before and plated in cytokine-free medium. APCs were irradiated with 30 Gy, pulsed with the peptide pool overnight and co-cultured with effector cells in X-VIVO supplemented with 5% human AB serum, 1 µg/ml CD28 monoclonal antibody and IL7 (5 ng/ml), IL15 (5 ng/ml), IL21 (10 ng/ml).

Assessment of T cell response

The percentage of T-cells responding to the WT1 peptide pool was measured by performing a 6 hours co-culture of the effector cells with autologous APCs (ratio of at least 1:1) pulsed with the desired antigen (WT1 peptide pool, WT1 subpools, WT1 individual peptides, unrelated peptide pool as control). Co-cultures were seeded in X-VIVO supplemented with 5% human AB serum and supplemented with the CD28 monoclonal antibody (1 µg/ml), Golgi Stop (BD) and CD107a-FITC antibody for assessment of degranulation. Cells were then fixed, permeabilized and stained intracellularly to determine the percentage of CD3⁺CD8⁺ or CD3⁺CD4⁺ cells expressing IFNγ and CD107a.

Mapping of immunogenic peptides

T-cells stimulated with the WT1 pool were seeded in different wells and co-cultured with autologous APCs loaded with one of each of the WT1 subpools at a ratio of at least 1:1. T-

cell responses to each subpool were measured as previously described by FACS analysis. Deconvolution of the mapping grid was essential to determine which shared peptides were eliciting a T cell response. Once determined the immunogenic peptides, T-cells were further stimulated with APCs loaded with the individual peptides to confirm their immunogenicity.

5 *Evaluation of T cell ability to recognize WT1-expressing cells*

WT1-specificity and HLA-restricted ability of T-cells to recognize target cells was measured with different experimental procedures. For T-cells derived from HD1, secretion of CD107a was determined by FACS analysis after 6 hours co-culture with target cells; for T cells derived from HD3, elimination index was calculated as the total number of target cells still present after 4 days co-culture with the WT1-specific T-cells divided by the total number of target cells alone; for T-cells derived from HD4, the percentage of CD33⁺ target cells (AML primary cells harbouring the HLA alleles of interest and as control, of AML primary cells not harbouring the specific HLA allele) still present after 3 days co-culture with CD3⁺ WT1-specific T cells was assessed by cytofluorimetric analysis.

15 *Assessment of T cell clonality*

In order to determine the clonality of the expanded WT1-specific T cells, the IO Test Beta Mark TCR V beta repertoire kit was used according to manufacturer's recommendations.

TCR repertoire sequencing

WT1-specific T cells were collected at different time points over the co-culture time frame and RNA was extracted by using the Arcturus Pico Pure RNA extraction kit. Complementarity determining region (CDR) 3 sequences of the WT1-specific T cells were amplified by using a modified RACE approach (Ruggiero,E. et al. Nat. Commun. 6,8081 (2015)). Samples were sequenced by using an IlluminaMiSeq sequencer and CDR3 clonotypes identified using the MiXCR software (Bolotin, DA et al. Nature Methods 12, 380–381 (2015)).

Lentiviral vectors

TCR α and β chain genes isolated from HD1 and HD3 were codon-optimized, cysteine-modified and cloned in a lentiviral vector (LV) under a bidirectional promoter. The amino acid (aa) and nucleotide (nt) sequences were:

		SEQ ID NO:
HD1	ATGGAAACCCTGCTGAAGGTGCTGAGCGGCACACTGCTGTGGCAGC	256

α chain – seq nt	TGACATGGGTCCGATCTCAGCAGCCTGTGCAGTCTCCTCAGGCCGT GATTCTGAGAGAAGGCGAGGACGCCGTGATCAACTGCAGCAGCTCT AAGGCCCTGTACAGCGTGCACTGGTACAGACAGAAGCACGGCGAGG CCCCTGTGTTCTCTGATGATCCTGCTGAAAGGCGGCGAGCAGAAGGG CCACGAGAAGATCAGCGCCAGCTTCAACGAGAAGAAGCAGCAGTCC AGCCTGTACCTGACAGCCAGCCAGCTGAGCTACAGCGGCACCTACT TTTGTGGCACCGCCTGGATCAACGACTACAAGCTGTCTTTCGGAGCC GGCACCACAGTGACAGTGCGGGCCAATATTCAGAACCCCGATCCTG CCGTGTACCAGCTGAGAGACAGCAAGAGCAGCGACAAGAGCGTGTG CCTGTTACCGACTTCGACAGCCAGACCAACGTGTCCCAGAGCAAG GACAGCGACGTGTACATACCGATAAGTGCGTGTGACATGCGGA GCATGGACTTCAAGAGCAACAGCGCCGTGGCCTGGTCCAACAAGAG CGATTTGCGCTGCGCCAACGCCTTCAACAACAGCATTATCCCCGAGG ACACATTCTTCCCAAGTCTTGAGAGCAGCTGCGACGTGAAGCTGGT GGAAAAGAGCTTCGAGACAGACACCAACCTGAACTTCCAGAACCTGA GCGTGATCGGCTTCAGAATCCTGCTGCTCAAGGTGGCCGGCTTCAA CCTGCTGATGACCCTGAGACTGTGGTCCAGC	
HD1 α chain – seq aa	METLLKVLSTLLWQLTWVRSQQPVQSPQAVILREGEDAVINCSSSKAL YSVHWYRQKHGEAPVFLMILLKGGEQKGHEKISASFNEKKQQSSLYLTA SQLSYSGTYFCGTAWINDYKLSFGAGTTVTVRANIQNPDPAVYQLRDSK SSDKSVCLFTDFDSQTNVSQSKSDSVYITDKCVLDMRSMDFKSNSAVA WSNKSDFACANAFNNSIIPEDTFFPSPESSCDVKLVEKSFETDTNLFQ NLSVIGFRILLKLVAGFNLLMTLRLWSS	257
HD1 β chain – seq nt	ATGGGATCTTGGACACTGTGTTGCGTGTCCCTGTGCATCCTGGTGGC CAAGCACACAGATGCCGGCGTGATCCAGTCTCCTAGACACGAAGTG ACCGAGATGGGCCAAGAAGTGACCCTGCGCTGCAAGCCTATCAGCG GCCACGATTACCTGTTCTGGTACAGACAGACCATGATGAGAGGCCTG GAACTGCTGATCTACTTCAACAACAACGTGCCCATCGACGACAGCGG CATGCCCCGAGGATAGATTCAGCGCCAAGATGCCCAACGCCAGCTTC AGCACCTGAAGATCCAGCCTAGCGAGCCCAGAGATAGCGCCCGTGT ACTTCTGCGCCAGCAGAAAGACAGGCGGCTACAGCAATCAGCCCCA GCACTTTGGAGATGGCACCCGGCTGAGCATCCTGGAAGATCTGAAG AACGTGTTCCACCTGAGGTGGCCGTGTTTCGAGCCTTCTGAGGCCG AGATCAGCCACACACAGAAAGCCACACTCGTGTGTCTGGCCACCGG CTTCTATCCCGATCACGTGGAACGTCTTGGTGGGTCAACGGCAAAG AGGTGCACAGCGGCGTCTGTACCGATCCTCAGCCTCTGAAAGAGCA GCCCCGCTCTGAACGACAGCAGATACTGCCTGAGCAGCAGACTGAGA GTGTCCGCCACCTTCTGGCAGAACCCAGAAACCACTTCAGATGCCA GGTGCAGTTCTACGGCCTGAGCGAGAACGATGAGTGGACCCAGGAT AGAGCCAAGCCTGTGACACAGATCGTGTCTGCCGAAGCCTGGGGCA GAGCCGATTGTGGCTTTACCAGCGAGAGCTACCAGCAGGGCGTGCT GTCTGCCACAATCCTGTACGAGATCCTGCTGGGCAAAGCCACTCTGT ACGCCGTGCTGGTGTCTGCCCTGGTGTGCTGATGGCCATGGTCAAGCG GAAGGATAGCAGGGGC	258
HD1 β chain – seq aa	MGSWTLCCVSLCILVAKHTDAGVIQSPRHEVTEMGQEVTLRCKPISGHD YLFWYRQTM MRGLELLIYFNNNVPIDDSGMPEDRFS AKMPNASFSTLKI QPSEPRDSAVYFCASRKTGGYSNQPHFGDGT RL SILEDLKNVFPPEV AVFEPSEAEISHTQKATLVCLATGFYPDHVELSWWVNGKEVHSGVCTD PQPLKEQPALNDSRYCLSSRLRV SATFWQNPRNHFR CQVQFYGLSEN DEWTQDRAKPVTQIVSAEAWGRADCGFTSESYQQGVLSATILYEILLGK ATLYAVLV SALVLMAMVKRKDSRG	259
HD3 α chain – seq nt	ATGATCAGCCTGAGAGTGCTGCTGGTCATCCTGTGGCTGCAGCTGT CTTGGGTCTGGTCCCAGCGGAAAGAGGTGGAACAGGACCCCGGAC CTTTCAATGTGCCTGAAGGCGCCACCGTGGCCTTCAACTGCACCTAC AGCAATAGCGCCAGCCAGAGCTTCTTCTGGTACAGACAGGACTGCC GGAAAGAACCCAAGCTGCTGATGAGCGTGTACAGCAGCGGCAACGA	260

	GGACGGCAGATTACAGCCCAGCTGAACAGAGCCAGCCAGTACATC AGCCTGCTGATCCGGGATAGCAAGCTGAGCGATAGCGCCACCTACC TGTGCGTGGTCAACCTGCTGTCTAATCAAGGCGGCAAGCTGATCTTC GGCCAGGGGCACAGAGCTGAGCGTGAAGCCCAACATTCAGAACCCCG ATCCTGCCGTGTACCAGCTGAGAGACAGCAAGAGCAGCGACAAGAG CGTGTGCCTGTTACCGACTTCGACAGCCAGACCAACGTGTCCCAG AGCAAGGACAGCGACGTGTACATCACCGATAAGTGCGTGCTGGACA TGCGGAGCATGGACTTCAAGAGCAACAGCGCCGTGGCCTGGTCCAA CAAGAGCGATTTGCGCTGCGCCAACGCCTTCAACAACAGCATTATCC CCGAGGACACATTCTTCCCAAGTCCTGAGAGCAGCTGCGACGTGAA GCTGGTGGAAAAGAGCTTCGAGACAGACACCAACCTGAACTTCCAG AACCTGTCCGTGATCGGCTTCCGGATCCTGCTGCTGAAAGTGGCCG GCTTCAACCTCCTGATGACCCTGAGACTGTGGTCCAGC	
HD3 α chain – seq aa	MISLRVLLVILWLQLSWVWSQRKEVEQDPGPFNVPEGATVAFNCTYSN SASQSFFWYRQDCRKEPKLLMSVYSSGNEDGRFTAQLNRASQYISLLIR DSKLSDSATYLCVVNLLSNQGGKLIFGQGTELSVKPNIQNPDPVYQLR DSKSSDKSVCLFTDFDSQTNVSQSKDSVDYITDKCVLDMRSMDFKSNS AVAWSNKSDFACANAFNNSIIPEDTFFPSPESSCDVKLVEKSFETDTNLN FQNLSVIGFRILLKLVAGFNLLMTLRLWSS	261
HD3 β chain – seq nt	ATGGGATGTAGACTTCTGTGTTGCGCCGTGCTGTGTCTGCTTGGAGC TGGCGAACTGGTGCCTATGGAAACCGGCGTGACCCAGACACCTAGA CACCTGGTCATGGGCATGACAAACAAGAAAAGCCTGAAGTGCGAGC AGCACCTGGGCCACAATGCCATGTACTGGTACAAGCAGAGCGCCAA GAAACCCCTGGAAGTATGTTCTGTGTACAGCCTGGAAGAGAGGGTC GAGAACAAACAGCGTGCCCAGCAGATTACGCCCTGAGTGCCCTAATA GCAGCCACCTGTTTCTGCATCTGCACACCCTGCAGCCTGAGGACTCT GCCCTGTATCTGTGTGCCAGCAGCCAGGACTACCTGGTGTCCAACG AGAAGCTGTTCTTCGGCAGCGGCACACAGCTGAGCGTGCTGGAAGA TCTGAAGAACGTGTTCCACCTGAGGTGGCCGTGTTTCGAGCCTTCTG AGGCCGAGATCAGCCACACACAGAAAGCCACACTCGTGTGTCTGGC CACCGGCTTCTATCCCGATCACGTGGAAGTGTCTTGGTGGGTCAACG GCAAAGAGGTGCACAGCGGCGTCTGTACCGATCCTCAGCCTCTGAA AGAGCAGCCCGCTCTGAACGACAGCAGATACTGCCTGAGCAGCAGA CTGAGAGTGTCCGCCACCTTCTGGCAGAACCCCAAGAAACCACTTCA GATGCCAGGTGCAGTTCTACGGCCTGAGCGAGAACGATGAGTGGAC CCAGGATAGAGCCAAGCCTGTGACACAGATCGTGTCTGCCGAAGCC TGGGGCAGAGCCGATTGTGGCTTTACCAGCGAGAGCTACCAGCAGG GCGTGCTGTCTGCCACAATCCTGTACGAGATCCTGCTGGGAAAAGC CACTCTGTACGCTGTGCTGGTGTCCGCTCTGGTGCTGATGGCCATG GTCAAGCGGAAGGATAGCAGGGGC	262
HD3 β chain – seq aa	MGCRLLCCAFLCLLGAGELVPMETGVTQTPRHLVMGMTNKKSLKCEQH LGHNAMEYWKQSAKKPLELMFVYSLEERVENNSVPSRFSPECNSSLH FLHLHTLQPEDSALYLCASSQDYLVSNELFFGSGTQLSVLEDLKNVFP EVAVFEPSEAEISHTQKATLVCLATGFYPDHVELSWVNGKEVHSGVC TDPQPLKEQPALNDSRYCLSSRLRVSATFWQNPRNHFRCQVQFYGLSE NDEWTQDRAKPVTVQIVSAEAWGRADCGFTSESYQQGVLSATILYEILLG KATLYAVLVLSALVLMAMVKRKDSRG	263

LVs were packaged by an integrase-competent third-generation construct and pseudotyped by the vesicular stomatitis virus (VSV) envelope. As control, we included the LV encoding for a WT1 126-134 TCR recognizing the RMFPNAPYL (SEQ ID NO: 255) peptide.

5 Vector transductions

For transduction with WT1-TCR lentiviral vector, T lymphocytes isolated from a healthy individual were activated and sorted using magnetic beads conjugated to antibodies to CD3 and CD28 (ClinExVivo CD3/CD28; Invitrogen), following the manufacturer instructions, and cultured in Iscove's Modified Dulbecco's Media (IMDM) (GIBCO-BRL) supplemented with penicillin, streptomycin, 10% FBS and 5 ng ml⁻¹ of each IL-7 and IL-15 (PeproTech). For transduction, T lymphocytes were plated at 2.5 × 10⁶ cells ml⁻¹ and infected with the LV for 24 h. Afterwards, T cells were cultured at 10⁶ cells ml⁻¹ and expanded. Transduction efficiency was determined by measuring the percentage of the CD3 T cells expressing the specific dextramers. Cells were sorted using APC or PE-fluorescently-labelled HLA-A*0201 dextramer specific for the VLDFAPPGA (SEQ ID NO: 157) or RMFPNAPYL (SEQ ID NO: 255) peptide (Immudex) using anti-APC or anti-PE microbeads (Miltenyi Biotec) following the manufacturer instructions.

Functional assays

The ability of HD1, HD3 and WT1 126-134 TCR-transferred T-cells to recognize WT1-expressing target cells was measured upon co-culture with (a) T2 cells either pulsed with the WT1 126-134 peptide or with the VLDFAPPGA (SEQ ID NO: 157) peptide (effector:target ratio=1:1); (b) K562 cells either wild type (K562) or genetically modified in order to express the HLA-A*0201 allele (effector:target ratio=1:1); (c) 3 different primary AML blasts selected according to the expression of the HLA-A*0201 allele and of the WT1 antigen (effector:target ratio=5:1). For the co-culture with T2 and K562 cell lines, we included untransduced T cells as control. After 3 days of culture, the percentage of target cells was assessed by cytofluorimetric analysis. The elimination index was calculated as follows: 1-(total number of target cells still present after 3 days co-culture with the WT1-specific T-cells/total number of target cells alone).

25

All publications mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described present invention will be apparent to those skilled in the art without departing from the scope and spirit of the present invention. Although the present invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in cell biology, immunology, immunotherapy, molecular biology, oncology, or related fields are intended to be within the scope of the following claims.

30

CLAIMS

1. A T-cell receptor (TCR), which binds to a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the TCR:

- 5 (i) comprises a CDR3 α comprising the amino acid sequence of CGTAWINDYKLSF (SEQ ID NO: 3) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRKTGGYSNQPQHF (SEQ ID NO: 8) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- 10 (ii) comprises a CDR3 α comprising the amino acid sequence of CVVNLLSNQGGKLIF (SEQ ID NO: 36) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSQDYLVSNKLFF (SEQ ID NO: 41) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- 15 (iii) comprises a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- 20 (iv) comprises a CDR3 α comprising the amino acid sequence of CAVRLSGSARQLTF (SEQ ID NO: 14) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- 25 (v) comprises a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLLGDEQYF (SEQ ID NO: 24) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- 30 (vi) comprises a CDR3 α comprising the amino acid sequence of CAYRSLKYGNKLVF (SEQ ID NO: 19) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLVALQGAGEQYF (SEQ ID NO: 30) or a variant thereof having up to three amino acid substitutions, additions or deletions;

- (vii) comprises a CDR3 α comprising the amino acid sequence of CATDAYSGNTPLVF (SEQ ID NO: 47) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRAAGLDTEAFF (SEQ ID NO: 57) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (viii) comprises a CDR3 α comprising the amino acid sequence of CATDAYSGNTPLVF (SEQ ID NO: 47) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASTQTPYEQYF (SEQ ID NO: 63) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (ix) comprises a CDR3 α comprising the amino acid sequence of CATDAYSGNTPLVF (SEQ ID NO: 47) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSTVGGEDYGYTF (SEQ ID NO: 69) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (x) comprises a CDR3 α comprising the amino acid sequence of CAVRAEIYNQGGKLIF (SEQ ID NO: 52) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRAAGLDTEAFF (SEQ ID NO: 57) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xi) comprises a CDR3 α comprising the amino acid sequence of CAVRAEIYNQGGKLIF (SEQ ID NO: 52) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASTQTPYEQYF (SEQ ID NO: 63) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xii) comprises a CDR3 α comprising the amino acid sequence of CAVRAEIYNQGGKLIF (SEQ ID NO: 52) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSTVGGEDYGYTF (SEQ ID NO: 69) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xiii) comprises a CDR3 α comprising the amino acid sequence of CAASMAGAGSYQLTF (SEQ ID NO: 75) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of

CAISVGQGALYEQYF (SEQ ID NO: 80) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xiv) comprises a CDR3 α comprising the amino acid sequence of CAASMAGAGSYQLTF (SEQ ID NO: 75) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSVARDRRNYGYTF (SEQ ID NO: 86) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xv) comprises a CDR3 α comprising the amino acid sequence of CAANNARLMF (SEQ ID NO: 92) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSDTRAREQFF (SEQ ID NO: 97) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xvi) comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xvii) comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSVGGSGSYNEQFF (SEQ ID NO: 169) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xviii) comprises a CDR3 α comprising the amino acid sequence of CAVEATDSWGKLQF (SEQ ID NO: 108) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xix) comprises a CDR3 α comprising the amino acid sequence of CAVEATDSWGKLQF (SEQ ID NO: 108) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSVGGSGSYNEQFF (SEQ ID NO: 169) or a variant thereof having up to three amino acid substitutions, additions or deletions;

- (xx) comprises a CDR3 α comprising the amino acid sequence of CAVRTSYDKVIF (SEQ ID NO: 113) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxi) comprises a CDR3 α comprising the amino acid sequence of CAVRTSYDKVIF (SEQ ID NO: 113) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSVGGSGSYNEQFF (SEQ ID NO: 169) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxii) comprises a CDR3 α comprising the amino acid sequence of CAVTVGNKLVF (SEQ ID NO: 175) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASRGWREQFF (SEQ ID NO: 180) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxiii) comprises a CDR3 α comprising the amino acid sequence of CAARSYNTDKLIF (SEQ ID NO: 186) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSWGYQETQYF (SEQ ID NO: 196) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxiv) comprises a CDR3 α comprising the amino acid sequence of CAARSYNTDKLIF (SEQ ID NO: 186) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPTGGEYYGYTF (SEQ ID NO: 202) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxv) comprises a CDR3 α comprising the amino acid sequence of CAARSYNTDKLIF (SEQ ID NO: 186) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSYPLRTGRYNSYNSPLHF (SEQ ID NO: 208) or a variant thereof having up to three amino acid substitutions, additions or deletions;
- (xxvi) comprises a CDR3 α comprising the amino acid sequence of CAASYNNARLMF (SEQ ID NO: 191) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of

CASSWGYQETQYF (SEQ ID NO: 196) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxvii) comprises a CDR3 α comprising the amino acid sequence of CAASYNNARLMF (SEQ ID NO: 191) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPTGGEYYGYTF (SEQ ID NO: 202) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxviii) comprises a CDR3 α comprising the amino acid sequence of CAASYNNARLMF (SEQ ID NO: 191) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSSYPLRTGRYNSYNSPLHF (SEQ ID NO: 208) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxix) comprises a CDR3 α comprising the amino acid sequence of CAASGGRDDKIIF (SEQ ID NO: 214) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSYSRTESTDTQYF (SEQ ID NO: 219) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxx) comprises a CDR3 α comprising the amino acid sequence of CAANNARLMF (SEQ ID NO: 92) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPGQHGELEFF (SEQ ID NO: 271) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxi) comprises a CDR3 α comprising the amino acid sequence of CAASATGNQFYF (SEQ ID NO: 266) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSDTRAREQFF (SEQ ID NO: 97) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxii) comprises a CDR3 α comprising the amino acid sequence of CAASATGNQFYF (SEQ ID NO: 266) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSPGQHGELEFF (SEQ ID NO: 271) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxiii) comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDSVSGNTIYF (SEQ ID NO: 163) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxiv) comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSVGGSGSYNEQFF (SEQ ID NO: 169) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxv) comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxvi) comprises a CDR3 α comprising the amino acid sequence of CATDGDSSYKLIF (SEQ ID NO: 277) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLGLSISQETQYF (SEQ ID NO: 288) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxvii) comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLGLSISQETQYF (SEQ ID NO: 288) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxviii) comprises a CDR3 α comprising the amino acid sequence of CAVEATDSWGKLQF (SEQ ID NO: 108) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CASSLGLSISQETQYF (SEQ ID NO: 288) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxix) comprises a CDR3 α comprising the amino acid sequence of CAVRTSYDKVIF (SEQ ID NO: 113) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of

CASSLGLSISQETQYF (SEQ ID NO: 288) or a variant thereof having up to three amino acid substitutions, additions or deletions.

(xxxx) comprises a CDR3 α comprising the amino acid sequence of CAERLNTDKLIF (SEQ ID NO: 103) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions;

(xxxxi) comprises a CDR3 α comprising the amino acid sequence of CAVEATDSWGKLQF (SEQ ID NO: 108) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions; or

(xxxxii) comprises a CDR3 α comprising the amino acid sequence of CAVRTSYDKVIF (SEQ ID NO: 113) or a variant thereof having up to three amino acid substitutions, additions or deletions, and a CDR3 β comprising the amino acid sequence of CSARDVLTGDYGYTF (SEQ ID NO: 282) or a variant thereof having up to three amino acid substitutions, additions or deletions.

2. A TCR according to claim 1 comprising the following CDR sequences:

(i) CDR1 α - KALYS (SEQ ID NO: 1),
 CDR2 α - LLKGGEQ (SEQ ID NO: 2),
 CDR3 α - CGTAWINDYKLSF (SEQ ID NO: 3),
 CDR1 β - SGHDY (SEQ ID NO: 6),
 CDR2 β - FNNNVP (SEQ ID NO: 7), and
 CDR3 β - CASRKTGGYSNQPQHF (SEQ ID NO: 8),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;

(ii) CDR1 α - NSASQS (SEQ ID NO: 34),
 CDR2 α - VYSSGN (SEQ ID NO: 35),
 CDR3 α - CVVNLLSNQGGKLIF (SEQ ID NO: 36),
 CDR1 β - LGHNA (SEQ ID NO: 39),
 CDR2 β - YSLEER (SEQ ID NO: 40), and
 CDR3 β - CASSQDYLVSNEKLFF (SEQ ID NO: 41),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (iii) CDR1 α - SSVPPY (SEQ ID NO: 12),
 CDR2 α - Y TSAATLV (SEQ ID NO: 13),
 5 CDR3 α - CAVRLSGSARQLTF (SEQ ID NO: 14),
 CDR1 β - SGHAT (SEQ ID NO: 22),
 CDR2 β - FQNNGV (SEQ ID NO: 23), and
 CDR3 β - CASSLLGDEQYF (SEQ ID NO: 24),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (iv) CDR1 α - SSVPPY (SEQ ID NO: 12),
 CDR2 α - Y TSAATLV (SEQ ID NO: 13),
 CDR3 α - CAVRLSGSARQLTF (SEQ ID NO: 14),
 CDR1 β - SGHTA (SEQ ID NO: 28),
 15 CDR2 β - FQGNSA (SEQ ID NO: 29), and
 CDR3 β - CASSLVALQGAGEQYF (SEQ ID NO: 30),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (v) CDR1 α - TSESDYY (SEQ ID NO: 17),
 20 CDR2 α - QEAYKQQN (SEQ ID NO: 18),
 CDR3 α - CAYRSLKYGNKLVF (SEQ ID NO: 19),
 CDR1 β - SGHAT (SEQ ID NO: 22),
 CDR2 β - FQNNGV (SEQ ID NO: 23), and
 CDR3 β - CASSLLGDEQYF (SEQ ID NO: 24),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (vi) CDR1 α - TSESDYY (SEQ ID NO: 17),
 CDR2 α - QEAYKQQN (SEQ ID NO: 18),
 CDR3 α - CAYRSLKYGNKLVF (SEQ ID NO: 19),
 30 CDR1 β - SGHTA (SEQ ID NO: 28),
 CDR2 β - FQGNSA (SEQ ID NO: 29), and
 CDR3 β - CASSLVALQGAGEQYF (SEQ ID NO: 30),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (vii) CDR1 α – TSINN (SEQ ID NO: 45),
 CDR2 α - IRSNERE (SEQ ID NO: 46),
 CDR3 α – CATDAYSGNTPLVF (SEQ ID NO: 47),
 CDR1 β - MNHNS (SEQ ID NO: 55),
 5 CDR2 β - SASEGT (SEQ ID NO: 56), and
 CDR3 β - CASRAAGLDTEAFF (SEQ ID NO: 57),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (viii) CDR1 α – TSINN (SEQ ID NO: 45),
 10 CDR2 α - IRSNERE (SEQ ID NO: 46),
 CDR3 α – CATDAYSGNTPLVF (SEQ ID NO: 47),
 CDR1 β – MNHNY (SEQ ID NO: 61),
 CDR2 β - SVGAGI (SEQ ID NO: 62), and
 CDR3 β - CASTQTPYEQYF (SEQ ID NO: 63),
 15 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (ix) CDR1 α – TSINN (SEQ ID NO: 45),
 CDR2 α - IRSNERE (SEQ ID NO: 46),
 CDR3 α – CATDAYSGNTPLVF (SEQ ID NO: 47),
 20 CDR1 β – SGHNS (SEQ ID NO: 67),
 CDR2 β - FNNNVP (SEQ ID NO: 68), and
 CDR3 β – CASSTVGGEDYGYTF (SEQ ID NO: 69),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- 25 (x) CDR1 α - DSAIYN (SEQ ID NO: 50),
 CDR2 α – IQSSQRE (SEQ ID NO: 51),
 CDR3 α - CAVRAEIYNQGGKLIF (SEQ ID NO: 52),
 CDR1 β - MNHNS (SEQ ID NO: 55),
 CDR2 β - SASEGT (SEQ ID NO: 56), and
 30 CDR3 β - CASRAAGLDTEAFF (SEQ ID NO: 57),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (xi) CDR1 α - DSAIYN (SEQ ID NO: 50),
 CDR2 α – IQSSQRE (SEQ ID NO: 51),
 35 CDR3 α - CAVRAEIYNQGGKLIF (SEQ ID NO: 52),

CDR1 β – MNHNY (SEQ ID NO: 61),
 CDR2 β - SVGAGI (SEQ ID NO: 62), and
 CDR3 β - CASTQTPYEQYF (SEQ ID NO: 63),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xii) CDR1 α - DSAIYN (SEQ ID NO: 50),
 CDR2 α – IQSSQRE (SEQ ID NO: 51),
 CDR3 α - CAVRAEIYNQGGKLIF (SEQ ID NO: 52),
 CDR1 β – SGHNS (SEQ ID NO: 67),
 CDR2 β - FNNNVP (SEQ ID NO: 68), and
 CDR3 β – CASSTVGGEDYGYTF (SEQ ID NO: 69),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xiii) CDR1 α - DSASNY (SEQ ID NO: 73),
 CDR2 α – IRSNVGE (SEQ ID NO: 74),
 CDR3 α - CAASMAGAGSYQLTF (SEQ ID NO: 75),
 CDR1 β - ENHRY (SEQ ID NO: 78),
 CDR2 β – SYGVKD (SEQ ID NO: 79), and
 CDR3 β - CAISVGQGALYEQYF (SEQ ID NO: 80),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xiv) CDR1 α - DSASNY (SEQ ID NO: 73),
 CDR2 α – IRSNVGE (SEQ ID NO: 74),
 CDR3 α - CAASMAGAGSYQLTF (SEQ ID NO: 75),
 CDR1 β – SGDLS (SEQ ID NO: 84),
 CDR2 β – YYNGEE (SEQ ID NO: 85), and
 CDR3 β – CASSVARDRRNYGYTF (SEQ ID NO: 86),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xv) CDR1 α - NSMFDY (SEQ ID NO: 90),
 CDR2 α – ISSIKDK (SEQ ID NO: 91),
 CDR3 α - CAANNARLMF (SEQ ID NO: 92),
 CDR1 β – SGHNS (SEQ ID NO: 95),
 CDR2 β – FNNNVP (SEQ ID NO: 96), and
 CDR3 β – CASSDTRAREQFF (SEQ ID NO: 97),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (xvi) CDR1 α - DSSSTY (SEQ ID NO: 101),
 CDR2 α – IFSNMDM (SEQ ID NO: 102),
 5 CDR3 α - CAERLNTDKLIF (SEQ ID NO: 103),
 CDR1 β – DFQATT (SEQ ID NO: 161),
 CDR2 β – SNEGSKA (SEQ ID NO: 162), and
 CDR3 β – CSARDSVSGNTIYF (SEQ ID NO: 163),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (xvii) CDR1 α - DSSSTY (SEQ ID NO: 101),
 CDR2 α – IFSNMDM (SEQ ID NO: 102),
 CDR3 α - CAERLNTDKLIF (SEQ ID NO: 103),
 CDR1 β – SQVTM (SEQ ID NO: 167),
 15 CDR2 β – ANQGSEA (SEQ ID NO: 168), and
 CDR3 β – CSVGGSGSYNEQFF (SEQ ID NO: 169),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (xviii) CDR1 α - DSVNN (SEQ ID NO:106),
 20 CDR2 α – IPSGT (SEQ ID NO: 107),
 CDR3 α - CAVEATDSWGKLQF (SEQ ID NO: 108),
 CDR1 β – DFQATT (SEQ ID NO: 161),
 CDR2 β – SNEGSKA (SEQ ID NO: 162), and
 CDR3 β – CSARDSVSGNTIYF (SEQ ID NO: 163),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (xix) CDR1 α - DSVNN (SEQ ID NO:106),
 CDR2 α – IPSGT (SEQ ID NO: 107),
 CDR3 α - CAVEATDSWGKLQF (SEQ ID NO: 108),
 30 CDR1 β – SQVTM (SEQ ID NO: 167),
 CDR2 β – ANQGSEA (SEQ ID NO: 168), and
 CDR3 β – CSVGGSGSYNEQFF (SEQ ID NO: 169),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

- (xx) CDR1 α - DSASNY (SEQ ID NO: 111),
 CDR2 α – IRSNVGE (SEQ ID NO: 112),
 CDR3 α - CAVRTSYDKVIF (SEQ ID NO: 113),
 CDR1 β – DFQATT (SEQ ID NO: 161),
 5 CDR2 β – SNEGSKA (SEQ ID NO: 162), and
 CDR3 β – CSARDSVSGNTIYF (SEQ ID NO: 163),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (xxi) CDR1 α - DSASNY (SEQ ID NO: 111),
 10 CDR2 α – IRSNVGE (SEQ ID NO: 112),
 CDR3 α - CAVRTSYDKVIF (SEQ ID NO: 113),
 CDR1 β – SQVTM (SEQ ID NO: 167),
 CDR2 β – ANQGSEA (SEQ ID NO: 168), and
 CDR3 β – CSVGGSGSYNEQFF (SEQ ID NO: 169),
 15 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (xxii) CDR1 α - VGISA (SEQ ID NO: 173),
 CDR2 α – LSSGK (SEQ ID NO: 174),
 CDR3 α - CAVTVGNKLVF (SEQ ID NO: 175),
 20 CDR1 β – MNHNS (SEQ ID NO: 178),
 CDR2 β – SASEGT (SEQ ID NO: 179), and
 CDR3 β – CASRGWREQFF (SEQ ID NO: 180),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- 25 (xxiii) CDR1 α - VGISA (SEQ ID NO: 184),
 CDR2 α – LSSGK (SEQ ID NO: 185),
 CDR3 α - CAARSYNTDKLIF (SEQ ID NO: 186),
 CDR1 β – SGHTS (SEQ ID NO: 194),
 CDR2 β – YDEGEE (SEQ ID NO: 195), and
 30 CDR3 β – CASSWGYQETQYF (SEQ ID NO: 196),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;
- (xxiv) CDR1 α - VGISA (SEQ ID NO: 184),
 CDR2 α – LSSGK (SEQ ID NO: 185),
 35 CDR3 α - CAARSYNTDKLIF (SEQ ID NO: 186),

CDR1 β – KGHSH (SEQ ID NO: 200),
 CDR2 β – LQKENI (SEQ ID NO: 201), and
 CDR3 β – CASSPTGGEYYGYTF (SEQ ID NO: 202),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxv) CDR1 α - VGISA (SEQ ID NO: 184),
 CDR2 α – LSSGK (SEQ ID NO: 185),
 CDR3 α - CAARSYNTDKLIF (SEQ ID NO: 186),
 CDR1 β – MNHEY (SEQ ID NO: 206),
 CDR2 β – SVGAGI (SEQ ID NO: 207), and
 CDR3 β – CASSSYPLRTGRYNSYNSPLHF (SEQ ID NO: 208),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxvi) CDR1 α - NSMFDY (SEQ ID NO: 189),
 CDR2 α – ISSIKDK (SEQ ID NO: 190),
 CDR3 α - CAASYNNAARLMF (SEQ ID NO: 191),
 CDR1 β – SGHTS (SEQ ID NO: 194),
 CDR2 β – YDEGEE (SEQ ID NO: 195), and
 CDR3 β – CASSWGYQETQYF (SEQ ID NO: 196),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxvii) CDR1 α - NSMFDY (SEQ ID NO: 189),
 CDR2 α – ISSIKDK (SEQ ID NO: 190),
 CDR3 α - CAASYNNAARLMF (SEQ ID NO: 191),
 CDR1 β – KGHSH (SEQ ID NO: 200),
 CDR2 β – LQKENI (SEQ ID NO: 201), and
 CDR3 β – CASSPTGGEYYGYTF (SEQ ID NO: 202),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxviii) CDR1 α - NSMFDY (SEQ ID NO: 189),
 CDR2 α – ISSIKDK (SEQ ID NO: 190),
 CDR3 α - CAASYNNAARLMF (SEQ ID NO: 191),
 CDR1 β – MNHEY (SEQ ID NO: 206),
 CDR2 β – SVGAGI (SEQ ID NO: 207), and

CDR3 β – CASSSYPLRTGRYNSYNSPLHF (SEQ ID NO: 208),
or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxix) CDR1 α - NSMFDY (SEQ ID NO: 212),
5 CDR2 α – ISSIKDK (SEQ ID NO: 213),
CDR3 α - CAASGGRDDKIIF (SEQ ID NO: 214),
CDR1 β – MNHEY (SEQ ID NO: 217),
CDR2 β – SVGAGI (SEQ ID NO: 218), and
CDR3 β – CASSYSRTESTDTQYF (SEQ ID NO: 219),

10 or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxx) CDR1 α - NSMFDY (SEQ ID NO: 90),
CDR2 α – ISSIKDK (SEQ ID NO: 91),
CDR3 α - CAANNARLMF (SEQ ID NO: 92),
15 CDR1 β – SGHRS (SEQ ID NO: 269),
CDR2 β – YFSETQ (SEQ ID NO: 270), and
CDR3 β – CASSPGQHGEFF (SEQ ID NO: 271),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

20 (xxxi) CDR1 α - NSMFDY (SEQ ID NO: 264),
CDR2 α – ISSIKDK (SEQ ID NO: 265),
CDR3 α - CAASATGNQFYF (SEQ ID NO: 266),
CDR1 β – SGHNS (SEQ ID NO: 95),
CDR2 β – FNNNVP (SEQ ID NO: 96), and
25 CDR3 β – CASSDTRAREQFF (SEQ ID NO: 97),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxxii) CDR1 α - NSMFDY (SEQ ID NO: 264),
CDR2 α – ISSIKDK (SEQ ID NO: 265),
30 CDR3 α - CAASATGNQFYF (SEQ ID NO: 266),
CDR1 β – SGHRS (SEQ ID NO: 269),
CDR2 β – YFSETQ (SEQ ID NO: 270), and
CDR3 β – CASSPGQHGEFF (SEQ ID NO: 271),

or variants thereof each having up to three amino acid substitutions, additions or deletions;

35

(xxxiii) CDR1 α - TSINN (SEQ ID NO: 275),
 CDR2 α – IRSNERE (SEQ ID NO: 276),
 CDR3 α - CATDGDSSYKLIF (SEQ ID NO: 277),
 CDR1 β – DFQATT (SEQ ID NO: 161),
 5 CDR2 β – SNEGSKA (SEQ ID NO: 162), and
 CDR3 β – CSARDSVSGNTIYF (SEQ ID NO: 163),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxxiv) CDR1 α - TSINN (SEQ ID NO: 275),
 10 CDR2 α – IRSNERE (SEQ ID NO: 276),
 CDR3 α - CATDGDSSYKLIF (SEQ ID NO: 277),
 CDR1 β – SQVTM (SEQ ID NO: 167),
 CDR2 β – ANQGSEA (SEQ ID NO: 168), and
 CDR3 β – CSVGGSGSYNEQFF (SEQ ID NO: 169),
 15 or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxxv) CDR1 α - TSINN (SEQ ID NO: 275),
 CDR2 α – IRSNERE (SEQ ID NO: 276),
 CDR3 α - CATDGDSSYKLIF (SEQ ID NO: 277),
 20 CDR1 β – DFQATT (SEQ ID NO: 280),
 CDR2 β – SNEGSKA (SEQ ID NO: 281), and
 CDR3 β – CSARDVLTGDYGYTF (SEQ ID NO: 282),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;

25 (xxxvi) CDR1 α - TSINN (SEQ ID NO: 275),
 CDR2 α – IRSNERE (SEQ ID NO: 276),
 CDR3 α - CATDGDSSYKLIF (SEQ ID NO: 277),
 CDR1 β – SGHDY (SEQ ID NO: 286),
 CDR2 β – FNNNVP (SEQ ID NO: 287), and
 30 CDR3 β – CASSLGLSISQETQYF (SEQ ID NO: 288),
 or variants thereof each having up to three amino acid substitutions, additions or deletions;

(xxxvii) CDR1 α - DSSSTY (SEQ ID NO: 101),
 CDR2 α – IFSNMDM (SEQ ID NO: 102),
 35 CDR3 α - CAERLNTDKLIF (SEQ ID NO: 103),

CDR1 β – SGHDY (SEQ ID NO: 286),
 CDR2 β – FNNNVP (SEQ ID NO: 287), and
 CDR3 β – CASSLGLSISQETQYF (SEQ ID NO: 288),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxxviii) CDR1 α - DSVNN (SEQ ID NO: 106),
 CDR2 α – IPSGT (SEQ ID NO: 107),
 CDR3 α - CAVEATDSWGKLQF (SEQ ID NO: 108),
 CDR1 β – SGHDY (SEQ ID NO: 286),
 CDR2 β – FNNNVP (SEQ ID NO: 287), and
 CDR3 β – CASSLGLSISQETQYF (SEQ ID NO: 288),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxxix) CDR1 α - DSASNY (SEQ ID NO: 111),
 CDR2 α – IRSNVGE (SEQ ID NO: 112),
 CDR3 α - CAVRTSYDKVIF (SEQ ID NO: 113),
 CDR1 β – SGHDY (SEQ ID NO: 286),
 CDR2 β – FNNNVP (SEQ ID NO: 287), and
 CDR3 β – CASSLGLSISQETQYF (SEQ ID NO: 288),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxxx) CDR1 α - DSSSTY (SEQ ID NO: 101),
 CDR2 α – IFSNMDM (SEQ ID NO: 102),
 CDR3 α - CAERLNTDKLIF (SEQ ID NO: 103),
 CDR1 β – DFQATT (SEQ ID NO: 280),
 CDR2 β – SNEGSKA (SEQ ID NO: 281), and
 CDR3 β – CSARDVLTGDYGYTF (SEQ ID NO: 282),
 or variants thereof each having up to three amino acid substitutions, additions or
 deletions;

(xxxxi) CDR1 α - DSVNN (SEQ ID NO: 106),
 CDR2 α – IPSGT (SEQ ID NO: 107),
 CDR3 α - CAVEATDSWGKLQF (SEQ ID NO: 108),
 CDR1 β – DFQATT (SEQ ID NO: 280),
 CDR2 β – SNEGSKA (SEQ ID NO: 281), and
 CDR3 β – CSARDVLTGDYGYTF (SEQ ID NO: 282),

or variants thereof each having up to three amino acid substitutions, additions or deletions; or

(xxxxii) CDR1 α - DSASNY (SEQ ID NO: 111),

CDR2 α - IRSNVGE (SEQ ID NO: 112),

5 CDR3 α - CAVRTSYDKVIF (SEQ ID NO: 113),

CDR1 β - DFQATT (SEQ ID NO: 280),

CDR2 β - SNEGSKA (SEQ ID NO: 281), and

CDR3 β - CSARDVLTGDYGYTF (SEQ ID NO: 282),

10 or variants thereof each having up to three amino acid substitutions, additions or deletions.

3. A TCR according to claim 1 or 2 comprising:

(i) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 4 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 9 or a variant thereof having at least 75% sequence identity thereto;

(ii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 37 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 42 or a variant thereof having at least 75% sequence identity thereto;

20 (iii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 15 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 25 or a variant thereof having at least 75% sequence identity thereto;

25 (iv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 15 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 31 or a variant thereof having at least 75% sequence identity thereto;

30 (v) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 20 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 25 or a variant thereof having at least 75% sequence identity thereto;

- (vi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 20 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 31 or variants thereof having at least 75% sequence identity thereto;
- 5 (vii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 48 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 58 or a variant thereof having at least 75% sequence identity thereto;
- 10 (viii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 48 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 64 or a variant thereof having at least 75% sequence identity thereto;
- 15 (ix) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 48 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 70 or a variant thereof having at least 75% sequence identity thereto;
- 20 (x) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 53 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 58 or a variant thereof having at least 75% sequence identity thereto;
- (xi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 53 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 64 or a variant thereof having at least 75% sequence identity thereto;
- 25 (xii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 53 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 70 or a variant thereof having at least 75% sequence identity thereto;
- 30 (xiii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 76 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 81 or a variant thereof having at least 75% sequence identity thereto;

- (xiv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 76 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 87 or a variant thereof having at least 75% sequence identity thereto;
- 5 (xv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 93 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 98 or a variant thereof having at least 75% sequence identity thereto;
- 10 (xvi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 104 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 164 or a variant thereof having at least 75% sequence identity thereto;
- 15 (xvii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 104 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 170 or a variant thereof having at least 75% sequence identity thereto;
- 20 (xviii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 109 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 164 or a variant thereof having at least 75% sequence identity thereto;
- (xix) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 109 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 170 or a variant thereof having at least 75% sequence identity thereto;
- 25 (xx) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 114 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 164 or a variant thereof having at least 75% sequence identity thereto;
- 30 (xxi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 114 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 170 or a variant thereof having at least 75% sequence identity thereto;

- (xxii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 176 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 181 or a variant thereof having at least 75% sequence identity thereto;
- 5 (xxiii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 187 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 197 or a variant thereof having at least 75% sequence identity thereto;
- 10 (xxiv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 187 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 203 or a variant thereof having at least 75% sequence identity thereto;
- 15 (xxv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 187 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 209 or a variant thereof having at least 75% sequence identity thereto;
- 20 (xxvi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 192 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 197 or a variant thereof having at least 75% sequence identity thereto;
- (xxvii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 192 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 203 or a variant thereof having at least 75% sequence identity thereto;
- 25 (xxviii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 192 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 209 or a variant thereof having at least 75% sequence identity thereto;
- 30 (xxix) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 215 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 220 or a variant thereof having at least 75% sequence identity thereto;

- (xxx) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 93 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 272 or a variant thereof having at least 75% sequence identity thereto;
- 5 (xxxi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 267 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 98 or a variant thereof having at least 75% sequence identity thereto;
- 10 (xxxii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 267 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 272 or a variant thereof having at least 75% sequence identity thereto;
- 15 (xxxiii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 278 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 164 or a variant thereof having at least 75% sequence identity thereto;
- 20 (xxxiv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 278 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 170 or a variant thereof having at least 75% sequence identity thereto;
- (xxxv) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 278 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 283 or a variant thereof having at least 75% sequence identity thereto;
- 25 (xxxvi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 278 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 289 or a variant thereof having at least 75% sequence identity thereto;
- 30 (xxxvii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 104 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 289 or a variant thereof having at least 75% sequence identity thereto;

- (xxxviii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 109 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 289 or a variant thereof having at least 75% sequence identity thereto;
- 5 (xxxix) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 114 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 289 or a variant thereof having at least 75% sequence identity thereto;
- 10 (xxxx) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 104 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 283 or a variant thereof having at least 75% sequence identity thereto;
- 15 (xxxxi) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 109 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 283 or a variant thereof having at least 75% sequence identity thereto; or
- 20 (xxxixii) an α chain variable domain comprising the amino acid sequence of SEQ ID NO: 114 or a variant thereof having at least 75% sequence identity thereto; and a β chain variable domain comprising the amino acid sequence of SEQ ID NO: 283 or a variant thereof having at least 75% sequence identity thereto.
4. A TCR according to any preceding claim comprising:
- (i) an α chain comprising the amino acid sequence of SEQ ID NO: 5 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 10, SEQ ID NO: 11 and variants of SEQ ID NOs: 10 and 11 having at least 75% sequence identity thereto;
- 25 (ii) an α chain comprising the amino acid sequence of SEQ ID NO: 38 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 43, SEQ ID NO: 44 and variants of SEQ ID NOs: 43 and 44 having at least 75% sequence identity thereto;
- 30

- (iii) an α chain comprising the amino acid sequence of SEQ ID NO: 16 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 26, SEQ ID NO: 27 and variants of SEQ ID NOs: 26 and 27 having at least 75% sequence identity thereto;
- (iv) an α chain comprising the amino acid sequence of SEQ ID NO: 16 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 32, SEQ ID NO: 33 and variants of SEQ ID NOs: 32 and 33 having at least 75% sequence identity thereto;
- (v) an α chain comprising the amino acid sequence of SEQ ID NO: 21 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 26, SEQ ID NO: 27 and variants of SEQ ID NOs: 26 and 27 having at least 75% sequence identity thereto;
- (vi) an α chain comprising the amino acid sequence of SEQ ID NO: 21 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 32, SEQ ID NO: 33 and variants of SEQ ID NOs: 32 and 33 having at least 75% sequence identity thereto;
- (vii) an α chain comprising the amino acid sequence of SEQ ID NO: 49 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 59, SEQ ID NO: 60 and variants of SEQ ID NOs: 59 and 60 having at least 75% sequence identity thereto;
- (viii) an α chain comprising the amino acid sequence of SEQ ID NO: 49 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 65, SEQ ID NO: 66 and variants of SEQ ID NOs: 65 and 66 having at least 75% sequence identity thereto;
- (ix) an α chain comprising the amino acid sequence of SEQ ID NO: 49 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 71, SEQ ID

NO: 72 and variants of SEQ ID NOs: 71 and 72 having at least 75% sequence identity thereto;

(x) an α chain comprising the amino acid sequence of SEQ ID NO: 54 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 59, SEQ ID NO: 60 and variants of SEQ ID NOs: 59 and 60 having at least 75% sequence identity thereto;

(xi) an α chain comprising the amino acid sequence of SEQ ID NO: 54 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 65, SEQ ID NO: 66 and variants of SEQ ID NOs: 65 and 66 having at least 75% sequence identity thereto;

(xii) an α chain comprising the amino acid sequence of SEQ ID NO: 54 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 71, SEQ ID NO: 72 and variants of SEQ ID NOs: 71 and 72 having at least 75% sequence identity thereto;

(xiii) an α chain comprising the amino acid sequence of SEQ ID NO: 77 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 82, SEQ ID NO: 83 and variants of SEQ ID NOs: 82 and 83 having at least 75% sequence identity thereto;

(xiv) an α chain comprising the amino acid sequence of SEQ ID NO: 77 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 88, SEQ ID NO: 89 and variants of SEQ ID NOs: 88 and 89 having at least 75% sequence identity thereto;

(xv) an α chain comprising the amino acid sequence of SEQ ID NO: 94 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 99, SEQ ID NO: 100 and variants of SEQ ID NO: 99, SEQ ID NO: 100 having at least 75% sequence identity thereto;

- (xvi) an α chain comprising the amino acid sequence of SEQ ID NO: 105 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 165, SEQ ID NO: 166 and variants of SEQ ID NOs: 165 and 166 having at least 75% sequence identity thereto;
- (xvii) an α chain comprising the amino acid sequence of SEQ ID NO: 105 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 171, SEQ ID NO: 172 and variants of SEQ ID NOs: 171 and 172 having at least 75% sequence identity thereto;
- (xviii) an α chain comprising the amino acid sequence of SEQ ID NO: 110 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 165, SEQ ID NO: 166 and variants of SEQ ID NOs: 165 and 166 having at least 75% sequence identity thereto;
- (xix) an α chain comprising the amino acid sequence of SEQ ID NO: 110 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 171, SEQ ID NO: 172 and variants of SEQ ID NOs: 171 and 172 having at least 75% sequence identity thereto;
- (xx) an α chain comprising the amino acid sequence of SEQ ID NO: 160 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 165, SEQ ID NO: 166 and variants of SEQ ID NOs: 165 and 166 having at least 75% sequence identity thereto;
- (xxi) an α chain comprising the amino acid sequence of SEQ ID NO: 160 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 171, SEQ ID NO: 172 and variants of SEQ ID NOs: 171 and 172 having at least 75% sequence identity thereto;
- (xxii) an α chain comprising the amino acid sequence of SEQ ID NO: 177 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 182, SEQ ID

NO: 183 and variants of SEQ ID NOs: 182 and 183 having at least 75% sequence identity thereto;

(xxiii) an α chain comprising the amino acid sequence of SEQ ID NO: 188 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 198, SEQ ID NO: 199 and variants of SEQ ID NOs: 198 and 199 having at least 75% sequence identity thereto;

(xxiv) an α chain comprising the amino acid sequence of SEQ ID NO: 188 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 204, SEQ ID NO: 205 and variants of SEQ ID NOs: 204 and 205 having at least 75% sequence identity thereto;

(xxv) an α chain comprising the amino acid sequence of SEQ ID NO: 188 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 210, SEQ ID NO: 211 and variants of SEQ ID NOs: 210 and 211 having at least 75% sequence identity thereto;

(xxvi) an α chain comprising the amino acid sequence of SEQ ID NO: 193 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 198, SEQ ID NO: 199 and variants of SEQ ID NOs: 198 and 199 having at least 75% sequence identity thereto;

(xxvii) an α chain comprising the amino acid sequence of SEQ ID NO: 193 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 204, SEQ ID NO: 205 and variants of SEQ ID NOs: 204 and 205 having at least 75% sequence identity thereto;

(xxviii) an α chain comprising the amino acid sequence of SEQ ID NO: 193 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 210, SEQ ID NO: 211 and variants of SEQ ID NOs: 210 and 211 having at least 75% sequence identity thereto;

- (xxix) an α chain comprising the amino acid sequence of SEQ ID NO: 216 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 221, SEQ ID NO: 222 and variants of SEQ ID NOs: 221 and 222 having at least 75% sequence identity thereto;
- (xxx) an α chain comprising the amino acid sequence of SEQ ID NO: 94 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 273, SEQ ID NO: 274 and variants of SEQ ID NOs: 273 and 274 having at least 75% sequence identity thereto;
- (xxxi) an α chain comprising the amino acid sequence of SEQ ID NO: 268 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 99, SEQ ID NO: 100 and variants of SEQ ID NOs: 99 and 100 having at least 75% sequence identity thereto;
- (xxxii) an α chain comprising the amino acid sequence of SEQ ID NO: 268 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 273, SEQ ID NO: 274 and variants of SEQ ID NOs: 273 and 274 having at least 75% sequence identity thereto;
- (xxxiii) an α chain comprising the amino acid sequence of SEQ ID NO: 279 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 165, SEQ ID NO: 166 and variants of SEQ ID NOs: 165 and 166 having at least 75% sequence identity thereto;
- (xxxiv) an α chain comprising the amino acid sequence of SEQ ID NO: 279 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 171, SEQ ID NO: 172 and variants of SEQ ID NOs: 171 and 172 having at least 75% sequence identity thereto;
- (xxxv) an α chain comprising the amino acid sequence of SEQ ID NO: 279 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 284, SEQ ID

NO: 285 and variants of SEQ ID NOs: 284 and 285 having at least 75% sequence identity thereto;

(xxxvi) an α chain comprising the amino acid sequence of SEQ ID NO: 279 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 290, SEQ ID NO: 291 and variants of SEQ ID NOs: 290 and 291 having at least 75% sequence identity thereto;

(xxxvii) an α chain comprising the amino acid sequence of SEQ ID NO: 105 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 290, SEQ ID NO: 291 and variants of SEQ ID NOs: 290 and 291 having at least 75% sequence identity thereto;

(xxxviii) an α chain comprising the amino acid sequence of SEQ ID NO: 110 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 290, SEQ ID NO: 291 and variants of SEQ ID NOs: 290 and 291 having at least 75% sequence identity thereto;

(xxxix) an α chain comprising the amino acid sequence of SEQ ID NO: 160 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 290, SEQ ID NO: 291 and variants of SEQ ID NOs: 290 and 291 having at least 75% sequence identity thereto;

(xxxx) an α chain comprising the amino acid sequence of SEQ ID NO: 105 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 284, SEQ ID NO: 285 and variants of SEQ ID NOs: 284 and 285 having at least 75% sequence identity thereto;

(xxxxi) an α chain comprising the amino acid sequence of SEQ ID NO: 110 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 284, SEQ ID NO: 285 and variants of SEQ ID NOs: 284 and 285 having at least 75% sequence identity thereto; or

(xxxii) an α chain comprising the amino acid sequence of SEQ ID NO: 160 or a variant thereof having at least 75% sequence identity thereto; and a β chain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 284, SEQ ID NO: 285 and variants of SEQ ID NOs: 284 and 285 having at least 75% sequence identity thereto.

5
10
15
5. A T-cell receptor (TCR), which binds to a Wilms tumour 1 protein (WT1) peptide when presented by a major histocompatibility complex (MHC), wherein the WT1 peptide comprises an amino acid sequence selected from the group consisting of EPASQHTLRSG (SEQ ID NO: 123), YESDNHTTPIL (SEQ ID NO: 126), NHTTPILCGAQYRIH (SEQ ID NO: 127), QCLSAFTVHFSGQFT (SEQ ID NO: 118), EDPMGQQGSLGEQQY (SEQ ID NO: 119), SQLECMTWNQMNLGA (SEQ ID NO: 120), APVLDFAAPPGA (SEQ ID NO: 117), NQMNLGATLKG (SEQ ID NO: 250), DPGGIWAKLGAAEAS (SEQ ID NO: 251), NHTTPILCGAQYRIH (SEQ ID NO: 252), KRHQRRHTGVKPFQC (SEQ ID NO: 253) PSCQKKFARSDELVR (SEQ ID NO: 254) and variants thereof each having up to three amino acid substitutions, additions or deletions.

6. A TCR according to any preceding claim which binds to an MHC I and/or MHC II peptide complex.

7. A TCR according to any preceding claim, which is restricted to a human leukocyte antigen (HLA) allele, preferably wherein the TCR is restricted to a HLA-A or a HLA-B allele, more preferably wherein the TCR is restricted to a HLA-A allele selected from the group consisting of HLA-A*0201, HLA-A*0101, HLA-A*2402 and HLA-A*0301 or wherein the TCR is restricted to a HLA-B allele selected from the group consisting of HLA-B*0702, HLA-B*3501 and HLA-B*3502.

8. A TCR according to:

25 a. (i), (ii), (xv), (xvi), (xxix), (xxx)-(xxxiii), (xxxv) or (xxxx) of any one of claims 1-4, which is restricted to HLA-A*0201;

b. (vii) of any one of claims 1-4, which is restricted to HLA-B*3502; or

c. any of (xiii)-(xiv) of any one of claims 1-4, which is restricted to HLA-B*3501.

9. A TCR according to any preceding claim comprising one or more mutations at the α chain/ β chain interface, such that when the α chain and the β chain are expressed in a T-cell, the frequency of mispairing between said chains and endogenous TCR α and β chains is reduced.

- 10.** A TCR according to claim 9, wherein the one or more mutations introduce a cysteine residue into the constant region domain of each of the α chain and the β chain, wherein the cysteine residues are capable of forming a disulphide bond between the α chain and the β chain.
- 5 **11.** A TCR according to any preceding claim, which comprises a murinized constant region.
- 12.** A TCR according to any preceding claim, wherein the TCR is a soluble TCR.
- 13.** An isolated polynucleotide encoding the α chain of a T-cell receptor (TCR) according to any preceding claim, and/or the β chain of a TCR according to any preceding claim.
- 10 **14.** An isolated polynucleotide according to claim 13, wherein the polynucleotide encodes the α chain linked to the β chain.
- 15.** An isolated polynucleotide according to claim 13 or 14, which encodes one or more short interfering RNA (siRNA) or other agents capable of reducing or preventing expression of one or more endogenous TCR gene.
- 15 **16.** A vector comprising a polynucleotide according to any one of claims 13-15.
- 17.** A vector according to claim 16 comprising a polynucleotide, which encodes one or more CD3 chains, CD8, a suicide gene, and/or a selectable marker.
- 18.** A cell comprising a TCR according to any one of claims 1-11, a polynucleotide according to any one of claims 13-15 or a vector according to claim 16 or 17,
- 20 optionally wherein the cell further comprises a vector which encodes one or more CD3 chains, CD8, a suicide gene and/or a selectable marker.
- 19.** A cell according to claim 18, wherein the cell is a T-cell, a lymphocyte, or a stem cell, optionally wherein the T-cell, the lymphocyte, or the stem cell is selected from the group consisting of CD4 cells, CD8 cells, naive T-cells, memory stem T-cells, central memory T-
- 25 cells, double negative T-cells, effector memory T-cells, effector T-cells, Th0 cells, Tc0 cells, Th1 cells, Tc1 cells, Th2 cells, Tc2 cells, Th17 cells, Th22 cells, gamma/delta T-cells, natural killer (NK) cells, natural killer T (NKT) cells, hematopoietic stem cells and pluripotent stem cells.
- 20.** A cell according to claim 19, wherein the cell is a T-cell which has been isolated from
- 30 a subject.

21. A cell according to any one of claims 18-20, wherein an endogenous gene encoding a TCR α chain and/or an endogenous gene encoding a TCR β chain is disrupted, preferably such that the endogenous gene encoding a TCR α chain and/or the endogenous gene encoding a TCR β chain is not expressed,

5 optionally wherein the endogenous gene encoding a TCR α chain and/or the endogenous gene encoding a TCR β chain is disrupted by insertion of an expression cassette comprising a polynucleotide sequence encoding a TCR of any one of claims 1-11,

further optionally wherein one or more endogenous genes encoding an MHC is disrupted,

10 further optionally wherein an endogenous gene involved in persistence, expansion, activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other T-cell functions is disrupted, preferably wherein the endogenous gene involved in persistence, expansion, activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other T-cell functions is selected from the group consisting of *PD1*, *TIM3*, *LAG3*, *2B4*, *KLRG1*, *TGFbR*, *CD160* and *CTLA4*.

15 **22.** A method of preparing a cell, which comprises the step of introducing a vector according to claim 16 and/or 17 into a cell *in vitro*, *ex vivo* or *in vivo*, e.g. by transfection or transduction.

23. A method of preparing a cell according to claim 22, which comprises the step of T-cell editing, which comprises disrupting an endogenous gene encoding a TCR α chain and/or an endogenous gene encoding a TCR β chain with an artificial nuclease, preferably
20 wherein the artificial nuclease is selected from the group consisting of zinc finger nucleases (ZFN), transcription activator-like effector nucleases (TALEN) and CRISPR/Cas systems.

24. A method of preparing a cell according to claim 23, which comprises the step of targeted integration of an expression cassette into the endogenous gene encoding the TCR
25 α chain and/or the endogenous gene encoding the TCR β chain disrupted by the artificial nuclease, wherein the expression cassette comprises a polynucleotide sequence encoding a TCR of any one of claims 1-11.

25. A method of preparing a cell according to any one of claims 22-24, which comprises the step of disrupting one or more endogenous genes encoding an MHC, preferably wherein
30 the cell prepared by the method is a non-alloreactive universal T-cell.

26. A method of preparing a cell according to any one of claims 22-25, which comprises the step of disrupting one or more endogenous genes to modify the persistence, expansion,

activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other T-cell functions, preferably wherein the method comprises the step of targeted integration of an expression cassette into an endogenous gene involved in persistence, expansion, activity, resistance to exhaustion/senescence/inhibitory signals, homing capacity, or other T-cell functions disrupted by an artificial nuclease, wherein the expression cassette comprises a polynucleotide sequence encoding a TCR of any one of claims 1-11, preferably wherein the endogenous gene is selected from the group consisting of *PD1*, *TIM3*, *LAG3*, *2B4*, *KLRG1*, *TGFbR*, *CD160* and *CTLA4*.

27. A cell according to any one of claims 18-21 or a cell prepared by the method of any one of claims 22-26 for use in adoptive cell transfer, preferably adoptive T-cell transfer, optionally wherein the adoptive T-cell transfer is allogenic adoptive T-cell transfer, autologous adoptive T-cell transfer, or universal non-alloreactive adoptive T-cell transfer.

28. A chimeric molecule comprising a TCR of any one of claims 1-11, or a portion thereof, conjugated to a non-cellular substrate, a toxin and/or an antibody, optionally wherein the non-cellular substrate is selected from the group consisting of nanoparticles, exosomes and other non-cellular substrates.

29. A TCR according to any one of claims 1-11, an isolated polynucleotide according to any one of claims 13-15, a vector according to claim 16 or 17, a cell according to any one of claims 18-21, a cell prepared by the method of any one of claims 22-26, or a chimeric molecule according to claim 28 for use in therapy.

30. A TCR according to any one of claims 1-11, an isolated polynucleotide according to any one of claims 13-15, a vector according to claim 16 or 17, a cell according to any one of claims 18-21, a cell prepared by the method of any one of claims 22-26, or a chimeric molecule according to claim 28 for use in treating and/or preventing a disease associated with expression of WT1, optionally wherein the disease associated with expression of WT1 is a proliferative disorder, preferably wherein the proliferative disorder is a hematological malignancy or a solid tumor, preferably wherein the hematological malignancy is selected from the group consisting of acute myeloid leukemia (AML), chronic myeloid leukemia (CML), lymphoblastic leukemia, myelodysplastic syndromes, lymphoma, multiple myeloma, non Hodgkin lymphoma, and Hodgkin lymphoma; or preferably wherein the solid tumor is selected from the group consisting of lung cancer, breast cancer, oesophageal cancer, gastric cancer, colon cancer, cholangiocarcinoma, pancreatic cancer, ovarian cancer, head and neck cancers, synovial sarcoma, angiosarcoma, osteosarcoma, thyroid cancer, endometrial cancer, neuroblastoma, rhabdomyosarcoma, liver cancer, melanoma, prostate

cancer, renal cancer, soft tissue sarcoma, urothelial cancer, biliary cancer, glioblastoma, mesothelioma, cervical cancer, and colorectal cancer.

31. A method for treating and/or preventing a disease associated with expression of WT1, which comprises the step of administering a TCR according to any one of claims 1-11,
 5 an isolated polynucleotide according to any one of claims 13-15, a vector according to claim 16 or 17, a cell according to any one of claims 18-21, a cell prepared by the method of any one of claims 22-26, or a chimeric molecule according to claim 28 to a subject in need thereof.

32. The method of claim 31, wherein the disease associated with expression of WT1 is a
 10 proliferative disorder, preferably wherein the proliferative disorder is a hematological malignancy or a solid tumor, preferably wherein the hematological malignancy is selected from the group consisting of acute myeloid leukemia (AML), chronic myeloid leukemia (CML), lymphoblastic leukemia, myelodysplastic syndromes, lymphoma, multiple myeloma, non Hodgkin lymphoma, and Hodgkin lymphoma; or preferably wherein the solid tumor is
 15 selected from the group consisting of lung cancer, breast cancer, oesophageal cancer, gastric cancer, colon cancer, cholangiocarcinoma, pancreatic cancer, ovarian cancer, head and neck cancers, synovial sarcoma, angiosarcoma, osteosarcoma, thyroid cancer, endometrial cancer, neuroblastoma, rhabdomyosarcoma, liver cancer, melanoma, prostate cancer, renal cancer, soft tissue sarcoma, urothelial cancer, biliary cancer, glioblastoma,
 20 cervical cancer, mesothelioma and colorectal cancer.


33. An isolated immunogenic WT1 peptide comprising an amino acid sequence selected from the group consisting of EPASQHTLRSG (SEQ ID NO: 123) YESDNHTTPIL (SEQ ID NO: 126), NHTTPILCGAQYRIH (SEQ ID NO: 127), QCLSAFTVHFSGQFT (SEQ ID NO: 118), EDPMGQQGSLGEQQY (SEQ ID NO: 119), SQLECMTWNQMNLGA (SEQ ID NO: 120), APVLDFAPPGA (SEQ ID NO: 117), NQMNLGATLKG (SEQ ID NO: 250), DPGGIWAKLGAAEAS (SEQ ID NO: 251), NHTTPILCGAQYRIH (SEQ ID NO: 252), KRHQRRHTGVKPFQC (SEQ ID NO: 253), PSCQKKFARSDELVR (SEQ ID NO: 254) and variants thereof each having up to three amino acid substitutions, additions or deletions.

Figure 1


a

S2

Unrelated pool




WT1




S6

Unrelated pool




WT1




b

S2

Unrelated pool




WT1




S7

Unrelated pool




WT1




c

S3

Unrelated pool




WT1




S4

Unrelated pool




WT1




d

S3

Unrelated pool




WT1




S4

Unrelated pool




WT1



e


S3




f

S3

Unrelated pool




WT1




S4

Unrelated pool




WT1




89

S3

Unrelated pool




WT1




S4

Unrelated pool




WT1




h

S4




S5




i

S4

Unrelated pool




WT1




S5

Unrelated pool




WT1




j

S4










S5






a
Figure 2


	Donor #1		Donor #2		Donor #3		Donor #4		Donor #5		Donor #6		Donor #7		Donor #8		Donor #9		Donor #10	
	IFN γ	CD107a	IFN γ	CD107a	IFN γ	CD107a	IFN γ	CD107a	IFN γ	CD107a	IFN γ	CD107a	IFN γ	CD107a	IFN γ	CD107a	IFN γ	CD107a	IFN γ	CD107a
SP1	19.3	8.1	1.4	ND	0.14	1.32	1.68	4.95	0.26	2.55	0.69	2.62	1.11	5.36	5.8	4.28	21.5	28.4	0.20	1.00
SP2	9.5	10.9	0.81	ND	0.2	0.98	1.73	4.95	0.37	3.06	0.83	2.39	1.25	5.31	6.24	6.00	21.3	26.2	0.10	0.30
SP3	7.04	9.17	1.48	ND	0.51	2.91	4.07	10.8	0.42	2.76	0.64	2.49	0.98	5.38	9.16	7.49	23.3	30.1	0.07	0.21
SP4	63.8	83.6	2.01	ND	13.4	22.9	6.94	12.2	0.45	2.29	47.5	65.3	38.4	50.3	6.04	4.67	24.2	29.6	20.4	49.4
SP5	65.5	81.5	1.24	ND	17.2	22.1	10.4	47.3	54.8	59.4	58.2	64.8	42.9	52.8	6.55	5.16	27.8	36.3	36.1	51.2
SP6	9.23	49.6	5.71	ND	0	1.23	11.6	49.7	0.62	6.55	0.91	2.68	1.22	7.55	6.37	5.84	22.3	32.4	0	0.94
SP7	9.81	12.1	2.39	ND	0.12	2.11	1.79	5.23	0.56	3.96	0.45	2.90	13.6	17.7	5.15	4.78	19.9	26.5	0	0.09
SP8	14.9	24.3	1.56	ND	0.25	3.08	1.45	4.57	0.29	2.77	0.78	1.78	13.6	16.9	7.19	5.81	18.2	26.3	0.12	0.84
SP9	9.91	61.9	0.84	ND	0.21	1.13	2.3	5.89	0.59	3.66	0.74	4.42	2.73	7.34	7.35	5.79	20.5	27.4	0.29	0.58
SP10	6.97	9.83	0.74	ND	0.73	1.91	1.63	4.54	0.44	2.53	0.55	2.36	1.27	4.26	6.17	5.89	18.3	30.1	0	0.20
SP11	10.9	9.54	1.91	ND	0.73	2.2	1.81	4.46	4.26	6.53	0.83	2.04	2.11	5.34	6.39	4.76	21.0	28.1	0.09	0.19
SP12	9.75	8.69	2.02	ND	0.3	1.79	2.13	5.32	3.58	6.48	0.54	2.25	1.54	5.85	16.70	15.80	22.8	28.0	0	0.92
SP13	5.98	8.54	1.44	ND	0	1.14	1.98	4.74	0.49	2.94	0.37	1.80	1.17	6.72	5.93	4.61	28.0	31.3	0.08	0.26
SP14	6.14	11.9	1.66	ND	0	1.17	12.3	54.6	0.34	2.95	0.48	2.57	1.96	7.52	14.60	13.30	25.4	31.1	0.43	0.69
SP15	12.8	58	0.9	ND	1.37	2.73	2.45	4.98	0.68	3.15	1.23	7.19	3.67	11.7	9.15	7.32	21.0	29.6	0.08	0.70
SP16	78.8	83.6	5.42	ND	19.9	23	1.73	4.66	0.32	2.69	56.5	65.9	48.2	54.6	7.30	5.80	25.7	31.5	27.7	56.7
SP17	8.08	8.55	4.64	ND	0.29	1.15	1.99	4.47	0.23	2.15	0.67	2.31	1.36	5.04	6.30	5.22	20.7	28.9	0	0.37
SP18	9.67	6.93	0.84	ND	0.37	2.01	4.24	8.48	0.61	2.23	0.83	1.79	1.20	5.26	7.57	4.95	14.8	22.2	0	0.57
SP19	9.6	7.26	1.14	ND	0	1.12	2.04	4.23	0.50	2.78	0.60	2.00	1.49	5.18	6.69	6.25	21.1	26.0	0	0.42
SP20	9.31	45.9	7.44	ND	0.85	2.67	1.92	4.18	0.42	2.95	0.33	3.02	2.00	13.00	7.17	4.86	22.9	28.3	0.44	0.89
SP21	9.97	6.84	0.85	ND	0.25	0.99	4.53	7.16	47.1	58.7	0.49	2.04	0.53	6.41	7.54	6.72	32.2	41.5	0	0.31
SP22	10.3	7.49	2.67	ND	0	0.78	1.6	3.62	2.34	5.26	0.50	2.82	0.81	5.86	9.62	5.97	20.4	25.5	0.19	0.45
SP23	8.25	5.18	4.13	ND	0.12	1.06	1.7	4.12	0.25	2.51	0.58	2.00	1.07	5.26	7.22	6.12	17.9	25.3	0.21	0.32
SP24	8.22	8.5	1.27	ND	0	1.72	1.98	4.37	0.50	2.54	0.73	2.83	0.66	5.42	7.37	5.70	15.1	25.7	0	0.17











b**Subpool 4****Subpool 5****Subpool 16**

C

d**Subpool 4****Subpool 5****Subpool 16**



e





f**Subpool 5****Subpool 11****Subpool 12****Subpool 21****Subpool 22**

69



Subpool 4



Subpool 5




Subpool 16




h


Subpool 4




Subpool 5




Subpool 7




Subpool 8




Subpool 16



Subpool 20




i




j


Subpool 5



Subpool 13



Subpool 21









k


Figure 3**a****Peptide 40****Peptide 41****Peptide 42****Peptide 43**

b


Peptide 42




Peptide 54




Peptide 77














Peptide 90




Peptide 138



c


d**Peptide 15****Peptide 16****Peptide 17****Peptide 18****Peptide 132**

e




f


Peptide 63




CD107a




Peptide 101




CD107a




Peptide 107




CD107a




Peptide 108




CD107a




Peptide 113




CD107a




Peptide 119




CD107a




Peptide 120




CD107a




89



h



i





j



Peptide 17

Unrelated Peptide



EBV-HLA*A24:02





EBV-HLA*A24:02



EBV-HLA*A68:01





EBV-HLA*A68:01





EBV-HLA*B35:02

HLA*C04:02





EBV-HLA*B35:02

HLA*C04:02





EBV-HLA*B40:01

HLA*C03:04





EBV-HLA*B40:01



HLA*C03:04





EBV-HLA*C03:04





EBV-HLA*C03:04














EBV-HLA*C04:01



EBV-HLA*C04:01





k**Peptide 101****Unrelated Peptide****EBV-HLA *A11:01****EBV-HLA *A11:01****EBV-HLA *B50:01**
EBV-HLA *C6:02**EBV-HLA *B50:01**
EBV-HLA *C6:02**EBV-HLA *B35:01**
HLA *C04:01**EBV-HLA *B35:01**
HLA *C04:01**EBV-HLA *C04:01****EBV-HLA *C04:01****EBV-HLA *C06:02****EBV-HLA *C06:02**




m

Donor	Peptide	Sequence
HD1	40	AAQWAPVLDFAPPGA
	41	APVLDFAPPGASAYG
HD2	54	QCLSAFTVHFSGQFT
	77	EDPMGQQGSLGEQQY
	90	SQLECMTWNQMNLGA
HD3	40	AAQWAPVLDFAPPGA
	41	APVLDFAPPGASAYG
	Specific Nonamer	VLDFAPPGA
HD4	17	TCVPEPASQHTLRSG
	18	EPASQHTLRSGPGCL
	99	HSTGYESDNHTTPIL
	100	YESDNHTTPILCGAQ
HD5	101	NHTTPILCGAQYRIH
HD6	Specific Nonamer	VLDFAPPGA
HD7	40	AAQWAPVLDFAPPGA
	41	APVLDFAPPGASAYG
	91	CMTWNQMNLGATLKG
	92	NQMNLGATLKGVAAG
HD8	24	DPGGIWAKLGAAEAS
HD9	101	NHTTPILCGAQYRIH
	125	KRHQRRHTGVKPFQC
	137	PSCQKKFARSDELVR
HD10	Specific Nonamer	VLDFAPPGA


Figure 4**b**

C

WT1-specific T cells+
AML HLA-B35*02 blasts



WT1-specific T cells+
AML unrelated blasts



E:T=10:1

Figure 5







Figure 6**Dominant clonotypes:** **α -chain****TRAV30*01-TRAJ20*01****CDR3: CGTAWINDYKLSF** **β -chain****TRBV12-4*01- TRBD2*01 -TRBJ1-5*01****CDR3: CASRKTGGYSNQPQHF**

b**Dominant clonotypes:****α-chain****TRAV8-4*01-TRAJ22*01****CDR3: CAVRLSGSARQLTF****TRAV38-2/DV8*01-TRAJ47*01****CDR3: CAYRSLKYGNKLVF****β-chain****TRBV11-2*01- TRBD2*02-TRBJ2-7*01****CDR3: CASSLLGDEQYF****TRBV7-2- TRBD1*01- TRBJ2-7*01****CDR3: CASSLVALQGAGEQYF**

C

Dominant clonotypes:

α -chain


TRAV12-1*01-TRAJ23*01

CDR3: CVVNLLSNQGGLIF

β -chain

TRBV4-3*01- TRBD2- TRBJ1-4*01

CDR3: CASSQDYLVSNKLFF

d

Dominant clonotypes:

α-chain (peptide 17)

TRAV17*01-TRAJ29*01

CDR3: CATDAYSGNTPLVF

α-chain (peptide 99-100)

TRAV21*01-TRAJ23*01

CDR3: CAVRAEIYNQGGLIF

β-chain (peptide 17)

TRBV6-1*01- TRBD1*01-TRBJ1-1*01

CDR3: CASRAAGLDTEAFF


β-chain (peptide 99-100)


TRBV6-6*01- TRBD1/TRBD2- TRBJ2-7*01


CDR3: CASTQTPYEQYF

TRBV12-3*01- TRBD2*02- TRBJ1-2*01

CDR3: CASSTVGGEDYGYTF

e**Dominant clonotypes:****α-chain****TRAV13-1*02-TRAJ28*01****CDR3: CAASMAGAGSYQLTF****β-chain****TRBV10-3*01- TRBD1*01-TRBJ2-7*01****CDR3: CAISVGQGALYEQYF****TRBV9*01- TRBD1*01- TRBJ1-2*01****CDR3: CASSVARDRRNYGYTF**

f**Dominant clonotypes:** **α -chain****TRAV29/DV5*01-TRA31*01****CDR3: CAANNARLMF****TRAV29/DV5*01-TRAJ49*01****CDR3: CAASATGNQFYF** **β -chain****TRBV12-3*01 - TRBD1*01 -TRBJ2-1*01****CDR3: CASSDTRAREQFF****TRBV5-1*01 - TRBD1*01 -TRBJ2-2*01****CDR3: CASSPGQHGELEFF**



Dominant clonotypes:

α-chain (overlapping portion of peptide 40 and 41)
TRAV5*01-TRAJ34*01
CDR3: CAERLNTDKLIF

TRAV17*01 -TRAJ12*01
CDR3: CATDGDSSYKLIF

α-chain (overlapping portion of peptide 91 and 92)
TRAV22*01-TRAJ24*02
CDR3: CAVEATDSWGKLQF


TRAV13-1*02 -TRAJ50*01
CDR3: CAVRTSYDKVIF


β-chain (overlapping portion of peptide 40 and 41)
TRBV20-1*01 - TRBD1*01- TRBJ1-3*01
CDR3: CSARDSVSGNTIYF

TRBV20-1*01 - TRBD1*01- TRBJ1-2*01
CDR3: CSARDVLTGDYGYTF

β-chain (overlapping portion of peptide 91 and 92)
TRBV29-1*01 - TRBD2*02 - TRBJ2-1*01
CDR3: CSVGGSGSYNEQFF

TRBV12-4*01 - TRBD2- TRBJ2-5*01
CDR3: CASSLGLSISQETQYF

h**Dominant clonotypes:** **α -chain****TRAV41*01-TRAJ47*01****CDR3: CAVTVGNKLVF** **β -chain****TRBV6-1*01- TRBD1*01-TRBJ2-1*01****CDR3: CASRGWREQFF**



Dominant clonotypes:

α -chain (peptide 101)

TRAV41*01-TRAJ34*01

CDR3: CAARSYNTDKLIF

α -chain (peptide 125)

TRAV29/DV5*01-TRAJ31*01

CDR3: CAASYNNARLMF

α -chain (peptide 137)

TRAV29/DV5*01-TRAJ31*01

CDR3: CAASYNNARLMF

β -chain (peptide 101)

TRBV5-8*01- TRBD1*01-TRBJ2-5*01

CDR3: CASSWGYQETQYF

β -chain (peptide 125)

TRBV18*01- TRBD1*01- TRBJ1-2*01


CDR3: CASSPTGGEYYGYTF

β -chain (peptide 137)

TRBV6-5*01- TRBD1*01- TRBJ1-6*01

CDR3: CASSSYPLRTGRYNSYNSPLHF

j



Dominant clonotypes:

α -chain

TRAV29/DV5*01-TRAJ30*01

CDR3: CAASGGRDDKIIF

β -chain

TRBV6-5*01- TRBD1*01- TRBJ2-3*01

CDR3: CASSYSRTESTDTQYF

Figure 7

