

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2014/016362 A1

(43) International Publication Date
30 January 2014 (30.01.2014)

(51) International Patent Classification:
A61K 39/12 (2006.01) *A61K 39/295* (2006.01)

(74) Agent: **COMMANDER, Paul, M., B.**; Sanofi Pasteur, Intellectual Property Department, 2 avenue Pont Pasteur, F-69367 Lyon Cedex 07 (FR).

(21) International Application Number:

PCT/EP2013/065669

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
24 July 2013 (24.07.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
12305908.1 24 July 2012 (24.07.2012) EP
12305911.5 25 July 2012 (25.07.2012) EP

(71) Applicant (for all designated States except US): **SANOFI PASTEUR** [FR/FR]; 2 avenue Pont Pasteur, F-69367 LYON Cedex (07) (FR).

(72) Inventors; and

(71) Applicants (for US only): **YAO, Jiansheng** [CA/CA]; 15, Partridge Lane, Toronto, M1T 3C5 (CA). **GIRERD-CHAMBAZ, Yves** [FR/FR]; 22, chemin de la Pra, F-69510 Messimy (FR). **LEGASTELOIS, Isabelle** [FR/FR]; 392, rue de la Joannas, F-69700 Saint Andeol le Château (FR). **MANTEL, Nathalie** [FR/FR]; 84, rue de la Garenne, F-69005 Lyon (FR). **BARBAN, Véronique** [FR/FR]; 15, rue Sainte Marguerite, F-69110 Sainte-Foy-Les Lyon (FR). **LANG, Jean** [FR/FR]; 118, route de Saint Priest, F-69780 Moins (FR). **GUY, Bruno** [FR/FR]; 2, rue Casimir Perrier, F-69002 Lyon (FR).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))



WO 2014/016362 A1

(54) Title: VACCINE COMPOSITIONS FOR PREVENTION AGAINST DENGUE VIRUS INFECTION

(57) Abstract: The present invention relates to vaccine compositions that are useful in a method of protecting a human subject against dengue disease.

VACCINE COMPOSITIONS FOR PREVENTION AGAINST DENGUE VIRUS INFECTION**FIELD OF THE INVENTION**

The present invention relates to vaccine compositions and uses of such compositions in a method of protecting a human subject against dengue disease.

BACKGROUND

Dengue is the second most important infectious tropical disease after malaria with approximately one-half of the world's population living in areas where there is a risk of epidemic transmission. There are estimated to be 50-100 million cases of dengue disease every year resulting in 500,000 patients being hospitalized for dengue hemorrhagic fever (DHF) and resulting in approximately 25,000 deaths.

Dengue disease infections are endemic in more than 100 tropical countries and dengue hemorrhagic fever (DHF) has been documented in 60 of these countries (Gubler, 2002, *TRENDS in Microbiology*, 10: 100-103).

Dengue disease is caused by four antigenically distinct, but closely related dengue virus serotypes of the flavivirus genus (Gubler et al., 1988, in: *Epidemiology of arthropod-borne viral disease*. Monath TPM, editor, Boca Raton (FL): CRC Press: 223-60; Kautner et al., 1997, *J. of Pediatrics*, 131 : 516-524; Rigau-Perez et al., 1998, *Lancet*, 352: 971-977; Vaughn et al., 1997, *J. Infect. Dis.*, 176: 322-30).

Dengue disease is usually transmitted by injection of the dengue virus during the blood meal of an *Aedes aegypti* mosquito infected by the virus. After an incubation period of 4-10 days, the illness begins abruptly and is followed by three phases: febrile (2 to 7 days), critical (24-48 hours - during which severe complications may occur) and recovery (48-72 hours). During the critical phase, life threatening complications such as hemorrhages, shock and acute organ impairment may occur. A proper management of these unpredictable outcomes can reduce the case fatality rate. Cure of dengue fever is complete after 7 to 10 days, but prolonged asthenia is normal. Reduced leukocyte and platelet numbers are frequently observed.

Dengue haemorrhagic fever (DHF) is a potentially deadly complication of dengue virus infection. DHF is characterized by a high fever and symptoms of dengue disease, but with extreme lethargy and drowsiness. Increased vascular permeability and abnormal homeostasis can lead to a decrease in blood volume, hypotension, and in severe cases, hypovolemic shock and internal bleeding. Two factors appear to play a major role in the occurrence of DHF - rapid viral replication with a high level of viremia (the severity of the disease being associated with the level of viremia; Vaughn et al., 2000, *J. Inf. Dis.*, 181: 2-

9) and a major inflammatory response with the release of high levels of inflammatory mediators (Rothman and Ennis, 1999, *Virology*, 257: 1-6; Alan L. Rothman. 2011, *Nature Reviews Immunology*, 11: 532-543)). The mortality rate for DHF can reach 10% without treatment, but is < 1 % in most centres with access to treatment.

5 Dengue shock syndrome (DSS) is a common progression of DHF and is frequently fatal. DSS results from generalized vasculitis leading to plasma leakage into the extravascular space. DSS is characterized by rapid and poor volume pulse, hypotension, cold extremities, and restlessness.

10 In Asia, DHF and DSS are observed primarily in children, with approximately 90% of those with DHF being less than 15 years of age (Malavige et al., 2004, *Postgrad Med. J.*, 80: 588-601; Meulen et al., 2000, *Trop. Med. Int. Health*, 5:325-9). In contrast, outbreaks in the Caribbean and Central America have predominantly affected adults (Malavige et al., 2004, *Postgrad Med. J.*, 80: 588-601).

15 The four serotypes of dengue virus possess approximately 60-80% sequence homology. Infection with one dengue serotype provides durable homologous immunity but limited heterologous immunity (Sabin, 1952, *Am. J. Trop. Med. Hyg.*, 1: 30-50). Accordingly, an individual that has been infected with one serotype of dengue may subsequently become infected with a different serotype. In the past, it has been considered that a second infection arising from a different dengue virus serotype is 20 theoretically a risk factor for the development of DHF, since the majority of patients that exhibit DHF have been previously exposed to at least one of the other four serotypes of dengue viruses.

25 To date, there is no specific treatment for dengue disease. Treatment for dengue disease is symptomatic, with bed rest, control of the fever and pain through antipyretics and analgesics, and adequate drinking. The treatment of DHF requires balancing of liquid losses, replacement of coagulation factors and the infusion of heparin.

30 Since dengue prevention measures, such as mosquito control and personal protection from bites, are limited in efficacy, difficult to enforce and expensive, a safe and efficacious dengue vaccine would be the best mode of prevention. However, there is no licensed vaccine of this type that is currently available.

It is therefore desirable to develop a vaccine composition that demonstrates efficacy when used in a method of protecting a human subject against dengue disease.

SUMMARY OF THE INVENTION

The present invention relates to a dengue virus serotype 2 vaccine composition comprising:

5

- (i) a dengue antigen selected from the group consisting of:
 - (a) a live attenuated dengue virus;
 - (b) an inactivated dengue virus;
 - (c) a live attenuated or inactivated chimeric dengue virus;
 - (d) a dengue virus-like particle (VLP); and
 - (e) a combination of two or more of (a) to (d);

10

or

15

- (ii) a nucleic acid construct or viral vector which is able to express in a human cell a dengue antigen which is a dengue VLP;

wherein said dengue antigen comprises a polypeptide having at least 90% identity to SEQ ID NO: 12.

20

The present invention further relates to a vaccine composition comprising a dengue antigen of serotype 2 selected from the group consisting of: (a) a live attenuated dengue virus; (b) an inactivated dengue virus; (c) a live attenuated or inactivated chimeric dengue virus; or (d) a combination of two or more of (a) to (c); wherein said dengue antigen comprises a nucleotide sequence encoding a protein comprising a polypeptide or polypeptides as defined in the claims.

25

A vaccine composition comprising a dengue antigen of serotype 2 selected from the group consisting of: (a) a live attenuated dengue virus; (b) an inactivated dengue virus; (c) a live attenuated or inactivated chimeric dengue virus; or (d) a combination of two or more of (a) to (c); wherein said dengue antigen comprises a nucleotide sequence having at least 90% sequence identity to a sequence selected from the group consisting of the RNA equivalent of SEQ ID NO: 1, the RNA equivalent of SEQ ID NO: 4, the RNA equivalent of SEQ ID NO: 5, the RNA equivalent of SEQ ID NO: 6, the RNA equivalent of SEQ ID NO: 7 and SEQ ID NO: 25.

30

35

The present invention further relates to pharmaceutical formulation comprising a vaccine composition of the present invention and a pharmaceutically acceptable carrier, diluent or excipient.

5 The present invention further relates to a vaccine composition of the present invention for use in therapy.

The present invention further relates to a vaccine composition of the present invention for use in a method of protecting a human subject against dengue disease caused by a 10 dengue virus of serotype 2.

The present invention further relates to a vaccine composition of the present invention for use in a method of generating neutralising antibodies against a dengue virus of serotype 2.

15

The present invention further relates to vaccine composition of the present invention which comprises a dengue antigen of serotype 1, a dengue antigen of serotype 2, a dengue antigen of serotype 3 and a dengue antigen of serotype 4 for use in a method of generating neutralising antibodies against the four serotypes of dengue.

20

The present invention further relates to the use of a vaccine composition of the present invention for the manufacture of a medicament for protecting a human subject against dengue disease caused by a dengue virus of serotype 2.

25

The present invention further relates to a method of protecting a human subject against dengue disease caused by a dengue virus of serotype 2, wherein said method comprises administering to said subject an effective amount of a composition according to the present invention.

30

The present invention further relates to a kit comprising a composition according to the present invention and instructions for the use of said composition in a method of protecting a human subject against dengue disease caused by a dengue virus of serotype 2.

35

The present invention relates to a vaccine composition for use in a method of protecting a human subject against dengue disease, wherein said composition comprises:

5 (i) a dengue antigen selected from the group consisting of:

- (a) a live attenuated dengue virus;
- (b) an inactivated dengue virus;
- (c) a live attenuated or inactivated chimeric dengue virus;
- (d) a dengue virus-like particle (VLP); and
- (e) a combination of two or more of (a) to (d);

or

10 (iii) a nucleic acid construct or viral vector which is able to express in a human cell a dengue antigen which is a dengue VLP.

15 The present invention further relates to the use of a vaccine composition of the present invention for the manufacture of a medicament for protecting a human subject against dengue disease.

The present invention further relates to a method of protecting a human subject against dengue disease, wherein said method comprises administering to said human subject an effective amount of a composition according to the present invention.

20 Additionally, the present invention relates to a kit comprising a composition according to the present invention and instructions for the use of said composition in a method of protecting a human subject against dengue disease.

25 **Description of the Figure**

Figure 1 illustrates the construction of the YF-VAX cDNA by RT-PCR and cloning

Definitions

30 The term "Dengue disease", as used herein, refers to the clinical symptoms exhibited by an individual following infection by any one of the four Dengue virus serotypes. Since 1970, clinical dengue has been classified according to World Health Organization guidelines as (i) dengue fever or (ii) dengue hemorrhagic fever (World Health Organization. Dengue hemorrhagic fever: Diagnosis, treatment, prevention and 35 control 2nd Ed. Geneva: WHO, 1997; ISBN 92 4 154500 3). In 2009, the WHO issued new guidelines that classify clinical dengue as (i) dengue with or without warning signs or (ii)

severe dengue. Both classifications are shown in Figures 1 & 2 of Srikiatkachorn et al., Clin. Infect. Dis. (2011) 53(6): 563. According to the earlier classification, dengue fever is characterized by at least two symptoms selected from headache, arthralgia, retro-orbital pain, rash, myalgia, hemorrhagic manifestations, and leucopenia, together with supportive 5 serology or occurrence at the same location and time as other confirmed dengue cases. Progression to Dengue hemorrhagic fever is confirmed when fever, hemorrhagic manifestations, thrombocytopenia and evidence of plasma leakage are all observed. According to the more recent classification, diagnosis of dengue requires the presence of 10 fever and at least two clinical symptoms selected from nausea, vomiting, rash, aches and pains, a positive tourniquet test, or any warning signs selected from abdominal pain and tenderness, persistent vomiting, clinical fluid accumulation, mucosal bleed, lethargy or restlessness, liver enlargement greater than 2 cm or an increase in hematocrit concurrent with a rapid decrease in platelet count. Severe dengue is diagnosed when any of the 15 following events are observed: severe plasma leakage leading to shock or respiratory distress, severe bleeding as evaluated by clinicians or severe organ involvement

The term "Dengue hemorrhagic fever or DHF", as used herein, refers to virologically-confirmed dengue disease wherein fever, hemorrhagic manifestations, thrombocytopenia and evidence of plasma leakage are all observed. DHF, as used herein, may be further defined on the basis of its severity. For instance, DHF may be 20 defined as being of Grade I, Grade II, Grade III or Grade IV (World Health Organization. Dengue hemorrhagic fever: Diagnosis, treatment, prevention and control 2nd Ed. Geneva: WHO, 1997; ISBN 92 4 154500 3). Grade I is defined as fever accompanied by non-specific constitutional symptoms; the only haemorrhagic manifestation is a positive tourniquet test and/or easy bruising. Grade II is defined as spontaneous bleeding in 25 addition to the manifestations of Grade I patients, usually in the form of skin or other haemorrhages. Grade III is defined as circulatory failure manifested by a rapid, weak pulse and narrowing of pulse pressure or hypotension, with the presence of cold clammy skin and restlessness. Grade IV is defined as profound shock with undetectable blood pressure or pulse. As would be understood by a person of skill in the art, in the practice of 30 the present invention, e.g. a method of protecting against DHF, said DHF need not be virologically-confirmed.

The term "virologically-confirmed dengue", as used herein, refers to an acute febrile episode which is confirmed to be induced by a dengue virus, e.g. by reverse transcriptase polymerase chain reaction (RT-PCR) or by a dengue non-structural 1 (NS1) 35 protein enzyme-linked immunosorbent assay (ELISA). In the RT-PCR method, serum samples are tested according to the method of Callahan et al, J. Clin. Microbiol. (2001)

39: 4119. Briefly, RNA is extracted from the serum to discard potential Taq polymerase inhibitors or interfering factors, using a commercial kit. Then an RT-PCR reaction is carried out with serotype specific primers from the dengue NS5 gene sequence. Results are expressed as a concentration of \log_{10} GEQ (genome equivalent)/mL, by comparison with standards containing known concentrations of viral genomic serotype-specific nucleic acid sequences integrated into plasmids. In the ELISA method, 50 μ L of patient serum, a positive control, a negative control, or a cut-off control are diluted 1:2 in sample diluent and combined with 100 μ L of diluted horseradish peroxidase (HRP)-labeled anti-NS1 monoclonal Ab (MAb). The diluted serum and conjugate are added to capture anti-NS1 MAb-coated microwells, and plates are incubated for 90 minutes at 37°C. Capture MAb/NS1/HRP-labeled-MAb complexes are formed when NS1 is present in the serum. Complexes are detected via a colorimetric reaction in positive wells which is induced by adding 160 μ L of 3,3',5,5' tetramethylbenzidine (TMB) substrate and incubating for 30 minutes at room temperature in the dark. The reaction is stopped with the addition of 100 μ L of stop solution (1N H₂SO₄) and the plate is read. A sample ratio is determined for each sample by dividing the average optical density (OD) of the test sample by the average OD of the cut-off control (tested in quadruplicate). Sample ratios of <0.5, 0.5-
15 <1.0, and \geq 1 are indicative of negative, equivocal, and positive results, respectively.

The term "severe virologically-confirmed dengue", as used herein, refers to dengue haemorrhagic fever (DHF) as defined by the 1997 WHO classification and further characterized by the following additional list of symptoms: haemorrhage requiring blood transfusion, objective evidence of capillary permeability, signs of circulatory failure or visceral manifestations.

The term "dengue shock syndrome", as used herein, refers to the most severe complications of DHF as defined above. According to the 1997 WHO classification, DSS corresponds to DHF of Grades III and IV.

The term "dengue fever viruses", "dengue viruses" and "DEN" are used interchangeably. They refer to positive single-strand RNA viruses belonging to the Flavivirus genus of the family of flaviviridae. There are four different serotypes of dengue virus (serotypes 1, 2 3 and 4), which possess approximately 60-80% sequence homology. The organization of the genome comprises the following elements: a 5' non-coding region (NCR), a region encoding structural proteins (capsid (C), pre-membrane (prM) and envelope (E)) and a region encoding non-structural proteins (NS1-NS2A-NS2B-NS3-NS4A-NS4B-NS5) and a 3' NCR. The dengue viral genome encodes an uninterrupted coding region which is translated into a single polyprotein which undergoes post-translational processing. The sub-sequences included in the prM-E sequences may be

numbered in various ways: (i) the total prM-E protein sequence is numbered from position 1 to position 661, with the preM protein sequence designated as position 1 to position 90/91, the M protein sequence designated as position 91/92 to position 166 and the E protein sequence designated as position 167 to position 661; (ii) the prM and M protein sequences are numbered together, i.e. from position 1 to position 166 of the total sequence and E is numbered separately from position 1 to position 495; (iii) the prM, M and E sequences are numbered separately, i.e. prM is numbered from position 1 to 90/91, M is numbered from 1 to 75/76 and E from position 1 to position 495. In the present disclosure the E protein is always numbered from position 1 to position 495. For example, a residue designated herein as E-154 refers to position 154 of the E protein.

In the context of the present invention, "vaccinal dengue virus" refers to a virus which is capable of inducing neutralizing antibodies against the dengue virus serotype from which the vaccinal dengue virus is derived, by the administration of such vaccinal dengue virus to an immunocompetent subject. Examples of vaccinal dengue viruses which may be used in a method of the present invention include inactivated dengue viruses, live attenuated dengue viruses and live attenuated or inactivated chimeric dengue viruses. Serotypes of vaccinal dengue viruses for use in the present invention include serotypes 1, 2, 3, and 4. Preferably a vaccinal dengue virus for use in the present invention is a live attenuated chimeric dengue virus.

The expression "inactivated virus", as used herein, refers to a virus that is incapable of replication to any significant degree in cells permissive for replication of the corresponding wild type virus. Viruses may be inactivated by a number of means well known to those skilled in the art. Examples of methods for inactivating a virus include chemical treatments, or radiation treatments (including heat or electromagnetic radiation typically in the forms of X-ray or ultraviolet radiation).

The term "inactivated dengue virus", as used herein refers to an inactivated wild-type virus containing all the dengue structural proteins (env, premembrane/membrane and capsid proteins) and inactivated viral RNA. An inactivated dengue virus may also refer to an inactivated chimeric dengue virus. Inactivated dengue viruses are for instance described in United States Patent No. 6,254,873.

The term "live attenuated virus or LAV", as used herein, refers to a virus which is not able to induce a disease state characterised by the same sets of symptoms associated with the corresponding wild-type virus. Examples of live attenuated viruses are well known in the art. A live attenuated virus may be prepared from a wild-type virus, for example, by recombinant DNA technology, site directed mutagenesis, genetic

manipulation, serial passages on replication-competent cells, chemical mutagenesis treatment or electromagnetic radiation.

The term "live attenuated dengue virus", as used herein, refers to a live dengue virus derived from a virulent wild-type dengue virus by genetic modification resulting in 5 attenuation of virulence and an inability to induce a disease state characterised by the same sets of symptoms associated with the corresponding wild type dengue virus. Examples of live attenuated dengue viruses useful in the practice of the present invention include VDV-1, VDV-2, and the strains described for example in applications WO 02/66621, WO 00/57904, WO 00/57908, WO 00/57909, WO 00/57910, WO 02/0950075 10 and WO 02/102828. Live attenuated dengue viruses of serotype 1 which may be used in the method of the invention include VDV-1. Live attenuated dengue viruses of serotype 2 which may be used in the method of the invention include VDV-2, and LAV-2.

"VDV" and "Vero dengue vaccine" are used interchangeably herein and designate a live attenuated dengue virus capable of replication in Vero cells and capable of inducing 15 a specific humoral response, including the induction of neutralizing antibodies, in a human.

The DEN-1 16007/PDK13 strain, also called "LAV1", is derived from wild-type DEN-1 (dengue virus serotype 1) 16007 strain which has undergone 11 passages through primary dog kidney (PDK) cells (DEN-1 16007/PDK11). LAV1 has been described in 20 patent application EP1 159968 in the name of Mahidol University and has been filed with the National Microorganisms Cultures Collection (CNCM) under number I-2480. "VDV-1" is a virus derived from LAV1 by subsequent adaptation to Vero cells; in this regard, the RNA from LAV1 has been extracted and purified before being transfected into Vero cells. The VDV-1 strain has subsequently been obtained by plate purification and amplification 25 in Vero cells. The VDV-1 strain has 14 additional mutations in comparison with the DEN-1 16007/PDK13 strain (13 passes through PDK cells). A process for preparing and characterizing the VDV-1 strain has been described in international patent application filed under number WO06/134433 in the names of Sanofi-Pasteur and the Center for Disease Control and Prevention.

The DEN-2 16681/PDK53 strain, also known as "LAV2", has been obtained from 30 wild-type strain DEN-2 (dengue virus serotype 2) 16681 which has undergone 50 passes through PDK cells (DEN-2 16681/PDK50). LAV2 has been described in in patent application EP1159968 in the name of Mahidol University and has been filed with the National Microorganisms Cultures Collection (CNCM) under number 1-2481. "VDV-2" is a 35 strain derived from LAV2 by subsequent adaptation to Vero cells; in this regard, the RNA from LAV2 has been extracted and purified before being transfected in Vero cells. The

VDV-2 strain has subsequently been obtained by plate purification and amplification in Vero cells. The VDV-2 strain has 10 additional mutations in comparison with the DEN-2 16681/PDK53 strain (53 passes through PDK cells), including 4 silent mutations. A process for preparing and characterizing the VDV-2 strain has been described in the 5 international patent application filed under number WO06/134443 in the names of Sanofi-Pasteur and the Center for Disease Control and Prevention. The complete nucleic acid sequence of the VDV-2 strain is as shown in SEQ ID NO: 24. The sequence of the E protein of the VDV-2 strain is as shown in SEQ ID NO: 26 and the sequence of the M protein of the VDV-2 strain is as shown in the SEQ ID NO: 27.

10 The VDV 1 and 2 strains are prepared by amplification in Vero cells. The viruses produced are harvested and clarified from cell debris by filtration. The DNA is digested by treatment with enzymes. Impurities are eliminated by ultrafiltration. Infectious titers may be increased by a concentration method. After adding a stabilizer, the strains are stored in lyophilized or frozen form before use and then reconstituted when needed.

15 In the context of the invention, "dengue chimera or chimeric dengue virus" means a recipient flavivirus in which the genetic backbone has been modified by exchanging the sequences encoding the prM and E proteins of the recipient flavivirus by the corresponding sequences of a dengue virus. Typically, the recipient flavivirus may be attenuated. The recipient flavivirus may be a yellow fever (YF) virus such as the 20 attenuated YF 17D, YF 17DD and YF 17D204 (YF-VAX®) viruses; in that case, such chimeras are referred to as YF/dengue chimeras. The recipient flavivirus may also be a dengue virus and in that case, it is referred to as dengue/dengue chimera, the dengue virus serotype characteristic of the prM and E proteins being identical or different from the recipient dengue virus serotype characteristic of the genetic backbone. When the 25 serotypes are identical, the recipient dengue virus and the dengue virus from which the prM and E protein encoding sequences originate, are two different virus strains of the same serotype. For use in the present invention, chimeric dengue viruses are typically YF/dengue chimeras. Chimeric dengue viruses are preferably inactivated or live attenuated chimeric dengue viruses. Advantageously, the recipient flavivirus of a live 30 attenuated chimeric dengue virus of the present invention is YF 17D or YF 17D204 (YF-VAX®). According to one embodiment dengue chimera is an inactivated virus. According to an alternative embodiment the dengue chimera is a live attenuated virus. Dengue Chimera that can be used in a vaccine composition of the present invention include Chimerivax™ Dengue Serotype 1 (also known as CYD-1), Chimerivax™ Dengue 35 Serotype 2 (also known as CYD-2), Chimerivax™ Dengue Serotype 3 (also known as CYD-3) and Chimerivax™ Dengue Serotype 4 (also known as CYD-4).

Examples of chimeric dengue viruses useful in the practice of the present invention include the dengue/YF chimeric viruses described in patent application WO 98/37911 and dengue/dengue fever chimeras such as those described in patent applications WO 96/40933 and WO 01/60847.

5 In one embodiment, the chimeric YF/dengue virus comprises the genomic backbone of the attenuated yellow fever virus strain YF17D (Theiler M. and Smith H.H., 1937, J.Exp.Med., 65: 767-786), e.g. viruses YF17D/DEN-1, YF17D/DEN-2, YF17D/DEN-3 and YF17D/DEN-4. Examples of YF17D strains which may be used include YF17D204 (YF-VAX(R), Sanofi-Pasteur, Swiftwater, PA, USA; Stamaril(R), Sanofi-Pasteur, Marcy 10 l'Etoile, France; ARILVAX(TM), Chiron, Speke, Liverpool, UK; FLAVIMUN(R), Berna Biotech, Bern, Switzerland; YF17D-204 France (X15067, X15062); YF17D-204,234 US (Rice et al., 1985, Science, 229: 726-733), or the related strains YF17DD (Genbank access number U17066), YF17D-213 (Genbank access number U17067) and the strains 15 YF17DD described by Galler et al. (1998, Vaccines, 16(9/10): 1024-1028). In another embodiment, the chimeric YF/dengue virus comprises the genomic backbone of the attenuated yellow fever virus strain YF17D204 (YF-VAX®).

One example of a chimeric dengue virus particularly suitable for use in the practice of the present invention is a "Chimerivax dengue virus". As used herein, a "Chimerivax dengue virus", is a live attenuated chimeric YF/dengue virus which comprises the genomic 20 backbone of a YF17D or YF17D204 (YF-VAX®) virus in which the nucleic acid sequences encoding the pre-membrane (prM) and envelope (E) proteins have been replaced by nucleic acid sequences encoding the corresponding structural proteins of a dengue virus. A preferred chimeric dengue virus for use in the present invention is a live attenuated chimeric YF/dengue virus which comprises the genomic backbone of a YF17D virus in 25 which the nucleic acid sequences encoding the pre-membrane (prM) and envelope (E) proteins have been replaced by nucleic acid sequences encoding the corresponding structural proteins of a dengue virus. A preferred chimeric dengue virus for use in the present invention is a live attenuated chimeric YF/dengue virus which comprises the genomic backbone of a YF17D204 (YF-VAX®) virus in which the nucleic acid sequences 30 encoding the pre-membrane (prM) and envelope (E) proteins have been replaced by nucleic acid sequences encoding the corresponding structural proteins of a dengue virus. Construction of such Chimerivax viruses may be achieved in accordance with, or in substantial 35 accordance with, the teaching of Chambers, et al. (1999, J. Virology 73(4):3095-3101). The particular Chimerivax (CYD) dengue viruses described in the examples have been generated by using prM and E sequences from strains DEN 1 PU0359 (TYP1 140), DEN2 PUO218, DEN3 PaH881/88 and DEN 4 1228 (TVP 980) and

the genomic backbone of YF17D virus. Those particular Chimerivax strains are referred to herein (see the present examples) as "CYD-1", "CYD-2", "CYD-3" and "CYD-4" respectively. The preparation of these particular CYD-1, CYD-2, CYD-3 and CYD-4 strains has been described in detail in international patent applications WO 98/37911, WO 5 03/101397, WO 07/021672, WO 08/007021, WO 08/047023 and WO 08/065315, to which reference may be made for a precise description of the processes for their preparation. Alternatively, other dengue fever virus strains may be used as a source of nucleic acids to facilitate construction of chimeric viruses useful in the practice of the present invention, for example in the construction of other Chimerivax dengue serotype 1 (CYD-1), Chimerivax 10 dengue serotype 2 (CYD-2), Chimerivax dengue serotype 3 (CYD-3) and Chimerivax dengue serotype 4 (CYD-4) strains. Advantageously, a vaccine composition of the present invention, e.g. a chimeric dengue virus, of serotype 2 may comprise prM-E sequences having at least 90%, at least 95%, at least 98% or at least 99% identity to the prM-E sequences from the serotype 2 strains LAV-2, BID-V585, PR/DB023 or MD1280 as 15 described in the examples or may comprise prM-E sequences having at least 90%, at least 95%, at least 98% or at least 99% identity to the prM-E sequence shown in SEQ ID NO: 2. Advantageously, a vaccine composition, e.g. a chimeric dengue virus, of serotype 2 for use in the method of the present invention may comprise prM-E sequences from the serotype 2 strains LAV-2, BID-V585, PR/DB023 or MD1280 or the prM-E sequence from 20 SEQ ID NO: 2 as described in the examples. When the recipient genomic backbone of such chimeric dengue viruses is derived from YF-VAX®, such strains are referred to herein as CYD-LAV, CYD-BID, CYD-PR and CYD-MD. A vaccine composition of the present invention comprising chimeric dengue virus of serotype 2 generated using the prM-E sequences of the serotype 2 strains LAV-2 (SEQ ID NO: 8), BID-V585 (SEQ ID 25 NO: 9), PR/DB023 (SEQ ID NO: 10), MD1280 (SEQ ID NO: 11) or SEQ ID NO: 2, or generated using prM-E sequences having at least 90%, at least 95%, at least 98% or at least 99% identity to the prM-E sequences from the serotype 2 strains LAV-2, BID-V585, PR/DB023, MD1280 or the prM-E sequence from SEQ ID NO: 2 may advantageously be used in combination with CYD-1, CYD-3 and CYD-4 in a vaccine composition according to 30 the present invention. Examples of chimeric dengue virus of serotype 2 generated using the prM-E sequences of the serotype 2 strains LAV-2 (SEQ ID NO: 8), PR/DB023 (SEQ ID NO: 10) and MD1280 (SEQ ID NO: 11) include CYD-LAV, CYD-PR and CYD-MD respectively.

An alternative embodiment of chimeric dengue virus usable in the method of 35 protection of the invention is a recipient flavivirus in which the genetic backbone has been modified by exchanging (i) the sequence encoding the E protein of the recipient flavivirus

by the corresponding sequence of a dengue virus and (ii) the sequence encoding the prM protein of the recipient flavivirus by the corresponding sequence of a non-dengue flavivirus, e.g. a JEV virus. Typically, the said chimeric virus may be a live attenuated virus or an inactivated virus. Examples of such chimeric dengue viruses are described in

5 WO2011/138586.

A vaccinal dengue virus of serotype 1 for use in a vaccine composition of the present invention may, for example, be the strain VDV1, CYD-1 or a YF17D/DEN-1 chimeric virus comprising the prM and E amino acid sequences of the DEN-1 16007/PDK13 strain. A vaccinal dengue virus of serotype 2 for use in the method of the present invention may, for example, be the strain VDV2, CYD-2, a YF17D/DEN-2 chimeric virus comprising the prM and E amino acid sequences of the DEN-2 16681/PDK53 strain, a chimeric virus comprising the prM and E amino acid sequences of the DEN-2 strains LAV-2, BID-V585, PR/DB023 or MD1280 or a chimeric virus comprising prM-E sequences having at least 90%, at least 95%, at least 98% or at least 99% identity to the prM-E sequences from the serotype 2 strains LAV-2, BID-V585, PR/DB023 or MD1280 or at least 90%, at least 95%, at least 98% or at least 99% identity to the prM-E sequence in SEQ ID NO: 2. A vaccinal dengue virus of serotype 3 for use in the method of present invention may, for example, be CYD-3 or an alternative YF17D/DEN-3 chimeric virus. An example of a vaccinal dengue virus of serotype 4 is CYD-4 or an alternative YF17D/DEN-4 chimeric virus.

A composition of the present invention comprises at least one dengue antigen. Typically a composition of the present invention comprises a dengue antigen, e.g. a vaccinal dengue virus, of each of serotypes 1, 2, 3 and 4. Dengue antigens, e.g. vaccinal dengue viruses, of the present invention of each serotype may be as described herein.

25 For instance, a composition of the present invention may advantageously comprise any one of the following combinations of dengue antigens: i) a dengue antigen comprising the prM and E sequences of CYD-1, a dengue antigen comprising the prM and E sequences of CYD-LAV, a chimeric dengue virus comprising the prM and E amino acid sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4; ii) a dengue 30 antigen comprising the prM and E sequences of CYD-1, a dengue antigen comprising the prM and E sequences of CYD-BID, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4; (iii) a dengue antigen comprising the prM and E sequences of CYD-1, a dengue antigen comprising the prM and E sequences of CYD-PR, a dengue antigen comprising the prM 35 and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4; (iv) a dengue antigen comprising the prM and E sequences of CYD-1, a

dengue antigen comprising the prM and E sequences of CYD-MD, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4;. For instance, a composition of the present invention may also advantageously comprise any one of the following combinations of dengue antigens:

- 5 i) CYD-1, CYD-LAV, CYD-3 and CYD-4; ii) CYD-1, CYD-BID, CYD-3 and CYD-4; (iii) CYD-1, CYD-PR, CYD-3 and CYD-4 or (iv) CYD-1, CYD-MD, CYD-3 and CYD-4. A composition of the present invention may also advantageously comprise the following combination of dengue antigens: i) a dengue antigen comprising the prM and E sequences of CYD-1, VDV2, a dengue antigen comprising the prM and E sequences of
- 10 CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4. For instance, a composition of the present invention may advantageously comprise CYD-1, VDV-2, CYD-3 and CYD-4. A composition of the present invention, as described herein, may advantageously comprise a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 15) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2. A composition of the present invention, as described herein, may advantageously comprise a dengue antigen of serotype 2 which comprises a sequence having at least 90% identity to the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2. For example, said sequence may be at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2.
- 20
- 25

The term "virus-like particles or VLPs", as used herein, refers to virus particles that do not contain replicative genetic material but present at their surface a dengue E protein in a repetitive ordered array similar to the virion structure. Typically, dengue VLPs also

- 30 contain dengue prM and/or M, and E proteins. VLPs may be produced *in vitro* (Zhang et al, J. Virol. (2011) 30 (8):333). VLPs may also be produced *in vivo*. To that end, nucleic acid constructs (e.g. DNA or RNA constructs) encoding prM and E dengue proteins may be introduced into a cell of a subject, e.g. a human subject, via methods known in the art, e.g. via use of a viral vector. Any viral vector may be used provided it is able to contain and express both prM and E dengue virus sequences. Non-limiting examples of viral vectors that may be used in the method of the present invention include the poxviruses (e.g. the attenuated pox Ankara virus) and the measles virus. For use in the present invention, a particular category of viral vector expressing VLPs *in vivo* includes replication-35 deficient pseudoinfectious (PIV) viruses, e.g. according to the Replivax™ technology. (Rumyantsev AA, et al. Vaccine. 2011 Jul 18;29(32):5184-94).
- 35

The term "replication-defective pseudo-infectious virus", as used herein, refers to a virion particle that is replication-defective *in vivo*, owing to the absence in their genome of an essential sequence of the replicative cycle, for example the sequence encoding a capsid protein. However, the virion particles can propagate in a culture of helper cells that provide for the essential sequence(s) *in trans*. Replication-deficient pseudoinfectious viruses for use in the present invention include any virus according to the above definition which is capable of expressing the prM and E proteins of a dengue virus of any serotype. Examples include replication defective flavivirus / dengue chimeras such as replication defective West Nile virus / dengue, Japanese Encephalitis virus / dengue and YF / dengue chimeras.

The ability of a vaccine composition of the present invention to provoke an immune response in a subject (i.e. induce the production of neutralizing antibodies) can be assessed, for example, by measuring the neutralizing antibody titre raised against the dengue virus serotype(s) comprised within the composition. The neutralizing antibody titre may be measured by the Plaque Reduction Neutralization Test (PRNT₅₀) test. Briefly, neutralizing antibody titre is measured in sera collected from vaccinated subjects at least 28 days following administration of a vaccine composition of the present invention. Serial, two-fold dilutions of sera (previously heat-inactivated) are mixed with a constant challenge-dose of each dengue virus of serotype 1, 2, 3 or 4 as appropriate (expressed as PFU/mL). The mixtures are inoculated into wells of a microplate with confluent Vero cell monolayers. After adsorption, cell monolayers are incubated for a few days. The presence of dengue virus infected cells is indicated by the formation of infected foci and a reduction in virus infectivity due to the presence of neutralising antibodies in the serum samples can thus be detected. The reported value (end point neutralization titre) represents the highest dilution of serum at which ≥ 50 % of dengue challenge virus (in foci counts) is neutralized when compared to the mean viral focus count in the negative control wells (which represents the 100% virus load). The end point neutralization titres are presented as continuous values. The lower limit of quantification (LLOQ) of the assay is 10 (1/dil). It is commonly considered that seroconversion occurs when the titer is superior or equal to 10 (1/dil). As PRNT tests may slightly vary from a laboratory to another the LLOQ may also slightly vary. Accordingly, in a general manner, it is considered that seroconversion occurs when the titre is superior or equal to the LLOQ of the test. Neutralising antibody titres were considered in the following references, but the authors did not establish a correlate of protection (Guirakhoo et al, J. Virol. (2004) 78 (9): 4761; Libraty et al, PLoS Medicine (2009) 6 (10); Gunther et al, Vaccine (2011) 29: 3895) and Endy et al, J. Infect. Dis. (2004), 189(6): 990-1000).

The term "CCID₅₀" refers to the quantity of virus (e.g. vaccinal virus) infecting 50% of the cell culture. The CCID₅₀ assay is a limit dilution assay with statistical titer calculation (*Morrison D et al J Infect Dis. 2010; 201(3):370-7*)).

The term "human subject" is intended to mean males and females of various ages.

5 Preferably a human subject according to the present invention is less than 18 years of age or less than 12 years of age. For example, a human subject according to the present invention may be 0-17 years of age, 0-11 years of age, 4-17 years of age, 4-11 years of age, 4-6 years of age, 6-8 years of age, 8-10 years of age, 2-8 years of age, 2-11 years of age, 2-14 years of age, 9-16 years of age, 12-17 years of age or 18-45 years of age. More
10 preferably, a human subject according to the present invention is 4-11 years of age, 2-14 years of age or 9-16 years of age. A human subject according to the present invention may be at least 9 months old or less than 9 months old. For instance a human subject according to the present invention may be 9 months to 16 years of age, 9 months to 14 years of age, 9 months to 11 years of age or 9 months to 8 years of age. A human subject
15 according to the present invention may be at least 9 months old, with no history of severe allergy to any component of the vaccine composition as defined herein, no congenital or acquired immune deficiency, no symptomatic HIV infection and said subject should not be pregnant or breast feeding.

As used herein, the expression "flavivirus-naïve subject" refers to a subject who
20 has not been infected by a flavivirus nor previously immunized with a flavivirus vaccine, i.e. a serum sample taken from said subject will produce a negative result in a flavivirus ELISA or PRNT assay.

As used herein, the expression "dengue-naïve subject" refers to a subject who has not been infected by a dengue virus nor previously immunized with a dengue vaccine, i.e.
25 a serum sample taken from said subject will produce a negative result in a dengue ELISA or PRNT assay.

As used herein, the expression "flavivirus-immune subject" refers to a subject who has been infected or immunized by a flavivirus before administration of the vaccine composition of the invention, i.e. a serum sample taken from said subject will produce a positive result in a flavivirus ELISA or PRNT assay.
30

As used herein, the expression "dengue-immune subject" refers to a subject who has been infected by a dengue virus or immunized by a dengue vaccine before administration of the vaccine composition of the present invention, i.e. a serum sample taken from said subject will produce a positive result in a dengue ELISA or PRNT assay.

35 In accordance with the present invention, a "method of protecting", as used herein, results in a reduction in the severity or in the likelihood of developing dengue disease in a

human subject exposed to a dengue virus. Advantageously, said reduction is statistically significant. For example, a method of protecting, according to the present invention, may result in a reduction in at least one symptom of dengue disease as defined herein or a reduction in a combination of any two or more of those symptoms. The protection may 5 result in any one or more of the following:

- (i) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, symptomatic virologically-confirmed dengue disease caused by dengue virus of any serotype;
- (ii) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, symptomatic virologically-confirmed dengue disease caused by dengue virus of any one of serotypes 1, 3 or 4;
- (iii) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, symptomatic dengue disease caused by dengue virus of any serotype;
- 15 (iv) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, symptomatic dengue disease caused by dengue virus of any one of serotypes 1, 3 or 4;
- (v) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, severe virologically-confirmed dengue caused by dengue virus of any serotype;
- 20 (vi) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, severe dengue disease caused by dengue virus of any serotype;
- (vii) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, dengue hemorrhagic fever cases of Grades I to IV caused by dengue virus of any serotype;
- 25 (viii) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, DHF cases of Grade I caused by dengue virus of any serotype;
- (ix) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, DHF cases of Grade II caused by dengue virus of any serotype;
- 30 (x) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, DHF cases of Grade III caused by dengue virus of any serotype;

- (xi) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, DHF cases of Grade IV caused by dengue virus of any serotype;
- 5 (xii) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, fever or a reduction in the mean duration and/or intensity of fever;
- 10 (xiii) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, plasma leakage as defined by a change in haematocrit or a reduction in the mean value for plasma leakage as defined by a change in haematocrit;
- (xiv) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, thrombocytopenia or a reduction in the mean value for thrombocytopenia;
- 15 (xv) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, increases in the level of liver enzymes including alanine aminotransferase (ALT) and aspartate aminotransferase (AST);
- (xvi) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, hospitalization due to virologically-confirmed dengue disease caused by dengue virus of any serotype;
- 20 (xvii) a statistically significant reduction in the incidence or likelihood of, e.g. the prevention of, hospitalization due to dengue disease caused by dengue virus of any serotype;
- (xviii) a statistically significant reduction in the length of hospital stay due to virologically-confirmed dengue disease.
- 25 (xix) a statistically significant reduction in the length of hospital stay due to dengue disease.

The duration and intensity of fever are monitored and recorded according to standard hospital procedures. In a human subject, a fever (i.e. a febrile episode) is defined as the observance of two temperature readings of at least 37.5°C measured twice over an interval of at least 4 hours. Measurement of haematocrit, thrombocytopenia and hepatic enzyme levels are standard tests well-known to the person of skill in the art, for example as described in the pharmacopea.

Protection against dengue disease, for example as defined in points (i) to (xix) above, may be demonstrated in respect of dengue disease caused by a particular dengue virus serotype. For example, protection against dengue disease, as defined herein, may be demonstrated in respect of dengue disease caused by a dengue virus of serotype 1, a

dengue virus of serotype 2, a dengue virus of serotype 3 or a dengue virus of serotype 4. Advantageously, protection against dengue disease, as defined herein, may be demonstrated in respect of dengue disease caused by, for example, dengue virus of serotype 1 or serotype 3, dengue virus of serotype 1 or serotype 4, dengue virus of serotype 3 or serotype 4, dengue virus of serotype 1 or serotype 2, dengue virus of serotype 2 or serotype 3, dengue virus of serotype 2 or serotype 4, dengue virus of serotype 1, 2 or 3, dengue virus of serotype 1, 3 or 4, dengue virus of serotype 2, 3 or 4 or dengue virus of serotype 1, 2, 3 or 4.

Protection against dengue disease, as defined herein, may advantageously be demonstrated in particular sub-groups of human subjects. For instance, protection against dengue disease may advantageously be demonstrated in a human subject who is less than 18 years of age or less than 12 years of age. For example, a human subject according to the present invention may be 0-17 years of age, 0-11 years of age, 4-17 years of age, 4-11 years of age, 4-6 years of age, 6-8 years of age, 8-10 years of age, 2-8 years of age, 2-11 years of age, 2-14 years of age, 9-16 years of age, 12-17 years of age or 18-45 years of age. More preferably, a human subject according to the present invention is 4-11 years of age, 2-14 years of age or 9-16 years of age. A human subject according to the present invention may be at least 9 months old or less than 9 months old. For instance a human subject according to the present invention may be 9 months to 16 years of age, 9 months to 14 years of age, 9 months to 11 years of age or 9 months to 8 years of age. A human subject according to the present invention may be at least 9 months old, with no history of severe allergy to any component of the vaccine composition as defined herein, no congenital or acquired immune deficiency, no symptomatic HIV infection and said subject should not be pregnant or breast feeding.

Protection against dengue disease, as defined herein, may advantageously be demonstrated in particular countries, areas or regions of the world. For instance, protection against dengue disease may advantageously be demonstrated in a dengue endemic area. For instance, a dengue endemic area according to the present invention in which protection may be demonstrated may comprise those American countries or parts thereof which fall within the tropics and sub-tropics. A dengue endemic area in which protection may be demonstrated according to the present invention may thus comprise any one or more of the following: Brazil, Venezuela, Colombia, Ecuador, Peru, Bolivia, Paraguay, Panama, Costa Rica, Nicaragua, Honduras, El Salvador, Guatemala, Belize, Mexico, the USA and the islands of the Caribbean. In a particular embodiment, a dengue endemic area of the present invention in which protection may be demonstrated may consist of the following: Brazil, Colombia, Honduras, Mexico and Puerto Rico. A dengue

endemic area in which protection may be demonstrated according to the present invention may also include south Asian and Oceania countries within the tropics and sub-tropics. A dengue endemic area according to the present invention in which protection may be demonstrated may thus consist of any one or more of the following: India, Myanmar (Burma), Thailand, Laos, Vietnam, Cambodia, Indonesia, Malaysia, Singapore, the Philippines, Taiwan, Papua New Guinea and Australia. In a dengue endemic area in which protection may be demonstrated according to the present invention, a particular serotype, strain or genotype of wild type dengue virus may be the dominant circulating strain. For example, a dengue virus of serotype 2 may be characterised as having an Asian I or an Asian/American genotype. Asian/American genotype strains are characterised by at least one of, at least two of, at least three of, at least four of, at least five of or all six of the following residues Arg, Asn, Asp, Thr, Gly and His at positions prM-16, E-83, E-203, E-226, E-228 and E-346 respectively (wherein prM-16 designates position 16 of the prM protein and E-83 etc. designates position 83 of the E protein). Asian I genotype strains are characterised by at least one of, at least two of, at least three of, at least four of, at least five of or all six of the following residues Ile, Lys, Asn, Arg, Glu and Tyr at positions prM-16, E-83, E-203, E-226, E-228 and E-346 respectively (see Table 1 of Hang et al., PLoS NTD, 4(7): e757). A preferred dengue endemic area in which protection may be demonstrated according to the present invention is one in which a dengue virus having an Asian/American genotype is the dominant circulating strain, i.e. at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or 100% of the cases of dengue disease in said dengue endemic area are caused by dengue virus having an Asian/American genotype. A preferred dengue endemic area in which protection may be demonstrated according to the present invention is one in which a dengue virus of any one or more of serotypes 1, 3 or 4 is/are the dominant circulating serotype(s), i.e. at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or 100% of the cases of dengue disease are caused by dengue virus of serotypes 1, 3 or 4.

The term "RNA equivalent" of a given DNA sequence, as used herein, refers to a sequence wherein deoxythymidines have been replaced by uridines. Since the DNA sequences in question constitute the cDNA sequences of the dengue viruses, the equivalent RNA sequences constitute the positive strand RNA of those dengue viruses.

Overview of Several Embodiments

The present inventors have, for the first time, demonstrated the efficacy of a vaccine composition in protecting a human subject against dengue disease.

The present invention relates to a dengue virus serotype 2 vaccine composition comprising:

5 (i) a dengue antigen selected from the group consisting of:

(a) a live attenuated dengue virus;

(b) an inactivated dengue virus;

(c) a live attenuated or inactivated chimeric dengue virus;

(d) a dengue virus-like particle (VLP); and

10 (e) a combination of two or more of (a) to (d);

or

15 (ii) a nucleic acid construct or viral vector which is able to express in a human cell a dengue antigen which is a dengue VLP;

wherein said dengue antigen comprises a polypeptide having at least 90% identity to SEQ ID NO: 12.

In preferred embodiments, said polypeptide has at least 92%, at least 94%, at least 96%, at least 98%, at least 99%, at least 99.5% identity or 100% identity with SEQ ID NO: 12.

Preferably, said dengue antigen is selected from the group consisting of a live attenuated dengue virus and a live attenuated or inactivated chimeric dengue virus. Preferably, said dengue antigen is selected from the group consisting of a live attenuated dengue virus and a live attenuated chimeric dengue virus. Preferably, said dengue antigen is a live attenuated chimeric dengue virus.

Preferably said dengue antigen according to the present invention comprises a polypeptide having at least 90% identity to SEQ ID NO: 12, for example at least 92%, at least 94%, at least 96%, at least 98%, at least 99%, at least 99.5% identity or 100% identity to SEQ ID NO: 12 over the full length of SEQ ID NO: 12.

Preferably, said dengue antigen does not comprise the prM-E sequence of CYD-2, as defined herein.

Preferably, said vaccine composition does not comprise CYP-2.

Preferably, said dengue antigen comprises a polypeptide which comprises a valine residue at the position within the polypeptide that corresponds to position 251 of SEQ ID NO: 12.

Preferably, said dengue antigen comprises a polypeptide which comprises a methionine residue at the position within the polypeptide that corresponds to position 6 of SEQ ID NO: 12.

5 Preferably, said dengue antigen comprises a polypeptide which comprises a valine residue at the position within the polypeptide that corresponds to position 129 of SEQ ID NO: 12.

Preferably, said dengue antigen comprises a polypeptide which comprises an isoleucine residue at the position within the polypeptide that corresponds to position 129 of SEQ ID NO: 12.

10 Preferably, said dengue antigen comprises a polypeptide which comprises an isoleucine residue at the position within the polypeptide that corresponds to position 141 of SEQ ID NO: 12.

15 Preferably, said dengue antigen comprises a polypeptide which comprises an isoleucine residue at the position within the polypeptide that corresponds to position 164 of SEQ ID NO: 12.

Preferably, said dengue antigen comprises a polypeptide which comprises an aspartate residue at the position within the polypeptide that corresponds to position 203 of SEQ ID NO: 12.

20 Preferably, said dengue antigen comprises a polypeptide which comprises an asparagine residue at the position within the polypeptide that corresponds to position 203 of SEQ ID NO: 12.

Preferably, said dengue antigen comprises a polypeptide which comprises a threonine residue at the position within the polypeptide that corresponds to position 226 of SEQ ID NO: 12.

25 Preferably, said dengue antigen comprises a polypeptide which comprises a glycine residue at the position within the polypeptide that corresponds to position 228 of SEQ ID NO: 12.

30 Preferably, said dengue antigen comprises a polypeptide which comprises an isoleucine residue at the position within the polypeptide that corresponds to position 308 of SEQ ID NO: 12.

Preferably, said dengue antigen comprises a polypeptide which comprises a valine residue at the position within the polypeptide that corresponds to position 308 of SEQ ID NO: 12.

35 Preferably, said dengue antigen comprises a polypeptide which comprises a threonine residue at the position within the polypeptide that corresponds to position 478 of SEQ ID NO: 12.

Preferably, said dengue antigen comprises a polypeptide which comprises a valine residue at the position within the polypeptide that corresponds to position 484 of SEQ ID NO: 12.

5 Preferably, said dengue antigen comprises a polypeptide which comprises an isoleucine residue at the position within the polypeptide that corresponds to position 484 of SEQ ID NO: 12.

Preferably, said dengue antigen comprises a polypeptide which comprises a isoleucine residue at the position within the polypeptide that corresponds to position 485 of SEQ ID NO: 12.

10 Preferably, said dengue antigen comprises a polypeptide which comprises a alanine residue at the position within the polypeptide that corresponds to position 491 of SEQ ID NO: 12.

Preferably, said dengue antigen comprises a polypeptide having at least 90% sequence identity to SEQ ID NO: 3.

15 In preferred embodiments, said polypeptide has at least 92%, at least 94%, at least 96%, at least 98%, at least 99%, at least 99.5% identity or 100% identity with SEQ ID NO: 3.

20 Preferably, a composition of the present invention comprises a polypeptide having at least 90%, at least 92%, at least 94%, at least 96%, at least 98%, at least 99%, at least 99.5% identity or 100% identity with SEQ ID NO: 3. Preferably a dengue antigen according to the present invention comprises a polypeptide having at least 90% identity to SEQ ID NO: 12, for example at least 92%, at least 94%, at least 96%, at least 98%, at least 99%, at least 99.5% identity or 100% identity to SEQ ID NO: 12 over the full length of SEQ ID NO: 12. Preferably, said dengue antigen comprises a polypeptide which 25 comprises a glycine residue at the position within the polypeptide that corresponds to position 15 of SEQ ID NO: 3.

Preferably, said dengue antigen comprises a polypeptide which comprises a serine residue at the position within the polypeptide that corresponds to position 15 of SEQ ID NO: 3.

30 Preferably, said dengue antigen comprises a polypeptide which comprises a leucine residue at the position within the polypeptide that corresponds to position 24 of SEQ ID NO: 3.

35 Preferably, said dengue antigen comprises a polypeptide which comprises an isoleucine residue at the position within the polypeptide that corresponds to position 39 of SEQ ID NO: 3.

Preferably, said dengue antigen comprises a polypeptide which comprises a methionine residue at the position within the polypeptide that corresponds to position 39 of SEQ ID NO: 3.

5 Preferably, said dengue antigen comprises a polypeptide which comprises a valine residue at the position within the polypeptide that corresponds to position 120 of SEQ ID NO: 3.

Preferably, said dengue antigen comprises a polypeptide which comprises an alanine residue at the position within the polypeptide that corresponds to position 120 of SEQ ID NO: 3.

10 Preferably, said dengue antigen comprises a polypeptide which comprises a threonine residue at the position within the polypeptide that corresponds to position 125 of SEQ ID NO: 3.

15 Preferably, the polypeptides as defined herein (as comprised within the dengue antigens as comprised within the vaccine compositions of the present invention) comprise a threonine residue at the position within the polypeptide that corresponds to position 125 of SEQ ID NO: 3 and a valine residue at the position within the polypeptide that corresponds to position 417 of SEQ ID NO: 3.

20 Preferably, the polypeptides which are encoded by the nucleotide sequences as defined herein (i.e as comprised within the vaccine compositions of the present invention) comprise a leucine residue at the position within the polypeptide that corresponds to position 24 of SEQ ID NO: 3, a threonine residue at the position within the polypeptide that corresponds to position 125 of SEQ ID NO: 3 and a valine residue at the position within the polypeptide that corresponds to position 417 of SEQ ID NO: 3.

25 Preferably, a polypeptide (as comprised within a dengue antigen as comprised within a vaccine composition of the invention) comprises (i) the sequence as set forth in SEQ ID NO: 13 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 13; (ii) the sequence as set forth in SEQ ID NO: 14 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 14; (iii) the sequence as set forth in SEQ ID NO: 15 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 15; (iv) the sequence as set forth in SEQ ID NO: 16 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 16; (v) the sequence as set forth in SEQ ID NO: 18 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 18; or (vi) the sequence as set forth in SEQ ID NO: 26 or a sequence having

at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 26. Preferably when said sequences comprise an amino acid substitution, said sequences have at least 1 and no more than 4 amino acid substitutions, preferably at least 1 and no more than 3 amino acid substitutions, preferably 1 or 2 amino acid substitutions, preferably 1 amino acid substitution. Preferably at most two, preferably one preferably none of the substitutions are high impact amino acid substitutions (i.e. achieving a score of > 25 in the impact scoring method disclosed in Example 2); preferably at most three, preferably two, preferably one, preferably none of the substitutions are median impact amino acid substitutions (i.e. achieving a score of >10 to 10 in the impact scoring method disclosed in Example 2); preferably at most five, preferably four, preferably three, preferably two, preferably one, preferably none of the substitutions are low impact amino acid substitutions (i.e. achieving a score of >0 to 10 in the impact scoring method disclosed in Example 2); preferably all said substitutions are no impact amino acid substitutions (i.e. achieving a score of 0 in the impact scoring method disclosed in Example 2). Preferably said substitutions do not occur at the positions within said sequences corresponding to positions 226, 228 and 251 of SEQ ID NO: 12. Preferably a dengue antigen comprising said polypeptide leads to a balanced immune response when used in the context of a tetravalent composition. Preferably when a vaccine composition comprising a dengue antigen comprising said polypeptide further comprises a dengue antigen of serotypes 1, 3 and 4 as defined herein, said vaccine composition produces a balanced immune response when administered to a mammal, preferably a human.

Preferably said dengue antigen comprises a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO: 13; SEQ ID NO: 14, SEQ ID NO: 15 SEQ ID NO: 16; SEQ ID NO: 18 and SEQ ID NO: 26.

Preferably said dengue antigen comprises a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO: 13 and SEQ ID NO: 16.

Preferably a dengue antigen (as comprised within a vaccine composition of the invention) comprises a polypeptide comprising: (i) the sequence as set forth in SEQ ID NO: 19 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 19; (ii) the sequence as set forth in SEQ ID NO: 20 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 20; (iii) the sequence as set forth in SEQ ID NO: 21 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 21; (iv) the sequence as set forth in SEQ ID NO: 22 or a sequence having at least 1 and

no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 22; (v) the sequence as set forth in SEQ ID NO: 23 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 23 or (vi) the sequence as set forth in SEQ ID NO: 27 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 27. Preferably when said sequences comprise an amino acid substitution, said sequences have at least 1 and no more than 4 amino acid substitutions, preferably at least 1 and no more than 3 amino acid substitutions, preferably 1 or 2 amino acid substitutions, preferably 1 amino acid substitution. Preferably at most two, preferably one preferably none of the substitutions are high impact amino acid substitutions (i.e. achieving a score of > 25 in the impact scoring method disclosed in Example 2); preferably at most three, preferably two, preferably one, preferably none of the substitutions are median impact amino acid substitutions (i.e. achieving a score of >10 to 25 in the impact scoring method disclosed in Example 2); preferably at most five, preferably four, preferably three, preferably two, preferably one, preferably none of the substitutions are low impact amino acid substitutions (i.e. achieving a score of >0 to 10 in the impact scoring method disclosed in Example 2); preferably all said substitutions are no impact amino acid substitutions (i.e. achieving a score of 0 in the impact scoring method disclosed in Example 2). Preferably said substitutions do not occur at the position within said sequences corresponding to position 125 of SEQ ID NO: 3. Preferably a dengue antigen comprising said polypeptide leads to a balanced immune response when used in the context of a tetravalent composition. Preferably when a vaccine composition comprising a dengue antigen comprising said polypeptide further comprises a dengue antigen of serotypes 1, 3 and 4 as defined herein, said vaccine composition produces a balanced immune response when administered to a mammal, preferably a human.

Preferably a dengue antigen (as comprised within a vaccine composition of the invention) comprises a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO: 19; SEQ ID NO: 20, SEQ ID NO: 21; SEQ ID NO: 22; SEQ ID NO: 23 and SEQ ID NO: 27.

30 Preferably a dengue antigen (as comprised within a vaccine composition of the invention) comprises:

35 i) a polypeptide having the sequence as set forth in SEQ ID NO: 13 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 13; and

a polypeptide having the sequence as set forth in SEQ ID NO: 19 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 19;

- 5 ii) a polypeptide having the sequence as set forth in SEQ ID NO: 14 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 14; and a polypeptide having the sequence as set forth in SEQ ID NO: 20 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 20;
- 10 iii) a polypeptide having the sequence as set forth in SEQ ID NO: 15 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 15; and a polypeptide having the sequence as set forth in SEQ ID NO: 21 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 21;
- 15 iv) a polypeptide having the sequence as set forth in SEQ ID NO: 16 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 16; and a polypeptide having the sequence as set forth in SEQ ID NO: 22 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 22;
- 20 v) a polypeptide having the sequence as set forth in SEQ ID NO: 18 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 18; and a polypeptide having the sequence as set forth in SEQ ID NO: 23 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 23; or
- 25 vi) a polypeptide having the sequence as set forth in SEQ ID NO: 26 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 26; and a polypeptide having the sequence as set forth in SEQ ID NO: 27 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 27.

30 Preferably when said sequences comprise an amino acid substitution, said sequences have at least 1 and no more than 4 amino acid substitutions, preferably at least 1 and no more than 3 amino acid substitutions, preferably 1 or 2 amino acid

substitutions, preferably 1 amino acid substitution. Preferably at most two, preferably one preferably none of the substitutions are high impact amino acid substitutions (i.e. achieving a score of >25 in the impact scoring method disclosed in Example 2); preferably at most three, preferably two, preferably one, preferably none of the substitutions are 5 median impact amino acid substitutions (i.e. achieving a score of >10 to 25 in the impact scoring method disclosed in Example 2); preferably at most five, preferably four, preferably three, preferably two, preferably one, preferably none of the substitutions are low impact amino acid substitutions (i.e. achieving a score of >0 to 10 in the impact scoring method disclosed in Example 2); preferably all said substitutions are no impact 10 amino acid substitutions (i.e. achieving a score of 0 in the impact scoring method disclosed in Example 2). Preferably said substitutions do not occur at the positions within said sequences corresponding to positions 226, 228 and 251 of SEQ ID NO: 12 and the position corresponding to position 125 of SEQ ID NO: 3. Preferably a dengue antigen comprising said polypeptides leads to a balanced immune response when used in the 15 context of a tetravalent composition. Preferably when a vaccine composition comprising a dengue antigen comprising said polypeptides further comprises a dengue antigen of serotypes 1, 3 and 4 as defined herein, said vaccine composition produces a balanced immune response when administered to a mammal, preferably a human.

Preferably a dengue antigen (as comprised within a vaccine composition of the 20 invention) comprises: i) a polypeptide of SEQ ID NO: 13 and a polypeptide of SEQ ID NO: 19; ii) a polypeptide of SEQ ID NO: 14 and a polypeptide of SEQ ID NO: 20; iii) a polypeptide of SEQ ID NO: 15 and a polypeptide of SEQ ID NO: 21; iv) a polypeptide of SEQ ID NO: 16 and a polypeptide of SEQ ID NO: 22; v) a polypeptide of SEQ ID NO: 18 and a polypeptide of SEQ ID NO: 23 or vi) a polypeptide of SEQ ID NO: 26 and a 25 polypeptide of SEQ ID NO: 27.

Preferably said dengue antigen comprises a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO: 8; SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11.

Preferably said dengue antigen comprises a polypeptide comprising a sequence 30 selected from the group consisting of SEQ ID NO: 8 and SEQ ID NO: 11.

Preferably a composition of the present invention comprises a dengue antigen selected from the group consisting of: (a) a live attenuated dengue virus; (b) an inactivated dengue virus; (c) a live attenuated or inactivated chimeric dengue virus; or (d) a combination of two or more of (a) to (c); wherein said dengue antigen comprises a 35 nucleotide sequence encoding a polypeptide comprising a polypeptide as defined herein.

The present invention is also directed to a vaccine composition comprising a dengue antigen of serotype 2 selected from the group consisting of: (a) a live attenuated dengue virus; (b) an inactivated dengue virus; (c) a live attenuated or inactivated chimeric dengue virus; or (d) a combination of two or more of (a) to (c); wherein said dengue antigen comprises a nucleotide sequence having at least 90% sequence identity to a sequence selected from the group consisting of the RNA equivalent of SEQ ID NO: 1, the RNA equivalent of SEQ ID NO: 4, the RNA equivalent of SEQ ID NO: 5, the RNA equivalent of SEQ ID NO: 6, the RNA equivalent of SEQ ID NO: 7 and SEQ ID NO: 25. References to at least 90% sequence identity to the RNA equivalent of SEQ ID NO: 1, the RNA equivalent of SEQ ID NO: 4, the RNA equivalent of SEQ ID NO: 5, the RNA equivalent of SEQ ID NO: 6, the RNA equivalent of SEQ ID NO: 7 or SEQ ID NO: 25 may preferably be read herein as at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, at least 99.5% or 100% sequence identity. When nucleotide sequences of this embodiment of the invention encode polypeptides comprising one or more amino acid substitutions with respect to the polypeptides encoded by SEQ ID NO: 1, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and SEQ ID NO: 25, preferably at most two, preferably one preferably none of the substitutions are high impact amino acid substitutions (i.e. achieving a score of >25 in the impact scoring method disclosed in Example 2); preferably at most three, preferably two, preferably one, preferably none of the substitutions are median impact amino acid substitutions (i.e. achieving a score of >10 to 25 in the impact scoring method disclosed in Example 2); preferably at most five, preferably four, preferably three, preferably two, preferably one, preferably none of the substitutions are low impact amino acid substitutions (i.e. achieving a score of >0 to 10 in the impact scoring method disclosed in Example 2); preferably all said substitutions are no impact amino acid substitutions (i.e. achieving a score of 0 in the impact scoring method disclosed in Example 2). Preferably said substitutions do not occur at the positions within said polypeptides corresponding to positions 226, 228 and 251 of SEQ ID NO: 12 and the positions within said polypeptides corresponding to positions 24 and 125 of SEQ ID NO: 3. Preferably the vaccine compositions comprising a dengue antigen of serotype 2 of this embodiment of the invention lead to a balanced immune response when used in the context of a tetravalent composition. Preferably a dengue antigen of serotype 2 according to this embodiment of the invention further comprises a dengue antigen of serotype 1, a dengue antigen of serotype 3 and a dengue antigen of serotype 4 as described elsewhere herein. Preferably when a vaccine composition comprising a dengue antigen of serotype 2 according to this embodiment of the invention further comprises a dengue antigen of serotypes 1, 3 and 4

as defined herein, said vaccine composition produces a balanced immune response when administered to a mammal, preferably a human. When a vaccine composition of the present invention comprises a dengue antigen of serotype 2 which comprises a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 7, said 5 vaccine composition is preferably either: (i) a vaccine composition comprising a dengue antigen comprising a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 7, wherein said dengue antigen is not CYD-MD or (ii) a vaccine composition comprising a dengue antigen which is CYD-MD. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising 10 a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 7, wherein said vaccine composition does not comprise prM-E sequences from MD1280 or (ii) a vaccine composition comprising prM-E sequences from MD1280. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a nucleotide sequence having at least 90% identity to the RNA 15 equivalent of SEQ ID NO: 7, wherein said vaccine composition does not comprise a dengue antigen comprising the M and E sequences of CYD-MD or (ii) a vaccine composition comprising a dengue antigen comprising the M and E sequences of CYD-MD. Said vaccine composition is also preferably either (i) a vaccine composition comprising a dengue antigen comprising a nucleotide sequence having at least 90% 20 identity to the RNA equivalent of SEQ ID NO: 7, wherein said vaccine composition does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequences of MD1280 (SEQ ID NO: 11) or (ii) a vaccine composition comprising a chimeric dengue virus of serotype 2 generated using the prM-E sequences of MD1280 (SEQ ID NO: 11). Said vaccine composition is also preferably either (i) a vaccine composition comprising a 25 dengue antigen comprising a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 7, wherein said vaccine composition does not comprise a dengue antigen comprising the prM and E sequences of CYD-MD or (ii) a vaccine composition comprising a dengue antigen comprising the prM and E sequences of CYD-MD. Said vaccine composition is also preferably either (i) a vaccine composition comprising a dengue antigen comprising a nucleotide sequence having at least 90% 30 identity to the RNA equivalent of SEQ ID NO: 7, wherein said vaccine composition does not comprise a dengue antigen comprising a polypeptide of SEQ ID NO: 16 and a polypeptide of SEQ ID NO: 22 (or a dengue antigen comprising a nucleotide sequence encoding a protein comprising said polypeptides) or (ii) a vaccine composition comprising a dengue antigen comprising a polypeptide of SEQ ID NO: 16 and a polypeptide of SEQ 35 ID NO: 22 (or a dengue antigen comprising a nucleotide sequence encoding a protein

comprising said polypeptides). Preferably, a vaccine composition of the present invention which comprises a dengue antigen of serotype 2 which comprises a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 7 does not comprise: (i) a chimeric virus comprising the prM and E amino acid sequences of MD1280 or (ii) a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-MD (SEQ ID NO: 11). When a vaccine composition of the present invention comprises a dengue antigen of serotype 2 which comprises a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 4, said vaccine composition is preferably either: (i) a vaccine composition comprising a dengue antigen comprising a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 4, wherein said dengue antigen is not CYD-LAV or (ii) a vaccine composition comprising a dengue antigen which is CYD-LAV. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 4, wherein said vaccine composition does not comprise prM-E sequences from LAV2 or (ii) a vaccine composition comprising prM-E sequences from LAV2. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 4, wherein said vaccine composition does not comprise a dengue antigen comprising the M and E sequences of CYD-LAV or (ii) a vaccine composition comprising a dengue antigen comprising the M and E sequences of CYD-LAV. Said vaccine composition is also preferably either (i) a vaccine composition comprising a dengue antigen comprising a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 4, wherein said vaccine composition does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequences of LAV2 (SEQ ID NO: 8) or (ii) a vaccine composition comprising a chimeric dengue virus of serotype 2 generated using the prM-E sequences of LAV2 (SEQ ID NO: 8). Said vaccine composition is also preferably either (i) a vaccine composition comprising a dengue antigen comprising a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 4, wherein said vaccine composition does not comprise a dengue antigen comprising the prM and E sequences of CYD-LAV or (ii) a vaccine composition comprising a dengue antigen comprising the prM and E sequences of CYD-LAV. Said vaccine composition is also preferably either (i) a vaccine composition comprising a dengue antigen comprising a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 4, wherein said vaccine composition does not comprise a dengue antigen comprising a polypeptide of SEQ ID NO: 13 and a polypeptide of SEQ ID NO: 19 (or a dengue antigen comprising a nucleotide sequence

encoding a protein comprising said polypeptides) or (ii) a vaccine composition comprising a dengue antigen comprising a polypeptide of SEQ ID NO: 13 and a polypeptide of SEQ ID NO: 19 (or a dengue antigen comprising a nucleotide sequence encoding a protein comprising said polypeptides). Preferably, a vaccine composition of the present invention which comprises a dengue antigen of serotype 2 which comprises a nucleotide sequence having at least 90% identity to the RNA equivalent of SEQ ID NO: 4 does not comprise: (i) a chimeric virus comprising the prM and E amino acid sequences of LAV2 or (ii) a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-LAV (SEQ ID NO: 8).

The present invention is also directed to a vaccine composition comprising a dengue antigen of serotype 2 selected from the group consisting of: (a) a live attenuated dengue virus; (b) an inactivated dengue virus; (c) a live attenuated or inactivated chimeric dengue virus; or (d) a combination of two or more of (a) to (c); wherein said dengue antigen comprises a nucleotide sequence having at least 1 and no more than 20 nucleotide substitutions with respect to a sequence selected from the group consisting of the RNA equivalent of SEQ ID NO: 1, the RNA equivalent of SEQ ID NO: 4, the RNA equivalent of SEQ ID NO: 5, the RNA equivalent of SEQ ID NO: 6, the RNA equivalent of SEQ ID NO: 7 and SEQ ID NO: 25. When nucleotide sequences of this embodiment of the invention encode polypeptides comprising one or more amino acid substitutions with respect to the polypeptides encoded by SEQ ID NO: 1, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and SEQ ID NO: 25, preferably at most two, preferably one preferably none of the substitutions are high impact amino acid substitutions (i.e. achieving a score of >25 in the impact scoring method disclosed in Example 2); preferably at most three, preferably two, preferably one, preferably none of the substitutions are median impact amino acid substitutions (i.e. achieving a score of >10 to 25 in the impact scoring method disclosed in Example 2); preferably at most five, preferably four, preferably three, preferably two, preferably one, preferably none of the substitutions are low impact amino acid substitutions (i.e. achieving a score of >0 to 10 in the impact scoring method disclosed in Example 2); preferably all said substitutions are no impact amino acid substitutions (i.e. achieving a score of 0 in the impact scoring method disclosed in Example 2). Preferably said substitutions do not occur at the positions within said polypeptides corresponding to positions 226, 228 and 251 of SEQ ID NO: 12 and the positions within said polypeptides corresponding to positions 24 and 125 of SEQ ID NO: 3. Preferably the vaccine compositions comprising a dengue antigen of serotype 2 of this embodiment of the invention lead to a balanced immune response when used in the context of a tetravalent composition. Preferably a dengue antigen of serotype 2 according to this embodiment of the invention further comprises a dengue antigen of serotype 1, a

dengue antigen of serotype 3 and a dengue antigen of serotype 4 as described elsewhere herein. Preferably when a vaccine composition comprising a dengue antigen of serotype 2 according to this embodiment of the invention further comprises a dengue antigen of serotypes 1, 3 and 4 as defined herein, said vaccine composition produces a balanced 5 immune response when administered to a mammal, preferably a human.

When a vaccine composition of the present invention comprises a dengue antigen of serotype 2 as defined herein which comprises a polypeptide having at least 90%, at least 92%, at least 94%, at least 96%, at least 98%, at least 99%, at least 99.5% or 100% identity to SEQ ID NO : 12, wherein said polypeptide comprises an isoleucine residue at the position within the polypeptide that corresponds to position 485 of SEQ ID

NO : 12 or an alanine residue at the position within the polypeptide that corresponds to position 491 of SEQ ID NO : 12, said vaccine composition preferably (i) is not CYD-MD; (ii) does not comprise the prM-E sequence from MD1280; (iii) does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequences of MD1280

(SEQ ID NO: 11); (iv) does not comprise a dengue antigen comprising the prM and E sequences of CYD-MD; (v) does not comprise a dengue antigen comprising a polypeptide of SEQ ID NO: 16 and a polypeptide of SEQ ID NO: 22 (or a dengue antigen comprising a nucleotide sequence encoding a protein comprising said polypeptides); (vi) does not comprise a chimeric virus comprising the prM and E amino acid sequences of MD1280;

(vii) does not comprise a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-MD (SEQ ID NO: 11) and/or (viii) does not comprise a dengue antigen of serotype 2 which comprises the M and E sequences of CYD-MD. When a vaccine composition of the present invention comprises a dengue antigen of serotype 2 as defined herein which comprises a polypeptide having at least 90%, at least 92%, at least 94%, at

least 96%, at least 98%, at least 99%, at least 99.5% or 100% identity to SEQ ID NO : 12, wherein said polypeptide comprises a methionine residue at the position within the polypeptide that corresponds to position 6 of SEQ ID NO : 12 or a threonine residue at the position within the polypeptide that corresponds to position 478 of SEQ ID NO : 12, said vaccine composition preferably (i) is not CYD-LAV; (ii) does not comprise the prM-E sequence from LAV2; (iii) does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequences of LAV2 (SEQ ID NO: 8); (iv) does not comprise a dengue antigen comprising the prM and E sequences of CYD-LAV; (v) does not comprise a dengue antigen comprising a polypeptide of SEQ ID NO: 13 and a polypeptide of SEQ

ID NO: 19 (or a dengue antigen comprising a nucleotide sequence encoding a protein comprising said polypeptides); (vi) does not comprise a chimeric virus comprising the prM and E amino acid sequences of LAV2 ; (vii) does not comprise a dengue antigen of

30

35

serotype 2 which comprises the prM-E sequence of CYD-LAV (SEQ ID NO: 8) and/or (viii) does not comprise a dengue antigen of serotype 2 which comprises the M and E sequences of CYD-LAV.

Preferably said dengue antigen comprises a polypeptide which comprises no more than 1, no more than 2, no more than 3, no more than 4, no more than 5, nor more than 6, no more than 7, no more than 8, no more than 9, no more than 10, no more than 11 or no more than 12 minor amino acid residues, wherein a minor amino acid residue at a given position of a prM-E or E sequence is defined as an amino acid that appears in less than 15% of dengue virus prM-E or E sequences of serotype 2 at that position.

Preferably the dengue disease according to the present invention is virologically-confirmed dengue disease.

Preferably a human subject according to the present invention is less than 18 years of age or less than 12 years of age. For example, a human subject according to the present invention may be 0-17 years of age, 0-11 years of age, 4-17 years of age, 4-11 years of age, 4-6 years of age, 6-8 years of age, 8-10 years of age, 2-8 years of age, 2-11 years of age, 2-14 years of age, 9-16 years of age, 12-17 years of age or 18-45 years of age. More preferably, a human subject according to the present invention is 4-11 years of age, 2-14 years of age or 9-16 years of age. A human subject according to the present invention may be at least 9 months old or less than 9 months old. For instance a human subject according to the present invention may be 9 months to 16 years of age, 9 months to 14 years of age, 9 months to 11 years of age or 9 months to 8 years of age. A human subject according to the present invention may be at least 9 months old, with no history of severe allergy to any component of the vaccine composition as defined herein, no congenital or acquired immune deficiency, no symptomatic HIV infection and said subject should not be pregnant or breast feeding.

A human subject to which a vaccine composition of the present invention is to be administered is preferably a person at risk of infection, for instance a person travelling in an area where dengue fever is present, i.e. a dengue endemic area, or a resident of such an area. Preferably a human subject of the present invention resides in a dengue endemic area. Dengue endemic areas according to the present invention include most of the tropics and sub-tropics, for instance any country identified as an endemic country by the WHO. For instance, a dengue endemic area according to the present invention may comprise those American countries or parts thereof which fall within the tropics and sub-tropics. A dengue endemic area according to the present invention may thus comprise any one or more of the following: Brazil, Venezuela, Colombia, Ecuador, Peru, Bolivia, Paraguay, Panama, Costa Rica, Nicaragua, Honduras, El Salvador, Guatemala, Belize,

Mexico, the USA and the islands of the Caribbean. In a particular embodiment, a dengue endemic area of the present invention may consist of the following: Brazil, Colombia, Honduras, Mexico and Puerto Rico. A dengue endemic area according to the present invention may also include south Asian and Oceania countries within the tropics and sub-tropics. A dengue endemic area according to the present invention may thus consist of any one or more of the following: India, Myanmar (Burma), Thailand, Laos, Vietnam, Cambodia, Indonesia, Malaysia, Singapore, the Philippines, Taiwan, Papua New Guinea and Australia. In a dengue endemic area according to the present invention, a particular serotype, strain or genotype of wild type dengue virus may be the dominant circulating strain. For example, a dengue virus of serotype 2 may be characterised as having an Asian I or an Asian/American genotype. Asian/American genotype strains are characterised by at least one of, at least two of, at least three of, at least four of, at least five of or all six of the following residues Arg, Asn, Asp, Thr, Gly and His at positions prM-16, E-83, E-203, E-226, E-228 and E-346 respectively (wherein prM-16 designates position 16 of the prM protein and E-83 etc. designates position 83 of the E protein). Asian I genotype strains are characterised by at least one of, at least two of, at least three of, at least four of, at least five of or all six of the following residues Ile, Lys, Asn, Arg, Glu and Tyr at positions prM-16, E-83, E-203, E-226, E-228 and E-346 respectively (see Table 1 of Hang et al., PLoS NTD, 4(7): e757). A preferred dengue endemic area according to the present invention is one in which a dengue virus having an Asian/American genotype is the dominant circulating strain, i.e. at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or 100% of the cases of dengue disease in said dengue endemic area are caused by dengue virus having an Asian/American genotype. A preferred dengue endemic area according to the present invention is one in which a dengue virus of any one or more of serotypes 1, 3 or 4 is/are the dominant circulating serotype(s), i.e. at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or 100% of the cases of dengue disease are caused by dengue virus of serotypes 1, 3 or 4.

A vaccine composition of the present invention may be administered to a flavivirus immune subject, for example a dengue-immune subject, or a vaccine composition of the present invention may be administered to a flavivirus-naïve subject. Advantageously, a vaccine composition of the present invention is administered to a flavivirus-immune subject, for example a dengue-immune subject.

Preferably, a composition according to the present invention, e.g. a composition for use in a method according to the present invention, reduces the likelihood or severity of DHF. A reduction in the likelihood of DHF (i.e. a reduction in the probability of contracting

DHF) may be measured by comparing the number of cases of DHF in a group of subjects who have received a vaccine composition according to the present invention and the number of cases of DHF in a control group of subjects who have not received a vaccine composition according to the present invention. A reduction in the severity of DHF may be 5 determined by calculating the number of subjects displaying DHF of each of Grades I, II, III or IV in a group of subjects who have received a vaccine composition according to the present invention and comparing those numbers to the equivalent numbers from a control group of subjects who have not received a vaccine composition according to the present invention. For instance, a composition for use in a method according to the present 10 invention preferably reduces the number of cases of Grade I DHF, the number of cases of Grade II DHF, the number of cases of Grade III DHF and/or the number of cases of Grade IV DHF in those subjects receiving the vaccine, when compared to the equivalent number of cases Grade I DHF, Grade II DHF, Grade III DHF and Grade IV DHF occurring in a control group of subjects who have not received a vaccine composition according to the 15 present invention.

Preferably, a composition according to the present invention, e.g. a composition for use in a method according to the present invention, reduces the incidence or likelihood of symptomatic virologically-confirmed dengue disease. Advantageously, a composition according to the present invention, e.g. a composition for use in a method according to the 20 present invention, reduces the incidence or likelihood of symptomatic virologically-confirmed dengue disease caused by dengue virus of serotypes 1, 3 or 4. Advantageously, a composition according to the present invention, e.g. a composition for use in a method according to the present invention, reduces the incidence or likelihood of symptomatic virologically-confirmed dengue disease caused by dengue virus of serotypes 25 1, 2, 3 or 4. Preferably, a composition according to the present invention, e.g. a composition for use in a method according to the present invention, reduces the rate of hospitalization due to virologically-confirmed dengue disease, i.e. reduces the incidence of hospitalized virologically-confirmed dengue disease. For instance, a composition according to the present invention, e.g. a composition for use in a method according to the 30 present invention, reduces the rate of hospitalization due to virologically-confirmed dengue disease caused by dengue virus of serotypes 1, 3 or 4, i.e. reduces the incidence of hospitalized virologically-confirmed dengue disease caused by dengue virus of serotypes 1, 3 or 4.

Preferably, a composition according to the present invention, e.g. a composition for use in a method according to the present invention, reduces the incidence or likelihood 35 of dengue disease. Advantageously, a composition according to the present invention,

e.g. a composition for use in a method according to the present invention, reduces the incidence or likelihood of dengue disease caused by dengue virus of serotypes 1, 3 or 4. Advantageously, a composition according to the present invention, e.g. a composition for use in a method according to the present invention, reduces the incidence or likelihood of 5 dengue disease caused by dengue virus of serotypes 1, 2, 3 or 4. Preferably, a composition according to the present invention, e.g. a composition for use in a method according to the present invention, reduces the rate of hospitalization due to dengue disease, i.e. reduces the incidence of hospitalized dengue disease. For instance, a composition according to the present invention, e.g. a composition for use in a method 10 according to the present invention, reduces the rate of hospitalization due to dengue disease caused by dengue virus of serotypes 1, 3 or 4, i.e. reduces the incidence of hospitalized dengue disease caused by dengue virus of serotypes 1, 3 or 4.

A vaccine composition according to the present invention may be administered in multiple doses. Doses of a vaccine composition according to the present invention may be 15 administered in an initial vaccination regimen followed by booster vaccinations. For example, a vaccine composition according to the present invention may be administered in one, two or three doses or more than three doses, e.g. four doses. Preferably, the first dose and the third dose are to be administered approximately twelve months apart. For example, an initial vaccination regimen according to the present invention is administered 20 in three doses, wherein the first and third doses of said vaccination regimen are to be administered approximately twelve months apart. Advantageously, a vaccine composition according to the present invention is to be administered in a first dose, a second dose and a third dose. In such an embodiment, said first dose and said third dose may be administered approximately twelve months apart. For instance, a vaccine composition of 25 the present invention may be administered in a first dose, a second dose and a third dose, wherein said second dose is to be administered about six months after said first dose and wherein said third dose is to be administered about twelve months after said first dose. Alternatively, the three doses may be administered at zero months, at about three to four months (e.g. at about three-and-a-half months) and at about twelve months (i.e. a regimen 30 wherein the second dose of the composition is administered at about three-and-a-half months after the first dose, and wherein the third dose of the composition is administered at about twelve months after the first dose).

A vaccine composition according to the present invention may be administered in two doses. Preferably, the first dose and the second dose are to be administered 35 approximately about six to twelve months after the first dose months apart. Preferably, the

second dose is to be administered at eight months after the first dose. Preferably the second dose is administered at about eight-and-a-half to nine months after the first dose.

A vaccine composition according to the present invention may be administered in a single dose.

5 Dengue disease, as defined herein, may be caused by any one of two serotypes of a dengue virus. For example, dengue disease is preferably caused by a dengue virus of serotype 1 or serotype 3, a dengue virus of serotype 1 or serotype 4, a dengue virus of serotype 3 or serotype 4, a dengue virus of serotype 1 or serotype 2, a dengue virus of serotype 2 or serotype 3, a dengue virus of serotype 2 or serotype 4. Dengue disease, as
10 defined herein, is preferably caused by any one of three serotypes of a dengue virus. For example, dengue disease is preferably caused by a dengue virus of serotype 1, 2 or 3, a dengue virus of serotype 1, 3 or 4, a dengue virus of serotype 1, 2 or 4, a dengue virus of serotype 2, 3 or 4. In another embodiment, dengue disease is caused by a dengue virus of serotype 1, a dengue virus of serotype 2, a dengue virus of serotype 3 or a dengue
15 virus of serotype 4.

A vaccine composition according to the present invention, e.g. for use in a method according to the present invention preferably comprises a dengue antigen of serotype 1, a dengue antigen of serotype 2, a dengue antigen of serotype 3 and a dengue antigen of serotype 4. Such a composition may be described herein as a tetravalent composition.
20 For instance, a composition of the present invention, e.g. for use in a method of protecting according to the present invention, may advantageously comprise any one of the following combinations of dengue antigens of serotypes 1, 2, 3 and 4: i) a dengue antigen comprising the prM and E sequences of CYD-1, a dengue antigen comprising the prM and E sequences of CYD-LAV, a chimeric dengue virus comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4; ii) a dengue antigen comprising the prM and E sequences of CYD-1, a dengue antigen comprising the prM and E sequences of CYD-BID, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4; (iii) a dengue antigen comprising the prM and E sequences of CYD-1, a dengue
25 antigen comprising the prM and E sequences of CYD-PR, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4; (iv) a dengue antigen comprising the prM and E sequences of CYD-1, a dengue antigen comprising the prM and E sequences of CYD-MD, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4;. For instance, a composition of the present invention may
30 also advantageously comprise any one of the following combinations of dengue antigens:
35

i) CYD-1, CYD-LAV, CYD-3 and CYD-4; ii) CYD-1, CYD-BID, CYD-3 and CYD-4; (iii) CYD-1, CYD-PR, CYD-3 and CYD-4 or (iv) CYD-1, CYD-MD, CYD-3 and CYD-4. A composition of the present invention may also advantageously comprise the following combination of dengue antigens: i) a dengue antigen comprising the prM and E sequences of CYD-1, VDV2, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4. For instance, a composition of the present invention may advantageously comprise CYD-1, VDV-2, CYD-3 and CYD-4. A composition of the present invention, as described herein, may advantageously comprise a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2. A composition of the present invention, as described herein, may advantageously comprise a dengue antigen of serotype 2 which comprises a sequence having at least 90% identity to the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2. For example, said sequence may be at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2.

A vaccine composition according to the present invention, e.g. for use in a method according to the present invention, preferably comprises a dengue antigen of serotype 1, a dengue antigen of serotype 2, a dengue antigen of serotype 3 and a dengue antigen of serotype 4. Such a composition may be described herein as a tetravalent composition. For instance, a composition of the present invention, e.g. for use in a method of protecting according to the present invention, may advantageously comprise any one of the following combinations of dengue antigens of serotypes 1, 2, 3 and 4: i) a dengue antigen comprising the M and E sequences of CYD-1, a dengue antigen comprising the M and E sequences of CYD-LAV, a chimeric dengue virus comprising the M and E sequences of CYD-3 and a dengue antigen comprising the M and E sequences of CYD-4; ii) a dengue antigen comprising the M and E sequences of CYD-1, a dengue antigen comprising the M and E sequences of CYD-BID, a dengue antigen comprising the M and E sequences of CYD-3 and a dengue antigen comprising the M and E sequences of CYD-4; (iii) a dengue antigen comprising the M and E sequences of CYD-1, a dengue antigen comprising the M and E sequences of CYD-PR, a dengue antigen comprising the M and E sequences of CYD-3 and a dengue antigen comprising the M and E sequences of CYD-4; (iv) a dengue antigen comprising the M and E sequences of CYD-1, a dengue antigen comprising the M and E sequences of CYD-MD, a dengue antigen comprising the M and E sequences of

CYD-3 and a dengue antigen comprising the M and E sequences of CYD-4;. For instance, a composition of the present invention may also advantageously comprise any one of the following combinations of dengue antigens: i) CYD-1, CYD-LAV, CYD-3 and CYD-4; ii)

5 CYD-1, CYD-BID, CYD-3 and CYD-4; (iii) CYD-1, CYD-PR, CYD-3 and CYD-4 or (iv)

CYD-1, CYD-MD, CYD-3 and CYD-4. A composition of the present invention may also advantageously comprise the following combination of dengue antigens: i) a dengue antigen comprising the M and E sequences of CYD-1, VDV2, a dengue antigen comprising the M and E sequences of CYD-3 and a dengue antigen comprising the M and E sequences of CYD-4. For instance, a composition of the present invention may

10 advantageously comprise CYD-1, VDV-2, CYD-3 and CYD-4. A composition of the present invention, as described herein, may advantageously comprise a dengue antigen of serotype 2 which comprises the E sequence of CYD-LAV (SEQ ID NO: 13), CYD-BID (SEQ ID NO: 14), CYD-PR (SEQ ID NO: 15) CYD-MD (SEQ ID NO: 16) or SEQ ID NO:

15 18. In certain embodiments, a composition of the present invention which comprises a

dengue antigen comprising the sequence as set forth in SEQ ID NO: 18 is not a vaccine composition of serotype 2 comprising the prM-E sequence from SEQ ID NO: 2. In certain

embodiments, a composition of the present invention is either a composition comprising a dengue antigen comprising the sequence as set forth in SEQ ID NO: 18, wherein said

composition is not a vaccine composition of the present invention comprising chimeric

20 dengue virus of serotype 2 generated using the prM-E sequence of SEQ ID NO: 2, or it is a composition comprising chimeric dengue virus of serotype 2 generated using the prM-E sequence of SEQ ID NO: 2. A composition of the present invention, as described herein,

may advantageously comprise a dengue antigen of serotype 2 which comprises a sequence having at least 90% identity to the E sequence of CYD-LAV (SEQ ID NO: 13),

25 CYD-BID (SEQ ID NO: 14), CYD-PR (SEQ ID NO: 15) CYD-MD (SEQ ID NO: 16) or SEQ ID NO: 18. For example, said sequence may be at least 91%, at least 92%, at least 93%,

at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to the E sequence of CYD-LAV (SEQ ID NO: 13), CYD-BID (SEQ ID NO: 14),

CYD-PR (SEQ ID NO: 15) CYD-MD (SEQ ID NO: 16) or SEQ ID NO: 18.

30 A composition of the present invention, as described herein, (e.g. a tetravalent formulation, e.g. for use in a method of the present invention), may advantageously comprise a dengue antigen of serotype 2 which comprises a polypeptide selected from the group consisting of SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22 or SEQ ID NO: 23. When a vaccine composition of the present invention comprises a

35 dengue antigen of serotype 2 which comprises a polypeptide having the sequence of SEQ ID NO: 19, said vaccine composition is preferably either: (i) a vaccine composition

comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 19, wherein said vaccine composition does not comprise CYD-LAV or (ii) a vaccine composition comprising CYD-LAV. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 19, wherein said vaccine composition does not comprise a dengue antigen comprising the M and E sequences of CYD-LAV or (ii) a vaccine composition comprising a dengue antigen comprising the M and E sequences of CYD-LAV. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 19, wherein said vaccine composition does not comprise a dengue antigen comprising the prM and E sequences of CYD-LAV or (ii) a vaccine composition comprising a dengue antigen comprising the prM and E sequences of CYD-LAV. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 19, wherein said vaccine composition does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequence of LAV-2 or (ii) a vaccine composition comprising a chimeric dengue virus of serotype 2 generated using the prM-E sequence of LAV-2 (SEQ ID NO: 8). Preferably, a vaccine composition of the present invention which comprises a polypeptide having the sequence of SEQ ID NO: 19 does not comprise: (i) a chimeric virus comprising the prM and E amino acid sequences of LAV-2; (ii) a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-LAV (SEQ ID NO: 8) or (iii) a dengue antigen comprising the prM-E sequence from LAV-2. When a vaccine composition of the present invention comprises a dengue antigen of serotype 2 which comprises a polypeptide having the sequence of SEQ ID NO: 21, said vaccine composition is preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 21, wherein said vaccine composition does not comprise CYD-PR or (ii) a vaccine composition comprising CYD-PR. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 21, wherein said vaccine composition does not comprise a dengue antigen comprising the prM and E sequences of CYD-PR or (ii) a vaccine composition comprising a dengue antigen comprising the prM and E sequences of CYD-PR. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 21, wherein said vaccine composition does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequence of PR/DB023 (SEQ ID NO: 10) or (ii) a vaccine composition

comprising a chimeric dengue virus of serotype 2 generated using the prM-E sequence of PR/DB023 (SEQ ID NO: 10). Preferably, a vaccine composition of the present invention which comprises a polypeptide having the sequence of SEQ ID NO: 21 does not comprise: (i) a chimeric virus comprising the prM and E amino acid sequences of PR/DB023; (ii) a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-PR (SEQ ID NO: 10) or (iii) a dengue antigen comprising the prM-E sequence from PR/DB023. When a vaccine composition of the present invention comprises a dengue antigen of serotype 2 which comprises a polypeptide having the sequence of SEQ ID NO: 22, said vaccine composition is preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 22, wherein said vaccine composition does not comprise CYD-MD or (ii) a vaccine composition comprising CYD-MD. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 22, wherein said vaccine composition does not comprise a dengue antigen comprising the prM and E sequences of CYD-MD or (ii) a vaccine composition comprising a dengue antigen comprising the prM and E sequences of CYD-MD. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 22, wherein said vaccine composition does not comprise a dengue antigen comprising the M and E sequences of CYD-MD or (ii) a vaccine composition comprising a dengue antigen comprising the M and E sequences of CYD-MD. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 22, wherein said vaccine composition does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequence of MD1280 (SEQ ID NO: 11) or (ii) a vaccine composition comprising a chimeric dengue virus of serotype 2 generated using the prM-E sequence of MD1280 (SEQ ID NO: 11). Preferably, a vaccine composition of the present invention which comprises a polypeptide having the sequence of SEQ ID NO: 22 does not comprise: (i) a chimeric virus comprising the prM and E amino acid sequences of MD1280; (ii) a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-MD (SEQ ID NO: 11) or (iii) a dengue antigen comprising the prM-E sequence from MD1280. When a vaccine composition of the present invention comprises a dengue antigen of serotype 2 which comprises a polypeptide having the sequence of SEQ ID NO: 23, said vaccine composition is preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 23, wherein said vaccine composition does not comprise a chimeric dengue virus of serotype

2 generated using the prM-E sequence SEQ ID NO: 2 or (ii) a vaccine composition comprising a chimeric dengue virus of serotype 2 generated using the prM-E sequence of SEQ ID NO: 2. Preferably, a vaccine composition of the present invention which comprises a polypeptide having the sequence of SEQ ID NO: 23 does not comprise: (i) a dengue antigen of serotype 2 which comprises SEQ ID NO: 2 or (ii) a dengue antigen comprising the prM-E sequence from SEQ ID NO: 2.

Preferably a dengue antigen of serotype 2 of the present invention further comprises a polypeptide selected from the group consisting of SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16 or SEQ ID NO: 18. For instance, said dengue

10 antigen of serotype 2 preferably comprises: i) a polypeptide of SEQ ID NO: 13 and a polypeptide of SEQ ID NO: 19; ii) a polypeptide of SEQ ID NO: 14 and a polypeptide of SEQ ID NO: 20; iii) a polypeptide of SEQ ID NO: 15 and a polypeptide of SEQ ID NO: 21;

15 iv) a polypeptide of SEQ ID NO: 16 and a polypeptide of SEQ ID NO: 22; or v) a polypeptide of SEQ ID NO: 18 and a polypeptide of SEQ ID NO: 23. When a vaccine

composition of the present invention comprises a dengue antigen of serotype 2 which comprises a polypeptide having the sequence of SEQ ID NO: 13 and a polypeptide having the sequence of SEQ ID NO: 19, said vaccine composition is preferably either: (i) a

20 vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 13 and a polypeptide having the sequence of SEQ ID NO: 19, wherein said vaccine composition does not comprise CYD-LAV or (ii) a vaccine

composition comprising CYD-LAV. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 13 and a polypeptide having the sequence of SEQ ID NO: 19, wherein said vaccine composition does not comprise a dengue antigen comprising the

25 prM and E sequences of CYD-LAV or (ii) a vaccine composition comprising a dengue antigen comprising the prM and E sequences of CYD-LAV. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 13 and a polypeptide having the sequence of SEQ ID NO: 19, wherein said vaccine composition does not comprise a

30 dengue antigen comprising the M and E sequences of CYD-LAV or (ii) a vaccine composition comprising a dengue antigen comprising the M and E sequences of CYD-LAV. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 13 and a polypeptide having the sequence of SEQ ID NO: 19, wherein said vaccine

35 composition does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequence of LAV-2 or (ii) a vaccine composition comprising a chimeric dengue

virus of serotype 2 generated using the prM-E sequence of LAV-2 (SEQ ID NO: 8). Preferably, a vaccine composition of the present invention which comprises a polypeptide having the sequence of SEQ ID NO: 13 and a polypeptide having the sequence of SEQ ID NO: 19 does not comprise: (i) a chimeric virus comprising the prM and E amino acid sequences of LAV-2; (ii) a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-LAV (SEQ ID NO: 8) or (iii) a dengue antigen comprising the prM-E sequence from LAV-2. When a vaccine composition of the present invention comprises a dengue antigen of serotype 2 which comprises a polypeptide having the sequence of SEQ ID NO: 15 and a polypeptide having the sequence of SEQ ID NO: 21, said vaccine composition is preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 15 and a polypeptide having the sequence of SEQ ID NO: 21, wherein said vaccine composition does not comprise CYD-PR or (ii) a vaccine composition comprising CYD-PR. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen a polypeptide having the sequence of SEQ ID NO: 15 and comprising a polypeptide having the sequence of SEQ ID NO: 21, wherein said vaccine composition does not comprise a dengue antigen comprising the prM and E sequences of CYD-PR or (ii) a vaccine composition comprising a dengue antigen comprising the prM and E sequences of CYD-PR. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 15 and a polypeptide having the sequence of SEQ ID NO: 21, wherein said vaccine composition does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequence of PR/DB023 (SEQ ID NO: 10) or (ii) a vaccine composition comprising a chimeric dengue virus of serotype 2 generated using the prM-E sequence of PR/DB023 (SEQ ID NO: 10). Preferably, a vaccine composition of the present invention which comprises a polypeptide having the sequence of SEQ ID NO: 15 and a polypeptide having the sequence of SEQ ID NO: 21 does not comprise: (i) a chimeric virus comprising the prM and E amino acid sequences of PR/DB023; (ii) a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-PR (SEQ ID NO: 10) or (iii) a dengue antigen comprising the prM-E sequence from PR/DB023. When a vaccine composition of the present invention comprises a dengue antigen of serotype 2 which comprises a polypeptide having the sequence of SEQ ID NO: 16 and a polypeptide having the sequence of SEQ ID NO: 22, said vaccine composition is preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 16 and a polypeptide having the sequence of SEQ ID NO: 22, wherein said vaccine composition does not comprise CYD-MD or (ii) a vaccine

composition comprising CYD-MD. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a polypeptide having the sequence of SEQ ID NO: 16 and a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 22, wherein said vaccine composition does not comprise a dengue antigen comprising the prM and E sequences of CYD-MD or (ii) a vaccine composition comprising a dengue antigen comprising the prM and E sequences of CYD-MD. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 16 and a polypeptide having the sequence of SEQ ID NO: 22, wherein said vaccine composition does not comprise a dengue antigen comprising the M and E sequences of CYD-MD or (ii) a vaccine composition comprising a dengue antigen comprising the M and E sequences of CYD-MD. Said vaccine composition is also preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 16 and a polypeptide having the sequence of SEQ ID NO: 22, wherein said vaccine composition does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequence of MD1280 (SEQ ID NO: 11) or (ii) a vaccine composition comprising a chimeric dengue virus of serotype 2 generated using the prM-E sequence of MD1280 (SEQ ID NO: 11). Preferably, a vaccine composition of the present invention which comprises a polypeptide having the sequence of SEQ ID NO: 16 and a polypeptide having the sequence of SEQ ID NO: 22 does not comprise: (i) a chimeric virus comprising the prM and E amino acid sequences of MD1280; (ii) a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-MD (SEQ ID NO: 11) or (iii) a dengue antigen comprising the prM-E sequence from MD1280. When a vaccine composition of the present invention comprises a dengue antigen of serotype 2 which comprises a polypeptide having the sequence of SEQ ID NO: 18 and a polypeptide having the sequence of SEQ ID NO: 23, said vaccine composition is preferably either: (i) a vaccine composition comprising a dengue antigen comprising a polypeptide having the sequence of SEQ ID NO: 18 and a polypeptide having the sequence of SEQ ID NO: 23, wherein said vaccine composition does not comprise a chimeric dengue virus of serotype 2 generated using the prM-E sequence of SEQ ID NO: 2 or (ii) a vaccine composition comprising a chimeric dengue virus of serotype 2 generated using the prM-E sequence of SEQ ID NO: 2. Preferably, a vaccine composition of the present invention which comprises a polypeptide having the sequence of SEQ ID NO: 18 and a polypeptide having the sequence of SEQ ID NO: 23 does not comprise: (i) a dengue antigen of serotype 2 which comprises SEQ ID NO: 2 or (ii) a dengue antigen comprising the prM-E sequence from SEQ ID NO: 2.

A composition of the present invention, as described herein (e.g. a tetravalent formulation, e.g. for use a method of the present invention), may advantageously comprise a dengue antigen of serotype 2 which comprises a polypeptide having at least 90% identity to SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22 or SEQ ID NO: 23. Preferably said dengue antigen of serotype 2 further comprises a polypeptide having at least 90% identity to SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16 or SEQ ID NO: 18. For instance, said dengue antigen of serotype 2 preferably comprises: i) a polypeptide having at least 90% sequence identity to SEQ ID NO: 13 and a polypeptide having at least 90% sequence identity to SEQ ID NO: 19; ii) a polypeptide having at least 90% sequence identity to SEQ ID NO: 14 and a polypeptide having at least 90% sequence identity to SEQ ID NO: 20; iii) a polypeptide having at least 90% sequence identity to SEQ ID NO: 15 and a polypeptide having at least 90% sequence identity to SEQ ID NO: 21; iv) a polypeptide having at least 90% sequence identity to SEQ ID NO: 16 and a polypeptide having at least 90% sequence identity to SEQ ID NO: 22; or v) a polypeptide having at least 90% sequence identity to SEQ ID NO: 18 and a polypeptide having at least 90% sequence identity to SEQ ID NO: 23. In preferred embodiments, the references herein to at least 90% identity may be read as at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity to the given sequence.

The dengue antigens of serotype 2 as described in the preceding paragraphs may advantageously be combined with any of the dengue antigens of serotypes 1, 3 and 4 as described elsewhere herein to form a tetravalent formulation comprising a dengue antigen of serotype 1, a dengue antigen of serotype 2, a dengue antigen of serotype 3 and a dengue antigen of serotype 4. For instance, the dengue antigens of serotypes 1, 3 and 4 may each be independently selected from the group consisting of a live attenuated dengue virus, an inactivated dengue virus, a live attenuated or inactivated chimeric dengue virus or a dengue virus-like particle (VLP). Preferably, said dengue antigens of serotype 1, 3 and 4 are each independently selected from the group consisting of a live attenuated dengue virus and a live attenuated chimeric dengue virus. Preferably, said dengue antigens of serotypes 1, 3 and 4 are live attenuated chimeric dengue viruses of serotypes 1, 3 and 4 respectively. Preferably, said live attenuated chimeric dengue viruses of serotypes 1, 3 and 4 each comprise one or more proteins from a dengue virus and one or more proteins from a different flavivirus. For example, each of said live attenuated chimeric dengue viruses of serotypes 1, 3 and 4 is advantageously a YF/Dengue chimera. Preferably, said dengue antigens of serotypes 1, 3 and 4 are each a live attenuated chimeric dengue virus in which the genetic backbone of a recipient

flavivirus has been modified by exchanging the sequences encoding the prM and E proteins of the recipient flavivirus with the corresponding sequences of a dengue virus. Preferably said recipient flavivirus is a yellow fever virus. For example, in an advantageous embodiment, said live attenuated chimeric dengue viruses of serotypes 1, 5 3 and 4 are respectively a Chimerivax dengue serotype 1 strain (i.e. a CYD-1 strain), a Chimerivax dengue serotype 3 strain (i.e. a CYD-3 strain) and a Chimerivax dengue serotype 4 strain (i.e. a CYD-4 strain).

It is an aim of the present inventors to provide an optimized tetravalent dengue vaccine composition (i.e. vaccine composition comprising a dengue antigen of each of 10 serotypes 1, 2, 3 and 4) which provides an improved neutralising antibody response against dengue virus of serotype 2 when compared with the neutralising antibody response generated by CYD-1, CYD-2, CYD-3 and CYD-4 as defined in Example 1.

Accordingly, in one aspect, the present invention advantageously provides a vaccine composition which comprises a dengue antigen of each of serotypes 1, 2, 3 and 15 4, wherein said dengue antigens of serotypes 1, 3 and 4 are each a live attenuated chimeric dengue virus and said dengue antigen of serotype 2 is a live attenuated dengue virus which comprises a nucleic acid sequence having at least 90% sequence identity to the sequence as set forth in SEQ ID NO: 24.

Accordingly, in another aspect, the present invention advantageously provides a 20 vaccine composition which comprises a dengue antigen of serotype 1, a dengue antigen of serotype 2, a dengue antigen of serotype 3 and a dengue antigen of serotype 4, wherein:

- i) said dengue antigen of serotype 1 is a YF/dengue chimeric dengue virus (i.e. a recipient yellow fever virus in which the genetic backbone of the YF virus has 25 been modified by exchanging the sequences encoding the prM and E proteins of the YF virus by the corresponding sequences of a dengue serotype 1 virus);
- ii) said dengue antigen of serotype 2 is a live attenuated dengue virus of serotype 2 which comprises a nucleic acid sequence having at least 90% sequence 30 identity to the sequence as set forth in SEQ ID NO: 24;
- iii) said dengue antigen of serotype 3 is a YF/dengue chimeric dengue virus (i.e. a recipient yellow fever virus in which the genetic backbone of the YF virus has been modified by exchanging the sequences encoding the prM and E proteins of the YF virus by the corresponding sequences of a dengue serotype 3 virus) and
- iv) said dengue antigen of serotype 4 is a YF/dengue chimeric dengue virus (i.e. a recipient yellow fever virus in which the genetic backbone of the YF virus has 35

been modified by exchanging the sequences encoding the prM and E proteins of the YF virus by the corresponding sequences of a dengue serotype 4 virus).

Preferably, said recipient YF virus (which forms the genetic backbone of the YF/dengue chimeric viruses of serotypes 1, 3 and 4) is an attenuated YF virus. For example, said recipient YF virus may be an attenuated YF virus selected from the group consisting of YF 17D, YF 17DD and YF 17D204. Preferably, the YF/dengue chimeric viruses of serotypes 1, 3 and 4 are respectively a Chimerivax dengue serotype 1 (i.e. a CYD-1), a Chimerivax dengue serotype 3 (i.e. a CYD-3) and a Chimerivax dengue serotype 4 (i.e. a CYD-4).

A reference herein to a nucleic acid sequence having at least 90% sequence identity to the sequence as set forth in SEQ ID NO: 24 may preferably be read as a nucleic acid sequence having at least 92%, at least 94%, at least 96%, at least 98%, at least 99% or 100% sequence identity to the sequence as set forth in SEQ ID NO: 24. Preferably the nucleotides at the positions within said nucleic acid sequences (that have at least 90% sequence identity to the sequence as set forth in SEQ ID NO: 24) which correspond to positions 736, 1619, 4723, 5062, 9191, 10063, 10507, 57, 524, 2055, 2579, 4018, 5547, 6599 and 8571 of SEQ ID NO: 24 are not mutated. Advantageously, a dengue antigen of serotype 2 which is a live attenuated dengue virus for use in a composition of the present invention (for example for use in combination with a dengue antigen of serotypes 1, 3 and 4 as described above and elsewhere herein (e.g. dengue antigens of serotypes 1, 3 and 4 which are live attenuated chimeric dengue viruses, e.g. YF/dengue chimeric dengue viruses)) is a live attenuated dengue virus which comprises a nucleic acid sequence having 100% sequence identity to the sequence as set forth in SEQ ID NO: 24 or a live attenuated dengue virus which comprises at least one and no more than 20 nucleotide substitutions when compared with the sequence as set forth in SEQ ID NO: 24. Preferably said live attenuated dengue virus comprises at least one and no more than 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3 or 2 nucleotide substitutions when compared with the sequence as set forth in SEQ ID NO: 24. Preferably the nucleotides at the positions within said nucleic acid sequences which correspond to positions 736, 1619, 4723, 5062, 9191, 10063, 10507, 57, 524, 2055, 2579, 4018, 5547, 6599 and 8571 of SEQ ID NO: 24 are not mutated. Advantageously, a dengue antigen of serotype 2 which is a live attenuated dengue virus for use in a composition of the present invention comprises a nucleic acid sequence that has no more than 20 base mutations, deletions or insertions when compared with the sequence as set forth in SEQ ID NO: 24. In certain cases said live attenuated dengue virus of serotype 2 comprises a nucleic acid sequence that has no more than 15 or even no more than 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 base

mutations, deletions or insertions when compared with the sequence as set forth in SEQ ID NO: 24. Preferably the nucleotides at the positions within said nucleic acid sequence that correspond to positions 736, 1619, 4723, 5062, 9191, 10063, 10507, 57, 524, 2055, 2579, 4018, 5547, 6599 and 8571 of SEQ ID NO: 24 are not mutated.

5 It is also preferred that a dengue antigen of serotype 2 for use in a vaccine composition of the present invention (e.g. a dengue antigen which is a live attenuated dengue virus or a live attenuated chimeric dengue virus of serotype 2) is capable of inducing neutralizing antibodies in humans and is capable of inducing a balanced immune response when used in the context of a tetravalent dengue vaccine composition. It is also
10 preferred that a dengue antigen of serotype 2 for use in a vaccine composition of the present invention (e.g. a dengue antigen which is a live attenuated dengue virus or a live attenuated chimeric dengue virus of serotype 2) for use in a vaccine composition of the invention results in low or absent viremia in humans. It is also preferred that a dengue antigen of serotype 2 for use in a tetravalent vaccine composition of the present invention
15 (e.g. a dengue antigen which is a live attenuated dengue virus or a live attenuated chimeric dengue virus of serotype 2) provides an improved neutralising antibody response against dengue virus of serotype 2 when compared with the neutralising antibody response generated by CYD-1, CYD-2, CYD-3 and CYD-4 as defined in Example 1.

Advantageously, a composition for use in the present invention comprises a
20 dengue antigen of each of serotypes 1, 2, 3 and 4, wherein: (i) said dengue antigen of serotype 1 is a live attenuated chimeric dengue virus other than CYD-1 or said dengue antigen of serotype 1 is CYD-1; (ii) said dengue antigen of serotype 2 is a live attenuated dengue virus other than VDV-2 or said dengue antigen of serotype 2 is VDV-2; (iii) said dengue antigen of serotype 3 is a live attenuated chimeric dengue virus other than CYD-3
25 or said dengue antigen of serotype 3 is CYD-3 and (iv) said dengue antigen of serotype 4 is a live attenuated chimeric dengue virus other than CYD-4 or said dengue antigen of serotype 4 is CYD-4. In this context, the VDV-2 strain is the strain derived from the DEN-2 16681/PDK53 strain (LAV2) by subsequent adaptation to Vero cells, wherein said VDV-2 strain has 10 additional mutations in comparison with the DEN-2 16681/PDK53 strain
30 including four silent mutations.

Advantageously, a composition for use in the present invention comprises a
dengue antigen of each of serotypes 1, 2, 3 and 4, wherein said dengue antigens of serotypes 1, 3 and 4 are each a live attenuated chimeric dengue virus and said dengue antigen of serotype 2 is a live attenuated dengue virus which comprises a nucleic acid
35 sequence having at least 90% sequence identity to the sequence as set forth in SEQ ID NO: 24 and wherein said dengue antigens of serotypes 1, 2, 3 and 4 are not CYD-1, VDV-

2, CYD-3 and CYD-4 respectively or a dengue antigen comprising the M and E sequences of CYD-1, VDV2, a dengue antigen comprising the M and E sequences of CYD-3 and a dengue antigen comprising the M and E sequences of CYD-4 respectively.

Advantageously, a composition for use in the present invention comprises a dengue antigen of each of serotypes 1, 2, 3 and 4, wherein: (i) said dengue antigen of serotype 1 is a live attenuated chimeric dengue virus other than CYD-1 or said dengue antigen of serotype 1 is CYD-1; (ii) said dengue antigen of serotype 2 is a live attenuated dengue virus other than VDV-2 or said dengue antigen of serotype 2 is VDV-2; (iii) said dengue antigen of serotype 3 is a live attenuated chimeric dengue virus other than CYD-3 or said dengue antigen of serotype 3 is CYD-3 and (iv) said dengue antigen of serotype 4 is a live attenuated chimeric dengue virus other than CYD-4 or said dengue antigen of serotype 4 is CYD-4. In this context, the VDV-2 strain is the strain comprising the nucleic acid sequence as set forth in SEQ ID NO: 24.

A preferred vaccine composition according to the present invention, e.g. for use in a method according to the present invention, comprises a dengue antigen of serotype 1, a dengue antigen of serotype 2, a dengue antigen of serotype 3 and a dengue antigen of serotype 4, wherein:

i) said dengue antigen of serotype 1 is a YF/dengue chimeric dengue virus other than a CYD-1 or said dengue antigen of serotype 1 is a CYD-1;

ii) said dengue antigen of serotype 2 is a live attenuated dengue virus of serotype 2 which comprises a nucleic acid sequence having at least 90% sequence identity to the sequence as set forth in SEQ ID NO: 24, wherein said dengue antigen of serotype 2 is not a live attenuated dengue virus of serotype 2 which comprises a nucleic acid sequence having 100% sequence identity to the sequence as set forth in SEQ ID NO: 24 or said dengue antigen of serotype 2 is a live attenuated dengue virus of serotype 2 which comprises a nucleic acid sequence having 100% sequence identity to the sequence as set forth in SEQ ID NO: 24;

iii) said dengue antigen of serotype 3 is a YF/dengue chimeric dengue virus other than a CYD-3 or said dengue antigen of serotype 3 is a CYD-3; and

iv) said dengue antigen of serotype 4 is a YF/dengue chimeric dengue virus other than a CYD-4 or said dengue antigen of serotype 4 is a CYD-4.

Advantageously, a dengue antigen of serotype 2 which is a live attenuated chimeric dengue virus for use in a vaccine composition of the present invention (for example for use in combination with a dengue antigen of serotypes 1, 3 and 4 as described above and elsewhere herein (e.g. dengue antigens of serotypes 1, 3 and 4

which are YF/dengue chimeric dengue viruses)) comprises a nucleic acid sequence having at least 90% identity to the sequence as set forth in SEQ ID NO: 25. Preferably said nucleic acid sequence has at least 92%, at least 94%, at least 96%, at least 98%, at least 99% or 100% sequence identity to the sequence as set forth in SEQ ID NO: 25.

5 Preferably the nucleotides at the positions within said nucleic acid sequence which correspond to positions 524, 736, 1619 and 2055 of SEQ ID NO: 24 are not mutated (i.e. maintain the nucleotide appearing in SEQ ID NO: 24 at those positions).

Advantageously, a dengue antigen of serotype 2 which is a chimeric dengue virus for use in a vaccine composition of the present invention (for example for use in combination with a dengue antigen of serotypes 1, 3 and 4 as described above and elsewhere herein (e.g. dengue antigens of serotypes 1, 3 and 4 which are YF/dengue chimeric dengue viruses)) comprises a prM-E sequence having at least 90%, at least 95%, at least 98%, at least 99% or 100% identity to the prM-E sequence from the LAV-2 strain (i.e. the RNA equivalent of SEQ ID NO: 4). Preferably the nucleotides at the positions within said prM-E sequence which correspond to positions 524, 736, 1619 and 2055 of the RNA equivalent of SEQ ID NO: 24 are not mutated (i.e. maintain the nucleotide appearing in the RNA equivalent of SEQ ID NO: 24 at those positions).

Advantageously, a dengue antigen of serotype 2 which is a chimeric dengue virus for use in a vaccine composition of the present invention (for example for use in combination with a dengue antigen of serotypes 1, 3 and 4 as described above and elsewhere herein (e.g. dengue antigens of serotypes 1, 3 and 4 which are YF/dengue chimeric dengue viruses)) comprises a prM-E sequence having at least 90%, at least 95%, at least 98%, at least 99% or 100% identity to the prM-E sequence from the MD1280 strain (i.e. the RNA equivalent of SEQ ID NO: 7).

25 A composition of the present invention, as described herein, may advantageously comprise a dengue antigen selected from the group consisting of: (a) a live attenuated dengue virus; (b) an inactivated dengue virus; (c) a live attenuated or inactivated chimeric dengue virus and (d) a combination of two or more of (a) to (c), wherein said dengue antigen comprises a nucleotide sequence selected from the group consisting of SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7 and SEQ ID NO: 1.

30 A composition of the present invention, as described herein, may advantageously comprise a dengue antigen selected from the group consisting of: (a) a live attenuated dengue virus; (b) an inactivated dengue virus; (c) a live attenuated or inactivated chimeric dengue virus and (d) a combination of two or more of (a) to (c), wherein said dengue antigen comprises a nucleotide sequence encoding M and E sequences as described herein.

For instance, a composition of the present invention, e.g. for use in a method of protecting according to the present invention, may advantageously comprise any one of the following combinations of dengue antigens of serotypes 1, 2, 3 and 4: i) CYD-1, CYD-LAV, CYD-3 and CYD-4; ii) CYD-1, CYD-BID, CYD-3 and CYD-4; (iii) CYD-1, CYD-PR, CYD-3 and CYD-4 or (iv) CYD-1, CYD-MD, CYD-3 and CYD-4. A composition of the present invention may also advantageously comprise the following combination of dengue antigens: i) a dengue antigen comprising the prM and E sequences of CYD-1, VDV2, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4. For instance, a composition of the present invention may advantageously comprise CYD-1, VDV-2, CYD-3 and CYD-4. A composition of the present invention, as described herein, may advantageously comprise a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2. A composition of the present invention, as described herein, may advantageously comprise a dengue antigen of serotype 2 which comprises a sequence having at least 90% identity to the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2. For example, said sequence may be at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2.

A vaccine composition for use the present invention, e.g. for use in a method according to the present invention preferably comprises a dengue antigen which is a vaccinal dengue virus. Such vaccinal dengue viruses include, for example, inactivated viruses, live attenuated viruses and live attenuated chimeric dengue viruses. Preferably, the vaccinal dengue viruses are live attenuated chimeric dengue viruses. Preferably, a live attenuated chimeric dengue virus according to the present invention comprises one or more proteins from a dengue virus and one or more proteins from a different flavivirus. Advantageously, said different flavivirus is a yellow fever virus, for example a yellow fever virus of strain YF 17D. Preferably a chimeric dengue virus according to the present invention comprises the prM-E amino acid sequences of a dengue virus, for example a chimeric dengue virus according to the present invention comprises a yellow fever virus genome whose prM-E whose prM-E sequence has been substituted with the prM-E sequence of a dengue virus. Advantageously, a vaccine composition according to the present invention, e.g. for use in a method of the present invention, comprises CYD-1, CYD-2, CYD-3 and CYD-4. A composition of the present invention may advantageously

comprise any one of the following combinations of dengue antigens i) a dengue antigen comprising the prM and E sequences of CYD-1, a dengue antigen comprising the prM and E sequences of CYD-LAV, a chimeric dengue virus comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4; ii) a dengue antigen comprising the prM and E sequences of CYD-1, a dengue antigen comprising the prM and E sequences of CYD-BID, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4; (iii) a dengue antigen comprising the prM and E sequences of CYD-1, a dengue antigen comprising the prM and E sequences of CYD-PR, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4; (iv) a dengue antigen comprising the prM and E sequences of CYD-1, a dengue antigen comprising the prM and E sequences of CYD-MD, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4;. For instance, a composition of the present invention may also advantageously comprise any one of the following combinations of dengue antigens: i) CYD-1, CYD-LAV, CYD-3 and CYD-4; ii) CYD-1, CYD-BID, CYD-3 and CYD-4; (iii) CYD-1, CYD-PR, CYD-3 and CYD-4 or (iv) CYD-1, CYD-MD, CYD-3 and CYD-4. A composition of the present invention may also advantageously comprise the following combination of dengue antigens: i) a dengue antigen comprising the prM and E sequences of CYD-1, VDV2, a dengue antigen comprising the prM and E sequences of CYD-3 and a dengue antigen comprising the prM and E sequences of CYD-4. For instance, a composition of the present invention may advantageously comprise CYD-1, VDV-2, CYD-3 and CYD-4. A composition of the present invention, as described herein, may advantageously comprise a dengue antigen of serotype 2 which comprises the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2. Advantageously, a vaccine composition of the present invention, e.g. a chimeric dengue virus, of serotype 2 may comprise prM-E sequences having at least 90%, at least 95%, at least 98% or at least 99% identity to the prM-E sequences from the serotype 2 strains LAV-2, BID-V585, PR/DB023 or MD1280 as described in the examples or may comprise prM-E sequences having at least 90%, at least 95%, at least 98% or at least 99% identity to the prM-E sequence shown in SEQ ID NO: 2. Advantageously, a vaccine composition, e.g. a chimeric dengue virus, of serotype 2 for use in the method of the present invention may comprise prM-E sequences from the serotype 2 strains LAV-2, BID-V585, PR/DB023 or MD1280 or the prM-E sequence from SEQ ID NO: 2 as described in the examples. When the recipient genomic backbone of such chimeric dengue viruses is derived from YF-VAX®, such strains are referred to

herein as CYD-LAV, CYD-BID, CYD-PR and CYD-MD. A vaccine composition of the present invention comprising chimeric dengue virus of serotype 2 generated using the prM-E sequences of the serotype 2 strains LAV-2 (SEQ ID NO: 8), BID-V585 (SEQ ID NO: 9), PR/DB023 (SEQ ID NO: 10), MD1280 (SEQ ID NO: 11) or SEQ ID NO: 2, or 5 generated using prM-E sequences having at least 90%, at least 95%, at least 98% or at least 99% identity to the prM-E sequences from the serotype 2 strains LAV-2, BID-V585, PR/DB023, MD1280 or the prM-E sequence from SEQ ID NO: 2 may advantageously be used in combination with CYD-1, CYD-3 and CYD-4 in a vaccine composition according to the present invention. A composition of the present invention, as described herein, may 10 advantageously comprise a dengue antigen of serotype 2 which comprises a sequence having at least 90% identity to the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2. For example, said sequence may be at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical 15 to the prM-E sequence of CYD-LAV (SEQ ID NO: 8), CYD-BID (SEQ ID NO: 9), CYD-PR (SEQ ID NO: 10) CYD-MD (SEQ ID NO: 11) or SEQ ID NO: 2.

The exact quantity of a vaccinal dengue virus of the present invention to be administered may vary according to the age and the weight of the patient being vaccinated, the frequency of administration as well as the other ingredients (e.g. 20 adjuvants) in the composition. The quantity of a live attenuated dengue virus of each of serotypes 1 to 4 comprised in a vaccine composition of the present invention lies within a range of from about 10^3 to about 10^7 CCID₅₀. Generally, the quantity of a live attenuated dengue virus of each of serotypes 1 to 4 comprised in a vaccine composition of the present invention lies within a range of from about 10^3 to about 10^6 CCID₅₀, for example 25 within a range of from about 5×10^3 to about 5×10^5 CCID₅₀, for example within a range of from about 1×10^4 to about 1×10^5 CCID₅₀, for example about 10^5 CCID₅₀. The quantity of a live attenuated dengue virus of each of serotypes 1 to 4 comprised in a vaccine composition of the present invention may also lie within a range of from about 10^4 to about 30 10^7 CCID₅₀, for example about 10^6 CCID₅₀. The quantity of a live attenuated dengue virus of each of serotypes 1 to 4 comprised in a tetravalent composition of the present invention may be equal. For example a tetravalent composition of the present invention may comprise about 10^5 CCID₅₀ of each live attenuated dengue virus of serotypes 1 to 4. Alternatively, a tetravalent composition of the present invention may comprise about 10^6 CCID₅₀ of each live attenuated dengue virus of serotypes 1 to 4. Generally, the quantity of 35 an inactivated dengue virus of each of serotypes 1 to 4 comprised in a composition of the present invention lies within a range of from about 10^4 to about 10^8 CCID₅₀ equivalent,

preferably within a range of from about 5×10^4 to about 5×10^7 CCID₅₀ equivalent, preferably within a range of from about 1×10^4 to about 1×10^6 CCID₅₀ equivalent, preferably about 10^5 CCID₅₀ equivalent. Generally, the quantity of a VLP of each of serotypes 1 to 4 comprised in the composition lies within a range of from about 100ng to about 100 μ g of VLP, preferably within a range of from about 100ng to about 50 μ g, preferably within a range of from about 100ng to about 20 μ g, preferably about 1 μ g to 10 μ g. The amount of VLP can be determined by ELISA. Advantageously, a vaccine composition according to the present invention comprises an effective amount of a dengue antigen as defined herein.

10 A vaccine composition according to the present invention may further comprise a pharmaceutically acceptable carrier or excipient. A pharmaceutically acceptable carrier or excipient according to the present invention means any solvent or dispersing medium etc., commonly used in the formulation of pharmaceuticals and vaccines to enhance stability, sterility and deliverability of the active agent and which does not produce any secondary 15 reaction, for example an allergic reaction, in humans. The excipient is selected on the basis of the pharmaceutical form chosen, the method and the route of administration. Appropriate excipients, and requirements in relation to pharmaceutical formulation, are described in "Remington's Pharmaceutical Sciences" (19th Edition, A.R. Gennaro, Ed., Mack Publishing Co., Easton, PA (1995)). Particular examples of pharmaceutically 20 acceptable excipients include water, phosphate-buffered saline (PBS) solutions and a 0.3% glycine solution. A vaccine composition according to the present invention may advantageously comprise 0.4% saline and 2.5% human serum albumin (HSA).

25 A vaccine composition for use in a method of the present invention may optionally contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, human serum albumin, essential amino acids, nonessential amino acids, L-30 arginine hydrochlorate, saccharose, D-trehalose dehydrate, sorbitol, tris (hydroxymethyl) aminomethane and/or urea. In addition, the vaccine composition may optionally comprise pharmaceutically acceptable additives including, for example, diluents, binders, stabilizers, and preservatives. Preferred stabilizers are described in WO 2010/003670.

35 A vaccine composition of the present invention may comprise a dengue antigen which is a dengue immunoprotein. A dengue immunoprotein, as used herein, is a dengue envelope (E) protein, or derivative or fragment thereof, that when administered to an immunocompetent subject induces neutralizing antibodies against a dengue virus of

serotype 1, 2, 3 or 4. Dengue immunoproteins include native and derivatized forms of dengue E proteins, including chemical conjugates, immunological fragments, and fusion proteins thereof.

Dengue immunoproteins, or derivatives or fragments thereof may be conjugated to carrier molecules. Such conjugation may be achieved by chemical conjugation techniques or through the recombinant expression of fusion proteins comprising the dengue immunoproteins or derivatives or fragments thereof and the carrier molecule. Examples of carrier molecules which may be used for preparing conjugates include diphtheria toxoid, tetanus toxoid, fragment C of tetanus toxin, mutants of diphtheria toxin including CRM 5 197, CRM 176, CRM228, CRM 45, CRM 9, CRM 45, CRM 102, CRM 103 and CRM 107, pneumococcal pneumolysin, OMPC, heat shock proteins, pertussis proteins, pneumococcal surface protein PspA or the toxin A or B of *Clostridium difficile*.

A vaccine composition of the present invention may comprise one or more adjuvants to enhance the immunogenicity of the dengue antigens. Those skilled in the art 15 will be able to select an adjuvant which is appropriate in the context of this invention. An adjuvant is preferably used in a vaccine composition of the invention comprising an inactivated virus or a VLP or a dengue structural protein. An adjuvant may be used in a vaccine composition of the invention comprising a live attenuated virus, as long as said adjuvant does not impact replication.

20 Suitable adjuvants include an aluminum salt such as aluminum hydroxide gel, aluminum phosphate or alum, but may also be a salt of calcium, magnesium, iron or zinc. Further suitable adjuvants include an insoluble suspension of acylated tyrosine or acylated sugars, cationically or anionically derivatized saccharides, or polyphosphazenes. Alternatively, the adjuvant may be an oil-in-water emulsion adjuvant (EP 0 399 843B), as 25 well as combinations of oil-in-water emulsions and other active agents (WO 95/17210; WO 98/56414; WO 99/12565 and WO 99/11241). Other oil emulsion adjuvants have been described, such as water-in-oil emulsions (U.S. Pat. No. 5,422, 109; EP 0 480 982 B2) and water-in-oil-in-water emulsions (U.S. Pat. No. 5,424,067; EP 0 480 981 B). Examples of such adjuvants include MF59, AF03 (WO 2007/006939), AF04 (WO 2007/080308), 30 AF05, AF06 and derivatives thereof. The adjuvant may also be a saponin, lipid A or a derivative thereof, an immunostimulatory oligonucleotide, an alkyl glucosamide phosphate, an oil in water emulsion or combinations thereof. Examples of saponins include Quil A and purified fragments thereof such as QS7 and QS21.

35 As appreciated by skilled artisans, a vaccine composition of the present invention is suitably formulated to be compatible with the intended route of administration. Examples of suitable routes of administration include for instance intramuscular,

transcutaneous, subcutaneous, intranasal, oral or intradermal. Advantageously, the route of administration is subcutaneous.

The vaccine compositions of the present invention may be administered using conventional hypodermic syringes or safety syringes such as those commercially available from Becton Dickinson Corporation (Franklin Lakes, NJ, USA) or jet injectors. For intradermal administration, conventional hypodermic syringes may be employed using the Mantoux technique or specialized intradermal delivery devices such as the BD Soluvia(TM) microinjection system (Becton Dickinson Corporation, Franklin Lakes, NJ, USA), may be used.

10 The volume of a vaccine composition of the present invention administered will depend on the method of administration. In the case of subcutaneous injections, the volume is generally between 0.1 and 1.0 ml, preferably approximately 0.5 ml.

15 Optionally, booster administrations of a vaccine composition according to the present invention may be used, for example between six months and ten years, for example six months, one year, three years, five years or ten years after initial immunization (i.e. after administration of the last dose scheduled in the initial immunization regimen).

20 According to one embodiment, the invention also provides a kit comprising a vaccine composition of the invention and instructions for the use of said vaccine composition in a method of protecting a human subject against dengue disease. The kit can comprise at least one dose (typically in a syringe) of any vaccine composition contemplated herein. According to one embodiment the kit may comprises a multi-dose formulation (typically in a vial) of any vaccine composition as described herein. The kit further comprises a leaflet mentioning the use of the said vaccine composition for the 25 prevention of dengue disease or the use of the said vaccine for the prophylaxis of dengue disease. The leaflet may further mention the vaccination regimen and the human subject population to be vaccinated.

30 The efficacy of a vaccine composition of the present invention in reducing the likelihood or severity of dengue disease may be measured in a number of ways. For instance the efficacy of a vaccine composition of the present invention in reducing the likelihood or severity of symptomatic virologically-confirmed dengue disease may be calculated by measuring after the administration of at least one dose of said vaccine composition (e.g. after administration of one, two or three doses of said vaccine composition):

35 (i) the percentage of symptomatic virologically-confirmed dengue cases caused by dengue virus of any serotype;

- (ii) the percentage of severe virologically-confirmed dengue cases caused by dengue virus of any serotype;
- (iii) the percentage of dengue hemorrhagic fever cases of Grades I to IV caused by dengue virus of any serotype;
- 5 (iv) the percentage of DHF cases of Grade I caused by dengue virus of any serotype;
- (v) the percentage of DHF cases of Grade II caused by dengue virus of any serotype;
- 10 (vi) the percentage of DHF cases of Grade III caused by dengue virus of any serotype;
- (vii) the percentage of DHF cases of Grade IV caused by dengue virus of any serotype;
- (viii) the annual incidence rate of hospitalized virologically-confirmed dengue caused by dengue virus of any serotype; and/or
- 15 (ix) the length of hospital stay for symptomatic virologically-confirmed dengue cases caused by dengue virus of any serotype;

in a group of subjects that has received said vaccine composition and comparing those measurements with the equivalent measurements from a control group of subjects that has not received said vaccine composition, wherein the subjects in both said groups are resident in a Dengue endemic region. A statistically significant reduction in any one or more of (i) to (ix) in the vaccinated group of subjects when compared with the unvaccinated control group of subjects is indicative of the efficacy of a vaccine composition according to the present invention. In a preferred embodiment, the efficacy of a vaccine composition according to the present invention is demonstrated by a statistically significant reduction of one or more of the measures as described above, wherein the DHF cases or dengue cases are caused by dengue virus of serotypes 1, 3 or 4.

The efficacy of a vaccine composition according to the present invention in reducing the severity or likelihood of dengue disease may also be calculated by measuring after the administration of at least one dose of said vaccine composition (e.g. 30 after administration of one, two or three doses of said vaccine composition):

- (i) the mean duration and/or intensity of fever;
- (ii) the mean value for plasma leakage as defined by a change in haematocrit;
- (iii) the mean value for thrombocytopenia (platelet count); and/or
- (iv) the mean value of the level of liver enzymes including alanine aminotransferase (ALT) and aspartate aminotransferase (AST);

in a group of subjects that has received said vaccine composition and who have developed virologically-confirmed dengue disease and comparing those measurements with the equivalent measurements from a control group of subjects that has not received said vaccine composition and who have developed virologically-confirmed dengue disease. A statistically significant reduction in any one or more of (i) to (v) in the vaccinated group of subjects who have developed virologically-confirmed dengue disease when compared with the control group of subjects who have developed virologically-confirmed dengue disease is indicative of the efficacy of a vaccine composition according to the present invention in reducing the severity or likelihood of dengue disease.

Typically the efficacy of the method of protection of the invention against a dengue disease, as measured e.g. by the method described in example 1 ($VE=100*(1-ID_{CYD}/ID_{Control})$, where ID is the incidence density (i.e., the number of human subjects with virologically-confirmed dengue divided by the number of person-years at risk) in each group), is at least 50%, preferably at least 60%, wherein said dengue disease is caused by serotype 1, 3 or 4. The efficacy of the method of protection being advantageously at least 70%, preferably 80% against a dengue disease caused by serotype 3 or 4. The efficacy of the method of protection being advantageously at least 90% against dengue disease caused by serotype 4.

Percent identity between two amino acid sequences or two nucleotide sequences is determined by standard alignment algorithms such as, for example, Basic Local Alignment Tool (BLAST) described in Altschul et al. (1990) *J. Mol. Biol.*, 215: 403-410, the algorithm of Needleman et al. (1970) *J. Mol. Biol.*, 48: 444-453; the algorithm of Meyers et al. (1988) *Comput. Appl. Biosci.*, 4: 11-17; or Tatusova et al. (1999) *FEMS Microbiol. Lett.*, 174: 247-250, etc. Such algorithms are incorporated into the BLASTN, BLASTP and "BLAST 2 Sequences" programs (see www.ncbi.nlm.nih.gov/BLAST). When utilizing such programs, the default parameters can be used. For example, for nucleotide sequences the following settings can be used for "BLAST 2 Sequences" : program BLASTN, reward for match 2, penalty for mismatch-2, open gap and extension gap penalties 5 and 2 respectively, gap x~dropoff 50, expect 10, word size 11, filter ON. For amino acid sequences the following settings can be used for "BLAST 2 Sequences" : program BLASTP, matrix BLOSUM62, open gap and extension gap penalties 11 and 1 respectively, gap x~dropoff 50, expect 10, word size 3, filter ON.

It is understood that the various features and preferred embodiments of the present invention as disclosed herein may be combined together.

Throughout this application, various references are cited. The disclosures of these references are hereby incorporated by reference into the present disclosure.

The present invention will be further illustrated by the following examples. It should be understood however that the invention is defined by the claims, and that these examples are given only by way of illustration of the invention and do not constitute in any way a limitation thereof.

5

EXAMPLES

Example 1: One year follow-up in Thailand of patients vaccinated with a tetravalent dengue vaccine (TDV) composition comprising Chimerivax™ DEN-1, DEN-2, DEN-3 and DEN-4

10

Methods

Study design and participants

15

An observer-blind, randomised, controlled, monocentre, Phase IIb trial of the efficacy of the tetravalent Chimerivax™ vaccine (i.e. a tetravalent vaccine comprising the particular CYD-1 strain generated from the prM and E sequences of DEN1 PU0359 (TYP 1 140), the particular CYD-2 strain generated from the prM and E sequences of DEN2 PUO218, the particular CYD-3 strain generated from the prM and E sequences of DEN3 PaH881/88 and the particular CYD-4 strain generated from the prM and E sequences of DEN4 1228 (TVP 980), see WO 03/101397 and Guy *et al.*, Vaccine (2011), 29(42): 7229-41) against virologically-confirmed dengue disease is conducted. 4002 schoolchildren aged 4-11 years who are in good health based on medical history and physical examination are enrolled into the trial. The study is conducted at Ratchaburi Regional

20

Hospital (RRH), Ratchaburi province, Muang district, Thailand. Children with acute febrile illness at enrolment, those with congenital or acquired immunodeficiency, and those receiving immunosuppressive therapy or other prohibited treatments or vaccines are excluded. Participants are randomly assigned 2:1 to receive three doses of dengue vaccine or a control product at Months 0, 6 and 12.

25

30

Products

Each of the chimeric viruses are produced and cultured on Vero cells as described in WO 03/10197, Guy *et al*, Hum. Vaccines (2010) 6 (9): 696; Guy *et al*, Vaccine (2010) 28: 632; Guirakhoo *et al*, J. Virol. (2000) 74 : 5477 ; Guirakhoo *et al*, J. Virol. (2001) 75 (16) : 7290 ; Guirakhoo *et al*, Virol. (June 20, 2002) 298: 146; and Guirakhoo *et al*, J. Virol. (2004) 78 (9): 4761.

The vaccine is presented as a lyophilized powder (previously stored at temperature of between 2°C and 8°C), which is reconstituted with 0.5 mL of solvent for injection (0.4% NaCl containing 2.5% human serum albumin). As reconstituted, each 0.5 mL dose of vaccine contains $5 \pm 1 \log_{10}$ CCID₅₀ of each chimeric dengue serotype (1, 2, 3 and 4) and excipients (essential amino acids, non-essential amino acids, L-arginine chlorhydrate, saccharose, D-trehalose dehydrate, sorbitol, tris (hydroxymethyl) aminoethane and urea). The control product is inactivated rabies vaccine (Verorab®, Sanofi Pasteur, Lyon France) for the first injection of the first 50 children randomised to the control group, and 0.9% NaCl saline placebo for all other injections. All products are injected subcutaneously into the deltoid region of the upper arm.

Assessments

All children are actively followed to detect acute febrile illness based on daily surveillance of school registers during school terms for absenteeism (followed by phone calls or home visits to absentees), and twice-weekly home visits, phone calls or mobile phone text-messages throughout school holidays. In any case of febrile illness (defined as illness with two temperature readings of $\geq 37.5^{\circ}\text{C}$ at least 4 hours apart) parents are asked to take their child to RRH for diagnosis and treatment. The surveillance system also captures spontaneous consultations at RRH. Consecutive febrile episodes separated by a symptom-free interval of ≥ 14 days are considered as separate episodes. Paired serum samples are collected at presentation (i.e., acute sample, collected no later than 7 days after fever onset) and 7–14 days later (convalescent sample) and sent to Sanofi Pasteur's Global Clinical Immunology (GCI) laboratory (Swiftwater, PA, USA) and to the Centre for Vaccine Development (CVD, Mahidol University, Thailand). Acute samples are screened for the presence of flavivirus using an initial RT-PCR assay which detects the presence of any flavivirus (using primers composed of highly conserved flavivirus sequences). Positive samples are tested for wild-type dengue virus with a serotype-specific quantitative RT-PCR, as described herein. In parallel, all acute samples are tested for the presence of dengue NS1 antigen using commercial ELISA kit (Platelia™, Bio-Rad Laboratories, Marnes-La-Coquette, France). A virologically-confirmed episode of dengue disease is defined as a positive result in either the serotype-specific RT-PCR, or the NS1 antigen ELISA.

Active surveillance is maintained until each participant has been followed for at least 13 months after the third vaccination and until the Independent Data Monitoring Committee (IDMC) confirms that ≥ 27 cases have occurred in the per-protocol (PP) population.

All serious adverse events (SAE) are documented until the sixth month after the last vaccination, and thereafter any fatal SAE or vaccine-related SAE.

Dengue immune responses are assessed in the first 300 enrolled children at RRH in sera collected at enrolment and 28 days after each injection. Sera are also 5 prospectively collected from all participants on Day 28 after the third injection to assess immune responses in children with virologically-confirmed dengue occurring from this timepoint. Sera are sent to GCI for measurement of serotype-specific neutralizing antibody titres against the CYD parental dengue viruses using the plaque-reduction neutralization test (PRNT₅₀) as described herein. The assay's quantitation limit is 10 10 (1/dil). Samples below this value are assigned the titre 5 and considered seronegative.

Statistical analysis

The primary objective is to determine vaccine efficacy (VE) against cases of symptomatic, virologically-confirmed dengue occurring more than 28 days after the third vaccination among children enrolled and vaccinated as planned, according to the 15 equation: $VE = 100 * (1 - ID_{CYD}/ID_{Control})$, where ID is the incidence density (i.e., the number of children with virologically-confirmed dengue divided by the number of person-years at risk) in each group. With an assumed disease incidence of 1·3%, a true VE of 70%, a minimum follow-up period of 1 year after the third vaccination, and a per protocol (PP) 20 subject attrition rate of 7·5%/year, 4002 subjects assigned with a 2:1 ratio to dengue vaccine or control are needed to demonstrate, with more than 82% power, and 95% confidence, that VE is not nul. Analyses are based on the two-sided 95% confidence interval (CI) of VE, calculated using the Exact method (Breslow NE, Day NE. Statistical Methods in Cancer Research, Volume II – The Design and Analysis of Cohort Studies. International Agency for Research on Cancer (IARC scientific publication No. 82), Lyon, 25 France). The primary analysis is performed on the PP population, i.e. those who satisfy the enrolment criteria, who correctly receive all three doses of the assigned vaccine at Months 0, 6 (± 15 days), and 12 (± 30 days), and for whom group allocation is not unmasked. This analysis is repeated on the full analysis set for efficacy, in those who receive three injections. As a secondary objective, VE against dengue is determined 30 before completion of the 3-dose vaccination regimen. In an analysis defined after unblinding, VE against each serotype individually is investigated. Analyses for safety and immunogenicity endpoints are descriptive, using 95%CI.

Results

Of the 4002 children enrolled, 95.9% complete the vaccinations and 91.8% are included in the per protocol (PP) analysis set for efficacy. Vaccine and control groups are comparable for age and gender. More than 90% of those sampled at baseline are positive for antibodies against dengue or JEV.

5 *Efficacy*

During the study, 131 dengue cases (131 children had 136 episodes) are virologically-confirmed. Of these, 77 occur more than 28 days after the third injection in the PP population and are included in the primary analysis: 45 cases occurred during 2522 person-years at risk in the vaccine group, while 32 cases occurred during 1251 person-years at risk in the control group. The corresponding vaccine efficacy is 30·2% (95%CI: -13·4–56·6). This finding is confirmed in the full analysis set (see Table 1 below). Efficacy after at least one injection is 33·4% (95%CI: 4·1–53·5) and after at least two injections is 35·3% (95%CI: 3·3–56·5).

15 **Table 1: Serotype-specific and overall efficacy of CYD tetravalent dengue vaccine against virologically-confirmed dengue disease**

	Dengue vaccine		Control		Efficacy	
	Person-years at risk	Cases or Episodes*	Person-years at risk	Cases or Episodes*	%	(95% CI)
>28 days after 3 injections (per-protocol analysis)						
Cases	2522	45	1251	32	30·2	(-13·4–56·6)
Serotype 1 episodes	2536	9	1251	10	55·6	(-21·6–84·0)
Serotype 2 episodes	2510	31	1250	17	9·2	(-75·3–51·3)
Serotype 3 episodes	2541	1	1257	2	75·3	(-375·0–99·6)
Serotype 4 episodes	2542	0	1263	4	100	(24·8–100)
NS1 Antigen positive only episodes	2542	4	1265	0	ND	ND
>28 days after 3 injections (Full analysis set)						
Cases	2620	46	1307	34	32·5	(-8·5–57·6)
Serotype 1 episodes	2633	9	1308	10	55·3	(-22·5–83·9)
Serotype 2 episodes	2608	32	1307	19	15·6	(-57·6–53·6)
Serotype 3 episodes	2638	1	1312	2	75·1	(-378–99·6)
Serotype 4 episodes	2641	0	1320	4	100	(-24·3–100)
NS1 Antigen positive only episodes	2640	4	1322	0	ND	ND
>28 days after at least 1 injection (Full analysis set)						
Cases	5089	75	2532	56	33·4	(4·1–53·5)
Serotype 1 episodes	5139	14	2564	18	61·2	(17·4–82·1)
Serotype 2 episodes	5107	51	2560	26	1·7	(-64·3–39·8)

64

Serotype 3 episodes	5144	4	2565	10	80·1	(30·9–95·4)
Serotype 4 episodes	5149	1	2577	5	90·0	(10·5–99·8)
NS1 Antigen positive only episodes	5147	5	2579	1	-150·	(-11750–72·0)

Data are number except where indicated. ND: not determined. *A 'case' was defined as a first episode of dengue fever virologically-confirmed by either serotype-specific PCRs, or NS1 antigen ELISA. Serotype-specific efficacy was calculated including all episodes of that serotype; 5 children with two virologically confirmed dengue episodes during the study were therefore included twice in the serotype-specific analysis.

Post-hoc analyses reveal differing efficacy by serotype (see Table 1). Efficacy against DENV1, DENV3, and DENV4 after at least one injection is in the range 61·2%–90·0%, compared with 1·7% against DENV2. Efficacy against DENV1, DENV3, and 5 DENV4 after three injections is in the range 55·3%–100%, compared with 15·6% against DENV2.

In those subjects that acquired virologically-confirmed dengue, a statistically significant reduction in the annual incidence rate of hospitalization was observed in the vaccinated group when compared with the control group. The relative risk (RR) after three 10 doses was 0·523 (see Table 2).

Table 2: Incidence of hospitalized virologically-confirmed dengue during the trial

Time period	CYD Dengue Vaccine Group (N=2666)				Control Group (N=1331)				Relative Risk RR (95%CI)
	M	Cases	Incidence Rate (95%CI)	n Occurrences	M	Cases	Incidence Rate (95%CI)	n Occurrences	
Year 1	2666	8	0.3 (0.1; 0.6)	8	1331	7	0.5 (0.2; 1.1)	7	0.571 (0.181, 1.85)
Year 2	2557	24	0.9 (0.5; 1.3)	24	1282	23	1.7 (1.0; 2.5)	23	0.523 (0.283, 0.970)

Year 1 = D0 to injection 3 ; Year 2 = Injection 3 to the end of Active Phase

15

Table 3: Rate of hospitalisation by serotype

	Vaccinee Group (%)	Control Group (%)
Serotype 1	8/14 (57.1)	9/18 (50%)
Serotype 2	20/52 (38.5)	15/27 (55.6)
Serotype 3	1/4 (25)	3/11 (27.3)
Serotype 4	0/1	2/5 (40)

No serotype	3/5 (60)	1/1 (100)
NS1 +ve		
Total	32/76 (42.1)	30/62 (48.4)

Immunogenicity

Geometric mean titres (GMT) of neutralising antibodies against dengue serotypes 1–4 on Day 28 after the third injection in the per-protocol analysis set are, respectively, 146 (95%CI: 98.5–217), 310 (224–431), 405 (307–534), and 155 (123–196) in the vaccine group. In the control group these values are 23.9 (14.0–40.9), 52.2 (26.8–102), 48.9 (25.5–93.9), and 19.4 (11.6–32.2). Post one year GMTs are respectively 76.5; 122; 94 and 153 for serotypes 1, 2, 3 and 4.

Safety

There are 584 SAEs during this phase of the study: 366 are reported by 11.8% (315/2666) of participants in the vaccine group, and 218 are reported by 13.2% (176/1331) of participants in the control group. There are no vaccine-related SAEs in the dengue group and there is one in the control group. SAEs observed are medical conditions consistent with the age group and showed no clustering within the 7- or 28-day post-vaccination periods.

Virologically-confirmed dengue cases occurring as a breakthrough in vaccinees were not more serious than those cases occurring in the control group.

Sequence of the prM-E region of circulating wild type serotype 2 strain in the trial

The nucleotide and amino acid sequence of the prM-E region of the wild type serotype 2 strain that causes the DEN-2 cases in the trial is determined. These are set out below as SEQ ID NO: 1 and SEQ ID NO: 2 respectively. The E and the M amino acid sequences of the serotype 2 strain that causes the DEN-2 cases in the trial are described in SEQ ID NOs: 18 and 23 respectively.

25

>nucleotide sequence (SEQ ID NO: 1)

ttccatctaaccacacgcaacggagaaccacacatgatcgctggatacaggagaaaggga
aaagtcttctgttcaaaacagaggatggtgtgaacatgtgcaccctcatggctatggacct
tggtaatttgtgaagacacaatcacgtacaagtgtccttctcaggcagaatgagcca
30 gaagacatagactgttggtgcaactccacgtccacgtggtaacctatggacactgtacca

ctacgggagaacataggagagaaaaaagatcagtggcactcgccatgtggaaatggg
 actggagacgcgaaccgaaacatggatgtcatcagaagggcttggaaacatgcccagaga
 attgaaacttggatcctgagacatccaggcttaccataatggcagaatcctggcataca
 5 ccataggaacgacacatccagagactcctgatccatcctactgacagctgtcgctcc
 ttcaatgacaatgcgttgcataggaatataatcataatagagactttgttagaagggttcagga
 ggaagttgggttgcataatgtcttagaacatggaaagctgtgacgcgatggcaaaaaaca
 aaccaacattggatttgcactgataaaaaacggaagccaaacagcctgcccacccctaaggaa
 gtactgcatagaagcaaaactaaccaacacaacagaatcccgtgcccacacaaggaa
 10 gaacccagcctaaaagaagagcaggacaagaggttcgtctgcaaacactccatggtagaca
 gaggatggggaaatggatgtggattattggaaagggaggcattgtgacctgtgctatgtt
 cacatgcaaaaagaacatggaaagggaaaatcgtgcaaccagaaaaacttggaaatacaccatt
 gtggtaacacccactcactcaggagaagagcatgcggcggaaatgacacaggaaacacggca
 15 aggaaatcaaagtaacaccacagacttccatcacagaagcagaactgacaggttatggcac
 cgtcacatggagtgcctcccgagaacacaggcctcgacttcaatgagatgggtttgctgcag
 atggaaaataaagcttggctgtgcataaggcaatggtttagacactgcccattaccatggc
 tgcccgagcggataaacaagaatcaaattggatacagaaagaaacattggtcacttcaa
 20 aaatccccatgcgaagaaacaggatgttggtttagatcccaagaagggccatgcata
 acagcactcacaggagccacagaaatccaaatgtcgtaggaaactgctcttcactggac
 atctcaagtgcaggctgagaatggacaagctacagctaaaggaatgtcataactctatgt
 cacagggaaagttaaagttgtgaaggaaatagcagaaacacaatggaaacgatagttatc
 25 aagtgcaatataatgaaaggggacggctccatgtaaaattcctttgagataatggattgg
 aaaaaagatatgtcttaggcccctgatcacagtcaacccaaattgttaacagaaaaagacag
 cccagtcaacatagaagcagaacccatccattcgagacagttacatcatcataggagtagag
 ccgggacaactgaagctcaactggcaagaaagggaaatgttcatcgccaaatgtttgaga
 30 caacgatgagagggggcgaagagaatggccatttgggtgacacagcctggacttcggatc
 cctgggaggagtgttacatctataggaaaagctctccaccaagtcttggagcgtatcat
 gggcgtccctcagtgggtttcatggaccatgaaaatcctcataggagtcattatcacat
 ggataggaatgaactcacgcagcacctcaactgtctgtcactggacttggggaaatgt
 gacactgtatttaggatcatggcaggcc

>amino acid sequence (SEQ ID NO: 2)

FHLTTRNGEPHMIVG1QEKGKSLLFKTEDGVNMCLTMAMDLGELCEDTITYKCPLLRQNEP
 EDIDWCNSTSTWVTYGTCTTGEHRREKRSVALVPHVMGLETRTETWMSSEGAWKHAQR
 IETWILRHPGFTIMAAILAYTIGTHFQRVLIFILLTAVAPSMTMRCIGISNRDFVEGVSG
 35 GSWVDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCIEAKLTNTTTESRCPTQG
 EPSLKEEQDKRFVCKHSMVDRGWNGCGLFGKGGIVTCAMFTCKKNMEGKIVQOPENLEYTI
 VVTPHSGEEHAVGNDTGKHGKEIKVTPQSSITEAEALTGYGTVMECSPRTGLDFNEMVLLQ
 MENKAWLVRQWFLDLPLPWLPAGDKQESNWIQKETLVTFKNPHAKKQDVFVVLGSQEGAMH
 TALTGATEIQMSSGNLLFTGHLKCRRLMDKLQLKGMSYSMCTGKFKVVKEIAETQHGTIVI
 40 RVQYEGDGSPCKIPFEIMDLEKRYVLGRLITVNPIVTEKDSPVNIEAEPPFGDSYIIIGVE
 PGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGGVFTSIGKALHQVFGAIY
 GAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVSLVGVIVTLYLGVMVQA

>amino acid sequence (SEQ ID NO: 18)

45 MRCIGISNRDFVEGVSGGSWVDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCI
 EAKLTNTTTESRCPTQGEPSLKEEQDKRFVCKHSMVDRGWNGCGLFGKGGIVTCAMFTCK
 KNMEGKIVQOPENLEYTI VVTPHSGEEHAVGNDTGKHGKEIKVTPQSSITEAEALTGYGTVM
 ECSPRTGLDFNEMVLLQ MENKAWLVRQWFLDLPLPWLPAGDKQESNWIQKETLVTFKNPH

AKKQDVVVLGSQEGAMHTALTGATEIQMSSGNLLFTGHLKCRLRMDKLQLKGMSYSMCTGK
FKVVKEIAETQHGTIVIRVQYEGDGSPCKIPFEIMDLEKRYVLGRLITVNPIVTEKDSPVN
IEAEPPFGDSYIIGVEPGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGG
VFTSIGKALHQVFGAIYGAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVSLVLGIVTLY
5 LGVMVQA

>amino acid sequence (SEQ ID NO: 23)

SVALVPHVMGLETRTETWMSSEGAWKHAQRIETWILRHPGFTIMAAILAYTIGTTHFQRV
LIFILLTAVAPSMT

10

Discussion

The main finding from this study is that a safe, efficacious vaccine against dengue based on the chimeric CYD viruses is possible. Estimated efficacy against DENV1, 3 and 4 is in a range consistent with the 70% hypothesis and is statistically significant after at 15 least one vaccination. Efficacy in a range consistent with the 70% hypothesis is not observed against DENV2. Since DENV2 is the prevalent serotype in this study, overall vaccine efficacy is diminished in this setting.

The vaccine's safety and reactogenicity profile is good, and no vaccine-related SAEs and no safety signals are identified during the review of AEs and SAEs collected 20 from over two years of active follow-up of more than 2600 vaccinees. Theoretical safety concerns associated with the potential enhancement of the rate or severity of dengue disease by an incomplete immune response against the four serotypes of dengue have previously hampered vaccine development. In this trial, the absence of disease enhancement in the presence of an incomplete immune response against the circulating 25 DENV2 viruses is an important and reassuring finding. For instance, cases in vaccinees do not differ from cases in controls in terms of factors such as the duration of fever or in terms of the classical clinical signs of dengue such as bleeding, plasma leakage or thrombocytopenia. Furthermore, severe dengue was not more frequent among vaccinees than controls at any point during the trial).

30 It was also demonstrated that, in those subjects that acquired virologically-confirmed dengue, a statistically significant reduction in the annual incidence rate of hospitalization was observed in the vaccinated group when compared with the control group. This reduction was seen in those subjects that acquired virologically-confirmed dengue of serotype 2 (see Table 3).

35 The results observed in respect of DENV2 may be explained by a number of contributing factors. For instance, there is a possible antigenic mismatch between the CYD2 vaccine virus and the DENV2 virus that causes disease in the trial. In the 1990s,

the Asian 1 genotype of DENV2 emerged in South-East Asia, replacing the previously dominant Asian/American lineage of viruses. Several mutations identified in Domain 2 of the E protein (E83, and in particular E226 and E228) are suggestive of changing viral fitness and antigenicity. The donor wild-type virus for the CYD2 vaccine (and the challenge strain used in the PRNT50) was a clinical isolate from Bangkok in 1980 (Guirakhoo F et al., J Virol 2000, 74: 5477–85). While this virus is also classified as belonging to the Asian 1 genotype, the above-mentioned key amino acid residues in this virus (and thus in CYD2) correspond to those of the Asian/American genotype (Hang et al PLoS Negl Trop Dis. 2010 Jul 20;4(7):e757).

. Additionally, there are two extremely rare mutations in the prM-E sequence of the CYD2 vaccine that may also contribute to a mismatched immune response. These mutations are at positions prM24 and E251 (Guirakhoo et al, J. Virol. (2004) 78 (9): 4761).

The results observed against DENV2 are not associated with an absence of immunogenicity in the PRNT₅₀ assay. Neutralising antibody responses after vaccination against DENV2 are higher than those against DENV1 and DENV3.

In conclusion, the present study constitutes the first ever demonstration that a safe and efficacious dengue vaccine is possible and represents a major milestone in dengue vaccine development.

20 **Example 2: Identification of optimized dengue vaccinal strains of serotype 2**

The objective of the present example is to identify dengue virus strains of serotype 2 which provide the basis for generating optimized dengue vaccine compositions against dengue virus of serotype 2, wherein said optimized dengue vaccine compositions provide improved efficacy in comparison to Chimerivax™ CYD-2 when used in a method according to the present invention.

Criteria determining the selection of optimized strains for the determination of a universal dengue 2 antigen include: (i) recently circulating strain; (ii) balanced selection between Asian and American strains; (iii) an optimized strain should have a prM-E sequence that is as similar as possible to a calculated global consensus sequence generated by aligning the available prM-E sequences of dengue viruses of serotype 2; (iv) amino acid variations that are predicted to impact antibody recognition should be avoided; (v) rare amino acids at a particular positions in the prM and E sequences should be avoided, especially in the E protein ectodomain (a rare amino acid at a particular position is defined as a amino acid that appears at that position in less than 15% of the aligned sequences); (vi) optimized strains for which some previous laboratory experience exists

are preferred and (vii) a dengue antigen that leads to a balanced immune response in a tetravalent composition.

Criteria determining the selection of optimized strains for a local dengue 2 antigen (i.e. that is especially effective against a wild type dengue virus circulating in a particular area)

5 are criteria (i) and (vii).

Methods

Databases

Sequence are retrieved from the National Center for Biotechnology Information (NCBI) Dengue virus variation database

10 (www.ncbi.nlm.nih.gov/genomes/VirusVariation/Database/nph-select.cgi?tax_id=12637).

Sequence analyses

Sequence alignments are performed using the MUSCLE algorithm (Edgar, R. C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res, 32(5):1792-1797).

15 Sequence alignment outputs are generated in Vector NTi version 9, module AlignX (Invitrogen). Sequence similarity searches are carried out using the BLAST algorithm (Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990) Basic local alignment search tool. J Mol Biol, 215(3):403-410).

Sequence numbering for prM-E sequences

The sub-sequences included in the prM-E sequences may be numbered in various ways: (i) the total prM-E protein sequence is numbered from position 1 to position 661, with the preM protein sequence designated as position 1 to position 90/91, the M protein sequence designated as position 91/92 to position 166 and the E protein sequence 25 designated as position 167 to position 661; (ii) the prM and M protein sequences are numbered together, i.e. from position 1 to position 166 of the total sequence and E is numbered separately from position 1 to position 495; (iii) the prM, M and E sequences are numbered separately, i.e. prM is numbered from position 1 to 90/91, M is numbered from 1 to 75/76 and E from position 1 to position 495.

Results

Public sequences retrieval

All available dengue virus serotype 2 full length prM and E protein sequences are downloaded from the NCBI Dengue database. Download of sequences takes place on two separate occasions - on 4 October 2010 and in 2011. On the first occasion 669 sequences are downloaded and on the second occasion approximately 3200 sequences are downloaded.

Global consensus sequence generation

On each occasion, all retrieved protein sequences are aligned to generate a global consensus sequence for the prM and E proteins of dengue virus of serotype 2. By definition, the global consensus sequence is an artificial sequence containing the most frequently encountered amino acid at each position. The global consensus sequences for the 2010 alignment and the 2011 alignment only differ by two amino acids. In the 2010 alignment, the global consensus sequence contains isoleucine and valine at positions 129 and 308 respectively of the E protein (by reference to the 1-495 E sequence numbering) and, by contrast, in the 2011 alignment, the global consensus sequence contains valine and isoleucine at positions 129 and 308 respectively of the E protein (by reference to the 1-495 E sequence numbering). The differences in the 2010 and 2011 global consensus sequences is explained by the fact that the respective percentages of strains containing valine or isoleucine at those positions is close to 50%. The global consensus sequence for the prM-E sequence is therefore represented as follows:

fhltrnghphmivgrqekgksllfktdgvmctlmaidlgeledtitykcp11rqnep
edidcwnststwvtygtcttgehrrekrsvvalvhvlgmletrtetwmssegawkhvqr
ietwilrhpgftimaailaytigtthfqralifilltavapsmtMRCIGISNRDFVEGVSG
GSWVDIVLHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCIEAKLTNTTCSRPTQG
EPSLNEEQDKRFVCKHSMVDRGWGNCGLFGKGGIVTCAMFTCKKNMEGK**X**VOPENLEYTI
VITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAELTGYGTVTMECSPTGLDFNEMVLLQ
MEDKAWLVHRQWFLDLPLPWLPGADTQGSNWIQKETLVTFKNPHAKQDVVVLGSQEGAMH
TALTGATEIQMSSGNLLFTGHLKCRMDKLQLKGMSYSMCTGKFK**Z**VKEIAETQHGTIVI
RVQYEGDGSCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVNIEAEPPFGDSYIIIGVE
PGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGGVFTSIGKALHQVFGAIY
GAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVSLVGVVTLYLGVMVQA (SEQ ID
NO: 3)

35

The global consensus sequence for the E sequence is represented as follows:

MRCIGISNRDFVEGVSGGSWVDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCI
EAKLTNTTTESRCPTQGEPSLNEEQDKRKFVCKHSMVDRGWNGCGLFGKGGIVTCAMFTCK
KNMEGK**X**VQOPENLEYTIVITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAELTGYGTVM
ECSPTRTGLDFNEMVLLQMEDKAFLVHRQWFLDLPLPWLGADTQGSNWIQKETLVTFKNPH
5 AKKQDVVVLGSQEGAMHTALTGATEIQMSSGNLLFTGHLKRLRMDKLQLKGMSYSMCTGK
FK**Z**VKEIAETQHGTIVIRVQYEGDGSPCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVN
IEAEPFGDSYIIIGVEPGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGG
VFTSIGKALHQVFGAIYGAASFSGVSWTMKILIGVIITWIGMNSRSTSLSVSLVLGVVTLY
LGVMVQA (SEQ ID NO: 12)

10

In the above sequence, the global consensus prM sequence is shown in lower case letters and the E sequence is shown in upper case letters. The amino acid positions denoted as X (position 129 of the E sequence) and Z (position 308 of the E sequence) are each independently Val or Ile, i.e. the proportion of aligned amino acid sequences including Val or Ile at those positions is close to 50%.

Determination of minor amino acid residues and analysis of the Chimerivax™ CYD2 sequence

A list of variable amino acid positions is established from the global alignment containing all amino acid positions varying in at least 5% of the aligned sequences. In addition, any amino acid from the sequence of the prM and E proteins of Chimerivax™ CYD2 that do not match the global consensus sequence are also identified. The results are shown in Table 4 (N.B., in the table, the prM and M protein sequences are numbered together, i.e. from position 1 to position 166 of the total sequence and E is numbered separately from position 1 to position 495).

25


prM				E			
Position	Consensus	%	CYD	%	Other variants		
15	G	76	S	24		I<1	
16	R	76			A16; D4		
24	L	99	V	<1			
29	D	91			N8; E-V-H<1		
31	V	94			T4; I2; M-D<1		
39	I	58	M	40		L2	
52	K	91			N9; T<1		
55	L	93			F7; R<1		
57	R	93			K8		
82	T	90			A9; S1; I-V<1		
120	V	55	N	43		A45	
125	T	99	I	<1		N-S<1	
127	I	94			V6; F<1		
134	T	95			A5; I-S<1		
148	H	90			Y9; N-D<1		
152	A	70			V28; T1; I<1		
							
52	Q	83				H15; E2; L<1	
61	I	93				V6; M-K-F-T<1	
71	E	76				A19; D5; P<1	
83	N	73				K25; V1; S-A-D<1	
91	V	67				I31; L1	
129	I	50	V	50		F-T<1	
131	Q	83				L17; E-H-P<1	
141	I	72	V	28		L<1	
149	H	80				N19; Y-R-Q-P-S-T<1	
160	K	94				Q3; E2; M1; R-N<1	
162	I	94				V6; L<1	
164	I	55	V	45			
203	D	49	N	46		E4; S1; K-G<1	
226	T	84				K16; I-E-P<1	
228	G	86				E14	
251	V	99	F	<1		I<1	
308	V	52				I48; L<1	
340	M	80				T19; I-A-L<1	
346	H	74				Y26; Q-L<1	
359	T	95				A4; I2; M-P<1	
462	I	78				V22; T<1	
484	V	69				I31; F-A-L-T<1	
485	V	94				I6; L<1	
491	V	62				A38; G-L<1	

Table 4: Dengue virus serotype 2 variable residues and CYD2 comparison

5

A total of 41 amino acid positions are identified in the prM and E sequences which either vary from the global consensus sequence in at least 5% of the aligned sequences and/or differ from the sequence of the prM and E proteins in CYD2. Ten amino acid positions in the sequence of the prM and E proteins in CYD2 differ from the global consensus sequence (5 positions in E, 2 positions in M and 3 in its precursor part, see 10 Table 4). Five out of the ten differing residues present a variation distribution close to 50:50, suggesting a naturally variable position. Only three positions in the CYD2 prM-E sequence appear as very minor variants (pr-24 Val, M-125 Ile and E-251 Phe).

Impact analysis of variations in the E and M proteins

15 To gain further insight into the variable positions, changes in the E protein ectodomain (amino acids 1-395), the most important domain for the seroneutralisation by the immune system are further analysed.

Using information available from a published 3D structure of the soluble ectodomain of the E protein of a dengue virus of serotype 2 (Modis, Y., et al. (2003) Proc Natl Acad Sci U S A, 100(12):6986-6991), a 3D model of the Dengue virus particle surface is reconstructed. This allows a fine tuned assessment of the accessibility of each amino acid from the E ectodomain, which in turn is used in association with the variability level and the nature of the amino acid change to assess a potential impact of CYD2 variations on antibody recognition.

The analysis demonstrates that two variations in the Chimerivax™ CYD2 sequence from the global consensus sequence (Val 141 and Val 164 of the E protein) are completely buried in the 3D structure and so cannot directly interact with an antibody at the surface of the virion. Position 129 of the E protein is a 50:50 variable amino acid position between Val (Chimerivax™ CYD2) and Ile (global consensus sequence) and the substitution is also a fully conservative change. The potential impact of these variations is therefore considered as very limited.

The variation at position 203 of the E protein (Asn in Chimerivax™ CYD2 and Asp in the global consensus sequence) could potentially have an impact (well exposed residue, change of charge) but the distribution of the variation among strains is close to 50:50, suggesting a naturally variable position.

The variation at position 251 of the E protein of Chimerivax™ CYD2 (Phe in Chimerivax™ CYD2 and Val in the global consensus sequence) is extremely rare among retrieved strains. Such a variation could have some impact on recognition by an antibody, as it is rare, rather well exposed at the surface of the virion (29%) and corresponds to a non-conservative amino acid change.

The modeling analysis described above identifies two other position variations in the E protein that could have a potential impact on antibody recognition (positions 226 and 228), although Chimerivax™ CYD2 does not vary from the global consensus sequence at those positions. Therefore in identifying optimised serotype 2 strains, variations from the global consensus sequence at those positions (i.e. Thr at position 226 and glycine at position 228) are preferably avoided for a universal dengue 2 vaccine.

Without being bound by theory, the present inventors consider that the impact of amino acid variations in dengue virus sequences can also be assessed using a scoring method which takes into account a number of relevant factors. In particular this method takes into account the genome location of the variation (G), the nature of the amino acid change (B), 3D mapping (M) and known variants at the position in question (DB), wherein the score is calculated as $G \times B \times M \times DB$. A score of 0 would be classified as no expected impact, a score of >0 to 10 would be classified as a low expected impact, a

score of >10 to 25 would be classified as a median expected impact and a score of >25 would be classified as a high expected impact.

The genome location (G) score is 0 if the amino acid is located in the M part of the prM/M protein (i.e. position 92 to 166 of the prM/M sequence) or in position 396 to 495 of the E protein. The genome location score is 1 if the amino acid is located in prM part of the prM/M protein (i.e. position 1 to 91 of the prM/M sequence) or in position 1 to 395 of the E protein.

The score related to the nature of the amino acid change (B) is calculated as $B = 100 - [(Blosum95 \text{ score} + 6) \times 10]$, wherein the Blosum95 score for different amino acid substitutions is as shown in Table 5 below.

Table 5

	A	R	N	D	C	Q	E	G	H	I	L	K	M	F	P	S	T	W	Y	V	B	Z	X	*
A		-2	-2	-3	-1	-1	-1	-3	-2	-2	-1	-2	-3	-1	1	0	-4	-3	-1	-3	-1	-3	-1	-6
R	-2		-1	-3	-5	0	-1	-4	-1	-4	-3	2	-2	-4	-3	-2	-2	-4	-3	-4	-2	-1	-2	-6
N	-2	-1		1	-4	0	-1	-1	0	-4	-5	0	-3	-4	-3	0	-1	-5	-3	-4	4	-1	-2	-6
D	-3	-3	1		-5	-1	1	-2	-2	-5	-5	-2	-5	-5	-3	-1	-2	-6	-5	-5	4	0	-2	-6
C	-1	-5	-4	-5		-4	-6	-5	-5	-2	-3	-5	-3	-3	-5	-2	-2	-4	-4	-2	-4	-5	-3	-6
Q	-1	0	0	-1	-4		2	-3	1	-4	-3	1	-1	-4	-2	-1	-1	-3	-3	-3	-1	4	-1	-6
E	-1	-1	-1	1	-6	2		-3	-1	-4	-4	0	-3	-5	-2	-1	-2	-5	-4	-3	0	4	-2	-6
G	-1	-4	-1	-2	-5	-3	-3		-3	-6	-5	-3	-4	-5	-4	-1	-3	-5	-5	-5	-2	-3	-3	-6
H	-3	-1	0	-2	-5	1	-1	-3		-4	-4	-1	-3	-2	-3	-2	-2	-3	1	-4	-1	0	-2	-6
I	-2	-4	-4	-5	-2	-4	-4	-6	-4		1	-4	1	-1	-4	-3	-2	-4	-2	3	-5	-4	-2	-6
L	-2	-3	-5	-5	-3	-3	-4	-5	-4	1		-3	2	0	-4	-3	-2	-3	-2	0	-5	-4	-2	-6
K	-1	2	0	-2	-5	1	0	-3	-1	-4	-3		-2	-4	-2	-1	-1	-5	-3	-3	-1	0	-1	-6
M	-2	-2	-3	-5	-3	-1	-3	-4	-3	1	2		-2		-1	-3	-1	-2	-3	0	-4	-2	-2	-6
F	-3	-4	-4	-5	-3	-4	-5	-5	-2	-1	0	-4		-1	-5	-3	-3	0	3	-2	-5	-4	-2	-6
P	-1	-3	-3	-3	-5	-2	-2	-4	-3	-4	-4	-2	-3	-5		-2	-2	-5	-5	-4	-3	-2	-3	-6
S	1	-2	0	-1	-2	-1	-1	-1	-2	-3	-3	-1	-3	-3	-2	1	-4	-3	-3	-1	-1	-1	-6	
T	0	-2	-1	-2	-2	-1	-2	-3	-2	-2	-2	-1	-1	-3	-2	1	-4	-3	-1	-1	-2	-1	-6	
W	-4	-4	-5	-6	-4	-3	-5	-5	-3	-4	-3	-5	-2	0	-5	-4	-4	2	-3	-6	-4	-4	-4	-6
Y	-3	-3	-3	-5	-4	-3	-4	-5	1	-2	-2	-3	-3	3	-5	-3	-3	2	-3	-3	-4	-4	-2	-6
V	-1	-4	-4	-5	-2	-3	-3	-5	-4	3	0	-3	0	-2	-4	-3	-1	-3	-3		-5	-3	-2	-6
B	-3	-2	4	4	-4	-1	0	-2	-1	-5	-5	-1	-4	-5	-3	-1	-1	-6	-4	-5	0	-2	-6	
Z	-1	-1	-1	0	-5	4	4	-3	0	-4	-4	0	-2	-4	-2	-1	-2	-4	-4	-3	0	1	-1	-6
X	-1	-2	-2	-2	-3	-1	-2	-3	-2	-2	-2	-1	-2	-2	-3	-1	-1	-4	-2	-2	-2	-1	1	-6
*	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	-6	

B = Asx, Z = Glx, X = Any and * = Stop

The M value depends on whether the amino acid is or is not located at the prM/E interface. For example, for CYD2 as used in Example 1, the amino acids that are located at the interface are prM residues 6, 7, 39, 40, 46-54, 56, 59-65, 67, 74 and 77 and E residues 64-72, 82-84, 101-104, 106-108 and 244-247. Where an amino acid is located at the interface, M equals 1. Where an amino acid is not located at the interface, M = Y x

SAS %. Y is 1 if the amino acid is located in an "up" position (i.e. directed towards the external environment); Y is 0.5 if the amino acid is located on the "side" of the molecule (i.e. the amino acid is neither directed towards the external environment nor towards the capsid) and Y is 0 if the amino acid is located in a "down" position (i.e. directed towards the capsid). The solvent accessibility surface % (SAS %) value is generated using the Discovery Studio 3D modeling software (Accelrys, Inc., CA, USA).

The DB value is 0 when the amino acid substitution results in an amino acid at the substitution position which is the most common amino acid at that position in the dengue sequences present in the GenBank database (<http://www.ncbi.nlm.nih.gov>). The DB value is 0.25 when the amino acid substitution results in an amino acid at the substitution position which is found in more than 5% of the dengue sequences present in the database (but is not the most common amino acid at that position). The DB value is 0.50 when the amino acid substitution results in an amino acid at the substitution position which is found in less than 5% of the dengue sequences present in the database (except unique substitutions). The DB value is 1 when the substitution amino acid is unique.

During replication, viruses may acquire a mutation leading to an amino acid substitution. The above-mentioned method provides a means to determine the effect of such mutations on the progeny of the mutated viruses.

Preferred sequences (i.e. sequences that are considered to be satisfactorily close to the identified consensus sequence) may have: (i) at most two, preferably one or no high-impact amino acid substitutions; (ii) at most three, preferably two or one, or no median impact amino acid substitutions; and/or (iii) at most five, four, three, two or one low impact amino acid substitutions.

Identification of optimized serotype 2 strains

Optimised serotype 2 strains are identified on the basis of the selection criteria described above.

A BLAST search is conducted to identify the strain having the closest sequence to the prM-E global consensus sequence in all of the available sequences. No sequence that is 100% identical to the prM-E global consensus sequence is found, but the best hit is a sequence from strain BID-V585 (NCBI Protein ID no. ACA58343; Genome ID no. EU529706; isolated from Puerto Rico in 2006) which shows only one variation from the global consensus sequence, at position 91 (Val in the global consensus sequence and Ile in BID-V585). The BID-V585 prM-E sequence contains 13 variations from the Chimerivax™ CYD-2 prM-E sequence.

A further strain selection is made so as to provide geographical balance in strain origin. Therefore a recently isolated Asian strain showing a good score in the BLAST analysis (strain MD-1280; NCBI Protein ID no. CAR65175; Genome ID no. FM21043; isolated from Viet Nam in 2004) is selected. Despite showing 6 variations with the global consensus sequence across prM-E, 3 of the 6 variations are identified as versatile positions naturally varying in more than 30% of the strains. The MD-1280 prM-E sequence contains 15 variations from the Chimerivax™ CYD-2 prM-E sequence.

A further strain selection is made on the basis of a large amount of previously accumulated experience with the strain. It is the PDK53-16681 strain, also known as the LAV-2 strain, a live-attenuated virus derived from Dengue serotype 2 16681 strain from Mahidol University (NCBI Protein ID no. AAA73186; Genome ID no. M84728; isolated from Thailand in 1964; Blok,J., et al. (1992); Virology 187 (2), 573-590). The LAV-2 prM-E sequence contains 10 variations from the global consensus sequence and 13 variations from the Chimerivax™ CYD-2 prM-E sequence.

A further strain selected on the basis of the above-mentioned criteria is strain PR/DB023 (NCBI Protein ID no. AEN71248; Genome ID no. JF804036; isolated from Puerto Rico in 2007). The PR/DB023 prM-E sequence contains 3 variations from the global consensus sequence and 13 variations from the Chimerivax™ CYD-2 prM-E sequence.

None of the selected strains contain the rare amino acids present in the Chimerivax™ CYD-2 prM-E sequence, i.e. Val at pr-24, Ile at M-125 and Phe at E-251.

PrM to E nucleotide sequences of the four selected strains

25 >LAV-2 prME nucleotide sequence (SEQ ID NO: 4)

ttccatctaaccacacgtaacggagaaccacacatgatcgtagcagcacaagagaaaaggaa
aaagtcttctgttaaaacagaggtggcgtgaacatgtgtaccctcatggccatggacact
tggtaattgtgtgaagacacaatcacgtacaagtgtccctctcaggcagaatgagcca
gaagacatagactgttggtgcaactctacgtccacgtggtaacttatggacgtgtacca
ccatggagaacatagaagagaaaaagatcagtggcactcgttccacatgtggaaatggg
actggagacacgaactgaaacatggatgtcatcagaaggggctggaaacatgtccagaga
attgaaacttggatcttggagacatccagggttccacatgtggcagaatcctggcataca
ccataggaacgacacattccaaagagccctgatttcatcttactgacagctgtcactcc
ttcaatgacaATGCGTTGCATAGAATGTCAAATAGAGACTTGTGGAAGGGTTTCAGGA
30 GGAAGCTGGTTGACATAGTCTTAGAACATGGAAGCTGTGACGACGATGGCAAAAACA
AACCAACATTGGATTGGAACTGATAAAAACAGAAAGCCAAACAGCCTGCCACCTAACGGAA
35 GTACTGTATAGAGGCAAAGCTAACCAACACAACAGAAATCTCGCTGCCAACACAAAGGG
GAACCCAGCCTAAATGAAGAGCAGGACAAAGGTTCGTCTGCAAACACTCCATGGTAGACA
GAGGATGGGAAATGGATGTGGACTATTGGAAAGGGAGGCATTGTGACCTGTGCTATGTT
40 CAGATGCAAAAGAACATGGAAGGAAAGTTGTGCAACCAGAAAACCTGGAATACACCATT

GTGATAACACCTCACTCAGGGGAAGAGCATGCAGTCGGAAATGACACAGGAAAACATGGCA
 AGGAAATCAAATAACACCAACAGAGTCCATCACAGAAGCAGAATTGACAGGTTATGGCAC
 TGTACAATGGAGTGCCTCCAAGAACGGGCCTCGACTTCAATGAGATGGTGTGCTGCAG
 ATGGAAAATAAAGCTTGGCTGGTCACAGGCAATGGTCTAGACCTGCCGTACCATGGT
 5 TGCCCGGAGCGGACACACAAGGGTCAAATTGGATACAGAAAGAGACATTGGTCACTTCAA
 AAATCCCCATGCGAAGAACAGGATGTTGTTAGGATCCAAAGAAGGGCCATGCAC
 ACAGCACTTACAGGGGCACAGAAATCCAATGTCATCAGGAAACTACTCTCACAGGAC
 ATCTCAAGTGCAGGCTGAGAATGGACAAGCTACAGCTAAAGGAATGTCATACTCTATGTG
 CACAGGAAAGTTAAAGTTGAAGGAAATAGCAGAAACACAATGGAACAATAGTTATC
 10 AGAGTGCAATATGAAGGGGACGGCTCCATGCAAGATCCCTTTGAGATAATGGATTGG
 AAAAAAGACATGTCTAGGTCGCCTGATTACAGTCACCCAAATTGTGACAGAAAAAGATAG
 CCCAGTCACATAGAACAGAACCTCCATTGGAGACAGCTACATCATAGGAGTAGAG
 CCGGGACAACACTGAAGCTCAACTGTTAAGAAAGGAAGTTCTATCGGCCAAATGTTGAGA
 CAACAATGAGGGGGCGAAGAGAATGGCCATTTAGGTGACACAGCCTGGGATTGGATC
 15 CTTGGGAGGAGTGTACATCTATAGGAAAGGCTCTCCACCAAGTCTTGGAGCAATCTAT
 GGAGCTGCCTTCAGTGGGTTTCATGGACTATGAAAATCCTCATAGGAGTCATTATCACAT
 GGATAGGAATGAATTCACGCAGCACCTCACTGTCGTGACACTAGTATTGGTGGGAATTGT
 GACACTGTATTGGGAGTCATGGTGCAGGCC
UPPERCASE: E coding sequence; lowercase: prM coding sequence

20

> BID/V585 – prME nucleotide sequence (SEQ ID NO: 5)

ttccatttaaccacacgtaatggagaaccacacatgatcggtggtaggcaagagaaaaggaa
 aaagtcttctgtttaaaacacagaggatgggttaacatgtgcaccctcatggccatagaccc
 25 tggtgaattgtgtgaagatacaatcacgtacaatgtgcggccctcctcaggcaaaatgaacca
 gaagacatagattgttggcaactctacgtccacatggtaacttatggacatgtacca
 ccacaggagaacacagaagagaaaaagatcagtggcaactcggtccacatgtgggcatgg
 actggagacacgaactgaaacatggatgtcatcagaagggcctggaaacatgtttcagaga
 attgaaacctggatcttggagacatccaggcttaccataatggcagcaatcctggcatata
 30 ccataggaacgacacattccaaaggctctgatcttactgacagccgttgc
 ttcaatgacaATGCCTTGCATAGGAATATCAAATAGAGACTTCGTAGAAGGGTTTCAGGA
 GGAAGTTGGGTTGACATAGTCTTAGAACATGGAAGTTGTGACGACGATGGAAAAATA
 AACCAACATTGGATTTCAGTAAACAGAACAGCCAAACACCTGCCACTCTAAGGAA
 GTACTGTATAGAAGCAAAGCTGACCAATACAACACAGAACACTCGTGTGCCAACACAAGGG
 35 GAACCCAGTCTAAATGAAGAGCAGGACAAAAGGTTCATCTGCAAACACTCCATGGTAGACA
 GAGGATGGGAAATGGATGTGGATTATTGGAAAGGGAGGCATTGTGACCTGTGCTATGTT
 CACATGCAAAAGAACATGGAAGGAAAAGTCGTGCAGCCAGAAAATCTGGAATACACCATC
 GTGATAACACCTCACTCAGGAGAAGAGCACGCTGTAGGTAATGACACAGGAAAGCATGGCA
 AGGAAATCAAATAACACCAACAGAGCTCCATCACAGAAGCAGAACTGACAGGCTATGGCAC
 40 TGTACAGATGGAGTGCCTCCGAGAACGGCCTCGACTTCAATGAGATGGTACTGCTGCAG
 ATGGAAGACAAAGCTTGGCTGGTCACAGGCAATGGTCTAGACCTGCCGTACCATGGC
 TACCCGGAGCGGACACACAAGGATCAAATTGGATACAGAAAGAGACGTTGGTCACTTCAA
 AAATCCCCACGCGAAGAACAGGACGTCGTTAGGATCTCAAGAACGGCCATGCAC
 ACGGCACCTACAGGGGCACAGAAATCCAGATGTCATCAGGAAACTACTGTCACAGGAC
 ATCTCAAGTGTAGGCTGAGAATGGACAAATTACAGCTAAAGGAATGTCATACTCTATGTG
 45 TACAGGAAAGTTAAAATTGTGAAGGAAATAGCAGAAACACAATGGAACAATAGTTATC
 AGAGTACAATATGAAGGGGACGGCTCCATGTAAGATTCTTTGAGATAATGGATTGG
 AAAAAAGACACGTCCTAGGTCGCCTGATTACAGTGAACCCAATCGTAACAGAAAAAGATAG
 CCCAGTCACATAGAACAGAACCTCCATTGGAGACAGCTACATCATAGGAGTAGAG
 CCGGGACAATTGAAACTCAATTGGTCAAGAACAGGAAAGTCCATTGGCCAAATGTTGAGA

CAACAATGAGAGGAGCGAAGAGAATGGCCATTTAGGTGACACAGCCTGGGATTTGGATC
 CCTGGGAGGAGTGTACATCTATAGGAAAGGCTCTCCACCAAGTTTCGGAGCAATCTAT
 GGGGCTGCTTAGTGGGTCTCATGGACTATGAAAATCCTCATAGGAGTTATTATCACAT
 GGATAGGAATGAATTACGTAGCACCTCACTGTCTGTCACTAGTATTGGTGGGAGTCGT
 5 GACACTGTACTTGGGGTTATGGTGCAGGCT

>PR/DB023 prME nucleotide sequence (SEQ ID NO: 6)

ttccathtaaccacacgtaatggagaaccacacatgatcgttgttaggcaagagaaaaggga
 aaagtcttctgttcaaaacagaggatggtgttaacatgtgtaccctcatggccatagacct

10 tggtgaattgtgtgaagatacaatcacgtacaagtgcggccctcctcaggcaaaatgaacca
 gaagacatagattgttgtgcaactctacgtccacatggtaacttatgggacatgtacca
 ccacaggagaacacagaagagaaaaagatcagtggcactcgttccacatgtgggcatgg
 actggagacacgaactgaaacatggatgtcatcagaaggcccctggaaacatgttcagaga
 attgaaacctggatattgagacatccaggcttaccataatggcagcaatcctggcatata

15 ccataggaacgacacatccaaaggcgtctgatcttactgacagcgcgtcgctcc
 ttcaatgacaATGCGTTGCATAGGAATATCAAATAGAGACTTCGTAGAAGGGGTTTCAGGA
 GGAAGTTGGGTTGACATAGTCTAGAACATGGAAGTTGTGACGACGATGGCAAAAATA
 AACCAACATTGGATTTGAACTGATAAAAACAGAACGCCAACACTGCCACTCTAAGGAA

20 GTACTGTATAGAAGCTGACCAATACAACACAGAACACTCGTGCCTAACACAAAGGG
 GAACCCAGTCTAAATGAAGAGCAGGACAAAAGGTTCATCTGCAAACACTCCATGGTAGACA
 GAGGATGGGAAATGGATGTGGATTATTGGAAAAGGAGGATTGTAAACCTGTGCTATGTT
 CACATGCAAAAGAACATGGAAGGAAAAGTTGTGCTGCCAGAAAATCTGGAATACACCATC
 GTGATAACACCTCACTCAGGAGAAGAGCACGCTGTAGGTAATGACACAGGAAAACATGGCA
 AGGAAATTAAAATAACACCACAGAGTTCCATCACAGAACAGAACACTGACAGGGCTATGGCAC

25 TGTCACGATGGAGTGTCTCCGAGAACGGGCTCGACTTCATGAGATGGTGTGCTGCAG
 ATGGAAGACAAAGCCTGGCTGGCACAGGCAATGGTCTAGATCTGCCGTACCATGGC
 TACCCGGAGCGGACACACAAGGATCAAATTGGATACAGAAAGAGACGTTGGTCACTTCAA
 AAATCCCCACGCGAAGAACAGGACGTCGTTAGGATCTCAAGAAGGGCCATGCAC
 ACGGCACCTACAGGGCCACAGAAATCCAGATGTCACTCAGGAAACTACTGTTCACAGGAC

30 ATCTCAAGTGTAGGCTGAGAATGGACAAATTACAGCTTAAAGGAATGTCAACTCTATGTG
 TACAGGAAAGTTAAAATTGTGAAGGAAATAGCAGAAACACACATGGAACAAATAGTTATC
 AGAGTACAATATGAAGGGGACGGCTCTCCATGTAAGATTCTTTGAGATAATGGATTAG
 AAAAAGACACGTCCTAGGTCGCCTGATTACAGTGAACCCAATCGTAACAGAAAAAGATAG
 CCCAGTCAACATAGAACGAGAACCTCCATTGGAGACAGCTACATCATCAGGAGTAGAG

35 CCGGGACAATTGAAACTCAATTGGTCAAGAACAGGAAGTTCCATTGGCAAATGTTGAGA
 CAACAATGAGAGGAGCGAAGAGAATGGCCATTTAGGTGACACAGCCTGGGATTTGGATC
 CCTGGGAGGAGTGTACATCTATAGGAAAGGCTCTCCACCAAGTTTCGGAGCAATCTAT
 GGGGCTGCTTAGTGGGTCTCATGGACTATGAAAATCCTCATAGGAGTTATCATCACAT
 GGATAGGAATGAATTACGTAGCACCTCACTGTCTGTCACTAGTATTGGTGGGAGTCGT

40 GACACTGTACTTGGGGTTATGGTGCAGGCT

>MD1280 prME nucleotide sequence (SEQ ID NO: 7)

ttccathtaaccacacgaaatggagaaccacacatgatcgttgttaggcaagagaaaaggga
 aaagccttctgttaaaacagaggatggtgtgaacatgtgtaccctcatggccattgtatct

45 tggtgaattgtgtgaagatacaatcacgtacaagtgcggccctcctcaggcagaatgaacca
 gaagatagattgttgtgcaactccacgtccacatggtaacttatgggacgtgtacca
 ccacaggagaacacagaagagaaaaagatcagtggcactcgttccacatgtgggtatgg
 actggagacacgaactgaaacatggatgtcgtcagaaggcccctggaaacacgcgtcagaga

attgaaacttgatcttggatccaggcttaccataatggcagcaatcctggcatata
 ccgttaggaacgacacattccaaaggccctgatttcatcttactggcagctgtcgctcc
 ttcaatgacaATGC GTTGCATAGGAATATCAAATAGAGACTTGTAGAAGGGTTTCAGGA
 GGAAGCTGGTTGACATAGTCTAGAACATGGAAGTTGTGACGACAATGGCAAAAATA
 5 AACCAACACTGGATTTGAAGTGTGACAAAAACAGAACGCAAACAAACCTGCCACTCTAAGGAA
 GTACTGTATAGAGGCAAAGCTGACCAATACAACAAACAGAACACTCGTTGCCAACACAAGGG
 GAACCCAGTCTAAATGAAGAGCAGGACAAAAGGTTCGTCTGCAAACACTCCATGGTAGACA
 GAGGATGGGAAATGGATGTGGATTATTGGAAAGGGAGGCATTGTGACCTGTGCTATGTT
 CACATGCAAAAAGAACATGGAAGGAAAATCGTCAACCAGAAAATTGGAATACACCATC
 10 GTGATAACACCTCACTCAGGAGAACGACGCTGTAGGTAATGACACAGGAAAACATGGTA
 AGGAAATTAAAATAACACCACAGAGTTCCATCACAGAACGAGACTGACAGGGCTATGGCAC
 AGTCACGATGGAGTGCCTCCGAGAACGGGCTTGACTCAATGAGATGGTGCCTGCGAG
 ATGGAAGATAAAGCTTGCTGGTGCACAGGCAATGGTCCTAGACCTGCCGTACCATGGC
 TACCCGGAGCGGACACACAAGGATCAAATTGGATACAGAAAGAGACATTGGTCACTTCAA
 15 AAATCCCCACCGGAAGAACGAGGATGTCGTTTTAGGATCTCAAGAACGGAGCCATGCAC
 ACGGCACTCACAGGGCACAGAAATCCAGATGTCATCAGGAAACTTACTATTACAGGAC
 ATCTCAAATGCAGGCTGAGAATGGACAAACTACAGCTCAAAGGAATGTCATACTCTATGTG
 TACAGGAAAGTTAAAATTGTGAAGGAAATAGCAGAAACACAACATGGAACAATAGTTATC
 AGAGTACAATAATGAAGGAGACGGCTCTCCATGTAAGATCCCTTTGAAATAATGGATTGG
 20 AAAAAAGACATGTCCTAGGTCGCTGATTACAGTTAACAGATCGTAAACAGAAAAGATAG
 CCCAGTCACATAGAACGAGAACCTCCATTGGAGACAGCTACATCATTATAGGAGTAGAG
 CCGGGACAATTGAAACTCACTGGTCAAGAACAGGAAGTTCCATCGGCCAAATGTTGAGA
 CGACAATGAGAGGAGCAAAGAGAACATGGCCATTAGGTGACACAGCCTGGGATTTGGATC
 TCTGGGAGGAGTGTACATCTATAGGAAAGGCTCTCCACCAAGTTTCGGAGCAATCTAT
 25 GGGCCTGCCTTAGTGGGTTCATGGACTATGAAAATCCTCATAGGAGTCATCATCACAT
 GGATAGGAATGAATTACGTAGCACCTCACTGTCTGTGCACTAGTATTGGTGGGAATCAT
 AACACTGTACTTGGGAGCTATGGTGCAGGCT

Corresponding protein prM to E sequences of the four selected strains

30

>LAV2 prME protein sequence (SEQ ID NO: 8)

fhltrngephmivsrqekgksllfktevgvnmc1mamdlgelcedtitykcpl1rqnep
 edidcwnststwvtygtcttmgehrrekrsvvalvhvlgmletrtetwmssegawkhvqr
 ietwilrhpgftmmaailaytigtthfqralifilltavtpsmtMRCIGMSNRDFVEGVSG
 35 GSWDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCIEAKLTNTTRESRCPTQG
 EPSLNEEQDKRFVCKHSMVDRGWNGCGLFGKGGIVTCAMFRCKKNMEGKVVQOPENLEYTI
 VITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAEALTGYGTVTMECSPRTGLDFNEMVLLQ
 MENKAWLVHRQWFLDLPLPWLPAGDTQGSNWIQKETLVTFKNPHAKQDVFVVLGSQEGAMH
 TALTGATEIQMSSGNLLFTGHLKCRMDKLQLKGMSYSMCTGKFVVKEIAETQHGTIVI
 40 RVQYEGDGSPCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVNIEAEPPFGDSYIIIGVE
 PGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGGVFTSIGKALHQVFGAIY
 GAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVTLVGIITLYLGVMVQA

45

>LAV2 E protein sequence (SEQ ID NO: 13)

MRCIGMSNRDFVEGVSGGSWVDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCI
 EAKLTNTTRESRCPTQGEPSLNEEQDKRFVCKHSMVDRGWNGCGLFGKGGIVTCAMFRCK
 KNMEGKVVQOPENLEYTIVITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAEALTGYGTVT
 ECSPRTGLDFNEMVLLQMENTKAWLVHRQWFLDLPLPWLPAGDTQGSNWIQKETLVTFKNPH

AKKQDVVVLGSQEGAMHTALTGATEIQMSSGNLLFTGHLKRLRMDKLQLKGMSYSMCTGK
 FKVVKEIAETQHGTIVIRVQYEGDGSPCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVN
 IEAEPFGDSYIIIGVEPGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGG
 VFTSIGKALHQVFGAIYGAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVTLVLGIVTLY
 5 LGVMVQA

>LAV2 M protein sequence (SEQ ID NO: 19)

svalvhvgmgletrtewmssegawkhvqrietwilrhpgftmmaailaytigtthfqra
 lifilltavtpsm

10

>BID/V585 prME protein sequence (SEQ ID NO: 9)

fhltrngephmivgrqekgksllfktdgvmctlmaidlgelcedtitykcpllrqne
 edidcwnststwvtygtcttgehrrekrsvlphvgmgletrtewmssegawkhvqr
 ietwilrhpgftimaailaytigtthfqralifilltavapsmtMRCIGISNRDFVEGVSG
 15 GSWDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCIEAKLTNTTTESRCPTQG
 EPSLNEEQDKRFICKHSMVDRGWNGCGLFGKGGIVTCAMFTCKKNMEGKVVQOPENLEYTI
 VITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAELTGYGTVMCESPRTGLDFNEMVLLQ
 MEDKAWLVRQWFLDLPLPWLPAGDTQGSNWIQKETLVTFKNPHAKQDVVVLGSQEGAMH
 TALTGATEIQMSSGNLLFTGHLKRLRMDKLQLKGMSYSMCTGKFKIVKEIAETQHGTIVI
 20 20 RVQYEGDGSPCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVNIEAEPFGDSYIIIGVE
 PGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGGVFTSIGKALHQVFGAIY
 GAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVSLVLGVVTLYLGVMVQA

>BID/V585 E protein sequence (SEQ ID NO: 14)

25 MRCIGISNRDFVEGVSGGSWVDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCI
 EAKLTNTTTESRCPTQGEPSLNEEQDKRFICKHSMVDRGWNGCGLFGKGGIVTCAMFTCK
 KNMEGKVVQOPENLEYTIITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAELTGYGTVM
 ECSPRTGLDFNEMVLLQMEDKAWLVRQWFLDLPLPWLPAGDTQGSNWIQKETLVTFKNPH
 AKKQDVVVLGSQEGAMHTALTGATEIQMSSGNLLFTGHLKRLRMDKLQLKGMSYSMCTGK
 30 FKIVKEIAETQHGTIVIRVQYEGDGSPCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVN
 IEAEPFGDSYIIIGVEPGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGG
 VFTSIGKALHQVFGAIYGAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVSLVLGVVTLY
 LGVMVQA

35

>BID/V585 M protein sequence (SEQ ID NO: 20)

svalvhvgmgletrtewmssegawkhvqrietwilrhpgftimaailaytigtthfqra
 lifilltavapsmt

>PR/DB023 prME protein sequence (SEQ ID NO: 10)

40 fhltrngephmivgrqekgksllfktdgvmctlmaidlgelcedtitykcpllrqne
 edidcwnststwvtygtcttgehrrekrsvlphvgmgletrtewmssegawkhvqr
 ietwilrhpgftimaailaytigtthfqralifilltavapsmtMRCIGISNRDFVEGVSG
 GSWDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCIEAKLTNTTTESRCPTQG
 EPSLNEEQDKRFICKHSMVDRGWNGCGLFGKGGIVTCAMFTCKKNMEGKVVLPENLEYTI
 45 VITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAELTGYGTVMCESPRTGLDFNEMVLLQ
 MEDKAWLVRQWFLDLPLPWLPAGDTQGSNWIQKETLVTFKNPHAKQDVVVLGSQEGAMH

TALTGATEIQMSSGNLLFTGHLKCRRLMDKLQLKGMSYSMCTGFKIVKEIAETQHGTIVI
 RVQYEGDGSPCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVNIEAEPPFGDSYIIIGVE
 PGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGGVFTSIGKALHQVFGAIY
 GAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVSLVGVVTLYLGVMVQA

5

>PR/DB023 E protein sequence (SEQ ID NO: 15)

MRCIGISNRDFVEGSGGSWVDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCI
 EAKLTNTTTESRCPTQGEPSLNEEQDKRFICKHSMVDRGWNGCGLFGKGGIVTCAMFTCK
 KNMEGKVVL PENLEYTIVITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAEALTGYGTVM
 10 ECSPRTGLDFNEMVLLQMEDKAWLVRQWFLDLPLPWLPAGDTQGSNWIQKETLVTFKNPH
 AKKQDVVVLGSQEGAMHTALTGATEIQMSSGNLLFTGHLKCRRLMDKLQLKGMSYSMCTGK
 FKIVKEIAETQHGTIVIRVQYEGDGSPCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVN
 15 IEAEPPFGDSYIIIGVEPGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGG
 VFTSIGKALHQVFGAIYGAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVSLVGVVTLY
 LGVMVQA

>PR/DB023 M protein sequence (SEQ ID NO : 21)

svalvhvlgmgletrtewmssegawkhvqrietwilrhpgftimaailaytigtthfqra
 lifilltavapsmt

20

>MD1280 prME protein sequence (SEQ ID NO: 11)

fhlttrngephmivgrqekgksllfktdedgvnmctlmaidlgelcedtitykcpllrqnepe
 edidcwcnststwvtygtcttgehrrekrsvvalvhvlgmgletrtewmssegawkhqqr
 ietwilrhpgftimaailaytvgthfqralifillaavapsmtMRCIGISNRDFVEGVSG
 25 GSWVDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCIEAKLTNTTTESRCPTQG
 EPSLNEEQDKRFVCKHSMVDRGWNGCGLFGKGGIVTCAMFTCKKNMEGKIVQOPENLEYTI
 VITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAEALTGYGTVMECSPRTGLDFNEMVLLQ
 MEDKAWLVRQWFLDLPLPWLPAGDTQGSNWIQKETLVTFKNPHAKKQDVVVLGSQEGAMH
 TALTGATEIQMSSGNLLFTGHLKCRRLMDKLQLKGMSYSMCTGFKIVKEIAETQHGTIVI
 30 RVQYEGDGSPCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVNIEAEPPFGDSYIIIGVE
 PGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGGVFTSIGKALHQVFGAIY
 GAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVSLVGIITLYLGAMVQA

>MD1280 E protein sequence (SEQ ID NO: 16)

35 MRCIGISNRDFVEGSGGSWVDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCI
 EAKLTNTTTESRCPTQGEPSLNEEQDKRFVCKHSMVDRGWNGCGLFGKGGIVTCAMFTCK
 KNMEGKIVQOPENLEYTIVITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAEALTGYGTVM
 ECSPRTGLDFNEMVLLQMEDKAWLVRQWFLDLPLPWLPAGDTQGSNWIQKETLVTFKNPH
 AKKQDVVVLGSQEGAMHTALTGATEIQMSSGNLLFTGHLKCRRLMDKLQLKGMSYSMCTGK
 40 FKIVKEIAETQHGTIVIRVQYEGDGSPCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVN
 IEAEPPFGDSYIIIGVEPGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGG
 VFTSIGKALHQVFGAIYGAAFSGVSWTMKILIGVIITWIGMNSRSTSLSVSLVGIITLY
 LGAMVQA

45

>MD1280 M protein sequence (SEQ ID NO: 22)

svalvhvlgmgletrtewmssegawkhqrietwilrhpgftimaailaytvgthfqra
 lifillaavapsmt

>Consensus M sequence (SEQ ID NO: 17)

svalvpvhvgmgletrtetwmssegawkhvqrietwilrhpqftimailaytigtthfqralifilltavapsmt

5 **Example 3: Construction of the cDNA clones corresponding to the optimized serotype 2 chimeric viruses and production of the encoded viruses**

Construction of chimeric dengue viruses corresponding to the optimized serotype 2 strains is achieved using the Chimerivax™ technology substantially in accordance with the teaching of Chambers, et al. (1999, *J. Virology* 73(4):3095-3101). Reference may also 10 be made to international patent applications WO 98/37911, WO 03/101397, WO 07/021672, WO 08/007021, WO 08/047023 and WO 08/065315, which detail the analogous processes used to construct CYD-1, CYD2, CYD-3 and CYD-4. Briefly, however, chimeric dengue viruses corresponding to the optimized serotype 2 strains are 15 constructed as follows (N.B. the optimized chimeric dengue viruses are constructed using the genomic backbone of YF strain YF17D204 (YF-VAX(R), Sanofi-Pasteur, Swiftwater, PA, USA)).

Construction of plasmid pSP1101

Construction of the YF-VAX cDNA clone - pJSY2284.1 (pACYC YF-Vax 5-3)

A full-length infectious cDNA clone of YF-VAX is constructed. The full-length 20 infectious cDNA clone is based on the sequence of YF-VAX. A low copy number plasmid pACYC177 (New England Biolabs, Inc., Ipswich, MA, USA) is used to assemble the full-length cDNA clone.

A DNA sequence named as SP6 YF-Vax 5-3 is synthesized by GeneArt®. The 25 sequence of SP6 YF-Vax 5-3 is designed in a way to facilitate an easy assembly of a full-length YF-Vax cDNA clone. The sequence is 2897 bp long and comprises the Xma I-SP6 promoter, the YF-Vax 5'UTR, the capsid, prM, M, part of E which extends to the Apa I site followed by unique sites Mlu I-Sap I-Ngo MI-Aat II-Cla I for assembly, part of NS5 and further extended to 3' UTR followed by an Nru I site, which is used for run-off. This synthesized DNA sequence is flanked by EcoR V and Xho I sites. After digestion with 30 EcoR V/Xho I, this DNA fragment is then cloned into the Aat II/Xho I sites of low copy number plasmid pACYC177 to replace the 1615bp Aat II/Xho I fragment. The resulting plasmid pJSY2284.1 (pACYC YF-Vax 5-3) is confirmed by sequence analysis.

RT-PCR and cloning of the YF-Vax cDNA fragments spanning from the sites Apa I, Mlu I, Sap I, Ngo M1, Aat II and Cla I and assembly of a full-length infectious cDNA clone of YF-vax (pJSY2374.5)

The yellow fever vaccine YF-VAX is grown in Vero cells, and the virus particles are concentrated. The viral RNA of YF-VAX is extracted from the concentrated virus and the cDNA copy is made by reverse transcriptase. Five cDNA fragments as shown herein are PCR amplified, TOPO cloned, sequenced and compared to the sequence of YF-VAX 2003. The PCR errors found in each fragment are corrected by either site-directed mutagenesis or fragment switching. There are too many sequence differences found in Ngo M1-Aat II fragment after TOPO cloning, and therefore, this fragment is synthesized by GeneArt®. After final sequence confirmation, the five DNA fragments; Apa I-Mlu I, Mlu I-Sap I, Sap I-Ngo M1, Ngo M1-Aat II, and Aat II-Cla I are isolated and stepwise cloned into the unique sites Apa I, Mlu I, Sap I, Ngo M1, Aat II and Cla I in the plasmid pJSY2284.1 to obtain plasmid pJSY2374.5, which is confirmed to contain the correct sequence of YF-VAX full-length cDNA.

Construction of cDNA for optimized chimeric dengue virus derived from the LAV2 strain (pSP1101)

The strategy is to replace the prM and E genes of the YF-VAX® vaccine strain in the pJSY2374.5 plasmid containing the YF-VAX genome with those of the LAV2 strain, as done previously to build the CYD-1, CYD-2, CYD-3 and CYD-4 dengue vaccines, using the Chimerivax™ technology. The resulting plasmid is pSP1101.

In pJSY2374, restriction sites used for cloning are Xma I and Mlu I. These sites are located upstream and downstream of a 3000 bp fragment which contains: the SP6 promoter, YF17D 5'UTR, YF17D-capsid, YF17D-prM, YF17D-E and the N terminus of YF17D-NS1. A sequence corresponding to this fragment but instead containing the prM and E genes of LAV2 flanked by Xma I and Mlu I sites is synthesized by GeneArt® and cloned into plasmid pMK-RQ (GeneArt®, Life Technologies Ltd, Paisley, U.K.) to create plasmid pMK-RQ-Seq1. Plasmid pJSY2374.5 and pMK-RQ-Seq1 are digested by Xma I and Mlu I. The Xma I-Mlu I fragment from pMK-RQ-Seq1 is then inserted into plasmid pJSY2374.5 to form plasmid pSP1101. XL-10 Gold Ultracompetent bacteria (Agilent Technologies, CA, USA) are used for transformation, as they are suitable for large plasmids. In a second step, positive clones are transferred into One Shot® TOP10 *E. coli* (Life Technologies Ltd, Paisley, U.K.), which allows the amplification of large size plasmids in significant amounts.

Plasmid pSP1101 thus allows the expression of LAV2 strain prM and E proteins with a YF-VAX replication engine. The resulting chimeric virus is designated CYD-LAV. Sequencing analysis shows no mutation as compared to the original sequences.

Construction of corresponding plasmids for strains BID-V585, PR/DB023 and MD1280

An analogous strategy to that described above is used to build the plasmids corresponding to the serotype 2 strains BID-V585, PR/DB023 and MD1280. These plasmids are designated pSP1102 (BID-V585), pSP1103 (PR/DB023) and pSP1104 (MD1280). The resulting chimeric viruses generated from those plasmids are designated CYD-BID, CYD-PR and CYD-MD. Sequence analysis of the generated plasmids shows no mutations compared to the original sequences.

Generation of chimeric viruses from plasmids pSP1101, pSP1102, pSP1103 and pSP1104

10 10 *In vitro* transcription of RNA and generation of viruses is carried out as previously described (Guirakhoo F et al. J. Virol. 2001; 75:7290-304).

Example 4: Evaluation of the immunogenicity and viremia of the optimized serotype 2 chimeric viruses in a monkey model

15 15 Evaluation of immunogenicity and viremia in monkeys

Design of the study

Four groups each containing four Cynomolgus monkeys are defined. The four groups receive the following formulations (containing $5 \log_{10}$ CCID₅₀ of each CYD dengue serotype):

20 20 1. Control tetravalent formulation, i.e. a formulation comprising CYD-1, CYD-2, CYD-3 and CYD-4.

2. CYD-LAV tetravalent formulation, i.e. a formulation comprising CYD-1, CYD-3, CYD-4 and CYD-LAV.

3. CYD-MD tetravalent formulation, i.e. a formulation comprising CYD-1, CYD-3, CYD-4 and CYD-MD.

25 25 4. CYD-PR tetravalent formulation, i.e. a formulation comprising CYD-1, CYD-3, CYD-4 and CYD-PR.

Monkeys receive two doses two months apart, as previously described (Guy B et al., Am J Trop Med Hyg. 2009; 80(2):302-11).

30 30 **Results**

Immunogenicity (SN₅₀ neutralizing response) and viremia are determined as described in the Materials and Methods section of Guy B., et al., Am. J. Trop. Med. Hyg. 2009; 80(2): 302-11.

Table 6: SN₅₀ neutralizing responses in monkeys immunized with optimized chimeric dengue serotype 2 viruses

		PD1				PD2			
		DEN1	DEN2	DEN3	DEN4	DEN1	DEN2	DEN3	DEN4
control CYD TV	responders	4/4	2/4	1/4	4/4	4/4	1/4	4/4	4/4
	GMT	27	5	7	636	71	8	35	425
CYD-LAV TV	responders	4/4	2/4	4/4	4/4	4/4	4/4	4/4	4/4
	GMT	95	63	19	477	189	95	80	477
CYD-MD TV	responders	4/4	4/4	3/4	4/4	4/4	4/4	4/4	4/4
	GMT	33	100	8	63	35	63	16	100
CYD-PR TV	responders	3/4	4/4	1/4	4/4	4/4	4/4	4/4	4/4
	GMT	27	63	7	168	109	84	38	212

5 PD: Post-dose; TV: tetravalent formulation

No serotype 2 viremia is observed, regardless of the serotype 2 chimeric virus administered. In respect of immunogenicity responses against DEN2, the tetravalent formulations comprising CYD-LAV, CYD-MD and CYD-PR demonstrate a higher response 10 (both GMTs and number of responding animals) than the control formulation (see Table 6).

Example 5. Assessment of tetravalent dengue vaccine formulations in flavivirus-naïve adults in Mexico.

15

The objective of the present study was to compare the immunogenicity and viremia of a blended tetravalent dengue vaccine comprising CYD-1 (i.e. the particular Chimerivax dengue serotype 1 (CYD-1) strain generated from the prM and E sequences of DEN1 PU0359 (TYP 1 140)), VDV2, CYD-3 (i.e. the particular Chimerivax dengue serotype 3 CYD-3) strain generated from the prM and E sequences of DEN3 PaH881/88) and CYD- 20 4 (i.e. the particular Chimerivax dengue serotype 4 (CYD-4) strain generated from the prM and E sequences of DEN4 1228 (TVP 980)) with the immunogenicity and viremia of a tetravalent dengue vaccine comprising CYD-1, CYD-2 (i.e. the particular Chimerivax dengue serotype 2 (CYD-2) strain generated from the prM and E sequences of DEN2 PUO218), CYD-3 and CYD-4. See Example 1 for more detail concerning the particular CYD-1, CYD-2, CYD-3 and CYD-4 used in this study.

25 The relevant nucleotide and protein sequences of the VDV2 strain are as follows:

>VDV2 nucleotide sequence (SEQ ID NO: 24)

30 AGUUGUUAGUCUACGUGGACCGACAAAGACAGAUUCUUUGAGGGAGCUAGCUAAUGUAG UUCUAACAGUUUUUUAAUAGAGAGCAGAUCUCUGAUGAAUACCAACGGAAAAGGGCAA

AAACACGCCUUUCAAUAUGCUGAAACGCGAGAGAAACCGCGUGUCGACUGUGCAACAGCUG
 ACAAAAGAGAUUCUCACUUGGAUGCUGCAGGGACGAGGACAUAAAACUGUUCAUGGCC
 UGGUGGCGUUCUUCGUUUCUAAACAAUCCACCAACAGCAGGGAUUUGAAGAGAUGGG
 AACAAUUAACAAAGCUAUUAUGUUUUGAGAGGGUUCAGGAAGAGAGAUUGGAAGG
 5 AUGCUGAACAUUCAUGAAUAGGAGACGCAGAUCUGCAGGAUGAUCAUUAUGCUGAUUCCAA
 CAGUGAUGGCGUUCUUAACCACACGUACGGAGAACCAUGAUCGUCAGCAGACA
 AGAGAAAGGGAAAAGUCUUCGUUUAAAACAGAGGUUGGCGUGAACAUUGUGUACCCUCAUG
 GCCAUGGACCUUGGUGAAUUGUGUGAAGACACAAUCACGUACAAGUGUCCCCUUCUCAGGC
 AGAAUGAGCCAGAACAUAGACUGUUGGUGCAACCUACGUCCACGUGGGUAACUUAUGG
 10 GACGUGUACCACCAUGGGAGAACAUAGAAGAGAAAAAGAUUCAGUGGCACUCGUUCCACAU
 GUGCGAAUGGGACUGGAGACACGAACUGAAACAUGGAUGUCAUCAGAAGGGGCCUGGAAAC
 AUGUCCAGAGAAUUGAACUUGGAUCUUGAGACAUCCAGGUUCACCAUGAUGGCAGCAAU
 CCUGGCAUACACCAUAGAACGACACAUUCCAAAGAGGCCUGAUUUUCAUCUUACUGACA
 GCUGUCACUCCUCAUAGACAAUGCUGAUAGGUAAAAGAGACAUUUGUGGAAG
 15 GGGUUUCAGGAGGAAGCUGGGUUGACAUAGCUUAGAACAUAGGAAGCUGUGACGACGAU
 GGCAAAAAACAAACCAACAUUGGAAUUGAACUGUAACAGAACAGCCAAACAGCCUGCC
 ACCCUAAGGAAGUACUGUAUAGAGGCAAAGCUAACCAAACACAACAGAACUUCUGCUGCC
 CAACACAAGGGGAACCCAGCCUAAAUGAAGAGCAGGACAAAAGGUUCGUCUGCAAACACUC
 CAUGGUAGACAGAGGAUGGGAAUUGGAUGGUGACAUUUGGAAAGGGAGGCAUUGUGACC
 20 UGUGCUAUGUUCAGAUGAAAAAGAACAUAGGAAGGAAAAGUUGUGCAACCAGAAAACUUGG
 AAUACACCAUUGUGUAACACCUCACUCAGGGGAAGAGCAUGCAGUCGGAAAUGACACAGG
 AAAACAUGGCAAGGAAAUCAAAUAACACCACAGAGUUCUCAUCACAGAAGCAGAAUUGACA
 GGUUAUGGCACUGUCACAAUGGAGUGCUCUCCAAGAACGGCCUCGACUUCAUUGAGAUGG
 UGUUGCUGCAGAUGGAAAAGCUUUGGUGGCACAGGCAAUUGGUUCCUAGACCUGCC
 25 GUUACCAUGGUUGCCGGACACACAAGAGUAAAUGGAUACAGAAGGGAGACAUUG
 GUCACUUUCAAAAUCCCAUUGCAGAAACAGGAUGUUGUUGUUUAGGAUCCCAAGAAG
 GGGCCAUGCACACAGCACUUACAGGGCCACAGAAAUCCAAUGUCAUCAGGAAACUUACU
 CUUCACAGGACACUCAAGUGCAGGCUGAGAAUUGGACAAGCUACAGCUAAAGGAAUGUCA
 UACUCUAUGUGCACAGGAAAGUAAAAGUUGUGAAGGAAUAGCAGAAACACAACAUUGGAA
 30 CAAUAGUUAUCAGAGGCAAAUUGAAGGGACGGCUCUCCAUGCAAGAUCCUUUUGAGAU
 AAUGGAUUUGGAAAAAGACAUGUCUUAGGUUGCUCUUAAGUCAACCCAAUUGUGACA
 GAAAAAGAUAGCCCAGUCAACAUAGAAGCAGAACCUCCAUUUGGAGACAGCUACAUCAUCA
 UAGGAGUAGAGCCGGACAACUGAAGCUAACUGGUUUAAGAAAGGAAGUUCUACUGGCCA
 AAUGUUUGAGACAACAAUAGAGGGGGCGAAGAGAAUUGGCCAUUUUAGGUGACACAGCCUGG
 35 GAUUUUGGAUCCUUGGGAGGAGUGUUUACAUCAUAGGAAGGCUCUCCACCAAGUCUUUG
 GAGCAUCUAUGGAGCUGCCUUCAGUGGGUUUCAUGGACUAUGAAAUCCUAUAGGAGU
 CAUUAUCACAUGGAUAGGAAUUGAACACGCAGCACCUACUGCUGUGACACUAGUAUUG
 GUGGGAAUUGUGACACUGUAUUUGGGAGUCAUGGUGCAGGGCGAUAGUGGUUGCGUUGUGA
 GCUGGAAAAACAAAGAACUGAAUUGGGAGUGGAGUUCACAGACAACGUGCACAC
 40 AUGGACAGAACAAUACAAUCCACAGAAUCCCUUCAAAACUAGCUUCAGCUAUCCAG
 AAAGCCCAUGAAGAGGACAUUUGUGGAAUCCGCUCAGUAACAAGACUGGAGAACUGAUGU
 GGAAACAAAUAACACCAGAAUUGAACACAUUCAUCAGAAAAGAGGUGAAGUUAACUAU
 UAUGACAGGAGACAUCAAGGAUCAUGCAGGCAGGAAACGAUCUCUGCGGCCUAGCCC
 ACUGAGCUGAAGUAUCAUGGAAAACAUGGGCAAAGCAAAAGCUCUACAGAGUCUC
 45 AUAACCAGACCUUUCUCAUUGAUGGCCCCGAAACAGCAGAACAGCAGAACACACAAUAGAGC
 UUGGAAUUCGUUGGAAGUUGAAGACUAUGGCUUUGGAGUAUUCACCAACAAUUAUGGC
 AAAUUGAAAGAAAACAGGAUGUAUUCUGCGACUAAAACUCAUGUCAGCGGCCAUAAAAG
 ACAACAGAGCCGUCCAUGCCGAUAUUGGUUAUUGGAUAGAAAGUGCACUCAUAGACACAU
 GAAGAUAGAGAAAGCCUCUUCAUUGAAGUAAAACUGCCACUGGCCAAAUCACACACC
 50 CUCUGGAGCAAUGGAGUGCUAGAAAGUGAGAUGUAUUCCAAAGAAUCUCUGCGUGGACCA

UGUCUCAACACAACUAUAGACCAGGCUACCAUACACAAUAACAGGACCAUGGCAUCUAGG
UAAGCUUGAGAUGGACUUUGAUUUCUGUGAUGGAACAACAGUGGUAGUGACUGAGGACUGC
GGAAAAGAGGGACCCUCUUGAGAACAAACCACUGCCUCUGGAAAACUCAUAACAGAAUGGU
GCUGCCGAUCUUGCACAUUACCACCGCUAAGAUACAGAGGUGAGGAUGGGUGCUGGUACGG
5 GAUGGAAAUCAGACCAUUGAAGGAGAAAGAAGAGAAUUGGUCAACUCCUUGGUACACAGCU
GGACAUGGGCAGGUCGACAACUUUCACUAGGAGUCUUGGGAAUGGCAUUGUUCUCCUGGAGG
AAAUGCUUAGGACCCGAGUAGGAACGAAACAUGCAAUACUACUAGUUGCAGUUUCUUUJGU
GACAUUGAUCACAGGGAACAUUGGUUAGAGACCUGGGAGAGUGAUGGUUAUGGUAGGC
GCCACUAUGACGGAUGACAUAGGUUAUGGGCUGACUUACUUGCCCACUAGCAGCCUCA
10 AAGUCAGACCAACUUUUGCAGCUGGACUACUCUUGAGAAAGCUGACCUCCAAGGAAUUGAU
GAUGACUACUAUAGGAUUGUACUCCUCUCCAGAGCACCAUACCAGAGACCAUUCUUGAG
UUGACUGAUGCNUUAGCCUAGGAUGGUCCUAAAUGGUGAGAAUAUGGAAAAGU
AUCAAUUGGCAGUGACUAUCAUGGCUAUCUUGUGCGUCCAAACGCAUGUAAUACAAAA
CGCAUGGAAAGUGAGUUGCACAUAUUGGCAGUGGUCCUACUGUGUGCUACGUGCUCACUGGAC
15 UCCUCACAGCAAAAACAGAUUUGGAUACCAUAGCAUUGACGAUCAAAGGUUCUAAUCCAA
CAGCUAUUUUUUCUAACAACCCUCUCAAGAACCAAGCAGCAAGAAAAGGAGCUGGCCAUAAAUGA
GGCUAUCAUGGCAGUCGGGAUGGUGAGCAUUUAGCCAGUUCUCUCCUAAAUAUGUAAU
CCCAUGACAGGACCAUUAGUGGCUGGAGGGCUCCUACUGUGUGCUACGUGCUCACUGGAC
GAUCGGCCGAUUUGGAACUGGAGAGCAGCCGAUGUAAAUGGGAGACCAAGGCAGAGAU
20 AUCAGGAAGCAGUCCAUCUGUCAAACAAUACAGAAGAAGAUGGUAGCAUGUCGAUAAA
AAUGAAGAGGAAGAACAAACACUGACCAUACUAAUAGAACAGGAUUGCUGGUACUCAG
GACUUUUUCCUGUAUCAAUACCAUACACGGCAGCAGCAUGGUACCUUGGGAGUGAAGAA
ACAACGGGCCGGAGUAUUGGGAGUUCUUCACCCCCACCCAUUGGGAAAGGCUGAACUG
GAAGAUGGAGCCUAUAGAAUUAAGCAAAAAGGGAUUCUUGGAUUUCCAGAUCGGAGCCG
25 GAGUUUACAAAGAAGGAACAUUCCAUACAAUAGUGGCAUGUCACACGUGGCGCUGUUCUAAU
GCAUAAAGGAAGAGGAUUGAACCAACAUGGGCGACGUCAAGAAAGACCUAAUCAUAAU
GGAGGAGGCUGGAAGUAGAAGGAGAAUGGAAGGAAGGAGAAGAAGUCCAGGUAUUGGCAC
UGGAGCCUGGAAAAAAUCCAAGAGCCGUCAAACGAAACCUGGUCCUUCACACGC
CGGAACAAUAGGUGCUGUAUCUCUGGACUUUCUCCUGGAACGUCAGGAUCUCCAAUUAUC
30 GACAAAAAAGGAAAAGUUGUGGGCUUUUAGGUUAUGGUUAGGUUGGUUACAAGGAGUGGAGCAU
AUGUGAGUGCUAUAGCCCAGACUGAAAAAGCAUUGAAGACAACCCAGAGAUCGAAGAUC
CAUUUUCCGAAAGAGAACUGACCAUCAUGGACCUCCACCCAGGAGCAGGGAAAGACGAAG
AGAUACCUUCCGGCAUAGUCAGAGAACGUAAACGGGUUUGAGAACAUUAAUCUUGG
CCCCCACUAGAGUUGUGGCAGCUGAAAUGGAGGAAGCCCUUAGAGGACUUCACAAAGAUA
35 CCAGACCCCAGCCAUCAGAGCUGAGCACACCGGGGGAGAUUGUGGGACCUAAUGGUCAU
GCCACAUUUACCAUGAGGCUGCUAUCACCAGUUAGAGUGCCAAACUACAACCUGAUUAUC
UGGACGAAGCCCAUUUCACAGACCCAGCAAGUUAAGCAGCUAGAGGAAUACUCAACUCG
AGUGGAGAUGGGUGAGGCAGCUGGGAUUUUAUGACAGCCACUCCCCGGAGCAGAGAC
CCAUUUCCUCAGAGCAAUGCACCAUCAAGAAGAACAGGAAUCCUGAACCGCUCGU
40 GGAAUUCGGACAUGAAUGGUACGGAAUAAAAGGGAGACUGUUUGGUUCGUUCCAG
UAUAAAAGCAGGAAAAGUAUAGCAGCUUGCCUGAGGAAAAAUGGAAAGAAGUGAUACAA
CUCAGUAGGAAGACCUUUGAUUCUGAGUAUGUCAAGACUAGAACCAUAGAUUGGGACUUCG
UGGUUACACUGACAUUUCAGAAAUGGGUGCCAAUUUCAAGGCUGAGAGGGUUUAAGACCC
CAGACGCUGCAUGAAACCAGUCAUACUACAGAUGGUGAAGAGCAGGGUGAUUCUGGCAGGA
45 CCUAUGCCAGUGACCCACUCUAGUGCAGCACAAAGAACAGAGGGAGAAUAGGAAGAAUCCAA
AAAUGAGAAUGACCAUACAUACAGGAGAACUACUGGGAGAACACUGUGCAGAGAC
ACACUGGAAAGAACGUAAAAGCUCCUAGAUACAUCAACACGCCAGAACAGGAUCAUUCU
AGCAUGUUCGAACCAGAGCGUGAAAAGGUGGAUGCCAUUGAUGGGCAUACCGCUUGAGAG
GAGAAGCAAGGAAAACCUUUGUAGACUUAAUGAGAACAGGAGACCUACCAGUCUGGUUGGC
50 CUACAGAGUGGCAGCUGAAGGCAUCAACUACGCAAGACAGAACAGAAGGUGGUUUUGAUGGAGUC

AAGAACACCAAAUCCUAGAAGAAAAGCUGGAGUUGAAAUCUGGACAAAGAAGGGGAAA
GGAAGAAAUUGAAACCCAGAUGGUUGGAUGCUAGGAUCUUCUGACCCACUGGCGCUAAA
AGAAUUAAGGAAUUGCAGCCGGAAGAAAGUCUCUGACCCUGAACCACAGAAAUG
GGUAGGCUCCCAACCUUCAUGACUCAGAAGGCAAGAGACGCACUGGACAACUAGCAGUGC
5 UGCACACGGCUGAGGCAGGUGGAAGGGCGUACAACCAUGCUCUCAGUGAACUGCCGGAGAC
CCUGGAGACAUUGCUUUACUGACACUUCUGGUACAGUCACGGGAGGGAUUUUUAUC
UUGAUGAGCGCAAGGGCAUAGGGAGAUGACCCUGGGAAUGUGCUGCAUAUCACGGC
GCAUCCUCCUAUUGGUACGCACAAUACAGCCACACUGGAUAGCAGCUUCAUAUAUCUGGA
GUUUUUCUCAUAGUUUUGCUUUAUCCAGAACUGAAAAACAGAGAACACCCCAAGACAAC
10 CAACUGACCUACGUUGCUAUAGCCAUCUCACAGUGGUGGCCGCAACCAUGGCAAACGAGA
UGGGUUUCCUAGAAAAACGAAGAAAGAACUUCGGAUUGGGAGCAUUGCAACCCAGCAACC
CGAGAGCAACAUCUGGACAUAGAUACGUACGUCCUGCAUCAGCAUGGACGCUGUAUGCC
GCCACAACAUUUGUUACACCAAUGUUGAGACAUAGCAUUGAAAAUCCUCAGUGAAUGUGU
CCCUAACAGCUAUAGCCAACCAAGCCACAGGUAAAUGGGUCUCGGGAAAGGAUGGCCAUU
15 GUCAAAGAUGGACAUCGGAGUUCCCUUCUGCCAUGGAUGCUACUCACAAGUCAACCC
AUAAUCUCACAGCAGCUCUUUUCUUAUUGGUAGCACAUAUGCCAUCAUAGGGCAGGAC
UCCAAGCAAAGCAACCAGAGAACGUCAGAAAAGAGCAGCGGGGGCAUCAUGAAAAACCC
AACUGUCGAUGGAAUACAGUGAUUGACCUAGAACCUUUAUGAUCCAAAGUUUGAA
AAGCAGUUGGGACAAGUAUUGCUCUAGGUCCUCUGCGUGACCUAGUAUUGAUGAUGAGGA
20 CUACAUGGGCUCUGUGAGGCUUUAACCUUAGCUACCCGGGCCAUCUCCACAUUGUGGG
AGGAAAUCAGGGAGGUUUGGAACACUACCAUUGCGGUGUCAUAGGCUACAUUUUAGA
GGGAGUUACUUGGCCGGAGCUGGACUUCUUCUUAUUAUGAAGAACACAACCAAACCAA
GAAGGGGAACUGGCAACAUAGGAGAGACGUUGGAGAGAAAUGGAAAGCCGAUUGAACGC
AUUGGGAAAAGUGAAUCCAGAUCUACAAGAAAAGUGGAUCCAGGAAGUGGAUAGAAC
25 UUAGCAAAGAAGGCAUAAAAGAGGAGAACGGACCAUCACGCUGUGUCGCGAGGCUCAG
CAAAACUGAGAUGGUUCGUUGAGAGAACAUAGGUCACACCAGAAGGGAAAGUAGUGGACCU
CGGUUGUGGCAGAGGAGGCUGGUCAUACUAAUUGUGGAGGACUAAAGAAUGUAAGAGAAC
AAAGGCCUACAAAAGGAGGACCAGGACACGAAGAACCCACUCCCAUGUCAACAUUAGGG
GGAAUCUAGUGCGUCUCAAAGUGGAGUUGACGUUUUCUCAUCCGCCAGAAAAGUGUGA
30 CACAUUAUUGUGUGACAUAGGGAGUCAUCACCAAAUCCACAGUGGAAGCAGGACGAACA
CUCAGAGUCCUUAACUAGUAGAAAUAUUGGUAGAACACAACACUCAAAUUGCAUAAGG
UUCUCAACCCAUUAUAGCCCUCAGUCAUAGAAAAAAUUGGAAGCACUACAAAGGAAAUAUGG
AGGAGCCUUAUGUGAGGAUCCACUCUCAGAAACUCCACACAUGAGAUGUACUGGGUAUCC
AAUGCUUCCGGGAACAUAGUGUCAUCAGUGAACAUAGGUAGUUGAUAACAGAU
35 UUACAAUGAGAUACAAGAAAGCCACUUACGAGCCGGAGUUGACCUCCUGGAAGCGGAACCC
UAACAUUCCGGAUUGAAAGUGAGAUACCAACCUAGAUUAUUAUUGGGAAAAGAAUAGAAAAA
AUAAAAGCAAGAGCAUGAAACAUCAUAGGCACUAAUGAGACCAAGACCACCAUACAAAAGUGGG
CAUACCAUGGUAGCUAUGAAACAAACAGACUGGAUCAGCAUCAUCCCAUGGUCAACGGAGU
GGUCAGGCUGCUGACAAAACCUUUGGGACGUUGUCCCAUGGUGACACAGAUGGCAAUAGACA
40 GACACGACUCCAUUUGGACAACAGCGGUUUUUAAGAGAAAGUGGACACGAGAACCC
AACCGAAAGAAGGCACGAAGAAACUAAUAGAAAUAACAGCAGAGUGGCUUUUGGAAAGAAU
AGGGAAGAAAAGACACCCAGGAUGUGCACCAGAGAAGAAUUCACAAGAAAGGUGAGAAC
AAUGCAGCCUUGGGGGCAUUAUCUGAUGAGAACAGUGGAAGUCGGCACGUGAGGCUG
UUGAAGAUAGUAGGUUUUGGGACUGGUUGACAAGGAAAGGAAUCUCCACUUCUGAAGGAAA
45 GUGUGAAACAUGUGUGUACAACAUGAUGGGAAAAGAGAGAACAGUAGGGAAUUCGGC
AAGGCCAAAGGCAGCAGAGCCAUUAGGUACAUUGGUUGGACUUGGGAGCACGC
AAGGCCUAGGAUUCUAAAUGAAGAUACACUGGUUCUCCAGAGAGAACUCC
GGAGGAGAAGGGCUGCACAAGCUAGGUACAUUCUAAGAGACGUGAGCAAGAAAGAGGG
GGAGCAAUGUAUGCCGAUGACACCGCAGGAUGGGAUACAAAUAACACUAGAAGACCUAA
50 AAAAUGAAGAGAUGGUACAAACACACAUUGGAAGGAGAACACAAGAAACUAGCCGAGGCCAU

5 UUUCAAACUAACGUACCAAAACAAGGUGGUGCGUGUGCAAAGACCAACACCAAGAGGCACA
 GUAAUGGACAUCAUUACGAGAAGAGACCAAGAGGUAGUGGACAAGUUGGCACCUAUGGAC
 UCAAUACUUUCAACCAAAUAGGAAGGCCAACUAAUCAGACAGAUGGAGGGAGAAGGAGUCUU
 UAAAAGCAUUCAGCACCACAAUCACAGAAGAAAUCGCUGUGCAAACUGGUUAGCAAGA
 10 GUGGGCGCGAAAGGUUAUCAAGAAUGGCCAUCAGUGGAGAUGAUUGGUUGUGAAACCUU
 UAGAUGACAGGUUCGCAAGCGCUUUAACAGCUCUAAAUGACAUGGGAAAGAUUAGGAAAGA
 CAUACAACAAUGGGAACCUUCAAGAGGAUGGAAUGAUUGGACACAAGUGGCCUUCGUUCA
 CACCAUUUCCAUGAGUAAAUCAGAAAGACGGUCGCUACUCGUUGGUUCCAUGUAGAAACC
 AAGAUGAACUGAUUGGCAGAGCCGAAUCUCCCAAGGAGCAGGGUGGUCUUUGCGGGAGAC
 15 GCCUGUUUGGGAAAGUCUUACGCCAAAUGUGGAGCUUGAUGUACUUCACAGACGCGAC
 CUCAGGCUGGCGCAAUGCUUUUGCUCGGCAGUACCAUCACAUUGGUUCCAACAAGUC
 GAACAACCUGGUCCAUACAUGCUAAACAUGAAUGGAUGACAACGGAAGACAUGCUGACAGU
 CUGGAACAGGGUGUGGAAUCAAGAAAACCAUGGAUGGAAGACAAAACUCCAGUGGAAACA
 UGGGAGGAAAUCCAUACUUGGGAAAAGAGAACCCAAUGGUGGCUCAUUGAUUGGU
 20 UAACAAGCAGGCCACCUUGGCAAAGAACAUCCAAGCAGCAAUAACAUCAAGUUAGAUCCU
 UAUAGGCAAUGAAGAAUACACAGAUUACAUGCCAUCGAUGAAAAGAUUCAGAAGAGAG
 GAAGAACAGGAGUUCUGUGGUAGAAAGCAAACUAACAGAAACAAGGCUAGAAGUCAGG
 UCGGAUUAAGCCAUCAGUACGGAAAAACUAUGCUACCUGUGAGCCCCGUCCAAGGACGUUA
 AAAGAACAGCAGGCCAUCAAAAUGCCAUAGCUUGAGUAAACUAUGCAGCCUGUAGCUC
 25 CUGAGAAGGUGUAAAAAUCCGGAGGCCACAAACCAUGGAAGCUGUACGCAUGGCGUAGU
 GGACUAGCGGUUAGGGAGACCCCUCCUACAAACUAGCAGCAACAAUUGGGGCCAAGGC
 GAGAUGAACGUAGUCUCGCUUGGAAGGACUAGAGGUUAGAGGAGACCCCCCGAAACAAA
 AAACAGCAUAUUGACGCUGGGAAAGACCAGAGAACCCUGCUGUCUCCUCAGCAUCAU
 GCACAGAACGCCAGAAAUGGAUGGUGCUGUUGAAUCAACAGGUUC
>VDV2 prME nucleotide sequence (SEQ ID NO: 25)

UUCCAUUUACCACACGUAACGGAGAACCAACACAAUGAUCGUCAGCAGACAAGAGAAAGGGA
 AAAGUCUUCUGUUAAAACAGAGGUUGGGGUGAACAUGUGUACCCUCAUGGCCAUGGACCU
 UGGUGAAUUGUGUGAAGACACAAUCACGUACAAGUGUCCCUUCUCAGGCAGAAUGAGCCA
 30 GAAGACAUAGACUGUUGGUGCAACUCUACGUCCACGUGGUACUUAUGGGACGUGUACCA
 CCAUGGGAGAACAUAGAAGAGAAAAAGAACAGUGGCCACUCGUUCCACAUGUGCGAAUGGG
 ACUGGGAGACACGAACUGAAACAUGGAUGUCAUCAGAAGGGCCUGGAAACAUGUCCAGAGA
 AUUGAAACUUGGAUCUUGAGACAUCCAGGCUUCACCAUGAUGGCAGCAAUCCUGGCAUACA
 CCAUAGGAACGACACAUUCCAAAGAGCCUGAUUUUCAUCUUACUGACAGCUGUCACUCC
 35 UUCAAUGACAAUGCGUUGCAUAGGAAUGUAAAAGAGACUUUGUGGAAGGGGUUUCAGGA
 GGAAGCUGGGUUGACAUAGCUUAGAACAGAACUGGAAAGCUGUGACGACGAUGGCAAAAACA
 AACCAACAUUGGAUUUUGAACUGAUAAAACAGAACGCCAACAGCCUGCCACCCUAGGAA
 GUACUGUAUAGAGGCAAAGCUAACCAACACAACAGAAUCUCGCUUGCCAACACAAGGG
 GAACCCAGCCUAAAUGAAGAGCAGGACAAAAGGUUCGUCUGCAAACACUCCAUGGUAGACA
 40 GAGGAUGGGAAAUGGAUGUGGACUAUUUGGAAGGGAGGCAUUGUGACCUUGUGCUGU
 CAGAUGCAAAAGAACAGAACUGGAAGGAAAAGUUGUGCAACCAGAAAACUUGGAACACCAUU
 GUGAUAAACACCUCACUCAGGGGAAGAGCAUGCAGUGGGAAAUGACACAGGAAAACAUGGCA
 AGGAAAUCAAAACACACAGAGUCCAUACACAGAACAGCAGAAUUGACAGGUUAUGGCAC
 UGUACAAUUGGAGUGCUCUCCAAGAACGGGCCUCGACUUCUCAAUGAGAUGGUGUUGCUG
 45 AUGGAAAUAAGCUUUGGCUGGUGCACAGGCAAUGGUUCCUAGACCUGGCCGUUACCAUGGU
 UGCCCGGAGCGGACACACAAGAGUAAAUGGAUACAGAACGGAGACAUUGGUACUUUCAA
 AAAUCCCCAUGCGAAGAACAGGAUGUUGUUGUUUAGGAUCCCAAGAACGGGCCAUGC
 ACAGCACUUACAGGGCCACAGAAAUCAGAACAGAACUACUCUUCACAGGAC
 AUCUCAAGUGCAGGCUGAGAAUGGACAAGCUACAGCUAAAGGAAUGUCAUACUCU
 AUGUG

CACAGGAAAGUUAAAAGUUGUGAAGGAAAUAGCAGAAACACAACAUAGGAACAAUAGUAUC
 AGAGUGCAAUAGAAGGGACGGCUCUCCAUGCAAGAUCCUUUUGAGAUAAUGGAUUGG
 AAAAAAGACAUGUCUAGGUCGCCUGAUACAGUCAACCCAAUUGUGACAGAAAAAGAUAG
 5 CCCAGUCAACAUAGAAGCAGAACCUCCAUUUGGAGACAGCUACAUCAUCAUAGGAGUAGAG
 CCGGGACAACUGAAGCUAACUGGUUUAAGAAAGGAAGUUCUAUCGCCAAUUGUUGAGA
 CAACAAUGAGGGGGCGAAGAGAAUGGCCAUUUUAGGUGACACAGCCUGGGAUUUGGAUC
 CUUGGGAGGAGUGUUUACAUCAUAGGAAGGCUCUCCACCAAGUCUUUGGAGCAUCUAU
 GGAGCUGCCUUCAGUGGGUUUCAUGGACUAUGAAAUCCUCAUAGGAGUCAUUAUCACAU
 10 GGAUAGGAAUGAAUUCACGCAGCACUCACUGUCUGACACUAGUAUUGGUGGGAAUUGU
 GACACUGUAUUUGGGAGUCAUGGUGCAGGCC

>VDV2 E protein sequence (SEQ ID NO: 26)

MRCIGMSNRDFVEGVSGGSWVDIVLEHGSCVTTMAKNKPTLDFELIKTEAKQPATLRKYCI
 EAKLTNTTTESRCPTQGEPSLNEQDKRFLVCKHSMVDRGWNGCGLFGKGGIVTCAMFRCK
 15 KNMEGKVVQOPENLEYTIVITPHSGEEHAVGNDTGKHGKEIKITPQSSITEAELTGYGTVM
 ECSPTGLDFNEMVLLQMenKAWLVRQWFLDLPLPWLGADTQESNWIQKETLVTFKNPH
 AKKQDVVVLGSQEGAMHTALTGATEIQMSSGNLLFTGHLKRLRMDKLQLKGMSYSMCTGK
 FKVVKEIAETQHGTIVIRVQYEGDGSPCKIPFEIMDLEKRHVLGRLITVNPIVTEKDSPVN
 20 IEAEPFGDSYIIIGVEPGQLKLNWFKKGSSIGQMFETTMRGAKRMAILGDTAWDFGSLGG
 VFTSIGKALHQVFGAIYGAAFGSVSWMKILIGVIITWIGMNSRSTSLSVTLVLVGIVTLY
 LGVMVQA

>VDV2 M protein sequence (SEQ ID NO: 27)

SVALVPHVRMGLETRTEWMSSEGAWKHVQRIETWILRHPGFTMMAAILAYTIGTTHFQRA
 25 LIFILLTAVTPSMT

Study design

In an open, randomised, controlled, phase IIa trial, 150 healthy adults aged 18-45 years were enrolled at two centres in Mexico City, which is a dengue non-endemic area.
 30 Main exclusion criteria were: pregnancy or breast-feeding, human immunodeficiency virus, hepatitis B or C seropositivity, immunodeficiency or any other chronic illness that could interfere with the results, previous residence in or travel of >2 weeks to areas with high dengue endemicity, a history of flavivirus infection or previous vaccination against flavivirus disease. Women who were capable of conceiving were required to use an effective method of contraception or abstinence for at least 4 weeks before the first
 35 injection until at least four weeks after the last injection.

Participants were randomised into two groups and vaccinations were performed on Day 0 and Day 105 (\pm 15 days). The groups received the following formulations:

40 **Group 1:** Blended CYD/VDV2 tetravalent formulation, i.e. a formulation comprising CYD-1, CYD-3, CYD4 and VDV2.

Group 2: Control tetravalent formulation (CYD-TDV), i.e. CYD-1, CYD-2, CYD-3 and CYD-4.

5 The formulations contained 10^5 CCID₅₀ of each serotype of the CYD viruses and the formulation administered to Group 1 contained 10^4 CCID₅₀ of the VDV-2 virus.

Viremia

10 To evaluate the safety of the vaccines, the presence of CYD-1–4 or VDV-2 was assessed in serum collected 7, 14 and 21 days after each injection. Analyses were performed by the Global Clinical Immunology laboratory (Sanofi Pasteur, Swiftwater, PA, USA).

15 Analyses for CYD-1–4 viremia were performed in two steps, as previously described in Poo *et al.*, *Pediatr Infect Dis J* (2011) 30: e9. Briefly, a first, non-serotype-specific, reverse transcriptase-polymerase chain reaction (RT-PCR) was used to detect the presence of any of the four CYD viruses. Samples that were positive in this first test were then analysed using four CYD serotype-specific quantitative RT-PCRs. In the non-serotype-specific RT-PCR, RNA was extracted from the serum using a commercial kit and an RT-PCR was carried out with primers from the yellow fever core gene sequence. In the serotype-specific RT-PCRs, RNA was again extracted from the serum using a commercial kit and an RT-PCR was carried out with serotype-specific primers from the envelope non-structural protein 1 junction gene sequence for each serotype. A dengue RT-PCR for serotype 2 was performed in group 1 since the tetravalent blending formulation administered to this group contained the VDV-2 virus.

20 25 Immunogenicity

30 Antibody levels to each of the four dengue virus serotypes were determined by 50% plaque reduction neutralisation test on serum collected 28 days after each injection as well as on day 365 after the first injection. Briefly, serial 2-fold dilutions of heat-inactivated serum were mixed with a constant challenge dose of each dengue serotype DEN-1, -2, -3, or -4 (expressed as plaque forming unit [PFU]/mL). The mixtures were inoculated into wells of a 24-well plate of confluent VERO cell monolayers. After incubation for several days, dengue virus infection is indicated by formation of plaques. The neutralising antibody titre is calculated as the highest reciprocal dilution (1/dil) of serum at which $\geq 50\%$ reduction in viral plaque count is observed (PRNT50). The lower 35 limit of quantitation of the dengue PRNT50 is 10; samples with titres ≥ 10 were considered seropositive.

Results

Formulations were administered to participants in Groups 1 and 2 on day 0 and day 105 of the study. There were no marked differences between the two groups with regard to the injection site or systemic reactogenicity after either the first or the second vaccination. Viremia was assessed in serum collected 7, 14 and 21 days after each injection (Table 7). The neutralising antibody titres were measured 28 days after each injection and on day 365 after the first injection (Table 8).

10 **Table 7. Vaccine virus viremia 7, 14, or 21 days after first and second injections (n (%) with detectable and quantifiable viremia)**

	First injection		Second injection	
	Group 1 Blended CYD/VDV	Group 2 Tetraivalent CYD-TDV	Group 1 Blended CYD/VDV	Group 2 Tetraivalent CYD-TDV
Non-serotype specific				
N	29	31	28	29
Detectable viraemia	27 (93%)	25 (81%)	1 (4%)	1 (3%)
Quantifiable viraemia	1 (3%)	2 (6%)	0	0
DENV-1				
Detectable viraemia	1 (3%)	4 (13%)	0	0
Quantifiable viraemia	0	2 (7%)	0	0
DENV-2				
Detectable viraemia	0	2 (6%)	0	0
Quantifiable viraemia	0	0	0	0
DENV-3				
Detectable viraemia	8 (28%)	7 (23%)	1 (4%)	0
Quantifiable viraemia	0	0	0	0
DENV-4				
Detectable viraemia	24 (83%)	21 (68%)	0	0
Quantifiable viraemia	0	3 (1%)	0	0

After the first injection, detectable viremia, as determined by the non-serotype specific RT-PCR test, was observed in a similar proportion of participants in both groups (see Table 7). In the majority of cases, viremia was below the lower limit of quantitation. Analysis with the serotype-specific assays showed that CYD-4 was the most commonly detected serotype, followed by CYD-3. After the second injection of the blended CYD/VDV

vaccine in Group 1 or the CYD-TDV vaccine in Group 2, viremia was only detected in one participant per group by the non-serotype-specific assay.

Accordingly, there was no significant difference between the levels of viremia induced by the blended CYD/VDV and CYD-TDV.

5

Table 8. Geometric mean titres (95% confidence interval) of dengue antibodies 28 days after the first and second injections and 365 days after the first injection

	Group 1 CYD/VDV blended	Group 2 CYD-TDV
First injection		
Serotype 1	15 (9;28)	17 (10;31)
Serotype 2	17 (8;33)	32 (16;65)
Serotype 3	64 (31;133)	23 (13;39)
Serotype 4	552 (299;1019)	468 (226;968)
Second injection		
Serotype 1	54 (30;96)	28 (15;50)
Serotype 2	152 (79;293)	43 (23;79)
Serotype 3	127 (71;229)	46 (29;73)
Serotype 4	246 (159;382)	173 (97;307)
365 days post-dose 1		
Serotype 1	14 (9;22)	18 (10;30)
Serotype 2	55 (32;94)	16 (9;29)
Serotype 3	36 (20;64)	11 (7;16)
Serotype 4	103 (69;155)	72 (44;117)

It can be seen from Table 8 that the second injection of the blended CYD/VDV vaccine (Group 1) induced higher GMTs against serotype 2 of dengue virus than the CYD-TDV vaccine (Group 2). An improved response to serotype 2 in the blended CYD/VDV group was also observed 365 days after the first dose.

Furthermore, the second injection of the blended CYD/VDV vaccine (Group 1) resulted in an improved neutralising antibody response against all serotypes of dengue virus when compared with the group receiving the CYD-TDV vaccine (Group 2). Importantly, the blended CYD/VDV formulation group demonstrated a more persistent neutralising antibody response against dengue virus than the CYD-TDV group on day 365 after the first injection.

The example therefore shows that, overall, the blended CYD-1, 3, 4/VDV2 vaccine formulation induces stronger and longer lasting immune responses against the dengue virus serotypes than the CYD-TDV vaccine while showing a similar safety profile, as determined by the levels of viremia.

Sequence Listing

SEQ ID NO.	Sequence
1	prM+E CYD23 circulating strain nucleotide sequence
2	prM+E CYD23 circulating strain protein sequence
3	prM+E consensus serotype 2 protein sequence
4	prM+E LAV2 nucleotide sequence
5	prM+E BID/V585 nucleotide sequence
6	prM+E PR/DB023 nucleotide sequence
7	prM+E MD1280 nucleotide sequence
8	prM+E LAV2 protein sequence
9	prM+E BID/V585 protein sequence
10	prM+E PR/DB023 protein sequence
11	prM+E MD1280 protein sequence
12	E consensus serotype 2 protein sequence
13	E LAV2 protein sequence
14	E BID/V585 protein sequence
15	E PR/DB023 protein sequence
16	E MD1280 protein sequence
17	M consensus serotype 2 protein sequence
18	E CYD23 circulating strain protein sequence
19	M LAV2 protein sequence
20	M BID/V585 protein sequence
21	M PR/DB023 protein sequence
22	M MD1280 protein sequence
23	M CYD23 circulating strain protein sequence
24	Entire nucleotide sequence of VDV2 (RNA equivalent)
25	prM+E VDV2 nucleotide sequence (RNA equivalent)
26	E VDV2 protein sequence
27	M VDV2 protein sequence

5 In the listed nucleotide sequences, where a nucleotide sequence is DNA, the nucleotide T may be replaced with the nucleotide U to give the RNA equivalent of that DNA sequence. Similarly, where a nucleotide sequence is RNA, the nucleotide U may be replaced by the nucleotide T to give the equivalent DNA sequence. The DNA sequences listed above constitute the cDNA sequences of the noted dengue viruses and therefore the equivalent RNA sequences constitute the positive strand RNA of those dengue viruses.

10

CLAIMS

1. A dengue virus serotype 2 vaccine composition comprising:

5 (i) a dengue antigen selected from the group consisting of:
(a) a live attenuated dengue virus;
(b) an inactivated dengue virus;
(c) a live attenuated or inactivated chimeric dengue virus;
(d) a dengue virus-like particle (VLP); and
10 (e) a combination of two or more of (a) to (d);

or

15 (iii) a nucleic acid construct or viral vector which is able to express in a human cell a dengue antigen which is a dengue VLP;

wherein said dengue antigen comprises a polypeptide having at least 90% identity to SEQ ID NO: 12.

20 2. A composition as claimed in claim 1, wherein said polypeptide comprises a valine residue at the position within the polypeptide that corresponds to position 251 of SEQ ID NO: 12.

25 3. A composition as claimed in claim 1 or claim 2, wherein said polypeptide comprises a methionine residue at the position within the polypeptide that corresponds to position 6 of SEQ ID NO: 12.

30 4. A composition as claimed in any preceding claim, wherein said polypeptide comprises a valine residue at the position within the polypeptide that corresponds to position 129 of SEQ ID NO: 12.

35 5. A composition as claimed in any preceding claim, wherein said polypeptide comprises an isoleucine residue at the position within the polypeptide that corresponds to position 141 of SEQ ID NO: 12.

6. A composition as claimed in any preceding claim, wherein said polypeptide comprises an isoleucine residue at the position within the polypeptide that corresponds to position 164 of SEQ ID NO: 12.
- 5 7. A composition as claimed in any preceding claim, wherein said polypeptide comprises an aspartate residue at the position within the polypeptide that corresponds to position 203 of SEQ ID NO: 12.
- 10 8. A composition as claimed in any preceding claim, wherein said polypeptide comprises a threonine residue at the position within the polypeptide that corresponds to position 226 of SEQ ID NO: 12.
- 15 9. A composition as claimed in any preceding claim, wherein said polypeptide comprises a glycine residue at the position within the polypeptide that corresponds to position 228 of SEQ ID NO: 12.
10. A composition as claimed in any preceding claim, wherein said polypeptide comprises an isoleucine residue at the position within the polypeptide that corresponds to position 308 of SEQ ID NO: 12.
- 20 11. A composition as claimed in any preceding claim, wherein said polypeptide comprises a threonine residue at the position within the polypeptide that corresponds to position 478 of SEQ ID NO: 12.
- 25 12. A composition as claimed in any preceding claim, wherein said polypeptide comprises an isoleucine residue at the position within the polypeptide that corresponds to position 484 of SEQ ID NO: 12.
13. A composition as claimed in any preceding claim, wherein said polypeptide comprises an isoleucine residue at the position within the polypeptide that corresponds to position 485 of SEQ ID NO: 12.
- 30 14. A composition as claimed in any preceding claim, wherein said polypeptide comprises an alanine residue at the position within the polypeptide that corresponds to position 491 of SEQ ID NO: 12.

15. A composition as claimed in any preceding claim, wherein said dengue antigen comprises a polypeptide having at least 90% identity to SEQ ID NO: 3.

16. A composition as claimed in any one of claims 1 to 14, wherein said polypeptide
5 has at least 90% identity to SEQ ID NO: 3.

17. A composition as claimed in claim 15 or 16, wherein said polypeptide comprises a glycine residue at the position within the polypeptide that corresponds to position 15 of SEQ ID NO: 3.

10

18. A composition as claimed in any one of claims 15 to 17, wherein said polypeptide comprises a leucine residue at the position within the polypeptide that corresponds to position 24 of SEQ ID NO: 3.

15

19. A composition as claimed in any one of claims 15 to 18, wherein said polypeptide comprises an isoleucine residue at the position within the polypeptide that corresponds to position 39 of SEQ ID NO: 3.

20

20. A composition as claimed in any one of claims 15 to 19, wherein said polypeptide comprises a valine residue at the position within the polypeptide that corresponds to position 120 of SEQ ID NO: 3.

25

21. A composition as claimed in any one of claims 15 to 20, wherein said polypeptide comprises a threonine residue at the position within the polypeptide that corresponds to position 125 of SEQ ID NO: 3.

30

22. A composition as claimed in any one of claims 1 to 21, wherein said polypeptide comprises: (i) the sequence as set forth in SEQ ID NO: 13 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 13; (ii) the sequence as set forth in SEQ ID NO: 14 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 14; (iii) the sequence as set forth in SEQ ID NO: 15 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 15; (iv) the sequence as set forth in SEQ ID NO: 16 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 16; (v) the sequence as set forth in SEQ ID NO: 18 or a sequence

35

having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 18 or (vi) the sequence as set forth in SEQ ID NO: 26 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 26.

5

23. A composition as claimed in claim 1, wherein said dengue antigen comprises a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO: 13; SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 18 and SEQ ID NO: 26.

10

24. A composition as claimed in claim 23, wherein said dengue antigen comprises a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO: 13 and SEQ ID NO: 16.

15

25. A composition as claimed in claim 22, wherein said dengue antigen further comprises a polypeptide comprising: (i) the sequence as set forth in SEQ ID NO: 19 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 19; (ii) the sequence as set forth in SEQ ID NO: 20 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 20; (iii) the sequence as set forth in SEQ ID NO: 21 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 21; (iv) the sequence as set forth in SEQ ID NO: 22 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 22; (v) the sequence as set forth in SEQ ID NO: 23 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 23; (vi) the sequence as set forth in SEQ ID NO: 27 or a sequence having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 27.

20

25

30

26. A composition as claimed in claim 23, wherein said dengue antigen further comprises a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO: 19; SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23 and SEQ ID NO: 27.

35

27. A composition as claimed in claim 22, wherein said dengue antigen comprises:

- i) a polypeptide having the sequence as set forth in SEQ ID NO: 13 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 13; and a polypeptide having the sequence as set forth in SEQ ID NO: 19 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 19;
- ii) a polypeptide having the sequence as set forth in SEQ ID NO: 14 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 14; and a polypeptide having the sequence as set forth in SEQ ID NO: 20 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 20;
- iii) a polypeptide having the sequence as set forth in SEQ ID NO: 15 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 15; and a polypeptide having the sequence as set forth in SEQ ID NO: 21 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 21;
- iv) a polypeptide having the sequence as set forth in SEQ ID NO: 16 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 16; and a polypeptide having the sequence as set forth in SEQ ID NO: 22 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 22;
- v) a polypeptide having the sequence as set forth in SEQ ID NO: 18 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 18; and a polypeptide having the sequence as set forth in SEQ ID NO: 23 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 23; or
- vi) a polypeptide having the sequence as set forth in SEQ ID NO: 26 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 26; and

a polypeptide having the sequence as set forth in SEQ ID NO: 27 or a polypeptide having at least 1 and no more than 5 amino acid substitutions with respect to the sequence as set forth in SEQ ID NO: 27.

5 28. A composition as claimed in claim 26, wherein said dengue antigen comprises: i) a polypeptide of SEQ ID NO: 13 and a polypeptide of SEQ ID NO: 19; ii) a polypeptide of SEQ ID NO: 14 and a polypeptide of SEQ ID NO: 20; iii) a polypeptide of SEQ ID NO: 15 and a polypeptide of SEQ ID NO: 21; iv) a polypeptide of SEQ ID NO: 16 and a polypeptide of SEQ ID NO: 22; v) a polypeptide of SEQ ID NO: 18 and a polypeptide of SEQ ID NO: 23 or vi) a polypeptide of SEQ ID NO: 26 and a polypeptide of SEQ ID NO: 27.

10 29. A composition as claimed in claim 15 or claim 16, wherein said dengue antigen comprises a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO: 8; SEQ ID NO: 9, SEQ ID NO: 10 and SEQ ID NO: 11.

15 30. A composition as claimed in claim 29, wherein said dengue antigen comprises a polypeptide comprising a sequence selected from the group consisting of SEQ ID NO: 8; and SEQ ID NO: 11.

20 31. A composition as claimed in any preceding claim, wherein said dengue antigen is selected from the group consisting of: (a) a live attenuated dengue virus; (b) an inactivated dengue virus; (c) a live attenuated or inactivated chimeric dengue virus; or (d) a combination of two or more of (a) to (c); wherein said dengue antigen comprises a nucleotide sequence encoding a protein comprising a polypeptide as defined in any one of claims 1 to 30.

25 32. A vaccine composition comprising a dengue antigen of serotype 2 selected from the group consisting of: (a) a live attenuated dengue virus; (b) an inactivated dengue virus; (c) a live attenuated or inactivated chimeric dengue virus; or (d) a combination of two or more of (a) to (c); wherein said dengue antigen comprises a nucleotide sequence encoding a protein comprising the polypeptide or polypeptides as defined in any one of claims 1 to 30.

30 33. A vaccine composition comprising a dengue antigen of serotype 2 selected from the group consisting of: (a) a live attenuated dengue virus; (b) an inactivated dengue virus; (c) a live attenuated or inactivated chimeric dengue virus; or (d) a

combination of two or more of (a) to (c); wherein said dengue antigen comprises a nucleotide sequence having at least 90% sequence identity to a sequence selected from the group consisting of the RNA equivalent of SEQ ID NO: 1, the RNA equivalent of SEQ ID NO: 4, the RNA equivalent of SEQ ID NO: 5, the RNA equivalent of SEQ ID NO: 6, the RNA equivalent of SEQ ID NO: 7 and SEQ ID NO: 25.

34. A composition as claimed in any preceding claim, wherein said composition comprises a live attenuated chimeric dengue virus.

10

35. A composition as claimed in claim 34, wherein said composition comprises one or more proteins from a dengue virus and one or more proteins from a different flavivirus.

15

36. A composition as claimed in claim 35, wherein the different flavivirus is a yellow fever virus.

37. A composition as claimed in claim 36, wherein the yellow fever virus is YF-Vax.

20

38. A composition as claimed in any preceding claim, wherein the composition further comprises a dengue antigen of serotype 1, a dengue antigen of serotype 3 and a dengue antigen of serotype 4.

25

39. A composition as claimed in claim 38, wherein said dengue antigens of serotypes 1, 3 and 4 are each independently selected from the group consisting of a live attenuated dengue virus and a live attenuated chimeric dengue virus.

30

40. A composition as claimed in claim 39, wherein said dengue antigens of serotypes 1, 3 and 4 are each a live attenuated chimeric dengue virus in which the genetic backbone of a recipient flavivirus has been modified by exchanging the sequences encoding the prM and E proteins of the recipient flavivirus with the corresponding sequences of a dengue virus.

35

41. A composition as claimed in claim 40, wherein the recipient flavivirus is a yellow fever virus.

42. A composition as claimed in any one of claims 38 to 41, wherein said dengue antigens of serotypes 1, 3 and 4 are each a live attenuated chimeric dengue virus and the dengue antigen of serotype 2 is a live attenuated dengue virus of serotype 2 which comprises a nucleic acid sequence having at least 90% sequence identity to SEQ ID NO: 24.

5

43. A pharmaceutical formulation comprising a composition as claimed in any preceding claim and a pharmaceutically acceptable carrier, diluent or excipient.

10 44. A composition as claimed in any one of claims 1 to 42 for use in therapy.

45. A composition as claimed in any one of claims 1 to 42 for use in a method of protecting a human subject against dengue disease caused by a dengue virus of serotype 2.

15 46. A composition as claimed in any one of claims 38 to 42 for use in a method of protecting a human subject against dengue disease caused by a dengue virus of serotype 1, serotype 2, serotype 3 or serotype 4.

20 47. A method of protecting a human subject against dengue disease caused by a dengue virus of serotype 2, wherein said method comprises administering to said subject an effective amount of a composition according to any one of claims 1 to 42.

25 48. A method of protecting a human subject against dengue disease caused by a dengue virus of serotype 1, serotype 2, serotype 3 or serotype 4, wherein said method comprises administering to said subject an effective amount of a composition according to any one of claims 38 to 42.



30 49. A kit comprising a composition according to any one of claims 1 to 42 and instructions for use of said composition in a method of protecting a human subject against dengue disease caused by a dengue virus of serotype 2.

35 50. A kit comprising a composition according to any one of claims 38 to 42 and instructions for use of said composition in a method of protecting a human subject against dengue disease caused by a dengue virus of serotype 1, serotype 2, serotype 3 or serotype 4.


Figure

Figure 1: Construction of the YF-VAX cDNA by RT-PCT and cloning

1/1



Stepwise ligation of each fragment to pJSY2284.1 (pACYC YF-vax 5-3) to obtain a full-length cDNA clone of YF-vax



SEQUENCE LISTING

<110> Sanofi Pasteur
<120> Vaccine Compositions
<130> P39479WO
<150> EP12305911.5
<151> 2012-07-25
<150> EP12305908.1
<151> 2012-07-24
<160> 27
<170> PatentIn version 3.5
<210> 1
<211> 1983
<212> DNA
<213> Artificial Sequence
<220>
<223> clinical trial circulating strain
<400> 1
ttccatctaa ccacacgcaa cggagaacca cacatgatcg tcggtataca ggagaaaggg 60
aaaagtcttc tggtaaaaac agaggatggt gtgaacatgt gcaccctcat ggctatggac 120
cttggtaat tggtaagaacaatcag tacaagtgtc ctcttc tagtgcagaatgag 180
ccagaagaca tagactgttgcgtcaactcc acgtccacgt gggtaaccta tgggacctgt 240
accactacgg gagaacatag gagagaaaaa agatcagtgg cactcggtcc acatgtggaa 300
atgggactgg agacgcgaac cggaaacatgg atgtcatcag aaggggcttgcgtca 360
cagagaatttggat cctgagacat ccaggcttca ccataatggc agcaatcctg 420
gcatacacca taggaacgcac acatttccag agagtccttgcgtca 480
gtcgctcctt caatgacaat gcggtgcata ggaatatcaa atagagactt tggtaaggg 540
gtttcaggag gaagttgggt tgacatagtc ttggaaacatg gaagctgtgt gacgacgt 600
gcaaaaaaaaaca aaccaacatt ggatttcgaa ctgataaaaaa cggaaagccaa acagcctgcc 660
accctaagga agtactgcata agaagaaaaa ctaaccaaca caacaacaga atcccggtc 720
ccaaacacaag gggAACCCAG CCTAAAAGAA gagcaggaca agaggttcgt ctgcaaacac 780
tccatggtag acagaggatg gggaaatgga tgtggattat ttggaaaggg aggcatgt 840
acctgtgcta tggcacatg caaaaagaac atggaaaggaa aaatcgtcga accagaaaaac 900
ttggaaataca ccattgtgggt aacacccatc acggggaaag agcatgcggt cggaaatgac 960
acaggaaaaac acggcaagga aatcaaagta acaccacaga gttccatcac agaagcagaa 1020
ctgacaggat atggcaccgt cacgtggag tgctccccga gaacaggcct cgacttcaat 1080
gagatgggtgt tgctgcagat gggaaaataaa gcttggctgg tgcataggca atggttctca 1140
gacctgccat taccatggct gcccggagcg gataaacaag aatcaaattt gatacagaaaa 1200
gaaacattgg tcactttcaa aaatccccat gcgaaagaaac aggtgttgcgt tggttttagga 1260

eolf-seq1.txt

tcccaagaag	ggccatgca	tacagca	tc acaggagcc	cagaatcca	aatgtcgta	1320
ggaaacttgc	tcttca	tgg acatctca	ag tg	caggctga	aatggacaa	1380
aaaggaatgt	cata	cttat gt	gcacagga	aagttaaag	ttgtgaagga	1440
acacaacatg	gaacgat	agt tatt	caatatga	gggacggct	tccatgt	1500
attcctttg	agataatgg	tttggaaaaa	agatatgtct	taggcgcct	gatcacagtc	1560
aacccaattg	taacagaaaa	agacagccca	gtcaacatag	aagcagaacc	tccattcgga	1620
gacagttaca	tcatcatagg	agtagagccg	ggacaactga	agctcaactg	gttcaagaaa	1680
ggaagttcta	tcggccaaat	gttgagaca	acgatgagag	ggcgaagag	aatggccatt	1740
ttgggtgaca	cagcctgg	gatcc	ctgggaggag	tgttacatc	tataggaaaa	1800
gctctccacc	aagtcttgg	agcgatctat	ggggctgcct	tcagtgggt	ttcatggacc	1860
atgaaaatcc	tcataggagt	cattatcaca	tggataggaa	tgaactcacg	cagcacctca	1920
ctgtctgtgt	cactggta	ct	gtggaaatt	gtgacactgt	attaggagt	catggcag
gcc						1980
						1983

<210> 2
<211> 661
<212> PRT
<213> Artificial Sequence

<220>
<223> clinical trial circulating strain

<400> 2

Phe His Leu Thr Thr Arg Asn Gly Glu Pro His Met Ile Val Gly Ile
1 5 10 15

Gln Glu Lys Gly Lys Ser Leu Leu Phe Lys Thr Glu Asp Gly Val Asn
20 25 30

Met Cys Thr Leu Met Ala Met Asp Leu Gly Glu Leu Cys Glu Asp Thr
35 40 45

Ile Thr Tyr Lys Cys Pro Leu Leu Arg Gln Asn Glu Pro Glu Asp Ile
50 55 60

Asp Cys Trp Cys Asn Ser Thr Ser Thr Trp Val Thr Tyr Gly Thr Cys
65 70 75 80

Thr Thr Thr Gly Glu His Arg Arg Glu Lys Arg Ser Val Ala Leu Val
85 90 95

Pro His Val Gly Met Gly Leu Glu Thr Arg Thr Glu Thr Trp Met Ser
100 105 110

Ser Glu Gly Ala Trp Lys His Ala Gln Arg Ile Glu Thr Trp Ile Leu
115 120 125

eof-seq1.txt

Arg His Pro Gly Phe Thr Ile Met Ala Ala Ile Leu Ala Tyr Thr Ile
130 135 140

Gly Thr Thr His Phe Gln Arg Val Leu Ile Phe Ile Leu Leu Thr Ala
145 150 155 160

Val Ala Pro Ser Met Thr Met Arg Cys Ile Gly Ile Ser Asn Arg Asp
165 170 175

Phe Val Glu Gly Val Ser Gly Ser Trp Val Asp Ile Val Leu Glu
180 185 190

His Gly Ser Cys Val Thr Thr Met Ala Lys Asn Lys Pro Thr Leu Asp
195 200 205

Phe Glu Leu Ile Lys Thr Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys
210 215 220

Tyr Cys Ile Glu Ala Lys Leu Thr Asn Thr Thr Glu Ser Arg Cys
225 230 235 240

Pro Thr Gln Gly Glu Pro Ser Leu Lys Glu Glu Gln Asp Lys Arg Phe
245 250 255

Val Cys Lys His Ser Met Val Asp Arg Gly Trp Gly Asn Gly Cys Gly
260 265 270

Leu Phe Gly Lys Gly Ile Val Thr Cys Ala Met Phe Thr Cys Lys
275 280 285

Lys Asn Met Glu Gly Lys Ile Val Gln Pro Glu Asn Leu Glu Tyr Thr
290 295 300

Ile Val Val Thr Pro His Ser Gly Glu Glu His Ala Val Gly Asn Asp
305 310 315 320

Thr Gly Lys His Gly Lys Glu Ile Lys Val Thr Pro Gln Ser Ser Ile
325 330 335

Thr Glu Ala Glu Leu Thr Gly Tyr Gly Thr Val Thr Met Glu Cys Ser
340 345 350

Pro Arg Thr Gly Leu Asp Phe Asn Glu Met Val Leu Leu Gln Met Glu
355 360 365

Asn Lys Ala Trp Leu Val His Arg Gln Trp Phe Leu Asp Leu Pro Leu
370 375 380

Pro Trp Leu Pro Gly Ala Asp Lys Gln Glu Ser Asn Trp Ile Gln Lys
385 390 395 400

eolf-seq1.txt

Glu Thr Leu Val Thr Phe Lys Asn Pro His Ala Lys Lys Gln Asp Val
405 410 415

Val Val Leu Gly Ser Gln Glu Gly Ala Met His Thr Ala Leu Thr Gly
420 425 430

Ala Thr Glu Ile Gln Met Ser Ser Gly Asn Leu Leu Phe Thr Gly His
435 440 445

Leu Lys Cys Arg Leu Arg Met Asp Lys Leu Gln Leu Lys Gly Met Ser
450 455 460

Tyr Ser Met Cys Thr Gly Lys Phe Lys Val Val Lys Glu Ile Ala Glu
465 470 475 480

Thr Gln His Gly Thr Ile Val Ile Arg Val Gln Tyr Glu Gly Asp Gly
485 490 495

Ser Pro Cys Lys Ile Pro Phe Glu Ile Met Asp Leu Glu Lys Arg Tyr
500 505 510

Val Leu Gly Arg Leu Ile Thr Val Asn Pro Ile Val Thr Glu Lys Asp
515 520 525

Ser Pro Val Asn Ile Glu Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile
530 535 540

Ile Ile Gly Val Glu Pro Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys
545 550 555 560

Gly Ser Ser Ile Gly Gln Met Phe Glu Thr Thr Met Arg Gly Ala Lys
565 570 575

Arg Met Ala Ile Leu Gly Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly
580 585 590

Gly Val Phe Thr Ser Ile Gly Lys Ala Leu His Gln Val Phe Gly Ala
595 600 605

Ile Tyr Gly Ala Ala Phe Ser Gly Val Ser Trp Thr Met Lys Ile Leu
610 615 620

Ile Gly Val Ile Ile Thr Trp Ile Gly Met Asn Ser Arg Ser Thr Ser
625 630 635 640

Leu Ser Val Ser Leu Val Leu Val Gly Ile Val Thr Leu Tyr Leu Gly
645 650 655

Val Met Val Gln Ala
660

eolf-seq1.txt

<210> 3
<211> 661
<212> PRT
<213> Artificial Sequence

<220>
<223> prM+E universal serotype 2

<220>
<221> Variant
<222> (295)..(295)
<223> Xaa can be Val or Ile

<220>
<221> Variant
<222> (474)..(474)
<223> Xaa can be Val or Ile

<400> 3

Phe His Leu Thr Thr Arg Asn Gly Glu Pro His Met Ile Val Gly Arg
1 5 10 15

Gln Glu Lys Gly Lys Ser Leu Leu Phe Lys Thr Glu Asp Gly Val Asn
20 25 30

Met Cys Thr Leu Met Ala Ile Asp Leu Gly Glu Leu Cys Glu Asp Thr
35 40 45

Ile Thr Tyr Lys Cys Pro Leu Leu Arg Gln Asn Glu Pro Glu Asp Ile
50 55 60

Asp Cys Trp Cys Asn Ser Thr Ser Thr Trp Val Thr Tyr Gly Thr Cys
65 70 75 80

Thr Thr Thr Gly Glu His Arg Arg Glu Lys Arg Ser Val Ala Leu Val
85 90 95

Pro His Val Gly Met Gly Leu Glu Thr Arg Thr Glu Thr Trp Met Ser
100 105 110

Ser Glu Gly Ala Trp Lys His Val Gln Arg Ile Glu Thr Trp Ile Leu
115 120 125

Arg His Pro Gly Phe Thr Ile Met Ala Ala Ile Leu Ala Tyr Thr Ile
130 135 140

Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe Ile Leu Leu Thr Ala
145 150 155 160

Val Ala Pro Ser Met Thr Met Arg Cys Ile Gly Ile Ser Asn Arg Asp
165 170 175

Phe Val Glu Gly Val Ser Gly Gly Ser Trp Val Asp Ile Val Leu Glu
Page 5

180 185 190
eolf-seq1.txt

His Gly Ser Cys Val Thr Thr Met Ala Lys Asn Lys Pro Thr Leu Asp
195 200 205

Phe Glu Leu Ile Lys Thr Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys
210 215 220

Tyr Cys Ile Glu Ala Lys Leu Thr Asn Thr Thr Glu Ser Arg Cys
225 230 235 240

Pro Thr Gln Gly Glu Pro Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe
245 250 255

Val Cys Lys His Ser Met Val Asp Arg Gly Trp Gly Asn Gly Cys Gly
260 265 270

Leu Phe Gly Lys Gly Ile Val Thr Cys Ala Met Phe Thr Cys Lys
275 280 285

Lys Asn Met Glu Gly Lys Xaa Val Gln Pro Glu Asn Leu Glu Tyr Thr
290 295 300

Ile Val Ile Thr Pro His Ser Gly Glu Glu His Ala Val Gly Asn Asp
305 310 315 320

Thr Gly Lys His Gly Lys Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile
325 330 335

Thr Glu Ala Glu Leu Thr Gly Tyr Gly Thr Val Thr Met Glu Cys Ser
340 345 350

Pro Arg Thr Gly Leu Asp Phe Asn Glu Met Val Leu Leu Gln Met Glu
355 360 365

Asp Lys Ala Trp Leu Val His Arg Gln Trp Phe Leu Asp Leu Pro Leu
370 375 380

Pro Trp Leu Pro Gly Ala Asp Thr Gln Gly Ser Asn Trp Ile Gln Lys
385 390 395 400

Glu Thr Leu Val Thr Phe Lys Asn Pro His Ala Lys Lys Gln Asp Val
405 410 415

Val Val Leu Gly Ser Gln Glu Gly Ala Met His Thr Ala Leu Thr Gly
420 425 430

Ala Thr Glu Ile Gln Met Ser Ser Gly Asn Leu Leu Phe Thr Gly His
435 440 445

Leu Lys Cys Arg Leu Arg Met Asp Lys Leu Gln Leu Lys Gly Met Ser
Page 6

eolf-seq1.txt

450	455	460
Tyr Ser Met Cys Thr Gly Lys Phe Lys Xaa Val Lys Glu Ile Ala Glu		
465	470	475 480
Thr Gln His Gly Thr Ile Val Ile Arg Val Gln Tyr Glu Gly Asp Gly		
485	490	495
Ser Pro Cys Lys Ile Pro Phe Glu Ile Met Asp Leu Glu Lys Arg His		
500	505	510
Val Leu Gly Arg Leu Ile Thr Val Asn Pro Ile Val Thr Glu Lys Asp		
515	520	525
Ser Pro Val Asn Ile Glu Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile		
530	535	540
Ile Ile Gly Val Glu Pro Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys		
545	550	555 560
Gly Ser Ser Ile Gly Gln Met Phe Glu Thr Thr Met Arg Gly Ala Lys		
565	570	575
Arg Met Ala Ile Leu Gly Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly		
580	585	590
Gly Val Phe Thr Ser Ile Gly Lys Ala Leu His Gln Val Phe Gly Ala		
595	600	605
Ile Tyr Gly Ala Ala Phe Ser Gly Val Ser Trp Thr Met Lys Ile Leu		
610	615	620
Ile Gly Val Ile Ile Thr Trp Ile Gly Met Asn Ser Arg Ser Thr Ser		
625	630	635 640
Leu Ser Val Ser Leu Val Leu Val Gly Val Val Thr Leu Tyr Leu Gly		
645	650	655
Val Met Val Gln Ala		
660		
<210> 4		
<211> 1983		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> prM+E LAV2		
<400> 4		
ttccatata ccacacgtaa cggagaacca cacatgatcg tcagcagaca agagaaaggg		60
aaaagtcttc tggtaaaac agagggtggc gtgaacatgt gtaccctcat ggccatggac		120

eolf-seq1.txt

cttggtaat	tgtgtgaaga	cacaatcacg	tacaagtgtc	cccttcttag	gcagaatgag	180
ccagaagaca	tagactgtt	gtcaactct	acgtccacgt	ggtaactta	tggacgtgt	240
accaccatgg	gagaacatag	aagagaaaaa	agatca	gtgg	cactcg	300
atgggactgg	agacacgaac	tgaaacatgg	atgtcatcg	aagg	gaaacatgtc	360
cagagaattt	aaacttggat	cttggagacat	ccagg	cc	catgtgg	420
gcatacacca	taggaacgac	acatttccaa	agagcc	tta	actgacagct	480
gtcactcctt	caatgacaat	gcgttgcata	ggaatgtca	atagagactt	tgtgg	540
gtttcaggag	gaagctgggt	tgacatagtc	ttagaacatg	gaagctgtgt	gacgacgat	600
gcaaaaaaca	aaccaacatt	ggat	ttgaa	aa	cagaagccaa	660
accctaagga	agtactgtat	agaggcaag	ctaacc	aa	caacaacaga	720
ccaacacaag	gggaacccag	cctaaatgaa	gagcagg	aca	aaagg	780
tccatggtag	acagaggatg	gggaaatgga	tgtggactat	ttggaa	agg	840
acctgtgcta	tgttcagatg	caaaaagaac	atggaaggaa	aagt	ttgtgc	900
ttggaataca	ccattgtgat	aacac	tcac	tca	acc	960
acaggaaaac	atggcaagga	aatcaaata	acacc	aca	gttccat	1020
ttgacaggtt	atggcactgt	cacaatggag	tgct	ctccaa	gaacgg	1080
gagatgggt	tgctgcagat	ggaaaataaa	gcttgg	ctgg	tgcacagg	1140
gacctgccgt	taccatggtt	gcccgagcg	gacaca	caag	gttcaat	1200
gagacattgg	tcacttcaa	aaatccccat	gcgaaga	aaac	aggatgtgt	1260
tcccaagaag	gggcatgca	cacag	caactt	acaggg	cca	1320
ggaaacttac	tcttcacagg	acatctcaag	tgcagg	ctga	aatgg	1380
aaaggaatgt	catacttat	gtgcacagga	aagt	ttaaag	ttgtga	1440
acacaacatg	gaacaatagt	tatcagagtg	caat	atgaa	ggacgg	1500
atccc	ttt	agataatg	tttgg	aaaaa	agacatgt	1560
aacccaattt	tgacagaaaa	agatgccc	gtca	acat	aagcaga	1620
gacagctaca	tcatcatagg	at	tagagcc	ggaca	actg	1680
ggaagttcta	tcggccaaat	gtt	gagaca	acaat	gaggg	1740
ttaggtgaca	cagc	ctgg	gaga	tttggat	ttggagg	1800
gctctccacc	aagt	cttgg	agcaat	ctat	ggagctgc	1860
atgaaaatcc	tcatagg	gtt	atc	ttttacat	tatagg	1920
ctgtctgtga	cactagtatt	gg	tgg	aaatt	gtgacact	1980
gcc						1983

<210> 5
<211> 1983
<212> DNA

eolf-seq1.txt

<213> Artificial Sequence

<220>

<223> prM+E BID/V585

<400>	5	
ttccatttaa ccacacgtaa tggagaacca cacatgatcg ttggtaggca agagaaaggg		60
aaaagtcttc tggtaaaaac agaggatggt gttAACATGT gcaccctcat ggcattAGAC		120
cttggtaat tggtaaga tacaatcACG tacaAGTGCc ccctcCTcAG gcaAAATGAA		180
ccagaAGACA tagATTGTTG gtGCAACTCT acgtccACAT ggtaACTTA tggacATGT		240
accACCAcAG gagaACACAG aAGAGAAAAA agatCAGTGG cactCGTTCC acatGTggGC		300
atgggactgg agacacgAAC tggAAACATGG atgtcatcAG aaggGGGCTG gaaACATGTT		360
cagaGAATTG aaACCTGGAT cttgAGACAT ccaggCTTA ccataATGGC agcaATCCTG		420
gcataTACCA taggaACGAC acATTTCAA AGGGCTCTGA tcttcATTtT actgACAGCC		480
gttGCTCCTT caatgacaAT gcgttgcATA ggaATATCAA atAGAGACTT cgtAGAAAGGG		540
gtttcaggAG gaAGTTGGT tgACATAGTC ttGAACATG gaAGTTGTGt gacgacGATG		600
gcaAAAAATA aACCAACATT ggATTTGAA ctgataAAAAA cagaAGCCAA acaACCTGCC		660
actctaAGGA agtACTGTAT agaAGCAAG ctGACCAATA caACAACAGA atctCGTTGC		720
ccaacacaAG gggAACCCAG tctAAATGAA gagcAGGACA aaAGGTTCAT ctGCAAACAC		780
tccatGGTAG acAGAGGATG gggAAATGGA tggattAT ttggAAAGGG aggCATTGt		840
acctGTGCTA tggcacATG caAAAGAAAC atggAAAGGA aagtCGTGCa gCcAGAAAT		900
ctggAAATACA ccatCGTgAT aacACCTCAC tcaggAGAAG agcACGCTGT aggtAAATGAC		960
acaggAAAGC atggCAAGGA aatCAAATA acaccACAGA gctCCATCAC agaAGCAGAA		1020
ctgacaggCT atggCACTGT cacGATGGAG tgctCTCCGA gaACGGGCT CGACTTCAAT		1080
gagatGGTAC tgctGCAgAT ggaAGACAAA gcttGGCTGG tgCACAGGCA atggTTCCtA		1140
gacCTGCCGT taccatGGCT accCGGAGCG gacACACAAg gatCAAATTG gatacAGAAA		1200
gagacGTTGG tcactTTCAA aaATCCCCAC gcgaAGAAAC aggACGTCGT tgTTTtagGA		1260
tctcaAGAAg gggCCATGCA cacGGCACTT acAGGGGCCA cagaAAATCCA gatgtCATCA		1320
ggAAACTTAC tggcacAGG acatCTCAAG tggAGGCTGA gaATGGACAA attACAGTT		1380
aaAGGAATGT cataCTAT gtgtACAGGA aagtTTAAAA ttgtGAAGGA aatAGCAGAA		1440
acacaACATG gaACAATAGT tatCAGAGTA caATATGAAG gggACGGCTC tccatGTAAG		1500
attcCTTTG agataATGGA tttggAAAAA agACACGTCC taggtCGCCT gattACAGTG		1560
aACCCAAATCG taACAGAAAA agatAGCCCA gtcaACATAG aAGCAGAACc tccattCGGA		1620
gacAGCTACA tcatCATAGG agtagAGCCG ggacaATTGA aactCAATTG gttcaAGAAG		1680
ggaAGTTCCA ttggCCAAAT gttGAGACA ACAATGAGAG gagcGAAGAG aatGGCCATT		1740
ttaggTGACA cAGCCTGGGA tttggatCC ctgggAGGAG tggTTACATC tatAGGAAAG		1800
gctCTCCACC aagtTTTCGG agcaATCTAT gggGCTGCTT ttagtGGGgt ctcATGGACT		1860

eolf-seq1.txt

atgaaaatcc	tcataggagt	tattatcaca	tggataggaa	tgaattcacg	tagcacctca	1920
ctgtctgtgt	cactagtatt	ggtgggagtc	gtgacactgt	acttgggggt	tatggtgtag	1980
gct						1983
<210>	6					
<211>	1983					
<212>	DNA					
<213>	Artificial Sequence					
<220>						
<223>	prM+E PR/DB023					
<400>	6					
ttccatttaa	ccacacgtaa	tggagaacca	cacatgatcg	ttggtaggca	agagaaaaggg	60
aaaagtcttc	tgttcaaaac	agaggatggt	gttaacatgt	gtaccctcat	ggccatagac	120
cttggtaat	tgtgtgaaga	tacaatcacg	tacaagtgcc	ccctcctcag	gcaaaaatgaa	180
ccagaagaca	tagatttttg	gtgcaactct	acgtccacat	ggtaactta	tgggacatgt	240
accaccacag	gagaacacag	aagagaaaaa	agatcagtgg	cactcggtcc	acatgtggc	300
atggactgg	agacacgaaac	tgaaacatgg	atgtcatcg	aaggggcctg	gaaacatgtt	360
cagagaattg	aaacctggat	attgagacat	ccaggcttta	ccataatggc	agcaatcctg	420
gcatatacca	taggaacgac	acatttccaa	agggctctga	tcttcatttt	actgacagcc	480
gtcgctcctt	caatgacaat	gcgttgcata	ggaatatcaa	atagagactt	cgtagaaggg	540
gtttcaggag	gaagttgggt	tgacatagtc	ttagaacatg	gaagttgtgt	gacgacgatg	600
gcaaaaaata	aaccaacatt	ggattttgaa	ctgataaaaa	cagaagccaa	acaacctgccc	660
actctaagga	agtactgtat	agaagcaaag	ctgaccaata	caacaacaga	atctcggtgc	720
ccaacacaag	gggaacccag	tctaaatgaa	gagcaggaca	aaaggttcat	ctgcaaacac	780
tccatggtag	acagaggatg	gggaaatgga	tgtggattat	ttggaaaagg	aggcattgt	840
acctgtgcta	tgttcacatg	caaaaagaac	atggaaggaa	aagttgtgct	gccagaaaat	900
ctggaataca	ccatcggtat	aacacctcac	tcaggagaag	agcacgctgt	agtaatgac	960
acaggaaaac	atggcaagga	aattaaaata	acaccacaga	gttccatcac	agaagcagaa	1020
ctgacaggct	atggcactgt	cacgatggag	tgctctccga	gaacgggcct	cgacttcaat	1080
gagatggtgc	tgctgcagat	ggaagacaaa	gcctggctgg	tgcacaggca	atggttccta	1140
gatctgccgt	taccatggct	acccggagcg	gacacacaag	gatcaaattt	gatacagaaa	1200
gagacgttgg	tcactttcaa	aatccccac	gcgaagaaac	aggacgtcgt	tgttttagga	1260
tctcaagaag	gggccatgca	cacggcactt	acaggggcc	cagaaatcca	gatgtcatca	1320
ggaaacttac	tgttcacagg	acatctcaag	tgtaggctga	aatggacaaa	attacagctt	1380
aaaggaatgt	catactctat	gtgtacagga	aagttaaaa	ttgtgaagga	aatagcagaa	1440
acacaacatg	gaacaatagt	tatcagagta	caatatgaag	gggacggctc	tccatgtaa	1500
attccttttgc	agataatgga	tttagaaaaa	agacacgtcc	taggtcgcc	gattacagtg	1560

	eolf-seq1.txt					
aacccaatcg	taacagaaaa	agatagccca	gtcaacatag	aagcagaacc	tccattcgga	1620
gacagctaca	tcatcatagg	agttagagccg	ggacaattga	aactcaattg	gttcaagaag	1680
ggaagttcca	ttggccaaat	gttgagaca	acaatgagag	gagcgaagag	aatggccatt	1740
ttagggtaca	cagcctggga	tttggatcc	ctgggaggag	tgtttacatc	tatagaaag	1800
gctctccacc	aagtttcgg	agcaatctat	ggggctgctt	ttagtggggt	ctcatggact	1860
atgaaaatcc	tcataggagt	tatcatcaca	tggataggaa	tgaattcacg	tagcacctca	1920
ctgtctgtgt	cactagtatt	ggtgggagtc	gtgacactgt	acttgggggt	tatggtgcag	1980
gct						1983

<210> 7
 <211> 1983
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> prM+E MD1280

<400> 7	ttccattnaa	ccacacgaaa	tggagaacca	cacatgatcg	ttggcagaca	agagaaaggg	60
	aaaagccttc	tgtttaaaac	agaggatggt	gtgaacatgt	gtaccctcat	ggccattgtat	120
	cttggtgaat	tgtgtgaaga	tacaatcacg	tacaagtgcc	ccctcctcag	gcagaatgaa	180
	ccagaagata	tagatttttgc	gtgcaactcc	acgtccacat	ggtaactta	tggacgtgt	240
	accaccacag	gagaacacag	aagagaaaaa	agatcagtgg	cactcggtcc	acatgtgggt	300
	atgggactgg	agacacgaac	tgaacatgg	atgtcgtag	aaggggcctg	gaaacacgct	360
	cagagaatttgc	aaacttggat	cttgagacat	ccaggcttta	ccataatggc	agcaatcctg	420
	gcatataccg	taggaacgac	acatttccaa	agggccctga	tttcatctt	actggcagct	480
	gtcgctcctt	caatgacaat	gcgttgata	ggaatatcaa	atagagactt	tgtagaaggg	540
	gtttcaggag	gaagctgggt	tgacatagtc	ttagaacatg	gaagttgtgt	gacgacaatg	600
	gcaaaaaata	aaccaacact	ggattttgaa	ctgataaaaaa	cagaagccaa	acaacctgcc	660
	actctaagga	agtactgtat	agaggcaag	ctgaccaata	caacaacaga	atctcggtgc	720
	ccaacacaag	gggaacccag	tctaaatgaa	gagcaggaca	aaaggttcgt	ctgcaaacac	780
	tccatggtag	acagaggatg	ggaaaatgga	tgtggattat	ttggaaaggg	aggcattgtg	840
	acctgtgcta	tgttcacatg	caaaaagaac	atggaaggaa	aaatcgatca	accagaaaat	900
	ttgaaataca	ccatcgat	aacacccac	tcaggagaag	agcacgctgt	aggtaatgac	960
	acaggaaaac	atggtaagga	aattaaaata	acaccacaga	gttccatcac	agaagcagaa	1020
	ctgacaggct	atggcacagt	cacgatggag	tgctctccga	gaacgggcct	tgacttcaat	1080
	gagatggtgc	tgctgcagat	ggaagataaa	gcttggctgg	tgcacaggca	atggttccta	1140
	gacctgccgt	taccatggct	accggagcg	gacacacaag	gatcaaatttgc	gatacagaaa	1200
	gagacattgg	tcactttcaa	aaatccccac	gcgaagaagc	aggatgtcgt	tgttttagga	1260

eolf-seq1.txt

tctcaagaag gagccatgca cacggcactc acagggcca cagaaatcca gatgtcatca	1320
ggaaacttac tattcacagg acatctaaa tgcaggctga gaatggacaa actacagctc	1380
aaagaatgt catacttat gtgtacagga aagttaaaa ttgtgaagga aatagcagaa	1440
acacaacatg gaacaatagt tatcagagta caatatgaag gagacggctc tccatgtaag	1500
atcccctttg aaataatgga tttggaaaaa agacatgtct taggtgcgcct gattacagtt	1560
aatccgatcg taacagaaaa agatagccca gtcaacatag aagcagaacc tccattcgga	1620
gacagctaca tcattatagg agtagagccg ggacaattga aactcaactg gttcaagaaa	1680
ggaagttcca tcggccaaat gttgagacg acaatgagag gagcaaagag aatggccatt	1740
ttaggtgaca cagcctggta tttggatct ctgggaggag ttttacatc tatagggaaag	1800
gctctccacc aagtttcgg agcaatctat ggggctgcct ttagtgggt ttcattggact	1860
atgaaaatcc tcataggagt catcatcaca tggataggaa tgaattcacg tagcacctca	1920
ctgtctgtgt cactagtatt ggtggaaatc ataacactgt acttgggagc tatggtgagc	1980
gct	1983

<210> 8
 <211> 661
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> prM+E LAV2

<400> 8

Phe His Leu Thr Thr Arg Asn Gly Glu Pro His Met Ile Val Ser Arg
 1 5 10 15

Gln Glu Lys Gly Lys Ser Leu Leu Phe Lys Thr Glu Val Gly Val Asn
 20 25 30

Met Cys Thr Leu Met Ala Met Asp Leu Gly Glu Leu Cys Glu Asp Thr
 35 40 45

Ile Thr Tyr Lys Cys Pro Leu Leu Arg Gln Asn Glu Pro Glu Asp Ile
 50 55 60

Asp Cys Trp Cys Asn Ser Thr Ser Thr Trp Val Thr Tyr Gly Thr Cys
 65 70 75 80

Thr Thr Met Gly Glu His Arg Arg Glu Lys Arg Ser Val Ala Leu Val
 85 90 95

Pro His Val Gly Met Gly Leu Glu Thr Arg Thr Glu Thr Trp Met Ser
 100 105 110

Ser Glu Gly Ala Trp Lys His Val Gln Arg Ile Glu Thr Trp Ile Leu
 115 120 125

eof-seq1.txt

Arg His Pro Gly Phe Thr Met Met Ala Ala Ile Leu Ala Tyr Thr Ile
130 135 140

Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe Ile Leu Leu Thr Ala
145 150 155 160

Val Thr Pro Ser Met Thr Met Arg Cys Ile Gly Met Ser Asn Arg Asp
165 170 175

Phe Val Glu Gly Val Ser Gly Ser Trp Val Asp Ile Val Leu Glu
180 185 190

His Gly Ser Cys Val Thr Thr Met Ala Lys Asn Lys Pro Thr Leu Asp
195 200 205

Phe Glu Leu Ile Lys Thr Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys
210 215 220

Tyr Cys Ile Glu Ala Lys Leu Thr Asn Thr Thr Glu Ser Arg Cys
225 230 235 240

Pro Thr Gln Gly Glu Pro Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe
245 250 255

Val Cys Lys His Ser Met Val Asp Arg Gly Trp Gly Asn Gly Cys Gly
260 265 270

Leu Phe Gly Lys Gly Gly Ile Val Thr Cys Ala Met Phe Arg Cys Lys
275 280 285

Lys Asn Met Glu Gly Lys Val Val Gln Pro Glu Asn Leu Glu Tyr Thr
290 295 300

Ile Val Ile Thr Pro His Ser Gly Glu Glu His Ala Val Gly Asn Asp
305 310 315 320

Thr Gly Lys His Gly Lys Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile
325 330 335

Thr Glu Ala Glu Leu Thr Gly Tyr Gly Thr Val Thr Met Glu Cys Ser
340 345 350

Pro Arg Thr Gly Leu Asp Phe Asn Glu Met Val Leu Leu Gln Met Glu
355 360 365

Asn Lys Ala Trp Leu Val His Arg Gln Trp Phe Leu Asp Leu Pro Leu
370 375 380

Pro Trp Leu Pro Gly Ala Asp Thr Gln Gly Ser Asn Trp Ile Gln Lys
385 390 395 400

eof-seq1.txt

Glu Thr Leu Val Thr Phe Lys Asn Pro His Ala Lys Lys Gln Asp Val
405 410 415

Val Val Leu Gly Ser Gln Glu Gly Ala Met His Thr Ala Leu Thr Gly
420 425 430

Ala Thr Glu Ile Gln Met Ser Ser Gly Asn Leu Leu Phe Thr Gly His
435 440 445

Leu Lys Cys Arg Leu Arg Met Asp Lys Leu Gln Leu Lys Gly Met Ser
450 455 460

Tyr Ser Met Cys Thr Gly Lys Phe Lys Val Val Lys Glu Ile Ala Glu
465 470 475 480

Thr Gln His Gly Thr Ile Val Ile Arg Val Gln Tyr Glu Gly Asp Gly
485 490 495

Ser Pro Cys Lys Ile Pro Phe Glu Ile Met Asp Leu Glu Lys Arg His
500 505 510

Val Leu Gly Arg Leu Ile Thr Val Asn Pro Ile Val Thr Glu Lys Asp
515 520 525

Ser Pro Val Asn Ile Glu Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile
530 535 540

Ile Ile Gly Val Glu Pro Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys
545 550 555 560

Gly Ser Ser Ile Gly Gln Met Phe Glu Thr Thr Met Arg Gly Ala Lys
565 570 575

Arg Met Ala Ile Leu Gly Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly
580 585 590

Gly Val Phe Thr Ser Ile Gly Lys Ala Leu His Gln Val Phe Gly Ala
595 600 605

Ile Tyr Gly Ala Ala Phe Ser Gly Val Ser Trp Thr Met Lys Ile Leu
610 615 620

Ile Gly Val Ile Ile Thr Trp Ile Gly Met Asn Ser Arg Ser Thr Ser
625 630 635 640

Leu Ser Val Thr Leu Val Leu Val Gly Ile Val Thr Leu Tyr Leu Gly
645 650 655

Val Met Val Gln Ala
660

eolf-seq1.txt

<210> 9
<211> 661
<212> PRT
<213> Artificial Sequence

<220>
<223> prM+E BID/V585

<400> 9

Phe His Leu Thr Thr Arg Asn Gly Glu Pro His Met Ile Val Gly Arg
1 5 10 15

Gln Glu Lys Gly Lys Ser Leu Leu Phe Lys Thr Glu Asp Gly Val Asn
20 25 30

Met Cys Thr Leu Met Ala Ile Asp Leu Gly Glu Leu Cys Glu Asp Thr
35 40 45

Ile Thr Tyr Lys Cys Pro Leu Leu Arg Gln Asn Glu Pro Glu Asp Ile
50 55 60

Asp Cys Trp Cys Asn Ser Thr Ser Thr Trp Val Thr Tyr Gly Thr Cys
65 70 75 80

Thr Thr Thr Gly Glu His Arg Arg Glu Lys Arg Ser Val Ala Leu Val
85 90 95

Pro His Val Gly Met Gly Leu Glu Thr Arg Thr Glu Thr Trp Met Ser
100 105 110

Ser Glu Gly Ala Trp Lys His Val Gln Arg Ile Glu Thr Trp Ile Leu
115 120 125

Arg His Pro Gly Phe Thr Ile Met Ala Ala Ile Leu Ala Tyr Thr Ile
130 135 140

Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe Ile Leu Leu Thr Ala
145 150 155 160

Val Ala Pro Ser Met Thr Met Arg Cys Ile Gly Ile Ser Asn Arg Asp
165 170 175

Phe Val Glu Gly Val Ser Gly Gly Ser Trp Val Asp Ile Val Leu Glu
180 185 190

His Gly Ser Cys Val Thr Thr Met Ala Lys Asn Lys Pro Thr Leu Asp
195 200 205

Phe Glu Leu Ile Lys Thr Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys
210 215 220

Tyr Cys Ile Glu Ala Lys Leu Thr Asn Thr Thr Glu Ser Arg Cys
Page 15

eolf-seq1.txt

225 230 235 240

Pro Thr Gln Gly Glu Pro Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe
245 250 255

Ile Cys Lys His Ser Met Val Asp Arg Gly Trp Gly Asn Gly Cys Gly
260 265 270

Leu Phe Gly Lys Gly Gly Ile Val Thr Cys Ala Met Phe Thr Cys Lys
275 280 285

Lys Asn Met Glu Gly Lys Val Val Gln Pro Glu Asn Leu Glu Tyr Thr
290 295 300

Ile Val Ile Thr Pro His Ser Gly Glu Glu His Ala Val Gly Asn Asp
305 310 315 320

Thr Gly Lys His Gly Lys Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile
325 330 335

Thr Glu Ala Glu Leu Thr Gly Tyr Gly Thr Val Thr Met Glu Cys Ser
340 345 350

Pro Arg Thr Gly Leu Asp Phe Asn Glu Met Val Leu Leu Gln Met Glu
355 360 365

Asp Lys Ala Trp Leu Val His Arg Gln Trp Phe Leu Asp Leu Pro Leu
370 375 380

Pro Trp Leu Pro Gly Ala Asp Thr Gln Gly Ser Asn Trp Ile Gln Lys
385 390 395 400

Glu Thr Leu Val Thr Phe Lys Asn Pro His Ala Lys Lys Gln Asp Val
405 410 415

Val Val Leu Gly Ser Gln Glu Gly Ala Met His Thr Ala Leu Thr Gly
420 425 430

Ala Thr Glu Ile Gln Met Ser Ser Gly Asn Leu Leu Phe Thr Gly His
435 440 445

Leu Lys Cys Arg Leu Arg Met Asp Lys Leu Gln Leu Lys Gly Met Ser
450 455 460

Tyr Ser Met Cys Thr Gly Lys Phe Lys Ile Val Lys Glu Ile Ala Glu
465 470 475 480

Thr Gln His Gly Thr Ile Val Ile Arg Val Gln Tyr Glu Gly Asp Gly
485 490 495

Ser Pro Cys Lys Ile Pro Phe Glu Ile Met Asp Leu Glu Lys Arg His
Page 16

500 505 eolf-seq1.txt 510

Val Leu Gly Arg Leu Ile Thr Val Asn Pro Ile Val Thr Glu Lys Asp
515 520 525

Ser Pro Val Asn Ile Glu Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile
530 535 540

Ile Ile Gly Val Glu Pro Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys
545 550 555 560

Gly Ser Ser Ile Gly Gln Met Phe Glu Thr Thr Met Arg Gly Ala Lys
565 570 575

Arg Met Ala Ile Leu Gly Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly
580 585 590

Gly Val Phe Thr Ser Ile Gly Lys Ala Leu His Gln Val Phe Gly Ala
595 600 605

Ile Tyr Gly Ala Ala Phe Ser Gly Val Ser Trp Thr Met Lys Ile Leu
610 615 620

Ile Gly Val Ile Ile Thr Trp Ile Gly Met Asn Ser Arg Ser Thr Ser
625 630 635 640

Leu Ser Val Ser Leu Val Leu Val Gly Val Val Thr Leu Tyr Leu Gly
645 650 655

Val Met Val Gln Ala
660

<210> 10
<211> 661
<212> PRT
<213> Artificial Sequence

<220>
<223> prM+E PR/DB023

<400> 10

Phe His Leu Thr Thr Arg Asn Gly Glu Pro His Met Ile Val Gly Arg
1 5 10 15

Gln Glu Lys Gly Lys Ser Leu Leu Phe Lys Thr Glu Asp Gly Val Asn
20 25 30

Met Cys Thr Leu Met Ala Ile Asp Leu Gly Glu Leu Cys Glu Asp Thr
35 40 45

Ile Thr Tyr Lys Cys Pro Leu Leu Arg Gln Asn Glu Pro Glu Asp Ile
50 55 60

eof-seq1.txt

Asp Cys Trp Cys Asn Ser Thr Ser Thr Trp Val Thr Tyr Gly Thr Cys
65 70 75 80

Thr Thr Thr Gly Glu His Arg Arg Glu Lys Arg Ser Val Ala Leu Val
85 90 95

Pro His Val Gly Met Gly Leu Glu Thr Arg Thr Glu Thr Trp Met Ser
100 105 110

Ser Glu Gly Ala Trp Lys His Val Gln Arg Ile Glu Thr Trp Ile Leu
115 120 125

Arg His Pro Gly Phe Thr Ile Met Ala Ala Ile Leu Ala Tyr Thr Ile
130 135 140

Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe Ile Leu Leu Thr Ala
145 150 155 160

Val Ala Pro Ser Met Thr Met Arg Cys Ile Gly Ile Ser Asn Arg Asp
165 170 175

Phe Val Glu Gly Val Ser Gly Gly Ser Trp Val Asp Ile Val Leu Glu
180 185 190

His Gly Ser Cys Val Thr Thr Met Ala Lys Asn Lys Pro Thr Leu Asp
195 200 205

Phe Glu Leu Ile Lys Thr Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys
210 215 220

Tyr Cys Ile Glu Ala Lys Leu Thr Asn Thr Thr Glu Ser Arg Cys
225 230 235 240

Pro Thr Gln Gly Glu Pro Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe
245 250 255

Ile Cys Lys His Ser Met Val Asp Arg Gly Trp Gly Asn Gly Cys Gly
260 265 270

Leu Phe Gly Lys Gly Gly Ile Val Thr Cys Ala Met Phe Thr Cys Lys
275 280 285

Lys Asn Met Glu Gly Lys Val Val Leu Pro Glu Asn Leu Glu Tyr Thr
290 295 300

Ile Val Ile Thr Pro His Ser Gly Glu Glu His Ala Val Gly Asn Asp
305 310 315 320

Thr Gly Lys His Gly Lys Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile
325 330 335

eof-seq1.txt

Thr Glu Ala Glu Leu Thr Gly Tyr Gly Thr Val Thr Met Glu Cys Ser
340 345 350

Pro Arg Thr Gly Leu Asp Phe Asn Glu Met Val Leu Leu Gln Met Glu
355 360 365

Asp Lys Ala Trp Leu Val His Arg Gln Trp Phe Leu Asp Leu Pro Leu
370 375 380

Pro Trp Leu Pro Gly Ala Asp Thr Gln Gly Ser Asn Trp Ile Gln Lys
385 390 395 400

Glu Thr Leu Val Thr Phe Lys Asn Pro His Ala Lys Lys Gln Asp Val
405 410 415

Val Val Leu Gly Ser Gln Glu Gly Ala Met His Thr Ala Leu Thr Gly
420 425 430

Ala Thr Glu Ile Gln Met Ser Ser Gly Asn Leu Leu Phe Thr Gly His
435 440 445

Leu Lys Cys Arg Leu Arg Met Asp Lys Leu Gln Leu Lys Gly Met Ser
450 455 460

Tyr Ser Met Cys Thr Gly Lys Phe Lys Ile Val Lys Glu Ile Ala Glu
465 470 475 480

Thr Gln His Gly Thr Ile Val Ile Arg Val Gln Tyr Glu Gly Asp Gly
485 490 495

Ser Pro Cys Lys Ile Pro Phe Glu Ile Met Asp Leu Glu Lys Arg His
500 505 510

Val Leu Gly Arg Leu Ile Thr Val Asn Pro Ile Val Thr Glu Lys Asp
515 520 525

Ser Pro Val Asn Ile Glu Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile
530 535 540

Ile Ile Gly Val Glu Pro Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys
545 550 555 560

Gly Ser Ser Ile Gly Gln Met Phe Glu Thr Thr Met Arg Gly Ala Lys
565 570 575

Arg Met Ala Ile Leu Gly Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly
580 585 590

Gly Val Phe Thr Ser Ile Gly Lys Ala Leu His Gln Val Phe Gly Ala
595 600 605

eolf-seq1.txt

Ile Tyr Gly Ala Ala Phe Ser Gly Val Ser Trp Thr Met Lys Ile Leu
610 615 620

Ile Gly Val Ile Ile Thr Trp Ile Gly Met Asn Ser Arg Ser Thr Ser
625 630 635 640

Leu Ser Val Ser Leu Val Leu Val Gly Val Val Thr Leu Tyr Leu Gly
645 650 655

Val Met Val Gln Ala
660

<210> 11
<211> 661
<212> PRT
<213> Artificial Sequence

<220>
<223> prM+E MD1280

<400> 11

Phe His Leu Thr Thr Arg Asn Gly Glu Pro His Met Ile Val Gly Arg
1 5 10 15

Gln Glu Lys Gly Lys Ser Leu Leu Phe Lys Thr Glu Asp Gly Val Asn
20 25 30

Met Cys Thr Leu Met Ala Ile Asp Leu Gly Glu Leu Cys Glu Asp Thr
35 40 45

Ile Thr Tyr Lys Cys Pro Leu Leu Arg Gln Asn Glu Pro Glu Asp Ile
50 55 60

Asp Cys Trp Cys Asn Ser Thr Ser Thr Trp Val Thr Tyr Gly Thr Cys
65 70 75 80

Thr Thr Thr Gly Glu His Arg Arg Glu Lys Arg Ser Val Ala Leu Val
85 90 95

Pro His Val Gly Met Gly Leu Glu Thr Arg Thr Glu Thr Trp Met Ser
100 105 110

Ser Glu Gly Ala Trp Lys His Ala Gln Arg Ile Glu Thr Trp Ile Leu
115 120 125

Arg His Pro Gly Phe Thr Ile Met Ala Ala Ile Leu Ala Tyr Thr Val
130 135 140

Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe Ile Leu Leu Ala Ala
145 150 155 160

Val Ala Pro Ser Met Thr Met Arg Cys Ile Gly Ile Ser Asn Arg Asp
Page 20

eof-seq1.txt

165

170

175

Phe Val Glu Gly Val Ser Gly Gly Ser Trp Val Asp Ile Val Leu Glu
180 185 190

His Gly Ser Cys Val Thr Thr Met Ala Lys Asn Lys Pro Thr Leu Asp
195 200 205

Phe Glu Leu Ile Lys Thr Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys
210 215 220

Tyr Cys Ile Glu Ala Lys Leu Thr Asn Thr Thr Glu Ser Arg Cys
225 230 235 240

Pro Thr Gln Gly Glu Pro Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe
245 250 255

Val Cys Lys His Ser Met Val Asp Arg Gly Trp Gly Asn Gly Cys Gly
260 265 270

Leu Phe Gly Lys Gly Gly Ile Val Thr Cys Ala Met Phe Thr Cys Lys
275 280 285

Lys Asn Met Glu Gly Lys Ile Val Gln Pro Glu Asn Leu Glu Tyr Thr
290 295 300

Ile Val Ile Thr Pro His Ser Gly Glu Glu His Ala Val Gly Asn Asp
305 310 315 320

Thr Gly Lys His Gly Lys Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile
325 330 335

Thr Glu Ala Glu Leu Thr Gly Tyr Gly Thr Val Thr Met Glu Cys Ser
340 345 350

Pro Arg Thr Gly Leu Asp Phe Asn Glu Met Val Leu Leu Gln Met Glu
355 360 365

Asp Lys Ala Trp Leu Val His Arg Gln Trp Phe Leu Asp Leu Pro Leu
370 375 380

Pro Trp Leu Pro Gly Ala Asp Thr Gln Gly Ser Asn Trp Ile Gln Lys
385 390 395 400

Glu Thr Leu Val Thr Phe Lys Asn Pro His Ala Lys Lys Gln Asp Val
405 410 415

Val Val Leu Gly Ser Gln Glu Gly Ala Met His Thr Ala Leu Thr Gly
420 425 430

Ala Thr Glu Ile Gln Met Ser Ser Gly Asn Leu Leu Phe Thr Gly His
Page 21

435

440

445

Leu Lys Cys Arg Leu Arg Met Asp Lys Leu Gln Leu Lys Gly Met Ser
 450 455 460

Tyr Ser Met Cys Thr Gly Lys Phe Lys Ile Val Lys Glu Ile Ala Glu
 465 470 475 480

Thr Gln His Gly Thr Ile Val Ile Arg Val Gln Tyr Glu Gly Asp Gly
 485 490 495

Ser Pro Cys Lys Ile Pro Phe Glu Ile Met Asp Leu Glu Lys Arg His
 500 505 510

Val Leu Gly Arg Leu Ile Thr Val Asn Pro Ile Val Thr Glu Lys Asp
 515 520 525

Ser Pro Val Asn Ile Glu Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile
 530 535 540

Ile Ile Gly Val Glu Pro Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys
 545 550 555 560

Gly Ser Ser Ile Gly Gln Met Phe Glu Thr Thr Met Arg Gly Ala Lys
 565 570 575

Arg Met Ala Ile Leu Gly Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly
 580 585 590

Gly Val Phe Thr Ser Ile Gly Lys Ala Leu His Gln Val Phe Gly Ala
 595 600 605

Ile Tyr Gly Ala Ala Phe Ser Gly Val Ser Trp Thr Met Lys Ile Leu
 610 615 620

Ile Gly Val Ile Ile Thr Trp Ile Gly Met Asn Ser Arg Ser Thr Ser
 625 630 635 640

Leu Ser Val Ser Leu Val Leu Val Gly Ile Ile Thr Leu Tyr Leu Gly
 645 650 655

Ala Met Val Gln Ala
 660

<210> 12
 <211> 495
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> E universal

eof-seq1.txt

<220>
<221> Variant
<222> (129)..(129)
<223> Xaa can be Val or Ile

<220>
<221> Variant
<222> (308)..(308)
<223> Xaa can be Val or Ile

<400> 12

Met Arg Cys Ile Gly Ile Ser Asn Arg Asp Phe Val Glu Gly Val Ser
1 5 10 15

Gly Gly Ser Trp Val Asp Ile Val Leu Glu His Gly Ser Cys Val Thr
20 25 30

Thr Met Ala Lys Asn Lys Pro Thr Leu Asp Phe Glu Leu Ile Lys Thr
35 40 45

Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys Tyr Cys Ile Glu Ala Lys
50 55 60

Leu Thr Asn Thr Thr Glu Ser Arg Cys Pro Thr Gln Gly Glu Pro
65 70 75 80

Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe Val Cys Lys His Ser Met
85 90 95

Val Asp Arg Gly Trp Gly Asn Gly Cys Gly Leu Phe Gly Lys Gly
100 105 110

Ile Val Thr Cys Ala Met Phe Thr Cys Lys Lys Asn Met Glu Gly Lys
115 120 125

Xaa Val Gln Pro Glu Asn Leu Glu Tyr Thr Ile Val Ile Thr Pro His
130 135 140

Ser Gly Glu Glu His Ala Val Gly Asn Asp Thr Gly Lys His Gly Lys
145 150 155 160

Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile Thr Glu Ala Glu Leu Thr
165 170 175

Gly Tyr Gly Thr Val Thr Met Glu Cys Ser Pro Arg Thr Gly Leu Asp
180 185 190

Phe Asn Glu Met Val Leu Leu Gln Met Glu Asp Lys Ala Trp Leu Val
195 200 205

His Arg Gln Trp Phe Leu Asp Leu Pro Leu Pro Trp Leu Pro Gly Ala
210 215 220

eof-seq1.txt

Asp Thr Gln Gly Ser Asn Trp Ile Gln Lys Glu Thr Leu Val Thr Phe
225 230 235 240

Lys Asn Pro His Ala Lys Lys Gln Asp Val Val Val Leu Gly Ser Gln
245 250 255

Glu Gly Ala Met His Thr Ala Leu Thr Gly Ala Thr Glu Ile Gln Met
260 265 270

Ser Ser Gly Asn Leu Leu Phe Thr Gly His Leu Lys Cys Arg Leu Arg
275 280 285

Met Asp Lys Leu Gln Leu Lys Gly Met Ser Tyr Ser Met Cys Thr Gly
290 295 300

Lys Phe Lys Xaa Val Lys Glu Ile Ala Glu Thr Gln His Gly Thr Ile
305 310 315 320

Val Ile Arg Val Gln Tyr Glu Gly Asp Gly Ser Pro Cys Lys Ile Pro
325 330 335

Phe Glu Ile Met Asp Leu Glu Lys Arg His Val Leu Gly Arg Leu Ile
340 345 350

Thr Val Asn Pro Ile Val Thr Glu Lys Asp Ser Pro Val Asn Ile Glu
355 360 365

Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile Ile Ile Gly Val Glu Pro
370 375 380

Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys Gly Ser Ser Ile Gly Gln
385 390 395 400

Met Phe Glu Thr Thr Met Arg Gly Ala Lys Arg Met Ala Ile Leu Gly
405 410 415

Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly Gly Val Phe Thr Ser Ile
420 425 430

Gly Lys Ala Leu His Gln Val Phe Gly Ala Ile Tyr Gly Ala Ala Phe
435 440 445

Ser Gly Val Ser Trp Thr Met Lys Ile Leu Ile Gly Val Ile Ile Thr
450 455 460

Trp Ile Gly Met Asn Ser Arg Ser Thr Ser Leu Ser Val Ser Leu Val
465 470 475 480

Leu Val Gly Val Val Thr Leu Tyr Leu Gly Val Met Val Gln Ala
485 490 495

eof-seq1.txt

<210> 13
<211> 495
<212> PRT
<213> Artificial Sequence

<220>
<223> E LAV2

<400> 13

Met Arg Cys Ile Gly Met Ser Asn Arg Asp Phe Val Glu Gly Val Ser
1 5 10 15

Gly Gly Ser Trp Val Asp Ile Val Leu Glu His Gly Ser Cys Val Thr
20 25 30

Thr Met Ala Lys Asn Lys Pro Thr Leu Asp Phe Glu Leu Ile Lys Thr
35 40 45

Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys Tyr Cys Ile Glu Ala Lys
50 55 60

Leu Thr Asn Thr Thr Glu Ser Arg Cys Pro Thr Gln Gly Glu Pro
65 70 75 80

Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe Val Cys Lys His Ser Met
85 90 95

Val Asp Arg Gly Trp Gly Asn Gly Cys Gly Leu Phe Gly Lys Gly
100 105 110

Ile Val Thr Cys Ala Met Phe Arg Cys Lys Lys Asn Met Glu Gly Lys
115 120 125

Val Val Gln Pro Glu Asn Leu Glu Tyr Thr Ile Val Ile Thr Pro His
130 135 140

Ser Gly Glu Glu His Ala Val Gly Asn Asp Thr Gly Lys His Gly Lys
145 150 155 160

Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile Thr Glu Ala Glu Leu Thr
165 170 175

Gly Tyr Gly Thr Val Thr Met Glu Cys Ser Pro Arg Thr Gly Leu Asp
180 185 190

Phe Asn Glu Met Val Leu Leu Gln Met Glu Asn Lys Ala Trp Leu Val
195 200 205

His Arg Gln Trp Phe Leu Asp Leu Pro Leu Pro Trp Leu Pro Gly Ala
210 215 220

Asp Thr Gln Gly Ser Asn Trp Ile Gln Lys Glu Thr Leu Val Thr Phe
225 230 235 240

eof-seq1.txt

Lys Asn Pro His Ala Lys Lys Gln Asp Val Val Val Leu Gly Ser Gln
245 250 255

Glu Gly Ala Met His Thr Ala Leu Thr Gly Ala Thr Glu Ile Gln Met
260 265 270

Ser Ser Gly Asn Leu Leu Phe Thr Gly His Leu Lys Cys Arg Leu Arg
275 280 285

Met Asp Lys Leu Gln Leu Lys Gly Met Ser Tyr Ser Met Cys Thr Gly
290 295 300

Lys Phe Lys Val Val Lys Glu Ile Ala Glu Thr Gln His Gly Thr Ile
305 310 315 320

Val Ile Arg Val Gln Tyr Glu Gly Asp Gly Ser Pro Cys Lys Ile Pro
325 330 335

Phe Glu Ile Met Asp Leu Glu Lys Arg His Val Leu Gly Arg Leu Ile
340 345 350

Thr Val Asn Pro Ile Val Thr Glu Lys Asp Ser Pro Val Asn Ile Glu
355 360 365

Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile Ile Ile Gly Val Glu Pro
370 375 380

Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys Gly Ser Ser Ile Gly Gln
385 390 395 400

Met Phe Glu Thr Thr Met Arg Gly Ala Lys Arg Met Ala Ile Leu Gly
405 410 415

Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly Gly Val Phe Thr Ser Ile
420 425 430

Gly Lys Ala Leu His Gln Val Phe Gly Ala Ile Tyr Gly Ala Ala Phe
435 440 445

Ser Gly Val Ser Trp Thr Met Lys Ile Leu Ile Gly Val Ile Ile Thr
450 455 460

Trp Ile Gly Met Asn Ser Arg Ser Thr Ser Leu Ser Val Thr Leu Val
465 470 475 480

Leu Val Gly Ile Val Thr Leu Tyr Leu Gly Val Met Val Gln Ala
485 490 495

<210> 14
<211> 495

eolf-seq1.txt

<212> PRT
<213> Artificial Sequence

<220>
<223> E BID/V585

<400> 14

Met Arg Cys Ile Gly Ile Ser Asn Arg Asp Phe Val Glu Gly Val Ser
1 5 10 15

Gly Gly Ser Trp Val Asp Ile Val Leu Glu His Gly Ser Cys Val Thr
20 25 30

Thr Met Ala Lys Asn Lys Pro Thr Leu Asp Phe Glu Leu Ile Lys Thr
35 40 45

Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys Tyr Cys Ile Glu Ala Lys
50 55 60

Leu Thr Asn Thr Thr Glu Ser Arg Cys Pro Thr Gln Gly Glu Pro
65 70 75 80

Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe Ile Cys Lys His Ser Met
85 90 95

Val Asp Arg Gly Trp Gly Asn Gly Cys Gly Leu Phe Gly Lys Gly
100 105 110

Ile Val Thr Cys Ala Met Phe Thr Cys Lys Lys Asn Met Glu Gly Lys
115 120 125

Val Val Gln Pro Glu Asn Leu Glu Tyr Thr Ile Val Ile Thr Pro His
130 135 140

Ser Gly Glu Glu His Ala Val Gly Asn Asp Thr Gly Lys His Gly Lys
145 150 155 160

Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile Thr Glu Ala Glu Leu Thr
165 170 175

Gly Tyr Gly Thr Val Thr Met Glu Cys Ser Pro Arg Thr Gly Leu Asp
180 185 190

Phe Asn Glu Met Val Leu Leu Gln Met Glu Asp Lys Ala Trp Leu Val
195 200 205

His Arg Gln Trp Phe Leu Asp Leu Pro Leu Pro Trp Leu Pro Gly Ala
210 215 220

Asp Thr Gln Gly Ser Asn Trp Ile Gln Lys Glu Thr Leu Val Thr Phe
225 230 235 240

eof-seq1.txt

Lys Asn Pro His Ala Lys Lys Gln Asp Val Val Val Leu Gly Ser Gln
245 250 255

Glu Gly Ala Met His Thr Ala Leu Thr Gly Ala Thr Glu Ile Gln Met
260 265 270

Ser Ser Gly Asn Leu Leu Phe Thr Gly His Leu Lys Cys Arg Leu Arg
275 280 285

Met Asp Lys Leu Gln Leu Lys Gly Met Ser Tyr Ser Met Cys Thr Gly
290 295 300

Lys Phe Lys Ile Val Lys Glu Ile Ala Glu Thr Gln His Gly Thr Ile
305 310 315 320

Val Ile Arg Val Gln Tyr Glu Gly Asp Gly Ser Pro Cys Lys Ile Pro
325 330 335

Phe Glu Ile Met Asp Leu Glu Lys Arg His Val Leu Gly Arg Leu Ile
340 345 350

Thr Val Asn Pro Ile Val Thr Glu Lys Asp Ser Pro Val Asn Ile Glu
355 360 365

Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile Ile Ile Gly Val Glu Pro
370 375 380

Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys Gly Ser Ser Ile Gly Gln
385 390 395 400

Met Phe Glu Thr Thr Met Arg Gly Ala Lys Arg Met Ala Ile Leu Gly
405 410 415

Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly Gly Val Phe Thr Ser Ile
420 425 430

Gly Lys Ala Leu His Gln Val Phe Gly Ala Ile Tyr Gly Ala Ala Phe
435 440 445

Ser Gly Val Ser Trp Thr Met Lys Ile Leu Ile Gly Val Ile Ile Thr
450 455 460

Trp Ile Gly Met Asn Ser Arg Ser Thr Ser Leu Ser Val Ser Leu Val
465 470 475 480

Leu Val Gly Val Val Thr Leu Tyr Leu Gly Val Met Val Gln Ala
485 490 495

<210> 15

<211> 495

<212> PRT

<213> Artificial Sequence

eof-seq1.txt

<220>

<223> E PR/DB023

<400> 15

Met Arg Cys Ile Gly Ile Ser Asn Arg Asp Phe Val Glu Gly Val Ser
1 5 10 15

Gly Gly Ser Trp Val Asp Ile Val Leu Glu His Gly Ser Cys Val Thr
20 25 30

Thr Met Ala Lys Asn Lys Pro Thr Leu Asp Phe Glu Leu Ile Lys Thr
35 40 45

Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys Tyr Cys Ile Glu Ala Lys
50 55 60

Leu Thr Asn Thr Thr Glu Ser Arg Cys Pro Thr Gln Gly Glu Pro
65 70 75 80

Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe Ile Cys Lys His Ser Met
85 90 95

Val Asp Arg Gly Trp Gly Asn Gly Cys Gly Leu Phe Gly Lys Gly
100 105 110

Ile Val Thr Cys Ala Met Phe Thr Cys Lys Lys Asn Met Glu Gly Lys
115 120 125

Val Val Leu Pro Glu Asn Leu Glu Tyr Thr Ile Val Ile Thr Pro His
130 135 140

Ser Gly Glu Glu His Ala Val Gly Asn Asp Thr Gly Lys His Gly Lys
145 150 155 160

Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile Thr Glu Ala Glu Leu Thr
165 170 175

Gly Tyr Gly Thr Val Thr Met Glu Cys Ser Pro Arg Thr Gly Leu Asp
180 185 190

Phe Asn Glu Met Val Leu Leu Gln Met Glu Asp Lys Ala Trp Leu Val
195 200 205

His Arg Gln Trp Phe Leu Asp Leu Pro Leu Pro Trp Leu Pro Gly Ala
210 215 220

Asp Thr Gln Gly Ser Asn Trp Ile Gln Lys Glu Thr Leu Val Thr Phe
225 230 235 240

Lys Asn Pro His Ala Lys Lys Gln Asp Val Val Val Leu Gly Ser Gln
245 250 255

eolf-seq1.txt

Glu Gly Ala Met His Thr Ala Leu Thr Gly Ala Thr Glu Ile Gln Met
260 265 270

Ser Ser Gly Asn Leu Leu Phe Thr Gly His Leu Lys Cys Arg Leu Arg
275 280 285

Met Asp Lys Leu Gln Leu Lys Gly Met Ser Tyr Ser Met Cys Thr Gly
290 295 300

Lys Phe Lys Ile Val Lys Glu Ile Ala Glu Thr Gln His Gly Thr Ile
305 310 315 320

Val Ile Arg Val Gln Tyr Glu Gly Asp Gly Ser Pro Cys Lys Ile Pro
325 330 335

Phe Glu Ile Met Asp Leu Glu Lys Arg His Val Leu Gly Arg Leu Ile
340 345 350

Thr Val Asn Pro Ile Val Thr Glu Lys Asp Ser Pro Val Asn Ile Glu
355 360 365

Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile Ile Ile Gly Val Glu Pro
370 375 380

Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys Gly Ser Ser Ile Gly Gln
385 390 395 400

Met Phe Glu Thr Thr Met Arg Gly Ala Lys Arg Met Ala Ile Leu Gly
405 410 415

Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly Gly Val Phe Thr Ser Ile
420 425 430

Gly Lys Ala Leu His Gln Val Phe Gly Ala Ile Tyr Gly Ala Ala Phe
435 440 445

Ser Gly Val Ser Trp Thr Met Lys Ile Leu Ile Gly Val Ile Ile Thr
450 455 460

Trp Ile Gly Met Asn Ser Arg Ser Thr Ser Leu Ser Val Ser Leu Val
465 470 475 480

Leu Val Gly Val Val Thr Leu Tyr Leu Gly Val Met Val Gln Ala
485 490 495

<210> 16
<211> 495

<212> PRT

<213> Artificial Sequence

<220>

eolf-seq1.txt

<223> E MD1280

<400> 16

Met Arg Cys Ile Gly Ile Ser Asn Arg Asp Phe Val Glu Gly Val Ser
1 5 10 15

Gly Gly Ser Trp Val Asp Ile Val Leu Glu His Gly Ser Cys Val Thr
20 25 30

Thr Met Ala Lys Asn Lys Pro Thr Leu Asp Phe Glu Leu Ile Lys Thr
35 40 45

Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys Tyr Cys Ile Glu Ala Lys
50 55 60

Leu Thr Asn Thr Thr Glu Ser Arg Cys Pro Thr Gln Gly Glu Pro
65 70 75 80

Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe Val Cys Lys His Ser Met
85 90 95

Val Asp Arg Gly Trp Gly Asn Gly Cys Gly Leu Phe Gly Lys Gly
100 105 110

Ile Val Thr Cys Ala Met Phe Thr Cys Lys Lys Asn Met Glu Gly Lys
115 120 125

Ile Val Gln Pro Glu Asn Leu Glu Tyr Thr Ile Val Ile Thr Pro His
130 135 140

Ser Gly Glu Glu His Ala Val Gly Asn Asp Thr Gly Lys His Gly Lys
145 150 155 160

Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile Thr Glu Ala Glu Leu Thr
165 170 175

Gly Tyr Gly Thr Val Thr Met Glu Cys Ser Pro Arg Thr Gly Leu Asp
180 185 190

Phe Asn Glu Met Val Leu Leu Gln Met Glu Asp Lys Ala Trp Leu Val
195 200 205

His Arg Gln Trp Phe Leu Asp Leu Pro Leu Pro Trp Leu Pro Gly Ala
210 215 220

Asp Thr Gln Gly Ser Asn Trp Ile Gln Lys Glu Thr Leu Val Thr Phe
225 230 235 240

Lys Asn Pro His Ala Lys Lys Gln Asp Val Val Val Leu Gly Ser Gln
245 250 255

eof-seq1.txt

Glu Gly Ala Met His Thr Ala Leu Thr Gly Ala Thr Glu Ile Gln Met
260 265 270

Ser Ser Gly Asn Leu Leu Phe Thr Gly His Leu Lys Cys Arg Leu Arg
275 280 285

Met Asp Lys Leu Gln Leu Lys Gly Met Ser Tyr Ser Met Cys Thr Gly
290 295 300

Lys Phe Lys Ile Val Lys Glu Ile Ala Glu Thr Gln His Gly Thr Ile
305 310 315 320

Val Ile Arg Val Gln Tyr Glu Gly Asp Gly Ser Pro Cys Lys Ile Pro
325 330 335

Phe Glu Ile Met Asp Leu Glu Lys Arg His Val Leu Gly Arg Leu Ile
340 345 350

Thr Val Asn Pro Ile Val Thr Glu Lys Asp Ser Pro Val Asn Ile Glu
355 360 365

Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile Ile Ile Gly Val Glu Pro
370 375 380

Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys Gly Ser Ser Ile Gly Gln
385 390 395 400

Met Phe Glu Thr Thr Met Arg Gly Ala Lys Arg Met Ala Ile Leu Gly
405 410 415

Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly Gly Val Phe Thr Ser Ile
420 425 430

Gly Lys Ala Leu His Gln Val Phe Gly Ala Ile Tyr Gly Ala Ala Phe
435 440 445

Ser Gly Val Ser Trp Thr Met Lys Ile Leu Ile Gly Val Ile Ile Thr
450 455 460

Trp Ile Gly Met Asn Ser Arg Ser Thr Ser Leu Ser Val Ser Leu Val
465 470 475 480

Leu Val Gly Ile Ile Thr Leu Tyr Leu Gly Ala Met Val Gln Ala
485 490 495

<210> 17
<211> 75
<212> PRT
<213> Artificial Sequence

<220>
<223> M consensus

eof-seq1.txt

<400> 17

Ser Val Ala Leu Val Pro His Val Gly Met Gly Leu Glu Thr Arg Thr
1 5 10 15

Glu Thr Trp Met Ser Ser Glu Gly Ala Trp Lys His Val Gln Arg Ile
20 25 30

Glu Thr Trp Ile Leu Arg His Pro Gly Phe Thr Ile Met Ala Ala Ile
35 40 45

Leu Ala Tyr Thr Ile Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe
50 55 60

Ile Leu Leu Thr Ala Val Ala Pro Ser Met Thr
65 70 75

<210> 18

<211> 495

<212> PRT

<213> Artificial Sequence

<220>

<223> E circulating strain

<400> 18

Met Arg Cys Ile Gly Ile Ser Asn Arg Asp Phe Val Glu Gly Val Ser
1 5 10 15

Gly Gly Ser Trp Val Asp Ile Val Leu Glu His Gly Ser Cys Val Thr
20 25 30

Thr Met Ala Lys Asn Lys Pro Thr Leu Asp Phe Glu Leu Ile Lys Thr
35 40 45

Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys Tyr Cys Ile Glu Ala Lys
50 55 60

Leu Thr Asn Thr Thr Glu Ser Arg Cys Pro Thr Gln Gly Glu Pro
65 70 75 80

Ser Leu Lys Glu Glu Gln Asp Lys Arg Phe Val Cys Lys His Ser Met
85 90 95

Val Asp Arg Gly Trp Gly Asn Gly Cys Gly Leu Phe Gly Lys Gly Gly
100 105 110

Ile Val Thr Cys Ala Met Phe Thr Cys Lys Lys Asn Met Glu Gly Lys
115 120 125

Ile Val Gln Pro Glu Asn Leu Glu Tyr Thr Ile Val Val Thr Pro His
130 135 140

eof-seq1.txt

Ser Gly Glu Glu His Ala Val Gly Asn Asp Thr Gly Lys His Gly Lys
145 150 155 160

Glu Ile Lys Val Thr Pro Gln Ser Ser Ile Thr Glu Ala Glu Leu Thr
165 170 175

Gly Tyr Gly Thr Val Thr Met Glu Cys Ser Pro Arg Thr Gly Leu Asp
180 185 190

Phe Asn Glu Met Val Leu Leu Gln Met Glu Asn Lys Ala Trp Leu Val
195 200 205

His Arg Gln Trp Phe Leu Asp Leu Pro Leu Pro Trp Leu Pro Gly Ala
210 215 220

Asp Lys Gln Glu Ser Asn Trp Ile Gln Lys Glu Thr Leu Val Thr Phe
225 230 235 240

Lys Asn Pro His Ala Lys Lys Gln Asp Val Val Val Leu Gly Ser Gln
245 250 255

Glu Gly Ala Met His Thr Ala Leu Thr Gly Ala Thr Glu Ile Gln Met
260 265 270

Ser Ser Gly Asn Leu Leu Phe Thr Gly His Leu Lys Cys Arg Leu Arg
275 280 285

Met Asp Lys Leu Gln Leu Lys Gly Met Ser Tyr Ser Met Cys Thr Gly
290 295 300

Lys Phe Lys Val Val Lys Glu Ile Ala Glu Thr Gln His Gly Thr Ile
305 310 315 320

Val Ile Arg Val Gln Tyr Glu Gly Asp Gly Ser Pro Cys Lys Ile Pro
325 330 335

Phe Glu Ile Met Asp Leu Glu Lys Arg Tyr Val Leu Gly Arg Leu Ile
340 345 350

Thr Val Asn Pro Ile Val Thr Glu Lys Asp Ser Pro Val Asn Ile Glu
355 360 365

Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile Ile Ile Gly Val Glu Pro
370 375 380

Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys Gly Ser Ser Ile Gly Gln
385 390 395 400

Met Phe Glu Thr Thr Met Arg Gly Ala Lys Arg Met Ala Ile Leu Gly
405 410 415

eof-seq1.txt

Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly Gly Val Phe Thr Ser Ile
420 425 430

Gly Lys Ala Leu His Gln Val Phe Gly Ala Ile Tyr Gly Ala Ala Phe
435 440 445

Ser Gly Val Ser Trp Thr Met Lys Ile Leu Ile Gly Val Ile Ile Thr
450 455 460

Trp Ile Gly Met Asn Ser Arg Ser Thr Ser Leu Ser Val Ser Leu Val
465 470 475 480

Leu Val Gly Ile Val Thr Leu Tyr Leu Gly Val Met Val Gln Ala
485 490 495

<210> 19

<211> 75

<212> PRT

<213> Artificial Sequence

<220>

<223> M LAV2

<400> 19

Ser Val Ala Leu Val Pro His Val Gly Met Gly Leu Glu Thr Arg Thr
1 5 10 15

Glu Thr Trp Met Ser Ser Glu Gly Ala Trp Lys His Val Gln Arg Ile
20 25 30

Glu Thr Trp Ile Leu Arg His Pro Gly Phe Thr Met Met Ala Ala Ile
35 40 45

Leu Ala Tyr Thr Ile Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe
50 55 60

Ile Leu Leu Thr Ala Val Thr Pro Ser Met Thr
65 70 75

<210> 20

<211> 75

<212> PRT

<213> Artificial Sequence

<220>

<223> M BID/V585

<400> 20

Ser Val Ala Leu Val Pro His Val Gly Met Gly Leu Glu Thr Arg Thr
1 5 10 15

Glu Thr Trp Met Ser Ser Glu Gly Ala Trp Lys His Val Gln Arg Ile
20 25 30

eolf-seq1.txt

Glu Thr Trp Ile Leu Arg His Pro Gly Phe Thr Ile Met Ala Ala Ile
35 40 45

Leu Ala Tyr Thr Ile Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe
50 55 60

Ile Leu Leu Thr Ala Val Ala Pro Ser Met Thr
65 70 75

<210> 21

<211> 75

<212> PRT

<213> Artificial Sequence

<220>

<223> M PR/DB023

<400> 21

Ser Val Ala Leu Val Pro His Val Gly Met Gly Leu Glu Thr Arg Thr
1 5 10 15

Glu Thr Trp Met Ser Ser Glu Gly Ala Trp Lys His Val Gln Arg Ile
20 25 30

Glu Thr Trp Ile Leu Arg His Pro Gly Phe Thr Ile Met Ala Ala Ile
35 40 45

Leu Ala Tyr Thr Ile Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe
50 55 60

Ile Leu Leu Thr Ala Val Ala Pro Ser Met Thr
65 70 75

<210> 22

<211> 75

<212> PRT

<213> Artificial Sequence

<220>

<223> M MD1280

<400> 22

Ser Val Ala Leu Val Pro His Val Gly Met Gly Leu Glu Thr Arg Thr
1 5 10 15

Glu Thr Trp Met Ser Ser Glu Gly Ala Trp Lys His Ala Gln Arg Ile
20 25 30

Glu Thr Trp Ile Leu Arg His Pro Gly Phe Thr Ile Met Ala Ala Ile
35 40 45

Leu Ala Tyr Thr Val Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe
50 55 60

eolf-seq1.txt

Ile Leu Leu Ala Ala Val Ala Pro Ser Met Thr
65 70 75

<210> 23
<211> 75
<212> PRT
<213> Artificial Sequence

<220>
<223> M clinical trial circulating strain

<400> 23

Ser Val Ala Leu Val Pro His Val Gly Met Gly Leu Glu Thr Arg Thr
1 5 10 15

Glu Thr Trp Met Ser Ser Glu Gly Ala Trp Lys His Ala Gln Arg Ile
20 25 30

Glu Thr Trp Ile Leu Arg His Pro Gly Phe Thr Ile Met Ala Ala Ile
35 40 45

Leu Ala Tyr Thr Ile Gly Thr Thr His Phe Gln Arg Val Leu Ile Phe
50 55 60

Ile Leu Leu Thr Ala Val Ala Pro Ser Met Thr
65 70 75

<210> 24
<211> 10723
<212> RNA
<213> Artificial Sequence

<220>
<223> VDV2

<400> 24
aguuguuagu cuacguggac cgacaaagac agauucuuug agggagcuuaa gcucaaugua 60
guucuaacag uuuuuuuuauu agagagcaga ucucugauga auaaccaacg gaaaaaggcg 120
aaaaacacgc cuucaauau gcugaaacgc gagagaaacc gcgugucgac ugugcaacag 180
cugacaaaga gauucucacu ugaaugcug cagggacgag gaccauuuaa acuguucaug 240
gcccuggugg cguuccuucg uuuccuaaca aucccacca cagcagggau auugaagaga 300
uggggaacaa uuaaaaaauc aaaagcuauu aauguuuuga gaggguucag gaaagagauu 360
ggaaggaugc ugaacaucuu gaauaggaga cgcagaucug caggcaugau cauuaugcug 420
auuccaacag ugauggcgua ccauuuaacc acacguaacg gagaaccaca caugaucguc 480
agcagacaag agaaaggaa aagucuucug uuuaaaacag agguuggcgu gaacaugugu 540
acccucaugg ccauggaccu ugugaaauug ugugaagaca caaucacgua caagugucc 600
cuucucaggc agaaugagcc agaagacaua gacuguuggu gcaacucuac guccacgugg 660
guaacuuuaug ggacguguac caccauggga gaacauagaa gagaaaaaaag aucaguggca 720

eolf-seq1.txt	
cucguuccac	780
augugcgaau	
gggacuggag	
acacgaacug	
aaacauggau	
gucaucagaa	
ggggccugga	840
aacaugucca	
gagaauugaa	
acuuggaucu	
ugagacaucc	
aggcuucacc	
augauggcag	900
caauccuggc	
auacaccaua	
ggaacgacac	
auuuuccaaag	
agcccugauu	
uuucaucuuac	960
ugacagcugu	
cacuccuuca	
augacaaugc	
guugcauagg	
aaugucaaau	
agagacuuug	1020
uggaaggggu	
uucaggagga	
agcuggguug	
acauagucuu	
agaacaugga	
agcuguguga	1080
cgacgauggc	
aaaaaacaaa	
ccaacauugg	
auuuugaacu	
gauaaaaaca	
gaagccaaac	1140
agccugccac	
ccuaaggaag	
uacuguauag	
aggcaaagcu	
aaccaacaca	
acaacagaau	1200
cucgcugccc	
aacacaaggg	
gaacccagcc	
uaaaugaaga	
gcaggacaaa	
agguucgucu	1260
gcaaacacuc	
caugguagac	
agaggauggg	
gaaauggaugg	
uggacuauuu	
ggaaaggggag	1320
gcauugugac	
cugugcuau	
uucagaugca	
aaaagaacau	
ggaaggaaaa	
guugugcaac	1380
cagaaaacuu	
ggaauacacc	
auugugauaa	
caccucacuc	
aggggaagag	
caugcagucg	1440
gaaaugacac	
aggaaaacau	
ggcaaggaaa	
ucaaaauuaac	
accacagagu	
uccaucacag	1500
aagcagaauu	
gacagguuau	
ggcacuguca	
caauggagug	
cucuccaaga	
acgggccucg	1560
acuucaauga	
gaugguguug	
cugcagauug	
aaaauuaaagc	
uuggcuggug	
cacaggcaau	1620
gguuccuaga	
ccugccguua	
ccaugguugc	
ccggagcgga	
cacacaagag	
ucaaaauugga	1680
uacagaagga	
gacauugguc	
acuuucaaaa	
aucccauagc	
gaagaaacag	
gauguuguug	1740
uuuuaggauc	
ccaagaaggg	
gccaugcaca	
cagcacuuac	
aggggccaca	
gaaaucaaaa	1800
ugucaucagg	
aaacuuacuc	
uucacaggac	
aucucaugug	
caggcugaga	
auggacaagc	1860
uacagcucaa	
aggaauguca	
uacucuaugu	
gcacagggaaa	
guuuuaaguu	
gugaaggaaa	1920
uagcagaaac	
acaacaugga	
acaauaguua	
ucagagugca	
auaugaagggg	
gacggcucuc	1980
caugcaagau	
cccuuuugag	
auaauggauu	
uggaaaaaaag	
acaugucuuua	
ggucgccuga	2040
uuacagucaa	
cccaauuugug	
acagaaaaag	
auagcccagu	
caacauagaa	
gcagaaccuc	2100
cauuuggaga	
cagcuacauc	
aucauaggag	
uagagccggg	
acaacugaag	
cuacauggu	2160
uuuagaaagg	
aaguucuauc	
ggccaaauug	
uugagacaac	
aaugaggggg	
gccaaggagaa	2220
uggccauuuu	
aggugacaca	
gccugggauu	
uuggauccuu	
gggaggagug	
uuuacaucua	2280
uaggaaaggc	
ucuccaccaa	
gucuuuggag	
caaucuaugg	
agcugccuuc	
agugggguuu	2340
cauggacuau	
gaaaauccuc	
auaggaguca	
uuaucacaug	
gauaggaaug	
aauucacgca	2400
gcaccucacu	
gucugugaca	
cuaguauugg	
ugggaaauugu	
gacacuguau	
uugggaguca	2460
uggugcaggc	
cgauaguggu	
ugcguuguga	
gcuggaaaaaa	
caaagaacug	2520
aaauguggca	
gugggauuuu	
caucacagac	
aacgugcaca	
cauggacaga	
acaauacaaa	
uuccaaccag	2580
aauccccuiuc	
aaaacuagcu	
ucagcuaucc	
agaaagccca	
ugaagaggac	
auuuguggaa	2640
uccgcucagu	
aacaagacug	
gagaaucuga	
uguggaaaca	
aauaacacca	
gaauguaauc	2700
acauiucuauc	
agaaaaugag	
gugaaguuaa	
cuauuaugac	
aggagacauc	
aaaggaaauca	2760
ugcaggcagg	
aaaacgaucu	
cugcgccuc	
agcccacuga	
gcugaaguau	

eolf-seq1.txt		
ucauggaaaa	cauggggcaa	2820
agcaaaaaug	cucucuacag	
cucauugaug	agcagaaugc	2880
gccccgaaac	cccaacacaa	
aaagagcuug	gaauucguug	
gaaguugaag	acuauggcuu	2940
uggaguauuc	accaccaaua	
uauggcuaaa	uauggaaagaa	
aaacaggaug	uaauucugcga	3000
cucaaaacuc	augucagcgg	
ccauaaaaga	caacagagcc	
guccaugccg	auauggguua	3060
uuggauagaa	agugcacuca	
augacacaaug	augacauagag	
aaagccucuu	ucauugaagu	3120
uaaaaacugc	cacuggccaa	
aaucacacac	ccucuggagc	
aauggagugc	uagaaaguga	3180
gaugauaauu	ccaaagaauc	
ucgcuggacc	agugucucaa	
cacaacuaua	gaccaggcua	3240
ccauacacaa	auaacaggac	
cauggcaucu	agguaagcuu	
gagauggacu	uugauuucug	3300
ugauggaaca	acagugguag	
ugacugagga	cugcggaaau	
agaggacccu	cuuugagaac	3360
aaccacugcc	ucuggaaaac	
ucauaacaga	auggugcugc	
cgaucuugca	cauuaccacc	3420
gcuaagauac	agaggugagg	
augggugcug	guacgggaug	
gaaaucagac	cauugaagga	3480
gaaagaagag	aauuugguca	
acuccuuggu	cacagcugga	
caugggcagg	ucgacaacuu	3540
uucacuagga	gucuugggaa	
uggcauuguu	ccuggaggaa	
augcuuagga	cccagliagg	3600
aacgaaacau	gcaauacuac	
uaguugcagu	uucuuuugug	
acauugauca	cagggAACAU	3660
cagggAACAU	guccuuuaga	
gaccugggaa	gagugauggu	
uaugguaggc	uaugguaggc	
gccacuauga	cggaugacau	3720
agguauuggc	agguauuggc	
gugacuuau	uugcccuacu	
agcagccuuc	agcagccuuc	
aaagucagac	caacuuuugc	3780
agcuggacua	cucuugagaa	
cucuugagaa	agcugaccuc	
caaggaauug	caaggaauug	
augaugacua	cuauaggaau	3840
uguacuccuc	ucccagagca	
ucccagagca	ccauaccaga	
gaccuuucuu	gaccuuucuu	
gaguugacug	augcguuagc	3900
cuuaggcaug	cuuaggcaug	
augguucca	aaauggugag	
aaauauggaa	aaauauggaa	
aaguaucaau	uggcagugac	3960
uaucuaggc	uaucuaggc	
aucuugugcg	ucccaaacgc	
ucccaaacgc	agugauauua	
caaaaacgcau	ggaaagugag	4020
uugcacaaua	uugcacaaua	
uuggcagugg	uuggcagugg	
uguuccuuuc	cccacuguuc	
uuaacauucc	cacagcaaaa	4080
aacagauugg	aacagauugg	
auaccuuuag	auaccuuuag	
cauugacgau	caaaggucuc	
aauccaacag	caaauuuuicu	4140
aacaacccuc	aacaacccuc	
ucaagaacca	gcaagaaaaag	
gcaagaaaaag	gagcugggca	
uuaaaugagg	uuaaaugagg	4200
cuaucuaggc	agucgggaug	
gugagcauuu	gugagcauuu	
uagccaguuc	uicuccuaaaa	
uicuccuaaaa	aaugauauic	4260
ccaugacagg	ccaugacagg	
accuuuagug	gcuggagggc	
gcuggagggc	uccucacugu	
gugcuacug	gugcuacug	
cucacuggac	gauccggccga	4320
uuuggaacug	uuuggaacug	
gagagagcag	ccgaugucaa	
ccgaugucaa	augggaaagac	
caggcagaga	uaucaggaag	4380
caguuccauc	caguuccauc	
cugucaaua	caaauaucaga	
caaauaucaga	agaugguagc	
augucgauaa	aaaaugaaga	4440
ggaagaacaa	ggaagaacaa	
acacugacca	uacuauuag	
uacuauuag	aacaggauug	
cuggugaucu	caggacuuuu	4500
uccuguaucu	auaccaauca	
auaccaauca	cggcagcagc	
cggcagcagc	augguaccug	
ugggaaguga	agaaacaacg	4560
ggccggagua	ggccggagua	
uugugggaug	uuccuucacc	
uuccuucacc	cccacccaug	
ggaaaggcug	ggaaaggcug	4620
aacuggaaga	uggagccuau	
agaauuaagc	agaauuaagc	
aaaaagggau	ucuuggauau	
ucuuggauau	uccccagau	4680
uccccagau	gagccggagu	
uuacaaagaa	ggaacauucc	
ggaacauucc	auacaaugug	
gcaugucaca	gcaugucaca	
cguggcgcug	uucuuaugca	4740
uucuuaugca	uaaaggaaag	
aggauugaac	caacaugggc	
caacaugggc	ggacgucaag	
aaagaccuaa	aaagaccuaa	4800
uaucauaugg	aggaggcugg	
aggaggcugg	aaguuagaag	
aaguuagaag	gagaauuggaa	
gagaauuggaa	ggaaggagaa	

eolf-seq1.txt		
gaaguccagg	uaauuggcacu	4860
ggucuuuuca	aaaccaacgc	4920
acgucaggau	cuccaaauua	4980
guuguuacaa	ggaguggagc	5040
gacaacccag	agaucgaaga	5100
cacccaggag	cgaaaagac	5160
cggguuuga	gaacauuaau	5220
gccuuuagag	gacuuccaaau	5280
cgggagauug	uggaccuaau	5340
agagugccaa	acuacaaccu	5400
auagcagcua	gaggauacau	5460
augacagcca	cucccccggg	5520
gaugaagaaa	gagaaauccc	5580
uuuuaaggga	agacuguuug	5640
ugccugagga	aaaauggaaa	5700
uaugucaaga	cuagaacaa	5760
ggugccaaau	ucaaggcuga	5820
cuaacagaug	gaggguuaua	5880
gcagcacaaa	gaagagggag	5940
uacauugggg	aaccucugga	6000
cuccuagaua	acaucacac	6060
ggaaaaggugg	augccauuga	6120
guagacuua	ugagaagagg	6180
acgcagacag	aagguggugu	6240
gaagaaaacg	uggaaguuga	6300
agaugguugg	aaucuggaca	6360
gcagccgaa	gaaagucucu	6420
uucaugacuc	gacccugaac	6480
gcagguggaa	cuaaucacag	6540
ciuiiuacuga	ccacuggcgc	6600
aggggcauag	uacagucacg	6660
ugguacgcac	ggagggaua	6720
auaguuuuugc	aaauacagcc	6780
uacguuguca	acacuggaua	6840
uagccauccu	gcaguggug	
cacagacca	gcccgaacca	
uggcaaacga	gauggguuuc	

eolf-seq1.txt			
cuagaaaaaa	cgaagaaaga	ucucggauug	6900
aacauccugg	acauagaucu	acguccugca	6960
acaauuuguua	caccaaughuu	gagacauagc	7020
acagcuauag	ccaaccaagc	cacaguguua	7080
aagauggaca	ucggaguucc	ccuucucgccc	7140
acucucacag	cagcucuuuu	cuuauuggua	7200
caagcaaag	caaccagaga	agcucagaaa	7260
acugucgaug	gaauaacagu	gauugaccua	7320
aagcaguugg	gacaaguauu	gcuccuaguc	7380
acuacauggg	cucuguguga	ggcuuuuacc	7440
gaaggaaauc	cagggaggguu	uuggaacacu	7500
agagggaguu	acuuggccgg	agcuggacuu	7560
acaagaaggg	gaacuggcaa	cauaggagag	7620
aacgc当地	gaaaaaguga	auuccagauc	7680
agaaccuuag	caaagaagg	cauuaaaaga	7740
ggcucagcaa	aacugagaug	guucguugag	7800
guggaccucg	guuguggcag	aggaggcugg	7860
agagaaguca	aaggccuaac	aaaaggagga	7920
acauaugggu	gaaaucuagu	gcgucuucaa	7980
gaaaagugug	acacauuauu	gugugacaua	8040
gcaggacgaa	cacucagagu	ccuuaacuu	8100
uuuugcauua	agguucucaa	cccuaauaug	8160
caaaggaaau	auggaggagc	cuuagugagg	8220
auguacuggg	uauccaaugc	uuccgggaac	8280
auguugauca	acaguuuac	aaugagauac	8340
cucggaagcg	gaacccguaa	caucggauu	8400
gggaaaagaa	uagaaaaaaau	aaagcaagag	8460
cacccauaca	aaacgugggc	auaccauggu	8520
ucauccaugg	ucaacggagu	ggucaggcug	8580
gugacacaga	uggcaaughac	agacacgacu	8640
aaaguggaca	cgagaaccca	agaaccgaaa	8700
gcagaguggc	uuuggaaaga	auuagggaag	8760
gaaucacaa	gaaaggugag	aagcaaugca	8820
aaguggaagu	cggcacguga	ggcuguugaa	8880

	eolf-seq1.txt	
gaaaggaauc	uccaucuuga	8940
agagagaaga	aggaaagugu	9000
uggcuuggag	aguacaacau	9060
uucuccagag	gaucacacugg	9120
auucuaagag	ggagcaaugu	9180
gaaggagaac	augccgauga	9240
acaagaaacu	caccgcagga	9300
gugcgugugc	aaaccacacu	9360
caaagaggua	agaagaccua	9420
acagaagaaa	aaaaaaugaag	9480
uggccauca	agaagguaac	9540
uuaacagcuc	aaacucuuac	9600
agaggaugga	ggcugcggaaag	9660
augaaagacg	guuacuacu	9720
gcccgaauu	gggaaagauu	9780
uacgcccuaa	aggaaagaca	9840
gcuauuugcu	uacacauugg	9900
caugcuuac	guuccaaca	9960
auucaagaaa	gacaaacggaa	10020
uacuugggaa	gacaugcuga	10080
accuaggaa	cacagacgca	10140
accuaggca	accucaggcu	10200
agaacaucc	ggcugcggaa	10260
guucugugg	ggcugcggaa	10320
aaaaaacauu	gcuaccugug	10380
gguuaggggaa	agccccgucc	10440
agcuguaguc	aaggacguua	10500
ucgcuggaag	aaagaaguca	10560
gacuagaggu	ggggggccaa	10620
uagaggagac	ggcugcggaa	10680
ccccccgaaa	cauauugacg	10723
cauauugacg	cuggaaaga	
ccagagaucc	ccagagaucc	
ugcugucucc	ugcugucucc	
ucagcaucau	ucagcaucau	
uccaggcaca	uccaggcaca	
gaacgccaga	gaacgccaga	
aaauggaaug	aaauggaaug	
gugcuguuga	gugcuguuga	
aucaacaggu	aucaacaggu	
ucu	ucu	

<210> 25
 <211> 1983
 <212> RNA
 <213> Artificial Sequence

eolf-seq1.txt

<220>
<223> prM+E VDV2

<400> 25		
uuccauuuuaa ccacacguua cggagaacca cacaugaucg ucagcagaca agagaaaaggg	60	
aaaagucuuc uguuuuaaaac agagguuggc guagaacaugu guacccucau ggccauggac	120	
cuuggugaaau ugugugaaga cacaaucacg uacaaguguc cccuucucag gcagaaugag	180	
ccagaagaca uagacuguug gugcaacucu acguccacgu ggguaacuuua uggaacgugu	240	
accaccaugg gagaacauag aagagaaaaa agaucagugg cacucguucc acaugugcga	300	
augggacugg agacacgaac ugaaacaugg augucaucag aaggggcccug gaaacauguc	360	
cagagaauug aaacuuggau cuugagacau ccagguuca ccaugauuggc agcaauccug	420	
gcauacacca uaggaacgac acauuuccaa agagcccuga uuuucaucuu acugacagcu	480	
gucacuccuu caaugacaau gcguugcaua ggaaugucaa auagagacuu uguggaaggg	540	
guuucaggag gaagcugggu ugacauaguc uuagaacaug gaagcugugu gacgacgaug	600	
gcaaaaaaca aaccaacauu ggauuuugaa cugauaaaaa cagaagccaa acagccugcc	660	
acccuaagga aguacuguau agaggcaaag cuaaccaaca caacaacaga aucucgcugc	720	
ccaacacaag gggAACCCAG ccuaauugaa gagcaggaca aaagguucgu cugcaaacac	780	
uccaugguag acagaggaug gggaaauugga uguggacuau uuggaaaggg aggcauugug	840	
accugugcua uguucagaug caaaaagaac auggaaggaa aaguugugca accagaaaac	900	
uuggaaauaca ccauugugau aacaccucac ucagggaaag agcaugcagu cggaaauugac	960	
acaggaaaac auggcaagga aauaaaaaua acaccacaga guuccaucac agaagcagaa	1020	
uugacagguu auggcacugu cacaauuggag ugcucuccaa gaacgggcccug cgaauucaau	1080	
gagauggugu ugcugcagau ggaaaaauaaa gcuuggcugg ugcacaggca augguuccua	1140	
gaccugccgu uaccaugguu gcccggagcg gacacacaag agucaaauug gauacagaag	1200	
gagacauuugg ucacuuucaa aaaucuccau gcgaagaaac aggauguugu uguuuuagga	1260	
uccaagaag gggcaugca cacagcacuu acaggggcca cagaaaucca aaugucauca	1320	
ggaaacuuac uciuucacagg acaucucaag ugcaggcuga gaauggacaa gcuacagcuc	1380	
aaaggaaugu cauacucuau gugcacagga aaguuuuaag uugugaagga aauagcagaa	1440	
acacaacaug gaacaauagu uaucagagug caauaugaag gggacggcuc uccaugcaag	1500	
auccuuuug agauaaugga uuuggaaaaa agacaugucu uaggucgcccug gauuacaguc	1560	
aacccaaauug ugacagaaaa agauagccca guacaacauag aagcagaacc uccauuugga	1620	
gacagcuaca ucaucauagg aguagagccg ggacaacuga agcuacaug guuuuagaaa	1680	
ggaaguucua ucggccaaau guuugagaca acaauggagg gggcgaagag aauggccauu	1740	
uuaggugaca cagccuggga uuuuggaucc uugggaggag uguuuuacauc uauaggaaag	1800	
gcucuccacc aagucuuugg agcaaucuau ggagcugccu ucaguggggu uucauggacu	1860	
augaaaaucc ucauaggagu cauuaucaca uggauaggaa ugaauucacg cagcaccuca	1920	

eolf-seq1.txt

cugucuguga cacuaguauu ggugggaauu gugacacugu auuugggagu cauggugcag 1980
gcc 1983

<210> 26
<211> 495
<212> PRT
<213> Artificial Sequence

<220>
<223> E VDV2

<400> 26

Met Arg Cys Ile Gly Met Ser Asn Arg Asp Phe Val Glu Gly Val Ser
1 5 10 15

Gly Gly Ser Trp Val Asp Ile Val Leu Glu His Gly Ser Cys Val Thr
20 25 30

Thr Met Ala Lys Asn Lys Pro Thr Leu Asp Phe Glu Leu Ile Lys Thr
35 40 45

Glu Ala Lys Gln Pro Ala Thr Leu Arg Lys Tyr Cys Ile Glu Ala Lys
50 55 60

Leu Thr Asn Thr Thr Glu Ser Arg Cys Pro Thr Gln Gly Glu Pro
65 70 75 80

Ser Leu Asn Glu Glu Gln Asp Lys Arg Phe Val Cys Lys His Ser Met
85 90 95

Val Asp Arg Gly Trp Gly Asn Gly Cys Gly Leu Phe Gly Lys Gly
100 105 110

Ile Val Thr Cys Ala Met Phe Arg Cys Lys Lys Asn Met Glu Gly Lys
115 120 125

Val Val Gln Pro Glu Asn Leu Glu Tyr Thr Ile Val Ile Thr Pro His
130 135 140

Ser Gly Glu Glu His Ala Val Gly Asn Asp Thr Gly Lys His Gly Lys
145 150 155 160

Glu Ile Lys Ile Thr Pro Gln Ser Ser Ile Thr Glu Ala Glu Leu Thr
165 170 175

Gly Tyr Gly Thr Val Thr Met Glu Cys Ser Pro Arg Thr Gly Leu Asp
180 185 190

Phe Asn Glu Met Val Leu Leu Gln Met Glu Asn Lys Ala Trp Leu Val
195 200 205

His Arg Gln Trp Phe Leu Asp Leu Pro Leu Pro Trp Leu Pro Gly Ala
Page 44

eolf-seq1.txt

210 215 220
Asp Thr Gln Glu Ser Asn Trp Ile Gln Lys Glu Thr Leu Val Thr Phe
225 230 235 240
Lys Asn Pro His Ala Lys Lys Gln Asp Val Val Val Leu Gly Ser Gln
245 250 255
Glu Gly Ala Met His Thr Ala Leu Thr Gly Ala Thr Glu Ile Gln Met
260 265 270
Ser Ser Gly Asn Leu Leu Phe Thr Gly His Leu Lys Cys Arg Leu Arg
275 280 285
Met Asp Lys Leu Gln Leu Lys Gly Met Ser Tyr Ser Met Cys Thr Gly
290 295 300
Lys Phe Lys Val Val Lys Glu Ile Ala Glu Thr Gln His Gly Thr Ile
305 310 315 320
Val Ile Arg Val Gln Tyr Glu Gly Asp Gly Ser Pro Cys Lys Ile Pro
325 330 335
Phe Glu Ile Met Asp Leu Glu Lys Arg His Val Leu Gly Arg Leu Ile
340 345 350
Thr Val Asn Pro Ile Val Thr Glu Lys Asp Ser Pro Val Asn Ile Glu
355 360 365
Ala Glu Pro Pro Phe Gly Asp Ser Tyr Ile Ile Ile Gly Val Glu Pro
370 375 380
Gly Gln Leu Lys Leu Asn Trp Phe Lys Lys Gly Ser Ser Ile Gly Gln
385 390 395 400
Met Phe Glu Thr Thr Met Arg Gly Ala Lys Arg Met Ala Ile Leu Gly
405 410 415
Asp Thr Ala Trp Asp Phe Gly Ser Leu Gly Gly Val Phe Thr Ser Ile
420 425 430
Gly Lys Ala Leu His Gln Val Phe Gly Ala Ile Tyr Gly Ala Ala Phe
435 440 445
Ser Gly Val Ser Trp Thr Met Lys Ile Leu Ile Gly Val Ile Ile Thr
450 455 460
Trp Ile Gly Met Asn Ser Arg Ser Thr Ser Leu Ser Val Thr Leu Val
465 470 475 480
Leu Val Gly Ile Val Thr Leu Tyr Leu Gly Val Met Val Gln Ala

<210> 27
<211> 75
<212> PRT
<213> Artificial Sequence

<220>
<223> M VDV2

<400> 27

Ser Val Ala Leu Val Pro His Val Arg Met Gly Leu Glu Thr Arg Thr
1 5 10 15

Glu Thr Trp Met Ser Ser Glu Gly Ala Trp Lys His Val Gln Arg Ile
20 25 30

Glu Thr Trp Ile Leu Arg His Pro Gly Phe Thr Met Met Ala Ala Ile
35 40 45

Leu Ala Tyr Thr Ile Gly Thr Thr His Phe Gln Arg Ala Leu Ile Phe
50 55 60

Ile Leu Leu Thr Ala Val Thr Pro Ser Met Thr
65 70 75