



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/00	A2	(11) International Publication Number: WO 99/52519 (43) International Publication Date: 21 October 1999 (21.10.99)
<p>(21) International Application Number: PCT/US99/08056</p> <p>(22) International Filing Date: 14 April 1999 (14.04.99)</p> <p>(30) Priority Data: 60/081,645 14 April 1998 (14.04.98) US</p> <p>(71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit Street, Boston, MA 02114 (US).</p> <p>(72) Inventors: TSAI, Guochuan; 3 Rollins Court, Cambridge, MA 02139 (US). COYLE, Joseph; 15 Wellesley Road, Belmont, MA 02178 (US).</p> <p>(74) Agent: ELLISON, Eldora, L.; Fish & Richardson P.C., 601 Thirteenth Street, N.W., Washington, DC 20005 (US).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>	
<p>(54) Title: METHODS FOR TREATING NEUROPSYCHIATRIC DISORDERS</p> <p>(57) Abstract</p> <p>The invention provides methods for treating neuropsychiatric disorders such as schizophrenia, Alzheimer's Disease, autism, depression, benign forgetfulness, childhood learning disorders, close head injury, and attention deficit disorder. The methods entail administering to a patient diagnosed as having a neuropsychiatric disorder a pharmaceutical composition containing (i) a therapeutically effective amount of D-alanine (or a modified form thereof), provided that the composition is substantially free of D-cycloserine, and/or (ii) D-serine (or a modified form thereof), and/or (iii) 105 to 500 mg of D-cycloserine (or a modified form thereof), and/or (iv) N-methylglycine (or a modified form thereof).</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

METHODS FOR TREATING NEUROPSYCHIATRIC DISORDERSBackground of the Invention

Schizophrenia, Alzheimer's Disease, autism, depression, benign forgetfulness, childhood learning disorders, close head injury, and attention deficit disorder are examples of neuropsychiatric disorders. Autism, for example, is a developmental mental disorder characterized by autistic behavior, social failure, and language delay. Alzheimer's Disease is a form of dementia that typically involves progressive mental deterioration, manifested by memory loss, confusion, and disorientation. Alzheimer's Disease typically is treated by acetylcholine esterase inhibitors such as tacrine hydrochloride or donepezil. Attention Deficit Disorder is a disorder that is most prevalent in children and is associated with increased motor activity and a decreased attention span. Attention Deficit Disorder commonly is treated by administration of psychostimulants such as Ritalin or Dexedrin. Depression is a clinical syndrome that includes a persistent sad mood or loss of interest in activities, which persists for at least two weeks in the absence of treatment. Conventional therapeutics include serotonin uptake inhibitors (e.g., PROZAC™), monoamine oxidase inhibitors, and tricyclic antidepressants.

The term schizophrenia represents a group of neuropsychiatric disorders characterized by dysfunctions of the thinking process, such as delusions, hallucinations, and extensive withdrawal of the patient's interests from other people. Approximately one percent of the worldwide population is afflicted with schizophrenia, and this disorder is accompanied by high morbidity and mortality rates.

- 2 -

Conventional antipsychotic drugs, which act on the dopamine D₂ receptor, can be used to treat the positive symptoms of schizophrenia, such as delusion and hallucination. In general, conventional antipsychotic drugs and the new atypical antipsychotic drugs, which act on the dopamine D₂ and 5HT₂ serotonin receptor, are limited in their ability to treat cognitive deficits and negative symptoms such as affect blunting (i.e., lack of facial expressions), anergia, and social withdrawal.

10 Summary of the Invention

The invention derives from the discovery that neuropsychiatric disorders characterized by a deficit in neurotransmission via the NMDA receptor can be alleviated by a compound that acts as an agonist of the glycine site on the NMDA receptor or an inhibitor of glycine uptake. The compound is either a partial agonist such as D-cycloserine, which can be used at a dosage of 105-500 mg, or a full agonist (e.g., D-serine or D-alanine) that is selective for the NMDA receptor (compared to the inhibitory glycine receptor and other receptors), or a glycine uptake inhibitor (e.g., N-methylglycine). The invention therefore provides new methods for treating neuropsychiatric disorders in patients (i.e., humans). Examples of disorders that can be treated by the methods of the invention include schizophrenia, Alzheimer's Disease, autism, depression, benign forgetfulness, childhood learning disorders, close head injury, and attention deficit disorder. The methods entail administering to a patient diagnosed as suffering from such a neuropsychiatric disorder a pharmaceutical composition that contains a therapeutically effective amount of an agonist of the glycine site of the NMDA receptor or a glycine uptake inhibitor, which agonist is relatively selective for (a) the glycine site of the NMDA

- 3 -

receptor, compared with (b) the inhibitory glycine receptor and other receptors. The pharmaceutical composition may include, for example, (i) a therapeutically effective amount of D-alanine (wherein
5 the pharmaceutical composition is substantially free of D-cycloserine) and/or (ii) a therapeutically effective amount of D-serine, and/or (iii) D-cycloserine in an amount of 105-500 mg, and/or (iv) a therapeutically effective amount of N-methylglycine.

10 In variations of the methods described herein, D-serine, D-alanine, D-cycloserine, and/or N-methylglycine can be substituted with a salt, ester, or alkylated form of the amino acid, or a precursor of the amino acid that is converted (e.g., metabolized) into the amino acid *in*
15 *vivo* (e.g., D-phosphoserine, L-phosphoserine, or L-phosphoserine, N,N,N-trimethylglycine (betaine), or N,N-dimethylglycine).

Typically, a dosage of 100 μ g to 100 g (e.g., 1 mg to 100 g; 1 mg to 100 mg; 10 mg to 100 g; 10 mg to 10 g;
20 or 10 to 500 mg) is suitable for D-alanine, D-serine, and N-methylglycine. D-cycloserine is administered at a dosage of 105 to 500 mg. When the patient is treated with both D-serine and D-alanine, D-serine and D-alanine can be administered to the patient simultaneously or
25 sequentially, e.g., by formulating the D-serine and D-alanine as a single pharmaceutical composition or as two or more pharmaceutical compositions. Likewise, the patient can be treated with both D-serine and D-cycloserine, or D-serine and N-methylglycine, or D-
30 alanine and N-methylglycine, or D-cycloserine and N-methylglycine simultaneously or sequentially. In one, but not the only, suitable method of treatment, the pharmaceutical composition is administered to the patient at least once daily for at least one week. If desired,
35 the pharmaceutical composition can be administered to the

- 4 -

patient in more than one dose per day (e.g., 2, 3, or 4 doses). Generally, the patient is treated for at least one week; typically, the patient is treated for at least several weeks (e.g., at least 4, 6, or 8 weeks) or months (e.g., at least 4, 8, or 12 months). If necessary, the treatment can continue indefinitely to keep the patient's symptoms under control throughout his or her life.

If desired, a pharmaceutical composition containing D-alanine (substantially free of D-cycloserine), D-serine, D-cycloserine and/or N-methylglycine (or a modified version thereof, as described herein) can be administered to a patient suffering from schizophrenia along with, or in sequence with, an art-known drug for treating schizophrenia (e.g., olanzapine, clozapine, haloperidol, and the like). Similarly, D-alanine (typically substantially free of D-cycloserine), D-serine, D-cycloserine and/or N-methylglycine (or a modified version thereof, as described herein) can be used in combination with, or in sequence with, other art-known antipsychotics (e.g., "typical," "atypical," and depot antipsychotics for treating schizophrenia and other psychotic conditions), antidepressants (for treating depression), psychostimulants (for treating attention deficit disorder, depression, or learning disorders), or Alzheimer's disease therapeutics (for treating Alzheimer's disease). Such pharmaceutical compositions are included within the invention. In general, the antipsychotic, antidepressant, psychostimulant, or Alzheimer's disease therapeutic typically is administered at a dosage of 0.25-5000 mg/d (e.g., 5-1000 mg/d). "Typical" antipsychotics are conventional antipsychotics such as phenothiazine, butyrophenones, thioxanthenes, dibenzoxazepines, dihydroindolones, and diphenylbutylpiperidines. "Atypical" antipsychotics are

- 5 -

a new generation of antipsychotics which generally act on the dopamine D₂ and 5HT₂ serotonin receptor and have high levels of efficacy and a benign extrapyramidal symptom side effect profile. Examples of typical antipsychotics (and examples of suitable daily (d) dosages) include Chlorpromazine (5-2000 mg/d, e.g., 30-800 mg/d), Thioridazine (5-2000 mg/d, e.g., 20-800 mg/d), Mesoridazine (1-1000 mg/d, e.g., 30-400 mg/d), Fluphenazine (0.5-200 mg/d, e.g., 1-40 mg/d), Perphenazine (0.5-300 mg/d, e.g., 10-65 mg/d), Trifluoperazine (0.5-200 mg/d, e.g., 2-40 mg/d), Thiothixene (1-200 mg/d, e.g., 6-60 mg/d), Haloperidol (0.25-500 mg/d, e.g., 1-100 mg/d), Loxapine (1-1000 mg/d e.g., 20-250 mg/d), Molindone (1-1000 mg/d, e.g., 15-225 mg/d), Acetophenazine (10-2000 mg/d, e.g., 30-500 mg/d), Chlorprothixene (5-2000 mg/d, e.g., 30-500 mg/d), Droperidol (0.25-500 mg/d, e.g., 1-100 mg/d), Pimozide (0.25-500 mg/d, e.g., 1-100 mg/d). Examples of atypical antipsychotics (and examples of suitable daily dosages) include Clozapine (5-2000 mg/d, e.g., 12-900 mg/d), Risperidone (0.25-500 mg/d, e.g., 2-16 mg/d), Olanzapine (1-100 mg/d, e.g., 5-10 mg/d), and Quetiapine (1-2000 mg/d, e.g., 50-750 mg/d). Depot antipsychotics also can be used, e.g., Haloperidol decanoate (10-1000 mg/month, e.g., 100-450 mg/month), Fluphenazine decanoate (5-1000 mg/month, e.g., 25-150 mg/month), and Fluphenazine enanthate (5-1000 mg/month, e.g., 25-200 mg/month). Additional antipsychotics include Butaperazine (0.5-500 mg/d, e.g., 1-200 mg/d), Carphenazine, (0.5-3000 mg/d, e.g., 1-1000 mg/d), Remoxipride (0.5-5000 mg/d, e.g., 1-2000 mg/d), Piperacetazine (0.5-500 mg/d, e.g., 1-2000 mg/d), Sulpiride (0.5-5000 mg/d, e.g., 1-2000 mg/d), and Ziprasidone (0.5-500 mg/d, e.g., 1-200 mg/d). Examples of antidepressants that can be used include Amitriptyline

- 6 -

(5-1000 mg/d, e.g., 50-300 mg/d), Amoxapine (5-1000 mg/d, e.g., 50-600 mg/d), Bupropion (5-1000 mg/d, e.g., 200-450 mg/d), Bupropion SR (5-1000 mg/d, e.g., 150-400 mg/d), Clomipramine (5-1000 mg/d, e.g., 25-250 mg/d),

5 Desipramine (5-1000 mg/d, e.g., 100-300 mg/d), Doxepin (5-1000 mg/d, e.g., 75-300 mg/d), Fluoxetine (1-200 mg/d, e.g., 20-80 mg/d), Fluvoxamine (5-1000 mg/d, e.g., 50-300 mg/d), Imipramine (5-1000 mg/d, e.g., 75-300 mg/d), Maprotiline (5-1000, e.g., 75-225 mg/d), Mirtazapine (1-

10 200 mg/d, e.g., 15-45 mg/d), Nefazodone (5-1000 mg/d, e.g., 200-600 mg/d), Nortriptyline (5-1000 mg/d, e.g., 75-150 mg/d), Paroxetine (1-200 mg/d, e.g., 10-60 mg/d), Phenelzine (1-500 mg/d, e.g., 5-90 mg/d), Protriptyline (1-200 mg/d, e.g., 15-60 mg/d), Sertraline (5-1000 mg/d,

15 e.g., 50-200 mg/d), Tranylcypramine (1-200 mg/d, e.g., 30-60 mg/d), Trazodone (5-1000 mg/d, e.g., 150-600 mg/d), Trimipramine (5-1000 mg/d, e.g., 5-300 mg/d), Venlafaxine (5-1000 mg/d, e.g., 75-375 mg/d), and Venlafaxine XR (5-1000 mg/d, e.g., 75-225 mg/d). Psychostimulants that are

20 particularly useful for treating attention deficit disorder include Dextroamphetamine (0.5-200 mg/d, e.g., 5-40 mg/d), Methamphetamine (0.5-200 mg/d, e.g., 5-25 mg/d), Methylphenidate (0.5-200 mg/d, e.g., 10-40 mg/d), and Pemoline (5-500 mg/d, e.g., 37.5-112.5 mg/d).

25 Examples of Alzheimer's disease therapeutics that can be used in the invention include Donepezil (0.5-200 mg/d, e.g., 1-100 mg/d) and Tacrine (0.5-1000 mg/d, e.g., 10-500 mg/d). Thus, the invention also provides pharmaceutical compositions that contain D-alanine

30 (typically substantially free of D-cycloserine), D-serine, D-cycloserine and/or N-methylglycine (or a modified version thereof, as described herein) along with an antipsychotic, antidepressant, psychostimulant, or Alzheimer's disease therapeutic.

- 7 -

If desired, one can measure negative and/or positive and/or cognitive symptom(s) of schizophrenia before and after treatment of the patient. A reduction in such a symptom indicates that the patient's condition
5 has improved. Improvement in the symptoms of schizophrenia can be assessed using the Scales for the Assessment of Negative Symptoms (SANS) or Positive and Negative Syndrome Scale (PANSS) (see, e.g., Andreasen, 1983, *Scales for the Assessment of Negative Symptoms*
10 (SANS), Iowa City, Iowa and Kay et al., 1987, *Schizophrenia Bulletin* 13:261-276). Likewise, one can measure improvement of other neuropsychiatric disorders in patients who have been treated by the methods of the invention.

15 As used herein, the term "neuropsychiatric disorder" refers to a disease having a pathophysiological component of attenuated NMDA receptor-mediated neurotransmission. Examples of such disorders include schizophrenia, Alzheimer's disease, autism, depression,
20 benign forgetfulness, childhood learning disorders, close head injury, and attention deficit disorder.

As used herein, the term "schizophrenia" refers to a psychiatric disorder that includes at least two of the following: delusions, hallucinations, disorganized
25 speech, grossly disorganized or catatonic behavior, or negative symptoms. Patients can be diagnosed as schizophrenic using the DSM-IV criteria (APA, 1994, *Diagnostic and Statistical Manual of Mental Disorders* (Fourth Edition), Washington, DC).

30 The term "Alzheimer's Disease" refers to a progressive mental deterioration manifested by memory loss, confusion and disorientation beginning in late middle life and typically resulting in death in five to ten years. Pathologically, Alzheimer's Disease can be
35 characterized by thickening, conglutination, and

- 8 -

distortion of the intracellular neurofibrils,
neurofibrillary tangles and senile plaques composed of
granular or filamentous argentophilic masses with an
amyloid core. Methods for diagnosing Alzheimer's Disease
5 are known in the art. For example, the National
Institute of Neurological and Communicative Disorders and
Stroke-Alzheimer's Disease-and the Alzheimer's Disease
and Related Disorders Association (NINCDS-ADRDA) criteria
can be used to diagnose Alzheimer's Disease (McKhann et
10 al., 1984, Neurology 34:939-944). The patient's
cognitive function can be assessed by the Alzheimer's
Disease Assessment Scale-cognitive subscale (ADAS-cog;
Rosen et al., 1984, Am. J. Psychiatry 141:1356-1364).

As used herein, the term "autism" refers to a
15 state of mental introversion characterized by morbid
self-absorption, social failure, language delay, and
stereotyped behavior. Patients can be diagnosed as
suffering from autism by using the DSM-IV criteria.

As used herein, the term "depression" refers to a
20 clinical syndrome that includes a persistent sad mood or
loss of interest in activities, which lasts for at least
two weeks in the absence of treatment. The DSM-IV
criteria can be used to diagnose patients as suffering
from depression.

25 The term "benign forgetfulness," as used herein,
refers to a mild tendency to be unable to retrieve or
recall information that was once registered, learned, and
stored in memory (e.g., an inability to remember where
one placed one's keys or parked one's car). Benign
30 forgetfulness typically affects individuals after 40
years of age and can be recognized by standard assessment
instruments such as the Wechsler Memory Scale (Russell,
1975, J. Consult Clin. Psychol. 43:800-809).

As used herein, the term "childhood learning
35 disorders" refers to an impaired ability to learn, as

- 9 -

experienced by certain children. Such learning disorders can be diagnosed by using the DSM-IV criteria.

The term "close head injury," as used herein, refers to a clinical condition after head injury or
5 trauma which condition can be characterized by cognitive and memory impairment. Such a condition can be diagnosed as "amnesic disorder due to a general medical condition" according to DSM-IV.

The term "attention deficit disorder," as used
10 herein, refers to a disorder that is most commonly exhibited by children and which can be characterized by increased motor activity and a decreased attention span. The DSM-IV criteria can be used to diagnose attention deficit disorder.

15 The terms "D-serine" and "D-alanine" refer to the D isomers of the amino acids serine and alanine, respectively. As D isomers, rather than L isomers, these amino acids are not naturally found in proteins.

"Negative" symptoms of schizophrenia include
20 affect blunting, anergia, alogia and social withdrawal, which can be measured using SANS (the Scales for the Assessment of Negative Symptoms; see Andreasen, 1983, *Scales for the Assessment of Negative Symptoms (SANS)*, Iowa City, Iowa).

25 "Positive" symptoms of schizophrenia include delusion and hallucination, which can be measured using PANSS (the Positive and Negative Syndrome Scale; see Kay et al., 1987, *Schizophrenia Bulletin* 13:261-276).

"Cognitive" symptoms of schizophrenia include
30 impairment in obtaining, organizing, and using intellectual knowledge which can be measured by the Positive and Negative Syndrome Scale-cognitive subscale (PANSS-cognitive subscale) (Lindenmayer et al., 1994, *J. Nerv. Ment. Dis.* 182:631-638) or with cognitive tasks
35 such as the Wisconsin Card Sorting Test.

- 10 -

A "full" agonist of the NMDA receptor is a compound that produces a maximal response at full receptor occupancy.

A "partial" agonist of the NMDA receptor is a
5 compound that produces a lower maximal response at full receptor occupancy than do full agonists.

A "glycine uptake inhibitor of the NMDA receptor" is a compound that inhibits the re-uptake of glycine and increases the availability of glycine for the NMDA
10 receptor (e.g., N-methylglycine).

The invention offers several advantages over many art-known methods for treating neuropsychiatric disorders. For example, unlike many conventional antipsychotic therapeutics, D-serine, D-alanine, and N-
15 methylglycine can produce a desirable reduction in the positive, negative, and cognitive symptoms of schizophrenia. As shown by the examples set forth below, clinically significant improvement can be achieved even with patients who are poorly responsive to treatment by
20 conventional antipsychotics. In addition, no significant side effects were detected after treatment of schizophrenia patients with D-serine, D-alanine, or N-methylglycine. In contrast, conventional antipsychotics typically lead to tardive dyskinesia (irreversible,
25 involuntary movement disorder), extrapyramidal symptoms, and akathisia symptoms.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

30

Detailed Description

The invention provides methods for treating a patient diagnosed as suffering from a neuropsychiatric disorder having a deficit in neurotransmission via the NMDA receptor (e.g., schizophrenia, Alzheimer's Disease,

- 11 -

autism, depression, benign forgetfulness, childhood learning disorders, close head injury, and attention deficit disorder). As described above, a variety of methods for diagnosing these disorders are known to those
5 of skill in the art of clinical psychiatry, and any conventional diagnostic method can be used in conjunction with the invention.

The treatment method of the invention entails administering to a patient diagnosed as having a
10 neuropsychiatric disorder a pharmaceutical composition containing a therapeutically effective amount of (i) an agonist of the glycine site of the NMDA receptor, which agonist is relatively selective for (a) the glycine site of the NMDA receptor, compared with (b) an inhibitory
15 glycine receptor or any other receptor, or (ii) a glycine uptake inhibitor. For example, suitable pharmaceutical compositions may include (i) D-alanine substantially free of D-cycloserine and/or (ii) D-serine and/or (iii) N-methylglycine. D-serine and D-alanine are commercially
20 available (e.g., from Spectrum Quality Products, Inc., Gardena, CA). Where D-alanine is used, the pharmaceutical composition is "substantially free" of D-cycloserine, meaning that the composition lacks D-cycloserine, or D-cycloserine is not included at a level
25 sufficient to have a statistically significant effect upon the efficacy of the pharmaceutical composition, as determined by any method (e.g., by comparing PANSS and/or SANS scores before and after treatment of the patient). In general, this means that D-cycloserine is absent from
30 the pharmaceutical composition or present in an amount such that the patient receives less than 0.02 mg/day.

Treatment includes administering a therapeutically effective amount of a composition containing D-alanine (substantially free of D-cycloserine) and/or D-serine
35 and/or N-methylglycine to a patient in need of such

- 12 -

treatment, thereby treating the neuropsychiatric disorder. Such compositions typically contain from about 0.1 to 90% by weight (such as 1 to 20% or 1 to 10%) of D-alanine, D-serine, or N-methylglycine in a
5 pharmaceutically acceptable carrier. Regardless of the concentration of D-serine or D-alanine in the pharmaceutical composition, D-serine and/or D-alanine and/or N-methylglycine is administered to the patient at a dosage of 10 mg to 100 g. More typically, D-serine
10 and/or D-alanine and/or N-methylglycine is administered at a dosage of 100 mg to 10 g. Generally, treatment continues for at least several weeks to several years or life-long as needed.

In an alternative method for treating a
15 neuropsychiatric disorder in a patient, a pharmaceutical composition containing D-cycloserine in an amount equivalent to a dosage of 105 to 500 mg/day is administered to a patient in need of such treatment. For example, the dosage can be in an amount of 125 to 400 mg,
20 such as 150 to 300 mg (e.g., 175 mg, 200 mg, 225 mg, or 250 mg). D-cycloserine (D-4-amino-3-isoxazolidinone) is commercially available from Eli Lilly, Inc. (Indianapolis, IN). Generally, treatment continues for at least one week and can continue for several years or
25 life-long as needed to control the patient's symptoms.

In all of the methods of the invention, D-alanine, D-serine, and/or D-cycloserine and/or N-methylglycine can be substituted with a modified version of the amino acid, such as a salt, ester, alkylated form, or a precursor of
30 the amino acid. For example, the amino acid can be in the form of a sodium salt, potassium salt, calcium salt, magnesium salt, zinc salt, or ammonium salt. Such salt forms of D-serine, D-alanine, N-methylglycine and D-cycloserine can be made in accordance with conventional
35 methods (see, e.g., *Organic Chemistry*, pgs. 822-823,

- 13 -

Morrison and Boyd, ed., Fifth Edition, Allyn and Bacon, Inc., Newton, MA). Other modified forms of D-serine, D-alanine, N-methylglycine and D-cycloserine also can be used in the methods of the invention. For example, the
5 carboxy group of the amino acid can be converted to an ester group by reaction with an alcohol in accordance with standard esterification methods (*Id.* at 841-843). For example, alcohols having 1-20 carbon atoms can be used to produce an ester of D-serine, D-alanine, N-
10 methylglycine or D-cycloserine for use in the invention (e.g., methyl-, ethyl-, propyl-, isopropyl-, butyl-, isobutyl-, sec-butyl-, tert-butyl-, pentyl-, isopentyl-, tert-pentyl-, hexyl-, heptyl-, octyl-, decyl-, dodecyl-, tetradecyl-, hexadecyl-, octadecyl-, and phenyl-alcohols
15 can be used). In another variation, the amino group of the amino acid can be alkylated, using conventional methods, to produce a secondary or tertiary amino group by ammonolysis of halides or reductive amination (*Id.* at 939-948). For example, an alkyl group having 1-20
20 carbon atoms can be added to the amino acid to produce an alkylated amino acid (e.g., methyl-, ethyl-, propyl-, isopropyl-, butyl-, isobutyl-, sec-butyl-, tert-butyl-, pentyl-, isopentyl-, tert-pentyl-, hexyl-, heptyl-, octyl-, decyl-, dodecyl-, tetradecyl-, hexadecyl-, octadecyl-
25 and phenyl-groups can be added to the amino acid). D-phosphoserine and L-phosphoserine are examples of precursors of D-serine, and are commercially available (e.g., from Sigma Chemical, St. Louis, MO). N,N,N-trimethylglycine (betaine) and N,N-dimethylglycine are
30 examples of precursors of N-methylglycine.

In all of the methods of the invention, appropriate dosages of D-alanine, D-serine, D-cycloserine, or N-methylglycine (or modified versions thereof) can readily be determined by those of ordinary
35 skill in the art of medicine by monitoring the patient

- 14 -

for signs of disease amelioration or inhibition, and increasing or decreasing the dosage and/or frequency of treatment as desired.

The pharmaceutical compositions can be administered to the patient by any, or a combination, of several routes, such as oral, intravenous, trans-mucosal (e.g., nasal, vaginal, etc.), pulmonary, transdermal, ocular, buccal, sublingual, intraperitoneal, intrathecal, intramuscular, or long term depot preparation. Solid compositions for oral administration can contain suitable carriers or excipients, such as corn starch, gelatin, lactose, acacia, sucrose, microcrystalline cellulose, kaolin, mannitol, dicalcium phosphate, calcium carbonate, sodium chloride, lipids, alginic acid, or ingredients for controlled slow release. Disintegrators that can be used include, without limitation, micro-crystalline cellulose, corn starch, sodium starch glycolate and alginic acid. Tablet binders that may be used include, without limitation, acacia, methylcellulose, sodium carboxymethylcellulose, polyvinylpyrrolidone (Povidone), hydroxypropyl methylcellulose, sucrose, starch, and ethylcellulose.

Liquid compositions for oral administration prepared in water or other aqueous vehicles can include solutions, emulsions, syrups, and elixirs containing, together with the active compound(s), wetting agents, sweeteners, coloring agents, and flavoring agents. Various liquid and powder compositions can be prepared by conventional methods for inhalation into the lungs of the patient to be treated.

Injectable compositions may contain various carriers such as vegetable oils, dimethylacetamide, dimethylformamide, ethyl lactate, ethyl carbonate, isopropyl myristate, ethanol, polyols (glycerol, propylene glycol, liquid polyethylene glycol, and the

- 15 -

like). For intravenous injections, the compounds may be administered by the drip method, whereby a pharmaceutical composition containing the active compound(s) and a physiologically acceptable excipient is infused.

5 Physiologically acceptable excipients may include, for example, 5% dextrose, 0.9% saline, Ringer's solution or other suitable excipients. For intramuscular preparations, a sterile composition of a suitable soluble salt form of the compound can be dissolved and

10 administered in a pharmaceutical excipient such as Water-for-Injection, 0.9% saline, or 5% glucose solution, or depot forms of the compounds (e.g., decanoate, palmitate, undecylenic, enanthate) can be dissolved in sesame oil. Alternatively, the pharmaceutical composition can be

15 formulated as a chewing gum, lollipop, or the like.

EXAMPLES

The following examples demonstrate that D-alanine, D-serine, and N-methylglycine each can be used to treat a neuropsychiatric disorder in patients.

20 Patients

This study employed 37 patients who were diagnosed as having schizophrenia. All patients fulfilled the DSM-IV diagnosis of schizophrenia (APA, 1994, Diagnostic and Statistical Manual of Mental

25 Disorders, Fourth Edition, Washington, DC). All of the patients also fulfilled the criteria of primary deficit syndrome, with a SANS score of more than 40 (Kirkpatrick et al., 1989, Psychiatry Research 30:119-123; Andreasen, 1983, Scales for the Assessment of Negative Symptoms

30 (SANS), Iowa City, Iowa). All of the patients were poorly responsive to treatment by other antipsychotic drugs, and had been kept on a stable dose of an

- 16 -

antipsychotic drug for at least 3 months prior to enrollment in this study.

Assessments

Several scales were used to assess the severity of the disorder in each patient. At the beginning of the study (i.e., the baseline), the PANSS, SANS, and Global Assessment Scales (CGI) were used. Each scale also was completed at the end of each 2-week period throughout the study. These assessments were performed by a psychiatrist who was blind to the treatment assignment. The Wisconsin Card Sort Test was used to provide a cognitive rating of the patients; in general, schizophrenic patients perform poorly on this test. The Wisconsin Card Sort Test was administered only at the initiation of the study and at the end of the 6-week study. To measure side effects, the Simpson-Angus Scale was used to measure extrapyramidal symptoms (EPS; Simpson et al., 1970, *Acta Psychiatrica Scandinavia Suppl.* 212:11-19). The Abnormal Involuntary Movement Scale (AIMS) was used to measure dyskinesia (Simpson et al., 1970, *Acta Psychiatrica Scandinavia Suppl.* 212:11-19). The Barnes Scale was used to measure akathisia (Barnes, 1989, *Brit. J. Psychiatry* 154:672-676). The side effects of D-serine, D-alanine, and N-methylglycine treatments were assessed biweekly according to the UKU side effects rating scale (*Scandinavian Society of Psychopharmacology Committee of Clinical Investigation: The UKU side effect rating scale: scale for the registration of unwanted effects of psychotropics.* *Acta. Psychiatr. Scand.* 1987; *Suppl.* 334:81-94).

Treatment and Results

Using double-blind conditions, the patients were randomly assigned to receive placebo (fruit juice), D-

- 17 -

serine (30 mg/kg/day), D-alanine (60-100 mg/kg/day), or N-methylglycine (30 mg/kg/day) once a day by mouth for a period of 6 weeks. As indicated by the results shown in Table 1, treatment with D-serine, D-alanine, or N-
5 methylglycine improved the schizophrenic symptoms and cognitive deficit of the patients. More specifically, treatment with D-serine resulted in a 21% reduction of the negative symptoms (on the SANS scale), and it resulted in a 17% reduction of the positive symptoms (on
10 the PANSS-positive subscale). Treatment with D-alanine resulted in an 11% reduction of the negative symptoms and a 12% reduction of the positive symptoms. Treatment with N-methylglycine resulted in a 20% reduction of the negative symptoms and a 15% reduction of the positive
15 symptoms. These reductions in the negative and positive symptoms represented clinically significant improvement. Treatment with each of D-serine, D-alanine, and N-methylglycine also improved cognition, as measured using the PANSS-cognitive subscale and the Wisconsin Card Sort
20 Test. These results indicate that D-serine, D-alanine, and N-methylglycine are effective in treating schizophrenia even in patients who are poorly responsive to treatment by conventional antipsychotic drugs.

Using the UKU scale for rating side effects, no
25 side effects were noted after treatment with D-serine, D-alanine, or N-methylglycine. In addition, there was no newly emergent tardive dyskinesia or worsening of extrapyramidal or akathisia symptoms. Thus, D-serine, D-alanine, and N-methylglycine offer an advantage over many
30 conventional drugs for treating schizophrenia in that they do not cause significant side effects.

TABLE 1: Effects of D-serine, D-alanine, and N-methylglycine Treatment on Schizophrenia Patients

	D-serine	D-alanine	N-methylglycine	Placebo
<i>Clinical Symptoms</i>				
Negative Symptoms	-21%*	-12%*	-20%*	-1%
5 Positive Symptoms	-17%*	-11%*	-15%*	3%
CGI	4.8->2.6*	3.9->2.8*	4.2->2.7*	4.5->4.0
<i>Cognition</i>				
Cognitive symptoms	-12%*	-11%*	-12%*	1%
WCST	+0.9 (category)*	+0.5*	+0.7*	-0.5
10 <i>Side Effects</i>				
EPS	1.4-->1.7	3.1-->3.1	2.1-->2.1	3.3-->3.4
AIMS	0.3-->0.3	0.5-->0.1	0.4-->0.3	0.5-->0.9
Barnes	0.4-->0.8	0.4-->0.6	0.5-->0.6	0.9-->0.9

* Clinically significant improvement

15

Other Embodiments

It is to be understood that, while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

20

What is claimed is:

- 19 -

1. A method for treating a neuropsychiatric disorder characterized by attenuated NMDA neurotransmission in a patient, the method comprising administering to a patient diagnosed as suffering from
5 the neuropsychiatric disorder a pharmaceutical composition comprising a therapeutically effective amount of an agonist of the glycine site of an NMDA receptor or a glycine uptake inhibitor, wherein:

the agonist is selected from the group consisting
10 of D-alanine, a salt of D-alanine, an ester of D-alanine, alkylated D-alanine, a precursor of D-alanine, D-serine, a salt of D-serine, an ester of D-serine, alkylated D-serine, a precursor of D-serine, D-cycloserine, a salt of D-cycloserine, an ester of D-cycloserine, a precursor of
15 D-cycloserine, and alkylated D-cycloserine;

the pharmaceutical composition is substantially free of D-cycloserine when the agonist is D-alanine, a salt of D-alanine, an ester of D-alanine, an alkylated D-alanine, or a precursor of D-alanine; and

20 when the agonist is D-cycloserine, a salt of D-cycloserine, an ester of D-cycloserine, a precursor of D-cycloserine, or alkylated D-cycloserine, the pharmaceutical composition comprises an amount of the agonist equivalent to 105-500 mg of D-cycloserine.

25 2. The method of claim 1, wherein the neuropsychiatric disorder is schizophrenia.

3. The method of claim 1, wherein the neuropsychiatric disorder is Alzheimer's disease.

4. The method of claim 1, wherein the
30 neuropsychiatric disorder is autism.

- 20 -

5. The method of claim 1, wherein the neuropsychiatric disorder is depression.

6. The method of claim 1, wherein the neuropsychiatric disorder is benign forgetfulness.

5 7. The method of claim 1, wherein the neuropsychiatric disorder is a childhood learning disorder.

8. The method of claim 1, wherein the neuropsychiatric disorder is attention deficit disorder.

10 9. The method of claim 1, wherein the neuropsychiatric disorder is close head injury.

10. The method of claim 1, wherein the agonist is selected from the group consisting of D-alanine, a salt of D-alanine, an ester of D-alanine, alkylated D-alanine,
15 and a precursor of D-alanine.

11. The method of claim 10, wherein the D-alanine, salt of D-alanine, ester of D-alanine, alkylated D-alanine, or precursor of D-alanine is administered at a dosage equivalent to 10 mg to 100 g of D-alanine.

20 12. The method of claim 10, wherein the agonist is a D-alanine salt selected from the group consisting of a sodium salt, a potassium salt, a calcium salt, a magnesium salt, a zinc salt, and an ammonium salt of D-alanine.

25 13. The method of claim 10, wherein the agonist is an ester of D-alanine having an ester group with 1-20 carbon atoms.

- 21 -

14. The method of claim 10, wherein the agonist is an alkylated D-alanine having an alkyl group with 1-20 carbon atoms.

15. The method of claim 10, wherein the pharmaceutical composition further comprises D-serine.

16. The method of claim 1, wherein the agonist is selected from the group consisting of D-serine, a salt of D-serine, an ester of D-serine, alkylated D-serine, and a precursor of D-serine.

17. The method of claim 16, wherein the D-serine, salt of D-serine, ester of D-serine, precursor of D-serine, or alkylated D-serine is administered at a dosage equivalent to 10 mg to 100 g of D-serine.

18. The method of claim 16, wherein the agonist is a D-serine salt selected from the group consisting of a sodium salt, a potassium salt, a calcium salt, a magnesium salt, a zinc salt, and an ammonium salt of D-serine.

19. The method of claim 16, wherein the agonist is an ester of D-serine having an ester group with 1-20 carbon atoms.

20. The method of claim 16, wherein the agonist is an alkylated D-serine having an alkyl group with 1-20 carbon atoms.

21. The method of claim 1, wherein the agonist is selected from the group consisting of D-cycloserine, a salt of D-cycloserine, an ester of D-cycloserine, a precursor of D-cycloserine, and an alkylated D-cycloserine.

- 22 -

22. The method of claim 21, wherein the D-cycloserine, salt of D-cycloserine, ester of D-cycloserine, alkylated D-cycloserine, or precursor of D-cycloserine is administered in a dose equivalent to 125-
5 400 mg of D-cycloserine.

23. The method of claim 22, wherein the D-cycloserine, D-cycloserine salt, ester of D-cycloserine, alkylated D-cycloserine, or precursor of D-cycloserine is administered in a dose equivalent to 150-300 mg of D-
10 cycloserine.

24. The method of claim 21, wherein the pharmaceutical composition comprises a salt of D-cycloserine selected from the group consisting of a sodium salt, a potassium salt, a calcium salt, a
15 magnesium salt, a zinc salt, and an ammonium salt of D-cycloserine.

25. The method of claim 21, wherein the pharmaceutical composition comprises an ester of D-cycloserine having an ester group with 1-20 carbon atoms.

20 26. The method of claim 21, wherein the pharmaceutical composition comprises an alkylated D-cycloserine having an alkyl group with 1-20 carbon atoms.

27. The method of claim 21, wherein the pharmaceutical composition comprises a precursor of D-
25 cycloserine.

28. The method of claim 1, wherein the glycine uptake inhibitor is selected from the group consisting of N-methylglycine, a salt of N-methylglycine, an ester of N-methylglycine, and a precursor of N-methylglycine.

- 23 -

29. The method of claim 28, wherein the N-methylglycine, salt of N-methylglycine, ester of N-methylglycine, alkylated N-methylglycine, or precursor of N-methylglycine is administered at a dosage equivalent to
5 10 mg to 100 g of N-methylglycine.

30. The method of claim 28, wherein the glycine uptake inhibitor is an ester of N-methylglycine having an ester group with 1-20 carbon atoms.

31. The method of claim 28, wherein glycine
10 uptake inhibitor is an alkylated N-methylglycine having an alkyl group with 1-20 carbon atoms.

32. The method of claim 28, wherein the precursor is selected from the group consisting of N,N,N-trimethylglycine and N,N-dimethylglycine.

15 33. The method of claim 1, wherein the pharmaceutical composition is administered to the patient at least once daily for at least one week.

34. The method of claim 1, further comprising administering to the patient at least one therapeutic
20 selected from the group consisting of antipsychotics, antidepressants, psychostimulants, and Alzheimer's disease therapeutics.

35. A pharmaceutical composition comprising (i) at least one agonist of the glycine site of an NMDA
25 receptor or at least one glycine uptake inhibitor and (ii) at least one therapeutic agent selected from the group consisting of antipsychotics, antidepressants, psychostimulants, and Alzheimer's disease therapeutics, wherein:

- 24 -

the agonist is selected from the group consisting of D-alanine, a salt of D-alanine, an ester of D-alanine, alkylated D-alanine, a precursor of D-alanine, D-serine, a salt of D-serine, an ester of D-serine, alkylated D-serine, a precursor of D-serine, D-cycloserine, a salt of D-cycloserine, an ester of D-cycloserine, a precursor of D-cycloserine, and alkylated D-cycloserine; and

the pharmaceutical composition is substantially free of D-cycloserine when the agonist is D-alanine, a salt of D-alanine, an ester of D-alanine, an alkylated D-alanine, or a precursor of D-alanine; and

when the agonist is D-cycloserine, a salt of D-cycloserine, an ester of D-cycloserine, a precursor of D-cycloserine, or alkylated D-cycloserine, the pharmaceutical composition comprises an amount of the agonist equivalent to 105-500 mg of D-cycloserine.

36. The pharmaceutical composition of claim 35, wherein the glycine uptake inhibitor is selected from the group consisting of N-methylglycine, a salt of N-methylglycine, an ester of N-methylglycine, alkylated N-methylglycine, and a precursor of N-methylglycine.

37. The pharmaceutical composition of claim 36, wherein the therapeutic agent is an antipsychotic selected from the group consisting of typical antipsychotics, atypical antipsychotics, and depot antipsychotics.

38. The pharmaceutical composition of claim 36, wherein the therapeutic agent is selected from the group consisting of Chlorpromazine, Thioridazine, Mesoridazine, Fluphenazine, Perphenazine, Trifluoperazine, Thiothixene, Haloperidol, Loxapine, Molindone, Clozapine, Risperidone, Olanzapine, Quetiapine, Haloperidol decanoate,

- 25 -

Fluphenazine decanoate, Fluphenazine enanthate,
Amitriptyline, Amoxapine, Bupropion, Bupropion SR,
Clomipramine, Desipramine, Doxepin, Fluoxetine,
Fluvoxamine, Imipramine, Maprotiline, Mirtazapine,
5 Nefazodone, Nortriptyline, Paroxetine, Phenelzine,
Protriptyline, Sertraline, Tranylcypromine, Trazodone,
Trimipramine, Venlafaxine, Venlafaxine XR,
Dextroamphetamine, Methamphetamine, Methylphenidate,
Pemoline, Donepezil, Tacrine, Acetophenazine,
10 Chlorprothixene, Droperidol, Pimozide, Butaperazine,
Carphenazine, Remoxipride, Piperacetazine, Sulpiride, and
Ziprasidone.