(12) STANDARD PATENT (11) Application No. **AU 2019336588 B2**
(19) AUSTRALIAN PATENT OFFICE

(54) Title
KRAS G12C inhibitors encompassing a piperazine ring and use thereof in the treatment of cancer

(51) International Patent Classification(s)
C07D 471/04 (2006.01) ***A61K 45/06*** (2006.01)
A61K 31/519 (2006.01) ***A61P 35/00*** (2006.01)
A61K 31/551 (2006.01) ***C07D 487/04*** (2006.01)

(21) Application No: **2019336588** (22) Date of Filing: **2019.06.11**

(87) WIPO No: **WO20/050890**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/684,117	**2018.06.12**	**US**

(43) Publication Date: **2020.03.12**
(44) Accepted Journal Date: **2022.07.28**

(71) Applicant(s)
Amgen Inc.

(72) Inventor(s)
ALLEN, John Gordon;LANMAN, Brian Alan;CHEN, Jian;REED, Anthony B.;CEE, Victor J.;LIU, Longbin;LOPEZ, Patricia;WURZ, Ryan Paul;NGUYEN, Thomas T.;BOOKER, Shon;ALLEN, Jennifer Rebecca;CHU-MOYER, Margaret;AMEGADZIE, Albert;CHEN, Ning;GOODMAN, Clifford;LOW, Jonathan D.;MA, Vu Van;MINATTI, Ana Elena;NISHIMURA, Nobuko;PICKRELL, Alexander J.;WANG, Hui-Ling;SHIN, Youngsook;SIEGMUND, Aaron C.;YANG, Kevin C.;TAMAYO, Nuria A.;WALTON, Mary;XUE, Qiufen

(74) Agent / Attorney
Spruson & Ferguson, GPO Box 3898, Sydney, NSW, 2001, AU

(56) Related Art
WO 2015/054572 A1
WO 2017/201161 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
12 March 2020 (12.03.2020)

WIPO | PCT

(10) International Publication Number
WO 2020/050890 A3

(51) International Patent Classification:
C07D 471/04 (2006.01) *A61K 31/551* (2006.01)
C07D 487/04 (2006.01) *A61P 35/00* (2006.01)
A61K 31/519 (2006.01) *A61K 45/06* (2006.01)

(21) International Application Number:
PCT/US2019/036626

(22) International Filing Date:
11 June 2019 (11.06.2019)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/684,117 12 June 2018 (12.06.2018) US

(71) Applicant: **AMGEN INC.** [US/US]; One Amgen Center Drive, Thousand Oaks, California 91320 (US).

(72) Inventors: **ALLEN, John Gordon**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **LANMAN, Brian Alan**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **CHEN, Jian**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **REED, Anthony B.**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **CEE, Victor J.**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **LIU, Longbin**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **LOPEZ, Patricia**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **WURZ, Ryan Paul**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **NGUYEN, Thomas T.**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **BOOKER, Shon**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **ALLEN, Jennifer Rebecca**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **CHU-MOYER, Margaret**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **AMEGADZIE, Albert**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **CHEN, Ning**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **GOODMAN, Clifford**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **LOW, Jonathan D.**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **MA, Vu Van**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **MINATTI, Ana Elena**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **NISHIMURA, Nobuko**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **PICKRELL, Alexander J.**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **WANG, Hui-Ling**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **SHIN, Youngsook**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **SIEGMUND, Aaron C.**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **YANG, Kevin C.**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **TAMAYO, Nuria A.**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **WALTON, Mary**; One Amgen Center Drive, Thousand Oaks, California 91320 (US). **XUE, Qiufen**; One Amgen Center Drive, Thousand Oaks, California 91320 (US).

(74) Agent: **REIDY, Joseph F.**; One Amgen Center Drive, Thousand Oaks, California 91320 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:
— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
— *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

WO 2020/050890 A3

(54) Title: KRAS G12C INHIBITORS ENCOMPASSING A PIPERAZINE RING AND USE THEREOF IN THE TREATMENT OF CANCER



(57) Abstract: Provided herein are KRAS G12C inhibitors, such as [Insert Compound Here] composition of the same, and methods of using the same. These inhibitors are useful for treating a number of disorders, including pancreatic, colorectal, and lung cancers.

[Continued on next page]

Published:

— *with international search report (Art. 21(3))*

— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

(88) Date of publication of the international search report:
25 June 2020 (25.06.2020)

KRAS G12C INHIBITORS AND METHODS OF USING THE SAME

RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional patent application 62/684,117 filed on June 12, 2018, which specification is hereby incorporated herein by reference in its entirety for all purposes.

[0002] The present application is being filed along with a sequence listing in electronic format. The sequence listing is provided as a file entitled A-2263-WO-PCT_SeqList_061119_ST25.txt, created June 11, 2019, which is 15.5 kb in size. The information in the electronic format of the sequence listing is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0003] The present invention relates to compounds that inhibit the KRAS G12C protein; methods of treating diseases or conditions, such as cancer, using the compounds; and pharmaceutical compositions containing the compounds.

BACKGROUND

[0004] KRAS gene mutations are common in pancreatic cancer, lung adenocarcinoma, colorectal cancer, gall bladder cancer, thyroid cancer, and bile duct cancer. KRAS mutations are also observed in about 25% of patients with NSCLC, and some studies have indicated that KRAS mutations are a negative prognostic factor in patients with NSCLC. Recently, V-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog (KRAS) mutations have been found to confer resistance to epidermal growth factor receptor (EGFR) targeted therapies in colorectal cancer; accordingly, the mutational status of KRAS can provide important information prior to the prescription of TKI therapy. Taken together, there is a need for new medical treatments for patients with pancreatic cancer, lung adenocarcinoma, or colorectal cancer, especially those who have been diagnosed to have such cancers characterized by a KRAS mutation, and including those who have progressed after chemotherapy.

[0005] The compounds disclosed herein can be in the form of a pharmaceutically acceptable salt. The compounds provided can be formulated into a pharmaceutical formulation comprising a compound disclosed herein and a pharmaceutically acceptable excipient.

[0006] Also provided is a method of inhibiting KRAS G12C in a cell, comprising contacting the cell with a compound or composition disclosed herein. Further provided is a method of treating cancer in a subject comprising administering to the subject a therapeutically effective

2019336588 14 Jun 2022

amount of a compound or composition disclosed herein. In some embodiments, the cancer is lung cancer, pancreatic cancer, or colorectal cancer.

SUMMARY

[0006a] In a first aspect of the present invention, there is provided a compound having a structure selected from the formula

 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
	 ;
	 ;
 ;	 ;

2019336588 14 Jun 2022

	 ;
 ;	 ;
	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
	 ;
 ;	 ;
	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
	 ;
 ;	 ;
	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2hh

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2mm

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ; and
	

2019336588 14 Jun 2022

or a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof, or a pharmaceutically acceptable salt of the atropisomer thereof.

[0006b] In a second aspect of the present invention, there is provided a compound having a structure selected from the formula

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ; and
	

or a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof, or a pharmaceutically acceptable salt of the atropisomer thereof.

[0006c] In a third aspect of the present invention, there is provided a pharmaceutical composition comprising the compound of the first or second aspect and a pharmaceutically acceptable excipient.

2019336588 14 Jun 2022

[0006d] In a fourth aspect of the present invention, there is provided a method of inhibiting KRAS G12C in a cell, comprising contacting the cell with the compound first or second aspect or the composition of the third aspect.

[0006e] In a fifth aspect of the present invention, there is provided use of the compound of the first or second aspect or the composition of the third aspect in the manufacture of a medicament for inhibiting KRAS G12C in a cell.

[0006f] In a sixth aspect of the present invention, there is provided a method of treating cancer in a subject comprising administering to the subject a therapeutically effective amount of the compound of the first or second aspect or the composition of the third aspect, wherein the cancer is mediated by a KRAS G12C mutation.

[0006g] In a seventh aspect of the present invention, there is provided use of the compound of the first or second aspect or the composition of the third aspect in the manufacture of a medicament for treating cancer in a subject, wherein the cancer is mediated by a KRAS G12C mutation.

[0007] Disclosed herein is a compound having a structure selected from the formula

2019336588 14 Jun 2022

	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ; and

 ;	

or a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof, or a pharmaceutically acceptable salt of the atropisomer thereof.

[0008] Another aspect of the present invention provides a compound having a structure selected from the formula

 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ; and
	

or a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof, or a pharmaceutically acceptable salt of the atropisomer thereof.

[0009] Another aspect of the present invention provides various compounds, stereoisomers, atropisomers, pharmaceutically acceptable salts, pharmaceutically acceptable salts of the stereoisomers, and pharmaceutically acceptable salts of the atropisomers as described in the embodiments set forth below.

[0010] Another aspect of the present invention provides a pharmaceutical composition that includes the compound of any of the embodiments or the pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.

[0011] Another aspect of the present invention provides a method of treating cancer. Such methods include: administering to a patient in need thereof a therapeutically effective amount of the compound of any of the embodiments or a pharmaceutically acceptable salt thereof. In some such methods, the cancer is a solid tumor. In some such methods, the cancer is selected from the group consisting of breast cancer, colorectal cancer, skin cancer, melanoma, ovarian cancer, kidney cancer, lung cancer, non-small cell lung cancer, cancer of the appendix, lymphoma, non-Hodgkin's lymphoma, myeloma, multiple myeloma, leukemia, and acute myelogenous leukemia. In some other such methods, the cancer is multiple myeloma.

[0012] In another aspect, the method further includes administering to a patient in need thereof a therapeutically effective amount of one or more additional pharmaceutically active compounds. For example, in some such methods the one or more additional pharmaceutically active compounds is pembrolizumab. In others, the one or more additional pharmaceutically active compounds is niolumab. In still other such methods, the one or more additional pharmaceutically active compounds is AMG 404. In still other such methods, the one or more additional pharmaceutically active compounds is daratumumab. In still other such methods, the one or more additional pharmaceutically active compound is a MEK inhibitor. In still other such methods, the MEK inhibitor is tremetinib. In still other such methods, the one or more additional pharmaceutically active compounds is an immunomodulatory agent (IMiD).

[0013] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Methods and materials are described herein for use in the present disclosure; other, suitable methods and materials known in the art can also be used. The materials, methods, and examples are illustrative only and not intended to be limiting. All publications, patent

applications, patents, sequences, database entries, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control.

[0014] Other features and advantages of the disclosure will be apparent from the following detailed description and figures, and from the Claims.

DETAILED DESCRIPTION

Definitions

Abbreviations: The following abbreviations may be used herein:

AcOH	acetic acid
aq or aq.	aqueous
BOC or Boc	*tert*-butyloxycarbonyl
CPhos	2-dicyclohexylphosphino-2′,6′-bis(*N*,*N*-dimethylamino)biphenyl
cpme	cyclopentyl methyl ether
DCE	1,2-dichloroethane
DABCO	1,4-diazabicyclo[2.2.2]octane
DCM	dichloromethane
DMA	*N*,*N*-dimethylacetamide
DMAP	4-dimethylaminopyridine
DME	1,2-dimethoxyethane
DMF	*N*,*N*-dimethylformamide
DMSO	dimethyl sulfoxide
Dppf, DPPF or dppf	1,1'-bis(diphenylphosphino)ferrocene
eq or eq. or equiv.	equivalent
ESI or ES	electrospray ionization
Et	ethyl
Et_2O	diethyl ether
EtOAc	ethyl acetate
g	gram(s)
h	hour(s)
HBTU	*N*,*N*,*N*′,*N*′-tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uronium hexafluorophosphate, *O*-(benzotriazol-1-yl)-*N*,*N*,*N*′,*N*′-tetramethyluronium hexafluorophosphate
HPLC	high pressure liquid chromatography
iPr	isopropyl
iPr_2NEt or DIPEA	*N*-ethyl diisopropylamine (Hünig's base)
KHMDS	potassium hexamethyldisilazide
KOAc	potassium acetate

Lawesson's reagent	2,4-bis(4-methoxyphenyl)-2,4-dithioxo-1,3,2,4-dithiadiphosphetane, 2,4-bis-(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane 2,4-disulfide
LC MS, LCMS, LC-MS or LC/MS	liquid chromatography mass spectroscopy
LG	leaving group (e.g., halogen, mesylate, triflate)
LHMDS or LiHMDS	lithium hexamethyldisilazide
m/z	mass divided by charge
Me	methyl
MeCN	acetonitrile
MeOH	methanol
Met	metal species for cross-coupling (e.g., MgX, ZnX, SnR_3, SiR_3, $B(OR)_2$)
mg	milligrams
min	minutes
mL	milliliters
MS	mass spectra
NaHMDS	sodium hexamethyldisilazide
NBS	*N*-bromosuccinimide
n-BuLi	*n*-butyllithium
NCS	*N*-chlorosuccinimide
NMR	nuclear magnetic resonance
$Pd_2(dba)_3$	tris(dibenzylideneacetone)dipalladium(0)
$Pd(dppf)Cl_2 \cdot DCM$, $Pd(dppf)Cl_2$	[1,1′-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane
$Pd(PPh_3)_4$	tetrakis(triphenylphosphine)palladium(0)
Ph	phenyl
PR or PG or Prot. group	protecting group
rbf	round-bottomed flask
RP-HPLC	reverse phase high pressure liquid chromatography
RT or rt or r.t.	room temperature
sat. or satd.	saturated

SFC	supercritical fluid chromatography
SPhos Pd G3 or SPhos G3	(2-dicyclohexylphosphino-2′,6′-dimethoxybiphenyl) [2-(2′-amino-1,1′-biphenyl)]palladium(II) methanesulfonate
TBAF	tetra-*n*-butylammonium fluoride
TBTU	*N,N,N′,N′*-tetramethyl-*O*-(benzotriazol-1-yl)uronium tetrafluoroborate
t-BuOH	*tert*-butanol
TEA or Et_3N	trimethylamine
TFA	trifluoroacetic acid
THF	tetrahydrofuran
UV	ultraviolet

[0015] The use of the terms "a," "an," "the," and similar referents in the context of describing the invention (especially in the context of the claims) are to be construed to cover both the singular and the plural, unless otherwise indicated. Recitation of ranges of values herein merely are intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. The use of any and all examples, or exemplary language (e.g., "such as") provided herein, is intended to better illustrate the invention and is not a limitation on the scope of the invention unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

[0016] As used herein, the term "alkyl" refers to straight chained and branched C1-C_8 hydrocarbon groups, including but not limited to, methyl, ethyl, npropyl, ipropyl, nbutyl, secbutyl, tbutyl, npentyl, 2methylbutyl, 3methylbutyl, 2,2dimethylpropyl, nhexyl, 2methylpentyl, 3-methylpentyl, 4-methylpentyl, 2,2dimethylbutyl, 2,3dimethylbutyl, 3,3dimethylbutyl, and 2ethybutyl. The term C_{m-n} means the alkyl group has "m" to "n" carbon atoms. The term "alkylene" refers to an alkyl group having a substituent. An alkyl (e.g., methyl), or alkylene (e.g., -CH_2-), group can be substituted with one or more, and typically one to three, of independently selected, for example, halo, trifluoromethyl, trifluoromethoxy, hydroxy, alkoxy, nitro, cyano, alkylamino, C_1-$_6$alkyl, C_2-$_6$alkenyl, C_2-$_6$alkynyl, -NC, amino, -CO_2H, -CO_2C_1-C_6alkyl, -$OCOC_1$-C_6alkyl, C_3-C_{10} cycloalkyl, C_3-C_{10} heterocycloalkyl, C_5-

C_{10}aryl, and C_5-C_{10} heteroaryl. The term "haloalkyl" specifically refers to an alkyl group wherein at least one, e.g., one to six, or all of the hydrogens of the alkyl group are substituted with halo atoms.

[0017] The terms "alkenyl" and "alkynyl" indicate an alkyl group that further includes a double bond or a triple bond, respectively.

[0018] As used herein, the term "halo" refers to fluoro, chloro, bromo, and iodo. The term "alkoxy" is defined as -OR, wherein R is alkyl.

[0019] As used herein, the term "amino" or "amine" interchangeably refers to a $-NR_2$ group, wherein each R is, e.g., H or a substituent. In some embodiments, the amino group is further substituted to form an ammonium ion, e.g., NR_3^+. Ammonium moieties are specifically included in the definition of "amino" or "amine." Substituents can be, for example, an alkyl, alkoxy, cycloalkyl, heterocycloalkyl, amide, or carboxylate. An R group may be further substituted, for example, with one or more, e.g., one to four, groups selected from halo, cyano, alkenyl, alkynyl, alkyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, urea, carbonyl, carboxylate, amine, and amide. An "amide" or "amido" group interchangeably refers to a group similar to an amine or amino group but further including a C(O), e.g., $-C(O)NR_2$. Some contemplated amino or amido groups (some with optional alkylene groups, e.g., alkylene-amino, or alkylene-amido) include CH_2NH_2, $CH(CH_3)NH_2$, $CH(CH_3)_2NH_2$, $CH_2CH_2NH_2$, $CH_2CH_2N(CH_3)_2$, CH_2NHCH_3, $C(O)NHCH_3$, $C(O)N(CH_3)_2$, $CH_2C(O)NH$phenyl, $CH_2NHC(O)CH_3$, $CH_2NHCH_2CH_2OH$, $CH_2NHCH_2CO_2H$, $CH_2NH(CH_3)CH_2CO_2CH_3$,$CH_2NHCH_2CH_2OCH_3$, $CH_2NH(CH_3)CH_2CH_2OCH_3$, $CH_2NH(CH_3)CH_2C(O)N(CH_3)_2$, $CH_2NH(CH_3)CH_2C(O)NHCH_3$, CH_2CH_2CCH, CH_2NMe_2, $CH_2NH(CH_3)CH_2CH_2OH$, $CH_2NH(CH_3)CH_2CH_2F$, $CH_2N^+(CH_3)_3$, $CH_2NHCH_2CHF_2$, $CH_2NHCH_2CH_3$,


















[0020] Collectively, antibodies form a family of plasma proteins known as immunoglobulins and comprise of immunoglobulin domains. (Janeway et al., Immunobiology: The Immune System in Health and Disease, 4th ed., Elsevier Science Ltd./Garland Publishing, 1999. As used herein, the term "antibody" refers to a protein having a conventional immunoglobulin format, comprising heavy and light chains, and comprising variable and constant regions. For example, an antibody may be an IgG which is a "Y-shaped" structure of two identical pairs of polypeptide chains, each pair having one "light" (typically having a molecular weight of about 25 kDa) and one "heavy" chain (typically having a molecular weight of about 50-70 kDa). An antibody has a variable region and a constant region. In IgG formats, the variable region is generally about 100-110 or more amino acids, comprises three complementarity determining regions (CDRs), is primarily responsible for antigen recognition, and substantially varies among other antibodies that bind to different antigens. The constant region allows the antibody to recruit cells and molecules of the immune system. The variable region is made of the N-terminal regions of each light chain and heavy chain, while the constant region is made of the C-terminal portions of each of the heavy and light chains. (Janeway et al., "Structure of the Antibody Molecule and the Immunoglobulin Genes", Immunobiology: The Immune System in Health and Disease, 4th ed. Elsevier Science Ltd./Garland Publishing, (1999)).

[0021] The general structure and properties of CDRs of antibodies have been described in the art. Briefly, in an antibody scaffold, the CDRs are embedded within a framework in the heavy and light chain variable region where they constitute the regions largely responsible for antigen binding and recognition. A variable region typically comprises at least three heavy or light chain CDRs (Kabat et al., 1991, Sequences of Proteins of Immunological Interest, Public

Health Service N.I.H., Bethesda, Md.; see also Chothia and Lesk, 1987, J. Mol. Biol. 196:901-917; Chothia et al., 1989, Nature 342: 877-883), within a framework region (designated framework regions 1-4, FR1, FR2, FR3, and FR4, by Kabat et al., 1991; see also Chothia and Lesk, 1987, supra).

[0022] Antibodies can comprise any constant region known in the art. Human light chains are classified as kappa and lambda light chains. Heavy chains are classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. IgG has several subclasses, including, but not limited to IgG1, IgG2, IgG3, and IgG4. IgM has subclasses, including, but not limited to, IgM1 and IgM2. Embodiments of the present disclosure include all such classes or isotypes of antibodies. The light chain constant region can be, for example, a kappa- or lambda-type light chain constant region, e.g., a human kappa- or lambda-type light chain constant region. The heavy chain constant region can be, for example, an alpha-, delta-, epsilon-, gamma-, or mu-type heavy chain constant regions, e.g., a human alpha-, delta-, epsilon-, gamma-, or mu-type heavy chain constant region. Accordingly, in exemplary embodiments, the antibody is an antibody of isotype IgA, IgD, IgE, IgG, or IgM, including any one of IgG1, IgG2, IgG3 or IgG4.

[0023] The antibody can be a monoclonal antibody or a polyclonal antibody. In some embodiments, the antibody comprises a sequence that is substantially similar to a naturally-occurring antibody produced by a mammal, e.g., mouse, rabbit, goat, horse, chicken, hamster, human, and the like. In this regard, the antibody can be considered as a mammalian antibody, e.g., a mouse antibody, rabbit antibody, goat antibody, horse antibody, chicken antibody, hamster antibody, human antibody, and the like. In certain aspects, the antibody is a human antibody. In certain aspects, the antibody is a chimeric antibody or a humanized antibody. The term "chimeric antibody" refers to an antibody containing domains from two or more different antibodies. A chimeric antibody can, for example, contain the constant domains from one species and the variable domains from a second, or more generally, can contain stretches of amino acid sequence from at least two species. A chimeric antibody also can contain domains of two or more different antibodies within the same species. The term "humanized" when used in relation to antibodies refers to antibodies having at least CDR regions from a non-human source which are engineered to have a structure and immunological function more similar to true human antibodies than the original source antibodies. For example, humanizing can involve grafting a CDR from a non-human antibody, such as a mouse antibody, into a human antibody. Humanizing also can involve select amino acid substitutions to make a non-human sequence more similar to a human sequence.

[0024] An antibody can be cleaved into fragments by enzymes, such as, e.g., papain and pepsin. Papain cleaves an antibody to produce two Fab fragments and a single Fc fragment. Pepsin cleaves an antibody to produce a F(ab')2 fragment and a pFc' fragment. As used herein, the term "antigen binding antibody fragment refers to a portion of an antibody molecule that is capable of binding to the antigen of the antibody and is also known as "antigen-binding fragment" or "antigen-binding portion". In exemplary instances, the antigen binding antibody fragment is a Fab fragment or a F(ab')2 fragment.

[0025] The architecture of antibodies has been exploited to create a growing range of alternative formats that span a molecular-weight range of at least about 12–150 kDa and has a valency (n) range from monomeric (n = 1), to dimeric (n = 2), to trimeric (n = 3), to tetrameric (n = 4), and potentially higher; such alternative formats are referred to herein as "antibody protein products". Antibody protein products include those based on the full antibody structure and those that mimic antibody fragments which retain full antigen-binding capacity, e.g., scFvs, Fabs and VHH/VH (discussed below). The smallest antigen binding antibody fragment that retains its complete antigen binding site is the Fv fragment, which consists entirely of variable (V) regions. A soluble, flexible amino acid peptide linker is used to connect the V regions to a scFv (single chain fragment variable) fragment for stabilization of the molecule, or the constant (C) domains are added to the V regions to generate a Fab fragment [fragment, antigen-binding]. Both scFv and Fab fragments can be easily produced in host cells, e.g., prokaryotic host cells. Other antibody protein products include disulfide-bond stabilized scFv (ds-scFv), single chain Fab (scFab), as well as di- and multimeric antibody formats like dia-, tria- and tetra-bodies, or minibodies (miniAbs) that comprise different formats consisting of scFvs linked to oligomerization domains. The smallest fragments are VHH/VH of camelid heavy chain Abs as well as single domain Abs (sdAb). The building block that is most frequently used to create novel antibody formats is the single-chain variable (V)-domain antibody fragment (scFv), which comprises V domains from the heavy and light chain (VH and VL domain) linked by a peptide linker of ~15 amino acid residues. A peptibody or peptide-Fc fusion is yet another antibody protein product. The structure of a peptibody consists of a biologically active peptide grafted onto an Fc domain. Peptibodies are well-described in the art. See, e.g., Shimamoto et al., mAbs 4(5): 586-591 (2012).

[0026] Other antibody protein products include a single chain antibody (SCA); a diabody; a triabody; a tetrabody; bispecific or trispecific antibodies, and the like. Bispecific antibodies can be divided into five major classes: BsIgG, appended IgG, BsAb fragments, bispecific

fusion proteins and BsAb conjugates. See, e.g., Spiess et al., Molecular Immunology 67(2) Part A: 97-106 (2015).

[0027] As used herein, the term "aryl" refers to a C_{6-14} monocyclic or polycyclic aromatic group, preferably a C_{6-10} monocyclic or bicyclic aromatic group, or C_{10-14} polycyclic aromatic group. Examples of aryl groups include, but are not limited to, phenyl, naphthyl, fluorenyl, azulenyl, anthryl, phenanthryl, pyrenyl, biphenyl, and terphenyl. Aryl also refers to C_{10-14} bicyclic and tricyclic carbon rings, where one ring is aromatic and the others are saturated, partially unsaturated, or aromatic, for example, dihydronaphthyl, indenyl, indanyl, or tetrahydronaphthyl (tetralinyl). Unless otherwise indicated, an aryl group can be unsubstituted or substituted with one or more, and in particular one to four, groups independently selected from, for example, halo, C_{1-6}alkyl, C_{2-6}alkenyl, C_{2-6}alkynyl, $-CF_3$, $-OCF_3$, $-NO_2$, -CN, -NC, -OH, alkoxy, amino, $-CO_2H$, $-CO_2C_1$-C_6alkyl, $-OCOC_1$-C_6alkyl, C_3-C_{10} cycloalkyl, C_3-C_{10} heterocycloalkyl, C_5-C_{10}aryl, and C_5-C_{10} heteroaryl.

[0028] As used herein, the term "carbocyclic ring" refers to a monocyclic ring which only includes carbon atoms as ring members. Such rings may be fully saturated, partially saturated, or aromatic and may include 3 to 10 carbon atoms.

[0029] As used herein, the term "cycloalkyl" refers to a monocyclic or polycyclic non-aromatic carbocyclic ring, where the polycyclic ring can be fused, bridged, or spiro. The carbocyclic ring can have 3 to 10 carbon ring atoms. Contemplated carbocyclic rings include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, and cyclononyl.

[0030] As used herein, the term "heterocycloalkyl" means a monocyclic or polycyclic (e.g., bicyclic), saturated or partially unsaturated, ring system containing 3 or more (e.g., 3 to 12, 4 to 10, 4 to 8, or 5 to 7) total atoms, of which one to five (e.g., 1, 2, 3, 4, or 5) of the atoms are independently selected from nitrogen, oxygen, and sulfur. Nonlimiting examples of heterocycloalkyl groups include azetidinyl, pyrrolidinyl, piperidinyl, piperazinyl, dihydropyrrolyl, morpholinyl, thiomorpholinyl, dihydropyridinyl, oxacycloheptyl, dioxacycloheptyl, thiacycloheptyl, and diazacycloheptyl.

[0031] Unless otherwise indicated, a cycloalkyl or heterocycloalkyl group can be unsubstituted or substituted with one or more, and in particular one to four, groups. Some contemplated substituents include halo, C_{1-6}alkyl, C_{2-6}alkenyl, C_{2-6}alkynyl, $-OCF_3$, $-NO_2$, -CN, -NC, -OH, alkoxy, amino, $-CO_2H$, $-CO_2C_1$-C_6alkyl, $-OCOC_1$-C_8alkyl, C_3-C_{10} cycloalkyl, C_3-C_{10} heterocycloalkyl, C_5-C_{10}aryl, and C_5-C_{10} heteroaryl.

[0032] As used herein, the term "heteroaryl" refers to a monocyclic or polycyclic ring system (for example, bicyclic) containing one to three aromatic rings and containing one to four (e.g., 1, 2, 3, or 4) heteroatoms selected from nitrogen, oxygen, and sulfur in an aromatic ring. In certain embodiments, the heteroaryl group has from 5 to 20, from 5 to 15, from 5 to 10 ring, or from 5 to 7 atoms. Heteroaryl also refers to C_{10-14} bicyclic and tricyclic rings, where one ring is aromatic and the others are saturated, partially unsaturated, or aromatic. Examples of heteroaryl groups include, but are not limited to, furanyl, imidazolyl, isothiazolyl, isoxazolyl, oxadiazolyl, oxazolyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridyl, pyrimidinyl, pyrrolyl, thiadiazolyl, thiazolyl, thienyl, tetrazolyl, triazinyl, triazolyl, benzofuranyl, benzimidazolyl, benzoisoxazolyl, benzopyranyl, benzothiadiazolyl, benzothiazolyl, benzothienyl, benzothiophenyl, benzotriazolyl, benzoxazolyl, furopyridyl, imidazopyridinyl, imidazothiazolyl, indolizinyl, indolyl, indazolyl, isobenzofuranyl, isobenzothienyl, isoindolyl, isoquinolinyl, isothiazolyl, naphthyridinyl, oxazolopyridinyl, phthalazinyl, pteridinyl, purinyl, pyridopyridyl, pyrrolopyridyl, quinolinyl, quinoxalinyl, quiazolinyl, thiadiazolopyrimidyl, and thienopyridyl. Unless otherwise indicated, a heteroaryl group can be unsubstituted or substituted with one or more, and in particular one to four or one or two, substituents. Contemplated substituents include halo, C_{1-6}alkyl, C_{2-6}alkenyl, C_{2-6}alkynyl, $-OCF_3$, $-NO_2$, -CN, -NC, -OH, alkoxy, amino, $-CO_2H$, $-CO_2C_1$-C_6alkyl, $-OCOC_1$-C_6alkyl, C_3-C_{10} cycloalkyl, C_3-C_{10} heterocycloalkyl, C_5-C_{10}aryl, and C_5-C_{10} heteroaryl.

[0033] As used herein, the term Boc refers to the structure  .

[0034] As used herein, the term Cbz refers to the structure  .

[0035] As used herein, the term Bn refers to the structure  .

[0036] As used herein, the term trifluoroacetamide refers to the structure  .

[0037] As used herein, the term trityl refers to the structure .

[0038] As used herein, the term tosyl refers to the structure  .

[0039] As used herein, the term Troc refers to the structure  .

[0040] As used herein, the term Teoc refers to the structure  .

[0041] As used herein, the term Alloc refers to the structure  .

[0042] As used herein, the term Fmoc refers to the structure  .

COMPOUNDS OF THE DISCLOSURE

[0043] The compounds disclosed herein include all pharmaceutically acceptable isotopically-labeled compounds wherein one or more atoms of the compounds disclosed herein are replaced by atoms having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into the disclosed compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, chlorine, and iodine, such as ^{2}H, ^{3}H, ^{11}C, ^{13}C, ^{14}C, ^{13}N, ^{15}N, ^{15}O, ^{17}O, ^{18}O, ^{31}P, ^{32}P, ^{35}S, ^{18}F, ^{36}Cl, ^{123}I, and ^{125}I, respectively. These radiolabelled compounds could be useful to help determine or measure the effectiveness of the compounds, by characterizing, for example, the site or mode of action, or binding affinity to pharmacologically important site of action. Certain isotopically-labeled compounds of the disclosure, for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, i.e. ^{3}H, and carbon-14, i.e. ^{14}C, are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

[0044] Substitution with heavier isotopes such as deuterium, i.e. ^{2}H, may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased in vivo half-life or reduced dosage requirements, and hence are preferred in some circumstances.

[0045] Substitution with positron emitting isotopes, such as ^{11}C, ^{18}F, ^{15}O and ^{13}N, can be useful in Positron Emission Topography (PET) studies for examining substrate receptor occupancy. Isotopically-labeled compounds of structure (I) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the Preparations and Examples as set out below using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

[0046] Isotopically-labeled compounds as disclosed herein can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those

described in the accompanying examples and schemes using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

[0047] Certain of the compounds as disclosed herein may exist as stereoisomers (i.e., isomers that differ only in the spatial arrangement of atoms) including optical isomers and conformational isomers (or conformers). The compounds disclosed herein include all stereoisomers, both as pure individual stereoisomer preparations and enriched preparations of each, and both the racemic mixtures of such stereoisomers as well as the individual diastereomers and enantiomers that may be separated according to methods that are known to those skilled in the art. Additionally, the compounds disclosed herein include all tautomeric forms of the compounds.

[0048] Certain of the compounds disclosed herein may exist as atropisomers, which are conformational stereoisomers that occur when rotation about a single bond in the molecule is prevented, or greatly slowed, as a result of steric interactions with other parts of the molecule. The compounds disclosed herein include all atropisomers, both as pure individual atropisomer preparations, enriched preparations of each, or a non-specific mixture of each. Where the rotational barrier about the single bond is high enough, and interconversion between conformations is slow enough, separation and isolation of the isomeric species may be permitted. For example, groups such as, but not limited to, the following groups  ,  ,  ,  , and  may exhibit restricted rotation.

EMBODIMENTS

Embodiment 1

[0049] In one embodiment of the present invention, the present invention comprises a compound having a structure selected from the formula

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

	
	
	
	
	

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ; and
	

or a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof, or a pharmaceutically acceptable salt of the atropisomer thereof.

Embodiment 2

[0050] In another embodiment of the present invention, the present invention comprises a compound having a structure selected from the formula



 ;	 ;
 ;	 ;
 ;	 ; and
	

or a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof, or a pharmaceutically acceptable salt of the atropisomer thereof.

Embodiment 3

[0051] In another embodiment of the present invention, the present invention comprises the compound of any one of embodiments 1-2 in the form of a pharmaceutically acceptable salt.

Embodiment 4

[0052] In another embodiment of the present invention, the present invention comprises a pharmaceutical composition comprising the compound of any one of embodiments 1-3 and a pharmaceutically acceptable excipient.

Embodiment 5

[0053] In another embodiment of the present invention, the present invention comprises a method of inhibiting KRAS G12C in a cell, comprising contacting the cell with the compound of any one of embodiments 1-3 or the composition of embodiment 4.

Embodiment 6

[0054] In another embodiment of the present invention, the present invention comprises a method of treating cancer in a subject comprising administering to the subject a therapeutically effective amount of the compound of any one of embodiments 1-3 or the composition of embodiment 4.

Embodiment 7

[0055] In another embodiment of the present invention, the present invention comprises the method of embodiment 6, wherein the cancer is lung cancer, pancreatic cancer, or colorectal cancer.

Embodiment 8

[0056] In another embodiment of the present invention, the present invention comprises the method of embodiment 7, wherein the cancer is lung cancer.

Embodiment 9

[0057] In another embodiment of the present invention, the present invention comprises the method of embodiment 7, wherein the cancer is pancreatic cancer.

Embodiment 10

[0058] In another embodiment of the present invention, the present invention comprises the method of embodiment 7, wherein the cancer is colorectal cancer.

Embodiment 11

[0059] In another embodiment of the present invention, the present invention comprises the method of embodiment 6, further comprising administering to the patient in need thereof a

therapeutically effective amount of one or more additional pharmaceutically active compounds.

Embodiment 12

[0060] In another embodiment of the present invention, the present invention comprises the method of embodiment 11, wherein the one or more additional pharmaceutically active compounds is an anti-PD-1 antibody

Embodiment 13

[0061] In another embodiment of the present invention, the present invention comprises the method of embodiment 12, wherein the anti-PD-1 antibody is pembrolizumab.

Embodiment 14

[0062] In another embodiment of the present invention, the present invention comprises the method of embodiment 12 wherein the anti-PD-1 antibody is niolumab.

Embodiment 15

[0063] In another embodiment of the present invention, the present invention comprises the method of embodiment 11, wherein the one or more additional pharmaceutically active compounds is an MCl-1 inhibitor.

Embodiment 16

[0064] In another embodiment of the present invention, the present invention comprises the method of embodiment 11, wherein the one or more additional pharmaceutically active compounds is a MEK inhibitor.

Embodiment 17

[0065] In another embodiment of the present invention, the present invention comprises the method of embodiment 11, wherein the one or more additional pharmaceutically active compounds is daratumumab.

Embodiment 18

[0066] In another embodiment of the present invention, the present invention comprises the method of embodiment 11, wherein the one or more additional pharmaceutically active compounds is an immunomodulatory agent.

Embodiment 19

[0067] In another embodiment of the present invention, the present invention comprises the use of a compound according to any one of embodiments 1-3 for treating cancer in a subject.

Embodiment 20

[0068] In another embodiment of the present invention, the present invention comprises a compound according to any one of embodiments 1-3 in the preparation of a medicament for treating cancer.

Embodiment 21

[0069] In another embodiment of the present invention, the present invention comprises the compound according to embodiment 20, wherein the cancer isnon-small cell lung cancer.

SYNTHESIS OF DISCLOSED COMPOUNDS

[0070] Compounds as disclosed herein can be synthesized via a number of specific methods. The examples which outline specific synthetic routes, and the generic schemes below are meant to provide guidance to the ordinarily skilled synthetic chemist, who will readily appreciate that the solvent, concentration, reagent, protecting group, order of synthetic steps, time, temperature, and the like can be modified as necessary, well within the skill and judgment of the ordinarily skilled artisan.

Pharmaceutical compositions, dosing, and routes of administration

[0071] Also provided herein are pharmaceutical compositions that include a compound as disclosed herein, together with a pharmaceutically acceptable excipient, such as, for example, a diluent or carrier. Compounds and pharmaceutical compositions suitable for use in the present invention include those wherein the compound can be administered in an effective amount to achieve its intended purpose. Administration of the compound described in more detail below.

[0072] Suitable pharmaceutical formulations can be determined by the skilled artisan depending on the route of administration and the desired dosage. *See, e.g.*, Remington's Pharmaceutical Sciences, 1435-712 (18th ed., Mack Publishing Co, Easton, Pennsylvania, 1990). Formulations may influence the physical state, stability, rate of in vivo release and rate of in vivo clearance of the administered agents. Depending on the route of administration, a suitable dose may be calculated according to body weight, body surface areas or organ size. Further refinement of the calculations necessary to determine the appropriate treatment dose is routinely made by those of ordinary skill in the art without undue experimentation, especially

in light of the dosage information and assays disclosed herein as well as the pharmacokinetic data obtainable through animal or human clinical trials.

[0073] The phrases "pharmaceutically acceptable" or "pharmacologically acceptable" refer to molecular entities and compositions that do not produce adverse, allergic, or other untoward reactions when administered to an animal or a human. As used herein, "pharmaceutically acceptable" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such excipients for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the therapeutic compositions, its use in therapeutic compositions is contemplated. Supplementary active ingredients also can be incorporated into the compositions. In exemplary embodiments, the formulation may comprise corn syrup solids, high-oleic safflower oil, coconut oil, soy oil, L-leucine, calcium phosphate tribasic, L-tyrosine, L-proline, L-lysine acetate, DATEM (an emulsifier), L-glutamine, L-valine, potassium phosphate dibasic, L-isoleucine, L-arginine, L-alanine, glycine, L-asparagine monohydrate, L-serine, potassium citrate, L-threonine, sodium citrate, magnesium chloride, L-histidine, L-methionine, ascorbic acid, calcium carbonate, L-glutamic acid, L-cystine dihydrochloride, L-tryptophan, L-aspartic acid, choline chloride, taurine, m-inositol, ferrous sulfate, ascorbyl palmitate, zinc sulfate, L-carnitine, alpha-tocopheryl acetate, sodium chloride, niacinamide, mixed tocopherols, calcium pantothenate, cupric sulfate, thiamine chloride hydrochloride, vitamin A palmitate, manganese sulfate, riboflavin, pyridoxine hydrochloride, folic acid, beta-carotene, potassium iodide, phylloquinone, biotin, sodium selenate, chromium chloride, sodium molybdate, vitamin D3 and cyanocobalamin.

[0074] The compound can be present in a pharmaceutical composition as a pharmaceutically acceptable salt. As used herein, "pharmaceutically acceptable salts" include, for example base addition salts and acid addition salts.

[0075] Pharmaceutically acceptable base addition salts may be formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible. Examples of metals used as cations are sodium, potassium, magnesium, ammonium, calcium, or ferric, and the like. Examples of suitable amines include isopropylamine, trimethylamine, histidine, N,N'-dibenzylethylenediamine,

chloroprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine.

[0076] Pharmaceutically acceptable acid addition salts include inorganic or organic acid salts. Examples of suitable acid salts include the hydrochlorides, formates, acetates, citrates, salicylates, nitrates, phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include, for example, formic, acetic, citric, oxalic, tartaric, or mandelic acids, hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, trifluoroacetic acid (TFA), propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane 1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene 2-sulfonic acid, naphthalene 1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose 6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid.

[0077] Pharmaceutical compositions containing the compounds disclosed herein can be manufactured in a conventional manner, e.g., by conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. Proper formulation is dependent upon the route of administration chosen.

[0078] For oral administration, suitable compositions can be formulated readily by combining a compound disclosed herein with pharmaceutically acceptable excipients such as carriers well known in the art. Such excipients and carriers enable the present compounds to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by adding a compound as disclosed herein with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include, for example, fillers and cellulose preparations. If desired, disintegrating agents can be added. Pharmaceutically acceptable ingredients are well known for the various types of formulation

and may be for example binders (e.g., natural or synthetic polymers), lubricants, surfactants, sweetening and flavoring agents, coating materials, preservatives, dyes, thickeners, adjuvants, antimicrobial agents, antioxidants and carriers for the various formulation types.

[0079] When a therapeutically effective amount of a compound disclosed herein is administered orally, the composition typically is in the form of a solid (e.g., tablet, capsule, pill, powder, or troche) or a liquid formulation (e.g., aqueous suspension, solution, elixir, or syrup).

[0080] When administered in tablet form, the composition can additionally contain a functional solid and/or solid carrier, such as a gelatin or an adjuvant. The tablet, capsule, and powder can contain about 1 to about 95% compound, and preferably from about 15 to about 90% compound.

[0081] When administered in liquid or suspension form, a functional liquid and/or a liquid carrier such as water, petroleum, or oils of animal or plant origin can be added. The liquid form of the composition can further contain physiological saline solution, sugar alcohol solutions, dextrose or other saccharide solutions, or glycols. When administered in liquid or suspension form, the composition can contain about 0.5 to about 90% by weight of a compound disclosed herein, and preferably about 1 to about 50% of a compound disclosed herein. In one embodiment contemplated, the liquid carrier is non-aqueous or substantially non-aqueous. For administration in liquid form, the composition may be supplied as a rapidly-dissolving solid formulation for dissolution or suspension immediately prior to administration.

[0082] When a therapeutically effective amount of a compound disclosed herein is administered by intravenous, cutaneous, or subcutaneous injection, the composition is in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred composition for intravenous, cutaneous, or subcutaneous injection typically contains, in addition to a compound disclosed herein, an isotonic vehicle. Such compositions may be prepared for administration as solutions of free base or pharmacologically acceptable salts in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions also can be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations can optionally contain a preservative to prevent the growth of microorganisms.

[0083] Injectable compositions can include sterile aqueous solutions, suspensions, or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions, suspensions, or dispersions. In all embodiments the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must resist the contaminating action of microorganisms, such as bacteria and fungi, by optional inclusion of a preservative. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. In one embodiment contemplated, the carrier is non-aqueous or substantially non-aqueous. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size of the compound in the embodiment of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many embodiments, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

[0084] Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the embodiment of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0085] Slow release or sustained release formulations may also be prepared in order to achieve a controlled release of the active compound in contact with the body fluids in the GI tract, and to provide a substantially constant and effective level of the active compound in the blood plasma. For example, release can be controlled by one or more of dissolution, diffusion, and ion-exchange. In addition, the slow release approach may enhance absorption via saturable or limiting pathways within the GI tract. For example, the compound may be embedded for this purpose in a polymer matrix of a biological degradable polymer, a water-soluble polymer

or a mixture of both, and optionally suitable surfactants. Embedding can mean in this context the incorporation of micro-particles in a matrix of polymers. Controlled release formulations are also obtained through encapsulation of dispersed micro-particles or emulsified micro-droplets via known dispersion or emulsion coating technologies.

[0086] For administration by inhalation, compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant. In the embodiment of a pressurized aerosol, the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin, for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0087] The compounds disclosed herein can be formulated for parenteral administration by injection (e.g., by bolus injection or continuous infusion). Formulations for injection can be presented in unit dosage form (e.g., in ampules or in multidose containers), with an added preservative. The compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing, and/or dispersing agents.

[0088] Pharmaceutical formulations for parenteral administration include aqueous solutions of the compounds in water-soluble form. Additionally, suspensions of the compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils or synthetic fatty acid esters. Aqueous injection suspensions can contain substances which increase the viscosity of the suspension. Optionally, the suspension also can contain suitable stabilizers or agents that increase the solubility of the compounds and allow for the preparation of highly concentrated solutions. Alternatively, a present composition can be in powder form for constitution with a suitable vehicle (e.g., sterile pyrogen-free water) before use.

[0089] Compounds disclosed herein also can be formulated in rectal compositions, such as suppositories or retention enemas (e.g., containing conventional suppository bases). In addition to the formulations described previously, the compounds also can be formulated as a depot preparation. Such long-acting formulations can be administered by implantation (e.g., subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example,

as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0090] In particular, a compound disclosed herein can be administered orally, buccally, or sublingually in the form of tablets containing excipients, such as starch or lactose, or in capsules or ovules, either alone or in admixture with excipients, or in the form of elixirs or suspensions containing flavoring or coloring agents. Such liquid preparations can be prepared with pharmaceutically acceptable additives, such as suspending agents. A compound also can be injected parenterally, for example, intravenously, intramuscularly, subcutaneously, or intracoronarily. For parenteral administration, the compound is best used in the form of a sterile aqueous solution which can contain other substances, for example, salts, or sugar alcohols, such as mannitol, or glucose, to make the solution isotonic with blood.

[0091] For veterinary use, a compound disclosed herein is administered as a suitably acceptable formulation in accordance with normal veterinary practice. The veterinarian can readily determine the dosing regimen and route of administration that is most appropriate for a particular animal.

[0092] In some embodiments, all the necessary components for the treatment of KRAS-related disorder using a compound as disclosed herein either alone or in combination with another agent or intervention traditionally used for the treatment of such disease may be packaged into a kit. Specifically, the present invention provides a kit for use in the therapeutic intervention of the disease comprising a packaged set of medicaments that include the compound disclosed herein as well as buffers and other components for preparing deliverable forms of said medicaments, and/or devices for delivering such medicaments, and/or any agents that are used in combination therapy with the compound disclosed herein, and/or instructions for the treatment of the disease packaged with the medicaments. The instructions may be fixed in any tangible medium, such as printed paper, or a computer readable magnetic or optical medium, or instructions to reference a remote computer data source such as a world wide web page accessible via the internet.

[0100] A "therapeutically effective amount" means an amount effective to treat or to prevent development of, or to alleviate the existing symptoms of, the subject being treated. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. Generally, a "therapeutically effective dose" refers to that amount of the compound that results in achieving the desired

effect. For example, in one preferred embodiment, a therapeutically effective amount of a compound disclosed herein decreases KRAS activity by at least 5%, compared to control, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, or at least 90%.

[0101] The amount of compound administered can be dependent on the subject being treated, on the subject's age, health, sex, and weight, the kind of concurrent treatment (if any), severity of the affliction, the nature of the effect desired, the manner and frequency of treatment, and the judgment of the prescribing physician. The frequency of dosing also can be dependent on pharmacodynamic effects on arterial oxygen pressures. However, the most preferred dosage can be tailored to the individual subject, as is understood and determinable by one of skill in the art, without undue experimentation. This typically involves adjustment of a standard dose (e.g., reduction of the dose if the patient has a low body weight).

[0102] While individual needs vary, determination of optimal ranges of effective amounts of the compound is within the skill of the art. For administration to a human in the curative or prophylactic treatment of the conditions and disorders identified herein, for example, typical dosages of the compounds of the present invention can be about 0.05 mg/kg/day to about 50 mg/kg/day, for example at least 0.05 mg/kg, at least 0.08 mg/kg, at least 0.1 mg/kg, at least 0.2 mg/kg, at least 0.3 mg/kg, at least 0.4 mg/kg, or at least 0.5 mg/kg, and preferably 50 mg/kg or less, 40 mg/kg or less, 30 mg/kg or less, 20 mg/kg or less, or 10 mg/kg or less, which can be about 2.5 mg/day (0.5 mg/kg x 5kg) to about 5000 mg/day (50mg/kg x 100kg), for example. For example, dosages of the compounds can be about 0.1 mg/kg/day to about 50 mg/kg/day, about 0.05 mg/kg/day to about 10 mg/kg/day, about 0.05 mg/kg/day to about 5 mg/kg/day, about 0.05 mg/kg/day to about 3 mg/kg/day, about 0.07 mg/kg/day to about 3 mg/kg/day, about 0.09 mg/kg/day to about 3 mg/kg/day, about 0.05 mg/kg/day to about 0.1 mg/kg/day, about 0.1 mg/kg/day to about 1 mg/kg/day, about 1 mg/kg/day to about 10 mg/kg/day, about 1 mg/kg/day to about 5 mg/kg/day, about 1 mg/kg/day to about 3 mg/kg/day, about 3 mg/day to about 1500 mg/day, about 5 mg/day to about 1000 mg/day, about 10 mg/day to about 750 mg/day, about 3 mg/day to about 350 mg/day, or about 100 mg/day to about 250 mg/day. Such doses may be administered in a single dose or it may be divided into multiple doses.

Methods of using KRAS G12C inhibitors

[0103] The present disclosure provides a method of inhibiting RAS-mediated cell signaling comprising contacting a cell with an effective amount of one or more compounds disclosed herein. Inhibition of RAS-mediated signal transduction can be assessed and demonstrated by a wide variety of ways known in the art. Non-limiting examples include a showing of (a) a decrease in GTPase activity of RAS; (b) a decrease in GTP binding affinity or an increase in GDP binding affinity; (c) an increase in K off of GTP or a decrease in K off of GDP; (d) a decrease in the levels of signaling transduction molecules downstream in the RAS pathway, such as a decrease in pMEK, pERK, or pAKT levels; and/or (e) a decrease in binding of RAS complex to downstream signaling molecules including but not limited to Raf. Kits and commercially available assays can be utilized for determining one or more of the above.

[0104] The disclosure also provides methods of using the compounds or pharmaceutical compositions of the present disclosure to treat disease conditions, including but not limited to conditions implicated by G12C KRAS, HRAS or NRAS mutation (e.g., cancer).

[0105] In some embodiments, a method for treatment of cancer is provided, the method comprising administering an effective amount of any of the foregoing pharmaceutical compositions comprising a compound as disclosed herein to a subject in need thereof. In some embodiments, the cancer is mediated by a KRAS, HRAS or NRAS G12C mutation. In various embodiments, the cancer is pancreatic cancer, colorectal cancer or lung cancer. In some embodiments, the cancer is gall bladder cancer, thyroid cancer, and bile duct cancer.

[0106] In some embodiments the disclosure provides method of treating a disorder in a subject in need thereof, wherein the said method comprises determining if the subject has a KRAS, HRAS or NRAS G12C mutation and if the subject is determined to have the KRAS, HRAS or NRAS G12C mutation, then administering to the subject a therapeutically effective dose of at least one compound as disclosed herein or a pharmaceutically acceptable salt thereof.

[0107] The disclosed compounds inhibit anchorage-independent cell growth and therefore have the potential to inhibit tumor metastasis. Accordingly, another embodiment the disclosure provides a method for inhibiting tumor metastasis, the method comprising administering an effective amount a compound disclosed herein.

[0108] KRAS, HRAS or NRAS G12C mutations have also been identified in hematological malignancies (e.g., cancers that affect blood, bone marrow and/or lymph nodes). Accordingly, certain embodiments are directed to administration of a disclosed compounds (e.g., in the form

of a pharmaceutical composition) to a patient in need of treatment of a hematological malignancy. Such malignancies include, but are not limited to leukemias and lymphomas. For example, the presently disclosed compounds can be used for treatment of diseases such as Acute lymphoblastic leukemia (ALL), Acute myelogenous leukemia (AML), Chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), Chronic myelogenous leukemia (CML), Acute monocytic leukemia (AMoL) and/ or other leukemias. In other embodiments, the compounds are useful for treatment of lymphomas such as all subtypes of Hodgkins lymphoma or non-Hodgkins lymphoma. In various embodiments, the compounds are useful for treatment of plasma cell malignancies such as multiple myeloma, mantle cell lymphoma, and Waldenstrom's macroglubunemia.

[0109] Determining whether a tumor or cancer comprises a G12C KRAS, HRAS or NRAS mutation can be undertaken by assessing the nucleotide sequence encoding the KRAS, HRAS or NRAS protein, by assessing the amino acid sequence of the KRAS, HRAS or NRAS protein, or by assessing the characteristics of a putative KRAS, HRAS or NRAS mutant protein. The sequence of wild-type human KRAS, HRAS or NRAS is known in the art, (e.g. Accession No. NP203524).

[0110] Methods for detecting a mutation in a KRAS, HRAS or NRAS nucleotide sequence are known by those of skill in the art. These methods include, but are not limited to, polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) assays, polymerase chain reaction-single strand conformation polymorphism (PCR-SSCP) assays, real-time PCR assays, PCR sequencing, mutant allele-specific PCR amplification (MASA) assays, direct sequencing, primer extension reactions, electrophoresis, oligonucleotide ligation assays, hybridization assays, TaqMan assays, SNP genotyping assays, high resolution melting assays and microarray analyses. In some embodiments, samples are evaluated for G12C KRAS, HRAS or NRAS mutations by real-time PCR. In real-time PCR, fluorescent probes specific for the KRAS, HRAS or NRAS G12C mutation are used. When a mutation is present, the probe binds and fluorescence is detected. In some embodiments, the KRAS, HRAS or NRAS G12C mutation is identified using a direct sequencing method of specific regions (e.g., exon 2 and/or exon 3) in the KRAS, HRAS or NRAS gene. This technique will identify all possible mutations in the region sequenced.

[0111] Methods for detecting a mutation in a KRAS, HRAS or NRAS protein are known by those of skill in the art. These methods include, but are not limited to, detection of a KRAS,

HRAS or NRAS mutant using a binding agent (e.g., an antibody) specific for the mutant protein, protein electrophoresis and Western blotting, and direct peptide sequencing.

[0112] Methods for determining whether a tumor or cancer comprises a G12C KRAS, HRAS or NRAS mutation can use a variety of samples. In some embodiments, the sample is taken from a subject having a tumor or cancer. In some embodiments, the sample is a fresh tumor/cancer sample. In some embodiments, the sample is a frozen tumor/cancer sample. In some embodiments, the sample is a formalin-fixed paraffin-embedded sample. In some embodiments, the sample is a circulating tumor cell (CTC) sample. In some embodiments, the sample is processed to a cell lysate. In some embodiments, the sample is processed to DNA or RNA.

[0113] The disclosure also relates to a method of treating a hyperproliferative disorder in a mammal that comprises administering to said mammal a therapeutically effective amount of a compound as disclosed herein, or a pharmaceutically acceptable salt thereof. In some embodiments, said method relates to the treatment of a subject who suffers from a cancer such as acute myeloid leukemia, cancer in adolescents, adrenocortical carcinoma childhood, AIDS-related cancers (e.g. Lymphoma and Kaposi's Sarcoma), anal cancer, appendix cancer, astrocytomas, atypical teratoid, basal cell carcinoma, bile duct cancer, bladder cancer, bone cancer, brain stem glioma, brain tumor, breast cancer, bronchial tumors, Burkitt lymphoma, carcinoid tumor, atypical teratoid, embryonal tumors, germ cell tumor, primary lymphoma, cervical cancer, childhood cancers, chordoma, cardiac tumors, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myleoproliferative disorders, colon cancer, colorectal cancer, craniopharyngioma, cutaneous T-cell lymphoma, extrahepatic ductal carcinoma in situ (DCIS), embryonal tumors, CNS cancer, endometrial cancer, ependymoma, esophageal cancer, esthesioneuroblastoma, ewing sarcoma, extracranial germ cell tumor, extragonadal germ cell tumor, eye cancer, fibrous histiocytoma of bone, gall bladder cancer, gastric cancer, gastrointestinal carcinoid tumor, gastrointestinal stromal tumors (GIST), germ cell tumor, gestational trophoblastic tumor, hairy cell leukemia, head and neck cancer, heart cancer, liver cancer, Hodgkin lymphoma, hypopharyngeal cancer, intraocular melanoma, islet cell tumors, pancreatic neuroendocrine tumors, kidney cancer, laryngeal cancer, lip and oral cavity cancer, liver cancer, lobular carcinoma in situ (LCIS), lung cancer, lymphoma, metastatic squamous neck cancer with occult primary, midline tract carcinoma, mouth cancer, multiple endocrine neoplasia syndromes, multiple myeloma/plasma cell neoplasm, mycosis fungoides, myelodysplastic syndromes, myelodysplastic/myeloproliferative

neoplasms, multiple myeloma, merkel cell carcinoma, malignant mesothelioma, malignant fibrous histiocytoma of bone and osteosarcoma, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, neuroblastoma, non-hodgkin lymphoma, non-small cell lung cancer (NSCLC), oral cancer, lip and oral cavity cancer, oropharyngeal cancer, ovarian cancer, pancreatic cancer, papillomatosis, paraganglioma, paranasal sinus and nasal cavity cancer, parathyroid cancer, penile cancer, pharyngeal cancer, pleuropulmonary blastoma, primary central nervous system (CNS) lymphoma, prostate cancer, rectal cancer, transitional cell cancer, retinoblastoma, rhabdomyosarcoma, salivary gland cancer, skin cancer, stomach (gastric) cancer, small cell lung cancer, small intestine cancer, soft tissue sarcoma, T-Cell lymphoma, testicular cancer, throat cancer, thymoma and thymic carcinoma, thyroid cancer, transitional cell cancer of the renal pelvis and ureter, trophoblastic tumor, unusual cancers of childhood, urethral cancer, uterine sarcoma, vaginal cancer, vulvar cancer, or viral-induced cancer. In some embodiments, said method relates to the treatment of a non-cancerous hyperproliferative disorder such as benign hyperplasia of the skin (e. g., psoriasis), restenosis, or prostate (e. g., benign prostatic hypertrophy (BPH)).

[0114] In some embodiments, the methods for treatment are directed to treating lung cancers, the methods comprise administering an effective amount of any of the above described compound (or a pharmaceutical composition comprising the same) to a subject in need thereof. In certain embodiments the lung cancer is a non- small cell lung carcinoma (NSCLC), for example adenocarcinoma, squamous-cell lung carcinoma or large-cell lung carcinoma. In some embodiments, the lung cancer is a small cell lung carcinoma. Other lung cancers treatable with the disclosed compounds include, but are not limited to, glandular tumors, carcinoid tumors and undifferentiated carcinomas.

[0115] The disclosure further provides methods of modulating a G12C Mutant KRAS, HRAS or NRAS protein activity by contacting the protein with an effective amount of a compound of the disclosure. Modulation can be inhibiting or activating protein activity. In some embodiments, the disclosure provides methods of inhibiting protein activity by contacting the G12C Mutant KRAS, HRAS or NRAS protein with an effective amount of a compound of the disclosure in solution. In some embodiments, the disclosure provides methods of inhibiting the G12C Mutant KRAS, HRAS or NRAS protein activity by contacting a cell, tissue, or organ that expresses the protein of interest. In some embodiments, the disclosure provides methods of inhibiting protein activity in subject including but not limited to rodents

and mammal (e.g., human) by administering into the subject an effective amount of a compound of the disclosure. In some embodiments, the percentage modulation exceeds 25%, 30%, 40%, 50%, 60%, 70%, 80%, or 90%. In some embodiments, the percentage of inhibiting exceeds 25%, 30%, 40%, 50%, 60%, 70%, 80%, or 90%.

[0116] In some embodiments, the disclosure provides methods of inhibiting KRAS, HRAS or NRAS G12C activity in a cell by contacting said cell with an amount of a compound of the disclosure sufficient to inhibit the activity of KRAS, HRAS or NRAS G12C in said cell. In some embodiments, the disclosure provides methods of inhibiting KRAS, HRAS or NRAS G12C activity in a tissue by contacting said tissue with an amount of a compound of the disclosure sufficient to inhibit the activity of KRAS, HRAS or NRAS G12C in said tissue. In some embodiments, the disclosure provides methods of inhibiting KRAS, HRAS or NRAS G12C activity in an organism by contacting said organism with an amount of a compound of the disclosure sufficient to inhibit the activity of KRAS, HRAS or NRAS G12C in said organism. In some embodiments, the disclosure provides methods of inhibiting KRAS, HRAS or NRAS G12C activity in an animal by contacting said animal with an amount of a compound of the disclosure sufficient to inhibit the activity of KRAS, HRAS or NRAS G12C in said animal. In some embodiments, the disclosure provides methods of inhibiting KRAS, HRAS or NRAS G12C activity in a mammal by contacting said mammal with an amount of a compound of the disclosure sufficient to inhibit the activity of KRAS, HRAS or NRAS G12C in said mammal. In some embodiments, the disclosure provides methods of inhibiting KRAS, HRAS or NRAS G12C activity in a human by contacting said human with an amount of a compound of the disclosure sufficient to inhibit the activity of KRAS, HRAS or NRAS G12C in said human. The present disclosure provides methods of treating a disease mediated by KRAS, HRAS or NRAS G12C activity in a subject in need of such treatment.

Combination Therapy:

[0093] The present disclosure also provides methods for combination therapies in which an agent known to modulate other pathways, or other components of the same pathway, or even overlapping sets of target enzymes are used in combination with a compound of the present disclosure, or a pharmaceutically acceptable salt thereof. In one aspect, such therapy includes but is not limited to the combination of one or more compounds of the disclosure with chemotherapeutic agents, therapeutic antibodies, and radiation treatment, to provide a synergistic or additive therapeutic effect.

[0117] Many chemotherapeutics are presently known in the art and can be used in combination with the compounds of the disclosure. In some embodiments, the chemotherapeutic is selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti- hormones, angiogenesis inhibitors, and anti-androgens. Non-limiting examples are chemotherapeutic agents, cytotoxic agents, and non-peptide small molecules such as Gleevec® (Imatinib Mesylate), Kyprolis® (carfilzomib), Velcade® (bortezomib), Casodex (bicalutamide), Iressa® (gefitinib), Venclexta™ (venetoclax) and Adriamycin™, (docorubicin) as well as a host of chemotherapeutic agents. Non-limiting examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclosphosphamide (Cytoxan™); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphaoramide and trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine, chlorocyclophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabicin, carminomycin, carzinophilin, Casodex™, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo- L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, androgens such as calusterone, dromostanolone propionate, epitiostanol, mepitiostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfomithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin;

podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxanes, e.g. paclitaxel and docetaxel; retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[0118] Also included as suitable chemotherapeutic cell conditioners are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, (Nolvadex™), raloxifene, aromatase inhibiting 4(5)- imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY 117018, onapristone, and toremifene (Fareston); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; camptothecin-11 (CPT-11); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO).

[0119] Where desired, the compounds or pharmaceutical composition of the present disclosure can be used in combination with commonly prescribed anti-cancer drugs such as Herceptin®, Avastin®, Erbitux®, Rituxan®, Taxol®, Arimidex®, Taxotere®, ABVD, AVICINE, Abagovomab, Acridine carboxamide, Adecatumumab, 17-N-Allylamino-17-demethoxygeldanamycin, Alpharadin, Alvocidib, 3-Aminopyridine-2-carboxaldehyde thiosemicarbazone, Amonafide, Anthracenedione, Anti-CD22 immunotoxins, Antineoplastic, Antitumorigenic herbs, Apaziquone, Atiprimod, Azathioprine, Belotecan, Bendamustine, BIBW 2992, Biricodar, Brostallicin, Bryostatin, Buthionine sulfoximine, CBV (chemotherapy), Calyculin, cell-cycle nonspecific antineoplastic agents, Dichloroacetic acid, Discodermolide, Elsamitrucin, Enocitabine, Epothilone, Eribulin, Everolimus, Exatecan, Exisulind, Ferruginol, Forodesine, Fosfestrol, ICE chemotherapy regimen, IT-101, Imexon, Imiquimod, Indolocarbazole, Irofulven, Laniquidar, Larotaxel, Lenalidomide, Lucanthone, Lurtotecan, Mafosfamide, Mitozolomide, Nafoxidine, Nedaplatin, Olaparib, Ortataxel, PAC-1, Pawpaw, Pixantrone, Proteasome inhibitor, Rebeccamycin, Resiquimod, Rubitecan, SN-38, Salinosporamide A, Sapacitabine, Stanford V, Swainsonine, Talaporfin, Tariquidar, Tegafur-uracil, Temodar, Tesetaxel, Triplatin tetranitrate, Tris(2-chloroethyl)amine, Troxacitabine, Uramustine, Vadimezan, Vinflunine, ZD6126 or Zosuquidar.

[0120] This disclosure further relates to a method for using the compounds or pharmaceutical compositions provided herein, in combination with radiation therapy for inhibiting abnormal cell growth or treating the hyperproliferative disorder in the mammal. Techniques for administering radiation therapy are known in the art, and these techniques can be used in the combination therapy described herein. The administration of the compound of the disclosure in this combination therapy can be determined as described herein.

[0121] Radiation therapy can be administered through one of several methods, or a combination of methods, including without limitation external-beam therapy, internal radiation therapy, implant radiation, stereotactic radiosurgery, systemic radiation therapy, radiotherapy and permanent or temporary interstitial brachytherapy. The term "brachytherapy," as used herein, refers to radiation therapy delivered by a spatially confined radioactive material inserted into the body at or near a tumor or other proliferative tissue disease site. The term is intended without limitation to include exposure to radioactive isotopes (e.g. At-211, I-131, I-125, Y-90, Re-186, Re-188, Sm- 153, Bi-212, P-32, and radioactive isotopes of Lu). Suitable radiation sources for use as a cell conditioner of the present disclosure include both solids and liquids. By way of non-limiting example, the radiation source can be a radionuclide, such as I-125, I-131, Yb-169, Ir-192 as a solid source, I-125 as a solid source, or other radionuclides that emit photons, beta particles, gamma radiation, or other therapeutic rays. The radioactive material can also be a fluid made from any solution of radionuclide(s), e.g., a solution of I-125 or I-131, or a radioactive fluid can be produced using a slurry of a suitable fluid containing small particles of solid radionuclides, such as Au-198, Y-90. Moreover, the radionuclide(s) can be embodied in a gel or radioactive micro spheres.

[0122] The compounds or pharmaceutical compositions of the disclosure can be used in combination with an amount of one or more substances selected from anti-angiogenesis agents, signal transduction inhibitors, antiproliferative agents, glycolysis inhibitors, or autophagy inhibitors.

[0123] Anti-angiogenesis agents, such as MMP-2 (matrix-metalloproteinase 2) inhibitors, MMP-9 (matrix-metalloproteinase 9) inhibitors, and COX-11 (cyclooxygenase 11) inhibitors, can be used in conjunction with a compound of the disclosure and pharmaceutical compositions described herein. Anti-angiogenesis agents include, for example, rapamycin, temsirolimus (CCI-779), everolimus (RAD001), sorafenib, sunitinib, and bevacizumab. Examples of useful COX-II inhibitors include alecoxib, valdecoxib, and rofecoxib. Examples of useful matrix metalloproteinase inhibitors are described in WO 96/33172 WO 96/27583 European Patent

Publication EP0818442, European Patent Publication EP1004578 , WO 98/07697, WO 98/03516, WO 98/34918, WO 98/34915, WO 98/33768, WO 98/30566, European Patent Publication 606046, European Patent Publication 931 788, WO 90/05719, WO 99/52910, WO 99/52889, WO 99/29667, WO1999007675, European Patent Publication EP1786785, European Patent Publication No. EP1181017, United States Publication No. US20090012085, United States Publication US5863 949, United States Publication US5861 510, and European Patent Publication EP0780386, all of which are incorporated herein in their entireties by reference. Preferred MMP-2 and MMP-9 inhibitors are those that have little or no activity inhibiting MMP-1. More preferred, are those that selectively inhibit MMP-2 and/or AMP-9 relative to the other matrix- metalloproteinases (i. e., MAP-1, MMP-3, MMP-4, MMP-5, MMP-6, MMP- 7, MMP- 8, MMP-10, MMP-11, MMP-12, andMMP-13). Some specific examples of MMP inhibitors useful in the disclosure are AG-3340, RO 32-3555, and RS 13-0830.

[0124] The present compounds may also be used in co-therapies with other anti-neoplastic agents, such as acemannan, aclarubicin, aldesleukin, alemtuzumab, alitretinoin, altretamine, amifostine, aminolevulinic acid, amrubicin, amsacrine, anagrelide, anastrozole, ANCER, ancestim, ARGLABIN, arsenic trioxide, BAM 002 (Novelos), bexarotene, bicalutamide, broxuridine, capecitabine, celmoleukin, cetrorelix, cladribine, clotrimazole, cytarabine ocfosfate, DA 3030 (Dong-A), daclizumab, denileukin diftitox, deslorelin, dexrazoxane, dilazep, docetaxel, docosanol, doxercalciferol, doxifluridine, doxorubicin, bromocriptine, carmustine, cytarabine, fluorouracil, HIT diclofenac, interferon alfa, daunorubicin, doxorubicin, tretinoin, edelfosine, edrecolomab, eflornithine, emitefur, epirubicin, epoetin beta, etoposide phosphate, exemestane, exisulind, fadrozole, filgrastim, finasteride, fludarabine phosphate, formestane, fotemustine, gallium nitrate, gemcitabine, gemtuzumab zogamicin, gimeracil/oteracil/tegafur combination, glycopine, goserelin, heptaplatin, human chorionic gonadotropin, human fetal alpha fetoprotein, ibandronic acid, idarubicin, (imiquimod, interferon alfa, interferon alfa, natural, interferon alfa-2, interferon alfa-2a, interferon alfa-2b, interferon alfa-N1, interferon alfa-$_{n3}$, interferon alfacon-1, interferon alpha, natural, interferon beta, interferon beta-1a, interferon beta-1b, interferon gamma, natural interferon gamma-1a, interferon gamma-1b, interleukin-1 beta, iobenguane, irinotecan, irsogladine, lanreotide, LC 9018 (Yakult), leflunomide, lenograstim, lentinan sulfate, letrozole, leukocyte alpha interferon, leuprorelin, levamisole + fluorouracil, liarozole, lobaplatin, lonidamine, lovastatin, masoprocol, melarsoprol, metoclopramide, mifepristone, miltefosine, mirimostim,

mismatched double stranded RNA, mitoguazone, mitolactol, mitoxantrone, molgramostim, nafarelin, naloxone + pentazocine, nartograstim, nedaplatin, nilutamide, noscapine, novel erythropoiesis stimulating protein, NSC 631570 octreotide, oprelvekin, osaterone, oxaliplatin, paclitaxel, pamidronic acid, pegaspargase, peginterferon alfa-2b, pentosan polysulfate sodium, pentostatin, picibanil, pirarubicin, rabbit antithymocyte polyclonal antibody, polyethylene glycol interferon alfa-2a, porfimer sodium, raloxifene, raltitrexed, rasburiembodiment, rhenium Re 186 etidronate, RII retinamide, rituximab, romurtide, samarium (153 Sm) lexidronam, sargramostim, sizofiran, sobuzoxane, sonermin, strontium-89 chloride, suramin, tasonermin, tazarotene, tegafur, temoporfin, temozolomide, teniposide, tetrachlorodecaoxide, thalidomide, thymalfasin, thyrotropin alfa, topotecan, toremifene, tositumomab-iodine 131, trastuzumab, treosulfan, tretinoin, trilostane, trimetrexate, triptorelin, tumor necrosis factor alpha, natural, ubenimex, bladder cancer vaccine, Maruyama vaccine, melanoma lysate vaccine, valrubicin, verteporfin, vinorelbine, VIRULIZIN, zinostatin stimalamer, or zoledronic acid; abarelix; AE 941 (Aeterna), ambamustine, antisense oligonucleotide, bcl-2 (Genta), APC 8015 (Dendreon), cetuximab, decitabine, dexaminoglutethimide, diaziquone, EL 532 (Elan), EM 800 (Endorecherche), eniluracil, etanidazole, fenretinide, filgrastim SD01 (Amgen), fulvestrant, galocitabine, gastrin 17 immunogen, HLA-B7 gene therapy (Vical), granulocyte macrophage colony stimulating factor, histamine dihydrochloride, ibritumomab tiuxetan, ilomastat, IM 862 (Cytran), interleukin-2, iproxifene, LDI 200 (Milkhaus), leridistim, lintuzumab, CA 125 MAb (Biomira), cancer MAb (Japan Pharmaceutical Development), HER-2 and Fc MAb (Medarex), idiotypic 105AD7 MAb (CRC Technology), idiotypic CEA MAb (Trilex), LYM-1-iodine 131 MAb (Techniclone), polymorphic epithelial mucin-yttrium 90 MAb (Antisoma), marimastat, menogaril, mitumomab, motexafin gadolinium, MX 6 (Galderma), nelarabine, nolatrexed, P 30 protein, pegvisomant, pemetrexed, porfiromycin, prinomastat, RL 0903 (Shire), rubitecan, satraplatin, sodium phenylacetate, sparfosic acid, SRL 172 (SR Pharma), SU 5416 (SUGEN), TA 077 (Tanabe), tetrathiomolybdate, thaliblastine, thrombopoietin, tin ethyl etiopurpurin, tirapazamine, cancer vaccine (Biomira), melanoma vaccine (New York University), melanoma vaccine (Sloan Kettering Institute), melanoma oncolysate vaccine (New York Medical College), viral melanoma cell lysates vaccine (Royal Newcastle Hospital), or valspodar.

[0125] The compounds of the invention may further be used with VEGFR inhibitors. Other compounds described in the following patents and patent applications can be used in combination therapy: US 6,258,812, US 2003/0105091, WO 01/37820, US 6,235,764, WO

01/32651, US 6,630,500, US 6,515,004, US 6,713,485, US 5,521,184, US 5,770,599, US 5,747,498, WO 02/68406, WO 02/66470, WO 02/55501, WO 04/05279, WO 04/07481, WO 04/07458, WO 04/09784, WO 02/59110, WO 99/45009, WO 00/59509, WO 99/61422, US 5,990,141, WO 00/12089, and WO 00/02871.

[0126] In some embodiments, the combination comprises a composition of the present invention in combination with at least one anti-angiogenic agent. Agents are inclusive of, but not limited to, *in vitro* synthetically prepared chemical compositions, antibodies, antigen binding regions, radionuclides, and combinations and conjugates thereof. An agent can be an agonist, antagonist, allosteric modulator, toxin or, more generally, may act to inhibit or stimulate its target (e.g., receptor or enzyme activation or inhibition), and thereby promote cell death or arrest cell growth.

[0127] Exemplary anti-angiogenic agents include ERBITUX™ (IMC-C225), KDR (kinase domain receptor) inhibitory agents (e.g., antibodies and antigen binding regions that specifically bind to the kinase domain receptor), anti-VEGF agents (e.g., antibodies or antigen binding regions that specifically bind VEGF, or soluble VEGF receptors or a ligand binding region thereof) such as AVASTIN™ or VEGF-TRAP™, and anti-VEGF receptor agents (e.g., antibodies or antigen binding regions that specifically bind thereto), EGFR inhibitory agents (e.g., antibodies or antigen binding regions that specifically bind thereto) such as Vectibix (panitumumab), IRESSA™ (gefitinib), TARCEVA™ (erlotinib), anti-Ang1 and anti-Ang2 agents (e.g., antibodies or antigen binding regions specifically binding thereto or to their receptors, e.g., Tie2/Tek), and anti-Tie2 kinase inhibitory agents (e.g., antibodies or antigen binding regions that specifically bind thereto). The pharmaceutical compositions of the present invention can also include one or more agents (e.g., antibodies, antigen binding regions, or soluble receptors) that specifically bind and inhibit the activity of growth factors, such as antagonists of hepatocyte growth factor (HGF, also known as Scatter Factor), and antibodies or antigen binding regions that specifically bind its receptor "c-met".

[0128] Other anti-angiogenic agents include Campath, IL-8, B-FGF, Tek antagonists (Ceretti et al., U.S. Publication No. 2003/0162712; U.S. Patent No. 6,413,932), anti-TWEAK agents (e.g., specifically binding antibodies or antigen binding regions, or soluble TWEAK receptor antagonists; see, Wiley, U.S. Patent No. 6,727,225), ADAM distintegrin domain to antagonize the binding of integrin to its ligands (Fanslow et al., U.S. Publication No. 2002/0042368), specifically binding anti-eph receptor and/or anti-ephrin antibodies or antigen binding regions (U.S. Patent Nos. 5,981,245; 5,728,813; 5,969,110; 6,596,852; 6,232,447;

6,057,124 and patent family members thereof), and anti-PDGF-BB antagonists (e.g., specifically binding antibodies or antigen binding regions) as well as antibodies or antigen binding regions specifically binding to PDGF-BB ligands, and PDGFR kinase inhibitory agents (e.g., antibodies or antigen binding regions that specifically bind thereto).

[0129] Additional anti-angiogenic/anti-tumor agents include: SD-7784 (Pfizer, USA); cilengitide.(Merck KGaA, Germany, EPO 770622); pegaptanib octasodium, (Gilead Sciences, USA); Alphastatin, (BioActa, UK); M-PGA, (Celgene, USA, US 5712291); ilomastat, (Arriva, USA, US 5892112); emaxanib, (Pfizer, USA, US 5792783); vatalanib, (Novartis, Switzerland); 2-methoxyestradiol, (EntreMed, USA); TLC ELL-12, (Elan, Ireland); anecortave acetate, (Alcon, USA); alpha-D148 Mab, (Amgen, USA); CEP-7055,(Cephalon, USA); anti-Vn Mab, (Crucell, Netherlands) DAC:antiangiogenic, (ConjuChem, Canada); Angiocidin, (InKine Pharmaceutical, USA); KM-2550, (Kyowa Hakko, Japan); SU-0879, (Pfizer, USA); CGP-79787, (Novartis, Switzerland, EP 970070); ARGENT technology, (Ariad, USA); YIGSR-Stealth, (Johnson & Johnson, USA); fibrinogen-E fragment, (BioActa, UK); angiogenesis inhibitor, (Trigen, UK); TBC-1635, (Encysive Pharmaceuticals, USA); SC-236, (Pfizer, USA); ABT-567, (Abbott, USA); Metastatin, (EntreMed, USA); angiogenesis inhibitor, (Tripep, Sweden); maspin, (Sosei, Japan); 2-methoxyestradiol, (Oncology Sciences Corporation, USA); ER-68203-00, (IVAX, USA); Benefin, (Lane Labs, USA); Tz-93, (Tsumura, Japan); TAN-1120, (Takeda, Japan); FR-111142, (Fujisawa, Japan, JP 02233610); platelet factor 4, (RepliGen, USA, EP 407122); vascular endothelial growth factor antagonist, (Borean, Denmark); bevacizumab (pINN), (Genentech, USA); angiogenesis inhibitors, (SUGEN, USA); XL 784, (Exelixis, USA); XL 647, (Exelixis, USA); MAb, alpha5beta3 integrin, second generation, (Applied Molecular Evolution, USA and MedImmune, USA); gene therapy, retinopathy, (Oxford BioMedica, UK); enzastaurin hydrochloride (USAN), (Lilly, USA); CEP 7055, (Cephalon, USA and Sanofi-Synthelabo, France); BC 1, (Genoa Institute of Cancer Research, Italy); angiogenesis inhibitor, (Alchemia, Australia); VEGF antagonist, (Regeneron, USA); rBPI 21 and BPI-derived antiangiogenic, (XOMA, USA); PI 88, (Progen, Australia); cilengitide (pINN), (Merck KGaA, German; Munich Technical University, Germany, Scripps Clinic and Research Foundation, USA); cetuximab (INN), (Aventis, France); AVE 8062, (Ajinomoto, Japan); AS 1404, (Cancer Research Laboratory, New Zealand); SG 292, (Telios, USA); Endostatin, (Boston Childrens Hospital, USA); ATN 161, (Attenuon, USA); ANGIOSTATIN, (Boston Childrens Hospital, USA); 2-methoxyestradiol, (Boston Childrens Hospital, USA); ZD 6474, (AstraZeneca, UK); ZD 6126, (Angiogene Pharmaceuticals, UK);

PPI 2458, (Praecis, USA); AZD 9935, (AstraZeneca, UK); AZD 2171, (AstraZeneca, UK); vatalanib (pINN), (Novartis, Switzerland and Schering AG, Germany); tissue factor pathway inhibitors, (EntreMed, USA); pegaptanib (Pinn), (Gilead Sciences, USA); xanthorrhizol, (Yonsei University, South Korea); vaccine, gene-based, VEGF-2, (Scripps Clinic and Research Foundation, USA); SPV5.2, (Supratek, Canada); SDX 103, (University of California at San Diego, USA); PX 478, (ProlX, USA); METASTATIN, (EntreMed, USA); troponin I, (Harvard University, USA); SU 6668, (SUGEN, USA); OXI 4503, (OXiGENE, USA); o-guanidines, (Dimensional Pharmaceuticals, USA); motuporamine C, (British Columbia University, Canada); CDP 791, (Celltech Group, UK); atiprimod (pINN), (GlaxoSmithKline, UK); E 7820, (Eisai, Japan); CYC 381, (Harvard University, USA); AE 941, (Aeterna, Canada); vaccine, angiogenesis, (EntreMed, USA); urokinase plasminogen activator inhibitor, (Dendreon, USA); oglufanide (pINN), (Melmotte, USA); HIF-1alfa inhibitors, (Xenova, UK); CEP 5214, (Cephalon, USA); BAY RES 2622, (Bayer, Germany); Angiocidin, (InKine, USA); A6, (Angstrom, USA); KR 31372, (Korea Research Institute of Chemical Technology, South Korea); GW 2286, (GlaxoSmithKline, UK); EHT 0101, (ExonHit, France); CP 868596, (Pfizer, USA); CP 564959, (OSI, USA); CP 547632, (Pfizer, USA); 786034, (GlaxoSmithKline, UK); KRN 633, (Kirin Brewery, Japan); drug delivery system, intraocular, 2-methoxyestradiol, (EntreMed, USA); anginex, (Maastricht University, Netherlands, and Minnesota University, USA); ABT 510, (Abbott, USA); AAL 993, (Novartis, Switzerland); VEGI, (ProteomTech, USA); tumor necrosis factor-alpha inhibitors, (National Institute on Aging, USA); SU 11248, (Pfizer, USA and SUGEN USA); ABT 518, (Abbott, USA); YH16, (Yantai Rongchang, China); S-3APG , (Boston Childrens Hospital, USA and EntreMed, USA); MAb, KDR, (ImClone Systems, USA); MAb, alpha5 beta1, (Protein Design, USA); KDR kinase inhibitor, (Celltech Group, UK, and Johnson & Johnson, USA); GFB 116, (South Florida University, USA and Yale University, USA); CS 706, (Sankyo, Japan); combretastatin A4 prodrug, (Arizona State University, USA); chondroitinase AC, (IBEX, Canada); BAY RES 2690, (Bayer, Germany); AGM 1470, (Harvard University, USA, Takeda, Japan, and TAP, USA); AG 13925, (Agouron, USA); Tetrathiomolybdate, (University of Michigan, USA); GCS 100, (Wayne State University, USA) CV 247, (Ivy Medical, UK); CKD 732, (Chong Kun Dang, South Korea); MAb, vascular endothelium growth factor, (Xenova, UK); irsogladine (INN), (Nippon Shinyaku, Japan); RG 13577, (Aventis, France); WX 360, (Wilex, Germany); squalamine (pINN), (Genaera, USA); RPI 4610, (Sirna, USA); cancer therapy, (Marinova, Australia); heparanase inhibitors, (InSight, Israel); KL 3106, (Kolon, South Korea); Honokiol, (Emory University, USA); ZK CDK, (Schering AG, Germany); ZK Angio, (Schering AG,

Germany); ZK 229561, (Novartis, Switzerland, and Schering AG, Germany); XMP 300, (XOMA, USA); VGA 1102, (Taisho, Japan); VEGF receptor modulators, (Pharmacopeia, USA); VE-cadherin-2 antagonists , (ImClone Systems, USA); Vasostatin, (National Institutes of Health, USA);vaccine, Flk-1, (ImClone Systems, USA); TZ 93, (Tsumura, Japan); TumStatin, (Beth Israel Hospital, USA); truncated soluble FLT 1 (vascular endothelial growth factor receptor 1), (Merck & Co, USA); Tie-2 ligands, (Regeneron, USA); and, thrombospondin 1 inhibitor, (Allegheny Health, Education and Research Foundation, USA).

[0130] Autophagy inhibitors include, but are not limited to chloroquine, 3- methyladenine, hydroxychloroquine (Plaquenil™), bafilomycin A1, 5-amino-4- imidazole carboxamide riboside (AICAR), okadaic acid, autophagy-suppressive algal toxins which inhibit protein phosphatases of type 2A or type 1, analogues of cAMP, and drugs which elevate cAMP levels such as adenosine, LY204002, N6-mercaptopurine riboside, and vinblastine. In addition, antisense or siRNA that inhibits expression of proteins including but not limited to ATG5 (which are implicated in autophagy), may also be used.

[0131] Additional pharmaceutically active compounds/agents that can be used in the treatment of cancers and that can be used in combination with one or more compound of the present invention include: epoetin alfa; darbepoetin alfa; panitumumab; pegfilgrastim; palifermin; filgrastim; denosumab; ancestim; AMG 102; AMG 176; AMG 386; AMG 479; AMG 655; AMG 745; AMG 951; and AMG 706, or a pharmaceutically acceptable salt thereof.

[0132] In certain embodiments, a composition provided herein is conjointly administered with a chemotherapeutic agent. Suitable chemotherapeutic agents may include, natural products such as vinca alkaloids (e.g., vinblastine, vincristine, and vinorelbine), paclitaxel, epidipodophyllotoxins (e.g., etoposide and teniposide), antibiotics (e.g., dactinomycin (actinomycin D), daunorubicin, doxorubicin, and idarubicin), anthracyclines, mitoxantrone, bleomycins, plicamycin (mithramycin), mitomycin, enzymes (e.g., L-asparaginase which systemically metabolizes L-asparagine and deprives cells which do not have the capacity to synthesize their own asparagine), antiplatelet agents, antiproliferative/antimitotic alkylating agents such as nitrogen mustards (e.g., mechlorethamine, cyclophosphamide and analogs, melphalan, and chlorambucil), ethylenimines and methylmelamines (e.g., hexaamethylmelaamine and thiotepa), CDK inhibitors (e.g., seliciclib, UCN-01, P1446A-05, PD-0332991, dinaciclib, P27-00, AT-7519, RGB286638, and SCH727965), alkyl sulfonates (e.g., busulfan), nitrosoureas (e.g., carmustine (BCNU) and analogs, and streptozocin), trazenes-dacarbazinine (DTIC), antiproliferative/antimitotic antimetabolites such as folic acid

analogs (e.g., methotrexate), pyrimidine analogs (e.g., fluorouracil, floxuridine, and cytarabine), purine analogs and related inhibitors (e.g., mercaptopurine, thioguanine, pentostatin and 2-chlorodeoxyadenosine), aromatase inhibitors (e.g., anastrozole, exemestane, and letrozole), and platinum coordination complexes (e.g., cisplatin and carboplatin), procarbazine, hydroxyurea, mitotane, aminoglutethimide, histone deacetylase (HDAC) inhibitors (e.g., trichostatin, sodium butyrate, apicidan, suberoyl anilide hydroamic acid, vorinostat, LBH 589, romidepsin, ACY-1215, and panobinostat), mTor inhibitors (e.g., temsirolimus, everolimus, ridaforolimus, and sirolimus), KSP(Eg5) inhibitors (e.g., Array 520), DNA binding agents (e.g., Zalypsis), PI3K delta inhibitor (e.g., GS-1101 and TGR-1202), PI3K delta and gamma inhibitor (e.g., CAL-130), multi-kinase inhibitor (e.g., TG02 and sorafenib), hormones (e.g., estrogen) and hormone agonists such as leutinizing hormone releasing hormone (LHRH) agonists (e.g., goserelin, leuprolide and triptorelin), BAFF-neutralizing antibody (e.g., LY2127399), IKK inhibitors, p38MAPK inhibitors, anti-IL-6 (e.g., CNTO328), telomerase inhibitors (e.g., GRN 163L), aurora kinase inhibitors (e.g., MLN8237), cell surface monoclonal antibodies (e.g., anti-CD38 (HUMAX-CD38), anti-CS1 (e.g., elotuzumab), HSP90 inhibitors (e.g., 17 AAG and KOS 953), P13K / Akt inhibitors (e.g., perifosine), Akt inhibitor (e.g., GSK-2141795), PKC inhibitors (e.g., enzastaurin), FTIs (e.g., Zarnestra™), anti-CD138 (e.g., BT062), Torc1/2 specific kinase inhibitor (e.g., INK128), kinase inhibitor (e.g., GS-1101), ER/UPR targeting agent (e.g., MKC-3946), cFMS inhibitor (e.g., ARRY-382), JAK1/2 inhibitor (e.g., CYT387), PARP inhibitor (e.g., olaparib and veliparib (ABT-888)), BCL-2 antagonist. Other chemotherapeutic agents may include mechlorethamine, camptothecin, ifosfamide, tamoxifen, raloxifene, gemcitabine, navelbine, sorafenib, or any analog or derivative variant of the foregoing.

[0133] The compounds of the present invention may also be used in combination with radiation therapy, hormone therapy, surgery and immunotherapy, which therapies are well known to those skilled in the art.

[0134] In certain embodiments, a pharmaceutical composition provided herein is conjointly administered with a steroid. Suitable steroids may include, but are not limited to, 21-acetoxypregnenolone, alclometasone, algestone, amcinonide, beclomethasone, betamethasone, budesonide, chloroprednisone, clobetasol, clocortolone, cloprednol, corticosterone, cortisone, cortivazol, deflazacort, desonide, desoximetasone, dexamethasone, diflorasone, diflucortolone, difuprednate, enoxolone, fluazacort, flucloronide, flumethasone, flunisolide, fluocinolone acetonide, fluocinonide, fluocortin butyl, fluocortolone, fluorometholone, fluperolone acetate,

fluprednidene acetate, fluprednisolone, flurandrenolide, fluticasone propionate, formocortal, halcinonide, halobetasol propionate, halometasone, hydrocortisone, loteprednol etabonate, mazipredone, medrysone, meprednisone, methylprednisolone, mometasone furoate, paramethasone, prednicarbate, prednisolone, prednisolone 25-diethylaminoacetate, prednisolone sodium phosphate, prednisone, prednival, prednylidene, rimexolone, tixocortol, triamcinolone, triamcinolone acetonide, triamcinolone benetonide, triamcinolone hexacetonide, and salts and/or derivatives thereof.In a particular embodiment, the compounds of the present invention can also be used in combination with additional pharmaceutically active agents that treat nausea. Examples of agents that can be used to treat nausea include: dronabinol; granisetron; metoclopramide; ondansetron; and prochlorperazine; or a pharmaceutically acceptable salt thereof.

[0135] The compounds of the present invention may also be used in combination with an additional pharmaceutically active compound that disrupts or inhibits RAS-RAF-ERK or PI3K-AKT-TOR signaling pathways. In other such combinations, the additional pharmaceutically active compound is a PD-1 and PD-L1 antagonist. The compounds or pharmaceutical compositions of the disclosure can also be used in combination with an amount of one or more substances selected from EGFR inhibitors, MEK inhibitors, PI3K inhibitors, AKT inhibitors, TOR inhibitors, Mcl-1 inhibitors, BCL-2 inhibitors, SHP2 inhibitors, proteasome inhibitors, and immune therapies, including monoclonal antibodies, immunomodulatory imides (IMiDs), anti-PD-1, anti-PDL-1, anti-CTLA4, anti-LAG1, and anti-OX40 agents, GITR agonists, CAR-T cells, and BiTEs.

[0136] EGFR inhibitors include, but are not limited to, small molecule antagonists, antibody inhibitors, or specific antisense nucleotide or siRNA. Useful antibody inhibitors of EGFR include cetuximab (Erbitux), panitumumab (Vectibix), zalutumumab, nimotuzumab, and matuzumab. Small molecule antagonists of EGFR include gefitinib, erlotinib (Tarceva), and most recently, lapatinib (TykerB). See e.g., Yan L, et. al., *Pharmacogenetics and Pharmacogenomics In Oncology Therapeutic Antibody Development*, BioTechniques 2005; 39(4): 565-8, and Paez J G, et. al., *EGFR Mutations In Lung Cancer Correlation With Clinical Response To Gefitinib Therapy*, Science 2004; 304(5676): 1497-500.

[0137] Non-limiting examples of small molecule EGFR inhibitors include any of the EGFR inhibitors described in the following patent publications, and all pharmaceutically acceptable salts and solvates of said EGFR inhibitors: European Patent Application EP 520722, published Dec. 30, 1992; European Patent Application EP 566226, published Oct. 20, 1993; PCT

International Publication WO 96/33980, published Oct. 31, 1996; U.S. Pat. No. 5,747,498, issued May 5, 1998; PCT International Publication WO 96/30347, published Oct. 3, 1996; European Patent Application EP 787772, published Aug. 6, 1997; PCT International Publication WO 97/30034, published Aug. 21, 1997; PCT International Publication WO 97/30044, published Aug. 21, 1997; PCT International Publication WO 97/38994, published Oct. 23, 1997; PCT International Publication WO 97/49688, published Dec. 31, 1997; European Patent Application EP 837063, published Apr. 22, 1998; PCT International Publication WO 98/02434, published Jan. 22, 1998; PCT International Publication WO 97/38983, published Oct. 23, 1997; PCT International Publication WO 95/19774, published Jul. 27, 1995; PCT International Publication WO 95/19970, published Jul. 27, 1995; PCT International Publication WO 97/13771, published Apr. 17, 1997; PCT International Publication WO 98/02437, published Jan. 22, 1998; PCT International Publication WO 98/02438, published Jan. 22, 1998; PCT International Publication WO 97/32881, published Sep. 12, 1997; German Application DE 19629652, published Jan. 29, 1998; PCT International Publication WO 98/33798, published Aug. 6, 1998; PCT International Publication WO 97/32880, published Sep. 12, 1997; PCT International Publication WO 97/32880 published Sep. 12, 1997; European Patent Application EP 682027, published Nov. 15, 1995; PCT International Publication WO 97/02266, published Jan. 23, 197; PCT International Publication WO 97/27199, published Jul. 31, 1997; PCT International Publication WO 98/07726, published Feb. 26, 1998; PCT International Publication WO 97/34895, published Sep. 25, 1997; PCT International Publication WO 96/31510′, published Oct. 10, 1996; PCT International Publication WO 98/14449, published Apr. 9, 1998; PCT International Publication WO 98/14450, published Apr. 9, 1998; PCT International Publication WO 98/14451, published Apr. 9, 1998; PCT International Publication WO 95/09847, published Apr. 13, 1995; PCT International Publication WO 97/19065, published May 29, 1997; PCT International Publication WO 98/17662, published Apr. 30, 1998; U.S. Pat. No. 5,789,427, issued Aug. 4, 1998; U.S. Pat. No. 5,650,415, issued Jul. 22, 1997; U.S. Pat. No. 5,656,643, issued Aug. 12, 1997; PCT International Publication WO 99/35146, published Jul. 15, 1999; PCT International Publication WO 99/35132, published Jul. 15, 1999; PCT International Publication WO 99/07701, published Feb. 18, 1999; and PCT International Publication WO 92/20642 published Nov. 26, 1992. Additional non-limiting examples of small molecule EGFR inhibitors include any of the EGFR inhibitors described in Traxler, P., 1998, Exp. Opin. Ther. Patents 8(12):1599-1625.

[0138] Antibody-based EGFR inhibitors include any anti-EGFR antibody or antibody fragment that can partially or completely block EGFR activation by its natural ligand. Non-limiting examples of antibody-based EGFR inhibitors include those described in Modjtahedi, H., et al., 1993, Br. J. Cancer 67:247-253; Teramoto, T., et al., 1996, Cancer 77:639-645; Goldstein et al., 1995, Clin. Cancer Res. 1:1311-1318; Huang, S. M., et al., 1999, Cancer Res. 15:59(8):1935-40; and Yang, X., et al., 1999, Cancer Res. 59:1236-1243. Thus, the EGFR inhibitor can be monoclonal antibody Mab E7.6.3 (Yang, 1999 supra), or Mab C225 (ATCC Accession No. HB-8508), or an antibody or antibody fragment having the binding specificity thereof.

[0139] MEK inhibitors include, but are not limited to, tremetinib, CI-1040, AZD6244, PD318088, PD98059, PD334581, RDEA119, ARRY-142886, ARRY-438162, and PD-325901.

[0140] PI3K inhibitors include, but are not limited to, wortmannin, 17-hydroxywortmannin analogs described in WO 06/044453, 4-[2-(1H-Indazol-4-yl)-6-[[4-(methylsulfonyl)piperazin-1-yl]methyl]thieno[3,2-d]pyrimidin-4-yl]morpholine (also known as GDC 0941 and described in PCT Publication Nos. WO 09/036,082 and WO 09/055,730), 2-Methyl-2-[4-[3-methyl-2-oxo-8-(quinolin-3-yl)-2,3-dihydroimidazo[4,5-c]quinolin-1-yl]phenyl]propionitrile (also known as BEZ 235 or NVP-BEZ 235, and described in PCT Publication No. WO 06/122806), (S)-1-(4-((2-(2-aminopyrimidin-5-yl)-7-methyl-4-morpholinothieno[3,2-d]pyrimidin-6-yl)methyl)piperazin-1-yl)-2-hydroxypropan-1-one (described in PCT Publication No. WO 2008/070740), LY294002 (2-(4-Morpholinyl)-8-phenyl-4H-1-benzopyran-4-one available from Axon Medchem), PI 103 hydrochloride (3-[4-(4-morpholinylpyrido-[3′,2′:4,5]furo[3,2-d]pyrimidin-2-yl]phenol hydrochloride available from Axon Medchem), PIK 75 (N′-[(1E)-(6-bromoimidazo[1,2-a]pyridin-3-yl)methylene]-N,2-dimethyl-5-nitrobenzenesulfono-hydrazide hydrochloride available from Axon Medchem), PIK 90 (N-(7,8-dimethoxy-2,3-dihydro-imidazo[1,2-c]quinazolin-5-yl)-nicotinamide available from Axon Medchem), GDC-0941 bismesylate (2-(1H-Indazol-4-yl)-6-(4-methanesulfonyl-piperazin-1-ylmethyl)-4-morpholin-4-yl-thieno[3,2-d]pyrimidine bismesylate available from Axon Medchem), AS-252424 (5-[1-[5-(4-Fluoro-2-hydroxy-phenyl)-furan-2-yl]-meth-(Z)-ylidene]-thiazolidine-2,4-dione available from Axon Medchem), and TGX-221 (7-Methyl-2-(4-morpholinyl)-9-[1-(phenylamino)ethyl]-4H-pyrido-[1,2-a]pyrimidin-4-one available from Axon Medchem), XL-765, and XL-147. Other PI3K inhibitors include demethoxyviridin, perifosine, CAL101, PX-866, BEZ235, SF1126, INK1117, IPI-145, BKM120, XL147, XL765, Palomid 529,

GSK1059615, ZSTK474, PWT33597, IC87114, TG100-115, CAL263, PI-103, GNE-477, CUDC-907, and AEZS-136.

[0141] AKT inhibitors include, but are not limited to, Akt-1-1 (inhibits Akt1) (Barnett et al. (2005) *Biochem. J.*, 385 (Pt. 2), 399-408); Akt-1-1,2 (inhibits Ak1 and 2) (Barnett et al. (2005) *Biochem. J.* 385 (Pt. 2), 399-408); API-59CJ-Ome (e.g., Jin et al. (2004) *Br. J. Cancer* 91, 1808-12); 1-H-imidazo[4,5-c]pyridinyl compounds (e.g., WO05011700); indole-3-carbinol and derivatives thereof (e.g., U.S. Pat. No. 6,656,963; Sarkar and Li (2004) *J Nutr.* 134(12 Suppl), 3493S-3498S); perifosine (e.g., interferes with Akt membrane localization; Dasmahapatra et al. (2004) *Clin. Cancer Res.* 10(15), 5242-52, 2004); phosphatidylinositol ether lipid analogues (e.g., Gills and Dennis (2004) *Expert. Opin. Investig. Drugs* 13, 787-97); and triciribine (TCN or API-2 or NCI identifier: NSC 154020; Yang et al. (2004) *Cancer Res.* 64, 4394-9).

[0142] TOR inhibitors include, but are not limited to, AP-23573, CCI-779, everolimus, RAD-001, rapamycin, temsirolimus, ATP-competitive TORC1/TORC2 inhibitors, including PI-103, PP242, PP30 and Torin 1. Other TOR inhibitors in FKBP12 enhancer; rapamycins and derivatives thereof, including: CCI-779 (temsirolimus), RAD001 (Everolimus; WO 9409010) and AP23573; rapalogs, e.g. as disclosed in WO 98/02441 and WO 01/14387, e.g. AP23573, AP23464, or AP23841; 40-(2-hydroxyethyl)rapamycin, 40-[3-hydroxy(hydroxymethyl)methylpropanoate]-rapamycin (also called CC1779), 40-epi-(tetrazolyt)-rapamycin (also called ABT578), 32-deoxorapamycin, 16-pentynyloxy-32(S)-dihydrorapanycin, and other derivatives disclosed in WO 05005434; derivatives disclosed in U.S. Pat. No. 5,258,389, WO 94/090101, WO 92/05179, U.S. Pat. No. 5,118,677, U.S. Pat. No. 5,118,678, U.S. Pat. No. 5,100,883, U.S. Pat. No. 5,151,413, U.S. Pat. No. 5,120,842, WO 93/111130, WO 94/02136, WO 94/02485, WO 95/14023, WO 94/02136, WO 95/16691, WO 96/41807, WO 96/41807 and U.S. Pat. No. 5,256,790; phosphorus-containing rapamycin derivatives (e.g., WO 05016252); 4H-1-benzopyran-4-one derivatives (e.g., U.S. Provisional Application No. 60/528,340).

[0143] MCl-1 inhibitors include, but are not limited to, AMG-176, AMG-397, MIK665, and S63845. The myeloid cell leukemia-1 (MCL-1) protein is one of the key anti-apoptotic members of the B-cell lymphoma-2 (BCL-2) protein family. Over-expression of MCL-1 has been closely related to tumor progression as well as to resistance, not only to traditional chemotherapies but also to targeted therapeutics including BCL-2 inhibitors such as ABT-263.

[0144] SHP inhibitors include, but are not limited to, SHP099.

[0145] Proteasome inhibitors include, but are not limited to, Kyprolis®(carfilzomib), Velcade®(bortezomib), and oprozomib.

[0146] Immune therapies include, but are not limited to, anti-PD-1 agents, anti-PDL-1 agents, anti-CTLA-4 agents, anti-LAG1 agents, and anti-OX40 agents.

[0147] Monoclonal antibodies include, but are not limited to, Darzalex® (daratumumab), Herceptin® (trastuzumab), Avastin® (bevacizumab), Rituxan® (rituximab), Lucentis® (ranibizumab), and Eylea® (aflibercept).

[0148] Immunomodulatory agents (IMiDs) are a class of immunomodulatory drugs (drugs that adjust immune responses) containing an imide group. The IMiD class includes thalidomide and its analogues (lenalidomide, pomalidomide, and apremilast).

[0149] Exemplary anti-PD-1 antibodies and methods for their use are described by Goldberg et al., *Blood* 110(1):186-192 (2007), Thompson et al., *Clin. Cancer Res.* 13(6):1757-1761 (2007), and Korman et al., International Application No. PCT/JP2006/309606 (publication no. WO 2006/121168 A1), each of which are expressly incorporated by reference herein, include: Keytruda®(pembrolizumab), Opdivo®(niolumab), Yervoy™ (ipilimumab) or Tremelimumab (to CTLA-4), galiximab (to B7.1), BMS-936558 (to PD-1), MK-3475 (to PD-1), AMP224 (to B7DC), BMS-936559 (to B7-H1), MPDL3280A (to B7-H1), MEDI-570 (to ICOS), AMG557 (to B7H2), MGA271 (to B7H3), IMP321 (to LAG-3), BMS-663513 (to CD137), PF-05082566 (to CD137), CDX-1127 (to CD27), anti-OX40 (Providence Health Services), huMAbOX40L (to OX40L), Atacicept (to TACI), CP-870893 (to CD40), Lucatumumab (to CD40), Dacetuzumab (to CD40), Muromonab-CD3 (to CD3), Ipilumumab (to CTLA-4). Immune therapies also include genetically engineered T-cells (e.g., CAR-T cells) and bispecific antibodies (e.g., BiTEs).

[0150] In a particular embodiment, the compounds of the present invention are used in combination with an anti-PD-1 antibody, such as AMG 404. In a specific embodiment, the anti-PD-1 antibody (or antigen binding antibody fragment thereof) comprises 1, 2, 3, 4, 5, or all 6 the CDR amino acid sequences of SEQ ID NOs: 1-6 (representing HC CDR1, HC CDR2, HC CDR3, LC CDR1, LC CDR2, and LC CDR3, in that order). In specific embodiments, the anti-PD-1 antibody (or antigen binding antibody fragment thereof) comprises all 6 of the CDR amino acid sequences of SEQ ID NOs: 1-6. In other embodiments, the anti-PD-1 antibody (or antigen binding antibody fragment thereof) comprises (a) the heavy chain variable region

amino acid sequence in SEQ ID NO: 7, or a variant sequence thereof which differs by only one or two amino acids or which has at least or about 70% sequence identity, or (b) the light chain variable region amino acid sequence in SEQ ID NO: 8 or a variant sequence thereof which differs by only one or two amino acids or which has at least or about 70% sequence identity. In an exemplary embodiment, the anti-PD-1 antibody (or antigen binding antibody fragment thereof) comprises the heavy chain variable region amino acid sequence in SEQ ID NO: 7 and the light chain variable region amino acid sequence in SEQ ID NO: 8. In other embodiments, the anti-PD-1 antibody (or antigen binding antibody fragment thereof) comprises (a) the heavy chain amino acid sequence of SEQ ID NO: 9 or a variant sequence thereof which differs by only one or two amino acids or which has at least or about 70% sequence identity; or (b) the light chain amino acid sequence of SEQ ID NO: 10 or a variant sequence thereof which differs by only one or two amino acids or which has at least or about 70% sequence identity. In an exemplary embodiment, the anti-PD-1 antibody (or antigen binding antibody fragment thereof) comprises the heavy chain amino acid sequence of SEQ ID NO: 9 and the light chain amino acid sequence of SEQ ID NO: 10.

[0151] The present disclosure further provides nucleic acid sequences encoding the anti-PD-1 antibody (or an antigen binding portion thereof). In exemplary aspects, the antibody comprises 1, 2, 3, 4, 5, or all 6 CDRs encoded by the nucleic acid(s) of SEQ ID NOs: 11-16 (representing HC CDR1, HC CDR2, HC CDR3, LC CDR1, LC CDR2, and LC CDR3, in that order). In another exemplary aspect, the antibody comprises all 6 CDRs encoded by the nucleic acids of SEQ ID NOs: 11-16. In some embodiments, the anti-PD-1 antibody (or an antigen binding portion thereof) comprises (a) a heavy chain variable region encoded by SEQ ID NO: 17 or a variant sequence thereof which differs by only 1, 2, 3, 4, 5, or 6 nucleic acids or which has at least or about 70%, 85%, 90%, or 95% sequence identity, or (b) a light chain variable region encoded by SEQ ID NO: 18 or a variant sequence thereof which differs by only 1, 2, 3, 4, 5, or 6 nucleic acids or which has at least or about 70%, 85%, 90%, or 95% sequence identity. In an exemplary embodiment, the anti-PD-1 antibody (or an antigen binding portion thereof) comprises a heavy chain variable region encoded by SEQ ID NO: 17 and a light chain variable region encoded by SEQ ID NO: 18. In other embodiments, the anti-PD-1 antibody (or an antigen binding portion thereof) comprises (a) a heavy chain encoded by SEQ ID NO: 19 or a variant sequence thereof which differs by only 1, 2, 3, 4, 5, or 6 nucleic acids or which has at least or about 70%, 85%, 90%, or 95% sequence identity, or (b) a light chain encoded by SEQ ID NO: 20 or a variant sequence thereof which differs by only 1, 2, 3, 4, 5, or 6 nucleic acids or which has at least or about 70%, 85%, 90%, or 95% sequence identity. In an exemplary

embodiment, the anti-PD-1 antibody (or an antigen binding portion thereof) comprises a heavy chain encoded by SEQ ID NO: 19 and a light chain encoded by SEQ ID NO: 20.

[0152] GITR agonists include, but are not limited to, GITR fusion proteins and anti-GITR antibodies (e.g., bivalent anti-GITR antibodies), such as, a GITR fusion protein described in U.S. Pat. No. 6,111,090box.c, European Patent No.: 090505B1, U.S. Pat. No. 8,586,023, PCT Publication Nos.: WO 2010/003118 and 2011/090754, or an anti-GITR antibody described, e.g., in U.S. Pat. No. 7,025,962, European Patent No.: 1947183B1, U.S. Pat. No. 7,812,135, U.S. Pat. No. 8,388,967, U.S. Pat. No. 8,591,886, European Patent No.: EP 1866339, PCT Publication No.: WO 2011/028683, PCT Publication No.: WO 2013/039954, PCT Publication No.: WO2005/007190, PCT Publication No.: WO 2007/133822, PCT Publication No.: WO2005/055808, PCT Publication No.: WO 99/40196, PCT Publication No.: WO 2001/03720, PCT Publication No.: WO99/20758, PCT Publication No.: WO2006/083289, PCT Publication No.: WO 2005/115451, U.S. Pat. No. 7,618,632, and PCT Publication No.: WO 2011/051726.

[0153] The compounds described herein can be used in combination with the agents disclosed herein or other suitable agents, depending on the condition being treated. Hence, in some embodiments the one or more compounds of the disclosure will be co-administered with other agents as described above. When used in combination therapy, the compounds described herein are administered with the second agent simultaneously or separately. This administration in combination can include simultaneous administration of the two agents in the same dosage form, simultaneous administration in separate dosage forms, and separate administration. That is, a compound described herein and any of the agents described above can be formulated together in the same dosage form and administered simultaneously. Alternatively, a compound of the disclosure and any of the agents described above can be simultaneously administered, wherein both the agents are present in separate formulations. In another alternative, a compound of the present disclosure can be administered just followed by and any of the agents described above, or vice versa. In some embodiments of the separate administration protocol, a compound of the disclosure and any of the agents described above are administered a few minutes apart, or a few hours apart, or a few days apart.

[0154] As one aspect of the present invention contemplates the treatment of the disease/conditions with a combination of pharmaceutically active compounds that may be administered separately, the invention further relates to combining separate pharmaceutical compositions in kit form. The kit comprises two separate pharmaceutical compositions: a

compound of the present invention, and a second pharmaceutical compound. The kit comprises a container for containing the separate compositions such as a divided bottle or a divided foil packet. Additional examples of containers include syringes, boxes, and bags. In some embodiments, the kit comprises directions for the use of the separate components. The kit form is particularly advantageous when the separate components are preferably administered in different dosage forms (e.g., oral and parenteral), are administered at different dosage intervals, or when titration of the individual components of the combination is desired by the prescribing health care professional.

EXAMPLES

Section 1—Methods and Method Examples

Method 55

6-Chloro-7-(2-fluoro-6-hydroxyphenyl)-1-(3-methyl-5-(2-propanyl)-1,2-oxazol-4-yl)-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-piperazinyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one.



[0155] Step 1: 2,5,6-Trichloro-*N*-((5-isopropyl-3-methylisoxazol-4-yl)carbamoyl)nicotinamide. To a grey heterogeneous mixture of 2,5,6-trichloronicotinamide (**Intermediate P**, 2.5 g, 11 mmol) in THF (22 mL) was added oxalyl chloride, 2 M solution in DCM (5.8 mL, 11.6 mmol) at rt. The resulting yellow heterogeneous mixture was stirred and heated at 65 °C. After 4 h, the mixture was cooled to 0 °C and treated with a white suspension of 5-isopropyl-3-methylisoxazol-4-amine hydrochloride (1.96 g, 11.08 mmol, Enamine, Monmouth Junction, NJ, USA) and DIPEA (3.9 mL, 22.2 mmol) in THF (5 mL) and the mixture was stirred at 0 °C. After 5 min, the cooling bath was removed and the mixture was stirred at rt for 2 h. The mixture was concentrated *in vacuo* to give the crude product as yellow syrup. The residue was partitioned between EtOAc (100 mL) and saturated $NaHCO_3$ (100 mL) and the organic extract was washed with brine (1 × 100 mL) and dried over Na_2SO_4. The solution was filtered and concentrated *in vacuo* and the resulting residue was suspended in acetonitrile (20 mL), filtered and the solid was washed with acetonitrile (20 mL), and dried to give 2,5,6-trichloro-*N*-((5-isopropyl-3-methylisoxazol-4-yl)carbamoyl)nicotinamide (2.84 g, 7.26 mmol, 65.5 % yield) as an off-white solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 11.31 (br s, 1 H), 9.22 (br s, 1 H), 8.62 (s, 1 H), 3.02 - 3.16 (m, 1 H), 2.11 (s, 3 H), 1.23 (d, *J*=7.0 Hz, 6 H). *m/z* (ESI, +ve ion): 390.8 $(M+H)^+$.

[0156] Step 2: 6,7-Dichloro-1-(5-isopropyl-3-methylisoxazol-4-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a cooled mixture of 2,5,6-trichloro-*N*-((5-isopropyl-3-methylisoxazol-4-yl)carbamoyl)nicotinamide (2.84 g, 7.24 mmol) in THF (24 mL) at 0 °C was added dropwise KHMDS, 1 M solution in THF (14.5 mL, 14.5 mmol) and the mixture was stirred at 0 °C. After 30 min, the cooling bath was removed and the reddish brown homogeneous mixture was stirred at rt for 1 h. The mixture was quenched with satd. ammonium chloride (50 mL) and brine (50 mL) and extracted with EtOAc (2 × 50 mL). The organic extract was dried over Na_2SO_4 and the solution was filtered and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0–50% EtOAc/heptane) to provide 6,7-dichloro-1-(5-isopropyl-3-methylisoxazol-4-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (1.68 g, 66 % yield). This material was used without further purification in the following step. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 12.24 (s, 1 H), 8.56 (s, 1 H), 3.00-3.13 (m, 1 H), 2.04 (s, 3 H), 1.15 (dd, *J*=6.9, 4.3 Hz, 6 H). *m/z* (ESI, +ve ion): 355.0 $(M+H)^+$.

[0157] Step 3: *tert*-Butyl (*S*)-4-(6,7-dichloro-1-(5-isopropyl-3-methylisoxazol-4-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate. A solution of 6,7-dichloro-1-(5-isopropyl-3-methylisoxazol-4-yl)pyrido[2,3-*d*]pyrimidine-

2,4(1*H*,3*H*)-dione (1.69 g, 4.74 mmol), DIPEA (1.1 mL, 6.2 mmol), and phosphoryl trichloride (0.53 mL, 5.7 mmol) in acetonitrile (2 mL) was stirred at 80 °C for 1 h. The reaction mixture cooled to rt, DIPEA (3.4 mL, 19.4 mmol) and *tert*-butyl (*S*)-3-methylpiperazine-1-carboxylate (1.04 g, 5.21 mmol) were added and the reaction was stirred at rt for 30 min. The mixture was poured into cold, satd. $NaHCO_3$ (5 mL) and stirred vigorously for 10 min. The mixture was partitioned between EtOAc (100 mL), and satd. $NaHCO_3$ (75 mL), the organic layer was washed with satd. $NaHCO_3$ (75 mL). The organic extract was dried over Na_2SO_4, filtered and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0–5% MeOH/DCM) to provide *tert*-butyl (*S*)-4-(6,7-dichloro-1-(5-isopropyl-3-methylisoxazol-4-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (2.5 g, 98 % yield). *m/z* (ESI, +ve ion): 537.0 $(M+H)^+$.

[0158] Step 4: *tert*-Butyl (3*S*)-4-(6-chloro-7-(2-fluoro-6-hydroxyphenyl)-1-(5-isopropyl-3-methylisoxazol-4-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate. A mixture of *tert*-butyl (*S*)-4-(6,7-dichloro-1-(5-isopropyl-3-methylisoxazol-4-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (0.6 g, 1.1 mmol), (2-fluoro-5-hydroxyphenyl)boronic acid (261 mg, 1.68 mmol), $Pd(PPh_3)_4$ (0.13 g, 0.11 mmol) and sodium carbonate (0.36 g, 3.35 mmol) was purged with N2 followed by the addition of 1,4-dioxane (12 mL) and water (3 mL). The mixture was heated at 80 °C for 1 h then quenched with sat. $NaHCO_3$, extracted with EtOAc, dried over Na_2SO_4, filtered and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0–5% MeOH/DCM) to provide *tert*-butyl (3*S*)-4-(6-chloro-7-(2-fluoro-6-hydroxyphenyl)-1-(5-isopropyl-3-methylisoxazol-4-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (0.22 g, 0.37 mmol, 33 % yield) with some traces of ether byproduct. *m/z* (ESI, +ve ion): 613.0 $(M+H)^+$.

[0159] Step 5: 6-Chloro-7-(2-fluoro-6-hydroxyphenyl)-1-(3-methyl-5-(2-propanyl)-1,2-oxazol-4-yl)-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-piperazinyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of *tert*-butyl (3*S*)-4-(6-chloro-7-(2-fluoro-6-hydroxyphenyl)-1-(5-isopropyl-3-methylisoxazol-4-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (0.22 g, 0.37 mmol) in DCM (5 mL) at rt was added TFA (5 mL, 64.9 mmol) and the mixture was stirred at rt for 30 min. The mixture was concentrated *in vacuo* to afford 6-chloro-7-(2-fluoro-6-hydroxyphenyl)-1-(5-isopropyl-3-methylisoxazol-4-yl)-4-((*S*)-2-methylpiperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. The crude product was dissolved in DCM (15 mL), treated with DIPEA (0.26 mL, 1.5 mmol) and a solution of acryloyl

chloride (0.021 mL, 0.26 mmol) in DCM (1 mL) in small portions. After 30 min, the mixture was diluted with DCM, washed with satd. $NaHCO_3$, with satd. ammonium chloride. The organic extract was dried over Na_2SO_4, filtered and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0–5% MeOH/DCM) to provide 6-chloro-7-(2-fluoro-6-hydroxyphenyl)-1-(3-methyl-5-(2-propanyl)-1,2-oxazol-4-yl)-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-piperazinyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.16 g, 0.14 mmol, 37.4 % yield). ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 10.18 (br s, 1 H), 8.40 (br d, *J*=13.1 Hz, 1 H), 7.25-7.32 (m, 1 H), 6.79-6.91 (m, 1 H), 6.68-6.79 (m, 2 H), 6.20 (br d, *J*=16.6 Hz, 1 H), 5.74-5.79 (m, 1 H), 4.93 (br d, *J*=27.8 Hz, 1 H), 3.97-4.46 (m, 3 H), 3.36-3.88 (m, 2 H), 2.98-3.28 (m, 1 H), 2.85-2.97 (m, 1 H), 1.92 (br d, *J*=6.0 Hz, 3 H), 1.33 (dd, *J*=12.6, 6.6 Hz, 3 H), 1.10 (d, *J*=6.8 Hz, 3 H), 1.05 (dd, *J*=6.9, 2.6 Hz, 3 H). ^{19}F NMR (376 MHz, *DMSO-d*$_6$) δ -115.73 (br dd, *J*=82.4, 10.4 Hz, 1 F). *m/z* (ESI, +ve ion): 567.2 $(M+H)^+$.

Table 55: Compounds 55-50 to 55-64 were prepared following the procedure described in Method 55, Steps 1-5, above as follows:

Ex.#	Chemical Structure	Name	Method Changes	Reagent
55-50		6-Chloro-1-(4-chloro-2-isopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Intermediate 190** in **Step 1**	**Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC, **Step 4**: (2-fluorophenyl)boranediol, Combi-Blocks Inc
55-51		6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-	**Intermediate 163** in **Step 1**	**Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC, **Step 4**: (2-fluorophenyl)boranediol, Combi-Blocks Inc

		d]pyrimidin-2-one		
55-52		1-(4-Chloro-2-isopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-(2-fluorophenyl)-7-methoxy-pyrido[2,3-*d*]pyrimidin-2-one	**Intermediate 190** in **Step 1** and **Intermediate 191**	(2- **Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC, **Step 4**: (2-fluorophenyl)borane diol, Combi-Blocks Inc
55-53		6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-methylsulfonylphenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Intermediate 163** in **Step 1**	**Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC, **Step 4:** (2-methylsulfonylphenyl)boronic acid, Combi-Blocks Inc.
55-54		7-(2-Amino-6-fluoro-phenyl)-6-chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]pyrido[2,3-*d*]pyrimidin-2-one	**Intermediate 163** in **Step 1**	**Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC **Step 4:** (2-amino-6-fluorophenyl)boronic acid pinacol ester, CombiPhos Catalysts, Inc.

55-55		6-Chloro-1-(2-dimethylphosphorylphenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**None**	**Step 1:** (2-aminophenyl)dimethylphosphine oxide, AstaTech, Inc, **Step 4:** (2-fluorophenyl)boranediol, Combi-Blocks Inc.
55-56	 Single isomer (M)	7-(2-Amino-6-fluoro-phenyl)-6-chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]pyrido[2,3-*d*]pyrimidin-2-one	**Intermediate 163** in **Step 1**	**Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC, **Step 4:** (2-amino-6-fluorophenyl)boronic acid pinacol ester, CombiPhos Catalysts, Inc.
55-57	 Single isomer (P)	7-(2-Amino-6-fluoro-phenyl)-6-chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]pyrido[2,3-*d*]pyrimidin-2-one	**Intermediate 163** in **Step 1**	**Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC, **Step 4:** (2-amino-6-fluorophenyl)boronic acid pinacol ester, CombiPhos Catalysts, Inc.
55-58		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(4-ethyl-2-isopropyl-3-pyridyl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Intermediate 197** in **Step 1**	**Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 4:** 2-Fluorophenyl boronic acid, Combi-Blocks

55-59		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-methyl-pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** methylamine (Sigma-Aldrich Corporation); **Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (AstaTech, Inc.); **Step 4:** (2-fluorophenyl)boronic acid (Combi-Blocks Inc.)
55-60		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-isopropyl-pyrido[2,3-*d*]pyrimidin-2-one	**None**	**Step 1:** isopropylamine (Sigma-Aldrich Corporation); **Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (AstaTech, Inc.); **Step 4:** (2-fluorophenyl)boronic acid (Combi-Blocks Inc.)

Method 69

6-Chloro-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-piperazinyl)-7-(1-piperidinyl)-1-(2-(2-propanyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one.



[0160] Step 1: *tert*-Butyl (*S*)-4-(6-chloro-1-(2-isopropylphenyl)-2-oxo-7-(piperidin-1-yl)-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate. A 25-mL round-bottomed flask was charged with (*S*)-*tert*-butyl 4-(6,7-dichloro-1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (**Intermediate 61B**, 45 mg, 0.085 mmol) and piperidine (0.1 mL, 1.01 mmol, Spectrum Chemicals & Laboratory Products, Gardena, CA, USA). The reaction mixture was stirred and heated at 80 °C for 16 h. The reaction mixture was concentrated *in vacuo* to give crude (*S*)-*tert*-butyl 4-(6-chloro-1-(2-isopropylphenyl)-2-oxo-7-(piperidin-1-yl)-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (49 mg) as a yellow solid. *m/z* (ESI, +ve ion): 581.3 $(M+H)^+$. The crude yellow solid was used in next step without purification.

[0161] Step 2: 6-Chloro-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-piperazinyl)-7-(1-piperidinyl)-1-(2-(2-propanyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of (*S*)-*tert*-butyl 4-(6-chloro-1-(2-isopropylphenyl)-2-oxo-7-(piperidin-1-yl)-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (49 mg, 0.084 mmol) in DCM (1 mL) was treated with TFA (1 mL) at rt and stirred for 15 min. The reaction was concentrated *in vacuo* to afford (*S*)-6-chloro-1-(2-isopropylphenyl)-4-(2-methylpiperazin-1-yl)-7-(piperidin-1-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. *m/z* (ESI, +ve ion): 481.3 $(M+H)^+$.

[0162] A mixture of (*S*)-6-chloro-1-(2-isopropylphenyl)-4-(2-methylpiperazin-1-yl)-7-(piperidin-1-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (0.07 mL, 0.42 mmol) in DCM (1.0 mL) was added acryloyl chloride (0.26 M in DCM, 0.33 mL, 0.084 mmol) at 0 °C and stirred for 40 min at 0 °C. The resulting mixture was concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-50% of EtOAc/EtOH(3:1)/heptane) to provide pure (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-6-chloro-1-(2-isopropylphenyl)-7-(piperidin-1-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (28 mg, 0.053 mmol, 62.5 % yield) as a light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.85 - 8.03 (m, 1 H), 7.42 - 7.50 (m, 1 H), 7.34 - 7.42 (m, 1 H), 7.26 (t, J=7.5 Hz, 1 H), 7.05 (d, J=7.7 Hz, 1 H), 6.74 - 6.92 (m, 1 H), 6.19 (br dd, J=17.5, 4.5 Hz, 1 H), 5.71 - 5.78 (m, 1 H), 4.68 - 4.84 (m, 1 H), 4.21 - 4.45 (m, 1 H),

3.93 - 4.19 (m, 2 H), 3.35 - 3.66 (m, 2 H), 3.25 (br s, 4 H), 2.89 - 3.19 (m, 1 H), 2.45 - 2.48 (m, 1 H), 1.34 (br d, J=3.9 Hz, 3 H), 1.20 - 1.29 (m, 6 H), 1.08 (d, J=6.8 Hz, 3 H), 0.98 (br d, J=6.8 Hz, 3 H). *m/z* (ESI, +ve ion): 536.3 $(M+H)^+$.

Table 69: Compounds 69-10 to 69-12 were prepared following the procedure described in Method 69, Steps 1-2, above as follows:

Ex.#	Chemical Structure	Name	Method Changes	Reagent
69-10		*(M)*-6-Chloro-7-[2-(difluoromethyl)pyrrolidin-1-yl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** 2-(difluoromethyl) pyrrolidine hydrochloride (Advanced ChemBlocks Inc.)
69-11		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(1-piperidyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** [3-(prop-1-yn-1-yl)phenyl]boronic acid (Enamine Ltd.)
69-12		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-[(2*SR*)-2-methyl-1-piperidyl]pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** piperidine (Spectrum Chemicals & Laboratory Products)

Method 72

(*M*)-6-Fluoro-7-(5-fluoro-2-hydroxyphenyl)-1-(4-methyl-2-(2-propanyl)-3-pyridinyl)-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-piperazinyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0163] Step 1: (*M*)-7-Chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4- dione (Intermediate 72A). A mixture of atropisomers 7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Example 41, Step 3**, 648 g) was purified by SFC (AD, 150 x 50 mm, 5 μm, 50% MeOH/CO_2, 180 g/min, 102 bar) to obtain two peaks: Peak 1 (P isomer, 230.6 g, >99% ee) and Peak 2 (M isomer, 227.8 g, 97.1% ee, **Intermediate 72A**).

[0164] Step 2: (*M*)-4,7-Dichloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a suspension of (*M*)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 72A**, 2.51 g, 7.19 mmol) in a mixture of acetonitrile (11 mL) and DIPEA (1.9 mL, 11 mmol) was added phosphorous oxychloride (0.87 mL, 9.3 mmol). The mixture was heated at 80 °C for 90 min and then concentrated *in vacuo*. The crude residue was used without further purification in the following step. *m/z* (ESI, +ve ion): 367.0 $(M+H)^+$.

[0165] Step 3: (*M*,*S*)-*tert*-Butyl 4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (Intermediate 65A). A solution of (*M*)-4,7-dichloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (2.64 g, 7.19 mmol) in acetonitrile (11 mL) was cooled in an ice-water bath to 0 °C. DIPEA (3.8 mL, 22 mmol) was added, followed by (3*S*)-1-(*tert*-butoxycarbonyl)-3-methylpiperazine (1.8 g, 8.7 mmol, Ark Pharm, Inc., Libertyville, IL, USA). The mixture was allowed to warm to rt and stir for 18 h. The mixture was quenched with satd. $NaHCO_3$ (100 mL). The mixture was diluted with EtOAc (175 mL), and water (75 mL). The

aqueous layer was washed with EtOAc (2 × 100 mL). The combined organic phases were dried over $MgSO_4$ and then concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 10-50% EtOAc-EtOH (3:1)/heptane) to provide (*M*,*S*)-*tert*-butyl 4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (Intermediate 65A). *m/z* (ESI, +ve ion): 530.9 $(M+H)^+$.

[0166] Step 4: (*M*,*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. TFA (10 mL) was added to a solution of (*M*,*S*)-*tert*-butyl 4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (**Intermediate 65A**, 3.0 g, 5.7 mmol) in DCM (20 mL). The mixture was stirred for 20 min at rt and then concentrated *in vacuo*. The resulting residue was re-dissolved in DCM (40 mL) and cooled to 0 °C. DIPEA (5.0 mL, 28 mmol) and acryloyl chloride (0.46 mL, 5.7 mmol) were sequentially added and the mixture was stirred for 80 min. The reaction mixture was quenched at 0 °C by adding satd. $NaHCO_3$ (100 mL) and water (50 mL) and diluted with DCM (150 mL). The aqueous layer was extracted with DCM (2 × 50 mL). The combined organic extracts were dried over $MgSO_4$ and then concentrated *in vacuo* to give crude (*M*,*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one which was used without further purification in the following step. *m/z* (ESI, +ve ion): 485.0 $(M+H)^+$.

[0167] Step 5: (*M*)-6-Fluoro-7-(5-fluoro-2-hydroxyphenyl)-1-(4-methyl-2-(2-propanyl)-3-pyridinyl)-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-piperazinyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A mixture of (*M*,*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (120 mg, 0.25 mmol), (5-fluoro-2-hydroxy)phenylboronic acid (58 mg, 0.37 mmol, Combi-Blocks, San Diego, CA, USA), $Pd(dppf)Cl_2$ (18 mg, 0.025 mmol), and potassium acetate (120 mg, 1.2 mmol) in 1,4-dioxane (1.2 mL) and one drop of water was deoxygenated with nitrogen for 10 min. The mixture was stirred at 90 °C for 2 h, then was filtered through a plug of silica gel and then partitioned between water (50 mL) and EtOAc (50 mL). The organic phase was washed with water (50 mL) and then with brine (50 mL), dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-100% EtOAc-EtOH (3:1)/heptane) to provide (*M*)-6-fluoro-7-(5-fluoro-2-hydroxyphenyl)-1-(4-methyl-2-(2-propanyl)-3-pyridinyl)-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-

piperazinyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.96 (d, J = 6.8 Hz, 3 H), 1.08 (d, J = 6.8 Hz, 3 H), 1.33 (d, J = 6.8 Hz, 3 H), 1.94 (s, 3 H), 2.68 - 2.78 (m, 0.5 H), 3.06 - 3.19 (m, 0.5 H), 3.43 - 3.55 (m, 0.5 H), 3.59 - 3.80 (m, 1.5 H), 3.98 - 4.07 (m, 0.5 H), 4.10 - 4.19 (m, 0.5 H), 4.29 (br d, J = 13.5 Hz, 1.5 H), 4.40 (br d, J = 12.7 Hz, 0.5 H), 4.91 (br s, 1 H), 5.73 - 5.81 (m, 1 H), 6.21 (br d, J=16.4 Hz, 1 H), 6.80 - 6.93 (m, 2 H), 7.09 (dd, J = 9.4, 3.2 Hz, 1 H), 7.18 (td, J = 8.5, 3.1 Hz, 1 H), 7.26 (d, J = 5.0 Hz, 1 H), 8.26 - 8.37 (m, 1 H), 8.48 (d, J = 5.0 Hz, 1 H), 10.35 (s, 1 H). ^{19}F NMR (376 MHz, DMSO-d_6) δ –126.02 (s, 1 F), –125.16 (s, 1 F). *m/z* (ESI, +ve ion ion): 561.0 $(M+H)^+$.

Table 72: Compounds 72-20 to 72-23 were prepared following the procedure described in Method 72, Steps 1-5, above as follows:

Ex.#	Chemical Structure	Name	Method Changes	Reagent
72-20	 Single isomer (M)	*(M)*-6-Fluoro-7-(2-fluoro-6-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)-4-(3,3,5,5-tetradeuterio-4-prop-2-enoyl-piperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2-one	Tetrakis and sodium carbonate used in **Step 5**	**Step 3:** piperazine-3,3,5,5-d_4-*N*-t-Boc, C/D/N Isotopes Inc.
72-21	 Single isomer (M)	*(M)*-6-Fluoro-7-(2-fluoro-6-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)-4-(2,2,6,6-tetradeuterio-4-prop-2-enoyl-piperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2-one	Tetrakis and sodium carbonate used in **Step 5**	**Step 3:** piperazine-3,3,5,5-d_4-*N*-t-Boc, C/D/N Isotopes Inc.

72-22	 Single isomer (M)	*(M)*-6-Fluoro-7-(2-fluoro-6-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)-4-(2,2,3,3,5,5,6,6-octadeuterio-4-prop-2-enoyl-piperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2-one	Tetrakis and sodium carbonate used in **Step 5**	**Step 3:** *tert*-butyl piperazine-1-carboxylate-2,2,3,3,5,5,6,6-d_8, CDN ISOTOPES
72-23	 Single isomer (M)	*(M)*-7-(2-Dimethylphosphorylphenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	Tetrakis, sodium carbonate and **Intermediate 192,** used in **Step 5**	**Step 3:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC

Method 79

(*M*)-6-Chloro-7-(2-fluorophenyl)-1-(4-methyl-2-(2-propanyl)-3-pyridinyl)-4-((1-(2-propenoyl)-3-azetidinyl)oxy)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one





[0168] Step 1: (*M*)-*tert*-Butyl 3-((6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)oxy)azetidine-1-

carboxylate. A 50-mL round-bottomed flask was charged with (*M*)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 76A**, 256 mg, 0.60 mmol) and DIPEA (0.2 mL, 0.90 mmol) in acetonitrile (3 mL) followed by phosphorous oxychloride (0.1 mL, 0.9 mmol). The mixture was stirred and heated at 80 °C for 40 min. The reaction mixture was concentrated *in vacuo* to give (*M*)-4,6-dichloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as brown solid [*m/z* (ESI, +ve ion): 443 $(M+H)^+$] which was used in next step without purification.

[0169] A mixture of the above crude (*M*)-4,6-dichloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (267 mg, 0.6 mmol) in acetonitrile (3 mL) was treated with DIPEA (0.3 mL, 1.81 mmol) followed by 1-Boc-3-(hydroxy)azetidine (313 mg, 1.81 mmol, CNH Technologies, Inc., Woburn, MA, USA). The reaction mixture was stirred at rt for 24 h then concentrated *in vacuo* and purified by silica gel chromatography (eluent: 0-30% 3:1 EtOAc-EtOH/heptane) to provide (*M*)-*tert*-butyl 3-((6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)oxy)azetidine-1-carboxylate (51 mg, 0.09 mmol, 14.6 % yield) as a light yellow foam. *m/z* (ESI, +ve ion) 580.3 $(M+H)^+$.

[0170] **Step 2: (M)-4-((1-Acryloylazetidin-3-yl)oxy)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-d]pyrimidin-2(1H)-one**. To a solution of (M)-tert-butyl 3-((6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)oxy)azetidine-1-carboxylate (51 mg, 0.9 mmol) in DCM (1 mL) was treated with TFA (1 mL) at rt and stirred for 1 h. The reaction was concentrated to afford (M)-4-(azetidin-3-yloxy)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-d]pyrimidin-2(1H)-one as a yellow gum. m/z (ESI, +ve ion) 480.2 (M+H)+.

[0171] A mixture of the above (M)-4-(azetidin-3-yloxy)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-d]pyrimidin-2(1H)-one and DIPEA (0.08 mL, 0.44 mmol) in DCM (1 mL) was treated with acryloyl chloride (0.25 M in DCM, 0.3 mL, 0.08 mmol) at 0 °C and stirred for 5 min. The reaction mixture was concentrated in vacuo and the crude product was purified by silica gel chromatography (eluent: 0-50% EtOAc-EtOH (3:1)/heptane) to give (M)-4-((1-acryloylazetidin-3-yl)oxy)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-d]pyrimidin-2(1H)-one (22 mg, 0.041 mmol, 46.2 % yield) as a white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.83 (s, 1 H), 8.43 (d, J=4.8 Hz, 1 H), 7.47 - 7.64 (m, 1 H), 7.18 - 7.39 (m, 4 H), 6.32 - 6.51 (m, 1 H), 6.16 (dd, J=17.0, 2.1 Hz,

1 H), 5.62 - 5.84 (m, 2 H), 4.70 - 4.85 (m, 1 H), 4.54 (td, J=10.4, 3.2 Hz, 1 H), 4.44 (br dd, J=11.2, 6.8 Hz, 1 H), 4.23 - 4.37 (m, 1 H), 2.76 (dt, J=13.4, 6.8 Hz, 1 H), 1.96 (s, 3 H), 1.07 (d, J=6.8 Hz, 3 H), 0.94 (d, J=6.6 Hz, 3 H). 19F NMR (376 MHz, DMSO-d6) δ -114.69 (d, J=4.3 Hz, 1 F). m/z (ESI, +ve) 534.1 (M+H)+.

Table 79: Compounds 79-10 to 79-12 were prepared following the procedure described in Method 79, Steps 1-2, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
79-10	 Single isomer (*M*)	(*M*)-6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)-4-[(1*S*,5*R*)-8-prop-2-enoyl-3,8-diazabicyclo[3.2.1]octan-3-yl]pyrido[2,3-*d*]pyrimidin-2-one	Omit **Step 2**	**Step 1.2**: Intermediate 213
79-11	 Single isomer (*M*)	(*M*)-6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)-4-[(1*S*,4*S*)-5-prop-2-enoyl-2,5-diazabicyclo[2.2.2]octan-2-yl]pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1.2**: (1*S*,4*S*)-2-Boc-2,5-diazabicyclo(2.2.2)octane; Astatech, Inc., Bristol, PA
79-12	 Single isomer (*M*)	(*M*)-6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)-4-[(1*S*,5*R*)-3-prop-2-enoyl-3,9-diazabicyclo[3.3.1]nonan-9-yl]pyrido[2,3-*d*]pyrimidin-2-one	Omit **Step 2**	**Step 1.2**: Intermediate 214

Method 80

6-Chloro-1-(4-((dimethylamino)methyl)-2-methyl-6-(2-propanyl)phenyl)-7-(2-fluorophenyl)-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-piperazinyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0172] Step 1: *N*-((4-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)carbamoyl)-2,5,6-trichloronicotinamide. To a stirred solution of 2,5,6-trichloronicotinamide (**Intermediate P**, 1.3 g, 5.5 mmol) in THF (10 mL) was added oxalyl chloride (2 M in DCM, 4.2 mL, 8.4 mmol). After the addition was completed, the reaction mixture was stirred and heated at 65 °C for 2 h. The reaction mixture was cooled, concentrated *in vacuo* and the crude residue was dissolved in THF (10 mL) and a solution of 4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylaniline (Intermediate I-37, 2.3 g, 5.5 mmol) in THF (10 mL) was added. After the addition was completed, the solution was maintained at rt for 2 h. The mixture was concentrated *in vacuo* to provide crude *N*-((4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)carbamoyl)-2,5,6-trichloronicotinamide which was carried forward in the next step without purification. *m/z* (ESI, +ve ion): 668.0 $(M+H)^+$.

[0173] Step 2: 1-(4-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a stirred solution of *N*-((4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-

methylphenyl)carbamoyl)-2,5,6-trichloronicotinamide (3.7 g, 5.5 mmol) in THF (20 mL) at 0 °C was added 1 M KHMDS in THF (11 mL, 11 mmol). After 2 h, the reaction was quenched with satd. ammonium chloride and diluted with EtOAc. The aqueous layer was extracted with EtOAc and the combined organic extracts were dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-50% EtOAc/heptane) to provide 1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. *m/z* (ESI, +ve ion): 632.0 $(M+H)^+$.

[0174] Step 3: *tert*-Butyl (*S*)-4-(1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate. To a stirred solution of 1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (2.8 g, 4.5 mmol) in acetonitrile (20 mL) was added DIPEA (1.2 mL, 6.7 mmol), followed by phosphorus oxychloride (0.63 mL, 6.7 mmol). After the addition was completed, the mixture was stirred and heated at 60 °C for 3 h. The mixture was concentrated *in vacuo* to provide a crude residue which was dissolved in DCM. The solution was cooled to 0 °C and DIPEA (3.9 mL, 23 mmol) and *tert*-butyl (*S*)-3-methylpiperazine-1-carboxylate (0.90 g, 4.5 mmol) were added. After 2 h, the reaction was quenched with water, the aqueous layer was extracted with DCM and the combined organic extracts were dried over Na_2SO_4, then concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-50% EtOAc/heptane) to provide *tert*-butyl (*S*)-4-(1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 0.96 (d, *J* = 6.7 Hz, 3 H), 1.02 (d, *J* = 6.8 Hz, 3 H), 1.08 (s, 9 H), 1.31 (dd, *J* = 8.9, 7.1 Hz, 3 H), 1.45 (s, 9 H), 1.87 (d, *J* = 2.5 Hz, 3 H), 2.39 - 2.47 (m, 1 H), 2.96 - 3.19 (m, 2 H), 3.28 (s, 1 H), 3.59 - 3.74 (m, 1 H), 3.82 (br d, *J* = 12.0 Hz, 1 H), 3.88 - 4.00 (m, 1 H), 4.09 - 4.23 (m, 1 H), 4.83 (s, 3 H), 7.10 (s, 1 H), 7.31 (s, 1 H), 7.41 - 7.54 (m, 6 H), 7.69 (br d, *J* = 7.7 Hz, 4 H), 8.42 (d, *J* = 9.5 Hz, 1 H).

[0175] Step 4: *tert*-Butyl (*S*)-4-(1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate. A mixture of *tert*-butyl (*S*)-4-(1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (2.8 g, 3.4 mmol), (2-

fluorophenyl)boronic acid (0.71 g, 5.1 mmol, Combi-Blocks, San Diego, CA, USA), potassium acetate (1.7 g, 17 mmol), and $Pd(dppf)Cl_2$ (0.25 g, 0.34 mmol) in 1,4-dioxane (20 mL)/water (0.5 mL) was stirred and heated at 90 °C for 3 h. The mixture was cooled to rt and diluted with water. The aqueous mixture was extracted with EtOAc, the combined organic extracts were dried over Na_2SO_4 and concentrated *in vacuo* to provide *tert*-butyl (*S*)-4-(1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate which was used in the next step without further purification.

[0176] Step 5: (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of *tert*-butyl (*S*)-4-(1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (2.0 g, 2.3 mmol) in DCM (10 mL) was added TFA (3.5 mL, 46 mmol). The reaction was stirred at rt for 3 h and concentrated *in vacuo*. The residue was dissolved in DCM (10 mL), cooled to 0 °C, then treated with DIPEA (2.0 mL, 11 mmol) and acryloyl chloride (1.1 M in DCM, 2.1 mL, 2.3 mmol). The reaction was stirred at 0 °C for 2 h, the mixture was diluted with water and extracted with DCM. The organic layer was dried over Na_2SO_4 and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-70% EtOAc/heptane) to provide (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 0.88 - 0.94 (m, 3 H), 1.03 (s, 9 H), 1.24 - 1.28 (m, 3 H), 1.31 - 1.37 (m, 3 H), 1.87 (s, 3 H), 3.03 - 3.33 (m, 2 H), 3.58 - 3.83 (m, 2 H), 3.98 - 4.19 (m, 1 H), 4.22 - 4.48 (m, 2 H), 4.76 (s, 2 H), 4.93 (br s, 1 H), 5.76 (br d, *J* = 10.6 Hz, 1 H), 6.13 - 6.27 (m, 1 H), 6.79 - 6.94 (m, 1 H), 7.02 (br s, 1 H), 7.17 - 7.33 (m, 4 H), 7.35 - 7.55 (m, 7 H), 7.60 - 7.68 (m, 4 H), 8.44 (br s, 1 H). ^{19}F NMR (377 MHz, *DMSO-d*$_6$) δ -114.17 (s, 1 F).

[0177] Step 6: (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-(hydroxymethyl)-2-isopropyl-6-methylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Intermediate 80A). To a stirred solution of (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (270 mg, 0.33 mmol) in THF (5 mL) was added TBAF (1 M in THF, 0.33 mL, 0.33 mmol). The mixture was stirred at rt for 2 h and

concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-50% 3:1 EtOAc–EtOH/heptane) to provide (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-(hydroxymethyl)-2-isopropyl-6-methylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 80A**). ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 0.94 (d, *J* = 6.8 Hz, 3 H), 1.05 (d, *J* = 6.8 Hz, 3 H), 1.33 (d, *J* = 6.8 Hz, 3 H), 1.87 (s, 3 H), 2.51 - 2.58 (m, 1 H), 3.03 - 3.28 (m, 1 H), 3.41 - 3.86 (m, 2 H), 3.97 - 4.44 (m, 3 H), 4.46 (d, *J* = 5.8 Hz, 2 H), 4.92 (br s, 1 H), 5.14 (t, *J* = 5.8 Hz, 1 H), 5.71 - 5.81 (m, 1 H), 6.20 (br dd, *J* = 16.6, 3.3 Hz, 1 H), 6.79 - 6.94 (m, 1 H), 7.05 (s, 1 H), 7.16 (s, 1 H), 7.18 - 7.23 (m, 1 H), 7.24 - 7.36 (m, 2 H), 7.47 - 7.56 (m, 1 H), 8.43 (br d, *J* = 4.6 Hz, 1 H). ^{19}F NMR (376 MHz, *DMSO-d*$_6$) δ -114.26 (s, 1 F); *m/z* (ESI, +ve ion): 590.0 $(M+H)^+$.

[0178] Step 7: (*S*)-4-(4-(4-Acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-3-isopropyl-5-methylbenzaldehyde. To a solution of (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-(hydroxymethyl)-2-isopropyl-6-methylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 80A**, 100 mg, 0.17 mmol) in DCM (3.4 mL) at rt was added Dess-Martin periodinane (110 mg, 0.25 mmol). After 20 min, the reaction was quenched by addition of 1 N sodium thiosulfate (10 mL) and diluted with DCM (5 mL). The layers were partitioned and then the aqueous phase was washed with DCM (2 × 10 mL). The combined organic extracts were dried over anhydrous sodium sulfate, filtered through Celite®, and concentrated under reduced pressure to afford crude (*S*)-4-(4-(4-acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-3-isopropyl-5-methylbenzaldehyde as a light-yellow foam that was carried forward in the following step without purification. *m/z* (ESI, +ve ion): 588.0 $(M+H)^+$.

[0179] Step 8: 6-Chloro-1-(4-((dimethylamino)methyl)-2-methyl-6-(2-propanyl)phenyl)-7-(2-fluorophenyl)-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-piperazinyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of (*S*)-4-(4-(4-acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-3-isopropyl-5-methylbenzaldehyde (100 mg, 0.17 mmol) in THF (1.0 mL) at rt was added dimethyl amine solution (2 M in THF, 95 µL, 0.19 mmol), glacial acetic acid (10 µL, 0.17 mmol), and sodium triacetoxyborohydride (72 mg, 0.34 mmol). The resulting cloudy yellow mixture was stirred at rt. After 18 h, the reaction mixture was diluted with EtOAc (10 mL) and water (5 mL), then saturated aqueous sodium bicarbonate (5 mL) was added until the aqueous phase was neutralized. The layers were partitioned and the aqueous phase was extracted with

EtOAc (1 × 20 mL), then the combined organic extracts were dried over anhydrous magnesium sulfate, filtered, and concentrated to afford a light-yellow oil. The crude product was purified by silica gel chromatography (eluent: 0-20% 2 M NH3 in MeOH/DCM) to afford 6-chloro-1-(4-((dimethylamino)methyl)-2-methyl-6-(2-propanyl)phenyl)-7-(2-fluorophenyl)-4-((2*S*)-2-methyl-4-(2-propenoyl)-1-piperazinyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 0.92 (d, *J* = 6.8 Hz, 3 H), 1.05 (d, *J* = 6.8 Hz, 3 H), 1.31 - 1.36 (m, 3 H), 1.87 (s, 3 H), 2.14 (s, 6 H), 3.02 - 3.26 (m, 1 H), 3.35 (s, 2 H), 3.44 - 3.83 (m, 2 H), 3.97 - 4.47 (m, 3 H), 4.93 (br s, 1 H), 5.72 - 5.80 (m, 1 H), 6.13 - 6.28 (m, 1 H), 6.79 - 6.93 (m, 1 H), 7.02 (s, 1 H), 7.11 (s, 1 H), 7.14 - 7.36 (m, 4 H), 7.42 - 7.56 (m, 1 H), 8.43 (br s, 1 H). ^{19}F NMR (376 MHz, *DMSO-d*$_6$) δ -113.53 (s, 1 F). *m/z* (ESI, +ve ion): 617.2 $(M+H)^+$.

Table 80: Compounds 80-10 to 80-21 were prepared following the procedure described in Method 80, Steps 1-8, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
80-10		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-[2-isopropyl-6-(pyrrolidin-1-ylmethyl)phenyl]pyrido[2,3-*d*]pyrimidin-2-one	Step 1 (Intermediates 99B and **I-40**). Omit step 4. Used Pyridine. SO3 instead of DMP in step 7	Step 8: (pyrrolidine)
80-11		6-chloro-1-[2-[(3,3-difluoroazetidin-1-yl)methyl]-6-isopropyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	Step 1 (Intermediates 99B and **I-40**). Omit step 4. Used Pyridine. SO3 instead of DMP in step 7	Step 8: (3,3-difluoroazetidine hydrochloride)
80-12		1-[2-(azetidin-1-ylmethyl)-6-isopropyl-phenyl]-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	Step 1 (Intermediates 99B and **I-40**). Omit step 4. Used Pyridine. SO3 instead of DMP in step 7	Step 8: (azetidine)

80-13		6-Chloro-1-[2-[(dimethylamino)methyl]-6-isopropyl-phenyl]-7-(2-fluorophenyl)-4-[(2*S*)-2-methyl-4-prop-2-enoyl-piperazin-1-yl]pyrido[2,3-*d*]pyrimidin-2-one	Step 8: dimethylamine hydrochloride and no acetic acid	Step 7: **Intermediate 206**
80-14		6-Chloro-1-[2-[(dimethylamino)methyl]-6-isopropyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	Step 8: dimethylamine hydrochloride and no acetic acid	Step 1: **Intermediate I-40**; Step 3: (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (eNovation Chemicals LLC)
80-16		6-Chloro-1-[4-[(dimethylamino)methyl]-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	Step 8: dimethylamine hydrochloride and no acetic acid	Step 1: **Intermediate 207**; Step 3: (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (eNovation Chemicals LLC)
80-18		6-Chloro-1-[4-[(dimethylamino)methyl]-6-isopropyl-pyrimidin-5-yl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-	Step 8: dimethylamine hydrochloride and no acetic acid	**Step 1: Intermediate 208**; Step 3: (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (eNovation Chemicals LLC)

		fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one		
80-20		1-[4-[(Dimethylamino)methyl]-2-isopropyl-6-methyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 6:** 0 °C to room temp	**Step 1:** Intermediate S; **Step 3 (2):** *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (eNovation Chemicals LLC, Bridgewater, NJ, USA)
80-21		1-[4-[(Dimethylamino)methyl]-2-isopropyl-6-methyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 6:** 0 °C to room temp	**Step 1:** Intermediate S; **Step 3 (2):** *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (eNovation Chemicals LLC, Bridgewater, NJ, USA); **Step 4:** Intermediate Q

Method 92

Example 92-1: (*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one



Intermediate 73B

Intermediate 92A

Intermediate 92B

[0180] Step 1: (*M*)-*tert*-Butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a mixture of (*M*)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 73B**, 9.96 g, 27.3 mmol) and DIPEA (7.1 mL, 40.9 mmol) in acetonitrile (100 mL) was added phosphorous oxychloride (3.8 mL, 40.9 mmol) and the mixture was stirred at 80 °C for 1.5 h. The reaction mixture was concentrated *in vacuo* to give brown solid. The crude solid was used in next step without purification. *m/z* (ESI, +ve ion): 383.0 $(M+H)^+$.

[0181] The above crude solid and DIPEA (7.1 mL, 40.9 mmol) in DMF (60 mL) was treated with (2*R*,5*S*)-1-(*tert*-butoxycarbonyl)-2,5-dimethylpiperazine (6.43 g, 30 mmol, AstaTech, Inc.) and stirred at rt for 16 h. The mixture was treated with satd. $NaHCO_3$ (100 mL) and stirred at rt for 15 min. The resulting precipitate was collected by filtration, washed with water, and dried to give (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 92A**, 14.8 g, 26.4 mmol, 97 % yield). ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 - 8.50 (m, 2 H), 7.26 (d, *J*=5.0 Hz, 1 H), 4.12 - 4.87 (m, 2 H), 3.94 - 4.05 (m, 1 H), 3.89 (br d, *J*=7.9 Hz, 1 H), 3.67 (dd, *J*=13.8, 2.6 Hz, 1 H), 3.41 - 3.58 (m, 1 H), 2.61 (quin, *J*=6.7 Hz, 1

H), 1.94 (s, 3 H), 1.44 (s, 9 H), 1.30 (d, *J*=6.4 Hz, 3 H), 1.09 (br d, *J*=6.4 Hz, 3 H), 1.05 (d, *J*=6.8 Hz, 3 H), 1.02 (d, *J*=6.6 Hz, 3 H). *m/z* (ESI, +ve ion): 561.2 $(M+H)^+$.

[0182] Step 2: (*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A mixture of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 92A**, 0.82 g, 1.45 mmol) in DCM (7 mL) and TFA (7 mL) was stirred at rt for 30 min. The reaction was concentrated *in vacuo* to afford (*M*)-6,7-dichloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as yellow syrup. *m/z* (ESI, +ve ion): 461.2 $(M+H)^+$.

[0183] A mixture of (*M*)-6,7-dichloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (1.3 mL, 7.26 mmol) in DCM (7 mL) was treated with acryloyl chloride (0.12 mL, 1.45 mmol) at 0 °C and stirred for 40 min. The reaction mixture was concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-100% EtOAc-MeOH (9:1)/heptane) to provide pure (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 92B**; 0.53 g, 1.03 mmol, 71 % yield) as a light-yellow foam. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 - 8.54 (m, 2 H), 7.30 (d, *J*=5.0 Hz, 1 H), 6.82 (ddd, *J*=16.5, 14.0, 10.5 Hz, 1 H), 6.18 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.74 (dt, *J*=10.4, 2.7 Hz, 1 H), 4.78 - 4.91 (m, 1 H), 4.39 - 4.75 (m, 1 H), 3.97 - 4.16 (m, 1 H), 3.94 (br s, 1 H), 3.83 (br d, *J*=3.9 Hz, 1 H), 3.49 (br dd, *J*=13.9, 3.7 Hz, 1 H), 2.59 - 2.70 (m, 1 H), 1.97 (s, 3 H), 1.25 - 1.32 (m, 3 H), 1.09 - 1.20 (m, 3 H), 1.05 (dd, *J*=11.4, 6.6 Hz, 6 H). *m/z* (ESI, +ve ion): 515.2 $(M+H)^+$.

[0184] Step 3: (*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one. A mixture of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 92B**; 1.01 g, 1.97 mmol), tetrakis(triphenylphosphine)palladium (0) (0.23 g, 0.20 mmol, Sigma-Aldrich Corporation), 2-isopropylphenylboronic acid (0.49 g, 2.95 mmol, Combi-Blocks Inc.) and sodium carbonate (0.63 g, 5.9 mmol) in 1,4-dioxane (6 mL) and water (3 mL) was stirred and heated at 90 °C for 2 h. The mixture was treated with water (25 mL), extracted with EtOAc (2 × 50 mL), dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-70% of EtOAc-EtOH (3:1)/heptane) to provide

(*M*)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one (0.52 g, 0.86 mmol, 43.7 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.42 (d, *J*=5.2 Hz, 1 H), 8.35 (d, *J*=5.0 Hz, 1 H), 7.37 (d, *J*=3.9 Hz, 2 H), 7.22 (dt, *J*=8.2, 4.0 Hz, 1 H), 7.16 (d, *J*=5.0 Hz, 1 H), 7.00 (d, *J*=7.5 Hz, 1 H), 6.84 (ddd, *J*=20.3, 16.8, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 4.44 - 4.93 (m, 2 H), 3.39 - 4.32 (m, 4 H), 2.74 (br s, 1 H), 2.52 - 2.60 (m, 1 H), 1.89 (s, 3 H), 1.36 (dd, *J*=6.3, 3.8 Hz, 3 H), 1.26 (br dd, *J*=24.7, 6.6 Hz, 3 H), 0.80 - 1.10 (m, 12 H). *m/z* (ESI, +ve ion): 599.2 $(M+H)^+$.

Table 92: Compounds 92-2 to 92-73 were prepared following the procedure described in Method 92, Steps 1-3, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
92-2		(*M*)-7-(2-Amino-6-fluoro-phenyl)-6-chloro-1-(2-isopropyl-4-methyl-3-pyridyl)-4-[(3*R*)-3-methyl-4-prop-2-enoyl-piperazin-1-yl]pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (*R*)-1-*N*-Boc-2-methylpiperazine (Combi-Blocks Inc); **Step 3:** (2-amino-6-fluorophenyl)boronic acid pinacol ester (CombiPhos Catalysts)
92-3		(*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-methyl-3-thienyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (2-methylthiophen-3-yl)boronic acid (Combi-Blocks, Inc.)
92-4		(*M*)-6-Chloro-7-(2-cyclopropylphenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-	None	**Step 3:** 2-cyclopropylphenylboronic acid (CombiPhos Catalysts, Inc.)

		piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one		
92-5		(*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(4-methyl-3-thienyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 4-methyl-3-thiopheneboronic acid (Combi-Blocks, Inc.)
92-6		(*M*)-6-Chloro-7-(cyclohexen-1-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** cyclohexen-1-ylboronic acid (Combi-Blocks Inc.)
92-7		(*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2,5-dimethyl-3-thienyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2,5-dimethylthiophene-3-boronic acid (Combi-Blocks Inc.)

92-8		(*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-methylprop-1-enyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2,2-dimethylethenylboronic acid (Synthonix Inc.)
92-9		(*M*)-6-Chloro-7-(3,6-dihydro-2H-pyran-4-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 3,6-dihydro-2*H*-pyran-4-boronic acid pinacol ester (Synthonix Inc.)
92-10		(*M*)-6-Chloro-7-(cyclohepten-1-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 1-cycloheptenylboronic acid pinacol ester (Combi-Blocks Inc.)
92-11		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(1*H*-indol-7-yl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-	None	**Step 3:** 1*H*-Indol-7-ylboronic acid pinacol ester (Combi-Blocks, San Diego, CA, USA)

		d]pyrimidin-2-one		
92-12		*(M)*-7-(2-Aminophenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-aniline (Oakwood Products, Inc.)
92-13		*(M)*-7-(2-Amino-3-pyridyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (2-Aminopyridi*N*-3-yl)boronic acid pinacol ester(Combi-Blocks, San Diego, CA, USA)
92-14		*(M)*-6-Chloro-7-[2-(difluoromethoxy)phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (2-(Difluoromethoxy)phenyl)boronic acid (Oxchem Corporation)
92-15		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-	None	**Step 3:** 6-Methyl-1h-indazole-7-boronic acid pinacol ester (J & W Pharmlab, LLC)

		pyridyl)-7-(6-methyl-1*H*-indazol-7-yl)pyrido[2,3-*d*]pyrimidin-2-one		
92-16		*(M)*-6-Chloro-7-[2-(difluoromethyl)phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2-[2-(Difluoromethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (Enamine)
92-17		*(M)*-7-(2-Amino-3-fluoro-phenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2-Fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (HDH Pharma)
92-18		*(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]phenyl]acetamide	None	**Step 3:** (2-Acetylaminophenyl)boronic acid (Combi-Blocks, San Diego, CA, USA)

92-19		*(M)*-6-Chloro-7-[2-(dimethylamino)phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** Dimethyl[2-(4,4,5,5-tetramethyl-1,3,2-dioxaborlan-2-yl)-phenyl]amine (Fluorochem Limited)
92-20		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-oxoindolin-7-yl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3-dihydroindol-2-one (Combi-Blocks, San Diego, CA, USA)
92-21		*(M)*-2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]benzamide	None	**Step 3:** 2-aminocarbonylphenylboronic acid (Combi-Blocks, San Diego, CA, USA)
92-22		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-8*H*-pyrido[2,3-*d*]pyrimidine-2,7-dione	None	side product when using 2-aminocarbonylphenylboronic acid (Combi-Blocks, San Diego, CA, USA)

92-23		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-[2-(trifluoromethyl)phenyl]pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2-(trifluoromethyl)phenylboronic acid (Combi-Blocks, San Diego, CA, USA)
92-24		*(M)*-2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-benzamide	None	**Step 3:** 2-carbamoyl-6-fluorophenylboronic acid pinacol ester (Combi-Blocks, San Diego, CA, USA)
92-25		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-[2-(methylamino)phenyl]pyrido[2,3-*d*]pyrimidin-2-one	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 2-(methylamino)phenylboronic acid (Matrix Scientific)
92-26		*(M)*-7-(2-Amino-3,5-difluoro-phenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** (2-amino-3,5-difluorophenyl)boronic acid pinacol ester (AstaTech, Inc)

		2,3-*d*]pyrimidin-2-one		
92-27		*(M)*-2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-*N*-methyl-benzamide	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 2-(methylcarbamoyl)benzeneboronic acid (Combi-Blocks, San Diego, CA, USA)
92-28		*(M)*-2-[6-Chloro-4-[(2S,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-N,N-dimethyl-benzamide	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 2-(*N*,*N*-dimethylaminocarbonyl)phenylboronic acid, pinacol ester (Combi-Blocks, San Diego, CA, USA)
92-29		*(M)*-2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]benzoic acid	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** (2-carboxyphenyl)boronic acid (Combi-Blocks, San Diego, CA, USA)
92-30		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(1*H*-indazol-7-yl)-1-(2-isopropyl-4-	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 1*H*-benzimidazol-4-ylboronic acid (Combi-Blocks, San Diego, CA, USA)

		methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one		
92-31		*(M)*-7-(2-Amino-4,5-difluoro-phenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 2-amino-4,5-difluorophenylboronic acid (Combi-Blocks, San Diego, CA, USA)
92-32		*(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]phenyl]methanesulfonamide	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** (2-(methylsulfonamido)phenyl)boronic acid (Combi-Blocks, San Diego, CA, USA)
92-33		*(M)*-2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]benzenesulfonamide	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** (2-sulfamoylphenyl)boronic acid (Combi-Blocks, San Diego, CA, USA)

92-34		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluoro-5-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 2-fluoro-5-hydroxyphenylboronic acid (Combi-Blocks, San Diego, CA, USA)
92-35		*(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]phenyl]-2,2-dimethyl-propanamide	**Step 3**: $PdCl_2(dppf)$, KOAc	**Step 3:** 2-(2,2-dimethyl-propionylamino)phenylboronic acid (Boron Molecular Inc.)
92-36		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-nitrophenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 4,4,5,5-tetramethyl-2-(2-nitrophenyl)-1,3,2-dioxaborolane (Combi-Blocks, San Diego, CA, USA)
92-37		*(M)*-7-(5-Bromo-2-fluoro-phenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-	None	**Step 3:** 2-(5-bromo-2-fluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (Combi-Blocks, San Diego, CA, USA)

		pyridyl)pyrido[2,3-*d*]pyrimidin-2-one		
92-38		*(M)*-7-(2-Amino-3,4,5-trifluoro-phenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (2-amino-3,4,5-trifluorophenyl)boronic acid pinacol ester (Aurum Pharmatech LLC)
92-39		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluoro-5-prop-1-ynyl-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (2-fluoro-5-(prop-1-yn-1-yl)phenyl)boronic acid (ChemShuttle)
92-40		*(M)*-2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-benzonitrile	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 3-fluoro-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzonitrile (Combi-Blocks, San Diego, CA, USA)

92-41		*(M)*-7-(2-Acetylphenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (2-acetylphenyl)boronic acid (Aurum Pharmatech Inc.)
92-42		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-phenyl-pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** phenylboronic acid (Sigma-Aldrich Corporation)
92-43		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-[2-[(1RS)-1-hydroxyethyl]phenyl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (2-(1-hydroxyethyl)phenyl)boronic acid (HDH Pharma, Inc.)
92-44		*(M)*-6-Chloro-7-(5-chloro-2-fluoro-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-	None	**Step 3:** 5-chloro-2-fluorophenylboronic acid (Sigma-Aldrich Corporation)

		d]pyrimidin-2-one		
92-45		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** *o*-tolylboronic acid (Sigma-Aldrich Corporation)
92-46		*(M)*-6-Chloro-7-(3-chloro-2-fluoro-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2-fluoro-3-chlorophenylboronic acid (Combi-Blocks Inc.)
92-47		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-ethynylphenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (2-ethynylphenyl)boronic acid (Combi-Blocks Inc.)
92-48		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(3-ethynylphenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-	None	**Step 3:** (3-ethynylphenyl)boronic acid pinacol ester (Combi-Blocks Inc.)

		d]pyrimidin-2-one		
92-49		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-prop-1-ynylphenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** [3-(prop-1-yn-1-yl)phenyl]boronic acid (Enamine)
92-50		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-thienyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 3-thienylboronic acid (Maybridge Chemical Co., Ltd.)
92-51		*(M)*-6-Chloro-7-(2-chlorophenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2-chlorophenylboronic acid (Matrix Scientific)
92-52		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(3-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-	None	**Step 3:** 3-fluorophenylboronic acid (Steris Corporation)

		d]pyrimidin-2-one		
92-53		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(m-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (3-methylphenyl)boranediol (Sigma-Aldrich Corporation)
92-54		*(M)*-6-Chloro-7-(3-chlorophenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 3-chlorophenylboronic acid (AstaTech, Inc)
92-55		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-ethylphenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (2-ethylphenyl)boronic acid (Combi-Block Inc.)
92-56		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-[3-(trifluoromethyl)phenyl]pyrido[2,3-	None	**Step 3:** 3-(trifluoromethyl)phenylboronic acid (Sigma-Aldrich Corporation)

		d]pyrimidin-2-one		
92-57		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(4-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 4-fluorobenzeneboronic acid (Sigma-Aldrich Corporation)
92-58		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluoro-3-methyl-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2-fluoro-3-methylphenylboronic acid (Combi-Blocks Inc.)
92-59		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(3-fluoro-2-methyl-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 3-fluoro-2-methylphenylboronic acid (Combi-Blocks Inc.)
92-60		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluoro-5-methyl-phenyl)-1-(2-isopropyl-4-methyl-3-	None	**Step 3:** (2-fluoro-5-methylphenyl)boranediol (Sigma-Aldrich Corporation)

		pyridyl)pyrido[2,3-*d*]pyrimidin-2-one		
92-61		*(M)*-6-Chloro-7-(2,4-difluorophenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2,4-difluorophenylboronic acid pinacol ester (Sigma-Aldrich Corporation)
92-62		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(3-fluoro-2-methoxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 3-fluoro-2-methoxyphenylboronic acid (Asymchem Laboratories, Inc.)
92-63		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-[3-fluoro-2-(trifluoromethyl)phenyl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (3-fluoro-2-(trifluoromethyl)phenyl)boronic acid (Combi-Blocks)

92-64		*(M)*-6-Chloro-7-(2-chloro-3-fluoro-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2-chloro-3-fluorophenylboronic acid (Combi-Blocks Inc.)
92-65		*(M)*-6-Chloro-7-(2,5-difluorophenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2,5-difluorophenylboronic acid pinacol ester (Combi-Blocks Inc.)
92-66		*(M)*-6-Chloro-7-(2,3-difluorophenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2,3-difluorophenylboronic acid pinacol ester (Combi-Blocks Inc.)
92-67		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-methoxyphenyl)pyrido[2,3-	None	**Step 3:** 2-methoxyphenylboronic acid (TCI America)

		d]pyrimidin-2-one		
92-68		(*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(4-isopropylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 4-isopropylpyrimidine-5-boronic acid (Combi-Blocks Inc.)
92-69		*(M)*-6-Chloro-7-(2-cyclopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (2-cyclopropylpyridin-3-yl)boronic acid (Activate Scientific)
92-70		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-methoxy-1-naphthyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (3-methoxynaphthalen-1-yl)boronic acid (Ark Pharm, Inc.)
92-71		*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2,5-dimethylthiazol-4-yl)-1-(2-	None	**Step 3:** 2,5-dimethylthiazole-4-boronic acid pinacol ester (CombiPhos Catalysts, Inc.)

		isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one		
92-72		(M)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(4-methylthiazol-5-yl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3-thiazole (CombiPhos Catalysts, Inc.)

Method 93

Example 93-1: (*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-methyl-2-thienyl)pyrido[2,3-*d*]pyrimidin-2-one



Intermediate 92B

[0185] (*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-methyl-2-thienyl)pyrido[2,3-*d*]pyrimidin-2-one. A mixture of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Intermediate 92B; 0.15 g, 0.29 mmol), 3-methylthiophene-2-boronic acid (0.06 g, 0.44 mmol, Combi-Blocks Inc.), sodium carbonate (0.094 g, 0.89 mmol), and Pd(AmPhos)Cl_2 (0.013 g, 0.018 mmol, Sigma-Aldrich Corporation) in 1,4-dioxane (1 mL) and water (0.5 mL) was stirred and heated at 100 °C for 16 h. The crude material was purified by silica gel chromatography (eluent: 0-60% EtOAc-EtOH (3:1)/heptane) to provide (*M*)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-methyl-2-thienyl)pyrido[2,3-

d]pyrimidin-2-one (0.010 g, 0.018 mmol, 6.1 % yield) as yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.45 (d, *J*=4.8 Hz, 1 H), 8.39 (d, *J*=3.9 Hz, 1 H), 7.66 (d, *J*=5.0 Hz, 1 H), 7.23 (d, *J*=5.0 Hz, 1 H), 6.94 (d, *J*=5.2 Hz, 1 H), 6.83 (td, *J*=17.2, 10.7 Hz, 1 H), 6.19 (dd, *J*=16.7, 1.3 Hz, 1 H), 5.71 - 5.78 (m, 1 H), 4.41 - 4.94 (m, 2 H), 3.42 - 4.23 (m, 4 H), 2.63 - 2.74 (m, 1 H), 1.94 (s, 3 H), 1.83 (s, 3 H), 1.33 (t, *J*=6.5 Hz, 3 H), 1.21 (br dd, *J*=27.7, 6.5 Hz, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.94 (d, *J*=6.6 Hz, 3 H). *m/z* (ESI, +ve ion): 577.3 $(M+H)^+$.

Table 93: Compound 93-2 was prepared following the procedure described in Method 93 above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
93-2		(*M*)-6-chloro-7-[(1*SR*)-2,2-dimethylcyclopropyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	1,1-dimethylcyclopropyl-2-boronic acid pinacol ester (Aurum Pharmatech LLC)

Method 94

Example 94-1: (*M*)-4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one



[0186] Step 1: (*M*)-*tert*-Butyl (2*R*,5*S*)-4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of (*M*)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 72A**; 3.35 g, 9.61 mmol), phosphorous oxychloride (1.07 mL, 11.53 mmol), and DIPEA (5.02 mL, 28.8 mmol) in acetonitrile (24 mL) was stirred at 80 °C for 30 min. The reaction mixture was cooled to rt and added (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (2.26 g, 10.6 mmol). The reaction mixture was stirred at rt for 15 min. The reaction mixture was diluted with EtOAc (200 mL), washed with satd. $NaHCO_3$ (3 × 75 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-100% EtOAc/heptane) to provide (*M*)-*tert*-butyl (2*R*,5*S*)-4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 94A**; 2.2 g, 4.04 mmol, 42.0 % yield) as yellow-orange solid. ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 8.57 (d, *J*=5.0 Hz, 1 H), 7.79 (br d, *J*=7.5 Hz, 1 H), 7.13 - 7.16 (m, 1 H), 4.80 - 5.04 (m, 1 H), 4.32 - 4.64 (m, 1 H), 3.73 - 4.08 (m, 3 H), 3.43 - 3.66 (m, 1 H), 2.58 (dt, *J*=13.4, 6.6 Hz, 1 H), 2.04 (s, 3 H), 1.59 (s, 9 H), 1.39 - 1.48 (m, 3 H), 1.11 - 1.25 (m, 9 H). ^{19}F NMR (377 MHz, *CHLOROFORM-d*) δ -126.30 (br s, 1 F) -126.34 (br s, 1 F). *m/z* (ESI, +ve ion): 544.8 $(M+H)^+$.

[0187] Step 2: (*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A mixture of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 94A**; 8.16 g, 14.97 mmol) in DCM (30 mL) and TFA (30 mL) was stirred at rt for 30 min. The reaction mixture was concentrated *in vacuo* to provide (*M*)-7-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. The crude product was used in next step without purification. *m/z* (ESI, +ve ion) 445.1 $(M+H)^+$.

[0188] A mixture of (*M*) 7-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (13.1 mL, 74.9 mmol) in DCM (20 mL) was cooled to 0 °C. To the cooled mixture was added acryloyl chloride (1.22 mL, 14.97 mmol) and stirred for 1 h. The reaction mixture was quenched with satd. NH_4Cl (50 mL), extracted with DCM (2 × 50 mL)), dried over MgSO4, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-100% EtOAc-MeOH (9:1)/heptane) to provide (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 94B;** 5.62 g, 11.3 mmol, 75 % yield) as light-yellow foam. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.53 (d, *J*=5.0 Hz, 1 H), 8.44 (d, *J*=8.5 Hz, 1 H), 7.35 (br d, *J*=3.7 Hz, 1 H), 6.83 (td, *J*=16.9, 10.6 Hz, 1 H), 6.18 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.69 - 5.80 (m, 1 H), 4.65 - 4.92 (m, 2 H), 3.79 - 4.15 (m, 3 H), 3.07 - 3.69 (m, 1 H), 2.60 - 2.72 (m, 1 H), 1.98 (s, 3 H), 1.20 - 1.33 (m, 6 H), 1.06 (dd, *J*=13.1, 6.8 Hz, 6 H). *m/z* (ESI, +ve ion): 499.2 $(M+H)^+$.

[0189] Step 3: (*M*)-4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one. A mixture of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 94B**; 0.21 g, 0.42 mmol), 2-isopropylphenylboronic acid (0.10 g, 0.62 mmol, Combi-Blocks Inc.), sodium carbonate (0.13 g, 1.25 mmol), and tetrakis(triphenylphosphine)palladium(0) (0.048 g, 0.042 mmol, Sigma-Aldrich Corporation) in 1,4-dioxane (1.2 mL) and water (0.6 mL) was stirred and heated at 90 °C. The crude material was purified by silica gel chromatography (eluent: 0-50% of EtOAc-EtOH (3:1)/heptane) to provide (*M*)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-

isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one (0.037 g, 0.063 mmol, 15% yield) as off-white solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.37 (d, *J*=4.8 Hz, 1 H), 8.30 (dd, *J*=9.4, 3.2 Hz, 1 H), 7.38 - 7.44 (m, 2 H), 7.21 - 7.27 (m, 1 H), 7.18 (d, *J*=5.0 Hz, 1 H), 7.12 (d, *J*=7.7 Hz, 1 H), 6.77 - 6.92 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.73 - 5.78 (m, 1 H), 4.82 - 4.93 (m, 1 H), 4.43 - 4.81 (m, 1 H), 3.44 - 4.27 (m, 4 H), 2.69 (quin, *J*=6.4 Hz, 2 H), 1.92 (s, 3 H), 1.33 (t, *J*=6.3 Hz, 3 H), 1.18 - 1.30 (m, 3 H), 0.87 - 1.09 (m, 12 H). ^{19}F NMR (376 MHz, *DMSO-d*$_6$) δ -129.20 (d, *J*=10.4 Hz, 1 F). *m/z* (ESI, +ve ion): 582.8 $(M+H)^+$.

Table 94: Compounds 94-2 to 94-20 were prepared following the procedure described in Method 94, Steps 1-3, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
94-2	 Single isomer (M)	(*M*)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-[2-(hydroxymethyl)phenyl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	Step 3: 1,3-dihydro-1-hydroxy-2,1-benzoxaborole
94-3	 Single isomer (M)	(*M*)-7-(2,5-difluorophenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** (2,5-difluorophenyl)boronic acid (Combi-Blocks)

94-4		(*M*)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-5-methyl-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** (2-fluoro-5-methylphenyl)boranediol
94-5		(*M*)-7-(5-chloro-2-fluoro-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 5-chloro-2-fluorophenyl boronic acid
94-6		(*M*)- 7-(2-chloro-5-methoxy-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 2-chloro-5-methoxyphenyl boronic acid (Combi Blocks)

94-7		(*M*)-7-(5-cyclopropyl-2-fluoro-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 5-cyclopropyl-2-fluorophenyl boronic acid (Combi-Blocks)
94-8	 Single isomer (*M*)	(*M*)-4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 3**: $PdCl_2$(dppf), KOAc, 100 °C	**Step 3**: *o*-tolylboronic acid; Sigma-Aldrich, St. Louis, MO
94-9	 Single isomer (*M*)	(*M*)-7-(2,6-Dimethylphenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	Switch **Step 2** & **Step 3**; **Step 3**: $PdCl_2$(dppf), KOAc, 100 °C	**Step 3**: 2,6-dimethylphenyl boronic acid; Sigma-Aldrich, St. Louis, MO
94-10	 Single isomer (*M*)	(*M*)-4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-methyl-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-	Switch **Step 2** & **Step 3**; **Step 3**: $PdCl_2$(dppf), KOAc, 100 °C	**Step 3**: 2-fluoro-6-dimethylphenyl boronic acid; Combi-Blocks, Inc., San Diego, CA

		d]pyrimidin-2-one		
94-11	 Single isomer (*M*)	(*M*)-7-(2,6-Difluorophenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	Switch **Step 2** & **Step 3**; **Step 3**: $PdCl_2$(dppf), KOAc, 100 °C	**Step 3**: 2,6-difluorophenyl boronic acid; Sigma-Aldrich, St. Louis, MO
94-12	 Single isomer (*M*)	(*M*)-7-(2-Chloro-6-fluoro-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	Switch **Step 2** & **Step 3**; **Step 3**: $PdCl_2$(dppf), KOAc, 100 °C	**Step 3**: 2-chloro-6-fluorophenyl boronic acid; Sigma-Aldrich, St. Louis, MO
94-13		*(M)*-7-[2-(Difluoromethyl)phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2-[2-(difluoromethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (Enamine Ltd.)
94-14		*(M)*-7-(1*H*-Benzimidazol-4-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 1*H*-benzimidazol-4-ylboronic acid (Combi-Blocks, San

		6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one		Diego, CA, USA)
94-15		*(M)*-2-[4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]benzoic acid	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** (2-carboxyphenyl)boronic acid (Combi-Blocks, San Diego, CA, USA
94-16		*(M)*-2-[4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]benzoic acid	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** (2-carboxyphenyl)boronic acid (Combi-Blocks, San Diego, CA, USA
94-17		*(M)*-2-[4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]benzamide	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 2-aminocarbonylphenylboronic acid (Combi-Blocks, San Diego, CA, USA)
94-18		*(M)*-2-[4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-	**Step 3**: $Pd(dppf)Cl_2$ KOAc	**Step 3:** 2-(methylcarbamoyl)benzeneboronic acid (Combi-Blocks, San

		pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-*N*-methyl-benzamide		Diego, CA, USA)
94-19		*(M)*-2-[4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-benzamide	None	**Step 3:** 2-carbamoyl-6-fluorophenyl boronic acid pinacol ester (Combi-Blocks, San Diego, CA, USA)
94-20		*(M)*-7-(2-Aminophenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-aniline (Oakwood Products, Inc.)
94-21		*(M)*-4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-methoxy-1-naphthyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 3:** (3-methoxynaphthalen-1-yl)boronic acid (Ark Pharm, Inc.)

Method 95

Example 95-1: (*M*)-6-Chloro-7-cyclohexyl-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one



[0190] Step 1: (*M*)-*tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-cyclohexyl-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 92A**; 0.31 g, 0.55 mmol), Pd(AmPhos)Cl_2 (0.020 g, 0.028 mmol, Sigma-Aldrich Corporation), and cyclohexylzinc bromide, 0.5 M in THF (2.20 mL, 1.10 mmol, Rieke Metals, Inc.) in THF (1 mL) was stirred and heated at 80 °C for 4 h. The reaction mixture was purified by silica gel chromatography (eluent: 0-50% of EtOAc-EtOH (3:1)/heptane) to provide (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-cyclohexyl-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.13 g, 0.22 mmol, 39.7 % yield) as light-yellow syrupy solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.48 (d, *J*=5.0 Hz, 1 H), 8.22 (s, 1 H), 7.25 (d, *J*=5.0 Hz, 1 H), 4.76 (br d, *J*=1.5 Hz, 1 H), 4.30 (br d, *J*=38.8 Hz, 1 H), 4.09 (br d, *J*=13.7 Hz, 1 H), 3.76 (br dd, *J*=9.7, 2.5 Hz, 1 H), 3.68 (br d, *J*=13.5 Hz, 1 H), 3.34 - 3.56 (m, 1 H), 3.01 (tt, *J*=11.1, 3.0 Hz, 1 H), 2.67 (quin, *J*=6.7 Hz, 1 H), 1.85 (s, 3 H), 1.50 - 1.75 (m, 6 H), 1.44 (s, 9 H), 1.31 (d, *J*=6.6 Hz, 3 H), 1.19 - 1.29 (m, 4 H), 1.15 (br d, *J*=6.4 Hz, 3 H), 1.09 (d, *J*=6.6 Hz, 3 H), 0.97 (d, *J*=6.8 Hz, 3 H). *m/z* (ESI, +ve ion): 609.3 $(M+H)^+$.

[0191] Step 2: (*M*)-6-Chloro-7-cyclohexyl-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one. A mixture of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-cyclohexyl-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.13 g, 0.22 mmol) in DCM (2 mL) and TFA (2 mL) was stirred at rt for 20 min. The mixture was concentrated *in vacuo* to give (*M*)-6-chloro-7-cyclohexyl-4-((2*S*,5*R*)-2,5-

dimethylpiperazin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as orange syrup. *m/z* (ESI, +ve ion): 509.1 $(M+H)^+$.

[0192] A mixture of (*M*)-6-chloro-7-cyclohexyl-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (0.57 mL, 3.3 mmol) in DCM (2 mL) was treated with acryloyl chloride, 0.2 M solution in DCM (1.19 mL, 0.24 mmol) at 0 °C and stirred for 30 min. The reaction mixture was quenched with satd. NH_4Cl (50 mL), extracted with EtOAc (2 × 50 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-60% EtOAc-EtOH (3:1)/heptane) to provide (*M*)-6-chloro-7-cyclohexyl-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one (0.05 g, 0.089 mmol, 41 % yield) as white solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.48 (d, *J*=4.8 Hz, 1 H), 8.25 (d, *J*=4.1 Hz, 1 H), 7.25 (d, *J*=4.8 Hz, 1 H), 6.82 (td, *J*=17.5, 10.5 Hz, 1 H), 6.18 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.71 - 5.77 (m, 1 H), 4.40 - 4.89 (m, 2 H), 3.37 - 4.19 (m, 4 H), 3.02 (tt, *J*=11.0, 2.9 Hz, 1 H), 2.63 - 2.72 (m, 1 H), 1.86 (s, 3 H), 1.50 - 1.68 (m, 5 H), 1.13 - 1.34 (m, 9 H), 1.03 (dd, *J*=45.3, 6.7 Hz, 8 H). *m/z* (ESI, +ve ion): 563.3 $(M+H)^+$.

Table 95: Compounds 95-2 to 95-7 were prepared following the procedure described in Method 95, Steps 1-2, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
95-2		(*M*)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2,2-dimethylpropyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** neopentylzinc bromide, 0.5 M in THF (Rieke)

95-3		(*M*)-6-chloro-7-cyclopentyl-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** cyclopentylzinc bromide, 0.5 M in THF (Sigma-Aldrich Corporation)
95-4		(*M*)-6-chloro-7-cyclopropyl-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** cyclopropylzinc bromide, 0.5 M in THF (Sigma-Aldrich Corporation)
95-5		(*M*)-6-chloro-7-cyclobutyl-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** 1,1-dimethylcyclopropyl-2-boronic acid pinacol ester (Aurum Pharmatech LLC)
95-6		(*M*)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-propyl-pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** n-propylzinc bromide, 0.5 M in THF (Sigma-Aldrich Corporation)

95-7		(*M*)-6-chloro-7-(3,3-dimethylbut-1-ynyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** 3,3-dimethyl-1-butyne (Enamine)

Method 96

Example 96-1: (*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-methyl-2-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one



Intermediate 92B

[0193] **(*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-methyl-2-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one**. A mixture of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 92B**; 0.2 g, 0.39 mmol), 3-methyl-2-(tributylstannyl)pyridine (0.17 g, 0.44 mmol, Indofine Chemical Company, Inc), copper iodide (7.4 mg, 0.039 mmol, Sigma-Aldrich Corporation), and tetrakis(triphenylphosphine)palladium(0) (0.022 g, 0.019 mmol, Sigma-Aldrich Corporation) in 1,4-dioxane (2 mL) was stirred and heated in a Initiator microwave reactor (Personal Chemistry, Biotage AB, Inc., Upssala, Sweden) at 150 °C for 30 min. The crude material was purified by silica gel chromatography (eluent: 0-50% DCM-MeOH (4:1)/DCM) to provide (*M*)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-methyl-2-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one (0.031 g, 0.054 mmol, 14 % yield) as yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.47 (d, *J*=4.8 Hz, 1 H), 8.41 (d,

J=3.9 Hz, 1 H), 8.38 (d, *J*=5.0 Hz, 1 H), 7.70 (d, *J*=7.7 Hz, 1 H), 7.35 (dd, *J*=7.9, 4.8 Hz, 1 H), 7.19 (d, *J*=4.8 Hz, 1 H), 6.84 (td, *J*=17.1, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.1 Hz, 1 H), 5.73 - 5.78 (m, 1 H), 4.43 - 4.96 (m, 2 H), 4.13 - 4.27 (m, 1 H), 3.43 - 3.93 (m, 3 H), 2.72 (tt, *J*=6.6, 3.2 Hz, 1 H), 1.95 (s, 3 H), 1.91 (s, 3 H), 1.35 (t, *J*=6.2 Hz, 3 H), 1.18 - 1.28 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.92 (d, *J*=6.6 Hz, 3 H). *m/z* (ESI, +ve ion): 572.2 $(M+H)^+$.

Table 96: Compound 96-2 was prepared following the procedure described in Method 96 above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
96-2		(*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-methoxy-2-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	3-methoxy-2-(tributylstannyl)pyridine (Combi-Blocks Inc.)

Method 100

Example 100-1: 6-Chloro-1-(2,4-diisopropylpyrazol-3-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one



[0194] Step 1: 2,5,6-Trichloro-*N*-((1,4-diisopropyl-1H-pyrazol-5-yl)carbamoyl)nicotinamide. To a mixture of 2,5,6-trichloronicotinamide (**Intermediate P**; 1.4 g, 6.1 mmol) in THF (12 mL) was added oxalyl chloride, 2 M solution in DCM (3.3 mL, 6.5 mmol) at rt and the mixture was stirred and heated at 65 °C for 3 h. The mixture was cooled to 0 °C. To the cooled mixture was added a solution of 1,4-diisopropyl-1*H*-pyrazol-5-amine (**Intermediate 161**; 1.03 g, 6.13 mmol) in THF (5 mL) and the mixture was stirred at rt for 15 h. The reaction mixture was quenched with satd $NaHCO_3$ (100 mL), extracted with EtOAc (2 × 50 mL), washed with brine (1 × 50 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (eluent: 0-50% EtOAc-

EtOH (3:1)/heptane) to provide 2,5,6-trichloro-*N*-((1,4-diisopropyl-1*H*-pyrazol-5-yl)carbamoyl)nicotinamide (2.43 g, 5.79 mmol, 94 % yield) as orange syrup. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 11.36 (br s, 1 H), 9.39 (br s, 1 H), 8.63 (s, 1 H), 7.33 (s, 1 H), 4.37 (dt, *J*=13.3, 6.6 Hz, 1 H), 2.64 - 2.75 (m, 1 H), 1.33 (d, *J*=6.4 Hz, 6 H), 1.13 (d, *J*=7.0 Hz, 6 H). *m/z* (ESI, +ve ion): 417.6 $(M+H)^+$.

[0195] Step 2: 6,7-Dichloro-1-(1,4-diisopropyl-1H-pyrazol-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a cooled mixture of 2,5,6-trichloro-*N*-((1,4-diisopropyl-1*H*-pyrazol-5-yl)carbamoyl)nicotinamide (2.42 g, 5.78 mmol) in THF (19 mL) at 0 °C was added 1 M KHMDS in THF (14.5 mL, 14.5 mmol) dropwise and the mixture was stirred at 0 °C for 30 min. The mixture was quenched with satd. NH_4Cl (50 mL) and brine (50 mL), extracted with EtOAc (2 × 50 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (eluent: 0-50% EtOAc/heptane) to provide 6,7-dichloro-1-(1,4-diisopropyl-1H-pyrazol-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.16 g, 0.41 mmol, 7 % yield) as yellow syrupy solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 12.26 (br s, 1 H), 8.57 (s, 1 H), 7.52 (s, 1 H), 4.32 (spt, *J*=6.6 Hz, 1 H), 2.53 - 2.61 (m, 1 H), 1.27 (d, *J*=6.6 Hz, 3 H), 1.22 (d, *J*=6.4 Hz, 3 H), 1.04 (d, *J*=6.8 Hz, 3 H), 0.99 (d, *J*=6.8 Hz, 3 H). *m/z* (ESI, +ve ion): 381.6 $(M+H)^+$.

[0196] Step 3 : *tert*-Butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(1,4-diisopropyl-1*H*-pyrazol-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a mixture of 6,7-dichloro-1-(1,4-diisopropyl-1*H*-pyrazol-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.15 g, 0.39 mmol) and DIPEA (0.34 mL, 2.0 mmol) in acetonitrile (2.0 mL) was added phosphorus oxychloride (0.070 mL, 0.79 mmol) and the mixture was stirred and heated at 80 °C for 1 h. The mixture was cooled to 0 °C. To the cooled mixture was added DIPEA (0.34 mL, 1.96 mmol) followed by (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (0.13 g, 0.59 mmol, AstaTech Inc.) and the mixture was stirred at 0 °C for 1 h. The reaction mixture was quenched with satd $NaHCO_3$ (50 mL), extracted with EtOAc (2 × 50 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (eluent: 0-25% EtOAc-EtOH(3:1)/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(1,4-diisopropyl-1*H*-pyrazol-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.15 g, 0.26 mmol, 66.7 % yield) as orange solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.44 (d, *J*=45.0 Hz, 1 H), 7.47 (d, *J*=7.3 Hz, 1 H), 3.33 - 4.89 (m, 7 H), 2.38 - 2.47 (m, 1 H), 0.94 - 1.47 (m, 27 H). *m/z* (ESI, +ve ion): 577.7 $(M+H)^+$.

[0197] Step 4 : *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(1,4-diisopropyl-1H-pyrazol-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(1,4-diisopropyl-1H-pyrazol-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.15 g, 0.25 mmol), 2-fluorophenylboronic acid (0.053 g, 0.38 mmol, Combi-Blocks Inc.), sodium carbonate (0.080 g, 0.76 mmol), tetrakis(triphenylphosphine)palladium(0) (0.029 g, 0.025 mmol, Sigma-Aldrich Corporation) in 1,4-dioxane (1 mL) and water (0.25 mL) was stirred and heated at 80 °C for 2 h. The crude material was purified by silica gel chromatography (eluent: 0-25% EtOAc-EtOH (3:1)/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(1,4-diisopropyl-1*H*-pyrazol-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.16 g, 0.25 mmol, quantitative yield) as yellow syrup. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.42 (d, *J*=25.9 Hz, 1 H), 7.49 - 7.66 (m, 1 H), 7.21 - 7.41 (m, 4 H), 3.35 - 4.90 (m, 7 H), 2.39 - 2.47 (m, 1 H), 1.45 (s, 9 H), 1.09 - 1.39 (m, 12 H), 0.98 (dd, *J*=9.2, 6.9 Hz, 3 H), 0.93 (dd, *J*=6.8, 3.1 Hz, 3 H). ^{19}F NMR (376 MHz, *DMSO-d*$_6$) δ -114.44 - -114.14 (m, 1 F). *m/z* (ESI, +ve ion): 638.2 $(M+H)^+$.

[0198] Step 5 : 6-Chloro-1-(2,4-diisopropylpyrazol-3-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one. A mixture of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(1,4-diisopropyl-1*H*-pyrazol-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.16 g, 0.25 mmol) in DCM (2.5 mL) and TFA (2.5 mL) was stirred at rt for 1 h. The mixture was concentrated *in vacuo* to afford 6-chloro-1-(1,4-diisopropyl-1*H*-pyrazol-5-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as yellow syrup. The crude product was used in next step without purification. *m/z* (ESI, +ve ion): 539.3 $(M+H)^+$.

[0199] A mixture of 6-chloro-1-(1,4-diisopropyl-1*H*-pyrazol-5-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (0.66 mL, 3.78 mmol) in DCM (2.5 mL) was cooled to 0 °C. To the cooled mixture was added acryloyl chloride (0.020 mL, 0.25 mmol) dropwise and the mixture was stirred at 0 °C for 10 min. The reaction mixture was quenched with satd. NH_4Cl (50 mL), extracted with EtOAc (2 × 50 mL), washed with satd $NaHCO_3$ (1 × 50 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (eluent: 0-50% DCM-EtOH (4:1)/DCM) to provide 6-chloro-1-(2,4-diisopropylpyrazol-3-yl)-4-

[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one (0.093 g, 0.16 mmol, 63 % yield) as yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.38 - 8.50 (m, 1 H), 7.50 - 7.59 (m, 1 H), 7.22 - 7.42 (m, 4 H), 6.75 - 6.91 (m, 1 H), 6.19 (dd, *J*=16.8, 2.1 Hz, 1 H), 5.71 - 5.79 (m, 1 H), 4.41 - 4.97 (m, 2 H), 3.38 - 4.33 (m, 5 H), 2.39 - 2.48 (m, 1 H), 1.11 - 1.37 (m, 12 H), 0.99 (dd, *J*=9.4, 6.9 Hz, 3 H), 0.93 (dd, *J*=6.9, 2.8 Hz, 3 H). ^{19}F NMR (376 MHz, *DMSO-d*$_6$) δ -114.42 - -114.15 (m, 1 F). *m/z* (ESI, +ve ion): 592.3 $(M+H)^+$.

Table 100: Compounds 100-2 to 100-9 were prepared following the procedure described in Method 100, Steps 1-5, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
100-2		4-[6-Chloro-1-(4,6-diisopropylpyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-benzonitrile	None	**Step 1: Intermediate U** **Step 4:** 4-Cyano-2-fluorophenylboronic acid pinacol ester, Combi-Blocks, San Diego, CA, USA
100-3		2-[6-Chloro-1-(4,6-diisopropylpyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]benzonitrile	None	**Step 1: Intermediate U** **Step 4:** 2-cyanophenylboronic acid, pinacol ester, Combi-Blocks, San Diego, CA, USA
100-4		6-Chloro-1-(4,6-diisopropylpyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1: Intermediate U** **Step 4:** 2-isopropylphenyl boronic acid, Combi-Blocks, San Diego, CA, USA

100-5		6,7-Dichloro-1-(2,2-dimethylpropyl)-4-(4-prop-2-enoylpiperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2-one	Omit **step 4&5**	**Step 1:** Neopentylamine (TCI America) **Step 3:** 1-Boc-piperazine, (Sigma-Aldrich Corporation)
100-6		1-(2,2-dimethylpropyl)-4-(4-prop-2-enoylpiperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2-one	Add a reduction step (H_2, Pd/C, NaOMe, THF/MeOH, 55 psi, RT) before Step 3, Omit **step 4&5**	**Step 1:** Neopentylamine (TCI America) **Step 3:** 1-Boc-piperazine, (Sigma-Aldrich Corporation)
100-7		6-Chloro-1-(4,6-diisopropylpyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-methylsulfonylphenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 4**: 95 °C	**Step 1**: Intermediate U **Step 4**: (2-methylsulfonylphenyl) boronic acid (Sigma-Aldrich Corporation)
100-8	 First-eluting isomer	6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(3-isopropyl-5-methyl-4-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 3 (1):** Toluene, 50 °C; **Step 4:** $Pd(dppf)Cl_2$, KOAc; Chiral separation after **Step 5**	**Step 1: Intermediate 222**

100-9	 Second-eluting isomer	6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(3-isopropyl-5-methyl-4-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 3 (1):** Toluene, 50 °C; **Step 4:** $Pd(dppf)Cl_2$, KOAc; Chiral separation after **Step 5**	**Step 1: Intermediate 222**

Method 101

Example 101-1: 1-(3,5-Diisopropyl-4-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)pyrido[2,3-*d*]pyrimidin-2-one



[0200] Step 1: 2,6-Dichloro-*N*-((3,5-diisopropylpyridin-4-yl)carbamoyl)-5-fluoronicotinamide. To a mixture of 2,6-dichloro-5-fluoronicotinamide (**Intermediate S**; 1.58 g, 7.57 mmol) in THF (15 mL) was added oxalyl chloride, 2 M solution in DCM (3.99 mL, 7.97 mmol) at rt and the mixture was stirred and heated at 65 °C for 1 h. The mixture was cooled to rt. To the mixture was added a solution of 3,5-diisopropylpyridin-4-amine (**Intermediate 162**; 1.35 g, 7.57 mmol) in THF (8 mL) and the mixture was stirred at rt for 1 h. The mixture was quenched with satd $NaHCO_3$ (100 mL), extracted with EtOAc (3 × 50 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (eluent: 0-40% EtOAc-EtOH(3:1)/heptane) to provide 2,6-dichloro-*N*-((3,5-diisopropylpyridin-4-yl)carbamoyl)-5-fluoronicotinamide (0.54 g, 1.3 mmol, 17 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 11.39 (br s, 1 H), 9.40 - 9.81 (m, 1 H), 8.54 (br d, *J*=7.9 Hz, 1 H), 8.45 (s, 2 H), 3.09 (spt, *J*=6.7 Hz, 2 H), 1.22 (d, *J*=6.8 Hz, 12 H). ^{19}F NMR (376 MHz, *DMSO-d*$_6$) δ -122.48 - -121.38 (m, 1 F). *m/z* (ESI, +ve ion): 413.0 $(M+H)^+$.

[0201] Step 2: 7-Chloro-1-(3,5-diisopropylpyridin-4-yl)-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a cooled mixture of 2,6-dichloro-*N*-((3,5-diisopropylpyridin-4-yl)carbamoyl)-5-fluoronicotinamide (0.54 g, 1.3 mmol) in THF (6 mL) at 0 °C was added 1 M KHMDS in THF (2.58 mL, 2.58 mmol) dropwise and the mixture was stirred at rt for 3 h. The mixture was quenched with satd NH_4Cl (50 mL), extracted with EtOAc (3 × 50 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (eluent: 0-25% EtOAc-EtOH(3:1)/heptane) to provide 7-chloro-1-(3,5-diisopropylpyridin-4-yl)-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.23 g, 0.62 mmol, 48 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 12.28 (s, 1 H), 8.62 (s, 2 H), 8.50 (d, *J*=7.5 Hz, 1 H), 2.65 - 2.78 (m, 2 H), 1.15 (d, *J*=7.0 Hz, 6 H), 1.03 (d, *J*=6.8 Hz, 6 H). ^{19}F NMR (377 MHz, *DMSO-d*$_6$) δ -126.68 (s, 1 F). *m/z* (ESI, +ve ion): 377.0 $(M+H)^+$.

[0202] Step 3: *tert*-Butyl (2*R*,5*S*)-4-(7-chloro-1-(3,5-diisopropylpyridin-4-yl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a mixture of 7-chloro-1-(3,5-diisopropylpyridin-4-yl)-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.23 g, 0.61 mmol) and DIPEA (0.53 mL, 3.05 mmol) in acetonitrile (4 mL) was added phosphorus oxychloride (0.11 mL, 1.22 mmol) and the mixture was stirred and heated at 80 °C for 30 min. The reaction mixture was cooled to 0 °C. To the cooled mixture was added DIPEA (0.53 mL, 3.05 mmol) followed by (2*R*,5*S*)-*tert*-butyl 2,5-

dimethylpiperazine-1-carboxylate (0.2 g, 0.92 mmol, eNovation Chemicals LLC) and the mixture was stirred at rt for 2 h. The reaction mixture was quenched with satd. $NaHCO_3$ (100 mL), extracted with EtOAc (3 × 50 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (eluent: 0-25% EtOAc-EtOH(3:1)/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(7-chloro-1-(3,5-diisopropylpyridin-4-yl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.14 g, 0.25 mmol, 40.7 % yield) as orange solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.59 (d, *J*=12.6 Hz, 2 H), 8.38 (d, *J*=8.5 Hz, 1 H), 4.80 (br s, 1 H), 4.14 - 4.40 (m, 1 H), 4.03 (br d, *J*=13.5 Hz, 1 H), 3.84 (br d, *J*=12.4 Hz, 1 H), 3.66 (br d, *J*=13.3 Hz, 1 H), 3.50 (br dd, *J*=31.9, 13.5 Hz, 1 H), 2.40 - 2.49 (m, 2 H), 1.44 (s, 9 H), 1.30 (d, *J*=6.6 Hz, 3 H), 1.08 - 1.15 (m, 9 H), 1.04 (dd, *J*=17.2, 6.8 Hz, 6 H). ^{19}F NMR (376 MHz, *DMSO-d*$_6$) δ -127.96 (s, 1 F). *m/z* (ESI, +ve ion): 573.0 $(M+H)^+$.

[0204] Step 4: *tert*-Butyl (2*R*,5*S*)-4-(1-(3,5-diisopropylpyridin-4-yl)-6-fluoro-7-(2-fluoro-6-hydroxyphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of *tert*-butyl (2*R*,5*S*)-4-(7-chloro-1-(3,5-diisopropylpyridin-4-yl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.13 g, 0.23 mmol), (2-fluoro-6-hydroxyphenyl)boronic acid (0.073 g, 0.47 mmol, Wuxi), potassium acetate (0.115 g, 1.17 mmol), (1,1'-bis(diphenylphosphino)ferrocene)dichloropalladium-dichloromethane (1:1) (0.019 g, 0.023 mmol, Oakwood Products, Inc.) in 1,4-dioxane (2 mL) and water (0.01 mL) was stirred and heated at 90 °C for 14 h. The crude material was purified by silica gel chromatography (eluent: 0-50% EtOAc-EtOH (3:1)/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(1-(3,5-diisopropylpyridin-4-yl)-6-fluoro-7-(2-fluoro-6-hydroxyphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.054 g, 0.083 mmol, 35.5 % yield) as yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 10.17 (br s, 1 H), 8.49 (d, *J*=2.9 Hz, 2 H), 8.27 (d, *J*=9.1 Hz, 1 H), 7.21 - 7.29 (m, 1 H), 6.60 - 6.76 (m, 2 H), 4.72 - 4.91 (m, 1 H), 4.18 - 4.42 (m, 1 H), 4.10 (br d, *J*=13.9 Hz, 1 H), 3.77 - 3.88 (m, 1 H), 3.43 - 3.73 (m, 2 H), 2.42 - 2.49 (m, 2 H), 1.45 (s, 9 H), 0.93 - 1.36 (m, 18 H). ^{19}F NMR (377 MHz, *DMSO-d*$_6$) δ -115.75 (br d, *J*=4.3 Hz, 1 F), -128.53 (br s, 1 F). *m/z* (ESI, +ve ion): 649.1 $(M+H)^+$.

[0205] Step 5: 1-(3,5-Diisopropyl-4-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)pyrido[2,3-*d*]pyrimidin-2-one. A mixture of *tert*-butyl (2*R*,5*S*)-4-(1-(3,5-diisopropylpyridin-4-yl)-6-fluoro-7-(2-fluoro-6-hydroxyphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-

carboxylate (0.052 g, 0.077 mmol) in DCM (2 mL) and TFA (2 mL) was stirred at rt for 10 min. The mixture was concentrated *in vacuo* to afford 1-(3,5-diisopropylpyridin-4-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-6-fluoro-7-(2-fluoro-6-hydroxyphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as orange syrup. The crude product was used in next step without purification. *m/z* (ESI, +ve ion): 549.0 $(M+H)^+$.

[0206] A mixture of 1-(3,5-diisopropylpyridin-4-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-6-fluoro-7-(2-fluoro-6-hydroxyphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (0.2 mL, 1.16 mmol) in DCM (2 mL) was cooled to -20 °C. To the cooled mixture at -20 °C was added acryloyl chloride, 0.2 M solution in DCM (0.39 mL, 0.077 mmol) dropwise and the mixture was stirred at -20 °C for 30 min. The mixture was concentrated *in vacuo* to afford the crude material as a yellow solid. The residue was dissolved in THF (3 mL), treated with 2 N NaOH (0.4 mL, 0.8 mmol), and stirred at rt for 1 h. The reaction mixture was diluted with satd. NH_4Cl (50 mL), extracted with EtOAc (3 × 50 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (eluent 0-25% DCM-MeOH (4:1)/DCM) to provide 1-(3,5-diisopropyl-4-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)pyrido[2,3-*d*]pyrimidin-2-one (0.024 g, 0.040 mmol, 51.7 % yield) as yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 10.19 (br s, 1 H), 8.49 (d, *J*=3.1 Hz, 2 H), 8.30 (d, *J*=9.1 Hz, 1 H), 7.25 (q, *J*=8.1 Hz, 1 H), 6.62 - 6.90 (m, 3 H), 6.18 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.74 - 5.79 (m, 1 H), 4.42 - 4.93 (m, 2 H), 3.79 - 4.23 (m, 4 H), 2.41 - 2.48 (m, 2 H), 1.16 - 1.33 (m, 6 H), 1.11 (dd, *J*=6.7, 4.5 Hz, 6 H), 0.97 (br d, *J*=6.6 Hz, 6 H). ^{19}F NMR (377 MHz, *DMSO-d*$_6$) δ -115.75 (br s, 1 F), -128.46 (br s, 1 F). *m/z* (ESI, +ve ion): 603.0 $(M+H)^+$.

Table 101: Compounds 101-2 to 101-3 were prepared following the procedure described in Method 101, Steps 1-5, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
101-2		1-(4,6-Diisopropylpyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** Intermediate U **Step 4:** 2-fluorophenylboronic acid, Combi-Blocks, San Diego, CA, USA

101-3		1-(4,6-Diisopropylpyrimidin-5-yl)-4-[(2*R*,6*S*)-2,6-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** Intermediate U **Step 3:** *tert*-butyl *cis*-3,5-dimethylpiperazine-1-carboxylate, Combi-Blocks, San Diego, CA, USA) **Step 4:** 2-fluorophenylboronic acid, Combi-Blocks, San Diego, CA, USA

Method 102

Example 102-1: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,6-diisopropylphenyl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0207] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2,6-diisopropylphenyl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a solution of *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (0.050 g, 0.10 mmol, **Intermediate 165**) and DIEA (0.038 mL, 0.22 mmol) in THF (5 mL) was added phosgene solution, 15% in toluene (0.078 mL, 0.11 mmol) at 0 °C. This solution was stirred for 10 min followed by

addition of a solution of 2,6-diisopropylaniline (0.022 g, 0.13 mmol, **Intermediate 193**) in THF (1 mL) at 0 °C. The resulting mixture was stirred at room temp for 15 min at which point LC-MS analysis indicated consumption of starting material and the presence of *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(((2,6-diisopropylphenyl)carbamoyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate. The reaction was diluted with EtOAc (20 mL) and quenched with saturated aqueous $NaHCO_3$. The layers were partitioned and the organic phase was dried over anhydrous $MgSO_4$, filtered, and concentrated under reduced pressure. *m/z* (ESI, +ve ion): 683.6 $(M+H)^+$.

[0208] Step 2: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2,6-diisopropylphenyl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. *tert*-Butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(((2,6-diisopropylphenyl)carbamoyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate was re-dissolved in THF (3 mL) and toluene (3 mL). The resulting solution was cooled to 0 °C followed by addition of sodium *tert*-butoxide (0.015 mg, 0.156 μmol). The reaction was stirred at this temperature for 40 min , diluted with EtOAc, and washed with sat. aqueous $NaHCO_3$. The combined organics were purified by chromatography on silica gel using 0-40% EtOAc in heptane to afford *tert*-butyl (2*R*,5S)-4-(6-chloro-1-(2,6-diisopropylphenyl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.029 g, 0.045 mmol, 43.3 % yield) as a yellow solid. *m/z* (ESI, +ve ion): 648.2 $(M+H)^+$.

[0209] Step 3: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,6-diisopropylphenyl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2,6-diisopropylphenyl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.029 g, 0.045 mmol) in DCM (3 mL) was added tfa (2.0 mL, 26.0 mmol) and the resulting mixture was stirred at rt for 30 min. The reaction went to completion and concentrated to afford

[0210] 6-chloro-1-(2,6-diisopropylphenyl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one which was dissolved in DCM (3 mL) then acryloyl chloride (3.27 μl, 0.040 mmol) was added at rt. The reaction was stirred at rt for 15 min, washed with sat. $NaHCO_3$ and extracted with DCM. The combined organics were purified by HPLC to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,6-diisopropylphenyl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.010 g, 8.30 μmol, 18.56 % yield) as a light yellow solid. ^1H NMR (400 MHz, *DMSO-d*$_6$) δ 8.46 (s, 1 H), 7.43 -

7.57 (m, 1 H), 7.12 - 7.36 (m, 6 H), 6.75 - 6.94 (m, 1 H), 6.19 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.67 - 5.85 (m, 1 H), 4.69 - 4.96 (m, 2 H), 4.39 - 4.52 (m, 1 H), 4.14 (br t, *J*=11.9 Hz, 1 H), 3.78 - 3.99 (m, 2 H), 3.44 - 3.62 (m, 1 H), 1.30 - 1.37 (m, 3 H), 1.25 (br d, *J*=6.4 Hz, 2 H), 1.17 (d, *J*=6.8 Hz, 2 H), 1.03 - 1.10 (m, 6 H), 0.92 (dd, *J*=10.0, 6.8 Hz, 6 H). 601.6 $(M+H)^+$.

Table 102: Compounds 102-2 to 102-7 were prepared following the procedure described in Method 102, Steps 1-3, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
102-2		6-Chloro-1-(4,6-diisopropylpyrimidin-5-yl)-4-[(2*S*)-4-[(*E*)-4-(dimethylamino)but-2-enoyl]-2-methyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	Step 1: 4,6-Diisopropylpyrimidin-5-amine and *tert*-butyl (*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-3-methylpiperazine-1-carboxylate	Step 3: ([(1-Cyano-2-ethoxy-2-oxoethylidene)amino]oxy)dimethylamino(morpholin-4-yl)carbenium hexafluorophosphate and *trans*-4-dimethylaminocrotonic acid hydrochloride
102-3		6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Intermediate 194** in **Step 1**	No change
102-4		1-(6-Bromo-2,4-diisopropyl-3-pyridyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)py	**Intermediate 196** used in **Step 1**	No change

		rido[2,3-*d*]pyrimidin-2-one		
102-5		6-Chloro-1-(4-chloro-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 2:** Performed in Toluene	**Step 1:** **Intermediate 177**
102-6		6-Chloro-1-(4-chloro-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 2:** Performed in Toluene	**Step 1:** **Intermediate 168** and **Intermediate 177**
102-7		6-Chloro-1-(4-dimethylphosphoryl-2-isopropyl-6-methyl-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 1 (1):** 0 °C; **Step 2:** 1:1 THF–toluene, 0 °C	**Step 1 (2):** Intermediate 221

102-8		3-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-1-yl]-4-isopropyl-*N*,*N*-dimethyl-pyridine-2-carboxamide	None	Step 1: Intermediate 210

Method 103

Example 103-1:



[0211] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4-(dimethylamino)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of 3-isopropyl-N1,N1-dimethylbenzene-1,2-diamine (282 mg, 1.583 mmol, **Intermediate 163**) and DIEA (580 µl, 3.32 mmol) in THF (6 mL; 20:1 mL) at 20 °C was added a phosgene solution, 15% in toluene (1248 µl, 1.741 mmol) dropwise. A precipitate formed and the resulting mixture was stirred at 20 °C for 15 min then chilled to 0 °C. Once chilled, solid *tert*-butyl (2*R*,5*S*)-4-((2,5-

dichloro-6-(2-isopropylphenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (800 mg, 1.583 mmol, **Intermediate 170**) was added in one portion. The ice bath was removed and stirred for a total of 45 min at 20 °C. The reaction was returned to 0 °C, a sodium *tert*-butoxide solution, 2.0 M in THF (2374 µl, 4.75 mmol) was added, and then the ice bath removed. After 30 min, the reaction was partitioned between EtOAc (60 mL) and sat. NaHCO3 (20 mL). The organic was further washed with sat. NaCl (10 mL), dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (40 g) eluting products with a gradient of 10>35% EtOAc/EtOH (3:1 blend)/heptane to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4-(dimethylamino)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as white foam. *m/z* (ESI, +ve ion): 674.5 $(M+H)^+$.

124611-28

[0212] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-(dimethylamino)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4-(dimethylamino)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (760 mg, 1.127 mmol) was stirred in TFA (3 mL) at 20 °C for 30 min. The solvent was then removed under reduced pressure followed by an azeotrope with heptane (30 mL). The residue was then partitioned between EtOAc (30 mL) and 5% Na_2CO_3 (20 mL). The separated organic was washed with sat. NaCl (10 mL), dried over $MgSO_4$, then concentrated under reduced pressure to afford the intermediate. To the material was added DCM (10 mL) and DIEA (394 µl, 2.254 mmol), and the solution was chilled to 0 °C. To this solution was added a second solution of acryloyl chloride (92 µl, 1.127 mmol) in DCM (3 mL) was dropwise over a period of 1 min. The reaction was then directly purified by silica gel chromatography (40 g) eluting products with a gradient of 0>70% EtOAc/EtOH (3:1 blend)/heptane to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-(dimethylamino)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as white foam. *m/z* (ESI, +ve ion): 628.3 $(M+H)^+$.

Table 103: Compounds 103-2 to 103-7 were prepared following the procedure described in Method 103, Steps x-x, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent

103-2		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(4-isopropyl-1-methyl-6-oxo-pyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 1-1** & **1-2**: 0°C, phosgene added to **Intermediate 165**; **Step 1-4**: **Intermediate 234** added as a THF solution; **Step 1-4**: aqueous workup performed; **Step 1-5**: THF/toluene (1:1); **Step 2-1**: omit DCM; **Step 2-2**: RT.	**Intermediate 234** & **Intermediate 165**
103-3		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-[2-isopropyl-4-[(*S*)-methylsulfinyl]phenyl]pyrido[2,3-*d*]pyrimidin-2-one	None	**Intermediate 173** and **Intermediate 165**
103-4		6-Chloro-1-(3,5-diisopropyl-4-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Intermediate 162** and **Intermediate 165**
103-5		6-Chloro-1-[4-(diethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-isopropylphenyl	None	**Intermediate 170** and **Intermediate 164**

		)pyrido[2,3-*d*]pyrimidin-2-one		
103-6		6-chloro-1-[4-(diethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Intermediate 168** and **Intermediate 164**
103-7		6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Intermediate 168** and **Intermediate 163**

Method 106

Example 106-1: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0213] **Step 1: *N*-((2-Bromo-4,6-diisopropylpyrimidin-5-yl)carbamoyl)-2,5-dichloro-6-(2-fluorophenyl)nicotinamide**. To a 250-mL round-bottomed flask was added 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (2.95 g, 9.73 mmol, **Intermediate 99B**) and oxalyl chloride (7.29 mL, 14.59 mmol) in tetrahydrofuran (32.4 mL). The flask was fitted with a reflux condensor, and the mixture was stirred and heated at 80 °C for 45 min. The reaction mixture was concentrated in vacuo. The crude mixture was carried on without further purification.

[0214] To a 150-mL round-bottomed flask was added the crude (2,5-dichloro-6-(2-fluorophenyl)nicotinoyl)carbamoyl isocyanate from above in tetrahydrofuran (32.4 mL). The mixture was cooled to -10 °C before a solution of 2-bromo-4,6-diisopropylpyrimidin-5-amine (2.89 g, 11.19 mmol, **Intermediate I-38**) in THF (3 mL) was added. The reaction mixture was allowed to stir under an inert (N2) atmosphere for 1 h. The reaction mixture was concentrated *in vacuo*. The crude material was triturated with EtOAc and heptane, and the solids were collected by filtration. The solids were washed with heptane to afford *N*-((2-bromo-4,6-diisopropylpyrimidin-5-yl)carbamoyl)-2,5-dichloro-6-(2-fluorophenyl)nicotinamide as a tan solid (4.500 g, 7.91 mmol, 81 % yield). ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.48 (s, 1 H) 9.74

(br s, 1 H) 8.62 (s, 1 H) 7.55 - 7.64 (m, 2 H) 7.37 - 7.42 (m, 2 H) 3.21 - 3.30 (m, 2 H) 1.16 - 1.19 (m, 12 H). *m/z* (ESI, +ve ion): 568.0 (M+H)+.

[0215] Step 2: 1-(2-Bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (Intermediate 231). To a 250-mL round-bottomed flask was added *N*-((2-bromo-4,6-diisopropylpyrimidin-5-yl)carbamoyl)-2,5-dichloro-6-(2-fluorophenyl)nicotinamide (4.790 g, 8.41 mmol) in tetrahydrofuran (42.1 mL). Potassium bis(trimethylsilyl)amide (1.0 M solution in tetrahydrofuran, 10.52 mL, 10.52 mmol) was added dropwise via addition funnel into the reaction mixture over 5 min. The reaction mixture was allowed to stir under an inert (N2) atmosphere for 10 min. Additional potassium bis(trimethylsilyl)amide (0.5 eq; 5.5 mL) was added dropwise into the reaction mixture. The reaction mixture was quenched with sat. aq. NH_4Cl (50 mL), then the mixture was diluted with 3:1 EtOAc/MeOH and brine solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over MgSO4, filtered and concentrated in vacuo. The crude was triturated with EtOAc and heptane. The solids were washed with heptane and allowed to dry in a reduced-pressure oven (40°C) for 1 h to afford 1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione as a tan solid (4.26 g, 7.99 mmol, 95 % yield, **Intermediate 231**). ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.83 - 12.28 (m, 1 H) 8.57 (s, 1 H) 7.50 - 7.56 (m, 1 H) 7.31 (d, J=7.88 Hz, 2 H) 7.18 - 7.23 (m, 1 H) 2.94 - 3.02 (m, 2 H) 1.08 (d, J=6.63 Hz, 6 H) 0.91 (d, J=6.63 Hz, 6 H). m/z (ESI, +ve ion): 532.0 (M+H)+.

[0216] Step 3: *tert*-Butyl (2*R*,5*S*)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (Intermediate 232). Phosphorus(V) oxychloride (0.237 mL, 2.54 mmol) was added to a stirred mixture of 1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (1.04 g, 1.952 mmol, **Intermediate 231**) and *N*,*N*-diisopropylethylamine (0.477 mL, 2.73 mmol) in acetonitrile (8 mL). The reaction mixture was stirred at 80 °C for 40 min before being cooled to room temperature. Additional *N*,*N*-diisopropylethylamine (0.477 mL, 2.73 mmol) was added followed by (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (0.418 mL, 1.952 mmol). The reaction mixture was stirred at room temperature for 25 min. The reaction was quenched with water (150 mL) and extracted with EtOAc (200 mL). The organic layer was separated, washed with saturated sodium chloride (150 mL), dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-30% [(3:1)

EtOAc/EtOH]/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as a white solid (1.10 g, 1.509 mmol, 77 % yield, **Intermediate 232**). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.17 (1 H, s) 7.48 - 7.54 (1 H, m) 7.15 - 7.28 (3 H, m) 3.49 - 5.08 (6 H, m) 2.62 - 2.72 (2 H, m) 1.56 (9 H, s) 1.50 - 1.55 (3 H, m) 1.30 - 1.36 (3 H, m) 1.26 (6 H, dd, *J*=6.63, 3.32 Hz) 1.06 (6 H, t, *J*=6.84 Hz). m/z (ESI, +ve ion): 728.0 $(M+H)^+$.

[0217] Step 4: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. Trifluoroacetic acid (1.00 mL, 13.42 mmol) was added to a stirred solution of *tert*-butyl (2*R*,5*S*)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (300 mg, 0.411 mmol) in dichloromethane (2 mL). The reaction mixture was stirred at room temperature for 20 min. The reaction mixture was concentrated *in vacuo* to provide crude 1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one that was used without purification assuming 100% yield.

[0218] Acryloyl chloride (1.1 M in DCM, 0.487 mL, 0.535 mmol) was added to a stirred mixture of the crude 1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (259 mg, 0.412 mmol) and *N*,*N*-diisopropylethylamine (0.360 mL, 2.059 mmol) in dichloromethane (2 mL). The reaction mixture was stirred at room temperature for 30 min. The reaction mixture was diluted with DCM (60 mL) and washed with saturated aqueous NH_4Cl (50 mL). The organic layer was separated, dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-50% [(3:1) EtOAc/EtOH]/heptane) to provide 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a white solid (241 mg, 0.353 mmol, 86 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.17 (1 H, s) 7.48 - 7.54 (1 H, m) 7.16 - 7.28 (3 H, m) 6.56 - 6.74 (1 H, m) 6.40 - 6.49 (1 H, m) 5.82 - 5.88 (1 H, m) 3.49 - 5.23 (6 H, m) 2.60 - 2.72 (2 H, m) 1.44 - 1.56 (3 H, m) 1.29 - 1.40 (3 H, m) 1.23 - 1.29 (6 H, m) 1.03 - 1.09 (6 H, m). m/z (ESI, +ve ion): 682.0 $(M+H)^+$.

Table 106: Compounds 106-2 to 106-10 were prepared following the procedure described in Method 106, Steps 1-4, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
106-2		6-Chloro-1-(2,2-dimethylpropyl)-7-(2-fluorophenyl)-4-[(2*S*)-2-methyl-4-prop-2-enoyl-piperazin-1-yl]pyrido[2,3-*d*]pyrimidin-2-one	**Step 1:** RT	**Step 1:** neopentylamine (TCI America) **Step 3:** (3*S*)-1-(*tert*-butoxycarbonyl)-3-methylpiperazine (Combi-Blocks Inc.)
106-3		6-Chloro-1-(2,2-dimethylpropyl)-7-(2-fluorophenyl)-4-[(3*R*)-3-(hydroxymethyl)-4-prop-2-enoyl-piperazin-1-yl]pyrido[2,3-*d*]pyrimidin-2-one	**Step 1:** RT	**Step 1:** neopentylamine (TCI America) **Step 3:** (*R*)-1-*N*-Boc-2-hydroxymethyl-piperazine, (Sigma-Aldrich Corporation)
106-4		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(2-methoxy-2-methyl-propyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** 2-methoxy-2-methylpropan-1-amine, (Aurum Pharmatech)
106-5		6-Chloro-7-(2-fluorophenyl)-1-(2-methoxy-2-methyl-propyl)-4-(4-prop-2-enoylpiperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** 2-methoxy-2-methylpropan-1-amine, (Aurum Pharmatech) **Step 3:** 1-Boc-piperazine, (Sigma-Aldrich Corporation)
106-6		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-	**Step 1 (1):** 80 °C; **Step 1 (2):** 0 °C to rt; **Step 2:** 0 °C to rt; **Step 3:** 0 °C; **Step 4:** 0 °C	**Step 1 (1): Example 50, step 3**; **Step 1 (2):** Intermediate R; **Step 3 (2):** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate

		pyridyl)pteridin-2-one		(eNovation Chemicals LLC)
106-7		6-Chloro-4-[(2*R*,6*S*)-2,6-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pteridin-2-one	**Step 1 (1):** 80 °C; **Step 1 (2):** 0 °C; **Step 2:** 0 °C to rt; **Step 3:** 0 °C; **Step 4:** 0 °C	**Step 1 (1): Example 50, step 3**; **Step 1 (2):** Intermediate R; **Step 3 (2):** *cis*-3,5-dimethyl-piperazine-1-carboxylic acid *tert*-butyl ester (Combi-Blocks Inc, San Diego, CA, USA)
106-8		3-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-1-yl]-2,2-dimethyl-propanenitrile	None	**Step 1:** 3-Amino-2,2-dimethylpropanenitrile hydrochloride salt and DIPEA
106-9		3-[6-Chloro-7-(2-fluorophenyl)-4-[(2*S*)-2-methyl-4-prop-2-enoyl-piperazin-1-yl]-2-oxo-pyrido[2,3-*d*]pyrimidin-1-yl]-2,2-dimethyl-propanenitrile	None	**Step 1:** 3-Amino-2,2-dimethylpropanenitrile hydrochloride salt and DIPEA; **Step 3:** (3*S*)-1-(*tert*-Butoxycarbonyl)-3-methylpiperazine
106-10		3-[6-Chloro-7-(2-fluorophenyl)-2-oxo-4-(4-prop-2-enoylpiperazin-1-yl)pyrido[2,3-*d*]pyrimidin-1-yl]-2,2-dimethyl-propanenitrile	None	**Step 1:** 3-Amino-2,2-dimethylpropanenitrile hydrochloride salt and DIPEA; **Step 3:** *tert*-butyl piperazine-1-carboxylate

Method 107

Example 107-1: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Intermediate 232

[0219] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. Methylamine (2.0 M in tetrahydrofuran, 0.312 mL, 0.624 mmol) was added to a stirred solution of *tert*-butyl (2*R*,5*S*)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (91 mg, 0.125 mmol, Intermediate 232) in acetonitrile (0.5 mL). The reaction mixture was stirred at room temperature for 9 h. The reaction mixture was concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-50% [(3:1) EtOAc/EtOH]/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as a yellow solid (45 mg, 0.066 mmol, 53.1 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.13 (1 H, s) 7.45 - 7.52 (1 H, m) 7.21 - 7.29 (2 H, m) 7.14 - 7.20 (1 H, m) 3.53 - 5.12 (7 H, m) 3.05 (3 H, d, *J*=4.98 Hz) 2.49 - 2.61 (2 H, m) 1.57 (9 H, s) 1.44 - 1.54 (3 H, m) 1.30 - 1.36 (3 H, m) 1.21 (3 H, dd, *J*=6.34, 4.25 Hz) 1.01 (6 H, br t, *J*=7.26 Hz). m/z (ESI, +ve ion): 679.1 $(M+H)^+$.

[0220] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. Trifluoroacetic acid (0.5 mL, 6.71 mmol) was added to a stirred solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (45 mg, 0.066 mmol) in dichloromethane (1 mL). The reaction mixture was stirred at room temperature for 25 min. The reaction mixture was concentrated *in vacuo* to give crude 6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one that was used without purification assuming 100% yield.

[0221] Acryloyl chloride (1.1 M in DCM, 0.072 mL, 0.079 mmol) was added to a stirred mixture of the crude 6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (38 mg, 0.066 mmol) and *N*,*N*-diisopropylethylamine (0.046 mL, 0.262 mmol) in dichloromethane (0.5 mL). The reaction mixture was stirred at room temperature for 1 h. The reaction mixture was diluted with DCM (40 mL) and washed with saturated aqueous NH_4Cl (30 mL). The organic layer was separated, dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-50% [(3:1) EtOAc/EtOH]/heptane) to provide 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a light yellow solid (33 mg, 0.052 mmol, 79 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.15 (1 H, br s) 7.47 - 7.54 (1 H, m) 7.16 - 7.29 (3 H, m) 6.59 - 6.76 (1 H, m) 6.42 - 6.51 (1 H, m) 5.84 - 5.90 (1 H, m) 3.53 - 5.24 (6 H, m) 3.07 (3 H, d, *J*=4.80 Hz) 2.50 - 2.63 (2 H, m) 1.46 - 1.54 (3 H, m) 1.32 - 1.42 (3 H, m) 1.19 - 1.28 (6 H, m) 0.98 - 1.09 (6 H, m). m/z (ESI, +ve ion): 633.2 $(M+H)^+$.

Table 107: Compounds 107-2 to 107-5 were prepared following the procedure described in Method 107, Steps 1-2, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
107-2		6-Chloro-1-[2-(dimethylamino)-4,6-diisopropyl-pyrimidin-5-yl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 1:** Performed at 40 °C	**Step 1:** dimethylamine, 2.0 M in THF (Sigma-Aldrich Corporation)

107-3		6-Chloro-1-(4,6-diisopropyl-2-morpholino-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** morpholine (Sigma-Aldrich Corporation)
107-4		6-Chloro-1-[2-[2-(dimethylamino)ethylamino]-4,6-diisopropyl-pyrimidin-5-yl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** *N*,*N*-dimethylethylenediamine (Sigma-Aldrich Corporation)
107-5		1-[2-(Azetidin-1-yl)-4,6-diisopropyl-pyrimidin-5-yl]-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** azetidine (Combi-Blocks Inc.)

Method 108

Example 108-1: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-cyclopropyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Intermediate 175

[0222] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4-cyclopropyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. Cyclopropylzinc bromide (0.5 M in THF, 3.819 mL, 1.910 mmol) was added to a flask charged with *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4-chloro-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (409 mg, 0.637 mmol, **Intermediate 175**) and Xantphos Pd G3 (60.4 mg, 0.064 mmol) under an argon atmosphere. The reaction mixture was stirred at room temperature for 2 h. The reaction mixture was quenched with saturated aqueous NH_4Cl (75 mL) and extracted with EtOAc (125 ml). The organic layer was separated, washed with saturated aqueous sodium chloride (75 mL), dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-50% [(3:1) EtOAc/EtOH]/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4-cyclopropyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as an off-white solid (297 mg, 0.458 mmol, 72.0 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.96 (1 H, s) 8.10 - 8.14 (1 H, m) 7.40 - 7.47 (1 H, m) 7.09 - 7.25 (3 H, m) 3.45 - 5.09 (6 H, m) 2.71 - 2.81 (1 H, m) 1.42 - 1.60 (13 H, m) 1.22 - 1.32 (6 H, m) 1.02 - 1.14 (4 H, m) 0.91 - 1.00 (1 H, m) 0.78 - 0.86 (1 H, m). m/z (ESI, +ve ion): 648.0 $(M+H)^+$.

[0223] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-cyclopropyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. Trifluoroacetic acid (1.00 mL, 13.42 mmol) was added to a stirred solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4-cyclopropyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (297 mg, 0.458 mmol) in dichloromethane (1 mL). The reaction mixture was stirred at room temperature for 40 min. The reaction mixture was concentrated *in vacuo* to give crude 6-chloro-1-(4-cyclopropyl-6-isopropylpyrimidin-5-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-

fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one that was used without purification assuming 100% yield.

[0224] Acryloyl chloride (1.1 M in DCM, 0.500 mL, 0.550 mmol) was added to a stirred mixture of the crude 6-chloro-1-(4-cyclopropyl-6-isopropylpyrimidin-5-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (251 mg, 0.458 mmol) and *N*,*N*-diisopropylethylamine (0.400 mL, 2.290 mmol) in dichloromethane (2 mL). The reaction mixture was stirred at room temperature for 30 min. The reaction mixture was diluted with DCM (100 mL) and washed with saturated aqueous NaHCO3 (75 mL). The organic layer was separated, dried over MgSO4, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-50% [(3:1) EtOAc/EtOH]/heptane) to provide 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-cyclopropyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as an off-white solid (220 mg, 0.183 mmol, 80 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 9.01 - 9.03 (1 H, m) 8.16 - 8.19 (1 H, m) 7.46 - 7.52 (1 H, m) 7.14 - 7.30 (3 H, m) 6.58 - 6.74 (1 H, m) 6.46 (1 H, br t, J=15.03 Hz) 5.83 - 5.89 (1 H, m) 3.49 - 5.27 (6 H, m) 2.75 - 2.86 (1 H, m) 1.44 - 1.63 (5 H, m) 1.27 - 1.41 (6 H, m) 1.07 - 1.20 (4 H, m) 0.96 - 1.06 (1 H, m) 0.84 - 0.95 (1 H, m). m/z (ESI, +ve ion): 602.0 $(M+H)^+$.

Table 108: Compound 108-2 was prepared following the procedure described in Method 108, Steps 1-2, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
108-2		6-Chloro-1-(4-cyclopropyl-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1: Intermediate 176**

Method 109

Example 109-1: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-ethyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Intermediate 175

[0225] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4-ethyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. Triethylaluminum (1.0 M in heptane, 0.934 mL, 0.934 mmol) was added to a stirred mixture of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4-chloro-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (400 mg, 0.623 mmol, **Intermediate 175**) and tetrakis(triphenylphosphine) palladium (71.9 mg, 0.062 mmol) in tetrahydrofuran (3 mL) under an argon atmosphere. The reaction mixture was stirred at 70 °C for 22 h. The reaction mixture was quenched with saturated aqueous $NaHCO_3$ (50 mL) and extracted with EtOAc (75 mL). The organic layer was separated, washed with saturated aqueous sodium chloride (50 mL), dried over $MgSO_4$, filtered, and concentrated *in vacuo* to provide crude *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4-ethyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as a brown solid. m/z (ESI, +ve ion): 636.1 $(M+H)^+$. The crude product contained a large amount of the des-Boc product and was used without purification assuming 100% yield.

[0226] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-ethyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. Trifluoroacetic acid (2 mL, 26.8 mmol) was added to a stirred solution of the crude *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4-ethyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (396 mg, 0.622 mmol) in dichloromethane (2 mL). The reaction mixture was stirred at room temperature for 25 min. The reaction mixture was concentrated *in vacuo* to provide crude 6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-1-(4-ethyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (124607-19-1) that was used without purification assuming 100% yield.

[0227] Acryloyl chloride (1.1 M in DCM, 0.680 mL, 0.748 mmol) was added to a stirred mixture of the crude 6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-1-(4-ethyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (334 mg, 0.623 mmol) and *N*,*N*-diisopropylethylamine (0.544 mL, 3.12 mmol) in dichloromethane (2 mL). The reaction mixture was stirred at room temperature for 30 min. The reaction mixture was diluted with DCM (75 mL) and washed with saturated aqueous NH_4Cl (50 mL). The organic layer was separated, dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-70% [(3:1) EtOAc/EtOH]/heptane) to provide 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-ethyl-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a light yellow solid (231 mg, 0.196 mmol, 62.8 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.11 (1 H, s) 8.10 - 8.14 (1 H, m) 7.40 - 7.47 (1 H, m) 7.08 - 7.20 (3 H, m) 6.51 - 6.70 (1 H, m) 6.41 (1 H, br t, *J*=14.93 Hz) 5.77 - 5.85 (1 H, m) 3.44 - 5.21 (6 H, m) 2.65 - 2.77 (1 H, m) 2.48 - 2.59 (1 H, m) 2.36 - 2.49 (1 H, m) 1.40 - 1.52 (4 H, m) 1.30 - 1.36 (2 H, m) 1.14 - 1.28 (6 H, m) 1.02 - 1.09 (3 H, m). m/z (ESI, +ve ion): 590.2 $(M+H)^+$.

Table 109: Compound 109-2 was prepared following the procedure described in Method 109, Steps 1-2, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
109-2		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(4-ethyl-6-isopropyl-pyrimidin-5-yl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1: Intermediate 176**

Method 110

Example 110-1: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-neopentylpyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0228] Step 1: *tert*-Butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate: A mixture of *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 165**, 1.30 g, 2.70 mmol) and neopentylamine (1.27 mL, 10.80 mmol) was heated at 110 °C for 3 days. The reaction mixture was purified on silica gel chromatography using 0-5% MeOH in DCM to afford *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (1.10 g, 2.07 mmol, 77.0 % yield). *m/z* (ESI, +ve ion): 532.3 $(M+H)^+$.

[0229] Step 2: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-neopentyl-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate: A solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (0.065 g, 0.122 mmol) and 1,1'-carbonyldiimidazole (0.040 g, 0.244 mmol) in THF (1.0 mL) was stirred at 100 °C for 2 hours. The reaction was concentrated and residue purified on silica gel chromatography using 0-60% EtOAc/EtOH (3:1) in heptane to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-neopentyl-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.064 g, 0.115 mmol, 94 % yield). *m/z* (ESI, +ve ion): 558.4 $(M+H)^+$.

[0230] Step 3: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-neopentylpyrido[2,3-*d*]pyrimidin-2(1*H*)-one: TFA (0.160 mL, 2.150 mmol) was added to a solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-

neopentyl-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.060 g, 0.108 mmol) in DCM (4 mL). The resulting mixture was stirred at r.t. for 1 h and then concentrated under reduced pressure. The residue was suspended in DCM (4 mL), cooled to 0 °C, and treated with DIPEA (0.056 mL, 0.323 mmol) followed by acryloyl chloride (0.013 mL, 0.161 mmol). The reaction was warmed to r.t. and stirred for 10 min. The mixture was quenched with sat'd aqueous $NaHCO_3$ and extracted with DCM (2X). The combined organic layers were concentrated and the residue purified on silica gel using 0-50% EtOAc/EtOH (3:1) in heptane to provide 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-neopentylpyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.048 g, 0.094 mmol, 87 % yield). ^{1}H NMR (400 MHz, DMSO-d6) δ 8.33 (s, 1H), 7.53-7.65 (m, 2H), 7.37-7.46 (m, 2H), 6.76-6.87 (m, 1H), 6.09-6.25 (m, 1H), 5.76 (s, 1H), 4.37-4.88 (m, 2H), 4.19-4.30 (m, 1H), 4.03-4.18 (m, 2H), 3.72-3.93 (m, 2H), 3.45 (br d, J=12.02 Hz, 1H), 1.23-1.31 (m, 3H), 1.08-1.21 (m, 3H), 0.88 (s, 9H). ^{19}F NMR (376 MHz, DMSO-d6) δ -113.60 (s, 1F). *m/z* (ESI, +ve ion): 512.3 $(M+H)^+$.

Table 110: Compound 110-2 was prepared following the procedure described in Method 110, Steps 1-3, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
110-2		6-Chloro-1-cyclohexyl-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	1-amino cyclohexane

Method 119

Example 119-1: 4-((2*S*,5*R*)-4-(But-2-ynoyl)-2,5-dimethylpiperazin-1-yl)-6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Intermediate 185

[0231] Steps 1&2. 4-((2*S*,5*R*)-4-(But-2-ynoyl)-2,5-dimethylpiperazin-1-yl)-6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. 4-((2*S*,5*R*)-4-(But-2-ynoyl)-2,5-dimethylpiperazin-1-yl)-6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To *tert*-butyl (2*R*,5*S*)-4-(6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 185**, 0.12 g, 0.2 mmol) dissolved in dichloromethane (0.9 mL) was added trifluoroacetic acid (0.3 mL, 4.0 mmol). The reaction mixture was stirred at RT for 30 min. The solvent was removed *in vacuo* and to the residue was added DCM (1 mL), 1,1'-dimethyltriethylamine (0.13 mL, 0.73 mmol), [(dimethylamino)([1,2,3]triazolo[4,5-b]pyridin-3-yloxy))methylidene]dimethylazanium hexafluorophosphate (0.12 mL, 0.30 mmol), and 2-butynoic acid (20 mg, 0.24 mmol). The reaction mixture was stirred at RT for 3 h. The reaction mixture was partitioned between DCM and brine. The aqueous layer was extracted with DCM (2x). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The crude material was purified by chromatography through an ISCO 12 Gold column, eluting with a gradient of 0% to 45% 3:1 EtOAc/EtOH in heptane, to provide 4-((2*S*,5*R*)-4-(but-2-ynoyl)-2,5-dimethylpiperazin-1-yl)-6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (44 mg, 0.077 mmol, 38 % yield) as off-white solid. *m/z* (ESI, +ve ion): 571.3 $(M+H)^+$. ^1H NMR (400 MHz, CHLOROFORM-d) δ 8.57 (d, *J*=4.98 Hz, 1H), 7.84 (dd, *J*=4.15, 8.91 Hz, 1H), 7.40-7.51 (m, 1H), 7.27-7.30 (m, 1H), 7.08-7.23 (m, 3H), 4.79-5.12 (m, 2H), 4.06-4.41 (m, 2H), 3.41-3.92 (m, 2H), 2.76 (dt, *J*=5.08, 6.58 Hz, 1H), 2.08-2.10 (m, 3H), 2.04-2.06 (m, 3H), 1.42 (t, *J*=7.26 Hz, 3H), 1.24-1.3 (m, 6H), 1.11 (dd, *J*=2.07, 6.84 Hz, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.58 (dd, *J*=16.04, 40.31 Hz, 1F), -126.38 (br dd, *J*=26.44, 40.31 Hz, 1F).

Table 119: Compounds 119-2 was prepared following the procedure described in Method 119, Steps 1-2, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
119-2		4-[(2*S*,5*R*)-4-[4-(Dimethylamino)but-2-ynoyl]-2,5-dimethyl-piperazin-1-yl]-6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	No	**Step 2:** Intermediate 186

Method 121

Example 121-1: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(6-((dimethylamino)methyl)-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Intermediate **165**

[0232] Steps 1&2. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(6-((dimethylamino)methyl)-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a solution of *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (Intermediate 165, 460 mg, 0.96 mmol) and DIEA (350 µl, 2.01 mmol) in THF (2 mL) cooled to 0 °C was added phosgene solution, 15% in toluene (716 µl, 1.00 mmol) dropwise . This solution was stirred for 10 min followed by addition of a solution of 6-((dimethylamino)methyl)-2-isopropyl-4-methylpyridin-3-amine (**Intermediate 187**, 238 mg, 1.15 mmol) in THF (2 mL) at 0 °C. The resulting mixture was stirred at rt for 15 min.

[0233] The reaction mixture was partitioned between EtOAc and sat. $NaHCO_3$. The organic was dried over Na_2SO_4, concentrated then re-dissolved in DMF (4 mL) and treated with cesium carbonate (311 mg, 0.956 mmol). The reaction mixture was heated in a 80°C oil bath for 30 min. The reaction mixture was then partitioned between EtOAc and brine, the organic was dried over Na_2SO_4, and concentrated under reduced pressure. The crude material was purified by chromatography through a Biotage pre-packed silica gel column (40S), eluting with a gradient of 0% to 100% 3:1 EtOAc/EtOH, then with 20% MeOH in DCM to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(6-((dimethylamino)methyl)-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (62 mg, 0.092 mmol, 9.6 % yield) as off-white solid. *m/z* (ESI, +ve ion): 678.3 $(M+H)^+$.

[0234] Step 3&4. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(6-((dimethylamino)methyl)-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A RBF was loaded with crude *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(6-((dimethylamino)methyl)-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (62 mg, 0.091 mmol) and dichloromethane (0.5 ml). Trifluoroacetic acid (136 µl, 1.83 mmol) was added dropwise. After stirring at rt 30 min, the reaction mixture was concentrated *in vacuo* and the residue was re-dissolved in dichloromethane (0.5 ml). 1,1'-Dimethyltriethylamine (80 µl, 0.46 mmol) was added followed by dropwise addition of acryloyl chloride (100 µl, 0.11 mmol) at 0°C. The reaction was stirred at 0°C for 30 min and the reaction mixture was purified using a RediSep Gold (12 g) column and an eluent gradient of 0-20% 5% 2N NH3 in MeOH/MeOH in DCM to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(6-((dimethylamino)methyl)-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (9.5 mg, 16%) as white solid. *m/z* (ESI, +ve ion): 632.4 $(M+H)^+$. ^1H NMR (400 MHz, CHLOROFORM-d) δ 8.15 (s, 1H), 7.79-8.04 (m, 1H), 7.40-7.49 (m, 1H), 7.08-7.22 (m, 3H), 6.32-6.72 (m, 2H), 5.72-5.94 (m, 1H), 4.96-5.19 (m, 1H), 4.56-4.76 (m, 1H), 3.58-4.49 (m, 4H), 3.09 (dq, *J*=4.35, 7.39 Hz, 1H), 2.76-3.01 (m, 6H), 2.16 (br s, 2H), 1.53-1.55 (m, 3H), 1.44-1.48 (m, 3H), 1.33 (br d, *J*=6.84 Hz, 6H), 1.14-1.21 (m, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.77 (br dd, *J*=7.37, 11.70 Hz, 1F).

Table 121: Compounds 121-2 to 121-4 were prepared following the procedure described in Method 121, Steps 1-4, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
121-2		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-[2-isopropyl-4-methyl-6-(pyrrolidin-1-ylmethyl)-3-pyridyl]pyrido[2,3-*d*]pyrimidin-2-one	No	**Step 1:** Intermediate 188
121-3	 Single Isomer	6-Chloro-1-[4-[(dimethylamino)methyl]-2-isopropyl-6-methyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 2:** tBuOK in toluene	**Step 1: I-8**
121-4	 Single Isomer	6-Chloro-1-[4-[(dimethylamino)methyl]-2-isopropyl-6-methyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 2:** tBuOK in toluene	**Step 1: I-8**

Method 122

Example 122-1: *N*-(2-(4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-(dimethylamino)-2-isopropylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-7-yl)-3-fluorophenyl)-2,2,2-trifluoroacetamide



Example 55-54

[0235] To a solution of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-amino-6-fluorophenyl)-6-chloro-1-(4-(dimethylamino)-2-isopropylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.050 g, 0.081 mmol, **Example 55-54**) in DCM (3 mL) was added trifluoroacetic anhydride (10.27 µl, 0.073 mmol) and the resulting mixture was stirred at rt for 15 min. The reaction went to completion, concentrated and purified by HPLC to afford *N*-(2-(4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-(dimethylamino)-2-isopropylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-7-yl)-3-fluorophenyl)-2,2,2-trifluoroacetamide (0.0432 g, 0.030 mmol, 37.4 % yield) as a yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.59 (s, 1 H), 8.07 - 8.18 (m, 1 H), 7.59 (br dd, *J*=7.8, 6.5 Hz, 1 H), 7.37 (br d, *J*=8.9 Hz, 1 H), 7.17 - 7.30 (m, 1 H), 6.84 (dt, *J*=16.7, 10.7 Hz, 1 H), 6.57 - 6.72 (m, 1 H), 6.19 (dd, *J*=16.5, 2.2 Hz, 1 H), 5.74 (br d, *J*=2.3 Hz, 1 H), 4.81 - 5.05 (m, 1 H), 4.66 - 4.76 (m, 1 H), 4.45 - 4.57 (m, 1 H), 4.04 - 4.29 (m, 2 H), 3.67 - 3.79 (m, 1 H), 2.72 (s, 7 H), 1.21 - 1.31 (m, 4 H), 1.00 - 1.12 (m, 3 H), 0.92 (br dd, *J*=6.6, 2.5 Hz, 3 H), 0.78 (br d, *J*=6.6 Hz, 3 H). *m/z* (ESI, +ve ion): 714.4 $(M+H)^+$.

Table 122: Compounds 122-2 to 122-3 were prepared following the procedure described in Method 122 above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
122-2		N-[2-[6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]propanamide	propionyl chloride was used in **Step 1**	No change
122-3		N-[2-[6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]cyclopropanecarboxamide	cyclopropane carbonyl chloride was used in **Step 1**	No change

Method 123

4-((3*S*,5*R*)-4-Acryloyl-3,5-dimethylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluoro-6-hydroxyphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one 2,2,2-trifluoroacetate



[0236] Step 1: *N*-((6-(bis(4-Methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)carbamoyl)-2,6-dichloro-5-fluoronicotinamide. To a stirred solution of 2,6-dichloro-5-fluoronicotinamide (0.34 g, 1.62 mmol, **Intermediate P**) in tetrahydrofuran (5.4 ml) was added oxalyl chloride (2M in DCM) (0.97 ml, 1.94 mmol) dropwise. The reaction mixture was heated to 70 °C for 1 h, cooled to rt, concentrated *in vacuo*, then redissolved in 5 mL acetonitrile. To this was added a stirred solution of 6-isopropyl-N^2,N^2-bis(4-methoxybenzyl)-4-methylpyridine-2,5-diamine (0.66 g, 1.62 mmol, **Intermediate I-33**) in 3 mL acetonitrile at 0 °C for 20 min at rt. The mixture was partitioned between EtOAc and 4:1 brine:water, backextracted with EtOAc (1x), dried over MgSO4, filtered, and concentrated to provide *N*-

((6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)carbamoyl)-2,6-dichloro-5-fluoronicotinamide as a tan solid which was used without purification. *m/z* (ESI, +ve ion): 640.1 $(M+H)^+$.

[0237] Step 2: 1-(6-(bis(4-Methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-7-chloro-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a stirred solution of *N*-((6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)carbamoyl)-2,6-dichloro-5-fluoronicotinamide (1.04 g, 1.62 mmol) in THF at 0 °C was added potassium bis(trimethylsilyl)amide (1M in THF) (4.06 ml, 4.06 mmol). The solution was warmed slowly to rt and stirred for 20 min, then the reaction was quenched with ammonium chloride (aq), and partitioned between EtOAc and brine. The aqueous was backextracted with EtOAc (2x) and the combined organics were dried over MgSO4, filtered, and concentrated. The crude was purified using silica gel chromatography (eluent: 0-40% 3:1 [EtOAc/EtOH] in heptane to provide 1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-7-chloro-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.61 g, 1.01 mmol, 62 % yield) as an off white solid. *m/z* (ESI, +ve ion): 604.6 $(M+H)^+$.

[0238] Step 3: *tert*-Butyl (2*S*,6*R*)-4-(1-(6-(bis(4-Methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-7-chloro-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,6-dimethylpiperazine-1-carboxylate.

Phosphorous oxychloride (0.11 ml, 1.22 mmol) was added dropwise to a solution of 1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-7-chloro-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.61 g, 1.01 mmol) and 1,1'-dimethyltriethylamine (0.23 ml, 1.32 mmol) in acetonitrile (1 ml). The mixture was heated to 80 °C for 1 h. The reaction mixture was cooled to 0°C and *N*-ethyl-*N*-isopropylpropan-2-amine (0.53 ml, 3.0 mmol) was added followed by (2*R*,6*S*) -*tert*-butyl 2,6-dimethylpiperazine-1-carboxylate (0.228 ml, 1.064 mmol). This mixture was warmed to RT and stirred for 1h then was partitioned between EtOAc/NaHCO3 (aq) and the resulting biphasic mixture was separated. The aqueous layer was back-extracted with EtOAc (2x) and the organic extracts were dried over MgSO4, filtered, and concentrated *in vacuo*. The crude material was purified via silica gel chromatography (eluent: 0-60% 3:1 [EtOAc:EtOH] in heptane to provide *tert*-butyl (2*S*,6*R*)-4-(1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-7-chloro-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,6-dimethylpiperazine-1-carboxylate (0.66 g, 0.83 mmol, 82 % yield) as a white solid, containing residual impurities but which will be used as-is without further purification. *m/z* (ESI, +ve ion): 800.4 $(M+H)^+$.

[0239] Step 4: 4-((3*S*,5*R*)-4-Acryloyl-3,5-dimethylpiperazin-1-yl)-1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-7-chloro-6-fluoropyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a vial loaded with *tert*-butyl (2*S*,6*R*)-4-(1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-7-chloro-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,6-dimethylpiperazine-1-carboxylate (0.66 g, 0.83 mmol), was added dichloromethane (4.1 ml), and trifluoroacetic acid (1.2 ml, 16.5 mmol). After stirring at rt 1h the reaction mixture was concentrated *in vacuo*, and the residue was re-dissolved in dichloromethane (4.1 ml) and triethylamine (0.58 ml, 4.1 mmol) followed by dropwise addition of acryloyl chloride (1.12 ml, 1.2 mmol). The reaction was stirred at rt for 30min, was partitioned between EtOAc/NaHCO3, washed with NaHCO3 (2x), brine (1x), dried over MgSO4 then purified using silica gel chromatography (eluent: 0-60% 3:1 [EtOAc:EtOH] in heptane to provide 4-((3*S*,5*R*)-4-acryloyl-3,5-dimethylpiperazin-1-yl)-1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-7-chloro-6-fluoropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.30 g, 0.39 mmol, 47 % yield) as a white solid *m/z* (ESI, +ve ion): 754.0 $(M+H)^+$.

[0240] Step 5: 4-((3*S*,5*R*)-4-Acryloyl-3,5-dimethylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-fluoro-7-(2-fluoro-6-hydroxyphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A vial was charged with 4-((3*S*,5*R*)-4-acryloyl-3,5-dimethylpiperazin-1-yl)-1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-7-chloro-6-fluoropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.1 g, 0.13 mmol), potassium trifluoro(2-fluoro-6-hydroxyphenyl)borate (0.04 g, 0.16 mmol, **Intermediate Q**), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (ii) dichloromethane adduct (9.7 mg, 0.013 mmol), and potassium acetate (0.065 g, 0.66 mmol). The flask was evacuated and backfilled with N2 (3x) followed by addition of 1,4-dioxane (1 ml) and water (0.2 ml). The mixture was stirred at 90 °C for 18 h. The reaction mixture was cooled to rt, then filtered through a short plug of SiO2 (eluent: 3:1 EtOAc:EtOH). The filtrate was concentrated, redissolved in trifluoroacetic acid (290 µl, 4 mmol) and heated to 75 °C for 10 min, the solvent was concentrated *in vacuo*, the remaining residue was taken up with EtOAc, washed with NaHCO3, dried over MgSO4, adsorbed directly onto silica gel and purified via chromatography (eluent: 60% 3:1 [EtOAc/EtOH] in heptane followed by SFC purification to provide 4-((3*S*,5*R*)-4-acryloyl-3,5-dimethylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-fluoro-7-(2-fluoro-6-hydroxyphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-. ^{1}H NMR (CDCl3, 400MHz): δ = 9.79 (br s, 1H), 8.05 (d, J=9.7 Hz, 1H), 7.25-7.32 (m, 1H),

6.57-6.76 (m, 3H), 6.34-6.43 (m, 2H), 5.78 (dd, J=10.5, 1.8 Hz, 1H), 4.68 (br s, 2H), 4.50 (s, 2H), 4.29 (br d, J=13.5 Hz, 2H), 3.60 (br dd, J=13.5, 4.1 Hz, 2H), 2.56 (dt, J=13.3, 6.7 Hz, 1H), 1.90 (s, 3H), 1.46-1.61 (m, 6H), 1.14 (d, J=6.6 Hz, 3H), 0.97 (d, J=6.8 Hz, 3H). ^{19}F NMR (CDCl3, 376MHz): δ = -107.30 (d, *J*=82.4 Hz, 1F), -121.70 (d, *J*=83.2 Hz, 1F). *m/z* (ESI, +ve ion): 590.2 $(M+H)^+$.

Table 123: Compounds 123-2 to 123-6 were prepared following the procedure described in Method 123, Steps 1-5, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
123-2	 Single isomer (P)	1-(6-Amino-2-isopropyl-4-methyl-3-pyridyl)-4-[(3*S*,5*R*)-3,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	Step 1: **I-33**
123-3	 Single isomer (P)	1-(6-Amino-2-isopropyl-4-methyl-3-pyridyl)-4-[(3*S*,5*R*)-3,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1: I-33** **Step 3:** (2*R*,6*S*) -*tert*-butyl 2,6-dimethylpiperazine-1-carboxylate (ArkPharm) **Step 5:** (2-fluorophenyl)boranediol (Combi-Blocks)
123-4	 Single isomer (M)	1-(6-Amino-2-isopropyl-4-methyl-3-pyridyl)-4-[(3*S*,5*R*)-3,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1: I-33** **Step 3:** (2*R*,6*S*)-*tert*-butyl 2,6-dimethylpiperazine-1-carboxylate (ArkPharm) **Step 5:** (2-fluorophenyl)boranediol

				(Combi-Blocks)
123-5	 Single isomer (P)	7-(2-Amino-6-fluoro-phenyl)-1-(6-amino-2-isopropyl-4-methyl-3-pyridyl)-4-[(3*S*,5*R*)-3,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1: I-33** **Step 3:** (2*R*,6*S*)-*tert*-butyl 2,6-dimethylpiperazine-1-carboxylate (ArkPharm) **Step 5:** (2-amino-6-fluorophenyl)boronic acid pinacol ester (CombiPhos)
123-6	 Single isomer (M)	7-(2-Amino-6-fluoro-phenyl)-1-(6-amino-2-isopropyl-4-methyl-3-pyridyl)-4-[(3*S*,5*R*)-3,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1: I-33** **Step 3:** (2*R*,6*S*)-*tert*-butyl 2,6-dimethylpiperazine-1-carboxylate (ArkPharm) **Step 5:** (2-amino-6-fluorophenyl)boronic acid pinacol ester (CombiPhos)

Method 124

Example 124-1: 6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one



[0241] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2,4-diisopropylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. Phosphorous oxychloride (0.016 mL, 0.168 mmol) was added dropwise to a solution of 6,7-dichloro-1-(2,4-diisopropylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.055 g, 0.140 mmol, **Intermediate 195A**) and 1,1'-dimethyltriethylamine (0.032 mL, 0.182 mmol) in CH_3CN (20 mL) under N_2. This mixture was then heated to 80 °C for 30 min. LCMS showed chlorination completed. The reaction mixture was cooled to 0 °C and 3 equiv DIPEA were added followed by (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (0.033 mL, 0.154 mmol, AstaTech). This mixture was stirred with warming to rt over 30 min at which time LCMS showed conversion to desired product. The mixture was poured into cold satd. NaHCO3 solution and stirred vigorously for 10 min. The mixture was extracted with EtOAc, the combined organics were dried over Na_2SO_4, filtered, concentrated and purified by chromatography on silica gel using 0-70% EtOAc in heptane to afford a light yellow solid as *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2,4-diisopropylpyridin-3-yl)-2-oxo-1,2-

dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.051 g, 0.087 mmol, 61.9 % yield). *m/z* (ESI, +ve ion): 588.6 $(M+H)^+$.

[0242] Step 2. *tert*-Butyl (2*R*,5S)-4-(6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2,4-diisopropylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.051 g, 0.087 mmol),2-fluorophenylboronic acid (0.015 ml, 0.104 mmol, CombiBlocks), sodium carbonate (0.028 g, 0.260 mmol) and tetrakis (10.00 mg, 8.65 µmol) in 1,4-dioxane/water (6/1.5 mL) was heated at 90 C for 1 h. The reaction went to completion, quenched with sat. $NaHCO_3$ and extracted with EtOAc. The combined organics were purified by chromatography on silica gel using 0-50 % 3:1 (EtOAc/EtOH) inheptane to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as a light solid. Theoretical yield considered. *m/z* (ESI, +ve ion): 648.6 $(M+H)^+$.

[0243] Step 3. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.056 g, 0.086 mmol) in DCM (5 mL) was added TFA (2.0 mL, 26.0 mmol) and the resulting mixture was stirred at rt for 30 min. The reaction went to completion and concentrated to afford 6-chloro-1-(2,4-diisopropylpyridin-3-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. *m/z* (ESI, +ve ion): 548.6 $(M+H)^+$.

[0244] 6-chloro-1-(2,4-diisopropylpyridin-3-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one was dissolved in DCM (5 mL) then acryloyl chloride (7.01 µl, 0.086 mmol) was added at rt. The reaction was stirred at rt for 15 min, washed with sat. $NaHCO_3$ and extracted with DCM. The combined organics were purified by HPLC to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.023 g, 0.038 mmol, 44.2 % yield) as a white solid. ^1H NMR (400 MHz, *DMSO-d*$_6$) δ 8.42 - 8.51 (m, 2 H), 7.45 - 7.58 (m, 1 H), 7.21 - 7.38 (m, 3 H), 7.18 (td, *J*=7.4, 1.6 Hz, 1 H), 6.84 (td, *J*=15.8, 10.5 Hz, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.72 - 5.82 (m, 1 H), 4.73 - 4.97 (m, 2 H), 4.47 (br dd, *J*=6.1, 3.4 Hz, 1 H), 4.12 - 4.23 (m, 1 H), 3.83 - 3.96 (m, 2 H), 3.51 (br dd, *J*=13.6, 3.6 Hz,

1 H), 1.31 - 1.40 (m, 3 H), 1.26 (br d, *J*=6.6 Hz, 2 H), 1.18 (d, *J*=6.6 Hz, 2 H), 1.07 (dd, *J*=6.6, 4.4 Hz, 6 H), 0.93 (t, *J*=7.7 Hz, 6 H). *m/z* (ESI, +ve ion): 602.6 $(M+H)^+$.

Table 124: Compounds 124-2 to 124-17 were prepared following the procedure described in Method 124, Steps 1-3, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
124-2	 Single isomer Peak 1	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC, **Step 2:** o-tolylboronic acid, Sigma-Aldrich Corporation
124-3	 Single isomer Peak 1	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-[2-(trifluoromethyl)phenyl]pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC, **Step 2:** 2-(trifluoromethyl)phenylboronic acid, Combi-Blocks Inc.
124-4	 Single isomer Peak 1	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluoro-6-hydroxy-phenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** (2-fluoro-6-hydroxyphenyl)boronic acid, WuXi

124-5		1-(2,4-Diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	2,6-dichloro-5-fluoronicotinamide, **Intermediate S**	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** o-tolylboronic acid, Sigma-Aldrich Corporation
124-6	 Single isomer Peak 2	6-Chloro-7-(2-chlorophenyl)-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** 2-chlorophenylboronic acid, Matrix Scientific
124-7		6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-phenyl-pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** phenylboronic acid, Sigma-Aldrich Corporation
124-8		6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** [2-(1-methylethyl)phenyl]-boronic acid, Combi-Blocks Inc.

124-9	 Single isomer Peak 1	6-Chloro-1-(4-cyclopropyl-2-isopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**Intermediate 199A/ Peak 1** in **Step 1**	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech
124-10	 Single isomer Peak 2	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2S)-2-methyl-4-prop-2-enoyl-piperazin-1-yl]-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	(s)-4-*N*-Boc-2-methyl piperazine in **Step 1**	**Step 1:** (s)-4-*N*-Boc-2-methyl piperazine, Sigma-Aldrich Corporation, **Step 2:** o-tolylboronic acid, Sigma-Aldrich Corporation
124-11		6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(3R,5S)-3,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	1-((2R,6S)-2,6-dimethyl piperazin-1-yl)prop-2-en-1-one, **Intermediate 223 Step 3:** Omit	**Step 1:** 1-((2R,6S)-2,6-dimethylpiperazin-1-yl)prop-2-en-1-one, Syngene, **Step 2:** o-tolylboronic acid, Sigma-Aldrich Corporation
124-12		(*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-[2-fluoro-5-(trifluoromethoxy)phenyl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1: Intermediate 73B**; **Step 2:** 2-fluoro-5-(trifluoromethoxy) benzeneboronic acid (Combi-Blocks, Inc.)

124-13		(*M*)-6-Chloro-7-(4,5-difluoro-2-hydroxy-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1: Intermediate 73B**; **Step 2:** 4,5-difluoro-2-hydroxyphenylboronic acid (Combi-Blocks, Inc., San Diego, CA)
124-14		(*M*)-4-[*cis*-2,3-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	Step 1: **Intermediate 72A** and 2,3-dimethyl-piperazine-1-carboxylic acid *tert*-butyl ester (Enamine, Monmouth Jct., NJ); **Step 2:** (2-fluoro-6-hydroxyphenyl)boronic acid (Wuxi)
124-16		(*M*)-4-[*trans*-2,3-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	Step 1: **Intermediate 72A** and 2,3-dimethyl-piperazine-1-carboxylic acid *tert*-butyl ester (Enamine, Monmouth Jct., NJ); Step 2: 2,3-dimethyl-piperazine-1-carboxylic acid (Alchem Pharmtech, Inc.)

Method 125

Example 125-1: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0245] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2,4-diisopropylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. Phosphorous oxychloride (0.206 mL, 2.206 mmol) was added dropwise to a solution of 6,7-dichloro-1-(2,4-diisopropylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.723 g, 1.838 mmol, **Intermediate 195B**) and 1,1'-dimethyltriethylamine (0.417 mL, 2.390 mmol) in CH3CN (5 mL) under N_2. This mixture was then heated to 80 °C for 30 min. LCMS showed chlorination completed. The reaction mixture was cooled to 0 °C and 3 equiv DIPEA were added followed by (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (0.433 mL, 2.022 mmol, AstaTech). This mixture was stirred with warming to rt over 30 min at which time LCMS showed conversion to desired product. The mixture was poured into cold satd. NaHCO3 solution and stirred vigorously for 10 min. The mixture was extracted with EtOAc, the combined organics were dried over Na_2SO_4, filtered, concentrated and purified by chromatography on silica gel using 0-70% EtOAc in heptane to afford a light yellow solid as *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2,4-diisopropylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.83 g, 1.408 mmol, 77 % yield). *m/z* (ESI, +ve ion): 588.6 $(M+H)^+$.

[0246] Step 2. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-

carboxylate. A mixture of *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2,4-diisopropylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.280 g, 0.475 mmol),2-fluorophenylboronic acid (0.080 ml, 0.570 mmol, CombiBlocks), sodium carbonate (0.151 g, 1.425 mmol) and tetrakis (0.055 g, 0.047 mmol) in 1,4-dioxane/water (6/1.5 mL) was heated at 90 C for 1 h. The reaction went to completion, quenched with sat. $NaHCO_3$ and extracted with EtOAc. The combined organics were purified by chromatography on silica gel using 0-50 % 3:1 (EtOAc/EtOH) inheptane to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.200 g, 0.308 mmol, 64.9 % yield). *m/z* (ESI, +ve ion): 648.6 $(M+H)^+$.

[0247] Step 3. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.200 g, 0.308 mmol) in DCM (3 mL) was added TFA (2.0 mL, 26.0 mmol) and the resulting mixture was stirred at rt for 30 min. The reaction went to completion, concentrated to afford 6-chloro-1-(2,4-diisopropylpyridin-3-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one to be used as is. *m/z* (ESI, +ve ion): 548.6 $(M+H)^+$.

[0248] 6-chloro-1-(2,4-diisopropylpyridin-3-yl)-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one was dissolved in DCM (6 mL) then acryloyl chloride (0.025 mL, 0.308 mmol) was added at rt. The reaction was stirred at rt for 15 min, washed with sat. $NaHCO_3$ and extracted with DCM. The combined organics were purified by HPLC to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.130 g, 0.216 mmol, 70.0 % yield) as a white solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.20 - 8.61 (m, 2 H), 7.44 - 7.59 (m, 1 H), 7.23 - 7.37 (m, 3 H), 7.10 - 7.22 (m, 1 H), 6.84 (td, *J*=16.1, 10.4 Hz, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.76 (ddd, *J*=10.3, 4.5, 2.3 Hz, 1 H), 4.72 - 4.97 (m, 2 H), 4.43 - 4.55 (m, 1 H), 4.10 - 4.24 (m, 2 H), 3.83 - 3.97 (m, 2 H), 1.31 - 1.40 (m, 3 H), 1.26 (br d, *J*=6.6 Hz, 2 H), 1.18 (br d, *J*=6.8 Hz, 2 H), 1.08 (br d, *J*=5.6 Hz, 6 H), 0.94 (t, *J*=6.5 Hz, 6 H). *m/z* (ESI, +ve ion): 602.6 $(M+H)^+$.

Table 125: Compounds 125-2 to 125-11 were prepared following the procedure described in Method 125, Steps 1-3, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
125-2	 Single isomer Peak 2	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC, **Step 2:** o-tolylboronic acid, Sigma-Aldrich Corporation
125-3	 Single isomer Peak 2	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-[2-(trifluoromethyl)phenyl]pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, eNovation Chemicals LLC, **Step 2:** 2-(trifluoromethyl)phenylboronic acid, Combi-Blocks Inc.
125-4	 Single isomer Peak 2	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluoro-6-hydroxy-phenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** (2-fluoro-6-hydroxyphenyl)boronic acid, WuXi

125-5	 Single isomer Peak 2	1-(2,4-Diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**Step 1:** 7-chloro-1-(2,4-diisopropylpyridin-3-yl)-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** o-tolylboronic acid, Sigma-Aldrich Corporation
125-6	 Single isomer Peak 1	6-Chloro-7-(2-chlorophenyl)-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** 2-chlorophenylboronic acid, Matrix Scientific
125-7	 Single isomer Peak 1	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-phenyl-pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** phenylboronic acid, Sigma-Aldrich Corporation
125-8	 Single isomer Peak 1	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** 2-isopropylphenylboronic acid, Combi-Blocks Inc.

125-9	 Single isomer Peak 2	6-Chloro-1-(4-cyclopropyl-2-isopropyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**Intermediate 199B/ Peak 2 In Step 1**	**Step 1:** (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate, Astatech, **Step 2:** o-tolylboronic acid, Sigma-Aldrich Corporation
125-10	 Single isomer Peak 1	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(2S)-2-methyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	None	**Step 1:** (s)-4-*N*-Boc-2-methyl piperazine, Sigma-Aldrich Corporation, **Step 2:** o-tolylboronic acid, Sigma-Aldrich Corporation
125-11	 Single isomer Peak 1	6-Chloro-1-(2,4-diisopropyl-3-pyridyl)-4-[(3R,5S)-3,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	1-((2R,6S)-2,6-dimethylpiperazin-1-yl)prop-2-en-1-one, **Intermediate 223** in **Step 1** Omit **Step 3**	**Step 2:** o-tolylboronic acid, Sigma-Aldrich Corporation

Method 129

Example 129-1: ***(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]acetamide**



Intermediate 233

[0249] A mixture of 4-((2*S*,5*R*,*M*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-amino-6-fluorophenyl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.11 g, 0.18 mmol, **Intermediate 233**) in dichloromethane (0.2 mL) was added acetic acid anhydride (0.1 mL, 0.98 mmol) and stirred at room temperature for 1 h. The resulting mixture was diluted with EtOAc (10 mL) and washed with saturated sodiom bicarbonate solution (10 mL). The aqueous layer was extracted with EtOAc (10 mL x 2). The combined organic extracts were washed with saturated NaCl solution (20 mL) and dried over MgSO4. The solution was filtered and concentrated *in vacuo* to give *(M)-N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]acetamide (0.10 mg, 0.17 mmol, 91 % yield) as a light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.73 - 9.11 (m, 1 H), 8.43 - 8.51 (m, 1 H), 8.37 (d, *J*=5.0 Hz, 1 H), 7.36 - 7.81 (m, 2 H), 7.14 - 7.22 (m, 1 H), 6.94 - 7.10 (m, 1 H), 6.73 - 6.92 (m, 1 H), 6.14 - 6.26 (m, 1 H), 5.76 (dt, *J*=10.2, 3.0 Hz, 1 H), 4.44 - 4.96 (m, 2 H), 3.76 - 4.25 (m, 4 H), 2.56 - 2.63 (m, 1 H), 2.22 (s, 3 H), 1.89 - 2.04 (m, 3 H), 1.34 (br t, *J*=7.2 Hz, 3 H), 1.15 - 1.27 (m, 3 H), 1.01 - 1.13 (m, 3 H), 0.79 - 0.99 (m, 3 H). ^{19}F NMR (376 MHz, DMSO-d_6) δ -116.97 - 112.40 (m, 1 F). *m/z* (ESI, +ve ion): 632.2 $(M+H)^+$.

Table 129: Compounds 129-2 to 129-13 were prepared following the procedure described in Method 129 above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
129-2		*(M)-N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-	None	2,2,2-trifluoroacetic anhydride (Sigma-Aldrich Corporation)

		pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]-2,2,2-trifluoro-acetamide		
129-3		*(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]phenyl]-*N*-methyl-acetamide	None	*(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-[2-(methylamino)phenyl]pyrido[2,3-*d*]pyrimidin-2-one **(Ex. 92-25)**
129-4		*(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]propanamide	None	propionyl chloride (Sigma-Aldrich Corporation)
129-5		*(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-	None	*(M)*-7-(2-Aminophenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-

		yl]phenyl]-2,2,2-trifluoro-acetamide		methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one **(Ex. 92-12)** and 2,2,2-trifluoroacetic anhydride (Sigma-Aldrich Corporation)
129-6		*(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]phenyl]cyclopropanecarboxamide	None	*(M)*-7-(2-Aminophenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one **(Ex. 92-12)** and cyclopropane carbonyl chloride (Sigma-Aldrich Corporation)
129-7		*(M)*-Methyl-*N*-[2-[6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]phenyl]carbamate	None	*(M)*-7-(2-Aminophenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-

				d]pyrimidin-2-one **(Ex. 92-12)** and methyl chloroformate (Sigma-Aldrich Corporation)
129-8		*(M)*-Methyl *N*-[2-[6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]carbamate	None	methyl chloroformate (Sigma-Aldrich Corporation)
129-9		*(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]cyclopropanecarboxamide	None	cyclopropane carbonyl chloride (Sigma-Aldrich Corporation)
129-10		*(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-	None	difluoroacetic anhydride (Matrix Scientific)

		phenyl]-2,2-difluoro-acetamide		
129-11		*(M)*-*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]phenyl]-2,2-difluoro-acetamide	None	*(M)*-7-(2-Aminophenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one **(Ex. 92-12)** and difluoroacetic anhydride (Matrix Scientific)
129-12		*N*-[2-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-6-fluoro-phenyl]acetamide	None	*(M)*-7-(2-Amino-3-fluoro-phenyl)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one **(Ex. 92-17)**

129-13		4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-[2-fluoro-6-(prop-2-ynylamino)phenyl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	Propargyl bromide, (Sigma-Aldrich Corporation), K_2CO_3 and DMF

Method 132

Example 132-1: *(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(3-hydroxy-1-naphthyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one



Example 92-70

[0250] A mixture of (*M*)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-methoxyphenyl)pyrido[2,3-*d*]pyrimidin-2-one (0.31 g, 0.49 mmol) in 1,2-dichloroethane (10 mL) was treated with boron tribromide solution (1.0 M in hexanes, 2.5 mL, 2.5 mmol) 0 °C and stirred for 1 h. The reaction mixture was added saturated sodium bicarbonate solution (10 mL) and extracted with EtOAc (10 mL x 2). The combined organic extracts were dried over Na2SO4. The solution was filtered and concentrated *in vacuo* to give the crude material. The crude material was purified by silica gel chromatography (eluent: 0-100% of EtOAc-MeOH (4:1)/heptane) to provide *(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(3-hydroxy-1-naphthyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one (0.26 g, 0.42 mmol, 85 % yield) as an yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 9.96 (s, 1 H), 8.49 (d, *J*=3.1 Hz, 1 H), 8.30 (d, *J*=4.8 Hz, 1 H), 7.73 (d, *J*=8.3 Hz, 1 H), 7.39 (ddd, *J*=8.1, 6.2, 1.9 Hz, 1 H), 7.15 - 7.25

(m, 3 H), 7.12 (d, *J*=4.8 Hz, 1 H), 6.78 - 6.95 (m, 2 H), 6.20 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.71 - 5.82 (m, 1 H), 4.89 (br s, 1 H), 4.46 - 4.84 (m, 1 H), 3.47 - 4.30 (m, 4 H), 2.70 - 2.80 (m, 1 H), 1.94 (s, 3 H), 1.32 - 1.41 (m, 3 H), 1.18 - 1.32 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.92 (d, *J*=6.6 Hz, 3 H). *m/z* (ESI, +ve ion): 632.2 $(M+H)^+$.

Table 132: Compound 132-2 was prepared following the procedure described in Method 132 above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
132-2		*(M)*-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(3-hydroxy-1-naphthyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	None	*(M)*-4-[(2*S*,5*R*)-2,5-Dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(3-methoxy-1-naphthyl)pyrido[2,3-*d*]pyrimidin-2-one **(Ex. 94-21)**

Method 141

Example 141-1

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-((diethylamino)methyl)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0251] Step 1. (2*R*,5*S*)-*tert*-Butyl 4-(1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6-chloro-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-

***d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**. To a solution of 4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropylpyridin-3-amine (1.0 g, 2.47 mmol, **Intermediate 207**) and *N,N*-diisopropylethylamine (0.950 ml, 5.44 mmol, Aldrich, St. Louis, MO) in tetrahydrofuran (25 mL) was added phosgene solution, 15% in toluene (1.30 ml, 2.74 mmol, Aldrich, St. Louis, MO) dropwise via syringe. After stirring for 10 min at rt, the resulting mixture was cooled to 0 °C and (2*R*,5*S*)-*tert*-butyl 4-((2,5-dichloro-6-(2-isopropylphenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (1.50 g, 2.97 mmol, Intermediate 170) was added as a solid. After the addition, the ice bath was removed and the reaction was warmed to rt and stirred for 1 h. To this mixture was added sodium *tert*-butoxide solution, 2.0 M in tetrahydrofuran (3.71 ml, 7.41 mmol, Aldrich, St. Louis, MO) dropwise via syringe and the mixture was stirred for 30 min. The reaction was quenched with water (3 mL) and partitioned between EtOAc (40 mL) and water (20 mL). The aqueous layer was extracted with EtOAc (30 mL). The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. The crude product was purified by silica gel chromatography (eluent: 5-40% acetone/heptane) to provide (2*R*,5*S*)-*tert*-butyl 4-(1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6-chloro-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as a brown foam (1.6 g). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.63-8.68 (m, 1H), 8.01 (d, *J*=3.32 Hz, 1H), 7.61 (s, 1H), 7.49-7.59 (m, 4H), 7.29-7.43 (m, 7H), 7.24 (d, *J*=6.84 Hz, 2H), 7.09-7.17 (m, 1H), 4.78-4.93 (m, 1H), 4.60-4.73 (m, 1H), 4.44-4.59 (m, 1H), 4.22-4.37 (m, 1H), 4.13 (d, *J*=7.26 Hz, 1H), 3.73-3.92 (m, 2H), 3.34-3.60 (m, 1H), 2.60-2.75 (m, 1H), 2.39-2.55 (m, 1H), 1.54 (d, *J*=14.31 Hz, 15H), 1.38 (s, 3H), 1.18-1.27 (m, 5H), 0.99-1.09 (m, 13H). *m/z* (ESI) M+H: 899.4.

[0252] Step 2. (2*R*,5*S*)-*tert*-Butyl 4-(6-chloro-1-(2-(hydroxymethyl)-6-isopropylphenyl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a solution of (2*R*,5*S*)-*tert*-butyl 4-(1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6-chloro-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (1.3 g, 1.44 mmol) in tetrahydrofuran (15 mL) was added tetrabutylammonium fluoride, 1.0 M in THF (1.60 mL, 1.60 mmol, Aldrich, St. Louis, MO). The resulting mixture was stirred at rt under N_2 for 2 h. The reaction was concentrated. The crude product was purified by silica gel chromatography (eluent: 0-40% acetone/heptane) to provide (2*R*,5*S*)-*tert*-butyl 4-(6-chloro-1-(2-(hydroxymethyl)-6-isopropylphenyl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-

d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as light yellow foam (0.740 g). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.61 (d, *J*=4.98 Hz, 1H), 8.16 (s, 1H), 8.07-8.13 (m, 1H), 7.38 (s, 1H), 7.30-7.34 (m, 2H), 7.19 (s, 1H), 6.95-7.03 (m, 1H), 4.78-4.97 (m, 1H), 4.52-4.73 (m, 1H), 4.34-4.42 (m, 1H), 4.21-4.33 (m, 1H), 3.94-4.10 (m, 1H), 3.73-3.88 (m, 1H), 3.49-3.69 (m, 1H), 2.68-2.80 (m, 1H), 2.37-2.58 (m, 2H), 1.56 (s, 6H), 1.53 (s, 9H), 1.26-1.30 (m, 3H), 1.18-1.24 (m, 3H), 0.97-1.09 (m, 6H). *m/z* (ESI) M+H: 661.2.

[0253] Step 3. (2*R*,5*S*)-*tert*-Butyl 4-(6-chloro-1-(2-formyl-6-isopropylphenyl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a solution of (2*R*,5*S*)-*tert*-butyl 4-(6-chloro-1-(2-(hydroxymethyl)-6-isopropylphenyl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.845 g, 1.28 mmol) in dichloromethane (12 mL) was added Dess-Martin periodinane (0.813 g, 1.92 mmol, Aldrich, St. Louis, MO). The resulting mixture was stirred at rt under N_2 for 2 h. The reaction was quenched with sat. $Na_2S_2O_3$ (10 mL) and stirred for 30 min. The aqueous layer was extracted with DCM (2 x 20 mL). The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. The crude product was purified by silica gel chromatography (eluent: 5-40% acetone/heptane) to provide (2*R*,5*S*)-*tert*-butyl 4-(6-chloro-1-(2-formyl-6-isopropylphenyl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as a white solid (0.150 g). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 9.86 (d, *J*=3.11 Hz, 1H), 8.82-8.89 (m, 1H), 8.16 (s, 1H), 7.55 (s, 1H), 7.35-7.40 (m, 1H), 7.31 (s, 1H), 7.17 (s, 1H), 6.89-7.03 (m, 1H), 4.88-5.12 (m, 1H), 4.54-4.70 (m, 1H), 4.33-4.53 (m, 1H), 4.02-4.24 (m, 1H), 3.84-3.96 (m, 1H), 3.46-3.70 (m, 1H), 2.90-3.06 (m, 1H), 2.35-2.53 (m, 1H), 1.55 (s, 6H), 1.53 (s, 9H), 1.25-1.30 (m, 6H), 1.04-1.10 (m, 3H), 0.94-1.03 (m, 3H). *m/z* (ESI) M+H: 659.2.

[0254] Step 4. (2*R*,5*S*)-*tert*-Butyl 4-(6-chloro-1-(4-((diethylamino)methyl)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a solution of (2*R*,5*S*)-*tert*-butyl 4-(6-chloro-1-(2-formyl-6-isopropylphenyl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.150 g, 0.228 mmol) in dichloromethane (3 mL) was added diethylamine (0.100 mL, 0.943 mmol, Aldrich, St. Louis, MO) and 2 drops of AcOH. After stirred for 10 min, sodium triacetoxyborohydride (0.145 g, 0.683 mmol, Aldrich, St. Louis, MO) was added and the resulting mixture was continued to stir at rt under N_2 for 1 h. Reaction was quenched with sat. $NaHCO_3$ (2 mL) and partitioned

between DCM (30 mL) and sat. $NaHCO_3$ (10 mL). The aqueous layer was extracted with DCM (30 mL). The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. The crude product was purified by silica gel chromatography (eluent: 3-5% 2M NH_3 MeOH/DCM) to provide (2*R*,5*S*)-*tert*-butyl 4-(6-chloro-1-(4-((diethylamino)methyl)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as white foam (0.120 g). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.49-8.56 (m, 1H), 8.07-8.16 (m, 1H), 7.39-7.45 (m, 1H), 7.37 (s, 1H), 7.33 (s, 1H), 7.17 (s, 1H), 6.95-7.03 (m, 1H), 4.87-5.08 (m, 1H), 4.37-4.66 (m, 1H), 4.01-4.22 (m, 1H), 3.85-3.98 (m, 1H), 3.75-3.85 (m, 1H), 3.49-3.68 (m, 1H), 3.34-3.46 (m, 1H), 2.99-3.13 (m, 1H), 2.68-2.82 (m, 1H), 2.50-2.63 (m, 1H), 2.29-2.42 (m, 2H), 2.18-2.29 (m, 2H), 1.53 (s, 9H), 1.42-1.49 (m, 3H), 1.28 (br s, 6H), 1.23 (d, *J*=6.84 Hz, 3H), 1.04 (s, 3H), 0.89 (s, 3H), 0.83 (br d, *J*=4.35 Hz, 6H). *m/z* (ESI) M+H: 716.4.

[0255] Step 5. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-((diethylamino)methyl)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of (2*R*,5*S*)-*tert*-butyl 4-(6-chloro-1-(4-((diethylamino)methyl)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.120 g, 0.168 mmol) in dichloromethane (3 mL) was added trifluoroacetic acid (0.262 mL, 3.52 mmol, Aldrich, St. Louis, MO). The resulting mixture was stirred at rt for 2 h. The mixture was concentrated, redissolved in DCM (3 mL) and treated with *N,N*-diisopropylethylamine (0.120 mL, 0.687 mmol, Aldrich, St. Louis, MO). The mixture was cooled to 0 °C and treated with acryloyl chloride (0.170 mL, 0.187 mmol, 1.1 M, Aldrich, St. Louis, MO). After the addition, the ice bath was removed and the reaction was warmed up to rt and continued to stir for 30 min. The reaction was quenched with sat. $NaHCO_3$ (3 mL) and extracted with DCM (2 x 10 mL). The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. The crude product was purified by silica gel chromatography (eluent: 3-5% 2M NH_3 MeOH/DCM) to provide 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-((diethylamino)methyl)-2-isopropylpyridin-3-yl)-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a white solid (0.100 g, **Example 141-1**). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.49-8.56 (m, 1H), 8.09-8.17 (m, 1H), 7.35-7.47 (m, 2H), 7.30-7.35 (m, 1H), 7.14-7.22 (m, 1H), 6.96-7.04 (m, 1H), 6.52-6.72 (m, 1H), 6.34-6.47 (m, 1H), 5.75-5.85 (m, 1H), 5.03-5.19 (m, 1H), 4.33-4.55 (m, 1H), 3.88-4.12 (m, 2H), 3.67-3.82 (m, 1H), 3.35-3.47 (m, 1H), 3.00-3.12 (m, 1H), 2.67-2.82 (m, 1H), 2.52-2.64 (m, 1H), 2.30-2.41 (m, 2H),

2.21-2.29 (m, 2H), 1.46 (br d, *J*=6.43 Hz, 4H), 1.28 (br s, 6H), 1.23 (br d, *J*=6.63 Hz, 3H), 1.03 (br d, *J*=6.63 Hz, 6H), 0.83 (br d, *J*=3.73 Hz, 6H). m/z (ESI) M+H: 670.2.

Table 141: Compounds 141-2 to 141-3 were prepared following the procedure described in Method 141, Steps 1-5, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
141-2		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-[4-[[ethyl(isopropyl)amino]methyl]-2-isopropyl-3-pyridyl]-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one	None	*N*-Ethylisopropylamine (Aldrich, St. Louis, MO) instead of diethylamine in Step 4
141-3		6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-isopropylphenyl)-1-[2-isopropyl-4-(pyrrolidin-1-ylmethyl)-3-pyridyl]pyrido[2,3-*d*]pyrimidin-2-one	None	Pyrrolidine (Aldrich, St. Louis, MO) instead of diethylamine in Step 4

Method 144

Example 144-1: (*M*)-2-[4-[7-(2-Amino-6-fluoro-phenyl)-6-chloro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile



[0256] Step 1: (*M*)-2-[4-[6,7-Dichloro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile. A mixture of (*M*)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 73B**, 0.43 g, 1.16 mmol) and DIPEA (1.0 mL, 5.82 mmol) in acetonitrile (5.0 mL) was treated with phosphorous oxychloride (0.3 mL, 3.49 mmol) at room temperature. The mixture was stirred at 80 °C for 30 min. The reaction mixture was evaporated to dryness to give 4,6,7-trichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a orange oil. LC/MS purity: >95% (215 nm); >95% (254 nm). *m/z* (ESI, +ve ion): 383.1 $(M+H)^+$. The resulting brown solid was used in next step without purification.

[0257] To a mixture of the above crude 4,6,7-trichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one in acetonitrile (5.0 mL) was added a solution of 2-(piperazin-2-yl)acetonitrile dihydrochloride (0.35 g, 1.75 mmol) and DIPEA (1.0 mL, 5.82 mmol) in acetonitrile (2.0 mL). The resulting mixture was stirred at room temperature for 10 min and then treated with acryloyl chloride (1.1 M solution in DCM 1.6 mL, 1.75 mmol). The resulting mixture was stirred at room temperature for 10 min and evaporated to dryness. The resulting crude product was purified by silica gel chromatography (eluent: 0-100% of EtOAc-MeOH (9:1)/heptane) to provide (*M*)-2-[4-[6,7-dichloro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile (0.36 g, 0.68 mmol, 58.5 % yield) as an yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.66 (s, 1 H), 8.52 (d, *J*=5.0 Hz, 1 H), 7.34 (br s, 1 H), 6.73 - 7.00 (m, 1 H), 6.22 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.73 - 5.89 (m, 1 H), 4.81 - 5.02 (m, 1 H), 4.26 - 4.41 (m, 2 H), 3.79 - 4.09 (m, 3 H), 3.57 (d, *J*=11.0 Hz, 2 H), 2.97 - 3.07 (m, 1 H), 2.59 - 2.82 (m, 1 H), 1.91 - 2.04 (m, 3 H), 1.08 (dd, *J*=6.6, 3.5 Hz, 3 H), 1.02 (dd, *J*=6.6, 2.7 Hz, 3 H). *m/z* (ESI, +ve ion): 526.1$(M+H)^+$.

[0258] Step 2: (*M*)-2-[4-[7-(2-Amino-6-fluoro-phenyl)-6-chloro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile. A mixture of (*M*)-2-[4-[6,7-dichloro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile (0.11 g, 0.21 mmol), tetrakis(triphenylphosphine)palladium (0.02 g, 0.02 mmol), 2-fluoro-6-aminophenylboronic acid pinacol ester (0.07 g, 0.31 mmol, CombiPhos Catalysts, Inc.) and sodium carbonate (0.11 g, 1.04 mmol) in 1,4-dioxane (1.0 mL)/ water (0.5 mL) was stirred at 90° for 30 min. The resulting mixture was evaporated to dryness. The resulting crude product was purified by silica gel chromatography (eluent: 0-100% of EtOAc/heptane) to provide (*M*)-2-[4-[7-(2-amino-6-fluoro-phenyl)-6-chloro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-

pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile (0.06 g, 0.10 mmol, 48.1 % yield) as an yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.59 - 8.65 (m, 1 H), 8.36 - 8.44 (m, 1 H), 7.19 (d, *J*=5.0 Hz, 1 H), 7.06 (q, *J*=7.9 Hz, 1 H), 6.77 - 6.98 (m, 1 H), 6.41 - 6.49 (m, 1 H), 6.32 (br t, *J*=8.3 Hz, 1 H), 6.23 (dd, *J*=16.6, 1.7 Hz, 1 H), 5.81 (br d, *J*=11.8 Hz, 1 H), 5.10 - 5.18 (m, 1 H), 5.05 (br s, 1 H), 4.83 - 5.01 (m, 1 H), 4.02 - 4.48 (m, 3 H), 3.45 - 4.00 (m, 3 H), 3.13 - 3.21 (m, 1 H), 2.98 - 3.13 (m, 1 H), 2.63 - 2.93 (m, 1 H), 1.81 - 2.02 (m, 3 H), 1.03 - 1.12 (m, 3 H), 0.86 - 1.01 (m, 3 H). ^{19}F NMR (376 MHz, DMSO-d_6) δ -116.31 - -115.08 (m, 1 F). *m/z* (ESI, +ve ion): 601.2 $(M+H)^+$.

Table 144: Compounds 144-2 to 144-3 were prepared following the procedure described in Method 144, Steps 1-2 above, as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
144-2		(*M*)-2-[4-[6,7-Dichloro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile	Omit step 2	No change
144-3		(*M*)-2-[4-[6-Chloro-7-(2-fluoro-6-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile	**Step 4:** Pd(dppf)2 Cl2 and KOAc	**Step 2:** (2-fluoro-6-hydroxyphenyl)boronic acid (WuXi)

Method 145

Example 145-1: 2-[4-[6-Chloro-1-(2,2-dimethylpropyl)-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile



[0259] Step 1: 2,5-Dichloro-6-(2-fluorophenyl)-*N*-(neopentylcarbamoyl)- nicotinamide. A solution of 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (**Intermediate 99B**, 2.75 g, 9.65 mmol) in THF (20 mL) was treated with oxalyl chloride (2.0 M solution in DCM, 5.8 mL, 11.57 mmol). The mixture was stirred at 75 °C for 60 min. The mixture was cooled to room temperature and concentrated *in vacuo*. The above crude material was redissolved into THF (20 mL) and a mixture of neopentylamine (1.1 mL, 9.65 mmol) and huenigs base (3.4 mL, 19.29 mmol) in THF (10 mL) was added. The resulting mixture was stirred at room temperature for 3 days. The reaction mixture was then partitioned between water (50 mL) and EtOAc (50 mL). The aqueous phase was extracted with EtOAc (50 mL). The combined organic phases were dried over Na2SO4 and concentrated to give a crude material, which was purified by silica gel chromatography (eluent: 0-50% of EtOAc/heptane) to provide 2,5-dichloro-6-(2-fluorophenyl)-*N*-(neopentylcarbamoyl)nicotinamide (1.09 g, 2.74 mmol, 28.4 % yield) as a light yellow solid. *m/z* (ESI, +ve ion): 398.0 $(M+H)^+$.

[0260] Step 2: 6-Chloro-7-(2-fluorophenyl)-1-neopentylpyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A mixture of 2,5-dichloro-6-(2-fluorophenyl)-*N*-(neopentylcarbamoyl)-nicotinamide (1.09 g, 2.74 mmol) in THF (10 mL) at 0 °C was treated with potassium bis(trimethylsilyl)amide (1.0 M in THF, 5.5 mL, 5.50 mmol) and stirred at 0 °C for 15 min. The reaction mixture was quenched with saturated aqueous NH4Cl (10 mL) and extracted with EtOAc (20 mL x 2). The combined organic layers were washed with brine (50 mL), dried over Na2SO4, filtered, and concentrated *in vacuo*. The resulting crude product was sonicated in EtOAc (3 mL). The resulting solid was collected by filtration, washed with EtOAc and dried to give pure 6-chloro-7-(2-fluorophenyl)-1-neopentylpyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.75 g, 2.06 mmol, 75 % yield) as a light yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6)

δ 11.88 (s, 1 H), 8.44 (s, 1 H), 7.56 - 7.66 (m, 1 H), 7.45 - 7.53 (m, 1 H), 7.35 - 7.44 (m, 2 H), 4.07 (s, 2 H), 0.90 (s, 9 H). ^{19}F NMR (376 MHz, DMSO-d_6) δ -114.02 (s, 1 F). *m/z* (ESI, +ve ion): 362.2 $(M+H)^+$.

[0261] Step 3: 2-[4-[6-Chloro-1-(2,2-dimethylpropyl)-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile. A mixture of 6-chloro-7-(2-fluorophenyl)-1-neopentylpyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.35 g, 0.97 mmol) and n,n'-diisopropylethylamine (0.8 mL, 4.84 mmol) in acetonitrile (5.0 mL) was treated with phosphorous oxychloride (0.3 mL, 2.90 mmol) at room temperature. The mixture was stirred at 80 °C for 1 h. The reaction mixture was evaporated to dryness to give 4,6-dichloro-7-(2-fluorophenyl)-1-neopentylpyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a brown oil. *m/z* (ESI, +ve ion): 380.0 $(M+H)^+$. The resulting brown oil was used in next step without purification.

[0262] To the mixture of 4,6-dichloro-7-(2-fluorophenyl)-1-neopentylpyrido[2,3-*d*]pyrimidin-2(1*H*)-one in acetonitrile (5.0 mL) was added a solution of 2-(piperazin-2-yl)acetonitrile dihydrochloride (0.29 g, 1.45 mmol, Enamine Ltd.) and n,n'-diisopropylethylamine (0.8 mL, 4.84 mmol) in acetonitrile (2.0 mL) and stirred at rt for 10 min. The resulting mixture was then treated with acryloyl chloride (1.1 M in DCM, 1.3 mL, 1.45 mmol) and stirred at rt for 10 min. The reaction mixture was evaporated to dryness. The resulting crude product was purified by silica gel chromatography (0-100% of EtOAc-MeOH (9:1)/heptane) to provide 2-[4-[6-chloro-1-(2,2-dimethylpropyl)-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile (0.23 g, 0.43 mmol, 44.7 % yield) as an yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (s, 1 H), 7.49 - 7.70 (m, 2 H), 7.33 - 7.46 (m, 2 H), 6.73 - 7.01 (m, 1 H), 6.20 (dd, *J*=16.6, 2.1 Hz, 1 H), 5.79 (br d, *J*=11.8 Hz, 1 H), 4.75 - 5.01 (m, 1 H), 4.03 - 4.37 (m, 5 H), 3.42 - 4.01 (m, 3 H), 2.90 - 3.20 (m, 2 H), 0.88 (s, 9 H). ^{19}F NMR (376 MHz, DMSO-d_6) δ -113.65 (s, 1 F). *m/z* (ESI, +ve ion): 523.3 $(M+H)^+$.

Table 145: Compounds 145-2 to 145-3 were prepared following the procedure described in Method 145, Steps 1-3, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent

145-2		(*M*)-2-[4-[6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile	Omit Step 1 and Step 2	**Step 3:** (*M*)-6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione **(Intermediate 76A)**
145-3		2-[1-[6-Chloro-1-(2,2-dimethylpropyl)-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-4-prop-2-enoyl-piperazin-2-yl]acetonitrile	None	**Step 3:** 2-(4-Acryloylpiperazin-2-yl)acetonitrile **(Intermediate 218)**

Method 146

Example 146-1: (*P*)-(9*Z*)-19-Chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one and (*P*)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one



[0263] **Step 1: *N*-((2-Allyl-6-isopropylphenyl)carbamoyl)-2,5,6-trichloronicotinamide.** A 100-mL round-bottomed flask was charged with 2,5,6-trichloronicotinamide (1.89 g, 8.38 mmol, WuXi) and tetrahydrofuran (32 ml). Oxalyl chloride (2.0 M solution in DCM, 5.0 mL, 10.06 mmol) was added to the reaction mixture and stirred at 75 °C for 1.5 h. The mixture was allowed to cool to rt and concentrated *in vacuo*.

[0264] The resulting residue was redissolved into tetrahydrofuran (32 mL) and treated with 2-(prop-2-en-1-yl)-6-(propan-2-yl)aniline (1.5 mL, 8.38 mmol, Enamine Ltd.). The mixture was stirred at rt for 10 min. The reaction mixture was partitioned between water (50 mL) and EtOAc (50 mL). The aqueous phase was extracted with EtOAc (50 mL). The aqueous phase was then saturated with solid NaCl and further extracted with EtOAc (200 mL). The combined organic phases were dried over Na_2SO_4 and concentrated to give *N*-((2-allyl-6-isopropylphenyl)carbamoyl)-2,5,6-trichloronicotinamide (3.81 g, 8.94 mmol, 100 % yield) as a tan foam. *m/z* (ESI, +ve ion): 426.0 $(M+H)^+$.

[0265] **Step 2: 1-(2-Allyl-6-isopropylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione.** A mixture of *N*-((2-allyl-6-isopropylphenyl)carbamoyl)-2,5,6-trichloronicotinamide (3.58 g, 8.38 mmol) in tetrahydrofuran (27.9 mL) at 0 °C was treated with potassium bis(trimethylsilyl)amide (1.0 M in THF, 16.8 mL, 16.76 mmol) The resulting mixture was stirred at 0 °C for 10 min. The reaction mixture was quenched with saturated aqueous NH_4Cl (30 mL) and extracted with EtOAc (30 mL x 2). The organic layers were

combined, washed with brine (50 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The resulting crude product was sonicated in MeOH (20 mL). The resulting solid was collected by filtration, washed with MeOH and dried to give pure 1-(2-allyl-6-isopropylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (2.636 g, 6.75 mmol, 81 % yield) as a tan solid. LC/MS purity: 99% (215 nm); 99% (254 nm). ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.18 (s, 1 H), 8.56 (s, 1 H), 7.32 - 7.49 (m, 2 H), 7.20 (dd, *J*=7.3, 1.5 Hz, 1 H), 5.66 (ddt, *J*=16.9, 10.1, 6.8, 6.8 Hz, 1 H), 4.80 - 4.90 (m, 2 H), 3.10 (d, *J*=6.6 Hz, 2 H), 2.64 - 2.74 (m, 1 H), 1.08 (d, *J*=6.8 Hz, 3 H), 0.99 (d, *J*=6.6 Hz, 3 H). *m/z* (ESI, +ve ion): 390.0 $(M+H)^+$.

[0266] Step 3: (*P*)-1-(2-Allyl-6-isopropylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione and (*M*)-1-(2-Allyl-6-isopropylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A mixture of atropisomers 1-(2-allyl-6-isopropylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (2.15 g) was purified by SFC (WelkoSS, 5 µm, 21 x 250 mm two in series, total 50 cm, 15% MeOH/CO2, 80 mL/min, 90 bar) to obtain two peaks: Peak 1 (**Intermediate 146A**, (*P*)-isomer, 780 mg, >99%ee) and Peak 2 (**Intermediate 146B**, (*M*)-isomer, 850 mg, 95.3% ee).

[0267] Step 4: (*P*)-*tert*-Butyl (2*R*,5*S*)-4-(1-(2-allyl-6-isopropylphenyl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of (*P*)-1-(2-allyl-6-isopropylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (Peak 1) (780 mg, 2.00 mmol) and DIPEA (1.75 mL, 9.99 mmol) in acetonitrile (10 mL) was treated with phosphorous oxychloride (0.56 mL, 6.00 mmol) at room temperature. The mixture was stirred at 80 °C for 30 min. The reaction mixture was evaporated to dryness to give (*P*)-1-(2-allyl-6-isopropylphenyl)-4,6,7-trichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a brown oil. *m/z* (ESI, +ve ion): 408.0 $(M+H)^+$. The resulting brown solid was used in next step without purification.

[0268] The mixture of (*P*)-1-(2-allyl-6-isopropylphenyl)-4,6,7-trichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (1.75 mL, 9.99 mmol) in acetonitrile (10 mL) was added *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (578 mg, 2.70 mmol, AstaTech) and stirred at rt for 15 min. The resulting mixture was diluted with water (15 mL) and extracted with EtOAc (3 x 15 mL). The organic extract was dried over $MgSO_4$. The solution was filtered and concentrated *in vacuo* to give the crude material as a brown solid. The crude material brown solid which was purified by silica gel chromatography (0-50% of EtOAc/heptane) to provide (*P*)-*tert*-butyl (2*R*,5*S*)-4-(1-(2-allyl-6-isopropylphenyl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (696 mg, 1.19

mmol, 59.4 % yield) as a light yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.38 (s, 1 H), 7.29 - 7.46 (m, 2 H), 7.18 (dd, *J*=7.3, 1.2 Hz, 1 H), 5.66 (ddt, *J*=17.8, 9.2, 6.8, 6.8 Hz, 1 H), 4.82 - 4.86 (m, 1 H), 4.81 (s, 1 H), 4.75 (br s, 1 H), 4.16 - 4.42 (m, 1 H), 4.08 (br d, *J*=13.7 Hz, 1 H), 3.79 (br s, 1 H), 3.67 (br d, *J*=12.9 Hz, 1 H), 3.34 - 3.51 (m, 1 H), 2.93 - 3.11 (m, 2 H), 2.42 (dt, *J*=13.7, 6.8 Hz, 1 H), 1.44 (s, 9 H), 1.32 (d, *J*=6.6 Hz, 3 H), 1.13 (br d, *J*=6.6 Hz, 3 H), 1.04 (d, *J*=6.8 Hz, 3 H), 0.97 (d, *J*=6.8 Hz, 3 H). *m/z* (ESI, +ve ion): 526.2 $(M+H)^+$.

[0269] Step 5: (*P*)-*tert*-Butyl (2*R*,5*S*)-4-(1-(2-allyl-6-isopropylphenyl)-7-(2-allylphenyl)-6-chloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of (*P*)-*tert*-butyl (2*R*,5*S*)-4-(1-(2-allyl-6-isopropylphenyl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (250 mg, 0.43 mmol), 2-(2-allylphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (156 mg, 0.64 mmol), tetrakis(triphenylphosphine)palladium (49.3 mg, 0.04 mmol), and sodium carbonate (226 mg, 2.13 mmol) in 1,4-dioxane (1.5 mL)/ water (0.8 mL) was stirred at 90° for 1 h. The resulting mixture was evaporated to dryness. The resulting crude product was purified by silica gel chromatography (0-50% of EtOAc/heptane) to provide (*P*)-*tert*-butyl (2*R*,5*S*)-4-(1-(2-allyl-6-isopropylphenyl)-7-(2-allylphenyl)-6-chloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (169 mg, 0.25 mmol, 59.5 % yield) a light yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.38 (s, 1 H), 7.31 - 7.40 (m, 1 H), 7.19 - 7.31 (m, 4 H), 7.06 - 7.14 (m, 2 H), 5.60 - 5.75 (m, 1 H), 5.49 (ddt, *J*=16.9, 10.1, 6.7, 6.7 Hz, 1 H), 4.74 - 4.91 (m, 5 H), 4.22 - 4.43 (m, 1 H), 4.09 (br d, *J*=13.9 Hz, 1 H), 3.81 - 3.90 (m, 1 H), 3.70 (dd, *J*=13.5, 2.3 Hz, 1 H), 3.38 - 3.58 (m, 1 H), 2.90 - 3.14 (m, 4 H), 2.43 - 2.47 (m, 1 H), 1.45 (s, 9 H), 1.34 (d, *J*=6.4 Hz, 3 H), 1.16 (br d, *J*=6.4 Hz, 3 H), 1.03 (d, *J*=6.8 Hz, 3 H), 0.90 (d, *J*=6.8 Hz, 3 H). *m/z* (ESI, +ve ion): 668.2 $(M+H)^+$.

[0270] Step 6: (*P*)-*tert*-Butyl (2*R*,5*S*)-4-(26-chloro-16-isopropyl-22-oxo-21,22-dihydro-2(1,7)-pyrido[2,3-*d*]pyrimidina-1,3(1,2)-dibenzenacycloheptaphan-5-en-24-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of Grubbs Catalyst, 2nd generation (tricyclohexylphosphine[1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene][benzylidine]rut) (43 mg, 0.05 mmol, Sigma-Aldrich Corporation), (*P*)-*tert*-butyl (2*R*,5*S*)-4-(1-(2-allyl-6-isopropylphenyl)-7-(2-allylphenyl)-6-chloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (168 mg, 0.25 mmol) in dichloromethane (14 mL) was stirred at 45° for 1 h. The resulting mixture was evaporated to dryness. The resulting crude product was purified by silica gel chromatography (0-50% of EtOAc/heptane) to provide (*P*)-*tert*-butyl (2*R*,5*S*)-4-(26-chloro-16-isopropyl-22-

oxo-21,22-dihydro-2(1,7)-pyrido[2,3-*d*]pyrimidina-1,3(1,2)-dibenzenacycloheptaphan-5-en-24-yl)-2,5-dimethylpiperazine-1-carboxylate (120 mg, 0.19 mmol, 74.6 % yield) as a light yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.41 - 8.48 (m, 1 H), 7.22 - 7.40 (m, 5 H), 7.04 - 7.14 (m, 2 H), 4.86 - 4.95 (m, 1 H), 4.80 - 4.84 (m, 1 H), 4.17 - 4.36 (m, 1 H), 3.93 - 4.04 (m, 1 H), 3.85 - 3.91 (m, 1 H), 3.50 - 3.75 (m, 2 H), 3.22 - 3.29 (m, 1 H), 3.02 (br dd, *J*=15.8, 3.3 Hz, 1 H), 2.86 (br dd, *J*=15.0, 9.0 Hz, 1 H), 2.61 - 2.80 (m, 2 H), 2.52 - 2.57 (m, 1 H), 1.45 (s, 9 H), 1.27 (br d, *J*=5.0 Hz, 3 H), 1.07 (br d, *J*=6.8 Hz, 6 H), 0.93 (d, *J*=6.8 Hz, 3 H). *m/z* (ESI, +ve ion): 640.2 $(M+H)^+$.

[0271] Step 7: A mixture of (*P*)-(9*Z*)-19-Chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one and (*P*)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one. To a solution of (*P*)-*tert*-butyl (2*R*,5*S*)-4-(26-chloro-16-isopropyl-22-oxo-21,22-dihydro-2(1,7)-pyrido[2,3-*d*]pyrimidina-1,3(1,2)-dibenzenacycloheptaphan-5-en-24-yl)-2,5-dimethylpiperazine-1-carboxylate (120 mg, 0.19 mmol) in dichloromethane (2.0 mL) was treated with 2,2,2-trifluoroacetic acid (2.0 mL, 0.19 mmol) at rt and stirred for 1.5 h. The reaction went to completion and the resulting mixture was concentrated to afford (*P*)-26-chloro-24-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-16-isopropyl-21,22-dihydro-2(1,7)-pyrido[2,3-*d*]pyrimidina-1,3(1,2)-dibenzenacycloheptaphan-5-en-22-one, *m/z* (ESI, +ve ion): 540.2 $(M+H)^+$.

[0272] A mixture of the above crude product and DIPEA (0.2 mL, 0.94 mmol) in dichloromethane (2.0 mL) was added acryloyl chloride (1.1 M in DCM, 0.2 mL, 0.19 mmol) at 0 °C and stirred for 5 min. The resulting mixture was evaporated to dryness. The resulting crude product was purified by silica gel chromatography (0-100% of EtOAc/heptane) to provide a mixture of (*P*)-(9*Z*)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one and (*P*)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one (1:6 ratio, 94 mg, 0.16 mmol, 84 % yield) as a light yellow foam. ^{1}H NMR

(400 MHz, DMSO-d_6) δ 8.48 (br d, J=8.7 Hz, 1 H), 7.22 - 7.37 (m, 5 H), 7.05 - 7.17 (m, 2 H), 6.76 - 6.93 (m, 1 H), 6.19 (dd, J=16.7, 2.2 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 4.95 (br s, 1 H), 4.84 (br d, J=4.6 Hz, 1 H), 4.40 - 4.77 (m, 1 H), 3.56 - 4.18 (m, 4 H), 3.23 - 3.29 (m, 1 H), 3.03 (br dd, J=15.4, 3.2 Hz, 1 H), 2.83 - 2.93 (m, 1 H), 2.74 - 2.82 (m, 1 H), 2.64 - 2.73 (m, 1 H), 2.54 - 2.60 (m, 1 H), 1.27 (br dd, J=18.1, 6.5 Hz, 3 H), 1.05 - 1.21 (m, 6 H), 0.94 (d, J=6.8 Hz, 3 H). m/z (ESI, +ve ion): 594.2 $(M+H)^+$.

Table 146: Compounds 146-2 to 146-3 were prepared following the procedure described in Method 146, Steps 1-7, above as follows:

Ex. #	Chemical structure	Name	Method Changes	Reagent
146-2		(*M*)-(9*Z*)-19-Chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one, (*M*)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one	None	**Step 3: Intermediate 146B**

146-3	   	(*P*)-19-Chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one, (*P*)-(9Z)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one, (*M*)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one, (*M*)-(9Z)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one	Omit Step 3	**Step 1:** 2-allylaniline hydrochloride salt (TCI America)
146-4	 	20-chloro-23-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-12-oxa-1,24,27-triazapentacyclo[17.6.2.0~2,7~.0~13,18~.0~22,26~]heptacosa-2,4,6,9,13,15,17,19,21,23,26-undecaen-25-one	After **Step 5**, added an alkylation step (allyl chloride, K_2CO_3, CH_3CN, RT)	**Step 5:** 2-allyloxyphenylboronic acid (Combi-Blocks Inc.)

Table 147: Separated Compound Examples From Methods

Ex.#	Chemical Structure	Name	Separation conditions
55-51-1	 1st-eluting isomer	6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC:** (R,R) Whelk-01 (150 x 4.6 mm, 5 µm); Mobile Phase: 70:30 B: methanol with 0.2% TEA Flow Rate: 4.0 mL/min Oven/Column Temp.: 23 C Outlet Pressure: 100 bar
55-51-2	 2nd-eluting isomer	6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC:** (R,R) Whelk-01 (150 x 4.6 mm, 5 µm) Mobile Phase: 70:30 B: methanol with 0.2% TEA Flow Rate: 4.0 mL/min Oven/Column Temp.: 23 C Outlet Pressure: 100 bar
55-58-1	 1st-eluting isomer	6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(4-ethyl-2-isopropyl-3-pyridyl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC:** IA (150 x 4.6 mm, 5 µm) Mobile Phase: 80:20 B: ethanol Flow Rate: 4.0 mL/min Oven/Column Temp.: 23 C Outlet Pressure: 100 bar

55-58-2	 2nd-eluting isomer	6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(4-ethyl-2-isopropyl-3-pyridyl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC:** IA (150 x 4.6 mm, 5 µm) Mobile Phase: 80:20 B: ethanol Flow Rate: 4.0 mL/min Oven/Column Temp.: 23 C Outlet Pressure: 100 bar
80-10-1	 1st-eluting isomer	6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-[2-isopropyl-6-(pyrrolidin-1-ylmethyl)phenyl]pyrido[2,3-*d*]pyrimidin-2-one	**SFC** Whelk-01 (250 X 21 mm, 5µm), 45:55 methanol with 0.2% TEA:CO2, 50 g/min, 102 bar
80-10-2	 2nd-eluting isomer	6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-[2-isopropyl-6-(pyrrolidin-1-ylmethyl)phenyl]pyrido[2,3-*d*]pyrimidin-2-one	**SFC** Whelk-01 (250 X 21 mm, 5µm), 45:55 methanol with 0.2% TEA:CO2, 50 g/min, 102 bar
80-14-1	 1st-eluting isomer	6-chloro-1-[2-[(dimethylamino)methyl]-6-isopropyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** Chiralpak ID (250 x 20 mm, 220-nm), 45:55 methanol with 0.2% TEA:CO_2, 80 ml/min, 207 bar

80-14-2	 2nd-eluting isomer	6-chloro-1-[2-[(dimethylamino)methyl]-6-isopropyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** Chiralpak ID (250 x 20 mm, 220-nm), 45:55 methanol with 0.2% TEA:CO_2, 80 ml/min, 207 bar
80-16-1	 1st-eluting isomer	6-chloro-1-[4-[(dimethylamino)methyl]-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** OX-H (150 x 21 mm, 5 μm), 30:70 methanol with 0.2% TEA:CO_2, 70 ml/min
80-16-2	 2nd -eluting isomer	6-chloro-1-[4-[(dimethylamino)methyl]-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** OX-H (150 x 21 mm, 5 μm), 30:70 methanol with 0.2% TEA:CO_2, 70 ml/min
80-18-1	 1st-eluting isomer	6-chloro-1-[4-[(dimethylamino)methyl]-6-isopropyl-pyrimidin-5-yl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** OD (250 x 21 mm, 5 μm), 15:85 methanol with 0.2% TEA:CO_2, 80 ml/min, 102 bar followed by **SFC** OZ (250 x 21 mm, 5 μm), 30:70 methanol with 0.2% TEA:CO_2, 65 ml/min, 102 bar
80-18-2	 2nd -eluting isomer	6-chloro-1-[4-[(dimethylamino)methyl]-6-isopropyl-pyrimidin-5-yl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** OD (250 x 21 mm, 5 μm), 15:85 methanol with 0.2% TEA:CO_2, 80 ml/min, 102 bar

80-20-1	 1st-eluting isomer	1-[4-[(Dimethylamino)methyl]-2-isopropyl-6-methyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	Preparative Thar 200 SFC method: Column: (R,R) whelk-01 (250 X 21 mm, 5u) Mobile Phase: 75:25 (A:B) A: Liquid CO2 B: methanol with 0.2% TEA Flow Rate: 60 g/min Column/Oven temp.: ambient temperature 220 nm BPR = 102 bar
80-20-2	 2nd-eluting isomer	1-[4-[(Dimethylamino)methyl]-2-isopropyl-6-methyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	Preparative Thar 200 SFC method: Column: (R,R) whelk-01 (250 X 21 mm, 5u) Mobile Phase: 75:25 (A:B) A: Liquid CO2 B: methanol with 0.2% TEA Flow Rate: 60 g/min Column/Oven temp.: ambient temperature 220 nm BPR = 102 bar
80-21-1	 1st-eluting isomer	1-[4-[(Dimethylamino)methyl]-2-isopropyl-6-methyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** Whelk-01 (R,R) (250 X 30 mm, 5 μm) 30% MeOH (with 20 mM NH_3)/CO_2, 180 g/min, 102 bar
80-21-2		1-[4-[(Dimethylamino)methyl]-2-isopropyl-6-methyl-phenyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-	**SFC** Whelk-01 (R,R) (250 X 30 mm, 5 μm) 30% MeOH (with 20 mM NH_3)/CO_2, 180 g/min, 102 bar

	2nd-eluting isomer	phenyl)pyrido[2,3-*d*]pyrimidin-2-one	
100-1-1	 1st-eluting isomer	6-Chloro-1-(2,4-diisopropylpyrazol-3-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** OD-H column (5 µm, 2 × 21 × 250 mm, two columns in series, total length 50 cm) F=80 ml, 20% Methanol, 80% carbon dioxide, Back pressure= 90 bars
100-1-2	 2nd-eluting isomer	6-Chloro-1-(2,4-diisopropylpyrazol-3-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** OD-H column (5 µm, 2 × 21 × 250 mm, two columns in series, total length 50 cm) F=80 ml, 20% Methanol, 80% carbon dioxide, Back pressure= 90 bars
102-5-1	 1st-eluting isomer	6-Chloro-1-(4-chloro-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** ID, 21 × 250 mm, 5 µm, 30% MeOH/CO_2, 80 g/min, 102 bar
102-5-2	 2nd-eluting isomer	6-Chloro-1-(4-chloro-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** ID, 21 × 250 mm, 5 µm, 30% MeOH/CO_2, 80 g/min, 102 bar

102-6-1	 1st-eluting isomer	6-Chloro-1-(4-chloro-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** IF, 21 × 250 mm, 5 μm, 35% MeOH/CO_2, 80 mL/min, 90 bar
102-6-2	 2nd-eluting isomer	6-Chloro-1-(4-chloro-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** IF, 21 × 250 mm, 5 μm, 35% MeOH/CO_2, 80 mL/min, 90 bar
102-7-1	 1st-eluting isomer	6-Chloro-1-(4-dimethylphosphoryl-2-isopropyl-6-methyl-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC:** Chiralcel ID (5 μm, 21x250 mm) F=70ml, 30% Ethanol, 70% carbon dioxide, bpr = 102 bars
102-7-2	 2nd-eluting isomer	6-Chloro-1-(4-dimethylphosphoryl-2-isopropyl-6-methyl-phenyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC:** Chiralcel ID (5 μm, 21 x 250 mm) F=70ml, 30% Ethanol, 70% carbon dioxide, bpr = 102 bars

103-1-1	 1st Eluting Isomer	6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** Welko (RR) (5 μm, 21 x 250 mm), 80 ml/min, 50% MeOH (0.2% triethylamine), 50% carbon dioxide , bpr= 90 bars
103-1-2	 2nd Eluting Isomer	6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** Welko (RR) (5 μm, 21 x 250 mm), 80 ml/min, 50% MeOH (0.2% triethylamine), 50% carbon dioxide , bpr= 90 bars
103-6-1	 1st Eluting Peak	6-Chloro-1-[4-(diethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-isopropylphenyl)pyrido[2,3-d]pyrimidin-2-one	**SFC** OD-H (21 x 250 mm, 5 μm), 80 ml/min, 15% MeOH, 85% carbon dioxide, bpr= 90 bars
103-6-2	 2nd Eluting Peak	6-Chloro-1-[4-(diethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** OD-H (21 x 250 mm, 5 μm), 80 ml/min, 15% MeOH, 85% carbon dioxide, bpr= 90 bars

103-7-1	 1st Eluting Peak	6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** (R,R) Whelk-01 (21 x 250 mm, 5 μm), 80 ml/min, 35% MeOH (0.2% TEA), 65% carbon dioxide, bpr= 102 bars
103-7-2	 2nd Eluting Peak	6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** (R,R) Whelk-01 (21 x 250 mm, 5 μm), 80 ml/min, 35% MeOH (0.2% TEA), 65% carbon dioxide, bpr= 102 bars
103-8-1	 1st Eluting Peak	4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-(diethylamino)-2-isopropylpyridin-3-yl)-7-(o-tolyl)pyrido[2,3-d]pyrimidin-2(1H)-one	**SFC** OD (21 x 150 mm, 5 μm), 80 ml/min, 20% MeOH (0.2% TEA), 80% carbon dioxide, bpr= 102 bars
103-8-2	 2nd Eluting Peak	4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4-(diethylamino)-2-isopropylpyridin-3-yl)-7-(o-tolyl)pyrido[2,3-d]pyrimidin-2(1H)-one	**SFC** OD (21 x 150 mm, 5 μm), 80 ml/min, 20% MeOH (0.2% TEA), 80% carbon dioxide, bpr= 102 bars

106-6-1	 1st-eluting isomer	6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pteridin-2-one	**SFC:** Column: ID (250 X 21 mm, 5μm) Mobile Phase: 70:30 (A:B) A: Liquid CO2 B: MeOH Flow Rate: 80 g/min Column/Oven temp.: ambient temperature BPR = 187 bar
106-6-2	 2nd -eluting isomer	6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pteridin-2-one	**SFC:** Column: ID (250 X 21 mm, 5μm) Mobile Phase: 70:30 (A:B) A: Liquid CO2 B: MeOH Flow Rate: 80 g/min Column/Oven temp.: ambient temperature BPR = 187 bar
106-7-1	 1st-eluting isomer	6-Chloro-4-[(2*R*,6*S*)-2,6-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pteridin-2-one	**SFC:** Chiralcel AS-H (5um, 21x250 mm) F=80ml, 15% methanol, 85% carbon dioxide, back pressure = 90 bars
106-7-2	 2nd -eluting isomer	6-Chloro-4-[(2*R*,6*S*)-2,6-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pteridin-2-one	**SFC:** Chiralcel AS-H (5um, 21x250 mm) F=80ml, 15% methanol, 85% carbon dioxide, back pressure = 90 bars
108-1-1		6-Chloro-1-(4-cyclopropyl-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** IC, 21 × 250 mm, 5 μm, 45% MeOH/CO_2, 70 g/min, 102 bar

	1st-eluting isomer		
108-1-2	 2nd-eluting isomer	6-Chloro-1-(4-cyclopropyl-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** IC, 21 × 250 mm, 5 μm, 45% MeOH/CO_2, 70 g/min, 102 bar
108-2-1	 1st-eluting isomer	6-Chloro-1-(4-cyclopropyl-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** IA, 21 × 250 mm, 5 μm, 20% EtOH/CO_2, 80 mL/min, 90 bar
108-2-2	 2nd-eluting isomer	6-Chloro-1-(4-cyclopropyl-6-isopropyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** IA, 21 × 250 mm, 5 μm, 20% EtOH/CO_2, 80 mL/min, 90 bar
109-1-1	 1st-eluting isomer	6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(4-ethyl-6-isopropyl-pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** IC, 21 × 250 mm, 5 μm, 40% MeOH/CO_2, 80 g/min, 102 bar

109-1-2	 2nd-eluting isomer	6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(4-ethyl-6-isopropyl-pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** IC, 21 × 250 mm, 5 μm, 40% MeOH/CO_2, 80 g/min, 102 bar
109-2-1	 1st-eluting isomer	6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(4-ethyl-6-isopropyl-pyrimidin-5-yl)-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** IF, 30 × 150 mm, 5 μm, 40% isopropanol/CO_2, 100 g/min, 102 bar
109-2-2	 2nd-eluting isomer	6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(4-ethyl-6-isopropyl-pyrimidin-5-yl)-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC** IF, 30 × 150 mm, 5 μm, 40% isopropanol/CO_2, 100 g/min, 102 bar
122-2-1	 1st-eluting isomer	N-[2-[6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]propanamide	**SFC:** DIOL (3um, 2.1x100mm) Mobile Phase: A – CO2, B Methanol with 0.2% triethylamine, Isocratic B: 15%, Flow Rate: 0.8 ml/min, BPR=120bar

122-2-2	 2nd-eluting isomer	*N*-[2-[6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]propanamide	**SFC:** DIOL (3um, 2.1x100mm) Mobile Phase: A – CO2, B Methanol with 0.2% triethylamine, Isocratic B: 15%, Flow Rate: 0.8 ml/min, BPR=120bar
122-3-1	 1st-eluting isomer	*N*-[2-[6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]cyclopropanecarboxamide	**SFC:** WElko-1 RR (5um, 21x150 mm), F=80ml, 35% MeOH with 0.2% triethylamine, 65% carbon dioxide, 120 BAR
122-3-2	 2nd-eluting isomer	*N*-[2-[6-Chloro-1-[4-(dimethylamino)-2-isopropyl-3-pyridyl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-2-oxo-pyrido[2,3-*d*]pyrimidin-7-yl]-3-fluoro-phenyl]cyclopropanecarboxamide	**SFC:** WElko-1 RR (5um, 21x150 mm), F=80ml, 35% MeOH with 0.2% triethylamine, 65% carbon dioxide, 120 BAR
124-14-1	 1st-eluting isomer	4-[*cis*-2,3-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC:** ID (250 x 20 mm, 5 µm), 75:25 CO_2 and iPrOH, 85 g/min, 103 bar

124-14-2	 2nd-eluting isomer	4-[*cis*-2,3-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC:** ID (250 x 20 mm, 5 µm), 75:25 CO_2 and iPrOH, 85 g/min, 103 bar
124-16-1	 1st-eluting isomer	4-[*trans*-2,3-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC:** IF (250 x 20 mm, 5 µm), 60:40 CO_2 and iPrOH, 75 g/min, 103 bar; then **SFC:** ID (250 x 20 mm, 5 µm), 75:25 CO_2 and iPrOH, 85 g/min, 103 bar
124-16-2	 2nd-eluting isomer	4-[*trans*-2,3-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-7-(2-fluoro-6-hydroxy-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one	**SFC:** IF (250 x 20 mm, 5 µm), 60:40 CO_2 and iPrOH, 75 g/min, 103 bar
144-2-1	 1st -eluting isomer	(*M*)-2-[4-[6,7-Dichloro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile	**SFC** OD x 2 (250 x 21 mm, 5 µm and 150 x 21, 5 µm) with a mobile phase of 60% Liquid CO2 and 40% MeOH, flowrate of 50 mL/min, 102 bar
144-2-2	 2nd -eluting isomer	(*M*)-2-[4-[6,7-Dichloro-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile	**SFC** OD x 2 (250 x 21 mm, 5 µm and 150 x 21, 5 µm) with a mobile phase of 60% Liquid CO2 and 40% MeOH, flowrate of 50 mL/min, 102 bar

145-1-1	 1st-eluting isomer	2-[4-[6-Chloro-1-(2,2-dimethylpropyl)-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile	**SFC** AS (250 x 21 mm, 5 µm) with a mobile phase of 80% Liquid CO2 and 20% MeOH, flowrate of 80 mL/min, 102 bar
145-1-2	 2nd-eluting isomer	2-[4-[6-Chloro-1-(2,2-dimethylpropyl)-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile	**SFC** AS (250 x 21 mm, 5 µm) with a mobile phase of 80% Liquid CO2 and 20% MeOH, flowrate of 80 mL/min, 102 bar
145-2-1	 1st-eluting isomer	(*M*)-2-[4-[6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile	**SFC** OJ (250 x 21 mm, 5 µm) with a mobile phase of 90% Liquid CO2 and 10% MeOH, flowrate of 90 mL/min, 102 bar
145-2-2	 2nd-eluting isomer	(*M*)-2-[4-[6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-4-yl]-1-prop-2-enoyl-piperazin-2-yl]acetonitrile	**SFC** OJ (250 x 21 mm, 5 µm) with a mobile phase of 90% Liquid CO2 and 10% MeOH, flowrate of 90 mL/min, 102 bar
146-1-1	 1st-eluting isomer	19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one	**SFC** Welko SS (5 µm, 21x250 mm) F=80ml, 35% Methanol, 65% carbon dioxide , Back pressure= 90 bar

146-1-2	 2nd-eluting isomer	(9*Z*)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one	**SFC** Welko SS (5 µm, 21x250 mm) F=80ml, 35% Methanol, 65% carbon dioxide , Back pressure= 90 bar
146-2-1	 1st -eluting isomer	19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one	**SFC** Welko SS (5 µm, 21x250 mm) F=80ml, 35% Methanol, 65% carbon dioxide , Back pressure= 90 bar
146-2-2	 2nd-eluting isomer	(9*Z*)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-3-(2-propanyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one	**SFC** Welko SS (5 µm, 21x250 mm) F=80ml, 35% Methanol, 65% carbon dioxide , Back pressure= 90 bar
146-3-1	 1st -eluting isomer	19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one	**SFC** (R,R) Whelk-01 (250 X 21 mm, 5 µm) with a mobile phase of 60% Liquid CO2 and 40% MeOH using a flowrate of 80 mL/min, 102 bar
146-3-2		19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-	**SFC** (R,R) Whelk-01 (250 X 21 mm, 5 µm) with a mobile phase of 60% Liquid CO2 and 40% MeOH using a flowrate of 80 mL/min, 102 bar

	2nd-eluting isomer	2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one	
146-3-3	 3rd-eluting isomer	(9*Z*)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one	**SFC** (R,R) Whelk-01 (250 X 21 mm, 5 µm) with a mobile phase of 60% Liquid CO2 and 40% MeOH using a flowrate of 80 mL/min, 102 bar
146-3-4	 4th-eluting isomer	(9*Z*)-19-chloro-22-((2*S*,5*R*)-2,5-dimethyl-4-(2-propenoyl)-1-piperazinyl)-1,23,26-triazapentacyclo[16.6.2.0~2,7~.0~12,17~.0~21,25~]hexacosa-2,4,6,9,12,14,16,18,20,22,25-undecaen-24-one	**SFC** (R,R) Whelk-01 (250 X 21 mm, 5 µm) with a mobile phase of 60% Liquid CO2 and 40% MeOH using a flowrate of 80 mL/min, 102 bar

Table 148: Analytical Data for Method Table Examples

Ex. #	LRMS: *m/z* (ESI, +ve ion): (M+H)+	NMR
55-50	594.4	^{1}H NMR (DMSO-d_6) δ 8.45-8.59 (m, 2H), 7.48-7.59 (m, 2H), 7.19-7.39 (m, 3H), 6.75-6.93 (m, 1H), 6.11-6.25 (m, 1H), 5.72-5.80 (m, 1H), 4.83-5.00 (m, 1H), 4.72-4.83 (m, 1H), 4.44-4.53 (m, 1H), 4.11-4.29 (m, 1H), 3.78-3.98 (m, 2H), 2.80-2.94 (m, 1H), 1.32-1.38 (m, 2H), 1.22-1.29 (m, 2H), 1.16-1.22 (m, 2H), 1.09-1.14 (m, 3H), 0.96-1.04 (m, 3H)
55-51	604.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.49 (s, 1 H), 8.16 (dd, *J*=9.4, 5.7 Hz, 1 H), 7.48 - 7.59 (m, 1 H), 7.19 - 7.38 (m, 3 H), 6.74 - 6.90 (m, 1 H), 6.68 (dd, *J*=5.7, 3.2 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.71 - 5.81 (m, 1 H), 4.68 - 4.95 (m, 2 H), 4.46 (br dd, *J*=8.8, 3.6 Hz, 1 H), 3.94 - 4.30 (m, 2 H), 3.79 - 3.91 (m, 1 H), 2.68 (d, *J*=4.1 Hz, 6 H), 1.31 - 1.38 (m, 3 H), 1.10 - 1.22 (m, 3 H), 1.09 - 1.10 (m, 1 H), 1.03 (dd, *J*=6.6, 3.1 Hz, 3 H), 0.81 - 0.92 (m, 3 H)
55-51-1	604.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.44 (d, *J*=5.4 Hz, 1 H), 8.15 (d, *J*=5.8 Hz, 1 H), 7.55 (br d, *J*=8.3 Hz, 1 H), 7.29 - 7.37 (m, 2 H), 7.22 - 7.28 (m, 1 H), 6.81 (br dd, *J*=16.4, 10.4 Hz, 1 H), 6.67 (d, *J*=5.8 Hz, 1 H), 6.19 (br d, *J*=16.8

		Hz, 1 H), 5.68 - 5.84 (m, 1 H), 4.68 - 4.94 (m, 2 H), 4.42 - 4.54 (m, 1 H), 4.09 - 4.33 (m, 1 H), 3.70 - 3.97 (m, 2 H), 2.68 (s, 7 H), 1.32 - 1.38 (m, 2 H), 1.20 - 1.26 (m, 2 H), 1.15 (br d, *J*=6.8 Hz, 2 H), 1.04 (d, *J*=6.6 Hz, 3 H), 0.84 (d, *J*=6.6 Hz, 3 H)
55-51-2	604.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.49 (s, 1 H), 8.17 (d, *J*=5.8 Hz, 1 H), 7.49 - 7.62 (m, 1 H), 7.17 - 7.38 (m, 3 H), 6.73 - 6.91 (m, 1 H), 6.68 (d, *J*=5.8 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.75 (br dd, *J*=10.3, 2.6 Hz, 1 H), 4.90 (br d, *J*=4.6 Hz, 1 H), 4.73 (dd, *J*=4.0, 1.6 Hz, 1 H), 4.44 (br d, *J*=3.5 Hz, 1 H), 3.92 - 4.18 (m, 3 H), 3.86 (br d, *J*=3.1 Hz, 1 H), 2.67 (s, 6 H), 1.30 (dd, *J*=12.4, 6.6 Hz, 3 H), 1.20 (br d, *J*=6.6 Hz, 1 H), 1.11 (br d, *J*=6.6 Hz, 2 H), 1.03 (d, *J*=6.6 Hz, 3 H), 0.87 (d, *J*=6.4 Hz, 3 H)
55-52	591.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.63 (d, *J*=5.4 Hz, 1 H), 8.18 (d, *J*=5.6 Hz, 1 H), 7.56 - 7.67 (m, 2 H), 7.44 - 7.52 (m, 1 H), 7.26 - 7.35 (m, 2 H), 6.83 (td, *J*=15.8, 10.4 Hz, 1 H), 6.13 - 6.22 (m, 1 H), 5.71 - 5.79 (m, 1 H), 4.75 - 4.94 (m, 1 H), 4.05 - 4.19 (m, 1 H), 3.96 - 4.05 (m, 1 H), 3.74 - 3.92 (m, 2 H), 3.47 (s, 3 H), 2.77 - 2.94 (m, 1 H), 1.26 - 1.32 (m, 3 H), 1.23 (br d, *J*=6.6 Hz, 2 H), 1.16 (d, *J*=6.6 Hz, 5 H), 1.09 (br d, *J*=6.4 Hz, 3 H)
55-53	664.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.56 (dd, *J*=8.4, 4.7 Hz, 1 H), 8.44 (d, *J*=3.9 Hz, 1 H), 7.97 - 8.11 (m, 2 H), 7.81 - 7.88 (m, 1 H), 7.70 - 7.81 (m, 1 H), 7.39 (d, *J*=7.9 Hz, 1 H), 6.55 (dd, *J*=9.7, 5.8 Hz, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.70 - 5.82 (m, 1 H), 4.72 - 4.98 (m, 2 H), 4.47 - 4.53 (m, 1 H), 4.08 - 4.26 (m, 1 H), 3.84 - 3.93 (m, 2 H), 2.87 (s, 3 H), 2.74 (s, 6 H), 2.66 - 2.70 (m, 1 H), 1.25 - 1.37 (m, 5 H), 1.23 - 1.25 (m, 1 H), 0.94 (d, *J*=6.8 Hz, 3 H), 0.74 - 0.80 (m, 3 H)
55-54	618.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.43 - 8.50 (m, 1 H), 8.18 (d, *J*=5.6 Hz, 1 H), 7.03 - 7.17 (m, 1 H), 6.77 - 6.91 (m, 1 H), 6.69 (br d, *J*=3.9 Hz, 1 H), 6.45 - 6.55 (m, 1 H), 6.38 (br t, *J*=9.1 Hz, 1 H), 6.20 (dd, *J*=16.5, 2.0 Hz, 1 H), 5.73 - 5.80 (m, 1 H), 5.13 - 5.19 (m, 1 H), 4.71 - 4.92 (m, 2 H), 4.43 - 4.51 (m, 1 H), 4.03 - 4.19 (m, 1 H), 3.82 - 3.99 (m, 2 H), 2.69 (br d, *J*=7.0 Hz, 6 H), 2.62 (br s, 1 H), 1.27 - 1.36 (m, 3 H), 1.22 (br d, *J*=5.0 Hz, 2 H), 1.14 (br d, *J*=6.6 Hz, 2 H), 1.02 - 1.06 (m, 3 H), 0.80 (br dd, *J*=10.7, 6.9 Hz, 3 H)
55-55	594.2	^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 7.97 - 8.09 (m, 1 H), 7.64 (br t, *J*=7.7 Hz, 2 H), 7.46 - 7.55 (m, 1 H), 7.39 (br d, *J*=7.7 Hz, 2 H), 7.22 (s, 1 H), 7.03 - 7.17 (m, 2 H), 6.49 - 6.70 (m, 1 H), 6.32 - 6.44 (m, 1 H), 5.71 - 5.84 (m, 1 H), 5.05 (br d, *J*=1.7 Hz, 1 H), 4.29 - 4.50 (m, 1 H), 4.05 - 4.19 (m, 1 H), 3.61 - 3.95 (m, 2 H), 3.41 - 3.48 (m, 1 H), 1.58 - 1.72 (m, 6 H), 1.40 - 1.50 (m, 4 H), 1.28 - 1.39 (m, 2 H)

55-56	618.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.45 (s, 1 H), 8.18 (d, *J*=5.6 Hz, 1 H), 7.03 - 7.19 (m, 1 H), 6.83 (br dd, *J*=16.4, 10.6 Hz, 1 H), 6.68 (d, *J*=5.6 Hz, 1 H), 6.44 - 6.53 (m, 1 H), 6.28 - 6.41 (m, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.72 - 5.81 (m, 1 H), 5.16 (br s, 2 H), 4.62 - 4.96 (m, 2 H), 4.40 - 4.52 (m, 1 H), 3.84 - 4.18 (m, 3 H), 2.58 - 2.74 (m, 7 H), 1.32 (br dd, *J*=10.3, 6.5 Hz, 3 H), 1.08 - 1.24 (m, 3 H), 1.03 (br d, *J*=6.4 Hz, 3 H), 0.81 (br d, *J*=6.8 Hz, 3 H)
55-57	618.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.36 - 8.49 (m, 1 H), 8.18 (d, *J*=5.8 Hz, 1 H), 7.03 - 7.16 (m, 1 H), 6.82 (br dd, *J*=16.9, 9.8 Hz, 1 H), 6.68 (br d, *J*=5.8 Hz, 1 H), 6.53 (d, *J*=8.1 Hz, 1 H), 6.38 (br t, *J*=9.3 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 5.29 (br s, 1 H), 5.13 (br s, 1 H), 4.72 - 4.95 (m, 2 H), 4.44 - 4.54 (m, 1 H), 4.07 - 4.22 (m, 1 H), 3.82 - 3.98 (m, 2 H), 2.60 - 2.72 (m, 7 H), 1.33 (br s, 3 H), 1.12 - 1.25 (m, 3 H), 0.97 - 1.07 (m, 3 H), 0.78 (d, *J*=6.8 Hz, 3 H)
55-58	588.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.44 - 8.53 (m, 2 H), 7.52 (br d, *J*=7.0 Hz, 1 H), 7.16 - 7.35 (m, 4 H), 6.75 - 6.93 (m, 1 H), 6.20 (dd, *J*=16.7, 1.6 Hz, 1 H), 5.76 (s, 1 H), 4.72 - 4.97 (m, 2 H), 4.48 (br d, *J*=2.5 Hz, 1 H), 4.08 - 4.27 (m, 1 H), 3.80 - 3.98 (m, 2 H), 2.19 - 2.32 (m, 2 H), 1.31 - 1.39 (m, 3 H), 1.25 (br t, *J*=5.7 Hz, 1 H), 1.21 - 1.23 (m, 1 H), 1.18 (br t, *J*=7.2 Hz, 2 H), 1.07 (dd, *J*=6.5, 4.0 Hz, 3 H), 0.91 - 1.04 (m, 6 H)
55-58-1	589.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.49 (s, 1 H), 8.45 (d, *J*=5.0 Hz, 1 H), 7.48 - 7.59 (m, 1 H), 7.25 - 7.35 (m, 2 H), 7.22 (br d, *J*=5.0 Hz, 2 H), 6.76 - 6.92 (m, 1 H), 6.15 - 6.25 (m, 1 H), 5.76 - 5.80 (m, 1 H), 4.74 - 4.95 (m, 2 H), 4.43 - 4.53 (m, 1 H), 4.10 - 4.21 (m, 1 H), 3.84 - 3.99 (m, 2 H), 2.21 - 2.30 (m, 2 H), 1.34 (br t, *J*=7.2 Hz, 3 H), 1.22 - 1.27 (m, 2 H), 1.17 (br d, *J*=8.9 Hz, 2 H), 1.05 - 1.10 (m, 4 H), 0.94 - 1.01 (m, 5 H)
55-58-2	589.3	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.42 - 8.50 (m, 2 H), 7.48 - 7.58 (m, 1 H), 7.17 - 7.36 (m, 4 H), 6.75 - 6.92 (m, 1 H), 6.20 (dd, *J*=16.8, 1.9 Hz, 1 H), 5.75 - 5.80 (m, 1 H), 4.93 (s, 2 H), 4.44 - 4.53 (m, 1 H), 4.13 - 4.28 (m, 1 H), 3.78 - 3.97 (m, 2 H), 2.22 - 2.33 (m, 2 H), 1.33 - 1.38 (m, 3 H), 1.26 (br d, *J*=6.8 Hz, 2 H), 1.17 - 1.21 (m, 2 H), 1.04 - 1.09 (m, 4 H), 0.98 - 1.04 (m, 2 H), 0.91 - 0.96 (m, 3 H)
55-59	456.2	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 8.33 (d, *J*=2.7 Hz, 1 H), 7.56 - 7.68 (m, 2 H), 7.34 - 7.46 (m, 2 H), 6.81 (td, *J*=16.8, 10.6 Hz, 1 H), 6.17 (dd, *J*=16.7, 1.6 Hz, 1 H), 5.74 (br dd, *J*=10.6, 1.9 Hz, 1 H), 4.37 - 4.79 (m, 2 H), 3.69 - 4.13 (m, 4 H), 3.48 (s, 3 H), 1.27 (br t, *J*=5.7 Hz, 3 H), 1.10 - 1.23 (m, 3 H)
55-60	484.2	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 8.30 (s, 1 H), 7.56 - 7.67 (m, 2 H), 7.37 - 7.47 (m, 2 H), 6.81 (td, *J*=17.5, 10.6 Hz, 1 H), 6.17 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.69 - 5.75 (m, 1 H), 5.35 (quin, *J*=6.9 Hz, 1 H), 4.37 - 4.80 (m, 2 H), 3.38 -

		4.19 (m, 4 H), 1.46 (dd, *J*=6.8, 3.9 Hz, 6 H), 1.25 (t, *J*=6.4 Hz, 3 H), 1.10 - 1.22 (m, 3 H)
69-10	600.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.45 (1 H, dd, *J*=6.3, 4.9 Hz), 7.90 - 8.13 (1 H, m), 7.24 (1 H, dd, *J*=17.9, 5.1 Hz), 6.69 - 6.94 (1 H, m), 6.17 (1 H, dd, *J*=16.7, 2.0 Hz), 5.68 - 5.79 (1 H, m), 5.02 - 5.62 (1 H, m), 4.36 - 4.89 (2 H, m), 4.03 - 4.23 (1 H, m), 3.42 - 4.00 (6 H, m), 2.70 - 2.96 (2 H, m), 2.54 (1 H, m), 2.04 (2 H, m), 1.86 (3 H, s), 1.10 - 1.35 (6 H, m), 0.84 - 1.09 (6 H, m)
69-11	564.2	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 8.46 (d, *J*=4.8 Hz, 1 H), 8.02 (s, 1 H), 7.27 (d, *J*=5.0 Hz, 1 H), 6.82 (td, *J*=17.5, 10.4 Hz, 1 H), 6.17 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.67 - 5.83 (m, 1 H), 4.36 - 4.87 (m, 2 H), 3.36 - 4.17 (m, 4 H), 3.27 (br t, *J*=5.3 Hz, 4 H), 2.58 - 2.72 (m, 1 H), 1.94 (s, 3 H), 1.44 - 1.54 (m, 2 H), 1.30 - 1.41 (m, 4 H), 1.11 - 1.29 (m, 6 H), 1.08 (d, *J*=6.8 Hz, 3 H), 0.98 (d, *J*=6.6 Hz, 3 H)
69-12	578.3	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 8.36 - 8.52 (m, 1 H), 7.90 - 8.08 (m, 1 H), 7.23 (d, *J*=4.8 Hz, 1 H), 6.71 - 6.96 (m, 1 H), 6.17 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.60 - 5.86 (m, 1 H), 4.31 - 4.86 (m, 2 H), 3.50 - 4.25 (m, 6 H), 2.84 - 2.98 (m, 1 H), 2.65 - 2.77 (m, 1 H), 2.52 - 2.62 (m, 1 H), 1.83 - 2.01 (m, 3 H), 1.34 - 1.64 (m, 5 H), 1.16 - 1.33 (m, 6 H), 1.07 (br d, *J*=6.6 Hz, 3 H), 0.89 - 1.04 (m, 6 H)
72-20	550.6	^{1}H NMR (DMSO-d$_6$) δ 10.04-10.46 (m, 1H), 8.35-8.44 (m, 2H), 7.21-7.33 (m, 1H), 7.13-7.21 (m, 1H), 6.78-6.93 (m, 1H), 6.64-6.76 (m, 2H), 6.14-6.26 (m, 1H), 5.77-5.82 (m, 1H), 3.93-4.07 (m, 4H), 2.67-2.78 (m, 1H), 1.86-1.93 (m, 3H), 1.04-1.12 (m, 3H), 0.89-0.97 (m, 3H)
72-21	550.6	^{1}H NMR (DMSO-d$_6$) δ 9.96-10.32 (m, 1H), 8.23-8.48 (m, 2H), 7.24-7.33 (m, 1H), 7.15-7.20 (m, 1H), 6.79-6.90 (m, 1H), 6.66-6.76 (m, 2H), 6.16-6.23 (m, 1H), 5.74-5.79 (m, 1H), 3.70-3.93 (m, 4H), 2.65-2.78 (m, 1H), 1.85-1.94 (m, 3H), 1.07-1.11 (m, 3H), 0.86-0.98 (m, 3H)
72-22	555.4	^{1}H NMR (DMSO-d$_6$) δ 9.98-10.30 (m, 1H), 8.35-8.45 (m, 2H), 7.22-7.34 (m, 1H), 7.14-7.21 (m, 1H), 6.78-6.90 (m, 1H), 6.63-6.76 (m, 2H), 6.15-6.26 (m, 1H), 5.71-5.81 (m, 1H), 2.65-2.80 (m, 1H), 1.85-1.94 (m, 3H), 1.05-1.12 (m, 3H), 0.89-0.97 (m, 3H)
72-23	617.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.37 (d, *J*=4.8 Hz, 1 H), 8.24 - 8.31 (m, 1 H), 7.76 (br dd, *J*=11.9, 7.4 Hz, 1 H), 7.59 (br dd, *J*=9.2, 7.6 Hz, 2 H), 7.30 (br d, *J*=2.5 Hz, 1 H), 7.18 (d, *J*=4.8 Hz, 1 H), 6.71 - 6.96 (m, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.66 - 5.82 (m, 1 H), 4.71 - 4.97 (m, 2 H), 4.50 (br s, 1 H), 3.87 (br s, 4 H), 1.97 (s, 3 H), 1.57 (br d, *J*=10.6 Hz, 3 H), 1.26 - 1.35 (m, 3 H), 1.09 - 1.26 (m, 6 H), 1.05 (br d, *J*=6.6 Hz, 3 H), 0.81 - 0.91 (m, 3 H)
79-10	573.2	^{1}H NMR (500 MHz, DMSO-*d$_6$*) δ 8.50 (s, 1 H) 8.39 (d, *J* = 4.8 Hz, 1 H) 7.46 - 7.54 (m, 1 H) 7.18 - 7.33 (m, 4 H) 6.83 (dd, *J* = 16.7, 10.4 Hz, 1 H) 6.27 (dd, *J* = 16.7, 2.3

		Hz, 1 H) 5.77 - 5.82 (m, 1 H) 4.72 (br s, 2 H) 4.51 - 4.61 (m, 2 H) 4.10 - 4.17 (m, 2 H) 3.55 - 3.67 (m, 2 H) 2.69 (tt, *J* = 12.9, 6.6 Hz, 1 H) 1.95 (d, *J* = 5.2 Hz, 3 H) 1.75 - 1.84 (m, 2 H) 1.05 (d, *J* = 6.7 Hz, 3 H) 0.96 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ-112.45 - -112.58 (m, 1 F).
79-11	573.2	^{1}H NMR (400 MHz, $CDCl_3$) δ 8.47 (d, *J* = 5.0 Hz, 1 H) 8.23 - 8.37 (m, 1 H) 7.35 - 7.45 (m, 1 H) 7.03 - 7.18 (m, 4 H) 6.38 - 6.61 (m, 2 H) 5.74 - 5.85 (m, 1 H) 3.74 - 5.56 (m, 6 H) 2.66 - 2.81 (m, 1 H) 2.13 - 2.47 (m, 2 H) 1.98 - 2.10 (m, 5 H) 1.19 - 1.26 (m, 3 H) 1.00 - 1.07 (m, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -112.71 - -112.43 (m, 1 F).
79-12	586.8	^{1}H NMR (400 MHz, $CDCl_3$) δ 8.49 (d, *J* = 5.0 Hz, 1 H) 8.03 (s, 1 H) 7.37 - 7.45 (m, 1 H) 7.04 - 7.21 (m, 4 H) 6.57 - 6.69 (m, 1 H) 6.39 (dd, *J* = 16.8, 1.7 Hz, 1 H) 5.80 (dd, *J* = 10.6, 1.7 Hz, 1 H) 4.54 - 5.07 (m, 3 H) 3.57 - 4.40 (m, 2 H) 3.28 (br d, *J* = 12.6 Hz, 1 H) 2.63 - 2.80 (m, 1 H) 2.08 - 2.37 (m, 4 H) 2.01 - 2.06 (m, 3 H) 1.72 - 1.99 (m, 2 H) 1.23 (d, *J* = 6.6 Hz, 3 H) 1.06 (d, *J* = 6.8 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.54 - -112.44 (m, 1 F) -112.60 (s, 1 F).
80-10	643.3	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.02-8.08 (m, 1H), 7.22-7.41 (m, 4H), 7.03-7.20 (m, 3H), 6.52-6.70 (m, 1H), 6.39 (br t, J=14.82 Hz, 1H), 5.75-5.83 (m, 1H), 5.00-5.26 (m, 1H), 4.31-4.49 (m, 1H), 3.85-4.06 (m, 2H), 3.35-3.81 (m, 2H), 3.07-3.17 (m, 1H), 2.50-2.73 (m, 1H), 2.15 (br s, 4H), 1.34-1.50 (m, 9H), 1.25 (br t, J=7.15 Hz, 2H), 1.18 (d, J=6.63 Hz, 3H), 1.01-1.08 (m, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-*d*) δ -112.37 (m, 1F).
80-10-1	643.4	^{1}H NMR (400 MHz, DICHLOROMETHANE-d_2) δ 8.03 (s, 1H), 7.17-7.37 (m, 4H), 7.00-7.14 (m, 3H), 6.46-6.63 (m, 1H), 6.22 (br t, *J*=15.13 Hz, 1H), 5.67 (br t, *J*=8.71 Hz, 1H), 5.23-5.26 (m, 1H), 4.64-5.01 (m, 2H), 4.14-4.35 (m, 2H), 3.91-4.07 (m, 1H), 3.62-3.83 (m, 2H), 3.29-3.48 (m, 1H), 3.04 (br d, *J*=13.48 Hz, 1H), 2.52-2.66 (m, 1H), 1.99-2.00 (m, 1H), 1.97-2.11 (m, 3H), 1.36 (br dd, *J*=6.53, 11.51 Hz, 8H), 1.05 (d, *J*=6.84 Hz, 3H), 0.97 (d, *J*=6.84 Hz, 3H). ^{19}F NMR (376 MHz, DICHLOROMETHANE-d_2) δ -113.32 (s, 1F).
80-10-2	643.4	^{1}H NMR (400 MHz, DICHLOROMETHANE-d_2) δ 8.02 (s, 1H), 7.16-7.37 (m, 4H), 7.00-7.13 (m, 3H), 6.46-6.63 (m, 1H), 6.22 (br t, *J*=14.82 Hz, 1H), 5.67 (br t, *J*=8.40 Hz, 1H), 5.24 (br s, 1H), 5.00 (br s, 1H), 4.87 (br s, 1H), 4.28 (br d, *J*=11.61 Hz, 1H), 3.78-4.04 (m, 3H), 3.44 (d, *J*=13.27 Hz, 2H), 3.03 (br d, *J*=13.27 Hz, 1H), 2.44-2.60 (m, 1H), 2.06 (br s, 4H), 1.32 (br s, 8H), 1.05 (d, *J*=6.84 Hz, 3H), 0.98 (br d, *J*=6.84 Hz, 3H). ^{19}F NMR (376 MHz, DICHLOROMETHANE-d_2) δ -113.35 (s, 1F).
80-11	665.3	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.01 (br d, *J*=4.98 Hz, 1H), 7.27 (br s, 3H), 7.15-7.22 (m, 2H), 6.99-

		7.11 (m, 3H), 6.44-6.62 (m, 1H), 6.27-6.37 (m, 1H), 5.68-5.76 (m, 1H), 4.77-5.23 (m, 2H), 4.25-4.44 (m, 1H), 3.73-3.97 (m, 2H), 3.40-3.70 (m, 2H), 3.04-3.38 (m, 4H), 2.34-2.64 (m, 1H), 1.31-1.37 (m, 3H), 1.24 (br dd, *J*=3.63, 6.12 Hz, 3H), 1.09 (d, *J*=6.84 Hz, 3H), 0.94 (br dd, *J*=6.84, 11.61 Hz, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-*d*) δ -99.38 (m, 2F), -112.89 (s, 1F).
80-12	629.3	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.06-8.13 (m, 1H), 7.22-7.42 (m, 5H), 7.05-7.19 (m, 3H), 6.53-6.71 (m, 1H), 6.35-6.46 (m, 1H), 5.76-5.85 (m, 1H), 4.21-5.31 (m, 3H), 3.47-4.08 (m, 1H), 3.25-3.44 (m, 2H), 2.98-3.08 (m, 3H), 2.48-2.58 (m, 1H), 2.37 (br s, 2H), 1.86-1.98 (m, 2H), 1.31-1.47 (m, 6H), 1.19 (d, *J*=6.84 Hz, 3H), 0.98-1.05 (m, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.80 (s, 1F).
80-13	603.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.06 (s, 1H), 7.32-7.42 (m, 4H), 7.12-7.17 (m, 2H), 7.05-7.11 (m, 1H), 6.54-6.72 (m, 1H), 6.37-6.46 (m, 1H), 5.81 (dd, *J*=1.76, 10.47 Hz, 1H), 4.80-5.22 (m, 1H), 4.39-4.79 (m, 1H), 3.97-4.37 (m, 1H), 3.51-3.95 (m, 3H), 3.26-3.50 (m, 1H), 2.97-3.24 (m, 2H), 2.59 (br s, 1H), 2.00-2.07 (m, 6H), 1.48 (br d, *J*=7.67 Hz, 3H), 1.19 (d, *J*=6.84 Hz, 3H), 1.02 (d, *J*=6.84 Hz, 3H).
80-14	617.2	^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 8.04 - 8.11 (1 H, m), 7.32 - 7.45 (4 H, m), 7.05 - 7.18 (3 H, m), 6.50 - 6.73 (1 H, m), 6.35 - 6.46 (1 H, m), 5.81 (1 H, br t, *J*=8.4 Hz), 4.85 - 5.25 (2 H, m), 4.32 - 4.51 (1 H, m), 3.84 - 4.26 (2 H, m), 3.38 - 3.83 (2 H, m), 3.07 - 3.35 (2 H, m), 2.47 - 2.63 (1 H, m), 2.13 (2 H, br d, *J*=8.1 Hz), 2.03 (3 H, s), 1.28 - 1.50 (6 H, m), 1.19 (3 H, d, *J*=6.8 Hz), 1.01 (3 H, br t, *J*=6.4 Hz)
80-14-1	617.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.40 (1 H, d, *J*=4.6 Hz), 7.44 - 7.54 (1 H, m), 7.22 - 7.34 (5 H, m), 7.16 (1 H, td, *J*=7.4, 1.7 Hz), 6.83 (1 H, td, *J*=16.3, 10.5 Hz), 6.19 (1 H, dd, *J*=16.6, 2.3 Hz), 5.71 - 5.80 (1 H, m), 4.41 - 4.93 (2 H, m), 3.39 - 4.26 (4 H, m), 2.91 - 3.10 (2 H, m), 2.56 - 2.62 (1 H, m), 1.85 (6 H, s), 1.34 (3 H, dd, *J*=6.1, 4.3 Hz), 1.17 - 1.29 (3 H, m), 1.06 (3 H, d, *J*=6.8 Hz), 0.96 (3 H, d, *J*=6.8 Hz).
80-14-2	617.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.45 (1 H, d, *J*=1.7 Hz), 7.44 - 7.53 (1 H, m), 7.21 - 7.35 (5 H, m), 7.11 - 7.19 (1 H, m), 6.83 (1 H, dt, *J*=16.7, 11.0 Hz), 6.19 (1 H, dd, *J*=16.7, 2.4 Hz), 5.71 - 5.79 (1 H, m), 4.86 - 4.96 (1 H, m), 4.40 - 4.80 (1 H, m), 3.83 - 4.18 (4 H, m), 2.95 - 3.13 (2 H, m), 2.53 - 2.56 (1 H, m), 1.81 - 1.92 (6 H, m), 1.29 (3 H, br dd, *J*=12.0, 6.6 Hz), 1.14 - 1.24 (3 H, m), 1.06 (3 H, d, *J*=6.8 Hz), 0.96 (3 H, d, *J*=6.8 Hz)
80-16	618.2	^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 8.58 (1 H, d, *J*=5.0 Hz), 8.08 - 8.13 (1 H, m), 7.38 - 7.47 (1 H, m), 7.29 (1 H, d, *J*=4.8 Hz), 7.05 - 7.17 (3 H, m), 6.55 (1 H, br d,

		J=10.6 Hz), 6.41 (1 H, br t, *J*=14.9 Hz), 5.77 - 5.86 (1 H, m), 4.89 - 5.24 (2 H, m), 4.31 - 4.53 (1 H, m), 3.85 - 4.08 (2 H, m), 3.68 - 3.77 (1 H, m), 3.08 - 3.27 (2 H, m), 2.66 - 2.83 (1 H, m), 1.98 - 2.03 (6 H, m), 1.30 - 1.48 (6 H, m), 1.21 - 1.24 (3 H, m), 1.06 - 1.13 (3 H, m)
80-16-1	618.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.44 - 8.52 (2 H, m), 7.45 - 7.55 (1 H, m), 7.23 - 7.34 (3 H, m), 7.17 (1 H, td, *J*=7.4, 1.5 Hz), 6.76 - 6.91 (1 H, m), 6.19 (1 H, dd, *J*=16.6, 2.3 Hz), 5.71 - 5.80 (1 H, m), 4.85 - 4.99 (1 H, m), 4.40 - 4.80 (1 H, m), 3.46 - 4.18 (4 H, m), 3.04 (2 H, s), 2.65 - 2.76 (1 H, m), 1.90 (6 H, s), 1.28 - 1.37 (3 H, m), 1.14 - 1.27 (3 H, m), 1.07 (3 H, d, *J*=6.6 Hz), 0.96 (3 H, d, *J*=6.6 Hz)
80-16-2	618.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.42 - 8.52 (2 H, m), 7.45 - 7.54 (1 H, m), 7.24 - 7.39 (3 H, m), 7.14 - 7.22 (1 H, m), 6.83 (1 H, td, *J*=15.9, 10.5 Hz), 6.19 (1 H, dd, *J*=16.6, 2.3 Hz), 5.70 - 5.80 (1 H, m), 4.83 - 4.95 (1 H, m), 4.77 (1 H, br d, *J*=1.9 Hz), 3.43 - 4.26 (4 H, m), 3.00 - 3.13 (2 H, m), 2.73 (1 H, dt, *J*=13.3, 6.5 Hz), 1.92 (6 H, s), 1.31 - 1.39 (3 H, m), 1.15 - 1.28 (3 H, m), 1.07 (3 H, d, *J*=6.6 Hz), 0.96 (3 H, d, *J*=6.8 Hz)
80-18	619.3	^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 9.11 - 9.17 (1 H, m), 8.12 (1 H, br d, *J*=5.6 Hz), 7.38 - 7.47 (1 H, m), 7.06 - 7.20 (3 H, m), 6.52 - 6.69 (1 H, m), 6.35 - 6.47 (1 H, m), 5.76 - 5.86 (1 H, m), 4.81 - 5.26 (2 H, m), 3.59 - 3.96 (6 H, m), 2.76 (1 H, td, *J*=12.8, 6.6 Hz), 2.13 - 2.33 (6 H, m), 1.42 - 1.51 (3 H, m), 1.28 - 1.41 (3 H, m), 1.24 (3 H, d, *J*=6.6 Hz), 1.08 (3 H, br d, *J*=6.6 Hz)
80-18-1	619.3	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 9.03 (1 H, s), 8.43 (1 H, d, *J*=8.5 Hz), 7.47 - 7.56 (1 H, m), 7.25 - 7.35 (2 H, m), 7.16 - 7.24 (1 H, m), 6.83 (1 H, td, *J*=17.5, 10.6 Hz), 6.19 (1 H, dd, *J*=16.8, 2.3 Hz), 5.72 - 5.80 (1 H, m), 4.45 - 4.96 (2 H, m), 3.42 - 4.37 (4 H, m), 3.39 (1 H, d, *J*=13.7 Hz), 3.15 (1 H, d, *J*=13.7 Hz), 2.82 - 2.95 (1 H, m), 1.83 (6 H, s), 1.37 (3 H, d, *J*=6.6 Hz), 1.19 - 1.29 (3 H, m), 1.10 (3 H, d, *J*=6.6 Hz), 1.01 (3 H, d, *J*=6.6 Hz)
80-18-2	619.3	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 9.03 (1 H, s), 8.51 (1 H, s), 7.46 - 7.56 (1 H, m), 7.25 - 7.35 (2 H, m), 7.15 - 7.23 (1 H, m), 6.84 (1 H, dt, *J*=16.6, 11.0 Hz), 6.19 (1 H, dd, *J*=16.6, 2.3 Hz), 5.71 - 5.82 (1 H, m), 4.95 (1 H, br s), 4.40 - 4.82 (1 H, m), 3.51 - 4.20 (4 H, m), 3.38 (1 H, br d, *J*=13.3 Hz), 3.15 (1 H, d, *J*=13.3 Hz), 2.72 - 2.85 (1 H, m), 1.85 (6 H, s), 1.31 (3 H, br dd, *J*=10.8, 6.6 Hz), 1.13 - 1.24 (3 H, m), 1.08 (3 H, d, *J*=6.6 Hz), 0.98 (3 H, d, *J*=6.6 Hz)
80-20	615.0	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 0.93 (dd, *J* = 15.6, 6.84 Hz, 3 H), 1.05 (dd, *J* = 6.6, 4.6 Hz, 3 H), 1.23 - 1.34 (m, 6 H), 1.88 (d, *J* = 3.5 Hz, 3 H), 2.16 (s, 6 H), 2.52 - 2.57 (m, 1H), 3.39 (s, 2 H), 3.47 - 3.56 (m, 0.5 H), 3.80 - 3.90 (m, 2 H), 4.05 - 4.21 (m, 1.5 H), 4.38 - 4.96 (m, 2 H), 5.70 - 5.81 (m, 1 H), 6.18 (dd, *J* = 16.6, 2.1 Hz, 1 H), 6.76 - 6.92

		(m, 1 H), 7.06 (br d, *J* = 8.7 Hz, 1 H), 7.15 (br d, *J* = 10.4 Hz, 1 H), 7.19 - 7.28 (m, 2 H), 7.29 - 7.38 (m, 1 H), 7.48 - 7.60 (m, 1 H), 8.32 (dd, *J* = 9.6, 1.4 Hz, 1 H); $^{19}F\{^{1}H\}$ NMR (376 MHz, DMSO-d_6) δ –129.38 - –129.36 (m, 1F), –129.29 - –129.26 (m, 1 F), –113.36 (dd, *J* = 34.7, 2.6 Hz, 1 F), –113.22 (dd, *J* = 35.6, 4.4 Hz, 1 F)
80-20-1	615.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 0.91 (d, *J* = 6.8 Hz, 3 H), 1.05 (d, *J* = 6.8 Hz, 3 H), 1.25 (br s, 3 H), 1.28 - 1.33 (m, 3 H), 1.88 (s, 3 H), 2.16 (s, 6 H), 2.52 - 2.58 (m, 1 H), 3.38 (s, 2 H), 3.45 - 4.21 (m, 4 H), 4.40 - 4.79 (m, 1 H), 4.86 (br s, 1 H), 5.69 - 5.81 (m, 1 H), 6.18 (dd, *J* = 16.7, 2.2 Hz, 1 H), 6.83 (td, *J* = 17.0, 10.5 Hz, 1 H), 7.07 (s, 1 H), 7.13 (s, 1 H), 7.20 - 7.28 (m, 2 H), 7.29 - 7.37 (m, 1 H), 7.49 - 7.60 (m, 1 H), 8.32 (d, *J* = 9.7 Hz, 1 H); $^{19}F\{^{1}H\}$ NMR (376 MHz, DMSO-d_6) δ –129.32 (dd, *J* = 30.1, 3.8 Hz, 1 F), –113.36 (dd, *J* = 30.1, 3.8 Hz, 1 F)
80-20-2	615.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 0.95 (d, *J* = 6.8 Hz, 3 H), 1.05 (dd, *J* = 6.4, 2.9 Hz, 3 H), 1.13 (d, *J* = 15.1 Hz, 3 H), 1.29 (br dd, *J* = 10.5, 6.7 Hz, 3 H), 1.87 (s, 3 H), 2.16 (s, 6 H), 2.45 - 2.49 (m, 1 H), 3.38 (s, 2 H), 3.80 - 4.21 (m, 4 H), 4.47 (br d, *J* = 4.8 Hz, 0.5 H), 4.72 - 4.90 (m, 1.5 H), 5.72 - 5.78 (m, 1 H), 6.19 (dd, *J* = 16.6, 2.3 Hz, 1 H), 6.76 - 6.91 (m, 1 H), 7.05 (s, 1 H), 7.16 (s, 1 H), 7.20 - 7.28 (m, 2 H), 7.29 - 7.36 (m, 1 H), 7.49 - 7.59 (m, 1 H), 8.33 (d, *J* = 9.7 Hz, 1 H); $^{19}F\{^{1}H\}$ NMR (376 MHz, DMSO-d_6) δ –129.32 (dd, *J* = 36.0, 3.9 Hz, 1 F), –113.22 (dd, *J* = 36.0, 3.9 Hz, 1 F)
80-21	631.1	^{1}H NMR (400 MHz, DMSO-d_6) δ 0.88 - 0.96 (m, 3 H), 1.04 (dd, *J* = 6.7, 2.6 Hz, 3 H), 1.11 - 1.22 (m, 3 H), 1.27 - 1.32 (m, 3 H), 1.85 (d, *J* = 1.9 Hz, 3 H), 2.16 (s, 6 H), 3.35 (s, 2 H), 3.78 - 3.92 (m, 2 H), 3.97 - 4.21 (m, 2 H), 4.42 - 4.94 (m, 2 H), 5.66 - 5.82 (m, 1 H), 6.18 (dd, *J* = 16.6, 2.3 Hz, 1 H), 6.60 - 6.72 (m, 2 H), 6.76 - 6.91 (m, 1 H), 7.02 (br s, 1 H), 7.10 (br s, 1 H), 7.20 - 7.31 (m, 2 H), 8.24 - 8.30 (m, 1 H), 10.26 (br s, 1 H); $^{19}F\{^{1}H\}$ NMR (376 MHz, DMSO-d_6) δ –128.74 (s, 0.5 F), –128.57 (s, 0.5 F), –115.38 (s, 0.5 F), –115.22 (s, 0.5 F)
80-21-1	630.6	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.30 - 9.81 (br s, 1 H) 7.84 - 7.93 (m, 1 H) 7.24 - 7.29 (m, 1 H) 7.22 (s, 1 H) 6.51 - 6.71 (m, 3 H) 6.40 (br t, *J*=14.82 Hz, 1 H) 5.73 - 5.88 (m, 1 H) 4.80 - 5.22 (m, 2 H) 4.28 - 4.50 (m, 1 H) 3.89 - 4.08 (m, 2 H) 3.67 - 3.87 (m, 1 H) 3.43 - 3.54 (m, 3 H) 2.54 - 2.68 (m, 1 H) 2.32 (s, 6 H) 1.94 - 2.03 (m, 3 H) 1.41 - 1.48 (m, 3 H) 1.25 - 1.32 (m, 3 H) 1.19 (d, *J*=6.63 Hz, 3 H) 1.00 (br d, *J*=6.84 Hz, 3 H)
80-21-2	630.6	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.56 (br s, 1 H) 7.85 - 7.95 (m, 1 H) 7.28 (s, 1 H) 7.20 - 7.25 (m, 1 H) 6.52 - 6.70 (m, 3 H) 6.34 - 6.45 (m, 1 H) 5.76 - 5.85 (m, 1 H) 4.86 - 5.19 (m, 2 H) 4.33 - 4.51 (m, 1 H) 3.87 - 4.09

		(m, 2 H) 3.68 - 3.84 (m, 1 H) 3.48 (s, 3 H) 2.48 - 2.61 (m, 1 H) 2.32 (s, 6 H) 2.00 (br d, *J*=7.05 Hz, 3 H) 1.44 (br d, *J*=6.43 Hz, 3 H) 1.33 (br d, *J*=7.05 Hz, 3 H) 1.20 (d, *J*=6.84 Hz, 3 H) 1.01 (m, *J*=6.40 Hz, 3 H)
92-1	599.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.42 (d, *J*=5.2 Hz, 1 H), 8.35 (d, *J*=5.0 Hz, 1 H), 7.37 (d, *J*=3.9 Hz, 2 H), 7.22 (dt, *J*=8.2, 4.0 Hz, 1 H), 7.16 (d, *J*=5.0 Hz, 1 H), 7.00 (d, *J*=7.5 Hz, 1 H), 6.84 (ddd, *J*=20.3, 16.8, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 4.44 - 4.93 (m, 2 H), 3.39 - 4.32 (m, 4 H), 2.74 (br s, 1 H), 2.52 - 2.60 (m, 1 H), 1.89 (s, 3 H), 1.36 (dd, *J*=6.3, 3.8 Hz, 3 H), 1.26 (br dd, *J*=24.7, 6.6 Hz, 3 H), 0.80 - 1.10 (m, 12 H).
92-2	576.0	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.56 (s, 1 H), 8.36 - 8.41 (m, 1 H), 7.18 (br d, *J*=4.6 Hz, 1 H), 7.05 (q, *J*=7.9 Hz, 1 H), 6.82 (dd, *J*=16.7, 10.5 Hz, 1 H), 6.41 - 6.48 (m, 1 H), 6.32 (t, *J*=8.8 Hz, 1 H), 6.19 (br d, *J*=16.4 Hz, 1 H), 5.71 - 5.79 (m, 1 H), 5.04 - 5.17 (m, 2 H), 4.47 - 4.81 (m, 1 H), 4.39 (br d, *J*=9.3 Hz, 1 H), 4.15 - 4.26 (m, 1 H), 3.97 - 4.13 (m, 1 H), 3.56 - 3.93 (m, 3 H), 2.54 - 2.83 (m, 1 H), 1.84 - 2.04 (m, 3 H), 1.29 (br s, 3 H), 0.82 - 1.12 (m, 6 H)
92-3	577.3	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.44 (d, *J*=4.8 Hz, 1 H), 8.34 - 8.41 (m, 1 H), 7.31 (d, *J*=5.4 Hz, 1 H), 7.22 (dd, *J*=8.5, 5.2 Hz, 2 H), 6.75 - 6.92 (m, 1 H), 6.19 (br d, *J*=16.6 Hz, 1 H), 5.70 - 5.80 (m, 1 H), 4.44 - 4.90 (m, 2 H), 3.41 - 4.25 (m, 4 H), 2.65 - 2.77 (m, 1 H), 1.99 (s, 3 H), 1.93 (s, 3 H), 1.34 (t, *J*=5.8 Hz, 3 H), 1.23 (br dd, *J*=25.8, 6.9 Hz, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.91 (d, *J*=6.6 Hz, 3 H)
92-4	597.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.44 (d, *J*=2.7 Hz, 1 H), 8.37 (d, *J*=4.8 Hz, 1 H), 7.30 (td, *J*=7.6, 0.9 Hz, 1 H), 7.15 - 7.22 (m, 2 H), 6.99 - 7.04 (m, 1 H), 6.94 (d, *J*=7.7 Hz, 1 H), 6.84 (td, *J*=16.5, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.72 - 5.79 (m, 1 H), 4.82 - 4.94 (m, 1 H), 4.44 - 4.81 (m, 1 H), 3.43 - 4.24 (m, 4 H), 2.68 - 2.76 (m, 1 H), 1.91 (s, 3 H), 1.43 - 1.52 (m, 1 H), 1.31 - 1.38 (m, 3 H), 1.17 - 1.29 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.94 (d, *J*=6.6 Hz, 3 H), 0.61 - 0.73 (m, 2 H), 0.52 - 0.59 (m, 1 H), 0.47 (br s, 1 H)
92-5	576.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.43 (d, *J*=5.0 Hz, 1 H), 8.40 (d, *J*=5.2 Hz, 1 H), 7.90 (d, *J*=3.1 Hz, 1 H), 7.22 (d, *J*=4.8 Hz, 1 H), 7.16 (dd, *J*=3.1, 0.8 Hz, 1 H), 6.75 - 6.92 (m, 1 H), 6.19 (br d, *J*=16.6 Hz, 1 H), 5.71 - 5.79 (m, 1 H), 4.43 - 4.93 (m, 2 H), 3.41 - 4.25 (m, 4 H), 2.65 - 2.75 (m, 1 H), 1.92 (s, 3 H), 1.70 (s, 3 H), 1.34 (br t, *J*=5.8 Hz, 3 H), 1.17 - 1.27 (m, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.89 (br d, *J*=6.6 Hz, 3 H)
92-6	560.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.46 (d, *J*=4.8 Hz, 1 H), 8.30 (d, *J*=2.1 Hz, 1 H), 7.24 (d, *J*=5.0 Hz, 1 H), 6.82 (td, *J*=16.0, 10.5 Hz, 1 H), 6.33 (dt, *J*=3.6, 2.0 Hz, 1 H), 6.18 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.71 - 5.79 (m, 1 H), 4.78 - 4.90

		(m, 1 H), 4.40 - 4.78 (m, 1 H), 3.41 - 4.18 (m, 4 H), 2.58 - 2.69 (m, 1 H), 2.13 (br d, *J*=1.5 Hz, 2 H), 1.93 - 2.07 (m, 2 H), 1.90 (s, 3 H), 1.45 - 1.58 (m, 4 H), 1.30 (t, *J*=7.4 Hz, 3 H), 1.18 (br dd, *J*=29.0, 6.6 Hz, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.97 (d, *J*=6.6 Hz, 3 H)
92-7	590.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.44 (d, *J*=4.8 Hz, 1 H), 8.36 (d, *J*=5.4 Hz, 1 H), 7.22 (d, *J*=5.0 Hz, 1 H), 6.89 (s, 1 H), 6.76 - 6.88 (m, 1 H), 6.19 (dd, *J*=16.6, 1.7 Hz, 1 H), 5.72 - 5.78 (m, 1 H), 4.41 - 4.89 (m, 2 H), 3.40 - 4.25 (m, 4 H), 2.65 - 2.77 (m, 1 H), 2.35 (s, 3 H), 1.91 (d, *J*=7.0 Hz, 6 H), 1.34 (br t, *J*=5.8 Hz, 3 H), 1.18 - 1.27 (m, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.92 (d, *J*=6.6 Hz, 3 H)
92-8	534.8	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.45 (d, *J*=5.0 Hz, 1 H), 8.25 (d, *J*=2.3 Hz, 1 H), 7.24 (d, *J*=4.8 Hz, 1 H), 6.82 (td, *J*=16.2, 10.6 Hz, 1 H), 6.56 (s, 1 H), 6.18 (br d, *J*=16.6 Hz, 1 H), 5.70 - 5.78 (m, 1 H), 4.78 - 4.89 (m, 1 H), 4.39 - 4.77 (m, 1 H), 3.41 - 4.18 (m, 4 H), 2.61 - 2.70 (m, 1 H), 1.92 (s, 3 H), 1.88 (s, 3 H), 1.37 (s, 3 H), 1.30 (br t, *J*=7.6 Hz, 3 H), 1.12 - 1.22 (m, 3 H), 1.07 (d, *J*=6.8 Hz, 3 H), 0.93 (d, *J*=6.4 Hz, 3 H)
92-9	563.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.47 (d, *J*=4.8 Hz, 1 H), 8.33 (d, *J*=2.7 Hz, 1 H), 7.24 (d, *J*=4.8 Hz, 1 H), 6.83 (td, *J*=16.1, 10.5 Hz, 1 H), 6.54 - 6.59 (m, 1 H), 6.18 (dd, *J*=16.5, 2.0 Hz, 1 H), 5.72 - 5.77 (m, 1 H), 4.79 - 4.91 (m, 1 H), 4.39 - 4.78 (m, 1 H), 3.39 - 4.23 (m, 8 H), 2.59 - 2.69 (m, 1 H), 1.95 - 2.13 (m, 2 H), 1.90 (s, 3 H), 1.30 (t, *J*=7.3 Hz, 3 H), 1.13 - 1.23 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.96 (d, *J*=6.6 Hz, 3 H)
92-10	575.3	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.45 (d, *J*=4.8 Hz, 1 H), 8.28 (d, *J*=2.9 Hz, 1 H), 7.23 (d, *J*=5.0 Hz, 1 H), 6.83 (td, *J*=16.5, 10.6 Hz, 1 H), 6.37 (t, *J*=6.7 Hz, 1 H), 6.14 - 6.22 (m, 1 H), 5.71 - 5.78 (m, 1 H), 4.82 (dt, *J*=6.7, 3.5 Hz, 1 H), 4.40 - 4.78 (m, 1 H), 3.40 - 4.18 (m, 4 H), 2.58 - 2.68 (m, 1 H), 2.17 - 2.28 (m, 4 H), 1.90 (s, 3 H), 1.67 (quin, *J*=5.8 Hz, 2 H), 1.35 - 1.43 (m, 2 H), 1.30 (t, *J*=7.3 Hz, 3 H), 1.13 - 1.26 (m, 5 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.97 (d, *J*=6.6 Hz, 3 H)
92-11	596.2	^{1}H NMR (400 MHz, DMSO-*d*$_6$) δ 10.04 (br s, 1 H), 8.49 (br s, 2 H), 7.66 (br d, *J*=7.7 Hz, 1 H), 7.42 - 7.50 (m, 1 H), 7.26 (br d, *J*=4.4 Hz, 1 H), 7.15 (br s, 1 H), 7.07 (t, *J*=7.6 Hz, 1 H), 6.86 (td, *J*=17.0, 10.6 Hz, 1 H), 6.48 (br s, 1 H), 6.20 (br d, *J*=16.4 Hz, 1 H), 5.76 (br d, *J*=9.5 Hz, 1 H), 4.89 (br s, 1 H), 4.44 - 4.83 (m, 1 H), 3.44 - 4.26 (m, 4 H), 2.70 - 2.86 (m, 1 H), 1.98 (s, 3 H), 1.35 (br t, *J*=6.9 Hz, 3 H), 1.16 - 1.30 (m, 3 H), 1.09 (br d, *J*=6.6 Hz, 3 H), 0.92 (br d, *J*=6.6 Hz, 3 H)
92-12	572.2	^{1}H NMR (400 MHz, DMSO-*d*$_6$) δ 8.44 (br d, *J*=4.8 Hz, 1 H), 8.39 (br s, 1 H), 7.22 (br d, *J*=4.6 Hz, 1 H), 7.03 - 7.13 (m, 2 H), 6.76 - 6.93 (m, 1 H), 6.65 (br d, *J*=8.1 Hz, 1 H), 6.54 (br t, *J*=7.2 Hz, 1 H), 6.19 (br d, *J*=16.8 Hz, 1 H),

		5.75 (br d, *J*=7.7 Hz, 1 H), 5.09 (br s, 2 H), 4.43 - 4.94 (m, 2 H), 3.43 - 4.21 (m, 4 H), 2.65 - 2.80 (m, 1 H), 1.93 (s, 3 H), 1.33 (br t, *J*=6.3 Hz, 3 H), 1.16 - 1.28 (m, 3 H), 1.07 (br d, *J*=6.4 Hz, 3 H), 0.98 (br d, *J*=6.4 Hz, 3 H)
92-13	573.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.38 - 8.47 (m, 2 H), 7.98 (br d, *J*=3.5 Hz, 1 H), 7.36 (br d, *J*=7.0 Hz, 1 H), 7.20 (d, *J*=4.8 Hz, 1 H), 6.76 - 6.94 (m, 1 H), 6.58 (dd, *J*=7.0, 5.0 Hz, 1 H), 6.14 - 6.28 (m, 1 H), 5.83 (s, 2 H), 5.75 (br d, *J*=9.7 Hz, 1 H), 4.44 - 4.92 (m, 2 H), 3.45 - 4.25 (m, 4 H), 2.63 - 2.77 (m, 1 H), 1.94 (s, 3 H), 1.29 - 1.39 (m, 3 H), 1.13 - 1.28 (m, 3 H), 1.06 (br d, *J*=6.6 Hz, 3 H), 0.97 (br d, *J*=6.6 Hz, 3 H)
92-14	623.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 (br s, 1 H), 8.38 (br d, *J*=4.1 Hz, 1 H), 7.47 - 7.58 (m, 1 H), 7.23 - 7.35 (m, 2 H), 7.18 (br d, *J*=3.7 Hz, 2 H), 6.76 - 7.09 (m, 2 H), 6.19 (br d, *J*=16.6 Hz, 1 H), 5.75 (br d, *J*=8.5 Hz, 1 H), 4.42 - 4.94 (m, 2 H), 3.44 - 4.24 (m, 4 H), 2.69 (br d, *J*=8.3 Hz, 1 H), 1.92 (br s, 3 H), 1.33 (br s, 3 H), 1.16 - 1.28 (m, 3 H), 1.06 (br d, *J*=6.2 Hz, 3 H), 0.93 (br d, *J*=4.1 Hz, 3 H)
92-15	611.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 12.44 - 12.72 (m, 1 H), 8.51 - 8.61 (m, 1 H), 8.31 (dd, *J*=8.3, 4.8 Hz, 1 H), 8.02 (d, *J*=14.7 Hz, 1 H), 7.69 (dd, *J*=8.1, 5.6 Hz, 1 H), 7.07 - 7.21 (m, 1 H), 7.01 (dd, *J*=11.2, 8.3 Hz, 1 H), 6.78 - 6.95 (m, 1 H), 6.21 (br d, *J*=16.6 Hz, 1 H), 5.74 - 5.83 (m, 1 H), 4.47 - 4.99 (m, 2 H), 3.83 - 4.25 (m, 4 H), 2.63 - 2.86 (m, 1 H), 2.03 - 2.11 (m, 3 H), 1.90 - 1.98 (m, 3 H), 1.32 - 1.41 (m, 3 H), 1.21 - 1.31 (m, 3 H), 1.07 (dd, *J*=6.5, 4.3 Hz, 3 H), 0.73 - 0.99 (m, 3 H)
92-16	607.1	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 (d, *J*=4.4 Hz, 1 H), 8.37 (d, *J*=4.8 Hz, 1 H), 7.55 - 7.70 (m, 3 H), 7.36 (br d, *J*=6.6 Hz, 1 H), 7.18 (d, *J*=5.0 Hz, 1 H), 6.76 - 6.93 (m, 1 H), 6.38 - 6.72 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.76 (ddd, *J*=10.2, 5.1, 2.5 Hz, 1 H), 4.46 - 4.94 (m, 2 H), 3.76 - 4.24 (m, 4 H), 2.72 (dt, *J*=6.8, 3.2 Hz, 1 H), 1.91 (s, 3 H), 1.31 - 1.39 (m, 3 H), 1.16 - 1.29 (m, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.90 (d, *J*=6.6 Hz, 3 H)
92-17	590.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.36 - 8.45 (m, 2 H), 7.20 (d, *J*=4.8 Hz, 1 H), 7.05 (dd, *J*=10.7, 8.0 Hz, 1 H), 6.75 - 6.92 (m, 2 H), 6.54 (td, *J*=7.8, 5.1 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.69 - 5.83 (m, 1 H), 5.02 (s, 2 H), 4.81 - 4.92 (m, 1 H), 4.46 - 4.80 (m, 1 H), 3.45 - 4.23 (m, 4 H), 2.63 - 2.78 (m, 1 H), 1.94 (s, 3 H), 1.33 (br t, *J*=6.9 Hz, 3 H), 1.15 - 1.29 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.98 (d, *J*=6.6 Hz, 3 H)
92-18	614.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.98 - 9.29 (m, 1 H), 8.42 (br s, 1 H), 8.39 (br d, *J*=4.1 Hz, 1 H), 7.62 (br s, 1 H), 7.38 (br t, *J*=7.3 Hz, 1 H), 7.03 - 7.24 (m, 3 H), 6.72 - 6.97 (m, 1 H), 6.19 (br d, *J*=16.8 Hz, 1 H), 5.63 - 5.88 (m, 1 H), 4.46 - 5.00 (m, 2 H), 3.73 - 4.27 (m, 4 H), 2.66 - 2.81 (m, 1 H), 1.97 (br s, 3 H), 1.85 (br s, 3 H), 1.33 (br d,

		J=6.0 Hz, 3 H), 1.14 - 1.25 (m, 3 H), 1.06 (br d, *J*=6.2 Hz, 3 H), 0.93 (br d, *J*=5.8 Hz, 3 H)
92-19	600.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.38 (br d, *J*=4.8 Hz, 2 H), 7.32 (br t, *J*=6.9 Hz, 1 H), 7.19 (br s, 1 H), 7.06 (br d, *J*=8.1 Hz, 1 H), 6.70 - 6.98 (m, 3 H), 6.19 (dd, *J*=16.7, 2.0 Hz, 1 H), 5.67 - 5.84 (m, 1 H), 4.41 - 4.93 (m, 2 H), 3.38 - 4.33 (m, 4 H), 2.72 - 2.88 (m, 1 H), 2.44 (s, 6 H), 1.90 (br s, 3 H), 1.35 (br d, *J*=6.0 Hz, 3 H), 1.14 - 1.29 (m, 3 H), 1.05 - 1.14 (m, 3 H), 0.98 (br d, *J*=3.5 Hz, 3 H)
92-20	612.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 9.97 (br d, *J*=7.5 Hz, 1 H), 8.45 (s, 1 H), 8.39 (d, *J*=4.8 Hz, 1 H), 7.25 (br d, *J*=6.8 Hz, 1 H), 7.18 (d, *J*=5.0 Hz, 1 H), 6.94 - 7.03 (m, 2 H), 6.85 (td, *J*=16.1, 10.4 Hz, 1 H), 6.20 (dd, *J*=16.6, 2.1 Hz, 1 H), 5.72 - 5.81 (m, 1 H), 4.87 (br s, 1 H), 4.43 - 4.82 (m, 1 H), 3.79 - 4.24 (m, 4 H), 3.52 (br s, 2 H), 2.64 - 2.75 (m, 1 H), 1.95 (s, 3 H), 1.29 - 1.38 (m, 3 H), 1.13 - 1.27 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.98 (d, *J*=6.6 Hz, 3 H)
92-21	600.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.35 (d, *J*=4.8 Hz, 1 H), 8.32 (d, *J*=3.1 Hz, 1 H), 7.81 (br s, 1 H), 7.68 (br d, *J*=7.3 Hz, 1 H), 7.50 (quin, *J*=7.5 Hz, 2 H), 7.10 - 7.20 (m, 3 H), 6.75 - 6.92 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.75 (ddd, *J*=10.1, 5.5, 2.0 Hz, 1 H), 4.47 - 4.90 (m, 2 H), 3.75 - 4.22 (m, 4 H), 2.65 - 2.77 (m, 1 H), 1.92 (s, 3 H), 1.34 (br t, *J*=5.5 Hz, 3 H), 1.18 - 1.29 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.90 (br d, *J*=6.6 Hz, 3 H)
92-22	497.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 11.53 - 11.97 (br s, 1 H), 8.39 (d, *J*=4.8 Hz, 1 H), 7.81 (br s, 1 H), 7.18 (d, *J*=4.8 Hz, 1 H), 6.81 (td, *J*=16.5, 10.7 Hz, 1 H), 6.16 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.67 - 5.80 (m, 1 H), 4.28 - 4.79 (m, 2 H), 3.42 - 4.15 (m, 4 H), 2.56 - 2.66 (m, 1 H), 1.91 (s, 3 H), 1.15 - 1.30 (m, 6 H), 1.04 (dd, *J*=9.7, 6.8 Hz, 6 H)
92-23	625.1	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.50 (br s, 1 H), 8.35 (d, *J*=4.8 Hz, 1 H), 7.80 (d, *J*=7.9 Hz, 1 H), 7.72 - 7.77 (m, 1 H), 7.62 - 7.71 (m, 1 H), 7.36 (d, *J*=7.5 Hz, 1 H), 7.16 (d, *J*=4.8 Hz, 1 H), 6.75 - 6.92 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.76 (ddd, *J*=10.4, 5.3, 2.2 Hz, 1 H), 4.46 - 4.94 (m, 2 H), 3.41 - 4.28 (m, 4 H), 2.69 (br dd, *J*=16.1, 1.3 Hz, 1 H), 1.88 (s, 3 H), 1.32 - 1.40 (m, 3 H), 1.15 - 1.30 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.89 (br d, *J*=1.2 Hz, 3 H)
92-24	618.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.29 - 8.43 (m, 2 H), 7.80 - 7.98 (m, 1 H), 7.37 - 7.61 (m, 3 H), 7.17 - 7.33 (m, 1 H), 7.07 - 7.16 (m, 1 H), 6.71 - 6.94 (m, 1 H), 6.19 (dd, *J*=16.7, 2.0 Hz, 1 H), 5.75 (dt, *J*=10.3, 2.6 Hz, 1 H), 4.44 - 4.93 (m, 2 H), 3.40 - 4.30 (m, 4 H), 2.64 - 2.84 (m, 1 H), 1.79 - 1.95 (m, 3 H), 1.35 (br d, *J*=4.4 Hz, 3 H), 1.17 - 1.29 (m, 3 H), 1.04 - 1.12 (m, 3 H), 0.79 - 0.98 (m, 3 H)
92-25	586.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.31 - 8.50 (m, 2 H), 7.17 - 7.29 (m, 2 H), 7.14 (br d, *J*=7.5 Hz, 1 H), 6.85 (td, *J*=17.3, 10.5 Hz, 1 H), 6.52 - 6.65 (m, 2 H), 6.19 (dd,

		J=16.6, 2.3 Hz, 1 H), 5.68 - 5.82 (m, 1 H), 5.32 (br d, *J*=4.4 Hz, 1 H), 4.42 - 4.94 (m, 2 H), 3.45 - 4.24 (m, 4 H), 2.62 - 2.79 (m, 1 H), 2.47 (br s, 3 H), 1.95 (s, 3 H), 1.33 (t, *J*=6.9 Hz, 3 H), 1.15 - 1.28 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.96 (d, *J*=6.6 Hz, 3 H)
92-26	608.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.38 - 8.47 (m, 2 H), 7.54 - 7.60 (m, 1 H), 7.43 - 7.52 (m, 1 H), 7.19 - 7.23 (m, 1 H), 7.11 - 7.19 (m, 1 H), 6.85 (td, *J*=16.6, 10.6 Hz, 1 H), 6.69 (br d, *J*=7.9 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.70 - 5.82 (m, 1 H), 4.73 - 4.86 (m, 1 H), 4.50 (br d, *J*=1.7 Hz, 1 H), 4.06 - 4.26 (m, 2 H), 3.83 - 3.98 (m, 2 H), 2.63 - 2.79 (m, 1 H), 1.94 (s, 3 H), 1.29 - 1.38 (m, 3 H), 1.14 - 1.26 (m, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.98 (d, *J*=6.6 Hz, 3 H)
92-27	614.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.18 - 8.42 (m, 3 H), 7.55 - 7.73 (m, 1 H), 7.39 - 7.54 (m, 2 H), 7.16 (d, *J*=5.0 Hz, 2 H), 6.72 - 6.93 (m, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.68 - 5.82 (m, 1 H), 4.44 - 4.99 (m, 2 H), 3.73 - 4.26 (m, 4 H), 2.65 - 2.78 (m, 1 H), 2.60 (d, *J*=4.4 Hz, 3 H), 1.84 - 1.92 (m, 3 H), 1.35 (br d, *J*=6.4 Hz, 3 H), 1.20 - 1.31 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.88 (br d, *J*=5.2 Hz, 3 H)
92-28	628.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.33 - 8.47 (m, 2 H), 7.42 - 7.51 (m, 2 H), 7.20 - 7.41 (m, 3 H), 6.73 - 6.92 (m, 1 H), 6.19 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.66 - 5.84 (m, 1 H), 4.41 - 4.93 (m, 2 H), 3.67 - 4.20 (m, 4 H), 2.70 (s, 3 H), 2.58 (br d, *J*=5.6 Hz, 1 H), 2.32 (br dd, *J*=4.3, 2.4 Hz, 3 H), 1.92 (s, 3 H), 1.32 (br t, *J*=6.8 Hz, 3 H), 1.14 - 1.27 (m, 3 H), 1.01 (d, *J*=6.6 Hz, 3 H), 0.80 (br d, *J*=6.0 Hz, 3 H)
92-29	601.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 12.12 (br s, 1 H), 8.24 - 8.45 (m, 2 H), 7.93 (br d, *J*=7.5 Hz, 1 H), 7.39 - 7.68 (m, 2 H), 7.05 - 7.21 (m, 2 H), 6.83 (td, *J*=16.0, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.62 - 5.87 (m, 1 H), 4.40 - 5.00 (m, 2 H), 3.45 - 4.28 (m, 4 H), 2.62 - 2.77 (m, 1 H), 1.91 (s, 3 H), 1.33 (br t, *J*=6.9 Hz, 3 H), 1.16 - 1.27 (m, 3 H), 1.07 (br d, *J*=6.6 Hz, 3 H), 0.91 (br s, 3 H)
92-30	597.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 12.24 (br s, 1 H), 8.54 (s, 1 H), 8.42 (d, *J*=4.8 Hz, 1 H), 8.13 (s, 1 H), 7.88 (d, *J*=8.1 Hz, 1 H), 7.48 (br dd, *J*=7.2, 3.8 Hz, 1 H), 7.14 - 7.26 (m, 2 H), 6.86 (td, *J*=16.4, 10.6 Hz, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.71 - 5.83 (m, 1 H), 4.45 - 5.02 (m, 2 H), 3.73 - 4.31 (m, 4 H), 2.73 (td, *J*=6.6, 3.1 Hz, 1 H), 1.97 (s, 3 H), 1.35 (br t, *J*=7.6 Hz, 3 H), 1.16 - 1.28 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.92 (d, *J*=6.6 Hz, 3 H)
92-31	608.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.45 (d, *J*=5.0 Hz, 1 H), 8.42 (d, *J*=2.1 Hz, 1 H), 7.23 (d, *J*=5.0 Hz, 1 H), 7.13 (ddd, *J*=11.6, 9.0, 3.1 Hz, 1 H), 6.84 (td, *J*=17.0, 10.4 Hz, 1 H), 6.60 (dd, *J*=13.2, 7.4 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.69 - 5.83 (m, 1 H), 5.29 (s, 2 H), 4.44 - 4.94

		(m, 2 H), 3.45 - 4.22 (m, 4 H), 2.65 - 2.77 (m, 1 H), 1.93 (s, 3 H), 1.32 (br t, *J*=6.8 Hz, 3 H), 1.16 - 1.26 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.98 (d, *J*=6.6 Hz, 3 H)
92-32	650.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.92 (br s, 1 H), 8.44 (s, 1 H), 8.37 (d, *J*=5.0 Hz, 1 H), 7.44 (br d, *J*=2.9 Hz, 2 H), 7.23 (br s, 1 H), 7.17 (d, *J*=5.0 Hz, 1 H), 7.14 (br d, *J*=6.6 Hz, 1 H), 6.82 (br dd, *J*=16.8, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.62 - 5.87 (m, 1 H), 4.46 - 4.92 (m, 2 H), 3.78 - 4.21 (m, 4 H), 2.66 - 2.74 (m, 1 H), 2.63 (s, 3 H), 1.96 (s, 3 H), 1.32 (br t, *J*=7.3 Hz, 3 H), 1.14 - 1.25 (m, 3 H), 1.05 (d, *J*=6.8 Hz, 3 H), 0.95 (d, *J*=6.6 Hz, 3 H)
92-33	636.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (s, 1 H), 8.36 (d, *J*=4.8 Hz, 1 H), 7.91 - 7.98 (m, 1 H), 7.62 - 7.72 (m, 2 H), 7.21 - 7.33 (m, 1 H), 7.15 (d, *J*=5.0 Hz, 1 H), 6.79 - 6.90 (m, 1 H), 6.73 (s, 2 H), 6.19 (dd, *J*=16.7, 2.0 Hz, 1 H), 5.71 - 5.81 (m, 1 H), 4.45 - 4.99 (m, 2 H), 3.82 - 4.30 (m, 4 H), 2.63 - 2.74 (m, 1 H), 1.95 (s, 3 H), 1.33 (s, 3 H), 1.13 - 1.22 (m, 3 H), 1.05 (br d, *J*=6.4 Hz, 3 H), 0.88 (br d, *J*=6.4 Hz, 3 H)
92-34	591.0	^{1}H NMR (400 MHz, DMSO-d_6) δ 9.66 (br s, 1 H), 8.43 (d, *J*=3.7 Hz, 1 H), 8.41 (d, *J*=4.8 Hz, 1 H), 7.20 (d, *J*=4.8 Hz, 1 H), 7.08 (t, *J*=9.2 Hz, 1 H), 6.70 - 6.93 (m, 2 H), 6.53 (dd, *J*=5.7, 3.0 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.0 Hz, 1 H), 5.68 - 5.81 (m, 1 H), 4.42 - 4.95 (m, 2 H), 4.11 - 4.24 (m, 1 H), 3.43 - 3.94 (m, 3 H), 2.63 - 2.75 (m, 1 H), 1.93 (s, 3 H), 1.29 - 1.40 (m, 3 H), 1.17 - 1.27 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.97 (d, *J*=6.6 Hz, 3 H)
92-35	656.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 9.11 (br s, 1 H), 8.34 - 8.50 (m, 2 H), 7.36 - 7.52 (m, 1 H), 7.17 - 7.30 (m, 3 H), 7.11 (br d, *J*=7.5 Hz, 1 H), 6.74 - 6.95 (m, 1 H), 6.19 (br d, *J*=16.0 Hz, 1 H), 5.66 - 5.82 (m, 1 H), 4.41 - 5.02 (m, 2 H), 3.50 - 4.27 (m, 4 H), 2.61 - 2.72 (m, 1 H), 2.00 (br s, 3 H), 1.23 - 1.34 (m, 3 H), 1.01 - 1.10 (m, 9 H), 0.97 (br s, 9 H)
92-36	602.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (d, *J*=3.5 Hz, 1 H), 8.36 (d, *J*=4.8 Hz, 1 H), 8.14 (d, *J*=8.1 Hz, 1 H), 7.81 - 7.90 (m, 1 H), 7.68 - 7.77 (m, 1 H), 7.41 - 7.49 (m, 1 H), 7.17 (d, *J*=5.0 Hz, 1 H), 6.84 (td, *J*=17.5, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.69 - 5.82 (m, 1 H), 4.44 - 4.95 (m, 2 H), 3.45 - 4.23 (m, 4 H), 2.63 - 2.78 (m, 1 H), 1.90 (s, 3 H), 1.35 (br t, *J*=5.7 Hz, 3 H), 1.16 - 1.28 (m, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.88 (br d, *J*=6.6 Hz, 3 H)
92-37	653.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (d, *J*=2.5 Hz, 1 H), 8.41 (d, *J*=5.0 Hz, 1 H), 7.72 (ddd, *J*=8.9, 4.6, 2.7 Hz, 1 H), 7.42 (dd, *J*=6.3, 2.6 Hz, 1 H), 7.33 (t, *J*=9.3 Hz, 1 H), 7.21 (d, *J*=4.8 Hz, 1 H), 6.72 - 6.90 (m, 1 H), 6.19 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.69 - 5.82 (m, 1 H), 4.41 - 4.94 (m, 2 H), 3.45 - 4.22 (m, 4 H), 2.67 - 2.77 (m, 1 H), 1.94 (s, 3 H), 1.32 (br t, *J*=7.0 Hz, 3 H), 1.13 - 1.27 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.99 (d, *J*=6.6 Hz, 3 H)

92-38	626.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.36 - 8.50 (m, 2 H), 7.20 (d, *J*=5.0 Hz, 1 H), 6.68 - 6.98 (m, 2 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.75 (ddd, *J*=10.3, 5.1, 2.0 Hz, 1 H), 5.28 (s, 2 H), 4.45 - 4.92 (m, 2 H), 3.43 - 4.25 (m, 4 H), 2.60 - 2.73 (m, 1 H), 1.93 (s, 3 H), 1.29 - 1.36 (m, 3 H), 1.13 - 1.26 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.98 (d, *J*=6.6 Hz, 3 H)
92-39	613.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (d, *J*=2.7 Hz, 1 H), 8.41 (d, *J*=5.0 Hz, 1 H), 7.51 (ddd, *J*=8.5, 5.1, 2.2 Hz, 1 H), 7.30 (t, *J*=9.2 Hz, 1 H), 7.24 (dd, *J*=6.8, 2.1 Hz, 1 H), 7.20 (d, *J*=4.8 Hz, 1 H), 6.83 (td, *J*=16.4, 10.8 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.71 - 5.80 (m, 1 H), 4.40 - 4.98 (m, 2 H), 3.46 - 4.22 (m, 4 H), 2.65 - 2.76 (m, 1 H), 2.01 (s, 3 H), 1.93 (s, 3 H), 1.32 (br t, *J*=6.9 Hz, 3 H), 1.13 - 1.26 (m, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.97 (d, *J*=6.6 Hz, 3 H)
92-40	600.1	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.50 - 8.64 (m, 1 H), 8.34 - 8.42 (m, 1 H), 7.80 - 7.92 (m, 1 H), 7.65 - 7.79 (m, 2 H), 7.12 - 7.25 (m, 1 H), 6.83 (td, *J*=17.7, 10.4 Hz, 1 H), 6.12 - 6.27 (m, 1 H), 5.75 (br dd, *J*=10.6, 2.3 Hz, 1 H), 4.41 - 4.98 (m, 2 H), 3.43 - 4.37 (m, 4 H), 2.63 - 2.83 (m, 1 H), 1.95 (s, 3 H), 1.36 (br d, *J*=6.4 Hz, 3 H), 1.19 - 1.30 (m, 3 H), 1.03 - 1.14 (m, 3 H), 0.80 - 1.00 (m, 3 H)
92-41	599.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.36 (d, *J*=5.0 Hz, 2 H), 7.94 (dd, *J*=7.6, 0.9 Hz, 1 H), 7.53 - 7.68 (m, 2 H), 7.23 (dd, *J*=7.4, 0.9 Hz, 1 H), 7.17 (d, *J*=4.8 Hz, 1 H), 6.83 (td, *J*=17.3, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.68 - 5.84 (m, 1 H), 4.41 - 4.93 (m, 2 H), 3.41 - 4.26 (m, 4 H), 2.62 - 2.78 (m, 1 H), 2.35 (s, 3 H), 1.89 (s, 3 H), 1.31 - 1.37 (m, 3 H), 1.18 - 1.29 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.88 (br d, *J*=6.4 Hz, 3 H)
92-42	557.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.37 - 8.51 (m, 2 H), 7.38 - 7.56 (m, 5 H), 7.22 (d, *J*=4.8 Hz, 1 H), 6.84 (td, *J*=16.4, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.5, 2.0 Hz, 1 H), 5.72 - 5.81 (m, 1 H), 4.41 - 4.95 (m, 2 H), 3.39 - 4.35 (m, 4 H), 2.63 - 2.84 (m, 1 H), 1.94 (s, 3 H), 1.33 (t, *J*=7.0 Hz, 3 H), 1.14 - 1.26 (m, 3 H), 1.08 (d, *J*=6.6 Hz, 3 H), 0.98 (d, *J*=6.6 Hz, 3 H)
92-43	601.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.39 - 8.47 (m, 1 H), 8.35 (dd, *J*=6.4, 5.0 Hz, 1 H), 7.56 (d, *J*=8.1 Hz, 1 H), 7.41 (t, *J*=7.7 Hz, 1 H), 7.23 - 7.31 (m, 1 H), 7.16 (dd, *J*=4.9, 2.0 Hz, 1 H), 7.04 (d, *J*=7.7 Hz, 1 H), 6.71 - 6.93 (m, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.69 - 5.81 (m, 1 H), 4.95 (br d, *J*=4.1 Hz, 1 H), 4.42 - 4.88 (m, 2 H), 3.34 - 4.38 (m, 5 H), 2.63 - 2.79 (m, 1 H), 1.91 (d, *J*=3.9 Hz, 3 H), 1.30 - 1.38 (m, 3 H), 1.14 - 1.30 (m, 3 H), 1.05 (dd, *J*=10.2, 6.6 Hz, 3 H), 0.78 - 0.97 (m, 6 H)
92-44	609.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.50 (d, *J*=2.7 Hz, 1 H), 8.41 (d, *J*=4.8 Hz, 1 H), 7.60 (ddd, *J*=8.9, 4.3, 2.8 Hz, 1 H), 7.39 (t, *J*=9.2 Hz, 1 H), 7.31 (dd, *J*=6.0, 2.7 Hz, 1 H),

		7.21 (d, *J*=5.0 Hz, 1 H), 6.84 (td, *J*=16.5, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.70 - 5.81 (m, 1 H), 4.81 - 4.95 (m, 1 H), 4.43 - 4.80 (m, 1 H), 3.47 - 4.20 (m, 4 H), 2.63 - 2.78 (m, 1 H), 1.94 (s, 3 H), 1.33 (br t, *J*=6.9 Hz, 3 H), 1.13 - 1.25 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.98 (d, *J*=6.6 Hz, 3 H)
92-45	571.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.44 (d, *J*=3.9 Hz, 1 H), 8.37 (d, *J*=4.8 Hz, 1 H), 7.14 - 7.37 (m, 4 H), 7.08 (d, *J*=7.3 Hz, 1 H), 6.84 (td, *J*=17.3, 10.4 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.69 - 5.82 (m, 1 H), 4.42 - 4.97 (m, 2 H), 3.42 - 4.27 (m, 4 H), 2.72 (td, *J*=6.6, 3.2 Hz, 1 H), 1.96 (s, 3 H), 1.92 (s, 3 H), 1.34 (t, *J*=6.1 Hz, 3 H), 1.16 - 1.30 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.93 (d, *J*=6.8 Hz, 3 H)
92-46	609.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.50 (br s, 1 H), 8.40 (br d, *J*=4.6 Hz, 1 H), 7.71 (br t, *J*=7.3 Hz, 1 H), 7.32 (br t, *J*=7.8 Hz, 1 H), 7.16 - 7.25 (m, 2 H), 6.74 - 6.93 (m, 1 H), 6.19 (br d, *J*=16.4 Hz, 1 H), 5.75 (br d, *J*=8.9 Hz, 1 H), 4.43 - 4.96 (m, 2 H), 3.44 - 4.26 (m, 4 H), 2.70 (br s, 1 H), 1.93 (s, 3 H), 1.33 (br t, *J*=5.8 Hz, 3 H), 1.14 - 1.29 (m, 3 H), 1.06 (br d, *J*=6.4 Hz, 3 H), 0.96 (br d, *J*=6.2 Hz, 3 H)
92-47	581.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.47 (d, *J*=2.3 Hz, 1 H), 8.36 (d, *J*=5.0 Hz, 1 H), 7.50 - 7.60 (m, 1 H), 7.40 - 7.49 (m, 2 H), 7.18 (d, *J*=5.2 Hz, 2 H), 6.83 (td, *J*=16.4, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.66 - 5.86 (m, 1 H), 4.42 - 4.96 (m, 2 H), 3.74 - 4.22 (m, 5 H), 2.63 - 2.77 (m, 1 H), 1.93 (s, 3 H), 1.34 (br t, *J*=6.8 Hz, 3 H), 1.15 - 1.29 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.93 (d, *J*=6.6 Hz, 3 H)
92-48	581.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.42 - 8.49 (m, 2 H), 7.53 - 7.74 (m, 3 H), 7.40 - 7.49 (m, 1 H), 7.22 (d, *J*=4.8 Hz, 1 H), 6.84 (td, *J*=16.4, 10.8 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.68 - 5.82 (m, 1 H), 4.82 - 4.95 (m, 1 H), 4.44 - 4.80 (m, 1 H), 4.25 (s, 1 H), 3.46 - 4.18 (m, 4 H), 2.65 - 2.78 (m, 1 H), 1.93 (s, 3 H), 1.33 (br t, *J*=7.2 Hz, 3 H), 1.13 - 1.27 (m, 3 H), 1.08 (d, *J*=5.8 Hz, 3 H), 1.00 (d, *J*=6.6 Hz, 3 H)
92-49	595.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.42 - 8.49 (m, 2 H), 7.34 - 7.54 (m, 4 H), 7.22 (d, *J*=5.0 Hz, 1 H), 6.84 (td, *J*=16.5, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.70 - 5.80 (m, 1 H), 4.81 - 4.95 (m, 1 H), 4.41 - 4.80 (m, 1 H), 3.45 - 4.20 (m, 4 H), 2.64 - 2.77 (m, 1 H), 2.04 (s, 3 H), 1.93 (s, 3 H), 1.33 (t, *J*=7.0 Hz, 3 H), 1.14 - 1.27 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.99 (d, *J*=6.6 Hz, 3 H)
92-50	563.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.53 (d, *J*=4.8 Hz, 1 H), 8.40 (d, *J*=2.3 Hz, 1 H), 8.30 (dd, *J*=2.9, 1.2 Hz, 1 H), 7.56 (dd, *J*=5.0, 2.9 Hz, 1 H), 7.30 (br d, *J*=4.6 Hz, 1 H), 7.16 (dd, *J*=5.1, 1.1 Hz, 1 H), 6.83 (td, *J*=15.8, 10.5 Hz, 1 H), 6.18 (br d, *J*=16.6 Hz, 1 H), 5.70 - 5.82 (m, 1 H), 4.81 - 4.95 (m, 1 H), 4.39 - 4.78 (m, 1 H), 3.77 - 4.22 (m, 4 H),

		2.64 - 2.75 (m, 1 H), 1.95 (s, 3 H), 1.32 (t, *J*=7.3 Hz, 3 H), 1.13 - 1.26 (m, 3 H), 1.09 (d, *J*=6.8 Hz, 3 H), 0.96 (d, *J*=6.6 Hz, 3 H)
92-51	591.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (d, *J*=3.3 Hz, 1 H), 8.37 (d, *J*=4.8 Hz, 1 H), 7.51 - 7.56 (m, 1 H), 7.37 - 7.50 (m, 2 H), 7.23 (dd, *J*=7.5, 1.7 Hz, 1 H), 7.18 (d, *J*=5.0 Hz, 1 H), 6.83 (td, *J*=17.1, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.70 - 5.83 (m, 1 H), 4.44 - 4.96 (m, 2 H), 3.72 - 4.23 (m, 4 H), 2.65 - 2.74 (m, 1 H), 1.92 (s, 3 H), 1.30 - 1.37 (m, 3 H), 1.25 (br dd, *J*=6.3, 3.0 Hz, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.95 (d, *J*=6.4 Hz, 3 H)
92-52	575.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.36 - 8.56 (m, 2 H), 7.44 - 7.55 (m, 1 H), 7.39 (d, *J*=7.9 Hz, 1 H), 7.16 - 7.36 (m, 3 H), 6.84 (td, *J*=16.5, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.1 Hz, 1 H), 5.69 - 5.83 (m, 1 H), 4.82 - 4.95 (m, 1 H), 4.44 - 4.81 (m, 1 H), 3.45 - 4.20 (m, 4 H), 2.65 - 2.75 (m, 1 H), 1.94 (s, 3 H), 1.33 (t, *J*=6.9 Hz, 3 H), 1.12 - 1.26 (m, 3 H), 1.08 (d, *J*=6.6 Hz, 3 H), 0.98 (d, *J*=6.6 Hz, 3 H)
92-53	571.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.40 - 8.49 (m, 2 H), 7.25 - 7.34 (m, 4 H), 7.22 (d, *J*=5.0 Hz, 1 H), 6.84 (td, *J*=16.3, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.5, 2.0 Hz, 1 H), 5.70 - 5.81 (m, 1 H), 4.81 - 4.96 (m, 1 H), 4.40 - 4.79 (m, 1 H), 3.43 - 4.25 (m, 4 H), 2.64 - 2.78 (m, 1 H), 2.27 (s, 3 H), 1.93 (s, 3 H), 1.33 (t, *J*=7.2 Hz, 3 H), 1.13 - 1.26 (m, 3 H), 1.08 (d, *J*=6.6 Hz, 3 H), 1.01 (d, *J*=6.6 Hz, 3 H)
92-54	591.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.37 - 8.51 (m, 2 H), 7.40 - 7.57 (m, 4 H), 7.23 (d, *J*=5.0 Hz, 1 H), 6.84 (td, *J*=16.3, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.66 - 5.82 (m, 1 H), 4.82 - 4.97 (m, 1 H), 4.42 - 4.81 (m, 1 H), 3.45 - 4.22 (m, 4 H), 2.64 - 2.78 (m, 1 H), 1.93 (s, 3 H), 1.32 (br t, *J*=7.0 Hz, 3 H), 1.14 - 1.25 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 1.00 (d, *J*=6.8 Hz, 3 H)
92-55	585.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.43 (d, *J*=4.1 Hz, 1 H), 8.36 (d, *J*=4.8 Hz, 1 H), 7.31 - 7.41 (m, 1 H), 7.20 - 7.30 (m, 2 H), 7.17 (d, *J*=5.0 Hz, 1 H), 7.06 (d, *J*=6.8 Hz, 1 H), 6.84 (td, *J*=17.4, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.69 - 5.83 (m, 1 H), 4.43 - 4.97 (m, 2 H), 3.41 - 4.31 (m, 4 H), 2.64 - 2.79 (m, 1 H), 2.17 - 2.38 (m, 2 H), 1.90 (s, 3 H), 1.35 (br t, *J*=5.8 Hz, 3 H), 1.17 - 1.29 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.92 (d, *J*=6.6 Hz, 3 H), 0.82 (t, *J*=7.5 Hz, 3 H)
92-56	625.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.50 (d, *J*=2.5 Hz, 1 H), 8.45 (d, *J*=5.0 Hz, 1 H), 7.92 (d, *J*=7.9 Hz, 1 H), 7.83 (d, *J*=7.9 Hz, 1 H), 7.76 (s, 1 H), 7.65 - 7.74 (m, 1 H), 7.22 (d, *J*=5.0 Hz, 1 H), 6.84 (td, *J*=16.4, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.1 Hz, 1 H), 5.70 - 5.80 (m, 1 H), 4.83 - 4.98 (m, 1 H), 4.40 - 4.81 (m, 1 H), 3.43 - 4.20 (m, 4 H), 2.72 (td, *J*=6.6, 2.4 Hz, 1 H), 1.93 (s, 3 H), 1.28 - 1.38 (m, 3 H), 1.13 - 1.26 (m, 3 H), 1.08 (d, *J*=6.6 Hz, 3 H), 1.00 (d, *J*=6.6 Hz, 3 H)

92-57	575.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.42 - 8.48 (m, 2 H), 7.57 (dd, *J*=8.7, 5.4 Hz, 2 H), 7.28 (t, *J*=8.9 Hz, 2 H), 7.23 (d, *J*=4.8 Hz, 1 H), 6.84 (td, *J*=16.3, 10.7 Hz, 1 H), 6.19 (dd, *J*=16.6, 1.7 Hz, 1 H), 5.66 - 5.83 (m, 1 H), 4.82 - 4.99 (m, 1 H), 4.38 - 4.80 (m, 1 H), 3.69 - 4.22 (m, 4 H), 2.66 - 2.76 (m, 1 H), 1.94 (s, 3 H), 1.33 (br t, *J*=7.0 Hz, 3 H), 1.14 - 1.26 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.97 (d, *J*=6.6 Hz, 3 H)
92-58	589.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 (d, *J*=3.3 Hz, 1 H), 8.39 (d, *J*=4.8 Hz, 1 H), 7.38 (br t, *J*=7.3 Hz, 1 H), 7.19 (d, *J*=5.0 Hz, 1 H), 7.15 (t, *J*=7.6 Hz, 1 H), 7.01 (br t, *J*=6.6 Hz, 1 H), 6.84 (td, *J*=16.7, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.70 - 5.81 (m, 1 H), 4.81 - 4.93 (m, 1 H), 4.42 - 4.81 (m, 1 H), 3.44 - 4.23 (m, 4 H), 2.63 - 2.76 (m, 1 H), 2.24 (s, 3 H), 1.93 (s, 3 H), 1.33 (t, *J*=6.6 Hz, 3 H), 1.14 - 1.27 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.97 (d, *J*=6.8 Hz, 3 H)
92-59	589.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 (d, *J*=3.7 Hz, 1 H), 8.38 (d, *J*=4.8 Hz, 1 H), 7.20 - 7.33 (m, 2 H), 7.19 (d, *J*=5.2 Hz, 1 H), 6.95 (d, *J*=7.5 Hz, 1 H), 6.84 (td, *J*=16.9, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.68 - 5.83 (m, 1 H), 4.82 - 4.93 (m, 1 H), 4.44 - 4.80 (m, 1 H), 3.44 - 4.26 (m, 4 H), 2.68 - 2.78 (m, 1 H), 1.92 (s, 3 H), 1.87 (s, 3 H), 1.34 (br t, *J*=6.1 Hz, 3 H), 1.16 - 1.29 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.93 (d, *J*=6.6 Hz, 3 H)
92-60	589.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 (d, *J*=2.5 Hz, 1 H), 8.40 (d, *J*=4.8 Hz, 1 H), 7.26 - 7.39 (m, 1 H), 7.12 - 7.25 (m, 2 H), 7.00 (br d, *J*=6.6 Hz, 1 H), 6.83 (td, *J*=16.3, 10.6 Hz, 1 H), 6.18 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.65 - 5.82 (m, 1 H), 4.85 (br d, *J*=6.2 Hz, 1 H), 4.42 - 4.79 (m, 1 H), 3.53 - 4.22 (m, 4 H), 2.62 - 2.77 (m, 1 H), 2.24 (s, 3 H), 1.93 (s, 3 H), 1.32 (br t, *J*=6.8 Hz, 3 H), 1.14 - 1.25 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.99 (br d, *J*=6.6 Hz, 3 H)
92-61	593.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.47 (d, *J*=3.7 Hz, 1 H), 8.41 (d, *J*=5.0 Hz, 1 H), 7.34 - 7.42 (m, 1 H), 7.30 (td, *J*=8.3, 6.5 Hz, 1 H), 7.13 - 7.24 (m, 2 H), 6.83 (td, *J*=16.9, 10.4 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.69 - 5.81 (m, 1 H), 4.42 - 4.94 (m, 2 H), 3.42 - 4.27 (m, 4 H), 2.65 - 2.77 (m, 1 H), 1.93 (s, 3 H), 1.33 (br t, *J*=6.4 Hz, 3 H), 1.15 - 1.28 (m, 3 H), 1.04 - 1.09 (m, 3 H), 0.96 (d, *J*=6.6 Hz, 3 H)
92-62	605.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.44 (d, *J*=3.5 Hz, 1 H), 8.38 (d, *J*=4.8 Hz, 1 H), 7.35 (ddd, *J*=11.9, 8.3, 1.3 Hz, 1 H), 7.18 (d, *J*=5.0 Hz, 1 H), 7.14 (td, *J*=8.0, 5.0 Hz, 1 H), 6.92 (d, *J*=7.7 Hz, 1 H), 6.84 (td, *J*=16.9, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.71 - 5.80 (m, 1 H), 4.45 - 4.94 (m, 2 H), 3.76 - 4.26 (m, 4 H), 3.55 (s, 3 H), 2.65 - 2.74 (m, 1 H), 1.93 (s, 3 H), 1.34 (t, *J*=6.5 Hz, 3 H), 1.16 - 1.28 (m, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.93 (d, *J*=6.6 Hz, 3 H)

92-63	643.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.45 - 8.54 (m, 1 H), 8.37 (d, *J*=5.0 Hz, 1 H), 7.76 - 7.88 (m, 1 H), 7.56 - 7.68 (m, 1 H), 7.12 - 7.22 (m, 2 H), 6.75 - 6.91 (m, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.76 (ddd, *J*=10.3, 5.4, 2.2 Hz, 1 H), 4.44 - 4.95 (m, 2 H), 3.74 - 4.26 (m, 4 H), 2.64 - 2.79 (m, 1 H), 1.82 - 1.95 (m, 3 H), 1.30 - 1.39 (m, 3 H), 1.18 - 1.28 (m, 3 H), 1.07 (d, *J*=6.0 Hz, 3 H), 0.86 - 0.93 (m, 3 H)
92-64	609.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.51 (d, *J*=3.3 Hz, 1 H), 8.38 (d, *J*=4.8 Hz, 1 H), 7.43 - 7.56 (m, 2 H), 7.19 (d, *J*=4.8 Hz, 1 H), 7.11 (br d, *J*=7.0 Hz, 1 H), 6.74 - 6.93 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.70 - 5.80 (m, 1 H), 4.45 - 4.93 (m, 2 H), 3.44 - 4.23 (m, 4 H), 2.64 - 2.82 (m, 1 H), 1.92 (s, 3 H), 1.34 (br t, *J*=6.2 Hz, 3 H), 1.17 - 1.28 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.94 (br d, *J*=6.6 Hz, 3 H)
92-65	593.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (d, *J*=3.5 Hz, 1 H), 8.41 (d, *J*=5.0 Hz, 1 H), 7.34 - 7.44 (m, 2 H), 7.21 (d, *J*=5.0 Hz, 1 H), 7.03 - 7.14 (m, 1 H), 6.83 (td, *J*=16.8, 10.7 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.66 - 5.84 (m, 1 H), 4.40 - 4.96 (m, 2 H), 3.43 - 4.24 (m, 4 H), 2.64 - 2.76 (m, 1 H), 1.94 (s, 3 H), 1.33 (t, *J*=6.6 Hz, 3 H), 1.15 - 1.27 (m, 3 H), 1.06 (d, *J*=6.4 Hz, 3 H), 0.96 (d, *J*=6.6 Hz, 3 H)
92-66	593.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.50 (d, *J*=3.9 Hz, 1 H), 8.41 (d, *J*=4.8 Hz, 1 H), 7.49 - 7.64 (m, 1 H), 7.27 - 7.36 (m, 1 H), 7.21 (d, *J*=4.8 Hz, 1 H), 7.07 (br t, *J*=6.8 Hz, 1 H), 6.84 (td, *J*=17.0, 10.6 Hz, 1 H), 6.20 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.69 - 5.84 (m, 1 H), 4.88 (br dd, *J*=15.2, 6.5 Hz, 1 H), 4.43 - 4.81 (m, 1 H), 3.46 - 4.25 (m, 4 H), 2.65 - 2.79 (m, 1 H), 1.94 (s, 3 H), 1.34 (br t, *J*=6.4 Hz, 3 H), 1.14 - 1.29 (m, 3 H), 1.07 (d, *J*=6.0 Hz, 3 H), 0.97 (d, *J*=6.6 Hz, 3 H)
92-67	587.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.39 (d, *J*=4.6 Hz, 2 H), 7.37 - 7.50 (m, 1 H), 7.19 (d, *J*=5.0 Hz, 1 H), 7.10 (d, *J*=8.3 Hz, 1 H), 6.93 - 7.05 (m, 2 H), 6.84 (td, *J*=17.2, 10.6 Hz, 1 H), 6.20 (dd, *J*=16.7, 1.6 Hz, 1 H), 5.68 - 5.83 (m, 1 H), 4.44 - 4.94 (m, 2 H), 3.80 - 4.23 (m, 4 H), 3.70 (s, 3 H), 2.64 - 2.78 (m, 1 H), 1.94 (s, 3 H), 1.34 (br t, *J*=6.5 Hz, 3 H), 1.16 - 1.30 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.97 (br d, *J*=6.2 Hz, 3 H)
92-68	601.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 9.18 (s, 1 H), 8.58 (s, 1 H), 8.50 (d, *J*=6.2 Hz, 1 H), 8.38 (d, *J*=5.0 Hz, 1 H), 7.18 (d, *J*=5.0 Hz, 1 H), 6.72 - 6.94 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.69 - 5.84 (m, 1 H), 4.43 - 4.92 (m, 2 H), 3.42 - 4.34 (m, 4 H), 2.63 - 2.80 (m, 2 H), 1.91 (s, 3 H), 1.36 (dd, *J*=6.3, 3.4 Hz, 3 H), 1.19 - 1.31 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.97 (br d, *J*=6.4 Hz, 3 H), 0.90 (br d, *J*=4.6 Hz, 6 H)
92-69	598.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (d, *J*=3.1 Hz, 1 H), 8.42 - 8.46 (m, 1 H), 8.38 (d, *J*=4.8 Hz, 1 H), 7.44 (dd,

		J=7.7, 1.5 Hz, 1 H), 7.12 - 7.25 (m, 2 H), 6.84 (td, *J*=17.1, 10.4 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.1 Hz, 1 H), 5.76 (br dd, *J*=10.6, 2.7 Hz, 1 H), 4.82 - 4.96 (m, 1 H), 4.43 - 4.81 (m, 1 H), 3.77 - 4.25 (m, 4 H), 2.62 - 2.78 (m, 1 H), 1.93 (s, 3 H), 1.55 - 1.66 (m, 1 H), 1.34 (br t, *J*=6.3 Hz, 3 H), 1.17 - 1.28 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.92 (br d, *J*=6.6 Hz, 3 H), 0.89 (br s, 2 H), 0.67 - 0.79 (m, 2 H)
92-70	637.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (d, *J*=2.5 Hz, 1 H), 8.30 (d, *J*=4.8 Hz, 1 H), 7.87 (d, *J*=8.3 Hz, 1 H), 7.46 (dt, *J*=8.2, 4.1 Hz, 1 H), 7.42 (d, *J*=2.5 Hz, 1 H), 7.26 (d, *J*=3.7 Hz, 2 H), 7.11 (d, *J*=5.0 Hz, 1 H), 6.96 (d, *J*=2.7 Hz, 1 H), 6.85 (td, *J*=17.2, 10.6 Hz, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 4.84 - 5.00 (m, 1 H), 4.44 - 4.83 (m, 1 H), 3.97 - 4.29 (m, 2 H), 3.87 - 3.96 (m, 2 H), 3.86 (s, 3 H), 2.69 - 2.80 (m, 1 H), 1.93 (s, 3 H), 1.32 - 1.39 (m, 3 H), 1.19 - 1.31 (m, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.91 (d, *J*=6.6 Hz, 3 H)
92-71	592.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.44 (d, *J*=5.0 Hz, 1 H), 8.38 (d, *J*=5.2 Hz, 1 H), 7.23 (d, *J*=5.0 Hz, 1 H), 6.84 (td, *J*=17.4, 10.5 Hz, 1 H), 6.19 (br d, *J*=17.0 Hz, 1 H), 5.71 - 5.81 (m, 1 H), 4.42 - 4.96 (m, 2 H), 3.38 - 4.30 (m, 4 H), 2.65 - 2.78 (m, 1 H), 2.57 (s, 3 H), 1.92 (s, 6 H), 1.34 (t, *J*=6.1 Hz, 3 H), 1.14 - 1.30 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.92 (d, *J*=6.6 Hz, 3 H)
92-72	578.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 9.11 (s, 1 H), 8.41 - 8.48 (m, 2 H), 7.25 (d, *J*=5.0 Hz, 1 H), 6.84 (td, *J*=17.1, 10.8 Hz, 1 H), 6.18 (br d, *J*=16.6 Hz, 1 H), 5.68 - 5.81 (m, 1 H), 4.80 - 4.95 (m, 1 H), 4.39 - 4.80 (m, 1 H), 3.42 - 4.24 (m, 4 H), 2.63 - 2.76 (m, 1 H), 2.01 (s, 3 H), 1.95 (s, 3 H), 1.33 (t, *J*=6.4 Hz, 3 H), 1.13 - 1.28 (m, 3 H), 1.06 (d, *J*=6.2 Hz, 3 H), 0.92 (d, *J*=6.6 Hz, 3 H)
93-1	577.3	^{1}H NMR (400 MHz, *DMSO-d_6*) δ 8.45 (d, *J*=4.8 Hz, 1 H), 8.39 (d, *J*=3.9 Hz, 1 H), 7.66 (d, *J*=5.0 Hz, 1 H), 7.23 (d, *J*=5.0 Hz, 1 H), 6.94 (d, *J*=5.2 Hz, 1 H), 6.83 (td, *J*=17.2, 10.7 Hz, 1 H), 6.19 (dd, *J*=16.7, 1.3 Hz, 1 H), 5.71 - 5.78 (m, 1 H), 4.41 - 4.94 (m, 2 H), 3.42 - 4.23 (m, 4 H), 2.63 - 2.74 (m, 1 H), 1.94 (s, 3 H), 1.83 (s, 3 H), 1.33 (t, *J*=6.5 Hz, 3 H), 1.21 (br dd, *J*=27.7, 6.5 Hz, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.94 (d, *J*=6.6 Hz, 3 H)
93-2	549.2	^{1}H NMR (400 MHz, *DMSO-d_6*) δ 8.52 - 8.57 (m, 1 H), 8.21 - 8.27 (m, 1 H), 7.34 - 7.43 (m, 1 H), 6.75 - 6.90 (m, 1 H), 6.18 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.71 - 5.78 (m, 1 H), 3.34 - 4.91 (m, 6 H), 2.70 - 2.83 (m, 1 H), 2.13 - 2.35 (m, 1 H), 1.81 - 1.97 (m, 3 H), 1.28 - 1.35 (m, 3 H), 1.14 - 1.27 (m, 6 H), 1.07 - 1.13 (m, 3 H), 1.01 (dd, *J*=11.8, 6.6 Hz, 3 H), 0.79 - 0.90 (m, 1 H), 0.65 - 0.72 (m, 1 H), 0.62 (s, 3 H)
94-1	582.8	^{1}H NMR (400 MHz, *DMSO-d_6*) δ 8.37 (d, *J*=4.8 Hz, 1 H), 8.30 (dd, *J*=9.4, 3.2 Hz, 1 H), 7.38 - 7.44 (m, 2 H), 7.21 - 7.27 (m, 1 H), 7.18 (d, *J*=5.0 Hz, 1 H), 7.12 (d, *J*=7.7 Hz,

		1 H), 6.77 - 6.92 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.73 - 5.78 (m, 1 H), 4.82 - 4.93 (m, 1 H), 4.43 - 4.81 (m, 1 H), 3.44 - 4.27 (m, 4 H), 2.69 (quin, *J*=6.4 Hz, 2 H), 1.92 (s, 3 H), 1.33 (t, *J*=6.3 Hz, 3 H), 1.18 - 1.30 (m, 3 H), 0.87 - 1.09 (m, 12 H)
94-2	571.0	^{1}H NMR (400 MHz, DMSO-d6) δ 0.88 - 0.99 (m, 3 H) 1.03 - 1.10 (m, 3 H) 1.18 - 1.29 (m, 3 H) 1.30 - 1.37 (m, 3 H) 1.90 - 1.96 (m, 3 H) 2.64 - 2.77 (m, 1 H) 3.72 - 4.25 (m, 6 H) 4.39 - 4.89 (m, 2 H) 4.90 - 4.96 (m, 1 H) 5.70 - 5.85 (m, 1 H) 6.14 - 6.25 (m, 1 H) 6.74 - 6.93 (m, 1 H) 7.16 - 7.23 (m, 1 H) 7.24 - 7.36 (m, 2 H) 7.41 - 7.49 (m, 1 H) 7.51 - 7.57 (m, 1 H) 8.21 - 8.33 (m, 1 H) 8.36 - 8.46 (m, 1 H) ^{19}F NMR (376 MHz, DMSO-d6) δ -129.76 (br d, J=6.07 Hz, 1 F)
94-3	577.1	^{1}H NMR (400MHz, DMSO-d_6) δ = 8.52 - 8.36 (m, 2H), 7.60 - 7.38 (m, 3H), 7.23 (d, *J*=4.8 Hz, 1H), 7.13 - 7.03 (m, 1H), 6.90 - 6.77 (m, 1H), 6.19 (dd, *J*=2.2, 16.5 Hz, 1H), 5.79 - 5.71 (m, 1H), 4.87 (br s, 1H), 4.25 - 4.08 (m, 2H), 3.85 (br s, 2H), 1.99 - 1.92 (m, 4H), 1.31 (br t, *J*=7.4 Hz, 3H), 1.27 - 1.14 (m, 3H), 1.07 (d, *J*=6.8 Hz, 3H), 0.97 (d, *J*=6.6 Hz, 3H); ^{19}F NMR (376MHz, DMSO-d6) δ = -118.05 (dd, J=3.0, 17.8 Hz, 1F), -118.95 (br dd, J=18.2, 32.9 Hz, 1F), -128.78 (d, 1F)
94-4	573.0	^{1}H NMR (400MHz, DMSO-d6) δ = 8.44 (d, J=5.0 Hz, 1H), 8.34 (br d, J=9.5 Hz, 1H), 7.37 - 7.32 (m, 1H), 7.26 - 7.19 (m, 2H), 7.06 (br d, J=5.0 Hz, 1H), 6.81 (br dd, J=10.4, 17.0 Hz, 1H), 6.21 (s, 1H), 5.75 (s, 2H), 4.93 - 4.39 (m, 2H), 4.14 (br t, J=12.4 Hz, 1H), 3.85 (br s, 2H), 2.26 - 2.20 (m, 3H), 1.94 (s, 3H), 1.34 - 1.21 (m, 5H), 1.17 (br d, J=6.8 Hz, 2H), 1.08 (d, J=6.8 Hz, 3H), 1.01 (d, J=6.6 Hz, 3H); ^{19}F NMR (376MHz, DMSO-d6) δ = -118.76 (s, 1F), -128.99 (s, 1F)
94-5	593.0	^{1}H NMR (400MHz, DMSO-d6) δ = 8.46 - 8.37 (m, 2H), 7.63 (ddd, J=2.9, 4.3, 8.8 Hz, 1H), 7.43 (t, J=9.3 Hz, 1H), 7.30 (dd, J=2.7, 5.8 Hz, 1H), 7.23 (d, J=5.0 Hz, 1H), 6.90 - 6.77 (m, 1H), 6.19 (dd, J=2.3, 16.6 Hz, 1H), 5.78 - 5.73 (m, 2H), 4.94 - 4.78 (m, 1H), 4.13 (m, 2H), 3.91 - 3.82 (m, 2H), 2.75 - 2.68 (m, 1H), 1.94 (s, 3H), 1.34 - 1.28 (m, 3H), 1.24 (br d, J=6.6 Hz, 3H), 1.07 (d, J=6.6 Hz, 3H), 1.00 (d, J=6.6 Hz, 3H); ^{19}F NMR (376MHz, DMSO-d6) δ = -115.61 (s, 1F), -128.78 (s, 1F)
94-6	605.0	^{1}H NMR (400MHz, CHLOROFORM-d) δ = 8.49 (d, *J*=5.0 Hz, 1H), 7.81 (br d, *J*=8.5 Hz, 1H), 7.28 (d, *J*=8.7 Hz, 1H), 7.08 (br d, *J*=5.0 Hz, 1H), 6.88 (dd, *J*=2.9, 8.9 Hz, 1H), 6.72 - 6.50 (m, 2H), 6.43 - 6.33 (m, 1H), 5.82 - 5.74 (m, 1H), 5.19 - 4.83 (m, 2H), 4.55 - 4.25 (m, 1H), 4.08 - 3.75 (m, 3H), 3.68 (s, 3H), 2.82 - 2.61 (m, 1H), 2.03 (br s, 3H), 1.33 - 1.25 (m, 3H), 1.25 - 1.18 (m, 6H), 1.13 -

		1.00 (m, 3H); ^{19}F NMR (376MHz, CDCl3) δ = -125.80 (s, 1F)
94-7	598.6	^{1}H NMR (400MHz, DMSO-d_6) δ = 8.46 (d, *J*=4.8 Hz, 1H), 8.34 (br d, *J*=9.7 Hz, 1H), 7.33 - 7.17 (m, 3H), 6.90 - 6.77 (m, 2H), 6.19 (dd, *J*=2.1, 16.6 Hz, 1H), 5.78 - 5.73 (m, 1H), 4.98 - 4.42 (m, 2H), 4.14 (br t, *J*=13.1 Hz, 2H), 3.92 - 3.81 (m, 2H), 2.75 - 2.61 (m, 1H), 1.94 (s, 3H), 1.91 - 1.82 (m, 1H), 1.37 - 1.13 (m, 8H), 1.08 (d, *J*=6.6 Hz, 3H), 1.01 (d, *J*=6.6 Hz, 3H), 0.93 (dd, *J*=1.8, 8.2 Hz, 2H); ^{19}F NMR (376MHz, DMSO-d_6) δ = -117.96 (s, 1F), -128.53 (s, 1F)
94-8	555.4	^{1}H NMR (400 MHz, $CDCl_3$) δ 8.49 (d, *J* = 5.0 Hz, 1 H) 7.84 (br d, *J* = 9.3 Hz, 1 H) 7.29 - 7.35 (m, 1 H) 7.18 - 7.26 (m, 3 H) 7.08 (d, *J* = 5.0 Hz, 1 H) 6.51 - 6.71 (m, 1 H) 6.40 (br t, *J* = 15.1 Hz, 1 H) 5.76 - 5.84 (m, 1 H) 4.86 - 5.19 (m, 2 H) 4.35 - 4.53 (m, 1 H) 3.88 - 4.13 (m, 2 H) 3.68 - 3.83 (m, 1 H) 2.64 - 2.77 (m, 1 H) 2.01 - 2.05 (m, 6 H) 1.32 - 1.50 (m, 6 H) 1.22 (d, *J* = 6.6 Hz, 3 H) 0.98 - 1.05 (m, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -127.82 - -127-.88 (m, 1 F)
94-9	569.4	^{1}H NMR (400 MHz, $CDCl_3$) δ 8.46 (d, *J* = 5.0 Hz, 1 H) 7.86 (d, *J* = 8.5 Hz, 1 H) 7.16 - 7.23 (m, 1 H) 6.99 - 7.08 (m, 3 H) 6.51 - 6.72 (m, 1 H) 6.40 (br t, *J* = 15.1 Hz, 1 H) 5.76 - 5.86 (m, 1 H) 3.67 - 5.17 (m, 6 H) 2.66 - 2.80 (m, 1 H) 1.96 - 2.02 (m, 6 H) 1.89 (s, 3 H) 1.46 (br d, *J* = 6.6 Hz, 3 H) 1.33 - 1.52 (m, 3 H) 1.23 (d, *J* = 6.8 Hz, 3 H) 0.98 - 1.06 (m, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -128.31 (s, 1 F)
94-10	573.4	^{1}H NMR (400 MHz, $CDCl_3$) δ 8.48 (d, *J* = 4.8 Hz, 1 H) 7.87 (d, *J* = 8.5 Hz, 1 H) 7.27 - 7.32 (m, 1 H) 7.08 (d, *J* = 5.0 Hz, 1 H) 6.99 (d, *J* = 7.5 Hz, 1 H) 6.94 (t, *J* = 8.8 Hz, 1 H) 6.52 - 6.72 (m, 1 H) 6.40 (br t, *J* = 14.9 Hz, 1 H) 5.76 - 5.85 (m, 1 H) 3.67 - 5.18 (m, 6 H) 2.67 - 2.79 (m, 1 H) 1.99 - 2.03 (m, 3 H) 1.97 (br s, 3 H) 1.46 (br d, *J* = 6.6 Hz, 3 H) 1.34 - 1.53 (m, 3 H) 1.22 (d, *J* = 6.8 Hz, 3 H) 1.01 (br s, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -115.74 - -115.69 (m, 1 F) -127.11 - -127.73 (m, 1 F)
94-11	577.4	^{1}H NMR (400 MHz, $CDCl_3$) δ 8.40 (d, *J* = 4.8 Hz, 1 H) 7.79 (d, *J* = 8.3 Hz, 1 H) 7.25 - 7.36 (m, 1 H) 6.99 (d, *J* = 5.0 Hz, 1 H) 6.85 (t, *J* = 8.3 Hz, 2 H) 6.42 - 6.62 (m, 1 H) 6.31 (br t, *J* = 14.9 Hz, 1 H) 5.66 - 5.76 (m, 1 H) 3.56 - 5.08 (m, 6 H) 2.55 - 2.70 (m, 1 H) 1.90 - 1.94 (m, 3 H) 1.24 - 1.43 (m, 6 H) 1.14 (d, *J* = 6.8 Hz, 3 H) 0.93 - 0.99 (m, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.48 (br s, 1 F) -127.51 (q, *J* = 11.3 Hz, 1 F)
94-12	593.2	^{1}H NMR (400 MHz, $CDCl_3$) δ 8.48 (d, *J* = 5.0 Hz, 1 H) 7.89 (d, *J* = 8.3 Hz, 1 H) 7.35 (td, *J* = 8.3, 6.0 Hz, 1 H) 7.23 (d, *J* = 8.1 Hz, 1 H) 7.00 - 7.11 (m, 2 H) 6.51 - 6.72 (m, 1 H) 6.40 (br t, *J* = 15.0 Hz, 1 H) 5.76 - 5.85 (m, 1 H) 4.45 - 5.18 (m, 2 H) 3.68 - 4.45 (m, 4 H) 2.66 - 2.79 (m, 1

		H) 2.01 (br s, 3 H) 1.46 (br d, *J* = 6.4 Hz, 3 H) 1.33 - 1.52 (m, 3 H) 1.23 (d, *J* = 6.8 Hz, 3 H) 1.03 (br d, *J* = 5.2 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -111.80 (br d, *J* = 221.9 Hz, 1 F) -128.23 - -127.28 (m, 1 F)
94-13	591.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.41 (d, *J*=4.8 Hz, 1 H), 8.36 (dd, *J*=9.7, 3.9 Hz, 1 H), 7.58 - 7.71 (m, 3 H), 7.52 (br d, *J*=6.0 Hz, 1 H), 7.22 (d, *J*=5.0 Hz, 1 H), 6.78 - 6.94 (m, 1 H), 6.44 - 6.77 (m, 1 H), 6.19 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.71 - 5.81 (m, 1 H), 4.85 (dt, *J*=6.4, 3.4 Hz, 1 H), 4.44 - 4.81 (m, 1 H), 3.43 - 4.28 (m, 4 H), 2.64 - 2.77 (m, 1 H), 1.93 (s, 3 H), 1.33 (t, *J*=5.9 Hz, 3 H), 1.17 - 1.30 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.89 (br d, *J*=6.6 Hz, 3 H)
94-14	580.6	^{1}H NMR (400 MHz, DMSO-d_6) δ 10.19 (s, 1 H), 8.75 (d, *J*=4.8 Hz, 1 H), 8.27 - 8.48 (m, 1 H), 7.79 - 8.02 (m, 2 H), 6.99 - 7.70 (m, 3 H), 6.73 - 6.95 (m, 1 H), 6.11 - 6.27 (m, 1 H), 5.70 - 5.80 (m, 1 H), 4.91 (br s, 1 H), 4.43 - 4.84 (m, 1 H), 3.46 - 4.29 (m, 4 H), 2.70 - 2.85 (m, 1 H), 1.92 - 2.08 (m, 3 H), 1.34 (br t, *J*=7.0 Hz, 3 H), 1.16 - 1.28 (m, 3 H), 1.06 - 1.15 (m, 3 H), 0.80 - 1.05 (m, 3 H)
94-15	585.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 12.94 (br s, 1 H), 8.37 (d, *J*=5.0 Hz, 1 H), 8.22 (br d, *J*=9.5 Hz, 1 H), 7.91 (d, *J*=7.3 Hz, 1 H), 7.50 - 7.69 (m, 2 H), 7.14 - 7.32 (m, 2 H), 6.84 (ddd, *J*=19.0, 17.0, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.67 - 5.81 (m, 1 H), 4.40 - 4.95 (m, 2 H), 4.02 - 4.23 (m, 1 H), 3.77 - 3.93 (m, 2 H), 3.51 (br dd, *J*=13.4, 2.4 Hz, 1 H), 2.62 - 2.75 (m, 1 H), 1.92 (d, *J*=12.9 Hz, 3 H), 1.30 (br dd, *J*=9.3, 6.8 Hz, 3 H), 1.14 - 1.26 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.91 (br d, *J*=6.6 Hz, 3 H)
94-16	566.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.53 - 8.62 (m, 1 H), 8.40 (s, 1 H), 8.27 (d, *J*=7.7 Hz, 1 H), 7.90 - 8.01 (m, 1 H), 7.80 - 7.87 (m, 1 H), 7.74 (d, *J*=7.9 Hz, 1 H), 7.47 - 7.56 (m, 2 H), 7.33 (d, *J*=5.0 Hz, 1 H), 6.70 - 6.97 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.65 - 5.82 (m, 1 H), 4.64 - 4.99 (m, 2 H), 3.79 - 4.32 (m, 4 H), 2.62 - 2.79 (m, 1 H), 1.97 (s, 3 H), 1.32 (dd, *J*=10.6, 6.6 Hz, 3 H), 1.13 - 1.25 (m, 3 H), 1.08 (d, *J*=6.6 Hz, 3 H), 0.94 (d, *J*=6.6 Hz, 3 H)
94-17	584.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.38 (d, *J*=4.8 Hz, 1 H), 8.17 (d, *J*=9.5 Hz, 1 H), 7.86 (s, 1 H), 7.62 - 7.69 (m, 1 H), 7.47 - 7.55 (m, 2 H), 7.20 - 7.28 (m, 1 H), 7.10 - 7.20 (m, 2 H), 6.73 - 6.93 (m, 1 H), 6.18 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.71 - 5.79 (m, 1 H), 4.44 - 4.92 (m, 2 H), 3.43 - 4.25 (m, 4 H), 2.68 (td, *J*=6.3, 3.8 Hz, 1 H), 1.94 (s, 3 H), 1.31 (t, *J*=6.7 Hz, 3 H), 1.17 - 1.27 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.92 (br d, *J*=6.6 Hz, 3 H)
94-18	598.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.38 (d, *J*=4.8 Hz, 1 H), 8.32 (br d, *J*=4.6 Hz, 1 H), 8.19 (d, *J*=9.5 Hz, 1 H), 7.56 - 7.64 (m, 1 H), 7.47 - 7.55 (m, 2 H), 7.16 - 7.25 (m, 2 H), 6.73 - 6.94 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.69 - 5.82 (m, 1 H), 4.41 - 4.93 (m, 2 H), 3.43 - 4.23 (m, 4 H), 2.60 - 2.74 (m, 4 H), 1.93 (s, 3 H), 1.31 (br t, *J*=7.4 Hz, 3

		H), 1.16 - 1.28 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.89 (br d, *J*=6.4 Hz, 3 H)
94-19	602.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.36 (d, *J*=4.8 Hz, 1 H), 8.25 (br s, 1 H), 7.87 (br s, 1 H), 7.53 - 7.65 (m, 1 H), 7.36 - 7.53 (m, 2 H), 7.27 (br s, 1 H), 7.16 (br d, *J*=4.8 Hz, 1 H), 6.71 - 6.98 (m, 1 H), 6.19 (dd, *J*=16.7, 1.8 Hz, 1 H), 5.69 - 5.84 (m, 1 H), 4.40 - 4.91 (m, 2 H), 3.71 - 4.32 (m, 4 H), 2.64 - 2.78 (m, 1 H), 1.87 - 1.97 (m, 3 H), 1.34 (br d, *J*=6.0 Hz, 3 H), 1.18 - 1.29 (m, 3 H), 1.07 (br d, *J*=6.4 Hz, 3 H), 0.82 (br d, *J*=6.0 Hz, 3 H)
94-20	556.4	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.51 (d, *J*=5.0 Hz, 1 H), 8.24 (dd, *J*=11.8, 3.1 Hz, 1 H), 7.51 (dd, *J*=8.1, 1.7 Hz, 1 H), 7.28 (d, *J*=5.0 Hz, 1 H), 7.05 - 7.13 (m, 1 H), 6.77 - 6.92 (m, 1 H), 6.68 (d, *J*=7.9 Hz, 1 H), 6.56 (t, *J*=7.6 Hz, 1 H), 6.19 (br d, *J*=16.8 Hz, 1 H), 5.76 (s, 3 H), 4.43 - 4.94 (m, 2 H), 4.09 - 4.20 (m, 1 H), 3.43 - 3.92 (m, 3 H), 2.64 - 2.77 (m, 1 H), 1.95 (s, 3 H), 1.32 (t, *J*=6.9 Hz, 3 H), 1.17 - 1.27 (m, 3 H), 1.08 (d, *J*=6.8 Hz, 3 H), 0.97 (d, *J*=6.6 Hz, 3 H)
94-21	621.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.31 - 8.44 (m, 2 H), 7.88 (br d, *J*=8.3 Hz, 1 H), 7.45 - 7.56 (m, 3 H), 7.27 (br t, *J*=7.7 Hz, 1 H), 7.16 (br d, *J*=4.8 Hz, 1 H), 7.09 (s, 1 H), 6.75 - 6.94 (m, 1 H), 6.20 (br d, *J*=16.8 Hz, 1 H), 5.69 - 5.80 (m, 1 H), 4.92 (br dd, *J*=6.7, 4.0 Hz, 1 H), 4.46 - 4.84 (m, 1 H), 4.14 - 4.26 (m, 1 H), 3.87 (s, 6 H), 2.74 (br d, *J*=6.0 Hz, 1 H), 1.96 (s, 3 H), 1.34 (br t, *J*=6.8 Hz, 3 H), 1.18 - 1.31 (m, 3 H), 1.07 (br d, *J*=6.6 Hz, 3 H), 0.95 (br d, *J*=6.4 Hz, 3 H)
95-1	563.3	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.48 (d, *J*=4.8 Hz, 1 H), 8.25 (d, *J*=4.1 Hz, 1 H), 7.25 (d, *J*=4.8 Hz, 1 H), 6.82 (td, *J*=17.5, 10.5 Hz, 1 H), 6.18 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.71 - 5.77 (m, 1 H), 4.40 - 4.89 (m, 2 H), 3.37 - 4.19 (m, 4 H), 3.02 (tt, *J*=11.0, 2.9 Hz, 1 H), 2.63 - 2.72 (m, 1 H), 1.86 (s, 3 H), 1.50 - 1.68 (m, 5 H), 1.13 - 1.34 (m, 9 H), 1.03 (dd, *J*=45.3, 6.7 Hz, 8 H)
95-2	551.3	^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 8.55 (br d, *J*=3.9 Hz, 1 H), 7.99 (s, 1 H), 7.15 (br s, 1 H), 6.51 - 6.72 (m, 1 H), 6.40 (br t, *J*=15.1 Hz, 1 H), 5.75 - 5.86 (m, 1 H), 4.78 - 5.18 (m, 2 H), 3.36 - 4.54 (m, 4 H), 2.47 - 2.93 (m, 3 H), 2.03 (br s, 3 H), 1.40 - 1.50 (m, 3 H), 1.24 - 1.36 (m, 6 H), 1.12 (br s, 3 H), 0.78 (s, 9 H)
95-3	549.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.47 (d, *J*=4.8 Hz, 1 H), 8.24 (d, *J*=4.1 Hz, 1 H), 7.24 (d, *J*=4.8 Hz, 1 H), 6.82 (td, *J*=17.3, 10.6 Hz, 1 H), 6.13 - 6.22 (m, 1 H), 5.70 - 5.78 (m, 1 H), 4.38 - 4.89 (m, 2 H), 3.37 - 4.19 (m, 5 H), 2.60 - 2.72 (m, 1 H), 1.87 (s, 3 H), 1.69 - 1.84 (m, 2 H), 1.27 - 1.48 (m, 7 H), 1.23 (br d, *J*=6.4 Hz, 3 H), 1.16 (d, *J*=6.6 Hz, 2 H), 1.08 (d, *J*=6.6 Hz, 3 H), 0.95 (d, *J*=6.6 Hz, 3 H)

95-4	521.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.47 (d, *J*=4.8 Hz, 1 H), 8.23 (d, *J*=3.1 Hz, 1 H), 7.24 (d, *J*=5.0 Hz, 1 H), 6.82 (td, *J*=16.8, 10.6 Hz, 1 H), 6.12 - 6.22 (m, 1 H), 5.69 - 5.79 (m, 1 H), 4.39 - 4.89 (m, 2 H), 3.38 - 4.17 (m, 4 H), 2.59 (ddd, *J*=13.3, 6.7, 3.0 Hz, 1 H), 2.37 - 2.45 (m, 1 H), 1.85 (s, 3 H), 1.26 - 1.33 (m, 3 H), 1.18 (dd, *J*=28.4, 6.6 Hz, 3 H), 1.07 (d, *J*=6.8 Hz, 3 H), 0.91 - 0.99 (m, 5 H), 0.39 - 0.49 (m, 2 H)
95-5	535.3	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.48 (d, *J*=5.0 Hz, 1 H), 8.25 (d, *J*=1.7 Hz, 1 H), 7.26 (d, *J*=4.8 Hz, 1 H), 6.75 - 6.89 (m, 1 H), 6.18 (dd, *J*=16.7, 2.0 Hz, 1 H), 5.71 - 5.77 (m, 1 H), 4.77 - 4.89 (m, 1 H), 4.39 - 4.77 (m, 1 H), 3.41 - 4.16 (m, 5 H), 2.61 - 2.70 (m, 1 H), 2.02 - 2.22 (m, 2 H), 1.91 (s, 3 H), 1.74 - 1.89 (m, 3 H), 1.52 (ddt, *J*=12.0, 7.7, 3.9, 3.9 Hz, 1 H), 1.25 - 1.33 (m, 3 H), 1.17 (br dd, *J*=30.0, 6.5 Hz, 3 H), 1.08 (d, *J*=6.6 Hz, 3 H), 0.99 (d, *J*=6.8 Hz, 3 H)
95-6	523.3	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.47 (d, *J*=5.0 Hz, 1 H), 8.27 (d, *J*=2.5 Hz, 1 H), 7.24 (d, *J*=5.0 Hz, 1 H), 6.75 - 6.89 (m, 1 H), 6.18 (dd, *J*=16.7, 1.8 Hz, 1 H), 5.71 - 5.78 (m, 1 H), 4.81 (dt, *J*=6.5, 3.4 Hz, 1 H), 4.39 - 4.77 (m, 1 H), 3.40 - 4.18 (m, 4 H), 2.60 - 2.74 (m, 3 H), 1.89 (s, 3 H), 1.39 (sxt, *J*=7.3 Hz, 2 H), 1.26 - 1.33 (m, 3 H), 1.12 - 1.23 (m, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.97 (d, *J*=6.4 Hz, 3 H), 0.72 (t, *J*=7.4 Hz, 3 H)
95-7	561.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.48 (d, *J*=4.8 Hz, 1 H), 8.39 (s, 1 H), 7.26 (d, *J*=4.8 Hz, 1 H), 6.82 (dt, *J*=16.6, 11.2 Hz, 1 H), 6.18 (dd, *J*=16.7, 2.0 Hz, 1 H), 5.71 - 5.77 (m, 1 H), 4.83 (dt, *J*=6.2, 3.2 Hz, 1 H), 4.34 - 4.75 (m, 1 H), 3.44 - 4.16 (m, 4 H), 2.53 - 2.60 (m, 1 H), 1.94 (s, 3 H), 1.22 - 1.31 (m, 12 H), 1.07 - 1.18 (m, 3 H), 1.04 (d, *J*=6.6 Hz, 6 H)
96-1	572.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.47 (d, *J*=4.8 Hz, 1 H), 8.41 (d, *J*=3.9 Hz, 1 H), 8.38 (d, *J*=5.0 Hz, 1 H), 7.70 (d, *J*=7.7 Hz, 1 H), 7.35 (dd, *J*=7.9, 4.8 Hz, 1 H), 7.19 (d, *J*=4.8 Hz, 1 H), 6.84 (td, *J*=17.1, 10.5 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.1 Hz, 1 H), 5.73 - 5.78 (m, 1 H), 4.43 - 4.96 (m, 2 H), 4.13 - 4.27 (m, 1 H), 3.43 - 3.93 (m, 3 H), 2.72 (tt, *J*=6.6, 3.2 Hz, 1 H), 1.95 (s, 3 H), 1.91 (s, 3 H), 1.35 (t, *J*=6.2 Hz, 3 H), 1.18 - 1.28 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.92 (d, *J*=6.6 Hz, 3 H)
96-2	587.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.43 (d, *J*=2.7 Hz, 1 H), 8.36 (d, *J*=4.8 Hz, 1 H), 8.15 (dd, *J*=4.6, 0.8 Hz, 1 H), 7.54 - 7.59 (m, 1 H), 7.44 (dd, *J*=8.5, 4.6 Hz, 1 H), 7.17 (d, *J*=5.0 Hz, 1 H), 6.83 (td, *J*=16.3, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.7, 1.6 Hz, 1 H), 5.72 - 5.78 (m, 1 H), 4.82 - 4.94 (m, 1 H), 4.42 - 4.81 (m, 1 H), 3.78 - 4.21 (m, 4 H), 3.69 (s, 3 H), 2.62 - 2.70 (m, 1 H), 1.93 (s, 3 H), 1.33 (t, *J*=7.0 Hz, 3 H), 1.21 (br dd, *J*=29.1, 6.7 Hz, 3 H), 1.05 (d, *J*=6.6 Hz, 3 H), 0.95 (d, *J*=6.6 Hz, 3 H)

100-1	592.3	^{1}H NMR (400 MHz, *DMSO-d_6*) δ 8.38 - 8.50 (m, 1 H), 7.50 - 7.59 (m, 1 H), 7.22 - 7.42 (m, 4 H), 6.75 - 6.91 (m, 1 H), 6.19 (dd, *J*=16.8, 2.1 Hz, 1 H), 5.71 - 5.79 (m, 1 H), 4.41 - 4.97 (m, 2 H), 3.38 - 4.33 (m, 5 H), 2.39 - 2.48 (m, 1 H), 1.11 - 1.37 (m, 12 H), 0.99 (dd, *J*=9.4, 6.9 Hz, 3 H), 0.93 (dd, *J*=6.9, 2.8 Hz, 3 H)
100-1-1	592.3	^{1}H NMR (400 MHz, *DMSO-d_6*) δ 8.41 (d, *J*=8.1 Hz, 1 H), 7.51 - 7.58 (m, 1 H), 7.39 (s, 1 H), 7.29 - 7.37 (m, 2 H), 7.23 - 7.28 (m, 1 H), 6.83 (ddd, *J*=24.3, 16.6, 10.6 Hz, 1 H), 6.19 (br d, *J*=16.8 Hz, 1 H), 5.75 (ddd, *J*=10.3, 4.9, 2.3 Hz, 1 H), 4.46 - 4.93 (m, 2 H), 3.38 - 4.32 (m, 5 H), 2.44 (td, *J*=6.9, 2.3 Hz, 1 H), 1.35 (br d, *J*=6.4 Hz, 3 H), 1.19 - 1.29 (m, 6 H), 1.14 (d, *J*=6.4 Hz, 3 H), 1.00 (d, *J*=6.8 Hz, 3 H), 0.93 (d, *J*=7.0 Hz, 3 H)
100-1-2	592.3	^{1}H NMR (400 MHz, *DMSO-d_6*) δ 8.48 (s, 1 H), 7.50 - 7.58 (m, 1 H), 7.38 (s, 1 H), 7.29 - 7.37 (m, 2 H), 7.23 - 7.29 (m, 1 H), 6.82 (dt, *J*=16.6, 11.0 Hz, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.72 - 5.78 (m, 1 H), 4.40 - 4.95 (m, 2 H), 3.48 - 4.17 (m, 5 H), 2.41 - 2.48 (m, 1 H), 1.29 - 1.35 (m, 3 H), 1.26 (d, *J*=6.4 Hz, 3 H), 1.11 - 1.23 (m, 6 H), 0.97 (d, *J*=6.8 Hz, 3 H), 0.93 (d, *J*=6.8 Hz, 3 H)
100-2	629.4	^{1}H NMR (500 MHz, DMSO-*d_6*) δ 9.06 (s, 1 H) 8.49 (d, *J*=7.01 Hz, 1 H) 7.52 (q, *J*=6.62 Hz, 1 H) 7.26 - 7.33 (m, 2 H) 7.17 - 7.20 (m, 1 H) 6.19 (br d, *J*=16.87 Hz, 1 H) 5.74 - 5.78 (m, 1 H) 4.88 (br d, *J*=13.62 Hz, 1 H) 4.46 - 4.52 (m, 1 H) 4.13 - 4.26 (m, 2 H) 3.75 - 3.94 (m, 4 H) 2.65 - 2.81 (m, 1 H) 1.34 (t, *J*=5.84 Hz, 3 H) 1.17 - 1.20 (m, 2 H) 1.07 (dd, *J*=6.55, 3.05 Hz, 6 H) 0.93 (br d, *J*=6.62 Hz, 6 H)
100-3	611.4	^{1}H NMR (500 MHz, DMSO-*d_6*) δ 9.09 (s, 1 H) 8.63 (d, *J*=4.80 Hz, 1 H) 8.00 (d, *J*=7.79 Hz, 1 H) 7.85 (t, *J*=7.72 Hz, 1 H) 7.71 (t, *J*=7.72 Hz, 1 H) 7.47 (d, *J*=7.79 Hz, 1 H) 6.83 - 6.94 (m, 1 H) 6.24 (dd, *J*=16.61, 2.08 Hz, 1 H) 5.80 (ddd, *J*=10.25, 5.97, 2.08 Hz, 1 H) 4.90 - 5.00 (m, 1 H) 4.83 (br s, 1 H) 4.24 (q, *J*=13.75 Hz, 1 H) 3.85 - 4.02 (m, 2 H) 3.49 - 3.60 (m, 1 H) 2.71 - 2.82 (m, 2 H) 1.37 - 1.43 (m, 3 H) 1.22 - 1.32 (m, 3 H) 1.12 (d, *J*=6.62 Hz, 6 H) 0.95 - 1.02 (m, 6 H)
100-4	628.4	^{1}H NMR (500 MHz, DMSO-*d_6*) δ 9.02 (s, 1 H) 8.47 (br s, 1 H) 7.39 (d, *J*=4.02 Hz, 2 H) 7.20 - 7.26 (m, 1 H) 7.00 (br d, *J*=7.53 Hz, 1 H) 6.79 - 6.89 (m, 1 H) 6.19 (dd, *J*=16.61, 2.34 Hz, 1 H) 5.74 - 5.78 (m, 1 H) 4.90 (br s, 1 H) 4.79 (br s, 1 H) 4.49 (br s, 1 H) 4.17 (br d, *J*=12.59 Hz, 1 H) 3.77 - 3.98 (m, 2 H) 2.64 - 2.85 (m, 2 H) 1.17 - 1.39 (m, 7 H) 1.02 - 1.12 (m, 9 H) 0.87 - 0.99 (m, 9 H)
100-5	424.1	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 7.96 (s, 1H), 6.51-6.67 (m, 1H), 6.31-6.45 (m, 1H), 5.79 (dd, *J*=1.87, 10.57 Hz, 1H), 4.30 (s, 2H), 3.81-3.88 (m, 8H), 1.00 (s, 9H).

100-6	356.3	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.61 (dd, *J*=1.87, 4.56 Hz, 1H), 7.94 (dd, *J*=1.76, 7.98 Hz, 1H), 7.08 (dd, *J*=4.56, 8.09 Hz, 1H), 6.60 (dd, *J*=10.57, 16.79 Hz, 1H), 6.37 (dd, *J*=1.87, 16.79 Hz, 1H), 5.78 (dd, *J*=1.76, 10.47 Hz, 1H), 4.41 (br s, 2H), 3.74-3.96 (m, 8H), 1.00 (s, 9H).
100-7	664.2	^{1}H NMR (400 MHz, *CDCl$_3$*) δ 9.07 (d, *J*=2.28 Hz, 1H), 8.01-8.19 (m, 2H), 7.63-7.74 (m, 2H), 7.18-7.25 (m, 1H), 6.50-6.74 (m, 1H), 6.40 (t, *J*=29.90 Hz, 1H), 5.76-5.86 (m, 1H), 3.57-5.30 (m, 6H), 2.82-2.96 (m, 1H), 2.53-2.66 (m, 4H), 0.91-1.61 (m, 18H).
100-8	574.6	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 1.03 (d, *J* = 6.8 Hz, 3 H), 1.12 (d, *J* = 6.8 Hz, 3 H), 1.16 - 1.27 (m, 3 H), 1.33 (t, *J* = 6.1 Hz, 3 H), 1.90 (s, 3 H), 2.56 - 2.65 (m, 1 H), 3.44 - 4.96 (m, 6 H), 5.67 - 5.82 (m, 1 H), 6.19 (dd, *J* = 16.7, 2.0 Hz, 1 H), 6.83 (td, *J* = 16.9, 10.6 Hz, 1 H), 7.18 - 7.37 (m, 3 H), 7.43 - 7.62 (m, 1 H), 8.38 (s, 1 H), 8.47 (d, *J* = 4.4 Hz, 1 H), 8.52 (s, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-*d$_6$*) δ –113.93 (s, 1 F)
100-9	574.6	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 1.00 (d, *J* = 6.8 Hz, 3 H), 1.11 (d, *J* = 6.8 Hz, 3 H), 1.13 - 1.27 (m, 3 H), 1.29 - 1.38 (m, 3 H), 1.93 (s, 3 H), 2.55 - 2.63 (m, 1 H), 3.44 - 4.97 (m, 6 H), 5.70 - 5.80 (m, 1 H), 6.19 (dd, *J* = 16.7, 2.0 Hz, 1 H), 6.83 (td, *J* = 15.9, 10.5 Hz, 1 H), 7.17 - 7.37 (m, 3 H), 7.47 - 7.57 (m, 1 H), 8.39 (s, 1 H), 8.48 (d, *J* = 3.5 Hz, 1 H), 8.50 (s, 1 H); ^{19}F NMR (376 MHz, DMSO-*d$_6$*) δ –114.03 (s, 1 F)
101-1	603.0	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 10.19 (br s, 1 H), 8.49 (d, *J*=3.1 Hz, 2 H), 8.30 (d, *J*=9.1 Hz, 1 H), 7.25 (q, *J*=8.1 Hz, 1 H), 6.62 - 6.90 (m, 3 H), 6.18 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.74 - 5.79 (m, 1 H), 4.42 - 4.93 (m, 2 H), 3.79 - 4.23 (m, 4 H), 2.41 - 2.48 (m, 2 H), 1.16 - 1.33 (m, 6 H), 1.11 (dd, *J*=6.7, 4.5 Hz, 6 H), 0.97 (br d, *J*=6.6 Hz, 6 H); ^{19}F NMR (377 MHz, *DMSO-d$_6$*) δ -115.75 (br s, 1 F), -128.46 (br s, 1 F)
101-2	588.2	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 9.03 (s, 1 H) 8.30 (dd, *J*=9.43, 4.87 Hz, 1 H) 7.48 (q, *J*=6.98 Hz, 1 H) 7.16 - 7.30 (m, 3 H) 6.71 - 6.84 (m, 1 H) 6.12 (br d, *J*=17.00 Hz, 1 H) 5.69 (br d, *J*=10.16 Hz, 1 H) 4.67 - 4.89 (m, 2 H) 4.06 - 4.19 (m, 1 H) 3.74 - 3.85 (m, 2 H) 2.65 (dq, *J*=12.75, 6.39 Hz, 2 H) 1.27 (t, *J*=5.91 Hz, 3 H) 1.19 (br d, *J*=6.63 Hz, 2 H) 1.12 (d, *J*=6.63 Hz, 2 H) 1.02 (dd, *J*=6.63, 1.87 Hz, 6 H) 0.87 (d, *J*=6.63 Hz, 6 H)
101-3	588.2	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 9.15 (s, 1 H) 8.27 (d, *J*=10.16 Hz, 1 H) 7.61 (q, *J*=7.19 Hz, 1 H) 7.27 - 7.43 (m, 3 H) 6.96 (dd, *J*=16.59, 10.37 Hz, 1 H) 6.32 (dd, *J*=16.69, 2.18 Hz, 1 H) 5.86 (dd, *J*=10.26, 1.97 Hz, 1 H) 5.12 (br s, 2 H) 4.39 (br d, *J*=13.06 Hz, 1 H) 4.12 (br d, *J*=13.48 Hz, 1 H) 3.66 (br d, *J*=12.23 Hz, 1 H) 3.19 - 3.31 (m, 1 H)

		2.72 - 2.85 (m, 2 H) 1.54 (d, *J*=6.84 Hz, 6 H) 1.16 (d, *J*=6.63 Hz, 6 H) 0.99 (d, *J*=6.63 Hz, 6 H)
102-1	601.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.46 (s, 1 H), 7.43 - 7.57 (m, 1 H), 7.12 - 7.36 (m, 6 H), 6.75 - 6.94 (m, 1 H), 6.19 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.67 - 5.85 (m, 1 H), 4.69 - 4.96 (m, 2 H), 4.39 - 4.52 (m, 1 H), 4.14 (br t, *J*=11.9 Hz, 1 H), 3.78 - 3.99 (m, 2 H), 3.44 - 3.62 (m, 1 H), 1.30 - 1.37 (m, 3 H), 1.25 (br d, *J*=6.4 Hz, 2 H), 1.17 (d, *J*=6.8 Hz, 2 H), 1.03 - 1.10 (m, 6 H), 0.92 (dd, *J*=10.0, 6.8 Hz, 6 H)
102-2	646.8	^{1}H NMR (400 MHz, DMSO-d6) δ 0.88 - 1.00 (m, 6 H) 1.04 - 1.18 (m, 6 H) 1.31 - 1.42 (m, 3 H) 2.06 - 2.28 (m, 6 H) 2.65 - 2.82 (m, 2 H) 3.02 - 3.17 (m, 2 H) 3.37 - 3.89 (m, 3 H) 3.95 - 4.19 (m, 1 H) 4.22 - 4.45 (m, 2 H) 4.85 - 5.18 (m, 1 H) 6.53 - 6.83 (m, 2 H) 7.07 - 7.24 (m, 1 H) 7.26 - 7.40 (m, 2 H) 7.43 - 7.62 (m, 1 H) 8.32 - 8.69 (m, 1 H) 8.88 - 9.20 (m, 1 H) ^{19}F NMR (376 MHz, DMSO-d6) δ -113.35 (s, 1 F)
102-3	602.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.44 - 8.51 (m, 2 H), 7.46 - 7.56 (m, 1 H), 7.25 - 7.35 (m, 3 H), 7.14 - 7.21 (m, 1 H), 6.78 - 6.91 (m, 1 H), 6.20 (dd, *J*=16.5, 2.2 Hz, 1 H), 5.74 - 5.80 (m, 1 H), 4.73 - 4.94 (m, 2 H), 4.44 (s, 1 H), 4.11 - 4.24 (m, 1 H), 3.84 - 3.96 (m, 2 H), 2.61 - 2.70 (m, 2 H), 1.34 (t, *J*=6.6 Hz, 3 H), 1.26 (br d, *J*=7.0 Hz, 1 H), 1.18 (d, *J*=6.8 Hz, 2 H), 1.07 (dd, *J*=6.5, 3.8 Hz, 6 H), 0.93 (td, *J*=6.2, 2.9 Hz, 6 H)
102-4	682.4	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.49 (d, *J*=3.1 Hz, 1 H), 7.53 (d, *J*=4.8 Hz, 2 H), 7.26 - 7.39 (m, 2 H), 7.14 - 7.24 (m, 1 H), 6.74 - 6.91 (m, 1 H), 6.20 (dd, *J*=16.9, 2.2 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 4.73 - 4.95 (m, 1 H), 4.44 - 4.53 (m, 1 H), 4.11 - 4.28 (m, 1 H), 3.78 - 3.96 (m, 2 H), 3.49 (br dd, *J*=14.1, 2.3 Hz, 1 H), 1.34 (br t, *J*=5.8 Hz, 3 H), 1.22 - 1.28 (m, 3 H), 1.18 (br d, *J*=6.8 Hz, 2 H), 1.06 (td, *J*=7.3, 3.8 Hz, 6 H), 0.86 - 0.98 (m, 6 H)
102-5	596.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.96 (1 H, s) 8.15 (1 H, s) 7.44 - 7.51 (1 H, m) 7.13 - 7.28 (3 H, m) 6.55 - 6.71 (1 H, m) 6.38 - 6.48 (1 H, m) 5.80 - 5.87 (1 H, m) 3.45 - 5.24 (6 H, m) 2.86 - 2.97 (1 H, m) 1.41 - 1.56 (4 H, m) 1.36 (2 H, br d, *J*=6.84 Hz) 1.25 - 1.34 (3 H, m) 1.12 (3 H, dd, *J*=8.71, 6.84 Hz)
102-5-1	596.0	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.96 (1 H, s) 8.15 (1 H, s) 7.44 - 7.51 (1 H, m) 7.13 - 7.28 (3 H, m) 6.55 - 6.71 (1 H, m) 6.38 - 6.48 (1 H, m) 5.80 - 5.87 (1 H, m) 3.49 - 5.24 (6 H, m) 2.87 - 2.97 (1 H, m) 1.42 - 1.55 (4 H, m) 1.36 (2 H, d, *J*=6.84 Hz) 1.31 (3 H, d, *J*=6.84 Hz) 1.14 (3 H, d, *J*=6.63 Hz)
102-5-2	596.0	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.96 (1 H, s) 8.14 (1 H, s) 7.44 - 7.51 (1 H, m) 7.13 - 7.28 (3 H, m) 6.55 - 6.71 (1 H, m) 6.38 - 6.48 (1 H, m) 5.80 - 5.87 (1 H, m) 3.45 - 5.24 (6 H, m) 2.86 - 2.97 (1 H, m) 1.49 (4 H, td,

		J=15.76, 6.63 Hz) 1.35 (2 H, d, *J*=6.63 Hz) 1.30 (3 H, d, *J*=6.63 Hz) 1.11 (3 H, d, *J*=6.84 Hz)
102-6	592.1	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.87 (1 H, s) 8.10 (1 H, s) 7.25 - 7.32 (1 H, m) 7.14 - 7.21 (2 H, m) 7.03 - 7.08 (1 H, m) 6.49 - 6.66 (1 H, m) 6.32 - 6.42 (1 H, m) 5.74 - 5.81 (1 H, m) 3.41 - 5.19 (6 H, m) 2.78 - 2.92 (1 H, m) 1.99 - 2.04 (3 H, m) 1.38 - 1.50 (4 H, m) 1.28 - 1.34 (2 H, m) 1.20 - 1.26 (3 H, m) 1.02 (3 H, br d, *J*=6.63 Hz)
102-6-1	592.1	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.91 (1 H, s) 8.14 (1 H, s) 7.29 - 7.35 (1 H, m) 7.19 - 7.24 (2 H, m) 7.08 (1 H, d, *J*=7.46 Hz) 6.52 - 6.69 (1 H, m) 6.40 (1 H, br t, *J*=15.13 Hz) 5.78 - 5.84 (1 H, m) 3.44 - 5.20 (6 H, m) 2.84 - 2.95 (1 H, m) 2.05 (3 H, s) 1.41 - 1.53 (4 H, m) 1.33 - 1.37 (2 H, m) 1.25 - 1.30 (3 H, m) 1.03 - 1.08 (3 H, m)
102-6-2	592.1	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.91 (1 H, s) 8.13 (1 H, s) 7.29 - 7.35 (1 H, m) 7.19 - 7.25 (2 H, m) 7.10 (1 H, br d, *J*=7.46 Hz) 6.52 - 6.69 (1 H, m) 6.36 - 6.45 (1 H, m) 5.78 - 5.84 (1 H, m) 3.44 - 5.22 (6 H, m) 2.82 - 2.94 (1 H, m) 2.06 (3 H, s) 1.40 - 1.53 (4 H, m) 1.30 - 1.36 (2 H, m) 1.27 (3 H, d, *J*=6.84 Hz) 1.05 (3 H, d, *J*=6.84 Hz)
102-7	650.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 0.95 - 1.06 (m, 3 H), 1.06 - 1.11 (m, 3 H), 1.15 - 1.20 (m, 3 H), 1.28 - 1.35 (m, 3 H), 1.66 (d, *J* = 13.3 Hz, 6 H), 1.92 - 2.01 (m, 3 H), 2.56 - 2.64 (m, 1 H), 3.45 - 4.96 (m, 6 H), 5.71 - 5.79 (m, 1 H), 6.19 (dd, *J* = 16.7, 2.2 Hz, 1 H), 6.76 - 6.91 (m, 1 H), 7.16 - 7.36 (m, 3 H), 7.44 - 7.57 (m, 2 H), 7.57 - 7.67 (m, 1 H), 8.47 (s, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –113.91 (s, 1 F), –113.81 (s, 1 F); ^{31}P{^{1}H} NMR (162 MHz, DMSO-d_6) δ 32.40 (s, 1 P)
102-7-1	649.6	^{1}H NMR (400 MHz, DMSO-d_6) δ 1.01 (d, *J* = 6.8 Hz, 3 H), 1.05 (d, *J* = 6.2 Hz, 3 H), 1.08 (d, *J* = 6.8 Hz, 3 H), 1.17 (br d, *J* = 6.8 Hz, 3 H), 1.66 (d, *J* = 13.3 Hz, 6 H), 1.95 (s, 3 H), 2.55 - 2.64 (m, 1 H), 3.51 (dd, *J* = 14.2, 3.4 Hz, 0.5 H), 3.83 - 3.97 (m, 2 H), 4.07 - 4.22 (m, 1.5 H), 4.40 - 4.52 (m, 0.5 H), 4.72 - 4.97 (m, 1.5 H), 5.70 - 5.81 (m, 1 H), 6.19 (dd, *J* = 16.7, 2.4 Hz, 1 H), 6.83 (td, *J* = 16.0, 10.6 Hz, 1 H), 7.19 - 7.35 (m, 3 H), 7.46 - 7.57 (m, 2 H), 7.62 (d, *J* = 11.8 Hz, 1 H), 8.47 (d, *J*=1.87 Hz, 1 H); ^{19}F{^{1}H} NMR (377 MHz, DMSO-d_6) δ –113.82 (s, 1 F); ^{31}P{^{1}H} NMR (162 MHz, DMSO-d_6) δ 32.41 (s, 1 P)
102-7-2	649.6	^{1}H NMR (400 MHz, DMSO-d_6) δ 0.97 (d, *J* = 6.8 Hz, 3 H), 1.03 - 1.08 (m, 6 H), 1.40 - 1.46 (m, 3 H), 1.66 (d, *J* = 13.3 Hz, 6 H), 1.98 (s, 3 H), 2.55 - 2.63 (m, 1 H), 3.44 - 3.55 (m, 0.5 H), 3.76 - 4.00 (m, 2 H), 4.10 - 4.22 (m, 1.5 H), 4.42 - 4.51 (m, 0.5 H), 4.72 - 4.95 (m, 1.5 H), 5.69 - 5.81 (m, 1 H), 6.19 (dd, *J* = 16.6, 2.1 Hz, 1 H), 6.75 - 6.92 (m, 1 H), 7.20 - 7.36 (m, 3 H), 7.47 - 7.63 (m, 3 H), 8.47 (s, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –113.92 (s, 1 F); ^{31}P{^{1}H} NMR (162 MHz, DMSO-d_6) δ 32.43 (s, 1 P)

102-8	632.2	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.30 (br dd, *J*=4.77, 12.44 Hz, 1H), 7.50 (s, 1H), 7.33-7.43 (m, 1H), 7.20 (br s, 2H), 7.04-7.17 (m, 2H), 6.26-6.38 (m, 1H), 5.65-5.80 (m, 1H), 5.01-5.14 (m, 1H), 4.27-4.47 (m, 1H), 3.76-3.94 (m, 1H), 3.54-3.74 (m, 1H), 3.23-3.41 (m, 1H), 2.93-3.00 (m, 1H), 2.66-2.83 (m, 2H), 1.33 (br s, 6H), 1.23-1.30 (m, 6H), 1.11-1.22 (m, 6H)
103-1	628.3	
103-1-1	628.4	^{1}H NMR (400 MHz, *CDCl$_3$*) δ 8.26 (br d, *J*=5.60 Hz, 1H), 8.08 (s, 1H), 7.30-7.42 (m, 2H), 7.16-7.22 (m, 1H), 7.00 (d, *J*=7.67 Hz, 1H), 6.52-6.68 (m, 2H), 6.40 (br t, *J*=14.82 Hz, 1H), 5.77-5.84 (m, 1H), 3.40-5.23 (m, 6H), 2.69 (s, 6H), 2.49-2.65 (m, 2H), 0.88-1.52 (m, 18H)
103-1-2	628.4	^{1}H NMR (400 MHz, *CDCl$_3$*) δ 8.29 (d, *J*=5.60 Hz, 1H), 8.06-8.12 (m, 1H), 7.30-7.41 (m, 2H), 7.19 (t, *J*=7.26 Hz, 1H), 7.00 (d, *J*=7.67 Hz, 1H), 6.52-6.69 (m, 2H), 6.40 (br t, *J*=15.45 Hz, 1H), 5.77-5.84 (m, 1H), 3.43-5.26 (m, 6H), 2.69 (br s, 6H), 2.51-2.63 (m, 2H), 0.94-1.45 (m, 18H)
103-2	591.8	^{1}H NMR (400 MHz, *CDCl$_3$*) δ 8.14 (s, 1 H) 8.06 (s, 1 H) 7.40 - 7.48 (m, 1 H) 7.30 (td, *J* = 7.4, 1.9 Hz, 1 H) 7.16 - 7.23 (m, 1 H) 7.13 (t, *J* = 9.1 Hz, 1 H) 6.51 - 6.69 (m, 1 H) 6.32 - 6.44 (m, 1 H) 5.74 - 5.83 (m, 1 H) 4.83 - 5.16 (m, 2 H) 4.25 - 4.48 (m, 1 H) 3.79 - 4.07 (m, 2 H) 3.48 - 3.51 (m, 3 H) 3.44 - 3.71 (m, 1 H) 2.65 - 2.79 (m, 1 H) 1.33 - 1.48 (m, 6 H) 1.22 (d, *J* = 6.6 Hz, 3 H) 1.06 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -112.44 - -112.19 (m, 1 F)
103-3	622.1	^{1}H NMR (400 MHz, *CDCl3*) δ 8.09 (s, 1H), 7.21-7.91 (m, 4H), 7.13-7.19 (m, 2H), 7.09 (t, *J*=18.20 Hz, 1H), 6.50-6.71 (m, 1H), 6.40 (t, *J*=29.60 Hz, 1H), 5.80 (t, *J*=18.00 Hz, 1H), 4.19-5.25 (m, 3H), 3.35-4.11 (m, 3H), 2.76-2.81 (m, 3H), 2.58-2.76 (m, 1H), 1.31-1.55 (m, 6H), 1.19-1.27 (m, 3H), 1.02-1.09 (m, 3H). ^{19}F NMR (376 MHz, *CDCl3*) δ -112.37 (s, 1F), -112.42 (s, 1F), -112.40 (br dd, *J*=16.91, 23.84 Hz, 1F)
103-4	603.2	^{1}H NMR (400 MHz, *CDCl3*) δ 8.53 (s, 1H), 8.52 (s, 1H), 8.10-8.14 (m, 1H), 7.38-7.46 (m, 1H), 7.07-7.18 (m, 3H), 6.52-6.70 (m, 1H), 6.40 (t, *J*=29.90 Hz, 1H), 5.80 (t, *J*=18.00 Hz, 1H), 4.26-5.22 (m, 3H), 3.44-4.08 (m, 3H), 2.45-2.68 (m, 2H), 1.31-1.50 (m, 6H), 1.25 (dd, *J*=2.28, 6.43 Hz, 6H), 1.01-1.07 (m, 6H). ^{19}F NMR (376 MHz, *CDCl3*) δ -113.19 (s, 1F), -113.21 (s, 1F)
103-5	656.2	^{1}H NMR (400 MHz, *CDCl3*) δ 8.27-8.36 (m, 1H), 8.04-8.14 (m, 1H), 7.29-7.42 (m, 2H), 7.17 (t, *J*=7.26 Hz, 1H), 7.04 (t, *J*=6.74 Hz, 1H), 6.52-6.71 (m, 2H), 6.35-6.45 (m, 1H), 5.77-5.84 (m, 1H), 3.27-5.33 (m, 6H), 2.98 (br s, 4H), 2.57 (br s, 2H), 1.32-1.62 (m, 6H), 1.23-1.31 (m,

		7H), 0.91-1.05 (m, 7H), 0.86-0.91 (m, 3H), 0.67-0.84 (m, 6H)
103-6	628.3	^{1}H NMR (400 MHz, *CDCl3*) δ 8.32-8.38 (m, 1H), 8.03-8.12 (m, 1H), 7.27-7.32 (m, 1H), 7.14-7.21 (m, 2H), 7.05-7.11 (m, 1H), 6.51-6.73 (m, 2H), 6.40 (br t, *J*=15.03 Hz, 1H), 5.76-5.83 (m, 1H), 4.16-5.32 (m, 3H), 3.27-4.10 (m, 3H), 2.90-3.05 (m, 4H), 2.52-2.67 (m, 1H), 1.99 (s, 3H), 1.35-1.67 (m, 6H), 1.15-1.35 (m, 6H), 0.86-0.99 (m, 4H), 0.68-0.85 (m, 6H)
103-6-1	656.2	^{1}H NMR (400 MHz, *CDCl3*) δ 8.30 (br d, *J*=4.98 Hz, 1H), 8.10 (br d, *J*=4.56 Hz, 1H), 7.29-7.40 (m, 2H), 7.17 (t, *J*=6.74 Hz, 1H), 7.04 (br d, *J*=7.88 Hz, 1H), 6.49-6.73 (m, 2H), 6.40 (br t, *J*=15.13 Hz, 1H), 5.80 (br t, *J*=8.91 Hz, 1H), 4.25-5.23 (m, 3H), 3.25-4.22 (m, 3H), 2.88-3.08 (m, 4H), 2.49-2.74 (m, 2H), 1.36-1.59 (m, 6H), 1.21-1.29 (m, 3H), 0.62-1.11 (m, 15H)
103-6-2	656.2	^{1}H NMR (400 MHz, *CDCl3*) δ 8.33 (d, *J*=5.60 Hz, 1H), 8.04-8.11 (m, 1H), 7.29-7.41 (m, 2H), 7.17 (t, *J*=7.36 Hz, 1H), 7.03 (d, *J*=7.46 Hz, 1H), 6.53-6.70 (m, 2H), 6.40 (br t, *J*=14.93 Hz, 1H), 5.77-5.83 (m, 1H), 3.54-5.33 (m, 6H), 2.98 (br s, 4H), 2.58 (br s, 2H), 1.23-1.44 (m, 9H), 0.67-1.05 (m, 15H)
103-7	600.3	
103-7-1	600.3	^{1}H NMR (400 MHz, *CDCl3*) δ 8.29 (br d, *J*=5.80 Hz, 1H), 8.08 (s, 1H), 7.28-7.34 (m, 1H), 7.17-7.23 (m, 2H), 7.10 (d, *J*=7.26 Hz, 1H), 6.50-6.70 (m, 2H), 6.40 (br t, *J*=14.82 Hz, 1H), 5.77-5.83 (m, 1H), 3.42-5.21 (m, 6H), 2.63-2.76 (m, 7H), 2.04 (s, 3H), 1.27-1.50 (m, 6H), 1.20-1.27 (m, 3H), 0.93 (br d, *J*=6.63 Hz, 3H)
103-7-2	600.2	^{1}H NMR (400 MHz, *CDCl3*) δ 8.32 (d, *J*=5.81 Hz, 1H), 8.06-8.11 (m, 1H), 7.27-7.34 (m, 1H), 7.16-7.23 (m, 2H), 7.08-7.12 (m, 1H), 6.52-6.70 (m, 2H), 6.40 (br t, *J*=15.34 Hz, 1H), 5.77-5.83 (m, 1H), 3.44-5.26 (m, 6H), 2.70 (s, 6H), 2.52-2.66 (m, 1H), 2.02 (s, 3H), 1.26-1.47 (m, 6H), 1.23 (d, *J*=6.63 Hz, 3H), 0.94 (br d, *J*=6.22 Hz, 3H)
103-8-1	628.4	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.26-8.38 (m, 1H), 8.10 (br d, *J*=6.84 Hz, 1H), 7.28-7.33 (m, 1H), 7.07-7.22 (m, 3H), 6.52-6.71 (m, 2H), 6.34-6.52 (m, 1H), 5.80 (br t, *J*=9.02 Hz, 1H), 3.64-5.17 (m, 6H), 2.90-3.06 (m, 4H), 2.61 (tdd, *J*=6.45, 13.09, 19.62 Hz, 1H), 1.99 (s, 3H), 1.34-1.59 (m, 6H), 1.25 (d, *J*=6.63 Hz, 3H), 0.93 (d, *J*=6.84 Hz, 3H), 0.70-0.85 (m, 6H)
103.8-2	628.4	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.36 (d, *J*=5.60 Hz, 1H), 8.02-8.14 (m, 1H), 7.27-7.34 (m, 1H), 7.14-7.22 (m, 2H), 7.00-7.14 (m, 1H), 6.71 (d, *J*=5.60 Hz, 1H), 6.51-6.67 (m, 1H), 6.34-6.51 (m, 1H), 5.73-5.83 (m, 1H), 3.46-5.31 (m, 6H), 2.99 (br d, *J*=6.63 Hz, 4H), 2.50-

		2.60 (m, 1H), 1.99 (s, 3H), 1.18-1.45 (m, 9H), 0.96 (br d, *J*=6.43 Hz, 3H), 0.69-0.89 (m, 6H)
106-1	682.0	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.17 (1 H, s) 7.48 - 7.54 (1 H, m) 7.16 - 7.28 (3 H, m) 6.56 - 6.74 (1 H, m) 6.40 - 6.49 (1 H, m) 5.82 - 5.88 (1 H, m) 3.49 - 5.23 (6 H, m) 2.60 - 2.72 (2 H, m) 1.44 - 1.56 (3 H, m) 1.29 - 1.40 (3 H, m) 1.23 - 1.29 (6 H, m) 1.03 - 1.09 (6 H, m)
106-2	498.2	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 7.93 (s, 1H), 7.44-7.53 (m, 2H), 7.27-7.34 (m, 1H), 7.14-7.19 (m, 1H), 6.46-6.71 (m, 1H), 6.30-6.43 (m, 1H), 5.71-5.82 (m, 1H), 4.39-4.74 (m, 2H), 4.34 (br s, 2H), 3.53-4.19 (m, 4H), 2.97-3.20 (m, 1H), 1.24 (s, 3H), 0.95 (s, 9H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.58 (s, 1F)
106-3	514.3	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.04-8.38 (m, 1H), 7.38-7.56 (m, 2H), 7.26-7.32 (m, 1H), 7.18 (t, *J*=9.12 Hz, 1H), 6.47-6.73 (m, 1H), 6.37 (br d, *J*=16.38 Hz, 1H), 5.68-5.84 (m, 1H), 4.45-4.87 (m, 2H), 4.28 (br d, *J*=8.50 Hz, 2H), 3.57-3.89 (m, 4H), 3.31-3.49 (m, 1H), 3.07 (tt, *J*=3.63, 7.36 Hz, 1H), 1.39 (d, *J*=6.63 Hz, 2H), 0.90-0.98 (m, 9H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.40 (br s, 1F)
106-4	528.2	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 7.94 (s, 1H), 7.40-7.56 (m, 2H), 7.26-7.32 (m, 1H), 7.18 (t, *J*=9.12 Hz, 1H), 6.48-6.60 (m, 1H), 6.25-6.41 (m, 1H), 5.69-5.80 (m, 1H), 4.80-4.99 (m, 2H), 3.59-4.62 (m, 6H), 3.11 (s, 3H), 1.35-1.37 (m, 3H), 1.23-1.26 (m, 3H), 1.19 (s, 6H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.72 (br d, *J*=9.54 Hz, 1F)
106-5	500.1	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.00 (s, 1H), 7.46-7.58 (m, 2H), 7.28-7.36 (m, 1H), 7.21 (t, *J*=9.12 Hz, 1H), 6.61 (dd, *J*=10.47, 16.69 Hz, 1H), 6.39 (dd, *J*=1.87, 16.79 Hz, 1H), 5.80 (dd, *J*=1.76, 10.47 Hz, 1H), 4.57 (br s, 2H), 3.75-4.08 (m, 8H), 3.15 (s, 3H), 1.23 (s, 6H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.74 (s, 1F)
106-6	576.0	^{1}H NMR (400 MHz, DMSO-d_6) 8. .42 (d, *J*=4.77 Hz, 1 H), 7.50 - 7.63 (m, 1 H), 7.29 - 7.41 (m, 3 H), 7.20 (dd, *J*=4.46, 3.42 Hz, 1 H), 6.77 - 7.04 (m, 1 H), 6.13 - 6.31 (m, 1 H), 5.69 - 6.02 (m, 1 H), 5.10 - 5.66 (m, 1 H), 4.54 - 4.97 (m, 2 H), 4.30 (br d, *J*=13.48 Hz, 1 H), 3.54 - 3.99 (m, 2 H), 2.73-2.96 (m, 1 H), 1.92 - 2.05 (m, 3 H), 1.24 - 1.52 (m, 6 H), 1.08 (br d, *J*=6.63 Hz, 3 H), 0.98 (br d, *J*=6.63 Hz, 3 H)

106-6-1	576.0	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.42 (d, J=4.77 Hz, 1 H), 7.51 - 7.66 (m, 1 H), 7.27 - 7.42 (m, 3 H), 7.20 (d, J=4.98 Hz, 1 H), 6.77 - 7.00 (m, 1 H), 6.21 (br d, J=16.59 Hz, 1 H), 5.43 - 6.01 (m, 2 H), 4.21 - 5.33 (m, 3 H), 3.41 - 4.10 (m, 2 H), 2.71 - 2.89 (m, 1 H), 2.01 (br s, 3 H), 1.18 - 1.53 (m, 6 H), 1.07 (br s, 3 H), 0.98 (d, J=6.63 Hz, 3 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –113.62 (s, 1 F)
106-6-2	576.0	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.42 (d, J=4.77 Hz, 1 H), 7.52 - 7.62 (m, 1 H), 7.28 - 7.42 (m, 3 H), 7.20 (d, J=4.98 Hz, 1 H), 6.76 - 7.01 (m, 1 H), 6.12 - 6.30 (m, 1 H), 5.42 - 5.96 (m, 2 H), 4.20 - 5.30 (m, 3 H), 3.39 - 4.11 (m, 2 H), 2.81 - 2.96 (m, 1 H), 1.97 (br s, 3 H), 1.20 - 1.53 (m, 6 H), 1.08 (d, J=6.63 Hz, 3 H), 0.96 (br s, 3 H) ; ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –113.80 (s, 1 F)
106-7	576.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 0.96 (br s, 3 H), 1.04 - 1.12 (m, 3 H), 1.28 - 1.42 (m, 3 H), 1.52 (br s, 3 H), 1.90 - 2.09 (m, 3 H), 2.77 - 2.90 (m, 1 H), 3.09 - 3.23 (m, 1 H), 3.51 - 3.62 (m, 1 H), 4.13 - 4.27 (m, 1 H), 4.36 - 4.50 (m, 1 H), 5.33 - 5.53 (m, 1 H), 5.63 - 5.75 (m, 1 H), 5.77 - 5.85 (m, 1 H), 6.27 (br d, J = 17.8 Hz, 1 H), 6.95 (dd, J = 16.5, 10.5 Hz, 1 H), 7.20 (d, J = 4.8 Hz, 1 H), 7.26 - 7.39 (m, 3 H), 7.52 - 7.62 (m, 1 H), 8.41 (d, J = 5.0 Hz, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –113.77 (s, 1 F), –113.56 (s, 1 F)
106-7-1	576.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 0.96 (br s, 3 H), 1.07 (d, J = 6.6 Hz, 3 H), 1.33 (br s, 3 H), 1.52 (br s, 3 H), 1.98 (s, 3 H), 2.72 - 2.96 (m, 1 H), 3.09 - 3.23 (m, 1 H), 3.47 - 3.65 (m, 1 H), 4.07 - 4.28 (m, 1 H), 4.31 - 4.52 (m, 1 H), 5.33 - 5.56 (m, 1 H), 5.61 - 5.75 (m, 1 H), 5.77 - 5.85 (m, 1 H), 6.27 (br d, J = 16.6 Hz, 1 H), 6.95 (dd, J = 16.6, 10.4 Hz, 1 H), 7.20 (d, J = 4.8 Hz, 1 H), 7.25 - 7.42 (m, 3 H), 7.51 - 7.63 (m, 1 H), 8.41 (d, J = 5.0 Hz, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –113.78 (s, 1 F), –113.56 (s, 1 F)
106-7-2	576.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 0.95 (br s, 3 H), 1.07 (br d, J = 6.6 Hz, 3 H), 1.33 (br s, 3 H), 1.52 (br s, 3 H), 1.98 (s, 3 H), 2.74 - 2.93 (m, 1 H), 3.09 - 3.25 (m, 1 H), 3.56 (br dd, J = 9.6, 2.8 Hz, 1 H), 4.12 - 4.26 (m, 1 H), 4.33 - 4.50 (m, 1 H), 5.34 - 5.53 (m, 1 H), 5.61 - 5.74 (m, 1 H), 5.79 (dd, J = 10.4, 2.1 Hz, 1 H), 6.27 (br d, J = 16.6 Hz, 1 H), 6.95 (br dd, J = 16.6, 10.4 Hz, 1 H), 7.19 (br d, J = 5.0 Hz, 1 H), 7.26 - 7.40 (m, 3 H), 7.52 - 7.61 (m, 1 H), 8.41 (d, J = 5.0 Hz, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –113.78 (s, 1 F), –113.57 (s, 1 F)
106-8	523.1	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.37 (br s, 1 H), 7.55 - 7.72 (m, 2 H), 7.34 - 7.46 (m, 2 H), 6.81 (td, J=16.0, 10.6 Hz, 1 H), 6.17 (br d, J=16.4 Hz, 1 H), 5.74 (br d, J=9.5 Hz, 1 H), 4.77 (br d, J=16.4 Hz, 2 H), 4.33 - 4.52 (m, 2 H),

		4.01 - 4.19 (m, 2 H), 3.73 - 3.91 (m, 2 H), 1.36 (s, 6 H), 1.29 (br t, *J*=5.7 Hz, 3 H), 1.07 - 1.23 (m, 3 H)
106-9	509.1	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.37 (br s, 1 H), 7.57 - 7.71 (m, 2 H), 7.34 - 7.46 (m, 2 H), 6.76 - 6.93 (m, 1 H), 6.19 (br d, *J*=16.4 Hz, 1 H), 5.75 (br d, *J*=10.8 Hz, 1 H), 4.88 (br s, 1 H), 4.44 (s, 2 H), 4.18 - 4.40 (m, 2 H), 3.94 - 4.16 (m, 1 H), 3.36 - 3.78 (m, 2 H), 2.99 - 3.25 (m, 1 H), 1.37 (s, 6 H), 1.29 (br d, *J*=6.4 Hz, 3 H)
106-10	495.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.47 (s, 1 H), 7.55 - 7.72 (m, 2 H), 7.34 - 7.49 (m, 2 H), 6.83 (dd, *J*=16.7, 10.5 Hz, 1 H), 6.10 - 6.27 (m, 1 H), 5.70 - 5.81 (m, 1 H), 4.45 (s, 2 H), 3.97 (br d, *J*=5.2 Hz, 4 H), 3.71 - 3.89 (m, 4 H), 1.37 (s, 6 H)
107-1	633.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.13 (1 H, s) 7.42 - 7.53 (1 H, m) 7.15 - 7.29 (3 H, m) 6.57 - 6.74 (1 H, m) 6.39 - 6.49 (1 H, m) 5.81 - 5.88 (1 H, m) 3.51 - 5.23 (7 H, m) 3.05 (3 H, d, *J*=4.77 Hz) 2.48- 2.61 (2 H, m) 1.44 - 1.52 (3 H, m) 1.28 - 1.39 (3 H, m) 1.21 (6 H, dd, *J*=6.53, 4.66 Hz) 0.98 - 1.05 (6 H, m)
107-2	647.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.05 - 8.09 (1 H, m) 7.40 - 7.47 (1 H, m) 7.15 - 7.23 (2 H, m) 7.12 (1 H, t, *J*=9.12 Hz) 6.51 - 6.69 (1 H, m) 6.34 - 6.44 (1 H, m) 5.76 - 5.83 (1 H, m) 3.47 - 5.17 (6 H, m) 3.21 (6 H, s) 2.44 - 2.60 (2 H, m) 1.39 - 1.46 (3 H, m) 1.25 - 1.33 (3 H, m) 1.14 - 1.19 (6 H, m) 0.95 - 1.01 (6 H, m)
107-3	689.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.11 - 8.15 (1 H, m) 7.46 - 7.53 (1 H, m) 7.22 - 7.28 (2 H, m) 7.15 - 7.21 (1 H, m) 6.56 - 6.74 (1 H, m) 6.39 - 6.49 (1 H, m) 5.81 - 5.88 (1 H, m) 3.80 - 3.94 (8 H, m) 3.51 - 5.24 (6 H, m) 2.51 - 2.69 (2 H, m) 1.43 - 1.52 (3 H, m) 1.29 - 1.38 (3 H, m) 1.20 (6 H, dd, *J*=6.63, 4.77 Hz) 0.99 - 1.05 (6 H, m)
107-4	690.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.12 - 8.18 (1 H, m) 7.45 - 7.55 (1 H, m) 7.15 - 7.30 (3 H, m) 6.58 - 6.76 (1 H, m) 6.41 - 6.51 (1 H, m) 5.83 - 5.90 (1 H, m) 3.53 - 5.59 (9 H, m) 2.37 - 2.85 (10 H, m) 1.45 - 1.53 (3 H, m) 1.31 - 1.41 (3 H, m) 1.21 (6 H, dd, *J*=6.63, 3.94 Hz) 0.97 - 1.04 (6 H, m)
107-5	659.5	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.15 (1 H, br s) 7.47 - 7.55 (1 H, m) 7.23 - 7.29 (2 H, m) 7.19 (1 H, t, *J*=9.12 Hz) 6.58 - 6.77 (1 H, m) 6.42 - 6.52 (1 H, m) 5.84 - 5.91 (1 H, m) 3.53 - 5.25 (10 H, m) 2.49 - 2.71 (2 H, m) 2.35 - 2.44 (2 H, m) 1.31 - 1.54 (6 H, m) 1.20 - 1.27 (6 H, m) 0.99 - 1.07 (6 H, m)
108-1	602.0	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.01 - 9.03 (1 H, m) 8.16 - 8.19 (1 H, m) 7.46 - 7.52 (1 H, m) 7.14 - 7.30 (3 H, m) 6.58 - 6.74 (1 H, m) 6.46 (1 H, br t, *J*=15.03 Hz) 5.83 - 5.89 (1 H, m) 3.49 - 5.27 (6 H, m) 2.75 - 2.86 (1 H, m) 1.44 - 1.63 (5 H, m) 1.27 - 1.41 (6 H, m) 1.07 - 1.20 (4 H, m) 0.96 - 1.06 (1 H, m) 0.84 - 0.95 (1 H, m)

108-1-1	602.0	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.01 (1 H, s) 8.18 (1 H, s) 7.45 - 7.52 (1 H, m) 7.14 - 7.29 (3 H, m) 6.57 - 6.74 (1 H, m) 6.45 (1 H, br t, *J*=14.72 Hz) 5.83 - 5.89 (1 H, m) 3.48 - 5.25 (6 H, m) 2.73 - 2.84 (1 H, m) 1.46 - 1.63 (5 H, m) 1.36 - 1.42 (2 H, m) 1.26 - 1.33 (4 H, m) 1.08 - 1.17 (4 H, m) 0.93 - 1.03 (1 H, m) 0.81 - 0.89 (1 H, m)
108-1-2	602.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.99 (1 H, s) 8.13 - 8.17 (1 H, m) 7.45 - 7.51 (1 H, m) 7.13 - 7.28 (3 H, m) 6.56 - 6.72 (1 H, m) 6.44 (1 H, br t, *J*=15.03 Hz) 5.81 - 5.87 (1 H, m) 3.50 - 5.27 (6 H, m) 2.75 - 2.86 (1 H, m) 1.53 - 1.61 (1 H, m) 1.43 - 1.53 (4 H, m) 1.26 - 1.38 (6 H, m) 1.11 - 1.18 (1 H, m) 1.08 (3 H, d, *J*=6.84 Hz) 0.96 - 1.05 (1 H, m) 0.84 - 0.92 (1 H, m)
108-2	598.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.91 (1 H, s) 8.09 - 8.14 (1 H, m) 7.26 - 7.31 (1 H, m) 7.15 - 7.21 (2 H, m) 7.04 - 7.09 (1 H, m) 6.50 - 6.67 (1 H, m) 6.38 (1 H, br t, *J*=14.93 Hz) 5.75 - 5.81 (1 H, m) 3.37 - 5.21 (6 H, m) 2.67 - 2.78 (1 H, m) 2.00 (3 H, s) 1.38 - 1.55 (5 H, m) 1.19 - 1.36 (6 H, m) 0.70 - 1.09 (6 H, m)
108-2-1	598.1	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.91 (1 H, s) 8.13 (1 H, s) 7.26 - 7.31 (1 H, m) 7.16 - 7.21 (2 H, m) 7.06 (1 H, br d, *J*=7.88 Hz) 6.50 - 6.67 (1 H, m) 6.38 (1 H, br t, *J*=15.45 Hz) 5.75 - 5.82 (1 H, m) 3.38 - 5.15 (6 H, m) 2.67 - 2.78 (1 H, m) 2.00 (3 H, s) 1.40 - 1.53 (5 H, m) 1.31 - 1.37 (2 H, m) 1.15 - 1.27 (4 H, m) 0.88 - 1.07 (5 H, m) 0.71 - 0.79 (1 H, m)
108-2-2	598.3	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.93 (1 H, s) 8.11 - 8.15 (1 H, m) 7.29 - 7.34 (1 H, m) 7.19 - 7.24 (2 H, m) 7.10 (1 H, d, *J*=7.67 Hz) 6.53 - 6.69 (1 H, m) 6.41 (1 H, br t, *J*=14.93 Hz) 5.78 - 5.84 (1 H, m) 3.48 - 5.25 (6 H, m) 2.68 - 2.82 (1 H, m) 2.03 (3 H, s) 1.39 - 1.57 (5 H, m) 1.30 - 1.35 (2 H, m) 1.17 - 1.28 (4 H, m) 0.91 - 1.11 (5 H, m) 0.74 - 0.84 (1 H, m)
109-1	590.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.11 (1 H, s) 8.10 - 8.14 (1 H, m) 7.40 - 7.47 (1 H, m) 7.08 - 7.20 (3 H, m) 6.51 - 6.70 (1 H, m) 6.41 (1 H, br t, *J*=14.93 Hz) 5.77 - 5.85 (1 H, m) 3.44 - 5.21 (6 H, m) 2.65 - 2.77 (1 H, m) 2.48 - 2.59 (1 H, m) 2.36 - 2.49 (1 H, m) 1.40 - 1.52 (4 H, m) 1.30 - 1.36 (2 H, m) 1.14 - 1.28 (6 H, m) 1.02 - 1.09 (3 H, m)
109-1-1	590.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.19 (1 H, s) 8.20 (1 H, s) 7.47 - 7.56 (1 H, m) 7.17 - 7.28 (3 H, m) 6.61 - 6.77 (1 H, m) 6.49 (1 H, br t, *J*=15.03 Hz) 5.86 - 5.93 (1 H, m) 3.54 - 5.30 (6 H, m) 2.72 - 2.85 (1 H, m) 2.44 - 2.69 (2 H, m) 1.48 - 1.60 (4 H, m) 1.41 (2 H, br d, *J*=6.84 Hz) 1.32 (3 H, d, *J*=6.84 Hz) 1.25 (3 H, t, *J*=7.57 Hz) 1.14 (3 H, d, *J*=6.63 Hz)
109-1-2	590.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.19 (1 H, s) 8.20 (1 H, s) 7.48 - 7.55 (1 H, m) 7.16 - 7.28 (3 H, m) 6.59 - 6.77 (1 H, m) 6.49 (1 H, br t, *J*=14.82 Hz) 5.85 - 5.93 (1

		H, m) 3.51 - 5.28 (6 H, m) 2.74 - 2.85 (1 H, m) 2.44 - 2.69 (2 H, m) 1.49 - 1.61 (4 H, m) 1.42 (2 H, d, *J*=6.84 Hz) 1.31 (3 H, d, *J*=6.63 Hz) 1.27 (3 H, t, *J*=7.57 Hz) 1.12 (3 H, d, *J*=6.84 Hz)
109-2	586.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.06 (1 H, s) 8.11 (1 H, s) 7.26 - 7.32 (1 H, m) 7.15 - 7.21 (2 H, m) 6.99 - 7.05 (1 H, m) 6.50 - 6.67 (1 H, m) 6.38 (1 H, br t, *J*=14.93 Hz) 5.75 - 5.82 (1 H, m) 3.43 - 5.17 (6 H, m) 2.62 - 2.75 (1 H, m) 2.45 - 2.58 (1 H, m) 2.30 - 2.45 (1 H, m) 1.98 (3 H, s) 1.39 - 1.50 (4 H, m) 1.29 - 1.35 (2 H, m) 1.11 - 1.26 (6 H, m) 0.97 - 1.04 (3 H, m)
109-2-1	586.1	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.06 (1 H, s) 8.11 (1 H, s) 7.26 - 7.32 (1 H, m) 7.15 - 7.21 (2 H, m) 7.00 - 7.04 (1 H, m) 6.50 - 6.67 (1 H, m) 6.33 - 6.43 (1 H, m) 5.76 - 5.82 (1 H, m) 3.43 - 5.17 (6 H, m) 2.63 - 2.74 (1 H, m) 2.45 - 2.57 (1 H, m) 2.31 - 2.45 (1 H, m) 1.98 (3 H, s) 1.39 - 1.50 (4 H, m) 1.32 (2 H, br d, *J*=6.84 Hz) 1.21 (3 H, d, *J*=6.63 Hz) 1.13 (3 H, t, *J*=7.57 Hz) 0.98 - 1.03 (3 H, m)
109-2-2	586.3	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 9.06 (1 H, s) 8.11 (1 H, s) 7.26 - 7.32 (1 H, m) 7.14 - 7.21 (2 H, m) 7.03 (1 H, d, *J*=7.05 Hz) 6.50 - 6.67 (1 H, m) 6.38 (1 H, br t, *J*=15.24 Hz) 5.75 - 5.82 (1 H, m) 3.43 - 5.17 (6 H, m) 2.63 - 2.74 (1 H, m) 2.45 - 2.57 (1 H, m) 2.28 - 2.43 (1 H, m) 1.98 (3 H, s) 1.37 - 1.50 (4 H, m) 1.31 (2 H, br d, *J*=6.84 Hz) 1.20 (3 H, d, *J*=6.84 Hz) 1.16 (3 H, t, *J*=7.46 Hz) 0.99 (3 H, d, *J*=6.84 Hz)
110-1	512.3	^{1}H NMR (400 MHz, DMSO-*d*$_6$) δ 8.33 (s, 1H), 7.53-7.65 (m, 2H), 7.37-7.46 (m, 2H), 6.76-6.87 (m, 1H), 6.09-6.25 (m, 1H), 5.76 (s, 1H), 4.37-4.88 (m, 2H), 4.19-4.30 (m, 1H), 4.03-4.18 (m, 2H), 3.72-3.93 (m, 2H), 3.45 (br d, *J*=12.02 Hz, 1H), 1.23-1.31 (m, 3H), 1.08-1.21 (m, 3H), 0.88 (s, 9H). ^{19}F NMR (376 MHz, DMSO-*d*$_6$) δ -113.60 (s, 1F)
110-2	524.3	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 7.97 (s, 1H), 7.49-7.60 (m, 2H), 7.20-7.36 (m, 2H), 6.49-6.69 (m, 1H), 6.31-6.43 (m, 1H), 5.73-5.81 (m, 1H), 5.03-5.15 (m, 1H), 4.99 (br s, 1H), 4.79 (br s, 0.5H), 4.26-4.50 (m, 1H), 4.07-4.23 (m, 0.5H), 3.33-4.01 (m, 3H), 2.54-2.70 (m, 2H), 1.83 (br d, *J*=12.44 Hz, 2H), 1.60-1.77 (m, 3H), 1.32-1.45 (m, 6H), 1.24-1.32 (m, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-*d*) δ -112.30 (s, 1F)
119-1	571.3	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.57 (d, *J*=4.98 Hz, 1H), 7.84 (dd, *J*=4.15, 8.91 Hz, 1H), 7.40-7.51 (m, 1H), 7.27-7.30 (m, 1H), 7.08-7.23 (m, 3H), 4.79-5.12 (m, 2H), 4.06-4.41 (m, 2H), 3.41-3.92 (m, 2H), 2.76 (dt, *J*=5.08, 6.58 Hz, 1H), 2.08-2.10 (m, 3H), 2.04-2.06 (m, 3H), 1.42 (t, *J*=7.26 Hz, 3H), 1.24-1.3 (m, 6H), 1.11 (dd, *J*=2.07, 6.84 Hz, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.58 (dd, *J*=16.04, 40.31 Hz, 1F), -126.38 (br dd, *J*=26.44, 40.31 Hz, 1F)

119-2	614.2	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.52 (d, *J*=4.77 Hz, 1H), 7.82 (dd, *J*=4.46, 9.02 Hz, 1H), 7.39-7.49 (m, 1H), 7.26-7.30 (m, 1H), 7.05-7.22 (m, 3H), 4.75-5.20 (m, 2H), 4.23-4.46 (m, 1H), 4.18 (br d, *J*=14.31 Hz, 1H), 4.06 (br d, *J*=14.10 Hz, 1H), 3.83-3.99 (m, 2H), 3.50 (d, *J*=3.11 Hz, 1H), 2.64-2.75 (m, 1H), 2.36 (d, *J*=3.94 Hz, 6H), 2.04 (d, *J*=2.70 Hz, 3H), 1.43 (dd, *J*=1.45, 6.84 Hz, 3H), 1.31 (d, *J*=6.84 Hz, 3H), 1.24 (d, *J*=6.84 Hz, 3H), 1.08 (dd, *J*=2.07, 6.63 Hz, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.45 (dd, *J*=19.07, 41.61 Hz, 1F), -126.70 (dd, *J*=28.61, 41.62 Hz, 1F)
121-1	632.4	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.15 (s, 1H), 7.79-8.04 (m, 1H), 7.40-7.49 (m, 1H), 7.08-7.22 (m, 3H), 6.32-6.72 (m, 2H), 5.72-5.94 (m, 1H), 4.96-5.19 (m, 1H), 4.56-4.76 (m, 1H), 3.58-4.49 (m, 4H), 3.09 (dq, *J*=4.35, 7.39 Hz, 1H), 2.76-3.01 (m, 6H), 2.16 (br s, 2H), 1.53-1.55 (m, 3H), 1.44-1.48 (m, 3H), 1.33 (br d, *J*=6.84 Hz, 6H), 1.14-1.21 (m, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.77 (br dd, *J*=7.37, 11.70 Hz, 1F)
121-2	658.3	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 7.94-8.10 (m, 1H), 7.28-7.40 (m, 1H), 6.97-7.16 (m, 4H), 6.42-6.65 (m, 1H), 6.27-6.40 (m, 1H), 5.73 (br t, *J*=8.60 Hz, 1H), 4.80-5.14 (m, 2H), 4.16-4.43 (m, 1H), 3.58-4.00 (m, 5H), 2.44-2.67 (m, 5H), 1.93 (s, 3H), 1.74 (br d, *J*=2.70 Hz, 3H), 1.54 (br s, 3H), 1.31-1.42 (m, 3H), 1.13 (dd, *J*=4.35, 6.43 Hz, 3H), 0.89-1.01 (m, 4H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.71--111.79 (m, 1F)
121-3	631.6	^{1}H NMR (400MHz, CDCl3) δ = 8.07 (br s, 1H), 7.48 - 7.28 (m, 1H), 7.22 - 7.02 (m, 5H), 6.71 - 6.50 (m, 1H), 6.47 - 6.33 (m, 1H), 5.79 (br d, *J*=9.3 Hz, 1H), 5.28 - 4.85 (m, 2H), 4.56 - 4.17 (m, 1H), 4.08 - 3.62 (m, 3H), 3.58 - 3.28 (m, 2H), 2.60 - 2.38 (m, 1H), 2.25 (s, 6H), 1.98 (br s, 3H), 1.43 (br d, *J*=5.8 Hz, 3H), 1.37 - 1.23 (m, 3H), 1.19 (br d, *J*=6.6 Hz, 3H), 1.01 (br d, *J*=6.6 Hz, 3H); ^{19}F NMR (376MHz, CDCl3) δ = -111.70 (s, 1F)
121-4	631.6	^{1}H NMR (400MHz, CDCl3) δ = 8.17 - 8.11 (m, 1H), 7.60 - 7.38 (m, 1H), 7.29 - 7.12 (m, 5H), 6.78 - 6.59 (m, 1H), 6.54 - 6.42 (m, 1H), 5.92 - 5.83 (m, 1H), 5.34 - 4.89 (m, 2H), 4.60 - 4.30 (m, 1H), 4.17 - 3.70 (m, 3H), 3.70 - 3.36 (m, 2H), 2.66 (td, *J*=6.8, 13.6 Hz, 1H), 2.34 (s, 6H), 2.05 (s, 3H), 1.55 - 1.47 (m, 3H), 1.42 - 1.33 (m, 3H), 1.25 (br d, *J*=6.8 Hz, 3H), 1.05 (br d, *J*=6.8 Hz, 3H); ^{19}F NMR (376MHz, CDCl3) δ = -111.77 (s, 1F)
122-1	714.4	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.59 (s, 1 H), 8.07 - 8.18 (m, 1 H), 7.59 (br dd, *J*=7.8, 6.5 Hz, 1 H), 7.37 (br d, *J*=8.9 Hz, 1 H), 7.17 - 7.30 (m, 1 H), 6.84 (dt, *J*=16.7, 10.7 Hz, 1 H), 6.57 - 6.72 (m, 1 H), 6.19 (dd, *J*=16.5, 2.2 Hz, 1 H), 5.74 (br d, *J*=2.3 Hz, 1 H), 4.81 - 5.05 (m, 1 H), 4.66 - 4.76 (m, 1 H), 4.45 - 4.57 (m, 1 H), 4.04 - 4.29 (m, 2 H),

		3.67 - 3.79 (m, 1 H), 2.72 (s, 7 H), 1.21 - 1.31 (m, 4 H), 1.00 - 1.12 (m, 3 H), 0.92 (br dd, *J*=6.6, 2.5 Hz, 3 H), 0.78 (br d, *J*=6.6 Hz, 3 H)
122-2	675.4	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.98 - 9.25 (m, 1 H), 8.36 - 8.59 (m, 1 H), 8.04 - 8.16 (m, 1 H), 7.37 - 7.49 (m, 1 H), 7.03 - 7.17 (m, 1 H), 6.78 - 6.93 (m, 1 H), 6.63 (dd, *J*=15.3, 6.0 Hz, 1 H), 6.20 (br d, *J*=16.6 Hz, 1 H), 5.76 (br dd, *J*=10.0, 2.1 Hz, 1 H), 4.72 - 4.98 (m, 2 H), 4.44 - 4.54 (m, 1 H), 4.01 - 4.21 (m, 2 H), 3.79 - 3.95 (m, 2 H), 2.70 - 2.77 (m, 6 H), 2.12 - 2.27 (m, 1 H), 1.91 - 2.11 (m, 2 H), 1.18 - 1.41 (m, 4 H), 1.01 - 1.17 (m, 3 H), 0.86 - 0.99 (m, 4 H), 0.78 - 0.85 (m, 2 H), 0.75 (br d, *J*=6.4 Hz, 2 H)
122-2-1	674.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.99 - 9.13 (m, 1 H), 8.38 - 8.56 (m, 1 H), 8.10 (dd, *J*=10.8, 5.8 Hz, 1 H), 7.35 - 7.48 (m, 2 H), 7.08 (q, *J*=8.6 Hz, 1 H), 6.73 - 6.92 (m, 1 H), 6.59 - 6.66 (m, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.71 - 5.81 (m, 1 H), 4.72 - 4.99 (m, 2 H), 4.12 - 4.24 (m, 1 H), 3.83 - 3.97 (m, 3 H), 2.75 (s, 6 H), 1.98 - 2.12 (m, 1 H), 1.23 - 1.39 (m, 3 H), 1.12 - 1.23 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.86 - 0.94 (m, 3 H), 0.82 (t, *J*=7.5 Hz, 2 H), 0.75 (d, *J*=6.6 Hz, 3 H)
122-2-2	675.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 9.19 (br d, *J*=11.6 Hz, 1 H), 8.53 (br d, *J*=5.2 Hz, 1 H), 8.12 (d, *J*=5.8 Hz, 1 H), 7.37 - 7.52 (m, 1 H), 7.31 - 7.36 (m, 1 H), 7.12 (t, *J*=8.9 Hz, 1 H), 6.79 - 6.91 (m, 1 H), 6.65 (d, *J*=5.8 Hz, 1 H), 6.20 (br d, *J*=16.4 Hz, 1 H), 5.72 - 5.83 (m, 1 H), 4.72 - 4.99 (m, 1 H), 4.43 - 4.54 (m, 1 H), 4.02 - 4.19 (m, 1 H), 3.89 (br s, 1 H), 3.81 (br d, *J*=14.7 Hz, 1 H), 2.73 (s, 6 H), 2.19 (dt, *J*=15.7, 7.0 Hz, 2 H), 1.95 - 2.02 (m, 2 H), 1.27 (br dd, *J*=14.8, 6.7 Hz, 3 H), 1.13 (br dd, *J*=11.7, 6.5 Hz, 2 H), 1.03 - 1.07 (m, 2 H), 0.92 (d, *J*=6.6 Hz, 3 H), 0.71 - 0.88 (m, 5 H)
122-3	687.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 9.18 - 9.70 (m, 1 H), 8.50 (d, *J*=10.4 Hz, 1 H), 8.03 - 8.14 (m, 1 H), 7.26 - 7.50 (m, 2 H), 7.03 - 7.17 (m, 1 H), 6.83 (br dd, *J*=16.1, 10.3 Hz, 1 H), 6.59 - 6.72 (m, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.71 - 5.83 (m, 1 H), 4.73 - 4.98 (m, 2 H), 4.47 - 4.56 (m, 1 H), 4.00 - 4.25 (m, 2 H), 3.79 (br d, *J*=13.1 Hz, 1 H), 2.72 - 2.78 (m, 6 H), 2.68 (br d, *J*=1.7 Hz, 1 H), 1.37 (br dd, *J*=5.9, 3.4 Hz, 1 H), 1.19 - 1.34 (m, 4 H), 1.11 - 1.17 (m, 2 H), 1.02 - 1.11 (m, 3 H), 0.91 (br d, *J*=6.4 Hz, 4 H), 0.74 (dd, *J*=6.6, 2.3 Hz, 3 H)
122-3-1	687.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 9.27 - 9.46 (m, 1 H), 8.49 (s, 1 H), 8.40 (d, *J*=4.8 Hz, 1 H), 8.10 (dd, *J*=11.8, 5.8 Hz, 1 H), 7.40 - 7.50 (m, 1 H), 7.01 - 7.13 (m, 1 H), 6.78 - 6.91 (m, 1 H), 6.63 (dd, *J*=5.8, 3.5 Hz, 1 H), 6.20 (dd, *J*=16.8, 2.3 Hz, 1 H), 5.71 - 5.83 (m, 1 H), 4.71 - 4.99 (m, 2 H), 4.11 - 4.26 (m, 1 H), 4.00 - 4.09 (m, 1 H), 3.81 - 3.97 (m, 2 H), 3.68 - 3.79 (m, 1 H), 2.77 (s, 7 H), 1.18 - 1.42 (m, 5 H), 1.02 - 1.12 (m, 2 H), 0.89 - 0.97 (m, 3 H),

		0.74 (d, *J*=6.4 Hz, 3 H), 0.60 - 0.71 (m, 2 H), 0.40 - 0.56 (m, 1 H)
122-3-2	687.4	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 9.59 (br d, *J*=13.9 Hz, 1 H), 8.51 (s, 1 H), 8.12 (d, *J*=5.8 Hz, 1 H), 7.40 - 7.50 (m, 1 H), 7.27 - 7.35 (m, 1 H), 7.12 (t, *J*=8.8 Hz, 1 H), 6.77 - 6.92 (m, 1 H), 6.66 (d, *J*=5.6 Hz, 1 H), 6.20 (br d, *J*=16.0 Hz, 1 H), 5.66 - 5.82 (m, 1 H), 4.72 - 4.98 (m, 2 H), 4.01 - 4.18 (m, 2 H), 3.91 (br s, 1 H), 3.79 (br d, *J*=13.7 Hz, 1 H), 2.74 (s, 7 H), 1.27 (br dd, *J*=12.6, 6.4 Hz, 5 H), 1.03 - 1.19 (m, 3 H), 0.91 (d, *J*=6.6 Hz, 3 H), 0.73 (d, *J*=6.6 Hz, 3 H), 0.51 - 0.68 (m, 2 H), 0.30 - 0.41 (m, 1 H)
123-1	590.2	^{1}H NMR (CDCl3, 400MHz): δ = 9.79 (br s, 1H), 8.05 (d, J=9.7 Hz, 1H), 7.25-7.32 (m, 1H), 6.57-6.76 (m, 3H), 6.34-6.43 (m, 2H), 5.78 (dd, J=10.5, 1.8 Hz, 1H), 4.68 (br s, 2H), 4.50 (s, 2H), 4.29 (br d, J=13.5 Hz, 2H), 3.60 (br dd, J=13.5, 4.1 Hz, 2H), 2.56 (dt, J=13.3, 6.7 Hz, 1H), 1.90 (s, 3H), 1.46-1.61 (m, 6H), 1.14 (d, J=6.6 Hz, 3H), 0.97 (d, J=6.8 Hz, 3H). ^{19}F NMR (CDCl3, 376MHz): δ = -107.30 (d, *J*=82.4 Hz, 1F), -121.70 (d, *J*=83.2 Hz, 1F)
123-2	590.2	^{1}H NMR (400MHz, CDCl3) δ = 9.79 (br s, 1H), 8.05 (d, *J*=9.7 Hz, 1H), 7.32 - 7.25 (m, 1H), 6.76 - 6.57 (m, 3H), 6.43 - 6.34 (m, 2H), 5.78 (dd, *J*=1.8, 10.5 Hz, 1H), 4.68 (br s, 2H), 4.50 (s, 2H), 4.29 (br d, *J*=13.5 Hz, 2H), 3.60 (br dd, *J*=4.1, 13.5 Hz, 2H), 2.56 (td, *J*=6.7, 13.3 Hz, 1H), 1.90 (s, 3H), 1.58 (d, *J*=7.0 Hz, 3H), 1.52 (d, *J*=7.0 Hz, 3H), 1.14 (d, *J*=6.6 Hz, 3H), 0.97 (d, *J*=6.8 Hz, 3H); ^{19}F NMR (376MHz, CDCl3) δ = -107.30 (d, *J*=82.4 Hz, 1F), -121.70 (d, *J*=83.2 Hz, 1F)
123-3	574.1	^{1}H NMR (400MHz, CDCl3) δ = 7.99 (d, *J*=9.1 Hz, 1H), 7.52 - 7.40 (m, 1H), 7.34 (dt, *J*=1.7, 7.4 Hz, 1H), 7.23 - 7.11 (m, 2H), 6.69 - 6.60 (m, 1H), 6.52 (s, 1H), 6.32 (s, 1H), 5.79 (dd, *J*=1.9, 10.4 Hz, 1H), 4.84 - 4.39 (m, 4H), 4.30 (br d, *J*=13.5 Hz, 2H), 3.57 (ddd, *J*=4.3, 4.4, 13.4 Hz, 2H), 2.63 - 2.52 (m, 1H), 1.91 (s, 3H), 1.64 - 1.50 (m, 6H) 1.17 (d, *J*=6.8 Hz, 3H), 1.00 (d, *J*=6.6 Hz, 3H); ^{19}F NMR (376MHz, CDCl3) δ = -112.60 (d, *J*=40.7 Hz, 1F), -125.87 - -127.76 (m, 1F)
123-4	574.1	^{1}H NMR (400MHz, CDCl3) δ = 7.99 (d, *J*=9.1 Hz, 1H), 7.50 - 7.29 (m, 2H), 7.24 - 7.10 (m, 2H), 6.71 - 6.57 (m, 1H), 6.43 (dd, *J*=2.0, 16.7 Hz, 1H), 6.31 (s, 1H), 5.80 (dd, *J*=1.9, 10.4 Hz, 1H), 4.71 (br s, 2H), 4.45 (br s, 2H), 4.30 (br d, *J*=13.3 Hz, 2H), 3.57 (td, *J*=4.2, 13.3 Hz, 2H), 2.57 (s, 1H), 1.90 (s, 3H), 1.57 (t, *J*=6.7 Hz, 6H), 1.16 (d, *J*=6.6 Hz, 3H), 0.99 (d, *J*=6.8 Hz, 3H); ^{19}F NMR (376MHz, CDCl3) δ = -112.60 (d, *J*=41.6 Hz, 1F), -127.13 (d, *J*=41.6 Hz, 1F)
123-5	589.1	^{1}H NMR (400MHz, CDCl3) δ = 8.00 (d, *J*=9.3 Hz, 1H), 7.14 (br d, *J*=6.4 Hz, 1H), 6.64 (dd, *J*=10.5, 16.7 Hz, 1H), 6.51 - 6.39 (m, 3H), 6.34 (s, 1H), 5.79 (dd, *J*=1.7, 10.4 Hz, 1H), 4.83 - 4.40 (m, 6H), 4.31 (br d, *J*=12.9 Hz, 2H), 3.59

		(ddd, *J*=3.9, 4.0, 13.4 Hz, 2H), 2.68 - 2.55 (m, 1H), 1.92 (s, 3H), 1.60 (br d, *J*=6.8 Hz, 3H), 1.55 (d, *J*=6.8 Hz, 3H), 1.17 (br d, *J*=6.6 Hz, 3H), 0.99 (br d, *J*=6.6 Hz, 3H); ^{19}F NMR (376MHz, CDCl3) δ = -110.99 (d, *J*=56.4 Hz, 1F), -123.22 (br d, *J*=56.4 Hz, 1F)
123-6	589.1	^{1}H NMR (400MHz, CDCl3) δ = 8.00 (d, *J*=9.3 Hz, 1H), 7.14 (dt, *J*=6.4, 8.1 Hz, 1H), 6.68 - 6.60 (m, 1H), 6.51 - 6.39 (m, 3H), 6.33 (s, 1H), 5.79 (dd, *J*=1.9, 10.6 Hz, 1H), 4.84 - 4.45 (m, 6H), 4.31 (br d, *J*=13.1 Hz, 2H), 3.59 (ddd, *J*=4.0, 4.1, 13.4 Hz, 2H), 2.70 - 2.52 (m, 1H), 1.91 (s, 3H), 1.60 (d, *J*=6.8 Hz, 3H), 1.55 (d, *J*=6.8 Hz, 3H), 1.16 (d, *J*=6.8 Hz, 3H), 0.98 (d, *J*=6.6 Hz, 3H); ^{19}F NMR (376MHz, CDCl3) δ = -110.99 (d, *J*=56.4 Hz, 1F), -123.22 (br d, *J*=56.4 Hz, 1F)
124-1	602.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.42 - 8.51 (m, 2 H), 7.45 - 7.58 (m, 1 H), 7.21 - 7.38 (m, 3 H), 7.18 (td, *J*=7.4, 1.6 Hz, 1 H), 6.84 (td, *J*=15.8, 10.5 Hz, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.72 - 5.82 (m, 1 H), 4.73 - 4.97 (m, 2 H), 4.47 (br dd, *J*=6.1, 3.4 Hz, 1 H), 4.12 - 4.23 (m, 1 H), 3.83 - 3.96 (m, 2 H), 3.51 (br dd, *J*=13.6, 3.6 Hz, 1 H), 1.31 - 1.40 (m, 3 H), 1.26 (br d, *J*=6.6 Hz, 2 H), 1.18 (d, *J*=6.6 Hz, 2 H), 1.07 (dd, *J*=6.6, 4.4 Hz, 6 H), 0.93 (t, *J*=7.7 Hz, 6 H)
124-2	599.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.33 - 8.60 (m, 2 H), 7.18 - 7.37 (m, 4 H), 7.02 (d, *J*=7.3 Hz, 1 H), 6.76 - 6.93 (m, 1 H), 6.20 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.76 (ddd, *J*=10.3, 5.7, 2.3 Hz, 1 H), 4.72 - 4.96 (m, 2 H), 4.48 (dd, *J*=5.1, 1.3 Hz, 1 H), 4.11 - 4.24 (m, 1 H), 3.83 - 3.99 (m, 2 H), 3.46 - 3.57 (m, 1 H), 1.95 (s, 3 H), 1.30 - 1.38 (m, 3 H), 1.25 (br d, *J*=6.6 Hz, 2 H), 1.18 (d, *J*=6.6 Hz, 2 H), 1.07 (t, *J*=6.3 Hz, 6 H), 0.93 (t, *J*=6.5 Hz, 6 H)
124-3	653.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.59 (br d, *J*=4.6 Hz, 1 H), 8.43 (d, *J*=5.2 Hz, 1 H), 7.78 (br dd, *J*=19.3, 7.7 Hz, 2 H), 7.64 - 7.71 (m, 1 H), 7.34 (br d, *J*=8.1 Hz, 1 H), 7.24 (d, *J*=5.0 Hz, 1 H), 6.72 - 6.90 (m, 1 H), 6.14 - 6.25 (m, 1 H), 5.76 (ddd, *J*=10.4, 5.9, 2.0 Hz, 1 H), 4.72 - 5.01 (m, 2 H), 4.43 - 4.55 (m, 1 H), 4.09 - 4.34 (m, 1 H), 3.94 - 4.09 (m, 1 H), 3.76 - 3.93 (m, 2 H), 3.39 - 3.64 (m, 1 H), 1.28 - 1.41 (m, 3 H), 1.13 - 1.28 (m, 2 H), 1.06 (br d, *J*=5.8 Hz, 7 H), 0.92 (br d, *J*=6.6 Hz, 6 H)
124-4	619.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 9.92 - 10.32 (m, 1 H), 8.45 (d, *J*=5.0 Hz, 2 H), 7.19 - 7.29 (m, 2 H), 6.84 (td, *J*=15.7, 10.9 Hz, 1 H), 6.60 - 6.72 (m, 2 H), 6.20 (dd, *J*=16.8, 2.3 Hz, 1 H), 5.71 - 5.80 (m, 1 H), 4.73 - 4.96 (m, 2 H), 4.43 - 4.55 (m, 1 H), 4.08 - 4.24 (m, 2 H), 3.84 - 3.97 (m, 2 H), 1.30 - 1.37 (m, 3 H), 1.27 (br d, *J*=6.4 Hz, 2 H), 1.19 (br d, *J*=6.6 Hz, 2 H), 1.07 (dd, *J*=6.6, 4.8 Hz, 6 H), 0.93 (br s, 6 H)

124-5	583.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.48 (d, *J*=5.2 Hz, 1 H), 8.33 (d, *J*=9.5 Hz, 1 H), 7.31 - 7.39 (m, 1 H), 7.22 - 7.29 (m, 3 H), 7.12 - 7.19 (m, 1 H), 6.78 - 6.95 (m, 1 H), 6.20 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.72 - 5.81 (m, 1 H), 4.72 - 4.96 (m, 2 H), 4.48 (br dd, *J*=5.0, 2.3 Hz, 1 H), 4.11 - 4.23 (m, 1 H), 3.82 - 3.94 (m, 2 H), 2.60 - 2.67 (m, 1 H), 1.98 (s, 3 H), 1.29 - 1.37 (m, 3 H), 1.26 (br d, *J*=6.8 Hz, 2 H), 1.18 (d, *J*=6.8 Hz, 2 H), 1.07 (d, *J*=6.8 Hz, 6 H), 0.91 (br t, *J*=7.7 Hz, 6 H)
124-6	618.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.51 (s, 1 H), 8.45 (d, *J*=5.0 Hz, 1 H), 7.51 - 7.59 (m, 1 H), 7.36 - 7.49 (m, 2 H), 7.27 (d, *J*=5.2 Hz, 1 H), 7.17 (dd, *J*=7.5, 1.7 Hz, 1 H), 6.71 - 6.94 (m, 1 H), 6.20 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 4.73 - 4.97 (m, 2 H), 4.42 - 4.54 (m, 1 H), 4.10 - 4.21 (m, 1 H), 3.80 - 3.98 (m, 2 H), 2.60 - 2.67 (m, 1 H), 1.34 (t, *J*=6.8 Hz, 3 H), 1.25 (br d, *J*=6.6 Hz, 2 H), 1.18 (br d, *J*=6.6 Hz, 2 H), 1.07 (t, *J*=6.1 Hz, 6 H), 0.95 (t, *J*=7.6 Hz, 6 H)
124-7	584.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.52 (d, *J*=5.2 Hz, 1 H), 8.45 (d, *J*=1.5 Hz, 1 H), 7.47 - 7.52 (m, 2 H), 7.39 - 7.46 (m, 3 H), 7.33 (d, *J*=5.0 Hz, 1 H), 6.67 - 6.98 (m, 1 H), 6.20 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.64 - 5.83 (m, 1 H), 4.71 - 4.97 (m, 2 H), 4.40 - 4.55 (m, 1 H), 4.11 - 4.24 (m, 1 H), 3.80 - 3.97 (m, 2 H), 2.62 - 2.72 (m, 2 H), 1.34 (t, *J*=6.5 Hz, 3 H), 1.25 (br d, *J*=6.6 Hz, 1 H), 1.18 (d, *J*=6.8 Hz, 2 H), 1.08 (t, *J*=6.4 Hz, 6 H), 0.94 (d, *J*=6.8 Hz, 3 H), 0.90 (d, *J*=6.8 Hz, 3 H)
124-8	626.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.43 (d, *J*=5.0 Hz, 2 H), 7.38 (d, *J*=3.9 Hz, 2 H), 7.17 - 7.29 (m, 2 H), 6.95 - 7.04 (m, 1 H), 6.76 - 6.92 (m, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.74 - 5.82 (m, 1 H), 4.72 - 4.97 (m, 2 H), 4.44 - 4.55 (m, 1 H), 4.11 - 4.28 (m, 1 H), 3.78 - 3.98 (m, 2 H), 1.35 (br d, *J*=6.8 Hz, 3 H), 1.14 - 1.30 (m, 4 H), 1.01 - 1.13 (m, 10 H), 0.87 - 0.99 (m, 10 H)
124-9	596.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.47 (s, 1 H), 8.35 (d, *J*=5.2 Hz, 1 H), 7.29 - 7.36 (m, 1 H), 7.18 - 7.28 (m, 2 H), 7.10 (d, *J*=7.7 Hz, 1 H), 6.81 - 6.92 (m, 1 H), 6.78 (d, *J*=5.2 Hz, 1 H), 6.20 (dd, *J*=16.8, 2.3 Hz, 1 H), 5.73 - 5.80 (m, 1 H), 4.71 - 4.96 (m, 2 H), 4.43 - 4.54 (m, 1 H), 4.15 (br dd, *J*=13.2, 8.4 Hz, 1 H), 3.82 - 3.98 (m, 2 H), 2.66 - 2.77 (m, 1 H), 1.98 (s, 3 H), 1.43 - 1.53 (m, 1 H), 1.33 (t, *J*=6.5 Hz, 3 H), 1.25 (br d, *J*=6.4 Hz, 2 H), 1.18 (d, *J*=6.8 Hz, 2 H), 1.07 (d, *J*=6.6 Hz, 3 H), 0.91 (d, *J*=6.6 Hz, 3 H), 0.78 - 0.87 (m, 1 H), 0.69 - 0.77 (m, 2 H)
124-10	584.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.39 - 8.50 (m, 2 H), 7.29 - 7.34 (m, 1 H), 7.20 - 7.27 (m, 3 H), 6.99 (d, *J*=7.7 Hz, 1 H), 6.81 - 6.96 (m, 1 H), 6.22 (br d, *J*=17.0 Hz, 1 H), 5.74 - 5.82 (m, 1 H), 4.98 (br d, *J*=3.1 Hz, 1 H), 4.26 - 4.46 (m, 2 H), 4.01 - 4.22 (m, 1 H), 3.63 - 3.88 (m, 2 H), 3.46 - 3.59 (m, 1 H), 3.08 - 3.21 (m, 1 H), 2.62 - 2.71 (m,

		1 H), 1.95 (s, 3 H), 1.35 (d, *J*=6.6 Hz, 3 H), 1.07 (d, *J*=6.6 Hz, 6 H), 0.93 (d, *J*=6.6 Hz, 6 H)
124-11	598.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.54 (s, 1 H), 8.45 (d, *J*=5.0 Hz, 1 H), 7.29 - 7.34 (m, 1 H), 7.22 - 7.25 (m, 1 H), 7.12 - 7.22 (m, 2 H), 7.01 (d, *J*=6.8 Hz, 1 H), 6.77 - 6.88 (m, 1 H), 6.21 (dd, *J*=16.6, 2.5 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 4.55 - 4.66 (m, 2 H), 4.35 (br d, *J*=13.1 Hz, 2 H), 3.66 (br dd, *J*=13.8, 3.2 Hz, 2 H), 2.63 - 2.73 (m, 2 H), 1.95 (s, 3 H), 1.41 (br d, *J*=5.0 Hz, 6 H), 1.07 (d, *J*=6.6 Hz, 6 H), 0.93 (d, *J*=6.4 Hz, 6 H)
124-12	659.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.50 (1 H, d, *J*=3.5 Hz), 8.41 (1 H, d, *J*=4.8 Hz), 7.54 - 7.61 (1 H, m), 7.46 - 7.53 (1 H, m), 7.25 (1 H, br s), 7.20 (1 H, d, *J*=4.8 Hz), 6.84 (1 H, td, *J*=16.7, 10.6 Hz), 6.19 (1 H, dd, *J*=16.7, 2.0 Hz), 5.72 - 5.80 (1 H, m), 4.83 - 4.94 (1 H, m), 4.43 - 4.81 (1 H, m), 4.11 - 4.21 (1 H, m), 3.83 - 3.95 (2 H, m), 3.50 (1 H, br dd, *J*=13.8, 3.4 Hz), 2.65 - 2.78 (1 H, m), 1.94 (3 H, s), 1.33 (3 H, t, *J*=6.7 Hz), 1.15 - 1.27 (3 H, m), 1.06 (3 H, d, *J*=6.6 Hz), 0.96 (3 H, d, *J*=6.6 Hz)
124-13	609.2	^{1}H NMR (400 MHz, DMSO-*d*$_6$) δ 8.41 (d, *J*=4.98 Hz, 1H), 8.37 (s, 1H), 7.20 (d, *J*=4.98 Hz, 1H), 6.94 (br t, *J*=10.06 Hz, 1H), 6.79-6.89 (m, 1H), 6.68-6.79 (m, 1H), 6.19 (dd, *J*=2.18, 16.69 Hz, 1H), 5.71-5.79 (m, 1H), 4.42-4.91 (m, 2H), 3.75-4.21 (m, 4H), 3.47 (br d, *J*=3.73 Hz, 1H), 2.64-2.75 (m, 1H), 1.92 (s, 3H), 1.31 (br t, *J*=7.36 Hz, 3H), 1.14-1.27 (m, 3H), 1.06 (d, *J*=6.63 Hz, 3H), 0.99 (d, *J*=6.63 Hz, 3H)
124-14	575.2	^{1}H NMR (400 MHz, DMSO-d$_6$) δ 10.20 (br s, 1H), 8.46 (dd, *J*=4.35, 9.12 Hz, 1H), 8.39 (dd, *J*=1.87, 4.56 Hz, 1H), 7.27 (q, *J*=7.88 Hz, 1H), 7.18 (d, *J*=4.56 Hz, 1H), 6.64-6.81 (m, 3H), 6.17 (br d, *J*=16.59 Hz, 1H), 5.72 (br d, *J*=11.61 Hz, 1H), 4.59-4.70 (m, 1H), 4.47-4.57 (m, 1H), 4.20-4.37 (m, 1H), 4.07-4.14 (m, 1H), 3.79-3.90 (m, 1H), 3.69-3.78 (m, 1H), 2.55-2.87 (m, 1H), 1.83-1.97 (m, 3H), 1.42 (br dd, *J*=2.90, 6.43 Hz, 3H), 1.24-1.29 (m, 3H), 1.06 (t, *J*=6.63 Hz, 3H), 0.93 (dd, *J*=6.63, 13.68 Hz, 3H)
124-14-1	575.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 10.19 (1 H, br s), 8.46 (1 H, d, *J*=9.1 Hz), 8.39 (1 H, d, *J*=4.8 Hz), 7.23 - 7.32 (1 H, m), 7.18 (1 H, d, *J*=5.0 Hz), 6.64 - 6.81 (3 H, m), 6.17 (1 H, dd, *J*=16.7, 2.2 Hz), 5.72 (1 H, dd, *J*=10.4, 2.3 Hz), 4.65 (1 H, quin, *J*=6.2 Hz), 4.52 (1 H, quin, *J*=6.4 Hz), 4.29 (1 H, ddd, *J*=12.5, 8.3, 3.8 Hz), 4.05 - 4.15 (1 H, m), 3.79 - 3.89 (1 H, m), 3.68 - 3.78 (1 H, m), 2.81 (1 H, quin, *J*=6.7 Hz), 1.85 (3 H, s), 1.40 (6 H, dd, *J*=11.0, 6.6 Hz), 1.07 (3 H, d, *J*=6.6 Hz), 0.95 (3 H, d, *J*=6.8 Hz)
124-14-2	575.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 10.17 (1 H, s), 8.45 (1 H, d, *J*=9.1 Hz), 8.38 (1 H, d, *J*=4.8 Hz), 7.22 - 7.32 (1 H, m), 7.18 (1 H, d, *J*=4.8 Hz), 6.62 - 6.82 (3 H, m), 6.16 (1 H, dd, *J*=16.6, 2.3 Hz), 5.72 (1 H, dd, *J*=10.4, 2.3 Hz), 4.57 - 4.68 (1 H, m), 4.46 - 4.56 (1 H, m), 4.26 (1 H, ddd,

		J=12.4, 8.2, 3.8 Hz), 4.02 - 4.13 (1 H, m), 3.79 - 3.90 (1 H, m), 3.67 - 3.78 (1 H, m), 2.53 - 2.61 (1 H, m), 1.95 (3 H, s), 1.41 (6 H, dd, *J*=13.1, 6.6 Hz), 1.05 (3 H, d, *J*=6.6 Hz), 0.91 (3 H, d, *J*=6.6 Hz)
124-16	575.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 10.20 (br d, *J*=3.73 Hz, 1H), 8.38 (d, *J*=4.77 Hz, 1H), 8.18-8.34 (m, 1H), 7.27 (q, *J*=7.81 Hz, 1H), 7.19 (br d, *J*=4.56 Hz, 1H), 6.77-6.90 (m, 1H), 6.64-6.75 (m, 2H), 6.18 (ddd, *J*=1.97, 6.38, 16.74 Hz, 1H), 5.76 (br d, *J*=10.99 Hz, 1H), 4.59-4.75 (m, 1H), 4.24-4.40 (m, 2H), 3.57-3.76 (m, 2H), 3.35-3.45 (m, 1H), 2.63-2.77 (m, 1H), 1.91 (br s, 3H), 1.36-1.44 (m, 3H), 1.22-1.34 (m, 3H), 1.07 (d, *J*=6.63 Hz, 3H), 0.93 (br d, *J*=6.63 Hz, 3H)
124-16-1	575.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 10.20 (br s, 1H), 8.39 (d, *J*=4.98 Hz, 1H), 8.18-8.33 (m, 1H), 7.23-7.32 (m, 1H), 7.19 (d, *J*=4.77 Hz, 1H), 6.77-6.91 (m, 1H), 6.65-6.76 (m, 2H), 6.18 (ddd, *J*=1.97, 6.27, 16.64 Hz, 1H), 5.76 (br d, *J*=11.82 Hz, 1H), 3.99-4.77 (m, 4H), 3.61-3.71 (m, 1H), 3.35-3.47 (m, 1H), 2.72 (td, *J*=6.66, 13.42 Hz, 1H), 1.91 (s, 3H), 1.41 (d, *J*=6.43 Hz, 3H), 1.16-1.34 (m, 3H), 1.07 (d, *J*=6.84 Hz, 3H), 0.93 (d, *J*=6.84 Hz, 3H)
124-16-2	575.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 10.19 (br s, 1H), 8.38 (d, *J*=4.77 Hz, 1H), 8.20-8.34 (m, 1H), 7.23-7.33 (m, 1H), 7.18 (d, *J*=4.77 Hz, 1H), 6.84 (dt, *J*=10.57, 17.00 Hz, 1H), 6.63-6.76 (m, 2H), 6.18 (ddd, *J*=2.18, 6.17, 16.74 Hz, 1H), 5.76 (br d, *J*=11.61 Hz, 1H), 4.42-4.93 (m, 2H), 4.23-4.38 (m, 2H), 3.58-4.13 (m, 2H), 2.62-2.75 (m, 1H), 1.92 (d, *J*=2.07 Hz, 3H), 1.37 (dd, *J*=2.28, 6.63 Hz, 3H), 1.20-1.35 (m, 3H), 1.04-1.12 (m, 3H), 0.94 (dd, *J*=2.49, 6.63 Hz, 3H)
125-1	603.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.40 - 8.52 (m, 2 H), 7.50 (br s, 1 H), 7.25 - 7.36 (m, 3 H), 7.14 - 7.20 (m, 1 H), 6.84 (td, *J*=16.5, 10.6 Hz, 1 H), 6.20 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.71 - 5.81 (m, 1 H), 4.74 - 4.96 (m, 2 H), 4.44 - 4.54 (m, 1 H), 4.17 (br d, *J*=14.3 Hz, 1 H), 3.82 - 3.95 (m, 2 H), 3.50 (dd, *J*=13.9, 2.9 Hz, 1 H), 1.31 - 1.38 (m, 3 H), 1.26 (br d, *J*=6.6 Hz, 2 H), 1.18 (d, *J*=6.6 Hz, 2 H), 1.07 (dd, *J*=6.7, 3.0 Hz, 6 H), 0.93 (dd, *J*=6.6, 2.9 Hz, 6 H)
125-2	598.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.39 - 8.50 (m, 2 H), 7.18 - 7.37 (m, 4 H), 7.02 (d, *J*=7.0 Hz, 1 H), 6.72 - 6.91 (m, 1 H), 6.20 (dd, *J*=16.8, 2.3 Hz, 1 H), 5.76 (ddd, *J*=10.3, 5.5, 2.3 Hz, 1 H), 4.72 - 4.97 (m, 2 H), 4.43 - 4.54 (m, 1 H), 4.12 - 4.25 (m, 1 H), 3.79 - 3.98 (m, 2 H), 3.46 - 3.56 (m, 1 H), 1.95 (s, 3 H), 1.30 - 1.38 (m, 3 H), 1.26 (br d, *J*=6.6 Hz, 2 H), 1.18 (d, *J*=6.8 Hz, 2 H), 1.07 (dd, *J*=6.6, 4.4 Hz, 6 H), 0.93 (t, *J*=5.8 Hz, 6 H)
125-3	652.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.58 (br d, *J*=1.5 Hz, 1 H), 8.43 (d, *J*=5.2 Hz, 1 H), 7.73 - 7.83 (m, 2 H), 7.65 - 7.71 (m, 1 H), 7.28 - 7.40 (m, 1 H), 7.23 (d, *J*=5.2 Hz, 1

		H), 6.82 (br dd, *J*=16.7, 10.3 Hz, 1 H), 6.20 (dd, *J*=16.4, 1.7 Hz, 1 H), 5.71 - 5.81 (m, 1 H), 4.71 - 5.01 (m, 2 H), 4.46 - 4.57 (m, 1 H), 4.10 - 4.35 (m, 1 H), 3.96 - 4.08 (m, 1 H), 3.75 - 3.93 (m, 2 H), 3.40 - 3.64 (m, 1 H), 1.34 (br d, *J*=17.4 Hz, 4 H), 1.13 - 1.27 (m, 2 H), 1.06 (br d, *J*=5.2 Hz, 6 H), 0.92 (br d, *J*=6.6 Hz, 6 H)
125-4	618.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 9.91 - 10.30 (m, 1 H), 8.46 (d, *J*=5.0 Hz, 2 H), 7.18 - 7.29 (m, 2 H), 6.76 - 6.92 (m, 1 H), 6.61 - 6.75 (m, 2 H), 6.20 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.75 - 5.81 (m, 1 H), 4.74 - 4.97 (m, 2 H), 4.45 - 4.52 (m, 1 H), 4.10 - 4.26 (m, 1 H), 3.80 - 3.95 (m, 2 H), 3.45 - 3.56 (m, 1 H), 1.31 - 1.39 (m, 3 H), 1.27 (br d, *J*=6.6 Hz, 2 H), 1.20 (br d, *J*=6.6 Hz, 2 H), 1.07 (dd, *J*=6.7, 4.0 Hz, 6 H), 0.93 (br s, 6 H)
125-5	583.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.47 (d, *J*=5.2 Hz, 1 H), 8.33 (d, *J*=9.5 Hz, 1 H), 7.30 - 7.38 (m, 1 H), 7.21 - 7.30 (m, 3 H), 7.11 - 7.19 (m, 1 H), 6.85 (br d, *J*=10.8 Hz, 1 H), 6.20 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.76 (ddd, *J*=7.8, 4.9, 2.6 Hz, 1 H), 4.72 - 4.95 (m, 2 H), 4.43 - 4.54 (m, 1 H), 4.16 (br dd, *J*=13.6, 5.9 Hz, 1 H), 3.83 - 3.94 (m, 2 H), 2.61 - 2.67 (m, 1 H), 1.98 (s, 3 H), 1.32 (br t, *J*=7.5 Hz, 3 H), 1.26 (br d, *J*=6.8 Hz, 2 H), 1.18 (d, *J*=6.6 Hz, 2 H), 1.07 (dd, *J*=6.7, 2.4 Hz, 6 H), 0.91 (br t, *J*=5.3 Hz, 6 H)
125-6	618.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.51 (s, 1 H), 8.45 (d, *J*=5.0 Hz, 1 H), 7.51 - 7.58 (m, 1 H), 7.37 - 7.51 (m, 2 H), 7.26 (d, *J*=5.0 Hz, 1 H), 7.17 (dd, *J*=7.4, 1.6 Hz, 1 H), 6.83 (s, 1 H), 6.20 (dd, *J*=16.5, 2.4 Hz, 1 H), 5.75 - 5.79 (m, 1 H), 4.74 - 4.97 (m, 2 H), 4.44 - 4.53 (m, 1 H), 4.11 - 4.22 (m, 2 H), 3.81 - 3.98 (m, 2 H), 1.34 (t, *J*=6.8 Hz, 3 H), 1.26 (br d, *J*=6.4 Hz, 2 H), 1.18 (br d, *J*=6.8 Hz, 2 H), 1.07 (dd, *J*=6.6, 4.6 Hz, 6 H), 0.95 (t, *J*=6.7 Hz, 6 H)
125-7	584.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.53 (d, *J*=5.0 Hz, 1 H), 8.45 (d, *J*=2.1 Hz, 1 H), 7.47 - 7.52 (m, 2 H), 7.40 - 7.46 (m, 3 H), 7.31 (d, *J*=5.2 Hz, 1 H), 6.77 - 6.95 (m, 1 H), 6.20 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 4.74 - 4.95 (m, 2 H), 4.48 (br dd, *J*=5.1, 1.6 Hz, 1 H), 4.08 - 4.25 (m, 2 H), 3.79 - 3.96 (m, 2 H), 1.34 (t, *J*=6.7 Hz, 3 H), 1.25 (br d, *J*=6.6 Hz, 2 H), 1.18 (d, *J*=6.8 Hz, 2 H), 1.08 (dd, *J*=6.5, 5.3 Hz, 6 H), 0.92 (dd, *J*=10.6, 6.8 Hz, 6 H
125-8	627.2	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.45 (br d, *J*=4.8 Hz, 2 H), 7.38 (d, *J*=3.9 Hz, 2 H), 7.28 (br d, *J*=4.6 Hz, 1 H), 7.20 - 7.26 (m, 1 H), 6.99 (br d, *J*=7.9 Hz, 1 H), 6.77 - 6.91 (m, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 4.73 - 4.96 (m, 2 H), 4.45 - 4.57 (m, 1 H), 4.11 - 4.31 (m, 1 H), 3.78 - 3.99 (m, 2 H), 3.46 - 3.59 (m, 1 H), 2.63 - 2.81 (m, 1 H), 1.31 - 1.39 (m, 3 H), 1.14 - 1.30 (m, 4 H), 1.07 (br t, *J*=6.7 Hz, 9 H), 0.91 - 0.99 (m, 9 H)
125-9	596.6	^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.45 (s, 1 H), 8.36 (d, *J*=5.0 Hz, 1 H), 7.29 - 7.35 (m, 1 H), 7.25 (br d, *J*=9.5 Hz, 2 H), 7.07 (d, *J*=7.5 Hz, 1 H), 6.78 - 6.87 (m, 2 H), 6.20

		(dd, *J*=16.5, 2.4 Hz, 1 H), 5.72 - 5.81 (m, 1 H), 4.72 - 4.95 (m, 2 H), 4.45 - 4.54 (m, 1 H), 4.12 - 4.24 (m, 1 H), 3.80 - 3.94 (m, 2 H), 2.69 - 2.80 (m, 1 H), 1.98 (s, 3 H), 1.38 - 1.46 (m, 1 H), 1.34 (br t, *J*=7.2 Hz, 3 H), 1.26 (br d, *J*=6.6 Hz, 2 H), 1.18 (br d, *J*=6.8 Hz, 2 H), 1.09 (d, *J*=6.6 Hz, 3 H), 0.94 (d, *J*=6.6 Hz, 3 H), 0.74 - 0.84 (m, 1 H), 0.63 - 0.73 (m, 2 H)
125-10	585.2	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.45 (d, *J*=5.2 Hz, 2 H), 7.19 - 7.35 (m, 4 H), 6.99 (d, *J*=7.5 Hz, 1 H), 6.80 - 6.94 (m, 1 H), 6.16 - 6.27 (m, 1 H), 5.76 - 5.81 (m, 1 H), 4.97 (br s, 1 H), 4.26 - 4.46 (m, 2 H), 4.00 - 4.22 (m, 1 H), 3.62 - 3.87 (m, 2 H), 3.45 - 3.58 (m, 1 H), 3.07 - 3.21 (m, 1 H), 2.62 - 2.72 (m, 1 H), 1.95 (s, 3 H), 1.36 (d, *J*=6.6 Hz, 3 H), 1.08 (d, *J*=6.6 Hz, 6 H), 0.93 (d, *J*=6.6 Hz, 6 H)
125-11	598.6	^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.54 (s, 1 H), 8.45 (d, *J*=5.2 Hz, 1 H), 7.29 - 7.35 (m, 1 H), 7.22 - 7.25 (m, 1 H), 7.11 - 7.22 (m, 2 H), 7.01 (d, *J*=6.8 Hz, 1 H), 6.82 (dd, *J*=16.6, 10.6 Hz, 1 H), 6.21 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.72 - 5.79 (m, 1 H), 4.53 - 4.67 (m, 2 H), 4.35 (br d, *J*=13.3 Hz, 2 H), 3.66 (br dd, *J*=13.9, 2.9 Hz, 2 H), 2.63 - 2.72 (m, 1 H), 1.95 (s, 3 H), 1.41 (br d, *J*=5.2 Hz, 6 H), 1.07 (d, *J*=6.6 Hz, 7 H), 0.93 (d, *J*=6.6 Hz, 6 H)
129-1	632.2	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 8.73 - 9.11 (m, 1 H), 8.43 - 8.51 (m, 1 H), 8.37 (d, *J*=5.0 Hz, 1 H), 7.36 - 7.81 (m, 2 H), 7.14 - 7.22 (m, 1 H), 6.94 - 7.10 (m, 1 H), 6.73 - 6.92 (m, 1 H), 6.14 - 6.26 (m, 1 H), 5.76 (dt, *J*=10.2, 3.0 Hz, 1 H), 4.44 - 4.96 (m, 2 H), 3.76 - 4.25 (m, 4 H), 2.56 - 2.63 (m, 1 H), 2.22 (s, 3 H), 1.89 - 2.04 (m, 3 H), 1.34 (br t, *J*=7.2 Hz, 3 H), 1.15 - 1.27 (m, 3 H), 1.01 - 1.13 (m, 3 H), 0.79 - 0.99 (m, 3 H)
129-2	686.2	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 11.15 (br s, 1 H), 8.57 (s, 1 H), 8.38 (d, *J*=4.8 Hz, 1 H), 7.50 - 7.63 (m, 1 H), 7.34 (br t, *J*=8.8 Hz, 1 H), 7.14 - 7.23 (m, 2 H), 6.75 - 6.92 (m, 1 H), 6.19 (br d, *J*=16.6 Hz, 1 H), 5.75 (br d, *J*=10.4 Hz, 1 H), 4.44 - 5.04 (m, 2 H), 3.46 - 4.29 (m, 4 H), 2.39 - 2.47 (m, 1 H), 1.73 - 2.00 (m, 3 H), 1.29 (br t, *J*=7.3 Hz, 3 H), 0.95 - 1.12 (m, 6 H), 0.83 (d, *J*=6.6 Hz, 3 H)
129-3	628.2	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 8.55 (br s, 1 H), 8.35 (br s, 1 H), 7.43 - 7.57 (m, 2 H), 7.37 (br s, 2 H), 7.20 (br d, *J*=3.3 Hz, 1 H), 6.68 - 6.95 (m, 1 H), 6.19 (br d, *J*=16.6 Hz, 1 H), 5.67 - 5.86 (m, 1 H), 4.36 - 5.03 (m, 2 H), 3.49 - 4.31 (m, 4 H), 2.66 - 2.84 (m, 1 H), 2.61 (br s, 3 H), 2.41 (br s, 3 H), 1.90 (br s, 3 H), 1.32 (br d, *J*=5.4 Hz, 3 H), 1.08 - 1.23 (m, 3 H), 1.02 (br d, *J*=11.8 Hz, 3 H), 0.83 (br s, 3 H)
129-4	646.2	^{1}H NMR (400 MHz, DMSO-*d$_6$*) δ 9.11 (br d, *J*=2.5 Hz, 1 H), 8.42 - 8.54 (m, 1 H), 8.37 (d, *J*=5.0 Hz, 1 H), 7.31 - 7.81 (m, 2 H), 7.17 (d, *J*=4.8 Hz, 1 H), 6.94 - 7.11 (m, 1 H), 6.72 - 6.93 (m, 1 H), 6.19 (br d, *J*=16.8 Hz, 1 H), 5.67 - 5.83 (m, 1 H), 4.45 - 5.05 (m, 2 H), 3.44 - 4.26 (m, 4 H),

		2.54 - 2.94 (m, 1 H), 2.01 - 2.24 (m, 2 H), 1.81 - 2.00 (m, 3 H), 1.25 - 1.37 (m, 3 H), 1.11 - 1.24 (m, 3 H), 0.99 - 1.11 (m, 3 H), 0.80 - 0.99 (m, 6 H)
129-5	668.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.52 (s, 1 H), 8.40 (d, *J*=4.8 Hz, 1 H), 7.48 - 7.60 (m, 2 H), 7.42 (br t, *J*=7.5 Hz, 1 H), 7.35 (br d, *J*=7.9 Hz, 1 H), 7.26 (br d, *J*=7.7 Hz, 1 H), 7.18 (br d, *J*=5.0 Hz, 1 H), 6.75 - 6.92 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.75 (br d, *J*=10.6 Hz, 1 H), 4.44 - 5.02 (m, 2 H), 3.75 - 4.29 (m, 4 H), 2.55 - 2.61 (m, 1 H), 1.96 (s, 3 H), 1.23 - 1.34 (m, 3 H), 1.00 - 1.15 (m, 6 H), 0.96 (br d, *J*=6.4 Hz, 3 H)
129-6	640.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 9.49 (br s, 1 H), 8.44 (s, 1 H), 8.39 (d, *J*=5.0 Hz, 1 H), 7.58 (br d, *J*=5.2 Hz, 1 H), 7.30 - 7.43 (m, 1 H), 7.08 - 7.24 (m, 3 H), 6.85 (td, *J*=16.5, 10.6 Hz, 1 H), 6.20 (dd, *J*=16.7, 2.0 Hz, 1 H), 5.68 - 5.83 (m, 1 H), 4.45 - 4.95 (m, 2 H), 3.79 - 4.23 (m, 4 H), 2.67 (br s, 1 H), 1.98 (s, 3 H), 1.64 (br d, *J*=2.1 Hz, 1 H), 1.27 - 1.37 (m, 3 H), 1.12 - 1.26 (m, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.92 (d, *J*=6.6 Hz, 3 H), 0.60 - 0.76 (m, 2 H), 0.54 (br s, 2 H)
129-7	630.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.84 (s, 1 H), 8.42 (s, 1 H), 8.38 (d, *J*=5.0 Hz, 1 H), 7.61 (br d, *J*=8.1 Hz, 1 H), 7.34 - 7.43 (m, 1 H), 7.08 - 7.30 (m, 3 H), 6.72 - 6.99 (m, 1 H), 6.19 (br dd, *J*=16.7, 2.0 Hz, 1 H), 5.69 - 5.82 (m, 1 H), 4.42 - 4.92 (m, 2 H), 3.72 - 4.27 (m, 4 H), 3.55 (s, 3 H), 2.70 (br dd, *J*=13.8, 7.2 Hz, 1 H), 1.96 (s, 3 H), 1.32 (br t, *J*=6.7 Hz, 3 H), 1.16 - 1.26 (m, 3 H), 1.06 (br d, *J*=6.6 Hz, 3 H), 0.80 - 0.97 (m, 3 H)
129-8	648.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.78 - 8.87 (m, 1 H), 8.42 - 8.47 (m, 1 H), 8.36 (d, *J*=4.8 Hz, 1 H), 7.35 - 7.48 (m, 2 H), 7.16 (d, *J*=4.8 Hz, 1 H), 6.75 - 7.07 (m, 2 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.69 - 5.83 (m, 1 H), 4.44 - 4.92 (m, 2 H), 3.76 - 4.27 (m, 4 H), 3.54 (s, 3 H), 2.59 - 2.70 (m, 1 H), 1.96 (s, 3 H), 1.30 - 1.37 (m, 3 H), 1.16 - 1.29 (m, 3 H), 1.01 - 1.12 (m, 3 H), 0.80 - 0.97 (m, 3 H)
129-9	658.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 9.46 (br d, *J*=6.0 Hz, 1 H), 8.48 (s, 1 H), 8.37 (d, *J*=4.8 Hz, 1 H), 7.42 (br d, *J*=1.5 Hz, 2 H), 7.17 (d, *J*=4.8 Hz, 1 H), 6.98 - 7.11 (m, 1 H), 6.70 - 6.92 (m, 1 H), 6.20 (br d, *J*=16.8 Hz, 1 H), 5.69 - 5.83 (m, 1 H), 4.40 - 5.03 (m, 2 H), 3.80 - 4.23 (m, 4 H), 2.55 (br d, *J*=6.8 Hz, 1 H), 2.01 (s, 3 H), 1.51 - 1.61 (m, 1 H), 1.28 - 1.38 (m, 3 H), 1.14 - 1.23 (m, 3 H), 1.03 (br d, *J*=6.6 Hz, 3 H), 0.79 (br d, *J*=6.6 Hz, 3 H), 0.58 - 0.74 (m, 3 H), 0.45 (br d, *J*=4.1 Hz, 1 H)
129-10	668.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 10.29 (br s, 1 H), 8.52 (s, 1 H), 8.31 - 8.41 (m, 1 H), 7.48 - 7.60 (m, 1 H), 7.21 - 7.33 (m, 2 H), 7.11 - 7.19 (m, 1 H), 6.84 (td, *J*=16.2, 10.5 Hz, 1 H), 5.94 - 6.43 (m, 2 H), 5.75 (br d, *J*=10.4 Hz, 1 H), 4.42 - 5.03 (m, 2 H), 3.41 - 4.31 (m, 4 H), 2.52 - 2.58 (m,

		1 H), 1.96 (s, 3 H), 1.27 - 1.41 (m, 3 H), 0.99 - 1.26 (m, 6 H), 0.80 - 0.96 (m, 3 H)
129-11	650.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 10.24 - 10.54 (m, 1 H), 8.39 (d, *J*=4.8 Hz, 2 H), 7.40 - 7.58 (m, 2 H), 7.28 - 7.38 (m, 1 H), 7.17 (br d, *J*=4.6 Hz, 2 H), 6.73 - 6.94 (m, 1 H), 5.84 - 6.43 (m, 2 H), 5.75 (br dd, *J*=10.4, 2.3 Hz, 1 H), 4.39 - 5.11 (m, 2 H), 3.66 - 4.26 (m, 4 H), 2.56 - 2.73 (m, 1 H), 1.96 (s, 3 H), 1.26 - 1.42 (m, 3 H), 1.07 (br d, *J*=6.6 Hz, 6 H), 0.95 (br d, *J*=6.2 Hz, 3 H)
129-12	631.6	^{1}H NMR (400 MHz, DMSO-d_6) δ 9.50 (br s, 1 H), 8.42 - 8.50 (m, 1 H), 8.39 (d, *J*=4.8 Hz, 1 H), 7.26 - 7.41 (m, 2 H), 7.18 (br d, *J*=4.8 Hz, 1 H), 6.95 (br d, *J*=0.8 Hz, 1 H), 6.84 (td, *J*=16.2, 10.6 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.75 (br dd, *J*=10.4, 2.3 Hz, 1 H), 4.46 - 5.01 (m, 2 H), 3.41 - 4.23 (m, 4 H), 2.67 (br s, 1 H), 1.94 (br s, 3 H), 1.80 (br s, 3 H), 1.32 (br d, *J*=6.6 Hz, 3 H), 1.10 - 1.25 (m, 3 H), 1.07 (br d, *J*=6.6 Hz, 3 H), 0.96 (br d, *J*=6.0 Hz, 3 H)
129-13	612.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.42 (d, *J*=5.0 Hz, 1 H), 8.33 (dd, *J*=9.3, 2.5 Hz, 1 H), 7.25 - 7.34 (m, 1 H), 7.23 (d, *J*=5.0 Hz, 1 H), 6.76 - 6.92 (m, 1 H), 6.57 (d, *J*=8.5 Hz, 1 H), 6.52 (t, *J*=8.9 Hz, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.72 - 5.81 (m, 1 H), 5.65 (br s, 1 H), 4.43 - 4.91 (m, 2 H), 4.16 (br t, *J*=12.8 Hz, 1 H), 3.79 - 3.93 (m, 2 H), 3.71 (dd, *J*=6.0, 2.1 Hz, 2 H), 3.51 (br dd, *J*=13.6, 3.4 Hz, 1 H), 3.04 (t, *J*=2.2 Hz, 1 H), 2.63 - 2.76 (m, 1 H), 1.96 (s, 3 H), 1.32 (t, *J*=6.8 Hz, 3 H), 1.18 - 1.29 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.93 (d, *J*=6.4 Hz, 3 H)
132-1	623.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 9.96 (s, 1 H), 8.49 (d, *J*=3.1 Hz, 1 H), 8.30 (d, *J*=4.8 Hz, 1 H), 7.73 (d, *J*=8.3 Hz, 1 H), 7.39 (ddd, *J*=8.1, 6.2, 1.9 Hz, 1 H), 7.15 - 7.25 (m, 3 H), 7.12 (d, *J*=4.8 Hz, 1 H), 6.78 - 6.95 (m, 2 H), 6.20 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.71 - 5.82 (m, 1 H), 4.89 (br s, 1 H), 4.46 - 4.84 (m, 1 H), 3.47 - 4.30 (m, 4 H), 2.70 - 2.80 (m, 1 H), 1.94 (s, 3 H), 1.32 - 1.41 (m, 3 H), 1.18 - 1.32 (m, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 0.92 (d, *J*=6.6 Hz, 3 H)
132-2	607.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 10.01 (br s, 1 H), 8.28 - 8.42 (m, 2 H), 7.74 (br d, *J*=7.9 Hz, 1 H), 7.34 - 7.51 (m, 2 H), 7.25 (br s, 1 H), 7.11 - 7.21 (m, 2 H), 7.02 (br s, 1 H), 6.74 - 6.95 (m, 1 H), 6.20 (br d, *J*=16.6 Hz, 1 H), 5.67 - 5.86 (m, 1 H), 4.85 - 4.97 (m, 1 H), 4.42 - 4.84 (m, 1 H), 3.49 - 4.28 (m, 4 H), 2.71 - 2.86 (m, 1 H), 1.96 (br s, 3 H), 1.35 (br s, 3 H), 1.19 - 1.31 (m, 3 H), 1.07 (br d, *J*=6.0 Hz, 3 H), 0.94 (br d, *J*=5.6 Hz, 3 H)
141-1	632.2	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.49-8.56 (m, 1H), 8.09-8.17 (m, 1H), 7.35-7.47 (m, 2H), 7.30-7.35 (m, 1H), 7.14-7.22 (m, 1H), 6.96-7.04 (m, 1H), 6.52-6.72 (m, 1H), 6.34-6.47 (m, 1H), 5.75-5.85 (m, 1H), 5.03-5.19 (m, 1H), 4.33-4.55 (m, 1H), 3.88-4.12 (m, 2H), 3.67-3.82 (m, 1H), 3.35-3.47 (m, 1H), 3.00-3.12 (m, 1H), 2.67-2.82 (m, 1H), 2.52-2.64 (m, 1H), 2.30-2.41 (m, 2H), 2.21-2.29 (m,

		2H), 1.46 (br d, *J*=6.43 Hz, 4H), 1.28 (br s, 6H), 1.23 (br d, *J*=6.63 Hz, 3H), 1.03 (br d, *J*=6.63 Hz, 6H), 0.83 (br d, *J*=3.73 Hz, 6H)
141-2	684.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.50-8.59 (m, 1H), 8.13 (br s, 1H), 7.47-7.58 (m, 1H), 7.37 (d, *J*=7.46 Hz, 1H), 7.33 (s, 1H), 7.14-7.21 (m, 1H), 6.95-7.05 (m, 1H), 6.51-6.72 (m, 1H), 6.35-6.47 (m, 1H), 5.76-5.86 (m, 1H), 4.95-5.21 (m, 1H), 4.32-4.54 (m, 1H), 3.90-4.12 (m, 2H), 3.69-3.86 (m, 1H), 3.38-3.55 (m, 1H), 3.01-3.15 (m, 1H), 2.78-2.92 (m, 1H), 2.67-2.75 (m, 1H), 2.57 (br d, *J*=1.45 Hz, 1H), 2.20-2.38 (m, 2H), 1.55-1.65 (m, 3H), 1.47 (br d, *J*=6.63 Hz, 7H), 1.31-1.37 (m, 2H), 1.17-1.29 (m, 6H), 1.00-1.07 (m, 6H), 0.86-0.92 (m, 4H)
141-3	668.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.51-8.57 (m, 1H), 8.12 (br s, 1H), 7.35-7.41 (m, 1H), 7.34 (s, 2H), 7.14-7.22 (m, 1H), 6.96-7.02 (m, 1H), 6.53-6.72 (m, 1H), 6.35-6.47 (m, 1H), 5.76-5.87 (m, 1H), 5.00-5.19 (m, 1H), 4.32-4.57 (m, 1H), 3.85-4.12 (m, 2H), 3.68-3.78 (m, 1H), 3.23-3.42 (m, 2H), 2.68-2.85 (m, 1H), 2.51-2.63 (m, 1H), 2.31 (br d, *J*=5.18 Hz, 3H), 1.61 (br d, *J*=5.60 Hz, 3H), 1.46 (br d, *J*=6.43 Hz, 3H), 1.31 (br s, 3H), 1.28 (br s, 6H), 1.23 (br d, *J*=6.63 Hz, 3H), 1.04 (br d, *J*=6.84 Hz, 6H)
144-1	601.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.59 - 8.65 (m, 1 H), 8.36 - 8.44 (m, 1 H), 7.19 (d, *J*=5.0 Hz, 1 H), 7.06 (q, *J*=7.9 Hz, 1 H), 6.77 - 6.98 (m, 1 H), 6.41 - 6.49 (m, 1 H), 6.32 (br t, *J*=8.3 Hz, 1 H), 6.23 (dd, *J*=16.6, 1.7 Hz, 1 H), 5.81 (br d, *J*=11.8 Hz, 1 H), 5.10 - 5.18 (m, 1 H), 5.05 (br s, 1 H), 4.83 - 5.01 (m, 1 H), 4.02 - 4.48 (m, 3 H), 3.45 - 4.00 (m, 3 H), 3.13 - 3.21 (m, 1 H), 2.98 - 3.13 (m, 1 H), 2.63 - 2.93 (m, 1 H), 1.81 - 2.02 (m, 3 H), 1.03 - 1.12 (m, 3 H), 0.86 - 1.01 (m, 3 H)
144-2	526.1	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.66 (s, 1 H), 8.52 (d, *J*=5.0 Hz, 1 H), 7.34 (br s, 1 H), 6.73 - 7.00 (m, 1 H), 6.22 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.73 - 5.89 (m, 1 H), 4.81 - 5.02 (m, 1 H), 4.26 - 4.41 (m, 2 H), 3.79 - 4.09 (m, 3 H), 3.57 (d, *J*=11.0 Hz, 2 H), 2.97 - 3.07 (m, 1 H), 2.59 - 2.82 (m, 1 H), 1.91 - 2.04 (m, 3 H), 1.08 (dd, *J*=6.6, 3.5 Hz, 3 H), 1.02 (dd, *J*=6.6, 2.7 Hz, 3 H)
144-2-1	526.0	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.65 (s, 1 H), 8.48 (d, *J*=4.8 Hz, 1 H), 7.26 (d, *J*=5.0 Hz, 1 H), 6.87 (br s, 1 H), 6.22 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.80 (br d, *J*=12.0 Hz, 1 H), 4.90 (br s, 1 H), 4.23 - 4.39 (m, 2 H), 3.46 - 4.10 (m, 4 H), 3.28 (br s, 1 H), 2.98 - 3.09 (m, 1 H), 2.65 - 2.77 (m, 1 H), 1.91 (s, 3 H), 1.06 (d, *J*=6.6 Hz, 3 H), 1.01 (d, *J*=6.6 Hz, 3 H)
144-2-2	526.0	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.65 (s, 1 H), 8.49 (d, *J*=4.8 Hz, 1 H), 7.26 (d, *J*=4.8 Hz, 1 H), 6.73 - 6.98 (m, 1 H), 6.22 (dd, *J*=16.7, 2.0 Hz, 1 H), 5.80 (br d, *J*=10.8 Hz, 1 H), 4.82 - 5.00 (m, 1 H), 4.20 - 4.40 (m, 2 H), 3.71 - 4.16 (m, 2 H), 3.36 - 3.70 (m, 2 H), 3.20 - 3.29 (m, 1 H),

		2.97 - 3.11 (m, 1 H), 2.54 - 2.60 (m, 1 H), 1.98 (s, 3 H), 1.05 (d, *J*=6.6 Hz, 3 H), 1.00 (d, *J*=6.6 Hz, 3 H)
144-3	602.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 10.03 - 10.16 (m, 1 H), 8.59 (s, 1 H), 8.38 (d, *J*=4.6 Hz, 1 H), 7.43 - 7.61 (m, 1 H), 7.20 - 7.27 (m, 1 H), 7.17 (br d, *J*=4.6 Hz, 1 H), 6.77 - 6.99 (m, 1 H), 6.59 - 6.74 (m, 2 H), 6.22 (dd, *J*=16.7, 1.6 Hz, 1 H), 5.81 (br d, *J*=11.8 Hz, 1 H), 4.82 - 5.01 (m, 1 H), 4.17 - 4.57 (m, 3 H), 3.87 - 4.12 (m, 1 H), 3.46 - 3.82 (m, 2 H), 2.99 - 3.14 (m, 1 H), 2.59 - 2.86 (m, 1 H), 1.85 - 1.96 (m, 3 H), 1.06 (dd, *J*=6.4, 3.5 Hz, 3 H), 0.92 (br d, *J*=6.4 Hz, 3 H)
145-1	523.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (s, 1 H), 7.49 - 7.70 (m, 2 H), 7.33 - 7.46 (m, 2 H), 6.73 - 7.01 (m, 1 H), 6.20 (dd, *J*=16.6, 2.1 Hz, 1 H), 5.79 (br d, *J*=11.8 Hz, 1 H), 4.75 - 5.01 (m, 1 H), 4.03 - 4.37 (m, 5 H), 3.42 - 4.01 (m, 3 H), 2.90 - 3.20 (m, 2 H), 0.88 (s, 9 H)
145-1-1	523.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (s, 1 H), 7.50 - 7.68 (m, 2 H), 7.32 - 7.45 (m, 2 H), 6.85 (br d, *J*=11.0 Hz, 1 H), 6.20 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.79 (br d, *J*=11.0 Hz, 1 H), 4.75 - 5.00 (m, 1 H), 4.09 - 4.33 (m, 5 H), 3.37 - 3.99 (m, 3 H), 2.91 - 3.13 (m, 2 H), 0.88 (s, 9 H)
145-1-2	523.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (s, 1 H), 7.52 - 7.68 (m, 2 H), 7.35 - 7.46 (m, 2 H), 6.74 - 6.95 (m, 1 H), 6.20 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.74 - 5.86 (m, 1 H), 4.76 - 5.02 (m, 1 H), 3.99 - 4.41 (m, 5 H), 3.36 - 3.91 (m, 3 H), 2.88 - 3.12 (m, 2 H), 0.88 (s, 9 H)
145-2	586.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.61 - 8.70 (m, 1 H), 8.50 (br d, *J*=4.8 Hz, 1 H), 7.48 - 7.58 (m, 1 H), 7.42 (br s, 1 H), 7.17 - 7.36 (m, 3 H), 6.76 - 7.01 (m, 1 H), 6.23 (dd, *J*=16.5, 2.0 Hz, 1 H), 5.81 (br d, *J*=12.0 Hz, 1 H), 4.84 - 5.03 (m, 1 H), 4.26 - 4.50 (m, 3 H), 3.71 - 4.17 (m, 3 H), 3.57 (d, *J*=11.2 Hz, 3 H), 2.02 (br d, *J*=19.5 Hz, 3 H), 1.12 (br d, *J*=6.6 Hz, 3 H), 0.95 - 1.02 (m, 3 H)
145-2-1	586.1	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.63 (s, 1 H), 8.40 (d, *J*=5.0 Hz, 1 H), 7.45 - 7.59 (m, 1 H), 7.14 - 7.35 (m, 4 H), 6.73 - 7.01 (m, 1 H), 6.23 (dd, *J*=16.6, 2.1 Hz, 1 H), 5.71 - 5.87 (m, 1 H), 4.94 (br d, *J*=2.7 Hz, 1 H), 4.28 - 4.48 (m, 2 H), 3.47 - 4.15 (m, 4 H), 3.07 (br d, *J*=8.1 Hz, 2 H), 2.70 - 2.82 (m, 1 H), 1.91 (s, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 0.93 (d, *J*=6.8 Hz, 3 H)
145-2-2	586.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.63 (s, 1 H), 8.39 (d, *J*=4.8 Hz, 1 H), 7.46 - 7.59 (m, 1 H), 7.14 - 7.36 (m, 4 H), 6.75 - 7.01 (m, 1 H), 6.23 (dd, *J*=16.6, 1.9 Hz, 1 H), 5.81 (br d, *J*=10.4 Hz, 1 H), 4.82 - 5.01 (m, 1 H), 4.28 - 4.47 (m, 2 H), 3.43 - 4.18 (m, 4 H), 3.00 - 3.17 (m, 2 H), 2.61 - 2.73 (m, 1 H), 1.96 (s, 3 H), 1.06 (br d, *J*=6.6 Hz, 3 H), 0.95 (br d, *J*=6.6 Hz, 3 H)
145-3	523.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.38 (s, 1 H), 7.51 - 7.67 (m, 2 H), 7.34 - 7.46 (m, 2 H), 6.70 - 6.91 (m, 1 H),

		6.18 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.77 (dd, *J*=10.5, 2.0 Hz, 1 H), 5.07 - 5.23 (m, 1 H), 4.32 - 4.48 (m, 1 H), 4.20 - 4.28 (m, 1 H), 4.13 (br d, *J*=12.6 Hz, 3 H), 3.56 - 3.90 (m, 2 H), 3.33 - 3.44 (m, 1 H), 3.05 - 3.20 (m, 1 H), 2.79 - 3.02 (m, 1 H), 0.88 (s, 9 H)
146-1	594.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (br d, *J*=8.7 Hz, 1 H), 7.22 - 7.37 (m, 5 H), 7.05 - 7.17 (m, 2 H), 6.76 - 6.93 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.72 - 5.80 (m, 1 H), 4.95 (br s, 1 H), 4.84 (br d, *J*=4.6 Hz, 1 H), 4.40 - 4.77 (m, 1 H), 3.56 - 4.18 (m, 4 H), 3.23 - 3.29 (m, 1 H), 3.03 (br dd, *J*=15.4, 3.2 Hz, 1 H), 2.83 - 2.93 (m, 1 H), 2.74 - 2.82 (m, 1 H), 2.64 - 2.73 (m, 1 H), 2.54 - 2.60 (m, 1 H), 1.27 (br dd, *J*=18.1, 6.5 Hz, 3 H), 1.05 - 1.21 (m, 6 H), 0.94 (d, *J*=6.8 Hz, 3 H)
146-1-1	594.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 7.91 - 8.05 (m, 1 H), 7.17 - 7.35 (m, 5 H), 7.03 - 7.14 (m, 2 H), 6.52 - 6.71 (m, 1 H), 6.33 - 6.47 (m, 1 H), 5.75 - 5.87 (m, 1 H), 5.06 - 5.38 (m, 1 H), 5.02 (br d, *J*=7.5 Hz, 1 H), 4.80 - 4.91 (m, 1 H), 4.31 - 4.50 (m, 1 H), 3.83 - 4.14 (m, 3 H), 3.60 - 3.73 (m, 1 H), 3.18 - 3.32 (m, 1 H), 3.05 - 3.16 (m, 1 H), 2.88 - 3.05 (m, 2 H), 2.76 (dt, *J*=13.6, 6.7 Hz, 1 H), 1.42 (br d, *J*=6.6 Hz, 2 H), 1.36 (br t, *J*=6.2 Hz, 2 H), 1.20 (d, *J*=6.8 Hz, 5 H), 1.00 (d, *J*=6.8 Hz, 3 H)
146-1-2	594.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 7.90 - 8.01 (m, 1 H), 7.53 (br d, *J*=6.2 Hz, 1 H), 7.13 - 7.38 (m, 6 H), 6.49 - 6.68 (m, 1 H), 6.29 - 6.47 (m, 1 H), 5.70 - 5.82 (m, 1 H), 5.33 (br s, 1 H), 5.02 - 5.22 (m, 2 H), 4.91 - 5.02 (m, 1 H), 4.25 - 4.50 (m, 1 H), 3.83 - 4.14 (m, 3 H), 3.57 - 3.70 (m, 1 H), 3.20 (br d, *J*=18.2 Hz, 1 H), 2.77 - 2.98 (m, 2 H), 2.59 - 2.75 (m, 1 H), 1.30 - 1.44 (m, 4 H), 1.12 - 1.26 (m, 5 H), 0.94 (d, *J*=6.8 Hz, 3 H)
146-2	594.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.21 (d, *J*=14.3 Hz, 1 H), 7.22 - 7.36 (m, 5 H), 7.10 (dd, *J*=6.2, 2.7 Hz, 1 H), 7.04 (d, *J*=7.5 Hz, 1 H), 6.74 - 6.92 (m, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.70 - 5.82 (m, 1 H), 4.86 (br d, *J*=3.9 Hz, 1 H), 4.77 - 4.80 (m, 1 H), 4.72 (br d, *J*=19.3 Hz, 1 H), 4.40 - 4.58 (m, 1 H), 4.14 - 4.25 (m, 1 H), 3.55 - 3.91 (m, 2 H), 3.26 (br dd, *J*=14.4, 3.4 Hz, 1 H), 3.01 (br dd, *J*=15.9, 3.2 Hz, 1 H), 2.85 - 2.95 (m, 1 H), 2.79 (dq, *J*=12.6, 6.5 Hz, 1 H), 2.62 - 2.71 (m, 1 H), 2.51 - 2.57 (m, 1 H), 1.42 (br t, *J*=5.9 Hz, 3 H), 1.34 (br dd, *J*=10.9, 6.7 Hz, 3 H), 1.08 (d, *J*=6.8 Hz, 3 H), 0.96 (br dd, *J*=6.7, 3.8 Hz, 3 H)
146-2-1	594.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.03 - 8.18 (m, 1 H), 7.23 - 7.34 (m, 4 H), 7.19 (d, *J*=7.5 Hz, 1 H), 7.09 (dd, *J*=7.2, 1.6 Hz, 1 H), 7.03 (br d, *J*=7.5 Hz, 1 H), 6.52 - 6.74 (m, 1 H), 6.31 - 6.46 (m, 1 H), 5.81 (br t, *J*=9.2 Hz, 1 H), 4.89 - 5.21 (m, 2 H), 4.69 - 4.87 (m, 2 H), 3.95 - 4.66 (m, 2 H), 3.56 - 3.87 (m, 2 H), 3.24 (br dd, *J*=15.3, 2.5 Hz, 1 H), 2.94 - 3.12 (m, 2 H), 2.76 - 2.92 (m, 2 H), 1.46 -

		1.57 (m, 6 H), 1.22 (br d, *J*=6.8 Hz, 3 H), 0.97 - 1.05 (m, 3 H)
146-2-2	594.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 7.98 - 8.13 (m, 1 H), 7.45 - 7.61 (m, 1 H), 7.13 - 7.39 (m, 6 H), 6.48 - 6.71 (m, 1 H), 6.30 - 6.45 (m, 1 H), 5.79 (br t, *J*=9.4 Hz, 1 H), 4.98 - 5.24 (m, 2 H), 4.70 - 4.96 (m, 2 H), 4.18 - 4.57 (m, 2 H), 3.50 - 3.79 (m, 3 H), 3.22 (br d, *J*=14.9 Hz, 1 H), 2.79 - 2.95 (m, 2 H), 2.56 - 2.76 (m, 1 H), 1.42 - 1.64 (m, 6 H), 1.23 (br d, *J*=6.8 Hz, 3 H), 0.99 (br t, *J*=6.2 Hz, 3 H)
146-3	552.2	^{1}H NMR (400 MHz, DMSO-d_6) δ 8.10 - 8.30 (m, 1 H), 7.00 - 7.58 (m, 8 H), 6.75 - 6.96 (m, 1 H), 6.03 - 6.29 (m, 2 H), 5.75 (br d, *J*=8.9 Hz, 2 H), 4.81 - 4.88 (m, 1 H), 4.40 - 4.78 (m, 2 H), 3.44 - 4.06 (m, 4 H), 2.71 - 3.13 (m, 2 H), 2.71 - 2.72 (m, 1 H), 1.31 - 1.48 (m, 3 H), 1.09 - 1.30 (m, 3 H)
146-3-1	552.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 7.90 - 7.99 (m, 1 H), 7.24 - 7.32 (m, 6 H), 7.17 (br d, *J*=7.5 Hz, 1 H), 7.01 - 7.09 (m, 1 H), 6.48 - 6.67 (m, 1 H), 6.28 - 6.42 (m, 1 H), 5.71 - 5.81 (m, 1 H), 4.78 - 5.38 (m, 4 H), 3.49 - 4.10 (m, 4 H), 3.24 (br d, *J*=13.3 Hz, 1 H), 3.03 - 3.16 (m, 1 H), 2.89 - 3.03 (m, 2 H), 1.30 - 1.39 (m, 4 H), 1.18 (br d, *J*=6.8 Hz, 2 H)
146-3-2	552.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.04 (d, *J*=19.3 Hz, 1 H), 7.26 - 7.32 (m, 4 H), 7.21 - 7.24 (m, 2 H), 7.18 (d, *J*=7.5 Hz, 1 H), 7.04 (dd, *J*=7.3, 1.0 Hz, 1 H), 6.49 - 6.70 (m, 1 H), 6.32 - 6.44 (m, 1 H), 5.78 (br t, *J*=8.9 Hz, 1 H), 4.61 - 5.16 (m, 4 H), 4.24 - 4.54 (m, 1 H), 3.72 (br d, *J*=1.7 Hz, 1 H), 3.50 - 3.69 (m, 1 H), 3.17 - 3.30 (m, 1 H), 3.05 - 3.15 (m, 1 H), 2.90 - 3.04 (m, 2 H), 1.60 (br d, *J*=6.6 Hz, 1 H), 1.46 - 1.55 (m, 6 H)
146-3-3	552.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 7.91 - 8.03 (m, 1 H), 7.48 (br d, *J*=7.3 Hz, 2 H), 7.24 - 7.38 (m, 4 H), 7.05 - 7.22 (m, 2 H), 6.47 - 6.67 (m, 1 H), 6.30 - 6.43 (m, 1 H), 5.70 - 5.82 (m, 1 H), 5.32 (br s, 1 H), 4.91 - 5.25 (m, 2 H), 4.26 - 4.47 (m, 1 H), 3.35 - 4.11 (m, 5 H), 3.19 (br d, *J*=16.8 Hz, 1 H), 2.91 - 3.05 (m, 1 H), 2.75 (br d, *J*=14.3 Hz, 1 H), 1.31 - 1.38 (m, 3 H), 1.13 - 1.26 (m, 3 H)
146-3-4	552.2	^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 7.98 - 8.13 (m, 1 H), 7.43 - 7.52 (m, 2 H), 7.26 - 7.38 (m, 4 H), 7.14 - 7.22 (m, 2 H), 6.48 - 6.69 (m, 1 H), 6.31 - 6.44 (m, 1 H), 5.78 (br t, *J*=9.1 Hz, 1 H), 5.07 - 5.29 (m, 2 H), 4.76 - 4.92 (m, 1 H), 4.44 - 4.76 (m, 1 H), 4.27 - 4.41 (m, 1 H), 3.71 (br s, 1 H), 3.50 - 3.68 (m, 2 H), 3.20 (br d, *J*=17.0 Hz, 1 H), 2.90 - 3.07 (m, 1 H), 2.73 (br dd, *J*=13.8, 2.0 Hz, 1 H), 1.60 (br d, *J*=6.6 Hz, 2 H), 1.46 - 1.55 (m, 5 H)
146-4	568.3	^{1}H NMR (400 MHz, CHLOROFORM-d) δ 7.89-8.04 (m, 1H), 7.25-7.41 (m, 4H), 7.17 (br s, 1H), 6.99-7.13 (m, 3H), 6.26-6.65 (m, 2H), 5.70-5.81 (m, 1H), 5.37-5.54 (m,

		1H), 4.73-5.11 (m, 3H), 4.27-4.46 (m, 2H), 3.59-4.24 (m, 4H), 2.92-3.24 (m, 2H), 1.32-1.46 (m, 6H)

Section 2—Individual Examples

Example 150

1-((2*R*,5*S*)-4-(6-Chloro-7-(2-fluorophenyl)-2-imino-1-neopentyl-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one:



[0273] Step 1: *tert*-Butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate: A mixture of *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 165**, 1.30 g, 2.70 mmol) and neopentylamine (1.27 mL, 10.80 mmol) was heated at 110 °C for 3 days. The reaction mixture was purified on silica gel chromatography using 0-5% MeOH in DCM to afford *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (1.10 g, 2.07 mmol, 77.0 % yield). *m/z* (ESI, +ve ion): 532.3 $(M+H)^+$.

[0274] Step 2: *tert*-Butyl (2*R*,5*S*)-4-((*E*)-(5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(cyanoimino)methyl)-2,5-dimethylpiperazine-1-carboxylate: To a solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (0.600 g, 1.128 mmol) and triethylamine (0.792 mL, 5.64 mmol) in THF (3 mL) at r.t. was added

cyanogen bromide solution (1.503 mL, 4.51 mmol) dropwise. The reaction mixture was stirred at r.t. for 30 min. The reaction was diluted with water and extracted with EtOAc (3x). The combined organics were dried (Na_2SO_4) and concentrated to afford crude *tert*-butyl (2*R*,5*S*)-4-((*E*)-(5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(cyanoimino)methyl)-2,5-dimethylpiperazine-1-carboxylate (0.170 g, 0.305 mmol, 27.1 % yield). The crude was used in the next step without further purification. *m/z* (ESI, +ve ion): 557.3 $(M+H)^+$.

[0275] Step 3: 1-((2*R*,5*S*)-4-(6-Chloro-7-(2-fluorophenyl)-2-imino-1-neopentyl-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one: TFA (0.803 mL, 10.77 mmol) was added to a solution of *tert*-butyl (2*R*,5*S*)-4-((*E*)-(5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(cyanoimino)methyl)-2,5-dimethylpiperazine-1-carboxylate (0.30 g, 0.538 mmol) in dichloromethane (5 mL). The reaction was stirred at rt for 1 hour and concentrated under reduced pressure. The residue was suspended in DCM (5 mL), cooled to 0 °C, and treated with TEA (0.078 mL, 0.538 mmol) followed by acryloyl chloride (0.044 mL, 0.538 mmol). The reaction was warmed to r.t. and stirred for 10 min. The mixture was quenched with sat'd aqueous $NaHCO_3$ and extracted with DCM (2X). The combined organic layers were concentrated and the residue purified on silica gel using 0-100% EtOAc/EtOH (3:1) in heptane to afford the interemediate *N*-((*E*)-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)(5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)methylene)cyanamide (0.160 g, 0.313 mmol, 58.1 % yield). To this intermediate was added MeOH (4 mL) and 5N HCl (1.077 mL, 5.38 mmol) and reaction stirred at 70 °C for 1 hour. The reaction was cool to r.t., pH adjusted to 10 with 2N NaOH solution and extracted with EtOAc (2X). The organic was dried and the crude product was purified with ISCO reverse phase hplc eluting with 10-70% CH3CN (0.1% TFA) in water (0.1% TFA) to afford 1-((2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-2-imino-1-neopentyl-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one (0.102 g, 0.200 mmol, 37.1 % yield). *m/z* (ESI, +ve ion): 511.3 $(M+H)^+$. ^1H NMR (400 MHz, DICHLOROMETHANE-d_2) δ 7.72 (s, 1H), 7.32-7.51 (m, 2H), 7.22 (t, J=7.12 Hz, 1H), 7.12 (t, J=9.13 Hz, 1H), 6.40-6.63 (m, 1H), 6.11-6.28 (m, 1H), 5.63 (br d, J=9.12 Hz, 1H), 5.25-5.35 (m, 1H), 4.80 (br s, 1H), 4.50 (br s, 1H), 4.32 (s, 2H), 4.12-4.24 (m, 1H), 3.50-3.78 (m, 3H), 1.11-1.27 (m, 6H), 0.84-0.94 (m, 9H). ^{19}F NMR (376 MHz, DICHLOROMETHANE-d_2) δ -112.65 (s, 1F).

Example 151

1-((2*R*,5*S*)-4-((*Z*)-6-Chloro-7-(2-fluorophenyl)-2-(methylimino)-1-neopentyl-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one 2,2,2-trifluoroacetate



3378631 3380038

[0276] To a solution of 1-((2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-2-imino-1-neopentyl-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one, **Example 150**, (0.050 g, 0.098 mmol) in THF (4 mL) was added sodium hydride (0.012 ml, 0.294 mmol). After evolution of hydrogen gas, iodomethane (0.056 ml, 0.391 mmol) was added and the reaction stirred at r.t. for 1 hour and at 50 °C for 4 hours. The reaction was cooled, diluted with water and extracted with EtOAc (3X). The organic was concentrated and the residue purified by a reverse phase HPLC eluting with 10-70% CH3CN (0.1% TFA) in water (0.1% TFA) to afford 1-((2*R*,5*S*)-4-((*Z*)-6-chloro-7-(2-fluorophenyl)-2-(methylimino)-1-neopentyl-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one 2,2,2-trifluoroacetate, as TFA salt. *m/z* (ESI, +ve ion): 525.4 $(M+H)^+$. ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 10.18-10.34 (m, 1H), 8.09 (s, 1H), 7.50-7.60 (m, 2H), 7.35 (t, J=7.17 Hz, 1H), 7.21-7.25 (m, 1H), 6.49-6.67 (m, 1H), 6.37-6.47 (m, 1H), 5.80-5.88 (m, 1H), 5.53-5.75 (m, 1H), 4.92-5.17 (m, 2H), 4.30-4.59 (m, 2H), 3.63-3.85 (m, 2H), 3.16 (br s, 3H), 1.50-1.71 (m, 3H), 1.12-1.40 (m, 3H), 0.93 (s, 9H). ^{19}F NMR (376 MHz, CHLOROFORM-*d*) δ -112.23 (s, 1F).

Example 152

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-isopropylphenyl)-7-phenyl-5,6,7,8-tetrahydropyrido[3,4-d]pyrimidin-2(1*H*)-one



[0277] Step 1: 3-((2-Isopropylphenyl)amino)isonicotinamide: A mixture of [2-(1-methylethyl)phenyl]-boronic acid (8.97 g, 54.7 mmol), [2-(1-methylethyl)phenyl]-boronic acid (8.97 g, 54.7 mmol), copper chloride (0.541 g, 5.47 mmol) and TEA (2.54 ml, 18.23 mmol) was purged with N_2 followed by the addition of MeOH (100 mL) and the resulting mixture was stirred at r.t. overnight. The reaction was quenched with a 9:1 sat. NH_4Cl/NH_4OH, and extracted with DCM. The combined organics were dried over Na2SO4, filtered, concentrated and purified by chromatography on silica gel using 0-5% MeOH in DCM to afford 3-((2-isopropylphenyl)amino)isonicotinamide (0.97 g, 3.80 mmol, 10.42 % yield) as a yellow solid. m/z (ESI, +ve ion): 256 $(M+H)^+$. ^1H NMR (400 MHz, *DMSO-d6*) δ 9.65 (s, 1H), 8.32 - 8.38 (m, 1H), 8.20 (s, 1H), 7.96 (d, *J*=5.2 Hz, 1H), 7.78 (br s, 1H), 7.60 (d, *J*=5.2 Hz, 1H), 7.39 (dd, *J*=7.6, 1.6 Hz, 1H), 7.28 - 7.32 (m, 1H), 7.15 - 7.25 (m, 2H), 3.04 - 3.19 (m, 1H), 1.19 (d, *J*=6.8 Hz, 6H).

[0278] Step 2: 1-(2-Isopropylphenyl)pyrido[3,4-d]pyrimidine-2,4(1*H*,3*H*)-dione: A mixture of 3-((2-isopropylphenyl)amino)isonicotinamide (1.51 g, 5.91 mmol), pyridine (1.435

ml, 17.74 mmol) and CDI (2.88 g, 17.74 mmol) in MeCN (100 mL) was heated at 85 °C overnight. The mixture was quenched with sat. $NaHCO_3$, and extracted with EtOAc. The combined organics were dried over Na_2SO_4, filtered, concentrated and purified by chromatography on silica gel using 0-50% EtOAc in heptane to afford 1-(2-isopropylphenyl)pyrido[3,4-d]pyrimidine-2,4(1*H*,3*H*)-dione (1.3 g, 4.62 mmol, 78 % yield) as a white solid. m/z (ESI, +ve ion): 281.8 $(M+H)^+$. ^1H NMR (400 MHz, *DMSO-d*$_6$) δ 12.03 (s, 1H), 8.48 (d, *J*=5.2 Hz, 1H), 7.92 (d, *J*=5.0 Hz, 1H), 7.70 (s, 1H), 7.54 - 7.66 (m, 2H), 7.42 - 7.47 (m, 1H), 7.32 - 7.41 (m, 1H), 2.71 - 2.85 (m, 1H), 1.13 (d, *J*=6.8 Hz, 3H), 1.05 (d, *J*=6.8 Hz, 3H).

[0279] Step 3: *tert*-Butyl (2*R*,5*S*)-4-(1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[3,4-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate: A suspension of 1-(2-isopropylphenyl)pyrido[3,4-d]pyrimidine-2,4(1*H*,3*H*)-dione (0.186 g, 0.661 mmol) in toluene (3 mL) was added 1,1'-dimethyltriethylamine (1.155 mL, 6.61 mmol) and phosphorous oxychloride (0.308 mL, 3.31 mmol) and the resulting mixture was heated at 80 °C. After 5 min, the mixture went into solution and the heating continued for 30 min. LCMS showed complete conversion to desired intermediate. The reaction mixture was cooled to 0 °C and 10 equiv DIEA was added followed by (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (0.213 mL, 0.992 mmol). This mixture was stirred with warming to r.t. over 1 h at which time LCMS showed conversion to desired product. The mixture was poured into cold satd. $NaHCO_3$ solution and stirred vigorously for 10 min. The mixture was extracted with EtOAc, the combined organics were dried over Na_2SO_4, filtered, concentrated and purified by chromatography on silica gel using 0-40% EtOAc in heptane to afford *tert*-butyl (2*R*,5*S*)-4-(1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[3,4-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.281 g, 0.588 mmol, 89 % yield) as a light yellow foam. m/z (ESI, +ve ion): 477.8 $(M+H)^+$. ^1H NMR (400 MHz, *DMSO-d*$_6$) δ 8.39 (dd, *J*=5.4, 2.9 Hz, 1H), 7.72 - 7.80 (m, 2H), 7.50 - 7.65 (m, 2H), 7.38 - 7.47 (m, 1H), 7.21 - 7.31 (m, 1H), 4.72 - 4.85 (m, 1H), 4.23 - 4.43 (m, 1H), 4.00 - 4.14 (m, 2H), 3.67 - 3.81 (m, 2H), 3.40 - 3.60 (m, 1H), 1.45 (s, 9H), 1.31 (dd, *J*=9.1, 6.6 Hz, 3H), 1.15 - 1.21 (m, 3H), 1.11 (dd, *J*=6.6, 4.8 Hz, 3H), 1.00 (dd, *J*=6.8, 4.8 Hz, 3H).

[0280] Step 4: 7-Benzyl-4-((2*S*,5*R*)-4-(*tert*-butoxycarbonyl)-2,5-dimethylpiperazin-1-yl)-1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[3,4-d]pyrimidin-7-ium: A mixture of *tert*-butyl (2*R*,5*S*)-4-(1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[3,4-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.36 g, 0.754 mmol) and (bromomethyl)benzene (0.193

mL, 1.131 mmol) in acetone (20 mL) was heated to reflux for 1 h. Some product was observed, mostly starting material. More (bromomethyl)benzene (0.193 mL, 1.131 mmol) was added and the resulting mixture was heated to reflux overnight. The mixture was brought to r.t., concentrated and purified by chromatography on a small amount of silica gel using 0-10% MeOH in DCM to afford 7-benzyl-4-((2*S*,5*R*)-4-(*tert*-butoxycarbonyl)-2,5-dimethylpiperazin-1-yl)-1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[3,4-d]pyrimidin-7-ium (0.401 g, 0.705 mmol, 94 % yield) as a yellow solid. m/z (ESI, +ve ion): 567.8 $(M+H)^+$. ^1H NMR (400 MHz, *DMSO-d*$_6$) δ 8.84 (br d, *J*=6.8 Hz, 1H), 8.30 - 8.40 (m, 2H), 7.56 - 7.65 (m, 2H), 7.34 - 7.48 (m, 6H), 7.25 (d, *J*=8.1 Hz, 1H), 5.87 - 5.90 (m, 2H), 4.66 - 4.76 (m, 1H), 4.24 - 4.46 (m, 1H), 4.01 - 4.15 (m, 1H), 3.66 - 3.87 (m, 2H), 3.41 - 3.61 (m, 1H), 2.58 - 2.66 (m, 1H), 1.45 (s, 9H), 1.34 (t, *J*=5.5 Hz, 3H), 1.18 (dd, *J*=14.5, 6.8 Hz, 3H), 1.08 (d, *J*=6.8 Hz, 3H), 0.95 (dd, *J*=10.8, 6.8 Hz, 3H).

[0281] Step 5: *tert*-Butyl (2*R*,5*S*)-4-(7-benzyl-1-(2-isopropylphenyl)-2-oxo-1,2,5,6,7,8-hexahydropyrido[3,4-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate: 7-benzyl-4-((2*S*,5*R*)-4-(*tert*-butoxycarbonyl)-2,5-dimethylpiperazin-1-yl)-1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[3,4-d]pyrimidin-7-ium was dissolved in 80% aq. MeOH (20 mL) and brought to 0 °C. Then, $NaBH_4$ (0.570 g, 15.08 mmol) was added and the resulting mixture was heated to reflux for 20 min. The reaction went to completion, brought to r.t., carefully quenched with sat. $NaHCO_3$ and extracted with DCM. The combined organics were dried over Na2SO4, filtered and concentrated to afford *tert*-butyl (2*R*,5*S*)-4-(7-benzyl-1-(2-isopropylphenyl)-2-oxo-1,2,5,6,7,8-hexahydropyrido[3,4-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. m/z (ESI, +ve ion): 571.8 $(M+H)^+$.

[0282] Step 6: *tert*-Butyl (2*R*,5*S*)-4-(1-(2-isopropylphenyl)-2-oxo-1,2,5,6,7,8-hexahydropyrido[3,4-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate: To a solution of *tert*-butyl (2*R*,5*S*)-4-(7-benzyl-1-(2-isopropylphenyl)-2-oxo-1,2,5,6,7,8-hexahydropyrido[3,4-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.280 g, 0.490 mmol) in MeOH (15 mL) was added palladium 10 wt. % on activated carbon (0.365 g, 0.343 mmol) and ammonium formate (0.309 g, 4.90 mmol) and the resulting mixture was heated to reflux. After 20 min the starting material was consumed and desired mass was observed. The mixture was brought to r.t., filtered through celite, concentrated and purified by chromatography on silica gel using 0-100% 3:1EtOAc/EtOH in heptane to afford *tert*-butyl (2*R*,5*S*)-4-(1-(2-isopropylphenyl)-2-oxo-1,2,5,6,7,8-hexahydropyrido[3,4-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.105 g, 0.218 mmol, 44.5 % yield) as a white solid.

m/z (ESI, +ve ion): 481.8 $(M+H)^+$. ^1H NMR (400 MHz, *DMSO-d_6*) δ 7.39 - 7.51 (m, 2H), 7.25 - 7.34 (m, 1H), 7.13 - 7.18 (m, 1H), 7.03 - 7.11 (m, 1H), 4.38 - 4.53 (m, 1H), 4.12 - 4.38 (m, 1H), 3.93 - 4.06 (m, 1H), 3.58 - 3.74 (m, 2H), 3.40 - 3.49 (m, 2H), 2.98 - 3.16 (m, 2H), 2.73 - 2.94 (m, 2H), 2.36 - 2.46 (m, 2H), 1.41 - 1.47 (m, 9H), 1.10 - 1.19 (m, 8H), 1.03 - 1.10 (m, 4H).

[0283] Steps 7 & 8: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-isopropylphenyl)-7-phenyl-5,6,7,8-tetrahydropyrido[3,4-d]pyrimidin-2(1*H*)-one, 3386982: A solution of *tert*-butyl (2*R*,5*S*)-4-(1-(2-isopropylphenyl)-2-oxo-1,2,5,6,7,8-hexahydropyrido[3,4-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.060 g, 0.125 mmol) in acetonitrile (2 mL) was added to a stirring mixture of 2-(trimethylsilyl)phenyl triflate (0.056 g, 0.187 mmol) and cesium fluoride (0.057 g, 0.374 mmol) in acetonitrile (2 mL). The reaction mixture was stirred at r.t. for 1 hour, concentrated, diluted with water and extracted with EtOAc. The organic was concentrated to give a crude *tert*-butyl (2*R*,5*S*)-4-(1-(2-isopropylphenyl)-2-oxo-7-phenyl-1,2,5,6,7,8-hexahydropyrido[3,4-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. *m/z* (ESI, +ve ion): 558.3 $(M+H)^+$. This crude intermediate was dissolved in DCM (2 mL) and treated with TFA (0.288 mL, 3.74 mmol). The resulting mixture was stirred at r.t. for 1 hour and then concentrated *in vacuo*. The residue was suspended in DCM (2 mL) and treated with TEA (0.087 mL, 0.623 mmol) followed by acryloyl chloride (0.020 mL, 0.249 mmol). The reaction was stirred at r.t. for 10 minutes, quenched with water, and extracted with DCM. The organic was concentrated and the residue purified with ISCO using 0-100% EtOAc/EtOH (3:1) in heptane to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-isopropylphenyl)-7-phenyl-5,6,7,8-tetrahydropyrido[3,4-d]pyrimidin-2(1*H*)-one (0.008 g, 7.82 μmol, 6.28 % yield) as a mixture of atropisomers. *m/z* (ESI, +ve ion): 512.4 $(M+H)^+$. ^1H NMR (400 MHz, CHLOROFORM-*d*) δ 7.44-7.50 (m, 2H), 7.28-7.36 (m, 1H), 7.22 (t, *J*=7.88 Hz, 2H), 7.10-7.16 (m, 1H), 6.86 (t, *J*=7.26 Hz, 1H), 6.68 (br d, *J*=7.67 Hz, 2H), 6.35 (br t, *J*=15.96 Hz, 1H), 5.75 (br t, *J*=10.26 Hz, 1H), 4.85-5.02 (m, 1H), 4.19-4.44 (m, 1H), 3.89-4.05 (m, 1H), 3.75-3.89 (m, 1H), 3.49-3.74 (m, 4H), 3.30-3.46 (m, 1H), 3.03-3.28 (m, 1H), 2.80-2.93 (m, 1H), 2.54-2.80 (m, 2H), 1.77 (td, *J*=6.63, 13.27 Hz, 1H), 1.17-1.31 (m, 12H).

Example 153

7-(2,4-Difluorophenyl)-1-(4,6-diisopropylpyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-6-fluoro-pyrido[2,3-*d*]pyrimidin-2-one



[0284] Step 1. 2,6-Dichloro-N-((4,6-diisopropylpyrimidin-5-yl)carbamoyl)-5-fluoronicotinamide. To a 250-mL round-bottomed flask was added 2,6-dichloro-5-

fluoronicotinamide (**Intermediate S**, 3.55 g, 16.7 mmol) and oxalyl chloride (12.5 mL, 25.1 mmol) in tetrahydrofuran (33.5 mL). The flask was fitted with a Findenser, and the mixture was stirred and heated at 80 °C for 1 h. The reaction mixture was concentrated in vacuo. The crude mixture was carried into the next step of the synthesis, without further purification to prevent decomposition.

[0285] To a 250-mL round-bottomed flask was added (2,6-dichloro-5-fluoronicotinoyl)carbamoyl isocyanate (crude material from previous step) in tetrahydrofuran (33.5 mL). Then a solution of 4,6-diisopropylpyrimidin-5-amine (**Intermediate U**, 3.1 g, 17.5 mmol) in THF (10 mL) was added dropwise into the reaction mixture. The mixture was allowed to stir under an inert (N2) atmosphere for 1.5 h. The reaction mixture was concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (220 grams), eluting with a gradient of 0-40% EtOAc/heptane, to afford 2,6-dichloro-*N*-((4,6-diisopropylpyrimidin-5-yl)carbamoyl)-5-fluoronicotinamide (4.2 g, 10.3 mmol, 61.7 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.41 (br s, 1 H) 9.66 (br s, 1 H) 8.99 (s, 1 H) 8.55 (br d, *J*=7.88 Hz, 1 H) 3.21 - 3.28 (m, 2 H) 1.17 (d, *J*=6.63 Hz, 12 H). *m/z* (ESI, +ve ion): 414.0 $(M+H)^+$.

[0286] Step 2. 7-Chloro-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a 250-mL round-bottomed flask was added 2,6-dichloro-*N*-((4,6-diisopropylpyrimidin-5-yl)carbamoyl)-5-fluoronicotinamide (4.0 g, 9.6 mmol) in tetrahydrofuran (48.3 mL). The reaction mixture was cooled to 0 °C with a wet ice/water bath. Then potassium bis(trimethylsilyl)amide, 1M solution in tetrahydrofuran (12.0 mL, 12.0 mmol) was added via an addition funnel dropwise to the reaction mixture over 5 min. The ice bath was removed and the reaction mixture was allowed to slowly warm to ambient temperature, while stirred under an inert (N_2) atmosphere for 1 h. More KHMDS (0.5 eq; 6 mL) was added dropwise to the reaction mixture. The reaction mixture was quenched with sat. aq. NH_4Cl (50 mL), then the mixture was diluted with 3:1 EtOAc/MeOH and brine solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. This afforded 7-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (2.5 g, 6.8 mmol, 70.9 % yield) as a tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.03 - 12.52 (m, 1 H) 8.97 - 9.23 (m, 1 H) 8.25 - 8.58 (m, 1 H) 2.80 (dt, *J*=13.22, 6.56 Hz, 2 H) 0.96 (d, *J*=6.63 Hz, 6 H) 0.85 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 378.0 $(M+H)^+$.

[0287] Step 3. 7-(2,4-Difluorophenyl)-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a 100-mL round-bottomed flask was added 7-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.2 g, 0.6 mmol) and potassium acetate (0.195 g, 1.985 mmol) in 1,4-dioxane (3.3 mL). The reaction mixture was deoxygenated by bubbling (N_2) gas into the mixture for 5 min. Then (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.048 g, 0.066 mmol) was added to the reaction mixture. The mixture was stirred and heated at 95 °C for 10 min. Then 2,4-difluorobenzeneboronic acid (0.136 g, 0.860 mmol), followed by water (0.05 mL) was added to the reaction mixture. The reaction mixture was allowed to stir at 95 °C for 16 h. The reaction mixture was diluted with sat. aq. NH_4Cl and EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-90% EtOAc/heptane, to afford 7-(2,4-difluorophenyl)-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.2 g, 0.4 mmol, 74.7 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.27 (s, 1 H) 9.13 (s, 1 H) 8.47 (d, *J*=8.50 Hz, 1 H) 7.37 - 7.45 (m, 1 H) 7.19 - 7.33 (m, 2 H) 2.97 (spt, *J*=6.57 Hz, 2 H) 1.10 (d, *J*=6.63 Hz, 6 H) 0.93 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 456.0 $(M+H)^+$.

[0288] Step 4. *tert*-Butyl (2*R*,5*S*)-4-(7-(2,4-difluorophenyl)-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added 7-(2,4-difluorophenyl)-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.2 g, 0.4 mmol) and n,n'-diisopropylethylamine (0.1 mL, 0.5 mmol) in acetonitrile (7.9 mL). Then phosphorous oxychloride (0.1 mL, 0.5 mmol) was added slowly to the reaction mixture. The flask was fitted with a Findenser, then the mixture was stirred and heated at 80 °C, while under an inert (N2) atmosphere for 45 min. The reaction mixture was removed from the heat bath and allowed to cool to ambient temperature.

[0289] The reaction mixture was cooled to 0 °C with a wet ice/water bath, while stirred. Then DIPEA (0.3 mL) was added slowly to the reaction mixture. Then a mixture of (2*R*,5*S*)-1-Boc-2,5-dimethylpiperazine (0.1 g, 0.5 mmol, Astatech, San Diego, CA, USA) in MeCN (3 mL) was added slowly to the reaction mixture. The ice bath was removed and the reaction mixture was allowed to slowly warm to ambient temperature over 1 h. The mixture was diluted

with EtOAc and sat. aq. NH_4Cl, then the layers were separated. The aqueous layer was extracted with EtOAc and brine. The combined organic extracts were dried over MgSO4, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through an Interchim (25 micron) silica-gel column (120 grams), eluting with a gradient of 0-100 % EtOAc/heptane, to afford *tert*-butyl (2R,5S)-4-(7-(2,4-difluorophenyl)-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.2 g, 0.3 mmol, 83 % yield) as tan solid. *m/z* (ESI, +ve ion): 652.1 $(M+H)^+$.

[0290] Step 5. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2,4-difluorophenyl)-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoropyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(7-(2,4-difluorophenyl)-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.2 g, 0.3 mmol) and trifluoroacetic acid (0.1 mL, 1.918 mmol) in dichloromethane (1.9 mL). The reaction mixture was stirred and heated at 38 °C for 2.5 h, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0291] The previous residue was diluted with dichloromethane (1.9 mL), then the mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.8 mL, 4.6 mmol) was added to the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (0.1 mL, 0.3 mmol) was added to the mixture dropwise. The mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-5% MeOH/DCM. This afforded 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2,4-difluorophenyl)-1-(4,6-diisopropylpyrimidin-5-yl)-6-fluoropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.1 g, 0.2 mmol, 60.3 % yield) as light-yellow solid. ^1H NMR (400 MHz, DMSO-d_6) δ 9.10 (s, 1 H) 8.38 (dd, *J*=9.64, 5.29 Hz, 1 H) 7.21 - 7.45 (m, 3 H) 6.77 - 6.91 (m, 1 H) 6.19 (br d, *J*=18.24 Hz, 1 H) 5.72 - 5.79 (m, 1 H) 4.72 - 4.98 (m, 2 H) 4.13 - 4.25 (m, 1 H) 3.85 (br s, 2 H) 2.66 - 2.77 (m, 2 H) 1.33 (t, *J*=6.01 Hz, 3 H) 1.25 (br d, *J*=6.63 Hz, 2 H) 1.18 (br d, *J*=6.63 Hz, 2 H) 1.09 (dd, *J*=6.63, 2.28 Hz, 6 H) 0.94 (d, *J*=6.43 Hz, 6 H). *m/z* (ESI, +ve ion): 606.4 $(M+H)^+$.

Example 154

6-Chloro-1-(4,6-diisopropyl-2-methoxy-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one



[0292] Step 1. 2,5,6-Trichloro-N-((4,6-diisopropyl-2-methoxypyrimidin-5-yl)carbamoyl)nicotinamide. To a 150-mL round-bottomed flask was added 2,5,6-trichloronicotinamide (**Intermediate P**, 0.7 g, 3.1 mmol) and oxalyl chloride (2.3 mL, 4.6 mmol) in tetrahydrofuran (6.2 mL). The flask was fitted with a Findenser, and the mixture was stirred and heated at 80 °C for 45 min. The reaction mixture was concentrated *in vacuo*. The crude mixture was carried into the next step of the synthesis, without further purification to prevent decomposition.

[0293] To a 100-mL round-bottomed flask was added (2,5,6-trichloronicotinoyl)carbamoyl isocyanate (crude material from previous step) in tetrahydrofuran (6.2 mL). The reaction

mixture was cooled to -10 °C with a dilute acetone/dry ice bath, then a solution of 4,6-diisopropyl-2-methoxypyrimidin-5-amine (**Intermediate I-36**, 0.6 g, 3.2 mmol) in THF (5 mL) was added to the reaction mixture. The mixture was allowed to stir under an inert (N2) atmosphere for 1 h. The reaction mixture was concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (220 grams), eluting with a gradient of 0-40% EtOAc/heptane, to provide 2,5,6-trichloro-*N*-((4,6-diisopropyl-2-methoxypyrimidin-5-yl)carbamoyl)nicotinamide as off-white solid. *m/z* (ESI, +ve ion): 460.1 $(M+H)^+$.

[0294] Step 2. 6,7-Dichloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a 250-mL round-bottomed flask was added 2,5,6-trichloro-*N*-((4,6-diisopropyl-2-methoxypyrimidin-5-yl)carbamoyl)nicotinamide (1.0 g, 2.2 mmol) in tetrahydrofuran (11.4 mL). Then potassium bis(trimethylsilyl)amide, 1M solution in tetrahydrofuran (2.8 mL, 2.8 mmol) was added via syringe dropwise to the reaction mixture. The reaction mixture was allowed to stir under an inert (N_2) atmosphere for 30 min. The reaction mixture was quenched with sat. aq. NH_4Cl (10 mL), then the mixture was diluted with 3:1 EtOAc/MeOH and brine solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through an Interchim (25 micron) silica-gel column (80 grams), eluting with a gradient of 0-40% EtOAc/heptane This afforded 6,7-dichloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.2 g, 0.5 mmol) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.17 - 12.42 (m, 1 H) 8.49 - 8.74 (m, 1 H) 3.98 (s, 3 H) 2.82 - 2.92 (m, 2 H) 1.08 (d, *J*=6.63 Hz, 6 H) 0.98 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 424.1 $(M+H)^+$.

[0295] Step 3. *tert*-Butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added 6,7-dichloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.2 g, 0.5 mmol) and n,n'-diisopropylethylamine (0.1 mL, 0.7 mmol) in acetonitrile (7.9 mL). Then phosphorous oxychloride (0.1 mL, 0.7 mmol) was added slowly to the reaction mixture. The flask was fitted with a Findenser, then the mixture was stirred and heated at 80 °C, while under an inert (N2) atmosphere for 45 min. The reaction mixture was removed from the heat bath and allowed to cool to ambient temperature. Then set aside for later use.

[0296] The previous reaction mixture was cooled to 0 °C with a wet ice/water bath, while stirred. Then DIPEA (1.5 mL) was added slowly to the reaction mixture. Then a mixture of (2*R*,5*S*)-1-Boc-2,5-dimethylpiperazine (0.1 g, 0.7 mmol, Astatech, San Diego, CA, USA) in MeCN (3 mL) was added slowly to the reaction mixture. The ice bath was removed and the overall mixture was allowed to slowly warm to ambient temperature over 1 h. The mixture was diluted with EtOAc and sat. aq. NH_4Cl, then the layers were separated. The aqueous layer was extracted with EtOAc and brine. The combined organic extracts were dried over MgSO4, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (120 grams), eluting with a gradient of 0-100 % EtOAc/heptane to afford *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.1 g, 0.2 mmol, 37.5 % yield) as tan solid. *m/z* (ESI, +ve ion): 620.1 $(M+H)^+$.

[0297] **Step 4. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one.** To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.1 g, 0.2 mmol) and trifluoroacetic acid (0.1 mL, 2.1 mmol) in 1,2-dichloroethane (4.0 mL). The reaction mixture was stirred and heated at 70 °C for 1 h, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification

[0298] The previous residue was diluted with DCM (4 mL), then the reaction mixture was cooled to 0 °C with a wet ice/water bath. Then N,N'-diisopropylethylamine (0.4 mL, 2.6 mmol) was added to the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (0.1 mL, 0.2 mmol) was added to the mixture dropwise. The mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (40 grams), eluting with a gradient of 0-5% MeOH/DCM. This afforded 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.1 g, 0.2 mmol, 92 % yield) as tan solid. *m/z* (ESI, +ve ion): 574.1 $(M+H)^+$.

[0299] Step 5. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.1 g, 0.191 mmol) and potassium acetate (0.1 g, 0.5 mmol) in 1,4-dioxane (3.3 mL). The reaction mixture was deoxygenated by bubbling (N_2) gas into the mixture for 5 min. Then (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.1 g, 0.1 mmol) was added to the reaction mixture. The mixture was stirred and heated at 95 °C for 10 min. Then 2-fluorophenylboronic acid (0.1 g, 0.2 mmol), followed by water (0.1 mL) was added to the reaction mixture. The overall reaction mixture was allowed to stir at 95 °C for 16 h. The reaction mixture was diluted with sat. aq. NH_4Cl and EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through an Interchim (15 micron) silica-gel column, eluting with a gradient of 0-4% MeOH/DCM, to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-methoxypyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.1 g, 0.1 mmol, 24.71 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.47 (d, *J*=3.94 Hz, 1 H) 7.50 - 7.56 (m, 1 H) 7.28 - 7.35 (m, 2 H) 7.19 - 7.24 (m, 1 H) 6.83 (td, *J*=17.00, 10.57 Hz, 1 H) 6.19 (dd, *J*=16.79, 1.87 Hz, 1 H) 5.73 - 5.78 (m, 1 H) 4.72 - 4.94 (m, 2 H) 4.13 - 4.24 (m, 1 H) 3.91 (s, 3 H) 3.83 - 3.88 (m, 1 H) 3.17 (d, *J*=5.18 Hz, 1 H) 2.58 - 2.71 (m, 2 H) 1.34 (t, *J*=5.80 Hz, 3 H) 1.25 (br d, *J*=6.63 Hz, 2 H) 1.18 (d, *J*=6.84 Hz, 2 H) 1.06 (dd, *J*=6.53, 3.84 Hz, 6 H) 0.92 (dd, *J*=6.53, 2.80 Hz, 6 H). *m/z* (ESI, +ve ion): 634.4 $(M+H)^+$.

Example 155

5-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-1-yl]-4,6-diisopropyl-pyrimidine-2-carbonitrile



[0300] Step 1. 2,5-Dichloro-N-((2-cyano-4,6-diisopropylpyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide. To a 100-mL round-bottomed flask was added 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (**Intermediate 99B**, 0.55 g, 1.81 mmol) and oxalyl chloride (1.36 mL, 2.72 mmol) in tetrahydrofuran (9.07 mL). The flask was fitted with a Findenser, and mixture was stirred and heated at 80 °C for 45 min. The reaction mixture was concentrated *in vacuo*. The crude mixture was carried into the next step of the synthesis, without further purification to prevent decomposition.

[0301] To a 150-mL round-bottomed flask was added (2,5-Dichloro-6-(2-fluorophenyl)nicotinoyl)carbamoyl isocyanate (crude material from previous step) in tetrahydrofuran (9.07 mL). The reaction mixture was cooled to -10 °C with a dilute acetone/dry ice bath, then a solution of 5-amino-4,6-diisopropylpyrimidine-2-carbonitrile (**Intermediate 219**, 0.38 g, 1.90 mmol) in THF (3 mL) was added to the reaction mixture. The mixture was

allowed to stir under an inert (N2) atmosphere for 1 h. The reaction mixture was concentrated *in vacuo*. The crude material was triturated from EtOAc and heptane, then the solids were collected by filtration. The solids were washed with heptane. This afforded 2,5-dichloro-*N*-((2-cyano-4,6-diisopropylpyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide (0.75 g, 1.45 mmol, 80 % yield) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.57 (br s, 1 H) 9.96 (br s, 1 H) 8.63 (s, 1 H) 7.52 - 7.64 (m, 2 H) 7.37 - 7.43 (m, 2 H) 1.20 (br d, *J*=6.63 Hz, 14 H). *m/z* (ESI, +ve ion): 515.0 $(M+H)^+$.

[0302] Step 2. 5-(6-Chloro-7-(2-fluorophenyl)-2,4-dioxo-3,4-dihydropyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidine-2-carbonitrile. To a 100-mL round-bottomed flask was added 2,5-dichloro-*N*-((2-cyano-4,6-diisopropylpyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide (0.72 g, 1.39 mmol) in tetrahydrofuran (6.9 mL). Then potassium bis(trimethylsilyl)amide, 1M solution in tetrahydrofuran (1.7 mL, 1.74 mmol) was added via an addition funnel dropwise to the reaction mixture over 5 min. The reaction mixture was allowed to stir under an inert (N_2) atmosphere for 1 h. More KHMDS (0.5 eq; 0.6 mL) was added dropwise to the reaction mixture, until SM was mostly consumed. The reaction mixture was quenched with sat. aq. NH_4Cl (50 mL), then the mixture was diluted with 3:1 EtOAc/MeOH and brine solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. This afforded 5-(6-chloro-7-(2-fluorophenyl)-2,4-dioxo-3,4-dihydropyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidine-2-carbonitrile (0.65 g, 1.35 mmol, 97 % yield) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.24 (br s, 1 H) 8.59 (s, 1 H) 7.49 - 7.54 (m, 1 H) 7.25 - 7.34 (m, 2 H) 7.16 - 7.25 (m, 1 H) 3.00 - 3.21 (m, 2 H) 1.09 - 1.15 (m, 6 H) 0.94 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 479.0 $(M+H)^+$.

[0303] Step 3. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-cyano-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added 5-(6-chloro-7-(2-fluorophenyl)-2,4-dioxo-3,4-dihydropyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidine-2-carbonitrile (0.60 g, 1.25 mmol) and N,N'-diisopropylethylamine (0.28 mL, 1.62 mmol) in acetonitrile (6.26 mL). Then phosphorous oxychloride (0.14 mL, 1.50 mmol) was added slowly to the reaction mixture. The flask was fitted with a Findenser, then the mixture was stirred and heated at 80 °C, while under an inert (N2) atmosphere for 45 min. Another aliquot of N,N'-diisopropylethylamine (0.28 mL, 1.62 mmol) and phosphorous oxychloride (0.14 mL, 1.50 mmol) was added to the reaction mixture, then allowed the mixture

to stir 10 min. The reaction mixture was removed from the heat bath and allowed to cool to ambient temperature.

[0304] The previous reaction mixture was cooled to 0 °C with a wet ice/water bath, while stirred. Then DIPEA (2.4 mL) was added slowly to the mixture. Then a mixture of (2*R*,5*S*)-1-Boc-2,5-dimethylpiperazine (0.33 g, 1.56 mmol, Astatech, San Diego, CA, USA) in MeCN (3 mL) was added slowly to the reaction mixture. The ice bath was removed and the overall mixture was allowed to slowly warm to ambient temperature over 1 h. Another aliquot of DIPEA (0.6 mL) and (2*R*,5*S*)-1-Boc-2,5-dimethylpiperazine (0.33 g, 1.56 mmol) in MeCN (2 mL) was added to the mixture, then allowed the mixture to stir an additional 10 min. The mixture was diluted with EtOAc and sat. aq. NH_4Cl, then the layers were separated. The aqueous layer was extracted with EtOAc. The combined organic extracts were dried over MgSO4, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (40 grams), eluting with a gradient of 0-50 % EtOAc/heptane, to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-cyano-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.53 g, 0.79 mmol, 63.7 % yield) as a light yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (s, 1 H) 7.50 - 7.56 (m, 1 H) 7.27 - 7.35 (m, 2 H) 7.17 - 7.24 (m, 1 H) 4.86 (br s, 1 H) 4.26 - 4.43 (m, 1 H) 4.22 (br d, *J*=13.48 Hz, 1 H) 3.86 (br d, *J*=13.68 Hz, 1 H) 3.72 (br d, *J*=13.06 Hz, 1 H) 3.43 - 3.61 (m, 1 H) 2.75 - 2.93 (m, 2 H) 1.45 (s, 9 H) 1.35 - 1.39 (m, 3 H) 1.18 (br d, *J*=6.63 Hz, 3 H) 1.10 (dd, *J*=6.63, 2.70 Hz, 6 H) 0.95 (dd, *J*=6.63, 2.49 Hz, 6 H). *m/z* (ESI, +ve ion): 675.1 $(M+H)^+$.

[0305] Step 4. 5-(4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidine-2-carbonitrile. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-cyano-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.53 g, 0.78 mmol) and trifluoroacetic acid (0.58 mL, 7.85 mmol) in 1,2-dichloroethane (3.9 mL). The reaction mixture was stirred and heated at 30 °C for 1 h, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0306] The previous residue was diluted with DCM (4 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then N,N'-diisopropylethylamine (1.64 mL, 9.42 mmol)

was added to the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (0.06 mL, 0.785 mmol) was added to the mixture dropwise. The mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (40 grams), eluting with a gradient of 0-5% MeOH/$CHCl_3$. This afforded 5-(4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidine-2-carbonitrile (0.28 g, 0.44 mmol, 56.7 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.51 (d, *J*=6.66 Hz, 1 H) 7.48 - 7.55 (m, 1 H) 7.18 - 7.33 (m, 3 H) 6.77 - 6.90 (m, 1 H) 6.19 (br d, *J*=16.79 Hz, 1 H) 5.72 - 5.78 (m, 1 H) 4.75 - 4.99 (m, 2 H) 4.19 - 4.36 (m, 1 H) 3.79 - 3.96 (m, 2 H) 2.76 - 2.93 (m, 2 H) 1.33 - 1.40 (m, 3 H) 1.23 - 1.29 (m, 2 H) 1.20 (br d, *J*=6.84 Hz, 2 H) 1.07 - 1.13 (m, 6 H) 0.93 - 0.99 (m, 6 H). *m/z* (ESI, +ve ion): 629.1 $(M+H)^+$.

Example 156

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0307] Step 1. 2,5,6-Trichloro-N-((4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)carbamoyl)nicotinamide. To a 100-mL round-bottomed flask was added 2,5,6-trichloronicotinamide (**Intermediate P**, 1.00 g, 4.44 mmol) and oxalyl chloride (3.33 mL, 6.65 mmol) in tetrahydrofuran (22.1 mL). The flask was fitted with a Findenser, then the mixture was stirred and heated at 80 °C for 45 min. The reaction mixture was concentrated *in vacuo*.

The crude mixture was carried into the next step of the synthesis, without further purification to prevent decomposition.

[0308] To a 150-mL round-bottomed flask was added (2,5,6-trichloronicotinoyl)carbamoyl isocyanate (crude material from previous step) in tetrahydrofuran (22.1 mL). The reaction mixture was cooled to -10 °C with a dilute acetone/dry ice bath, then a solution of 4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-amine (**Intermediate 220**, 1.32 g, 4.66 mmol) in THF (5 mL) was added to the reaction mixture. The mixture was allowed to stir under an inert (N2) atmosphere for 1 h. The reaction mixture was concentrated *in vacuo*. The crude material was triturated from DCM and heptane and the solids were collected by filtration. The solids were washed with heptane. This afforded 2,5,6-trichloro-*N*-((4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)carbamoyl)nicotinamide (2.30 g, 4.28 mmol, 97 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.52 (s, 1 H) 9.88 (br s, 1 H) 8.69 (s, 1 H) 5.75 (s, 2 H) 3.60 - 3.66 (m, 2 H) 1.68 - 1.80 (m, 2 H) 1.21 (d, *J*=6.84 Hz, 12 H) 1.00 - 1.05 (m, 3 H). *m/z* (ESI, +ve ion): 536.0 $(M+H)^+$.

[0309] Step 2. 6,7-Dichloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a 250-mL round-bottomed flask was added 2,5,6-trichloro-*N*-((4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)carbamoyl)nicotinamide (2.30 g, 4.28 mmol) in tetrahydrofuran (48.3 mL). The reaction mixture was cooled to 0 °C in a wet ice/water bath. Then potassium bis(trimethylsilyl)amide, 1M solution in tetrahydrofuran (5.36 mL, 5.36 mmol) was added via an addition funnel, dropwise to the reaction mixture over 5 min. The ice bath was removed and the reaction mixture was allowed to slowly warm to ambient temperature, while stirred under an inert (N_2) atmosphere for 1 h. More KHMDS (0.5 eq; 2 mL) was added dropwise to the reaction mixture. The reaction mixture was quenched with sat. aq. NH_4Cl (50 mL), then the mixture was diluted with 3:1 EtOAc/MeOH and brine solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The residue was diluted with DCM (50 mL), then agitated 5 min. The mixture was filtered and the filtrate was collected, then concentrated *in vacuo*. This afforded 6,7-dichloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (1.82 g, 3.65 mmol, 85 % yield) as a tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.32 - 12.59 (m, 1 H) 8.50 - 8.81 (m, 1 H) 3.67 - 3.74 (m, 2 H) 3.13 (dt, *J*=13.22, 6.56 Hz, 2 H) 1.78 - 1.87 (m, 2 H) 1.23 - 1.24 (m, 3 H) 1.11 - 1.14 (m, 6 H) 1.03 (s, 6 H). *m/z* (ESI, +ve ion): 500.0 $(M+H)^+$.

[0310] Step 3. 6,7-Dichloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a 100-mL round-bottomed flask was added 6,7-dichloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.45 g, 0.89 mmol) and sodium methoxyethoxide (0.44 g, 4.50 mmol) in 1,4-dioxane (4.0 mL). The reaction mixture was stirred and heated at 95 °C for 16 h. The flask was removed from the heat bath and the reaction mixture was allowed to cool to ambient temperature. The reaction mixture was concentrated *in vacuo*. The residue was diluted with DCM and filtered through a fine-fritted funnel and the filtrate (desired material) was collected. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (40 grams), eluting with a gradient of 0-45% EtOAc/heptane, to afford 6,7-dichloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.12 g, 0.26 mmol, 29.9 % yield) as white solid. 1H NMR (400 MHz, DMSO-*d6*) δ 12.29 (s, 1 H) 8.59 (s, 1 H) 4.45 - 4.55 (m, 2 H) 3.71 - 3.77 (m, 2 H) 3.33 - 3.35 (m, 3 H) 2.81 - 2.92 (m, 2H) 1.07 (d, *J*=6.63 Hz, 6 H) 0.97 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 468.0 $(M+H)^+$.

[0311] Step 4. 6-Chloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a 100-mL round-bottomed flask was added 6,7-dichloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.24 g, 0.51 mmol) and potassium acetate (0.15 g, 1.53 mmol) in 1,4-dioxane (3.31 mL). The reaction mixture was deoxygenated by bubbling (N_2) gas into the mixture for 5 min. Then (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.03 g, 0.05 mmol) was added to the reaction mixture. The reaction mixture was stirred and heated at 95 °C for 10 min. Then 2-fluorophenylboronic acid (0.08 g, 0.61 mmol), followed by water (0.05 mL) was added to the reaction mixture. The overall mixture was allowed to stir at 95 °C for 2 h. The reaction mixture was diluted with sat. aq. NH_4Cl and EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (40 grams), eluting with a gradient of 0-10% MeOH/DCM, to afford 6-chloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.22 g, 0.42 mmol, 83 % yield) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.24 (br s, 1 H) 8.57 (s, 1 H) 7.47 - 7.55 (m, 1 H) 7.25 - 7.35 (m, 2 H) 7.17 - 7.23 (m, 1 H) 4.39 - 4.49 (m, 2 H) 3.63 - 3.73 (m, 2 H)

3.29 - 3.30 (m, 3 H) 2.80 - 2.94 (m, 2 H) 1.08 (s, 6 H) 0.91 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 528.0 $(M+H)^+$.

[0312] Step 5. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added 6-chloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.22 g, 0.41 mmol) and N,N'-diisopropylethylamine (0.09 mL, 0.54 mmol) in acetonitrile (2.08 mL). Then phosphorous oxychloride (0.04 mL, 0.50 mmol) was added slowly to the reaction mixture. The flask was fitted with a Findenser, then the mixture was stirred and heated at 80 °C, while under an inert (N2) atmosphere for 45 min. The reaction mixture was removed from the heat bath and allowed to cool to ambient temperature.

[0313] The reaction mixture was cooled to 0 °C with a wet ice/water bath, while stirred. Then DIPEA (0.5 mL) was added slowly to the reaction mixture. Then a mixture of (2*R*,5*S*)-1-Boc-2,5-dimethylpiperazine (0.11 g, 0.52 mmol) in MeCN (1 mL) was added slowly to the reaction mixture. The ice bath was removed and the overall mixture was allowed to slowly warm to ambient temperature over 1 h. Another aliquot of DIPEA (0.6 mL) and (2*R*,5*S*)-1-Boc-2,5-dimethylpiperazine (0.11 g, 0.52 mmol) in MeCN (2 mL) was added to the mixture, then allowed the mixture to stir an additional 10 min. The mixture was diluted with EtOAc and sat. aq. NH_4Cl, then the layers were separated. The aqueous layer was extracted with EtOAc. The combined organic extracts were dried over MgSO4, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (40 grams), eluting with a gradient of 0-5% MeOH/DCM, to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.24 g, 0.33 mmol, 80 % yield) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.44 (s, 1 H) 7.48 - 7.56 (m, 1 H) 7.31 (br d, *J*=7.88 Hz, 2 H) 7.17 - 7.25 (m, 1 H) 4.82 (br s, 1 H) 4.34 - 4.49 (m, 2 H) 4.15 (br d, *J*=13.48 Hz, 1 H) 3.86 (br s, 1 H) 3.50 - 3.75 (m, 5 H) 2.91 - 3.07 (m, 2 H) 1.44 - 1.47 (m, 6 H) 1.39 (s, 9 H) 1.32 - 1.36 (m, 2 H) 1.15 - 1.20 (m, 3 H) 1.04 - 1.12 (m, 10 H). *m/z* (ESI, +ve ion): 724.1 $(M+H)^+$.

[0314] Step 6. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-

d]**pyrimidin-2(1*H*)-one**. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.24 g, 0.33 mmol) and trifluoroacetic acid (0.24 mL, 3.31 mmol) in 1,2-dichloroethane (3.92 mL). The reaction mixture was stirred and heated at 70 °C for 2 h, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification

[0315] The previous residue was diluted with DCM (4 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then N,N'-diisopropylethylamine (0.69 mL, 3.98 mmol) was added to the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (0.02 mL, 0.3 mmol was added to the mixture dropwise. The mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (40 grams), eluting with a gradient of 0-4% MeOH/$CHCl_3$. This afforded 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(2-methoxyethoxy)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.07 g, 0.10 mmol, 32.0 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (d, *J*=4.35 Hz, 1 H) 7.50 - 7.57 (m, 1 H) 7.20 - 7.36 (m, 3 H) 6.84 (td, *J*=17.10, 10.57 Hz, 1 H) 6.20 (dd, *J*=16.59, 1.87 Hz, 1 H) 5.73 - 5.79 (m, 1 H) 4.74 - 4.95 (m, 2 H) 4.38 - 4.55 (m, 3 H) 4.12 - 4.28 (m, 1 H) 3.80 - 3.95 (m, 2 H) 3.66 - 3.71 (m, 2 H) 3.29 - 3.31 (m, 3 H) 2.55 - 2.70 (m, 2 H) 1.31 - 1.39 (m, 3 H) 1.26 (br d, *J*=6.63 Hz, 1 H) 1.19 (d, *J*=6.63 Hz, 2 H) 1.06 (dd, *J*=6.63, 3.52 Hz, 6 H) 0.92 (br d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 678.4 $(M+H)^+$.

Example 157

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-(2-(dimethylamino)ethoxy)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0316] Step 1. 2,5-Dichloro-*N*-((4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide. To a 100-mL round-bottomed flask was added 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (**Intermediate 99B**, 1.00 g, 3.30 mmol) and oxalyl chloride (2.47 mL, 4.95 mmol) in tetrahydrofuran (16.49 mL). The flask was fitted

with a Findenser, and mixture was stirred and heated at 80 °C for 45 min. The reaction mixture was concentrated *in vacuo*. The crude mixture was carried into the next step of the synthesis, without further purification to prevent decomposition.

[0317] To a 100-mL round-bottomed flask was added (2,5-dichloro-6-(2-fluorophenyl)nicotinoyl)carbamoyl isocyanate (crude material from previous step) in tetrahydrofuran (16.55 mL). The reaction mixture was cooled to -10 °C with a dilute acetone/dry ice bath, then a solution of 4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-amine (**Intermediate 220**, 0.98 g, 3.46 mmol) in THF (3 mL) was added to the reaction mixture. The reaction mixture was allowed to stir under an inert (N2) atmosphere for 1 h. The reaction mixture was concentrated *in vacuo*. The crude material was triturated from EtOAc and heptane, then the solids were collected by filtration. The solids were washed with heptane. This afforded 2,5-dichloro-*N*-((4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide (1.705 g, 2.86 mmol, 87 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.56 (br s, 1 H) 9.94 (br s, 1 H) 8.64 (br s, 1 H) 7.62 (s, 1 H) 7.57 (br s, 1 H) 7.40 (br s, 2 H) 3.57 - 3.80 (m, 3 H) 1.79 (br s, 2 H) 1.22 (br s, 12 H) 1.03 (br s, 4 H). *m/z* (ESI, +ve ion): 596.0 $(M+H)^+$.

[0318] **Step 2. 6-Chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione**. To a 150-mL round-bottomed flask was added 2,5-dichloro-*N*-((4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide (1.48 g, 2.48 mmol) in tetrahydrofuran (12.44 mL). Then potassium bis(trimethylsilyl)amide, 1M solution in tetrahydrofuran (3.11 mL, 3.10 mmol) was added via an addition funnel dropwise to the reaction mixture over 5 min. The reaction mixture was allowed to stir under an inert (N_2) atmosphere for 1 h. More KHMDS (0.5 eq; 2 mL) was added dropwise to the reaction mixture. The reaction mixture was quenched with sat. aq. NH_4Cl (50 mL), then the mixture was diluted with 3:1 EtOAc/MeOH and brine solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. This afforded 6-chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (1.34 g, 2.39 mmol, 97 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.60 - 8.64 (m, 1 H) 7.48 - 7.54 (m, 1 H) 7.23 - 7.33 (m, 3 H) 7.15 - 7.21 (m, 1 H) 3.61 - 3.67 (m, 2 H) 3.13 - 3.19 (m, 2 H) 1.66 - 1.74 (m, 2 H) 1.21 - 1.23 (m, 3 H) 1.13 (d, *J*=6.63 Hz, 6 H) 0.96 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 560.0 $(M+H)^+$.

[0319] Step 3. 6-Chloro-1-(2-(2-(dimethylamino)ethoxy)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A resealable vial was charged with 6-chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.53 g, 0.94 mmol) and N,N'-dimethylethanolamine (1.91 mL, 18.93 mmol) in 1,4-dioxane (9.46 mL). The vial was sealed, then the reaction mixture was stirred and heated at 90 °C for 16 h. The reaction mixture was concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (80 grams), eluting with a gradient of 0-20% MeOH/DCM, to afford 6-chloro-1-(2-(2-(dimethylamino)ethoxy)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.255 g, 0.471 mmol, 49.8 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.03 - 12.41 (m, 1 H) 8.57 (s, 1 H) 7.48 - 7.55 (m, 1 H) 7.26 - 7.35 (m, 2 H) 7.17 - 7.23 (m, 1 H) 4.39 (t, *J*=5.91 Hz, 2 H) 2.88 (quin, *J*=6.63 Hz, 2 H) 2.64 (br t, *J*=5.80 Hz, 2 H) 2.22 (s, 6 H) 1.07 (d, *J*=6.63 Hz, 6 H) 0.91 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 541.1 $(M+H)^+$.

[0320] Step 4. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-(2-(dimethylamino)ethoxy)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added 6-chloro-1-(2-(2-(dimethylamino)ethoxy)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.22 g, 0.40 mmol) and N,N'-diisopropylethylamine (0.18 mL, 1.05 mmol) in acetonitrile (4.07 mL). Then phosphorous oxychloride (0.04 mL, 0.48 mmol) was added to the reaction mixture, then the mixture was heated and stirred at 80 °C for 15 min, while under an inert (N_2) atmosphere. The reaction mixture was removed from the heat bath and allowed to cool to rt.

[0321] The previous reaction mixture was cooled to 0 °C with a wet ice/water bath. Then DIPEA (2 mL) was added dropwise to the reaction mixture. Then a solution of (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (0.13 g, 0.61 mmol) in MeCN (1 mL) was added dropwise to the reaction mixture. The ice bath was removed and the mixture was allowed to warm to rt over 20 min. More DIPEA (2 mL) and (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (0.13 g, 0.61 mmol) was added to the reaction mixture and stirred an additional 10 min. The reaction mixture was diluted with EtOAc and sat. aq. NH_4Cl. The aqueous layer was extracted with EtOAc. The combined organic extracts were dried over MgSO4, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (80 grams),

eluting with a gradient of 0-25% 2M NH3·MeOH in CH2CL2, to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-(2-(dimethylamino)ethoxy)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as tan solid. The crude material (0.450 grams) was carried into the next step of the synthesis, without further purification. *m/z* (ESI, +ve ion): 737.2 $(M+H)^+$.

[0322] Step 5. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-(2-(dimethylamino)ethoxy)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-(2-(dimethylamino)ethoxy)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.30 g, 0.40 mmol) and trifluoroacetic acid (0.30 mL, 4.07 mmol) in 1,2-dichloroethane (4.07 mL). The reaction mixture was stirred and heated at 60 °C for 2 h, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0323] The previous residue was diluted with DCM (4 mL), the reaction mixture was cooled to 0 °C with a wet ice bath. Then N,N'-diisopropylethylamine (0.85 mL, 4.88 mmol) was added to the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (1.1M in THF) (0.37 mL, 0.40 mmol) was added to the mixture dropwise. The reaction mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (40 grams), eluting with a gradient of 0-25% MeOH/DCM. This afforded 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-(2-(dimethylamino)ethoxy)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.110 g, 0.159 mmol, 39.1 % yield) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (br d, *J*=4.35 Hz, 1 H) 7.48 - 7.58 (m, 1 H) 7.20 - 7.36 (m, 3 H) 6.76 - 6.91 (m, 1 H) 6.16 - 6.23 (m, 1 H) 5.71 - 5.80 (m, 1 H) 4.81 - 4.93 (m, 1 H) 4.57 - 4.63 (m, 2 H) 4.42 - 4.53 (m, 1 H) 4.13 - 4.24 (m, 1 H) 3.79 - 3.94 (m, 2 H) 3.39 - 3.45 (m, 2 H) 2.77 (s, 6 H) 2.61 - 2.73 (m, 2 H) 1.34 (br t, *J*=5.70 Hz, 3 H) 1.23 - 1.23 (m, 2 H) 1.16 - 1.20 (m, 2 H) 1.07 (dd, *J*=6.53, 4.25 Hz, 6 H) 0.93 (dd, *J*=6.53, 3.63 Hz, 6 H). . *m/z* (ESI, +ve ion): 691.3 $(M+H)^+$.

Example 158

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0324] **Step 1. 2,5-Dichloro-N-((4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide**. To a 100-mL round-bottomed flask was added 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (**Intermediate 99B**, 1.00 g, 3.30 mmol) and oxalyl chloride (2.47 mL, 4.95 mmol) in tetrahydrofuran (16.49 mL). The flask was fitted with a Findenser, and mixture was stirred and heated at 80 °C for 45 min. The reaction mixture was concentrated *in vacuo*. The crude mixture was carried into the next step of the synthesis, without further purification to prevent decomposition.

[0325] To a 100-mL round-bottomed flask was added (2,5-dichloro-6-(2-fluorophenyl)nicotinoyl)carbamoyl isocyanate (crude material from previous step) in tetrahydrofuran (16.49 mL). The reaction mixture was cooled to -10 °C with a dilute acetone/dry ice bath, then a solution of 4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-amine

(Intermediate 220, 0.98 g, 3.46 mmol) in THF (3 mL) was added to the reaction mixture. The reaction mixture was allowed to stir under an inert (N2) atmosphere for 1 h. The reaction mixture was concentrated *in vacuo*. The crude material was triturated from EtOAc and heptane, then the solids were collected by filtration. The solids were washed with heptane. This afforded 2,5-dichloro-*N*-((4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide (1.70 g, 2.86 mmol, 87 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.56 (br s, 1 H) 9.94 (br s, 1 H) 8.64 (br s, 1 H) 7.62 (s, 1 H) 7.57 (br s, 1 H) 7.40 (br s, 2 H) 3.57 - 3.80 (m, 3 H) 1.79 (br s, 2 H) 1.22 (br s, 12 H) 1.03 (br s, 4 H). *m/z* (ESI, +ve ion): 596.0 $(M+H)^+$.

[0326] Step 2. 6-Chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a 150-mL round-bottomed flask was added 2,5-dichloro-*N*-((4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide (1.48 g, 2.48 mmol) in tetrahydrofuran (12.41 mL). Then potassium bis(trimethylsilyl)amide, 1M solution in tetrahydrofuran (3.10 mL, 3.10 mmol) was added via an addition funnel dropwise to the reaction mixture over 5 min. The reaction mixture was allowed to stir under an inert (N_2) atmosphere for 1 h. More KHMDS (0.5 eq; 2 mL) was added dropwise to the reaction mixture. The reaction mixture was quenched with sat. aq. NH_4Cl (50 mL), then the mixture was diluted with 3:1 EtOAc/MeOH and brine solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. This afforded 6-chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (1.34 g, 2.39 mmol, 97 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.60 - 8.64 (m, 1 H) 7.48 - 7.54 (m, 1 H) 7.23 - 7.33 (m, 3 H) 7.15 - 7.21 (m, 1 H) 3.61 - 3.67 (m, 2 H) 3.13 - 3.19 (m, 2 H) 1.66 - 1.74 (m, 2 H) 1.21 - 1.23 (m, 3 H) 1.13 (d, *J*=6.63 Hz, 6 H) 0.96 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 560.0 $(M+H)^+$.

[0327] Step 3. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added 6-chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.35 g, 0.62 mmol) and N,N'-diisopropylethylamine (0.14 mL, 0.81 mmol) in acetonitrile (3.12 mL). Then phosphorous oxychloride (0.07 mL, 0.75 mmol) was added slowly to the reaction mixture. The flask was

fitted with a Findenser, then the mixture was stirred and heated at 80 °C, while under an inert (N2) atmosphere for 45 min. Another aliquot of N,N'-diisopropylethylamine (0.14 mL, 0.81 mmol) and phosphorous oxychloride (0.07 mL, 0.75 mmol) was added to the reaction mixture, then allowed the mixture to stir an additional 10 min. The reaction mixture was removed from the heat bath and allowed to cool to ambient temperature.

[0328] The previous reaction mixture was cooled to 0 °C with a wet ice/water bath, while stirred. Then DIPEA (2 mL) was added slowly to the mixture. Then a mixture of (2*R*,5*S*)-1-Boc-2,5-dimethylpiperazine (0.33 g, 0.78 mmol) in MeCN (1 mL) was added slowly to the reaction mixture. The ice bath was removed and the overall mixture was allowed to slowly warm to ambient temperature over 10 min. The mixture was diluted with EtOAc and sat. aq. NH_4Cl, then the layers were separated. The aqueous layer was extracted with EtOAc. The combined organic extracts were dried over MgSO4, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (40 grams), eluting with a gradient of 0-8% MeOH/DCM, to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.42 g, 0.55 mmol, 89 % yield) as tan solid. *m/z* (ESI, +ve ion): 756.0 $(M+H)^+$.

[0329] Step 4. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.20 g, 0.26 mmol) and trifluoroacetic acid (0.19 mL, 2.64 mmol) in 1,2-dichloroethane (2.03 mL). The reaction mixture was stirred and heated at 60 °C for 2 h, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0330] The previous residue was diluted with DCM (4 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then N,N'-diisopropylethylamine (0.55 mL, 3.17 mmol) was added to the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (0.02 mL, 0.26 mmol) was added to the mixture dropwise. The mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in*

vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (25 grams), eluting with a gradient of 0-100% EtOAc/heptane. This afforded 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.023 g, 0.032 mmol, 12.25 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.51 (d, *J*=7.46 Hz, 1 H) 7.48 - 7.55 (m, 1 H) 7.17 - 7.34 (m, 3 H) 6.84 (ddd, *J*=19.95, 16.84, 10.47 Hz, 1 H) 6.19 (br d, *J*=16.79 Hz, 1 H) 5.73 - 5.79 (m, 1 H) 4.76 - 4.98 (m, 2 H) 4.14 - 4.34 (m, 2 H) 3.80 - 3.94 (m, 2 H) 3.60 - 3.66 (m, 2 H) 2.79 - 2.96 (m, 2 H) 1.71 (sxt, *J*=7.51 Hz, 2 H) 1.33 - 1.40 (m, 3 H) 1.18 - 1.29 (m, 3 H) 1.10 - 1.14 (m, 6 H) 0.90 - 1.00 (m, 9 H). *m/z* (ESI, +ve ion): 710.0 $(M+H)^+$.

Example 159

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0331] Step 1. 6-Chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a 100-mL round-bottomed flask was added 1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 231**, 0.17 g, 0.31

mmol), sodium carbonate (0.10 g, 0.95 mmol) and vinylboronic acid pinacol ester (0.10 mL, 0.63 mmol) in 1,4-dioxane (2.32 mL)-water (0.55 mL) (4:1). The reaction mixture was deoxygenated by bubbling argong (gas) into the mixture for 5 min. Then (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.02 g, 0.03 mmol) was added to the reaction mixture. The reaction mixture was stirred and heated at 90 °C, while under an inert (N_2) atmosphere for 16 h. The reaction mixture was allowed to cool to rt, then the mixture was diluted was EtOAc and sat. aq. NH_4Cl. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (25 grams), eluting with a gradient of 0-50% EtOAc/heptane, to afford 6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.10 g, 0.21 mmol, 68.6 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.28 (s, 1 H) 8.59 (s, 1 H) 7.47 - 7.53 (m, 1 H) 7.25 - 7.33 (m, 2 H) 7.18 (td, *J*=7.36, 1.66 Hz, 1 H) 6.80 (dd, *J*=17.21, 10.37 Hz, 1 H) 6.57 (dd, *J*=17.21, 2.07 Hz, 1 H) 5.72 - 5.78 (m, 1 H) 2.90 - 3.01 (m, 2 H) 1.10 (d, *J*=6.63 Hz, 6 H) 0.94 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 480.0 $(M+H)^+$.

[0332] Step 2. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added 6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.10 g, 0.20 mmol) and N,N-diisopropylethylamine (0.04 mL, 0.27 mmol) in acetonitrile (2.08 mL). Then phosphorous oxychloride (0.02 mL, 0.25 mmol) was added to the reaction mixture, then the mixture was heated and stirred at 80 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was removed from the heat bath and allowed to cool to rt.

[0333] The reaction mixture was cooled to 0 °C with a wet ice/water bath. Then DIPEA (0.7 mL) was added dropwise to the reaction mixture. Then a solution of (2*R*,5*S*)-1-Boc-2,5-dimethylpiperazine (0.067 g, 0.313 mmol) in MeCN (1 mL) was added dropwise to the reaction mixture. The ice bath was removed and the mixture was allowed to warm to rt. More DIPEA (0.7 mL) and (2*R*,5*S*)-1-Boc-2,5-dimethylpiperazine (0.067 g, 0.313 mmol) was added to the reaction mixture and stirred an additional 10 min. The reaction mixture was diluted with EtOAc and sat. aq. NH_4Cl. The aqueous layer was extracted with EtOAc. The combined organic extracts were dried over MgSO4, filtered and concentrated *in vacuo*. The crude

material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (25 grams), eluting with a gradient of 0-50% EtOAc in CH2CL2, to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.132 g, 0.195 mmol, 94 % yield) as light-yellow solid. *m/z* (ESI, +ve ion): 676.2 $(M+H)^+$.

[0334] Step 3. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.13 g, 0.19 mmol) and trifluoroacetic acid (0.14 mL, 1.92 mmol) in 1,2-dichloroethane (1.92 mL). The reaction mixture was stirred and heated at 60 °C for 2 h, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0335] The previous residue was diluted with DCM (4 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then N,N'-diisopropylethylamine (0.40 mL, 2.30 mmol) was added to the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (0.01 mL, 0.19 mmol) was added to the mixture dropwise. The mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (25 grams), eluting with a gradient of 0-100% EtOAc/DCM. This afforded 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.082 g, 0.130 mmol, 67.7 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.50 (d, *J*=4.77 Hz, 1 H) 7.49 - 7.55 (m, 1 H) 7.19 - 7.35 (m, 3 H) 6.76 - 6.90 (m, 2 H) 6.56 (dd, *J*=17.31, 1.97 Hz, 1 H) 6.20 (dd, *J*=16.69, 1.97 Hz, 1 H) 5.72 - 5.79 (m, 2 H) 4.73 - 4.99 (m, 2 H) 4.21 (q, *J*=14.31 Hz, 1 H) 3.81 - 3.95 (m, 2 H) 2.64 - 2.78 (m, 2 H) 1.36 (t, *J*=5.60 Hz, 3 H) 1.27 (br d, *J*=6.63 Hz, 2 H) 1.20 (d, *J*=6.84 Hz, 2 H) 1.10 (dd, *J*=6.63, 3.11 Hz, 6 H) 0.95 (br d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 630.2 $(M+H)^+$.

Example 160

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-((dimethylamino)methyl)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Intermediate 235



[0336] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-((dimethylamino)methyl)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(3-chloro-2-(2-fluorophenyl)-8-(2-formyl-4,6-diisopropylpyrimidin-5-yl)-7-oxo-7,8-dihydro-1,6-naphthyridin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 235**, 0.16 g, 0.24 mmol) and dimethylamine (2.0M in THF) (0.18 mL, 0.36 mmol) in tetrahydrofuran (2.43 mL). Then glacial acetic acid (1.2 μL, 0.024 mmol), followed by sodium cyanoborohydride (0.03 mL, 0.73 mmol) was added to the reaction mixture. The overall reaction mixture was allowed to stir at rt, while under an inert (N_2) atmosphere for 30 min. The reaction mixture was quenched with MeOH (0.5 mL), then allowed the mixture to stir 5 min. Then the mixture was diluted with EtOAc and sat. aq.

$NaHCO_3$. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material (0.195 grams) was carried into the next step of the synthesis, without further purification. *m/z* (ESI, +ve ion): 707.2 $(M+H)^+$.

[0337] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-((dimethylamino)methyl)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-((dimethylamino)methyl)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.17 g, 0.24 mmol) and trifluoroacetic acid (0.17 mL, 2.40 mmol) in 1,2-dichloroethane (2.40 mL). The reaction mixture was stirred and heated at 60 °C for 20 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0338] The previous residue was diluted with DCM (4 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then N,N'-diisopropylethylamine (0.50 mL, 2.88 mmol) was added to the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (1.1M in THF) (0.21 mL, 0.24 mmol) was added to the reaction mixture. The mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-25% 2M NH3 in MeOH/DCM. This afforded 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-((dimethylamino)methyl)-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.083 g, 0.126 mmol, 52.2 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (d, *J*=4.98 Hz, 1 H) 7.52 (q, *J*=7.12 Hz, 1 H) 7.18 - 7.34 (m, 3 H) 6.84 (td, *J*=17.41, 10.37 Hz, 1 H) 6.20 (br d, *J*=17.62 Hz, 1 H) 5.76 (br d, *J*=10.37 Hz, 1 H) 4.76 - 4.97 (m, 2 H) 4.46 - 4.58 (m, 1 H) 4.04 - 4.28 (m, 2 H) 3.98 (br s, 2 H) 3.69 - 3.95 (m, 2 H) 3.15 - 3.21 (m, 2 H) 2.66 - 2.80 (m, 2 H) 1.35 (br t, *J*=5.39 Hz, 3 H) 1.15 - 1.29 (m, 6 H) 1.09 (dd, *J*=6.53, 2.80 Hz, 6 H) 0.95 (br d, *J*=6.22 Hz, 6 H). *m/z* (ESI, +ve ion): 661.1 $(M+H)^+$.

Example 161

1-[2-(Azetidin-1-ylmethyl)-4,6-diisopropyl-pyrimidin-5-yl]-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one



Intermediate 235



[0339] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(1-(2-(azetidin-1-ylmethyl)-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(3-chloro-2-(2-fluorophenyl)-8-(2-formyl-4,6-diisopropylpyrimidin-5-yl)-7-oxo-7,8-dihydro-1,6-naphthyridin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 235**, 0.200 g, 0.295 mmol) and azetidine (0.02 mL, 0.443 mmol) in tetrahydrofuran (2.95 mL). Then glacial acetic acid (1.7 µl, 0.03 mmol), followed by sodium cyanoborohydride (0.056 g, 0.886 mmol) was added to the reaction mixture. The overall reaction mixture was allowed to stir at rt, while under an inert (N_2) atmosphere for 30 min. The reaction mixture was quenched with MeOH (0.5 mL) and allowed the mixture to stir 5 min. Then the mixture was diluted with EtOAc and sat. aq. $NaHCO_3$. The

layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material (0.212 grams) was carried into the next step of the synthesis, without further purification. *m/z* (ESI, +ve ion): 719.3 $(M+H)^+$.

[0340] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-(azetidin-1-ylmethyl)-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(1-(2-(azetidin-1-ylmethyl)-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.18 g, 0.25 mmol) and trifluoroacetic acid (0.18 mL, 2.5 mmol) in 1,2-dichloroethane (2.4 mL). The reaction mixture was stirred and heated at 60 °C for 20 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0341] The previous residue was diluted with DCM (4 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then N,N'-diisopropylethylamine (0.5 mL, 3.0 mmol) was added to the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (1.1M in THF) (0.2 mL, 0.25 mmol) was added to the mixture dropwise. The reaction mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-25% 2M NH3-MeOH/DCM, This afforded 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-(azetidin-1-ylmethyl)-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.008 g, 0.012 mmol, 4.75 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (d, *J*=4.77 Hz, 1 H) 7.49 - 7.61 (m, 1 H) 7.32 (s, 1 H) 7.26 - 7.30 (m, 1 H) 7.21 (br d, *J*=7.67 Hz, 1 H) 6.74 - 6.94 (m, 1 H) 6.16 - 6.23 (m, 1 H) 5.67 - 5.81 (m, 1 H) 4.76 - 4.94 (m, 1 H) 4.41 - 4.55 (m, 1 H) 4.12 - 4.28 (m, 1 H) 3.87 (br s, 2 H) 3.62 - 3.74 (m, 2 H) 3.45 - 3.55 (m, 1 H) 2.68 (br s, 4 H) 2.32 - 2.48 (m, 1 H) 1.79 - 2.05 (m, 2 H) 1.35 (br t, *J*=5.80 Hz, 2 H) 1.26 (br d, *J*=7.26 Hz, 1 H) 1.19 (br d, *J*=7.46 Hz, 2 H) 1.02 - 1.13 (m, 6 H) 0.99 (s, 1 H) 0.93 (br dd, *J*=6.43, 3.32 Hz, 7 H). *m/z* (ESI, +ve ion): 673.3 $(M+H)^+$.

Example 162

6-Chloro-1-[4,6-diisopropyl-2-(morpholinomethyl)pyrimidin-5-yl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one



Intermediate 235



[0342] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(morpholinomethyl)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(3-chloro-2-(2-fluorophenyl)-8-(2-formyl-4,6-diisopropylpyrimidin-5-yl)-7-oxo-7,8-dihydro-1,6-naphthyridin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 235**, 0.21 g, 0.29 mmol) and morpholine (0.03 mL, 0.44 mmol) in tetrahydrofuran (2.95 mL). Then glacial acetic acid (1.7 µl, 0.03 mmol), followed by sodium cyanoborohydride (0.056 g, 0.88 mmol) was added to the reaction mixture. The overall reaction mixture was allowed to stir at rt, while under an inert (N_2) atmosphere for 30 min. The reaction mixture was quenched with MeOH (0.5 mL) and allowed the mixture to stir 5 min. Then the mixture was diluted with EtOAc and sat. aq. $NaHCO_3$. The

layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material (0.263 grams) was carried into the next step of the synthesis, without further purification. *m/z* (ESI, +ve ion): 749.3 $(M+H)^+$.

[0343] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(morpholinomethyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(morpholinomethyl)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.220 g, 0.294 mmol) and trifluoroacetic acid (0.2 mL, 2.94 mmol) in 1,2-dichloroethane (2.4 mL). The reaction mixture was stirred and heated at 60 °C for 20 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0344] The previous residue was diluted with DCM (4 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then N,N'-diisopropylethylamine (0.61 mL, 3.52 mmol) was added to the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (1.1M in THF) (0.26 mL, 0.294 mmol) was added to the mixture dropwise. The mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-25% 2M NH3-MeOH/DCM. This afforded 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(morpholinomethyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.085 g, 0.121 mmol, 41.2 % yield) as white solid. ^1H NMR (500 MHz, DMSO-d_6) δ 8.48 (d, *J*=6.62 Hz, 1 H) 7.52 (q, *J*=7.14 Hz, 1 H) 7.24 - 7.34 (m, 2 H) 7.16 - 7.21 (m, 1 H) 6.79 - 6.90 (m, 1 H) 6.16 - 6.22 (m, 1 H) 5.73 - 5.78 (m, 1 H) 4.83 - 4.93 (m, 1 H) 4.14 - 4.26 (m, 1 H) 3.86 - 3.94 (m, 1 H) 3.74 (s, 2 H) 3.54 - 3.59 (m, 4 H) 3.17 (d, *J*=5.32 Hz, 2 H) 2.61 - 2.75 (m, 2 H) 2.52 - 2.60 (m, 4 H) 1.31 - 1.37 (m, 3 H) 1.21 - 1.28 (m, 2 H) 1.18 (d, *J*=6.75 Hz, 2 H) 1.03 - 1.11 (m, 6 H) 0.92 (br d, *J*=6.49 Hz, 6 H). *m/z* (ESI, +ve ion): 703.1 $(M+H)^+$.

Example 163

6-Chloro-1-[4,6-diisopropyl-2-(pyrrolidin-1-ylmethyl)pyrimidin-5-yl]-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one



Intermediate 235



[0345] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(pyrrolidin-1-ylmethyl)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(3-chloro-2-(2-fluorophenyl)-8-(2-formyl-4,6-diisopropylpyrimidin-5-yl)-7-oxo-7,8-dihydro-1,6-naphthyridin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 235**, 0.200 g, 0.295 mmol) and pyrrolidine (0.03 mL, 0.443 mmol) in tetrahydrofuran (2.95 mL). Then glacial acetic acid (1.7 µl, 0.030 mmol), followed by sodium cyanoborohydride (0.056 g, 0.886 mmol) was added to the reaction mixture. The overall reaction mixture was allowed to stir at rt, while under an inert (N_2) atmosphere for 30 min. The reaction mixture was quenched with MeOH (0.5 mL) and allowed the mixture to stir 5 min. Then the mixture was diluted with EtOAc and sat. aq. $NaHCO_3$. The layers were separated

and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material (0.238 grams) was carried into the next step of the synthesis, without further purification. *m/z* (ESI, +ve ion): 733.3 $(M+H)^+$.

[0346] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(pyrrolidin-1-ylmethyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(pyrrolidin-1-ylmethyl)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.190 g, 0.259 mmol) and trifluoroacetic acid (0.19 mL, 2.59 mmol) in 1,2-dichloroethane (2.40 mL). The reaction mixture was stirred and heated at 60 °C for 20 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0347] The previous residue was diluted with DCM (4 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then N,N'-diisopropylethylamine (0.54 mL, 3.11 mmol) was added to the reaction mixture and stirred 2 min. Then acryloyl chloride (1.1M in THF) (0.23 mL, 0.259 mmol) was added to the mixture dropwise. The mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-25% 2M NH3-MeOH/DCM. This afforded 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-(pyrrolidin-1-ylmethyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.060 g, 0.087 mmol, 33.7 % yield) as light-yellow solid. ^{1}H NMR (500 MHz, DMSO-d_6) δ 8.48 (d, *J*=6.49 Hz, 1 H) 7.44 - 7.55 (m, 1 H) 7.24 - 7.34 (m, 2 H) 7.18 (td, *J*=7.46, 1.69 Hz, 1 H) 6.79 - 6.89 (m, 1 H) 6.19 (dt, *J*=16.64, 2.12 Hz, 1 H) 5.76 (ddd, *J*=10.35, 6.07, 2.27 Hz, 1 H) 4.83 - 4.93 (m, 1 H) 4.78 (br s, 1 H) 4.10 - 4.25 (m, 1 H) 3.78 - 3.91 (m, 4 H) 2.59 - 2.74 (m, 6 H) 1.61 - 1.69 (m, 4 H) 1.30 - 1.38 (m, 3 H) 1.21 - 1.29 (m, 2 H) 1.18 (d, *J*=6.75 Hz, 2 H) 1.02 - 1.11 (m, 6 H) 0.89 - 0.96 (m, 6 H). *m/z* (ESI, +ve ion): 687.4 $(M+H)^+$.

Example 164-1 (Peak 1)

6-Chloro-1-(6-cyclopropyl-2-isopropyl-4-methyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one



[0348] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(6-cyclopropyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.275 g, 0.393 mmol, **Intermediate 202**) in tetrahydrofuran (2.86 mL). The reaction mixture was degassed by bubbling argon (gas) into the mixture for 5 min. Then cyclopropylzinc bromide (0.5M in THF) (1.37 mL, 0.687 mmol) and xantphos pd g3 (0.011 g, 0.012 mmol) was added to the reaction mixture. The overall reaction mixture was allowed to stir under an inert (N_2) atmosphere at rt for 30 min. The reaction mixture was quenched with sat. aq. NH_4Cl and the mixture was diluted with EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and

concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (25 grams), eluting with a gradient of 0-40% EtOAc in CH2CL2, to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(6-cyclopropyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.235 g, 0.355 mmol, 90 % yield) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.42 (d, *J*=6.22 Hz, 1 H) 7.48 - 7.56 (m, 1 H) 7.20 - 7.35 (m, 3 H) 7.04 (d, *J*=5.80 Hz, 1 H) 4.81 (br s, 1 H) 4.36 (br s, 1 H) 4.00 - 4.16 (m, 1 H) 3.85 (br s, 1 H) 3.69 (br d, *J*=13.48 Hz, 1 H) 3.43 - 3.59 (m, 1 H) 2.54 - 2.64 (m, 1 H) 1.99 - 2.05 (m, 1 H) 1.87 (d, *J*=12.44 Hz, 3 H) 1.44 (s, 9 H) 1.30 - 1.38 (m, 3 H) 1.11 - 1.20 (m, 4 H) 0.86 - 1.02 (m, 9 H). *m/z* (ESI, +ve ion): 661.2 $(M+H)^+$.

[0349] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(6-cyclopropyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(6-cyclopropyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.230 g, 0.348 mmol) and trifluoroacetic acid (0.25 mL, 3.48 mmol) in 1,2-dichloroethane (3.48 mL). The reaction mixture was stirred and heated at 60 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0350] The previous residue was diluted with 1,2-dichloroethane (3.48 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.72 mL, 4.17 mmol) was added to the reaction mixture and stirred 2 min. Then acryloyl chloride (0.31 mL, 0.348 mmol) was added to the mixture dropwise and stirred under an inert (N_2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-5% MeOH/DCM. The racemic mixture of atropisomers were separated by SFC; (Column: IC, 5 μm, 21 x 250 mm, F=80 mL/min, 40% MeOH/60% Carbon dioxide) This afforded Peak 1: 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(6-cyclopropyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.060 g, 0.098 mmol, 28.0 % yield) as off-white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.44 (d, *J*=3.11 Hz,

1 H) 7.49 - 7.56 (m, 1 H) 7.21 - 7.35 (m, 3 H) 7.04 (s, 1 H) 6.83 (td, *J*=16.90, 10.57 Hz, 1 H) 6.19 (dd, *J*=16.59, 2.07 Hz, 1 H) 5.72 - 5.78 (m, 1 H) 4.84 (br s, 1 H) 4.77 (br s, 1 H) 4.10 - 4.20 (m, 1 H) 3.79 - 3.92 (m, 2 H) 3.45 - 3.53 (m, 1 H) 2.55 - 2.65 (m, 1 H) 1.90 - 2.05 (m, 1 H) 1.87 (s, 3 H) 1.29 - 1.36 (m, 3 H) 1.24 (br d, *J*=6.63 Hz, 1 H) 1.17 (d, *J*=6.63 Hz, 2 H) 0.98 - 1.03 (m, 3 H) 0.91 (br d, *J*=6.63 Hz, 7 H). *m/z* (ESI, +ve ion): 615.2 $(M+H)^+$.

Example 164-2 (Peak 2)

[0351] The racemic mixture of atropisomers were separated by SFC; (Column: IC, 5 µm, 21 x 250 mm, F=80 mL/min, 40% MeOH/60% Carbon dioxide) This afforded Peak 2: 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(6-cyclopropyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.055 g, 0.089 mmol, 25.7 % yield) as off-white solid.

Example 165-1 (Peak 1)

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4,6-dimethylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0352] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4,6-dimethylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.250 g, 0.357 mmol, **Intermediate 202**) and sodium carbonate (0.114 g, 1.071 mmol) in 1,4-dioxane (1.0 mL). The reaction mixture was degassed by bubbling argon (gas) into the mixture for 5 min. Then xantphos pd g3 (0.034 g, 0.036 mmol), water (0.2 mL) and methylboronic acid (1.069 g, 17.86 mmol) was added to the reaction mixture. The overall reaction mixture was stirred and heated at 100 °C for 2 h. The reaction mixture was cooled to rt, then the mixture was diluted with sat. aq. $NaHCO_3$ and EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (25 grams), eluting with a gradient of 0-100% EtOAc in CH2CL2, to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4,6-dimethylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.100 g, 0.157 mmol, 44.1 % yield) as light-yellow solid. *m/z* (ESI, +ve ion): 635.2 $(M+H)^+$.

[0353] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4,6-dimethylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4,6-dimethylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.100 g, 0.157 mmol) and trifluoroacetic acid (0.1 mL, 1.574 mmol) in 1,2-dichloroethane (1.5 mL). The reaction mixture was stirred and heated at 60 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0354] The previous residue was diluted with 1,2-dichloroethane (1.5 mL), then cooled the reaction mixture to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.3 mL, 1.889 mmol) was added to the reaction mixture and stirred 2 min. Then acryloyl chloride (0.14 mL, 0.157 mmol) was added to the mixture dropwise and stirred under an inert (N_2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts

were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-5% MeOH/DCM. The racemic mixture was separated by SFC; (Column, IA, 5 µm, 21 x 250 mm, F=80 mL/min, 15% Isopropanol/85% Carbon dioxide) This afforded the separated Atropisomers, as Peak 1 (M) 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4,6-dimethylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.030 g, 0.051 mmol, 32.3 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.38 (d, *J*=3.11 Hz, 1 H) 7.42 - 7.48 (m, 1 H) 7.14 - 7.27 (m, 3 H) 6.96 (s, 1 H) 6.76 (td, *J*=16.79, 10.37 Hz, 1 H) 6.12 (dd, *J*=16.69, 1.76 Hz, 1 H) 5.65 - 5.71 (m, 1 H) 4.78 (br s, 1 H) 4.70 (br s, 1 H) 4.03 - 4.14 (m, 1 H) 3.73 - 3.87 (m, 2 H) 2.55 - 2.62 (m, 1 H) 2.32 - 2.38 (m, 3 H) 1.81 (s, 3 H) 1.26 (t, *J*=6.63 Hz, 3 H) 1.18 (br d, *J*=6.63 Hz, 2 H) 1.11 (br d, *J*=6.84 Hz, 2 H) 0.98 (d, *J*=6.63 Hz, 3 H) 0.88 (d, *J*=6.63 Hz, 3 H). *m/z* (ESI, +ve ion): 589.1 $(M+H)^+$.

Example 165-2 (Peak 2)

[0355] The racemic mixture was separated by SFC; (Column, IA, 5 µm, 21 x 250 mm, F=80 mL/min, 15% Isopropanol/85% Carbon dioxide) This afforded Peak 2 (P) 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4,6-dimethylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.035 g, 0.059 mmol, 37.7 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.47 (s, 1 H) 7.48 - 7.56 (m, 1 H) 7.22 - 7.34 (m, 3 H) 7.04 (s, 1 H) 6.83 (ddd, *J*=16.59, 13.89, 10.57 Hz, 1 H) 6.19 (dd, *J*=16.69, 2.18 Hz, 1 H) 5.71 - 5.78 (m, 1 H) 4.86 (br s, 1 H) 4.76 (br s, 1 H) 4.08 - 4.20 (m, 1 H) 3.79 - 3.96 (m, 2 H) 2.61 - 2.70 (m, 1 H) 2.39 - 2.45 (m, 3 H) 1.91 (s, 3 H) 1.33 (br t, *J*=6.32 Hz, 3 H) 1.23 (br d, *J*=6.43 Hz, 2 H) 1.16 (br d, *J*=6.84 Hz, 2 H) 1.04 (d, *J*=6.63 Hz, 3 H) 0.92 (d, *J*=6.63 Hz, 3 H). *m/z* (ESI, +ve ion): 589.1 $(M+H)^+$.

Example 166-1 (Peak 1)

6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(6-ethyl-2-isopropyl-4-methyl-3-pyridyl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one



[0356] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(6-ethyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.230 g, 0.329 mmol, **Intermediate 202**) in tetrahydrofuran (1.6 mL). The reaction mixture was degassed by bubbling argon (gas) into the mixture for 5 min. Then triethylaluminum (1.0M in THF) (0.8 mL, 0.821 mmol) and tetrakis (0.038 g, 0.033 mmol) was added to the reaction mixture. The mixture was stirred and heated at 60 °C for 16 h, while under an inert (N_2) atmosphere. The reaction was quench with sat. aq. $NaHCO_3$ and diluted with mixture with EtOAc and brine. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*.

The crude material (0.250 g) was carried into the next step of the synthesis, without further purification. *m/z* (ESI, +ve ion): 649.3 $(M+H)^+$.

[0357] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(6-ethyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(6-ethyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.213 g, 0.328 mmol) and trifluoroacetic acid (0.2 mL, 3.28 mmol) in 1,2-dichloroethane (1.5 mL). The reaction mixture was stirred and heated at 60 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated *in vacuo*. This material was carried directly into the next step of the synthesis, without further purification.

[0358] The previous residue was diluted with 1,2-dichloroethane (1.5 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.6 mL, 3.94 mmol) was added to the reaction mixture and stirred 2 min. Then acryloyl chloride (0.29 mL, 0.328 mmol) was added to the mixture dropwise and stirred under an inert (N_2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through an Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-5% MeOH/DCM. The racemates were separated by SFC; (Column, IA, 5 μm, 21 x 250 mm, F= 80 mL/min, 15% MeOH/85: Carbon dioxide). This afforded the separated Atropisomers as Peak 1: 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(6-ethyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.040 g, 0.066 mmol, 20.21 % yield) as off-white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 (d, *J*=2.90 Hz, 1 H) 7.47 - 7.57 (m, 1 H) 7.20 - 7.35 (m, 3 H) 7.05 (s, 1 H) 6.78 - 6.90 (m, 1 H) 6.20 (dd, *J*=16.59, 1.87 Hz, 1 H) 5.72 - 5.79 (m, 1 H) 4.86 (br s, 1 H) 4.77 (br s, 1 H) 4.10 - 4.22 (m, 1 H) 3.80 - 3.95 (m, 2 H) 3.50 (br d, *J*=10.37 Hz, 1 H) 2.55 - 2.75 (m, 3 H) 1.90 (s, 3 H) 1.34 (br t, *J*=6.74 Hz, 3 H) 1.16 - 1.29 (m, 6 H) 1.06 (d, *J*=6.63 Hz, 3 H) 0.97 (d, *J*=6.63 Hz, 3 H). *m/z* (ESI, +ve ion): 603.2 $(M+H)^+$.

Example 166-2 (Peak 2)

[0359] The racemates were separated by SFC; (Column, IA, 5 μm, 21 x 250 mm, F= 80 mL/min, 15% MeOH/85: Carbon dioxide). This afforded the separated Atropisomers as Peak

2: 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(6-ethyl-2-isopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.030 g, 0.050 mmol, 15.16 % yield) as off-white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (s, 1 H) 7.49 - 7.56 (m, 1 H) 7.22 - 7.35 (m, 3 H) 7.06 (s, 1 H) 6.78 - 6.89 (m, 1 H) 6.20 (dd, *J*=16.59, 2.07 Hz, 1 H) 5.73 - 5.79 (m, 1 H) 4.87 (br s, 1 H) 4.76 (br s, 1 H) 4.14 (br d, *J*=13.06 Hz, 1 H) 3.81 - 3.96 (m, 2 H) 3.40 - 3.53 (m, 1 H) 2.62 - 2.74 (m, 3 H) 1.89 - 1.97 (m, 3 H) 1.34 (br t, *J*=6.22 Hz, 3 H) 1.16 - 1.27 (m, 6 H) 1.05 (d, *J*=6.84 Hz, 3 H) 0.94 (d, *J*=6.63 Hz, 3 H). *m/z* (ESI, +ve ion): 603.2 $(M+H)^+$.

Example 167

6-Chloro-1-(4,6-diisopropyl-2-methyl-pyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2-one



[0360] Step 1: tert-Butylbutyl (2R,5S)-4-(6-chloro-1-(4,6-diisopropyl-2-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-

fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.300 g, 0.411 mmol, **Intermediate 232**) and sodium carbonate (0.131 g, 1.234 mmol) in 1,4-dioxane (1.6 mL). The reaction mixture was degassed by bubbling argon (gas) into the mixture for 5 min. Then xantphos pd g3 (0.020 g, 0.021 mmol), water (0.4 mL) and methylboronic acid (1.2 mL, 20.57 mmol) was added into the reaction mixture. The overall reaction mixture was stirred and heated at 80 °C for 8 h. The reaction mixture was cooled to rt, then the mixture was diluted with sat. aq. $NaHCO_3$ and EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (25 grams), eluting with a gradient of 0-50% EtOAc in CH2CL2, to provide tert-butyl (2R,5S)-4-(6-chloro-1-(4,6-diisopropyl-2-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.207 g, 0.312 mmol, 76 % yield) as light-yellow solid. *m/z* (ESI, +ve ion): 664.2 $(M+H)^+$.

[0361] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(6-chloro-1-(4,6-diisopropyl-2-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.200 g, 0.301 mmol) and trifluoroacetic acid (0.2 mL, 3.01 mmol) in 1,2-dichloroethane (3.0 mL). The reaction mixture was stirred and heated at 60 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated in vacuo. This material was carried directly into the next step of the synthesis, without further purification.

[0362] The previous residue was diluted with 1,2-dichloroethane (3.0 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.6 mL, 3.61 mmol) was added into the reaction mixture and stirred 2 min. Then acryloyl chloride (0.2 mL, 0.301 mmol) was added into the mixture dropwise and stirred under an inert (N_2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (25 grams), eluting with a gradient of 0-8% MeOH/DCM. This afforded 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-

methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.130 g, 0.210 mmol, 69.8 % yield) as off-white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.42 (d, J=4.15 Hz, 1 H) 7.41 - 7.49 (m, 1 H) 7.19 - 7.29 (m, 2 H) 7.10 - 7.17 (m, 1 H) 6.77 (td, J=17.00, 10.57 Hz, 1 H) 6.13 (dd, J=16.69, 1.97 Hz, 1 H) 5.65 - 5.73 (m, 1 H) 4.76 - 4.87 (m, 1 H) 4.71 (br s, 1 H) 4.03 - 4.22 (m, 1 H) 3.65 - 3.87 (m, 2 H) 2.50 - 2.69 (m, 5 H) 1.16 - 1.38 (m, 5 H) 1.12 (br d, J=6.84 Hz, 2 H) 1.00 (br d, J=3.73 Hz, 3 H) 0.99 (br d, J=3.73 Hz, 3 H) 0.82 - 0.90 (m, 6 H). m/z (ESI, +ve ion): 618.0 $(M+H)^+$.

Example 168

6-Chloro-1-(2-cyclopropyl-4,6-diisopropyl-pyrimidin-5-yl)-4-[(2S,5R)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2-one



[0363] Step 1. *tert*-Butyl (2R,5S)-4-(6-chloro-1-(2-cyclopropyl-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-

carboxylate (0.300 g, 0.411 mmol, **Intermediate 232**) in tetrahydrofuran (4.1 mL). The reaction mixture was degassed by bubbling argon (gas) into the mixture for 5 min. Then cyclopropylzinc bromide (0.5M in THF) (1.4 mL, 0.720 mmol) and xantphos pd g3 (0.012 g, 0.012 mmol) was added into the reaction mixture. The overall reaction mixture was allowed to stir under an inert (N_2) atmosphere at rt for 30 min. The reaction mixture was quenched with sat. aq. NH_4Cl and diluted the mixture with EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (25 grams), eluting with a gradient of 0-40% EtOAc in CH2CL2, to provide tert-butyl (2R,5S)-4-(6-chloro-1-(2-cyclopropyl-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.275 g, 0.398 mmol, 97 % yield) as tan solid. *m/z* (ESI, +ve ion): 690.2 $(M+H)^+$.

[0364] Step 2. 4-((2S,5R)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-cyclopropyl-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(6-chloro-1-(2-cyclopropyl-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.265 g, 0.384 mmol) and trifluoroacetic acid (0.2 mL, 3.84 mmol) in 1,2-dichloroethane (3.8 mL). The reaction mixture was stirred and heated at 60 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated in vacuo. This material was carried directly into the next step of the synthesis, without further purification.

[0365] The previous residue was diluted with 1,2-dichloroethane (3.8 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.8 mL, 4.61 mmol) was added into the reaction mixture and stirred 2 min. Then acryloyl chloride (0.3 mL, 0.384 mmol) was added into the mixture dropwise and stirred under an inert (N_2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-8% MeOH/DCM. This afforded 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-cyclopropyl-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.180 g,

0.279 mmol, 72.8 % yield) as off-white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.41 (d, J=3.94 Hz, 1 H) 7.39 - 7.54 (m, 1 H) 7.19 - 7.34 (m, 2 H) 7.10 - 7.19 (m, 1 H) 6.77 (td, J=16.90, 10.57 Hz, 1 H) 6.13 (dd, J=16.59, 2.07 Hz, 1 H) 5.64 - 5.73 (m, 1 H) 4.75 - 4.87 (m, 1 H) 4.71 (br s, 1 H) 4.01 - 4.22 (m, 1 H) 3.70 - 3.92 (m, 2 H) 2.47 - 2.63 (m, 2 H) 2.04 - 2.13 (m, 1 H) 1.07 - 1.33 (m, 7 H) 0.90 - 1.02 (m, 10 H) 0.85 (dd, J=6.53, 3.01 Hz, 6 H). m/z (ESI, +ve ion): 644.2 $(M+H)^+$.

Example 169

6-Chloro-4-[(2S,5R)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-ethyl-4,6-diisopropyl-pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2-one



[0366] Step 1. *tert*-Butyl (2R,5S)-4-(6-chloro-1-(2-ethyl-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.200 g, 0.274 mmol, **Intermediate 232**) in tetrahydrofuran (1.6 mL). The reaction mixture was

degassed by bubbling argon (gas) into the mixture for 5 min. Then triethylaluminum (1.0M in THF) (0.6 mL, 0.686 mmol) and tetrakis (0.032 g, 0.027 mmol) was added into the reaction mixture. The mixture was stirred and heated at 60 °C for 16 h, while under an inert (N_2) atmosphere. The reaction was quench with sat. aq. $NaHCO_3$ and diluted with mixture with EtOAc and brine. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was carried into the next step of the synthesis, without further purification. *m/z* (ESI, +ve ion): 678.2 $(M+H)^+$.

[0367] Step 2. 4-((2S,5R)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-ethyl-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(6-chloro-1-(2-ethyl-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.036 g, 0.053 mmol) and trifluoroacetic acid (0.1 mL, 0.531 mmol) in 1,2-dichloroethane (1.5 mL). The reaction mixture was stirred and heated at 60 °C for 30 min, while under an inert (N2) atmosphere. The reaction mixture was concentrated in vacuo. This material was carried directly into the next step of the synthesis, without further purification.

[0368] The previous residue was diluted with 1,2-dichloroethane (0.5 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.1 mL, 0.637 mmol) was added into the reaction mixture and stirred 2 min. Then acryloyl chloride (0.1 mL, 0.053 mmol) was added into the mixture dropwise and stirred under an inert (N2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. NaHCO3, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts were dried over MgSO4, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-8% MeOH/DCM. This afforded 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-ethyl-4,6-diisopropylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.033 g, 0.052 mmol, 98 % yield) as off-white solid. 1H NMR (400 MHz, DMSO-d6) δ 8.48 (d, J=4.15 Hz, 1 H) 7.49 - 7.55 (m, 1 H) 7.26 - 7.34 (m, 2 H) 7.20 (td, J=7.36, 1.66 Hz, 1 H) 6.77 - 6.89 (m, 1 H) 6.19 (dd, J=16.59, 2.28 Hz, 1 H) 5.73 - 5.78 (m, 1 H) 4.87 (br s, 1 H) 4.78 (br s, 1 H) 4.10 - 4.25 (m, 1 H) 3.79 - 3.95 (m, 2 H) 2.87 (q, J=7.67 Hz, 2 H) 2.61 - 2.73 (m, 2

H) 1.34 (t, J=5.91 Hz, 3 H) 1.26 - 1.30 (m, 4 H) 1.24 (br d, J=3.32 Hz, 3 H) 1.07 (dd, J=6.74, 3.84 Hz, 6 H) 0.93 (dd, J=6.63, 3.32 Hz, 6 H). m/z (ESI, +ve ion): 632.2 (M+H)+.

Example 170

6-Chloro-1-[4,6-diisopropyl-2-[(4-methylpiperazin-1-yl)methyl]pyrimidin-5-yl]-4-[(2S,5R)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2-one



Intermediate 235



[0369] Step 1. *tert*-Butyl (2R,5S)-4-(6-chloro-1-(4,6-diisopropyl-2-((4-methylpiperazin-1-yl)methyl)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate.

[0370] To a 50-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(3-chloro-2-(2-fluorophenyl)-8-(2-formyl-4,6-diisopropylpyrimidin-5-yl)-7-oxo-7,8-dihydro-1,6-naphthyridin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 235**, 0.265 g, 0.391 mmol) and 1-methylpiperazine (0.1 mL, 0.587 mmol) in tetrahydrofuran (3.9 mL). Then glacial acetic acid (2.3 µl, 0.039 mmol), followed by sodium cyanoborohydride (0.074 g, 1.174

mmol) was added into the reaction mixture. The overall reaction mixture was allowed to stir at rt, while under an inert (N_2) atmosphere for 30 min. The reaction mixture was quenched with MeOH (1 mL) and allowed the mixture to stir 5 min. Then the mixture was diluted with EtOAc and sat. aq. $NaHCO_3$. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material (0.250 g) was carried into the next step of the synthesis, without further purification. *m/z* (ESI, +ve ion): 762.2 $(M+H)^+$.

[0371] Step 2. 4-((2S,5R)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-((4-methylpiperazin-1-yl)methyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(6-chloro-1-(4,6-diisopropyl-2-((4-methylpiperazin-1-yl)methyl)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.250 g, 0.328 mmol) and trifluoroacetic acid (0.2 mL, 3.28 mmol) in 1,2-dichloroethane (1.5 mL). The reaction mixture was stirred and heated at 60 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated in vacuo. This material was carried directly into the next step of the synthesis, without further purification.

[0372] The previous residue was diluted with 1,2-dichloroethane (1.5 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.6 mL, 3.94 mmol) was added into the reaction mixture and stirred 2 min. Then acryloyl chloride (0.2 mL, 0.328 mmol) was added into the mixture dropwise and stirred under an inert (N_2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (25 grams), eluting with a gradient of 0-5% MeOH/DCM. This afforded 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropyl-2-((4-methylpiperazin-1-yl)methyl)pyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.050 g, 0.070 mmol, 21.29 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.48 (d, *J*=4.77 Hz, 1 H) 7.48 - 7.55 (m, 1 H) 7.16 - 7.33 (m, 3 H) 6.84 (td, *J*=17.31, 10.78 Hz, 1 H) 6.19 (br d, *J*=16.59 Hz, 1 H) 5.75 (br d, *J*=10.16 Hz, 1 H) 4.88 (br s, 1 H) 4.78 (br s, 1 H) 4.43 - 4.54 (m, 1 H) 4.13 - 4.26 (m, 1 H) 3.79 - 3.93 (m, 2 H) 3.70 - 3.76 (m, 2 H) 3.39 - 3.55 (m, 1 H) 2.53 - 2.78 (m, 6 H) 2.28 - 2.37 (m, 3 H) 2.15 (s, 3 H) 1.34 (br t, *J*=5.49 Hz, 3

H) 1.26 (br d, *J*=6.43 Hz, 1 H) 1.19 (br d, *J*=6.63 Hz, 2 H) 1.02 - 1.10 (m, 6 H) 0.92 (br d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 716.3 $(M+H)^+$.

Example 171

6-Chloro-4-[(2S,5R)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-[2-(3-furyl)-4,6-diisopropyl-pyrimidin-5-yl]pyrido[2,3-d]pyrimidin-2-one



[0373] Step 1. *tert*-Butyl (2R,5S)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-(furan-3-yl)-4,6-diisopropylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (Intermediate 232, 0.080 g, 0.110 mmol) and sodium carbonate (0.035 g, 0.329 mmol) in 1,4-dioxane (2.4 mL). The reaction mixture was degassed by bubbling argon (gas) into the mixture for 5 min. Then xantphos pd g3 (5.20 mg, 5.49 μmol), water (0.6 mL) and 3-furanboronic acid (0.1 mL, 0.274 mmol) was added into the reaction mixture. The overall reaction mixture was stirred and

heated at 80 °C for 16h. The reaction mixture was cooled to rt, then the mixture was diluted with sat. aq. $NaHCO_3$ and EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (15 grams), eluting with a gradient of 0-25% EtOAc in CH2CL2, to provide tert-butyl (2R,5S)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-(furan-3-yl)-4,6-diisopropylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.070 g, 0.098 mmol, 89 % yield) as tan solid. *m/z* (ESI, +ve ion): 716.2 $(M+H)^+$.

[0374] Step 2. 4-((2S,5R)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-(furan-3-yl)-4,6-diisopropylpyrimidin-5-yl)pyrido[2,3-d]pyrimidin-2(1H)-one. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-(furan-3-yl)-4,6-diisopropylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.070 g, 0.098 mmol) and trifluoroacetic acid (0.1 mL, 0.977 mmol) in 1,2-dichloroethane (1.7 mL). The reaction mixture was stirred and heated at 60 °C for 20 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated in vacuo. This material was carried directly into the next step of the synthesis, without further purification.

[0375] The previous residue was diluted with DCM (4 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.2 mL, 1.173 mmol) was added into the reaction mixture and allowed the mixture to stir 2 min. Then acryloyl chloride (7.9 µl, 0.098 mmol) was added into the mixture dropwise over 2 min. The mixture was diluted with DCM and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (25 grams), eluting with a gradient of 0-65% EtOAc/DCM. This afforded 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-(furan-3-yl)-4,6-diisopropylpyrimidin-5-yl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.010 g, 0.015 mmol, 15.27 % yield) as white solid. ^{1}H NMR (500 MHz, DMSO-d_6) δ 8.49 (d, *J*=6.36 Hz, 1 H) 8.39 (d, *J*=0.78 Hz, 1 H) 7.79 (t, *J*=1.69 Hz, 1 H) 7.46 - 7.53 (m, 1 H) 7.19 - 7.33 (m, 3 H) 7.02 (d, *J*=1.17 Hz, 1 H) 6.79 - 6.90 (m, 1 H) 6.17 - 6.22 (m, 1 H) 5.73 - 5.78 (m, 1 H) 4.84 - 4.94 (m, 1 H) 4.14 - 4.26 (m, 1 H) 3.81 - 3.95 (m, 2 H) 2.65 - 2.77 (m, 2 H) 2.54 (s, 2 H) 1.30 - 1.40 (m,

3 H) 1.16 - 1.28 (m, 3 H) 1.12 (br d, *J*=4.02 Hz, 3 H) 1.11 (br d, *J*=4.15 Hz, 3 H) 0.94 - 1.01 (m, 6 H). *m/z* (ESI, +ve ion): 670.2 $(M+H)^+$.

Example 172

6-Chloro-1-(4,6-diisopropylpyrimidin-5-yl)-4-[(2S,5R)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-[2-(trifluoromethyl)phenyl]pyrido[2,3-d]pyrimidin-2-one



[0376] Step 1. 6-Chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(2-(trifluoromethyl)phenyl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione. To a 100-mL round-bottomed flask was added 6,7-dichloro-1-(4,6-diisopropylpyrimidin-5-yl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione (0.300 g, 0.761 mmol, **Intermediate 205**) and potassium acetate (0.224 g, 2.283 mmol) in 1,4-dioxane (3.8 mL). The reaction mixture was deoxygenated by bubbling (N_2) gas into the mixture 5 min. Then (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.056 g, 0.076 mmol) was added into the reaction mixture. The reaction mixture was stirred and heated at 80 °C for 10 min. Then a solution of (2-(trifluoromethyl)phenyl)boronic acid (0.217 g, 1.141 mmol) in 1,4-dioxane (1 mL) was added into the mixture, followed by water (0.1 mL). The resulting reaction mixture was heated and stirred at 95 °C for 16 h. The reaction mixture was allowed to cool to ambient temperature, then the reaction mixture was diluted with sat. aq. NH_4Cl and EtOAc. The aqueous layer was extracted with EtOAc and brine. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Redi-Sep pre-packed silica gel column (40 g),

eluting with a gradient of 0-30% EtOAc in CH2CL2, to provide 6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(2-(trifluoromethyl)phenyl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione (0.163 g, 0.323 mmol, 42.5 % yield) as white solid. *m/z* (ESI, +ve ion): 504.1 $(M+H)^+$.

[0377] Step 2. 4-((2S,5R)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(2-(trifluoromethyl)phenyl)pyrido[2,3-d]pyrimidin-2(1H)-one. To a 100-mL round-bottomed flask was added 6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(2-(trifluoromethyl)phenyl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione (0.150 g, 0.298 mmol) and n,n-diisopropylethylamine (0.1 mL, 0.387 mmol) in acetonitrile (1.4 mL). Then phosphorous oxychloride (0.1 mL, 0.357 mmol) was added into the reaction mixture, then the mixture was heated and stirred at 80 °C for 30 min, while under an inert (N_2) atmosphere. Another aliquot of n,n-diisopropylethylamine (0.1 mL, 0.387 mmol) and phosphorous oxychloride (0.1 mL, 0.357 mmol) was added into the mixture, then allowed the mixture to stir an additional 10 min. The reaction mixture was removed from the heat bath and allowed to cool to rt.

[0378] The reaction mixture was cooled to 0 °C with a wet ice/water bath. Then DIPEA (3 mL) was added dropwise into the reaction mixture. Then a solution of 1-((2R,5S)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one (0.233 g, 0.595 mmol, **Intermediate 160**) in MeCN (10 mL) was added dropwise into the reaction mixture. The ice bath was removed and the mixture was allowed to warm to rt over 20 min. More DIPEA (3 mL) and 1-((2R,5S)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one (0.233 g, 0.595 mmol, **Intermediate 160**) was added into the reaction mixture and stirred an additional 10 min. The reaction mixture was diluted with EtOAc and sat. aq. NH_4Cl. The aqueous layer was extracted with EtOAc. The combined organic extracts were dried over MgSO4, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (80 grams), eluting with a gradient of 0-80% EtOAc in CH2CL2, to provide 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(2-(trifluoromethyl)phenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.020 g, 0.031 mmol, 10.27 % yield) as white solid. ^{1}H NMR (500 MHz, DMSO-d_6) δ 9.02 (s, 1 H) 8.60 (br s, 1 H) 7.81 (d, *J*=7.78 Hz, 1 H) 7.77 (t, *J*=7.56 Hz, 1 H) 7.66 - 7.71 (m, 1 H) 7.35 (br s, 1 H) 6.82 (br dd, *J*=16.67, 10.45 Hz, 1 H) 6.19 (dd, *J*=16.67, 2.27 Hz, 1 H) 5.74 - 5.78 (m, 1 H) 4.64 - 5.02 (m, 2 H) 4.09 (br d, *J*=5.32 Hz, 1 H) 4.04 (br s, 1 H) 3.88 (br s, 2 H)

2.64 - 2.83 (m, 2 H) 1.30 - 1.38 (m, 3 H) 1.21 - 1.30 (m, 2 H) 1.19 (br s, 1 H) 1.07 (br s, 6 H) 0.92 (br s, 6 H). *m/z* (ESI, +ve ion): 654.2 $(M+H)^+$.

Example 173-1 (Peak 1)

6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(6-isopropenyl-2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one.



[0379] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-(prop-1-en-2-yl)pyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.420 g, 0.600 mmol, **Intermediate 202**) and sodium carbonate (0.191 g, 1.800 mmol) in 1,4-dioxane (2.4 mL). The reaction mixture was degassed by bubbling argon (gas) into the mixture for 5 min. Then xantphos pd g3 (0.028 g, 0.030 mmol), water (0.6 mL) and 2-isopropenyl boronic acid pinacol ester (0.2 mL, 1.500 mmol) was added into the reaction mixture. The overall reaction mixture was stirred and heated at 100 °C for 16 h. The reaction mixture was cooled

to rt, then the mixture was diluted with sat. aq. $NaHCO_3$ and EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (40 grams), eluting with a gradient of 0-20% EtOAc in CH2CL2, to provide tert-butyl (2R,5S)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-(prop-1-en-2-yl)pyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.200 g, 0.302 mmol, 50.4 % yield) as white solid. *m/z* (ESI, +ve ion): 661.3 $(M+H)^+$.

[0380] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-(prop-1-en-2-yl)pyridin-3-yl)pyrido[2,3-d]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-(prop-1-en-2-yl)pyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.200 g, 0.302 mmol) and trifluoroacetic acid (0.2 mL, 3.02 mmol) in 1,2-dichloroethane (3.0 mL). The reaction mixture was stirred and heated at 60 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated in vacuo. This material was carried directly into the next step of the synthesis, without further purification.

[0381] The previous residue was diluted with 1,2-dichloroethane (3.0 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.6 mL, 3.63 mmol) was added into the reaction mixture and stirred 2 min. Then acryloyl chloride (0.2 mL, 0.302 mmol) was added into the mixture dropwise and stirred under an inert (N_2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-6% MeOH/DCM. The racemic mixture was separated by SFC; (Column, IC, 5 μm, 21 x 250 mm, F=80 mL/min, 30% Isopropanol/70% Carbon dioxide). This afforded the separated Atropisomers, which were arbitrarily assigned as (Peak 1) 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-(prop-1-en-2-yl)pyridin-3-yl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.060 g, 0.098 mmol, 32.2 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.47 (d, *J*=3.52 Hz, 1 H) 7.46 - 7.56 (m, 1 H) 7.43 (s, 1 H) 7.22 - 7.35 (m, 3 H) 6.84 (td, *J*=16.74, 10.47 Hz, 1 H) 6.20 (dd, *J*=16.79, 1.87 Hz, 1 H) 5.97 (s, 1 H) 5.73 - 5.79 (m, 1 H) 5.30 (s, 1 H) 4.86

(br s, 1 H) 4.78 (br s, 1 H) 4.12 - 4.23 (m, 1 H) 3.80 - 4.00 (m, 2 H) 2.66 - 2.75 (m, 1 H) 2.15 (s, 3 H) 1.95 (s, 3 H) 1.32 - 1.39 (m, 3 H) 1.24 - 1.28 (m, 2 H) 1.17 - 1.21 (m, 2 H) 1.09 (d, *J*=6.84 Hz, 3 H) 0.99 (d, *J*=6.63 Hz, 3 H). *m/z* (ESI, +ve ion): 615.2 $(M+H)^+$.

Example 173-2 (peak 2)

[0382] The racemic mixture from example 173-1 was separated by SFC; (Column, IC, 5 µm, 21 x 250 mm, F=80 mL/min, 30% Isopropanol/70% Carbon dioxide). This afforded the separated Atropisomers, which were arbitrarily assigned as (Peak 2) 1H NMR (500 MHz, DMSO-*d*6) δ 8.48 (d, *J*=3.76 Hz, 1 H) 7.49 - 7.54 (m, 1 H) 7.44 (s, 1 H) 7.23 - 7.33 (m, 3 H) 6.83 (td, *J*=17.06, 10.51 Hz,1 H) 6.19 (br d, *J*=16.74 Hz, 1 H) 5.96 (s, 1 H) 5.73 - 5.78 (m, 1 H) 5.29 (s, 1 H) 4.83 - 4.91 (m, 1 H) 4.76 (br s, 1 H) 4.10 - 4.21 (m, 2 H) 3.80 - 3.97 (m, 2 H) 2.64 - 2.71 (m, 1 H) 2.14 (s, 3 H) 1.97 (s, 3 H) 1.30 - 1.37 (m, 3 H) 1.24 (d, *J*=6.49 Hz, 1 H) 1.16 (d, *J*=6.75 Hz, 2 H) 1.07 (d, *J*=6.75 Hz, 3 H) 0.95 (d, *J*=6.75 Hz, 3 H). *m/z* (ESI, +ve ion): 615.4 $(M+H)^+$.

Example 174

6-Chloro-1-(4,6-diisopropylpyrimidin-5-yl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(5-hydroxy-2-methyl-phenyl)pyrido[2,3-*d*]pyrimidin-2-one



[0383] Step 1. 6-Chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(5-methoxy-2-methylphenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a 100-mL round-bottomed flask was added 6,7-dichloro-1-(4,6-diisopropylpyrimidin-5-yl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione (0.400 g, 1.015 mmol, **Intermediate 205**) and potassium acetate (0.299 g, 3.04 mmol) in 1,4-dioxane (7.6 mL). The reaction mixture was deoxygenated by bubbling (N_2) gas into the mixture 5 min. Then PdCl2(dppf) (0.074 g, 0.101 mmol) was added into the reaction mixture. The mixture was stirred and heated at 105 °C for 10 min. Then a solution of (5-methoxy-2-methylphenyl)boronic acid (0.3 mL, 2.029 mmol) in 1,4-dioxane (1 mL) was added into the mixture, followed by water (1.9 mL). The resulting reaction mixture was heated and stirred at 105 °C for 3 h. The reaction mixture was allowed to cool to ambient temperature, then diluted the mixture with sat. aq. NH_4Cl and EtOAc. The aqueous layer was extracted with EtOAc and brine. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Redi-Sep pre-packed silica gel column (40 g), eluting with a gradient of 0-20% EtOAc in CH2CL2, to provide 6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(5-methoxy-2-methylphenyl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione (0.450 g, 0.938 mmol, 92 % yield) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.28 (s, 1 H) 9.07 (s, 1 H) 8.57 (s, 1 H) 7.16 (d, *J*=8.50 Hz, 1 H) 6.85 - 6.93 (m, 1 H) 6.51 (d, *J*=2.70 Hz, 1 H) 3.63 - 3.67 (m, 3 H) 2.98 (quin, *J*=6.63 Hz, 2 H) 1.85 (s, 3 H) 1.09 (d, *J*=6.63 Hz, 6 H) 0.95 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 480.1 $(M+H)^+$.

[0384] Step 2. 4-((2*S*,5*R*)-4-((11-Boraneyl)carbonyl)-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(5-methoxy-2-methylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added 6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(5-methoxy-2-methylphenyl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione (0.450 g, 0.938 mmol) and n,n-diisopropylethylamine (0.2 mL, 1.219 mmol) in acetonitrile (6.2 mL). Then phosphorous oxychloride (0.1 mL, 1.125 mmol) was added into the reaction mixture, then the mixture was heated and stirred at 80 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was removed from the heat bath and allowed to cool to rt.

[0385] The reaction mixture was cooled to 0 °C with a wet ice/water bath. Then DIPEA (2.5 mL) was added dropwise into the reaction mixture. Then a solution of (2r,5s)-1-boc-2,5-dimethylpiperazine (0.301 g, 1.406 mmol) in MeCN (4 mL) was added dropwise into the reaction mixture. The ice bath was removed and the mixture was allowed to warm to rt. The

reaction mixture was diluted with DCM and sat. aq. NH_4Cl. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (120 grams), eluting with a gradient of 0-10% MeOH in CH2CL2, to provide tert-butyl (2R,5S)-4-(6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(5-methoxy-2-methylphenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.600 g, 0.887 mmol, 95 % yield) as tan solid. *m/z* (ESI, +ve ion): 676.3 $(M+H)^+$.

[0386] Step 3. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(5-hydroxy-2-methylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(5-methoxy-2-methylphenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.600 g, 0.887 mmol) in 1,2-dichloroethane (17.7 mL). Then boron tribromide (1.0M in DCM) (4.4 mL, 4.44 mmol) was added dropwise into the reaction mixture, while the mixture was under an inert (N_2) atmosphere. The resulting reaction mixture was allowed to stir 30 min. The reaction mixture was quenched with the addition of MeOH (3 mL) and allowed the mixture to stir 10 min. Then the reaction mixture was diluted with DCM and sat. aq. $NaHCO_3$. The aqueous layer was extracted with DCM and the combined organic extracts were dried over MgSO4, filtered and concentrated in vacuo. This afforded tert-butyl (2R,5S)-4-(6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(5-hydroxy-2-methylphenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as tan solid. The crude mixture (0.499 grams) was carried into the next step of the synthesis, without further purification. *m/z* (ESI, +ve ion): 562.2 $(M-BOC+H)^+$.

[0387] Step 4. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(5-hydroxy-2-methylphenyl)pyrido[2,3-d]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added 6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-4-((2S,5R)-2,5-dimethylpiperazin-1-yl)-7-(5-hydroxy-2-methylphenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.499 g, 0.888 mmol) and n,n'-diisopropylethylamine (0.2 mL, 1.332 mmol) in 1,2-dichloroethane (4.4 mL). Then acryloyl chloride (1.1M in DCM) (0.8 mL, 0.888 mmol) was added slowly into the reaction mixture. The overall reaction mixture was allowed to stir under an inert (N_2) atmosphere 10 min. The reaction mixture was quenched with sat. aq. NH_4Cl, then diluted the mixture with DCM and

water. The layers were separated and the aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Redi-Sep pre-packed silica gel column (80 g), eluting with a gradient of 0-8% MeOH/DCM, to provide 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(4,6-diisopropylpyrimidin-5-yl)-7-(5-hydroxy-2-methylphenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.175 g, 0.284 mmol, 32.0 % yield) as off-white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 9.40 (s, 1 H) 9.04 (s, 1 H) 8.45 (d, *J*=3.32 Hz, 1 H) 7.02 (d, *J*=8.29 Hz, 1 H) 6.84 (td, *J*=16.64, 10.68 Hz, 1 H) 6.70 (dd, *J*=8.29, 2.49 Hz, 1 H) 6.40 (d, *J*=2.49 Hz, 1 H) 6.19 (dd, *J*=16.59, 2.07 Hz, 1 H) 5.75 (br dd, *J*=10.68, 3.42 Hz, 1 H) 4.87 (br s, 1 H) 4.13 - 4.25 (m, 1 H) 3.79 - 3.94 (m, 2 H) 3.16 - 3.18 (m, 1 H) 2.63 - 2.79 (m, 2 H) 1.82 (s, 3 H) 1.34 (br t, *J*=5.80 Hz, 3 H) 1.25 (br d, *J*=6.63 Hz, 2 H) 1.18 (br d, *J*=6.63 Hz, 2 H) 1.08 (dd, *J*=6.32, 4.66 Hz, 6 H) 0.95 (br t, *J*=5.70 Hz, 6 H). *m/z* (ESI, +ve ion): 616.2 $(M+H)^+$.

Example 175-1 (peak 1)

1-[6-(Azetidin-1-yl)-2-isopropyl-4-methyl-3-pyridyl]-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one



[0388] Step 1. 1-(6-(Azetidin-1-yl)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a 50-mL round-bottomed flask was added 1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione (0.350 g, 0.695 mmol, **Intermediate 236**) and azetidine (0.3 mL, 6.95 mmol) in 1,4-dioxane (2.8 mL). The reaction mixture was stirred and heated at 70°C while under an inert (N_2) atmosphere overnight. The mixture was allowed to cool to ambient temperature, then diluted the mixture with EtOAc and water. The aqueous layer was extracted with EtOAc (3x) and the combined organic extracts were dried over MgSO4, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Redi-Sep pre-packed silica gel column (40 g), eluting with a gradient of 0-10% EtOAc in CH2CL2, to provide 1-(6-(azetidin-1-yl)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidine-

2,4(1H,3H)-dione (0.240 g, 0.500 mmol, 72.0 % yield) as yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.12 (s, 1 H) 8.53 (s, 1 H) 7.43 - 7.56 (m, 1 H) 7.19 - 7.38 (m, 3 H) 6.10 (s, 1 H) 3.85 - 4.04 (m, 4 H) 2.68 - 2.74 (m, 1 H) 2.23 - 2.32 (m, 2 H) 1.88 (s, 3 H) 1.03 (d, *J*=6.84 Hz, 3 H) 0.88 (d, *J*=6.63 Hz, 3 H). *m/z* (ESI, +ve ion): 480.0 $(M+H)^+$.

[0389] Step 2. *tert*-Butyl (2*R*,5*S*)-4-(1-(6-(azetidin-1-yl)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate.

[0390] To a 100-mL round-bottomed flask was added 1-(6-(azetidin-1-yl)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione (0.480 g, 1.000 mmol) and n,n-diisopropylethylamine (0.2 ml, 1.300 mmol) in acetonitrile (6.6 ml). Then phosphorous oxychloride (0.1 ml, 1.200 mmol) was added into the reaction mixture, then the mixture was heated and stirred at 80 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was removed from the heat bath and allowed to cool to rt.

[0391] The reaction mixture was cooled to 0 °C with a wet ice/water bath. Then DIPEA (2.5 mL) was added dropwise into the reaction mixture. Then a solution of (2r,5s)-1-boc-2,5-dimethylpiperazine (0.322 g, 1.500 mmol) in MeCN (4 mL) was added dropwise into the reaction mixture. The ice bath was removed and the mixture was allowed to warm to rt. The reaction mixture was diluted with DCM and sat. aq. NH_4Cl. The aqueous layer was extracted with DCM. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (120 grams), eluting with a gradient of 0-30% EtOAc in CH_2CL_2, to provide tert-butyl (2R,5S)-4-(1-(6-(azetidin-1-yl)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.385 g, 0.569 mmol, 56.9 % yield) as yellow solid. *m/z* (ESI, +ve ion): 676.2 $(M+H)^+$.

[0392] Step 3. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(6-(azetidin-1-yl)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(1-(6-(azetidin-1-yl)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.381 g, 0.563 mmol) and trifluoroacetic acid (0.4 mL, 5.63 mmol) in 1,2-dichloroethane (3.0 mL). The reaction mixture was stirred and heated at 60 °C for 30 min, while under an inert (N_2)

atmosphere. The reaction mixture was concentrated in vacuo. This material was carried directly into the next step of the synthesis, without further purification.

[0393] The previous residue was diluted with 1,2-dichloroethane (3.0 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (1.1 mL, 6.76 mmol) was added into the reaction mixture and stirred 2 min. Then acryloyl chloride (0.5 mL, 0.563 mmol) was added into the mixture dropwise and stirred under an inert (N_2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-6% MeOH/DCM. The racemates were separated by SFC; (Column, ID, 5 μm, 21 x 250 mm, F= 60 mL/min, 40% EtOH/60%: Carbon dioxide). This afforded the separated Atropisomers as Peak 1: 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(6-(azetidin-1-yl)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.118 g, 0.187 mmol, 33.2 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.44 (s, 1 H) 7.43 - 7.60 (m, 1 H) 7.23 - 7.38 (m, 3 H) 6.82 (dt, *J*=16.53, 11.32 Hz, 1 H) 6.11 - 6.22 (m, 2 H) 5.75 (dt, *J*=12.75, 2.23 Hz, 1 H) 4.84 (br s, 1 H) 4.41 - 4.65 (m, 1 H) 4.11 (br t, *J*=13.16 Hz, 2 H) 3.90 - 3.94 (m, 4 H) 3.79 - 3.87 (m, 1 H) 2.28 (quin, *J*=7.31 Hz, 2 H) 1.77 - 1.86 (m, 3 H) 1.31 (br t, *J*=7.05 Hz, 3 H) 1.22 (br d, *J*=6.63 Hz, 2 H) 1.10 - 1.15 (m, 3 H) 1.00 (d, *J*=6.63 Hz, 3 H) 0.87 (d, *J*=6.63 Hz, 3 H). *m/z* (ESI, +ve ion): 630.2 $(M+H)^+$.

Example 175-2 (peak 2)

[0394] The racemates were separated by SFC; (Column, ID, 5 μm, 21 x 250 mm, F= 60 mL/min, 40% EtOH/60%: Carbon dioxide). This afforded the separated Atropisomers as Peak 2: 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(6-(azetidin-1-yl)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.120 g, 0.190 mmol, 33.8 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.42 (s, 1 H) 7.43 - 7.61 (m, 1 H) 7.20 - 7.39 (m, 3 H) 6.83 (td, *J*=16.27, 10.57 Hz, 1 H) 6.19 (br d, *J*=16.38 Hz, 1 H) 6.11 (s, 1 H) 5.74 (br d, *J*=9.95 Hz, 1 H) 4.82 (br s, 1 H) 4.54 - 4.67 (m, 1 H) 4.12 (br t, *J*=12.96 Hz, 2 H) 3.89 - 3.95 (m, 4 H) 3.85 (br s, 1 H) 2.28 (quin, *J*=7.20 Hz, 2 H) 1.79 (s, 3 H) 1.31 (br t, *J*=7.05 Hz, 3 H) 1.24 (br d, *J*=6.01 Hz, 2 H) 1.10 - 1.15 (m, 3 H) 0.99 - 1.03 (m, 3 H) 0.90 (br d, *J*=6.43 Hz, 3 H). *m/z* (ESI, +ve ion): 630.2 $(M+H)^+$.

Example 176-1 (peak 1)

6-Chloro-1-(2,6-diisopropyl-4-methyl-3-pyridyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2-one



[0395] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2,6-diisopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.400 g, 0.571 mmol, **Intermediate 202**) in tetrahydrofuran (2.8 mL). The reaction mixture was degassed by bubbling argon (gas) into the mixture for 5 min. Then 2-propylzinc bromide (1.0M in THF) (3.0 mL, 3.03 mmol) and xantphos pd g3 (0.016 g, 0.017 mmol) was added into the reaction mixture. The overall reaction mixture was allowed to stir under an inert (N_2) atmosphere at rt for 16 h. The reaction mixture was quenched with sat. aq. NH_4Cl and diluted the mixture with EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in

vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (25 grams), eluting with a gradient of 0-40% EtOAc in CH2CL2, to provide tert-butyl (2R,5S)-4-(6-chloro-1-(2,6-diisopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.253 g, 0.381 mmol, 66.8 % yield) as tan solid. *m/z* (ESI, +ve ion): 663.2 $(M+H)^+$.

[0396] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,6-diisopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(6-chloro-1-(2,6-diisopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.253 g, 0.381 mmol) and trifluoroacetic acid (0.2 mL, 3.81 mmol) in 1,2-dichloroethane (3.8 mL). The reaction mixture was stirred and heated at 60 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated in vacuo. This material was carried directly into the next step of the synthesis, without further purification.

[0397] The previous residue was diluted with 1,2-dichloroethane (3.8 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.8 mL, 4.58 mmol) was added into the reaction mixture and stirred 2 min. Then acryloyl chloride (0.3 mL, 0.381 mmol) was added into the mixture dropwise and stirred under an inert (N_2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-6% MeOH/DCM. The racemates were separated by SFC; (Column, Chiralpak AD-H, 5 μm, 2 x 25 cm, F= 80 mL/min, 25% Isopropanol/75%: Carbon dioxide). This afforded the separated Atropisomers as Peak 1: 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,6-diisopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.070 g, 0.113 mmol, 29.7 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 (d, *J*=2.90 Hz, 1 H) 7.48 - 7.55 (m, 1 H) 7.20 - 7.34 (m, 3 H) 7.05 (s, 1 H) 6.83 (td, *J*=16.74, 10.47 Hz, 1 H) 6.19 (dd, *J*=16.69, 2.18 Hz, 1 H) 5.72 - 5.78 (m, 1 H) 4.85 (br s, 1 H) 4.77 (br s, 1 H) 4.42 - 4.52 (m, 1 H) 4.09 - 4.19 (m, 1 H) 3.80 - 3.93 (m, 2 H) 2.96 (quin, *J*=6.89 Hz, 1 H) 2.61 - 2.69

(m, 1 H) 1.91 (s, 3 H) 1.33 (t, *J*=6.95 Hz, 3 H) 1.15 - 1.28 (m, 9 H) 1.06 (d, *J*=6.63 Hz, 3 H) 0.97 (d, *J*=6.63 Hz, 3 H). *m/z* (ESI, +ve ion): 617.2 $(M+H)^+$.

Example 176-2 (peak 2)

[0398] The racemates were separated by SFC; (Column, Chiralpak AD-H, 5 µm, 2 x 25 cm, F= 80 mL/min, 25% Isopropanol/75%: Carbon dioxide). This afforded the separated Atropisomers as Peak 2: 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,6-diisopropyl-4-methylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.054 g, 0.087 mmol, 22.94 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ ppm 8.42 - 8.50 (m, 1 H) 7.46 - 7.56 (m, 1 H) 7.20 - 7.35 (m, 3 H) 7.06 (s, 1 H) 6.77 - 6.89 (m, 1 H) 6.19 (dd, *J*=16.69, 2.18 Hz, 1 H) 5.71 - 5.79 (m, 1 H) 4.86 (br s, 1 H) 4.76 (br s, 1 H) 4.43 - 4.52 (m, 1 H) 4.08 - 4.21 (m, 1 H) 3.75 - 3.96 (m, 2 H) 2.96 (spt, *J*=6.91 Hz, 1 H) 2.61 - 2.68 (m, 1 H) 1.93 (s, 3 H) 1.33 (br t, *J*=6.22 Hz, 3 H) 1.24 (dd, *J*=6.84, 1.04 Hz, 7 H) 1.16 (br d, *J*=6.63 Hz, 2 H) 1.02 - 1.08 (m, 3 H) 0.94 (d, *J*=6.84 Hz, 3 H). *m/z* (ESI, +ve ion): 617.2 $(M+H)^+$.

Example 177-1 (peak 1)

6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-vinyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one



[0399] Step 1. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-vinylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 50-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.420 g, 0.600 mmol, **Intermediate 202**) and sodium carbonate (0.191 g, 1.800 mmol) in 1,4-dioxane (2.4 mL). The reaction mixture was degassed by bubbling argon (gas) into the mixture for 5 min. Then xantphos pd g3 (0.028 g, 0.030 mmol), water (0.6 mL) and vinylboronic acid pinacol ester (0.2 mL, 1.500 mmol) was added into the reaction mixture. The overall reaction mixture was stirred and heated at 100 °C for 16 h. The reaction mixture was cooled to rt, then the mixture was diluted with sat. aq. $NaHCO_3$ and EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over

$MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (25 grams), eluting with a gradient of 0-30% EtOAc in CH2CL2, to provide tert-butyl (2R,5S)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-vinylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.290 g, 0.448 mmol, 74.7 % yield) as tan solid. *m/z* (ESI, +ve ion): 647.2 $(M+H)^+$.

[0400] Step 2. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-vinylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a 100-mL round-bottomed flask was added tert-butyl (2R,5S)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-vinylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.290 g, 0.448 mmol) and trifluoroacetic acid (0.3 mL, 4.48 mmol) in 1,2-dichloroethane (2.0 mL). The reaction mixture was stirred and heated at 60 °C for 10 min, while under an inert (N_2) atmosphere. The reaction mixture was concentrated in vacuo. This material was carried directly into the next step of the synthesis, without further purification.

[0401] The previous residue was diluted with 1,2-dichloroethane (2.2 mL), then the reaction mixture was cooled to 0 °C with a wet ice bath. Then n,n'-diisopropylethylamine (0.9 mL, 5.38 mmol) was added into the reaction mixture and stirred 2 min. Then acryloyl chloride (0.4 mL, 0.448 mmol) was added into the mixture dropwise and stirred under an inert (N_2) atmosphere for 10 min. The mixture was diluted with EtOAc and sat. aq. $NaHCO_3$, then the layers were separated. The aqueous layer was extracted with EtOAc, then the combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (15 micron) silica-gel column (40 grams), eluting with a gradient of 0-6% MeOH/DCM. The racemates were separated by SFC; (Column, AD, 5 μm, 20 x 250 mm, F= 70 mL/min, 20% Isopropanol/80%: Carbon dioxide). This afforded the separated Atropisomers as Peak 1: 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-vinylpyridin-3-yl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.050 g, 0.083 mmol, 18.56 % yield) as white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.47 (d, *J*=3.73 Hz, 1 H) 7.49 - 7.55 (m, 1 H) 7.22 - 7.34 (m, 4 H) 6.73 - 6.90 (m, 2 H) 6.17 - 6.27 (m, 2 H) 5.73 - 5.79 (m, 1 H) 5.46 (d, *J*=11.40 Hz, 1 H) 4.87 (br s, 1 H) 4.78 (br s, 1 H) 4.13 - 4.22 (m, 1 H) 3.81 - 3.93 (m, 2 H) 2.66 - 2.75 (m, 1 H) 1.94 (s, 3 H) 1.34 (t, *J*=6.53 Hz, 3 H) 1.26 (br d, *J*=6.63 Hz, 2 H) 1.19 (br d,

J=6.63 Hz, 2 H) 1.08 (d, *J*=6.63 Hz, 3 H) 0.98 (d, *J*=6.63 Hz, 3 H). *m/z* (ESI, +ve ion): 601.2 $(M+H)^+$.

Example 177-2 (peak 2)

[0402] The racemates were separated by SFC; (Column, AD, 5 μm, 20 x 250 mm, F= 70 mL/min, 20% Isopropanol/80%: Carbon dioxide). This afforded the separated Atropisomers as Peak 2: 4-((2S,5R)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methyl-6-vinylpyridin-3-yl)pyrido[2,3-d]pyrimidin-2(1H)-one (0.060 g, 0.100 mmol, 22.28 % yield) as white solid. ^{1}H NMR (500 MHz, DMSO-d_6) δ ppm 8.48 (d, *J*=3.89 Hz, 1 H) 7.49 - 7.54 (m, 1 H) 7.23 - 7.33 (m, 4 H) 6.73 - 6.88 (m, 2 H) 6.21 (t, *J*=18.13 Hz, 2 H) 5.73 - 5.78 (m, 1 H) 5.44 - 5.48 (m, 1 H) 4.82 - 4.91 (m, 1 H) 4.76 (br s, 1 H) 4.11 - 4.21 (m, 1 H) 3.80 - 3.96 (m, 2 H) 3.41 - 3.58 (m, 1 H) 2.64 - 2.70 (m, 1 H) 1.95 (s, 3 H) 1.30 - 1.37 (m, 3 H) 1.24 (d, *J*=6.75 Hz, 1 H) 1.16 (d, *J*=6.75 Hz, 2 H) 1.06 (d, *J*=6.75 Hz, 3 H) 0.95 (d, *J*=6.75 Hz, 3 H). *m/z* (ESI, +ve ion): 601.2 $(M+H)^+$.

Example 178



[0403] 4-((2*S*,5*R*)-4-((*Z*)-4-(Dimethylamino)but-2-enoyl)-2,5-dimethylpiperazin-1-yl)-6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. 4-((2*S*,5*R*)-4-(4-(Dimethylamino)but-2-ynoyl)-2,5-dimethylpiperazin-1-yl)-6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 119-2, 64 mg, 0.10 mmol) in methanol (1.7 mL) was treated with Lindlar catalyst (5.6 mg, 0.021 mmol) and purged with hydrogen 4 x 30 psi and then allowed to stir under 30 psi hydrogen overnight. Added more Lindlar catalyst (5.6 mg, 0.021 mmol) and stirred under 35 psi for 24 h to get full conversion. The reaction mixture was filtered through a plug of celite, washed with MeOH, and concentrated to dryness on the rotavap. Purification of the crude by chromatography on silica gel using an ISCO Combiflash

RF (12 g RediSep Gold column, using a gradient of 0-100% 3:1 EtOAc/EtOH in heptane, then switch to 20% MeOH in DCM, afforded 4-((2*S*,5*R*)-4-((Z)-4-(dimethylamino)but-2-enoyl)-2,5-dimethylpiperazin-1-yl)-6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (5.5 mg, 8.9 μmol, 8.6 % yield) as off-white solid. *m/z* (ESI, +ve ion): 616.4 $(M+H)^+$. ^1H NMR (400 MHz, CHLOROFORM-d) δ 8.52 (d, *J*=4.77 Hz, 1H), 7.82 (dd, *J*=4.46, 9.02 Hz, 1H), 7.39-7.49 (m, 1H), 7.27-7.29 (m, 1H), 7.05-7.22 (m, 3H), 4.75-5.20 (m, 2H), 4.23-4.46 (m, 1H), 4.18 (br d, *J*=14.31 Hz, 1H), 4.06 (br d, *J*=14.10 Hz, 1H), 3.83-3.99 (m, 2H), 3.74 (s, 1H), 3.50 (d, *J*=3.11 Hz, 2H), 2.64-2.75 (m, 1H), 2.36-2.37 (d, *J*=4.0 Hz, 6H), 1.48 (d, *J*=6.63 Hz, 3H), 1.43 (dd, *J*=1.45, 6.84 Hz, 3H), 1.31 (d, *J*=6.84 Hz, 3H), 1.24 (d, *J*=6.84 Hz, 3H), 1.08 (dd, *J*=2.07, 6.63 Hz, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.45 (dd, *J*=19.07, 41.61 Hz, 1F), -126.70 (dd, *J*=28.61, 41.62 Hz, 1F).

Example 179

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethylpyridin-4-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one.



[0404] Steps 1&2. 6-Chloro-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethylpyridin-4-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A 50-mL round-bottomed flask was charged with 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (**Intermediate 99B**, 0.45 g, 1.58 mmol) and tetrahydrofuran (5 ml). Oxalyl dichloride (0.95 ml, 1.9 mmol) was added and the mixture was

stirred at 65 °C for 1 h. 5-Isopropyl-2,3-dimethylpyridin-4-amine (**Intermediate 189**, 0.28 g, 1.7 mmol) dissolved in 2 mL of THF was then added, follwed by *N,N*-diisopropylethylamine (0.69 ml, 3.9 mmol). The reaction mixture was stirred at rt for 30 mins and then heated at 65°C for 30 min. The reaction mixture was partitioned between water (30 mL) and EtOAc (50 mL). The aqueous phase was extracted with EtOAc (50 mL). The combined organic phases were dried and concentrated to give a tan solid.

[0405] To the residue 2,5-dichloro-6-(2-fluorophenyl)-*N*-((5-isopropyl-2,3-dimethylpyridin-4-yl)carbamoyl)nicotinamide (0.73 g, 1.54 mmol, 97% yield) dissloved in THF (10 mL) was added sodium *tert*-butoxide (0.30 g, 3.2 mmol). The resulting suspension was stirred for 5 h then diluted with water and extratcted with EtOAc (2x). The combined organic phases were dried (Na_2SO_4), concentrated and the residue purified by chromatography on silica gel eluting with EtOAc in heptane (30 % - 50 %) to afford 6-chloro-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethylpyridin-4-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.16 g, 0.36 mmol, 23 % yield) as white solid. *m/z* (ESI, +ve ion): 439.1 $(M+H)^+$.

[0406] **Steps 3&4. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethylpyridin-4-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**. Phosphoryl trichloride (40 ul, 0.43 mmol) was added dropwise to a solution of 6-chloro-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethylpyridin-4-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.16 g, 0.36 mmol) and N-ethyl-*N*-isopropylpropan-2-amine (0.087 ml, 0.50 mmol) in CH_3CN (1.2 mL) under N_2. This mixture was then heated to 80 °C for 1 h. Added more $POCl_3$ (10 uL, 0.1 mmol, 0.3 eq) and stirred at 80 °C for another hour. The reaction mixture was cooled to 0 °C and N-ethyl-*N*-isopropylpropan-2-amine (0.17 ml, 1.0 mmol) was added followed by *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (Astatech, 0.084 g, 0.39 mmol). This mixture was stirred at rt over 30 min. and then was poured into cold saturated $NaHCO_3$ solution and stirred vigorously for 10 min. The mixture was extracted with EtOAc, the combined organics were dried over Na_2SO_4, filtered, concentrated and purified by chromatography on silica gel using 0-40% 3:1 EtOAc/EtOH in heptane to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethylpyridin-4-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.125 g, 0.197 mmol, 55.0 % yield) as light-yellow solid. *m/z* (ESI, +ve ion): 635.2 $(M+H)^+$.

[0407] **Step 5&6. 6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethyl-4-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one.**

To *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethylpyridin-4-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (125 mg, 0.197 mmol) dissolved in dichloromethane (1 ml) was added trifluoroacetic acid (0.33 ml, 4.4 mmol). The reaction mixture was stirred at RT for 1 h then concentrated *in vacuo* and the residue was re-dissolved in dichloromethane (1 mL) and 1,1'-dimethyltriethylamine (0.10 ml, 0.59 mmol) was added followed by dropwise addition of acryloyl chloride (0.018 ml, 0.22 mmol) at 0°C. The reaction was stirred at 0°C for 30 min. The reaction mixture was purified by chromatography on silica gel using a RediSep Gold (24 g Gold) column, eluting with a gradient of 0% to 50% 3:1 EtOAc/EtOH in hetane, to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethylpyridin-4-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (54 mg, 0.046 mmol, 46 % yield) as white solid. *m/z* (ESI, +ve ion): 589.1 $(M+H)^+$.

Example 179-1

[0408] SFC conditions: Sample from above was purified via preparative SFC using an (R,R) Whelk-01 (250 x 21 mm, 5u) column, a mobile phase of 30% MeOH/CO_2 at 100 bar and at a flowrate of 80 g/min to generate 18 mg of peak 1 with an ee of >99% (chemcial purity 99.74%) and 9 mg of peak 2 with an ee of 99.34% (chemical purity 99.61%).

[0409] **6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethyl-4-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one (3380251, M isomer).** ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.42 (s, 1H), 8.10 (s, 1H), 7.36-7.49 (m, 1H), 7.07-7.19 (m, 3H), 6.51-6.73 (m, 1H), 6.40 (br t, *J*=14.82 Hz, 1H), 5.75-5.87 (m, 1H), 4.87-5.24 (m, 2H), 4.22-4.54 (m, 1H), 3.80-4.10 (m, 2H), 3.72 (br d, *J*=13.89 Hz, 1H), 3.45-3.77 (m, 1H), 3.43-3.56 (m, 1H), 2.52 (s, 3H), 1.89-1.99 (m, 3H), 1.40-1.48 (m, 3H), 1.32 (br d, *J*=6.84 Hz, 3H), 1.22 (d, *J*=6.84 Hz, 3H), 1.08 (d, *J*=6.84 Hz, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.70--112.47 (m, 1F). *m/z* (ESI, +ve ion): 589.1 $(M+H)^+$.

Example 179-2

[0410] **6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-1-(5-isopropyl-2,3-dimethyl-4-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one (3380252, P isomer).** ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.40 (s, 1H), 8.04-8.13 (m, 1H), 7.37-7.47 (m, 1H), 7.05-7.20 (m, 3H), 6.48-6.73 (m, 1H), 6.40 (br t, *J*=15.03 Hz, 1H),

5.73-5.88 (m, 1H), 4.84-5.26 (m, 2H), 4.21-4.56 (m, 1H), 3.40-4.11 (m, 3H), 2.53 (s, 3H), 1.88-1.98 (m, 3H), 1.38-1.50 (m, 3H), 1.30 (br d, *J*=6.84 Hz, 3H), 1.21 (d, *J*=6.84 Hz, 3H), 1.04 (d, *J*=6.84 Hz, 3H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.75--112.67 (m, 1F). *m/z* (ESI, +ve ion): 589.1 $(M+H)^+$.

Example 180



[0411] 4-(4-Acryloylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-hydroxy-2-methylpropyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of 4-(4-acryloylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-methoxy-2-methylpropyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (24 mg, 0.048 mmol) in 1,1-dichloroethane (240 µl) was added boron tribromide, 1.0m solution in methylene chloride (53 µl, 0.053 mmol) at 0°C. The reaction mixture was stirred at 0°C for 15 min (no change), then heated to 60°C and stirred for 5 h. The crude product was purified by ISCO chromatography through a Redi-Sep pre-packed silica gel column (column size: 24 g, flow rate: 20 mL/min, eluent: 0-80% 3:1 EtOAc/EtOH in heptane for 35 min) to give 4-(4-acryloylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-hydroxy-2-methylpropyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (5.8 mg, 0.012 mmol, 25% yield) as off-white solid. *m/z* (ESI, +ve ion): 486.2 $(M+H)^+$. ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.07 (s, 1H), 7.43-7.56 (m, 2H), 7.26-7.35 (m, 2H), 6.57 (dd, *J*=10.57, 16.59 Hz, 1H), 6.36 (dd, *J*=1.66, 16.79 Hz, 1H), 5.77 (dd, *J*=1.87, 10.57 Hz, 1H), 4.51 (s, 2H), 3.92-3.98 (m, 4H), 3.59-3.70 (m, 1H), 3.02-3.12 (m, 1H), 1.40 (d, *J*=6.84 Hz, 6H), 1.24 (s, 6H). ^{19}F NMR (376 MHz, CHLOROFORM-d) δ -112.83 (s, 1F).

Example 181

1-(4-(3-Chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-7-(morpholinomethyl)-1,6-naphthyridin-5-yl)piperazin-1-yl)prop-2-en-1-one.



[0412] Step 1. Methyl 5-(4-(*tert*-butoxycarbonyl)piperazin-1-yl)-3-chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridine-8-carboxylate. A mixture of methyl 3,5-dichloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridine-8-carboxylate (1.50 g, 3.80 mmol, Intermediate 85A), *tert*-butyl piperazine-1-carboxylate (2.83 g, 15.18 mmol, Aldrich), potassium carbonate (10.49 g, 76 mmol, Aldrich) and sodium sulfate (10.78 g, 76 mmol) in CH_3CN (30 mL) was heated at 85C for 2 h. The reaction went to completion, brought to rt, washed with water and extracted with EtOAc. The combined organics were dried over Na_2SO_4 , filtered, concentrated and purified by chromatography on silica gel using 0-50% EtOAc in heptane to afford methyl 5-(4-(*tert*-butoxycarbonyl)piperazin-1-yl)-3-chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridine-8-carboxylate (1.09 g, 2.000 mmol, 52.7 % yield) as a yellow solid. *m/z* (ESI, +ve ion): 545.2 $(M+H)^+$.

[0413] Step 2. *tert*-Butyl 4-(8-bromo-3-chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridin-5-yl)piperazine-1-carboxylate. To a solution of methyl 5-(4-(*tert*-butoxycarbonyl)piperazin-1-yl)-3-chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridine-8-carboxylate (1.09 g, 2.000 mmol) in EtOH (15 mL) was added KOH (2.469 g, 44.0 mmol, Aldrich) and the resulting mixture was heated at 85 C for 20 min. The reaction went to completion and concentrated to afford 5-(4-(*tert*-butoxycarbonyl)piperazin-1-yl)-3-

chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridine-8-carboxylic acid to be used as is. *m/z* (ESI, +ve ion): 531.2 $(M+H)^+$.

[0414] To 5-(4-(*tert*-butoxycarbonyl)piperazin-1-yl)-3-chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridine-8-carboxylic acid was added CH_3CH (30 mL) and water (15 mL) followed by the addition of LiOH (6.56 g, 274 mmol, Aldrich) and NBS (7.12 g, 40.0 mmol, Aldrich) at rt and the resulting mixture was stirred for 7 min. The reaction went to completion, washed with sat. $NaHCO_3$ and extracted with DCM. The combined organics were dried over Na_2SO_4, filtered, concentrated and purified by chromatography on silica gel using 0-50% EtOAc in heptane to afford *tert*-butyl 4-(8-bromo-3-chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridin-5-yl)piperazine-1-carboxylate (1.0 g, 1.767 mmol, 88 % yield) as a yellow solid. *m/z* (ESI, +ve ion): 567.2 $(M+H)^+$.

[0415] Step 3. *tert*-Butyl 4-(3-chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-7-(methoxymethyl)-1,6-naphthyridin-5-yl)piperazine-1-carboxylate. A mixture of *tert*-butyl 4-(8-bromo-3-chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridin-5-yl)piperazine-1-carboxylate (1.0 g, 1.767 mmol), cesium carbonate (2.303 g, 7.07 mmol, Aldrich), (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.078 ml, 0.106 mmol, Aldrich) and [2-(1-methylethyl)phenyl]-boronic acid (0.870 ml, 5.30 mmol, Combi-Blocks) was purged with N_2 followed by the addition of 1,4-dioxane/water (4/0.4 mL) and the resulting mixture was heated at 85 C for 1 h. The reaction went to completion, brought to rt, washed with sat. $NaHCO_3$ and extracted with EtOAC. The combined organics were purified on silica gel using 0-50% EtOAc in heptane to afford *tert*-butyl 4-(3-chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-7-(methoxymethyl)-1,6-naphthyridin-5-yl)piperazine-1-carboxylate (0.701 g, 1.158 mmol, 65.5 % yield) as a yellow-greenish oil. *m/z* (ESI, +ve ion): 604.6 $(M+H)^+$.

[0416] Step 4. (3-Chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-5-(piperazin-1-yl)-1,6-naphthyridin-7-yl)methanol. To a solution of *tert*-butyl 4-(3-chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-7-(methoxymethyl)-1,6-naphthyridin-5-yl)piperazine-1-carboxylate (0.701 g, 1.158 mmol) in DCM (20 mL) at 0C was added bbr3 (5.79 ml, 5.79 mmol, Aldrich) dropwise. After complete addition, the ice bath was removed and the resulting suspension was stirred at rt for 2 h. The reaction was brought to 0C, carefully basified with sat. $NaHCO_3$ and extracted with DCM. The combined organics were purified on silica gel using 0-10% MeOH in DCM to afford (3-chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-5-(piperazin-1-yl)-1,6-naphthyridin-7-yl)methanol (0.23 g, 0.468 mmol, 40.4 % yield) as a yellow solid. *m/z* (ESI, +ve ion): 490.6 $(M+H)^+$.

[0417] Step 5. 3-Chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-5-(piperazin-1-yl)-1,6-naphthyridine-7-carbaldehyde. To a suspension of (3-chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-5-(piperazin-1-yl)-1,6-naphthyridin-7-yl)methanol (0.23 g, 0.468 mmol, 40.4 % yield) in DCM (20 mL) was added Dess-martinperiodinane (0.737 g, 1.738 mmol, Aldrich) and the resulting mixture was stirred at rt for 2 h. The reaction went to completion, washed with sat. $NaHCO_3$, extracted with DCM and concentrated to afford 3-chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-5-(piperazin-1-yl)-1,6-naphthyridine-7-carbaldehyde as a yellow solid to be used as is. *m/z* (ESI, +ve ion): 488.6 $(M+H)^+$.

[0418] Step 6. 1-(4-(3-Chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-7-(morpholinomethyl)-1,6-naphthyridin-5-yl)piperazin-1-yl)prop-2-en-1-one. To a solution of 3-chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-5-(piperazin-1-yl)-1,6-naphthyridine-7-carbaldehyde (0.22 g, 0.450 mmol) in chloroform (15 mL) were added morpholine (0.078 ml, 0.900 mmol, Aldrich) and acetic acid (0.026 ml, 0.450 mmol, Aldrich) and the resulting mixture was stirred at rt for 10 min then sodium triacetoxyborohydride (0.381 g, 1.800 mmol, Aldrich) was added and the stirring continued for 1 h more. The reaction went to completion, brought to 0C and carefully basified with sat. $NaHCO_3$. The mixture was extracted with DCM, dried over Na_2SO_4, filtered and concentrated and purified by chromatography to afford 4-((3-chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-5-(piperazin-1-yl)-1,6-naphthyridin-7-yl)methyl)morpholine (0.050 g, 0.089 mmol, 19.84 % yield) as a light solid. *m/z* (ESI, +ve ion): 560 $(M+H)^+$.

[0419] 4-((3-chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-5-(piperazin-1-yl)-1,6-naphthyridin-7-yl)methyl)morpholine (0.050 g, 0.089 mmol, 19.84 % yield) was dissolved in DCM (5 mL) followed by the addition of acryloyl chloride (8.94 µl, 0.110 mmol, Aldrich) and the resulting mixture was stirred at rt for 30 min. The reaction went to completion, washed with sat. $NaHCO_3$, extracted with DCM and purified by chromatography on silica gel using 0-10% MeOH in DCM to afford 1-(4-(3-chloro-2-(2-fluorophenyl)-8-(2-isopropylphenyl)-7-(morpholinomethyl)-1,6-naphthyridin-5-yl)piperazin-1-yl)prop-2-en-1-one (0.039 g, 0.064 mmol, 14.11 % yield) as a yellow solid. *m/z* (ESI, +ve ion): 613.6 $(M+H)^+$. ^1H NMR (CHLOROFORM-d) δ: 8.38-8.45 (m, 1H), 7.31-7.47 (m, 4H), 7.01-7.19 (m, 4H), 6.62-6.76 (m, 1H), 6.33-6.44 (m, 1H), 5.72-5.83 (m, 1H), 3.84-4.08 (m, 7H), 3.62-3.74 (m, 4H), 3.50-3.60 (m, 3H), 3.38-3.47 (m, 1H), 2.38-2.66 (m, 4H), 1.04-1.12 (m, 3H), 0.93-0.99 (m, 3H).

Example 182

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(6-(azetidin-1-yl)-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one.



[0420] Step 1. *tert*-Butyl (2*R*,5*S*)-4-((((6-bromo-2,4-diisopropylpyridin-3-yl)carbamoyl)imino)(2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)methyl)-2,5-dimethylpiperazine-1-carboxylate. To a solution of 6-bromo-2,4-diisopropylpyridin-3-amine (0.217 g, 0.844 mmol, **Intermediate 196**) and DIEA (0.309 ml, 1.772 mmol, Aldrich) in THF (10 mL) at rt was added dropwise phosgene solution, 15% in toluene (0.662 ml, 0.928 mmol, Aldrich) and the resulting mixture was stirred at rt for 10 min then the mixture was brought to 0C and solid *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (0.487 g, 1.013 mmol, **Intermediate 165**) was added in three portions. The ice-bath was removed and after stirring for 15 min at rt *tert*-butyl (2*R*,5*S*)-4-((((6-bromo-2,4-diisopropylpyridin-3-yl)carbamoyl)imino)(2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)methyl)-2,5-dimethylpiperazine-1-carboxylate was observed. *m/z* (ESI, +ve ion): 764.4 $(M+H)^+$.

[0421] Step 2. *tert*-Butyl (2*R*,5*S*)-4-(1-(6-bromo-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. Then sodium *tert*-butoxide (0.243 g, 2.53 mmol, Aldrich) was added and the resulting mixture was stirred at rt for 20 min. The reaction went to completion, washed with water, extracted with EtOAc and purified by chromatography on

silica gel using 0-40% EtOAc in heptane to afford *tert*-butyl (2*R*,5*S*)-4-(1-(6-bromo-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.213 g, 0.293 mmol, 34.7 % yield) as a light yellow solid. *m/z* (ESI, +ve ion): 728.4 $(M+H)^+$.

[0422] Step 3. *tert*-Butyl (2*R*,5*S*)-4-(1-(6-(azetidin-1-yl)-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of *tert*-butyl (2*R*,5*S*)-4-(1-(6-bromo-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.085 g, 0.117 mmol), 1,1'-bis(diphenylphosphino)ferrocene (5.18 mg, 9.34 μmol, Strem), palladium (ii) acetate (1.048 mg, 4.67 μmol, Strem) and sodium *tert*-butoxide (0.017 g, 0.175 mmol, Aldrich) was purged with N_2 followed by the addition of azetidine (7.33 μl, 0.128 mmol, Combi-Blocks) and toluene (4 mL). The resulting mixture was heated at 90C for 17 h. The starting material was consumed and desired mass was observed with other peaks. The reaction was brought to rt, quenched with sat. $NaHCO_3$ and extracted with EtOAc. The combined organics were purified by HPLC to afford *tert*-butyl (2*R*,5*S*)-4-(1-(6-(azetidin-1-yl)-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.035 g, 0.050 mmol, 42.6 % yield) as a light yellow solid. *m/z* (ESI, +ve ion): 703.4 $(M+H)^+$.

[0423] Step 4. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(6-(azetidin-1-yl)-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of *tert*-butyl (2*R*,5*S*)-4-(1-(6-(azetidin-1-yl)-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.035 g, 0.050 mmol) in DCM (3 mL) was added tfa (2.0 mL, 26.0 mmol) and the resulting mixture was stirred at rt for 30 min. The reaction went to completion, concentrated and used as is in the next step. 1-(6-(azetidin-1-yl)-2,4-diisopropylpyridin-3-yl)-6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. *m/z* (ESI, +ve ion): 603.4 $(M+H)^+$.

[0424] 1-(6-(azetidin-1-yl)-2,4-diisopropylpyridin-3-yl)-6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one was dissolved in DCM (3 mL) then acryloyl chloride (4.04 μl, 0.050 mmol, Aldrich) was added at rt. The reaction was stirred at rt for 15 min, washed with sat. $NaHCO_3$ and extracted with DCM. The combined organics were purified by chromatography on silica gel using 0-5% MeOH in DCM

to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(6-(azetidin-1-yl)-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a light yellow solid. *m/z* (ESI, +ve ion): 657.4 $(M+H)^+$. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 7.94 - 8.04 (m, 1 H), 7.69 - 7.78 (m, 1 H), 7.38 - 7.48 (m, 1 H), 7.20 - 7.28 (m, 2 H), 7.11 - 7.19 (m, 1 H), 6.97 - 7.11 (m, 1 H), 6.16 (br d, *J*=16.8 Hz, 1 H), 5.71 - 5.75 (m, 1 H), 4.64 - 5.00 (m, 1 H), 4.17 - 4.38 (m, 1 H), 3.79 - 3.93 (m, 1 H), 3.57 - 3.75 (m, 2 H), 3.40 - 3.55 (m, 1 H), 3.14 - 3.28 (m, 2 H), 2.96 - 3.15 (m, 2 H), 1.24 (br dd, *J*=10.9, 6.7 Hz, 6 H), 1.16 (br s, 5 H), 1.02 - 1.11 (m, 4 H), 0.89 - 1.00 (m, 7 H).

[0425] The racemic mixture was separeted under the following conditions: SFC. IE (5um, 21x250 mm two in series, total 50cm) F=65ml, 35%; ethanol, 65% carbon dioxide , Back pressure= 90 bar.

Example 182-1

[0426] 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(6-(azetidin-1-yl)-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one

m/z (ESI, +ve ion): 657.6 $(M+H)^+$

^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.44 (s, 1 H), 7.48 - 7.58 (m, 1 H), 7.26 - 7.38 (m, 2 H), 7.16 - 7.24 (m, 1 H), 6.75 - 6.90 (m, 1 H), 6.19 (dd, *J*=16.6, 2.3 Hz, 1 H), 6.10 (s, 1 H), 5.71 - 5.80 (m, 1 H), 4.72 - 4.94 (m, 2 H), 4.50 (br d, *J*=2.7 Hz, 1 H), 4.08 - 4.19 (m, 1 H), 3.89 - 3.99 (m, 4 H), 3.83 - 3.88 (m, 1 H), 2.26 - 2.33 (m, 2 H), 1.29 - 1.36 (m, 3 H), 1.22 - 1.27 (m, 4 H), 1.00 - 1.07 (m, 7 H), 0.88 (dd, *J*=6.4, 4.8 Hz, 7 H).

Example 182-2

[0427] 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(6-(azetidin-1-yl)-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one

m/z (ESI, +ve ion): 657.6 $(M+H)^+$

^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.45 (s, 1 H), 7.49 - 7.57 (m, 1 H), 7.28 - 7.38 (m, 2 H), 7.17 - 7.25 (m, 1 H), 6.76 - 6.92 (m, 1 H), 6.19 (dd, *J*=16.7, 2.2 Hz, 1 H), 6.11 (s, 1 H), 5.72 - 5.79 (m, 1 H), 4.70 - 4.91 (m, 2 H), 4.41 - 4.51 (m, 1 H), 4.08 - 4.18 (m, 1 H), 3.89 - 3.99 (m, 4 H), 3.82 - 3.88 (m, 1 H), 2.24 - 2.32 (m, 2 H), 1.28 - 1.35 (m, 3 H), 1.25 (s, 4 H), 0.99 - 1.07 (m, 7 H), 0.91 (d, *J*=6.8 Hz, 3 H), 0.86 (br d, *J*=6.6 Hz, 4 H).

Example 183-1

[0428] (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (first eluting compound)

Example 183-2

[0429] (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (second eluting compound)



[0430] Step 1: *N*-((6-(Bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)carbamoyl)-2,5,6-trichloronicotinamide. To a solution of 2,5,6-trichloronicotinamide (187 mg, 0.831 mmol, intermediate P) in 1,2-dichloroethane (10 mL) under N_2 was added oxalyl chloride, 2.0 M solution in methylene chloride (0.457 mL, 0.914 mmol). The resulting mixture was then stirred at 80 °C for 1 hour. Then, the mixture was cooled to room temperature and a solution of 6-isopropyl-N2,N2-bis(4-methoxybenzyl)-4-methylpyridine-2,5-diamine (337 mg, 0.831 mmol, **intermediate I-33**) in MeCN (5 mL) was added. The resulting mixture was then stirred at room temperature for 1 hour. Then, the mixture was quenched with saturated NH_4Cl (5 mL) and was then diluted with saturated $NaHCO_3$ (25 mL). The mixture was then extracted with EtOAc (2 x 100 mL). The combined organic extracts were then dried over $MgSO_4$ and concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc:EtOH (3:1)/heptane) provided N-((6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)carbamoyl)-2,5,6-

trichloronicotinamide as a yellow solid which was used in the next step without purification requirement. m/z (ESI, +ve ion) 656.0 (M+H).

[0431] Step 2: 1-(6-(Bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a solution of N-((6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)carbamoyl)-2,5,6-trichloronicotinamide (546 mg, 0.831 mmol) in tetrahydrofuran (15 mL) at 0 °C under N_2 was added potassium bis(trimethylsilyl)amide, 1M solution in tetrahydrofuran (1.66 mL, 1.66 mmol) dropwise. After addition, the mixture was then stirred at 0 °C for 30 mins. Then, the mixture was quenched with saturated NH_4Cl (5 mL) and was then diluted with saturated $NaHCO_3$ (16 mL). The mixture was then extracted with EtOAc (2 x 100 mL). The combined organic extracts were then dried over $MgSO_4$ and concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc:EtOH (3:1)/heptane) provided 1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (476 mg, 0.767 mmol, 92 % yield) as a yellow solid. m/z (ESI, +ve ion) 620.0 (M+H)

[0432] Step 3: *tert*-Butyl (*S*)-4-(1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate. To a solution of 1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (471 mg, 0.759 mmol) in acetonitrile (8 mL) was added 1,1'-dimethyltriethylamine (0.464 mL, 2.66 mmol) and phosphorus oxide chloride (0.177 mL, 1.90 mmol). The resulting mixture was then stirred at 70°C under N_2 for 30 mins. Then, the mixture was cooled to room temperature and a solution of (3s)-1-(*tert*-butoxycarbonyl)-3-methylpiperazine (0.304 mL, 1.52 mmol, Ark Pharm, Inc., Arlington Heights, IL, USA) and 1,1'-dimethyltriethylamine (0.464 mL, 2.66 mmol) in MeCN (4 mL) was added. The resulting mixture was then stirred at room temperature under N_2 for 30 mins. The mixture was then diluted with water (20 mL) and was then extracted with EtOAc (2 x 50 mL). The combined organic extracts were then dried over $MgSO_4$ and concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-80% EtOAc:EtOH (3:1)/heptane) provided *tert*-butyl (*S*)-4-(1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate as a yellow solid which was used in the next step without purification requirement. m/z (ESI, +ve ion) 746.0 (M+H).

[0433] Step 4: *tert*-Butyl (*S*)-4-(1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate. To a solution of *tert*-butyl (*S*)-4-(1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (609 mg, 0.759 mmol) in 1,4-dioxane (7 mL) was added (2-fluorophenyl)boranediol (0.212 mL, 1.52 mmol, Combi-Blocks Inc., San Diego, CA, USA), (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.056 mL, 0.076 mmol), and potassium acetate (0.237 mL, 3.79 mmol). The resulting mixture was then stirred at 80 °C under N_2 for 1 hour. Then, the mixture was diluted with saturated $NaHCO_3$ (10 mL) and was then extracted with EtOAc (2 x 50 mL). The combined organic extracts were then dried over $MgSO_4$ and concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc:EtOH (3:1)/heptane) provided *tert*-butyl (*S*)-4-(1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (654 mg, 0.758 mmol, 100 % yield) as a yellow solid. m/z (ESI, +ve ion) 862.2 (M+H).

[0434] Step 5: (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one

[0435] Step 5-1: (*S*)-1-(6-(Bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-4-(2-methylpiperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one trifluoracetic acid. To a solution of *tert*-butyl (*S*)-4-(1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (654 mg, 0.758 mmol) in dichloromethane (5 mL) was added trifluoroacetic acid (1.130 mL, 15.17 mmol). The resulting mixture was then stirred at room temperature for 1 hour. Then, the mixture was concentrated and dried *in vacuo* provided (*S*)-1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-4-(2-methylpiperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one trifluoroacetic acid which was used in the next step without purification requirement. m/z (ESI, +ve ion) 762.2 (M+H).

[0436] Step 5-2: (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of (*S*)-1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-4-(2-

methylpiperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one trifluoroacetic acid in dichloromethane (7 mL) at 0 °C under N_2 was added potassium carbonate (0.230 mL, 3.79 mmol), 1,1'-dimethyltriethylamine (1.986 mL, 11.37 mmol), and a solution of 2-propenoyl chloride (0.062 mL, 0.758 mmol) in DCM (0.3 mL). The resulting mixture was then stirred at 0 °C for 15 mins. The mixture was then quenched with saturated $NaHCO_3$ (10 mL) and was then extracted with EtOAc (2 x 30 mL). The combined organic extracts were then dried over $MgSO_4$ and concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc:EtOH (3:1)/heptane) provided (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one which was used in the next step without purification requirement. m/z (ESI, +ve ion) 816.5 (M+H).

[0437] Step 5-3: (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-1-(6-(bis(4-methoxybenzyl)amino)-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (568 mg, 0.696 mmol) in trifluoroacetic acid (3.6 mL, 48.7 mmol) was subjected to a microwave irradiation at 100 °C for 10 mins. Then, the mixture was concentrated under reduced pressure. Then, the residue was dissolved in DCM (20 mL) and saturated $NaHCO_3$ (50 mL) was added slowly at 0 °C. After addition, the mixture was then stirred at room temperature for 5 mins. The organic layer was collected, dried over $MgSO_4$, and concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-10% NH_3 2M in MeOH/DCM) provided (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (165 mg, 0.143 mmol, 41.2 % yield) as a yellow solid. m/z (ESI, +ve ion) 576.0 (M+H). ^{1}H NMR (DMSO-d6) δ: 8.40 (br s, 1H), 7.48-7.60 (m, 1H), 7.22-7.36 (m, 3H), 6.78-6.93 (m, 1H), 6.15-6.25 (m, 2H), 5.70-5.84 (m, 3H), 4.79-5.03 (m, 1H), 4.22-4.49 (m, 2H), 3.96-4.19 (m, 1H), 3.40-3.83 (m, 2H), 3.01-3.29 (m, 2H), 1.73 (s, 3H), 1.29-1.35 (m, 3H), 0.99 (d, J=6.8 Hz, 3H), 0.86-0.89 (m, 3H).

[0438] Step 6: (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (first eluting atropisomer, 3371969) and (S)-4-(4-acryloyl-2-methylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (second eluting atropisomer). An atropisomeric mixture of (*S*)-4-(4-acryloyl-2-

methylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (155 mg) was separated by preparative Thar 200 SFC method, column: AS (150 X 30 mm, 5u), mobile phase: 80:20 (A:B) with A: Liquid CO2 and B: methanol(NH3), flow rate: 180 g/min; column/oven temp.: ambient temperature provided (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (61 mgs, first eluting atropisomer, 3371969) as a yellow solid. m/z (ESI, +ve ion) 576.0 (M+H). ^{1}H NMR (DMSO-d6) δ: 8.41 (br d, J=3.7 Hz, 1H), 7.48-7.58 (m, 1H), 7.21-7.36 (m, 3H), 6.78-6.94 (m, 1H), 6.16-6.25 (m, 2H), 5.76-5.80 (m, 2H), 5.73-5.76 (m, 1H), 4.83-5.00 (m, 1H), 4.21-4.49 (m, 2H), 3.97-4.19 (m, 1H), 3.38-3.83 (m, 2H), 3.00-3.30 (m, 2H), 1.72 (s, 3H), 1.31 (d, J=6.6 Hz, 3H), 0.99 (d, J=6.6 Hz, 3H), 0.87 (d, J=6.6 Hz, 3H). ^{19}F NMR (DMSO-d6) δ: -114.13 (s, 1F) and (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-1-(6-amino-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (63 mgs, second eluting atropisomer, 3371970) as a yellow solid. m/z (ESI, +ve ion) 576.0 (M+H). ^{1}H NMR (DMSO-d6) δ: 8.39 (br d, J=4.8 Hz, 1H), 7.48-7.58 (m, 1H), 7.22-7.36 (m, 3H), 6.77-6.93 (m, 1H), 6.15-6.26 (m, 2H), 5.76-5.79 (m, 2H), 5.74-5.76 (m, 1H), 4.89 (br s, 1H), 4.21-4.45 (m, 2H), 3.96-4.19 (m, 1H), 3.40-3.81 (m, 2H), 2.99-3.30 (m, 2H), 1.73 (s, 3H), 1.32 (d, J=6.6 Hz, 3H), 0.99 (d, J=6.6 Hz, 3H), 0.87 (d, J=6.6 Hz, 3H). ^{19}F NMR (DMSO-d6) δ: -114.09 (s, 1F)

Example 184

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-amino-6-fluorophenyl)-1-(4-((dimethylamino)methyl)-2-isopropyl-6-methylphenyl)-6-fluoropyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0439] Step 1: *N*-((4-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)carbamoyl)-2,6-dichloro-5-fluoronicotinamide. To a solution of 2,6-dichloro-5-fluoronicotinamide (998 mg, 4.78 mmol, **intermediate S**) in tetrahydrofuran (15 mL) was added oxalyl chloride, 2.0M in DCM (2.45 mL, 4.89 mmol) dropwise. The resulting mixture was then stirred at 65 °C under N_2 for 3 hours. Then, the mixture was cooled to room temperature and was then concentrated under reduced pressure to afford a light-yellow syrup which was redissolved in tetrahydrofuran (15 mL). Then, 4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylaniline (2.12 g, 5.08 mmol, intermediate I-37) in tetrahydrofuran (15 mL) was added to the solution. The resulting mixture was then stirred at room temperature for 30 min. Then, the mixture was concentrated under reduced pressure provided N-((4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)carbamoyl)-2,6-dichloro-5-fluoronicotinamide which was used in the next step without purification requirement. m/z (ESI, +ve ion) 652.2 (M+H).

[0440] Step 2: 1-(4-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-7-chloro-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a stirred solution of crude N-((4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)carbamoyl)-2,6-dichloro-5-fluoronicotinamide (2.9 g, 4.44 mmol) in tetrahydrofuran (16.0mL) at 0 °C was added potassium bis(trimethylsilyl)amide (8.89 mL, 8.89 mmol). After the mixture was stirred at room temperature for 2 hours, an additional potassium bis(trimethylsilyl)amide (1.0 mL, 1.0 mmol) was added. After 30 mins, an additional potassium bis(trimethylsilyl)amide (1.0 mL, 1.0mmol) was added. The resulting mixture was then stirred at room temperature for 15 min. The reaction was quenched by saturated NH_4Cl (75 mL) and diluted with EtOAc (50 mL). The mixture was then extracted with EtOAc (1 x 30 mL). The organic layer was collected, dried over $MgSO_4$, and concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-40% EtOAc/heptane) provided 1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-7-chloro-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (2.365 g, 3.84 mmol, 86 % yield) as a light-yellow solid. m/z (ESI, +ve ion) 616.0 (M+H). ^{1}H NMR (400 MHz, DMSO-d6) δ 0.96 (d, J = 6.8 Hz, 3 H), 1.05 (d, J = 6.8 Hz, 3 H), 1.08 (s, 9 H), 1.96 (s, 3 H), 2.67 (sept, J = 6.8 Hz, 1 H), 4.84 (s, 2 H), 7.12 (s, 1 H), 7.34 (s, 1 H), 7.40 – 7.53 (m, 6 H), 7.69 (dd, J = 6.2, 1.2 Hz, 4 H), 8.46 (d, J = 7.5 Hz, 1 H), 12.18 (br s, 1 H).

[0441] Step 3: *tert*-Butyl (2*R*,5*S*)-4-(1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-7-chloro-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a solution of 1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-7-chloro-6-fluoropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (2.37 g, 3.84 mmol) in acetonitrile (20 mL) under N_2 was DIPEA (1.00 mL, 5.76 mmol) and $POCl_3$ (0.537 mL, 5.76 mmol) at room temperature. The resulting mixture was then stirred at 60 °C for 1 hour 20 mins. The mixture was then cooled to room temperature and was concentrated under reduced pressure. The crude mixture was then dissolved in dichloromethane (20 mL) and was cooled to 0 °C. Then, DIPEA (3.34 mL, 19.19 mmol) was added, followed by *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (1.65 g, 7.68 mmol, eNovation Chemicals LLC, Bridgewater, NJ, USA). The resulting mixture was then stirred at room temperature for 2.5 hours. Then, the mixture was diluted with water (150 mL) and EtOAc (50 mL). The mixture was then extracted with EtOAc (2 x 30 mL). The combined organic extracts were then dried over $MgSO_4$ and concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-50% EtOAc/heptane)

provided *tert*-butyl (2*R*,5*S*)-4-(1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-7-chloro-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (2.78 g, 3.42 mmol, 89 % yield) as a tan foam. m/z (ESI, +ve ion) 812.1 (M+H). ^{1}H NMR (400 MHz, DMSO-d6) δ 0.96 (dd, J = 13.5, 6.8 Hz, 3 H), 1.02 (dd, J = 6.8, 3.4 Hz, 3 H), 1.08 (s, 9 H), 1.10 - 1.13 (m, 2 H), 1.28 (br dd, J = 10.2, 6.6 Hz, 3 H), 1.44 (s, 9 H), 1.87 (s, 3 H), 2.34 - 2.47 (m, 1 H), 3.42 - 3.59 (m, 1 H), 3.62 - 3.71 (m, 1 H), 3.76 - 3.88 (m, 1 H), 3.89 - 4.09 (m, 1 H), 4.13 - 4.25 (m, 1 H), 4.26 - 4.39 (m, 1 H), 4.74 - 4.80 (m, 1 H), 4.83 (s, 2 H), 7.11 (br d, J = 6.4 Hz, 1 H), 7.31 (br d, J = 10.0 Hz, 1 H), 7.42 - 7.54 (m, 6 H), 7.65 - 7.75 (m, 4 H), 8.35 (dd, J = 12.9, 8.3 Hz, 1 H).

[0442] Step 4: *tert*-Butyl (2*R*,5*S*)-4-(7-(2-amino-6-fluorophenyl)-1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of *tert*-butyl (2*R*,5*S*)-4-(1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-7-chloro-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (450 mg, 0.554 mmol), (2-amino-6-fluorophenyl)boronic acid (129 mg, 0.831 mmol, Aces Pharma, Princeton, NJ , USA), potassium carbonate (383 mg, 2.77 mmol), tetrakis(triphenylphosphine)palladium(0) (64.0 mg, 0.055 mmol) in dioxane (4.4 mL) and water (1.1 mL) was deoxygenated by bubbling nitrogen gas through the solvents for 10 mins. Then, the mixture was stirred at 90 °C overnight. Then, the mixture was cooled to room temperature and was concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc/heptane) provided *tert*-butyl (2*R*,5*S*)-4-(7-(2-amino-6-fluorophenyl)-1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (513.4 mg, 0.579 mmol, 94 % yield) as a goldenrod-colored. m/z (ESI, +ve ion) 887.1 (M+H).

[0443] Step 5: 2-Methyl-2-propanyl (2*R*,5*S*)-4-(7-(2-(bis(((2-methyl-2-propanyl)oxy)carbonyl)amino)-6-fluorophenyl)-6-fluoro-1-(2-methyl-4-((((2-methyl-2-propanyl)(diphenyl)silyl)oxy)methyl)-6-(2-propanyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethyl-1-piperazinecarboxylate. To a solution of *tert*-butyl (2*R*,5*S*)-4-(7-(2-amino-6-fluorophenyl)-1-(4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylphenyl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (490 mg, 0.552 mmol) in dichloromethane (4 mL) was added di-*tert*-butyl dicarbonate (241 mg, 1.11 mmol),

4-dimethylaminopyridine (135 mg, 1.11 mmol), and triethylamine (0.155 mL, 1.11 mmol). The resulting mixture was then stirred at room temperature for 4.5 hours. Then, the mixture was concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc/heptane) provided 2-methyl-2-propanyl (2*R*,5*S*)-4-(7-(2-(bis(((2-methyl-2-propanyl)oxy)carbonyl)amino)-6-fluorophenyl)-6-fluoro-1-(2-methyl-4-((((2-methyl-2-propanyl)(diphenyl)silyl)oxy)methyl)-6-(2-propanyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethyl-1-piperazinecarboxylate (470 mg, 0.432 mmol, 78 % yield) as a yellow solid. m/z (ESI, +ve ion) 1087.2 (M+H).

[0444] Step 6: 2-Methyl-2-propanyl (2*R*,5*S*)-4-(7-(2-(bis(((2-methyl-2-propanyl)oxy)carbonyl)amino)-6-fluorophenyl)-6-fluoro-1-(4-(hydroxymethyl)-2-methyl-6-(2-propanyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethyl-1-piperazinecarboxylate. To a solution of 2-methyl-2-propanyl (2*R*,5*S*)-4-(7-(2-(bis(((2-methyl-2-propanyl)oxy)carbonyl)amino)-6-fluorophenyl)-6-fluoro-1-(2-methyl-4-((((2-methyl-2-propanyl)(diphenyl)silyl)oxy)methyl)-6-(2-propanyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethyl-1-piperazinecarboxylate (465 mg, 0.428 mmol) in tetrahydrofuran (3.5 mL) at 0 °C was added tetra-n-butylammonium fluoride 1M in THF (0.428 mL, 0.428 mmol). The resulting mixture was then stirred at room temperature for 4.5 hours and was then concentrated under reduced pressure. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc/heptane) provided 2-methyl-2-propanyl (2*R*,5*S*)-4-(7-(2-(bis(((2-methyl-2-propanyl)oxy)carbonyl)amino)-6-fluorophenyl)-6-fluoro-1-(4-(hydroxymethyl)-2-methyl-6-(2-propanyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethyl-1-piperazinecarboxylate (125 mg, 0.147 mmol, 34 % yield) as a yellow solid. m/z (ESI, +ve ion) 749.2 (M+H).

[0445] Step 7: 2-Methyl-2-propanyl (2*R*,5*S*)-4-(7-(2-(bis(((2-methyl-2-propanyl)oxy)carbonyl)amino)-6-fluorophenyl)-1-(4-((dimethylamino)methyl)-2-methyl-6-(2-propanyl)phenyl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethyl-1-piperazinecarboxylate. To a solution of 2-methyl-2-propanyl (2*R*,5*S*)-4-(7-(2-(bis(((2-methyl-2-propanyl)oxy)carbonyl)amino)-6-fluorophenyl)-6-fluoro-1-(4-(hydroxymethyl)-2-methyl-6-(2-propanyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethyl-1-piperazinecarboxylate (125 mg, 0.147 mmol) in dichloromethane (1.5 mL) at 0 °C under N_2 was added triethylamine (0.062 mL, 0.442 mmol) followed by a solution of methanesulfonyl chloride (0.011 mL, 0.147 mmol) in DCM (0.2 mL). The resulting mixture was then stirred at 0 °C for 1 hour. Then, dimethylamine (2M in THF) (0.736 mL, 1.47 mmol)

and triethylamine (0.062 mL, 0.442 mmol) were added. The mixture was stirred at room temperature for 30 mins. Then, the mixture was diluted with saturated $NaHCO_3$ (5 mL) and was then extracted with EtOAc (2 x 20 mL). Chromatographic purification of the residue (silica gel, 0%-20% NH3 2M in MeOH/DCM) provided 2-methyl-2-propanyl (2*R*,5*S*)-4-(7-(2-(bis(((2-methyl-2-propanyl)oxy)carbonyl)amino)-6-fluorophenyl)-1-(4-((dimethylamino)methyl)-2-methyl-6-(2-propanyl)phenyl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethyl-1-piperazinecarboxylate (98 mg, 0.112 mmol, 76 % yield) as a yellow solid. m/z (ESI, +ve ion) 876.2 (M+H).

[0446] Step 8: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-amino-6-fluorophenyl)-1-(4-((dimethylamino)methyl)-2-isopropyl-6-methylphenyl)-6-fluoropyrido[2,3-*d*]pyrimidin-2(1*H*)-one trifluoroacetic acid. To a solution of 2-methyl-2-propanyl (2*R*,5*S*)-4-(7-(2-(bis(((2-methyl-2-propanyl)oxy)carbonyl)amino)-6-fluorophenyl)-1-(4-((dimethylamino)methyl)-2-methyl-6-(2-propanyl)phenyl)-6-fluoro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethyl-1-piperazinecarboxylate (98 mg, 0.112 mmol) in dichloromethane (1.0 mL) was added trifluoroacetic acid (0.167 mL, 2.237 mmol). The resulting mixture was then stirred at room temperature for 40 mins. Then, the mixture was concentrated and dried *in vacuo*. Then, the residue was dissolved in DCM (20 mL) and saturated $NaHCO_3$ (40 mL) was added. The mixture was then stirred at room temperature for 15 mins. The organic layer was then collected and aqueous layer was extracted with EtOAc (2 x 20 mL). The combined organic extracts were then dried over $MgSO_4$, concentrated, and dried *in vacuo*. The residue was then dissolved in dichloromethane (1 mL). The mixture was then cooled to 0 °C under N_2 and triethylamine (0.015 mL, 0.106 mmol) was added followed by a solution of acryloyl chloride (8.6 µl, 0.106 mmol) in DCM (0.2 mL) dropwise. After stirring at 0 °C for 5 mins, triethylamine (0.075 mL) and potassium carbonate (0.032 mL, 0.530 mmol) were added. The mixture was then stirred for 25 mins. The mixture was then quenched with saturated $NaHCO_3$ (5 mL) and was extracted with EtOAc (3 x 15 mL). The combined organic extracts were then dried over $MgSO_4$ and concentrated. Chromatographic purification of the residue (silica gel, 0%-20% NH_3 2M in MeOH/DCM) then by purified via preparative HPLC (Phenomenex Gemini C18 column, 150×30 mm, 10u, 110 A, 10-100% 0.1% TFA in MeCN/H_2O) provided 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-amino-6-fluorophenyl)-1-(4-((dimethylamino)methyl)-2-isopropyl-6-methylphenyl)-6-fluoropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (2.3 mg, 1.83 µmol, 1.7 % yield), TFA salt as a yellow solid. m/z (ESI, +ve ion) 630.2 (M+H). ^{19}F NMR (METHANOL-d4) δ: -76.95 (s, 1F),

-128.59 (dd, J=32.5, 4.8 Hz, 1F), -130.14 (br d, J=32.1 Hz, 1F). ^{1}H NMR (400 MHz, METHANOL-d4) δ 8.17-8.25 (m, 1H), 7.31-7.36 (m, 1H), 7.21-7.25 (m, 1H), 6.75-6.92 (m, 3H), 6.57 (td, J=2.67, 5.65 Hz, 1H), 6.30 (ddd, J=1.76, 6.69, 16.64 Hz, 1H), 5.83 (ddd, J=1.76, 6.17, 10.52 Hz, 1H), 4.87-5.15 (m, 2H), 4.19-4.59 (m, 2H), 3.89-4.11 (m, 2H), 3.83 (br s, 2H), 2.58-2.66 (m, 1H), 2.42-2.58 (m, 6H), 1.99-2.04 (m, 3H), 1.45-1.51 (m, 3H), 1.19 (dd, J=2.49, 6.84 Hz, 3H), 0.99-1.06 (m, 3H), 0.91 (br t, J=6.74 Hz, 3H).

Example 185

1-((2*R*,5*S*)-4-(6-Chloro-7-(2-fluorophenyl)-2-imino-1-(2-isopropylphenyl)-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one



[0447] Step 1: *tert*-Butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropylphenyl)amino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate. A vial was charged with *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 165**, 1 g, 2.1 mmol), copper(I) iodide (40 mg, 0.2 mmol), and potassium phosphate (1.1 g, 5.2 mmol). The vial was evacuated and backfilled with nitrogen. DMSO (4 ml) were added to the reaction mixture, followed by addition of 2-(methylethyl)phenylamine (365 µl, 2.7 mmol) added. The reaction mixture was heated to 80 °C. After 1 hour, the reaction mixture was diluted with water and extracted with EtOAc. The organic phase was washed with 1M LiCl solution and dried over $MgSO_4$. The filtrate was concentrated *in vacuo* to give the

crude material. Purification by preparative SFC (Chiralcel IC column (5 μm, 21 x 250 mm), F = 80 mL, (20% MeOH with 0.2% NEt_3)/80% CO_2) afforded *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropylphenyl)amino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate. *m/z* (ESI, +ve ion): 580.0 $(M+H)^+$.

[0448] Step 2: *tert*-Butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropylphenyl)amino)pyridin-3-yl)(cyanoimino)methyl)-2,5-dimethylpiperazine-1-carboxylate. Triethylamine (0.2 mL, 1.5 mmol) and cyanogen bromide (3 M in DCM; 0.29 mL, 0.86 mmol) were added to a solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropylphenyl)amino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (50 mg, 0.086 mmol) in tetrahydrofuran (1.5 mL). The reaction mixture was allowed to stir overnight. The reaction mixture was filtered and the filtrate was washed with aqueous saturated $NaHCO_3$ solution, dried over $MgSO_4$ and concentrated under reduced pressure. The crude product was taken onto the next step without further purification. *m/z* (ESI, +ve ion): 604.6 $(M)^+$.

[0449] Step 3: 1-((2*R*,5*S*)-4-(6-Chloro-7-(2-fluorophenyl)-2-imino-1-(2-isopropylphenyl)-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one. Hydrochloric acid (5N, 0.13 mL, 0.66 mmol) was added to a solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropylphenyl)amino)pyridin-3-yl)(cyanoimino)methyl)-2,5-dimethylpiperazine-1-carboxylate (40 mg, 0.066 mmol) in MeOH (1 mL). The reaction mixture was stirred for 10 min at rt and then heated to 72 °C for one hour. The reaction mixture was cooled to rt, neutralized with aqueous saturated $NaHCO_3$ solution and extracted with EtOAc. The organic extract was washed with brine and dried over $MgSO_4$. The solution was filtered and concentrated *in vacuo* to give the crude material. Residual solvents were removed from the crude residue by azeotropic distillation with toluene (3 mL). The residue was dissolved in DCM (2 mL) and the solution was cooled to 10 °C . 1,1'-Dimethyltriethylamine (0.023 mL, 0.13 mmol) added, followed by solution of 2-propenoyl chloride (1.1 M in DCM; 0.06 mL, 0.066 mmol). After 5 min, the reaction mixture was concentrated under reduced pressure. The residue was dissolved in MeOH (1 mL) and treated with hydrochloric acid (5N, 0.3 mL). The reaction mixture was heated to 72 °C for 30 min and then neutralized with aqueous saturated $NaHCO_3$ solution (5 mL). The reaction mixture was extracted with EtOAc and washed with brine. The organic phase was separated and dried over $MgSO_4$. The solvent was removed under reduced pressure and the crude material was absorbed onto silica gel. Purification by silica gel chromatography (eluent: 10-100% of EtOAc/heptane,

followed by 100% (3:1 EtOH in EtOAc), followed by 100% (20 % (2 M NH3/MeOH)in DCM)) provided 1-((2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-2-imino-1-(2-isopropylphenyl)-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one (10 mg, 9 μmol, 13 % yield). *m/z* (ESI, +ve ion): 559.3 $(M+H)^+$. ^{1}H NMR (400 MHz, DMSO-*d*6) δ 0.94 - 1.01 (m, 3 H) 1.14 - 1.17 (m, 3 H) 1.22 - 1.31 (m, 3 H) 1.47 (br s, 3 H) 2.52 - 2.66 (m, 1 H) 3.38 - 4.29 (m, 4 H) 4.52 - 5.16 (m, 2 H) 5.78 (br d, *J*=10.37 Hz, 1 H) 6.21 (dt, *J*=16.64, 2.15 Hz, 1 H) 6.65 - 6.98 (m, 1 H) 7.14 - 7.64 (m, 8 H) 8.43 - 8.69 (m, 1 H) 9.04 (br s, 1 H)

Example 186

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0450] Step 1: *tert*-Butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropylphenyl)amino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate. The synthesis of the title compound is described in Step 1 of Example 185.

[0451] Step 2: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate.

[0452] A solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropylphenyl)amino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (0.065 g, 0.11 mmol) and 1,1'-carbonyldiimidazole (0.036 g, 0.22 mmol, Acros Organics) in 2-Me-tetrahydrofuran (1 mL) was heated to 100 °C for 12 h. The reaction mixture was cooled

to rt and absorbed onto a pad of silica gel. Purification by silica gel chromatography (eluent: 10-100% of EtOAc/heptane) afforded *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (25 mg, 0.04 mmol, 36% yield). *m/z* (ESI, +ve ion): 605.6 $(M+H)^+$.

[0453] Step 3: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. TFA (0.1 mL) was added to a solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate in (1 mL) at rt. After 1 h, the reaction mixture was concentrated under reduced pressure and residual solvents were removed from the crude residue by azeotropic distillation with toluene (2 mL). The crude material was dissolved in DCM (2 mL) and the solution was cooled to 10 °C, followed by the addition of 1,1'-dimethyltriethylamine (0.02 mL, 0.1 mmol) and 2-propenoyl chloride (1.1 M in DCM; 0.067 mL, 0.074 mmol). After 10 min, the reaction mixture was diluted with DCM and treated with aqueous saturtaed $NaHCO_3$ (5 mL) and brine. The organic phase was separated, dried over MgSO4 and filtered. The crude material was absorbed onto silica gel and purified by siliga gel chromatography (eluent: 10-100% EtOAc/heptane) to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (10 mg, 0.018 mmol, 36 % yield). *m/z* (ESI, +ve ion): 560.2 $(M+H)^+$. ^1H NMR (400 MHz, DMSO-*d6*) δ 0.92 - 1.01 (m, 3 H) 1.07 (d, *J*=6.84 Hz, 3 H) 1.17 - 1.37 (m, 6 H) 2.53 - 2.61 (m, 1 H) 3.39 - 4.27 (m, 4 H) 4.42 - 4.95 (m, 2 H) 5.69 - 5.80 (m, 1 H) 6.19 (br d, *J*=16.79 Hz, 1 H) 6.73 - 6.92 (m, 1 H) 7.14 (d, *J*=7.67 Hz, 1 H) 7.18 - 7.37 (m, 5 H) 7.38 - 7.44 (m, 1 H) 7.46 - 7.55 (m, 1 H) 8.35 - 8.47 (m, 1 H)

Example 187

***N*-[5-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-1-yl]-4,6-diisopropyl-pyrimidin-2-yl]-*N*-methyl-acetamide**



Intermediate 232

[0454] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. Methylamine (2.0 M in tetrahydrofuran, 0.312 mL, 0.624 mmol) was added to a stirred solution of *tert*-butyl (2*R*,5*S*)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (91 mg, 0.125 mmol, **Intermediate 232**) in acetonitrile (0.5 mL). The reaction mixture was stirred at room temperature for 9 h. The reaction mixture was concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-50% [(3:1) EtOAc/EtOH]/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as a yellow solid (45 mg, 0.066 mmol, 53.1 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.13 (1 H, s) 7.45 - 7.52 (1 H, m) 7.21 - 7.29 (2 H, m) 7.14 - 7.20 (1 H, m) 3.53 - 5.12 (7 H, m) 3.05 (3 H, d, *J*=4.98 Hz) 2.49 - 2.61 (2 H, m) 1.57 (9 H, s) 1.44 - 1.54 (3 H, m) 1.30 - 1.36 (3 H, m) 1.21 (3 H, dd, *J*=6.34, 4.25 Hz) 1.01 (6 H, br t, *J*=7.26 Hz). m/z (ESI, +ve ion): 679.1 $(M+H)^+$.

[0455] Step 2: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(*N*-methylacetamido)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(methylamino)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-

dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (154 mg, 0.227 mmol), *N*,*N*-diisopropylethylamine (0.198 mL, 1.134 mmol), and 4-(*N*,*N*-dimethylamino)-pyridine (1.385 mg, 0.011 mmol) were mixed in toluene (1 mL) in a sealed vial. Acetyl chloride (0.036 mL, 0.453 mmol) was added, and the reaction mixture was stirred at 80 °C for 18 h. The reaction mixture was diluted with EtOAc (60 mL) and washed with saturated aqueous NH_4Cl (40 mL). The organic layer was separated, washed with brine (40 mL), dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-50% [(3:1) EtOAc/EtOH]/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(*N*-methylacetamido)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as a yellow solid (135 mg, 0.187 mmol, 83 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.19 (1 H, s) 7.47 - 7.54 (1 H, m) 7.15 - 7.29 (3 H, m) 3.59 (3 H, s) 3.55 - 5.12 (6 H, m) 2.73 (2 H, dq, *J*=13.01, 6.58 Hz) 2.62 (3 H, s) 1.59 (9 H, s) 1.49 - 1.57 (3 H, m) 1.32 - 1.39 (3 H, m) 1.27 (6 H, dd, *J*=6.63, 2.49 Hz) 1.07 (6 H, t, *J*=6.84 Hz). m/z (ESI, +ve ion): 721.3 $(M+H)^+$.

[0456] Step 3: *N*-(5-(4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidin-2-yl)-*N*-methylacetamide. Trifluoroacetic acid (0.5 mL, 6.71 mmol) was added to a stirred solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(*N*-methylacetamido)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (135 mg, 0.187 mmol) in dichloromethane (1 mL). The reaction mixture was stirred at room termperature for 20 min. The reaction mixture was concentrated *in vacuo* to give crude *N*-(5-(6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidin-2-yl)-*N*-methylacetamide that was used without purification assuming 100% yield.

[0457] Acryloyl chloride (1.1 M in DCM, 0.204 mL, 0.224 mmol) was added to a stirred mixture of the crude *N*-(5-(6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidin-2-yl)-*N*-methylacetamide (116 mg, 0.187 mmol) and *N*,*N*-diisopropylethylamine (0.163 mL, 0.934 mmol) in dichloromethane (1 mL). The reaction mixture was stirred at room temperature for 45 min. The reaction mixture was diluted with DCM (50 mL) and washed with saturated aqueous NH_4Cl (30 mL). The organic layer was separated, dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-100% EtOAc/heptane) to provide *N*-(5-(4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-

6-chloro-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidin-2-yl)-*N*-methylacetamide as a light yellow solid (95 mg, 0.141 mmol, 75 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.19 (1 H, s) 7.47 - 7.55 (1 H, m) 7.16 - 7.28 (3 H, m) 6.59 - 6.76 (1 H, m) 6.42 - 6.52 (1 H, m) 5.84 - 5.91 (1 H, m) 3.59 (3 H, s) 3.52 - 5.26 (6 H, m) 2.65 - 2.78 (2 H, m) 2.61 (3 H, s) 1.47 - 1.58 (4 H, m) 1.37 - 1.42 (2 H, m) 1.27 (6 H, dd, *J*=6.63, 2.28 Hz) 1.03 - 1.11 (6 H, m). m/z (ESI, +ve ion): 675.3 $(M+H)^+$.

Example 188

***N*-[5-[6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(2-fluorophenyl)-2-oxo-pyrido[2,3-*d*]pyrimidin-1-yl]-4,6-diisopropyl-pyrimidin-2-yl]acetamide**



Intermediate 232



[0458] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-((4-methoxybenzyl)amino)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. 4-Methoxybenzylamine (0.283 mL, 2.147 mmol) was added to a stirred solution of *tert*-butyl (2*R*,5*S*)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (313 mg, 0.429 mmol, **Intermediate 232**) in acetonitrile (1.5 mL). The reaction mixture was stirred at room temperature for 20 h before being warmed to 40 °C and stirred for another 7 h. The reaction mixture was concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-50% [(3:1) EtOAc/EtOH]/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-

diisopropyl-2-((4-methoxybenzyl)amino)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as a light yellow solid (179 mg, 0.228 mmol, 53.1 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.05 (1 H, m) 7.25 - 7.45 (3 H, m) 7.04 - 7.19 (3 H, m) 6.79 - 6.93 (2 H, m) 3.76 (3 H, s) 3.43 - 5.31 (8 H, m) 2.40 - 2.57 (2 H, m) 1.48 (9 H, s) 1.37 - 1.44 (3 H, m) 1.18 - 1.27 (3 H, m) 1.07 - 1.17 (6 H, m) 0.88 - 0.99 (6 H, m). m/z (ESI, +ve ion): 785.2 $(M+H)^+$.

Step 2: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(*N*-(4-methoxybenzyl)acetamido)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-((4-methoxybenzyl)amino)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (169 mg, 0.215 mmol), *N*,*N*-diisopropylethylamine (0.188 mL, 1.076 mmol), and 4-(*N*,*N*-dimethylamino)-pyridine (2 mg, 0.016 mmol) were mixed in toluene (1 mL) in a sealed vial. Acetyl chloride (0.034 mL, 0.430 mmol) was added, and the reaction mixture was stirred at 80 °C for 16 h. The reaction mixture was diluted with EtOAc (50 mL) and washed with saturated aqueous NH_4Cl (40 mL). The organic layer was separated, washed with brine (40 mL), dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-70% EtOAc/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(*N*-(4-methoxybenzyl)acetamido)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as a light yellow solid (108 mg, 0.131 mmol, 60.7 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.08 (1 H, s) 7.36 - 7.44 (1 H, m) 7.33 (2 H, d, *J*=8.71 Hz) 7.05 - 7.15 (3 H, m) 6.69 - 6.73 (2 H, m) 5.22 (2 H, s) 3.72 (3 H, s) 3.35 - 5.01 (6 H, m) 2.55 - 2.67 (2 H, m) 2.51 (3 H, s) 1.48 (9 H, s) 1.38 - 1.46 (3 H, m) 1.20 - 1.28 (3 H, m) 1.15 (6 H, dd, *J*=6.63, 2.70 Hz) 0.93 (6 H, t, *J*=7.26 Hz). m/z (ESI, +ve ion): 827.3 $(M+H)^+$.

[0459] Step 3: *N*-(5-(4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidin-2-yl)acetamide. Trifluoroacetic acid (0.5 mL, 6.71 mmol) and trifluoromethane sulfonic acid (0.05 mL, 0.563 mmol) were added to a stirred solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-(*N*-(4-methoxybenzyl)acetamido)pyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (100 mg, 0.121 mmol) in dichloromethane (0.5 mL). The reaction mixture was stirred at room

temperature for 7 h. The reaction mixture was diluted with DCM (40 mL) and washed with saturated aqueous $NaHCO_3$ (25 mL). The organic layer was separated, dried over $MgSO_4$, filtered, and concentrated *in vacuo* to give crude *N*-(5-(6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidin-2-yl)acetamide that was used without purification assuming 100% yield.

[0460] Acryloyl chloride (1.1 M in DCM, 0.131 mL, 0.144 mmol) was added to a stirred soluiton of the crude *N*-(5-(6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidin-2-yl)acetamide (73 mg, 0.120 mmol) in dichloromethane (0.5 mL). The reaction mixture was stirred at room temperaturefor 2 h. The reaction mixture was concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-50% [(3:1) EtOAc/EtOH]/heptane) to provide *N*-(5-(4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-2-oxopyrido[2,3-*d*]pyrimidin-1(2*H*)-yl)-4,6-diisopropylpyrimidin-2-yl)acetamide as a white solid (20 mg, 0.030 mmol, 25.2 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.17 (1 H, s) 7.97 (1 H, s) 7.46 - 7.53 (1 H, m) 7.14 - 7.28 (3 H, m) 6.57 - 6.74 (1 H, m) 6.40 - 6.50 (1 H, m) 5.82 - 5.89 (1 H, m) 3.50 - 5.24 (6 H, m) 2.59 - 2.73 (5 H, m) 1.45 - 1.56 (4 H, m) 1.38 (2 H, br d, *J*=6.84 Hz) 1.23 (6 H, dd, *J*=6.63, 2.70 Hz) 0.99 - 1.07 (6 H, m). m/z (ESI, +ve ion): 661.2 $(M+H)^+$.

Example 189

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0094] **Step 1: *tert*-Butyl (2*R*,5*S*)-4-(1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**. A mixture of *tert*-butyl (2*R*,5*S*)-4-(1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.228 g, 0.284

mmol, **Intermediate 211**), (2-fluorophenyl)boronic acid (0.060 g, 0.43 mmol; Combi-Blocks, Inc., San Diego, CA), $PdCl_2$(dppf) (0.021 g, 0.028 mmol), and potassium acetate (0.084 g, 0.85 mmol) in 1,4-dioxane (2.6 mL)/water (0.26 mL) was sparged with nitrogen then stirred at 100 °C for 1 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–60% [(3:1) EtOAc/EtOH]/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (252 mg, 0.292 mmol, >99% yield) as a brown solid. MS (ESI, +ve) *m/z*: 863.1 $(M + 1)^+$.

[0461] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.252 g, 0.292 mmol) in 2,2,2-trifluoroacetic acid (3.33 g, 29.2 mmol) was stirred at RT for 15 min. The reaction mixture was concentrated.

[0462] A yellow solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.700 mL, 0.350 mmol), and DIPEA (0.153 mL, 0.876 mmol) in dichloromethane (1.5 mL) was stirred at RT for 15 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*.

[0463] A solution of the resulting oil in 2,2,2-trifluoroacetic acid (3.33 g, 29.2 mmol) was heated in the microwave at 100 °C for 5 min, 120 °C for 10 min, then 100 °C for 30 min. The reaction mixture was concentrated. The crude product in DCM was loaded onto the column and was purified via automated flash chromatography (silica gel, 0–100% [(3:1) EtOAc/EtOH]/heptane then 0–5% [2 M ammonia in MeOH]/DCM) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (11 mg, 0.019 mmol, 7% yield) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.54 (s, 1 H) 8.08 - 8.13 (m, 1 H) 7.41 - 7.48 (m, 1 H) 7.23 - 7.30 (m, 1 H) 7.16 - 7.22 (m, 1 H) 7.08 - 7.16 (m, 1 H) 6.50 - 6.71 (m, 1 H) 6.33 - 6.46 (m, 1 H) 5.75 - 5.85 (m, 1 H) 4.89 - 5.19 (m, 2 H) 4.79 - 4.88 (m, 2 H) 4.28 - 4.53 (m, 1

H) 3.38 - 4.01 (m, 3 H) 2.48 - 2.65 (m, 1 H) 1.30 - 1.52 (m, 6 H) 1.15 - 1.20 (m, 3 H) 0.99 - 1.05 (m, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.26 - -112.15 (m, 1 F). MS (ESI, +ve) *m/z*: 576.8 $(M + 1)^+$.

Example 189-1

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0464] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (139 mg, 0.241 mmol) were separated by chiral SFC: ID 250 x 21 mm, 5 μm, 45% MeOH/CO_2, 70 g/min, 204 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (58 mg, 0.10 mmol, 42% yield) (first eluting isomer) as a yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.57 (s, 1 H) 8.11 (s, 1 H) 7.40 - 7.50 (m, 1 H) 7.23 - 7.29 (m, 1 H) 7.08 - 7.21 (m, 2 H) 6.50 - 6.69 (m, 1 H) 6.34 - 6.45 (m, 1 H) 5.76 - 5.85 (m, 1 H) 4.90 - 5.21 (m, 1 H) 4.74 (s, 2 H) 3.39 - 4.56 (m, 5 H) 2.49 - 2.63 (m, 1 H) 1.33 - 1.52 (m, 6 H) 1.19 (d, *J* = 6.6 Hz, 3 H) 1.03 (br d, *J* = 6.4 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.21 - -112.13 (m, 1 F). MS (ESI, +ve) *m/z*: 577.1 $(M + 1)^+$.

Example 189-2

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0465] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (139 mg, 0.241 mmol) were separated by chiral SFC: ID 250 x 21 mm, 5 μm, 45% MeOH/CO_2, 70 g/min, 204 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (58 mg, 0.10 mmol, 42% yield) (second eluting isomer) as a yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.56 (s, 1 H) 8.08 - 8.12 (m, 1 H) 7.41 - 7.49 (m, 1 H) 7.24 - 7.30 (m, 1 H) 7.17 - 7.22 (m, 1 H) 7.10 - 7.16 (m, 1 H) 6.51 - 6.70 (m, 1 H) 6.35 - 6.45 (m, 1 H) 5.77 - 5.85 (m, 1 H) 4.87 - 5.20 (m, 1 H) 4.76 (br s, 2 H) 3.39 - 4.51 (m, 5 H) 2.50 - 2.66 (m, 1 H) 1.30 - 1.51 (m, 6 H) 1.18 (d, *J* = 6.6 Hz, 3 H) 1.01 (d, *J* = 6.8 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.22 - -112.11 (m, 1 F). MS (ESI, +ve) *m/z*: 577.1 $(M + 1)^+$.

Example 190

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluoro-6-hydroxyphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0466] **Step 1: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one**. A solution of *tert*-butyl (2*R*,5*S*)-4-(1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.194 g, 0.241 mmol, **Intermediate 211**) and 2,2,2-trifluoroacetic acid (2.75 g, 24.1 mmol) in dichloromethane (1.2 mL) was stirred at RT for 15 min. The reaction mixture was concentrated.

[0467] A yellow solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.579 mL, 0.290 mmol), and DIPEA (0.126 mL, 0.724 mmol) in dichloromethane (1.2 mL) was stirred at RT for 30 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo* to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.171 g, 0.226 mmol). MS (ESI, +ve) *m/z*: 756.7 $(M + 1)^+$.

[0468] **Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one**. A solution of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.171 g, 0.226

mmol) and trifluoromethane sulfonic acid (0.020 mL, 0.226 mmol) in 2,2,2-trifluoroacetic acid (1.8 mL, 22.6 mmol) was stirred at RT for 30 min, 50 °C for 30 min, then 80 °C for 5 min. The reaction mixture was concentrated, diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo* to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.100 g, 0.193 mmol). MS (ESI, +ve) *m/z*: 516.8 $(M + 1)^+$.

[0469] Step 3: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluoro-6-hydroxyphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A mixture of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.100 g, 0.193 mmol), potassium F-phenol-trifluoroborate (0.063 g, 0.29 mmol, **Intermediate Q**), $PdCl_2$(dppf) (0.014 g, 0.019 mmol), and potassium acetate (0.057 g, 0.58 mmol) in 1,4-dioxane (1.8 mL)/water (0.18 mL) was sparged with nitrogen then stirred at 100 °C for 1.5 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–100% [(3:1) EtOAc/EtOH]/heptane: elutes slowly) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloro-7-(2-fluoro-6-hydroxyphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (36 mg, 0.061 mmol, 31% yield) as a light brown solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.47 - 8.53 (m, 1 H) 8.10 - 8.20 (m, 1 H) 7.35 - 7.63 (m, 1 H) 7.21 - 7.31 (m, 1 H) 6.73 - 6.79 (m, 1 H) 6.47 - 6.72 (m, 2 H) 6.31 - 6.44 (m, 1 H) 5.75 - 5.87 (m, 1 H) 4.79 - 5.23 (m, 4 H) 4.18 - 4.68 (m, 2 H) 3.75 - 4.04 (m, 2 H) 2.58 - 2.72 (m, 1 H) 1.33 - 1.49 (m, 6 H) 1.21 - 1.25 (m, 3 H) 1.19 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -111.92 (br s, 1 F). MS (ESI, +ve) *m/z*: 592.8 $(M + 1)^+$.

Example 191-1

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



Intermediate 215
single isomer

single isomer

[0470] A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.048 g, 0.073 mmol, **Intermediate 215**) in 2,2,2-trifluoroacetic acid (0.560 mL, 7.31 mmol) was stirred at RT for 10 min. The reaction mixture was concentrated.

[0471] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.175 mL, 0.088 mmol), and DIPEA (0.051 mL, 0.292 mmol) in dichloromethane (0.7 mL) was stirred at RT for 10 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (42 mg, 0.069 mmol, 94% yield) as a light yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.13 (s, 1 H) 7.42 - 7.50 (m, 1 H) 7.17 - 7.24 (m, 2 H) 7.09 - 7.17 (m, 1 H) 6.51 - 6.71 (m, 1 H) 6.36 - 6.45 (m, 1 H) 5.75 - 5.85 (m, 1 H) 3.43 - 5.24 (m, 6 H) 2.63 - 2.77 (m, 1 H) 2.20 - 2.27 (m, 3 H) 1.29 - 1.53 (m, 6 H) 1.22 (d, *J* = 6.8 Hz, 3 H) 1.04 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.95 - -112.86 (m, 1 F). MS (ESI, +ve) *m/z*: 610.0 $(M + 1)^+$.

Example 191-2

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



Intermediate 204
single isomer

single isomer

[0472] A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.103 g, 0.157 mmol, **Intermediate 204**) in 2,2,2-trifluoroacetic acid (1.20 mL, 15.7 mmol) was stirred at RT for 10 min. The reaction mixture was concentrated.

[0473] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.345 mL, 0.173 mmol), and DIPEA (0.082 mL, 0.471 mmol) in dichloromethane (0.8 mL) was stirred at RT for 15 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–60% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (47 mg, 0.077 mmol, 49% yield) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.13 (s, 1 H) 7.42 - 7.50 (m, 1 H) 7.17 - 7.24 (m, 2 H) 7.10 - 7.17 (m, 1 H) 6.51 - 6.70 (m, 1 H) 6.41 (br t, *J* = 14.7 Hz, 1 H) 5.77 - 5.85 (m, 1 H) 3.42 - 5.19 (m, 6 H) 2.63 - 2.75 (m, 1 H) 2.19 - 2.25 (m, 3 H) 1.31 - 1.51 (m, 6 H) 1.23 (d, *J* = 6.8 Hz, 3 H) 1.07 (br d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.96 - -112.83 (m, 1 F). MS (ESI, +ve) *m/z*: 610.1 $(M + 1)^+$.

Example 192

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-amino-6-fluorophenyl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0474] Step 1: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.723 g, 0.899 mmol, **Intermediate 211**) and 2,2,2-trifluoroacetic acid (10 g, 90 mmol) in dichloromethane (5 mL) was stirred at RT for 15 min. The reaction mixture was concentrated.

[0475] A yellow solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 2.16 mL, 1.08 mmol), and DIPEA (0.470 mL, 2.70 mmol) in dichloromethane (5 mL) was stirred at RT for 30 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo* to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one. MS (ESI, +ve) *m/z*: 756.8 $(M + 1)^+$.

[0476] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.682 g, 0.900

mmol) and trifluoromethane sulfonic acid (0.400 mL, 4.50 mmol) in 2,2,2-trifluoroacetic acid (6.9 mL, 90 mmol) was stirred at RT for 1 h. The reaction mixture was concentrated, diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo* to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.500 g, 0.966 mmol, >99% yield) as a light yellow solid. MS (ESI, +ve) *m/z*: 516.8 $(M + 1)^+$.

[0477] Step 3: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-amino-6-fluorophenyl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A mixture of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.055 g, 0.11 mmol), (2-amino-6-fluorophenyl)boronic acid pinacol ester (0.038 mL, 0.16 mmol, CombiPhos Catalysts, Inc., Trenton, NJ), $PdCl_2$(dppf) (8 mg, 11 μmol), and potassium acetate (0.031 g, 0.32 mmol) in 1,4-dioxane (1 mL)/water (0.1 mL) was sparged with nitrogen then stirred at 100 °C for 1.5 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–10% [2 M ammonia in MeOH]/DCM then 0–100% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-amino-6-fluorophenyl)-1-(4-amino-6-isopropylpyrimidin-5-yl)-6-chloropyrido[2,3-*d*]pyrimidin-2(1*H*)-one (3 mg, 5.1 μmol, 5% yield) as a yellow film. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.54 - 8.61 (m, 1 H) 8.13 - 8.22 (m, 1 H) 7.11 - 7.19 (m, 1 H) 6.35 - 6.72 (m, 4 H) 5.77 - 5.86 (m, 1 H) 4.80 - 5.23 (m, 2 H) 4.67 - 4.79 (m, 2 H) 4.17 - 4.57 (m, 3 H) 3.78 - 4.07 (m, 1 H) 3.74 (s, 1 H) 2.50 - 2.79 (m, 1 H) 1.38 - 1.53 (m, 6 H) 1.20 (d, J = 6.6 Hz, 3 H) 0.97 - 1.06 (m, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -111.95 (br d, J = 23.4 Hz, 1 F). MS (ESI, +ve) *m/z*: 591.8 $(M + 1)^+$.

Example 193

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0478] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.103 g, 0.157 mmol, Intermediate 204), methylzinc chloride (2 M in THF, 0.094 mL, 0.188 mmol; Sigma-Aldrich, St. Louis, MO), and [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) chloride (6 mg, 8 μmol) in 1,4-dioxane (0.8 mL) was sparged with nitrogen then stirred at 50 °C for 16 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% [(3:1) EtOAc/EtOH]/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (41 mg, 0.064 mmol, 41% yield). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.11 (s, 1 H) 7.39 - 7.47 (m, 1 H) 7.15 - 7.23 (m, 2 H) 7.09 - 7.14 (m, 1 H) 3.41 - 5.10 (m, 6 H) 2.70 (s, 3 H) 2.18 (s, 3 H) 1.51 (s, 9 H) 1.45 (br s, 3 H) 1.29 (d, J = 6.8 Hz, 3 H) 1.21 (d, J = 6.8 Hz, 3 H) 1.05 (d, J = 6.8 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.70 (br s, 1 F). MS (ESI, +ve) *m/z*: 636.2 $(M + 1)^+$.

[0479] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.041 g, 0.064 mmol) in 2,2,2-trifluoroacetic acid (0.49 mL, 6.4 mmol) was stirred at RT for 10 min. The reaction mixture was concentrated.

[0480] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.14 mL, 0.071 mmol), and DIPEA (0.034 mL, 0.19 mmol) in dichloromethane (0.3 mL) was stirred at RT for 15 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (50 mg, 0.085 mmol, >99% yield) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.11 (s, 1 H) 7.40 - 7.48 (m, 1 H) 7.15 - 7.23 (m, 2 H) 7.12 (t, *J* = 9.1 Hz, 1 H) 6.51 - 6.70 (m, 1 H) 6.33 - 6.45 (m, 1 H) 5.76 - 5.84 (m, 1 H) 3.44 - 5.19 (m, 6 H) 2.70 (s, 3 H) 2.61 - 2.69 (m, 1 H) 2.19 (s, 3 H) 1.30 - 1.51 (m, 6 H) 1.27 (d, *J* = 6.8 Hz, 3 H) 1.05 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.73 - -112.64 (m, 1 F). MS (ESI, +ve) *m/z*: 590.2 $(M + 1)^+$.

Example 194

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0481] **Step 1: 6,7-Dichloro-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione**. To a solution of 2,5,6-trichloronicotinamide (2.43 g, 10.8 mmol; **Intermediate P**) in THF (20 mL) was added oxalyl chloride (5.38 mL, 10.8 mmol), and the resulting mixture was stirred at 65 °C under a reflux condenser and drying tube for 2 h.

[0482] The reaction mixture was concentrated, dissolved in THF (20 mL), and cooled to 0 °C; a solution of 4-isopropyl-6-methylpyrimidin-5-amine (1.48 g, 9.79 mmol; **Intermediate I-5**) in THF (10 mL) was added, and the resulting mixture was stirred at 0 °C for 1 h. The reaction was diluted with EtOAc, quenched with brine and saturated aqueous NH_4Cl, extracted with EtOAc, and concentrated *in vacuo*. The crude product was purified via automated flash chromatography (silica gel, 0–5% MeOH/DCM) to give 2,5,6-trichloro-*N*-((4-isopropyl-6-methylpyrimidin-5-yl)carbamoyl)nicotinamide (3.24 g, 8.05 mmol, 82% yield) as a brown solid.

[0483] To a solution of 2,5,6-trichloro-*N*-((4-isopropyl-6-methylpyrimidin-5-yl)carbamoyl)nicotinamide (3.24 g, 7.95 mmol) in THF (40 mL) in a water-bath was added dropwise potassium bis(trimethylsilyl)amide (1 M in THF, 14.3 mL, 14.3 mmol). The mixture was stirred for 10 min; the water bath was removed, and the reaction mixture was stirred at RT for 30 min. The reaction mixture was quenched with saturated aqueous NH_4Cl and was extracted with EtOAc. The combined organics were dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The product was azeotropically dried to give 6,7-dichloro-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. ^{1}H NMR (DMSO-d_6) δ 12.20-12.47 (m, 1H), 8.90-9.18 (m, 1H), 8.46-8.67 (m, 1H), 2.82-3.09 (m, 1H), 2.19-2.28 (m, 3H), 1.06-1.10 (m, 3H), 0.98-1.04 (m, 3H). MS (ESI, +ve) *m/z*: 366.0 $(M + 1)^+$.

[0484] **Step 2: 6-Chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione**. A mixture of 6,7-dichloro-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.350 g, 0.956 mmol), (2-fluorophenyl)boronic acid (0.201 g, 1.43 mmol; Combi-Blocks, Inc., San Diego, CA), $PdCl_2$(dppf) (0.070 g, 0.096 mmol), and potassium acetate (0.281 g, 2.87 mmol) in 1,4-dioxane (8.7 mL)/water (0.87 mL) was sparged with nitrogen then stirred at 100 °C for 16 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% EtOAc/heptane) to give 6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-

d]pyrimidine-2,4(1*H*,3*H*)-dione (200 mg, 0.470 mmol, 49% yield) as an amber oil. MS (ESI, +ve) *m/z*: 425.9 $(M + 1)^+$.

[0485] Step 3: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of 6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.200 g, 0.470 mmol), phosphoryl trichloride (0.053 mL, 0.56 mmol), and DIPEA (0.491 mL, 2.82 mmol) in acetonitrile (1.2 mL) was stirred at 80 °C for 30 min. The reaction mixture was removed from the heating block, and a solution of *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (0.151 g, 0.704 mmol; eNovation Chemicals LLC, Bridgewater, NJ) in acetonitrile (1.2 mL) was added; the reaction mixture was stirred at RT for 1 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was loaded onto the column and was purified via automated flash chromatography (silica gel, 0–70% [(3:1) EtOAc/EtOH]/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (112 mg, 0.180 mmol, 38% yield) as an amber oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.03 (s, 1 H) 8.12 (s, 1 H) 7.39 - 7.47 (m, 1 H) 7.14 - 7.20 (m, 2 H) 7.07 - 7.14 (m, 1 H) 4.81 - 5.05 (m, 1 H) 4.31 - 4.81 (m, 1 H) 3.78 - 4.23 (m, 3 H) 3.44 - 3.66 (m, 1 H) 2.68 - 2.80 (m, 1 H) 2.25 (d, *J* = 5.8 Hz, 3 H) 1.52 (s, 9 H) 1.29 (d, *J* = 6.8 Hz, 3 H) 1.21 - 1.28 (m, 6 H) 1.06 (dd, *J* = 11.8, 6.6 Hz, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -112.68 (br d, *J* = 5.2 Hz, 1 F). MS (ESI, +ve) *m/z*: 621.8 $(M + 1)^+$.

[0486] Step 4: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.112 g, 0.180 mmol) in 2,2,2-trifluoroacetic acid (1.38 mL, 18.0 mmol) was stirred at RT for 30 min. The reaction mixture was concentrated *in vacuo*.

[0487] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.360 mL, 0.180 mmol), and DIPEA (0.125 mL, 0.720 mmol) in dichloromethane (0.9 mL) was stirred at RT for 1 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was

separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–100% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (97 mg, 0.17 mmol, 94% yield) as a yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.03 (s, 1 H) 8.13 (s, 1 H) 7.39 - 7.48 (m, 1 H) 7.07 - 7.21 (m, 3 H) 6.50 - 6.71 (m, 1 H) 6.41 (br t, *J* = 14.8 Hz, 1 H) 5.76 - 5.86 (m, 1 H) 4.33 - 5.22 (m, 3 H) 3.45 - 4.08 (m, 3 H) 2.65 - 2.81 (m, 1 H) 2.21 - 2.28 (m, 3 H) 1.31 - 1.48 (m, 6 H) 1.23 (dd, *J* = 6.8, 3.5 Hz, 3 H) 1.02 - 1.09 (m, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -112.71 - -112.60 (m, 1 F). MS (ESI, +ve) *m/z*: 575.8 $(M + 1)^+$.

Example 194-1

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0488] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (97 mg, 0.168 mmol) were separated by chiral SFC: Chirakpak IC, ID 250 x 2 mm, 5 μm, 45% methanol/CO_2, 70 mL/min, 220 nm, 185 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (31 mg, 0.054 mmol, 32% yield; first eluting isomer) as a light yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.03 (s, 1 H) 8.13 (s, 1 H) 7.40 - 7.48 (m, 1 H) 7.15 - 7.21 (m, 2 H) 7.07 - 7.15 (m, 1 H) 6.52 - 6.71 (m, 1 H) 6.34 - 6.46 (m, 1 H) 5.76 - 5.86 (m, 1 H) 5.02 - 5.20 (m, 1 H) 4.31 - 4.53 (m, 1 H) 4.00 - 4.13 (m, 1 H) 3.45 - 4.00 (m, 3 H) 2.66 - 2.80 (m, 1 H) 2.21 - 2.27 (m, 3 H) 1.31 - 1.52 (m, 6 H) 1.24 (d, *J* = 6.6 Hz, 3 H) 1.07

(d, J = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.71 - -112.58 (m, 1 F). MS (ESI, +ve) m/z: 575.9 $(M + 1)^+$.

Example 194-2

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0489] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (97 mg, 0.168 mmol) were separated by chiral SFC: Chirakpak IC, ID 250 x 2 mm, 5 μm, 45% methanol/CO_2, 70 mL/min, 220 nm, 185 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (10 mg, 0.017 mmol, 10% yield; second eluting isomer) as a light yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.11 (s, 1 H) 8.19 (s, 1 H) 7.45 - 7.54 (m, 1 H) 7.22 - 7.28 (m, 2 H) 7.14 - 7.22 (m, 1 H) 6.57 - 6.80 (m, 1 H) 6.40 - 6.54 (m, 1 H) 5.80 - 5.93 (m, 1 H) 5.11 - 5.29 (m, 1 H) 4.40 - 4.62 (m, 1 H) 3.50 - 4.20 (m, 4 H) 2.75 - 2.87 (m, 1 H) 2.26 - 2.36 (m, 3 H) 1.38 - 1.60 (m, 7 H) 1.30 (d, J = 6.8 Hz, 3 H) 1.12 (d, J = 6.6 Hz, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -112.70 - -112.61 (m, 1 F). MS (ESI, +ve) m/z: 575.9 $(M + 1)^+$.

Example 195

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)-6-methylpyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0490] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)-6-methyl-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.103 g, 0.157 mmol; **Intermediate 204**), methylzinc chloride (2 M in THF, 0.094 mL, 0.188 mmol; Sigma-Aldrich, St. Louis, MO), and [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) chloride (6 mg, 8 μmol) in 1,4-dioxane (0.8 mL) was sparged with nitrogen then stirred at 50 °C for 16 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% [(3:1) EtOAc/EtOH]/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)-6-methyl-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (30 mg, 0.049 mmol, 31% yield). ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.92 (s, 1 H) 7.36 - 7.44 (m, 1 H) 7.06 - 7.21 (m, 3 H) 3.47 - 5.05 (m, 6 H) 2.75 (s, 3 H) 2.28 (d, *J* = 1.5 Hz, 3 H) 2.24 (s, 3 H) 1.51 (s, 9 H) 1.44 (br s, 3 H) 1.28 (d, *J* = 6.8 Hz, 3 H) 1.22 (d, *J* = 6.8 Hz, 3 H) 1.05 (d, *J* = 6.8 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -114.75 (br s, 1 F). MS (ESI, +ve) *m/z*: 616.3 $(M + 1)^+$.

[0491] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)-6-methylpyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-

yl)-6-methyl-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.030 g, 0.049 mmol) in 2,2,2-trifluoroacetic acid (0.37 mL, 4.9 mmol) was stirred at RT for 10 min. The reaction mixture was concentrated.

[0492] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.11 mL, 0.054 mmol), and DIPEA (0.025 mL, 0.15 mmol) in dichloromethane (0.24 mL) was stirred at RT for 15 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-1-(4-isopropyl-2,6-dimethylpyrimidin-5-yl)-6-methylpyrido[2,3-*d*]pyrimidin-2(1*H*)-one (10 mg, 0.018 mmol, 36% yield) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.86 - 7.94 (m, 1 H) 7.36 - 7.44 (m, 1 H) 7.07 - 7.21 (m, 3 H) 6.51 - 6.72 (m, 1 H) 6.33 - 6.47 (m, 1 H) 5.74 - 5.84 (m, 1 H) 3.46 - 5.24 (m, 6 H) 2.70 (s, 3 H) 2.64 - 2.69 (m, 1 H) 2.29 (br s, 3 H) 2.14 - 2.22 (m, 3 H) 1.27 - 1.48 (m, 6 H) 1.21 (d, *J* = 6.8 Hz, 3 H) 1.04 (br d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -114.83 - -114.51 (m, 1 F). MS (ESI, +ve) *m/z*: 570.2 $(M + 1)^+$.

Example 196

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-(dimethylamino)-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0493] **Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-(dimethylamino)-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-

dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.103 g, 0.157 mmol; **Intermediate 204**) and dimethylamine (2 M in THF, 1.18 mL, 2.35 mmol; Sigma-Aldrich, St. Louis, MO) was stirred at 50 °C for 4 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% [(3:1) EtOAc/EtOH]/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-(dimethylamino)-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (101 mg, 0.152 mmol, 97% yield) as a yellow oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.06 (s, 1 H) 7.39 - 7.47 (m, 1 H) 7.15 - 7.25 (m, 2 H) 7.12 (t, *J* = 9.1 Hz, 1 H) 3.46 - 5.03 (m, 6 H) 2.46 - 2.57 (m, 1 H) 2.03 (s, 3 H) 1.51 (s, 9 H) 1.43 (br s, 3 H) 1.23 - 1.33 (m, 9 H) 1.17 (d, *J* = 6.6 Hz, 3 H) 1.00 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.86 (br s, 1 F). MS (ESI, +ve) *m/z*: 665.2 $(M + 1)^+$.

[0494] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-(dimethylamino)-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-(dimethylamino)-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.101 g, 0.152 mmol) in 2,2,2-trifluoroacetic acid (1.16 mL, 15.2 mmol) was stirred at RT for 10 min. The reaction mixture was concentrated.

[0495] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.33 mL, 0.17 mmol), and DIPEA (0.079 mL, 0.46 mmol) in dichloromethane (0.8 mL) was stirred at RT for 15 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-(dimethylamino)-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (97 mg, 0.157 mmol, >99% yield) as a yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.07 (s, 1 H) 7.40 - 7.48 (m, 1 H) 7.16 - 7.26 (m, 2 H) 7.12 (t, *J* = 9.1 Hz, 1 H) 6.50 - 6.71 (m, 1 H) 6.39 (br t, *J* = 15.2 Hz, 1 H) 5.74 - 5.86 (m, 1 H) 3.45 - 5.20 (m, 6 H) 3.19 (s, 6 H) 2.43 - 2.55 (m, 1 H) 1.99 - 2.06 (m, 3 H) 1.29 - 1.46 (m, 6 H)

1.17 (d, J = 6.8 Hz, 3 H) 1.00 (d, J = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.90 - -112.77 (m, 1 F). MS (ESI, +ve) m/z: 619.3 $(M + 1)^+$.

Example 197-1

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0496] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of 1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.556 g, 1.10 mmol; **Intermediate 216**), phosphoryl trichloride (0.123 mL, 1.32 mmol), DIPEA (0.768 mL, 4.41 mmol) in acetonitrile (5.5 mL) was stirred at 80 °C for 30 min.

[0497] The solution was cooled in an ice bath, then *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (0.236 g, 1.102 mmol; eNovation Chemicals LLC, Bridgewater, NJ) was added, and the dark red solution was stirred at RT for 5 min. The reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% EtOAc/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (423 mg, 0.603 mmol, 55% yield) as a light yellow solid. ^{1}H NMR (400 MHz, CDCl) δ 8.12 (s, 1 H) 7.43 - 7.50 (m, 1 H) 7.17 - 7.24 (m, 2 H) 7.10 - 7.17 (m, 1 H) 3.43 - 5.04 (m, 6 H) 2.61 - 2.74 (m, 1 H) 2.23 (s, 3 H) 1.51 (s, 9 H) 1.41 - 1.50 (m, 3 H) 1.26 - 1.28 (m, 3

H) 1.22 (d, J = 6.8 Hz, 3 H) 1.03 (d, J = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.94 (br s, 1 F). MS (ESI, +ve) *m/z*: 702.0 $(M + 1)^+$.

[0498] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.101 g, 0.144 mmol) in 2,2,2-trifluoroacetic acid (1.10 mL, 14.4 mmol) was stirred at RT for 10 min. The reaction mixture was concentrated.

[0499] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.317 mL, 0.158 mmol), and DIPEA (0.075 mL, 0.43 mmol) in dichloromethane (0.72 mL) was stirred at RT for 15 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (72 mg, 0.11 mmol, 76% yield) as a light yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.12 (s, 1 H) 7.42 - 7.52 (m, 1 H) 7.17 - 7.24 (m, 2 H) 7.10 - 7.17 (m, 1 H) 6.48 - 6.69 (m, 1 H) 6.41 (br t, J = 15.1 Hz, 1 H) 5.77 - 5.85 (m, 1 H) 3.41 - 5.18 (m, 6 H) 2.60 - 2.75 (m, 1 H) 2.19 - 2.26 (m, 3 H) 1.30 - 1.52 (m, 6 H) 1.22 (d, J = 6.6 Hz, 3 H) 1.04 (d, J = 6.8 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.97 - -112.88 (m, 1 F). MS (ESI, +ve) *m/z*: 656.0 $(M + 1)^+$.

Example 197-2

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0500] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of 1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.525 g, 1.04 mmol; **Intermediate 217**), phosphoryl trichloride (0.116 mL, 1.25 mmol), and DIPEA (0.725 mL, 4.16 mmol) in acetonitrile (5.2 mL) was stirred at 80 °C for 30 min.

[0501] The solution was cooled in an ice bath, then *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (0.223 g, 1.040 mmol; eNovation Chemicals LLC, Bridgewater, NJ) was added, and the dark red solution was stirred at RT for 5 min. The reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–70% EtOAc/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (317 mg, 0.452 mmol, 44% yield). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.13 (s, 1 H) 7.43 - 7.51 (m, 1 H) 7.18 - 7.24 (m, 2 H) 7.11 - 7.18 (m, 1 H) 3.42 - 5.05 (m, 6 H) 2.68 (quin, *J* = 6.8 Hz, 1 H) 2.22 (s, 3 H) 1.52 (s, 9 H) 1.43 - 1.50 (m, 3 H) 1.29 (d, *J* = 6.6 Hz, 3 H) 1.24 (d, *J* = 6.6 Hz, 3 H) 1.07 (d, *J* = 6.8 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.93 (br s, 1 F). MS (ESI, +ve) *m/z*: 702.0 $(M + 1)^+$.

[0502] **Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one**. A solution of *tert*-butyl (2*R*,5*S*)-4-(1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.043 g, 0.061 mmol) in 2,2,2-trifluoroacetic acid (0.47 mL, 6.1 mmol) was stirred at RT for 10 min. The reaction mixture was concentrated.

[0503] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.14 mL, 0.067 mmol), and DIPEA (0.032 mL, 0.18 mmol) in dichloromethane (0.31 mL) was stirred at RT for 15 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (30 mg, 0.046 mmol, 75% yield) as a light yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.13 (s, 1 H) 7.42 - 7.51 (m, 1 H) 7.17 - 7.24 (m, 2 H) 7.09 - 7.17 (m, 1 H) 6.50 - 6.69 (m, 1 H) 6.40 (br t, *J* = 15.3 Hz, 1 H) 5.76 - 5.86 (m, 1 H) 4.33 - 5.22 (m, 3 H) 3.43 - 4.09 (m, 3 H) 2.60 - 2.74 (m, 1 H) 2.16 - 2.26 (m, 3 H) 1.30 - 1.52 (m, 6 H) 1.23 (d, *J* = 6.8 Hz, 3 H) 1.06 (br d, *J* = 6.4 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.94 - -112.86 (m, 1 F). MS (ESI, +ve) *m/z*: 656.0 $(M + 1)^+$.

Example 198

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-cyclopropyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0504] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-cyclopropyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.102 g, 0.155 mmol; **Intermediate 204**), cyclopropylboronic acid (0.040 g, 0.47 mmol; Oakwood Chemical, Estill, SC), $PdCl_2$(dppf) (0.011 g, 0.016 mmol), and potassium carbonate 2 M aq.; 0.233 mL, 0.466 mmol) in 1,4-dioxane (0.4 mL) was stirred at 100 °C for 1 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–70% [(3:1) EtOAc/EtOH]/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-cyclopropyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (56 mg, 0.085 mmol, 54% yield) as an amber oil. MS (ESI, +ve) *m/z*: 662.2 $(M + 1)^+$.

[0505] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-cyclopropyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-cyclopropyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.058 g, 0.088 mmol) and 2,2,2-trifluoroacetic acid (0.67 mL, 8.8 mmol) in DCM (2 mL) was stirred at RT for 15 min. The reaction mixture was concentrated.

[0506] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.210 mL, 0.105 mmol), and DIPEA (0.061 mL, 0.350 mmol) in dichloromethane (0.876 mL) was stirred at RT for 30 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 40–70% [(3:1) EtOAc/EtOH]/heptane then isocratic 60% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-cyclopropyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (40 mg, 0.065 mmol, 74% yield) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.10 (s, 1 H) 7.41 -

7.49 (m, 1 H) 7.15 - 7.22 (m, 2 H) 7.08 - 7.15 (m, 1 H) 6.51 - 6.71 (m, 1 H) 6.40 (br t, J = 14.8 Hz, 1 H) 5.74 - 5.85 (m, 1 H) 3.44 - 5.20 (m, 6 H) 2.54 - 2.67 (m, 1 H) 2.17 - 2.25 (m, 1 H) 2.16 (s, 3 H) 1.30 - 1.48 (m, 6 H) 1.12 - 1.20 (m, 5 H) 0.98 - 1.05 (m, 5 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.77 - -112.69 (m, 1 F). MS (ESI, +ve) *m/z*: 616.1 $(M + 1)^+$.

Example 199

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-ethyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0507] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-ethyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A mixture of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.100 g, 0.152 mmol; **Intermediate 204**), diethyl zinc (1 M in hexanes, 0.305 mL, 0.305 mmol; Sigma-Aldrich, St. Louis, MO), and tetrakis (0.018 g, 0.015 mmol) in 1,4-dioxane (0.8 mL) was sparged with nitrogen then stirred at 70 °C for 18 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–100% EtOAc/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-ethyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (50 mg, 0.077 mmol, 51% yield) as an amber oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.10 (s, 1 H) 7.39 - 7.47 (m, 1 H) 7.14 - 7.22 (m, 2 H) 7.07 - 7.14 (m, 1 H) 3.46 - 5.06 (m, 6 H) 2.94 (q, J = 7.6 Hz, 2 H) 2.60 - 2.74 (m, 1 H) 2.19 (s, 3 H) 1.51 (s, 9 H) 1.36 (t, J = 7.6 Hz, 3 H) 1.23 - 1.31 (m, 6

H) 1.21 (d, J = 6.6 Hz, 3 H) 1.05 (d, J = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.55 (br s, 1 F). MS (ESI, +ve) *m/z*: 650.2 $(M + 1)^+$.

[0508] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-ethyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-ethyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.050 g, 0.077 mmol) in 2,2,2-trifluoroacetic acid (0.59 mL, 7.7 mmol) was stirred at RT for 10 min. The reaction mixture was concentrated.

[0509] A solution of the resulting oil, acryloyl chloride (0.5 in DCM, 0.17 mL, 0.085 mmol), and DIPEA (0.040 mL, 0.23 mmol) in dichloromethane (0.4 mL) was stirred at RT for 15 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–90% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-ethyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (51 mg, 0.084 mmol, >99% yield) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.11 (s, 1 H) 7.38 - 7.49 (m, 1 H) 7.15 - 7.23 (m, 2 H) 7.07 - 7.15 (m, 1 H) 6.50 - 6.71 (m, 1 H) 6.40 (br t, J = 14.9 Hz, 1 H) 5.75 - 5.86 (m, 1 H) 3.44 - 5.23 (m, 6 H) 2.95 (q, J = 7.7 Hz, 2 H) 2.60 - 2.72 (m, 1 H) 2.20 (s, 3 H) 1.37 (t, J = 7.7 Hz, 3 H) 1.30 - 1.50 (m, 6 H) 1.22 (d, J = 6.8 Hz, 3 H) 1.05 (d, J = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.57 - -112.48 (m, 1 F). MS (ESI, +ve) *m/z*: 604.3 $(M + 1)^+$.

Example 200

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-fluoro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0510] Step 1: 6-Chloro-1-(2-fluoro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A solution of 1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.080 g, 0.16 mmol; **Intermediate 217**) and potassium fluoride, anhydrous (0.046 g, 0.79 mmol) in dimethyl sulfoxide (0.8 mL) was stirred in the microwave at 170 °C for 2 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–60% EtOAc/heptane) to give 6-chloro-1-(2-fluoro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (125 mg, 0.282 mmol) as a white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.73 (br s, 1 H) 8.62 (s, 1 H) 7.42 - 7.50 (m, 1 H) 7.09 - 7.24 (m, 3 H) 2.73 - 2.86 (m, 1 H) 2.31 (s, 3 H) 1.24 (d, *J* = 6.8 Hz, 3 H) 1.08 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -45.06 (s, 1 F) -113.22 (s, 1 F). MS (ESI, +ve) *m/z*: 444.0 $(M + 1)^+$.

[0511] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-fluoro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of 6-chloro-1-(2-fluoro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.095 g, 0.21 mmol), phosphoryl trichloride (0.024 mL, 0.26 mmol), and DIPEA (0.097 mL, 1.1 mmol) in acetonitrile (1.1 mL) was stirred at 80 °C for 30 min.

[0512] The reaction mixture was removed from the heating block, and 1-((2*R*,5*S*)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one, TFA salt (0.265 g, 0.428 mmol; **Intermediate 160**) was added to the reaction mixture; this was stirred at RT for 10 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–90% [(3:1) EtOAc/EtOH]/heptane then 30–50% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-fluoro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (22 mg, 0.037 mmol, 17% yield) as a yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.13 (s, 1 H) 7.42 - 7.50 (m, 1 H) 7.17 - 7.23 (m, 2 H) 7.09 - 7.17 (m, 1 H) 6.50 - 6.71 (m, 1 H) 6.35 - 6.47 (m, 1 H) 5.75 - 5.88 (m, 1 H) 3.43 - 5.21 (m, 6 H) 2.63 - 2.77 (m, 1 H) 2.20 - 2.25 (m, 3 H) 1.32 - 1.51 (m, 6 H) 1.19 - 1.25 (m, 3 H) 1.02 - 1.08 (m, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -46.32 - -46.38 (m, 1 F) -112.78 - 112.87 (m, 1 F). MS (ESI, +ve) *m/z*: 594.0 $(M + 1)^+$.

Example 201

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-(prop-1-en-2-yl)pyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0513] **Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-(prop-1-en-2-yl)pyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.309 g, 0.471 mmol; Intermediate 204), 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (0.133

mL, 0.706 mmol; Combi-Blocks, Inc., San Diego, CA), $PdCl_2$(dppf) (0.034 g, 0.047 mmol), and sodium carbonate (2 M, aq., 0.706 mL, 1.41 mmol) in 1,4-dioxane (2.4 mL) was sparged in nitrogen then was stirred at 80 °C for 2 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–60% EtOH/[heptane/EtOAc (2:1)]) to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-(prop-1-en-2-yl)pyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (336 mg, 0.507 mmol, >99% yield) as an amber oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.11 (s, 1 H) 7.37 - 7.46 (m, 1 H) 7.14 - 7.21 (m, 2 H) 7.11 (t, *J* = 9.2 Hz, 1 H) 6.43 - 6.45 (m, 1 H) 5.47 - 5.49 (m, 1 H) 3.46 - 5.07 (m, 6 H) 2.68 (spt, *J* = 6.7 Hz, 1 H) 2.24 (s, 3 H) 2.22 (s, 3 H) 1.51 (s, 9 H) 1.25 - 1.27 (m, 6 H) 1.20 -1.23 (m, 3 H) 1.06 (d, *J* = 6.8 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.81 (br s, 1 F). MS (ESI, +ve) *m/z*: 662.2 $(M + 1)^+$.

[0514] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-(prop-1-en-2-yl)pyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-(prop-1-en-2-yl)pyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.100 g, 0.151 mmol) in 2,2,2-trifluoroacetic acid (1.16 mL, 15.1 mmol) was stirred at RT for 15 min. The reaction mixture was concentrated.

[0515] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.362 mL, 0.181 mmol), and DIPEA (0.105 mL, 0.604 mmol) in dichloromethane (1.5 mL) was stirred at RT for 30 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–90% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-(prop-1-en-2-yl)pyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (66 mg, 0.11 mmol, 71% yield) as a light yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.11 (s, 1 H) 7.37 - 7.48 (m, 1 H) 7.14 - 7.22 (m, 2 H) 7.05 - 7.14 (m, 1 H) 6.50 - 6.71 (m, 1 H) 6.31 - 6.47 (m, 2 H) 5.75 - 5.86 (m, 1 H) 5.47 - 5.50 (m, 1 H) 4.17 - 5.22 (m, 3 H) 3.44 - 4.08 (m, 3 H) 2.63 - 2.74 (m, 1 H) 2.25 (s, 3 H) 2.22 (s, 3 H)

1.31 - 1.50 (m, 6 H) 1.23 (d, J = 6.6 Hz, 3 H) 1.06 (d, J = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.88 - -112.71 (m, 1 F). MS (ESI, +ve) m/z: 616.1 $(M + 1)^+$.

Example 202

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-vinylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



Intermediate 204
single isomer

single isomer

single isomer

[0516] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-vinylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.170 g, 0.259 mmol; Intermediate 204), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (0.060 g, 0.39 mmol; AstaTech, Inc., Bristol, PA), $PdCl_2$(dppf) (0.019 g, 0.026 mmol), and sodium carbonate (2 M aq., 0.388 mL, 0.777 mmol) in 1,4-dioxane (1.3 mL) was sparged in nitrogen then was stirred at 80 °C for 1 h then at 100 °C for 1 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% EtOH/[heptane/EtOAc (2:1)]) to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-vinylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.131 g, 0.202 mmol, 78% yieldas a light yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.11 (s, 1 H) 7.38 - 7.46 (m, 1 H) 7.13 - 7.21 (m, 2 H) 7.06 - 7.13 (m, 1 H) 6.81 - 6.92 (m, 1 H) 6.64 (dd, J = 17.2, 2.1 Hz, 1 H) 5.64 - 5.71 (m, 1 H) 3.46 - 5.05 (m, 6 H) 2.70 (spt, J = 6.6 Hz, 1 H) 2.22 (s, 3 H) 1.51 (s, 9 H) 1.42 - 1.49

(m, 3 H) 1.29 (d, J = 6.8 Hz, 3 H) 1.21 - 1.24 (m, 3 H) 1.07 (d, J = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.80 (br s, 1 F). MS (ESI, +ve) m/z: 648.2 $(M + 1)^+$.

[0517] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-vinylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-vinylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.131 g, 0.202 mmol) and 2,2,2-trifluoroacetic acid (1.55 mL, 20.2 mmol) in DCM (2 mL) was stirred at RT for 15 min. The reaction mixture was concentrated.

[0518] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.485 mL, 0.243 mmol), and DIPEA (0.141 mL, 0.808 mmol) in dichloromethane (2.0 mL) was stirred at RT for 30 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-vinylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (97 mg, 0.16 mmol, 80% yield) as a light yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.12 (s, 1 H) 7.38 - 7.47 (m, 1 H) 7.14 - 7.22 (m, 2 H) 7.07 - 7.14 (m, 1 H) 6.81 - 6.91 (m, 1 H) 6.51 - 6.70 (m, 2 H) 6.40 (br t, J = 14.7 Hz, 1 H) 5.76 - 5.85 (m, 1 H) 5.66 - 5.72 (m, 1 H) 4.29 - 5.21 (m, 3 H) 3.45 - 4.08 (m, 3 H) 2.63 - 2.76 (m, 1 H) 2.22 (s, 3 H) 1.29 - 1.51 (m, 6 H) 1.23 (d, J = 6.6 Hz, 3 H) 1.07 (br d, J = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.83 - -112.69 (m, 1 F). MS (ESI, +ve) m/z: 602.2 $(M + 1)^+$.

Example 203

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,4-diisopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



Intermediate 204
single isomer

single isomer

single isomer

[0519] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2,4-diisopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.166 g, 0.253 mmol; **Intermediate 204**), isopropylzinc bromide (0.5 M in THF, 0.607 mL, 0.303 mmol; Sigma-Aldrich, St. Louis, MO), and XantPhos Pd G3 (0.013 g, 0.013 mmol) in 1,4-dioxane (1.3 mL) was sparged in nitrogen then was stirred at 50 °C for 1 h. More isopropylzinc bromide (0.5 M in THF, 0.607 mL, 0.303 mmol) was added, and the reaction mixture was then stirred at 80 °C for 2 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% EtOH/[heptane/EtOAc (2:1)]) to give impure *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2,4-diisopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (70 mg, 0.11 mmol, 42% yield) as an off-white solid. MS (ESI, +ve) *m/z*: 664.3 $(M + 1)^+$.

[0520] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,4-diisopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2,4-diisopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-

carboxylate (0.070 g, 0.11 mmol) and 2,2,2-trifluoroacetic acid (0.81 mL, 11 mmol) in DCM (2 mL) was stirred at RT for 15 min. The reaction mixture was concentrated.

[0521] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.25 mL, 0.13 mmol), and DIPEA (0.073 mL, 0.42 mmol) in dichloromethane (1.1 mL) was stirred at RT for 30 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product in methanol was purified via preparative HPLC (Phenomenex Kinetex EVO C_{18} column, 150 × 21.2 mm, Axia packed, 5 μm; 50-70% 0.1% TFA in MeCN/H_2O, 35 mL/min, 13 min); the pure fractions were diluted with EtOAc (75 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 50 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo* to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2,4-diisopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (23 mg, 0.037 mmol, 35% yield) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.10 (s, 1 H) 7.39 - 7.47 (m, 1 H) 7.14 - 7.22 (m, 2 H) 7.07 - 7.14 (m, 1 H) 6.52 - 6.70 (m, 1 H) 6.40 (br t, *J* = 14.9 Hz, 1 H) 5.75 - 5.85 (m, 1 H) 4.25 - 5.25 (m, 3 H) 3.39 - 4.08 (m, 3 H) 3.09 - 3.21 (m, 1 H) 2.58 - 2.71 (m, 1 H) 2.19 (s, 3 H) 1.31 - 1.49 (m, 12 H) 1.21 (d, *J* = 6.6 Hz, 3 H) 1.04 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.30 - -112.15 (m, 1 F). MS (ESI, +ve) *m/z*: 618.0 $(M + 1)^+$.

Example 204

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-ethynyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0522] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-((trimethylsilyl)ethynyl)pyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.233 g, 0.355 mmol; **Intermediate 204**), ethynyltrimethylsilane (0.098 mL, 0.71 mmol; Matrix Scientific, Columbia, SC), DIPEA (0.248 mL, 1.42 mmol), copper(I) iodide (14 mg, 0.071 mmol), and $PdCl_2$(dppf) (0.026 g, 0.035 mmol) in DMF (3.6 mL) was sparged with nitrogen then was stirred at 80 °C for 3 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% [(3:1) EtOAc/EtOH]/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-

fluorophenyl)-1-(4-isopropyl-6-methyl-2-((trimethylsilyl)ethynyl)pyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (188 mg, 0.262 mmol, 74% yield) as an amber oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.08 - 8.13 (m, 1 H) 7.40 - 7.49 (m, 1 H) 7.15 - 7.22 (m, 2 H) 7.08 - 7.15 (m, 1 H) 3.43 - 5.05 (m, 6 H) 2.65 - 2.77 (m, 1 H) 2.22 (s, 3 H) 1.51 (s, 9 H) 1.26 - 1.30 (m, 6 H) 1.23 (d, J = 6.8 Hz, 3 H) 1.07 (d, J = 6.8 Hz, 3 H) 0.29 (s, 8 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.89 (s, 1 F). MS (ESI, +ve) *m/z*: 718.1 $(M + 1)^+$.

[0523] Step 2: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-ethynyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(4-isopropyl-6-methyl-2-((trimethylsilyl)ethynyl)pyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.188 g, 0.262 mmol) and potassium carbonate (7 mg, 0.05 mmol) in methanol (3 mL) was stirred at 22 °C for 3 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% [(3:1) EtOAc/EtOH]/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-ethynyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (82 mg, 0.13 mmol, 49% yield) as an amber oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.12 (s, 1 H) 7.39 - 7.49 (m, 1 H) 7.08 - 7.21 (m, 3 H) 4.91 - 5.06 (m, 1 H) 4.36 - 4.65 (m, 1 H) 4.05 - 4.23 (m, 1 H) 3.77 - 3.98 (m, 2 H) 3.44 - 3.64 (m, 1 H) 3.09 (s, 1 H) 2.66 - 2.77 (m, 1 H) 2.23 (s, 3 H) 1.51 (s, 9 H) 1.20 - 1.32 (m, 9 H) 1.07 (d, J = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.79 (br s, 1 F). MS (ESI, +ve) *m/z*: 646.1 $(M + 1)^+$.

[0524] Step 3: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-ethynyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-ethynyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.082 g, 0.13 mmol) and 2,2,2-trifluoroacetic acid (0.97 mL, 13 mmol) in DCM (2 mL) was stirred at RT for 15 min. The reaction mixture was concentrated.

[0525] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.31 mL, 0.15 mmol), and DIPEA (0.088 mL, 0.51 mmol) in dichloromethane (1.3 mL) was stirred at RT for 30 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-ethynyl-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (59 mg, 0.098 mmol, 77% yield) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.13 (s, 1 H) 7.40 - 7.49 (m, 1 H) 7.08 - 7.23 (m, 3 H) 6.51 - 6.70 (m, 1 H) 6.40 (br t, *J* = 14.7 Hz, 1 H) 5.76 - 5.86 (m, 1 H) 4.33 - 5.21 (m, 3 H) 3.44 - 4.07 (m, 3 H) 3.10 (s, 1 H) 2.65 - 2.78 (m, 1 H) 2.21 - 2.26 (m, 3 H) 1.29 - 1.51 (m, 6 H) 1.24 (d, *J* = 6.6 Hz, 3 H) 1.07 (br d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.92 - -112.59 (m, 1 F). MS (ESI, +ve) *m/z*: 600.0 $(M + 1)^+$.

Example 205

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Intermediate 212

[0526] **Step 1: *tert*-Butyl (2*S*,5*R*)-4-(6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate.** A solution of 6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.439 g, 0.967 mmol, **Intermediate 212**), phosphoryl trichloride (0.108 mL, 1.16 mmol), and DIPEA (0.168 mL, 0.967 mmol) in acetonitrile (5 mL) was stirred under an air condensor and drying tube at 80 °C for 30 min. The reaction mixture was cooled to RT, and *tert*-butyl (2*S*,5*R*)-2,5-dimethylpiperazine-1-carboxylate (0.207 g, 0.967 mmol; eNovation Chemicals LLC, Bridgewater, NJ) was added; the solution was stirred a RT for 30 min. The reaction mixture was diluted with EtOAc (200

mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo* . The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–100% EtOAc/heptane) to give *tert*-butyl (2*S*,5*R*)-4-(6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (258 mg, 0.397 mmol, 41% yield) as a yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.11 (d, *J* = 2.3 Hz, 1 H) 8.13 (d, *J* = 1.9 Hz, 1 H) 7.38 - 7.47 (m, 1 H) 7.05 - 7.18 (m, 3 H) 4.16 - 5.08 (m, 3 H) 3.76 - 4.08 (m, 2 H) 3.45 - 3.65 (m, 1 H) 2.70 - 2.82 (m, 1 H) 2.58 - 2.70 (m, 1 H) 1.52 (s, 9 H) 1.24 - 1.37 (m, 12 H) 1.17 (dd, *J* = 9.1, 6.8 Hz, 3 H) 1.04 (dd, *J* = 6.7, 4.7 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -113.34 - -113.24 (m, 1 F). MS (ESI, +ve) *m/z*: 650.2 $(M + 1)^+$.

[0527] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of *tert*-butyl (2*S*,5*R*)-4-(6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.258 g, 0.397 mmol) and 2,2,2-trifluoroacetic acid (3.04 mL, 39.7 mmol) in DCM (2 mL) was stirred at RT for 15 min. The reaction mixture was concentrated.

[0528] A solution of the resulting oil, acryloyl chloride (0.5 M in DCM, 0.952 mL, 0.476 mmol), and DIPEA (0.276 mL, 1.59 mmol) in dichloromethane (4 mL) was stirred at RT for 10 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–100% [(3:1) EtOAc/EtOH]/heptane) to give 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (145 mg, 0.240 mmol, 61% yield) as an amber oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.11 - 9.13 (m, 1 H) 8.14 (s, 1 H) 7.38 - 7.47 (m, 1 H) 7.06 - 7.19 (m, 3 H) 6.52 - 6.70 (m, 1 H) 6.41 (br t, *J* = 14.9 Hz, 1 H) 5.77 - 5.85 (m, 1 H) 3.80 - 5.21 (m, 6 H) 2.70 - 2.85 (m, 1 H) 2.55 - 2.70 (m, 1 H) 1.23 - 1.53 (m, 15 H) 1.14 - 1.21 (m, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -113.43 - -113.09 (m, 1 F). MS (ESI, +ve) *m/z*: 604.1 $(M + 1)^+$.

Example 205-1

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0529] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.145 g, 0.240 mmol) were separated by chiral SFC: Chirakpak IC, ID 150 x 30 mm, 5 μm, 60% methanol/CO_2, 80 mL/min. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (42.0 mg, 0.070 mmol, 29% yield; first eluting isomer) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.12 (s, 1 H) 8.13 (s, 1 H) 7.38 - 7.47 (m, 1 H) 7.06 - 7.19 (m, 3 H) 6.50 - 6.70 (m, 1 H) 6.41 (br t, *J* = 15.2 Hz, 1 H) 5.77 - 5.88 (m, 1 H) 4.91 - 5.23 (m, 2 H) 4.33 - 4.51 (m, 1 H) 3.68 - 4.11 (m, 3 H) 2.70 - 2.84 (m, 1 H) 2.56 - 2.70 (m, 1 H) 1.40 - 1.52 (m, 5 H) 1.30 - 1.38 (m, 4 H) 1.26 (d, *J* = 7.0 Hz, 3 H) 1.16 (br d, *J* = 6.8 Hz, 3 H) 1.02 - 1.07 (m, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -113.28 (s, 1 F). MS (ESI, +ve) *m/z*: 604.1 $(M + 1)^+$.

Example 205-2

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0530] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.145 g, 0.240 mmol) were separated by chiral SFC: Chirakpak IC, ID 150 x 30 mm, 5 μm, 60% methanol/CO_2, 80 mL/min. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (43 mg, 0.071 mmol, 30% yield; second eluting isomer) as an off-white solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.12 (s, 1 H) 8.14 (s, 1 H) 7.38 - 7.49 (m, 1 H) 7.05 - 7.19 (m, 3 H) 6.51 - 6.70 (m, 1 H) 6.41 (br t, *J* = 14.7 Hz, 1 H) 5.75 - 5.86 (m, 1 H) 4.91 - 5.22 (m, 2 H) 4.37 - 4.52 (m, 1 H) 3.69 - 4.14 (m, 3 H) 2.69 - 2.83 (m, 1 H) 2.57 - 2.69 (m, 1 H) 1.41 - 1.54 (m, 5 H) 1.30 - 1.37 (m, 4 H) 1.22 - 1.28 (m, 3 H) 1.14 - 1.20 (m, 3 H) 1.00 - 1.08 (m, 3 H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -113.30 - -113.20 (m, 1 F). MS (ESI, +ve) *m/z*: 604.1 $(M + 1)^+$.

Examples 206-1 & 206-2

4-(*cis*-4-Acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomers



Step 1: 4-(*cis*-4-Acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A solution of 6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.439 g, 0.967 mmol; **Intermediate 212**), phosphoryl trichloride (0.108 mL, 1.16 mmol), and DIPEA (0.168 mL, 0.967 mmol) in acetonitrile (5 mL) was stirred under an air condensor and drying tube at 80 °C for 30 min. The reaction mixture was cooled to RT and 1-(*cis*-2,6-dimethylpiperazin-1-yl)prop-2-en-1-one (0.644 g, 0.967 mmol; **Intermediate 223**) was added; the solution was stirred a RT for 30 min. The reaction mixture was diluted with EtOAc (200 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 20–100% [(3:1) EtOAc/EtOH]/heptane) to give 4-(*cis*-4-acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (120 mg, 0.199 mmol, 21% yield). MS (ESI, +ve) *m/z*: 604.1 $(M + 1)^+$.

[0531] Step 2: 4-(*cis*-4-Acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomers. Atropisomers of 4-(*cis*-4-acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.120 g, 0.199 mmol) were separated by chiral SFC: Welko SS, ID 250 x 21 mm, 5 μm, 35% methanol/CO_2, 80 mL/min, 90 bar. This gave 4-(*cis*-4-acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (30 mg, 0.050 mmol, 25% yield; first eluting isomer) [^{1}H NMR (400 MHz, $CDCl_3$) δ 9.12 (s, 1 H) 8.35 (s, 1 H) 7.38 - 7.48 (m, 1 H) 7.06 - 7.19 (m, 3 H) 6.64 (dd, *J* = 16.7, 10.5 Hz, 1 H) 6.44 (dd, *J* = 16.7, 2.0 Hz, 1 H) 5.76 - 5.85 (m, 1 H) 4.60 - 4.83 (m, 2 H) 4.34 (br d, *J* = 12.9 Hz, 2 H) 3.69 (dd, *J* = 13.5, 5.0 Hz, 2 H) 2.71 - 2.83 (m, 1 H) 2.58 - 2.71 (m, 1 H) 1.59 (d, *J* = 1.9 Hz, 3 H) 1.57 (d, *J* = 1.9 Hz, 3 H) 1.35 (d, *J* = 6.8 Hz, 3 H) 1.26 (d, *J* = 6.8 Hz, 3 H) 1.17 (d, *J* = 6.8 Hz, 3 H) 1.04 (d, *J* = 7.0 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -113.33 (s, 1 F). MS (ESI, +ve) *m/z*: 604.1 $(M + 1)^+$] as a dark yellow solid and 4-(*cis*-4-acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (30 mg, 0.050 mmol, 25% yield; second eluting isomer) [^{1}H NMR (400 MHz, $CDCl_3$) δ 9.12 (s, 1 H) 8.35 (s, 1 H) 7.39 - 7.47 (m, 1 H) 7.05 - 7.19 (m, 3 H) 6.64 (dd, *J* = 16.7, 10.5 Hz, 1 H) 6.44 (dd, *J* = 16.7, 2.0 Hz, 1 H) 5.78 -

5.84 (m, 1 H) 4.65 - 4.83 (m, 2 H) 4.34 (br d, *J* = 13.7 Hz, 2 H) 3.69 (dd, *J* = 13.4, 4.9 Hz, 2 H) 2.71 - 2.81 (m, 1 H) 2.58 - 2.69 (m, 1 H) 1.59 (d, *J* = 1.9 Hz, 3 H) 1.57 (d, *J* = 1.9 Hz, 3 H) 1.35 (d, *J* = 6.8 Hz, 3 H) 1.26 (d, *J* = 6.8 Hz, 3 H) 1.17 (d, *J* = 6.8 Hz, 3 H) 1.04 (d, *J* = 7.0 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -113.33 (s, 1 F). MS (ESI, +ve) *m/z*: 604.1 $(M + 1)^+$] as a dark yellow solid.

Example 207

(*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-[norcaran-1-yl]pyrido[2,3-*d*]pyrimidin-2-one



[0532] Step 1: (*M*)- *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(cyclohex-1-en-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A yellow heterogeneous mixture of *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 92A**, 0.50 g, 0.89 mmol), cyclohexen-1-ylboronic acid (0.17 g, 1.34 mmol), sodium carbonate, anhydrous, powder (0.28 g, 2.67 mmol), and tetrakis(triphenylphosphine) palladium(0) (0.10 g, 0.09 mmol) in 1,4-dioxane (3.0 mL) and water (1.5 mL) was stirred and heated at 90 °C for 3.5 h. The reaction mixture was allowed to cool to room temperature and evaporated to dryness. The resulting crude product was purified by silica gel chromatography (eluent: 0-100% of EtOAc/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(cyclohex-1-en-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.26 g, 0.43 mmol, 48.8 % yield) as white syrupy solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.44 - 8.48 (m, 1 H), 8.27 (s, 1 H), 7.22 - 7.25 (m, 1 H), 6.33 (t, J=3.7 Hz, 1 H), 4.78 (br s, 1 H), 4.18 - 4.41 (m, 1 H), 3.36 - 4.09 (m, 4 H), 2.55 - 2.68 (m, 1 H), 2.14 (br d, J=1.9 Hz, 2 H), 2.05 (br s, 1 H), 1.96 (br d, J=2.1 Hz, 1 H), 1.85 - 1.91 (m, 3 H), 1.47 - 1.56 (m, 4 H), 1.44 (s, 9 H), 1.31 (d, J=6.4 Hz, 3 H), 1.13 (br d, J=6.4 Hz, 3 H), 1.05 (d, J=6.6 Hz, 3 H), 0.94 - 0.99 (m, 3 H). *m/z* (ESI, +ve ion): 607.3 $(M+H)^+$.

[0533] **Step 2 : (*M*)-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-[norcaran-1-yl]pyrido[2,3-*d*]pyrimidin-2-one**. To a stirred yellow clear solution of trimethylsulfoxonium iodide (0.11 g, 0.51 mmol) in dimethyl sulfoxide (1 mL) was added potassium *tert*-butoxide solution (1 M in THF, 0.51 mL, 0.51 mmol) and the resulting colorless clear mixture was stirred at room temperature. After 40 munites, to the above colorless clear mixture was added a solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(cyclohex-1-en-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.26 g, 0.43 mmol) in tetrahydrofuran (2 mL) and the resulting red clear mixture was stirred at room temperature. After 21.5 hours, the reaction mixture was quenched with sat'd NH4Cl (50 mL) and extracted with EtOAc (2 × 50 mL). The combined organic extracts were dried over MgSO4. The solution was filtered and concentrated *in vacuo* to give the crude material as a yellow syrup. The resulting crude product was purified by silica gel chromatography (eluent: 0-100% of EtOAc-EtOH (3:1)/heptane) to provide *tert*-butyl (2*R*,5*S*)-4-(7-(bicyclo[4.1.0]heptan-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.06 g, 0.10 mmol, 23.70 % yield) as white syrupy solid, which was including triphenylphosphine oxdie and byproducts. *m/z* (ESI, +ve ion): 621.3 $(M+H)^+$. The material was used in the next step without further purification.

[0534] To a colorless clear solution of *tert*-butyl (2*R*,5*S*)-4-(7-(bicyclo[4.1.0]heptan-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.06 g, 0.10 mmol) in dichloromethane (2.0 mL) was added trifluoroacetic acid (2.0 mL) and the resulting yellow clear mixture was stirred at room temperature. After 20 minutes, the mixture was concentrated *in vacuo* to give 7-(bicyclo[4.1.0]heptan-1-yl)-6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as yellow syrup. *m/z* (ESI, +ve ion): 521.2 $(M+H)^+$.

[0535] The above yellow syrup was dissolved in dichloromethane (2.0 mL) at 0 °C and treated with DIPEA (0.25 mL, 1.46 mmol), followed by acryloyl chloride (9.49 µl, 0.12 mmol) dropwise. The resulting mixture was stirred at 0 °C for 10 minutes and diluted with EtOAc (50 mL) and sat'd NH4Cl (50 mL). The aqueous layer was extracted with EtOAc (2 × 50 mL). The combined organic extracts were washed with satd NaHCO3 (1 × 50 mL) and dried over Na2SO4. The solution was filtered and concentrated *in vacuo* to give the crude material as a yellow syrup. The resulting crude product was purified by silica gel chromatography (eluent:

0-100% of EtOAc-EtOH (3:1)/heptane) to provide a light-yellow syrupy solid. The above solid was further purified by reverse-phase preparative HPLC (using a Phenimenex Gemini column, 5 μm, C18, 110 Å, AXIA, 150 × 30 mm, 0.1% TFA in $CH_3CN/H2O$, gradient 10% to 90% over 15 min). The pure fractions were combined, neutralized with sat'd NaHCO3 (50 mL), and extracted with DCM (2 × 50 mL). The combined organic extracts were dried over Na2SO4. The solution was filtered and concentrated *in vacuo* to give (*M*)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-[norcaran-1-yl]pyrido[2,3-*d*]pyrimidin-2-one (0.01 g, 7.65 μmol, 15.7 % yield) as yellow solid: ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.47 (d, *J*=4.8 Hz, 1 H), 8.22 - 8.29 (m, 1 H), 7.22 - 7.27 (m, 1 H), 6.75 - 6.89 (m, 1 H), 6.18 (dd, *J*=16.7, 2.2 Hz, 1 H), 5.70 - 5.77 (m, 1 H), 4.39 - 4.88 (m, 2 H), 3.44 - 4.17 (m, 4 H), 2.57 - 2.65 (m, 1 H), 2.18 (t, *J*=7.4 Hz, 1 H), 1.83 - 1.93 (m, 4 H), 1.70 - 1.81 (m, 2 H), 1.57 - 1.66 (m, 1 H), 1.44 - 1.51 (m, 1 H), 1.28 (br dd, *J*=9.5, 2.7 Hz, 5 H), 1.10 - 1.20 (m, 4 H), 1.06 (dd, *J*=12.2, 6.8 Hz, 3 H), 0.95 (br dd, *J*=17.8, 6.6 Hz, 3 H), 0.70 - 0.77 (m, 1 H), 0.52 (t, *J*=5.4 Hz, 1 H). *m/z* (ESI, +ve ion): 574.6 $(M+H)^+$.

Example 208

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1H)-one



[0536] Step 1: 6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a stirring solution of 6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (700 mg, 1.483 mmol, **Method 103 using Intermediate 173**) in DCM (12 mL) was added *m*-chloroperoxybenzoic acid (77wt%, 512 mg, 2.22 mmol) at 20 °C. After 15 min, the reaction was then washed with sat. NaHCO3 (2 mL), dried over MgSO4, then purified by silica gel chromatography (4 g) eluting product with a gradient of 15>40% EtOAc/EtOH (3:1 blend)/heptane to afford 6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione as white solid. ^{1}H NMR (400 MHz, *CDCl*$_3$) δ 8.58 (s, 1H), 8.57 (br s, 1H), 8.01 (d, *J*=2.28 Hz, 1H), 7.85 (dd, *J*=1.87, 8.29 Hz, 1H), 7.39-7.46 (m, 1H), 7.36 (d, *J*=8.29 Hz, 1H), 7.06-7.19 (m, 3H), 3.09-3.11 (m, 3H), 2.81 (spt, *J*=41.90 Hz, 1H), 1.24 (d, *J*=6.84 Hz, 3H), 1.09 (d, *J*=6.84 Hz, 3H). *m/z* (ESI, +ve ion): 488.0 $(M+H)^+$.

[0537] Step 2: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-(methylsulfonyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of 6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidine-2,4(1H,3H)-dione (660 mg, 1.353 mmol)in MeCN (6 mL) at 20 °C was added DIEA (354 µl, 2.029 mmol) followed by phosphorus(V) oxide chloride (189 µl, 2.029 mmol). After 45 min, the reaction was concentrated under reduced pressure, then azeotroped once with dry toluene (10 mL). The residue was dissolved in THF (6 mL), chilled to 0 °C, DIEA added (1 mL) followed by (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (435 µl, 2.029 mmol). After 10 min, the reaction was partitioned between EtOAc (30 mL) and sat. NaHCO3 (15 mL). The organic was dried over MgSO4, concentrated under reduced pressure, then purified via silica gel chromatography (40 g) eluting product with a gradient of 5>50% EtOAc/EtOH (3:1 blend) to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-(methylsulfonyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as white solid. *m/z* (ESI, +ve ion): 684.0 $(M+H)^+$.

[0538] Step 3: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 208). A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-(methylsulfonyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (400 mg, 0.585 mmol) was stirred in TFA for 10 min at

RT. The reaction was concentrated under reduced pressure and azeotroped once with dry toluene (10 mL). The residue was then dissolved in DCM (5 mL), chilled to 0 °C under nitrogen, and DIEA (1 mL) added. To the reaction was added a solution of 2-propenoyl chloride (52.2 µl, 0.643 mmol) in DCM (1 mL). After 5 min, the reaction was partitioned between DCM (5 mL) and sat. NaHCO3 (10 mL). The aqueous was further extracted with DCM (2 mL). The combined organics were then dried over MgSO4, concentrated under reduced pressure, then purified via silica gel chromatography (40 g) eluting products with a gradient of 5>80% EtOAc/EtOH (3:1 blend) to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 208**) as white solid. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.10 (s, 1H), 7.96-8.01 (m, 1H), 7.80-7.88 (m, 1H), 7.38-7.47 (m, 1H), 7.30-7.37 (m, 1H), 7.07-7.20 (m, 3H), 6.50-6.70 (m, 1H), 6.33-6.46 (m, 1H), 5.75-5.84 (m, 1H), 4.86-5.25 (m, 2H), 4.26-4.53 (m, 1H), 3.36-4.06 (m, 3H), 3.08-3.12 (m, 3H), 2.61-2.80 (m, 1H), 1.30-1.54 (m, 6H), 1.20-1.29 (m, 6H). *m/z* (ESI, +ve ion): 638.2 $(M+H)^+$.

Example 208-1

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



Example 208 Single Isomer

[0539] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1H)-one (**Example 208,** 200 mg, 0.157 mmol) were separated by chiral SFC: ID 150 x 21 mm, 5 µm, 60% MeOH/CO_2, 80 g/min, 102 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 208-1**), first eluting isomer, as white fluffy solid from lyophilization. ^{1}H NMR (400 MHz, *DMSO-d6*) δ 8.43 (d, *J*=6.84 Hz, 1H), 7.94 (d, *J*=1.87 Hz,

1H), 7.84 (dd, *J*=1.87, 8.29 Hz, 1H), 7.48-7.56 (m, 2H), 7.21-7.34 (m, 3H), 6.77-6.92 (m, 1H), 6.20 (d, *J*=15.76 Hz, 1H), 5.72-5.80 (m, 1H), 4.44-4.92 (m, 2H), 3.73-4.31 (m, 4H), 3.31 (s, 3H), 2.64-2.77 (m, 1H), 1.36 (d, *J*=6.43 Hz, 3H), 1.24 (dd, *J*=6.63, 26.54 Hz, 3H), 1.14 (d, *J*=6.84 Hz, 3H), 1.04 (d, *J*=6.84 Hz, 3H). ^{19}F NMR (376 MHz, DMSO-d_6) δ -113.55 (s, 1F), -113.57 (s, 1F). MS (ESI, +ve) *m/z*: 638.2 $(M + 1)^+$.

Example 208-2

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



Example 208 **Single Isomer**

[0540] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one **(Example 208,** 200 mg, 0.157 mmol) were separated by chiral SFC: ID 150 x 21 mm, 5 µm, 60% MeOH/CO_2, 80 g/min, 102 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one **(Example 208-2)**, second eluting isomer, as a white fluffy solid from lyophilization. ^{1}H NMR (400 MHz, *DMSO-d_6*) δ 8.47 (d, *J*=1.66 Hz, 1H), 7.94 (d, *J*=1.66 Hz, 1H), 7.82 (dd, *J*=2.28, 8.29 Hz, 1H), 7.46-7.54 (m, 2H), 7.18-7.34 (m, 3H), 6.75-6.92 (m, 1H), 6.19 (d, *J*=15.34 Hz, 1H), 5.71-5.79 (m, 1H), 4.44-4.96 (m, 2H), 3.46-4.22 (m, 4H), 3.30 (s, 3H), 2.58-2.77 (m, 1H), 1.32 (t, *J*=14.30 Hz, 3H), 1.23 (dd, *J*=7.05, 26.95 Hz, 3H), 1.12 (d, *J*=6.84 Hz, 3H), 1.04 (d, *J*=6.63 Hz, 3H). ^{19}F NMR (376 MHz, *DMSO-d_6*) δ -113.74 (s, 1F), -113.75 (s, 1F). MS (ESI, +ve) *m/z*: 638.2 $(M + 1)^+$.

Example 209-1 and Example 209-2

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, 2 single isomers



[0541] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one **(Example 103-3)** were separated by chiral SFC: Step 1 chiral SFC AD-H, 30 x 250 mm, 5 μm, 20% IPA/CO_2, 150 g/min, 178 bar. Step 2: chiral SFC Welko (*R,R*) 21 x 250 mm, 5 μm, 60% MeOH/CO_2, 64 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one **(Example 209-1)**, single isomer, this was the first eluting peak of step 2, and was isolated as a white fluffy solid from lyophilization. ^{1}H NMR (400 MHz, *CDCl$_3$*) δ 8.01-8.18 (m, 1H), 7.61-7.68 (m, 1H), 7.58-7.61 (m, 1H), 7.29-7.48 (m, 2H), 7.13-7.20 (m, 2H), 7.09 (t, *J*=18.50 Hz, 1H), 6.50-6.72 (m, 1H), 6.39 (t, *J*=30.30 Hz, 1H), 5.80 (dd, *J*=8.09, 16.59 Hz, 1H), 4.42-5.27 (m, 2H), 3.42-4.42 (m, 4H), 2.75-2.82 (m, 3H), 2.60-2.72 (m, 1H), 1.19-1.45 (m, 9H), 1.01-1.10 (m, 3H). ^{19}F NMR (376 MHz, *CDCl$_3$*) δ -112.42 (s, 1F), -112.45 (s, 1F). MS (ESI, +ve) *m/z*: 622.3 $(M + 1)^+$. The separation also provided 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one **(Example 209-2)**, single isomer, this was the second eluting peak of step 2, and isolated as white fluffy solid from lyophilization. ^{1}H NMR (400 MHz, *CDCl$_3$*) δ 8.09 (br s, 1H), 7.54-7.90 (m, 2H), 7.30-7.48 (m, 2H), 7.14-7.20 (m, 2H), 7.10 (t, *J*=18.00 Hz, 1H), 6.49-6.71 (m, 1H), 6.33-6.47 (m, 1H), 5.72-5.89 (m, 1H), 3.35-5.21 (m, 6H), 2.78 (s, 3H), 2.63-2.75 (m, 1H), 1.17-1.48 (m, 9H), 1.01-1.09 (m, 3H). ^{19}F NMR (376 MHz, *CDCl$_3$*) δ -112.34 (s, 1F), -112.35 (s, 1F), -112.39 (s, 1F). MS (ESI, +ve) *m/z*: 622.2 $(M + 1)^+$.

Example 209-3

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



Example 103-3 Single Isomer

[0542] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 103-3**) were separated by chiral SFC: AD-H, 30 x 250 mm, 5 μm, 20% IPA/CO_2, 150 g/min, 178 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 209-3**), single isomer, this was the third eluting peak (peak 1 and peak 2 overlap), and was isolated as a white fluffy solid from lyophilization. ^{1}H NMR (400 MHz, *CDCl*$_3$) δ 8.08 (s, 1H), 7.88 (d, *J*=1.87 Hz, 1H), 7.38-7.47 (m, 1H), 7.35 (dd, *J*=1.87, 7.88 Hz, 1H), 7.21-7.25 (m, 1H), 7.13-7.18 (m, 2H), 7.09 (t, *J*=18.70 Hz, 1H), 6.51-6.70 (m, 1H), 6.40 (t, *J*=30.10 Hz, 1H), 5.80 (dd, *J*=8.09, 17.83 Hz, 1H), 3.44-5.26 (m, 6H), 2.78 (s, 3H), 2.59-2.71 (m, 1H), 1.30-1.48 (m, 6H), 1.25 (d, *J*=6.63 Hz, 3H), 1.07 (d, *J*=6.63 Hz, 3H) [33 H]. ^{19}F NMR (376 MHz, *CDCl*$_3$) δ -112.37 (s, 1F), -112.41 (s, 1F), -112.43 (s, 1F). MS (ESI, +ve) *m/z*: 622.3 $(M + 1)^+$.

Example 209-4

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0543] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 103-3**) were separated by chiral SFC: AD-H, 30 x 250 mm, 5 μm, 20% IPA/CO_2, 150 g/min, 178 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 209-4**), single isomer. This was the forth eluting peak (peak 1 and peak 2 overlap), and was isolated as a white fluffy solid from lyophilization. ^{1}H NMR (400 MHz, *CDCl*$_3$) δ 8.09 (s, 1H), 7.86 (s, 1H), 7.33-7.47 (m, 2H), 7.21-7.25 (m, 1H), 7.14-7.19 (m, 2H), 7.10 (t, *J*=18.70 Hz, 1H), 6.49-6.74 (m, 1H), 6.40 (t, *J*=32.30 Hz, 1H), 5.80 (t, *J*=17.60 Hz, 1H), 3.38-5.19 (m, 6H), 2.78 (s, 3H), 2.71 (br s, 1H), 1.18-1.54 (m, 9H), 1.05 (d, *J*=7.46 Hz, 3H) [33 H]. ^{19}F NMR (376 MHz, *CDCl*$_3$) δ -112.34 (s, 1F), -112.39 (s, 1F). MS (ESI, +ve) *m/z*: 622.3 $(M + 1)^+$.

Example 210

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one.



Intermediate 92A Step 1



Example 210

[0544] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfonyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A suspension of *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (250 mg, 0.445 mmol, **Intermediate 92A**), 2-(methylsulfonyl)phenyl)boronic acid (Sigma-Aldrich, St. Louis, MO, USA, 134 mg, 0.668 mmol), tetrakis(triphenylphosphine)palladium(0) (51.5 mg, 0.045 mmol), sodium carbonate (142 mg, 1.336 mmol) in 1,4-dioxane (2 mL) and water (1 mL) was sparged with argon for 1 min then heated to 95 °C for 40 min. The reaction was then extracted with EtOAc (20 mL), dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (24 g) eluting products with a gradient of 5>60% EtOAc/EtOH (3:1 blend)/heptane to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfonyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as colorless solid. *m/z* (ESI, +ve ion): 681.2 $(M+H)^+$.

[0545] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-

2(1*H*)-one (Example 210). A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfonyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (135 mg, 0.198 mmol) was stirred in TFA (2 mL) for 15 min at 20 °C. The reaction was then concentrated under reduced pressure, and the residue was dissolved in DCM (5 mL) with DIEA (0.25 mL) added. The solution was chilled to 0 °C and acryloyl chloride (19.3 µl, 0.238 mmol) added. The cooling bath was removed and the solution stirred at 20 °C for 1 hr. The reaction was then washed with sat. NaHCO3 (5 mL). The organic was dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (4 g) eluting products with a gradient of 15>100% EtOAc/EtOH (3:1 blend)/heptane to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 210**) as white solid. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.45 (d, *J*=4.77 Hz, 1H), 8.14 (s, 1H), 8.01-8.07 (m, 1H), 7.60-7.73 (m, 2H), 7.20-7.51 (m, 1H), 7.04 (d, *J*=4.98 Hz, 1H), 6.31-6.71 (m, 2H), 5.75-5.86 (m, 1H), 3.51-5.23 (m, 6H), 2.57-2.72 (m, 1H), 2.47-2.54 (m, 3H), 1.90-2.17 (m, 3H), 0.92-1.55 (m, 12H). *m/z* (ESI, +ve ion): 635.2 $(M+H)^+$.

Example 211-1

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0546] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylthio)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A suspension of *tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (350 mg, 0.623 mmol, **Intermediate 92A**), (2-fluoro-6-(methylthio)phenyl)boronic acid (Aurum Pharmatech LLC, Franklin Park, NJ, USA, 128 mg, 0.686 mmol), tetrakis(triphenylphosphine)palladium(0) (72.0 mg, 0.062 mmol), sodium carbonate (198 mg, 1.87 mmol) in 1,4-dioxane (4 mL) and water (2 mL) was sparged with argon for 1 min then heated to 95 °C for 60 min. The reaction was then extracted with EtOAc (20 mL), dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (24 g) eluting products with a gradient of 5>60% EtOAc/EtOH (3:1 blend)/heptane to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylthio)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as white solid. ^{1}H NMR (400 MHz, *CDCl*$_3$) δ 8.47 (d, *J*=3.73 Hz, 1H), 8.15 (d, *J*=1.87 Hz, 1H), 7.30-7.39 (m, 1H), 7.04-7.11 (m, 2H), 6.90 (q, *J*=24.70 Hz, 1H), 4.16-5.05 (m, 3H), 3.44-3.96 (m, 4H), 2.63-2.92 (m, 1H), 2.32 (d, *J*=14.51 Hz, 3H), 2.02 (d, *J*=35.66 Hz, 3H), 1.51 (s, 9H), 1.28-1.50 (m, 8H), 1.01-1.16 (m, 3H). ^{19}F NMR (376 MHz, *CDCl*$_3$) δ -114.43 (s, 1F), -115.25 (s, 1F). *m/z* (ESI, +ve ion): 667.3 $(M+H)^+$.

[0547] Step 2: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylthio)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (210 mg, 0.315 mmol) in DCM (5 mL) at 20 °C was added dropwise a solution of m-chloroperoxobenzoic acid (77wt%, 78 mg, 0.346 mmol) in DCM (1 mL). The reaction was then washed with 5% Na2CO3 (5 mL). The aqueous was then further extracted with DCM. The combined organics were then dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (12 g) eluting products with a gradient of 15>70% EtOAc/EtOH (3:1 blend)/heptane to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as colorless film. *m/z* (ESI, +ve ion): 683.0 $(M+H)^+$.

[0548] **Step 3: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 211-1).** A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (170 mg, 0.249 mmol) in TFA was stirred for 15 min at 20 °C. The reaction was then concentrated under reduced pressure, and the residue was dissolved in DCM (5 mL), DIEA (1 mL) followed by the addition of acryloyl chloride (30.3 µl, 0.373 mmol) at 20 °C. The solution was stirred at RT for 1 hr, then washed with sat. Na_2CO_3 (5 mL). The separated aqueous was further extracted with DCM (10 mL). The organic was dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (4 g) eluting products with a gradient of 15>100% EtOAc/EtOH (3:1 blend)/heptane to afford (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 211-1**) as white solid. ^{1}H NMR (400 MHz, *CDCl*$_3$) δ 8.47-8.61 (m, 1H), 8.11-8.23 (m, 1H), 7.46-7.95 (m, 2H), 7.28-7.32 (m, 1H), 7.06-7.20 (m, 1H), 6.33-6.69 (m, 2H), 5.75-5.98 (m, 1H), 3.67-5.19 (m, 6H), 2.50-2.96 (m, 1H), 1.90-2.19 (m, 6H), 1.13-1.47 (m, 12H). ^{19}F NMR (376 MHz, *CDCl*$_3$) δ -112.28 (d, *J*=13.88 Hz, 1F), -112.63 (d, *J*=20.81 Hz, 1F). *m/z* (ESI, +ve ion): 637.1 $(M+H)^+$.

Example 211-2

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0549] Atropisomers of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 211-1**) were separated by chiral SFC: OD, 21 x 150 mm, 5 µm, 15% EtOH/CO_2, 80 g/min, 102 bar. This gave (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-

methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 211-2**), single isomer, first eluting peak, and isolated as white fluffy powder from loyphilization. ^{1}H NMR (600 MHz, *CDCl3*) δ 8.48 (d, *J*=5.20 Hz, 1H), 8.18 (s, 1H), 7.91 (d, *J*=7.94 Hz, 1H), 7.67-7.73 (m, 1H), 7.29 (t, *J*=16.70 Hz, 1H), 7.13 (br s, 1H), 6.53-6.69 (m, 1H), 6.37-6.45 (m, 1H), 5.78-5.85 (m, 1H), 3.48-5.18 (m, 6H), 2.55-2.65 (m, 1H), 2.10-2.15 (m, 6H), 1.13-1.53 (m, 12H). *m/z* (ESI, +ve ion): 637.3 $(M+H)^+$.

Example 211-3

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0550] Atropisomers of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 211-1**) were separated by chiral SFC: OD, 21 x 150 mm, 5 μm, 15% EtOH/CO_2, 80 g/min, 102 bar. This gave (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfinyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 211-3**), single isomer, second eluting peak, and isolated as white fluffy powder from loyphilization. ^{1}H NMR (600 MHz, *CDCl3*) δ 8.47 (d, *J*=4.65 Hz, 1H), 8.10-8.22 (m, 1H), 7.44-7.97 (m, 2H), 7.22-7.33 (m, 2H), 7.00-7.14 (m, 1H), 6.50-6.71 (m, 1H), 6.33-6.46 (m, 1H), 5.73-5.87 (m, 1H), 3.37-5.18 (m, 6H), 2.61-2.90 (m, 1H), 1.91-2.11 (m, 6H), 1.15-1.58 (m, 12H). *m/z* (ESI, +ve ion): 637.3 $(M+H)^+$.

Example 212-1

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0551] Step 1: (*M*)-*tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylthio)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A suspension of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 92A**, 1.0 g, 1.78 mmol), 2-(methylthio)phenylboronic acid (Sigma-Aldrich Corp., St. Louis, MO, USA, 0.449 g, 2.67 mmol), tetrakis(triphenylphosphine)palladium(0) (0.103 g, 0.089 mmol), sodium carbonate (0.224 ml, 5.34 mmol) in 1,4-dioxane (8 mL) and water (4 mL) was sparged with argon for 1 min then heated to 95 °C for 30 min. The reaction was then extracted with EtOAc (60 mL), dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (40 g) eluting products with a gradient of 5>60% EtOAc/EtOH (3:1 blend)/heptane to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylthio)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as yellow tar. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.47 (br s, 1H), 8.12 (s, 1H), 7.31-7.73 (m, 3H), 7.19 (ddd, *J*=1.45, 7.46, 14.51 Hz, 1H), 7.03 (dd, *J*=0.83, 7.26 Hz, 1H), 3.42-5.02 (m, 6H), 2.66-2.86 (m, 1H), 2.28-2.32 (m, 3H), 2.01-2.10 (m, 3H), 1.49-

1.55 (m, 9H), 1.43-1.49 (m, 3H), 1.21-1.35 (m, 6H), 0.99-1.15 (m, 3H). *m/z* (ESI, +ve ion): 649.1 $(M+H)^+$.

[0552] Step 2: (*M*)-*tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (Intermediate 212-1b). To a stirring solution of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylthio)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (1.0 g, 1.54 mmol) in DCM (10 mL) at 20 °C was added dropwise a solution of *m*-chloroperoxobenzoic acid (77wt%, 0.345 g, 1.54 mmol) in DCM (5 mL). After 5 min, the reaction was then washed with 5% $Na2CO_3$ (10 mL). The aqueous was then further extracted with DCM. The combined organics were then dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (40 g) eluting products with a gradient of 15>70% EtOAc/EtOH (3:1 blend)/heptane to give (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as colorless film. ^1H NMR (400 MHz, $CDCl_3$) δ 8.48 (d, *J*=4.56 Hz, 1H), 8.16-8.20 (m, 1H), 8.03-8.14 (m, 1H), 7.63-7.72 (m, 1H), 7.54 (dd, *J*=7.05, 15.13 Hz, 1H), 7.37-7.48 (m, 1H), 7.00-7.20 (m, 1H), 3.76-5.06 (m, 6H), 2.49-2.91 (m, 1H), 1.94-2.17 (m, 7H), 1.51-1.53 (m, 9H), 1.27-1.50 (m, 10H), 1.13-1.21 (m, 3H). *m/z* (ESI, +ve ion): 665.3 $(M+H)^+$.

[0553] Step 3: (*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 212-1). A solution of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.65 g, 0.977 mmol) in TFA (2 mL) was stirred for 15 min at 20 °C. The reaction was then concentrated under reduced pressure, and the residue was dissolved in DCM (5 mL) with DIEA (1 mL) added. The solution was chilled to 15 °C and acryloyl chloride (0.119 ml, 1.46 mmol) added. The cooling bath was removed and the solution stirred at 20 °C for 15 min. The reaction was then washed with sat. Na_2CO_3 (5 mL). The separated aqeous was futher extracted with DCM (10 mL). The organic was dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (40 g) eluting products with a gradient of 15>100% EtOAc/EtOH (3:1 blend)/heptane to give as colorless film. The film was then dissolved in 1:1 MeCN/water (2 mL), frozen, then lyophilized to afford (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-

yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 212-1**) as white fluffy solid. *m/z* (ESI, +ve ion): 619.3 $(M+H)^+$.

Example 212-2

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0554] Atropisomers of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 212-1**) were separated by chiral SFC: OX, 21 x 250 mm, 5 μm, 45% $MeOH/CO_2$, 80 g/min, 102 bar. This gave (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1H)-one (**Example 212-2**), first eluting peak, isolated as a white fluffy powder from loyphilization. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.53 (d, *J*=3.11 Hz, 1H), 8.39 (d, *J*=4.98 Hz, 1H), 8.00 (d, *J*=7.88 Hz, 1H), 7.71-7.79 (m, 1H), 7.61-7.68 (m, 1H), 7.47 (d, *J*=7.67 Hz, 1H), 7.26 (d, *J*=4.98 Hz, 1H), 6.77-6.92 (m, 1H), 6.20 (dd, *J*=1.87, 16.59 Hz, 1H), 5.72-5.81 (m, 1H), 4.82-4.98 (m, 1H), 4.41-4.81 (m, 1H), 3.48-4.27 (m, 4H), 2.59-2.71 (m, 1H), 2.07 (s, 3H), 2.01 (s, 3H), 1.35 (t, *J*=12.20 Hz, 3H), 1.22 (dd, *J*=6.43, 27.78 Hz, 3H), 1.02 (d, *J*=6.63 Hz, 3H), 0.83 (d, *J*=5.60 Hz, 3H). *m/z* (ESI, +ve ion): 619.1 $(M+H)^+$.

Example 212-3

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0555] Atropisomers of of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(2-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 212-1**) were separated by chiral SFC: OX, 21 x 250 mm, 5 μm, 45% MeOH/CO_2, 80 g/min, 102 bar. This gave (*M*)-6-chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)-7-(2-methylsulfinyl)phenyl]pyrido[2,3-*d*]pyrimidin-2-one (**Example 212-3**), second eluting peak, isolated as white fluffy powder from loyphilization. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.46 (d, *J*=7.88 Hz, 1H), 8.41 (d, *J*=3.94 Hz, 1H), 7.97 (d, *J*=8.29 Hz, 1H), 7.71-7.78 (m, 1H), 7.61-7.68 (m, 1H), 7.47 (d, *J*=7.26 Hz, 1H), 7.22 (t, *J*=9.30 Hz, 1H), 6.76-6.93 (m, 1H), 6.22 (dd, *J*=1.87, 16.59 Hz, 1H), 5.72-5.82 (m, 1H), 3.72-4.96 (m, 6H), 2.75-2.88 (m, 1H), 2.07 (s, 3H), 1.91 (s, 3H), 1.35-1.42 (m, 3H), 1.26 (dd, *J*=6.43, 23.84 Hz, 3H), 1.10 (d, *J*=6.63 Hz, 3H), 1.03 (dd, *J*=2.07, 6.43 Hz, 3H). *m/z* (ESI, +ve ion): 619.1 $(M+H)^+$.

Example 213

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-d]pyrimidin-2(1*H*)-one



[0556] **Step 1: 4-Iodo-2-isopropyl-6-methylaniline.** To a biphasic solution of 2-isopropyl-6-methylaniline (Enamine, Monmouth Jct., NJ, USA, 2.6 g, 17.42 mmol) and sodium bicarbonate, powder (2.93 g, 34.8 mmol) in DCM (20 mL) and water (20 mL) at 20 °C was added iodine (4.42 g, 17.4 mmol). After 10 min, the organic was pushed through a plug of silica (40 g) eluting products with 20% EtOH/EtOAc (1:3 blend)/heptane to afford 4-iodo-2-isopropyl-6-methylaniline (4.26 g, 15.5 mmol, 89 % yield) as an amber oil. ^{1}H NMR (400 MHz, *CDCl*$_3$) δ 7.28 (d, *J*=1.66 Hz, 1H), 7.24 (d, *J*=1.87 Hz, 1H), 3.40-3.99 (bs, 2H), 2.83 (spt, *J*=40.80 Hz, 1H), 2.13 (s, 3H), 1.24 (d, *J*=6.84 Hz, 6H). *m/z* (ESI, +ve ion): 276.1 $(M+H)^+$.

[0557] **Step 2: 2-Isopropyl-6-methyl-4-(methylthio)aniline.** To a stirring solution of 4-iodo-2-isopropyl-6-methylaniline (4.2 g, 15.3 mmol) in THF (15 mL) at 0 °C under nitrogen

was added isopropylmagnesium chloride, 2.0 M in THF (16.0 ml, 32.1 mmol) dropwise at a rate not to exceed an internal temp of 10 °C. The solution was stirred for 5 min, cooled to -70 °C, then n-butyllithium, 2.5 M in hexanes (10.7 ml, 26.7 mmol) added dropwise at a rate not to exceed an internal temp of -60 °C. The solution was stirred for 10 min, then dimethyl disulfide (2.77 ml, 30.5 mmol) added. The cooling bath was removed and reaction quenched with sat. NH_4Cl (20 mL) at -20 °C. The suspension was then partitioned between EtOAc (120 mL) and water (20 mL). The organic was further extracted with EtOAc (20 mL). The combined organics were then dried over MgSO4, filtered, then concentrated under reduced pressure to afford 2-isopropyl-6-methyl-4-(methylthio)aniline (3.6 g) as tan oil. *m/z* (ESI, +ve ion): 196.3 $(M+H)^+$.

[0558] Step 3: 2-Isopropyl-6-methyl-4-(methylsulfinyl)aniline. To a stirring solution of crude (50%) 2-isopropyl-6-methyl-4-(methylthio)aniline (3.6 g, 18.4 mmol) in DCM (30 mL) at 20 °C was added *m*-chloroperoxybenzoic acid (77wt%, 2.06 g, 9.22 mmol). After 5 min, the reaction was washed with 0.5 M NaOH (20 mL), and the organic directly purified by silica gel chromatography (40 g) eluting products with a gradient of 0>50% EtOAc/EtOH (3:1 blend)/heptane to give 2-isopropyl-6-methyl-4-(methylsulfinyl)aniline (1.0 g, 4.73 mmol, 25.7 % yield) as tan oil. ^1H NMR (400 MHz, *CDCl3*) δ 7.33 (d, *J*=1.66 Hz, 1H), 7.25 (d, *J*=1.45 Hz, 1H), 3.95 (br s, 2H), 2.85-2.98 (m, 1H), 2.69 (s, 3H), 2.23 (s, 3H), 1.27-1.31 (m, 6H). *m/z* (ESI, +ve ion): 212.3 $(M+H)^+$.

[0559] Step 4: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (Intermediate 213a). To a diluted solution of phosgene solution, 15% in toluene (3.34 ml, 4.67 mmol) in THF (10 mL) at 0 °C was added a second solution of *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 165**, 1.5 g, 3.12 mmol) and DIEA (2.17 ml, 12.5 mmol) in THF (12 mL). After 5 min, a solution of 2-isopropyl-6-methyl-4-(methylsulfinyl)aniline (0.988 g, 4.67 mmol) in THF (8 mL) was added. The reaction was then partitioned between EtOAc (80 mL) and sat. $NaHCO_3$ (20 mL). The organic was washed with sat. NaCl (10 mL), dried over $MgSO_4$, then concentrated under reduced pressure. The residue was then dissolved in toluene (15 mL). The solution was chilled to 10 °C under nitrogen and then sodium *tert*-butoxide (0.374 g, 3.89 mmol) added. After 20 min, the reaction was then partitioned between EtOAc (60 mL) and sat. $NaHCO_3$ (20 mL). The organic was washed with sat. NaCl (10 mL), dried over $MgSO_4$, concentrated under reduced

pressure, then purified by silica gel chromatography (40 g) eluting products with a gradient of 0>70% EtOAc/EtOH (3:1 blend)/heptane to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 213a**) as yellow foam. ^{1}H NMR (400 MHz, *CDCl*$_3$) δ 8.10 (s, 1H), 7.48-7.69 (m, 1H), 7.06-7.46 (m, 5H), 4.23-5.08 (m, 2H), 3.43-4.07 (m, 4H), 2.77 (s, 3H), 2.53-2.69 (m, 1H), 2.05-2.10 (m, 3H), 1.50-1.55 (m, 9H), 1.40-1.49 (m, 3H), 0.98-1.27 (m, 9H). ^{19}F NMR (376 MHz, *CDCl*$_3$) δ -112.48 (s, 1F), -112.57 (s, 1F), -112.62 (s, 1F). *m/z* (ESI, +ve ion): 682.1 $(M+H)^+$.

[0560] Step 5: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 213). A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 213a**, 390 mg, 0.572 mmol) in TFA (2 mL) was stirred for 15 min at 20 °C. The reaction was then concentrated under reduced pressure, and the residue was dissolved in DCM (5 mL) with DIEA (1 mL) added. To the solution was added acryloyl chloride (69.7 µl, 0.857 mmol). After 15 min, the reaction was then washed with sat. Na_2CO_3 (5 mL). The separated aqeous was futher extracted with DCM (10 mL). The combined organics were then dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (40 g) eluting products with a gradient of 15>100% EtOAc/EtOH (3:1 blend)/heptane to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 213**) as tan foam. *m/z* (ESI, +ve ion): 636.1 $(M+H)^+$.

Example 213-1

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0561] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 213**) were separated by were separated by chiral Whelk O1(S,S), 20 x 250 mm, 5 μm, 35% MeOH/CO_2, 70 g/min, 164 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer (**Example 213-1**), the first eluting peak, isolated as a white fluffy powder from loyphilization. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.49 (s, 1H), 8.11 (d, *J*=5.60 Hz, 1H), 7.51-7.56 (m, 1H), 7.35-7.46 (m, 2H), 7.06-7.19 (m, 3H), 6.50-6.72 (m, 1H), 6.40 (t, *J*=31.50 Hz, 1H), 5.80 (dd, *J*=8.09, 17.42 Hz, 1H), 3.41-5.23 (m, 6H), 2.77 (s, 3H), 2.64 (spt, *J*=42.70 Hz, 1H), 2.05-2.11 (m, 3H), 0.98-1.50 (m, 12H). ^{19}F NMR (376 MHz, *CDCl3*) δ -112.53 (s, 1F), -112.56 (s, 1F), -112.58 (s, 1F). *m/z* (ESI, +ve ion): 636.2 $(M+H)^+$.

Example 213-2

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer, single isomer



Single Isomer

[0562] Atropisomers 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 213**) were separated by chiral SFC Whelk O1(S,S), 20 x 250 mm, 5 µm, 35% MeOH/CO_2, 70 g/min, 164 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 213-2**), single isomer, the second eluting peak, and isolated as white fluffy powder from loyphilization. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.10 (s, 1H), 7.67 (s, 1H), 7.37-7.47 (m, 1H), 7.05-7.24 (m, 4H), 6.50-6.70 (m, 1H), 6.39 (t, *J*=30.70 Hz, 1H), 5.80 (t, *J*=18.50 Hz, 1H), 3.45-5.20 (m, 6H), 2.77 (s, 3H), 2.54-2.68 (m, 1H), 2.03-2.07 (m, 3H), 1.01-1.49 (m, 12H). ^{19}F NMR (376 MHz, *CDCl3*) δ -112.43 (s, 1F), -112.49 (s, 1F). *m/z* (ESI, +ve ion): 636.2 $(M+H)^+$.

Example 213-3

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



Single Isomer

[0563] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one **(Example 213)** were separated by chiral SFC Step 1: Whelk O1(S,S), 20 x 250 mm, 5 μm, 35% MeOH/CO_2, 70 g/min, 164 bar. Peaks 3 and 4 co-eluted and were then futher purified by Step 2: Whelk O1(R,R), 20 x 250 mm, 5 μm, 30% MeOH/CO_2, 70 g/min, 196 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one **(Example 213-3)**, single isomer, the first eluting peak from step 2, isolated as a white fluffy powder from loyphilization. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.10 (s, 1H), 7.49-7.69 (m, 1H), 7.36-7.46 (m, 2H), 7.05-7.17 (m, 3H), 6.49-6.71 (m, 1H), 6.39 (t, *J*=31.10 Hz, 1H), 5.80 (t, *J*=17.40 Hz, 1H), 3.47-5.22 (m, 6H), 2.77 (s, 3H), 2.54-2.66 (m, 1H), 2.02-2.12 (m, 3H), 1.03-1.47 (m, 12H). ^{19}F NMR (376 MHz, *CDCl3*) δ -112.43 (s, 1F), -112.49 (s, 1F), -112.52 (s, 1F), -112.55 (s, 1F), -112.57 (s, 1F). *m/z* (ESI, +ve ion): 636.2 $(M+H)^+$.

Example 213-4

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer, single isomer



Single Isomer

[0564] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 213**) were separated by chiral SFC Step 1: Whelk O1(S,S), 20 x 250 mm, 5 µm, 35% MeOH/CO_2, 70 g/min, 164 bar. Peaks 3 and 4 co-eluted and were then futher purified by Step 2: Whelk O1(*R*,*R*), 20 x 250 mm, 5 µm, 30% MeOH/CO_2, 70 g/min, 196 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 213-4**), single isomer, the second eluting peak from step 2, isolated as white fluffy powder from loyphilization. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.09 (d, *J*=4.98 Hz, 1H), 7.63-7.68 (m, 1H), 7.38-7.47 (m, 1H), 6.99-7.26 (m, 4H), 6.51-6.70 (m, 1H), 6.39 (t, *J*=31.30 Hz, 1H), 5.81 (dd, *J*=7.26, 16.79 Hz, 1H), 3.43-5.18 (m, 6H), 2.77 (s, 3H), 2.59-2.71 (m, 1H), 2.05 (s, 3H), 0.99-1.48 (m, 12H). ^{19}F NMR (376 MHz, *CDCl3*) δ -112.54 (s, 1F). *m/z* (ESI, +ve ion): 636.2 $(M+H)^+$.

Example 214

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0565] Step 1: (*M*)-*tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylthio)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A suspension of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 92A**, 2.0 g, 3.56 mmol), 2-fluoro-6-(methylthio)phenyl)boronic acid (0.861 g, 4.63 mmol, Aurum Pharmatech LLC, Franklin Park, NJ, USA, 128 mg, 0.686 mmol), tetrakis(triphenylphosphine)palladium(0) (0.206 g, 0.178 mmol), sodium carbonate (1.133 g, 10.69 mmol) in 1,4-dioxane (20 mL) and water (10 mL) was sparged with argon for 1 min then heated to 95 °C for 60 min. The reaction was then extracted with EtOAc (150 mL), dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (80 g) eluting products with a gradient of 10>60% EtOAc/EtOH (3:1 blend)/heptane to give (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylthio)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as white foam. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.46 (d, *J*=4.56 Hz, 1H), 8.15 (d, *J*=1.66 Hz, 1H), 7.30-7.38 (m, 1H), 7.02-7.10 (m,

2H), 6.86-6.95 (m, 1H), 3.41-5.03 (m, 6H), 2.64-2.92 (m, 1H), 2.28-2.37 (m, 3H), 1.93-2.08 (m, 3H), 1.50-1.52 (m, 9H), 1.42-1.49 (m, 3H), 1.20-1.40 (m, 6H), 1.00-1.14 (m, 3H). ^{19}F NMR (376 MHz, *CDCl3*) δ -114.43 (s, 1F), -115.22 (s, 1F). *m/z* (ESI, +ve ion): 667.3 $(M+H)^+$.

[0566] Step 2: (*M*)-*tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylthio)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (850 mg, 1.27 mmol) in DCM (8 mL) at 20 °C was added *m*-chloroperoxybenzoic acid (600 mg, 2.68 mmol) (2x300 mg portions). After 20 min, the reaction was then extracted with sat. Na_2CO_3 (5 mL). The organic was dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (40 g) eluting products with a grandient of 10>60% EtOAc/EtOH (3:1 blend)/heptane to afford (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as white foam. *m/z* (ESI, +ve ion): 699.2 $(M+H)^+$.

[0567] Step 3: (*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 214). A solution of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (370 mg, 0.529 mmol) in TFA (4 mL) was stirred for 15 min at 20 °C. The reaction was then concentrated under reduced pressure, and the residue was dissolved in DCM (5 mL) with DIEA (462 µl, 2.65 mmol) added. The solution was chilled to 15 °C and acryloyl chloride (64.5 µl, 0.794 mmol) added. After 20 min, the reaction was then washed with sat. Na_2CO_3 (5 mL). The separated aqeous was futher extracted with DCM (10 mL). The organic was dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (24 g) eluting products with a gradient of 15>100% EtOAc/EtOH (3:1 blend)/heptane to give (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one **(Example 214)** as colorless foam. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.43-8.51 (m, 1H), 8.12-8.18 (m, 1H), 7.82-7.90 (m, 1H), 7.62-7.71 (m, 1H), 7.45 (t, *J*=17.20 Hz, 1H), 7.02-7.12 (m, 1H), 6.51-6.70 (m, 1H), 6.33-6.46 (m, 1H), 5.75-5.85 (m, 1H), 4.17-5.21 (m, 3H), 3.51-3.99 (m, 3H), 2.61-3.00

(m, 1H), 2.57 (s, 3H), 1.93-2.15 (m, 3H), 1.42-1.49 (m, 3H), 0.99-1.32 (m, 9H). ^{19}F NMR (376 MHz, *CDCl3*) δ -111.28 (s, 1F), -111.31 (s, 1F). *m/z* (ESI, +ve ion): 653.3 $(M+H)^+$.

Example 214-1

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0568] Atropisomers of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one **(Example 214)** were separated by chiral SFC Step 1: OX, 21 x 250 mm, 5 μm, 30% MeOH/CO_2, 80 g/min, 185 bar. (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one **(Example 214-1)** was the first eluting peak, and isolated as a white fluffy solid from lyophilization. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.46 (d, *J*=4.98 Hz, 1H), 8.14 (s, 1H), 7.86 (d, *J*=8.29 Hz, 1H), 7.61-7.69 (m, 1H), 7.44 (dd, *J*=8.71, 17.00 Hz, 1H), 7.04 (d, *J*=4.77 Hz, 1H), 6.49-6.68 (m, 1H), 6.39 (t, *J*=29.40 Hz, 1H), 5.75-5.84 (m, 1H), 4.17-5.24 (m, 3H), 4.00-4.06 (m, 1H), 3.88-3.99 (m, 1H), 3.50-3.78 (m, 1H), 2.60-2.74 (m, 1H), 2.56 (s, 3H), 2.11 (s, 3H), 1.45 (t, *J*=13.70 Hz, 3H), 1.34 (dd, *J*=6.63, 34.83 Hz, 3H), 1.19 (d, *J*=6.84 Hz, 3H), 0.99 (d, *J*=6.43 Hz, 3H). ^{19}F NMR (376 MHz, *CDCl3*) δ -111.29 (d, *J*=10.41 Hz, 1F). *m/z* (ESI, +ve ion): 653.0 $(M+H)^+$.

Example 214-2

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0569] Atropisomers of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 214**) were separated by chiral SFC Step 1: OX, 21 x 250 mm, 5 μm, 30% MeOH/CO_2, 80 g/min, 185 bar. (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-6-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 214-2**) was the second eluting peak, and isolated as a white fluffy solid from lyophilization. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.45 (d, *J*=4.35 Hz, 1H), 8.16 (d, *J*=9.74 Hz, 1H), 7.87 (d, *J*=8.09 Hz, 1H), 7.63-7.71 (m, 1H), 7.45 (t, *J*=17.60 Hz, 1H), 7.07 (d, *J*=3.32 Hz, 1H), 6.51-6.73 (m, 1H), 6.40 (t, *J*=29.00 Hz, 1H), 5.80 (t, *J*=19.90 Hz, 1H), 3.29-5.20 (m, 6H), 2.92 (spt, *J*=39.40 Hz, 1H), 2.57 (s, 3H), 1.96 (s, 3H), 1.43-1.62 (m, 6H), 1.26 (br d, *J*=6.01 Hz, 3H), 1.19 (d, *J*=6.22 Hz, 3H). ^{19}F NMR (376 MHz, *CDCl3*) δ -111.73 (br d, *J*=13.01 Hz, 1F). *m/z* (ESI, +ve ion): 653.0 $(M+H)^+$.

Example 215

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Example 215

[0570] Step 1: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfinyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 213a**, 310 mg, 0.454 mmol) in DCM (3 mL) at 0 °C was added *m*-chloroperoxybenzoic acid (77wt%, 107 mg, 0.477 mmol) in DCM. After 1 hr, the reaction was then washed with 5% Na2CO3 (1 mL), brine (1 mL), dried over MgSO4, filtered, then concentrated to dryness under reduced pressure to afford *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate as white solid. *m/z* (ESI, +ve ion): 698.0 $(M+H)^+$.

[0571] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 215). A solution of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (320 mg, 0.458 mmol) in TFA (2 mL) was stirred for 15 min at 20 °C. The reaction was then concentrated

under reduced pressure, and the residue was dissolved in DCM (3 mL), DIEA (200 µl, 1.146 mmol)) followed by the addition of acryloyl chloride (48.4 µl, 0.596 mmol) at 20 °C. The solution was stirred at 20 °C for 1 hr. The reaction was then partitioned between EtOAc (10 mL) and sat. $NaHCO_3$ (3 mL). The separated aqeous was futher extracted with EtOAc (3 mL). The organic was dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (4 g) eluting products with a gradient of 15>80% EtOAc/EtOH (3:1 blend)/heptane to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 215**) as white solid. ^{1}H NMR (400 MHz, *CDCl*$_3$) δ 8.09-8.13 (m, 1H), 7.66-7.85 (m, 2H), 7.37-7.48 (m, 1H), 7.06-7.21 (m, 3H), 6.50-6.70 (m, 1H), 6.34-6.46 (m, 1H), 5.76-5.85 (m, 1H), 4.19-5.23 (m, 3H), 3.44-4.08 (m, 3H), 3.09 (s, 3H), 2.56-2.73 (m, 1H), 2.06-2.13 (m, 3H), 0.99-1.53 (m, 12H). *m/z* (ESI, +ve ion): 652.1 $(M+H)^+$.

Example 215-1

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



[0572] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 215**) were separated by chiral SFC Step 1: OX-H, 21 x 250 mm, 5 µm, 45% MeOH/CO_2, 80 g/min, 120 bar. This gave 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 215-1**), single isomer, was the first eluting peak, and isolated as a white fluffy solid from lyophilization. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (d, *J*=3.32 Hz, 1H), 7.74-7.78 (m, 2H), 7.47-7.56 (m, 1H), 7.21-7.35 (m, 3H), 6.76-6.91 (m, 1H), 6.20 (dd, *J*=2.07, 16.17 Hz, 1H), 5.72-5.80 (m, 1H), 4.36-4.97 (m, 2H), 3.48-4.25 (m, 4H), 3.29 (s, 3H), 2.59-2.70 (m, 1H), 2.04 (s, 3H), 1.35 (dd, *J*=6.01, 12.02 Hz, 3H), 1.21 (dd, *J*=6.22, 29.23 Hz, 3H), 1.10 (d, *J*=7.26 Hz, 3H), 0.99 (d, *J*=6.63 Hz,

3H). ^{19}F NMR (376 MHz, DMSO-d_6) δ -113.97 (s, 1F), -113.98 (s, 1F). *m/z* (ESI, +ve ion): 652.1 $(M+H)^+$.

Example 215-2

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, single isomer



single isomer

[0573] Atropisomers of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 215**) were separated by chiral SFC Step 1: OX-H, 21 x 250 mm, 5 μm, 45% MeOH/CO_2, 80 g/min, 120 bar. This gave 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-6-methyl-4-(methylsulfonyl)phenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 215-2**), single isomer, was the second eluting peak, and isolated as a white fluffy solid from lyophilization. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (d, *J*=4.15 Hz, 1H), 7.76 (d, *J*=5.39 Hz, 2H), 7.47-7.56 (m, 1H), 7.17-7.36 (m, 3H), 6.76-6.94 (m, 1H), 6.22 (d, *J*=17.00 Hz, 1H), 5.69-5.80 (m, 1H), 4.40-4.98 (m, 2H), 3.46-4.27 (m, 4H), 3.29 (s, 3H), 2.60-2.73 (m, 1H), 2.02 (s, 3H), 1.34 (dd, *J*=6.63, 13.06 Hz, 3H), 1.21 (dd, *J*=6.63, 29.02 Hz, 3H), 1.11 (d, *J*=6.84 Hz, 3H), 1.02 (d, *J*=6.43 Hz, 3H). ^{19}F NMR (376 MHz, DMSO-d_6) δ -113.88 (s, 1F), -113.88 (s, 1F). *m/z* (ESI, +ve ion): 652.1 $(M+H)^+$.

Example 217

5-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-8-(2-fluorophenyl)-1-neopentyl-1,2-dihydro-3*H*-pyrido[2,3-*e*][1,4]diazepin-3-one.



[0574] Step 1: *tert*-Butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)((2-chloroacetyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (Intermediate 217a). To a stirring biphasic solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Example 150**, Step 1; 650 mg, 1.22 mmol) in DCM (5 mL) and 1 M sodium hydroxide (195 µl, 1.95 mmol) at 0 °C was added a solution of 2-chloroacetyl chloride (107 µl, 1.34 mmol) in DCM (1 mL). After 10 min, the aqueous phase was discarded and the organic was washed with sat. NaCl (5 mL). The organic was then purified by silica gel chromatography (24 g) eluting products with a gradient of 0>30% EtOAc/heptane to afford *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)((2-chloroacetyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate as a white foam. *m/z* (ESI, +ve ion): 608.3 $(M+H)^+$.

[0575] Step 2: *tert*-Butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-neopentyl-3-oxo-2,3-dihydro-1H-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)((2-chloroacetyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (230 mg, 0.378 mmol) in THF (3 mL) at 0 °C under nitrogen was added sodium hydride, 60% dispersion in mineral oil (60.5 mg, 1.51 mmol). After 10 min, the cooling bath was removed and the suspension was stirred for 2 hrs at 20 °C. The reaction was chilled to 0 °C, then carefully quenched with sat. NH4Cl (1 mL). The reaction was then partitioned between EtOAc (20 mL) and water (2 mL). The organic was washed with sat. NaCl (2 mL),

dried over MgSO4, and then purified by silica gel chromatography (12 g) eluting products with EtOAc/EtOH (3:1 blend)/heptanes to afford *tert*-butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-neopentyl-3-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate as colorless film. ^{1}H NMR (400 MHz, *CDCl$_3$*) δ 7.41-7.52 (m, 3H), 7.27-7.30 (m, 1H), 7.18 (dd, *J*=9.12, 18.24 Hz, 1H), 2.90-5.08 (m, 10H), 1.48 (s, 9H), 1.20-1.42 (m, 6H), 0.87 (br s, 9H). *m/z* (ESI, +ve ion): 572.3 $(M+H)^+$.

[0576] Step 3: 5-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-8-(2-fluorophenyl)-1-neopentyl-1,2-dihydro-3H-pyrido[2,3-*e*][1,4]diazepin-3-one (Example 217). A solution of *tert*-butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-neopentyl-3-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (60 mg, 0.105 mmol) in TFA (2 mL) was stirred for 15 min at RT. The solvent was then removed under reduced pressure, and the resulting residue was then partitioned between EtOAc (20 mL) and 5% Na2CO3 (5 mL). The organic was then washed with sat. NaCl (2 mL), dried over MgSO4, then concentrated under reduced pressure. The residue was then dissolved in DCM (4 mL) and DIPEA (55.0 µl, 0.315 mmol) was added. The stirring solution was then chilled to 0 °C under nitrogen and a second solution of acryloyl chloride (9.41 µl, 0.115 mmol) in DCM (1 mL) was added dropwise. The cooling bath was removed, stirred for 10 min, then the reaction was washed with sat. NaHCO3 (2 mL). The separated organic was then directly purified by silica gel chromatography (4 g) eluting products with a gradient of 10>50% EtOAc/EtOH (3:1 blend)/heptane to affor a light yellow foam. The material was then dissolved in 1:1 MeCN/water (2 mL), frozen, and lyophilized to afford afford 5-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-8-(2-fluorophenyl)-1-neopentyl-1,2-dihydro-3*H*-pyrido[2,3-*e*][1,4]diazepin-3-one (**Example 217**) as light yellow fluffy powder. ^{1}H NMR (400 MHz, *DMSO-d$_6$*) δ 8.00 (br s, 1H), 7.48-7.62 (m, 2H), 7.34-7.41 (m, 2H), 6.67-6.89 (m, 1H), 6.12-6.20 (m, 1H), 5.73 (br d, *J*=10.37 Hz, 1H), 3.21-4.90 (m, 10H), 1.04-1.49 (m, 6H), 0.84 (br s, 9H). ^{19}F NMR (376 MHz, DMSO-*d$_6$*) δ -114.06 (s, 1F). *m/z* (ESI, +ve ion): 526.3 $(M+H)^+$.

Example 218

5-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-1,2-dihydro-3*H*-pyrido[2,3-*e*][1,4]diazepin-3-one



[0577] Step 1: *tert*-Butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)((2-chloroacetyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 237**) (0.5 g, 0.840 mmol) and DIPEA (0.161 ml, 0.924 mmol) in DCM (10 mL) at 0 °C under argon was added a second solution of chloroacetyl chloride (0.053 ml, 0.672 mmol) in DCM (0.5 mL). After 5 min, the reaction was then directly purified by silica gel chromatography (12 g) eluting products with a gradient of 10>40% EtOAc/heptane to afford *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)((2-chloroacetyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate as white solid. *m/z* (ESI, +ve ion): 671.3 $(M+H)^+$.

[0578] Step 2: *tert*-Butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-3-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)((2-chloroacetyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (225 mg, 0.335 mmol) in THF (5 mL) at -70 °C under argon was added sodium bis(trimethylsilyl)amide, 1 M in THF (670 µl, 0.670 mmol). After 5 min, the reaction was then quenched with sat. NH_4Cl (0.5 mL). The reaction was then partitioned between EtOAc (10 mL) and water (1 mL). The organic was washed with sat. NaCl (1 mL), dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (12 g) eluting products with a gradient of 10>60% EtOAc/EtOH (3:1 blend)/heptane to afford *tert*-butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-

1-(2-isopropyl-4-methylpyridin-3-yl)-3-oxo-2,3-dihydro-1H-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate as yellow fim. *m/z* (ESI, +ve ion): 635.3 $(M+H)^+$.

[0579] Step 3: 5-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-1,2-dihydro-3*H*-pyrido[2,3-*e*][1,4]diazepin-3-one (Example 218). A solution of *tert*-butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-3-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (65 mg, 0.102 mmol) in TFA was stirred for 5 min at RT. The solvent was then removed under reduced pressure, and the residue was partitioned between EtOAc (10 mL) and 5% Na_2CO_3 (5 mL). The organic was washed with sat. NaCl (1 mL), dried over $MgSO_4$, then concentrated under reduced pressure. The resulting residue was then dissolved in DCM (2 mL), DIEA (17.9 µl, 0.102 mmol) added, chilled to 0 °C under nitrogen, and a second solution of acryloyl chloride (8.31 µl, 0.102 mmol) in DCM (0.5 mL) was added dropwise. After 5 min, the reaction was then directly purified by silica gel chromatography (4 g) eluting products with a gradient of 10>70% EtOAc/EtOH (3:1 blend)/heptane to afford a yellow oil. The oil was then suspended in MeCN/water (1:1; 1.5 mL), frozen, and lyophilized to afford 5-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-1,2-dihydro-3H-pyrido[2,3-*e*][1,4]diazepin-3-one **(Example 218)** as an off white fluffy powder. *m/z* (ESI, +ve ion): 589.3 $(M+H)^+$. ^1H NMR (400 MHz, *DMSO-d_6*) δ 8.29 (d, *J*=4.77 Hz, 1H), 8.03 (br s, 1H), 7.34-7.48 (m, 1H), 7.04-7.27 (m, 4H), 6.73-6.89 (m, 1H), 6.17 (br d, *J*=16.79 Hz, 1H), 5.70-5.77 (m, 1H), 3.32-4.93 (m, 8H), 2.82-3.19 (m, 1H), 2.08 (br d, *J*=6.22 Hz, 3H), 1.09-1.36 (m, 9H), 0.84-0.92 (m, 3H).

Example 219

5-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-1,3-dihydro-2*H*-pyrido[2,3-*e*][1,4]diazepin-2-one



[0580] Step 1: *tert*-Butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (Intermediate 237). A 500 mL round bottom flask was flushed with argon then charged with diacetoxypalladium (1.68 g, 7.48 mmol), (9,9-dimethyl-9*H*-xanthene-4,5-diyl)bis(diphenylphosphane) (6.49 g, 11.2 mmol), cesium carbonate (12.2 g, 37.4 mmol), and 1,4-dioxane (60 mL). The suspension was then sparged with nitrogen for 2 min, stirred for 10 min at 20 °C, then *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 165**, 6.0 g, 12.5 mmol) and 2-isopropyl-4-methylpyridin-3-amine (**Intermediate R**, 2.34 g, 15.6 mmol) were added. The suspension was then heated to 100 °C. After 45 min, the reaction was partitioned between EtOAc (200 mL) and water (100 mL). The organic was then dried over $MgSO_4$, and concentrated under reduced pressure. The resulting residue was dissolved in EtOAc (150 mL), and SiliaMetS® Thiol by Silicycle (40 g, 1.3 mmol/g). The slurry was stirred for 90 min at RT. The solvent was removed under reduced pressure, and then purified by silica gel chromatography (4 g) eluting products with a gradient of 10>50% EtOAc/EtOH (3:1 blend)/heptane to afford *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 237**) as brown foam. *m/z* (ESI, +ve ion): 595.4 $(M+H)^+$.

[0581] Step 2: *tert*-Butyl (2*R*,5*S*)-4-((*E*)-(5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)((2-methoxy-2-oxoethyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-

6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 237**, 400 mg, 0.672 mmol) and DIPEA (235 µl, 1.34 mmol) in 1,2-dichloroethane (8 mL) was added methyl bromoacetate (623 µl, 6.72 mmol). The solution was stirred at 85 °C in a 20 mL reaction vessel equipped with a pressure relief cap for 16 hrs. The reaction was then directly purified by silica gel chromatography (12 g) eluting products with a gradient of 0>50% EtOAc/EtOH (3:1 blend)/heptane to afford *tert*-butyl (2*R*,5*S*)-4-((*E*)-(5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)((2-methoxy-2-oxoethyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate as a white solid. *m/z* (ESI, +ve ion): 667.3 $(M+H)^+$.

[0582] Step 3: 2-(((*E*)-((2*S*,5*R*)-4-(*tert*-Butoxycarbonyl)-2,5-dimethylpiperazin-1-yl)(5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)methylene)amino)acetic acid. A suspension of *tert*-butyl (2*R*,5*S*)-4-((*E*)-(5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)((2-methoxy-2-oxoethyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (125 mg, 0.187 mmol) and lithium hydroxide, monohydrate (9.83 mg, 0.234 mmol) in THF (1 mL), MeOH (0.5 mL), water (0.5 mL) was stirred for 18 hrs at RT. The reaction was then partitioned between DCM (20 ml) and sat. $NaHCO_3$ (10 mL). The aqueous was then further extracted with DCM (5 mL). The combined organics were washed with sat. NaCl (5 mL), dried over $MgSO_4$, fitlered, then concentrated under reduced pressure to afford as yellow solid. This material was then further purified by silica gel chromatography (4 g) eluting products with 100% EtOAc/EtOH (3:1 blend) to afford 2-(((*E*)-((2*S*,5*R*)-4-(*tert*-butoxycarbonyl)-2,5-dimethylpiperazin-1-yl)(5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)methylene)amino)acetic acid as white solid. *m/z* (ESI, +ve ion): 653.3 $(M+H)^+$.

[0583] Step 4: *tert*-Butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of 2-(((*E*)-((2*S*,5*R*)-4-(*tert*-butoxycarbonyl)-2,5-dimethylpiperazin-1-yl)(5-chloro-6-(2-fluorophenyl)-2-((2-isopropyl-4-methylpyridin-3-yl)amino)pyridin-3-yl)methylene)amino)acetic acid (45 mg, 0.069 mmol) and DIEA (15.0 µl, 0.086 mmol in DCM (2 mL) at 0 °C under nitrogen was added phosphorus oxychloride (6.42 µl, 0.069 mmol). After 10 min, the reaction was then partitioned between EtOAc (10 mL) and sat. $NaHCO_3$ (5 mL). The organic was further washed with sat. NaCl (2 mL), then dried over $MgSO_4$. The organic layer was then concentrated under reduced pressure, and purified by silica gel chromatography (4 g) eluting the product with a gradient of 10>50%

EtOAc/EtOH (3:1 blend)/heptane to afford *tert*-butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate as colorless film. *m/z* (ESI, +ve ion): 635.3 $(M+H)^+$.

[0584] Step 5: 5-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-1,3-dihydro-2*H*-pyrido[2,3-*e*][1,4]diazepin-2-one (Example 219). A solution of *tert*-butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (6 mg, 9.45 μmol) in TFA (1 mL) was stirred for 15 min at RT. The solvent was then removed under reduced pressure, and the residue was azeotropted with tolune (2x5 mL). The residue was then dissolved in DCM (1 mL), DIEA (16.5 μL, 0.094 mmol) added, and chilled to 0 °C under an atmosphere of argon. To this was added a second solution of acryloyl chloride (0.767 μl, 9.45 μmol) in DCM (0.5 mL). The reaction was then directly purified by silica gel chromatography (4 g) eluting the products with a gradient of 10>60% EtOAc/EtOH (3:1 blend)/heptane to afford 5-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-8-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-1,3-dihydro-2*H*-pyrido[2,3-*e*][1,4]diazepin-2-one **(Example 219)** as a colorless film._*m/z* (ESI, +ve ion): 589.1 $(M+H)^+$. ^{1}H NMR (400 MHz, *CDCl3*) δ 8.41-8.64 (m, 1H), 7.96-8.09 (m, 1H), 7.37-7.48 (m, 1H), 7.07-7.23 (m, 4H), 6.27-6.75 (m, 2H), 5.76 (br d, *J*=10.37 Hz, 1H), 4.17-5.12 (m, 3H), 2.50-3.99 (m, 6H), 1.91-2.35 (m, 3H), 1.47-1.61 (m, 3H), 1.08-1.46 (m, 6H), 0.81-0.95 (m, 3H). ^{19}F NMR (376 MHz, *CDCl3*) δ -112.62 (s, 1F).

Example 220

5-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-8-(2-fluorophenyl)-1-neopentyl-1,3-dihydro-2H-pyrido[2,3-*e*][1,4]diazepin-2-one



[0585] Step 1: *tert*-Butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)((2-methoxy-2-oxoethyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate. A solution of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Example 150**, Step 1, 400 mg, 0.752 mmol), methyl bromoacetate (348 µl, 3.76 mmol), DIPEA (197 µl, 1.13 mmol) in DMF (3 mL) was heated to 85 °C for 16 hrs. The reaction miture was then partitioned between EtOAc (20 mL) and sat. $NaHCO_3$ (20 mL). The organic was then further washed with water (10 mL) and sat. NaCl (5 mL), then dried over $MgSO_4$. The organic layer was concentrated under reduced pressure, and then purified by silica gel chromatography (24 g) eluting the product with a gradient of 10>50% EtOAc/EtOH (3:1 blend)/heptane to afford *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)((2-methoxy-2-oxoethyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate as white solid. *m/z* (ESI, +ve ion): 604.4 $(M+H)^+$.

[0586] Step 2: 2-((((2*S*,5*R*)-4-(*tert*-Butoxycarbonyl)-2,5-dimethylpiperazin-1-yl)(5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)methylene)amino)acetic acid. A suspension of *tert*-butyl (2*R*,5*S*)-4-((5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)((2-methoxy-2-oxoethyl)imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (310 mg, 0.513 mmol) and lithium hydroxide, monohydrate (23.7 mg, 0.564 mmol) in MeOH (5 mL) and water (1 mL) was stirred at RT for 3 days. The reaction mixture was then partitioned between DCM (20 ml) and sat. $NaHCO_3$ (10 mL). The aqueous layer was further extracted with DCM (5 mL). The combined organics were washed with sat. NaCl (5 mL), dried over MgSO4, fitlered, and then concentrated under reduced pressure to afford 2-((((2*S*,5*R*)-4-(*tert*-butoxycarbonyl)-2,5-dimethylpiperazin-1-yl)(5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)methylene)amino)acetic acid as white solid. *m/z* (ESI, +ve ion): 590.4 $(M+H)^+$.

[0587] Step 3: *tert*-Butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-neopentyl-2-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate. To a stirring solution of 2-((((2*S*,5*R*)-4-(*tert*-butoxycarbonyl)-2,5-dimethylpiperazin-1-yl)(5-chloro-6-(2-fluorophenyl)-2-(neopentylamino)pyridin-3-yl)methylene)amino)acetic acid (300 mg, 0.508 mmol) in 1,2-dichloroethane (3 mL) at RT was added DIPEA (266 µl, 1.53 mmol) followed by phosphorus oxychloride (95 µl, 1.02 mmol). After 15 min, the reaction was then partitioned between EtOAc (20 mL) and sat. $NaHCO_3$ (20 mL). The organic was then washed with sat. NaCl (5 mL), dried over $MgSO_4$, concentrated under reduced pressure, and then

purified by silica gel chromatography (12 g) eluting products with a gradient of 10>40% EtOAc/EtOH (3:1 blend)/heptane to afford *tert*-butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-neopentyl-2-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate as white solid. *m/z* (ESI, +ve ion): 572.2 $(M+H)^+$.

[0588] Step 4: *tert*-Butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-neopentyl-2-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (Example 220). A solution of *tert*-butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-neopentyl-2-oxo-2,3-dihydro-1*H*-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (40 mg, 0.070 mmol) in TFA (1 ml) was stirred for 15 min at RT. The solvent was then removed under reduced pressure, and the residue was azeotropted with toluene (2x5 mL). The resulting residue was then dissolved in DCM (1 mL), DIPEA (12.2 µl, 0.070 mmol) added, and the solution was then chilled to 0 °C under an atmosphere of argon. To this solution was added a second solution of acryloyl chloride (5.68 µl, 0.070 mmol) in DCM (0.5 mL). After 5 min, the reaction was then directly purified by silica gel chromatography (4 g) eluting products with a gradient of 10>60% EtOAc/EtOH (3:1 blend)/heptane to afford the product as colorless film. The film was then suspended in MeCN/water (1:1; 2 mL), frozen, then lyophilized to afford *tert*-butyl (2*R*,5*S*)-4-(7-chloro-8-(2-fluorophenyl)-1-neopentyl-2-oxo-2,3-dihydro-1H-pyrido[2,3-*e*][1,4]diazepin-5-yl)-2,5-dimethylpiperazine-1-carboxylate **(Example 220l)** as a white fluffy powder. *m/z* (ESI, +ve ion): 526.6 $(M+H)^+$. ^1H NMR (400 MHz, $CDCl_3$) δ 7.85-7.98 (m, 1H), 7.43-7.54 (m, 2H), 7.27-7.35 (m, 1H), 7.21 (t, *J*=18.00 Hz, 1H), 6.45-6.69 (m, 1H), 6.28-6.40 (m, 1H), 5.69-5.79 (m, 1H), 3.02-5.05 (m, 10H), 1.36-1.55 (m, 3H), 0.93-1.26 (m, 3H), 0.72-0.78 (m, 9H). ^{19}F NMR (376 MHz, $CDCl_3$) δ -113.20 (s, 1F), -113.23 (s, 1F), -113.30 (s, 1F).

Example 221

4-(4-Acryloylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carboxamide



[0589] **Step 1: 2-Cyano-*N*-(2-isopropyl-4-methylpyridin-3-yl)acetamide.** To a solution of 2-isopropyl-4-methylpyridin-3-amine (4.18 g, 27.8 mmol, **Intermediate R**) and cyano acetic acid (3.31 g, 39 mmol, Sigma-Aldrich, St. Louis, MO) in DCE (25 mL) was added 1-propanephosphonic acid cyclic anhydride, 50 wt. % solution in ethyl acetate (25.1 mL, 36 mmol, Sigma-Aldrich, St. Louis, MO) and *i*Pr$_2$EtN (9.72 mL, 57 mmol). The mixture was heated to 50 °C for 30 min. The mixture was cooled to rt, added 30 mL ethyl acetate and 30 mL of satd. $NaHCO_3$ and stirred for 5 min. The organic layer was separated, washed with brine (20 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The yellow solid obtained was triturated in heptane:EtOAc (5:1) (20 mL), filtered off, and dried under vacuum. 2-Cyano-*N*-(2-isopropyl-4-methylpyridin-3-yl)acetamide was obtained as a light yellow solid. m/z (ESI, +ve) 218.0 $(M+H)^+$.

[0590] **Step 2: 2,5-Dichloro-6-(2-fluorophenyl)nicotinoyl chloride.** To a solution of 2,5-dichloro-6-(2-fluorophenyl)nicotinic acid (7.1 g, 25 mmol, **Intermediate 99A**) in DCM (50 mL) was added oxalyl chloride, 2.0 M solution in methylene chloride (15.4 mL, 31 mmol, Sigma-Aldrich, St. Louis, MO), followed by several drops of DMF. The mixture was stirred at RT for 45 min, evaporated, dried *in vacuo* to provide 2,5-dichloro-6-(2-fluorophenyl)nicotinoyl chloride, that was carried forward into the next step.

[0591] **Step 3: 6-Chloro-7-(2-fluorophenyl)-4-hydroxy-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carbonitrile.** To a 250-mL round-bottomed flask was added 2-cyano-*N*-(2-isopropyl-4-methylpyridin-3-yl)acetamide

(4.97 g, 22.9 mmol) in THF (30 mL). The reaction mixture was cooled to 0 °C and slowly added sodium hydride (3.66 g, 91 mmol, 60%, Sigma-Aldrich, St. Louis, MO) portion wise. The mixture was then stirred at rt for 20 min and cooled to 0 °C and slowly added 2,5-dichloro-6-(2-fluorophenyl)nicotinoyl chloride (7.31 g, 24.0 mmol) in THF (20 mL). After stirring at rt for 10 min, the mixture was heated in 55 °C for 1 h. The reaction mixture was allowed to cool to rt and quenched with saturated aqueous NH_4Cl (10 mL) and extracted with EtOAc (10 mL). The organic layer was separated, washed with brine (5 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The orange solid obtained was triturated in heptane:EtOAc (5:1) (10 mL), filtered off, and dried under vacuum. 6-Chloro-7-(2-fluorophenyl)-4-hydroxy-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carbonitrile was obtained as an orange solid. m/z (ESI, +ve) 448.9 $(M+H)^+$.

[0592] Step 4: *tert*-Butyl 4-(6-chloro-3-cyano-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydro-1,8-naphthyridin-4-yl)piperazine-1-carboxylate. To a solution of 6-chloro-7-(2-fluorophenyl)-4-hydroxy-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carbonitrile (7.71 g, 17.2 mmol) and *i*Pr_2EtN (9 mL, 51.5 mmol) in ACN (40 mL) at 0 °C was added phosphorus oxychloride (2.1 mL, 22.3 mmol, Sigma-Aldrich, St. Louis, MO), and the resulting solution was stirred at 80 °C for 30 min. The reaction mixture was cooled down to 0 °C and *i*Pr_2EtN (9 mL, 51.5 mmol) was added followed by *tert*-butyl piperazine-1-carboxylate (4.80 g, 25.8 mmol, Sigma-Aldrich, St. Louis, MO) in ACN (6 mL). After stirring at rt for 20 min, satd $NaHCO_3$ (10 mL) and EtOAc (10mL) were added. The organic layer was taken, washed with water, dried over Na_2SO_4, filtered, and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, 0–25% EtOAc:EtOH (3:1) in heptane) provided *tert*-butyl 4-(6-chloro-3-cyano-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydro-1,8-naphthyridin-4-yl)piperazine-1-carboxylate as a light orange solid: m/z (ESI, +ve) 617.0 $(M+H)^+$.

[0593] Step 5: 6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-4-(piperazin-1-yl)-1,2-dihydro-1,8-naphthyridine-3-carboxamide. To a 50-mL round-bottomed flask was added *tert*-butyl 4-(6-chloro-3-cyano-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydro-1,8-naphthyridin-4-yl)piperazine-1-carboxylate (133 mg, 0.22 mmol), DCM (2 mL), and trifluoroacetic acid (1 mL). The reaction mixture was stirred at rt for 30 min and the solvent was removed under vacuum. The orange residue obtained was suspended in H_2SO_4 95% (1 mL) and heated at 70 °C for 16 h. The reaction mixture was allowed to cool to rt and slowly poured over ice water. The solution was neutralized with NaOH

(2 N) and extracted with EtOAc. The organic layer was taken, washed with water, dried over Na_2SO_4, filtered, and concentrated *in vacuo*. 6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-4-(piperazin-1-yl)-1,2-dihydro-1,8-naphthyridine-3-carboxamide was obtained as a light yellow solid. m/z (ESI, +ve) 535.1 $(M+H)^+$.

[0594] Step 6: 4-(4-Acryloylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carboxamide. To a 50-mL round bottomed flask was added 6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-4-(piperazin-1-yl)-1,2-dihydro-1,8-naphthyridine-3-carboxamide (110 mg, 0.21 mmol), DCM (5 mL) and iPr$_2$EtN (0.14 mL, 0.82 mmol). The reaction mixture was cooled to 0 °C and acryloyl chloride (0.45 mL, 0.23 mmol, 0.5 M in DCM) was added. The reaction mixture was stirred at 0 °C for 60 min and quenched with saturated aqueous $NaHCO_3$ (5 mL) and extracted with EtOAc (10 mL). The organic layer was separated, washed with brine (5 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, 0–60% EtOAc:EtOH (3:1) in heptane) provided 4-(4-acryloylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carboxamide as a light yellow solid. ^{1}H NMR (400 MHz, *DMSO-d_6*) δ 8.43 (s, 1 H), 8.41 (d, *J*=4.77 Hz, 1 H), 7.87 (s, 1 H), 7.66 (s, 1 H), 7.44 - 7.56 (m, 1 H), 7.17 - 7.33 (m, 4 H), 6.90 (dd, *J*=16.59, 10.37 Hz, 1 H), 6.18 (dd, *J*=16.59, 2.49 Hz, 1 H), 5.71 - 5.80 (m, 1 H), 3.86 (br s, 4 H), 3.25 - 3.30 (m, 4 H), 2.64 (quin, *J*=6.84 Hz, 1 H), 1.90 (s, 3 H), 1.07 (d, *J*=6.63 Hz, 3 H), 0.91 (d, *J*=6.63 Hz, 3 H); m/z (ESI, +ve) 589.0 $(M+H)^+$.

Example 222:

(*M*)-4-(5-Acryloyl-2,5-diazabicyclo[4.1.0]heptan-2-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Examples 222, 222-1, and 222-2)



[0595] Step 1: (*M*)-*tert*-Butyl 5-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-diazabicyclo[4.1.0]heptane-2-carboxylate. DIPEA (0.19 mL, 1.1 mmol) and $POCl_3$ (0.10 mL, 1.11 mmol) were sequentially added to a room temperature solution of (*M*)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 73B**, 270 mg, 0.74 mmol) in acetonitrile (3.5 mL). The homogenous bright-yellow solution was heated at 60 °C. After 2 h, the reaction mixture was concentrated under reduced pressure and azeotropically dried with heptane. The resulting dark-orange residue was dissolved in DCM (2.0 mL) and cooled to 0 °C. DIPEA (0.65 mL, 3.7 mmol) was added, followed by a solution of *tert*-butyl 2,5-diazabicyclo[4.1.0]heptane-2-carboxylate (Aurum Pharmatech LLC, Franklin Park, NJ, USA, 250 mg, 1.3 mmol) in DCM (1.5 mL). After 18 h, the reaction was diluted with EtOAc (70 mL) and water (70 mL). The layers were partitioned and the aqueous phase was extracted with EtOAc (2 x 30 mL). The combined organic extracts were dried over anhydrous magnesium sulfate, filtered, and concentrated to afford an orange oil which was purified by silica gel chromatography (eluent: 0–50% (3:1 EtOAc–EtOH)/heptane) to afford (*M*)-*tert*-butyl 5-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-diazabicyclo[4.1.0]heptane-2-carboxylate (390 mg, 0.72 mmol, 97% yield) as a bright yellow-orange solid.

[0596] Step 2: (*M*)-*tert*-Butyl 5-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-diazabicyclo[4.1.0]heptane-2-carboxylate. A biphasic mixture of (*M*)-*tert*-butyl 5-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-diazabicyclo[4.1.0]heptane-2-carboxylate (390 mg, 0.72 mmol), 2-fluorophenylboronic acid (0.15 mg, 1.1 mmol, Combi-Blocks Inc., San Diego, CA, USA), Pd(dppf)Cl_2 (53 mg, 0.072 mmol), and potassium acetate (0.23 mg, 3.6 mmol) in 1,4-dioxane (4.2 mL)/water (0.10 mL) was deoxygenated with bubbling nitrogen gas for 10 min, then heated at 90 °C overnight. The reaction mixture was filtered through Celite and diluted with EtOAc (100 mL). The organic phase was sequentially washed with saturated aqueous ammonium chloride (100 mL), water (100 mL), and brine (100 mL), then dried over anhydrous magnesium sulfate, filtered, and concentrated to afford crude (*M*)-*tert*-butyl 5-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-diazabicyclo[4.1.0]heptane-2-carboxylate as a brown foam which was carried forward in the next step without purification, assuming theoretical yield.

[0597] Step 3: (*M*)-4-(5-Acryloyl-2,5-diazabicyclo[4.1.0]heptan-2-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 222). A solution of (*M*)-*tert*-butyl 5-(6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-diazabicyclo[4.1.0]heptane-2-carboxylate (440 mg, 0.72 mmol) in DCM (4.0 mL) was treated with trifluoroacetic acid (2.0 mL, 26 mmol). After 15 min, the mixture was concentrated *in vacuo*, then the crude intermediate was redissolved in DCM (8.0 mL) and cooled to 0 °C. Sequentially, DIPEA (1.4 mL, 8.0 mmol) and acryloyl chloride solution (0.26 M in DCM, 3.0 mL, 0.77 mmol) were added and the mixture was allowed to warm to ambient temperature. After 45 min, the reaction was quenched with sat. aq. sodium bicarbonate (15 mL) and diluted with DCM (15 mL). The layers were partitioned and then the aqueous layer was extracted with DCM (2 x 40 mL). The combined organic extracts were dried over anhydrous magnesium sulfate, filtered, and concentrated to afford a dark orange-brown residue which was purified by silica gel chromatography (eluent: 0-100% (3:1 EtOAc–EtOH)/heptane) to afford (*M*)-4-(5-acryloyl-2,5-diazabicyclo[4.1.0]heptan-2-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 222**, 360 mg, 0.64 mmol, 88% yield) as a mixture of piperazine diastereomers. ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.91 (t, J = 5.8 Hz, 3 H), 1.06 (d, J = 6.6 Hz, 3 H), 1.45 - 1.54 (m, 1 H), 1.85 - 2.00 (m, 3 H), 2.62

(sept, J = 6.6 Hz, 0.5 H), 2.79 (sept, J = 6.6 Hz, 0.5 H), 3.35 - 4.08 (m, 6 H), 4.56 - 4.81 (m, 1 H), 5.75 - 5.83 (m, 1 H), 6.16 - 6.32 (m, 1 H), 6.96 (ddd, J = 16.9, 10.4, 3.8 Hz, 1 H), 7.15 - 7.36 (m, 4 H), 7.46 - 7.56 (m, 1 H), 8.38 (dd, J = 4.8, 1.2 Hz, 1 H), 9.09 (s, 1 H); $^{19}F\{^{1}H\}$ NMR (376 MHz, DMSO-d_6) δ –114.20 (s, 1 F), –114.18 (s, 1 F); LRMS: (ESI, +ve ion) m/z 559.1 $(M+H)^+$.

[0598] Step 4: (*M*)-4-((1*R*,6*S*)-5-Acryloyl-2,5-diazabicyclo[4.1.0]heptan-2-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 222-1) and (*M*)-4-((1*S*,6*R*)-5-acryloyl-2,5-diazabicyclo[4.1.0]heptan-2-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 222-2). (*M*)-4-(5-Acryloyl-2,5-diazabicyclo[4.1.0]heptan-2-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (350 mg, 0.62 mmol) was dissolved in MeOH (4.0 mL) and DCM (1.0 mL) and purified by preparative SFC (Chiralcel ID column (5 um, 21 x 250 mm), F = 80 mL, 50% MeOH/50% CO_2) to afford (*M*)-4-((1*R*,6*S*)-5-acryloyl-2,5-diazabicyclo[4.1.0]heptan-2-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 222-1**, 150 mg, 0.26 mmol) as a tan solid (1st-eluting isomer) and (*M*)-4-((1*S*,6*R*)-5-acryloyl-2,5-diazabicyclo[4.1.0]heptan-2-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 222-2**, 130 mg, 0.24 mmol) as a tan solid (2nd-eluting isomer). **Example 222-1**: ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.92 (d, J = 6.6 Hz, 3 H), 1.06 (d, J = 6.6 Hz, 3 H), 1.10 - 1.17 (m, 1 H), 1.50 (q, J = 6.4 Hz, 1 H), 1.85 - 1.91 (m, 3 H), 2.71 - 2.87 (m, 1 H), 3.44 - 4.81 (m, 6 H), 5.76 - 5.84 (m, 1 H), 6.19 - 6.30 (m, 1 H), 6.96 (dd, J = 16.8, 10.4 Hz, 1 H), 7.14 - 7.34 (m, 4 H), 7.47 - 7.55 (m, 1 H), 8.38 (d, J = 5.0 Hz, 1 H), 9.09 (s, 1 H); $^{19}F\{^{1}H\}$ NMR (376 MHz, DMSO-d_6) δ –114.17 (s, 1 F); LRMS: (ESI, +ve ion) m/z 559.1 $(M+H)^+$. **Example 222-2**: ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.91 (d, J = 6.8 Hz, 3 H), 1.06 (d, J = 6.8 Hz, 3 H), 1.08 - 1.12 (m, 1 H), 1.50 (q, J = 6.4 Hz, 1 H), 1.92 - 2.02 (m, 3 H), 2.62 (sept, J = 6.6 Hz, 1 H), 3.43 - 4.78 (m, 6 H) 5.76 - 5.83 (m, 1 H), 6.16 - 6.29 (m, 1 H), 6.95 (dd, J = 16.8, 10.4 Hz, 1 H), 7.13 - 7.35 (m, 4 H), 7.46 - 7.55 (m, 1 H), 8.37 (d, J = 4.8 Hz, 1 H), 9.09 (s, 1 H); $^{19}F\{^{1}H\}$ NMR (376 MHz, DMSO-d_6) δ –114.11 (s, 1 F); LRMS: (ESI, +ve ion) m/z 559.3 $(M+H)^+$.

Example 223:

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-5-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Example 223)



Intermediate 92B **Example 223**

[0599] A biphasic mixture of (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 92B**, 220 mg, 0.43 mmol), 2-(2-fluoro-5-(methylsulfonyl)phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (Aurum Pharmatech LLC, Franklin Park, NJ, USA, 220 mg, 0.73 mmol), $Pd(dppf)Cl_2$ (31 mg, 0.043 mmol), and potassium acetate (210 mg, 2.1 mmol) in 1,4-dioxane (2.2 mL)/water (200 μL) was deoxygenated with bubbling nitrogen gas for 10 min, then heated at 90 °C overnight. The reaction mixture was filtered through Celite using EtOAc (100 mL). The organic phase was sequentially washed with 50% saturated aqueous ammonium chloride (2 x 100 mL) and brine (100 mL), then dried over anhydrous magnesium sulfate, filtered, and concentrated to afford a gold-orange oil which was purified by silica gel chromatography (eluent: 10–100% 3:1 EtOAc–EtOH/heptane) to afford (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluoro-5-(methylsulfonyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Example 223**, 210 mg, 0.32 mmol, 75% yield) as an orange solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 1.00 (br d, J = 6.6 Hz, 3 H), 1.07 (d, J = 6.6 Hz, 3 H), 1.13 - 1.28 (m, 3 H), 1.34 (br t, J = 6.6 Hz, 3 H), 1.94 (s, 3 H), 2.68 - 2.76 (m, 1 H), 3.21 (s, 3 H), 3.46 - 4.97 (m, 6 H), 5.75 (s, 1 H), 6.19 (br d, J = 17.2 Hz, 1 H), 6.84 (td, J = 16.5, 10.6 Hz, 1 H), 7.19 (d, J = 4.8 Hz, 1 H), 7.65 (t, J = 9.1 Hz, 1 H), 7.84 (br d, J = 4.6 Hz, 1 H), 8.04 - 8.18 (m, 1 H), 8.40 (d, J = 4.8 Hz, 1 H), 8.52 (d, J = 2.9 Hz, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –106.12 - –106.13 (m, 1 F); LRMS: (ESI, +ve ion) m/z 652.4 $(M+H)^+$.

Example 224:

4-((3*R*,5*S*)-4-Acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridin-2(1*H*)-one (Examples 224, 224-1, and 224-2)



[0600] Step 1: 4-((3*R*,5*S*)-4-Acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridin-2(1*H*)-one (Example 224). A room-temperature mixture of 6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridine-2,4(1*H*,3*H*)-dione (**Intermediate 111A**, 260 mg, 0.62 mmol) and DIPEA (0.18 mL, 1.1 mmol) in acetonitrile (5.0 mL) was treated with phosphorus oxychloride (0.058 mL, 0.63 mmol). The resulting homogenous brown mixture was heated at 80 °C for 3 h. The reaction mixture was concentrated *in vacuo* and the resulting brown residue was dissolved in acetonitrile (4.0 mL) and cooled to 0 °C in an ice-water bath. DIPEA (1.1 mL, 6.2 mmol) was added, followed by drop-wise addition of a suspension of 1-((2*R*,6*S*)-2,6-dimethylpiperazin-1-yl)prop-2-en-1-one 2,2,2-trifluoroacetate (**Intermediate 223**, 240 mg, 0.85 mmol) in acetonitrile (2.0 mL). After 1 h, the reaction mixture was diluted with EtOAc (30 mL) and washed with 50% saturated aqueous ammonium chloride (100 mL). The layers were partitioned and the aqueous phase was extracted with EtOAc (2 × 30 mL). The combined organic extracts were dried over anhydrous magnesium sulfate, filtered, and concentrated to afford a red-brown residue which was purified by silica gel chromatography (eluent: 0–100% 3:1 EtOAc–EtOH/heptane) to afford 4-((3*R*,5*S*)-4-acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridin-2(1*H*)-one (**Example 224**, 190 mg, 0.32 mmol, 52% yield) as a rust-colored solid (mixture of atropisomers). ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.82 - 0.89 (m, 3 H), 0.96 (br d, J = 6.4 Hz, 3 H), 1.07 (d, J = 6.6 Hz, 3 H), 1.25 (br s, 3 H), 1.99 (d, J = 2.7 Hz, 3 H), 2.77 - 2.95 (m, 1 H), 3.34 - 3.90 (m, 2 H), 4.46 - 5.25 (m, 3 H), 5.32 - 5.68 (m, 1 H), 5.71 - 5.81 (m, 1 H), 6.21 (dd, J = 16.5, 2.4 Hz, 1 H), 6.87 (dd, J = 16.7, 10.5 Hz, 1 H), 7.20 (d, J = 4.8 Hz, 1 H), 7.27 - 7.42 (m, 3 H), 7.50 - 7.64 (m, 1 H), 8.41 (d, J = 4.8 Hz, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –113.7 (s, 1 F); LRMS: (ESI, +ve ion) 576.2 m/z $(M+H)^+$.

[0601] Step 2: Chiral separation of 4-((3*R*,5*S*)-4-acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridin-2(1*H*)-one (Examples 224-1 and 224-2). 4-((3*R*,5*S*)-4-Acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridin-2(1*H*)-one (**Example 224**, 170 mg, 0.29 mmol) was dissolved in MeOH (8.0 mL) and DCM (2.0 mL) and purified by preparative SFC (Chiralcel AS-H column (5 um, 21 x 250 mm), F = 80 mL, 15% *i*PrOH/85% CO_2) to afford 4-((3*R*,5*S*)-4-acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridin-2(1*H*)-one (**Example 224-1**, 50 mg, 0.089 mmol, 61% yield) as an orange solid (single isomer, 1st-eluting isomer) and 4-((3*R*,5*S*)-4-acryloyl-3,5-dimethylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridin-2(1*H*)-one (53 mg, 0.093 mmol, 64% yield) as an orange solid (single isomer, 2nd-eluting isomer). **Example 224-1**: ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.96 (br d, J = 6.4 Hz, 3 H), 1.07 (d, J = 6.6 Hz, 3 H), 1.37 (br s, 6 H), 1.99 (s, 3 H), 2.86 (sept, J = 6.6 Hz, 1 H), 3.31 - 5.71 (m, 6 H), 5.73 - 5.85 (m, 1 H), 6.21 (dd, J = 16.5, 2.4 Hz, 1 H), 6.87 (dd, J = 16.6, 10.4 Hz, 1 H), 7.20 (d, J = 4.8 Hz, 1 H), 7.27 - 7.42 (m, 3 H), 7.51 - 7.66 (m, 1 H), 8.41 (d, J = 5.0 Hz, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –113.68 (s, 1 F); LRMS: (ESI, +ve ion) 576.2 *m/z* $(M+H)^+$. **Example 224-2**: ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.96 (br d, J = 6.4 Hz, 3 H), 1.07 (d, J = 6.8 Hz, 3 H), 1.37 (br s, 6 H), 1.99 (s, 3 H), 2.86 (sept, J = 6.6 Hz, 1 H), 3.38 - 3.72 (m, 2 H), 4.59 - 4.79 (m, 2 H), 4.85 - 5.15 (m, 1 H), 5.43 - 5.68 (m, 1 H), 5.70 - 5.80 (m, 1 H), 6.21 (dd, J = 16.5, 2.4 Hz, 1 H), 6.87 (dd, J = 16.7, 10.5 Hz, 1 H), 7.20 (d, J = 5.0 Hz, 1 H), 7.26 - 7.42 (m, 3 H), 7.51 - 7.61 (m, 1 H), 8.41 (d, J = 4.8 Hz, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –113.68 (s, 1 F); LRMS: (ESI, +ve ion) 576.2 *m/z* $(M+H)^+$.

Example 225:

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-6-methylpteridin-2(1*H*)-one (Examples 225, 225-1, and 225-2)



[0602] Step 1: 7-(2-Fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-6-methylpteridine-2,4(1*H*,3*H*)-dione. A nitrogen-sparged mixture of 6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridine-2,4(1*H*,3*H*)-dione (**Intermediate 111A**, 500 mg, 1.2 mmol), trimethylboroxine (0.33 mL, 2.3 mmol), potassium carbonate (330 mg, 2.3 mmol), and $Pd(dppf)Cl_2$ (86 mg, 0.12 mmol) in 1,4-dioxane (5.0 mL)/water (0.50 mL) was heated in a Biotage Initiator+ microwave at 100 °C for 1.5 h. The reaction mixture was diluted with water (50 mL) and extracted with EtOAc (2 × 50 mL). Sat. aq. sodium bicarbonate (30 mL) was added to the aqueous phase which was further extracted with EtOAc (2 × 50 mL). The combined organic extracts were dried over anhydrous magnesium suflate, filtered, and concentrated *in vacuo* to give crude 7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-6-methylpteridine-2,4(1*H*,3*H*)-dione as a burgendy-colored foam which was carried forward in the next step without purification, assuming theoretical yield.

[0603] Step 2: 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-6-methylpteridin-2(1*H*)-one. A room-temperature mixture of crude 7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-6-methylpteridine-2,4(1*H*,3*H*)-dione (480 mg, 1.2 mmol) and DIPEA (0.41 mL, 2.3 mmol) in acetonitrile (7.8

mL) was treated with phosphorus oxychloride (0.22 mL, 2.3 mmol). The resulting homogenous brown mixture was heated at 80 °C for 2 h, then additional phosphorus oxychloride (0.10 mL, 1.1 mmol) was added and the dark brown mixture heated at 80 °C for 2 h, after which time the reaction mixture was cooled to 0 °C. 1-((2*R*,5*S*)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one 2,2,2-trifluoroacetate (**Intermediate 160**, 2.1 g, 2.9 mmol) was added as a solution in acetonitrile (5.0 mL) and the reaction mixture stirred slowly warming to room temp overnight. The reaction mixture was diluted with EtOAc (30 mL) and washed with 50% saturated aqueous ammonium chloride (100 mL). The layers were partitioned and the aqueous phase was washed with EtOAc (2 × 30 mL). The combined organic extracts were dried over anhydrous magnesium sulfate, filtered, and concentrated to afford a red-brown residue that was purified by silica gel chromatography (eluent: 0–100% 3:1 EtOAc–EtOH/heptane) and preparative SFC (Step I, Princeton DEAP (5 um, 21 x 250 mm), F = 80mL, 15% MeOH/85% CO_2, back pressure = 90 bars, 1.0 mL injection; Step II, Welko RR (5 um, 21 x 250 mm), F = 80mL, 35% MeOH/65% CO_2, back pressure = 90 bars, 1.0 mL injection) to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-6-methylpteridin-2(1*H*)-one (**Example 225**, 130 mg, 0.23 mmol) as a mixture of atropisomers. ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.87 - 0.99 (m, 3 H), 1.08 (br d, *J* = 6.6 Hz, 3 H), 1.16 - 1.50 (m, 6 H), 1.95 (br d, *J* = 17.6 Hz, 3 H), 2.45 (s, 3 H), 3.65 - 3.84 (m, 1 H), 3.95 (br d, *J* = 13.3 Hz, 1 H), 4.28 (br d, *J* = 13.3 Hz, 1 H), 4.52 - 4.63 (m, 1 H), 4.77 - 4.98 (m, 1 H), 5.15 - 5.38 (m, 1 H), 5.75 (dt, *J* = 10.3, 2.6 Hz, 1 H), 5.85 - 6.04 (m, 1 H), 6.19 (br d, *J* = 17.4 Hz, 1 H), 6.78 - 6.97 (m, 1 H), 7.15 - 7.23 (m, 1 H), 7.24 - 7.39 (m, 3 H), 7.48 - 7.59 (m, 1 H), 8.40 (d, *J* = 4.8 Hz, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ –115.43 - –115.26 (m, 1 F); LRMS: (ESI, +ve ion) 555.6 *m/z* $(M+H)^+$.

[0604] Step 3: Chiral separation of 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-6-methylpteridin-2(1*H*)-one (Examples 225-1 and 225-2). 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-6-methylpteridin-2(1*H*)-one (**Example 225**, 120 mg, 0.22 mmol) was dissolved in 1:1 DCM–MeOH (25 mL) and purified by preparative SFC (Chiralcel AD-H (5 um, 21 x 250 mm), F = 80mL, 20% *i*PrOH/80% CO_2, back pressure = 90 bars, 2.0 mL injections) to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-6-methylpteridin-2(1*H*)-one (**Example 225-1**, 42 mg, 0.076 mmol, 73% yield) as a yellow-tan solid (single isomer, 1st-eluting peak) and 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-

7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-6-methylpteridin-2(1*H*)-one (**Example 225-2**, 42 mg, 0.076 mmol, 73% yield) as a yellow-tan solid (single isomer, 2nd-eluting peak). **Example 225-1**: ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.95 (d, J = 6.6 Hz, 3 H), 1.00 - 1.12 (m, 3 H), 1.18 - 1.51 (m, 6 H), 1.97 (br s, 3 H), 2.45 (s, 3 H), 2.64 - 2.75 (m, 1 H), 3.31 - 3.56 (m, 1 H), 3.67 - 3.83 (m, 1 H), 3.95 (br d, J = 13.5 Hz, 0.5 H), 4.24 - 4.33 (m, 0.5 H), 4.59 (br s, 0.5 H), 4.79 - 4.96 (m, 1 H), 5.16 - 5.38 (m, 0.5 H), 5.69 - 5.80 (m, 1 H), 5.84 - 6.04 (m, 0.5 H), 6.19 (br d, J = 16.8 Hz, 1.5 H), 6.76 - 6.98 (m, 1 H), 7.19 (d, J = 4.8 Hz, 1 H), 7.25 - 7.39 (m, 3 H), 7.50 - 7.58 (m, 1 H), 8.40 (d, J = 5.0 Hz, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ −115.32 - −115.26 (m, 1 F); LRMS: (ESI, +ve ion) 556.3 *m/z* (M+ H)$^+$. **Example 225-2**: ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.93 (br d, J = 6.2 Hz, 3 H), 1.08 (d, J = 6.8 Hz, 3 H), 1.20 - 1.50 (m, 6 H), 1.93 (br s, 3 H), 2.45 (s, 3 H), 2.77 - 2.86 (m, 1 H), 3.31 - 3.55 (m, 1 H), 3.64 - 3.83 (m, 1 H), 3.95 (br d, J = 14.3 Hz, 0.5 H), 4.20 - 4.35 (m, 0.5 H), 4.58 (br s, 0.5 H), 4.75 - 4.95 (m, 1 H), 5.15 - 5.37 (m, 0.5 H), 5.69 - 5.80 (m, 1 H), 5.86 - 6.04 (m, 0.5 H), 6.19 (br d, J = 16.8 Hz, 1.5 H), 6.77 - 6.97 (m, 1 H), 7.20 (d, J = 5.0 Hz, 1 H), 7.27 - 7.38 (m, 3 H), 7.49 - 7.58 (m, 1 H), 8.40 (d, J = 4.8 Hz, 1 H); ^{19}F{^{1}H} NMR (376 MHz, DMSO-d_6) δ −115.42 - −115.36 (m, 1 F); LRMS: (ESI, +ve ion) 556.3 *m/z* (M+ H)$^+$.

Example 226

***(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(5-ethynyl-2-fluoro-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one.**



[0605] Step 1: *(M)*- 7-(5-Bromo-2-fluorophenyl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A mixture of (*M*)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione

(**Intermediate 73B**, 300 mg, 0.82 mmol), tetrakis(triphenylphosphine)palladium (285 mg, 0.25 mmol), 2-(5-bromo-2-fluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (494 mg, 1.64 mmol) and sodium carbonate(435 mg, 4.11 mmol) in 1,4-dioxane (3 mL)/ water (1.5 mL) was stirred at 90 °C for 1 h. The reaction mixture was diluted with EtOAc (10 mL)/water (10 mL). The aqueous was added sat'd sodium chloride (5 mL) and extracted with EtOAc (10 mLx2). The combined organic extracts were dried over $MgSO_4$. The solution was filtered and concentrated *in vacuo* to give the crude material. The crude material brown oil was purified by scilica gel chromatography (eluent: 0-50% of EtOAc/heptane) to provide (*M*)-7-(5-bromo-2-fluorophenyl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (229 mg, 0.45 mmol, 55.3 % yield) as a light yellow foam. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.25 (br s, 1 H), 8.59 (s, 1 H), 8.44 (d, *J*=4.8 Hz, 1 H), 7.70 (ddd, *J*=8.8, 4.6, 2.6 Hz, 1 H), 7.43 (dd, *J*=6.2, 2.5 Hz, 1 H), 7.32 (t, *J*=9.2 Hz, 1 H), 7.22 (d, *J*=5.0 Hz, 1 H), 2.91 (quin, *J*=6.6 Hz, 1 H), 2.03 (s, 3 H), 1.04 - 1.12 (m, 3 H), 0.97 (d, *J*=6.6 Hz, 3 H). ^{19}F NMR (376 MHz, DMSO-d_6) δ -116.25 (s, 1 F). *m/z* (ESI, +ve ion): 442.2 $(M+H)^+$.

[0606] Step 2: *(M)*- 6-Chloro-7-(2-fluoro-5-((trimethylsilyl)ethynyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A mixture of (*M*)-7-(5-bromo-2-fluorophenyl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (182 mg, 0.36 mmol), (trimethylsilyl)acetylene (0.7 mL, 0.47 mmol), triethylamine, anhydrous (0.6 mL, 4.27 mmol), copper(i) iodide (21 mg, 0.11 mmol) and *trans*-dichlorobis(triphenyl-phosphine)palladium (ii) (76 mg, 0.11 mmol) in ethyl acetate (1.0 mL) was stirred at 50 °C for 18 h. The reaction mixture was concentrated *in vacuo* and purified by scilica gel chromatography (eluent: 0-50% of EtOAc/heptane) to provide (*M*)-6-chloro-7-(2-fluoro-5-((trimethylsilyl)ethynyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (60 mg, 0.12 mmol, 31.9 % yield) as a light yellow foam. *m/z* (ESI, +ve ion): 521.2 $(M+H)^+$.

[0607] Step 3: *(M)*-6-Chloro-7-(2,2-dimethyl-1-piperidyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one. A mixture of (*M*)-6-chloro-7-(2-fluoro-5-((trimethylsilyl)ethynyl)phenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (60 mg, 0.12 mmol) and potassium carbonate (32 mg, 0.23 mmol) in methanol (1 mL) was stirred at 25 °C for 30 min. The reaction mixture was diluted with satd NH_4Cl (3 mL) and extracted with EtOAc (2 x 10 mL). The combined organic extracts were dried over $MgSO_4$. The solution was filtered and concentrated *in vacuo* to give (*M*)-6-chloro-7-(5-ethynyl-2-fluorophenyl)-1-(2-isopropyl-

4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (47 mg, 0.11 mmol, 91 % yield) as a brown foam. *m/z* (ESI, +ve ion): 449.0 $(M+H)^+$.

[0608] Step 4: *(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(5-ethynyl-2-fluoro-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one. A mixture of (*M*)-6-chloro-7-(5-ethynyl-2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (47 mg, 0.11 mmol) and n,n'-diisopropylethylamine (0.1 mL, 0.52 mmol) in acetonitrile (1 mL) was treated with phosphorous oxychloride (0.03 mL, 0.31 mmol) at room temperature. The mixture was stirred at 80 °C for 1 h. The reaction mixture was evaporated to dryness to give (*M*)-4,6-dichloro-7-(5-ethynyl-2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a brown solid. *m/z* (ESI, +ve ion): 467.0 $(M+H)^+$. The resulting brown solid was used in next step without purification.

[0609] To the mixture of (*M*)-4,6-dichloro-7-(5-ethynyl-2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (0.1 mL, 0.52 mmol) in acetonitrile (1 mL) was added a mixture of 1-((2*R*,5*S*)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one TFA salt (148 mg, 0.524 mmol) and DIPEA (0.1 mL, 0.52 mmol) in acetonitrile (0.5 mL) and stirred at rt for 5 min. The resulting mixture was evaporated to dryness. The resulting crude product was purified by by scilica gel chromatography (eluent: 0-100% of MeOH-EtOAc (5:95)/heptane) to provide *(M)*-6-Chloro-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-7-(5-ethynyl-2-fluoro-phenyl)-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one (19 mg, 30% yield) as an yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.50 (d, *J*=2.7 Hz, 1 H), 8.43 (d, *J*=4.8 Hz, 1 H), 7.64 (ddd, *J*=8.7, 4.9, 2.2 Hz, 1 H), 7.31 - 7.42 (m, 2 H), 7.24 (d, *J*=5.0 Hz, 1 H), 6.75 - 6.95 (m, 1 H), 6.20 (dd, *J*=16.7, 2.4 Hz, 1 H), 5.69 - 5.85 (m, 1 H), 4.83 - 4.95 (m, 1 H), 4.40 - 4.81 (m, 1 H), 4.25 (s, 1 H), 3.79 - 4.22 (m, 4 H), 2.69 - 2.83 (m, 1 H), 1.96 (s, 3 H), 1.30 - 1.38 (m, 3 H), 1.14 - 1.28 (m, 4 H), 1.08 (d, *J*=6.6 Hz, 3 H), 1.00 (d, *J*=6.6 Hz, 3 H). ^{19}F NMR (376 MHz, DMSO-d_6) δ -111.42 (s, 1 F). *m/z* (ESI, +ve ion): 599.2 $(M+H)^+$.

Example 227

***(M)*-6-Chloro-7-(2,2-dimethyl-1-piperidyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one.**



[0610] Step 1: *(M)*-6-Chloro-7-(2,2-dimethylpiperidin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A glass microwave reaction vessel was charged with 2,2-dimethylpiperidine (511 mg, 4.52 mmol), 6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 73B**, 550 mg, 1.51 mmol) and DIPEA (1.3 mL, 7.53 mmol) in DMF (1.0 mL). The reaction mixture was stirred and heated in an Initiator microwave reactor (Personal Chemistry, Biotage AB, Inc., Upssala, Sweden) at 110 °C for 6 h and then 120 °C for 3 h. The reaction mixture was diluted with water (5 mL) and extracted with EtOAc (3 x 5 mL). The organic extracts were combined and dried over $MgSO_4$. The solution was filtered and concentrated *in vacuo* to give the crude material as a brown solid. The crude material brown solid was purified by silica gel chromatography (eluent: 0-50% of EtOAc/heptane) to provide *(M)*-6-chloro-7-(2,2-dimethylpiperidin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (276 mg, 0.624 mmol, 41.5 % yield) as a light yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.90 (s, 1 H), 8.49 (d, *J*=4.8 Hz, 1 H), 8.18 (s, 1 H), 7.28 (d, *J*=4.8 Hz, 1 H), 3.10 - 3.23 (m, 1 H), 2.91 - 3.02 (m, 1 H), 2.74 - 2.86 (m, 1 H), 2.02 (s, 3 H), 1.43 - 1.61 (m, 4 H), 1.31 (br t, *J*=4.8 Hz, 2 H), 1.08 (d, *J*=6.6 Hz, 3 H), 1.03 (d, *J*=6.6 Hz, 3 H), 0.81 (s, 3 H), 0.69 (s, 3 H). *m/z* (ESI, +ve ion): 442.2 $(M+H)^+$.

[0611] Step 2: *(M)*-6-Chloro-7-(2,2-dimethyl-1-piperidyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one. A mixture of *(M)*-6-chloro-7-(2,2-dimethylpiperidin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (252 mg, 0.570 mmol) and DIPEA (0.5 mL, 2.85 mmol) in acetonitrile (2.0 mL) was treated with phosphorous oxychloride (0.2 mL, 1.71 mmol) at room temperature. The mixture was stirred at 80 °C for 1 h. The reaction mixture was evaporated to dryness to give (*M*)-4,6-dichloro-7-(2,2-dimethylpiperidin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a brown solid. *m/z* (ESI, +ve ion): 460.2 $(M+H)^+$. The resulting brown solid was used in next step without purification.

[0612] The mixture of (*M*)-4,6-dichloro-7-(2,2-dimethylpiperidin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (0.5 mL, 2.85 mmol) in acetonitrile (2.0 mL) was added a solution of 1-((2*R*,5*S*)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one TFA salt (483 mg, 1.71 mmol) and DIPEA (0.5 mL, 2.85 mmol) in MeCN (1.0 mL) and stirred at rt for 5 min. The resulting mixture was diluted with water (5 mL) and extracted with EtOAc (3 x 10 mL). The organic extracts were combined and dried over $MgSO_4$. The solution was filtered and concentrated *in vacuo* to give the crude material as a brown solid. The crude material brown solid which was absorbed onto a plug of silica gel and purified by silica gel chromatography (eluent: 0-100% of MeOH-EtOAc (5:95)/heptane) to provide *(M)*-6-chloro-7-(2,2-dimethyl-1-piperidyl)-4-[(2*S*,5*R*)-2,5-dimethyl-4-prop-2-enoyl-piperazin-1-yl]-1-(2-isopropyl-4-methyl-3-pyridyl)pyrido[2,3-*d*]pyrimidin-2-one (134 mg, 0.23 mmol, 39.7 % yield) as a light yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.49 (br t, *J*=5.3 Hz, 1 H), 7.85 - 8.17 (m, 1 H), 7.32 (br s, 1 H), 6.67 - 6.93 (m, 1 H), 6.17 (br d, *J*=16.6 Hz, 1 H), 5.73 (br d, *J*=8.1 Hz, 1 H), 4.36 - 4.95 (m, 3 H), 3.67 - 4.17 (m, 4 H), 2.80 - 3.45 (m, 2 H), 2.60 - 2.77 (m, 1 H), 1.94 (br d, *J*=7.7 Hz, 3 H), 1.47 - 1.67 (m, 6 H), 1.19 - 1.34 (m, 6 H), 1.12 (tt, *J*=12.4, 6.3 Hz, 6 H), 1.01 (br d, *J*=6.4 Hz, 3 H), 0.70 - 0.84 (m, 2 H). *m/z* (ESI, +ve ion): 592.2 $(M+H)^+$.

Example 228

4-((*S*)-4-Acryloyl-2-methylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(5-methyl-2-oxoindolin-4-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0613] **Step 1: (*M*)-*tert*-Butyl (*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate**. A 250-mL round-bottomed flask was charged with (*M*)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 73B**, 5.9 g, 16 mmol) and DIPEA (4.2 mL, 24 mmol) in acetonitrile (65 mL) followed by phosphorous oxychloride (1.8 mL, 19.2 mmol). The mixture was stirred at 80 °C for 1 h. The reaction mixture was concentrated in vacuo to give a brown solid. The crude solid was used in the next step without purification. *m/z* (ESI, +ve ion): 383.0 $(M+H)^+$.

[0614] The above crude solid and DIPEA (4.2 mL, 24 mmol) in DMF (50 mL) was treated with (3*S*)-1-(*tert*-butoxycarbonyl)-3-methylpiperazine (4.81 g, 24 mmol, Ark Pharm, Inc.) and stirred at rt for 30 min. The resulting mixture was added to ice water (100 mL) and stirred for 15 min. The resulting precipitate was collected by filtration, washed with water, and dried to give (*M*)-*tert*-butyl (*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-

dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (**Intermediate 73C**, 7.71 g, 14.1 mmol, 88 % yield). ^{1}H NMR (400 MHz, DMSO-d_6) δ ppm 8.39 - 8.54 (m, 2 H), 7.25 (d, *J*=5.0 Hz, 1 H), 4.87 (br d, *J*=1.0 Hz, 1 H), 4.15 (br d, *J*=13.3 Hz, 1 H), 3.95 (br dd, *J*=5.9, 4.0 Hz, 1 H), 3.83 (br d, *J*=13.3 Hz, 1 H), 3.71 (br t, *J*=11.0 Hz, 1 H), 2.96 - 3.27 (m, 2 H), 2.57 - 2.64 (m, 1 H), 1.94 (s, 3 H), 1.45 (s, 9 H), 1.32 (d, *J*=6.6 Hz, 3 H), 1.06 (d, *J*=6.8 Hz, 3 H), 1.01 (d, *J*=6.6 Hz, 3 H). *m/z* (ESI, +ve ion): 547.3 $(M+H)^+$.

[0615] Step 2: (*M*,*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of (*M*)-*tert*-butyl (*S*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (**Intermediate 73C**, 2.71 g, 4.95 mmol) in DCM (25 mL) was treated with TFA (10 mL, 134 mmol) at rt and stirred for 20 min. The reaction was concentrated in vacuo to afford (*M*,*S*)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-4-(2-methylpiperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a brown solid. *m/z* (ESI, +ve ion): 447.2 $(M+H)^+$.

[0616] A mixture of (*M*,*S*)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-4-(2-methylpiperazin-1-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (4.3 mL, 24.8 mmol) in DCM (25 mL) was treated with acryloyl chloride (0.36 mL, 4.5 mmol) at 0 °C and stirred for 20 min. The reaction mixture was concentrated in vacuo. The crude product was purified by silica gel chromatography (eluent: 0-60% EtOAc-EtOH(3:1)/heptane) to provide pure (*M*,*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 73D**; 2.28 g, 4.55 mmol, 92 % yield) as a light-yellow foam. ^{1}H NMR (400 MHz, *DMSO-d_6*) δ ppm 8.50 (d, *J*=4.8 Hz, 2 H), 7.30 (d, *J*=4.8 Hz, 1 H), 6.77 - 6.94 (m, 1 H), 6.20 (br d, *J*=16.0 Hz, 1 H), 5.76 (dd, *J*=10.5, 2.4 Hz, 1 H), 4.92 (br s, 1 H), 3.96 - 4.44 (m, 3 H), 3.54 - 3.85 (m, 2 H), 3.00 - 3.24 (m, 1 H), 2.59 - 2.70 (m, 1 H), 1.97 (s, 3 H), 1.30 (br d, *J*=6.6 Hz, 3 H), 1.07 (d, *J*=6.6 Hz, 3 H), 1.02 (d, *J*=6.6 Hz, 3 H). *m/z* (ESI, +ve ion): 501.2 $(M+H)^+$.

[0617] Step 3: 4-((*S*)-4-Acryloyl-2-methylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(5-methyl-2-oxoindolin-4-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a pressure vial was charged (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 73D**, 0.2 g, 0.399 mmol), 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (**Intermediate 209**, 0.163 g, 0.598 mmol), tetrakis(triphenylphosphine)palladium(0) (0.046 g, 0.040 mmol), sodium carbonate (0.127 g, 1.20 mmol), and 1,4-dioxane (5 mL). The vial was

purged with N_2 for 3 min, sealed and then heated at 100°C for 24 h. LCMS showed mostly SM. To the reaction mixture was added potassium acetate (0.117 g, 1.20 mmol),Pd(dppf)Cl_2 CH_2Cl_2 adduct (0.033 g, 0.040 mmol) and water (0.3 mL). The reaction was heated at 90°C for 1.5 h. The reaction mixture was partitioned between EtOAc and water. The separated organic layer was washed with brine, dried over Na_2SO_4 and then filtered. The filtrate was concentrated and purified by silica gel chromatography (0-90% EtOAc/heptane) to afford 4-((*S*)-4-acryloyl-2-methylpiperazin-1-yl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)-7-(5-methyl-2-oxoindolin-4-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.042 g, 0.069 mmol, 17.2 % yield) as light brown solid. ^{1}H NMR (CHLOROFORM-*d*) δ: 8.47 (d, *J*=4.8 Hz, 1H), 8.14 (s, 1H), 7.70 (br s, 1H), 7.03-7.13 (m, 2H), 6.79 (d, *J*=7.9 Hz, 1H), 6.62 (br s, 1H), 6.38-6.46 (m, 1H), 5.80-5.85 (m, 1H), 4.32-5.14 (m, 3H), 3.54-4.10 (m, 3H), 3.09-3.39 (m, 1H), 2.88-3.06 (m, 2H), 2.70-2.84 (m, 1H), 2.00 (s, 6H), 1.53 (br d, *J*=0.8 Hz, 3H), 1.22 (d, *J*=6.8 Hz, 3H), 1.03 (br d, *J*=6.6 Hz, 3H). *m/z* (ESI, +ve ion): 612.0 $(M+H)^+$.

Example 229

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0618] Step 1. 1-(2-Amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a pressure vial was charged 1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 231**, 0.28 g, 0.53 mmol), copper (I) oxide (3.76 mg, 0.158 mmol), ammonium hydroxide solution, 28.0-30%, NH_3 basis (3.57 ml, 26.3 mmol) and methanol (3.5 mL). The vial was sealed and heated at 70 °C for 19 h. The product mixture

was concentrated to dryness. The resulted solid residue was dissolved in EtOAc and partitioned with saturated NH_4Cl. The separated organic layer was washed with water and brine, dried over Na_2SO_4 and then filtered. The filtrate was concentrated and purified by silica gel chromatography to afford 1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.164 g, 0.350 mmol, 66 % yield) as a white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.86 (d, *J*=6.63 Hz, 6 H) 1.02 (d, *J*=6.63 Hz, 6 H) 2.68 (quin, *J*=6.69 Hz, 2 H) 6.58 (s, 2 H) 7.21 (td, *J*=7.52, 1.55 Hz, 1 H) 7.28 - 7.35 (m, 2 H) 7.48 - 7.56 (m, 1 H) 8.54 (s, 1 H) 12.15 (s, 1 H).

[0619] Step 2. *tert*-Butyl (2*R*,5*S*)-4-(1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a mixture of 1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.100 g, 0.213 mmol) and *N*-ethyl-*N*-isopropylpropan-2-amine (0.111 ml, 0.640 mmol) in acetonitrile (5 ml) was added phosphoryl trichloride (0.040 ml, 0.427 mmol) at RT and the reaction was heated at 80°C for 30 min. Additional phosphoryl trichloride (0.040 ml, 0.427 mmol) was added and heated for an additional 30 min. The mixture was concentrated to give the crude 1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-4,6-dichloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as brown oil which was used in the next step without purification.

[0620] Crude 1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-4,6-dichloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one was dissolvd in DMF (2 mL). To this solution was added *N*-ethyl-*N*-isopropylpropan-2-amine (0.111 ml, 0.640 mmol) followed by (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (0.050 ml, 0.235 mmol, eNovation Chemicals, LLC). The mixture was stirred at RT for 1 h. Water was added to quench the reaction, and the resulting suspension was filtered. The recovered solid was purified by silica gel chromatography (0-100% (3:1 EtOAc:EtOH)/heptane) to afford *tert*-butyl (2*R*,5*S*)-4-(1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.05 g, 0.075 mmol, 35 % yield) as light yellow solid. *m/z* (ESI, +ve ion): 665.2 $(M+H)^+$.

[0621] Step 3. 4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. To a solution of *tert*-butyl (2*R*,5*S*)-4-(1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.050 g, 0.075 mmol) in DCM (3 mL) was added TFA (0.5

mL, 6.49 mmol). The mixture was stirred at RT for 30 min. The mixture was concentrated to afford 1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as light yellow oil that was used without purification.

[0622] Crude 1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one was dissolved in DCM (3 mL) and cooled with an ice bath. To this solution was added diisopropylethylamine (0.10 mL, 0.57 mmol) followed by acryloyl chloride. 0.258 M in DCM (0.262 mL, 0.068 mmol). The reaction was stirred at 0°C for 10 min, then concentrated to remove the solvent. The residue was purified by silica gel chromatography (0-60% (3:1 EtOAc:EtOH)/heptane) to afford 4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-1-(2-amino-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (0.024 g, 0.039 mmol, 52 % yield) as an off-white solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 0.86 (dd, *J*=6.84, 3.11 Hz, 6 H) 1.00 (dd, *J*=6.63, 4.35 Hz, 6 H) 1.25 (br s, 3 H) 1.32 (t, *J*=6.84 Hz, 3 H) 2.40 - 2.48 (m, 2 H) 3.31 - 3.53 (m, 1 H) 3.81 - 3.92 (m, 2 H) 4.10 - 4.19 (m, 1 H) 4.43 - 4.93 (m, 2 H) 5.71 - 5.80 (m, 1 H) 6.18 (dd, *J*=16.69, 2.18 Hz, 1 H) 6.50 (s, 2 H) 6.83 (td, *J*=16.17, 10.57 Hz, 1 H) 7.21 - 7.27 (m, 1 H) 7.28 - 7.36 (m, 2 H) 7.50 - 7.57 (m, 1 H) 8.44 (s, 1 H). *m/z* (ESI, +ve ion): 618.6 $(M+H)^+$.

Section 3—Synthesis of Intermediates

Intermediate I-5

4-Isopropyl-6-methylpyrimidin-5-amine



[0623] **Step 1: 4-Chloro-6-(prop-1-en-2-yl)pyrimidin-5-amine**. A mixture of 4,6-dichloro-5-aminopyrimidine (5.00 g, 30.5 mmol), 2-isopropenylboronic acid, pincol ester (6.15 g, 36.6 mmol, Combi-Blocks, San Diego, CA), $Pd(dppf)Cl_2$ (2.23 g, 3.05 mmol), and sodium carbonate (9.69 g, 91 mmol) in a mixture of 1,4-dioxane (40 mL) and water (10 mL) was sparged with N_2 then heated at 95 °C for 2 h. The mixture was cooled to rt, satd. aq. $NaHCO_3$ was added, and the resulting mixture was extracted with EtOAc. The combined extracts were concentrated *in vacuo*, and the residue was purified by silica gel chromatography (eluent: 0-

40% EtOAc/heptane) to provide 4-chloro-6-(prop-1-en-2-yl)pyrimidin-5-amine (1.24 g, 7.31 mmol, 24 % yield) as a yellow oil. *m/z* (ESI, +ve ion): 170.0 $(M+H)^+$.

[0624] **Step 2: 4-Methyl-6-(prop-1-en-2-yl)pyrimidin-5-amine**. A mixture of 4-chloro-6-(prop-1-en-2-yl)pyrimidin-5-amine (2.0 g, 11.79 mmol), methylboronic acid (3.53 g, 59.0 mmol, Combi-Blocks, San Diego, CA), $Pd(dppf)Cl_2$ (0.863 g, 1.179 mmol) and sodium carbonate (6.25 g, 59.0 mmol) in 1,4-dioxane (40 mL) and water (10 mL) was sparged with N_2, then heated at 95 °C for 17 h. The mixture was then cooled to rt, diluted with satd. aq. $NaHCO_3$, and extracted with EtOAc. The combined extracts were concentrated *in vacuo*, and the residue was purified by silica gel chromatography (eluent: 0-5% MeOH/DCM) to provide 4-methyl-6-(prop-1-en-2-yl)pyrimidin-5-amine (1.3 g, 8.71 mmol, 73.9 % yield). *m/z* (ESI, +ve ion): 150.2 $(M+H)^+$.

[0625] **Step 3: 4-Isopropyl-6-methylpyrimidin-5-amine (Intermediate I-5)**. Palladium 10 wt. % on activated carbon (0.522 g, 0.491 mmol) was added to a solution of 4-methyl-6-(prop-1-en-2-yl)pyrimidin-5-amine (1.464 g, 9.81 mmol) in EtOH (30 mL), and the resulting suspension was stirred under hydrogen gas (30 psi) for 1.5 h. The resulting mixture was subsequently filtered through Celite, and the filtrate was concentrated *in vacuo* to provide 4-isopropyl-6-methylpyrimidin-5-amine (**Intermediate I-5**, 1.48 g, 9.79 mmol, 100 % yield) as a brown oil. ^1H NMR (400 MHz, *DMSO-d_6*) δ: 7.96-8.30 (m, 1H), 4.94-5.11 (m, 2H), 3.12-3.27 (m, 1H), 2.22-2.31 (m, 3H), 1.13-1.16 (m, 6H). *m/z* (ESI, +ve ion): 152.2 $(M+H)^+$.

Intermediate I-33

6-Isopropyl-N^2,N^2-bis(4-methoxybenzyl)-4-methylpyridine-2,5-diamine



[0626] **Step 1: 2-(Bis(4-methoxybenzyl)amino)-4-methyl-5-nitropyridine 1-oxide.** A mixture of 2-chloro-4-methyl-5-nitropyridine 1-oxide (2.3 g, 12.20 mmol; product of step 2 in

preparation of **Intermediate I-32**), bis(4-methoxybenzyl)-amine (3.45 g, 13.42 mmol), and sodium carbonate (anhydrous, powder; 2.6 g, 24.39 mmol) in toluene (70 mL) was stirred at 70 °C for 16 h. Additional sodium carbonate (anhydrous, powder; 2.6 g, 24.39 mmol) and bis(4-methoxybenzyl)-amine (3.45 g, 13.42 mmol) were added, and the resulting mixture was stirred at 90 °C for 2 h. The reaction mixture was then filtered, and the filter cake washed with DCM (3 x 100 mL). The combined filtrates were concentrated *in vacuo* and the residue chromatographically purified (silica gel, 0%-10% MeOH/DCM) to provide 2-(bis(4-methoxybenzyl)amino)-4-methyl-5-nitropyridine 1-oxide (2.75 g, 6.72 mmol, 55 % yield) as a brown solid. ^{1}H NMR (*CDCl$_3$*) δ: 9.11 (s, 1H), 7.26 (d, J=8.7 Hz, 4H), 6.92 (d, J=8.7 Hz, 4H), 6.60 (s, 1H), 4.77 (s, 4H), 3.87 (s, 6H), 2.59 (s, 3H). m/z (ESI, +ve ion): 410.0 $(M+H)^+$.

[0627] Step 2: 6-Chloro-*N*,*N*-bis(4-methoxybenzyl)-4-methyl-5-nitropyridin-2-amine. Phosphorus oxychloride (12.29 ml, 132 mmol) was added, dropwise, to a solution of 2-(bis(4-methoxybenzyl)amino)-4-methyl-5-nitropyridine 1-oxide (2.7 g, 6.59 mmol) in *N,N'*-diisopropylethylamine (27.6 mL, 158 mmol) at 0 °C , and the resulting mixture was heated to 70 °C and stirred for 10 mins. The reaction mixture was then concentrated *in vacuo* and the residue diluted with ice water (20 mL) and EtOAc (100 mL). The resulting mixture was stirred as saturated aq. $NaHCO_3$ (50 mL) was slowly added until a pH of 10-11 was achieved. The resulting mixture was then stirred at room temperature for 30 mins before the organic layer was separated, dried over MgSO4, and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc/heptane) provided 6-chloro-*N,N*-bis(4-methoxybenzyl)-4-methyl-5-nitropyridin-2-amine (1.62 g, 3.77 mmol, 57 % yield) as a yellow solid. ^{1}H NMR (METHANOL-d4) δ: 7.15 (br d, J=8.7 Hz, 4H), 6.87 (d, J=8.7 Hz, 4H), 6.44 (s, 1H), 4.72 (s, 4H), 3.77 (s, 6H), 2.21 (s, 3H).. m/z (ESI, +ve ion): 427.9 $(M+H)^+$.

[0628] Step 3: *N*,*N*-Bis(4-Methoxybenzyl)-4-methyl-5-nitro-6-(prop-1-en-2-yl)pyridin-2-amine. A mixture of 6-chloro-*N,N*-bis(4-methoxybenzyl)-4-methyl-5-nitropyridin-2-amine (1.59 g, 3.72 mmol), 2-isopropenylboronic acid, pincol ester (1.873 g, 11.15 mmol), cesium carbonate (3.63 g, 11.15 mmol), and *trans*-dichlorobis(triphenyl-phosphine)palladium (II) (0.261 g, 0.372 mmol) in 1,2-dimethoxyethane (20 mL) and water (4.00 mL) was sparged with $N_{2(g)}$ for 5 min and then stirred at 80 °C for 3 h. Additional 2-isopropenylboronic acid, pincol ester (1.873 g, 11.15 mmol), cesium carbonate (3.63 g, 11.2 mmol), and *trans*-dichlorobis(triphenyl-phosphine)palladium (II) (0.261 g, 0.372 mmol) were then added, and the resulting mixture was stirred at 80 °C for 1 hour. After cooling to rt, saturated aq. $NaHCO_3$ (30 mL) was added, and the resulting mixture was extracted with EtOAc (2 x 120 mL). The

combined extracts were dried over $MgSO_4$ and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, 0%-100% EtOAc:EtOH (3:1)/heptane) provided *N,N*-bis(4-methoxybenzyl)-4-methyl-5-nitro-6-(prop-1-en-2-yl)pyridin-2-amine (1.00 g, 2.31 mmol, 62 % yield) as a yellow solid. m/z (ESI, +ve ion): 434.0 $(M+H)^+$.

[0629] Step 4: 6-Isopropyl-N^2,N^2-bis(4-methoxybenzyl)-4-methylpyridine-2,5-diamine (Intermediate I-33). A mixture of *N,N*-bis(4-methoxybenzyl)-4-methyl-5-nitro-6-(prop-1-en-2-yl)pyridin-2-amine (1.00 g, 2.31 mmol) and palladium, 10 wt. % on activated carbon (0.1 g) in ethanol (15 mL) and ethyl acetate (15 mL) was then stirred under $H_{2(g)}$ (40 psi) at rt for 1.5 hours, then under $H_{2(g)}$ (30 psi) at rt for 16 hours. The mixture was subsequently filtered through Celite (washing with EtOAc (2 x 100 mL)), and the combined filtrates were concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, 0%-10% ammonia in MeOH 2M/heptane) provided 6-isopropyl-N^2,N^2-bis(4-methoxybenzyl)-4-methylpyridine-2,5-diamine (**Intermediate I-33**, 340 mg, 0.838 mmol, 36 % yield) as a yellow oil. ^1H NMR ($CDCl_3$) δ: 7.26 (br d, J=7.3 Hz, 4H), 6.90 (br d, J=7.9 Hz, 4H), 6.13-6.26 (m, 1H), 4.56-4.83 (m, 4H), 3.87 (s, 8H), 3.02-3.19 (m, 1H), 2.17 (br d, J=10.4 Hz, 3H), 1.34 (br d, J=5.6 Hz, 6H). m/z (ESI, +ve ion): 406.1 $(M+H)^+$.

Intermediate I-36

4,6-Diisopropyl-2-methoxypyrimidin-5-amine



Intermediate I-36

[0630] Step 1: 4,6-Diisopropyl-2-(propylthio)pyrimidin-5-amine. A mixture of 4,6-dichloro-2-(propylthio)pyrimidin-5-amine (2.45 g, 10.29 mmol) and tetrahydrofuran (20.6 mL) was spared with argon gas for 5 min, then 2-propylzinc bromide (0.5 M in THF, 61.7 mL, 30.9 mmol) and Xantphos Pd G3 (Sigma-Aldrich, St. Louis, MO; 0.293 g, 0.309 mmol) were

sequentially added. The resulting mixture was heated to 60 °C and stirred for 2 h, then cooled to rt. Sat. aq. NH_4Cl (30 mL) was added, and the resulting mixture was stirred for 10 min, then diluted with EtOAc and brine (10 mL). The organic layer was separated, and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered, and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, 0-30% EtOAc/heptane) provided 4,6-diisopropyl-2-(propylthio)pyrimidin-5-amine (1.75 g, 6.91 mmol, 67 % yield) as a tan oil. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 4.80 (s, 2 H) 3.13 - 3.25 (m, 2 H) 2.91 - 3.05 (m, 2 H) 1.66 (sxt, *J*=7.26 Hz, 2 H) 1.12 (d, *J*=6.63 Hz, 12 H) 0.96 (t, *J*=7.36 Hz, 3 H). m/z (ESI, +ve ion): 254.1 (M+H)+.

[0631] Step 2: 4,6-Diisopropyl-2-(propylsulfonyl)pyrimidin-5-amine. A mixture of 4,6-diisopropyl-2-(propylthio)pyrimidin-5-amine (1.56 g, 6.16 mmol) and 3-chloroperoxybenzoic acid (3.45 g, 15.4 mmol) in THF (30.8 mL) was stirred at rt for 45 min, then cooled to 0 °C. Sat. aq. $NaHCO_3$ was added slowly (3 x 10 mL portions) over 20 min. The resulting mixture was diluted with 4:1 EtOAc/MeOH and brine (20 mL), and the organic layer was separated. The aqueous layer was extracted with EtOAc, and the combined organic extracts were then dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The residue was diluted with DMF (20 mL) and agitated by sonication to provide a homogeneous mixture. The mixture was slowly poured into water (100 mL), and the precipitated solid was collected by filtration, then washed with water and dried in a reduced-pressure oven at 45 °C overnight to afforded 4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-amine (1.65 g, 5.76 mmol, 94 % yield) as tan solid. ^{1}H NMR (400 MHz, DMSO-*d*$_6$) δ 6.08 (br s, 2 H) 3.48 - 3.68 (m, 3 H) 1.79 (br s, 2 H) 1.29 (br s, 12 H) 1.07 (br s, 4 H). m/z (ESI, +ve ion): 286.2 (M+H)+.

[0632] Step 3: 4,6-Diisopropyl-2-methoxypyrimidin-5-amine (Intermediate I-36). A mixture of 4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-amine (0.300 g, 1.05 mmol) and sodium methoxide (0.5 M in MeOH, 3.4 mL, 1.68 mmol) in *N, N*-dimethylformamide (3.5 mL) was stirred in a sealed vial at 90 °C for 16 h. Additional sodium methoxide (0.5 M in MeOH, 3.4 mL, 1.68 mmol) was subsequently added, and the resulting mixture was stirred at 90 °C for an additional 16 h. After cooling, the reaction mixture was partitioned between EtOAc and sat. aq. NH_4Cl. The organic layer was separated, and the aqueous layer was extracted with EtOAc. The combined organic extracts were then dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The residue was triturated with EtOAc and heptane, and the collected solids were washed with heptane and dried to provide 4,6-diisopropyl-2-methoxypyrimidin-5-amine

(**Intermediate I-36**, 0.095 g, 0.454 mmol, 43 % yield) as tan oil. m/z (ESI, +ve ion): 210.2 (M+H)+.

Intermediate I-37

4-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylaniline



Intermediate I-37

[0633] Step 1: 4-Iodo-2-isopropyl-6-methylaniline. To a suspension of 2-isopropyl-6-methylaniline (3.2 mL, 20 mmol, Advanced Chemblocks Inc., Burlingame, CA, USA), and sodium bicarbonate (3.4 g, 40 mmol) in DCM (20 mL) and water (20 mL) was added iodine (5.4 g, 21 mmol) in three portions. After 90 min, the 1 N sodium thiosulfate (30 mL) was added, and the resulting mixture was partitioned between DCM (30 mL) and water (10 mL). The organic layer was separated, and the aqueous layer was extracted with DCM (100 mL). The combined organic layers were then sequentially washed with brine (300 mL), dried over magnesium sulfate, filtered, and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, eluent: 0-30% EtOAc in heptanes) afforded 4-iodo-2-isopropyl-6-methylaniline as a red-purple oil. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 1.11 (d, *J* = 6.8 Hz, 6 H), 2.04 (s, 3 H), 2.97 (sept, *J* = 6.8 Hz, 1 H), 4.73 (s, 2 H), 7.08 – 7.10 (s, 1 H), 7.10 - 7.13 (m, 1 H); *m/z* (ESI, +ve ion) 275.9 $(M+H)^+$.

[0634] Step 2: 4-Amino-3-isopropyl-5-methylbenzonitrile. A nitrogen-sparged suspension of 4-iodo-2-isopropyl-6-methylaniline (2.2 g, 7.9 mmol), XantPhos Pd G3 (170 mg, 0.20 mmol), potassium acetate (580 mg, 5.9 mmol), and potassium hexacyanoferrate(II) trihydrate (5.0 g, 12 mmol) in 1,4-dioxane (20 mL) and water (20 mL) was vigorously stirred at 100 °C for 4 h. The reaction mixture was then partitioned between water (200 mL) and EtOAc (150 mL). The organic layer was separated, and the aqueous layer was extracted with EtOAc (2 × 70 mL). The combined organic extracts were then dried over magnesium sulfate,

filtered, and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, eluent: 0-40% EtOAc in heptanes) afforded 4-amino-3-isopropyl-5-methylbenzonitrile as an amber oil. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 1.09 - 1.18 (m, 6 H), 2.08 (s, 3 H), 3.02 (sept, *J* = 6.6 Hz, 1 H), 5.59 (br s, 2 H), 7.20 (br s, 1 H), 7.22 (br s, 1 H); *m/z* (ESI, +ve ion) 175.1 $(M+H)^+$.

[0635] Step 3: 4-Amino-3-isopropyl-5-methylbenzaldehyde. Diisobutylaluminum hydride (1 M in toluene, 17 mL, 17 mmol) was added, dropwise, to a room temperature solution of 4-amino-3-isopropyl-5-methylbenzonitrile (1.2 g, 6.9 mmol) in THF (34 mL). After 20 min, the reaction mixture was cooled to 0 °C and 1 M aq. Rochelle salt (35 mL) was added. The resulting mixture was stirred for 45 min, and the organic layer was then separated. The aqueous layer was extracted with EtOAc (1 × 70 mL), and the combined organic extracts were then sequentially washed with brine (300 mL), dried over magnesium sulfate, filtered through Celite®, and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, eluent: 0-15% EtOAc in heptanes) afforded 4-amino-3-isopropyl-5-methylbenzaldehyde as a thick yellow oil. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 1.17 (d, *J* = 6.6 Hz, 6 H), 2.14 (s, 3 H), 3.06 (spt, *J* = 6.6 Hz, 1 H), 5.75 (s, 2 H), 7.35 (s, 1 H), 7.44 (s, 1 H), 9.59 (s, 1 H); *m/z* (ESI, +ve ion) 178.1 $(M+H)^+$.

[0636] Step 4: (4-Amino-3-isopropyl-5-methylphenyl)methanol. Calcium borohydride bis(tetrahydrofuran) (1.8 g, 8.2 mmol) was added to a stirred solution of 4-amino-3-isopropyl-5-methylbenzaldehyde (970 mg, 5.5 mmol) in THF (8 mL), and the resulting mixture was stirred at rt for 2 h. Saturated aqueous NH4Cl (30 mL) was then slowly added, and the resulting mixture was extracted with (3 × 20 mL). The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated *in vacuo* to give crude (4-amino-3-isopropyl-5-methylphenyl)methanol, which was used without purification. *m/z* (ESI, +ve ion) 180.2 $(M+H)^+$.

[0637] Step 5: 4-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylaniline (Intermediate I-37). DIPEA (1.4 mL, 8.2 mmol) and *tert*-butyl(chloro)diphenylsilane (1.7 mL, 6.6 mmol) were sequentially added to a solution of (4-amino-3-isopropyl-5-methylphenyl)methanol (710 mg, 4.0 mmol) in DCM (10 mL), and the resulting mixture was stirred at room temperature for 4 h. The reaction mixture was then concentrated *in vacuo* and the residue chromatographically purified (silica gel, eluent: 0-40% EtOAc in heptanes) to provide 4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropyl-6-methylaniline (**Intermediate I-37**). *m/z* (ESI, +ve ion) 418.1 $(M+H)^+$.

Intermediate I-38

2-Bromo-4,6-diisopropylpyrimidin-5-amine



Intermediate I-38

[0638] A mixture of 4,6-diisopropylpyrimidin-5-amine (1 g, 5.58 mmol, **Intermediate U**), 1-bromopyrrolidine-2,5-dione (1.191 g, 6.69 mmol), and DMF (5 mL) was heated at 70 °C in a sealed pressure vial for 4 h, then cooled to rt and diluted with water. The resulting mixture was extracted with EtOAc, and the combined extracts were washed with brine, dried over Na_2SO_4, filtered, and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, 0-40% EtOAc/heptane) furnished 2-bromo-4,6-diisopropylpyrimidin-5-amine (**Intermediate I-38**) as brown oil. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 1.12 (d, *J*=6.63 Hz, 12 H) 3.12 - 3.27 (m, 2 H) 5.27 (s, 2 H). *m/z* (ESI, +ve ion): 258.0/260.0 $(M+H)^+$

Intermediate I-40

2-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-6-isopropylaniline



Intermediate I-40

[0639] **Step 1**: **(2-Amino-3-bromophenyl)methanol**. Borane tetrahydrofuran complex (1.0 M solution in THF, 23.4 mL, 23.4 mmol) was added, dropwise over 20 min, to a mixture of 2-amino-3-bromobenzoic acid (2.02 g, 9.34 mmol, Sigma-Aldrich, St. Louis, MO, USA) and THF (30 mL) at 0 °C. The resulting mixture was allowed to warm to rt and stir for 30 min before being stirred at 70 °C for 20 h. The reaction mixture was subsequently cooled to 0 °C, and MeOH (~5 mL) and ice-water (30 mL) were sequentially added. The aqueous layer was saturated with solid NaCl, and the resulting mixture was extracted with EtOAc (2 x 40 mL).

The combined extracts were dried over sodium sulfate, filtered and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, 10-50% EtOAc in heptane) furnished (2-amino-3-bromophenyl)methanol as a white solid (1.83 g, 9.04 mmol, 97% yield). m/z (ESI, +ve ion): 184.0 $(M+H)^+$.

[0640] Step 2: 2-Bromo-6-(((*tert*-butyldiphenylsilyl)oxy)methyl)aniline. *tert*-Butyldiphenylsilyl chloride (2.0 mL, 7.66 mmol, Sigma-Aldrich) was added to a mixture of (2-amino-3-bromophenyl)methanol (1.19 g, 5.89 mmol), DIPEA (3.6 mL, 20.6 mmol), DMAP (0.035 g, 0.28 mmol), and DCM (20 mL). The resulting mixture was stirred at rt for 24 h, then concentrated *in vacuo*. Chromatographic purification of the residues (silica gel, 0-50% EtOAc in heptane) provided 2-bromo-6-(((*tert*-butyldiphenylsilyl)oxy)methyl)aniline (2.42 g, 5.50 mmol, 93% yield) as a clear oil. m/z (ESI, +ve ion): 462.2 $(M+Na)^+$.

[0641] Step 3: 2-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-6-(prop-1-en-2-yl)aniline. A mixture of 2-bromo-6-(((*tert*-butyldiphenylsilyl)oxy)methyl)aniline (2.22 g, 5.04 mmol), (4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isopropene (1.23 mL, 6.55 mmol, Combi-Blocks, Inc.), bis-(di-*tert*-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II) (0.432 g, 0.610 mmol, Sigma-Aldrich), sodium carbonate (2 M aq solution; 6.30 mL, 12.60 mmol), and 1,4-dioxane (25 mL) was sparged with $Ar_{(g)}$ for 5 min, then stirred at 90 °C for 3 h. After cooling to rt, water (30 mL) was added, and the resulting mixture was extracted with EtOAc (2 x 30 mL). The combined extracts were dried over sodium sulfate, filtered, and concentrated *in vacuo*. Chromatographic purification of the residue (silica gel, 0-20% EtOAc in heptane) gave 2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-(prop-1-en-2-yl)aniline (1.80 g, 4.49 mmol, 89% yield) as a blue-green tinged oil.

[0642] Step 4: 2-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-6-isopropylaniline (Intermediate I-40). A mixture of 2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-(prop-1-en-2-yl)aniline (1.80 g, 4.48 mmol), palladium, 10 wt.%(dry basis) on activated carbon (wet, Degussa type E101 NE/W; 0.477 g, 0.224 mmol), ammonium formate (2.83 g, 44.8 mmol), and EtOH (20 mL) was stirred at 70 °C for 30 min. The mixture was then cooled to rt and filtered through a pad of diatomaceous earth. The filtrate was concentrated *in vacuo*, and the residue was partitioned between EtOAc (30 mL) and water (30 mL). The organic layer was separated and sequentially washed with water (10 mL), dried over sodium sulfate, filtered, and concentrated *in vacuo* to afford 2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylaniline (**Intermediate I-40**, 1.73 g, 4.29 mmol, 96% yield) as clear paste. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.71 (4 H, dd, *J*=7.9, 1.5 Hz), 7.37 - 7.46 (6 H, m), 7.13 (1 H, dd, *J*=7.7, 1.2 Hz), 6.74 - 6.79

(1 H, m), 6.66 - 6.72 (1 H, m), 4.75 (2 H, s), 4.42 (2 H, br d, *J*=1.5 Hz), 2.98 (1 H, dt, *J*=13.5, 6.8 Hz), 1.31 (6 H, d, *J*=6.6 Hz), 1.07 (9 H, s). m/z (ESI, +ve ion): 426.2 (M+Na).

Example 50, Step 3

3,6-Dichloro-5-(2-fluorophenyl)pyrazine-2-carboxamide





Example 50, Step 3

[0643] Step 1: Methyl 3-amino-6-chloro-5-(2-fluorophenyl)pyrazine-2-carboxylate. A mixture of methyl 3-amino-5,6-dichloropyrazine-2-carboxylate (10.0 g, 45 mmol, Ark Pharm, Inc., Arlington Heights, IL), (2-fluorophenyl)boronic acid (6.93 g, 49.5 mmol, Combi-Blocks, San Diego, CA, USA) and potassium carbonate (13.1 g, 95 mmol) in a 10:1 mixture of DME/water (220 mL) was degassed with nitrogen for 5 min and then tetrakis(triphenylphosphine)palladium(0) (1.04 g, 0.90 mmol) was added. The reaction mixture was stirred at 90 °C for 16 h, then was allowed to cool to room temperature and partitioned between EtOAc (200 mL) and 1 N HCl (200 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated to provide methyl 3-amino-6-chloro-5-(2-fluorophenyl)pyrazine-2-carboxylate. *m/z* (ESI, +ve): 282.1 $(M+H)^+$.

[0644] Step 2: Methyl 3,6-dichloro-5-(2-fluorophenyl)pyrazine-2-carboxylate. To a solution of methyl 3-amino-6-chloro-5-(2-fluorophenyl)pyrazine-2-carboxylate (22.0 g, 78 mmol) in *t*-BuOH (220 mL) was added isoamyl nitrite (15.8 mL,117 mmol) and copper chloride (12.6 g, 94 mmol). The resulting mixture was stirred at 60 °C for 16 h and then was partitioned between water (1 L) and EtOAc (2 L). The organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was purified by silica gel chromatography (eluent: 0-5% EtOAc/hexane) to provide methyl 3,6-dichloro-5-(2-fluorophenyl)pyrazine-2-carboxylate. *m/z* (ESI, +ve ion): 300.9 $(M+H)^+$.

[0645] **Step 3: 3,6-Dichloro-5-(2-fluorophenyl)pyrazine-2-carboxamide.** To a solution of methyl 3,6-dichloro-5-(2-fluorophenyl)pyrazine-2-carboxylate (13.5 g, 44.8 mmol) in THF (150 mL) was added ammonium hydroxide solution (30%, 150 mL, 44.8 mmol). The resulting mixture was heated at 50 °C for 4 h, then was allowed to cool to room temperature and partitioned between water and EtOAc (500 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was purified by silica gel chromatography (eluent: 0-30% EtOAc/hexanes) to provide 3,6-dichloro-5-(2-fluorophenyl)pyrazine-2-carboxamide (**Example 50, Step 3**). ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.32 (br s, 1 H), 8.16 (br s, 1 H), 7.64-7.77 (m, 2 H), 7.41-7.47 (m, 2 H). *m/z* (ESI, +ve): 286.0 $(M+H)^+$.

Intermediate P

2,5,6-Trichloronicotinamide



Intermediate P

[0646] 1,1'-Carbonyldiimidazole (40 g, 247 mmol) was added in portions to 2,5,6-trichloronicotinic acid (50.7 g, 224 mmol, Combi-Blocks, San Diego, CA, USA) in THF (400 mL), allowing gas evolution to cease between additions. The resulting mixture was stirred for 5 min and then was degassed with house vacuum and flushed with nitrogen (x2). The resulting mixture was heated to 50 °C for 60 min, then diluted with toluene (100 mL) and concentrated to half volume. The resulting mixture was cooled to 0 °C and ammonium hydroxide (60 mL, 437 mmol) was added slowly via syringe. The reaction was stirred for 10 min at room temperature, diluted with EtOAc (200 mL) and washed with water (3 x 100 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was suspended in 9:1 heptane/EtOAc (300 mL) and filtered. The filtered solids were collected and the remaining mother liquor was partially evaporated to half volume, cooled to 0 °C, and filtered. The two crops of filtered solids were combined to provide 2,5,6-trichloronicotinamide (**Intermediate P**, 46.2 g, 92% yield).

Intermediate Q

(2-Fluoro-6-hydroxyphenyl)potassium trifluoroborate



Intermediate Q

[0647] A solution of potassium fluoride (44.7 g, 770 mmol) in water (75 mL) was added to a suspension of (2-fluoro-6-hydroxyphenyl)boronic acid (30 g, 192 mmol, Combi-Blocks, San Diego, CA, USA) in acetonitrile (750 mL). The mixture was stirred for 2 min and then a solution of L-(+)-tartaric acid (72.2 g, 481 mmol) in THF (375 mL) was added over a 10 min period via addition funnel. The mixture was stirred vigorously with a mechanical stirrer for 1 h, and then the resulting suspension was filtered, and the filtered solids were washed with a small amount of THF. The solids were discarded and the filtrate was partially concentrated until solids started to precipitate out of solution. The mixture was then cooled to -20 °C and stirred for 16 h. The reaction was slowly warmed and 2-propanol (20 mL) was added. The resulting suspension was filtered and the filtered solids were washed with 2-propanol to provide 27.5 g of solid. The filtrate was again partially concentrated until a suspension formed and then was cooled to -20 °C and stirred for an additional 20 min. The resulting suspension was diluted with 2-propanol and filtered, and the filtered solids were washed with 2-propanol. The two batches of solids were combined to provide 2-fluoro-6-hydroxyphenyl)potassium trifluoroborate (**Intermediate Q**, 37.9 g, 90% yield). ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.07 (q, J = 14.7 Hz, 1 H) 6.93 (q, J = 7.5 Hz, 1 H) 6.30-6.38 (m, 2 H).

Intermediate R

2-Isopropyl-4-methylpyridin-3-amine



Intermediate R

[0648] To a slurry of 3-amino-2-bromo-4-picoline (360 mg, 1.9 mmol, Combi-Blocks, San Diego, CA, USA) in THF (4 mL) was added [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (79 mg, 0.10 mmol). The resulting slurry was deoxygenated with argon for 2 min and then 2-

propylzinc bromide (0.5 M solution in THF, 5.40 mL, 2.7 mmol) was added. The resulting solution was heated at 60 °C for 17 h, then the heating was stopped and the reaction was allowed to cool to room temperature. The reaction mixture was quenched with water (10 mL) and 1 N NaOH solution (20 mL) and then was extracted with EtOAc (2x). The combined organic layers were dried over anhydrous sodium sulfate and concentrated. The residue was purified by silica gel chromatography (eluent: 0-15% MeOH/DCM) to provide 2-isopropyl-4-methylpyridin-3-amine (**Intermediate R**, 284 mg, 98% yield). ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.66 (d, J = 4.6 Hz, 1 H), 6.78 (d, J = 4.8 Hz, 1 H), 4.72 (br s, 2 H), 3.14-3.25 (m, 1 H), 2.08 (s, 3 H), 1.14 (d, J = 6.8 Hz, 6 H). *m/z* (ESI, +ve ion): 151.1 $(M+H)^+$.

Intermediate U

4,6-Diisopropylpyrimidin-5-amine



Intermediate U

[0649] A solution of 4,6-dichloro-5-aminopyrimidine (3.00 g, 18.29 mmol, Combi-Blocks Inc., San Diego, CA, USA) in THF (18 mL) was deoxygenated by bubbling argon into the mixture for 5 min. 2-Isopropylzinc bromide (0.5 M solution in THF, 91.0 mL, 45.5 mmol) was added via syringe followed by XantPhos Pd G3 (434 mg, 0.46 mmol, Sigma-Aldrich, St. Louis, MO, USA). The resulting mixture was stirred at room temperature for 16 h and then was filtered through a pad of Celite. The filter cake was rinsed with EtOAc, and the filtrate was collected and concentrated to afford 4,6-diisopropylpyrimidin-5-amine (**Intermediate U**, 3.45 g). This material was used without further purification in the following step. *m/z* (ESI, +ve ion): 180.2 $(M+H)^+$.

Intermediates 61A and 61B

6,7-Dichloro-1-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione and (*S*)-*tert*-Butyl 4-(6,7-dichloro-1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate



[0650] Step 1: 2,5,6-Trichloro-*N*-((2-isopropylphenyl)carbamoyl)nicotinamide. A suspension mixture of 2,5,6-trichloronicotinamide (**Intermediate P**, 6.7 g, 29.7 mmol) in 1,2-dichloroethane (100 mL) was treated with oxalyl chloride (3.0 mL, 35.7 mmol) at rt. The resulting reaction mixture was stirred at 80 °C for 30 min then the white suspension was evaporated to give a slurry. The slurry was treated with acetonitrile (100 mL) and then with 2-isopropylaniline (4.6 mL, 32.7 mmol, Sigma-Aldrich Corporation, St. Louis, MO, USA) at rt. The mixture was stirred for 15 min and the white solid was collected by filtration, washed with acetonitrile and dried to give pure 2,5,6-trichloro-*N*-((2-isopropylphenyl)carbamoyl)nicotinamide (8.55 g, 22.1 mmol, 74.4 % yield) as a white solid. *m/z* (ESI, +ve ion): 386.0 $(M+H)^+$.

[0651] Step 2: 6,7-Dichloro-1-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (Intermediate 61A). A mixture of 2,5,6-trichloro-*N*-((2-isopropylphenyl)carbamoyl)nicotinamide (8.55 g, 22.1 mmol) in THF (74 mL) at 0 °C was treated with KHDMS (1M solution in THF, 44.3 mL, 44.3 mmol). The mixture was stirred at 0 °C for 10 min and at rt for 30 min. The reaction mixture was quenched with satd. ammonium chloride (100 mL) and extracted with EtOAc (200 mL). The organic layer was separated, washed with brine (150 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The resulting crude product was sonicated in MeOH (20 mL), filtered, and washed with MeOH and dried to give pure 6,7-dichloro-1-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 61A**, 7.17 g, 20.5 mmol, 92 % yield) as a tan solid. ^{1}H NMR (400 MHz,

DMSO-d$_6$) δ 12.10 (s, 1 H), 8.52 (s, 1 H), 7.38 - 7.60 (m, 2 H), 7.12 - 7.38 (m, 2 H), 2.74 (dt, J=13.5, 6.8 Hz, 1 H), 1.08 (d, J=6.8 Hz, 3 H), 1.03 (d, J=6.8 Hz, 3 H).). *m/z* (ESI, +ve ion): 350.0 $(M+H)^+$.

[0652] Step 3: (*S*)-*tert*-Butyl 4-(6,7-dichloro-1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (Intermediate 61B). To a mixture of 6,7-dichloro-1-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 61A**, 1.03 g, 2.94 mmol) and DIPEA (1.5 mL, 8.82 mmol) in acetonitrile (19.6 mL) was added phosphorus oxychloride (1.4 mL, 8.82 mmol) at rt and heated at 80 °C for 30 min. The mixture was concentrated *in vacuo* to give the crude 4,6,7-trichloro-1-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a brown solid. *m/z* (ESI, +ve ion): 368.0 $(M+H)^+$. The crude material was used in next step without purification.

[0653] To a mixture of the above 4,6,7-trichloro-1-(2-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (1.08 g, 2.94 mmol) and DIPEA(1.5 mL, 8.82 mmol) in DMF (14.7 mL) was added (*S*)-4-*N*-Boc-2-methyl piperazine (0.88 g, 4.41 mmol) and stirred at rt for 10 min. Ice water (10 mL) was added and stirred for 15 min. The resulting precipitate was collected by filtration, washed with water, and dried to give (*S*)-*tert*-butyl 4-(6,7-dichloro-1-(2-isopropylphenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (**Intermediate 61B**, 1.52 g, 2.85 mmol, 97 % yield) as a yellow solid. *m/z* (ESI, +ve ion): 532.0 $(M+H)^+$.

Intermediates S and 72A

2,6-Dichloro-5-fluoronicotinamide and 7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0654] Step 1: 2,6-Dichloro-5-fluoronicotinamide (Intermediate S). To a mixture of 2,6-dichloro-5-fluoro-nicotinic acid (4.0 g, 19.1 mmol, AstaTech Inc., Bristol, PA) in dichloromethane (48 mL) was added oxalyl chloride (2M solution in DCM, 11.9 mL, 23.8 mmol), followed by a catalytic amount of DMF (0.05 mL). The reaction was stirred at room temperature overnight and then was concentrated. The residue was dissolved in 1,4-dioxane (48 mL) and cooled to 0 °C. Ammonium hydroxide solution (28.0-30% NH3 basis, 3.6 mL, 28.6 mmol) was added slowly via syringe. The resulting mixture was stirred at 0 °C for 30 min and then was concentrated. The residue was diluted with a 1:1 mixture of EtOAc/heptane and agitated for 5 min, then was filtered. The filtered solids were discarded, and the remaining mother liquor was partially concentrated to half volume and filtered. The filtered solids were washed with heptane and dried in a reduced-pressure oven (45 °C) overnight to provide 2,6-dichloro-5-fluoronicotinamide (**Intermediate S**, 2.00 g, 50% yield). ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.23 (d, J = 7.9 Hz, 1 H) 8.09 (br s, 1 H) 7.93 (br s, 1 H). m/z (ESI, +ve ion): 210.9 $(M+H)^+$.

[0655] Step 2: 2,6-Dichloro-5-fluoro-*N*-((2-isopropyl-4-methylpyridin-3-yl)carbamoyl)nicotinamide. To an ice-cooled slurry of 2,6-dichloro-5-fluoronicotinamide (**Intermediate S**, 5.0 g, 23.9 mmol) in THF (20 mL) was added oxalyl chloride (2 M solution in DCM, 14.4 mL, 28.8 mmol) slowly via syringe. The resulting mixture was heated at 75 °C for 1 h, then heating was stopped, and the reaction was concentrated to half volume. After

cooling to 0 °C, THF (20 mL) was added, followed by a solution of 2-isopropyl-4-methylpyridin-3-amine (**Intermediate R**, 3.59 g, 23.92 mmol) in THF (10 mL), dropwise via cannula. The resulting mixture was stirred at 0 °C for 1 h and then was quenched with a 1:1 mixture of brine and saturated aqueous ammonium chloride. The mixture was extracted with EtOAc (3x) and the combined organic layers were dried over anhydrous sodium sulfate and concentrated to provide 2,6-dichloro-5-fluoro-*N*-((2-isopropyl-4-methylpyridin-3-yl)carbamoyl)nicotinamide (9.2 g, 100% yield). This material was used without further purification in the following step. *m/z* (ESI, +ve ion): 385.1(M+H)$^+$.

[0656] Step 3: 7-Chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To an ice-cooled solution of 2,6-dichloro-5-fluoro-*N*-((2-isopropyl-4-methylpyridin-3-yl)carbamoyl)nicotinamide (9.2 g, 24.0 mmol) in THF (40 mL) was added KHMDS (1 M solution in THF, 50.2 mL, 50.2 mmol) slowly via syringe. The ice bath was removed and the resulting mixture was stirred for 40 min at room temperature. The reaction was quenched with saturated aqueous ammonium chloride and extracted with EtOAc (3x). The combined organic layers were dried over anhydrous sodium sulfate and concentrated. The residue was purified by silica gel chromatography (eluent: 0-50% 3:1 EtOAc-EtOH/heptane) to provide 7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (5.0 g, 60% yield). ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.27 (br s, 1H), 8.48-8.55 (m, 2 H), 7.29 (d, J = 4.8 Hz, 1 H), 2.87 (quin, J = 6.6 Hz, 1 H), 1.99-2.06 (m, 3 H), 1.09 (d, J = 6.6 Hz, 3 H), 1.01 (d, J = 6.6 Hz, 3 H). ^{19}F NMR (376 MHz, DMSO-d_6) δ: -126.90 (s, 1 F). *m/z* (ESI, +ve ion): 349.1 (M+H)$^+$.

[0657] Step 4: 7-Chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (Intermediate 72A). A mixture of atropisomers 7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Example 41**, Step 3, 648 g) was purified by SFC (AD, 150 x 50 mm, 5 μm, 50% MeOH/CO_2, 180 g/min, 102 bar) to obtain two peaks: Peak 1 (P isomer, 230.6 g, >99% ee) and Peak 2 (M isomer, 227.8 g, 97.1% ee, **Intermediate 72A**).

Intermediates 73A and 73B

(*P*)- and (*M*)-6,7-Dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0658] Step 1: 2-Isopropyl-4-methylpyridin-3-amine (Intermediate R). To a slurry of 3-amino-2-bromo-4-picoline (360 mg, 1.9 mmol, Combi-Blocks, San Diego, CA, USA) in THF (4 mL) was added [1,1′-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane (79 mg, 0.10 mmol). The resulting slurry was deoxygenated with argon for 2 min and then 2-propylzinc bromide (0.5 M solution in THF, 5.40 mL, 2.7 mmol) was added. The resulting solution was heated at 60 °C for 17 h, then the heating was stopped and the reaction was allowed to cool to room temperature. The reaction mixture was quenched with water (10 mL) and 1 N NaOH solution (20 mL) and then was extracted with EtOAc (2x). The combined organic layers were dried over anhydrous sodium sulfate and concentrated. The residue was purified by silica gel chromatography (eluent: 0-15% MeOH/DCM) to provide 2-isopropyl-4-methylpyridin-3-amine (284 mg, 98% yield). ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.66 (d, J = 4.6 Hz, 1 H), 6.78 (d, J = 4.8 Hz, 1 H), 4.72 (br s, 2 H), 3.14-3.25 (m, 1 H), 2.08 (s, 3 H), 1.14 (d, J = 6.8 Hz, 6 H). *m/z* (ESI, +ve ion): 151.1 $(M+H)^+$.

[0659] Step 2: 2,5,6-Trichloro-*N*-((2-isopropyl-4-methylpyridin-3-yl)carbamoyl)nicotinamide. To a -78 °C slurry of 2,5,6-trichloronicotinamide (**Intermediate P**, 3.10 g, 13.8 mmol) in THF (46 mL) was added oxalyl chloride (2 M solution in DCM, 7.4 mL, 14.7 mmol) slowly via syringe. The resulting slurry was heated at 60 °C for 3.5 h, then heating was stopped and the reaction was cooled to -78 °C. Triethylamine (6.0 mL, 42.6 mmol) was added followed by a solution of 2-isopropyl-4-methylpyridin-3-amine (**Intermediate R**,

2.12 g, 14.1 mmol) via cannula. The resulting slurry was warmed to room temperature and stirred for 1 h, then was partitioned between water (120 mL) and EtOAc (175 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was suspended in 9:1 heptane/EtOAc and filtered. The filtered solids were collected to provide 2,5,6-trichloro-*N*-((2-isopropyl-4-methylpyridin-3-yl)carbamoyl)nicotinamide (4.71 g, 85% yield). ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.31 (s, 1 H), 9.54 (s, 1 H), 8.66 (s, 1 H), 8.34 (d, J = 4.8 Hz, 1 H), 7.16 (d, J = 5.0 Hz, 1 H), 3.24-3.33 (m, 1 H), 2.22 (s, 3 H), 1.17 (d, J = 6.6 Hz, 6 H). *m/z* (ESI, +ve ion): 400.9 $(M+H)^+$.

[0660] Step 3: 6,7-Dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A mixture of atropisomers 6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (55.1 g) was purified by SFC (AD, 250 x 50 mm, 5 μm, 50% MeOH/CO2, 180 g/min, 102 bar) to obtain two peaks: Peak 1 (**Intermediate 73A**, (*P*)-isomer, 22.1 g, >99% ee) and Peak 2 ((*M*)-isomer, 23.2 g, >99% ee). Peak 2 was the desired material (**Intermediate 73B**).

Intermediate 76A

(*M*)-6-Chloro-7-(2-fluorophenyl)-4-hydroxy-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Intermediate 73B Intermediate 76A

[0661] A mixture of (*M*)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 73B**, 4.38 g, 12 mmol), 2-fluorophenylboronic acid (2.35 g, 16.8 mmol), (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.351 g, 0.48 mmol), potassium acetate (5.1 g, 52.0 mmol) in 1,4-dioxane (30 mL) and water (1 mL) was stirred and heated at 85 °C for 15 h. The mixture was cooled to rt and diluted with EtOAc and water. The organic layer was washed with brine, dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The residue was purified by silica gel chromatography (eluent: 0-40% EtOAc/heptane) to provide (*M*)-6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 76A**) as white solids. *m/z* (ESI, +ve ion): 424.9 $(M+H)^+$.

Intermediate 85A

Methyl 3,5-dichloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridine-8-carboxylate



Intermediate 85A

[0662] Step 1: Methyl 2,3-dichloro-7-(methoxymethyl)-5-oxo-5,6-dihydro-1,6-naphthyridine-8-carboxylate. A mixture of 2,5,6-trichloronicotinamide (**Intermediate P**, 5.0 g, 22 mmol), 4-methoxy-3-oxo-butanoic acid methyl ester (4.86 mL, 33.3 mmol), copper(I) iodide (0.42 g, 2.22 mmol) and cesium carbonate (14.45 g, 44.4 mmol) was purged with N_2 followed by the addition of 1,4-dioxane (110 mL) and the reaction mixture was heated at 80 °C under nitrogen for 16 h. The mixture was quenched with 9:1 sat. NH_4Cl/NH_4OH and extracted with EtOAc. The combined organics were concentrated to give methyl 2,3-dichloro-7-(methoxymethyl)-5-oxo-5,6-dihydro-1,6-naphthyridine-8-carboxylate (4.0 g, 12.6 mmol, 56.9 % yield). *m/z* (ESI, +ve ion): 317.0 $(M+H)^+$. The crude material was used as is in the subsequent step.13

[0663] Step 2: Methyl 3-chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-5-oxo-5,6-dihydro-1,6-naphthyridine-8-carboxylate. A mixture of methyl 2,3-dichloro-7-(methoxymethyl)-5-oxo-5,6-dihydro-1,6-naphthyridine-8-carboxylate (4.0 g, 12.6 mmol), (2-fluorophenyl)boronic acid (2.29 g, 16.4 mmol, Combi-Blocks Inc.), palladium tetrakis (1.46 g, 1.26 mmol) and sodium carbonate (4.01 g, 37.8 mmol) in 1,4-dioxane/water (30/7.5 mL) was heated at 85 °C for 45 min. The reaction was quenched with saturated $NaHCO_3$ and extracted with EtOAc. The crude product was purified by silica gel chromatography (eluent: 0-50% EtOAc:EtOH (3:1)/heptane) to provide methyl 3-chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-5-oxo-5,6-dihydro-1,6-naphthyridine-8-carboxylate (4.0 g, 10.6 mmol, 84 % yield) as a yellow solid. *m/z* (ESI, +ve ion): 377.0 $(M+H)^+$.

[0664] Step 3: Methyl 3,5-dichloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridine-8-carboxylate (Intermediate 85A). A solution of methyl 3-chloro-2-(2-fluorophenyl)-7-(methoxymethyl)-5-oxo-5,6-dihydro-1,6-naphthyridine-8-carboxylate (0.30 g, 0.80 mmol) and phosphoryl trichloride (5.0 mL, 54 mmol) was heated at 90 °C for 1 h. The reaction was concentrated in vacuo and the residue was diluted with EtOAc, washed with sat. $NaHCO_3$ and brine. The combined organics were concentrated to afford methyl 3,5-dichloro-2-(2-fluorophenyl)-7-(methoxymethyl)-1,6-naphthyridine-8-carboxylate (**Intermediate 85A**, 0.3 g, 0.759 mmol, 95 % yield) as a brown solid. *m/z* (ESI, +ve ion): 395.0 $(M+H)^+$.

Intermediate 92B

4-((2*S*,5*R*,*M*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0665] Step 1: (*2R,5S,M*)-*tert*-Butyl 4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A 250-mL round-bottomed flask was charged with (*M*)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 73B**, 6.65 g, 18.2 mmol) and DIPEA (4.8 mL, 27.3 mmol) in acetonitrile (91 mL) followed by phosphorous oxychloride (2.6 mL, 27.3 mmol). The resulting mixture was stirred at 80 °C for 30 min and then concentrated *in vacuo* to give (*M*)-4,6,7-trichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as a brown solid. The crude brown solid was used in next step without purification. *m/z* (ESI, +ve): 383.0 $(M+H)^+$.

[0666] To a mixture of crude (*M*)-4,6,7-trichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (4.8 mL, 27.3 mmol) in DMF (50 mL) was added (*2R,5S*)-1-Boc-2,5-dimethylpiperazine (4.29 g, 20.03 mmol, AstaTech Inc., Bristol, PA) and the mixture was stirred at rt for 15 min. The mixture was added to ice water (80 mL) and stirred for 15 min. The resulting precipitate was collected by filtration, washed with water, and dried to give *tert*-butyl (*2R,5S,M*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-

oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (4.70 g, 8.37 mmol, 46.0 % yield) as a yellow solid. The filtrate was extracted with EtOAc (2 × 100 mL). The combined organic extracts were dried over $MgSO_4$. The solution was filtered and concentrated *in vacuo* to give additional title compound (5.51 g, 9.81 mmol, 53.9 % yield) as a yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.59 (br d, J=4.8 Hz, 1 H), 8.54 (s, 1 H), 7.40 (br d, J=2.7 Hz, 1 H), 4.87 (br s, 1 H), 4.23 - 4.44 (m, 1 H), 4.01 - 4.09 (m, 1 H), 3.95 (br s, 1 H), 3.73 (br dd, J=13.7, 2.5 Hz, 1 H), 3.46 - 3.65 (m, 1 H), 2.67 - 2.76 (m, 1 H), 2.04 (s, 3 H), 1.45 - 1.57 (m, 9 H), 1.36 (d, J=6.6 Hz, 3 H), 1.08 - 1.18 (m, 9 H). *m/z* (ESI, +ve): 561.2 $(M+H)^+$.

[0667] Step 2: 4-((2*S*,5*R*,*M*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (Intermediate 92B). To a solution of *tert*-butyl (2*R*,5*S*,*M*)-4-(6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (5.51 g, 9.81 mmol) in DCM (20 mL) was added trifluoroacetic acid (10 mL, 134 mmol) at rt and the mixture was stirred for 1 h. After the reaction was complete, the mixture was concentrated *in vacuo* to afford (*M*)-6,7-dichloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, m/z (ESI, +ve): 461.2 $(M+H)^+$.

[0668] To the above (*M*)-6,7-dichloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and *N*,*N*'-diisopropylethylamine (8.6 mL, 49.1 mmol) in DCM (20 mL) was added acryloyl chloride (0.8 mL, 9.81 mmol) at 0 °C and the mixture was stirred for 1 h. The reaction mixture was diluted with DCM (50 mL) and washed with sat. ammonium chloride solution (50 mL). To the aqueous was added sat'd sodium chloride (25 mL) and the mixture was extracted with DCM (50 mL × 2). The organic extracts were combined and dried over $MgSO_4$. The solution was filtered and concentrated *in vacuo* to give a brown oil. The crude brown oil was adsorbed onto a plug of silica gel and purified by silica gel chromatography (eluent: 0-10% of MeOH/DCM) to give (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 92B**, 4.66 g, 9.04 mmol, 92 % yield) as a light yellow foam. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 - 8.54 (m, 2 H), 7.30 (d, J=5.0 Hz, 1 H), 6.82 (ddd, J=16.5, 14.0, 10.5 Hz, 1 H), 6.18 (dd, J=16.7, 2.2 Hz, 1 H), 5.74 (dt, J=10.4, 2.7 Hz, 1 H), 4.78 - 4.91 (m, 1 H), 4.39 - 4.75 (m, 1 H), 3.97 - 4.16 (m, 1 H), 3.94 (br s, 1 H), 3.83 (br d, J=3.9 Hz, 1 H), 3.49 (br dd, J=13.9, 3.7 Hz, 1 H), 2.59 - 2.70 (m, 1

H), 1.97 (s, 3 H), 1.25 - 1.32 (m, 3 H), 1.09 - 1.20 (m, 3 H), 1.05 (dd, J=11.4, 6.6 Hz, 6 H). *m/z* (ESI, +ve): 515.2 $(M+H)^+$.

Intermediate 94B

(*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Intermediate 72A | Intermediate 94A | Intermediate 94B

[0669] Step 1: (*M*)-*tert*-Butyl (2*R*,5*S*)-4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of (*M*)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (**Intermediate 72A**; 3.35 g, 9.61 mmol), phosphorous oxychloride (1.07 mL, 11.53 mmol), and DIPEA (5.02 mL, 28.8 mmol) in acetonitrile (24 mL) was stirred at 80 °C for 30 min. The reaction mixture was cooled to rt and added (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (2.26 g, 10.6 mmol). The reaction mixture was stirred at rt for 15 min. The reaction mixture was diluted with EtOAc (200 mL), washed with satd. $NaHCO_3$ (3 × 75 mL), dried over Na_2SO_4, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-100% EtOAc/heptane) to provide (*M*)-*tert*-butyl (2*R*,5*S*)-4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 94A**; 2.2 g, 4.04 mmol, 42.0 % yield) as yellow-orange solid. ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 8.57 (d, *J*=5.0 Hz, 1 H), 7.79 (br d, *J*=7.5 Hz, 1 H), 7.13 - 7.16 (m, 1 H), 4.80 - 5.04 (m, 1 H), 4.32 - 4.64 (m, 1 H), 3.73 - 4.08 (m, 3 H), 3.43 - 3.66 (m, 1 H), 2.58 (dt, *J*=13.4, 6.6 Hz, 1 H), 2.04 (s, 3 H), 1.59 (s, 9 H), 1.39 - 1.48 (m, 3 H), 1.11 - 1.25 (m, 9 H). ^{19}F NMR (377 MHz, *CHLOROFORM-d*) δ -126.30 (br s, 1 F) -126.34 (br s, 1 F). *m/z* (ESI, +ve ion): 544.8 $(M+H)^+$.

[0670] Step 2: (*M*)-4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A mixture of (*M*)-*tert*-butyl (2*R*,5*S*)-4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate

(**Intermediate 94A**; 8.16 g, 14.97 mmol) in DCM (30 mL) and TFA (30 mL) was stirred at rt for 30 min. The reaction mixture was concentrated *in vacuo* to provide (*M*)-7-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. The crude product was used in next step without purification. *m/z* (ESI, +ve ion) 445.1 $(M+H)^+$.

[0671] A mixture of (*M*) 7-chloro-4-((2*S*,5*R*)-2,5-dimethylpiperazin-1-yl)-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one and DIPEA (13.1 mL, 74.9 mmol) in DCM (20 mL) was cooled to 0 °C. To the cooled mixture was added acryloyl chloride (1.22 mL, 14.97 mmol) and stirred for 1 h. The reaction mixture was quenched with satd. NH_4Cl (50 mL), extracted with DCM (2 × 50 mL)), dried over MgSO4, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-100% EtOAc-MeOH (9:1)/heptane) to provide (*M*)-4-((2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 94B**; 5.62 g, 11.3 mmol, 75 % yield) as light-yellow foam. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.53 (d, *J*=5.0 Hz, 1 H), 8.44 (d, *J*=8.5 Hz, 1 H), 7.35 (br d, *J*=3.7 Hz, 1 H), 6.83 (td, *J*=16.9, 10.6 Hz, 1 H), 6.18 (dd, *J*=16.6, 2.3 Hz, 1 H), 5.69 - 5.80 (m, 1 H), 4.65 - 4.92 (m, 2 H), 3.79 - 4.15 (m, 3 H), 3.07 - 3.69 (m, 1 H), 2.60 - 2.72 (m, 1 H), 1.98 (s, 3 H), 1.20 - 1.33 (m, 6 H), 1.06 (dd, *J*=13.1, 6.8 Hz, 6 H). *m/z* (ESI, +ve ion): 499.2 $(M+H)^+$.

Intermediate 99 A and B

2,5-Dichloro-6-(2-fluorophenyl)nicotinamide



[0672] **Step 1: 2,5-Dichloro-6-(2-fluorophenyl)nicotinic acid**. A mixture of 2,5,6-trichloronicotinic acid (1.03 g, 4.54 mmol, Combi-Blocks, San Diego, CA), palladium tetrakis (0.131 g, 0.114 mmol), (2-fluorophenyl)boronic acid (0.699 g, 5.0 mmol, TCI America, Portland, OR), and sodium carbonate (2M in water, 6.82 mL, 13.6 mmol) in 1,4-dioxane (11 mL) was sparged with nitrogen and heated to 80 °C for 1 h followed by 90 °C for 5 h. The reaction mixture was diluted with EtOAc (150 mL), washed with 1 N aqueous citric acid (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in*

vacuo to give 2,5-dichloro-6-(2-fluorophenyl)nicotinic acid (**Intermediate 99A**, 1.27 g, 4.43 mmol, 97% yield) as an amber oil. *m/z* (ESI, +ve ion): 285.8 $(M+H)^+$.

[0673] **Step 2: 2,5-Dichloro-6-(2-fluorophenyl)nicotinamide**. A solution of 2,5-dichloro-6-(2-fluorophenyl)nicotinic acid (**Intermediate 99A**, 1.27 g, 4.43 mmol) in sulfurous dichloride (13 mL, 177 mmol) was stirred at 70 °C for 30 min. The reaction mixture was concentrated *in vacuo* to give a dark brown oil. The oil was dissolved in 1,4-dioxane (8.9 mL) and treated with ammonium hydroxide (30% aq., 3.5 mL, 89 mmol) and the mixture was stirred at rt for 5 min. The reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (3 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-80% EtOAc/heptane) to provide crude product as an off-white solid. The solid was stirred in EtOH (8 mL) at rt for 15 min and filtered to give 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (**Intermediate 99B**, 0.449 g, 1.58 mmol, 36% yield) as a white solid. *m/z* (ESI, +ve ion): 284.8 $(M+H)^+$.

Intermediate 106A

7-Bromo-6-chloro-1-(2-isopropyl-6-methylphenyl)quinazoline-2,4(1*H*,3*H*)-dione



[0674] **Step 1: 4-Bromo-5-chloro-2-fluorobenzamide.** A mixture of 4-bromo-5-chloro-2-fluorobenzoic acid (**Oxchem Corp.,** Wood Dale, IL, USA; 23.3 g, 92 mmol) in thionyl chloride (67 mL, 0.92 mol) was stirred at 70 °C for 1 h. The reaction mixture was then concentrated *in vacuo*, and the residue was taken up in 1,4-dioxane (200 mL), treated with ammonium hydroxide (30% aqueous, 82 mL, 0.64 mol), and stirred at rt for 15 min. The reaction mixture

was concentrated *in vacuo* to give 4-bromo-5-chloro-2-fluorobenzamide: *m/z* (ESI, +ve ion): 251.8 $(M+H)^+$.

[0675] Step 2: 4-Bromo-5-chloro-2-fluoro-*N*-((2-isopropyl-6-methylphenyl)carbamoyl) benzamide. Oxalyl chloride (2 M in DCM, 8.9 mL, 17.8 mmol) was added over a 2 min period to a suspension of 4-bromo-5-chloro-2-fluorobenzamide (3.0 g, 11.9 mmol) in 1,2-dichloroethane (30 mL) under nitrogen atmosphere. The suspension was stirred at rt for 5 min and then heated to 70 °C for 75 min. The reaction mixture was concentrated *in vacuo* to afford a white solid. This solid was suspended in acetonitrile (15 mL), and a solution of 2-(1-methylethyl)-6-methylaniline (2.13 g, 14.3 mmol, Advanced Chemblocks Inc., Burlingame, CA) in acetonitrile (5 mL) was added dropwise at rt. The precipitate was filtered off, washed with heptane, and dried under reduced pressure to give 4-bromo-5-chloro-2-fluoro-*N*-((2-isopropyl-6-methylphenyl)carbamoyl)benzamide. *m/z* (ESI, +ve ion): 427.0 and 428.9 $(M+H)^+$.

[0676] Step 3: 7-Bromo-6-chloro-1-(2-isopropyl-6-methylphenyl)quinazoline-2,4(1*H*,3*H*)-dione (Intermediate 106A). KHMDS (1 M in THF, 20.6 mL, 20.6 mmol) was added portion-wise over a period of 10 min to a suspension of 4-bromo-5-chloro-2-fluoro-*N*-((2-isopropyl-6-methylphenyl)carbamoyl)benzamide (4.4 g, 10.3 mmol) in THF (30 mL) cooled to 0 °C under nitrogen atmosphere. The reaction mixture was allowed to warm to rt and stirred for 18 h. The reaction mixture was then partitioned between EtOAc (100 mL) and satd. $NaHCO_3$ (50 mL). The organic layer was washed with brine (20 mL), dried over $MgSO_4$, and concentrated *in vacuo* to give 7-bromo-6-chloro-1-(2-isopropyl-6-methylphenyl)quinazoline-2,4(1*H*,3*H*)-dione (Intermediate 106A). *m/z* (ESI, +ve ion): 407.0 and 409.0 $(M+H)^+$.

Intermediate 111A

6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridine-2,4(1*H*,3*H*)-dione



[0677] Step 1: Methyl 3-amino-6-chloro-5-(2-fluorophenyl)pyrazine-2-carboxylate. A mixture of methyl 3-amino-5,6-dichloropyrazine-2-carboxylate (10.0 g, 45 mmol, Ark Pharm, Inc., Arlington Heights, IL), (2-fluorophenyl)boronic acid (6.93 g, 49.5 mmol, Combi-Blocks, San Diego, CA, USA) and potassium carbonate (13.1 g, 95 mmol) in a 10:1 mixture of DME/water (220 mL) was degassed with nitrogen for 5 min and then tetrakis(triphenylphosphine)palladium(0) (1.04 g, 0.90 mmol) was added. The reaction mixture was stirred at 90 °C for 16 h, then was allowed to cool to room temperature and partitioned between EtOAc (200 mL) and 1 N HCl (200 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated to provide methyl 3-amino-6-chloro-5-(2-fluorophenyl)pyrazine-2-carboxylate. *m/z* (ESI, +ve): 282.1 $(M+H)^+$.

[0678] Step 2: Methyl 3,6-dichloro-5-(2-fluorophenyl)pyrazine-2-carboxylate. To a solution of methyl 3-amino-6-chloro-5-(2-fluorophenyl)pyrazine-2-carboxylate (22.0 g, 78 mmol) in *t*-BuOH (220 mL) was added isoamyl nitrite (15.8 mL,117 mmol) and copper chloride (12.6 g, 94 mmol). The resulting mixture was stirred at 60 °C for 16 h and then was partitioned between water (1 L) and EtOAc (2 L). The organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was purified by silica gel chromatography (eluent: 0-5% EtOAc/hexane) to provide methyl 3,6-dichloro-5-(2-fluorophenyl)pyrazine-2-carboxylate. *m/z* (ESI, +ve ion): 300.9 $(M+H)^+$.

[0679] Step 3: 3,6-Dichloro-5-(2-fluorophenyl)pyrazine-2-carboxamide. To a solution of methyl 3,6-dichloro-5-(2-fluorophenyl)pyrazine-2-carboxylate (13.5 g, 44.8 mmol) in THF (150 mL) was added ammonium hydroxide solution (30%, 150 mL, 44.8 mmol). The resulting mixture was heated at 50 °C for 4 h, then was allowed to cool to room temperature and partitioned between water and EtOAc (500 mL). The organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was purified by silica gel chromatography (eluent: 0-30% EtOAc/hexanes) to provide 3,6-dichloro-5-(2-fluorophenyl)pyrazine-2-carboxamide. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.32 (br s, 1 H), 8.16 (br s, 1 H), 7.64-7.77 (m, 2 H), 7.41-7.47 (m, 2 H). *m/z* (ESI, +ve): 286.0 $(M+H)^+$.

[0680] Step 4: 3,6-Dichloro-5-(2-fluorophenyl)-*N*-((2-isopropyl-4-methylpyridin-3-yl)carbamoyl)pyrazine-2-carboxamide. A mixture of 3,6-dichloro-5-(2-fluorophenyl)pyrazine-2-carboxamide (**Example 50, Step 3**, 1.45 g, 5.06 mmol) and oxalyl chloride (2 M in DCM, 2.8 mL, 5.6 mmol) in THF (25 mL) was heated to 80 °C for 2 h. The reaction mixture was then cooled to 0 °C and 2-isopropyl-4-methylpyridin-3-amine (**Intermediate R**, 0.8 g, 5.34 mmol) was added. The cold bath was removed and stirring was continued at rt for 45 min. The reaction mixture was concentrated *in vacuo*, the residue was taken up in EtOAc, and sonicated. The precipitate was filtered off and dried to give 3,6-dichloro-5-(2-fluorophenyl)-*N*-((2-isopropyl-4-methylpyridin-3-yl)carbamoyl)pyrazine-2-carboxamide as a light yellow solid. *m/z* (ESI, +ve ion): 461.8 and 463.8 $(M+H)^+$.

[0681] Step 5: 6-Chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridine-2,4(1*H*,3*H*)-dione (Intermediate 111A). KHMDS (1 M in THF, 5.3 mL, 5.3 mmol) was added to a solution of 3,6-dichloro-5-(2-fluorophenyl)-*N*-((2-isopropyl-4-methylpyridin-3-yl)carbamoyl)pyrazine-2-carboxamide (1.22 g, 2.64 mmol) in THF (15 mL) at 0 °C. The reaction mixture was allowed to warm to rt over a period of 1 h. The reaction mixture was then diluted with EtOAc and washed with satd. ammonium chloride. The organic layer was dried over Na_2SO_4, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent: 0-50% EtOAc/heptane) to provide 6-chloro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)pteridine-2,4(1*H*,3*H*)-dione (**Intermediate 111A**) as a light yellow solid: H NMR (400 MHz, *DMSO-d*$_6$) δ 12.40 (s, 1 H), 8.44 (br d, *J*=4.6 Hz, 1 H), 7.52 - 7.62 (m, 1 H), 7.28 - 7.39 (m, 3 H), 7.21 (br d, *J*=4.6 Hz, 1 H), 3.03 (dt, *J*=12.7, 6.4 Hz, 1 H), 2.07 (s, 3 H), 1.07 (br d, *J*=6.6 Hz, 3 H), 0.95 (br d, *J*=6.4 Hz, 3 H). *m/z* (ESI, +ve) 425.9 and 427.9 $(M+H)^+$.

Intermediate 160

1-((2*R*,5*S*)-2,5-Dimethylpiperazin-1-yl)prop-2-en-1-one 2,2,2-trifluoroacetate



Intermediate 160

[0682] To a solution of (2*S*,5*R*)-1-Boc-2,5-dimethylpiperazine (3.5 g, 16.3 mmol, Astatech) in dichloromethane (35 ml), anhydrous triethylamine (4.59 ml, 32.7 mmol) was added and the mixture was cooled to 0 °C in ice - water bath. Acryloyl chloride (1.46 ml, 18.0 mmol) was added dropwise over ~5 min at which point the reaction mixture became yellow and viscous and formation of a white precipitate was observed. Water (10 ml) was added and the mixture was removed from the ice bath and was allowed to stir for 10 min. The organic layer was separated, quickly washed with 2N HCl (40 ml), water and brine, filtered through pad of MgSO4 and concentrated to afford crude *tert*-butyl (2*S*,5*R*)-4-acryloyl-2,5-dimethylpiperazine-1-carboxylate (4.4 g) as yellow oil. This material was redissolved in DCM (40 ml) and TFA (12.6 ml, 163 mmol) was added and the mixture was stirred at rt for 4 h at which point complete deprotection was observed. The mixture was concentrated under reduced pressure and dried under vacuum to afford 1-((2*R*,5*S*)-2,5-dimethylpiperazin-1-yl)prop-2-en-1-one 2,2,2-trifluoroacetate (**Intermediate 160**, ~9 g) as an oil. Analysis by qNMR using benzyl benzoate as internal standard showed 40.2 wt% purity. The material was used without further purification. ^{1}H NMR (400 MHz, DMSO-d_6) δ 9.03 (br s, 1H), 8.95 (br s, 1H), 6.77 (dd, *J*=10.6, 16.8 Hz, 1H), 6.16 (dd, *J*=2.3, 16.8 Hz, 1H), 5.74 (dd, *J*=2.3, 10.6 Hz, 1H), 4.69 - 4.56 (m, 1H), 4.09 - 3.94 (m, 1H), 3.70 - 3.57 (m, 1H), 3.45 - 3.19 (m, 2H), 2.98 - 3.09 (m, 1H), 1.25 (d, *J*=6.8 Hz, 3H), 1.20 (d, *J*=6.8 Hz, 3H). m/z (ESI, +ve ion): 169.3 (M+H).

Intermediate 161

1,4-Diisopropyl-1H-pyrazol-5-amine



Intermediate 161

[0683] To a mixture of 2-formyl-3-methylbutanenitrile (1.5 ml, 13.5 mmol)<autotext key="18F898A2" name="[Reactants]" index="1" field="Reactants" type="field" length="53" /> in EtOH (45 mL)<autotext key="18F898A3" name="[Solvents]" index="1" field="Solvents" type="field" length="17" /> was added isopropylhydrazine (1.0 g, 13.5 mmol, Matrix Scientific)<autotext key="18F898A4" name="[Reactants]" index="2" field="Reactants" type="field" length="40" /> and the mixture <autotext key="" name="124102-8-02" type="SampleID" length="11" />was stirred and heated at 85 °C for 21 h. The mixture was concentrated *in vacuo*. The residue was treated with <autotext key="18F898A9" name="[FORM]" type="lookup" length="3" />satd $NaHCO_3$<autotext key="18F898AB" name="[WASHES]" type="lookup" length="11" /> (50 mL), extracted with EtOAc<autotext key="18F898AC" name="[SOLVENTS]" type="lookup" length="5" /> (2 × 50 mL), dried over Na_2SO_4<autotext key="18F898AF" name="[DESSICANT]" type="lookup" length="6" />, filtered and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (eluent: 0-50% EtOAc/heptane) to provide 1,4-diisopropyl-1H-pyrazol-5-amine (**Intermediate 161**, 1.04 g, 6.21 mmol, 46.0 % yield)<autotext key="18F898B6" name="[PRODUCTS]" index="1" field="Products" type="field" length="70" /> as light-yellow<autotext key="18F898B7" name="[COLOR]" type="lookup" length="12" /> syrupy solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 6.95 (s, 1 H), 4.65 (s, 2 H), 4.37 (spt, *J*=6.5 Hz, 1 H), 2.68 (spt, *J*=6.8 Hz, 1 H), 1.25 (d, *J*=6.6 Hz, 6 H), 1.08 (d, *J*=6.8 Hz, 6 H). *m/z* (ESI, +ve ion): 168.2 $(M+H)^+$.

Intermediate 162

3,5-Diisopropylpyridin-4-amine



Intermediate 162

[0684] To a mixture of 3,5-dibromo-4-amino-pyridine (5.0 g, 19.9 mmol, AstaTech, Inc.)<autotext key="197D23A4" name="[Reactants]" index="1" field="Reactants" type="field" length="51" />, Cphos (0.868 g, 1.989 mmol)<autotext key="197D23A5" name="[Reactants]" index="2" field="Reactants" type="field" length="27" />, palladium acetate (0.223 g, 0.994 mmol)<autotext key="197D23A6" name="[Reactants]" index="3" field="Reactants" type="field" length="44" />, and THF (40 mL)<autotext key="197D23A7" name="[Solvents]" index="1" field="Solvents" type="field" length="25" /> was added 2-

propylzinc bromide (1 M in THF) (80 mL, 80 mmol, Rieke Metals, Inc.)<autotext key="197D23A8" name="[Reactants]" index="4" field="Reactants" type="field" length="50" /> dropwise and the mixture was stirred at rt for 2 h. The reaction mixture was quenched with 5 N NaOH<autotext key="197C0DC7" name="[WASHES]" type="lookup" length="8" /> (100 mL), extracted with EtOAc<autotext key="197C0DC8" name="[SOLVENTS]" type="lookup" length="5" /> (2 × 50 mL), dried over Na_2SO_4<autotext key="197C0DCB" name="[DESSICANT]" type="lookup" length="6" />, filtered and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (eluent 0-30% DCM-MeOH (4:1)/DCM) to provide 3,5-diisopropylpyridin-4-amine (**Intermediate 162**, 1.4 g, 7.8 mmol, 39.4 % yield)<autotext key="197C7844" name="[Products]" index="2" field="Products" type="field" length="66" /> as orange<autotext key="197C7845" name="[COLOR]" type="lookup" length="6" /> syrup. <autotext key="197C7846" name="[FORM]" type="lookup" length="5" /> ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 7.88 (s, 2 H), 5.47 - 5.60 (m, 2 H), 3.00 (spt, *J*=6.8 Hz, 2 H), 1.17 (d, *J*=6.8 Hz, 12 H). *m/z* (ESI, +ve ion): 179.1 $(M+H)^+$.

Intermediate 163

2-Isopropyl-*N*^{4},*N*^{4}-dimethylpyridine-3,4-diamine



[0685] Step 1: 2-Chloro-*N*,*N*-dimethyl-3-nitropyridin-4-amine. To a stirring solution of 2,4-dichloro-3-nitropyridine (Ark Pharm, Inc., Arlington Heights, IL, 4.9 g, 25.4 mmol) in MeCN (20 mL) and triethylamine (3.93 ml, 27.9 mmol) at 0 °C under nitrogen was added dimethylamine, 2 M in THF (12.70 ml, 25.4 mmol) dropwise at a rate not to exceed an internal temp of 10 °C. After addition the ice bath was removed and the suspension stirred for 10 min reaching 20 °C. The reaction was then partitioned between EtOAc (100 mL) and sat. NaCl (25 mL). The organic was dried over MgSO4, filtered, and concentrated under reduced pressure.

The resulting solid was dissoved in DCM, dry silica added (30 g), then concentrated under reduced pressure. The products were then separated by silica gel chromatography (120 g) eluting products with a gradient of 15>60% EtOAc/heptane to afford 2-chloro-N,N-dimethyl-3-nitropyridin-4-amine as yellow solid. ^{1}H NMR (400 MHz, *CDCl$_3$*) δ 8.02 (d, *J*=6.01 Hz, 1H), 6.66 (d, *J*=6.22 Hz, 1H), 3.03 (s, 6H). *m/z* (ESI, +ve ion): 202.1 $(M+H)^+$.

[0686] Step 2: *N,N*-Dimethyl-3-nitro-2-(prop-1-en-2-yl)pyridin-4-amine. A suspension of 2-chloro-N,N-dimethyl-3-nitropyridin-4-amine (3.7 g, 18.35 mmol) in 1,4-dioxane (30 mL) and water (15 mL) was sparged with argon for 5 min followed by the addition of (4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isopropene (7.71 g, 45.9 mmol), (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.336 g, 0.459 mmol), and sodium carbonate (5.84 g, 55.1 mmol). After 5 min of additional argon sparging, the suspension was heated to refflux. After 2 hrs, the reaction was then partitioned between EtOAc (100 mL), sat. NaHCO3 (30 mL), and water (30 mL). The organic was then dried over $MgSO_4$, concentrated under reduced pressure, then purified by silica gel chromatography (80 g) eluting products with a gradient of 10>30% EtOAc/heptane to afford *N,N*-dimethyl-3-nitro-2-(prop-1-en-2-yl)pyridin-4-amine as yellow solid. ^{1}H NMR (400 MHz, *CDCl$_3$*) δ 8.23 (d, *J*=6.01 Hz, 1H), 6.64 (d, *J*=6.01 Hz, 1H), 5.18-5.22 (m, 1H), 5.02 (s, 1H), 2.99 (s, 6H), 2.15 (s, 3H). *m/z* (ESI, +ve ion): 208.3 $(M+H)^+$.

[0687] Step 3: 2-isopropyl-N^4,N^4-dimethylpyridine-3,4-diamine (Intermediate 163). A hydrogenation apparatus was set up with 250 mL glass cylinder, which was flushed with N_2, and then charged with palladium 10 wt. % on activated carbon (0.976 g, 0.917 mmol). A solution of *N,N*-dimethyl-3-nitro-2-(prop-1-en-2-yl)pyridin-4-amine (3.8 g, 18.34 mmol) in ethanol (40 mL) was then transferred into the glass cylinder. The cylinder was sealed, purged with H_2, and then vented - this procedure was repeated twice. Hydrogen was then charged into the cylinder to 45 psi. The reaction mixture was stirred under H_2 for 3 hr. The reaction was then filtered through a bed of Celite, concentrated under reduced pressure, then purified by silica gel chromatography (80 g) eluting products with a gradient of 10>30% Solvent A: EtOAc/EtOH (3:1 blend with 4% by volume of 2 M NH3 in MeOH)/Solvent B: heptane to afford 2-isopropyl-N^4,N^4-dimethylpyridine-3,4-diamine **(Intermediate 163)** as colorless oil. ^{1}H NMR (400 MHz, *CDCl$_3$*) δ 8.00 (d, *J*=5.18 Hz, 1H), 6.74 (d, *J*=5.39 Hz, 1H), 3.82 (br s, 2H), 3.04 (td, *J*=6.82, 13.53 Hz, 1H), 2.70 (s, 6H), 1.31 (d, *J*=6.84 Hz, 6H). *m/z* (ESI, +ve ion): 180.2 $(M+H)^+$.

Intermediate 164

***N*⁴,*N*⁴-Diethyl-2-isopropylpyridine-3,4-diamine**



[0688] Step 1: 2-Chloro-*N*,*N*-diethyl-3-nitropyridin-4-amine. To a stirring solution of 2,4-dichloro-3-nitropyridine (4.0 g, 20.73 mmol) in MeCN (16 mL; 4:1) and triethylamine (2.91 ml, 20.7 mmol) at 0 °C under nitrogen was added diethylamine (2.37 ml, 22.8 mmol) in THF (10 mL) dropwise. After the addition was complete, the ice bath was removed and the suspension was stirred for 3 hrs. The reaction was then partitioned between EtOAc (100 mL) and sat. NaCl (25 mL). The organic was dried over MgSO4, filtered, and concentrated under reduced pressure, then purified by silica gel chromatography (120 g) elutiing products with a gradient of 15>60% EtOAc/heptane to afford 2-chloro-*N*,*N*-diethyl-3-nitropyridin-4-amine as yellow solid. ^{1}H NMR (400 MHz, *CDCl*$_3$) δ 8.00 (d, *J*=6.01 Hz, 1H), 6.69 (d, *J*=6.22 Hz, 1H), 3.31 (q, *J*=7.12 Hz, 4H), 1.20 (t, *J*=7.05 Hz, 6H). *m/z* (ESI, +ve ion): 230.2 $(M+H)^+$.

[0689] Step 2: *N*,*N*-Diethyl-3-nitro-2-(prop-1-en-2-yl)pyridin-4-amine. A suspension of 2-chloro-*N*,*N*-diethyl-3-nitropyridin-4-amine (2.2 g, 9.58 mmol) in 1,4-dioxane (20 mL) and water (10 mL) was sparged with argon for 5 min followed by the addition of (4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isopropene (3.22 g, 19.2 mmol), (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.175 g, 0.239 mmol), and sodium carbonate (3.05 g, 28.7 mmol). After 5 min of additional sparging, the suspension was heated to refflux for 2 hrs. The reaction was then partitioned between EtOAc (60 mL) and sat. NaHCO3 (20 mL). The organic was then dried over MgSO4, concentrated under reduced pressure, then purified by silica gel chromatography (40 g) eluting products with a gradient of 10>60% EtOAc/heptane to afford *N*,*N*-diethyl-3-nitro-2-(prop-1-en-2-yl)pyridin-4-amine as yellow oil. ^{1}H NMR (400 MHz, *CDCl*$_3$) δ 8.23 (d, *J*=6.22 Hz, 1H), 6.70 (d, *J*=6.01 Hz, 1H),

5.21 (s, 1H), 5.04 (s, 1H), 3.27 (q, *J*=7.05 Hz, 4H), 2.14 (s, 3H), 1.17 (dd, *J*=6.63, 13.68 Hz, 6H). *m/z* (ESI, +ve ion): 236.2 $(M+H)^+$.

[0690] **Step 3: N^4,N^4-Diethyl-2-isopropylpyridine-3,4-diamine (Intermediate 164):** A hydrogenation apparatus was set up with a 250 mL glass cylinder, was sparged with N_2, and was then filled with palladium 10 wt. % on activated carbon (0.249 g, 0.234 mmol). A solution of *N,N*-diethyl-3-nitro-2-(prop-1-en-2-yl)pyridin-4-amine (2.2 g, 9.35 mmol) in ethanol (40 mL) was then transferred into the glass cylinder. The cylinder was sealed, purged with H_2, and then vented - this procedure was repeated twice. Hydrogen was then charged into the cylinder to 45 psi. The reaction mixture was started stirring under H_2 for 6 hrs. The reaction was then filtered through a bed of Celite, and the filtrated concentrated under reduced pressure to afford N^4,N^4-diethyl-2-isopropylpyridine-3,4-diamine (**Intermediate 164**) as amber oil. ^{1}H NMR (400 MHz, *CDCl$_3$*) δ 7.98 (d, *J*=5.18 Hz, 1H), 6.76 (d, *J*=5.18 Hz, 1H), 3.92 (br s, 2H), 2.93-3.09 (m, 5H), 1.31 (d, *J*=6.84 Hz, 6H), 1.01 (t, *J*=7.05 Hz, 6H). *m/z* (ESI, +ve ion): 208.3 $(M+H)^+$.

Intermediate 165

***tert*-Butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate**



[0691] A 3 neck flask equipped with overhead stirrer, thermocouple, and nitrogen inlet was charged with *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (Astatech, Bristol, PA, USA, 78 g, 365 mmol) and THF (400 mL). The resulting solution was degassed and purged with argon then cooled to 0 °C. Isopropylmagnesium chloride (183 mL, 365 mmol) (2M in THF) was added dropwise via an addition funnel to the solution. The mixture was stirred for 1 hr at 0 °C. Zinc chloride (192 mL, 365 mmol) (1.9M in 2-Me THF) was added dropwise via an addition funnel. The mixture was warmed to RT and stirred for 12 hours. In a separate flask was dissolved 2,5-dichloro-6-(2-fluorophenyl)nicotinonitrile (**Intermediate 101A**, 65 g, 243 mmol) in THF (400 mL). The orange solution was added to the reaction flask dropwise via an addition funnel. The mixture was stirred overnight at RT. The hazy reaction mixture

was cooled to 0 °C. Ten volumes of ethyl acetate and 10 volumes of 15 wt% NH_4Cl (aq) were added. The biphasic mixture was stirred for 1 hour and the aqueous layer was separated. The organic layer was washed with 15 wt% sat. NaCl. The organic layer was concentrated to a dark orange oil. Isopropanol was added and evaporated several times. Eight volumes of 1:1 isopropanol/water was added to the crude oil. The resulting solids were filtered through a glass frit. The filter cake was washed with 25% IPA/water solution. The solid was dried under vacuum with nitrogen sweep to afford *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 165**). ^{1}H NMR (500 MHz, DMSO-d_6) δ 8.03 - 8.19 (m, 1 H), 7.57 - 7.63 (m, 1 H), 7.36 - 7.56 (m, 4 H), 4.18 (br s, 1 H), 3.62 (br s, 1 H), 3.39 (br d, *J*=13.75 Hz, 1 H), 3.32 (s, 2 H), 3.02 - 3.23 (m, 1 H), 1.35 -1.45 (m, 9 H), 1.09 - 1.21 (m, 4 H), 1.05 (br dd, *J*=12.52, 6.16 Hz, 2 H). *m/z* (ESI, +ve ion): 481.2 $(M+H)^+$.

Intermediate 168

***tert*-Butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(o-tolyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate**





Intermediate 168

[0692] Step 1: 2,5-Dichloro-6-(*o*-tolyl)nicotinonitrile. A 3 neck flask equipped with an overhead stirrer, thermocouple, condensor and nitrogen inlet was charged with 2,5,6-trichloronicotinonitrile (Syngene, 70 g, 337 mmol), o-tolylboronic acid (Combi-Blocks, San Diego, CA, USA, 48.2 g, 354 mmol), K_2CO_3 (140 g, 1012 mmol) and $PdCl_2$(dppf) (12.35 g, 16.87 mmol). Nitrogen sparged 1,4-dioxane (700 mL) was added to the flask followed by nitrogen sparged water (233 mL). The reaction flask was purged with argon, stirred for 10 min, and then warmed to 65 °C for 1 hour. The reaction was cooled to RT and stirred overnight. The aqueous layer was separated from the reaction mixture. The organic layer was diluted

with ethyl acetate and water. The aqueous layer was separated. The organic layer was washed with brine. The organic layer was dried over sodium sulfate and concentrated to dryness. The residue was taken up in neat DCM and filtered through a silica gel pad washing with DCM. The filtrate was concentrated to a reddish oil under reduced pressure. The oil was dissolved in 500 mL isopropanol. The resulting red solution was charged with crystalline seed and the slurry was stirred for 4 hours at RT. The slurry was filtered, washed with minimal cold IPA, then dried under vacuum with nitrogen sweep to afford 2,5-dichloro-6-(*o*-tolyl)nicotinonitrile as an off white solid. ^{1}H NMR (500 MHz, DMSO-*d*6) δ 8.92 (s, 1 H), 7.27 - 7.44 (m, 4 H), 2.11 (s, 3 H). *m/z* (ESI, +ve ion): 263.2 $(M+H)^+$.

[0693] Step 2: *tert*-Butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(*o*-tolyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (Intermediate 168). A 3L (3 neck) round bottom flask equiped with an overhead stirrer, thermocouple, and nitrogen inlet was charged with *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (Astatech, Bristol, PA, USA,, 57.2 g, 267 mmol) and THF (300 mL). The resulting solution was degassed, purged with argon, then cooled to 0 °C. To the solution was charged isopropylmagnesium chloride (133 mL, 267 mmol) (2M in THF) dropwise via addition funnel and the mixture was stirred for 1 hr at 0 °C. Zinc chloride (140 mL, 267 mmol) (1.9M in THF) was added dropwise via addition funnel. The reaction was then allowed to warm to RT and stir overnight. A solution of 2,5-dichloro-6-(*o*-tolyl)nicotinonitrile (46.8 g, 178 mmol) in THF (300 mL) was charged to reaction flask dropwise via addition funnel. The reaction was then stirred overnight at RT. The reaction was then quenched into a stirring biphasic mixture of 15 wt% NH_4Cl (500mL) and 500mL EtOAc. The mixture was stirred for 10 min and the aqueous layer was separated. The organic layer was washed with 15 wt% sat. NaCl. The organic layer was concentrated to orange oil under reduced pressure. A 3 neck flask equipped with overhead stirrer, thermocouple and nitrogen inlet was charged with fine filter frit and vacuum port. The crude oil was dissolved in 3 volumes of IPA (250mL), and then filtered to remove inorganic salts. The orange solution was seeded with a crystal and stirred for 18 hrs at RT. The solids were isolated by filtration, and washed with cold IPA. The solids were then further purified by a plug of silica gel and washing with 100% EtOH/EtOAc (3:1 blend) to afford *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(*o*-tolyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (Intermediate 168) as a white solid. *m/z* (ESI, +ve ion): 477.2 $(M+H)^+$.

Intermediate 170

***tert*-Butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-isopropylphenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate**





Intermediate 170

[0694] Step 1: 2,5-Dichloro-6-(2-isopropylphenyl)nicotinonitrile. To a 2-L three neck round bottom flask equipped with condenser with a nitrogen inlet, an overhead stirrer and thermocouple was added 2,5,6-trichloronicotinonitrile (90 g, 434 mmol), 2-(1-methylethyl)phenyl]-boronic acid (74.7 ml, 456 mmol), potassium carbonate (180 g, 1302 mmol), and (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (15.80 g, 21.69 mmol). The flask was purged and filled with argon 3 times. Water (300 ml) and 1,4-dioxane (900 ml) were added by canulation. The reaction was heated to 65 °C and stirred for 1.5 hrs. The reaction was then diluted with water and extracted three times with ethyl acetate. The organics were dried over sodium sulfate and concetrated under reduced pressure. The dried organics were dissolved in DCM and filtered through sand and silica gel. The filtrate was evaporated. The residue was taken up in isopropanol (600 mL). The resulting solid product was then isolated by filtration to afford 2,5-dichloro-6-(2-isopropylphenyl)nicotinonitrile as white solid. ^{1}H NMR (500 MHz, DMSO-*d6*) δ 8.91 (s, 1 H), 7.47 - 7.52 (m, 2 H), 7.29 - 7.35 (m, 1 H), 7.21 (d, *J*=7.40 Hz, 1 H), 2.54 - 2.62 (m, 1 H), 1.11 (br d, *J*=6.75 Hz, 6 H). *m/z* (ESI, +ve ion): 291.2 $(M+H)^+$.

[0695] Step 2: *tert*-Butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-isopropylphenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate. A 3 neck flask equipped with overhead stirrer, thermocouple and nitrogen inlet was charged with *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (Astatech, Bristol, PA, USA, 44.2 g, 206 mmol) and THF

(250 mL). The resulting solution was degassed and purged with argon and cooled to 0 °C. Isopropylmagnesium chloride (103 mL, 206 mmol) (2M in THF) was added dropwise via addition funnel into the solution. The mixture was stirred for 1 hr at 0 °C. Zinc chloride (108 mL, 206 mmol) (1.9M in 2-Me THF) was added dropwise via addition funnel. The mixture was allowed to warm to RT and was stirred for 12 hours. In a separate flask was dissolved 2,5-dichloro-6-(2-isopropylphenyl)nicotinonitrile (40 g, 137 mmol) in THF (250 mL). The mixture was added to the reaction flask dropwise via addition funnel. The mixture was stirred at RT for 18 hrs. The hazy reaction mixture was cooled to 0 °C and charged with 10 volumes of ethyl acetate and 10 volumes of 15 wt% NH_4Cl (aq). The biphasic mixture was stirred for 1 hour and the aqueous layer was separated. The organics were washed with 15 wt% brine. The organics were concentrated to a dark orange oil. Isopropanol was added and evaporated several times. Eight volumes of 1:1 isopropanol/water were added to the crude oil. The resulting solid was isolated by filtration to afford ***tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-isopropylphenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (Intermediate 170)** as an off white solid. *m/z* (ESI, +ve ion): 505.2 $(M+H)^+$.

Intermediate 175

***tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4-chloro-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**



[0696] **Step 1-2: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4-chloro-6-isopropylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**. Synthesized analogous to Steps 1 and 2 in Method 102 using **Intermediate 177** in Step 1 and performing Step 2 in toluene. ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.93 (1 H, s) 8.12 (1 H, s) 7.40 - 7.48 (1 H, m) 7.09 - 7.25 (3 H, m) 3.46 - 5.07 (6 H, m) 2.85 - 2.94 (1 H, m) 1.51 (9 H, s) 1.42 - 1.50 (3 H, m) 1.23 - 1.32 (6 H, m) 1.06 - 1.13 (3 H, m). m/z (ESI, +ve ion): 642.2 $(M+H)^+$.

Intermediate 176

***tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4-chloro-6-isopropylpyrimidin-5-yl)-2-oxo-7-(*o*-tolyl)-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**



[0697] **Steps 1-2: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4-chloro-6-isopropylpyrimidin-5-yl)-2-oxo-7-(*o*-tolyl)-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**. Synthesized analogous to Steps 1 and 2 in Method 102 using **Intermediate 168** and **Intermediate 177** in Step 1 and performing Step 2 in toluene. ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.87 (1 H, s) 8.11 (1 H, s) 7.26 - 7.31 (1 H, m) 7.15 - 7.21 (2 H, m) 7.03 - 7.08 (1 H, m) 3.43 - 5.04 (6 H, m) 2.82 - 2.91 (1 H, m) 2.00 - 2.04 (3 H, m) 1.49 (9 H, s) 1.41 - 1.51 (3 H, m) 1.21 - 1.31 (6 H, m) 1.00 - 1.05 (3 H, m). m/z (ESI, +ve ion): 638.2 $(M+H)^+$.

Intermediate 177

4-Chloro-6-isopropylpyrimidin-5-amine



Intermediate 177

[0698] 4-Chloro-6-isopropylpyrimidin-5-amine. 2-Propylzinc bromide (1.0 M in THF, 13.42 ml, 13.42 mmol) was added to a flask charged with 4,6-dichloro-5-aminopyrimidine (2.00 g, 12.20 mmol) and Xantphos Pd G3 (0.578 g, 0.610 mmol) under an argon atmosphere. The reaction mixture was stirred at room temperature for 4 h. The reaction mixture was diluted with EtOAc (125 mL) and quenched with saturated aqueous NH_4Cl (75 mL). The organic layer was separated, washed with saturated aqueous sodium chloride (75 mL), dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (eluent 0-30% EtOAc/heptane) to provide 4-chloro-6-isopropylpyrimidin-5-amine as a yellow solid (**Intermediate 177,** 1.18 g, 6.88 mmol, 56.4 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 8.39 (1 H, s) 4.09 (2 H, br s) 3.02 (1 H, spt, *J*=6.74 Hz) 1.31 (6 H, d, *J*=6.84 Hz). m/z (ESI, +ve ion): 172.1 $(M+H)^+$.

Intermediate 185

***tert*-Butyl (2*R*,5*S*)-4-(6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**



Intermediate 72A **Intermediate 185**

[0699] Step 1. 4,7-Dichloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-d]pyrimidin-2(1*H*)-one. To a RBF was added 7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (4.81 g, 13.8 mmol, **Intermediate 72A**), toluene (55 ml), and hunig's base (4.8 ml, 27.6 mmol). To this was added phosphorous oxychloride (2.57 ml, 27.6 mmol) dropwise at rt and the reaction was stirred for 50 minutes at 50 °C at which point LC/MS showed near complete consumption of SM and desired product. The reaction was concentrated under reduced pressure and used directly in the next step. ◻

[0700] Step 2. *tert*-Butyl (2*R*,5*S*)-4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a flask charged with 4,7-dichloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (5.2 g, 14.2 mmol) was added DMF (118 ml) followed by (2*R*,5*S*)-*tert*-butyl 2,5-dimethylpiperazine-1-carboxylate (4.55 ml, 21.2 mmol). To the solution was added hunig's base (12.4 ml, 70.8 mmol) dropwise, with stirring. Upon complete addition, the reaction was quenched by the slow addition of water then further diluted with water and EtOAc before transferring to a separated funnel. The phases were mixed and the organic layer was separated. The organic phase was washed with 1M LiCl and brine then dried over magnesium sulfate. The material was purified via ISCO (120g, dry load, 0-80% 3:1 EtOAc/EtOH in heptane) to afford *tert*-butyl (2*R*,5*S*)-4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (6.87 g, 12.60 mmol, 89 % yield). *m/z* (ESI, +ve ion): 545.2 $(M+H)^+$.

[0701] Step 3. *tert*-Butyl (2*R*,5*S*)-4-(6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A resealable vial was charged with potassium acetate (0.35 g, 3.6 mmol), *tert*-butyl (2*R*,5*S*)-4-(7-chloro-6-fluoro-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.39 g, 0.72 mmol) and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (ii) dichloromethane adduct (0.052 g, 0.072 mmol). The vial was sealed and evacuated / backfilled with Nitrogen. Dioxane (4.5 ml) was added followed by water (0.2 ml). The cap was replaced and the reaction was heated to 90 °C for 2 minutes and (2-fluorophenyl)boronic acid (0.2 g, 1.43 mmol) were added and the reaction was stirred at 90 °C for 2 hours. The reaction was poured into a separated funnel containing water and EtOAc. The phases were mixed and the organic phase was separated, washed with brine, dried over magnesium sulfate and concentrated under reduced pressure. The crude residue was purified via ISCO (40 g, 0-60% 3:1 EtOAc/EtOH in heptane) to afford *tert*-butyl (2*R*,5*S*)-4-(6-fluoro-7-(2-fluorophenyl)-1-(2-isopropyl-4-methylpyridin-3-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 185**, 0.262 g, 0.433 mmol, 60.6 % yield) as off-white solid. *m/z* (ESI, +ve ion): 605.3 $(M+H)^+$.

Intermediate 186

4-(Dimethylamino)but-2-ynoic acid



[0702] To a solution of 1-dimethylamino-2-propyne (3.0 g, 36.1 mmol) in THF (30 mL) cooled in a -78 °C bath was slowly added butyl lithium, 2.5 M in hexanes (14.4 mL, 36.1 mmol) dropwise. The solution was stirred at -78 °C for 1 h. The reaction mixture was then quenched with crushed dry ice (carbon dioxide (1.59 g, 36.1 mmol) and the reaction mixture was removed from the ice bath and allowed to warm to r.t. The reaction mixture was concentrated to dryness, treated with water (50 mL), extracted with EtOAc (30 mL) and the aqueous was again concentrated on the rotovap. The crude residue was taken up in MeOH and filtered and the concentrated residue was dried on the lyophilizer overnight affording 4-(dimethylamino)but-2-ynoic acid (5.67 g, 44.6 mmol, 123 % yield, 75% pure) as a light yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 3.15 (2 H, s), 2.15 (6 H, s). *m/z* (ESI, +ve ion): 128.1 $(M+H)^+$.

Intermediate 187

6-((Dimethylamino)methyl)-2-isopropyl-4-methylpyridin-3-amine



[0703] **Step 1. 6-Bromo-2-isopropyl-4-methylpyridin-3-amine.** To 2-isopropyl-4-methylpyridin-3-amine (**Intermediate R**, 4 g, 26.6 mmol) dissolved in *N,N*-dimethylformamide (81 ml) at 0°C was added bromosuccinimide (4.98 g, 28.0 mmol) in one portion. The reaction mixture was stirred at 0°C for 1 h. Water was added to the resulting reaction mixture to stop the reaction. The aqueous layer was extracted with ethyl acetate (2x). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The crude material was purified by chromatography through an ISCO 24 Gold column, eluting with a gradient of 0% to 25% 3:1 EtOAc/EtOH in heptane, to provide 6-bromo-2-isopropyl-4-

methylpyridin-3-amine (5 g, 21.82 mmol, 82 % yield) as orange oil. ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 7.00 (s, 1H), 3.58 (br s, 2H), 2.96 (td, *J*=6.74, 13.48 Hz, 1H), 2.13 (s, 3H), 1.28 (d, *J*=6.63 Hz, 6H). *m/z* (ESI, +ve ion): 228.9 $(M+H)^+$.

[0704] **Step 2. 5-Amino-6-isopropyl-4-methylpicolinonitrile.** A 5 mL microwave vial containing was charged with a magnetic stir bar, 6-bromo-2-isopropyl-4-methylpyridin-3-amine (1.16 g, 5.06 mmol), tetrakis(triphenylphosphine)palladium(0) (0.585 g, 0.506 mmol), dicyanozinc (0.832 g, 7.09 mmol). The vial was evacuated and flushed with nitrogen. *N,N*-dimethylformamide (13 ml) was added the reaction mixture was stirred and heated in a Emrys Optmizer microwave reactor (Personal Chemistry, Biotage AB, Inc., Upssala, Sweden) at 110°C for 30 min. The reaction mixture was partitioned between EtOAc and brine. The layers were separated and the aqueous layer was further extracted with EtOAc (2x). The combined organic extracts were dried over sodium sulfate, filtered, and concentrated. The crude material was purified on a 40 g HP-GOLD column eluting with a gradient of 0-30% 3:1 EtOAc/EtOH in heptane over 30 min. to afford 5-amino-6-isopropyl-4-methylpicolinonitrile as a dark-red oil. *m/z* (ESI, +ve ion): 176.1 $(M+H)^+$.

[0705] **Step 3. 5-Amino-6-isopropyl-4-methylpicolinaldehyde (Intermediate 239).** A nitrogen-flushed 100 mL round-bottomed flask containing 5-amino-6-isopropyl-4-methylpicolinonitrile (1 g, 5.7 mmol) was charged with a magnetic stir bar and tetrahydrofuran (19 mL). diisobutylaluminium hydride (45.7 ml, 45.7 mmol) was added dropwise at room temp and the resulting orange solution stirred for 30 min. The reaction mixture was then cooled to 0 °C in an ice-water bath and quenched by adding 1 M Rochelle salt (40 mL) dropwise to the reaction mixture (hydrogen gas was generated) and diluted with EtOAc (100 mL). The resulting biphasic mixture was stirred for 15 min at room temp and was then filtered through a glass frit and washed with EtOAc. The filtrate was transferred to a separatory funnel and the layers were partitioned. The aqueous phase was extracted with EtOAc (2 x 70 mL) and the combined organic extracts were dried over anhydrous sodium sulfate, filtered, and concentrated. The crude material was dissolved in DCM and then purified on a 40 g HP-GOLD column eluting with a gradient of 0-40% 3:1 EtOAc/EtOH in heptane over 30 min, to afford 5-amino-6-isopropyl-4-methylpicolinaldehyde (**Intermediate 239** 0.18 g, 1.0 mmol, 18 % yield) as a thick yellow oil which was carried forward in the next step without further purification. *m/z* (ESI, +ve ion): 179.1 $(M+H)^+$.

[0706] **Step 4. 6-((Dimethylamino)methyl)-2-isopropyl-4-methylpyridin-3-amine.** A nitrogen-purged 100 mL round-bottomed flask containing 5-amino-6-isopropyl-4-

methylpicolinaldehyde (**Intermediate 239,** 180 mg, 1.0 mmol) and a magnetic stir bar was charged with tetrahydrofuran (5 mL), dimethylamine solution 2.0m in THF (0.56 mL, 1.1 mmol), acetic acid, glacial (0.058 mL, 1.0 mmol), and sodium triacetoxyborohydride (428 mg, 2.02 mmol). The resulting cloudy mixture was stirred at RT for 2 h then quenched by adding water (~20 mL) and saturated aqueous sodium bicarbonate (~20 mL) and the mixture was diluted with EtOAc (~50 mL). The biphasic mixture was transferred to a separatory funnel and the layers were partitioned. The aqueous phase was extracted with EtOAc until the desired mass was no longer detected in the aqueous phase. The combined organic extracts were dried over magnesium sulfate, filtered, and concentrated to afford crude 6-((dimethylamino)methyl)-2-isopropyl-4-methylpyridin-3-amine (**Intermediate 187**) as a light-yellow oil ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 6.92 (s, 1H), 3.53 (br s, 2H), 3.46 (s, 2H), 2.96-3.10 (m, 1H), 2.27 (s, 6H), 2.16 (s, 3H), 1.29 (d, *J*=6.63 Hz, 6H). *m/z* (ESI, +ve ion): 208.3 $(M+H)^+$.

Intermediate 188

2-Isopropyl-4-methyl-6-(pyrrolidin-1-ylmethyl)pyridin-3-amine



Intermediate 239 Intermediate 188

[0707] A nitrogen-purged 50 mL round-bottomed flask containing 5-amino-6-isopropyl-4-methylpicolinaldehyde (**Intermediate 239,** 265 mg, 1.49 mmol) and a magnetic stir bar was charged with tetrahydrofuran (5 mL), pyrrolidine (137 µl, 1.64 mmol), acetic acid, glacial (85 µl, 1.49 mmol), and sodium triacetoxyborohydride (630 mg, 2.97 mmol). The resulting cloudy mixture was stirred at RT for 18 h. The reaction was quenched by adding water (~10 mL) and saturated aqueous sodium bicarbonate (~10 mL) and the mixture was diluted with EtOAc (~40 mL). The biphasic mixture was transferred to a separatory funnel and the layers were partitioned. The aqueous phase was extracted with EtOAc until the desired mass was no longer detected in the aqueous phase. The combined organic extracts were dried over magnesium sulfate, filtered, and concentrated to afford crude 2-isopropyl-4-methyl-6-(pyrrolidin-1-ylmethyl)pyridin-3-amine as a light-yellow oil (**Intermediate 188**). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 6.93 (s, 1H), 3.66 (s, 2H), 3.52 (br s, 2H), 2.93-3.11 (m, 1H), 2.60 (br s, 4H), 2.15 (s, 3H), 1.78 (td, *J*=3.32, 6.63 Hz, 4H), 1.29 (d, *J*=6.84 Hz, 6H). *m/z* (ESI, +ve ion): 234.2 $(M+H)^+$.

Intermediate 189

5-Isopropyl-2,3-dimethylpyridin-4-amine



[0708] **Step 1. 2,3-Dimethyl-5-(prop-1-en-2-yl)pyridin-4-amine.** To a RBF was added 5-bromo-2,3-dimethylpyridin-4-amine (ACES Pharma, 1.05 g, 5.22 mmol), 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (0.965 g, 5.74 mmol) and AmPhos (0.074 g, 0.10 mmol). The flask was evacuated under vacuum and then nitrogen was flushed through it for 1 min. 1,4-dioxane (17 mL) and aqueous potassium phosphate (3.33 g, 15.7 mmol in 6 mL of water) was added and the reaction was warmed to 90 °C for 3 h. Added more AmPhos (100 mg, 0.25 eq) and stirred at 90 °C for another 16 h. The mixture was partitioned between EtOAc and brine. The aqueous layer was back extracted with EtOAc (3×) and the combined organics was dried (Na2SO4) and concentrated. The crude was purified by chromatography through a Biotage 40 g ultra column, eluting with a gradient of 0% to 50% EtOAc in heptane, to provide 2,3-dimethyl-5-(prop-1-en-2-yl)pyridin-4-amine (0.27 g, 1.6 mmol, 32 % yield) as light-green oil. *m/z* (ESI, +ve ion): 163.2 $(M+H)^+$.

[0709] **Step 2. 5-Isopropyl-2,3-dimethylpyridin-4-amine.** A mixture of 2,3-dimethyl-5-(prop-1-en-2-yl)pyridin-4-amine (0.27 g, 1.6 mmol) in ethanol (5.5 mL) under nitrogen was treated with palladium 10 wt. % on activated carbon (0.089 g, 0.083 mmol) in a 75-mL glass reaction vessel, the solution was purged 5×40 psi with hydrogen and allowed to stir under an atmosphere of hydrogen at ca. 30-40 psi for 3 h. The reaction mixture was filtered through a pad of celite, washed with EtOH and concentrated to dryness affording 5-isopropyl-2,3-dimethylpyridin-4-amine (**Intermediate 189,** 0.29 g, 1.7 mmol, 104 % yield) as orange viscous oil. *m/z* (ESI, +ve ion): 165.2 $(M+H)^+$.

Intermediate 190

4-Chloro-2-isopropylpyridin-3-amine



Intermediate 190

[0710] A mixture of 2,4-dichloro-3-aminopyridine (5.00 ml, 30.7 mmol, Combi-Blocks) and 1,1'-bis(diphenylphosphino)ferrocene-palladium dichloride (2.244 g, 3.07 mmol) was purged with N2 followed by the addition of THF (100 mL) and 2-propylzinc bromide (46.0 ml, 46.0 mmol). The resulting mixture was heated at 60 °C. After 2 h the reaction went to completion. The reaction was brought to rt, washed with sat. $NaHCO_3$ and extracted with EtOAc. The combined organics were purified by chromatography on silica gel using 0-30% EtOAc in heptane to afford 4-chloro-2-isopropylpyridin-3-amine (**Intermediate 190,** 3.45 g, 20.22 mmol, 65.9 % yield) as a light yellow oil. ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 7.92 (d, *J*=5.2 Hz, 1 H), 7.07 (d, *J*=5.4 Hz, 1 H), 4.09 (br s, 2 H), 2.86 - 3.22 (m, 1 H), 1.32 (d, *J*=6.6 Hz, 6 H). m/z (ESI, +ve ion): 170 (M+H).

Intermediates 240 and 191

6,7-Dichloro-1-(4-chloro-2-isopropylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione

6-Chloro-1-(4-chloro-2-isopropylpyridin-3-yl)-7-methoxypyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



Intermediate 240 Intermediate 191

[0711] To a solution of 6,7-dichloro-1-(4-chloro-2-isopropylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.500 g, 1.297 mmol, **Intermediate 240**) in dry MeOH (15 mL) at rt was added sodium methoxide (0.700 g, 12.97 mmol) and the resulting mixture was heated at 50C for 2 h. The reaction went to completion and concentrated.The white residue was washed with water and extracted with EtOAc. The combined organics were dried over Na_2SO_4, filtered and concentrated to afford 6-chloro-1-(4-chloro-2-isopropylpyridin-3-yl)-7-

methoxypyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione, **Intermediate 191**. ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 8.58 (d, *J*=5.2 Hz, 1 H), 8.36 (s, 1 H), 7.34 (d, *J*=5.2 Hz, 1 H), 3.61 (s, 3 H), 2.88 (br d, *J*=6.6 Hz, 1 H), 1.22 (br d, *J*=6.6 Hz, 3 H), 1.20 - 1.21 (m, 1 H), 1.13 (br d, *J*=6.6 Hz, 3 H). m/z (ESI, +ve ion): 380.6 (M+H).

Intermediate 192

Dimethyl(2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)phosphine oxide



[0712] Step 1. (2-Iodophenyl)dimethylphosphine oxide. A mixture of (2-aminophenyl)dimethylphosphine oxide (1.500 ml, 8.87 mmol, AstaTech, Inc.), copper(i) iodide (0.602 ml, 17.73 mmol) and 1,1-dimethylethylnitrite (1.829 ml, 17.73 mmol) in MeCN (25 mL) was heated at 60C overnight. The reaction went to completion, poured into water and extracted with EtOAc. The combined organics were dried over Na_2SO_4, filtered, concentrated and purified by chromatography on silica gel using 0-50% EtOAc in heptane to afford (2-iodophenyl)dimethylphosphine oxide (0.88 g, 3.14 mmol, 35.4 % yield) as brown solid. m/z (ESI, +ve ion): 280.8 (M+H).

[0713] Step 2. A mixture of (2-iodophenyl)dimethylphosphine oxide (0.88 g, 3.14 mmol), bis(pinacolato)diboron (1.037 g, 4.09 mmol), (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.115 ml, 0.157 mmol) andpotassium acetate (0.617 g, 6.28 mmol) was purged with N_2 followed by the addition of 1,4-dioxane and the resulting mixture was heated at 90 C for 6 h. The reaction was washed with sat. $NaHCO_3$, extracted with EtOAc, DCM and purified by chromatography on silica gel using 0-10% MeOH in DCM to afford a brown oil as dimethyl(2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)phosphine oxide (**Intermediate 192,** 0.15 g, 0.536 mmol, 17.04 % yield). 1H NMR (400 MHz, *CHLOROFORM-d*) δ 7.92 - 8.03 (m, 1 H), 7.51 (br d, *J*=6.6 Hz, 3 H), 1.82 - 1.93 (m, 6 H), 1.39 (s, 12 H). m/z (ESI, +ve ion): 380.6 (M+H).

Intermediate 193

2,6-Diisopropylaniline



Intermediate 193

[0714] **Step 1. 2,6-Di(prop-1-en-2-yl)aniline.** A mixture of 2,6-dibromoaniline (5.00 ml, 19.93 mmol, Ark Pharma),2-isopropenylboronic acid, pincol ester (8.37 g, 49.8 mmol, Combi-Blocks), sodium carbonate (12.67 g, 120 mmol) and tetrakis (2.303 g, 1.993 mmol) in 1,4-dioxane/water (24/6 mL) was heated at 90 C for 24 h. The reaction went to completion, quenched with sat. $NaHCO_3$ and extracted with EtOAc. The combined organics were purified by chromatography on silica gel using 0-50 % EtOAc in heptane to afford 2,6-di(prop-1-en-2-yl)aniline (1.31 g, 7.56 mmol, 37.9 % yield) as a greenish oil with another impurity. m/z (ESI, +ve ion): 174 (M+H).

[0715] **Step 2. 2,6-Diisopropylaniline.** A mixture of 2,6-di(prop-1-en-2-yl)aniline (1.31 g, 7.56 mmol), ammonium formate (15.02 ml, 302 mmol) and palladium hydroxide carbon (2.65 ml, 3.78 mmol) was purged with N_2, followed by the addition of EtOH (30 mL). The resulting mixture was heated to reflux for 1 h. The reaction went to completion and filtered through celite. The filtrate was washed with sat. $NaHCO_3$, water, brine, dried over Na_2SO_4, and concentrated. The crude product was chomatographed on silica gel using 0-70% EtOAc in heptane to afford 2,6-diisopropylaniline (**Intermediate 193**) as a brown oil. 1H NMR (400 MHz, *CHLOROFORM-d*) δ 7.04 (d, *J*=7.7 Hz, 2 H), 6.77 - 6.86 (m, 1 H), 2.88 - 3.01 (m, 2 H), 1.27 (d, *J*=6.8 Hz, 12 H). NH2 not observed. m/z (ESI, +ve ion): 178 (M+H).

Intermediate 194

2,4-Diisopropylpyridin-3-amine



Intermediate 194

[0716] **Step 1. 2,4-Di(prop-1-en-2-yl)pyridin-3-amine.** A mixture of 2,4-dichloro-3-aminopyridine (4.40 ml, 27.0 mmol, Combi-Blocks) ,4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (10.89 g, 64.8 mmol, Combi-Blocks), sodium carbonate (14.30 g, 135 mmol) and chloro(2-dicyclohexylphosphino-2',4',6'-tri-i-propyl-1,1'-biphenyl)[2-(2-aminoethyl)phenyl] palladium(ii) methyl-t-butylether adduct (2.232 ml, 2.70 mmol) in 1,4-dioxane/water (100/25 mL) was heated at 90 C for 6 h. The reaction went to completion,

quenched with sat. $NaHCO_3$ and extracted with EtOAc. The combined organics were purified by chromatography on silica gel using 0-80 EtOAc in heptane to afford 2,4-di(prop-1-en-2-yl)pyridin-3-amine (4.5 g, 25.8 mmol, 96 % yield) as a yellow oil. m/z (ESI, +ve ion): 175 (M+H).

[0717] **Step 2. 2,4-Diisopropylpyridin-3-amine.** Through a mixture of 2,4-di(prop-1-en-2-yl)pyridin-3-amine (4.5 g, 25.8 mmol) and palladium 10 wt. % on activated carbon (0.229 ml, 2.58 mmol) in EtOH (100 mL) was bubbled H_2 through a hydrogenation kit at 30 psi for 1 h. The reaction went to completion, filtered through celite and concentrated to afford 2,4-diisopropylpyridin-3-amine (**Intermediate 194,** 4.0 g, 22.44 mmol, 87 % yield) as a bright yellow oil. 1H NMR (400 MHz, *CHLOROFORM-d*) δ 8.05 (d, *J*=5.2 Hz, 1 H), 6.96 (br d, *J*=5.0 Hz, 1 H), 3.72 (br d, *J*=7.0 Hz, 2 H), 3.09 (s, 1 H), 2.71 - 2.96 (m, 1 H), 1.35 (d, *J*=6.8 Hz, 6 H), 1.27 (d, *J*=6.8 Hz, 6 H). m/z (ESI, +ve ion): 179 (M+H).

Intermediate 195

6,7-Dichloro-1-(2,4-diisopropylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0718] Same as **Intermediate 61A** but using 2,4-diisopropylpyridin-3-amine **(Intermediate 194)** instead of 2-isopropylaniline in Step 1. 1H NMR (400 MHz, *DMSO-d6*) δ 11.94 - 12.45 (m, 1 H), 8.59 (s, 1 H), 8.36 (s, 1 H), 7.39 (d, *J*=5.2 Hz, 1 H), 2.83 - 2.91 (m, 1 H), 2.70 - 2.79 (m, 1 H), 1.05 - 1.13 (m, 6 H), 0.98 (d, *J*=6.6 Hz, 6 H). m/z (ESI, +ve ion): 392.6 (M+H).

Intermediate 196

6-Bromo-2,4-diisopropylpyridin-3-amine



Intermediate 194 **Intermediate 195**

[0719] To a solution of 2,4-diisopropylpyridin-3-amine (0.500 g, 2.80 mmol, **Intermediate 194)** in DMF (10 mL) at 0C was added NBS (0.499 g, 2.80 mmol) in one portion and the resulting mixture was stirred at this temperature for 30 min. The reaction went to

completion.The mixture was washed with water (60 mL), extracted with EtOAc. The combined organics were dried over Na_2SO_4, filtered, concentrated and purified by chromatography on silica gel using 0-40% EtOAc in heptane to afford 6-bromo-2,4-diisopropylpyridin-3-amine (**Intermediate 195,** 0.31 g, 1.205 mmol, 43.0 % yield) as an orange oil. 1H NMR (400 MHz, *CHLOROFORM-d*) δ 7.04 (s, 1 H), 2.93 - 3.09 (m, 1 H), 2.71 - 2.90 (m, 1 H), 1.29 (d, *J*=6.6 Hz, 6 H), 1.25 (d, *J*=6.8 Hz, 6 H). NH2 not observed. m/z (ESI, +ve ion): 258.8 (M+H).

Intermediate 197

4-Ethyl-2-isopropylpyridin-3-amine



Intermediate 190 **Intermediate 197**

[0720] A mixture of 4-chloro-2-isopropylpyridin-3-amine (0.800 g, 4.69 mmol, **Intermediate 190**) ,ethylboronic acid (0.416 g, 5.63 mmol, FSSI), sodium carbonate (2.484 g, 23.44 mmol) and chloro(2-dicyclohexylphosphino-2',4',6'-tri-i-propyl-1,1'-biphenyl)[2-(2-aminoethyl)phenyl] palladium(ii) methyl-t-butylether adduct (0.388 ml, 0.469 mmol) in 1,4-dioxane/water (100/25 mL) was heated at 90 C for 17 h. The reaction was quenched with sat. $NaHCO_3$ and extracted with EtOAc. The combined organics were purified by chromatography on silica gel using 0-80 EtOAc in heptane to afford 4-ethyl-2-isopropylpyridin-3-amine (**Intermediate 197,** 0.093 g, 0.566 mmol, 12.08 % yield) as a green oil. 1H NMR (400 MHz, *DMSO-d6*) δ 7.71 (d, *J*=4.6 Hz, 1 H), 6.79 (d, *J*=4.8 Hz, 1 H), 4.73 (s, 2 H), 2.43 - 2.49 (m, 3 H), 1.15 (d, *J*=6.8 Hz, 9 H). m/z (ESI, +ve ion): 165.2 (M+H).

Intermediate 198

4-Cyclopropyl-2-isopropylpyridin-3-amine



Intermediate 190 **Intermediate 198**

[0721] A mixture of4-chloro-2-isopropylpyridin-3-amine (0.900 g, 5.27 mmol, **Intermediate 190**), cyclopropaneboronic acid (0.680 ml, 7.91 mmol, Small Molecules, Inc.), sodium carbonate (2.236 g, 21.10 mmol) and chloro(2-dicyclohexylphosphino-2',4',6'-tri-i-

propyl-1,1'-biphenyl)[2-(2-aminoethyl)phenyl] palladium(ii) methyl-t-butylether adduct (0.436 ml, 0.527 mmol) in 1,4-dioxane/water (100/25 mL) was heated at 90 C for 24 h. The reaction was washed with sat. $NaHCO_3$, extracted with EtOAc and purified by chromatography on silica gel using 0-5% MeOH in DCM to afford 4-cyclopropyl-2-isopropylpyridin-3-amine (**Intermediate 198,** 0.74 g, 4.20 mmol, 80 % yield) as a yellow oil. ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 7.97 (d, *J*=5.0 Hz, 1 H), 6.77 (d, *J*=4.8 Hz, 1 H), 3.94 (br s, 2 H), 1.63 - 1.70 (m, 1 H), 1.37 - 1.43 (m, 1 H), 1.32 (d, *J*=6.8 Hz, 6 H), 0.98 (dd, *J*=8.4, 1.8 Hz, 2 H), 0.64 (dd, *J*=5.4, 1.5 Hz, 2 H). m/z (ESI, +ve ion): 177 (M+H).

Intermediate 199

6-Chloro-1-(4-cyclopropyl-2-isopropylpyridin-3-yl)-7-(*o*-tolyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione.



[0722] Same as **Example 153** up to step 3 but using 4-cyclopropyl-2-isopropylpyridin-3-amine (**Intermediate 198**) in Step 1 and o-tolylboronic acid is Step 3 . ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 12.22 (br s, 1 H), 8.56 (s, 1 H), 8.38 (d, *J*=5.0 Hz, 1 H), 7.31 (s, 1 H), 7.21 - 7.27 (m, 2 H), 7.06 (d, *J*=7.5 Hz, 1 H), 6.81 (d, *J*=5.2 Hz, 1 H), 2.85 - 3.01 (m, 1 H), 1.94 (s, 3 H), 1.64 - 1.77 (m, 1 H), 1.10 (d, *J*=6.6 Hz, 3 H), 0.93 (d, *J*=6.6 Hz, 3 H), 0.84 - 0.90 (m, 2 H), 0.70 - 0.79 (m, 2 H). m/z (ESI, +ve ion): 447.2 (M+H).

Intermediate 200

6-Chloro-1-(2,4-diisopropylpyridin-3-yl)-7-(o-tolyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0723] Same as **Example 153** up to step 3 using 2,4-di(prop-1-en-2-yl)pyridin-3-amine (**Intermediate 194**) in Step 1 and o-tolylboronic acid is Step 3. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 12.09 - 12.37 (m, 1 H), 8.56 (s, 1 H), 8.48 (d, *J*=5.2 Hz, 1 H), 7.17 - 7.34 (m, 4 H), 6.98 (d, *J*=7.3 Hz, 1 H), 2.88 (quin, *J*=6.6 Hz, 1 H), 2.71 - 2.81 (m, 1 H), 1.92 (s, 3 H), 1.09 (d, *J*=6.6 Hz, 6 H), 0.94 (d, *J*=6.6 Hz, 6 H). m/z (ESI, +ve ion): 448.6 (M+H).

Intermediate 201

6-bromo-2-isopropyl-4-methylpyridin-3-amine



Intermediate 201

[0724] To a 150-mL round-bottomed flask was added 2-isopropyl-4-methylpyridin-3-amine (3.00 g, 19.97 mmol) and n-bromosuccinimide (4.44 g, 24.96 mmol) in N, N-dimethylformamide (24.96 mL). The reaction mixture was stirred and heated at 60 °C for 1.5 h, while under an inert atmosphere. The reaction mixture was poured into water (50 mL), then agitated (sonicator) 5 min. The mixture was diluted with EtOAc and the layers were separated. The aqueous layer was extracted with EtOAc and water. The combined organics were washed with sat. aq. $NaHCO_3$. The organics were collected, dried over $MgSO_4$, filtered and concentrated *in vacuo*, to afford 6-bromo-2-isopropyl-4-methylpyridin-3-amine (**Intermediate 201**, 3.27 g, 14.30 mmol, 72% Yield) as tan oil. *m/z* (ESI, +ve ion): 230.1 (M+H)+. *(Note: The crude oil was carried into the next step of the synthesis, without further purification).

Intermediate 202

tert-Butyl (2*R*,5*S*)-4-(1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate



Intermediate 236

[0725] To a 250-mL round-bottomed flask was added 1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione (2.00 g, 3.97 mmol) and N,N-diisopropylethylamine (0.90 mL, 5.16 mmol) in acetonitrile (19.85 mL). Then phosphorous oxychloride (0.44 mL, 4.76 mmol) was added to the reaction mixture, then the mixture was heated and stirred at 80 °C for 30 min, while under an inert (N_2) atmosphere. The reaction mixture was removed from the heat bath and allowed to cool to rt.

[0726] The reaction mixture was cooled to 0 °C with a wet ice/water bath. Then DIPEA (10 mL) was added dropwise to the reaction mixture. Then a solution of (2r,5s)-1-boc-2,5-dimethylpiperazine (1.27 g, 5.96 mmol) in MeCN (4 mL) was added dropwise to the reaction mixture. The ice bath was removed and the mixture was allowed to warm to rt. More DIPEA (2 mL) and (2r,5s)-1-boc-2,5-dimethylpiperazine (1.27 g, 5.96 mmol) was added to the reaction mixture and stirred an additional 10 min. The reaction mixture was diluted with EtOAc and sat. aq. NH_4Cl. The aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (120 grams), eluting with a gradient of 0-50% EtOAc in CH2CL2, to afford tert-butyl (2R,5S)-4-(1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (1.16 g, 1.66 mmol, 41.9 % yield, **Intermediate 202**) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.44 (d, *J*=4.98 Hz, 1 H) 7.48 - 7.56 (m, 2 H) 7.23 - 7.36 (m, 3 H) 4.83 (br s, 1 H) 4.37 (br s, 1 H) 4.14 (br d, *J*=13.48 Hz, 1 H) 3.86 (br s, 1 H) 3.71 (br d, *J*=13.89 Hz, 1 H) 2.61 - 2.73 (m, 1 H) 1.89 - 1.99 (m, 3 H) 1.45 (s, 9 H) 1.35 (d, *J*=6.63 Hz, 3 H) 1.10 - 1.24 (m, 4 H) 1.04 (dd, *J*=6.74, 2.80 Hz, 3 H) 0.93 (dd, *J*=8.50, 6.84 Hz, 3 H). *m/z* (ESI, +ve ion): 699.0 (M+H)+.

Intermediate 205

6,7-Dichloro-1-(4,6-diisopropylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0727] Step 1. 2,5,6-Trichloro-N-((4,6-diisopropylpyrimidin-5-yl)carbamoyl)nicotinamide. To a 250-mL round-bottomed flask was added 2,5,6-trichloronicotinamide (**Intermediate P**, 4.450 g, 19.74 mmol) and oxalyl chloride (14.80 mL, 29.6 mmol) in tetrahydrofuran (39.5 mL). The flask was fitted with a Findenser, and the mixture was stirred and heated at 80 °C for 1 h. The reaction mixture was concentrated in vacuo. The crude mixture was carried into the next step of the synthesis, without further purification to prevent decomposition.

[0728] To a 250-mL round-bottomed flask was added (2,5,6-trichloronicotinoyl)carbamoyl isocyanate (crude material from previous step) in tetrahydrofuran (39.5 mL). Then a solution of 4,6-diisopropylpyrimidin-5-amine (**Intermediate U**, 3.72 g, 20.72 mmol) in THF (10 mL) was added dropwise into the reaction mixture. The mixture was allowed to stir under an inert (N2) atmosphere for 1.5 h. The reaction mixture was diluted with DCM (50 mL) and heptane (200 mL), then allowed the mixture to stir 30 min. The mixture was filtered and the filtrate (desired material) was collected, then concentrated *in vacuo*. The residue was triturated from a small amount of EtOAc (20 mL) and heptane (200 mL). The precipitate was collected by filtration and the solids were washed with heptane, to afford 2,5,6-trichloro-*N*-((4,6-

diisopropylpyrimidin-5-yl)carbamoyl)nicotinamide (6.310 g, 14.65 mmol, 74.2 % yield) as light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 11.35 - 11.49 (m, 1 H) 9.60 - 9.87 (m, 1 H) 8.91 - 9.13 (m, 1 H) 8.56 - 8.78 (m, 1 H) 3.23 - 3.30 (m, 2 H) 1.17 (d, *J*=6.84 Hz, 12 H). *m/z* (ESI, +ve ion): 430.1 (M+H)+.

[0729] Step 2. 6,7-Dichloro-1-(4,6-diisopropylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (Intermediate 205). To a 250-mL round-bottomed flask was added 2,5,6-trichloro-*N*-((4,6-diisopropylpyrimidin-5-yl)carbamoyl)nicotinamide (3.125 g, 7.26 mmol) in tetrahydrofuran (36.3 mL). The reaction mixture was cooled to 0 °C in a wet ice/water bath. Then potassium bis(trimethylsilyl)amide, 1M solution in tetrahydrofuran (9.07 mL, 9.07 mmol) was added via an addition funnel dropwise into the reaction mixture over 5 min. The ice bath was removed and the reaction mixture was allowed to slowly warm to ambient temperature, while stirred under an inert (N_2) atmosphere for 1 h. More KHMDS (0.5 eq; 6 mL) was added dropwise into the reaction mixture, until SM was mostly consumed **(Note: While adding reagent, carefully monitor reaction as side-product will form if too much KHMDS is added at this step)* The reaction mixture was quenched with sat. aq. NH_4Cl (50 mL), then the mixture was diluted with 3:1 EtOAc/MeOH and brine solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. This afforded 6,7-dichloro-1-(4,6-diisopropylpyrimidin-5-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (2.326 g, 5.90 mmol, 81 % yield, **Intermediate 205**) as a light-yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.33 (br s, 1 H) 9.18 (s, 1 H) 8.60 (s, 1 H) 2.95 (spt, *J*=6.60 Hz, 2 H) 1.09 (d, *J*=6.63 Hz, 6 H) 0.99 (d, *J*=6.63 Hz, 6 H). *m/z* (ESI, +ve ion): 394.1 (M+H)+.

Intermediate 206

(*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-(hydroxymethyl)-6-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, first eluting isomer



[0730] Step 1: *N*-((2-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)carbamoyl)-2,5,6-trichloronicotinamide. To a mixture of 2,5,6-trichloronicotinamide (**Intermediate P**, 0.70 g, 3.12 mmol) in THF (22 mL) was added oxalyl chloride, 2 M solution in DCM (1.71 mL, 3.43 mmol) at rt. The resulting yellow heterogeneous mixture was stirred and heated at 70 °C. After 50 min, the mixture was concentrated and the residue redissolved in THF (8 mL). 2-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-6-isopropylaniline (**Intermediate I-40**, 1.31 g, 3.25 mmol) was added as a solution in THF (7 mL), and the mixture was stirred at rt for 20 min. The mixture was evaporated and the residue was purified by chromatography on silica gel (0-40$ ethyl acetate in heptane) to give *N*-((2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)carbamoyl)-2,5,6-trichloronicotinamide (1.88 g, 2.88 mmol, 98 % yield) as a white solid ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 11.19 (1 H, br s), 9.19 - 9.59 (1 H, m), 8.50 (1 H, s), 7.60 - 7.65 (4 H, m), 7.41 -

7.48 (6 H, m), 7.36 - 7.40 (2 H, m), 7.29 - 7.33 (1 H, m), 4.75 (2 H, s), 3.07 - 3.16 (1 H, m), 1.17 (6 H, d, *J*=6.8 Hz), 1.03 (9 H, s). *m/z* (ESI, +ve ion): 678.2 $(M+Na)^+$.

[0731] Step 2: 1-(2-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. To a cooled mixture of *N*-((2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)carbamoyl)-2,5,6-trichloronicotinamide (1.88 g, 2.87 mmol) in THF (15 mL) at 0 °C was added dropwise KHMDS, 1 M solution in THF (5.74 mL, 5.74 mmol) and the mixture was stirred at rt. After 40 min the mixture was cooled to 0 °C and KHMDS, 1 M solution in THF (1.0 mL, 1.0 mmol) was added. The mixture was stirred at rt 10 min and then quenched with half-satd. ammonium chloride (30 mL) and extracted with EtOAc (2 × 20 mL). The organic phase was washed with water (30 mL) and dried by passage through a Chem Elut extraction cartridge (10 mL, 1219-8007) eluting with ethyl acetate (2 x 10 mL). The filtrate was evaporated and the reside was purified by silica gel chromatography (0-40 % ethyl acetate in heptane) to provide 1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (1.15 g, 65 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 8.35 (s, 1H), 7.48-7.56 (m, 5H), 7.36-7.47 (m, 5H), 7.29-7.35 (m, 4H), 4.42-4.52 (m, 2H), 2.50-2.60 (m, 1H), 1.18 (d, *J*=6.84 Hz, 3H), 1.10 (d, *J*=6.84 Hz, 3H), 0.94 (s, 9H). *m/z* (ESI, +ve ion): 640.2 $(M+Na)^+$.

[0732] Step 3: *tert*-Butyl (*S*)-4-(1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate. A solution of 1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (0.509 g, 0.823 mmol), DIPEA (0.187 mL, 1.07 mmol), and phosphoryl trichloride (0.092 ml, 0.987 mmol) in acetonitrile (5 ml) was stirred at 80 °C for 0.5 h. The reaction mixture cooled to rt, DIPEA (0.43 mL, 2.47 mmol) and *tert*-butyl (*S*)-3-methylpiperazine-1-carboxylate (0.181 g, 0.905 mmol) were added and the reaction was stirred at rt for 10 min. The mixture was poured into cold, satd. $NaHCO_3$ (20 mL) and stirred vigorously for 10 min. The mixture was partitioned between EtOAc (100 mL), and satd. $NaHCO_3$ (75 mL), the organic layer was washed with satd. $NaHCO_3$ (75 mL). The organic extract was dried by passage through a Chem Elut extraction cartridge (10 mL, 1219-8007) eluting with ethyl acetate (2 x 10 mL). The filtrate was evaporated and the reside was purified by silica gel chromatography (10-60 % ethyl acetate in heptane) to provide *tert*-butyl (*S*)-4-(1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-

methylpiperazine-1-carboxylate (0.584 g, 0.729 mmol, 89 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 7.84 (s, 1H), 7.52-7.56 (m, 2H), 7.45-7.51 (m, 3H), 7.28-7.44 (m, 7H), 4.63-5.03 (m, 1H), 4.45-4.57 (m, 2H), 3.73-4.36 (m, 4H), 3.40-3.71 (m, 1H), 2.72-3.32 (m, 2H), 2.40 (dt, *J*=2.90, 6.84 Hz, 1H), 1.51 (d, *J*=1.45 Hz, 9H), 1.31 (br d, *J*=6.63 Hz, 3H), 1.17 (dd, *J*=2.07, 6.63 Hz, 3H), 1.11 (dd, *J*=2.90, 6.84 Hz, 3H), 0.89 (s, 9H). *m/z* (ESI, +ve ion): 800.2 $(M+H)^+$.

[0733] Step 4: *tert*-Butyl (*S*)-4-(1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate. A mixture of *tert*-butyl (*S*)-4-(1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (0.57 g, 0.712 mmol), 2-fluorophenyl boronic acid (Comi-Blocks, Inc., 119 mg, 0.852 mmol), Pd(dppf)Cl_2 (0.037 g, 0.050 mmol) and potassium acetate (0.358 g, 3.64 mmol) was purged with argon followed by the addition of 1,4-dioxane (7.5 mL) and a drop of water. The mixture was heated at 90 °C for 2 h. Fluorophenyl boronic acid (Comi-Blocks, Inc., 76.4 mg) and Pd(dppf)Cl_2 (29.9 mg) was added and the reaction was continued overnight. The mixture was cooled and fluorophenyl boronic acid (Comi-Blocks, Inc., 49.8 mg), Pd(dppf)Cl_2 (13.7 mg) and potassium acetate (98.8 mg) was added, the mixture was degassed by bubbling argon, a drop of water was added and the mixture was heated to 90 C for one hour. The mixture was cooled and filtered through diaomaceous earth. The filtrate was dried by passage though a ChemElute extraction cartridge (5 mL, 1219-0006) eluting with ethyl acetate. The solvent was evaporated and the residue was purified by chromatography on silica gel (20-80% ethyl acetate in heptane to provide *tert*-butyl (*S*)-4-(1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate as a light yellow foam (0.38 g, 0.44 mmol, 61 % yield). ^{1}H NMR (400 MHz, CHLOROFORM-*d*) δ 7.93 (d, *J*=2.28 Hz, 1H), 7.51-7.56 (m, 2H), 7.45-7.50 (m, 3H), 7.38-7.43 (m, 2H), 7.28-7.37 (m, 4H), 7.23-7.26 (m, 1H), 7.16-7.22 (m, 2H), 7.04-7.11 (m, 3H), 4.64-4.99 (m, 1H), 4.54-4.60 (m, 1H), 4.39-4.45 (m, 1H), 3.77-4.37 (m, 4H), 3.44-3.73 (m, 1H), 3.08-3.34 (m, 1H), 2.47-2.57 (m, 1H), 1.52 (s, 9H), 1.37 (d, *J*=6.84 Hz, 3H), 1.18 (br d, *J*=6.84 Hz, 3H), 1.01 (dd, *J*=2.90, 6.84 Hz, 3H), 0.91 (s, 9H). *m/z* (ESI, +ve ion): 860.4 $(M+H)^+$.

[0734] Step 5: (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-(hydroxymethyl)-6-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one. A 250-mL

round-bottomed flask was charged with *tert*-butyl (*S*)-4-(1-(2-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylphenyl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-3-methylpiperazine-1-carboxylate (0.376 g, 0.437 mmol) and DCM (3 mL). TFA (1.5 mL, 19.5 mmol) was added and the mixture was stirred at rt for 25 min. The mixture was concentrated and redissolved in DCM (4 mL). The mxiture was cooled in an ice water bath. DIPEA (0.382 mL, 2.19 mmol) was added followed by acryloyl chloride (1.78 mL, 0.437 mmol). The mixture was stirred at 0 °C for 10 min. The mixture was concentrated and the crude product was purified by chromatography on silica gel (EtOAc in heptanes 30 % - 100 %) to afford 0.251 g of off-white foam.

[0735] The material was dissolved in THF (3 mL) and tetrabutylammonium fluoride (0.481 mL, 0.481 mmol) was added. The mixture was stirred at rt for 3 h. The mixture was concentrated and the crude product was purified by chromatography on silica gel (3:1 EtOAc/EtOH in heptanes 30 % - 100 %) to provide (*S*)-4-(4-acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-(hydroxymethyl)-6-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one as an off-white solid (0.0102 g, 0.018 mmol, 4 %). ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 8.10 (1 H, s), 7.35 - 7.45 (4 H, m), 7.05 - 7.19 (3 H, m), 6.53 - 6.73 (1 H, m), 6.37 - 6.47 (1 H, m), 5.82 (1 H, d, *J*=10.4 Hz), 4.54 - 5.22 (2 H, m), 4.37 - 4.53 (1 H, m), 4.36 (2 H, br s), 3.87 (1 H, br s), 3.63 (1 H, br s), 2.98 - 3.37 (1 H, m), 2.54 (1 H, br s), 2.11 - 2.46 (1 H, m), 1.65 (3 H, br s), 1.22 - 1.28 (1 H, m), 1.18 (3 H, d, *J*=6.8 Hz), 0.96 - 1.03 (3 H, m). *m/z* (ESI, +ve ion): 576.2 $(M+H)^+$.

[0736] **Step 6: (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-(hydroxymethyl)-6-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one, first eluting isomer (Intermediate 206).** The mixture was purified by SFC (Chiralpak IE (150 x 20 mm, 5 μm), 50% MeOH /CO_2, 60 mL/min, 165 bar), to provide (*S*)-4-(4-Acryloyl-2-methylpiperazin-1-yl)-6-chloro-7-(2-fluorophenyl)-1-(2-(hydroxymethyl)-6-isopropylphenyl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 206**, as the first-eluting isomer. ^{1}H NMR (400 MHz, METHANOL-d_4) δ 8.43 (br d, *J*=4.56 Hz, 1H), 7.34-7.50 (m, 4H), 7.10-7.25 (m, 3H), 6.84 (dt, *J*=10.68, 16.53 Hz, 1H), 6.32 (br d, *J*=16.59 Hz, 1H), 5.83 (dd, *J*=1.87, 10.57 Hz, 1H), 5.05 (br s, 1H), 4.42-4.62 (m, 2H), 4.34-4.41 (m, 1H), 4.24-4.30 (m, 1H), 4.04-4.23 (m, 1H), 3.58-3.90 (m, 2H), 3.33-3.43 (m, 1H), 3.17-3.29 (m, 1H), 2.51-2.66 (m, 1H), 1.49 (br d, *J*=6.63 Hz, 3H), 1.15 (d, *J*=6.84 Hz, 3H), 1.00 (d, *J*=6.84 Hz, 3H). *m/z* (ESI, +ve ion): 576.2 $(M+H)^+$. The second eluting isomer was also characterized. ^{1}H NMR (400 MHz, METHANOL-d_4) δ 8.43 (br d, *J*=4.35 Hz, 1H), 7.34-7.50 (m, 4H), 7.10-7.26

(m, 3H), 6.77-6.92 (m, 1H), 6.31 (br dd, *J*=3.01, 16.90 Hz, 1H), 5.83 (dd, *J*=1.87, 10.57 Hz, 1H), 5.09 (br s, 1H), 4.55 (br d, *J*=5.81 Hz, 1H), 4.45 (br d, *J*=4.56 Hz, 1H), 4.34-4.41 (m, 1H), 4.23-4.32 (m, 1H), 4.03-4.21 (m, 1H), 3.82-3.95 (m, 1H), 3.55-3.79 (m, 1H), 3.34-3.44 (m, 1H), 3.16-3.28 (m, 1H), 2.51-2.66 (m, 1H), 1.48 (br d, *J*=6.43 Hz, 3H), 1.16 (d, *J*=6.84 Hz, 3H), 1.00 (d, *J*=6.84 Hz, 3H). *m/z* (ESI, +ve ion): 576.2 $(M+H)^+$.

Intermediate 207

4-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-2-isopropylpyridin-3-amine



[0737] Step 1. Methyl 3-amino-2-isopropylisonicotinate. A 250-mL round-bottomed flask was charged with 3-amino-2-chloro-4-(methoxycarbonyl)pyridine (3.09 g, 16.6 mmol, Combi-Block, Inc.), methanesulfonato[9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene][2'-amino-1,1'-biphenyl]palladium(ii) dichloromethane adduct (Xantophos G3) (0.862 g, 0.835 mmol,) and THF (50 mL) under argon. 2-Propylzinc bromide (0.5 M in THF, 43.1 mL, 21.5 mmol, Aldrich) was added slowly via addition funnel. The addition funnel was swapped with a Findensor condenser and the mixture was stirred at 50 °C for 20 min. The mixture was cooled to rt then placed in an ice water bath. Water (~80 mL) and EtOAc (~50 mL) were added and the mixture was filtered through fine frit glass filter funnel (eluent:EtOAc) to remove insoluble emulsions. The phases were separated and the aqueous phase was extracted with EtOAc (2 x 50 mL). The combined organic phases were dried over sodium sulfate, filtered and concentrated. A chromatographic purification (silica gel, 5-60% EtOAc/heptane) obtained methyl 3-amino-2-isopropylisonicotinate (2.31 g, 11.9 mmol, 72% yield) as as a yellow oil. ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 7.94 (1 H, d, *J*=5.2 Hz), 7.50 (1 H, d, *J*=5.2 Hz), 5.87 (2 H, br s), 3.91 (3 H, s), 3.08 (1 H, dt, *J*=13.5, 6.7 Hz), 1.32 (6 H, d, *J*=6.8 Hz). m/z (ESI, +ve ion): 195.2 [M+H].

[0738] Step 2. 4-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-2-isopropylpyridin-3-amine. A 100-mL round-bottomed flask was charged with methyl 3-amino-2-isopropylisonicotinate (2.20 g, 11.3 mmol) and THF (20 mL). Lithium borohydride (2 M in THF, 8.49 mL, 17.0 mmol, Aldrich) was added dropwise. The mixture was stirred at rt for 15 min. Then methanol (3.67 mL, 91 mmol) was added slowly and the mixture was stirred at rt for 18 h. Saturated

NH_4Cl (~20 mL) was added slowly. The aqueous phase was extracted with EtOAc (2 x 20 mL). The combined organic phases were dried over Na_2SO_4, filtered and concentrated. This material was dissolved in DCM (20 mL). Benzene, 1,1'-[chloro(1,1-dimethylethyl)silylene]bis- (3.83 mL, 14.7 mmol), DIPEA (6.92 mL, 39.6 mmol) and DMAP (0.0707 g, 0.579 mmol) were added and the mixture was stirred at rt for 20 h. The mixture was washed with water. The organic phase was dried over sodium sulfate, filtered and concentrated *in vacuo*. A chromatographic purification (silica gel, 0-40% EtOAc/heptane) afforded 4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-2-isopropylpyridin-3-amine (**Intermediate 207**, 1.31 g, 3.25 mmol, 29% yield). ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 7.97 (1 H, d, *J*=4.8 Hz), 7.68 (4 H, dd, *J*=8.1, 1.5 Hz), 7.37 - 7.43 (6 H, m), 6.73 (1 H, d, *J*=4.8 Hz), 4.68 (2 H, s), 4.19 (2 H, br s), 3.04 - 3.14 (1 H, m), 1.34 (6 H, d, *J*=6.6 Hz), 1.07 (9 H, s). m/z (ESI, +ve ion): 405.2 [M+H].

Intermediate 208

4-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-6-isopropylpyrimidin-5-amine



[0739] **Step 1. 4-Chloro-6-isopropylpyrimidin-5-amine**. A 250-mL round-bottomed flask was charged with 4,6-dichloro-5-aminopyrimidine (3.01 ml, 18.4 mmol) and THF (20 mL). XantPhos Pd G3 (0.957 g, 0.926 mmol) was added and 2-propylzinc bromide (1.0 M in THF, 19.3 ml, 19.3 mmol, Tieki Metal) was added dropwise via addition funnel. The mixture was stirred at rt for 20 h. Additional 2-propylzinc bromide (10 mL) was added slowly and the mixture was stirred at rt for 3 h. The reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (3 × 75 mL). The organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. A

chromatographic purification (silica gel, 0–40% EtOAc/heptane) afforded 4-chloro-6-isopropylpyrimidin-5-amine (0.912 g, 5.31 mmol, 29 % yield) as a yellow solid. ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 8.39 (1 H, s), 4.10 (2 H, br s), 3.02 (1 H, spt, *J*=6.7 Hz), 1.31 (6 H, d, *J*=6.8 Hz). m/z (ESI, +ve ion): 172.1 [M+1].

[0740] Step 2. Methyl 5-amino-6-isopropylpyrimidine-4-carboxylate. 4-Chloro-6-isopropylpyrimidin-5-amine (0.75 g, 4.4 mmol), TEA (1.83 mL, 13.1 mmol), $PddppfCl_2$ (0.327 g, 0.447 mmol, Strem) were suspended in MeOH (15 mL). The suspention was divided into 4 reaction glass tube and stirred at 50 °C under 20 psi CO for 18 h using Endeavor reactor (Biotage). The reaction mixture was combined and concentrated. A chromatographic purification (silica gel, 20-80% EtOAc/heptane) afforded methyl 5-amino-6-isopropylpyrimidine-4-carboxylate (0.871, g 4.46 mmol, 100% yield) as light yellow solid. ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 8.64 (1 H, s), 5.87 (2 H, br s), 4.00 (3 H, s), 3.10 (1 H, dt, *J*=13.4, 6.7 Hz), 1.34 (6 H, d, *J*=6.8 Hz). m/z (ESI, +ve ion): 196.1 [M+H].

[0741] Step 3. 4-(((*tert*-Butyldiphenylsilyl)oxy)methyl)-6-isopropylpyrimidin-5-amine. To a stirred solution of methyl 5-amino-6-isopropylpyrimidine-4-carboxylate (0.870 g, 4.46 mmol) in THF (10 mL) was added a solution of lithium borohydride (2 M in THF, 3.34 mL, 6.68 mmol, Aldrich) dropwise. After the addition was completed, the reaction mixture was stirred for 15 min. Then methanol (1.44 mL, 35.7 mmol) was added slowly. The reaction mixture was then stirred at RT for 18 h. 10 mL of saturated NH_4Cl was added slowly. The aqueous phase was extracted with EtOAc (3 x 10 mL). The combined organic phases were dried over Na_2SO_4, filtered and concentrated. This material was dissolved into DCM (10 mL). *tert*-butylchlorodiphenylsilane (1.51 mL, 5.79 mmol), DIPEA (2.72 mL, 15.6 mmol) and DMAP (0.031 g, 0.25 mmol) were added and the mixture was stirred at rt for 20 h. The reaction mixture was washed with water and dried over Na_2SO_4, filtered and concentrated. A chromatographic puridication (silica gel, 0-40% EtOAc/heptane) afforded 4-(((*tert*-butyldiphenylsilyl)oxy)methyl)-6-isopropylpyrimidin-5-amine (**Intermediate 208**, 1.12 g, 2.76 mmol, 62% yield) as a clear oil. ^{1}H NMR (400 MHz, *CHLOROFORM-d*) δ 8.47 (1 H, s), 7.64 - 7.69 (4 H, m), 7.42 - 7.47 (2 H, m), 7.36 - 7.41 (4 H, m), 4.91 (2 H, s), 4.55 (2 H, br s), 3.06 (1 H, quin, *J*=6.7 Hz), 1.32 (6 H, d, *J*=6.8 Hz), 1.09 (9 H, s). m/z (ESI, +ve ion): 406.1 [M+H].

Intermediate 209

5-Methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one



[0742] To a pressure vial was added 4-bromo-5-methylindolin-2-one (0.5 g, 2.21 mmol, ChemShuttle, Hayward, CA), bis(pinacolato)diboron (1.12 g, 4.42 mmol, Matrix Scientific, Columbia, SC), [1,1'-bis(diphenylphosphino)ferrocene]-dichloropalladium(ii), complex with dichloromethane (0.181 g, 0.221 mmol, Sigma-Aldrich, St. Louis, MO), potassium acetate (0.651 g, 6.64 mmol) and 1,4-dioxane (10 mL). The vial was purged with N_2 for 3 min, sealed and then heated at 100°C for 18 h. LCMS showed convesion of the starting material was complete. The reaction mixture was partitioned between EtOAc and water. The separated organic layer was washed with brine, dried over Na2SO4 and then filtered. The filtrate was concentrated and purified by silica gel chromatography (0-40% (3:1 EtOAc:EtOH)/heptane) to afford 5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indolin-2-one (**Intermediate 209**, 0.365 g, 1.336 mmol, 60.4 % yield) as an off-white solid. 1H NMR (400 MHz, DMSO-d_6) δ 1.36 (s, 12 H) 2.41 (s, 3 H) 3.51 (s, 2 H) 6.80 (d, *J*=7.67 Hz, 1 H) 7.02 (d, *J*=7.88 Hz, 1 H) 10.26 (s, 1 H); m/z (ESI, +ve ion): 274.1 (M+H).

Intermediate 210

3-Amino-4-isopropyl-*N*,*N*-dimethylpicolinamide



[0743] **Step 1: 3-Amino-4-chloro-*N*,*N*-dimethylpicolinamide.** To a 20 mL microwave vial was added palladium (II) acetate (0.070 g, 0.312 mmol, Strem Chemicals, Newburyport, MA), 9,9-dimethyl-4,5-bis(bis[3,5-dimethyl-4-methoxyphenyl]phosphino)xanthene (0.355 g,

0.613 mmol, Strem Chemicals, Newburyport, MA) and 4-(dimethylamino) pyridine (1.5 g, 12.3 mmol, Aldrich, St. Louis, MO). The vial was purged with Ar followed by adding 2,4-dichloro-3-aminopyridine (1.0 g, 6.13 mmol, Combi-Blocks, San Diego, CA) in toluene (10 mL) and dimethylamine solution, 2 M in THF (6.13 mL, 12.3 mmol, Aldrich, St. Louis, MO). The vial was capped and underwent microwave heating at 100 °C for 30 min. The reaction was partitioned between EtOAc (60 mL) and water (30 mL). The aqueous layer was extracted with EtOAc (50 mL). The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. The crude product was purified by silica gel chromatography (eluent: 5-40% acetone/heptane) to provide 3-amino-4-chloro-*N*,*N*-dimethylpicolinamide as a light brown oil (0.620 g). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 7.86 (d, *J*=4.98 Hz, 1H), 7.23 (d, *J*=4.98 Hz, 1H), 5.20 (br s, 2H), 3.14 (s, 3H), 3.13 (s, 3H). *m/z* (ESI) M+H: 200.2.

[0744] Step 2: 3-Amino-*N*,*N*-dimethyl-4-(prop-1-en-2-yl)picolinamide. To a 150 mL round bottom flask was added 3-amino-4-chloro-*N*,*N*-dimethylpicolinamide (1.2 g, 6.01 mmol), chloro(2-dicyclohexylphosphino-2',4',6'-tri-i-propyl-1,1'-biphenyl)[2-(2-aminoethyl)phenyl] palladium(II) methyl-*t*-butylether adduct (0.350 g, 0.423 mmol, Strem Chemicals, Newburyport, MA), sodium carbonate (1.6 g, 15.1 mmol, Aldrich, St. Louis, MO), 1,4-dioxane (40 mL), water (10 mL), and 2-isopropenylboronic acid, pincol ester (4.04 g, 24.04 mmol, Aurum Pharmatech, Franklin Park, NJ). The resulting mixture was heated at 95 °C under N_2 for 2 h. The reaction was cooled to rt and partitioned between EtOAc (60 mL) and water (20 mL). The aqueous layer was extracted with EtOAc (30 mL). The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. The crude product was purified by silica gel chromatography (eluent: 5-40% acetone/heptane) to provide 3-amino-*N*,*N*-dimethyl-4-(prop-1-en-2-yl)picolinamide as a light brown oil (1.2 g). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 7.93 (d, *J*=4.77 Hz, 1H), 6.95 (d, *J*=4.77 Hz, 1H), 5.40 (t, *J*=1.45 Hz, 1H), 5.15 (s, 1H), 4.88 (br s, 2H), 3.14 (d, *J*=4.35 Hz, 6H), 2.07 (s, 3H). m/z (ESI) M+H: 206.2

[0745] Step 3: 3-Amino-4-isopropyl-*N*,*N*-dimethylpicolinamide. A 250 mL hydrogenation apparatus was pre-treated with N_2. Palladium 10 wt. % on activated carbon (0.311 g, 0.292 mmol, Aldrich, St. Louis, MO) was added. A solution of 3-amino-*N*,*N*-dimethyl-4-(prop-1-en-2-yl)picolinamide (1.2 g, 5.85 mmol) in ethanol (50 mL) was then added. The cylinder was sealed, purged with H_2 then vented and this procedure was repeated. Fresh H_2 was then introduced to 30 psi. The reaction mixture was stirred under H_2 for 2 h. Reaction mixture was filtered through a plug of celite, and rinsed with MeOH. The resulting filtrate was concentrated to provide 3-amino-4-isopropyl-*N*,*N*-dimethylpicolinamide as brown

solid (1.16 g, **Intermediate 210**). ^{1}H NMR (400 MHz, CHLOROFORM-d) δ 7.96 (d, *J*=4.98 Hz, 1H), 7.06 (d, *J*=4.98 Hz, 1H), 4.73 (br s, 2H), 3.12 (br d, *J*=18.24 Hz, 6H), 2.90 (s, 1H), 1.28 (d, *J*=6.84 Hz, 6H). *m/z* (ESI) M+H: 208.3.

Intermediate 211

***tert*-Butyl (2*R*,5*S*)-4-(1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**



[0746] Step 1: 6-Chloro-N^4,N^4-bis(4-methoxybenzyl)pyrimidine-4,5-diamine. An amber solution of 4,6-dichloropyrimidin-5-amine (1.7 g, 10.4 mmol; CombiBlocks, Inc., San Diego, CA), bis(4-methoxybenzyl)amine (2.67 g, 10.4 mmol; Sigma-Aldrich, St. Louis, MO), and Et_3N (2.89 mL, 20.7 mmol) in isopropanol (10 mL) was stirred in the microwave at 120 °C for 1 h then at 160 °C for 1 h. The reaction mixture was diluted with EtOAc (200 mL), added to a separatory funnel, and washed with water (3 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% EtOAc/heptane) to give 6-chloro- N^4,N^4-bis(4-methoxybenzyl)pyrimidine-4,5-diamine (2.11 g, 5.48 mmol, 53% yield) as an amber oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.12 (s, 1 H) 7.14 (br d, *J* = 8.1 Hz, 4 H) 6.85 (br d, *J* = 8.5 Hz, 4 H) 4.47 (br s, 4 H) 3.91 (br s, 2 H) 3.80 (s, 6 H). MS (ESI, +ve) *m/z*: 385.0 $(M + 1)^+$.

[0747] Step 2: 6-Isopropyl-N^4,N^4-bis(4-methoxybenzyl)pyrimidine-4,5-diamine. A solution of 6-chloro- N^4,N^4-bis(4-methoxybenzyl)pyrimidine-4,5-diamine (2.11 g, 5.48 mmol), isopropylzinc bromide in THF (11.5 mL, 5.75 mmol), and XantPhos Pd G3 (0.283 g, 0.274 mmol) in 1,4-dioxane (5 mL) was stirred at RT for 2 h. More isopropylzinc bromide in THF

(11.5 mL, 5.75 mmol) was added, and the reddish solution was stirred at 50 °C for 16 h. The reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% EtOAc/heptane) to give 6-isopropyl- N^4,N^4-bis(4-methoxybenzyl)pyrimidine-4,5-diamine (1.24 g, 3.15 mmol, 58% yield) as a brown oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.39 (s, 1 H) 7.13 (d, J = 8.5 Hz, 4 H) 6.82 (d, J = 8.7 Hz, 4 H) 4.32 (s, 4 H) 3.79 (s, 6 H) 3.67 (s, 2 H) 3.01 (spt, J = 6.7 Hz, 1 H) 1.30 (d, J = 6.8 Hz, 6 H). MS (ESI, +ve) *m/z*: 393.0 $(M + 1)^+$.

[0748] Step 3: [1-(4-(bis(4-Methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. Oxalyl chloride (2 M in DCM, 1.73 mL, 3.46 mmol) was added to a mixture of 2,5,6-trichloronicotinamide (0.710 g, 3.15 mmol, **Intermediate P**) in tetrahydrofuran (16 mL); the mixture was stirred at 65 °C for 30 min. The reaction mixture was poured into a flask containing 6-isopropyl-N4,N4-bis(4-methoxybenzyl)pyrimidine-4,5-diamine (1.236 g, 3.15 mmol), and the reaction mixture was stirred at RT for 15 min. The reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo* to give N-((4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)carbamoyl)-2,5,6-trichloronicotinamide (2.30 g, 3.57 mmol, >99% yield) as an amber foam. MS (ESI, +ve) *m/z*: 642.7/644.8 $(M + 1)^+$.

[0749] A solution of *N*-((4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)carbamoyl)-2,5,6-trichloronicotinamide (2.03 g, 3.15 mmol) and potassium bis(trimethylsilyl)amide (1 M in THF, 6.30 mL, 6.30 mmol) in tetrahydrofuran (16 mL) was stirred at RT for 2 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–70% EtOAc/heptane) to give 1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (1.22 g, 2.00 mmol, 64% yield) as a light yellow amorphous solid and a mixture of atopisomers (first eluting product with desired mass). ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.76 (s, 1 H) 8.16 (br. s., 1 H) 8.13 (s, 1 H) 6.93 - 6.97 (m, 4 H) 6.69 - 6.74 (m, 4 H) 4.83 (d, J = 16.8 Hz, 2 H) 4.46 (d, J

= 16.8 Hz, 2 H) 3.76 (s, 6 H) 2.35 - 2.47 (m, 1 H) 1.18 (d, *J* = 6.6 Hz, 3 H) 0.99 (d, *J* = 6.6 Hz, 3 H). MS (ESI, +ve) *m/z*: 606.8 $(M + 1)^+$.

[0750] Step 4: *tert*-Butyl (2*R*,5*S*)-4-(1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of 1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloropyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (1.22 g, 2.00 mmol), phosphoryl trichloride (0.224 mL, 2.40 mmol), and DIPEA (1.05 mL, 6.01 mmol) in acetonitrile (5 mL) was stirred at 80 °C for 30 min. The reaction mixture was removed from the reaction block, and a solution of *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (0.515 g, 2.40 mmol; eNovation Chemicals LLC, Bridgewater, NJ) in acetonitrile (5 mL) was added; the reaction mixture was stirred at RT for 1 h. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was loaded onto the column and was purified via automated flash chromatography (silica gel, 0–90% [(3:1) EtOAc/EtOH]/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(1-(4-(bis(4-methoxybenzyl)amino)-6-isopropylpyrimidin-5-yl)-6,7-dichloro-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 211**, 1.15 g, 1.43 mmol, 71% yield) as a yellow solid and a mixture of atropisomers. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.72 (s, 1 H) 7.61 - 7.71 (m, 1 H) 6.94 (t, *J* = 8.4 Hz, 4 H) 6.62 - 6.73 (m, 4 H) 5.05 (br d, *J* = 16.8 Hz, 1 H) 4.88 (br d, *J* = 16.6 Hz, 1 H) 4.47 - 4.69 (m, 1 H) 4.30 (dd, *J* = 16.5, 12.1 Hz, 2 H) 3.75 (s, 3 H) 3.74 (s, 3 H) 3.25 - 3.94 (m, 5 H) 2.25 - 2.42 (m, 1 H) 1.50 (s, 9 H) 1.26 (t, *J* = 7.2 Hz, 3 H) 1.16 - 1.24 (m, 6 H) 0.85 - 1.00 (m, 3 H). MS (ESI, +ve) *m/z*: 802.7 $(M + 1)^+$.

Intermediate 212

6-Chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0751] **Step 1: *N*-Benzyl-3,5-dichloropyridazin-4-amine**. A solution of 3,4,5-trichloropyridazine (2.04 g, 11.1 mmol; CombiBlocks, Inc., San Diego, CA) and benzylamine (3.65 mL, 33.4 mmol; Sigma-Aldrich, St. Louis, MO) in isopropanol (10 mL) was heated in the microwave at 120 °C for 20 min when a solid had formed (run four times) and two isomers with product mass were observed. The four reaction mixtures were cooled to RT, suspended in heptane, and filtered to give an off-white solid (mostly the undesired isomer). The solute was concentrated to give a reddish oil; the resulting oil (in DCM/MeOH) was loaded onto the column and was purified via automated flash chromatography (silica gel, 0–40% EtOAc/heptane) to give *N*-benzyl-3,5-dichloropyridazin-4-amine (3.92 g, 15.4 mmol, 35% yield) as a reddish oil which became a yellowish waxy solid overnight. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.63 (s, 1 H) 7.29 - 7.43 (m, 5 H) 5.22 - 5.29 (m, 1 H) 4.93 (d, J = 6.0 Hz, 2 H). MS (ESI, +ve) *m/z*: 254.0 $(M + 1)^+$.

[0752] **Step 2: *N*-Benzyl-3,5-di(prop-1-en-2-yl)pyridazin-4-amine**. A mixture (separated into 4 equal reactions, ~1 g each) of *N*-benzyl-3,5-dichloropyridazin-4-amine (3.92 g, 15.4 mmol), 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane (11.6 mL, 61.8 mmol), PdCl2(dppf) (0.791 g, 1.081 mmol), and sodium carbonate (2 M aq., 30.9 mL, 61.8 mmol) in 1,4-dioxane (39 mL) was sparged with nitrogen then was heated in the microwave at 160 °C for 1 h. Each reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel,

and washed with saturated, aqueous sodium bicarbonate (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4. The combined crude products were adsorbed onto silica and were purified via automated flash chromatography (silica gel, 0–80% [(3:1) EtOAc/EtOH]/heptane) to give *N*-benzyl-3,5-di(prop-1-en-2-yl)pyridazin-4-amine (2.24 g, 8.44 mmol, 54.7 % yield) as a brown oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.46 (s, 1 H) 7.27 - 7.37 (m, 3 H) 7.11 - 7.19 (m, 2 H) 5.51 (t, *J*=1.6 Hz, 1 H) 5.35 (t, *J*=1.6 Hz, 1 H) 5.23 (s, 1 H) 5.15 (s, 1 H) 4.90 (br t, *J* = 5.6 Hz, 1 H) 4.39 (d, *J* = 6.2 Hz, 2 H) 2.16 (s, 3 H) 2.05 (s, 3 H). MS (ESI, +ve) *m/z*: 266.1 $(M + 1)^+$.

[0753] **Step 3: 3,5-Diisopropylpyridazin-4-amine**. Palladium hydroxide on carbon, wet (1.19 g, 0.844 mmol) was added to a solution of *N*-benzyl-3,5-di(prop-1-en-2-yl)pyridazin-4-amine (2.24 g, 8.44 mmol) in ethanol (42 mL) in a hydrogenation vial at RT under a blanket of nitrogen. The reaction mixture was stirred at RT under hydrogen (50 psig) for 3 h (pressure @ 27 psig). The reaction mixture was stirred again at RT under hydrogen (50 psig) for 2.5 d (pressure @ 30 psig). The mixture was filtered through a pad of Celite, covered with sand, washed with MeOH (2 x 100 mL), and concentrated to give an off-white solid.

[0754] Palladium hydroxide on carbon, wet (1.19 g, 0.844 mmol) was added to a solution of the off-white solid in EtOH (42 mL) and was stirred under hydrogen (42 psig) at RT for 20 h (pressure @ 36 psig). The reaction mixture was charged with more palladium hydroxide on carbon, wet (1.185 g, 0.844 mmol) and was stirred under hydrogen (50 psig) for 5 d (pressure @ 42 psig). The mixture was filtered through a pad of Celite, covered with sand, washed with MeOH (3 x 100 mL), and concentrated to give 3,5-diisopropylpyridazin-4-amine (1.407 g, 7.85 mmol, 93 % yield) as an off-white solid. The crude product in DCM/MeOH was loaded onto the column and was purified via automated flash chromatography (silica gel, 0–100% MeOH/DCM) to give 3,5-diisopropylpyridazin-4-amine (0.923 g, 5.15 mmol, 61% yield) as a light yellow waxy solid. The solid was dissolved in THF (5 mL) and concentrated (2×) to remove any excess methanol. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.62 (s, 1 H) 4.25 (br s, 2 H) 3.08 (spt, *J* = 6.8 Hz, 1 H) 2.85 (spt, *J* = 6.9 Hz, 1 H) 1.42 (d, *J* = 6.8 Hz, 6 H) 1.30 (d, *J* = 6.8 Hz, 6 H).

[0755] **Step 4: 6-Chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione**. A solution of 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (1.07 g, 3.75 mmol, **Intermediate 99B**) and oxalyl chloride (2 M in DCM, 2.07 mL, 4.13 mmol) in tetrahydrofuran (13 mL) was stirred at 65 °C for 1 h. The reaction mixture was removed from the heating block, and 3,5-diisopropylpyridazin-4-amine

(0.673 g, 3.75 mmol) in THF (10 mL) was added; the solution became a white slurry, then DIEA (1.31 mL, 7.51 mmol) was added; the reaction mixture was stirred at RT for 5 min. The reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–100% EtOAc/heptane) to give 2,5-dichloro-*N*-((3,5-diisopropylpyridazin-4-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide (1.09 g, 2.21 mmol, 59% yield) as a colorless oil. MS (ESI, +ve) *m/z*: 490.0 $(M + 1)^+$.

[0756] A solution of 2,5-dichloro-*N*-((3,5-diisopropylpyridazin-4-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide (0.985 g, 2.01 mmol) and potassium bis(trimethylsilyl)amide (1 M in THF, 4.02 mL, 4.02 mmol) in tetrahydrofuran (10 mL) (light yellow solid formed) was stirred at 0 °C for 5 min then was allowed to warm to RT and stir for 15 min. The reaction mixture was diluted with EtOAc (200 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo* . The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% [(3:1) EtOAc/EtOH]/heptane) to give impure 6-chloro-1-(3,5-diisopropylpyridazin-4-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione **(Intermediate 212,** 878 mg, 1.93 mmol, 96% yield) as a colorless oil. MS (ESI, +ve) *m/z*: 454.1 $(M + 1)^+$.

Intermediate 213

1-((1*R*,5*S*)-3,8-Diazabicyclo[3.2.1]octan-8-yl)prop-2-en-1-one



Intermediate 213

[0757] **1-((1*R*,5*S*)-3,8-Diazabicyclo[3.2.1]octan-8-yl)prop-2-en-1-one**. A solution of 3-Boc-3,8-diazabicyclo[3.2.1]octane (0.50 g, 2.36 mmol; CombiBlocks, Inc., San Diego, CA), acryloyl chloride (0.5 M in DCM, 5.65 mL, 2.83 mmol), and DIEA (1.23 mL, 7.07 mmol) in dichloromethane (24 mL) was stirred at RT for 30 min. The reaction mixture was washed with saturated, aqueous sodium bicarbonate (75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica

and was purified via automated flash chromatography (silica gel, 0–100% EtOAc/heptane) to give *tert*-butyl (1*R*,5*S*)-8-acryloyl-3,8-diazabicyclo[3.2.1]octane-3-carboxylate (535 mg, 2.01 mmol, 85% yield) as a light yellow oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 6.43 - 6.52 (m, 1 H) 6.36 - 6.42 (m, 1 H) 5.73 (dd, *J* = 10.0, 2.5 Hz, 1 H) 4.69 - 4.83 (m, 1 H) 4.27 (br d, *J* = 16.0 Hz, 1 H) 3.68 - 4.03 (m, 2 H) 2.84 - 3.21 (m, 2 H) 1.71 - 2.01 (m, 4 H) 1.46 (s, 9 H). MS (ESI, +ve) *m/z*: 289.0 $(M + Na)^+$.

[0758] A solution of *tert*-butyl (1*R*,5*S*)-8-acryloyl-3,8-diazabicyclo[3.2.1]octane-3-carboxylate (535 mg, 2.01 mmol) and TFA (2.72 mL, 35.3 mmol) in dichloromethane (24 mL) was stirred at RT for 15 min. The reaction mixture was concentrated to give 1-((1*R*,5*S*)-3,8-diazabicyclo[3.2.1]octan-8-yl)prop-2-en-1-one 2,2,2-trifluoroacetate (**Intermediate 213,** 0.921 g, 3.29 mmol, >99% yield) as an amber oil which was used as is. ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.57 - 9.12 (m, 2 H) 6.48 (s, 1 H) 6.46 (d, *J* = 4.1 Hz, 1 H) 5.92 - 5.97 (m, 1 H) 4.98 (br s, 1 H) 4.55 (br s, 1 H) 3.17 - 3.55 (m, 4 H) 2.13 - 2.47 (m, 4 H). MS (ESI, +ve) *m/z*: 167.0 $(M + 1)^+$.

Intermediate 214

1-(3,9-Diazabicyclo[3.3.1]nonan-3-yl)prop-2-en-1-one



Intermediate 214

[0759] **1-(3,9-Diazabicyclo[3.3.1]nonan-3-yl)prop-2-en-1-one**. A solution of *tert*-butyl 3,9-diazabicyclo[3.3.1]nonane-9-carboxylate (0.581 g, 2.57 mmol; Enamine LLC, Monmouth Jct., NJ), acryloyl chloride (0.250 mL, 3.08 mmol), and DIPEA (1.34 mL, 7.70 mmol) in dichloromethane (13 mL) was stirred at RT for 30 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–90% EtOAc/heptane) to give *tert*-butyl 3-acryloyl-3,9-diazabicyclo[3.3.1]nonane-9-carboxylate (587 mg, 2.09 mmol, 82% yield) as a colorless oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 6.50 - 6.62 (m, 1 H) 6.30 (dd, *J* = 16.8, 1.9 Hz, 1 H) 5.71 (dd, *J* = 10.6, 1.9 Hz, 1 H) 4.57 - 4.65 (m, 1 H) 4.29 (br s, 1 H) 4.17 (br s, 1 H) 3.88 -

3.96 (m, 1 H) 3.39 - 3.48 (m, 1 H) 2.92 - 3.06 (m, 1 H) 1.66 - 1.99 (m, 5 H) 1.49 - 1.56 (m, 1 H) 1.48 (s, 9 H). MS (ESI, +ve) *m/z*: 303.0 $(M + Na)^+$.

[0760] A solution of the resulting oil in 2,2,2-trifluoroacetic acid (29.3 g, 257 mmol) was stirred at RT for 15 min. The reaction mixture was concentrated to give 1-(3,9-diazabicyclo[3.3.1]nonan-3-yl)prop-2-en-1-one 2,2,2-trifluoroacetate (**Intermediate 214,** 818 mg, 2.78 mmol, >99% yield) as a colorless oil which was used as is. MS (ESI, +ve) *m/z*: 181.1 $(M + 1)^+$.

Intermediates 204 & 215

***tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (two atropisomers)**



[0761] **Step 1: 2-Chloro-4-isopropyl-6-methylpyrimidin-5-amine**. To a 1-L three-necked RBF was added 2,4-dichloro-6-methylpyrimidin-5-amine (16.1 g, 90 mmol; CombiBlocks, Inc., San Diego, CA) and [(4,5-bis(diphenylphosphino)-9,9-dimethylxanthene)-2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate (4.28 g, 4.51 mmol) in tetrahydrofuran (180 mL) under argon. 2-Propylzinc bromide (0.5 M in tetrahydrofuran, 198 mL, 99 mmol; Sigma-Aldrich, St. Louis, MO) was added over 5 min (no exotherm observed). The reaction mixture

was then stirred at 23 °C for 90 min. To the reaction mixture was added ice (~15 g) and Celite (~50 g) with stirring, and the mixture was filtered through a fine fritted glass filter funnel (eluent: EtOAc) to remove insoluble emulsions. The combined organic phases were dried over sodium sulfate, filtered, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–70% EtOAc/heptane) to give 2-chloro-4-isopropyl-6-methylpyrimidin-5-amine (8.83 g, 47.6 mmol, 53% yield) as yellow oil. ^{1}H NMR (400 MHz, DMSO-d_6) δ 1.12 (d, J = 6.63 Hz, 6 H) 2.27 (s, 3 H) 3.20 (spt, J = 6.70 Hz, 1 H) 5.25 (s, 2 H). MS (ESI, +ve) m/z: 186.0 $(M + 1)^+$.

[0762] Step 2: 6-Chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A solution of 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (9.41 g, 33.0 mmol, **Intermediate 99B**) and oxalyl chloride (2 M in DCM, 18.2 mL, 36.3 mmol) in tetrahydrofuran (100 mL) was stirred under an air condensor and drying tube at 65 °C for 1 h. The solution was cooled to RT; 2-chloro-4-isopropyl-6-methylpyrimidin-5-amine (6.13 g, 33.0 mmol) in THF (5 mL) at RT was added, and the yellow solution was stirred at RT for 30 min. The reaction mixture was concentrated, diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (3 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo* to give impure product. The crude product in DCM was loaded onto the column and was purified via automated flash chromatography (silica gel, 0–40% EtOAc/heptane) to give 2,5-dichloro-*N*-((2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide (12.2 g, 24.6 mmol, 75% yield) as a white solid. MS (ESI, +ve) m/z: 496.0/497.9 $(M + 1)^+$.

[0763] A solution of 2,5-dichloro-*N*-((2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)carbamoyl)-6-(2-fluorophenyl)nicotinamide (12.2 g, 24.6 mmol) and potassium bis(trimethylsilyl)amide (1 M in THF, 49.2 mL, 49.2 mmol) in tetrahydrofuran (100 mL) was stirred at 0 °C for 20 min. The reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% EtOAc/heptane) to give 6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (3.18 g, 6.91 mmol, 28% yield) as an off-white solid and slighly impure 6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-

dione (5.89 g, 12.8 mmol, 52% yield) as a light yellow solid. ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.21 (br s, 1 H) 8.63 (s, 1 H) 7.42 - 7.52 (m, 1 H) 7.08 - 7.24 (m, 3 H) 2.78 (spt, *J* = 6.7 Hz, 1 H) 2.30 (s, 3 H) 1.24 (d, *J* = 6.6 Hz, 3 H) 1.09 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -113.25 (s, 1 F). MS (ESI, +ve) *m/z*: 460.0 $(M + 1)^+$.

[0764] Step 3: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. A solution of 6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (5.89 g, 12.8 mmol), phosphoryl trichloride (1.43 mL, 15.4 mmol), and DIPEA (8.92 mL, 51.2 mmol) in acetonitrile (64 mL) was stirred at 80 °C for 30 min.

[0765] The solution was cooled in an ice bath, then *tert*-butyl (2*R*,5*S*)-2,5-dimethylpiperazine-1-carboxylate (2.74 g, 12.80 mmol, AstaTech, Inc., Bristol, PA) was added, and the dark red solution was stirred at RT for 5 min. The reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–80% EtOAc/heptane) to give *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (6.13 g, 9.34 mmol, 73% yield) as an amber foam and a mixture of atropisomers. MS (ESI, +ve) *m/z*: 656.0 $(M + 1)^+$.

[0766] Step 4: *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (separated atropisomers). Atropisomers of *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (6.13 g, 9.34 mmol) were separated via chiral SFC (IC, 21 x 250 mm, 5 µm, 40% MeOH/CO_2, 80 mL/min); this gave *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (first eluting isomer, 2.70 g, 4.12 mmol, 44% yield, **Intermediate 204**) [^{1}H NMR (400 MHz, $CDCl_3$) δ 8.13 (s, 1 H) 7.41 - 7.49 (m, 1 H) 7.16 - 7.24 (m, 2 H) 7.09 - 7.16 (m, 1 H) 3.45 - 5.07 (m, 6 H) 2.62 - 2.77 (m, 1 H) 2.22 (s, 3 H) 1.51 (s, 9 H) 1.43 - 1.49 (m, 3 H) 1.29 (d, *J* = 6.8 Hz, 3 H) 1.23 (d, *J* = 6.8 Hz, 3 H) 1.07 (d, *J* = 6.6

Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.91 (br s, 1 F). MS (ESI, +ve) *m/z*: 656.0 (M + 1)$^+$.] as a yellow solid and *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(2-chloro-4-isopropyl-6-methylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (second eluting isomer, 3.54 g, 5.39 mmol, 58% yield, **Intermediate 215**) [^{1}H NMR (400 MHz, $CDCl_3$) δ 8.13 (s, 1 H) 7.42 - 7.49 (m, 1 H) 7.17 - 7.24 (m, 2 H) 7.10 - 7.17 (m, 1 H) 3.41 - 5.07 (m, 6 H) 2.70 (quin, *J* = 6.6 Hz, 1 H) 2.24 (s, 3 H) 1.51 (s, 9 H) 1.43 - 1.50 (m, 3 H) 1.28 (d, *J* = 6.8 Hz, 3 H) 1.22 (d, *J* = 6.8 Hz, 3 H) 1.04 (d, *J* = 6.8 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -112.93 (br s, 1 F). MS (ESI, +ve) *m/z*: 656.0 (M + 1)$^+$.].

Intermediates 216 & 217

1-(2-Bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (two atropisomers)



[0767] Step 1: 2-Bromo-4-isopropyl-6-methylpyrimidin-5-amine. A solution of 4-isopropyl-6-methylpyrimidin-5-amine (2.01 g, 13.3 mmol, **Intermediate I-5**), acetic acid (0.076 mL, 1.33 mmol), and NBS (3.55 g, 20.0 mmol) in tetrahydrofuran (67 mL) was stirred at 65 °C for 15 min. The reaction mixture was diluted with EtOAc (300 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (3 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product in DCM was loaded onto the column and was purified via automated flash chromatography (silica gel, 0–50% EtOAc/heptane) to give 2-bromo-4-isopropyl-6-methylpyrimidin-5-amine (2.07 g, 9.01 mmol, 68% yield) as an amber oil. ^{1}H NMR (400 MHz, $CDCl_3$) δ 3.59 (br s, 2 H) 2.94 (spt, *J*= 6.8 Hz, 1 H) 2.38 (s, 3 H) 1.28 (d, *J* = 6.6 Hz, 6 H). MS (ESI, +ve) *m/z*: 230.0/232.1 (M + 1)$^+$.

[0768] Step 2: 1-(2-Bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione. A solution of 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (2.57 g, 9.01 mmol, **Intermediate 99B**) and oxalyl chloride (2 M in DCM, 4.95 mL, 9.91 mmol) in tetrahydrofuran (45 mL) was stirred under an air condensor and drying tube at 65 °C for 45 min; more oxalyl chloride (2 M in DCM, 4 mL, 8 mmol) was then added, and the reaction mixture was stirred for 15 min. The solution was concentrated, dissolved in tetrahydrofuran (45 mL), and cooled in an ice bath; the solution was added to 2-bromo-4-isopropyl-6-methylpyrimidin-5-amine (2.07 g, 9.01 mmol) in THF (5 mL) at RT; the yellow solution was stirred at RT for 5 min. The reaction mixture was concentrated, diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous sodium bicarbonate (3 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo* to give impure product as an amber oil. The crude product in DCM was loaded onto the column and was purified via automated flash chromatography (silica gel, 0–50% EtOAc/heptane) to give *N*-((2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)carbamoyl)-2,5-dichloro-6-(2-fluorophenyl)nicotinamide (3.42 g, 6.32 mmol, 70% yield) as a light yellow foam. MS (ESI, +ve) *m/z*: 541.8 $(M + 1)^+$.

[0769] A solution of *N*-((2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)carbamoyl)-2,5-dichloro-6-(2-fluorophenyl)nicotinamide (3.42 g, 6.32 mmol) and potassium bis(trimethylsilyl)amide (1 M in THF, 12.6 mL, 12.6 mmol) in tetrahydrofuran (32 mL) was stirred at 0 °C for 10 min. The reaction mixture was diluted with EtOAc (150 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (2 × 100 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–50% EtOAc/heptane) to give 1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (1.95 g, 3.86 mmol, 61% yield) as a light yellow solid. MS (ESI, +ve) *m/z*: 505.9 $(M + 1)^+$.

[0770] Step 3: 1-(2-Bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (separated atropisomers). Atropisomers of 1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (1.95 g, 3.86 mmol) were separated via SFC (OD 250 x 21 mm, 5 µm, 20% MeOH/CO_2, 80 g/min, 102 bar). This gave 1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (first eluting isomer, 556 mg, 1.10 mmol, 29% yield, **Intermediate 216**) [^{1}H

NMR (400 MHz, $CDCl_3$) δ 8.62 (s, 1 H) 7.42 - 7.50 (m, 1 H) 7.10 - 7.24 (m, 3 H) 2.75 (spt, *J* = 6.7 Hz, 1 H) 2.28 (s, 3 H) 1.23 (d, *J* = 6.6 Hz, 3 H) 1.07 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -113.27 (s, 1 F). MS (ESI, +ve) *m/z*: 506.0 $(M + 1)^+$.] as a beige solid and 1-(2-bromo-4-isopropyl-6-methylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (second eluting isomer, 511 mg, 1.01 mmol, 26% yield, **Intermediate 217**) [^1H NMR (400 MHz, $CDCl_3$) δ 8.61 (s, 1 H) 7.42 - 7.50 (m, 1 H) 7.10 - 7.24 (m, 3 H) 2.75 (spt, *J* = 6.7 Hz, 1 H) 2.28 (s, 3 H) 1.23 (d, *J* = 6.8 Hz, 3 H) 1.07 (d, *J* = 6.6 Hz, 3 H). ^{19}F NMR (377 MHz, $CDCl_3$) δ -113.27 (s, 1 F). MS (ESI, +ve) *m/z*: 506.0 $(M + 1)^+$.] as a beige solid.

Intermediate 218

Rac-2-(4-Acryloylpiperazin-2-yl)acetonitrile



Intermediate 218

[0771] A white suspension solution of 2-(piperazin-2-yl)acetonitrile dihydrochloride (287 mg, 1.45 mmol, Enamine Ltd.) in DCM (7 mL) was treated with n,n'-diisopropylethylamine (0.5 mL, 2.90 mmol) at room temperature. The resulting clear solution was cooled to 0 °C and then treated with acryloyl chloride (1.1 M in DCM, 1.3 mL, 1.45 mmol) and stirred for 5 min. The reaction mixture was evaporated to dryness to provide a mixture of the desired 2-(4-acryloylpiperazin-2-yl)acetonitrile and undesired 2-(1,4-diacryloylpiperazin-2-yl)acetonitrile as 1:1 ratio. The crude material was used in next step without purification.

Intermediate 219

5-Amino-4,6-diisopropylpyrimidine-2-carbonitrile



Intermediate I-38

Intermediate 219

[0772] To a 100-mL round-bottomed flask was added 2-bromo-4,6-diisopropylpyrimidin-5-amine (0.500 g, 1.937 mmol, **Intermediate I-38**), potassium hexacyanoferrate(ii) trihydrate (3.27 g, 7.75 mmol, Sigma-Aldrich, St. Louis, MO, USA), water (6.46 mL) and potassium acetate (0.570 g, 5.81 mmol) in 1,4-dioxane (6.46 mL). The reaction mixture was deoxygenated by bubbling Argon gas into the mixture for 5 min. Then methanesulfonato(2-

dicyclohexylphosphino-2',4',6'-tri-i-propyl-1,1'-biphenyl)(2'-amino-1,1'-biphenyl-2-yl)palladium(ii) (0.164 g, 0.194 mmol, Strem Chemicals, Inc., Newburyport, MA, USA) was added to the reaction mixture. The mixture was heated and stirred at 100 °C for 1 h, while under an inert (N_2) atmosphere. The flask was removed from the heat bath and allowed the reaction mixture to cool to ambient temperature. The mixture was diluted with sat. aq. $NaHCO_3$ and EtOAc. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over MgSO4, filtered and concentrated *in vacuo*. The crude material was triturated from EtOAc and heptane. The precipitate was collected by filtration and the solids were washed with heptane. This afforded 5-amino-4,6-diisopropylpyrimidine-2-carbonitrile (0.210 g, 1.028 mmol, 53.1 % yield, **Intermediate 219**) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 6.18 (s, 2 H) 3.21 - 3.29 (m, 2 H) 1.15 (d, *J*=6.63 Hz, 12 H). *m/z* (ESI, +ve ion): 205.3 (M+H)+.

Intermediate 220

4,6-Diisopropyl-2-(propylsulfonyl)pyrimidin-5-amine



Step 1 Step 2 Intermediate 220

[0773] Step 1. 4,6-Diisopropyl-2-(propylthio)pyrimidin-5-amine. To a 250-mL round-bottomed flask was added 4,6-dichloro-2-(propylthio)pyrimidin-5-amine (8.00 g, 33.6 mmol, Synthonix, Wake Forest, NC, USA) in tetrahydrofuran (67.2 mL). The mixture was deoxygenated by bubbling Argon (gas) into the mixture 5 min. Then 2-propylzinc bromide (0.5M in THF) (94 mL, 94 mmol), followed by xantphos pd g3 (0.956 g, 1.008 mmol) was added to the reaction mixture. The reaction mixture was heated to 60 °C and allowed the mixture to stir under an inert (N2) atmosphere for 2 h. The reaction mixture was quenched with sat. aq. NH4Cl (50 mL) and allowed the mixture to stir 10 min. The mixture was diluted with EtOAc and Brine solution (50 mL). The layers were separated and aqueous layer was extracted with EtOAc. The combined organic extracts were dried over MgSO4, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (220 grams), eluting with a gradient of 0-30% EtOAc/heptane, to provide 4,6-diisopropyl-2-(propylthio)pyrimidin-5-amine (6.148 g, 24.26 mmol, 72.2 % yield) as tan oil. 1H NMR (400 MHz, DMSO-d6) δ

4.80 (s, 2 H) 3.13 - 3.25 (m, 2 H) 2.91 - 3.05 (m, 2 H) 1.66 (sxt, J=7.26 Hz, 2 H) 1.12 (d, J=6.63 Hz, 12 H) 0.96 (t, J=7.36 Hz, 3 H). m/z (ESI, +ve ion): 254.1 (M+H)+.

[0774] Step 2. 4,6-Diisopropyl-2-(propylsulfonyl)pyrimidin-5-amine. To a 150-mL round-bottomed flask was added 4,6-diisopropyl-2-(propylthio)pyrimidin-5-amine (1.560 g, 6.16 mmol) and 3-chloroperoxybenzoic acid (3.45 g, 15.39 mmol, Sigma-Aldrich, St. Louis, MO, USA) in tetrahydrofuran (30.8 mL). The reaction mixture was allowed to stir under an inert (N2) atmosphere for 45 min. The reaction mixture was cooled to 0 °C with a wet ice/water bath. Then sat. aq. NaHCO3 was added slowly (3 x 10 mL portions) over 20 min. The mixture was diluted with 4:1 EtOAc/MeOH and brine solution (20 mL). The aqueous layer was extracted with EtOAc and the organic extracts were combined, dried over MgSO4, filtered and concentrated in vacuo. The crude material (3.118 grams) was diluted with DMF (20 mL), then agitated with sonicator until mixture became homogeneous. The mixture was slowly poured into water (100 mL) and the precipitate was collected by filtration. The solids were washed with water and allowed to dry in a reduced-pressure oven (45 °C) overnight. This afforded 4,6-diisopropyl-2-(propylsulfonyl)pyrimidin-5-amine (1.645 g, 5.76 mmol, 94 % yield, **Intermediate 220**) as tan solid. 1H NMR (400 MHz, DMSO-d6) δ 6.08 (br s, 2 H) 3.48 - 3.68 (m, 3 H) 1.79 (br s, 2 H) 1.29 (br s, 12 H) 1.07 (br s, 4 H). m/z (ESI, +ve ion): 286.3 (M+H)+.

Intermediate 221

(4-Amino-3-isopropyl-5-methylphenyl)dimethylphosphine oxide



Intermediate 221

[0775] Step 1: 4-Iodo-2-isopropyl-6-methylaniline. Iodine (9.3 g, 37 mmol) was slowly added to a biphasic mixture of 2-(1-methylethyl)-6-methylaniline (Advanced ChemBlocks Inc., Burlingame, CA, USA, 5.5 mL, 35 mmol) and sodium bicarbonate (5.9 g, 70 mmol) in DCM–water (1:1, 70 mL) at room temperature. After 1 h, the reaction was quenched by adding 1 N aqueous sodium thiosulfate (100 mL). Additional DCM (70 mL) was added and the biphasic mixture stirred vigorously for 30 min at room temperature. The layers were partitioned and the aqueous layer was washed with DCM (1 x 100 mL). The combined organic extracts were dried over anhydrous magnesium sulfate, filtered, and concentrated *in vacuo* to afford

crude 4-iodo-2-isopropyl-6-methylaniline (9.4 g, 98 % yield) as a dark purple oil that was carried forward in the next step without purification. LRMS: (ESI, +ve ion) *m/z* 275.9 $(M+H)^+$.

[0776] Step 2: (4-Amino-3-isopropyl-5-methylphenyl)dimethylphosphine oxide (Intermediate 221). A nitrogen-sparged mixture of 4-iodo-2-isopropyl-6-methylaniline (500 mg, 1.8 mmol), dimethylphosphine oxide (170 mg, 2.2 mmol), potassium phosphate, tribasic (420 mg, 2.0 mmol), palladium(II) acetate (20 mg, 0.091 mmol), and Xantphos (53 mg, 0.091 mmol) in DMF (3.2 mL) was stirred 120 °C for 18 h. The reaction mixture was filtered through a pad of Celite and rinsed with EtOAc. The filtrate was concentrated *in vacuo* and then redissolved in DCM (50 mL). The solution was washed with 1 M LiCl (2 × 50 mL) and the aqueous layer was back-extracted with EtOAc (1 × 50 mL). The combined organic extracts were dried over anhydrous magnesium sulfate, filtered, and concentrated to afford (4-amino-3-isopropyl-5-methylphenyl)dimethylphosphine oxide (**Intermediate 221**, 440 mg, 100% yield) as a thick red-brown oil. ^{1}H NMR (400 MHz, DMSO-d_6) δ 1.16 (d, J = 6.6 Hz, 6 H), 1.52 (d, J = 13.1 Hz, 6 H), 2.11 (s, 3 H), 3.04 (sept, J = 6.6 Hz, 1 H), 5.07 (s, 2 H), 7.14 (d, J = 11.4 Hz, 1 H), 7.23 (d, J = 12.1 Hz, 1 H); ^{31}P{^{1}H} NMR (162 MHz, DMSO-d_6) δ 31.71 (s, 1 P); LRMS: (ESI, +ve ion) *m/z* 226.1 $(M+H)^+$.

Intermediate 222

3-Isopropyl-5-methylpyridin-4-amine



Intermediate 222

[0777] Step 1: 3-Methyl-5-(prop-1-en-2-yl)pyridin-4-amine. A nitrogen-sparged mixture of sodium carbonate (7.7 g, 72 mmol), $Pd(Amphos)Cl_2$ (0.64 g, 0.90 mmol), 3-bromo-5-methyl-4(1*H*)-pyridinimine (5.1 g, 27 mmol), and 2-isopropenylboronic acid, pincol ester (Combi-Blocks Inc., San Diego, CA, USA, 6.8 mL, 36 mmol) in dioxane–water (2:1, 90 mL) was heated at 90 °C for 15 h. EtOAc (100 mL) and water (100 mL) were added and the layers were partitioned. The aqueous layer was washed with EtOAc (2 x 50 mL). The combined organic extracts were sequentially washed with 25% saturated sodium bicarbonate (100 mL) and water (100 mL), then treated with 1 N HCl (2 × 150 mL). The combined acidic aqueous washes were collected and the organic phase was discarded. The aqueous was treated with 5 N NaOH (200 mL) and extracted with EtOAc (2 x 100 mL). The combined organic extracts were

dried over anhydrous sodium sulfate, filtered, and then concentrated *in vacuo* to give 3-methyl-5-(prop-1-en-2-yl)pyridin-4-amine (3.7 g, 92% yield) as a purple-brown oil which was carried forward in the next step without purification. ^{1}H NMR (400 MHz, DMSO-d_6) δ 2.00 (s, 3 H), 2.03 (s, 3 H), 4.97 - 4.99 (m, 1 H), 5.25 - 5.29 (m, 1 H), 5.33 (br s, 2 H), 7.77 (s, 1 H), 7.83 (s, 1 H); LRMS: (ESI, +ve ion) *m/z* 149.0 $(M+H)^+$.

[0778] Step 2: 3-Isopropyl-5-methylpyridin-4-amine (Intermediate 222). Crude 3-methyl-5-(prop-1-en-2-yl)pyridin-4-amine (3.7 g, 25 mmol) was dissolved in ethanol (60 mL) and then palladium 10 wt. % on activated carbon (11 g, 5.0 mmol) and ammonium formate (17 g, 270 mmol) were added. The resulting black solution was heated at 80 °C for 9 h, then filtered through a pad of Celite using EtOAc. The filtrate was concentrated to ~50 mL *in vacuo* and then 5 N sodium hydroxide (100 mL), water (200 mL), and MTBE (150 mL) were added and the phases mixed and separated. The aqueous layer was extracted with additional MTBE (3 × 100 mL) and the organic layers were combined, dried over anhydrous magnesium sulfate, and concentrated *in vacuo* to give 3-isopropyl-5-methylpyridin-4-amine (**Intermediate 222**, 3.7 g, 100% yield) as a clear oil that later solidified to an opaque off-white wax. ^{1}H NMR (400 MHz, DMSO-d_6) δ 1.17 (d, J = 6.8 Hz, 6 H), 2.01 (s, 3 H), 2.95 - 3.06 (m, 1 H), 5.43 (br s, 2 H), 7.78 (s, 1 H), 7.88 (s, 1 H); LRMS: (ESI, +ve ion) *m/z* 151.1 $(M+H)^+$.

Intermediate 223

1-(*cis*-2,6-Dimethylpiperazin-1-yl)prop-2-en-1-one 2,2,2-trifluoroacetate



Intermediate 223

[0779] Step 1: *tert*-Butyl 4-acryloyl-*cis*-3,5-dimethylpiperazine-1-carboxylate. To a solution of *tert*-butyl *cis*-3,5-dimethylpiperazine-1-carboxylate (Combi-Blocks Inc., San Diego, CA, USA, 130 g, 580 mmol, 1.0 equiv) and triethylamine (100 mL, 700 mmol, 1.2 equiv) in tetrahydrofuran (1250 mL, 10 mL/g) was added acryloyl chloride (56 mL, 700 mmol, 1.2 equiv) at 0 °C. The reaction mixture was stirred for 1 h at 0 °C. The reaction mixture was diluted with ethyl acetate (2.0 L) and washed with water (2 x 500 mL). The organic layer was washed with 1 N aqueous HCl (500 mL), saturated $NaHCO_3$ (500 mL), and brine (500 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated *in vacuo*. The crude

material was adsorbed to a plug of silica gel and purified by silica gel column chromatography (30–40% EtOAc/hexane) to provide the title compound as off-white solid (140 g, 91%). MS (ESI, +ve ion) *m/z*: 269.0 [M+1]; ^{1}H NMR (400 MHz, DMSO-d_6) δ 6.78 – 6.72 (ddd, *J* = 16.7, 10.4, 1.5 Hz, 1H), 6.15 – 6.11 (dt, *J* = 16.6, 1.9 Hz, 1H), 5.70 – 5.67 (dt, *J* = 10.6, 1.9 Hz, 1H), 4.35 – 4.25 (m, 2H), 3.82 (d, *J* = 12.5 Hz, 2H), 3.01 – 2.95 (m, 2H), 1.43 – 1.42 (d, *J* = 1.6 Hz, 9H), 1.19 – 1.14 (m, 6H).

[0780] Step 2: 1-(*cis*-2,6-Dimethylpiperazin-1-yl)prop-2-en-1-one 2,2,2-trifluoroacetate (Intermediate 223). To a solution of *tert*-butyl 4-acryloyl-*cis*-3,5-dimethylpiperazine-1-carboxylate (140 g, 530 mmol, 1.0 equiv) in dichloromethane (1.4 L, 10 mL/g) was added trifluoroacetic acid (510 mL, 6900 mmol, 13 equiv) at 0 °C. The reaction mixture was stirred for 1 h at 0 °C and concentrated *in vacuo*. The crude material was stirred with diethyl ether (1.0 L) for 30 min at room temperature. The precipitated solid was filtered and washed with *n*-hexane (500 mL) and dried to afford 1-(*cis*-2,6-dimethylpiperazin-1-yl)prop-2-en-1-one 2,2,2-trifluoroacetate (**Intermediate 223**, 140 g, 96%) as off-white solid. MS (ESI, +ve ion) *m/z*: 169.0 [M+1]; ^{1}H NMR (400 MHz, DMSO-d_6) δ 9.59 (br, 1H), 9.17 (br, 1H), 6.82 – 6.75 (ddt, *J* = 17.0, 10.8, 2.1 Hz, 1H), 6.21 – 6.16 (dq, *J* = 16.5, 2.3 Hz, 1H), 5.77 – 5.74 (dq, *J* = 10.6, 2.8, 2.4 Hz, 1H), 4.60 (t, *J* = 6.9 Hz, 2H), 3.40 – 3.38 (dd, *J* = 13.4, 5.7 Hz, 2H), 3.14 – 3.11 (dd, *J* = 13.2, 5.5 Hz, 2H), 1.31 – 1.29 (m, 6H).

Intermediate 224

6-Chloro-1-(4-chloro-2-isopropylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0781] Same as **Example 153** up tp Step 3 using 4-chloro-2-isopropylpyridin-3-amine (**Intermediate 19o**) in Step 1 and (2-fluorophenyl) boronic acid in Step 3. ^{1}H NMR (DMSO-d_6) δ: 12.20-12.54 (m, 1H), 8.60-8.62 (m, 1H), 8.57-8.59 (m, 1H), 7.56-7.60 (m, 1H), 7.49-7.56 (m, 1H), 7.26-7.35 (m, 2H), 7.19-7.26 (m, 1H), 3.07-3.16 (m, 1H), 1.10-1.14 (m, 3H), 0.99-1.03 (m, 3H). MS (ESI, +ve) *m/z*: 444.6 $(M + 1)^+$.

Intermediate 225

6,7-Dichloro-1-(4-(dimethylamino)-2-isopropylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0782] Same as **Intermediate 61A** using 2-isopropyl,N^4,N^4-dimethylpyridine-3,4-diamine (**Intermediate 163**) instead of 2-isopropylaniline in Step 1. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 12.32 (br s, 1 H), 8.59 (s, 1 H), 8.28 (d, *J*=5.8 Hz, 1 H), 6.77 (d, *J*=5.8 Hz, 1 H), 2.69 (s, 7 H), 1.05 (d, *J*=6.6 Hz, 3 H), 0.95 (d, *J*=6.6 Hz, 3 H). MS (ESI, +ve) *m/z*: 393.6 $(M + 1)^+$.

Intermediate 226

4-((2*S*,5*R*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(4-(dimethylamino)-2-isopropylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



[0783] Same as **Intermediate 92B** using 2-isopropyl, N^4,N^4-dimethylpyridine-3,4-diamine (**Intermediate 163**). 1H NMR (400 MHz, *DMSO-d6*) δ 8.54 (s, 1 H), 8.42 (d, *J*=9.3 Hz, 1 H), 8.25 (dd, *J*=9.7, 5.8 Hz, 1 H), 6.75 (br d, *J*=5.8 Hz, 1 H), 6.18 (dd, *J*=16.8, 2.3 Hz, 1 H), 5.72 - 5.78 (m, 1 H), 4.60 - 4.96 (m, 2 H), 4.33 - 4.52 (m, 1 H), 3.92 - 4.25 (m, 2 H), 3.83 (br s, 2 H), 2.68 (br d, *J*=6.0 Hz, 6 H), 1.34 (br d, *J*=6.4 Hz, 3 H), 1.23 - 1.30 (m, 3 H), 1.11 (br d, *J*=6.6 Hz, 3 H), 1.01 - 1.05 (m, 3 H). MS (ESI, +ve) *m/z*: 543.6 $(M + 1)^+$.

Intermediate 227

6,7-Dichloro-1-(2-(dimethylphosphoryl)phenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0784] Same as **Intermediate 61A** using (2-aminophenyl)dimethylphosphine oxide (AstaTech, Inc.) instead of 2-isopropylaniline. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ ppm 11.78 - 12.20 (m, 1 H), 8.53 (s, 1 H), 7.82 - 7.90 (m, 1 H), 7.70 - 7.77 (m, 1 H), 7.60 - 7.67 (m, 1 H), 7.46 (dd, *J*=7.9, 2.9 Hz, 1 H), 1.60 (d, *J*=13.5 Hz, 3 H), 1.53 (d, *J*=13.5 Hz, 3 H). MS (ESI, +ve) *m/z*: 384 $(M + 1)^+$.

Intermediate 228

***tert*-Butyl (2*R*,5*S*)-4-(1-(6-bromo-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**



Intermediate 165 **Intermediate 228**

[0785] To a solution of 6-bromo-2,4-diisopropylpyridin-3-amine (0.217 g, 0.844 mmol, **Intermediate 196**)and DIPEA (0.309 ml, 1.772 mmol) in THF (10 mL) at rt was added dropwise phosgene solution, 15% in toluene (0.662 ml, 0.928 mmol) and the resulting mixture was stirred at rt for 10 min then the mixture was brought to 0C and solid *tert*-butyl (2*R*,5*S*)-4-((2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)(imino)methyl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 165,** 0.487 g, 1.013 mmol) was added in three portions. The ice-

bath was removed and after stirring for 15 min at rt *tert*-butyl (2*R*,5*S*)-4-((((6-bromo-2,4-diisopropylpyridin-3-yl)carbamoyl)imino)(2,5-dichloro-6-(2-fluorophenyl)pyridin-3-yl)methyl)-2,5-dimethylpiperazine-1-carboxylate was observed then sodium *tert*-butoxide (0.243 g, 2.53 mmol) was added and the resulting mixture was stirred at rt for 20 min. The reaction went to completion, washed with water, extracted with EtOAc and purified by chromatography on silica gel using 0-40% EtOAc in heptane to afford *tert*-butyl (2*R*,5*S*)-4-(1-(6-bromo-2,4-diisopropylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 228,** 0.213 g, 0.293 mmol, 34.7 % yield) as a light yellow solid. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 8.46 (s, 1 H), 7.52 (d, *J*=4.4 Hz, 2 H), 7.27 - 7.37 (m, 2 H), 7.16 - 7.24 (m, 1 H), 4.79 - 4.89 (m, 1 H), 4.23 - 4.43 (m, 1 H), 4.16 (br d, *J*=13.5 Hz, 1 H), 3.78 - 3.92 (m, 1 H), 3.66 - 3.76 (m, 1 H), 1.46 (s, 9 H), 1.35 (br d, *J*=6.8 Hz, 2 H), 1.26 (br s, 4 H), 1.15 - 1.21 (m, 3 H), 1.02 - 1.09 (m, 6 H), 0.87 - 0.96 (m, 6 H). MS (ESI, +ve) *m/z*: 728.4 $(M + 1)^+$.

Intermediate 229

6-Chloro-1-(4-ethyl-2-isopropylpyridin-3-yl)-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0786] Same as **Example 153** up to Step 3 using **Intermediate 197** in Step 1 and (2-fluorophenyl) boronic acid in Step 3. ^{1}H NMR (400 MHz, *DMSO-d*$_6$) δ 12.03 - 12.45 (m, 1 H), 8.56 (s, 1 H), 8.48 (d, *J*=5.0 Hz, 1 H), 7.21 - 7.33 (m, 4 H), 7.16 - 7.21 (m, 1 H), 2.89 (quin, *J*=6.5 Hz, 1 H), 2.35 - 2.44 (m, 2 H), 1.09 (d, *J*=6.8 Hz, 3 H), 1.00 (t, *J*=7.6 Hz, 3 H), 0.95 (d, *J*=6.6 Hz, 3 H). MS (ESI, +ve) *m/z*: 439.1 $(M + 1)^+$.

Intermediate 233

4-((2*S*,5*R*,*M*)-4-Acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-amino-6-fluorophenyl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one



Intermediate 92B Intermediate 233

[0787] A mixture of 4-((2*S*,5*R*,*M*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-6,7-dichloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 92B**, 3.07 g, 5.96 mmol), tetrakis(triphenylphosphine)palladium (0.34 g, 0.30 mmol), (2-amino-6-fluorophenyl)boronic acid pinacol ester (1.55 g, 6.55 mmol, CombiPhos, Trenton, NJ) and sodium carbonate, anhydrous, powder (3.16 g, 29.8 mmol) in 1,4-dioxane (20 mL)/water (10 mL) was stirred at 90 °C for 40 min. To the resulting mixture was added water (25 mL) and the mixture was extracted with EtOAc (2 × 50 mL). The organic extracts were combined and dried over $MgSO_4$. The solution was filtered and concentrated *in vacuo* to give the crude product as a yellow solid. The resulting crude product was adsorbed onto a plug of silica gel and purified by silica gel chromatography (eluent: 0-10% EtOAc (with 10 % MeOH)/heptane) to give 4-((2*S*,5*R*,*M*)-4-acryloyl-2,5-dimethylpiperazin-1-yl)-7-(2-amino-6-fluorophenyl)-6-chloro-1-(2-isopropyl-4-methylpyridin-3-yl)pyrido[2,3-*d*]pyrimidin-2(1*H*)-one (**Intermediate 233,** 1.83 g, 3.09 mmol, 52.0 % yield) as a yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.43 (s, 1 H), 8.39 (d, J=4.8 Hz, 1 H), 7.19 (d, J=4.8 Hz, 1 H), 7.00 - 7.11 (m, 1 H), 6.82 (br dd, J=16.6, 10.6 Hz, 1 H), 6.45 (d, J=8.3 Hz, 1 H), 6.27 - 6.35 (m, 1 H), 6.19 (dd, J=16.6, 2.3 Hz, 1 H), 5.76 (ddd, J=10.1, 5.5, 2.2 Hz, 1 H), 5.07 - 5.19 (m, 2 H), 4.45 - 4.90 (m, 2 H), 3.47 - 4.24 (m, 4 H), 2.60 - 2.88 (m, 1 H), 1.85 - 1.99 (m, 3 H), 1.30 - 1.39 (m, 3 H), 1.16 - 1.29 (m, 3 H), 1.03 - 1.11 (m, 3 H), 0.87 - 1.03 (m, 3 H). ^{19}F NMR (376 MHz, DMSO-d_6) δ -116.01 - -115.34 (m, 1 F). *m/z* (ESI, +ve): 590.2 $(M+H)^+$.

Intermediate 234

5-Amino-6-isopropyl-3-methylpyrimidin-4(3H)-one



Intermediate 234

[0788] A solution of 5-amino-6-chloro-3-methyl-3,4-dihydropyrimidin-4-one (0.414 g, 2.59 mmol; Enamine LLC, Monmouth Jct., NJ), XantPhos Pd G3 (0.268 g, 0.259 mmol), and isopropylzinc bromide (0.5 M in THF, 5.71 mL, 2.85 mmol) in 1,4-dioxane (3 mL) was stirred at 50 °C for 30 min. The reaction mixture was diluted with EtOAc (100 mL), added to a separatory funnel, and washed with saturated, aqueous ammonium chloride (2 × 75 mL); the organic layer was separated, dried over anhydrous Na_2SO_4, and concentrated *in vacuo*. The crude product was adsorbed onto silica and was purified via automated flash chromatography (silica gel, 0–90% [(3:1) EtOAc/EtOH]/heptane) to give 5-amino-6-isopropyl-3-methylpyrimidin-4(3H)-one (34 mg, 0.20 mmol, 8% yield, **Intermediate 234**) as a light brown film. ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.67 (s, 1 H) 3.94 (br s, 2 H) 3.52 (s, 3 H) 2.85 (spt, J = 6.8 Hz, 1 H) 1.22 (d, J = 6.8 Hz, 6 H). MS (ESI, +ve) *m/z*: 168.1 $(M + 1)^+$.

Intermediate 235

tert-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-formyl-4,6-diisopropylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-d]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate



[0789] **Step 1.** ***tert*-Butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate**. To a 250-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(1-(2-bromo-4,6-diisopropylpyrimidin-5-yl)-6-chloro-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (**Intermediate 232**, 2.352 g, 3.23 mmol), sodium carbonate (1.026 g, 9.68 mmol) and vinylboronic acid pinacol ester (1.09 mL, 6.45 mmol) in 1,4-dioxane (25.8 mL)-water (6.45 mL) (4:1). The reaction mixture was deoxygenated by bubbling argong (gas) into the mixture for 5 min. Then (1,1'-bis(diphenylphosphino) ferrocene) dichloropalladium (0.236 g, 0.323 mmol) was added to the reaction mixture. The reaction mixture was stirred and heated at 90 °C, while under an inert (N_2) atmosphere for 16 h. The reaction mixture was allowed to cool to rt, then the mixture was diluted was EtOAc and sat. aq. $NaHCO_3$. The layers were separated and

the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a Interchim (25 micron) silica-gel column (200 grams), eluting with a gradient of 0-70% EtOAc/DCM, to provide *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (1.579 g, 2.335 mmol, 72.4 % yield) as tan solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.46 (s, 1 H) 7.48 - 7.54 (m, 1 H) 7.18 - 7.34 (m, 3 H) 6.79 (dd, *J*=17.21, 10.37 Hz, 1 H) 6.55 (dd, *J*=17.21, 2.07 Hz, 1 H) 5.71 - 5.76 (m, 1 H) 4.85 (br s, 1 H) 4.26 - 4.42 (m, 1 H) 4.17 (br d, *J*=14.31 Hz, 1 H) 3.81 - 3.92 (m, 1 H) 3.71 (br d, *J*=12.65 Hz, 1 H) 3.44 - 3.61 (m, 1 H) 2.62 - 2.76 (m, 2 H) 1.45 (s, 9 H) 1.36 (d, *J*=6.63 Hz, 3 H) 1.16 - 1.20 (m, 3 H) 1.07 - 1.11 (m, 6 H) 0.91 - 0.97 (m, 6 H). *m/z* (ESI, +ve ion): 676.2 $(M+H)^+$.

[0790] Step 2. *tert*-Butyl (2*R*,5*S*)-4-(6-chloro-7-(2-fluorophenyl)-1-(2-formyl-4,6-diisopropylpyrimidin-5-yl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate. To a 100-mL round-bottomed flask was added *tert*-butyl (2*R*,5*S*)-4-(6-chloro-1-(4,6-diisopropyl-2-vinylpyrimidin-5-yl)-7-(2-fluorophenyl)-2-oxo-1,2-dihydropyrido[2,3-*d*]pyrimidin-4-yl)-2,5-dimethylpiperazine-1-carboxylate (0.300 g, 0.444 mmol) in acetone (3.70 mL)-water (0.74 mL) (5:1). To this solution was added potassium osmate(vi) (0.016 g, 0.044 mmol) and solid 4-methylmorpholine 4-oxide (0.182 g, 1.553 mmol). The overall reaction mixture was allowed to stir under an inert (N_2) atmosphere, while at rt for 15 min. The reaction mixture was quenched with solid sodium sulfite (40 mg) and allowed the mixture to stir 5 min. The reaction mixture was partially concentrated (to remove acetone) *in vacuo*. The mixture was diluted with EtOAc and brine soln. The layers were separated and the aqueous layer was extracted with EtOAc. The organics were combined, dried over $MgSO_4$, filtered and concentrated *in vacuo*. The crude residue (0.362 grams) was carried immediately into the next step of the synthesis, without further purification.

[0791] The crude diol was diluted with THF (15 mL) then sodium (meta)periodate (0.190 g, 0.887 mmol), followed by water (0.1 mL) was added to the mixture. The resulting reaction mixture was allowed to stir under an inert (N_2) atmosphere for 2 h. The reaction mixture was diluted with a mixture of EtOAc/heptane (1:1) (40 mL). The mixture was agitated with sonicator 1 min. The mixture was filtered and the filtrate was collected, then the mixture was diluted with sat. aq. $NaHCO_3$ (40 mL. The layers were separated and the aqueous layer was extracted with EtOAc (3x). The combined organic extracts were dried over $MgSO_4$, filtered

and concentrated *in vacuo*. This afforded *tert*-butyl (2*R*,5*S*)-4-(3-chloro-2-(2-fluorophenyl)-8-(2-formyl-4,6-diisopropylpyrimidin-5-yl)-7-oxo-7,8-dihydro-1,6-naphthyridin-5-yl)-2,5-dimethylpiperazine-1-carboxylate (0.190 g, 0.281 mmol, 63.2 % yield, **Intermediate 235**) as a light-yellow solid. This material was carried into the next step of the synthesis, without further purification. ^{1}H NMR (400 MHz, DMSO-d_6) δ 9.99 (s, 1 H) 8.55 (s, 1 H) 7.50 - 7.62 (m, 1 H) 7.21 - 7.40 (m, 3 H) 4.92 (br s, 1 H) 4.32 - 4.51 (m, 1 H) 4.27 (br d, *J*=13.48 Hz, 1 H) 3.85 - 3.99 (m, 1 H) 3.78 (br d, *J*=13.06 Hz, 1 H) 3.48 - 3.67 (m, 1 H) 2.73 - 2.98 (m, 2 H) 1.51 (s, 9 H) 1.41 - 1.45 (m, 3 H) 1.23 - 1.27 (m, 3 H) 1.15 - 1.21 (m, 6 H) 1.00 - 1.08 (m, 6 H). *m/z* (ESI, +ve ion): 678.1 $(M+H)^+$.

Intermediate 236

1-(6-Bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione





[0792] Step 1. *N*-((6-Bromo-2-isopropyl-4-methylpyridin-3-yl)carbamoyl)-2,5-dichloro-6-(2-fluorophenyl)nicotinamide.

[0793] To a 250-mL round-bottomed flask was added 2,5-dichloro-6-(2-fluorophenyl)nicotinamide (**Intermediate 99B**, 6.00 g, 19.78 mmol) and oxalyl chloride (14.8 mL, 29.7 mmol) in tetrahydrofuran (39.6 mL). The flask was fitted with a Findenser, and the reaction mixture was stirred and heated at 80 °C for 45 min. The reaction mixture was concentrated in vacuo. The crude mixture was carried into the next step of the synthesis, without further purification to prevent decomposition.

[0794] To a 150-mL round-bottomed flask was added (2,5-dichloro-6-(2-fluorophenyl)nicotinoyl)carbamoyl isocyanate (crude material from previous step) in tetrahydrofuran (39.6 mL). The reaction mixture was cooled to -10 °C with a dilute acetone/dry ice bath, then a solution of 6-bromo-2-isopropyl-4-methylpyridin-3-amine (5.21 g, 22.75 mmol, **Intermediate 201**) in THF (3 mL) was added into the reaction mixture. The mixture was allowed to stir under an inert (N2) atmosphere for 1 h. The reaction mixture was concentrated in vacuo. The crude material was triturated from EtOAc and heptane, then the solids were collected by filtration. The solids were washed with heptane. This afforded N-((6-bromo-2-isopropyl-4-methylpyridin-3-yl)carbamoyl)-2,5-dichloro-6-(2-fluorophenyl)nicotinamide (6.880 g, 12.74 mmol, 64.4 % yield) as tan solid. *m/z* (ESI, +ve ion): 539.0 $(M+H)^+$.

[0795] Step 2. 1-(6-Bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione.

To a 250-mL round-bottomed flask was added N-((6-bromo-2-isopropyl-4-methylpyridin-3-yl)carbamoyl)-2,5-dichloro-6-(2-fluorophenyl)nicotinamide (5.00 g, 9.26 mmol) in tetrahydrofuran (63.3 mL). Then potassium bis(trimethylsilyl)amide, 1m solution in tetrahydrofuran (11.5 mL, 11.57 mmol) was added via an addition funnel dropwise into the reaction mixture over 5 min. The reaction mixture was allowed to stir under an inert (N_2) atmosphere for 3 h. More KHMDS (0.5 eq; 8 mL) was added dropwise into the reaction mixture. The reaction mixture was quenched with sat. aq. NH_4Cl (50 mL), then diluted the mixture with 3:1 EtOAc/MeOH and brine solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were dried over $MgSO_4$, filtered and concentrated in vacuo. The crude material was absorbed onto a plug of silica gel and purified by chromatography through a ISCO (300 gram column), eluting with a gradient of 0-20% EtOAc in CH_2CL_2, to provide 1-(6-bromo-2-isopropyl-4-methylpyridin-3-yl)-6-chloro-7-(2-fluorophenyl)pyrido[2,3-d]pyrimidine-2,4(1H,3H)-dione (2.05 g, 4.07 mmol, Intermediate 236) as light-yellow solid. *m/z* (ESI, +ve ion): 503.0 $(M+H)^+$.

Intermediate 240

6,7-dichloro-1-(4-chloro-2-isopropylpyridin-3-yl)pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione



[0796] **Intermediate 240** was made same as **Intermediate 61A** but using **Intermediate 190** instead of 2-isopropylaniline. ^{1}H NMR (DMSO-d_6) δ: 12.34-12.56 (m, 1H), 8.66-8.70 (m, 1H), 8.61-8.64 (m, 1H), 7.62-7.68 (m, 1H), 1.10-1.15 (m, 3H), 1.05-1.09 (m, 3H). m/z (ESI, +ve ion): 385 (M+H).

BIOLOGICAL ANALYSIS

Section 4--Biochemical and cellular activity of compounds

[0797] For compounds in the "Biological Data" table below, the following assay conditions were employed:

Coupled Nucleotide Exchange Assay: Purified GDP-bound KRAS protein (aa 1-169), containing both G12C and C118A amino acid substitutions and an *N*-terminal His-tag, was pre-incubated in assay buffer (25 mM HEPES pH 7.4, 10 mM $MgCl_2$, and 0.01% Triton X-100) with a compound dose-response titration for either 5 min or 2 hours (see Table 15). Following compound pre-incubation, purified SOS protein (aa 564-1049) and GTP (Roche 10106399001) were added to the assay wells and incubated for an additional 30 min (for 5 min compound pre-incubation) or 1 hour (for 2 hour compound pre-incubation). To determine the extent of inhibition of SOS-mediated nucleotide exchange, purified GST-tagged cRAF (aa 1-149), nickel chelate AlphaLISA acceptor beads (PerkinElmer AL108R), and AlphaScreen glutathione donor beads (PerkinElmer 6765302) were added to the assay wells and incubated for 10 minutes. The assay plates were then read on a PerkinElmer EnVision Multilabel Reader, using AlphaScreen® technology, and data were analyzed using a 4-parameter logistic model to calculate IC_{50} values.

[0798] **Phospho-ERK1/2 MSD Assay**: MIA PaCa-2 (ATCC® CRL-1420™) and A549 (ATCC® CCL-185™) cells were cultured in RPMI 1640 Medium (ThermoFisher Scientific 11875093) containing 10% fetal bovine serum (ThermoFisher Scientific 16000044) and 1x penicillin-streptomycin-glutamine (ThermoFisher Scientific 10378016). Sixteen hours prior

to compound treatment, MIA PaCa-2 or A549 cells were seeded in 96-well cell culture plates at a density of 25,000 cells/well and incubated at 37°C, 5% CO2. A compound dose-response titration was diluted in growth media, added to appropriate wells of a cell culture plate, and then incubated at 37°C, 5% CO2 for 2 or 4 hours (see Table 15). Following compound treatment, cells were stimulated with 10 ng/mL EGF (Roche 11376454001) for 10 min, washed with ice-cold Dulbecco's phosphate-buffered saline, no Ca2+ or Mg2+ (ThermoFisher Scientific 14190144), and then lysed in RIPA buffer (50 mM Tris-HCl pH 7.5, 1% Igepal, 0.5% sodium deoxycholate, 150 mM NaCl, and 0.5% sodium dodecyl sulfate) containing protease inhibitors (Roche 4693132001) and phosphatase inhibitors (Roche 4906837001). Phosphorylation of ERK1/2 in compound-treated lysates was assayed using Phospho-ERK1/2 Whole Cell Lysate kits (Meso Scale Discovery K151DWD) according to the manufacturer's protocol. Assay plates were read on a Meso Scale Discovery Sector Imager 6000, and data were analyzed using a 4-parameter logistic model to calculate IC50 values.

Biological Data

Ex.#	Coupled exchange IC50 (5 min; μM)	p-ERK IC_{50} (2 h, MiaPaCa-2; μM)	p-ERK IC_{50} (2 h, A549; μM)
55-50	0.09	0.023	> 7.5
55-51	0.19	0.079	> 7.5
55-51-1	0.298	0.157	> 7.5
55-51-2	0.07	0.037	> 7.5
55-52	10.215	10	> 7.5
55-53	0.356	-	-
55-54	0.032	0.016	> 7.5
55-55	3.14	-	-
55-56	0.023	0.01	> 7.5
55-57	0.168	0.177	> 7.5
55-58	0.052	0.056	> 7.5
55-58-1	0.03	0.107	> 7.5
55-58-2	0.092	0.101	> 7.5
55-59	174	10	> 7.5
55-60	7.85	7.77	> 7.5
69-10	11	-	-

69-11	1.14	-	-
69-12	2.88	-	-
72-20	0.382	0.223	> 7.5
72-21	0.468	0.306	> 7.5
72-22	0.408	-	-
72-23	0.647	-	-
79-10	2.78	-	-
79-11	26.7	-	-
79-12	5.46	-	-
80-10	0.077	0.063	> 7.5
80-10-1	2.04	-	-
80-10-2	0.053	0.04	> 7.5
80-11	0.078	0.222	> 7.5
80-12	0.064	0.074	> 7.5
80-13	0.043	0.053	> 7.5
80-14-1	1.26	-	-
80-14-2	0.061	0.031	> 7.5
80-16-1	0.061	0.041	> 7.5
80-16-2	0.903	-	-
80-18-1	1.39	-	-
80-18-2	0.11	0.044	> 7.5
80-20	0.763	0.235	> 7.5
80-20-1	13.9	-	-
80-20-2	0.368	0.295	> 7.5
80-21	0.25	0.138	> 7.5
80-21-1	0.949	-	-
80-21-2	0.061	0.072	> 7.5
92-1	0.016	0.014	> 7.5
92-2	0.024	0.035	> 7.5
92-3	0.095	0.094	> 7.5
92-4	0.044	0.042	> 7.5
92-5	0.094	0.119	> 7.5

92-6	0.174	0.237	> 7.5
92-7	0.031	0.259	> 7.5
92-8	0.985	-	-
92-9	8.53	-	-
92-10	0.059	0.367	> 7.5
92-11	0.086	0.262	> 7.5
92-12	0.019	0.022	> 7.5
92-13	0.201	0.251	> 7.5
92-14	0.121	0.074	> 7.5
92-15	0.094	0.125	> 7.5
92-16	0.043	0.015	> 7.5
92-17	0.042	0.018	> 7.5
92-18	0.085	0.17	> 7.5
92-19	0.044	0.027	> 7.5
92-20	0.403	-	-
92-21	0.051	0.637	> 7.5
92-22	2.94	-	-
92-23	0.038	0.036	> 7.5
92-24	0.019	0.116	> 7.5
92-25	0.04	0.016	> 7.5
92-26	0.101	0.023	> 7.5
92-27	0.149	0.235	> 7.5
92-28	0.123	0.11	> 7.5
92-29	0.139	10	> 7.5
92-30	0.479	-	-
92-31	0.014	0.022	> 7.5
92-32	0.075	0.129	> 7.5
92-33	0.113	0.216	> 7.5
92-34	0.054	0.06	> 7.5
92-35	0.758	-	---
92-36	0.275	0.133	1.87
92-37	0.048	0.06	> 7.5

92-38	0.042	0.042	> 7.5
92-39	0.032	0.145	> 7.5
92-40	0.244	0.14	> 7.5
92-41	0.117	0.075	3.75
92-42	0.117	0.08	> 7.5
92-43	0.054	0.076	> 7.5
92-44	0.04	0.098	> 7.5
92-45	0.025	0.023	> 7.5
92-46	0.023	0.032	> 7.5
92-47	0.057	0.049	> 7.5
92-48	0.247	0.365	> 7.5
92-49	0.123	0.328	> 7.5
92-50	1.1	-	-
92-51	0.033	0.03	> 7.5
92-52	0.161	0.196	> 7.5
92-53	0.135	0.309	> 7.5
92-54	0.169	0.217	> 7.5
92-55	0.021	0.018	> 7.5
92-56	0.591	-	-
92-57	0.126	0.126	> 7.5
92-58	0.074	0.023	> 7.5
92-59	0.028	0.034	> 7.5
92-60	0.02	0.018	> 7.5
92-61	0.008	0.027	> 7.5
92-62	0.01	0.075	> 7.5
92-63	0.039	0.052	> 7.5
92-64	0.039	0.044	> 7.5
92-65	0.066	0.046	> 7.5
92-66	0.034	0.033	> 7.5
92-67	0.13	0.062	> 7.5
92-68	3.31	-	-
92-69	1.36	-	-

92-70	0.6	-	-
92-71	1.36	-	-
92-72	8.87	-	-
93-1	0.055	0.097	> 7.5
93-2	0.305	0.558	> 7.5
94-1	0.077	0.107	> 7.5
94-2	0.322	0.204	> 7.5
94-3	0.684	-	-
94-4	0.222	0.218	> 7.5
94-5	0.708	0.265	> 7.5
94-6	0.468	-	-
94-7	0.976	-	-
94-8	0.255	0.131	> 7.5
94-9	0.469	-	-
94-10	0.137	0.079	> 7.5
94-11	0.844	-	-
94-12	0.184	0.128	> 7.5
94-13	0.387	-	-
94-14	38.2	-	-
94-15	0.963	-	-
94-16	73.7	-	-
94-17	0.39	-	-
94-18	0.749	-	-
94-19	0.067	0.518	> 7.5
94-20	0.452	-	-
94-21	32.4	-	-
95-1	0.053	0.318	> 7.5
95-2	1.04	-	-
95-3	0.631	-	-
95-4	0.571	-	-
95-5	0.304	0.611	> 7.5
95-6	0.823	-	-

95-7	34	-	-
96-1	0.345	-	-
96-2	1.22	-	-
100-1	0.084	0.123	> 7.5
100-1-1	0.061	0.091	> 7.5
100-1-2	0.062	0.098	> 7.5
100-2	0.324	0.298	> 7.5
100-3	0.763	-	-
100-4	0.006	0.106	> 7.5
100-5	58.3	-	-
100-6	250	-	-
100-7	5.097	-	-
100-8	0.05	0.042	> 7.5
100-9	1.15	-	-
101-1	0.134	0.116	> 7.5
101-2	0.16	0.116	> 7.5
101-3	0.195	0.208	> 7.5
102-1	0.071	0.395	5.245
102-2	1.23	-	-
102-3	0.037	0.032	> 7.5
102-4	0.079	0.262	4.73
102-5	0.071	0.084	> 7.5
102-5-1	0.031	0.058	> 7.5
102-5-2	0.173	0.873	> 7.5
102-6	0.066	0.063	> 7.5
102-6-1	0.094	0.056	> 7.5
102-6-2	0.726	-	-
102-7	0.376	-	-
102-7-1	0.181	1.063	> 7.5
102-7-2	5.28	-	-
102-8	184	-	-
103-1	0.105	0.107	> 7.5

103-1-1	1.26	-	-
103-1-2	0.053	0.039	> 7.5
103-2	0.084	0.04	> 7.5
103-3	0.677	0.143	> 7.5
103-4	0.057	0.062	> 7.5
103-5	0.2	0.507	> 7.5
103-6	0.08	0.474	> 7.5
103-6-1	1.15	-	-
103-6-2	0.112	0.26	5.19
103-7	0.143	0.128	3.75
103-7-1	0.653	-	-
103-7-2	0.088	0.052	> 7.5
106-1	0.08	0.096	4.545
106-2	1.01	5.31	> 7.5
106-3	8.34	10	> 7.5
106-4	5.04	2.45	> 7.5
106-5	6.13	1.46	> 7.5
106-6	0.322	-	-
106-6-1	3.67	-	-
106-6-2	0.059	0.301	> 7.5
106-7	0.301	-	-
106-7-1	1.98	-	-
106-7-2	0.129	0.271	> 7.5
106-8	3.35	1.88	> 7.5
106-9	3.7	2.24	> 7.5
106-10	12.1	-	-
107-1	0.026	0.064	5.57
107-2	0.098	0.299	1.25
107-3	0.114	0.144	3.735
107-4	0.051	0.085	> 7.5
107-5	0.103	0.3	3.255
108-1	0.009	0.039	> 7.5

108-1-1	0.006	0.021	> 7.5
108-1-2	0.008	0.054	> 7.5
108-2	0.049	0.032	> 7.5
108-2-1	0.035	0.046	> 7.5
108-2-2	0.181	0.148	> 7.5
109-1	0.194	0.052	> 7.5
109-1-1	0.06	0.045	> 7.5
109-1-2	0.209	0.08	> 7.5
109-2	0.107	0.098	> 7.5
109-2-1	0.059	0.065	> 7.5
109-2-2	0.226	-	-
110-1	0.57	1.12	> 7.5
110-2	13.4	-	-
119-1	24	-	-
119-2	0.33	-	-
121-1	0.195	0.086	> 7.5
121-2	0.096	0.081	> 7.5
121-3	0.066	0.027	> 7.5
121-4	1.48	-	-
122-1	0.455	-	-
122-2	0.357	1.21	-
122-2-1	12.3	-	-
122-2-2	0.088	0.63	> 7.5
122-3	0.503	-	-
122-3-1	6.82	-	-
122-3-2	0.179	0.145	> 7.5
123-1	1.25	-	-
123-2	0.4	0.26	> 7.5
123-3	14	-	-
123-4	1	-	-
123-5	7.53	10	-
123-6	0.231	0.133	> 7.5

124-1	0.027	0.024	> 7.5
124-2	0.033	0.032	> 7.5
124-3	0.028	0.032	> 7.5
124-4	0.022	0.016	> 7.5
124-5	0.439	-	-
124-6	0.037	0.062	> 7.5
124-7	0.109	0.093	> 7.5
124-8	0.034	0.24	6.355
124-9	0.122	0.169	> 7.5
124-10	0.046	0.049	> 7.5
124-11	0.075	0.098	> 7.5
124-12	0.313	0.231	> 7.5
124-13	0.068	0.094	> 7.5
124-14-1	22.2	-	-
124-14-2	59.5	-	-
124-16-1	1.12	-	-
124-16-2	7.37	-	-
125-1	0.028	0.055	> 7.5
125-2	0.042	0.038	> 7.5
125-3	0.04	0.041	> 7.5
125-4	0.038	0.023	> 7.5
125-5	0.46	-	-
125-6	0.038	0.105	> 7.5
125-7	0.085	0.137	> 7.5
125-8	0.113	0.137	5.05
125-9	0.035	0.048	> 7.5
125-10	0.032	0.068	> 7.5
125-11	0.091	0.187	> 7.5
129-1	0.054	0.124	> 7.5
129-2	0.065	0.027	> 7.5
129-3	0.352	-	-
129-4	0.053	0.058	> 7.5

129-5	0.19	0.087	> 7.5
129-6	0.06	0.058	> 7.5
129-7	0.064	0.025	> 7.5
129-8	0.056	0.023	> 7.5
129-9	0.008	0.048	> 7.5
129-10	0.041	0.049	> 7.5
129-11	0.152	0.247	> 7.5
129-12	0.167	0.39	> 7.5
129-13	0.067	0.054	> 7.5
132-1	0.053	0.46	> 7.5
132-2	0.57	-	-
141-1	0.268	-	-
141-2	0.279	-	-
141-3	0.197	0.331	> 7.5
144-1	0.166	0.23	> 7.5
144-2	24.9	-	-
144-2-1	63.7	-	-
144-2-2	9.13	3.1	4
144-3	0.166	2.41	> 7.5
145-1	2.07	10	5.75
145-1-1	1.71	1.3	5.53
145-1-2	1.15	1.05	3.94
145-2	0.344	1.68	> 7.5
145-2-1	0.294	0.356	> 7.5
145-2-2	0.238	0.213	> 7.5
145-3	3.82	10	4.41
146-1	12.8	-	-
146-1-1	9.43	-	-
146-1-2	53.8	-	-
146-2	0.028	0.038	6.51
146-2-1	0.025	0.024	-
146-2-2	0.061	0.096	5.475

146-3	7.03	4.26	> 7.5
146-3-1	204	-	-
146-3-2	4.4	1.05	> 7.5
146-3-3	57.3	-	-
146-3-4	7.96	1.26	> 7.5
146-4	8.13	4.19	> 7.5
150	0.313	-	-
151	0.348	-	-
152	132	-	-
153	0.38	0.264	> 7.5
154	0.062	0.049	6.96
155	0.104	0.063	2.59
156	0.369	0.148	> 7.5
157	0.064	0.07	> 7.5
158	0.189	0.191	3.835
159	0.046	0.119	3.035
160	0.081	0.058	> 7.5
161	0.294	0.061	> 7.5
162	0.133	0.078	> 7.5
163	0.127	0.055	> 7.5
164-1	0.066	0.198	6.05
164-2	3.44	-	-
165-1	0.073	0.049	> 7.5
165-2	1.61	-	-
166-1	0.049	0.118	> 7.5
166-2	1.032	-	-
167	0.052	0.083	> 7.5
168	0.096	0.198	4.345
169	0.55	-	-
170	0.064	0.077	> 7.5
171	0.206	-	-
172	0.08	0.099	> 7.5

173-1	0.128	0.229	5.97
173-2	5.25	-	-
174	0.069	0.082	> 7.5
175-1	1.16	-	-
175-2	0.071	0.171	> 7.5
176-1	0.131	0.347	5.755
176-2	2.43	-	-
177-1	0.031	0.063	3.34
177-2	1.26	-	-
178	3.77	-	-
179-1	0.048	0.049	> 7.5
179-2	2.42	-	-
179	0.069	0.069	> 7.5
180	85.1	10	-
181	4.53	10	> 7.5
182-1	0.14	0.319	6.62
182-2	0.187	0.365	5.62
182	8.794	0.239	3.52
183-1	0.61	-	-
183-2	0.058	0.043	> 7.5
184	2.81	-	-
185	0.055	0.08	> 7.5
186	0.093	0.199	> 7.5
187	0.09	0.081	0.317
188	0.038	0.04	4.975
189-1	0.036	0.072	> 7.5
189-2	14.7	-	-
189	0.158	0.209	> 7.5
190	0.126	1.705	> 7.5
191-1	2.99	-	-
191-2	0.079	0.079	> 7.5
192	0.029	0.161	> 7.5

193	0.056	0.055	> 7.5
194-1	0.067	0.033	> 7.5
194-2	1.45	-	-
194	0.186	0.048	> 7.5
195	0.488	-	-
196	0.048	0.097	> 7.5
197-1	1.59	-	-
197-2	0.01	0.091	> 7.5
198	0.075	0.136	5.96
199	0.132	0.087	> 7.5
200	0.03	0.094	> 7.5
201	0.075	0.205	5.05
202	0.071	0.075	> 7.5
203	0.223	0.243	> 7.5
204	0.066	0.091	3.01
205-1	0.098	0.07	> 7.5
205-2	0.078	0.091	> 7.5
205	0.08	0.095	> 7.5
206-1	0.143	0.1	> 7.5
206-2	0.098	0.139	> 7.5
207	0.932	-	-
208-1	2.56	-	-
208-2	0.117	0.106	> 7.5
208	0.231	0.14	> 7.5
209-1	0.088	0.116	> 7.5
209-2	1.34	-	-
209-3	0.115	0.107	> 7.5
210	0.049	0.039	> 7.5
211-1	0.113	0.128	> 7.5
211-2	0.05	0.106	> 7.5
211-3	0.084	0.253	> 7.5
212-1	0.256	0.705	> 7.5

212-2	0.104	0.237	> 7.5
212-3	0.703	-	-
213-1	1.5	-	-
213-2	0.072	0.069	> 7.5
213-3	0.066	0.074	> 7.5
213-4	1.51	-	-
213	0.182	0.104	> 7.5
214-1	0.025	0.03	> 7.5
214-2	2.84	-	-
214	0.02	0.025	> 7.5
215-1	3.79	-	-
215-2	0.055	0.086	> 7.5
215	0.076	0.275	-
216	2.01	-	-
217	1.56	0.972	> 7.5
218	0.094	0.035	> 7.5
219	250	-	-
220	250	-	-
221	1.26	-	-
222-1	0.95	-	-
222-2	0.014	0.048	2.52
222	0.205	0.057	2.71
223	0.18	0.659	> 7.5
224-1	23.3	-	-
224-2	1.29	-	-
224	7.62	-	-
225-1	7.62	-	-
225-2	0.501	-	-
225	4.43	-	-
226	0.04	0.063	> 7.5
227	3.57	-	-
228	0.769	-	-

229	0.071	0.032	> 7.5

(-) denotes not tested

[0799] The present invention is described in connection with preferred embodiments. However, it should be appreciated that the invention is not limited to the disclosed embodiments. It is understood that, given the description of the embodiments of the invention herein, various modifications can be made by a person skilled in the art. Such modifications are encompassed by the claims below.

2019336588 14 Jun 2022

What is claimed is:

1. A compound having a structure selected from the formula


;


;


;


;


;


;


;


;

2019336588 14 Jun 2022

	 ;
	 ;
 ;	 ;
	 ;
 ;	 ;

2019336588 14 Jun 2022

	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

 ;	 ;
	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
	 ;

2019336588 14 Jun 2022

 ;	 ;
	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
	 ;

2019336588 14 Jun 2022

 ;	 ;
	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ;
 ;	 ; and
	

or a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof, or a pharmaceutically acceptable salt of the atropisomer thereof.

2. A compound having a structure selected from the formula

 ;	 ;

 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;
 ;	 ;

2019336588 14 Jun 2022

 ;	 ; and
	

or a stereoisomer thereof, an atropisomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable salt of the stereoisomer thereof, or a pharmaceutically acceptable salt of the atropisomer thereof.

3. The compound of any one of claims 1-2 in the form of a pharmaceutically acceptable salt.

4. A pharmaceutical composition comprising the compound of any one of claims 1-3 and a pharmaceutically acceptable excipient.

5. A method of inhibiting KRAS G12C in a cell, comprising contacting the cell with the compound of any one of claims 1-3 or the composition of claim 4.

6. Use of the compound of any one of claims 1-3 or the composition of claim 4 in the manufacture of a medicament for inhibiting KRAS G12C in a cell.

7. A method of treating cancer in a subject comprising administering to the subject a therapeutically effective amount of the compound of any one of claims 1-3 or the composition of claim 4, wherein the cancer is mediated by a KRAS G12C mutation.

2019336588 14 Jun 2022

8. Use of the compound of any one of claims 1-3 or the composition of claim 4 in the manufacture of a medicament for treating cancer in a subject, wherein the cancer is mediated by a KRAS G12C mutation.

9. The method of claim 7 or the use of claim 8, wherein the cancer is non-small cell lung cancer, small intestine cancer, appendix cancer, colorectal cancer, endometrial cancer, pancreatic cancer, skin cancer, gastric cancer, nasal cavity cancer, or bile duct cancer.

10. The method or use of claim 9, wherein the cancer is non-small cell lung cancer.

11. The method or use of claim 9, wherein the cancer is pancreatic cancer.

12. The method or use of claim 9, wherein the cancer is colorectal cancer.

Amgen Inc.
Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON

SEQUENCE LISTING

<110> ALLEN, John Gordon
LANMAN, Brian Alan
CHEN, Jian
REED, Anthony B.
CEE, Victor J.
LIU, Longbin
LOPEZ, Patricia
WURZ, Ryan Paul
NGUYEN, Thomas T.
BOOKER, Shon
ALLEN, Jennifer Rebecca
CHU-MOYER, Margaret
AMEGADZIE, Albert
CHEN, Ning
GOODMAN, Clifford
LOW, Jonathan D.
MA, Vu Van
MINATTI, Ana Elena
NISHIMURA, Nobuko
PICKRELL, Alexander J.
WANG, Hui-Ling
SHIN, Youngsook
SIEGMUND, Aaron C.
YANG, Kevin C.
TAMAYO, Nuria A.
WALTON, Mary
XUE, Qiufen

<120> INHIBITORS OF KRAS G12C AND METHODS OF USING THE SAME

<130> A-2263-WO-PCT

<140> PCT/USXXXX/XXXXXX
<141> 2019-06-11

<150> 62/684,117
<151> 2018-06-12

<160> 20

<170> PatentIn version 3.5

<210> 1

<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 1

Ser Tyr Asp Met Ser
1 5

<210> 2
<211> 16
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 2

Leu Ile Ser Gly Gly Gly Ser Gln Thr Tyr Tyr Ala Glu Ser Val Lys
1 5 10 15

<210> 3
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 3

Pro Ser Gly His Tyr Phe Tyr Ala Met Asp Val
1 5 10

<210> 4
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 4

Arg Ala Ser Gln Gly Ile Ser Asn Trp Leu Ala
1 5 10

<210> 5
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 5

Ala Ala Ser Ser Leu Gln Ser
1 5

<210> 6
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 6

Gln Gln Ala Glu Ser Phe Pro His Thr
1 5

<210> 7
<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 7

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Asp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ser Leu Ile Ser Gly Gly Gly Ser Gln Thr Tyr Tyr Ala Glu Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Phe Cys
85 90 95

Ala Ser Pro Ser Gly His Tyr Phe Tyr Ala Met Asp Val Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 8
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 8

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Asn Trp
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Phe Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Glu Ser Phe Pro His
85 90 95

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 9
<211> 472
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 9

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp
1 5 10 15

Leu Arg Gly Ala Arg Cys Glu Val Gln Leu Leu Glu Ser Gly Gly Gly
20 25 30

Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly
35 40 45

Phe Thr Phe Ser Ser Tyr Asp Met Ser Trp Val Arg Gln Ala Pro Gly
50 55 60

Lys Gly Leu Glu Trp Val Ser Leu Ile Ser Gly Gly Gly Ser Gln Thr
65 70 75 80

Tyr Tyr Ala Glu Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn
85 90 95

Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
100 105 110

Thr Ala Val Tyr Phe Cys Ala Ser Pro Ser Gly His Tyr Phe Tyr Ala
115 120 125

Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser
130 135 140

Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr
145 150 155 160

Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro
165 170 175

Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val
180 185 190

His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser
195 200 205

Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile
210 215 220

Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val
225 230 235 240

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
245 250 255

Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
260 265 270

Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
275 280 285

Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
290 295 300

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Cys Glu Glu Gln
305 310 315 320

Tyr Gly Ser Thr Tyr Arg Cys Val Ser Val Leu Thr Val Leu His Gln
325 330 335

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
340 345 350

Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
355 360 365

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
370 375 380

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser

385 390 395 400

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
405 410 415

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
420 425 430

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
435 440 445

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
450 455 460

Ser Leu Ser Leu Ser Pro Gly Lys
465 470

<210> 10
<211> 236
<212> PRT
<213> Artificial Sequence

<220>
<223> Synthetic peptide

<400> 10

Met Asp Met Arg Val Pro Ala Gln Leu Leu Gly Leu Leu Leu Leu Trp
1 5 10 15

Leu Arg Gly Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
20 25 30

Val Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser
35 40 45

Gln Gly Ile Ser Asn Trp Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys
50 55 60

Ala Pro Lys Leu Leu Ile Phe Ala Ala Ser Ser Leu Gln Ser Gly Val
65 70 75 80

Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
85 90 95

Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
100 105 110

Ala Glu Ser Phe Pro His Thr Phe Gly Gly Gly Thr Lys Val Glu Ile
115 120 125

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp
130 135 140

Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn
145 150 155 160

Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu
165 170 175

Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp
180 185 190

Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr
195 200 205

Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser
210 215 220

Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

225 230 235

<210> 11
<211> 15
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic polynucleotide

<400> 11
agctatgaca tgagc
15

<210> 12
<211> 51
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic polynucleotide

<400> 12
cttattagtg gtggtggtag tcaaacatac tacgcagaat ccgtgaaggg c
51

<210> 13
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic polynucleotide

<400> 13
cccagtggcc actacttcta cgctatggac gtc
33

<210> 14
<211> 33
<212> DNA

<213> Artificial Sequence

<220>
<223> Synthetic polynucleotide

<400> 14
cgggcgagtc agggtattag caactggtta gcc
33

<210> 15
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic polynucleotide

<400> 15
gctgcatcca gtttgcaaag t
21

<210> 16
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic polynucleotide

<400> 16
caacaggctg aaagtttccc tcacact
27

<210> 17
<211> 360
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic polynucleotide

<400> 17

gaggtgcagc tgttggagtc tgggggaggc ttggtacagc ctgggggtc cctgagactc 60

tcctgtgcag cctctggatt cacctttagc agctatgaca tgagctgggt ccgccaggct 120

ccagggaagg ggctggaatg ggtctcactt attagtggtg gtggtagtca aacatactac 180

gcagaatccg tgaagggccg gttcaccatc tccagagaca attccaagaa cacgctgtat 240

ctgcaaatga acagcctgag agccgaggac acggccgtat atttctgtgc gtcccccagt 300

ggccactact tctacgctat ggacgtctgg ggccaaggga ccacggtcac cgtctcctca 360

<210> 18
<211> 321
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic polynucleotide

<400> 18
gacatccaga tgacccagtc tccatcttcc gtgtctgcat ctgttggaga cagagtcacc 60

atcacttgtc gggcgagtca gggtattagc aactggttag cctggtatca gcagaaacca 120

gggaaagccc ctaagctcct gatctttgct gcatccagtt tgcaaagtgg ggtcccatca 180

aggttcagcg gcagtggatc tgggacagat ttcaccctca ccatcagcag cctgcagcct 240

gaagattttg caacttacta ttgtcaacag gctgaaagtt tccctcacac tttcggcgga 300

g g g a c c a a g g t g g a g a t c a a a 321

<210> 19
<211> 1416
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic polynucleotide

<400> 19
atggacatga gggtgcccgc tcagctcctg gggctcctgc tgctgtggct gagaggtgcg
60

cgctgtgagg tgcagctgtt ggagtctggg ggaggcttgg tacagcctgg ggggtccctg
120

agactctcct gtgcagcctc tggattcacc tttagcagct atgacatgag ctgggtccgc
180

caggctccag ggaaggggct ggaatgggtc tcacttatta gtggtggtgg tagtcaaaca
240

tactacgcag aatccgtgaa gggccggttc accatctcca gagacaattc caagaacacg
300

ctgtatctgc aaatgaacag cctgagagcc gaggacacgg ccgtatattt ctgtgcgtcc
360

cccagtggcc actacttcta cgctatggac gtctggggcc aagggaccac ggtcaccgtc
420

tcctcagcct ccaccaaggg cccatcggtc ttcccctgg caccctcctc caagagcacc
480

tctgggggca cagcggccct gggctgcctg gtcaaggact acttccccga accggtgacg
540

gtgtcgtgga actcaggcgc cctgaccagc ggcgtgcaca ccttcccggc tgtcctacag
600

tcctcaggac tctactccct cagcagcgtg gtgaccgtgc cctccagcag cttgggcacc
660

cagacctaca tctgcaacgt gaatcacaag cccagcaaca ccaaggtgga caagaaagtt
720

gagcccaaat cttgtgacaa aactcacaca tgcccaccgt gcccagcacc tgaactcctg 780

gggggaccgt cagtcttcct cttcccccca aaacccaagg acaccctcat gatctcccgg 840

acccctgagg tcacatgcgt ggtggtggac gtgagccacg aagaccctga ggtcaagttc 900

aactggtacg tggacggcgt ggaggtgcat aatgccaaga caaagccgtg cgaggagcag 960

tacggcagca cgtaccgttg cgtcagcgtc ctcaccgtcc tgcaccagga ctggctgaat 1020

ggcaaggagt acaagtgcaa ggtgtccaac aaagccctcc cagccccat cgagaaaacc 1080

atctccaaag ccaaagggca gccccgagaa ccacaggtgt acaccctgcc cccatcccgg 1140

gaggagatga ccaagaacca ggtcagcctg acctgcctgg tcaaaggctt ctatcccagc 1200

gacatcgccg tggagtggga gagcaatggg cagccggaga acaactacaa gaccacgcct 1260

cccgtgctgg actccgacgg ctccttcttc ctctatagca agctcaccgt ggacaagagc 1320

aggtggcagc aggggaacgt cttctcatgc tccgtgatgc atgaggctct gcacaaccac 1380

tacacgcaga agagcctctc cctgtctccg ggtaaa 1416

<210> 20
<211> 708
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic polynucleotide

<400> 20
atggacatga gggtgcccgc tcagctcctg gggctcctgc tgctgtggct gagaggtgcg
60

cgctgtgaca tccagatgac ccagtctcca tcttccgtgt ctgcatctgt tggagacaga
120

gtcaccatca cttgtcgggc gagtcagggt attagcaact ggttagcctg gtatcagcag
180

aaaccaggga aagcccctaa gctcctgatc tttgctgcat ccagtttgca aagtggggtc
240

ccatcaaggt tcagcggcag tggatctggg acagatttca ccctcaccat cagcagcctg
300

cagcctgaag attttgcaac ttactattgt caacaggctg aaagtttccc tcacactttc
360

ggcggaggga ccaaggtgga gatcaaacga acggtggctg caccatctgt cttcatcttc
420

ccgccatctg atgagcagtt gaaatctgga actgcctctg ttgtgtgcct gctgaataac
480

ttctatccca gagaggccaa agtacagtgg aaggtggata acgccctcca atcgggtaac
540

tcccaggaga gtgtcacaga gcaggacagc aaggacagca cctacagcct cagcagcacc
600

ctgacgctga gcaaagcaga ctacgagaaa cacaaagtct acgcctgcga agtcacccat
660

cagggcctga gctcgcccgt cacaaagagc ttcaacaggg gagagtgt
708