

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number

WO 2017/192867 A1

(43) International Publication Date
09 November 2017 (09.11.2017)

(51) International Patent Classification:

A61K 38/00 (2006.01) *C07K 16/24* (2006.01)
A61K 39/395 (2006.01) *C07K 16/28* (2006.01)

Published:

— with international search report (Art. 21(3))

(21) International Application Number:

PCT/US2017/031089

(22) International Filing Date:

04 May 2017 (04.05.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/331,813 04 May 2016 (04.05.2016) US

(71) Applicants: **MILLENNIUM PHARMACEUTICALS, INC.** [US/US]; 40 Landsdowne Street, Cambridge, Massachusetts 02139 (US). **ICAHN SCHOOL OF MEDICINE AT MOUNT SINAI** [US/US]; One Gustave L. Levy Place, Box 1675, New York, New York 10029 (US).



(72) Inventors: **LISSOOS, Trevor Warren**; c/o Takeda Pharmaceuticals North America, Inc., One Takeda Parkway, Deerfield, Illinois 60015 (US). **COLOMBEL, Jean-Fredéric**; c/o Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, Box 1675, New York, New York 10029 (US).

(74) Agent: **CONNARN, Kristin A.** et al.; McDermott Will & Emery LLP, 500 North Capitol Street, N.W., Washington, District of Columbia 20001-1531 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: TRIPLE COMBINATION THERAPY FOR TREATING INFLAMMATORY BOWEL DISEASE

(57) Abstract: The invention features new a method for treating inflammatory bowel disease, including Crohn's Disease (CD) or ulcerative colitis in a human patient, comprising administering to a patient a combination therapy comprising an anti- $\alpha 4\beta 7$ antibody, a TNF α antagonist, and an immunomodulator.

TRIPLE COMBINATION THERAPY FOR TREATING
INFLAMMATORY BOWEL DISEASE

RELATED APPLICATIONS

5 This application claims the benefit of U.S. Provisional Application No. 62/331,813 filed on May 4, 2016. The entire contents of the foregoing application are hereby incorporated by reference.

BACKGROUND

10 Inflammatory bowel disease (IBD) is a chronic, relapsing, inflammatory disorder of the gastrointestinal (GI) tract that includes 2 entities, namely ulcerative colitis (UC) and Crohn's disease (CD). In contrast to the diffuse, superficial, continuous inflammation limited to the colon in UC, the inflammation of CD is focal, may be transmural, and can involve any segment of the GI tract from mouth to anus. The prevalence of CD is approximately 150/100,000 of the

15 United States (US) population. The characteristic pathology involves a chronic inflammatory infiltrate consisting of neutrophils and macrophages. Hallmarks of CD include granulomatous inflammation and aphthous ulceration. CD is neither medically or surgically curable at the current time.

Untreated CD patients commonly develop structural complications that include stricture and fistula formation, which commonly result in hospitalizations and surgery over time. When examining 306 CD patients from the US, Thia et al. (2010, *Gastroenterology* 139:1147-55) showed that the cumulative risk of developing CD complications of either stricturing or penetrating disease was 33.7% at 5 years and 50.8% at 20 years after establishment of CD diagnosis. The natural course of the disease can range from an indolent course with prolonged periods of remission to aggressive, incapacitating disease associated with structural bowel wall damage. Certain patients have aggressive CD, which could be defined as having a high relapse rate, development of penetrating disease, need for repeat surgery, or multiple admissions for flares. The pathophysiology of IBD is complex as multiple environmental factors interact with the genotype of an individual to cause expression of disease. Published risk factors for aggressive disease include involvement of the upper GI tract and ileum, penetrating disease, early age at diagnosis, smoking, extensive ulceration of the mucosa, high titers of certain serum antibodies, and mutations of the nucleotide-binding oligomerization domain-containing protein 2 (NOD2) gene. The ability to more readily predict which patients will be at higher risk for complications and target more intensive, early treatment to that

group would be invaluable. The Crohn's Disease Personalized Risk and Outcome Prediction Tool (PROSPECT), which incorporates clinical, serological, and genetic variables, has been developed by Siegel et al. (*Aliment. Pharmacol. Ther.* 2016, 43:262-71; *Inflamm. Bowel Dis.* 2011, 17:30-38). This tool is used to predict the probability of developing complications

5 related to CD, specifically strictures or internal penetrating disease. The output of the predictive model allows risk stratification of patients from low to medium to high risk of complications within a 3-year period.

Treatment with TNF-antagonist therapy substantially improved the care of patients with CD refractory to corticosteroids, immunomodulators, and mesalamine. In 2010, the SONIC trial 10 demonstrated that combination therapy with the TNF antagonist infliximab and immunomodulator azathioprine further significantly improved treatment efficacy. About 40% of patients, however, will not achieve remission after initiation of a combined TNF antagonist and immunomodulator. Thus, there remains an urgent unmet medical need for a combination therapy that can improve remission rates and modify disease progression, 15 particularly for patients who are at higher risk for CD complications.

SUMMARY OF THE INVENTION

The invention relates to methods of treating patient having inflammatory bowel disease (e.g., Crohn's disease (CD) or ulcerative colitis (UC)) using combination therapies 20 that can improve remission rates and modify disease progression.

In one aspect, the patient is at moderate to high risk of complication from the inflammatory bowel disease. In an embodiment, the patient is a high risk CD patient.

In one aspect, the invention provides a method of treating a patient having Crohn's disease using a triple combination therapy comprising a humanized anti- $\alpha 4\beta 7$ antibody, 25 adalimumab, and an immunomodulator, and the triple combination therapy is administered to the patient according to the following dosing regimen: an initial dose of 300 mg of humanized anti- $\alpha 4\beta 7$ antibody as an intravenous infusion, followed by a second subsequent dose of 300 mg of humanized anti- $\alpha 4\beta 7$ antibody as an intravenous infusion at about two weeks after the initial dose; followed by a third subsequent dose of 300 mg of the humanized 30 anti- $\alpha 4\beta 7$ antibody as an intravenous infusion at about six weeks after the initial dose; an initial dose of 160 mg of adalimumab as a subcutaneous injection, followed by a second subsequent dose of 80 mg of adalimumab as a subcutaneous injection, followed by a third subsequent dose of 40 mg of adalimumab at about four weeks after the initial dose; and an initial dose of 15 mg of immunomodulator orally; wherein the dosing regimen results in

endoscopic remission and further wherein the anti- α 4 β 7 antibody comprises an antigen binding region of nonhuman origin and at least a portion of an antibody of human origin, the humanized antibody has binding specificity for the α 4 β 7 complex, and the antigen-binding region comprises the CDRs:

5 Light chain: CDR1 SEQ ID NO:7
 CDR2 SEQ ID NO:8 and
 CDR3 SEQ ID NO:9; and
 Heavy chain: CDR1 SEQ ID NO:4
 CDR2 SEQ ID NO:5 and
10 CDR3 SEQ ID NO:6.

In one embodiment, the humanized anti- α 4 β 7 antibody is administered to the patient over about 30 minutes.

In one embodiment, the humanized anti- α 4 β 7 antibody has a heavy chain variable region sequence of amino acids 20 to 140 of SEQ ID NO:1. In one embodiment, the 15 humanized anti- α 4 β 7 antibody has a light chain variable region sequence of amino acids 20 to 131 of SEQ ID NO:2. In yet another embodiment, the humanized anti- α 4 β 7 antibody has a heavy chain comprising amino acids 20 to 470 of SEQ ID NO:1 and a light chain comprising amino acids 20 to 238 of SEQ ID NO:2. In one embodiment, the humanized anti- α 4 β 7 antibody is vedolizumab.

20 In one embodiment, the immunomodulator is methotrexate.

In one aspect, the method further comprises administering a subsequent dose of 300 mg of the humanized anti- α 4 β 7 antibody as an intravenous infusion at about eight weeks after the third subsequent dose.

25 In one aspect, the method further comprises administering a subsequent dose of 108 mg of the humanized anti- α 4 β 7 antibody as subcutaneous injection at about every two, three or four weeks after the third subsequent dose.

In one aspect, concomitant corticosteroid use by the patient is tapered to discontinuation during the triple combination therapy.

30 In one aspect, the invention provides a method of treating a patient having an inflammatory bowel disease, comprising administering to the patient vedolizumab, adalimumab, and methotrexate according to a triple combination therapy wherein the triple combination therapy comprises administering vedolizumab at a dose of 300 mg at weeks 0, 2, and 6, followed by administration every 8 weeks thereafter; subcutaneously administering a 160 mg dose of adalimumab at week 0, an 80 mg dose of adalimumab at week 2, and a 40 mg

dose of adalimumab at week 4 and every two weeks thereafter; and administering methotrexate. In one embodiment, adalimumab is discontinued at week 26. In one embodiment, methotrexate is administered orally. In one embodiment, methotrexate is administered at a dose of 15 mg. In one embodiment, methotrexate is discontinued at week

5 34. In one embodiment, the inflammatory bowel disease is Crohn's disease. In one embodiment, the inflammatory bowel disease is moderately to severely active Crohn's disease. In one embodiment, the patient is a high risk CD patient. In one embodiment, the patient is selected as a high risk CD patient prior to administration of the triple combination therapy. In one embodiment, the inflammatory bowel disease is ulcerative colitis. In one embodiment the ulcerative colitis is moderately to severely active ulcerative colitis.

10 In one aspect, the invention provides a method of treating a patient having an inflammatory bowel disease, comprising administering to the patient vedolizumab, infliximab, and methotrexate according to a triple combination therapy. The triple combination therapy can comprise administering vedolizumab at a dose of 300 mg at weeks 15 0, 2, and 6, followed by administration every 8 weeks thereafter; intravenously administering infliximab at a dose of 5mg/kg at weeks 0, 2 and 6, and then every eight weeks thereafter; and administering methotrexate. In one embodiment, methotrexate is administered orally. In one embodiment, methotrexate is administered at a dose of 15 mg. In one embodiment, methotrexate is discontinued at week 34. In one embodiment, the inflammatory bowel 20 disease is Crohn's disease. In one embodiment, the inflammatory bowel disease is moderately to severely active Crohn's disease. In one embodiment, the patient is a high risk CD patient. In one embodiment, the patient is selected as a high risk CD patient prior to administration of the triple combination therapy. In one embodiment, the inflammatory bowel disease is ulcerative colitis. In one embodiment, the ulcerative colitis is moderately to 25 severely active ulcerative colitis.

25 In one aspect, the invention provides a method of treating a patient having an inflammatory bowel disease, comprising administering to the patient vedolizumab, golimumab, and methotrexate according to a triple combination therapy. The triple combination therapy can comprise administering vedolizumab at a dose of 300 mg at weeks 30 0, 2, and 6, followed by administration every 8 weeks thereafter; subcutaneously administering golimumab at a dose of 200 mg at week 0, 100 mg at week 2, and 100 mg every four weeks thereafter; and administering methotrexate. In one embodiment, methotrexate is administered orally. In one embodiment, methotrexate is administered at a dose of 15 mg. In one embodiment, methotrexate is discontinued at week 34. In one

embodiment, the inflammatory bowel disease is Crohn's disease. In one embodiment, the inflammatory bowel disease is moderately to severely active Crohn's disease. In one embodiment, the patient is a high risk CD patient. In one embodiment, the patient is selected as a high risk CD patient prior to administration of the triple combination therapy. In one embodiment, the inflammatory bowel disease is ulcerative colitis. In one embodiment, the ulcerative colitis is moderately to severely active ulcerative colitis.

In one aspect, the invention provides a method of treating a patient having an inflammatory bowel disease, comprising administering to the patient vedolizumab, certolizumab pegol, and methotrexate according to a triple combination therapy. The triple combination therapy can comprise administering vedolizumab at a dose of 300 mg at weeks 0, 2, and 6, followed by administration every 8 weeks thereafter; subcutaneously administering certolizumab at a dose of 400 mg at week 0, 400 mg at weeks 2 and 4, and optionally 400 mg every four weeks thereafter; and administering methotrexate. In one embodiment, methotrexate is administered orally. In one embodiment, methotrexate is administered at a dose of 15 mg. In one embodiment, methotrexate is discontinued at week 34. In one embodiment, the inflammatory bowel disease is Crohn's disease. In one embodiment, the inflammatory bowel disease is moderately to severely active Crohn's disease. In one embodiment, the patient is a high risk CD patient. In one embodiment, the patient is selected as a high risk CD patient prior to administration of the triple combination therapy. In one embodiment, the inflammatory bowel disease is ulcerative colitis. In one embodiment, the ulcerative colitis is moderately to severely active ulcerative colitis.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a schematic illustrating an overview of the study design.

25

DETAILED DESCRIPTION

The present invention relates to a method of treating inflammatory bowel disease (e.g., Crohn's Disease or ulcerative colitis) with a triple combination therapy. The method comprises administering a triple combination therapy, comprising an anti-integrin $\alpha 4\beta 7$ (e.g., vedolizumab), a TNF α antagonist (e.g., adalimumab), and an immunomodulator (e.g., methotrexate), to a patient with Crohn's Disease, particularly a patient at high risk for developing complications.

In one aspect, the patient is administered the triple combination therapy through one

or more administration routes. In some embodiments, the anti-integrin is administered as an infusion, the TNF antagonist as a subcutaneous injection, and the immunomodulator orally.

In some embodiments, the anti-integrin component of the triple combination therapy is an antibody, such as an anti- $\alpha 4\beta 7$ antibody. In some embodiments, the anti- $\alpha 4\beta 7$ antibody 5 is a humanized antibody, e.g., a humanized antibody with the epitopic specificity of Act-1 mouse monoclonal antibody. In some embodiments, the anti- $\alpha 4\beta 7$ antibody is vedolizumab.

In some embodiments, the TNF antagonist is adalimumab. In some embodiments, the TNF antagonist is infliximab, golimumab, or certolizumab pegol.

In some embodiments, the immunomodulator is methotrexate. In some embodiments, 10 the immunomodulator is azathioprine or 6-mercaptopurine.

Definitions

The term “combination therapy”, as used herein, refers to the administration of two or more therapeutic substances, e.g., an anti- $\alpha 4\beta 7$ antibody and another drug. The other drug(s) may be administered concomitant with, prior to, or following the administration of an 15 anti- $\alpha 4\beta 7$ antibody. The term “triple combination therapy” as used herein, refers to the administration of at least three therapeutic substances, e.g., an anti- $\alpha 4\beta 7$ antibody, an anti-TNF antibody, and methotrexate.

The terms “TNF α antagonist” or “TNF antagonist”, as used interchangeably herein, refer to a therapeutic agent that inhibits, antagonizes, and/or neutralizes 20 TNF α activity. Examples of a TNF antagonist include, but are not limited to, adalimumab, infliximab, golimumab, certolizumab pegol, and etanercept.

The term “high risk Crohn’s Disease (CD) patient” refers to a patient having Crohn’s Disease who has at least a 20% chance of a disease complication by year 2 following 25 diagnosis as determined by the Crohn’s Disease Personalized Risk and Outcome Prediction Tool (PROSPECT) predictive tool (Siegel, et al., *Aliment. Pharmacol. Ther.* 2016; 43:262-271). In one embodiment, the methods disclosed herein are used to treat a patient in this particular subpopulation.

Alternatively, the term “high risk Crohn’s Disease (CD) patient” refers to a patient having Crohn’s Disease who has a risk of complications as defined by the 2014 American 30 Gastroenterological Association (AGA) CD Clinical Care Pathway Criteria (Sandborn, *Gastroenterology* 2014, 147:702-705). The criteria assess the severity of the current and prior disease burden, such as initial diagnosis <30 years of age, extensive anatomic involvement, perianal and/or severe rectal disease, deep ulcers, prior surgical resection or

stricturing and/or penetrating behavior. In one embodiment, the methods disclosed herein are used to treat a patient whose CD has at least one of these criteria. In one embodiment, the methods disclosed herein are not used to treat a patient who has had prior surgical resection.

“Baseline” as used herein describes a value of a parameter which is measured prior to the initial dose of a treatment. It can refer to a measurement of a sample obtained the same day, the day before, during the week before initial treatment, i.e., at a time period before the first dose when little change is expected until after the first dose and values of the measurement obtained after the first dose can be compared to this baseline value to represent the change caused by the dose.

10 The term “pharmaceutical formulation” refers to a preparation that contains an $\alpha 4\beta 7$ antagonist, such as an anti- $\alpha 4\beta 7$ antibody, in such form as to permit the biological activity of the antibody to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered.

15 The cell surface molecule, “ $\alpha 4\beta 7$ integrin,” or “ $\alpha 4\beta 7$,” is a heterodimer of an α_4 chain (CD49D, ITGA4) and a β_7 chain (ITGB7). Each chain can form a heterodimer with an alternative integrin chain, to form $\alpha_4\beta_1$ or $\alpha_E\beta_7$. Human α_4 and β_7 genes (GenBank (National Center for Biotechnology Information, Bethesda, MD) RefSeq Accession numbers NM_000885 and NM_000889, respectively) are expressed by B and T lymphocytes, particularly memory CD4+ lymphocytes. Typical of many integrins, $\alpha 4\beta 7$ can exist in either 20 a resting or activated state. Ligands for $\alpha 4\beta 7$ include vascular cell adhesion molecule (VCAM), fibronectin and mucosal addressin (MAdCAM (e.g., MAdCAM-1)).

25 An “ $\alpha 4\beta 7$ antagonist” is a molecule which antagonizes, reduces or inhibits the function of $\alpha 4\beta 7$ integrin. Such antagonist may antagonize the interaction of $\alpha 4\beta 7$ integrin with one or more of its ligands. An $\alpha 4\beta 7$ antagonist may bind either chain of the heterodimer or a complex requiring both chains of the $\alpha 4\beta 7$ integrin, or it may bind a ligand, such as MAdCAM. An $\alpha 4\beta 7$ antagonist may be an antibody which performs such binding function, such as an anti- $\alpha 4\beta 7$ -integrin antibody or “anti- $\alpha 4\beta 7$ antibody”. In some embodiments, an $\alpha 4\beta 7$ antagonist, such as an anti- $\alpha 4\beta 7$ antibody, has “binding specificity for the $\alpha 4\beta 7$ complex” and binds to $\alpha 4\beta 7$, but not to $\alpha 4\beta 1$ or $\alpha E\beta 7$.

30 The term “antibody” or “antibodies” herein is used in the broadest sense and specifically covers full length antibody, antibody peptide(s) or immunoglobulin(s), monoclonal antibodies, chimeric antibodies (including primatized antibodies), polyclonal antibodies, human antibodies, humanized antibodies and antibodies from non-human species,

including human antibodies derived from a human germline immunoglobulin sequence transduced into the non-human species, e.g., mouse, sheep, chicken or goat, recombinant antigen binding forms such as monobodies and diabodies, multispecific antibodies (e.g. bispecific antibodies) formed from at least two full length antibodies (e.g., each portion 5 comprising the antigen binding region of an antibody to a different antigen or epitope), and individual antigen binding fragments of any of the foregoing, e.g., of an antibody or the antibody from which it is derived, including dAbs, Fv, scFv, Fab, F(ab')₂, Fab'.

The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies 10 comprising the population are identical and/or bind the same epitope. The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method.

“Antigen binding fragments” of an antibody preferably comprise at least the variable 15 regions of the heavy and/or light chains of an anti- $\alpha 4\beta 7$ antibody. For example, an antigen binding fragment of vedolizumab can comprise amino acid residues 20-131 of the humanized light chain sequence of SEQ ID NO:2 and amino acid residues 20-140 of the humanized heavy chain sequence of SEQ ID NO:1. Examples of such antigen binding fragments include Fab fragments, Fab' fragments, Fv fragments, scFv and F(ab')₂ fragments. Antigen binding 20 fragments of an antibody can be produced by enzymatic cleavage or by recombinant techniques. For instance, papain or pepsin cleavage can be used to generate Fab or F(ab')₂ fragments, respectively. Antibodies can also be produced in a variety of truncated forms using antibody genes in which one or more stop codons have been introduced upstream of the natural stop site. For example, a recombinant construct encoding the heavy chain of an 25 F(ab')₂ fragment can be designed to include DNA sequences encoding the CH_I domain and hinge region of the heavy chain. In one aspect, antigen binding fragments inhibit binding of $\alpha 4\beta 7$ integrin to one or more of its ligands (e.g. the mucosal addressin MAdCAM (e.g., MAdCAM-1), fibronectin).

A “therapeutic monoclonal antibody” is an antibody used for therapy of a human 30 subject. Therapeutic monoclonal antibodies disclosed herein include anti- $\alpha 4\beta 7$ antibodies.

Antibody “effector functions” refer to those biological activities attributable to the Fc region (a native sequence Fc region or amino acid sequence variant Fc region) of an antibody. Examples of antibody effector functions include C1q binding; complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC);

phagocytosis; down regulation of cell surface receptors (e.g. B cell receptor; BCR), and the like. To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as those described in U.S. Pat. Nos. 5,500,362 or 5,821,337 may be performed.

Depending on the amino acid sequence of the constant domain of their heavy chains, 5 full length antibodies can be assigned to different “classes”. There are five major classes of full length antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into “subclasses” (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the different classes of antibodies are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of 10 different classes of antibodies are well known.

The “light chains” of antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

The term “hypervariable region” when used herein refers to the amino acid residues 15 of an antibody which are responsible for antigen binding. The hypervariable region generally comprises amino acid residues from a “complementarity determining region” or “CDR” (e.g. residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National 20 Institutes of Health, Bethesda, Md. (1991)) and/or those residues from a “hypervariable loop” (e.g. residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). “Framework Region” or “FR” residues are those 25 variable domain residues other than the hypervariable region residues as herein defined. The hypervariable region or the CDRs thereof can be transferred from one antibody chain to another or to another protein to confer antigen binding specificity to the resulting (composite) antibody or binding protein.

“Humanized” forms of non-human (e.g., rodent) antibodies are chimeric antibodies 30 that contain minimal sequence derived from the non-human antibody. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human antibody are replaced by corresponding non-human

residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. For further details, see Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 5 (1992).

An “affinity matured” antibody has one or more alterations in one or more hypervariable regions thereof which result an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). In one aspect, affinity matured antibodies will have nanomolar or even picomolar affinities for the 10 target antigen. Affinity matured antibodies are produced by procedures known in the art. Marks *et al.* *Bio/Technology* 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by: Barbas *et al.* *Proc Nat. Acad. Sci, USA* 91:3809-3813 (1994); Schier *et al.* *Gene* 169:147-155 15 (1995); Yelton *et al.* *J. Immunol.* 155:1994-2004 (1995); Jackson *et al.*, *J. Immunol.* 154(7):3310-9 (1995); and Hawkins *et al.*, *J. Mol. Biol.* 226:889-896 (1992).

An “isolated” antibody is one which has been identified and separated and/or recovered from a component of its natural environment. In certain embodiments, the antibody will be purified (1) to greater than 95% by weight of protein as determined by the Lowry method, and alternatively, more than 99% by weight, (2) to a degree sufficient to 20 obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or non-reducing conditions using Coomassie blue or silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural 25 environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

“Treatment” refers to therapeutic treatment. Those in need of treatment include those already with disease. Hence, the patient, e.g., human, to be treated herein may have been diagnosed as suffering from a disease, such as an inflammatory bowel disease, e.g., Crohn's Disease or ulcerative colitis. The terms “patient” and “subject” are used interchangeably 30 herein.

“Prevention” refers to a treatment that results in the absence or reduction in the severity of an adverse event. In a population of patients, when treatment typically results in a certain percentage of adverse events, or a certain percentage of adverse events that are severe, but a treatment administered for prevention purposes instead results in a lower percentage of

adverse events (i.e., a lower or reduced risk of adverse events) or a lower percentage of adverse events that are severe (i.e., a lower or reduced risk that the adverse event is severe).

An anti- α 4 β 7 antibody, vedolizumab, a humanized monoclonal antibody that has binding specificity for the α 4 β 7 integrin, is already indicated for the treatment of patients with moderately to severely active ulcerative colitis (UC) and Crohn's disease (CD).

Vedolizumab has a novel gut-selective mechanism of action. By binding to cell surface-expressed α 4 β 7, vedolizumab is an α 4 β 7 antagonist and blocks a subset of memory gut-homing T lymphocytes from interacting with mucosal addressin cell adhesion molecule-1 (MAdCAM-1) expressed on endothelial cells.

Several factors are associated with accelerated clearance of antibodies including the presence of anti-drug antibodies, sex, body size, concomitant immunosuppressant use, disease type, albumin concentration, and degree of systemic inflammation. Furthermore, a consistent relationship between efficacy and exposure, in distinction to drug dose, has been observed for many of these agents, such that higher trough drug concentrations are associated with greater efficacy. Differences in drug clearance may be an important explanation for this observation. For example, cancer patients undergo immunosuppressive treatment of the tumor and treatment for infection. Therefore, an understanding of the determinants of clearance for therapeutic antibodies in transplant patients may result in optimization of drug regimens.

In previous studies, single-dose pharmacokinetics, pharmacodynamics (α 4 β 7 receptor saturation), safety, and tolerability of vedolizumab were investigated over a dose range of 0.2 to 10 mg/kg in healthy volunteers (intravenous [IV] infusion) (unpublished data). After reaching peak concentrations, vedolizumab serum concentrations fell in a generally biexponential fashion until concentrations reached approximately 1 to 10 ng/mL. Thereafter, concentrations appeared to fall in a nonlinear fashion. The multiple-dose pharmacokinetics and pharmacodynamics of vedolizumab have been investigated following IV infusions of 0.5 and 2 mg/kg in patients with CD and infusion of 2, 6, and 10 mg/kg in patients with UC. Vedolizumab pharmacokinetics was generally linear following an IV infusion over the dose range of 2 to 10 mg/kg in patients with UC. After multiple-dose administration, rapid and near complete α 4 β 7 receptor saturation was achieved following the initial dose of vedolizumab.

The efficacy and safety of vedolizumab induction and maintenance therapy were demonstrated in patients with CD in the GEMINI 2 (ClinicalTrials.gov number, NCT00783692) and GEMINI 3 (ClinicalTrials.gov number, NCT01224171) trials. The

exposure-response (efficacy) relationships of vedolizumab in patients with CD for induction and maintenance therapy have been presented elsewhere.

The TNF antagonist may be Adalimumab. Adalimumab is a recombinant human IgG1 monoclonal antibody specific for human TNF- α . Adalimumab binds specifically to TNF- α and blocks its interaction with the p55 and p75 cell surface TNF- α receptors. Adalimumab also lyses surface TNF- α expressing cells in vitro in the presence of complement. Adalimumab does not bind or inactivate lymphotoxin (TNF- β). TNF- α is a naturally occurring cytokine that is involved in normal inflammatory and immune responses. TNF- α plays an important role in both the pathologic inflammation and the joint destruction that are hallmarks of inflammatory disease. Adalimumab also modulates biological responses that are induced or regulated by TNF- α , including changes in the levels of adhesion molecules responsible for leukocyte migration.

Adalimumab is approved globally for the treatment of moderately to severely active CD in patients who have had an inadequate response to conventional therapy. Additionally, adalimumab is approved for reducing signs and symptoms and inducing clinical remission in these patients if they have also lost response to or are intolerant to infliximab.

Adalimumab 40 mg weekly or every other week has been shown to be effective for maintaining clinical remission in patients with CD who have responded to adalimumab induction therapy. Adalimumab has also been shown to have corticosteroid-sparing effects in patients with CD (Behm BW BS. Cochrane Database of Systematic Reviews 2008(1)).

The immunomodulator may be an antimetabolite. Methotrexate is an antimetabolite that is approved for the treatment of certain neoplastic diseases, severe psoriasis, and adult rheumatoid arthritis. Methotrexate was originally developed as an antineoplastic, and is an inhibitor of the enzyme dihydrofolate reductase, which catalyzes a reaction in the production of tetrahydrofolic acid. This pathway is essential to the synthesis of the purine and pyrimidine components of deoxyribonucleic acid (DNA), and thus the drug slows cellular division. Although the mechanism of action of low-dose methotrexate in the treatment of autoimmune disease is not well understood, there is evidence supporting both immunosuppressive and anti-inflammatory effects.

30

Treatment of Inflammatory Bowel Disease with Triple Combination Therapy

The invention relates to a method of treating inflammatory bowel disease (e.g., Crohn's Disease (CD) or ulcerative colitis) in a patient using a triple combination therapy. The human patient may be an adult (e.g., 18 years or older), an adolescent, or a child. The

triple combination therapy may improve remission rates and modify disease progression by including agents with different mechanisms of action, particularly for patients who are at higher risk for CD complications. Patients who respond to treatment with infliximab or adalimumab have been shown to have a decrease in MAdCAM and an increase in $\alpha 4\beta 7$ -positive cells (Biancheri P, et al. Inflamm Bowel Dis 2013;19(2):259-64.). Thus, their vedolizumab pathway is active, and the addition of vedolizumab to the therapy regimen can provide clinically relevant synergy.

In one embodiment, the triple combination therapy of the invention is used to treat patients who are at high risk for developing complications from inflammatory bowel disease. For example, patients at high risk for developing complications from CD, or aggressive CD, may include those patients with involvement of the upper GI tract and ileum, penetrating disease, early age at diagnosis, smoking, extensive ulceration of the mucosa, high titers of serum antibodies, use of corticosteroids at diagnosis, and mutations of the NOD2 gene. Patients at high risk for developing complications of CD may be defined as described above for the PROSPECT algorithm or the AGA criteria. Patients identified as high risk CD patients can be targeted for triple combination therapy. The patient targeted for triple combination therapy may have a Simple Endoscopic Score for Crohn's Disease (SES-CD) of ≥ 7 or ≥ 4 if isolated ileal disease. The patient targeted for triple combination therapy may have active disease as judged by the Crohn's Disease Activity Index (CDAI) of ≥ 220 . The CDAI score weighs factors including the number of liquid or very soft stools, the severity of abdominal pain, general well being, extra-intestinal manifestations of the disease, such as arthritis, iritis, erythema, fistula or abscess or fever, whether the patient is taking an antidiarrheal medication, abdominal mass, hematocrit and body weight.

Anti-Integrin

An anti-integrin, such as an anti- $\alpha 4\beta 7$ antibody, is administered in an effective amount which inhibits binding of $\alpha 4\beta 7$ integrin to a ligand thereof. For therapy, an effective amount will be sufficient to achieve the desired effect (e.g., inducing endoscopic remission, inducing mucosal healing, achieving clinical remission, achieving biomarker remission). An $\alpha 4\beta 7$ antagonist, such as an anti- $\alpha 4\beta 7$ antibody may be administered in a unit dose or multiple doses. The dosage can be determined by methods known in the art and can be dependent, for example, upon the individual's age, sensitivity, tolerance and overall well-being. Examples of modes of administration include topical routes such as nasal or inhalational or transdermal administration, enteral routes, such as through a feeding tube or suppository, and parenteral routes, such as intravenous, intramuscular, subcutaneous,

intraarterial, intraperitoneal, or intravitreal administration. Suitable dosages for antibodies can be from about 0.1 mg/kg body weight to about 10.0 mg/kg body weight per treatment, for example about 2 mg/kg to about 7 mg/kg, about 3 mg/kg to about 6 mg/kg, or about 3.5 to about 5 mg/kg. In particular embodiments, the dose administered is about 0.3 mg/kg, about 5 mg/kg, about 1 mg/kg, about 2 mg/kg, about 3 mg/kg, about 4 mg/kg, about 5 mg/kg, about 6 mg/kg, about 7 mg/kg, about 8 mg/kg, about 9 mg/kg, or about 10 mg/kg. In some embodiments, vedolizumab is administered at a dose of 50 mg, 75 mg, 100 mg, 300 mg, 450 mg, 500 mg or 600 mg. In some embodiments, vedolizumab is administered at a dose of 108 mg, 90 to 120 mg, 216 mg, 160 mg, 165 mg, 155 to 180 mg, 170 mg or 180 mg. In some 10 embodiments, the anti- $\alpha 4\beta 7$ antibody, such as vedolizumab is provided as a dry, lyophilized formulation which can be reconstituted with a liquid, such as sterile water, for administration. Administration of a reconstituted formulation can be by parenteral injection by one of the routes described above. An intravenous injection can be by infusion, such as by further dilution with sterile isotonic saline, buffer, e.g., phosphate-buffered saline or Ringer's 15 (lactated or dextrose) solution. In some embodiments, the anti- $\alpha 4\beta 7$ antibody is administered by subcutaneous injection, e.g., a dose of about 108 mg or about 165 mg or about 216 mg, at about every two, three or four weeks after the start of therapy or after the third subsequent dose.

The anti-integrin, which is an anti- $\alpha 4\beta 7$ antibody, can bind to an epitope on the $\alpha 4$ 20 chain (e.g., humanized MAb 21.6 (Bendig *et al.*, U.S. Pat. No. 5,840,299), on the $\beta 7$ chain (e.g., FIB504 or a humanized derivative (e.g., Fong *et al.*, U.S. Pat. No. 7,528,236)), or to a combinatorial epitope formed by the association of the $\alpha 4$ chain with the $\beta 7$ chain. AMG-181 or other antibodies described in US 2010/0254975 are anti- $\alpha 4\beta 7$ antibodies. In one aspect, the antibody binds a combinatorial epitope on the $\alpha 4\beta 7$ complex, but does not bind an 25 epitope on the $\alpha 4$ chain or the $\beta 7$ chain unless the chains are in association with each other. The association of $\alpha 4$ integrin with $\beta 7$ integrin can create a combinatorial epitope for example, by bringing into proximity residues present on both chains which together comprise the epitope or by conformationally exposing on one chain, e.g., the $\alpha 4$ integrin chain or the $\beta 7$ integrin chain, an epitopic binding site that is inaccessible to antibody binding in the 30 absence of the proper integrin partner or in the absence of integrin activation. In another aspect, the anti- $\alpha 4\beta 7$ antibody binds both the $\alpha 4$ integrin chain and the $\beta 7$ integrin chain, and thus, is specific for the $\alpha 4\beta 7$ integrin complex. Such antibodies can bind $\alpha 4\beta 7$ but not bind $\alpha 4\beta 1$, and/or not bind $\alpha E\beta 7$, for example. In another aspect, the anti- $\alpha 4\beta 7$ antibody binds to

the same or substantially the same epitope as the Act-1 antibody (Lazarovits, A. I. *et al.*, *J. Immunol.*, 133(4): 1857-1862 (1984), Schweighoffer *et al.*, *J. Immunol.*, 151(2): 717-729, 1993; Bednarczyk *et al.*, *J. Biol. Chem.*, 269(11): 8348-8354, 1994). Murine ACT-1 Hybridoma cell line, which produces the murine Act-1 monoclonal antibody, was deposited 5 under the provisions of the Budapest Treaty on Aug. 22, 2001, on behalf Millennium Pharmaceuticals, Inc., 40 Landsdowne Street, Cambridge, Mass. 02139, U.S.A., at the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209, U.S.A., under Accession No. PTA-3663. In another aspect, the anti- $\alpha 4\beta 7$ antibody is a human antibody or an $\alpha 4\beta 7$ binding protein using the CDRs provided in U.S. Patent 10 Application Publication No. 2010/0254975.

In one aspect, the anti-integrin is an anti-MAdCAM antibody (see e.g., US Patent No. 8,277,808, PF-00547659 or antibodies described in WO2005/067620), or an engineered form of a ligand, such as a MAdCAM-Fc chimera such as described in US Patent No. 7,803,904.

In one aspect, the anti-integrin inhibits binding of $\alpha 4\beta 7$ to one or more of its ligands 15 (e.g. the mucosal addressin, e.g., MAdCAM (e.g., MAdCAM-1), fibronectin, and/or vascular addressin (VCAM)). Primate MAdCAMs are described in the PCT publication WO 96/24673, the entire teachings of which are incorporated herein by this reference. In another aspect, the anti-integrin inhibits binding of $\alpha 4\beta 7$ to MAdCAM (e.g., MAdCAM-1) and/or fibronectin without inhibiting the binding of VCAM.

20 In one aspect, the anti-integrin, e.g., an anti- $\alpha 4\beta 7$ antibody, has the binding specificity, e.g., comprises the complementarity determining regions of mouse Act-1 antibody. For example, an anti- $\alpha 4\beta 7$ antibody comprises a heavy chain that contains the 3 heavy chain complementarity determining regions (CDRs, CDR1, SEQ ID NO:4, CDR2, SEQ ID NO:5 and CDR3, SEQ ID NO:6) of the mouse Act-1 antibody and suitable human 25 heavy chain framework regions; and also comprises a light chain that contains the 3 light chain CDRs (CDR1, SEQ ID NO:7, CDR2, SEQ ID NO:8 and CDR3, SEQ ID NO:9) of the mouse Act-1 antibody and suitable human light chain framework regions. In some embodiments the anti- $\alpha 4\beta 7$ antibody is an IgG1 isotype.

30 In one aspect, the anti-integrin for use in the treatments are humanized versions of the mouse Act-1 antibody. Suitable methods for preparing humanized antibodies are well-known in the art. Generally, the humanized anti- $\alpha 4\beta 7$ antibody will contain a heavy chain that contains the 3 heavy chain complementarity determining regions (CDRs, CDR1, SEQ ID NO:4, CDR2, SEQ ID NO:5 and CDR3, SEQ ID NO:6) of the mouse Act-1 antibody and suitable human heavy chain framework regions; and also contain a light chain that contains

the 3 light chain CDRs (CDR1, SEQ ID NO:7, CDR2, SEQ ID NO:8 and CDR3, SEQ ID NO:9) of the mouse Act-1 antibody and suitable human light chain framework regions. The humanized Act-1 antibody can contain any suitable human framework regions, including consensus framework regions, with or without amino acid substitutions. For example, one or 5 more of the framework amino acids can be replaced with another amino acid, such as the amino acid at the corresponding position in the mouse Act-1 antibody. The human constant region or portion thereof, if present, can be derived from the κ or λ light chains, and/or the γ (e.g., γ 1, γ 2, γ 3, γ 4), μ , α (e.g., α 1, α 2), δ or ϵ heavy chains of human antibodies, including allelic variants. A particular constant region (e.g., IgG1), variant or portions thereof can be 10 selected in order to tailor effector function. For example, a mutated constant region (variant) can be incorporated into a fusion protein to minimize binding to Fc receptors and/or ability to fix complement (see e.g., Winter *et al.*, GB 2,209,757 B; Morrison *et al.*, WO 89/07142; Morgan *et al.*, WO 94/29351, Dec. 22, 1994). Humanized versions of Act-1 antibody were described in PCT publications nos. WO98/06248 and WO07/61679, the entire teachings of 15 each of which are incorporated herein by this reference. Treatment methods using anti- α 4 β 7 integrin antibodies are described in publication nos. U.S. 2005/0095238, U.S. 2005/0095238, WO2012151248 and WO 2012/151247.

In one aspect, the anti-integrin is the anti- α 4 β 7 antibody, vedolizumab. Vedolizumab IV (also called MLN0002, ENTYVIOTM or KYNTELESTM) is a humanized antibody (Ig) G1 20 mAb directed against the human lymphocyte integrin α 4 β 7. The α 4 β 7 integrin mediates lymphocyte trafficking to GI mucosa, gut-associated lymphoid tissue (GALT) and mesenteric lymph nodes through adhesive interaction with mucosal addressin cell adhesion molecule-1 (MAdCAM-1), which is expressed on the endothelium of mesenteric lymph nodes and GI 25 mucosa. Vedolizumab binds the α 4 β 7 integrin, antagonizes its adherence to MAdCAM-1 and as such, impairs the migration of naïve T cells to the GALT and mesenteric lymph nodes and gut homing leukocytes into GI mucosa.

In another aspect, the anti-integrin is a humanized anti- α 4 β 7 antibody for use in the treatment and comprises a heavy chain variable region comprising amino acids 20 to 140 of SEQ ID NO:1, and a light chain variable region comprising amino acids 20 to 131 of SEQ ID 30 NO:2 or amino acids 1 to 112 of SEQ ID NO:3. If desired, a suitable human constant region(s) can be present. For example, the humanized anti- α 4 β 7 antibody can comprise a heavy chain that comprises amino acids 20 to 470 of SEQ ID NO:1 and a light chain comprising amino acids 1 to 219 of SEQ ID NO:3. In another example, the humanized anti- α 4 β 7 antibody can comprise a heavy chain that comprises amino acids 20 to 470 of SEQ ID

NO:1 and a light chain comprising amino acids 20 to 238 of SEQ ID NO:2. Vedolizumab is cataloged under Chemical Abstract Service (CAS, American Chemical Society) Registry number 943609-66-3).

Substitutions to the humanized anti- α 4 β 7 antibody sequence can be, for example, 5 mutations to the heavy and light chain framework regions, such as a mutation of isoleucine to valine on residue 2 of SEQ ID NO:10; a mutation of methionine to valine on residue 4 of SEQ ID NO:10; a mutation of alanine to glycine on residue 24 of SEQ ID NO:11; a mutation of arginine to lysine at residue 38 of SEQ ID NO:11; a mutation of alanine to arginine at residue 40 of SEQ ID NO:11; a mutation of methionine to isoleucine on residue 48 of SEQ 10 ID NO:11; a mutation of isoleucine to leucine on residue 69 of SEQ ID NO:11; a mutation of arginine to valine on residue 71 of SEQ ID NO:11; a mutation of threonine to isoleucine on residue 73 of SEQ ID NO:11; or any combination thereof; and replacement of the heavy chain CDRs with the CDRs (CDR1, SEQ ID NO:4, CDR2, SEQ ID NO:5 and CDR3, SEQ ID NO:6) of the mouse Act-1 antibody; and replacement of the light chain CDRs with the 15 light chain CDRs (CDR1, SEQ ID NO:7, CDR2, SEQ ID NO:8 and CDR3, SEQ ID NO:9) of the mouse Act-1 antibody.

The dose can be administered to the patient over about 20 minutes, about 25 minutes, about 30 minutes, about 35 minutes, or about 40 minutes. In some embodiments, the dose can be administered to the patient for up to two hours.

20 TNF Antagonist

The invention features a triple combination therapy which includes administration of a TNF antagonist for the treatment of an inflammatory bowel disease. A TNF antagonist is administered in an effective amount which inhibits binding TNF- α and blocks its interaction with the p55 and p75 cell surface TNF- α receptors. Suitable TNF antagonists include, for 25 example, adalimumab, infliximab, etanercept, golimumab, certolizumab pegol, biosimilars thereof and equivalents thereof. For therapy, an effective amount will be sufficient to achieve the desired effect (e.g., inducing endoscopic remission, inducing mucosal healing, achieving clinical remission, achieving biomarker remission). A TNF antagonist, such as an anti-TNF- α antibody (e.g., adalimumab) may be administered in a unit dose or multiple doses. The 30 dosage can be determined by methods known in the art and can be dependent, for example, upon the individual's age, sensitivity, tolerance and overall well-being. Examples of modes of administration include topical routes such as nasal or inhalational or transdermal administration, enteral routes, such as through a feeding tube or suppository, and parenteral routes, such as intravenous, intramuscular, subcutaneous, intraarterial, intraperitoneal, or

intravitreal administration. Suitable dosages for antibodies can be from about 0.1 mg/kg body weight to about 10.0 mg/kg body weight per treatment, for example about 1 mg/kg to about 8 mg/kg, about 2 mg/kg to about 7 mg/kg, about 3 mg/kg to about 6 mg/kg, or about 3.5 to about 5 mg/kg. In particular embodiments, the dose administered is about 0.3 mg/kg, 5 about 0.5 mg/kg, about 1 mg/kg, about 2 mg/kg, about 3 mg/kg, about 4 mg/kg, about 5 mg/kg, about 6 mg/kg, about 7 mg/kg, about 8 mg/kg, about 9 mg/kg, or about 10 mg/kg. In some embodiments, the TNF antagonist is administered at a dose of 40 mg, 50 mg, 75 mg, 80 mg, 100 mg, 150 mg, 160 mg, 175 mg, 200 mg, 300 mg, 400 mg, 450 mg, 500 mg or 600 mg. In some embodiments, the TNF antagonist is administered at a dose of 40 mg, 80 mg, 10 160 mg, 80 to 200 mg, 170 mg, 165 mg, 155 mg, 150 to 180 mg, 140 mg or 130 mg.

In one embodiment, the triple combination therapy of the invention includes the use of adalimumab for the treatment of an inflammatory bowel disease. Adalimumab may be administered to a patient having an inflammatory bowel disease, such as Crohn's disease or ulcerative colitis. In one embodiment, adalimumab is subcutaneously administered to a 15 subject having Crohn's disease or ulcerative colitis at an initial dose of 160 mg at week 0, followed by a second dose of 80 mg at week 2, followed by a 40 dose at week 4 and every two weeks thereafter. In one embodiment, the triple combination therapy comprises discontinuing adalimumab at week 26 of treatment in the patient.

In one embodiment, the triple combination therapy of the invention includes the use 20 of infliximab for the treatment of an inflammatory bowel disease, such as Crohn's disease or ulcerative colitis. In one embodiment, infliximab is intravenously administered to a subject having Crohn's disease or ulcerative colitis at a dose of 5 mg/kg at weeks 0, 2 and 6 weeks, and then every 8 weeks thereafter. In one embodiment, the triple combination therapy comprises discontinuing infliximab at week 26 of treatment in the patient.

In one embodiment, the triple combination therapy of the invention includes the use 25 of golimumab for the treatment of an inflammatory bowel disease, such as Crohn's disease or ulcerative colitis. In one embodiment, golimumab is subcutaneously administered to a subject having Crohn's disease or ulcerative colitis at a dose of 200 mg at week 0, followed by a 100 mg dose a week 2, followed by a 100 mg dose every 4 weeks thereafter. In one 30 embodiment, the triple combination therapy comprises discontinuing golimumab at week 26 of treatment in the patient.

In one embodiment, the triple combination therapy of the invention includes the use of certolizumab pegol for the treatment of an inflammatory bowel disease, such as Crohn's disease or ulcerative colitis. In one embodiment, certolizumab pegol is subcutaneously

administered to a subject having Crohn's disease or ulcerative colitis at a dose of 400 mg at week 0, followed by 400 mg doses at weeks 2 and 4. If a response occurs, then 400 mg of certolizumab pegol is administered every four weeks thereafter. In one embodiment, the triple combination therapy comprises discontinuing certolizumab pegol at week 26 of

5 treatment in the patient.

Immunomodulators

An immunomodulator is administered in an effective amount to reduce immunogenicity. Examples of suitable immunomodulators include fluorouracil, cladribine, capecitabine, methotrexate, pemetrexed, mercaptopurine, hydroxyurea, gemcitabine,

10 fludarabine, pralatrexate, clofarabine, cytarabine, decitabine, floxuridine, nelarabine, and thioguanine. In one embodiment the immunomodulator is methotrexate. For therapy, an effective amount will be sufficient to achieve the desired effect (e.g., inducing endoscopic remission, inducing mucosal healing, achieving clinical remission, achieving biomarker remission). An immunomodulator, such as an antimetabolite (e.g., methotrexate) may be

15 administered in a unit dose or multiple doses. The dosage can be determined by methods known in the art and can be dependent, for example, upon the individual's age, sensitivity, tolerance and overall well-being. Examples of modes of administration include oral administration, topical routes such as nasal or inhalational or transdermal administration, enteral routes, such as through a feeding tube or suppository, and parenteral routes, such as

20 intravenous, intramuscular, subcutaneous, intraarterial, intraperitoneal, or intravitreal administration. Suitable dosages for immunomodulators can be a dose of 5 mg, 10 mg, 15 mg, 20 mg, 25 mg, 30 mg, or 45 mg. In some embodiments, methotrexate is administered at a dose of 10 mg, 15 mg, 20 mg, 10 to 30 mg, 25 mg, 30 mg, or 45 mg.

25 The dose of the immunomodulator, e.g., methotrexate, can be administered to the patient daily, weekly, or monthly. In some embodiments, the dose can be administered to the patient orally weekly for up to 34 weeks.

The triple combination therapy, such as anti-integrin/TNF antagonist/immunomodulator, may be administered to an individual (e.g., a human) in series, in parallel, or simultaneously. The triple combination therapy may be administered to an 30 individual by a single route of administration, by two different routes of administration, or by three different routes of administration.

In one embodiment, the dosing regimen comprises administration of 300 mg of the anti-integrin by intravenous infusion at Weeks 0, 2, 6, and then every 8 weeks thereafter; administration of 160 mg of the TNF antagonist, e.g., adalimumab, by subcutaneous injection

at Week 0, administration of 80 mg of the TNF antagonist, *e.g.*, adalimumab, at Week 2, and then 40 mg of the TNF antagonist, *e.g.*, adalimumab, every 2 weeks thereafter; and oral administration of 15 mg of the immunomodulator, *e.g.*, methotrexate, weekly.

In one embodiment, the method comprises administering an effective amount of the 5 triple combination therapy to a patient. If the triple combination therapy, or any component of the triple combination therapy, is in a formulation which is in a solid, *e.g.*, dry state, the process of administration can comprise a step of converting the formulation to a liquid state. In one aspect, a dry formulation can be reconstituted, *e.g.*, by a liquid as described above, for use in injection, *e.g.* intravenous, intramuscular or subcutaneous injection. In another aspect, 10 a solid or dry formulation can be administered topically, *e.g.*, in a patch, cream, aerosol or suppository.

In one embodiment, the present invention provides a method for treating a high risk CD patient with triple combination therapy. The method comprises the steps of 15 administering an initial 300 mg dose of an anti-integrin to a high risk CD patient, administering a subsequent anti-integrin 300 mg dose two weeks after the initial dose, administering a second subsequent anti-integrin 300 mg dose six weeks after the initial dose, and subsequent 300 mg doses every eight weeks thereafter, administering an initial 160 mg dose of a TNF antagonist, followed by a subsequent 80 mg dose of TNF antagonist two weeks after the initial dose, and subsequent 40 mg doses of TNF antagonist every two weeks 20 thereafter, and administering a 15 mg methotrexate dose weekly beginning at Week 0.

The invention provides a triple combination therapy for use in treating an 25 inflammatory bowel disease, such as CD or UC, in a patient having high risk of complications related to the inflammatory bowel disease, such as CD or UC, the use comprising administering an initial dose of anti-integrin, two weeks after the initial dose, and six weeks after the initial dose, and every eight weeks thereafter for a total of 102 weeks; administering an initial dose of TNF antagonist, half the initial dose two weeks after the initial dose, and a quarter of the initial dose every two weeks thereafter for a total of 26 weeks; and administering a weekly dose of an immunomodulator for 34 weeks. In some 30 embodiments, the anti-integrin is vedolizumab. In some embodiments the TNF modulator is adalimumab. In some embodiments the immunomodulator is methotrexate. In some embodiments the triple combination therapy includes administration of vedolizumab, adalimumab, and methotrexate. In some embodiments the triple combination therapy includes vedolizumab, a TNF antagonist, and methotrexate. In some embodiments the triple combination therapy includes vedolizumab, a TNF antagonist, and an immunomodulator. In

some embodiments the triple combination therapy includes an anti- α 4 β 7 antibody, a TNF antagonist, and methotrexate.

In one embodiment, the dosing regimen comprises administration of 300 mg of vedolizumab by intravenous infusion at Weeks 0, 2, 6, and then every 8 weeks thereafter; 5 administration of 160 mg of adalimumab by subcutaneous injection at Week 0, administration of 80 mg of adalimumab at Week 2, and then 40 mg of the adalimumab every 2 weeks thereafter; and oral administration of 15 mg of methotrexate weekly.

In some embodiments, the IBD, e.g., CD, has an exacerbation after 26 weeks, e.g., as defined by a CDAI increase of >70 from the prior visit on two occasions two weeks apart and 10 disease activity as seen by colonoscopy and a CRP value above normal (> 5 mg/L); or fecal calprotectin >250 μ g/g. In an embodiment of exacerbation of IBD, 300 mg vedolizumab may be administered every 4 weeks.

The triple combination therapy may further comprise a corticosteroid tapering regimen, such as beginning about 1 to 3 weeks or about two weeks after starting triple 15 combination therapy. In some embodiments, a high risk IBD patient may be undergoing prior therapy with corticosteroids, such as prednisone, e.g. 10 to 20 mg/day, 20 to 40 mg/day, 25 to 35 mg/day or about 30 mg/day, or budesonide, e.g., 2 to 12 mg/day, 3 to 10 mg/day or about 9 mg/day, at diagnosis or at Baseline. For prednisone, the dose may be reduced at a rate of 5 mg/ week until 5 mg/day is reached, then the dose of prednisone is reduced at a rate 20 of 2.5 mg/week until discontinuation. For budesonide, the dose is reduced at a rate of 3 mg every 2 weeks until discontinuation.

In some embodiments, the triple combination therapy of a high risk patient results in 25 endoscopic remission of the inflammatory bowel disease. “Endoscopic remission” as used herein, refers to a condition with a low endoscopic score. An example of a method to assess the endoscopic score in ulcerative colitis is flexible sigmoidoscopy. The endoscopic score in ulcerative colitis can be the Mayo subscore. An example of a method to assess the endoscopic score in Crohn’s disease is ileocolonoscopy. The endoscopic score in Crohn’s disease can be the simple endoscopic score for Crohn’s Disease (SES-CD). The SES-CD can include measures such as the size of ulcers, the amount of ulcerated surface, the amount of 30 affected surface and whether and to what extent the alimentary canal is narrowed.

In an embodiment, a high risk Crohn’s disease triple combination therapy patient achieves endoscopic remission as defined by a SES-CD score of 0 to 2 by week 26, or by weeks 22 to 30 or weeks 24 to 28 of treatment. In other embodiments, the triple combination

therapy of the invention results in a degree of response or remission selected from the group consisting of SES-CD ≤ 4 and reduction from baseline SES-CD of at least 2 points and no individual SES-CD subscore >1 at Week 26; 50% reduction in SES-CD from Baseline at week 26; deep remission (defined as CDAI<150 and SES-CD 0-2) at week 26; clinical 5 remission (defined as CDAI<150) and endoscopic response as a measure of mucosal healing (defined as 50% reduction in SES-CD from Baseline) at week 26; clinical remission (CDAI score <150) at Weeks 10 and 26; clinical response (defined as ≥ 100 -point decrease in CDAI score) at Weeks 10 and 26; clinical remission (defined as CDAI<150) and C-reactive protein (CRP) <5 mg/L (in subjects with elevated CRP at Baseline) at Weeks 26, 52, 78, and 102; 10 and discontinuation of Baseline corticosteroid use and clinical remission (CDAI score <150) at Weeks 10, 26, and 102.

In some embodiments, a high risk Crohn's disease triple combination therapy patient achieves long-term benefit. In one embodiment, the triple combination therapy of the invention results in maintaining clinical remission (defined as CDAI <150) at Weeks 52, 78, 15 and 102. In another embodiment, the triple combination therapy of the invention results in maintaining endoscopic remission defined as SES-CD 0-2 at Week 102. In another embodiment, the triple combination therapy of the invention results in maintaining deep remission (defined as CDAI <150 and SES-CD 0-2) at Week 102. In other embodiments, the triple combination therapy of the invention results in a degree of long term benefit selected 20 from the group consisting of maintaining endoscopic healing defined as SES-CD ≤ 4 and reduction from baseline SES-CD of at least 2 points and no individual SES-CD subscore >1 at Week 102; maintaining endoscopic response defined as 50% reduction in SES-CD from Baseline at Week 102; maintaining clinical remission (defined as CDAI <150) and endoscopic response as a measure of mucosal healing (defined as 50% reduction in SES-CD 25 from baseline) at Week 102; achieving normalization of CRP defined as <5 mg/L at Week 26 (in those elevated at Baseline); and achieving a Patient-reported outcome 2 (PRO-2) score ≤ 75 (a subset of the CDAI score and derived from a combination of liquid or very soft stools (multiplied by 2) and abdominal pain severity ((on a scale of 0 to 3) multiplied by 5) over averaged over the prior 7 days) and an SES-CD ≤ 4 and a reduction from baseline SES-CD of 30 at least 2 points and no individual SES-CD subscore >1 at Weeks 26 and 102.

High Risk Crohn's Disease Patients

In one embodiment, the triple combination therapy disclosed herein is used to treat a subpopulation of Crohn's disease patients who are at risk for developing complications following diagnosis. Such high risk patients may be identified using predictive methods

known in the art, including, for example, the PROSPECT predictive method. As described in Siegel et al. (2016) *Aliment Pharmacol Ther* 43: 262–271, the PROSPECT predictive test is based on clinical, serologic and genetic variables of the patient, and is used to determine the likelihood that a CD patient will have a disease complication by year 2 of diagnosis. A high 5 risk CD patient is one who has at least a 20% chance of such a complication by year 2, as determined by, for example, the PROSPECT predictive method.

High Risk Ulcerative Colitis Disease Patients

In one embodiment, the triple combination therapy disclosed herein is used to treat a subpopulation of UC disease patients who are at risk for developing complications following 10 diagnosis. Such high risk UC patients are understood to be at an increased risk for colectomy. High risk UC patients may also be identified by their failure to maintain steroid-induced remission.

The invention will be more fully understood by reference to the following examples. They should not, however, be construed as limiting the scope of the invention. All literature 15 and patent citations are incorporated herein by reference.

EXEMPLIFICATION

Example 1

A phase 4, open-label, multicenter study in subjects with newly-diagnosed CD at 20 higher risk for complications due to aggressive disease is designed to evaluate the safety, tolerability, and clinical activity of triple combination therapy (300 mg vedolizumab IV, 160/80/40 mg adalimumab SC, and 15 mg oral methotrexate). A predictive tool incorporating clinical, serological, and genetic variables will be used to stratify which subjects are at higher risk for developing complications. The study will run over a 26-week treatment period for 25 induction of endoscopic remission and mucosal healing followed by efficacy and safety of vedolizumab IV monotherapy to maintain remission for 76 weeks, for a total treatment period of 102 weeks.

Patients are eligible for the study if they have been diagnosed with CD within the previous 18 months, and are naïve to biologics. Patients who meet all eligibility criteria and 30 provide written informed consent are enrolled in this study.

Patients are excluded from the study if they are at low risk for CD complications as calculated by the CD PROSPECT predictive tool, have any evidence of active infection during screening, have a history of any bacterial, viral, and other infection due to opportunistic pathogens, have had any prior exposure to any TNF antagonist biologic

therapy, or to vedolizumab, natalizumab, efalizumab, or rituximab, have any prior CD-related surgery or CD complication requiring surgery, have a history of tuberculosis, lymphoma, lymphoproliferative disease, or congestive heart failure or unstable angina. Patients with a medical history that contradicts the use of methotrexate or adalimumab or indicates

5 intolerance to methotrexate or adalimumab are also excluded.

The study consists of up to a 4-week screening period, a 26-week combination treatment period, an additional 76-week vedolizumab IV monotherapy treatment period, and a 26-week follow-up period following last dose (with final efficacy evaluations at week 120).

The duration of the study from Screening to final efficacy visit at Week 120 will be

10 approximately 124 weeks. All subjects will participate in safety follow-up telephone calls 18 and 26 weeks after last dose, for a total follow-up of 128 weeks. Subjects who discontinue early will also participate in safety follow-up telephone calls 18 and 26 weeks after their last dose. The primary endpoint will be subjects achieving endoscopic remission (defined as Simple Endoscopic Score for Crohn's Disease (SES-CD) 0-2) at Week 26. The SES-CD

15 evaluates 4 endoscopic variables (ulcer size, proportion of the surface area that is ulcerated, proportion of the surface area affected, and stenosis) by scoring each variable on a scale from 0 to 3 where higher scores indicate more severe disease, in 5 colonic segments (ileum, right colon, transverse colon, left colon, and rectum). The score for each endoscopic variable is the sum of the values obtained for each segment. The SES-CD Total is the sum of the 4

20 endoscopic variable scores from 0 to 56.

Vital signs, physical and neurological examinations, adverse event (AE) assessments, and laboratory values (chemistry, hematology, and urinalysis) are obtained to evaluate the safety and tolerability of the triple combination therapy. To exclude patients with progressive multifocal leukoencephalopathy (PML), a Risk Assessment and Minimization for PML

25 (RAMP) questionnaire is administered at Screening and prior to dosing at each visit, to evaluate symptoms suggestive of PML.

SEQUENCE LISTING

SEQ ID NO:1

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly
5 1 5 10 15
Val His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
20 25 30
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Gly Ser Gly Tyr Thr Phe
35 40 45
10 Thr Ser Tyr Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu
50 55 60
Glu Trp Ile Gly Glu Ile Asp Pro Ser Glu Ser Asn Thr Asn Tyr Asn
65 70 75 80
Gln Lys Phe Lys Gly Arg Val Thr Leu Thr Val Asp Ile Ser Ala Ser
15 85 90 95
Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
100 105 110
Tyr Tyr Cys Ala Arg Gly Gly Tyr Asp Gly Trp Asp Tyr Ala Ile Asp
115 120 125
20 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
130 135 140
Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly
145 150 155 160

Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
165 170 175

Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
180 185 190

5 Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
195 200 205

Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn
210 215 220

Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro
10 225 230 235 240

Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
245 250 255

Leu Ala Gly Ala Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
260 265 270

15 Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
275 280 285

Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly
290 295 300

Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn
20 305 310 315 320

Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp
325 330 335

Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro

340 345 350

Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu

355 360 365

5 Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn

370 375 380

Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile

385 390 395 400

Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr

10 405 410 415

Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys

420 425 430

Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys

435 440 445

15 Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu

450 455 460

Ser Leu Ser Pro Gly Lys

465 470

20 SEQ ID NO:2

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly

1 5 10 15

Val His Ser Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val
 20 25 30
 Thr Pro Gly Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu
 35 40 45
 5 Ala Lys Ser Tyr Gly Asn Thr Tyr Leu Ser Trp Tyr Leu Gln Lys Pro
 50 55 60
 Gly Gln Ser Pro Gln Leu Leu Ile Tyr Gly Ile Ser Asn Arg Phe Ser
 65 70 75 80
 Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 10 85 90 95
 Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys
 100 105 110
 Leu Gln Gly Thr His Gln Pro Tyr Thr Phe Gly Gln Gly Thr Lys Val
 115 120 125
 Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro
 130 135 140
 Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu
 145 150 155 160
 20 Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn
 165 170 175
 Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser

180 185 190

Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala

195 200 205

Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly

5 210 215 220

Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

225 230 235

SEQ ID NO:3

10 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Ala Lys Ser

20 25 30

Tyr Gly Asn Thr Tyr Leu Ser Trp Tyr Leu Gln Lys Pro Gly Gln Ser

15 35 40 45

Pro Gln Leu Leu Ile Tyr Gly Ile Ser Asn Arg Phe Ser Gly Val Pro

50 55 60

Asp Arg Phe Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile

65 70 75 80

20 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Leu Gln Gly

85 90 95

Thr His Gln Pro Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys

100 105 110
Arg Ala Asp Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
115 120 125
Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
5 130 135 140
Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
145 150 155 160
Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
165 170 175
10 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
180 185 190
Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
195 200 205
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
15 210 215

SEQ ID NO:4

Ser Tyr Trp Met His

1 5

20

SEQ ID NO:5

Glu Ile Asp Pro Ser Glu Ser Asn Thr Asn Tyr Asn Gln Lys Phe Lys

1 5 10 15

Gly

SEQ ID NO:6

5 Gly Gly Tyr Asp Gly Trp Asp Tyr Ala Ile Asp Tyr

1 5 10

SEQ ID NO:7

Arg Ser Ser Gln Ser Leu Ala Lys Ser Tyr Gly Asn Thr Tyr Leu Ser

10 1 5 10 15

SEQ ID NO:8

Gly Ile Ser Asn Arg Phe Ser

1 5

15

SEQ ID NO:9

Leu Gln Gly Thr His Gln Pro Tyr Thr

1 5

20 SEQ ID NO:10

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
 20 25 30

Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 5 35 40 45

Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
 50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80

10 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Ala
 85 90 95

Leu Gln Thr Pro Gln Thr Phe Gly Gln Gly Lys Val Glu Ile Lys
 100 105 110

15 SEQ ID NO:11

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30

20 Ala Met His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Met
 35 40 45

Gly Trp Ile Asn Ala Gly Asn Gly Asn Thr Lys Tyr Ser Gln Lys Phe

50 55 60

Gln Gly Arg Val Thr Ile Thr Arg Asp Thr Ser Ala Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys

5 85 90 95

Ala Arg Gly Gly Tyr Tyr Gly Ser Gly Ser Asn Tyr Trp Gly Gln Gly

100 105 110

Thr Leu Val Thr Val Ser Ser

115

CLAIMS

1. A method of treating a patient having Crohn's disease (CD), wherein the method comprises the step of:
 - administering to a human patient identified as a high risk CD patient, a triple combination therapy comprising a humanized anti- α 4 β 7 antibody, adalimumab, and an immunomodulator,
 - wherein the triple combination therapy is administered to the patient according to the following dosing regimen:
 - a. an initial dose of 300 mg of humanized anti- α 4 β 7 antibody as an intravenous infusion, followed by a second subsequent dose of 300 mg of humanized anti- α 4 β 7 antibody as an intravenous infusion at about two weeks after the initial dose; followed by a third subsequent dose of 300 mg of the humanized anti- α 4 β 7 antibody as an intravenous infusion at about six weeks after the initial dose;
 - b. an initial dose of 160 mg of adalimumab as a subcutaneous injection, followed by a second subsequent dose of 80 mg of adalimumab as a subcutaneous injection, followed by a third subsequent dose of 40 mg of adalimumab at about four weeks after the initial dose; and
 - c. an initial dose of 15 mg of immunomodulator orally wherein the dosing regimen results in endoscopic remission, further wherein the anti- α 4 β 7 antibody comprises an antigen binding region of nonhuman origin and at least a portion of an antibody of human origin, wherein the humanized antibody has binding specificity for the α 4 β 7 complex, wherein the antigen-binding region comprises the CDRs:

Light chain: CDR1 SEQ ID NO:7
CDR2 SEQ ID NO:8 and
CDR3 SEQ ID NO:9; and

Heavy chain: CDR1 SEQ ID NO:4
CDR2 SEQ ID NO:5 and
CDR3 SEQ ID NO:6.
2. The method of claim 1, wherein the humanized anti- α 4 β 7 antibody is administered to the patient over about 30 minutes.

3. The method of claim 1 or 2, wherein the humanized anti- $\alpha 4\beta 7$ antibody has a heavy chain variable region sequence of amino acids 20 to 140 of SEQ ID NO:1.
4. The method of any one of the preceding claims, wherein the humanized anti- $\alpha 4\beta 7$ antibody has a light chain variable region sequence of amino acids 20 to 131 of SEQ ID NO:2.
5. The method of any one of the preceding claims, wherein the humanized anti- $\alpha 4\beta 7$ antibody has a heavy chain comprising amino acids 20 to 470 of SEQ ID NO:1 and a light chain comprising amino acids 20 to 238 of SEQ ID NO:2.
6. The method of any one of the preceding claims, wherein the anti- $\alpha 4\beta 7$ antibody is vedolizumab.
7. The method of any one of the preceding claims, wherein the immunomodulator is methotrexate.
8. The method of any one of the preceding claims, wherein the high risk CD patient additionally is administered a dose of corticosteroid.
9. The method of claim 8, wherein the corticosteroid is prednisone or budesonide.
10. The method of claim 8 or 9, further comprising tapering the corticosteroid dose to discontinuation.
11. The method of any one of the preceding claims, further comprising administration of a subsequent dose of 300 mg of the humanized anti- $\alpha 4\beta 7$ antibody as an intravenous infusion at about four weeks after the third subsequent dose.
12. The method of any one of claims 1 to 10, further comprising administration of a subsequent dose of 300 mg of the humanized anti- $\alpha 4\beta 7$ antibody as an intravenous infusion at about eight weeks after the third subsequent dose.
13. The method of claim 11 or 12, further comprising administration of a subsequent dose

of 300 mg of the humanized anti- α 4 β 7 antibody as an intravenous infusion at about every eight weeks thereafter.

14. The method of any one of claims 1 to 13, further comprising administration of a subsequent dose of 300 mg of the humanized anti- α 4 β 7 antibody as an intravenous infusion at about every four weeks after 26 weeks of therapy.
15. The method of any one of the claims 1 to 10, further comprising administration of a subsequent dose of 108 mg of the humanized anti- α 4 β 7 antibody as a subcutaneous injection at about every two or four weeks after the third subsequent dose.
16. A method of treating a patient having an inflammatory bowel disease, comprising administering to the patient vedolizumab, adalimumab, and methotrexate according to a triple combination therapy, wherein the triple combination therapy comprises
administering vedolizumab at a dose of 300 mg at weeks 0, 2, and 6,
followed by administration every 8 weeks thereafter;
subcutaneously administering a 160 mg dose of adalimumab at week 0,
an 80 mg dose of adalimumab at week 2, and a 40 mg dose of adalimumab at
week 4 and every two weeks thereafter; and
administering methotrexate.
17. The method of claim 16, wherein adalimumab is discontinued at week 26.
18. A method of treating a patient having an inflammatory bowel disease, comprising administering to the patient vedolizumab, infliximab, and methotrexate according to a triple combination therapy, wherein the triple combination therapy comprises
administering vedolizumab at a dose of 300 mg at weeks 0, 2, and 6,
followed by administration every 8 weeks thereafter;
intravenously administering infliximab at a dose of 5 mg/kg at weeks
0, 2 and 6, and then every 8 weeks thereafter; and
administering methotrexate.
19. The method of any one of claims 16-18, wherein the methotrexate is administered orally.

20. The method of claim 19, wherein the methotrexate is administered at a dose of 15 mg.
21. The method of any one of claims 16-20, wherein the methotrexate is discontinued at week 34.
22. The method of any one of claims 16-21, wherein the inflammatory bowel disease is Crohn's disease.
23. The method of claim 22, wherein the Crohn's disease is moderately to severely active Crohn's disease.
24. The method of claim 22 or 23, wherein the patient is a high risk Crohn's disease patient.
25. The method of claim 24, wherein the patient is selected as a high risk Crohn's disease patient prior to administration of the triple combination therapy.
26. The method of any one of claims 16-21, wherein the inflammatory bowel disease is ulcerative colitis.
27. The method of claim 26, wherein the ulcerative colitis is moderately to severely active ulcerative colitis.
28. A method of treating a patient having an inflammatory bowel disease, comprising administering to the patient vedolizumab, golimumab, and methotrexate according to a triple combination therapy, wherein the triple combination therapy comprises
administering vedolizumab at a dose of 300 mg at weeks 0, 2, and 6, followed by administration every 8 weeks thereafter;
subcutaneously administering golimumab at a dose of 200 mg at week 0, 100 mg at week 2, and 100 mg every four weeks thereafter; and
administering methotrexate.
29. The method of claim 28, wherein the methotrexate is administered orally.
30. The method of claim 28, wherein the methotrexate is administered at a dose of 15 mg.

31. The method of any one of claims 28-30, wherein the methotrexate is discontinued at week 34.
32. The method of any one of claims 28-31, wherein the inflammatory bowel disease is Crohn's disease.
33. The method of claim 32, wherein the Crohn's disease is moderately to severely active Crohn's disease.
34. The method of claim 32 or 33, wherein the patient is a high risk Crohn's disease patient.
35. The method of claim 34, wherein the patient is selected as a high risk Crohn's disease patient prior to administration of the triple combination therapy.
36. The method of any one of claims 28-31, wherein the inflammatory bowel disease is ulcerative colitis.
37. The method of claim 36, wherein the ulcerative colitis is moderately to severely active ulcerative colitis.
38. A method of treating a patient having an inflammatory bowel disease, comprising administering to the patient vedolizumab, certolizumab pegol, and methotrexate according to a triple combination therapy, wherein the triple combination therapy comprises
 - administering vedolizumab at a dose of 300 mg at weeks 0, 2, and 6, followed by administration every 8 weeks thereafter;
 - subcutaneously administering certolizumab at a dose of 400 mg at week 0, 400 mg at weeks 2 and 4, and optionally 400 mg every four weeks thereafter; and
 - administering methotrexate.
39. The method of claim 38, wherein the methotrexate is administered orally.
40. The method of claim 38, wherein the methotrexate is administered at a dose of 15 mg.

41. The method of any one of claims 38-40, wherein the methotrexate is discontinued at week 34.
42. The method of any one of claims 38-41, wherein the inflammatory bowel disease is Crohn's disease.
43. The method of claim 42, wherein the Crohn's disease is moderately to severely active Crohn's disease.
44. The method of claim 42 or 43, wherein the patient is a high risk Crohn's disease patient.
45. The method of claim 44, wherein the patient is selected as a high risk Crohn's disease patient prior to administration of the triple combination therapy.
46. The method of any one of claims 38-41, wherein the inflammatory bowel disease is ulcerative colitis.
47. The method of claim 46, wherein the ulcerative colitis is moderately to severely active ulcerative colitis.
48. The method of any one of claims 16-47, wherein a corticosteroid dose administered to the patient is tapered to discontinuation.
49. A method of treating a patient having an inflammatory bowel disease, comprising administering to the patient vedolizumab, adalimumab, and methotrexate according to a triple combination therapy, wherein the triple combination therapy comprises
administering vedolizumab at a dose of 300 mg at weeks 0, 2, 6, 14, 22, and 30, followed by administration every 4 weeks thereafter;
subcutaneously administering a 160 mg dose of adalimumab at week 0, an 80 mg dose of adalimumab at week 2, and a 40 mg dose of adalimumab at week 4 and every two weeks thereafter; and
administering methotrexate.
50. The method of claim 49, wherein adalimumab is discontinued at week 26.

51. A method of treating a patient having an inflammatory bowel disease, comprising administering to the patient vedolizumab, infliximab, and methotrexate according to a triple combination therapy, wherein the triple combination therapy comprises
 - administering vedolizumab at a dose of 300 mg at weeks 0, 2, 6, 14, 22 and 30, followed by administration every 4 weeks thereafter;
 - intravenously administering infliximab at a dose of 5 mg/kg at weeks 0, 2 and 6, and then every 8 weeks thereafter; and
 - administering methotrexate.
52. The method of any one of claims 49-51, wherein the methotrexate is administered orally.
53. The method of claim 52, wherein the methotrexate is administered at a dose of 15 mg.
54. The method of any one of claims 49-53, wherein the methotrexate is discontinued at week 34.
55. The method of any one of claims 49-54, wherein the inflammatory bowel disease is Crohn's disease.
56. The method of claim 55, wherein the Crohn's disease is moderately to severely active Crohn's disease.
57. The method of claim 55 or 56, wherein the patient is a high risk Crohn's disease patient.
58. The method of claim 57, wherein the patient is selected as a high risk Crohn's disease patient prior to administration of the triple combination therapy.
59. The method of any one of claims 49-58, wherein a corticosteroid dose administered to the patient is tapered to discontinuation.




FIG. 1

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/031089

A. CLASSIFICATION OF SUBJECT MATTER
 INV. A61K38/00 A61K39/395 C07K16/24 C07K16/28
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>JUSTIN CÔTÉ-DAIGNEAULT ET AL: "Biologics in inflammatory bowel disease: what are the data?", UNITED EUROPEAN GASTROENTEROLOGY JOURNAL UEG WEEK 2015 ORAL PRESENTATIONS, vol. 3, no. 5, 9 June 2015 (2015-06-09), pages 419-428, XP055393744, ISSN: 2050-6406, DOI: 10.1177/2050640615590302 figure A11; example A11; table A11</p> <p>-----</p> <p style="text-align: center;">-/-</p>	1-59



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
25 July 2017	02/08/2017

Name and mailing address of the ISA/
 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040,
 Fax: (+31-70) 340-3016

Authorized officer

Fellows, Edward

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/031089

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>GLEN S. HAZLEWOOD ET AL: "Comparative Effectiveness of Immunosuppressants and Biologics for Inducing and Maintaining Remission in Crohn's Disease: A Network Meta-analysis", <i>GASTROENTEROLOGY</i>, vol. 148, no. 2, 1 February 2015 (2015-02-01), pages 344-354.e5, XP055393745, AMSTERDAM, NL ISSN: 0016-5085, DOI: 10.1053/j.gastro.2014.10.011 figure A11; example A11; table A11 -----</p>	1-59
X	<p>LEON P MCLEAN ET AL: "Vedolizumab for the treatment of ulcerative colitis and Crohn's disease", <i>IMMUNOTHERAPY</i>, vol. 4, no. 9, 1 September 2012 (2012-09-01), pages 883-898, XP055393761, GB ISSN: 1750-743X, DOI: 10.2217/imt.12.85 figure A11; example A11; table A11 -----</p>	1-59
1		