



(86) Date de dépôt PCT/PCT Filing Date: 2013/03/14
(87) Date publication PCT/PCT Publication Date: 2013/09/19
(45) Date de délivrance/Issue Date: 2020/06/23
(85) Entrée phase nationale/National Entry: 2014/09/11
(86) N° demande PCT/PCT Application No.: US 2013/031217
(87) N° publication PCT/PCT Publication No.: 2013/138560
(30) Priorités/Priorities: 2012/03/15 (US61/611,428);
2012/10/18 (US61/715,327)

(51) Cl.Int./Int.Cl. A61K 31/4985 (2006.01),
A61K 35/00 (2006.01)

(72) Inventeurs/Inventors:
XU, SHUICHAN, US;
HEGE, KRISTEN MAE, US;
RAYMON, HEATHER, US;
TSUJI, TOSHIYA, US;
SAPINOSO, LISA, US

(73) Propriétaire/Owner:
SIGNAL PHARMACEUTICALS, LLC, US

(74) Agent: SMART & BIGGAR LLP

(54) Titre : TRAITEMENT DU CANCER AVEC DES INHIBITEURS DE LA KINASE TOR

(54) Title: TREATMENT OF CANCER WITH TOR KINASE INHIBITORS

(57) Abrégé/Abstract:

Provided herein are methods for treating or preventing ETS overexpressing castration-resistant prostate cancer, comprising administering an effective amount of a TOR kinase inhibitor to a patient having ETS overexpressing castration-resistant prostate cancer.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number

WO 2013/138560 A1

(43) International Publication Date
19 September 2013 (19.09.2013)

(51) International Patent Classification:
A61K 31/4985 (2006.01) *A61K 35/00* (2006.01)

(21) International Application Number:
PCT/US2013/031217

(22) International Filing Date:
14 March 2013 (14.03.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/611,428 15 March 2012 (15.03.2012) US
61/715,327 18 October 2012 (18.10.2012) US

(71) Applicant: SIGNAL PHARMACEUTICALS, LLC [US/US]; 10300 Campus Point Drive, Suite 100, San Diego, CA 92121 (US).

(72) Inventors: XU, Shuichan; 9650 Deer Trail Place, San Diego, CA 92127 (US). HEGE, Kristen, Mae; 616 Howard Ave., Burlingame, CA 94010 (US). RAYMON, Heather; 3520 Vista de la Orilla, San Diego, CA 92117 (US). TSUJI, Toshiya; 8155 Cargill Av #34, San Diego, CA 92122 (US). SAPINOSO, Lisa; 14597 Via Bettone, San Diego, CA 92127 (US).

(74) Agents: INSOGNA, Anthony, M. et al.; Jones Day, 222 East 41st Street, New York, NY 10017-6702 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2013/138560 A1

(54) Title: TREATMENT OF CANCER WITH TOR KINASE INHIBITORS

(57) Abstract: Provided herein are methods for treating or preventing ETS overexpressing castration-resistant prostate cancer, comprising administering an effective amount of a TOR kinase inhibitor to a patient having ETS overexpressing castration-resistant prostate cancer.

TREATMENT OF CANCER WITH TOR KINASE INHIBITORS

[0001] This application claims the benefit of U.S. Provisional Application No. 61/611,428, filed March 15, 2012 and claims the benefit of U.S. Provisional Application No. 61/715,327, filed October 18, 2012.

1. **FIELD**

[0002] Provided herein are methods for treating or preventing ETS overexpressing castration-resistant prostate cancer, comprising administering an effective amount of a TOR kinase inhibitor to a patient having ETS overexpressing castration-resistant prostate cancer.

2. **BACKGROUND**

[0003] The connection between abnormal protein phosphorylation and the cause or consequence of diseases has been known for over 20 years. Accordingly, protein kinases have become a very important group of drug targets. *See Cohen, Nature*, 1:309-315 (2002). Various protein kinase inhibitors have been used clinically in the treatment of a wide variety of diseases, such as cancer and chronic inflammatory diseases, including diabetes and stroke. *See Cohen, Eur. J. Biochem.*, 268:5001-5010 (2001), *Protein Kinase Inhibitors for the Treatment of Disease: The Promise and the Problems*, Handbook of Experimental Pharmacology, Springer Berlin Heidelberg, 167 (2005).

[0004] The protein kinases are a large and diverse family of enzymes that catalyze protein phosphorylation and play a critical role in cellular signaling. Protein kinases may exert positive or negative regulatory effects, depending upon their target protein. Protein kinases are involved in specific signaling pathways which regulate cell functions such as, but not limited to, metabolism, cell cycle progression, cell adhesion, vascular function, apoptosis, and angiogenesis. Malfunctions of cellular signaling have been associated with many diseases, the most characterized of which include cancer and diabetes. The regulation of signal transduction by cytokines and the association of signal molecules with

protooncogenes and tumor suppressor genes have been well documented. Similarly, the connection between diabetes and related conditions, and deregulated levels of protein kinases, has been demonstrated. See e.g., Sridhar *et al.* *Pharmaceutical Research*, 17(11):1345-1353 (2000). Viral infections and the conditions related thereto have also been associated with the regulation of protein kinases. Park *et al.* *Cell* 101 (7): 777-787 (2000).

[0005] Because protein kinases regulate nearly every cellular process, including metabolism, cell proliferation, cell differentiation, and cell survival, they are attractive targets for therapeutic intervention for various disease states. For example, cell-cycle control and angiogenesis, in which protein kinases play a pivotal role are cellular processes associated with numerous disease conditions such as but not limited to cancer, inflammatory diseases, abnormal angiogenesis and diseases related thereto, atherosclerosis, macular degeneration, diabetes, obesity, and pain.

[0006] Protein kinases have become attractive targets for the treatment of cancers. Fabbro *et al.*, *Pharmacology & Therapeutics* 93:79-98 (2002). It has been proposed that the involvement of protein kinases in the development of human malignancies may occur by: (1) genomic rearrangements (e.g., BCR-ABL in chronic myelogenous leukemia), (2) mutations leading to constitutively active kinase activity, such as acute myelogenous leukemia and gastrointestinal tumors, (3) deregulation of kinase activity by activation of oncogenes or loss of tumor suppressor functions, such as in cancers with oncogenic RAS, (4) deregulation of kinase activity by over-expression, as in the case of EGFR and (5) ectopic expression of growth factors that can contribute to the development and maintenance of the neoplastic phenotype. Fabbro *et al.*, *Pharmacology & Therapeutics* 93:79-98 (2002).

[0007] The elucidation of the intricacy of protein kinase pathways and the complexity of the relationship and interaction among and between the various protein kinases and kinase pathways highlights the importance of developing pharmaceutical agents capable of acting as protein kinase modulators, regulators or inhibitors that have beneficial

activity on multiple kinases or multiple kinase pathways. Accordingly, there remains a need for new kinase modulators.

[0008] The protein named mTOR (mammalian target of rapamycin), which is also called FRAP, RAFT1 or RAPT1, is a 2549-amino acid Ser/Thr protein kinase, that has been shown to be one of the most critical proteins in the mTOR/PI3K/Akt pathway that regulates cell growth and proliferation. Georgakis and Younes *Expert Rev. Anticancer Ther.* 6(1):131-140 (2006). mTOR exists within two complexes, mTORC1 and mTORC2. While mTORC1 is sensitive to rapamycin analogs (such as temsirolimus or everolimus), mTORC2 is largely rapamycin-insensitive. Notably, rapamycin is not a TOR kinase inhibitor. Several mTOR inhibitors have been or are being evaluated in clinical trials for the treatment of cancer. Temsirolimus was approved for use in renal cell carcinoma in 2007 and sirolimus was approved in 1999 for the prophylaxis of renal transplant rejection. Everolimus was approved in 2009 for renal cell carcinoma patients that have progressed on vascular endothelial growth factor receptor inhibitors, in 2010 for subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis (TS) in patients who require therapy but are not candidates for surgical resection, and in 2011 for progressive neuroendocrine tumors of pancreatic origin (PNET) in patients with unresectable, locally advanced or metastatic disease. There remains a need for additional TOR kinase inhibitors.

[0009] Citation or identification of any reference in Section 2 of this application is not to be construed as an admission that the reference is prior art to the present application.

3. SUMMARY

[0010] Provided herein are methods for treating or preventing E-twenty six (ETS) overexpressing castration-resistant prostate cancer comprising administering an effective amount of a TOR kinase inhibitor to a patient having ETS overexpressing castration-resistant prostate cancer.

[0011] In certain embodiments, provided herein are methods for improving the Prostate-Specific Antigen Working Group 2 (PSAWG2) Criteria for prostate cancer of a

patient, comprising administering an effective amount of a TOR kinase inhibitor to a patient having ETS overexpressing castration-resistant prostate cancer.

[0012] In some embodiments, the TOR kinase inhibitor is a compound as described herein.

[0012a] The present application as claimed relates to:

- use of 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof in the manufacture of a medicament for treating E-twenty six (ETS) overexpressing castration-resistant prostate cancer; and

- use of 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof for treating E-twenty six (ETS) overexpressing castration-resistant prostate cancer.

[0013] The present embodiments can be understood more fully by reference to the detailed description and examples, which are intended to exemplify non-limiting embodiments.

4. DETAILED DESCRIPTION

4.1 DEFINITIONS

[0014] An "alkyl" group is a saturated, partially saturated, or unsaturated straight chain or branched non-cyclic hydrocarbon having from 1 to 10 carbon atoms, typically from 1 to 8 carbons or, in some embodiments, from 1 to 6, 1 to 4, or 2 to 6 or carbon atoms. Representative alkyl groups include -methyl, -ethyl, -n-propyl, -n-butyl, -n-pentyl and -n-hexyl; while saturated branched alkyls include -isopropyl, -*sec*-butyl, -isobutyl, -*tert*-butyl, -isopentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2,3-dimethylbutyl and the like. Examples of unsaturated alkyl groups include, but are not limited to, vinyl, allyl, -CH=CH(CH₃), -CH=C(CH₃)₂, -C(CH₃)=CH₂, -C(CH₃)=CH(CH₃), -C(CH₂CH₃)=CH₂, -C≡CH, -C≡C(CH₃), -C≡C(CH₂CH₃), -CH₂C≡CH, -CH₂C≡C(CH₃) and -CH₂C≡C(CH₇CH₃), among others. An alkyl group can be substituted or unsubstituted. In certain embodiments,

when the alkyl groups described herein are said to be "substituted," they may be substituted with any substituent or substituents as those found in the exemplary compounds and embodiments disclosed herein, as well as halogen (chloro, iodo, bromo, or fluoro); hydroxyl; alkoxy; alkoxyalkyl; amino; alkylamino; carboxy; nitro; cyano; thiol; thioether; imine; imide; amidine; guanidine; enamine; aminocarbonyl; acylamino; phosphonato; phosphine; thiocarbonyl; sulfonyl; sulfone; sulfonamide; ketone; aldehyde; ester; urea; urethane; oxime; hydroxyl amine; alkoxyamine; aralkoxyamine; N-oxide;

hydrazine; hydrazide; hydrazone; azide; isocyanate; isothiocyanate; cyanate; thiocyanate; B(OH)₂, or O(alkyl)aminocarbonyl.

[0015] An “alkenyl” group is a straight chain or branched non-cyclic hydrocarbon having from 2 to 10 carbon atoms, typically from 2 to 8 carbon atoms, and including at least one carbon-carbon double bond. Representative straight chain and branched (C₂-C₈)alkenyls include -vinyl, -allyl, -1-butenyl, -2-butenyl, -isobutylenyl, -1-pentenyl, -2-pentenyl, -3-methyl-1-butenyl, -2-methyl-2-butenyl, -2,3-dimethyl-2-butenyl, -1-hexenyl, -2-hexenyl, -3-hexenyl, -1-heptenyl, -2-heptenyl, -3-heptenyl, -1-octenyl, -2-octenyl, -3-octenyl and the like. The double bond of an alkenyl group can be unconjugated or conjugated to another unsaturated group. An alkenyl group can be unsubstituted or substituted.

[0016] A “cycloalkyl” group is a saturated, partially saturated, or unsaturated cyclic alkyl group of from 3 to 10 carbon atoms having a single cyclic ring or multiple condensed or bridged rings which can be optionally substituted with from 1 to 3 alkyl groups. In some embodiments, the cycloalkyl group has 3 to 8 ring members, whereas in other embodiments the number of ring carbon atoms ranges from 3 to 5, 3 to 6, or 3 to 7. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, 1-methylcyclopropyl, 2-methylcyclopentyl, 2-methylcyclooctyl, and the like, or multiple or bridged ring structures such as adamantyl and the like. Examples of unsaturated cycloalkyl groups include cyclohexenyl, cyclopentenyl, cyclohexadienyl, butadienyl, pentadienyl, hexadienyl, among others. A cycloalkyl group can be substituted or unsubstituted. Such substituted cycloalkyl groups include, by way of example, cyclohexanone and the like.

[0017] An “aryl” group is an aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring (e.g., phenyl) or multiple condensed rings (e.g., naphthyl or anthryl). In some embodiments, aryl groups contain 6-14 carbons, and in others from 6 to 12 or even 6 to 10 carbon atoms in the ring portions of the groups. Particular aryls include phenyl, biphenyl, naphthyl and the like. An aryl group can be substituted or unsubstituted.

The phrase “aryl groups” also includes groups containing fused rings, such as fused aromatic-aliphatic ring systems (e.g., indanyl, tetrahydronaphthyl, and the like).

[0018] A “heteroaryl” group is an aryl ring system having one to four heteroatoms as ring atoms in a heteroaromatic ring system, wherein the remainder of the atoms are carbon atoms. In some embodiments, heteroaryl groups contain 5 to 6 ring atoms, and in others from 6 to 9 or even 6 to 10 atoms in the ring portions of the groups. Suitable heteroatoms include oxygen, sulfur and nitrogen. In certain embodiments, the heteroaryl ring system is monocyclic or bicyclic. Non-limiting examples include but are not limited to, groups such as pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl, pyrrolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, thiophenyl, benzothiophenyl, furanyl, benzofuranyl (for example, isobenzofuran-1,3-diimine), indolyl, azaindolyl (for example, pyrrolopyridyl or 1H-pyrrolo[2,3-b]pyridyl), indazolyl, benzimidazolyl (for example, 1H-benzo[d]imidazolyl), imidazopyridyl (for example, azabenzimidazolyl, 3H-imidazo[4,5-b]pyridyl or 1H-imidazo[4,5-b]pyridyl), pyrazolopyridyl, triazolopyridyl, benzotriazolyl, benzoxazolyl, benzothiazolyl, benzothiadiazolyl, isoxazolopyridyl, thianaphthalenyl, purinyl, xanthinyl, adeninyl, guaninyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, quinoxalinyl, and quinazolinyl groups.

[0019] A “heterocyclyl” is an aromatic (also referred to as heteroaryl) or non-aromatic cycloalkyl in which one to four of the ring carbon atoms are independently replaced with a heteroatom from the group consisting of O, S and N. In some embodiments, heterocyclyl groups include 3 to 10 ring members, whereas other such groups have 3 to 5, 3 to 6, or 3 to 8 ring members. Heterocyclyls can also be bonded to other groups at any ring atom (*i.e.*, at any carbon atom or heteroatom of the heterocyclic ring). A heterocyclylalkyl group can be substituted or unsubstituted. Heterocyclyl groups encompass unsaturated, partially saturated and saturated ring systems, such as, for example, imidazolyl, imidazolinyl and imidazolidinyl groups. The phrase heterocyclyl includes fused ring species, including those comprising fused aromatic and non-aromatic groups, such as, for example, benzotriazolyl, 2,3-dihydrobenzo[1,4]dioxinyl, and benzo[1,3]dioxolyl. The phrase also

includes bridged polycyclic ring systems containing a heteroatom such as, but not limited to, quinuclidyl. Representative examples of a heterocyclyl group include, but are not limited to, aziridinyl, azetidinyl, pyrrolidyl, imidazolidinyl, pyrazolidinyl, thiazolidinyl, tetrahydrothiophenyl, tetrahydrofuranyl, dioxolyl, furanyl, thiophenyl, pyrrolyl, pyrrolinyl, imidazolyl, imidazolinyl, pyrazolyl, pyrazolinyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl, thiazolinyl, isothiazolyl, thiadiazolyl, oxadiazolyl, piperidyl, piperazinyl, morpholinyl, thiomorpholinyl, tetrahydropyranyl (for example, tetrahydro-2H-pyranyl), tetrahydrothiopyranyl, oxathiane, dioxy, dithianyl, pyranyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, triazinyl, dihydropyridyl, dihydrotiinyl, dihydrotithionyl, homopiperazinyl, quinuclidyl, indolyl, indolinyl, isoindolyl, azaindolyl (pyrrolopyridyl), indazolyl, indolizinyl, benzotriazolyl, benzimidazolyl, benzofuranyl, benzothiophenyl, benzthiazolyl, benzoxadiazolyl, benzoxazinyl, benzodithiinyl, benzoxathiinyl, benzothiazinyl, benzoxazolyl, benzothiazolyl, benzothiadiazolyl, benzo[1,3]dioxolyl, pyrazolopyridyl, imidazopyridyl (azabenzimidazolyl; for example, 1H-imidazo[4,5-b]pyridyl, or 1H-imidazo[4,5-b]pyridin-2(3H)-onyl), triazolopyridyl, isoxazolopyridyl, purinyl, xanthinyl, adeninyl, guaninyl, quinolinyl, isoquinolinyl, quinolizinyl, quinoxalinyl, quinazolinyl, cinnolinyl, phthalazinyl, naphthyridinyl, pteridinyl, thianaphthalenyl, dihydrobenzothiazinyl, dihydrobenzofuranyl, dihydroindolyl, dihydrobenzodioxinyl, tetrahydroindolyl, tetrahydroindazolyl, tetrahydrobenzimidazolyl, tetrahydrobenzotriazolyl, tetrahydropyrrolopyridyl, tetrahydropyrazolopyridyl, tetrahydroimidazopyridyl, tetrahydrotriazolopyridyl, and tetrahydroquinolinyl groups. Representative substituted heterocyclyl groups may be mono- substituted or substituted more than once, such as, but not limited to, pyridyl or morpholinyl groups, which are 2-, 3-, 4-, 5-, or 6-substituted, or disubstituted with various substituents such as those listed below.

[0020] An “cycloalkylalkyl” group is a radical of the formula: -alkyl-cycloalkyl, wherein alkyl and cycloalkyl are defined above. Substituted cycloalkylalkyl groups may be substituted at the alkyl, the cycloalkyl, or both the alkyl and the cycloalkyl portions of the group. Representative cycloalkylalkyl groups include but are not limited to

cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, and cyclohexylpropyl. Representative substituted cycloalkylalkyl groups may be mono-substituted or substituted more than once.

[0021] An “aralkyl” group is a radical of the formula: -alkyl-aryl, wherein alkyl and aryl are defined above. Substituted aralkyl groups may be substituted at the alkyl, the aryl, or both the alkyl and the aryl portions of the group. Representative aralkyl groups include but are not limited to benzyl and phenethyl groups and fused (cycloalkylaryl)alkyl groups such as 4-ethyl-indanyl.

[0022] A “heterocyclalkyl” group is a radical of the formula: -alkyl-heterocyclyl, wherein alkyl and heterocyclyl are defined above. Substituted heterocyclalkyl groups may be substituted at the alkyl, the heterocyclyl, or both the alkyl and the heterocyclyl portions of the group. Representative heterocyclalkyl groups include but are not limited to 4-ethyl-morpholinyl, 4-propylmorpholinyl, furan-2-yl methyl, furan-3-yl methyl, pyrdine-3-yl methyl, (tetrahydro-2H-pyran-4-yl)methyl, (tetrahydro-2H-pyran-4-yl)ethyl, tetrahydrofuran-2-yl methyl, tetrahydrofuran-2-yl ethyl, and indol-2-yl propyl.

[0023] A “halogen” is fluorine, chlorine, bromine or iodine.

[0024] A “hydroxyalkyl” group is an alkyl group as described above substituted with one or more hydroxy groups.

[0025] An “alkoxy” group is -O-(alkyl), wherein alkyl is defined above.

[0026] An “alkoxyalkyl” group is -(alkyl)-O-(alkyl), wherein alkyl is defined above.

[0027] An “amino” group is a radical of the formula: -NH₂.

[0028] An “alkylamino” group is a radical of the formula: -NH-alkyl or -N(alkyl)₂, wherein each alkyl is independently as defined above.

[0029] A “carboxy” group is a radical of the formula: -C(O)OH.

[0030] An “aminocarbonyl” group is a radical of the formula: -C(O)N(R[#])₂, -C(O)NH(R[#]) or -C(O)NH₂, wherein each R[#] is independently a substituted or unsubstituted alkyl, cycloalkyl, aryl, aralkyl, heterocyclyl or heterocyclyl group as defined herein.

[0031] An “acylamino” group is a radical of the formula: -NHC(O)(R[#]) or -N(alkyl)C(O)(R[#]), wherein each alkyl and R[#] are independently as defined above.

[0032] An “alkylsulfonylamino” group is a radical of the formula: -NHSO₂(R[#]) or -N(alkyl)SO₂(R[#]), wherein each alkyl and R[#] are defined above.

[0033] A “urea” group is a radical of the formula: -N(alkyl)C(O)N(R[#])₂, -N(alkyl)C(O)NH(R[#]), -N(alkyl)C(O)NH₂, -NHC(O)N(R[#])₂, -NHC(O)NH(R[#]), or -NH(CO)NHR[#], wherein each alkyl and R[#] are independently as defined above.

[0034] When the groups described herein, with the exception of alkyl group are said to be “substituted,” they may be substituted with any appropriate substituent or substituents. Illustrative examples of substituents are those found in the exemplary compounds and embodiments disclosed herein, as well as halogen (chloro, iodo, bromo, or fluoro); alkyl; hydroxyl; alkoxy; alkoxyalkyl; amino; alkylamino; carboxy; nitro; cyano; thiol; thioether; imine; imide; amidine; guanidine; enamine; aminocarbonyl; acylamino; phosphonato; phosphine; thiocarbonyl; sulfonyl; sulfone; sulfonamide; ketone; aldehyde; ester; urea; urethane; oxime; hydroxyl amine; alkoxyamine; aralkoxyamine; N-oxide; hydrazine; hydrazide; hydrazone; azide; isocyanate; isothiocyanate; cyanate; thiocyanate; oxygen (=O); B(OH)₂, O(alkyl)aminocarbonyl; cycloalkyl, which may be monocyclic or fused or non-fused polycyclic (e.g., cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl), or a heterocyclyl, which may be monocyclic or fused or non-fused polycyclic (e.g., pyrrolidyl, piperidyl, piperazinyl, morpholinyl, or thiazinyl); monocyclic or fused or non-fused polycyclic aryl or heteroaryl (e.g., phenyl, naphthyl, pyrrolyl, indolyl, furanyl, thiophenyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, triazolyl, tetrazolyl, pyrazolyl, pyridinyl, quinolinyl, isoquinolinyl, acridinyl, pyrazinyl, pyridazinyl, pyrimidinyl, benzimidazolyl, benzothiophenyl, or benzofuranyl) aryloxy; aralkyloxy; heterocyclyloxy; and heterocyclalkoxy.

[0035] As used herein, the term “pharmaceutically acceptable salt(s)” refers to a salt prepared from a pharmaceutically acceptable non-toxic acid or base including an inorganic acid and base and an organic acid and base. Suitable pharmaceutically acceptable base

addition salts of the TOR kinase inhibitors include, but are not limited to metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from lysine, N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. Suitable non-toxic acids include, but are not limited to, inorganic and organic acids such as acetic, alginic, anthranilic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethenesulfonic, formic, fumaric, furoic, galacturonic, gluconic, glucuronic, glutamic, glycolic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phenylacetic, phosphoric, propionic, salicylic, stearic, succinic, sulfanilic, sulfuric, tartaric acid, and p-toluenesulfonic acid. Specific non-toxic acids include hydrochloric, hydrobromic, phosphoric, sulfuric, and methanesulfonic acids.

Examples of specific salts thus include hydrochloride and mesylate salts. Others are well-known in the art, see for example, *Remington's Pharmaceutical Sciences*, 18th eds., Mack Publishing, Easton PA (1990) or *Remington: The Science and Practice of Pharmacy*, 19th eds., Mack Publishing, Easton PA (1995).

[0036] As used herein and unless otherwise indicated, the term "clathrate" means a TOR kinase inhibitor, or a salt thereof, in the form of a crystal lattice that contains spaces (e.g., channels) that have a guest molecule (e.g., a solvent or water) trapped within or a crystal lattice wherein a TOR kinase inhibitor is a guest molecule.

[0037] As used herein and unless otherwise indicated, the term "solvate" means a TOR kinase inhibitor, or a salt thereof, that further includes a stoichiometric or non-stoichiometric amount of a solvent bound by non-covalent intermolecular forces. In one embodiment, the solvate is a hydrate.

[0038] As used herein and unless otherwise indicated, the term "hydrate" means a TOR kinase inhibitor, or a salt thereof, that further includes a stoichiometric or non-stoichiometric amount of water bound by non-covalent intermolecular forces.

[0039] As used herein and unless otherwise indicated, the term "prodrug" means a TOR kinase inhibitor derivative that can hydrolyze, oxidize, or otherwise react under


biological conditions (*in vitro* or *in vivo*) to provide an active compound, particularly a TOR kinase inhibitor. Examples of prodrugs include, but are not limited to, derivatives and metabolites of a TOR kinase inhibitor that include biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate analogues. In certain embodiments, prodrugs of compounds with carboxyl functional groups are the lower alkyl esters of the carboxylic acid. The carboxylate esters are conveniently formed by esterifying any of the carboxylic acid moieties present on the molecule. Prodrugs can typically be prepared using well-known methods, such as those described by *Burger's Medicinal Chemistry and Drug Discovery* 6th ed. (Donald J. Abraham *ed.*, 2001, Wiley) and *Design and Application of Prodrugs* (H. Bundgaard *ed.*, 1985, Harwood Academic Publishers Gmfh).

[0040] As used herein and unless otherwise indicated, the term "stereoisomer" or "stereomerically pure" means one stereoisomer of a TOR kinase inhibitor that is substantially free of other stereoisomers of that compound. For example, a stereomerically pure compound having one chiral center will be substantially free of the opposite enantiomer of the compound. A stereomerically pure compound having two chiral centers will be substantially free of other diastereomers of the compound. A typical stereomerically pure compound comprises greater than about 80% by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomers of the compound, greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound, greater than about 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomers of the compound, or greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound. The TOR kinase inhibitors can have chiral centers and can occur as racemates, individual enantiomers or diastereomers, and mixtures thereof. All such isomeric forms are included within the embodiments disclosed herein, including mixtures thereof. The use of

stereomerically pure forms of such TOR kinase inhibitors, as well as the use of mixtures of those forms are encompassed by the embodiments disclosed herein. For example, mixtures comprising equal or unequal amounts of the enantiomers of a particular TOR kinase inhibitor may be used in methods and compositions disclosed herein. These isomers may be asymmetrically synthesized or resolved using standard techniques such as chiral columns or chiral resolving agents. *See, e.g.*, Jacques, J., *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley-Interscience, New York, 1981); Wilen, S. H., *et al.*, *Tetrahedron* 33:2725 (1977); Eliel, E. L., *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); and Wilen, S. H., *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN, 1972).

[0041] It should also be noted the TOR kinase inhibitors can include E and Z isomers, or a mixture thereof, and cis and trans isomers or a mixture thereof. In certain embodiments, the TOR kinase inhibitors are isolated as either the cis or trans isomer. In other embodiments, the TOR kinase inhibitors are a mixture of the cis and trans isomers.

[0042] “Tautomers” refers to isomeric forms of a compound that are in equilibrium with each other. The concentrations of the isomeric forms will depend on the environment the compound is found in and may be different depending upon, for example, whether the compound is a solid or is in an organic or aqueous solution. For example, in aqueous solution, pyrazoles may exhibit the following isomeric forms, which are referred to as tautomers of each other:



[0043] As readily understood by one skilled in the art, a wide variety of functional groups and other structures may exhibit tautomerism and all tautomers of the TOR kinase inhibitors are within the scope of the present invention.

[0044] It should also be noted the TOR kinase inhibitors can contain unnatural proportions of atomic isotopes at one or more of the atoms. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125

(¹²⁵I), sulfur-35 (³⁵S), or carbon-14 (¹⁴C), or may be isotopically enriched, such as with deuterium (²H), carbon-13 (¹³C), or nitrogen-15 (¹⁵N). As used herein, an “isotopologue” is an isotopically enriched compound. The term “isotopically enriched” refers to an atom having an isotopic composition other than the natural isotopic composition of that atom. “Isotopically enriched” may also refer to a compound containing at least one atom having an isotopic composition other than the natural isotopic composition of that atom. The term “isotopic composition” refers to the amount of each isotope present for a given atom. Radiolabeled and isotopically enriched compounds are useful as therapeutic agents, e.g., cancer and inflammation therapeutic agents, research reagents, e.g., binding assay reagents, and diagnostic agents, e.g., in vivo imaging agents. All isotopic variations of the TOR kinase inhibitors as described herein, whether radioactive or not, are intended to be encompassed within the scope of the embodiments provided herein. In some embodiments, there are provided isotopologues of the TOR kinase inhibitors, for example, the isotopologues are deuterium, carbon-13, or nitrogen-15 enriched TOR kinase inhibitors.

[0045] “Treating” as used herein, means an alleviation, in whole or in part, of ETS overexpressing castration-resistant prostate cancer, or a symptom thereof, or slowing, or halting of further progression or worsening of ETS overexpressing castration-resistant prostate cancer.

[0046] “Preventing” as used herein, means the prevention of the onset, recurrence or spread, in whole or in part, of ETS overexpressing castration-resistant prostate cancer, or a symptom thereof.

[0047] The term “effective amount” in connection with an TOR kinase inhibitor means an amount capable of alleviating, in whole or in part, symptoms associated with ETS overexpressing castration-resistant prostate cancer, or slowing or halting further progression or worsening of those symptoms, or treating or preventing ETS overexpressing castration-resistant prostate cancer. The effective amount of the TOR kinase inhibitor, for example in a pharmaceutical composition, may be at a level that will exercise the desired effect; for example, about 0.005 mg/kg of a subject’s body weight to about 100 mg/kg of a patient’s

body weight in unit dosage for both oral and parenteral administration. As will be apparent to those skilled in the art, it is to be expected that the effective amount of a TOR kinase inhibitor disclosed herein may vary depending on the severity of the indication being treated.

[0048] The terms "patient" and "subject" as used herein include an animal, including, but not limited to, an animal such as a cow, monkey, horse, sheep, pig, chicken, turkey, quail, cat, dog, mouse, rat, rabbit or guinea pig, in one embodiment a mammal, in another embodiment a human. In one embodiment, a "patient" or "subject" is a human having ETS overexpressing castration-resistant prostate cancer. In one embodiment, a patient is a human having histologically or cytologically-confirmed ETS overexpressing castration-resistant prostate cancer, including subjects who have progressed on (or not been able to tolerate) standard anticancer therapy or for whom no standard anticancer therapy exists.

[0049] In the context of ETS overexpressing castration-resistant prostate cancer, treatment may be assessed by inhibition of disease progression, inhibition of tumor growth, reduction of primary and/or secondary tumor(s), relief of tumor-related symptoms, improvement in quality of life, inhibition of tumor secreted factors (including prostate specific antigen or PSA), delayed appearance of primary and/or secondary tumor(s), slowed development of primary and/or secondary tumor(s), decreased occurrence of primary and/or secondary tumor(s), slowed or decreased severity of secondary effects of disease, arrested tumor growth and/or regression of tumors, among others. In certain embodiments, treatment of ETS overexpressing castration-resistant prostate cancer may be assessed by the inhibition of phosphorylation of S6RP, 4E-BP1 and/or AKT in circulating blood and/or tumor cells and/or skin biopsies or tumor biopsies/aspirates, before, during and/or after treatment with a TOR kinase inhibitor. In other embodiments, treatment of ETS overexpressing castration-resistant prostate cancer may be assessed by the inhibition of DNA-dependent protein kinase (DNA-PK) activity in skin samples and/or tumor biopsies/aspirates, such as by assessment of the amount of pDNA-PK S2056 as a biomarker

for DNA damage pathways before, during, and/or after TOR kinase inhibitor treatment. In one embodiment, the skin sample is irradiated by UV light. In the extreme, complete inhibition, is referred to herein as prevention or chemoprevention. In this context, the term “prevention” includes either preventing the onset of clinically evident ETS overexpressing castration-resistant prostate cancer altogether or preventing the onset of a preclinically evident stage of ETS overexpressing castration-resistant prostate cancer. Also intended to be encompassed by this definition is the prevention of transformation into malignant cells or to arrest or reverse the progression of premalignant cells to malignant cells. This includes prophylactic treatment of those at risk of developing ETS overexpressing castration-resistant prostate cancer.

4.2 BRIEF DESCRIPTION OF THE DRAWINGS

[0050] FIG. 1 A-C provides a clonogenic growth inhibition assay for Compound 1 with ETS positive and ETS negative cell lines.

[0051] FIG. 2 provides a caspase analysis apoptosis assay for Compound 1 with ETS positive and ETS negative cell lines.

[0052] FIG. 3 provides a cell invasion assay for ETS positive and ETS negative cell lines.


[0053] FIG. 4 provides a COMET assay for the potentiation of Compound 1 on DNA damage in ETS positive cells.

[0054] FIG. 5. provides anti-tumor activity of Compound 1 in a castration resistant LNCaP (LNCaP-HR) prostate cancer model.

4.3 TOR KINASE INHIBITORS

[0055] The compounds provided herein are generally referred to as “TOR kinase inhibitor(s).” In a specific embodiment, the TOR kinase inhibitors do not include rapamycin or rapamycin analogs (rapalogs).

[0056] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (I):



(I)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

X, Y and Z are at each occurrence independently N or CR³, wherein at least one of X, Y and Z is N and at least one of X, Y and Z is CR³;

-A-B-Q- taken together form -CHR⁴C(O)NH-, -C(O)CHR⁴NH-, -C(O)NH-, -CH₂C(O)O-, -C(O)CH₂O-, -C(O)O- or C(O)NR³;

L is a direct bond, NH or O;

R¹ is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted C₂₋₈alkenyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl;

R² is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl;

R³ is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclalkyl, -NHR⁴ or -N(R⁴)₂; and

R⁴ is at each occurrence independently substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[0057] In one embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -CH₂C(O)NH-.

[0058] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -C(O)CH₂NH-.

[0059] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -C(O)NH-.

[0060] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -CH₂C(O)O-.

[0061] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -C(O)CH₂O-.

[0062] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -C(O)O-.

[0063] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -C(O)NR³-.

[0064] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein Y is CR³.

[0065] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein X and Z are N and Y is CR³.

[0066] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein X and Z are N and Y is CH.

[0067] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein X and Z are CH and Y is N.

[0068] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein Y and Z are CH and X is N.

[0069] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein X and Y are CH and Z is N.

[0070] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein R¹ is substituted aryl, such as substituted phenyl.

[0071] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein R¹ is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl or substituted or unsubstituted naphthyl.

[0072] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted quinoline, substituted or unsubstituted pyridine, substituted or unsubstituted pyrimidine, substituted or unsubstituted indole, or substituted or unsubstituted thiophene.

[0073] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein R¹ is H.

[0074] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein R² is substituted C₁₋₈alkyl.

[0075] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein R² is methyl or ethyl substituted with substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[0076] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein R² is substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl.

[0077] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[0078] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein R² is H.

[0079] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein L is a direct bond.

[0080] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -C(O)NH-, X and Z are N and Y is CH, R¹ is substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl, L is a direct bond, and R² is substituted or unsubstituted C₁₋₈alkyl.

[0081] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -C(O)NH-, X and Z are N and Y is CH, R¹ is substituted or unsubstituted aryl, L is a direct bond, and R² is substituted or unsubstituted C₁₋₈alkyl.

[0082] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -C(O)NH-, X and Z are N and Y is CH, R¹ is substituted or unsubstituted aryl, and R² is C₁₋₈alkyl substituted with one or more substituents selected from alkoxy, amino, hydroxy, cycloalkyl, or heterocyclalkyl.

[0083] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -C(O)NH-, X and Z are N and Y is CH, R¹ is substituted or unsubstituted aryl, and R² is substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[0084] In another embodiment, the TOR kinase inhibitors of formula (I) are those wherein -A-B-Q- taken together form -C(O)NH-, X and Z are N and Y is CH, R¹ is substituted phenyl, L is a direct bond, and R² is substituted C₁₋₈alkyl.

[0085] In another embodiment, the TOR kinase inhibitors of formula (I) do not include compounds wherein X and Z are both N and Y is CH, -A-B-Q- is -C(O)NH-, L is a direct bond, R¹ is substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl, and R² is C₁₋₈alkyl substituted with substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl.

[0086] In another embodiment, the TOR kinase inhibitors of formula (I) do not include compounds wherein X and Z are both N and Y is CH, -A-B-Q- is -C(O)NH-, L is a direct bond, R¹ is phenyl, naphthyl, indanyl or biphenyl, each of which may be optionally substituted with one or more substituents independently selected from the group consisting substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted C₂₋₈alkenyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl.

[0087] In another embodiment, the TOR kinase inhibitors of formula (I) do not include compounds wherein X and Z are both N and Y is CH, -A-B-Q- is -C(O)NH-, L is a direct bond, R¹ is phenyl, naphthyl or biphenyl, each of which may be optionally substituted with one or more substituents each independently selected from the group consisting of C₁₋₄alkyl, amino, aminoC₁₋₁₂alkyl, halogen, hydroxy, hydroxyC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, -CF₃, C₁₋₁₂alkoxy, aryloxy, arylC₁₋₁₂alkoxy, -CN, -OCF₃, -COR_g, -COOR_g, -CONR_gR_h, -NR_gCOR_h, -SO₂R_g, -SO₃R_g or -SO₂NR_gR_h, wherein each R_g and R_h are independently selected from the group consisting of hydrogen, C₁₋₄alkyl, C₃₋₆cycloalkyl, aryl, arylC₁₋₆alkyl, heteroaryl or heteroarylC₁₋₆alkyl; or A is a 5- to 6-membered monocyclic heteroaromatic ring having from one, two, three or four heteroatoms independently selected from the group consisting of N, O and S, that monocyclic heteroaromatic ring may be optionally substituted with one or more substituents each independently selected from the group consisting of C₁₋₆alkyl, amino, aminoC₁₋₁₂alkyl, halogen, hydroxy, hydroxyC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, C₁₋₁₂alkoxy, aryloxy, aryl C₁₋₁₂alkoxy, -CN, -CF₃, -OCF₃, -COR_i, -COOR_i, -CONR_iR_j, -NR_iCOR_j, -NR_iSO₂R_j, -SO₂R_i, -SO₃R_i or -SO₂NR_iR_j, wherein each R_i and R_j are independently selected from the group consisting of hydrogen, C₁₋₄ alkyl, C₃₋₆cycloalkyl, aryl, arylC₁₋₆alkyl, heteroaryl or heteroarylC₁₋₆alkyl; or A is a 8- to 10 membered bicyclic heteroaromatic ring from one, two, three or four heteroatoms selected from the group consisting of N, O and S, and may be optionally substituted with one, two or three substituents each independently selected from the group consisting of C₁₋₆alkyl, amino, aminoC₁₋₁₂alkyl, halogen, hydroxy, hydroxyC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, C₁₋₁₂alkoxy, aryloxy, aryl C₁₋₁₂alkoxy, -CN, -CF₃, -OCF₃, -COR_k, -COOR_k, -CONR_kR_l, -NR_kCOR_l, -NR_kSO₂R_l, -SO₂R_k, -SO₃R_k or -SO₂NR_kR_l, wherein each R_k and R_l are independently selected from the group consisting of hydrogen, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, aryl, arylC₁₋₆alkyl, heteroaryl or heteroarylC₁₋₆alkyl, and R² is C₁₋₈alkyl substituted with substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl.

[0088] In another embodiment, the TOR kinase inhibitors of formula (I) do not include compounds wherein X and Y are both N and Z is CH, -A-B-Q- is -C(O)NH-, L is a

direct bond, R¹ is substituted or unsubstituted phenyl or substituted or unsubstituted heteroaryl, and R² is substituted or unsubstituted methyl, unsubstituted ethyl, unsubstituted propyl, or an acetamide.

[0089] In another embodiment, the TOR kinase inhibitors of formula (I) do not include compounds wherein X and Y are both N and Z is CH, -A-B-Q- is -C(O)NH-, L is a direct bond, R¹ is substituted or unsubstituted phenyl or substituted or unsubstituted heteroaryl, and R² is an acetamide.

[0090] In another embodiment, the TOR kinase inhibitors of formula (I) do not include compounds wherein X is N and Y and Z are both CH, -A-B-Q- is -C(O)NH-, L is a direct bond, R¹ is a (2,5'-Bi-1H-benzimidazole)-5-carboxamide, and R² is H.

[0091] In another embodiment, the TOR kinase inhibitors of formula (I) do not include compounds wherein one of X and Z is CH and the other is N, Y is CH, -A-B-Q- is -C(O)NH-, L is a direct bond, R¹ is unsubstituted pyridine, and R² is H, methyl or substituted ethyl.

[0092] In another embodiment, the TOR kinase inhibitors of formula (I) do not include compounds wherein X and Z are both N and Y is CH, -A-B-Q- is -C(O)NH-, R¹ is H, C₁₋₈alkyl, C₂₋₈alkenyl, aryl or cycloalkyl, and L is NH.


[0093] In another embodiment, the TOR kinase inhibitors of formula (I) do not include compounds wherein X and Z are both N and Y is CH, -A-B-Q- is -C(O)NR³-, R² is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted phenyl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclylalkyl, and L is NH.

[0094] In another embodiment, the TOR kinase inhibitors of formula (I) do not include compounds wherein R¹ is a substituted or unsubstituted oxazolidinone.

[0095] In another embodiment, the TOR kinase inhibitors of formula (I) do not include one or more of the following compounds: 1,7-dihydro-2-phenyl-8H-Purin-8-one, 1,2-dihydro-3-phenyl-6H-Imidazo[4,5-c]-1,2,4-triazin-6-one, 1,3-dihydro-6-(4-pyridinyl)-2H-Imidazo[4,5-b]pyridin-2-one, 6-(1,3-benzodioxol-5-yl)-1,3-dihydro-1-[(1S)-1-phenylethyl]-2H-Imidazo[4,5-b]pyrazin-2-one, 3-[2,3-dihydro-2-oxo-3-(4-

pyridinylmethyl)-1H-imidazo[4,5-b]pyrazin-5-yl]-Benzamide, 1-[2-(dimethylamino)ethyl]-1,3-dihydro-6-(3,4,5-trimethoxyphenyl)-2H-Imidazo[4,5-b]pyrazin-2-one, N-[5-(1,1-dimethylethyl)-2-methoxyphenyl]-N'-[4-(1,2,3,4-tetrahydro-2-oxopyrido[2,3-b]pyrazin-7-yl)-1-naphthalenyl]-Urea, N-[4-(2,3-dihydro-2-oxo-1H-imidazo[4,5-b]pyridin-6-yl)-1-naphthalenyl]-N'-[5-(1,1-dimethylethyl)-2-methoxyphenyl]-Urea, 1,3-dihydro-5-phenyl-2H-Imidazo[4,5-b]pyrazin-2-one, 1,3-dihydro-5-phenoxy-2H-Imidazo[4,5-b]pyridin-2-one, 1,3-dihydro-1-methyl-6-phenyl-2H-Imidazo[4,5-b]pyridin-2-one, 1,3-dihydro-5-(1H-imidazol-1-yl) 2H-Imidazo[4,5-b]pyridin-2-one, 6-(2,3-dihydro-2-oxo-1H-imidazo[4,5-b]pyridin-6-yl)-8-methyl-2(1H)-Quinolinone and 7,8-dihydro-8-oxo-2-phenyl-9H-purine-9-acetic acid.

[0096] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (Ia):



(Ia)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

L is a direct bond, NH or O;

Y is N or CR³;

R¹ is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted C₂₋₈alkenyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl;

R² is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl;

R^3 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclalkyl, $-NHR^4$ or $-N(R^4)_2$; and

R^4 is at each occurrence independently substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[0097] In one embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R^1 is substituted aryl, such as substituted phenyl.

[0098] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R^1 is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl or substituted or unsubstituted naphthyl.

[0099] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R^1 is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted quinoline, substituted or unsubstituted pyridine, substituted or unsubstituted pyrimidine, substituted or unsubstituted indole, or substituted or unsubstituted thiophene.

[0100] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R^1 is H.

[0101] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R^2 is substituted C_{1-8} alkyl.

[0102] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R^2 is methyl or ethyl substituted with substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[0103] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R^2 is substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl.

[0104] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R^2 is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00105] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R² is H.

[00106] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein Y is CH.

[00107] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein L is a direct bond.


[00108] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R¹ is substituted or unsubstituted aryl and R² is unsubstituted C₁₋₈alkyl.

[00109] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R¹ is substituted or unsubstituted aryl and R² is C₁₋₈alkyl substituted with one or more substituents selected from alkoxy, amino, hydroxy, cycloalkyl, or heterocyclylalkyl.

[00110] In another embodiment, the TOR kinase inhibitors of formula (Ia) are those wherein R¹ is substituted or unsubstituted aryl and R² is substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclylalkyl.

[00111] In another embodiment, the TOR kinase inhibitors of formula (Ia) do not include compounds wherein Y is CH, L is a direct bond, R¹ is substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl, and R² is C₁₋₈alkyl substituted with substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl.

[00112] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (Ib):



(Ib)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

L is a direct bond, NH or O;

R^1 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted C_{2-8} alkenyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl; and

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00113] In one embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R^1 is substituted aryl, such as substituted phenyl.

[00114] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R^1 is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl or substituted or unsubstituted naphthyl.

[00115] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R^1 is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted quinoline, substituted or unsubstituted pyridine, substituted or unsubstituted pyrimidine, substituted or unsubstituted indole, or substituted or unsubstituted thiophene.

[00116] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R^1 is H.

[00117] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R^2 is substituted C_{1-8} alkyl.

[00118] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R^2 is methyl or ethyl substituted with substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00119] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R^2 is substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl.

[00120] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R^2 is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00121] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R² is H.


[00122] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein L is a direct bond.

[00123] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R¹ is substituted or unsubstituted aryl and R² is unsubstituted C₁₋₈alkyl.

[00124] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R¹ is substituted or unsubstituted aryl and R² is C₁₋₈alkyl substituted with one or more substituents selected from alkoxy, amino, hydroxy, cycloalkyl, or heterocyclylalkyl.

[00125] In another embodiment, the TOR kinase inhibitors of formula (Ib) are those wherein R¹ is substituted or unsubstituted aryl and R² is substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclylalkyl.

[00126] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (Ic):



(Ic)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

L is a direct bond, NH or O;

R¹ is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted C₂₋₈alkenyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclylalkyl; and

R² is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclylalkyl.

[00127] In one embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R¹ is substituted aryl, such as substituted phenyl.

[00128] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R¹ is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl or substituted or unsubstituted naphthyl.

[00129] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted quinoline, substituted or unsubstituted pyridine, substituted or unsubstituted pyrimidine, substituted or unsubstituted indole, or substituted or unsubstituted thiophene.

[00130] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R¹ is H.

[00131] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R² is substituted C₁₋₈alkyl.

[00132] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R² is methyl or ethyl substituted with substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00133] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R² is substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl.

[00134] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00135] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R² is H.


[00136] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein L is a direct bond.

[00137] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R¹ is substituted or unsubstituted aryl and R² is unsubstituted C₁₋₈alkyl.

[00138] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R¹ is substituted or unsubstituted aryl and R² is C₁₋₈alkyl substituted with one or more substituents selected from alkoxy, amino, hydroxy, cycloalkyl, or heterocyclylalkyl.

[00139] In another embodiment, the TOR kinase inhibitors of formula (Ic) are those wherein R¹ is substituted or unsubstituted aryl and R² is substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclylalkyl.

[00140] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (Id):



(Id)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

L is a direct bond, NH or O;

R¹ is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted C₂₋₈alkenyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclylalkyl; and

R² is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclylalkyl.

[00141] In one embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R¹ is substituted aryl, such as substituted phenyl.

[00142] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R¹ is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl or substituted or unsubstituted naphthyl.

[00143] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted quinoline, substituted or unsubstituted pyridine, substituted or unsubstituted pyrimidine, substituted or unsubstituted indole, or substituted or unsubstituted thiophene.

[00144] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R¹ is H.

[00145] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R² is substituted C₁₋₈alkyl.

[00146] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R² is methyl or ethyl substituted with substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00147] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R² is substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl.

[00148] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00149] In another embodiment, the Heteroaryl Compounds of formula (Id) are those wherein R² is H.


[00150] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein L is a direct bond.

[00151] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R¹ is substituted or unsubstituted aryl and R² is unsubstituted C₁₋₈alkyl.

[00152] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R¹ is substituted or unsubstituted aryl and R² is C₁₋₈alkyl substituted with one or more substituents selected from alkoxy, amino, hydroxy, cycloalkyl, or heterocyclalkyl.

[00153] In another embodiment, the TOR kinase inhibitors of formula (Id) are those wherein R¹ is substituted or unsubstituted aryl and R² is substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclylalkyl.

[00154] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (Ie):



(Ie)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

L is a direct bond, NH or O;

R¹ is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted C₂₋₈alkenyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclylalkyl; and

R² is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclylalkyl.

[00155] In one embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R¹ is substituted aryl, such as substituted phenyl.

[00156] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R¹ is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl or substituted or unsubstituted naphthyl.

[00157] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted

quinoline, substituted or unsubstituted pyridine, substituted or unsubstituted pyrimidine, substituted or unsubstituted indole, or substituted or unsubstituted thiophene.

[00158] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R¹ is H.

[00159] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R² is substituted C₁₋₈alkyl.

[00160] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R² is methyl or ethyl substituted with substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00161] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R² is substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl.

[00162] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00163] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R² is H.


[00164] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein L is a direct bond.

[00165] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R¹ is substituted or unsubstituted aryl and R² is unsubstituted C₁₋₈alkyl.

[00166] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R¹ is substituted or unsubstituted aryl and R² is C₁₋₈alkyl substituted with one or more substituents selected from alkoxy, amino, hydroxy, cycloalkyl, or heterocyclalkyl.

[00167] In another embodiment, the TOR kinase inhibitors of formula (Ie) are those wherein R¹ is substituted or unsubstituted aryl and R² is substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00168] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (If):



(If)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

L is a direct bond, NH or O;

R¹ is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted C₂₋₈alkenyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl; and

R² is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00169] In one embodiment, the TOR kinase inhibitors of formula (If) are those wherein R¹ is substituted aryl, such as substituted phenyl.

[00170] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R¹ is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl or substituted or unsubstituted naphthyl.

[00171] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted quinoline, substituted or unsubstituted pyridine, substituted or unsubstituted pyrimidine, substituted or unsubstituted indole, or substituted or unsubstituted thiophene.

[00172] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R¹ is H.

[00173] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R² is substituted C₁₋₈alkyl.

[00174] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R² is methyl or ethyl substituted with substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00175] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R² is substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl.

[00176] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00177] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R² is H.


[00178] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein L is a direct bond.

[00179] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R¹ is substituted or unsubstituted aryl and R² is unsubstituted C₁₋₈alkyl.

[00180] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R¹ is substituted or unsubstituted aryl and R² is C₁₋₈alkyl substituted with one or more substituents selected from alkoxy, amino, hydroxy, cycloalkyl, or heterocyclalkyl.

[00181] In another embodiment, the TOR kinase inhibitors of formula (If) are those wherein R¹ is substituted or unsubstituted aryl and R² is substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00182] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (Ig):



(Ig)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

L is a direct bond, NH or O;

R¹ is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted C₂₋₈alkenyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl; and

R² is H, substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00183] In one embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R¹ is substituted aryl, such as substituted phenyl.

[00184] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R¹ is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl or substituted or unsubstituted naphthyl.

[00185] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted quinoline, substituted or unsubstituted pyridine, substituted or unsubstituted pyrimidine, substituted or unsubstituted indole, or substituted or unsubstituted thiophene.

[00186] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R¹ is H.

[00187] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R² is substituted C₁₋₈alkyl.

[00188] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R² is methyl or ethyl substituted with substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00189] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R² is substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclalkyl.

[00190] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00191] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R² is H.

[00192] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein L is a direct bond.

[00193] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R¹ is substituted or unsubstituted aryl and R² is unsubstituted C₁₋₈alkyl.

[00194] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R¹ is substituted or unsubstituted aryl and R² is C₁₋₈alkyl substituted with one or more substituents selected from alkoxy, amino, hydroxy, cycloalkyl, or heterocyclalkyl.

[00195] In another embodiment, the TOR kinase inhibitors of formula (Ig) are those wherein R¹ is substituted or unsubstituted aryl and R² is substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl.

[00196] Representative TOR kinase inhibitors of formula (I) include:

(S)-1-(1-hydroxy-3-methylbutan-2-yl)-6-phenyl-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
1-((tetrahydro-2H-pyran-4-yl)methyl)-6-(3,4,5-trimethoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

(R)-6-(naphthalen-1-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(3-methoxybenzyl)-6-(4-(methylsulfonyl)phenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-1-(1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(4-hydroxyphenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-6-(naphthalen-1-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-1-(1-hydroxy-3-methylbutan-2-yl)-6-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (R)-1-(1-hydroxy-3-methylbutan-2-yl)-6-phenyl-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (R)-1-(1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-1-(1-hydroxy-3-methylbutan-2-yl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (R)-1-(1-hydroxy-3-methylbutan-2-yl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (R)-1-(1-hydroxy-3-methylbutan-2-yl)-6-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-benzyl-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(4-methoxybenzyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (R)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-isopropyl-6-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-cyclohexyl-6-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 5-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-isobutyl-6-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(2-hydroxyethyl)-6-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(5-isopropyl-2-methoxyphenyl)-1-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (R)-1-(1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-c]pyridin-2(3H)-one;

(S)-1-(1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-c]pyridin-2(3H)-one;
 3-(1-phenylethyl)-5-(quinolin-5-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;
 (R)-3-(1-phenylethyl)-5-(quinolin-5-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;
 (R)-6-(5-isopropyl-2-methoxyphenyl)-1-(3-methylbutan-2-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-6-(5-isopropyl-2-methoxyphenyl)-1-(tetrahydrofuran-3-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-6-(5-isopropyl-2-methoxyphenyl)-1-(3-methylbutan-2-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-cyclopentyl-6-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (R)-6-(5-isopropyl-2-methoxyphenyl)-1-(tetrahydrofuran-3-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(cyclopropylmethyl)-6-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(cyclopentylmethyl)-6-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(cyclohexylmethyl)-6-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(5-isopropyl-2-methoxyphenyl)-1-neopentyl-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-isopropyl-6-(3-isopropylphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-isopropyl-6-(2-methoxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-3-(1-hydroxy-3-methylbutan-2-yl)-5-(5-isopropyl-2-methoxyphenyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;
 (R)-1-(2-hydroxy-1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-1-(2-hydroxy-1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-benzhydryl-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-1-(1-phenylpropyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

(R)-1-(1-phenylpropyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(5-isopropyl-2-methoxyphenyl)-1-(tetrahydro-2H-pyran-3-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(3-methoxybenzyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (R)-1-methyl-3-(1-phenylethyl)-5-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (S)-1-methyl-3-(1-phenylethyl)-5-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(cyclopentylmethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-(2-fluorophenyl)ethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-(4-fluorophenyl)ethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-cyclopentyl-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-(3-fluorophenyl)ethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-(3-methoxyphenyl)ethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-(4-methoxyphenyl)ethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(quinolin-5-yl)-1-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(quinolin-5-yl)-1-(tetrahydro-2H-pyran-3-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-((1s,4s)-4-hydroxycyclohexyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-((1r,4r)-4-hydroxycyclohexyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(isoquinolin-5-yl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 (R)-1-(1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;
 1-(1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;
 1-isopropyl-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-(4-chlorophenyl)ethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-(4-(methylsulfonyl)phenyl)ethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-(pyridin-4-yl)ethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 5-methyl-1-((S)-1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 5-methyl-1-((R)-1-phenylethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-phenylethyl)-6-(quinolin-4-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(3-fluorophenyl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(2-fluorophenyl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-phenylethyl)-6-(quinolin-6-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(piperidin-4-ylmethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-(pyridin-2-yl)ethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-(pyridin-3-yl)ethyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-((1s,4s)-4-(hydroxymethyl)cyclohexyl)-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 N-(4-(2-oxo-3-(1-phenylethyl)-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)phenyl)methanesulfonamide;
 6-(3-(methylsulfonyl)phenyl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(3-aminophenyl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(3-(dimethylamino)phenyl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-phenyl-6-(quinolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(1-phenylethyl)-6-(4-(trifluoromethyl)phenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 N-(3-(2-oxo-3-(1-phenylethyl)-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)phenyl)methanesulfonamide;
 6-(4-(methylsulfonyl)phenyl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 3-(1-phenylethyl)-5-(quinolin-5-yl)oxazolo[5,4-b]pyrazin-2(3H)-one;
 1-(cyclopentylmethyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one
 6-(4-hydroxyphenyl)-1-isopropyl-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(4-hydroxyphenyl)-1-isobutyl-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(4-hydroxyphenyl)-1-((tetrahydro-2H-pyran-3-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(cyclohexylmethyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 5-(3-Hydroxyphenyl)-3-(2-methoxyphenyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;
 4-(3-(3-Methoxybenzyl)-2-oxo-2,3-dihydrooxazolo[5,4-b]pyrazin-5-yl)-N-methylbenzamide;

1-Cyclopentyl-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-Cyclohexyl-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 4-(3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)benzamide;
 Methyl 4-(3-(cyclohexylmethyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)benzoate;
 1-(Cyclohexylmethyl)-6-(pyridin-4-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 4-(3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)-N-methylbenzamide;
 1-(Cyclohexylmethyl)-6-(4-(hydroxymethyl)phenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(Cyclohexylmethyl)-6-(pyridin-3-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)benzonitrile;
 1-(Cyclohexylmethyl)-6-(1H-indol-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 4-(3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)-N-isopropylbenzamide;
 1-(2-Hydroxyethyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(Cyclohexylmethyl)-6-(1H-indol-6-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 3-(3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)benzamide;
 6-(4-(Aminomethyl)phenyl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 6-(4-Hydroxyphenyl)-1-((1-methylpiperidin-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;;
 4-(3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)benzonitrile;
 1-((1s,4s)-4-Hydroxycyclohexyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 1-(Cyclohexylmethyl)-6-(pyridin-2-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;
 4-(3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)-N-ethylbenzamide;
 1-(Cyclohexylmethyl)-6-(4-(2-hydroxypropan-2-yl)phenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(4-hydroxy-2-methylphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

4-(3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-5-yl)benzoic acid;

6-(4-Hydroxyphenyl)-1-(2-methoxyethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-(3-methoxypropyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(4-Hydroxyphenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-phenethyl-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-((1r,4r)-4-Hydroxycyclohexyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-1,2,4-Triazol-3-yl)phenyl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-phenyl-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(1H-pyrazol-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(1H-pyrazol-4-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(1-oxoisooindolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(3-(1H-Tetrazol-5-yl)phenyl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(2-oxoindolin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(1H-indazol-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(6-methoxypyridin-3-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-(piperidin-4-ylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(((1r,4r)-4-Aminocyclohexyl)methyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(6-hydroxypyridin-3-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(2-methoxypyridin-4-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

4-((3-((1*r*,4*r*)-4-Hydroxycyclohexyl)-2-oxo-2,3-dihydro-1*H*-imidazo[4,5-*b*]pyrazin-5-yl)benzamide;

2-(4-(3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1*H*-imidazo[4,5-*b*]pyrazin-5-yl)phenyl)acetic acid;

2-(4-(3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1*H*-imidazo[4,5-*b*]pyrazin-5-yl)phenyl)acetamide;

1-(Cyclohexylmethyl)-6-(2-oxoindolin-6-yl)-1*H*-imidazo[4,5-*b*]pyrazin-2(3*H*)-one;

4-(3-(Cyclohexylmethyl)-2-oxo-2,3-dihydro-1*H*-imidazo[4,5-*b*]pyrazin-5-yl)-3-methylbenzoic acid;

N-Methyl-4-(2-oxo-3-((tetrahydro-2*H*-pyran-4-yl)methyl)-2,3-dihydro-1*H*-imidazo[4,5-*b*]pyrazin-5-yl)benzamide;

4-(2-oxo-3-((Tetrahydro-2*H*-pyran-4-yl)methyl)-2,3-dihydro-1*H*-imidazo[4,5-*b*]pyrazin-5-yl)benzamide;

7-(4-Hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-*b*]pyrazin-2(1*H*)-one;

6-(4-(2-Hydroxypropan-2-yl)phenyl)-1-((tetrahydro-2*H*-pyran-4-yl)methyl)-1*H*-imidazo[4,5-*b*]pyrazin-2(3*H*)-one;

6-(1*H*-Indol-5-yl)-1-((tetrahydro-2*H*-pyran-4-yl)methyl)-1*H*-imidazo[4,5-*b*]pyrazin-2(3*H*)-one;

6-(4-(4*H*-1,2,4-Triazol-3-yl)phenyl)-1-((tetrahydro-2*H*-pyran-4-yl)methyl)-1*H*-imidazo[4,5-*b*]pyrazin-2(3*H*)-one;

6-(1*H*-Benzo[*d*]imidazol-5-yl)-1-(cyclohexylmethyl)-1*H*-imidazo[4,5-*b*]pyrazin-2(3*H*)-one;

4-(2-oxo-3-(2-(Tetrahydro-2*H*-pyran-4-yl)ethyl)-2,3-dihydro-1*H*-imidazo[4,5-*b*]pyrazin-5-yl)benzamide;

6-(3-(2*H*-1,2,3-Triazol-4-yl)phenyl)-1-(cyclohexylmethyl)-1*H*-imidazo[4,5-*b*]pyrazin-2(3*H*)-one;

6-(4-(1*H*-Imidazol-1-yl)phenyl)-1-(cyclohexylmethyl)-1*H*-imidazo[4,5-*b*]pyrazin-2(3*H*)-one;

6-(4-(1H-1,2,4-Triazol-3-yl)phenyl)-1-((1r,4r)-4-hydroxycyclohexyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(2H-tetrazol-5-yl)phenyl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(2-hydroxypyridin-4-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-1,2,4-Triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-Imidazol-2-yl)phenyl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-1,2,3-Triazol-1-yl)phenyl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(2-Hydroxypropan-2-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(4-(5-methyl-1H-1,2,4-triazol-3-yl)phenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-Pyrazol-3-yl)phenyl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-Pyrazol-4-yl)phenyl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-(Aminomethyl)-1H-1,2,4-triazol-3-yl)phenyl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one hydrochloride;

1-(Cyclohexylmethyl)-6-(4-(5-(trifluoromethyl)-1H-1,2,4-triazol-3-yl)phenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-((1r,4r)-4-methoxycyclohexyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-((tetrahydrofuran-2-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(3-(1H-1,2,4-Triazol-3-yl)phenyl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-((1r,4r)-4-(Hydroxymethyl)cyclohexyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-((1s,4s)-4-methoxycyclohexyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-((1r,4r)-4-(methoxymethyl)cyclohexyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(1-Methyl-1H-pyrazol-4-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(((1r,4r)-4-Hydroxycyclohexyl)methyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-((tetrahydrofuran-3-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(((1s,4s)-4-Hydroxycyclohexyl)methyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(1H-Benzo[d]imidazol-5-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one hydrochloride;

6-(4-(5-(Morpholinomethyl)-1H-1,2,4-triazol-3-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-(3-(2-oxopyrrolidin-1-yl)propyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-(2-morpholinoethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one hydrochloride;

1-(Cyclohexylmethyl)-6-(4-(oxazol-5-yl)phenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(2-Methyl-1H-benzo[d]imidazol-5-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one hydrochloride;

6-(4-(5-(Methoxymethyl)-1H-1,2,4-triazol-3-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-((1s,4s)-4-(Hydroxymethyl)cyclohexyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(3-Methyl-1H-pyrazol-4-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(1H-Pyrazol-4-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(2-Amino-1H-benzo[d]imidazol-5-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one di hydrochloride;

6-(4-(5-(2-Hydroxypropan-2-yl)-1H-1,2,4-triazol-3-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-Isopropyl-1H-1,2,4-triazol-3-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

4-(2-Methoxy-1-(2-morpholinoethyl)-1H-imidazo[4,5-b]pyrazin-6-yl)benzamide hydrochloride;

4-(1-((1s,4s)-4-Hydroxycyclohexyl)-2-methoxy-1H-imidazo[4,5-b]pyrazin-6-yl)benzamide;

6-(4-Hydroxyphenyl)-1-((1s,4s)-4-(methoxymethyl)cyclohexyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(3H-imidazo[4,5-b]pyridin-6-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(2-(2,2-Dimethyltetrahydro-2H-pyran-4-yl)ethyl)-6-(4-hydroxyphenyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-Pyrazol-1-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(4H-1,2,4-Triazol-3-yl)phenyl)-1-(2-morpholinoethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-Benzo[d]imidazol-2-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-Imidazol-2-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one hydrochloride;

6-(4-(5-(Hydroxymethyl)-1H-1,2,4-triazol-3-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-Imidazol-5-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one hydrochloride;

6-(4-Hydroxyphenyl)-1-((5-oxopyrrolidin-2-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(4,5-Dimethyl-1H-imidazol-2-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-1,2,4-Triazol-5-yl)phenyl)-1-(((1s,4s)-4-methoxycyclohexyl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-1,2,4-Triazol-5-yl)phenyl)-1-(((1r,4r)-4-methoxycyclohexyl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(6-(1H-1,2,4-Triazol-3-yl)pyridin-3-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-1,2,4-Triazol-3-yl)phenyl)-1-(2-(2-oxopyrrolidin-1-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-((dimethylamino)methyl)-1H-1,2,4-triazol-3-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-(pyrrolidin-2-ylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one hydrochloride;

6-(2-Aminobenzimidazol-5-yl)-1-(cyclohexylmethyl)-4-imidazolino[4,5-b]pyrazin-2-one di hydrochloride;

6-(2-(Dimethylamino)-1H-benzo[d]imidazol-5-yl)-1-((tetrahydro-2H-pyran-4-yl) methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-(piperidin-3-ylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-1-(2-(piperidin-1-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one hydrochloride;

1-(Cyclohexylmethyl)-6-(2-(methylamino)pyrimidin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(3-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(2-(2-methoxyethylamino)pyrimidin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-((methylamino)methyl)-1H-1,2,4-triazol-3-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-Oxopyrrolidin-2-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-methyl-1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(1H-imidazol-2-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-1-(2-methyl-2-morpholinopropyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(4H-1,2,4-Triazol-3-yl)phenyl)-1-(1-morpholinopropan-2-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(Pyrrolidin-2-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-(aminomethyl)-1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(5-(Hydroxymethyl)thiophen-2-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

(1r,4r)-4-(6-(4-Hydroxyphenyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-1-yl)cyclohexanecarboxamide;

(1s,4s)-4-(6-(4-Hydroxyphenyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-1-yl)cyclohexanecarboxamide;

6-(4-(5-methyl-1H-1,2,4-triazol-3-yl)phenyl)-1-(2-morpholinoethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-Oxopyrrolidin-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(Pyrrolidin-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(1H-benzo[d]imidazol-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(3-(Hydroxymethyl)thiophen-2-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(5-(2-Hydroxyethyl)thiophen-2-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(pyrimidin-5-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(6-Fluoropyridin-3-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(6-Aminopyridin-3-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-methyl-1H-imidazol-2-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-Methyl-1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(2-oxopyrrolidin-1-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(6-(Methylamino)pyridin-3-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(2-aminopyrimidin-5-yl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(2-hydroxypropan-2-yl)phenyl)-1-(((1r,4r)-4-methoxycyclohexyl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-hydroxyphenyl)-1-((1-methylpiperidin-3-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

1-(cyclohexylmethyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(hydroxymethyl)thiophen-2-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(1H-benzo[d]imidazol-6-yl)-1-(((1r,4r)-4-methoxycyclohexyl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(4,5-dimethyl-1H-imidazol-2-yl)phenyl)-1-(2-morpholinoethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-1-(2-morpholino-2-oxoethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyridin-2(3H)-one;

(R)-6-(4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

(S)-6-(4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

(1r,4r)-4-(6-(4-(2-hydroxypropan-2-yl)phenyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-1-yl)cyclohexanecarboxamide;

6-(3-Methyl-4-(1H-1,2,4-Triazol-3-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-B]pyrazin-2(3H)-one;

6-(4-(1H-imidazol-2-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(5-(Aminomethyl)-1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(1H-benzo[d]imidazol-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(2-Aminopyrimidin-5-yl)-1-(cyclohexylmethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-Hydroxyphenyl)-1-((1-methylpiperidin-2-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one hydrochloride;

6-(3-Methyl-4-(1H-1,2,4-Triazol-3-yl)phenyl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-B]pyrazin-2(3H)-one;

1-(Cyclohexylmethyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(6-(2-Hydroxypropan-2-yl)pyridin-3-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(6-(2-Hydroxypropan-2-yl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

6-(4-(4H-1,2,4-Triazol-3-yl)phenyl)-1-(2-morpholino-2-oxoethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

(R)-6-(4-(4H-1,2,4-Triazol-3-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(R)-6-(4-(1H-1,2,4-Triazol-3-yl)phenyl)-1-(1-phenylethyl)-1H-imidazo[4,5-B]pyrazin-2(3H)-one;


(S)-6-(4-(4H-1,2,4-Triazol-3-yl)phenyl)-1-(1-phenylethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one;

(1r,4r)-4-(6-(4-(2-Hydroxypropan-2-yl)phenyl)-2-oxo-2,3-dihydro-1H-imidazo[4,5-b]pyrazin-1-yl)cyclohexanecarboxamide; and

6-(4-(5-Methyl-1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-1H-imidazo[4,5-b]pyrazin-2(3H)-one,

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof.

[00197] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (II):



(II)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

R¹ is substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl;

-X-A-B-Y- taken together form -N(R²)CH₂C(O)NH-, -N(R²)C(O)CH₂NH-, -N(R²)C(O)NH-, -N(R²)C=N-, or -C(R²)=CHNH-;

L is a direct bond, NH or O;

R² is substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl; and

R³ and R⁴ are independently H or C₁₋₈alkyl.

[00198] In one embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)CH₂C(O)NH-.

[00199] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)CH₂NH-.

[00200] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH-.

[00201] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C=N-.

[00202] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -C(R²)=CHNH-.

[00203] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein L is a direct bond.

[00204] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein R¹ is substituted aryl, such as substituted phenyl.

[00205] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted pyridine, substituted or unsubstituted indole or substituted or unsubstituted quinoline.

[00206] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein R¹ is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclopentyl.

[00207] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH- and R¹ is substituted aryl, such as phenyl.

[00208] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH- and R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted pyridine, substituted or unsubstituted indole or substituted or unsubstituted quinoline.

[00209] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH- and R¹ is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclopentyl.

[00210] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein R² is substituted C₁₋₈alkyl, such as -CH₂C₆H₅.

[00211] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein R² is unsubstituted C₁₋₈alkyl, such as unsubstituted methyl.

[00212] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00213] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein R² is substituted aryl, such as halo, haloalkyl or alkoxy substituted phenyl.

[00214] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein R² is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclohexyl or substituted or unsubstituted cycloheptyl.

[00215] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein R² is substituted heterocyclalkyl, such as substituted piperidine.

[00216] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein R³ and R⁴ are H.

[00217] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH- and R² is unsubstituted aryl, such as unsubstituted phenyl.

[00218] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH-, R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted pyridine, and R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00219] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH-, R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted pyridine, R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl, and R³ and R⁴ are H.

[00220] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH-, L is a direct bond, R¹ is

substituted or unsubstituted heteroaryl, such as substituted or unsubstituted pyridine, R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl, and R³ and R⁴ are H.

[00221] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH-, R¹ is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl, and R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00222] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH-, R¹ is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl, R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl, and R³ and R⁴ are H.

[00223] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH-, L is a direct bond, R¹ is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl, R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl, and R³ and R⁴ are H.

[00224] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH-, R¹ is substituted or unsubstituted heteroaryl, L is a direct bond and R² is substituted or unsubstituted C₁₋₈alkyl or substituted or unsubstituted cycloalkyl.

[00225] In another embodiment, the TOR kinase inhibitors of formula (II) are those wherein -X-A-B-Y- taken together form -N(R²)C(O)NH-, R¹ is substituted or unsubstituted aryl, L is a direct bond and R² is substituted or unsubstituted C₁₋₈alkyl or substituted or unsubstituted cycloalkyl.

[00226] In another embodiment, the TOR kinase inhibitors of formula (II) do not include 8,9-dihydro-8-oxo-9-phenyl-2-(3-pyridinyl)-7H-purine-6-carboxamide, 8,9-dihydro-8-oxo-9-phenyl-2-(3-pyridinyl)-7H-purine-6-carboxamide, 8,9-dihydro-8-oxo-9-phenyl-2-(3-pyridinyl)-7H-purine-6-carboxamide, 2-(4-cyanophenyl)-8-oxo-9-phenyl-8,9-dihydro-


7H-purine-6-carboxamide, 2-(4-nitrophenyl)-8-oxo-9-phenyl-8,9-dihydro-7H-purine-6-carboxamide, 9-benzyl-2-(4-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide, 2-methyl-8-oxo-9-phenyl-8,9-dihydro-7H-purine-6-carboxamide, 9-benzyl-9H-purine-2,6-dicarboxamide, 9-[2,3-bis[(benzoyloxy)methyl]cyclobutyl]-2-methyl-9H-Purine-6-carboxamide, 9-benzyl-2-methyl-9H-purine-6-carboxamide, 9-(2-hydroxyethyl)-2-methyl-9H-purine-6-carboxamide, 9-(2-hydroxyethyl)-2-(trifluoromethyl)-9H-purine-6-carboxamide, 9-(2-hydroxyethyl)-2-(prop-1-enyl)-9H-purine-6-carboxamide, 9-(2-hydroxyethyl)-2-phenyl-9H-purine-6-carboxamide, 9-(3-hydroxypropyl)-2-methyl-9H-purine-6-carboxamide, 9-(3-hydroxypropyl)-2-(trifluoromethyl)-9H-purine-6-carboxamide, 2-methyl-9-phenylmethyl-9H-purine-6-carboxamide or 2-methyl-9- β -D-ribofuranosyl-9H-purine-6-carboxamide.

[00227] In another embodiment, the TOR kinase inhibitors of formula (II) do not include compounds wherein R² is a substituted furanoside.

[00228] In another embodiment, the TOR kinase inhibitors of formula (II) do not include compounds wherein R² is a substituted or unsubstituted furanoside.

[00229] In another embodiment, the TOR kinase inhibitors of formula (II) do not include (2'R)-2'-deoxy-2'-fluoro-2'-C-methyl nucleosides.

[00230] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (IIa):



and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl;

R^2 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl; and

R^3 and R^4 are independently H or C_{1-8} alkyl.

[00231] In one embodiment, the TOR kinase inhibitors of formula (IIa) are those wherein R^1 is substituted aryl, substituted or unsubstituted heteroaryl, such as substituted phenyl.

[00232] In another embodiment, the TOR kinase inhibitors of formula (IIa) are those wherein R^1 is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted pyridine, substituted or unsubstituted indole or substituted or unsubstituted quinoline.

[00233] In another embodiment, the TOR kinase inhibitors of formula (IIa) are those wherein R^1 is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclopentyl.

[00234] In another embodiment, the TOR kinase inhibitors of formula (IIa) are those wherein R^2 is substituted C_{1-8} alkyl, such as $-CH_2C_6H_5$.

[00235] In another embodiment, the TOR kinase inhibitors of formula (IIa) are those wherein R^2 is unsubstituted C_{1-8} alkyl, such as unsubstituted methyl.

[00236] In another embodiment, the TOR kinase inhibitors of formula (IIa) are those wherein R^2 is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00237] In another embodiment, the TOR kinase inhibitors of formula (IIa) are those wherein R^2 is substituted aryl, such as halo, haloalkyl or alkoxy substituted phenyl.

[00238] In another embodiment, the TOR kinase inhibitors of formula (IIa) are those wherein R² is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclohexyl or substituted or unsubstituted cycloheptyl.

[00239] In another embodiment, the TOR kinase inhibitors of formula (IIa) are those wherein R² is substituted heterocyclalkyl, such as substituted piperidine.

[00240] In another embodiment, the TOR kinase inhibitors of formula (IIa) are those wherein R³ and R⁴ are H.


[00241] In another embodiment, the TOR kinase inhibitors of formula (IIa) do not include 8,9-dihydro-8-oxo-9-phenyl-2-(3-pyridinyl)-7H-Purine-6-carboxamide, 8,9-dihydro-8-oxo-9-phenyl-2-(3-pyridinyl)-7H-Purine-6-carboxamide, 8,9-dihydro-8-oxo-9-phenyl-2-(3-pyridinyl)-7H-Purine-6-carboxamide, 2-(4-cyanophenyl)-8-oxo-9-phenyl-8,9-dihydro-7H-purine-6-carboxamide, 2-(4-nitrophenyl)-8-oxo-9-phenyl-8,9-dihydro-7H-purine-6-carboxamide, 9-benzyl-2-(4-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide, 9-phenylmethyl-9H-purine-2,6-dicarboxamide, or 2-methyl-8-oxo-9-phenyl-8,9-dihydro-7H-purine-6-carboxamide.

[00242] In another embodiment, the TOR kinase inhibitors of formula (IIa) do not include compounds wherein R² is a substituted furanoside.

[00243] In another embodiment, the TOR kinase inhibitors of formula (IIa) do not include compounds wherein R² is a substituted or unsubstituted furanoside.

[00244] In another embodiment, the TOR kinase inhibitors of formula (IIa) do not include (2'R)-2'-deoxy-2'-fluoro-2'-C-methyl nucleosides.

[00245] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (IIb):



(IIb)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

$\text{---X} \begin{smallmatrix} \nearrow \\ \searrow \end{smallmatrix} \text{---Y---}$ is $-\text{C}(\text{R}^2)=\text{CH}-\text{NH}-$ or $-\text{N}(\text{R}^2)-\text{CH}=\text{N}-$;

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl;

R^2 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl; and

R^3 and R^4 are independently H or C_{1-8} alkyl.

[00246] In one embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R^1 is substituted aryl, such as substituted phenyl.

[00247] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R^1 is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted pyridine, substituted or unsubstituted indole or substituted or unsubstituted quinoline.

[00248] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R^1 is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclopentyl.

[00249] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R² is substituted C₁₋₈alkyl, such as -CH₂C₆H₅.

[00250] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R² is unsubstituted C₁₋₈alkyl, such as unsubstituted methyl.

[00251] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00252] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R² is substituted aryl, such as halo, haloalkyl or alkoxy substituted phenyl.

[00253] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R² is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclohexyl or substituted or unsubstituted cycloheptyl.

[00254] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R² is substituted heterocyclalkyl, such as substituted piperidine.

[00255] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R³ and R⁴ are H.

[00256] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein $\text{---X} \overbrace{\text{---}}^{\text{dashed line}} \text{---Y---}$ is $-\text{C}(\text{R}^2)=\text{CH}-\text{NH}-$ and R² is substituted aryl, such as substituted phenyl.

[00257] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein $\text{---X} \overbrace{\text{---}}^{\text{dashed line}} \text{---Y---}$ is $-\text{N}(\text{R}^2)-\text{CH}=\text{N}-$ and R² is substituted aryl, such as substituted phenyl.

[00258] In another embodiment, the TOR kinase inhibitors of formula (IIb) are those wherein R¹ is substituted aryl, such as phenyl, and R² is substituted aryl, such as substituted phenyl.

[00259] In another embodiment, the TOR kinase inhibitors of formula (IIb) do not include 9-benzyl-9H-purine-2,6-dicarboxamide, 9-[2,3-bis[(benzoyloxy)methyl]cyclobutyl]-2-methyl-9H-Purine-6-carboxamide, 9-benzyl-2-methyl-9H-purine-6-carboxamide, 9-(2-hydroxyethyl)-2-methyl-9H-purine-6-carboxamide, 9-(2-hydroxyethyl)-2-(trifluoromethyl)-

9H-purine-6-carboxamide, 9-(2-hydroxyethyl)-2-(prop-1-enyl)-9H-purine-6-carboxamide, 9-(2-hydroxyethyl)-2-phenyl-9H-purine-6-carboxamide, 9-(3-hydroxypropyl)-2-methyl-9H-purine-6-carboxamide, 9-(3-hydroxypropyl)-2-(trifluoromethyl)-9H-purine-6-carboxamide, 9-phenylmethyl-9H-purine-2,6-dicarboxamide, 2-methyl-9-phenylmethyl-9H-purine-6-carboxamide or 2-methyl-9- β -D-ribofuranosyl-9H-purine-6-carboxamide.

[00260] In another embodiment, the TOR kinase inhibitors of formula (IIb) do not include compounds wherein R² is substituted cyclobutyl when $\text{---X} \begin{array}{c} \diagup \\ \diagdown \end{array} \text{---Y---}$ is -N(R²)-CH=N-.


[00261] In another embodiment, the TOR kinase inhibitors of formula (IIb) do not include compounds wherein R² is a substituted furanoside when $\text{---X} \begin{array}{c} \diagup \\ \diagdown \end{array} \text{---Y---}$ is -N(R²)-CH=N-.

[00262] In another embodiment, the TOR kinase inhibitors of formula (IIb) do not include compounds wherein R² is substituted pyrimidine when $\text{---X} \begin{array}{c} \diagup \\ \diagdown \end{array} \text{---Y---}$ is -C(R²)=CH-NH-.

[00263] In another embodiment, the TOR kinase inhibitors of formula (IIb) do not include compounds wherein R² is substituted oxetane when $\text{---X} \begin{array}{c} \diagup \\ \diagdown \end{array} \text{---Y---}$ is -N(R²)-CH=N-.

[00264] In another embodiment, the TOR kinase inhibitors of formula (IIb) do not include compounds wherein R² is substituted cyclopentyl or a heterocyclopentyl when $\text{---X} \begin{array}{c} \diagup \\ \diagdown \end{array} \text{---Y---}$ is -N(R²)-CH=N-.

[00265] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (IIc):



and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

R¹ is substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl;

R² is substituted or unsubstituted C₁₋₈alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl; and

R³ and R⁴ are independently H or C₁₋₈alkyl.

[00266] In one embodiment, the TOR kinase inhibitors of formula (IIc) are those wherein R¹ is substituted aryl, such as substituted phenyl.

[00267] In another embodiment, the TOR kinase inhibitors of formula (IIc) are those wherein R¹ is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted pyridine, substituted or unsubstituted indole or substituted or unsubstituted quinoline.

[00268] In another embodiment, the TOR kinase inhibitors of formula (IIc) are those wherein R¹ is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclopentyl.

[00269] In another embodiment, the TOR kinase inhibitors of formula (IIc) are those wherein R² is substituted C₁₋₈alkyl, such as -CH₂C₆H₅.

[00270] In another embodiment, the TOR kinase inhibitors of formula (IIc) are those wherein R² is unsubstituted C₁₋₈alkyl, such as unsubstituted methyl.

[00271] In another embodiment, the TOR kinase inhibitors of formula (IIc) are those wherein R² is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.


[00272] In another embodiment, the TOR kinase inhibitors of formula (IIc) are those wherein R² is substituted aryl, such as halo, haloalkyl or alkoxy substituted phenyl.

[00273] In another embodiment, the TOR kinase inhibitors of formula (IIc) are those wherein R² is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclohexyl or substituted or unsubstituted cycloheptyl.

[00274] In another embodiment, the TOR kinase inhibitors of formula (IIc) are those wherein R² is substituted heterocyclalkyl, such as substituted piperidine.

[00275] In another embodiment, the TOR kinase inhibitors of formula (IIc) are those wherein R³ and R⁴ are H.

[00276] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (IId):



(IId)

and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl;

R^2 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or substituted or unsubstituted heterocyclalkyl; and

R^3 and R^4 are independently H or C_{1-8} alkyl.

[00277] In one embodiment, the TOR kinase inhibitors of formula (IId) are those wherein R^1 is substituted aryl, such as substituted phenyl.

[00278] In another embodiment, the TOR kinase inhibitors of formula (IId) are those wherein R^1 is substituted or unsubstituted heteroaryl, such as substituted or unsubstituted pyridine, substituted or unsubstituted indole or substituted or unsubstituted quinoline.

[00279] In another embodiment, the TOR kinase inhibitors of formula (IId) are those wherein R^1 is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclopentyl.

[00280] In another embodiment, the TOR kinase inhibitors of formula (IId) are those wherein R^2 is substituted C_{1-8} alkyl, such as $-CH_2C_6H_5$.

[00281] In another embodiment, the TOR kinase inhibitors of formula (IId) are those wherein R^2 is unsubstituted C_{1-8} alkyl, such as unsubstituted methyl.

[00282] In another embodiment, the TOR kinase inhibitors of formula (IId) are those wherein R^2 is substituted or unsubstituted aryl, such as substituted or unsubstituted phenyl.

[00283] In another embodiment, the TOR kinase inhibitors of formula (IId) are those wherein R^2 is substituted aryl, such as halo, haloalkyl or alkoxy substituted phenyl.

[00284] In another embodiment, the TOR kinase inhibitors of formula (IId) are those wherein R^2 is substituted or unsubstituted cycloalkyl, such as substituted or unsubstituted cyclohexyl or substituted or unsubstituted cycloheptyl.

[00285] In another embodiment, the TOR kinase inhibitors of formula (IId) are those wherein R^2 is substituted heterocyclalkyl, such as substituted piperidine.

[00286] In another embodiment, the TOR kinase inhibitors of formula (IIId) are those wherein R³ and R⁴ are H.

[00287] Representative TOR kinase inhibitors of formula (IV) include:

9-benzyl-8-oxo-2-(pyridin-3-yl)-8,9-dihydro-7H-purine-6-carboxamide;
 N-methyl-8-oxo-9-phenyl-2-(pyridin-3-yl)-8,9-dihydro-7H-purine-6-carboxamide;
 8-oxo-9-phenyl-2-(pyridin-2-yl)-8,9-dihydro-7H-purine-6-carboxamide;
 2-(2-chloropyridin-3-yl)-8-oxo-9-phenyl-8,9-dihydro-7H-purine-6-carboxamide;
 2-(2-methoxypyridin-3-yl)-8-oxo-9-phenyl-8,9-dihydro-7H-purine-6-carboxamide;
 N,N-dimethyl-8-oxo-9-phenyl-2-(pyridin-3-yl)-8,9-dihydro-7H-purine-6-carboxamide;
 9-methyl-8-oxo-2-(pyridin-3-yl)-8,9-dihydro-7H-purine-6-carboxamide;
 2-(4-hydroxyphenyl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(3-hydroxyphenyl)-8-oxo-9-0-tolyl-8,9-dihydro-7H-purine-6-carboxamide;
 2-(1H-indol-4-yl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(1H-indol-6-yl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(3-hydroxyphenyl)-9-(4-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(2-hydroxypyridin-4-yl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-
 carboxamide;
 9-(2-chlorophenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 9-(2-fluorophenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 9-(2,6-difluorophenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 9-cycloheptyl-8-oxo-2-(pyridin-3-yl)-8,9-dihydro-7H-purine-6-carboxamide;
 9-(2-methoxyphenyl)-8-oxo-2-(quinolin-5-yl)-8,9-dihydro-7H-purine-6-carboxamide;
 2-cyclopentyl-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 9-(2-methoxyphenyl)-8-oxo-2-(3-(trifluoromethyl)phenyl)-8,9-dihydro-7H-purine-6-
 carboxamide;
 9-(2-methoxyphenyl)-2-(6-methoxypyridin-3-yl)-8-oxo-8,9-dihydro-7H-purine-6-
 carboxamide;

2-(3-hydroxyphenyl)-8-oxo-9-(4-(trifluoromethyl)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;

9-benzyl-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-hydroxyphenyl)-8-oxo-9-(2-(trifluoromethoxy)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;

9-(2,4-dichlorophenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-methoxyphenyl)-2-(3-nitrophenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-cyanophenyl)-8-oxo-9-phenyl-8,9-dihydro-7H-purine-6-carboxamide;

9-(3-fluorophenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-methoxyphenyl)-8-oxo-2-(2-(trifluoromethyl)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;

2-(5-fluoropyridin-3-yl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(1-benzylpiperidin-4-yl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

benzyl 4-(6-carbamoyl-8-oxo-2-(pyridin-3-yl)-7H-purin-9(8H)-yl)piperidine-1-carboxylate;

9-cyclohexyl-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-methoxyphenyl)-8-oxo-2-(3-(trifluoromethoxy)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;

9-phenyl-2-(pyridin-3-yl)-9H-purine-6-carboxamide;

6-oxo-8-phenyl-2-(pyridin-3-yl)-5,6,7,8-tetrahydropteridine-4-carboxamide;

6-oxo-8-phenyl-2-(pyridin-4-yl)-5,6,7,8-tetrahydropteridine-4-carboxamide;

2-(3-aminophenyl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-hydroxyphenyl)-9-(2-methoxyphenyl)-9H-purine-6-carboxamide;

9-Cyclopentyl-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-tert-Butyl-2-(3-hydroxy-phenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

[2-(3-Hydroxyphenyl)-9-(2-methoxyphenyl)-8-oxo(7-hydropurin-6-yl)]-N-methylcarboxamide;

2-phenyl-5H-pyrrolo[3,2-d]pyrimidine-4-carboxamide;
 [2-(3-Hydroxyphenyl)-9-(2-methoxyphenyl)-8-oxo(7-hdropurin-6-yl)]-N,N-dimethyl carboxamide;
 2-(3-Hydroxyphenylamino)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(4-Hydroxyphenylamino)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 9-(trans-4-Hydroxycyclohexyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 9-(trans-4-Hydroxycyclohexyl)-8-oxo-2-(pyridin-3-yl)-8,9-dihydro-7H-purine-6-carboxamide;
 9-(trans-4-Hydroxycyclohexyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 9-(trans-4-Hydroxycyclohexyl)-8-oxo-2-(pyridin-3-yl)-8,9-dihydro-7H-purine-6-carboxamide;
 2-(3-Hydroxyphenylamino)-9-(2-methoxyphenyl)-9H-purine-6-carboxamide;
 9-Isopropyl-2-(3-hydroxy-phenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 Methyl 4-(6-carbamoyl-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purin-2-yl) benzoate;
 2-(2-Chloro-3-hydroxyphenyl)-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carbox amide;
 2-(3-Cyanophenyl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(2-Hydroxyphenylamino)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(3-Hydroxyphenyl)-9-(4-methoxy-2-methylphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(3-Hydroxyphenyl)-8-oxo-9-(2-(trifluoromethyl)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;
 2-(4-Cyano-phenyl)-9-(2-methoxy-phenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 4-[6-Carbamoyl-9-(2-methoxy-phenyl)-8-oxo-8,9-dihydro-7H-purin-2-yl]-benzoic acid;

Methyl 3-(6-carbamoyl-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purin-2-yl)benzoate; 3-(6-Carbamoyl-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purin-2-yl)benzoic acid; 2-(3-Hydroxyphenyl)-9-(2-isopropylphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide; 2-(1H-Indazol-6-yl)-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carboxamide; 2-(4-Carbamoylphenyl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide; 9-(2-Ethylphenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide; 9-(2,5-Dichlorophenyl)-2-(3-hydroxyphenyl)-8-oxo-7-hdropurine-6-carboxamide; 2-(3-Carbamoylphenyl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide; 9-(2,6-Dichlorophenyl)-2-(3-hydroxyphenyl)-8-oxo-7-hdropurine-6-carboxamide; 2-(2-Hydroxyphenyl)-9-(2-methoxyphenyl)purine-6-carboxamide; 2-(1H-Indazol-5-yl)-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carboxamide; 9-(2,3-Dichlorophenyl)-2-(3-hydroxyphenyl)-8-oxo-7-hdropurine-6-carboxamide; 2-[4-(Hydroxymethyl)phenyl]-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carbox-amide; 2-[3-(Hydroxymethyl)phenyl]-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carbox-amide; 9-(2-Methoxyphenyl)-8-oxo-2-(pyridin-4-yl)-8,9-dihydro-7H-purine-6-carboxamide; 2-(4-Fluoro-3-hydroxyphenyl)-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carbox-amide; 2-(2-Fluoro-3-hydroxyphenyl)-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carbox-amide; 2-[4-(1-Hydroxy-isopropyl)phenyl]-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carboxamide; 2-[3-(1-Hydroxy-isopropyl)phenyl]-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carboxamide; 9-(2-Methoxyphenyl)-2-(2-nitrophenyl)-8-oxo-7-hdropurine-6-carboxamide; 9-(2-Methoxyphenyl)-2-(4-nitrophenyl)-8-oxo-7-hdropurine-6-carboxamide; 9-(2-Methoxyphenyl)-2-(2-nitrophenyl)-8-oxo-7-hdropurine-6-carboxamide; 9-(2,4-Difluorophenyl)-2-(3-hydroxyphenyl)-8-oxo-7-hdropurine-6-carboxamide;

9-(2-Methoxyphenyl)-2-{3-[(methylsulfonyl)amino]phenyl}-8-oxo-7-hdropurine-6-carboxamide;

9-(4-Chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-8-oxo-7-hdropurine-6-carboxamide;

9-(2-Chlorophenyl)-8-oxo-2-(3-pyridyl)-7-hdropurine-6-carboxamide;

8-Oxo-2-(3-pyridyl)-9-[2-(trifluoromethyl)phenyl]-7-hdropurine-6-carboxamide;

9-(3-Chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-8-oxo-7-hdropurine-6-carboxamide;

9-(2-Fluoro-3-trifluoromethylphenyl)-2-(3-hydroxyphenyl)-8-oxo-7-hdropurine-6-carboxamide;

9-(2, 3, 4-Trifluorophenyl)-2-(3-hydroxyphenyl)-8-oxo-7-hdropurine-6-carboxamide;

2-(1H-Benzo[d]imidazol-6-yl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-[3-(Acetylamino)phenyl]-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carboxamide;

2-(3-hydroxyphenyl)-8-(2-methoxyphenyl)-6-oxo-5,6,7,8-tetrahydropteridine-4-carboxamide;

9-(2-Methoxyphenyl)-8-oxo-2-pyrazol-4-yl-7-hdropurine-6-carboxamide;

9-(2-Methoxyphenyl)-8-oxo-2-pyrazol-3-yl-7-hdropurine-6-carboxamide;

9-(4-Aminocyclohexyl)-2-(3-hydroxyphenyl)-8-oxo-7-hdropurine-6-carboxamide;

2-[3-(Difluoromethyl)phenyl]-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carboxamide;

2-[5-(Difluoromethyl)-2-fluorophenyl]-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carboxamide;

2-(1H-benzo[d]imidazol-4-yl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(6-Hydroxypyridin-3-yl)-8-oxo-9-(2-(trifluoromethyl)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;

2-(1H-benzo[d]imidazol-6-yl)-9-(2-fluorophenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-Benzimidazol-6-yl-8-oxo-9-[2-(trifluoromethyl)phenyl]-7-hdropurine-6-carboxamide;

2-(5-Chloropyridin-3-yl)-8-oxo-9-(2-(trifluoromethyl)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;

trans-4-(6-Carbamoyl-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purin-2-ylamino)cyclohexyl carbamate;

(R)-9-(2-Methoxyphenyl)-8-oxo-2-(pyrrolidin-3-ylamino)-8,9-dihydro-7H-purine-6-carboxamide;

(S)-9-(2-Methoxyphenyl)-8-oxo-2-(pyrrolidin-3-ylamino)-8,9-dihydro-7H-purine-6-carboxamide;

(cis)-4-(6-Carbamoyl-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purin-2-ylamino)cyclohexyl carbamate;

2-(trans-4-Hydroxycyclohexylamino)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(4-Chloropyridin-3-yl)-8-oxo-9-(2-(trifluoromethyl)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;

2-(cis-4-Hydroxycyclohexylamino)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(4-((1H-Imidazol-1-yl)methyl)phenylamino)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(4-Hydroxypyridin-3-yl)-8-oxo-9-(2-(trifluoromethyl)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;

(R)-9-(2-Methoxyphenyl)-8-oxo-2-(pyrrolidin-2-ylmethylamino)-8,9-dihydro-7H-purine-6-carboxamide;

(S)-9-(2-Methoxyphenyl)-8-oxo-2-(pyrrolidin-2-ylmethylamino)-8,9-dihydro-7H-purine-6-carboxamide;

2-(4-(1H-1,2,4-Triazol-3-yl)phenyl)-9-(2-methoxyphenyl)-8-oxo-7-hydropurine-6-carboxamide;

2-(2-Hydroxyethylamino)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide ;

9-(2-Methoxyphenyl)-8-oxo-2-(2-(trifluoromethyl)-1H-benzo[d]imidazol-6-yl)-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-(1H-1,2,4-Triazol-3-yl)phenyl)-9-(2-methoxyphenyl)-8-oxo-7-hdropurine-6-carboxamide;

9-(Biphenyl-2-yl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(4-(1H-1,2,4-Triazol-3-yl)phenyl)-9-(2-fluorophenyl)-8-oxo-7-hdropurine-6-carboxamide;

2-(4-(1H-1,2,4-Triazol-3-yl)phenyl)-9-(2-isopropylphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-Methoxyphenyl)-2-(2-methyl-1H-benzo[d]imidazol-6-yl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-(Hydroxymethyl)phenylamino)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(2-(Hydroxymethyl)phenylamino)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-tert-Butylphenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-Hydroxyphenyl)-8-oxo-9-(2-phenoxyphenyl)-8,9-dihydro-7H-purine-6-carboxamide;

2-(1H-Benzo[d]imidazol-6-yl)-9-(2-isopropylphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(1H-Indazol-4-yl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(2-Hydroxypyridin-3-yl)-8-oxo-9-(2-(trifluoromethyl)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;

2-(1H-Imidazo[4,5-b]pyridin-6-yl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(4-(1H-Imidazol-1-yl)phenyl)-9-(2-isopropylphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-Cyclohexylphenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(4-(1H-Imidazol-2-yl)phenyl)-9-(2-isopropylphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(1H-Benzo[d]imidazol-1-yl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(1H-Imidazo[4,5-b]pyridin-6-yl)-9-(2-isopropylphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-Isopropylphenyl)-8-oxo-2-(1H-pyrrolo[2,3-b]pyridin-5-yl)-8,9-dihydro-7H-purine-6-carboxamide;

2-(1H-Imidazo[4,5-b]pyridin-6-yl)-8-oxo-9-(2-(trifluoromethyl)phenyl)-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-Methoxyphenyl)-2-(2-(methylthio)-1H-benzo[d]imidazol-5-yl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(1H-Indol-5-yl)-9-(2-isopropylphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(Cyclohexylmethyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(2,3-Dihydro-1H-inden-1-yl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-Hydroxyphenyl)-9-isobutyl-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(trans-4-Methoxycyclohexyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(cis-4-Methoxycyclohexyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-Hydroxyphenyl)-8-oxo-9-(5,6,7,8-tetrahydronaphthalen-1-yl)-8,9-dihydro-7H-purine-6-carboxamide;

2-(4-(1H-1,2,4-Triazol-3-yl)phenyl)-9-cyclohexyl-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-Hydroxyphenyl)-9-(1H-indol-4-yl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-Fluoro-3-methoxyphenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-Fluoro-5-methoxyphenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-Cyclohexyl-2-(1H-imidazo[4,5-b]pyridin-6-yl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-Hydroxyphenyl)-8-oxo-9-(tetrahydro-2H-pyran-4-yl)-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-Hydroxyphenyl)-8-oxo-9-((tetrahydro-2H-pyran-4-yl)methyl)-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-Cyclopentylphenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-Hydroxyphenyl)-8-oxo-9-(piperidin-4-yl)-8,9-dihydro-7H-purine-6-carboxamide;

9-(2-Fluoro-4-methoxyphenyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(1H-benzo[d]imidazol-6-yl)-9-cyclohexyl-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-Benzimidazol-6-yl-9-(trans-4-methoxycyclohexyl)-8-oxo-7-hydropurine-6-carboxamide;

2-(4-(Aminomethyl)phenyl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-Hydroxyphenyl)-9-(cis-4-(methoxymethyl)cyclohexyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

9-(trans-4-Aminocyclohexyl)-2-(3-hydroxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-Hydroxyphenyl)-9-(2-isobutylphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;


(R)-2-(3-Hydroxyphenyl)-8-oxo-9-(tetrahydrofuran-3-yl)-8,9-dihydro-7H-purine-6-carboxamide;

(S)-2-(3-Hydroxyphenyl)-8-oxo-9-(tetrahydrofuran-3-yl)-8,9-dihydro-7H-purine-6-carboxamide;

2-(3-(Aminomethyl)phenyl)-9-(2-methoxyphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;

2-(4-(1H-1,2,3-Triazol-5-yl)phenyl)-9-(2-isopropylphenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(4-(1H-1,2,4-Triazol-3-yl)phenyl)-9-(cis-4-methoxycyclohexyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(1H-Benzo[d]imidazol-6-yl)-9-(cis-4-methoxycyclohexyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(1H-Imidazo[4,5-b]pyridin-6-yl)-9-(cis-4-methoxycyclohexyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide;
 2-(3-Hydroxyphenyl)-9-((1r,4r)-4-(methoxymethyl)cyclohexyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide; and
 9-(2-Isopropylphenyl)-2-(4-(5-methyl-4H-1,2,4-triazol-3-yl)phenyl)-8-oxo-8,9-dihydro-7H-purine-6-carboxamide,
 and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof.

[00288] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (III):



and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:


R¹ is substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;


R^3 and R^4 are each independently H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclalkyl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkylalkyl, or R^3 and R^4 , together with the atoms to which they are attached, form a substituted or unsubstituted cycloalkyl or substituted or unsubstituted heterocyclyl;

or R^2 and one of R^3 and R^4 , together with the atoms to which they are attached, form a substituted or unsubstituted heterocyclyl,

wherein in certain embodiments, the TOR kinase inhibitors do not include the compounds depicted below, namely:




6-(4-hydroxyphenyl)-4-(3-methoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;



6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;


or,



(R)-6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-3-(cyclohexylmethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.


[00289] In some embodiments of compounds of formula (III), R¹ is substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl. In one embodiment, R¹ is phenyl, pyridyl, pyrimidyl, benzimidazolyl, indolyl, indazolyl, 1H-pyrrolo[2,3-b]pyridyl, 1H-imidazo[4,5-b]pyridyl, 1H-imidazo[4,5-b]pyridin-2(3H)-onyl, 3H-imidazo[4,5-b]pyridyl, or pyrazolyl, each optionally substituted. In some embodiments, R¹ is phenyl substituted with one or more substituents independently selected from the group consisting of substituted or unsubstituted C₁₋₈ alkyl (for example, methyl), substituted or unsubstituted heterocyclyl (for example, substituted or unsubstituted triazolyl or pyrazolyl), halogen (for example, fluorine), aminocarbonyl, cyano, hydroxyalkyl (for example, hydroxypropyl), and hydroxy. In other embodiments, R¹ is pyridyl substituted with one or more substituents independently selected from the group consisting of substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted heterocyclyl (for example, substituted or unsubstituted triazolyl), halogen, aminocarbonyl, cyano, hydroxyalkyl, -OR, and -NR₂, wherein each R is independently H, or a substituted or unsubstituted C₁₋₄ alkyl. In yet other embodiments, R¹ is 1H-pyrrolo[2,3-b]pyridyl or benzimidazolyl, each optionally substituted with one or more substituents independently selected from the group consisting of substituted or unsubstituted C₁₋₈ alkyl, and -NR₂, wherein each R is independently H, or a substituted or unsubstituted C₁₋₄ alkyl.

[00290] In some embodiments of compounds of formula (III), R^1 is



wherein R is at each occurrence independently H, or a substituted or unsubstituted C_{1-4} alkyl (for example, methyl); R' is at each occurrence independently a substituted or unsubstituted C_{1-4} alkyl, halogen (for example, fluorine), cyano, -OR, or -NR₂; m is 0-3; and n is 0-3. It will be understood by those skilled in the art that any of the substituents R' may be attached to any suitable atom of any of the rings in the fused ring systems. It will also be understood by those skilled in the art that the connecting bond of R^1 (designated by the bisecting wavy line) may be attached to any of the atoms in any of the rings in the fused ring systems.


[00291] In some embodiments of compounds of formula (III), R¹ is



wherein R is at each occurrence independently H, or a substituted or unsubstituted C₁₋₄ alkyl; R' is at each occurrence independently a substituted or unsubstituted C₁₋₄ alkyl, halogen, cyano, -OR, or -NR₂; m is 0-3; and n is 0-3.


[00292] In some embodiments of compounds of formula (III), R² is H, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted C₁₋₄ alkyl-heterocyclyl, substituted or unsubstituted C₁₋₄ alkyl-aryl, or substituted or unsubstituted C₁₋₄ alkyl-cycloalkyl. For example, R² is H, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, *tert*-butyl, n-pentyl, isopentyl, cyclopentyl, cyclohexyl, tetrahydrofuranyl, tetrahydropyranyl, (C₁₋₄ alkyl)-phenyl, (C₁₋₄ alkyl)-cyclopropyl, (C₁₋₄ alkyl)-cyclobutyl, (C₁₋₄ alkyl)-cyclopentyl, (C₁₋₄ alkyl)-cyclohexyl, (C₁₋₄ alkyl)-pyrrolidyl, (C₁₋₄ alkyl)-piperidyl, (C₁₋₄ alkyl)-piperazinyl, (C₁₋₄ alkyl)-morpholinyl, (C₁₋₄ alkyl)-tetrahydrofuranyl, or (C₁₋₄ alkyl)-tetrahydropyranyl, each optionally substituted.

[00293] In other embodiments, R^2 is H, C_{1-4} alkyl, $(C_{1-4}\text{alkyl})(\text{OR})$,



wherein R is at each occurrence independently H, or a substituted or unsubstituted C_{1-4} alkyl (for example, methyl); R' is at each occurrence independently H, -OR, cyano, or a substituted or unsubstituted C_{1-4} alkyl (for example, methyl); and p is 0-3.


[00294] In some such embodiments, R^2 is H, C_{1-4} alkyl, $(C_{1-4}\text{alkyl})(\text{OR})$,



wherein R is at each occurrence independently H, or a substituted or unsubstituted C_{1-2} alkyl; R' is at each occurrence independently H, -OR, cyano, or a substituted or unsubstituted C_{1-2} alkyl; and p is 0-1.

[00295] In some other embodiments of compounds of formula (III), R^2 and one of R^3 and R^4 together with the atoms to which they are attached form a substituted or

unsubstituted heterocyclyl. For example, in some embodiments, the compound of formula (III) is



wherein R is at each occurrence independently H, or a substituted or unsubstituted C₁₋₄ alkyl; R" is H, OR, or a substituted or unsubstituted C₁₋₄ alkyl; and R¹ is as defined herein.

[00296] In some embodiments of compounds of formula (III), R³ and R⁴ are both H. In others, one of R³ and R⁴ is H and the other is other than H. In still others, one of R³ and R⁴ is C₁₋₄ alkyl (for example, methyl) and the other is H. In still others, both of R³ and R⁴ are C₁₋₄ alkyl (for example, methyl).

[00297] In some such embodiments described above, R¹ is substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl. For example, R¹ is phenyl, pyridyl, pyrimidyl, benzimidazolyl, indolyl, indazolyl, 1H-pyrrolo[2,3-b]pyridyl, 1H-imidazo[4,5-b]pyridyl, 1H-imidazo[4,5-b]pyridin-2(3H)-onyl, 3H-imidazo[4,5-b]pyridyl, or pyrazolyl, each optionally substituted. In some embodiments, R¹ is phenyl substituted with one or more substituents independently selected from the group consisting of substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted heterocyclyl, halogen, aminocarbonyl, cyano, hydroxyalkyl and hydroxy. In others, R¹ is pyridyl substituted with one or more substituents independently selected from the group

consisting of cyano, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted heterocyclyl, hydroxyalkyl, halogen, aminocarbonyl, -OR, and -NR₂, wherein each R is independently H, or a substituted or unsubstituted C₁₋₄ alkyl. In others, R¹ is 1H-pyrrolo[2,3-b]pyridyl or benzimidazolyl, optionally substituted with one or more substituents independently selected from the group consisting of substituted or unsubstituted C₁₋₈ alkyl, and -NR₂, wherein R is independently H, or a substituted or unsubstituted C₁₋₄ alkyl

[00298] In certain embodiments, the compounds of formula (III) have an R¹ group set forth herein and an R² group set forth herein.

[00299] In some embodiments of compounds of formula (III), the compound at a concentration of 10 μM inhibits mTOR, DNA-PK, or PI3K or a combination thereof, by at least about 50%. Compounds of formula (III) may be shown to be inhibitors of the kinases above in any suitable assay system.

[00300] Representative TOR kinase inhibitors of formula (III) include:
 6-(1H-pyrrolo[2,3-b]pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 6-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((*trans*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((*cis*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((*trans*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((*trans*-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((*cis*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((*trans*-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((*cis*-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((*cis*-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((*trans*-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((*cis*-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((*cis*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-isopropyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((*cis*-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-((*cis*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-ethyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(*trans*-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-isopropyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-ethyl-6-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(3-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(3-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(*cis*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(3-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-4-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-(2-methoxyethyl)-6-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(3-(1H-1,2,4-triazol-5-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

5-(8-(2-methoxyethyl)-6-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-4-methylpicolinamide;

3-(6-oxo-8-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)benzamide;

3-(6-oxo-8-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)benzonitrile;

5-(8-(*trans*-4-methoxycyclohexyl)-6-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-4-methylpicolinamide;

6-(1H-imidazo[4,5-b]pyridin-6-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(1H-indazol-6-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-((1R,3S)-3-methoxycyclopentyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-((1S,3R)-3-methoxycyclopentyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-((1R,3R)-3-methoxycyclopentyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-((1S,3S)-3-methoxycyclopentyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-ethyl-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(1H-pyrrolo[2,3-b]pyridin-5-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(1H-indol-6-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(1H-indol-5-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-((1R,3S)-3-methoxycyclopentyl)methyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-((1S,3R)-3-methoxycyclopentyl)methyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(3-fluoro-2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(3-fluoro-2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-methoxycethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

3,3-dimethyl-6-(4-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((1R,3S)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((1S,3R)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(((1S,3S)-3-methoxycyclopentyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(((1R,3R)-3-methoxycyclopentyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((1S,3S)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((1R,3R)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(((1R,3S)-3-methoxycyclopentyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(((1S,3R)-3-methoxycyclopentyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(3-fluoro-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(3-fluoro-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7'-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1'-(tetrahydro-2H-pyran-4-yl)methyl)-1'H-spiro[cyclopentane-1,2'-pyrazino[2,3-b]pyrazin]-3'(4'H)-one;

7'-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1'-(tetrahydro-2H-pyran-4-yl)methyl)-1'H-spiro[cyclobutane-1,2'-pyrazino[2,3-b]pyrazin]-3'(4'H)-one;

4-(cyclopropylmethyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7'-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1'H-spiro[cyclopentane-1,2'-pyrazino[2,3-b]pyrazin]-3'(4'H)-one;

7'-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1'H-spiro[cyclobutane-1,2'-pyrazino[2,3-b]pyrazin]-3'(4'H)-one;

7'-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1'H-spiro[cyclopropane-1,2'-pyrazino[2,3-b]pyrazin]-3'(4'H)-one;

(R)-6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-((tetrahydrofuran-2-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(S)-6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-((tetrahydrofuran-2-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(1H-indazol-5-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-(6-oxo-8-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)benzamide;

4-(2-methoxyethyl)-3,3-dimethyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-ethyl-3,3-dimethyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

3,3-dimethyl-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(R)-6-(6-(1-hydroxyethyl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

3,3-dimethyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)-4-methylpyridin-3-yl)-4-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)-4-methylpyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

3,3-dimethyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

3,3-dimethyl-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)-2-methylpyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)-2-methylpyridin-3-yl)-4-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(S)-6-(6-(1-hydroxyethyl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

3,3-dimethyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,3-dimethyl-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(4-(2-hydroxypropan-2-yl)phenyl)-4-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(4-(2-hydroxypropan-2-yl)phenyl)-4-((*trans*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-(*cis*-4-methoxycyclohexyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-(*trans*-4-methoxycyclohexyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(4-(2-hydroxypropan-2-yl)phenyl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-(2-methoxyethyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

9-(6-(4H-1,2,4-triazol-3-yl)-3-pyridyl)-6,11,4a-trihydromorpholino[4,3-e]pyrazino[2,3-b]pyrazin-5-one;

6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

5-(8-(*cis*-4-methoxycyclohexyl)-6-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-6-methylpicolinonitrile;

6-(6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-3-(2-methoxyacetyl)-6,11,4a-trihydropiperazino[1,2-e]pyrazino[2,3-b]pyrazin-5-one;

9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-6,11,4a-trihydropiperazino[1,2-e]pyrazino[2,3-b]pyrazin-5-one;

9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-3-(2-methoxyethyl)-6,11,4a-trihydropiperazino[1,2-e]pyrazino[2,3-b]pyrazin-5-one;

4-(cyclopentylmethyl)-6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

9-(6-(4H-1,2,4-triazol-3-yl)-2-methyl-3-pyridyl)-6,11,4a-trihydromorpholino[4,3-e]pyrazino[2,3-b]pyrazin-5-one;

4-(*trans*-4-hydroxycyclohexyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-(*cis*-4-hydroxycyclohexyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((tetrahydrofuran-3-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-(cyclopentylmethyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-neopentyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-isobutyl-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

3-methyl-6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(piperidin-4-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-3-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

8-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)(3aS,2R)-2-methoxy-5,10,3a-trihdropyrazino[2,3-b]pyrrolidino[1,2-e]pyrazin-4-one;

8-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)(2R,3aR)-2-methoxy-5,10,3a-trihdropyrazino[2,3-b]pyrrolidino[1,2-e]pyrazin-4-one;

8-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)(2S,3aR)-2-methoxy-5,10,3a-trihdropyrazino[2,3-b]pyrrolidino[1,2-e]pyrazin-4-one;

8-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)(2S,3aS)-2-methoxy-5,10,3a-trihdropyrazino[2,3-b]pyrrolidino[1,2-e]pyrazin-4-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(3-methoxypropyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

(S)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((tetrahydrofuran-2-yl)methyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

(R)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((tetrahydrofuran-2-yl)methyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-3-methyl-6,11,4a-trihydropiperazino[1,2-e]pyrazino[2,3-b]pyrazin-5-one;

9-(4-(4H-1,2,4-triazol-3-yl)phenyl)-6,11,4a-trihydromorpholino[4,3-c]pyrazino[2,3-b]pyrazin-5-one;

9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-6,11,4a-trihydropiperidino[1,2-e]pyrazino[2,3-b]pyrazin-5-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(*cis*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(2-morpholinoethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-phenethyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-(cyclohexylmethyl)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((*trans*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((*cis*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(R)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(tetrahydrofuran-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(S)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(tetrahydrofuran-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-phenyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(S)-6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3-methyl-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

9-[6-(1-hydroxy-isopropyl)-3-pyridyl]-6,11,4a-trihydromorpholino[4,3-c]pyrazino[2,3-b]pyrazin-5-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(2-amino-7-methyl-1H-benzo[d]imidazol-5-yl)-4-(3-(trifluoromethyl)benzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(3-(trifluoromethyl)benzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

9-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-6,11,4a-trihydromorpholino[4,3-e]pyrazino[2,3-b]pyrazin-5-one;

6-(4-methyl-2-(methylamino)-1H-benzo[d]imidazol-6-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

8-(4-(4H-1,2,4-triazol-3-yl)-2-methylphenyl)-5,10,3a-trihydropyrazino[2,3-b]pyrrolidino[1,2-e]pyrazin-4-one;

6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-ethyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;


6-(4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(3-(trifluoromethyl)benzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(4-methyl-1H-benzo[d]imidazol-6-yl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

6-(4-(2-hydroxypropan-2-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one; and
 6-(4-(1H-1,2,4-triazol-5-yl)phenyl)-4-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one,
 and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof.

[00301] In one embodiment, the TOR kinase inhibitors include compounds having the following formula (IV):



(IV)


and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof, wherein:

R^1 is substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted aryl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, or substituted or unsubstituted heterocyclylalkyl;

R^2 is H, substituted or unsubstituted C_{1-8} alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted heterocyclylalkyl, substituted or unsubstituted aralkyl, or substituted or unsubstituted cycloalkylalkyl;


R^3 is H, or a substituted or unsubstituted C_{1-8} alkyl,

wherein in certain embodiments, the TOR kinase inhibitors do not include 7-(4-hydroxyphenyl)-1-(3-methoxybenzyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one, depicted below:




[00302] In some embodiments of compounds of formula (IV), R¹ is substituted or unsubstituted aryl or substituted or unsubstituted heteroaryl. For example, R¹ is phenyl, pyridyl, pyrimidyl, benzimidazolyl, 1H-pyrrolo[2,3-b]pyridyl, indazolyl, indolyl, 1H-imidazo[4,5-b]pyridyl, 1H-imidazo[4,5-b]pyridin-2(3H)-onyl, 3H-imidazo[4,5-b]pyridyl, or pyrazolyl, each optionally substituted. In some embodiments, R¹ is phenyl substituted with one or more substituents independently selected from the group consisting of substituted or unsubstituted C₁₋₈ alkyl (for example, methyl), substituted or unsubstituted heterocyclyl (for example, a substituted or unsubstituted triazolyl or pyrazolyl), aminocarbonyl, halogen (for example, fluorine), cyano, hydroxyalkyl and hydroxy. In other embodiments, R¹ is pyridyl substituted with one or more substituents independently selected from the group consisting of substituted or unsubstituted C₁₋₈ alkyl (for example, methyl), substituted or unsubstituted heterocyclyl (for example, a substituted or unsubstituted triazolyl), halogen, aminocarbonyl, cyano, hydroxyalkyl (for example, hydroxypropyl), -OR, and -NR₂, wherein each R is independently H, or a substituted or unsubstituted C₁₋₄ alkyl. In some embodiments, R¹ is 1H-pyrrolo[2,3-b]pyridyl or benzimidazolyl, optionally substituted with one or more substituents independently selected from the group consisting of substituted or unsubstituted C₁₋₈ alkyl, and -NR₂, wherein R is independently H, or a substituted or unsubstituted C₁₋₄ alkyl.

[00303] In some embodiments, R¹ is




[00304] In some embodiments of compounds of formula (IV), R¹ is



wherein R is at each occurrence independently H, or a substituted or unsubstituted C₁₋₄ alkyl; R' is at each occurrence independently a substituted or unsubstituted C₁₋₄ alkyl, halogen, cyano, -OR or -NR₂; m is 0-3; and n is 0-3.


[00305] In some embodiments of compounds of formula (IV), R² is H, substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclyl, substituted or unsubstituted C₁₋₄ alkyl-heterocyclyl, substituted or unsubstituted C₁₋₄ alkyl-aryl, or substituted or unsubstituted C₁₋₄ alkyl-cycloalkyl. For example, R² is H, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, *tert*-butyl, n-pentyl, isopentyl, cyclopentyl, cyclohexyl, tetrahydrofuran, tetrahydropyran, (C₁₋₄ alkyl)-phenyl, (C₁₋₄ alkyl)-cyclopropyl, (C₁₋₄ alkyl)-cyclobutyl, (C₁₋₄ alkyl)-cyclopentyl, (C₁₋₄ alkyl)-cyclohexyl, (C₁₋₄ alkyl)-pyrrolidyl, (C₁₋₄ alkyl)-piperidyl, (C₁₋₄ alkyl)-piperazinyl, (C₁₋₄ alkyl)-morpholinyl, (C₁₋₄ alkyl)-tetrahydrofuran, or (C₁₋₄ alkyl)-tetrahydropyran, each optionally substituted.

[00306] In other embodiments, R² is H, C₁₋₄ alkyl, (C₁₋₄alkyl)(OR),



wherein R is at each occurrence independently H, or a substituted or unsubstituted C₁₋₄ alkyl (for example, methyl); R' is at each occurrence independently H, -OR, cyano, or a substituted or unsubstituted C₁₋₄ alkyl (for example, methyl); and p is 0-3.

[00307] In other embodiments of compounds of formula (IV), R² is H, C₁₋₄ alkyl, (C₁₋₄alkyl)(OR),



wherein R is at each occurrence independently H, or a substituted or unsubstituted C₁₋₂ alkyl; R' is at each occurrence independently H, -OR, cyano, or a substituted or unsubstituted C₁₋₂ alkyl; and p is 0-1.

[00308] In other embodiments of compounds of formula (IV), R³ is H.

[00309] In some such embodiments described herein, R¹ is substituted or unsubstituted aryl, or substituted or unsubstituted heteroaryl. For example, R¹ is phenyl, pyridyl, pyrimidyl, benzimidazolyl, 1H-pyrrolo[2,3-b]pyridyl, indazolyl, indolyl, 1H-imidazo[4,5-b]pyridine, pyridyl, 1H-imidazo[4,5-b]pyridin-2(3H)-onyl, 3H-imidazo[4,5-b]pyridyl, or pyrazolyl, each optionally substituted. In some embodiments, R¹ is phenyl substituted with one or more substituents independently selected from the group consisting of substituted or unsubstituted C₁₋₈ alkyl, substituted or unsubstituted heterocyclyl, aminocarbonyl, halogen, cyano, hydroxyalkyl and hydroxy. In others, R¹ is pyridyl substituted with one or more substituents independently selected from the group consisting of C₁₋₈ alkyl, substituted or unsubstituted heterocyclyl, halogen, aminocarbonyl, cyano, hydroxyalkyl, -OR, and -NR₂, wherein each R is independently H, or a substituted or unsubstituted C₁₋₄ alkyl. In still others, R¹ is 1H-pyrrolo[2,3-b]pyridyl or benzimidazolyl, optionally substituted with one or more substituents independently selected from the group consisting of substituted or unsubstituted C₁₋₈ alkyl, and -NR₂, wherein R is independently H, or a substituted or unsubstituted C₁₋₄ alkyl.

[00310] In certain embodiments, the compounds of formula (IV) have an R¹ group set forth herein and an R² group set forth herein.

[00311] In some embodiments of compounds of formula (IV), the compound at a concentration of 10 µM inhibits mTOR, DNA-PK, PI3K, or a combination thereof by at least about 50%. Compounds of formula (IV) may be shown to be inhibitors of the kinases above in any suitable assay system.

[00312] Representative TOR kinase inhibitors of formula (IV) include:
 7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-((*trans*-4-methoxycyclohexyl)methyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((*cis*-4-methoxycyclohexyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(1H-pyrrolo[2,3-b]pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-((*cis*-4-methoxycyclohexyl)methyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 1-ethyl-7-(1H-pyrrolo[3,2-b]pyridin-5-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((*cis*-4-methoxycyclohexyl)methyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(1H-benzo[d]imidazol-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(1H-pyrrolo[2,3-b]pyridin-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((*trans*-4-methoxycyclohexyl)methyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((*trans*-4-hydroxycyclohexyl)methyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((*cis*-4-hydroxycyclohexyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-((*cis*-4-hydroxycyclohexyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(tetrahydro-2H-pyran-4-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(2-methoxyethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-ethyl-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-((*cis*-4-hydroxycyclohexyl)methyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(tetrahydro-2H-pyran-4-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(1H-indol-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-((*trans*-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((*cis*-4-hydroxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(*trans*-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-isopropyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(*trans*-4-hydroxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-isopropyl-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-ethyl-7-(5-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(2-hydroxypyridin-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-isopropyl-7-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

5-(8-isopropyl-7-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-4-methylpicolinamide;

7-(1H-indazol-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(2-aminopyrimidin-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(2-aminopyridin-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(methylamino)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-hydroxypyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(4-(1H-pyrazol-3-yl)phenyl)-1-(2-methoxyethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(1H-indazol-4-yl)-1-(2-methoxyethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(1H-indazol-6-yl)-1-(2-methoxyethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(pyrimidin-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-methoxypyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

1-(2-methoxyethyl)-7-(1H-pyrrolo[2,3-b]pyridin-5-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

1-ethyl-7-(1H-pyrrolo[2,3-b]pyridin-5-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

1-ethyl-7-(1H-indazol-4-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(pyridin-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-aminopyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

1-methyl-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

2-(2-hydroxypropan-2-yl)-5-(8-(*trans*-4-methoxycyclohexyl)-7-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)pyridine 1-oxide;

4-methyl-5-(7-oxo-8-((tetrahydro-2H-pyran-4-yl)methyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)picolinamide;

5-(8-((*cis*-4-methoxycyclohexyl)methyl)-7-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-4-methylpicolinamide;

7-(1H-pyrazol-4-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

1-(*trans*-4-methoxycyclohexyl)-7-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

3-((7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-2-oxo-3,4-dihdropyrazino[2,3-b]pyrazin-1(2H)-yl)methyl)benzonitrile;

1-((*trans*-4-methoxycyclohexyl)methyl)-7-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

3-(7-oxo-8-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)benzamide;

5-(8-((*trans*-4-methoxycyclohexyl)methyl)-7-oxo-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)-4-methylpicolinamide;

3-((7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-2-oxo-3,4-dihdropyrazino[2,3-b]pyrazin-1(2H)-yl)methyl)benzonitrile;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((1R,3R)-3-methoxycyclopentyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((1S,3R)-3-methoxycyclopentyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((1S,3S)-3-methoxycyclopentyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((1R,3S)-3-methoxycyclopentyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(1H-indazol-6-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(2-morpholinoethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-(*trans*-4-hydroxycyclohexyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-(*cis*-4-hydroxycyclohexyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(2-morpholinoethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-isopropyl-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(1H-imidazo[4,5-b]pyridin-6-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-((*cis*-4-methoxycyclohexyl)methyl)-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-(*trans*-4-hydroxycyclohexyl)-7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-(*cis*-4-hydroxycyclohexyl)-7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

4-(7-oxo-8-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-5,6,7,8-tetrahydropyrazino[2,3-b]pyrazin-2-yl)benzamide;

7-(1H-indazol-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(1H-pyrrolo[2,3-b]pyridin-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-((1S,3R)-3-methoxycyclopentyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-((1R,3R)-3-methoxycyclopentyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-((1R,3S)-3-methoxycyclopentyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-((1S,3S)-3-methoxycyclopentyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(1H-indol-5-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-ethyl-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(1H-indol-6-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(4-(2-hydroxypropan-2-yl)phenyl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(tetrahydro-2H-pyran-4-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-((*trans*-4-methoxycyclohexyl)methyl)-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((*cis*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

1-(2-methoxyethyl)-7-(4-methyl-2-(methylamino)-1H-benzo[d]imidazol-6-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(7-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 1-(2-methoxyethyl)-7-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 1-benzyl-7-(2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(3-fluoro-4-(4H-1,2,4-triazol-3-yl)phenyl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(3-fluoro-4-(4H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(3-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 1-(*trans*-4-methoxycyclohexyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(*trans*-4-methoxycyclohexyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(5-fluoro-2-methyl-4-(4H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(3-fluoro-2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 1-(2-methoxyethyl)-7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((*trans*-4-methoxycyclohexyl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 1-(cyclopentylmethyl)-7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;
 7-(4-(2-hydroxypropan-2-yl)phenyl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(S)-7-(6-(1-hydroxyethyl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(R)-7-(6-(1-hydroxyethyl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(4-(2-hydroxypropan-2-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(4-(trifluoromethyl)benzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(3-(trifluoromethyl)benzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(3-methoxypropyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(4-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(2-methoxyethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(4-methyl-2-(methylamino)-1H-benzo[d]imidazol-6-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(2-amino-4-methyl-1H-benzo[d]imidazol-6-yl)-1-((tetrahydro-2H-pyran-4-yl)methyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

7-(2-methyl-6-(4H-1,2,4-triazol-3-yl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(R)-7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3-methyl-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one;

(S)-7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3-methyl-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-3,3-dimethyl-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-amino-4-methyl-1H-benzo[d]imidazol-6-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(6-(2-hydroxypropan-2-yl)pyridin-3-yl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(2-methyl-4-(1H-1,2,4-triazol-3-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
7-(4-(1H-1,2,4-triazol-5-yl)phenyl)-1-(2-(tetrahydro-2H-pyran-4-yl)ethyl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one;
1-(1-hydroxypropan-2-yl)-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one; and
1-(2-hydroxyethyl)-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihdropyrazino[2,3-b]pyrazin-2(1H)-one,
and pharmaceutically acceptable salts, clathrates, solvates, stereoisomers, tautomers, and prodrugs thereof.

4.4 METHODS FOR MAKING TOR KINASE INHIBITORS

[00313] The TOR kinase inhibitors can be obtained via standard, well-known synthetic methodology, see *e.g.*, March, J. Advanced Organic Chemistry; Reactions Mechanisms, and Structure, 4th ed., 1992. Starting materials useful for preparing compounds of formula (III) and intermediates therefore, are commercially available or can be prepared from commercially available materials using known synthetic methods and reagents.

[00314] Particular methods for preparing compounds of formula (I) are disclosed in U.S. Patent No. 7,981,893, issued July 19, 2011. Particular methods for preparing compounds of formula (II) are disclosed in U.S.

Patent No. 7,968,556, issued June 28, 2011.

Particular methods for preparing compounds of formula (III) and (IV) are disclosed in U.S. Patent No. 8,110,578, issued February 7, 2012, and U.S. Publication No. 2011/0137028, filed October 25, 2010.

4.5 METHODS OF USE

[00315] Provided herein are methods for treating or preventing E-twenty six (ETS) overexpressing castration-resistant prostate cancer, comprising administering an effective amount of a TOR kinase inhibitor to a patient having ETS overexpressing castration-resistant prostate cancer. In certain embodiments, a TOR kinase inhibitor is administered to a patient who has locally advanced, recurrent or metastatic, ETS overexpressing castration-resistant prostate cancer not amenable to curative surgical resection. In another embodiment, a TOR kinase inhibitor is administered to a patient who has received at least one prior line of platinum based chemotherapy. In some embodiments, a TOR kinase inhibitor is administered to a patient who has a tumor showing DNA-PK overexpression.

[00316] In some embodiments, provided herein are methods for treating E-twenty six (ETS) overexpressing castration-resistant prostate cancer, the methods comprising administering an effective amount of a TOR kinase inhibitor to a patient having E-twenty six (ETS) overexpressing castration-resistant prostate cancer, wherein the treatment results in one or more of inhibition of disease progression, inhibition of tumor growth, reduction of primary tumor, relief of tumor-related symptoms, inhibition of tumor secreted factors (including tumor secreted hormones, such as those that contribute to carcinoid syndrome), delayed appearance of primary or secondary tumors, slowed development of primary or secondary tumors, decreased occurrence of primary or secondary tumors, slowed or decreased severity of secondary effects of disease, arrested tumor growth and regression of tumors, increased Time To Progression (TTP), increased Progression Free Survival (PFS), and/or increased Overall Survival (OS), among others.

[00317] In certain embodiments, provided herein are methods for improving the Prostate-Specific Antigen Working Group 2 (PSAWG2) Criteria for prostate cancer (*see*

Scher, H., Halab, S., Tannock, S., Morris, M., Sternberg, C. N., et al. Design and end points of clinical trials for patients with progressive prostate cancer and castrate levels of testosterone: Recommendations of the Prostate Cancer Clinical Trials Working Group. *J Clin Oncol.* 2008; (26) 148-1159) of a patient, comprising administering an effective amount of a TOR kinase inhibitor to a patient having ETS overexpressing castration-resistant prostate cancer.

[00318] In one embodiment, provided herein are methods for inhibiting phosphorylation of S6RP, 4E-BP1 and/or AKT in a patient having ETS overexpressing castration-resistant prostate cancer, comprising administering an effective amount of a TOR kinase inhibitor to said patient. In some such embodiments, the inhibition of phosphorylation is assessed in a biological sample of the patient, such as in circulating blood and/or tumor cells, skin biopsies and/or tumor biopsies or aspirate. In such embodiments, the amount of inhibition of phosphorylation is assessed by comparison of the amount of phospho- S6RP, 4E-BP1 and/or AKT before and after administration of the TOR kinase inhibitor. In certain embodiments, provided herein are methods for measuring inhibition of phosphorylation of S6RP, 4E-BP1 or AKT in a patient having ETS overexpressing castration-resistant prostate cancer, comprising administering an effective amount of a TOR kinase inhibitor to said patient, measuring the amount of phosphorylated S6RP, 4E BP1 and/or AKT in said patient, and comparing said amount of phosphorylated S6RP, 4E BP1 and/or AKT to that of said patient prior to administration of an effective amount of a TOR kinase inhibitor.

[00319] In certain embodiments, provided herein are methods for inhibiting phosphorylation of S6RP, 4E-BP1 and/or AKT in a biological sample of a patient having ETS overexpressing castration-resistant prostate cancer, comprising administering an effective amount of a TOR kinase inhibitor to said patient and comparing the amount of phosphorylated S6RP, 4E-BP1 and/or AKT in a biological sample of a patient obtained prior to and after administration of said TOR kinase inhibitor, wherein less phosphorylated S6RP, 4E-BP1 and/or AKT in said biological sample obtained after administration of said

TOR kinase inhibitor relative to the amount of phosphorylated S6RP, 4E-BP1 and/or AKT in said biological sample obtained prior to administration of said TOR kinase inhibitor indicates inhibition.

[00320] In one embodiment, provided herein are methods for inhibiting DNA-dependent protein kinase (DNA-PK) activity in a patient having ETS overexpressing castration-resistant prostate cancer, comprising administering an effective amount of a TOR kinase inhibitor to said patient. In some embodiments, DNA-PK inhibition is assessed in the skin of the patient having ETS overexpressing castration-resistant prostate cancer, in one example in a UV light-irradiated skin sample of said patient. In another embodiment, DNA-PK inhibition is assessed in a tumor biopsy or aspirate of a patient having ETS overexpressing castration-resistant prostate cancer. In one embodiment, inhibition is assessed by measuring the amount of phosphorylated DNA-PK S2056 (also known as pDNA-PK S2056) before and after administration of the TOR kinase inhibitor. In certain embodiments, provided herein are methods for measuring inhibition of phosphorylation of DNA-PK S2056 in a skin sample of a patient having ETS overexpressing castration-resistant prostate cancer, comprising administering an effective amount of a TOR kinase inhibitor to said patient, measuring the amount of phosphorylated DNA-PK S2056 present in the skin sample and comparing said amount of phosphorylated DNA-PK S2056 to that in a skin sample from said patient prior to administration of an effective amount of a TOR kinase inhibitor. In one embodiment, the skin sample is irradiated with UV light.

[00321] In certain embodiments, provided herein are methods for inhibiting DNA-dependent protein kinase (DNA-PK) activity in a skin sample of a patient having ETS overexpressing castration-resistant prostate cancer, comprising administering an effective amount of a TOR kinase inhibitor to said patient and comparing the amount of phosphorylated DNA-PK in a biological sample of a patient obtained prior to and after administration of said TOR kinase inhibitor, wherein less phosphorylated DNA-PK in said biological sample obtained after administration of said TOR kinase inhibitor relative to the

amount of phosphorylated DNA-PK in said biological sample obtained prior to administration of said TOR kinase inhibitor indicates inhibition.

[00322] In some embodiments, the TOR kinase inhibitor is a compound as described herein. In one embodiment, the TOR kinase inhibitor is Compound 1 (a TOR kinase inhibitor set forth herein having molecular formula C₁₆H₁₆N₈O). In one embodiment, Compound 1 is 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one.

[00323] A TOR kinase inhibitor can be combined with radiation therapy or surgery. In certain embodiments, a TOR kinase inhibitor is administered to patient who is undergoing radiation therapy, has previously undergone radiation therapy or will be undergoing radiation therapy. In certain embodiments, a TOR kinase inhibitor is administered to a patient who has undergone tumor removal surgery.

[00324] Further provided herein are methods for treating patients who have been previously treated for ETS overexpressing castration-resistant prostate cancer, but are non-responsive to standard therapies, as well as those who have not previously been treated. Further provided herein are methods for treating patients who have undergone surgery in an attempt to treat the condition at issue, as well as those who have not. Because patients with ETS overexpressing castration-resistant prostate cancer may have heterogenous clinical manifestations and varying clinical outcomes, the treatment given to a patient may vary, depending on his/her prognosis. The skilled clinician will be able to readily determine without undue experimentation specific secondary agents, types of surgery, and types of non-drug based standard therapy that can be effectively used to treat an individual patient with ETS overexpressing castration-resistant prostate cancer.

[00325] In one embodiment, the ETS overexpressing castration-resistant prostate cancer is that in which the PI3K/mTOR pathway is activated. In certain embodiments, the ETS overexpressing castration-resistant prostate cancer is that in which the PI3K/mTOR pathway is activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

4.6 PHARMACEUTICAL COMPOSITIONS AND ROUTES OF ADMINISTRATION

[00326] Provided herein are compositions, comprising an effective amount of a TOR kinase inhibitor, and compositions comprising an effective amount of a TOR kinase inhibitor and a pharmaceutically acceptable carrier or vehicle. In some embodiments, the pharmaceutical compositions described herein are suitable for oral, parenteral, mucosal, transdermal or topical administration.

[00327] The TOR kinase inhibitors can be administered to a patient orally or parenterally in the conventional form of preparations, such as capsules, microcapsules, tablets, granules, powder, troches, pills, suppositories, injections, suspensions and syrups. Suitable formulations can be prepared by methods commonly employed using conventional, organic or inorganic additives, such as an excipient (*e.g.*, sucrose, starch, mannitol, sorbitol, lactose, glucose, cellulose, talc, calcium phosphate or calcium carbonate), a binder (*e.g.*, cellulose, methylcellulose, hydroxymethylcellulose, polypropylpyrrolidone, polyvinylpyrrolidone, gelatin, gum arabic, polyethyleneglycol, sucrose or starch), a disintegrator (*e.g.*, starch, carboxymethylcellulose, hydroxypropylstarch, low substituted hydroxypropylcellulose, sodium bicarbonate, calcium phosphate or calcium citrate), a lubricant (*e.g.*, magnesium stearate, light anhydrous silicic acid, talc or sodium lauryl sulfate), a flavoring agent (*e.g.*, citric acid, menthol, glycine or orange powder), a preservative (*e.g.*, sodium benzoate, sodium bisulfite, methylparaben or propylparaben), a stabilizer (*e.g.*, citric acid, sodium citrate or acetic acid), a suspending agent (*e.g.*, methylcellulose, polyvinyl pyrrolidone or aluminum stearate), a dispersing agent (*e.g.*, hydroxypropylmethylcellulose), a diluent (*e.g.*, water), and base wax (*e.g.*, cocoa butter, white petrolatum or polyethylene glycol). The effective amount of the TOR kinase inhibitor in the pharmaceutical composition may be at a level that will exercise the desired effect; for example, about 0.005 mg/kg of a patient's body weight to about 10 mg/kg of a patient's body weight in unit dosage for both oral and parenteral administration.

[00328] The dose of a TOR kinase inhibitor to be administered to a patient is rather widely variable and can be subject to the judgment of a health-care practitioner. In general,

the TOR kinase inhibitors can be administered one to four times a day in a dose of about 0.005 mg/kg of a patient's body weight to about 10 mg/kg of a patient's body weight in a patient, but the above dosage may be properly varied depending on the age, body weight and medical condition of the patient and the type of administration. In one embodiment, the dose is about 0.01 mg/kg of a patient's body weight to about 5 mg/kg of a patient's body weight, about 0.05 mg/kg of a patient's body weight to about 1 mg/kg of a patient's body weight, about 0.1 mg/kg of a patient's body weight to about 0.75 mg/kg of a patient's body weight, about 0.25 mg/kg of a patient's body weight to about 0.5 mg/kg of a patient's body weight, or about 0.007 mg/kg of a patient's body weight to about 1.7 mg/kg of patient's body weight. In one embodiment, one dose is given per day. In another embodiment, two doses are given per day. In any given case, the amount of the TOR kinase inhibitor administered will depend on such factors as the solubility of the active component, the formulation used and the route of administration.

[00329] In another embodiment, provided herein are methods for the treatment or prevention of ETS overexpressing castration-resistant prostate cancer, comprising the administration of about 0.375 mg/day to about 750 mg/day, about 0.75 mg/day to about 375 mg/day, about 3.75 mg/day to about 75 mg/day, about 7.5 mg/day to about 55 mg/day, about 18 mg/day to about 37 mg/day, about 0.5 mg/day to about 60 mg/day, or about 0.5 mg/day to about 128 mg/day of a TOR kinase inhibitor to a patient in need thereof. In another embodiment, provided herein are methods for the treatment or prevention of ETS overexpressing castration-resistant prostate cancer, comprising the administration of about 0.5 mg/day to about 1200 mg/day, about 10 mg/day to about 1200 mg/day, about 100 mg/day to about 1200 mg/day, about 400 mg/day to about 1200 mg/day, about 600 mg/day to about 1200 mg/day, about 400 mg/day to about 800 mg/day or about 600 mg/day to about 800 mg/day of a TOR kinase inhibitor to a patient in need thereof. In a particular embodiment, the methods disclosed herein comprise the administration of 0.5 mg/day, 1 mg/day, 2 mg/day, 4 mg/day, 8 mg/day, 16 mg/day, 20 mg/day, 25 mg/day,

30 mg/day, 45 mg/day, 60 mg/day, 90 mg/day, 120 mg/day or 128 mg/day of a TOR kinase inhibitor to a patient in need thereof.

[00330] In another embodiment, provided herein are unit dosage formulations that comprise between about 0.1 mg and about 2000 mg, about 1 mg and 200 mg, about 35 mg and about 1400 mg, about 125 mg and about 1000 mg, about 250 mg and about 1000 mg, or about 500 mg and about 1000 mg of a TOR kinase inhibitor.

[00331] In a particular embodiment, provided herein are unit dosage formulation comprising about 0.1 mg, 0.25 mg, 0.5 mg, 1 mg, 5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 45 mg, 50 mg, 60 mg, 75 mg, 100 mg, 125 mg, 150 mg, 200 mg, 250 mg, 300 mg, 400 mg, 600 mg or 800 mg of a TOR kinase inhibitor.

[00332] In another embodiment, provided herein are unit dosage formulations that comprise 0.1 mg, 0.25 mg, 0.5 mg, 1 mg, 2.5 mg, 5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 35 mg, 50 mg, 70 mg, 100 mg, 125 mg, 140 mg, 175 mg, 200 mg, 250 mg, 280 mg, 350 mg, 500 mg, 560 mg, 700 mg, 750 mg, 1000 mg or 1400 mg of a TOR kinase inhibitor. In a particular embodiment, provided herein are unit dosage formulations that comprise 5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 45 mg or 60 mg of a TOR kinase inhibitor.

[00333] A TOR kinase inhibitor can be administered once, twice, three, four or more times daily.

[00334] A TOR kinase inhibitor can be administered orally for reasons of convenience. In one embodiment, when administered orally, a TOR kinase inhibitor is administered with a meal and water. In another embodiment, the TOR kinase inhibitor is dispersed in water or juice (e.g., apple juice or orange juice) and administered orally as a suspension. In another embodiment, when administered orally, a TOR kinase inhibitor is administered in a fasted state.

[00335] The TOR kinase inhibitor can also be administered intradermally, intramuscularly, intraperitoneally, percutaneously, intravenously, subcutaneously, intranasally, epidurally, sublingually, intracerebrally, intravaginally, transdermally, rectally, mucosally, by inhalation, or topically to the ears, nose, eyes, or skin. The mode of

administration is left to the discretion of the health-care practitioner, and can depend in-part upon the site of the medical condition.

[00336] In one embodiment, provided herein are capsules containing a TOR kinase inhibitor without an additional carrier, excipient or vehicle.

[00337] In another embodiment, provided herein are compositions, comprising an effective amount of a TOR kinase inhibitor and a pharmaceutically acceptable carrier or vehicle, wherein a pharmaceutically acceptable carrier or vehicle can comprise an excipient, diluent, or a mixture thereof. In one embodiment, the composition is a pharmaceutical composition.

[00338] The compositions can be in the form of tablets, chewable tablets, capsules, solutions, parenteral solutions, troches, suppositories and suspensions and the like.

Compositions can be formulated to contain a daily dose, or a convenient fraction of a daily dose, in a dosage unit, which may be a single tablet or capsule or convenient volume of a liquid. In one embodiment, the solutions are prepared from water-soluble salts, such as the hydrochloride salt. In general, all of the compositions are prepared according to known methods in pharmaceutical chemistry. Capsules can be prepared by mixing a TOR kinase inhibitor with a suitable carrier or diluent and filling the proper amount of the mixture in capsules. The usual carriers and diluents include, but are not limited to, inert powdered substances such as starch of many different kinds, powdered cellulose, especially crystalline and microcrystalline cellulose, sugars such as fructose, mannitol and sucrose, grain flours and similar edible powders.

[00339] Tablets can be prepared by direct compression, by wet granulation, or by dry granulation. Their formulations usually incorporate diluents, binders, lubricants and disintegrators as well as the compound. Typical diluents include, for example, various types of starch, lactose, mannitol, kaolin, calcium phosphate or sulfate, inorganic salts such as sodium chloride and powdered sugar. Powdered cellulose derivatives are also useful. In one embodiment, the pharmaceutical composition is lactose-free. Typical tablet binders are substances such as starch, gelatin and sugars such as lactose, fructose, glucose and the like.

Natural and synthetic gums are also convenient, including acacia, alginates, methylcellulose, polyvinylpyrrolidine and the like. Polyethylene glycol, ethylcellulose and waxes can also serve as binders.

[00340] A lubricant might be necessary in a tablet formulation to prevent the tablet and punches from sticking in the die. The lubricant can be chosen from such slippery solids as talc, magnesium and calcium stearate, stearic acid and hydrogenated vegetable oils. Tablet disintegrators are substances that swell when wetted to break up the tablet and release the compound. They include starches, clays, celluloses, algins and gums. More particularly, corn and potato starches, methylcellulose, agar, bentonite, wood cellulose, powdered natural sponge, cation-exchange resins, alginic acid, guar gum, citrus pulp and carboxymethyl cellulose, for example, can be used as well as sodium lauryl sulfate. Tablets can be coated with sugar as a flavor and sealant, or with film-forming protecting agents to modify the dissolution properties of the tablet. The compositions can also be formulated as chewable tablets, for example, by using substances such as mannitol in the formulation.

[00341] When it is desired to administer a TOR kinase inhibitor as a suppository, typical bases can be used. Cocoa butter is a traditional suppository base, which can be modified by addition of waxes to raise its melting point slightly. Water-miscible suppository bases comprising, particularly, polyethylene glycols of various molecular weights are in wide use.

[00342] The effect of the TOR kinase inhibitor can be delayed or prolonged by proper formulation. For example, a slowly soluble pellet of the TOR kinase inhibitor can be prepared and incorporated in a tablet or capsule, or as a slow-release implantable device. The technique also includes making pellets of several different dissolution rates and filling capsules with a mixture of the pellets. Tablets or capsules can be coated with a film that resists dissolution for a predictable period of time. Even the parenteral preparations can be made long-acting, by dissolving or suspending the TOR kinase inhibitor in oily or emulsified vehicles that allow it to disperse slowly in the serum.

4.7 KITS

[00343] In certain embodiments, provided herein are kits comprising a TOR kinase inhibitor.

[00344] In other embodiments, provide herein are kits comprising a TOR kinase inhibitor and means for monitoring patient response to administration of said TOR kinase inhibitor. In certain embodiments, the patient has ETS overexpressing castration-resistant prostate cancer. In particular embodiments, the patient response measured is inhibition of disease progression, inhibition of tumor growth, reduction of primary and/or secondary tumor(s), relief of tumor-related symptoms, improvement in quality of life, delayed appearance of primary and/or secondary tumors, slowed development of primary and/or secondary tumors, decreased occurrence of primary and/or secondary tumors, slowed or decreased severity of secondary effects of disease, arrested tumor growth or regression of tumor.

[00345] In other embodiments, provided herein are kits comprising a TOR kinase inhibitor and means for measuring the amount of inhibition of phosphorylation of S6RP, 4E-BP1 and/or AKT in a patient. In certain embodiments, the kits comprise means for measuring inhibition of phosphorylation of S6RP, 4E-BP1 and/or AKT in circulating blood or tumor cells and/or skin biopsies or tumor biopsies/aspirates of a patient. In certain embodiments, provided herein are kits comprising a TOR kinase inhibitor and means for measuring the amount of inhibition of phosphorylation as assessed by comparison of the amount of phospho- S6RP, 4E-BP1 and/or AKT before, during and/or after administration of the TOR kinase inhibitor. In certain embodiments, the patient has ETS overexpressing castration-resistant prostate cancer.

[00346] In other embodiments, provided herein are kits comprising a TOR kinase inhibitor and means for measuring the amount of inhibition of DNA-dependent protein kinase (DNA-PK) activity in a patient. In certain embodiments, the kits comprise means for measuring the amount of inhibition of DNA-dependent protein kinase (DNA-PK) activity in a skin sample and/or a tumor biopsy/aspirate of a patient. In one embodiment, the kits comprise a means for measuring the amount of pDNA-PK S2056 in a skin sample and/or a

tumor biopsy/aspirate of a patient. In one embodiment, the skin sample is irradiated by UV light. In certain embodiments, provided herein are kits comprising a TOR kinase inhibitor and means for measuring the amount of inhibition of DNA-dependent protein kinase (DNA-PK) activity before, during and/or after administration of the TOR kinase inhibitor. In certain embodiments, provided herein are kits comprising a TOR kinase inhibitor and means for measuring the amount of phosphorylated DNA-PK S2056 before, during and/or after administration of the TOR kinase inhibitor. In certain embodiments, the patient has ETS overexpressing castration-resistant prostate cancer.

[00347] In certain embodiments, the kits provided herein comprise an amount of a TOR kinase inhibitor effective for treating or preventing ETS overexpressing castration-resistant prostate cancer. In certain embodiments, the kits provided herein comprise a TOR kinase inhibitor having molecular formula C₁₆H₁₆N₈O. In certain embodiments, the kits provided herein comprise Compound 1.

[00348] In certain embodiments, the kits provided herein further comprise instructions for use, such as for administering a TOR kinase inhibitor and/or monitoring patient response to administration of a TOR kinase inhibitor.

5. EXAMPLES

5.1 BIOLOGICAL EXAMPLES

5.1.1 Biochemical assays

[00349] **mTOR HTR-FRET Assay.** The following is an example of an assay that can be used to determine the TOR kinase inhibitory activity of a test compound. TOR kinase inhibitors were dissolved in DMSO and prepared as 10 mM stocks and diluted appropriately for the experiments. Reagents were prepared as follows:

[00350] “Simple TOR buffer” (used to dilute high glycerol TOR fraction): 10 mM Tris pH 7.4, 100 mM NaCl, 0.1% Tween-20TM, 1 mM DTT. Invitrogen mTOR (cat#PV4753) was diluted in this buffer to an assay concentration of 0.200 µg/mL.

[00351] ATP/Substrate solution: 0.075 mM ATP, 12.5 mM MnCl₂, 50 mM Hepes, pH 7.4, 50 mM β-GOP, 250 nM Microcystin LR, 0.25 mM EDTA, 5 mM DTT, and 3.5 µg/mL GST-p70S6.

[00352] Detection reagent solution: 50 mM HEPES, pH 7.4, 0.01% Triton X-100, 0.01% BSA, 0.1 mM EDTA, 12.7 µg/mL Cy5-αGST Amersham (Cat#PA92002V), 9 ng/mL α-phospho p70S6 (Thr389) (Cell Signaling Mouse Monoclonal #9206L), 627 ng/mL α-mouse Lance EuTM (Perkin Elmer Cat#AD0077).

[00353] To 20 µL of the Simple mTor buffer is added 0.5 µL of test compound in DMSO. To initiate the reaction 5 µL of ATP/Substrate solution was added to 20 µL of the Simple TOR buffer solution (control) and to the compound solution prepared above. The assay was stopped after 60 min by adding 5 µL of a 60 mM EDTA solution; 10 µL of detection reagent solution was then added and the mixture was allowed to sit for at least 2 hours before reading on a Perkin-Elmer EnvisionTM Microplate Reader set to detect LANCE Eu TR-FRET (excitation at 320 nm and emission at 495/520 nm).

[00354] TOR kinase inhibitors were tested in the mTor HTR-FRET assay and were found to have activity therein, with certain compounds having an IC₅₀ below 10 µM in the assay, with some compounds having an IC₅₀ between and 0.005 nM and 250 nM, others having an IC₅₀ between and 250 nM and 500 nM, others having an IC₅₀ between 500 nM and 1 µM, and others having an IC₅₀ between 1 µM and 10 µM.

[00355] DNA-PK assay. DNA-PK assays were performed using the procedures supplied in the PromegaTM DNA-PK assay kit (catalog # V7870). DNA-PK enzyme was purchased from Promega (Promega cat#V5811).

[00356] Selected TOR kinase inhibitors as described herein have, or are expected to have, an IC₅₀ below 10 µM in this assay, with some TOR kinase inhibitors as described herein having an IC₅₀ below 1 µM, and others having an IC₅₀ below 0.10 µM.

5.1.2 Cell based assays

[00357] Clonogenic Assay: Growth Inhibition Assay for ETS Positive and ETS Negative Lines.

[00358] In general, cells (LNCaP, VCaP, DU145 or PC3 prostate cancer cell lines, or PC3-ERG, an isogenic model of ETS on PC3 background) were treated with compound (for example, Compound 1 (a TOR kinase inhibitor set forth herein having molecular formula C₁₆H₁₆N₈O), negative control DMSO, or Olaparib) at a range of concentrations (for example, 10, 3, 1, 0.3 0.1, 0.03 and/or 0.01 μ M) for a time period of for example, 4 h. Then, cells were washed three times with PBS, trypsinized, and replated onto 6 well plates at desired cell densities. Every 3 days the media was replaced with fresh media containing compound. After typically 10-18 days, the cells were fixed and surviving colonies were stained with 0.1% crystal violet and counted by Image QuantTM (BD biosciences).

[00359] In FIG. 1A-C, the effect of compound treatment on Caspase activity is shown. As can be seen, ETS overexpression sensitizes PC3 cells to inhibition by Compound 1.

[00360] Caspase Analysis: Apoptosis Assay for ETS Positive and ETS Negative Lines.

[00361] Increasing concentrations of Compound 1 were spotted via an acoustic dispenser (EDC ATS-100) into an empty 384-well plate in a 10-point serial dilution fashion (3-fold dilution) in duplicate within the plate. Cells were then directly seeded at desired densities to the Compound 1-spotted 384-well plates. Cells were cultured for 48 hours at 37 °C/5% CO₂ and were assessed via Caspase 3/7-Glo (Promega Corporation, Madison, WI) and read for luminescence. In FIG. 2, Caspase activity of single treatment with Compound 1 in ETS-positive (VCaP and LNCaP) and ETS-negative (PC3 and DU145) prostate cancer cell lines is shown and shows the increased apoptosis in ETS positive cell lines.

[00362] Cell Invasion Assay for ETS Positive and ETS Negative Lines.

[00363] Cells (LNCaP, VCaP, DU145 or PC3 prostate cancer cell lines) were treated with Compound 1 (DMSO, 10 μ M Compound 1 or 10 μ M Olaparib) for 48 hours. Then

cells were trypsinized and re-seeded onto BD biocoat matrigel invasion chamber insert with 8.0 μ M pores of a 24-well culture plate in serum free media. Cells were then attracted to the lower chamber by the addition of complete media as a chemoattractant. After 48 hours incubation at 37 °C/5% CO₂, the non-invading cells and EC matrix were gently removed with a cotton swab. Invasive cells, which were located on the lower side of the membrane, were stained with crystal violet, air dried and photographed (FIG. 3). As can be seen, treatment with Compound 1 resulted in reduced cell invasion.

[00364] DNA damage assays: COMET assay.

[00365] Generally, prostate cancer cell lines, for example, VCAP and LNCAP, are seeded at for example, 250,000 cells/mL, in a 6-well plate 24 hours prior to treatment with Compound 1 or vehicle control. After for example 48 hours, cells are trypsinized, harvested by centrifugation and re-suspended in PBS. Cell counts are normalized to 1 x 10⁵ cells/mL. Suspended cells (25 μ L) are mixed with 250 μ L 1.0% ultrapure low melting point agarose (Invitrogen) made in 1x Tris-Borate buffer. The agarose-cell mixture is dropped onto slides allowed to solidify at 4 °C in the dark for 20 minutes before immersion in COMET™ assay lysis solution (Trevigen, Gaithersburg, MD) at 4 °C in the dark for 45 minutes. Excess buffer is removed and slides are submerged in freshly prepared neutral solution (Tris Base 60.57 g, sodium acetate 204.12 g, dissolved in 450 ml of dH₂O, adjusted to pH = 9.0 with glacial acetic acid.) at room temperature in the dark for 40 minutes. Slides are washed twice by immersing in 1 X TBE buffer prior to neutral electrophoresis at 20 volts for 60 minutes. Slides are fixed in 70% ethanol for 5 minutes. Following air drying of the agarose, slides are stained with SYBR Green dye (Invitrogen) and images are collected with a 10x and 40x objective lens. COMET tail moments are assessed using COMETscore.v1.5 image processing software as described by the manufacturer with greater than 100 cells analyzed in triplicate experiments. Data (see FIG. 4) shows that Compound 1 potentiates DNA damage in ETS positive cells.

5.1.3 In vivo Assays:**[00366] Castration-Resistant LNCaP (LNCaP-HR) Prostate Cancer *In Vivo* Xenograft Model**

[00367] A xenograft study was conducted with castration resistant LNCaP-HR tumor-bearing mice. Castrated male SCID mice were inoculated subcutaneously with LNCaP-HR cells in the flank region above the right hind leg. Following inoculation of animals, the tumors were allowed to grow to about 325 mm³ prior to randomization. On Day 26 following tumor cell inoculation, the mice bearing LNCaP-HR tumors ranging between 98 and 530 mm³ were pooled together and randomized into various treatment groups.

Compound 1 was formulated in 0.5% CMC and 0.25% Tween 80 in water (as a suspension). The animals were orally administered vehicle (CMC-Tween) or Compound 1 once daily (QD) for up to 15 days. Doses of Compound 1 ranged between 1 and 5 mg/kg. The positive control MDV-3100 (50 mg/kg, Q4D) was administered via oral route. MDV-3100 was formulated in 1% CMC, 0.1% Tween 80 and 5% dimethyl sulfoxide (DMSO) in water (as a suspension). Tumors were measured twice a week using calipers and tumor volumes were calculated using the formula of W² x L / 2 (wherein "W" is tumor width and "L" is tumor length). Statistical analysis was performed using a one-way analysis of variance (ANOVA) followed by Dunnett's post-hoc comparison with the vehicle-treated control group. Compound 1 significantly inhibited LNCaP-HR prostate cancer tumor growth (see FIG. 5).

5.1.4 Clinical Study**[00368] Phase 1A/1B, Multi-Center, Open-Label, Dose Finding Study to Assess the Safety, Tolerability, Pharmacokinetics and Preliminary Efficacy of Compound 1 Administered Orally to Subjects with ETS Overexpressing Castration-Resistant Prostate Cancer**

[00369] Compound 1 will be administered orally to subjects with ETS overexpressing castration-resistant prostate cancer. The safety and tolerability of Compound 1 in humans, as well as the efficacy, will be evaluated in this study. The study will be conducted in two

parts: dose escalation (Part A) and dose expansion (Part B). Subjects will be enrolled sequentially in Part A. Enrollment in Part B will be stratified by tumor type.

[00370] Compound 1 will be available in three strengths (0.25 mg, 1.0 mg and 5.0 mg) presented in gelatin capsules containing only the active pharmaceutical ingredient. The capsules will be packaged in high density polyethylene (HDPE) bottles, fitted with induction seals and child-resistant polypropylene closures. Research pharmacists will repackage and dispense appropriately for each subject.

[00371] Between 30 and 60 subjects will be enrolled in Part A, designed to establish initial toxicity.

[00372] Part B will consist of approximately 100 subjects with prespecified types of tumors, including E-twenty six (ETS) overexpressing castration-resistant prostate cancer, to further assess the safety profile of Compound 1 and provide efficacy information. Tumor response rate will be assessed by tumor type and dose level. The Part B population will be defined by the efficacy seen during Part A and by data from ongoing preclinical studies.

[00373] The overall study design will be comprised of a Screening Period (Day -28 to Day 1), a Treatment and Evaluation Period (28-day QD (and/or BID) cycles until tumor progression, unacceptable toxicity or subject/physician decision to discontinue administration of Compound 1) and an End of Treatment and Follow-up Period (end of treatment procedures within 21 days of last dose; follow-up for 28 days after last dose for final safety assessment).

[00374] Subjects will start Compound 1 QD or BID dosing (or other suitable regimen) on Cycle 1 Day 1 and receive daily treatment in 28-day cycles. Compound 1 may be discontinued when there is evidence of tumor progression, but subjects can continue to receive study drug as long as the Investigator considers they are deriving benefit. Compound 1 administration will be discontinued when there is unacceptable toxicity, or the subject decides to withdraw from the study.

[00375] Compound 1 will be administered orally either once or twice daily (or other suitable dosing regimen) with no rest period between cycles. Each QD dose will be taken in

the morning with at least 200 mL of water, with the subject having fasted overnight (minimum of 6 hours). Food intake will be delayed until at least 90 minutes after dosing on the days Compound 1 is taken at home. On clinic visit days, the morning Compound 1 dose will be administered in the clinic after any predose tests have been completed. Food may be taken after all fasting tests have been completed but in no case earlier than 90 minutes after dosing (3 hours after dosing on Day 15). For subjects receiving Compound 1 QD where troublesome related GI symptoms, fatigue or other symptoms persist beyond the end of Cycle 1, dosing may be moved to later in the day providing the subject can maintain a 3-hour separation between Compound 1 administration and the last intake of food and a 90-minute delay before ingesting further food. Compound 1 may be taken up to 12 hours late if dosing has been delayed on a single day; otherwise that dose should be omitted.

[00376] Compound 1 will be administered initially as a QD regimen.

[00377] Doses will be administered in an escalating manner following satisfactory review of safety data from the lower doses. There will be a minimum of 28 days after the first dose has been administered to the last subject between dose escalations. Within each cohort, enrollment will be staggered so that there is a minimum of 24 hours between Cycle 1 Day 1 for each subject in order to evaluate initial toxicity.

[00378] Each cycle of Compound 1 lasts 28 days and there is no rest period between cycles. Subjects may continue to receive Compound 1 for as long as they derive benefit from treatment, as judged by the Investigator. Compound 1 administration will be discontinued when there is evidence of disease progression, unacceptable toxicity or when either the subject or Investigator decides to discontinue it.

[00379] In Part A, cohorts of subjects will initially receive QD ascending doses of Compound 1 to measure PK and to identify the MTD. In Part A, 0.5 mg QD is the Compound 2 starting dose. A modified accelerated titration design (Simon, R., Freidlin, B., Rubinstein, L., et al. Accelerated titration designs for Phase I clinical trials in oncology, *J Nat Canc Institute* 1997;89, (15): 1138-1147) will be used to establish initial toxicity. During the accelerated phase, initial cohorts of one subject will be given Compound 1 at

dose increments of 100% until the first instance of first-Cycle grade 2 or higher toxicity suspected to be drug-related, at which point the accelerated phase will stop and this particular cohort will be expanded to a total of 6 subjects. Subsequently, a standard escalation dosing schedule with approximately 50% dose increments and 6 subjects per cohort will be initiated in order to establish the NTD and MTD. Smaller increments and additional subjects within a dose cohort may also be evaluated, if necessary, based on toxicity, PK/PD results or tumor biopsy findings.

[00380] Based on interim PK and PD results from initial dose cohorts, a twice-daily (BID) dosing regimen will also be evaluated in part A. This will be initiated in cohorts of 6 subjects at or below a total daily dose level already shown to be tolerable, but divided into two equal doses administered approximately 12 hours apart. Thereafter, dose escalation for QD and BID dosing cohorts may occur independently. Intermittent dosing schedules of comparable or lower dose intensity than continuous daily dosing may also be considered for evaluation. .

[00381] A dose will be considered to be non-tolerated if 2 or more out of 6 evaluable subjects in a dose cohort experience DLT during Cycle 1. When a NTD is defined, dose escalation will be stopped. The MTD will be defined as the last dose tested below the NTD with 0 or 1 out of 6 evaluable subjects experiencing DLT during Cycle 1. An intermediate dose (*i.e.*, one between the NTD and the last dose level before the NTD) or additional subjects within any dose cohort may be required to more precisely determine the MTD more precisely, as may alternate regimens if emerging PK-PD results suggest these may be appropriate.

[00382] In Part B, subjects may start Compound 1 on a QD or BID regimen at the MTD and/or lower dose levels based on safety, PK and PD data from Part A. In Part B, approximately 100 subjects will be evaluated for safety and antitumor activity after every two cycles of therapy.

[00383] All subjects who receive at least one dose of Compound 1 will be evaluable for safety. In Part A, a subject evaluable for dose-limiting toxicity (DLT) is defined as one who, in the first 28 days after Cycle 1 dosing began, either (a) received at least 21 of the

planned 28 doses of Compound 1 at the cohort-specified dose and has sufficient data for safety evaluation by the SRC, or (b) experienced study drug-related DLT. Non-evaluable subjects will be replaced in the dosing cohort. In Part B, an efficacy evaluable subject for tumor response is defined as one who received at least one cycle of Compound 1, and have baseline and at least one post-baseline efficacy assessment.

[00384] In Parts A and B, dose reductions are permitted in any cycle, including Cycle 1. Dose reductions that occur in Cycle 1 during Part A will constitute DLT, but subjects will be allowed to continue on study drug at the reduced dose. National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) Version 4, 2009 will be used to grade AEs.

[00385] When a dose reduction is indicated, the next lower dose level will be on a QD or BID schedule will be selected. For BID dose reductions below the starting dose of 10 mg BID, 8 mg BID and 4 mg BID will be selected. Two dose reductions are allowed. Additional PK evaluations may be conducted at modified dose level(s) in order to characterize intrasubject PK profiles with alternate doses.

[00386] In Part A, intrasubject dose escalation beyond the dose initially assigned to a subject is not permitted in Cycle 1. Those continuing to take Compound 1 beyond Cycle 1 may, following approval by the SRC, have the dose level increased providing the alternative dose level has been shown to be well tolerated by at least one cohort of other subjects in this study. In these instances, additional PK evaluation at the higher dose level may be conducted. In Part B, no dose escalation beyond the MTD is allowed.

[00387] The primary objectives of this Phase 1a/1b study are to determine the safety, tolerability, NTD and MTD of Compound 1 when administered orally to adult subjects and to determine the PK characteristics of oral Compound 1. The secondary objectives are to evaluate the extent of inhibition of phosphorylation of S6RP and/or 4E-BP1 for mTORC1 activity and AKT and/or other relevant biomarkers for mTORC2 activity in blood, skin and/or tumor biopsies/aspirates and to explore the antitumor activity of Compound 1 at selected dose levels/regimens by tumor type. Further secondary objectives are to evaluate

the inhibition of DNA-PK activity in skin samples irradiated by UV light and/or tumor biopsies/aspirates using pDNA-PK S2056 and other relevant biomarkers for DNA damage pathways before and during Compound 1 treatment.

[00388] In the following, statistical analyses will be performed by study phase, dose level, dosing regimen and tumor cohort as needed or applicable.

[00389] The study population definitions are as follows: (a) Intent-to-Treat (ITT) Population – All subjects who take at least one dose of Compound 1; (b) Safety Population – All subjects who take at least one dose of Compound 1, which is the same as ITT population for this study; (c) Efficacy Evaluable (EE) Population – All ITT subjects who meet eligibility criteria, complete at least one cycle of Compound 1, and have baseline and at least one valid post-baseline efficacy assessment.

[00390] Subject enrollment will be curtailed when up to 20 evaluable subjects have been enrolled in each tumor type and dose level/regimen. In Part B as a whole, sample sizes are not based on statistical calculation but rather on clinical empirical and practical considerations traditionally used for Phase 1 studies of this kind.

[00391] All efficacy evaluable subjects in the Part B portion will be included for efficacy analysis. Efficacy will be analyzed by each tumor type once all subjects have withdrawn from the study or completed 6 cycles. Two-sided ninety-five percent confidence intervals of the response rate will be provided by tumor type. A case-by-case description of all subjects who exhibited a complete or partial response during the Part A segment will be provided. A descriptive analysis of other evidence of anti-tumor activity will be provided based on clinical, radiographic, and biologic assessments of efficacy.

[00392] Subjects will be evaluated for efficacy during even-numbered cycles. The primary efficacy variable is response rate. Tumor response will be based on the Prostate-Specific Antigen Working Group 2 (PSAWG2) Criteria for prostate cancer. Other supplementary efficacy variables, including CTC assessments, will be summarized using frequency tabulations for categorical variables or descriptive statistics for continuous variables.

[00393] For both the dose escalation and dose expansion parts of this protocol, inclusion criteria are: (a) Understand and voluntarily sign an informed consent document before any study-related assessments/procedures are conducted; (b) Men and women, 18 years or older, with histological or cytological confirmation of ETS overexpressing castration-resistant prostate cancer, including those who have progressed on (or not been able to tolerate) standard anticancer therapy or for whom no other conventional therapy exists; (c) Consent to screening tumor biopsy (Part A optional; Part B mandatory except as specified for individual tumor types below); (d) ECOG PS of 0 or 1; (e) The following laboratory values: (1) Absolute neutrophil count (ANC) $\geq 1.5 \times 10^9/L$; (2) Hemoglobin (Hgb) $\geq 9 \text{ g/dL}$; (3) Platelets (plt) $\geq 100 \times 10^9/L$; (4) Potassium within normal range, or correctable with supplements; (5) AST/SGOT and ALT/SGPT $\leq 2.5 \times$ Upper Limit of Normal (ULN) or $\leq 5.0 \times$ ULN if liver tumor is present; (6) Serum total bilirubin $\leq 1.5 \times$ ULN; (7) Serum creatinine $\leq 1.5 \times$ ULN, or 24-hr clearance $\geq 50 \text{ mL/min}$; and (8) Negative serum or urine pregnancy test within 72 hrs before starting study treatment in females of childbearing potential; and (f) Able to adhere to the study visit schedule and other protocol requirements.

[00394] For the dose expansion part (Part B) of this protocol, inclusion criteria are: (a) Subject consent to retrieve formalin-fixed, paraffin-embedded (FFPE) archival tumor tissue, either in tumor blocks or sectioned/mounted specimens; and (b) Histologically-confirmed ETS overexpressing castration-resistant prostate cancer (metastatic adenocarcinoma of the prostate with measurable or unmeasurable disease; Evidence of TMPRSS2-ERG or other ETS gene fusion or ETS overexpression; unresponsive or refractory to hormone therapy with disease progression as determined by PSAWG2 criteria; serum testosterone $< 50 \text{ ng/dL}$; must have had orchiectomy or is currently receiving an LHRH agonist/antagonist; hormonal therapies (other than below), drugs known to decrease PSA (e.g., megestrol acetate, Saw Palmetto) and systemic corticosteroid discontinued at least 4 weeks prior to starting Compound 1; finasteride, bicalutamide and nilutamide discontinued at least 6 weeks prior to Compound 1; bisphosphonates or denosumab are

allowed in stable doses but therapy may not be initiated within 4 weeks prior to starting Compound 1; and optional paired tumor biopsies if tumor is accessible).

[00395] For both the dose escalation and dose expansion parts of this protocol, exclusion criteria are: (a) Symptomatic central nervous system metastases; (b) Known acute or chronic pancreatitis; (c) Any peripheral neuropathy \geq NCI CTCAE grade 2; (d) Persistent diarrhea or malabsorption \geq NCI CTCAE grade 2, despite medical management. Impaired ability to swallow; (e) Impaired cardiac function or clinically significant cardiac diseases; (f) Diabetes mellitus on active treatment; (g) Other concurrent severe and/or uncontrolled concomitant medical conditions (e.g. active or uncontrolled infection) that could cause unacceptable safety risks or compromise compliance with the protocol; (h) Prior systemic cancer-directed treatments or investigational modalities \leq 5 half lives or 4 weeks, whichever is shorter, prior to starting study drug or who have not recovered from side effects of such therapy; (i) Major surgery \leq 2 weeks prior to starting study drug or who have not recovered from side effects of such therapy; (j) Pregnancy or breast feeding; (k) Adults of reproductive potential not employing two forms of birth control; (l) Known HIV infection; (m) Known chronic hepatitis B or C virus (HBV/HCV) infection, unless this is comorbidity in subjects with HCC; (n) Any significant medical condition, laboratory abnormality, or psychiatric illness, including the inability to swallow capsules, that would prevent subjects from participating in the study; (o) Any condition including the presence of laboratory abnormalities, which places subjects at unacceptable risk if they were to participate in the study; (p) Any condition that confounds the ability to interpret study data; or (q) Concurrent active second malignancy for which the subject is receiving therapy, excluding non-melanomatous skin cancer or carcinoma in situ of the cervix.

[00396] For the dose expansion part (Part B) of this protocol, exclusion criteria are: Prior treatment with agents targeting both mTOR complexes (dual TORC1+TORC2 inhibitors) and/or PI3K/AKT pathways. However, prior treatment with isolated TORC1 inhibitors (e.g., rapalogs) is allowed in both parts of this study.

[00397] In some embodiments, patients undergoing the clinical protocol provided herein will show a positive tumor response, such as inhibition of tumor growth or a reduction in tumor size. In certain embodiments, patients undergoing the clinical protocol provided herein will achieve a Response Evaluation Criteria in Solid Tumors (for example, RECIST 1.1) of complete response, partial response or stable disease after administration of an effective amount of Compound 1. In certain embodiments, patients undergoing the clinical protocol provided herein will show increased survival without tumor progression. In some embodiments, patients undergoing the clinical protocol provided herein will show inhibition of disease progression, inhibition of tumor growth, reduction of primary tumor, relief of tumor-related symptoms, inhibition of tumor secreted factors (including tumor secreted hormones, such as those that contribute to carcinoid syndrome), delayed appearance of primary or secondary tumors, slowed development of primary or secondary tumors, decreased occurrence of primary or secondary tumors, slowed or decreased severity of secondary effects of disease, arrested tumor growth and regression of tumors, increased Time To Progression (TTP), increased Progression Free Survival (PFS), and/or increased Overall Survival (OS), among others.

[00398] The embodiments disclosed herein are not to be limited in scope by the specific embodiments disclosed in the examples which are intended as illustrations of a few aspects of the disclosed embodiments and any embodiments that are functionally equivalent are encompassed by the present disclosure. Indeed, various modifications of the embodiments disclosed herein are in addition to those shown and described herein will become apparent to those skilled in the art and are intended to fall within the scope of the appended claims.

CLAIMS:

1. Use of 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof in the manufacture of a medicament for treating E-twenty six (ETS) overexpressing castration-resistant prostate cancer.

2. The use of claim 1, wherein the E-twenty six (ETS) overexpressing castration-resistant prostate cancer is that in which the PI3K/mTOR pathway is activated.

3. The use of claim 2, wherein the E-twenty six (ETS) overexpressing castration-resistant prostate cancer is that in which the PI3K/mTOR pathway is activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

4. The use of any one of claims 1 to 3, wherein said medicament is suitable to be administered in an amount of about 0.5 mg/day to about 128 mg/day of 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof.

5. The use of claim 4, wherein said medicament is suitable to be administered in an amount of 0.5 mg/day, 1 mg/day, 2 mg/day, 4 mg/day, 8 mg/day, 16 mg/day, 20 mg/day, 30 mg/day, 45 mg/day, 60 mg/day, 90 mg/day, 120 mg/day or 128 mg/day of 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof.

6. The use of any one of claims 1 to 3, wherein said medicament is suitable to be administered in a unit dosage form comprising 0.25 mg, 1.0 mg, 5.0 mg, 7.5 mg, or 10 mg of 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof.

7. Use of 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer

or tautomer thereof in the manufacture of a medicament for improving the Prostate-Specific Antigen Working Group 2 (PSAWG2) Criteria for prostate cancer of a patient.

8. The use of claim 7, wherein the E-twenty six (ETS) overexpressing castration-resistant prostate cancer is that in which the PI3K/mTOR pathway is activated.

9. The use of claim 8, wherein the E-twenty six (ETS) overexpressing castration-resistant prostate cancer is that in which the PI3K/mTOR pathway is activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

10. Use of 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof for treating E-twenty six (ETS) overexpressing castration-resistant prostate cancer.

11. The use of claim 10, wherein the E-twenty six (ETS) overexpressing castration-resistant prostate cancer is that in which the PI3K/mTOR pathway is activated.

12. The use of claim 11, wherein the E-twenty six (ETS) overexpressing castration-resistant prostate cancer is that in which the PI3K/mTOR pathway is activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

13. The use of any one of claims 10 to 12, wherein 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof is for administration in an amount of about 0.5 mg/day to about 128 mg/day.

14. The use of claim 13, wherein 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof is for administration in an amount of 0.5 mg/day, 1 mg/day, 2 mg/day, 4 mg/day, 8 mg/day, 16 mg/day, 20 mg/day, 30 mg/day, 45 mg/day, 60 mg/day, 90 mg/day, 120 mg/day or 128 mg/day.

15. The use of any one of claims 10 to 12, wherein 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof is for administration in a unit dosage form comprising 0.25 mg, 1.0 mg, 5.0 mg, 7.5 mg, or 10 mg.

16. Use of 1-ethyl-7-(2-methyl-6-(1H-1,2,4-triazol-3-yl)pyridin-3-yl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one or a pharmaceutically acceptable salt, stereoisomer or tautomer thereof for improving the Prostate-Specific Antigen Working Group 2 (PSAWG2) Criteria for prostate cancer of a patient.

17. The use of claim 16, wherein the E-twenty six (ETS) overexpressing castration-resistant prostate cancer is that in which the PI3K/mTOR pathway is activated.

18. The use of claim 17, wherein the E-twenty six (ETS) overexpressing castration-resistant prostate cancer is that in which the PI3K/mTOR pathway is activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

1/6




FIG. 1A

2/6




FIG. 1B




FIG. 1C

3/6




FIG. 2




FIG. 3 (n=2)

5/6




FIG. 4

6/6




FIG. 5