



(86) Date de dépôt PCT/PCT Filing Date: 2011/06/23  
(87) Date publication PCT/PCT Publication Date: 2012/04/26  
(85) Entrée phase nationale/National Entry: 2013/04/10  
(86) N° demande PCT/PCT Application No.: JP 2011/064487  
(87) N° publication PCT/PCT Publication No.: 2012/053255  
(30) Priorités/Priorities: 2010/10/20 (JP2010-235203);  
2010/12/20 (JP2010-282938); 2010/12/20 (US61/424885)

(51) Cl.Int./Int.Cl. *G01N 33/50* (2006.01),  
*G01N 33/15* (2006.01), *G01N 33/68* (2006.01),  
*G01N 33/92* (2006.01)  
(71) Demandeur/Applicant:  
EISAI R&D MANAGEMENT CO., LTD., JP  
(72) Inventeurs/Inventors:  
SATO, YOSHIAKI, JP;  
AOSHIMA, KEN, JP;  
BERNIE, FRANCOIS, JP  
(74) Agent: TORYS LLP

(54) Titre : BIOMARQUEUR DE LA MALADIE D'ALZHEIMER OU D'UN TROUBLE COGNITIF LEGER  
(54) Title: BIOMARKER FOR ALZHEIMER'S DISEASE OR MILD COGNITIVE IMPAIRMENT

(57) **Abrégé/Abstract:**

Desmosterol alone, the combination of desmosterol and amyloid beta, the combination of desmosterol and gelsolin, and the combination of desmosterol, amyloid beta, and gelsolin can be used as in-blood biomarkers for Alzheimer's disease or mild cognitive impairment. Provided are: a method using these biomarkers for evaluating the effect of candidate substances for treatment agents for Alzheimer's disease or mild cognitive impairment; a method for assisting in the diagnosis of Alzheimer's disease or mild cognitive impairment; and a method for diagnosing Alzheimer's disease or mild cognitive impairment.



## (12) 特許協力条約に基づいて公開された国際出願

(19) 世界知的所有権機関  
国際事務局(43) 国際公開日  
2012年4月26日(26.04.2012)(10) 国際公開番号  
**WO 2012/053255 A1**

- (51) 国際特許分類:  
G01N 33/50 (2006.01) G01N 33/68 (2006.01)  
G01N 33/15 (2006.01) G01N 33/92 (2006.01)
- (21) 国際出願番号: PCT/JP2011/064487
- (22) 国際出願日: 2011年6月23日(23.06.2011)
- (25) 国際出願の言語: 日本語
- (26) 国際公開の言語: 日本語
- (30) 優先権データ:  
特願 2010-235203 2010年10月20日(20.10.2010) JP  
61/424885 2010年12月20日(20.12.2010) US  
特願 2010-282938 2010年12月20日(20.12.2010) JP
- (71) 出願人 (米国を除く全ての指定国について):  
エーザイ・アール・アンド・ディー・マネジメント株式会社 (Eisai R&D Management Co., Ltd.)  
[JP/JP]; 〒1128088 東京都文京区小石川四丁目6番10号 Tokyo (JP).
- (72) 発明者; および
- (75) 発明者/出願人 (米国についてのみ): 佐藤 義明 (SATO Yoshiaki) [JP/JP]; 〒3002635 茨城県つくば市東光台5丁目1番地3 エーザイ株式会社筑波研究所内 Ibaraki (JP). 青島 健 (AOSHIMA Ken) [JP/JP]; 〒3002635 茨城県つくば市東光台5丁目1番地3 エーザイ株式会社筑波研究所内 Ibaraki (JP). ベルニエ フランソワ (BERNIE Francois) [CA/JP]; 〒3002635 茨城県つくば市東光台5丁目1番地3 エーザイ株式会社筑波研究所内 Ibaraki (JP).
- (74) 代理人: 長谷川 芳樹, 外 (HASEGAWA Yoshiki et al.); 〒1000005 東京都千代田区丸の内二丁目1
- 番1号丸の内 MY PLAZA (明治安田生命ビル) 9階 創英国際特許法律事務所 Tokyo (JP).
- (81) 指定国 (表示のない限り、全ての種類の国内保護が可能): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) 指定国 (表示のない限り、全ての種類の広域保護が可能): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), ユーラシア (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), ヨーロッパ (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- 規則 4.17 に規定する申立て:  
— 不利にならない開示又は新規性喪失の例外に関する申立て (規則 4.17(v))
- 添付公開書類:  
— 国際調査報告 (条約第 21 条(3))

(54) Title: BIOMARKER FOR ALZHEIMER'S DISEASE OR MILD COGNITIVE IMPAIRMENT

(54) 発明の名称: アルツハイマー病又は軽度認知症のバイオマーカー

(57) Abstract: Desmosterol alone, the combination of desmosterol and amyloid beta, the combination of desmosterol and gelsolin, and the combination of desmosterol, amyloid beta, and gelsolin can be used as in-blood biomarkers for Alzheimer's disease or mild cognitive impairment. Provided are: a method using these biomarkers for evaluating the effect of candidate substances for treatment agents for Alzheimer's disease or mild cognitive impairment; a method for assisting in the diagnosis of Alzheimer's disease or mild cognitive impairment; and a method for diagnosing Alzheimer's disease or mild cognitive impairment.

(57) 要約: デスモステロール単独、デスモステロール及びアミロイドベータの組み合わせ、デスモステロール及びゲルゾリンの組み合わせ並びにデスモステロール、アミロイドベータ及びゲルゾリンの組み合わせは、アルツハイマー病又は軽度認知症の血中バイオマーカーとして利用可能である。これらのバイオマーカーを利用した、アルツハイマー病又は軽度認知症治療剤の候補物質の効果を評価する方法、アルツハイマー病又は軽度認知症の診断を補助する方法及びアルツハイマー病又は軽度認知症を診断する方法が提供される。

WO 2012/053255 A1

**DESCRIPTION****Title of Invention**

BIOMARKER FOR ALZHEIMER'S DISEASE OR MILD COGNITIVE IMPAIRMENT

**Technical Field**

5 [0001] The present invention relates to a biomarker for Alzheimer's disease or mild cognitive impairment. More specifically, the present invention relates to a method for evaluating the effect of a candidate for a therapeutic agent for Alzheimer's disease or mild cognitive impairment, a method for aiding the diagnosis of Alzheimer's disease or mild cognitive impairment, and a method for diagnosing Alzheimer's disease or mild cognitive impairment.

**Background Art**

10 [0002] A biomarker is very useful for diagnosing and monitoring the progress of a disease, and is important for screening patients, monitoring side effects, aiding for the selection of an appropriate treatment and discovering a new drug. In Alzheimer's disease, A $\beta$  (amyloid beta) 1-42 and phosphorylated tau protein in the cerebrospinal fluid are currently considered to be the most useful biomarkers (Non Patent Literatures 1 and 2).

**Citation List****Non Patent Literature**

20 [0003] Non Patent Literature 1: Ibach et al., "Cerebrospinal fluid tau and beta-amyloid in Alzheimer patients, disease controls and an age-matched random sample," *Neurobiol. Aging.*, 2006; vol. 27 (no. 9): p.1202-1211.

Non Patent Literature 2: Bouwman et al., "CSF biomarker levels in early and late onset Alzheimer's disease," *Neurobiol. Aging.*, 2009; vol. 30 (no. 12): p.1895-1901.

25 Non Patent Literature 3: Koelsch et al. "Alterations of cholesterol precursor levels in Alzheimer's disease," *Biochim. Biophys. Acta.*, 2010; vol. 1801 (no. 8): p.945-950.

Non Patent Literature 4: Guentert et al., "Plasma Gelsolin is Decreased and Correlates with Rate of Decline in Alzheimer's Disease," *J. Alzheimer's. Dis.*, 2010; vol. 21 (no. 2): p.585-596.

**Summary of Invention****30 Technical Problem**

[0004] However, for the above biomarkers, use of the cerebrospinal fluid is required, and, in particular, collecting the cerebrospinal fluid every time the time-dependent change of Alzheimer pathology is monitored imposes an enormous burden on the patient, and thus a

low invasive biomarker using blood or the like, has been demanded.

#### **Solution to Problem**

5 [0005] Desmosterol is a precursor of cholesterol. In Alzheimer's disease, it is thought that the disorder of cholesterol metabolism has occurred, and documented that the desmosterol level in the cerebrospinal fluid of patients with **Alzheimer's disease** is lower than that of healthy elderly people, while the desmosterol level in the plasma remains unchanged (Non Patent Literature 3). The present inventors found that, inconsistent with this finding, the desmosterol level in plasma of patients with Alzheimer's disease is lower than that of healthy elderly people, and the desmosterol in plasma can be used as a biomarker for Alzheimer's disease. Further, the present inventors found that desmosterol alone enables diagnosis of the mild cognitive impairment, manifesting the mild cognitive dysfunction which is a pre-existing disease of Alzheimer's disease.

10 [0006] Furthermore, the present inventors found that the combination of desmosterol and amyloid beta in blood when used as a biomarker enables more accurate diagnosis of Alzheimer's disease or mild cognitive impairment than when desmosterol alone in blood is used as a biomarker.

15 [0007] Furthermore, the present inventors found that the combination of desmosterol and gelsolin in blood, or the combination of desmosterol, amyloid beta and gelsolin in blood, when used as a biomarker, enables more accurate diagnosis of Alzheimer's disease or mild cognitive impairment than when desmosterol alone in blood is used as a biomarker. It is known that the plasma gelsolin level is low in patients with Alzheimer's disease (Non Patent Literature 4).

20 [0008] Specifically, the present invention provides the following [1] to [28].

[1] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

25 a step of measuring a desmosterol level in blood of a non-human animal;

a step of administering a candidate to the non-human animal;

a step of measuring a desmosterol level in blood of the non-human animal after the candidate is administered; and

30 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the non-human animal after the candidate is administered is higher than the level before the candidate is administered.

[2] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level in blood of a first non-human animal;

a step of administering a candidate to a second non-human animal;

5 a step of measuring a desmosterol level in blood of the second non-human animal after the candidate is administered; and

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the second non-human animal after the candidate is administered is higher than the desmosterol level in blood of the first non-human animal to which the candidate is not administered.

10

[3] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level in a blood sample of a subject before a candidate is administered;

15

a step of measuring a desmosterol level in the blood sample of the subject after the candidate is administered; and

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered.

20

[4] A method for aiding diagnosis of Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level in a blood sample of a subject; and

25

a step of providing an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value.

[5] A method for diagnosing Alzheimer's disease or mild cognitive impairment, the method comprising:

30

a step of measuring a desmosterol level in blood of a subject; and

a step of identifying Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value.

[6] A method for selecting a patient which is likely to benefit from a therapeutic effect

of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level in a blood sample of a subject; and

5 a step of providing an indicator of a patient to be administered with a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value.

[7] A method for predicting a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level in a blood sample of a subject; and

10 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value.

[8] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

15 a step of measuring a desmosterol level and an amyloid beta level in blood of a non-human animal;

a step of administering a candidate to the non-human animal;

a step of measuring a desmosterol level and an amyloid beta level in blood of the non-human animal after the candidate is administered; and

20 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the non-human animal after the candidate is administered is higher than the level before the candidate is administered, and  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the non-human animal after the candidate is administered is higher than that before the candidate is administered.

25

[9] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and an amyloid beta level in blood of a first non-human animal;

30

a step of administering a candidate to a second non-human animal;

a step of measuring a desmosterol level and an amyloid beta level in blood of the second non-human animal after the candidate is administered; and

a step of providing an indicator of being potential for the candidate to be effective

5 in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the second non-human animal after the candidate is administered is higher than the desmosterol level in blood of the first non-human animal to which the candidate is not administered, and  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the second non-human animal after the candidate is administered is higher than  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the first non-human animal to which the candidate is not administered.

[10] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

10 a step of measuring a desmosterol level and an amyloid beta level in a blood sample of a subject before a candidate is administered;

a step of measuring a desmosterol level and an amyloid beta level in the blood sample of the subject after the candidate is administered; and

15 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered, and  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the subject after the candidate is administered is higher than that before the candidate is administered.

20 [11] A method for aiding diagnosis of Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and an amyloid beta level in a blood sample of a subject; and

25 a step of providing an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the subject is lower than a reference value.

[12] A method for diagnosing Alzheimer's disease or mild cognitive impairment, the method comprising:

30 a step of measuring a desmosterol level and an amyloid beta level in blood of a subject; and

a step of identifying Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the subject is lower than a reference value.

[13] A method for selecting a patient which is likely to benefit from a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

5 a step of measuring a desmosterol level and an amyloid beta level in a blood sample of a subject; and

a step of providing an indicator of a patient to be administered with a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value.

10 [14] A method for predicting a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and an amyloid beta level in a blood sample of a subject; and

15 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value.

[15] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

20 a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood of a non-human animal;

a step of administering a candidate to the non-human animal;

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood of the non-human animal after the candidate is administered; and

25 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the non-human animal after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the non-human animal after the candidate is administered is higher than that before the candidate is administered, and the gelsolin level in blood of the non-human animal after the candidate is administered is higher than the gelsolin level in blood before the candidate is administered.

30

[16] A method for evaluating an effect of a candidate for a therapeutic agent to treat

Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood of a first non-human animal;

a step of administering a candidate to a second non-human animal;

5 a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood of the second non-human animal after the candidate is administered; and

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the second non-human animal after the candidate is administered is higher than the desmosterol level in blood of the first non-human animal to which the candidate is not administered,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the second non-human animal after the candidate is administered is higher than  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the first non-human animal to which the candidate is not administered, and the gelsolin level in blood of the second non-human animal after the candidate is administered is higher than the gelsolin level in blood of the first non-human animal to which the candidate is not administered.

[17] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

20 a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in a blood sample of a subject before a candidate is administered,

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in the blood sample of the subject after the candidate is administered,

25 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject after the candidate is administered is higher than the desmosterol level before the candidate is administered,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject after the candidate is administered is higher than that before the candidate is administered, and the gelsolin level in blood of the subject after the candidate is administered is higher than the gelsolin level in blood before the candidate is administered.

30 [18] A method for aiding diagnosis of Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in a blood sample of a subject; and

a step of providing an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

5 [19] A method for diagnosing Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood of a subject; and

10 a step of identifying Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

[20] A method for selecting a patient which is likely to benefit from a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

15 a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in a blood sample of a subject; and

20 a step of providing an indicator of a patient to be administered with a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

[21] A method for predicting a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

25 a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in a blood sample of a subject, and

30 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

[22] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and a gelsolin level in blood of a non-human animal;

a step of administering a candidate to the non-human animal;

5 a step of measuring a desmosterol level and a gelsolin level in blood of the non-human animal after the candidate is administered; and

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the non-human animal after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered, and the gelsolin level in blood of the non-human animal after the candidate is administered is higher than the gelsolin level in blood before the candidate is administered.

[23] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

15 a step of measuring a desmosterol level and a gelsolin level in blood of a first non-human animal;

a step of administering a candidate to a second non-human animal;

a step of measuring a desmosterol level and a gelsolin level in blood of the second non-human animal after the candidate is administered; and

20 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the second non-human animal after the candidate is administered is higher than the desmosterol level in blood of the first non-human animal to which the candidate is not administered, and the gelsolin level in blood of the second non-human animal after the candidate is administered is higher than the gelsolin level in blood of the first non-human animal to which the candidate is not administered.

[24] A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and a gelsolin level in a blood sample of a subject before a candidate is administered;

30 a step of measuring a desmosterol level and a gelsolin level in the blood sample of the subject after the candidate is administered; and

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in

blood of the subject after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered, and the gelsolin level in blood of the subject after the candidate is administered is higher than the gelsolin level in blood before the candidate is administered.

5 [25] A method for aiding diagnosis of Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and a gelsolin level in a blood sample of a subject; and

10 a step of providing an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

[26] A method for diagnosing Alzheimer's disease or mild cognitive impairment, the method comprising:

15 a step of measuring a desmosterol level and a gelsolin level in blood of a subject; and

a step of identifying Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

20 [27] A method for selecting a patient which is likely to benefit from a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and a gelsolin level in a blood sample of a subject; and

25 a step of providing an indicator of a patient to be administered with a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

[28] A method for predicting a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

30 a step of measuring a desmosterol level and a gelsolin level in a blood sample of a subject; and

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in

blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

#### **Advantageous Effects of Invention**

5 [0009] Using a desmosterol level in blood, or the like, as an indicator, it is possible to diagnose accurately the progress of Alzheimer's disease or mild cognitive impairment, it is also possible to evaluate accurately an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, it is further possible to select a patient which is likely to benefit from the therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, and it is furthermore possible to predict the therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment. The desmosterol and the like, of the present invention are the biomarkers in blood, and it is thus possible to diagnose accurately Alzheimer's disease or mild cognitive impairment without collecting the cerebrospinal fluid.

#### **Brief Description of Drawings**

15 [0010] [Figure 1] Figure 1 is a chart showing the Receiver Operating Characteristic Curves of desmosterol,  $A\beta_{x-42}/A\beta_{x-40}$  and the combination thereof in healthy elderly people and patients with Alzheimer's disease.

[Figure 2] Figure 2 is a chart showing the Receiver Operating Characteristic Curves of desmosterol,  $A\beta_{x-42}/A\beta_{x-40}$  and the combination thereof in healthy elderly people and patients with mild cognitive impairment.

20 [Figure 3] Figure 3 is a chart showing the Receiver Operating Characteristic Curves of desmosterol/cholesterol,  $A\beta_{x-42}/A\beta_{x-40}$ , gelsolin and the combination thereof in healthy elderly people and patients with Alzheimer's disease.

[Figure 4] Figure 4 is a chart showing the Receiver Operating Characteristic Curves of desmosterol/cholesterol,  $A\beta_{x-42}/A\beta_{x-40}$ , gelsolin and the combination thereof in healthy elderly people and patients with mild cognitive impairment.

25 [Figure 5] Figure 5 is a chart showing the Receiver Operating Characteristic Curves of desmosterol/cholesterol, gelsolin and the combination thereof in healthy elderly people and patients with Alzheimer's disease.

30 [Figure 6] Figure 6 is a chart showing the Receiver Operating Characteristic Curves of desmosterol/cholesterol, gelsolin and the combination thereof in healthy elderly people and patients with mild cognitive impairment.

[Figure 7] Figure 7 is a chart showing desmosterol levels in healthy elderly people and

patients with mild cognitive impairment. In the chart, the thick bar represents the median value. The fold of change in patients with mild cognitive impairment against healthy elderly people is 0.7368.

[Figure 8] Figure 8 is a chart showing the Receiver Operating Characteristic Curves of desmosterol in healthy elderly people and patients with mild cognitive impairment.

#### **Description of Embodiments**

[0011] The method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention targets a non-human animal or a human. It is preferable that the non-human animal be a mammal. It is preferable that the non-human animal be a model animal with Alzheimer's disease or mild cognitive impairment. Examples of the model animal with Alzheimer's disease or mild cognitive impairment include an animal with an increased production of amyloid beta by the modification of amyloid precursor protein (APP) or presenilin 2. More specifically, an example includes Tg2576, a genetically modified mouse overexpressing APP. It is preferable that the human be a patient with Alzheimer's disease or mild cognitive impairment.

[0012] The effect of a candidate can be evaluated when a desmosterol level in blood of a non-human animal or a human of the same individual is measured before and after the candidate is administered and the levels are compared.

[0013] The desmosterol level in blood used herein can be measured by the liquid chromatography/mass spectrometry (LC/MS). More specifically, it can be measured under the conditions described in Examples. The desmosterol level in blood may be a concentration of desmosterol, or desmosterol/cholesterol, which is the ratio of desmosterol to cholesterol, with a concentration of desmosterol being preferable. Also, the desmosterol level in blood can be measured by the Enzyme-Linked ImmunoSorbent Assay (ELISA) or gas chromatography/mass spectrometry (GC-MS). The desmosterol level in blood is preferably a level in plasma.

[0014] The candidate drug efficacy can be evaluated based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered.

[0015] Further, using non-human animals of different individuals, the effect of a candidate can be evaluated by measuring desmosterol levels in blood of one or a plurality of individuals

in one group without administering the candidate (control group), measuring desmosterol levels in blood of one or a plurality of individuals of the other group after the candidate is administered (candidate-administered group), and comparing the levels of those groups. In this instance, the candidate drug efficacy can be evaluated based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood is higher in the candidate-administered group than the desmosterol level in blood in the control group.

[0016] The method for aiding diagnosis of Alzheimer's disease or mild cognitive impairment of the present invention provides an indicator for diagnosis by measuring a desmosterol level in a blood sample of a subject which is suspected to have Alzheimer's disease or mild cognitive impairment and comparing a reference value therewith. More specifically, the method of the present invention provides an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value. A reference value of desmosterol level in blood can be determined by referring to the values of a patient definitely diagnosed with Alzheimer's disease or mild cognitive impairment and of a healthy person.

[0017] The method for diagnosing Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring a desmosterol level of a subject which is suspected to have Alzheimer's disease or mild cognitive impairment and comparing a reference value therewith. More specifically, when the desmosterol level in blood of the subject is lower than a reference value, the subject is identified as having Alzheimer's disease or mild cognitive impairment. The reference value of desmosterol level in blood is as described above.

[0018] The method for selecting a patient which is likely to benefit from the therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring a desmosterol level in blood of a subject which is diagnosed with Alzheimer's disease or mild cognitive impairment and comparing a reference value therewith. More specifically, the selection can be made based on an indicator of a patient to be administered with the candidate for the therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood is as described above.

[0019] The method for predicting the therapeutic effect of a candidate for a therapeutic

agent to treat Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring a desmosterol level in blood of a subject which is diagnosed with Alzheimer's disease or mild cognitive impairment and comparing a reference value therewith. More specifically, the prediction can be made based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood is as described above.

5 [0020] The candidate for the therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention is not particularly limited, and is preferably a modulator of the cholesterol metabolism pathway or a modulator based on A $\beta$  hypothesis, more preferably a  $\beta$ -secretase inhibitor, a  $\gamma$ -secretase inhibitor, a  $\gamma$ -secretase modulator, anti-A $\beta$  antibody, anti-A $\beta$  oligomer antibody and the like.

10 [0021] In the above methods, a desmosterol level alone in blood may be used as a biomarker, or may be used in combination with other biomarkers. More accurate evaluation or the like, can be carried out when a desmosterol level in blood is combined with other biomarkers. In particular, the combination with an amyloid beta level in blood, the combination with gelsolin level in blood, or the combination with an amyloid beta level and a gelsolin level in blood enables a significantly accurate evaluation or the like, to be carried out, hence preferable. Hereinbelow, embodiments using the combination with these

15

20 biomarkers are described in detail.

[0022] The method, which uses a desmosterol level and amyloid beta level in blood as the biomarkers, is described.

[0023] The method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention targets a non-human animal or a human. Specifics are as described above.

25 [0024] The effect of a candidate can be evaluated when a desmosterol level and an amyloid beta level in blood of a non-human animal or a human of the same individual are measured before and after the candidate is administered and the levels are compared.

[0025] The method for measuring a desmosterol level in blood is as described above.

30 [0026] The amyloid beta level in blood can be measured, for example, by known methods such as ELISA or LC/MS. It is desirable that the amyloid beta be measured in both A $\beta$ x-40 and A $\beta$ x-42. A $\beta$ x-40 means A $\beta$ 1-40 and a peptide having the N-terminus thereof is partially deleted, and A $\beta$ x-42 means A $\beta$ 1-42 and a peptide having the N-terminus thereof is

partially deleted. The amyloid beta level in blood may be a concentration of A $\beta$ x-42 in blood, or may be A $\beta$ x-42/A $\beta$ x-40, which is the ratio of A $\beta$ x-42 to A $\beta$ x-40, with A $\beta$ x-42/A $\beta$ x-40 being preferable. The amyloid beta level in blood is preferably a level in plasma.

5 [0027] The candidate drug efficacy can be evaluated based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered, and A $\beta$ x-42 or A $\beta$ x-42/A $\beta$ x-40 in blood after the candidate is administered is higher than that before the candidate is administered. The A $\beta$ x-42/A $\beta$ x-40 in blood means a ratio of the A $\beta$ x-42 level in blood to the A $\beta$ x-40 level in blood.

10 [0028] Also, using non-human animals of different individuals, the effect of a candidate can be evaluated by measuring desmosterol levels and amyloid beta levels in blood of one or a plurality of individuals in one group without administering the candidate (control group), measuring desmosterol levels and amyloid beta levels in blood of one or a plurality of individuals of the other group after the candidate is administered (candidate-administered group), and comparing the levels of those groups. In this instance, the candidate drug efficacy can be evaluated based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the candidate-administered group is higher than the level of the control group, and A $\beta$ x-42 or A $\beta$ x-42/A $\beta$ x-40 in blood of the candidate-administered group is higher than that of the control group.

15 [0029] The method for aiding diagnosis of Alzheimer's disease or mild cognitive impairment of the present invention provides an indicator for diagnosis by measuring a desmosterol level and an amyloid beta level in a blood sample of a subject which is suspected to have Alzheimer's disease and comparing reference values therewith. More specifically, the method of the present invention provides an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and A $\beta$ x-42 or A $\beta$ x-42/A $\beta$ x-40 in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood and the reference value of A $\beta$ x-42 or A $\beta$ x-42/A $\beta$ x-40 in blood can be determined by referring to the value of a patient definitely diagnosed with Alzheimer's disease or mild cognitive impairment.

[0030] The method for diagnosing Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring a desmosterol level and an amyloid beta level in blood of a subject which is suspected to have Alzheimer's disease or mild cognitive impairment and comparing reference values therewith. More specifically, when  
5 the desmosterol level in blood of the subject is lower than a reference value, and  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value, the subject is identified as having Alzheimer's disease or mild cognitive impairment. The reference value of desmosterol level in blood and the reference value of  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood are as described above.

[0031] The method for selecting a patient which is likely to benefit from the therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring a desmosterol level and an amyloid beta level in blood of a subject which is diagnosed with Alzheimer's disease or mild cognitive impairment and comparing reference values therewith. More specifically,  
15 the selection can be made based on an indicator of a patient to be administered with the candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood and the reference value of  $A\beta_{x-42}$  or  
20  $A\beta_{x-42}/A\beta_{x-40}$  in blood are as described above.

[0032] The method for predicting the therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring a desmosterol level and an amyloid beta level in blood of a subject which is diagnosed with Alzheimer's disease or mild cognitive impairment and  
25 comparing reference values therewith. More specifically, the prediction can be made based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood and the  
30 reference value of  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood are as described above.

[0033] The candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention is not particularly limited, and is preferably as described above.

[0034] The method, which uses a desmosterol level, an amyloid beta level and a gelsolin level in blood as the biomarkers, is described.

[0035] The method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention targets a non-human animal or a human. Specifics are as described above.

[0036] The effect of a candidate can be evaluated when a desmosterol level, an amyloid beta level and a gelsolin level in blood of a non-human animal or a human of the same individual are measured before and after the candidate is administered and the levels are compared.

[0037] The method for measuring a desmosterol level and an amyloid beta level in blood is as described above.

[0038] The gelsolin level in blood can be measured, for example, by known methods such as ELISA. The gelsolin level in blood is preferably a level in plasma.

[0039] The candidate drug efficacy can be evaluated based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered,  $A\beta_x-42$  or  $A\beta_x-42/A\beta_x-40$  in blood after the candidate is administered is higher than that before the candidate is administered, and the gelsolin level in blood after the candidate is administered is higher than the gelsolin level in blood before the candidate is administered.

[0040] Also, using non-human animals of different individuals, the effect of a candidate can be evaluated by measuring desmosterol levels, amyloid beta levels and gelsolin levels in blood of one or a plurality of individuals in one group without administering the candidate (control group), measuring desmosterol levels, amyloid beta levels and gelsolin levels in blood of one or a plurality of individuals of the other group after the candidate is administered (candidate-administered group), and comparing the levels of those groups. In this instance, the candidate drug efficacy can be evaluated based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the candidate-administered group is higher than the level of the control group,  $A\beta_x-42$  or  $A\beta_x-42/A\beta_x-40$  in blood of the candidate-administered group is higher than that of the control group, and the gelsolin level in blood of the candidate-administered group is higher than the level of the control group.

[0041] The method for aiding diagnosis of Alzheimer's disease or mild cognitive

impairment of the present invention provides an indicator for diagnosis by measuring a desmosterol level, an amyloid beta level and a gelsolin level in a blood sample of a subject which is suspected to have Alzheimer's disease or mild cognitive impairment and comparing reference values therewith. More specifically, the method of the present invention provides an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood and the reference value of  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood and the reference value of gelsolin level in blood can be determined by referring to the values of a patient definitely diagnosed with Alzheimer's disease or mild cognitive impairment.

[0042] The method for diagnosing Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring desmosterol in blood, an amyloid beta level and a gelsolin level in blood of a subject which is suspected to have Alzheimer's disease or mild cognitive impairment and comparing reference values therewith. More specifically, when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value, the subject is identified as having Alzheimer's disease or mild cognitive impairment. The reference value of desmosterol level in blood, the reference value of  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood and the reference value of gelsolin level in blood are as described above.

[0043] The method for selecting a patient which is likely to benefit from the therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring desmosterol in blood, an amyloid beta level and a gelsolin level in blood of a subject which is diagnosed with Alzheimer's disease or mild cognitive impairment and comparing reference values therewith. More specifically, the selection can be made based on an indicator of a patient to be administered with the candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood, the reference value of  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood and the reference value of gelsolin level in blood are as described

above.

5 [0044] The method for predicting the therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring desmosterol, an amyloid beta level and a gelsolin level in blood of a subject which is diagnosed with Alzheimer's disease and comparing reference values therewith. More specifically, the prediction can be made based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood, the reference value of  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood and the reference value of gelsolin level in blood are as described above.

10 [0045] The candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention is not particularly limited, and is preferably as described above.

[0046] The method, which uses a desmosterol level and a gelsolin level in blood as the biomarkers, is described.

15 [0047] The method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention targets a non-human animal or a human. Specifics are as described above.

[0048] The effect of a candidate can be evaluated when a desmosterol level and a gelsolin level in blood of a non-human animal or a human of the same individual are measured before and after the candidate is administered and the levels are compared.

20 [0049] The method for measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood is as described above.

25 [0050] The candidate drug efficacy can be evaluated based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered, and the gelsolin level in blood after the candidate is administered is higher than the gelsolin level in blood before the candidate is administered.

30 [0051] Also, using non-human animals of different individuals, the effect of a candidate

can be evaluated by measuring desmosterol levels and gelsolin levels in blood of one or a plurality of individuals in one group without administering the candidate (control group), measuring desmosterol levels and gelsolin levels in blood of one or a plurality of individuals of the other group after the candidate is administered (candidate-administered group), and comparing the levels of those groups. In this instance, the candidate drug efficacy can be evaluated based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the candidate-administered group is higher than the level of the control group, and the gelsolin level in blood of the candidate-administered group is higher than the level of the control group.

[0052] The method for aiding diagnosis of Alzheimer's disease or mild cognitive impairment of the present invention provides an indicator for diagnosis by measuring a desmosterol level and a gelsolin level in a blood sample of a subject which is suspected to have Alzheimer's disease or mild cognitive impairment and comparing reference values therewith. More specifically, the method of the present invention provides an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood and the reference value of gelsolin level in blood can be determined by referring to the values of a patient definitely diagnosed with Alzheimer's disease or mild cognitive impairment.

[0053] The method for diagnosing Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring a desmosterol level in blood and a gelsolin level in blood of a subject which is suspected to have Alzheimer's disease or mild cognitive impairment and comparing reference values therewith. More specifically, when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value, the subject is identified as having Alzheimer's disease or mild cognitive impairment. The reference value of desmosterol level in blood and the reference value of gelsolin level in blood are as described above.

[0054] The method for selecting a patient which is likely to benefit from the therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring a desmosterol level in blood and a gelsolin level in blood of a subject which is diagnosed with Alzheimer's disease

or mild cognitive impairment and comparing reference values therewith. More specifically, the selection can be made based on an indicator of a patient to be administered with the candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood and the reference value of gelsolin level in blood are as described above.

[0055] The method for predicting the therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention makes identification by measuring a desmosterol level and a gelsolin level in blood of a subject which is diagnosed with Alzheimer's disease and comparing reference values therewith. More specifically, the prediction can be made based on an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value. The reference value of desmosterol level in blood and the reference value of gelsolin level in blood are as described above.

[0056] The candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment of the present invention is not particularly limited, and preferably as described above.

#### **Examples**

[0057] Example 1: Measurement of desmosterol levels in blood of healthy elderly people and patients with Alzheimer's disease

#### Sample

Plasma of healthy elderly people (10 cases) and patients with Alzheimer's disease (10 cases) were used as samples.

[0058] Measurement of desmosterol

To 25  $\mu$ l of plasma were added as the internal standards 100  $\mu$ L of 500 ng/ml of desmosterol-d6 (purchased from Avanti Polar Lipids, Inc.) and 100  $\mu$ L of 200  $\mu$ g/ml of cholesterol-d7 (purchased from KANTO CHEMICAL CO., INC.), and 100  $\mu$ L of a 50% potassium hydroxide aqueous solution was further added thereto, and the incubation was carried out at 70°C for 60 minutes. Subsequently, 2 mL of hexane and 0.5 mL of phosphate buffered saline (pH 6.8) were added thereto, stirred and centrifuged, and the hexane layer

was collected. The extraction operation was repeated twice. After combining the hexane layers obtained, the solvent was dried by evaporation using a nitrogen gas. The pellet was dissolved in ethanol and the solution was subjected to liquid chromatography/atmospheric pressure chemical ionization mass spectrometry (LC/APCI-MS).

5 [0059] The apparatus for LC/APCI-MS used was an LC-20AD system (manufactured by Shimadzu Corporation) equipped with an autosampler SIL-20AC, a column oven CTO-20AC and a quadrupole mass spectrometer LCMS-2010EV. The column temperature was set at 50°C, using YMC-Pack Pro C18 RS column (purchased from YMC Co., Ltd.) having an internal diameter of 4.6 mm and a length of 250 mm as a column. The  
10 mobile phase used was a mixed solvent of water and methanol (flow rate 1 ml/min.). More specifically, solution A (water:methanol = 50:50) and solution B (methanol) were used, with B being 85% during 0 to 45 minutes, 100% during 45 to 55 minutes, and 85% during 55 to 70 minutes.

15 [0060] The MS analysis was carried out in Selective Ion Monitoring (SIM) mode, using, as the monitoring ions, m/z 369.3, 376.3, 367.3 and 373.3 (corresponding to cholesterol, cholesterol-d7, desmosterol and desmosterol-d6, respectively). The concentration of each sample was calculated based on the calibration curve of the subject to be measured.

20 [0061] The results are shown in the following Tables 1 and 2. The desmosterol levels in plasma of the patients with Alzheimer's disease were found to have been lower than the level of healthy elderly people.

[0062] [Table 1]

Healthy elderly people

Sex	Age	Desmosterol (ng/mL)
F	63	931
M	72	865
F	66	880
M	68	886
M	71	854
F	60	1261
M	61	500
M	69	795
F	74	621
F	71	889
Average	67.5	848

[0063] [Table 2]

Patients with Alzheimer's disease

Sex	Age	Desmosterol (ng/mL)
M	80	367
M	70	519
M	77	367
M	84	590
F	86	284
F	70	337
F	75	427
F	71	692
M	74	322
F	90	433
Average	77.7	434

5 [0064] Example 2: Measurement of desmosterol levels and amyloid beta levels in blood of healthy elderly people and patients with Alzheimer's disease

Sample

Plasma of healthy elderly people (10 cases) and patients with Alzheimer's disease (10 cases) were used as samples.

[0065] Measurement of desmosterol

10 The measurement was carried out by the same method as in Example 1.

[0066] Measurement of A $\beta$ x-40 and A $\beta$ x-42

15 The concentrations of A $\beta$ x-40 and A $\beta$ x-42 in blood were measured using a WAKO A $\beta$ x-40 ELISA kit (Wako Pure Chemical Industries, Ltd.) and a WAKO A $\beta$ x-42 High sensitivity ELISA kit (Wako Pure Chemical Industries, Ltd.) in accordance with the manuals of the kits.

[0067] The results are shown in the following Tables 3 and 4.

[0068] [Table 3]

Healthy elderly people

Sex	Age	Desmosterol (ng/mL)	x-40 (pM)	x-42 (pM)	x-42/x-40
F	63	931	88	6.7	0.076
M	72	865	77	7.7	0.101
F	66	880	61	7.3	0.120
M	68	886	92	8.9	0.096
M	71	854	98	8.8	0.090
F	60	1261	67	7.2	0.107
M	61	500	94	8.5	0.091
M	69	795	61	5.9	0.097
F	74	621	64	5.5	0.086
F	71	889	53	5.1	0.097
Average	67.5	848	75.5	7.2	0.096

[0069] [Table 4]

5

Patients with Alzheimer's disease

Sex	Age	Desmosterol (ng/mL)	x-40 (pM)	x-42 (pM)	x-42/x-40
M	80	367	96	8.7	0.090
M	70	519	99	7.7	0.078
M	77	367	88	5.3	0.060
M	84	590	88	5.3	0.060
F	86	284	99	8.2	0.083
F	70	337	55	4.3	0.078
F	75	427	78	6.0	0.076
F	71	692	34	1.7	0.049
M	74	322	87	7.6	0.088
F	90	433	84	7.2	0.085
Average	77.7	434	80.9	6.2	0.075

[0070] Statistical analysis

Using the above data, the Receiver Operating Characteristic Curve was used to evaluate the diagnosis accuracy of each measured value and the combinations thereof. The Receiver Operating Characteristic Curve of each of desmosterol and  $A\beta_{x-42}/A\beta_{x-40}$  was drawn using ROC, a basic package for R. For the combination of desmosterol and  $A\beta_{x-42}/A\beta_{x-40}$ , a machine learning model called SVM (Support Vector Machine), (R package e1071) was used. The evaluation was carried out by random sampling of 10 samples, as a set of training examples, from the total of 20 samples consisting of 10 samples of the healthy elderly people group and 10 samples of the group of patients with Alzheimer's disease, with the remaining 10 samples being a set of test examples. Also, the Receiver

Operating Characteristic Curves of the prediction results of the test data were drawn using the ROCR package (Figure 1). Desmosterol alone had AUC of 0.96, A $\beta$ x-42/A $\beta$ x-40 had AUC of 0.91, and the combination of desmosterol and A $\beta$ x-42/A $\beta$ x-40 had AUC of 1.

[0071] Example 3: Measurement of desmosterol levels and amyloid beta levels in blood of healthy elderly people and patients with mild cognitive impairment

Sample

Plasma of healthy elderly people (42 cases) and patients with mild cognitive impairment (26 cases) were used as samples.

[0072] Measurement of desmosterol

The measurement was carried out by the same method as in Example 1.

[0073] Measurement of A $\beta$ x-40 and A $\beta$ x-42

The measurement was carried out by the same method as in Example 2.

[0074] Statistical analysis

Using the above data, the Receiver Operating Characteristic Curve was used as in Example 2 to evaluate the diagnosis accuracy of each measured value and the combinations thereof. The evaluation was carried out by random sampling of 34 samples, as a set of training examples, from the total of 68 samples consisting of 42 samples of the healthy elderly people group and 26 samples of the group of patients with mild cognitive impairment, with the remaining 34 samples being a set of test examples. Also, the Receiver Operating Characteristic Curves of the prediction results of the test data were drawn using the ROCR package (Figure 2). Desmosterol alone had AUC of 0.71, A $\beta$ x-42/A $\beta$ x-40 had AUC of 0.84, and the combination of desmosterol and A $\beta$ x-42/A $\beta$ x-40 had AUC of 0.92.

[0075] Example 4: Measurement of desmosterol levels, amyloid beta levels and gelsolin levels in blood of healthy elderly people, patients with Alzheimer's disease and patients with mild cognitive impairment

Sample

Plasma of healthy elderly people (37 cases), patients with Alzheimer's disease (39 cases) and patients with mild cognitive impairment (26 cases) were used as samples.

[0076] Measurement of desmosterol

The measurement was carried out by the same method as in Example 1.

[0077] Measurement of A $\beta$ x-40 and A $\beta$ x-42

The measurement was carried out by the same method as in Example 2.

[0078] Measurement of gelsolin

The concentration of gelsolin in blood was measured using a gelsolin ELISA kit for human (USCN Life Science Inc.) in accordance with the manual of the kit.

[0079] Statistical analysis

Using the above data, the Receiver Operating Characteristic Curve was used as in  
 5 Example 2 to evaluate the diagnosis accuracy of each measured value and the combinations thereof. The evaluation was carried out by random sampling of 38 samples, as a set of training examples, from the total of 76 samples consisting of 37 samples of the healthy elderly people group and 39 samples of the patients with Alzheimer's disease group, with the remaining 38 samples being a set of test examples. The Receiver Operating Characteristic  
 10 Curves of the prediction results of the test data were drawn using the ROCR package (Figure 3). Desmosterol/cholesterol had AUC of 0.80, A $\beta$ x-42/A $\beta$ x-40 had AUC of 0.88, gelsolin had AUC of 0.88, and the combination of three had AUC of 0.98. The combination of three had a sensitivity of 90% when the specificity is 100%.

[0080] Also, the evaluation was carried out by random sampling of 32 samples, as a set of  
 15 training examples, from the total of 63 samples consisting of 37 samples of the healthy elderly people group and 26 samples of the patients with mild cognitive impairment group, with the remaining 31 samples being a set of test examples. The Receiver Operating Characteristic Curves of the prediction results of the test data were drawn using the ROCR package (Figure 4). Desmosterol/cholesterol had AUC of 0.71, A $\beta$ x-42/A $\beta$ x-40 had AUC  
 20 of 0.81, gelsolin had AUC of 0.80, and the combination of three had AUC of 0.86. The combination of three had a sensitivity of 85% when the specificity is 85%.

[0081] Example 5: Measurement of desmosterol levels and gelsolin levels in blood of healthy elderly people, patients with Alzheimer's disease and patients with mild cognitive impairment

25 Sample

Plasma of healthy elderly people (42 cases), patients with Alzheimer's disease (41 cases) and patients with mild cognitive impairment (26 cases) were used as samples.

[0082] Measurement of desmosterol

The measurement was carried out by the same method as in Example 1.

30 [0083] Measurement of gelsolin

The measurement was carried out by the same method as in Example 4.

[0084] Statistical analysis

Using the above data, the Receiver Operating Characteristic Curve was used as in

Example 2 to evaluate the diagnosis accuracy of each measured value and the combinations thereof. The evaluation was carried out by random sampling of 42 samples, as a set of training examples, from the total of 83 samples consisting of 42 samples of the healthy elderly people group and 41 samples of the patients with Alzheimer's disease group, with the remaining 41 samples being a set of test examples. The Receiver Operating Characteristic Curves of the prediction results of the test data were drawn using the ROCR package (Figure 5). Desmosterol/cholesterol had AUC of 0.80, gelsolin had AUC of 0.88, and the combination of both had AUC of 0.96.

[0085] Also, the evaluation was carried out by random sampling of 34 samples, as a set of training examples, from the total of 68 samples consisting of 42 samples of the healthy elderly people group and 26 samples of the patients with mild cognitive impairment group, with the remaining 34 samples being a set of test examples. The Receiver Operating Characteristic Curves of the prediction results of the test data were drawn using the ROCR package (Figure 6). Desmosterol/cholesterol had AUC of 0.71, gelsolin had AUC of 0.80 and the combination of both had AUC of 0.82.

[0086] Example 6: Measurement of desmosterol levels in blood of healthy elderly people and patients with mild cognitive impairment

Sample

Plasma of healthy elderly people (42 cases) and patients with mild cognitive impairment (26 cases) were used as samples.

[0087] Measurement of desmosterol

The measurement was carried out by the same method as in Example 1. The desmosterol levels in blood of healthy elderly people and patients with mild cognitive impairment were shown in Figure 7. The average plasma desmosterol level of healthy elderly people was 1046 ng/ml and the average plasma desmosterol level of patients with mild cognitive impairment was 770 ng/ml. Significant difference was found between both groups ( $p < 0.01$ ; t-test).

[0088] Statistical analysis

Using the above data, the Receiver Operating Characteristic Curve was used as in Example 2 to evaluate the diagnosis accuracy. The evaluation was carried out by random sampling of 34 samples, as a set of training examples, from the total of 68 samples consisting of 42 samples of the healthy elderly people group and 26 samples of the patients with mild cognitive impairment group, with the remaining 34 samples being a set of test examples.

The Receiver Operating Characteristic Curves of the prediction results of the test data were drawn using the ROCR package (Figure 8). AUC was 0.72.

**CLAIMS**

1. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:
- 5 a step of measuring a desmosterol level in blood of a non-human animal;  
a step of administering a candidate to the non-human animal;  
a step of measuring a desmosterol level in blood of the non-human animal after the candidate is administered; and
- 10 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the non-human animal after the candidate is administered is higher than the level before the candidate is administered.
2. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:
- 15 a step of measuring a desmosterol level in blood of a first non-human animal;  
a step of administering a candidate to a second non-human animal;  
a step of measuring a desmosterol level in blood of the second non-human animal after the candidate is administered; and
- 20 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the second non-human animal after the candidate is administered is higher than the desmosterol level in blood of the first non-human animal to which the candidate is not administered.
3. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:
- 25 a step of measuring a desmosterol level in a blood sample of a subject before a candidate is administered;  
a step of measuring a desmosterol level in the blood sample of the subject after the candidate is administered; and
- 30 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered.
4. A method for aiding diagnosis of Alzheimer's disease or mild cognitive

impairment, the method comprising:

a step of measuring a desmosterol level in a blood sample of a subject; and

a step of providing an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value.

5

5. A method for diagnosing Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level in blood of a subject; and

a step of identifying Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value.

10

6. A method for selecting a patient which is likely to benefit from a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level in a blood sample of a subject; and

a step of providing an indicator of a patient to be administered with a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value.

15

7. A method for predicting a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level in a blood sample of a subject; and

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value.

20

8. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and an amyloid beta level in blood of a non-human animal;

a step of administering a candidate to the non-human animal;

a step of measuring a desmosterol level and an amyloid beta level in blood of the non-human animal after the candidate is administered; and

30

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the non-human animal after the candidate is administered is higher than the level

before the candidate is administered, and  $A\beta_x-42$  or  $A\beta_x-42/A\beta_x-40$  in blood of the non-human animal after the candidate is administered is higher than that before the candidate is administered.

5 9. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and an amyloid beta level in blood of a first non-human animal;

a step of administering a candidate to a second non-human animal;

10 a step of measuring a desmosterol level and an amyloid beta level in blood of the second non-human animal after the candidate is administered; and

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the second non-human animal after the candidate is administered is higher than the desmosterol level in blood of the first non-human animal to which the candidate is not administered, and  $A\beta_x-42$  or  $A\beta_x-42/A\beta_x-40$  in blood of the second non-human animal after the candidate is administered is higher than  $A\beta_x-42$  or  $A\beta_x-42/A\beta_x-40$  in blood of the first non-human animal to which the candidate is not administered.

15

10. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

20

a step of measuring a desmosterol level and an amyloid beta level in a blood sample of a subject before a candidate is administered;

a step of measuring a desmosterol level and an amyloid beta level in the blood sample of the subject after the candidate is administered; and

25

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered, and  $A\beta_x-42$  or  $A\beta_x-42/A\beta_x-40$  in blood of the subject after the candidate is administered is higher than that before the candidate is administered.

30

11. A method for aiding diagnosis of Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and an amyloid beta level in a blood sample of a subject; and

a step of providing an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value.

5 12. A method for diagnosing Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and an amyloid beta level in blood of a subject; and

10 a step of identifying Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value.

13. A method for selecting a patient which is likely to benefit from a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

15 a step of measuring a desmosterol level and an amyloid beta level in a blood sample of a subject; and

a step of providing an indicator of a patient to be administered with a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value.

20

14. A method for predicting a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and an amyloid beta level in a blood sample of a subject; and

25 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value.

30 15. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood of a non-human animal;

a step of administering a candidate to the non-human animal;

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood of the non-human animal after the candidate is administered; and

5 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the non-human animal after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the non-human animal after the candidate is administered is higher than that before the candidate is administered, and the gelsolin level in blood of the non-human animal after the candidate is administered is higher than the gelsolin level in blood before the candidate is administered.

10

16. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood of a first non-human animal;

15 a step of administering a candidate to a second non-human animal;

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood of the second non-human animal after the candidate is administered; and

20 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the second non-human animal after the candidate is administered is higher than the desmosterol level in blood of the first non-human animal to which the candidate is not administered,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the second non-human animal after the candidate is administered is higher than  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the first non-human animal to which the candidate is not administered, and the gelsolin level in blood of the second non-human animal after the candidate is administered is higher than the gelsolin level in blood of the first non-human animal to which the candidate is not administered.

25

17. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

30 a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in a blood sample of a subject before a candidate is administered,

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in the blood sample of the subject after the candidate is administered,

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject after the candidate is administered is higher than the desmosterol level before the candidate is administered,  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the subject after the candidate is administered is higher than that before the candidate is administered, and the gelsolin level in blood of the subject after the candidate is administered is higher than the gelsolin level in blood before the candidate is administered.

18. A method for aiding diagnosis of Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in a blood sample of a subject; and

a step of providing an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

19. A method for diagnosing Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in blood of a subject; and

a step of identifying Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

20. A method for selecting a patient which is likely to benefit from a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in a blood sample of a subject; and

a step of providing an indicator of a patient to be administered with a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta x-42$  or  $A\beta x-42/A\beta x-40$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

21. A method for predicting a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level, an amyloid beta level and a gelsolin level in a blood sample of a subject, and

5 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value,  $A\beta_{x-42}$  or  $A\beta_{x-42}/A\beta_{x-40}$  in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

10 22. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and a gelsolin level in blood of a non-human animal;

a step of administering a candidate to the non-human animal;

15 a step of measuring a desmosterol level and a gelsolin level in blood of the non-human animal after the candidate is administered; and

20 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the non-human animal after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered, and the gelsolin level in blood of the non-human animal after the candidate is administered is higher than the gelsolin level in blood before the candidate is administered.

23. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

25 a step of measuring a desmosterol level and a gelsolin level in blood of a first non-human animal;

a step of administering a candidate to a second non-human animal;

a step of measuring a desmosterol level and a gelsolin level in blood of the second non-human animal after the candidate is administered; and

30 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the second non-human animal after the candidate is administered is higher than the desmosterol level in blood of the first non-human animal to which the candidate is not

administered, and the gelsolin level in blood of the second non-human animal after the candidate is administered is higher than the gelsolin level in blood of the first non-human animal to which the candidate is not administered.

24. A method for evaluating an effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and a gelsolin level in a blood sample of a subject before a candidate is administered;

a step of measuring a desmosterol level and a gelsolin level in the blood sample of the subject after the candidate is administered; and

a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject after the candidate is administered is higher than the desmosterol level in blood before the candidate is administered, and the gelsolin level in blood of the subject after the candidate is administered is higher than the gelsolin level in blood before the candidate is administered.

25. A method for aiding diagnosis of Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and a gelsolin level in a blood sample of a subject; and

a step of providing an indicator for potentially having Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

26. A method for diagnosing Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and a gelsolin level in blood of a subject; and

a step of identifying Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

27. A method for selecting a patient which is likely to benefit from a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and a gelsolin level in a blood sample of a

subject; and

5 a step of providing an indicator of a patient to be administered with a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

28. A method for predicting a therapeutic effect of a candidate for a therapeutic agent to treat Alzheimer's disease or mild cognitive impairment, the method comprising:

a step of measuring a desmosterol level and a gelsolin level in a blood sample of a subject; and

10 a step of providing an indicator of being potential for the candidate to be effective in treating Alzheimer's disease or mild cognitive impairment when the desmosterol level in blood of the subject is lower than a reference value, and the gelsolin level in blood of the subject is lower than a reference value.

15






Fig.1


Fig.2



**Fig.3**



**Fig.4**









Fig.5

**Fig.6**



**Fig.7**





**Fig.8**