



(51) International Patent Classification:

A61K 31/439 (2006.01) C07D 261/20 (2006.01)
A61K 31/423 (2006.01) C07D 453/02 (2006.01)
A61P 25/28 (2006.01)

(21) International Application Number:

PCT/US2016/036689

(22) International Filing Date:

9 June 2016 (09.06.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/173,717 10 June 2015 (10.06.2015) US

(71) Applicant: **FORUM PHARMACEUTICALS, INC.**
[US/US]; 225 Second Avenue, Waltham, MA 02451 (US).

(72) Inventors: **ACHARYA, Raksha**; 2 Glen Ora Drive, Bedford, MA 01730 (US). **BURNETT, Duane, A.**; 19 Sherman Bridge Road, Wayland, MA 01778 (US). **BURSAVICH, Matthew, Gregory**; 172 Damon Road, Needham, MA 02494 (US). **COOK, Andrew, Simon**; 9 October Lane, Stow, MA 01775 (US). **HARRISON, Bryce, Alden**; 7 Old Wood Road, Framingham, MA 01701 (US). **MCRINER, Andrew, J.**; 31 Lynde Street, Melrose, MA 02176 (US).

(74) Agents: **SHARER, Paul, L.** et al.; Jones Day, 250 Vesey Street, New York, NY 10281-1047 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: AMINO BENZISOXAZOLE COMPOUNDS AS AGONISTS OF A7-NICOTINIC ACETYLCHOLINE RECEPTORS

(57) Abstract: The present invention relates to novel aminobenzisoxazole compounds, and pharmaceutical compositions of the same, that are suitable as agonists or partial agonists of $\alpha 7$ -nAChR, and methods of preparing these compounds and compositions, and the use of these compounds and compositions in methods of maintaining, treating and/or improving cognitive function. In particular, methods of administering the compound or composition to a patient in need thereof, for example a patient with a cognitive deficiency and/or a desire to enhance cognitive function, that may derive a benefit therefrom.



**AMINO BENZISOXAZOLE COMPOUNDS AS AGONISTS OF
 $\alpha 7$ -NICOTINIC ACETYLCHOLINE RECEPTORS**

CROSS REFERENCE TO RELATED APPLICATION

[0001] This application claims the benefit of priority from U.S. Provisional Application No. 62/173,717, filed June 10, 2015. The foregoing related application, in its entirety, is incorporated herein by reference.

FIELD OF THE INVENTION

[0002] The present invention relates to novel aminobenzisoxazole compounds, and pharmaceutical compositions of the same, that are suitable as agonists or partial agonists of the $\alpha 7$ -nicotinic acetylcholine receptor, and methods of preparing these compounds and compositions, and the use of these compounds and compositions in methods of maintaining, treating and/or improving cognitive function. In particular, methods of administering the compound or composition to a patient in need thereof, for example a patient with a cognitive deficiency and/or a desire to enhance cognitive function, that may derive a benefit therefrom.

BACKGROUND OF THE INVENTION

[0003] The prevalence of cognitive disease, for example dementia in North America, is approximately 6 to 10% of the population, with Alzheimer's disease accounting for a substantial portion of these cases. Many forms of cognitive disease represent a steadily growing medical and social problem of our aging societies around the world. Some believe the main pathological features may relate to intraneuronal neurofibrillary tangles, formation of amyloid beta plaques and/or neurodegeneration of mainly cholinergic and, in later stages, also serotonergic, noradrenergic, and other neurons, resulting in deficiencies of acetylcholine and other neurotransmitters. Some theories suggest that the gradual development of an acetylcholine signaling deficiency may be responsible for the early clinical manifestations of cognitive disease. Consequently, some believe that compounds that improve cholinergic functioning, such as acetylcholine esterase inhibitors may ameliorate the cognitive deficits in patients with cognitive disease. The most widely used acetylcholine esterase inhibitor is donepezil hydrochloride (Aricept®).


[0004] Nicotinic acetylcholine receptors (nAChR) form a large family of ion channels which are activated by the messenger acetylcholine which is produced in the body (Galzi and Changeux, *Neuropharmacol.* 1995, 34, 563-582). A functional nAChR consists of five subunits which may be different (certain combinations of $\alpha 1-9$ and $\beta 1-4, \gamma, \delta, \epsilon$ subunits) or identical ($\alpha 7-9$). This leads to the formation of a diversity of subtypes which differ in the distribution in the muscles, the nervous system and other organs (McGehee and Role, *Annu. Rev. Physiol.* 1995, 57, 521-546). Activation of nAChR

leads to influx of cations into the cell and to stimulation of nerve cells or muscle cells. Selective activation of individual nAChR subtypes restricts this stimulation to the cell types which have a corresponding subtype and is thus able to avoid unwanted side effects such as, for example, stimulation of nAChR in the muscles. Clinical experiments with nicotine and experiments in various animal models indicate that central nicotinic acetylcholine receptors are involved in learning and memory processes (e.g. Rezvani and Levin, *Biol. Psychiatry* 2001, 49, 258-267). Nicotinic acetylcholine receptors of the $\alpha 7$ subtype ($\alpha 7$ nAChR) have a particularly high concentration in regions of the brain which are important for learning and memory, such as the hippocampus and the cerebral cortex (Séguéla et al., *J. Neurosci.* 1993, 13, 596-604). The $\alpha 7$ nAChR has a particularly high permeability for calcium ions, modulates neurotransmission, influences the growth of axons and, in this way, modulates neuronal plasticity (Broide and Leslie, *Mol. Neurobiol.* 1999, 20, 1-16).

[0005] WO 2003/055878 describes a variety of agonists of the $\alpha 7$ nAChR said to be useful for improving cognition. WO 2003/055878 suggests that certain agonists of the $\alpha 7$ nAChR are useful for improving perception, concentration, learning or memory, especially after cognitive impairments like those occurring for example in situations/diseases/syndromes such as mild cognitive impairment, age-associated learning and memory impairments, age-associated memory loss, Alzheimer's disease, schizophrenia and certain other cognitive disorders.

BRIEF SUMMARY OF THE INVENTION

[0006] An aspect of the invention provides an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib):



wherein:

R^1 , R^2 , R^3 , and R^4 independently represent -H, -D, halogen radical, -CN, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, an unbranched $-OC_1$ - C_4 -alkyl, a branched or cyclic $-OC_3$ - C_4 -alkyl, $-N(R^5)(R^6)$, $-(CO)N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-SO_2C_1$ - C_4 -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1$ - C_4 -alkyl, an aryl radical, or a heteroaryl radical; wherein the alkyl portion of the unbranched C_1 - C_4 -alkyl radical, the branched C_3 - C_4 -alkyl radical, the C_3 - C_6 -cycloalkyl radical, the unbranched $-OC_1$ - C_4 -alkyl, the branched or cyclic $-OC_3$ - C_4 -alkyl, the $-SO_2C_1$ - C_4 -alkyl, the $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, or the $-N(R^5)SO_2C_1$ - C_4 -alkyl, may be independently substituted with up to 5 radical

substituents comprising: $-D$, halogen radical, $=O$, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1-C_4$ -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1-C_4$ -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1-C_4$ -alkyl, $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an unbranched C_1-C_6 -alkyl radical, a branched C_3-C_6 -alkyl radical, a C_3-C_6 -cycloalkyl radical, a C_1-C_6 -hydroxyalkyl radical, a C_1-C_2 -haloalkyl radical, or $-OC_1-C_2$ -haloalkyl radical; and wherein the aryl radical or the heteroaryl radical may be independently substituted with up to 5 radical substituents comprising: $-D$, halogen radical, $-CN$, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1-C_4$ -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1-C_4$ -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1-C_4$ -alkyl, $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an unbranched C_1-C_6 -alkyl radical, a branched C_3-C_6 -alkyl radical, a C_3-C_6 -cycloalkyl radical, a C_1-C_6 -hydroxyalkyl radical, a C_1-C_2 -haloalkyl radical, or $-OC_1-C_2$ -haloalkyl radical;

R^5 and R^6


independently represent $-H$; an unbranched C_1-C_6 -alkyl radical, a branched C_3-C_6 -alkyl radical; a C_3-C_6 -cycloalkyl radical; or the $N(R^5)(R^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a C_2-C_6 -alkyl di-radical or a (3-6 membered)-heteroalkyl di-radical; wherein the (3-6 membered)-heteroalkyl di-radical comprises at least one ring atom selected from the group consisting of oxygen, nitrogen, and sulfur, with the proviso that when the at least one ring atom is nitrogen, the nitrogen is independently substituted with $-H$, an unbranched C_1-C_4 -alkyl radical, a branched C_3-C_4 -alkyl radical, a C_3-C_4 -cycloalkyl radical, $-(CO)$ -unbranched C_1-C_4 -alkyl, $-(CO)$ -branched C_3-C_4 -alkyl, $-(SO_2)$ -unbranched C_1-C_4 -alkyl, or $-(SO_2)$ -branched C_3-C_4 -alkyl, and with the further proviso that when the at least one ring atom is sulfur, the sulfur may be independently substituted with 0 to 2 $=O$; wherein the C_2-C_6 -alkyl di-radical or the alkyl portion of said (3-6 membered)-heteroalkyl di-radical may be independently substituted with up to 5 radical substituents comprising: $-D$, halogen radical, $=O$, an unbranched C_1-C_6 -alkyl radical, or a branched C_3-C_6 -alkyl radical; and

m


independently represents an integer from 1 to 6;

or a pharmaceutically acceptable salt thereof.

[0007] An aspect of the invention relates to the aminobenzisoxazole compound represented by Formula (Ia):



[0008] An aspect of the invention relates to the aminobenzisoxazole compound represented by Formula (Ib):



[0009] An aspect of the invention relates to a single stereoisomer of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof.

[0010] An aspect of the invention relates to a single enantiomer or a single diastereomer of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof.

[0011] An aspect of the invention relates to a pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0012] An aspect of the invention relates to a method comprising administering to a patient in need thereof an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0013] Another aspect of the invention provides a method of treating a patient in need thereof, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0014] Another aspect of the invention provides a method of maintaining, treating, curing and/or improving at least one cognitive function in a patient in need thereof, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound

represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0015] Another aspect of the invention provides a method of maintaining, treating, curing and/or improving at least one cognitive function in a patient in need thereof, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0016] Another aspect of the invention provides a method of treating a patient diagnosed as having a cognitive impairment, comprising: administering to the an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient an effective dose of a pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0017] Another aspect of the invention provides a method of treating a patient in need thereof, comprising: administering to the patient, for example, a patient diagnosed with having a cognitive impairment, Limited Cognitive Impairment, Mild Cognitive Impairment, Alzheimer's disease, and/or schizophrenia, an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent; such that the patient may derive a benefit therefrom.

[0018] Another aspect of the invention provides a method of treating one or more symptoms associated with a cognitive impairment, comprising administering to a patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent; wherein the patient suffers from, or has been diagnosed as having, a cognitive impairment.

[0019] Another aspect of the invention provides a method of improving cognition of a patient in need thereof, comprising: administering to the patient an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole

compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0020] Another aspect of the invention provides a method of improving cognition in a patient suffering from a cognitive impairment, such as a cognitive impairment associated with either schizophrenia or Alzheimer's disease, for example mild Alzheimer's disease, moderate Alzheimer's disease, severe Alzheimer's disease, or mild-to-moderate Alzheimer's disease, comprising administering an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0021] Another aspect of the invention provides a method of treating a patient suffering from, diagnosed with having, or suffers from one or more symptoms associated with, a cognitive impairment, for example, Alzheimer's disease, dementia of an Alzheimer's type, MCI, LCI, or schizophrenia, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent. For example, the method of treating a patient suffering from, diagnosed with having, or suffers from one or more symptoms associated with, a cognitive impairment, may provide said patient at least one of the following: (i) treats, minimizes progression of, prevents the deterioration of, or reduces the rate of deterioration of, one or more symptoms associated with the cognitive impairment; (ii) treats the cognitive impairment; (iii) improves cognition in said cognitively impaired patient; (iv) improves one or more behavioral symptoms associated with the cognitive impairment; (v) provides a pro-cognitive effect; (vi) provides a pro-cognitive effect in at least one of the following: visual motor, learning, delayed memory, or executive function, or (vii) provides a positive effect on clinical function in said cognitively impaired patient.

[0022] Another aspect of the invention provides a method of treating a patient previously treated, or currently being treated, with an AChEI, that is suffering from, or has been diagnosed with having, a cognitive impairment, for example, Alzheimer's disease, dementia of an Alzheimer's type, MCI, LCI, or schizophrenia, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluents;

wherein the method improves one or more symptoms associated with the cognitive impairment in the previously, or currently, AChEI treated patient.

[0023] Another aspect of the invention provides a method of treating a patient suffering from, or diagnosed with having a cognitive impairment, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent; wherein the method provides a positive effect on cognition or a positive effect on clinical function in said cognitively impaired patient, and wherein said patient has been previously treated or is currently being treated with an AChEI.

[0024] Another aspect of the invention provides a method of improving cognition in a patient diagnosed as having a probable cognitive disease, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient an effective dose of a pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0025] Another aspect of the invention provides a method of improving or substantially improving one or more symptoms in a cognitive disease patient, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient an effective dose of a pharmaceutical composition comprising the effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0026] Another aspect of the invention provides a method of slowing the rate of deterioration of at least one symptom in a cognitive disease patient, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient the pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0027] Another aspect of the invention provides a method of treating one or more symptoms associated with a cognitive disease in a patient suffering therefrom, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient an effective dose of a pharmaceutical composition comprising the aminobenzisoxazole compound represented by

Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent

[0028] Another aspect provides a method of minimizing or substantially halting the rate of progression of one or more cognitive diseases in a patient suffering from a cognitive disease, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient an effective dose of a pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0029] Another aspect of the invention provides a method of substantially stopping or reversing progression of one or more cognitive diseases, in a patient suffering therefrom, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient an effective dose of a pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0030] Another aspect of the invention provides a method of treating dementia, comprising: administering to a patient in need thereof an effective amount of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient an effective dose of a pharmaceutical composition comprising the effective amount of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent; wherein said effective amount is administered in an effective dose.

[0031] Another aspect of the invention provides a method of treating dementia, comprising: administering to a patient in need thereof an effective amount of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient an effective dose of a pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0032] Another aspect of the invention provides a method of treating dementia, comprising: administering to a patient in need thereof an effective amount of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, wherein the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, is administered in the form of a pharmaceutical composition comprising at least one pharmaceutically acceptable carrier, excipient or diluent.


[0033] Another aspect of the invention provides a method of treating dementia, comprising: administering to a patient in need thereof an effective amount of an aminobenzisoxazole compound

represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient an effective dose of a pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent; wherein the pharmaceutical composition is in the form of a tablet.

[0034] Another aspect of the invention provides a method of treating a patient having a cognitive disease and being administered an acetylcholine esterase inhibitor, comprising: administering to a patient in need thereof an effective amount of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient an effective dose of a pharmaceutical composition comprising an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent; wherein the treatment comprises halting the administration of the acetylcholine esterase inhibitor prior to treating with the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof.

DETAILED DESCRIPTION OF THE INVENTION

[0035] An embodiment of the present invention provides an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib):



wherein:

R^1 , R^2 , R^3 , and R^4 independently represent $-H$, $-D$, halogen radical, $-CN$, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, an unbranched $-OC_1$ - C_4 -alkyl, a branched or cyclic $-OC_3$ - C_4 -alkyl, $-N(R^5)(R^6)$, $-(CO)N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-SO_2C_1$ - C_4 -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1$ - C_4 -alkyl, an aryl radical, or a heteroaryl radical; wherein the alkyl portion of the unbranched C_1 - C_4 -alkyl radical, the branched C_3 - C_4 -alkyl radical, the C_3 - C_6 -cycloalkyl radical, the unbranched $-OC_1$ - C_4 -alkyl, the branched or cyclic $-OC_3$ - C_4 -alkyl, the $-SO_2C_1$ - C_4 -alkyl, the $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, or the $-N(R^5)SO_2C_1$ - C_4 -alkyl, may be independently substituted with up to 5 radical substituents comprising: $-D$, halogen radical, $=O$, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1$ - C_4 -alkyl,

$-\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CH}_2)_m\text{SO}_2\text{C}_1\text{-C}_4\text{-alkyl}$, $-(\text{CH}_2)_m\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$,
 $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1\text{-C}_4\text{-alkyl}$, $-(\text{CO})(\text{CH}_2)_m\text{R}^5$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, an unbranched $\text{C}_1\text{-C}_6\text{-alkyl}$ radical, a branched $\text{C}_3\text{-C}_6\text{-alkyl}$ radical, a $\text{C}_3\text{-C}_6\text{-cycloalkyl}$ radical, a $\text{C}_1\text{-C}_6\text{-hydroxyalkyl}$ radical, a $\text{C}_1\text{-C}_2\text{-haloalkyl}$ radical, or $-\text{OC}_1\text{-C}_2\text{-haloalkyl}$ radical; and wherein the aryl radical or the heteroaryl radical may be independently substituted with up to 5 radical substituents comprising: $-\text{D}$, halogen radical, $-\text{CN}$, $-\text{OR}^5$, $-(\text{CH}_2)_m\text{OR}^5$, $-\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{SO}_2\text{C}_1\text{-C}_4\text{-alkyl}$, $-\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CH}_2)_m\text{SO}_2\text{C}_1\text{-C}_4\text{-alkyl}$, $-(\text{CH}_2)_m\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1\text{-C}_4\text{-alkyl}$, $-(\text{CO})(\text{CH}_2)_m\text{R}^5$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, an unbranched $\text{C}_1\text{-C}_6\text{-alkyl}$ radical, a branched $\text{C}_3\text{-C}_6\text{-alkyl}$ radical, a $\text{C}_3\text{-C}_6\text{-cycloalkyl}$ radical, a $\text{C}_1\text{-C}_6\text{-hydroxyalkyl}$ radical, a $\text{C}_1\text{-C}_2\text{-haloalkyl}$ radical, or $-\text{OC}_1\text{-C}_2\text{-haloalkyl}$ radical;

R^5 and R^6

independently represent $-\text{H}$; an unbranched $\text{C}_1\text{-C}_6\text{-alkyl}$ radical, a branched $\text{C}_3\text{-C}_6\text{-alkyl}$ radical, a $\text{C}_3\text{-C}_6\text{-cycloalkyl}$ radical; or the $\text{N}(\text{R}^5)(\text{R}^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a $\text{C}_2\text{-C}_6\text{-alkyl}$ di-radical or a (3-6 membered)-heteroalkyl di-radical; wherein the (3-6 membered)-heteroalkyl di-radical comprises at least one ring atom selected from the group consisting of oxygen, nitrogen, and sulfur, with the proviso that when the at least one ring atom is nitrogen, the nitrogen is independently substituted with $-\text{H}$, an unbranched $\text{C}_1\text{-C}_4\text{-alkyl}$ radical, a branched $\text{C}_3\text{-C}_4\text{-alkyl}$ radical, a $\text{C}_3\text{-C}_4\text{-cycloalkyl}$ radical, $-(\text{CO})\text{-unbranched C}_1\text{-C}_4\text{-alkyl}$, $-(\text{CO})\text{-branched C}_3\text{-C}_4\text{-alkyl}$, $-(\text{SO}_2)\text{-unbranched C}_1\text{-C}_4\text{-alkyl}$, or $-(\text{SO}_2)\text{-branched C}_3\text{-C}_4\text{-alkyl}$, and with the further proviso that when the at least one ring atom is sulfur, the sulfur may be independently substituted with 0 to 2 $=\text{O}$; wherein the $\text{C}_2\text{-C}_6\text{-alkyl}$ di-radical or the alkyl portion of said (3-6 membered)-heteroalkyl di-radical may be independently substituted with up to 5 radical substituents comprising: $-\text{D}$, halogen radical, $=\text{O}$ an unbranched $\text{C}_1\text{-C}_6\text{-alkyl}$ radical, or a branched $\text{C}_3\text{-C}_6\text{-alkyl}$ radical; and

m

independently represents an integer from 1 to 6;

or a pharmaceutically acceptable salt thereof.

[0036] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^1 , R^2 , R^3 , and R^4 independently representing $-\text{H}$, $-\text{D}$, halogen radical, $-\text{CN}$, an unbranched $\text{C}_1\text{-C}_3\text{-alkyl}$ radical, a branched $\text{C}_3\text{-C}_4\text{-alkyl}$ radical, a $\text{C}_3\text{-C}_5\text{-cycloalkyl}$ radical, an unbranched $-\text{OC}_1\text{-C}_4\text{-alkyl}$, a branched or cyclic $-\text{OC}_3\text{-C}_4\text{-alkyl}$, $-\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, $-\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CH}_2)_m\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, $-(\text{CH}_2)_m\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, an aryl radical, or a heteroaryl radical; wherein the alkyl portion of the unbranched $\text{C}_1\text{-C}_3\text{-alkyl}$ radical, the branched $\text{C}_3\text{-C}_4\text{-alkyl}$ radical, the $\text{C}_3\text{-C}_5\text{-cycloalkyl}$ radical, the

unbranched $-\text{OC}_1\text{-C}_4\text{-alkyl}$, the branched or cyclic $-\text{OC}_3\text{-C}_4\text{-alkyl}$, the $-\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, the $-(\text{CH}_2)_m\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, or the $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-\text{D}$, halogen radical, $=\text{O}$, $-\text{OR}^5$, $-(\text{CH}_2)_m\text{OR}^5$, $-\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, $-\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CH}_2)_m\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, $-(\text{CH}_2)_m\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, $-(\text{CO})(\text{CH}_2)_m\text{R}^5$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, an unbranched $\text{C}_1\text{-C}_4\text{-alkyl}$ radical, a branched $\text{C}_3\text{-C}_4\text{-alkyl}$ radical, a $\text{C}_3\text{-C}_5\text{-cycloalkyl}$ radical, a $\text{C}_1\text{-C}_4\text{-hydroxyalkyl}$ radical, a $\text{C}_1\text{-C}_2\text{-haloalkyl}$ radical, or $-\text{OC}_1\text{-C}_2\text{-haloalkyl}$ radical; and wherein the aryl radical or the heteroaryl radical may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-\text{D}$, halogen radical, $-\text{CN}$, $-\text{OR}^5$, $-(\text{CH}_2)_m\text{OR}^5$, $-\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, $-\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CH}_2)_m\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, $-(\text{CH}_2)_m\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1\text{-C}_2\text{-alkyl}$, $-(\text{CO})(\text{CH}_2)_m\text{R}^5$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, an unbranched $\text{C}_1\text{-C}_4\text{-alkyl}$ radical, a branched $\text{C}_3\text{-C}_4\text{-alkyl}$ radical, a $\text{C}_3\text{-C}_5\text{-cycloalkyl}$ radical, a $\text{C}_1\text{-C}_4\text{-hydroxyalkyl}$ radical, a $\text{C}_1\text{-C}_2\text{-haloalkyl}$ radical, or $-\text{OC}_1\text{-C}_2\text{-haloalkyl}$ radical; and wherein R^5 and R^6 may independently represent $-\text{H}$, an unbranched $\text{C}_1\text{-C}_6\text{-alkyl}$ radical, such as $-\text{CH}_3$ or $-\text{CH}_2\text{CH}_3$, a branched $\text{C}_3\text{-C}_6\text{-alkyl}$ radical, such as $-\text{CH}(\text{CH}_3)_2$, or a $\text{C}_3\text{-C}_6\text{-cycloalkyl}$ radical, such as a cyclopropyl radical, or the $\text{N}(\text{R}^5)(\text{R}^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a $\text{C}_2\text{-C}_6\text{-alkyl}$ di-radical, such as a $\text{C}_2\text{-C}_5\text{-alkyl}$ di-radical, or a (3-6 membered)-heteroalkyl di-radical, such as a (3-5 membered)-heteroalkyl di-radical; and wherein m may independently represents an integer from 1 to 6, for example, an integer from 1 to 4, such as 1 to 2, 2 to 3, or 3 to 4.

[0037] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^1 and R^2 independently representing $-\text{H}$, $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{CN}$, an unbranched $\text{C}_1\text{-C}_3\text{-alkyl}$ radical, a branched $\text{C}_3\text{-C}_4\text{-alkyl}$ radical, a $\text{C}_3\text{-C}_4\text{-cycloalkyl}$ radical, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{CF}_3$, an unbranched $-\text{OC}_1\text{-C}_3\text{-alkyl}$, a branched or cyclic $-\text{OC}_3\text{-alkyl}$, $-\text{OCF}_3$, $-\text{SO}_2\text{CH}_3$, $-\text{SO}_2\text{N}(\text{CH}_3)_2$, or $-\text{N}(\text{R}^5)\text{SO}_2\text{CH}_3$; wherein the alkyl portion of the unbranched $\text{C}_1\text{-C}_3\text{-alkyl}$ radical, the branched $\text{C}_3\text{-C}_4\text{-alkyl}$ radical, the $\text{C}_3\text{-C}_4\text{-cycloalkyl}$ radical, the unbranched $-\text{OC}_1\text{-C}_3\text{-alkyl}$, or the branched or cyclic $-\text{OC}_3\text{-alkyl}$, may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{OR}^5$, $=\text{O}$, $-\text{CH}_3$, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{CF}_3$, cyclopropyl radical, cyclobutyl radical, or $-\text{OCF}_3$; and wherein R^5 may independently represent $-\text{H}$, $-\text{CH}_3$, or $-\text{CH}_2\text{CH}_3$.

[0038] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^1 and R^2 independently representing $-\text{H}$, $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, a cyclopropyl radical, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{CF}_3$, $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}(\text{CH}_3)_2$, $-\text{OCF}_3$, $-\text{SO}_2\text{CH}_3$, $-\text{SO}_2\text{N}(\text{CH}_3)_2$, or $-\text{N}(\text{H})\text{SO}_2\text{CH}_3$.

[0039] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^1 and R^2 independently representing $-\text{H}$, $-\text{D}$, or halogen radical, for

example, -F, -Cl, or -Br. For example, in certain embodiments, R^1 and R^2 may independently represent -H, -D, -F, or -Cl, such as -H, -D, or -F. In certain embodiments, R^1 and R^2 may independently represent -H or -D. In certain embodiments, R^1 may independently represent -H or -D, and R^2 may independently represent -F or -Cl, such as -F. In certain embodiments, R^1 may independently represent -F or -Cl, such as -F, and R^2 may independently represent -H or -D.

[0040] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^3 independently representing -F, -Cl, -Br, -CN, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, an unbranched $-OC_1$ - C_4 -alkyl, a branched or cyclic $-OC_3$ - C_4 -alkyl, $-N(R^5)(R^6)$, $-(CO)N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-SO_2C_1$ - C_4 -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1$ - C_4 -alkyl, an aryl radical, or a heteroaryl radical; wherein the alkyl portion of the unbranched C_1 - C_4 -alkyl radical, the branched C_3 - C_4 -alkyl radical, the C_3 - C_6 -cycloalkyl radical, the unbranched $-OC_1$ - C_4 -alkyl, the branched or cyclic $-OC_3$ - C_4 -alkyl, the $-SO_2C_1$ - C_4 -alkyl, the $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, or the $-N(R^5)SO_2C_1$ - C_4 -alkyl, may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: -D, -F, -Cl, -Br, =O, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1$ - C_4 -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1$ - C_4 -alkyl, $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an unbranched C_1 - C_6 -alkyl radical, a branched C_3 - C_6 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, a C_1 - C_6 -hydroxyalkyl radical, a C_1 - C_2 -haloalkyl radical, or $-OC_1$ - C_2 -haloalkyl radical; and wherein the aryl radical or the heteroaryl radical may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: -D, -F, -Cl, -Br, -CN, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1$ - C_4 -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1$ - C_4 -alkyl, $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an unbranched C_1 - C_6 -alkyl radical, a branched C_3 - C_6 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, a C_1 - C_6 -hydroxyalkyl radical, a C_1 - C_2 -haloalkyl radical, or $-OC_1$ - C_2 -haloalkyl radical; and wherein R^5 and R^6 may independently represent -H, an unbranched C_1 - C_6 -alkyl radical, such as $-CH_3$ or $-CH_2CH_3$, a branched C_3 - C_6 -alkyl radical, such as $-CH(CH_3)_2$, or a C_3 - C_6 -cycloalkyl radical, such as a cyclopropyl radical, or the $N(R^5)(R^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a C_2 - C_6 -alkyl di-radical, such as a C_2 - C_5 -alkyl di-radical, or a (3-6 membered)-heteroalkyl di-radical, such as a (3-5 membered)-heteroalkyl di-radical; and wherein m may independently represents an integer from 1 to 6, for example, an integer from 1 to 4, such as 1 to 2, 2 to 3, or 3 to 4.

[0041] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^3 independently representing -F, -Cl, -Br, -CN, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_4 -cycloalkyl radical, an unbranched $-OC_1$ - C_4 -alkyl, a branched or cyclic $-OC_3$ - C_4 -alkyl, $-OCHF_2$, $-OCH_2F$, $-OCF_3$, $-OCH_2CF_3$, $-N(R^5)(R^6)$, $-(CO)N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-SO_2CH_3$, $-SO_2N(R^5)(R^6)$, $-CH_2CH_2SO_2C_1$ - C_4 -alkyl, or

$-\text{N}(\text{R}^5)\text{SO}_2\text{CH}_3$, an aryl radical, or a heteroaryl radical; wherein the alkyl portion of the unbranched $\text{C}_1\text{-C}_4$ -alkyl radical, the branched $\text{C}_3\text{-C}_4$ -alkyl radical, the $\text{C}_3\text{-C}_4$ -cycloalkyl radical, the unbranched $-\text{OC}_1\text{-C}_4$ -alkyl, the branched or cyclic $-\text{OC}_3\text{-C}_4$ -alkyl, or the $-\text{CH}_2\text{CH}_2\text{SO}_2\text{C}_1\text{-C}_4$ -alkyl, may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $=\text{O}$, $-\text{OR}^5$, $-(\text{CH}_2)_m\text{OR}^5$, $-\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{SO}_2\text{C}_1\text{-C}_4$ -alkyl, $-\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CH}_2)_m\text{SO}_2\text{C}_1\text{-C}_4$ -alkyl, $-(\text{CH}_2)_m\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1\text{-C}_4$ -alkyl, $-(\text{CO})(\text{CH}_2)_m\text{R}^5$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, an unbranched $\text{C}_1\text{-C}_6$ -alkyl radical, a branched $\text{C}_3\text{-C}_6$ -alkyl radical, a $\text{C}_3\text{-C}_6$ -cycloalkyl radical, a $\text{C}_1\text{-C}_6$ -hydroxyalkyl radical, a $\text{C}_1\text{-C}_2$ -haloalkyl radical, or $-\text{OC}_1\text{-C}_2$ -haloalkyl radical; and wherein the aryl radical or the heteroaryl radical may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{CN}$, $-\text{OR}^5$, $-(\text{CH}_2)_m\text{OR}^5$, $-\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{SO}_2\text{C}_1\text{-C}_4$ -alkyl, $-\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CH}_2)_m\text{SO}_2\text{C}_1\text{-C}_4$ -alkyl, $-(\text{CH}_2)_m\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1\text{-C}_4$ -alkyl, $-(\text{CO})(\text{CH}_2)_m\text{R}^5$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, an unbranched $\text{C}_1\text{-C}_6$ -alkyl radical, a branched $\text{C}_3\text{-C}_6$ -alkyl radical, a $\text{C}_3\text{-C}_6$ -cycloalkyl radical, a $\text{C}_1\text{-C}_6$ -hydroxyalkyl radical, a $\text{C}_1\text{-C}_2$ -haloalkyl radical, or $-\text{OC}_1\text{-C}_2$ -haloalkyl radical; and wherein R^5 and R^6 may independently represent $-\text{H}$, an unbranched $\text{C}_1\text{-C}_6$ -alkyl radical, such as $-\text{CH}_3$ or $-\text{CH}_2\text{CH}_3$, a branched $\text{C}_3\text{-C}_6$ -alkyl radical, such as $-\text{CH}(\text{CH}_3)_2$, or a $\text{C}_3\text{-C}_6$ -cycloalkyl radical, such as a cyclopropyl radical, or the $\text{N}(\text{R}^5)(\text{R}^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a $\text{C}_2\text{-C}_6$ -alkyl di-radical, such as a $\text{C}_2\text{-C}_5$ -alkyl di-radical, or a (3-6 membered)-heteroalkyl di-radical, such as a (3-5 membered)-heteroalkyl di-radical; and wherein m may independently represents an integer from 1 to 6, for example, an integer from 1 to 4, such as 1 to 2, 2 to 3, or 3 to 4.

[0042] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^3 independently representing $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{CN}$, $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, cyclopropyl radical, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{CF}_3$, $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}(\text{CH}_3)_2$, $-\text{O-cyclopropyl}$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{F}$, $-\text{OCF}_3$, $-\text{OCH}_2\text{CF}_3$, $-\text{SO}_2\text{CH}_3$, a phenyl radical or a heteroaryl radical, such as an N-pyrazole radical, a furan radical, a thiophene radical, an imidazole radical, an oxazole radical, a thiazole radical, a pyridyl radical, a pyrazine radical, a pyrimidine radical, or an oxadiazole radical; wherein the phenyl radical or the heteroaryl radical, such as the N-pyrazole radical, the furan radical, the thiophene radical, the imidazole radical, the oxazole radical, the thiazole radical, the pyridyl radical, the pyrazine radical, the pyrimidine radical, or the oxadiazole radical, may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{CN}$, $-\text{OR}^5$, $-(\text{CH}_2)_m\text{OR}^5$, $-\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CO})(\text{CH}_2)_m\text{R}^5$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, an unbranched $\text{C}_1\text{-C}_6$ -alkyl radical, a branched $\text{C}_3\text{-C}_6$ -alkyl radical, a $\text{C}_3\text{-C}_6$ -cycloalkyl radical, a $\text{C}_1\text{-C}_6$ -hydroxyalkyl radical, a $\text{C}_1\text{-C}_2$ -haloalkyl radical, or $-\text{OC}_1\text{-C}_2$ -haloalkyl radical; and wherein R^5 and R^6 may independently represent $-\text{H}$, an unbranched $\text{C}_1\text{-C}_6$ -alkyl radical, such as $-\text{CH}_3$ or $-\text{CH}_2\text{CH}_3$, a branched $\text{C}_3\text{-C}_6$ -alkyl radical, such as $-\text{CH}(\text{CH}_3)_2$, or a $\text{C}_3\text{-C}_6$ -cycloalkyl radical, such as a cyclopropyl radical, or the $\text{N}(\text{R}^5)(\text{R}^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a $\text{C}_2\text{-C}_6$ -alkyl di-radical, such as a $\text{C}_2\text{-C}_5$ -alkyl di-radical, or a (3-6 membered)-heteroalkyl di-radical, such as a (3-5 membered)-heteroalkyl di-radical; and wherein m may independently represents an integer from 1 to 6, for example, an integer from 1 to 4, such as 1 to 2, 2 to 3, or 3 to 4.

$\text{CH}(\text{CH}_3)_2$, or a C_3 - C_6 -cycloalkyl radical, such as a cyclopropyl radical, or the $\text{N}(\text{R}^5)(\text{R}^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a C_2 - C_6 -alkyl di-radical, such as a C_2 - C_5 -alkyl di-radical, or a (3-6 membered)-heteroalkyl di-radical, such as a (3-5 membered)-heteroalkyl di-radical; and wherein m may independently represents an integer from 1 to 6, for example, an integer from 1 to 4, such as 1 to 2, 2 to 3, or 3 to 4.

[0043] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^3 independently representing $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{CN}$, $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, cyclopropyl radical, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{CF}_3$, $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}(\text{CH}_3)_2$, $-\text{O}$ -cyclopropyl, $-\text{OCF}_3$, $-\text{SO}_2\text{CH}_3$, a phenyl radical or a heteroaryl radical, such as an N-pyrazole radical, a furan radical, a thiophene radical, an imidazole radical, an oxazole radical, a thiazole radical, a pyridyl radical, a pyrazine radical, a pyrimidine radical, or an oxadiazole radical; wherein the phenyl radical or the heteroaryl radical, such as the N-pyrazole radical, the furan radical, the thiophene radical, the imidazole radical, the oxazole radical, the thiazole radical, the pyridyl radical, the pyrazine radical, the pyrimidine radical, or the oxadiazole radical, may be substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{CN}$, $-\text{OR}^5$, $-\text{CH}_3$, cyclopropyl radical, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{CF}_3$, $-\text{OCF}_3$, or $-\text{OCH}_2\text{CF}_3$. For example, in certain embodiments, R^3 may independently represent $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{CN}$, $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, cyclopropyl radical, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{CF}_3$, $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}(\text{CH}_3)_2$, $-\text{O}$ -cyclopropyl, or $-\text{OCF}_3$. In certain embodiments, R^3 may independently represent $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{CH}_3$, or $-\text{OCH}_3$, such as R^3 may independently represent $-\text{Cl}$, $-\text{CH}_3$, or $-\text{OCH}_3$.

[0044] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^4 independently representing $-\text{H}$, $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{CN}$, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, an unbranched $-\text{OC}_1$ - C_4 -alkyl, a branched or cyclic $-\text{OC}_3$ - C_4 -alkyl, $-\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-\text{SO}_2\text{C}_1$ - C_4 -alkyl, $-\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CH}_2)_m\text{SO}_2\text{C}_1$ - C_4 -alkyl, $-(\text{CH}_2)_m\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1$ - C_4 -alkyl, an aryl radical, or a heteroaryl radical; wherein the alkyl portion of the unbranched C_1 - C_4 -alkyl radical, the branched C_3 - C_4 -alkyl radical, the C_3 - C_6 -cycloalkyl radical, the unbranched $-\text{OC}_1$ - C_4 -alkyl, the branched or cyclic $-\text{OC}_3$ - C_4 -alkyl, the $-\text{SO}_2\text{C}_1$ - C_4 -alkyl, the $-(\text{CH}_2)_m\text{SO}_2\text{C}_1$ - C_4 -alkyl, or the $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1$ - C_4 -alkyl, may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $=\text{O}$, $-\text{OR}^5$, $-(\text{CH}_2)_m\text{OR}^5$, $-\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{SO}_2\text{C}_1$ - C_4 -alkyl, $-\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CH}_2)_m\text{SO}_2\text{C}_1$ - C_4 -alkyl, $-(\text{CH}_2)_m\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1$ - C_4 -alkyl, $-(\text{CO})(\text{CH}_2)_m\text{R}^5$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{OCF}_3$ and an unbranched C_1 - C_6 -alkyl radical, a branched C_3 - C_6 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, a C_1 - C_6 -hydroxyalkyl radical, a C_1 - C_2 -haloalkyl radical, or $-\text{OC}_1$ - C_2 -haloalkyl radical; and wherein aryl radical or the heteroaryl radical may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$,

$-\text{CN}$, $-\text{OR}^5$, $-(\text{CH}_2)_m\text{OR}^5$, $-\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{SO}_2\text{C}_1\text{-C}_4\text{-alkyl}$,
 $-\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CH}_2)_m\text{SO}_2\text{C}_1\text{-C}_4\text{-alkyl}$, $-(\text{CH}_2)_m\text{SO}_2\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{N}(\text{R}^5)\text{SO}_2\text{C}_1\text{-C}_4\text{-alkyl}$,
 $-(\text{CO})(\text{CH}_2)_m\text{R}^5$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{OCF}_3$, an unbranched $\text{C}_1\text{-C}_6\text{-alkyl}$ radical, a branched $\text{C}_3\text{-C}_6\text{-alkyl}$
 radical, a $\text{C}_3\text{-C}_6\text{-cycloalkyl}$ radical, a $\text{C}_1\text{-C}_6\text{-hydroxyalkyl}$ radical, a $\text{C}_1\text{-C}_2\text{-haloalkyl}$ radical, or $-\text{OC}_1\text{-C}_2\text{-haloalkyl}$ radical; and wherein R^5 and R^6 may independently represent $-\text{H}$, an unbranched $\text{C}_1\text{-C}_6\text{-alkyl}$
 radical, such as $-\text{CH}_3$ or $-\text{CH}_2\text{CH}_3$, a branched $\text{C}_3\text{-C}_6\text{-alkyl}$ radical, such as $-\text{CH}(\text{CH}_3)_2$, or a $\text{C}_3\text{-C}_6\text{-cycloalkyl}$ radical, such as a cyclopropyl radical, or the $\text{N}(\text{R}^5)(\text{R}^6)$ moiety forms a cycle, wherein
 R^5 and R^6 taken together represent a $\text{C}_2\text{-C}_6\text{-alkyl}$ di-radical, such as a $\text{C}_2\text{-C}_5\text{-alkyl}$ di-radical, or a (3-
 6 membered)-heteroalkyl di-radical, such as a (3-5 membered)-heteroalkyl di-radical; and wherein m
 may independently represents an integer from 1 to 6, for example, an integer from 1 to 4, such as 1 to
 2, 2 to 3, or 3 to 4.

[0045] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia)
 or Formula (Ib), may comprise R^4 independently representing $-\text{H}$, $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{CN}$, an unbranched
 $\text{C}_1\text{-C}_3\text{-alkyl}$ radical, a branched $\text{C}_3\text{-C}_4\text{-alkyl}$ radical, a $\text{C}_3\text{-C}_4\text{-cycloalkyl}$ radical, unbranched $-\text{OC}_1\text{-C}_3\text{-alkyl}$,
 a branched or cyclic $-\text{OC}_3\text{-C}_4\text{-alkyl}$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{F}$, $-\text{OCF}_3$, $-\text{OCH}_2\text{CF}_3$; wherein the alkyl
 portion of the unbranched $\text{C}_1\text{-C}_3\text{-alkyl}$ radical, the branched $\text{C}_3\text{-C}_4\text{-alkyl}$ radical, the $\text{C}_3\text{-C}_4\text{-cycloalkyl}$
 radical, the unbranched $-\text{OC}_1\text{-C}_3\text{-alkyl}$, or the branched or cyclic $-\text{OC}_3\text{-C}_4\text{-alkyl}$, may be
 independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or
 up to 3 radical substituents, comprising: $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $=\text{O}$, $-\text{OR}^5$, $-(\text{CH}_2)_m\text{OR}^5$, $-\text{N}(\text{R}^5)(\text{R}^6)$,
 $-\text{NR}^5(\text{CO})(\text{R}^6)$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)(\text{R}^6)$, $-(\text{CO})(\text{CH}_2)_m\text{R}^5$, $-(\text{CO})\text{N}(\text{R}^5)(\text{R}^6)$, $-\text{OCF}_3$, an unbranched $\text{C}_1\text{-C}_6\text{-alkyl}$
 radical, a branched $\text{C}_3\text{-C}_6\text{-alkyl}$ radical, a $\text{C}_3\text{-C}_6\text{-cycloalkyl}$ radical, a $\text{C}_1\text{-C}_6\text{-hydroxyalkyl}$ radical,
 a $\text{C}_1\text{-C}_2\text{-haloalkyl}$ radical, or $-\text{OC}_1\text{-C}_2\text{-haloalkyl}$ radical; and wherein R^5 and R^6 may independently
 represent $-\text{H}$, an unbranched $\text{C}_1\text{-C}_6\text{-alkyl}$ radical, such as $-\text{CH}_3$ or $-\text{CH}_2\text{CH}_3$, a branched $\text{C}_3\text{-C}_6\text{-alkyl}$
 radical, such as $-\text{CH}(\text{CH}_3)_2$, or a $\text{C}_3\text{-C}_6\text{-cycloalkyl}$ radical, such as a cyclopropyl radical, or the
 $\text{N}(\text{R}^5)(\text{R}^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a $\text{C}_2\text{-C}_6\text{-alkyl}$ di-radical,
 such as a $\text{C}_2\text{-C}_5\text{-alkyl}$ di-radical, or a (3-6 membered)-heteroalkyl di-radical, such as a (3-5
 membered)-heteroalkyl di-radical; and wherein m may independently represents an integer from 1 to
 6, for example, an integer from 1 to 4, such as 1 to 2, 2 to 3, or 3 to 4.

[0046] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia)
 or Formula (Ib), may comprise R^4 independently representing $-\text{H}$, $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{CN}$, $-\text{CH}_3$,
 cyclopropyl radical, cyclobutyl radical, $-\text{CH}_2\text{F}$, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}(\text{CH}_3)_2$,
 $-\text{O-cyclopropyl}$, $-\text{OCHF}_2$, $-\text{OCH}_2\text{F}$, $-\text{OCF}_3$, or $-\text{OCH}_2\text{CF}_3$. For example, in certain embodiments, R^4
 may independently represent $-\text{H}$, $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{CN}$, $-\text{CH}_3$, cyclopropyl radical, cyclobutyl radical,
 $-\text{CH}_2\text{F}$, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}(\text{CH}_3)_2$, $-\text{O-cyclopropyl}$, or $-\text{OCF}_3$, such as R^4 may
 independently represent $-\text{H}$, $-\text{D}$, $-\text{F}$, $-\text{Cl}$, $-\text{CH}_3$, $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, or $-\text{CF}_3$.

[0047] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia)
 or Formula (Ib), may comprise: R^1 and R^2 independently representing $-\text{H}$ or $-\text{D}$, R^2 and R^4

independently representing $-H$ or $-D$, R^1 and R^4 independently representing $-H$ or $-D$, or R^1 , R^2 , and R^4 independently representing $-H$ or $-D$; and R^3 independently representing $-H$, $-D$, $-F$, $-Cl$, $-Br$, $-CN$, an unbranched C_1 - C_3 -alkyl radical, for example, $-CH_3$ or $-CH_2CH_3$, a branched C_3 - C_4 -alkyl radical, a cyclopropyl radical, a cyclobutyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, an unbranched $-OC_1$ - C_3 -alkyl, such as $-OCH_3$ or $-OCH_2CH_3$, a branched or cyclic $-OC_3$ - C_4 -alkyl, such as $-OCH(CH_3)_2$ or $-O$ -cyclopropyl, $-OCHF_2$, $-OCH_2F$, $-OCF_3$, $-OCH_2CF_3$, $-SO_2CH_3$, a phenyl radical or a heteroaryl radical, such as an N-pyrazole radical, a furan radical, a thiophene radical, an imidazole radical, an oxazole radical, a thiazole radical, a pyridyl radical, a pyrazine radical, a pyrimidine radical, or an oxadiazole radical; wherein the alkyl portion of the unbranched C_1 - C_3 -alkyl radical, branched C_3 - C_4 -alkyl radical, unbranched $-OC_1$ - C_3 -alkyl, or the branched or cyclic $-OC_3$ - C_4 -alkyl, may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-D$, $-F$, $-Cl$, $=O$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-CH_3$, $-CH_2CH_3$, cyclopropyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCF_3$, or $-OCH_2CF_3$; and wherein the phenyl radical or the heteroaryl radical may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-D$, $-F$, $-Cl$, $-Br$, $-CN$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-CH_3$, $-CH_2CH_3$, cyclopropyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCF_3$, or $-OCH_2CF_3$.

[0048] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^1 independently representing $-H$, $-D$, or a halogen radical, for example $-H$, $-D$, $-F$, or $-Cl$, such as $-H$, $-D$, or $-F$; R^2 independently representing $-H$, $-D$, or a halogen radical, for example $-H$, $-D$, $-F$, or $-Cl$, such as $-H$, $-D$, or $-F$; R^3 independently representing $-H$, $-D$, $-F$, $-Cl$, $-Br$, $-CN$, an unbranched C_1 - C_3 -alkyl radical, for example, $-CH_3$ or $-CH_2CH_3$, $-CH(CH_3)_2$, a cyclopropyl radical, a cyclobutyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, an unbranched $-OC_1$ - C_3 -alkyl, such as $-OCH_3$ or $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-O$ -cyclopropyl, $-OCHF_2$, $-OCH_2F$, $-OCF_3$, $-OCH_2CF_3$, $-SO_2CH_3$, a phenyl radical or a heteroaryl radical, such as an N-pyrazole radical or an oxadiazole radical; and R^4 independently representing $-H$, $-D$, $-F$, $-Cl$, $-CN$, an unbranched C_1 - C_3 -alkyl radical, for example, $-CH_3$ or $-CH_2CH_3$, $-CH(CH_3)_2$, a cyclopropyl radical, a cyclobutyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, an unbranched $-OC_1$ - C_3 -alkyl, such as $-OCH_3$ or $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-O$ -cyclopropyl, $-OCHF_2$, $-OCH_2F$, $-OCF_3$, $-OCH_2CF_3$; wherein the alkyl portion of the unbranched C_1 - C_3 -alkyl radical, $-CH(CH_3)_2$, unbranched $-OC_1$ - C_3 -alkyl, $-OCH(CH_3)_2$, or $-O$ -cyclopropyl, may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-D$, $-F$, $-Cl$, $=O$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-CH_3$, $-CH_2CH_3$, cyclopropyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCF_3$, or $-OCH_2CF_3$; and wherein the phenyl radical or the heteroaryl radical may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-D$, $-F$, $-Cl$, $-Br$, $-CN$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-CH_3$, $-CH_2CH_3$, cyclopropyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCF_3$, or $-OCH_2CF_3$.

[0049] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^1 independently representing $-H$, $-D$, or a halogen radical, for example $-H$, $-D$, $-F$, or $-Cl$, such as $-H$, $-D$, or $-F$; R^2 independently representing $-H$, $-D$, or a halogen radical, for example $-H$, $-D$, $-F$, or $-Cl$, such as $-H$, $-D$, or $-F$; R^3 independently representing $-H$, $-D$, $-F$, $-Cl$, $-Br$, $-CN$, an unbranched C_1 - C_3 -alkyl radical, for example, $-CH_3$ or $-CH_2CH_3$, $-CH(CH_3)_2$, a cyclopropyl radical, a cyclobutyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, an unbranched $-OC_1$ - C_3 -alkyl, such as $-OCH_3$ or $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-O$ -cyclopropyl, $-OCHF_2$, $-OCH_2F$, $-OCF_3$, $-OCH_2CF_3$, or $-SO_2CH_3$; and R^4 independently representing $-H$, $-D$, $-F$, $-Cl$, $-CN$, an unbranched C_1 - C_3 -alkyl radical, for example, $-CH_3$ or $-CH_2CH_3$, $-CH(CH_3)_2$, a cyclopropyl radical, a cyclobutyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, an unbranched $-OC_1$ - C_3 -alkyl, such as $-OCH_3$ or $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-O$ -cyclopropyl, $-OCHF_2$, $-OCH_2F$, $-OCF_3$, $-OCH_2CF_3$; wherein the alkyl portion of the unbranched C_1 - C_3 -alkyl radical, $-CH(CH_3)_2$, the cyclopropyl radical, the cyclobutyl radical, the unbranched $-OC_1$ - C_3 -alkyl, $-OCH(CH_3)_2$, or $-O$ -cyclopropyl, may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-D$, $-F$, $-Cl$, $=O$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-CH_3$, $-CH_2CH_3$, cyclopropyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCF_3$, or $-OCH_2CF_3$.

[0050] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^1 independently representing $-H$, $-D$, or a halogen radical, for example $-H$, $-D$, $-F$, or $-Cl$, such as $-H$, $-D$, or $-F$; R^2 independently representing $-H$, $-D$, or a halogen radical, for example $-H$, $-D$, $-F$, or $-Cl$, such as $-H$, $-D$, or $-F$; R^3 independently representing $-H$, $-D$, $-F$, $-Cl$, $-Br$, $-CN$, $-CH_3$, $-CH_2CH_3$, $-CH(CH_3)_2$, a cyclopropyl radical, a cyclobutyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, or $-O$ -cyclopropyl, or $-OCF_3$; and R^4 independently representing $-H$, $-D$, $-F$, $-Cl$, $-CN$, $-CH_3$, $-CH_2CH_3$, a cyclopropyl radical, a cyclobutyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, or $-O$ -cyclopropyl, $-OCF_3$.

[0051] For example, in certain embodiments, R^1 may independently represent $-H$, $-D$, $-F$, or $-Cl$, such as $-H$, $-D$, or $-F$; R^2 may independently represent $-H$, $-D$, $-F$, or $-Cl$, such as $-H$, $-D$, or $-F$; R^3 may independently represent $-F$, $-Cl$, $-Br$, $-CN$, $-CH_3$, $-CH_2CH_3$, a cyclopropyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, or $-O$ -cyclopropyl, or $-OCF_3$, such as $-F$, $-Cl$, $-Br$, $-CH_3$, or $-OCH_3$, or such as $-Cl$, $-CH_3$, or $-OCH_3$; and R^4 may independently represent $-H$, $-D$, $-F$, $-Cl$, $-CN$, $-CH_3$, $-CH_2CH_3$, a cyclopropyl radical, a cyclobutyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, or $-O$ -cyclopropyl, $-OCF_3$, such as $-H$, $-D$, $-F$, $-Cl$, $-CH_3$, $-OCH_3$, $-OCH_2CH_3$, or $-CF_3$.

[0052] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), at least one of R^1 , R^2 , R^3 , and R^4 does not independently represent $-H$. For example, in certain embodiments, R^1 independently represents $-H$, and at least one of R^2 , R^3 , and R^4 does not independently represent $-H$; R^2 independently represents $-H$, and at least one of R^1 , R^3 , and R^4 does

not independently represent $-H$; R^3 independently represents $-H$, and at least one of R^1 , R^2 , and R^4 does not independently represent $-H$; or R^4 independently represents $-H$, and at least one of R^1 , R^2 , and R^3 does not independently represent $-H$.

[0053] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise R^5 , R^6 , or both R^5 and R^6 , independently representing $-H$; an unbranched C_1 - C_6 -alkyl radical, such as $-CH_3$ or $-CH_2CH_3$, a branched C_3 - C_6 -alkyl radical, such as $-CH(CH_3)_2$; or a C_3 - C_6 -cycloalkyl radical, such as a cyclopropyl radical or a cyclobutyl radical. For example, R^5 and R^6 may independently represent $-H$, $-CH_3$, $-CH_2CH_3$, $-CH(CH_3)_2$, a cyclopropyl radical, or a cyclobutyl radical, such as independently represent $-H$, $-CH_3$, or $-CH_2CH_3$.

[0054] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise an $N(R^5)(R^6)$ moiety, wherein the $N(R^5)(R^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a C_2 - C_6 -alkyl di-radical or a (3-6 membered)-heteroalkyl di-radical; wherein the (3-6 membered)-heteroalkyl di-radical comprises at least one ring atom selected from the group consisting of oxygen, nitrogen, and sulfur, with the proviso that when the at least one ring atom is nitrogen, the nitrogen is independently substituted with $-H$, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_4 -cycloalkyl radical, $-(CO)$ -unbranched C_1 - C_4 -alkyl, $-(CO)$ -branched C_3 - C_4 -alkyl, $-(SO_2)$ -unbranched C_1 - C_4 -alkyl, or $-(SO_2)$ -branched C_3 - C_4 -alkyl, and with the further proviso that when the at least one ring atom is sulfur, the sulfur may be independently substituted with 0 to 2 $=O$; wherein the C_2 - C_6 -alkyl di-radical or the alky portion of said (3-6 membered)-heteroalkyl di-radical may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-D$, halogen radical, $=O$, an unbranched C_1 - C_6 -alkyl radical, or a branched C_3 - C_6 -alkyl radical.

[0055] In certain embodiments, the $N(R^5)(R^6)$ moiety may form a cycle, wherein R^5 and R^6 taken together represent a C_2 - C_6 -alkyl di-radical, such as a C_2 - C_5 -alkyl di-radical or C_3 - C_4 -alkyl di-radical; wherein the C_2 - C_6 -alkyl di-radical, such as a C_2 - C_5 -alkyl di-radical or C_3 - C_4 -alkyl di-radical, may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: $-D$, halogen radical, $=O$, an unbranched C_1 - C_6 -alkyl radical, or a branched C_3 - C_6 -alkyl radical. For example, the $N(R^5)(R^6)$ moiety may form a cycle, wherein R^5 and R^6 taken together represent a C_2 -alkyl di-radical, a C_3 -alkyl di-radical, C_4 -alkyl di-radical, or C_5 -alkyl di-radical, such as a C_2 -alkyl di-radical.

[0056] In certain embodiments, the $N(R^5)(R^6)$ moiety may, for example, form a cycle wherein the R^5 and R^6 taken together represent a (3-6 membered)-heteroalkyl di-radical, such as (4-5 membered)-heteroalkyl di-radical; wherein the (3-6 membered)-heteroalkyl di-radical comprises at least one ring atom selected from the group consisting of oxygen, nitrogen, and sulfur, with the proviso that when the at least one ring atom is nitrogen, the nitrogen is independently substituted with $-H$; an unbranched C_1 - C_4 -alkyl radical, such as $-CH_3$, $-CH_2CH_3$, or $-CH_2CH_2CH_3$, a branched C_3 - C_4 -alkyl radical, such as $-CH(CH_3)_2$; a C_3 - C_4 -cycloalkyl radical; $-(CO)$ -unbranched C_1 - C_4 -alkyl;

–(CO)–branched C₃–C₄–alkyl; –(SO₂)–unbranched C₁–C₄–alkyl; or –(SO₂)–branched C₃–C₄–alkyl; and with the further proviso that when the at least one ring atom is sulfur, the sulfur may be independently substituted with 0 to 2 =O; wherein the alky portion of said (3-6 membered)-heteroalkyl di-radical may be independently substituted with up to 5 radical substituents, for example, up to 4 radical substituents or up to 3 radical substituents, comprising: –D, halogen radical, =O, an unbranched C₁–C₆–alkyl radical, or a branched C₃–C₆–alkyl radical. For example, the N(R⁵)(R⁶) moiety may form a cycle, wherein R⁵ and R⁶ taken together represent a (4-5 membered)-heteroalkyl di-radical, wherein the (4-5 membered)-heteroalkyl di-radical comprises at least one ring atom selected from the group consisting of oxygen or nitrogen, with the proviso that when the at least one ring atom is nitrogen, the nitrogen is independently substituted with –H; –CH₃, –CH₂CH₃, –CH(CH₃)₂, a cyclopropyl radical. –(CO)CH₃, –(CO)CH₂CH₃, –(SO₂)CH₃, or –(SO₂)CH₂CH₃.

[0057] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise racemic mixture of enantiomers, a mixture of diastereomers, a single enantiomer, or a single diastereomer, of the compound, or a pharmaceutically acceptable salt thereof. In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), may comprise a mixture of tautomers, substantially a single tautomer form, or a single tautomer form, such as a tautomer contained within the aminobenzisoxazole ring system or a tautomer resulting from one or more substituents substituted on the aminobenzisoxazole ring system, for example, a tautomer may be contained within the aminobenzisoxazole ring system or one or more substituents substituted on the aminobenzisoxazole ring system containing a heteroaryl ring nitrogen adjacent to a heteroaryl ring carbon substituted with a hydroxyl group.

[0058] The chemical names and structure diagrams used herein to describe the compounds of the present invention, supra and infra, were created with the use of ChemBioDraw Ultra® Version 12.0 (available from CambridgeSoft Corp., Cambridge, Mass.).

[0059] In certain embodiments, specific examples of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib) may include, collectively or individually, the compounds listed below, and single enantiomers and pharmaceutically acceptable salts thereof:

N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 7-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 5-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 6-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 N-(quinuclidin-3-yl)-6-(trifluoromethyl)benzo[d]isoxazol-3-amine;
 6,7-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 4-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
3-(quinuclidin-3-ylamino)benzo[d]isoxazole-6-carbonitrile;
6-(methylsulfonyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-(tert-butyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
N-(quinuclidin-3-yl)-6-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
4-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
6-chloro-7-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
4-fluoro-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
4,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-fluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-chloro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-chloro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-isopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-7-carbonitrile;
7-chloro-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-(1H-pyrazol-1-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-(5-methyl-1,3,4-oxadiazol-2-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-methoxy-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6,7-dimethyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
6-bromo-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine.

[0060] In certain embodiments, specific examples of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib) may include, collectively or individually, the compounds listed below, and single enantiomers and pharmaceutically acceptable salts thereof:

6-cyclopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

6-ethyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
6-chloro-7-cyclobutyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5-fluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-methoxy-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-methyl-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
6-chloro-5,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-(difluoromethyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-isopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-cyclopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-ethoxy-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-ethoxy-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-methoxy-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5,6-difluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-5-fluoro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
6-chloro-5-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5,7-difluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6,7-dichloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-ethoxy-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
5,7-difluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine.

[0061] In certain embodiments, specific examples of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib) may include, collectively or individually, the single enantiomers listed below, and pharmaceutically acceptable salts thereof:

(*R*)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

(*R*)-6-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-N-(quinuclidin-3-yl)-6-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*S*)-N-(quinuclidin-3-yl)-6-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*R*)-6,7-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6,7-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-4-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-4-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-6-carbonitrile;
(*S*)-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-6-carbonitrile;
(*R*)-6-(methylsulfonyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-(methylsulfonyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-(tert-butyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-(tert-butyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-N-(quinuclidin-3-yl)-6-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
(*S*)-N-(quinuclidin-3-yl)-6-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
(*R*)-7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-4-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-4-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

(S)-6-chloro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(S)-6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(R)-6-chloro-7-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-chloro-7-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-4-fluoro-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-4-fluoro-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-4,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-4,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-7-fluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-7-fluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-7-chloro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-7-chloro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-7-chloro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-7-chloro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-isopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-isopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-chloro-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-7-carbonitrile;
(S)-6-chloro-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-7-carbonitrile;
(R)-7-chloro-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-7-chloro-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-(1H-pyrazol-1-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-(1H-pyrazol-1-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-(5-methyl-1,3,4-oxadiazol-2-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-(5-methyl-1,3,4-oxadiazol-2-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-7-methoxy-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-7-methoxy-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-chloro-7-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-chloro-7-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6,7-dimethyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6,7-dimethyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-chloro-7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-chloro-7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-chloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-chloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

(*R*)-6-bromo-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
(*S*)-6-bromo-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine.

[0062] In certain embodiments, specific examples of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib) may include, collectively or individually, the single enantiomers listed below, and pharmaceutically acceptable salts thereof:

(*R*)-6-cyclopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-cyclopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-ethyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-ethyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-cyclobutyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-cyclobutyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5-fluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5-fluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methoxy-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-methoxy-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methyl-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*S*)-6-methyl-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-5,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-5,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-(difluoromethyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-(difluoromethyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-isopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-isopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-cyclopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-cyclopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-ethoxy-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-ethoxy-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-ethoxy-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-ethoxy-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methoxy-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*S*)-6-methoxy-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*R*)-5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

(*R*)-5,6-difluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5,6-difluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-5-fluoro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-5-fluoro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-5-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-5-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5,7-difluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5,7-difluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6,7-dichloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6,7-dichloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-ethoxy-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-ethoxy-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5,7-difluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
(*S*)-5,7-difluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine.

[0063] In certain embodiments, specific examples of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib) may include, collectively or individually, the single enantiomers listed below, and pharmaceutically acceptable salts thereof:

(*R*)-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6,7-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
(*R*)-6-chloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine.

[0064] In certain embodiments, specific examples of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib) may include, collectively or individually, the single enantiomers listed below, and pharmaceutically acceptable salts thereof:

(*R*)-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6,7-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine; and
(*R*)-6-chloro-7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine.

[0065] In certain embodiments, specific examples of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib) may include, collectively or individually, the single enantiomers listed below, and pharmaceutically acceptable salts thereof:

(*S*)-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
(*R*)-6-chloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine.

[0066] In certain embodiments, the aminobenzisoxazole compounds of the present invention represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, may be more potent against $\alpha 7$ nAChR (according to the $\alpha 7$ nAChR Binding Assay (K_i)) than against a 5-HT₃ serotonin receptor (according to the [³H]BRL 43694 competition binding (K_i)). For example, the aminobenzisoxazole compounds of the present invention represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, may be at least 1.5 times more potent against $\alpha 7$ nAChR than against a 5-HT₃ serotonin receptor, as determined by the $\alpha 7$ nAChR Binding Assay and the [³H]BRL 43694 competition binding assay, respectively, such as at least 2 times more potent, at least 3 times more potent, at least 4 times more potent, at least 5 times more potent, at least 6 times more potent, at least 7 times more potent, at least 8 times more potent, at least 9 times more potent, at least 10 times more potent, at least 15 times more potent, at least 20 times more potent, or at least 25 times more potent against $\alpha 7$ nAChR than against a 5-HT₃ serotonin receptor, as determined by the $\alpha 7$ nAChR Binding Assay and the [³H]BRL 43694 competition binding assay, respectively.

[0067] As used herein, the term “treating” (or “treat” or “treatment”), unless otherwise specified, includes the generally accepted meaning which encompasses improving, modifying, decreasing, prohibiting, preventing, restraining, minimizing, slowing, halting, stopping, curing, and/or reversing a symptom associated with a disease and/or a disease. Treatment may include both therapeutic and prophylactic administration. For example, treatment of a cognitive impairment, in a patient diagnosed as having a cognitive impairment, may include, but is not limited to, curing the cognitive impairment, preventing the deterioration of one or more symptoms associated with the cognitive impairment;

improving cognition in a patient suffering from the cognitive impairment, slowing the progression of the cognitive impairment and/or modifying the cognitive impairment.

[0068] As used herein, the term “effective dose” (or “dose”), unless otherwise specified, is understood to include a therapeutically acceptable dose, a therapeutically acceptable amount, a therapeutically effective dose, a therapeutically effective amount, a pharmaceutically acceptable dose, a pharmaceutically acceptable amount, a pharmaceutically effective dose, or a pharmaceutically effective amount.

[0069] As used herein, the term “cognitive impairment,” unless otherwise specified, includes at least one of the following: Limited Cognitive Impairment (LCI), Mild Cognitive Impairment (MCI), Alzheimer’s disease (or dementia of an Alzheimer’s-type) or a particular stage of Alzheimer’s disease, inclusive of pre-Alzheimer’s disease, early Alzheimer’s disease, mild Alzheimer’s disease, moderate Alzheimer’s disease, severe Alzheimer’s disease, pre-Alzheimer’s-to-mild Alzheimer’s disease, mild-to-moderate Alzheimer’s disease, moderate-to-severe Alzheimer’s disease, schizophrenia (for example, paranoid type schizophrenia, disorganized type schizophrenia, catatonic type schizophrenia, undifferentiated type schizophrenia), schizophreniform disorder, schizoaffective disorder, delusional disorder, positive symptoms of schizophrenia, negative symptoms of schizophrenia, or schizophrenia with dementia.

[0070] Alzheimer’s disease may include, unless otherwise specified, any of the sub-diagnostic categories used to characterize the type or degree of cognitive impairment in a patient for treatment purposes. A commonly referenced diagnostic scale for characterizing the degree of cognitive impairment for a patient with Alzheimer’s disease includes the 3-stage Alzheimer Disease Model. The 3-stages consist of: mild stage (also referred to as “early Alzheimer’s disease” or “mild Alzheimer’s disease” or “early stage Alzheimer’s disease” or “mild dementia of an Alzheimer’s-type”), moderate stage (also referred to as “middle Alzheimer’s disease” or “moderate Alzheimer’s disease” or “middle stage Alzheimer’s disease” or “moderate dementia of an Alzheimer’s-type”), and severe stage (also referred to as “late Alzheimer’s disease” or “severe Alzheimer’s disease” or “late stage Alzheimer’s disease” or “severe dementia of an Alzheimer’s-type”). For patients with a condition that has not progressed to the point of mild stage Alzheimer’s disease, they may be diagnosed as having pre-Alzheimer’s disease. It is also not uncommon for treatment purposes to characterize stages together, such as pre-Alzheimer’s disease-to-mild stage Alzheimer’s disease, mild-to-moderate Alzheimer’s disease, or moderate-to-severe Alzheimer’s disease. Another useful diagnostic scale that is used in characterizing the degree of cognitive impairment for a patient having Alzheimer’s disease is the Seven Stage Alzheimer’s Disease Model (sometimes known as the “Seven Stage Global Deterioration Scale” or the “Reisberg Scale”). This diagnostic scale divides the progression of the cognitive disorder associated with Alzheimer’s disease as follows: Stage 1-no Alzheimer’s disease (generally characterized by absence of impairment, no impairment, or normal function), Stage 2-pre-Alzheimer’s disease (generally characterized by minimal impairment, normal

forgetfulness, or very mild cognitive decline), Stage 3-early-stage Alzheimer's disease (generally characterized by a noticeable cognitive decline, early confusional/mild cognitive impairment, or mild cognitive decline), Stage 4-early-stage/mild Alzheimer's disease (also referred to as late confusional/mild Alzheimer's, and generally characterized by moderate cognitive decline), Stage 5-middle-stage/moderate Alzheimer's (also referred to as early dementia/moderate Alzheimer's disease and generally characterized by moderately severe cognitive decline), Stage 6-middle dementia/moderately severe Alzheimer's disease (also referred to as middle-stage/moderate to late-stage/severe Alzheimer's disease and generally characterized by severe cognitive decline), and Stage 7-late-stage/severe Alzheimer's disease (also referred to as severe dementia or failure-to-thrive, and generally characterized by very severe cognitive decline). It is also not uncommon for treatment purposes to characterize stages together, such as pre-Alzheimer's disease-to-mild stage Alzheimer's disease, mild-to-moderate Alzheimer's disease, or moderate-to-severe Alzheimer's disease. As used herein, unless otherwise specified, Alzheimer's disease includes all of the above named diagnostic categories or disease characterizations. It is also not uncommon for a physician to categorize any one or more of the above noted states of Alzheimer's disease as being probable, for example, probable mild-to-moderate Alzheimer's disease or probable severe Alzheimer's disease, when their diagnosis does not include, for example a physical biopsy or other definitive analysis.

[0071] Mild Cognitive Impairment (MCI) is considered by some to be an intermediate stage between normal aging and the onset of Alzheimer's disease. For example, MCI may be characterized by persistent forgetfulness, but may lack some or many of the more debilitating symptoms of Alzheimer's disease. Another set of criteria that may characterize a patient as having mild cognitive impairment suitable for treatment includes a patient that meets the following: 1) memory complaints corroborated by an informant, 2) objective memory impairment for age and education, 3) normal general cognitive function, 4) intact activities of daily living, and 5) the patient does not meet criteria for dementia. In general, a patient characterized as having mild cognitive impairment may not yet have a clinical cognitive deficit. Mild cognitive impairment may also be distinguished from senile dementia in that mild cognitive impairment involves a more persistent and troublesome problem of memory loss for the age of the patient. On the clinical diagnostic scale, mild cognitive impairment is followed, in increased severity, by Alzheimer's disease.

[0072] Limited Cognitive Impairment (LCI) describes a cognitive impairment (*i.e.*, symptoms or conditions), which precedes mild cognitive impairment on a clinical diagnostic scale, and includes any chronic or temporary impairment in cognition, learning or memory that prevents or reduces the ability of a patient from achieving their individual potential in these areas. For example, LCIs may include minor impairments to memory associated with focus and concentration (*e.g.*, accuracy and speed of learning and recalling information), working memory (*e.g.*, used in decision making and problem solving), cognition, focus, mental quickness, and mental clarity.

[0073] The term “stereoisomer” refers to a molecule capable of existing in more than one spatial atomic arrangement for a given atomic connectivity (e.g., enantiomers, meso compounds, and diastereomers). As used herein, the term “stereoisomer” means either or both enantiomers and diastereomers.

[0074] The aminobenzisoxazole compounds of the present invention represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, may contain one or more stereogenic centers. Accordingly, compounds of this invention can exist as either individual stereoisomers or mixtures of two or more stereoisomers. A compound of the present invention will include both mixtures (e.g., racemic mixtures) and also individual respective stereoisomers that are substantially free from another possible stereoisomer. The term “substantially free of other stereoisomers” as used herein means less than 25% of other stereoisomers, less than 10% of other stereoisomers, less than 5% of other stereoisomers, less than 2% of other stereoisomers, or less than “X”% of other stereoisomers (wherein X is a number between 0 and 100, inclusive) are present.

[0075] The aminobenzisoxazole compounds of the present invention represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, may contain one or more tautomeric forms. Accordingly, compounds of this invention can exist as either individual tautomers or mixtures of tautomeric forms. A compound of the present invention will include both mixtures (e.g., mixtures of tautomeric forms) and also individual respective tautomers that are substantially free from another possible tautomer.

[0076] The aminobenzisoxazole compounds of the present invention represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, may contain one or more geometric isomers. Accordingly, compounds of this invention can exist as either geometric isomers or mixtures of geometric isomers. A compound of the present invention will include both mixtures (e.g., mixtures of geometric isomers) and also individual respective geometric isomers that are substantially free from another possible geometric isomer.

[0077] The term “haloalkyl” refers to an alkyl group having from 1 to 5 halogen substituents independently selected from -F, -Cl, -Br, and -I. For example, a haloalkyl may represent a -CF₃ group, a -CCl₃ group, a -CH₂CF₃ group, or a -CF₂CF₃ group.

[0078] The term “heteroaryl” refers to an aromatic ring system comprising at least one or more hetero- ring atoms, such as two, three, four, or five hetero- ring atoms, independently selected from N, O, and S. Suitable heteroaryl groups may include a single ring, for example, thienyl, pyridyl, thiazolyl, pyrazinyl, pyrimidyl, imidazolyl, furanyl, isothiazolyl, pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxazolyl, pyrrolyl, pyridazinyl, triazinyl, oxadiazolyl, and furazanlyl. Suitable heteroaryl groups may include a fused ring system, for example, a six-six fused ring system, a six-five fused ring system, or a five-six fused ring system, such as benzothienyl, quinolyl, benzofuranyl, benzothiazolyl, benzisothiazolyl, benzisoxazolyl, benzimidazolyl, indolyl, benzoxazolyl, isoquinolinyl, cinnolinyl,

indazolyl, indoliziny, phthalaziny, isoindolyl, puriny, benzofurazany, benzothiopheny, benzothiazolyl, quinazoliny, quinoxaliny, naphthridiny, and furopyridiny.

[0079] Suitable "heterocycloalkyl" groups include those having at least one or more hetero- ring atoms, such as two or three hetero- ring atoms, independently selected from at least one ring atom selected from the group consisting of oxygen, nitrogen, and sulfur, with the proviso that when the at least one ring atom is nitrogen, the nitrogen is independently substituted with -H, an unbranched C₁-C₄-alkyl radical, a branched C₃-C₄-alkyl radical, a C₃-C₄-cycloalkyl radical, -(CO)-unbranched C₁-C₄-alkyl, -(CO)-branched C₃-C₄-alkyl, -(SO₂)-unbranched C₁-C₄-alkyl, or -(SO₂)-branched C₃-C₄-alkyl, and with the further proviso that when the at least one ring atom is sulfur, the sulfur may be independently substituted with 0 to 2 =O. Suitable heterocycloalkyl groups may include, for example, tetrahydrofurano, tetrahydropyrano, morpholino, pyrrolidino, piperidino, piperazino, azetidino, azetidinono, oxindolo, oxetano, dihydroimidazolo, and pyrrolidinono.

[0080] The pharmaceutically acceptable salt of the aminobenzisoxazole compounds represented by Formula (Ia) or Formula (Ib), according to the present invention may be acid addition salts with inorganic or organic acids. Specific examples of these salts include acid addition salts with, for instance, mineral acids such as hydrochloric acid, hydrobromic acid, hydriodic acid, sulfuric acid, nitric acid or phosphoric acid; organic acids, for example carboxylic acids or sulfonic acids, such as formic acid, acetic acid, propionic acid, oxalic acid, malonic acid, succinic acid, fumaric acid, maleic acid, lactic acid, malic acid, tartaric acid, citric acid, benzoic acid, p-toluenesulfonic acid, benzenesulfonic acid, naphthalenedisulfonic acid, isethionic acid, glucuronic acid, gluconic acid, methanesulfonic acid or ethanesulfonic acid; or acidic amino acids such as aspartic acid or glutamic acid.

[0081] In certain embodiments, a pharmaceutical composition may comprise an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0082] In certain embodiments, the aminobenzisoxazole compounds represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, are suitable for use as medicaments for the treatment and/or prophylaxis of diseases in humans and/or animals.

[0083] In certain embodiments, the invention relates to a method comprising administering to a patient in need thereof an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0084] In certain embodiments, the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, act as ligands, in particular as α 7-nAChR agonists.

[0085] In certain embodiments, a method of treating a patient in need thereof, comprising administering an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof. In certain embodiments, a method of treating a patient in need thereof, comprising administering a pharmaceutical composition comprising an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof. For example, the patient may suffer from a cognitive impairment or suffers from one or more symptoms associated with a cognitive impairment, such as Limited Cognitive Impairment (LCI), Mild Cognitive Impairment (MCI), Alzheimer's disease, dementia of an Alzheimer's-type, schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, positive symptoms of schizophrenia, negative symptoms of schizophrenia, or schizophrenia with dementia.

[0086] In certain embodiments, the aminobenzisoxazole compounds represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, can, because of their pharmacological properties, be employed alone or in combination with other active ingredients for the treatment and/or prevention of cognitive impairments, for example, Alzheimer's disease or schizophrenia. Because of their selective effect as $\alpha 7$ -nAChR agonists, the aminobenzisoxazole compounds represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, are particularly suitable for improving cognition, providing procognitive effects, improving perception, improving concentration, improving learning or memory, improving one or more aspects of cognition, e.g., one or more of: executive function, memory (e.g., working memory), social cognition, visual learning, verbal learning and speed of processing, especially after or associated with cognitive impairments like those occurring for example in situations/diseases/syndromes such as mild cognitive impairment, age-associated learning and memory impairments, age-associated memory loss, vascular dementia, craniocerebral trauma, stroke, dementia occurring after strokes (post-stroke dementia), post-traumatic brain syndrome, general concentration impairments, concentration impairments in children with learning and memory problems, attention deficit hyperactivity disorder, Alzheimer's disease, Lewy body dementia, dementia with degeneration of the frontal lobes, including Pick's syndrome, Parkinson's disease, dyskinesias associated with dopamine agonist therapy in Parkinson's Disease, progressive nuclear palsy, dementia with corticobasal degeneration, amyotrophic lateral sclerosis (ALS), Huntington's disease, multiple sclerosis, thalamic degeneration, Creutzfeld-Jakob dementia, HIV dementia, schizophrenia (e.g., paranoid type, disorganized type, catatonic type, and undifferentiated type), schizophreniform disorder, schizoaffective disorder, delusional disorder, positive symptoms of schizophrenia, negative symptoms of schizophrenia, schizophrenia with dementia, Korsakoff's psychosis, depression, anxiety, mood and affective disorders, traumatic brain injury, withdrawal symptoms associated with smoking cessation and dependent drug cessation, Gilles de la Tourette's Syndrome, age-related macular degeneration, glaucoma, neurodegeneration associated with glaucoma, treatment (including amelioration, prevention or delay of progression) of sleep

disorders (e.g., narcolepsy, excessive daytime sleepiness, nocturnal sleep disruption and/or cataplexy), treatment (including amelioration, prevention or delay) of progression of fatigue, or use for facilitation of emergence from general anesthesia.

[0087] In certain embodiments, the aminobenzisoxazole compounds represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, can be employed alone or in combination with other active ingredients for the prophylaxis and treatment of acute and/or chronic pain (for a classification, see “Classification of Chronic Pain, Descriptions of Chronic Pain Syndromes and Definitions of Pain Terms”, 2nd edition, Meskey and Begduk, editors; IASP Press, Seattle, 1994), especially for the treatment of cancer-induced pain and chronic neuropathic pain like, for example, that associated with diabetic neuropathy, postherpetic neuralgia, peripheral nerve damage, central pain (for example as a consequence of cerebral ischaemia) and trigeminal neuralgia, and other chronic pain such as, for example, lumbago, backache, or rheumatic pain. In addition, these active ingredients are also suitable for the therapy of primary acute pain of any origin and of secondary states of pain resulting therefrom, and for the therapy of states of pain which were formerly acute and have become chronic.

[0088] In certain embodiments, the invention relates to a method comprising administering to a patient in need thereof, such as a patient suffering from, or diagnosed as having, a cognitive impairment or having one or more symptoms associated with a cognitive impairment, an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent. For example, the method may treat and/or improve the one or more symptoms associated with a cognitive impairment and/or the cognitive impairment.

[0089] A certain embodiment of the present invention provides a method of improving one or more cognitive symptoms, improving one or more behavioral symptoms, or both, associated with a cognitive impairment, comprising: administering to a patient in need thereof an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0090] In a certain embodiment of the present invention, the method provides a pro-cognitive effect in a patient suffering from, or diagnosed as having, a cognitive disease or dementia, comprising: administering to a patient in need thereof an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of the

aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent; wherein the method provides at least one of the following: visual motor, learning, delayed memory, or executive function; for example provides a pro-cognitive effect, exclusive of attention, in said patient; for example provides a pro-cognitive effect in at least one of the following: visual motor, learning, delayed memory, or executive function.

[0091] A certain embodiment of the present invention provides a method of treating a patient with a cognitive disease, comprising: administering to the patient a daily dose of a pharmaceutical composition comprising an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0092] In a certain embodiment of the present invention, the method provides a pro-cognitive effect in a patient suffering from, or diagnosed as having, schizophrenia, for example, paranoid type schizophrenia, disorganized type schizophrenia, catatonic type schizophrenia, undifferentiated type schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, positive symptoms of schizophrenia, negative symptoms of schizophrenia, or schizophrenia with dementia, comprising: administering to a patient in need thereof an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to a patient in need thereof, a pharmaceutical composition comprising an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluents; wherein the method provides at least one of the following: visual motor, learning, delayed memory, or executive function; for example provides a pro-cognitive effect, exclusive of attention, in said patient; for example provides a pro-cognitive effect in at least one of the following: visual motor, learning, delayed memory, or executive function.

[0093] In an embodiment of the present invention, any one of the above-noted embodiments, includes wherein the daily dose is an initial daily dose.

[0094] In a certain embodiment of the present invention provides a method of improving cognition of a patient in need thereof, comprising: administering to the patient an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluents.

[0095] In a certain embodiment of the present invention provides a method of treating or improving one or more symptoms associated with a cognitive disease and/or a cognitive impairment in a patient in need thereof, comprising: administering to the patient an effective dose of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically

acceptable salt thereof; or administering to the patient a pharmaceutical composition comprising the aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier, excipient or diluent.

[0096] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes treating a symptom associated with a cognitive disease.

[0097] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes improving a symptom associated with a cognitive disease.

[0098] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes preventing progression of a cognitive disease.

[0099] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the patient has been diagnosed as having a cognitive disease.

[00100] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the patient has been diagnosed as having Alzheimer's disease.

[00101] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes treating a symptom associated with Alzheimer's disease.

[00102] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes improving a symptom associated with Alzheimer's disease.

[00103] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes preventing progression of Alzheimer's disease.

[00104] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the patient has been diagnosed as having mild-to-moderate Alzheimer's disease.

[00105] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes treating a symptom associated with schizophrenia.

[00106] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes improving a symptom associated with schizophrenia.

[00107] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes preventing progression of schizophrenia.

[00108] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the patient has been diagnosed as having schizophrenia.

[00109] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes treating a symptom associated with positive symptoms of schizophrenia.

[00110] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes improving a symptom associated with positive symptoms of schizophrenia.

[00111] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes preventing progression of positive symptoms of schizophrenia.

[00112] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes the patient has been diagnosed as having positive symptoms of schizophrenia.

[00113] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes treating a symptom associated with negative symptoms of schizophrenia.

[00114] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes improving a symptom associated with negative symptoms of schizophrenia.

[00115] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes preventing progression of negative symptoms of schizophrenia.

[00116] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes the patient has been diagnosed as having negative symptoms of schizophrenia.

[00117] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes treating a symptom associated with schizophrenia with dementia.

[00118] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes improving a symptom associated with schizophrenia with dementia.

[00119] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes preventing progression of schizophrenia with dementia.

[00120] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes the patient has been diagnosed as having schizophrenia with dementia.

[00121] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the method specifically includes the patient has been diagnosed as having a disease associated with chronic inflammation, including atherosclerosis, rheumatoid arthritis and inflammatory bowel diseases.

[00122] In an embodiment of the present invention, any one of the above-noted embodiments, wherein the pharmaceutical composition is in the form of a tablet.

[00123] **Pharmaceutical Compositions**

[00124] In certain embodiments, the invention also includes pharmaceutical preparations which, besides inert, nontoxic, pharmaceutically suitable excipients, adjuvants and carriers, contain one or more aminobenzisoxazole compounds represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, or consist of one or more aminobenzisoxazole compounds represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, and processes for producing these preparations.

[00125] An aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, may be formulated for administration in solid or liquid form. For example, an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, may be formulated for administration in a capsule, a tablet, or a powder form. For example, an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, may be formulated alone or as part of a pharmaceutical composition, suitable for oral administration, such as in a capsule or tablet, intravenous administration, parenteral administration, topical administration, or transdermal administration, such as in a patch, to a patient in need thereof.

[00126] An aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, may be administered as a pharmaceutical composition, for example, in the presence of carriers, adjuvants, excipients, diluents, fillers, buffers, stabilizers, preservatives, lubricants, and the like, for example, administered as a pharmaceutical composition (e.g., formulation) comprising at least an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, together with one or more pharmaceutically acceptable carriers, adjuvants, excipients, diluents, or other materials well known to those skilled in the art. As used herein, the term “pharmaceutically acceptable”, unless otherwise specified, includes the generally accepted meaning which encompasses combinations, compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for consumption by humans without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[00127] Suitable pharmaceutically acceptable carriers, adjuvants, excipients, and diluents, can include, but are not limited to, lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum, acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water syrup, methyl cellulose, methyl and propyl hydroxybenzoates, talc, magnesium stearate, and mineral oil. Pharmaceutically acceptable carriers, adjuvants and vehicles that may be used in the pharmaceutical compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts,

colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[00128] The formulations can additionally include, but are not limited to, pharmaceutically acceptable lubricating agents, glidants, wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents, and/or flavoring agents. The pharmaceutical compositions of the present invention may be formulated so as to provide quick release, immediate release, sustained release, or delayed release of an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, after administration to the patient by employing procedures well-known in the art.

[00129] Another embodiment of the invention further comprises methods of making Pharmaceutical Composition, comprising admixing at least an aminobenzisoxazole compound represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, together with one or more pharmaceutically acceptable carriers, excipients, buffers, adjuvants, stabilizers, or other materials.

[00130] In certain embodiments, the aminobenzisoxazole compounds represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, are to be present in these preparations in a concentration of from 0.1 to 99.5% by weight, preferably from 0.5 to 95% by weight, of the complete mixture. Besides the aminobenzisoxazole compounds represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, the pharmaceutical preparations may also contain other active pharmaceutical ingredients.

[00131] In certain embodiments, the novel active ingredients can be converted in a known manner into conventional formulations such as tablets, coated tablets, pills, granules, aerosols, syrups, emulsions, suspensions and solutions, using inert, nontoxic, pharmaceutically suitable excipients or solvents. In these cases, the therapeutically active compound should in each case be present in a concentration of about 0.5 to 90% by weight of the entire mixture, i.e., in amounts which are sufficient to reach the stated dose range.

[00132] In certain embodiments, the formulations are produced, for example, by extending the active ingredients with solvents and/or excipients, where appropriate with use of emulsifiers and/or dispersants, it being possible for example when water is used as diluent where appropriate to use organic solvents as auxiliary solvents.

[00133] In certain embodiments, administration may take place in a conventional way, for example, orally, transdermally or parenterally, especially perlingually or intravenously. In certain embodiments, administration may also take place by inhalation through the mouth or nose, for example, with the aid of a spray, or topically via the skin.

[00134] In certain embodiments, the aminobenzisoxazole compounds represented by Formula (Ia) or Formula (Ib), or a pharmaceutically acceptable salt thereof, may be administered in amounts of

about 0.01 to 10 mg/kg, on oral administration, for example, about 0.05 to 5 mg/kg, of body weight to achieve effective results.

[00135] EXAMPLES

[00136] Analytical instrument model:

Table 1

LCMS	Shimadzu UFLC MS: LCMS-2020 Agilent Technologies 1200 series MS: Agilent Technologies 6110 Agilent Technologies 1200 series MS: LC/MSD VL Agilent Technologies 1100 Agilent Technologies 1260
NMR	BRUKER ADVANCE III/400 (400 MHz) BRUKER ADVANCE 400 (400MHz) BRUKER DMX300 (300MHz)
Prep-HPLC	Gilson GX-281 systems: instruments GX-A, GX-B, GX-C, GX-D, GX-E, GX-F, GX-G and GX-H
GCMS	SHIMADZU GCMS-QP2010 Ultra
Analytical cSFC	Agilent Technologies 1290 Infinity
Prep-cSFC	Waters SFC Prep 80

[00137] LCMS:

[00138] LCMS Conditions A ("LCMS (A)"): Instrument: Shimadzu LCMS 2020; Mobile phase A: 4L H₂O \ 1.5 mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 10-80AB_4MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 10%-80%; Column: Boston Green ODS 2.1x30 mm, 3 μm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00139] LCMS Conditions B ("LCMS (B)"): Instrument: Agilent 1200 Series; Mobile phase A: 4L H₂O \ 1.5 ml TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 5-95AB_R_2W; Flow Rate: 1.5 mL/min.; Gradient: 5%-95%; Column: Chromolith@Flash RP-18e 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00140] LCMS Conditions C ("LCMS (C)"): Instrument: Agilent 1200 Series; Mobile phase A: 4L H₂O \ 2 mL NH₃H₂O; Mobile phase B: Acetonitrile; Method name: 5-95CD_4.5MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 5%-95%; Column: Chromolith@Flash RP-18e 25x2 mm; Column temperature 50 °C; Wavelength: 220 nm & 254 nm.

[00141] LCMS Conditions D ("LCMS (D)"): Instrument: Agilent 1200 Series; Mobile phase A: 4L H₂O \ 1.5 mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 5-

95AB_R_4MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 5%-95%; Column: Chromolith@Flash RP-18e 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00142] LCMS Conditions E ("LCMS (E)"): Instrument: Agilent 1200 Series; Mobile phase A: 4L H₂O \ 1.5 mL TFA, Mobile phase B: 4L ACN\0.75 mL TFA; Method name: 5-95AB_R; Flow Rate: 1.5 mL/min. ; Gradient: 5%-95%; Column: Chromolith@Flash RP-18e 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00143] LCMS Conditions F ("LCMS (F)"): Instrument: Agilent 1200 Series; Mobile phase A: 4L H₂O \ 2 mL NH₃H₂O, Mobile phase B: Acetonitrile; Method name: 5-95CD_2MIN_2W; Flow Rate: 1.2 mL/min.; Gradient: 5%-95%; Column: XBridge Shield RP-18 2.1x50 mm, 5 µm; Column temperature: 30 °C; Wavelength: 220 nm & 254 nm.

[00144] LCMS Conditions G ("LCMS (G)"): Instrument: Agilent 1200 Series; Mobile phase A: 4L H₂O \ 2 mL NH₃H₂O, Mobile phase B: Acetonitrile; Method name: 10-80CD_4MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 10%-80%; Column: XBridge C-18 2.1x50 mm, 5µm; Column temperature: 40 °C; Wavelength: 220 nm & 254 nm.

[00145] LCMS Conditions H ("LCMS (H)"): Instrument: Agilent 1200 Series; Mobile phase A: 4L H₂O \ 1.5 mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 10-80AB_4MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 10%-80%; Column: Xtimate C-18, 2.1x30 mm, 3µm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00146] LCMS Conditions I ("LCMS (I)"): Instrument: Agilent 1200 Series; Mobile phase A: 4L H₂O \ 2 mL NH₃H₂O, Mobile phase B: Acetonitrile; Method name: 0-60CD_4.5MIN_2W; Flow Rate: 0.8 ml/min.; Gradient: 0%-60%; Column: XBridge Shield RP-18 2.1x50 mm, 5µm; Column temperature 50 °C; Wavelength: 220 nm & 254 nm.

[00147] LCMS Conditions J ("LCMS (J)"): Instrument: Agilent 1200 Series; Mobile phase A: 4L H₂O \ 2mL NH₃H₂O, Mobile phase B: Acetonitrile; Method name: 10-80CD_2MIN_POS_2W; Flow Rate: 1.2ml/min.; Gradient: 10%-80%; Column: Xbridge C-18 2.1x50 mm, 5µm; Column temperature: 40 °C; Wavelength: 220 nm & 254 nm.

[00148] LCMS Conditions K ("LCMS (K)"): Instrument: Shimadzu LCMS 2020; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 0-30AB_2MIN_2W; Flow Rate: 1.2 mL/min.; Gradient: 0%-30%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00149] LCMS Conditions L ("LCMS (L)"): Instrument: Shimadzu LCMS 2020; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 0-30AB_4MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 0%-30%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00150] LCMS Conditions M ("LCMS (M)"): Instrument: Shimadzu LCMS 2020; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 0-

60AB_2MIN_2W; Flow Rate: 1.2 mL/min.; Gradient: 0%-60%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00151] LCMS Conditions N ("LCMS (N)"): Instrument: Shimadzu LCMS 2020; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 0-60AB_4MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 0%-60%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00152] LCMS Conditions O ("LCMS (O)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 2mL NH₃H₂O, Mobile phase B: CAN; Method name: 0-30CD_2MIN_POS_2W; Flow Rate: 1.0 mL/min.; Gradient: 0%-30%; Column: Xbridge C18 2.1x50 mm, 5µm; Column temperature: 40 °C; Wavelength: 220 nm & 254 nm.

[00153] LCMS Conditions P ("LCMS (P)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 2mL NH₃H₂O, Mobile phase B: CAN; Method name: 0-60CD_2MIN_POS_2W; Flow Rate: 1.0 mL/min.; Gradient: 0%-60%; Column: Xbridge C18 2.1x50 mm, 5µm; Column temperature: 40 °C; Wavelength: 220 nm & 254 nm.

[00154] LCMS Conditions Q ("LCMS (Q)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 2mL NH₃H₂O, Mobile phase B: CAN; Method name: 0-60CD_4MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 0%-60%; Column: Xbridge C18 2.1x50 mm, 5µm; Column temperature: 40 °C; Wavelength: 220 nm & 254 nm.

[00155] LCMS Conditions R ("LCMS (R)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 10-80AB_2MIN_2W; Flow Rate: 1.2 mL/min.; Gradient: 10%-80%; Column: Xtimate C18, 2.1x30mm, 3µm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00156] LCMS Conditions S ("LCMS (S)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 2mL NH₃H₂O, Mobile phase B: CAN; Method name: 30-90CD_4MIN_POS_2W; Flow Rate: 0.8 mL/min.; Gradient: 30%-90%; Column: Xbridge C18 2.1x50 mm, 5µm; Column temperature: 40 °C; Wavelength: 220 nm & 254 nm.

[00157] LCMS Conditions T ("LCMS (T)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 5-95AB_15MIN_YMC; Flow Rate: 1.0 mL/min.; Gradient: 5%-95%; Column: YMC-Pack ODS-A 5µm 150x4.6mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00158] LCMS Conditions U ("LCMS (U)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 0-30AB_2MIN_2W; Flow Rate: 1.2 mL/min.; Gradient: 0%-30%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00159] LCMS Conditions V ("LCMS (V)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 0-

30AB_4MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 0%-30%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00160] LCMS Conditions W ("LCMS (W)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 0-60AB_2MIN_2W; Flow Rate: 1.2 mL/min.; Gradient: 0%-60%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00161] LCMS Conditions X ("LCMS (X)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 0-60AB_4MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 0%-60%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00162] LCMS Conditions Y ("LCMS (Y)"): Instrument: Shimadzu LCMS 2020; Mobile phase A: 4L H₂O \ 1.5 ml TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 5-95AB_R_2W; Flow Rate: 1.5 mL/min.; Gradient: 5%-95%; Column: Chromolith@Flash RP-18e 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00163] LCMS Conditions Z ("LCMS (Z)"): Instrument: Shimadzu LCMS 2020; Mobile phase A: 4L H₂O \ 1.5 mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 5-95AB_R_4MIN_2W; Flow Rate: 0.8 mL/min.; Gradient: 5%-95%; Column: Chromolith@Flash RP-18e 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00164] LCMS Conditions AA ("LCMS (AA)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 2mL NH₃ H₂O, Mobile phase B: ACN; Method name: 10-80CD_2MIN_NEG; Flow Rate: 1.2 mL/min.; Gradient: 10%-80%; Column: Xbridge C18 2.1x50 mm, 5µm; Column temperature: 40 °C; Wavelength: 220 nm & 254 nm.

[00165] LCMS Conditions BB ("LCMS (BB)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 0-60AB_R_2W; Flow Rate: 1.5 mL/min.; Gradient: 0%-60%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00166] LCMS Conditions CC ("LCMS (CC)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 0-30AB_R_2W; Flow Rate: 1.5 mL/min.; Gradient: 0%-30%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00167] LCMS Conditions DD ("LCMS (DD)"): Instrument: Agilent 1200 Series LCMS; Mobile phase A: 4L H₂O \ 1.5mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: 10-80AB_R_2W; Flow Rate: 1.5 mL/min.; Gradient: 10%-80%; Column: Chromolith@Flash RP-18E 25x2 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00168] LCMS Conditions EE ("LCMS (EE)"): Instrument: Agilent 1200 Series; Mobile phase A: 1L H₂O \ 0.375mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: WUXIAB00;

Flow Rate: 0.6 -1.0mL/min; Gradient: 0%-80%-100%; Column: Agilent 5 TC-C18 50x2.1 mm;
Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00169] LCMS Conditions FF ("LCMS (FF)"): Instrument: Agilent 1200 Series; Mobile phase A: 1L H₂O \ 0.375mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: WUXIAB01;
Flow Rate: 0.8 -1.0mL/min; Gradient: 1%-90%-100%; Column: Agilent 5 TC-C18 50x2.1 mm;
Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00170] LCMS Conditions GG ("LCMS (GG)"): Instrument: Agilent 1200 Series; Mobile phase A: 1L H₂O \ 0.375mL TFA, Mobile phase B: 4L ACN \ 0.75 mL TFA; Method name: WUXIAB10;
Flow Rate: 0.8 -1.0mL/min; Gradient: 10%-100%; Column: Agilent 5 TC-C18 50x2.1 mm; Column temperature: 50 °C; Wavelength: 220 nm & 254 nm.

[00171] LCMS Conditions 1 ("LCMS (1)"): Instrument: Agilent 1100 Bin. Pump: G1312A, degasser; autosampler, ColCom, DAD: Agilent G1315B, 210 nm, MSD: Agilent LC/MSD G1956B ESI, pos/neg 100-800; MS parameters: Source: ESI, Capillary voltage: 3000V, Drying gas flow: 12 L/min., Nebulizer Pressure 60 psig, Drying Gas Temperature: 350°C, Fragmentor 70, MS scan: MS range 100-800 (positive and negative mode), Flow into MS 0.4 mL/min.; Mobile phase A: 95% acetonitrile + 5% 10 mM ammonium bicarbonate in water; Mobile phase B: 10 mM ammonium bicarbonate in water pH = 9.0; Flow Rate: 0.8 mL/min; Linear Gradient: t=0 min 5% A, t = 3.5 min 98% A, t=6 min 98% A; Column: Phenomenex Gemini NX (C18, 50×2.0 mm, particle size: 3 μm); Column temperature: 25°C; Detection DAD: Wavelength 220-320 nm.

[00172] LCMS Conditions 2 ("LCMS (2)"): Instrument Apparatus: Agilent 1260 Bin. Pump: G1312B, degasser; autosampler, ColCom, DAD: Agilent G1315D, 220-320 nm, MSD: Agilent LC/MSD G6130B ESI, pos/neg 100-800, ELSD Alltech 3300 gas flow 1.5 mL/min., Gas Temperature: 40°C; MS parameters: Source: ESI, Capillary voltage: 3000V, Drying gas flow: 12 L/min., Nebulizer Pressure 60 psig, Drying Gas Temperature: 350°C, Fragmentor 70, MS scan: MS range 100-800 (positive and negative mode), Flow into MS 0.4 mL/min.; Mobile phase A: 0.1% formic acid in acetonitrile; Mobile phase B: 0.1% formic acid in water; Flow Rate: 1 mL/min; Linear gradient: t=0 min 5% A, t=1.6min 98% A, t=3 min 98% A; Column: Waters XSelect (C18, 30x2.1 mm, particle size 3.5μm); Column temperature: 35°C; Detection DAD: Wavelength 220-320 nm.

[00173] GCMS:


[00174] GCMS Conditions Instrument: SHIMADZU GCMS-QP2010 Ultra; Carrier gas: He; Column Flow: 1.5mL/min; Injector: 250 °C; Split Ratio:100:1; Column: HP-5MS 15mx0.25mmx0.25um; FILM From: 40 °C (holding 3min) to 250 °C (holding 3min) at the rate of 25°C/min.

[00175] cSFC Analytical:

[00176] cSFC Analytical Conditions: Flow rate: 3mL/min; Wavelength: 220 nm; and Column temperature: 35°C, were used for each of the specified conditions below:

- [00177] cSFC Analytical Conditions A ("cSFC analytical (A)"): Column: Chiralpak OD-3 100×4.6mm I.D., 3μm; Mobile phase: ethanol (0.05% diethylamine ("DEA")) in CO₂ from 5% to 40%.
- [00178] cSFC Analytical Conditions B ("cSFC analytical (B)"): Column: Chiralpak OD-3 100×4.6mm I.D., 3μm; Mobile phase: methanol (0.05% DEA) in CO₂ from 5% to 40%.
- [00179] cSFC Analytical Conditions C ("cSFC analytical (C)"): Column: Chiralpak OD-3 100×4.6mm I.D., 3μm; Mobile phase: 40% ethanol (0.05% DEA) in CO₂.
- [00180] cSFC Analytical Conditions D ("cSFC analytical (D)"): Column: Chiralpak AY-3 100×4.6mm I.D., 3μm; Mobile phase: ethanol (0.05% DEA) in CO₂ from 5% to 40%.
- [00181] cSFC Analytical Conditions E ("cSFC analytical (E)"): Column: Chiralpak OJ-3 100×4.6mm I.D., 3μm; Mobile phase: ethanol (0.05% DEA) in CO₂ from 5% to 40%.
- [00182] cSFC Analytical Conditions F ("cSFC analytical (F)"): Column: Chiralpak OJ-3 100×4.6mm I.D., 3μm; Mobile phase: methanol (0.05% DEA) in CO₂ from 5% to 40%.
- [00183] cSFC Analytical Conditions G ("cSFC analytical (G)"): Column: Chiralpak AD-3 100×4.6mm I.D., 3μm; Mobile phase: ethanol (0.05% DEA) in CO₂ from 5% to 40%.
- [00184] cSFC Analytical Conditions H ("cSFC analytical (H)"): Column: Chiralpak AD-3 100×4.6mm I.D., 3μm; Mobile phase: methanol (0.05% DEA) in CO₂ from 5% to 40%.
- [00185] For each final compound prepared below that indicates the presence of a salt associated with the final compound (i.e., a salt complex), the specific molar equivalence of salt included in the final compound, unless specified, was not determined.


[00186] **General Procedure A1: Synthesis of N-hydroxyimidoyl chloride.**



- [00187] A mixture of aldehyde (1 eq.), hydroxylamine hydrochloride (1.3-2 eq.) and triethylamine (2 eq.) in dichloromethane (1.2-2.5 mL/mmol aldehyde) was stirred at room temperature for 16 hours. On completion, the reaction mixture was diluted with water and extracted with dichloromethane (3 × 20 mL). The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate and concentrated in vacuo to give the oxime intermediate. This intermediate was either purified by silica gel chromatography or used without further purification in the next step.
- [00188] To a solution of oxime intermediate (1 eq.) in dichloromethane (10 mL) at 0 °C was added a solution of *N*-chlorosuccinimide (1.2 eq.) in *N,N*-dimethylformamide (0.5 mL). The mixture was stirred at 30 °C for 1 hour. On completion, the reaction mixture was diluted with water and extracted with dichloromethane (3 × 10 mL). The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate, filtered and concentrated in vacuo to give N-

hydroxyimidoyl chloride intermediate, which was used crude in the next reaction without further purification.


[00189] General Procedure A2: Synthesis of *N*-hydroxyimidoyl chloride.




[00190] To a solution of aldehyde in ethanol/water (8/1, v/v) at room temperature was added hydroxylamine hydrochloride (2 eq.) and sodium acetate (3 eq.). The reaction was stirred for 1-2 hour until TLC showed the reaction was complete. The mixture was concentrated in vacuo, and the residue was triturated from water, collected by filtration, washed with water and dried in vacuo to afford the oxime product, which was used as such in the next step.

[00191] To a solution of oxime in *N,N*-dimethylformamide at room temperature was added *N*-chlorosuccinimide (1 eq.). The reaction was stirred for 1 or more hours until TLC showed the reaction was complete. The solution was diluted with ethyl acetate and water and filtered through Celite to remove particles. The layers were separated, and the organic layer was washed with water and brine (2x), dried with sodium sulfate, filtered, and concentrated in vacuo to afford the *N*-hydroxybenzimidoyl chloride product, which was used as such in the next step.


[00192] Example 1A: 7-chlorobenzo[d]isoxazol-3-amine (A-1)




[00193] To a solution of *N*-hydroxyacetamide (3.6 g, 48 mmol) in dry *N,N*-dimethylformamide (60 mL) at room temperature was added potassium *t*-butoxide (5.4 g, 48 mmol). After stirring for 30 minutes, 3-chloro-2-fluorobenzonitrile (5.0 g, 32 mmol) was added, and stirring was continued for another 4.5 hours. On completion, the reaction mixture was poured into a mixture of brine (60 mL) and ethyl acetate (60 mL). The organic phase was separated, dried over anhydrous sodium sulfate, filtered and concentrated to dryness. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 3:1] to afford compound **A-1** (3.9 g, 73% yield) as a pale yellow solid. ¹H-NMR (DMSO-*d*₆, 400 MHz): δ 7.82-7.80 (d, *J*=7.6 Hz, 1H), 7.65-7.63 (d, *J*=7.2 Hz, 1H), 7.30-7.26 (m, 1H), 6.61 (s, 2H).

[00194] Example 2A: 6-chlorobenzo[d]isoxazol-3-amine (A-2)

[00195] To a solution of *N*-hydroxyacetamide (3.1 g, 41 mmol) in dry *N,N*-dimethylformamide (60 mL) at room temperature was added potassium *t*-butoxide (4.6 g, 41 mmol). After stirring for 30 minutes, 4-chloro-2-nitrobenzonitrile (5.0 g, 27 mmol) was added, and stirring was continued for another 4.5 hours. On completion, the reaction mixture was poured into a mixture of brine (60 mL) and ethyl acetate (60 mL). The organic phase was separated, dried over anhydrous sodium sulfate, filtered and concentrated to dryness. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 3:1] to afford compound **A-2** (3.1 g, 66% yield) as a pale yellow solid. ¹H-NMR (DMSO-*d*₆, 400 MHz): δ 7.84-7.82 (d, *J*=8.8 Hz, 1H), 7.65-7.64 (d, *J*=1.2 Hz, 1H), 7.33-7.31 (dd, *J*=1.2 Hz, *J*=8.8 Hz, 1H), 6.52 (s, 2H).

[00196] Example 3A: 5-chlorobenzo[d]isoxazol-3-amine (A-3)


[00197] To a solution of *N*-hydroxyacetamide (2.2 g, 29 mmol) in dry *N,N*-dimethylformamide (30 mL) at room temperature was added potassium *t*-butoxide (3.2 g, 29 mmol). After stirring for 30 minutes, 5-chloro-2-fluorobenzonitrile (3.0 g, 19 mmol) was added, and stirring was continued for another 4.5 hours. On completion, the reaction mixture was quenched with water and extracted with ethyl acetate (3 × 20 mL). The combined organic layers were washed with water and brine (5 × 50 mL), dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 1:1] to afford compound **A-3** (1.5 g, 31% yield) as a white solid.

[00198] Example 4A: 6-cyclopropylbenzo[d]isoxazol-3-amine (A-4)

[00199] To a solution of 6-bromobenzo[d]isoxazol-3-amine (1.0 g, 4.7 mmol) and cyclopropylboronic acid (0.81 g, 9.4 mmol) in a mixture of toluene (10 mL) and water (1.0 mL) under nitrogen at room temperature were added potassium phosphate (1.0 g, 9.4 mmol), palladium acetate (53 mg, 0.2 mmol) and tricyclohexylphosphine (0.11 g, 0.4 mmol). The resulting mixture was stirred at 100 °C for 6 hours, then diluted with water (10 mL) and extracted with ethyl acetate (3 × 10 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated in vacuo. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 5:1]


to afford compound **A-4** (0.59 g, 72% yield) as a off-white solid. LCMS (B): t_R =0.684 min., (ES⁺) m/z (M+H)⁺ = 175.1. ¹H-NMR (CD₃OD, 400 MHz): δ 7.40-7.38 (d, J=8.0 Hz, 1H), 7.09 (s, 1H), 6.99-6.97 (d, J=8.4 Hz, 1H), 4.37 (s, 2H), 2.05-2.01 (m, 1H), 1.10-1.05 (m, 2H), 0.81-0.77 (m, 2H).

[00200] Example 5A: 6-methoxybenzo[d]isoxazol-3-amine (A-5)




[00201] To a solution of *N*-hydroxyacetamide (3.2 g, 42 mmol) in dry *N,N*-dimethylformamide (50 mL) at room temperature was added potassium *t*-butoxide (6.6 g, 59 mmol). After stirring for 30 minutes, 4-methoxy-2-nitrobenzonitrile (3.0 g, 17 mmol) was added, and stirring was continued for another 16 hours. On completion, the reaction mixture was quenched with water and extracted with ethyl acetate (3 × 100 mL). The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate, concentrated in vacuo and purified by silica gel chromatography [petroleum ether: ethyl acetate = 1:1] to give compound **A-5** (2.0 g, 72% yield) as a yellow solid. LCMS (J): (ES⁺) m/z (M+H)⁺ = 165.1, t_R = 1.164 min.

[00202] Example 6A: 3,4-dichloro-2-fluorobenzaldehyde oxime (A-6)




[00203] A mixture of 3,4-dichloro-2-fluorobenzaldehyde (2.0 g, 9.5 mmol), hydroxylamine hydrochloride (0.98 g, 14 mmol) and triethylamine (2.0 mL, 14 mmol) in dichloromethane (10 mL) was stirred at room temperature for 12 hours. On completion, the reaction mixture was diluted with water and extracted with dichloromethane (3 × 20 mL). The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate and concentrated in vacuo to give compound **A-6** (1.5 g, white solid, 69% yield). LCMS (B): (ES⁺) m/z (M+H)⁺ = 208.0. t_R =0.81 min.

[00204] Example 7A: 3,4-dichloro-2-fluorobenzonitrile (A-7)



[00205] To a solution of compound **A-6** (1.5 g, 7.2 mmol) in toluene (50 mL) was added slowly thionyl chloride (0.90 g, 7.6 mmol). The mixture was stirred at 120 °C for 2 hours, then concentrated in vacuo. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1] to give compound **A-7** (0.80 g, 58% yield) as a white solid. ¹H-NMR (CD₃OD, 400 MHz): δ 7.81-7.73 (m, 1H), 7.62-7.60 (m, 1H).

[00206] Example 8A: 6,7-dichlorobenzo[d]isoxazol-3-amine (A-8)

[00207] To a solution of *N*-hydroxyacetamide (1.4 g, 19 mmol) in dry *N,N*-dimethylformamide (20 mL) at room temperature was added potassium *t*-butoxide (2.1 g, 19 mmol). After stirring for 0.5 hour, compound **A-7** was added, and stirring was continued for another 12 hours. On completion the reaction mixture was poured into brine and extracted with ethyl acetate (2× 25 mL). The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate and concentrated in vacuo. The resulting solid was recrystallized from dichloromethane / petroleum ether to give compound **A-8** (1.0 g, 50% yield) as a white solid. LCMS (B): (ES⁺) *m/z* (M+H)⁺ = 203.1. *t*R=0.73 min.

[00208] Example 9A: 2-chloro-6-fluoro-N-hydroxybenzimidoyl chloride (A-10)

[00209] Following general procedure A1, compound **A-10** was prepared from 2-chloro-6-fluorobenzaldehyde:

[00210] Compound **A-9** (790 mg, white solid, 72% yield) was prepared from 2-chloro-6-fluorobenzaldehyde (1.0 g, 6.37 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1].

[00211] Compound **A-10** (0.3 g, white solid, crude) was prepared from compound **A-9** (0.4 g, 2.3 mmol). TLC [petroleum ether: ethyl acetate = 5:1]: *R*_f = 0.7.


[00212] Example 10A: 2,4-difluoro-N-hydroxybenzimidoyl chloride (A-12)

[00213] Following general procedure A1, compound **A-12** was prepared from 2,4-difluorobenzaldehyde:

[00214] Compound **A-11** (2.0 g, white solid, 90% yield) was prepared from 2,4-difluorobenzaldehyde (2.0 g, 14 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1]. LCMS (B): (ES⁺) *m/z* (M+H)⁺ = 158.1, *t*R=0.598 min.

[00215] Compound **A-12** (1.4 g, white solid, crude) was prepared from compound **A-11** (1.0 g, 6.4 mmol). TLC [petroleum ether: ethyl acetate = 5:1]: *R*_f = 0.6.

[00216] Example 11A: 4-cyano-2-fluoro-N-hydroxybenzimidoyl chloride (A-14)




[00217] Following general procedure A1, compound **A-14** was prepared from 3-fluoro-4-formylbenzonitrile:

[00218] Compound **A-13** (2.7 g, white solid, 82% yield) was prepared from 3-fluoro-4-formylbenzonitrile (3.0 g, 20 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1]. LCMS (B): (ES⁺) m/z (M+H)⁺ = 165.1, tR=0.538 min.

[00219] Compound **A-14** (0.30 g, white solid, crude) was prepared from compound **A-13** (0.40 g, 2.4 mmol). TLC [petroleum ether: ethyl acetate = 5:1]: R_f = 0.6.

[00220] **Example 12A: 2-fluoro-N-hydroxy-4-(methylsulfonyl)benzimidoyl chloride (A-16)**




[00221] Following general procedure A1, compound **A-16** was prepared from 2-fluoro-4-(methylsulfonyl) benzaldehyde:


[00222] Compound **A-15** (1.8 g, white solid, 81% yield) was prepared from 2-fluoro-4-(methylsulfonyl) benzaldehyde (2.0 g, 9.9 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 8:1]. LCMS (B): (ES⁺) m/z (M+H)⁺ = 218.1, tR=0.798 min.

[00223] Compound **A-16** (0.6 g, white solid, crude) was prepared from compound **A-15** (1.0 g, 4.6 mmol). TLC [petroleum ether: ethyl acetate = 10:1]: R_f = 0.5.


[00224] **Example 13A: 1-(tert-butyl)-3-fluorobenzene (A-17)**



[00225] To 40% fluoroboric acid (40 mL) was added 3-tert-butylaniline (5.5 g, 37 mmol) portion-wise. The mixture was stirred at 25 °C for 0.5 hour and then cooled to 0 °C. Sodium nitrite (4.1 g, 59 mmol) in water (10 mL) was added, and the solution was stirred at 0 °C for 0.5 hour and at 25 °C for 20 mins, resulting in formation of a solid. The solid was collected by filtration, washed with 40% fluoroboric acid (10 mL), ethanol (5 mL) and n-hexane (10 mL), dried, dissolved in tetrahydrofuran (60 mL) and stirred at 40 °C for 12 hours. On completion, the mixture was concentrated, and the residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 1:0] to afford compound **A-17** (4.0 g, 71% yield) as a yellow oil.

[00226] Example 14A: 4-(tert-butyl)-2-fluorobenzaldehyde (A-18)


[00227] To a solution of compound **A-17** (1.9 g, 12 mmol) in anhydrous tetrahydrofuran (25 mL) at -70 °C was added dropwise sec-butyllithium (12 mL, 1.3 N in pentane, 15 mmol). The reaction was stirred at -70 °C for 1h. Then *N,N*-dimethylformamide (3.6 g, 49 mmol) was added, and stirring was continued at -70 °C for another 1h. On completion, the mixture was poured into aqueous ammonium chloride (20 mL) and extracted with ethyl acetate (3 × 20 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 15:1] to give compound **A-18** (2.0 g, 91% yield) as a yellow oil.

[00228] Example 15A: 4-(tert-butyl)-2-fluoro-N-hydroxybenzimidoyl chloride (A-20)


[00229] Following general procedure A1, compound **A-20** was prepared from compound **A-18**:

[00230] Compound **A-19** (0.35 g, white solid, 73% yield) was prepared from compound **A-18** (0.45 g, 2.5 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1]. LCMS (DD): (ES⁺) *m/z* (M+H)⁺ = 196.2, *t*R=0.835 min.


[00231] Compound **A-20** (0.30 g, white solid, crude) was prepared from compound **A-19** (0.35 g, 1.8 mmol).

[00232] Example 16A: (4,5-dichloro-2-fluorophenyl)methanol (A-21)

[00233] To a solution of 4, 5-dichloro-2-fluoro-benzoic acid (5.0 g, 24 mmol) in tetrahydrofuran (50 mL) at 0 °C was added borane-tetrahydrofuran complex (1 M, 60 mL, 60 mmol) dropwise over 30 minutes. The resulting solution was stirred at 30 °C for 14 hours until TLC analysis showed the reaction was complete. The reaction was quenched carefully with methanol (10 mL) and then concentrated in vacuo to give compound **A-21** (4.50 g, 85% yield) as a yellow oil, which was used in the next step without further purification. LCMS (DD): (ES⁺) *m/z* (M+H)⁺ = 176.9, *t*R=0.859 min.

[00234] Example 17A: 4,5-dichloro-2-fluorobenzaldehyde (A-22)


[00235] A mixture of compound **A-21** (2.8 g, 14 mmol) and manganese dioxide (12 g, 143 mmol) in dichloromethane (40 mL) was stirred at 30 °C for 16 hours until TLC analysis showed the starting material was consumed. The reaction mixture was filtered, and the filtrate was concentrated in vacuo. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 5:1] to afford compound **A-22** (2.0 g, 73% yield) as a white solid.

[00236] Example 18A: 4,5-dichloro-2-fluoro-N-hydroxybenzimidoyl chloride (A-24)

[00237] Following general procedure A1, compound **A-24** was prepared from compound **A-22**:

[00238] Compound **A-23** (3.0 g, white solid, 80% yield) was prepared from compound **A-22** (3.5 g, 18 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1]. LCMS (DD): (ES⁺) m/z (M+H)⁺ = 207.9, tR=0.931 min.


[00239] Compound **A-24** (0.30 g, white solid, crude) was prepared from compound **A-23** (0.30 g, 1.4 mmol).

[00240] Example 19A: 2-fluoro-N-hydroxy-4-(trifluoromethoxy)benzimidoyl chloride (A-26)

[00241] Following general procedure A1, compound **A-26** was prepared from 2-fluoro-4-(trifluoromethoxy) benzaldehyde:

[00242] Compound **A-25** (0.93 g, white solid, crude) was prepared from 2-fluoro-4-(trifluoromethoxy)benzaldehyde (1.0 g, 4.8 mmol) and hydroxylamine hydrochloride (1.0 g, 14 mmol) using a mixture of ethanol (12 mL) and water (2.4 mL) without triethylamine as the solvent instead of dichloromethane. The reaction time was 2.5 hours. The product was used directly for the next step without further purification. LCMS (DD): (ES⁺) m/z (M+H)⁺ = 224.0, tR=0.953 min.


[00243] Compound **A-26** (0.63 g of yellow oil, crude) was prepared from compound **A-25** (0.55 g, 2.5 mmol) and *N*-chlorosuccinimide (0.40g, 3.0 mmol). The reaction time was 3 hours. LCMS (B): tR=0.836 min, (ES⁺) m/z (M+H)⁺ = 258.0.

[00244] Example 20A: 2,3-difluoro-N-hydroxybenzimidoyl chloride (A-28)

[00245] Following general procedure A1, compound A-28 was prepared from 2,3-difluorobenzaldehyde:

[00246] Compound A-27 (1.0 g, white solid, 90% yield) was prepared from 2,3-difluorobenzaldehyde (1.0 g, 7.0 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1].


[00247] Compound A-28 (1.2 g, white solid, crude) was prepared from compound A-27 (1.0 g, 6.4 mmol).

[00248] Example 21A: 2,5-difluoro-N-hydroxybenzimidoyl chloride (A-30)

[00249] Following general procedure A1, compound A-30 was prepared from 2,5-difluorobenzaldehyde:

[00250] Compound A-29 (1.0 g, white solid, crude) was prepared from 2,5-difluorobenzaldehyde (1.0 g, 7.0 mmol) with a reaction time of 14 hours and used in next step without further purification. LCMS (B): (ES⁺) m/z (M+H)⁺ = 158.0, tR = 0.559 min.


[00251] Compound A-30 (1.0 g, white solid, crude) was prepared from compound A-29 (1.0 g, 6.4 mmol) with a reaction time of 12 hours. [petroleum ether: ethyl acetate = 5:1]; R_f = 0.7.

[00252] Example 22A: 2,6-difluoro-N-hydroxybenzimidoyl chloride (A-32)

[00253] Following general procedure A1, compound A-32 was prepared from 2,6-difluorobenzaldehyde:

[00254] Compound A-31 (1.1 g, white solid, 99% yield) was prepared from 2,4-difluorobenzaldehyde (1.0 g, 7.0 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 158.1, tR = 0.838 min.


[00255] Compound A-32 (0.60 g, white solid, crude) was prepared from compound A-31 (0.50 g, 3.2 mmol). The reaction time was 16 hours. TLC [petroleum ether: ethyl acetate = 10:1]; R_f = 0.45.

[00256] Example 23A: 2,3-difluoro-N-hydroxy-4-methylbenzimidoyl chloride (A-34)

[00257] Following general procedure A1, compound A-34 was prepared from 2,3-difluoro-4-methylbenzaldehyde:

[00258] Compound A-33 (1.0 g, white solid, 91% yield) was prepared from 2,3-difluoro-4-methylbenzaldehyde (1.0 g, 6.4 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 172.1, tR=1.279 min.


[00259] Compound A-34 (0.60 g, white solid, crude) was prepared from compound A-33 (0.50 g, 3.2 mmol). The reaction time was 16 hours. TLC [petroleum ether: ethyl acetate = 10:1]: R_f = 0.5.

[00260] Example 24A: 2,3,4-trifluoro-N-hydroxybenzimidoyl chloride (A-36)


[00261] Following general procedure A1, compound A-36 was prepared from 2, 3, 4-trifluorobenzaldehyde:

[00262] Compound A-35 (0.80 g, white solid, 73% yield) was prepared from 2,3,4-trifluorobenzaldehyde (1.0 g, 6.3 mmol) (reaction time was 12 hours), and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1]. LCMS (B): (ES⁺) m/z (M+H)⁺ = 176.1, tR=0.68 min.


[00263] Compound A-36 (0.35 g, white solid, crude) was prepared from compound A-35 (0.40 g, 2.3 mmol) with a reaction time of 12 hours. TLC [petroleum ether: ethyl acetate = 7:1]: R_f = 0.6.

[00264] Example 25A: (2,5-difluoro-4-methylphenyl)methanol (A-37)

[00265] To a solution of 4, 5-dichloro-2-fluoro-benzoic acid (2.75 g, 16 mmol) in tetrahydrofuran (30 mL) at 0 °C was added borane-tetrahydrofuran complex (1 M, 40 mL, 40 mmol) drop-wise over 30 minutes. The resulting solution was stirred at 30 °C for 2.5 hours. On completion, the reaction was quenched carefully with methanol (10 mL) and then concentrated in vacuo to give compound A-37 (2.0 g, crude) as a yellow solid, which was used in next step without further purification. LCMS (J): (ES⁺) m/z (M+H)⁺ = 159.1, tR=1.12 min.

[00266] Example 26A: 2, 5-difluoro-4-methylbenzaldehyde (A-38)


[00267] A mixture of compound **A-37** (2.5 g, 16 mmol) and manganese dioxide (14 g, 0.16 mol) in dichloromethane (20 mL) was stirred at 30°C for 46 hours. On completion, the reaction mixture was filtered, and the filtrate was concentrated in vacuo to give compound **A-38** (2.0 g, crude) as a white solid. LCMS (J): (ES+) m/z (M+H) $^+$ = 157.1, t_R = 1.28 min.

[00268] Example 27A: 2,5-difluoro-N-hydroxy-4-methylbenzimidoyl chloride (A-40)


[00269] Following general procedure A1, compound **A-40** was prepared from compound **A-38**.

[00270] Compound **A-39** (2.3 g, crude) was prepared from compound **A-38** (2.5 g, 16 mmol). The reaction time was 15 hours.

[00271] Compound **A-40** (1.2 g, white solid, crude) was prepared from compound **A-39** (1.0 g, 5.8 mmol). The reaction time was 14 hours. TLC [petroleum ether: ethyl acetate = 8:1]: R_f = 0.5.

[00272] Example 28A: 4-chloro-2-fluoro-3-methylbenzaldehyde (A-41)

[00273] To a solution of 1-chloro-3-fluoro-2-methylbenzene (10 g, 69 mmol) in anhydrous tetrahydrofuran (20 mL) at -78°C was added lithium diisopropylamide (2 M in tetrahydrofuran, 0.10 mol, 50 mL). The reaction was stirred for 0.5 hr, then *N,N*-dimethylformamide (0.21 mol, 15 g) was added, and stirring was continued -78°C for 1.5 hrs. On completion, the reaction mixture was quenched with water and extracted with ethyl acetate (3×200 mL). The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate and concentrated in vacuo to give **A-41** (12 g, crude) as a white solid.


[00274] Example 29A: 4-chloro-2-fluoro-N-hydroxy-3-methylbenzimidoyl chloride (A-43)

[00275] Following general procedure A1, compound **A-43** was prepared from compound **A-41**.

[00276] Compound **A-42** (1.1 g crude) was prepared from compound **A-41** (12 g, 70 mmol). ¹H-NMR (CD₃OD, 400 MHz): δ 11.68 (m, 1 H), 8.21-8.19 (m, 1H), 7.61-7.57 (m, 1H), 7.34-7.32(m, 1H), 2.28 (s, 3H).

[00277] Compound **A-43** (0.52 g, white solid, crude) was prepared from compound **A-42** (0.50 g, 2.7 mmol). The reaction time was 14 hours. TLC [petroleum ether: ethyl acetate = 8:1]: R_f = 0.5.

[00278] **Example 30A: 4-chloro-2-fluoro-N-hydroxy-3-methoxybenzimidoyl chloride (A-45)**




[00279] Following general procedure A1, compound **A-45** was prepared from 4-chloro-2-fluoro-3-methoxybenzaldehyde:

[00280] Compound **A-44** (0.49 g, white solid, crude) was prepared from 4-chloro-2-fluoro-3-methoxybenzaldehyde (0.50 g, 2.7 mmol) with a reaction time of 15 hours.


[00281] Compound **A-45** (0.47 g, white solid, crude) was prepared from compound **A-44** (0.45 g, 2.2 mmol) with a reaction time of 16 hours. TLC [petroleum ether: ethyl acetate = 5:1]: R_f = 0.7.

[00282] **Example 31A: 4-chloro-2-fluoro-3-(trifluoromethyl)benzaldehyde (A-46)**



[00283] To a solution of 1-chloro-3-fluoro-2-(trifluoromethyl)benzene (0.50 g, 2.5 mmol) at -70 °C was added dropwise lithium diisopropylamide (2 M in tetrahydrofuran, 3.8 mmol, 1.9 mL). The reaction was stirred at -70 °C for 0.5 hour, then *N,N*-dimethylformamide (0.55 g, 7.6 mmol) was added slowly, and stirring was continued at -70 °C for another hour. The reaction was quenched with water (10 mL) and extracted with ethyl acetate (3 x 10 mL). The combined organic layers were concentrated in vacuo, and the residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1] to give **A-46** (0.40 g, 70% yield) as a yellow solid. ¹H-NMR (CDCl₃, 400 MHz): δ 10.28 (s, 1H), 7.95-7.90 (t, J=10.0 Hz, 1H), 7.40-7.37 (d, J=11.2 Hz, 1H).

[00284] **Example 32A: 4-chloro-2-fluoro-N-hydroxy-3-(trifluoromethyl)benzimidoyl chloride (A-48)**




[00285] Following general procedure A1, compound **A-48** was prepared from compound **A-46**:

[00286] Compound **A-47** (0.40 g, yellow solid, 62% yield) was prepared from compound **A-46** (0.60 g, 2.7 mmol) with a reaction time of 14 hours and used in next step without further purification.


[00287] Compound **A-48** (0.40 g, white solid, crude) was prepared from compound **A-47** (0.38 g, 1.6 mmol). TLC [petroleum ether: ethyl acetate = 8:1]. Rf = 0.6.

[00288] **Example 33A:** (3-bromo-4-chloro-2-fluorophenyl)methanediol (**A-49**)




[00289] To a solution of 2-bromo-1-chloro-3-fluorobenzene (6.0 g, 29 mmol) in tetrahydrofuran (30 mL) at -70 °C was added dropwise lithium diisopropylamide (2 M in tetrahydrofuran/n-heptane, 22 mL, 43 mmol). The resulting solution was stirred at -70 °C for 0.5 hour. Then *N,N*-dimethylformamide (4.2 g, 57 mmol) was added slowly, and stirring was continued at -70 °C for another 0.5 hour. On completion of the reaction by TLC [petroleum ether: ethyl acetate = 10:1, Rf=0.6], the reaction was quenched with saturated aqueous ammonium chloride (2 mL) at 0 °C and extracted with acetate ethyl (3 × 20 mL). The combined organic extracts were concentrated to give compound **A-49** (4.0 g, 58% yield) as a yellow solid. ¹H-NMR (CD₃OD, 400 MHz): δ 7.64-7.57 (m, 1H), 7.21-7.17 (m, 1H), 6.02 (s, 1H).

[00290] **Example 34A:** (3-bromo-4-chloro-2-fluorophenyl)methanol (**A-50**)



[00291] To a solution of compound **A-49** (2.0 g, 8.4 mmol) in methanol (10 mL) at 0 °C was added sodium borohydride (0.64 g, 17 mmol). The reaction was allowed to warm to room temperature and stirred for another 1 hour, until TLC showed the reaction was complete [petroleum ether: ethyl acetate = 10:1, Rf=0.4]. The reaction was quenched at 0 °C with 2 N hydrochloric acid (2 mL) and extracted with ethyl acetate (3 × 200 mL). The combined organic phase was dried over anhydrous sodium sulfate and concentrated to afford compound **A-50** (2.0 g, crude) as a yellow oil. ¹H-NMR (CD₃OD, 400 MHz): δ 7.50-7.46 (m, 1H), 7.17-7.13 (m, 1H), 4.89 (s, 2H).


[00292] **Example 35A:** (4-chloro-3-cyclopropyl-2-fluorophenyl)methanol (**A-51**)



[00293] To a solution of compound **A-50** (2.0 g, 8.4 mmol) in dioxane (8.0 mL) and water (1.0 mL) was added potassium phosphate (5.3 g, 25 mmol), [1,1'-bis(diphenylphosphino)ferrocene] dichloropalladium(II) (0.61 g, 0.84 mmol) and cyclopropylboronic acid (1.1 g, 13 mmol). The mixture was stirred at 80 °C for 4 hours, then concentrated and purified by silica gel chromatography


[petroleum ether: ethyl acetate = 5:1] to afford compound **A-51** (1.2 g, 72% yield over two steps) as a yellow oil. GCMS: tR=9.103 min, 199.9, (EI) m/z (M)⁺.

[00294] Example 36A: 4-chloro-3-cyclopropyl-2-fluorobenzaldehyde (A-52)



[00295] A mixture of compound **A-51** (1.2 g, 6.0 mmol) and manganese dioxide (11 g, 0.12 mol) in toluene (40 mL) was stirred at 100 °C for 16 hours until TLC analysis showed the starting material was consumed. The reaction mixture was filtered, and the filtrate was concentrated in vacuo. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 5:1] to afford compound **A-52** (0.6 g, crude) as a yellow oil. LCMS (B): (ES⁺) m/z (M+H)⁺ = 199.0, tR = 0.827 min.

[00296] Example 37A: 4-chloro-3-cyclopropyl-2-fluoro-N-hydroxybenzimidoyl chloride (A-54)




[00297] Following general procedure A1, compound **A-54** was prepared from compound **A-52**:

[00298] Compound **A-53** (0.45 g, white solid, 93% yield over two steps) was prepared from compound **A-52** (0.45 g, 2.3 mmol). ¹H-NMR (CDCl₃, 400 MHz): δ 9.23 (s, 1H), 8.47 (s, 1H), 7.15-7.12 (d, J=8.4 Hz, 1H), 6.87-6.83 (d, J=8.0 Hz, 1H), 2.16-2.07 (m, 1H), 1.05-0.97 (m, 2H), 0.73-0.68 (m, 2H).

[00299] Compound **A-54** (0.50 g, colorless oil, crude) was prepared from compound **A-53** (0.40 g, 1.7 mmol) with a reaction time of 16 hours. TLC [petroleum ether: ethyl acetate = 10:1]: R_f = 0.5.


[00300] Example 38A: 4-chloro-2,6-difluoro-N-hydroxybenzimidoyl chloride (A-56)




[00301] Following general procedure A1, compound **A-56** was prepared from 4-chloro-2,6-difluorobenzaldehyde:

[00302] Compound **A-55** (0.50 g, white solid, 92% yield) was prepared from 4-chloro-2,6-difluorobenzaldehyde (0.50 g, 2.8 mmol) with a reaction time of 12 hours and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1]. LCMS (B): tR=0.678 min., 192.1 m/z (M+1); ¹H-NMR (CDCl₃, 400 MHz): δ 8.27 (s, 1H), 8.21 (s, 1H), 7.03-7.00 (m, 2H).

[00303] Compound **A-56** (0.80 g of colorless oil, crude) was prepared from compound **A-55** (0.80 g, 4.2 mmol) with a reaction time of 16 hours. TLC [petroleum ether: ethyl acetate = 5:1]: R_f = 0.70.

[00304] Example 39A: 2,4-dichloro-6-fluorobenzaldehyde (A-57)


[00305] To a solution of 1,3-dichloro-5-fluorobenzene (2.0 g, 12 mmol) in tetrahydrofuran (20 mL) at -70 °C was added dropwise lithium diisopropylamide (2 M in tetrahydrofuran, 18 mmol., 9.1 mL). The reaction was stirred at -70 °C for 0.5 hr, then *N,N*-dimethylformamide (1.8 g, 24 mmol) was added slowly, and stirring was continued at -70 °C for another 0.5 hr. The reaction was quenched with water (20 mL) and extracted with acetate ethyl (3 x 20 mL). The combined organic layers were concentrated in vacuo and purified by silica gel chromatography [petroleum ether: ethyl acetate = 30 : 1] to give **A-57** (1.5 g, 64% yield) as a yellow solid. ¹H-NMR (CDCl₃, 400 MHz): δ 10.41 (s, 1H), 7.934 (s, 1H), 7.18-7.16 (d, J=10.0 Hz, 1H).

[00306] Example 40A: 2,4-dichloro-6-fluoro-N-hydroxybenzimidoyl chloride (A-59)

[00307] Following general procedure A1, compound **A-59** was prepared from compound **A-57**:

[00308] Compound **A-58** (0.75 g, white solid, 90% yield) was prepared from compound **A-57** (0.77 g, 7.0 mmol) with a reaction time of 3 hours and used in the next step without further purification. LCMS (B): (ES⁺) m/z (M+H)⁺ = 207.9, t_R=0.628 min.

[00309] Compound **A-59** (0.35 g, white solid, crude) was prepared from compound **A-58** (0.30 g, 1.4 mmol) with a reaction time of 12 hours. TLC [petroleum ether: ethyl acetate = 5:1]: R_f = 0.8.


[00310] Example 41A: 2,3-difluoro-N-hydroxy-4-methoxybenzimidoyl chloride (A-61)

[00311] Following general procedure A1, compound **A-61** was prepared from 2,3-difluoro-4-methoxybenzaldehyde:

[00312] Compound **A-60** (0.50 g, white solid, 92% yield) was prepared from 2,3-difluoro-4-methoxybenzaldehyde (0.50 g, 2.9 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 8:1].


[00313] Compound **A-61** (0.50 g, white solid, crude) was prepared from compound **A-60** (0.40 g, 2.1 mmol). TLC [petroleum ether: ethyl acetate = 3:1]: R_f = 0.5.

[00314] Example 42A: 3-chloro-2-fluoro-4-methoxybenzaldehyde (A-62)



[00315] To a solution of 2-chloro-1-fluoro-3-methoxybenzene (1.5 g, 9.3 mmol) in tetrahydrofuran (10 mL) at -78 °C was added lithium diisopropylamide (2M in tetrahydrofuran, 7.0 mL, 14 mmol). The reaction was stirred for 0.5 hour, then *N,N*-dimethylformamide (2.1 g, 28 mmol) was added, and stirring was continued at -78 °C for 2.5 hours. On completion, the reaction mixture was quenched with water (20 mL) and extracted with ethyl acetate (3 × 20 mL). The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate and concentrated in vacuo to give compound **A-62** (0.40 g, 23% yield) as a white solid.

[00316] **Example 43A:** 3-chloro-2-fluoro-*N*-hydroxy-4-methoxybenzimidoyl chloride (**A-64**)




[00317] Following general procedure A1, compound **A-64** was prepared from compound **A-62**:

[00318] Compound **A-63** (0.43 g, white solid, crude) was prepared from compound **A-62** (0.40 g, 2.1 mmol) with a reaction time is 12 hours.


[00319] Compound **A-64** (0.45 g, white solid, crude) was prepared from compound **A-63** (0.43 g, 2.1 mmol) with a reaction time is 15 hours. TLC [petroleum ether: ethyl acetate = 5:1]: *R_f* = 0.7.

[00320] **Example 44A:** 3-chloro-2-fluoro-4-methylbenzaldehyde (**A-65**)



[00321] To a solution of 2-chloro-1-fluoro-3-methylbenzene (1.0 g, 6.9 mmol) in tetrahydrofuran (30 mL) at -70 °C was added dropwise *n*-butyllithium (2.5 N in hexane, 3.3 mL, 8.3 mmol). The reaction was stirred at -70 °C for 0.5 h, then *N,N*-dimethylformamide (2.0 g, 28 mmol) was added, and stirring was continued at -70 °C for another 0.5 h. On completion, the mixture was poured into ammonium chloride (40 mL) solution and extracted with ethyl acetate (3 × 50 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated in vacuo to give compound **A-65** (1.2 g, 86% yield) as yellow solid, which was used in the next step without further purification. LCMS (J): (ES⁺) *m/z* (M+H)⁺ = 173.0, *t_R* = 0.508 min.

[00322] **Example 45A:** 3-chloro-2-fluoro-*N*-hydroxy-4-methylbenzimidoyl chloride (**A-67**)




[00323] Following general procedure A1, compound **A-67** was prepared from compound **A-65**:

[00324] Compound **A-66** (1.1 g, white solid, 85% yield) was prepared from compound **A-65** (1.2 g, 7.0 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 8:1]. ¹H-NMR (CD₃OD, 400 MHz): δ 8.20 (s, 1H), 7.61-7.57 (t, J=7.2 Hz, 1H), 7.09-7.07 (d, J=8.0 Hz, 1H).


[00325] Compound **A-67** (0.60 g, white solid, crude) was prepared from compound **A-66** (0.50 g, 2.7 mmol). TLC [petroleum ether: ethyl acetate = 3:1]: R_f = 0.5.

[00326] **Example 46A: 1-bromo-2-fluoro-4-isopropoxybenzene (A-68)**




[00327] A mixture of 4-bromo-3-fluorophenol (5.0 g, 26 mmol), 2-bromopropane (6.4 g, 52 mmol) and potassium carbonate (29 g, 0.21 mol) in *N,N*-dimethylformamide (50 mL) was stirred at 80 °C for 16 hrs. On completion, the reaction was diluted with water (50 mL) and extracted with ethyl acetate (3 x 50 mL). The combined organic layers were concentrated in vacuo, and the residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 30:1] to give compound **A-68** (5.0 g, 82% yield) as colourless oil. ¹H-NMR (CDCl₃, 400 MHz): δ 7.42-7.38 (t, J=8.8 Hz, 1H), 6.71-6.68 (dd, J=10.8 Hz, J=2.8 Hz, 1H), 6.62-6.59 (dd, J=8.8 Hz, J=1.6 Hz, 1H), 4.55-4.46 (m, 1H), 1.36-1.34 (d, J=6.0 Hz, 6H).

[00328] **Example 47A: 2-fluoro-4-isopropoxybenzaldehyde (A-69)**



[00329] To a solution of **A-68** (1.5 g, 6.4 mmol) in tetrahydrofuran (20 mL) at -70 °C was added dropwise *n*-butyllithium (2.5 M in hexanes, 3.1 mL, 7.8 mmol). The reaction was stirred at -70 °C for 0.5 hr., then *N,N*-dimethylformamide (1.4 g, 19 mmol) was added slowly, and stirring was continued at -70 °C for another 0.5 hr. On completion, the reaction was quenched slowly with saturated aqueous ammonium chloride (30 mL) and extracted with ethyl acetate (3 x 30 mL). The combined organic phases were concentrated in vacuo to give compound **A-69** (1.0 g, 85% yield) as a light yellow solid.

[00330] **Example 48A: 2-fluoro-N-hydroxy-4-isopropoxybenzimidoyl chloride (A-71)**




[00331] Following general procedure A1, compound **A-71** was prepared from compound **A-69**:

[00332] Compound **A-70** (0.90 g, light yellow solid, 83% yield) was prepared from compound **A-69** (1.0 g, 5.5 mmol) with a reaction time of 14 hours and purified by silica gel chromatography [petroleum ether: ethyl acetate = 20:1]. ¹H-NMR (CDCl₃, 400 MHz): δ 8.31 (s, 1H), 7.83 (s, 1H),

7.67-7.62 (t, J=8.4 Hz, 1H), 6.72-6.69 (dd, J=8.4 Hz, J=2.4 Hz, 1H), 6.64-6.60 (m, 1H), 4.62-4.52 (m, 1H), 1.38-1.37 (m, 6H).


[00333] Compound **A-71** (0.80 g, yellow solid, crude) was prepared from compound **A-70** (0.60 g, 3.0 mmol) with a reaction time of 16 hours. TLC [petroleum ether: ethyl acetate = 5:1]: R_f = 0.5.

[00334] **Example 49A: 4-ethoxy-2-fluorobenzaldehyde (A-72)**



[00335] To a mixture of 2-fluoro-4-hydroxybenzaldehyde (3.0 g, 21 mmol) and iodoethane (5.0 g, 32 mmol) in *N,N*-dimethylformamide (0.5 mL) was added potassium carbonate (4.4 g, 32 mmol). The reaction mixture was stirred at 50 °C for 16 hours. On completion, the reaction mixture was diluted with water and extracted with ethyl acetate (3 × 150 mL). The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate, filtered and concentrated in vacuo to give compound **A-72** (3.4 g, crude) as a yellow solid. LCMS (B): (ES⁺) m/z (M+H)⁺ = 169.2, t_R=0.649 min.

[00336] **Example 50A: 4-ethoxy-2-fluoro-N-hydroxybenzimidoyl chloride (A-74)**




[00337] Following general procedure A1, compound **A-74** was prepared from compound **A-72**:

[00338] Compound **A-73** (1.0 g, white solid, 92% yield over two steps) was prepared from compound **A-72** (1.0 g, 6.0 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1]. LCMS (B): (ES⁺) m/z (M+H)⁺ = 184.2, t_R=0.757 min.

[00339] Compound **A-74** (0.60 g, white solid, crude) was prepared from compound **A-73** (0.50 g, 2.9 mmol) with a reaction time of 16 hours. TLC [petroleum ether: ethyl acetate = 5:1]: R_f = 0.7.

[00340] **Example 51A: 6-chloro-2-fluoro-3-formylbenzonitrile (A-75)**



[00341] To a mixture of 2-chloro-6-fluorobenzonitrile (5.0 g, 32 mmol) in anhydrous tetrahydrofuran (50 mL) at -70 °C under nitrogen was added dropwise lithium diisopropylamide (2.0 M in tetrahydrofuran/n-heptane, 24 mL, 48 mmol). The mixture was stirred at -70 °C for 0.5 hour, and *N,N*-dimethylformamide (3.5 g, 48 mmol) was added dropwise. The reaction was stirred at -70 °C for another 0.5 hour, then quenched with saturated ammonium chloride solution (500 mL) at 0 °C and extracted with ethyl acetate (2 × 200 mL). The combined organic layers were dried with anhydrous sodium sulfate, filtered and concentrated in vacuo. The residue was purified by silica gel

chromatography [petroleum ether: ethyl acetate = 30:1] to give compound **A-75** (1.0 g, 17% yield) as a yellow solid. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 10.33 (s, 1H), 8.07 (t, $J=8.0$ Hz, 1H), 7.50 (d, $J=8.4$ Hz, 1H).

[00342] **Example 52A: 4-chloro-3-cyano-2-fluoro-N-hydroxybenzimidoyl chloride (A-77)**




[00343] Following general procedure A1, compound **A-77** was prepared from compound **A-75**:

[00344] Compound **A-76** (0.36 g, white solid, 33% yield) was prepared from compound **A-75** (0.20 g, 1.1 mmol, 5 batches) with a reaction time of 2 hours and 0 °C and purified by prep-TLC [petroleum ether: ethyl acetate = 5:1]. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 8.31 (s, 1H), 8.00 (t, $J=8.0$ Hz, 1H), 7.71 (s, 1H), 7.36 (d, $J=8.8$ Hz, 1H).

[00345] Compound **A-77** (0.50 g of yellow oil, crude) was prepared from compound **A-76** (0.31 g, 1.3 mmol), in dichloromethane and stirred at 20 °C for 12 hours. TLC [petroleum ether: ethyl acetate = 5:1]: R_f = 0.61.

[00346] **Example 53A: 3-chloro-2,4-difluoro-N-hydroxybenzimidoyl chloride (A-79)**




[00347] Following general procedure A1, compound **A-79** was prepared from 3-chloro-2,4-difluorobenzaldehyde:

[00348] Compound **A-78** (0.90 g, white solid, 83% yield) was prepared from 3-chloro-2,4-difluorobenzaldehyde (1.0 g, 5.7 mmol) and used in the next step without further purification.

[00349] Compound **A-79** (0.80 g, white solid, crude) was prepared from compound **A-78** (0.90 g, 4.7 mmol) with a reaction of 16 hours. TLC [petroleum ether: ethyl acetate = 8:1]: R_f = 0.75.

[00350] **Example 54A: 2-fluoro-4-(1H-pyrazol-1-yl)benzaldehyde (A-80)**



[00351] A mixture of 2,4-difluorobenzaldehyde (14 g, 1.0 mol), 1H-pyrazole (6.8 g, 1.0 mmol) and potassium carbonate (14 g, 1.0 mmol) in *N,N*-dimethylacetamide (200 mL) was stirred at 200 °C for 10 mins. TLC [petroleum ether: ethyl acetate = 5:1] showed starting material consumed and two new spots formed. The reaction mixture was diluted with water (200 mL) and extracted with ethyl acetate (2 x 150 mL). The combined organic phases were washed with water (5 x 150 mL) and brine (150 mL), dried over sodium sulfate, filtered and concentrated. The residue was purified by

Prep-HPLC (TFA, Instrument: HPLC-B, column: Phenomenex Synergi Max-RP 250x50mmx10 μ m. Condition: 0.1%TFA-ACN) and lyophilization to give compound **A-80** (3.5 g, 18% yield) as a white solid. LCMS (D): (ES⁺) m/z (M+H)⁺ = 191.1, tR = 1.271 min.

[00352] **Example 55A:** 2-fluoro-N-hydroxy-4-(1H-pyrazol-1-yl)benzimidoyl chloride (**A-82**)




[00353] Following general procedure A1, compound **A-82** was prepared from compound **A-80**:

[00354] Compound **A-81** (0.75 g, white solid, 90% yield) was prepared from compound **A-80** (1.0 g, 5.3 mmol) with a reaction time of 3 hours and purified by silica gel chromatography [petroleum ether: ethyl acetate = 10:1]. LCMS (Y): (ES⁺) m/z (M+H)⁺ = 206.2, tR = 0.666 min.


[00355] Compound **A-82** (0.40 g, white solid, crude) was prepared from compound **A-81** (0.30 g, 1.5 mmol) with a reaction time of 12 hours. [petroleum ether: ethyl acetate = 5:1]: Rf = 0.62.

[00356] **Example 56A:** methyl 4-bromo-3-fluorobenzoate (**A-83**)




[00357] To a solution of 4-bromo-3-fluoro-benzoic acid (10 g, 46 mmol) in methanol (120 mL) was added thionyl chloride (33 g, 0.28 mol). The reaction was stirred at 80 °C for 5 hours, then concentrated in vacuo to give compound **A-83** (11 g, crude) as yellow solid, which was used in the next step without further purification. TLC [petroleum ether:ethyl acetate = 10:1]: Rf = 0.80.

[00358] **Example 57A:** 4-bromo-3-fluorobenzohydrazide (**A-84**)



[00359] A mixture of compound **A-83** (11 g, crude) and hydrazine hydrate (23 g, 0.46 mol) in methanol (200 mL) was stirred at 70 °C for 12 hours. On completion, the mixture was concentrated in vacuo, diluted with ethyl acetate (60 mL), washed with water (50 mL) and brine (2 × 20 mL), dried over anhydrous sodium sulfate, filtered and concentrated in vacuo to afford compound **A-84** (5.0 g, crude) as a yellow solid.


[00360] **Example 58A:** 2-(4-bromo-3-fluorophenyl)-5-methyl-1,3,4-oxadiazole (**A-85**)



[00361] A mixture of compound **A-84** (4.0 g, crude) and triethylorthoacetate (2.5 g, 17 mmol) was stirred at 100 °C for 18 hours. On completion, the mixture was poured into water (40 mL) and


extracted with ethyl acetate (2 × 50 mL). The combined organic phase was washed with brine (2 × 20 mL), dried with anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 3:1] to give compound **A-85** (3.7 g, 39% yield over three steps) as a yellow solid. ¹H-NMR (CD₃OD, 400 MHz): δ 7.88-7.85 (m, 2H), 7.79-7.77 (m, 1H), 2.65 (m, 3H).

[00362] Example 59A: 2-(3-fluoro-4-vinylphenyl)-5-methyl-1,3,4-oxadiazole (A-86)




[00363] To a solution of compound **A-85** (3.3 g, 13 mmol), potassium trifluoro(vinyl)boranuide (2.1 g, 15 mmol) and cesium carbonate (13 g, 39 mmol) in tetrahydrofuran (9.0 mL) and water (1.0 mL) was added triphenylphosphine (0.34 g, 1.3 mmol) and palladium chloride (0.23 g, 1.3 mmol). The reaction was stirred at 80-90°C for 16 hours, then poured into water (20 mL) and extracted with ethyl acetate (2 × 20 mL). The combined organic phase was washed with brine, dried with anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 3:1] to give compound **A-86** (2.3 g, 88% yield) as a yellow solid. LCMS (B): (ES⁺) m/z (M+H)⁺ = 205.1, tR=0.730 min.

[00364] Example 60A: 2-fluoro-4-(5-methyl-1,3,4-oxadiazol-2-yl)benzaldehyde (A-87)



[00365] To a solution of compound **A-86** (1.5 g, 7.2 mmol), sodium periodate (3.1 g, 14 mmol) and 4-methylmorpholine (2.5 g, 21 mmol) in tetrahydrofuran (50 mL) and water (25 mL) was added osmium tetroxide (18 mg, 72 μmol). The reaction was stirred at 25 °C for 5 hours, then poured into aqueous sodium thiosulfate (20 mL) and extracted with ethyl acetate (2 × 20 mL). The combined organic phase was washed with brine, dried with anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 3:1] to give compound **A-87** (1.3 g, 88% yield) as a yellow solid. LCMS (M): (ES⁺) m/z (M+H)⁺ = 207.1, tR=0.705 min.

[00366] Example 61A: 2-fluoro-N-hydroxy-4-(5-methyl-1,3,4-oxadiazol-2-yl)benzimidoyl chloride (A-89)




[00367] Following general procedure A1, compound **A-89** was prepared from compound **A-87**:

[00368] Compound **A-88** (0.50 g, yellow solid, 34% yield) was prepared from compound **A-87** (1.4 g, 6.6 mmol) and purified by silica gel chromatography [petroleum ether: ethyl acetate = 2:1]. LCMS (B): (ES⁺) m/z (M+H)⁺ = 222.1, tR=0.657 min.


[00369] Compound **A-89** (0.60 g, yellow solid, crude) was prepared from compound **A-88** (0.50 g, 2.3 mmol). TLC [petroleum ether:ethyl acetate = 3:1]: R_f = 0.60.

[00370] **Example 62A: 2-fluoro-3-methoxy-4-methylbenzaldehyde (A-90)**



[00371] To a solution of 4-bromo-2-fluoro-3-methoxy-benzaldehyde (3.0 g, 13 mmol), 2,4,6-trimethyl-1,3,5,2,4,6-trioxatriborinane (2.1 g, 17 mmol) and potassium phosphate (5.5 g, 26 mmol) in water (10 mL) and tetrahydrofuran (30 mL) was added dicyclohexyl-[3-(2,4,6-triisopropylphenyl)phenyl]phosphane [2-(2-aminophenyl)phenyl]-chloro-palladium (0.51g, 0.64 mmol). The mixture was stirred at 80 °C for 12 hours. On completion, the mixture was poured into water (20 mL) and extracted with ethyl acetate (2 × 20 mL). The combined organic phase was washed with brine (2 × 20 mL), dried with anhydrous sodium sulfate, filtered and concentrated in vacuo to give compound **A-90** (1.60 g, crude) as a yellow solid, which was used in the next step without further purification. LCMS (B): (ES⁺) m/z (M+H)⁺ = 169.2, tR=0.640 min.

[00372] **Example 63A: 2-fluoro-N-hydroxy-3-methoxy-4-methylbenzimidoyl chloride (A-92)**




[00373] Following general procedure A1, compound **A-92** was prepared from compound **A-90**:

[00374] Compound **A-91** (0.70 g, yellow solid, 40% yield over two steps) was prepared from compound **A-90** (1.6 g, 9.5 mmol) with a reaction time of 12 hours and purified by silica gel chromatography [petroleum ether: ethyl acetate = 5:1]. LCMS (B): (ES⁺) m/z (M+H)⁺ = 184.2, tR=0.641 min.

[00375] Compound **A-92** (0.50 g, yellow solid, crude) was prepared from compound **A-91** (0.50 g, 2.7 mmol) with a reaction time of 12 hours and dichloromethane as the solvent. TLC [petroleum ether:ethyl acetate = 3:1]: R_f = 0.70.


[00376] **Example 64A: 1-chloro-2-ethoxy-3-fluorobenzene (A-93)**



[00377] A mixture of 2-chloro-6-fluoro-phenol (2.0 g, 14 mmol, iodoethane (4.3 g, 27 mmol) and potassium carbonate (3.8 g, 27 mmol) in *N,N*-dimethylformamide (20 mL) was stirred at 50 °C for

16 hrs. On completion, the reaction was diluted with water (30 mL) and extracted with methyl t-butyl ether (3 x 20 mL). The combined organic phases were washed with water (10 mL) and brine (10 mL), dried with anhydrous sodium sulfate and concentrated in vacuo. The residue was purified by silica gel chromatography [petroleum ether: ethyl acetate = 30:1] to give compound **A-93** (1.5 g, 63% yield) as light yellow oil.

[00378] Example 65A: 1-chloro-2-ethoxy-3-fluorobenzene (A-94)



[00379] To a solution of Compound **A-93** (1.0 g, 5.7 mmol) at $-70\text{ }^\circ\text{C}$ was added dropwise lithium diisopropylamide (2 M in tetrahydrofuran, 4.3 mL, 8.6 mmol). The reaction was stirred at $-70\text{ }^\circ\text{C}$ for 0.5 hr. Then *N,N*-dimethylformamide (1.7 g, 23 mmol) was added slowly, and stirring was continued at $-70\text{ }^\circ\text{C}$ for another 0.5 hr. The reaction was quenched with water (20 mL) and extracted with ethyl acetate (3 x 20 mL). The combined organic extracts were concentrated in vacuo and purified by silica gel chromatography [petroleum ether: ethyl acetate = 8:1] to give compound **A-94** (0.80 g, 69% yield) as a yellow solid.

[00380] Example 66A: 4-chloro-3-ethoxy-2-fluoro-N-hydroxybenzimidoyl chloride (A-96)




[00381] Following general procedure A1, compound **A-96** was prepared from compound **A-94**:

[00382] Compound **A-95** (90 mg, yellow solid, 84% yield) was prepared from compound **A-94** (0.10 g, 0.49 mmol), reaction was stirred at $20\text{ }^\circ\text{C}$, and purified by silica gel chromatography [petroleum ether: ethyl acetate = 20:1]. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 8.33 (s, 1H), 7.79 (s, 1H), 7.45-7.41 (dd, $J=7.2\text{ Hz}$, $J=1.2\text{ Hz}$, 1H), 7.20-7.17 (dd, $J=8.4\text{ Hz}$, $J=1.6\text{ Hz}$, 1H), 4.23-4.18 (dd, $J=14\text{ Hz}$, $J=7.2\text{ Hz}$, 1H), 1.47-1.44 (t, $J=7.2\text{ Hz}$, 1H).

[00383] Compound **A-96** (0.60 g, white solid, crude) was prepared from compound **A-95** (0.45g, 2.1 mmol), reaction was stirred at $20\text{ }^\circ\text{C}$. TLC [petroleum ether: ethyl acetate = 8:1]: $R_f = 0.52$.

[00384] Example 67A: 2-fluoro-3,4-dimethylbenzaldehyde (A-97)



[00385] To a mixture of tetramethylethylenediamine (2.1 g, 18 mmol) in anhydrous tetrahydrofuran (20 mL) at $-70\text{ }^\circ\text{C}$ under nitrogen was added tert-butyllithium (1.3 M in pentane solution, 14 mL, 18 mmol) dropwise. The resulting yellow solution was stirred at $-70\text{ }^\circ\text{C}$ for 0.5 hour.

Then a solution of 1-fluoro-2,3-dimethyl-benzene (2.0 g, 16 mmol) in anhydrous tetrahydrofuran (12 mL) was added dropwise. The mixture was stirred at -70 °C for 1 hour. Then a solution of *N,N*-dimethylformamide (1.3 g, 18 mmol) in anhydrous tetrahydrofuran (8.0 mL) was added, and the reddish-brown mixture was stirred at -70°C for an additional 1 hour. On completion, the reaction mixture was quenched with acetic acid (2.0 mL) and water (20 mL) at 0 °C and extracted with ethyl acetate (3 × 30 mL). The combined organic layers were washed with water (2 × 20 mL), dried with anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give compound **A-97** (1.5 g, 61% yield) as a yellow oil. TLC [petroleum ether: ethyl acetate = 5:1]: R_f = 0.61.

[00386] **Example 68A: 2-fluoro-*N*-hydroxy-3,4-dimethylbenzimidoyl chloride (**A-99**)**




[00387] Following general procedure A1, compound **A-99** was prepared from compound **A-97**:

[00388] Compound **A-98** (1.5 g, yellow solid, 90% yield) was prepared from compound **A-97** (1.5 g, 9.9 mmol) with a reaction time of 12 hours and purified by silica gel chromatography [petroleum ether: ethyl acetate = 50:1~5:1]. ¹H-NMR (CDCl₃, 400 MHz): δ 8.35 (s, 1H), 7.44 (t, J=8.0 Hz, 1H), 6.95 (d, J=8.0 Hz, 1H), 2.30 (s, 3H), 2.20(s, 3H).

[00389] Compound **A-99** (1.9 g, yellow solid, crude) was prepared from compound **A-98** (1.0 g, 6.0 mmol) using dichloromethane and a reaction time of 12 hours. TLC [petroleum ether: ethyl acetate = 10:1]: R_f = 0.43.

[00390] **Example 69A: 4-chloro-2,3-difluoro-*N*-hydroxybenzimidoyl chloride (compound-**A-101**)**




[00391] Following general procedure A2, compound **A-101** was prepared from 4-chloro-2,3-difluorobenzaldehyde:

[00392] Compound **A-100** (0.5 g, 92% yield, 11:1 mixture of (E)/(Z) isomers) was prepared as a white solid from 4-chloro-2,3-difluorobenzaldehyde (0.5 g, 2.8 mmol) using 9 mL of ethanol/water and a reaction time of 2 hours. LCMS (2): t_R=1.910 min., (ES⁺) m/z (M+H)⁺ = 191.9. ¹H NMR (300 MHz, DMSO-d₆, major isomer) δ 11.90 (s, 1H), 8.21 (s, 1H), 7.61 – 7.51 (m, 1H), 7.51 – 7.42 (m, 1H).

[00393] Compound **A-101** (99 mg, 95% yield) was prepared as a white solid from compound **A-100** (100 mg, 0.5 mmol) using 1 mL of *N,N*-dimethylformamide and a reaction time of 2 hours. ¹H NMR (300 MHz, DMSO-d₆) δ 12.91 (s, 1H), 7.61 – 7.53 (m, 2H).

[00394] **Example 70A: 4-chloro-2,5-difluoro-*N*-hydroxybenzimidoyl chloride (compound-**A-103**)**




[00395] Following general procedure A2, compound **A-103** was prepared from 4-chloro-2,5-difluorobenzaldehyde:

[00396] Compound **A-102** (0.5 g, 92% yield, 8:1 mixture of (E)/(Z) isomers) was prepared as a white solid from 4-chloro-2,5-difluorobenzaldehyde (0.5 g, 2.8 mmol) using 7 mL of ethanol/water and a reaction time of 2 hours. LCMS (2): $t_R=1.907$ min., (ES^+) m/z $(\text{M}+\text{H})^+ = 192.0$. ^1H NMR (300 MHz, DMSO- d_6 , major isomer) δ 11.89 (s, 1H), 8.15 (s, 1H), 7.84 – 7.58 (m, 2H).

[00397] Compound **A-103** (99 mg, 84% yield) was prepared as a white solid from compound **A-102** (100 mg, 0.5 mmol) using 1 mL of *N,N*-dimethylformamide and a reaction time of 16 hours. ^1H NMR (300 MHz, DMSO- d_6) δ 12.87 (s, 1H), 7.89 – 7.74 (m, 2H).

[00398] **Example 71A:** 4-bromo-2-fluoro-*N*-hydroxybenzimidoyl chloride (compound **A-105**)




[00399] Following general procedure A2, compound **A-105** was prepared from 4-bromo-2-fluorobenzaldehyde:

[00400] Compound **A-104** (2.5 g, 93% yield, 9:1 mixture of (E)/(Z) isomers) was prepared as a white solid from 4-bromo-2-fluorobenzaldehyde (2.5 g, 12.3 mmol) using 45 mL of ethanol/water and a reaction time of 2 hours. ^1H NMR (300 MHz, DMSO- d_6 , major isomer) δ 11.69 (s, 1H), 8.15 (s, 1H), 7.69 – 7.60 (m, 2H), 7.48 – 7.42 (m, 1H).

[00401] Compound **A-105** (2.3 g, 82% yield) was prepared as a white solid from compound **A-104** (2.4 g, 11.0 mmol) using 20 mL of *N,N*-dimethylformamide and a reaction time of 3 days. ^1H NMR (300 MHz, DMSO- d_6) δ 12.73 (s, 1H), 7.73 (dd, $J = 10.4, 1.8$ Hz, 1H), 7.66 – 7.52 (m, 2H).

[00402] **Example 72A:** 4-chloro-2-fluoro-*N*-hydroxybenzimidoyl chloride (compound **A-107**)




[00403] Following general procedure A2, compound **A-107** was prepared from 4-chloro-2-fluorobenzaldehyde:

[00404] Compound **A-106** (2.7 g, 96% yield) was prepared as a white solid from 4-chloro-2-fluorobenzaldehyde (2.6 g, 16.4 mmol) using 45 mL of ethanol/water and a reaction time of 1 hour. LCMS (2): $t_R=1.856$ min., (ES^+) m/z $(\text{M}+\text{H})^+ = 174.0$. ^1H NMR (300 MHz, DMSO- d_6) δ 11.83 (s, 1H), 8.16 (s, 1H), 7.87 – 7.63 (m, 1H), 7.62 – 7.41 (m, 1H), 7.31 (d, $J = 7.9$ Hz, 1H).

[00405] Compound **A-107** (2.7 g, 84% yield) was prepared as a white solid from **A-106** (2.7 g, 15.6 mmol) using 25 mL of *N,N*-dimethylformamide and a reaction time of 1.5 hours. ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.73 (s, 1H), 7.75 – 7.65 (m, 1H), 7.65 – 7.57 (m, 1H), 7.47 – 7.37 (m, 1H).


[00406] General Procedure B1: Synthesis and chiral separation of amino-benzisoxazoles.



[00407] To a solution of benzo[d]isoxazol-3-amine (1 eq.) and quinuclidin-3-one (1.1 eq.) in toluene (7 mL/mmol benzo[d]isoxazol-3-amine) at 25 °C was added portion-wise titanium(IV) isopropoxide (9 eq.). The resulting solution was stirred at 100 °C for 12 hours. On completion, the mixture was cooled to 0 °C, and ethanol (1 mL/mmol benzo[d]isoxazol-3-amine) was added via syringe, followed by sodium borohydride (3.7 eq.) in portions. The reaction was stirred at 25 °C for 3 hours, then quenched with saturated aqueous potassium carbonate solution, resulting in the formation of a solid. The mixture was filtered, and the filtrate was extracted with dichloromethane (5 × 50 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated in vacuo. The filter cake from the original filtration was slurried with methanol, and the mixture was filtered. The filtrate was directly evaporated to dryness. The combined residue from both batches was dissolved in 4N hydrochloric acid (20 mL) and stirred at room temperature for 4 hours. The mixture was made basic by addition of saturated potassium carbonate solution and extracted with dichloromethane (5 × 50 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated in vacuo. The residue was purified by prep-HPLC to give the racemic aminobenzoisoxazole product.

[00408] Chiral Separation: A solution of racemic aminobenzoisoxazole product in 3-5 mL of methanol was separated by cSFC (Waters SFC Prep 80, Column temperature: 25 °C, back pressure: 100 bar, and wavelength: 220 nm). Each set of collected fractions was concentrated at room temperature. The residue was dissolved in 0.2 M hydrochloric acid and lyophilized to give each enantiomer of the aminobenzoisoxazole product.


[00409] General Procedure C1: Synthesis of Aminobenzoisoxazoles.



[00410] N-hydroxybenzimidoyl chloride intermediate (1 eq.) in methanol (7 mL/mmol imidoyl chloride intermediate) was added dropwise over 30 min. to a solution of amine A-NH₂ (1.2-2 eq.) and triethylamine (2 eq) in methanol (5-10 mL/mmol imidoyl chloride intermediate) at room temperature. The resulting mixture was stirred at room temperature for 30 min. On completion, the reaction mixture was concentrated in vacuo and purified by prep-HPLC to give the N-hydroxyimide intermediate.

[00411] A mixture of N-hydroxyimide intermediate and base, in an appropriate solvent, was heated until the reaction was judged complete by LCMS. The mixture was filtered, concentrated in vacuum and purified by prep-HPLC to give the the aminobenzoisoxazole product.


[00412] **General Procedure C2: Synthesis of Aminobenzoisoxazoles.**



[00413] To a solution of amine A-NH₂ (1 eq.) and triethylamine (1 eq.) in methanol (3-5 mL/mmol amine A-NH₂) at room temperature was added dropwise a solution of -N-hydroxybenzimidoyl chloride (1 eq) in methanol (3-5 mL/mmol N-hydroxybenzimidoyl chloride). The mixture was stirred for 1 or more hours, then filtered and concentrated in vacuo. The residue was purified by silica gel column chromatography to afford the N-hydroxyimide intermediate.

[00414] A mixture of N-hydroxyimide intermediate and base, in an appropriate solvent, was heated until the reaction was judged complete by LCMS. The mixture was filtered, concentrated in vacuum and purified by silica gel column chromatography to give the aminobenzoisoxazole product.

[00415] **Example 1: N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine (*rac*-1)**



[00416] Following general procedure B1, *rac*-1 was prepared from benzo[d]isoxazol-3-amine (0.40 g, 3.0 mmol). The product was purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 150×25 mm, particle size: 10 μm; Mobile phase: 44-74% acetonitrile in H₂O (add 0.5% NH₃ · H₂O, v/v)] to give *rac*-1 (70 mg, 9% yield) as a yellow solid. LCMS (B): tR=1.179 min., (ES⁺) m/z (M+H)⁺ = 244.2.


[00417] Chiral Separation:

[00418] *rac*-1 (27 mg, 0.11 mmol) was separated by SFC (Column: Chiralpak AY-H -150×4.6mm, I.D., 5 μm; Mobile phase: ethanol (0.05% DEA) in CO₂) according to the chiral separation of general procedure B1 to give:

[00419] *N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer1 hydrochloride (compound 1-P1) (11 mg, 41% yield) as a white solid: cSFC analytical (A) tR=5.521 min., purity: 100.00%; LCMS (B): tR=0.115 min., (ES⁺) m/z (M+H)⁺ = 244.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.88-7.86 (d, J=7.6 Hz, 1H), 7.58-7.54 (m, 1H), 7.42-7.40 (d, J=8.4 Hz, 1H), 7.30-7.26 (t, J=7.6 Hz, 1H), 4.20-4.18 (m, 1H), 3.87-3.84 (m, 1H), 3.43-3.35 (m, 5H), 2.58-2.56 (m, 1H), 2.38-2.35 (m, 1H), 2.16-2.10 (m, 2H), 1.99-1.93 (m, 1H); and

[00420] *N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer2 hydrochloride (compound 1-P2) (11 mg, 41% yield) as a white solid: cSFC analytical (A) tR=7.926 min., purity: 98.82%; LCMS (J): tR=1.150 min., (ES⁺) m/z (M+H)⁺ = 244.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.90-7.88 (d, J=8.0 Hz, 1H), 7.61-7.57 (t, J=7.6 Hz, 1H), 7.45-7.43 (d, J=8.4 Hz, 1H), 7.32-7.29 (t, J=8.0 Hz, 1H), 4.20-4.18 (m, 1H), 3.90-3.86 (m, 1H), 3.48-3.33 (m, 5H), 2.58-2.56 (m, 1H), 2.38-2.36 (m, 1H), 2.16-2.10 (m, 2H), 1.97-1.96 (m, 1H).

[00421] **Example 2:** 7-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine (*rac*-2)



[00422] Following general procedure B1, *rac*-2 was prepared from A-1 (0.30 g, 1.8 mmol). The product was purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 150×25 mm, particle size: 10 μm; Mobile phase: 40-70% acetonitrile in H₂O (add 0.5% NH₃·H₂O, v/v)] to give *rac*-2 (170 mg, 34% yield) as an off-white solid. LCMS (B): tR=0.617 min., (ES⁺) m/z (M+H)⁺ = 278.1. ¹H-NMR (CD₃OD, 400 MHz): δ 7.81-7.78 (d, J=8.4 Hz, 1H), 7.55 (m, 1H), 7.34-7.32 (d, J=7.2 Hz, 1H), 4.18-4.15 (m, 1H), 3.90-3.84 (td, J=2.0 Hz, J=13.2 Hz, 1H), 3.44-3.38 (m, 4H), 3.29-3.26 (m, 1H), 2.55-2.53 (m, 1H), 2.35-2.34 (m, 1H), 2.16-2.12 (m, 2H), 2.09-1.97 (m, 1H).

[00423] Chiral Separation:


[00424] *rac*-2 (170 mg, 0.6 mmol) was separated by SFC (Column: Chiralpak AY-H -150×4.6mm, I.D., 5 μm; Mobile phase: ethanol (0.05% DEA) in CO₂) according to the chiral separation of general procedure B1 to give:

[00425] 7-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer1 hydrochloride (compound 2-P1) (40 mg, 47% yield) as a white solid: cSFC analytical (A) tR=6.113 min., purity: 100%; LCMS (B): tR=0.633 min., (ES⁺) m/z (M+H)⁺ = 278.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.80-7.78 (d, J=8.0 Hz, 1H), 7.61-7.59 (d, J=7.6 Hz, 1H), 7.69-7.67 (t, J=8.0 Hz, 1H), 4.19-4.17 (m, 1H), 3.89-3.83 (m, 1H), 3.44-3.35 (m, 4H), 3.28-3.27 (m, 1H), 2.56-2.54 (m, 1H), 2.34-2.33 (m, 2H), 2.14-2.10 (m, 1H), 2.08-1.96 (m, 1H); and

[00426] 7-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer2 hydrochloride (compound **2-P2**) (35 mg, 41% yield) as a white solid: cSFC analytical (A) tR=7.304 min., purity: 100.00%; LCMS (B): tR=0.636 min., (ES⁺) m/z (M+H)⁺ = 278.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.81-7.79 (d, J=8.0 Hz, 1H), 7.61-7.59 (d, J=7.2 Hz, 1H), 7.69-7.67 (t, J=8.0 Hz, 1H), 4.19-4.17 (m, 1H), 3.89-3.83 (m, 1H), 3.45-3.35 (m, 4H), 3.29-3.28 (m, 1H), 2.55-2.53 (m, 1H), 2.34-2.33 (m, 2H), 2.14-2.11 (m, 1H), 1.99-1.96 (m, 1H).

[00427] **Example 3:**

[00428] Preparation of 6-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine (**rac-3**)



[00429] Following general procedure B1, **rac-3** was prepared from **A-2** (0.30 g, 1.8 mmol). The product was purified by prep-HPLC [Instrument: PREP-A; Column: Phenomenex Gemini C18 150×21.2 mm, particle size: 5 μm; Mobile phase: 30-55% acetonitrile in H₂O (add 0.05% NH₃·H₂O, v/v)] to give **rac-3** (210 mg, 43% yield) as an off-white solid. LCMS (B): tR=0.631 min., (ES⁺) m/z (M+H)⁺ = 278.1. ¹H-NMR (CD₃OD, 400 MHz): δ 7.79-7.77 (d, J=8.0 Hz, 1H), 7.63-7.61 (d, J=7.6 Hz, 1H), 7.32-7.28 (m, 1H), 4.21-4.18 (m, 1H), 3.92-3.88 (m, 1H), 3.45-3.37 (m, 4H), 3.33-3.30 (m, 1H), 2.58-2.55 (m, 1H), 2.36-2.34 (m, 1H), 2.16-2.12 (m, 2H), 2.10-1.98 (m, 1H);


[00430] Chiral Separation:

[00431] **rac-3** (170 mg, 0.6 mmol) was separated by SFC (Column: Chiralpak AY-H -150×4.6mm, I.D., 5 μm; Mobile phase: ethanol (0.05% DEA) in CO₂) according to the chiral separation of general procedure B1 to give:

[00432] 6-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer1 hydrochloride (compound **3-P1**) (50 mg, 59% yield) as a white solid: cSFC analytical (A) tR=5.617 min., purity: 99.33%; LCMS (B): tR=0.683 min., (ES⁺) m/z (M+H)⁺ = 278.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.85-7.83 (d, J=8.8 Hz, 1H), 7.52-7.51 (d, J=1.6 Hz, 1H), 7.32-7.29 (dd, J=1.6 Hz, J=8.8 Hz, 1H), 4.20-4.18 (m, 1H), 3.87-3.84 (m, 1H), 3.43-3.35 (m, 5H), 2.58-2.56 (m, 1H), 2.38-2.35 (m, 1H), 2.16-2.10 (m, 2H), 1.99-1.93 (m, 1H); and

[00433] 6-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer2 hydrochloride (compound **3-P2**) (32 mg, 38% yield) as a white solid: cSFC analytical (A) tR=8.069 min., purity: 100.00%; LCMS (B): tR=0.681 min., (ES⁺) m/z (M+H)⁺ = 278.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.85-7.83 (d, J=8.4 Hz, 1H), 7.54-7.53 (d, J=1.6 Hz, 1H), 7.34-7.31 (dd, J=1.6 Hz, J=8.4 Hz, 1H), 4.18-4.16 (m, 1H), 3.90-3.83 (m, 1H), 3.36-3.33 (m, 4H), 3.46-3.37 (m, 1H), 2.56-2.54 (m, 1H), 2.36 (m, 1H), 2.15-2.11 (m, 2H), 1.98-1.97 (m, 1H).

[00434] Preparation of (*R*)-6-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine ((*R*)-**3**)




[00435] Following general procedure C2, compound (*R*)-3 was prepared from compound **A-107**:

[00436] Compound (*R*)-**A-107-1** (144 mg, 61% yield) was prepared as a white solid from **A-107** (165 mg, 0.8 mmol) and (*R*)-quinuclidin-3-amine (100 mg, 0.8 mmol) using 5 mL of methanol and a reaction time of 1 hour. The product was purified by silica gel column chromatography [chloroform: methanol = 1:0 to 17:3]. LCMS (1): *t*_R=2.965 min., (ES⁺) *m/z* (M+H)⁺ = 298.1.

[00437] To a solution of compound (*R*)-**A-107-1** (101 mg, 0.3 mmol) in *N,N*-dimethylacetamide (5 mL) was added potassium tertbutoxide (57 mg, 0.5 mmol). The mixture was stirred at room temperature for 20 hours. The solution was put on an SCX column and eluted with methanol. The product was eluted from the column using 3.5 M ammonia in methanol, concentrated and purified by silica gel column chromatography [chloroform: 7M NH₃ in methanol = 1/0 to 9/1]. The resulting product was lyophilized to afford:

[00438] Compound (*R*)-**3** (23 mg, 24% yield) as a white solid: cHPLC analytical [cHPLC analytical conditions: Column: Chiralcel OD-H, 250×4.6 mm, particle size: 5 μm; Flow: 1.0 mL/min; Column temp: 25 °C; Mobile phase: 0.1% diethylamine in Heptane/Ethanol = 8/2; detection: DAD (220-320 nm)] *t*_R=4.505 min., purity: 100%; LCMS (1): *t*_R=3.132 min., (ES⁺) *m/z* (M+H)⁺ = 278.0; ¹H NMR (400 MHz, CDCl₃), δ 7.48 – 7.40 (m, 2H), 7.24 – 7.19 (m, 1H), 4.28 (d, *J* = 4.6 Hz, 1H), 3.90 – 3.81 (m, 1H), 3.55 – 3.45 (m, 1H), 3.01 – 2.78 (m, 4H), 2.70 – 2.60 (m, 1H), 2.28 – 2.21 (m, 1H), 1.88 – 1.63 (m, 3H), 1.58 – 1.38 (m, 1H).

[00439] **Example 4: 5-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine (*rac*-4)**



[00440] Following general procedure B1, *rac*-4 was prepared from **A-3** (0.20 g, 1.2 mmol). The product was purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 150×25 mm, particle size: 10 μm; Mobile phase: 44-74% acetonitrile in H₂O (add 0.5% NH₃·H₂O, v/v)] to give *rac*-4 (45 mg, 14% yield) as a white solid. LCMS (J): *t*_R=1.346 min., (ES⁺) *m/z* (M+H)⁺ = 278.1.

[00441] Chiral Separation:


[00442] *rac*-4 (45 mg, 0.16 mmol) was separated by SFC (Column: Chiralpak AY-H -150×4.6mm, I.D., 5 μm; Mobile phase: ethanol (0.05% DEA) in CO₂) according to the chiral separation of general procedure B1 to give:

[00443] 5-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer1 hydrochloride (compound **4-P1**) (13 mg, 29% yield) as a white solid: cSFC analytical (D) *t*_R=5.053 min., purity:

100%; LCMS (B): tR=0.615 min., (ES⁺) m/z (M+H)⁺ = 278.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.94 (s, 1H), 7.56-7.50 (m, 1H), 7.44-7.38 (m, 1H), 4.19-4.17 (m, 1H), 3.89-3.83 (m, 1H), 3.49-3.36 (m, 5H), 2.56-2.55 (m, 1H), 2.36-2.36 (m, 1H), 2.13-2.12 (m, 2H), 1.99-1.96 (m, 1H); and

[00444] 5-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer2 hydrochloride (compound **4-P2**) (10 mg, 22% yield) as a white solid: cSFC analytical (D) tR=6.445 min., purity: 98.3%; LCMS (B): tR=0.616 min., (ES⁺) m/z (M+H)⁺ = 2278.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.92-7.92 (d, J=1.6 Hz, 1H), 7.57-7.55 (dd, J=8.8 Hz, J=1.6 Hz, 1H), 7.45-7.43 (d, J=8.8 Hz, 1H), 4.19-4.17 (m, 1H), 3.89-3.83 (m, 1H), 3.51-3.36 (m, 5H), 2.56-2.55 (m, 1H), 2.36-2.33 (m, 1H), 2.15-2.11 (m, 2H), 2.10-1.97 (m, 1H).

[00445] **Example 5:** 6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine (*rac*-**5**)



[00446] Following general procedure B1, *rac*-**5** was prepared from 6-methylbenzo[d]isoxazol-3-amine (0.50 g, 3.4 mmol) using 4 equivalents sodium borohydride and a reaction time of 16 hours for the first step. The product was purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 150×25 mm, particle size: 10 μm; Mobile phase: 44-74% acetonitrile in H₂O (add 0.5% NH₃·H₂O, v/v)] to give *rac*-**5** (0.70 g, 14% yield) as a white solid. LCMS (G): (ES⁺) m/z (M+H)⁺ = 258.0, tR=2.749 min.


[00447] Chiral Separation:

[00448] *rac*-**5** (0.10 g, 0.46 mmol) was separated by SFC (Column: Chiralpak AY-H -150×4.6mm, I.D., 5 μm; Mobile phase: ethanol (0.05% DEA) in CO₂) according to the chiral separation of general procedure B1 to give:

[00449] 6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer1 hydrochloride (compound **5-P1**) (40 mg, 40% yield) as a white solid: cSFC analytical (G) tR=2.61 min., purity: 96.98%; LCMS (J): tR=1.21 min., (ES⁺) m/z (M+H)⁺ = 258; ¹H-NMR (CD₃OD, 400 MHz): δ 7.72 (d, J=8.4 Hz, 1H), 7.24 (s, 1H), 7.14 (d, J=8.4 Hz, 1H), 4.17-4.15 (m, 1H), 3.88-3.82 (m, 1H), 3.48-3.27 (m, 5H), 2.56-2.53 (m, 1H), 2.50 (m, 3H), 2.37-2.34 (m, 1H), 2.15-2.09 (m, 2H), 1.98-1.92 (m, 1H); and

[00450] 6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer2 hydrochloride (compound **5-P2**) (45 mg, 45% yield) as a white solid: cSFC analytical (G) tR=3.05 min., purity: 96.94%; LCMS (J): tR=1.22 min., (ES⁺) m/z (M+H)⁺ = 258; ¹H-NMR (CD₃OD, 400 MHz): δ 7.71 (d, J=8.4 Hz, 1H), 7.25 (s, 1H), 7.14 (d, J=8.4 Hz, 1H), 4.17-4.16 (m, 1H), 3.87-3.82 (m, 1H), 3.48-3.27 (m, 5H), 2.56-2.53 (m, 1H), 2.50 (m, 3H), 2.37-2.34 (m, 1H), 2.15-2.09 (m, 2H), 1.98-1.93 (m, 1H).

[00451] **Example 6:** 6-cyclopropyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine (*rac*-**6**)



[00452] Following general procedure B1, *rac-6* was prepared from **A-4** (0.20 g, 1.1 mmol) using 10 eq. of titanium(IV) isopropoxide. The product was purified by prep-HPLC [Instrument: GX-I; Column: Xtimate C18 150x25 mm, particle size: 5 μm ; Mobile phase: 25-55% acetonitrile in H_2O (add 0.1% TFA, v/v)] to give *rac-6* (150 mg, 40% yield) as a white solid. LCMS (J): $t_R=1.425\text{ min.}$, $(\text{ES}^+) m/z (\text{M}+\text{H})^+ = 278.1$. $^1\text{H-NMR}$ (CD_3OD , 400 MHz): δ 7.70-7.68 (d, $J=8.4\text{ Hz}$, 1H), 7.06 (m, 1H), 7.00-6.98 (d, $J=8.4\text{ Hz}$, 1H), 3.79-3.77 (m, 1H), 3.39-3.35 (m, 1H), 2.98-2.95 (m, 1H), 2.88-2.81 (m, 3H), 2.77-2.72 (m, 1H), 2.21-2.19 (m, 1H), 2.07-2.03 (m, 2H), 1.97-1.95 (m, 1H), 1.78-1.76 (m, 1H), 1.55-1.49 (m, 1H), 1.07-1.05 (m, 2H), 0.79-0.77 (m, 2H);


[00453] Chiral Separation:

[00454] *rac-6* (60 mg, 0.2 mmol) was separated by SFC (Column: Chiralpak AD-3 -100x4.6mm, I.D., 3 μm ; Mobile phase: ethanol (0.05% DEA) in CO_2) according to the chiral separation of general procedure B1 to give:

[00455] 6-cyclopropyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer1 hydrochloride (compound **6-P1**) (21 mg, 67% yield) as a white solid: cSFC analytical (A) $t_R=1.010\text{ min.}$, purity: 98.24%; LCMS (B): $t_R=0.649\text{ min.}$, $(\text{ES}^+) m/z (\text{M}+\text{H})^+ = 284.2$; $^1\text{H-NMR}$ (CD_3OD , 400 MHz): δ 7.73-7.70 (m, 1H), 7.09 (s, 1H), 7.02-7.00 (d, $J=8.0\text{ Hz}$, 1H), 4.15-4.12 (m, 1H), 3.86-3.83 (m, 1H), 3.46-3.33 (m, 5H), 2.53-2.52 (m, 1H), 2.29 (m, 1H), 2.10-2.06 (m, 3H), 2.04 (m, 1H), 1.07-1.05 (m, 2H), 0.79-0.78 (m, 2H); and

[00456] 6-cyclopropyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer2 hydrochloride (compound **6-P2**) (22 mg, 73% yield) as a white solid: cSFC analytical (A) $t_R=1.633\text{ min.}$, purity: 95.20%; LCMS (B): $t_R=0.652\text{ min.}$, $(\text{ES}^+) m/z (\text{M}+\text{H})^+ = 284.2$; $^1\text{H-NMR}$ (CD_3OD , 400 MHz): δ 7.73-7.71 (d, $J=8.4\text{ Hz}$, 1H), 7.09 (s, 1H), 7.02-7.00 (d, $J=8.0\text{ Hz}$, 1H), 4.15-4.12 (m, 1H), 3.85-3.80 (m, 1H), 3.46-3.33 (m, 5H), 2.53-2.52 (m, 1H), 2.34-2.33 (m, 1H), 2.10-2.06 (m, 3H), 1.93 (m, 1H), 1.08-1.04 (m, 2H), 0.80-0.77 (m, 2H).

[00457] **Example 7:** 6-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine (*rac-7*)



[00458] Following general procedure B1, *rac-7* was prepared from compound **A-5** (0.50 g, 3.1 mmol). The first step was run at 110 $^\circ\text{C}$ for 48 hours, and the second step at 25 $^\circ\text{C}$ for 16 hours. The product was purified by prep-HPLC [Instrument: GX-B; Column: YMC-pack ODS-AQ 150x30mm, particle size: 5 μm ; Mobile phase: 10-40% acetonitrile in H_2O (add 0.1% TFA, v/v)] to give *rac-7* (0.60 g, 70% yield) as a white solid. LCMS (J): $t_R=1.135\text{ min.}$, $(\text{ES}^+) m/z (\text{M}+\text{H})^+ = 274.1$.


[00459] Chiral Separation:

[00460] *rac*-7 (0.30 g, 1.1 mmol) was separated by SFC (Column: Chiralpak AD-3 100×4.6mm, I.D., 10 μm; Mobile phase: 40% ethanol (0.05% DEA) in CO₂) according to the chiral separation of general procedure B1 to give:

[00461] 6-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer1 hydrochloride (compound 7-P1) (40 mg, 13% yield) as a white solid: cSFC analytical (B) tR=2.479 min., purity: 98.09%; LCMS (Q): tR=2.777 min., (ES⁺) m/z (M+H)⁺ = 274.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.73-7.71 (d, J=8.4 Hz, 1H), 6.97-6.96 (d, J=1.6 Hz, 1H), 6.91-6.88 (dd, J₁=8.8 Hz, J₂=2.0 Hz, 1H), 4.16-4.12 (m, 1H), 3.89(s, 3H), 3.85-3.82 (m, 1H), 3.51-3.37 (m, 4H), 3.33-3.30 (m, 1H), 2.54-2.52 (m, 1H), 2.36-2.35 (m, 1H), 2.15-2.09 (m, 2H), 1.96-1.92 (m, 1H); and

[00462] 6-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer2 hydrochloride (compound 7-P2) (40 mg, 13% yield) as a white solid: cSFC analytical (B) tR=2.771 min., purity: 97.76%; LCMS (Q): tR=2.770 min., (ES⁺) m/z (M+H)⁺ = 2740.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.75-7.73 (d, J=8.8 Hz, 1H), 6.97-6.96 (d, J=2.0 Hz, 1H), 6.91-6.88 (dd, J₁=8.8 Hz, J₂=2.0 Hz, 1H), 4.15-4.13 (m, 1H), 3.89(s, 3H), 3.85-3.82 (m, 1H), 3.52-3.37 (m, 4H), 3.33-3.30 (m, 1H), 2.55-2.52 (m, 1H), 2.39-2.35 (m, 1H), 2.15-2.08 (m, 2H), 1.98-1.92 (m, 1H).

[00463] **Example 8:** *N*-(quinuclidin-3-yl)-6-(trifluoromethyl)benzo[d]isoxazol-3-amine (*rac*-8)



[00464] Following general procedure B1, *rac*-8 was prepared from 6-(trifluoromethyl)benzo[d]isoxazol-3-amine (0.40 g, 2.0 mmol) using 4 equivalents of sodium borohydride and a reaction time of 16 hours for the first step. The product was purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 150×25 mm, particle size: 10 μm; Mobile phase: 28-58% acetonitrile in H₂O (add 0.5% NH₃·H₂O, v/v)] to give *rac*-8 (0.15 g, 24% yield) as a white solid. LCMS (G): (ES⁺) m/z (M+H)⁺ = 312.1, tR=2.84 min.

[00465] Chiral Separation:


[00466] *rac*-8 (0.12 g, 0.46 mmol) was separated by SFC (Column: Chiralpak AY-H -150×4.6mm, I.D., 5 μm; Mobile phase: ethanol (0.05% DEA) in CO₂) according to the chiral separation of general procedure B1 to give:

[00467] *N*-(quinuclidin-3-yl)-6-(trifluoromethyl)benzo[d]isoxazol-3-amine-enantiomer1 hydrochloride (compound 8-P1) (35 mg, 35% yield) as a white solid: cSFC analytical (G) tR=2.66 min., purity: 99.42%; LCMS (J): tR=1.42 min., (ES⁺) m/z (M+H)⁺ = 312.1; ¹H-NMR (CD₃OD, 400 MHz): δ 8.09-8.04 (m, 1H), 7.84 (s, 1H), 7.61 (d, J=8 Hz, 1H), 4.24-4.20 (m, 1H), 3.92-3.86 (m, 1H), 3.47-3.37 (m, 5H), 2.58-2.57 (m, 1H), 2.38-2.37 (m, 1H), 2.20-2.11 (m, 2H), 2.01-1.98 (m, 1H); and

[00468] *N*-(quinuclidin-3-yl)-6-(trifluoromethyl)benzo[d]isoxazol-3-amine-enantiomer2 hydrochloride (compound 8-P2) (45 mg, 45% yield) as a white solid: cSFC analytical (G) tR=3.70

min., purity: 99.39%; LCMS (J): $t_R=1.42$ min., (ES^+) m/z $(M+H)^+ = 312.1$; ^1H -NMR (CD_3OD , 400 MHz): δ 8.09-8.05 (m, 1H), 7.83 (s, 1H), 7.61 (d, $J=8$ Hz, 1H), 4.23-4.20 (m, 1H), 3.91-3.86 (m, 1H), 3.47-3.37 (m, 5H), 2.59-2.56 (m, 1H), 2.38-2.34 (m, 1H), 2.19-2.11 (m, 2H), 2.01-1.95 (m, 1H).

[00469] Example 9: 6,7-dichloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine (*rac*-9)



[00470] Following general procedure B1, *rac*-9 was prepared from **A-8** (0.26 g, 1.3 mmol) using 3 equivalents of sodium borohydride and a reaction time of 10 hours for the first step and 2 hours for the second. The product was purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 150×25 mm, particle size: 10 μ m; Mobile phase: 44-74% acetonitrile in H_2O (add 0.5% $NH_3 \cdot H_2O$, v/v)] to give *rac*-9 (0.10 g, 40% yield) as a white solid. LCMS (B): (ES^+) m/z $(M+H)^+ = 311.9$. $t_R=0.71$ min.


[00471] Chiral Separation:

[00472] *rac*-9 (0.10 g, 0.49 mmol) was separated by SFC (Column: Chiralpak AY-H -150×4.6mm, I.D., 5 μ m; Mobile phase: ethanol (0.05% DEA) in CO_2) according to the chiral separation of general procedure B1 to give:

[00473] 6,7-dichloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer1 hydrochloride (compound **9-P1**) (50 mg, 50% yield) as a white solid: cSFC analytical (G) $t_R=2.70$ min., purity: 98.53%; LCMS (B): $t_R=0.68$ min., (ES^+) m/z $(M+H)^+ = 312.1$; ^1H -NMR (CD_3OD , 400 MHz): δ 7.80 (d, $J=8.4$ Hz, 1H), 7.47 (d, $J=8.4$ Hz, 1H), 4.19-4.17 (m, 1H), 3.90-3.84 (m, 1H), 3.51-3.37 (m, 5H), 2.57-2.54 (m, 1H), 2.38-2.34 (m, 1H), 2.16-2.09 (m, 2H), 1.99-1.94 (m, 1H); and

[00474] 6,7-dichloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine-enantiomer2 hydrochloride (compound **9-P2**) (40 mg, 40% yield) as a white solid: cSFC analytical (G) $t_R=3.44$ min., purity: 99.72%; LCMS (B): $t_R=0.69$ min., (ES^+) m/z $(M+H)^+ = 312.1$; ^1H -NMR (CD_3OD , 400 MHz): δ 7.81 (d, $J=8.4$ Hz, 1H), 7.45 (d, $J=8$ Hz, 1H), 4.19-4.17 (m, 1H), 3.90-3.84 (m, 1H), 3.50-3.45 (m, 5H), 2.57-2.54 (m, 1H), 2.39-2.35 (m, 1H), 2.16-2.09 (m, 2H), 2.00-1.97 (m, 1H).

[00475] Example 10: (*R*)-4-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-10)



[00476] Following general procedure C1, compound (*R*)-10 was prepared from compound **A-10**:


[00477] Compound (*R*)-A-10-1 (0.1 g, white solid, 42% yield over two steps) was prepared from compound **A-10** (0.3 g, 1.46 mmol) and (*R*)-quinuclidin-3-amine (0.25 g, 2.0 mmol). The product was

purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 250×50 mm, particle size: 10 μm; Mobile phase: 22-52% acetonitrile in H₂O (add 0.5% NH₃·H₂O, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 298, tR=0.866 min.

[00478] A mixture of compound (*R*)-**A-10-1** (100 mg, 0.34 mmol) and potassium carbonate (186 mg, 1.4 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 120 °C for 2 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250x50 mm, particle size: 10 μm; Mobile phase: 38-68% acetonitrile in H₂O (add 0.05% ammonia-ACN, v/v)]. The resulting solid was dissolved in 0.2 M hydrochloric acid and again lyophilized to give:

[00479] Compound (*R*)-**10** (30 mg, 29% yield) as a white solid: cSFC analytical tR=2.97 min., purity: 96.01%; LCMS (EE): tR=2.55 min., 278.1 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.54-7.53 (t, J=8 Hz, 1H), 7.43 (d, J=8.4 Hz, 1H), 7.30 (d, J=8.4 Hz, 1H), 4.23-4.20 (m, 1H), 3.89-3.82 (m, 1H), 3.51-3.37 (m, 5H), 2.63-2.61 (m, 1H), 2.36-2.31 (m, 1H), 2.16-2.10 (m, 2H), 2.00-1.99 (m, 1H).

[00480] **Example 11:** (*R*)-6-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**11**)




[00481] Following general procedure C1, compound (*R*)-**11** was prepared from compound **A-12**:

[00482] Compound (*R*)-**A-12-1** (0.10 g, white solid, 31% yield over two steps) was prepared from compound **A-12** (0.20 g, 1.0 mmol) and (*R*)-quinuclidin-3-amine (0.25 g, 2.0 mmol), using *N,N*-dimethylformamide as the solvent instead of methanol. The product was purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 250×50 mm, particle size: 10 μm; Mobile phase: 19-49% acetonitrile in H₂O (add 0.5% NH₃·H₂O, v/v)]. LCMS (B): (ES⁺) m/z (M+H)⁺ = 282.1, tR=2.387 min.

[00483] A mixture of compound (*R*)-**A-12-1** (0.10 g, 0.36 mmol) and potassium tert-butoxide (60 mg, 0.53 mmol) in *N,N*-dimethylacetamide (5 mL) was stirred at room temperature for 1 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: YMC-Actus Pro C18 150x30 mm, particle size: 5 μm; Mobile phase: 12-42% acetonitrile in H₂O (add 0.1% TFA, v/v)]. The resulting solid was dissolved in 0.2 M hydrochloric acid and again lyophilized to give:

[00484] Compound (*R*)-**11** (10 mg, 10% yield) as a yellow solid: cSFC analytical (A) tR=4.76 min., purity: 100%; LCMS (J): tR=1.259 min., (ES⁺) m/z (M+H)⁺ = 262.2; ¹H-NMR (CD₃OD, 400 MHz): δ 7.88-7.84 (m, 1H), 7.25-7.23 (d, J=8.8 Hz, 1H), 7.13-7.09 (t, J=8.8 Hz, 1H), 4.16-4.15 (m, 1H), 3.89-3.83 (t, J=11.6 Hz, 1H), 3.42-3.38 (m, 4H), 3.29 (m, 1H), 2.55-2.54 (m, 1H), 2.36 (m, 1H), 2.13-2.12 (m, 2H), 1.96 (m, 1H).

[00485] Example 12: (*R*)-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-6-carbonitrile hydrochloride ((*R*)-12)



[00486] Following general procedure C1, compound (*R*)-12 was prepared from compound A-14:


[00487] Compound (*R*)-A-14-1 (0.15 g, white solid, 42% yield over two steps) was prepared from compound A-14 (0.30 g, 1.0 mmol) and (*R*)-quinuclidin-3-amine (0.25 g, 2.0 mmol). The product was purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 250×50 mm, particle size: 10 μm; Mobile phase: 19-49% acetonitrile in H₂O (add 0.5% NH₃·H₂O, v/v)]. LCMS (B): (ES⁺) *m/z* (M+H)⁺ = 289.2, *t*_R=0.928 min.

[00488] A mixture of compound (*R*)-A-14-1 (0.13 g, 0.45 mmol) and potassium carbonate (0.19 g, 1.4 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 110 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: YMC-Actus Pro C18 150x30 mm, particle size: 5 μm; Mobile phase: 17-74% acetonitrile in H₂O (add 0.5% HCl, v/v)]. The product was lyophilized to give:

[00489] Compound (*R*)-12 (30 mg, 23% yield) as a white solid: cSFC analytical (D) *t*_R=2.15 min., purity: 99.44%; LCMS (EE): *t*_R=2.39 min., 269.1 *m/z* (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 8.08 (d, *J*=8 Hz, 1H), 7.94 (s, 1H), 7.64-7.62 (dd, *J*=8 Hz, *J*=0.8 Hz, 1H), 4.22-4.20 (m, 1H), 3.91-3.85 (m, 1H), 3.51-3.35 (m, 5H), 2.58-2.56 (m, 1H), 2.37-2.35 (m, 1H), 2.16-2.10 (m, 2H), 1.99-1.97 (m, 1H).

[00490] Example 13:

[00491] Preparation: (*R*)-6-(methylsulfonyl)-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-13)



[00492] Following general procedure C1, compound (*R*)-13 was prepared from compound A-16:


[00493] Compound (*R*)-A-16-1 (0.35 g, pale yellow solid, 49% yield over two steps) was prepared from compound A-16 (0.30 g, 1.2 mmol) and (*R*)-quinuclidin-3-amine (0.18 g, 1.4 mmol), using *N,N*-dimethylformamide as the solvent instead of methanol. The product was purified by silica gel chromatography [DCM : MeOH = 5:1]. LCMS (B): (ES⁺) *m/z* (M+H)⁺ = 342.1, *t*_R=0.894 min.

[00494] A mixture of compound (*R*)-A-16-1 (0.10 g, 0.29 mmol) and potassium tert-butoxide (36 mg, 0.32 mmol) in tetrahydrofuran (5 mL) was stirred at room temperature for 6 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument:

GX-G; Column: Phenomenex Synergi Max-RP C18 250x80 mm, particle size: 10 μ m; Mobile phase: 1-31% acetonitrile in H₂O (add 0.2% TFA, v/v)]. The product was lyophilized, dissolved in 0.2 N hydrochloric acid and again lyophilized to give:

[00495] Compound (*R*)-13 (15 mg, 14% yield) as a white solid: cSFC analytical (A) t_R=6.79 min., purity: 100%; LCMS (W): t_R=0.740 min., (ES⁺) m/z (M+H)⁺ = 322.1; ¹H-NMR (CD₃OD, 400 MHz): δ 8.12-8.09 (d, J=8.4 Hz, 1H), 8.07 (s, 1H), 7.88-7.87 (dd, J=1.2 Hz, J=8.4 Hz, 1H), 4.21-4.19 (m, 1H), 3.90-3.85 (m, 1H), 3.45-3.34 (m, 5H), 3.21 (s, 3H), 2.57-2.54 (m, 1H), 2.36-2.35 (m, 1H), 2.14-2.10 (m, 2H), 2.04-1.93 (m, 1H).

[00496] Preparation: (*S*)-6-(methylsulfonyl)-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*S*)-13)




[00497] Following general procedure C1, compound (*S*)-13 was prepared from compound A-16:

[00498] Compound (*S*)-A-16-1 (0.15 g, pale yellow solid, 36% yield over two steps) was prepared from compound A-16 (0.30 g, 1.2 mmol) and (*S*)-quinuclidin-3-amine (0.18 g, 1.4 mmol), using N, N-dimethylformamide as the solvent instead of methanol. The product was purified by silica gel chromatography [DCM / MeOH = 5/1]. LCMS (B): (ES⁺) m/z (M+H)⁺ = 342.1, t_R=0.852 min.

[00499] A mixture of compound (*S*)-A-16-1 (0.30 g, 0.88 mmol) and potassium tert-butoxide (0.11 g, 0.97 mmol) in tetrahydrofuran (5 mL) was stirred at room temperature for 12 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-G; Column: Phenomenex Synergi Max-RP C18 250x80 mm, particle size: 10 μ m; Mobile phase: 1-31% acetonitrile in H₂O (add 0.2% TFA, v/v)]. The resulting solid was dissolved in 0.2 N hydrochloric acid and again lyophilized to give:

[00500] Compound (*S*)-13 (25 mg, 9% yield) as a white solid: cSFC analytical (A) t_R=8.80 min., purity: 100%; LCMS (W): t_R=0.708 min., (ES⁺) m/z (M+H)⁺ = 322.1; ¹H-NMR (CD₃OD, 400 MHz): δ 8.11-8.09 (d, J=8.4 Hz, 1H), 8.07 (s, 1H), 7.88-7.86 (d, J=8.4 Hz, 1H), 4.16-4.14 (m, 1H), 3.84-3.78 (m, 1H), 3.35-3.25 (m, 5H), 3.20 (s, 3H), 2.52-2.50 (m, 1H), 2.30-2.29 (m, 1H), 2.10-2.06 (m, 2H), 1.91-1.89 (m, 1H).

[00501] **Example 14:** (*R*)-6-(tert-butyl)-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-14)



[00502] Following general procedure C1, compound (*R*)-14 was prepared from compound A-20:


[00503] Compound (*R*)-**A-20-1** (60 mg, white solid, 14% yield over two steps) was prepared from compound **A-20** (0.30 g, 1.3 mmol) and (*R*)-quinuclidin-3-amine (0.16 g, 1.3 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Synergi C18 250x50 mm, particle size: 10 μ m; Mobile phase: 25-55% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 320.2, t_R=1.301 min.

[00504] A mixture of compound (*R*)-**A-20-1** (50 mg, 0.16 mmol) and potassium carbonate (66 mg, 0.48 mmol) in *N,N*-dimethylformamide (4.0 mL) was stirred at 120 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: YMC-Actus Pro C18 150x30 mm, particle size: 5 μ m; Mobile phase: 17-74% acetonitrile in H₂O (add 0.5% HCl, v/v)]. The product was lyophilized to give:

[00505] Compound (*R*)-**14** (6.5 mg, 14% yield) as a white solid: cSFC analytical (D) t_R=2.62 min., purity: 100%; LCMS (EE): t_R=2.99 min., 300.2 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.79-7.77 (d, J=8.4 Hz, 1H), 7.44-7.43 (s, 1H), 7.40-7.40 (m, 1H), 4.17-4.15 (m, 1H), 3.89-3.83 (t, J=11.6 Hz, 1H), 3.43-3.33 (m, 5H), 2.55-2.54 (m, 1H), 2.36 (m, 1H), 2.15-2.11 (m, 2H), 1.96 (m, 1H), 1.40-1.36 (m, 9H).

[00506] **Example 15:**

[00507] Preparation: (*R*)-5,6-dichloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**15**)




[00508] Following general procedure C1, compound (*R*)-**15** was prepared from compound **A-24**:

[00509] Compound (*R*)-**A-24-1** (30 mg, white solid, 22% yield over two steps) was prepared from compound **A-24** (0.10 g, 0.41 mmol) and (*R*)-quinuclidin-3-amine (62 mg, 0.49 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Synergi C18 250x50 mm, particle size: 10 μ m; Mobile phase: 25-55% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (X): (ES⁺) m/z (M+H)⁺ = 332.1, t_R=1.178 min.

[00510] A mixture of compound (*R*)-**A-24-1** (75 mg, 0.23 mmol) and potassium carbonate (94 mg, 0.68 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 110 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: YMC-Actus Pro C18 150x30 mm, particle size: 5 μ m; Mobile phase: 17-74% acetonitrile in H₂O (add 0.5% HCl, v/v)]. The product was lyophilized to give:

[00511] Compound (*R*)-**15** (30 mg, 43% yield) as a white solid: cSFC analytical t_R=5.47 min., purity: 100%; LCMS (X): t_R=0.16 min., 312.0 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 8.09 (s, 1H), 7.73 (s, 1H), 4.18-4.16 (m, 1H), 3.89-3.83 (m, 1H), 3.51-3.31 (m, 5H), 2.55-2.53 (m, 1H), 2.38-2.34 (m, 1H), 2.16-2.09 (m, 2H), 2.08-1.96 (m, 1H).

[00512] Preparation: (*S*)-5,6-dichloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*S*)-15)




[00513] Following general procedure C1, compound (*S*)-15 was prepared from compound A-24:

[00514] Compound (*S*)-A-24-1 (0.30 g, white solid, 44% yield over two steps) was prepared from compound A-24 (0.50 g, 2.1 mmol) and (*S*)-quinuclidin-3-amine (0.26 g, 2.1 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250x50 mm, particle size: 10 μ m; Mobile phase: 25-55% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 332.1, tR = 1.166 min.

[00515] A mixture of compound (*S*)-A-24-1 (0.20 g, 0.60 mmol) and potassium carbonate (0.25 g, 1.8 mmol) in *N,N*-dimethylacetamide (8.0 mL) was stirred at 110 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Synergi C18 150x30 mm, particle size: 4 μ m; Mobile phase: 25-55% acetonitrile in H₂O (add 0.5% HCl, v/v)]. The product was lyophilized to give:

[00516] Compound (*S*)-15 (0.12 g, 64% yield) as a white solid: cSFC analytical (D) tR = 1.977 min., purity: 99.16%; LCMS (B): tR = 0.683 min., (ES⁺) m/z (M+H)⁺ = 312.1; ¹H-NMR (CD₃OD, 400 MHz): δ 8.06 (s, 1H), 7.75 (s, 1H), 4.18-4.15 (m, 1H), 3.89-3.83 (m, 1H), 3.49-3.27 (m, 5H), 2.55-2.53 (m, 1H), 2.38-2.34 (m, 1H), 2.16-2.09 (m, 2H), 1.99-1.96 (m, 1H).

[00517] **Example 16:** (*R*)-*N*-(quinuclidin-3-yl)-6-(trifluoromethoxy)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-16)




[00518] Following general procedure C1, compound (*R*)-16 was prepared from compound A-26:

[00519] Compound (*R*)-A-26-1 (0.22 g, white solid, 30% yield over three steps) was prepared from compound A-26 (0.55 g, 2.5 mmol), triethylamine (0.42 g, 4.2 mmol) and (*R*)-quinuclidin-3-amine (0.26 g, 2.1 mmol), using *N,N*-dimethylformamide as the solvent instead of methanol. The reaction time was 10 hours. The product was purified by prep-HPLC [Instrument: GX-H; Column: Waters Xbridge C18 150x25 mm, particle size: 5 μ m; Mobile phase: 26-56% acetonitrile in H₂O (add 0.5% NH₃·H₂O, v/v)]. ¹H-NMR (CDCl₃, 400 MHz): δ 7.43 (t, J = 8.0 Hz, 1H), 7.05 (m, 2H), 5.63 (d, J = 8.0 Hz, 1H), 3.13-3.09 (m, 2H), 2.93-2.77 (m, 3H), 2.63-2.57 (m, 2H), 1.99-1.84 (m, 2H), 1.68-1.60 (m, 2H), 1.48-1.46 (m, 1H), 1.34-1.33 (m, 1H). LCMS (J): (ES⁺) m/z (M+H)⁺ = 348.0, tR = 1.052 min.

[00520] A mixture of compound (*R*)-**A-26-1** (0.25 g, 0.52 mmol) and potassium tert-butoxide (88 mg, 0.79 mmol) in *N,N*-dimethylacetamide (26 mL) was stirred at 25 °C for 2 hours. The mixture was diluted with ethyl acetate (260 mL), washed with brine (6 × 40 mL), dried with anhydrous sodium sulfate, filtered and concentrated in vacuo. The product was purified twice by prep-HPLC [Instrument: GX-D; Column: Boston Green ODS 150×30 mm, particle size: 5 μm; Mobile phase: 6-42% acetonitrile in H₂O (add 0.225% FA, v/v), and Instrument: GX-B; Column: Phenomenex Synergi C18 150×30 mm, particle size: 4 μm; Mobile phase: 23-53% acetonitrile in H₂O (add 0.1% TFA, v/v)], treated with 0.2 M hydrochloric acid and lyophilized to give:

[00521] Compound(*R*)-**16** (32 mg, 17% yield) as a yellow solid : cSFC analytical (D) tR=1.64 min., purity: 100%; LCMS (EE): tR=2.820 min., (ES⁺) m/z (M+H)⁺ = 328.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.95 (d, J=8.4 Hz, 1H), 7.38 (s, 1H), 7.19 (d, J=8.4 Hz, 1H), 4.14 (t, J=9.2 Hz, 1H), 3.82 (t, J=10.6 Hz, 1H), 3.47-3.28 (m, 5H), 2.51 (d, J=3.2 Hz, 1H), 2.34-2.32 (m, 1H), 2.13-2.07 (m, 2H), 1.95-1.92 (m, 1H).

[00522] **Example 17:** (*R*)-7-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**17**)




[00523] Following general procedure C1, compound (*R*)-**17** was prepared from compound **A-28**:

[00524] Compound (*R*)-**A-28-1** (0.20 g, white solid, 27% yield over two steps) was prepared from compound **A-28** (0.50 g, 2.6 mmol) and (*R*)-quinuclidin-3-amine (0.40 g, 3.1 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250x50 mm, particle size: 10 μm; Mobile phase: 20-50% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 282.1, tR= 0.999 min.

[00525] A mixture of compound (*R*)-**A-28-1** (0.20 g, 0.71 mmol) and potassium tert-butoxide (0.30 g, 2.0 mmol) in *N,N*-dimethylacetamide (5.0 mL) was stirred at 110 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: Phenomenex Synergi C18 150x30 mm, particle size: 4 μm; Mobile phase: 10-40% acetonitrile in H₂O (add 0.5% HCl, v/v)]. The product was lyophilized to give:

[00526] Compound (*R*)-**17** (0.10 g, 47% yield) as a white solid: cSFC analytical (D) tR=2.22 min., purity: 94%; LCMS (EE): tR=2.50 min., 262.1 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.69-7.67 (d, J=8.0 Hz, 1H), 7.39-7.34 (m, 1H), 7.30-7.26 (m, 1H), 4.21-4.18 (m, 1H), 3.91-3.85 (m, 1H), 3.49-3.31 (m, 5H), 2.58-2.56 (m, 1H), 2.40-2.35 (m, 1H), 2.16-2.10 (m, 2H), 1.99-1.97 (m, 1H).

[00527] **Example 18:** (*R*)-5-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**18**) and (*S*)-5-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*S*)-**18**)



[00528] Following general procedure C1, compound **(R)-18** was prepared from compound **A-30**:

[00529] Compound **(R)-A-30-1** (0.20 g, white solid, 27% yield over three steps) was prepared from compound **A-30** (0.50 g, 2.6 mmol) and **(R)**-quinuclidin-3-amine (0.40 g, 3.1 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250x50 mm, particle size: 10 μ m; Mobile phase: 20-50% acetonitrile in H₂O (add 0.5% ammonia, v/v)].

LCMS (J): (ES⁺) m/z (M+H)⁺ = 282.2, tR = 0.964 min.

[00530] A mixture of compound **(R)-A-30-1** (0.30 g, 1.1 mmol) and potassium carbonate (0.44 g, 3.2 mmol) in *N,N*-dimethylformamide (5.0 mL) was stirred at 110 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: Phenomenex Synergi C18 150x30 mm, particle size: 4 μ m; Mobile phase: 12-42% acetonitrile in H₂O (add 0.5% HCl, v/v)]. The product was lyophilized to give:

[00531] Compound **(R)-18** (0.15 g, 37% yield) as a white solid: cSFC analytical (D) tR = 2.027 min., chiral purity: 90%;

[00532] Chiral Purification:


[00533] A solution of **(R)-18** at 90% chiral purity (0.12 g, 0.46 mmol) in 3 mL of methanol was purified by cSFC (Instrument: SFC A; Column: AY-10 μ m; Mobile phase: 50% methanol (0.01% NH₃·H₂O) in CO₂) at room temperature. In addition to the major product compound **(R)-18**, the minor product compound **(S)-18** was also collected. Each set of collected fractions was concentrated at room temperature and lyophilized. The resulting solids were dissolved in 0.2 M hydrochloric acid and again lyophilized to give:

[00534] Compound **(R)-18** (80 mg, 59% yield) as a white solid : cSFC analytical (D) tR = 2.038 min., purity: 100%; LCMS (EE): tR = 2.464 min., (ES⁺) m/z (M+H)⁺ = 262.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.59-7.56 (dd, J = 8.0 Hz, J = 2.4 Hz, 1H), 7.47-7.44 (dd, J = 8.8 Hz, J = 4.0 Hz, 1H), 7.40-7.35 (m, 1H), 4.19-4.16 (m, 1H), 3.90-3.84 (m, 1H), 3.47-3.29 (m, 5H), 2.56-2.54 (m, 1H), 2.37-2.33 (m, 1H), 2.16-2.10 (m, 2H), 1.99-1.95 (m, 1H); and

[00535] Compound **(S)-18** (8.0 mg, 6% yield) as a white solid : cSFC analytical (D) tR = 2.673 min., purity: 97.2%; LCMS (EE): tR = 2.438 min., (ES⁺) m/z (M+H)⁺ = 262.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.61-7.58 (dd, J = 8.0 Hz, J = 2.4 Hz, 1H), 7.47-7.44 (dd, J = 8.4 Hz, J = 4.0 Hz, 1H), 7.40-7.35 (m, 1H), 4.19-4.16 (m, 1H), 3.90-3.83 (m, 1H), 3.48-3.30 (m, 5H), 2.56-2.54 (m, 1H), 2.39-2.33 (m, 1H), 2.18-2.09 (m, 2H), 1.98-1.96 (m, 1H);

[00536] cSFC analytical conditions: Column: Chiralcel AY-3 100x4.6mm, I.D., 3 μ m; Mobile phase: ethanol (0.05% DEA) in CO₂ from 5% to 40%; Flow rate: 3 mL/min.; Back pressure: 120 bar.

[00537] **Example 19:** (*R*)-4-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-19)




[00538] Following general procedure C1, compound (*R*)-19 was prepared from compound A-32:

[00539] Compound (*R*)-A-32-1 (0.60 g, white solid, crude) was prepared from compound A-32 (0.40 g, 2.1 mmol) and (*R*)-quinuclidin-3-amine (0.26 g, 2.1 mmol). The reaction time was 2 hours. LCMS (Q): (ES⁺) *m/z* (M+H)⁺ = 282.1, *t*R = 2.310 min.

[00540] A mixture of compound (*R*)-A-32-1 (0.30 g, 1.1 mmol) and potassium carbonate (0.44 g, 3.2 mmol) in *N,N*-dimethylformamide (10 mL) was stirred at 120 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 250x50, particle size: 10 μm; Mobile phase: 29-59% acetonitrile in H₂O (add 0.05% NH₃·H₂O, v/v)]. The resulting solid was dissolved in 0.2 N hydrochloric acid and again lyophilized to give:

[00541] Compound (*R*)-19 (90 mg, 28% yield over three steps) as a yellow solid: cSFC analytical (D) *t*R = 2.672 min., purity: 97.27%; LCMS (EE): *t*R = 2.383 min., (ES⁺) *m/z* (M+H)⁺ = 262.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.61-7.56 (m, 1H), 7.30-7.28 (d, *J* = 8.0 Hz, 1H), 7.03-7.98 (t, *J* = 10.0 Hz, 1H), 4.21-4.18 (m, 1H), 3.87-3.81 (m, 1H), 3.50-3.35 (m, 5H), 2.61-2.59 (m, 1H), 2.34-2.32 (m, 1H), 2.15-2.09 (m, 2H), 1.98-1.92 (m, 1H).

[00542] **Example 20:** (*R*)-7-fluoro-6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-20)



[00543] Following general procedure C1, compound (*R*)-20 was prepared from compound A-34:


[00544] Compound (*R*)-A-34-1 (0.11 g, white solid, 26% yield over two steps) was prepared from compound A-34 (0.30 g, 1.5 mmol) and (*R*)-quinuclidin-3-amine (0.18 g, 1.5 mmol). The product was purified by prep-HPLC [Instrument: GX-H; Waters Xbridge 150x25, particle size: 5 μm; Mobile phase: 20-50% acetonitrile in H₂O (add 0.05% NH₃·H₂O, v/v)]. LCMS (J): (ES⁺) *m/z* (M+H)⁺ = 296.1, *t*R = 1.108 min.

[00545] A mixture of compound (*R*)-A-34-1 (80 mg, 0.27 mmol) and potassium carbonate (0.11 g, 0.81 mmol) in *N,N*-dimethylformamide (10 mL) was stirred at 120 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250x50, particle size: 10 μm; Mobile phase: 30-60%

acetonitrile in H₂O (add 0.05% NH₃·H₂O, v/v)]. The resulting product was lyophilized, dissolved in 0.2 N hydrochloric acid and again lyophilized to give:

[00546] Compound **(R)**-**20** (30 mg, 36% yield) as a yellow solid: cSFC analytical (D) t_R=2.377 min., purity: 96.68%; LCMS (EE): t_R=2.659 min., (ES⁺) m/z (M+H)⁺ = 276.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.52-7.50 (d, J=8.0 Hz, 1H), 7.18-7.15 (t, J=7.6 Hz, 1H), 4.18-4.15 (m, 1H), 3.90-3.83 (m, 1H), 3.46-3.38 (m, 4H), 3.29-3.28 (m, 1H), 2.56-2.54 (m, 1H), 2.43-2.42 (m, 3H), 2.38-2.34 (m, 1H), 2.16-2.09 (m, 2H), 1.99-1.95 (m, 1H).

[00547] **Example 21:** *(R)*-6,7-difluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**21**)




[00548] Following general procedure C1, compound **(R)**-**21** was prepared from compound **A-36**:

[00549] Compound **(R)**-**A-36-1** (0.11 g, white solid, 77% yield over two steps) was prepared from compound **A-36** (0.10 g, 0.48 mmol) and *(R)*-quinuclidin-3-amine (0.12 g, 1.0 mmol) with a reaction time of 3 hours. The product was purified by prep-HPLC [Instrument: GX-H; Column: Waters Xbridge 150×25 mm, particle size: 5 μm; Mobile phase: 20-5% acetonitrile in H₂O (add 0.05% ammonia-ACN, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 300.1, t_R=1.1min.

[00550] A mixture of compound **(R)**-**A-36-1** (0.11 g, 0.45 mmol) and potassium carbonate (0.19 g, 1.4 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 120 °C for 2 hours. On completion, the reaction mixture was filtered. The reaction mixture was concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: YMC-Actus Pro C18 150×30 mm, particle size: 5 μm; Mobile phase: 17-74% acetonitrile in H₂O (add 0.5% HCl, v/v)]. The product was lyophilized to give:

[00551] Compound **(R)**-**21** (35 mg, 34% yield) as a white solid: cSFC analytical t_R=1.87 min., purity: 97.57%; LCMS (GG): t_R=1.74 min., 280.1 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.70-7.86 (m, 1H), 7.28-7.22 (m, 1H), 4.18-4.16 (m, 1H), 3.90-3.83 (m, 1H), 3.58-3.36 (m, 5H), 2.57-2.54 (m, 1H), 2.38-2.35 (m, 1H), 2.17-2.09 (m, 2H), 1.98-1.95 (m, 1H).

[00552] **Example 22:** *(R)*-5-fluoro-6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**22**)



[00553] Following general procedure C1, compound **(R)**-**22** was prepared from compound **A-40**:


[00554] Compound **(R)**-**A-40-1** (0.14 g, white solid, 16% yield over four steps) was prepared from compound **A-40** (0.50 g, 2.4 mmol) and *(R)*-quinuclidin-3-amine (0.31 g, 2.4 mmol). The reaction

time was 0.5 hours. The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Synergi C18 250x50 mm, particle size: 10 μ m; Mobile phase: 25-55% acetonitrile in H₂O (add 0.05% ammonia-ACN, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 296.2, tR=1.092 min.

[00555] A mixture of compound (*R*)-**A-40-1** (90 mg, 0.30 mmol) and potassium phosphate (0.19 g, 0.91 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 150 °C for 0.5 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: YMC-Actus Pro C18 150x30 mm, particle size: 4 μ m; Mobile phase: 25-55% acetonitrile in H₂O (add 0.1% TFA-ACN, v/v)]. The resulting solid was dissolved in 0.2 M hydrochloric acid and again lyophilized to give:

[00556] Compound (*R*)-**22** (35 mg, 37% yield) as a white solid: cSFC analytical (D) tR=5.47 min., purity: 98.28%; LCMS (FF): tR=2.07 min., 276.1 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.54 (d, J=8.8 Hz, 1H), 7.32 (d, J=5.6 Hz, 1H), 4.17-4.14 (m, 1H), 3.88-3.82 (m, 1H), 3.47-3.36 (m, 4H), 3.31-3.30 (m, 1H), 2.55-2.53 (m, 1H), 2.38-2.35 (m, 1H), 2.15-2.08 (m, 2H), 1.97-1.92 (m, 1H).

[00557] **Example 23:** (*R*)-6-chloro-7-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**23**)




[00558] Following general procedure C1, compound (*R*)-**23** was prepared from compound **A-43**:

[00559] Compound (*R*)-**A-43-1** (0.14 g, white solid, 20% yield over four steps) was prepared from compound **A-43** (0.45 g, 2.3 mmol) and (*R*)-quinuclidin-3-amine (0.29 g, 2.3 mmol). The reaction was stirred at 18 °C for 2 hours. The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Synergi C18 250x50 mm, particle size: 10 μ m; Mobile phase: 27-58% acetonitrile in H₂O (add 0.05% ammonia-ACN, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 312.2, tR=1.172 min.

[00560] A mixture of compound (*R*)-**A-43-1** (90 mg, 0.30 mmol) and potassium phosphate (0.19 g, 0.91 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 150 °C for 0.5 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-I; Column: Welch Ultimate AQ-C18 150x30mm, particle size: 5 μ m; Mobile phase: 30-60% acetonitrile in H₂O (add 0.1% TFA-ACN, v/v)]. The resulting solid was dissolved in 0.2 M hydrochloric acid and again lyophilized to give:

[00561] Compound (*R*)-**23** (55 mg, 65% yield) as a white solid: cSFC analytical (D) tR=2.74 min., purity: 100.00%; LCMS (FF): tR=2.30 min., 292.1 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.65 (d, J=8.4 Hz, 1H), 7.32 (d, J=8.4 Hz, 1H), 4.19-4.16 (m, 1H), 3.90-3.83 (m, 1H), 3.50-3.36 (m, 4H), 3.30-3.29 (m, 1H), 2.56-2.54 (m, 1H), 2.50 (m, 3H), 2.36-2.32 (m, 1H), 2.16-2.09 (m, 2H), 2.00-1.93 (m, 1H).

[00562] **Example 24:** (*R*)-6-chloro-7-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-24)




[00563] Following general procedure C1, compound (*R*)-24 was prepared from compound A-45:

[00564] Compound (*R*)-A-45-1 (0.15 g, white solid, 23% yield over three steps) was prepared from compound A-45 (0.45 g, 1.89 mmol) and (*R*)-quinuclidin-3-amine (0.24 g, 2.0 mmol) with a reaction time of 0.5 h. The product was purified by prep-HPLC [Instrument: GX-H; Column: Waters Xbridge 150×25 mm, particle size: 5 μm; Mobile phase: 20-50% acetonitrile in H₂O (add 0.5% NH₃·H₂O, v/v)]. LCMS (J): (ES⁺) *m/z* (M+H)⁺ = 328.1, *t*R=1.04 min.

[00565] A mixture of compound (*R*)-A-45-1 (0.14 g, 0.43 mmol) and potassium carbonate (0.19 g, 1.4 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 120 °C for 3 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: Phenomenex Gemini C18 250×50 mm, particle size: 10 μm; Mobile phase: 37-67% acetonitrile in H₂O (add 0.05% ammonia-ACN, v/v)]. The residue solid was dissolved in 0.2 N hydrochloric acid and lyophilized again to give:

[00566] Compound (*R*)-24 (45 mg, 33% yield) as a white solid: cSFC analytical *t*R=2.51 min., purity: 97%; LCMS (FF): *t*R=2.25 min., (ES⁺) *m/z* (M+H)⁺ = 308.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.48 (d, *J*=8.8 Hz, 1H), 7.30 (d, *J*=8.0 Hz, 1H), 4.20-4.16 (m, 4H), 3.90-3.83 (m, 1H), 3.50-3.31 (m, 5H), 2.57-2.55 (m, 1H), 2.36-2.35 (m, 1H), 2.16-2.10 (m, 2H), 1.98-1.94 (m, 1H).

[00567] **Example 25:** (*R*)-6-chloro-*N*-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-25)



[00568] Following general procedure C1, compound (*R*)-25 was prepared from compound A-48:


[00569] Compound (*R*)-A-48-1 (0.20 g, white solid, 39% yield over two steps) was prepared from compound A-48 (0.39 g, 1.4 mmol) and (*R*)-quinuclidin-3-amine (0.21 g, 1.7 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250×50 mm, particle size: 10 μm; Mobile phase: 30-60% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (J): (ES⁺) *m/z* (M+H)⁺ = 366.2, *t*R= 1.239 min.

[00570] A mixture of compound (*R*)-A-48-1 (0.15 g, 0.41 mmol) and potassium carbonate (0.17 g, 1.2 mmol) in *N,N*-dimethylformamide (4.0 mL) was stirred at 120 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-A;

Column: Phenomenex Synergi C18 150x30 mm, particle size: 4 μ m; Mobile phase: 25-55% acetonitrile in H₂O (add 0.5% HCl, v/v)]. The product was lyophilized to give:

[00571] Compound (*R*)-**25** (60 mg, 38% yield) as a white solid: cSFC analytical (D) t_R=1.977 min., purity: 100%; LCMS (EE): t_R=2.154 min., 346.1 m/z (M+1); ¹H-NMR (D₂O, 400 MHz): δ 7.66-7.63 (d, J=8.4 Hz, 1H), 7.25-7.23 (d, J=8.0 Hz, 1H), 4.00-3.98 (m, 1H), 3.75-3.69 (m, 1H), 3.36-3.15 (m, 5H), 2.40-2.39 (m, 1H), 2.18-1.85 (m, 4H).

[00572] **Example 26:** (*R*)-6-chloro-7-cyclopropyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine ((*R*)-**26**)




[00573] Following general procedure C1, compound (*R*)-**26** was prepared from compound **A-54**:

[00574] Compound (*R*)-**A-54-1** (0.25 g, white solid, 36% yield over two steps) was prepared from compound **A-54** (0.50 g, 2.0 mmol) and (*R*)-quinuclidin-3-amine (0.25 g, 2.0 mmol). The product was purified by prep-HPLC [Instrument: GX-H; Column: Phenomenex Gemini C18 250x50 mm, particle size: 10 μ m; Mobile phase: 27-57% acetonitrile in H₂O (add 0.5% NH₃OH · H₂O, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 338.1, t_R=1.159 min.

[00575] A mixture of compound (*R*)-**A-54-1** (0.15 g, 0.44 mmol) and potassium phosphate (0.28 g, 1.3 mmol) in *N,N*-dimethylformamide (20 mL) was stirred at 150 °C for 0.5 hour. On completion, the reaction mixture was filtered and concentrated in vacuo. The residue was purified by prep-HPLC [Instrument: GX-I; Column: Welch Ultimate AQ-C18 150x30 mm; particle size: 5 μ m; Mobile phase: 25-55% acetonitrile in H₂O (add 0.1% TFA, v/v)]. The resulting solid was dissolved in 0.2 N hydrochloric acid and again lyophilized to give:

[00576] Compound (*R*)-**26** (80 mg, 50% yield) as a white solid: cSFC analytical (D) t_R=3.728 min., purity: 99.57%; LCMS (GG): t_R=2.152 min., (ES⁺) m/z (M+H)⁺ = 318.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.18-7.13 (m, 2H), 4.22-4.20 (m, 1H), 3.89-3.83 (m, 1H), 3.50-3.36 (m, 5H), 2.63-2.60 (m, 1H), 2.35-2.31 (m, 1H), 2.21-2.10 (m, 3H), 2.00-1.98 (m, 1H), 1.12-1.07 (m, 2H), 0.95-0.93 (m, 2H).

[00577] **Example 27:** (*R*)-4-fluoro-6-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**27**)




[00578] Following general procedure C1, compound (*R*)-**27** was prepared from compound **A-56**:

[00579] Compound (*R*)-**A-56-1** (1.1 g, white solid, 90% yield over two steps) was prepared from compound **A-56** (0.80 g, 3.5 mmol) and (*R*)-quinuclidin-3-amine (0.45 g, 3.5 mmol). The product was purified by silica gel chromatography (dichloromethane/methanol=100:1~10:1). LCMS (J): tR=1.144 min., 316.0 m/z (M+1).

[00580] A mixture of compound (*R*)-**A-56-1** (0.30 g, 0.86 mmol) and potassium carbonate (0.35 g, 2.6 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 150 °C for 1 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: Phenomenex Synergi C18 150x30 mm, particle size: 4 µm; Mobile phase: 17-47% acetonitrile in H₂O (add 0.1%TFA-ACN, v/v)]. The resulting solid was dissolved in 0.2 M hydrochloric acid and again lyophilized again to give:

[00581] Compound (*R*)-**27** (70 mg, 14% yield) as a yellow solid: cSFC analytical (D) tR=1.1 min., purity: 98%; LCMS (EE): tR=2.101 min., 296.0 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.43 (s, 1H), 7.14-7.12 (d, J=9.2 Hz, 1H), 4.19-4.17 (m, 1H), 3.86-3.80 (m, 1H), 3.46-3.35 (m, 5H), 2.59-2.58 (m, 1H), 2.31 (m, 1H), 2.14-2.08 (m, 2H), 1.95 (m, 1H).

[00582] **Example 28:** (*R*)-4,6-dichloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**28**)




[00583] Following general procedure C1, compound (*R*)-**28** was prepared from compound **A-59**:

[00584] Compound (*R*)-**A-59-1** (0.20 g, white solid, 27% yield over two steps) was prepared from compound **A-59** (0.35 g, 1.4 mmol) and (*R*)-quinuclidin-3-amine (0.22 g, 1.7 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250x50 mm, particle size: 10 µm; Mobile phase: 22-52% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 332.1, tR= 1.125 min.

[00585] A mixture of compound (*R*)-**A-59-1** (0.10 g, 0.30 mmol) and potassium carbonate (0.12 g, 0.90 mmol) in *N,N*-dimethylformamide (4.0 mL) was stirred at 150 °C for 0.5 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Synergi C18 150x30 mm, particle size: 10 µm; Mobile phase: 16-46% acetonitrile in H₂O (add 0.5% HCl, v/v)]. The product was lyophilized to give:

[00586] Compound (*R*)-**28** (0.15 g, 37% yield) as a white solid: cSFC analytical (D) tR=3.25 min., purity: 99%; LCMS (EE): tR=2.776 min., 312.0 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.57-7.60 (d, J=1.6 Hz, 1H), 7.40-7.39 (d, J=1.2 Hz, 1H), 4.21-4.19 (m, 1H), 3.87-3.81 (m, 1H), 3.50-3.36 (m, 5H), 2.62-2.60 (m, 1H), 2.35-2.29 (m, 1H), 2.16-2.09 (m, 2H), 1.99-1.94 (m, 1H).

[00587] **Example 29:** (*R*)-7-fluoro-6-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**29**)




[00588] Following general procedure C1, compound (*R*)-29 was prepared from compound A-61:

[00589] Compound (*R*)-A-61-1 (0.12 g, white solid, 17% yield over two steps) was prepared from compound A-61 (0.50 g, 2.1 mmol) and (*R*)-quinuclidin-3-amine (0.26 g, 2.3 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Synergi C18 250x50 mm, particle size: 10 μ m; Mobile phase: 25-55% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 312.1, tR = 1.161 min.

[00590] A mixture of compound (*R*)-A-61-1 (0.10 g, 0.32 mmol) and potassium carbonate (0.13 g, 0.56 mmol) in *N,N*-dimethylformamide (4.0 mL) was stirred at 120 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: Phenomenex Synergi C18 100x21.2mm, particle size: 4 μ m; Mobile phase: 15-45% acetonitrile in H₂O (add 0.1% TFA, v/v)]. The resulting solid was dissolved in 0.2 M hydrochloric acid and again lyophilized to give:

[00591] Compound (*R*)-29 (15 mg, 16 % yield) as a yellow solid: cSFC analytical tR = 2.83 min., purity: 100.00%; LCMS (FF): tR = 2.08 min., 292.1 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.59-7.57 (m, 1H), 7.16-7.12 (m, 1H), 4.16-4.14 (m, 1H), 4.00-3.98 (m, 3H), 3.89-3.85 (m, 1H), 3.43-3.29 (m, 5H), 2.55-2.53 (m, 1H), 2.35-2.34 (m, 1H), 2.15-2.10 (m, 2H), 1.99-1.97 (m, 1H).

[00592] **Example 30:** (*R*)-7-chloro-6-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-30) and (*S*)-7-chloro-6-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*S*)-30)



[00593] Following general procedure C1, compound (*R*)-30 was prepared from compound A-64:

[00594] Compound (*R*)-A-64-1 (0.35 g, white solid, 51% yield over three steps) was prepared from compound A-64 (0.45 g, 1.9 mmol) and (*R*)-quinuclidin-3-amine (0.24 g, 2.0 mmol). The product was purified by prep-HPLC [Instrument: GX-H; Column: Waters Xbridge 150x25 mm, particle size: 5 μ m; Mobile phase: 20-5% acetonitrile in H₂O (add 0.05% ammonia-ACN, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 328.1, tR = 1.03 min.

[00595] A solution of compound (*R*)-A-64-1 (250 mg, 0.76 mmol) and potassium carbonate (186 mg, 1.35 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 120 °C for 1 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: Phenomenex Gemini C18 250x50 mm, particle size: 10 μ m; Mobile phase: 37-67%

acetonitrile in H₂O (add 0.05% ammonia-ACN, v/v)]. The residue was dissolved in 0.2 M hydrochloric acid and lyophilized to give:


[00596] Compound (*R*)-**30** (50 mg, 21% yield) as a white solid: cSFC analytical (D) t_R=1.045 min., chiral purity: 87%.

[00597] A solution of (*R*)-**30** at 87% chiral purity (50 mg, 0.46 mmol) in 3 mL of methanol was purified by cSFC (Instrument: SFC A; Column: AY-10 μm; Mobile phase: 50% methanol (0.01% NH₃·H₂O) in CO₂) at room temperature. In addition to the major product compound (*R*)-**30**, the minor product compound (*S*)-**30** was also collected. Each set of collected fractions was concentrated at room temperature and lyophilized. The resulting solids were dissolved in 0.2 M hydrochloric acid solution and again lyophilized to give:

[00598] Compound (*R*)-**30** (35 mg, 70%) as a white solid. cSFC analytical (D) t_R=1.085 min., purity: 100.00%; LCMS (FF): t_R=2.15 min., (ES⁺) m/z (M+H)⁺ = 308.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.78 (d, J=8.8 Hz, 1H), 7.13 (d, J=8.8 Hz, 1H), 4.16-4.14 (m, 1H), 4.01 (m, 3H), 3.88-3.83 (m, 1H), 3.51-3.30 (m, 5H), 2.55-2.54 (m, 1H), 2.36-2.35 (m, 1H), 2.19-2.09 (m, 2H), 1.99-1.93 (m, 1H); and

[00599] Compound (*S*)-**30** (4 mg, 10% yield) as a white solid: cSFC analytical (D) t_R=2.65 min., purity: 96.44%; LCMS (FF): t_R=2.15 min., (ES⁺) m/z (M+H)⁺ = 308.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.79 (d, J=8.8 Hz, 1H), 7.12 (d, J=8.8 Hz, 1H), 4.19-4.14 (m, 1H), 4.01 (s, 3H), 3.88-3.83 (m, 1H), 3.60-3.37 (m, 5H), 2.55-2.54 (m, 1H), 2.36-2.35 (m, 1H), 2.15-2.10 (m, 2H), 1.99-1.93 (m, 1H).

[00600] **Example 31:** (*R*)-7-chloro-6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**31**)




[00601] Following general procedure C1, compound (*R*)-**31** was prepared from compound **A-67**:

[00602] Compound (*R*)-**A-67-1** (0.10 g, white solid, 15% yield over two steps) was prepared from compound **A-67** (0.50 g, 2.3 mmol) and (*R*)-quinuclidin-3-amine (0.28 g, 2.3 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Synergi C18 250x50 mm, particle size: 10 μm; Mobile phase: 25-55% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 312.1, t_R = 1.152 min.

[00603] A mixture of compound (*R*)-**A-67-1** (80 mg, 0.26 mmol) and potassium carbonate (0.11 g, 0.77 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 120 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: Phenomenex Synergi C18 100x21.2mm, particle size: 4 μm; Mobile phase: 20-50% acetonitrile in H₂O (add 0.1%TFA, v/v)]. The resulting solid was dissolved in 0.2 M hydrochloric acid and again lyophilized to give:

[00604] Compound (*R*)-**31** (20 mg, 26 % yield) as a white solid: cSFC analytical t_R =2.90 min., purity: 94.51%; LCMS (FF): t_R =2.31 min., 292.1 m/z ($M+1$); ^1H -NMR (CD_3OD , 400 MHz): δ 7.68-7.66 (m, 1H), 7.26-7.24 (d, J =8.0 Hz, 1H), 4.18-4.16 (m, 1H), 3.89-3.84 (m, 1H), 3.45-3.33 (m, 5H), 2.56-2.54 (m, 1H), 2.53 (s, 3H), 2.35-2.35 (m, 1H), 2.16-2.12 (m, 2H), 1.99-1.96 (m, 1H).

[00605] **Example 32:** (*R*)-6-isopropoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**32**) and (*S*)-6-isopropoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*S*)-**32**)



[00606] Following general procedure C1, compound (*R*)-**32** was prepared from compound **A-71**:

[00607] Compound (*R*)-**A-71-1** (0.40 g, white solid, 68% yield over two steps) was prepared from compound **A-71** (0.40 g, 1.7 mmol) and (*R*)-quinuclidin-3-amine (0.26 g, 2.1 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250x50 mm, particle size: 10 μ m; Mobile phase: 22-52% acetonitrile in H_2O (add 0.5% ammonia, v/v)]. LCMS (J): (ES^+) m/z ($M+H$) $^+$ = 322.2, t_R = 1.166 min.

[00608] A mixture of compound (*R*)-**A-71-1** (0.16 g, 0.50 mmol) and potassium phosphate (0.32 g, 1.5 mmol) in *N,N*-dimethylformamide (4.0 mL) was stirred at 150 °C for 1 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Synergi C18 150x30 mm, particle size: 10 μ m; Mobile phase: 16-46% acetonitrile in H_2O (add 0.5% HCl, v/v)]. The product was lyophilized to give:

[00609] Compound (*R*)-**32** (80 mg, 53% yield) as a white solid: cSFC analytical (D) t_R =2.760 min., chiral purity: 90%;


[00610] Chiral Purification:

[00611] A solution of compound (*R*)-**32** at 90% chiral purity (80 mg, 0.27 mmol) in 4 mL of methanol was purified by cSFC (Instrument: SFC A; Column: AY-10 μ m; Mobile phase: 50% methanol (0.01% $NH_3 \cdot H_2O$) in CO_2) at room temperature. In addition to the major product compound (*R*)-**32**, the minor product compound (*S*)-**32** was also collected. Each set of collected fractions was concentrated at room temperature and lyophilized. The resulting solids were dissolved in 0.2 M hydrochloric acid and again lyophilized to give:

[00612] Compound (*R*)-**32** (60 mg, 67% yield) as a yellow solid : cSFC analytical (D) t_R =2.684 min., purity: 100%; LCMS (EE): t_R =2.264 min., (ES^+) m/z ($M+H$) $^+$ = 302.2; ^1H -NMR (CD_3OD , 400 MHz): δ 7.71-7.69 (d, J =8.4 Hz, 1H), 6.94-6.94 (d, J =1.6 Hz, 1H), 6.87-6.84 (dd, J =8.4 Hz, J =2.0 Hz, 1H), 4.74-4.68 (m, 1H), 4.14-4.12 (m, 1H), 3.87-3.81 (m, 1H), 3.49-3.30 (m, 5H), 2.54-2.52 (m, 1H), 2.36-2.34 (m, 1H), 2.15-2.08 (m, 2H), 1.97-1.95 (m, 1H), 1.38-1.36 (d, J =7.0 Hz, 6H); and

[00613] Compound (*S*)-**32** (6.0 mg, 7% yield) as a yellow solid : cSFC analytical (D) t_R =4.131 min., purity: 97.12%; LCMS (EE): t_R =2.281 min., (ES^+) m/z ($M+H$) $^+$ = 302.1; ^1H -NMR (CD_3OD , 400 MHz): δ 7.71-7.69 (d, J =8.4 Hz, 1H), 6.94-6.94 (d, J =1.6 Hz, 1H), 6.87-6.84 (dd, J =8.4 Hz, J =2.0 Hz, 1H), 4.74-4.68 (m, 1H), 4.14-4.12 (m, 1H), 3.87-3.82 (m, 1H), 3.47-3.28 (m, 5H), 2.54-2.52 (m, 1H), 2.36-2.32 (m, 1H), 2.15-2.08 (m, 2H), 1.98-1.92 (m, 1H), 1.38-1.36 (d, J =7.0 Hz, 6H).

[00614] **Example 33:** (*R*)-6-ethoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine ((*R*)-**33**) and (*S*)-6-ethoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine ((*S*)-**33**)



[00615] Following general procedure C1, compound (*R*)-**33** was prepared from compound **A-74**:

[00616] Compound (*R*)-**A-74-1** (0.30 g, white solid, 42% yield over two steps) was prepared from compound **A-74** (0.50 g, 2.3 mmol) and (*R*)-quinuclidin-3-amine (0.29 g, 2.3 mmol). The product was purified by base prep-HPLC [Instrument: GX-A; Column: Phenomenex Gemini C18 250×50 mm, particle size: 10 μ m; Mobile phase: 24-54% acetonitrile in H_2O (add 0.5% $NH_3 \cdot H_2O$, v/v)]. LCMS (J): (ES^+) m/z ($M+H$) $^+$ = 308.2, t_R =1.117 min.

[00617] A mixture of compound (*R*)-**A-74-1** (0.15 g, 0.49 mmol) and potassium carbonate (0.20 g, 1.5 mmol) in *N,N*-dimethylformamide (20 mL) was stirred at 150 °C for 2 hours. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: Welch Ultimate AQ-C18 150x30 mm, particle size: 5 μ m; Mobile phase: 17-47% acetonitrile in H_2O (add 0.1% TFA, v/v)]. The resulting solid was dissolved in 0.2 M hydrochloric acid and again lyophilized to give compound (*R*)-**33** (0.12 g, 81% chiral purity).


[00618] A solution of compound (*R*)-**33** at 81% chiral purity (0.12 g, 0.39 mmol) in 5 mL ethanol was purified by SFC (Column: Chiralpak AY-H -150×4.6mm, I.D., 5 μ m; Mobile phase: ethanol (0.05% DEA) in CO_2) at room temperature. In addition to the major product compound (*R*)-**33**, the minor product compound (*S*)-**33** was also collected. Each set of collected fractions was concentrated at room temperature. The residue was dissolved in 0.2 M hydrochloric acid and lyophilized to give:

[00619] Compound (*R*)-**33** (40 mg, 32% yield) as a yellow solid: cSFC analytical (A) t_R =2.73 min., purity: 100%; LCMS (GG): t_R =1.969 min., (ES^+) m/z ($M+H$) $^+$ = 288.1; ^1H -NMR (CD_3OD , 400 MHz): δ 7.70-7.68 (d, J =8.8 Hz, 1H), 6.94-6.93 (d, J =1.6 Hz, 1H), 6.89-6.87 (m, 1H), 4.14-4.09 (m, 3H), 3.87-3.81 (m, 1H), 3.47-3.36 (m, 4H), 3.33-3.30 (m, 1H), 2.54-2.52 (m, 1H), 2.36-2.34 (m, 1H), 2.15-2.08 (m, 2H), 1.97-1.95 (m, 1H), 1.46-1.43 (t, J =7.2 Hz, 3H); and

[00620] Compound (*S*)-**33** (10 mg, 8% yield) as a yellow solid: cSFC analytical (A) t_R =3.86 min., purity: 100%; LCMS (GG): t_R =1.935 min., (ES^+) m/z ($M+H$) $^+$ = 288.1; ^1H -NMR (CD_3OD , 400 MHz): δ 7.71-7.69 (d, J =8.4 Hz, 1H), 6.94-6.93 (d, J =1.6 Hz, 1H), 6.89-6.86 (m, 1H), 4.14-4.09 (m,

3H), 3.87-3.81 (m, 1H), 3.50-3.37 (m, 4H), 3.33-3.28 (m, 1H), 2.54-2.52 (m, 1H), 2.36-2.34 (m, 1H), 2.15-2.08 (m, 2H), 1.97-1.95 (m, 1H), 1.46-1.43(t, J=7.2 Hz, 3H).

[00621] Example 34: (*R*)-6-chloro-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-7-carbonitrile hydrochloride ((*R*)-34)




[00622] Following general procedure C1, compound (*R*)-34 was prepared from compound A-77:

[00623] Compound (*R*)-A-77-1 was prepared as follows: A solution of (*R*)-quinuclidin-3-amine (0.27 g, 2.2 mmol) in methanol (5 mL) was added dropwise to a solution of 4-chloro-3-cyano-2-fluoro-*N*-hydroxybenzimidoyl chloride compound A-77 (0.50 g, 2.2 mmol) in methanol (5 mL) at room temperature. The resulting mixture was stirred at room temperature for 2 hours. On completion, the reaction mixture was concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: Atlantis HILIC Silica C18 150×19 mm, particle size: 5 µm; Mobile phase: 11-41% acetonitrile in H₂O (add 0.1% TFA-ACN, v/v)] to give compound (*R*)-A-77-1 (0.20 g, 28% yield over two steps) as a yellow solid. LCMS (M): t_R=0.540 min., 323.1 m/z (M+1).

[00624] A solution of compound (*R*)-A-77-1 (0.18 g, 0.41 mmol) and potassium carbonate (0.17 g, 1.2 mmol) in *N,N*-dimethylformamide (2.0 mL) was stirred at 150 °C for 2 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-E; Column: Phenomenex Synergi C18 150×30 mm, particle size: 4 µm; Mobile phase: 12-42% acetonitrile in H₂O (add 0.05% HCl-ACN, v/v)]. The resulting solution was lyophilized to give:

[00625] Compound (*R*)-34 (20 mg, 16% yield) as a white solid: cSFC analytical (D) t_R=2.8 min., purity: 97%; LCMS (GG): t_R=1.942 min., 303.1 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 8.17 (d, J=8.8 Hz, 1H), 7.56 (d, J=8.4 Hz, 1H), 4.21-4.18 (m, 1H), 3.91-3.88 (m, 1H), 3.47-3.36 (m, 5H), 2.57-2.55 (m, 1H), 2.34 (m, 1H), 2.16-2.10 (m, 2H), 1.99-1.98 (m, 1H).

[00626] Example 35: (*R*)-7-chloro-6-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-35)



[00627] Following general procedure C1, compound (*R*)-35 was prepared from compound A-79:


[00628] Compound (*R*)-A-79-1 (0.30 g, white solid, 54% yield over two steps) was prepared from compound A-79 (0.40 g, 1.8 mmol) and (*R*)-quinuclidin-3-amine (0.22 g, 1.8 mmol). The product was purified by prep-HPLC [Instrument: GX-H; Column: Phenomenex Gemini C18 250×50 mm, particle

size: 10 μ m; Mobile phase: 24-54% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 316.1, tR= 1.084 min.

[00629] A mixture of compound (*R*)-**A-79-1** (0.10 g, 0.32 mmol) and potassium phosphate (0.20 g, 0.95 mmol) in *N,N*-dimethylformamide (10 mL) was stirred at 150 °C for 0.5 hour. On completion, the reaction mixture was filtered and concentrated in vacuo. The residue was purified by prep-HPLC [Instrument: GX-I; Column: Xtimate C18 150x25 mm; particle size: 5 μ m; Mobile phase: 12-42% acetonitrile in H₂O (add 0.1% TFA, v/v)]. The resulting solid was dissolved in 0.2 N hydrochloric acid and again lyophilized to give:

[00630] Compound (*R*)-**35** (70 mg, 66% yield) as a white solid: cSFC analytical (D) tR=2.375 min., purity: 98.24%; LCMS (FF): tR=2.306 min., (ES⁺) m/z (M+H)⁺ = 296.1; ¹H-NMR (CD₃OD, 400 MHz): δ 7.84-7.81 (m, 1H), 7.28-7.23 (t, J=9.0 Hz, 1H), 4.18-4.16 (m, 1H), 3.90-3.84 (m, 1H), 3.48-3.36 (m, 4H), 3.33-3.30 (m, 1H), 2.56-2.54 (m, 1H), 2.38-2.32 (m, 1H), 2.16-2.09 (m, 2H), 1.99-1.95 (m, 1H).

[00631] **Example 36:** (*R*)-6-(1H-pyrazol-1-yl)-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**36**)




[00632] Following general procedure C1, compound (*R*)-**36** was prepared from compound **A-82**:

[00633] Compound (*R*)-**A-82-1** (0.15 g, white solid, 27% yield over two steps) was prepared from compound **A-82** (0.30 g, 1.3 mmol) and (*R*)-quinuclidin-3-amine (0.16 g, 1.3 mmol). The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250x50 mm, particle size: 10 μ m; Mobile phase: 22-52% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 330.2, tR= 1.044 min.

[00634] A solution of compound (*R*)-**A-82-1** (0.10 g, 0.30 mmol) and potassium phosphate (0.19 g, 0.91 mmol) in *N,N*-dimethylformamide (5.0 mL) was stirred at 150 °C for 1 hour. On completion, the reaction mixture was filtered and concentrated in vacuo. The residue was purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Synergi C18 150x30 mm, particle size: 10 μ m; Mobile phase: 16-46% acetonitrile in H₂O (add 0.5% HCl, v/v)] to give:

[00635] Compound (*R*)-**36** (50 mg, 48% yield) as a yellow solid: cSFC analytical (D) tR=3.154 min., purity: 95.41%; LCMS (EE): tR=2.174 min., 310.1 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 8.40-8.39 (d, J=2.4 Hz, 1H), 7.99-7.97 (d, J=8.4 Hz, 1H), 7.86 (s, 1H), 7.81-7.77 (m, 2H), 6.66 (s, 1H), 4.20-4.18 (m, 1H), 3.91-3.79 (m, 1H), 3.53-3.36 (m, 5H), 2.57-2.56 (m, 1H), 2.41-2.31 (m, 1H), 2.18-2.10 (m, 2H), 2.00-1.94 (m, 1H).

[00636] **Example 37:** (*R*)-6-(5-methyl-1,3,4-oxadiazol-2-yl)-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**37**)




[00637] Following general procedure C1, compound **(R)-37** was prepared from compound **A-89**:

[00638] Compound **(R)-A-89-1** (0.78 g, yellow solid, crude) was prepared from compound **A-89** (0.58 g, 2.3 mmol) and *(R)*-quinuclidin-3-amine (0.28 g, 2.3 mmol). The crude was used for the next step without purification. LCMS (J): (ES⁺) *m/z* (M+H)⁺ = 346.2, *t*_R=0.931 min.

[00639] A solution of compound **(R)-A-89-1** (0.78 g, crude) and potassium phosphate (1.4 g, 6.8 mmol) in *N,N*-dimethylformamide (20 mL) was stirred at 150 °C for 1 hour. On completion, the reaction mixture was filtered and concentrated in vacuo. The residue was purified by prep-HPLC [Instrument: GX-B; Column: Welch Ultimate AQ-C18 150x30 mm, particle size: 5 μm; Mobile phase: 10-40% acetonitrile in H₂O (add 0.1%TFA, v/v)]. The resulting solid was dissolved in 0.2 N hydrochloric acid and again lyophilized to give:

[00640] Compound **(R)-37** (0.13 g, 16% yield over three steps) as a yellow solid: cSFC analytical(D) *t*_R=3.11 min., purity: 100%; LCMS (X): *t*_R=1.78 min., 326.1 *m/z* (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 8.08 (s, 1H), 8.04-8.02 (m, 1H), 7.99-7.97 (m, 1H), 4.23-4.20 (m, 1H), 3.93-3.86 (m, 1H), 3.51-3.33 (m, 5H), 2.57 (s, 3H), 2.68-2.57 (m, 1H), 2.38-2.37 (m, 1H), 2.17-2.13 (m, 2H), 2.11-1.98 (m, 1H).

[00641] **Example 38:** *(R)*-7-methoxy-6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride (**(R)-38**)




[00642] Following general procedure C1, compound **(R)-38** was prepared from compound **A-92**:

[00643] Compound **(R)-A-92-1** (0.13 g, yellow solid, 13% yield over two steps) was prepared from compound **A-92** (0.50 g, 2.3 mmol) and *(R)*-quinuclidin-3-amine (0.4 g, 3.2 mmol) with a reaction time of 4 hours. The product was purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250x50mm, particle size: 10 μm; Mobile phase: 21-51% acetonitrile in H₂O (add 0.05% ammonia, v/v)]. LCMS (J): (ES⁺) *m/z* (M+H)⁺ = 308.2, *t*_R=1.092 min.

[00644] A mixture of compound **(R)-A-92-1** (0.13 g, 0.41 mmol) and potassium phosphate (0.26 g, 1.2 mmol) in *N,N*-dimethylformamide (20 mL) was stirred at 150 °C for 1 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-B; Column: Welch Ultimate AQ-C18 150x30 mm, particle size: 5 μm; Mobile phase: 23-53% acetonitrile in H₂O (add 0.1%TFA, v/v)]. The resulting solution was dissolved in 0.2 N hydrochloric acid and again lyophilized to give:

[00645] Compound (*R*)-**38** (40 mg, 34% yield) as a white solid: cSFC analytical(D) t_R=2.64 min., purity: 97.49%; LCMS (X): t_R=1.90 min., 288.2 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.39-7.38 (d, J=7.2 Hz, 1H), 7.10-7.08 (d, J=7.2 Hz, 1H), 4.18-4.16 (m, 1H), 4.14 (s, 3H), 3.86-3.85 (m, 1H), 3.43-3.29 (m, 5H), 2.56 (m, 1H), 2.55 (m, 1H), 2.55 (s, 3H), 2.54-2.51 (m, 2H), 2.35 (m, 1H).

[00646] **Example 39:** (*R*)-6-chloro-7-ethoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**39**)




[00647] Following general procedure C1, compound (*R*)-**39** was prepared from compound **A-96**:

[00648] Compound (*R*)-**A-96-1** (0.30 g, white solid, 44% yield over two steps) was prepared from compound **A-96** (0.50 g, 2.0 mmol) and (*R*)-quinuclidin-3-amine (0.30 g, 2.4 mmol). The product was purified by prep-HPLC [Instrument: GX-H; Column: Phenomenex Gemini C18 250x50 mm, particle size: 10 μm; Mobile phase: 30-60% acetonitrile in H₂O (add 0.5% ammonia, v/v)]. LCMS (J): (ES⁺) m/z (M+H)⁺ = 342.1, t_R = 1.223 min.

[00649] A solution of compound (*R*)-**A-96-1** (0.20 g, 0.59 mmol) and potassium phosphate (0.37 g, 1.8 mmol) in *N,N*-dimethylacetamide (4.0 mL) was stirred at 150 °C for 1 hour. On completion, the reaction mixture was filtered and concentrated in vacuo. The residue was purified by prep-HPLC [Instrument: GX-A; Column: Phenomenex Synergi C18 150x30 mm, particle size: 4 μm; Mobile phase: 25-55% acetonitrile in H₂O (add 0.5% HCl, v/v)] to give:

[00650] Compound (*R*)-**39** (60 mg, 29% yield) as a yellow solid: cSFC analytical (D) t_R=2.600 min., purity: 96.70%; LCMS (EE): t_R = 2.12 min., 322.1 m/z (M+1); ¹H-NMR (CD₃OD, 400 MHz): δ 7.50-7.48 (d, J=8.4 Hz, 1H), 7.31-7.29 (d, J=8.0 Hz, 1H), 4.51-4.46 (dd, J=14 Hz, J=7.2 Hz, 1H), 4.18-4.16 (m, 1H), 3.89-3.83 (m, 1H), 3.47-3.31 (m, 5H), 2.57-2.54 (m, 1H), 2.36-2.32 (m, 1H), 2.16-2.10 (m, 2H), 1.98-1.96 (m, 1H), 1.45-1.42 (t, J=7.2 Hz, 3H).

[00651] **Example 40:** (*R*)-6,7-dimethyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*R*)-**40**) and (*S*)-6,7-dimethyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine hydrochloride ((*S*)-**40**)



[00652] Following general procedure C1, compound (*R*)-**40** was prepared from compound **A-99**:

[00653] Compound (*R*)-**A-99-1** (0.60 g, yellow solid, 48% yield over two steps) was prepared from compound **A-99** (1.9 g, 4.0 mmol) and (*R*)-quinuclidin-3-amine (0.50 g, 4.0 mmol) with a

reaction of 2 hours. The product was purified by prep-HPLC [Instrument: GX-B; Column: Phenomenex Gemini C18 250×50 mm, particle size: 10 μm; Mobile phase: 25-48% acetonitrile in H₂O (add 0.05% NH₃·H₂O, v/v)]. LCMS (M): tR=0.529 min., 292.2 m/z (M+1).

[00654] A solution of compound (*R*)-**A-99-1** (0.58 g, 1.8 mmol) and potassium carbonate (0.76 g, 5.5 mmol) in *N,N*-dimethylformamide (3.0 mL) was stirred at 150 °C for 6 hour. On completion, the reaction mixture was filtered, concentrated in vacuo and purified by prep-HPLC [Instrument: GX-H; Column: Phenomenex Gemini C18 150×25 mm, particle size: 10 μm; Mobile phase: 36-66% acetonitrile in H₂O (add 0.05% NH₃·H₂O-ACN, v/v)]. The resulting solution was lyophilized to give:

[00655] Compound (*R*)-**40** (210 mg, 42% yield) as a white solid: cSFC analytical (D) tR=2.7 min., chiral purity: 78%;

[00656] A solution of compound (*R*)-**40** at 78% chiral purity (210 mg, 0.77 mmol) in 4 mL of ethanol was purified by SFC (Instrument: SFC A; Column: AD-10 μm; Mobile phase: 50% ethanol (0.01% NH₃·H₂O) in CO₂) at room temperature. In addition to the major product compound (*R*)-**40**, the minor product compound (*S*)-**40** was also collected. Each set of collected fractions was concentrated at room temperature and lyophilized.


[00657] The resulting solids were re-purified by prep-HPLC [Instrument: GX-C; Column: Phenomenex Gemini C18 250×50 mm, particle size: 10 μm; Mobile phase: 46-76% acetonitrile in H₂O (add 0.05% NH₃·H₂O, v/v)]. The resulting solids was dissolved in 0.2 M hydrochloric acid solution and again lyophilized to give:

[00658] Compound (*R*)-**40** (100 mg, 48% yield) as a yellow solid : cSFC analytical (D) tR=2.789 min., purity: 99.85%; LCMS (GG): tR=1.976 min., (ES⁺) m/z (M+H)⁺ = 272.2; ¹H-NMR (CD₃OD, 400 MHz): δ7.56-7.54 (d, J=8.4 Hz, 1H), 7.12-7.10 (d, J=8.0 Hz, 1H), 4.17-4.16 (m, 1H), 3.88-3.85 (m, 1H), 3.42-3.36 (m, 5H), 2.56-2.53 (m, 1H), 2.42-2.39 (m, 7H), 2.14-1.96 (m, 3H); and

[00659] Compound (*S*)-**40** (20 mg, 9.5% yield) as a yellow solid : cSFC analytical (D) tR=3.726 min., purity: 98.48%; LCMS (GG): tR=1.953 min., (ES⁺) m/z (M+H)⁺ = 272.2; ¹H-NMR (CD₃OD, 400 MHz): δ7.56-7.54 (d, J=8.0 Hz, 1H), 7.11-7.09 (d, J=8.0 Hz, 1H), 4.16-4.15 (m, 1H), 3.86-3.81 (m, 1H), 3.41-3.31 (m, 5H), 2.53-2.52 (m, 1H), 2.40-2.37 (m, 7H), 2.11-1.94 (m, 3H).

[00660] **Example 41:**

[00661] Preparation: (*R*)-6-chloro-7-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine ((*R*)-**41**)



[00662] Following general procedure C2, compound (*R*)-**41** was prepared from compound **A-101**:

[00663] Compound (*R*)-**A-101-1** (88 mg, 70% yield) was prepared as a white solid from **A-101** (165 mg, 0.7 mmol) and (*R*)-quinuclidin-3-amine (63 mg, 0.5 mmol) using 5 mL of methanol and a

reaction time of 1 hour. The product purified by silica gel column chromatography [chloroform: methanol = 1:0 to 17:3]. LCMS (1): $t_R=3.163$ min., (ES^+) m/z $(M+H)^+ = 316.0$.


[00664] To a solution of compound **(R)-A-101-1** (103 mg, 0.3 mmol) in *N,N*-dimethylacetamide (5 mL) was added potassium tertbutoxide (37 mg, 0.3 mmol). The mixture was stirred at room temperature for 16 hours. Additional potassium tertbutoxide (9 mg, 0.1 mmol) was added, and the mixture was stirred for an additional 2 hours. The solution was put on an SCX column and eluted with methanol. The product was eluted from the column using 7 M ammonia in methanol, concentrated and purified by silica gel column chromatography [chloroform: 7M NH_3 in methanol = 99:1 to 9:1]. The resulting product was triturated with water, collected by filtration and dried in a vacuum oven to afford:

[00665] Compound **(R)-41** (32 mg, 32% yield) as a white solid: cHPLC analytical [Chiralcel OD-H, 250×4.6 mm, particle size: 5 μ m; Flow: 1.0 mL/min; Column temp: 25 °C; Mobile phase: 0.1% diethylamine in Heptane/Ethanol = 9/1; detection: DAD (220-320 nm)] $t_R=6.105$ min., purity: 100%; LCMS (1): $t_R=3.269$ min., (ES^+) m/z $(M+H)^+ = 296.0$; ^1H NMR (300 MHz, $CDCl_3$) δ 7.28 – 7.17 (m, 2H), 4.27 (d, $J = 5.3$ Hz, 1H), 3.90 – 3.79 (m, 1H), 3.54 – 3.42 (m, 1H), 2.96 – 2.74 (m, 4H), 2.67 – 2.56 (m, 1H), 2.27 – 2.19 (m, 1H), 1.86 – 1.62 (m, 3H), 1.56 – 1.42 (m, 1H).

[00666] The hydrochloride salt form of compound **(R)-41** was also prepared by dissolving the freebase form of compound **(R)-41** in 0.2 N hydrochloric acid and subjected to lyophilization to give:

[00667] Compound **(R)-41**-hydrochloride as a white solid: cSFC analytical (D) $t_R=2.31$ min., purity: 98.36%; LCMS (FF): $t_R=1.98$ min., 296.1 m/z $(M+1)$; ^1H -NMR (CD_3OD , 400 MHz): δ 7.67-7.72 (d, $J=8.4$ Hz 1H), 7.38-7.36 (dd, $J=5.6$ Hz, $J=8.4$ Hz, 1H), 4.18-4.16 (m, 1H), 3.88-3.82 (m, 1H), 3.41-3.34 (m, 5H), 2.55-2.53 (m, 1H), 2.37-2.30 (m, 1H), 2.18-2.04 (m, 2H), 2.00-1.93 (m, 1H).

[00668] Preparation: **(S)**-6-chloro-7-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine (**(S)-41**)



[00669] Following general procedure C2, compound **(S)-41** was prepared from compound **A-101**:

[00670] Compound **(S)-A-101-1** (171 mg, 74% yield) was prepared as a white solid from **A-101** (165 mg, 0.7 mmol) and **(S)**-quinuclidin-3-amine (92 mg, 0.7 mmol) using 6 mL of methanol and a reaction time of 1 hour. The product purified by silica gel column chromatography [chloroform: 7M NH_3 in methanol = 99:1 to 9:1]. LCMS (1): $t_R=2.847$ min., (ES^+) m/z $(M+H)^+ = 316.0$.


[00671] To a solution of compound **(S)-A-101-1** (171 mg, 0.5 mmol) in *N,N*-dimethylacetamide (5 mL) was added potassium tertbutoxide (73 mg, 0.7 mmol). The mixture was stirred at room temperature for 2 hours. Additional potassium tertbutoxide (15 mg, 0.1 mmol) was added, and the mixture was stirred for an additional 16 hours. The solution was put on an SCX column and eluted with methanol. The product was eluted from the column using 7 M ammonia in methanol,

concentrated and purified by silica gel column chromatography [chloroform: 7M NH₃ in methanol = 99/1 to 9/1]. The resulting product was further purified by preparative HPLC [Instrument: AT; Column: Phenomenex Gemini-NX C18 100x21.2 mm, particle size: 10 µm; Mobile phase A: 99% acetonitrile + 1% 10 mM ammonium bicarbonate in water pH=9.0, Mobile phase B: 10mM ammonium bicarbonate in water pH=9.0] and lyophilized to afford:

[00672] Compound (*S*)-**41** (45 mg, 28% yield) as a white solid: cHPLC analytical [Chiralcel OD-H, 250x4.6 mm, particle size: 5 µm; Flow: 1.0 mL/min; Column temp: 25 °C; Mobile phase: 0.1% diethylamine in Heptane/Ethanol = 9/1; detection: DAD (220-320 nm)] tR=17.874 min., purity: 98%; LCMS (1): tR=3.305 min., (ES⁺) m/z (M+H)⁺ = 296.0; ¹H NMR (300 MHz, CDCl₃) δ 7.28 – 7.18 (m, 2H), 4.30 (d, *J* = 5.2 Hz, 1H), 3.89 – 3.83 (m, 1H), 3.55 – 3.43 (m, 1H), 2.98 – 2.80 (m, 4H), 2.69 – 2.59 (m, 1H), 2.28 – 2.21 (m, 1H), 1.90 – 1.63 (m, 3H), 1.53 – 1.46 (m, 1H).

[00673] Example 42:

[00674] Preparation: (*R*)-6-chloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine ((*R*)-**42**)




[00675] Following general procedure C2, compound (*R*)-**42** was prepared from compound **A-103**:

[00676] Compound (*R*)-**A-103-1** ((88 mg, 70% yield) was prepared as a white solid from **A-103** (90 mg, 0.4 mmol) and (*R*)-quinuclidin-3-amine (50 mg, 0.4 mmol) using 4 mL of methanol and a reaction time of 1 hour. The product was purified by silica gel column chromatography [chloroform: methanol = 1:0 to 17:3]. LCMS (1): tR=2.788 min., (ES⁺) m/z (M+H)⁺ = 316.0.

[00677] To a solution of compound (*R*)-**A-103-1** (77 mg, 0.2 mmol) in N,N-dimethylacetamide (3 mL) was added potassium tertbutoxide (33 mg, 0.3 mmol). The mixture was stirred at room temperature for 2 hours. Additional potassium tertbutoxide (7 mg, 0.1 mmol) was added, and the mixture was stirred for an additional hour. The solution was put on an SCX column and eluted with methanol. The product was eluted from the column using 7 M ammonia in methanol, concentrated and purified by silica gel column chromatography [chloroform: 7M NH₃ in methanol = 99:1 to 9:1]. The resulting product was lyophilized to afford:

[00678] Compound (*R*)-**42** (30 mg, 41% yield) as a white solid: cHPLC analytical [Column: Chiralcel OD-H, 250x4.6 mm, particle size: 5 µm; Flow: 1.0 mL/min; Column temp: 25 °C; Mobile phase: 0.1% diethylamine in Heptane/Ethanol = 9/1; detection: DAD (220-320 nm)] tR=5.949 min., purity: 99%; LCMS (1): tR=3.275 min., (ES⁺) m/z (M+H)⁺ = 296.1; ¹H NMR (300 MHz, CDCl₃) δ 7.49 (d, *J* = 5.5 Hz, 1H), 7.27 (d, *J* = 7.8 Hz, 1H), 4.19 (d, *J* = 5.5 Hz, 1H), 3.89 – 3.78 (m, 1H), 3.53 – 3.42 (m, 1H), 2.96 – 2.79 (m, 4H), 2.67 – 2.56 (m, 1H), 2.26 – 2.18 (m, 1H), 1.85 – 1.63 (m, 3H), 1.56 – 1.42 (m, 1H).

[00679] Preparation: (*S*)-6-chloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine ((*S*)-42)




[00680] Following general procedure C2, compound (*S*)-42 was prepared from compound A-103:

[00681] Compound (*S*)-A-103-1 (167 mg, 35% yield) was prepared as a white solid from A-103 (185 mg, 0.8 mmol) and (*S*)-quinuclidin-3-amine (103 mg, 0.8 mmol) using 6 mL of methanol and a reaction time of 1 hour. The product was purified by silica gel column chromatography [chloroform: 7M NH₃ in methanol = 99:1 to 9:1] to afford (*S*)-A-103-1 (167 mg, 65% yield) as a white solid. LCMS (1): tR=2.806 min., (ES⁺) m/z (M+H)⁺ = 316.0.

[00682] To a solution of compound (*S*)-A-103-1 (167 mg, 0.5 mmol) in *N,N*-dimethylacetamide (5 mL) was added potassium tertbutoxide (71 mg, 0.6 mmol). The mixture was stirred at room temperature for 2 hours. Additional potassium tertbutoxide (15 mg, 0.1 mmol) was added, and the mixture was stirred for an additional 16 hours. The solution was put on an SCX column and eluted with methanol. The product was eluted from the column using 7 M ammonia in methanol, concentrated and purified by silica gel column chromatography [chloroform: 7M NH₃ in methanol = 99/1 to 9/1]. The resulting product was lyophilized to afford:

[00683] Compound (*S*)-42 (65 mg, 42% yield) as a white solid: cHPLC analytical [Column: Chiralcel OD-H, 250×4.6 mm, particle size: 5 μm; Flow: 1.0 mL/min; Column temp: 25 °C; Mobile phase: 0.1% diethylamine in Heptane/Ethanol = 9/1; detection: DAD (220-320 nm)] tR=9.752 min., purity: 98%; LCMS (1): tR=3.244 min., (ES⁺) m/z (M+H)⁺ = 296.0; ¹H NMR (300 MHz, CDCl₃) δ 7.49 (d, *J* = 5.5 Hz, 1H), 7.27 (d, *J* = 7.9 Hz, 1H), 4.21 (d, *J* = 5.1 Hz, 1H), 3.93 – 3.75 (m, 1H), 3.56 – 3.41 (m, 1H), 3.03 – 2.72 (m, 4H), 2.71 – 2.53 (m, 1H), 2.28 – 2.16 (m, 1H), 1.92 – 1.59 (m, 3H), 1.58 – 1.40 (m, 1H).

[00684] Example 43: (*R*)-6-bromo-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine ((*R*)-43)



[00685] Following general procedure C2, compound (*R*)-43 was prepared from compound A-105:

[00686] Compound (*R*)-A-105-1: To a solution of (*R*)-quinuclidin-3-amine (634 mg, 5.0 mmol) and triethylamine (533 mg, 5.3 mmol) in methanol (12 mL) at room temperature was added a solution of compound A-105 (1.3 g, 5.3 mmol) in methanol (25 mL) over 4 hours using a syringe pump. The mixture was stirred for an additional 12 hours, then filtered and concentrated in vacuo. The residue was purified by silica gel column chromatography [chloroform: 7M NH₃ in methanol = 99:1 to 9:1] to

afford (**R**)-**A-105-1** (1.2 g, 70% yield) as a white solid. LCMS (1): tR=2.753 min., (ES⁺) m/z (M+H)⁺ = 342.0/344.0.

[00687] To a solution of compound (**R**)-**A-105-1** (320 mg, 0.9 mmol) in dimethylsulfoxide (5 mL) was added potassium tertbutoxide (131 mg, 1.2 mmol). The mixture was stirred at room temperature for 1.5 hours. Additional potassium tertbutoxide (60 mg, 0.5 mmol) was added, and the mixture was stirred for an additional hour. Additional potassium tertbutoxide (40 mg, 0.4 mmol) was added, and the mixture was stirred for 30 min. The solution was put on an SCX column and eluted with methanol. The product was eluted from the column using 3.5 M ammonia in methanol, concentrated and purified by silica gel column chromatography [chloroform: 7M NH₃ in methanol = 1/0 to 9/1]. The resulting product was further purified by preparative HPLC [Instrument: AT; Column: Phenomenex Gemini-NX C18 100x21.2 mm, particle size: 10 µm; Mobile phase A: 99% acetonitrile + 1% 10 mM ammonium bicarbonate in water pH=9.0, Mobile phase B: 10mM ammonium bicarbonate in water pH=9.0] and lyophilized to afford:

[00688] Compound (**R**)-**43** (91 mg, 21% yield) as a white solid: LCMS (1): tR=3.222 min., (ES⁺) m/z (M+H)⁺ = 322.0/324.0; ¹H NMR (300 MHz, CDCl₃) δ 7.61 (dd, *J* = 1.3, 0.8 Hz, 1H), 7.41 – 7.32 (m, 2H), 4.29 (d, *J* = 5.0 Hz, 1H), 3.88 – 3.84 (m, 1H), 3.54 – 3.42 (m, 1H), 2.98 – 2.80 (m, 4H), 2.70 – 2.59 (m, 1H), 2.29 – 2.21 (m, 1H), 1.88 – 1.64 (m, 3H), 1.56 – 1.45 (m, 1H).

[00689] **Example 44:**

[00690] **Human α7 nAChR Binding Assay**

[00691] The ability of compounds to displace binding of radioactive ligands from human α7 nAChR was determined, as a measure of the affinity of the compounds for these ligand-gated ion channels. The [¹²⁵I]-αBungarotoxin competition binding assay was performed under contract by Cerep Poitiers, France following published the methods (Sharples *et al.*, J Neurosci. 2000; 20(8):2783-91). “SH-SY5Y cells stably expressing human α7 nicotinic acetylcholine receptors, grown to confluency in 175 cm² flasks, were washed briefly with warm PBS containing (in mM): (150 NaCl, 8 K₂HPO₄, 2 KH₂PO₄, pH 7.4, 37°C) and scraped into cold phosphate buffer. Cells were washed by centrifugation for 3 min at 500 × g and resuspended in 10 mL of ice-cold phosphate buffer. The suspension was homogenized for 10 sec using an Ultraturax and centrifuged for 30 min at 45,000 ×g. The pellet was resuspended in phosphate buffer (0.5 mL per original flask). SH-SY5Y membranes (30 µg protein) were incubated in a total volume of 2 mL in 50 mM phosphate buffer with 0.05 nM [¹²⁵I]-αBgt and serial dilutions of test compound. Nonspecific binding was determined in the presence of α-bungarotoxin (1 µM). Samples were incubated for 120 min at 37°C. The reaction was terminated by filtration through Whatman GFA/E filter paper (presoaked overnight in 0.3% polyethyleneimine in PBS), using a Brandel Cell Harvester. Each condition was measured in duplicate. Filters were counted for radioactivity using a scintillation counter. The results were

expressed as a percent inhibition of control specific binding obtained in the presence of the test compounds where Inhibition (%) = $100 - [(\text{measured specific binding}/\text{control specific binding}) \times 100]$.

[00692] The IC₅₀ values (concentration causing a half-maximal inhibition of control specific binding) and Hill coefficients (nH) were determined by non-linear regression analysis of the competition curves generated with mean replicate values using Hill equation:

$$Y = D + \left[\frac{A - D}{1 + (C/C_{50})^{nH}} \right]$$

where Y = specific binding, A = left asymptote of the curve, D = right asymptote of the curve, C = compound concentration, C₅₀ = IC₅₀, and nH = slope factor.

[00693] This analysis was performed using software developed at Cerep (Hill software) and validated by comparison with data generated by the commercial software SigmaPlot® 4.0 for Windows® (© 1997 by SPSS Inc.). The inhibition constants (K_i) were calculated using the Cheng Prusoff equation:

$$K_i = \frac{IC_{50}}{(1 + L/K_D)}$$

where L = concentration of radioligand in the assay, and K_D = affinity of the radioligand for the receptor.

[00694] A scatchard plot is used to determine the K_D. Results are provided in Table 2 (reported as h-α7 Ki (μM)).

[00695] [³H]BRL 43694 competition binding (h-5HT₃ Ki (μM))

[00696] [³H]BRL 43694 competition binding assay was performed under contract by Cerep Poitiers, France following the methods described in Hope, A.G *et al.*, “Characterization of a human 5-hydroxytryptamine 3 receptor type A (h5-HT₃R-AS) subunit stably expressed in HEK 293 cells,” Brit. J. Pharmacol., (1996) 118: 1237-1245.

[00697] In brief, Chinese Hamster Ovary (CHO) cells stably expressing human 5-HT₃ serotonin receptors, grown to confluence in 175 cm² flasks. Following aspiration of the culture medium, cells were harvested by mechanical agitation in ice cold PBS containing (in mM): (150 NaCl, 8 K₂HPO₄, 2 KH₂PO₄, pH 7.4, 37°C), centrifuged at 4,000 g for 10 min and subsequently stored as a cell pellet at -80 C. When required, the pellet was thawed and resuspended in ice cold homogenization buffer (Tris 50 mM, EGTA 5.0 mM, phenylmethylsulphonylfluoride 0.1 mM, pH 7.6) and homogenized. The homogenate was centrifuged at 48,000 g for 10 minutes at 40°C. The resulting pellet was resuspended in ice cold binding buffer comprising (in mM): NaCl 140, KCl 2.8, CaCl₂ 1.0; MgCl₂, 2.0; HEPES 10 (pH 7.4) and centrifuged as above. The pellet was resuspended in ice cold binding buffer and the protein concentration was determined by the method of Lowry *et al.*, “Protein measurement with the

Folin phenol reagent,” J. Biol. Chem., (1953) 193, 265-275). The membrane homogenate was adjusted to a protein concentration of approximately 600 mg/mL in binding buffer. Assay tubes were loaded with equal volumes of binding buffer containing [³H]BRL 43694 and test compound and 0.5 mL of membrane homogenate in a total reaction volume of 1 mL. Binding was initiated by the addition of the membrane homogenate and allowed to proceed for 120 min. at room temperature. Bound and free radioligand were separated by the addition of 3 mL of ice-cold binding buffer and immediate vacuum filtration through pre-soaked (0.1% (v/v) polyethyleneimine) Whatman GF/B filters. Filters were washed with a further 2 x 3 mL applications of binding buffer and counted for radioactivity using a scintillation counter.

[00698] The results were expressed as a percent inhibition of control specific binding obtained in the presence of the test compounds where Inhibition (%) = 100 – [(measured specific binding/control specific binding) x 100].

[00699] The IC₅₀ values (concentration causing a half-maximal inhibition of control specific binding) and Hill coefficients (nH) were determined by non-linear regression analysis of the competition curves generated with mean replicate values using Hill equation

$$Y = D + \left[\frac{A - D}{1 + (C/C_{50})^{nH}} \right]$$

where Y = specific binding, A = left asymptote of the curve, D = right asymptote of the curve, C = compound concentration, C₅₀ = IC₅₀, and nH = slope factor. This analysis was performed using software developed at Cerep (Hill software) and validated by comparison with data generated by the commercial software SigmaPlot® 4.0 for Windows® (© 1997 by SPSS Inc.).

[00700] The inhibition constants (K_i) were calculated using the Cheng Prusoff equation

$$K_i = \frac{IC_{50}}{(1 + L/K_D)}$$

where L = concentration of radioligand in the assay, and K_D = affinity of the radioligand for the receptor.

[00701] A scatchard plot is used to determine the K_D. Results are provided in Table 2 (reported as h-5HT₃ K_i (μM)).

[00702] For reference, the literature reported α7 nAChR agonist AQW051 has a K_i of 255 nM in the above described assay provided by Cerep (lit: K_i = 28 nM; radioligand binding assay using recombinantly expressed human α7-nAChR and [¹²⁵I] α-BTX radioligand; Feuerbach *et al.*, Br. J. Pharmacol., 2014, doi: 10.1111/bph.13001).

[00703] Oocyte Electrophysiology Screen (% ACh @ 10μM Oocyte)

[00704] The Oocyte Electrophysiology Screen studies were performed under contract by HiQScreen Geneva, Switzerland. All experiments were carried out at human α7 nAChRs transiently expressed in *Xenopus laevis* oocytes using the method of cDNA expression. Currents evoked by

acetylcholine or other agonist ligands were recorded using the standard two-electrode voltage-clamp configuration (TEVC). *X. laevis* oocytes were prepared and injected using standard procedures. Briefly, ovaries were harvested from *X. laevis* females that were deeply anesthetized and pithed following the animal rights rule from the Geneva canton. A small piece of ovary was isolated for immediate preparation while the remaining part was placed at 4°C in a sterile Barth solution containing in mM: NaCl 88, KCl 1, NaHCO₃ 2.4, HEPES 10, MgSO₄·7H₂O 0.82, Ca(NO₃)₂·4H₂O 0.33, CaCl₂·6H₂O 0.41, at pH 7.4, and supplemented with 20 µg/mL of kanamycin, 100 unit/mL penicillin and 100 µg/mL streptomycin. On the second day following dissociation, oocytes were injected with 2 ng of cDNA per oocyte containing the gene encoding human $\alpha 7$ nicotinic acetylcholine receptor subunits using an automated injector (Hogg et al., 2008). All recordings were performed at 18°C and cells were superfused with OR2 medium containing in mM: NaCl 82.5, KCl 2.5, HEPES 5, CaCl₂·2H₂O 2.5, pH 7.4. Cells were held at -80 mV. Data were filtered at 10 Hz, captured at 100 Hz and analyzed using proprietary data acquisition and analysis software running under Matlab (Mathworks Inc.).

[00705] Experimental protocol and analysis

[00706] After establishing a baseline transmembrane current, acetylcholine (ACh) was applied for 5 seconds at a concentration of 0.2 mM to establish a control ACh-evoked current response. Following a wash period of 90 s in OR2 medium (free of ACh), cells were then exposed for 30 s to the test compound applied at 0.01 mM. The same reference ACh test pulse was immediately given at the end of the compound exposure and again after 90 s of recovery in OR2 Medium (free of ACh or test compound). All data were determined in triplicate. The response evoked by the test compound was expressed as a percentage of that evoked by ACh:

$$\text{Response (\%ACh)} = 100 \times (I_{\text{test}} / I_{\text{ACh}})$$

where I_{test} is the peak inward current measured during exposure to 0.01 mM of test compound and I_{ACh} is the peak inward current measured in the presence of ACh.

[00707] Results are provided in Table 2 (reported as % ACh @ 10µM Oocyte).

Table 2:

Compound	h- $\alpha 7$ K _i (µM)	h-5HT ₃ K _i (µM)	% ACh @ 10µM Oocyte
1-P1	0.815	1.9	483
1-P2	5.05	0.069	206
2-P1	0.37	0.019	418
2-P2	3.5		363
3-P1	0.032	1.4	465
3-P2	0.4575	0.25	379
(R)-3	0.057		
4-P1	1		326
4-P2	16		17

5-P1	1	0.16	548
5-P2	0.159	1.85	645
6-P1	0.96	1.6	157
6-P2	0.41	>10	355
7-P1	0.36	0.74	326
7-P2	0.032	4.2	427
8-P1	6.8		26
8-P2	1.2		221
9-P1	0.0435	0.0595	
9-P2	0.29	0.026	
(R)-10	>30		
(R)-11	0.54	1.9	
(R)-12	0.74		
(R)-13	>30		
(S)-13	>30		
(R)-14	>30		
(R)-15	0.93		
(S)-15	52		
(R)-16	2.2		
(R)-17	1.7	0.24	
(R)-18	2		
(S)-18	24		
(R)-19	9.7		
(R)-20	0.22	0.58	
(R)-21	0.82	1.5	
(R)-22	0.14	3	
(R)-23	0.071	0.19	
(R)-24	0.14	0.4	
(R)-25	0.245	4.1	
(R)-26	11	1.2	
(R)-27	0.7		
(R)-28	14		
(R)-29	0.925	>10	
(R)-30	2.5	0.28	
(S)-30	7.4		
(R)-31	0.18	0.029	
(R)-32	4.3		
(S)-32	18		
(R)-33	0.75	>10	
(S)-33	5.8		
(R)-35	0.18	0.065	
(R)-36	0.7		
(R)-37	9		

(R)-39	0.12	6.9	
(R)-41	0.046	1.6333	
(S)-41	0.67		
(R)-42	0.26	1.55	
(S)-42	0.98		
(R)-43	0.044	0.93	

[00708] Example 45:

[00709] Novel Object Recognition Task:

[00710] The Novel Object Recognition (NOR) task is a behavioral assay commonly used to evaluate cognition, particularly recognition memory, in rodent models of CNS disorders. This test is based on the spontaneous tendency of rodents to spend more time exploring a novel object compared to a familiar one. The choice to explore the novel object reflects the use of learning and recognition memory. The assay is commonly used to evaluate potential therapeutic agents for Alzheimer's disease, other neurodegenerative diseases and psychiatric disorders.

[00711] Procedure:

[00712] Male Wistar rats (Harlan Laboratories) weighing 350-400 grams were housed under a reversed light cycle and are tested during the dark cycle. Testing was done under low lux conditions, measured to be ~2-7 lux under red light. Animals were habituated and weighed one day prior to testing. During habituation, animals were placed in a cylindrical arena and allowed to explore for 3 minutes. Training (T1) was conducted approximately 24 hours later, with one set of identical objects placed on opposite sides of the arena. Animals were allowed to explore the objects in 3-minute sessions. Animals were dosed with a designated treatment 15-60 minutes prior to testing depending on the pharmacokinetic profile of the compound before the start of T1. Drug or vehicle was dosed subcutaneously based on body weight at 5 mL/kg. Testing (T2) was done at 48 hours after T1. During testing, one familiar object is replaced with a novel object. Animals were allowed to explore both objects in 3-minute sessions.

[00713] Equipment Specification:

[00714] Animals were tracked using Noldus Ethovision XT (EthoVision XT version: 8.5, Noldus Inc. Wageningen, Netherlands) tracking software, using a 2 centimeter (cm) perimeter for each object as a separate zone. The test arena consisted of a cylinder, 80 cm diameter with 40 cm high walls of black acrylic that was opaque and matte. Objects were custom fabricated shapes (cone and bullet) similar in overall size (8cm high x 8cm diameter) and were counterbalanced between treatment groups.

[00715] Data Analysis and Statistics:

[00716] Contact time was defined as the amount of time (seconds) an animal spent within the 2 cm perimeter of an object. All animals that had ≤ 5 seconds total contact time were excluded from the

study. Statistical significance was determined using a Mann Whitney U-test and the criterion was set at $p < 0.05$.

[00717] Results:

[00718] Natural forgetting in an object recognition task in male Wistar rats ($n = 4-20/\text{group}$). Test compound was administered via sub-cutaneous administration 30 minutes before T1. Test compounds improved object recognition using a 48-hour retention interval (mean $\pm$ SEM). * $p < 0.05$ = novel (N) vs. familiar (F) object. Results are illustrated in Table 3.

Table 3:


Compound	Active doses (mg/kg)
3-P1	0.003, 0.01
(R)-41	0.003, 0.03, 0.1

[00719] All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

[00720] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

What is claimed is:

1. A compound represented by Formula (Ia) or Formula (Ib):



wherein:

R^1 , R^2 , R^3 , and R^4 independently represent $-H$, $-D$, halogen radical, $-CN$, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, an unbranched $-OC_1$ - C_4 -alkyl, a branched or cyclic $-OC_3$ - C_4 -alkyl, $-N(R^5)(R^6)$, $-(CO)N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-SO_2C_1$ - C_4 -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1$ - C_4 -alkyl, an aryl radical, or a heteroaryl radical; wherein the alkyl portion of the unbranched C_1 - C_4 -alkyl radical, the branched C_3 - C_4 -alkyl radical, the C_3 - C_6 -cycloalkyl radical, the unbranched $-OC_1$ - C_4 -alkyl, the branched or cyclic $-OC_3$ - C_4 -alkyl, the $-SO_2C_1$ - C_4 -alkyl, the $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, or the $-N(R^5)SO_2C_1$ - C_4 -alkyl, may be independently substituted with up to 5 radical substituents comprising: $-D$, halogen radical, $=O$, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1$ - C_4 -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1$ - C_4 -alkyl, $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an unbranched C_1 - C_6 -alkyl radical, a branched C_3 - C_6 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, a C_1 - C_6 -hydroxyalkyl radical, a C_1 - C_2 -haloalkyl radical, or $-OC_1$ - C_2 -haloalkyl radical; and wherein the aryl radical or the heteroaryl radical may be independently substituted with up to 5 radical substituents comprising: $-D$, halogen radical, $-CN$, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1$ - C_4 -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1$ - C_4 -alkyl, $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an unbranched C_1 - C_6 -alkyl radical, a branched C_3 - C_6 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, a C_1 - C_6 -hydroxyalkyl radical, a C_1 - C_2 -haloalkyl radical, or $-OC_1$ - C_2 -haloalkyl radical;

R^5 and R^6 independently represent $-H$; an unbranched C_1 - C_6 -alkyl radical, a branched C_3 - C_6 -alkyl radical; a C_3 - C_6 -cycloalkyl radical; or the $N(R^5)(R^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a C_2 - C_6 -alkyl di-radical or a (3-6 membered)-heteroalkyl di-radical; wherein the (3-6

membered)-heteroalkyl di-radical comprises at least one ring atom selected from the group consisting of oxygen, nitrogen, and sulfur, with the proviso that when the at least one ring atom is nitrogen, the nitrogen is independently substituted with $-H$, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_4 -cycloalkyl radical, $-(CO)$ -unbranched C_1 - C_4 -alkyl, $-(CO)$ -branched C_3 - C_4 -alkyl, $-(SO_2)$ -unbranched C_1 - C_4 -alkyl, or $-(SO_2)$ -branched C_3 - C_4 -alkyl, and with the further proviso that when the at least one ring atom is sulfur, the sulfur may be independently substituted with 0 to 2 $=O$; wherein the C_2 - C_6 -alkyl di-radical or the alkyl portion of said (3-6 membered)-heteroalkyl di-radical may be independently substituted with up to 5 radical substituents comprising: $-D$, halogen radical, $=O$, an unbranched C_1 - C_6 -alkyl radical, or a branched C_3 - C_6 -alkyl radical; and

m independently represents an integer from 1 to 6;
or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1, wherein R^1 independently represents $-H$, $-D$, or halogen radical.
3. The compound of claim 1 or claim 2, wherein R^1 independently represents $-H$, $-D$, $-F$, or $-Cl$.
4. The compound of any one of claims 1-3, wherein R^1 independently represents $-H$, $-D$, or $-F$.
5. The compound of any one of claims 1-4, wherein R^1 independently represents $-H$ or $-D$.
6. The compound of any one of claims 1-5, wherein R^2 independently represents $-H$, $-D$, or halogen radical.
7. The compound of any one of claims 1-6, wherein R^2 independently represents $-H$, $-D$, $-F$, or $-Cl$.
8. The compound of any one of claims 1-7, wherein R^2 independently represents $-H$ or $-D$.
9. The compound of any one of claims 1-7, wherein R^2 independently represents $-F$.
10. The compound of any one of claims 1-9, wherein R^3 independently represents $-H$, $-D$, $-F$, $-Cl$, $-Br$, $-CN$, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, an unbranched $-OC_1$ - C_4 -alkyl, a branched or cyclic $-OC_3$ - C_4 -alkyl, $-N(R^5)(R^6)$,

$-(CO)N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-SO_2C_1-C_4\text{-alkyl}$, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1-C_4\text{-alkyl}$,
 $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1-C_4\text{-alkyl}$, an aryl radical, or a heteroaryl radical; wherein the
 alkyl portion of the unbranched $C_1-C_4\text{-alkyl}$ radical, the branched $C_3-C_4\text{-alkyl}$ radical, the $C_3-C_6\text{-}$
 cycloalkyl radical, the unbranched $-OC_1-C_4\text{-alkyl}$, the branched or cyclic $-OC_3-C_4\text{-alkyl}$, the $-SO_2C_1\text{-}$
 $C_4\text{-alkyl}$, the $-(CH_2)_mSO_2C_1-C_4\text{-alkyl}$, or the $-N(R^5)SO_2C_1-C_4\text{-alkyl}$, may be independently substituted
 with up to 5 radical substituents comprising: $-D$, $-F$, $-Cl$, $-Br$, $=O$, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$,
 $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1-C_4\text{-alkyl}$, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1-C_4\text{-alkyl}$,
 $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1-C_4\text{-alkyl}$, $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an unbranched $C_1\text{-}$
 $C_6\text{-alkyl}$ radical, a branched $C_3-C_6\text{-alkyl}$ radical, a $C_3-C_6\text{-cycloalkyl}$ radical, a $C_1-C_6\text{-hydroxyalkyl}$
 radical, a $C_1-C_2\text{-haloalkyl}$ radical, or $-OC_1-C_2\text{-haloalkyl}$ radical; and wherein the aryl radical or the
 heteroaryl radical may be independently substituted with up to 5 radical substituents comprising: $-D$,
 $-F$, $-Cl$, $-Br$, $-CN$, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1-C_4\text{-}$
 $alkyl$, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1-C_4\text{-alkyl}$, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1-C_4\text{-alkyl}$,
 $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an unbranched $C_1-C_6\text{-alkyl}$ radical, a branched $C_3-C_6\text{-alkyl}$ radical,
 a $C_3-C_6\text{-cycloalkyl}$ radical, a $C_1-C_6\text{-hydroxyalkyl}$ radical, a $C_1-C_2\text{-haloalkyl}$ radical, or $-OC_1-C_2\text{-}$
 $haloalkyl$ radical.

11. The compound of any one of claims 1-10, wherein R^3 independently represents $-H$, $-D$, $-F$,
 $-Cl$, $-Br$, $-CN$, an unbranched $C_1-C_4\text{-alkyl}$ radical, a branched $C_3-C_4\text{-alkyl}$ radical, a $C_3-C_4\text{-cycloalkyl}$
 radical, an unbranched $-OC_1-C_4\text{-alkyl}$, a branched or cyclic $-OC_3-C_4\text{-alkyl}$, $-OCHF_2$, $-OCHF_2$,
 $-OCF_3$, $-OCH_2CF_3$, $-N(R^5)(R^6)$, $-(CO)N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-SO_2CH_3$, $-SO_2N(R^5)(R^6)$,
 $-CH_2CH_2SO_2C_1-C_4\text{-alkyl}$, or $-N(R^5)SO_2CH_3$, an aryl radical, or a heteroaryl radical; wherein the alkyl
 portion of the unbranched $C_1-C_4\text{-alkyl}$ radical, the branched $C_3-C_4\text{-alkyl}$ radical, the $C_3-C_4\text{-cycloalkyl}$
 radical, the unbranched $-OC_1-C_4\text{-alkyl}$, the branched or cyclic $-OC_3-C_4\text{-alkyl}$, or the $-CH_2CH_2SO_2C_1\text{-}$
 $C_4\text{-alkyl}$, may be independently substituted with up to 5 radical substituents comprising: $-D$, $-F$, $-Cl$,
 $-Br$, $=O$, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1-C_4\text{-alkyl}$,
 $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1-C_4\text{-alkyl}$, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1-C_4\text{-alkyl}$,
 $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an unbranched $C_1-C_6\text{-alkyl}$ radical, a branched $C_3-C_6\text{-alkyl}$ radical,
 a $C_3-C_6\text{-cycloalkyl}$ radical, a $C_1-C_6\text{-hydroxyalkyl}$ radical, a $C_1-C_2\text{-haloalkyl}$ radical, or $-OC_1-C_2\text{-}$
 $haloalkyl$ radical; and wherein the aryl radical or the heteroaryl radical may be independently
 substituted with up to 5 radical substituents comprising: $-D$, $-F$, $-Cl$, $-Br$, $-CN$, $-OR^5$, $-(CH_2)_mOR^5$,
 $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1-C_4\text{-alkyl}$, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1-C_4\text{-}$
 $alkyl$, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1-C_4\text{-alkyl}$, $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an
 unbranched $C_1-C_6\text{-alkyl}$ radical, a branched $C_3-C_6\text{-alkyl}$ radical, a $C_3-C_6\text{-cycloalkyl}$ radical, a $C_1-C_6\text{-}$
 $hydroxyalkyl$ radical, a $C_1-C_2\text{-haloalkyl}$ radical, or $-OC_1-C_2\text{-haloalkyl}$ radical.

12. The compound of any one of claims 1-11, wherein R^3 independently represents $-F$, $-Cl$, $-Br$, $-CN$, $-CH_3$, $-CH_2CH_3$, cyclopropyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-O$ -cyclopropyl, $-OCF_3$, $-SO_2CH_3$, a phenyl radical, or an N-pyrazole radical, or an oxadiazole radical; wherein the phenyl radical, the N-pyrazole radical, or the oxadiazole radical may be independently substituted with up to 5 radical substituents comprising: $-D$, $-F$, $-Cl$, $-Br$, $-CN$, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, an unbranched C_1 - C_6 -alkyl radical, a branched C_3 - C_6 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, a C_1 - C_6 -hydroxyalkyl radical, a C_1 - C_2 -haloalkyl radical, or $-OC_1$ - C_2 -haloalkyl radical.

13. The compound of any one of claims 1-12, wherein R^3 independently represents $-F$, $-Cl$, $-Br$, $-CN$, $-CH_3$, $-CH_2CH_3$, cyclopropyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-O$ -cyclopropyl, $-OCF_3$, $-SO_2CH_3$, a phenyl radical, an N-pyrazole radical, or an oxadiazole radical; wherein the phenyl radical, the N-pyrazole radical, or the oxadiazole radical may be substituted with up to 5 radical substituents comprising: $-D$, $-F$, $-Cl$, $-Br$, $-CN$, $-OR^5$, $-CH_3$, cyclopropyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCF_3$, or $-OCH_2CF_3$.

14. The compound of any one of claims 1-13, wherein R^3 independently represents $-F$, $-Cl$, $-Br$, $-CN$, $-CH_3$, $-CH_2CH_3$, cyclopropyl radical, $-CHF_2$, $-CH_2F$, $-CF_3$, $-OCH_3$, $-OCH_2CH_3$, $-OCH(CH_3)_2$, $-O$ -cyclopropyl, or $-OCF_3$.

15. The compound of any one of claims 1-14, wherein R^3 independently represents $-F$, $-Cl$, $-Br$, $-CH_3$, or $-OCH_3$.

16. The compound of any one of claims 1-15, wherein R^3 independently represents $-Cl$, $-CH_3$, or $-OCH_3$.

17. The compound of any one of claims 1-16, wherein R^4 independently represents $-H$, $-D$, $-F$, $-Cl$, $-Br$, $-CN$, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_6 -cycloalkyl radical, an unbranched $-OC_1$ - C_4 -alkyl, a branched or cyclic $-OC_3$ - C_4 -alkyl, $-N(R^5)(R^6)$, $-(CO)N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-SO_2C_1$ - C_4 -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1$ - C_4 -alkyl, an aryl radical, or a heteroaryl radical; wherein the alkyl portion of the unbranched C_1 - C_4 -alkyl radical, the branched C_3 - C_4 -alkyl radical, the C_3 - C_6 -cycloalkyl radical, the unbranched $-OC_1$ - C_4 -alkyl, the branched or cyclic $-OC_3$ - C_4 -alkyl, the $-SO_2C_1$ - C_4 -alkyl, the $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, or the $-N(R^5)SO_2C_1$ - C_4 -alkyl, may be independently substituted with up to 5 radical substituents comprising: $-D$, $-F$, $-Cl$, $-Br$, $=O$, $-OR^5$, $-(CH_2)_mOR^5$, $-N(R^5)(R^6)$, $-NR^5(CO)(R^6)$, $-(CH_2)_mN(R^5)(R^6)$, $-SO_2C_1$ - C_4 -alkyl, $-SO_2N(R^5)(R^6)$, $-(CH_2)_mSO_2C_1$ - C_4 -alkyl, $-(CH_2)_mSO_2N(R^5)(R^6)$, $-N(R^5)SO_2C_1$ - C_4 -alkyl, $-(CO)(CH_2)_mR^5$, $-(CO)N(R^5)(R^6)$, $-OCF_3$, and

unbranched C₁-C₆-alkyl radical, a branched C₃-C₆-alkyl radical, a C₃-C₆-cycloalkyl radical, a C₁-C₆-hydroxyalkyl radical, a C₁-C₂-haloalkyl radical, or -OC₁-C₂-haloalkyl radical; and wherein aryl radical or the heteroaryl radical may be independently substituted with up to 5 radical substituents comprising: -D, -F, -Cl, -Br, -CN, -OR⁵, -(CH₂)_mOR⁵, -N(R⁵)(R⁶), -NR⁵(CO)(R⁶), -(CH₂)_mN(R⁵)(R⁶), -SO₂C₁-C₄-alkyl, -SO₂N(R⁵)(R⁶), -(CH₂)_mSO₂C₁-C₄-alkyl, -(CH₂)_mSO₂N(R⁵)(R⁶), -N(R⁵)SO₂C₁-C₄-alkyl, -(CO)(CH₂)_mR⁵, -(CO)N(R⁵)(R⁶), -OCF₃, an unbranched C₁-C₆-alkyl radical, a branched C₃-C₆-alkyl radical, a C₃-C₆-cycloalkyl radical, a C₁-C₆-hydroxyalkyl radical, a C₁-C₂-haloalkyl radical, or -OC₁-C₂-haloalkyl radical.

18. The compound of any one of claims 1-17, wherein R⁴ independently represents -H, -D, -F, -Cl, -CN, an unbranched C₁-C₃-alkyl radical, a branched C₃-C₄-alkyl radical, a C₃-C₄-cycloalkyl radical, an unbranched -OC₁-C₃-alkyl, a branched or cyclic -OC₃-C₄-alkyl; wherein the alkyl portion of the unbranched C₁-C₃-alkyl radical, the branched C₃-C₄-alkyl radical, the C₃-C₄-cycloalkyl radical, the unbranched -OC₁-C₃-alkyl, or the branched or cyclic -OC₃-C₄-alkyl, may be independently substituted with up to 5 radical substituents comprising: -D, -F, -Cl, -Br, =O, -OR⁵, -(CH₂)_mOR⁵, -N(R⁵)(R⁶), -NR⁵(CO)(R⁶), -(CH₂)_mN(R⁵)(R⁶), -(CO)(CH₂)_mR⁵, -(CO)N(R⁵)(R⁶), -OCF₃, an unbranched C₁-C₆-alkyl radical, a branched C₃-C₆-alkyl radical, a C₃-C₆-cycloalkyl radical, a C₁-C₆-hydroxyalkyl radical, a C₁-C₂-haloalkyl radical, or -OC₁-C₂-haloalkyl radical.

19. The compound of any one of claims 1-18, wherein R⁴ independently represents -H, -D, -F, -Cl, -CN, -CH₃, cyclopropyl radical, cyclobutyl radical, -CH₂F, -CHF₂, -CF₃, -OCH₃, -OCH₂CH₃, -OCH(CH₃)₂, -O-cyclopropyl, or -OCF₃.

20. The compound of any one of claims 1-19, wherein R⁴ independently represents -H, -D, -F, -Cl, -CH₃, -OCH₃, -OCH₂CH₃, or -CF₃.

21. The compound of any one of claims 1-20, wherein R¹ and R² independently represent -H or -D.

22. The compound of any one of claims 1-20, wherein R² and R⁴ independently represent -H or -D.

23. The compound of any one of claims 1-20, wherein R¹ and R⁴ independently represent -H or -D.

24. The compound of any one of claims 1-20, wherein R^1 , R^2 , and R^4 independently represent $-H$ or $-D$.
25. The compound of any one of claims 1-24, wherein R^3 independently represents $-Cl$.
26. The compound of any one of claims 1-24, wherein R^3 independently represents $-CH_3$.
27. The compound of any one of claims 1-24, wherein R^3 independently represents $-OCH_3$.
28. The compound of any one of claims 25-27, wherein R^4 independently represents $-H$ or $-D$.
29. The compound of any one of claims 25-27, wherein R^4 independently represents $-F$.
30. The compound of any one of claims 25-27, wherein R^4 independently represents $-Cl$.
31. The compound of any one of claims 25-27, wherein R^4 independently represents $-CH_3$.
32. The compound of any one of claims 25-27, wherein R^4 independently represents $-OCH_3$.
33. The compound of any one of claims 25-27, wherein R^4 independently represents $-CF_3$.
34. The compound of any one of claims 1-33, wherein R^5 independently represents $-H$; an unbranched C_1 - C_6 -alkyl radical; a branched C_3 - C_6 -alkyl radical; a C_3 - C_6 -cycloalkyl radical.
35. The compound of any one of claims 1-34, wherein R^6 independently represents $-H$; an unbranched C_1 - C_6 -alkyl radical; a branched C_3 - C_6 -alkyl radical; a C_3 - C_6 -cycloalkyl radical.
36. The compound of any one of claims 1-33, wherein the $N(R^5)(R^6)$ moiety forms a cycle, wherein R^5 and R^6 taken together represent a C_2 - C_6 -alkyl di-radical or a (3-6 membered)-heteroalkyl di-radical; wherein the (3-6 membered)-heteroalkyl di-radical comprises at least one ring atom selected from the group consisting of oxygen, nitrogen, and sulfur, with the proviso that when the at least one ring atom is nitrogen, the nitrogen is independently substituted with $-H$, an unbranched C_1 - C_4 -alkyl radical, a branched C_3 - C_4 -alkyl radical, a C_3 - C_4 -cycloalkyl radical, $-(CO)$ -unbranched C_1 - C_4 -alkyl, $-(CO)$ -branched C_3 - C_4 -alkyl, $-(SO_2)$ -unbranched C_1 - C_4 -alkyl, or $-(SO_2)$ -branched C_3 - C_4 -alkyl, and with the further proviso that when the at least one ring atom is sulfur, the sulfur may be independently substituted with 0 to 2 $=O$; wherein the C_2 - C_6 -alkyl di-radical or the alky portion of said (3-6 membered)-heteroalkyl di-radical may be independently substituted with up to 5 radical

substituents comprising: -D, halogen radical, =O, an unbranched C₁-C₆-alkyl radical, or a branched C₃-C₆-alkyl radical.

37. The compound of any one of claims 1-33, wherein the N(R⁵)(R⁶) moiety forms a cycle, wherein R⁵ and R⁶ taken together represent a C₂-C₂-alkyl di-radical.
38. The compound of any one of claims 1-37, wherein m independently represents an integer from 1 to 4.
39. The compound of any one of claims 1-38, wherein m independently represents an integer from 1 to 2.
40. The compound of any one of claims 1-39, wherein the compound is represented by Formula (Ia).
41. The compound of any one of claims 1-39, wherein the compound is represented by Formula (Ib).
42. The compound of any one of claims 1-39, wherein the compound is the pharmaceutically acceptable salt thereof.
43. The compound of any one of claims 1-42, wherein the compound is a single enantiomer or a single diastereomer.
44. The compound of claim 43, wherein the compound is a single enantiomer.
45. The compound of claim 43, wherein the compound is a single diastereomer.
46. The compound of any one of claims 1-45, wherein the compound is selected from the group consisting of:
- N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 - 7-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 - 6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 - 5-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 - 6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 - 6-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
 - 6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

N-(quinuclidin-3-yl)-6-(trifluoromethyl)benzo[d]isoxazol-3-amine;
6,7-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
4-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
3-(quinuclidin-3-ylamino)benzo[d]isoxazole-6-carbonitrile;
6-(methylsulfonyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-(tert-butyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
N-(quinuclidin-3-yl)-6-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
4-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
6-chloro-7-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
4-fluoro-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
4,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-fluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-chloro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-chloro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-isopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-7-carbonitrile;
7-chloro-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-(1H-pyrazol-1-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-(5-methyl-1,3,4-oxadiazol-2-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-methoxy-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6,7-dimethyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
6-bromo-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
and single enantiomers and pharmaceutically acceptable salts thereof.

47. The compound of any one of claims 1-45, wherein the compound is selected from the group consisting of:

6-cyclopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-ethyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
6-chloro-7-cyclobutyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5-fluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-methoxy-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-methyl-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
6-chloro-5,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-(difluoromethyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-isopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-cyclopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-ethoxy-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
7-ethoxy-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-methoxy-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5,6-difluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-5-fluoro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
6-chloro-5-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
5,7-difluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6,7-dichloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
6-chloro-7-ethoxy-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
5,7-difluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine.

and single enantiomers and pharmaceutically acceptable salts thereof.

48. The compound of any one of claims 1-46, wherein the compound is selected from the group consisting of:

(*R*)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

(S)-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-5-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-5-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-N-(quinuclidin-3-yl)-6-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(S)-N-(quinuclidin-3-yl)-6-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(R)-6,7-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6,7-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-4-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-4-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-6-carbonitrile;
(S)-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-6-carbonitrile;
(R)-6-(methylsulfonyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-(methylsulfonyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6-(tert-butyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6-(tert-butyl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-5,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-5,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-N-(quinuclidin-3-yl)-6-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
(S)-N-(quinuclidin-3-yl)-6-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
(R)-7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-7-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-4-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-4-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-7-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-7-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(R)-6,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(S)-6,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

(*R*)-5-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-cyclopropyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-4-fluoro-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-4-fluoro-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-4,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-4,6-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-fluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-fluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-chloro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-chloro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-chloro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-chloro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-isopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-isopropoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-7-carbonitrile;
(*S*)-6-chloro-3-(quinuclidin-3-ylamino)benzo[d]isoxazole-7-carbonitrile;
(*R*)-7-chloro-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-chloro-6-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-(1H-pyrazol-1-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-(1H-pyrazol-1-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-(5-methyl-1,3,4-oxadiazol-2-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-(5-methyl-1,3,4-oxadiazol-2-yl)-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-methoxy-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-methoxy-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-ethoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6,7-dimethyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

(*S*)-6,7-dimethyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-5-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-5-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-bromo-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
(*S*)-6-bromo-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

and pharmaceutically acceptable salts thereof.

49. The compound of any one of claims 1-47, wherein the compound is selected from the group consisting of:

(*R*)-6-cyclopropoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-cyclopropoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-ethyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-ethyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-*N*-(quinuclidin-3-yl)-7-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-*N*-(quinuclidin-3-yl)-7-(trifluoromethoxy)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-cyclobutyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-cyclobutyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5-fluoro-6-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5-fluoro-6-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methoxy-7-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-methoxy-7-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methyl-*N*-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*S*)-6-methyl-*N*-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-5,7-difluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-5,7-difluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-(difluoromethyl)-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-(difluoromethyl)-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-isopropoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-isopropoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-cyclopropoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-cyclopropoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-ethoxy-6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-ethoxy-6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-ethoxy-6-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-7-ethoxy-6-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

(*R*)-6-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methoxy-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*S*)-6-methoxy-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*R*)-5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5,6-difluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5,6-difluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5,7-difluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-5-fluoro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-5-fluoro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-5-fluoro-N-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-5-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-5-fluoro-7-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5,7-difluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-5,7-difluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6,7-dichloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6,7-dichloro-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-ethoxy-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-7-ethoxy-5-fluoro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5,7-difluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
(*S*)-5,7-difluoro-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

and pharmaceutically acceptable salts thereof.

50. The compound of any one of claims 1-46 or 48, wherein the compound is selected from the group consisting of:

(*R*)-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-chloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-methoxy-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6,7-dichloro-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5-fluoro-6-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methyl-N-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;

(*R*)-6-chloro-7-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-*N*-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
(*R*)-6-chloro-5-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
and pharmaceutically acceptable salts thereof.

51. The compound of any one of claims 1-46 or 48, wherein the compound is selected from the group consisting of:

(*R*)-6-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6,7-dichloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-7-fluoro-6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-7-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-chloro-*N*-(quinuclidin-3-yl)-7-(trifluoromethyl)benzo[d]isoxazol-3-amine; and
(*R*)-6-chloro-7-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
and pharmaceutically acceptable salts thereof.

52. The compound of any one of claims 1-46 or 48, wherein the compound is selected from the group consisting of:

(*S*)-6-chloro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-6-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*S*)-6-methoxy-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
(*R*)-5-fluoro-6-methyl-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine; and
(*R*)-6-chloro-5-fluoro-*N*-(quinuclidin-3-yl)benzo[d]isoxazol-3-amine;
and pharmaceutically acceptable salts thereof.

53. A pharmaceutical composition, comprising:

- i) the compound, or pharmaceutically acceptable salt thereof, of any one of claims 1-52; and
- ii) at least one pharmaceutically acceptable carrier, excipient or diluent.

54. A method of treating a patient in need thereof, comprising administering to the patient the pharmaceutical composition of claim 53.

55. A method of treating a patient in need thereof, comprising administering to the patient the compound, or pharmaceutically acceptable salt thereof, of any one of claims 1-52.

56. A method of improving cognition of a patient in need thereof, comprising: administering to the patient the compound, or pharmaceutically acceptable salt thereof, of any one of claims 1-52.
57. A method of improving cognition of a patient in need thereof, comprising administering to the patient a pharmaceutical composition comprising:
- i) the compound, or pharmaceutically acceptable salt thereof, of any one of claims 1-52; and
 - ii) at least one pharmaceutically acceptable carrier, excipient or diluent.
58. A method of treating or improving one or more symptoms associated with a cognitive disease and/or a cognitive impairment in a patient in need thereof, comprising: administering to the patient the compound, or pharmaceutically acceptable salt thereof, of any one of claims 1-52.
59. A method of treating or improving one or more symptoms associated with a cognitive disease and/or a cognitive impairment in a patient in need thereof, comprising administering to the patient a pharmaceutical composition comprising:
- i) the compound, or pharmaceutically acceptable salt thereof, of any one of claims 1-52; and
 - ii) at least one pharmaceutically acceptable carrier, excipient or diluent.
60. The method of any one of claims 54-59, wherein the patient suffers from a cognitive impairment, suffers from a cognitive loss associated with a cognitive impairment, or suffers from one or more symptoms associated with a cognitive impairment.
61. The method of claim 60, wherein the cognitive impairment comprises Limited Cognitive Impairment (LCI), Mild Cognitive Impairment (MCI), Alzheimer's disease, dementia of an Alzheimer's-type, schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, positive symptoms of schizophrenia, negative symptoms of schizophrenia, or schizophrenia with dementia.
62. The method of claim 60, wherein the cognitive impairment is Limited Cognitive Impairment (LCI).
63. The method of claim 60, wherein the cognitive impairment is Mild Cognitive Impairment (MCI).
64. The method of claim 60, wherein the cognitive impairment is Alzheimer's disease.

65. The method of claim 60, wherein the cognitive impairment is dementia of an Alzheimer's-type.
66. The method of claim 60, wherein the cognitive impairment is schizophrenia.
67. The method of claim 60, wherein the cognitive impairment is schizophreniform disorder, schizoaffective disorder, or delusional disorder.
68. The method of claim 60, wherein the cognitive impairment comprises positive symptoms of schizophrenia.
69. The method of claim 60, wherein the cognitive impairment comprises negative symptoms of schizophrenia.
70. The method of claim 60, wherein the cognitive impairment is schizophrenia with dementia.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2016/036689

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 31/439; A61K 31/423; A61P 25/28; C07D 261/20; C07D 453/02 (2016.01)

CPC - A61K 31/423; A61K 31/439; C07D 261/20; C07D 453/02 (2016.05)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC - A61K 31/423; A61K 31/439; A61P 25/28; C07D 261/20; C07D 453/02

CPC - A61K 31/423; A61K 31/439; C07D 261/20; C07D 453/02

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC - 514/305; 514/379; 546/133; IPC - A61K 31/423; A61K 31/439; A61P 25/28; C07D 261/20; C07D 453/02; CPC - A61K 31/423; A61K 31/439; C07D 261/20; C07D 453/02 (keyword delimited)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PatBase, Google Patents, STN, Google Scholar, PubChem

Search terms used: aminobenzisoxazole, quinuclidine, nicotinic acetylcholine, amino, isoxazole, benzisoxazole

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2013/0231365 A1 (KOENIG) 05 September 2013 (05.09.2013) entire document	1-3
A	US 8,273,891 B2 (SCHUMACHER et al) 25 September 2012 (25.09.2012) entire document	1-3
A	US 7,902,217 B2 (XIE et al) 08 March 2011 (08.03.2011) entire document	1-3
A	US 6,911,543 B2 (WALKER et al) 28 June 2005 (28.06.2005) entire document	1-3
A	US 7,795,453 B2 (FLEBNER et al) 14 September 2010 (14.09.2010) entire document	1-3

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

26 July 2016

Date of mailing of the international search report

16 AUG 2016

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, VA 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Blaine R. Copenheaver

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2016/036689

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 4-70
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.