



(86) Date de dépôt PCT/PCT Filing Date: 2000/03/10
(87) Date publication PCT/PCT Publication Date: 2000/09/28
(45) Date de délivrance/Issue Date: 2010/09/21
(85) Entrée phase nationale/National Entry: 2001/09/14
(86) N° demande PCT/PCT Application No.: EP 2000/002117
(87) N° publication PCT/PCT Publication No.: 2000/056885
(30) Priorités/Priorities: 1999/03/18 (EP99105530.2);
1999/05/12 (EP99109503.5)

(51) Cl.Int./Int.Cl. *C12N 15/12* (2006.01),
A61K 38/17 (2006.01), *C07K 14/435* (2006.01),
C07K 16/18 (2006.01), *A61K 38/00* (2006.01)

(72) Inventeurs/Inventors:
STRITTMATTER, WOLFGANG, DE;
GUSSOW, DETLEF, DE;
HOFMANN, UWE, DE;
HEMBERGER, JURGEN, DE;
FOTEV, ZISI, DE;
SCHEUBLE, BERNHARD, DE

(73) Propriétaire/Owner:
MERCK PATENT GESELLSCHAFT MIT
BESCHRAENKTER HAFTUNG, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : PROTEINE PERMETTANT DE BLOQUER L'ADHESION DES PLAQUETTES
(54) Title: PROTEIN FOR BLOCKING PLATELET ADHESION

(57) Abrégé/Abstract:

A naturally occurring protein isolated from the saliva of the medicinal leech *Hirudo medicinalis* is described which strongly binds to collagen thus acting as an inhibitor of natural platelet adhesion to collagen. The protein has a molecular weight of about 12 000, an acidic isoelectric point and contains six cysteins. The protein was sequenced and the gene was cloned from a *H. medicinalis* cDNA-library. Procedures for producing such polypeptide by recombinant techniques are disclosed. The recombinant and the natural occurring proteins are potent inhibitors of collagen-dependent platelet adhesion and therefore useful for the therapeutic treatment of various conditions related to hard disease and diseases of the circulation system. Furthermore the protein is useful for coating natural or artificial collagen surfaces in order to render them nonadhesive for cells and prevent the activation of cells.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/435, 16/18, A61K 38/17	A1	(11) International Publication Number: WO 00/56885 (43) International Publication Date: 28 September 2000 (28.09.00)
(21) International Application Number: PCT/EP00/02117 (22) International Filing Date: 10 March 2000 (10.03.00) (30) Priority Data: 99105530.2 18 March 1999 (18.03.99) EP 99109503.5 12 May 1999 (12.05.99) EP (71) Applicant (for all designated States except US): MERCK PATENT GMBH [DE/DE]; Frankfurter Strasse 250, D-64293 Darmstadt (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): STRITTMATTER, Wolf- gang [DE/DE]; Heyerstr. 16, D-64372 Ober Ramstadt (DE). GÜSSOW, Detlef [DE/DE]; Grafenstr. 17, D-64283 Darmstadt (DE). HOFMANN, Uwe [DE/DE]; Hähnleiner Str. 42, D-64665 Alsbach (DE). HEMBERGER, Jürgen [DE/DE]; Kiebitzweg 2, D-63741 Aschaffenburg (DE). FOTEV, Zisi [DE/DE]; Im Höhsand 15, D-64404 Bick- enbach (DE). SCHEUBLE, Bernhard [DE/DE]; Schelleklin- genweg 5, D-64342 Balkhausen (DE).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: PROTEIN FOR BLOCKING PLATELET ADHESION (57) Abstract A naturally occurring protein isolated from the saliva of the medicinal leech <i>Hirudo medicinalis</i> is described which strongly binds to collagen thus acting as an inhibitor of natural platelet adhesion to collagen. The protein has a molecular weight of about 12 000, an acidic isoelectric point and contains six cysteins. The protein was sequenced and the gene was cloned from a H. medicinalis cDNA-library. Procedures for producing such polypeptide by recombinant techniques are disclosed. The recombinant and the natural occurring proteins are potent inhibitors of collagen-dependent platelet adhesion and therefore useful for the therapeutic treatment of various conditions related to hard disease and diseases of the circulation system. Furthermore the protein is useful for coating natural or artificial collagen surfaces in order to render them nonadhesive for cells and prevent the activation of cells.		

26474-568

1

Protein for blocking platelet adhesion

Summary of the invention

A naturally occurring protein isolated from the saliva of the medicinal leech *Hirudo medicinalis* is described which strongly binds to collagen thus acting as an inhibitor of natural platelet adhesion to collagen. The protein has a molecular weight of about 12000, an acidic isoelectric point and contains six cysteins. The protein was sequenced and the gene was cloned from a *H. medicinalis* cDNA-library. Procedures for producing such polypeptide by recombinant techniques are disclosed. The recombinant and the natural occurring proteins are potent inhibitors of collagen-dependent platelet adhesion and therefore useful for the therapeutic treatment of various conditions related to hard disease and diseases of the circulation system. Furthermore the protein is useful for coating natural or artificial collagen surfaces in order to render them nonadhesive for cells and prevent the activation of cells.

Thus, according to one aspect of the present invention, there is provided a polypeptide isolated from *H. medicinalis* having a molecular weight of about 12,000 $\pm$ 1 kD with the biological activity of an inhibitor of collagen-dependent platelet adhesion, which is at least 80% identical to the amino acid sequence of SEQ ID NO:2 over its entire length.

According to another aspect of the present invention, there is provided an isolated polynucleotide comprising a DNA sequence that is at least 80% identical to SEQ ID NO:1 over its entire length, wherein said polynucleotide encodes the polypeptide as described herein.

26474-568

1a

According to another aspect of the present invention, there is provided an expression vector comprising the DNA sequence as described herein.

According to another aspect of the present invention, there is provided a host cell comprising the expression vector as described herein.

According to another aspect of the present invention, there is provided an expression system comprising the host cell as described herein.

According to another aspect of the present invention, there is provided a process for producing a polypeptide comprising culturing the host cell as described herein under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture supernatant or cell residue.

According to another aspect of the present invention, there is provided an antibody immunospecific for the polypeptide as described herein.

According to another aspect of the present invention, there is provided a pharmaceutical composition comprising the polypeptide as described herein, together with a pharmaceutically acceptable carrier or excipient.

According to another aspect of the present invention, there is provided a medicament for the treatment of a thromboembolic disease comprising the polypeptide as described herein and a pharmaceutically acceptable carrier.

According to another aspect of the present invention, there is provided a use of the polypeptide as described herein for ex vivo coating of artificial surfaces.

26474-568

1b

According to another aspect of the present invention, there is provided a use of the polypeptide as described herein for ex vivo modifying intraocular lenses in order to lessen the thrombogenicity of the lens material.

5 According to another aspect of the present invention, there is provided a use of the antibody as described herein to identify ex vivo the polypeptide as described herein.

 According to another aspect of the present
10 invention, there is provided a method for determining whether a compound competes with the binding of the polypeptide as described herein to collagen, the method comprising applying said polypeptide to collagen in the absence and presence of said compound and determining the
15 binding rate of said polypeptide in the absence of said competing compound compared to the respective binding rate in the presence of said compound.

Field of invention

 In haemostasis or thrombosis, platelets adhere to
20 the cell-extracellular matrix of an injured vessel and cover the surface of the damaged area. Preventing this important initial step in the pathogenesis of thrombosis and arterial occlusion should be of therapeutic benefit in the effort of to prevent thrombotic diseases. Collagen is considered to
25 be the most thrombogenic surface component and has been shown to be a strong stimulant for platelet adhesion, aggregation and the release of their granules leading to the recruitment of (Ruggeri, Z.M. et al.; Seminars in Hematology, 1994, 31, 229-39) additional platelets to this
30 area to form aggregates or a thrombus. The initial contact of the platelets to the vessel surface is mediated by collagen bound von Willebrand Factor (vWF) and a specific

26474-568

1c

vWF receptor on platelets, the glycoprotein Ib-V-IX complex. In addition ADP, epinephrine and circulating clotting factors drive the further activation process of platelets while simultaneously an increase in thrombin activity

contributes to the formation of the cross-linked fibrin clot. Platelet-platelet aggregation supports this process and is mainly driven by fibrinogen as a mediator that bridges cells through the glycoprotein IIb/IIIa receptor.

- 5 This normal physiological response is quite critical in the course of the pathological process where platelets adhere to collagenes exposed in sclerotic lesions (Van der Rest M. et al.; FASEB Journal, 1991, 5, 2814-23) and start to build-up occlusions. Depending on the location and extent of the occlusion severe complications such as myocardial infarction, stroke, inflammation or pulmonary
10 embolism may be the severe outcome of this process.

As a direct acting antithrombotic agent heparin which blocks the thrombin activity, thus preventing the formation of fibrin rich thrombi, is the currently most well known drug used in anti-thrombotic interventions. Heparin is widely used in
15 indications such as: unstable angina and acute myocardial infarction. However despite the wide use several severe shortcomings such as intravenous application, requirement for anti-thrombin-III as a cofactor, reduced affinity for clot-bound thrombin, its inactivation by several plasma proteins, the occasional induction of thrombocytopenia and its biological heterogeneity remain
20 unresolved. As a consequence the results of using heparin in the clinical setting have not been overwhelming so far.

Recent development of low molecular weight heparin has contributed a version for subcutaneous application, however the therapeutic benefit over the standard
25 heparin has been modest. Unfortunately the same applies to the other directly acting antithrombins such as Hirudin, Hirulog and Warfarin. It turned out, that one of the major problems seems to be related to the increased production of thrombin under antithrombotic treatment (Rao, A.K et al., Circulation, 1996, 94, 389-2395).

- 30 Other recent strategies have therefore been focussed to the process of prothrombin activation which is driven by Factor Xa. The major challenge is the design of appropriate inhibitors directed to this factor. In summary one would

therefore argue that the full therapeutic potential of this type of intervention has not yet been realized.

Another pannel of therapeutics is represented by the thrombolytic regimens and has been focussed on the development of staphylokinase, streptokinase, urokinase type Plasminogen Activator, tissue type Plasminogen Activator and anisoylated-plasminogen-streptokinase activator complex. The differences in time necessary to inducing reperfusion is remarkable different for each of these thrombolytic agents, however the contribution in terms of reducing the overall mortality is equal for all the products. In addition reocclusion or prolonged bleeding are frequent complications. This might be due to relatively low specificity for fibrin and the short plasma half-life of these compounds. Currently various application regimens and combinations of different fibrinolytic principles are tested in order to overcome some of the current short comings in thrombolytic therapie. The improvement which are expected are however rather small.

Recently a new group of patients occured with probelms such as acute thrombotic occlusion and late restenosis due to procedures such as angioplasty, atherectomy, arterial grafting or vessel wall stenting. The possible therapeutic interventions comprise anti-platelet, antithrombotic and thrombolytic strategies. Various other agents such as ticlopdine acting as ADP antagonists or Calcium ionophore A-23187 and especially Asprin have a direct influence on the platelet function and have been suggested or used to prevent or minimize platelet aggregation. The new anti-platelet adhesion substance according to this invention could as well help to overcome these clinical complications when applied during surgery.

Another complication related to this topic arises if artificial surfaces come in contact with blood, then their is increased tendency to induce thrombotic events by activation of platelets and/or induction of coagulation. These effects may cause failure of vascular grafts, cardiac valves, stents, catheters or any other blood contacting device or material. The ability of the protein disclosed here to create non-thrombogenic surfaces may therefore be further exploited by immobilization of this protein to the materials and devices described above. Such a treatment should render such materials or devices biocompatible and thromboresistant.

Due to the limitations associated with the available antithrombotic agents there is an actual need for new alternative strategies and therapeutics.

Background of the invention

5

A potential for future improvements in the treatment of cardiovascular disorders may be contributed by approaches as disclosed in this invention which directly interfere with the collagen and/or vWF factor induced platelet adhesion.

Several novel inhibitors which prevent platelet adhesion are monoclonal
10 antibodies directed to vWF. It has as well been suggested that glycoprotein IIb/IIIa inhibitors may be beneficial in inhibiting platelet adhesion

Some of these inhibitors like the monoclonal Ab c7E3 have already been tested clinically while others like the KGD- and RGDF-inhibitors are still under study. However, the specificity of most of these new inhibitors is not very well studied ,
15 thus the spectrum of side effects that will be induced by using this inhibitors is still open and deserves careful examination.

A rich source for the screening of new compounds that interfere with collagen induced platelet adhesion is given in nature through blood-sucking animals. Several inhibitors have been isolated from nature as described in the literature: A
20 65 kD protein called Calin isolated from *Hirudo medicinalis* (US 5,587,360, WO 92/07005) (Munro, R., et al., Blood Coagulation and Fibrinolysis, 1991, 2, 179-184) and a 16kD (LAPP) protein isolated from the salivary glands of the leech *Haementeria officinalis* (US 5,324,715). Both proteins have been described as aggregation inhibitors as tested in static assays of collagen dependent platelet
25 aggregation.

Despite a proven in vitro activity LAPP failed to act in several well-established in vivo models (Schaffer L.W. et al.; Arterioscler. Thromb., 1993, 13, 1593-1601) and Connolly T.M. et al.; Thromb. Haemostas., 1993, 69, 589). The soft tick, *Ornithodoros moubata*, also contains an antiplatelet protein (Moubatin) which is
30 active in preventing collagen-stimulated platelet aggregation (Waxman, L. et al.; J. Biol. Chem., 1993, 268, 5445-49). Another inhibitor of platelet aggregation from a blood-sucking bug was disclosed in WO 9309137 by Noeske-Jungblut C. et al.. Smith et al. have isolated a 50 kDa protein from snake venom and a 19 kDa

protein was isolated from the saliva of *Triatoma pallidipennis*, a blood-sucking bug. The protein was found to contain a factor that specifically inhibits collagen-induced platelet aggregation. The 19 kDa protein named pallidipin inhibits collagen-mediated aggregation of platelets in plasma. No inhibition of aggregation
5 stimulated by other effectors (ADP, thrombin, thromboxane A2 mimetic U46619, phorbol ester) was detected. Pallidipin had no effect on platelet adhesion to collagen but inhibited ATP release from platelets. It interacted reversibly with platelets and may share with collagen a common target on them. The precise mechanism of action and therapeutic benefit of this protein is under investigation.

10 Gan et al. described Echistatin as an inhibitor binding to the fibrinogen receptor GP IIa/IIIb (J. Biol. Chem, 1988, 263, 19827-32).

Despite these exciting developments, the need continues to exist for supplying further anticoagulants and antithrombin which have increased efficacy in the inhibition of clot formation, vWF-induced platelet activation or endothelial cell
15 activation and which may be used pharmaceutically and produced in commercially feasible quantities.

Since none of the known proteins described so far has developed into a compound with ideal therapeutic profile the inventors of the present invention decided to go ahead with a new screening strategy in order to detect more
20 relevant proteins.

Description of the Invention

25 In this invention an inhibitor isolated from *H. medicinalis* is described which directly acts on collagen-platelet interaction thus inhibiting platelet activation and early platelet-platelet interaction.

Sofar, there has not been a positive example in the literature which indicates that by using a screening approach that would exclude aggregation inhibitors as
30 well as lytic proteins from a source of naturally occurring compounds one could identify new anti-adhesive mechanisms or compounds. However this strategy was used in this invention. Since at least six different platelet surface

glycoproteins are known to be involved in collagen adhesion and in addition several platelet derived compounds such as von Willebrand factor, fibronectin and thrombospondin have been shown to be involved as indirect mediators of collagen - platelet adhesion there has been little optimism in the beginning to
5 identify a new adhesion inhibitors.

Nevertheless, this approach was used to screen *Hirudo medicinalis* saliva knowing that not all the documented or unknown vWF related inhibitors as well as compounds directly acting on platelet receptors could be excluded. Thus the result of the screening was a surprise: A new protein named Saratin with anti-
10 adhesive activity for platelets which can be isolated from tissues and secretions of well investigated leech of the species *Hirudo medicinalis*.

The present invention comprises the active polypeptide Saratin isolated from the leech *Hirudo medicinalis*. The protein was isolated from saliva by a combination
15 of pressurized dialysis and at least one chromatographic step like anionic exchange chromatography and at least one reversed phase high performance chromatography (RP-HPLC) step. The pressure dialysis step turned out to be absolutely critical during the recovery of Saratin from saliva, since the strong concentration of saliva helped to overcome the otherwise tremendous loss of
20 bioactive Saratin. The isolated Saratin binds strongly to several collagens and blocks the adhesion of platelets to collagen coated such surfaces in a dose dependent fashion.

In order to optimize the screening cascade currently available techniques have
25 been developed to distinguish platelet adhesion and platelet aggregation: the ability of platelets to retard or stop flow through fibers, the contribution of platelets to in vitro clot formation, glass bead adhesion laboratories, or whole blood flows through the filter and platelet adhesion of anticoagulated platelet-rich plasma to filters composed of glass fibers or collagen under a regulated pressure gradient.

30

The protein (named Saratin) is characterized by the amino acid sequences depicted in sequence (SEQ. ID. NO. 2) and is constituted from 103 amino acids which make up a theoretical relative molecular weight of approximately 12068

dalton $\mp$ 1 kDa The protein exhibits a unique primary structure with no significant similarity to other previously described sequences. The protein is rich in aspartic and glutamic acid which contribute to the low isoelectric point of of pH 3.7 $\mp$ 0.5 of the molecule as measured by IEF-PAGE.

5

SDS-PAGE analysis demonstrated a strong shift in mobility upon reduction of the protein prior to electrophoresis, indicating posttranslational modifications. Sequencing of the polypeptide had revealed six cysteine molecules which could make-up post-translational modifications of the protein. Electrospray mass
10 spectrometry of Saratin revealed an actual molecular weight of 12061 indicating that up to three disulphide bonds are involved in the formation of the secondary structure of the native form of the protein.

The adhesion inhibitor according to the present invention is new because it differs from known aggregation inhibitors isolated from leeches especially from Calin or
15 LAPP in the molecular weight, isoelectric point and amino acid sequence and biological activities.

The present invention provides as well isolated DNA comprising a polynucleotide encoding the leech derived platelet adhesion inhibitor having the amino acid
20 sequence as shown for the protein. The nucleotide sequence representing the cDNA clone is shown in SEQ. ID. NO. 1, Position 1-63 of the nucleotide sequence represents a putative 21 amino acid leader sequence and position 64-372 contains an open reading frame coding for a polypeptide of 103 amino acid residues and an amino acid sequence as shown for the mature protein in SEQ.
25 ID. NO. 2.

The present invention also relates to recombinant vectors which include the synthetic gene coding for the leech-derived platelet adhesion inhibitor of the
30 present invention, and a host cell containing the recombinant vectors. Methods for recovering and isolating the expressed proteins have been based on tag-technologies or have been adapted from the purification scheme developed for the naturally occurring Saratin. Depending on the individual protocols used for

extracellular or intracellular expression in yeast cells, insect cells, baby hamster kidney cells and E. coli cells transformed with the appropriate vectors the steps for recovering the recombinant protein from the supernatant or sediments have to be adapted by techniques known to the expert. Excellent expression was found in
5 E. coli as a host, where periplasmatic expression was contributed by insertion of a pelB leader sequence. Products recovery from Escherichia coli (E. coli) was achieved (arround 5 mg/l) after osmolysis and centrifugation. Saccharomyces cerevisiae (S. cerevisiae) (> 10 mg/l culture broth) with the alternative yeast adopted vector was used in a paralelled experiment. The secreted material was
10 isolated by centrifugation. Purification was achieved by cross-flow filtration and ion exchange chromatography. In other expression approches using either COS cells or CHO cells product expression was arround 750 ng/ml. The purified recombinant material proved to be pure and homogeneous by electrophoretic and chromatographic analysis and identical to saliva derived Saratin as demonstrated
15 by amino acid sequencing and molecular mass determination .

The invention also comprises methods for purifying the active protein from crude saliva of the leech and measuring its activity against platelets by static and dynamic assay methods as well as the use of this method to isolate rekombinant protein.

20 Techniques for the production of Saratin, include the Examples 6, 7, 8 and 13 however the expression methods to be used are not restricted to these examples. For instance transgenic mice, or other organisms, including other mammals, may as well be used to express Saratin.

25 Proteins of the present invention include variants which conserve the activity of the disclosed sequences, including fragments or subunits, naturally occuring variants, allelic variants, randomly generated artificial mutants and intentional sequence variations such as adding which conserve activity. Fragments or subunits refers to to any portion of the sequence which contain fewer amino acids
30 than the complete protein e.g. partial sequences excluding portions of the N- and/or C-termini of the complete protein.

The invention further covers hybrid proteins, such as fusion proteins or proteins resulting from the expression of multiple genes within the expression vector, and

may include a polypeptide having the specific activity of a disclosed protein linked by peptide bonds to a second polypeptide. Notably other variants of the proteins of the present invention are included, especially any variants that differ from the isolated protein only by conservative amino acid substitution. Such conservative
5 amino acid substitutions are defined in Taylor et al., J. Mol. Biol., 1986, 188, 233.

Also included are methods for using the proteins to prevent or delay of platelet activation by inhibition of collagen-platelet interactions. The protein is useful in the prevention, prophylaxis, therapy and treatment of thrombotic diseases. Unlike all
10 these previously described proteins, which act at various surface proteins on the platelet, the protein from this invention has a unique mechanism of action. It binds tightly to the collagen surface and one mechanism of action is given by the coverage of specific collagen sides no longer available for interactions and binding of platelets. This type of new mechanism has the great advantage, that
15 the platelets stay functionally intact during the application of the protein, so that very low or even no bleeding tendency is expected from this treatment.

Another important area of use is the treatment of various surfaces with the protein to render them non-adhesive for platelets and thereby create blood-compatible
20 devices.

As indicated above, the polypeptides according to the present invention are suitable as pharmaceutically effective compounds in pharmaceutical compositions and combinations.

25 The pharmaceutical formulations according to the invention optionally may comprise additional active ingredients like Aspirin, anti-coagulants such as hirudin or heparin or thrombolytic agents such as plasminogen activator or streptokinase.

The novel polypeptide according to the invention may form pharmaceutically
30 acceptable salts with any non-toxic, organic or inorganic acid. Inorganic acids are, for example, hydrochloric, hydrobromic, sulphuric or phosphoric acid and acid metal salts such as sodium monohydrogen orthophosphate and potassium hydrogen sulfate. Examples for organic acids are the mono, di and tri carboxylic

acids such as acetic, glycolic, lactic, pyruvic, malonic, succinic, glutaric, fumaric, malic, tartaric, citric, ascorbic, maleic, hydroxymaleic, benzoic, hydroxybenzoic, phenylacetic, cinnamic, salicylic and sulfonic acids such as methane sulfonic acid. Salts of the carboxy terminal amino acid moiety include the non-toxic
5 carboxylic acid salts formed with any suitable inorganic or organic bases. These salts include, for example, alkali metals such as sodium and potassium, alkaline earth metals such as calcium and magnesium, light metals of Group IIIA including aluminium, and organic primary, secondary and tertiary amines such as trialkylamines, including triethylamine, procaine, dibenzylamine, 1-ethenamine,
10 N,N'-dibenzylethylene-diamine, dihydroabietylamine and N-alkylpiperidine.

As used herein, the term "pharmaceutically acceptable carrier" means an inert, non toxic solid or liquid filler, diluent or encapsulating material, not reacting adversely with the active compound or with the patient. Suitable, preferably liquid
15 carriers are well known in the art such as sterile water, saline, aqueous dextrose, sugar solutions, ethanol, glycols and oils, including those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil and mineral oil.

The formulations according to the invention may be administered as unit doses
20 containing conventional non-toxic pharmaceutically acceptable carriers, diluents, adjuvants and vehicles which are typical for parenteral administration.

The term "parenteral" includes herein subcutaneous, intravenous, intra-articular and intratracheal injection and infusion techniques. Also other administrations
25 such as oral administration and topical application are suitable. Parenteral compositions and combinations are most preferably administered intravenously either in a bolus form or as a constant fusion according to known procedures.

Tablets and capsules for oral administration contain conventional excipients such
30 as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, and wetting agents. The tablets may be coated according to methods well known in the art.

Oral liquid preparations may be in the form of aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or another suitable vehicle before use. Such liquid preparations may contain conventional additives like suspending agents, emulsifying agents, non-aqueous vehicles and preservatives.

Topical applications may be in the form of aqueous or oily suspensions, solutions, emulsions, jellies or preferably emulsion ointments.

Unit doses according to the invention may contain daily required amounts of the protein according to the invention, or sub-multiples thereof to make up the desired dose. The optimum therapeutically acceptable dosage and dose rate for a given patient (mammals, including humans) depends on a variety of factors, such as the activity of the specific active material employed, the age, body weight, general health, sex, diet, time and route of administration, rate of clearance, the object of the treatment, i. e., therapy or prophylaxis and the nature of the thrombotic disease to be treated, antiplatelet or anticoagulant activity.

Therefore, in compositions and combinations useful as antithrombotic in a treated patient (in vivo) a pharmaceutical effective daily dose of the peptides of this invention is between about 0.01 and 100 mg/kg body weight, preferably between 0.1 and 10 mg/kg body weight. According to the application form one single dose may contain between 0.5 and 10 mg of the thrombin inhibitor. To achieve an anticoagulant effect in extracorporeal blood a pharmaceutically effective amount of the inventive peptides is between 0.2 and 150 mg /l, preferably between 1mg and 20 mg/l extracorporeal blood.

It is also object of this invention to provide an implantable or extracorporeal medical device for use in contact with body fluids in order to render the device surface substantially thromboresistant, coated with an immobilized polypeptide as defined above and in the claims. The polypeptide according to the invention is immobilized on a medical device so as to render the surface biocompatible and thromboresistant. Such devices sometimes have surface properties which typically induce platelet aggregation, which is a disadvantage in their intended uses in implantable and extracorporeal devices in contact with blood or other body

fluids. Examples for such devices which are commonly made from plastics materials and synthetic fibres are prostheses, artificial organs, ocular lenses, sutures, artificial vascular segments, catheters, dialysers, tubes and vessels carrying blood.

- 5 Posterior capsule opacification (PCO) is a common complication after cataract extraction, despite the modern surgical techniques and lenses which are used for this procedure. PCO is caused by the proliferation and migration of lens epithelial cells across the posterior capsule thus reducing the visual acuity. Physical treatments as well as chemically modified lenses have been proposed to reduce
10 formation of PCO. Heparin lens coating or topical heparin eyedrops have been used to reduce PCO, indicating that thrombogenic mechanisms are involved in the formation of PCO.

- Saratin has been shown to have significant advantages over heparin in preventing and blocking thrombogenicity. It is therefore another feature of this
15 invention to provide a coating comprising Saratin which reduces thrombogenicity of the lens material used for refractive anterior or posterior chamber ocular implants which may be surgically implanted into the eye. This new type of coating avoids problems contributed by stimulated cell growth. In combination with other medicaments which are for instance conferring cell death, Saratin coating helps to
20 completely overcome posterior capsule opacification.

Brief Description of the Figures

Details of the figures are explained in examples 1 to 10.

Fig. 1

5 Separation of saliva components. Elution of Saratin is marked by *

a) shows the separation profile of saliva after DEAE anion exchange. Saratin fractions have been collected from peak 3 (Example 2).

10 b) shows re-chromatography of pooled fractions on Mono Q HR5/5. Samples have been collected from the last part of the major peak as indicates by bar (Example 2).

15 c) shows last chromatography step on semi-preparative analytical RP-HPLC of Saratin positive fractions collected from Mono Q HR5/5 (example 2). Active Saratin was recovered from the major peak (peak 3).

Fig. 2

20 SDS-PAGE of fractions collected from RP-HPLC. Saratin positive fractions are marked * (Examples 2 and 3).

Fig. 3

E. coli expression vector for Saratin (Example 7).

Fig. 4

25 Baculo donor plasmid for Saratin (Example 8)

Fig. 5

30 Whole blood has been exposed to an artifical collagen surface and platelets adhesion has been visualized by staining. Saratin has been used as an inhibitor (Protein #607) Example 9

Fig. 6

Inhibition of platelet adhesion on collagen type III coated coverslips under she conditions. Comparison of saliva and saratin. Example 9

5

Fig. 7

Saratin exhibits dose dependent inhibition of platelet binding adhesion to collagen type III coated coverslips under she conditions. Example 9

10

Fig. 8

Yeast expression vector for Saratin (Example 13)

15

Detailed description of the invention**Example 1****20 Screening adhesion inhibitors**

Platelet adhesion to collagen has been the functional background for screening saliva components. In addition four additional tests available for the assessment of various actions of antithrombotic drugs such as the AZOCOLL assay, amidase
25 activity assay, vWillebrand dependent binding assay and platelet aggregation assay have been used to exclude functional properties which ideally would not be linked to an adhesion inhibitor. While most of these assays used here are standard assays, the platelet adhesion assay had to be modified to fit our specific needs. In short: Horm collagen (Nycomed) has been coated to 96 well plates (Nunc) by
30 using acidified collagen at a concentration of 20 µg /ml and incubating the plates overnight. After washing the plates 3 times with PBS the residual surface of the wells have been blocked with 1% BSA. Platelets isolated freshly from human citrorylated blood have been added simultaneously with fractions derived from

the individual column steps. When required pre-activation of platelets has been performed by incubating platelets prior to use in TBS in the presence of bivalent magnesium ions. Raw saliva, standardized to a total protein concentration of 200 µg/ml, has been used as a standard to compare for inhibitory activity. The platelet inhibition assay turned out to be highly susceptible to buffer changes and elevated salt concentrations. Since all the samples have been processed by ion-exchanger chromatography the direct testing of the fractions in the inhibition assay turned out to be complicated and unreliable. Therefore all samples to be tested have been applied to a Centricon^{*} based concentration step which simultaneously reduced ionic strength and supports concentration prior to measurement.

Example 2

15 Purification of a natural inhibitor

The invention used saliva collected from *H. medicinalis* which is known to contain a number of bioactive proteins such as hirudin, elastase inhibitors, collagenases and platelet aggregation inhibitors such as calin (Munro, R. et al.; Blood Coagulation and Fibrinolysis, 1991, 2, 179-184) and LAPP (Schaffer, L.W. et al.; Arterioscler. Thromb., 1993, 13, 1593-1601). Besides this characterized proteins the majority of the roughly eighty proteins detectable by SDS-PAGE are still unknown. In the present invention the fractionated saliva and the screening strategy as described in Example 1 was used in order to screen for the novel proteins that interferes directly with platelet collagen interaction.

25

Separation of an adhesion inhibitor activity from raw saliva turned out to be a critical task, mainly due to the irreversible loss of most of the adhesion inhibitory activity in the first chromatographic step. Additives such as 12% ethanol and divalent cations as recommended by Munro, R. et al did not improve the situation. However the high salt concentration of the saliva in combination with the low overall protein concentration (190-250 µg/ml) asked for an initial concentration or buffer exchange step. Thus several strategies for enrichment, concentration or buffer exchange were explored. Traditional dialysis led to the complete loss of

30
*Trade-mark

activity. Most of the other standard techniques such as ion exchangers, affinity columns, size exclusion (loss was irrespective of the column resin) failed independent of the nature of the separation technology or buffer and additives used. Surprisingly it turned out, that pressurized dialysis of 500ml saliva was a successful method to concentrate the saliva proteins (about 30-40 times) and simultaneously get rid of the unwanted buffer components. Unexpectedly the saliva raw material processed in this way was an ideal starting material for further purification and the recovery of bioactivity anti-adhesive components out of the saliva was no longer a real problem. Since cation exchangers or affinity columns turned out to be an insufficient step in purification, weak anion exchanger such as DEAE-Fastflow^{*} or EMD-DEAE-Fractogel^{*} were used. 12% ethanol and divalent cations have been tested, however the results were similar with or without this additives. Further optimization of the chromatographic steps led to a sequence of DEAE-column, Mono Q-column and as a final step an reversed phase RP18 column. Optimization of the chromatographic conditions have been performed on a BiaCore^{*} chromatographic system using analytical columns available from Pharmacia. The gradients used to operate the DEAE-column, Mono Q-column^{*} and RP18 columns are given in Figures 1a, b, c. The BiaCore based separation has been scaled up by using FPLC techniques. The optimized running conditions have been directly converted into the a semi-preparative scale of separation by following the instruction of the manufacturer with the exception of the RP18 column step which was maintained with Biacore technique in order to minimize loss of purified material. Recovery of purified protein from the last chromatographic RP-step was done by speed vacuum centrifugation. Samples collected from the last RP-step have been collected and analysed by SDS-PAGE (Figure 2) Subsequently samples have been resuspended in PBS and used to perform analytical as well as functional tests. Typically a yield of about 750 µg/l Saratin was recovered from unprocessed saliva.

*Trade-mark

Example 3

Biochemical characterization

5 Purification of Saratin as disclosed in Example 1 led to an essentially pure protein with an apparent molecular weight around 21 kDa shown under reducing conditions in SDS-PAGE (Figure 2). The complete amino acid sequence was obtained by direct sequencing of the first 48 amino acids of the purified protein and was completed by the sequencing of several internal peptides generated via
10 enzymatic degradation. The complete protein sequence is disclosed in SEQ. ID. NO. 2

The protein is composed of 103 amino acids with a calculated molecular weight of 12067,9 and an actual molecular weight of 12061,9 as deduced by ESI-Mass spectrometry. This difference in theoretical and measured molecular weight
15 indicates that all the six cysteines identified in Saratin are involved in the formation of S-S bridges. This finding is supported by the observation of a strong change in mobility of the protein when chromatographed in SDS-PAGE under reducing or non-reducing conditions. Furthermore, the protein is rich in acidic acids such as Glu and Asp. Isoelectric focussing using IEF-PAGE (Immobiline) technique
20 revealed an isoelectric point of $pH\ 3.7 \pm 0.5$. In a comparative study we have used the purified leech protein as a reference and compared it with the physicochemical properties of recombinant Saratin derived from baculo, yeast and E. coli expression. All three proteins turned out to be identical in their properties.

Characterization of protein by SDS-PAGE visualized by Coomassie (Figure 2) or
25 silver-staining and/or Western blot analysis revealed that the protein was homogeneous and in a non-glycosylated form.

Example 4

30 mRNA Preparation and cDNA Synthesis

RNA was prepared from the medicinal leech, *Hirudo medicinalis*, using guanidinium thiocyanate method. mRNA was purified from total RNA using "Oligotex mRNA kit" (QIAGEN).

26474-568

- 13 -

cDNA was synthesized using "Marathon^{*} cDNA Amplification kit" (CLONTECH). The DNA sequence encoding *Saratin* was initially amplified using PCR oligonucleotide primers according to the instruction of the supplier. After cDNA synthesis, a universal adaptor was ligated to both ends of the cDNA. The sequences of the universal adaptor and the universal primers AP1 or AP2 was chosen according to the instructions of the supplier of said kit.

Example 5

10 Amplification and isolation of the *Saratin* gene by PCR

Degenerate primers have been synthesised based on the immediate N-terminus of the reverse translated *Saratin* amino acid sequence.

These primer01 and primer02 have been designed to hybridize specifically to the 5'-end of the *Saratin* cDNA. The primer design was based on the reverse translated amino acid sequence of the experimentally determined eight N-terminal amino acids of the purified *Saratin* protein. The two primers primer01 and primer02 were synthesized to keep the degeneracy as low as possible and to give the primers at the critical 3'-end a stretch of 8 basepairs of perfect match with the *Saratin* cDNA sequence in order to ensure efficient and specific amplification of the template cDNA. According to the DNA degenerate alphabet (IUPAC code) R = A or G; M = A or C; Y = C or T; and N = A or G or C or T.

A 3'-RACE PCR reaction was carried out using a mixture of the primer01 and primer02 and one universal primer AP1 or AP2. PCR products were cloned in TA cloning vector pCR2.1 or pCR^{*} Script SK(+) vector and sequenced. After sequencing several of the 3'-RACE PCR fragments obtained, the *Saratin* gene sequence was obtained, with the exception of the 5'-untranslated region, the signal peptide sequence and the sequence coding for the first 8 amino acids from the N-terminus of the mature protein. In order to obtain this missing *Saratin* cDNA sequence information, a 5'-RACE experiment was performed using a gene specific primer from the middle of the *Saratin* cDNA and one of the universal primers AP1 or AP2. After sequencing several of the resulting 5'-RACE PCR fragments, the complete *Saratin* gene sequence was obtained. Amplifying the

*Trade-mark

Saratin gene by PCR using gene specific, non-degenerate primers from both the 3'-end and the 5'-end of the Saratin gene, a full length of Saratin gene was obtained.

DNA Sequencing was performed with more than 15 different PCR clones. However only one significant change which caused amino acid sequence change in only one clone was found. It is very likely that this change was caused by PCR. Another five silent changes which do not cause amino acid sequence change were found. Thus it is unlikely that these changes were caused by PCR, because the same changes were found in different clones.

- 10 The Saratin gene ORF is 372bp and contains a 21 amino acid signal sequence and a 103 amino acid sequence coding for the mature protein. The amino acid sequence deduced from the PCR clone was found to be identical to the sequence obtained by sequencing of the natural saliva derived protein.

15 **Example 6**

Expression in COS cells and the detection of expressed protein

- 20 In order to express Saratin gene in mammalian cells such as COS or CHO, Saratin gene was cut from the vector pCR Script SK(+) using XhoI+XbaI and cloned into mammalian cell expression vector pCI-neo (Promega). pCI-neo was chosen because it contains T7 and T3 promoter for in vitro expression and neomycin resistant gene for G418 selection, and can be used in both COS and CHO cells.

- 25 In addition to the signal sequence and mature protein sequence, the insert contains a Koz. sequence at the 5'-end for efficient translation and the his-tag MRGS(H)₆ at the 3'-end (C-Terminal) for detection of protein expression, purification and concentration. The plasmid constructure was named pNC-31.

- 30 pNC-31 plasmid DNA was used for the transfection of COS cells. The COS cells growing at log phase were washed twice with PBS and dissolved in PBS at a concentration of 1×10^7 /ml. Then 12µg plasmid DNA (less than 50µl in H₂O or TE buffer) was added in 0.7-0.8 ml COS cells suspension and mixed in a

electroporation cuvette. Electroporation was performed at 1.9kv, 25 μ FD for 10 min and transferred to a 90mm plate. After adding 8 ml DMEM medium containing 10% FCS and antibiotics the cells were grown for three days. The supernatantes and the cells have been used for further isolation of protein and detection. The
5 expressed protein was detected by western blotting method using anti-MRGS(H)6 antibody. Purification was performed using chelators such as NTA or imido acetic acid immobilized on a column matrix and modified with metall ions such as Co, Ni, or Cu.

10 Example 7

Construction of the *E. coli* expression vector and expression

Because of the variable codon usage in different biological systems, some
15 codons are used very rarely in *E. coli*. In order to enable expression an optimized version in *E. coli* the gene had to be converted to *E. coli* codon usage according to standard procedures.

The expression in *E. coli* was performed using a modified version of the plasmid
20 pASK75, which carries the tet promoter region. (Skerra, A., et al. Gene, 1994, 151, 131 - 135). In short: The modification was made by cloning a new linker between the XbaI and the Hind III sites (Figure 3). The new linker contains the ompA leader sequence, another multiple cloning site and a 6xHis-tag instead of the strep-tag.

25

To construct the expression vector for Saratin it was necessary to introduce 5' Cla I and 3' Eco47III restrictionsites by PCR method.

Therefore the 5'-primer03 and the 3' primer04 have used. The PCR product first was cloned into the PCR II vector system (Invitrogen) and sequenced.

30 In a second step the Saratin gene was cloned into the modified pASK75 vector using the rectrictionsites 5'ClaI and 3' HindIII.

After expressing and proving the activity of this recombinant Saratin in a second PCR reaction the His-tag was removed and the start codon of the Saratin gene

26474-568

- 21 -

was directly fused to the omp A leader sequence. The 5'-primer05 and the 3'-primer06 for this PCR reaction are given in the sequence listing.

As an example for the expression in *E. coli* the expression vector pRG72 (Figure 3), which contains the structural gene of Saratin fused to the ompA leader sequence, was transformed into W3110 competent cells. Cells have been induced after they have been grown to mid-log phase. 1 hour thereafter the recombinant Sarastatin could be clearly detected.

10

Example 8

Construction of the Baculo Donor Plasmid and expression

15 For expression of Saratin in the Baculo virus expression system the Bac-To-BacTM Baculovirus Expression System from Gibco Life Technologies was used. To get a selection system the Honeybee melitin leader sequence was fused to the Saratin gene and to introduce the restriction sites 5' BamHI and 3' KpnI one single PCR reaction was carried out using the 5'-primer07 and the 3'-primer08.

20 The corresponding PCR product was cloned into the PCR II Vector (Invitrogen) and sequenced. Then the Melitin – Saratin fusion was cloned into the pFastBac^{*} vector using the restriction sites 5'BamHI and 3'KpnI resulting in pTD13 (Figure 4). Generation of recombinant baculoviruses and Saratin expression was performed with the Bac-To-Bac^{*} Expression System. The donor plasmid pTD13

25 was transformed into DH10Bac competent cells which contain the bacmid with a mini-attTn7 target site and the helper plasmid. The mini-Tn7 element on the donor plasmid ca transpose to the a mini-attTn7 target site on the bacmid in the presence of transposition proteins provided by the helper plasmid. Colonies containing recombinant bacmids were identified by disruption of the *lacZ* gene.

30 High molecular weight mini-prep DNA is prepared from selected *E. coli* clones containing the recombinant bacmid, and this DNA was then used to transfect insect cells. Detailed descriptions are included in the instruction manual of the expression kit.

*Trade-mark

Example 9

5 Platelet Adhesion to Collagen Under Flow Conditions (Dynamic assay)

In the platelet adhesion assay, whole human blood is perfused through a parallel flow chamber to examine the adhesive activity of platelets to a collagen coated coverslips under high shear flow (simulating in vivo arterial conditions) as
10 originally described by Sakariassen et al (Meth. Enz., 1988, 169, 37-70). Human placental collagen, type III (Sigma), solubilized in 50 mmol/L acetic acid was sprayed onto cleaned glass coverslips (18mm x 18mm) with a retouching airbrush. Coverslips were stored in PBS at 4°C overnight.

Fresh whole human blood (anticoagulated with low molecular weight heparin; 20
15 U/ml) was prewarmed at 37°C for 10 min before being used. Preparations of proteins according to this invention were pipetted onto coverslips (30 µl per coverslip) and incubated for 10 min in a humid chamber at room temperature before being inserted into the perfusion chamber. The blood was allowed to perfuse through the chamber (at 37°C for 5 min at a shear rate of
20 1300 s^{-1} .)

Subsequently, the coverslips are removed, washed in PBS and fixed in 0.25% gluteraldehyde for 30 min., and thereafter stained with May-Grünwald Giemsa. Figure 9 shows a typical expamle. Extensive coverage with stained platelets is seen in the untreated control surface. A comparable surface pretreated with
25 Saratin shows dramatically deminished (by 80%) platelet binding. Platelet adhesion was quantified with a light microscope as shown in Figure 5 (magnification x 1000) coupled to a computerized image analyzer (Leica). Results were expressed as the percentage of the surface covered with platelets and platelet aggregates.

30 Comparison of saliva inhibitory activity has been compared to purified protein as demonstrated in Figure 6. Platelet adhesion on collagen type III coated coverslips at a shear rate of 1300 s^{-1} with crude saliva (#616) produced an inhibition of about 48% compared to control. Purified protein (#607; Saratin)

demonstrated a reduction in platelet deposition of about 81% at standardized protein concentrations. The inhibition induced by Saratin increases in a dose-dependent manner with higher concentrations of purified protein as demonstrated in Figure 7.

5

Example 10

Immunization and antibodies

10 With the first lot of highly purified natural protein available, immunization of animals has been started right away. Immune sera were raised in rabbits and high titered reagents were available for further screening. Additional antisera became available when the peptide sequence of the complete protein was available and three synthetic peptides have been synthesized (amino-acid
15 sequence 83-103, 13-30, 58-69) coupled to KLH with a standard linker procedure and used for immunization. Three sera directed to an N-terminal peptide segment and two sera specific for C-terminal peptides have established. With the high titered immune sera available it has been possible to establish and use Elisa technology to monitor and quantitate purification of naturally purified as well as
20 recombinant protein. Thus the time consuming and rather work intensive platelet-inhibition assay could be replaced and was applied only to confirm the inhibitory potential of finally purified protein.

Example 11

25

Immuno-Assays for estimation of Saratin binding

Acidified Horm-collagen (Nycomed) has been used for the coating of 96-well micro-titer plates (Nunc) 50 μ l of the collagen solution (20 μ g/ml) has been used
30 to coat plates over night. Prior to testing the plates have been washed three times with PBS and have been incubated with a BSA solution (1%) in order to prevent non-specific adhesion. 50 μ l Saratin has been added in serial delution and have been incubated for one hour. Plates were washed three times prior to the

application of anti-Saratin antibody for detection. After an additional one our incubation step surplus antibody has been removed and a second biotin-labeled Antibody has been used for detection. Read-out has been performed via streptavidin-POD catalyzed colour reaction with substrates such as ODB-tablets
5 (Dako) measured at 490 nm.

Example 12

Competition assay for screening inhibitors

10

Recombinant tagged Saratin (His-tag) prepared as described in Example 7 has been compared with native untagged Saratin for collagen binding. Plates coated with acidified Horm-collagen (Nycomed) have been prepared as described in Example 11. Detection was performed with rabbit anti-Saratin antibodies. The
15 tagged and untagged Saratins showed identical binding properties. Alternatively the untagged Saratin version has been modified by biotinylation (Pierce, biotinylation kit) and compared to unmodified Saratin. Binding properties to collagen have been identical. Furthermore experiments have been performed using the biotinylated Saratin to cross-compete with unmodified Saratin, peptides,
20 saliva derived Saratin, complete saliva or antibodies directed to Saratin. Binding of the various „competitors“ to collagen has been tested by estimating the binding of biotinylated Saratin using a streptavidine-POD conjugate and ODB-substrate reaction for detection. This assay comprising biotinylated Saratin was typically used for the estimation of Saratin concentration in saliva (750µg/l), epitope
25 mapping of Saratin directed antibodies, evaluation of bioactive Saratin, mutated Saratin. In order to explore the potential of the assay blocking as well as none blocking anti-Saratin antibodies raised with specific Saratin peptides have been used.

Example 13

Yeast expression vector and expression

5 The pichia multi copy expression system (Invitrogen) has been used as a typical example for yeast expression. The construction of the yeast expression vector is shown in Figure 8. For generation of the expression vector for Saratin the PCR amplification method has been used to generate restriction ends (5' EcoR I, 3'Not I) compatible with the ligation into the appropriate vector (pPIC9K). The
10 following 5'-primer09 and the 3'-primer10.

Before transforming the *Pichia* spheroplasts the expression vector has been linearized with Sal I. Colonies have been screened for His⁺ Mut⁺-positiv mutants to ensure integration of the Saratin gene. Typical growing conditions have been: 28–30°C, up to an optical density of OD 2-6. Induction of expression was
15 performed after resuspending centrifuged cells in medium and addition of methanol up to a final concentration of 0,5% and maintaining this condition for an extented time periode which typically would be 24 hours. After a typical fermentation periode of 6 days the product was recovered from the supernatant and analysed by SDS-PAGE and ELISA.

WO 00/56885

PCT/EP00/02117

1

SEQUENCE LISTING

<110> Merck Patent GmbH

5 <120> Protein for blocking platelet adhesion

<130> Saratin Sequence

<140>

10 <141>

<160> 12

<170> PatentIn Ver. 2.1

15

<210> 1

<211> 375

<212> DNA

<213> Hirudo medicinalis

20

<220>

<221> CDS

<222> (64)..(372)

25 <400> 1

atgaagtatt tcttgatttc cttcctttgc ctcgcaagct tgctgatctc aactacttct 60

tca	gaa	gaa	cgt	gaa	gat	tgt	tgg	acg	ttt	tac	gcg	aac	aga	aaa	tat	108
	Glu	Glu	Arg	Glu	Asp	Cys	Trp	Thr	Phe	Tyr	Ala	Asn	Arg	Lys	Tyr	
30	1				5				10					15		

aca	gac	ttc	gat	aaa	tct	ttt	aag	aag	tcc	tct	gat	ctt	gac	gaa	tgc	156
Thr	Asp	Phe	Asp	Lys	Ser	Phe	Lys	Lys	Ser	Ser	Asp	Leu	Asp	Glu	Cys	
				20				25						30		

35

aaa	aaa	aca	tgt	ttc	aag	acg	gag	tac	tgc	tac	atc	gtt	ttt	gaa	gac	204
Lys	Lys	Thr	Cys	Phe	Lys	Thr	Glu	Tyr	Cys	Tyr	Ile	Val	Phe	Glu	Asp	
			35				40						45			

40

acg	gtc	aac	aag	gaa	tgt	tac	tac	aat	gtc	gtt	gat	ggc	gaa	gag	tta	252
Thr	Val	Asn	Lys	Glu	Cys	Tyr	Tyr	Asn	Val	Val	Asp	Gly	Glu	Glu	Leu	
		50				55					60					

gac	caa	gaa	aaa	ttt	gtt	gtc	gac	gaa	aac	ttc	acg	gaa	aat	tat	ttg	300
Asp	Gln	Glu	Lys	Phe	Val	Val	Asp	Glu	Asn	Phe	Thr	Glu	Asn	Tyr	Leu	
	65				70				75							

aca	gac	tgc	gag	ggc	aaa	gat	gca	ggc	aat	gag	gca	ggc	aca	ggc	gac	348
Thr	Asp	Cys	Glu	Gly	Lys	Asp	Ala	Gly	Asn	Ala	Ala	Gly	Thr	Gly	Asp	
50	80				85			90						95		

gag	tca	gat	gaa	gtt	gat	gaa	gat	taa	375
Glu	Ser	Asp	Glu	Val	Asp	Glu	Asp		
				100					

55

<210> 2

<211> 103

<212> PRT

60 <213> Hirudo medicinalis

WO 00/56885

PCT/EP00/02117

2

<400> 2
 Glu Glu Arg Glu Asp Cys Trp Thr Phe Tyr Ala Asn Arg Lys Tyr Thr
 1 5 10 15
 5 Asp Phe Asp Lys Ser Phe Lys Lys Ser Ser Asp Leu Asp Glu Cys Lys
 20 25 30
 Lys Thr Cys Phe Lys Thr Glu Tyr Cys Tyr Ile Val Phe Glu Asp Thr
 35 40 45
 10 Val Asn Lys Glu Cys Tyr Tyr Asn Val Val Asp Gly Glu Glu Leu Asp
 50 55 60
 Gln Glu Lys Phe Val Val Asp Glu Asn Phe Thr Glu Asn Tyr Leu Thr
 15 65 70 75 80
 Asp Cys Glu Gly Lys Asp Ala Gly Asn Ala Ala Gly Thr Gly Asp Glu
 85 90 95
 20 Ser Asp Glu Val Asp Glu Asp
 100
 25 <210> 3
 <211> 23
 <212> DNA
 <213> Artificial Sequence
 30 <220>
 <223> Description of Artificial Sequence:degenerate
 primer01
 <400> 3
 35 gargarmgng argaytggtg gac 23
 <210> 4
 <211> 23
 40 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence:degenerate
 45 primer02
 <400> 4
 gargarmgng argaytgctg gac 23
 50 <210> 5
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 55 <220>
 <223> Description of Artificial Sequence: primer03
 <400> 5
 60 gcatcgatgg aagaacgtga agac 24

WO 00/56885

PCT/EP00/02117

3

<210> 6
 <211> 23
 <212> DNA
 5 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: primer04

 10 <400> 6 23
 tagcgctttt gacgtcgtcg tca

 <210> 7
 15 <211> 26
 <212> DNA
 <213> Artificial Sequence

 <220>
 20 <223> Description of Artificial Sequence: primer05

 <400> 7 26
 gaagaatgca aggatgagga ttattg

 25
 <210> 8
 <211> 25
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> Description of Artificial Sequence: primer06

 <400> 8 25
 35 aagcttctag tcttcgtcaa cttcg

 <210> 9
 <211> 94
 40 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: primer07
 45
 <400> 9
 cggatccatg aaattcttag tcaacgttgc ccttgttttt atggtcgtat acatttctta 60
 catctatgcg gaagaacgtg aagattgttg gact 94

 50
 <210> 10
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 55
 <220>
 <223> Description of Artificial Sequence: primer08

 <400> 10 24
 60 ggtacctcac atatcttcat caac

WO 00/56885

PCT/EP00/02117

4

<210> 11
<211> 30
<212> DNA
5 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer09

10 <400> 11
gcatgcggcc gcctaattctt catcaacttc 30

<210> 12
15 <211> 27
<212> DNA
<213> Artificial Sequence

<220>
20 <223> Description of Artificial Sequence: primer10

<400> 12
gcatgaattc gaagaacgtg aagattg 27

25

26474-568

26

CLAIMS:

1. A polypeptide isolated from *H. medicinalis* having a molecular weight of about 12,000 $\pm$ 1 kD with the biological activity of an inhibitor of collagen-dependent platelet
5 adhesion, which is at least 80% identical to the amino acid sequence of SEQ ID NO:2 over its entire length.
2. The polypeptide of claim 1 having an isoelectric point of pH 3.7 $\pm$ 0.5.
3. The polypeptide of claim 1 or 2 having six
10 cysteine molecules capable of forming -S-S-bridges.
4. The polypeptide of any one of claims 1 to 3, named Saratin, consisting of the amino acid sequence of SEQ ID NO:2.
5. An isolated polynucleotide encoding the
15 polypeptide of claim 4.
6. An isolated polynucleotide comprising a DNA sequence that is at least 80% identical to SEQ ID NO:1 over its entire length, wherein said polynucleotide encodes the polypeptide of claim 1.
- 20 7. The isolated polynucleotide of claim 6 consisting of the nucleotide sequence of SEQ ID NO:1.
8. An expression vector comprising the DNA sequence of claim 6.
9. A host cell comprising the expression vector of
25 claim 8.
10. An expression system comprising the host cell of claim 9.

26474-568

27

11. A process for producing a polypeptide comprising culturing the host cell of claim 9 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture supernatant or
5 cell residue.
12. An antibody immunospecific for the polypeptide of claim 1.
13. A pharmaceutical composition comprising the polypeptide of claim 1, together with a pharmaceutically
10 acceptable carrier or excipient.
14. The pharmaceutical composition of claim 13 further comprising an anti-coagulant agent or a thrombolytic agent.
15. A medicament for the treatment of a thromboembolic disease comprising the polypeptide of claim 1 and a
15 pharmaceutically acceptable carrier.
16. A use of the polypeptide of claim 1 for ex vivo coating of artificial surfaces.
17. A use of the polypeptide of claim 1 for ex vivo modifying intraocular lenses in order to lessen the
20 thrombogenicity of the lens material.
18. A use of the antibody of claim 12 to identify ex vivo the polypeptide of claim 1.
19. A method for determining whether a compound competes with the binding of the polypeptide of claim 1 to
25 collagen, the method comprising applying said polypeptide to collagen in the absence and presence of said compound and determining the binding rate of said polypeptide in the absence of said competing compound compared to the respective binding rate in the presence of said compound.

26474-568

28

20. The method of claim 19, wherein said polypeptide is the polypeptide of claim 4.

21. The method of claim 19, wherein said competitor is an antibody.


5 22. The method of claim 19, wherein said competitor is an anti-Saratin antibody.

FETHERSTONHAUGH & CO.


OTTAWA, CANADA

PATENT AGENTS

1 / 9




2 / 9



WO 00/56885


PCT/EP00/02117

3 / 9





4/9


Fig. 2



5 / 9

**FIG. 4**

6 / 9



Shear rate: 1300 s^{-1}

Human Blood


Collagen Type III from Human Placenta

$30 \mu\text{l}/\text{coverslip}$

FIG. 5

7 / 9

FIG. 6




Collagen Type III from Human Placenta

Saliva # 616 0,11 mg/ml: 30 μ l on coverslip n = 4

Protein # 607 0,11 mg/ml: 30 μ l on coverslip n = 2

FIG. 7



Protein # 607

Collagen Type III from Human Placenta

9 / 9




FIG. 8