




- (51) **International Patent Classification:**
C07K 16/00 (2006.01) A61K 38/17 (2006.01)
A61K 38/00 (2006.01)
- (21) **International Application Number:**
PCT/US2014/072087
- (22) **International Filing Date:**
23 December 2014 (23.12.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/920,547 24 December 2013 (24.12.2013) US
- (71) **Applicants:** ARGEN-X N.V. [NL/NL]; Willemstraat 5, NL-4811 AH Breda (NL). **THE BOARD OF REGENTS OF THE UNIVERSITY OF TEXAS SYSTEM** [US/US]; 201 West 7th Street, Austin, TX 78701 (US).
- (72) **Inventors:** ULRICHTS, Peter; C/o ARGEN-X B.V., Willemstraat 5, NL-4811 AH Breda (NL). BLANCHETOT, Christophe; C/o ARGEN-X B.V., Willemstraat 5, NL-4811 AH Breda (NL). DREIER, Torsten; C/o ARGEN-X B.V., Willemstraat 5, NL-4811 AH Breda (NL). DE HAARD, Johannes; C/o ARGEN-X B.V., Willemstraat 5, NL-4811 AH Breda (NL). WARD OBER, E., Sally; C/o The Board of Regents of the University of Texas, 201 West 7th Street, Austin, TX 78701 (US). ONGENAE, Nicolas, G.H.; C/o ARGEN-X B.V., Willemstraat 5, NL-4811 AH Breda (NL).
- (74) **Agents:** VELEMA, James, H. et al.; Lathrop & Gage LLP, 28 State Street, Boston, MA 02109 (US).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:**
- with international search report (Art. 21(3))
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: FCRN ANTAGONISTS AND METHODS OF USE

FIG.1



(57) **Abstract:** Provided are novel FcRn antagonist compositions comprising a variant Fc region that binds specifically to FcRn with increased affinity and reduced pH dependence relative to the native Fc region. Also provided are FcRn antagonists with enhanced CD 16 binding affinity. Also provided are methods of treating antibody-mediated disorders (e.g. autoimmune diseases) using the these FcRn antagonist compositions, nucleic acids encoding the FcRn antagonist compositions, recombinant expression vectors and host cells for making the FcRn antagonist compositions, and pharmaceutical compositions comprising the FcRn antagonist compositions.

FCRN ANTAGONISTS AND METHODS OF USE

RELATED APPLICATIONS

This application claims priority to U.S. provisional application 61/920,547, filed
5 December 24, 2013, which is hereby incorporated by reference in its entirety.

BACKGROUND

Immunoglobulin gamma (IgG) antibodies play a key role in the pathology of many disorders, such as autoimmune diseases, inflammatory diseases, and disorders in which the
10 pathology is characterized by over-expression of IgG antibodies (e.g., hypergammaglobulinemia) (see e.g. Junghans, Immunologic Research 16 (1):29 (1997)).

The half-life of IgG in the serum is prolonged relative to the serum half-life of other plasma proteins (Roopenian et al., J. Immunology 170:3528 (2003); Junghans and Anderson, Proc. Natl. Acad. Sci. USA 93:5512 (1996)). This long half-life is due, in part, to the binding
15 of the Fc region of IgG to the Fc receptor, FcRn. Although FcRn was originally characterized as a neonatal transport receptor for maternal IgG, it also functions in adults to protect IgG from degradation. FcRn binds to pinocytosed IgG and protects the IgG from transport to degradative lysosomes by recycling it back to the extracellular compartment. This recycling is facilitated by the pH dependent binding of IgG to FcRn, where the
20 IgG/FcRn interaction is stronger at acidic endosomal pH than at extracellular physiological pH.

When the serum concentration of IgG reaches a level that exceeds available FcRn molecules, unbound IgG is not protected from degradative mechanisms and will consequently have a reduced serum half-life. Thus, inhibition of IgG binding to FcRn reduces the serum
25 half-life of IgG by preventing IgG endosomal recycling of IgG. Accordingly, agents that antagonize the binding of IgG to FcRn may be useful for regulating, treating or preventing antibody-mediated disorders, such as autoimmune diseases, inflammatory diseases, etc. One example of a method of antagonizing IgG Fc binding to FcRn involves the generation of blocking antibodies to FcRn (see e.g. WO2002/43658). Peptides have also been identified
30 that bind to and antagonize FcRn function (see e.g. US6,212,022 and US8,101,186). In addition, full-length IgG antibodies comprising variant Fc receptors with enhanced FcRn binding and decreased pH dependence have also been identified that antagonize FcRn binding to IgG (see e.g. 8,163,881). However, there is a need in the art for improved agents that antagonize FcRn binding to IgG for use in the treatment of antibody-mediated disorders.

SUMMARY

The present disclosure provides novel FcRn antagonist compositions. These compositions generally comprise a variant Fc region, or FcRn-binding fragment thereof, that binds specifically to FcRn with increased affinity and reduced pH dependence relative to the native Fc region. The invention is based, in part, on the surprising finding that an isolated variant Fc region (e.g., a variant Fc region comprising the amino acids Y, T, E, K, F, and Y at EU positions (EU numbering) 252, 254, 256, 433, 434, and 436 respectively) is a more efficacious FcRn antagonist *in vivo* than a full-length antibody comprising that variant Fc region. The FcRn antagonist compositions of the present disclosure are particularly useful for reducing the serum levels of Fc-containing agents (e.g., antibodies and immunoadhesins). Accordingly, the instant disclosure also provides methods of treating antibody-mediated disorders (e.g. autoimmune diseases) using the FcRn antagonist compositions disclosed herein. Also provided are nucleic acids encoding the FcRn antagonist compositions, recombinant expression vectors and host cells for making the FcRn antagonist compositions, and pharmaceutical compositions comprising the FcRn antagonist compositions.

The FcRn antagonists disclosed herein are particularly advantageous over previously described FcRn antagonist compositions and known treatments for antibody-mediated disorders. For example, the FcRn antagonists disclosed herein are smaller and more potent than intravenous gamma globulin (IVIG), the current treatment for many antibody-mediated disorders. Accordingly, the effective dose of the disclosed FcRn antagonists can be far less than that of IVIG. Moreover, IVIG is isolated and purified from human donors and, as a consequence, suffers from considerable batch-batch variation. The FcRn antagonists compositions disclosed herein can be recombinantly produced or chemically synthesized and, therefore, are far more homogeneous. As demonstrated herein, the FcRn antagonists disclosed herein are also surprisingly more efficacious than full-length IgG antibodies comprising variant Fc receptors, such as set forth in Vaccaro et al., Nature Biotech 23(9) 1283-1288 (1997).

Accordingly, in one aspect, the instant disclosure provides an isolated FcRn antagonist comprising a variant Fc region or FcRn-binding fragment thereof, wherein the Fc region or fragment comprises the amino acids Y, T, E, K, F, and Y at EU positions 252, 254, 256, 433, 434, and 436 respectively, and wherein the FcRn antagonist is not a full-length antibody.

In certain embodiments, the FcRn antagonist does not comprise an antibody variable region or a CH1 domain. In certain embodiments, the FcRn antagonist does not comprise a free cysteine residue. In certain embodiments, the Fc region is an IgG Fc region (e.g., a human IgG Fc region). In certain embodiments, the Fc region is an IgG1 Fc region (e.g., a human IgG1 Fc region). In certain embodiments, the Fc region is a chimeric Fc region.

In certain embodiments, the FcRn antagonist comprises the variant Fc region amino acid sequence set forth in SEQ ID NO:1. In certain embodiments, the FcRn antagonist comprises a variant Fc region wherein the amino acid sequence of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO: 1, 2, or 3. In certain embodiments, the FcRn antagonist consists of a variant Fc region wherein the amino acid sequence of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO: 2.

In certain embodiments, the FcRn antagonist comprises a variant Fc region that has altered (increased or decreased) affinity for an Fc receptor relative to the affinity of a wild-type IgG1 Fc region for the Fc gamma receptor. In certain embodiments, the variant Fc has increased affinity for CD16a.

In certain embodiments, the FcRn antagonist comprises a variant Fc region that does not comprise an N-linked glycan at EU position 297. In certain embodiments, the FcRn antagonist comprises a variant Fc region that comprises an afucosylated N-linked glycan at EU position 297. In certain embodiments, the FcRn antagonist comprises a variant Fc region that comprises an N-linked glycan having a bisecting GlcNac at EU position 297.

In certain embodiments, the FcRn antagonist comprises a variant Fc region linked to a half-life extender. In certain embodiments, the half-life extender is polyethylene glycol or human serum albumin. In certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules disclosed herein, wherein at least 50% (optionally, at least 60, 70, 80, 90, 95, or 99%) of the molecules comprise a variant Fc region, or FcRn-binding fragment thereof, having an afucosylated N-linked glycan. In certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules disclosed herein, wherein at least 50% (optionally, at least 60, 70, 80, 90, 95, or 99%) of the molecules comprise a variant Fc region, or FcRn-binding fragment thereof, comprising an N-linked glycan having a bisecting GlcNac.

In certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules as disclosed herein,

wherein greater than 95% the of the FcRn antagonist molecules in the composition are monomers (e.g., greater than 95, 96, 97, 98, 99, 99.1, 99.2, 99.3, 99.4, 99.5, 99.6, 99.7, 99.8, 99.9 %).

In certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules disclosed herein, wherein less than 5% the of the FcRn antagonist molecules in the composition are present in aggregates, (e.g., less than 5, 4, 3, 2, 1, 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.2, 0.1 %).

In certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules disclosed herein, wherein the composition is substantially free of FcRn antagonist molecule degradation products.

In another aspect, the instant disclosure provides pharmaceutical compositions comprising an FcRn antagonist or FcRn antagonist composition disclosed herein and a pharmaceutically acceptable carrier or excipient.

In another aspect, the instant disclosure provides a method of inhibiting FcRn function in a subject, the method comprising administering to the subject an effective amount of an FcRn-antagonist composition disclosed herein.

In another aspect, the instant disclosure provides a method of reducing the serum levels of an Fc-containing agent in subject that has been administered the Fc-containing agent, the method comprising administering to subject an effective amount of an FcRn-antagonist composition disclosed herein. In certain embodiments, the Fc-containing agent is an antibody or immunoadhesin. In certain embodiments, the Fc-containing agent is a therapeutic or diagnostic agent. In certain embodiments the Fc-containing agent is an imaging agent. In certain embodiments, the Fc-containing agent is an antibody drug conjugate.

In another aspect, the instant disclosure provides a method of treating an antibody-mediated disorder in a subject, the method comprising administering to the subject an effective amount of an FcRn-antagonist composition disclosed herein. In certain embodiments, the antibody-mediated disorder is hyperglobulinemia. In certain embodiments, the antibody-mediated disorder is a disease or disorder that is treatable using intravenous immunoglobulin (IVIG). In certain embodiments, the antibody-mediated disorder is a disease or disorder that is treatable using plasmapheresis and/or immunoadsorption.

In certain embodiments, the antibody-mediated disorder is an autoimmune disease. In certain embodiments, the autoimmune disease is selected from the group consisting of allogenic islet graft rejection, alopecia areata, ankylosing spondylitis, antiphospholipid

syndrome, autoimmune Addison's disease, Alzheimer's disease, antineutrophil cytoplasmic autoantibodies (ANCA), autoimmune diseases of the adrenal gland, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune myocarditis, autoimmune neutropenia, autoimmune oophoritis and orchitis, autoimmune thrombocytopenia, autoimmune urticaria,

5 Behcet's disease, bullous pemphigoid, cardiomyopathy, Castleman's syndrome, celiac spruce-dermatitis, chronic fatigue immune dysfunction syndrome, chronic inflammatory demyelinating polyneuropathy (CIDP), Churg-Strauss syndrome, cicatricial pemphigoid, CREST syndrome, cold agglutinin disease, Crohn's disease, dermatomyositis, dilated cardiomyopathy, discoid lupus, epidermolysis bullosa acquisita, essential mixed

10 cryoglobulinemia, factor VIII deficiency, fibromyalgia-fibromyositis, glomerulonephritis, Grave's disease, Guillain-Barre, Goodpasture's syndrome, graft-versus-host disease (GVHD), Hashimoto's thyroiditis, hemophilia A, idiopathic membranous neuropathy, idiopathic pulmonary fibrosis, idiopathic thrombocytopenia purpura (ITP), IgA neuropathy, IgM polyneuropathies, immune mediated thrombocytopenia, juvenile arthritis, Kawasaki's

15 disease, lichen plantus, lichen sclerosus, lupus erythematosus, Meniere's disease, mixed connective tissue disease, mucous membrane pemphigoid, multiple sclerosis, type 1 diabetes mellitus, Multifocal motor neuropathy (MMN), myasthenia gravis, paraneoplastic bullous pemphigoid, pemphigoid gestationis, pemphigus vulgaris, pemphigus foliaceus, pernicious anemia, polyarteritis nodosa, polychondritis, polyglandular syndromes,

20 polymyalgia rheumatica, polymyositis and dermatomyositis, primary agammaglobulinemia, primary biliary cirrhosis, psoriasis, psoriatic arthritis, relapsing polychondritis, Reynauld's phenomenon, Reiter's syndrome, rheumatoid arthritis, sarcoidosis, scleroderma, Sjorgen's syndrome, solid organ transplant rejection, stiff-man syndrome, systemic lupus erythematosus, takayasu arteritis, toxic epidermal necrolysis

25 (TEN), Stevens Johnson syndrome (SJS), temporal arteritis/giant cell arteritis, thrombotic thrombocytopenia purpura, ulcerative colitis, uveitis, dermatitis herpetiformis vasculitis, anti-neutrophil cytoplasmic antibody-associated vasculitides, vitiligo, and Wegner's granulomatosis.

In certain embodiments, the autoimmune disease is an autoimmune channelopathy. In

30 certain embodiments, the channelopathy is selected from the group consisting of autoimmune limbic encephalitis, epilepsy, neuromyelitis optica, Lambert-Eaton myasthenic syndrome, myasthenia gravis, anti- N-Methyl-D-aspartate (NMDA) receptor encephalitis, anti- α -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor encephalitis, Morvan syndrome, neuromyotonia, pediatric autoimmune neuropsychiatric disorders

associated with streptococcal infection (PANDAS), and Glycine receptor antibody-associated disorder.

In certain embodiments, the FcRn antagonist is administered to the subject simultaneously or sequentially with an additional therapeutic agent. In certain embodiments, the additional therapeutic agent is an anti-inflammatory agent. In certain embodiments, the additional therapeutic agent is rituximab, daclizumab, basiliximab, muromonab-cd3, infliximab, adalimumab, omalizumab, efalizumab, natalizumab, tocilizumab, eculizumab, golimumab, canakinumab, ustekinumab, or belimumab. In certain embodiments, the additional therapeutic agent is a leucocyte depleting agent. In certain embodiments, the additional therapeutic agent is a B-cell depleting agent. In certain embodiments, the B-cell depleting agent is an antibody, e.g., an antibody that specifically binds to CD10, CD19, CD20, CD21, CD22, CD23, CD24, CD37, CD53, CD70, CD72, CD74, CD75, CD77, CD79a, CD79b, CD80, CD81, CD82, CD83, CD84, CD85, or CD86.

In another aspect, the instant disclosure provides a nucleic acid molecule encoding an FcRn-antagonist disclosed herein. In another aspect, the instant disclosure provides an expression vector comprising a nucleic acid molecule encoding an FcRn-antagonist disclosed herein. In another aspect, the instant disclosure provides a host cell comprising an expression vector or nucleic acid encoding an FcRn-antagonist disclosed herein. In another aspect, the instant disclosure provides a method of producing an FcRn-antagonist, the method comprising culturing a host cell disclosed herein under conditions such that an FcRn-antagonist is expressed.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG.1 depicts the results of experiments to determine the effect of Fc-Abdeg and HEL-Abdeg on the serum levels of a tracer antibody (FR70-hIgG1) in cynomolgous monkey.

FIG.2 depicts the results of experiments to determine the effect of Fc-Abdeg and HEL-Abdeg on total IgG serum levels in cynomolgous monkey.

FIG.3 depicts the results of experiments the effect of Fc-Abdeg and HEL-Abdeg on albumin levels in cynomolgous monkey.

FIG.4 depicts the results of experiments to determine the effect of Fc-Abdeg and IVIG on the serum levels of a tracer antibody (FR70-hIgG1) in cynomolgous monkey.

FIG.5 depicts the results of ELISA assays comparing the affinity of Fc-Abdeg, Fc-Abdeg-POT and Fc-Abdeg-S239D/I332E for human CD16a.

FIG.6 depicts the results of ELISA assays comparing the affinity of Fc-Abdeg, Fc-Abdeg-POT and Fc-Abdeg-S239D/I332E for murine CD16-2.

FIG.7 depicts the results of experiments to determine the effect of Fc-Abdeg, Abdeg-POT and Fc-AbdegS239D/I332E on anti-CD20-induced ADCC-signal using the Promega's Raji-based ADCC reporter bioassay.

FIG.9 depicts the results of experiments to determine the effect of Fc-Abdeg and Abdeg- POT on anti-CD70-induced lysis of CD70+ U266 cells *in vitro*.

FIG.9 depicts the results of experiments to determine the effect of Fc-Abdeg, Fc-Abdeg-POT, Fc-Abdeg-S239D/I332E and IVIG on platelet levels in an acute murine model for immune thrombocytopenia.

FIG.10 depicts the result of an exemplary gel filtration purification of Fc-Abdeg.

DETAILED DESCRIPTION

The present disclosure provides novel FcRn antagonist compositions. These compositions generally comprise a variant Fc region, or FcRn-binding fragment thereof, that binds specifically to FcRn with increased affinity and reduced pH dependence relative to the native Fc region. The invention is based, in part, on the surprising finding that an isolated variant Fc region (e.g., a variant Fc region comprising the amino acids Y, T, E, K, F, and Y at EU positions 252, 254, 256, 433, 434, and 436 respectively) is a more efficacious FcRn antagonist *in vivo* than a full-length antibody comprising that variant Fc region. The FcRn antagonist compositions of the present disclosure are particularly useful for reducing the serum levels of Fc-containing agents (e.g., antibodies and immunoadhesins). Accordingly, the instant disclosure also provides methods of treating antibody-mediated disorder (e.g. autoimmune diseases) using the FcRn antagonist compositions disclosed herein. Also provided are nucleic acids encoding the FcRn antagonist compositions, recombinant expression vectors and host cells for making the FcRn antagonist compositions, and pharmaceutical compositions comprising the FcRn antagonist compositions.

I. Definitions

Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. The meaning and scope of the terms should be clear, however, in the event of any latent ambiguity, definitions provided herein take precedent over any dictionary or extrinsic definition. Further, unless otherwise required by context, singular terms shall

include pluralities and plural terms shall include the singular. Generally, nomenclature used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art.

5 In order that the present invention may be more readily understood, certain terms are first defined.

 As used herein the term "FcRn antagonist" refers to any agent comprising an Fc region (e.g., a variant Fc region disclosed herein) that binds specifically to FcRn through the Fc region and inhibits the binding of immunoglobulin to FcRn, with the proviso that the agent
10 is not a full length IgG antibody.

 As used herein, the term "Fc region" refers to the portion of a native immunoglobulin formed by the Fc domains of its two heavy chains. A native Fc region is homodimeric.

 As used herein, the term "variant Fc region" refers to an Fc region with one or more alteration relative to a native Fc region. Alteration can include amino acid substitutions,
15 additions and/or deletions, linkage of additional moieties, and/or alteration the native glycans. The term encompasses heterodimeric Fc regions where each of the constituent Fc domains is different. Examples of such heterodimeric Fc regions include, without limitation, Fc regions made using the "knobs and holes" technology as described in, for example, US 8216805, which is incorporated by reference herein in its entirety. The term also encompasses single
20 chain Fc regions where the constituent Fc domains are linked together by a linker moiety, as described in, for example, US20090252729A1 and US20110081345A1, which are each incorporated by reference herein in their entirety.

 As used herein, the term "Fc domain" refers to the portion of a single immunoglobulin heavy chain beginning in the hinge region just upstream of the papain cleavage site and
25 ending at the C-terminus of the antibody. Accordingly, a complete Fc domain comprises at least a portion of a hinge (e.g., upper, middle, and/or lower hinge region) domain, a CH2 domain, and a CH3 domain.

 As used herein the term "FcRn binding fragment" refers to a portion of an Fc region that is sufficient to confer FcRn binding.

30 As used herein, the term "EU position" refers to the amino acid position in the EU numbering convention for the Fc region described in Edelman, G.M. et al., Proc. Natl. Acad. USA, 63, 78-85 (1969) and Kabat et al, in "Sequences of Proteins of Immunological Interest", U.S. Dept. Health and Human Services, 5th edition, 1991.

As used herein, the term "CH1 domain" refers to the first (most amino terminal) constant region domain of an immunoglobulin heavy chain that extends from about EU positions 118-215. The CH1 domain is adjacent to the VH domain and amino terminal to the hinge region of an immunoglobulin heavy chain molecule, and does not form a part of the Fc region of an immunoglobulin heavy chain.

As used herein, the term "hinge region" refers to the portion of a heavy chain molecule that joins the CH1 domain to the CH2 domain. This hinge region comprises approximately 25 residues and is flexible, thus allowing the two N-terminal antigen binding regions to move independently. Hinge regions can be subdivided into three distinct domains: upper, middle, and lower hinge domains (Roux et al. J. Immunol. 161: 4083 (1998)). The FcRn antagonists of the instant disclosure can include all or a portion of a hinge region.

As used herein, the term "CH2 domain" refers to the portion of a heavy chain immunoglobulin molecule that extends from about EU positions 231-340.

As used herein, the term "CH3 domain" includes the portion of a heavy chain immunoglobulin molecule that extends approximately 110 residues from N-terminus of the CH2 domain, e.g., from about position 341-446 (EU numbering system).

As used herein, the term "FcRn" refers to a neonatal Fc receptor. Exemplary FcRn molecules include human FcRn encoded by the FCGRT gene as set forth in RefSeq NM_004107.

As used herein, the term "CD16" refers to FcγRIII Fc receptors that are required for Antibody-Dependent Cell-mediated Cytotoxicity (ADCC). Exemplary CD16 molecules include human CD16a as set forth in RefSeq NM_000569.

As used herein, the term "free cysteine" refers to native or engineered cysteine amino acid residue that exists in a substantially reduced form in a mature FcRn antagonist.

As used herein, the term "antibody" refers to immunoglobulin molecules comprising four polypeptide chains, two heavy (H) chains and two light (L) chains interconnected by disulfide bonds, as well as multimers thereof (e.g., IgM). Each heavy chain comprises a heavy chain variable region (abbreviated VH) and a heavy chain constant region. The heavy chain constant region comprises three domains, CH1, CH2 and CH3. Each light chain comprises a light chain variable region (abbreviated VL) and a light chain constant region. The light chain constant region comprises one domain (CL). The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FR).

As used herein the term “N-linked glycan” refers to the N-linked glycan attached to the nitrogen (N) in the side chain of asparagine in the sequon (i.e., Asn-X-Ser or Asn-X-Thr sequence, where X is any amino acid except proline) present in the CH2 domain of an Fc region. Such N-Glycans are fully described in, for example, Drickamer K, Taylor ME
5 (2006). Introduction to Glycobiology, 2nd ed., which is incorporated herein by reference in its entirety.

As used herein the term “afucosylated” refers to an N-linked glycan which lacks a core fucose molecule as described in US8067232, the contents of which is incorporated by reference herein in its entirety.

10 As used herein the term “bisecting GlcNAc” refers to an N-linked glycan having an N-acetylglucosamine (GlcNAc) molecule linked to a core mannose molecule, as described in US8021856, the contents of which is incorporated by reference herein in its entirety.

As used herein, the term “antibody-mediated disorder” refers to any disease or disorder caused or exacerbated by the presence of an antibody in a subject.

15 As used herein, the term “Fc-containing agent” is any molecule that comprises an Fc region.

As used herein, the term “leucocyte depleting agent” refers to an agent that reduces the number of leucocytes in a subject upon administration.

20 As used herein, the term “B-cell depleting agent” refers to an agent that reduces the number of B-cells in a subject upon administration.

As used herein, the term “T-cell depleting agent” refers to an agent that reduces the number of T-cells in a subject upon administration.

As used herein, the term “autoimmune channelopathy” refers to a diseases caused by autoantibodies against an ion channel subunit or a molecule that regulates the channel.

25 As used herein, the term "treat," "treating," and "treatment" refer to therapeutic or preventative measures described herein. The methods of "treatment" employ administration to a subject, an antibody or antigen binding fragment thereof of the present invention, for example, a subject having an IL-6-associated disease or disorder (e.g. inflammation and cancer) or predisposed to having such a disease or disorder, in order to prevent, cure, delay,
30 reduce the severity of, or ameliorate one or more symptoms of the disease or disorder or recurring disease or disorder, or in order to prolong the survival of a subject beyond that expected in the absence of such treatment. As used herein, the term "subject" includes any human or non-human animal.

As used herein, the term “immunoadhesin” refers to an antibody-like molecule, which comprises a functional domain of a binding protein (e.g., a receptor, ligand, or cell-adhesion molecule) with an Fc region.

5 **II. FcRn Antagonists**

In one aspect, the invention provides novel FcRn antagonist compositions. In general, these compositions comprise a variant Fc region, or FcRn-binding fragment thereof, that binds specifically to FcRn with increased affinity and reduced pH dependence relative to a native Fc region. These FcRn antagonists inhibit the binding of Fc-containing agents (e.g.,
10 antibodies and immunoadhesins) to FcRn *in vivo*, which results in an increased rate of degradation of the Fc-containing agents and, concomitantly, a reduced serum level of these agents.

The instant specification discloses, for the first time, that an isolated variant Fc region (e.g., a variant Fc region comprising the amino acids Y, T, E, K, F, and Y at EU positions
15 252, 254, 256, 433, 434, and 436 respectively) is a more efficacious FcRn antagonist *in vivo* than a full-length antibody comprising the same variant Fc region. Accordingly, in certain embodiments, the FcRn antagonist compositions are not full-length antibodies. In certain embodiments, the FcRn antagonist compositions do not comprise an antibody variable domain. In certain embodiments, the FcRn antagonist compositions do not comprise an
20 antibody variable domain or a CH1 domain. However, in certain embodiments, the FcRn antagonist compositions may comprise a variant Fc region linked to one or more additional binding domains or moieties, including antibody variable domains.

Any Fc region can be altered to produce a variant Fc region for use in the FcRn antagonist compositions disclosed herein. In general, an Fc region, or FcRn-binding
25 fragment thereof, is from a human immunoglobulin. It is understood, however, that the Fc region may be derived from an immunoglobulin of any other mammalian species, including for example, a Camelid species, a rodent (e.g. a mouse, rat, rabbit, guinea pig) or non-human primate (e.g. chimpanzee, macaque) species. Moreover, the Fc region or portion thereof may be derived from any immunoglobulin class, including IgM, IgG, IgD, IgA and IgE, and any
30 immunoglobulin isotype, including IgG1, IgG2, IgG3 and IgG4. In certain embodiments, the Fc region is an IgG Fc region (e.g., a human IgG region). In certain embodiments, the Fc region is an IgG1 Fc region (e.g., a human IgG1 region). In certain embodiments, the Fc region is a chimeric Fc region comprising portions of several different Fc regions. Suitable

examples of chimeric Fc regions are set forth in US20110243966A1, which is herein incorporated by reference in its entirety. A variety of Fc region gene sequences (e.g. human constant region gene sequences) are available in the form of publicly accessible deposits. It will be appreciated that the scope of this invention encompasses alleles, variants and

5 mutations of Fc regions.

An Fc region can be further truncated or internally deleted to produce a minimal FcRn-binding fragment thereof. The ability of an Fc-region fragment to bind to FcRn can be determined using any art recognized binding assay e.g., ELISA.

To enhance the manufacturability of the FcRn antagonists disclosed herein, it is
10 preferable that the constituent Fc regions do not do comprise any non-disulphide bonded cysteine residues. Accordingly, in certain embodiments the Fc regions do not comprise a free cysteine residue.

Any Fc variant, or FcRn-binding fragment thereof, that binds specifically to FcRn with increased affinity and reduced pH dependence relative to the native Fc region can be
15 used in the FcRn antagonist compositions disclosed herein. In certain embodiments, the variant Fc region comprises amino acid alterations, substitutions, insertions and/or deletions that confer the desired characteristics. In certain embodiments, the variant Fc region or fragment comprises the amino acids Y, T, E, K, F, and Y at EU positions 252, 254, 256, 433, 434, and 436 respectively. Non-limiting examples of amino acid sequences that can be used
20 in variant Fc regions are set forth in Table 1, herein. In certain embodiments, the amino acid sequence of the Fc domains of the variant Fc region comprises the amino acid sequence set forth in SEQ ID NO:1. In certain embodiments, the amino acid sequence of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO:1, 2, or 3. In certain embodiments an FcRn-antagonist consists of a variant Fc region, wherein the
25 amino acid sequence of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO:1, 2, or 3.

Table 1. Amino acid sequences of non-limiting examples of variant Fc regions

SEQ ID NO	Amino Acid Sequence
1	CPPCPAPELLGGPSVFLFPPKPKD <u>TL</u> <u>Y</u> <u>I</u> <u>T</u> <u>R</u> <u>E</u> PEVTCVVVDVSHEDPEVKFNWYVD GVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKT ISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENN YKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVVFSCSVMHEAL <u>K</u> <u>F</u> HYTQKSLSLSP

	G
2	DKTHTCPCPAPELLGGPSVFLFPPKPKDTL <u>Y</u> <u>I</u> <u>T</u> <u>R</u> <u>E</u> PEVTCVVVDVSHEDPEVKF NWTVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSVMSHEAL <u>K</u> <u>F</u> HYTQKS LSLSPGK
3	DKTHTCPCPAPELLGGPSVFLFPPKPKDTL <u>Y</u> <u>I</u> <u>T</u> <u>R</u> <u>E</u> PEVTCVVVDVSHEDPEVKF NWTVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSVMSHEAL <u>K</u> <u>F</u> HYTQKS LSLSPG
Amino acids at EU positions 252, 254, 256, 433, and 434 are underlined	

In certain embodiments, the variant Fc region has altered (e.g., increased or decreased) binding affinity for an additional Fc receptor. The variant Fc region can have altered (e.g., increased or decreased) binding affinity for one or more of Fcγ receptors e.g., FcγRI (CD64), FcγRIIA (CD32), FcγRIIB (CD32), FcγRIIIA (CD16a), and FcγRIIIB (CD16b). Any art recognized means of altering the affinity for an additional Fc receptor can be employed. In certain embodiments, the amino acid sequence of the variant Fc region is altered.

In certain embodiments, the variant Fc region comprises a non-naturally occurring amino acid residue at one or more positions selected from the group consisting of 234, 235, 236, 239, 240, 241, 243, 244, 245, 247, 252, 254, 256, 262, 263, 264, 265, 266, 267, 269, 296, 297, 298, 299, 313, 325, 326, 327, 328, 329, 330, 332, 333, and 334 as numbered by the EU index as set forth in Kabat. Optionally, the Fc region may comprise a non-naturally occurring amino acid residue at additional and/or alternative positions known to one skilled in the art (see, e.g., U.S. Pat. Nos. 5,624,821; 6,277,375; 6,737,056; PCT Patent Publications WO 01/58957; WO 02/06919; WO 04/016750; WO 04/029207; WO 04/035752 and WO 05/040217, the contents of which are incorporated by reference herein in their entirety).

In certain embodiments, the variant Fc region comprises at least one non-naturally occurring amino acid residue selected from the group consisting of 234D, 234E, 234N, 234Q, 234T, 234H, 234Y, 234I, 234V, 234F, 235A, 235D, 235R, 235W, 235P, 235S, 235N, 235Q,

235T, 235H, 235Y, 235I, 235V, 235F, 236E, 239D, 239E, 239N, 239Q, 239F, 239T, 239H, 239Y, 240I, 240A, 240T, 240M, 241W, 241 L, 241Y, 241E, 241R, 243W, 243L 243Y, 243R, 243Q, 244H, 245A, 247V, 247G, 252Y, 254T, 256E, 262I, 262A, 262T, 262E, 263I, 263A, 263T, 263M, 264L, 264I, 264W, 264T, 264R, 264F, 264M, 264Y, 264E, 265G, 265N, 5 265Q, 265Y, 265F, 265V, 265I, 265L, 265H, 265T, 266I, 266A, 266T, 266M, 267Q, 267L, 269H, 269Y, 269F, 269R, 296E, 296Q, 296D, 296N, 296S, 296T, 296L, 296I, 296H, 269G, 297S, 297D, 297E, 298H, 298I, 298T, 298F, 299I, 299L, 299A, 299S, 299V, 299H, 299F, 299E, 313F, 325Q, 325L, 325I, 325D, 325E, 325A, 325T, 325V, 325H, 327G, 327W, 327N, 327L, 328S, 328M, 328D, 328E, 328N, 328Q, 328F, 328I, 328V, 328T, 328H, 328A, 329F, 10 329H, 329Q, 330K, 330G, 330T, 330C, 330L, 330Y, 330V, 330I, 330F, 330R, 330H, 332D, 332S, 332W, 332F, 332E, 332N, 332Q, 332T, 332H, 332Y, and 332A as numbered by the EU index as set forth in Kabat. Optionally, the Fc region may comprise additional and/or alternative non-naturally occurring amino acid residues known to one skilled in the art (see, e.g., U.S. Pat. Nos. 5,624,821; 6,277,375; 6,737,056; PCT Patent Publications WO 01/58957; 15 WO 02/06919; WO 04/016750; WO 04/029207; WO 04/035752 and WO 05/040217, the contents of which are incorporated by reference herein in their entirety).

Other known Fc variants that may be used in the FcRn antagonists disclosed herein include without limitations those disclosed in Ghetie et al., 1997, *Nat. Biotech.* 15:637-40; Duncan et al, 1988, *Nature* 332:563-564; Lund et al., 1991, *J. Immunol.*, 147:2657-2662; 20 Lund et al, 1992, *Mol. Immunol.*, 29:53-59; Alegre et al, 1994, *Transplantation* 57:1537-1543; Hutchins et al., 1995, *Proc Natl. Acad Sci USA*, 92:11980-11984; Jefferis et al, 1995, *Immunol Lett.*, 44:111-117; Lund et al., 1995, *Faseb J.*, 9:115-119; Jefferis et al, 1996, *Immunol Lett.*, 54:101-104; Lund et al, 1996, *J. Immunol.*, 157:4963-4969; Armour et al., 1999, *Eur J Immunol* 29:2613-2624; Idusogie et al, 2000, *J. Immunol.*, 164:4178-4184; 25 Reddy et al, 2000, *J. Immunol.*, 164:1925-1933; Xu et al., 2000, *Cell Immunol.*, 200:16-26; Idusogie et al, 2001, *J. Immunol.*, 166:2571-2575; Shields et al., 2001, *J Biol. Chem.*, 276:6591-6604; Jefferis et al, 2002, *Immunol Lett.*, 82:57-65; Presta et al., 2002, *Biochem Soc Trans.*, 30:487-490; U.S. Pat. Nos. 5,624,821; 5,885,573; 5,677,425; 6,165,745; 6,277,375; 5,869,046; 6,121,022; 5,624,821; 5,648,260; 6,528,624; 6,194,551; 6,737,056; 30 6,821,505; 6,277,375; U.S. Patent Publication Nos. 2004/0002587 and PCT Publications WO 94/29351; WO 99/58572; WO 00/42072; WO 02/060919; WO 04/029207; WO 04/099249; WO 04/063351, the contents of which are incorporated by reference herein in their entirety.

In certain embodiments, the variant Fc region is a heterodimer, where the constituent Fc domains are different from each other. Methods of producing Fc heterodimers are known

in the art (see e.g., US 8216805, which is incorporated by reference herein in its entirety). In certain embodiments, the variant Fc region is a single chain Fc region, where the constituent Fc domains are linked together by a linker moiety. Methods of producing single chain Fc regions are known in the art (see e.g., US20090252729A1 and US20110081345A1, which are each incorporated by reference herein in their entirety).

It is believed that pathogenic IgG antibodies observed in autoimmune diseases are either the pathogenic triggers for these diseases or contribute to disease progression and mediate disease through the inappropriate activation of cellular Fc receptors. Aggregated autoantibodies and/or autoantibodies complexed with self antigens (immune complexes) bind to activating Fc receptors, causing numerous autoimmune diseases (which occur in part because of immunologically mediated inflammation against self tissues) (see e.g., Clarkson et al., NEJM 314(9), 1236-1239 (2013)); US20040010124A1; US20040047862A1; and US2004/0265321A1, which are each incorporated by reference herein in their entirety). Accordingly, to treat antibody-mediated disorders (e.g. autoimmune diseases), it would be advantageous to both remove the deleterious autoantibodies and to block the interaction of the immune complexes of these antibodies with activating Fc receptors (e.g., Fcγ receptors, such as CD16a).

Accordingly, in certain embodiments, the variant Fc region of the FcRn antagonist exhibits increased binding to CD16a (e.g., human CD16a). This is particularly advantageous in that it allows the FcRn antagonist to additionally antagonize the immune complex-induced inflammatory response of autoantibodies being targeted for removal by FcRn inhibition. Any art recognized means of increasing affinity for CD16a (e.g., human CD16a) can be employed. In certain embodiments, the FcRn-antagonist comprises a variant Fc-region comprising an N-linked glycan (e.g., at EU position 297). In this case it is possible to increase the binding affinity of the FcRn-antagonist for CD16a by altering the glycan structure. Alterations of the N-linked glycan of Fc regions are well known in the art. For example, afucosylated N-linked glycans or N-glycans having a bisecting GlcNac structure have been shown to exhibit increased affinity for CD16a. Accordingly, in certain embodiments, the N-linked glycan is afucosylated. Afucosylation can be achieved using any art recognized means. For example, an FcRn-antagonist can be expressed in cells lacking fucosyl transferase, such that fucose is not added to the N-linked glycan at EU position 297 of the variant Fc region (see e.g., US 8,067,232, the contents of which is incorporated by reference herein in its entirety). In certain embodiments, the N-linked glycan has a bisecting GlcNac structure. The bisecting GlcNac structure can be achieved using any art recognized means. For example, an FcRn-

antagonist can be expressed in cells expressing beta1-4-N-acetylglucosaminyltransferase III (GnTIII), such that bisecting GlcNac is added to the N-linked glycan at EU position 297 of the variant Fc region (see e.g., US 8021856, the contents of which is incorporated by reference herein in its entirety). Additionally or alternatively, alterations of the N-linked glycan structure can also be achieved by enzymatic means *in vitro*.

In certain embodiments, the instant disclosure provides FcRn-antagonist compositions wherein a portion of the FcRn-antagonist molecules contained therein comprise altered glycan structures. In certain embodiments, the FcRn-antagonist composition comprises a plurality of FcRn-antagonist molecules disclosed herein, wherein at least 50% (optionally, at least 60, 70, 80, 90, 95, or 99%) of the molecules comprise an Fc region or FcRn-binding fragment thereof having an afucosylated N-linked glycan. In certain embodiments, the FcRn-antagonist composition comprising a plurality of FcRn-antagonist molecules disclosed herein, wherein at least 50% (optionally, at least 60, 70, 80, 90, 95, or 99%) of the molecules comprise an Fc region or FcRn-binding fragment thereof comprising an N-linked glycan having a bisecting GlcNac.

In certain embodiments, the variant Fc region does not comprise an N-linked glycan. This can be achieved using any art recognized methods. For example, the Fc variant can be expressed in a cell that is incapable of N-linked glycosylation. Additionally or alternatively, the amino acid sequence of the Fc variant can be altered to prevent or inhibit N-linked glycosylation (e.g., by mutation of the NXT sequon). Alternatively, the Fc variant can be synthesized in an acellular system (e.g., chemically synthesized).

In certain embodiments, FcRn-antagonist molecules may be modified, e.g., by the covalent attachment of a molecule (e.g., a binding or imaging moiety) to the FcRn-antagonist such that covalent attachment does not prevent the FcRn-antagonist from specifically binding to FcRn. For example, but not by way of limitation, the FcRn-antagonist may be modified by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc.

In certain embodiments, the FcRn antagonist comprises a variant Fc region linked to a half-life extender. As used herein, the term "half-life extender" refers to any molecule that, when linked to an FcRn antagonist disclosed herein, increases the half-life of an FcRn antagonist. Any half-life extender may be linked (either covalently or non-covalently) to the FcRn antagonist. In certain embodiments, the half-life extender is polyethylene glycol or human serum albumin. In certain embodiments, the FcRn antagonist is linked to a binding

molecule that specifically binds to a half-life extender present in a subject, such as a blood-carried molecule or cell, such as serum albumin (e.g., human serum albumin), IgG, erythrocytes, etc.

The FcRn antagonists disclosed herein have excellent manufacturability. For example, as shown in Example 5 herein, they can be expressed at high levels in mammalian cells (e.g., at 6g/L in CHO cells in a 10 L stirred tank bioreactor). Moreover, after Protein A purification, the resultant purified FcRn antagonist composition has a very high percentage of FcRn antagonist monomers, and contains an extremely low level of FcRn antagonist protein aggregates and degradation products. Accordingly, in certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules as disclosed herein, wherein greater than 95% the of the FcRn antagonist molecules in the composition are monomers (e.g., greater than 95, 96, 97, 98, 99, 99.1, 99.2, 99.3, 99.4, 99.5, 99.6, 99.7, 99.8, 99.9 %). In certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules disclosed herein, wherein less than 5% the of the FcRn antagonist molecules in the composition are present in aggregates, (e.g., less than 5, 4, 3, 2, 1, 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.2, 0.1 %). In certain embodiments, the instant disclosure provides an FcRn antagonist composition comprising a plurality of FcRn antagonist molecules disclosed herein, wherein the composition is substantially free of FcRn antagonist molecule degradation products.

III. Uses of FcRn Antagonists

The FcRn antagonist compositions of the present disclosure are particularly useful for reducing the serum levels of Fc-containing agents (e.g., antibodies and immunoadhesins). Accordingly, in one aspect the instant disclosure provides a method of inhibiting FcRn function in a subject, the method generally comprising administering to the subject an effective amount of an FcRn antagonist composition (e.g., a pharmaceutical composition) disclosed herein.

The reduction of serum levels of Fc-containing agents (e.g., antibodies and immunoadhesins) is particularly applicable to the treatment of antibody-mediated disorders (e.g. autoimmune diseases). Accordingly, in one aspect the instant disclosure provides methods of treating antibody-mediated disorders (e.g. autoimmune diseases) using the FcRn antagonist compositions disclosed herein.

Any antibody-mediated disorder can be treated using the FcRn antagonist compositions disclosed herein. In certain embodiments, the antibody-mediated disorder is one that is amenable to treatment by IVIG. In certain embodiments, the antibody-mediated disorder is an autoimmune disease. Non-limiting autoimmune diseases include allogenic islet graft rejection, alopecia areata, ankylosing spondylitis, antiphospholipid syndrome, autoimmune Addison's disease, Alzheimer's disease, antineutrophil cytoplasmic autoantibodies (ANCA), autoimmune diseases of the adrenal gland, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune myocarditis, autoimmune neutropenia, autoimmune oophoritis and orchitis, autoimmune thrombocytopenia, autoimmune urticaria, Behcet's disease, bullous pemphigoid, cardiomyopathy, Castleman's syndrome, celiac spruce-dermatitis, chronic fatigue immune dysfunction syndrome, chronic inflammatory demyelinating polyneuropathy (CIDP), Churg-Strauss syndrome, cicatricial pemphigoid, CREST syndrome, cold agglutinin disease, Crohn's disease, dermatomyositis, discoid lupus, essential mixed cryoglobulinemia, factor VIII deficiency, fibromyalgia-fibromyositis, glomerulonephritis, Grave's disease, Guillain-Barre, Goodpasture's syndrome, graft-versus-host disease (GVHD), Hashimoto's thyroiditis, hemophilia A, idiopathic pulmonary fibrosis, idiopathic thrombocytopenia purpura (ITP), IgA neuropathy, IgM polyneuropathies, immune mediated thrombocytopenia, juvenile arthritis, Kawasaki's disease, lichen plantus, lupus erthematosus, Meniere's disease, mixed connective tissue disease, multiple sclerosis, type 1 diabetes mellitus, Multifocal motor neuropathy (MMN), myasthenia gravis, paraneoplastic bullous pemphigoid, pemphigus vulgaris, pemphigus foliaceus, pernicious anemia, polyarteritis nodosa, polychondritis, polyglandular syndromes, polymyalgia rheumatica, polymyositis and dermatomyositis, primary agammaglobulinemia, primary biliary cirrhosis, psoriasis, psoriatic arthritis, Reynauld's phenomenon, Reiter's syndrome, rheumatoid arthritis, sarcoidosis, scleroderma, Sjorgen's syndrome, solid organ transplant rejection, stiff-man syndrome, systemic lupus erythematosus, takayasu arteritis, toxic epidermal necrolysis (TEN), Stevens Johnson syndrome (SJS), temporal arteritis/giant cell arteritis, thrombotic thrombocytopenia purpura, ulcerative colitis, uveitis, dermatitis herpetiformis vasculitis, anti-neutrophil cytoplasmic antibody-associated vasculitides, vitiligo, and Wegner's granulomatosis.

In certain embodiments, the autoimmune disease is an autoimmune channelopathy. Non-limiting channelopathies include neuromyelitis optica, Lambert-Eaton myasthenic syndrome, myasthenia gravis, anti- N-Methyl-D-aspartate (NMDA) receptor encephalitis, anti- α -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor encephalitis,

Morvan syndrome, and Glycine receptor antibody-associated disorder.

The FcRn antagonist compositions of the instant disclosure are particularly suited to treating antibody-mediated disorders characterized by an over production of serum immunoglobulin. Accordingly, in certain embodiments, the FcRn antagonist compositions
5 are used to treat hypergammaglobulinemia.

The FcRn antagonist compositions can also be used in combination with one or more additional therapeutic agents. In certain embodiments, the additional therapeutic agent is an anti-inflammatory agent. Any inflammatory agent can be used in combination with the compositions disclosed herein. In certain embodiments, the therapeutic agent is rituximab,
10 daclizumab, basiliximab, muronomab-cd3, infliximab, adalimumab, omalizumab, efalizumab, natalizumab, tocilizumab, eculizumab, golimumab, canakinumab, ustekinumab, or belimumab. In certain embodiments, the additional therapeutic agent is leucocyte depleting agent (e.g., B-cell or T-cell depleting agent). Any leucocyte depleting agent can be used in combination with the FcRn antagonist compositions disclosed herein. In certain
15 embodiments, the leucocyte depleting agent is a B-cell depleting agent. In certain embodiments, the leucocyte depleting agent is an antibody against a cell surface marker. Suitable cell surface markers include, without limitation, CD10, CD19, CD20, CD21, CD22, CD23, CD24, CD37, CD53, CD70, CD72, CD74, CD75, CD77, CD79a, CD79b, CD80, CD81, CD82, CD83, CD84, CD85, or CD86. The FcRn antagonist and the additional
20 therapeutic agent(s) can be administered to the subject simultaneously or sequentially, via the same or different route(s) of administration.

The FcRn antagonist compositions of the instant disclosure are also well suited to rapidly reducing the serum levels of an Fc-containing agent in subject. Such rapid clearance is advantageous in cases where the Fc-containing agent is toxic (e.g., an antibody-drug
25 conjugate or an agent that is immunogenic) because it reduces the exposure of the subject to the drug. Rapid clearance is also advantageous in cases where the Fc-containing agent is an imaging agent that requires a low serum level of the agent to facilitate imaging. Accordingly, in certain embodiments, the FcRn antagonist compositions are used to reduce the serum levels of an Fc-containing agent in subject that has been administered the Fc-containing
30 agent. The serum levels of any Fc-containing agent (e.g., therapeutic or diagnostic agent) can be reduced using the FcRn antagonist compositions disclosed herein. Non limiting examples of Fc-containing agents include imaging agents (e.g., labeled antibodies), antibody drug conjugates, or immunogenic agents (e.g., non-human antibodies or immunoadhesins). The FcRn antagonist can be administered simultaneously with the Fc-containing agent or

sequentially (e.g., before or after the Fc-containing agent).

Furthermore, in diseases or conditions requiring administration of a therapeutic agent, the subject will often develop antibodies (e.g., anti-drug antibodies) against the therapeutic agent, which, in turn, prevent the therapeutic agent from being available for its intended therapeutic purpose or cause an adverse reaction in the subject. Accordingly, the FcRn antagonist compositions disclosed herein can also be used to remove antibodies (e.g., anti-drug antibodies) against the therapeutic agent that develop in a subject.

The FcRn antagonist compositions disclosed herein can also be used in combination with the therapeutic protein to enhance the benefit of the therapeutic protein by reducing the levels of IgG; wherein, IgG antibodies are responsible for the decreased bioavailability of a therapeutic protein. In certain embodiments the instant disclosure provides a method of treating a disorder resulting from an immune response to a clotting factor comprising administering to a subject a therapeutically effective amount of an FcRn antagonist compositions disclosed herein. Suitable clotting factors include, without limitation, fibrinogen, prothrombin, factor V, factor VII, factor VIII, factor IX, factor X, factor XI, factor XII, factor XIII, or von Willebrand's factor. This method may be used to regulate or treat, or prevent an immune response to a clotting factor in a patient suffering, e.g., from hemophilia A or hemophilia B. In certain embodiments, the method may be used to regulate or treat an immune response to, e.g., therapeutic erythropoietin in a patient suffering from pure red cell aplasia (PRCA).

FcRn is responsible for transporting maternal antibodies across the placenta to the fetus in a pregnant woman. Accordingly, if a pregnant female is administered an Fc-containing agent (e.g., a therapeutic antibody), the agent may come in contact with the fetus as a result of the FcRn-mediated transport across the placenta. To avoid any potential deleterious effect of the Fc-containing agent on fetal development, it would be advantageous to block FcRn function. Accordingly, the instant disclosure provides a method of preventing placental transfer of an Fc-containing agent (e.g., a therapeutic antibody) to the fetus in a pregnant woman, the method comprising administering to the woman an FcRn antagonist compositions disclosed herein, either simultaneously or sequentially (prior or post) with the Fc-containing agent.

The FcRn antagonist compositions disclosed herein can also be used to treat inflammatory disorders including, but not limited to, asthma, ulcerative colitis and inflammatory bowel syndrome allergy, including allergic rhinitis/sinusitis, skin allergies (urticaria/hives, angioedema, atopic dermatitis), food allergies, drug allergies, insect allergies,

mastocytosis, arthritis, including osteoarthritis, rheumatoid arthritis, and spondyloarthropathies.

Successful implementation of gene therapy for the treatment of a disease or condition may be hampered by the development of antibodies specific to the therapeutic protein encoded by the transgene as well as possibly to the vector used to deliver the transgene. Accordingly, the FcRn antagonist compositions disclosed herein can be administered in combination with gene therapy to enhance the benefit of the encoded therapeutic protein by reducing the levels of IgG. These methods are particularly useful in situations where IgG antibodies are responsible for the decreased bioavailability of a gene therapy vector or the encoded therapeutic protein. The gene therapy vector may be, e.g., a viral vector such as adenovirus and adeno-associated virus. Diseases that can be treated using gene therapy include, but are not limited to, cystic fibrosis, hemophilia, PRCA, muscular dystrophy, or lysosomal storage diseases, such as, e.g., Gaucher's disease and Fabry's disease.

One skilled in the art would be able, by routine experimentation, to determine what an effective, non-toxic amount of FcRn antagonist composition would be for the purpose of treating an antibody-mediated disorder. For example, a therapeutically active amount of a polypeptide may vary according to factors such as the disease stage (e.g., stage I versus stage IV), age, sex, medical complications (e.g., immunosuppressed conditions or diseases) and weight of the subject, and the ability of the antibody to elicit a desired response in the subject. The dosage regimen may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily, or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation. Generally, however, an effective dosage is expected to be in the range of about 0.1 to 10,000 mg/kg body weight per day e.g., about 1 to 1000, about 10-500, or about 50-250 or mg/kg body weight per day (e.g., about 70 mg/kg body weight per day).

IV. Pharmaceutical Compositions

In another aspect, the instant disclosure provides pharmaceutical compositions comprising an FcRn antagonist or FcRn antagonist composition disclosed herein and a pharmaceutically acceptable carrier or excipient. Examples of suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences by E. W. Martin. Examples of excipients can include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol,

propylene, glycol, water, ethanol, and the like. The composition can also contain pH buffering reagents, and wetting or emulsifying agents.

The pharmaceutical composition can be formulated for parenteral administration (e.g., intravenous or intramuscular) by bolus injection. Formulations for injection can be presented in unit dosage form, e.g., in ampoules or in multidose containers with an added preservative. The compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle, e.g., pyrogen free water.

FcRn antagonists may be linked to chelators such as those described in U.S. Pat. No.5,326,856. The peptide-chelator complex may then be radiolabeled to provide an imaging agent for diagnosis or treatment of diseases or conditions involving the regulation of IgG levels.

V. Production of FcRn Antagonists

In one aspect, the invention provides polynucleotides, vectors and host cells encoding the FcRn antagonists disclosed herein. Methods of making an FcRn antagonists comprising expressing these polynucleotides are also provided.

Polynucleotides encoding the FcRn antagonists disclosed herein are typically inserted in an expression vector for introduction into host cells that may be used to produce the desired quantity of the claimed FcRn antagonists. Accordingly, in certain aspects, the invention provides expression vectors comprising polynucleotides disclosed herein and host cells comprising these vectors and polynucleotides.

The term "vector" or "expression vector" is used herein for the purposes of the specification and claims, to mean vectors used in accordance with the present invention as a vehicle for introducing into and expressing a desired gene in a cell. As known to those skilled in the art, such vectors may easily be selected from the group consisting of plasmids, phages, viruses and retroviruses. In general, vectors compatible with the instant invention will comprise a selection marker, appropriate restriction sites to facilitate cloning of the desired gene and the ability to enter and/or replicate in eukaryotic or prokaryotic cells.

Numerous expression vector systems may be employed for the purposes of this invention. For example, one class of vector utilizes DNA elements, which are derived from animal viruses such as bovine papilloma virus, polyoma virus, adenovirus, vaccinia virus, baculovirus, retroviruses (RSV, MMTV or MOMLV) or SV40 virus. Others involve the use

of polycistronic systems with internal ribosome binding sites. Additionally, cells that have integrated the DNA into their chromosomes may be selected by introducing one or more markers that allow selection of transfected host cells. The marker may provide for prototrophy to an auxotrophic host, biocide resistance (e.g., antibiotics) or resistance to heavy metals such as copper. The selectable marker gene can either be directly linked to the DNA sequences to be expressed, or introduced into the same cell by cotransformation. Additional elements may also be needed for optimal synthesis of mRNA. These elements may include signal sequences, splice signals, as well as transcriptional promoters, enhancers, and termination signals.

More generally, once a vector or DNA sequence encoding an FcRn antagonist has been prepared, the expression vector may be introduced into an appropriate host cell. That is, the host cells may be transformed. Introduction of the plasmid into the host cell can be accomplished by various techniques well known to those of skill in the art. These include, but are not limited to, transfection (including electrophoresis and electroporation), protoplast fusion, calcium phosphate precipitation, cell fusion with enveloped DNA, microinjection, and infection with intact virus. See, Ridgway, A. A. G. "Mammalian Expression Vectors" Chapter 24.2, pp. 470-472 Vectors, Rodriguez and Denhardt, Eds. (Butterworths, Boston, Mass. 1988). Most preferably, plasmid introduction into the host is via electroporation. The transformed cells are grown under conditions appropriate to the production of the FcRn antagonist, and assayed for FcRn antagonist expression. Exemplary assay techniques include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), or fluorescence-activated cell sorter analysis (FACS), immunohistochemistry and the like.

As used herein, the term "transformation" shall be used in a broad sense to refer to the introduction of DNA into a recipient host cell that changes the genotype and consequently results in a change in the recipient cell.

Along those same lines, "host cells" refers to cells that have been transformed with vectors constructed using recombinant DNA techniques and encoding at least one heterologous gene. In descriptions of processes for isolation of polypeptides from recombinant hosts, the terms "cell" and "cell culture" are used interchangeably to denote the source of FcRn antagonist unless it is clearly specified otherwise. In other words, recovery of FcRn antagonist from the "cells" may mean either from spun down whole cells, or from the cell culture containing both the medium and the suspended cells.

In one embodiment, the host cell line used for FcRn antagonist expression is of mammalian origin; those skilled in the art can determine particular host cell lines, which are

best suited for the desired gene product to be expressed therein. Exemplary host cell lines include, but are not limited to, DG44 and DUXB11 (Chinese Hamster Ovary lines, DHFR minus), HELA (human cervical carcinoma), CVI (monkey kidney line), COS (a derivative of CVI with SV40 T antigen), R1610 (Chinese hamster fibroblast) BALBC/3T3 (mouse fibroblast), HAK (hamster kidney line), SP2/O (mouse myeloma), BFA-1c1BPT (bovine endothelial cells), RAJI (human lymphocyte), 293 (human kidney). In one embodiment, the cell line provides for altered glycosylation, e.g., afucosylation, of the FcRn antagonist expressed therefrom (e.g., PER.C6.RTM. (Crucell) or FUT8-knock-out CHO cell lines (PotelligentTM Cells) (Biowa, Princeton, N.J.)). In one embodiment NS0 cells may be used. CHO cells are particularly preferred. Host cell lines are typically available from commercial services, the American Tissue Culture Collection or from published literature.

In vitro production allows scale-up to give large amounts of the desired FcRn antagonist. Techniques for mammalian cell cultivation under tissue culture conditions are known in the art and include homogeneous suspension culture, e.g. in an airlift reactor or in a continuous stirrer reactor, or immobilized or entrapped cell culture, e.g. in hollow fibers, microcapsules, on agarose microbeads or ceramic cartridges. If necessary and/or desired, the solutions of polypeptides can be purified by the customary chromatography methods, for example gel filtration, ion-exchange chromatography, chromatography over DEAE-cellulose and/or (immuno-) affinity chromatography.

Genes encoding the FcRn antagonists of the invention can also be expressed in non-mammalian cells such as bacteria or yeast or plant cells. In this regard it will be appreciated that various unicellular non-mammalian microorganisms such as bacteria can also be transformed; i.e. those capable of being grown in cultures or fermentation. Bacteria, which are susceptible to transformation, include members of the enterobacteriaceae, such as strains of *Escherichia coli* or *Salmonella*; Bacillaceae, such as *Bacillus subtilis*; *Pneumococcus*; *Streptococcus*, and *Haemophilus influenzae*. It will further be appreciated that, when expressed in bacteria, the FcRn antagonists can become part of inclusion bodies. The FcRn antagonists must be isolated, purified and then assembled into functional molecules. In addition to prokaryotes, eukaryotic microbes may also be used. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among eukaryotic microorganisms although a number of other strains are commonly available.

In addition to cell-based expression systems, the FcRn antagonists can also be produced using acellular or chemically synthetic methods. In certain embodiments, the FcRn antagonists are produced by *in vitro* chemical synthesis.

IV. Exemplification

The present invention is further illustrated by the following examples, which should not be construed as further limiting. The contents of Sequence Listing, figures and all references, patents, and published patent applications cited throughout this application are expressly incorporated herein by reference.

Example 1: Effect of Fc-Abdeg on serum IgG levels in cynomolgus monkeys

The effect of a human anti-lysozyme IgG (HEL-Abdeg) and a human IgG Fc region (Fc-Abdeg), comprising the amino acids Y, T, E, K, F, and Y at EU positions 252, 254, 256, 433, 434, and 436, respectively (Fc-Abdeg; SEQ ID NO:2), on serum IgG levels of a tracer antibody was determined in cynomolgus monkeys. Specifically, cynomolgus monkeys were administered 1mg/kg of an anti-murine CD70 hIgG1 tracer antibody (FR70-hIgG1; Oshima et al., Int Immunol 10(4): 517-26 (1998)) by i.v. bolus injection. Animals were infused 5 minutes later with either 7 mg/kg Fc-Abdeg, 20 mg/kg HEL-Abdeg, or PBS (2 monkeys per group). Infusion was performed within 1 hour and animals were administered a volume of 10 ml/kg. Blood samples (3x 150 µl) were taken at 5 min prior to dosing ("pre-dose") and 5 min, 2h, 6h, 24h, 48h, 72h, 96h and 120h after completion of the infusion. Tracer levels were determined by performing a mCD70-binding ELISA and data were plotted relative to tracer levels at end of dosing (FIG.1). Total cynomolgus IgG levels were also determined (FIG.2). The results of these experiments show that Fc-Abdeg reduced tracer antibody more efficiently than equimolar amounts of HEL-Abdeg.

In addition to its key role in the IgG salvage pathway, FcRn is also involved in albumin homeostasis (Chaudhury et al., J Exp Med. 197(3):315-22 (2003). FcRn interacts with IgG-Fc and albumin at distinct sites and binding can happen concurrently (Andersen et al., Nat Commun. 3:610 (2012)). Conceptually, blockage of IgG recycling using Abdeg-modified molecules should not interfere with albumin-FcRn interaction. This hypothesis was confirmed in a mouse *in vivo* study, where the authors showed no influence of an Abdeg-equipped hIgG1 molecule on albumin levels (Patel et al., J Immunol 187(2): 1015-22 (2011)). In the experiment described above, albumin levels were also determined at day -3, day 3 and day 17 after the completion of the infusion. Analogous to the mouse study, no significant changes in albumin levels were observed after Fc-Abdeg or HEL-Abdeg treatment (see FIG.3).

In a subsequent experiment, the antibody-depleting potency of Fc-Abdeg was compared to IVIG. Specifically, cynomolgus monkeys were administered with 1 mg/kg tracer antibody (FR70-hIgG1) 2 days prior to dosing with 70 mg/kg Fc-Abdeg or 2 g/kg IVIG (2 monkeys per group). Infusion of Fc-Abdeg and IVIG was performed within 4 hours and animals were administered a volume of 20 ml/kg. Blood (3x 150 μ l) samples were taken 5 min prior to dosing ("pre-dose"), and 5 min, 2h, 6h, 24h, 48h, 72h, 96h, 120h, and 168h after completion of the infusion. Tracer levels were determined by mCD70-binding ELISA and plotted relative to pre-dose levels (FIG.4). In comparison with IVIG treatment at clinical dose (2 g/kg), 70 mg/kg Fc-Abdeg showed significantly enhanced kinetics of tracer clearance and was also able to clear more efficiently (>95% tracer clearance in 4 days for Abdeg versus ~75% in 7 days for IVIG).

Example 2: Effect of afucosylation on Fc-Abdeg affinity for human CD16a and murine CD16-2

The binding affinity of Fc-Abdeg for hCD16a was determined and compared to the afucosylated form (Fc-Abdeg-POT). In the same experiment an Fc-Abdeg variant showing improved affinity for all Fc γ Rs was included ("Fc-Abdeg-S239D/I332E). Specifically, a Maxisorp plate was coated with 100ng/well of Neutravidin Biotin-binding Protein (ThermoScientific, 31000) and incubated overnight at 4°C. The following day, the plate was blocked with PBS+1% casein for 2 hours at room temperature. Subsequently, 100 μ l/well of a 250ng/ml solution (dilution in PBS + 0.1% casein) of biotinylated hCD16a (Sino Biological Inc., 10389-H27H1-B) was added to the plate and incubated for 1 hour at room temperature prior to applying a concentration gradient of Fc-Abdeg or Fc-Abdeg-POT molecules (1 μ M-0.005 nM) for a further hour. Binding to hCD16a was detected using an HRP-conjugated polyclonal goat anti-human Fc antibody (Jackson ImmunoResearch, 109-035-008) (incubation 1 hour at RT, dilution 1/50,000 in PBS+0.1% casein), followed by addition of 100 μ l room temperature-equilibrated TMB (SDT-reagents #s TMB). Plates were incubated for 10 minutes prior to addition of 100 μ l 0.5N H₂SO₄ and OD450nm measurement. EC₅₀ values were determined using GraphPad Prism software. The results of these experiments, set forth in FIG.5, show that defucosylation of the Fc-Abdeg molecule results in a >30-fold increase in affinity for hCD16a (EC₅₀= 13nM for the Fc-Abdeg-POT vs. EC₅₀ >0.4 μ M for the fucosylated Fc-Abdeg). As expected, binding affinity of the Fc-Abdeg-S239D/I332E variant for hCD16a was increased compared to wild-type Fc-Abdeg (EC₅₀=6nM).

Using a similar experimental procedure as described above, the binding affinity for murine CD16-2 (Sino Biological Inc., 50036-M27H-B) was determined. The results of these experiments, set forth in FIG.6, again show an increased affinity of the afucosylated variant compared to the wild-type Fc-Abdeg ($EC_{50}=11\text{nM}$ vs. $EC_{50}>100\text{nM}$). The fold increase in affinity for mCD16-2 of the Fc-Abdeg-POT variant over the wild-type Fc-Abdeg is lower compared to that observed for binding to human CD16a. This effect was not observed for the Fc-Abdeg-S239D/I332E variant ($EC_{50}=2\text{nM}$), which has a similar fold increase in affinity over wild type Fc-Abdeg for both the human and murine CD16 ($EC_{50}=2\text{nM}$).

Autoantibodies complexed with self antigens bind to activating Fc γ Rs and thereby trigger the autoimmune diseases, which occur in part because of immunologically mediated inflammation against self tissue. The ability of Fc-Abdeg to antagonize the interaction of autoimmune antibodies and Fc γ RIII receptors on NK cells was evaluated in two ADCC-based assays.

Initially, an ADCC reporter bioassay (Promega, G7016) was used to analyze the competitive hCD16a binding potency of Fc-Abdeg, Fc-Abdeg-POT and Fc-Abdeg-S239D/I332E. Specifically, 10,000 CD20-expressing Raji cells (target cells) were incubated with 60,000 Jurkat cells expressing hCD16a (effector cells) in presence of 100ng/ml anti-CD20 antibody and increasing concentration of competitor. Cells were incubated for 6 hours at 37°C prior to measuring the bioluminescence signal, which is a measure of ADCC-activity. The luciferase signal was plotted relative to the signal obtained by 100ng/ml anti-CD20 in the absence of competitor (see FIG.7). These experiments demonstrate that both Fc-Abdeg-POT and Fc-Abdeg-S239D/I332E efficiently block the anti-CD20-induced ADCC signal, whilst incubation with wild-type Fc-Abdeg does not lead to competitive binding for hCD16a expressed on Jurkat cells.

In a next ADCC assay, inhibition of the lytic activity of an anti-hCD70 antibody (27B3-hIgG1) by Fc-Abdeg and Fc-Abdeg-POT was tested as a measure of competitive hCD16 binding. Specifically, about 50,000 hCD70-expressing U266 cells were spiked into about 300,000 freshly purified PBMCs from a healthy donor in the presence of 50 ng/ml of the anti-hCD70 antibody and a concentration-gradient of Fc-Abdeg, Fc-Abdeg-POT and IVIG. The U266 cells were incubated for two days, and subsequent cell lysis was analyzed by FACS using a marker specific for the U266 cells (CD28). The results of these experiments, set forth in FIG.8, show that the anti-CD70 antibody efficiently lyses U266 cells and that this depletion could be attenuated in a dose-dependent fashion by addition of Fc-

Abdeg-POT but not by wild-type Fc-Abdeg nor IVIG. These data demonstrate that Fc-Abdeg POT has enhanced competitive CD16a binding properties relative to wild-type Fc-Abdeg and IVIG.

5 **Example 3: Murine acute ITP model**

The therapeutic potency of Fc-Abdeg, Fc-Abdeg-POT, Fc-Abdeg-S239D/I332E molecules was tested in a mouse model of acute immune thrombocytopenia. Specifically, C57BL/6 mice were treated with IVIG (20mg/animal), Fc-Abdeg (1 mg/animal), Fc-Abdeg-POT (1 mg/animal), Fc-Abdeg-S239D/I332E (1mg/animal) or saline via the intraperitoneal
10 infusion (5 animals/group). Prior to treatment, a blood sample was withdrawn for a baseline measurement of platelet counts. One hour later, mice were treated with 5µg/animal of the anti-mouse platelet antibody MWReg30 (Nieswandt et al., Blood 94:684-93 (1999)). Platelet counts were monitored over 24 hours. Platelet counts were normalized relative to the initial counts for each mouse and platelets numbers were determined using flow cytometry via anti-
15 CD61 staining. The results of these experiments, set forth in FIG.9, demonstrate that pretreatment with Fc-Abdeg reduces MWReg30-induced thrombocytopenia with a similar potency compared to a 7-fold higher molar dose of IVIG, and further, that blockade of FcγRs by Fc-Abdeg POT and Fc-Abdeg-S239D/I332E had a synergistic beneficial effect in this model, as seen by the improved platelet counts at the 180 and 1440 minutes time points.

20

Example 4: Manufacturability Fc-Abdeg

Fc-Abdeg (comprising Fc domains having SEQ ID NO:2) was produced in CHO cells (Evitria, Switzerland) by transient transfection. Following transfection, high titers of Fc-Abdeg were detected in the supernatants (between 200 and 400 mg/ml). A similar favorable
25 production profile was seen when Fc-Abdeg was expressed from an expression construct stably integrated into the CHO GS-XCEED cell line (Lonza, Great-Britain). On average, stable transfectants yielded 3g/L and several clones were identified which produced up to 6g/L Fc-Abdeg in a 10 L stirred tank bioreactor.

The manufacturability of the Fc-Abdeg was further investigated by analysis of
30 aggregates and degradation products following protein A-purification of the aforementioned Fc-Abdeg production runs. Specifically, 137 µg of Fc-Abdeg was loaded on a Superdex 200 10/300 GL gelfiltration column (GE Healthcare) coupled to an ÄktaPurifier chromatography system. Results of this experiment, set forth in FIG.10, showed that only a very small percentage of Fc-Abdeg aggregates was observed (~0.5%), whilst no Fc-Abdeg degradation

products were detected. Additionally, applying various stress conditions (freeze-thaw, rotational or temperature stress) to the protein A-purified Fc-Abdeg did not lead to any apparent change in physicochemical and functional properties. Taken together, these data demonstrate the excellent manufacturability of the Fc-Abdeg.

We claim:

1. An isolated FcRn-antagonist comprising a variant Fc region, or FcRn-binding fragment thereof, wherein the amino acid sequence of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO:1, 2, or 3, and wherein the FcRn-antagonist does not comprise an antibody variable region or a CH1 domain.
2. The FcRn-antagonist of claims 1, wherein the variant Fc region has an increased affinity for an Fc gamma receptor relative to the affinity of a wild-type IgG1 Fc region for the Fc gamma receptor.
3. An isolated FcRn-antagonist comprising a variant Fc region, or FcRn-binding fragment thereof, wherein the Fc domains of the Fc region comprise the amino acids Y, T, E, K, F, and Y at EU positions 252, 254, 256, 433, 434, and 436 respectively, and wherein the Fc region has an increased affinity for an Fc gamma receptor relative to a the affinity of a wild-type IgG1 Fc region for the Fc gamma receptor.
4. The FcRn-antagonist of claim 3, wherein the Fc region is an IgG Fc region.
5. The FcRn-antagonist of claim 3, wherein the Fc region is an IgG1 Fc region.
6. The FcRn-antagonist of claim 3, wherein the Fc region is chimeric Fc region.
7. The FcRn-antagonist of claim 3, wherein the amino acid sequence of the Fc domains of the variant Fc region comprises the amino acid sequence set forth in SEQ ID NO:1, 2, or 3.
8. The FcRn-antagonist of claim 3, wherein the amino acid sequence of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO:1, 2, or 3.
9. The FcRn-antagonist of claim 3, which consists of a variant Fc region, wherein the amino acid sequence of the Fc domains of the variant Fc region consists of the amino acid sequence set forth in SEQ ID NO:1, 2, or 3.

10. The FcRn-antagonist of any one of claims 2-9, wherein the FcRn-antagonist does not comprise an antibody variable region or a CH1 domain.
11. The FcRn-antagonist of any one of claims 1-10, wherein the FcRn-antagonist does not comprise a free cysteine residue.
- 5 12. The FcRn-antagonist of any one claims 1-11, wherein the variant Fc region has increased affinity for CD16a.
13. The FcRn-antagonist of any one of claims 1-12, wherein the Fc domains of the variant Fc region do not comprise an N-linked glycan at EU position 297.
- 10 14. The FcRn-antagonist of any one of claims 1-12, wherein the Fc domains of the variant Fc region comprise an Fc domain having an afucosylated N-linked glycan at EU position 297.
15. The FcRn-antagonist of any one of claims 1-12, wherein the Fc domains of the variant Fc region comprise an N-linked glycan having a bisecting GlcNac at EU position 297 of the Fc domains.
- 15 16. The FcRn-antagonist of any one of the preceding claims, wherein the variant Fc region is linked to a half-life extender.
17. The FcRn-antagonist of claim 16, wherein the half-life extender is polyethylene glycol or human serum albumin.
- 20 18. An FcRn-antagonist composition comprising a plurality of FcRn-antagonist molecules of claim 14, wherein at least 50% (optionally, at least 60, 70, 80, 90, 95, or 99%) of the molecules comprise a variant Fc region, or FcRn-binding fragment thereof, comprising an afucosylated N-linked glycan.
- 25 19. An FcRn-antagonist composition comprising a plurality of FcRn-antagonist molecules of claim 15, wherein at least 50% (optionally, at least 60, 70, 80, 90, 95, or 99%) of the molecules comprise a variant Fc region or FcRn-binding fragment thereof, comprising an N-linked glycan having a bisecting GlcNac.

20. A pharmaceutical composition comprising the FcRn-antagonist or FcRn-antagonist composition of any one of the preceding claims and a pharmaceutically acceptable carrier or excipient.
21. A method of inhibiting FcRn function in a subject, the method comprising administering to the subject an effective amount of the pharmaceutical composition of claim 20.
22. A method of reducing the serum levels of an Fc-containing agent in subject that has been administered the Fc-containing agent, the method comprising simultaneously or sequentially administering to subject an effective amount of the pharmaceutical composition of claim 20.
23. The method of claim 22, wherein the Fc-containing agent is an antibody or immunoadhesin.
24. The method of claim 22, wherein the Fc-containing agent is a therapeutic or diagnostic agent.
25. The method of claim 22, wherein the Fc-containing agent is an imaging agent.
26. The method of claim 22, wherein the Fc-containing agent is an antibody drug conjugate.
27. A method of treating an antibody-mediated disorder in a subject, the method comprising administering to the subject an effective amount of the pharmaceutical composition of claim 20.
28. The method of claim 27, wherein the antibody-mediated disorder is an autoimmune disease.
29. The method of claim 28, wherein the autoimmune disease is selected from the group consisting of allogenic islet graft rejection, alopecia areata, ankylosing spondylitis, antiphospholipid syndrome, autoimmune Addison's disease, Alzheimer's disease, antineutrophil cytoplasmic autoantibodies (ANCA), autoimmune diseases of the adrenal gland, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune myocarditis, autoimmune neutropenia, autoimmune oophoritis and orchitis, autoimmune

thrombocytopenia, autoimmune urticaria, Behcet's disease, bullous pemphigoid, cardiomyopathy, Castleman's syndrome, celiac spruce-dermatitis, chronic fatigue immune disfunction syndrome, chronic inflammatory demyelinating polyneuropathy (CIDP), Churg-Strauss syndrome, cicatrical pemphigoid, CREST syndrome, cold agglutinin disease, Crohn's disease, dermatomyositis, dilated cardiomyopathy, discoid lupus, epidermolysis bullosa acquisita, essential mixed cryoglobulinemia, factor VIII deficiency, fibromyalgia-fibromyositis, glomerulonephritis, Grave's disease, Guillain-Barre, Goodpasture's syndrome, graft-versus-host disease (GVHD), Hashimoto's thyroiditis, hemophilia A, idiopathic membranous neuropathy, idiopathic pulmonary fibrosis, idiopathic thrombocytopenia purpura (ITP), IgA neuropathy, IgM polyneuropathies, immune mediated thrombocytopenia, juvenile arthritis, Kawasaki's disease, lichen plantus, lichen sclerosus, lupus erthematosus, Meniere's disease, mixed connective tissue disease, mucous membrane pemphigoid, multiple sclerosis, type 1 diabetes mellitus, Multifocal motor neuropathy (MMN), myasthenia gravis, paraneoplastic bullous pemphigoid, pemphigoid gestationis, pemphigus vulgaris, pemphigus foliaceus, pernicious anemia, polyarteritis nodosa, polychondritis, polyglandular syndromes, polymyalgia rheumatica, polymyositis and dermatomyositis, primary agammaglobulinemia, primary biliary cirrhosis, psoriasis, psoriatic arthritis, relapsing polychondritis, Reynauld's phenomenon, Reiter's syndrome, rheumatoid arthritis, sarcoidosis, scleroderma, Sjorgen's syndrome, solid organ transplant rejection, stiff-man syndrome, systemic lupus erythematosus, takayasu arteritis, toxic epidermal necrolysis (TEN), Stevens Johnson syndrome (SJS), temporal arteritis/giant cell arteritis, thrombotic thrombocytopenia purpura, ulcerative colitis, uveitis, dermatitis herpetiformis vasculitis, anti-neutrophil cytoplasmic antibody-associated vasculitides, vitiligo, and Wegner's granulomatosis.

30. The method of claim 27, wherein the antibody-mediated disorder is treatable using intravenous immunoglobulin (IVIG), plasmapheresis and/or immunoadsorption.

31. The method of claim 27, wherein the autoimmune disease is an autoimmune channelopathy.

32. The method of claim 31, wherein the channelopathy is selected from the group consisting of autoimmune limbic encephalitis, epilepsy, neuromyelitis optica, Lambert-Eaton myasthenic syndrome, myasthenia gravis, anti- N-Methyl-D-aspartate (NMDA) receptor encephalitis, anti- α -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor encephalitis, Morvan syndrome, neuromyotonia, pediatric autoimmune neuropsychiatric disorders associated with streptococcal infection (PANDAS), and Glycine receptor antibody-associated disorder.
33. The method of claim 27, wherein the antibody-mediated disorder is hyperglobulinemia.
34. The method of claim 27, wherein the FcRn antagonist is administered to the subject simultaneously or sequentially with an additional therapeutic agent.
35. The method of claim 34, wherein the additional therapeutic agent is an anti-inflammatory agent.
36. The method of claim 34, wherein the additional therapeutic agent is a leucocyte depleting agent.
37. The method of claim 36, wherein the leucocyte depleting agent is a B-cell depleting agent.
38. The method of claim 37, wherein the B-cell depleting agent is an antibody.
39. The method of claim 38, wherein the antibody specifically binds to CD10, CD19, CD20, CD21, CD22, CD23, CD24, CD37, CD53, CD70, CD72, CD74, CD75, CD77, CD79a, CD79b, CD80, CD81, CD82, CD83, CD84, CD85, or CD86.
40. The method of claim 34, wherein the additional therapeutic agent is rituximab, daclizumab, basiliximab, muronomab-CD3, infliximab, adalimumab, omalizumab, efalizumab, natalizumab, tocilizumab, eculizumab, golimumab, canakinumab, ustekinumab, belimumab, or a combination thereof.
41. A nucleic acid molecule encoding the FcRn-antagonist of any one of claims 1-19.

- 42. An expression vector comprising the nucleic acid molecule of claim 41.
- 43. A host cell comprising the expression vector of claim 42.
- 44. A method of producing an FcRn-antagonist, comprising culturing the host cell of claim 43 under conditions such that an FcRn-antagonist is expressed.

FIG.1




FIG.2




FIG.3





FIG.4






FIG.6




FIG.7




FIG.8




FIG.9





FIG. 10



INTERNATIONAL SEARCH REPORT

International application No

PCT/US2014/072087

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07K16/00 A61K38/00 A61K38/17
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2013/074598 A1 (MERCK SHARP & DOHME [US]; ZHA DONGXING [US]) 23 May 2013 (2013-05-23) cited in the application figures 1,2; sequence 22 page 33, lines 20-23 page 39, lines 11-13 -----	1-21, 27-44
X	WO 2006/130834 A2 (UNIV TEXAS [US]; WARD E SALLY [US]) 7 December 2006 (2006-12-07) page 17, lines 1-12 figures 3-6; examples -----	3-12,20, 41-44
X	WO 2009/100105 A2 (ATTOGEN INC [US]; CHANG XIAO-JIA [US]; SCHWERTSCHLAG ULLRICH S [US]) 13 August 2009 (2009-08-13) claim 26; figure 13c examples -----	3-9,20, 41-44
	-/-	

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

4 May 2015

Date of mailing of the international search report

12/05/2015

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Covone-van Hees, M

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2014/072087

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CARLOS VACCARO ET AL: "Engineering the Fc region of immunoglobulin G to modulate in vivo antibody levels", NATURE BIOTECHNOLOGY, vol. 23, no. 10, 1 October 2005 (2005-10-01), pages 1283-1288, XP55049342, ISSN: 1087-0156, DOI: 10.1038/nbt1143 the whole document	3-9, 20-27, 41-44
A	----- WO 2010/014909 A1 (SYNTONIX PHARMACEUTICALS INC [US]; MEZO ADAM R [US]; MCDONNELL KEVIN A) 4 February 2010 (2010-02-04) the whole document	1-44
A	----- WO 2009/131702 A2 (DYAX CORP [US]; SYNTONIX PHARMACEUTICALS INC [US]; TENHOOR CHRISTOPHER) 29 October 2009 (2009-10-29) the whole document	1-44
A	----- WO 2010/106180 A2 (LFB BIOTECHNOLOGIES [FR]; BEHRENS CHRISTIAN [FR]; JORIEUX SYLVIE [FR];) 23 September 2010 (2010-09-23) the whole document	1-44
A	----- US 6 277 375 B1 (WARD ELIZABETH SALLY [US]) 21 August 2001 (2001-08-21) the whole document	1-44
A	----- WO 02/060919 A2 (MEDIMMUNE INC [US]; DALL ACQUA WILLIAM [US]; JOHNSON LESLIE S [US]; WA) 8 August 2002 (2002-08-08) the whole document -----	1-44

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2014/072087

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2013074598	A1	23-05-2013	EP 2780463 A1	24-09-2014
			US 2014302028 A1	09-10-2014
			WO 2013074598 A1	23-05-2013

WO 2006130834	A2	07-12-2006	EP 1896503 A2	12-03-2008
			ES 2523666 T3	28-11-2014
			US 2007041907 A1	22-02-2007
			US 2012195892 A1	02-08-2012
			WO 2006130834 A2	07-12-2006

WO 2009100105	A2	13-08-2009	US 2011059091 A1	10-03-2011
			WO 2009100105 A2	13-08-2009

WO 2010014909	A1	04-02-2010	US 2010048488 A1	25-02-2010
			WO 2010014909 A1	04-02-2010

WO 2009131702	A2	29-10-2009	AU 2009238605 A1	29-10-2009
			CA 2722082 A1	29-10-2009
			CN 102149729 A	10-08-2011
			CN 104479017 A	01-04-2015
			EP 2310415 A2	20-04-2011
			JP 2011523351 A	11-08-2011
			KR 20110044729 A	29-04-2011
			NZ 589451 A	27-07-2012
			SG 196839 A1	13-02-2014
			TW 201010720 A	16-03-2010
			US 2009324614 A1	31-12-2009
			US 2013045218 A1	21-02-2013
			US 2013078262 A1	28-03-2013
			US 2014248287 A1	04-09-2014
			WO 2009131702 A2	29-10-2009

WO 2010106180	A2	23-09-2010	AU 2010224800 A1	13-10-2011
			CA 2755905 A1	23-09-2010
			CN 102405231 A	04-04-2012
			EP 2233500 A1	29-09-2010
			EP 2408813 A2	25-01-2012
			EP 2805969 A2	26-11-2014
			JP 2012524522 A	18-10-2012
			KR 20110128883 A	30-11-2011
			US 2012009188 A1	12-01-2012
			WO 2010106180 A2	23-09-2010

US 6277375	B1	21-08-2001	US 6277375 B1	21-08-2001
			US 2002098193 A1	25-07-2002

WO 02060919	A2	08-08-2002	AT 489395 T	15-12-2010
			CA 2431600 A1	08-08-2002
			DK 1355919 T3	14-03-2011
			EP 1355919 A2	29-10-2003
			EP 2341060 A1	06-07-2011
			EP 2354149 A1	10-08-2011
			EP 2357187 A1	17-08-2011
			JP 4336376 B2	30-09-2009
			JP 4336498 B2	30-09-2009
			JP 5265446 B2	14-08-2013
			JP 5444010 B2	19-03-2014
			JP 2005501514 A	20-01-2005

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2014/072087

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		JP 2008054679 A	13-03-2008
		JP 2009261394 A	12-11-2009
		JP 2010154855 A	15-07-2010
		JP 2014050385 A	20-03-2014
		PT 1355919 E	02-03-2011
		US 2003190311 A1	09-10-2003
		US 2006198840 A1	07-09-2006
		US 2007122403 A1	31-05-2007
		US 2010189718 A1	29-07-2010
		US 2011311454 A1	22-12-2011
		US 2013052135 A1	28-02-2013
		US 2013272964 A1	17-10-2013
		US 2014377181 A1	25-12-2014
		WO 02060919 A2	08-08-2002
