

(19) **DANMARK**

(10) **DK/EP 3020727 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 K 14/47 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-07-02**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-04-04**
- (86) Europæisk ansøgning nr.: **15196254.5**
- (86) Europæisk indleveringsdag: **2014-09-15**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-05-18**
- (30) Prioritet: **2013-09-16 EP 13184632** **2013-10-01 EP 13186939**
2013-11-26 US 201361909313 P
- (62) Stamansøgningsnr: **14184835.8**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **CeMM - Forschungszentrum für Molekulare Medizin GmbH, Lazarettgasse 14 / AKH BT 25.3, 1090 Wien, Østrig**
- (72) Opfinder: **Kralovics, Robert, Breitenfeldergasse 13/23, 1080 Wien, Østrig**
KLAMPFL, Thorsten, 41 Malvern Road, Cambridge, CB1 9LH, Storbritannien
Gisslinger, Heinz, Fasangartensiedlung 1/2, 1130 Wien, Østrig
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **Mutant calreticulin til diagnose af myeloide maligniteter**
- (56) Fremdragne publikationer:
WO-A1-2011/069004
M. OLAD NABI ET AL: "Novel mutations in the calreticulin gene core promoter and coding sequence in schizoaffective disorder", AMERICAN JOURNAL OF MEDICAL GENETICS PART B: NEUROPSYCHIATRIC GENETICS, vol. 9999B, 1 January 2009 (2009-01-01), pages n/a-n/a, XP055099786, ISSN: 1552-4841, DOI: 10.1002/ajmg.b.31036
BAXTER E. ET AL.: "Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders", THE LANCET, LANCET LIMITED. LONDON, GB, vol. 365, no. 9464, 19 March 2005 (2005-03-19), pages 1054-1061, XP027764644, ISSN: 0140-6736 [retrieved on 2005-03-19]
SCOTT LINDA M ET AL: "JAK2 exon 12 mutations in polycythemia vera and idiopathic erythrocytosis", NEW ENGLAND JOURNAL OF MEDICINE, MASSACHUSETTS MEDICAL SOCIETY, BOSTON, MA, US, vol. 356, no. 5, 1 February 2007 (2007-02-01), pages 459-468, XP002616834, ISSN: 1533-4406
DATABASE EMBL [Online] 26 May 2000 (2000-05-26), "CM4-NN1011-100300-109-d12 NN1011 Homo sapiens cDNA, mRNA sequence.", XP002719674, retrieved from EBI accession no. EM_EST:AW901361 Database accession no. AW901361
DATABASE EMBL [Online] 25 July 2003 (2003-07-25), "UI-HF-ES0-avy-I-14-0-UI.r1 NIH_MGC_213 Homo sapiens cDNA clone IMAGE:30565045 5', mRNA sequence.", XP002719675, retrieved from EBI accession no. EM_EST:CF129773 Database accession no. CF129773

Fortsættes ...

ELISA RUMI: "Familial chronic myeloproliferative disorders: the state of the art", HEMATOLOGICAL ONCOLOGY, vol. 26, no. 3, 1 September 2008 (2008-09-01), pages 131-138, XP055148140, ISSN: 0278-0232, DOI: 10.1002/hon.863

E. RUMI ET AL: "CALR exon 9 mutations are somatically acquired events in familial cases of essential thrombocythemia or primary myelofibrosis", BLOOD, vol. 123, no. 15, 19 February 2014 (2014-02-19), pages 2416-2419, XP055148139, ISSN: 0006-4971, DOI: 10.1182/blood-2014-01-550434

E. RUMI ET AL: "Clinical effect of driver mutations of JAK2, CALR, or MPL in primary myelofibrosis", BLOOD, vol. 124, no. 7, 1 July 2014 (2014-07-01), pages 1062-1069, XP055148138, ISSN: 0006-4971, DOI: 10.1182/blood-2014-05-578435

J. NANGALIA ET AL: "Somatic CALR Mutations in Myeloproliferative Neoplasms with Nonmutated JAK2", NEW ENGLAND JOURNAL OF MEDICINE, vol. 369, no. 25, 10 December 2013 (2013-12-10), pages 2391-2405, XP055148137, ISSN: 0028-4793, DOI: 10.1056/NEJMoa1312542

BIBI ASIMA ET AL: "Calreticulin is crucial for calcium homeostasis mediated adaptation and survival of thick ascending limb of Henle's loop cells under osmotic stress", INTERNATIONAL JOURNAL OF BIOCHEMISTRY & CELL BIOLOGY, vol. 43, no. 8, August 2011 (2011-08), pages 1187-1197, XP002733583, ISSN: 1357-2725

VANNUCCHI A M ET AL: "Calreticulin mutation-specific immunostaining in myeloproliferative neoplasms: pathogenetic insight and diagnostic value", LEUKEMIA (BASINGSTOKE), vol. 28, no. 9, September 2014 (2014-09), pages 1811-1818, XP002733584,

KLAMPFL THORSTEN ET AL: "Somatic Mutations of Calreticulin in Myeloproliferative Neoplasms", NEW ENGLAND JOURNAL OF MEDICINE, vol. 369, no. 25, December 2013 (2013-12), pages 2379-2390, XP002733585,

CLARK ROBERT A ET AL: "Regulation of calreticulin expression during induction of differentiation in human myeloid cells. Evidence for remodeling of the endoplasmic reticulum", JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 277, no. 35, 30 August 2002 (2002-08-30), pages 32369-32378, XP002733586, ISSN: 0021-9258

DESCRIPTION

[0001] The present invention relates to a method for diagnosing a myeloid malignancy comprising determining the presence of a mutant allele of the calreticulin gene. Also genomic sequences, cDNA sequences, mRNA sequences and protein sequences of the mutant calreticulin are subject of the present invention. Further, the invention relates to medical uses of inhibitors of mutant calreticulin.

[0002] Primary myelofibrosis (PMF), essential thrombocythemia (ET) and polycythemia vera (PV) are monoclonal hematological disorders that belong to the classical BCR-ABL negative myeloproliferative neoplasms (MPN) (Campbell & Green, 2006). Since the 2005 discovery of a somatic mutation in the JAK2 kinase gene, a tremendous progress has been made in molecular diagnosis, clinical management, treatment and molecular understanding of MPN. The valine to phenylalanine (V617F) mutation constitutively activates the Jak2 kinase resulting in increased phosphorylation of its substrates (Stat5, Stat3, Erk, etc.) and leading to increased cytokine responsiveness of myeloid cells (Baxter et al, 2005; James et al, 2005; Kralovics et al, 2005; Levine et al, 2005). Identification of additional mutations soon followed such as in JAK2 exon 12 in PV (Scott et al, 2007) and in the thrombopoietin receptor gene MPL in PMF and ET (Pardanani et al, 2006; Pikman et al, 2006). Although the three MPN disease entities differ in their clinical presentation, they share many molecular as well as clinical features. The JAK2-V617F mutation is present in about 95% of PV cases, 60% PMF and 50% of ET cases, respectively. Mutations in JAK2 exon 12 are specific to about 3% of PV cases whereas MPL mutations are restricted to the PMF (5%) and ET (3%). All three MPN entities are predisposed at a variable degree to thrombosis, bleeding and leukemic transformation (Sverdlow et al, 2008). Although patients may remain in the chronic phase of MPN for several years, disease progression occurs in a form of secondary myelofibrosis in PV and ET, development of accelerated phase with variable degree of pancytopenia followed by leukemic transformation affecting all three MPN entities (Sverdlow et al, 2008).

[0003] Somatic mutations accumulate during the entire clonal evolution of MPN hematopoietic stem cells. These acquired genetic alterations may be point mutations, chromosomal lesions and epigenetic defects and they all may contribute to the fitness of the evolving clone (Klampfl et al, 2011; Kralovics, 2008). These mutations may accelerate proliferation by various means, decrease differentiation potential of progenitors or render them less susceptible to apoptosis. Mutations affecting these mechanisms have been described in genes such as TET2 (Delhommeau et al, 2009), EZH2 (Ernst et al, 2010), DNMT3A (Stegemann et al, 2011), ASXL1 (Stein et al, 2011), and TP53 (Harutyunyan et al, 2011) in different types of myeloid malignancies including MPN (Milosevic & Kralovics, 2013). However, so far only JAK2 and MPL mutations are considered strongly MPN associated and they represent the most useful molecular markers of MPN.

[0004] Despite the progress made in the understanding of the molecular pathogenesis of MPN approximately half of the patients with PMF and ET lack a molecular marker for diagnosis as these patients are negative for both JAK2 and MPL mutations.

[0005] Thus, the technical problem underlying the present invention is the provision of means and methods for diagnosis of a myeloid malignancy.

[0006] Accordingly, the present invention relates to a method for assessing whether a patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy when said one or more mutant alleles of the calreticulin gene is present, wherein said one or more mutant alleles comprises a nucleic acid encoding a mutant calreticulin protein, wherein said mutant calreticulin protein is selected from the group consisting of
 1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;

2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4,
 3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
 4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
 5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
 6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),
- wherein said gene product is protein, and
- wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0007] The technical problem is solved by provision of the embodiments characterized in the claims.

[0008] The present invention solves the above identified technical problem since, as documented herein below and in the appended examples, it was surprisingly found that patients suffering from a myeloid malignancy, preferably primary myelofibrosis (PMF) and essential thrombocytemia (ET), have somatic mutations in the calreticulin (CALR) gene. Another surprising finding was that these myeloid cell specific somatic mutations in the CALR gene in patients with MPN strongly associate with those patients that are negative for both JAK2 and MPL mutations (the previously described disease causing mutations in MPN). As shown herein, CALR mutations are found in 88% of PMF cases, and in 68% of ET cases double negative for JAK2 and MPL. Thus, the present invention provides a reliable diagnosis of myeloid malignancies. The invention is especially useful for patients for which no reliable markers exist, such as patients which are negative for JAK2 and MPL mutations.

[0009] Moreover, it was found herein that the herein provided somatic mutations in the calreticulin (CALR) gene result in a C-terminus of the calreticulin protein which has completely different characteristics compared with the wild type calreticulin protein. It is believed that these different characteristics cause or contribute to the development of myeloid malignancy, preferably primary myelofibrosis (PMF) and essential thrombocytemia (ET).

[0010] All the mutations of CALR identified herein are in the last exon 9 encoding the C-terminal amino acids of the protein and are predominantly insertion/deletion mutations. The majority of the mutations were present in a heterozygous state and they cause a frameshift to an alternative reading frame (alternative frame 1 as shown in Figure 3A). This frameshift results in the replacement of the C-terminal negatively charged amino acids (aspartic and glutamic acid rich) of calreticulin by a predominantly positively charged polypeptide rich in arginine and methionine. In addition, the last 4 amino acids of calreticulin (KDEL (SEQ ID NO: 1331)) contain the endoplasmatic reticulum retention signal. This signal is absent in the mutant calreticulin suggesting that the mutant protein is less represented in the ER compared to the wild type protein. As the negatively charged C-terminus of calreticulin is a low affinity high capacity Ca²⁺ binding domain, it is believed that the Ca²⁺ binding function of the mutant protein is lost. It has been demonstrated herein that the predominant mutations of CALR are type 1 and type 2 mutations as defined herein; see Fig. 3E. These mutants and their use in accordance with the present invention is therefore preferred. Nucleic acid sequences encoding the C-terminus and the amino acid sequence of the C-terminus of type 1 and type 2 CALR mutations are shown in SEQ ID NO: 5 to 12. Further nucleic acids of type 1 and type 2 CALR mutations are disclosed herein.

[0011] The detection of the herein provided CALR mutations at the level of genomic DNA, RNA, cDNA and protein is useful for the diagnosis of a myeloid malignancy, for example, whether a patient has a myeloid malignancy, what type of myeloid malignancy, and specific features of the disease.

[0012] As used herein, "diagnosis" refers, inter alia, to the identification of the nature of an illness or the identification of a physiological or pathophysiological problem underlying a symptom. Thus "diagnosis of a myeloid malignancy" refers to determining (a) if a patient has a myeloid malignancy and/or (b) what *type(s)* of myeloid

malignancy and/or (c) features of the specific myeloid malignancy. Diagnosis can be performed e.g. based on examination of symptoms and/or complementary tests (e.g. cytogenetic or molecular tests).

[0013] The term "assessing whether a patient suffers from a myeloid malignancy" and "diagnosing myeloid malignancy" can be used interchangeably herein. The diagnosis can also comprise or relate to the assessment whether a patient is prone to suffering from a myeloid malignancy, i.e. whether the patient is at risk of developing a myeloid malignancy.

[0014] The present invention relates to a method for assessing whether a patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles comprises a nucleic acid encoding a mutant calreticulin protein, wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4,
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and

wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0015] A sample can be obtained from the patient. "Obtaining" encompasses receipt of a sample that is provided by a third party. For example, blood or bone marrow may be drawn from a patient, placed in appropriate receptacle, and then provided for analysis.

[0016] The present disclosure relates to a method for assessing whether a patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy, said method comprising

- obtaining a sample from said patient;
- determining the presence of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy when said one or more mutant alleles of the calreticulin gene is present.

[0017] In accordance with the present invention, a patient is assessed "positive" for a myeloid malignancy, if one or more mutant alleles of the calreticulin gene are present in a sample, preferably a blood sample, from said patient.

[0018] The term "myeloid malignancy" as used herein refers to clonal haematological diseases affecting the myeloid blood lineages including those with chronic and those with acute clinical course. Myeloid malignancies include myeloproliferative neoplasms, myelodysplastic syndromes and acute myeloid leukemias. It is preferred

herein that the myeloid malignancy is a myeloproliferative neoplasm, particularly primary myelofibrosis (PMF) or essential thrombocythemia (ET), or a myelodysplastic syndrome, particularly refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T).

[0019] Thus, the diagnosis of myeloid malignancy can be to further diagnose subtypes of disease. In further embodiments, the diagnosis utilizes additional tests in combination, such as blood chemistry, cytology, and genetic analysis. Depending on the nature of the myeloproliferative neoplasm, additional diagnostic tests may include red cell mass determination (for polycythemia), bone marrow aspirate and trephine biopsy, arterial oxygen saturation and carboxyhaemoglobin level, neutrophil alkaline phosphatase level, vitamin B12 (or B12 binding capacity) and serum urate. Genetic tests have proven to be increasingly important in diagnosis.

[0020] The following tests are traditionally done to diagnose the following diseases. See e.g. Vardiman, et al. (2009). "The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: Rationale and important changes". Blood 114 (5): 937-51.

Chronic myelogenous leukemia (CML)

[0021] With defining translocation t(9;22);Philadelphia chromosome, BCR-ABL translocation which has three breakpoints:

- u-BCR-ABL (p230): leads to CML with usual neutrophilia and basophilia
- minor-BCR-ABL (p190): leads to CML which has a tendency to become acute lymphoblastic leukemia (ALL) usually precursor B ALL and rarely precursor T ALL
- major-BCR-ABL (p210): normal usual breakpoint

Essential thrombocythemia (ET)

[0022] ET is associated with the JAK2V617F mutation in up to 55% of cases and with an MPL (thrombopoietin receptor) mutation in up to 5% of cases:

- Cellular phase - increased large megakaryocytes with fibrosis and little increase in other bone marrow elements
- Fibrotic phase - collagenous fibrosis with lack of marrow elements

[0023] These disorders are still being revised according to more specific genetic mutations and how often patients end in a fibrotic marrow event.

Polycythemia vera (PV)

[0024] PV is associated most often with the JAK2V617F mutation in greater than 95% of cases, whereas the remainder have a JAK2 exon 12 mutation:

- Cellular phase - increased megakaryocytes which cluster, reticulin fibrosis, later trichrome fibrosis, and increased myeloid and erythroid precursors
- Fibrotic phase - collagenous fibrosis with lack of marrow elements

Primary myelofibrosis (PMF)

[0025] PMF is associated with the JAK2V617F mutation in up to 50% of cases, the JAK2 exon 12 mutations in 1-2% of cases, and the MPL (thrombopoietin receptor) mutation in up to 5% of cases:

- Cellular phase - increased megakaryocytes which cluster, reticulin fibrosis, later trichrome (collagenous) fibrosis, and increased myeloid precursors
- Fibrotic phase - collagenous fibrosis with lack of marrow elements

[0026] **Refractory anemia with ring sideroblasts associated with marked thrombocytosis (RARS-T)** is often considered a myeloid malignancy. Diagnosis of RARS-T may traditionally involve hematology and cytology, analysis of bone marrow, and lack of karyotype abnormalities such as del (5q), t(3;3)(q21;q26) or inv(3)(q21;q26). See Broseus et al. "Clinical features and course of refractory anemia with ring sideroblast associated with marked thrombocytosis" *Haematologica* 9(7): 1036-1041 (2012).

[0027] While the type of myeloid malignancy guides diagnosis and treatment, individual malignancies may have specific mutations that further determine the prognosis and course of treatment. Genetic markers are particularly useful because they often illuminate the underlying pathogenesis of the disease.

[0028] The determination of the presence of one (or more) mutant alleles of the calreticulin gene or of a gene product thereof as described herein can be performed as a stand-alone analysis. Alternatively, this analysis can be followed or preceded by the analysis of other markers for myeloid malignancies, such as JAK2 and MPL mutations. For example, patients suspected to suffer from a myeloid malignancy, such as a myeloproliferative neoplasm (and in particular primary myelofibrosis (PMF) or essential thrombocythemia (ET)), can be tested first for a JAK2 mutation (in particular the V617F mutation). If they are tested negative for the JAK2 mutation they can be tested for mutant calreticulin. If they are then tested negative for mutant calreticulin, they can be tested for MPL mutations, e.g. mutations in exon 10 of the mpl gene. Of course, further markers can also be tested. Also different orders or modes of testing JAK2 mutations, mutant calreticulin and/or MPL mutations and, optionally, further markers are envisaged herein. For example, a positive JAK2 mutation test can be followed by a test for mutant calreticulin (and vice versa) for further diagnosis or prognostic assessment of the myeloid malignancy. Also simultaneous determination of such markers is envisaged, like the simultaneous test for JAK2 mutation(s) and mutant calreticulin (and, optionally, further markers), or the simultaneous test of JAK2 mutation(s), mutant calreticulin and MPL mutation(s) (and, optionally, further markers). Preferably, the patients (or a sample from the patients) suffering from a myeloid malignancy or being prone to suffering from a myeloid malignancy are negative for both JAK2 and MPL mutations, i.e. mutations of JAK2 and MPL are absent in patients assessed to suffer from a myeloid malignancy or being prone to suffering from a myeloid malignancy in accordance with the present invention. In other words, the patients (or a sample from the patients) assessed to suffer from a myeloid malignancy or being prone to suffering from a myeloid malignancy in accordance with the present invention have preferably wild-type JAK2 and MPL present. For further diagnosis, the use of further markers/tests is envisaged. For example, routine bone marrow testing can be used. Such further markers/testing, like bone marrow testing, may be used to validate e.g. a positive mutant calreticulin test or may follow e.g. a negative mutant calreticulin test.

[0029] Wild-type nucleic acid sequences and amino acid sequences of JAK2 and MPL are known and can be deduced from the respective databases, such as NCBI. Exemplary nucleic acid sequences and amino acid sequences of wild-type JAK2 are shown in NM_004972.3 (JAK2 cDNA) and NP_004963.1 (JAK2 protein), respectively. Exemplary nucleic acid sequences and amino acid sequences of wild-type MPL are shown in NM_005373.2 (MPL cDNA) and NP_005364.1 (MPL protein).

[0030] Mutations of JAK2 and MPL in myeloid malignancies have been described herein above. Such mutations are, for example, the V617F mutation of JAK2 (valine to phenylalanine mutation at position 617 of the amino acid sequence of JAK2), mutations in exon 12 of the nucleic acid sequence encoding JAK2 and/or mutations in exon 10

of MPL.

[0031] The valine to phenylalanine (V617F) mutation is disclosed in Baxter et al, 2005; James et al, 2005; Kralovics et al, 2005; Levine et al, 2005). Mutations in JAK2 exon 12 in PV and in the thrombopoietin receptor gene MPL in PMF and ET have been disclosed in Scott et al, 2007 and in Pardanani et al, 2006; Pikman et al, 2006, respectively. The presence of JAK2 and MPL mutations can be excluded by allele specific PCR for JAK2-V617F(ref) and by Sanger sequencing of exon 12 of JAK2 and exon 10 of MPL. An exemplary protocol that can be used in this context is disclosed in Kralovics R, Teo SS, Li S, Theodorides A, Buser AS, Tichelli A, Skoda RC. Acquisition of the V617F mutation of JAK2 is a late genetic event in a subset of patients with myeloproliferative disorders. Blood. 2006 Aug 15;108(4):1377-80. Epub 2006 May 4.

[0032] Accordingly, the present invention provides a novel myeloid malignancy patient group which is assessed to be positive for mutant calreticulin and negative for mutant JAK2 and mutant MPL (or, in other words, the novel myeloid malignancy patient group is assessed to be positive for mutant calreticulin and positive for wild-type JAK2 and wild-type MPL).

[0033] In a preferred embodiment, the methods of the present invention comprise

- a step of determining the presence of a wild type JAK2 protein or a wild type JAK2 nucleic acid in a sample from the patient; and/or
- a step of determining the presence of a wild type MPL protein or a wild type MPL nucleic acid in a sample from the patient.

[0034] In a particularly preferred embodiment, the methods of the present invention comprise

- a step of determining the presence of a wild type JAK2 protein or a wild type JAK2 nucleic acid in a sample from the patient; and
- a step of determining the presence of a wild type MPL protein or a wild type MPL nucleic acid in a sample from the patient.

[0035] The above steps of determining the presence of a wild type JAK2 protein or a wild type JAK2 nucleic acid in a sample from the patient; and/or determining the presence of a wild type MPL protein or a wild type MPL nucleic acid in a sample from the patient can be performed prior to or after the step of determining the presence of a gene product of one or more mutant alleles of the calreticulin gene as defined in the claims in a sample from said patient as provided and defined herein.

[0036] The present invention relates to a method for assessing whether a patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene as defined in the claims in a sample from said patient;
- determining the presence of a wild type JAK2 protein or a wild type JAK2 nucleic acid in a sample from the patient; and
- assessing that said patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy when said one or more mutant alleles of the calreticulin gene is present, wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0037] The present invention relates to a method for assessing whether a patient suffers from a myeloid

malignancy or is prone to suffering from a myeloid malignancy, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene as defined in the claims in a sample from said patient;
- determining the presence of a wild type MPL protein or a wild type MPL nucleic acid in a sample from said patient; and
- assessing that said patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy when said one or more mutant alleles of the calreticulin gene is present, wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0038] The present invention relates to a method for assessing whether a patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene as defined in the claims in a sample from said patient;
- determining the presence of a wild type JAK2 protein or a wild type JAK2 nucleic acid in a sample from said patient;
- determining the presence of a wild type MPL protein or a wild type MPL nucleic acid in a sample from said patient; and
- assessing that said patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy when said one or more mutant alleles of the calreticulin gene is present, wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0039] Preferably, the method of the invention relates solely to the assessment whether a patient suffers from a myeloid malignancy.

[0040] The present invention relates to a method for assessing whether a patient suffers from a myeloid malignancy, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene as defined in the claims in a sample from said patient; and
- assessing that said patient suffers from a myeloid malignancy when said one or more mutant alleles of the calreticulin gene is present, wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0041] The present invention relates to a method for assessing whether a patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene as defined in the claims in a sample from said patient; and
- assessing that said patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis when said one or more mutant alleles of the calreticulin gene is present, wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0042] The present invention relates to a method for assessing whether a patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene as defined in the claims in a sample from said patient; and
- assessing that said patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia when said one or more mutant alleles of the calreticulin gene is present, wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0043] The present invention relates to a method for assessing whether a patient suffers from refractory anemia with ringed sideroblasts and thrombocythemia or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene as defined in the claims in a sample from said patient; and
- assessing that said patient suffers from refractory anemia with ringed sideroblasts and thrombocythemia or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia when said one or more mutant alleles of the calreticulin gene is present, wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0044] The method provided herein comprises determining the presence of preferably solely one mutant allele of the calreticulin gene in a sample from the patient. Preferably, the method is an in vitro method. The herein provided and disclosed mutations of the calreticulin gene are somatic mutations. These mutations can be present in a homozygous state or a heterozygous state, preferably in a heterozygous state.

[0045] The one or more mutant alleles of the calreticulin gene can comprise a nucleic acid encoding a mutant calreticulin protein. The mutant calreticulin proteins disclosed and provided herein are characterized by a common C-terminal amino acid sequence. As it is evident, for example, from Table 2 in the Example, the C-termini of the mutant calreticulin proteins have a common minimum sequence. Said common minimum sequence is shown the amino acid sequence as depicted in SEQ ID NO. 4 and is encoded by nucleic acid molecules having a nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3.

[0046] Accordingly, the mutant calreticulin protein to be used in accordance with the present invention is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0047] In one embodiment, the mutant calreticulin protein to be used in accordance with the present invention is

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3; or
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4.

[0048] The present invention relates to a method for assessing whether a patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

(a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;

(b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4;

(c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;

(d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;

(e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and

(f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0049] The present invention relates to a method for assessing whether a patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;

4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0050] The present invention relates to a method for assessing whether a patient suffers from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T), said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0051] The mutant calreticulin proteins provided and to be used herein have characteristic C-termini, which are shown in SEQ ID NO:s 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, and 144. These C-termini comprise the amino acid sequence as shown in SEQ ID NO: 4.

[0052] The mutant calreticulin protein can, in accordance with the above, be selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, or 143;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136,

140, or 144;

3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0053] The present invention relates to a method for assessing whether a patient suffers from a primary myelofibrosis or is prone to suffering from primary myelofibrosis, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, or 143;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, or 144;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0054] The present invention relates to a method for assessing whether a patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, or 143;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, or 144;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0055] The present invention relates to a method for assessing whether a patient suffers from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T), said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, or 143;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, or 144;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0056] In one embodiment, the mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, or 143; and
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, or 144.

[0057] Herein, 36 types of mutant calreticulin protein have been identified (see Table 2 showing C-termini of the full-length mutant calreticulin proteins). These mutant proteins are unified by their common characteristic C-terminus as shown in SEQ ID NO. 4. The full-length sequences of the mutant calreticulin proteins are shown in SEQ ID NOs: 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, and 288.

[0058] Accordingly, the mutant calreticulin protein provided and to be used herein can be selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, or 287;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, or 288;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0059] The present invention relates to a method for assessing whether a patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, or 287;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, or 288;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0060] The present invention relates to a method for assessing whether a patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, or 287;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, or 288;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of

the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0061] The present invention relates to a method for assessing whether a patient suffers from refractory anemia with ringed sideroblasts and thrombocytopenia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocytopenia (RARS-T), said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from refractory anemia with ringed sideroblasts and thrombocytopenia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocytopenia (RARS-T) when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, or 287;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, or 288;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c);
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0062] In one embodiment, the mutant calreticulin protein provided and to be used herein can be selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, or 287; and
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, or 288.

[0063] It has been shown herein that the identified mutations occur in exon 9 of the calreticulin gene. The following relates therefore to the mutations in the wild-type calreticulin gene and in exon 9 thereof.

The wild-type calreticulin gene is well known. Its nucleic acid sequence and amino acid sequence can be obtained from databases like NCBI under accession number NG_029662.1 (gene) and NP_004334.1 (protein).

An exemplary nucleic acid sequence of the wild-type calreticulin gene is shown in SEQ ID NO: 289. The corresponding amino acid sequence is shown in SEQ ID NO: 290.

[0064] Accordingly, the wild-type calreticulin gene can comprise a sequence selected from the group consisting of:

- a) a nucleic acid encoding a polypeptide comprising an amino acid sequence as depicted in SEQ ID NO: 290;
- (b) a nucleic acid comprising a nucleotide sequence as depicted in SEQ ID NO: 289;
- (c) a nucleic acid hybridizing under stringent conditions to the complementary strand of the nucleic acid as defined in (a) or (b);
- (d) a nucleic acid comprising a nucleotide sequence with at least 70 % identity to the nucleotide sequence of the nucleic acids of any one of (a) to (c); and
- (e) a nucleic acid comprising a nucleotide sequence which is degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid of any one of (a) to (d).

[0065] The one or more mutant allele of the calreticulin gene can be in a region encompassing exon 9 of the above described calreticulin gene. The wild-type nucleic acid sequence of exon 9 of the calreticulin gene is shown in SEQ ID NO:435. The corresponding wild-type amino acid sequence is shown SEQ ID NO:436.

[0066] In accordance with the above, exon 9 of the wild-type calreticulin gene can comprise a sequence selected from the group consisting of:

- a) a nucleic acid encoding a polypeptide comprising an amino acid sequence as depicted in SEQ ID NO:436;
- (b) a nucleic acid comprising a nucleotide sequence as depicted in SEQ ID NO:435;
- (c) a nucleic acid hybridizing under stringent conditions to the complementary strand of the nucleic acid as defined in (a) or (b);
- (d) a nucleic acid comprising a nucleotide sequence with at least 70 % identity to the nucleotide sequence of the nucleic acids of any one of (a) to (c); and
- (e) a nucleic acid comprising a nucleotide sequence which is degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid of any one of (a) to (d).

[0067] As shown herein (see, for example, Table 2), the herein provided mutant alleles of the calreticulin genes have a frameshift mutation compared to the wild-type calreticulin gene. The frameshift mutation can be in exon 9 of the wild-type calreticulin gene. Due to the frameshift mutation, the open reading frame of the wild-type calreticulin gene is no longer used, but an alternative frame 1, which leads to the generation of the characteristic C-terminus of the mutant calreticulin proteins (the common minimum amino acid sequence of the mutant proteins is shown in SEQ ID NO. 4).

[0068] The frameshift mutation can be caused by the deletion of one or more nucleotides, by the insertion of two or more nucleotides or a combination of insertion and deletion of one or more nucleotides, provided that the mutant protein comprises the characteristic C-terminus (as shown in SEQ ID NO: 4) or a fragment thereof.

[0069] For example, the frameshift mutation is (or is caused by) the deletion of one nucleotide from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, or the insertion of two nucleotides into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0070] For example, $(1 + (3 \times n_0))$ nucleotides can be deleted from the calreticulin gene (or from exon 9 thereof), whereby n_0 can be any natural number including zero. Non-limiting examples of the number of nucleotides that can be deleted from the calreticulin gene (or from exon 9 thereof) to generate a nucleic acid encoding the herein provided mutant calreticulin proteins are 1, 4, 19, 22, 31, 34, 46, 52 nucleotides.

[0071] Likewise, the frameshift mutation can be (or can be caused by) the insertion of two nucleotides into the coding sequence of the wild-type calreticulin gene, particularly in exon 9 thereof. Accordingly, $(2 + (3 \times n_0))$ nucleotides can be inserted into the calreticulin gene (or into exon 9 thereof), whereby n_0 can be any natural number including zero. For example, 5 nucleotides can be inserted into the calreticulin gene (or into exon 9 thereof) to generate a nucleic acid encoding the herein provided mutant calreticulin proteins.

[0072] The frameshift mutation can also be caused by a combination of insertion and deletion of one or more nucleotides into/from the wild-type calreticulin gene (or into/from exon 9 thereof), provided that the resulting mutant protein comprises the characteristic C-terminus (as shown in SEQ ID NO: 4) or a fragment thereof.

[0073] For example, the frameshift mutation can be (or can be caused by) the deletion of one nucleotide from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, and by the insertion of six nucleotides into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0074] For example, the frameshift mutation can be (or can be caused by) the deletion of two nucleotides from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, and by the insertion of four nucleotides into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0075] For example, the frameshift mutation can be (or can be caused by) the deletion of three nucleotides from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, and by the insertion of five nucleotides into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0076] For example, the frameshift mutation can be (or can be caused by) the deletion of 12 nucleotides from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, and by the insertion of 5 nucleotides into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0077] For example, the frameshift mutation can be (or can be caused by) the deletion of 18 nucleotides from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, and by the insertion of 11 nucleotides into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0078] For example, the frameshift mutation can be (or can be caused by) the deletion of 18 nucleotides from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, and by the insertion of 14 nucleotides into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0079] For example, the frameshift mutation can be (or can be caused by) the deletion of 20 nucleotides from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, and by the insertion of 1 nucleotide into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0080] For example, the frameshift mutation can be (or can be caused by) the deletion of 28 nucleotides from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, and by the insertion of 6 nucleotides into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0081] For example, the frameshift mutation can be (or can be caused by) the deletion of 35 nucleotides from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, and by the insertion of 1

nucleotide into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0082] For example, the frameshift mutation can be (or can be caused by) the deletion of 36 nucleotides from the coding sequence of the wild-type calreticulin gene, particularly from exon 9 thereof, and by the insertion of 2 nucleotide into the coding sequence of the wild-type calreticulin gene, particularly into exon 9 thereof.

[0083] Further combinations of insertion/deletion inventions that result in the generation of the characteristic C-terminus of the mutant calreticulin proteins (the common minimum amino acid sequence of the mutant proteins is shown in SEQ ID NO. 4) or of a fragment thereof are readily conceivable.

[0084] Due to the above described insertions, deletions and combinations of insertions/deletions, a frameshift is introduced into the (coding sequence of the) wild-type calreticulin gene and particularly in exon 9 thereof. Accordingly, the mutant calreticulin protein disclosed herein and to be used in accordance with the present invention comprises a mutant amino acid stretch encoded by these mutant exon 9 sequences.

[0085] Accordingly, the mutant calreticulin protein can be selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0086] The present invention relates to a method for assessing whether a patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429,

431, 432, or 433;

2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0087] The present invention relates to a method for assessing whether a patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0088] The present invention relates to a method for assessing whether a patient suffers from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T), said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from refractory anemia with ringed sideroblasts and thrombocytopenia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocytopenia (RARS-T) when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein,

wherein said mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is a mutant of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0089] In one embodiment, the mutant calreticulin protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433; and
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434.

[0090] The presence of the one or more mutant alleles of the calreticulin gene is assessed on the protein level.

[0091] If the presence of the one or more mutant alleles of the calreticulin gene is to be assessed on the genomic level, the mutant allele can comprise or consist of DNA, preferably genomic DNA.

[0092] For example, the mutant allele can comprise a nucleic acid selected from the group consisting of

1. (a) a nucleic acid encoding a polypeptide comprising an amino acid sequence as depicted in SEQ ID NO: 4, 8,

- 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288; 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
2. (b) a nucleic acid comprising a nucleotide sequence as depicted in SEQ ID NO: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, 65, 69, 73, 77, 81, 85, 89, 93, 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189, 193, 197, 201, 205, 209, 213, 217, 221, 225, 229, 233, 237, 241, 245, 249, 253, 257, 261, 265, 269, 273, 277, 281, 285, 291, 295, 299, 303, 307, 311, 315, 319, 323, 327, 331, 335, 339, 343, 347, 351, 355, 359, 363, 367, 371, 375, 379, 383, 387, 391, 395, 399, 403, 407, 411, 415, 419, 423, 427, or 431;
 3. (c) a nucleic acid hybridizing under stringent conditions to the complementary strand of the nucleic acid as defined in (a) or (b) and encoding one or more mutant alleles of the calreticulin gene, said allele comprising a frameshift mutation in exon 9 compared to the wild-type calreticulin gene;
 4. (d) a nucleic acid comprising a nucleotide sequence with at least 70 % identity to the nucleotide sequence of the nucleic acids of any one of (a) to (c); and
 5. (e) a nucleic acid comprising a nucleotide sequence which is degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid of any one of (a) to (d).

[0093] The present invention relates to a method for assessing whether a patient suffers from a primary myelofibrosis or is prone to suffering from primary myelofibrosis, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein as defined herein above,

wherein said mutant allele comprises a nucleic acid selected from the group consisting of

1. (a) a nucleic acid encoding a polypeptide comprising an amino acid sequence as depicted in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288; 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
2. (b) a nucleic acid comprising a nucleotide sequence as depicted in SEQ ID NO: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, 65, 69, 73, 77, 81, 85, 89, 93, 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189, 193, 197, 201, 205, 209, 213, 217, 221, 225, 229, 233, 237, 241, 245, 249, 253, 257, 261, 265, 269, 273, 277, 281, 285, 291, 295, 299, 303, 307, 311, 315, 319, 323, 327, 331, 335, 339, 343, 347, 351, 355, 359, 363, 367, 371, 375, 379, 383, 387, 391, 395, 399, 403, 407, 411, 415, 419, 423, 427, or 431;
3. (c) a nucleic acid hybridizing under stringent conditions to the complementary strand of the nucleic acid as defined in (a) or (b) and encoding one or more mutant alleles of the calreticulin gene, said allele comprising a frameshift mutation in exon 9 compared to the wild-type calreticulin gene;
4. (d) a nucleic acid comprising a nucleotide sequence with at least 70 % identity to the nucleotide sequence of the nucleic acids of any one of (a) to (c); and
5. (e) a nucleic acid comprising a nucleotide sequence which is degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid of any one of (a) to (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0094] The present invention relates to a method for assessing whether a patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein as defined herein above,

wherein said mutant allele comprises a nucleic acid selected from the group consisting of

1. (a) a nucleic acid encoding a polypeptide comprising an amino acid sequence as depicted in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288; 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
2. (b) a nucleic acid comprising a nucleotide sequence as depicted in SEQ ID NO: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, 65, 69, 73, 77, 81, 85, 89, 93, 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189, 193, 197, 201, 205, 209, 213, 217, 221, 225, 229, 233, 237, 241, 245, 249, 253, 257, 261, 265, 269, 273, 277, 281, 285, 291, 295, 299, 303, 307, 311, 315, 319, 323, 327, 331, 335, 339, 343, 347, 351, 355, 359, 363, 367, 371, 375, 379, 383, 387, 391, 395, 399, 403, 407, 411, 415, 419, 423, 427, or 431;
3. (c) a nucleic acid hybridizing under stringent conditions to the complementary strand of the nucleic acid as defined in (a) or (b) and encoding one or more mutant alleles of the calreticulin gene, said allele comprising a frameshift mutation in exon 9 compared to the wild-type calreticulin gene;
4. (d) a nucleic acid comprising a nucleotide sequence with at least 70 % identity to the nucleotide sequence of the nucleic acids of any one of (a) to (c); and
5. (e) a nucleic acid comprising a nucleotide sequence which is degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid of any one of (a) to (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0095] The present invention relates to a method for assessing whether a patient suffers from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T), said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) when said one or more mutant alleles of the calreticulin gene is present,

wherein said one or more mutant alleles of the calreticulin gene comprises a nucleic acid encoding a mutant calreticulin protein as defined herein above,

wherein said mutant allele comprises a nucleic acid selected from the group consisting of

1. (a) a nucleic acid encoding a polypeptide comprising an amino acid sequence as depicted in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288; 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370,

- 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
2. (b) a nucleic acid comprising a nucleotide sequence as depicted in SEQ ID NO: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, 65, 69, 73, 77, 81, 85, 89, 93, 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189, 193, 197, 201, 205, 209, 213, 217, 221, 225, 229, 233, 237, 241, 245, 249, 253, 257, 261, 265, 269, 273, 277, 281, 285, 291, 295, 299, 303, 307, 311, 315, 319, 323, 327, 331, 335, 339, 343, 347, 351, 355, 359, 363, 367, 371, 375, 379, 383, 387, 391, 395, 399, 403, 407, 411, 415, 419, 423, 427, or 431;
 3. (c) a nucleic acid hybridizing under stringent conditions to the complementary strand of the nucleic acid as defined in (a) or (b) and encoding one or more mutant alleles of the calreticulin gene, said allele comprising a frameshift mutation in exon 9 compared to the wild-type calreticulin gene;
 4. (d) a nucleic acid comprising a nucleotide sequence with at least 70 % identity to the nucleotide sequence of the nucleic acids of any one of (a) to (c); and
 5. (e) a nucleic acid comprising a nucleotide sequence which is degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid of any one of (a) to (d),

wherein said gene product is protein, and wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0096] Any methods routinely employed for mutational analyses can be used in accordance with the present invention. The presence of the mutant allele on genomic level, can, for example, be determined by sequencing (such as Sanger sequencing e.g. bidirectional Sanger sequencing) and/or PCR-based detection strategies, such as PCR sizing assays (i.e. PCR followed by fragment analysis e.g. via agarose gel electrophoresis (like high-density agarose gel electrophoresis)).

[0097] Detection of a mutation in a nucleic acid can be performed by methods known in the art, including direct sequencing, restriction fragment length polymorphism identification (RFLPI) of genomic DNA, random amplified polymorphic detection (RAPD), amplified fragment length polymorphism detection (AFLPD), polymerase chain reaction (PCR), DNA sequencing, allele specific oligonucleotide (ASO) probes, hybridization to DNA microarrays or beads, high resolution melting (HRM), and TaqMan probe principle. The nucleic acid can be genomic DNA, amplified genomic DNA, mRNA, cDNA, or amplified cDNA.

[0098] Sequencing is typically performed on specifically amplified nucleic acids. Fragment size analysis typically uses differences in sizes of amplicons following PCR. High resolution melting (HRM) detects mutations in DNA by precisely measuring the melting point of double stranded DNA. Gundry et al., "Amplicon Melting Analysis with Labeled Primers: A Closed-Tube Method for Differentiating Homozygotes and Heterozygotes" *Clinical Chemistry* 49: 396-406 (2003). Typically the user will use PCR to amplify the DNA region in which their mutation of interest lies. The amplified DNA is then precisely heated from around 50°C up to around 95°C, until the strands separate. This process is typically monitored with fluorescent dyes.

[0099] One approach that can be employed herein uses fragment size analysis, followed or not by sequencing. As mentioned above, PCR assays using e.g. genomic DNA of mutant calreticulin as template can be used for amplification of the DNA. Subsequently the amplified DNA can be subject to fragment analysis e.g. via agarose gel electrophoresis.

[0100] Methods for determining the presence of the mutant allele on mRNA level or protein level are described further below.

[0101] For mRNA, many of the same methods as used for DNA can be performed after reverse transcription to generate cDNA. Other methods include RealTime PCR, Reverse Transcriptase PCR, Whole Transcriptome Shotgun Sequencing (RNAseq), in situ hybridization or micro-arrays. Real Time PCR simultaneously amplifies and detects a sequence of interest. The use of specific primers and fluorescent labels can distinguish between wild type and mutations.

[0102] Proteins can be analyzed by methods that include immunohistochemistry (IHC), immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or fluorescent activated cell sorting (FACS). Many methods monitor the binding of an antibody or set of antibodies to a protein of interest that detect differences between a wild type and mutant forms. Mass spectrometry detects differences in the size of a protein and its fragments that reveal information about the underlying sequence. For example, polyclonal antibodies that specifically bind to mutant calreticulin protein can be used, as shown in Example 2.

[0103] As described above, the present invention takes advantage of the determination of the presence of a gene product of one or more mutant alleles of the calreticulin gene in order to diagnose myeloid malignancy, wherein the gene product is protein.

[0104] Myeloid malignancies include myeloproliferative neoplasms and myelodysplastic syndromes. It is preferred herein that the myeloid malignancy is a myeloproliferative neoplasm, particularly primary myelofibrosis (PMF) or essential thrombocythemia (ET), or a myelodysplastic syndrome, particularly refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T).

[0105] If the gene product is mRNA, the presence or amount of said mRNA can be determined by routine techniques, such as RealTime PCR, ReverseTranscriptase PCR, Whole Transcriptome Shotgun Sequencing (RNAseq), sanger sequencing, in situ hybridization or micro-arrays.

[0106] Accordingly, the determination by PCR techniques such as RealTime PCR or ReverseTranscriptase PCR can further comprise the steps

1. (i) contacting the nucleic acid in the sample with one or two oligonucleotides; and
2. (ii) generating an amplification product containing the target sequence.

[0107] Exemplary mutation specific probes and primers are provided and used herein.

[0108] Exemplary oligonucleotides (primers) to be used in accordance with the present invention are

Forward: ACAACTTCCTCATCACCAACG (SEQ ID NO: 437)
and/or

Reverse: GGCCTCAGTCCAGCCCTG (SEQ ID NO: 438)

Forward: GGCAAGGCCCTGAGGTGT (SEQ ID NO: 439)
and/or

Reverse: GGCCTCAGTCCAGCCCTG (SEQ ID NO: 438)

[0109] Further suitable mutation specific probes and primers for use in the present invention can, for example, be derived from the cDNA sequences of the mutated calreticulin gene. Such cDNA sequences are provided and described below. Exemplary cDNA sequences that can be used in this context are shown in SEQ ID NO: 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42, 46, 50, 54, 58, 62, 66, 70, 74, 78, 82, 86, 90, 94, 98, 102, 106, 110, 114, 118, 122, 126, 130, 134, 138, 142, 146, 150, 154, 158, 162, 166, 170, 174, 178, 182, 186, 190, 194, 198, 202, 206, 210, 214, 218, 222, 226, 230, 234, 238, 242, 246, 250, 254, 258, 262, 266, 270, 274, 278, 282, or 286; 292, 296, 300, 304, 308, 312, 316, 320, 324, 328, 332, 336, 340, 344, 348, 352, 356, 360, 364, 368, 372, 376, 380, 384, 388, 392, 396, 400, 404, 408, 412, 416, 420, 424, 428, or 432.

[0110] Further exemplary cDNA sequences that can be used for the design of mutation specific probes and primers are depicted in the following table:

Sequences of mutation junctions in the cDNA sequence of CALR for the design of mutation specific probes or PCR

primers.

CALR mutation	cDNA junction sequences in mutated positions
Type 1	GAAGGACAAACAGGACGAGG AG CAGAGGACAAGGAGGATGAT (SEQ ID NO: 440)
Type 2	GAGGAGGAGGCAGAGGACA ATTGTC GGAGGATGATGAGGACAAAG (SEQ ID NO: 441)
Type 3	GGACAAACAGGACGAGGAGC AG AGGCAGAGGACAAGGAGGAT (SEQ ID NO: 442)
Type 4	CAGGACGAGGAGCAGAGGCT TAGG AGGAGGCAGAGGACAAGG (SEQ ID NO: 443)
Type 5	TGAAGGACAAACAGGACGAG GGC CAGAGGACAAGGAGGATGA (SEQ ID NO: 444)
Type 6	AGGACAAACAGGACGAGGAG CG GAGGCAGAGGACAAGGAGGA (SEQ ID NO: 445)
Type 7	CAGGACGAGGAGCAGAGGCT TAGG AGGATGATGAGGACAAAG (SEQ ID NO: 446)
Type 8	GGACGAGGAGCAGAGGCT TAA AGAGGAGGCAGAGGACAAGGAG (SEQ ID NO: 447)
Type 9	CAAGAAACGCAAAGAGGAG AG AGGCAGAGGACAAGGAGGAT (SEQ ID NO: 448)
Type 10	AGGAGGAGGAGGCAGAGGAC ATGTGTC GGAGGATGATGAGGACAAAG (SEQ ID NO: 449)
Type 11	AAGGACAAACAGGACGAGGACCAG AG GCAGAGGACAAGGAGGAT (SEQ ID NO: 450)
Type 12	CAAACAGGACGAGGAGCAGAGGAGGAGGAGGCAGAGGAC (SEQ ID NO: 451)
Type 13	AACAGGACGAGGAGCAGAGGCAGAGGAGGAGGCAGAGGACAAG (SEQ ID NO: 452)
Type 14	ACAGGACGAGGAGCAGAGGCTGAGGAGGAGGCAGAGGACAAG (SEQ ID NO: 453)
Type 15	CAGGACGAGGAGCAGAGGCT TAGG AGGAGGGAGAGGACAAGGAGGATGATG (SEQ ID NO: 454)
Type 16	CAGGACGAGGAGCAGAGGCT TC AGAGGAGGCAGAGGACAAGGAG (SEQ ID NO: 455)
Type 17	GGACGAGGAGCAGAGGCT TAA AGAGGAGGCAG TGG ACAAGGAGGATGATGAGG (SEQ ID NO: 456)
Type 18	GGACGAGGAGCAGAGGCT TAA AGAGGATGATGAGGACAAAGAT (SEQ ID NO: 457)
Type 19	GGAGCAGAGGCT TAA GGAGG AG AGGCAGAGGACAAGGAGGAT (SEQ ID NO: 458)
Type 20	GGCT TAA GGAGGAGGAAGAAGGGAGGAGGCAGAGGACAAGGA (SEQ ID NO: 459)
Type 21	GGCT TAA GGAGGAGGAAGAAG CGTTTAA AGAGGACAAGGAGGATGATGA (SEQ ID NO: 460)
Type 22	CT TAA GGAGGAGGAAGAAGAC CA ACGCAAAGAGGAGGAGGAGG (SEQ ID NO: 461)
Type 23	CT TAA GGAGGAGGAAGAAGACTGCGTGAGGAGGAGGAGGCAGAGGAC (SEQ ID NO: 462)
Type 24	CT TAA GGAGGAGGAAGAAGAC AGG AGGCAGAGGACAAGGAGG (SEQ ID NO: 463)
Type 25	TAAGGAGGAGGAAGAAGAC AAA AGGCAGAGGACAAGGAGGATG (SEQ ID NO: 464)
Type 26	TAAGGAGGAGGAAGAAGAC AAA ACGCAAAGAGGAGGAGGAG (SEQ ID NO: 465)
Type 27	AAGGAGGAGGAAGAAGAC AACTTGTTC GCAAAGAGGAGGAGGAGGCA (SEQ ID NO: 466)
Type 28	GGAAGAAGACAAGAAACGCA AA AGGAGGATGATGAGGACAAA (SEQ ID NO: 467)
Type 29	GAAGACAAGAAACGCAAAGAG CCCTCCTCTTGTCTA AGGAGGATGATGAGGACAAA (SEQ ID NO: 468)
Type 30	AGACAAGAAACGCAAAGAG GAACCATCCTTGTC GGAGGATGATGAGGACAAAGA (SEQ ID NO: 469)
Type 31	AGAGGAGGAGGAGGCAGAGGGCA ATTGTC GGAGGATGATGAGGACAAAG (SEQ ID NO: 470)
Type 32	GAGGAGGAGGAGGCAGAGGACTGTCT GGAGGATGATGAGGACAAAGA (SEQ ID NO: 471)
Type 33	GAGGAGGAGGCAGAGGACAAATGTCGGAGGATGATGAGGACAAAG (SEQ ID NO: 472)

CALR mutation	cDNA junction sequences in mutated positions
	472)
Type 34	AGGAGGAGGAGGCAGAGGAC ACTTGTCT GGAGGATGATGAGGACAAAGA (SEQ ID NO: 473)
Type 35	AGGAGGAGGAGGCAGAGGAC ATTTGTCT GGAGGATGATGAGGACAAAGA (SEQ ID NO: 474)
Type 36	AGGAGGAGGCAGAGGACA ACTTGTCT GGAGGATGATGAGGACAAAGA (SEQ ID NO: 475)

Bold letters indicate the borders of a deletion event; underlined letters indicate inserted sequences; Bold and italic letters indicate single nucleotide variants

[0111] The following relates to embodiments, wherein the gene product is a protein/polypeptide.

[0112] The gene product can comprise a polypeptide selected from the group consisting of

- a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;
- (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4;
- (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
- (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;
- (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
- (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0113] The present invention relates to a method for assessing whether a patient suffers from a primary myelofibrosis or is prone to suffering from primary myelofibrosis, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis when said gene product is present,

wherein said one or more mutant alleles comprises a nucleic acid encoding a mutant calreticulin protein as defined herein above,

wherein said gene product is selected from the group consisting of

- a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;
- (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4;
- (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
- (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;
- (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and

(f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0114] The present invention relates to a method for assessing whether a patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia when said gene product is present,

wherein said one or more mutant alleles comprises a nucleic acid encoding a mutant calreticulin protein as defined herein above,

wherein said gene product is selected from the group consisting of

a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;

(b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4;

(c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;

(d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;

(e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and

(f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0115] The present invention relates to a method for assessing whether a patient suffers from a refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T), said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T) when said gene product is present,

wherein said one or more mutant alleles comprises a nucleic acid encoding a mutant calreticulin protein as defined herein above,

wherein said gene product is selected from the group consisting of

a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;

(b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4;

(c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;

(d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to

the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;

(e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and

(f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0116] The gene product can comprise a polypeptide selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, 143, 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0117] The present invention relates to a method for assessing whether a patient suffers from a primary myelofibrosis or is prone to suffering from primary myelofibrosis, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from primary myelofibrosis or is prone to suffering from primary myelofibrosis when said gene product is present,

wherein said one or more mutant alleles comprises a nucleic acid encoding a mutant calreticulin protein as defined herein above,

wherein said gene product is a polypeptide selected from the group consisting of

(a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, or 143 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433;

(b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;

(c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;

(d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;

(e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and

fg) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0118] The present invention relates to a method for assessing whether a patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia, said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from essential thrombocythemia or is prone to suffering from essential thrombocythemia when said gene product is present,

wherein said one or more mutant alleles comprises a nucleic acid encoding a mutant calreticulin protein as defined herein above,

wherein said gene product is a polypeptide selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, or 143 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170,

- 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
 3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
 4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;
 5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
 6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0119] The present invention relates to a method for assessing whether a patient suffers from a refractory anemia with ringed sideroblasts and thrombocytopenia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocytopenia (RARS-T), said method comprising

- determining the presence of a gene product of one or more mutant alleles of the calreticulin gene in a sample from said patient; and
- assessing that said patient suffers from refractory anemia with ringed sideroblasts and thrombocytopenia (RARS-T) or is prone to suffering from refractory anemia with ringed sideroblasts and thrombocytopenia (RARS-T) when said gene product is present,

wherein said one or more mutant alleles comprises a nucleic acid encoding a mutant calreticulin protein as defined herein above,

wherein said gene product is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, or 143 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368,

369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433;

2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d),

wherein the presence or amount of said protein is determined by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, mass spectrometry, flow cytometry, or FACS.

[0120] If the gene product is protein, the presence or amount of said protein can be determined routine techniques, such as by immunohistochemistry (IHC), by immunoassay, gel- or blot-based methods, IHC, mass spectrometry, flow cytometry, or FACS.

[0121] As the CALR mutations cause a frameshift of the translated polypeptide, a characteristic C-terminal amino acid sequence is present in the mutated calreticulin proteins as described and provided herein. This characteristic amino acid sequence alters the overall charge of the protein. It also alters the migration of the mutated calreticulin during protein electrophoresis. One can take advantage of this difference in charge and/or in migration behaviour in order to determine the presence of a mutated calreticulin protein. For example, antibodies specific to mutant calreticulin protein can be used to identify said mutant protein e.g. by Western immunoblotting. Optionally, also antibodies specific to the wild type calreticulin protein can be used (in addition) as a control. Such antibodies can include polyclonal and monoclonal antibodies which can be prepared by routine techniques.

[0122] Preferably, the patient is a human patient. The patient can be suspected of suffering from a myeloid malignancy or he/she can be suspected of being prone to suffering from a myeloid malignancy.

[0123] The following relates to samples to be used in accordance with the present invention. The sample can be a bone marrow sample, a blood sample or a saliva sample. The sample is preferably a blood sample. The blood sample preferably comprises peripheral granulocytes. The sample can be obtained from a patient by routine techniques, for example, by biopsy.

[0124] The following relates to herein provided mutant calreticulin proteins.

[0125] The present invention relates to a protein selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and

6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0126] The present invention relates to a protein selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, or 143;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, or 144;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0127] The present invention relates to a protein selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, or 287;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, or 288;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0128] The present invention relates to a protein selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373,

- 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, or 434;
 3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted
 4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by said one or more mutant alleles of the calreticulin gene;
 5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
 6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0129] The meaning of the terms "polypeptide", "protein" and "nucleic acid sequence(s)/molecule(s)" are well known in the art and are used accordingly in context of the present invention. For example, "nucleic acid sequence(s)/molecule(s)" as used herein refer(s) to all forms of naturally occurring or recombinantly generated types of nucleic acids and/or nucleic acid sequences/molecules as well as to chemically synthesized nucleic acid sequences/molecules. This term also encompasses nucleic acid analogues and nucleic acid derivatives. The term "nucleic acid sequence(s)/molecule(s)" can refer to deoxyribonucleic acid (DNA) and ribonucleic acid (RNA). The "nucleic acid sequence(s)/molecule(s)" may be made by synthetic chemical methodology known to one of ordinary skill in the art, or by the use of recombinant technology, or may be isolated from natural sources, or by a combination thereof. The DNA and RNA may optionally comprise unnatural nucleotides and may be single or double stranded. "Nucleic acid sequence(s)/molecule(s)" also refers to sense and anti-sense DNA and RNA, that is, a nucleotide sequence which is complementary to a specific sequence of nucleotides in DNA and/or RNA. Furthermore, the term "nucleic acid sequence(s)/molecule(s)" may refer to DNA or RNA or hybrids thereof or any modification thereof that is known in the state of the art (see, e.g., US 5525711, US 4711955, US 5792608 or EP 302175 for examples of modifications). The nucleic acid molecule(s) may be single- or double-stranded, linear or circular, natural or synthetic, and without any size limitation. For instance, the nucleic acid molecule(s) may be genomic DNA, cDNA, mRNA, antisense RNA, or a DNA encoding such RNAs or chimeroplasts (Colestrauss, Science (1996), 1386-1389). Said nucleic acid molecule(s) may be in the form of a plasmid or of viral DNA or RNA. "Nucleic acid sequence(s)/molecule(s)" may also refer to (an) oligonucleotide(s), wherein any of the state of the art modifications such as phosphothioates or peptide nucleic acids (PNA) are included.

[0130] Nucleic acid sequence with a certain level of identity to the herein provided human sequences can be identified by the skilled person using methods known in the art, e.g. by using hybridization assays or by using alignments, either manually or by using computer programs such as those mentioned herein below in connection with the definition of the term "hybridization" and degrees of homology.

[0131] The nucleic acid sequence may be at least 70% identical to the nucleic acid sequence as shown in SEQ ID NO. 1. More preferably, the nucleic acid sequence is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% identical to the nucleic acid sequence as shown in SEQ ID NO. 1, wherein the higher values are preferred. Most preferably, the nucleic acid sequence is at least 99% identical to the nucleic acid sequence as shown in SEQ ID NO. 1.

[0132] The nucleic acid sequence may be at least 70% identical to the nucleic acid sequence as shown in SEQ ID NO. 2. More preferably, the nucleic acid sequence is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% identical to the nucleic acid sequence as shown in SEQ ID NO. 2, wherein the higher values are preferred. Most preferably, the nucleic acid sequence is at least 99% identical to the nucleic acid sequence as shown in SEQ ID NO. 2.

[0133] The nucleic acid sequence may be at least 70% identical to the nucleic acid sequence as shown in SEQ ID

NO. 3. More preferably, the nucleic acid sequence is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% identical to the nucleic acid sequence as shown in SEQ ID NO. 3, wherein the higher values are preferred. Most preferably, the nucleic acid sequence is at least 99% identical to the nucleic acid sequence as shown in SEQ ID NO. 3.

[0134] The nucleic acid sequence may be at least 70% identical to the nucleic acid sequence as shown in SEQ ID NO. 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, 143, 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433.

[0135] More preferably, the nucleic acid sequence is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% identical to the nucleic acid sequence as shown in SEQ ID NOs. 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, 143, 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433, wherein the higher values are preferred.

[0136] Most preferably, the nucleic acid sequence is at least 99% identical to the nucleic acid sequence as shown in SEQ ID NO. 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, 143, 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, or 433.

[0137] Hybridization assays for the characterization of nucleic acids with a certain level of identity to the nucleic acid sequences as provided herein are well known in the art; see e.g. Sambrook, Russell "Molecular Cloning, A Laboratory Manual", Cold Spring Harbor Laboratory, N.Y. (2001); Ausubel, "Current Protocols in Molecular Biology", Green Publishing Associates and Wiley Interscience, N.Y. (1989). The term "hybridization" or "hybridizes" as used herein may relate to hybridizations under stringent or non-stringent conditions. If not further specified, the conditions are preferably non-stringent. Said hybridization conditions may be established according to conventional protocols described, e.g., in Sambrook (2001) loc. cit.; Ausubel (1989) loc. cit., or Higgins and Hames (Eds.) "Nucleic acid hybridization, a practical approach" IRL Press Oxford, Washington DC, (1985). The setting of conditions is well within the skill of the artisan and can be determined according to protocols described in the art. Thus, the detection of only specifically hybridizing sequences will usually require stringent hybridization and washing conditions such as, for example, the highly stringent hybridization conditions of 0.1 x SSC, 0.1% SDS at 65°C or 2 x SSC, 60°C, 0.1 % SDS. Low stringent hybridization conditions for the detection of homologous or not exactly complementary sequences may, for example, be set at 6 x SSC, 1% SDS at 65°C. As is well known, the length of the probe and the composition of the nucleic acid to be determined constitute further parameters of the hybridization conditions. It is envisaged herein that a nucleic acid can be a primer or probe, for example, a nucleic acid hybridizing under stringent conditions to the complementary strand of the nucleic acid of a mutant calreticulin (or of a fragment thereof as defined herein) or of the nucleic acid encoding a mutant calreticulin protein (or encoding the C-terminus thereof) or of exon 9 of the mutant calreticulin and the like as defined and provided herein above. Primers and probes are often in the range of 10-30 nucleotides. Thus, the invention relates to a nucleic acid (like a primer or probe) hybridizing under stringent conditions to the complementary strand of the nucleic acid mutant calreticulin as defined and provided herein above, wherein said hybridizing nucleic acid is smaller than 50, 49, 48, 47, 46, 45, 44, 43, 42, 41, 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, or 20 nucleotides and is larger than 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 nucleotides.

[0138] Preferably, the nucleic acid has a length of 10 to 35 nucleotides, more preferably 15 to 25 nucleotides, particularly preferred a length of 18 to 21, e.g. 18, 19, 20 or 21 nucleotides.

[0139] In accordance with the present invention, the terms "homology" or "percent homology" or "identical" or "percent identity" or "percentage identity" or "sequence identity" in the context of two or more nucleic acid sequences refers to two or more sequences or subsequences that are the same, or that have a specified percentage of nucleotides that are the same (at least 70%, 75%, 80%, 85%, most preferably at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% identity, most preferably at least 99% identity), when compared and aligned for maximum correspondence over a window of comparison (preferably over the full length), or over a designated region as measured using a sequence comparison algorithm as known in the art, or by manual alignment and visual inspection. Sequences having, for example, 75% to 90% or greater sequence identity may be considered to be substantially identical. Such a definition also applies to the complement of a test sequence. Preferably the described identity exists over a region that is at least about 15 to 25 nucleotides in length, more preferably, over a region that is at least about 50 to 100 nucleotides in length and most preferably over the full length. Those having skill in the art will know how to determine percent identity between/among sequences using, for example, algorithms such as those based on CLUSTALW computer program (Thompson Nucl. Acids Res. 2 (1994), 4673-4680) or FASTDB (Brutlag Comp. App. Biosci. 6 (1990), 237-245), as known in the art.

[0140] Although the FASTDB algorithm typically does not consider internal non-matching deletions or additions in sequences, i.e., gaps, in its calculation, this can be corrected manually to avoid an overestimation of the % identity. CLUSTALW, however, does take sequence gaps into account in its identity calculations. Also available to those having skill in this art are the BLAST and BLAST 2.0 algorithms (Altschul, (1997) Nucl. Acids Res. 25:3389-3402; Altschul (1993) J. Mol. Evol. 36:290-300; Altschul (1990) J. Mol. Biol. 215:403-410). The BLASTN program for nucleic acid sequences uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=4, and a comparison of both strands. The BLOSUM62 scoring matrix (Henikoff (1989) PNAS 89:10915) uses alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

[0141] In order to determine whether a nucleotide residue in a nucleic acid sequence corresponds to a certain position in the nucleotide sequence of e.g. SEQ ID NOs: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63,

65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, 143, 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, and 433, respectively, the skilled person can use means and methods well-known in the art, e.g., alignments, either manually or by using computer programs such as those mentioned herein. For example, BLAST 2.0, which stands for Basic Local Alignment Search Tool BLAST (Altschul (1997), loc. cit.; Altschul (1993), loc. cit.; Altschul (1990), loc. cit.), can be used to search for local sequence alignments. BLAST, as discussed above, produces alignments of nucleotide sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST is especially useful in determining exact matches or in identifying similar sequences. The fundamental unit of BLAST algorithm output is the High-scoring Segment Pair (HSP). An HSP consists of two sequence fragments of arbitrary but equal lengths whose alignment is locally maximal and for which the alignment score meets or exceeds a threshold or cut-off score set by the user. The BLAST approach is to look for HSPs between a query sequence and a database sequence, to evaluate the statistical significance of any matches found, and to report only those matches which satisfy the user-selected threshold of significance. The parameter E establishes the statistically significant threshold for reporting database sequence matches. E is interpreted as the upper bound of the expected frequency of chance occurrence of an HSP (or set of HSPs) within the context of the entire database search. Any database sequence whose match satisfies E is reported in the program output.

[0142] Analogous computer techniques using BLAST (Altschul (1997), loc. cit.; Altschul (1993), loc. cit.; Altschul (1990), loc. cit.) are used to search for identical or related molecules in nucleotide databases such as GenBank or EMBL. This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

and it takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error; and at 70, the match will be exact. Similar molecules are usually identified by selecting those, which show product scores between 15 and 40, although lower scores may identify related molecules. Another example for a program capable of generating sequence alignments is the CLUSTALW computer program (Thompson (1994) Nucl. Acids Res. 2:4673-4680) or FASTDB (Brutlag (1990) Comp. App. Biosci. 6:237-245), as known in the art.

[0143] The explanations and definitions given herein above in respect of "homology/identity of nucleic acid sequences" apply, mutatis mutandis, to "amino acid sequences" of the herein provided mutant calreticulin proteins as depicted in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434, respectively, as explained below.

[0144] The polypeptide to be used in accordance with the present invention may have at least 70 % identity/similarity to the proteins having the amino acid sequence as, for example, depicted in SEQ ID NO: 4, respectively. More preferably, the polypeptide has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% identity/similarity to the proteins depicted in SEQ ID NO: 4, respectively, wherein the higher values are preferred. Most preferably, the polypeptide has at least 99% homology to the protein as depicted in 4.

[0145] The polypeptide to be used in accordance with the present invention may have at least 70 % identity/similarity to the proteins having the amino acid sequence as, for example, depicted in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434, respectively. More preferably, the polypeptide has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% or 98% identity/similarity to the proteins depicted in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434, respectively, wherein the higher values are preferred. Most preferably, the polypeptide has at least 99% homology to the protein as depicted in 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434, respectively.

[0146] Without deferring from the gist of the present invention also (a) (functional) fragment(s) or (a) (functional) derivative(s) of the herein provided polypeptides or proteins can be used, for example, (functional) fragment(s) or (functional) derivative(s) of the minimum C-terminus of the mutant calreticulin as shown in SEQ ID NO. 4. Also (a) (functional) fragment(s) or (a) (functional) derivative(s) of further herein provided mutant calreticulin polypeptides or proteins can be used, for example, (functional) fragment(s) or (functional) derivative(s) of the polypeptide(s) as shown in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434, respectively.

[0147] Thus, a (functional) fragment of the above polypeptide(s)/protein(s) provided herein and to be used in accordance with the present invention can be any of the above specific polypeptides as shown in any one of SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434, respectively, wherein one to 10 amino acids are deleted.

[0148] A (functional) derivative(s) of the above polypeptide(s)/protein(s) provided herein and to be used in accordance with the present invention can be any of the above specific polypeptides as shown in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434, respectively, wherein one to 10 amino acids are inserted, added or substituted.

[0149] Preferably, the deletion, insertion, addition and/or substitution of one to 10 amino acids is within the C-terminus of the herein provided mutant calreticulin, i.e. within the amino acid sequence of the polypeptide as shown in SEQ ID NOs: 4.

[0150] Preferably, the deletion, insertion, addition and/or substitution of one to 10 amino acids is within the C-terminus of the herein provided mutant calreticulin, i.e. within the amino acid sequence of the polypeptides as shown in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, respectively.

[0151] 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids can be deleted, inserted, added or substituted preferably within the C-terminus of the herein provided mutant calreticulin, i.e. within the amino acid sequence of the polypeptides as shown in SEQ ID NOs: 4.

[0152] 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids can be deleted preferably from the C-terminus of the herein provided mutant calreticulin, i.e. from the amino acid sequence of the polypeptide as shown in SEQ ID NOs: 4.

[0153] 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids can be deleted, inserted, added and/or substituted preferably within the C-terminus of the herein provided mutant calreticulin, i.e. within the amino acid sequence of the polypeptides as shown in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, respectively.

[0154] The term "one to 10 amino acids deleted" relates to (functional) fragments of the specific mutant calreticulin proteins provided herein.

[0155] A preferred (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention consists of from 15 to 25 contiguous amino acids. Accordingly, a (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention preferably consists of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 contiguous amino acids.

[0156] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention preferably consists of from 15 to 25 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 4.

[0157] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention preferably consists of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 4.

[0158] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention preferably consists of from 15 to 25 contiguous amino acids of the polypeptides as shown in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, respectively.

[0159] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention preferably consists of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 contiguous amino acids of the polypeptides as shown in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, respectively.

[0160] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41 and up to 42 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 4.

[0161] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42 and up to 43 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 32 or 112.

[0162] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance

with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43 and up to 44 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 8, 128, 132 or 144.

[0163] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, and up to 45 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 12, 44, 136 or 140.

[0164] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45 and up to 46 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 16 or 124.

[0165] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, and up to 47 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 24, 40, 76, 100, or 120.

[0166] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, and up to 48 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 28, 36, 72, 84, 96 or 116.

[0167] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 and up to 49 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 20, 48, 60, 64, 68, or 80.

[0168] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 and up to 50 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 52 or 56.

[0169] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51 and up to 52 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 92.

[0170] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 and up to 53 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 88 or 104.

[0171] A (functional) fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention can consist of at least 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53 and up to 54 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 108.

[0172] The fragment or derivative preferably has the same (or essentially the same) biological activity as the full length polypeptide from which it is derived, the full length polypeptide having the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294,

298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434. In this sense, the fragment or derivative is a "functional" fragment or derivative to be used herein.

[0173] The herein provided polypeptide (as shown, for example, in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434, respectively) may have one to 10 amino acids deleted, inserted, added and/or substituted provided that the polypeptide maintains essentially the biological activity which is characteristic of the polypeptides from which it is derived.

[0174] Preferably, any such deletions, insertions, additions and/or substitutions (in this context particularly substitutions) are conservative, i.e. amino acids are substituted by amino acids having the same or similar characteristics. For example, a hydrophobic amino acid will preferably be substituted by another hydrophobic amino acid and so on.

[0175] The "biological activity" characteristic of the herein provided polypeptides can be considered as an activity which is causative for the (development) of a myeloid malignancy as defined herein, such as a myeloproliferative neoplasm (particularly primary myelofibrosis and essential thrombocytemia).

[0176] The present invention also provides for inhibitors of mutant calreticulin. These inhibitors can be used as a medicament.

[0177] The term "antagonist of mutant calreticulin" or "inhibitor of mutant calreticulin" means in context of the present invention a compound capable of fully or partially preventing or reducing the physiologic activity and/or expression level of (a) mutant calreticulin. The terms "antagonist" or "inhibitor" are used interchangeably herein.

[0178] In the context of the present invention said antagonist may, therefore, prevent, reduce, inhibit or inactivate the physiological activity of a mutant calreticulin upon binding of said compound/substance (i.e. antagonist/inhibitor) to said mutant calreticulin. As used herein, the term "antagonist" also encompasses competitive antagonists, (reversible) non-competitive antagonists or irreversible antagonist, as described, inter alia, in Mutschler, "Arzneimittelwirkungen" (1986), Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, Germany. Such an inhibition can be measured by determining substrate turnover.

[0179] An "antagonist" or "inhibitor" of a mutant calreticulin may also be capable of preventing the function of a mutant calreticulin by preventing/reducing the expression of the nucleic acid molecule encoding for said mutant calreticulin. Thus, an antagonist/inhibitor of a mutant calreticulin may lead to a decreased expression level of the mutant calreticulin (e.g. decreased level of an mutant calreticulin mRNA and/or of mutant calreticulin protein); this may be reflected in a decreased mutant calreticulin activity. The decreased activity and/or expression level can be measured/detected by known methods which are also described herein.

[0180] An "antagonist/inhibitor of a mutant calreticulin" may, for example, interfere with transcription of (an) mutant calreticulin gene(s), processing (e.g. splicing, export from the nucleus and the like) of the gene product(s) (e.g. unspliced or partially spliced mRNA) and/or translation of the gene product (e.g. mature mRNA). The "antagonist/inhibitor of a mutant calreticulin" may also interfere with further modification (like glycosylation or phosphorylation) of the polypeptide/protein encoded by the mutant calreticulin gene(s) and thus completely or partially inhibit the activity of the a mutant calreticulin protein(s) as described herein above. Furthermore, the "antagonist/inhibitor of a mutant calreticulin" may interfere with interactions of the mutant calreticulin protein(s) with other proteins (thus, for example, interfering with the activity of complexes involving mutant calreticulin protein(s)) or, in general, with its synthesis, e.g. by interfering with upstream steps of mutant calreticulin expression or with signalling pathways in which the mutant calreticulin is involved. Depending on the mode of action, such antagonists may, for example, be denoted "sequestering antagonists" or "signalling antagonists".

[0181] In sum, the herein described mutant calreticulin antagonist/inhibitor will, accordingly, lead to a decrease or reduction of mutant calreticulin expression level and/or activity, and thereby reduce its contribution to the development or proliferation of a myeloid malignancy as defined herein.

[0182] The antagonist(s) may be shRNA (small hairpin RNA), siRNA (small interfering RNA), miRNA (microRNA), dsRNA (double stranded RNA), stRNA (small temporal RNA), antisense molecules, extracellular binding-partners, small (binding) molecules, aptamers, intramers, or antibody molecules such as a full antibody (immunoglobulin), a F(ab)-fragment, a F(ab)₂-fragment, a single-chain antibody, a chimeric antibody, a CDR-grafted antibody, a bivalent antibody-construct, a synthetic antibody, a bispecific single chain antibody or a cross-cloned antibody. The present invention relates to an siRNA or shRNA specifically targeting the nucleic acid encoding the mutant calreticulin protein(s), whereby the nucleic is especially mRNA as defined herein.

[0183] Up to 10 % of the contiguous bases of the herein provided siRNAs or shRNAs can be non-complementary. The siRNA can further comprise at least one base at the 5' end and/or at least one base at the 3' end.

[0184] Antagonist(s)/inhibitor(s) which are nucleic acids, such as siRNAs, shRNAs, antisense molecules and the like can readily be prepared by known techniques using, for example, the following target sequences. For example, siRNAs, shRNAs and the like to be employed herein can comprise or consist of an RNA sequence corresponding to one of the target sequences below. The term "RNA sequence corresponding to" means in this context that the RNA sequence is identical to one of the target sequences below with the exception that the thymidine (T) residues of the target sequence is replaced by a uracil (U) residue. The siRNA can consist of a nucleic acid molecule comprising at least ten contiguous bases. For example, the siRNA, shRNA and the like can comprise at least ten contiguous bases of an RNA sequence corresponding to one of the target sequences below as defined above. The siRNA, shRNA and the like can consist of ten contiguous bases of an RNA sequence corresponding to one of the target sequences below as defined above.

The siRNA, shRNA and the like can target one of the target sequences below. These sequences relate to SEQ ID NO: 476 to SEQ ID NO: 1309.

Type of CALR mutation	shRNA sequences covering the mutation site
Type 1	AAGGACAAACAGGACGAGGAG AGGACAAACAGGACGAGGAGC GGACAAACAGGACGAGGAGCA GACAAACAGGACGAGGAGCAG ACAAACAGGACGAGGAGCAGG CAAACAGGACGAGGAGCAGAG AAACAGGACGAGGAGCAGAGG AACAGGACGAGGAGCAGAGGA ACAGGACGAGGAGCAGAGGAC CAGGACGAGGAGCAGAGGACA AGGACGAGGAGCAGAGGACAA GGACGAGGAGCAGAGGACAAG GACGAGGAGCAGAGGACAAGG ACGAGGAGCAGAGGACAAGGA CGAGGAGCAGAGGACAAGGAG GAGGAGCAGAGGACAAGGAGG AGGAGCAGAGGACAAGGAGGA GGAGCAGAGGACAAGGAGGAT GAGCAGAGGACAAGGAGGATG AGCAGAGGACAAGGAGGATGA
Type 2	GAGGAGGAGGCAGAGGACAAT AGGAGGAGGCAGAGGACAATT GGAGGAGGCAGAGGACAATTG GAGGAGGCAGAGGACAATTGT AGGAGGCAGAGGACAATTGTC GGAGGCAGAGGACAATTGTCC

Type of CALR mutation	shRNA sequences covering the mutation site
	GAGGCAGAGGACAATTGTCGG AGGCAGAGGACAATTGTCGGA GGCAGAGGACAATTGTCGGAG GCAGAGGACAATTGTCGGAGG CAGAGGACAATTGTCGGAGGA AGAGGACAATTGTCGGAGGAT GAGGACAATTGTCGGAGGATG AGGACAATTGTCGGAGGATGA GGACAATTGTCGGAGGATGAT GACAATTGTCGGAGGATGATG ACAATTGTCGGAGGATGATGA CAATTGTCGGAGGATGATGAG AATTGTCGGAGGATGATGAGG ATTGTCGGAGGATGATGAGGA TTGTCGGAGGATGATGAGGAC TGTCGGAGGATGATGAGGACA GTCGGAGGATGATGAGGACAA TCGGAGGATGATGAGGACAAA CGGAGGATGATGAGGACAAAG
Type 3	GACAAACAGGACGAGGAGCAG ACAAACAGGACGAGGAGCAGA CAAACAGGACGAGGAGCAGAG AAACAGGACGAGGAGCAGAGG AACAGGACGAGGAGCAGAGGC ACAGGACGAGGAGCAGAGGCA CAGGACGAGGAGCAGAGGCAG AGGACGAGGAGCAGAGGCAGA GGACGAGGAGCAGAGGCAGAG GACGAGGAGCAGAGGCAGAGG ACGAGGAGCAGAGGCAGAGGA CGAGGAGCAGAGGCAGAGGAC GAGGAGCAGAGGCAGAGGACA AGGAGCAGAGGCAGAGGACAA GGAGCAGAGGCAGAGGACAAG GAGCAGAGGCAGAGGACAAGG AGCAGAGGCAGAGGACAAGGA GCAGAGGCAGAGGACAAGGAG CAGAGGCAGAGGACAAGGAGG AGAGGCAGAGGACAAGGAGGA
Type 4	AGGACGAGGAGCAGAGGCTTA GGACGAGGAGCAGAGGCTTAG GACGAGGAGCAGAGGCTTAGG ACGAGGAGCAGAGGCTTAGGA
	CGAGGAGCAGAGGCTTAGGAG GAGGAGCAGAGGCTTAGGAGG AGGAGCAGAGGCTTAGGAGGA GGAGCAGAGGCTTAGGAGGAG GAGCAGAGGCTTAGGAGGAGG AGCAGAGGCTTAGGAGGAGGC GCAGAGGCTTAGGAGGAGGCA CAGAGGCTTAGGAGGAGGCAG AGAGGCTTAGGAGGAGGCAGA GAGGCTTAGGAGGAGGCAGAG

Type of CALR mutation	shRNA sequences covering the mutation site
	AGGCTTAGGAGGAGGCAGAGG GGCTTAGGAGGAGGCAGAGGA GCTTAGGAGGAGGCAGAGGAC CTTAGGAGGAGGCAGAGGACA TTAGGAGGAGGCAGAGGACAA TAGGAGGAGGCAGAGGACAAG
Type 5	GAAGGACAAACAGGACGAGGG AAGGACAAACAGGACGAGGGG AGGACAAACAGGACGAGGGGC GGACAAACAGGACGAGGGGCA GACAAACAGGACGAGGGGCAG ACAAACAGGACGAGGGGCAGA CAAACAGGACGAGGGGCAGAG AAACAGGACGAGGGGCAGAGG AACAGGACGAGGGGCAGAGGA ACAGGACGAGGGGCAGAGGAC CAGGACGAGGGGCAGAGGACA AGGACGAGGGGCAGAGGACAA GGACGAGGGGCAGAGGACAAG GACGAGGGGCAGAGGACAAGG ACGAGGGGCAGAGGACAAGGA CGAGGGGCAGAGGACAAGGAG GAGGGGCAGAGGACAAGGAGG AGGGGCAGAGGACAAGGAGGA GGGGCAGAGGACAAGGAGGAT GGGCAGAGGACAAGGAGGATG
Type 6	GGACAAACAGGACGAGGAGCG GACAAACAGGACGAGGAGCGG ACAAACAGGACGAGGAGCGGA CAAACAGGACGAGGAGCGGAG AAACAGGACGAGGAGCGGAGG AACAGGACGAGGAGCGGAGGC ACAGGACGAGGAGCGGAGGCA
	CAGGACGAGGAGCGGAGGCAG AGGACGAGGAGCGGAGGCAGA GGACGAGGAGCGGAGGCAGAG GACGAGGAGCGGAGGCAGAGG ACGAGGAGCGGAGGCAGAGGA CGAGGAGCGGAGGCAGAGGAC GAGGAGCGGAGGCAGAGGACA AGGAGCGGAGGCAGAGGACAA GGAGCGGAGGCAGAGGACAAG GAGCGGAGGCAGAGGACAAGG AGCGGAGGCAGAGGACAAGGA GCGGAGGCAGAGGACAAGGAG CGGAGGCAGAGGACAAGGAGG
Type 7	AGGACGAGGAGCAGAGGCTTA GGACGAGGAGCAGAGGCTTAG GACGAGGAGCAGAGGCTTAGG ACGAGGAGCAGAGGCTTAGGA CGAGGAGCAGAGGCTTAGGAG GAGGAGCAGAGGCTTAGGAGG

Type of CALR mutation	shRNA sequences covering the mutation site
	AGGAGCAGAGGCTTAGGAGGA GGAGCAGAGGCTTAGGAGGAT GAGCAGAGGCTTAGGAGGATG AGCAGAGGCTTAGGAGGATGA GCAGAGGCTTAGGAGGATGAT CAGAGGCTTAGGAGGATGATG AGAGGCTTAGGAGGATGATGA GAGGCTTAGGAGGATGATGAG AGGCTTAGGAGGATGATGAGG GGCTTAGGAGGATGATGAGGA GCTTAGGAGGATGATGAGGAC CTTAGGAGGATGATGAGGACA TTAGGAGGATGATGAGGACAA TAGGAGGATGATGAGGACAAA
Type 8	GACGAGGAGCAGAGGCTTAAG ACGAGGAGCAGAGGCTTAAGA CGAGGAGCAGAGGCTTAAGAG GAGGAGCAGAGGCTTAAGAGG AGGAGCAGAGGCTTAAGAGGA GGAGCAGAGGCTTAAGAGGAG GAGCAGAGGCTTAAGAGGAGG AGCAGAGGCTTAAGAGGAGGC GCAGAGGCTTAAGAGGAGGCA CAGAGGCTTAAGAGGAGGCAG
	AGAGGCTTAAGAGGAGGCAGA GAGGCTTAAGAGGAGGCAGAG AGGCTTAAGAGGAGGCAGAGG GGCTTAAGAGGAGGCAGAGGA GCTTAAGAGGAGGCAGAGGAC CTTAAGAGGAGGCAGAGGACA TTAAGAGGAGGCAGAGGACAA TAAGAGGAGGCAGAGGACAAG AAGAGGAGGCAGAGGACAAGG AGAGGAGGCAGAGGACAAGGA
Type 9	AAGAAACGCAAAGAGGAGGAG AGAAACGCAAAGAGGAGGAGA GAAACGCAAAGAGGAGGAGAG AAACGCAAAGAGGAGGAGAGG AACGCAAAGAGGAGGAGAGGC ACGCAAAGAGGAGGAGAGGCA CGCAAAGAGGAGCAGAGGCAG GCAAAGAGGAGGAGAGGCAGA CAAAGAGCAGGAGAGGCAGAG AAAGAGGAGGAGAGGCAGAGG AAGAGGAGGAGAGGCAGAGGA AGAGCAGGAGAGGCAGAGGAC GAGGAGGAGAGGCAGAGGACA AGGAGGAGAGGCAGAGGACAA GGAGCAGAGGCAGAGGACAAG GAGGAGAGGCAGAGGACAAGG AGGAGAGGCAGAGGACAAGGA GGAGAGGCAGAGGACAAGGAG GAGAGGCAGAGGACAAGGAGG AGAGGCAGAGGACAAGGAGGA

Type of CALR mutation	shRNA sequences covering the mutation site
Type 10	GGAGGAGGAGGCAGAGGACAT GAGGAGGAGGCAGAGGACATG AGGAGGAGGCAGAGGACATGT GGAGGAGGCAGAGGACATGTG GAGGAGGCAGAGGACATGTGT AGGAGGCAGAGGACATGTGTC GGAGGCAGAGGACATGTGTCCG GAGGCAGAGGACATGTGTCCG AGGCAGAGGACATGTGTCCGA GGCAGAGGACATGTGTCCGAG GCAGAGGACATGTGTCCGAGG CAGAGGACATGTGTCCGAGGA AGAGGACATGTGTCCGAGGAT
	GAGGACATGTGTCCGAGGATG AGGACATGTGTCCGAGGATGA GGACATGTGTCCGAGGATGAT GACATGTGTCCGAGGATGATG ACATGTGTCCGAGGATGATGA CATGTGTCCGAGGATGATGAG ATGTGTCCGAGGATGATGAGG TGTGTCCGAGGATGATGAGGA GTGTCCGAGGATGATGAGGAC TGTCGGAGGATGATGAGGACA GTCGGAGGATGATGAGGACAA TCGGAGGATGATGAGGACAAA CGGAGGATGATGAGGACAAAG
Type 11	AAGGACAAACAGGACGAGGAC AGGACAAACAGGACGAGGACC GGACAAACAGGACGAGGACCA GACAAACAGGACGAGGACCAG ACAAACAGGACGAGGACCAGA CAAACAGGACGAGGACCAGAG AAACAGGACGAGGACCAGAGG AACAGGACGAGGACCAGAGGC ACAGGACGAGGACCAGAGGCA CAGGACGAGGACCAGAGGCAG AGGACGAGGACCAGAGGCAGA GGACGAGGACCAGAGGCAGAG GACGAGGACCAGAGGCAGAGG ACGAGGACCAGAGGCAGAGGA CGAGGACCAGAGGCAGAGGAC GAGGACCAGAGGCAGAGGACA AGGACCAGAGGCAGAGGACAA GGACCAGAGGCAGAGGACAAG GACCAGAGGCAGAGGACAAGG ACCAGAGGCAGAGGACAAGGA CCAGAGGCAGAGGACAAGGAG CAGAGGCAGAGGACAAGGAGG AGAGGCAGAGGACAAGGAGGA
Type 12	AAACAGGACGAGGAGCAGAGG AACAGGACGAGGAGCAGAGGA ACAGGACGAGGAGCAGAGGAG

Type of CALR mutation	shRNA sequences covering the mutation site
	CAGGACGAGGAGCAGAGGAGG AGGACGAGGAGCAGAGGAGGA GGACGAGGAGCAGAGGAGGAG GACGAGGAGCAGAGGAGGAGG
	ACGAGGAGCAGAGGAGGAGGA CGAGGAGCAGAGGAGGAGGAG GAGGAGCAGAGGAGGAGGAGG AGGACGAGGAGGAGGAGGAGGA GGAGCAGAGGAGGAGGAGGAG GAGCAGAGGAGGAGGAGGAGG AGCAGAGGAGGAGGAGGAGGC GCAGAGGAGGAGGAGGAGGCA CAGAGGAGGAGGAGGAGGCG AGAGGAGGAGGAGGAGGCGA GAGGAGGAGGAGGAGGCGAG AGGAGGAGGAGGAGGCGAGG GGAGGAGGAGGAGGCGAGGA
Type 13	ACAGGACGAGGAGCAGAGGCA CAGGACGAGGAGCAGAGGCAG AGGACGAGGAGCAGAGGCAGA GGACGAGGAGCAGAGGCAGAG GACGAGGAGCAGAGGCAGAGG ACCAGGAGCAGAGGCAGAGGA CCAGGAGCAGAGGCAGAGGAG GAGGAGCAGAGGCAGAGGAGG AGGAGCAGAGGCAGAGGAGGA GGAGCAGAGGCAGAGGAGGAG GAGCAGAGGCAGAGGAGGAGG AGCAGAGGCAGAGGAGGAGGC GCAGAGGCAGAGGAGGAGGCA CAGAGGCAGAGGAGGAGGCAG AGAGGCAGAGGAGGAGGCAGA GAGGCAGAGGAGGAGGCAGAG AGGCAGAGGAGGAGGCAGAGG GGCAGAGGAGGAGGCAGAGGA GCAGAGGAGGAGGCAGAGGAC CAGAGGAGGAGGCAGAGGACA AGAGGAGGAGGCAGAGGACAA
Type 14	CAGGACGAGGAGCAGAGGCTG AGGACGAGGAGCAGAGGCTGA GGACGAGGAGCAGAGGCTGAG GACGAGGAGCAGAGGCTGAGG ACGAGGAGCAGAGGCTGAGGA CGAGGAGCAGAGGCTGAGGAG GAGGAGCAGAGGCTGAGGAGG AGGAGCAGAGGCTGAGGAGGA GGAGCAGAGGCTGAGGAGGAG
	GAGCAGAGGCTGAGGAGGAGG AGCAGAGGCTGAGGAGGAGGC GCAGAGGCTGAGGAGGAGGCA CAGAGGCTGAGGAGGAGGCAG AGAGGCTGAGGAGGAGGCAGA

Type of CALR mutation	shRNA sequences covering the mutation site
	GAGGCTGAGGAGGAGGCAGAG AGGCTGAGGAGGAGGCAGAGG GGCTGAGGAGGAGGCAGAGGA GCTGAGGAGGAGGCAGAGGAC CTGAGGAGGAGGCAGAGGACA TGAGGAGGAGGCAGAGGACAA
Type 15	AGGACGAGGAGCAGAGGCTTA GGACGAGGAGCAGAGGCTTAG GACGAGGAGCAGAGGCTTAGG ACGAGGAGCAGAGGCTTAGGA CGAGGAGCAGAGGCTTAGGAG GAGGAGCAGAGGCTTAGGAGG AGGAGCAGAGGCTTAGGAGGA GGAGCAGAGGCTTAGGAGGAG GAGCAGAGGCTTAGGAGGAGG AGCAGAGGCTTAGGAGGAGGG GCAGAGGCTTAGCAGGAGGGA CAGAGGCTTAGCAGGAGGGAG AGAGGCTTAGGAGGAGGGAGA GAGGCTTAGGAGGAGGGAGAG AGGCTTAGGAGGAGGGAGAGG GGCTTAGGAGGAGGGAGAGGA GCTTAGGAGGAGGGAGAGGAC CTTAGGAGGAGGGAGAGGACA TTAGGAGGAGGGAGAGGACAA TAGGAGCAGGGAGAGGACAAG AGGAGGAGGGAGAGGACAAGG GGAGGAGGGAGAGGACAAGGA GAGGAGGGAGAGGACAAGGAG AGGAGGGAGAGGACAAGGAGG GGAGGGAGAGGACAAGGAGGA GAGGCAGAGGACAAGCAGGAT AGGGAGAGGACAAGCAGGATG GGGAGAGGACAAGGAGGATGA GGAGAGGACAAGGAGGATGAT GAGAGGACAAGGAGGATGATG
Type 16	AGGACGAGGAGCAGAGGCTTC GGACGAGGAGCAGAGGCTTCA
	GACGAGGAGCAGAGGCTTCAG ACGAGGAGCAGAGGCTTCAGA CGAGGAGCAGAGGCTTCAGAG GAGGAGCAGAGGCTTCAGAGG AGGAGCAGAGGCTTCAGAGGA GGAGCAGAGGCTTCAGAGGAG GAGCAGAGGCTTCAGAGGAGG AGCAGAGGCTTCAGAGGAGGC GCAGAGGCTTCAGAGGAGGCA CAGAGGCTTCAGAGGAGGCAG AGAGGCTTCAGAGGAGGCAGA GAGGCTTCAGAGGAGGCAGAG AGGCTTCAGAGGAGGCAGAGG GGCTTCAGAGGAGGCAGAGGA GCTTCAGAGGAGGCAGAGGAC CTTCAGAGGAGGCAGAGGACA

Type of CALR mutation	shRNA sequences covering the mutation site
	TTCAGAGGAGGCAGAGGACAA TCAGAGGAGGCAGAGGACAAG CAGAGGAGGCAGAGGACAAGG AGAGGAGGCAGAGGACAAGGA
Type 17	GACGAGGAGCAGAGGCTTAAG ACGAGGAGCAGAGGCTTAAGA CGAGGAGCAGAGGCTTAAGAG GAGGAGCAGAGGCTTAAGAGG AGGAGCAGAGGCTTAAGAGGA GGAGCAGAGGCTTAAGAGGAG GAGCAGAGGCTTAAGAGGAGG AGCAGAGGCTTAAGAGGAGGC GCAGAGGCTTAAGAGGAGGCA CAGAGGCTTAAGAGGAGGCAG AGAGGCTTAAGAGGAGGCAGT GAGGCTTAAGAGGAGGCAGTG AGGCTTAAGAGGAGGCAGTGG GGCTTAAGAGGAGGCAGTGGG GCTTAAGAGGAGGCAGTGGAC CTTAAGAGGAGGCAGTGGACA TTAAGAGGAGGCAGTGGACAA TAAGAGGAGGCAGTGGACAAG AAGAGGAGGCAGTGGACAAGG AGAGGAGGCAGTGGACAAGGA GAGGAGGCAGTGGACAAGGAG AGGAGGCAGTGGACAAGGAGG GGAGGCAGTGGACAAGGAGGA
	GAGGCAGTGGACAAGGAGGAT AGGCAGTGGACAAGGAGGATG GGCAGTGGACAAGGAGGATGA GCAGTGGACAAGGAGGATGAT CAGTGGACAAGGAGGATGATG AGTGGACAAGGAGGATGATGA GTGGACAAGGAGGATGATGAG TGGACAAGGAGGATGATGAGG
Type 18	GACGAGGAGCAGAGGCTTAAG ACGAGGAGCAGAGGCTTAAGA CGAGGAGCAGAGGCTTAAGAG GAGGAGCAGAGGCTTAAGAGG AGGAGCAGAGGCTTAAGAGGA GGAGCAGAGGCTTAAGAGGAT GAGCAGAGGCTTAAGAGGATG AGCAGAGGCTTAAGAGGATGA GCAGAGGCTTAAGAGGATGAT CAGAGGCTTAAGAGGATGATG AGAGGCTTAAGAGGATGATGA GAGGCTTAAGAGGATGATGAG AGGCTTAAGAGGATGATGAGG GGCTTAAGAGGATGATGAGGA GCTTAAGAGGATGATGAGGAC CTTAAGAGGATGATGAGGACA TTAAGAGGATGATGAGGACAA TAAGAGGATGATGAGGACAAA AACAGGATGATGAGGACAAA

Type of CALR mutation	shRNA sequences covering the mutation site
	TAAGGATGATGAGGACAAAGA AGAGGATGATGAGGACAAAGA
Type 19	GAGCAGAGGCTTAAGGAGGAG AGCAGAGGCTTAAGGAGGAGA GCAGAGGCTTAAGGAGGAGAG CAGAGGCTTAAGGAGGAGAGG AGAGGCTTAAGGAGGAGAGGC GAGGCTTAAGGAGGAGAGGCA AGGCTTAAGGAGGAGAGGCAG GGCTTAAGGAGGAGAGGCAGA GCTTAAGGAGGAGAGGCAGAG CTTAAGGAGGAGAGGCAGAGG TTAAGGAGGAGAGGCAGAGGA TAAGGAGGAGAGGCAGAGGAC AAGGAGGAGAGGCAGAGGACA AGGAGGAGAGGCAGAGGACAA GGAGGAGAGGCAGAGGACAAG
	GAGGAGAGGCAGAGGACAAGG AGGAGAGGCAGAGGACAAGGA GGAGAGGCAGAGGACAAGGAG GAGAGGCAGAGGACAAGGAGG AGAGGCAGAGGACAAGGAGGA
Type 20	GCTTAAGCAGGAGGAAGAAGG CTTAAGCAGGAGGAAGAAGGG TTAAGCAGGAGGAAGAAGGGA TAAGCAGGAGGAAGAAGGGAG AAGGAGGAGGAAGAAGGGAGG AGGAGGAGGAAGAAGGGAGGA GGAGCAGGAAGAAGGGAGGAG GAGGAGGAAGAAGGGAGGAGG AGGAGGAAGAAGGGAGGAGGC GGAGGAAGAAGGGAGGAGGCA GAGGAAGAAGGGAGGAGGCAG AGGAAGAAGGGAGGAGGCAGA GGAAGAAGGGAGGAGGCAGAG GAAGAAGGGAGGAGGCAGAGG AAGAAGGGAGGAGGCAGAGGA AGAAGGGAGGAGGCAGAGGAC GAAGGGAGGAGGCAGAGGACA AAGGGAGGAGGCAGAGGACAA AGGGAGGAGGCAGAGGACAAG GGGAGGAGGCAGAGGACAAGG
Type 21	GCTTAAGGAGGAGGAAGAAGC CTTAAGGAGGAGGAAGAAGCG TTAAGGAGGAGGAAGAAGCGT TAAGGAGGAGGAAGAAGCGTT AAGGAGGAGGAAGAAGCGTTT AGGAGGAGGAAGAAGCGTTTA GGAGGAGGAAGAAGCGTTTAA GAGGAGGAAGAAGCGTTTAAG AGGAGGAAGAAGCGTTTAAGA GGAGGAAGAAGCGTTTAAGAG GAGGAGGAAGAAGCGTTTAAGG

Type of CALR mutation	shRNA sequences covering the mutation site
	GAGGAAAGAAAGCGTTTAAAGAGG AGGAAGAAGCGTTTAAAGAGGA GGAAGAAGCGTTTAAAGAGGAC GAAGAAGCGTTTAAAGAGGACA AAGAAGCGTTTAAAGAGGACAA AGAAGCGTTTAAAGAGGACAAG GAAGCGTTTAAAGAGGACAAGG AAGCGTTTAAAGAGGACAAGGA
	AGCGTTTAAAGAGGACAAGGAG GCGTTTAAAGAGGACAAGGAGG CGTTTAAAGAGGACAAGGAGGA GTTTAAAGAGGACAAGGAGGAT TTTAAAGAGGACAAGGAGGATG TTAAGAGGACAAGGAGGATGA TAAGAGGACAAGGAGGATGAT AAGAGGACAAGGAGGATGATG
Type 22	TTAAGGAGGAGGAAGAAGACA TAAGGAGGAGGAAGAAGACAA AAGGAGGAGGAAGAAGACAAC AGGAGGAGGAAGAAGACAACG GGAGGAGGAAGAAGACAACGC GAGGAGGAAGAAGACAACGCA AGGAGGAAGAAGACAACGCAA GGAGGAAGAAGACAACGCAAA GAGGAAGAAGACAACGCAAAAG AGGAAGAAGACAACGCAAAAG GGAAGAAGACAACGCAAAAGAG GAAGAAGACAACGCAAAAGAGG AAGAAGACAACGCAAAAGAGGA AGAAGACAACGCAAAAGAGGAG GAAGACAACGCAAAAGAGGAGG AAGACAACGCAAAAGAGGAGGA AGACAACGCAAAAGAGGAGGAG GACAACGCAAAAGAGGAGGAGG ACAACGCAAAAGAGGAGGAGGA CAACGCAAAAGAGGAGGAGGAG
Type 23	TTAAGGAGGAGGAAGAAGACT TAAGGAGGAGGAAGAAGACTG AAGGAGGAGGAAGAAGACTGC AGGAGGAGGAAGAAGACTGCG GGAGGAGGAAGAAGACTGCGT GAGGAGGAAGAAGACTGCGTG AGGAGGAAGAAGACTGCGTGA GGAGGAAGAAGACTGCGTGAG GAGGAAGAAGACTGCGTGAGG AGGAAGAAGACTGCGTGAGGA GGAAGAAGACTGCGTGAGGAG GAAGAAGACTGCGTGAGGAGG AAGAAGACTGCGTGAGGAGGA AGAAGACTGCGTGAGGAGGAG GAAGACTGCGTGAGGAGGAGG
	AAGACTGCGTGAGGAGGAGGA

Type of CALR mutation	shRNA sequences covering the mutation site
	AGACTGCGTGAGGAGGAGGAG GACTGCGTGAGGAGGAGGAGG ACTGCGTGAGGAGGAGGAGGC CTGCGTGAGGAGGAGGAGGCA TGCGTGAGGAGGAGGAGGCAG GCGTGAGGAGGAGGAGGCAGA CGTGAGGAGGAGGAGGCAGAG GTGAGGAGGAGGAGGCAGAGG TGAGGAGGAGGAGGCAGAGGA
Type 24	TTAAGGAGGAGGAAGAAGACA TAAGGAGGAGGAAGAAGACAG AAGGAGGAGGAAGAAGACAGG AGGAGGAGGAAGAAGACAGGA GGAGGAGGAAGAAGACAGGAG GAGGAGGAAGAAGACAGGAGG AGGAGGAAGAAGACAGGAGGC GGAGGAAGAAGACAGGAGGCA GAGGAAGAAGACAGGAGGCAG AGGAAGAAGACAGGAGGCAGA GGAAGAAGACAGGAGGCAGAG GAAGAAGACAGGAGGCAGAGG AAGAAGACAGGAGGCAGAGGA AGAAACACAGGAGGCAGAGGAC GAAGACAGGAGGCAGAGGACA AAGACAGGAGGCAGAGGACAA AGACAGGAGGCAGAGGACAAG GACAGGAGGCAGAGGACAAGG ACAGCAGGCAGAGGACAAGGA CAGGAGGCAGAGGACAAGGAG
Type 25	AAGGAGGAGGAAGAAGACAAA AGGAGGAGGAAGAAGACAAAA GGAGGAGGAAGAAGACAAAAG GAGGAGGAAGAAGACAAAAGG AGGAGGAAGAAGACAAAAGGC GGAGGAAGAAGACAAAAGGCA GAGGAAGAAGACAAAAGGCAG AGGAAGAAGACAAAAGGCAGA GGAAGAAGACAAAAGGCAGAG GAAGAAGACAAAAGGCAGAGG AAGAAGACAAAAGGCAGAGGA AGAAGACAAAAGGCAGAGGAC GAAGACAAAAGGCAGAGGACA
	AAGACAAAAGGCAGAGGACAA AGACAAAAGGCAGAGGACAAG GACAAAAGGCAGAGGACAAGG ACAAAAGGCAGAGGACAAGGA CAAAAGGCAGAGGACAAGGAG AAAAGGCAGAGGACAAGGAGG AAAGGCAGAGGACAAGGAGGA AAGGCAGAGGACAAGGAGGAT
Type 26	AAGGAGCAGGAAGAAGACAAA AGGAGGAGGAAGAAGACAAAA

Type of CALR mutation	shRNA sequences covering the mutation site
	GGAGGAGGAAGAAGACAAAAA GAGGAGGAAGAAGACAAAAAC AGGAGGAAGAAGACAAAAACG GGAGGAAGAAGACAAAAACGC GAGGAAGAAGACAAAAACGCA AGGAAGAAGACAAAAACGCAA GGAAGAAGACAAAAACGCAAA GAAGAAGACAAAAACGCAAAG AAGAAGACAAAAACGCAAAGA AGAAGACAAAAACGCAAAGAG GAAGACAAAAACGCAAAGAGG AAGACAAAAACGCAAAGAGGA AGACAAAAACGCAAAGAGGAG GACAAAAACGCAAAGAGGAGG ACAAAAACGCAAAGAGGAGGA CAAAAACGCAAAGAGGAGGAG AAAAACGCAAAGAGGAGGAGG AAAACGCAAAGAGGAGGAGGA AAACGCAAAGAGGAGGAGGAG
Type 27	AGGAGGAGGAAGAAGACAAGT GGAGGAGGAAGAAGACAAGTG GAGGAGGAAGAAGACAAGTGT AGGAGGAAGAAGACAAGTGTT GGAGGAAGAAGACAAGTGTTT GAGGAAGAAGACAAGTGTTTC AGGAAGAAGACAAGTGTTTCG GGAAGAAGACAAGTGTTTCGC GAAGAAGACAAGTGTTTCGCA AAGAAGACAAGTGTTTCGCAA AGAAGACAAGTGTTTCGCAAA GAAGACAAGTGTTTCGCAAAG AAGACAAGTGTTTCGCAAAGA AGACAAGTGTTTCGCAAAGAG
	GACAAGTGTTTCGCAAAGAGG ACAAGTGTTTCGCAAAGAGGA CAAGTGTTTCGCAAAGAGGAG AAGTGTTTCGCAAAGAGGAGG AGTGTTTCGCAAAGAGGAGGA GTGTTTCGCAAAGAGGAGGAG TGTTTCGCAAAGAGGAGGAGG GTTTCGCAAAGAGGAGGAGGA TTTCGCAAAGAGGAGGAGGAG TTCGCAAAGAGGAGGAGGAGG TCGCAAAGAGGAGGAGGAGGC
Type 28	GAAGAAGACAAGAAACGCAAA AAGAAGACAAGAAACGCAAAA AGAAGACAAGAAACGCAAAAG GAAGACAAGAAACGCAAAAGG AAGACAAGAAACGCAAAAGGA AGACAAGAAACGCAAAAGGAG GACAAGAAACGCAAAAGGAGG ACAAGAAACGCAAAAGGAGGA CAAGAAACGCAAAAGGAGGAT AAGAAACGCAAAAGGAGGATG

Type of CALR mutation	shRNA sequences covering the mutation site
	AGAAAACGCAAAAGGAGGATGA GAAAACGCAAAAGGAGGATGAT AAACGCAAAAGGAGGATGATG AACGCAAAAGGAGGATGATGA ACGCAAAAGGAGGATGATGAG CGCAAAAGGAGGATGATGAGG GCAAAAGGAGGATGATGAGGA CAAAAGGAGGATGATGAGGAC AAAAGGAGGATGATGAGGACA AAAGGAGGATGATGAGGACAA
Type 29	AAGACAAGAAACGCAAAGAGC AGACAAGAAACGCAAAGAGCC GACAAGAAACGCAAAGAGCCT ACAAGAAACGCAAAGAGCCTC CAAGAAACGCAAAGAGCCTCC AAGAAACGCAAAGAGCCTCCT AGAAACGCAAAGAGCCTCCTC GAAACGCAAAGAGCCTCCTCT AAACGCAAAGAGCCTCCTCTT AACGCAAAGAGCCTCCTCTTT ACGCAAAGAGCCTCCTCTTTG CGCAAAGAGCCTCCTCTTTGT
	GCAAAGAGCCTCCTCTTTGTC CAAAGAGCCTCCTCTTTGTCT AAAGAGCCTCCTCTTTGTCTA AAGAGCCTCCTCTTTGTCTAA AGAGCCTCCTCTTTGTCTAAG GAGCCTCCTCTTTGTCTAAGG AGCCTCCTCTTTGTCTAAGGA GCCTCCTCTTTGTCTAAGGAG CCTCCTCTTTGTCTAAGGAGG CTCCTCTTTGTCTAAGGAGGA TCCTCTTTGTCTAAGGAGGAT CCTCTTTGTCTAAGGAGGATG CTCTTTGTCTAAGGAGGATGA TCTTTGTCTAAGGAGGATGAT CTTTGTCTAAGGAGGATGATG TTTGTCTAAGGAGGATGATGA TTGTCTAAGGAGGATGATGAG TGTCTAAGGAGGATGATGAGG GTCTAAGGAGGATGATGAGGA TCTAAGGAGGATGATGAGGAC CTAAGGAGGATGATGAGGACA TAAGGAGGATGATGAGGACAA
Type 30	GACAAGAAACGCAAAGAGGAC ACAAGAAACGCAAAGAGGACC CAAGAAACGCAAAGAGGACCA AAGAAACGCAAAGAGGACCAT AGAAACGCAAAGAGGACCATC GAAACGCAAAGAGGACCATCC AAACGCAAAGAGGACCATCCT AACGCAAAGAGGACCATCCTT ACGCAAAGAGGACCATCCTTG CGCAAAGAGGACCATCCTTGT

Type of CALR mutation	shRNA sequences covering the mutation site
	GCAAAGAGGACCATCCTTGTC CAAAGAGGACCATCCTTGTCG AAAGAGGACCATCCTTGTCGG AAGAGGACCATCCTTGTCGGA AGAGGACCATCCTTGTCGGAG GAGGACCATCCTTGTCGGAGG AGGACCATCCTTGTCGGAGGA GGACCATCCTTGTCGGAGGAT GACCATCCTTGTCGGAGGATG ACCATCCTTGTCGGAGGATGA CCATCCTTGTCGGAGGATGAT
	CATCCTTGTCGGAGGATGATG ATCCTTGTCGGAGGATGATGA TCCTTGTCGGAGGATGATGAG CCTTGTCGGAGGATGATGAGG CTTGTCGGAGGATGATGAGGA TTGTCGGAGGATGATGAGGAC TGTCGGAGGATGATGAGGACA GTCGGAGGATGATGAGGACAA TCGGAGGATGATGAGGACAAA CGGAGGATGATGAGGACAAAG
Type 31	AGAGGAGGAGGAGGCAGAGGG GAGGAGCAGGAGGCAGAGGGC AGGAGGAGGAGGCAGAGGGCA GGAGGAGGAGGCAGAGGGCAA GAGGAGGAGGCAGAGGGCAAT AGGAGGAGGCAGAGGGCAATT GGAGGAGGCAGAGGGCAATTG GAGGAGGCAGAGGGCAATTGT AGGAGGCAGAGGGCAATTGTC GGAGGCAGAGGGCAATTGTCG GAGGCAGAGGGCAATTGTCGG AGGCAGAGGGCAATTGTCGGA GGCAGAGGGCAATTGTCGGAG GCAGAGGGCAATTGTCGGAGG CAGAGGGCAATTGTCGGAGGA AGAGGGCAATTGTCGGAGGAT GAGGGCAATTGTCGGAGGATG AGGGCAATTGTCGGAGGATGA GGGCAATTGTCGGAGGATGAT GGCAATTGTCGGAGGATGATG GCAATTGTCGGAGGATGATGA CAATTGTCGGAGGATGATGAG AATTGTCGGAGGATGATGAGG ATTGTCGGAGGATGATGAGGA TTGTCGGAGGATGATGAGGAC TGTCGGAGGATGATGAGGACA GTCGGAGGATGATGAGGACAA TCGGAGGATGATGAGGACAAA CGGAGGATGATGAGGACAAAG
Type 32	AGGAGGAGGAGGCAGAGGACT GGAGGAGGAGGCAGAGGACTG GAGGAGGAGGCAGAGGACTGT AGGAGGAGGCAGAGGACTGTC

Type of CALR mutation	shRNA sequences covering the mutation site
	GGAGGAGGCAGAGGACTGTCG GAGGAGGCAGAGGACTGTCGG AGGAGGCAGAGGACTGTCGGA GGAGGCAGAGGACTGTCGGAG GAGGCAGAGGACTGTCGGAGG AGGCAGAGGACTGTCGGAGGA GGCAGAGGACTGTCGGAGGAT GCAGAGGACTGTCGGAGGATG CAGAGGACTGTCGGAGGATGA AGAGGACTGTCGGAGGATGAT GAGGACTGTCGGAGGATGATG AGGACTGTCGGAGGATGATGA GGACTGTCGGAGGATGATGAG GACTGTCGGAGGATGATGAGG ACTGTCGGAGGATGATGAGGA CTGTCGGAGGATGATGAGGAC TGTCGGAGGATGATGAGGACA GTCCGAGGATGATGAGGACAA TCGGAGGATGATGAGGACAAA CGGAGGATGATGAGGACAAAG
Type 33	GAGGAGGAGGCAGAGGACAAA AGGAGGAGGCAGAGGACAAAT GGAGGAGGCAGAGGACAAATG GAGGAGGCAGAGGACAAATGT AGGAGGCAGAGGACAAATGTC GGAGGCAGAGGACAAATGTCG GAGGCAGAGGACAAATGTCGG AGGCAGAGGACAAATGTCGGA GGCAGAGGACAAATGTCGGAG GCAGAGGACAAATGTCGGAGG CAGAGGACAAATGTCGGAGGA AGAGGACAAATGTCGGAGGAT GAGGACAAATGTCGGAGGATG AGGACAAATGTCGGAGGATGA GGACAAATGTCGGAGGATGAT GACAAATGTCGGAGGATGATG ACAAATGTCGGAGGATGATGA CAAATGTCGGAGGATGATGAG AAATGTCGGAGGATGATGAGG AATGTCGGAGGATGATGAGGA ATGTCGGAGGATGATGAGGAC TGTCGGAGGATGATGAGGACA GTCGGAGGATGATGAGGACAA
	TCGGAGGATGATGAGGACAAA CGGAGGATGATGAGGACAAAG
Type 34	GGAGGAGGAGGCAGAGGACAC GAGGAGGAGGCAGAGGACACT AGGAGGAGGCAGAGGACACTT GGAGGAGGCAGAGGACACTTG GAGGAGGCAGAGGACACTTGT AGGAGGCAGAGGACACTTGTC GGAGGCAGAGGACACTTGTCG

Type of CALR mutation	shRNA sequences covering the mutation site
	GAGGCAGAGGACACTTGTCCG AGGCAGAGGACACTTGTCCGA GGCAGAGGACACTTGTCCGAG GCAGAGGACACTTGTCCGAGG CAGAGGACACTTGTCCGAGGA AGAGGACACTTGTCCGAGGAT GAGGACACTTGTCCGAGGATG AGGACACTTGTCCGAGGATGA GGACACTTGTCCGAGGATGAT GACACTTGTCCGAGGATGATG ACACTTGTCCGAGGATGATGA CACTTGTCCGAGGATGATGAG ACTTGTCCGAGGATGATGAGG CTTGTCCGAGGATGATGAGGA TTGTCCGAGGATGATGAGGAC TGTCGGAGGATGATGAGGACA GTCGGAGGATGATGAGGACAA TCGGAGGATGATGAGGACAAA CGGAGGATGATGAGGACAAAG
Type 35	GGAGGAGGAGGCAGAGGACAT GAGGAGGAGGCAGAGGACATT AGGAGGAGGCAGAGGACATTT GGAGGAGGCAGAGGACATTTG GAGGAGGCAGAGGACATTTGT AGGAGGCAGAGGACATTTGTC GGAGGCAGAGGACATTTGTCTG GAGGCAGAGGACATTTGTCCG AGGCAGAGGACATTTGTCCGA GGCAGAGGACATTTGTCCGAG GCAGAGGACATTTGTCCGAGG CAGAGGACATTTGTCCGAGGA AGAGGACATTTGTCCGAGGAT GAGGACATTTGTCCGAGGATG AGGACATTTGTCCGAGGATGA
	GGACATTTGTCCGAGGATGAT GACATTTGTCCGAGGATGATG ACATTTGTCCGAGGATGATGA CATTTGTCCGAGGATGATGAG ATTTGTCCGAGGATGATGAGG TTTGTCCGAGGATGATGAGGA TTGTCCGAGGATGATGAGGAC TGTCGGAGGATGATGAGGACA GTCGGAGGATGATGAGGACAA TCGGAGGATGATGAGGACAAA CGGAGGATGATGAGGACAAAG
Type 36	GGAGGAGGCAGAGGACAAGTG GAGGAGGCAGAGGACAAGTGT AGGAGGCAGAGGACAAGTGTC GGAGGCAGAGGACAAGTGTCG GAGGCAGAGGACAAGTGTCGG AGGCAGAGGACAAGTGTCGGA GGCAGAGGACAAGTGTCGGAG GCAGAGGACAAGTGTCGGAGG CAGAGGACAAGTGTCGGAGGA

Type of CALR mutation	shRNA sequences covering the mutation site
	AGAGGACAAGTGTCTGGAGGAT GAGGACAAGTGTCTGGAGGATG AGGACAAGTGTCTGGAGGATGA GGACAAGTGTCTGGAGGATGAT GACAAGTGTCTGGAGGATGATG ACAAGTGTCTGGAGGATGATGA CAAGTGTCTGGAGGATGATGAG AAGTGTCTGGAGGATGATGAGG AGTGTCTGGAGGATGATGAGGA GTGTCTGGAGGATGATGAGGAC TGTCTGGAGGATGATGAGGACA GTCCTGGAGGATGATGAGGACAA TCCTGGAGGATGATGAGGACAAA CGGAGGATGATGAGGACAAAG

[0185] Herein contemplated are antibodies that specifically bind to the above provided mutant calreticulin protein(s). Such antibodies can be used for diagnostic and therapeutic purposes in accordance with the present invention. For example, antibodies raised against the C-terminal unique polypeptide of mutated calreticulin offers a diagnostic test for myeloid malignancy. Also detection of peptides derived from this unique C-terminus by mass spectrometry offers a diagnostic test for myeloid malignancy. Preferably, such antibodies are inhibitors of mutant calreticulin.

[0186] For example, antibodies to be used herein can specifically bind to the following mutant calreticulin protein(s) shown in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434, respectively. Particularly, such antibodies can specifically bind to the C-terminus of the mutant calreticulin protein(s), for example, to proteins as shown in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, respectively.

[0187] It is envisaged herein that the antibodies can specifically bind to (functional) fragments or (functional) derivatives of the mutant calreticulin proteins as defined herein, for example also to polypeptides having at least 70 % or more identity to herein provided mutant calreticulin protein(s).

[0188] Accordingly, the present invention relates to the use of these antibodies in the methods of the present invention. Therefore, the present invention relates to the use of the herein above described antibody/antibodies specifically binding to or specifically recognizing one or more of the herein described and provided mutant calreticulin proteins polypeptides for assessing whether a patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy.

[0189] The present invention also relates to an antibody/antibodies specifically binding to the C-terminus of the mutant calreticulin protein(s) as shown in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, respectively, or the above composition comprising said antibody/antibodies for the preparation of a diagnostic kit for use in the methods of the present invention.

[0190] The antibody may be a polyclonal antibody, a monoclonal antibody, a full antibody (immunoglobulin), a F(ab)-fragment, a F(ab)₂-fragment, a single-chain antibody, a chimeric antibody, a CDR-grafted antibody, a bivalent antibody-construct, a bispecific single chain antibody, a synthetic antibody or a cross-cloned antibody and the like.

[0191] Polyclonal or monoclonal antibodies or other antibodies (derived therefrom) can be routinely prepared using, inter alia, standard immunization protocols; see Ed Harlow, David Lane, (December 1988), *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory; or Ed Harlow, David Lane, (December 1998), *Portable Protocols (Using Antibodies): A Laboratory Manual* 2nd edition, Cold Spring Harbor Laboratory.

[0192] For example, immunization may involve the intraperitoneal or subcutaneous administration of the mutant calreticulin protein/polypeptide (and/or fragments, isoforms, homologues and so on) as defined herein to a mammal (e.g. rodents such as mice, rats, hamsters and the like). Preferably, fragments of the mutant protein/polypeptide are used, wherein the fragment preferably bears the C-terminus (or a fragment thereof) as defined herein.

[0193] A preferred fragment of the above mentioned polypeptides (i.e. the mutant calreticulin protein/polypeptide(s)) provided herein and to be used in accordance with the present invention consists of from 15 to 25 contiguous amino acids. Accordingly, a fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention preferably consists of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 contiguous amino acids. A fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention preferably consists of from 15 to 25 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 4. A fragment of the above mentioned polypeptides provided herein and to be used in accordance with the present invention preferably consists of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 4.

[0194] As proof of principle, a fragment (RRKMSPARPTSCREACLQGWTEA) was used to generate polyclonal antibodies that bind specifically to mutant calreticulin antibody; see Example 2. This fragment consists of 24 contiguous amino acids of the amino acid sequence shown in SEQ ID NO: 4. These polyclonal antibodies can be used as a research tool and/or in the diagnostic methods provided herein. This fragment of the mutant calreticulin protein (or other fragments provided and defined herein) can also be used as a vaccine, as it is described herein further below.

[0195] Methods for the preparation and screening of antibodies that specifically bind to or specifically recognize the mutant polypeptides are known in the art. For example, antibodies recognizing the mutant protein may be affinity purified. ELISA is commonly used for screening sera and/or assaying affinity column fractions. Western Blots can be used to demonstrate that the antibody can detect the actual protein of interest and to evaluate whether the antibody only recognizes the protein of interest, or if it cross-reacts with other proteins.

[0196] A person skilled in the art is in the position to apply and to adapt the teaching of these documents for the generation and validation of antibodies specifically binding to or specifically recognizing the polypeptides as defined herein in context of the present invention.

[0197] The following relates to the use of the herein provided mutant calreticulin proteins as a vaccine. Thus, the mutant calreticulin proteins act as antigens. Therefore, the terms "mutant calreticulin protein" and "mutant calreticulin protein antigens" and the like can be used interchangeably herein in the following.

[0198] In accordance with the above, the herein provided mutant calreticulin proteins can be used as a vaccine. In other words, herein provided mutant calreticulin proteins can be used in active immunization. Thus, the present invention relates to mutant calreticulin proteins as defined and provided herein (or nucleic acids (or vectors comprising same)) encoding mutant calreticulin proteins as defined and provided herein for use as vaccine. Mutant calreticulin proteins, fragments and derivatives thereof have been described herein above in great detail. These explanations and definitions apply, mutatis mutandis, in this context. Useful as vaccine are, in particular, proteins comprising or consisting of the C-terminus of the herein provided mutant calreticulin proteins as shown in SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, respectively, as well as of fragments thereof. Particularly useful in this context are mutant calreticulin proteins that comprise or consist of the minimum C-terminus as shown in SEQ ID NO: 4 or a fragment thereof.

[0199] The present invention relates to a protein for use as vaccine, wherein the protein is selected from the group

consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2 or 3;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by one or more mutant alleles of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a) or (d).

[0200] The present invention relates to a protein for use as vaccine, wherein the protein is selected from the group consisting of

1. (a) a protein encoded by a nucleic acid molecule having the nucleic acid sequence as depicted in SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, or 143;
2. (b) a protein comprising the amino acid sequence as shown in SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, or 144;
3. (c) a protein as defined in (a) or (b) wherein one to 10 amino acids are deleted, inserted, added or substituted;
4. (d) a protein comprising an amino acid sequence encoded by a nucleic acid hybridizing under stringent conditions to the complementary strand of nucleic acid molecules as defined in (a) or (c) and wherein the protein is encoded by one or more mutant alleles of the calreticulin gene;
5. (e) a protein having at least 70 % identity to the protein of any one of (a) to (d); and
6. (f) a protein comprising an amino acid sequence encoded by a nucleic acid being degenerate as a result of the genetic code to the nucleotide sequence of a nucleic acid as defined in (a)(d).

[0201] The present invention relates to a protein for use as vaccine as defined herein above, wherein the protein consists of 15 to 25 contiguous amino acids of the protein as shown in SEQ ID NO: 4.

[0202] Preferably, a fragment of the protein shown in SEQ ID NO: 4 comprising or consisting of 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 amino acids of the protein as shown in SEQ ID NO: 4 is used as vaccine. Particularly preferred are fragments of the protein shown in SEQ ID NO: 4 comprising or consisting of 15 to 25 contiguous amino acids of the protein as shown in SEQ ID NO: 4 for use as vaccine. Accordingly, a fragment of the protein shown in SEQ ID NO: 4 comprising or consisting of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 contiguous amino acids of the protein as shown in SEQ ID NO: 4 is used as vaccine.

[0203] Such fragments or derivatives of the protein shown in SEQ ID NO: 4 can be coupled to proteins such as keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA), or bacterial toxoids (e.g. tetanus toxoid, diphtheria toxoid) and used for immunization.

[0204] The vaccine can be used in the treatment of myeloid malignancies as defined herein.

[0205] Accordingly, the present invention relates to a vaccine for use in a method for prophylactic or therapeutic treatment of a myeloid malignancy as defined herein, the method comprising administering an effective amount of the vaccine as defined herein above to a patient. In other words, the present invention relates to a mutant calreticulin protein as defined herein for use in a method for prophylactic or therapeutic treatment of a myeloid malignancy as defined herein, the method comprising administering an effective amount of the mutant calreticulin protein as defined herein above to a patient.

[0206] For example, while a patient would not normally react to CALR, because of self-tolerance, many of the exon 9 mutations of CALR cause frameshifts such that the C-terminal portion of mutant CALR differs from the wild type and is therefore not subject to self-tolerance. Immunization with the mutant polypeptide would generate an immune response against mutant CALR. Accordingly, a vaccine can be used therapeutically to target the existing cancer.

[0207] A vaccine could also be used prophylactically. For example, it is known that cancers can mutate and evolve to evade the host immune response and anticancer treatment. Thus, a patient with a myeloid malignancy that has no CALR mutation, or a single CALR mutation, may later develop additional CALR mutations. Immunization against other CALR mutant forms can, therefore, select against such CALR mutants.

[0208] CALR mutants can also be used to generate antibodies in vitro or in another animal for use in therapy in the patient. Such an approach is particularly useful because the antibodies can be produced against epitopes tolerated in the patient. Such antibodies are also useful for therapy because the titre can be precisely controlled, and the antibody may also be conjugated to toxins or radionuclides for targeted therapy.

[0209] An exemplary protocol for performing active immunization (or the use of the vaccines provided herein above) is described in the following:

For example, mice can be immunized with a vaccine (for example a peptide derived from the protein shown in SEQ ID NO: 4 or a peptide which is a fragment of the protein shown in SEQ ID NO: 4, wherein the peptide is coupled with KLH or BSA) before transplantation of bone marrow cells expressing mutant or wild type calreticulin. The immunized recipient mice can be sublethally or lethally irradiated to promote engraftment or not irradiated at all. The transplanted cells can be a mixture of cells expressing either mutated calreticulin or wild type calreticulin. The immunized mice will be followed after transplantation. If an immune response is elicited against mutant calreticulin expressing cells, the engraftment will preferably occur with wild type calreticulin expressing cells. Control mice without immunisation will be used. Alternatively, immunization can be performed after engraftment of mice with a 50:50 (or other ratio) mixture of wild type/mutated calreticulin expressing cells. If an immune response is elicited against mutant calreticulin expressing cells, the 50:50 (or other) ratio will change in favour of the wild type calreticulin expressing cells. Control mice without immunization will be used for comparison.

[0210] The present invention relates to the use of a mutant calreticulin protein as antigen as provided herein and defined herein above and, optionally, an adjuvant, for the manufacture of a vaccine composition for the treatment or prevention of myeloid malignancy as defined herein.

[0211] The mutant calreticulin protein can be produced recombinantly (i.e. produced in appropriate host cells) or synthetic (i.e. chemically synthesized). Recombinant production of mutant calreticulin protein is described herein. For example, recombinant production can be achieved using any one of the molecular cloning and recombinant expression techniques known in the art. For example, a nucleic acid molecule encoding mutant calreticulin protein can be introduced into an appropriate host cell, such as a bacterium, a yeast cell (e.g., a *Pichia* cell), an insect cell or a mammalian cell (e.g., CHO cell). The encoding nucleic acid molecule can be placed in an operable linkage to a promoter capable of effecting the expression of the mutant calreticulin protein antigen in the host cell. mutant calreticulin protein, which is expressed by the host cell, can be readily purified using routine protein purification techniques.

[0212] For example, the nucleotide sequence as set forth in SEQ ID NO: 1, 2 or 3 or a nucleic acid sequence

encoding the mutant calreticulin protein antigen shown in SEQ ID NO: 4 or encoding fragment thereof, such as a protein consisting of 15 to 25 contiguous amino acids of the protein shown in SEQ ID NO: 4, can be cloned in an expression vector and placed in an operable linkage to a temperature sensitive promoter. The expression vector can be introduced into *Escherichia coli* and the antigen can be expressed upon heat induction. The cells can be lysed and the inclusion bodies where the antigen accumulates are separated by centrifugation. The recombinant protein in the inclusion bodies is solubilized using SDS or other solubilization agents known in the art such as urea, guanidine hydrochloride, sodium cholate, taurocholate, and sodium deoxycholate. In accordance with the present invention, a purified recombinant mutant calreticulin protein is combined with a pharmaceutically acceptable carrier to form a vaccine composition.

[0213] The present invention provides an immunogenic composition for conferring protection in a patient against myeloid malignancy, the composition comprising a mutant calreticulin protein. The composition can be formulated as a vaccine for *in vivo* administration to a patient. The composition can comprise an adjuvant, such as aluminum hydroxide or aluminum phosphate.

[0214] Further, the present invention provides an immunogenic composition comprising a mutant calreticulin protein as defined herein above for use as a medicament. A composition comprising a mutant calreticulin protein as defined herein above can be used for the manufacture of a medicament for immunizing a host or patient against disease a myeloid malignancy.

[0215] The present invention is directed to a vaccine composition containing a mutant calreticulin protein antigen as defined herein (or "a mutant calreticulin protein vaccine"), that is suitable for administration to patients and is capable of protecting patients against a myeloid malignancy.

[0216] The term "a pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, adjuvants, stabilizing agents, diluents, preservatives, antibacterial and antifungal agents, isotonic agents, adsorption delaying agents, and the like. Diluents can include water, saline, dextrose, ethanol, glycerol, and the like. Isotonic agents can include sodium chloride, dextrose, mannitol, sorbitol, and lactose, among others. Stabilizers include albumin, among others.

[0217] Adjuvants suitable for use in a vaccine composition in accordance with the present invention include, but are not limited to several adjuvant classes such as; mineral salts, e.g., Alum, aluminum hydroxide, aluminum phosphate and calcium phosphate; surface-active agents and microparticles, e.g., nonionic block polymer surfactants (e.g., cholesterol), virosomes, saponins (e.g., Quil A, QS-21 and GPI-0100), proteosomes, immune stimulating complexes, cochleates, quaternary amines (dimethyl diocetadecyl ammonium bromide (DDA)), avridine, vitamin A, vitamin E; bacterial products such as the RIBI adjuvant system (Ribi Inc.), cell wall skeleton of *Mycobacterum phlei* (Detox®), muramyl dipeptides (MDP) and tripeptides (MTP), monophosphoryl lipid A, *Bacillus Calmette-Guerin*, heat labile *E. coli* enterotoxins, cholera toxin, trehalose dimycolate, CpG oligodeoxynucleotides; cytokines and hormones, e.g., interleukins (IL-1, IL-2, IL-6, IL-12, IL-15, IL-18), granulocyte-macrophage colony stimulating factor, dehydroepiandrosterone, 1,25-dihydroxy vitamin D3; polyanions, e.g., dextran; polyacrylics (e.g., polymethylmethacrylate, Carbopol 934P); carriers e.g., tetanus toxoid, diphtheria toxoid, cholera toxin B subunit, mutant heat labile enterotoxin of enterotoxigenic *E. coli* (rmLT), heat shock proteins; oil-in-water emulsions e.g., AMPHIGEN® (Hydronics, USA); and water-in-oil emulsions such as, e.g., Freund's complete and incomplete adjuvants.

[0218] The herein provided mutant calreticulin protein antigen and the pharmaceutically acceptable carrier can be combined in any convenient and practical manner to form a vaccine composition, e.g., by admixture, solution, suspension, emulsification, encapsulation, absorption and the like, and can be made in formulations such as tablets, capsules, powder, syrup, suspensions that are suitable for injections, implantations, inhalations, ingestions or the like. Preferably, the vaccine is formulated such that it can be administered to patients by injection in a dose of about 0.1 to 5 ml, or preferably about 0.5 to 2.5 ml, or even more preferably, in a dose of about 1 ml. When appropriate, the pharmaceutical compositions of the present invention should be made sterile by well-known procedures.

[0219] The amount of herein provided mutant calreticulin protein antigen in the vaccines should be immunizing-

effective and is generally in the range of 0.5 -1000 pg per dose.

[0220] The amount of adjuvants suitable for use in the vaccines depends upon the nature of the adjuvant used. For example, when Quil A and cholesterol are used as adjuvant, Quil A is generally in an amount of about 1-1000 Pg per dose; and cholesterol is generally in an amount of about 1-1000 Pg per dose.

[0221] In accordance with the present disclosure, a vaccine can be administered by any known routes, including the oral, intranasal, mucosal, topical, transdermal, and parenteral (e.g., intravenous, intraperitoneal, intradermal, subcutaneous or intramuscular). Administration can also be achieved using needle-free delivery devices. Administration can be achieved using a combination of routes, e.g., first administration using a parental route and subsequent administration using a mucosal route. Preferred routes of administration include subcutaneous and intramuscular administrations.

[0222] The present invention provides combination vaccines and methods for protecting patients by administering such combination vaccines.

[0223] The present invention provides an immunogenic agent, wherein the immunogenic agent is mutant calreticulin protein as defined and provided herein above. Said agent is effective to induce an immunogenic response against mutant calreticulin proteins in a patient. Therefore, the immunogenic agent can be used in the treatment or prevention of a myeloid malignancy as defined herein.

[0224] Further, the present invention provides a conjugate comprising an immunogenic agent linked to a carrier protein. The immunogenic agent is mutant calreticulin protein as defined and provided herein above. The carrier protein (such as serum albumin among others) can enhance the immune response. The conjugate can be a fusion protein comprising an immunogenic agent (i.e. the mutant calreticulin protein as defined and provided herein above) fused to a carrier protein. The agent can also be linked to the carrier protein by chemical cross-linking. The agent can be linked or fused to the amino terminus of the carrier protein. The agent can be linked or fused to the carboxyl of the carrier protein. The agent can be linked or fused internally to the carrier protein. Multiple repeats or multimers of the agent can be present in a conjugate, such as a fusion protein. The agent can be part of a longer polypeptide that includes the agent with other amino acids. The agent can be a component of a particle. The particle can be a liposome or a microparticle. The agent can be emulsified or encapsulated in the particle, such as a liposome or a microparticle.

[0225] The presence of a unique C-terminal polypeptide in mutated calreticulin proteins as provided herein offers the opportunity to target the mutant protein leaving the wild type protein intact. As the amino acid sequence derived from the -1 reading frame of calreticulin exon 9 encodes a peptide not showing homology to any other vertebrate protein, inhibitors as defined above (like antibodies (preferably inhibitory antibodies), siRNA, shRNAs or small molecule drugs) can be generated against it with a therapeutic effect. For example, the mutant calreticulin protein C-terminal amino acid sequence (such as the mutant protein derived from the alternative reading frame of exon 9) can be used for generation of polyclonal and monoclonal antibodies (preferably inhibitory antibodies). The mutant calreticulin as provided herein is therefore a valuable target for immunotherapy in myeloid malignancies. Alternatively, immunization of patients with the mutant peptide or recombinant mutant proteins can be used in therapeutic intervention of myeloid malignancy.

[0226] It has been shown that calreticulin is secreted from the cells and can be detected in serum. Calreticulin is also trafficked to the cellular surface where it provides an "eat me signal" for phagocytosis of apoptotic cells.

[0227] Accordingly, the present invention provides for an inhibitor of a mutant calreticulin for use as a medicament. Further, the present invention relates to the mutant calreticulin protein as provided herein, antibodies specifically binding thereto (preferably inhibitory antibodies), nucleic acids as provided herein (particularly nucleic acids encoding mutant calreticulin protein), the siRNA as provided herein for use as a medicament. The terms "medicament" and "pharmaceutical composition" are used interchangeably herein. Accordingly, definitions and explanations provided herein in relation to "pharmaceutical compositions", apply, mutatis mutandis, to the term "medicament".

[0228] The present invention relates to an inhibitor of a mutant calreticulin for use in the treatment of a myeloid malignancy.

[0229] The present invention provides a method for treating a myeloid malignancy patient comprising administering an effective amount of an inhibitor of a mutant calreticulin to the patient.

[0230] The patient to be treated can be a patient assessed "positive" in accordance with the present invention, i.e. a patient a sample of whom has been determined to have one or more mutant alleles of the calreticulin gene present.

[0231] Accordingly, the present invention relates to an inhibitor of a mutant calreticulin for use in the treatment of a myeloid malignancy, whereby a patient is to be treated, who suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy, whereby one or more mutant alleles of the calreticulin gene (or of a gene product thereof) in a sample from said patient is determined to be present.

[0232] All definitions and explanations given herein above in relation to the determination of the presence of one or more mutant alleles of the calreticulin gene (or of a gene product thereof) in a sample from said patient apply, mutatis mutandis, in this context.

[0233] The present disclosure provides a method for treating a myeloid malignancy patient comprising administering an effective amount of an inhibitor of a mutant calreticulin to the patient, the method further comprising assessing that the patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy, comprising

- determining the presence of one or more mutant alleles of the calreticulin gene (or of a gene product thereof) in a sample from said patient; and
- assessing that said patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy when said one or more mutant alleles of the calreticulin gene (or a gene product thereof) is present.

[0234] All definitions and explanations given herein above in relation to the determination of the presence of one or more mutant alleles of the calreticulin gene (or of a gene product thereof) in a sample from said patient apply, mutatis mutandis, in this context.

[0235] The mutant calreticulin can be a mutant calreticulin protein as defined herein above. If the mutant calreticulin is a mutant calreticulin protein, the inhibitor can be an antibody (preferably an inhibitory antibody), extracellular binding partners, small binding molecules, aptamers, or intramers.

[0236] The mutant calreticulin can be a nucleic acid encoding a mutant calreticulin protein as defined herein above. If the mutant calreticulin is such a nucleic acid (preferably mRNA as provided and defined above), the inhibitor can be siRNA, miRNA, dsRNA, shRNA, stRNA, and antisense molecules.

[0237] As mentioned above, myeloid malignancy includes a myeloproliferative neoplasm or a myelodysplastic syndrome. An exemplary myeloproliferative neoplasm is primary myelofibrosis (PMF). Preferred herein is the therapy of a myeloproliferative neoplasm, particularly that of primary myelofibrosis (PMF) or of essential thrombocythemia (ET). An exemplary myelodysplastic syndrome subject to therapeutic intervention in accordance with the present invention is refractory anemia with ringed sideroblasts and thrombocythemia (RARS-T).

[0238] The terms "treatment", "treating" and the like are used herein to generally mean obtaining a desired pharmacological and/or physiological effect. The effect may be prophylactic in terms of completely or partially preventing a disease or symptom thereof and/or may be therapeutic in terms of partially or completely curing a disease and/or adverse effect attributed to the disease. The term "treatment" as used herein covers any treatment

of a disease in a subject and includes: (a) preventing a disease related in a subject which may be predisposed to the disease; (b) inhibiting the disease, i.e. arresting its development; or (c) relieving the disease, i.e. causing regression of the disease.

[0239] An "individual", "patient" or "subject" for the purposes of the present invention includes both humans and other animals, particularly mammals, and other organisms. Thus, the methods are applicable to both human therapy and veterinary applications. Preferably, the "individual", "patient" or "subject" is a mammal, and most preferably the "individual", "patient" or "subject" is human.

[0240] The inhibitor of a mutant calreticulin may be administered as a single agent (i.e. in form of a monotherapy) or in form of a combination therapy, for example, conventional therapies like hydroxyurea or interferon alpha therapy.

[0241] The pharmaceutical composition will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient, the site of delivery of the pharmaceutical composition, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" of the pharmaceutical composition for purposes herein is thus determined by such considerations.

[0242] The skilled person knows that the effective amount of pharmaceutical composition administered to an individual will, inter alia, depend on the nature of the compound.

[0243] For example, if said inhibitor is a small molecule, the total (pharmaceutically) effective amount of the inhibitor in the pharmaceutical composition administered orally per dose will be in the range of about 50 mg inhibitor per day to 1000 mg inhibitor per day of patient, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 50 mg inhibitor per day, and most preferably for humans between about 50mg and 600 mg inhibitor per day. For example, an inhibitor may be administered at a dose of 15 mg/kg body weight per day. If given continuously, the inhibitor is typically administered at a dose rate of about 50 mg per day to about 600 mg per day. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect. The particular amounts may be determined by conventional tests which are well known to the person skilled in the art. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect. The particular amounts may be determined by conventional tests which are well known to the person skilled in the art.

[0244] The administration of the herein provided compositions may, inter alia, comprise an administration twice daily, every day, every other day, every third day, every fourth day, every fifth day, once a week, once every second week, once every third week, once every month, etc.

[0245] For example, if said compound is a (poly)peptide or protein the total pharmaceutically effective amount of pharmaceutical composition administered parenterally per dose will be in the range of about 1 µg protein /kg/day to 15 mg protein /kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg protein /kg/day, and most preferably for humans between about 0.01 and 1 mg protein /kg/day. If given continuously, the pharmaceutical composition is typically administered at a dose rate of about 1 µg/kg/hour to about 50 µg/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect. The particular amounts may be determined by conventional tests which are well known to the person skilled in the art.

[0246] Pharmaceutical compositions of the invention may be administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), buccally, or as an oral or nasal spray.

[0247] Pharmaceutical compositions of the invention preferably comprise a pharmaceutically acceptable carrier. By "pharmaceutically acceptable carrier" is meant a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

[0248] The pharmaceutical composition is also suitably administered by sustained release systems. Suitable examples of sustained-release compositions include semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. et al., *Biopolymers* 22:547-556 (1983)), poly (2-hydroxyethyl methacrylate) (R. Langer et al., *J. Biomed. Mater. Res.* 15:167-277 (1981), and R. Langer, *Chem. Tech.* 12:98-105 (1982)), ethylene vinyl acetate (R. Langer et al., *Id.*) or poly-D-(-)-3-hydroxybutyric acid (EP 133,988). Sustained release pharmaceutical composition also include liposomally entrapped compound. Liposomes containing the pharmaceutical composition are prepared by methods known per se: DE 3,218,121; Epstein et al., *Proc. Natl. Acad. Sci. (USA)* 82:3688-3692 (1985); Hwang et al., *Proc. Natl. Acad. Sci. (USA)* 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal therapy.

[0249] For parenteral administration, the pharmaceutical composition is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation.

[0250] Generally, the formulations are prepared by contacting the components of the pharmaceutical composition uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes. The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) (poly)peptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

[0251] The components of the pharmaceutical composition to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutic components of the pharmaceutical composition generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0252] The components of the pharmaceutical composition ordinarily will be stored in unit or multidose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized compound(s) using bacteriostatic Water-for-Injection.

[0253] The purpose of treatment for polycythemia vera is to reduce the number of extra blood cells. Treatment of polycythemia vera may include, phlebotomy, chemotherapy with or without phlebotomy, biologic therapy using

interferon alfa or pegylated interferon alpha and low-dose aspirin.


[0254] The treatment of primary myelofibrosis in patients without signs or symptoms is usually watchful waiting. Patients with primary myelofibrosis may have signs or symptoms of anemia. Anemia is usually treated with transfusion of red blood cells to relieve symptoms and improve quality of life. In addition, anemia may be treated with erythropoietic growth factors, prednisone, danazol, thalidomide, lenalidomide, or pomalidomide. Treatment of primary myelofibrosis in patients with other signs or symptoms may include targeted therapy with ruxolitinib (a JAK1 and JAK2 inhibitor), chemotherapy, donor stem cell transplant, thalidomide, lenalidomide, or pomalidomide, splenectomy, radiation therapy to the spleen, lymph nodes, or other areas outside the bone marrow where blood cells are forming, biologic therapy using interferon alfa or erythropoietic growth factors, or the inclusion in a clinical trial of other targeted therapy drugs.

[0255] Treatment of essential thrombocythemia in patients younger than 60 years who have no signs or symptoms and an acceptable platelet count is usually watchful waiting. In some cases, the patient can take aspirin to help prevent blood clots. Treatment of other patients may include Chemotherapy, hydroxyurea, Anagrelide therapy, biologic therapy using interferon alfa or pegylated interferon alpha, platelet apheresis.

[0256] The JAK-binding inhibitor ruxolitinib shows promise for curative and supportive treatment. Ruxolitinib has been approved by the Food and Drug Administration) for use in the treatment of high and intermediate risk myelofibrosis in 2011; see Tefferi March 22, 2012; Blood: 119 (12) Also Ostojic reports that ruxolitinib is used in the therapy of myelofibrosis; see Ostojic Therapeutics and Clinical Risk Management 2012:8 95-103.

[0257] JAK inhibitors that are currently used in clinical trials for myeloproliferative neoplasms include, besides ruxolitinib, SAR302503, CYT387, lestaurtinib, SB1518, AZD1480, BMS911543, LY2784544, NS-018, and XL019; see Tefferi March 22, 2012; Blood: 119 (12).

[0258] An exemplary formula of ruxolitinib ((3R)-3-cyclopentyl-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)pyrazol-1-yl]propanenitrile; trade name Jakafi, Jakavi) is shown below:



[0259] Refractory anemia with ringed sideroblast and thrombocytosis may require blood transfusions and other supportive therapy to remedy anemia, including high doses of pyridoxine (Vitamin B6). Bone marrow transplant is also an option. RARS-T may also progress to leukemia.

[0260] The use of above therapies is contemplated for patients diagnosed positive or negative for the presence of mutant calreticulin in accordance with the present invention, either alone or in combination with therapies (e.g. antibodies) specifically targeting the mutant calreticulin. Accordingly, therapies (e.g. antibodies) that target mutant CALR, can likewise be useful in treatment if used as monotherapy or in combination with other therapies. For example, interferon alfa therapy can be used to treat patients with MPN (like essential thrombocythemia patients)

diagnosed positive for the presence of mutant calreticulin in accordance with the present invention.

[0261] If, for example, the patient is tested positive for the presence of mutant calreticulin and (a) JAK2 mutation(s), the use of JAK inhibitor(s) (like ruxolitinib) is contemplated herein. Depending on clinical parameters, (e.g. age, prognosis of the patient) also further therapies, like stem cell transplantation can be used to treat e.g. a patient tested positive for the presence of mutant calreticulin.

[0262] Inhibitors for use in accordance with the present invention are described and provided herein. Also the use of inhibitors yet to be generated or known compounds to be tested for their inhibiting activity is envisaged in context of the present invention.

[0263] Therefore, the present invention provides a method for assessing the activity of a candidate molecule suspected of being an inhibitor of a mutant calreticulin as defined and provided herein comprising the steps of:

1. (a) contacting a cell, tissue or a non-human animal comprising a mutant calreticulin with said candidate molecule;
2. (b) detecting a decrease in activity of said mutant calreticulin; and
3. (c) selecting a candidate molecule that decreases activity of said mutant calreticulin;
wherein a decrease of the activity is indicative for the capacity of the selected molecule to be useful in the treatment of myeloid malignancy as defined herein.

[0264] Also a decrease in the (expression) level can indicate useful inhibitors.

[0265] The present invention relates to a method for assessing the activity of a candidate molecule suspected of being an inhibitor of a mutant calreticulin as defined and provided herein comprising the steps of:

1. (a) contacting a cell, tissue or a non-human animal comprising a mutant calreticulin with said candidate molecule;
2. (b) detecting a decrease in the (expression) level of said mutant calreticulin; and
3. (c) selecting a candidate molecule that decreases the (expression) level of said mutant calreticulin;
wherein a decrease of the (expression) level is indicative for the capacity of the selected molecule to be useful in the treatment of myeloid malignancy as defined herein.

[0266] The mutant calreticulin can be any of the mutant calreticulin proteins/polypeptides as defined herein above or any of the nucleic acids (particularly mRNAs) as defined herein, which encode the mutant calreticulin proteins/polypeptides.

[0267] The mutant calreticulin proteins can comprise or consist of the amino acid sequence of the polypeptides shown in 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430, and 434, respectively. Also the use of fragments or derivatives of these proteins as defined above is envisaged in this context.

[0268] The mutant calreticulin nucleic acids can comprise or consist of the nucleic acids shown in 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, 143, 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183,

185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, and 433, respectively.

[0269] Further, the present invention provides the use of (a) nucleic acid(s) (e.g. oligonucleotide(s) like primer(s)/primer pair(s) or probe(s) as described below) or antibody capable of detecting the presence of one or more mutant alleles of the calreticulin gene or the presence or amount of a gene product of one or more mutant alleles of the calreticulin gene as defined herein for the methods of the present invention, i.e. for assessing whether a patient suffers from a myeloid malignancy or is prone to suffering from a myeloid malignancy.

[0270] The oligonucleotide(s) may be about 15 to 100 nucleotides in length. A person skilled in the art is, based on his general knowledge and the teaching provided herein, easily in the position to identify and/or prepare (a) an oligo- or polynucleotide capable of detecting the presence of one or more mutant alleles of the calreticulin gene or the presence or amount of a gene product of one or more mutant alleles of the calreticulin gene as defined herein. These oligo- or polynucleotides may be used as probe(s) or primers in the methods provided herein. Often primers and/or probes have a length of 10 to 30 nucleotides. Accordingly, the invention relates to (a) nucleic acid(s) (in particular (a) primer(s) or (a) probe(s)) capable of detecting the presence of one or more mutant alleles of the calreticulin gene or the presence or amount of a gene product of one or more mutant alleles of the calreticulin gene as defined herein, wherein said nucleic acid is smaller than 50, 49, 48, 47, 46, 45, 44, 43, 42, 41, 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, or 20 nucleotides and is larger than 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 nucleotides. Preferably, the nucleic acid has a length of 10 to 35 nucleotides, more preferably 15 to 25 nucleotides, particularly preferred a length of 18 to 21, e.g. 18, 19, 20 or 21 nucleotides. These nucleic acid(s) may hybridize under stringent conditions to the complementary strand of the nucleic acid mutant calreticulin as defined and provided herein above.

[0271] A skilled person will know, for example, computer programs which may be useful for the identification of corresponding probes/primers to be used herein. For example, the nucleic acid sequence(s) of exemplary coding sequences as disclosed herein (SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, 143, 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432, and 433, respectively) may be used in this context.

[0272] Also the sequences disclosed in the following table may be used in this context:

Sequences of mutation junctions in the cDNA sequence of CALR for the design of mutation specific probes or PCR primers.

CALR mutation	cDNA junction sequences in mutated positions
Type 1	GAAGGACAAACAGGACGAGG AG CAGAGGACAAGGAGGATGAT
Type 2	GAGGAGGAGGCAGAGGACA ATTGTC GGAGGATGATGAGGACAAAG
Type 3	GGACAAACAGGACGAGGAGC AG AGGCAGAGGACAAGGAGGAT
Type 4	CAGGACGAGGAGCAGAGGCT TAG GAGGAGGCAGAGGACAAGG
Type 5	TGAAGGACAAACAGGACGAG GGG CAGAGGACAAGGAGGATGA
Type 6	AGGACAAACAGGACGAGGAG CG GAGGCAGAGGACAAGGAGGA
Type 7	CAGGACGAGGAGCAGAGGCT TAG GAGGATGATGAGGACAAAG
Type 8	GGACGAGGAGCAGAGGCT TAA GAGGAGGCAGAGGACAAGGAG
Type 9	CAAGAAACGCAAAGAGGAGG AG AGGCAGAGGACAAGGAGGAT
Type 10	AGGAGGAGGAGGCAGAGGAC ATGTGT CGGAGGATGATGAGGACAAAG
Type 11	AAGGACAAACAGGACGAGGACC AG AGGCAGAGGACAAGGAGGAT
Type 12	CAAACAGGACGAGGAGCAGAGG AGG AGGAGGAGGCAGAGGAC
Type 13	AACAGGACGAGGAGCAGAGGC AG AGGAGGAGGCAGAGGACAAG
Type 14	ACAGGACGAGGAGCAGAGGCT TG AGGAGGAGGCAGAGGACAAG
Type 15	CAGGACGAGGAGCAGAGGCT TAG GAGGAGG G AGAGGACAAGGAGGATGATG
Type 16	CAGGACGAGGAGCAGAGGCT TCAG AGGAGGCAGAGGACAAGGAG
Type 17	GGACGAGGAGCAGAGGCT TAA GAGGAGGCAG TGG ACAAGGAGGATGATGAGG
Type 18	GGACGAGGAGCAGAGGCT TAA GAGGATGATGAGGACAAAGAT
Type 19	GGAGCAGAGGCTTAAGGAGG AG AGGCAGAGGACAAGGAGGAT
Type 20	GGCTTAAGGAGGAGGAAGA AGG AGGAGGCAGAGGACAAGGA
Type 21	GGCTTAAGGAGGAGGAAGA AGCGTTTAA GAGGACAAGGAGGATGATGA
Type 22	CTTAAGGAGGAGGAAGAAGAC CAAC GCAAAGAGGAGGAGGAGG
Type 23	CTTAAGGAGGAGGAAGAAGAC CTGCGT GAGGAGGAGGAGGCAGAGGAC
Type 24	CTTAAGGAGGAGGAAGAAGAC CAG GAGGCAGAGGACAAGGAGG
Type 25	TAAGGAGGAGGAAGAAGACAA AA AGGCAGAGGACAAGGAGGATG
Type 26	TAAGGAGGAGGAAGAAGACAA AAA ACGCAAAGAGGAGGAGGAG
Type 27	AAGGAGGAGGAAGAAGACAA GTTGTTT CGCAAAGAGGAGGAGGAGGCA
Type 28	GGAAGAAGACAAGAAACGCA AA AGGAGGATGATGAGGACAAA
Type 29	GAAGACAAGAAACGCAAAGAG GCCTCCTCTTTGTCT AAGGAGGATGATGAGGACAAA
Type 30	AGACAAGAAACGCAAAGAGG ACCATCCTTGT CGGAGGATGATGAGGACAAAGA
Type 31	AGAGGAGGAGGAGGCAGAGG GCAATTGT CGGAGGATGATGAGGACAAAG
Type 32	GAGGAGGAGGAGGCAGAGGAC CTGT CGGAGGATGATGAGGACAAAGA
Type 33	GAGGAGGAGGCAGAGGACAA ATTGT CGGAGGATGATGAGGACAAAG
Type 34	AGGAGGAGGAGGCAGAGGAC ACTTGT CGGAGGATGATGAGGACAAAGA
Type 35	AGGAGGAGGAGGCAGAGGAC ATTTGT CGGAGGATGATGAGGACAAAGA
Type 36	AGGAGGAGGCAGAGGACAAG TGT CGGAGGATGATGAGGACAAAGA

Bold letters indicate the borders of a deletion event; underlined letters indicate inserted sequences; Bold and italic letters indicate single nucleotide variants

[0273] Exemplary oligonucleotides (primers) provided herein and to be used in accordance with the present invention are

Forward: ACAACTTCCTCATACCAACG (SEQ ID NO: 437)

and/or

Reverse: GGCCTCAGTCCAGCCCTG (SEQ ID NO: 438)

Forward: GGCAAGGCCCTGAGGTGT (SEQ ID NO: 439)

and/or

Reverse: GGCCTCAGTCCAGCCCTG (SEQ ID NO: 438)

[0274] In one embodiment, the present invention relates to transgenic cell or a transgenic non-human animal having the nucleic acid as described and explained herein (or a vector comprising same), e.g. a nucleic acid comprising at least one or more mutant alleles of the calreticulin gene as defined herein. Such transgenic cell(s) or a transgenic non-human animal(s) can be used for screening and/or validation of a medicament for the treatment of a myeloid malignancy.

[0275] The term "cell" as used in this context may also comprise a plurality of cells as well as cells comprised in a tissue. The cell to be used in the screening or validation method may be obtained from samples from a (transgenic) non-human animal or human suffering from a myeloid malignancy. The tumor cell or cell may also be obtained from patient samples (e.g. biopsies), in particular a biopsy/biopsies from a patient/subject suffering from a myeloid malignancy. Accordingly, the cell may be a human cell. Again, such a cell to be used in the present screening or validation methods may be comprised in a tissue or tissue sample, like in a sample biopsy.

[0276] The used non-human animal or cell may be transgenic or non transgenic. "Transgenic" in this context particularly means that at least one of the mutant calreticulins as described or defined herein is (over-) expressed and/or that the activity of at least one of the mutant calreticulins is present (or increased).

[0277] A preferred (transgenic) non-human animal or (transgenic) cell in context of the invention suffers from a myeloid malignancy.

[0278] The term "transgenic non-human animal" or "transgenic cell" as used herein refers to an non-human animal or cell, not being a human, that comprises genetic material different from the genetic material of a corresponding wild-type animal/cell. "Genetic material" in this context may be any kind of a nucleic acid molecule, or analogues thereof, for example a nucleic acid molecule, or analogues thereof as defined herein. "Different" in this context means additional or fewer genetic material with respect to the genome of the wild-type animal/cell and/or rearranged genetic material, i.e. genetic material present at a different locus of the genome with respect to the genome of the wild-type animal/cell. An overview of examples of different expression systems to be used for generating transgenic cell/animal is, for instance, contained in Methods in Enzymology 153 (1987), 385-516, in Bitter et al. (Methods in Enzymology 153 (1987), 516-544) and in Sawers et al. (Applied Microbiology and Biotechnology 46 (1996), 1-9), Billman-Jacobe (Current Opinion in Biotechnology 7 (1996), 500-4), Hockney (Trends in Biotechnology 12 (1994), 456-463), Griffiths et al., (Methods in Molecular Biology 75 (1997), 427-440).

[0279] In a preferred embodiment, the (transgenic) non-human animal or (transgenic) cell is or is derived from a mammal. Non-limiting examples of the (transgenic) non-human animal or derived (transgenic) cell are selected from the group consisting of a mouse, a rat, a rabbit, and a guinea pig.

[0280] The present invention also relates to a vector comprising the nucleic acid molecule of the present invention.

[0281] Many suitable vectors are known to those skilled in molecular biology, the choice of which would depend on the function desired and include plasmids, cosmids, viruses, bacteriophages and other vectors used conventionally in genetic engineering. Methods which are well known to those skilled in the art can be used to construct various plasmids and vectors; see, for example, the techniques described in Sambrook et al. (loc cit.) and Ausubel, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. (1989), (1994). Alternatively, the polynucleotides and vectors of the invention can be reconstituted into liposomes for delivery to

target cells. As discussed in further details below, a cloning vector was used to isolate individual sequences of DNA. Relevant sequences can be transferred into expression vectors where expression of a particular polypeptide is required. Typical cloning vectors include pBluescript SK, pGEM, pUC9, pBR322 and pGBT9. Typical expression vectors include pTRE, pCAL-n-EK, pESP-1, pOP13CAT.

[0282] Preferably said vector comprises a nucleic acid sequence which is a regulatory sequence operably linked to said nucleic acid sequence defined herein.

[0283] The term "regulatory sequence" refers to DNA sequences, which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism. In prokaryotes, control sequences generally include promoter, ribosomal binding site, and terminators. In eukaryotes generally control sequences include promoters, terminators and, in some instances, enhancers, transactivators or transcription factors. The term "control sequence" is intended to include, at a minimum, all components the presence of which are necessary for expression, and may also include additional advantageous components.

[0284] The term "operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences. In case the control sequence is a promoter, it is obvious for a skilled person that double-stranded nucleic acid is preferably used.

[0285] Thus, the recited vector is preferably an expression vector. An "expression vector" is a construct that can be used to transform a selected host and provides for expression of a coding sequence in the selected host. Expression vectors can for instance be cloning vectors, binary vectors or integrating vectors. Expression comprises transcription of the nucleic acid molecule preferably into a translatable mRNA. Regulatory elements ensuring expression in prokaryotes and/or eukaryotic cells are well known to those skilled in the art. In the case of eukaryotic cells they comprise normally promoters ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Possible regulatory elements permitting expression in prokaryotic host cells comprise, e.g., the P_L , *lac*, *trp* or *tac* promoter in *E. coli*, and examples of regulatory elements permitting expression in eukaryotic host cells are the *AOX1* or *GAL1* promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells.

[0286] Beside elements, which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. Furthermore, depending on the expression system used leader sequences capable of directing the polypeptide to a cellular compartment or secreting it into the medium may be added to the coding sequence of the recited nucleic acid sequence and are well known in the art; see also the appended Examples. The leader sequence(s) is (are) assembled in appropriate phase with translation, initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein, or a portion thereof, into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product; see supra. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3 (In-vitrogene), pEF-DHFR, pEF-ADA or pEF-neo (Mack et al. PNAS (1995) 92, 7021-7025 and Raum et al. Cancer Immunol Immunother (2001) 50(3), 141-150) or pSPORT1 (GIBCO BRL).

[0287] Preferably, the expression control sequences will be eukaryotic promoter systems in vectors capable of transforming of transfecting eukaryotic host cells, but control sequences for prokaryotic hosts may also be used. Once the vector has been incorporated into the appropriate host, the host is maintained under conditions suitable for high level expression of the nucleotide sequences, and as desired, the collection and purification of the polypeptide of the invention may follow; see, e.g., the appended examples.

[0288] An alternative expression system, which can be used to express a cell cycle interacting protein is an insect system. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The coding sequence of a recited nucleic acid molecule may be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of said coding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein coat. The recombinant viruses are then used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which the protein of the invention is expressed (Smith, J. Virol. 46 (1983), 584; Engelhard, Proc. Natl. Acad. Sci. USA 91 (1994), 3224-3227).

[0289] Additional regulatory elements may include transcriptional as well as translational enhancers. Advantageously, the above-described vectors of the invention comprise a selectable and/or scorable marker.

[0290] Selectable marker genes useful for the selection of transformed cells and, e.g., plant tissue and plants are well known to those skilled in the art and comprise, for example, antimetabolite resistance as the basis of selection for dhfr, which confers resistance to methotrexate (Reiss, Plant Physiol. (Life Sci. Adv.) 13 (1994), 143-149); npt, which confers resistance to the aminoglycosides neomycin, kanamycin and paromycin (Herrera-Estrella, EMBO J. 2 (1983), 987-995) and hygromycin (Marsh, Gene 32 (1984), 481-485). Additional selectable genes have been described, namely trpB, which allows cells to utilize indole in place of tryptophan; hisD, which allows cells to utilize histinol in place of histidine (Hartman, Proc. Natl. Acad. Sci. USA 85 (1988), 8047); mannose-6-phosphate isomerase which allows cells to utilize mannose (WO 94/20627) and ODC (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue, 1987, In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory ed.) or deaminase from *Aspergillus terreus* which confers resistance to Blastidicin S (Tamura, Biosci. Biotechnol. Biochem. 59 (1995), 2336-2338).

[0291] Useful scorable markers are also known to those skilled in the art and are commercially available. Advantageously, said marker is a gene encoding luciferase (Giacomin, Pl. Sci. 116 (1996), 59-72; Scikantha, J. Bact. 178 (1996), 121), green fluorescent protein (Gerdes, FEBS Lett. 389 (1996), 44-47) or β -glucuronidase (Jefferson, EMBO J. 6 (1987), 3901-3907). This embodiment is particularly useful for simple and rapid screening of cells, tissues and organisms containing a recited vector.

[0292] As described above, the recited nucleic acid molecule can be used alone or as part of a vector to express the polypeptide of the invention in cells, for, e.g., purification but also for gene therapy purposes. The nucleic acid molecules or vectors containing the DNA sequence(s) encoding any one of the above described polypeptide of the invention is introduced into the cells which in turn produce the polypeptide of interest. Gene therapy, which is based on introducing therapeutic genes into cells by ex-vivo or in-vivo techniques is one of the most important applications of gene transfer. Suitable vectors, methods or gene-delivery systems for in-vitro or in-vivo gene therapy are described in the literature and are known to the person skilled in the art; see, e.g., Giordano, Nature Medicine 2 (1996), 534-539; Schaper, Circ. Res. 79 (1996), 911-919; Anderson, Science 256 (1992), 808-813; Verma, Nature 389 (1994), 239; Isner, Lancet 348 (1996), 370-374; Muhlhauser, Circ. Res. 77 (1995), 1077-1086; Onodera, Blood 91 (1998), 30-36; Verma, Gene Ther. 5 (1998), 692-699; Nabel, Ann. N.Y. Acad. Sci. 811 (1997), 289-292; Verzeletti, Hum. Gene Ther. 9 (1998), 2243-51; Wang, Nature Medicine 2 (1996), 714-716; WO 94/29469; WO 97/00957, US 5,580,859; US 5,589,466; or Schaper, Current Opinion in Biotechnology 7 (1996), 635-640. The recited nucleic acid molecules and vectors may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g., adenoviral, retroviral) into the cell. Preferably, said cell is a germ line cell, embryonic cell, or egg cell or derived there from, most preferably said cell is a stem cell. An example for an embryonic stem cell can be, inter alia, a stem cell as described in Nagy, Proc. Natl. Acad. Sci. USA 90 (1993), 8424-8428.

[0293] The invention also provides for a host transformed or transfected with a vector of the invention. Said host may be produced by introducing the above described vector of the invention or the above described nucleic acid molecule of the invention into the host. The presence of at least one vector or at least one nucleic acid molecule in the host may mediate the expression of a gene encoding the above described single chain antibody constructs.

[0294] The described nucleic acid molecule or vector of the invention, which is introduced in the host may either

integrate into the genome of the host or it may be maintained extrachromosomally.
The host can be any prokaryotic or eukaryotic cell.

[0295] The term "prokaryote" is meant to include all bacteria, which can be transformed or transfected with DNA or RNA molecules for the expression of a protein of the invention. Prokaryotic hosts may include gram negative as well as gram positive bacteria such as, for example, *E. coli*, *S. typhimurium*, *Serratia marcescens* and *Bacillus subtilis*. The term "eukaryotic" is meant to include yeast, higher plant, insect and preferably mammalian cells. Depending upon the host employed in a recombinant production procedure, the protein encoded by the polynucleotide of the present invention may be glycosylated or may be non-glycosylated. Especially preferred is the use of a plasmid or a virus containing the coding sequence of the polypeptide of the invention and genetically fused thereto an N-terminal FLAG-tag and/or C-terminal His-tag. Preferably, the length of said FLAG-tag is about 4 to 8 amino acids, most preferably 8 amino acids. An above described polynucleotide can be used to transform or transfect the host using any of the techniques commonly known to those of ordinary skill in the art. Furthermore, methods for preparing fused, operably linked genes and expressing them in, e.g., mammalian cells and bacteria are well-known in the art (Sambrook, loc cit.).

[0296] Preferably, said the host is a bacterium or an insect, fungal, plant or animal cell.

[0297] It is particularly envisaged that the recited host may be a mammalian cell. Particularly preferred host cells comprise CHO cells, COS cells, myeloma cell lines like SP2/0 or NS/0. As illustrated in the appended examples, particularly preferred are CHO-cells as hosts.

[0298] More preferably said host cell is a human cell or human cell line, e.g. per.c6 (Kroos, *Biotechnol. Prog.*, 2003, 19:163-168).

[0299] In a further embodiment, the present invention thus relates to a process for the production of a polypeptide of the invention, said process comprising culturing a host of the invention under conditions allowing the expression of the polypeptide of the invention and recovering the produced polypeptide from the culture.

[0300] The transformed hosts can be grown in fermentors and cultured according to techniques known in the art to achieve optimal cell growth. The polypeptide of the invention can then be isolated from the growth medium, cellular lysates, or cellular membrane fractions. The isolation and purification of the, e.g., microbially expressed polypeptides of the invention may be by any conventional means such as, for example, preparative chromatographic separations and immunological separations such as those involving the use of monoclonal or polyclonal antibodies directed, e.g., against a tag of the polypeptide of the invention or as described in the appended examples.

[0301] The conditions for the culturing of a host, which allow the expression are known in the art to depend on the host system and the expression system/vector used in such process. The parameters to be modified in order to achieve conditions allowing the expression of a recombinant polypeptide are known in the art. Thus, suitable conditions can be determined by the person skilled in the art in the absence of further inventive input.

[0302] Once expressed, the polypeptide of the invention can be purified according to standard procedures of the art, including ammonium sulfate precipitation, affinity columns, column chromatography, gel electrophoresis and the like; see, Scopes, "Protein Purification", Springer-Verlag, N.Y. (1982). Substantially pure polypeptides of at least about 90 to 95% homogeneity are preferred, and 98 to 99% or more homogeneity are most preferred, for pharmaceutical uses. Once purified, partially or to homogeneity as desired, the polypeptide of the invention may then be used therapeutically (including extracorporeally) or in developing and performing assay procedures. Furthermore, examples for methods for the recovery of the polypeptide of the invention from a culture are described in detail in the appended examples.

[0303] Furthermore, the present invention provides a kit useful for carrying out the methods of the invention, the kit comprising a nucleic acid or an antibody capable of detecting the presence of a one or more mutant alleles of the calreticulin gene or the presence or amount of a gene product of one or more mutant alleles of the calreticulin gene

as defined herein. Also envisaged herein is the use of the herein described kit for carrying out the herein provided methods.

[0304] For example, said kit may comprise (a) compound(s) required for specifically determining the presence of one or more mutant alleles of the calreticulin gene or the presence or amount of a gene product of one or more mutant alleles of the calreticulin gene as defined herein. Moreover, the present invention also relates to the use of (a) compound(s) required for specifically determining the presence of one or more mutant alleles of the calreticulin gene or the presence or amount of a gene product of one or more mutant alleles of the calreticulin gene as defined herein for the preparation of a kit for carrying out the methods of this invention. On the basis of the teaching of this invention, the skilled person knows which compound(s) is (are) required for specifically determining the presence of one or more mutant alleles of the calreticulin gene or the presence or amount of a gene product of one or more mutant alleles of the calreticulin gene as defined herein. For example, such compound(s) may be (a) "binding molecule(s)", like, for example, (a) antibody. Particularly, such compound(s) may be (a) (nucleotide) probe(s), (a) primer(s) (pair(s)), (an) antibody(ies) and/or (an) aptamer(s) specific for at least one mutant allele of the calreticulin gene or for a gene product of at least one mutant alleles of the calreticulin gene as defined herein. The kit (to be prepared in context) of this invention may be a diagnostic kit. As mentioned above, the determination of the presence of one (or more) mutant alleles of the calreticulin gene or of a gene product thereof as described herein can be performed as a stand-alone analysis. Alternatively, this analysis can be followed or preceded by the analysis of other markers for myeloid malignancies, such as JAK2 and MPL mutations. Also simultaneous determination of such markers is envisaged, like the simultaneous test for JAK2 mutation(s) and mutant calreticulin (and, optionally, further markers), or the simultaneous test of JAK2 mutation(s), mutant calreticulin and MPL mutation(s) (and, optionally, further markers). According (a) kit(s) (or uses of such kits) is/are envisaged herein that provide means for such subsequent or simultaneous tests. For example, said kit may comprise, in addition to (a) compound(s) required for specifically determining the presence (or amount) of one or more mutant alleles of the calreticulin gene (or of a gene product thereof), (a) compound(s) required for specifically determining the presence as JAK2 and/or MPL mutations (and optionally further markers), e.g. (a) antibody(ies), (a) (nucleotide) probe(s), (a) primer(s) (pair(s)), (an) antibody(ies) and/or (an) aptamer(s) specific that allow the specific detection of JAK2 and MPL mutations (and optionally further markers).

[0305] The kit (to be prepared in context) of this invention may further comprise or be provided with (an) instruction manual(s). For example, said instruction manual(s) may guide the skilled person (how) to determine the presence of one or more mutant alleles of the calreticulin gene or the presence or amount of a gene product of one or more mutant alleles of the calreticulin gene as defined herein and/or (how) to diagnose myeloid malignancy. Said instruction manual(s) may comprise guidance to use or apply the herein provided methods or uses. The kit (to be prepared in context) of this invention may further comprise substances/chemicals and/or equipment suitable/required for carrying out the methods and uses of this invention. For example, such substances/chemicals and/or equipment are solvents, diluents and/or buffers for stabilizing and/or storing (a) compound(s) required for specifically determining the presence of one or more mutant alleles of the calreticulin gene or the presence or amount of a gene product of one or more mutant alleles of the calreticulin gene as defined herein.

[0306] As used herein, the terms "comprising" and "including" or grammatical variants thereof are to be taken as specifying the stated features, integers, steps or components but do not preclude the addition of one or more additional features, integers, steps, components or groups thereof. This term encompasses the terms "consisting of" and "consisting essentially of."

Thus, the terms "comprising"/"including"/"having" mean that any further component (or likewise features, integers, steps and the like) can be present.

[0307] The term "consisting of" means that no further component (or likewise features, integers, steps and the like) can be present.

[0308] The term "consisting essentially of" or grammatical variants thereof when used herein are to be taken as specifying the stated features, integers, steps or components but do not preclude the addition of one or more additional features, integers, steps, components or groups thereof but only if the additional features, integers, steps, components or groups thereof do not materially alter the basic and novel characteristics of the claimed composition,

device or method.

[0309] Thus, the term "consisting essentially of" means that specific further components (or likewise features, integers, steps and the like) can be present, namely those not materially affecting the essential characteristics of the composition, device or method. In other words, the term "consisting essentially of" (which can be interchangeably used herein with the term "comprising substantially"), allows the presence of other components in the composition, device or method in addition to the mandatory components (or likewise features, integers, steps and the like), provided that the essential characteristics of the device or method are not materially affected by the presence of other components.

[0310] The term "method" refers to manners, means, techniques and procedures for accomplishing a given task including, but not limited to, those manners, means, techniques and procedures either known to, or readily developed from known manners, means, techniques and procedures by practitioners of the chemical, biological and biophysical arts.

[0311] As used herein, the term "isolated" refers to a composition that has been removed from its in-vivo location. Preferably the isolated compositions or compounds of the present invention are substantially free from other substances (e.g., other proteins or other compounds) that are present in their in-vivo location (i.e. purified or semi-purified compositions or compounds.)

[0312] As used herein the term "about" refers to $\pm 10\%$.

[0313] The present invention is further described by reference to the following non-limiting figures and examples.

[0314] The Figures show:

Figure 1.

Validation of *CALR* gene mutations in patients 191 and 296 by Sanger sequencing

Depicted are Sanger sequencing traces. Boxes around the sequence letters mark the bases that are lost due to the deletion events. Fig. 1 shows SEQ ID NOs: 1324 to 1327.

Figure 2.

Frequency of *CALR* mutations in myeloid malignancies

A: Distribution of JAK2, MPL and *CALR* mutations in the three classical MPN disease entities. B: The frequency of patients with mutant *CALR* in different myeloid malignancies, is indicated by the dark bars. N, total numbers of patients analysed for a specific disease entity. C: Distribution of JAK2, MPL, *CALR* and SF3B1 mutations in 24 patients with refractory anemia with ring sideroblasts associated with marked thrombocytosis. PV, polycythemia vera; ET, essential thrombocythemia; PMF primary myelofibrosis; dnAML, de novo AML; CML, chronic myeloid leukaemia; MDS, myelodysplastic syndrome; CMML, chronic myelomonocytic leukaemia; RARS-T, refractory anemia with ring sideroblasts associated with marked thrombocytosis.

Figure 3.

Mutational pattern of *CALR* mutations in MPN patients.

The wide black bar represents exon 9 of *CALR*, the narrow bar the 3' UTR of the gene, the thin line intronic and intergenic regions.

A: indicated are the cDNA sequence in the beginning and end of exon 9. Below the cDNA sequence are the amino acid sequences derived from the three alternative reading frames. **B:** The three reading frames result in different peptide compositions, especially with respect to the charge of amino acids. **C:** Summary of all mutations detected in MPN patients and their position within *CALR* exon 9. Bars indicate deletion events, letters inserted sequences. Independent insertions and deletions are depicted above the exon 9 scheme, combined insertion/deletion events below. **D:** The specific peptide makeup of wild type *CALR* and of the two most frequently detected types of mutations. **B, D:** Each box represents an amino acid. Black boxes with '-' sign are negatively charged amino acids, boxes with '+' sign are positively charged amino acids. Crossed boxes represent stop codons. **E:** Relative frequencies of all 36 mutation types observed in *CALR*.

Fig. 3 shows SEQ ID NOs: 1328 to 1339.

Figure 4.**Association of CALR mutations with uniparental disomies and clonal hierarchies in patients with multiple somatic mutations**

Panel A: Affymetrix SNP 6.0 array data is shown for three samples that had more than 50% burden of *CALR* mutations. Each dot represents a single SNP. The x-axis shows the genomic position, the y-axis depicts the allelic status of the SNP. (heterozygous status = allelic difference of 0). The array data shows 19p uniparental disomy of different sizes in the three samples. Boxes indicate the genomic region of the uniparental disomies. To the right of each allelic difference plot, the results of the *CALR* exon 9 fragment analysis for the same sample is shown. In each case the burden of the mutant allele is higher than the wild type allele burden. Panel B: Clonal hierarchies derived from the analysis of hematopoietic progenitor colonies. Patient H_0191 (top) harbored somatic mutations in a total of 4 genes. As shown in the bar chart, 51% of the colonies had mutations in *CALR*, *GAB2* and *METTL11B*. The other half of the colonies (48%) had mutations in all four genes indicating that the mutation in *PHF16* was acquired later and gave rise to a subclone. Additionally, one colony harbored a one base pair deletion in *CALR*, different from the 52 base pair deletion that was observed in the granulocyte sample and all other colonies of this patient. As this colony had none of the other mutations observed in the patient it represents an independent clone although this conclusion is based only on a single colony. Patient H_0296 (bottom) had somatic mutations in *CALR*, *PIK3R1* and *C100RF71*. All colonies analyzed for this patient had all three mutations. One colony showed an 18 base pair deletion in *CALR* in addition to the 1 base pair deletion observed in this patient. Both mutations were on the same allele.

Figure 5.**Clinical significance of CALR mutations**

Outcome estimates in patients with essential thrombocythemia or primary myelofibrosis stratified according to their somatic mutation. Panel A: Kaplan-Meier analysis of overall survival in patients with primary myelofibrosis. Subgroups were compared using the log-rank test. Patients with myelofibrosis carrying a somatic mutation of *CALR* had a better overall survival than those with JAK2-V617F [median value 21.4 years (95% CI, 17.1-22.9) vs. 11.0 years (95% CI, 7.8-14.4), respectively; $P < 0.001$] or *MPL* mutation [median value 8.2 years (95% CI, 2.0-not reached); $P < 0.001$], while no difference was observed between the latter 2 subgroups. Panel B: Kaplan-Meier analysis of overall survival in patients with essential thrombocythemia. Subgroups were compared using the log-rank test. Median value for overall survival was not reached in any subgroup. Patients with *CALR* mutation had better overall survival compared with those carrying JAK2-V617F. The 10-year overall survival was 96.9% (95% CI, 91.7-98.8%) in the former vs. 91.1% (95% CI, 87.1-93.9%) in the latter ($P = 0.043$). Panel C: Cumulative incidence of thrombosis in patients with essential thrombocythemia. Death in absence of the event of interest was considered as a competing event, and subgroups were compared with the Pepe and Mori's test. Patients carrying a *CALR* exon 9 mutation had a lower cumulative incidence of thrombosis compared with those carrying JAK2-V617F: the 10-year cumulative incidence was 11.0% (95% CI, 6.3-17.1%) in the former vs. 21.0% (95% CI, 16.6-25.7%) in the latter ($P = 0.003$).

Figure 6.**Functional analysis of CALR type 1 mutation**

Panel A: Cell viability of Ba/F3 cells expressing an empty vector (GFP), wild type *CALR* (*CALR* wt-GFP) or mutant *CALR* (*CALR* de152-GFP) was assessed after 72 hours in presence of increasing interleukin-3 concentration. RLU, relative luminescence units. Error bars represent standard error of the mean. Panel B: Cell proliferation of Ba/F3 cells expressing an empty vector (GFP), wild type *CALR* (*CALR* wt-GFP) or mutant *CALR* (*CALR* de152-GFP), in the absence of interleukin-3 was determined for 7 days. Error bars represent standard error of the mean. Panel C: shows the activation of STAT5 in response to interleukin-3. Ba/F3 cells expressing the empty vector (GFP), the wild type *CALR* (wt), or the *CALR* mutant (de152) were starved for 4 hours in serum free medium without interleukin-3 and were then stimulated for 20 minutes with 100 pg/ml and 1 ng/ml of interleukin-3, as indicated. Western blot was performed on the cell lysates with antibodies against pYSTAT5, STAT5, and *CALR*. An antibody against GAPDH was used as loading control. Panel D: Immunofluorescence was performed against *CALR* (third panel) and an endoplasmic reticulum specific marker (Calnexin, second panel), in HEK293T transfected with the respective constructs. The wild type *CALR* co-localizes with calnexin, in the endoplasmic reticulum (last panel). However, the mutant *CALR* is not constrained within the endoplasmic reticulum. The nucleus is stained with the dye DAPI (first panel).

Figure 7.**The sensitivity of Ba/F3 cells to SAR302503.**

Ba/F3 cells expressing the empty vector (GFP), CALR wild type (CALR wt-GFP) or CALR mutant (CALR de152-GFP) were analyzed by defining cell viability after 48 hours in presence of decreasing SAR302503. For relative viability, relative luminescence units were normalized to the DMSO control. Error bars represent standard error of the mean.

Figure 8.

Western blot analysis was performed probing the sera of the four immunized mice, against lysates from HEK293T cells expressing the CALR mutant del52. The figure shows that the sera from all four immunized mice did not have any antibodies against the mutant calreticulin peptide before immunization - p (pre-immunized lanes). They generated specific antibody after 2 booster doses.

Figure 9.

Western blot analysis was performed probing the sera of the four immunized mice, against lysates from HEK293T cells expressing the CALR Δexon9. The figure shows that the sera from all four immunized mice did not have any antibodies against the CALR Δexon9 in both the pre-immunized sera (p) or after booster doses.

Figure 10.

Western blot analysis was performed probing the sera of the four immunized mice, against lysates from HEK293T cells expressing the CALR mutant del52. The figure shows that the sera of all four immunized mice had more specific antibody against calreticulin mutant after the third booster dose.

The Examples illustrate the invention.

Example 1: Somatic mutations of calreticulin in primary myelofibrosis and essential thrombocythemia

Material and Methods

Patient sampling

[0315] We studied patients with Philadelphia chromosome-negative myeloproliferative neoplasms followed at the Medical University of Vienna, Austria, and the Department of Hematology Oncology, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy. The investigations have been approved by the ethics committees of both institutions, and all patients provided written informed consent. Diagnosis of polycythemia vera, essential thrombocythemia and primary myelofibrosis was done according to the 2008 criteria of the World Health Organization (Swerdlow et al, 2008).

Diagnostic criteria

[0316] Diagnosis of polycythemia vera, essential thrombocythemia and primary myelofibrosis was done according to the criteria in use at the time of the first observation, as previously reported (Passamonti et al. 2004). In 2002 the criteria of the World Health Organization (WHO) were adopted (Vardiman et al, 2002) and in 2008 their revision was implemented (Swerdlow et al, 2008). *JAK2* and *MPL* mutation analyses were performed as previously described (Passamonti et al, 2010a; Rumi et al, 2013; Passamonti et al, 2011) in Pavia and as outlined below in Vienna. Bone marrow fibrosis was assessed semi-quantitatively following the European consensus guidelines (Thiele et al, 2005).

Thrombotic events were defined as described in detail previously (Marchioli et al, 2013). Patients were treated according to the recommendations that have been formalized by the European Leukemia Net in 2011 (Barbui et al, 2011)

Assessment of JAK2-V617F, JAK2-exon 12 and MPL-W515L mutations in Vienna

[0317] An allele specific PCR assay was used to detect the JAK2-V617F mutation. A primer mix consisting of 4 μ M of a common forward primer (gtttcttAGTGCATCTTTATTATGGCAGA (SEQ ID NO: 1340)), 2 μ M of a reverse primer specific for the wild type allele (TTACTCTCGTCTCCACAGAC (SEQ ID NO: 1341)) and 2 μ M of a reverse primer specific for the mutant allele (aaaTTACTCTCGTCTCCACAGAA (SEQ ID NO: 1342)) was prepared. The two reverse primers were fluorescently labeled with 6 - carboxyfluorescein (6-FAM) on the 5' end. A PCR reaction was set up using the AmpliTaq Gold DNA Polymerase with Gold Buffer and $MgCl_2$ (Applied Biosystems) containing 1.1 μ l of the 10x PCR GOLD buffer, 0.66 μ l of 25 mM $MgCl_2$, 0.44 μ l of 2.5 mM dNTPs, 1.4 μ l of the primer mix, 0.05 μ l of the AmpliTaq Gold polymerase (5 U/ μ l), 6.36 μ l H_2O and 1 μ l genomic DNA (10 ng/ μ l). PCR was performed as follows: 95°C for 5 min - 36 x (94°C for 30 sec, 62.2°C for 30 sec, 72°C for 30 sec) - 72°C for 15 min - 8°C hold. The PCR products were sized on a 3130x1 Genetic Analyzer (Applied Biosystems) and the data were analyzed using Gene Mapper software (Applied Biosystems).

[0318] The assay used to detect mutations in JAK2 exon 12 was reported before (Li et al, 2008)

For testing for the MPL-W515L the following allele specific PCR assay was used. A primer mix consisting of 8 μ M of a common forward primer (GTTTCTTCCGAAGTCTGACCCTTTTG (SEQ ID NO: 1343)), 4 μ M of a reverse primer specific for the wild type allele (GTAGTGTGCAGGAACTGCC (SEQ ID NO: 1344)) and 4 μ M of a reverse primer specific for the mutant allele (AAAGTAGTGTGCAGGAACTGCA (SEQ ID NO: 1345)) was prepared. The two reverse primers were fluorescently labeled with 6-carboxyfluorescein (6-FAM) on the 5' end. A PCR reaction was set up as described above for JAK2-V617F mutation. PCR was performed as follows: 94°C for 10 min - 30 x (94°C for 30 sec, 62.2°C for 30 sec, 72°C for 30 sec) - 72°C for 15 min - 8°C hold. The PCR products were sized on a 3130x1 Genetic Analyzer (Applied Biosystems) and the data were analyzed using Gene Mapper software (Applied Biosystems).

Whole exome sequencing

[0319] Genomic DNA was isolated from peripheral blood granulocytes (tumor tissue) and $CD3^+$ T-lymphocytes (control tissue) according to standard procedures. DNA libraries were generated using the NEB Next DNA Sample prep kit (reagent set) (New England Biolabs, Ipswich, MA) and whole-exome enrichment was performed using the Sure Select Human All Exon kit (Agilent, Santa Clara, CA) according to the manufacturers instructions. The libraries were sequenced on an Illumina HiSeq2000 system (Illumina, San Diego, CA) following the manufacturers recommendations. See Supplementary Table 1 for details on whole-exome sequencing parameters.

Supplementary Table 1

sample_ID	gDNA source	sequencing parameters	total reads	mean coverage	exonic bases covered at least			
					2X	10X	20X	30X
H_0010B_GD	granulocytes	52bp PE, 51bp PE	440,507,278	230	93.92%	82.95%	80.45%	78.86%
H_0010B_TD	T-lymphocytes	70bp SR, 75bp PE	305,708,053	187	95.67%	87.27%	82.73%	79.58%
H_0191B_GD	granulocytes	70bp SR, 51bp PE	329,139,581	163	95.93%	88.71%	84.26%	81.04%
H_0191A_TD	T-lymphocytes	70bp SR, 75bp PE	300,420,257	173	95.55%	88.26%	84.01%	80.81%
H_0202B_GD	granulocytes	70bp SR,	286,094,080	149	95.41%	87.81%	83.32%	79.84%

sample_ID	gDNA source	sequencing parameters	total reads	mean coverage	exonic bases covered at least			
					2X	10X	20X	30X
		51bp PE						
H 0202B_TD	T-lymphocytes	70bp SR, 75bp PE	262,563,166	160	95.25%	87.80%	83.44%	80.01%
H_0296C_GD	granulocytes	70bp SR, 51bp PE	279,769,005	138	95.17%	87.45%	82.68%	78.78%
H 0296C_TD	T-lymphocytes	70bp SR, 75bp PE	266,487,695	158	95.34%	87.79%	83.36%	79.83%
H_0333B_GD	granulocytes	70bp SR, 51bp PE	293,076,867	156	95.65%	88.23%	83.79%	80.41%
H_0333B_TD	T-lymphocytes	70bp SR, 75bp PE	260,476,597	155	94.99%	87.47%	83.03%	79.50%
H_0386B_GD	granulocytes	70bp SR, 51bp PE	285,286,604	148	95.58%	88.00%	83.48%	79.96%
H_0386B_TD	T-lymphocytes	70bp SR, 75bp PE	266,364,404	162	95.18%	87.69%	83.39%	79.98%
minimum			260,476,597	138	93.92%	82.95%	80.45%	78.78%
maximum			440,507,278	230	95.93%	88.71%	84.26%	81.04%
average			297,991,132	165	95.30%	87.45%	83.16%	79.88%

gDNA, genomic DNA; bp, base pairs; PE, paired-end; SR single read

Supplementary Table 2: Primers used to analyze the mutational status of CALR

application	targeted CALR exon	forward primer sequence	reverse primer sequence
Sanger sequencing / PCR product subcloning	9	ACAACTTCCTCATCACCAACG (SEQ ID NO: 437)	GGCCTCAGTCCAGCCCTG (SEQ ID NO: 438)
	9	GGCAAGGCCCTGAGGTGT (SEQ ID NO: 439)	GGCCTCAGTCCAGCCCTG (SEQ ID NO: 438)
PCR fragment analysis	1	GTCAGGTTGTTTGAGAGGC (SEQ ID NO: 1310)	GCTAACCCCTAACTCCCGCC (SEQ ID NO: 1311)
Sanger sequencing	2	GGATCTCCTTTCTGTCCCC (SEQ ID NO: 1312)	CCACCTGTCCTCCTCCAAG (SEQ ID NO: 1313)
Sanger sequencing	3	GAGGACAGGTGGAGGAAGTG (SEQ ID NO: 1314)	AAATTGTTGCTGGGACTTATTC (SEQ ID NO: 1315)
Sanger sequencing	4	CAGACCCGAGTTGAAGAACC (SEQ ID NO: 1316)	AGAAGGAAGAAGGTGAGCGG (SEQ ID NO: 1317)
Sanger sequencing	5	CTGATCAACAAGGACATCCG (SEQ ID NO: 1318)	CTCGGGCTTCTTAGCATCAG (SEQ ID NO: 1319)
Sanger sequencing	6-7	AAGCCTGAGGTTGGTGTTTG (SEQ ID NO: 1320)	CTCACCTGGGGTGCCTACC (SEQ ID NO: 1321)
Sanger sequencing	8	GTGTCAGCGGTGTTCTTG (SEQ ID NO: 1322)	TTAAGCCTCTGCTCCTCGTC (SEQ ID NO: 1323)
Sanger sequencing			

Whole-exome Sequencing analysis

[0320] The sequencing reads were aligned against the human reference genome (hg18) using BWA v0.5.9 (Li & Durbin, 2009). The genome analysis toolkit (GATK) v1.5 (McKenna et al, 2010) was used to post-process the alignments according to the best practices guidelines v3 of GATK. The coverage data presented in Supplementary Table 1 were calculated from the post-processed alignment files using the CalculateHsMetrics.jar script from Picard (<http://picard.sourceforge.net>). The post-processed alignment files were further analyzed by two variant callers:

1. 1. GATK's Unified Genotyper (DePristo et al, 2011) was used to call single nucleotide variants and small insertions/deletions from the granulocyte DNA samples. The preliminary variant lists were further processed using the Variant Quality Score Recalibrator (GATK) to generate recalibrated variant lists. The variants were annotated using ANNOVAR version 2012-05-25 (Wang et al, 2010). We filtered for variants that are found in coding exons and that affect the amino acid composition of the protein, as well as for variants at splice sites.
2. 2. The Varscan 2.3.2 tool (Koboldt et al, 2012) was used to call somatic variants comparing post-processed alignments from the granulocyte DNA sample with the alignments from the T-lymphocyte sample of the same patient. Varscan was used according to the programmer's instructions. Samtools 0.1.18 (Li et al, 2009) generated the mpileup files needed as the input for Varscan. Varscan hit lists were annotated by ANNOVAR and filtered as described above. Intersecting the variant lists retrieved from the two variant calling pipelines, as follows, generated final variant lists. GATK provides a score for the likelihood of a variant to be a true variant, which is the VQSLOD score. Varscan provides a p-value for a variant to be somatic. Basic requirements for final single nucleotide variant (SNV) calling were that the variants had to be called by both variant-calling pipelines and that the variant in the granulocyte sample was not classified as "low quality" by GATK. From all SNVs falling into this category, all those were called which had a VQSLOD score > 0 and a somatic p-value of < 0.05. We also called variants with a VQSLOD [-2;0] but required a somatic p-value of <0.01 for those. Insertion / deletion variant calling is more complex than SNV calling. In order not to miss true variants, we just required an insertion / deletion to be found by both pipelines. No further quality measures or p-values were required to call these variants.

Sanger Sequencing

[0321] Primers for Sanger Sequencing were designed using the PrimerZ tool (<http://ncbi36.genepipe.ncgm.sinica.edu.tw/primerz/beginDesign.do>) or the Primer3 tool (<http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi/>). Primer sequences are listed in Supplementary Table 2.

[0322] PCRs were performed using the AmpliTaq Gold 360 Mastermix (Applied Biosystems / Life Technologies, Paisley, UK).

[0323] A touchdown program was used for the PCR: 95 °C, 5 min - 10x (94 °C, 30 sec - 67 °C, 30 sec [-1 °C per cycle] - 72 °C, 30sec.) - 29x (94 °C, 30 sec - 57 °C, 30 sec - 72 °C, 30 sec) - 72 °C, 10 min. 10 °C, hold. For Sanger Sequencing the BigDye Terminator v3.1 Cycle Sequencing kit (Applied Biosystems) was used with the following program: 96 °C, 1 min. - 25x (96 °C, 10 sec. - 50 °C, 5 sec. - 60 °C, 4 min.) - 10 °C hold. Sequencing traces were read on a 3130x1 Genomic Analyzer (Genetic Analyzer) ((Applied Biosystems). Sequence analysis was done using the Sequencher Software 4.9 (Gene Codes, Ann Arbor, MI)

PCR fragment analysis for detection of CALR-exon9 mutations

[0324] Primers were designed for CALR exon 9 and the forward primer was 6-FAM labeled (Supplementary Table 2). PCR was performed as follows: 95 °C, 10 min. - 10x (94 °C, 15 sec. - 55 °C, 15 sec. - 72 °C, 30 sec.) - 20x (89 °C, 15 sec. - 55 °C, 15 sec. - 72 °C, 30 sec.) - 72 °C, 20 min - 10 °C, hold. PCR products were diluted 1:25 in water and sized on a 3130x1 Genomic Analyzer (Genetic Analyzer) (Applied Biosystems). The results were analyzed

using the Gene Mapper software version 4.0 (Applied Biosystems).

PCR product subcloning

[0325] PCR products were subcloned with the TOPO TA cloning kit (Invitrogen / Life Technologies, Paisley, UK) following the manufacturer's instructions using TOP-10 bacteria. Single bacterial colonies were picked the following day and expanded in an over-night culture. Plasmids were extracted with the QIAprep Spin Mini Prep kit (Qiagen, Hilden, Germany). A sequencing reaction was set up using the BigDye Terminator v3.1 Cycle Sequencing kit (Applied Biosystems): 50 - 200 ug plasmid, 4ul Primer, 1ul BigDye Terminator mix, 1ul sequencing buffer and HPLC water up to 10ul. The sequencing program was 96 °C, 5 min. - 25x (96 °C, 1 min. - 50 °C, 5 sec. - 60 °C, 4 min.) - 10 °C hold.

SNP microarray analysis

[0326] DNA samples were processed and hybridized to Genome-Wide Human SNP 6.0 arrays (Affymetrix) according to the protocol supplied by the manufacturer. The raw data was analyzed by Genotyping Console version 3.0.2 software (Affymetrix). The samples were assessed for chromosomal aberrations (deletions, gains and acquired uniparental disomies) as implemented in the Genotyping Console software.

Cloning of CALR exon 9 mutations

[0327] The wild type and mutant CALR was amplified from the clone purchased from Source Biosciences and cloned into the XhoI and EcoRI sites in the retroviral construct pMSCV-IRES-GFP. The wild type CALR was amplified using the following primers - (FP - ATGCCTCGAGCCGCCACCATGCTGCTATCCGTGCCGCTGCTGCTC (SEQ ID NO: 1346) and RP- ATGCGAATTCCTACAGCTCGTCCTTGGCCTGGCC (SEQ ID NO: 1347)).

[0328] The mutant CALR was amplified in two fragments followed by nested PCR -

(FP1 -ATGCCTCGAGCCGCCACCATGCTGCTATCCGTGCCGCTGCTGCTC (SEQ ID NO: 1346), RP1 - CCTCATCATCCTCCTTGTCTCTGCTCCTCGTCCTG (SEQ ID NO: 1348),

FP2 - CAGGACGAGGAGCAGAGGACAAGGAGGATGATGAGG (SEQ ID NO: 1349), RP2 - ATGCCCGCGGCTAGGCCTCAGTCCAGCCCTGGAGG (SEQ ID NO: 1350))

Virus production and transduction

[0329] The retrovirus was generated and cells were transduced as described before (Zuber et al, 2013). Briefly, PlatE cells (75% confluent in a 10 cm dish) were transfected with 20 µg of the respective viral vector using the Calcium Phosphate Transfection Kit (Sigma #CAPHOS). The medium was changed after 24 hours and viral supernatant was collected at 36, 40, 44 and 60 hours after transfection. 1 million Ba/F3 cells were transduced with the fresh viral supernatant, in 6 well plates by spinoculation (4 µg/ml of polybrene, 1350g, 30 minutes at 32°C), at every virus collection point. The cells were analyzed for transduction efficiency by flow cytometry, 48 hours after the final transduction step. The GFP positive cells were sorted by FACS and the sorting efficiency was analyzed by flow cytometry.

Proliferation and viability assays

[0330] To assess the viability of transduced Ba/F3 cells in presence of interleukin-3, cells were plated in 96-well plates at 5000 cells per well in triplicates on a dilution series of interleukin-3 (highest dose 25 ng/ml). After 72 hours cell viability was determined by the CellTiter-Glo® Luminescent Cell Viability Assay (Promega).

[0331] To determine cell proliferation in the absence of interleukin-3, Ba/F3 cells were plated in 12-well plates at 1000000 cells per well in triplicates and cultured for 7 days in complete RPMI (with 10% FCS, Pen/Strep and L-glutamine) without interleukin-3. Every 24 hours cell number was assessed using CASY® Cell Counter (Roche Innovatis).

[0332] To define the sensitivity to the inhibitor SAR302503 (Sanofi), Ba/F3 cells were plated in 96-well plates at 25000 cells per well in triplicates on a dilution series of SAR302503 (highest concentration 40 µM) and in presence of 10 ng/ml interleukin-3. After 48 hours cell viability was determined by the CellTiter-Glo® Luminescent Cell Viability Assay (Promega).

Interleukin-3 stimulation and western blotting

[0333] Ba/F3 cells were cultured in complete RPMI (10% FCS, Pen/Strep, L-glutamine) in the presence of 1 ng/ml of interleukin-3. Cells were starved in serum free medium without interleukin-3 for 4 hours. Stimulation was performed with respective concentration of interleukin-3, for 20 minutes. The cells were pelleted and protein extraction was done as described before (Corvinus et al, 2005). Briefly, complete whole cell extract buffer (containing the protease and phosphatase inhibitors) was added to the cells, and lysis was done by 3 consecutive freeze-thaw cycles with liquid Nitrogen. Lysates were collected after centrifugation for 20 minutes at 20,000 g. Protein concentration was measured using the Bradford reagent. Western blot was performed by standard techniques and 50 µg of protein was loaded per well. The following antibodies were used - pYStat5 (Invitrogen, #71-6900), Stat5 (Santa Cruz, sc-836), Calreticulin (Millipore, MABT145), GAPDH (Santa Cruz, sc-32233), anti-rabbit HRP (GE, NA934) and anti-mouse HRP (GE, NA931).

Immunofluorescence

[0334] HEK293T cells were seeded on glass coverslips coated with 0.1% gelatin transfected with the CMV-CALR wt and CMV-CALR del52 plasmids by lipofection (Invitrogen), according to the manufacturer's instructions, for 24 hours. ER staining was visualized using anti-Calnexin (ab31290, abcam); secondary anti-mouse AlexaFluor 546 (Invitrogen). Anti-Calreticulin antibody (MABT145, Millipore) was used to stain CALR; secondary anti-rabbit AlexaFluor 594 (Invitrogen). Slides were visualized using an LSM780 (Carl Zeiss, Germany) with a GaAsP multi-detector unit and two PMTs. Pinhole was set to 1AU on each channel. Images were taken sequentially, and channels selected, to reduce overlap. Images were taken at 100x and analyzed with ImageJ (NIH, open source.).

Statistical analysis

[0335] Statistical analysis was performed with the use of standard methods. Hypothesis testing was carried out with a non-parametric approach. All tests were two-tailed and P-values were considered significant when lower than 0.05. Microsoft Office Excel (Copyright Microsoft Corp), Stata 11.2 (Copyright StataCorp LP) and R 2.15.2 (R Core Team, 2012) were used for data management and analysis. The cumulative incidence of thrombotic complications was estimated with a competing risk approach according to the Kalbfleisch-Prentice method (Kalbfleisch & Prentice, 1980). Death in absence of the event of interest was considered as a competing event.

Results

Whole exome sequencing reveals recurrent Mutations of CALR in PMF (Primary myelofibrosis)

[0336] Genomic DNA from peripheral blood granulocytes (tumor tissue) and CD3⁺ T-lymphocytes (control tissue) from 6 patients with PMF was analyzed using whole exome sequencing. Independent validation of the detected variants using classical Sanger sequencing confirmed somatic mutations in between two and twelve genes per patient. The only recurrently affected gene was *CALR* encoding calreticulin. Two patients harbored somatic deletions in exon 9 of *CALR*. PCR product subcloning and sequencing revealed that patient 191 had a 52 base pair deletion and patient 296 harbored a one base pair deletion (Figure 1). As the 52 base pair deletion in patient 191 was incorrectly annotated as a one base pair deletion coupled with a single nucleotide variant by our variant calling analysis pipeline the sequence alignment of patient 191 was manually reviewed. A misalignment of the sequencing reads covering the site of mutation was observed. The incorrect alignment was due to a repetitive element in the affected genomic region. Following up on this finding the sequence alignments for the remaining four patients were investigated and a recurrent 5 bp insertion was detected in all 4 patients. The mutations of *CALR* found in the patients by whole exome sequencing were confirmed to be somatic by Sanger sequencing of the matched T-lymphocyte DNA samples. In summary, all six PMF patients analyzed by whole exome sequencing harbored somatic insertion or deletion mutations in exon 9 of *CALR*.

Frequency of CALR exon 9 mutations in myeloproliferative neoplasm (MPN) patients

[0337] In order to estimate the prevalence of *CALR* mutations in MPN a cohort of 896 MPN patients was screened for insertion and deletion mutations in *CALR* exon 9 using high-resolution sizing of fluorescent dye-labelled PCR products. This cohort included 382 patients with polycythemia vera (PV), 311 with essential thrombocythemia (ET) and 203 with primary myelofibrosis (PMF) (Table 1). 150 samples harboring insertions or deletions in *CALR* (17%) were identified. The mutations have been independently validated by Sanger sequencing. In PV no *CALR* mutations were observed. In ET and PMF, 78 (25%) and 72 (35%) patients had mutations in *CALR*, respectively (Table 1). All patients were genotyped for the JAK2-V617F mutation. PV patients negative for this mutation were tested for mutations in JAK2 exon 12. ET and PMF patients with wildtype JAK2 were tested for mutations in *MPL* exon 10.

[0338] The distribution of *JAK2*, *MPL* and *CALR* mutations in the three MPN disease entities is depicted in Figure 2A. All patients with mutant *CALR* had wild type *JAK2* and *MPL*. Therefore, mutations in *CALR* significantly associate with ET ($P=9.33 \times 10^{-45}$) and PMF ($P=1.71 \times 10^{-44}$) that are wild type for *JAK2* and *MPL* mutations. A total of 67 MPN patients tested wild type for *JAK2* and *MPL* as well as for *CALR* exon 9. Of these "triple-negative" cases, 19 patients were Sanger sequenced for mutations in all 9 exons of *CALR* but no mutations were detected.

[0339] As *CALR* mutations were highly associated with *JAK2* and *MPL* wild type ET or PMF, another 211 patients falling into this disease category were analyzed. In total, of 289 *JAK2/MPL* wild type ET patients 195 had mutant *CALR* (67%). Of the combined 120 *JAK2/MPL* wild type PMF patients, 105 had a mutation in *CALR* (88%). In 150 patients with mutant *CALR* for whom we had matched T-lymphocyte DNA available, the mutations were somatic.

Table 1. Comparison of *JAK2*, *MPL*, and *CALR* mutations in the three subtypes of MPN (number of patients are shown).

MPN subtype	N	JAK2 mutant	MPL mutant	CALR mutant	JAK2/MPL/CALR wild type
ET	311	184	11	78	38
PMF	203	108	13	72	10
PV	382	363	0	0	19
total:	896	655	24	150	67

Frequency of CALR mutations in other myeloid malignancies

[0340] To investigate if *CALR* mutations are present in other myeloid malignancies we screened 254 patients with *de novo* AML, 45 with chronic myeloid leukemia, 73 with myelodysplastic syndrome, 64 with chronic myelomonocytic leukemia and 24 with refractory anemia with ringsideroblasts associated with marked thrombocytosis for mutations in *CALR* exon 9. While the vast majority of these patients had wild type *CALR* exon 9, three patients with refractory anemia with ring sideroblasts associated with marked thrombocytosis harbored mutations in *CALR* (Figure 2B), all of them having wild-type *JAK2* and *MPL* (Figure 2C). Mutations in the gene encoding splicing factor 3B, subunit 1 (SF3B1) co-occurred with mutations in all three genes. Out of 524 healthy subjects one had a 3 bp in-frame deletion in *CALR*.

***CALR* frameshift mutations substitute the C-terminal amino acid sequence with a novel peptide derived from an alternative reading frame**

[0341] A total of 36 different types of mutations in *CALR* including insertions, deletions, combinations of deletions and insertions, as well as combinations of insertions/deletions with single nucleotide variants was detected. All observed mutations result in a frameshift to the alternative reading frame 1 of *CALR* (Figure 3A). This leads to a marked change in the amino acid composition of the C-terminus of the mutant *CALR* protein (Figure 3B). The C-terminus derived from exon 9 in wild type *CALR* is highly negatively charged, whereas the peptides of the mutants derived from the alternative reading frame 1 are positively charged. As the alternative reading frame 2 has a number of stop codons, frameshift mutations into this frame would result in a premature termination of translation and truncation of the protein (Figure 3B). No frameshift mutations into the alternative reading frame 2 were observed in the MPN patients studied. The 36 types of mutations are depicted in Figure 3C. Due to the different sizes and positions of the mutations gains and losses of variable numbers of amino acids are found at the C-terminal region of the protein (Table 2). All frameshift mutations generate a novel C-terminal amino acid sequence (Table 2). Full-length protein sequences of all mutants are shown in SEQ ID NO: 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288. The cDNA sequences around the mutation junctions are provided in Table 3.

Table 2. C-terminal amino acid sequences of insertion/deletion frameshift mutations of *CALR* found in MPN patients. Table 2 discloses SEQ ID NOs 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 4, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, and 144, respectively, in order of appearance.

Type 1	TRRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 2	NCRRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 3	QRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 4	RRRQRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 5	GQRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 6	RRQRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 7	RRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 8	RRQRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 9	RQRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 10	MCRRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 11	DQRQRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 12	RRRRQRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 13	QRRRQRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 14	RRRQRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 15	RRRERTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 16	QRRQRTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 17	RRQWTRMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-
Type 18	RMMRTKMRMRMRRTTRRKMRKMSPARPRTSCREACLQGWTEA-

Type 19	RQRTRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 20	GRRQRTRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 21	AFKRTRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 22	NAKRRRRQRTRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 23	CVRRRRQRTRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 24	RRQRTRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 25	RQRTRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 26	NAKRRRRQRTRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 27	CFAKRRRRQRTRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 28	RRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 29	PPLCLRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 30	DHPCRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 31	GNCRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 32	CRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 33	CRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 34	TCRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 35	ICRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-
Type 36	CRRMMRTKMRMRMRRTTRKMRRKMSPARPRTSCREACLOGWTEA-

Table 3. Sequences of mutation junctions in the cDNA sequence of CALR for the design of mutation specific probes or PCR primers. Table 7 discloses SEQ ID NOS 440-475, respectively, in order of appearance.

CALR mutation	cDNA junction sequences in mutated positions
Type 1	GAAGGACAAACAGGACGAGG AG CAGAGGACAAGGAGGATGAT
Type 2	GAGGAGGAGGCAGAGGACAATTGTCGGAGGATGATGAGGACAAAG
Type 3	GGACAAACAGGACGAGGAGC AG AGGCAGAGGACAAGGAGGAT
Type 4	CAGGACGAGGAGCAGAGGCT TAG GAGGAGGCAGAGGACAAGG
Type 5	TGAAGGACAAACAGGACGAG GG GCAGAGGACAAGGAGGATGA
Type 6	AGGACAAACAGGACGAGGAGC CG AGGCAGAGGACAAGGAGGA
Type 7	CAGGACGAGGAGCAGAGGCT TAG GAGGATGATGAGGACAAAG
Type 8	GGACGAGGAGCAGAGGCTTA AG AGGAGGCAGAGGACAAGGAG
Type 9	CAAGAAACGCAAAGAGGAG AG AGGCAGAGGACAAGGAGGAT
Type 10	AGGAGGAGGAGGCAGAGGAC ATGTGTC GGAGGATGATGAGGACAAAG
Type 11	AAGGACAAACAGGACGAGGAC CCAG AGGCAGAGGACAAGGAGGAT
Type 12	CAAACAGGACGAGGAGCAGAG GG AGGAGGAGGAGGCAGAGGAC
Type 13	AACAGGACGAGGAGCAGAGG CA GAGGAGGAGGCAGAGGACAAG
Type 14	ACAGGACGAGGAGCAGAGGCT TG AGGAGGAGGCAGAGGACAAG
Type 15	CAGGACGAGGAGCAGAGGCT TAG GAGGAGGGAGAGGACAAGGAGGATGATG
Type 16	CAGGACGAGGAGCAGAGGCT TC AGAGGAGGCAGAGGACAAGGAG
Type 17	GGACGAGGAGCAGAGGCTTA AG AGGAGGCAGTGGACAAGGAGGATGATGAGG
Type 18	GGACGAGGAGCAGAGGCTTA AG AGGATGATGAGGACAAAGAT
Type 19	GGAGCAGAGGCTTAAGGAG AG AGGCAGAGGACAAGGAGGAT
Type 20	GGCTTAAGGAGGAGGAAGA AGG AGGAGGCAGAGGACAAGGA
Type 21	GGCTTAAGGAGGAGGAAGA AGCGTTTA AGAGGACAAGGAGGATGATGA
Type 22	CTTAAGGAGGAGGAAGA ACA CGCAAAGAGGAGGAGGAGG

CALR mutation	cDNA junction sequences in mutated positions
Type 23	CTTAAGGAGGAGGAAGAAGAC <u>CTGCGT</u> GAGGAGGAGGAGGCAGAGGAC
Type 24	CTTAAGGAGGAGGAAGAAGAC <u>C</u> AGGAGGCAGAGGACAAGGAGG
Type 25	TAAGGAGGAGGAAGAAGACAA <u>AA</u> AGGCAGAGGACAAGGAGGATG
Type 26	TAAGGAGGAGGAAGAAGACAA <u>AAA</u> ACGCAAAGAGGAGGAGGAG
Type 27	AAGGAGGAGGAAGAAGACAA <u>GTGTTT</u> CGCAAAGAGGAGGAGGAGGCA
Type 28	GGAAGAAGACAAGAAACGCA <u>AA</u> AGGAGGATGATGAGGACAAA
Type 29	GAAGACAAGAAACGCAAAGAG <u>GCCTCCTCTTTGTCT</u> AAGGAGGATGATGAGGACAAA
Type 30	AGACAAGAAACGCAAAGAGG <u>ACCATCCTTGTCT</u> GGAGGATGATGAGGACAAAGA
Type 31	AGAGGAGGAGGAGGCAGAGG <u>GCAATTGTCT</u> CGGAGGATGATGAGGACAAAG
Type 32	GAGGAGGAGGAGGCAGAGGAC <u>CTGTCT</u> GGAGGATGATGAGGACAAAGA
Type 33	GAGGAGGAGGCAGAGGACAAAT <u>GTCT</u> CGGAGGATGATGAGGACAAAG
Type 34	AGGAGGAGGAGGCAGAGGAC <u>ACTTGTCT</u> GGAGGATGATGAGGACAAAGA
Type 35	AGGAGGAGGAGGCAGAGGAC <u>ATTTGTCT</u> GGAGGATGATGAGGACAAAGA
Type 36	AGGAGGAGGCAGAGGACA <u>ACTTGTCT</u> GGAGGATGATGAGGACAAAGA

Bold letters indicate the borders of a deletion event; underlined letters indicate inserted sequences; Bold and italic letters indicate single nucleotide variants

[0342] Myeloid cell specific somatic mutations in the CALR gene were identified in patients with MPN. The mutations strongly associate with those patients that are negative for both JAK2 and MPL mutations (the previously described disease causing mutations in MPN). As CALR mutations are found in 88% of PMF cases, and in 67% of ET cases double negative for JAK2 and MPL, it is believed that CALR mutations are filling in a large diagnostic gap for JAK2/MPL negative ET and PMF. Therefore, detection of CALR mutations at the level of genomic DNA, RNA or cDNA offers an important diagnostic test for MPN.

[0343] All identified mutations of CALR are in last exon 9 encoding the C-terminal amino acids of the protein and are predominantly insertion/deletion mutations. The vast majority of mutations were present in a heterozygous state and they cause a frameshift to the same alternative reading frame. This frameshift results in the replacement of the C-terminal negatively charged amino acids (aspartic and glutamic acid rich) of calreticulin by a predominantly positively charged polypeptide rich in arginine and methionine. In addition, the last 4 amino acids of calreticulin (KDEL (SEQ ID NO: 1331)) contain the endoplasmic reticulum retention signal. This signal is absent in the mutant calreticulin suggesting that the mutant protein is less represented in the ER compared to the wild type protein. As the negatively charged C-terminus of calreticulin is the low affinity high capacity Ca²⁺ binding domain, it is believed that the Ca²⁺ binding function of the mutant protein is lost.

[0344] From all CALR mutated cases, mutations of type 1 (52 base pair deletion) and type 2 (5 base pair insertion) were representing 53% and 32%, respectively (Figure 3D,E). The other mutation types were observed at much lower frequencies, many detected only in a single patient (Figure 3E). As all 36 mutation types frameshift to alternative reading frame 1, the resulting mutant CALR proteins share a common novel amino acid sequence at the C-terminus (Table 2). The C-terminal peptide derived from alternative reading frame 1 contains a number of positively charged amino acids, whereas the wild type CALR C-terminus is largely negatively charged (Figure 3B). In addition, the wild type calreticulin contains the endoplasmic reticulum retention motif at the C-terminal end (KDEL (SEQ ID NO: 1331) amino acid sequence). The C-terminal KDEL end (SEQ ID NO: 1331) is lost in all mutant variants (Table 2). Depending on the type of mutation, the mutant proteins retain variable amounts of the negatively charged amino acids of wild type calreticulin. The 52 base pair deletions of type 1 eliminate almost all negatively charged amino acids, whereas the 5 base pair insertions of type 2 retain approximately half of negatively charged amino acids (Figure 3D). Given these differences, we hypothesized that type 1 and type 2 mutations may be associated with qualitatively different phenotypes. Accordingly, we found that the type 1 deletions were significantly more frequent in primary myelofibrosis compared to essential thrombocythemia (P=0.0007). In addition, we

detected only 3 patients homozygous for CALR mutations associated with uniparental disomy of chromosome 19p and all 3 cases had a 5 base pair insertion of type 2 (Figure 4A).

CALR mutations are acquired early in the clonal evolution and mutant clones are stable

[0345] In order to investigate whether mutations in CALR are acquired early or late in the clonal history of a patient, we analyzed hematopoietic progenitor colonies from two patients for which we had mutational profiles from whole exome sequencing. The clonal hierarchies of patients H_0191 and H_0296 are shown in Figure 4B. We conclude that mutations in CALR were acquired early in the major clones of the two patients analyzed.

[0346] For 24 patients with mutant CALR we had follow-up samples available, all of which tested positive for the mutation.

Clinical significance of CALR mutations

[0347] Overall we studied 1215 patients with essential thrombocythemia or primary myelofibrosis Table 4.. Of those, 63.4% carried JAK2-V617F, 4.4% carried activating mutations of MPL exon 10, 23.5% carried mutations of CALR exon 9, and only 8.8% had none of the previous clonal markers. Of note, most of these latter patients clustered in the essential thrombocythemia subgroup. We used the Wilcoxon rank-sum test to compare hematologic values in patients carrying different mutant genes. Within patients with essential thrombocythemia, those carrying CALR mutation had lower hemoglobin level, lower white blood cell count, and higher platelet count at diagnosis compared with patients carrying mutant JAK2 ($P<0.001$ in all instances). Within patients with primary myelofibrosis, those carrying a CALR mutation had a lower white blood cell count ($P=0.027$) and a higher platelet count ($P<0.001$) than patients with mutant JAK2. Overall survival and risk of thrombosis were analyzed only in patients carrying a mutation in JAK2, MPL, or CALR, i.e., having a clonal marker. Assuming that mutation status did not change with time, we did all analyses since initial diagnosis. Since 6 patients were excluded due to inadequate follow-up, a total of 1102 patients were examined. The median follow-up for the whole cohort of patients with any of the three genes mutated was 5.7 years (range 0-31 years). As shown in Figure 5A, there was a significant difference in overall survival between the 3 subgroups of patients with primary myelofibrosis ($P<0.001$). Those carrying a somatic mutation of CALR had a better overall survival than those with JAK2 ($P<0.001$) or MPL mutation ($P<0.001$), while no difference was observed between the latter two subgroups. In patients with essential thrombocythemia, who have much longer overall survivals, there was a significant difference only between CALR mutated and JAK2 mutated patients ($P=0.043$, Figure 5B). In a Cox regression multivariate analysis of overall survival including type of myeloid neoplasm (essential thrombocythemia versus primary myelofibrosis), type of mutant gene, and patient cohort (Pavia versus Vienna) as covariates, the first two factors were found to be independent prognostic factors. As expected, primary myelofibrosis was associated with shorter overall survival as compared with essential thrombocythemia (hazard ratio of death 7.1, $P<0.001$, CI 4.9-10.2). In addition, the type of mutant gene had an independent effect on survival. In fact, as compared with patients carrying a CALR mutation, both those with JAK2 (hazard ratio 3.1, $P<0.001$, CI 2.0-4.7) and those carrying an MPL mutation (hazard ratio 3.5, $P<0.001$, CI 1.8-6.7) had a higher risk of death. The cumulative incidence of thrombosis in essential thrombocythemia was calculated with a competing risk approach with death from all causes as a competing event, and is reported in Table 5 while the actual curves are shown in Figure 5C. Patients with essential thrombocythemia carrying a CALR mutation had a lower risk of thrombosis than patients carrying a JAK2 mutation ($P=0.003$), while no significant difference was found between the other subgroups. It should be noted that the subgroup of patients carrying an MPL mutation was small.

Table 4: Cohort descriptives for the 1215 patients used to estimate clinical significance of CALR mutations

Diagnosis	JAK2 mutated (n=770)	MPL exon 10 mutated (n=53)	CALR mutated (n=285)	JAK2/MPL/CALR wild type (n=107)	All patients (n=1215)
Essential thrombocythemia	581 (65%)	35 (3.9%)	186 (20.8%)	92 (10.3%)	894
Primary	189	18	99	15	321

Diagnosis	JAK2 mutated (n=770)	MPL exon 10 mutated (n=53)	CALR mutated (n=285)	JAK2/MPL/CALR wild type (n=107)	All patients (n=1215)
myelofibrosis	(58.9%)	(5.6%)	(30.8%)	(4.7%)	

Table 5: Cumulative incidence of thrombosis comparing patients with mutant JAK2, MPL and CALR

Cumulative incidence of thrombosis	At 5 years	At 10 years	At 15 years
JAK2-mutated	13% (CI 10-16.4)	21% (CI 16.6-25.7)	27.1% (CI 21.4-33)
MPL-mutated	9.3% (CI 2.3-22.3)	9.3% (CI 2.3-22.3)	17.6% (CI 4.4-38.1)
CALR-mutated	6.3% (CI 3.2-10.8)	11% (CI 6.3-17.1)	12.8% (CI 7.3-20)

Functional analysis of the type 1 CALR mutation

[0348] In order to study the functional effects of mutant CALR we cloned the cDNA of the wild type CALR and the type 1 mutation (52 base pair deletion) into the retroviral expression vector pMSCV-IRES-GFP. After retroviral production and transfection of the CALR cDNAs into the interleukin-3-dependent murine cell line Ba/F3 we sorted the transgene positive cells by flow cytometry for GFP. Next we measured interleukin-3-dependent proliferation of cells. Cells expressing the type 1 CALR mutation exhibited interleukin-3-independent growth and hypersensitivity to interleukin-3 (Figure 6A). When we measured the proliferation of cells in the absence of interleukin-3, only the CALR 52 base pair deletion mutation exhibited significant accumulation of cells (Figure 6B). To investigate if the interleukin-3-independence in the CALR 52 base pair deletion mutant cells is caused by activation of JAK-STAT signaling, we determined the sensitivity of cells to the JAK2 kinase inhibitor SAR302503. As shown in Figure 7 both wild type and the 52 base pair deletion mutant of CALR showed similar sensitivity to SAR302503 suggesting that the interleukin-3-independent growth of the mutant CALR cells is dependent on JAK2 or a JAK family kinase targeted by SAR302503. To confirm this hypothesis, we examined the phosphorylation of STAT5 in the presence and absence of interleukin-3 in the control and CALR transfected cell lines. We detected increased phosphorylation of STAT5 in the absence of interleukin-3 in the 52 base pair deletion mutant of CALR and at 0.1 ng/ml interleukin-3 concentration (Figure 6C). Thus, increased activation of JAK-STAT signaling is likely responsible for the cytokine-independent growth of cells expressing the 52 base pair deletion mutant of CALR.

[0349] Immunofluorescence microscopy was used to determine the localization of wild type and type 1 mutant CALR. Upon overexpression in HEK cells, the wild type CALR colocalized with the endoplasmic reticulum (stained with calnexin). In case of the type 1 mutant CALR, this colocalization was less prominent (most likely due to the absence of the KDEL sequence (SEQ ID NO: 1331) from the C-terminus of the mutant CALR (Figure 6D).

[0350] We have identified somatic mutations in the CALR gene in patients with primary myelofibrosis and essential thrombocythemia. CALR mutations are mutually exclusive with mutations in both JAK2 and MPL. No CALR mutations were found in polycythemia vera, a myeloproliferative neoplasm that is specifically associated with JAK2 mutations. CALR mutations are the second most frequent mutation after JAK2 in myeloproliferative neoplasms. We have also studied patients with other myeloid neoplasms, and found CALR mutations only in 12.5% of cases with refractory anemia with ring sideroblasts associated with marked thrombocytosis, a typical myelodysplastic/myeloproliferative neoplasm (Malcovati et al, 2009) This strongly supports a causal relationship between CALR mutations and excessive platelet production.

[0351] As CALR mutations are found in about 73% of patients that do not carry alterations of JAK2 and MPL, we believe they are filling in the current molecular diagnostic gap in myeloproliferative neoplasms. Altogether, only less

than 10% of our patients with essential thrombocythemia or primary myelofibrosis do not carry a somatic mutation of JAK2, MPL or CALR. In some of these subjects, the mutated clone might be too small to be detected with the current approaches. Rare mutant driver genes may play a role in other patients, while some patients might not have a clonal disease at all. This is particularly true for patients with a clinical diagnosis of essential thrombocythemia, as differential diagnosis between clonal and reactive thrombocytosis may be difficult without a clonal marker (Schafer, 2004). Overall, the assessment of CALR mutations markedly improves the current diagnostic approach for essential thrombocythemia or primary myelofibrosis, and should be included in the WHO criteria for these disorders (Sverdlow et al, 2008)

[0352] All the mutations of CALR we identified are insertion/deletion mutations in the last exon encoding the C-terminal amino acids of the protein. The most mutations are present in a heterozygous state and cause a frameshift to a specific alternative reading frame. This frameshift results in the replacement of the C-terminal negatively charged amino acids of calreticulin by a positively charged polypeptide rich in arginine and methionine. The last 4 amino acids of calreticulin (KDEL (SEQ ID NO: 1331)) contain the endoplasmic reticulum retention signal. This signal is absent in the mutant calreticulin. Consequently mutant calreticulin has an altered subcellular localization. As the negatively charged C-terminus of calreticulin is the low-affinity high-capacity Ca²⁺ binding domain, the Ca²⁺ binding function of the mutant protein may be impaired. The presence of the peptide sequence derived from the alternative reading frame at the C-terminus of mutated CALR offers an opportunity for immunologic targeting as it represents a cancer specific epitope.

[0353] To further analyze the oncogenic capability of the mutant calreticulin, we generated Ba/F3 cells with overexpression of the wild type and the type 1 mutant CALR (52 base pair deletion - del52). Interestingly, the CALR del52 Ba/F3 cells showed cytokine independent proliferation. However, the growth of Ba/F3 cells expressing wild type and mutant calreticulin was suppressed equally upon treatment with a JAK2 kinase inhibitor, suggesting the requirement of the JAK-STAT pathway in the mutant calreticulin-induced cytokine independence. In accordance, we could detect increased phosphorylation of STAT5 in del52 Ba/F3 cells, both in the absence and presence of interleukin-3 stimulation. Calreticulin/Ca²⁺/calmodulin has been previously shown to modulate the activity of Stats. Calreticulin complex with ERp57, in endoplasmic reticulum, suppresses the phosphorylation and transcriptional activity of Stat3 in mouse embryonic fibroblasts (Coe et al, 2010). Moreover, inhibition of the Ca²⁺/calmodulin dependent kinase II gamma results in reduced levels of phosphorylated Stat1, Stat3 and Stat5 (Si & Collins 2008). Interestingly, overexpression of calreticulin attenuates interferon alpha induced Stat1 phosphorylation, resulting in interferon resistance (Yue et al, 2012). Further studies are required to elucidate the mechanism of the activation of JAK-STAT pathway by the mutant calreticulin in myeloid cells. The involvement of the JAK-STAT signaling pathway in CALR positive patients may also explain the effectivity of JAK2 inhibitor therapy in primary myelofibrosis. However, our results indicate that the JAK2 inhibitors may not be selective for cells expressing the mutated CALR compared to the CALR wild type cells.

[0354] Although our analyses of clinical outcome are retrospective, they strongly suggest that CALR positive myeloproliferative neoplasms have a more benign clinical course than the corresponding disorders associated with JAK2 or MPL mutation. Due to the small number of MPL mutated patients, the more reliable comparisons are those between JAK2 mutated and CALR mutated patients. Our observations clearly show that CALR mutated patients have lower risk of thrombosis and better overall survival than JAK2 mutated patients. The lower incidence of thromboembolic complications might be related to the fact that CALR mutated patients had lower hemoglobin levels and lower white blood cell counts. A better overall survival was observed both in patients with primary myelofibrosis and those with essential thrombocythemia, although it was much more relevant in the former, confirming previous findings on patients with and without JAK2 mutations (Campbell et al, 2006b; Rumi et al, 2013). From a practical point of view, the different impact of mutant genes might be incorporated in existing prognostic scoring systems for primary myelofibrosis and essential thrombocythemia (Passamonti 2010b, Passamonti 2012)_ and may also guide therapeutic decision-making. More specifically, CALR molecular characterization should become an essential component of future clinical management of essential thrombocythemia and primary myelofibrosis.

Example 2: Generation of CALR mutant specific antibodies in mice

[0355] The *CALR* mutations associated with MPN occur in the last exon of the gene (exon 9). These mutations are insertions and/or deletions that result in a 'frameshift' mutation to a very specific alternative reading frame, leading to synthesis of a novel C-terminal peptide in the mutant. As all the mutations result in generation of the same alternative reading frame, the C-terminal peptide has the same sequence in all the *CALR* mutants.

A synthetic peptide with the c-terminal end sequence of the mutant calreticulin protein (Sequence - RRKMSPARPRTSCREACLGWTEA-), conjugated to the Keyhole Limpet Hemocyanin (KLH) was used to immunize four wild type C57B1/6 mice.

[0356] The mice received 3 booster doses after the primary immunization. The sera of the mice was tested (pre-immune and after boosters) for the presence mutant calreticulin specific antibodies by western blot analysis of lysates from HEK cells that over-expressed the *CALR* del52 and the artificially generated *CALR* mutant that lacks the exon 9 (Δ exon9, which lacks the mutant peptide). The lysates were run on 8% polyacrylamide gels and probed with the mouse serum. Anti-mouse antibody conjugated to HRP (GE NA931) was used as secondary antibody. After the second booster, the sera from all four mice had *CALR* mutant specific antibodies that detected the *CALR* del52 mutant (Figure 8), but not the control exon 9 deleted *CALR* (Figure 9). Anti- calreticulin antibody (Millipore MABT145) was used as positive control (Pos), which recognizes all three forms of calreticulin - wild type, mutant del 52 and deleted exon 9. The signal, from the sera of all four mice, was stronger after the third booster was applied (Figure 10).

[0357] The C-terminal peptide of the mutant calreticulin (mentioned above) is immunogenic and can successfully be used to generate specific antibodies, in particular polyclonal antibodies against the mutant calreticulin. These antibodies can be used as research reagents as well as for diagnostic purposes as disclosed here.

[0358] The present invention refers to the following nucleotide and amino acid sequences:

The sequences provided herein are in part available in the NCBI database and can be retrieved from [www at ncbi.nlm.nih.gov/sites/entrez?db=gene](http://www.ncbi.nlm.nih.gov/sites/entrez?db=gene); The sequences also relate to annotated and modified sequences. The present invention also provides techniques and methods wherein homologous sequences, and variants of the concise sequences provided herein are used. Preferably, such "variants" are genetic variants.

SEQ ID No. 1: Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 18

SEQ ID No. 2: Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 18

SEQ ID No. 3: Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 18

SEQ ID No. 4: Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 18

SEQ ID No. 5: Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 1

SEQ ID No. 6: Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 1

SEQ ID No. 7: Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 1

SEQ ID No. 8: Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 1

SEQ ID No. 9: Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 2

SEQ ID No. 10: Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 2

SEQ ID No. 11 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 2

SEQ ID No. 12 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 2

SEQ ID No. 13 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 3

SEQ ID No. 14 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 3

SEQ ID No. 15 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 3

[illegible]

[illegible]

SEQ ID No.81 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 21

SEQ ID No.82 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 21

SEQ ID No.83 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 21

SEQ ID No.84 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 21

SEQ ID No.85 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 22

SEQ ID No.86 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 22

SEQ ID No.87 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 22

SEQ ID No.88 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 22

SEQ ID No.89 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 23

SEQ ID No.90 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 23

SEQ ID No.91 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 23

SEQ ID No.92 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 23

SEQ ID No.93 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 24

SEQ ID No.94 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 24

SEQ ID No.95 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 24

SEQ ID No.96 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 24

SEQ ID No.97 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 25

SEQ ID No.98 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 25

SEQ ID No.99 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 25

SEQ ID No. 100 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 25

SEQ ID No.101 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 26

SEQ ID No. 102 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 26

SEQ ID No. 103 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 26

SEQ ID No. 104 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 26

SEQ ID No. 105 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 27

SEQ ID No.106 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 27

SEQ ID No. 107 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 27

SEQ ID No.108 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 27

SEQ ID No. 109 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 28

SEQ ID No.110 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 28

SEQ ID No.111 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 28

SEQ ID No.112 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 28

SEQ ID No. 113 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 29

SEQ ID No. 114 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 29

SEQ ID No. 115 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 29

SEQ ID No.116 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 29

SEQ ID No. 117 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 30

SEQ ID No.118 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 30

SEQ ID No. 119 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 30

SEQ ID No. 120 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 30

SEQ ID No.121 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 31

SEQ ID No.122 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 31

SEQ ID No.123 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 31

SEQ ID No. 124 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 31

SEQ ID No.125 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 32

SEQ ID No. 126 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 32

SEQ ID No. 127 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 32

SEQ ID No. 128 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 32

SEQ ID No.129 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 33

SEQ ID No. 130 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 33

SEQ ID No.131 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 33

SEQ ID No. 132 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 33

SEQ ID No.133 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 34

SEQ ID No. 134 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 34

SEQ ID No. 135 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 34

SEQ ID No. 136 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 34

SEQ ID No.137 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 35

SEQ ID No.138 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 35

SEQ ID No.139 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 35

SEQ ID No. 140 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 35

SEQ ID No.141 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 36

SEQ ID No. 142 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 36

SEQ ID No. 143 Nucleotide sequence encoding the C-terminus of homo sapiens calreticulin mutant type 36

SEQ ID No. 144 Amino acid sequence of the C-terminus of homo sapiens calreticulin mutant type 36

SEQ ID No. 145 Nucleotide sequence encoding homo sapiens calreticulin mutant type 1

SEQ ID No. 146 Nucleotide sequence encoding homo sapiens calreticulin mutant type 1

SEQ ID No. 147 Nucleotide sequence encoding homo sapiens calreticulin mutant type 1

SEQ ID No. 148 Amino acid sequence of homo sapiens calreticulin mutant type 1

SEQ ID No. 149 Nucleotide sequence encoding homo sapiens calreticulin mutant type 2

SEQ ID No.150 Nucleotide sequence encoding homo sapiens calreticulin mutant type 2

SEQ ID No. 151 Nucleotide sequence encoding homo sapiens calreticulin mutant type 2

SEQ ID No. 152 Amino acid sequence of homo sapiens calreticulin mutant type 2

SEQ ID No.153 Nucleotide sequence encoding homo sapiens calreticulin mutant type 3

SEQ ID No. 154 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 3

SEQ ID No.155 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 3

SEQ ID No.156 Amino acid sequence of homo sapiens calreticulin mutant type 3

SEQ ID No.157 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 4

SEQ ID No.158 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 4

SEQ ID No.159 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 4

SEQ ID No.160 Amino acid sequence of homo sapiens calreticulin mutant type 4

SEQ ID No.161 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 5

SEQ ID No.162 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 5

SEQ ID No.163 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 5

SEQ ID No.164 Amino acid sequence of homo sapiens calreticulin mutant type 5

SEQ ID No.165 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 6

SEQ ID No.166 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 6

SEQ ID No.167 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 6

SEQ ID No.168 Amino acid sequence of homo sapiens calreticulin mutant type 6

SEQ ID No.169 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 7

SEQ ID No. 170 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 7

SEQ ID No. 171 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 7

SEQ ID No. 172 Amino acid sequence of homo sapiens calreticulin mutant type 7

SEQ ID No. 173 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 8

SEQ ID No. 174 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 8

SEQ ID No. 175 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 8

SEQ ID No. 176 Amino acid sequence of homo sapiens calreticulin mutant type 8

SEQ ID No. 177 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 9

SEQ ID No. 178 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 9

SEQ ID No. 179 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 9

SEQ ID No.180 Amino acid sequence of homo sapiens calreticulin mutant type 9

SEQ ID No.181 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 10

SEQ ID No.182 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 10

SEQ ID No.183 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 10

SEQ ID No.184 Amino acid sequence of homo sapiens calreticulin mutant type 10

SEQ ID No.185 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 11

SEQ ID No.186 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 11

SEQ ID No.187 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 11

SEQ ID No.188 Amino acid sequence of homo sapiens calreticulin mutant type 11

SEQ ID No.189 Nucleotide sequence encoding homo sapiens calreticulin mutant type 12

SEQ ID No. 190 Nucleotide sequence encoding homo sapiens calreticulin mutant type 12

SEQ ID No. 191 Nucleotide sequence encoding homo sapiens calreticulin mutant type 12

SEQ ID No. 192 Amino acid sequence of homo sapiens calreticulin mutant type 12

SEQ ID No. 193 Nucleotide sequence encoding homo sapiens calreticulin mutant type 13

SEQ ID No. 194 Nucleotide sequence encoding homo sapiens calreticulin mutant type 13

SEQ ID No. 195 Nucleotide sequence encoding homo sapiens calreticulin mutant type 13

SEQ ID No. 196 Amino acid sequence of homo sapiens calreticulin mutant type 13

SEQ ID No. 197 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 14

SEQ ID No.198 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 14

SEQ ID No. 199 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 14

SEQ ID No.200 Amino acid sequence of homo sapiens calreticulin mutant type 14

SEQ ID No.201 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 15

SEQ ID No.202 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 15

SEQ ID No.203 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 15

SEQ ID No.204 Amino acid sequence of homo sapiens calreticulin mutant type 15

SEQ ID No.205 Nucleotide sequence encoding homo sapiens calreticulin mutant type 16

SEQ ID No.206 Nucleotide sequence encoding homo sapiens calreticulin mutant type 16

SEQ ID No.207 Nucleotide sequence encoding homo sapiens calreticulin mutant type 16

SEQ ID No.208 Amino acid sequence of homo sapiens calreticulin mutant type 16

SEQ ID No.209 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 17

SEQ ID No.210 Nucleotide sequence encoding homo sapiens calreticulin mutant type 17

SEQ ID No.211 Nucleotide sequence encoding homo sapiens calreticulin mutant type 17

SEQ ID No.212 Amino acid sequence of homo sapiens calreticulin mutant type 17

SEQ ID No.213 Nucleotide sequence encoding homo sapiens calreticulin mutant type 18

SEQ ID No.214 Nucleotide sequence encoding homo sapiens calreticulin mutant type 18

SEQ ID No.215 Nucleotide sequence encoding the homo sapiens calreticulin mutant type 18

SEQ ID No.216 Amino acid sequence of homo sapiens calreticulin mutant type 18

SEQ ID No.217 Nucleotide sequence encoding homo sapiens calreticulin mutant type 19

SEQ ID No.218 Nucleotide sequence encoding homo sapiens calreticulin mutant type 19

SEQ ID No.219 Nucleotide sequence encoding homo sapiens calreticulin mutant type 19

SEQ ID No.220 Amino acid sequence of homo sapiens calreticulin mutant type 19

SEQ ID No.221 Nucleotide sequence encoding homo sapiens calreticulin mutant type 20

SEQ ID No.222 Nucleotide sequence encoding homo sapiens calreticulin mutant type 20

SEQ ID No.223 Nucleotide sequence encoding homo sapiens calreticulin mutant type 20

SEQ ID No.224 Amino acid sequence of homo sapiens calreticulin mutant type 20

SEQ ID No.225 Nucleotide sequence encoding homo sapiens calreticulin mutant type 21

SEQ ID No.226 Nucleotide sequence encoding homo sapiens calreticulin mutant type 21

SEQ ID No.227 Nucleotide sequence encoding homo sapiens calreticulin mutant type 21

SEQ ID No.228 Amino acid sequence of homo sapiens calreticulin mutant type 21

SEQ ID No.229 Nucleotide sequence encoding homo sapiens calreticulin mutant type 22

SEQ ID No.230 Nucleotide sequence encoding homo sapiens calreticulin mutant type 22

SEQ ID No.231 Nucleotide sequence encoding homo sapiens calreticulin mutant type 22

SEQ ID No.232 Amino acid sequence of homo sapiens calreticulin mutant type 22

SEQ ID No.233 Nucleotide sequence encoding homo sapiens calreticulin mutant type 23

SEQ ID No.234 Nucleotide sequence encoding homo sapiens calreticulin mutant type 23

SEQ ID No.235 Nucleotide sequence encoding homo sapiens calreticulin mutant type 23

SEQ ID No.236 Amino acid sequence of homo sapiens calreticulin mutant type 23

SEQ ID No.237 Nucleotide sequence encoding homo sapiens calreticulin mutant type 24

SEQ ID No.238 Nucleotide sequence encoding homo sapiens calreticulin mutant type 24

SEQ ID No.239 Nucleotide sequence encoding homo sapiens calreticulin mutant type 24

SEQ ID No.240 Amino acid sequence of homo sapiens calreticulin mutant type 24

SEQ ID No.241 Nucleotide sequence encoding homo sapiens calreticulin mutant type 25

SEQ ID No.242 Nucleotide sequence encoding homo sapiens calreticulin mutant type 25

SEQ ID No.243 Nucleotide sequence encoding homo sapiens calreticulin mutant type 25

SEQ ID No.244 Amino acid sequence of homo sapiens calreticulin mutant type 25

SEQ ID No.245 Nucleotide sequence encoding homo sapiens calreticulin mutant type 26

SEQ ID No.246 Nucleotide sequence encoding homo sapiens calreticulin mutant type 26

SEQ ID No.247 Nucleotide sequence encoding homo sapiens calreticulin mutant type 26

SEQ ID No.248 Amino acid sequence of homo sapiens calreticulin mutant type 26

SEQ ID No.249 Nucleotide sequence encoding homo sapiens calreticulin mutant type 27

SEQ ID No.250 Nucleotide sequence encoding homo sapiens calreticulin mutant type 27

SEQ ID No.251 Nucleotide sequence encoding homo sapiens calreticulin mutant type 27

SEQ ID No.252 Amino acid sequence of homo sapiens calreticulin mutant type 27

SEQ ID No.253 Nucleotide sequence encoding homo sapiens calreticulin mutant type 28

SEQ ID No.254 Nucleotide sequence encoding homo sapiens calreticulin mutant type 28

SEQ ID No.255 Nucleotide sequence encoding homo sapiens calreticulin mutant type 28

SEQ ID No.256 Amino acid sequence of homo sapiens calreticulin mutant type 28

SEQ ID No.257 Nucleotide sequence encoding homo sapiens calreticulin mutant type 29

SEQ ID No.258 Nucleotide sequence encoding homo sapiens calreticulin mutant type 29

SEQ ID No.259 Nucleotide sequence encoding homo sapiens calreticulin mutant type 29

SEQ ID No.260 Amino acid sequence of homo sapiens calreticulin mutant type 29

SEQ ID No.261 Nucleotide sequence encoding homo sapiens calreticulin mutant type 30

SEQ ID No.262 Nucleotide sequence encoding homo sapiens calreticulin mutant type 30

SEQ ID No.263 Nucleotide sequence encoding homo sapiens calreticulin mutant type 30

SEQ ID No.264 Amino acid sequence of homo sapiens calreticulin mutant type 30

SEQ ID No.265 Nucleotide sequence encoding homo sapiens calreticulin mutant type 31

SEQ ID No.266 Nucleotide sequence encoding homo sapiens calreticulin mutant type 31

SEQ ID No.267 Nucleotide sequence encoding homo sapiens calreticulin mutant type 31

SEQ ID No.268 Amino acid sequence of homo sapiens calreticulin mutant type 31

SEQ ID No.269 Nucleotide sequence encoding homo sapiens calreticulin mutant type 32

SEQ ID No.270 Nucleotide sequence encoding homo sapiens calreticulin mutant type 32

SEQ ID No.271 Nucleotide sequence encoding homo sapiens calreticulin mutant type 32

SEQ ID No.272 Amino acid sequence of homo sapiens calreticulin mutant type 32

SEQ ID No.273 Nucleotide sequence encoding homo sapiens calreticulin mutant type 33

SEQ ID No.274 Nucleotide sequence encoding homo sapiens calreticulin mutant type 33

SEQ ID No.275 Nucleotide sequence encoding homo sapiens calreticulin mutant type 33

SEQ ID No.276 Amino acid sequence of homo sapiens calreticulin mutant type 33

SEQ ID No.277 Nucleotide sequence encoding homo sapiens calreticulin mutant type 34

SEQ ID No.278 Nucleotide sequence encoding homo sapiens calreticulin mutant type 34

SEQ ID No.279 Nucleotide sequence encoding homo sapiens calreticulin mutant type 34

SEQ ID No.280 Amino acid sequence of homo sapiens calreticulin mutant type 34

SEQ ID No.281 Nucleotide sequence encoding homo sapiens calreticulin mutant type 35

SEQ ID No.282 Nucleotide sequence encoding homo sapiens calreticulin mutant type 35

SEQ ID No.283 Nucleotide sequence encoding homo sapiens calreticulin mutant type 35

SEQ ID No.284 Amino acid sequence of homo sapiens calreticulin mutant type 35

SEQ ID No.285 Nucleotide sequence encoding homo sapiens calreticulin mutant type 36

SEQ ID No.286 Nucleotide sequence encoding homo sapiens calreticulin mutant type 36

SEQ ID No.287 Nucleotide sequence encoding homo sapiens calreticulin mutant type 36

SEQ ID No.288 Amino acid sequence of homo sapiens calreticulin mutant type 36

SEQ ID No.289 Nucleotide sequence encoding homo sapiens calreticulin wild type

SEQ ID No.290 Amino acid sequence of homo sapiens calreticulin wild type

SEQ ID No.291 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 1

SEQ ID No.292 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 1

SEQ ID No.293 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 1

SEQ ID No.294 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 1

SEQ ID No.295 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 2

SEQ ID No.296 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 2

SEQ ID No.297 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 2

SEQ ID No.298 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 2

SEQ ID No.299 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 3

SEQ ID No.300 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 3

SEQ ID No.301 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 3

SEQ ID No.302 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 3

SEQ ID No.303 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 4

SEQ ID No.304 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 4

SEQ ID No.305 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 4

SEQ ID No.306 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 4

SEQ ID No.307 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 5

SEQ ID No.308 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 5

SEQ ID No.309 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 5

SEQ ID No.310 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 5

SEQ ID No.311 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 6

SEQ ID No.312 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 6

SEQ ID No.313 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 6

SEQ ID No.314 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 6

SEQ ID No.315 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 7

SEQ ID No.316 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 7

SEQ ID No.317 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 7

SEQ ID No.318 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 7

SEQ ID No.319 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 8

SEQ ID No.320 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 8

SEQ ID No.321 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 8

SEQ ID No.322 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 8

SEQ ID No.323 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 9

SEQ ID No.324 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 9

SEQ ID No.325 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 9

SEQ ID No.326 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 9

SEQ ID No.327 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 10

SEQ ID No.328 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 10

SEQ ID No.329 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 10

SEQ ID No.330 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 10

SEQ ID No.331 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 11

SEQ ID No.332 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 11

SEQ ID No.333 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 11

SEQ ID No.334 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 11

SEQ ID No.335 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 12

SEQ ID No.336 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 12

SEQ ID No.337 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 12

SEQ ID No.338 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 12

SEQ ID No.339 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 13

SEQ ID No.340 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 13

SEQ ID No.341 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 13

SEQ ID No.342 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 13

SEQ ID No.343 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 14

SEQ ID No.344 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 14

SEQ ID No.345 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 14

SEQ ID No.346 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 14

SEQ ID No.347 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 15

SEQ ID No.348 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 15

SEQ ID No.349 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 15

SEQ ID No.350 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 15

SEQ ID No.351 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 16

SEQ ID No.352 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 16

SEQ ID No.353 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 16

SEQ ID No.354 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 16

SEQ ID No.355 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 17

SEQ ID No.356 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 17

SEQ ID No.357 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 17

SEQ ID No.358 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 17

SEQ ID No.359 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 18

SEQ ID No.360 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 18

SEQ ID No.361 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 18

SEQ ID No.362 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 18

SEQ ID No.363 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 19

SEQ ID No.364 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 19

SEQ ID No.365 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 19

SEQ ID No.366 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 19

SEQ ID No.367 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 20

SEQ ID No.368 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 20

SEQ ID No.369 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 20

SEQ ID No.370 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 20

SEQ ID No.371 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 21

SEQ ID No.372 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 21

SEQ ID No.373 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 21

SEQ ID No.374 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 21

SEQ ID No.375 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 22

SEQ ID No.376 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 22

SEQ ID No.377 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 22

SEQ ID No.378 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 22

SEQ ID No.379 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 23

SEQ ID No.380 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 23

SEQ ID No.381 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 23

SEQ ID No.382 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 23

SEQ ID No.383 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 24

SEQ ID No.384 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 24

SEQ ID No.385 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 24

SEQ ID No.386 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 24

SEQ ID No.387 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 25

SEQ ID No.388 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 25

SEQ ID No.389 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 25

SEQ ID No.390 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 25

SEQ ID No.391 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 26

SEQ ID No.392 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 26

SEQ ID No.393 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 26

SEQ ID No.394 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 26

SEQ ID No.395 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 27

SEQ ID No.396 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 27

SEQ ID No.397 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 27

SEQ ID No.398 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 27

SEQ ID No.399 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 28

SEQ ID No.400 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 28

SEQ ID No.401 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 28

SEQ ID No.402 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 28

SEQ ID No.403 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 29

SEQ ID No.404 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 29

SEQ ID No.405 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 29

SEQ ID No.406 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 29

SEQ ID No.407 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 30

SEQ ID No.408 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 30

SEQ ID No.409 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 30

SEQ ID No.410 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 30

SEQ ID No.411 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 31

SEQ ID No.412 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 31

SEQ ID No.413 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 31

SEQ ID No.414 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 31

SEQ ID No.415 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 32

SEQ ID No.416 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 32

SEQ ID No.417 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 32

SEQ ID No.418 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 32

SEQ ID No.419 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 33

SEQ ID No.420 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 33

SEQ ID No.421 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 33

SEQ ID No.422 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 33

SEQ ID No.423 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 34

SEQ ID No.424 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 34

SEQ ID No.425 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 34

SEQ ID No.426 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 34

SEQ ID No.427 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 35

SEQ ID No.428 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 35

SEQ ID No.429 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 35

SEQ ID No.430 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 35

SEQ ID No.431 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 36

SEQ ID No.432 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 36

SEQ ID No.433 Nucleotide sequence encoding homo sapiens calreticulin exon 9 mutant type 36

SEQ ID No.434 Amino acid sequence of homo sapiens calreticulin exon 9 mutant type 36

SEQ ID No.435 Nucleotide sequence encoding homo sapiens calreticulin exon 9 wild type

SEQ ID No.436 Amino acid sequence of homo sapiens calreticulin exon 9 wild type

[0359] The following table provides an overview of herein provided and used SEQ ID NOs.

	SEQ ID NO			
	genomic DNA	copy(c)DNA	mRNA	protein
C-terminus (minimum sequence) of mutant calreticulin	1	2	3	4
C-termini of mutant calreticulin				
type 1	5	6	7	8
type 2	9	10	11	12
type 3	13	14	15	16
type 4	17	18	19	20
type 5	21	22	23	24
type 6	25	26	27	28
type 7	29	30	31	32
type 8	33	34	35	36
type 9	37	38	39	40
type 10	41	42	43	44
type 11	45	46	47	48
type 12	49	50	51	52
type 13	53	54	55	56
type 14	57	58	59	60
type 15	61	62	63	64
type 16	65	66	67	68
type 17	69	70	71	72
type 18	1	2	3	4
type 19	73	74	75	76
type 20	77	78	79	80
type 21	81	82	83	84
type 22	85	86	87	88
type 23	89	90	91	92
type 24	93	94	95	96
type 25	97	98	99	100
type 26	101	102	103	104
type 27	105	106	107	108
type 28	109	110	111	112
type 29	113	114	115	116
type 30	117	118	119	120
type 31	121	122	123	124
type 32	125	126	127	128
type 33	129	130	131	132
type 34	133	134	135	136
type 35	137	138	139	140

	SEQ ID NO			
	genomic DNA	copy(c)DNA	mRNA	protein
C-terminus (minimum sequence) of mutant calreticulin	1	2	3	4
type 36	141	142	143	144
full length mutant calreticulin				
type 1	145	146	147	148
type 2	149	150	151	152
type 3	153	154	155	156
type 4	157	158	159	160
type 5	161	162	163	164
type 6	165	166	167	168
type 7	169	170	171	172
type 8	173	174	175	176
type 9	177	178	179	180
type 10	181	182	183	184
type 11	185	186	187	188
type 12	189	190	191	192
type 13	193	194	195	196
type 14	197	198	199	200
type 15	201	202	203	204
type 16	205	206	207	208
type 17	209	210	211	212
type 18	213	214	215	216
type 19	217	218	219	220
type 20	221	222	223	224
type 21	225	226	227	228
type 22	229	230	231	232
type 23	233	234	235	236
type 24	237	238	239	240
type 25	241	242	243	244
type 26	245	246	247	248
type 27	249	250	251	252
type 28	253	254	255	256
type 29	257	258	259	260
type 30	261	262	263	264
type 31	265	266	267	268
type 32	269	270	271	272
type 33	273	274	275	276
type 34	277	278	279	280
type 35	281	282	283	284
type 36	285	286	287	288

	SEQ ID NO			
	genomic DNA	copy(c)DNA	mRNA	protein
C-terminus (minimum sequence) of mutant calreticulin	1	2	3	4
full length wild-type calreticulin	289			290
Exon 9 of mutant calreticulin				
type 1	291	292	293	294
type 2	295	296	297	298
type 3	299	300	301	302
type 4	303	304	305	306
type 5	307	308	309	310
type 6	311	312	313	314
type 7	315	316	317	318
type 8	319	320	321	322
type 9	323	324	325	326
type 10	327	328	329	330
type 11	331	332	333	334
type 12	335	336	337	338
type 13	339	340	341	342
type 14	343	344	345	346
type 15	347	348	349	350
type 16	351	352	353	354
type 17	355	356	357	358
type 18	359	360	361	362
type 19	363	364	365	366
type 20	367	368	369	370
type 21	371	372	373	374
type 22	375	376	377	378
type 23	379	380	381	382
type 24	383	384	385	386
type 25	387	388	389	390
type 26	391	392	393	394
type 27	395	396	397	398
type 28	399	400	401	402
type 29	403	404	405	406
type 30	407	408	409	410
type 31	411	412	413	414
type 32	415	416	417	418
type 33	419	420	421	422
type 34	423	424	425	426
type 35	427	428	429	430

	SEQ ID NO			
	genomic DNA	copy(c)DNA	mRNA	protein
C-terminus (minimum sequence) of mutant calreticulin	1	2	3	4
type 36	431	432	433	434
Exon 9 of wild-type calreticulin	435			436

References

[0360]

Barbui T, Barosi G, Birgegard G, et al. Philadelphia-negative classical myeloproliferative neoplasms: critical concepts and management recommendations from European LeukemiaNet. *J Clin Oncol* 2011;29:761-70.

Baxter EJ, Scott LM, Campbell PJ, East C, Fourouclas N, Swanton S, Vassiliou GS, Bench AJ, Boyd EM, Curtin N, Scott MA, Erber WN, Green AR (2005) Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet* 365(9464): 1054-1061

Campbell PJ, Green AR (2006a) The myeloproliferative disorders. *N Engl J Med* 355(23): 2452-2466

Campbell PJ, Griesshammer M, Dohner K, et al. V617F mutation in JAK2 is associated with poorer survival in idiopathic myelofibrosis. *Blood* 2006b;107:2098-100.

Coe H, Jung J, Groenendyk J, Prins D, Michalak M. ERp57 modulates STAT3 signaling from the lumen of the endoplasmic reticulum. *J Biol Chem* 2010;285:6725-38.

Corvinus FM, Orth C, Moriggl R, et al. Persistent STAT3 activation in colon cancer is associated with enhanced cell proliferation and tumor growth. *Neoplasia* 2005;7:545-55.

Delhommeau F, Dupont S, Della Valle V, James C, Trannoy S, Masse A, Kosmider O, Le Couedic JP, Robert F, Alberdi A, Lecluse Y, Plo I, Dreyfus FJ, Marzac C, Casadevall N, Lacombe C, Romana SP, Dessen P, Soulier J, Viguie F, Fontenay M, Vainchenker W, Bernard OA (2009) Mutation in TET2 in myeloid cancers. *N Engl J Med* 360(22): 2289-2301

DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del Angel G, Rivas MA, Hanna M, McKenna A, Fennell TJ, Kernysky AM, Sivachenko AY, Cibulskis K, Gabriel SB, Altshuler D, Daly MJ (2011) A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet* 43(5): 491-498

Ernst T, Chase AJ, Score J, Hidalgo-Curtis CE, Bryant C, Jones AV, Waghorn K, Zoi K, Ross FM, Reiter A, Hochhaus A, Drexler HG, Duncombe A, Cervantes F, Oscier D, Boulwood J, Grand FH, Cross NC (2010) Inactivating mutations of the histone methyltransferase gene EZH2 in myeloid disorders. *Nat Genet* 42(8): 722-726

Harutyunyan A, Klampfl T, Cazzola M, Kralovics R (2011) p53 lesions in leukemic transformation. *N Engl J Med* 364(5): 488-490

Kalbfleisch JD, Prentice RL. The statistical analysis of failure time data. New York: Wiley; 1980.

James C, Ugo V, Le Couedic JP, Staerk J, Delhommeau F, Lacout C, Garcon L, Raslova H, Berger R, Bennaceur-Griscelli A, Villeval JL, Constantinescu SN, Casadevall N, Vainchenker W (2005) A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. *Nature* 434(7037): 1144-1148

Klampfl T, Harutyunyan A, Berg T, Gisslinger B, Schalling M, Bagienski K, Olcaydu D, Passamonti F, Rumi E, Pietra

D, Jager R, Pieri L, Guglielmelli P, Iacobucci I, Martinelli G, Cazzola M, Vannucchi AM, Gisslinger H, Kralovics R (2011) Genome integrity of myeloproliferative neoplasms in chronic phase and during disease progression. *Blood* 118(1): 167-176

Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding L, Wilson RK (2012) VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res* 22(3): 568-576

Kralovics R (2008) Genetic complexity of myeloproliferative neoplasms. *Leukemia* 22(10): 1841-1848

Kralovics R, Passamonti F, Buser AS, Teo SS, Tiedt R, Passweg JR, Tichelli A, Cazzola M, Skoda RC (2005) A gain-of-function mutation of JAK2 in myeloproliferative disorders. *N Engl J Med* 352(17): 1779-1790

Levine RL, Wadleigh M, Cools J, Ebert BL, Wernig G, Huntly BJ, Boggon TJ, Wlodarska I, Clark JJ, Moore S, Adelsperger J, Koo S, Lee JC, Gabriel S, Mercher T, D'Andrea A, Frohling S, Dohner K, Marynen P, Vandenberghe P, Mesa RA, Tefferi A, Griffin JD, Eck MJ, Sellers WR, Meyerson M, Golub TR, Lee SJ, Gilliland DG (2005) Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis. *Cancer Cell* 7(4): 387-397

Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25(14): 1754-1760

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25(16): 2078-2079

Li S, Kralovics R, De Libero G, Theodorides A, Gisslinger H, Skoda RC. Clonal heterogeneity in polycythemia vera patients with JAK2 exon12 and JAK2-V617F mutations. *Blood* 2008;111:3863-6.

Malcovati L, Della Porta MG, Pietra D, et al. Molecular and clinical features of refractory anemia with ringed sideroblasts associated with marked thrombocytosis. *Blood* 2009;114:3538-45.

Marchioli R, Finazzi G, Specchia G, et al. Cardiovascular events and intensity of treatment in polycythemia vera. *N Engl J Med* 2013;368:22-33.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel S, Daly M, DePristo MA (2010) The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 20(9): 1297-1303

Milosevic JD, Kralovics R (2013) Genetic and epigenetic alterations of myeloproliferative disorders. *Int J Hematol*

Pardanani AD, Levine RL, Lasho T, Pikman Y, Mesa RA, Wadleigh M, Steensma DP, Elliott MA, Wolanskyj AP, Hogan WJ, McClure RF, Litzow MR, Gilliland DG, Tefferi A (2006) MPL515 mutations in myeloproliferative and other myeloid disorders: a study of 1182 patients. *Blood* 108(10): 3472-3476

Passamonti F, Rumi E, Pungolino E, et al. Life expectancy and prognostic factors for survival in patients with polycythemia vera and essential thrombocythemia. *Am J Med.* 2004;117(10):755-61.

Passamonti F, Rumi E, Pietra D, et al. A prospective study of 338 patients with polycythemia vera: the impact of JAK2 (V617F) allele burden and leukocytosis on fibrotic or leukemic disease transformation and vascular complications. *Leukemia* 2010a;24:1574-9.

Passamonti F, Cervantes F, Vannucchi AM, et al. A dynamic prognostic model to predict survival in primary myelofibrosis: a study by the IWG-MRT (International Working Group for Myeloproliferative Neoplasms Research and Treatment). *Blood* 2010b;115:1703-8.

Passamonti F, Elena C, Schnittger S, et al. Molecular and clinical features of the myeloproliferative neoplasm associated with JAK2 exon 12 mutations. *Blood* 2011;117:2813-6.

Passamonti F, Thiele J, Girodon F, et al. A prognostic model to predict survival in 867 World Health Organization-defined essential thrombocythemia at diagnosis: a study by the International Working Group on Myelofibrosis

Research and Treatment. *Blood* 2012;120:1197-201.

Pikman Y, Lee BH, Mercher T, McDowell E, Ebert BL, Gozo M, Cuker A, Wernig G, Moore S, Galinsky I, DeAngelo DJ, Clark JJ, Lee SJ, Golub TR, Wadleigh M, Gilliland DG, Levine RL (2006) MPLW515L is a novel somatic activating mutation in myelofibrosis with myeloid metaplasia. *PLoS Med* 3(7): e270

R Core Team. R: A language and environment for statistical computing.: R Foundation for Statistical Computing, Vienna, Austria.; 2012.

Rumi E, Pietra D, Guglielmelli P, et al. Acquired copy-neutral loss of heterozygosity of chromosome 1p as a molecular event associated with marrow fibrosis in MPL-mutated myeloproliferative neoplasms. *Blood* 2013;121:4388-95.

Schafer AI. Thrombocytosis. *N Engl J Med* 2004;350:1211-9.

Scott LM, Tong W, Levine RL, Scott MA, Beer PA, Stratton MR, Futreal PA, Erber WN, McMullin MF, Harrison CN, Warren AJ, Gilliland DG, Lodish HF, Green AR (2007) JAK2 exon 12 mutations in polycythemia vera and idiopathic erythrocytosis. *N Engl J Med* 356(5): 459-468

Si J, Collins SJ. Activated Ca²⁺/calmodulin-dependent protein kinase IIgamma is a critical regulator of myeloid leukemia cell proliferation. *Cancer Res* 2008;68:3733-42.

Stegelmann F, Bullinger L, Schlenk RF, Paschka P, Griesshammer M, Biersch C, Kuhn S, Schauer S, Dohner H, Dohner K (2011) DNMT3A mutations in myeloproliferative neoplasms. *Leukemia* 25(7): 1217-1219

Stein BL, Williams DM, O'Keefe C, Rogers O, Ingersoll RG, Spivak JL, Verma A, Maciejewski JP, McDevitt MA, Moliterno AR (2011) Disruption of the ASXL1 gene is frequent in primary, post-essential thrombocytosis and post-polycythemia vera myelofibrosis, but not essential thrombocytosis or polycythemia vera: analysis of molecular genetics and clinical phenotypes. *Haematologica* 96(10): 1462-1469

Swerdlow S, Campo E, Harris N, Jaffe E, Pileri S, Stein H, Thiele J, Vardiman J (2008) WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, Lyon: International Agency for Research on Cancer.

Thiele J, Kvasnicka HM, Facchetti F, Franco V, van der Walt J, Orazi A. European consensus on grading bone marrow fibrosis and assessment of cellularity. *Haematologica* 2005;90:1128-32.

Wang K, Li M, Hakonarson H (2010) ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res* 38(16): e164

Vardiman JW, Harris NL, Brunning RD. The World Health Organization (WHO) classification of the myeloid neoplasms. *Blood* 2002;100:2292-302.

Yue X, Wang H, Zhao F, et al. Hepatitis B virus-induced calreticulin protein is involved in IFN resistance. *J Immunol* 2012;189:279-86.

Zuber J, McJunkin K, Fellmann C, et al. Toolkit for evaluating genes required for proliferation and survival using tetracycline-regulated RNAi. *Nature biotechnology* 2011;29:79-83.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US5525711A](#) [0129]
- [US4711955A](#) [0129]
- [US5792608A](#) [0129]
- [EP302175A](#) [0129]
- [US3773919A](#) [0248]
- [EP58481A](#) [0248]
- [EP133988A](#) [0248]
- [DE3218121](#) [0248]
- [EP52322A](#) [0248]
- [EP36676A](#) [0248]
- [EP86046A](#) [0248]
- [EP143949A](#) [0248]
- [EP142641A](#) [0248]
- [JP58118008A](#) [0248]
- [US4485045A](#) [0248]
- [US4544545A](#) [0248]
- [EP102324A](#) [0248]
- [WO9420627A](#) [0290]
- [WO9429469A](#) [0292]
- [WO9700957A](#) [0292]
- [US5580859A](#) [0292]
- [US5589466A](#) [0292]

Non-patent literature cited in the description

- **VARDIMAN et al.** The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: Rationale and important changes *Blood*, 2009, vol. 114, 5937-51 [0020]
- **BROSEUS et al.** Clinical features and course of refractory anemia with ring sideroblast associated with marked thrombocytosis *Haematologica*, 2012, vol. 9, 71036-1041 [0026]
- **KRALOVICS R TEO SSSI THEOCHARIDES ABUSER ASTICHELLI ASKODA RC** Acquisition of the V617F mutation of JAK2 is a late genetic event in a subset of patients with myeloproliferative disorders *Blood*, 2006, vol. 15;108, 41377-80 [0031]
- **GUNDRY et al.** Amplicon Melting Analysis with Labeled Primers: A Closed-Tube Method for Differentiating Homozygotes and Heterozygotes *Clinical Chemistry*, 2003, vol. 49, 396-406 [0098]
- **COLESTRAUSS** Science, 1996, 1386-1389 [0129]
- **SAMBROOKRUSSELL** Molecular Cloning, A Laboratory Manual Cold Spring Harbor Laboratory 20010000 [0137]
- **AUSUBEL** Current Protocols in Molecular Biology Green Publishing Associates and Wiley Interscience 19890000 [0137]
- Nucleic acid hybridization, a practical approach IRL Press Oxford 19850000 [0137]
- Thompson *Nucl. Acids Res.*, 1994, vol. 2, 4673-4680 [0139]
- Brutlag *Comp. App. Biosci.*, 1990, vol. 6, 237-245 [0139]
- **ALTSCHUL** *Nucl. Acids Res.*, 1997, vol. 25, 3389-3402 [0140]
- **ALTSCHUL** *J. Mol. Evol.*, 1993, vol. 36, 290-300 [0140]
- **ALTSCHUL** *J. Mol. Biol.*, 1990, vol. 215, 403-410 [0140]
- **HENIKOFF** *PNAS*, 1989, vol. 89, 10915- [0140]
- **THOMPSON** *Nucl. Acids Res.*, 1994, vol. 2, 4673-4680 [0142]

- **BRUTLAG**Comp. App. Biosci., 1990, vol. 6, 237-245 [0142]
- **MUTSCHLER**ArzneimittelwirkungenWissenschaftliche Verlagsgesellschaft mbH19860000 [0178]
- **Antibodies: A Laboratory Manual**Cold Spring Harbor Laboratory19881200 [0191]
- **Portable Protocols (Using Antibodies): A Laboratory Manual**Cold Spring Harbor Laboratory19981200 [0191]
- **SIDMAN, U. et al.**Biopolymers, 1983, vol. 22, 547-556 [0248]
- **R. LANGER et al.**J. Biomed. Mater. Res., 1981, vol. 15, 167-277 [0248]
- **R. LANGER**Chem. Tech., 1982, vol. 12, 98-105 [0248]
- **EPSTEIN et al.**Proc. Natl. Acad. Sci. (USA), 1985, vol. 82, 3688-3692 [0248]
- **HWANG et al.**Proc. Natl. Acad. Sci. (USA), 1980, vol. 77, 4030-4034 [0248]
- **for use in the treatment of high and intermediate risk myelofibrosis**Food and Drug Administration, 2011, [0256]
- **TEFFER**Blood, 2012, vol. 119, 12- [0256] [0257]
- **OSTOJIC**Therapeutics and Clinical Risk Management, 2012, vol. 8, 95-103 [0256]
- **Methods in Enzymology**, 1987, vol. 153, 385-516 [0278]
- **BITTER et al.**Methods in Enzymology, 1987, vol. 153, 516-544 [0278]
- **SAWERS et al.**Applied Microbiology and Biotechnology, 1996, vol. 46, 1-9 [0278]
- **BILLMAN-JACOB**Current Opinion in Biotechnology, 1996, vol. 7, 500-4 [0278]
- **HOCKNEY**Trends in Biotechnology, 1994, vol. 12, 456-463 [0278]
- **GRIFFITHS et al.**Methods in Molecular Biology, 1997, vol. 75, 427-440 [0278]
- **SAMBROOKAUSUBEL et al.**Current Protocols in Molecular BiologyGreen Publishing Associates and Wiley Interscience19890000 [0281]
- **MACK et al.**PNAS, 1995, vol. 92, 7021-7025 [0286]
- **RAUM et al.**Cancer Immunol Immunother, 2001, vol. 50, 3141-150 [0286]
- **SMITH**J. Virol., 1983, vol. 46, 584- [0288]
- **ENGELHARD**Proc. Nat. Acad. Sci. USA, 1994, vol. 91, 3224-3227 [0288]
- **REISS**Plant Physiol. (Life Sci. Adv.), 1994, vol. 13, 143-149 [0290]
- **HERRERA-ESTRELLA**EMBO J., 1983, vol. 2, 987-995 [0290]
- **MARSH**Gene, 1984, vol. 32, 481-485 [0290]
- **HARTMAN**Proc. Natl. Acad. Sci. USA, 1988, vol. 85, 8047- [0290]
- **MCCONLOGUE**Current Communications in Molecular Biology19870000 [0290]
- **TAMURA**Biosci. Biotechnol. Biochem., 1995, vol. 59, 2336-2338 [0290]
- **GIACOMINI**Pl. Sci., 1996, vol. 116, 59-72 [0291]
- **SCI KANTHA**J. Bact., 1996, vol. 178, 121- [0291]
- **GERDES**FEBS Lett, 1996, vol. 389, 44-47 [0291]
- **JEFFERSON**EMBO J., 1987, vol. 6, 3901-3907 [0291]
- **GIORDANO**Nature Medicine, 1996, vol. 2, 534-539 [0292]
- **SCHAPER**Circ. Res., 1996, vol. 79, 911-919 [0292]
- **ANDERSON**Science, 1992, vol. 256, 808-813 [0292]
- **VERMA**Nature, 1994, vol. 389, 239- [0292]
- **ISNER**Lancet, 1996, vol. 348, 370-374 [0292]
- **MUHLHAUSER**Circ. Res., 1995, vol. 77, 1077-1086 [0292]
- **ONODERA**Blood, 1998, vol. 91, 30-36 [0292]
- **VERMA**Gene Ther, 1998, vol. 5, 692-699 [0292]
- **NABEL**Ann. N.Y. Acad. Sci., 1997, vol. 811, 289-292 [0292]
- **VERZELETTI**Hum. Gene Ther., 1998, vol. 9, 2243-51 [0292]
- **WANG**Nature Medicine, 1996, vol. 2, 714-716 [0292]
- **SCHAPER**Current Opinion in Biotechnology, 1996, vol. 7, 635-640 [0292]
- **NAGY**Proc. Natl. Acad. Sci. USA, 1993, vol. 90, 8424-8428 [0292]
- **KROOS**Biotechnol. Prog., 2003, vol. 19, 163-168 [0298]
- **SCOPES**Protein PurificationSpringer-Verlag19820000 [0302]
- **BARBUI T BAROSI G BIRGE GARD G et al.**Philadelphia-negative classical myeloproliferative neoplasms: critical concepts and management recommendations from European LeukemiaNetJ Clin Oncol, 2011, vol. 29, 761-70 [0360]
- **BAXTER E JSCOTT L MCAMPBELL P JEAST C FOUROUCLAS N SWANTON S VASSILIOU G SBENCH**

- AJBOYD EMCURTIN N** Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders *Lancet*, 2005, vol. 365, 94641054-1061 [0360]
- **CAMPBELL PJGREEN AR** The myeloproliferative disorders *N Engl J Med*, 2006, vol. 355, 232452-2466 [0360]
 - **CAMPBELL PJGRIESSHAMMER MDOHNER K et al.** V617F mutation in JAK2 is associated with poorer survival in idiopathic myelofibrosis *Blood*, 2006, vol. 107, 2098-100 [0360]
 - **COE HJUNG JGROENENDYK JPRINS DMICHALAK MER** p57 modulates STAT3 signaling from the lumen of the endoplasmic reticulum *J Biol Chem*, 2010, vol. 285, 6725-38 [0360]
 - **CORVINUS FMORTH CMORIGGL R et al.** Persistent STAT3 activation in colon cancer is associated with enhanced cell proliferation and tumor growth *Neoplasia*, 2005, vol. 7, 545-55 [0360]
 - **DELHOMMEAU FDUPONT SDELLA VALLE VJAMES CTRANNOY SMASSE AKOSMIDER OLE COUEDIC JPROBERT FALBERDI A** Mutation in TET2 in myeloid cancers *N Engl J Med*, 2009, vol. 360, 222289-2301 [0360]
 - **DEPRISTO MABANKS EPOPLIN RGARIMELLA KVMAGUIRE JRHARTL CPHILIPPAKIS AADEL ANGEL GRIVAS MAHANNA MA** framework for variation discovery and genotyping using next-generation DNA sequencing data *Nat Genet*, 2011, vol. 43, 5491-498 [0360]
 - **ERNST TCHASE AJSCORE JHIDALGO-CURTIS CEBRYANT CJONES AVWAGHORN KZOI KROSS FMREITER A** inactivating mutations of the histone methyltransferase gene EZH2 in myeloid disorders *Nat Genet*, 2010, vol. 42, 8722-726 [0360]
 - **HARUTYUNYAN AKLAMPFL TCAZZOLA MKRALOVICS R** lesions in leukemic transformation, 2011, 53- [0360]
 - *N Engl J Med*, vol. 364, 5488-490 [0360]
 - **KALBFLEISCH JDPRENTICE RL** The statistical analysis of failure time data *Wiley* 19800000 [0360]
 - **JAMES CUGO VLE COUEDIC JPSTAERK JDELHOMMEAU FLACOUT CGARCON LRASLOVA HBERGER RBENNACEUR-GRISCELLI AA** unique clonal JAK2 mutation leading to constitutive signalling causes polycythemia vera *Nature*, 2005, vol. 434, 70371144-1148 [0360]
 - **KLAMPFL THARUTYUNYAN ABERG TGISSLINGER BSCHALLING MBAGIENSKI KOLCAYDU DPASSAMONTI FRUMI EPIETRA D** Genome integrity of myeloproliferative neoplasms in chronic phase and during disease progression *Blood*, 2011, vol. 118, 1167-176 [0360]
 - **KOBOLDT DCZHANG QLARSON DESHEN DMCLELLAN MDLIN LMILLER CAMARDIS ERDING LWILSON RK** VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing *Genome Res*, 2012, vol. 22, 3568-576 [0360]
 - **KRALOVICS R** Genetic complexity of myeloproliferative neoplasms *Leukemia*, 2008, vol. 22, 101841-1848 [0360]
 - **KRALOVICS RPASSAMONTI FBUSER ASTEO SSTIEDT RPASSWEG JRTICHELLI ACAZZOLA MSKODA RCA** gain-of-function mutation of JAK2 in myeloproliferative disorders *N Engl J Med*, 2005, vol. 352, 171779-1790 [0360]
 - **LEVINE RLWADLEIGH MCOOLS JEBERT BLWERNIG GHUNTLY BJBOGGON TJWLODARSKA ICLARK JJMOORE S** Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis *Cancer Cell*, 2005, vol. 7, 4387-397 [0360]
 - **LI HDURBIN R** Fast and accurate short read alignment with Burrows-Wheeler transform *Bioinformatics*, 2009, vol. 25, 141754-1760 [0360]
 - **LI HHANDSAKER BWYSOKER AFENNELL TRUAN JHOMER NMARTH GABECASIS GDURBIN R** The Sequence Alignment/Map format and SAMtools *Bioinformatics*, 2009, vol. 25, 162078-2079 [0360]
 - **LI SKRALOVICS RDE LIBERO GTHEOCHARIDES AGISSLINGER HSKODA RC** Clonal heterogeneity in polycythemia vera patients with JAK2 exon12 and JAK2-V617F mutations *Blood*, 2008, vol. 111, 3863-6 [0360]
 - **MALCOVATI LDELLA PORTA MGPIETRA D et al.** Molecular and clinical features of refractory anemia with ringed sideroblasts associated with marked thrombocytosis *Blood*, 2009, vol. 114, 3538-45 [0360]
 - **MARCHIOLI RFINAZZI GSPECCHIA G et al.** Cardiovascular events and intensity of treatment in polycythemia vera *N Engl J Med*, 2013, vol. 368, 22-33 [0360]
 - **MCKENNA AHANNA MBANKS ESIVACHENKO ACIBULSKIS KKERNYTSKY AGARIMELLA KALTSHULER DGABRIEL SDALY M** The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data *Genome Res*, 2010, vol. 20, 91297-1303 [0360]
 - **MILOSEVIC JDKRALOVICS R** Genetic and epigenetic alterations of myeloproliferative disorders *Int J Hematol*,

2013, [0360]

- **PARDANANI ADLEVINE RLLASHO TPIKMAN YMESA RAWADLEIGH MSTEENSMA DPELLIOTT MAWOLANSKYJ APHOGAN WJMPL** 515 mutations in myeloproliferative and other myeloid disorders: a study of 1182 patients *Blood*, 2006, vol. 108, 103472-3476 [0360]
- **PASSAMONTI FRUMI EPUNGOLINO E et al.** Life expectancy and prognostic factors for survival in patients with polycythemia vera and essential thrombocythemia *Am J Med.*, 2004, vol. 117, 10755-61 [0360]
- **PASSAMONTI FRUMI EPIETRA D et al.** A prospective study of 338 patients with polycythemia vera: the impact of JAK2 (V617F) allele burden and leukocytosis on fibrotic or leukemic disease transformation and vascular complications *Leukemia*, 2010, vol. 24, 1574-9 [0360]
- **PASSAMONTI FCERVANTES FVANNUCCHI AM et al.** A dynamic prognostic model to predict survival in primary myelofibrosis: a study by the IWG-MRT (International Working Group for Myeloproliferative Neoplasms Research and Treatment) *Blood*, 2010, vol. 115, 1703-8 [0360]
- **PASSAMONTI FELENA CSCHNITTGER S et al.** Molecular and clinical features of the myeloproliferative neoplasm associated with JAK2 exon 12 mutations *Blood*, 2011, vol. 117, 2813-6 [0360]
- **PASSAMONTI FTHIELE JGIRODON F et al.** A prognostic model to predict survival in 867 World Health Organization-defined essential thrombocythemia at diagnosis: a study by the International Working Group on Myelofibrosis Research and Treatment *Blood*, 2012, vol. 120, 1197-201 [0360]
- **PIKMAN YLEE BHMERCHER TMCOWELL EEBERT BLGOZO MCUKER AVERNIG GMOORE SGALINSKY IMPLW** 515L is a novel somatic activating mutation in myelofibrosis with myeloid metaplasia *PLoS Med*, 2006, vol. 3, 7e270- [0360]
- **R CORE** Team. R: A language and environment for statistical computing *R Foundation for Statistical Computing*, 2012, [0360]
- **RUMI EPIETRA DGUGLIELMELLI P et al.** Acquired copy-neutral loss of heterozygosity of chromosome 1p as a molecular event associated with marrow fibrosis in MPL-mutated myeloproliferative neoplasms *Blood*, 2013, vol. 121, 4388-95 [0360]
- **SCHAFER AI** Thrombocytosis *N Engl J Med*, 2004, vol. 350, 1211-9 [0360]
- **SCOTT LMTONG WLEVINE RLSCOTT MABEER PASTRATTON MRFUTREAL PAERBER WNMCMULLIN MFHARRISON CN** JAK2 exon 12 mutations in polycythemia vera and idiopathic erythrocytosis *N Engl J Med*, 2007, vol. 356, 5459-468 [0360]
- **SI JCOLLINS SJ** Activated Ca²⁺/calmodulin-dependent protein kinase IIgamma is a critical regulator of myeloid leukemia cell proliferation *Cancer Res*, 2008, vol. 68, 3733-42 [0360]
- **STEGELMANN FBULLINGER LSCHLENK RFPASCHKA PGRIESSHAMMER MBLERSCH CKUHN SSCHAUER SDOHNER HDOHNER KDNMT3A** mutations in myeloproliferative neoplasms *Leukemia*, 2011, vol. 25, 71217-1219 [0360]
- **STEIN BLWILLIAMS DMO'KEEFE CROGERS OINGERSOLL RGSPIVAK JLVERMA AMACIEJEWSKI JPMCDEVITT MAMOLITERNO AR** Disruption of the ASXL1 gene is frequent in primary, post-essential thrombocytosis and post-polycythemia vera myelofibrosis, but not essential thrombocytosis or polycythemia vera: analysis of molecular genetics and clinical phenotypes *Haematologica*, 2011, vol. 96, 101462-1469 [0360]
- **SWERDLOW SCAMPO EHARRIS NJAFFE EPILERI SSTEIN HTHIELE JVARDIMAN J** WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, Lyon *International Agency for Research on Cancer*, 2008, [0360]
- **THIELE JKVASNICKA HMFACCHETTI FFRANCO VVAN DER WALT JORAZI A** European consensus on grading bone marrow fibrosis and assessment of cellularity *Haematologica*, 2005, vol. 90, 1128-32 [0360]
- **WANG KLI MHAKONARSON HANNOVAR**: functional annotation of genetic variants from high-throughput sequencing data *Nucleic Acids Res*, 2010, vol. 38, 16e164- [0360]
- **VARDIMAN JWHARRIS NLBRUNNING RD** The World Health Organization (WHO) classification of the myeloid neoplasms *Blood*, 2002, vol. 100, 2292-302 [0360]
- **YUE XWANG HZHAO F et al.** Hepatitis B virus-induced calreticulin protein is involved in IFN resistance *J Immunol*, 2012, vol. 189, 279-86 [0360]
- **ZUBER JMCJUNKIN KFELLMANN C et al.** Toolkit for evaluating genes required for proliferation and survival using tetracycline-regulated RNAi *Nature biotechnology*, 2011, vol. 29, 79-83 [0360]

P a t e n t k r a v

1. Fremgangsmåde til vurdering af, om en patient lider af en myeloid malignitet eller er tilbøjelig til at lide af en myeloid malignitet, hvilken fremgangsmåde
5 omfatter

- at bestemme tilstedeværelsen af et genprodukt af et eller flere mutante alleler af calreticulin-genet i en prøve fra patienten; og

- at vurdere, at patienten lider af en myeloid malignitet eller er tilbøjelig til at lide af en myeloid malignitet, når genproduktet er til stede,

10 hvor det ene eller de flere mutante alleler omfatter nukleinsyre, der koder for et mutant calreticulin-protein, hvor det mutante calreticulin-protein er udvalgt fra gruppen bestående af

(a) et protein, der er kodet af et nukleinsyremolekyle med nukleinsyresekvensen som gengivet i SEQ ID NO: 1, 2 eller 3;

15 (b) et protein omfattende aminosyresekvensen som vist i SEQ ID NO: 4,

(c) et protein som defineret i (a) eller (b), hvor en til 10 aminosyrer er deleteret, insereret, tilsat eller substitueret

(d) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, der hybridiserer under stringente forhold til den komplementære streng af nukleinsyremolekyler som defineret i (a) eller (c), og hvor proteinet er en mutant
20 af calreticulin-genet;

(e) et protein med mindst 70 % identitet med proteinet i et hvilket som helst af (a) til (d); og

(f) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, som er degenereret som følge af den genetiske kode til nukleotidsekvensen af en nukleinsyre som defineret i (a) eller (d),
25

hvor genproduktet er protein, og

hvor tilstedeværelsen eller mængden af proteinet bestemmes ved immunhistokemi (IHC), by immunoassay, gel- eller blotbaserede fremgangsmåder, massespektrometri, flowcytometri eller FACS.
30

2. Fremgangsmåde ifølge krav 1, hvor det mutante calreticulin-protein er udvalgt fra gruppen bestående af

(a) et protein, der er kodet af et nukleinsyremolekyle med nukleinsyresekvensen som gengivet i SEQ ID NO: 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21,
35 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49,

50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, 143, 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432 eller 433; og

(b) et protein omfattende aminosyresekvensen som vist i SEQ ID NO: 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430 eller 434.

3. Fremgangsmåde ifølge krav 1 eller 2, hvor det mutante allel omfatter en nukleinsyre udvalgt fra gruppen bestående af

(a) en nukleinsyre, der koder for et polypeptid omfattende en aminosyresekvens som gengivet i SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302,

306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430 eller 434; og

- 5 (b) en nukleinsyre omfattende en nukleotidsekvens som gengivet i SEQ ID NO: 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45, 49, 53, 57, 61, 65, 69, 73, 77, 81, 85, 89, 93, 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189, 193, 197, 201, 205, 209, 213, 217, 221, 225, 229, 233, 237, 241, 245, 249, 253, 257, 261, 265, 269, 273, 277, 281, 285, 291, 295, 299, 303, 307, 311, 315, 319, 323, 327, 10 331, 335, 339, 343, 347, 351, 355, 359, 363, 367, 371, 375, 379, 383, 387, 391, 395, 399, 403, 407, 411, 415, 419, 423, 427 eller 431.

4. Fremgangsmåde ifølge et hvilket som helst af kravene 1 to 3, hvor genproduktet omfatter et polypeptid udvalgt fra gruppen bestående af

- 15 (a) et protein, der er kodet af et nukleinsyremolekyle med nukleinsyresekvensen som gengivet i SEQ ID NO: 1, 2 eller 3;
 (b) et protein omfattende aminosyresekvensen som vist i SEQ ID NO: 4;
 (c) et protein som defineret i (a) eller (b), hvor en til 10 aminosyrer er deleteret, insereret, tilsat eller substitueret;
 20 (d) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, der hybridiserer under stringente forhold til den komplementære streng af nukleinsyremolekyler som defineret i (a) eller (c), og hvor proteinet er en mutant af calreticulin-genet;
 (e) et protein med mindst 70 % identitet med proteinet i et hvilket som helst af
 25 (a) til (d); og
 (f) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, som er degenereret som følge af den genetiske kode til nukleotidsekvensen af en nukleinsyre som defineret i (a) eller (d).

5. Fremgangsmåde ifølge et hvilket som helst af kravene 1 to 4, hvor genproduktet omfatter et polypeptid udvalgt fra gruppen bestående af

- (a) et protein, der er kodet af et nukleinsyremolekyle med nukleinsyresekvensen som gengivet i SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 35 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101,

- 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142, 143, 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286, 287, 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432 eller 433;
- (b) et protein omfattende aminosyresekvensen som vist i SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284, 288, 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430 eller 434;
- (c) et protein som defineret i (a) eller (b), hvor en til 10 aminosyrer er deleteret, insereret, tilsat eller substitueret;
- (d) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, der hybridiserer under stringente forhold til den komplementære streng af nukleinsyremolekyler som defineret i (a) eller (c), og hvor proteinet er en mutant af calreticulin-genet
- (e) et protein med mindst 70 % identitet med proteinet i et hvilket som helst af (a) til (d); og
- (f) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, som er degenereret som følge af den genetiske kode til nukleotidsekvensen af en nukleinsyre som defineret i (a) eller (d).

6. Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 5, hvor prøven er en knoglemarvsprøve eller en blodprøve.

5 **7.** Fremgangsmåde ifølge krav 6, hvor knoglemarvsprøven opnås ved hjælp af biopsi.

8. Protein, der er udvalgt fra gruppen bestående af

10 (a) et protein, der er kodet af et nukleinsyremolekyle med nukleinsyresekvensen som gengivet i SEQ ID NO: 1, 2 eller 3;

(b) et protein omfattende aminosyresekvensen som vist i SEQ ID NO: 4;

(c) et protein som defineret i (a) eller (b), hvor en til 10 aminosyrer er deleteret, insereret, tilsat eller substitueret;

15 (d) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, der hybridiserer under stringente forhold til den komplementære streng af nukleinsyremolekyler som defineret i (a) eller (c), og hvor proteinet er en mutant af calreticulin-genet;

(e) et protein med mindst 70 % identitet med proteinet i et hvilket som helst af (a) til (d); og

20 (f) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, som er degenereret som følge af den genetiske kode til nukleotidsekvensen af en nukleinsyre som defineret i (a) eller (d).

9. Protein ifølge krav 8, hvor proteinet er udvalgt fra gruppen bestående af

25 (a) et protein, der er kodet af et nukleinsyremolekyle med nukleinsyresekvensen som gengivet i SEQ ID NO: 1, 2, 3, 5, 6, 7, 9, 10, 11, 13, 14, 15, 17, 18, 19, 21, 22, 23, 25, 26, 27, 29, 30, 31, 33, 34, 35, 37, 38, 39, 41, 42, 43, 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 97, 98, 99, 101, 30 102, 103, 105, 106, 107, 109, 110, 111, 113, 114, 115, 117, 118, 119, 121, 122, 123, 125, 126, 127, 129, 130, 131, 133, 134, 135, 137, 138, 139, 141, 142 eller 143;

(b) et protein omfattende aminosyresekvensen som vist i SEQ ID NO: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 35 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140 eller 144;

(c) et protein som defineret i (a) eller (b), hvor en til 10 aminosyrer er deleteret, insereret, tilsat eller substitueret;

(d) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, der hybridiserer under stringente forhold til den komplementære streng af nukleinsyremolekyler som defineret i (a) eller (c), og hvor proteinet kodes af et eller flere mutante alleler af calreticulin-genet;

(e) et protein med mindst 70 % identitet med proteinet i et hvilket som helst af (a) til (d); og

(f) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, som er degenereret som følge af den genetiske kode til nukleotidsekvensen af en nukleinsyre som defineret i (a) eller (d).

10. Protein ifølge krav 8 eller 9, hvor proteinet er udvalgt fra gruppen bestående af

(a) et protein, der er kodet af et nukleinsyremolekyle med nukleinsyresekvensen som gengivet i SEQ ID NO: 291, 292, 293, 295, 296, 297, 299, 300, 301, 303, 304, 305, 307, 308, 309, 311, 312, 313, 315, 316, 317, 319, 320, 321, 323, 324, 325, 327, 328, 329, 331, 332, 333, 335, 336, 337, 339, 340, 341, 343, 344, 345, 347, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 361, 363, 364, 365, 367, 368, 369, 371, 372, 373, 375, 376, 377, 379, 380, 381, 383, 384, 385, 387, 388, 389, 391, 392, 393, 395, 396, 397, 399, 400, 401, 403, 404, 405, 407, 408, 409, 411, 412, 413, 415, 416, 417, 419, 420, 421, 423, 424, 425, 427, 428, 429, 431, 432 eller 433;

(b) et protein omfattende aminosyresekvensen som vist i SEQ ID NO: 294, 298, 302, 306, 310, 314, 318, 322, 326, 330, 334, 338, 342, 346, 350, 354, 358, 362, 366, 370, 374, 378, 382, 386, 390, 394, 398, 402, 406, 410, 414, 418, 422, 426, 430 eller 434;

(c) et protein som defineret i (a) eller (b), hvor en til 10 aminosyrer er deleteret, insereret, tilsat eller substitueret;

(d) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, der hybridiserer under stringente forhold til den komplementære streng af nukleinsyremolekyler som defineret i (a) eller (c), og hvor proteinet er en mutant af calreticulin-genet;

(e) et protein med mindst 70 % identitet med proteinet i et hvilket som helst af (a) til (d); og

(f) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, som er degenereret som følge af den genetiske kode til nukleotidsekvensen af en nukleinsyre som defineret i (a) eller (d).

5 **11.** Protein ifølge et hvilket som helst af kravene 8 til 10, hvor proteinet er udvalgt fra gruppen bestående af

10 (a) et protein, der er kodet af et nukleinsyremolekyle med nukleinsyresekvensen som gengivet i SEQ ID NO: 145, 146, 147, 149, 150, 151, 153, 154, 155, 157, 158, 159, 161, 162, 163, 165, 166, 167, 169, 170, 171, 173, 174, 175, 177, 178, 179, 181, 182, 183, 185, 186, 187, 189, 190, 191, 193, 194, 195, 197, 198, 199, 201, 202, 203, 205, 206, 207, 209, 210, 211, 213, 214, 215, 217, 218, 219, 221, 222, 223, 225, 226, 227, 229, 230, 231, 233, 234, 235, 237, 238, 239, 241, 242, 243, 245, 246, 247, 249, 250, 251, 253, 254, 255, 257, 258, 259, 261, 262, 263, 265, 266, 267, 269, 270, 271, 273, 274, 275, 277, 278, 279, 281, 282, 283, 285, 286 eller 287;

15 (b) et protein omfattende aminosyresekvensen som vist i SEQ ID NO: 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188, 192, 196, 200, 204, 208, 212, 216, 220, 224, 228, 232, 236, 240, 244, 248, 252, 256, 260, 264, 268, 272, 276, 280, 284 eller 288;

20 (c) et protein som defineret i (a) eller (b), hvor en til 10 aminosyrer er deleteret, insereret, tilsat eller substitueret;

25 (d) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, der hybridiserer under stringente forhold til den komplementære streng af nukleinsyremolekyler som defineret i (a) eller (c), og hvor proteinet er en mutant af calreticulin-genet;

 (e) et protein med mindst 70 % identitet med proteinet i et hvilket som helst af (a) til (d); og

30 (f) et protein omfattende en aminosyresekvens, der kodes af en nukleinsyre, som er degenereret som følge af den genetiske kode til nukleotidsekvensen af en nukleinsyre som defineret i (a) eller (d).

12. Vaccine omfattende proteinet ifølge krav 8 eller 9.

35 **13.** Protein ifølge krav 8 eller 9, såsom et protein omfattende 15 til 25 sammenhængende aminosyrer af proteinet som vist i SEQ ID NO: 4, til anvendelse

som vaccine.

14. Immunogent middel, hvor det immunogene middel er et mutant calreticulin-protein ifølge et hvilket som helst af kravene 8 til 11.

5

15. Konjugat omfattende det immunogene middel ifølge krav 14, bundet til et bæreprøtein.

16. Antistof, der binder specifikt til den C-terminale ende af mutant(e) calreticulin-protein(er) som vist i SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140 eller 144.

10

17. Immunogent middel ifølge krav 14 eller antistof ifølge krav 16 til anvendelse ved behandling af en myeloid malignitet.

15

18. Anvendelse af et antistof, der binder specifikt til proteinet ifølge et hvilket som helst af kravene 8 til 11 til vurdering af, om en patient lider af en myeloid malignitet eller er tilbøjelig til at lide af en myeloid malignitet.

20

19. Antistof, der binder specifikt til den C-terminale ende af mutant(e) calreticulin-protein(er) som vist i SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140 eller 144 til fremstilling af et diagnosekit til anvendelse i fremgangsmåden ifølge et hvilket som helst af kravene 1 til 7.

25

20. Anvendelse ifølge krav 18, hvor antistoffet binder specifikt til den C-terminale ende af mutant(e) calreticulin-protein(er) som vist i SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140 eller 144.

30

21. Antistof ifølge krav 16, eller antistof til anvendelse ifølge krav 17, eller anvendelse ifølge krav 18 eller 20, hvor antistoffet er et polyklonalt antistof, et monoklonalt antistof, et komplet antistof (immunoglobulin), et F(ab)-fragment, et F(ab)₂-fragment, et enkeltkædet antistof, et kimært antistof, et CDR-podet antistof, et bivalent antistof-konstrukt, et bispecifikt enkeltkædet antistof, et

35

syntetisk antistof eller et krydsklonet antistof.

- 5 **22.** Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 7, anvendelse ifølge et hvilket som helst af kravene 18, 20 og 21 eller antistof til anvendelse ifølge krav 17 eller 21 eller immunogent middel til anvendelse ifølge krav 17, hvor den myeloide malignitet er en myeloproliferativ neoplasme, eller hvor den myeloide malignitet er et myelodysplastisk syndrom.
- 10 **23.** Fremgangsmåde ifølge krav 22, eller anvendelse ifølge krav 22, eller antistof til anvendelse ifølge krav 22 eller immunogent middel til anvendelse ifølge krav 22, hvor den myeloproliferative neoplasme er primær myelofibrose (PMF), eller hvor den myeloproliferative neoplasme er essentiel trombocytæmi (ET).
- 15 **24.** Fremgangsmåde ifølge krav 22 eller anvendelse ifølge krav 22 eller antistof til anvendelse ifølge krav 22 eller immunogent middel til anvendelse ifølge krav 22, hvor det myelodysplastiske syndrom er refraktær anæmi med ring-sideroblaster og trombocytæmi (RARS-T).
- 20 **25.** Protein ifølge krav 8 eller 9 eller antistof ifølge krav 16 eller 21 til anvendelse som et medikament.
- 25 **26.** Proces til fremstilling af polypeptidet ifølge et hvilket som helst af kravene 8 til 11, hvilken proces omfatter
at dyrke værtsceller omfattende en nukleinsyre udvalgt fra gruppen bestående af
- 30 (a) en nukleinsyre, der koder for et polypeptid omfattende en aminosyresekvens som gengivet i SEQ ID NO: 4;
(b) en nukleinsyre omfattende en nukleotidsekvens som gengivet i SEQ ID NO: 2;
(c) en nukleinsyre, der hybridiserer under stringente forhold til den komplementære streng af nukleinsyren som defineret i (a) eller (b), og hvor nukleinsyren omfatter en frameshift-mutation i exon 9 sammenlignet med vildtype-calreticulin-genet;

(d) en nukleinsyre omfattende en nukleotidsekvens med mindst 70 % identitet med nukleotidsekvensen af nukleinsyrerne ifølge et hvilket som helst af (a) til (c); og

5 (e) en nukleinsyre omfattende en nukleotidsekvens, som er degenereret som som følge af den genetiske kode til nukleotidsekvensen af en nukleinsyre ifølge et hvilket som helst af (a) til (d); eller

at dyrke værtsceller omfattende en vektor omfattende nukleinsyren under betingelser, der muliggør ekspression af polypeptidet, og udvinde det fremstillede polypeptid fra kulturen.

10

27. Transgen celle eller transgent ikke-menneskeligt dyr, der omfatter og/eller eksprimerer en nukleinsyre eller en vektor omfattende nukleinsyren, hvor nukleinsyren omfatter mindst et eller flere mutante alleler af calreticulin-genet som defineret i et hvilket som helst af kravene 1 til 3.

DRAWINGS

Figure 1.




Figure 2A




Figure 2B




Figure 2C




Figure 3.




Figure 3C




Figure 3D




Figure 3E




Figure 4A.




Figure 4B





Figure 5A



Figure 5B

Figure 5C

Number at risk by mutation type:

CALR	186	115	63	27	9	3	0
JAK2	575	267	116	62	25	5	0
MPL	35	21	13	8	4	2	0

Figure 6A




Figure 6B





Figure 6C

Figure 6D




Figure 7




Figure 8




Figure 9




Figure 10

