



(86) Date de dépôt PCT/PCT Filing Date: 2004/02/10
(87) Date publication PCT/PCT Publication Date: 2004/09/02
(85) Entrée phase nationale/National Entry: 2005/08/10
(86) N° demande PCT/PCT Application No.: US 2004/003926
(87) N° publication PCT/PCT Publication No.: 2004/074243
(30) Priorité/Priority: 2003/02/14 (60/447,613) US

(51) Cl.Int.⁷/Int.Cl.⁷ C07D 401/04, A61K 31/53, A61P 25/00,
C07D 403/04


(71) Demandeur/Applicant:
WYETH, US

(72) Inventeurs/Inventors:
BERNOTAS, RONALD CHARLES, US;
YAN, YINFA, US;
ROBICHAUD, ALBERT JEAN, US;
LIU, GUANGCHENG, US

(74) Agent: TORYS LLP

(54) Titre : HETEROCYCLYL-3-SULFONYLINDAZOLES UTILISES COMME LIGANDS DE LA 5-HYDROXYTRYPTAMINE-6

(54) Title: HETEROCYCLYL-3-SULFONYLINDAZOLES AS 5-HYDROXYTRYPTAMINE-6 LIGANDS



(57) Abrégé/Abstract:

The invention provides compounds of formula (I) and the use thereof in the therapeutic treatment of disorders related to or affected by the 5-HT6 receptor.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
2 September 2004 (02.09.2004)

PCT

(10) International Publication Number
WO 2004/074243 A3

(51) International Patent Classification⁷: **C07D 401/04**,
403/04, A61K 31/53, A61P 25/00

(21) International Application Number:
PCT/US2004/003926

(22) International Filing Date: 10 February 2004 (10.02.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/447,613 14 February 2003 (14.02.2003) US

(71) Applicant (for all designated States except US): **WYETH**
[US/US]; Five Giralda Farms, Madison, NJ 07940 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BERNOTAS, Ronald, Charles** [US/US]; 908 Morgan Drive, Royersford, PA 19468 (US). **YAN, Yinfu** [CN/US]; 362 Wren Lane, Bedminster, NJ 07921 (US). **ROBICHAUD, Albert, Jean** [US/US]; 3 Riverview Court, Ringoes, NJ 08551 (US). **LIU, Guangcheng** [CN/US]; 7 Greenfield Drive, North, Princeton Junction, NJ 08550 (US).

(74) Agent: **LENCES, Barbara, L.**; Wyeth, Patent Law Department, Five Giralda Farms, Madison, NJ 07940 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

(88) Date of publication of the international search report:
2 December 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HETEROCYCLYL-3-SULFONYLINDAZOLES AS 5-HYDROXYTRYPTAMINE-6 LIGANDS



(I)

(57) Abstract: The invention provides compounds of formula (I) and the use thereof in the therapeutic treatment of disorders related to or affected by the 5-HT₆ receptor.

WO 2004/074243 A3

**HETEROCYCLYL-3-SULFONYLINDAZOLES AS 5-HYDROXYTRYPTAMINE-6
LIGANDS**

5

BACKGROUND OF THE INVENTION

Serotonin (5-Hydroxytryptamine)(5-HT) receptors play a critical role in many physiological and behavioral functions in humans and animals. These functions are mediated through various 5-HT receptors distributed throughout the body. There are now approximately fifteen different human 5-HT receptor subtypes that have been cloned, many with well-defined roles in humans. One of the most recently identified 5-HT receptor subtypes is the 5-HT₆ receptor, first cloned from rat tissue in 1993 (Monsma, F. J.; Shen, Y.; Ward, R. P.; Hamblin, M. W. *Molecular Pharmacology* 1993, 43, 320-327) and subsequently from human tissue (Kohen, R.; Metcalf, M. A.; Khan, N.; Druck, T.; Huebner, K.; Sibley, D. R. *Journal of Neurochemistry* 1996, 66, 47-56). The receptor is a G-protein coupled receptor (GPCR) positively coupled to adenylate cyclase (Ruat, M.; Traiffort, E.; Arrang, J-M.; Tardivel-Lacombe, L.; Diaz, L.; Leurs, R.; Schwartz, J-C. *Biochemical Biophysical Research Communications* 1993, 193, 268-276). The receptor is found almost exclusively in the central nervous system (CNS) areas both in rat and in human. *In situ* hybridization studies of the 5-HT₆ receptor in rat brain using mRNA indicate principal localization in the areas of 5-HT projection including striatum, nucleus accumbens, olfactory tubercle, and hippocampal formation (Ward, R. P.; Hamblin, M. W.; Lachowicz, J. E.; Hoffman, B. J.; Sibley, D. R.; Dorsa, D. M. *Neuroscience* 1995, 64, 1105-1111).

There are many potential therapeutic uses for 5-HT₆ ligands in humans based on direct effects and on indications from available scientific studies. These studies include the localization of the receptor, the affinity of ligands with known *in vivo* activity, and various animal studies conducted so far.

One potential therapeutic use of modulators of 5-HT₆ receptor function is in the enhancement of cognition and memory in human diseases such as Alzheimer's. The high levels of receptor found in important structures in the forebrain, including the caudate/putamen, hippocampus, nucleus accumbens, and cortex

suggest a role for the receptor in memory and cognition since these areas are known to play a vital role in memory (Gerard, C.; Martres, M.-P.; Lefevre, K.; Miquel, M.C.; Verge, D.; Lanfumey, R.; Doucet, E.; Hamon, M.; El Mestikawy, S. *Brain Research*, 1997, 746, 207-219). The ability of known 5-HT6 receptor ligands to enhance
5 cholinergic transmission also supported the potential cognition use (Bentley, J. C.; Boursson, A.; Boess, F. G.; Kone, F. C.; Marsden, C. A.; Petit, N.; Sleight, A. J. *British Journal of Pharmacology*, 1999, 126(7), 1537-1542). Studies have found that a known 5-HT6 selective antagonist significantly increased glutamate and aspartate levels in the frontal cortex without elevating levels of noradrenaline, dopamine, or 5-
10 HT. This selective elevation of neurochemicals known to be involved in memory and cognition strongly suggests a role for 5-HT6 ligands in cognition (Dawson, L. A.; Nguyen, H. Q.; Li, P. *British Journal of Pharmacology*, 2000, 130(1), 23-26). Animal studies of memory and learning with a known selective 5-HT6 antagonist found some positive effects (Rogers, D. C.; Hatcher, P. D.; Hagan, J. J. *Society of Neuroscience, Abstracts* 2000, 26, 680). Further support for the role of a selective 5-HT6 ligand in
15 cognition can be found in Woolley, M. L.; Marsden, C. A.; Sleight, A. J.; and Fone, K. C. F., *Psychopharmacology*, 2003, 170(4), 358-367.

A related potential therapeutic use for 5-HT6 ligands is the treatment of attention deficit disorders (ADD, also known as Attention Deficit Hyperactivity
20 Disorder or ADHD) in both children and adults. Because 5-HT6 antagonists appear to enhance the activity of the nigrostriatal dopamine pathway and because ADHD has been linked to abnormalities in the caudate (Ernst, M; Zametkin, A. J.; Matochik, J. H.; Jons, P. A.; Cohen, R. M. *Journal of Neuroscience* 1998, 18(15), 5901-5907), 5-HT6 antagonists may attenuate attention deficit disorders.

25 Early studies examining the affinity of various CNS ligands with known therapeutic utility or a strong structural resemblance to known drugs suggests a role for 5-HT6 ligands in the treatment of schizophrenia and depression. For example, clozapine (an effective clinical antipsychotic) has high affinity for the 5-HT6 receptor subtype. Also, several clinical antidepressants have high affinity for the receptor as
30 well and act as antagonists at this site (Branchek, T. A.; Blackburn, T. P. *Annual Reviews in Pharmacology and Toxicology* 2000, 40, 319-334).

Further, recent *in vivo* studies in rats indicate 5-HT6 modulators may be useful in the treatment of movement disorders including epilepsy (Stean, T.;

Routledge, C.; Upton, N. *British Journal of Pharmacology* 1999, 127 Proc. Supplement 131P and Routledge, C.; Bromidge, S. M.; Moss, S. F.; Price, G. W.; Hirst, W.; Newman, H.; Riley, G.; Gager, T.; Stean, T.; Upton, N.; Clarke, S. E.; Brown, A. M. *British Journal of Pharmacology* 2000, 130(7), 1606-1612).

5 Taken together, the above studies strongly suggest that compounds which are 5-HT6 receptor modulators, i.e. ligands, may be useful for therapeutic indications including: the treatment of diseases associated with a deficit in memory, cognition, and learning such as Alzheimer's and attention deficit disorder; the treatment of personality disorders such as schizophrenia; the treatment of behavioral disorders,
10 e.g., anxiety, depression and obsessive compulsive disorders; the treatment of motion or motor disorders such as Parkinson's disease and epilepsy; the treatment of diseases associated with neurodegeneration such as stroke and head trauma; or withdrawal from drug addiction including addiction to nicotine, alcohol, and other substances of abuse.

15 Therefore, it is an object of this invention to provide compounds which are useful as therapeutic agents in the treatment of a variety of central nervous system disorders related to or affected by the 5-HT6 receptor.


It is another object of this invention to provide therapeutic methods and pharmaceutical compositions useful for the treatment of central nervous system
20 disorders related to or affected by the 5-HT6 receptor.

It is a feature of this invention that the compounds provided may also be used to further study and elucidate the 5-HT6 receptor.

SUMMARY OF THE INVENTION

25

The present invention provides a compound of formula I



wherein

5

A is C, CR₈ or N;

R₁ is H, halogen, CN, COR₉, OCO₂R₁₀, CO₂R₁₁, CONR₁₂R₁₃, SO_xR₁₄,

NR₁₅R₁₆, OR₁₇ or a C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl or heteroaryl group each optionally substituted;

10

R₂ is an optionally substituted C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl, or heteroaryl group or an optionally substituted 8- to 13-membered bicyclic or tricyclic ring system having a N atom at the bridgehead and optionally containing 1, 2 or 3 additional heteroatoms selected from N, O or S ;

R₃ is H or a C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl or heteroaryl group each optionally substituted;

15

R₄ is H or a C₁-C₆alkyl or C₃-C₇cycloalkyl group each optionally substituted;

R₅, R₆ and R₇ are each independently H or a C₁-C₆alkyl or C₃-C₇cycloalkyl group each optionally substituted;

m and p are each independently an integer of 1, 2 or 3;

n is an integer of 1 or 2;

20

R₈ is H, OH or an optionally substituted C₁-C₆alkoxy group;

R₉, R₁₀, R₁₁ and R₁₇ are each independently H or a C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₃-C₆cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;

25

R₁₂, R₁₃, R₁₅ and R₁₆ are each independently H or an optionally substituted C₁-C₄alkyl group or R₁₂ and R₁₃ or R₁₅ and R₁₆ may be taken together with the atom to which they are attached to form a 5- to 7-membered

ring optionally containing another heteroatom selected from O, NR₁₈
 or SO_x;
 R₁₄ is a C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₃-C₆cycloalkyl,
 cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;
 5 R₁₈ is H or a C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₃-C₇cycloalkyl,
 cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;
 x is 0 or an integer of 1 or 2; and

--- represents a single bond or a double bond; or

the stereoisomers thereof or the pharmaceutically acceptable salts thereof.


10 The present invention also provides methods and compositions useful in the
 treatment of central nervous system disorders.

DETAILED DESCRIPTION OF THE INVENTION

15

The 5-hydroxytryptamine-6 (5-HT₆) receptor is one of the most recent
 receptors to be identified by molecular cloning. Its ability to bind a wide range of
 therapeutic compounds used in psychiatry, coupled with its intriguing distribution in
 the brain has stimulated significant interest in new compounds which are capable of
 20 interacting with or affecting said receptor. Significant efforts are being made to
 understand the possible role of the 5-HT₆ receptor in psychiatry, cognitive
 dysfunction, motor function and control, memory, mood and the like. To that end,
 compounds which demonstrate a binding affinity for the 5-HT₆ receptor are earnestly
 sought both as an aid in the study of the 5-HT₆ receptor and as potential therapeutic
 25 agents in the treatment of central nervous system disorders, for example see C.
 Reavill and D. C. Rogers, Current Opinion in Investigational Drugs, 2001, 2(1):104-
 109, Pharma Press Ltd.

Surprisingly, it has now been found that heterocyclyl-3-sulfonylindazoles of
 formula I demonstrate 5-HT₆ affinity along with significant sub-type selectivity.
 30 Advantageously, said formula I indazoles are effective therapeutic agents for the
 treatment of central nervous system (CNS) disorders associated with or affected by
 the 5-HT₆ receptor. Accordingly, the present invention provides 1-heterocyclyl-3-
 sulfonylindazole compounds of formula I



wherein

5 A is C, CR₈ or N;

R₁ is H, halogen, CN, COR₉, OCO₂R₁₀, CO₂R₁₁, CONR₁₂R₁₃, SO_xR₁₄,
NR₁₅R₁₆, OR₁₇ or a C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl or heteroaryl
group each optionally substituted;

10 R₂ is an optionally substituted C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl, or heteroaryl
group or an optionally substituted 8- to 13-membered bicyclic or
tricyclic ring system having a N atom at the bridgehead and optionally
containing 1, 2 or 3 additional heteroatoms selected from N, O or S;

R₃ is H or a C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl or heteroaryl group each
optionally substituted;

15 R₄ is H or a C₁-C₆alkyl or C₃-C₇cycloalkyl group each optionally substituted;

R₅, R₆ and R₇ are each independently H or a C₁-C₆alkyl or C₃-C₇cycloalkyl
group each optionally substituted;

m and p are each independently an integer of 1, 2 or 3;

n is an integer of 1 or 2;

20 R₈ is H, OH or an optionally substituted C₁-C₆alkoxy group;

R₉, R₁₀, R₁₁ and R₁₇ are each independently H or a C₁-C₆alkyl, C₂-C₆alkenyl,
C₂-C₆alkynyl, C₃-C₆cycloalkyl, cycloheteroalkyl, aryl or heteroaryl
group each optionally substituted;

25 R₁₂, R₁₃, R₁₅ and R₁₆ are each independently H or an optionally substituted
C₁-C₄alkyl group or R₁₂ and R₁₃ or R₁₅ and R₁₆ may be taken together
with the atom to which they are attached to form a 5- to 7-membered

ring optionally containing another heteroatom selected from O, NR₁₈ or SO_x;

R₁₄ is a C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₃-C₆cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;


5 R₁₈ is H or a C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₃-C₇cycloalkyl, cycloheteroalkyl, aryl or heteraryl group each optionally substituted;

x is 0 or an integer of 1 or 2; and

--- represents a single bond or a double bond; or


the stereoisomes thereof or the pharmaceutically acceptable salts thereof.

10 As used in the specification and claims, the term halogen designates F, Cl, Br or I and the term cycloheteroalkyl designates a five- to seven-membered cycloalkyl ring system containing 1 or 2 heteroatoms, which may be the same or different, selected from N, O or S and optionally containing one double bond. Exemplary of the cycloheteroalkyl ring systems included in the term as designated herein are the
15 following rings wherein X is NR, O or S; and R is H or an optional substituent as described hereinbelow:



Similarly, as used in the specification and claims, the term heteroaryl
20 designates a five- to ten-membered aromatic ring system containing 1, 2 or 3 heteroatoms, which may be the same or different, selected from N, O or S. Such heteroaryl ring systems include pyrrolyl, azolyl, oxazolyl, thiazolyl, imidazolyl, furyl, thienyl, quinoliny, isoquinoliny, indolyl, benzothienyl, benzofuranyl, benzisoxazolyl or the like. The term aryl designates a carbocyclic aromatic ring system such as
25 phenyl, naphthyl, anthracenyl or the like. The term haloalkyl as used herein designates a C_nH_{2n+1} group having from one to 2n+1 halogen atoms which may be the same or different and the term haloalkoxy as used herein designates an OC_nH_{2n+1} group having from one to 2n+1 halogen atoms which may be the same or different.

Exemplary of the 8- to 13-membered bicyclic or tricyclic ring systems having a N atom at the bridgehead and optionally containing 1, 2 or 3 additional heteroatoms selected from N, O or S included in the term as designated herein are the following ring systems wherein W_2 is NR, O or S; and R is H or an optional substituent as described hereinbelow:



In the specification and claims, when the terms C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_7 cycloalkyl, cycloheteroalkyl, aryl or heteroaryl are designated as being optionally substituted, the substituent groups which are optionally present may be one or more of those customarily employed in the development of pharmaceutical compounds, or the modification of such compounds, to influence their structure/activity, persistence, absorption, stability or other beneficial property. Specific examples of such substituents include halogen atoms, nitro, cyano, thiocyanato, cyanato, hydroxyl, alkyl, haloalkyl, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, formyl, alkoxycarbonyl, carboxyl, alkanoyl, alkylthio,

alkylsulphanyl, alkylsulphonyl, carbamoyl, alkylaminocarbonyl, phenyl and substituted phenyl (where the substituents are as defined herein), phenoxy, benzyl, benzyloxy, heteroaryl, indolyl, heterocyclyl (e.g., heteroaryl, cycloheteroalkyl) or cycloalkyl groups, preferably halogen atoms or lower alkyl or lower alkoxy groups. Typically, 0-
5 3 substituents may be present. When any of the foregoing substituents represents or contains an alkyl substituent group, this may be linear or branched and may contain up to 12, preferably up to 6, more preferably up to 4 carbon atoms.

Pharmaceutically acceptable salts may be any acid addition salt formed by a compound of formula I and a pharmaceutically acceptable acid such as phosphoric,
10 sulfuric, hydrochloric, hydrobromic, citric, maleic, malonic, mandelic, succinic, fumaric, acetic, lactic, nitric, sulfonic, p-toluene sulfonic, methane sulfonic acid or the like.

Compounds of the invention include esters, carbamates or other conventional prodrug forms, which in general, are functional derivatives of the compounds of the
15 invention and which are readily converted to the inventive active moiety *in vivo*. Correspondingly, the method of the invention embraces the treatment of the various conditions described hereinabove with a compound of formula I or with a compound which is not specifically disclosed but which, upon administration, converts to a compound of formula I *in vivo*. Also included are metabolites of the compounds of
20 the present invention defined as active species produced upon introduction of these compounds into a biological system.

Compounds of the invention may exist as one or more stereoisomers. The various stereoisomers include enantiomers, diastereomers, atropisomers and geometric isomers. One skilled in the art will appreciate that one stereoisomer may
25 be more active or may exhibit beneficial effects when enriched relative to the other stereoisomer(s) or when separated from the other stereoisomer(s). Additionally, the skilled artisan knows how to separate, enrich or selectively prepare said stereoisomers. Accordingly, the present invention comprises compounds of formula I, the stereoisomers thereof and the pharmaceutically acceptable salts thereof. The
30 compounds of the invention may be present as a mixture of stereoisomers, individual stereoisomers, or as an optically active or enantiomerically pure form.

Examples of R_3 are hydrogen, phenyl and C_1 - C_4 alkyl which may be substituted by optionally substituted phenyl, e.g., R_3 is methyl, phenyl or 3-chlorobenzyl. An example of A is N. Examples of R_2 are phenyl, 1-naphthyl, 2-naphthyl, cyanophenyl, 2, 3, or 4-halophenyl, 2-phenethyl, 2-chloro-4-fluorophenyl, aminophenyl and thienyl. Examples of R_4 are H, methyl, ethyl, propyl and phenethyl.

Preferred compounds of the invention are those compounds of formula I wherein n is 1. Another group of preferred compounds are those compounds of formula I wherein R_4 is H or an optionally substituted C_1 - C_4 alkyl group. A further group of preferred compounds of the invention are those compounds of formula I wherein R_2 is an optionally substituted phenyl, naphthyl or heteroaryl group. Also preferred are those compounds of formula I wherein R_5 , R_6 and R_7 are H.

More preferred compounds of the invention are those compounds of formula I wherein A is N and R_4 is H or an optionally substituted C_1 - C_4 alkyl group. Another group of more preferred compounds are those compounds of formula I wherein A is N; n is 1 and R_2 is an optionally substituted phenyl, naphthyl or heteroaryl group. Further more preferred compounds of the invention are those compounds of formula I wherein the 6- or 7-membered azacyclic ring is attached to the indazole in the 5 or 7 position.

Among the preferred compounds of the invention are:


- 20 5-(4-methylpiperazin-1-yl)-3-(1-naphthylsulfonyl)-1H-indazole;
- 3-(1-naphthylsulfonyl)-5-piperazin-1-yl-1H-indazole;
- 1-(3-chlorobenzyl)-5-(4-methylpiperazin-1-yl)-3-(1-naphthylsulfonyl)-1H-indazole;
- 7-(4-methylpiperazin-1-yl)-3-(phenylsulfonyl)-1H-indazole;
- 3-(phenylsulfonyl)-7-(4-propylpiperazin-1-yl)-1H-indazole;
- 25 3-(phenylsulfonyl)-7-piperazin-1-yl-1H-indazole;
- 5-(4-methylpiperazin-1-yl)-3-(phenylsulfonyl)-1H-indazole;
- 3-(phenylsulfonyl)-5-(4-propylpiperazin-1-yl)-1H-indazole;
- 3-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole;
- 5-piperazin-1-yl-3-[(3-cyanophenyl)sulfonyl]-1H-indazole;
- 30 7-(4-methylpiperazin-1-yl)-3-[(2-phenethyl)sulfonyl]-1H-indazole;
- 3-(1-naphthylsulfonyl)-7-(4-propylpiperazin-1-yl)-1H-indazole;
- 5-(4-phenethylpiperazin-1-yl)-3-(1-naphthylsulfonyl)-1H-indazole;
- 5-(4-methylpiperazin-1-yl)-3-(2-naphthylsulfonyl)-1H-indazole;

- 3-[(2-chloro-4-fluorophenyl)sulfonyl]-5-piperazin-1-yl-1*H*-indazole;
 1-methyl-3-(phenylsulfonyl)-5-piperazin-1-yl-1*H*-indazole;
 1-phenyl-3-(phenylsulfonyl)-5-piperazin-1-yl-1*H*-indazole;
 1-methyl-3-(phenylsulfonyl)-7-piperazin-1-yl-1*H*-indazole;
 5 1-phenyl-3-(phenylsulfonyl)-7-(piperazin-1-yl)-1*H*-indazole;
 7-piperazin-1-yl-3-[(3-fluorophenyl)sulfonyl]-1*H*-indazole;
 3-[(4-fluorophenyl)sulfonyl]-7-(4-methylpiperazin-1-yl)-1*H*-indazole;
 3-[(2-chlorophenyl)sulfonyl]-7-piperazin-1-yl-1*H*-indazole;
 3-[(4-aminophenyl)sulfonyl]-7-(piperazin-1-yl)-1*H*-indazole;
 10 5-piperazin-1-yl-3-[(3-fluorophenyl)sulfonyl]-1*H*-indazole;
 3-[(4-fluorophenyl)sulfonyl]-5-(4-methylpiperazin-1-yl)-1*H*-indazole;
 3-[(2-chlorophenyl)sulfonyl]-5-(4-propylpiperazin-1-yl)-1*H*-indazole;
 3-[(4-aminophenyl)sulfonyl]-5-piperazin-1-yl-1*H*-indazole;
 3-[(5-chlorothien-2-yl)sulfonyl]-5-piperazin-1-yl-1*H*-indazole;
 15 3-[(5-chlorothien-2-yl)sulfonyl]-7-piperazin-1-yl-1*H*-indazole;
 4-chloro-3-(phenylsulfonyl)-7-piperazin-1-yl-1*H*-indazole;
 5-fluoro-3-(phenylsulfonyl)-7-piperazin-1-yl-1*H*-indazole;
 6-fluoro-3-(phenylsulfonyl)-7-piperazin-1-yl-1*H*-indazole;
 4-chloro-3-(phenylsulfonyl)-5-piperazin-1-yl-1*H*-indazole;
 20 7-fluoro-3-(phenylsulfonyl)-5-piperazin-1-yl-1*H*-indazole;
 6-fluoro-3-(phenylsulfonyl)-5-piperazin-1-yl-1*H*-indazole;
 6-(4-phenethylpiperazin-1-yl)-3-(phenylsulfonyl)-1*H*-indazole;
 6-(4-methylpiperazin-1-yl)-3-(phenylsulfonyl)-1*H*-indazole;
 6-(4-propylpiperazin-1-yl)-3-(phenylsulfonyl)-1*H*-indazole;
 25 3-(phenylsulfonyl)-6-piperazin-1-yl-1*H*-indazole;
 4-piperazin-1-yl-3-(phenylsulfonyl)-1*H*-indazole;
 4-(4-methylpiperazin-1-yl)-3-(phenylsulfonyl)-1*H*-indazole;
 4-(4-propylpiperazin-1-yl)-3-(phenylsulfonyl)-1*H*-indazole;
 3-(phenylsulfonyl)-4-piperazin-1-yl-1*H*-indazole;
 30 3-[(5-chlorothien-2-yl)sulfonyl]-5-piperidin-1-yl-1*H*-indazole;
 3-[(5-chlorothien-2-yl)sulfonyl]-7-piperidin-1-yl-1*H*-indazole;
 1-methyl-3-(phenylsulfonyl)-5-piperidin-1-yl-1*H*-indazole;
 3-[(3-fluorophenyl)sulfonyl]-5-piperidin-1-yl-1*H*-indazole;

- 1-methyl-3-(phenylsulfonyl)-7-(4-methylpiperidin-1-yl)-1*H*-indazole;
 3-[(3-fluorophenyl)sulfonyl]-7-piperidin-1-yl-1*H*-indazole;
 3-[(3-fluorophenyl)sulfonyl]-5-piperidin-1-yl-1*H*-indazole;
 3-(phenylsulfonyl)-5-piperidin-1-yl-1*H*-indazole;
 5 1-methyl-3-[(3-fluorophenyl)sulfonyl]-7-piperidin-1-yl-1*H*-indazole;
 3-(phenylsulfonyl)-7-piperidin-1-yl-1*H*-indazole;
 the stereoisomers thereof; or
 the pharmaceutically acceptable salts thereof.


10 This invention also provides processes for preparing compounds of formula (I) or salts thereof, which processes include one of the following

- (a) diazotising and cyclising a compound of formula (VIIA):



wherein the dotted line, m, n, p, A, R₁, R₂, R₄, R₅, R₆ and R₇ are as defined herein,
 15 e.g. with NaNO₂/acid to give a compound of formula (I) wherein R₃ is hydrogen,
 or

- (b) alkylating or arylating a protected compound of formula (IA)




wherein the dotted line, m, n, p, A, R₁, R₂, R₅, R₆ and R₇ are as defined herein and P is a protecting group or R₄ where R₄ is as defined herein excepting hydrogen, with an alkylating or arylating agent of formula R₃-LG where LG is a leaving group and R₃ is a C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl or heteroaryl group each optionally substituted;
 5 and if required removing the protecting group, to give a corresponding compound of formula (I) where R₃ is a C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl or heteroaryl group each optionally substituted;

or

(c) alkylating a compound of formula (IB)


10



wherein the dotted line, m, n, p, A, R₁, R₂, R₃, R₅, R₆ and R₇ are as defined herein, with an alkylating agent of formula R₄-LG where LG is a leaving group and R₄ is a C₁-C₆alkyl or C₃-C₇cycloalkyl group each optionally substituted, to give a corresponding compound of formula (I) where R₄ is a C₁-C₆alkyl or C₃-C₇cycloalkyl group each optionally substituted;


or

(d) reacting a compound of formula (XII):



20


wherein m, R₁, R₂ and R₃ are as defined herein, and LG' is a leaving group such as Cl, Br, I or an activated hydroxyl group such as triflate, with a compound of formula (V):



(V)

wherein n , p , R_4 , R_5 , R_6 and R_7 are as defined herein, in the presence of a catalyst, e.g., a palladium or nickel catalyst, to give a corresponding compound of formula (I); or

5 (e) reducing a compound of formula (XII):




(XVI)

wherein m , n , p , R_1 , R_2 , R_5 , R_6 and R_7 are as defined herein, P is a protecting group or R_4 as defined in claim 1 and P' is a protecting group or R_3 as defined herein, if required removing any protecting group, to give a corresponding compound of formula (I) wherein A is CH ;

or

15 (f) acidifying a compound of formula (XII):



(XV)

wherein m , n , p , R_1 , R_2 , R_5 , R_6 and R_7 are as defined herein, P is a protecting group or R_4 as defined herein and P' is a protecting group or R_3 as defined herein, if

required removing any protecting group, to give a corresponding compound of formula (I) wherein A is C and ---- represents a double bond;


or

- 5 (g) converting a basic compound of formula (I) as defined herein to an acid addition salt thereof or vice versa.

or

- (h) separating a stereoisomer of a compound of formula (I) from a mixture thereof.

- 10 In particular compounds of the invention may be conveniently prepared using conventional synthetic methods and, if required, standard separation or isolation techniques. For example, compounds of formula I wherein A is N; R₃ is H; and ---- represents a single bond (Ia) may be prepared by reacting a halonitrobenzene compound of formula II with a chloromethylsulfone of formula III in the presence of
15 strong base, such as KO-t-Bu or KOH, to give the benzylsulfonyl compound of formula IV; reacting said formula IV compound with a piperazine or homopiperazine of formula V in the presence of a base such as K₂CO₃ to give the compound of formula VI; reacting said formula VI compound with a reducing agent such as Sn, Fe or Zn in the presence of an acid to give the corresponding amine of formula VII; and
20 reacting said amine with NaNO₂ in the presence of an acid to give the desired compound of formula Ia. The reaction is illustrated in flow diagram I wherein Hal represents Cl or F.



5

Compounds of formula Ia wherein R_3 is other than H may be prepared using conventional alkylation/deprotection procedures. For example, compounds of formula I wherein A is N; R_3 is other than H; and --- represents a single bond (Ib) may be prepared by reacting a protected compound of formula VIII with an alkylating agent of formula IX in the presence of a base and a solvent optionally in the presence of a phase-transfer agent to give the protected alkylated compound of formula X and deprotecting said formula X compound to give the desired compound


10

of formula Ib wherein R_4 is H. Alternatively, for those compounds wherein R_3 is an aryl or heteroaryl group, the agent of formula IX may be an aryl or heteroaryl boronic acid and may be coupled with the formula VIII indazole in the presence of a catalyst, e.g., $\text{Cu}(\text{OCOCH}_3)_2$ to give the corresponding protected compound of formula X

5 wherein R_3 is an aryl or heteroaryl group. Optionally the formula Ib compound may be reacted with an alkylating agent of formula XI under standard alkylation conditions to give the compound of formula Ib wherein R_4 is other than H. If desired, the sequence may be reversed by deprotecting the formula VIII compound to give the compound of formula Ia wherein R_3 and R_4 are H (Ic) and alkylating the formula Ic

10 compound with the formula XI alkylating agent to give the compound of formula Id. Optionally, the formula Id compound may be reacted with a formula IX alkylating agent, as described hereinabove, to give the compound of formula Ib. The reactions are shown in flow diagram II wherein P is a protecting group and LG is a leaving group such as Cl, Br, I, OH, $\text{B}(\text{OH})_2$, tosyl, mesyl or the like.


15

FLOW DIAGRAM II


5

Alternatively, compounds of formula Ib may be prepared directly from an indazole compound of formula XII by coupling said formula XII compound with the appropriate piperazine or homopiperazine compound of formula V in the presence of a catalyst such as a palladium or nickel catalyst. The reaction is shown in flow diagram III wherein LG' is a leaving group such as Cl, Br, I or an activated hydroxyl group such as CF₃SO₃ (triflate).

10


FLOW DIAGRAM III

- 5 Compounds of formula VIII may also be prepared by reacting a protected 3-iodoindazole of formula XIII with a sodium sulfinate of formula XIV in the presence of copper iodide and dimethyl formamide (DMF) to give the desired compound of formula VIII or by reacting the formula XIII compound with a thiol of formula XV in the presence of copper iodide and a base, such as K₂CO₃, to give the thio ether of
- 10 formula XVI and oxidizing said formula XVI compound to give the desired compound of formula VIII. The reactions are shown in flow diagram IV wherein P represents a protecting group.

FLOW DIAGRAM IV


Compounds of formula XIII wherein R_1 is H (XIIIa) may be prepared by reacting a dihalobenzaldehyde of formula XVII with the piperazine or homopiperazine compound of formula V wherein R_4 is a protecting group (Va) to give the ortho halobenzaldehyde compound of formula XVIII, reacting the formula XVIII compound with hydrazine hydrate to give the indazole of formula XIX and reacting the formula XIX indazole with iodine in the presence of a base, such as KOH, to give the desired formula XIIIa compounds. The reactions are shown in flow diagram V wherein P is a protecting group and Hal is F, Cl, Br, or I, preferably F.

10

FLOW DIAGRAM V

Compounds of formula XIII wherein n is 1 and R_5 , R_6 and R_7 are H (XIIIb) may be prepared by reacting an indazole amine of formula XX with di(2-chloroethyl)amine hydrochloride to give the corresponding piperazinyl indazole of formula XXI, protecting said formula XXI compound and reacting the protected formula XXI compound with iodine in the presence of a base, such as KOH, to give the desired


compound of formula XIIIb. The reactions are shown in flow diagram VI whereing P represents a protecting group.



5


Compounds of XIIIa and XIIIb may be converted to compounds of formula Ia, Ib, or Ic as shown hereinabove in flow diagrams II and III.

Corresponding compounds of the invention wherein A is CR₈ may be obtained, for example, by lithiating a protected bromoindazole of formula XXII, and reacting the lithiated compound *in situ* with an N-protected azacyclic ketone of formula XXIII to give the protected hydroxy compound of formula XXIV, said hydroxy compound may then be dehydrated to give the protected compound of formula XXV. Catalytic hydrogenation and subsequent deprotection of said formula XXV compound gives the desired compounds of formula I wherein A is CR₈; R₃ and R₄ are H; and --- represents a single bond (Id). The reaction sequence is shown in flow diagram VII wherein P and P' each independently represent a protecting group.

FLOW DIAGRAM VII

5 Using the procedures shown in flow diagram II hereinabove, the compounds of formulas XXIV, XXV and Id may be readily converted to compounds of formula I wherein A is CR₈ and R₃ or R₄ are other than H.

Alternatively, compounds of formula I wherein A is C or CR₈ and R₃ is H (Ie) may be prepared by reacting a chloromethylsulfonyl compound of formula III with a
 10 nitrobenzene derivative of formula XXVI in the presence of a strong base, such as KO-*t*-Bu or KOH, to provide the compound of formula XXVII; reducing said formula XVII compound with a reducing agent such as Sn, Fe or Zn in the presence of an acid to give the corresponding amine of formula XXVIII; and reacting said amine with NaNO₂ in the presence of an acid to give the desired indazole compound of formula
 15 Ie. The reactions are shown in flow diagram VIII.

FLOW DIAGRAM VIII

5 Compounds of formula Ie may be converted to the corresponding compounds of formula I wherein A is C or CR₈ and R₃ is other than H by reacting said formula Ie compound with an alkylating agent as shown hereinabove in flow diagram II.

Protecting groups suitable for use in the reactions shown hereinabove include t-butyloxycarbonyl, benzyl, acetyl, benzyloxycarbonyl, or any conventional group
 10 known to protect a basic nitrogen in standard synthetic procedures.

Advantageously, the formula I compounds of the invention are useful for the treatment of CNS disorders relating or affected by 5-HT₆ receptor including motor, mood, personality, behavioral, psychiatric, cognitive, neurodegenerative, or the like disorders, for example Alzheimer's disease, Parkinson's disease, attention deficit
 15 disorder, anxiety, epilepsy, depression, obsessive compulsive disorder, sleep disorders, neurodegenerative disorders (such as head trauma or stroke), feeding disorders (such as anorexia or bulimia), schizophrenia, memory loss, disorders

associated with withdrawal from drug or nicotine abuse, or the like or certain gastrointestinal disorders such as irritable bowel syndrome. Accordingly, the present invention provides a method for the treatment of a disorder of the central nervous system related to or affected by the 5-HT₆ receptor in a patient in need thereof which comprises providing said patient a therapeutically effective amount of a compound of formula I as described hereinabove. The compounds may be provided by oral or parenteral administration or in any common manner known to be an effective administration of a therapeutic agent to a patient in need thereof.

The term "providing" as used herein with respect to providing a compound or substance embraced by the invention, designates either directly administering such a compound or substance, or administering a prodrug, derivative or analog which forms an equivalent amount of the compound or substance within the body.

The therapeutically effective amount provided in the treatment of a specific CNS disorder may vary according to the specific condition(s) being treated, the size, age and response pattern of the patient, the severity of the disorder, the judgment of the attending physician or the like. In general, effective amounts for daily oral administration may be about 0.01 to 1,000 mg/kg, preferably about 0.5 to 500 mg/kg and effective amounts for parenteral administration may be about 0.1 to 100 mg/kg, preferably about 0.5 to 50 mg/kg.

In actual practice, the compounds of the invention are provided by administering the compound or a precursor thereof in a solid or liquid form, either neat or in combination with one or more conventional pharmaceutical carriers or excipients. Accordingly, the present invention provides a pharmaceutical composition which comprises a pharmaceutically acceptable carrier and an effective amount of a compound of formula I as described hereinabove.


Solid carriers suitable for use in the composition of the invention include one or more substances which may also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aides, binders, tablet-disintegrating agents or encapsulating materials. In powders, the carrier may be a finely divided solid which is in admixture with a finely divided compound of formula I. In tablets, the formula I compound may be mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. Said powders and tablets may contain up to 99% by weight of the formula I compound.

Solid carriers suitable for use in the composition of the invention include calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

- 5 Any pharmaceutically acceptable liquid carrier suitable for preparing solutions, suspensions, emulsions, syrups and elixirs may be employed in the composition of the invention. Compounds of formula I may be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, or a pharmaceutically acceptable oil or fat, or a mixture thereof. Said liquid
- 10 composition may contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, coloring agents, viscosity regulators, stabilizers, osmo-regulators, or the like. Examples of liquid carriers suitable for oral and parenteral administration include water (particularly containing additives as
- 15 above, e.g., cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g., glycols) or their derivatives, or oils (e.g., fractionated coconut oil and arachis oil). For parenteral administration the carrier may also be an oily ester such as ethyl oleate or isopropyl myristate.
- 20 Compositions of the invention which are sterile solutions or suspensions are suitable for intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions may also be administered intravenously. Inventive compositions suitable for oral administration may be in either liquid or solid composition form.

25 For a more clear understanding, and in order to illustrate the invention more clearly, specific examples thereof are set forth hereinbelow. The following examples are merely illustrative and are not to be understood as limiting the scope and underlying principles of the invention in any way.


30 Unless otherwise stated, all parts are parts by weight. The term NMR designates proton nuclear magnetic resonance. The terms THF, DMF and DMSO designate tetrahydrofuran, dimethyl formamide and dimethylsulfoxide, respectively. In the chemical drawings, the term Ph represents a phenyl group.

EXAMPLE 1**Preparation of 1-Benzyl-4-(4-nitrophenyl)piperazine**

5

A stirred solution of 1-benzylpiperazine (8.81 g, 50.0 mmol), 4-fluoronitrobenzene (5.31 mL, 50.0 mmol), and K_2CO_3 (6.90 g, 50.0 mmol) in ethanol is heated at reflux temperature under nitrogen for 18 h, cooled, diluted with water, and extracted with CH_2Cl_2 . The combined extracts are dried over $MgSO_4$ and concentrated *in vacuo* to give a solid residue. The solid is triturated with 20:80 ethyl acetate:hexanes and filtered. The filtercake is air-dried to afford the title compound as orange crystals, 8.45 g (57% yield), mp 218-219°C, characterized by NMR and mass spectral analyses.

15

EXAMPLE 2**Preparation of 1-Benzyl-4-{4-nitro-3-[(phenylsulfonyl)methyl]phenyl}piperazine**

20

A stirred solution of 1-benzyl-4-(4-nitrophenyl)piperazine (5.95 g, 20.0 mmol) and chloromethylphenylsulfone (3.82 g, 20.0 mmol) in dry THF under nitrogen at -60°C is treated with 1.0M KO-t-Bu in THF (44.0 mL, 44.0 mmol), warmed to -20°C over a 1 h period, quenched with acetic acid and treated sequentially with water, saturated aqueous $NaHCO_3$ and ether. The phases are separated and the aqueous phase is extracted with ether. The combined ethers are washed with water and brine, dried over $MgSO_4$ and concentrated *in vacuo*. The resultant residue is

30


chromatographed (silica gel, 1:1 and 1:0 ethyl acetate:hexanes as eluent) to give the title compound as a yellow solid, 7.52 g (83% yield), mp 145-146°C, characterized by NMR and mass spectral analyses.

5

EXAMPLE 3

Preparation of 4-(4-Benzylpiperazin-1-yl)-2-[(phenylsulfonyl)methyl]aniline

10



15


A mixture of 1-benzyl-4-{4-nitro-3-[(phenylsulfonyl)methyl]phenyl}piperazine (6.77 g, 15.0 mmol) and granular tin (7.48 g, 63.0 mmol) in methanol and conc. hydrochloric acid is heated at 45°C for 4 h, stirred at ambient temperature for 18 h, carefully poured into saturated aqueous NaHCO₃, treated with ether and stirred for 0.5 h. The phases are separated and the aqueous phase is extracted sequentially with ether and CH₂Cl₂. The extracts and organic phase are combined, dried over MgSO₄ and concentrated *in vacuo*. The resultant residue is dissolved in CH₂Cl₂ and filtered through Celite[®]. The filtrate is concentrated *in vacuo* to afford the title compound as a pale yellow solid, 6.11 g (97% yield), mp 141-143°C, characterized by NMR and mass spectral analyses

25

EXAMPLE 4

Preparation of 5-(4-Benzylpiperazin-1-yl)-3-(phenylsulfonyl)-1H-indazole hydrochloride

30




A stirred solution of 4-(4-benzylpiperazin-1-yl)-2-[(phenylsulfonyl)methyl]-
 5 aniline (0.371 g, 0.88 mmol) in 4.0 M aqueous hydrochloric acid is cooled in an ice
 bath, treated dropwise with NaNO_2 (91 mg, 1.32 mmol) in water, stirred for 40 min,
 treated with 2.5 M aqueous NaOH to pH ~ 14 and filtered. The filtercake is dissolved
 in CH_2Cl_2 , and chromatographed (silica gel, ethyl acetate as eluent) to afford the free
 indazole of the title product as a yellow solid (276 mg, 64%). The solid is dissolved in
 10 a mixture of ethanol and CH_2Cl_2 , treated with 4.0 M HCl in dioxane, and concentrated
in vacuo to give a solid residue. The residue is triturated with ethyl acetate to afford
 the title product as an off-white solid, 278 mg (57% yield), mp $177-180^\circ\text{C}$, identified
 by NMR and mass spectral analyses.

15


EXAMPLE 5

Preparation of 5-(4-Benzylpiperazin-1-yl)-3-[(3-fluorophenyl)sulfonyl]-1H-indazole hydrochloride


20



Using essentially the same procedures described hereinabove for Examples
 25 2 - 4 and employing chloromethyl-(3-fluorophenyl)sulfone, the title compound is
 obtained as an off-white solid, mp $157-160^\circ\text{C}$, identified by NMR and mass spectral
 analyses.

EXAMPLE 6**5 Preparation of 1-Benzyl-4-(3-methoxy-4-nitrophenyl)piperazine**

A stirred solution of 1-benzylpiperazine (3.53 g, 20.0 mmol), 4-chloro-3-methoxy-nitrobenzene (3.75 g, 20.0 mmol), and K_2CO_3 (2.76 g, 20.0 mmol) in DMF is heated at 85° to 100°C under nitrogen for 23 h, cooled to room temperature, treated with 2M aqueous hydrochloric acid, diluted with water and extracted with ether. The combined extracts are washed with brine, dried over $MgSO_4$ and concentrated *in vacuo*. The resultant residue is chromatographed (silica gel, 30:70 ethyl acetate:hexanes and 50:50 ethyl acetate:hexanes as eluent) to afford the title compound as a yellow-orange oil 2.64 g (57% yield), identified by NMR analysis.

EXAMPLE 7**20 Preparation of 5-(4-Benzylpiperazin-1-yl)-6-methoxy-3-(phenylsulfonyl)-1H-indazole hydrochloride**

25

Using essentially the same procedures described hereinabove for Examples 2 - 4 and employing 1-benzyl-4-(2-methoxy-4-nitrophenyl)piperazine as starting


substrate, the title compound is obtained as a tan solid, 70 mg, mp 185-187°C, identified by NMR and mass spectral analyses.

5

EXAMPLE 8

Preparation of 5-(4-Benzylpiperazin-1-yl)-6-methoxy-1-phenyl-3-(phenylsulfonyl)-1H-indazole hydrochloride

10




A mixture of 5-(4-benzylpiperazin-1-yl)-6-methoxy-3-(phenylsulfonyl)-1H-indazole (448 mg, 0.97 mmol), phenylboronic acid (242 mg, 1.94 mmol), copper(II) acetate (271 mg, 1.50 mmol), pyridine (160 mg, 2.00 mmol) and 4Å molecular sieves (1.0 g) in methylene chloride is stirred at room temperature for 48 h and filtered. The filtercake is washed with THF. The filtrates are combined and concentrated *in vacuo*. The resultant residue is chromatographed (silica gel, 60:40 and 80:20 ethyl acetate:hexanes as eluents) to afford the free amine of the title product as a white solid, 130 mg (25% yield). The free amine is dissolved in ethanol, treated with 4.0 M HCl in dioxane, and concentrated *in vacuo* to a solid residue. This solid is triturated with ethyl acetate to afford the title product as a white solid, mp: 210-212°C, identified by NMR and mass spectral analyses.


25

EXAMPLE 9**Preparation of 6-Methoxy-1-phenyl 3-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole hydrochloride**

5



A mixture of 5-(4-benzylpiperazin-1-yl)-6-methoxy-1-phenyl-3-(phenylsulfonyl)-1H-indazole (190 mg, 0.353 mmol) and 1-chloroethylchloroformate (250 mg, 1.77 mmol) in 1,2-dichloroethane is heated at reflux temperature for 4 h, cooled to room temperature and concentrated to dryness. The resultant residue is dissolved in ethanol, heated at reflux temperature for 16 h and concentrated *in vacuo* to give the title compound as a white solid, 130 mg, (82% yield). This compound is purified by recrystallization from ethanol/ether to afford a white solid, mp 167-170°C, identified by NMR and mass spectral analyses.


EXAMPLE 10**Preparation of 6-Methoxy-3-(phenylsulfonyl)-5-(piperazin-1-yl)-1-[4-(trifluoromethyl)phenyl]-1H-indazole hydrochloride**

Using essentially the same procedures described in Examples 8 and 9 hereinabove and employing 4-(trifluoromethyl)phenyl boronic acid, the title product is obtained as a white solid, mp 175-177°C, identified by NMR and mass spectral analyses.

5

EXAMPLE 11

Preparation of 5-(4-Benzylpiperazin-1-yl)-1-methyl-3-(phenylsulfonyl)-1H-indazole




A solution of 5-(4-benzylpiperazin-1-yl)-3-(phenylsulfonyl)-1H-indazole (433 mg, 1.00 mmol) in THF at room temperature is treated with NaH (80 mg, 60% in mineral oil, 2.00 mmol), stirred for 0.5 h, treated with methyl iodide (142 mg, 1.00 mmol), stirred for 16 h, and diluted with water and EtOAc. The organic phase is separated, dried over MgSO₄, and concentrated *in vacuo*. The resultant residue is chromatographed (SiO₂, 90:10 EtOAc:hexanes as eluent) to afford the title compound as a white solid, 190 mg (43% yield), mp 195-197°C, identified by mass spectral and NMR analyses.

20

EXAMPLE 12

Preparation of 5-(4-Benzylpiperazin-1-yl)-1-ethyl-3-(phenylsulfonyl)-1H-indazole dihydrochloride

25




A mixture of 5-(4-benzylpiperazin-1-yl)-3-(phenylsulfonyl)-1H-indazole (400
 5 mg, 0.92 mmol), bromoethane (108 mg, 1.00 mmol) and K_2CO_3 (276 mg, 2.00 mmol)
 in DMF is stirred for 16 h at room temperature and diluted water and EtOAc. The
 organic phase is separated, dried over $MgSO_4$, and concentrated *in vacuo*. The
 resultant residue is chromatographed (SiO_2 , 80:20, then 90:10 EtOAc:hexanes as
 eluent) to afford the free amine of the title product as a white solid, 230 mg. A portion
 10 of this solid (115 mg, 0.25 mmol) is dissolved in ethanol, treated with 4M HCl in
 dioxane and concentrated *in vacuo* to give the title compound as a white solid, mp
 >250°C, identified by mass spectral and NMR analyses.

EXAMPLES 13 - 93

15


Preparation of 3-Arylsulfonyl- -5-piperaziny lindazole Derivatives




20

Using essentially the same procedures described in Examples 2 - 12
 hereinabove and employing the appropriate 4-nitrophenylpiperazine as substrate and
 suitable chloromethylarylsulfone as reactant, the compounds shown on Table I are
 obtained and identified by HNMR and mass spectral analyses.


25

TABLE I


Ex No	R2	R3	R4	mp °C
13	phenyl	H	methyl	>250
14	1-naphthyl	H	methyl	>250
15	5-chlorophenyl	H	methyl	>250**
16	phenyl	H	methyl	208-210
17	4-bromophenyl	H	methyl	>250**
18	2-fluorophenyl	H	methyl	160-161
19	3-fluorophenyl	H	methyl	>250
20	phenyl	phenyl	benzyl	148-151
21	phenyl	4-trifluoromethylphenyl	benzyl	169-171
22	phenyl	4-ethoxyphenyl	benzyl	140-144
23	phenyl	3-fluorophenyl	benzyl	146-149
24	phenyl	3-methoxyphenyl	benzyl	135-138
25	phenyl	4-fluorophenyl	benzyl	206-208
26	phenyl	3-chloro-4-fluorophenyl	benzyl	157-161
27	phenyl	4-methylthiophenyl	benzyl	156-160
28	phenyl	3-trifluoromethylphenyl	benzyl	146-150
29	phenyl	3,4-dimethoxyphenyl	benzyl	>200
30	phenyl	4-iodophenyl	benzyl	196-199
31	phenyl	2-phenethyl	H	127-130
32	phenyl	phenyl	H	>200

TABLE I, cont.

Ex No	R2	R3	R4	mp °C
33	phenyl	4-trifluoromethylphenyl	H	170-172
34	phenyl	4-ethoxyphenyl	H	>200
35	phenyl	3-fluorophenyl	H	139-142
36	phenyl	3-methoxyphenyl	H	132-135
37	phenyl	4-fluorophenyl	H	159-161
38	phenyl	3-chloro-4-fluorophenyl	H	>200
39	phenyl	4-methylthiophenyl	H	160-164
40	phenyl	3-trifluoromethylphenyl	H	139-142
41	phenyl	3,4-dimethoxyphenyl	H	156-158
42	phenyl	4-iodophenyl	H	>200
43	phenyl	ethyl	H	>200
44	phenyl	methyl	H	>200
45	1-naphthyl	1-propyl	methyl	**
46	1-naphthyl	ethyl	methyl	**
47	1-naphthyl	methyl	methyl	**
48	1-naphthyl	1-butyl	methyl	**
49	1-naphthyl	3-chlorobenzyl	methyl	**
50	1-naphthyl	2-propyl	methyl	**
51	1-naphthyl	1-(2-methylpropyl)	methyl	**
52	1-naphthyl	benzyl	methyl	**
53	1-naphthyl	H	H	--

TABLE I, cont.


Ex No	R2	R3	R4	mp °C
54	2-thienyl	H	methyl	**
55	3-methylphenyl	H	methyl	**
56	4-methylphenyl	H	methyl	**
57	benzyl	H	methyl	**
58	2-fluorophenyl	H	methyl	**
59	3-fluorophenyl	H	methyl	**
60	4-fluorophenyl	H	methyl	**
61	3-cyanophenyl	H	methyl	**
62	4-cyanophenyl	H	methyl	**
63	3-methoxyphenyl	H	methyl	**
64	4-methoxyphenyl	H	methyl	**
65	2-chlorophenyl	H	methyl	**
66	3-chlorophenyl	H	methyl	**
67	4-chlorophenyl	H	methyl	**
68	2,4-difluorophenyl	H	methyl	**
69	3,4-difluorophenyl	H	methyl	**
70	5-chlorothien-2-yl	H	methyl	**
71	4-isopropylphenyl	H	methyl	**
72	2-naphthyl	H	methyl	**
73	3-chloro-4-fluorophenyl	H	methyl	**
74	4-n-butylphenyl	H	methyl	**
75	4-t-butylphenyl	H	methyl	**

TABLE I, cont.

Ex No	R2	R3	R4	mp °C
76	4-aminophenyl	H	methyl	**
77	5-chloro-2-methoxyphenyl	H	methyl	**
78	2,5-dichlorophenyl	H	methyl	**
79	3,5-dichlorophenyl	H	methyl	**
80	4-methylsulfonylphenyl	H	methyl	**
81	4-bromophenyl	H	methyl	**
82	3,5-bis(trifluoromethyl)phenyl	H	methyl	**
83	2-methylphenyl	H	methyl	**
84	2-cyanophenyl	H	methyl	**
85	4-n-propylphenyl	H	methyl	**
86	3-(difluoromethoxy)phenyl	H	methyl	**
87	4-acetylphenyl	H	methyl	**
88	2,3,4-trifluorophenyl	H	methyl	**
90	2,4,5-trichlorophenyl	H	methyl	**
91	2,6-dichlorophenyl	H	methyl	**
92	2,5-dichlorothien-3-yl	H	methyl	**
93	2-propyl	H	methyl	**

5


** Free Amine

EXAMPLE 94**Preparation of 1-(5-Chloro-2-nitrophenyl)-4-methylpiperazine**

5

A mixture of 4-chloro-2-fluoro-nitrobenzene (28.5 mmol), 1-methylpiperazine (28.5 mmol) and K₂CO₃ (37.0 mmol) in DMF is stirred at 100° C for 16 h and diluted with water and EtOAc. The phases are separated and the aqueous phase is extracted with EtOAc. The organic phase and the extracts are combined, washed sequentially with water and brine, dried over Na₂SO₄ and dried *in vacuo*. The resultant residue is chromatographed (SiO₂, 10% methanol in EtOAc as eluent) to afford the title product.

15

EXAMPLE 95**Preparation of 1-{5-Chloro-2-nitro-3-[[4-fluorophenyl)sulfonyl]methyl}phenyl}-4-methylpiperazine**

20

A stirred mixture of 1-(5-chloro-2-nitrophenyl)-4-methylpiperazine (1.0 mmol) and 4-fluorobenzylsulfonyl chloride (1.0 mmol) in THF is treated with 2.2 mL of a 1M KOt-Bu (2.2 mmol) solution in THF at -78° C, warmed to room temperature over a 2 h period, quenched with acetic acid and concentrated *in vacuo*. The resultant residue is partitioned between EtOAc and aqueous Na₂CO₃. The phases are separated and

25


the aqueous phase is extracted with EtOAc. The organic phase is combined with the extracts, dried over Na₂SO₄ and concentrated to dryness *in vacuo* to afford the title product.

5

EXAMPLE 96

Preparation of 4-Chloro-2-[(4-fluorophenyl)sulfonyl]methyl}-6-(4-methylpiperazin-1-yl)aniline

10




A solution of 4-chloro-2-[(4-fluorophenyl)sulfonyl]methyl}-6-(4-methylpiperazin-1-yl)aniline (1.0 mmol) in methanol is treated with Sn foil (593.3 mg), followed by 5.8 mL of 12 M HCl, heated at 60° C for 4 h (until reaction is complete by thin layer chromatography), cooled to 0° C, treated with solid NaOH to pH >10 and extracted with CH₂Cl₂. The extracts are combined, washed with water and concentrated to dryness *in vacuo* to give the title compound.

20

EXAMPLE 97

Preparation of 5-Chloro-3[(4-fluorophenyl)sulfonyl]-7-(4-methylpiperazin-1-yl)-1H-indazole

25




A solution of NaNO_2 (2.0 mg) in water at 0° is treated dropwise with a solution of 4-chloro-2-[[[(4-fluorophenyl)sulfonyl]methyl]-6-(4-methylpiperazin-1-yl)aniline (102.2 mg, 0.25 mmol) in 1M HCl, allowed to warm to room temperature over a 2 h period, treated with saturated Na_2CO_3 to pH >10 and filtered. The filtercake is washed with water and dried in vacuo to give the title compound, identified by mass spectral and NMR analyses.

10


EXAMPLE 98-101**Preparation of 3 (Arylsulfonyl)-7-(4-methylpiperazin-1-yl)-1H-indazole Derivatives**

15



Using essentially the same procedures described hereinabove in Examples 95-97 and employing the appropriate arylsulfonyl chloride, the compounds shown in Table II are obtained and identified by mass spectral and NMR analyses.

20


TABLE II

Ex. No.	R1	R2	% Yield (Over 3 Steps)
98	Cl	4-(isopropyl)phenyl	74
99	Cl	3-(trifluoromethyl)phenyl	65
100	Cl	1-naphthyl	7.7
101	H	1-naphthyl	--

5

EXAMPLE 102Preparation of 1-Fluoro-3-[4-(*t*-butoxycarbonyl)piperazin-1-yl]benzaldehyde

10




15 A stirred solution of 2,6-difluorobenzaldehyde 5 (10.0 g, 70 mmol) in *N,N*-dimethylformamide is treated with *tert*-butyl piperazine-1-carboxylate (Boc) 6 (14.4 g, 77.0 mmol) and potassium carbonate (11.7 g, 85.0 mmol), stirred at 80 °C for 16 h, cooled to ambient temperature, diluted with water and extracted with methylene chloride. The extracts are combined, washed sequentially with water and brine, dried over sodium sulfate, and concentrated *in vacuo* to afford a crude oil. This oil is purified by column chromatography (silica gel, 15:85 ethyl acetate/hexanes) to afford

the title compound as a yellow solid, 15.5g (71% yield), identified by NMR and mass spectral analyses.

EXAMPLE 103

5

Preparation of 4-[4-(t-Butoxycarbonyl)piperazin-1-yl]-1H-indazole




A solution of 7 (20.0 g, 64.9 mmol) and hydrazine monohydrate (30 mL) in methyl sulfoxide is stirred at 95 °C for 96 h, cooled to room temperature, partitioned between saturated sodium bicarbonate and ether. The phases are separated and the aqueous phase is extracted with ether. The extracts are combined with the organic phase, washed sequentially with water and brine, dried over sodium sulfate and concentrated *in vacuo* to a crude residue. This residue is purified by column chromatography (silica gel, 10:90 to 30:70 ethyl acetate/hexanes) to afford the title compound as a light yellow solid, 12.3 g (57% yield), identified by NMR and mass spectral analyses.

EXAMPLE 104

20


Preparation of 4-[4-(t-Butoxycarbonyl)piperazin-1-yl]-3-iodo-1H-indazole




A solution of 4-[4-(t-butoxycarbonyl)piperazin-1-yl]-1H-indazole (5.00 g, 16.5 mmol) in *N,N*-dimethylformamide is treated with powdered potassium hydroxide

(1.85 g, 33.07 mmol) and iodine (4.61 g, 18.2 mmol), stirred at room temperature for 16 h, diluted with ethyl acetate and quenched with 15% aqueous sodium metabisulfite. The phases are separated and the aqueous phase is extracted with ethyl acetate. The extracts and the organic phase are combined, washed with brine, dried over sodium sulfate and concentrated *in vacuo* to afford a crude solid. This solid is purified by column chromatography (silica gel, 10:90 to 15:85 to 20:80 ethyl acetate/hexanes) to afford the title product as a yellow solid, 2.40 g (35% yield) identified by HNMR and mass spectral analyses.

10

EXAMPLE 105Preparation of 4-[4-(t-Butoxycarbonyl)piperazin-1-yl]-3-(phenylsulfonyl)-1H-indazole


A mixture of 4-[4-(t-butoxycarbonyl)piperazin-1-yl]-3-iodo-1H-indazole (0.110 g, 0.257 mmol), sodium benzene sulfinate (0.068 g, 0.411 mmol), copper(I) iodide (0.073 g, 0.386 mmol) and *N,N*-dimethylformamide is added to a screw-capped test-tube with Teflon-lined septum. The tube is evacuated and backfilled with argon. This procedure is repeated twice, then the tube is heated to 125 °C for 4 h. The reaction mixture is cooled, partitioned between ethyl acetate and water and filtered through a pad of diatomaceous earth. The filtrate is separated. The organic phase is washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The resulting crude material is purified by column chromatography (silica gel, 40:60 ethyl acetate/hexanes) to afford the title compound as a pale yellow foam, 0.022 g (18% yield), identified by NMR and mass spectral analyses.

EXAMPLE 106Preparation of 3-(Phenylsulfonyl)-4-piperazin-1-yl-1H-indazole Hydrochloride

10 A solution of 4-[4-(t-butoxycarbonyl)piperazin-1-yl]-3-(phenylsulfonyl)-1H-indazole (0.022 g, 0.050 mmol) in methanol is treated with a 2 M solution of HCl in ether (0.4 mL), heated at reflux temperature for 1 h, cooled to room temperature and evaporated to dryness. The residue is re-dissolved in methanol and concentrated. Repeating this procedure three more times affords the title product as a white solid, 0.009 g (40% yield), identified by NMR and mass spectral analyses.

EXAMPLE 107

15

Preparation of 4-[4-t-Butoxycarbonyl)piperazin-1-yl]-3-(1-naphthylthio)-1H-indazole


20 A mixture of 4-[4-(t-butoxycarbonyl)piperazin-1-yl]-3-iodo-1H-indazole (0.068 g, 0.159 mmol) copper(I) iodide (0.005 g, 0.024 mmol) and potassium carbonate (0.044 g, 0.318 mmol) are added to a screw-capped test tube with a Teflon-lined septum. The tube is evacuated and backfilled with argon and this procedure repeated twice more. The reaction mixture is treated sequentially with 2-propanol

(1.5 mL), ethylene glycol (19 μ L, 0.318 mmol) and 1-naphthalenethiol (0.051 g, 0.318 mmol), heated at 130°–140° C for 8 h, cooled to room temperature, diluted with ethyl acetate and filtered through diatomaceous earth. The filtrate is concentrated under reduced pressure. The resultant crude material is purified by flash chromatography (silica gel, 40:60 ethyl acetate/hexanes) to afford the title compound as a colorless oil, 0.033 g (45% yield), identified by NMR and mass spectral analyses.

EXAMPLE 108


10

Preparation of 4-[4-(t-Butoxycarbonyl)piperazin-1-yl]-3-(1-naphthylsulfonyl)-1H-indazole



15 A stirred solution of 4-[4-t-butoxycarbonyl)piperazin-1-yl]-3-(1-naphthylthio)-1H-indazole (0.030 g, 0.065 mmol) in glacial acetic acid is treated with 35% aqueous hydrogen peroxide (0.127 g, 1.30 mmol) stirred for 24 h, quenched with 10% aqueous sodium metabisulfite solution (1 mL) and concentrated *in vacuo*. The resulting solid is directly added to a suspension of 10% palladium-on-carbon (3 mg) in methanol (10 mL) and the mixture is shaken on a Parr hydrogenator under a 40 psi hydrogen atmosphere for 2 h and filtered through diatomaceous earth. The filtrate is concentrated *in vacuo*. Purification of the resulting crude material by flash chromatography (silica gel, 30:70 ethyl acetate/hexanes) affords the title compound as a colorless oil, 0.011 g (35% yield), identified by NMR and mass spectral analyses.

25

EXAMPLE 109Preparation of 3-(1-Naphthylsulfonyl)-4-piperazin-1-yl-1H-indazole Hydrochloride


Using essentially the same procedure described in Example 106 hereinabove and employing 4-[4-(t-butoxycarbonyl)piperazin-1-yl]-3-(1-naphthylsulfonyl)-1H-indazole as substrate, the title product is obtained as a light yellow solid (74% yield), mp 220°C dec, identified by NMR and mass spectral analyses.

10

EXAMPLES 110-114Preparation of 3-(Arylsulfonyl)-4-piperazin-1-yl-1H-indazole Hydrochloride

15


Derivatives



Using essentially the same procedures described in Examples 102-109 hereinabove and employing the appropriate arylthiol or arylsulfinate, the compounds shown on Table III are obtained and identified by NMR and mass spectral analyses.


20

TABLE III



Ex. No.	R2	mp °C
110	4-methylphenyl	190-192
111	4-chlorophenyl	230 dec
112	3-fluorophenyl	180-183
113	4-methoxyphenyl	178-182
114	2-naphthyl	200-202

5

EXAMPLE 115Preparation of 6-[4-t-Butoxycarbonyl)piperazin-1-yl]-1H-indazole


10

15 A stirred solution of 6-aminoindazole (13.3 g, 100 mmol) in *n*-butanol is treated with bis(2-chloroethyl)amine hydrochloride (28.6 g, 160 mmol) and potassium carbonate (34.6 g, 250 mmol), heated to reflux temperature for 72 h, cooled to room temperature and concentrated *in vacuo*. A stirred solution of the resultant crude material in 1,4-dioxane at 0° C is treated with 1 N sodium hydroxide solution (200 mL) and di-*tert*-butyldicarbonate (34.9 g, 160 mmol), stirred at ambient temperature for 24 h and concentrated *in vacuo* to afford a crude oil. This oil is partitioned between


ethyl acetate and water. The organic layer is separated, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. Further purification of this resultant residue by flash chromatography (silica gel, 40:60 ethyl acetate/hexanes) affords the title compound as a colorless oil, 11.5 g (38% yield) identified by NMR and mass spectral analyses.

EXAMPLE 116

Preparation of 6-[4-t-Butoxycarbonyl)piperazin-1-yl]-3-iodo-1H-indazole




A stirred solution of 6-[4-(tert-butoxycarbonyl)piperazin-1-yl]-1H-indazole (7.20 g, 23.8 mmol) in *N,N*-dimethylformamide at 0 °C is treated with powdered potassium hydroxide (5.40 g, 95.4 mmol) followed by the dropwise addition of a solution of iodine (10.9 g, 42.9 mmol) in *N,N*-dimethylformamide, stirred at ambient temperatures for 16 h, diluted with ethyl acetate and quenched with 10% aqueous sodium metabisulfite solution. The phases are separated. The aqueous phase is extracted with ethyl acetate. The extracts are combined with the organic phase, washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The resulting crude material is purified by flash chromatography (silica gel, 30:70 ethyl acetate/hexanes) to afford the title compound as a yellow solid, 11.5 g (38% yield) identified by NMR and mass spectral analyses.

EXAMPLE 117Preparation of 6-[4-t-Butoxycarbonyl]piperazin-1-yl]-3-(phenylsulfonyl)-1H-indazole

Using essentially the same procedure described in Example 105 hereinabove and employing of 6-[4-t-butoxycarbonyl]piperazin-1-yl]-3-iodo-1H-indazole and sodium benzenesulfinate affords the title compound as a colorless foam (80% yield), identified by NMR and mass spectral analyses.


10

EXAMPLE 118Preparation of 3-(Phenylsulfonyl)- 6-piperazin-1-yl]-1H-indazole Hydrochloride


20 Using essentially the same procedure described in Example 106 hereinabove and employing 6-[4-t-butoxycarbonyl]piperazin-1-yl]-3-(phenylsulfonyl)-1H-indazole as substrate, the title compound is obtained as an off-white solid (81% yield), identified by NMR and mass spectral analyses.

EXAMPLES 119-124Preparation of 3-(Arylsulfonyl)-6-piperazin-1-yl-1H-indazole Hydrochloride Compounds

5




Using essentially the same procedures described in Examples 117 and 118
 10 hereinabove and employing the appropriate arylsulfinate, the compounds shown in Table IV are obtained and identified by NMR and mass spectral analyses.

TABLE IV

15


Ex. No.	R2	mp °C
119	4-methylphenyl	172-175
120	4-(trifluoromethyl)phenyl	189-191
121	4-chlorophenyl	175-178
122	3-fluorophenyl	260 dec
123	4-methoxyphenyl	165-170
124	2-naphthyl	260 dec

EXAMPLE 125Preparation of 6-[4-t-Butoxycarbonyl)piperazin-1-yl]-3-(1-naphthylthio)-1H-indazole

Using essentially the same procedure described in Example 107 hereinabove and employing 6-[4-t-butoxycarbonyl)piperazin-1-yl]-3-iodo-1H-indazole as substrate,
 10 the title compound is obtained as a colorless oil (78% yield), identified by NMR and mass spectral analyses.

EXAMPLE 126

15 Preparation of 6-[4-t-Butoxycarbonyl)piperazin-1-yl]-3-(1-naphthylsulfonyl)-1H-indazole




20

A stirred solution of 6-[4-t-butoxycarbonyl)piperazin-1-yl]-3-(1-naphthylthio)-1H-indazole (0.240 g, 0.521 mmol) in methylene chloride is treated with meta-chloroperoxybenzoic acid (mcpba) (77% pure, 0.350 g, 1.56 mmol) and sodium

bicarbonate (0.219 g, 2.61 mmol), stirred at ambient temperatures for 16 h and partitioned between ethyl acetate and water. The phases are separated and the aqueous phase is extracted with methylene chloride. The extracts and the organic phase are combined, washed sequentially with 10% aqueous sodium bicarbonate solution and brine, dried over sodium sulfate and concentrated *in vacuo* to afford a crude oil. This oil is added to a suspension of 10% palladium-on-carbon (40 mg) in methanol and shaken on a Parr hydrogenator under a 40 psi hydrogen atmosphere for 2 h and filtered through diatomaceous earth. The filtrate is concentrated *in vacuo*. The resulting crude material is purified by flash chromatography (silica gel, 50:50 ethyl acetate/hexanes) to afford the title compound as a colorless oil, 0.106 g (41% yield), identified by NMR and mass spectral analyses.

EXAMPLE 127

15 Preparation of 3-(1-Naphthylsulfonyl)-6-piperazin-1-yl-1H-indazole Hydrochloride



20 Using essentially the same procedure described in Example 106 hereinabove and employing of 6-[4-t-butoxycarbonyl]piperazin-1-yl)-3-(1-naphthylsulfonyl)-1H-indazole as substrate, the title product is obtained as a white solid (98% yield), mp 205-210 °C, identified by NMR and mass spectral analyses.

25

EXAMPLE 128**Comparative Evaluation of 5-HT6 Binding Affinity of Test Compounds**

5 The affinity of test compounds for the serotonin 5-HT6 receptor is evaluated in the following manner. Cultured Hela cells expressing human cloned 5-HT6 receptors are harvested and centrifuged at low speed (1,000 x g) for 10.0 min to remove the culture media. The harvested cells are suspended in half volume of fresh physiological phosphate buffered saline solution and recentrifuged at the same
10 speed. This operation is repeated. The collected cells are then homogenized in ten volumes of 50 mM Tris.HCl (pH 7.4) and 0.5 mM EDTA. The homogenate is centrifuged at 40,000 x g for 30.0 min and the precipitate is collected. The obtained pellet is resuspended in 10 volumes of Tris.HCl buffer and recentrifuged at the same speed. The final pellet is suspended in a small volume of Tris.HCl buffer and the
15 tissue protein content is determined in aliquots of 10-25 μ l volumes. Bovine Serum Albumin is used as the standard in the protein determination according to the method described in Lowry et al., J. Biol. Chem., 193:265 (1951). The volume of the suspended cell membranes is adjusted to give a tissue protein concentration of 1.0 mg/ml of suspension. The prepared membrane suspension (10 times concentrated)
20 is aliquoted in 1.0 ml volumes and stored at -70° C until used in subsequent binding experiments.

 Binding experiments are performed in a 96 well microtiter plate format, in a total volume of 200 μ l. To each well is added the following mixture: 80.0 μ l of incubation buffer made in 50 mM Tris.HCl buffer (pH 7.4) containing 10.0 mM MgCl₂
25 and 0.5 mM EDTA and 20 μ l of [³H]-LSD (S.A., 86.0 Ci/mmol, available from Amersham Life Science), 3.0 nM. The dissociation constant, K_D of the [³H]LSD at the human serotonin 5-HT6 receptor is 2.9 nM, as determined by saturation binding with increasing concentrations of [³H]LSD. The reaction is initiated by the final addition of 100.0 μ l of tissue suspension. Nonspecific binding is measured in the presence of
30 10.0 μ M methiothepin. The test compounds are added in 20.0 μ l volume.

 The reaction is allowed to proceed in the dark for 120 min at room temperature, at which time, the bound ligand-receptor complex is filtered off on a 96 well unifier with a Packard Filtermate[®] 196 Harvester. The bound complex caught

on the filter disk is allowed to air dry and the radioactivity is measured in a Packard TopCount[®] equipped with six photomultiplier detectors, after the addition of 40.0 μ l Microscint[®]-20 scintillant to each shallow well. The unifilter plate is heat-sealed and counted in a PackardTopCount[®] with a tritium efficiency of 31.0%.

5 Specific binding to the 5-HT₆ receptor is defined as the total radioactivity bound less the amount bound in the presence of 10.0 μ M unlabeled methiothepin. Binding in the presence of varying concentrations of test compound is expressed as a percentage of specific binding in the absence of test compound. The results are plotted as log % bound versus log concentration of test compound. Nonlinear
10 regression analysis of data points with a computer assisted program Prism[®] yields both the IC₅₀ and the K_i values of test compounds with 95% confidence limits.

The amount of displacement by the test compound is given in percent (%) inhibition and is derived from the following equation:

15

$$\% \text{ inhibition} = \left(1 - \frac{B_0 - \text{NSB}}{\text{TB} - \text{NSB}} \right) 100$$

where B₀ is the amount of CPM bound in the presence of the testing agent. NSB
20 represents the CPM bound in the presence of a saturating concentration of a displacer and TB represents the total amount of CPM bound at zero (0) concentration of test compound.

Alternatively, a linear regression line of data points is plotted, from which the IC₅₀ value is determined and the K_i value is determined based upon the following
25 equation:

$$K_i = \text{IC}_{50} / (1 + L/K_D)$$

where L is the concentration of the radioactive ligand used and K_D is the dissociation constant of the ligand for the receptor, both expressed in nM.

Using this assay, the % inhibition and K_i values are determined and
30 compared to those values obtained by representative compounds known to demonstrate binding to the 5-HT₆ receptor. The data are shown in TableV, below.

TABLE V

Ex. No.	5-HT6 Binding Ki (nM)	% Inhibition at 1μM
4	86	--
5	--	53
7	113	--
8	399	--
9	97	--
10	224	--
11	--	67
12	64	--
13	2	--
14	1	--
15	2	--
16	13	--
17	7	--
18	6	--
19	2	--
20	--	63
21	--	22
22	--	25
23	--	5
24	--	23
25	--	74
26	--	8
27	--	13
28	136	--
29	--	24
30	--	30
31	3	--
32	18	--
33	19	--
34	135	--

TABLE V, cont.

EX. No.	5-HT₆ Binding K_i (nM)	% Inhibition at 1μM
35	38	--
36	20	--
37	4	--
38	73	--
39	69	--
40	62	--
41	13	--
42	117	--
43	3	--
44	--	67
45	2	--
46	2	--
47	2	--
48	4	--
49	6	--
50	2	--
51	1	--
52	2	--
53	1	--
54	8	--
55	1	--
56	5	--
58	3	--
59	2	--
60	5	--
63	1	--
64	12	--
65	4	--
66	1	--


TABLE V, cont.

EX. No.	5-HT6 Binding Ki (nM)	% Inhibition at 1μM
67	8	--
68	14	--
69	28	--
70	3	--
71	4	--
72	2	--
74	26	--
77	2	--
80	62	--
81	7	--
83	4	--
85	10	--
86	1	--
89	10	--
91	6	--
92	1	--
97	12	--
98	22	--
99	20	--
100	5	--
101	5	--

Comparative Examples	5-HT6 binding Ki (nM)
Clozapine	6.0
Loxapine	41.4
Bromocriptine	23.0
Methiothepin	8.3
Mianserin	44.2
Olanzapine	19.5

CLAIMS:

1. A compound of formula I



wherein

A is C, CR₈ or N;

10 R₁ is H, halogen, CN, COR₉, OCO₂R₁₀, CO₂R₁₁, CONR₁₂R₁₃, SO_xR₁₄, NR₁₅R₁₆, OR₁₇ or a C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl or heteroaryl group each optionally substituted;

15 R₂ is an optionally substituted C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl, or heteroaryl group or an optionally substituted 8- to 13-membered bicyclic or tricyclic ring system having a N atom at the bridgehead and optionally containing 1, 2 or 3 additional heteroatoms selected from N, O or S ;

R₃ is H or a C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl or heteroaryl group each optionally substituted;

R₄ is H or a C₁-C₆alkyl or C₃-C₇cycloalkyl group each optionally substituted;

20 R₅, R₆ and R₇ are each independently H or a C₁-C₆alkyl or C₃-C₇cycloalkyl group each optionally substituted;

m and p are each independently an integer of 1, 2 or 3;

n is an integer of 1 or 2;

R₈ is H, OH or an optionally substituted C₁-C₆alkoxy group;

25 R₉, R₁₀, R₁₁ and R₁₇ are each independently H or a C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₃-C₆cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;

5 R_{12} , R_{13} , R_{15} and R_{16} are each independently H or an optionally substituted
 C_1 - C_4 alkyl group or R_{12} and R_{13} or R_{15} and R_{16} may be taken together
 with the atom to which they are attached to form a 5- to 7-membered
 ring optionally containing another heteroatom selected from O, NR_{18}
 or SO_x ;
 R_{14} is a C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_6 cycloalkyl,
 cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;
 R_{18} is H or a C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_7 cycloalkyl,
 cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;
 10 x is 0 or an integer of 1 or 2; and
 --- represents a single bond or a double bond; or a
 stereoisomer thereof or a pharmaceutically acceptable salt thereof.

- 15 2. A compound according to claim 1 wherein n is 1.
3. A compound according to claim 1 or claim 2 wherein R_4 is H or an
 optionally substituted C_1 - C_4 alkyl group.
- 20 4. A compound according to any one of claims 1 to 3 wherein R_2 is an
 optionally substituted phenyl, naphthyl or heteroaryl group.
5. A compound according to any one of claims 1 to 4 wherein the 6- or 7-
 membered azacyclic ring is attached to the indazole in the 5 or 7 position.
- 25 6. A compound according to any one of claims 1 to 5 wherein A is N.
7. A compound according to any one of claims 1 to 6 wherein R_3 is
 selected from hydrogen, phenyl and C_1 - C_4 alkyl which may be substituted by
 optionally substituted phenyl.
- 30 8. A compound according to any one of claims 1 to 7 wherein m is 1 and
 R_1 is hydrogen or halogen.

9. A compound according to claim 1 which is one of the following:
- 5-(4-methylpiperazin-1-yl)-3-(1-naphthylsulfonyl)-1H-indazole;
 - 3-(1-naphthylsulfonyl)-5-piperazin-1-yl-1H-indazole;
 - 1-(3-chlorobenzyl)-5-(4-methylpiperazin-1-yl)-3-(1-naphthylsulfonyl)-1H-indazole;
 - 5 7-(4-methylpiperazin-1-yl)-3-(phenylsulfonyl)-1H-indazole;
 - 3-(phenylsulfonyl)-7-(4-propylpiperazin-1-yl)-1H-indazole;
 - 3-(phenylsulfonyl)-7-piperazin-1-yl-1H-indazole;
 - 5-(4-methylpiperazin-1-yl)-3-(phenylsulfonyl)-1H-indazole;
 - 3-(phenylsulfonyl)-5-(4-propylpiperazin-1-yl)-1H-indazole;
 - 10 3-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole;
 - 5-piperazin-1-yl-3-[(3-cyanophenyl)sulfonyl]-1H-indazole;
 - 7-(4-methylpiperazin-1-yl)-3-[(2-ethylphenyl)sulfonyl]-1H-indazole;
 - 3-(1-naphthylsulfonyl)-7-(4-propylpiperazin-1-yl)-1H-indazole;
 - 5-(4-phenethylpiperazin-1-yl)-3-(1-naphthylsulfonyl)-1H-indazole;
 - 15 5-(4-methylpiperazin-1-yl)-3-(2-naphthylsulfonyl)-1H-indazole;
 - 3-[(2-chloro-4-fluorophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
 - 1-methyl-3-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole;
 - 1-phenyl-3-(phenylsulfonyl)-5-piperazin-1-yl-1H-indazole;
 - 1-methyl-3-(phenylsulfonyl)-7-piperazin-1-yl-1H-indazole;
 - 20 1-phenyl-3-(phenylsulfonyl)-7-(piperazin-1-yl)-1H-indazole;
 - 7-piperazin-1-yl-3-[(3-fluorophenyl)sulfonyl]-1H-indazole;
 - 3-[(4-fluorophenyl)sulfonyl]-7-(4-methylpiperazin-1-yl)-1H-indazole;
 - 3-[(2-chlorophenyl)sulfonyl]-7-piperazin-1-yl-1H-indazole;
 - 3-[(4-aminophenyl)sulfonyl]-7-(piperazin-1-yl)-1H-indazole;
 - 25 5-piperazin-1-yl-3-[(3-fluorophenyl)sulfonyl]-1H-indazole;
 - 3-[(4-fluorophenyl)sulfonyl]-5-(4-methylpiperazin-1-yl)-1H-indazole;
 - 3-[(2-chlorophenyl)sulfonyl]-5-(4-propylpiperazin-1-yl)-1H-indazole;
 - 3-[(4-aminophenyl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
 - 3-[(5-chlorothiophen-2-yl)sulfonyl]-5-piperazin-1-yl-1H-indazole;
 - 30 3-[(5-chlorothiophen-2-yl)sulfonyl]-7-piperazin-1-yl-1H-indazole;
 - 4-chloro-3-(phenylsulfonyl)-7-piperazin-1-yl-1H-indazole;
 - 5-fluoro-3-(phenylsulfonyl)-7-piperazin-1-yl-1H-indazole;
 - 6-fluoro-3-(phenylsulfonyl)-7-piperazin-1-yl-1H-indazole;

- 4-chloro-3-(phenylsulfonyl)-5-piperazin-1-yl-1*H*-indazole;
 7-fluoro-3-(phenylsulfonyl)-5-piperazin-1-yl-1*H*-indazole;
 6-fluoro-3-(phenylsulfonyl)-5-piperazin-1-yl-1*H*-indazole;
 6-(4-phenethylpiperazin-1-yl)-3-(phenylsulfonyl)-1*H*-indazole;
 5 6-(4-methylpiperazin-1-yl)-3-(phenylsulfonyl)-1*H*-indazole;
 6-(4-propylpiperazin-1-yl)-3-(phenylsulfonyl)-1*H*-indazole;
 3-(phenylsulfonyl)-6-piperazin-1-yl-1*H*-indazole;
 4-piperazin-1-yl-3-(phenylsulfonyl)-1*H*-indazole;
 4-(4-methylpiperazin-1-yl)-3-(phenylsulfonyl)-1*H*-indazole;
 10 4-(4-propylpiperazin-1-yl)-3-(phenylsulfonyl)-1*H*-indazole;
 3-(phenylsulfonyl)-4-piperazin-1-yl-1*H*-indazole;
 3-[(5-chlorothien-2-yl)sulfonyl]-5-piperidin-1-yl-1*H*-indazole;
 3-[(5-chlorothien-2-yl)sulfonyl]-7-piperidin-1-yl-1*H*-indazole;
 1-methyl-3-(phenylsulfonyl)-5-piperidin-1-yl-1*H*-indazole;
 15 3-[(3-fluorophenyl)sulfonyl]-5-piperidin-1-yl-1*H*-indazole;
 1-methyl-3-(phenylsulfonyl)-7-(4-methylpiperidin-1-yl)-1*H*-indazole;
 3-[(3-fluorophenyl)sulfonyl]-7-piperidin-1-yl-1*H*-indazole;
 3-[(3-fluorophenyl)sulfonyl]-5-piperidin-1-yl-1*H*-indazole;
 3-(phenylsulfonyl)-5-piperidin-1-yl-1*H*-indazole;
 20 1-methyl-3-[(3-fluorophenyl)sulfonyl]-7-piperidin-1-yl-1*H*-indazole;
 3-(phenylsulfonyl)-7-piperidin-1-yl-1*H*-indazole;
 or a stereoisomer thereof; or a pharmaceutically acceptable salt thereof.

10. A method for the treatment of a central nervous system disorder related to or
 25 affected by the 5-HT₆ receptor in a patient in need thereof which comprises providing
 to said patient a therapeutically effective amount of a compound of formula I as
 claimed in any one of claims 1 to 9 or a stereoisomer thereof; or a pharmaceutically
 acceptable salt thereof.

30 11. A method according to claim 10 wherein said disorder is a motor
 disorder, anxiety disorder or cognitive disorder.

12. A method according to claim 10 wherein said disorder is a neurodegenerative disorder.


13. A method according to claim 11 wherein said disorder is selected from the group consisting of: attention deficit disorder; obsessive compulsive disorder; withdrawal from drug, alcohol or nicotine addiction; schizophrenia; depression; and Alzheimer's disease.

14. A method according to claim 12 wherein said disorder is selected from the group consisting of: stroke; head trauma; and neuropathic pain.

15. A pharmaceutical composition which comprises a pharmaceutically acceptable carrier and a compound of formula I as claimed in any one of claims 1 to 9 or a stereoisomer thereof; or a pharmaceutically acceptable salt thereof.


16. A process for preparing a compound of formula (I) according to claim 1 which comprises one of the following

(a) diazotising and cyclising a compound of formula (VIIA):



20 wherein the dotted line, m, n, p, A, R₁, R₂, R₄, R₅, R₆ and R₇ are as defined in claim 1, e.g. with NaNO₂/acid to give a compound of formula (I) wherein R₃ is hydrogen, or


(b) alkylating or arylating a protected compound of formula (IA)



wherein the dotted line, m, n, p, A, R₁, R₂, R₅, R₆ and R₇ are as defined in claim 1 and P is a protecting group or R₄ where R₄ is as defined in claim 1 excepting hydrogen, with an alkylating or arylating agent of formula R₃-LG where LG is a leaving group and R₃ is a C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl or heteroaryl group each optionally substituted; and if required removing the protecting group, to give a corresponding compound of formula (I) where R₃ is a C₁-C₆alkyl, C₃-C₇cycloalkyl, aryl or heteroaryl group each optionally substituted;

10 or


(c) alkylating a compound of formula (IB)




15 wherein the dotted line, m, n, p, A, R₁, R₂, R₃, R₅, R₆ and R₇ are as defined in claim 1, with an alkylating agent of formula R₄-LG where LG is a leaving group and R₄ is a C₁-C₆alkyl or C₃-C₇cycloalkyl group each optionally substituted, to give a corresponding compound of formula (I) where R₄ is a C₁-C₆alkyl or C₃-C₇cycloalkyl group each optionally substituted;

20 or

(d) reacting a compound of formula (XII):




wherein m , R_1 , R_2 and R_3 are as defined in claim 1, and LG' is a leaving group such as Cl, Br, I or an activated hydroxyl group such as triflate, with a compound of
5 formula (V):



wherein n , p , R_4 , R_5 , R_6 and R_7 are as defined in claim 1, in the presence of a catalyst, e.g., a palladium or nickel catalyst, to give a corresponding compound of formula (I);

10 or


(e) reducing a compound of formula (XII):



wherein m , n , p , R_1 , R_2 , R_5 , R_6 and R_7 are as defined in claim 1, P is a protecting
15 group or R_4 as defined in claim 1 and P' is a protecting group or R_3 as defined in claim 1, if required removing any protecting group, to give a corresponding compound of formula (I) wherein A is CH;

or

20 (f) acidifying a compound of formula (XII):




wherein m, n, p, R₁, R₂, R₅, R₆ and R₇ are as defined in claim 1, P is a protecting group or R₄ as defined in claim 1 and P' is a protecting group or R₃ as defined in claim 1, if required removing any protecting group, to give a corresponding compound of formula (I) wherein A is C and ---- represents a double bond;
or

(g) converting a basic compound of formula (I) as defined in claim 1 to an acid addition salt thereof or vice versa.

10 or

(h) separating a stereoisomer of a compound of formula (I) from a mixture thereof.



(I)