




(72) HILLMAN, JENNIFER L., US

(72) GOLI, SURYA K., US

(71) INCYTE PHARMACEUTICALS, INC., US

(51) Int.Cl.⁶ C12N 15/12, C07K 14/825, C12Q 1/68, C07K 16/18,
A61K 38/17, C12N 15/11, C12N 5/10

(30) 1997/01/17 (08/785,530) US

(54) **NOUVELLE METALLOTHIONEINE HUMAINE**

(54) **NOVEL HUMAN METALLOTHIONEIN**

```

          9      18      27      36      45      54
5' NNC TCC CCT GAG TAG AAA AGC AGC CGC AGT TTG GNC GCT CCA CCA CGN CGT CCG

          63      72      81      90      99      108
GGT GGG CCT AGC AGT CGC TCC ATT TAT CGC TTG AGA TCT CCA GCC TTA CCG CCG

          117     126     135     144     153     162
CTC GAA ATG GAC CCC AAC TGC TCC TGC ACC ACT GGT GTC TCC TGC GCC TGC ACC
  M  D  P  N  C  S  C  T  T  G  V  S  C  A  C  T

          171     180     189     198     207     216
GGC TCC TGC AAG TGC AAA GAG TGC AAA TGC ACC TYC TGC AAG AAG AGC TGC TGC
G  S  C  K  C  K  E  C  K  C  T  X  C  K  K  S  C  C

          225     234     243     252     261     270
TCC TGC TGC CCC GTG GGC TGT GGC AAG TGT GGC CAC GGC TGT GTC TGC AAA GGG
S  C  C  P  V  G  C  A  K  C  A  E  G  C  V  C  K  G

          279     288     297     306     315     324
ACG TTG GAG AAC TCC AGC TGC TGT GGC TGA TGT GGG AAC AGC TCT TCT CCC AGA
T  L  E  N  C  S  C  C  A

          333     342     351     360     369     378
TGT TAA TAG AAC AAG CTG CAC AAC CTG GAT TTT TTT TCA ATA CGA TAC TGA GCC
3'

```

(57) L'invention concerne une métallothionéine humaine (HMBP-1), ainsi que des polynucléotides qui identifient et codent pour la HMBP-1. Elle concerne également des vecteurs d'expression et des cellules hôtes mis au point par génie génétique, qui comprennent les séquences d'acide nucléique codant pour la HMBP-1, ainsi qu'un procédé permettant de produire la HMBP-1. Elle concerne aussi des agonistes, anticorps ou antagonistes qui se lient spécifiquement avec la HMBP-1, et leur utilisation pour prévenir et traiter les maladies associées à l'expression de la HMBP-1. Elle concerne en outre l'utilisation de molécules antisens par rapport aux polynucléotides codant pour la HMBP-1 pour traiter les maladies associées à l'expression de la HMBP-1. Elle concerne enfin des analyses diagnostiques qui utilisent le polynucléotide, ou des fragments ou le complément de celui-ci, et les anticorps qui se fixent spécifiquement sur la HMBP-1.

(57) The present invention provides a human metallothionein (HMBP-I) and polynucleotides which identify and encode HMBP-I. The invention also provides genetically engineered expression vectors and host cells comprising the nucleic acid sequences encoding HMBP-I and a method for producing HMBP-I. The invention also provides for agonists, antibodies, or antagonists specifically binding HMBP-I, and their use, in the prevention and treatment of diseases associated with expression of HMBP-I. Additionally, the invention provides for the use of antisense molecules to polynucleotides encoding HMBP-I for the treatment of diseases associated with the expression of HMBP-I. The invention also provides diagnostic assays which utilize the polynucleotide, or fragments or the complement thereof, and antibodies specifically binding HMBP-I.



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/825, C12N 15/11, 5/10, A61K 38/17, C07K 16/18, C12Q 1/68		A3	(11) International Publication Number: WO 98/31795
			(43) International Publication Date: 23 July 1998 (23.07.98)
(21) International Application Number: PCT/US97/24129		(81) Designated States: AT, AU, BR, CA, CH, CN, DE, DK, ES, FI, GB, IL, JP, KR, MX, NO, NZ, RU, SE, SG, US, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 30 December 1997 (30.12.97)		Published <i>With international search report.</i>	
(30) Priority Data: 08/785,530 17 January 1997 (17.01.97) US		(88) Date of publication of the international search report: 11 September 1998 (11.09.98)	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/785,530 (CIP) Filed on 17 January 1997 (17.01.97)			
(71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive #12, Mountain View, CA 94040 (US). GOLI, Surya, K. [IN/US]; 620 Iris Avenue #338, Sunnyvale, CA 94086 (US).			
(74) Agent: BILLINGS, Lucy, J.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).			

(54) Title: NOVEL HUMAN METALLOTHIONEIN

```

          9      18      27      36      45      54
5' MNC TCC CCT GAG TAG AAA AGC AGC CGC AGT TTG GNC GCT CCA CCA CGN CGT CCG

          63      72      81      90      99      108
GGT GGG CCT AGC AGT CGC TCC ATT TAT CGC TTG AGA TCT CCA GCC TTA CCG CGG

          117     126     135     144     153     162
CTC GAA ATG GAC CCC AAC TGC TCC TGC ACC ACT GGT GTC TCC TGC GCC TGC ACC
M   D   P   N   C   S   C   T   T   G   V   S   C   A   C   T

          171     180     189     198     207     216
GGC TCC TGC AAG TGC AAA GAG TGC AAA TGC ACC TYC TGC AAG AAG AGC TGC TGC
G   S   C   K   C   K   E   C   K   C   T   X   C   K   K   S   C   C

          225     234     243     252     261     270
TCC TGC TGC CCC GTG GGC TGT GCC AAG TGT GCC CAC GGC TGT GTC TGC AAA GGG
S   C   C   P   V   G   C   A   K   C   A   H   G   C   V   C   K   G

          279     288     297     306     315     324
ACG TTG GAG AAC TGC AGC TGC TGT GCC TGA TGT GGG AAC AGC TCT TCT CCC AGA
T   L   E   N   C   S   C   C   A

          333     342     351     360     369     378
TGT TAA TAG AAC AAG CTG CAC AAC CTG GAT TTT TTT TCA ATA CGA TAC TGA GCC
3'

```

(57) Abstract

The present invention provides a human metallothionein (HMBP-I) and polynucleotides which identify and encode HMBP-I. The invention also provides genetically engineered expression vectors and host cells comprising the nucleic acid sequences encoding HMBP-I and a method for producing HMBP-I. The invention also provides for agonists, antibodies, or antagonists specifically binding HMBP-I, and their use, in the prevention and treatment of diseases associated with expression of HMBP-I. Additionally, the invention provides for the use of antisense molecules to polynucleotides encoding HMBP-I for the treatment of diseases associated with the expression of HMBP-I. The invention also provides diagnostic assays which utilize the polynucleotide, or fragments or the complement thereof, and antibodies specifically binding HMBP-I.

NOVEL HUMAN METALLOTHIONEIN

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of a novel
5 metallothionein and to the use of these sequences in the diagnosis, prevention, and treatment of heavy metal toxicity, cancer, inflammatory disease and immune disorders.

BACKGROUND ART

The metallothioneins (MTs) are a group of small (61 amino acids), cysteine rich proteins that bind heavy metals such as cadmium, zinc, mercury, lead, and copper and are thought to play
10 a role in metal detoxification or the metabolism and homeostasis of metals. They are present in a wide variety of eucaryotes including invertebrates, vertebrates, plants, and fungi. The primary structure of the vertebrate MTs is strongly conserved between species, particularly the positions of the cysteine residues which serve to chelate heavy metal ions via thiolate complexes.

At least two closely related but chromatographically distinct isoforms of MT, MT-I and
15 MT-II, have been observed in every vertebrate species examined. All isoforms of MT sequenced thus far share 20 cysteine residues found in the same positions and lack aromatic amino acids or leucine (Schmidt, C.J. and Hamer, D.H. (1983) *Gene* 24:137-46; Schmidt, C.J. et al. (1985) *J. Biol Chem.* 260(12):7731-37). MT-II isoforms differ from MT-I only in the presence of an aspartate residue at position 10 or 11 while MT-I contains either glycine or valine at these
20 positions. Recently, a third class of MT, MT-0, has been discovered in human liver that is characterized by a negatively charged amino acid at position 8 (glutamate) and a lysine substitution at the highly conserved Glu₂₃ (Soumilion, A. Et al. (1992) *Eur. J. Biochem.* 209:999-1004). In humans, only a single sequence of MT-0 and MT-II have been found, while at least six isoforms of MT-I have been identified. These various isoforms of MT-I differ only by a
25 few residues distributed throughout the molecule. The localization and identification of 12 functional MT genes on human chromosome 16 indicates that several additional MT isoforms exist.

The reason for multiple human MT genes is unclear. It is possible that different isoforms are responsible for binding different metals, or that different control sequences for the
30 transcription of these various isoforms respond to different stimuli, such as heavy metals, glucocorticoids, or enterotoxin (Schmidt, et al. *supra*). A third possibility is that multiple isoforms could play different roles in metal homeostasis during development or in different tissues.

WO 98/31795

PCT/US97/24129

2

Acute or chronic exposure to heavy metals such as lead, arsenic, mercury or cadmium leads to a variety of diseases and disorders involving neuromuscular, CNS, cardiovascular, and gastrointestinal effects. MTs may play a role in the prevention or alleviation of these conditions. In addition, MTs are transcriptionally regulated by glucocorticoids which suggests that MTs have
5 a direct role in the effects of glucocorticoids to treat inflammatory disease, immune disorders, and cancer.

Discovery of new proteins related to metallothioneins, and the polynucleotides that encode them, satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in diagnosing and treating heavy metal toxicity, cancer, inflammatory disease and immune
10 disorders.

DISCLOSURE OF THE INVENTION

The present invention features a novel human metallothionein hereinafter designated HMBP-I (heavy metal binding protein) and characterized as having similarity to MT-I from both human and animal sources. Accordingly, the invention features a substantially purified HMBP-I
15 having the amino acid sequence shown in SEQ ID NO:1.

One aspect of the invention features isolated and substantially purified polynucleotides that encode HMBP-I. In a particular aspect, the polynucleotide is the nucleotide sequence of SEQ ID NO:2.

The invention also relates to a polynucleotide sequence comprising the complement of
20 SEQ ID NO:2 or variants thereof. In addition, the invention features polynucleotide sequences which hybridize under stringent conditions to SEQ ID NO:2.

The invention additionally features nucleic acid sequences encoding polypeptides, oligonucleotides, peptide nucleic acids (PNA), fragments, portions or antisense molecules thereof, and expression vectors and host cells comprising polynucleotides that encode HMBP-I.
25 The present invention also features antibodies which bind specifically to HMBP-I, and pharmaceutical compositions comprising substantially purified HMBP-I. The invention also features the use of agonists and antagonists of HMBP-I.

BRIEF DESCRIPTION OF DRAWINGS

Figure 1 shows the amino acid sequence (SEQ ID NO:1) and nucleic acid sequence (SEQ
30 ID NO:2) of HMBP-I. The alignment was produced using MacDNASIS PRO™ software (Hitachi Software Engineering Co., Ltd., San Bruno, CA).

Figure 2 shows the amino acid sequence alignments among HMBP-I (SEQ ID NO:1), and three metallothionein-I isoforms from man (hMT-Ib; GI 386962; SEQ ID NO:3 and hMT-Ie; GI

WO 98/31795

PCT/US97/24129

3

386865; SEQ ID NO:4), and monkey kidney cells (pmkMT1; GI 342385; SEQ ID NO:5), and one metallothionein-II isoform from rodent (chMT-II; GI 305363; SEQ ID NO:6). The alignment was produced using the multisequence alignment program of DNASTAR™ software (DNASTAR Inc, Madison WI).

5 Figures 3A, 3B, 3C, 3D, and 3E show the hydrophobicity plots (MacDNASIS PRO software) for HMBP-I, SEQ ID NO: 1; hMT-Ib, SEQ ID NO:3; hMT-Ie, SEQ ID NO:4; pmkMT1, SEQ ID NO:5; and chMT-II, SEQ ID NO:6, respectively. The positive X axis reflects amino acid position, and the negative Y axis, hydrophobicity.

MODES FOR CARRYING OUT THE INVENTION

10 Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

15 It must be noted that as used herein and in the appended claims, the singular forms “a”, “an”, and “the” include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to “a host cell” includes a plurality of such host cells, reference to the “antibody” is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

20 Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices, and materials are now described. All publications mentioned herein are incorporated herein by
25 reference for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are reported in the publications which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

30 “Nucleic acid sequence” as used herein refers to an oligonucleotide, nucleotide, or polynucleotide, and fragments or portions thereof, and to DNA or RNA of genomic or synthetic origin which may be single- or double-stranded, and represent the sense or antisense strand. Similarly, “amino acid sequence” as used herein refers to an oligopeptide, peptide, polypeptide,

WO 98/31795

PCT/US97/24129

4

or protein sequence, and fragments or portions thereof, and to naturally occurring or synthetic molecules.

Where "amino acid sequence" is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms, such as

5 "polypeptide" or "protein" are not meant to limit the amino acid sequence to the complete, native amino acid sequence associated with the recited protein molecule.

"Peptide nucleic acid", as used herein, refers to a molecule which comprises an oligomer to which an amino acid residue, such as lysine, and an amino group have been added. These small molecules, also designated anti-gene agents, stop transcript elongation by binding to their
10 complementary strand of nucleic acid (Nielsen, P.E. et al. (1993) Anticancer Drug Des. 8:53-63).

HMBP-I, as used herein, refers to the amino acid sequences of substantially purified HMBP-I obtained from any species, particularly mammalian, including bovine, ovine, porcine, murine, equine, and preferably human, from any source whether natural, synthetic, semi-synthetic, or recombinant.

15 "Consensus", as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, or which has been extended using XL-PCR™ (Perkin Elmer, Norwalk, CT) in the 5' and/or the 3' direction and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte clone using the GELVIEW™ Fragment Assembly system (GCG, Madison, WI), or which has been both extended and
20 assembled.

A "variant" of HMBP-I, as used herein, refers to an amino acid sequence that is altered by one or more amino acids. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine. More rarely, a variant may have "nonconservative" changes, e.g., replacement of a
25 glycine with a tryptophan. Similar minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, DNASTAR software.

A "deletion", as used herein, refers to a change in either amino acid or nucleotide
30 sequence in which one or more amino acid or nucleotide residues, respectively, are absent.

An "insertion" or "addition", as used herein, refers to a change in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid or nucleotide residues, respectively, as compared to the naturally occurring molecule.

5

A "substitution", as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

The term "biologically active", as used herein, refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise,

5 "immunologically active" refers to the capability of the natural, recombinant, or synthetic HMBP-I, or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The term "agonist", as used herein, refers to a molecule which, when bound to HMBP-I, causes a change in HMBP-I which modulates the activity of HMBP-I. Agonists may include
10 proteins, nucleic acids, carbohydrates, or any other molecules which bind to HMBP-I.

The terms "antagonist" or "inhibitor", as used herein, refer to a molecule which, when bound to HMBP-I, blocks or modulates the biological or immunological activity of HMBP-I. Antagonists and inhibitors may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to HMBP-I.

15 The term "modulate", as used herein, refers to a change or an alteration in the biological activity of HMBP-I. Modulation may be an increase or a decrease in protein activity, a change in binding characteristics, or any other change in the biological, functional or immunological properties of HMBP-I.

The term "mimetic", as used herein, refers to a molecule, the structure of which is
20 developed from knowledge of the structure of HMBP-I or portions thereof and, as such, is able to effect some or all of the actions of metalloprotein-like molecules.

The term "derivative", as used herein, refers to the chemical modification of a nucleic acid encoding HMBP-I or the encoded HMBP-I. Illustrative of such modifications would be replacement of hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative would
25 encode a polypeptide which retains essential biological characteristics of the natural molecule.

The term "substantially purified", as used herein, refers to nucleic or amino acid sequences that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% free from other components with which they are naturally associated.

30 "Amplification" as used herein refers to the production of additional copies of a nucleic acid sequence and is generally carried out using polymerase chain reaction (PCR) technologies well known in the art (Dieffenbach, C.W. and G.S. Dveksler (1995) PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainville, NY).

WO 98/31795

PCT/US97/24129

6

The term "hybridization", as used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

The term "hybridization complex", as used herein, refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary
5 G and C bases and between complementary A and T bases; these hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., membranes, filters, chips,
10 pins or glass slides to which cells have been fixed for in situ hybridization).

The terms "complementary" or "complementarity", as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, for the sequence "A-G-T" binds to the complementary sequence "T-C-A". Complementarity between two single-stranded molecules may be "partial", in which only some
15 of the nucleic acids bind, or it may be complete when total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands.

20 The term "homology", as used herein, refers to a degree of complementarity. There may be partial homology or complete homology (i.e., identity). A partially complementary sequence is one that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid; it is referred to using the functional term "substantially homologous." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined
25 using a hybridization assay (Southern or northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or probe will compete for and inhibit the binding (i.e., the hybridization) of a completely homologous sequence or probe to the target sequence under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that
30 the binding of two sequences to one another be a specific (i.e., selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% identity); in the absence of non-specific binding, the probe will not hybridize to the second non-complementary target

WO 98/31795

PCT/US97/24129

7

sequence.

As known in the art, numerous equivalent conditions may be employed to comprise either low or high stringency conditions. Factors such as the length and nature (DNA, RNA, base composition) of the sequence, nature of the target (DNA, RNA, base composition, presence in solution or immobilization, etc.), and the concentration of the salts and other components (e.g., the presence or absence of formamide, dextran sulfate and/or polyethylene glycol) are considered and the hybridization solution may be varied to generate conditions of either low or high stringency different from, but equivalent to, the above listed conditions.

The term "stringent conditions", as used herein, is the "stringency" which occurs within a range from about $T_m - 5^\circ\text{C}$ (5°C below the melting temperature (T_m) of the probe) to about 20°C to 25°C below T_m . As will be understood by those of skill in the art, the stringency of hybridization may be altered in order to identify or detect identical or related polynucleotide sequences.

The term "antisense", as used herein, refers to nucleotide sequences which are complementary to a specific DNA or RNA sequence. The term "antisense strand" is used in reference to a nucleic acid strand that is complementary to the "sense" strand. Antisense molecules may be produced by any method, including synthesis by ligating the gene(s) of interest in a reverse orientation to a viral promoter which permits the synthesis of a complementary strand. Once introduced into a cell, this transcribed strand combines with natural sequences produced by the cell to form duplexes. These duplexes then block either the further transcription or translation. In this manner, mutant phenotypes may be generated. The designation "negative" is sometimes used in reference to the antisense strand and "positive" is sometimes used in reference to the sense strand.

The term "portion", as used herein, with regard to a protein (as in "a portion of a given protein") refers to fragments of that protein. The fragments may range in size from four amino acid residues to the entire amino acid sequence minus one amino acid. Thus, a protein "comprising at least a portion of the amino acid sequence of SEQ ID NO:1" encompasses the full-length human HMBP-I and fragments thereof.

"Transformation", as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. It may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method is selected based on the host cell being transformed and may include, but is not limited to, viral

WO 98/31795

PCT/US97/24129

8

infection, electroporation, lipofection, and particle bombardment. Such "transformed" cells include stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome. They also include cells which transiently express the inserted DNA or RNA for limited periods of time.

5 The term "antigenic determinant", as used herein, refers to that portion of a molecule that makes contact with a particular antibody (i.e., an epitope). When a protein or fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to a given region or three-dimensional structure on the protein; these regions or structures are referred to as antigenic determinants. An antigenic
10 determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The terms "specific binding" or "specifically binding", as used herein, in reference to the interaction of an antibody and a protein or peptide, mean that the interaction is dependent upon the presence of a particular structure (i.e., the antigenic determinant or epitope) on the protein; in
15 other words, the antibody is recognizing and binding to a specific protein structure rather than to proteins in general. For example, if an antibody is specific for epitope "A", the presence of a protein containing epitope A (or free, unlabeled A) in a reaction containing labeled "A" and the antibody will reduce the amount of labeled A bound to the antibody.

The term "sample", as used herein, is used in its broadest sense. A biological sample
20 suspected of containing nucleic acid encoding HMBP-I or fragments thereof may comprise a cell, chromosomes isolated from a cell (e.g., a spread of metaphase chromosomes), genomic DNA (in solution or bound to a solid support such as for Southern analysis), RNA (in solution or bound to a solid support such as for northern analysis), cDNA (in solution or bound to a solid support), an extract from cells or a tissue, and the like.

25 The term "correlates with expression of a polynucleotide", as used herein, indicates that the detection of the presence of ribonucleic acid that is similar to SEQ ID NO:2 by northern analysis is indicative of the presence of mRNA encoding HMBP-I in a sample and thereby correlates with expression of the transcript from the polynucleotide encoding the protein.

"Alterations" in the polynucleotide of SEQ ID NO: 2, as used herein, comprise any
30 alteration in the sequence of polynucleotides encoding HMBP-I including deletions, insertions, and point mutations that may be detected using hybridization assays. Included within this definition is the detection of alterations to the genomic DNA sequence which encodes HMBP-I (e.g., by alterations in the pattern of restriction fragment length polymorphisms capable of

WO 98/31795

PCT/US97/24129

9

hybridizing to SEQ ID NO:2), the inability of a selected fragment of SEQ ID NO: 2 to hybridize to a sample of genomic DNA (e.g., using allele-specific oligonucleotide probes), and improper or unexpected hybridization, such as hybridization to a locus other than the normal chromosomal locus for the polynucleotide sequence encoding HMBP-I (e.g., using fluorescent in situ

5 hybridization [FISH] to metaphase chromosomes spreads).

As used herein, the term "antibody" refers to intact molecules as well as fragments thereof, such as Fa, F(ab')₂, and Fv, which are capable of binding the epitopic determinant. Antibodies that bind HMBP-I polypeptides can be prepared using intact polypeptides or fragments containing small peptides of interest as the immunizing antigen. The polypeptide or
10 peptide used to immunize an animal can be derived from the transition of RNA or synthesized chemically, and can be conjugated to a carrier protein, if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin and thyroglobulin. The coupled peptide is then used to immunize the animal (e.g., a mouse, a rat, or a rabbit).

The term "humanized antibody", as used herein, refers to antibody molecules in which
15 amino acids have been replaced in the non-antigen binding regions in order to more closely resemble a human antibody, while still retaining the original binding ability.

THE INVENTION

The invention is based on the discovery of a novel human metallothionein, (HMBP-I), the polynucleotides encoding HMBP-I, and the use of these compositions for the diagnosis,
20 prevention, or treatment of heavy metal toxicity, cancer, inflammatory disease and immune disorders.

Nucleic acids encoding the human HMBP-I of the present invention were first identified in Incyte Clone 977981 from the breast tissue cDNA library BRSTNOT02 through a computer-generated search for amino acid sequence alignments. A consensus sequence, SEQ ID
25 NO:2, was derived from the following overlapping and/or extended nucleic acid sequences: Incyte Clones 726198 (SYNOOAT01), 882343 (THYRNOT02), and 977891 (BRSTNOT02).

In one embodiment, the invention encompasses a polypeptide comprising the amino acid sequence of SEQ ID NO:1, as shown in Figure 1. HMBP-I is 61 amino acids in length and, as shown in Figure 2, has chemical and structural homology with human hMT-Ib (GI 386962; SEQ
30 ID NO:3), human hMT-Ie (GI 386865; SEQ ID NO:4), monkey kidney pmkMT1 (GI 342385; SEQ ID NO:5), and chinese hamster chMT-II (GI 305363; SEQ ID NO:6). In particular, HMBP-I shares 87%, 86%, 87%, and 82% identity with hMT-1b, hMT-1e, pmkMT1, and chMT-II, respectively. The 20 cysteine residues found in all knownMTs are conserved in

HMBP-1 as well. A strong preference for C or G in the third position of the triplet nucleotide codons for cysteine is also a characteristic of MT encoding sequences. This is apparent in the polynucleotide sequence for HMBP-I shown in Figure 1. HMBP-I also contains no aromatic amino acids, although it does contain a single leucine residue at position 54. Since HMBP-1
5 contains G₁₀, V₁₁ and G₂₃, HMBP-I appears to be an isoform of MT-I. As illustrated by Figures 3A, 3B, 3C, 3D, and 3E, HMBP-I and the four MTs shown have rather similar hydrophobicity plots. Northern analysis shows that this sequence is most prominently expressed in prostate (20/196), lung (14/196), and colon (14/196), but it is present in a wide variety of other normal and diseased tissues as well.

10 The invention also encompasses HMBP-I variants. A preferred HMBP-I variant is one having at least 80%, and more preferably 90%, amino acid sequence similarity to the HMBP-I amino acid sequence (SEQ ID NO:1). A most preferred HMBP-I variant is one having at least 95% amino acid sequence similarity to SEQ ID NO:1.

The invention also encompasses polynucleotides which encode HMBP-I. Accordingly,
15 any nucleic acid sequence which encodes the amino acid sequence of HMBP-I can be used to generate recombinant molecules which express HMBP-I. In a particular embodiment, the invention encompasses the polynucleotide comprising the nucleic acid sequence of SEQ ID NO:2 as shown in Figure 1.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the
20 genetic code, a multitude of nucleotide sequences encoding HMBP-I, some bearing minimal homology to the nucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the
25 nucleotide sequence of naturally occurring HMBP-I, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode HMBP-I and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring HMBP-I under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide
30 sequences encoding HMBP-I or its derivatives possessing a substantially different codon usage. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence

WO 98/31795

PCT/US97/24129

II

encoding HMBP-I and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences, or portions thereof, which encode HMBP-I and its derivatives, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents that are well known in the art at the time of the filing of this application. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding HMBP-I or any portion thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed nucleotide sequences, and in particular, those shown in SEQ ID NO:2, under various conditions of stringency. Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex or probe, as taught in Wahl, G.M. and S.L. Berger (1987; Methods Enzymol. 152:399-407) and Kimmel, A.R. (1987; Methods Enzymol. 152:507-511), and may be used at a defined stringency.

Altered nucleic acid sequences encoding HMBP-I which are encompassed by the invention include deletions, insertions, or substitutions of different nucleotides resulting in a polynucleotide that encodes the same or a functionally equivalent HMBP-I. The encoded protein may also contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent HMBP-I. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the biological activity of HMBP-I is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid; positively charged amino acids may include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; phenylalanine and tyrosine.

Also included within the scope of the present invention are alleles of the genes encoding HMBP-I. As used herein, an "allele" or "allelic sequence" is an alternative form of the gene which may result from at least one mutation in the nucleic acid sequence. Alleles may result in altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions, or substitutions of nucleotides.

WO 98/31795

PCT/US97/24129

12

Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

Methods for DNA sequencing which are well known and generally available in the art may be used to practice any embodiments of the invention. The methods may employ such
5 enzymes as the Klenow fragment of DNA polymerase I, Sequenase® (US Biochemical Corp, Cleveland, OH), Taq polymerase (Perkin Elmer), thermostable T7 polymerase (Amersham, Chicago, IL), or combinations of recombinant polymerases and proofreading exonucleases such as the ELONGASE Amplification System marketed by Gibco BRL (Gaithersburg, MD). Preferably, the process is automated with machines such as the Hamilton Micro Lab 2200
10 (Hamilton, Reno, NV), Peltier Thermal Cycler (PTC200; MJ Research, Watertown, MA) and the ABI 377 DNA sequencers (Perkin Elmer).

The nucleic acid sequences encoding HMBP-I may be extended utilizing a partial nucleotide sequence and employing various methods known in the art to detect upstream sequences such as promoters and regulatory elements. For example, one method which may be
15 employed, "restriction-site" PCR, uses universal primers to retrieve unknown sequence adjacent to a known locus (Sarkar, G. (1993) PCR Methods Applic. 2:318-322). In particular, genomic DNA is first amplified in the presence of primer to linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round
20 of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR may also be used to amplify or extend sequences using divergent primers based on a known region (Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186). The primers may be designed using OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth,
25 MN), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Another method which may be used is capture PCR which involves PCR amplification of
30 DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA (Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119). In this method, multiple restriction enzyme digestions and ligations may also be used to place an engineered double-stranded sequence into an unknown portion of the DNA molecule before performing

PCR.

Another method which may be used to retrieve unknown sequences is that of Parker, J.D. et al. (1991; Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PromoterFinder™ libraries to walk in genomic DNA (Clontech, Palo Alto, CA). This
5 process avoids the need to screen libraries and is useful in finding intron/exon junctions. When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Also, random-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a
10 full-length cDNA. Genomic libraries may be useful for extension of sequence into the 5' and 3' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation,
15 four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity may be converted to electrical signal using appropriate software (e.g. Genotyper™ and Sequence Navigator™, Perkin Elmer) and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially
20 preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode HMBP-I, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of HMBP-I in appropriate host cells. Due to
25 the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express HMBP-I.

As will be understood by those of skill in the art, it may be advantageous to produce HMBP-I-encoding nucleotide sequences possessing non-naturally occurring codons. For
30 example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter HMBP-I encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene
5 fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, or introduce mutations, and so forth.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding HMBP-I may be ligated to a heterologous sequence to encode a fusion
10 protein. For example, to screen peptide libraries for inhibitors of HMBP-I activity, it may be useful to encode a chimeric HMBP-I protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between the HMBP-I encoding sequence and the heterologous protein sequence, so that HMBP-I may be cleaved and purified away from the heterologous moiety.

15 In another embodiment, sequences encoding HMBP-I may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M.H. et al. (1980) Nucl. Acids Res. Symp. Ser. 215-223, Horn, T. et al. (1980) Nucl. Acids Res. Symp. Ser. 225-232). Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of HMBP-I, or a portion thereof. For example, peptide synthesis can be performed
20 using various solid-phase techniques (Roberge, J.Y. et al. (1995) Science 269:202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer).

The newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) Proteins, Structures and
25 Molecular Principles, WH Freeman and Co., New York, NY). The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; Creighton, supra). Additionally, the amino acid sequence of HMBP-I, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

30 In order to express a biologically active HMBP-I, the nucleotide sequences encoding HMBP-I or functional equivalents, may be inserted into appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence.

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding HMBP-I and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described in

5 Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, and Ausubel, F.M. et al. (1989) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY.

A variety of expression vector/host systems may be utilized to contain and express sequences encoding HMBP-I. These include, but are not limited to, microorganisms such as
10 bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems.

15 The "control elements" or "regulatory sequences" are those non-translated regions of the vector--enhancers, promoters, 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be
20 used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the Bluescript® phagemid (Stratagene, LaJolla, CA) or pSport1™ plasmid (Gibco BRL) and the like may be used. The baculovirus polyhedrin promoter may be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO; and storage protein genes) or from plant viruses (e.g., viral promoters or leader
25 sequences) may be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding HMBP-I, vectors based on SV40 or EBV may be used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected depending upon the
30 use intended for HMBP-I. For example, when large quantities of HMBP-I are needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional E. coli cloning and expression vectors such as Bluescript® (Stratagene), in which the sequence

encoding HMBP-I may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Promega, Madison, WI) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

10 In the yeast, Saccharomyces cerevisiae, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) Methods Enzymol. 153:516-544.

In cases where plant expression vectors are used, the expression of sequences encoding HMBP-I may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or Murry, L.E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, NY; pp. 191-196.

An insect system may also be used to express HMBP-I. For example, in one such system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in Spodoptera frugiperda cells or in Trichoplusia larvae. The sequences encoding HMBP-I may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of HMBP-I will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, S. frugiperda cells or Trichoplusia larvae in which HMBP-I may be expressed (Engelhard, E.K. et al. (1994) Proc. Nat. Acad. Sci. 91:3224-3227).

In mammalian host cells, a number of viral-based expression systems may be utilized. In

cases where an adenovirus is used as an expression vector, sequences encoding HMBP-I may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing HMBP-I in infected host cells

5 (Logan, J. and Shenk, T. (1984) *Proc. Natl. Acad. Sci.* 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding HMBP-I. Such signals include the ATG initiation codon and adjacent
10 sequences. In cases where sequences encoding HMBP-I, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading
15 frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162).

20 In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding
25 and/or function. Different host cells such as CHO, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express HMBP-I may be transformed using
30 expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance

to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These
5 include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) Cell 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1980) Cell 22:817-23) genes which can be employed in tk⁻ or apt⁻ cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. 77:3567-70);
10 npt, which confers resistance to the aminoglycosides neomycin and G-418 (Colbere-Garapin, F. et al (1981) J. Mol. Biol. 150:1-14) and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, supra). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman, S.C. and R.C. Mulligan
15 (1988) Proc. Natl. Acad. Sci. 85:8047-51). Recently, the use of visible markers has gained popularity with such markers as anthocyanins, β glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C.A. et al. (1995) Methods Mol. Biol. 55:121-131).

20 Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding HMBP-I is inserted within a marker gene sequence, recombinant cells containing sequences encoding HMBP-I can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding
25 HMBP-I under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells which contain the nucleic acid sequence encoding HMBP-I and express HMBP-I may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and
30 protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

The presence of polynucleotide sequences encoding HMBP-I can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or portions or fragments of

polynucleotides encoding HMBP-I. Nucleic acid amplification based assays involve the use of oligonucleotides or oligomers based on the sequences encoding HMBP-I to detect transformants containing DNA or RNA encoding HMBP-I. As used herein "oligonucleotides" or "oligomers" refer to a nucleic acid sequence of at least about 10 nucleotides and as many as about 60

5 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20-25 nucleotides, which can be used as a probe or amplimer.

A variety of protocols for detecting and measuring the expression of HMBP-I, using either polyclonal or monoclonal antibodies specific for the protein are known in the art.

Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and
10 fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on HMBP-I is preferred, but a competitive binding assay may be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990; Serological Methods, a Laboratory Manual, APS Press, St Paul, MN) and Maddox, D.E. et al. (1983; J. Exp. Med. 158:1211-1216).

15 A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding HMBP-I include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding HMBP-I, or any portions thereof may
20 be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits (Pharmacia & Upjohn, (Kalamazoo, MI); Promega (Madison WI); and U.S. Biochemical Corp., Cleveland, OH).

25 Suitable reporter molecules or labels, which may be used, include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding HMBP-I may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The
30 protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode HMBP-I may be designed to contain signal sequences which direct secretion of HMBP-I through a prokaryotic or eukaryotic cell membrane.

Other recombinant constructions may be used to join sequences encoding HMBP-I to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the 5 FLAGS extension/affinity purification system (Immunex Corp., Seattle, WA). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and HMBP-I may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing HMBP-I and a 10 nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography as described in Porath, J. et al. (1992, Prot. Exp. Purif. 3: 263-281) while the enterokinase cleavage site provides a means for purifying HMBP-I from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D.J. et al. (1993; DNA 15 Cell Biol. 12:441-453).

In addition to recombinant production, fragments of HMBP-I may be produced by direct peptide synthesis using solid-phase techniques Merrifield J. (1963) J. Am. Chem. Soc. 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide 20 Synthesizer (Perkin Elmer). Various fragments of HMBP-I may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

THERAPEUTICS

Chemical and structural homology exists among HMBP-I and human, monkey, and rodent MTs. In addition, HMBP-I is expressed in a variety of both normal and diseased tissues. Thus 25 HMBP-I appears to play a role in the diagnosis or treatment of heavy metal toxicity, inflammatory disease, immune disorders, and cancer.

Therefore, in one embodiment, HMBP-I or a fragment or derivative thereof may be administered to a subject to treat heavy metal toxicity. Such conditions and diseases may include, but are not limited to, myopathy, encephalopathy, renal nephropathy or necrosis, liver 30 necrosis or cirrhosis, anemia, myocardial damage, and pneumonitis or any other condition or disease caused by exposure to heavy metals. Heavy metals may include, but are not limited to, lead, arsenic, mercury, cadmium and copper.

In another embodiment, HMBP-I or a fragment or derivative thereof may be administered

to a subject to treat immune disorders. Such conditions and diseases may include, but are not limited to, bronchial asthma, chronic obstructive pulmonary disease, pneumonia, multiple sclerosis, rheumatoid arthritis, inflammatory bowel disease, chronic hepatitis, and cerebral edema.

5 In another embodiment, HMBP-I or a fragment or derivative thereof may be administered to a subject to treat cancer. Cancers may include, but are not limited to, leukemias, lymphomas, and cancers of the liver, lung, prostate, ovaries, pancreas, stomach, bladder, gall bladder, and breast.

In another embodiment, a vector capable of expressing HMBP-I, or a fragment or a
10 derivative thereof, may also be administered to a subject to treat the conditions or diseases resulting from heavy metal toxicity which are listed above.

In another embodiment, a vector capable of expressing HMBP-I or a fragment or derivative thereof may be administered to a subject to treat immune disorders which are listed above.

15 In another embodiment, a vector capable of expressing HMBP-I or a fragment or derivative thereof may be administered to a subject to treat cancers which are listed above.

Antibodies which are specific for HMBP-I may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express HMBP-I.

20 In other embodiments, HMBP-I or a vector expressing HMBP-I may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above.

25 Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

There may be situations in which it would be desirable to use antagonists or inhibitors of HMBP-I for clinical purposes. These antagonists or inhibitors may be produced using methods which are generally known in the art. In particular, purified HMBP-I may be used to produce
30 antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind HMBP-I.

Antibodies to HMBP-I may be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single

chain, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies, (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others, may be immunized by injection with HMBP-I or any fragment or
 5 oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli
 10 Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the peptides, fragments, or oligopeptides used to induce antibodies to HMBP-I have an amino acid sequence consisting of at least five amino acids, and more preferably at least 10 amino acids. It is also preferable that they are identical to a portion of the amino acid sequence of the natural protein, and they may contain the entire amino acid sequence
 15 of a small, naturally occurring molecule. Short stretches of HMBP-I amino acids may be fused with those of another protein such as keyhole limpet hemocyanin and antibody produced against the chimeric molecule.

Monoclonal antibodies to HMBP-I may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but
 20 are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique (Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. 80:2026-2030; Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120).

In addition, techniques developed for the production of "chimeric antibodies", the splicing
 25 of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used (Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; Takeda, S. et al. (1985) Nature 314:452-454). Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce HMBP-I-specific single chain
 30 antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton D.R. (1991) Proc. Natl. Acad. Sci. 88:11120-3).

Antibodies may also be produced by inducing in vivo production in the lymphocyte

23

population or by screening recombinant immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature (Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. 86: 3833-3837; Winter, G. et al. (1991) Nature 349:293-299).

Antibody fragments which contain specific binding sites for HMBP-I may also be
5 generated. For example, such fragments include, but are not limited to, the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse, W.D. et al. (1989) Science 254:1275-1281).

10 Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between HMBP-I and its specific antibody. A two-site, monoclonal-based immunoassay utilizing
15 monoclonal antibodies reactive to two non-interfering HMBP-I epitopes is preferred, but a competitive binding assay may also be employed (Maddox, supra).

In another embodiment of the invention, the polynucleotides encoding HMBP-I, or any fragment thereof, or antisense molecules, may be used for therapeutic purposes. In one aspect, antisense to the polynucleotide encoding HMBP-I may be used in situations in which it would be
20 desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding HMBP-I. Thus, antisense molecules may be used to modulate HMBP-I activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligomers or larger fragments, can be designed from various locations along the coding or control regions of sequences encoding
25 HMBP-I.

Expression vectors derived from retro viruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids may be used for delivery of nucleotide sequences to the targeted organ, tissue or cell population. Methods which are well known to those skilled in the art can be used to construct recombinant vectors which will express antisense molecules complementary to
30 the polynucleotides of the gene encoding HMBP-I. These techniques are described both in Sambrook et al. (supra) and in Ausubel et al. (supra).

Genes encoding HMBP-I can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide or fragment thereof which

encodes HMBP-I. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases.

Transient expression may last for a month or more with a non-replicating vector and even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing antisense molecules, DNA, RNA, or PNA, to the control regions of the gene encoding HMBP-I, i.e., the promoters, enhancers, and introns. Oligonucleotides derived from the transcription initiation site, e.g., between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature (Gee, J.E. et al. (1994) In: Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, NY). The antisense molecules may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples which may be used include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding HMBP-I.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Antisense molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription

of DNA sequences encoding HMBP-I. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize antisense RNA constitutively or inducibly can be introduced into cell lines, cells, or tissues.

5 RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to the addition of flanking sequences at the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such
10 as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced
15 into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection and by liposome injections may be achieved using methods which are well known in the art.

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits,
20 monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of HMBP-I, antibodies to HMBP-I, mimetics, agonists, antagonists, or inhibitors of HMBP-I. The
25 compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

30 The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution,

Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of HMBP-I, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example HMBP-I or fragments thereof, antibodies of HMBP-I, agonists, antagonists or inhibitors of

HMBP-I, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from 0.1 to 10,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

25 DIAGNOSTICS

In another embodiment, antibodies which specifically bind HMBP-I may be used for the diagnosis of conditions or diseases characterized by expression of HMBP-I, or in assays to monitor patients being treated with HMBP-I, agonists, antagonists or inhibitors. The antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for HMBP-I include methods which utilize the antibody and a label to detect HMBP-I in human body fluids or extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by joining them, either covalently or non-covalently, with a reporter molecule. A wide variety of reporter molecules which are known

in the art may be used, several of which are described above.

A variety of protocols including ELISA, RIA, and FACS for measuring HMBP-I are known in the art and provide a basis for diagnosing altered or abnormal levels of HMBP-I expression. Normal or standard values for HMBP-I expression are established by combining
5 body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to HMBP-I under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, but preferably by photometric means. Quantities of HMBP-I expressed in subject, control and disease, samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values
10 establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding HMBP-I may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, antisense RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of HMBP-I may be
15 correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of HMBP-I, and to monitor regulation of HMBP-I levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding HMBP-I or closely related
20 molecules, may be used to identify nucleic acid sequences which encode HMBP-I. The specificity of the probe, whether it is made from a highly specific region, e.g., 10 unique nucleotides in the 5' regulatory region, or a less specific region, e.g., especially in the 3' coding region, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low) will determine whether the probe identifies only naturally occurring sequences encoding
25 HMBP-I, alleles, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably contain at least 50% of the nucleotides from any of the HMBP-I encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and derived from the nucleotide sequence of SEQ ID NO:2 or from genomic sequence including promoter, enhancer
30 elements, and introns of the naturally occurring HMBP-I.

Means for producing specific hybridization probes for DNAs encoding HMBP-I include the cloning of nucleic acid sequences encoding HMBP-I or HMBP-I derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, commercially available, and

WO 98/31795

PCT/US97/24129

30

may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, radionuclides such as ^{32}P or ^{35}S , or enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding HMBP-I may be used for the diagnosis of conditions or diseases which are associated with expression of HMBP-I. Examples of such conditions or diseases include heavy metal toxicities such as myopathy, encephalopathy, renal nephropathy or necrosis, liver necrosis or cirrhosis; immune disorders such as bronchial asthma, chronic obstructive pulmonary disease, and pneumonia; and cancer such as leukemias, lymphomas, and cancers of the liver, lung, and prostate. The polynucleotide sequences encoding HMBP-I may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; or in dip stick, pin, ELISA or chip assays utilizing fluids or tissues from patient biopsies to detect altered HMBP-I expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding HMBP-I may be useful in assays that detect activation or induction of various cancers, particularly those mentioned above. The nucleotide sequences encoding HMBP-I may be labeled by standard methods, and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the biopsied or extracted sample is significantly altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding HMBP-I in the sample indicates the presence of the associated disease. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

In order to provide a basis for the diagnosis of disease associated with expression of HMBP-I, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, which encodes HMBP-I, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with those from an experiment where a known amount of a

WO 98/31795

PCT/US97/24129

31

substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for disease. Deviation between standard and subject values is used to establish the presence of disease.

Once disease is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to evaluate whether the level of expression in the patient begins to approximate that which is observed in the normal patient. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding HMBP-I may involve the use of PCR. Such oligomers may be chemically synthesized, generated enzymatically, or produced from a recombinant source. Oligomers will preferably consist of two nucleotide sequences, one with sense orientation (5'→3') and another with antisense (3'←5'), employed under optimized conditions for identification of a specific gene or condition. The same two oligomers, nested sets of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for detection and/or quantitation of closely related DNA or RNA sequences.

Methods which may also be used to quantitate the expression of HMBP-I include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated (Melby, P.C. et al. (1993) J. Immunol. Methods, 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 229-236). The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In another embodiment of the invention, the nucleic acid sequences which encode HMBP-I may also be used to generate hybridization probes which are useful for mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome or to a specific region of the chromosome using well known techniques. Such

32.

techniques include FISH, FACS, or artificial chromosome constructions, such as yeast artificial chromosomes, bacterial artificial chromosomes, bacterial P1 constructions or single chromosome cDNA libraries as reviewed in Price, C.M. (1993) *Blood Rev.* 7:127-134, and Trask, B.J. (1991) *Trends Genet.* 7:149-154.

5 FISH (as described in Verma et al. (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York, NY) may be correlated with other physical chromosome mapping techniques and genetic map data. Examples of genetic map data can be found in the 1994 Genome Issue of *Science* (265:1981f). Correlation between the location of the gene encoding HMBP-I on a physical chromosomal map and a specific disease, or predisposition to a
10 specific disease, may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier, or affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques such as linkage analysis using established chromosomal markers may be used for extending genetic
15 maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms, or parts thereof, by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or
20 syndrome has been crudely localized by genetic linkage to a particular genomic region, for example, AT to 11q22-23 (Gatti, R.A. et al. (1988) *Nature* 336:577-580), any sequences mapping to that area may represent associated or regulatory genes for further investigation. The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. among normal, carrier, or affected individuals.

25 In another embodiment of the invention, HMBP-I, its catalytic or immunogenic fragments or oligopeptides thereof, can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes, between HMBP-I and the agent being tested, may be measured.

30 Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, as applied to HMBP-I large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or

33

some other surface. The test compounds are reacted with HMBP-I, or fragments thereof, and washed. Bound HMBP-I is then detected by methods well known in the art. Purified HMBP-I can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding HMBP-I specifically compete with a test compound for binding HMBP-I. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with HMBP-I.

In additional embodiments, the nucleotide sequences which encode HMBP-I may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

The examples below are provided to illustrate the subject invention and are not included for the purpose of limiting the invention.

INDUSTRIAL APPLICABILITY

I BRSTNOT01 cDNA Library Construction

The breast (BRSTNOT02) cDNA library was constructed from breast tissue excised from a 55-year-old Caucasian female (specimen #0005, Mayo Clinic, Rochester, MN) during unilateral extended simple mastectomy. The patient was diagnosed with invasive grade 4 mammary adenocarcinoma of mixed lobular and ductal type which involved all 4 quadrants of the left breast. Twelve axillary lymph nodes were found positive for tumor. Prior to surgery, the patient was diagnosed with cerebrovascular disease, atherosclerosis, paroxysmal supraventricular tachycardia, and gastrointestinal hemorrhage. The patient had a history of alcohol and tobacco abuse.

The frozen tissue was homogenized and lysed using a Brinkmann Homogenizer Polytron PT-3000 (Brinkmann Instruments, Westbury, NJ) in guanidinium isothiocyanate solution. The lysate was centrifuged over a 5.7 M CsCl cushion using an Beckman SW28 rotor in a Beckman L8-70M Ultracentrifuge (Beckman Instruments) for 18 hours at 25,000 rpm at ambient temperature. The RNA was extracted with acid phenol pH 4.7, precipitated using 0.3 M sodium acetate and 2.5 volumes of ethanol, resuspended in RNase-free water, and DNase treated at 37°C. Extraction and precipitation were repeated as before. The mRNA was isolated with the Qiagen Oligotex kit (QIAGEN, Inc.; Chatsworth, CA) and used to construct the cDNA library.

The mRNA was handled according to the recommended protocols in the SuperScript Plasmid System for cDNA Synthesis and Plasmid Cloning (Cat. #18248-013; Gibco/BRL, Gaithersburg, MD). cDNAs were fractionated on a Sepharose CL4B column (Cat. #275105-01; Pharmacia), and those cDNAs exceeding 400 bp were ligated into pSport I. The plasmid pSport I was subsequently transformed into DH12S competent cells (Cat. #18312-017; Gibco/BRL).

II Isolation and Sequencing of cDNA Clones

Plasmid DNA was released from the cells and purified using the REAL Prep 96 Plasmid Kit (Catalog #26173; QIAGEN, Inc.). This kit enabled the simultaneous purification of 96 samples in a 96-well block using multi-channel reagent dispensers. The recommended protocol was employed except for the following changes: 1) the bacteria were cultured in 1 ml of sterile Terrific Broth (Catalog #22711, LIFE TECHNOLOGIES™, Gaithersburg, MD) with carbenicillin at 25 mg/L and glycerol at 0.4%; 2) after inoculation, the cultures were incubated for 19 hours and at the end of incubation, the cells were lysed with 0.3 ml of lysis buffer; and 3) following isopropanol precipitation, the plasmid DNA pellet was resuspended in 0.1 ml of distilled water. After the last step in the protocol, samples were transferred to a 96-well block for storage at 4° C.

The cDNAs were sequenced by the method of Sanger et al. (1975, J. Mol. Biol. 94:441f), using a Hamilton Micro Lab 2200 (Hamilton, Reno, NV) in combination with Peltier Thermal Cyclers (PTC200 from MJ Research, Watertown, MA) and Applied Biosystems 377 DNA Sequencing Systems. Reading frames were determined.

III Homology Searching of cDNA Clones and Their Deduced Proteins

Each cDNA was compared to sequences in GenBank using a search algorithm developed by Applied Biosystems and incorporated into the INHERIT™ 670 sequence analysis system. In this algorithm, Pattern Specification Language (TRW Inc, Los Angeles, CA) was used to determine regions of homology. The three parameters that determine how the sequence comparisons run were window size, window offset, and error tolerance. Using a combination of these three parameters, the DNA database was searched for sequences containing regions of homology to the query sequence, and the appropriate sequences were scored with an initial value. Subsequently, these homologous regions were examined using dot matrix homology plots to distinguish regions of homology from chance matches. Smith-Waterman alignments were used to display the results of the homology search.

Peptide and protein sequence homologies were ascertained using the INHERIT- 670 sequence analysis system using the methods similar to those used in DNA sequence homologies.

35

Pattern Specification Language and parameter windows were used to search protein databases for sequences containing regions of homology which were scored with an initial value. Dot-matrix homology plots were examined to distinguish regions of significant homology from chance matches.

5 BLAST, which stands for Basic Local Alignment Search Tool (Altschul, S.F. (1993) J. Mol. Evol. 36:290-300; Altschul et al. (1990) J. Mol. Biol. 215:403-410), was used to search for local sequence alignments. BLAST produces alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST is especially useful in determining exact matches or in identifying homologs. BLAST is
10 useful for matches which do not contain gaps. The fundamental unit of BLAST algorithm output is the High-scoring Segment Pair (HSP).

An HSP consists of two sequence fragments of arbitrary but equal lengths whose alignment is locally maximal and for which the alignment score meets or exceeds a threshold or cutoff score set by the user. The BLAST approach is to look for HSPs between a query sequence
15 and a database sequence, to evaluate the statistical significance of any matches found, and to report only those matches which satisfy the user-selected threshold of significance. The parameter E establishes the statistically significant threshold for reporting database sequence matches. E is interpreted as the upper bound of the expected frequency of chance occurrence of an HSP (or set of HSPs) within the context of the entire database search. Any database sequence
20 whose match satisfies E is reported in the program output.

IV Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound (Sambrook et al., supra).

25 Analogous computer techniques using BLAST (Altschul, S.F. 1993 and 1990, supra) are used to search for identical or related molecules in nucleotide databases such as GenBank or the LIFESEQ™ database (Incyte Pharmaceuticals). This analysis is much faster than multiple, membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or homologous.

30 The basis of the search is the product score which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

100

The product score takes into account both the degree of similarity between two sequences and the

36

length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error; and at 70, the match will be exact. Homologous molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

5 The results of northern analysis are reported as a list of libraries in which the transcript encoding HMBP-I occurs. Abundance and percent abundance are also reported. Abundance directly reflects the number of times a particular transcript is represented in a cDNA library, and percent abundance is abundance divided by the total number of sequences examined in the cDNA library.

10 V Extension of HMBP-I-Encoding Polynucleotides to Full Length or to Recover Regulatory Sequences

Full length HMBP-I-encoding nucleic acid sequence (SEQ ID NO:2) is used to design oligonucleotide primers for extending a partial nucleotide sequence to full length or for obtaining 5' or 3', intron or other control sequences from genomic libraries. One primer is synthesized to
15 initiate extension in the antisense direction (XLR) and the other is synthesized to extend sequence in the sense direction (XLF). Primers are used to facilitate the extension of the known sequence "outward" generating amplicons containing new, unknown nucleotide sequence for the region of interest. The initial primers are designed from the cDNA using OLIGO 4.06 (National Biosciences), or another appropriate program, to be 22-30 nucleotides in length, to have a GC
20 content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations is avoided.

The original, selected cDNA libraries, or a human genomic library are used to extend the sequence; the latter is most useful to obtain 5' upstream regions. If more extension is necessary
25 or desired, additional sets of primers are designed to further extend the known region.

By following the instructions for the XL-PCR kit (Perkin Elmer) and thoroughly mixing the enzyme and reaction mix, high fidelity amplification is obtained. Beginning with 40 pmol of each primer and the recommended concentrations of all other components of the kit, PCR is performed using the Peltier Thermal Cycler (PTC200; M.J. Research, Watertown, MA) and the
30 following parameters:

Step 1	94° C for 1 min (initial denaturation)
Step 2	65° C for 1 min
Step 3	68° C for 6 min
Step 4	94° C for 15 sec

37

	Step 5	65° C for 1 min
	Step 6	68° C for 7 min
	Step 7	Repeat step 4-6 for 15 additional cycles
	Step 8	94° C for 15 sec
5	Step 9	65° C for 1 min
	Step 10	68° C for 7:15 min
	Step 11	Repeat step 8-10 for 12 cycles
	Step 12	72° C for 8 min
	Step 13	4° C (and holding)

10

A 5-10 μ l aliquot of the reaction mixture is analyzed by electrophoresis on a low concentration (about 0.6-0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products are selected and removed from the gel. Further purification involves using a commercial gel extraction method such as

15 QIAQuick™ (QIAGEN Inc., Chatsworth, CA). After recovery of the DNA, Klenow enzyme is used to trim single-stranded, nucleotide overhangs creating blunt ends which facilitate religation and cloning.

After ethanol precipitation, the products are redissolved in 13 μ l of ligation buffer, 1 μ l T4-DNA ligase (15 units) and 1 μ l T4 polynucleotide kinase are added, and the mixture is

20 incubated at room temperature for 2-3 hours or overnight at 16° C. Competent *E. coli* cells (in 40 μ l of appropriate media) are transformed with 3 μ l of ligation mixture and cultured in 80 μ l of SOC medium (Sambrook et al., supra). After incubation for one hour at 37° C, the whole transformation mixture is plated on Luria Bertani (LB)-agar (Sambrook et al., supra) containing 2x Carb. The following day, several colonies are randomly picked from each plate and cultured

25 in 150 μ l of liquid LB/2x Carb medium placed in an individual well of an appropriate, commercially-available, sterile 96-well microtiter plate. The following day, 5 μ l of each overnight culture is transferred into a non-sterile 96-well plate and after dilution 1:10 with water, 5 μ l of each sample is transferred into a PCR array.

For PCR amplification, 18 μ l of concentrated PCR reaction mix (3.3x) containing 4 units

30 of rTth DNA polymerase, a vector primer, and one or both of the gene specific primers used for the extension reaction are added to each well. Amplification is performed using the following conditions:

	Step 1	94° C for 60 sec
	Step 2	94° C for 20 sec
35	Step 3	55° C for 30 sec
	Step 4	72° C for 90 sec
	Step 5	Repeat steps 2-4 for an additional 29 cycles
	Step 6	72° C for 180 sec

Step 7 4° C (and holding) **38**

Aliquots of the PCR reactions are run on agarose gels together with molecular weight markers. The sizes of the PCR products are compared to the original partial cDNAs, and
5 appropriate clones are selected, ligated into plasmid, and sequenced.

VI Labeling and Use of Hybridization Probes

Hybridization probes derived from SEQ ID NO:2 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA
10 fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 (National Biosciences), labeled by combining 50 pmol of each oligomer and 250 &Ci of [³²P] adenosine triphosphate (Amersham) and T4 polynucleotide kinase (DuPont NEN®, Boston, MA). The labeled oligonucleotides are substantially purified with Sephadex G-25 superfine resin column (Pharmacia & Upjohn). A portion containing 10⁷ counts per minute of each of the sense
15 and antisense oligonucleotides is used in a typical membrane based hybridization analysis of human genomic DNA digested with one of the following endonucleases (Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II; DuPont NEN®).

The DNA from each digest is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham, NH). Hybridization is carried
20 out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR™ film (Kodak, Rochester, NY) is exposed to the blots in a Phosphorimager cassette (Molecular Dynamics, Sunnyvale, CA) for several hours, hybridization patterns are compared visually.

25 VII Antisense Molecules

Antisense molecules to the HMBP-I-encoding sequence, or any part thereof, is used to inhibit in vivo or in vitro expression of naturally occurring HMBP-I. Although use of antisense oligonucleotides, comprising about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA fragments. An oligonucleotide based on the coding
30 sequences of HMBP-I, as shown in Figure 1, is used to inhibit expression of naturally occurring HMBP-I. The complementary oligonucleotide is designed from the most unique 5' sequence as shown in Figure 1 and used either to inhibit transcription by preventing promoter binding to the upstream nontranslated sequence or translation of an HMBP-I-encoding transcript by preventing

the ribosome from binding. Using an appropriate portion of the signal and 5' sequence of SEQ ID NO:2, an effective antisense oligonucleotide includes any 15-20 nucleotides spanning the region which translates into the signal or 5' coding sequence of the polypeptide as shown in Figure 1.

5 VIII Expression of HMBP-I

Expression of HMBP-I is accomplished by subcloning the cDNAs into appropriate vectors and transforming the vectors into host cells. In this case, the cloning vector, pSport, previously used for the generation of the cDNA library is used to express HMBP-I in *E. coli*. Upstream of the cloning site, this vector contains a promoter for β -galactosidase, followed by
10 sequence containing the amino-terminal Met, and the subsequent seven residues of β -galactosidase. Immediately following these eight residues is a bacteriophage promoter useful for transcription and a linker containing a number of unique restriction sites.

Induction of an isolated, transformed bacterial strain with IPTG using standard methods produces a fusion protein which consists of the first eight residues of β -galactosidase, about 5 to
15 15 residues of linker, and the full length protein. The signal residues direct the secretion of HMBP-I into the bacterial growth media which can be used directly in the following assay for activity.

IX Demonstration of HMBP-I Activity

HMBP-I is measured by its ability to confer resistance to heavy metal toxicity when
20 expressed in a heavy metal sensitive cell line. Mouse C127 cells, which are sensitive to high concentrations of cadmium and will not grow in the presence of 10 μ M CdCl₂, are first transfected with the HMBP-I gene using bovine papilloma virus as the vector (Schmidt, et al., supra). The cells are then grown in the presence of 0-50 μ M CdCl₂ and compared with untransformed cells. The rate of cell growth in the presence of varying Cd concentrations is a
25 measure of the level of HMBP-I present in the transformed cells.

X Production of HMBP-I Specific Antibodies

HMBP-I that is substantially purified using PAGE electrophoresis (Sambrook, supra), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols. The amino acid sequence deduced from SEQ ID NO:2 is analyzed using
30 DNASTAR software (DNASTAR Inc) to determine regions of high immunogenicity and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions, is described by Ausubel et al. (supra), and others.

Typically, the oligopeptides are 15 residues in length, synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry, and coupled to keyhole limpet hemocyanin (KLH, Sigma, St. Louis, MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS; Ausubel et al., supra). Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity, for example, by binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radioiodinated, goat anti-rabbit IgG.

XI Purification of Naturally Occurring HMBP-I Using Specific Antibodies

Naturally occurring or recombinant HMBP-I is substantially purified by immunoaffinity chromatography using antibodies specific for HMBP-I. An immunoaffinity column is constructed by covalently coupling HMBP-I antibody to an activated chromatographic resin, such as CnBr-activated Sepharose (Pharmacia & Upjohn). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing HMBP-I is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of HMBP-I (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/HMBP-I binding (eg, a buffer of pH 2-3 or a high concentration of a chaotrope, such as urea or thiocyanate ion), and HMBP-I is collected.

XII Identification of Molecules Which Interact with HMBP-I

HMBP-I or biologically active fragments thereof are labeled with ¹²⁵I Bolton-Hunter reagent (Bolton et al. (1973) Biochem. J. 133: 529). Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled HMBP-I, washed and any wells with labeled HMBP-I complex are assayed. Data obtained using different concentrations of HMBP-I are used to calculate values for the number, affinity, and association of HMBP-I with the candidate molecules.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are

WO 98/31795

PCT/US97/24129

41

intended to be within the scope of the following claims.

WO 98/31795

PCT/US97/24129

42

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: INCYTE PHARMACEUTICALS, INC.
- (ii) TITLE OF THE INVENTION: A NOVEL HUMAN METALLOTHIONINE
- (iii) NUMBER OF SEQUENCES: 6
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Incyte Pharmaceuticals, Inc.
 - (B) STREET: 3174 Porter Drive
 - (C) CITY: Palo Alto
 - (D) STATE: CA
 - (E) COUNTRY: USA
 - (F) ZIP: 94304
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ for Windows Version 2.0
- (vi) CURRENT APPLICATION DATA:
 - (A) PCT APPLICATION NUMBER: To Be Assigned
 - (B) FILING DATE: Herewith
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 08/785,530
 - (B) FILING DATE: 17-JAN-1997
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Billings, Lucy J.
 - (B) REGISTRATION NUMBER: 36,749
 - (C) REFERENCE/DOCKET NUMBER: PF-0194 PCT
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 650-855-0555
 - (B) TELEFAX: 650-845-4166
 - (C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 61 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (vii) IMMEDIATE SOURCE:
 - (A) LIBRARY: Consensus
 - (B) CLONE: 977891
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

WO 98/31795

PCT/US97/24129

43

Met	Asp	Pro	Asn	Cys	Ser	Cys	Thr	Thr	Gly	Val	Ser	Cys	Ala	Cys	Thr
1				5					10					15	
Gly	Ser	Cys	Lys	Cys	Lys	Glu	Cys	Lys	Cys	Thr	Xaa	Cys	Lys	Lys	Ser
			20					25					30		
Cys	Cys	Ser	Cys	Cys	Pro	Val	Gly	Cys	Ala	Lys	Cys	Ala	His	Gly	Cys
		35					40					45			
Val	Cys	Lys	Gly	Thr	Leu	Glu	Asn	Cys	Ser	Cys	Cys	Ala			
	50					55					60				

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 376 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Consensus
- (B) CLONE: 977891

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CTCCCCTGAG	TAGAAAAGCA	GCCGCAGTTT	GGNCGCTCCA	CCACGNCGTC	CGGGTGGGCC	60
TAGCAGTCGC	TCCATTTATC	GCTTGAGATC	TCCAGCCTTA	CCGCGGCTCG	AAATGGACCC	120
CAACTGCTCC	TGCACCACTG	GTGTCTCCTG	CGCCTGCACC	GGCTCCTGCA	AGTGCAAAGA	180
GTGCAAATGC	ACCTYCTGCA	AGAAGAGCTG	CTGCTCCTGC	TGCCCCGTGG	GCTGTGCCAA	240
GTGTGCCCCAC	GGCTGTGTCT	GCAAAGGGAC	GTTGGAGAAC	TGCAGCTGCT	GTGCCTGATG	300
TGGGAACAGC	TCTTCTCCCA	GATGTTAATA	GAACAAGCTG	CACAACCTGG	ATTTTTTTTC	360
AATACGATAC	TGAGCC					376

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 61 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: GenBank
- (B) CLONE: 386962

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met	Asp	Pro	Asn	Cys	Ser	Cys	Thr	Thr	Gly	Gly	Ser	Cys	Ala	Cys	Ala
1				5					10					15	
Gly	Ser	Cys	Lys	Cys	Lys	Glu	Cys	Lys	Cys	Thr	Ser	Cys	Lys	Lys	Cys
			20					25					30		
Cys	Cys	Ser	Cys	Cys	Pro	Val	Gly	Cys	Ala	Lys	Cys	Ala	Gln	Gly	Cys
		35					40					45			
Val	Cys	Lys	Gly	Ser	Ser	Glu	Lys	Cys	Arg	Cys	Cys	Ala			
	50					55					60				

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 61 amino acids
- (B) TYPE: amino acid

WO 98/31795

PCT/US97/24129

44

- (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
(A) LIBRARY: GenBank
(B) CLONE: 386865

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Asp	Pro	Asn	Cys	Ser	Cys	Ala	Thr	Gly	Gly	Ser	Cys	Thr	Cys	Ala
1				5					10					15	
Gly	Ser	Cys	Lys	Cys	Lys	Glu	Cys	Lys	Cys	Thr	Ser	Cys	Lys	Lys	Ser
			20					25					30		
Cys	Cys	Ser	Cys	Cys	Pro	Val	Gly	Cys	Ala	Lys	Cys	Ala	Gln	Gly	Cys
		35					40					45			
Val	Cys	Lys	Gly	Ala	Ser	Glu	Lys	Cys	Ser	Cys	Cys	Ala			
50						55					60				

- (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 61 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
(A) LIBRARY: GenBank
(B) CLONE: 342385

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met	Asp	Pro	Asn	Cys	Ser	Cys	Ala	Thr	Gly	Val	Ser	Cys	Thr	Cys	Ala
1				5					10					15	
Asp	Ser	Cys	Lys	Cys	Lys	Glu	Cys	Lys	Cys	Thr	Ser	Cys	Lys	Lys	Ser
			20					25					30		
Cys	Cys	Ser	Cys	Cys	Pro	Val	Gly	Cys	Ala	Lys	Cys	Ala	Gln	Gly	Cys
		35					40					45			
Val	Cys	Lys	Gly	Ala	Ser	Glu	Lys	Cys	Asn	Cys	Cys	Ala			
50						55					60				

- (2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 61 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (vii) IMMEDIATE SOURCE:
(A) LIBRARY: GenBank
(B) CLONE: 305363

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met	Asp	Pro	Asn	Cys	Ser	Cys	Ala	Thr	Asp	Gly	Ser	Cys	Ser	Cys	Ser
1				5					10					15	
Gly	Ser	Cys	Lys	Cys	Lys	Glu	Cys	Lys	Cys	Thr	Thr	Cys	Lys	Lys	Ser
			20					25					30		
Cys	Cys	Ser	Cys	Cys	Pro	Val	Gly	Cys	Ala	Lys	Cys	Ser	Gln	Gly	Cys
		35					40					45			

WO 98/31795

PCT/US97/24129

45

Val Cys Lys Glu Ala Ser Glu Lys Cys Ser Cys Cys Ala
50 55 60

What is claimed is:

1. A substantially purified metallothionein comprising the amino acid sequence of SEQ ID NO:1 or fragments thereof.
2. An isolated and purified polynucleotide sequence encoding the metallothionein of claim 1.
3. A polynucleotide sequence which hybridizes under stringent conditions to the polynucleotide sequence of claim 2.
4. A hybridization probe comprising the polynucleotide sequence of claim 2.
5. An isolated and purified polynucleotide sequence comprising SEQ ID NO:2 or variants thereof.
6. A polynucleotide sequence which is complementary to the polynucleotide sequence of claim 2 or variants thereof.
7. A hybridization probe comprising the polynucleotide sequence of claim 6.
8. An expression vector containing the polynucleotide sequence of claim 2.
9. A host cell containing the vector of claim 8.
10. A method for producing a polypeptide comprising the amino acid sequence of SEQ ID NO:1 the method comprising the steps of:
 - a) culturing the host cell of claim 9 under conditions suitable for the expression of the polypeptide; and
 - b) recovering the polypeptide from the host cell culture.
11. A pharmaceutical composition comprising a substantially purified metallothionein having an amino acid sequence of claim 1 in conjunction with a suitable pharmaceutical carrier.
12. A purified antibody which binds specifically to the polypeptide of claim 1.
13. A purified agonist which specifically binds to and modulates the activity of the polypeptide of claim 1.
14. A purified antagonist which specifically binds to and modulates the activity of the polypeptide of claim 1.
15. A method for treating disorders caused by heavy metal toxicity comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 11.
16. A method for treating immune disorders comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 11.

WO 98/31795

PCT/US97/24129

47

17. A method for treating cancer comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 11.

18. A method for detection of polynucleotides encoding metallothionein in a biological sample comprising the steps of:

- 5 a) hybridizing the polynucleotide of claim 6 to nucleic acid material of a biological sample, thereby forming a hybridization complex; and
- b) detecting said hybridization complex, wherein the presence of said complex correlates with the presence of a polynucleotide encoding metallothionein in said biological sample.

3

1	M	D	P	N	C	S	C	T	T	G	V	S	C	A	C	T	G	S	C	K	C	K	E	C	K	C	T	X	C	K	K	S	C	C	S	C	P	V	G	HMBP-I
1	M	D	P	N	C	S	C	T	T	G	G	S	C	A	C	A	G	S	C	K	C	K	E	C	K	C	T	S	C	K	K	C	C	S	C	P	V	G	g386962	
1	M	D	P	N	C	S	C	A	T	G	G	S	C	T	C	A	G	S	C	K	C	K	E	C	K	C	T	S	C	K	K	S	C	C	P	V	G	g386865		
1	M	D	P	N	C	S	C	A	T	G	V	S	C	T	C	A	D	S	C	K	C	K	E	C	K	C	T	S	C	K	K	S	C	C	P	V	G	g342385		
1	M	D	P	N	C	S	C	A	T	D	G	S	C	S	C	S	G	S	C	K	C	K	E	C	K	C	T	T	C	K	K	S	C	C	P	V	G	g305363		
41	C	A	K	C	A	H	G	C	V	C	K	G	T	L	E	N	C	S	C	C	C	A																		HMBP-I
41	C	A	K	C	A	Q	G	C	V	C	K	G	S	S	E	K	C	R	C	C	C	A																	g386962	
41	C	A	K	C	A	Q	G	C	V	C	K	G	A	S	E	K	C	S	C	C	C	A																	g386865	
41	C	A	K	C	A	Q	G	C	V	C	K	G	A	S	E	K	C	N	C	C	C	A																	g342385	
41	C	A	K	C	S	Q	G	C	V	C	K	E	A	S	E	K	C	S	C	C	C	A																	g305363	

FIGURE 2

WO 98/31795

PCT/US97/24129

3/5




FIGURE 3A




FIGURE 3B

WO 98/31795

PCT/US97/24129

4/5




FIGURE 3C




FIGURE 3D

5/5




FIGURE 3E