



(43) International Publication Date
27 September 2012 (27.09.2012)

(10) International Publication Number
WO 2012/128965 A2

(51) International Patent Classification:
C07D 239/24 (2006.01)

(21) International Application Number:
PCT/US2012/028322

(22) International Filing Date:
8 March 2012 (08.03.2012)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/466,276 22 March 2011 (22.03.2011) US

(71) Applicant (for all designated States except US): **RE-
GENTS OF THE UNIVERSITY OF MINNESOTA**
[US/US]; 1000 Westgate Drive, Suite 160, St. Paul, Min-
nesota 55114-8658 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **VINCE, Robert**
[US/US]; 782 Hilltop Road, St. Paul, Minnesota 55108
(US). **VARTAK, Ashish P.** [IN/US]; 1390 Carling Drive,
#107, St. Paul, Minnesota 55108 (US).

(74) Agents: **PERDOK SHONKA, Monique M.** et al.;
Schwegman, Lundberg & Woessner, P.A., P.O. Box 2938,
Minneapolis, Minnesota 55402 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report (Rule 48.2(g))

(54) Title: METHOD FOR PREPARATION OF 5-SUBSTITUTED PYRIMIDINES

(57) Abstract: The present invention provides a method to prepare 5-substituted pyrimidines by reacting a 5-acylpyrimidine with a suitable nucleophile to afford a 5-methanolpyrimidine, such as flurprimidol.



WO 2012/128965 A2


METHOD FOR THE PREPARATION OF 5-SUBSTITUTED PYRIMIDINES

Cross-Reference to Related Applications

5 This application claims priority to U.S. Provisional Patent Application Serial No. 61/466,276 filed March 22, 2011, which application is incorporated herein by reference.


Background of the Invention

10 The 5-pyrimidinemethanol, flurprimidol (**1**, α -isopropyl- α -[p-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol, CAS # 56425-91-3) is a plant-growth regulator currently marketed under the brand names Cutless® and Topflor® as well as a mixture with various other agrochemicals. In the United States, it is manufactured by SePRO Corporation (Carmel, IN). Flurprimidol
15 was first described in US Patent 3,967,949. Its structure is depicted below:



Flurprimidol, **1**


Flurprimidol (**1**) is currently manufactured by the methodology illustrated in Scheme 1.



Scheme 1

Briefly, p-(trifluoromethoxy)bromobenzene (**2**) is converted to p-(trifluoromethoxy) phenylmagnesium bromide and reacted with isobutyronitrile to
25 afford isopropyl p-(trifluoromethoxy)phenyl ketone (**3**). The reaction of **3** with


5-bromopyrimidine **4** in the presence of *n*-BuLi affords **1**. This reaction proceeds through the intermediacy of the lithiated intermediate **5**. The conversion of 5-bromopyrimidine (**4**) to the 5-lithiopyrimidine (**5**) occurs without significant degradation only at or below -80°C . The cryogenic operation required to maintain such reaction conditions is expensive. The limited stability of **5** mandates the presence of **3** during the addition of *n*-butyllithium, which introduces impurities such as **6** and **7** in the final product.




Summary of the Invention

The present invention provides a method for preparing 5-acylpyrimidines comprising:


(a) reacting a compound of formula (II):




wherein X is halo or a sulfonate leaving group, such as tosyl or mesyl, with a compound of formula (III), (IV) or (V):




wherein R^1 is an organic group, M is a suitable metal cation, Y is $(\text{C}_1\text{-C}_4)\text{alkoxy}$, each X is halo, and x, y and z are selected so that the net charge on compound (V) is 0, in the presence of an effective catalytic amount of a complexed metal, to yield a compound of the formula (VI):



(b) oxidizing compound (VI) to yield compound (VII):




The present invention also comprises the step of reacting compound (VII) with a suitable nucleophile, such as $(R^2)MX$ or $(R^2)_2M$ wherein R^2 is an organic group, M is a single metal cation or a mixed metal cation and X is halo, or mixed halo, to yield a 5-methanolpyrimidine compound of formula (I):



Such compounds include the fungicides and/or plant growth retardants flurprimidol ($R^1 = 4$ -trifluoromethoxyphenyl, $R^2 =$ isopropyl), ancymidol ($R^1 = 4$ -methoxyphenyl, $R^2 =$ cyclopropyl), fenarimol ($R^1 = 4$ -chlorophenyl, $R^2 = 2$ -chlorophenyl), triarimol ($R^1 =$ phenyl, $R^2 = 3,4$ -dichlorophenyl), and nuarimol ($R^1 = 4$ -fluorophenyl, $R^2 = 2$ -chlorophenyl).

Novel 5-acylpyrimidine intermediates (VII) employed in the synthesis are also within the scope of the invention, as are other useful 5-methanolpyrimidines of general structure (I), and uses therefore. Such uses include plant growth control/limitation of plants such as turfgrass and ornamental trees and shrubs, as well as herbicides.


Thus, the present invention also includes 5-substituted pyrimidines of formula (IX):



wherein Y is O or $(H)_2$, and Z is (C_1-C_4) trifluoroalkyl, (C_1-C_4) alkoxy or halo, including compounds **8**, **12**, **18** and **23**.

Detailed Description of the Invention

The present invention provides a means to bypass the problematic chemical transformations in the currently-utilized method for 5-methanolpyrimidine synthesis. Shown in Scheme 2, for the retro-synthesis of flurprimidol, the sequence of alkyl group introduction is altered so that a 5-metallopyrimidine intermediate is not necessary. For example, the ketone intermediate necessary for an improved method to prepare flurprimidol (**1**) is 4-trifluoromethylphenyl-5-pyrimidyl ketone (**8**), wherein (isopropyl)M (**9**) can be isopropyl lithium, an isopropyl Grignard reagent, and the like.




Scheme 2


Therefore, the synthetic route of the present invention begins with a novel synthesis of a 5-acylpyrimidine **VII**. Ketone **VII** is then reacted with a suitable nucleophile, such as (**9**) to afford a 5-methanolpyrimidine such as flurprimidol. Traditionally, such transformations have been inefficient due to enolization of the ketone and free-radical mediated reduction of the C=O bond, both reducing yields. However, the Grignard reagent in a suitable organic solvent such as THF or an ether, can be mixed with one equivalent of LiCl and, subsequently, 50-100 mol-% ZnCl₂ or other metal halide as discussed below to form the lithium complex of a magnesio-zincate which appears to possess much reduced basicity and therefore leading to nucleophilic attack as the predominant reaction pathway.

The present method comprises the following steps:


Step 1 A transition-state metal mediated C–C bond coupling between a 5-sulfonyl or halopyrimidine (**II**) and an appropriate organometallic or organoboron reagent (**III**) to afford a 5-substituted pyrimidine (**VI**);



Step 2 Oxidation of the activated benzylic carbon of VI to afford a 5-acylpyrimidine (VII); and



5 Step 3 Addition of a nucleophile to the ketone to afford the target pyrimidinemethanol (I)



Step 1. Synthesis of III and IV.

Compound III can be an organometallic reagent with the metal being
 10 Mg(II), Zn(II), Mn(II), Cu(I), Sn(IV) or Li(I). Compound IV can also include mixed organometallic complexes such as $\text{R}_1\text{CH}_2(\text{Li}^+)_x(\text{M}^{++})_y\text{X}^-_z$ (where x, y and z are such that the net charge of the species = 0); for example, if M^{++} is zinc, the mixed metal complex is lithium zincate.

Preparation of compounds III or IV can be accomplished through one of
 15 the following methods.

Method 1) An alkyl halide $\text{R}_1\text{CH}_2\text{X}$ is reacted with a suitable form of a metal in an inert solvent at a temperature range between -20°C and 90°C . An activating agent such as I_2 or dibromoethane may be added. This procedure affords $\text{R}_1\text{CH}_2\text{MX}$.

20 Method 2) An alkyl halide $\text{R}_1\text{CH}_2\text{X}$ is reacted as per Method 1, except that LiCl is mixed with the metal in an inert solvent before addition of the alkyl halide. This procedure affords $\text{R}_1\text{-CH}_2\text{M(Li)XCl}$. LiCl may be substituted with LiBr or LiI.

Method 3) An alkyl halide $\text{R}_2\text{CH}_2\text{X}$ is treated with an organometallic
 25 reagent such as *n*-BuLi, *s*-BuLi, *t*-BuLi or *i*-PrMgX in the presence of a

complexing agent such as tetramethyl ethylenediamine (TMEDA). Additives such as MgX_2 or ZnX_2 may be added to generate a mixed organometallic reagent.

A "suitable form of metal" is a form that reacts with R_2CH_2X to preferentially afford R_2CH_2MX , comprising a "suitable metal cation," over the Würtz coupling product, $R_2CH_2CH_2R_2$. For example, commercial zinc dust reacts with 4-(OCF₃)BnBr to afford 80% of (4-(OCF₃)-PhCH₂)₂ and is hence an unsuitable form of zinc. When the zinc dust is heated with I₂ (1 wt%) and dibromoethane (2 wt%) in THF at reflux until ethylene gas evolves, the resulting active form of zinc dust reacts with 4-(OCF₃)BnBr to afford >98% of 4-(OCF₃)BnZnBr.

The catalyst utilized for the coupling of compounds **III**, **IV** or **V** with **II** may be derived from palladium, nickel or iron in any oxidation state. Examples are (Ph₃P)₄Pd, (Ph₃P)₂PdCl₂, (Ph₃P)₂NiCl₂ and Fe(acac)₂. Catalyst loading is from 0.1 to 10 mol% of **II**. The reaction temperature ranges from -10 °C in case of palladium complexes to room temperature in case of the nickel and iron complexes. Reaction time is between 48 h for the palladium-catalyzed reactions to 15 minutes for the nickel-catalyzed reaction. Precise conditions, solvents, method of isolation and yields are given in Table 1. In all cases, the reaction can be quenched with saturated aqueous NH₄Cl, washed with sat. NaHCO₃, the organic layer dried with MgSO₄ and evaporated to a pale yellow to brown gum, which was in all cases found to be >95% pure **VI** by NMR and carried to Stage 2 without further purification. Table 1 describes the yields obtained with specific conditions and catalysts.

Entry	(III)	(II)	Additive 100 mol%	Catalyst	Cat Mol%	Solvent	T (°C)	Time (hours)	Yield (VI)
1	R ₁ -MgBr*	5-bromo	LiCl	(Ph ₃ P) ₄ Pd	5	THF	rt	8	ca.20%
2	R ₁ -MgBr*	5-bromo	ZnCl ₂	(Ph ₃ P) ₄ Pd	5	THF	rt	24	46%
3	R ₁ -MgBr*	5-bromo	ZnCl ₂	(Ph ₃ P) ₄ Pd	5	THF	0	24	40%
4	R ₁ -ZnBr*	5-bromo	No	(Ph ₃ P) ₄ Pd	5	THF	0 to rt	24	73%
5	R ₁ -ZnBr*	5-bromo	LiCl	(Ph ₃ P) ₄ Pd	5	THF	rt	4	90%
6	R ₁ -ZnBr*	5-bromo	LiCl	(Ph ₃ P) ₂ NiCl ₂	5	THF	rt	0.25	95%
7	R ₁ -ZnBr*	5-bromo	LiCl	(Ph ₃ P) ₂ NiCl ₂	0.5	THF	rt	1	98%
8	R ₁ -ZnBr*	5-bromo	LiCl	(Ph ₃ P) ₂ NiCl ₂	0.1	THF	rt	2	93%
9	R ₁ -ZnBr*	5-iodo	LiCl	(Ph ₃ P) ₂ NiCl ₂	0.1	THF	rt	2	85%
10	R ₁ -ZnBr*	5-chloro	LiCl	(Ph ₃ P) ₂ NiCl ₂	0.1	THF	rt	24	12%
11	R ₁ -ZnBr*	5-bromo	LiCl	Fe(acac) ₂	5	THF	rt	48	35%
12	R ₁ -ZnBr*	5-bromo	LiCl	Fe(acac) ₂	10	THF	Rt	48	28%
13	R ₁ -ZnBr*	5-bromo	LiCl	Fe(acac) ₂	10	DME	90	3	74%
14	R ₁ -ZnBr*	5-bromo	LiCl	Fe(acac) ₂	1	DME	90	3	88%
15	R ₁ -ZnBr*	5-bromo	No	(Ph ₃ P) ₂ NiCl ₂	0.1	THF	rt	14	71%
16	R ₁ -ZnBr*	5-bromo	LiCl	(Ph ₃ P) ₂ NiCl ₂	0.1	Ether	rt	2	94%

Entry	(III)	(II)	Additive 100 mol%	Catalyst	Cat Mol%	Solvent	T (°C)	Time (hours)	Yield (VI)
17	R ₁ -ZnBr [*]	5-bromo	LiCl	(Ph ₃ P) ₂	0.1	di- <i>n</i> -bu ether	Rt	48	NR
18	R ₁ -ZnCl [*]	5-bromo	LiCl	(Ph ₃ P) ₂ NiCl ₂	0.1	THF	Rt	5	88%
19	R ₁ -ZnCl [†]	5-bromo	LiCl	(Ph ₃ P) ₂ NiCl ₂	0.1	THF	0 °C	5	54%
20	R ₁ -ZnCl [‡]	5-bromo	LiCl	(Ph ₃ P) ₂ NiCl ₂	0.1	THF	0 °C	5	99%
21	R ₁ -ZnCl [‡]	5-bromo	LiCl	(Ph ₃ P) ₂ NiCl ₂	0.1	THF	0 °C	5	92%

Table 1. Conditions scan for stage 1. * R₁ = 4-trifluoromethoxyphenyl, † R₁ = 4-methoxyphenyl, ‡ R₁ = 4-chlorophenyl. Yields refer to isolated yields. LiCl was dried under hyvac for 15 min at 150 °C. ZnCl₂ was melted under hyvac and congealed with vigorous shaking to afford a dry powder. Zn dust was heated with I₂ (1 wt%) and dibromoethane (2 wt%) at reflux in the solvent until ethylene gas evolved and then cooled to rt before use for formation of R₁-ZnX.

Nickel(II)chloride (as its bis-triphenylphosphine complex) is a surprisingly active catalyst for this particular benzylic sp^3 -aryl sp^2 coupling reaction. An organozincate, prepared in the absence of magnesium(II) is the most suitable benzylic coupling partner. The nickel(II) catalyst may be substituted with

5 Fe(acac)₂, but the reactions requires higher temperature (and therefore change of solvent to DME) to go to completion. Palladium catalysts are also active, but are impractical due to their cost (2000-times more expensive than nickel catalysts). Diethyl ether is also a suitable solvent for the reaction. When the stoichiometry of reactants II and III is held in strict 1:1 stoichiometry, the oily residue after workup is

10 essentially pure **VI** with only traces of the Würtz coupling product described earlier. The residue is utilized directly for the oxidation in Step 2. In case the stoichiometry is not rigidly 1:1, the presence of traces of **II** and **III** do not affect the outcome of the manipulations in Step II.

15 Step 2. Oxidation of **VI** to pyrimidyl ketone (**VII**).

The benzylic methylene of compound **VI** is susceptible to oxidation. This oxidation may be conducted utilizing a number of known oxidizing agents. The oxidant may be a sole, stoichiometric oxidant, such as SeO₂, KMnO₄ and CrO₃, or may be a catalytic oxidant combined with a 'sacrificial' oxidant, such as the

20 combination of SeO₂ and *t*-butylhydroperoxide, *N*-hydroxyphthalimide and *t*-butyl hydroperoxide, RuO₂ and commercial bleach. Alternatively, benzylic halogenations may be effected with NBS (N-bromosuccinimide), NCS (N-chlorosuccinimide), Br₂, chloramine-T, etc. in the presence or absence of a radical initiator like azobisisobutyronitrile (AIBN), dibenzoylperoxide and irradiation with any suitable

25 wavelength of light. The halogenated products are carried forward into an oxidation, for example, with nitric acid in AcOH, which affords the ketone **VII**.

Amongst stoichiometric oxidants, KMnO₄ is an oxidant of choice because of its low cost and the environmentally benign nature of the MnO₂ by-product. Conditions for KMnO₄ oxidation are tailored to the reactivity of the 5-substituted

30 pyrimidine **VI**. For example, while a temperature of 80 °C is required for the oxidation of 4-(trifluoromethoxy)benzyl substituted **VI**, the oxidation of 4-


(methoxy)substituted **VI** occurs at room temperature. A suitable organic solvent such as ethyl acetate may be optionally employed, but is not necessary.

After consumption of **VI**, the reaction mixture is extracted with a solvent such as ethyl acetate and the organic layer can be washed with dilute sodium

- 5 bisulfite, brine, dehydrated and evaporated. The residue is utilized directly in Step 3.

Step 3. Conversion of acylpyrimidine (**VII**) to pyrimidinemethanol (**I**).


- The reaction of nucleophiles such as organolithiums (R_2Li) and
10 organomagnesiums (R_2MgX) with 5-acylpyrimidines such as **VII** occurs in low yields (<10%). Competing reactions are enolization leading to recovery of **VII** after workup and β -hydride transfer leading to recovery of **VIII** after workup.



- To obtain maximal conversion of **VII** to **I**, the reactivity of the nucleophile is
15 modulated by the addition of LiCl and 50–100 mol% of either ZnX_2 , MnX_2 , LnX_3 where Ln is a lanthanide and X is halo, preferably chloro. The effective nucleophile in such as case may be represented as $(Li(R_2-ZnX)MgX_2)_y \cdot solvate_z$, where the solvent is an ether such as THF or diethyl ether and where y and z are such that all the low-lying *d*-orbitals of the transition metal are filled.

- 20 When duly optimized, a 100% conversion of **VII** can be obtained and the isolated yield is as high as 98%. Isolation of **I** can be effected by extraction of the reaction mixture of Step 3 with saturated aqueous NH_4Cl , evaporation of the organic layer and recrystallization of the semisolid residue from a suitable solvent or solvent combination.

25

Experimental section.**EXAMPLE 1. SYNTHESIS OF FLURPRIMIDOL**Scheme 3. Synthesis of flurprimidol (**12**).

5

Stage 1. Synthesis of 5-[4-(trifluoromethoxy)benzyl] pyrimidine (**12**)(A) From 4-(trifluoromethoxy)benzyl bromide (**10a**)

A suspension of zinc dust (15 g, excess) in THF (100 mL) was treated with LiCl (10.00 g, 238 mmol) and 1,2-dibromoethane (0.5 mL, catalytic). The warm mixture thus formed was treated dropwise with a solution of 4-(trifluoromethoxy)benzyl bromide (**10a**) (50.00 g, 209 mmol) in THF (200 mL) over a period of 3 hours. The resulting mixture was stirred for an additional hour, affording a solution of the lithium zincate **11**, and then treated with $(\text{Ph}_3\text{P})_2\text{NiCl}_2$ (300 mg, 0.1 mol or 0.001 equivalents). The red suspension was treated drop wise with a solution of 5-bromopyrimidine (**4**) (33.23 g, 200 mmol) in THF (50 mL) over a period of one hour. The mixture was stirred for another 10 minutes and then poured into 300 mL of saturated aq. NH_4Cl . The aqueous layer was washed with 200 mL of THF-ether (1:1), the combined organic layers washed with saturated aqueous NaHCO_3 , filtered, dried (MgSO_4) and evaporated under reduced pressure to afford a pale yellow oily residue weighing 51.00 g (96%) and was found to be pure 5-[4-(trifluoromethoxy)benzyl]pyrimidine (**12**). ^1H NMR (400 MHz, CDCl_3) δ ppm 9.26 (s, 1H, pyrimidine), 8.84 (s, 2H, pyrimidine), 7.19 (s, 4H, Ar, no para-pattern observed), 4.06 (s, 2H, methylene); ^{13}C NMR (100 MHz, CDCl_3) 156.3, 157.1, 147.4, 139.6, 134.8, 131, 121.7, 35.0. ESI-MS (low-res) 255.0 ($\text{M}+\text{H}$) $^+$.

(B) From 4-(trifluoromethoxy)benzyl chloride (**10b**)

25

To zinc dust (1.30 g, 20 mmol) in THF (20 mL) was added I₂ (10 mg) and dibromoethane (100 mg). The suspension was heated to reflux and then cooled to rt before being treated with **10b**. The resulting suspension was heated to reflux for 6 h and then cooled to rt. An iodine titration indicated 94% yield of the organozinc reagent. (Ph₃P)₂NiCl₂ (50 mg, cat) was added resulting in a bright red suspension, which was treated with **4** (2.97 g, 18.8 mmol), resulting in discharge of the red color. The mixture was extracted with NH₄Cl, saturated NaHCO₃ and the organic layer was evaporated under reduced pressure to afford a pale yellow oily residue (4.4 g, 92%) which was found to be pure **12**.

10

Step 2. Synthesis of 4-trifluoromethoxyphenyl-5-pyrimidyl ketone (**8**).

(A) By NBS and nitric acid-AcOH mediated oxidation

To a solution of **12** (25.00 g, 98.4 mmol) in EtOAc (55 mL) was added NBS (17.8 g, 100 mmol). The mixture was flushed with argon gas and then irradiated with a 300 W tungsten-filament lamp for 90 minutes, during which, all of IV was consumed according to TLC (R_f of IV = 0.35 in CH₂Cl₂-EtOAc, 3:1). The mixture was washed thrice with water (20 mL), dried, and evaporated to afford a crude brown oil weighing 35 g (108% mass recovery). The crude gum was dissolved in AcOH (50 mL) and nitric acid (25 mL) was added. The mixture was heated at 50 °C until evolution of a brown gas completely ceased, and the poured into 10% aq. sodium chloride and 20 g of ice. The mixture was extracted with EtOAc (50 mL, 3x), the EtOAc extracts dried (MgSO₄) and evaporated to a brown gum. This gum was triturated with hexanes (100 mL) to afford a brown solid, which was stirred in boiling hexanes-EtOAc-EtOH (10:1:1, 200 mL) until complete solution occurred. The solution was allowed to cool in the refrigerator overnight and the needles formed were filtered, crushed, and dried under vacuum to afford pure **8** (15.71 g, 60%). The filtrates contained negligible amounts of V. mp = 56–58 °C; ¹H NMR (400 MHz, CDCl₃) δ ppm 9.18 (s, 1H, pyrimidine), 9.02 (s, 1H, pyrimidine), 7.40 (d, 2H, Ar), 7.18 (d, 2H, Ar). ¹³C NMR (100 MHz, CDCl₃) δ ppm 213.8 (ketone) 158.2, 156.1 (pyrimidine), 136.4, 136.1, 135.0, 127.9, 121.8 (Ar).


30

(B) By *N*-hydroxyphthalimide and *tert*-butylhydroperoxide mediated oxidation.

A solution of **12** (2.54 g, 10 mmol) in *tert*-butyl hydroperoxide (10 mL) was treated with *N*-hydroxyphthalimide (10 mg, catalytic). The mixture was irradiated with 300 W lamp while being cooled in an ice-water bath. After 6 h, the mixture was evaporated under reduced pressure and partitioned between EtOAc and brine. The EtOAc layer was treated with activated charcoal, filtered and evaporated under reduced pressure. The residue was triturated with hexanes:ether (2:1) and the supernatant was decanted and evaporated to afford **8** as an oil (2.60 g, 98% yield). The white solid left behind after trituration was found to be mostly *N*-hydroxyphthalimide.

(C) By KMnO_4 mediated oxidation.


To solution of **12** (2.54 g, 10 mmol) in EtOAc (10 mL) was added water (20 mL) and KMnO_4 (6.32 g, 40 mmol). The mixture was heated to reflux for 1 h and then filtered through celite. The organic layer was evaporated under reduced pressure to afford an oil (2.40 g, 88%) that was found by NMR to be pure **8**.

Step 3. Conversion of 5-acylpyrimidine **8** to flurprimidol **1**.

A solution of **8** (14.0 g, 53.23 mmol) in THF (100 mL) was treated with LiCl (2.23 g, 53.23 mmol), ZnCl_2 (3.61 g, 0.5 equiv), followed by commercial isopropyl magnesium chloride (2M in THF, 26.6 mL) dropwise at 0 °C. The mixture was stirred for an extra 30 minutes and then extracted with sat. aq. NH_4Cl (25 mL, 2x). The organic layer was evaporated to a pale yellow gum, which was dissolved in toluene (10 mL) and hexane (10 mL). This solution upon cooling in the refrigerator afforded white needles. Hexane (30 mL) was added to complete crystallization (TLC of the mother-liquor showed absence of **1**) and the filtered. The cake was

dried in air until constant weight and was found to be pure flurprimidol (**1**). The yield was 15.14 g (94%). M.p. = 95–97 °C; ¹H NMR (600 MHz, CDCl₃) δ ppm 9.06 (s, 1H, ar), 8.82 (s, 1H, Ar), 7.52 (d, 2H, *J* = 1.8 Hz, Ar), 7.19 (d, 2H, *J* = 1.9 Hz, Ar), 2.86 (dq, 1H, *J*₁ = 5.7 Hz, *J*₂ = 4.3 Hz, CH), 0.93 (d, 6H, *J* = 4.5 Hz, (CH(CH₃)₂)). ¹³C NMR (125 MHz, CDCl₃) δ ppm 157.0, 154.6, 143.41, 138.9, 127.1, 121.0 (Ar), 78.34 (C), 35.0 (CH), 16.8, 16.6 (CH(CH₃)₂).

EXAMPLE 2. SYNTHESIS OF ANCYMIDOL (**15**).



Scheme 4. Synthesis of ancymidol (**15**).

Step 1. Synthesis of 4-(methoxybenzyl)pyrimidine (**17**).

A suspension of zinc dust (1300 mg, 20 mmol), LiCl (420 mg, 10 mmol), I₂ (10 mg), dibromoethane (50 mg) in THF (20 mL) was heated to reflux for 15 min. and then treated with 4-methoxybenzyl chloride (**16**) (1.56 g, 10 mmol). Heating was continued for 6 h and the suspension was then cooled to rt before being treated (Ph₃P)₂NiCl₂ (50 mg). The resulting red suspension was treated with **4** (1.56 g, 10 mmol) and the mixture was stirred for an extra 10 min. before being washed with NH₄Cl and saturated NaHCO₃. The organic layer was evaporated under reduced pressure to afford a pale yellow oil (3.03 g, 94%), which was found to be pure **17** by NMR. ¹H NMR (300 MHz, CDCl₃) δ ppm 9.01 (s, 1H), 8.55 (s, 2H), 7.95 (d, 2H), 7.39 (d, 2H), 4.12 (s, 2H), 3.52 (s, 3H).

Step 2. Synthesis of 4-(methoxybenzyl)-5-pyrimidyl ketone (**18**).


A solution of **17** (2.00 g, 10 mmol) in EtOAc (10 mL) was treated with KMnO₄ (3.18 g, 20 mmol) in water (20 mL). The biphasic mixture was stirred at rt

for 12 h, at which point the purple color was discharged. The mixture was filtered through celite and the organic layer was evaporated to afford **18** as an oil (2.04 g, 94 %). ¹H NMR (300 MHz, CDCl₃) δ ppm 9.78 (s, 1H), 9.14 (s, 2H), 7.83 (d, 2H), 7.08 (d, 2H), 3.74 (s, 3H). ¹³C NMR (75 MHz, CDCl₃) δ ppm 194.3, 158.3, 158.0, 138.5, 136.6, 128.4, 118.0, 116.8, 63.1.

Stage 3. Synthesis of Ancymidol (**15**).

To a solution of cyclopropylmagnesium bromide (0.5 M in THF, 9 mL, 4.5 mmol) was added ZnCl₂ (612 mg, 4.5 mmol) and LiCl (0.189 mg, 4.5 mmol). To this clear yellow solution was added **18** (1.00 g, 4.5 mmol) in THF and the mixture was stirred for 4 h at rt before being washed with sat. NH₄Cl. The organic layer was diluted with EtOAc and washed with sat. NaHCO₃ before being dried and evaporated to a brown gum, which was dissolved in toluene and poured over a plug of silica gel (20 g). The plug was washed with hexane-EtOAc (100 mL). The eluants were evaporated to a yellow oil that solidified upon trituration with hexanes. The solid was filtered and dried to afford 374 mg (31%) of **15** as a white solid. Mp = 108-110 °C; ¹H NMR (300 MHz, CDCl₃) δ ppm 9.06 (s, 1H), 8.48 (s, 2H), 7.82 (d, 2H), 7.03 (d, 2H), 3.99 (s, 3H), 1.08 (m, 1H), 0.50-0.31 (m, 4H).

EXAMPLE 3. SYNTHESIS OF FENARIMOL (**19**).



Scheme 5. Synthesis of fenarimol (**19**)

Step 1. Synthesis of 5-(4-chlorobenzyl)pyrimidine (**22**).

A mixture of Zn dust (13 g, 100 mmol), LiCl (4.2 g, 100 mmol), I₂ (250 mg), dibromoethane (2.5 g) in THF (100 mL) was heated to reflux for 15 min. and then

treated with 4-chlorobenzyl chloride (8.1 g, 50 mmol). The suspension was heated to reflux for 48 h and then cooled to rt. An iodometric titration indicated a 65% yield of the zincate. $(\text{Ph}_3\text{P})_2\text{NiCl}_2$ (200 mg) was then added and the red suspension treated with **4** (5.13 g, 32.5 mmol). After 30 min, the pale green suspension was
5 extracted with sat. NH_4Cl (200 mL), diluted with EtOAc (300 mL) and extracted with sat. NaHCO_3 (200 mL). The organic layer was evaporated under reduced pressure to afford a semisolid weighing 6.07 g (60% yield) that was identified by NMR to be **22** and the "homocoupled" product, 1,2-di-(4,4'-dichloro)phenyl ethane in a 3:2 ratio. This residue was directly subjected to Step 2 without further
10 purification.

Step 2. Synthesis of 4-chlorophenyl-5-pyrimidyl ketone (**23**).

A mixture of the residue of Step 1 containing **22** (5.00 g, 5 mmol based on NMR purity), EtOAc (50 mL), water (100 mL) and KMnO_4 (5 g, excess) was
15 heated to reflux for 6 h and then filtered through celite. The filtrate was diluted with brine (50 mL) and the organic layer was separated, washed with 5% sodium bisulfate, sat. NaHCO_3 (50 mL, 3 times) and evaporated to a tan solid. This solid was stirred in hexanes and then filtered to afford pure **23** as an off-white solid. (4.5 g, 85%) $\text{Mp} = 88\text{--}90\text{ }^\circ\text{C}$; ^1H NMR (300 MHz, CDCl_3) δ ppm 9.81 (s, 1H), 9.20 (s,
20 2H), 7.92 (d, 2H), 7.26 (d, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ ppm 190.4, 158.6, 158.1, 138.4, 132.1, 127.5, 118.7, 118.0.

Step 3. Synthesis of fenarimol (**19**).

To a suspension of magnesium turnings (1.00 g, excess) in THF (20 mL)
25 was added 1,2-dibromoethane (0.05 mL). After 5 min, *o*-bromochlorobenzene (960 mg, 5 mmol) was added in one portion, causing slight warming as the oxidative addition initiated. After 90 min. a menthol-phenanthroline titration of this solution indicated 100% insertion. This grey solution was treated with ZnCl_2 (680 mg, 5 mmol) and then with **23** (1.1 g, 5 mmol). The resulting yellow-brown solution was
30 stirred at rt for 30 min during which all of **23** was consumed according to TLC. The mixture was washed with NH_4Cl , sat. aq. NaHCO_3 , and the organic layer was

evaporated under reduced pressure to a brown gum. This residue was subjected to chromatography over 15 g of silica gel, eluting with 3:1 hexane-EtOAc going to 1:1 hexane-EtOAc. Evaporation of the eluant between 100 mL and 250 mL afforded **19** as a white solid (330 mg, 20%). Mp = 115-116 °C; ¹H NMR (300 MHz, CDCl₃) δ
5 ppm 9.00 (s, 1H), 8.48 (s, 2H), 8.20–7.08 (m, 8H, Ar); ¹³C NMR (75 MHz, CDCl₃) δ 160.1, 156.7, 138.7, 138.1, 137.5, 132.8, 132.3, 128.9, 128.4, 128.2, 127.4, 118.4, 118.5, 80.1.

In compounds **I**, and **III-VII**, R¹ and R² can individually be organic groups or "radicals" that are substituted or unsubstituted, including alkyl, aryl, aralkyl
10 cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl or the like. R¹ and R² each can include about 1-20 carbon atoms. R¹ and/or R² can include double or triple bonds. R¹ and/or R² can also include heteroatoms.

R¹ and R² can each be substituted or unsubstituted as those terms are defined
15 herein. Substituted alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, and cycloalkenylalkyl groups as well as other substituted groups also include groups in which one or more bonds to a hydrogen atom are replaced by one or more bonds, including double or triple bonds to a heteroatom such as, but not limited to, oxygen in carbonyl (oxo), carboxyl, ester, amide, imide, urethane, and
20 urea groups; and nitrogen in imines, hydroxyimines, oximes, hydrazones, amidines, guanidines, and nitriles.

In general, "substituted" refers to an organic group as defined herein such as R¹ or R² in which one or more bonds to a hydrogen atom contained therein are replaced by one or more bonds to a non-hydrogen atom such as, but not limited to,
25 a halogen (i.e., F, Cl, Br, and I); an oxygen atom in groups such as hydroxyl groups, alkoxy groups, aryloxy groups, aralkyloxy groups, oxo(carbonyl) groups, carboxyl groups including carboxylic acids, carboxylates, and carboxylate esters; a sulfur atom in groups such as thiol groups, alkyl and aryl sulfide groups, sulfoxide groups, sulfone groups, sulfonyl groups, and sulfonamide groups; a nitrogen atom in groups
30 such as amines, hydroxylamines, nitriles, nitro groups, N-oxides, hydrazides, azides, and enamines; and other heteroatoms in various other groups.

Non-limiting examples of substituents that can be bonded to a substituted carbon (or other) atom include halo (F, Cl, Br, I), OR', e.g., alkoxy, OC(O)N(R')₂, CN, NO, NO₂, ONO₂, azido, haloalkyl, haloalkoxy, e.g., OCF₃, O (oxo), S (thiono), C(O), S(O), methylenedioxy, ethylenedioxy, N(R')₂, SR', SOR', SO₂R', SO₂N(R')₂,
 5 SO₃R', C(O)R', C(O)C(O)R', C(O)CH₂C(O)R', C(S)R', C(O)OR', OC(O)R', C(O)N(R')₂, OC(O)N(R')₂, C(S)N(R')₂, (CH₂)₀₋₂N(R')C(O)R', (CH₂)₀₋₂N(R')N(R')₂, N(R')N(R')C(O)R', N(R')N(R')C(O)OR', N(R')N(R')CON(R')₂, N(R')SO₂R', N(R')SO₂N(R')₂, N(R')C(O)OR', N(R')C(O)R', N(R')C(S)R', N(R')C(O)N(R')₂, N(R')C(S)N(R')₂, N(COR')COR', N(OR')R', C(=NH)N(R')₂, C(O)N(OR')R', or
 10 C(=NOR')R' wherein R' can be hydrogen or a carbon-based moiety, and wherein the carbon-based moiety can itself be further substituted.

When a substituent is monovalent, such as, for example, F or Cl, it is bonded to the atom it is substituting by a single bond. When a substituent is more than monovalent, such as O, which is divalent, it can be bonded to the atom it is
 15 substituting by more than one bond, i.e., a divalent substituent is bonded by a double bond; for example, a C substituted with O forms a carbonyl group, C=O, which can also be written as "CO", "C(O)", or "C(=O)", wherein the C and the O are double bonded. When a carbon atom is substituted with a double-bonded oxygen (=O) group, the oxygen substituent is termed an "oxo" group. When a divalent
 20 substituent such as NR is double-bonded to a carbon atom, the resulting C(=NR) group is termed an "imino" group. When a divalent substituent such as S is double-bonded to a carbon atom, the results C(=S) group is termed a "thiocarbonyl" group.

Alternatively, a divalent substituent such as O, S, C(O), S(O), or S(O)₂ can be connected by two single bonds to two different carbon atoms. For example, O, a
 25 divalent substituent, can be bonded to each of two adjacent carbon atoms to provide an epoxide group, or the O can form a bridging ether group, termed an "oxy" group, between adjacent or non-adjacent carbon atoms, for example bridging the 1,4-carbons of a cyclohexyl group to form a [2.2.1]-oxabicyclo system. Further, any substituent can be bonded to a carbon or other atom by a linker, such as (CH₂)_n or
 30 (CR'₂)_n wherein n is 1, 2, 3, or more, and each R' is independently selected.

C(O) and S(O)₂ groups can be bound to one or two heteroatoms, such as nitrogen, rather than to a carbon atom. For example, when a C(O) group is bound to one carbon and one nitrogen atom, the resulting group is called an "amide" or "carboxamide." When a C(O) group is bound to two nitrogen atoms, the functional
5 group is termed a urea. When a S(O)₂ group is bound to one carbon and one nitrogen atom, the resulting unit is termed a "sulfonamide." When a S(O)₂ group is bound to two nitrogen atoms, the resulting unit is termed a "sulfamate."

Substituted ring groups such as substituted cycloalkyl, aryl, heterocyclyl and heteroaryl groups also include rings and fused ring systems in which a bond to a
10 hydrogen atom is replaced with a bond to a carbon atom. Therefore, substituted cycloalkyl, aryl, heterocyclyl and heteroaryl groups can also be substituted with alkyl, alkenyl, and alkynyl groups as defined herein.

As to any of the R¹ or R² other groups described herein, which contain one or more substituents, it is understood, of course, that such groups do not contain any
15 substitution or substitution patterns which are sterically impractical and/or synthetically non-feasible. In addition, the compounds of this disclosed subject matter include all stereochemical isomers arising from the substitution of these compounds.

By "chemically feasible" is meant a bonding arrangement or a compound
20 where the generally understood rules of organic structure are not violated; for example a structure within a definition of a claim that would contain in certain situations a pentavalent carbon atom that would not exist in nature would be understood to not be within the claim. The structures disclosed herein, in all of their embodiments are intended to include only "chemically feasible" structures, and any
25 recited structures that are not chemically feasible, for example in a structure shown with variable atoms or groups, are not intended to be disclosed or claimed herein.

When a substituent is specified to be an atom or atoms of specified identity, "or a bond", a configuration is referred to when the substituent is "a bond" that the groups that are immediately adjacent to the specified substituent are directly
30 connected to each other in a chemically feasible bonding configuration.

An "acyl" group ($R^1C(O)$) as the term is used herein refers to a group containing a carbonyl moiety wherein the group is bonded, to a moiety such as a pyrimidine ring, via the carbonyl carbon atom. The carbonyl carbon atom is also bonded to R^1 . A nicotinoyl group (pyridyl-3-carbonyl) group is an example of an acyl group within the meaning herein. Other examples include acetyl, i-butyryl, benzoyl, phenylacetyl, pyridylacetyl, cinnamoyl, and acryloyl groups and the like.

The term "alkoxy" refers to an oxygen atom connected to an alkyl group, including a cycloalkyl group, as are defined above. Examples of linear alkoxy groups include but are not limited to methoxy, ethoxy, propoxy, butoxy, pentyloxy, hexyloxy, and the like. Examples of branched alkoxy include but are not limited to isopropoxy, sec-butoxy, tert-butoxy, isopentyloxy, isohexyloxy, and the like. Examples of cyclic alkoxy include but are not limited to cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, and the like. An alkoxy group can include one to about 12-20 carbon atoms bonded to the oxygen atom, e.g., is (C_1 - C_4)alkoxy and can further include double or triple bonds, and can also include heteroatoms. For example, an allyloxy group is an alkoxy group within the meaning herein. A methoxyethoxy group is also an alkoxy group within the meaning herein, as is a methylenedioxy group in a context where two adjacent atoms of a structures are substituted therewith.

The terms "aryloxy" and "arylalkoxy" refer to, respectively, an aryl group bonded to an oxygen atom and an aralkyl group bonded to the oxygen atom at the alkyl moiety. Examples include but are not limited to phenoxy, naphthyloxy, and benzyloxy.

The terms "halo" or "halogen" or "halide" by themselves or as part of another substituent mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom, preferably, fluorine, chlorine, or bromine.

A "haloalkyl" group includes mono-halo alkyl groups, poly-halo alkyl groups wherein all halo atoms can be the same or different, and per-halo alkyl groups, wherein all hydrogen atoms are replaced by halogen atoms, such as fluoro. Examples of (C_1 - C_4)haloalkyl include trifluoromethyl, 1,1-dichloroethyl, 1,2-dichloroethyl, 1,3-dibromo-3,3-difluoropropyl, perfluorobutyl, and the like.

A "haloalkoxy" group includes mono-halo alkoxy groups, poly-halo alkoxy groups wherein all halo atoms can be the same or different, and per-halo alkoxy groups, wherein all hydrogen atoms are replaced by halogen atoms, such as fluoro. Examples of haloalkoxy include trifluoromethoxy, 1,1-dichloroethoxy, 1,2-dichloroethoxy, 1,3-dibromo-3,3-difluoropropoxy, perfluorobutoxy, and the like.

The term "(C_x-C_y)perfluoroalkyl," wherein $x < y$, means an alkyl group with a minimum of x carbon atoms and a maximum of y carbon atoms, wherein all hydrogen atoms are replaced by fluorine atoms. Preferred is -(C₁-C₆)perfluoroalkyl, more preferred is -(C₁-C₃)perfluoroalkyl, most preferred is -CF₃.

The term "(C_x-C_y)perfluoroalkylene," wherein $x < y$, means an alkyl group with a minimum of x carbon atoms and a maximum of y carbon atoms, wherein all hydrogen atoms are replaced by fluorine atoms. Preferred is -(C₁-C₆)perfluoroalkylene, more preferred is -(C₁-C₃)perfluoroalkylene, most preferred is -CF₂-.

Alkyl groups include straight chain and branched alkyl groups having from 1 to about 20 carbon atoms, and typically from 1 to 12 carbons or, in some embodiments, from 1 to 8 carbon atoms. Examples of straight chain alkyl groups include those with from 1 to 8 carbon atoms such as methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl, and n-octyl groups. Examples of branched alkyl groups include, but are not limited to, isopropyl, iso-butyl, sec-butyl, t-butyl, neopentyl, isopentyl, and 2,2-dimethylpropyl groups. As used herein, the term "alkyl" encompasses n-alkyl, isoalkyl, and anteisoalkyl groups as well as other branched chain forms of alkyl. Representative substituted alkyl groups can be substituted one or more times with any of the groups listed above, for example, halo, amino, hydroxy, cyano, carboxy, nitro, thio, alkoxy, and halogen groups.

Cycloalkyl groups are C₃-C₁₂ cyclic alkyl groups such as, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl groups. In some embodiments, the cycloalkyl group can have 3 to about 8-12 ring members, whereas in other embodiments the number of ring carbon atoms range from 3 to 4, 5, 6, or 7. Cycloalkyl groups further include polycyclic cycloalkyl groups such as, but not limited to, norbornyl, adamantyl, bornyl, camphenyl,

isocamphenyl, and carenyl groups, and fused rings such as, but not limited to, decalanyl, and the like. Cycloalkyl groups also include rings that are substituted with straight or branched chain alkyl groups as defined herein. Representative substituted cycloalkyl groups can be mono-substituted or substituted more than
5 once, such as, but not limited to, 2,2-, 2,3-, 2,4- 2,5- or 2,6-disubstituted cyclohexyl groups or mono-, di- or tri-substituted norbornyl or cycloheptyl groups, which can be substituted with, for example, amino, hydroxy, cyano, carboxy, nitro, thio, alkoxy, and halogen groups. The term "cycloalkenyl" alone or in combination denotes a cyclic alkenyl group.

10 Unless specifically indicated to the contrary, the cycloalkyl ring can be substituted with as many as n-1 substituents wherein n is the size of the carbocyclic ring with, for example, alkyl, alkenyl, alkynyl, amino, aryl, hydroxy, cyano, carboxy, heteroaryl, heterocyclyl, nitro, thio, alkoxy, and halogen groups, or other groups as are listed above.

15 (Cycloalkyl)alkyl groups, also denoted cycloalkylalkyl, are alkyl groups as defined herein in which a hydrogen or carbon bond of the alkyl group is replaced with a bond to a cycloalkyl group as defined above.

Alkenyl groups include straight and branched chain alkyl groups as defined above, except that at least one double bond exists between two carbon atoms. Thus,
20 alkenyl groups have from 2 to about 20 carbon atoms, and typically from 2 to 12 carbons or, in some embodiments, from 2 to 8 carbon atoms. Examples include, but are not limited to vinyl, $-\text{CH}=\text{CH}(\text{CH}_3)$, $-\text{CH}=\text{C}(\text{CH}_3)_2$, $-\text{C}(\text{CH}_3)=\text{CH}_2$, $-\text{C}(\text{CH}_3)=\text{CH}(\text{CH}_3)$, $-\text{C}(\text{CH}_2\text{CH}_3)=\text{CH}_2$, butadienyl, pentadienyl, and hexadienyl among others.

25 Cycloalkenyl groups include cycloalkyl groups having at least one double bond between 2 carbons. Thus for example, cycloalkenyl groups include but are not limited to cyclohexenyl, cyclopentenyl, and cyclohexadienyl groups. Cycloalkenyl groups can have from 3 to about 8-12 ring members, whereas in other embodiments the number of ring carbon atoms range from 3 to 5, 6, or 7.

30 Cycloalkenyl groups further include polycyclic cycloalkyl groups such as, but not limited to, norbornyl, adamantyl, bornyl, camphenyl, isocamphenyl, and

carenyl groups, and fused rings such as, but not limited to, decaliny, and the like, provided they include at least one double bond within a ring. Cycloalkenyl groups also include rings that are substituted with straight or branched chain alkyl groups as defined above.

- 5 (Cycloalkenyl)alkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of the alkyl group is replaced with a bond to a cycloalkenyl group as defined above.

Alkynyl groups include straight and branched chain alkyl groups, except that at least one triple bond exists between two carbon atoms. Thus, alkynyl groups
10 have from 2 to about 20 carbon atoms, and typically from 2 to 12 carbons or, in some embodiments, from 2 to 8 carbon atoms. Examples include, but are not limited to $-\text{C}\equiv\text{CH}$, $-\text{C}\equiv\text{C}(\text{CH}_3)$, $-\text{C}\equiv\text{C}(\text{CH}_2\text{CH}_3)$, $-\text{CH}_2\text{C}\equiv\text{CH}$, $-\text{CH}_2\text{C}\equiv\text{C}(\text{CH}_3)$, and $-\text{CH}_2\text{C}\equiv\text{C}(\text{CH}_2\text{CH}_3)$ among others.

The term "heteroalkyl" by itself or in combination with another term means,
15 unless otherwise stated, a straight or branched chain alkyl group consisting of the stated number of carbon atoms and one or two heteroatoms selected from the group consisting of non-peroxide O, N(R), and S, and wherein the nitrogen and sulfur atoms may be optionally oxidized and the nitrogen heteroatom may be optionally quaternized. R can be H, alkyl, aryl, aralkyl or a suitable protecting group. The
20 heteroatom(s) may be placed at any position of the heteroalkyl group, including between the rest of the heteroalkyl group and the fragment to which it is attached, as well as attached to the most distal carbon atom in the heteroalkyl group. Examples include: $-\text{CH}_2-\text{CH}_2\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{NH}-\text{CH}_3$, $-\text{CH}_2-\text{S}-\text{CH}_2-\text{CH}_3$, $-\text{CH}_2\text{CH}_2-\text{S}(=\text{O})-\text{CH}_3$, and $-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-\text{CH}_3$. Up to two heteroatoms
25 may be consecutive, such as, for example, $-\text{CH}_2-\text{NH}-\text{OCH}_3$, or $-\text{CH}_2-\text{CH}_2-\text{S}-\text{S}-\text{CH}_3$.

A "heterocycloalkyl" ring is a cycloalkyl ring containing at least one heteroatom. A heterocycloalkyl ring can also be termed a "heterocyclyl," described below.

The term "heteroalkenyl" by itself or in combination with another term
30 means, unless otherwise stated, a stable straight or branched chain monounsaturated or di-unsaturated hydrocarbon group consisting of the stated number of carbon

atoms and one or two heteroatoms selected from the group consisting of non-peroxide O, N(R'), and S, and wherein the nitrogen and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. Up to two heteroatoms may be placed consecutively. Examples include -CH=CH-O-CH₃,
5 -CH=CH-CH₂-OH, -CH₂-CH=N-OCH₃, -CH=CH-N(CH₃)-CH₃,
-CH₂-CH=CH-CH₂-SH, and and -CH=CH-O-CH₂CH₂-O-CH₃.

Aryl groups are (C₆-C₁₂)cyclic aromatic hydrocarbons that do not contain heteroatoms in the ring. Thus aryl groups include, but are not limited to, phenyl, azulenyl, heptalenyl, biphenyl, indacenyl, fluorenyl, phenanthrenyl, triphenylenyl,
10 pyrenyl, naphthacenyl, chrysenyl, biphenylenyl, anthracenyl, and naphthyl groups. In some embodiments, aryl groups contain about 6 to about 14 carbons in the ring portions of the groups. Aryl groups can be unsubstituted or substituted, as defined herein. Representative substituted aryl groups can be mono-substituted or substituted more than once, such as, but not limited to, 2-, 3-, 4-, 5-, or 6-substituted
15 phenyl or 2-8 substituted naphthyl groups, which can be substituted with carbon or non-carbon groups such as those listed above.

Aralkyl groups are alkyl groups as defined above in which a hydrogen or carbon bond of an alkyl group is replaced with a bond to an aryl group as defined above. Representative aralkyl groups include benzyl and phenylethyl groups and
20 fused (cycloalkylaryl)alkyl groups such as 4-ethyl-indanyl. Aralkenyl group are alkenyl groups as defined above in which a hydrogen or carbon bond of an alkyl group is replaced with a bond to an aryl group as defined above.

The term "heteroaryl" refers to a cyclic aromatic hydrocarbon containing at least one heteroatom, e.g., N, S or non-peroxide O, in the ring. Heteroaralkyl refers
25 to a heteroaromatic ring connected to an alkyl moiety, e.g., pyrid-2-yl methyl.

All chiral, diastereomeric, racemic forms of a structure are intended, unless a particular stereochemistry or isomeric form is specifically indicated. Compounds used in the present invention can include enriched or resolved optical isomers at any or all asymmetric atoms as are apparent from the depictions, at any degree of
30 enrichment. Both racemic and diastereomeric mixtures, as well as the individual optical isomers can be isolated or synthesized so as to be substantially free of their

enantiomeric or diastereomeric partners, and these are all within the scope of the invention.

Selected substituents within the compounds described herein are present to a recursive degree. In this context, "recursive substituent" means that a substituent
5 may recite another instance of itself or of another substituent that itself recites the first substituent. Because of the recursive nature of such substituents, theoretically, a large number may be present in any given claim. One of ordinary skill in the art of medicinal chemistry and organic chemistry understands that the total number of such substituents is reasonably limited by the desired properties of the compound
10 intended. Such properties include, by of example and not limitation, physical properties such as molecular weight, solubility or log P, application properties such as activity against the intended target, and practical properties such as ease of synthesis.

Recursive substituents are an intended aspect of the disclosed subject matter.
15 One of ordinary skill in the art of medicinal and organic chemistry understands the versatility of such substituents. To the degree that recursive substituents are present in a claim of the disclosed subject matter, the total number should be determined as set forth above.


A "salt" as is well known in the art includes an organic compound such as a
20 carboxylic acid, a sulfonic acid, or an amine, in ionic form, in combination with a counterion. For example, acids in their anionic form can form salts with cations such as metal cations, for example sodium, potassium, and the like; with ammonium salts such as NH_4^+ or the cations of various amines, including tetraalkyl ammonium salts such as tetramethylammonium, or other cations such as trimethylsulfonium,
25 and the like.

All patents cited herein are incorporated by reference herein as though fully set forth.

What is Claimed is:

1. A method for preparing a 5-acylpyrimidine comprising:


5 (a) reacting a pyrimidine of formula (II):




wherein X is halo or a sulfonate leaving group, and a compound of formula (III), (IV) or (V),




10 wherein R^1 is an organic group, M is a metal cation, each X is halo, Y is (C₁-C₄)alkoxy and x, y and z are selected so that the net charge of the compound (V) is 0, wherein the coupling is carried out in the presence of an effective catalytic amount of a complexed metal, to yield a compound of formula (IV);



15 (b) oxidizing the compound of formula (VI) to yield a 5-acylpyrimidine compound of formula (VII):




2. The method of claim 1 further comprising reacting the compound of formula
20 VII with a compound of the formula $(Li(R^2-ZnX)MgX_2)_y \cdot$ solvate wherein R^2 is an organic group and each X is halo, and x and y are selected so that the low-lying d-orbitals of the transition metal are filled, to yield a compound of formula (I):



3. The method of claim 1 wherein X is halo.
- 5 4. The method of claim 3 wherein X is Br or Cl.
5. The method of claim 1 wherein M in compound (III) is Mg(II), Zn(II), Mn(II), Cr(I), Sn(IV) or Li(I).
- 10 6. The method of claim 1 wherein Y in formula (IV) is methoxy.
7. The method of claim 1 wherein M^{++} in compound (V) is Zn^{++} .
8. The method of claim 1 wherein R^1 is aryl or (substituted)aryl.
- 15 9. The method of claim 8 wherein R^1 is (substituted)phenyl.
10. The method of claim 1 wherein the metal in the complexed metal is palladium, nickel or iron.
- 20 11. The method of claim 10 wherein the metal is complexed with triphenylphosphine or acetylacetone.
12. The method of claim 11 wherein the catalyst is $(Ph_3P)_4Pd$, $(PPh_3)_2NiCl_2$ or
- 25 Fe(acac)₂.
13. The method of claim 2 wherein the compound of formula (VI) is oxidized with SeO_2 , $KMnO_4$ or CrO_3 .

14. The method of claim 1 wherein the solvate is THF or diethyl ether.
15. The method of claim 1 wherein R^1 is phenyl, 4-trifluorophenyl, 4-methoxyphenyl or 4-halophenyl or 4-trifluoromethylphenyl.
16. The method of claim 15 wherein R^1 is 4-chlorophenyl.
17. The method of claim 16 wherein R^2 is isopropyl, halo-substituted phenyl or cyclopropyl.
18. The method of claim 17 wherein R^2 is 2-chlorophenyl or 3,5-dichlorophenyl.
19. The method of claim 2 wherein R^1 is 4-trifluoromethylphenyl and R^2 is isopropyl.
20. The method of claim 2 wherein R^1 is 4-chlorophenyl or 4-fluorophenyl and R^2 is 2-chlorophenyl.
21. The method of claim 2 wherein R^1 is 4-methoxyphenyl and R^2 is cyclopropyl.
22. The method of claim 2 wherein R^1 is phenyl and R^2 is 3,4-dichlorophenyl.
23. A compound of formula (IX):



wherein Y is O or (H)₂, Z is (C₁-C₄)perfluoroalkyl, (C₁-C₄)alkoxy or halo.

24. The compound of claim 23 wherein Z is F₃C-.
25. The compound of claim 23 wherein Z is Cl.
- 5 26. The compound of claim 23 wherein R is methoxy.