



HU000027088T2


(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 027 088**(13) **T2****EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 11 290269**(22) A bejelentés napja: **2011. 06. 14.**(96) Az európai bejelentés bejelentési száma:
EP 20110290269(97) Az európai bejelentés közzétételi adatai:
EP 2404600 A1 **2012. 01. 11.**(97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2404600 B1 **2015. 05. 27.**(51) Int. Cl.: **A61K 3140/45** (2006.01)**A61K 31/55** (2006.01)**A61K 45/06** (2006.01)**A61P 9/04** (2006.01)(30) Elsőbbségi adatok:
1002525 **2010. 06. 15.** **FR**(72) Feltalálók(k):
Thuillez, Christian, 76000 Rouen (FR)
Mulder, Paulus, 76130 Mont Saint Aignan (FR)
Vilaine, Jean-Paul, 92290 Chatenay Malabry (FR)
Fratacci, Marie-Dominique, 78390 Bois d'Arcy (FR)
Lerebours-Piegeonniere, Guy, 92300 Levallois-Perret (FR)
Feldmann, Luc, 75017 Paris (FR)
Roussel, Jérôme, 78220 Viroflay (FR)(73) Jogosult(ak):
Les Laboratoires Servier, 92284 Suresnes Cedex (FR)(74) Képviselő:
dr. Török Ferenc, DANUBIA Szabadalmi és Jogi Iroda Kft., Budapest(54) **Szinuszos If-áram inhibitor és az angiotenzin konverzió enzim inhibitor kombinációjának alkalmazása megtartott szisztolés funkcióval rendelkező szívelégtelenség kezelésére**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.


The present invention relates to the use of the association of:

- ivabradine, or 3-{3-[[[(7S)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]-methyl](methyl)amino]propyl}-7,8-dimethoxy-1,3,4,5-tetrahydro-2H-3-benzazepin-2-one, of formula (I):



and its addition salts with a pharmaceutically acid, their hydrates and crystalline forms, or

- *N*-{[(7S)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]methyl}-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3H-3-benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine of formula (II)



and its addition salts with a pharmaceutically acid, their hydrates and crystalline forms, with perindopril, or one of its addition salts with a pharmaceutically acceptable base, their hydrates or crystalline forms, in obtaining medicaments intended for the treatment of heart failure with preserved systolic function.

Ivabradine, and its addition salts with a pharmaceutically acceptable acid, and more especially its hydrochloride, their hydrates and crystalline forms, and *N*-{[(7S)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]methyl}-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3H-3-benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine, and its addition salts with a pharmaceutically acceptable acid, and more especially its hydrochloride and its fumarate, their hydrates and crystalline forms, are selective and specific sinus node I_f current inhibitors and have very valuable pharmacological and therapeutic properties, especially negative chronotropic (heart-rate-reducing) properties, making those compounds useful in treating, preventing and improving the prognosis of various cardiovascular diseases associated with myocardial ischaemia such as angina pectoris, myocardial infarction and associated rhythm disturbances, and also in various pathologies involving rhythm disturbances, especially supraventricular rhythm disturbances, and in chronic heart failure.

The preparation and therapeutic use of ivabradine and its addition salts with a pharmaceutically acceptable acid, more especially its hydrochloride, have been described in European patent specification EP 0 534 859.

The preparation and therapeutic use of *N*-{[(7*S*)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl)methyl}-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3*H*-3-benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine and its addition salts with a pharmaceutically acceptable acid, and more especially its hydrochloride and its fumarate, have been described in European patent specification EP 2 036 892.

Perindopril is an angiotensin-converting enzyme inhibitor.

Angiotensin-converting enzyme inhibitors are one of the major therapeutic classes in the treatment of arterial hypertension. They act principally by inhibiting the synthesis of angiotensin II and by blocking the breakdown of bradykinin.

In addition to the lowering of arterial pressure, they have been shown to improve the morbidity (myocardial infarction, cerebral vascular accidents) and cardiovascular mortality of hypertensive patients, diabetic patients and patients with pre-existing coronary disease.

Amongst the pharmaceutically acceptable acids there may be mentioned, without implying any limitation, hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, acetic acid, trifluoroacetic acid, lactic acid, pyruvic acid, malonic acid, succinic acid, glutaric acid, fumaric acid, tartaric acid, maleic acid, citric acid, ascorbic acid, oxalic acid, methanesulphonic acid, benzenesulphonic acid, camphoric acid, pamoic acid and 1,5-naphthalenedisulphonic acid.

The Applicant has discovered that the association of:

- ivabradine, or 3-{3-[[[(7*S*)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl)methyl](methyl)amino]propyl}-7,8-dimethoxy-1,3,4,5-tetrahydro-2*H*-3-benzazepin-2-one, or
- *N*-{[(7*S*)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl)methyl}-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3*H*-3-benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine,

and perindopril has valuable properties allowing its use in the treatment of heart failure with preserved systolic function.

Heart failure due to systolic dysfunction of the left ventricle is not the only form of heart failure. Increasingly often, patients with heart failure have an ejection fraction which is greater than 40 %. The proportion of heart failure referred to as "diastolic heart failure" (or rather "heart failure with preserved systolic function") increases with age. It currently accounts for 30 to 40 % of hospital admissions for heart failure and, after the age of 80, its frequency exceeds that of heart failures due to systolic dysfunction. Diastolic heart failures generally feature both prolonged ventricular relaxation and a reduction in the distensibility of the left ventricle chamber. The basic causes are ischaemic, hypertensive and elderly-patient cardiopathies. Predisposing factors are age, sex (female), diabetes, obesity and arterial hypertension. Concentric remodelling of the left ventricle, with or without hypertrophy, consistently gives rise to disruption of diastolic function. In most cases a triggering factor is found to be the cause of a congestive attack. The frequency of "diastolic" heart failure increases with age. Its physiopathology remains complex and merits being better understood by clinicians.

No treatment has hitherto demonstrated efficacy in this pathology, the mortality (50 % at 4 years) of which corresponds to that of systolic heart failure.

The Applicant has discovered that the use of the association of ivabradine or *N*-{[(7*S*)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl)methyl}-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3*H*-3-

benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine and perindopril makes it possible to obtain pharmacological effects that are superior to those observed when using either ivabradine or *N*-{[(7*S*)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]methyl}-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3*H*-3-benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine on their own or perindopril on its own. The use of the association of ivabradine or *N*-{[(7*S*)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]methyl}-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3*H*-3-benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine and perindopril moreover makes it possible for the observed physiological parameters to return to values very close to normal. These observations make it possible to envisage using the association of ivabradine or *N*-{[(7*S*)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]methyl}-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3*H*-3-benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine and perindopril in the treatment of heart failure having preserved systolic function.

Perindopril is used as such or in the form of one of its addition salts with a pharmaceutically acceptable acid or base, and more especially its *tert*-butylamine or arginine salts, their hydrates and crystalline forms.

The present invention relates also to pharmaceutical compositions comprising as active ingredients:

- ivabradine, or one of its hydrates, crystalline forms, and addition salts with a pharmaceutically acceptable acid and more especially its hydrochloride, or *N*-{[(7*S*)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]methyl}-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3*H*-3-benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine or one of its addition salts with a pharmaceutically acceptable acid, and more especially its hydrochloride and its fumarate, their hydrates or crystalline forms, and
- perindopril, or one of its addition salts with a pharmaceutically acceptable base, and more especially its *tert*-butylamine or arginine salts, their hydrates or crystalline forms,

for use in the treatment of heart failure with preserved systolic function.

The pharmaceutical compositions that may be used are those that are suitable for oral, parenteral or nasal administration, tablets, dragées, sublingual tablets, capsules, lozenges, suppositories, creams, ointments, dermal gels etc. and also pharmaceutical compositions having programmed, delayed, prolonged or deferred release.

Besides ivabradine or *N*-{[(7*S*)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]methyl}-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3*H*-3-benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine and perindopril, said pharmaceutical compositions comprise one or more excipients or carriers selected from diluents, lubricants, binders, disintegration agents, absorbents, colourants, sweeteners etc..

By way of non-limiting example there may be mentioned :

- as *diluents*: lactose, dextrose, sucrose, mannitol, sorbitol, cellulose, glycerol,
- as *lubricants*: silica, talc, stearic acid and its magnesium and calcium salts, polyethylene glycol,

- * *as binders*: magnesium aluminium silicate, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose and polyvinylpyrrolidone,
- * *as disintegrants*: agar, alginic acid and its sodium salt, effervescent mixtures.

The useful dosage varies according to the sex, age and weight of the patient, the administration route, the nature of the disorder and of any associated treatments and ranges from 2.5 to 30 mg of ivabradine per 24 hours, and more preferably from 5 to 15 mg per day, and even more preferably from 10 to 15 mg per day. The dose of N-[[[(7S)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]methyl]-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3H-3-benzazepin-3-yl)-N-methyl-3-oxo-1-propanamine (hereinafter referred to as compound A) may vary from 5 to 100 mg per day. The dose of perindopril may be less than that used when it is administered on its own.

The daily dose of perindopril will preferably be between 1 and 10 mg inclusive.

The Examples that follow illustrate the invention.

List of abbreviations used

dP/dt _{max}	:	maximum increase in pressure per second
dP/dt _{min}	:	maximum reduction in pressure per second
HF	:	heart failure
LVEDP	:	Left Ventricular End Diastolic Pressure
LVEDPVR	:	Left Ventricular End Diastolic Pressure Volume Relation
LVESP	:	Left Ventricular End Systolic Pressure
LVESPVR	:	Left Ventricular End Systolic Pressure Volume Relation
LV	:	left ventricle

Pharmacological tests:

Heart failure is induced in rats by ligation of the left coronary artery (the control animals undergo an operation but are not ligated), which causes ischaemia of part of the wall of the left ventricle. The animals recuperate for 7 days and then, for 12 weeks, they are given either 3 mg/kg of compound A, or 0.4 mg/kg of perindopril, or perindopril and compound A concomitantly.

Twelve weeks after the operation it is found that the animals having undergone coronary ligation develop heart failure that is both systolic (ejection anomaly) and diastolic (filling anomaly).

In those animals, compound A, on its own or in association with perindopril, makes possible a significant reduction in heart rate (Table 1 and Figure 1).

Table 1

		HF (untreated)		HF + A	HF + perindopril	HF + A + perindopril
Heart rate (bpm)	Duration of treatment	4 weeks	372.5	349.3	388.2	352.8
		12 weeks	387.2	342.7 [†]	387.7	353.1 [†]

[†] p<0.05 vs HF

Co-treatment with perindopril and compound A makes possible a significant increase in the shortening fraction of the left ventricle, that is to say an improvement in its contractility (Table 2 and Figure 2). Consequently, heart rate is improved compared to animals with heart failure that are not given treatment.

Table 2

	HF (untreated)	HF + A	HF + perindopril	HF + A + perindopril
shortening fraction (% of the diameter of the LV)	14.4	18.0	17.1	22.3 [†]
heart rate (mL/min)	114	127	142 [†]	140 [†]

[†] p<0.05 vs HF

As Table 3 shows (Figure 3), the various systolic and diastolic parameters are modified by heart failure. The left ventricle contracts less well (dP/dt_{max} and LVESPVR significantly lower in the HF animals than in the healthy controls), which indicates systolic impairment. There is a major deterioration in diastolic function: the pressure inside the ventricle at the end of diastole is raised (LVEDP), the relaxation time (tau) is lengthened and the compliance (ability of the ventricle to distend) is low (LVEDPVR increased).

Table 3

	Control	HF (untreated)	HF + A	HF + perindopril	HF + A + perindopril
LVESP (mm Hg)	140	120	118	99	105
dP/dt _{max} (10 ³ mm Hg/s)	9.92	6.89*	6.78	5.97	7.69
LVESPVR (mm Hg/RVU)	26.4	11.1*	16.1 [†]	16.4 [†]	15.6 [†]
LVEDP (mm Hg)	1.86	9.43*	4.89 [†]	5.17 [†]	3.32 [†]
dP/dt _{min} (-10 ³ mm Hg/s)	10.24	5.66*	5.87	5.11	6.19
tau (ms)	3.54	12.64*	8.37 [†]	7.31 [†]	6.05 [†]
LVEDPVR (mm Hg/RVU)	0.84	6.93*	2.70 [†]	2.36 [†]	1.37 ^{††}

* p<0.05 vs control; [†] p<0.05 vs HF; ^{††} p<0.05 vs HF+A and vs HF+perindopril

It is found that treatment of the animals which have heart failure, whether with perindopril on its own or with compound A on its own, improves systolic function, which can be seen from the LVESPVR, the only load-independent parameter.

The end diastolic pressure and relaxation time are clearly improved by perindopril on its own or by compound A on its own, and a tendency to a further reduction in those two parameters is noted when the two substances are administered together. The compliance of the left ventricle (measured by LVEDPVR), the

only load-independent parameter, is very clearly improved by perindopril and by compound A. Surprisingly, this effect is significantly increased when the animals are given the two treatments concomitantly.

In fact, the association of compound A and perindopril makes it possible to significantly improve the compliance, which returns to a level close to that of the control animals.

The association of perindopril and N-[[[(7S)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]methyl]-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3H-3-benzazepin-3-yl)-N-methyl-3-oxo-1-propanamine accordingly makes it possible to improve the deterioration in diastolic function.

This effect on systolic function and diastolic function was then studied using the association of perindopril and a different I_f current inhibitor, ivabradine.

It is found that treatment with perindopril on its own or in association with ivabradine improves the systolic function (Table 4a and Figure 4a).

With respect to the diastolic dysfunction, treatment with perindopril and ivabradine is clearly more effective than perindopril on its own (the effect of ivabradine on its own is comparable to that of perindopril on its own, cf. Table 4b and Figure 4b). The compliance of the left ventricle is returned to a level similar to that of the healthy animals.

Table 4a

	Control	HF (untreated)	HF + perindopril	HF + ivabradine + perindopril
LVEBP (mm Hg)	163	134*	102 [†]	100 [†]
dP/dt _{max} (10 ³ mm Hg/s)	10.11	7.68*	6.08 [†]	6.10 [†]
LVEBPVR (mm Hg/RVU)	20.2	6.6*	14.5 [†]	12.6 ^{††}
LVEDP (mm Hg)	3.29	13.93*	6.88 [†]	5.01 [†]
dP/dt _{min} (-10 ³ mm Hg/s)	10.63	5.54*	4.99	4.97
tau (ms)	3.21	14.29*	10.92 [†]	8.52 [†]
LVEDPVR (mm Hg/RVU)	0.79	4.06*	2.25 [†]	1.15 ^{††}

* p<0.05 vs control; [†] p<0.05 vs HF; ^{††} p<0.05 vs HF+A and vs HF+perindopril

Table 4b

	Control	HF (untreated)	HF + ivabradine
LVEBPVR (mm Hg/RVU)	35.53	9.66*	20.63* [†]
LVEDPVR (mm Hg/RVU)	0.85	5.33*	1.87* [†]

* p<0.05 vs control; [†] p<0.05 vs HF

These experiments show that, in a model of heart failure, the association of ivabradine or N-[[[(7S)-3,4-dimethoxybicyclo[4.2.0]octa-1,3,5-trien-7-yl]methyl]-3-(7,8-dimethoxy-1,2,4,5-tetrahydro-3H-3-

benzazepin-3-yl)-*N*-methyl-3-oxo-1-propanamine and perindopril makes possible an improvement in diastolic function which is greater than that obtained with one of those two treatments used on its own, this improvement allowing a return to normal diastolic function.

Pharmaceutical compositions:

- 5 Formula for the preparation of 1000 tablets each containing as active ingredients 7.5 mg of ivabradine and 2 mg of perindopril *tert*-butylamine:

Ivabradine hydrochloride	8.085 g
Perindopril <i>tert</i> -butylamine	2 g
Lactose monohydrate	62 g
10 Magnesium stearate	1.3 g
Povidone	9 g
Anhydrous colloidal silica	0.3 g
Cellulose sodium glycolate	30 g
Stearic acid	2.6 g

- 15 Formula for the preparation of 1000 tablets each containing as active ingredients 10 mg of compound A and 2 mg of perindopril *tert*-butylamine:

Compound A fumarate	12.48 g
Perindopril <i>tert</i> -butylamine	2 g
Lactose monohydrate	62 g
20 Magnesium stearate	1.3 g
Povidone	9 g
Anhydrous colloidal silica	0.3 g
Cellulose sodium glycolate	30 g
Stearic acid	2.6 g

- 25 Other examples of pharmaceutical compositions according to the invention are given hereinbelow, without implying any limitation:

Example	Ivabradine (mg)	Compound A (mg)	Perindopril <i>tert</i> -butylamine salt (mg)	Perindopril arginine salt (mg)
1	10	-	2	-
2	15	-	4	-
3	10	-	-	2.5
4	15	-	-	5
5	-	60	2	-
6	-	80	4	-
7	-	60	-	2.5
8	-	80	-	5

Szabadalmi igénypontok

1. Asszociátuma a következőknek:

- ivabradin vagy 3-{3-[[[(7S)-3,4-dimetoxibiciklo[4.2.0]okta-1,3,5-trien-7-il]-metil](metil)amino]propil}-7,8-dimetoxi-1,3,4,5-tetrahidro-2H-3-benzazepin-2-on vagy gyógyszerészetileg elfogadható savakkal képzett addíciós sói egyike, hidrátjai és kristályformái,

vagy
- N-[[[(7S)-3,4-dimetoxibiciklo[4.2.0]okta-1,3,5-trien-7-il]metil]-3-(7,8-dimetoxi-1,2,4,5-tetrahidro-3H-3-benzazepin-3-il)-N-metil-3-oxo-1-propánamin vagy gyógyszerészetileg elfogadható savakkal képzett addíciós sói egyike, hidrátjai és kristályformái,

és perindopril vagy gyógyszerészetileg elfogadható bázisokkal képzett addíciós sói egyike, hidrátjai és kristályformái,

megtartott szisztolés funkcióval rendelkező szívelégtelenség kezelésében történő alkalmazásra.

2. Az 1. igénypont szerinti asszociátum az 1. igénypont szerinti kezelési eljárásban történő alkalmazásra, azzal jellemezve, hogy az ivabradin hidroklorid formában van, vagy ennek hidrátjai és kristályformái egyike.

3. Az 1. igénypont szerinti asszociátum az 1. igénypont szerinti kezelési eljárásban történő alkalmazásra, azzal jellemezve, hogy az N-[[[(7S)-3,4-dimetoxibiciklo[4.2.0]okta-1,3,5-trien-7-il]metil]-3-(7,8-dimetoxi-1,2,4,5-tetrahidro-3H-3-benzazepin-3-il)-N-metil-3-oxo-1-propánamin hidroklorid vagy fumarát formában van, vagy ezek hidrátjai és kristályformái egyike.

4. Az 1-3. igénypontok bármelyike szerinti asszociátum az 1. igénypont szerinti kezelési eljárásban történő alkalmazásra, azzal jellemezve, hogy a perindopril terc-butilamin- vagy argininsó formában van, vagy ezek hidrátjai és kristályformái egyike.

5. Az 1. igénypont szerinti asszociátum az 1. igénypont szerinti kezelési eljárásban történő alkalmazásra, azzal jellemezve, hogy az ivabradin hidroklorid formában van, vagy ennek hidrátjai és kristályformái egyike, és a perindopril terc-butilamin- vagy argininsó formában van, vagy ezek hidrátjai és kristályformái egyike.

6. Az 1. igénypont szerinti asszociátum az 1. igénypont szerinti kezelési eljárásban történő alkalmazásra, azzal jellemezve, hogy az N-[[[(7S)-3,4-dimetoxibiciklo[4.2.0]okta-1,3,5-trien-7-il]metil]-3-(7,8-dimetoxi-1,2,4,5-tetrahidro-3H-3-benzazepin-3-il)-N-metil-3-oxo-1-propánamin hidroklorid vagy fumarát formában van, vagy ezek hidrátjai és kristályformái egyike, és a perindopril terc-butilamin- vagy argininsó formában van, vagy ezek hidrátjai és kristályformái egyike.

7. Gyógyszerészeti készítmény, amely tartalmazza hatóanyagként a következőket:

- ivabradin, amely hidroklorid formában van, vagy ennek hidrátjai és kristályformái egyike, és
- a perindopril, amely terc-butilamin- vagy argininsó formában van, vagy ezek hidrátjai és kristályformái egyike,

magukban vagy kombinációban egy vagy több gyógyszerészetileg elfogadható segédanyaggal, megtartott szisztolés funkcióval rendelkező szívelégtelenség kezelésében történő alkalmazásra.

8. Gyógyszerészeti készítmény, amely tartalmazza hatóanyagként a következőket:

- N-[[[(7S)-3,4-dimetoxibicyclo[4.2.0]okta-1,3,5-trien-7-il]metil]-3-(7,8-dimetoxi-1,2,4,5-tetrahidro-3H-3-benzazepin-3-il)-N-metil-3-oxo-1-propánamin, amely hidroklorid vagy fumarát formában van, vagy ezek hidrátjai és kristályformái egyike, és

- a perindopril, amely terc-butilamin- vagy argininsó formában van, vagy ezek hidrátjai és kristályformái egyike,

magukban vagy kombinációban egy vagy több gyógyszerészetileg elfogadható segédanyaggal, megtartott szisztolés funkcióval rendelkező szívelégtelenség kezelésében történő alkalmazásra.

Figure 1




Figure 2




Figure 3




Figure 4a




Figure 4b

