



(51) International Patent Classification:

C07D 405/10 (2006.01) *C07D 413/14* (2006.01)
C07D 405/12 (2006.01) *C07D 417/14* (2006.01)
C07D 405/14 (2006.01) *A61K 31/397* (2006.01)
C07D 307/84 (2006.01) *A61K 31/496* (2006.01)
C07D 401/10 (2006.01) *A61P 31/14* (2006.01)
C07D 401/14 (2006.01)

(21) International Application Number:

PCT/US2016/018171

(22) International Filing Date:

17 February 2016 (17.02.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/118,057 19 February 2015 (19.02.2015) US

(71) Applicant: **BRISTOL-MYERS SQUIBB COMPANY**
[US/US]; Route 206 and Province Line Road, Princeton,
New Jersey 08543 (US).(72) Inventors: **WANG, Tao**; c/o Bristol-Myers Squibb Com-
pany, 5 Research Parkway, Wallingford, Connecticut
06492 (US). **YIN, Zhiwei**; c/o Bristol-Myers Squibb Com-
pany, 5 Research Parkway, Wallingford, Connecticut
06492 (US). **KADOW, John F.**; 9 Quarry Run, Walling-
ford, CT 06429 (US).(74) Agents: **VOLLES, Warren** et al.; Bristol-Myers Squibb
Company, Route 206 and Province Line Road, Princeton,
New Jersey 08543 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))

(54) Title: BENZOFURANS SUBSTITUTED WITH SECONDARY BENZAMIDE AS HCV INHIBITORS



I

(57) Abstract: Compounds of Formula I, including their salts, as well as compositions and methods of using the compounds are set forth. The compounds have activity against hepatitis C virus (HCV) and are useful in treating those infected with HCV.

BENZOFURANS SUBSTITUTED WITH SECONDARY BENZAMIDE AS HCV
INHIBITORS

CROSS REFERENCE TO RELATED APPLICATION

5 This application claims the benefit of U.S. Provisional Application Serial No. 62/118,057 filed February 19, 2015 which is herein incorporated by reference in its entirety.

FIELD OF THE INVENTION

10 The invention relates to novel compounds, including their salts, which have activity against hepatitis C virus (HCV) and which are useful in treating those infected with HCV. The invention also relates to compositions and methods of making and using these compounds.

BACKGROUND OF THE INVENTION

15 Hepatitis C virus (HCV) is a major human pathogen, infecting an estimated 170 million persons worldwide - roughly five times the number infected by human immunodeficiency virus type 1. A substantial fraction of these HCV infected individuals develop serious progressive liver disease, including cirrhosis and hepatocellular carcinoma (Lauer, G. M.; Walker, B. D. *N. Engl. J. Med.* 2001, 345, 41-52).

20 HCV is a positive-stranded RNA virus. Based on a comparison of the deduced amino acid sequence and the extensive similarity in the 5'-untranslated region, HCV has been classified as a separate genus in the Flaviviridae family. All members of the Flaviviridae family have enveloped virions that contain a positive stranded RNA genome encoding all known virus-specific proteins via translation of a single, uninterrupted, open reading frame.

25 Considerable heterogeneity is found within the nucleotide and encoded amino acid sequence throughout the HCV genome. At least six major genotypes have been characterized, and more than 50 subtypes have been described. The major genotypes of HCV differ in their distribution worldwide, and the clinical significance of the genetic

heterogeneity of HCV remains elusive despite numerous studies of the possible effect of genotypes on pathogenesis and therapy.

The single strand HCV RNA genome is approximately 9500 nucleotides in length and has a single open reading frame (ORF) encoding a single large polypeptide of about 3000 amino acids. In infected cells, this polypeptide is cleaved at multiple sites by cellular and viral proteases to produce the structural and non-structural (NS) proteins. In the case of HCV, the generation of mature non-structural proteins (NS2, NS3, NS4A, NS4B, NS5A, and NS5B) is effected by two viral proteases. The first one is believed to be a metalloprotease and cleaves at the NS2-NS3 junction; the second one is a serine protease contained within the N-terminal region of NS3 (also referred to as NS3 protease) and mediates all the subsequent cleavages downstream of NS3, both in cis, at the NS3-NS4A cleavage site, and in trans, for the remaining NS4A-NS4B, NS4B-NS5A, NS5A-NS5B sites. The NS4A protein appears to serve multiple functions, acting as a cofactor for the NS3 protease and possibly assisting in the membrane localization of NS3 and other viral replicase components. The complex formation of the NS3 protein with NS4A seems necessary to the processing events, enhancing the proteolytic efficiency at all of the sites. The NS3 protein also exhibits nucleoside triphosphatase and RNA helicase activities. NS5B (also referred to as HCV polymerase) is a RNA-dependent RNA polymerase that is involved in the replication of HCV. The HCV NS5B protein is described in "Structural Analysis of the Hepatitis C Virus RNA Polymerase in Complex with Ribonucleotides (Bressanelli; S. et al., *Journal of Virology* 2002, 3482-3492; and Defrancesco and Rice, *Clinics in Liver Disease* 2003, 7, 211-242).


Currently, the most effective HCV therapy employs a combination of alpha-interferon and ribavirin, leading to sustained efficacy in 40% of patients (Poynard, T. et al. *Lancet* 1998, 352, 1426-1432). Recent clinical results demonstrate that pegylated alpha-interferon is superior to unmodified alpha-interferon as monotherapy (Zeuzem, S. et al. *N. Engl. J. Med.* 2000, 343, 1666-1672). However, even with experimental therapeutic regimens involving combinations of pegylated alpha-interferon and ribavirin, a substantial fraction of patients do not have a sustained reduction in viral load. Thus, there is a clear and important need to develop effective therapeutics for treatment of HCV infection.

HCV-796, an HCV NS5B inhibitor, has shown an ability to reduce HCV RNA levels in patients. The viral RNA levels decreased transiently and then rebounded during dosing when treatment was with the compound as a single agent but levels dropped more robustly when combined with the standard of care which is a form of interferon and ribavirin. The development of this compound was suspended due to hepatic toxicity observed during extended dosing of the combination regimens. U.S. patent 7,265,152 and the corresponding PCT patent application WO2004/041201 describe compounds of the HCV-796 class. Other compounds have been disclosed; see for example, WO2009/101022, as well as WO 2012/058125.

What is therefore needed in the art are additional compounds which are novel and effective against hepatitis C. Additionally, these compounds should provide advantages for pharmaceutical uses, for example, with regard to one or more of their mechanism of action, binding, inhibition efficacy, target selectivity, solubility, safety profiles, or bioavailability. Also needed are new formulations and methods of treatment which utilize these compounds.

SUMMARY OF THE INVENTION

One aspect of the invention is a compound of Formula I, including pharmaceutically acceptable salts thereof:



I

wherein R^1 , R^2 are each independently selected from the group of hydrogen, halo, nitro, alkyl, cycloalkyl, haloalkyl, aminoalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyl, alkoxy, OR^{17} , cycloalkoxy, amino, alkylamino, dialkylamino, alkylcarboxamido, alkoxycarboxamido, alkoxyalkylcarboxamido, and Ar^1 ;

R³ is selected from the group of cyano, alkoxy carbonyl, (cycloalkyl)oxy carbonyl, (alkylsulfonyl)aminocarbonyl, CONR¹¹R¹², (R¹¹)(R¹²)NCONH, triazolyl, thiazolyl, and tetrazolyl;

R⁴ is phenyl substituted with 0-2 halo substituents;

5 R⁵ is selected from the group of hydrogen, alkyl, alkylsulfonyl, alkylcarbonyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, cyanoalkyl, cyanoalkenyl, cyanoalkynyl, cycloalkyl, halocycloalkyl, (haloalkyl)cycloalkyl, (halocycloalkyl)cycloalkyl, cyanocycloalkyl, (cyanoalkyl)cycloalkyl, (cyanocycloalkyl)cycloalkyl, hydroxycycloalkyl, (hydroxyalkyl)cycloalkyl, (hydroxycycloalkyl)cycloalkyl, alkoxycycloalkyl, (alkoxyalkyl)cycloalkyl, and (alkoxycycloalkyl)cycloalkyl;

10 R⁶ is selected from the group of hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, cyanoalkyl, cyanoalkenyl, cyanoalkynyl, cycloalkyl, halocycloalkyl, (haloalkyl)cycloalkyl, (halocycloalkyl)cycloalkyl, cyanocycloalkyl, (cyanoalkyl)cycloalkyl, (cyanocycloalkyl)cycloalkyl, hydroxycycloalkyl, (hydroxyalkyl)cycloalkyl, (hydroxycycloalkyl)cycloalkyl, alkoxycycloalkyl, (alkoxyalkyl)cycloalkyl, and (alkoxycycloalkyl)cycloalkyl;

15 R⁷R⁸N taken together is azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, dihydroindolyl, or isoindoliny, and is substituted with 0-2 substituents selected from the group of alkyl, hydroxyalkyl, alkoxyalkyl, hydroxy, alkoxy, carboxy, alkoxy carbonyl, dialkylcarboxamido, alkylcarbonylamino, alkoxy carbonylamino, pyridiny and phenyl, and is further substituted with one or more additional substituent(s) selected from the group of halo, haloalkyl, cyano, cyanoalkyl, cyanoalkenyl, 20 cyanocycloalkyl, cyanocycloalkenyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, urea, Ar² and phenyl groups, wherein said Ar² and phenyl groups are substituted with one or more substituents selected from the group of haloalkyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, 25 sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, and urea, and further wherein said pyridiny is substituted with one or more substituents selected from the group of halo, alkyl, haloalkyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl,

alkynyl, cycloalkenyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidinyl, guanidiny, carbamide, and urea;

or

R^7R^8N taken together is selected from the group of azepanyl, 1,4-diazepanyl, 1,4-oxazepanyl, 1,4-thiazepanyl, 1,1-dioxo 1,4-thiazepanyl, azocanyl, 1,4-diazocanyl, 1,5-diazocanyl, 1,4-oxazocanyl, 1,5-oxazocanyl, 1,4-thiazocanyl, 1,5-thiazocanyl, 1,1-dioxo 1,4-thiazocanyl, and 1,1-dioxo 1,5-thiazocanyl, and is substituted with 0-2 substituents selected from the group of halo, haloalkyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, alkyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, aminocarbonyl, alkoxy carbonyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidinyl, guanidiny, carbamide, urea, and Ar^3 ;

or

R^7R^8N taken together is selected from the group of azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, dihydroindolyl, isoindoliny, azepanyl, 1,4-diazepanyl, 1,4-oxazepanyl, 1,4-thiazepanyl, 1,1-dioxo 1,4-thiazepanyl, azocanyl, 1,4-diazocanyl, 1,5-diazocanyl, 1,4-oxazocanyl, 1,5-oxazocanyl, 1,4-thiazocanyl, 1,5-thiazocanyl, 1,1-dioxo 1,4-thiazocanyl, and 1,1-dioxo 1,5-thiazocanyl, and is substituted with a methylene group which is substituted with 0-2 substituents selected from the group of halo, haloalkyl, cyano, cyanoalkyl, cyanoalkenyl, cyanocycloalkyl, cyanocycloalkenyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidinyl, guanidiny, carbamide, urea, and Ar^3 ;

R^{11} is hydrogen, alkyl, or cycloalkyl;

R^{12} is hydrogen, alkyl, or cycloalkyl;

or R^{11} and R^{12} taken together with the nitrogen to which they are attached is azetidiny, pyrrolidiny, piperidiny, piperaziny, or morpholiny;

R^{17} is selected from the group of haloalkyl, cyanoalkyl, (cycloalkyl)alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, (R^{18})alkyl, (Ar^4)alkyl, alkynyl, and aminocycloalkyl;

R^{18} is selected from the group of $CONH_2$, H_2NCONH , dibenzylamino, phthalimido, amino, alkylamino, dialkylamino, azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny where azetidiny, pyrrolidiny, piperidiny, piperaziny, and morpholiny, and is substituted with 0-3 alkyl or alkoxy carbonyl substituents;

R²⁰ is hydrogen, halo, alkyl, or alkoxy;

R²¹ is hydrogen, halo, alkyl, or alkoxy;

Ar¹ is selected from the group of phenyl, naphthalenyl, quinoliny, isoquinoliny, quinoxaliny, quinazoliny, pyridiny, furanyl, thienyl, pyrroly, pyrazoly, isoxazoly, isothiazoly, imidazoly, oxazoly, thiazoly, oxadiazoly, oxadiathiazoly, triazoly, tetrazoly, pyraziny, pyrimidiny, pyridaziny, triazoly, indoly, azaindoly, indazoly, azaindazoly, benzoxazoly, benzoisoxazoly, benzoisorhiazoly, benzimidazoly, and benzothiazoly, and is substituted with 0-2 substituents selected from the group of halo, alkyl, cycloalkyl, haloalkyl, alkoxyalkyl, hydroxy, alkoxy, amino, alkylamino, dialkylamino, aminocarbonyl, pyridiny, phenyl, halophenyl, alkylphenyl, and alkoxyphenyl;

Ar² is selected from the group of naphthalenyl, quinoliny, isoquinoliny, quinoxaliny, quinazoliny, pyridiny, furanyl, thienyl, pyrroly, pyrazoly, isoxazoly, isothiazoly, imidazoly, oxazoly, thiazoly, oxadiazoly, oxadiathiazoly, triazoly, tetrazoly, pyraziny, pyridaziny, triazoly, indoly, azaindoly, indazoly, azaindazoly, benzoxazoly, benzoisoxazoly, benzoisorhiazoly, benzimidazoly, and benzothiazoly, and is substituted with 0-2 substituents selected from the group of halo, alkyl, cycloalkyl, haloalkyl, alkoxyalkyl, hydroxy, alkoxy, amino, alkylamino, dialkylamino, aminocarbonyl, halophenyl, alkylphenyl, alkoxyphenyl and Ar³;

Ar³ is selected from the group of phenyl, naphthalenyl, quinoliny, isoquinoliny, quinoxaliny, quinazoliny, pyridiny, furanyl, thienyl, pyrroly, pyrazoly, isoxazoly, isothiazoly, imidazoly, oxazoly, thiazoly, oxadiazoly, oxadiathiazoly, triazoly, tetrazoly, pyraziny, pyrimidiny, pyridaziny, triazoly, indoly, azaindoly, indazoly, azaindazoly, benzoxazoly, benzoisoxazoly, benzoisorhiazoly, benzimidazoly, and benzothiazoly, and is substituted with 0-2 substituents selected from the group of halo, alkyl, cycloalkyl, haloalkyl, alkoxyalkyl, hydroxy, alkoxy, amino, alkylamino, dialkylamino, aminocarbonyl, pyridiny, phenyl, halophenyl, alkylphenyl, and alkoxyphenyl; and

Ar⁴ is selected from the group of furanyl, thienyl, pyrroly, pyrazoly, isoxazoly, isothiazoly, imidazoly, oxadiazoly, thiadiazoly, triazoly, pyridiny, indoly, and phenyl, and is substituted with 0-2 substituents selected from the group of halo, alkyl, haloalkyl, hydroxyl, and alkoxy.

The invention also relates to pharmaceutical compositions comprising a compound of Formula I, including pharmaceutically acceptable salts thereof, and a pharmaceutically acceptable carrier, excipient and/or diluent.

5 In addition, the invention provides one or more methods of treating hepatitis C infection comprising administering a therapeutically effective amount of a compound of Formula I to a patient.

Also provided as part of the invention are one or more methods for making the
10 compounds of Formula I.

The present invention is directed to these, as well as other important ends, hereinafter described.

15 DETAILED DESCRIPTION OF THE EMBODIMENTS

The singular forms “a”, “an”, and “the” include plural reference unless the context clearly dictates otherwise.

Unless otherwise specifically set forth elsewhere in the application, the following terms can be used herein and shall have the following meanings: “Hydrogen” or “H”
20 refers to hydrogen, including its isotopes, such as deuterium. “Halo” means fluoro, chloro, bromo, or iodo. “Alkyl” means a straight or branched alkyl group composed of 1 to 6 carbons. “Alkenyl” means a straight or branched alkyl group composed of 2 to 6 carbons with at least one double bond. “Cycloalkyl” means a monocyclic ring system composed of 3 to 7 carbons. “Hydroxyalkyl,” “alkoxy” and other terms with a substituted
25 alkyl moiety include straight and branched isomers composed of 1 to 6 carbon atoms for the alkyl moiety. “Halo” includes all halogenated isomers from monohalo substituted to perhalo substituted in substituents defined with halo, for example, “Haloalkyl” and “haloalkoxy”, “halophenyl”, “halophenoxy.” “Aryl” means a monocyclic or bicyclic aromatic hydrocarbon groups having 6 to 12 carbon atoms, or a bicyclic fused ring
30 system wherein one or both of the rings is a phenyl group. Bicyclic fused ring systems consist of a phenyl group fused to a four- to six-membered aromatic or non-aromatic carbocyclic ring. Representative examples of aryl groups include, but are not limited to, indanyl, indenyl, naphthyl, phenyl, and tetrahydronaphthyl. “Heteroaryl” means a 5 to 7

membered monocyclic or 8 to 11 membered bicyclic aromatic ring system with 1-5 heteroatoms independently selected from nitrogen, oxygen, and sulfur. Parenthetic and multiparenthetic terms are intended to clarify bonding relationships to those skilled in the art. For example, a term such as ((R)alkyl) means an alkyl substituent further substituted with the substituent R. Substituents which are illustrated by chemical drawing to bond at variable positions on a multiple ring system (for example a bicyclic ring system) are intended to bond to the ring where they are drawn to append.


The invention includes all pharmaceutically acceptable salt forms of the compounds. Pharmaceutically acceptable salts are those in which the counter ions do not contribute significantly to the physiological activity or toxicity of the compounds and as such function as pharmacological equivalents. These salts can be made according to common organic techniques employing commercially available reagents. Some anionic salt forms include acetate, acistrate, besylate, bromide, camsylate, chloride, citrate, fumarate, glucuronate, hydrobromide, hydrochloride, hydroiodide, iodide, lactate, maleate, mesylate, nitrate, pamoate, phosphate, succinate, sulfate, tartrate, tosylate, and xinofoate. Some cationic salt forms include ammonium, aluminum, benzathine, bismuth, calcium, choline, diethylamine, diethanolamine, lithium, magnesium, meglumine, 4-phenylcyclohexylamine, piperazine, potassium, sodium, tromethamine, and zinc.

Some of the compounds of the invention possess asymmetric carbon atoms. The invention includes all stereoisomeric forms, including enantiomers and diastereomers as well as mixtures of stereoisomers such as racemates. Some stereoisomers can be made using methods known in the art. Stereoisomeric mixtures of the compounds and related intermediates can be separated into individual isomers according to methods commonly known in the art. The use of wedges or hashes in the depictions of molecular structures in the following schemes and tables is intended only to indicate relative stereochemistry, and should not be interpreted as implying absolute stereochemical assignments.

The invention is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include deuterium and tritium. Isotopes of carbon include ^{13}C and ^{14}C . Isotopically-

labeled compounds of the invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described herein, using an appropriate isotopically-labeled reagent in place of the non-labeled reagent otherwise employed. Such compounds can have a variety of potential uses, for example as standards and reagents in determining biological activity. In the case of stable isotopes, such compounds can have the potential to favorably modify biological, pharmacological, or pharmacokinetic properties.

As set forth above, the invention is directed to one or more compounds of Formula I, including pharmaceutically acceptable salts thereof:



I

wherein R¹, R² are each independently selected from the group of hydrogen, halo, nitro, alkyl, cycloalkyl, haloalkyl, aminoalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyl, alkoxy, OR¹⁷, cycloalkoxy, amino, alkylamino, dialkylamino, alkylcarboxamido, alkoxyalkylcarboxamido, and Ar¹;

R³ is selected from the group of cyano, alkoxycarbonyl, (cycloalkyl)oxy carbonyl, (alkylsulfonyl)aminocarbonyl, CONR¹¹R¹², (R¹¹)(R¹²)NCONH, triazolyl, thiazolyl, and tetrazolyl;

R⁴ is phenyl substituted with 0-2 halo substituents;

R⁵ is selected from the group of hydrogen, alkyl, alkylsulfonyl, alkylcarbonyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, cyanoalkyl, cyanoalkenyl, cyanoalkynyl, cycloalkyl, halocycloalkyl, (haloalkyl)cycloalkyl, (halocycloalkyl)cycloalkyl, cyanocycloalkyl, (cyanoalkyl)cycloalkyl, (cyanocycloalkyl)cycloalkyl, hydroxycycloalkyl, (hydroxyalkyl)cycloalkyl, (hydroxycycloalkyl)cycloalkyl, alkoxyalkyl, (alkoxyalkyl)cycloalkyl, and (alkoxycycloalkyl)cycloalkyl;

R⁶ is selected from the group of hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, cyanoalkyl,

cyanoalkenyl, cyanoalkynyl, cycloalkyl, halocycloalkyl, (haloalkyl)cycloalkyl, (halocycloalkyl)cycloalkyl, cyanocycloalkyl, (cyanoalkyl)cycloalkyl, (cyanocycloalkyl)cycloalkyl, hydroxycycloalkyl, (hydroxyalkyl)cycloalkyl, (hydroxycycloalkyl)cycloalkyl, alkoxy cycloalkyl, (alkoxyalkyl)cycloalkyl, and
 5 (alkoxycycloalkyl)cycloalkyl;

R^7R^8N taken together is azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, dihydroindoly, or isoindoliny, and is substituted with 0-2 substituents selected from the group of alkyl, hydroxyalkyl, alkoxyalkyl, hydroxy, alkoxy, carboxy, alkoxy carbonyl, dialkylcarboxamido, alkylcarbonylamino, alkoxy carbonylamino,
 10 pyridiny and phenyl, and is further substituted with one or more additional substituent(s) selected from the group of halo, haloalkyl, cyano, cyanoalkyl, cyanoalkenyl, cyanocycloalkyl, cyanocycloalkenyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, urea, Ar^2 and phenyl groups, wherein said Ar^2 and phenyl groups are
 15 substituted with one or more substituents selected from the group of haloalkyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, and urea, and further wherein said pyridiny is substituted with one or more substituents selected from the group of
 20 halo, alkyl, haloalkyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, and urea;
 or

R^7R^8N taken together is selected from the group of azepanyl, 1,4-diazepanyl, 1,4-oxazepanyl, 1,4-thiazepanyl, 1,1-dioxo 1,4-thiazepanyl, azocanyl, 1,4-diazocanyl, 1,5-diazocanyl, 1,4-oxazocanyl, 1,5-oxazocanyl, 1,4-thiazocanyl, 1,5-thiazocanyl, 1,1-dioxo 1,4-thiazocanyl, and 1,1-dioxo 1,5-thiazocanyl, and is substituted with 0-2 substituents selected from the group of halo, haloalkyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, alkyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, aminocarbonyl, alkoxy carbonyl, ester,
 25 amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, urea, and Ar^3 ;
 or

R⁷R⁸N taken together is selected from the group of azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, dihydroindolyl, isoindoliny, azepany, 1,4-diazepany, 1,4-oxazepany, 1,4-thiazepany, 1,1-dioxo 1,4-thiazepany, azocany, 1,4-diazocany, 1,5-diazocany, 1,4-oxazocany, 1,5-oxazocany, 1,4-thiazocany, 1,5-thiazocany, 1,1-dioxo 1,4-thiazocany, and 1,1-dioxo 1,5-thiazocany, and is substituted with a methylene group which is substituted with 0-2 substituents selected from the group of halo, haloalkyl, cyano, cyanoalkyl, cyanoalkenyl, cyanocycloalkyl, cyanocycloalkenyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, urea, and Ar³;

R¹¹ is hydrogen, alkyl, or cycloalkyl;

R¹² is hydrogen, alkyl, or cycloalkyl;

or R¹¹ and R¹² taken together with the nitrogen to which they are attached is azetidiny, pyrrolidiny, piperidiny, piperaziny, or morpholiny;

R¹⁷ is selected from the group of haloalkyl, cyanoalkyl, (cycloalkyl)alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, (R¹⁸)alkyl, (Ar⁴)alkyl, alkynyl, and aminocycloalkyl;

R¹⁸ is selected from the group of CONH₂, H₂NCONH, dibenzylamino, phthalimido, amino, alkylamino, dialkylamino, azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny where azetidiny, pyrrolidiny, piperidiny, piperaziny, and morpholiny, and is substituted with 0-3 alkyl or alkoxy carbonyl substituents;

R²⁰ is hydrogen, halo, alkyl, or alkoxy;

R²¹ is hydrogen, halo, alkyl, or alkoxy;

Ar¹ is selected from the group of phenyl, naphthalenyl, quinoliny, isoquinoliny, quinoxaliny, quinazoliny, pyridiny, furany, thienyl, pyrroly, pyrazolyl, isoxazolyl, isothiazolyl, imidazolyl, oxazolyl, thiazolyl, oxadiazolyl, oxadiathiazolyl, triazolyl, tetrazolyl, pyraziny, pyrimidiny, pyridaziny, triazolyl, indolyl, azaindolyl, indazolyl, azaindazolyl, benzoxazolyl, benzoisoxazolyl, benzoisorhiazolyl, benzimidazolyl, and benzothiazolyl, and is substituted with 0-2 substituents selected from the group of halo, alkyl, cycloalkyl, haloalkyl, alkoxyalkyl, hydroxy, alkoxy, amino, alkylamino, dialkylamino, aminocarbonyl, pyridiny, phenyl, halophenyl, alkylphenyl, and alkoxyphenyl;

Ar² is selected from the group of naphthalenyl, quinoliny, isoquinoliny, quinoxaliny, quinazoliny, pyridiny, furany, thieny, pyrroly, pyrazoly, isoxazoly, isothiazoly, imidazoly, oxazoly, thiazoly, oxadiazoly, oxadiathiazoly, triazoly, tetrazoly, pyraziny, pyridaziny, triazoly, indoly, azaindoly, indazoly, azaindazoly, benzoxazoly, benzoisoxazoly, benzoisorhiazoly, benzimidazoly, and benzothiazoly, and is substituted with 0-2 substituents selected from the group of halo, alkyl, cycloalkyl, haloalkyl, alkoxyalkyl, hydroxy, alkoxy, amino, alkylamino, dialkylamino, aminocarbonyl, halophenyl, alkylphenyl, alkoxyphenyl and Ar³;

Ar³ is selected from the group of phenyl, naphthalenyl, quinoliny, isoquinoliny, quinoxaliny, quinazoliny, pyridiny, furany, thieny, pyrroly, pyrazoly, isoxazoly, isothiazoly, imidazoly, oxazoly, thiazoly, oxadiazoly, oxadiathiazoly, triazoly, tetrazoly, pyraziny, pyrimidiny, pyridaziny, triazoly, indoly, azaindoly, indazoly, azaindazoly, benzoxazoly, benzoisoxazoly, benzoisorhiazoly, benzimidazoly, and benzothiazoly, and is substituted with 0-2 substituents selected from the group of halo, alkyl, cycloalkyl, haloalkyl, alkoxyalkyl, hydroxy, alkoxy, amino, alkylamino, dialkylamino, aminocarbonyl, pyridiny, phenyl, halophenyl, alkylphenyl, and alkoxyphenyl; and

Ar⁴ is selected from the group of furany, thieny, pyrroly, pyrazoly, isoxazoly, isothiazoly, imidazoly, oxadiazoly, thiadiazoly, triazoly, pyridiny, indoly, and phenyl, and is substituted with 0-2 substituents selected from the group of halo, alkyl, haloalkyl, hydroxyl, and alkoxy.

It is preferred that R¹ and R² are each hydrogen.

It is also preferred that R³ is -CONR¹¹,R¹². R¹¹ and R¹² are each independently preferred to be hydrogen or alkyl.

In addition, it is preferred that R⁴ is a phenyl group substituted with one fluoro (F) group.


It is further preferred that R⁵ is alkyl or haloalkyl, and R⁶ is alkylsulfonyl.


It is also preferred that R^7R^8N taken together is selected from the group of azetidiny, pyrrolidiny, piperidiny, and piperaziny.


In a further embodiment, it is preferred that R^{20} and R^{21} are each hydrogen.

5

Also preferred are compounds of Formula I, including pharmaceutically acceptable salts thereof, which are selected from the group of:







Pharmaceutical Compositions and Methods of Treatment

The compounds according to the various embodiments herein set forth demonstrate activity against HCV NS5B, and can be useful in treating HCV and HCV infection. Therefore, another aspect of the invention is a composition comprising a
5 compound of Formula I, or a pharmaceutically acceptable salt(s) thereof, and a pharmaceutically acceptable carrier, excipient and/or diluent.

Another aspect of the invention is a composition further comprising an additional compound having anti-HCV activity.

10

Another aspect of the invention is a composition where the compound having anti-HCV activity is an interferon or a ribavirin. Another aspect of the invention is wherein the interferon is selected from interferon alpha 2B, pegylated interferon alpha, consensus interferon, interferon alpha 2A, interferon lambda, and lymphoblastoid interferon tau.

15

Another aspect of the invention is a composition where the compound having anti-HCV activity is a cyclosporin. Another aspect of the invention is where the cyclosporin is cyclosporin A.

Another aspect of the invention is a composition where the compound having anti-HCV activity is selected from the group consisting of interleukin 2, interleukin 6, interleukin 12, a compound that enhances the development of a type 1 helper T cell response, interfering RNA, anti-sense RNA, Imiquimod, ribavirin, an inosine 5'-monophosphate dehydrogenase inhibitor, amantadine, and rimantadine.

25

Another aspect of the invention is a composition where the compound having anti-HCV activity is effective to inhibit the function of a target selected from HCV metalloprotease, HCV serine protease, HCV polymerase, HCV helicase, HCV NS4B protein, HCV entry, HCV assembly, HCV egress, HCV NS5A protein, IMPDH, and a
30 nucleoside analog for the treatment of an HCV infection.

Another aspect of the invention is a composition comprising a compound of Formula I, or a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable carrier, excipient and/or diluent, an interferon and ribavirin.

- 5 Another aspect of the invention is a method of inhibiting the function of the HCV replicon comprising contacting the HCV replicon with a compound of Formula I or a pharmaceutically acceptable salt thereof.

- 10 Another aspect of the invention is a method of inhibiting the function of the HCV NS5B protein comprising contacting the HCV NS5B protein with a compound of Formula I or a pharmaceutically acceptable salt thereof.

- 15 Another aspect of the invention is a method of treating an HCV infection in a patient comprising administering to the patient a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt thereof. In another embodiment the compound is effective to inhibit the function of the HCV replicon. In another embodiment the compound is effective to inhibit the function of the HCV NS5B protein.

- 20 Another aspect of the invention is a method of treating an HCV infection in a patient comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof, in conjunction with (prior to, after, or concurrently) another compound having anti-HCV activity.

- 25 Another aspect of the invention is the method wherein the other compound having anti-HCV activity is an interferon or a ribavirin.

- 30 Another aspect of the invention is the method where the interferon is selected from interferon alpha 2B, pegylated interferon alpha, consensus interferon, interferon alpha 2A, interferon lambda, and lymphoblastoid interferon tau.

Another aspect of the invention is the method where the other compound having anti-HCV activity is a cyclosporin.

Another aspect of the invention is the method where the cyclosporin is cyclosporin A.

5 Another aspect of the invention is the method where the other compound having anti-HCV activity is selected from interleukin 2, interleukin 6, interleukin 12, a compound that enhances the development of a type 1 helper T cell response, interfering RNA, anti-sense RNA, Imiqimod, ribavirin, an inosine 5'-monophosphate dehydrogenase inhibitor, amantadine, and rimantadine.

10

Another aspect of the invention is the method wherein the other compound having anti-HCV activity is effective to inhibit the function of a target selected from the group consisting of HCV metalloprotease, HCV serine protease, HCV polymerase, HCV helicase, HCV NS4B protein, HCV entry, HCV assembly, HCV egress, HCV NS5A
15 protein, IMPDH, and a nucleoside analog for the treatment of an HCV infection.

20

Another aspect of the invention is the method wherein the other compound having anti-HCV activity is effective to inhibit the function of target in the HCV life cycle other than the HCV NS5B protein.

25

“Therapeutically effective” means the amount of agent required to provide a meaningful patient benefit as understood by practitioners in the field of hepatitis and HCV infection.

“Patient” means a person infected with the HCV virus and suitable for therapy as understood by practitioners in the field of hepatitis and HCV infection.

30

“Treatment,” “therapy,” “regimen,” “HCV infection,” and related terms are used as understood by practitioners in the field of hepatitis and HCV infection.

The compounds of this invention are generally given as pharmaceutical compositions comprised of a therapeutically effective amount of a compound or its pharmaceutically acceptable salt and a pharmaceutically acceptable carrier and can

contain conventional excipients. Pharmaceutically acceptable carriers are those conventionally known carriers having acceptable safety profiles. Compositions encompass all common solid and liquid forms including for example capsules, tablets, lozenges, and powders as well as liquid suspensions, syrups, elixers, and solutions.

- 5 Compositions are made using common formulation techniques, and conventional excipients (such as binding and wetting agents) and vehicles (such as water and alcohols) are generally used for compositions. See, for example, *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Easton, PA, 17th edition, 1985.

- 10 Solid compositions are normally formulated in dosage units and compositions providing from about 1 to 1000 mg of the active ingredient per dose are preferred. Some examples of dosages are 1 mg, 10 mg, 100 mg, 250 mg, 500 mg, and 1000 mg. Generally, other agents will be present in a unit range similar to agents of that class used clinically. Typically, this is 0.25-1000 mg/unit.

15

Liquid compositions are usually in dosage unit ranges. Generally, the liquid composition will be in a unit dosage range of 1-100 mg/mL. Some examples of dosages are 1 mg/mL, 10 mg/mL, 25 mg/mL, 50 mg/mL, and 100 mg/mL. Generally, other agents will be present in a unit range similar to agents of that class used clinically.

- 20 Typically, this is 1-100 mg/mL.

The invention encompasses all conventional modes of administration; oral and parenteral methods are preferred. Generally, the dosing regimen will be similar to other agents used clinically. Typically, the daily dose will be 1-100 mg/kg body weight daily.

- 25 Generally, more compound is required orally and less parenterally. The specific dosing regimen, however, will be determined by a physician using sound medical judgment.

- The invention also encompasses methods where the compound is given in combination therapy. That is, the compound can be used in conjunction with, but
30 separately from, other agents useful in treating hepatitis and HCV infection. In these combination methods, the compound will generally be given in a daily dose of 1-100 mg/kg body weight daily in conjunction with other agents. The other agents generally

will be given in the amounts used therapeutically. The specific dosing regimen, however, will be determined by a physician using sound medical judgment.

Some examples of compounds suitable for compositions and methods are listed in
5 Table 1.

Table 1

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
NIM811		Cyclophilin Inhibitor	Novartis
Zadaxin		Immuno-modulator	Sciclone
Suvus		Methylene blue	Bioenvision
Actilon (CPG10101)		TLR9 agonist	Coley
Batabulin (T67)	Anticancer	β -tubulin inhibitor	Tularik Inc., South San Francisco, CA
ISIS 14803	Antiviral	antisense	ISIS Pharmaceuticals Inc, Carlsbad, CA/Elan Pharmaceuticals Inc., New York, NY
Summetrel	Antiviral	antiviral	Endo Pharmaceuticals Holdings Inc., Chadds Ford, PA
GS-9132 (ACH- 806)	Antiviral	HCV Inhibitor	Achillion / Gilead
Pyrazolopyrimidine compounds and salts From WO- 2005047288 26 May 2005	Antiviral	HCV Inhibitors	Arrow Therapeutics Ltd.

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
Levovirin	Antiviral	IMPDH inhibitor	Ribapharm Inc., Costa Mesa, CA
Merimepodib (VX-497)	Antiviral	IMPDH inhibitor	Vertex Pharmaceuticals Inc., Cambridge, MA
XTL-6865 (XTL-002)	Antiviral	monoclonal antibody	XTL Biopharmaceuticals Ltd., Rehovot, Israel
Telaprevir (VX-950, LY-570310)	Antiviral	NS3 serine protease inhibitor	Vertex Pharmaceuticals Inc., Cambridge, MA/ Eli Lilly and Co. Inc., Indianapolis, IN
HCV-796	Antiviral	NS5B Replicase Inhibitor	Wyeth / Viropharma
NM-283	Antiviral	NS5B Replicase Inhibitor	Idenix / Novartis
GL-59728	Antiviral	NS5B Replicase Inhibitor	Gene Labs / Novartis
GL-60667	Antiviral	NS5B Replicase Inhibitor	Gene Labs / Novartis
2'C MeA	Antiviral	NS5B Replicase Inhibitor	Gilead
PSI 6130	Antiviral	NS5B Replicase Inhibitor	Roche
R1626	Antiviral	NS5B Replicase Inhibitor	Roche
2'C Methyl adenosine	Antiviral	NS5B Replicase Inhibitor	Merck
JTK-003	Antiviral	RdRp inhibitor	Japan Tobacco Inc., Tokyo, Japan

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
Levovirin	Antiviral	ribavirin	ICN Pharmaceuticals, Costa Mesa, CA
Ribavirin	Antiviral	ribavirin	Schering-Plough Corporation, Kenilworth, NJ
Viramidine	Antiviral	Ribavirin Prodrug	Ribapharm Inc., Costa Mesa, CA
Heptazyme	Antiviral	ribozyme	Ribozyme Pharmaceuticals Inc., Boulder, CO
BILN-2061	Antiviral	serine protease inhibitor	Boehringer Ingelheim Pharma KG, Ingelheim, Germany
SCH 503034	Antiviral	serine protease inhibitor	Schering Plough
Zadazim	Immune modulator	Immune modulator	SciClone Pharmaceuticals Inc., San Mateo, CA
Ceplene	Immunomodulator	immune modulator	Maxim Pharmaceuticals Inc., San Diego, CA
CellCept	Immunosuppressant	HCV IgG immuno-suppressant	F. Hoffmann-La Roche LTD, Basel, Switzerland
Civacir	Immunosuppressant	HCV IgG immuno-suppressant	Nabi Biopharmaceuticals Inc., Boca Raton, FL
Albuferon - α	Interferon	albumin IFN- α 2b	Human Genome Sciences Inc., Rockville, MD

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
Infergen A	Interferon	IFN alfacon-1	InterMune Pharmaceuticals Inc., Brisbane, CA
Omega IFN	Interferon	IFN- ω	Intarcia Therapeutics
IFN- β and EMZ701	Interferon	IFN- β and EMZ701	Transition Therapeutics Inc., Ontario, Canada
Rebif	Interferon	IFN- β 1a	Serono, Geneva, Switzerland
Roferon A	Interferon	IFN- α 2a	F. Hoffmann-La Roche LTD, Basel, Switzerland
Intron A	Interferon	IFN- α 2b	Schering-Plough Corporation, Kenilworth, NJ
Intron A and Zadaxin	Interferon	IFN- α 2b/ α 1-thymosin	RegeneRx Biopharma. Inc., Bethesda, MD/ SciClone Pharmaceuticals Inc, San Mateo, CA
Rebetron	Interferon	IFN- α 2b/ribavirin	Schering-Plough Corporation, Kenilworth, NJ
Actimmune	Interferon	INF- γ	InterMune Inc., Brisbane, CA
Interferon- β	Interferon	Interferon- β -1a	Serono
Multiferon	Interferon	Long lasting IFN	Viragen/ Valentis
Wellferon	Interferon	Lympho-blastoid IFN- α n1	GlaxoSmithKline plc, Uxbridge, UK
Omniferon	Interferon	natural IFN- α	Viragen Inc., Plantation, FL

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
Pegasys	Interferon	PEGylated IFN- α 2a	F. Hoffmann-La Roche LTD, Basel, Switzerland
Pegasys and Ceplene	Interferon	PEGylated IFN- α 2a/immune modulator	Maxim Pharmaceuticals Inc., San Diego, CA
Pegasys and Ribavirin	Interferon	PEGylated IFN- α 2a/ribavirin	F. Hoffmann-La Roche LTD, Basel, Switzerland
PEG-Intron	Interferon	PEGylated IFN- α 2b	Schering-Plough Corporation, Kenilworth, NJ
PEG-Intron / Ribavirin	Interferon	PEGylated IFN- α 2b/ribavirin	Schering-Plough Corporation, Kenilworth, NJ
IP-501	Liver protection	antifibrotic	Indevus Pharmaceuticals Inc., Lexington, MA
IDN-6556	Liver protection	caspase inhibitor	Idun Pharmaceuticals Inc., San Diego, CA
ITMN-191 (R-7227)	Antiviral	serine protease inhibitor	InterMune Pharmaceuticals Inc., Brisbane, CA
GL-59728	Antiviral	NS5B Replicase Inhibitor	Genelabs
ANA-971	Antiviral	TLR-7 agonist	Anadys
Boceprevir	Antiviral	serine protease inhibitor	Schering Plough
TMS-435	Antiviral	serine protease inhibitor	Tibotec BVBA, Mechelen, Belgium

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
BI-201335	Antiviral	serine protease inhibitor	Boehringer Ingelheim Pharma KG, Ingelheim, Germany
MK-7009	Antiviral	serine protease inhibitor	Merck
PF-00868554	Antiviral	replicase inhibitor	Pfizer
ANA598	Antiviral	Non-Nucleoside NS5B Polymerase Inhibitor	Anadys Pharmaceuticals, Inc., San Diego, CA, USA
IDX375	Antiviral	Non-Nucleoside Replicase Inhibitor	Idenix Pharmaceuticals, Cambridge, MA, USA
BILB 1941	Antiviral	NS5B Polymerase Inhibitor	Boehringer Ingelheim Canada Ltd R&D, Laval, QC, Canada
PSI-7851	Antiviral	Nucleoside Polymerase Inhibitor	Pharmasset, Princeton, NJ, USA
PSI-7977	Antiviral	Nucleotide NS5B Polymerase Inhibitor	Pharmasset, Princeton, NJ, USA
VCH-759	Antiviral	NS5B Polymerase Inhibitor	ViroChem Pharma
VCH-916	Antiviral	NS5B Polymerase Inhibitor	ViroChem Pharma
GS-9190	Antiviral	NS5B Polymerase Inhibitor	Gilead
Peg-interferon lambda	Antiviral	Interferon	ZymoGenetics/Bristol-Myers Squibb

Synthesis Methods

The compounds can be made by methods known in the art, including those described below. Some reagents and intermediates are known in the art. Other reagents and intermediates can be made by methods known in the art using commercially available materials. The variables (e.g. numbered "R" substituents) used to describe the synthesis of the compounds are intended only to illustrate how to make and are not to be confused with variables used in the claims or in other sections of the specification. Abbreviations used within the schemes generally follow conventions used in the art.


Abbreviations used in the schemes generally follow conventions used in the art. Chemical abbreviations used in the specification and examples are defined as follows: "NaHMDS" for sodium bis(trimethylsilyl)amide; "DMF" for N,N-dimethylformamide; "MeOH" for methanol; "NBS" for N-bromosuccinimide; "Ar" for aryl; "TFA" for trifluoroacetic acid; "LAH" for lithium aluminum hydride; "DMSO" for dimethylsulfoxide; "h" for hours; "rt" for room temperature or retention time (context will dictate); "min" for minutes; "EtOAc" for ethyl acetate; "THF" for tetrahydrofuran; "EDTA" for ethylenediaminetetraacetic acid; "Et₂O" for diethyl ether; "DMAP" for 4-dimethylaminopyridine; "DCE" for 1,2-dichloroethane; "ACN" for acetonitrile; "DME" for 1,2-dimethoxyethane; "HOBt" for 1-hydroxybenzotriazole hydrate; "DIEA" for diisopropylethylamine.

For the section of compounds in the 0000 series all Liquid Chromatography (LC) data were recorded on a Shimadzu LC-10AS or LC-20AS liquid chromatograph using a SPD-10AV or SPD-20A UV-Vis detector and Mass Spectrometry (MS) data were determined with a Micromass Platform for LC in electrospray mode.

HPLC Method (i.e., compound isolation). Compounds purified by preparative HPLC were diluted in methanol (1.2 mL) and purified using a Shimadzu LC-8A or LC-10A or Dionex APS-3000 or Waters Acquity™ automated preparative HPLC system.

Examples:

Preparation of 1001:



Step 1: Palladium on carbon (0.032 g, 10%) was added into a solution of Compound 1 (1 g) in ethyl acetate (80 mL). The reaction was carried out under hydrogen atmosphere with hydrogen balloon at room temperature for 16 hours. Solid was filtered away, and, organic solution was concentrated to give crude Compound 2 which was used as was.

Compound 2	
MS (M+H) ⁺ Calcd.	301.1
MS (M+H) ⁺ Observ.	301.2
Retention Time	2.02 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min

Wavelength	220
Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 30x2, 3u

Step 2: Acetaldehyde (0.264 g) was added into a solution of Compound 2 (0.9 g) in methanol (10 mL). The mixture was stirred at room temperature for 1 hour, before sodium cyanotrihydroborate (0.377 g) was added in. The reaction was carried out at room temperature for 16 hours. After removal of solvents under vacuum, the residue was purified by silica gel chromatography.

Compound 3	
MS (M+H) ⁺ Calcd.	357.2
MS (M+H) ⁺ Observ.	357.2
Retention Time	2.26 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 30x2, 3u

Step 3: 1,1,1-trifluoro-N-phenyl-N-((trifluoromethyl)sulfonyl)methanesulfonamide (1.053 g) and triethylamine (0.596 g) were added into a solution of Compound 3 (0.7 g) in acetonitrile (15 mL). The reaction was heated at 85°C for 16 hours. After removal of solvents under vacuum, the residue was purified by silica gel chromatography.

Compound 4	
MS (M+H) ⁺ Calcd.	489.1
MS (M+H) ⁺ Observ.	489.1
Retention Time	2.44 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

- Step 4: NaHCO₃ (0.112 g) and tetrakis(triphenylphosphine)palladium(0) (0.118 g) were added into a solution of Compound 4 (0.5 g) and (3-(methoxycarbonyl)phenyl)boronic acid (0.221 g) in dioxane (5 mL) and water (2 mL). The reaction was heated at 85°C for 40 hours. After removal of solvents under vacuum, the residue was purified by preparative HPLC system.

Compound 5	
MS (M+H) ⁺ Calcd.	475.2
MS (M+H) ⁺ Observ.	475.3
Retention Time	2.59 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min

Wavelength	220
Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 30x2, 3u

Step 5: Potassium carbonate (0.218 g) was added into a solution of Compound 5 (0.250 g) in methanol (10 mL) and water (5 mL). The reaction was heated at 95°C for 16 hours. After removal of solvents under vacuum, the residue was purified by preparative HPLC

5 system.


Compound 6	
MS (M+H) ⁺ Calcd.	461.2
MS (M+H) ⁺ Observ.	461.1
Retention Time	1.77 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Step 6: iPr₂NEt (5.61 mg) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (10.73 mg) were added into a solution of Compound 6 (0.010 g) and 3,3-difluoroazetidine (2.425 mg) in DMF (1 mL). The reaction was stirred at room temperature for 1 hour. After removal of solvents under vacuum, the residue was purified by preparative HPLC system.

10

1001	
MS (M+H) ⁺ Calcd.	536.2
MS (M+H) ⁺ Observ.	536.2
Retention Time	1.97 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of 2001 and 3003:



5

Step 1: Palladium on carbon (0.028 g, 10%) was added into a solution of Compound 7 (1.3 g) in ethyl acetate (120 mL). The reaction was carried out under hydrogen atmosphere with hydrogen balloon at room temperature for 48 hours. Solid was filtered away, and, organic solution was concentrated to give crude Compound 8 which was used as was.

10

Compound 8	
MS (M+H) ⁺ Calcd.	461.2
MS (M+H) ⁺ Observ.	461.2
Retention Time	1.87 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 30x2, 3u

- Step 2: A solution of Compound 8 (1.0 g) and acetaldehyde (0.124 g) in methanol (10 mL) was stirred at room temperature for 1 hour, before sodium cyanotrihydroborate (0.164 g) was added. The reaction was carried out at room temperature for 16 hours.
- 5 After 100 mL of EtOAc was added into the solution, the mixture was washed with water (2 x 30 mL), brine (20 mL). Then the organic layer was dried over MgSO₄ and concentrated to give crude product Compound 9 which was used as was.

Compound 9	
MS (M+H) ⁺ Calcd.	489.2
MS (M+H) ⁺ Observ.	489.2
Retention Time	2.16 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100

Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 30x2, 3u

- Step 3: TFA (0.963 g) was added into a solution of Compound 9 (0.825 g) in CH₂Cl₂ (5 mL). The reaction was stirred at room temperature for 24 hours. After removal of solvents, the residue was washed with 2 mL of water and solid was collected as the
- 5 desired product Compound 10 which was used as was.

Compound 10	
MS (M+H) ⁺ Calcd.	433.2
MS (M+H) ⁺ Observ.	433.1
Retention Time	2.37 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100
Gradient Time	4 min
Flow Rate	0.8 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 50x2, 3u

- Step 4: iPr₂NEt (0.017 g) and 2-(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yl)-1,1,3,3-tetramethylisouronium hexafluorophosphate(V) (0.032 g) were added into a solution of
- 10 Compound 10 (0.028 g) and piperazine-2-carbonitrile (8.64 mg) in DMF (1 mL). The reaction was stirred at room temperature for 16 hours. After removal of solvents under vacuum, the residue was purified by preparative HPLC system.


2001	
MS (M+H) ⁺ Calcd.	526.2
MS (M+H) ⁺ Observ.	526.1
Retention Time	2.07 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

- Step 5: Pyridine (4.52 mg) and methanesulfonyl chloride (6.54 mg) were added into a solution of Compound 2001 (0.010 g) in CH₂Cl₂ (1 mL). The reaction was stirred at room temperature for 16 hours. After removal of solvents under vacuum, the residue was
- 5 purified by preparative HPLC system.

3003	
MS (M+H) ⁺ Calcd.	682.2
MS (M+H) ⁺ Observ.	682.4
Retention Time	2.82 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100
Gradient Time	4 min
Flow Rate	0.8 mL/min
Wavelength	220

Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 50x2, 3u

Preparation of intermediate 13:



5

Step 1: 2-Iodoethane (0.236 g) and Cs₂CO₃ (0.492 g) were added into a solution of methyl 3-(2-(4-fluorophenyl)-3-(methylcarbamoyl)-6-(methylsulfonamido)benzofuran-5-yl)benzoate (0.25 g) in DMF (5 mL). The reaction was stirred at room temperature for 16 hours. After removal of solvents under vacuum, the residue was purified by preparative

10 HPLC system.

Compound 12	
MS (M+H) ⁺ Calcd.	525.1
MS (M+H) ⁺ Observ.	525.0
Retention Time	2.04 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Step 2: K₂CO₃ (0.020 g) was added into a solution of Compound 12 (0.025 g) in methanol (3 mL) and water (1 mL). The reaction was heated at 70°C for 1 hour. After


removal of solvents under vacuum, the residue was purified by preparative HPLC system to give Compound 13.

Compound 13	
MS (M+H) ⁺ Calcd.	511.1
MS (M+H) ⁺ Observ.	511.0
Retention Time	1.95 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um


5 General procedure of amide formation, preparation of 3001 and 3002:

iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 7 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was

10 isolated by preparative HPLC system.


 3001	
MS (M+H) ⁺ Calcd.	704.3
MS (M+H) ⁺ Observ.	704.4

Retention Time	1.81 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 30x2, 3u



MS (M-H) ⁺ Calcd.	702.3
MS (M-H) ⁺ Observ.	702.6
Retention Time	1.81 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 30x2, 3u

Preparation of Intermediate 11:



- 5 Step 1: Compound 14 was prepared via the same process of synthesizing Compound 12, using 1-fluoro-2-iodoethane as starting material.

Compound 14	
MS (M+H) ⁺ Calcd.	543.1
MS (M+H) ⁺ Observ.	543.3
Retention Time	3.03 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 30x2, 3u

- 10 Step 2: Compound 15 was prepared via the same process of synthesizing Compound 13, using Compound 14 as starting material.

Compound 15	
MS (M+H) ⁺ Calcd.	529.1
MS (M+H) ⁺ Observ.	529.1
Retention Time	3.38 min




LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	0
Final % B	100
Gradient Time	4 min
Flow Rate	0.8 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 X 50mm 3um




General procedure of amide formation, preparation of 4001-4027:



- 5 iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 15 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.



LC Condition A	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 X 30mm 3um




LC Condition B	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Acetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Acetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 30x2, 3u



Cmpd #	LC Method	Structure	MS (M+H) ⁺ Calcd.	MS (M+H) ⁺ Observ.	Retention Time (min)
4001	A		693.2	693.2	2.02
4002	A		637.2	637.3	1.70
4003	A		675.2	675.3	2.01




Cmpd #	LC Method	Structure	MS (M+H) ⁺ Calcd.	MS (M+H) ⁺ Observ.	Retention Time (min)
4004	A		710.2	710.3	1.91
4005	A		769.3	769.4	2.17
4006	B		725.2	725.5	1.71




Cmpd #	LC Method	Structure	MS (M+H) ⁺ Calcd.	MS (M+H) ⁺ Observ.	Retention Time (min)
4007	B		749.2	749.6	1.78
4008	A		741.2	741.2	2.00




Cmpd #	LC Method	Structure	MS (M+H) ⁺ Calcd.	MS (M+H) ⁺ Observ.	Retention Time (min)
4010	A		755.3	755.2	2.05
4011	B		760.3	760.6	2.10



Cmpd #	LC Method	Structure	MS (M+H) ⁺ Calcd.	MS (M+H) ⁺ Observ.	Retention Time (min)
4012	A		766.2	766.2	2.11
4013	B		709.2	709.5	1.82
4014	B		710.2	710.6	1.65

Cmpd #	LC Method	Structure	MS (M+H) ⁺ Calcd.	MS (M+H) ⁺ Observ.	Retention Time (min)
4015	B		650.2	650.5	1.78
4016	A		701.3	701.2	1.80

Cmpd #	LC Method	Structure	MS (M+H) ⁺ Calcd.	MS (M+H) ⁺ Observ.	Retention Time (min)
4017	B		625.2	625.4	1.46
4018	B		732.2	732.5	1.67
4019	A		661.2	661.2	1.90

Cmpd #	LC Method	Structure	MS (M+H) ⁺ Calcd.	MS (M+H) ⁺ Observ.	Retention Time (min)
4020	A		600.2	600.2	1.91
4021	B		607.2	607.4	1.63
4022	A		765.2	765.3	2.19

Cmpd #	LC Method	Structure	MS (M+H) ⁺ Calcd.	MS (M+H) ⁺ Observ.	Retention Time (min)
4023	A		665.2	665.3	2.02
4024	A		600.2	600.2	1.93
4025	A		654.1	654.2	2.11

Cmpd #	LC Method	Structure	MS (M+H) ⁺ Calcd.	MS (M+H) ⁺ Observ.	Retention Time (min)
4026	A		665.2	665.3	2.00
4027	A		697.2	697.3	1.93

Biological Methods

The compound demonstrated activity against HCV NS5B as determined in the following HCV RdRp assays.

HCV NS5B RdRp cloning, expression, and purification. The cDNA encoding NS5B proteins of HCV genotype 1b (Con1), a genotype 1b variant with amino acid 316 mutated from cysteine to asparagine, and genotype 2a (JFH-1), were cloned into the pET21a expression vector. Each untagged protein was expressed with an 18 amino acid C-terminal truncation to enhance the solubility. The E. coli competent cell line BL21(DE3) was used for expression of the protein. Cultures were grown at 37 °C for ~ 4 hours until the cultures reached an optical density of 2.0 at 600 nm. The cultures were cooled to 20 °C and induced with 1 mM IPTG. Fresh ampicillin was added to a final concentration of 50 µg/mL and the cells were grown overnight at 20 °C.

Cell pellets (3L) were lysed for purification to yield 15-24 mgs of purified NS5B. The lysis buffer consisted of 20 mM Tris-HCl, pH 7.4, 500 mM NaCl, 0.5% triton X-100, 1 mM DTT, 1mM EDTA, 20% glycerol, 0.5 mg/mL lysozyme, 10 mM MgCl₂, 15 ug/mL deoxyribonuclease I, and Complete TM protease inhibitor tablets (Roche). After addition of the lysis buffer, frozen cell pellets were resuspended using a tissue homogenizer. To reduce the viscosity of the sample, aliquots of the lysate were sonicated on ice using a microtip attached to a Branson sonicator. The sonicated lysate was centrifuged at 100,000 x g for 30 minutes at 4 °C and filtered through a 0.2 µm filter unit (Corning).

The protein was purified using two sequential chromatography steps: Heparin sepharose CL-6B and polyU sepharose 4B. The chromatography buffers were identical to the lysis buffer but contained no lysozyme, deoxyribonuclease I, MgCl₂ or protease inhibitor and the NaCl concentration of the buffer was adjusted according to the requirements for charging the protein onto the column. Each column was eluted with a NaCl gradient which varied in length from 5-50 column volumes depending on the

column type. After the final chromatography step, the resulting purity of the enzyme is >90% based on SDS-PAGE analysis. The enzyme was aliquoted and stored at -80 °C.

HCV NS5B RdRp enzyme assay. An on-bead solid phase homogeneous assay was used in a 384-well format to assess NS5B inhibitors (Wang Y-K, Rigat K, Roberts S, and Gao M (2006) Anal Biochem, 359: 106-111). The biotinylated oligo dT₁₂ primer was captured on streptavidin-coupled imaging beads (GE, RPNQ0261) by mixing primer and beads in 1X buffer and incubating at room temperature for three hours. Unbound primer was removed after centrifugation. The primer-bound beads were resuspended in 3X reaction mix (20 mM Hepes buffer, pH 7.5, dT primer coupled beads, poly A template, ³H-UTP, and RNase inhibitor (Promega N2515)). Compounds were serially diluted 1:3 in DMSO and aliquoted into assay plates. Equal volumes (5 µL) of water, 3X reaction mix, and enzyme in 3X assay buffer (60 mM Hepes buffer, pH 7.5, 7.5 mM MgCl₂, 7.5 mM KCl, 3 mM DTT, 0.03 mg/mL BSA, 6 % glycerol) were added to the diluted compound on the assay plate. Final concentration of components in 384-well assay: 0.36 nM template, 15 nM primer, 0.29 µM ³H-UTP (0.3 µCi), 1.6 U/µL RNase inhibitor, 7 nM NS5B enzyme, 0.01 mg/mL BSA, 1 mM DTT, and 0.33 µg/µL beads, 20 mM Hepes buffer, pH 7.5, 2.5 mM MgCl₂, 2.5 mM KCl, and 0.1 % DMSO.

Reactions were allowed to proceed for 24 hours at 30° C and terminated by the addition of 50 mM EDTA (5 µL). After incubating for at least 15 minutes, plates were read on an Amersham LEADseeker multimodality imaging system.

IC₅₀ values for compounds were determined using ten different [I]. IC₅₀ values were calculated from the inhibition using the four-parameter logistic formula $y = A + ((B - A) / (1 + ((C/x)^D)))$, where A and B denote minimal and maximal % inhibition, respectively, C is the IC₅₀, D is hill slope and x represents compound concentration.

Cell lines. The cell lines used to evaluate compounds consist of a human hepatocyte derived cell line (Huh-7) that constitutively expresses a genotype 1b (Con-1) HCV replicon or a genotype 1b (Con-1) HCV replicon with an asparagine replacing the cysteine at amino acid 316, or a genotype 2a (JFH-1) replicon, containing a Renilla


luciferase reporter gene. These cells were maintained in Dulbecco's modified Eagle medium (DMEM) containing 10% FBS, 100 U/mL penicillin/streptomycin and 1.0 mg/mL G418.






5 *HCV replicon luciferase assay.* To evaluate compound efficacy, titrated compounds were transferred to sterile 384-well tissue culture treated plates, and the plates were seeded with HCV replicon cells (50 μ L at a density of 2.4×10^3 cells/well) in DMEM containing 4 % FBS (final DMSO concentration at 0.5 %). After 3 days incubation at 37°C, cells were analyzed for Renilla Luciferase activity using the EnduRen
 10 substrate (Promega cat #E6485) according to the manufacturer's directions. Briefly, the EnduRen substrate was diluted in DMEM and then added to the plates to a final concentration of 7.5 μ M. The plates were incubated for at least 1 h at 37°C then read on a Viewlux Imager (PerkinElmer) using a luminescence program. The 50% effective concentration (EC₅₀) was calculated using the four-parameter logistic formula noted
 15 above.

To assess cytotoxicity of compounds, Cell Titer-Blue (Promega) was added to the EnduRen-containing plates and incubated for at least 4 hrs at 37°C. The fluorescence
 20 signal from each well was read using a Viewlux Imager. All CC₅₀ values were calculated using the four-parameter logistic formula.





Compound EC₅₀ data is expressed as A: < 100 nM; B = 100-1000 nM; C > 1000 nM). Representative data for compounds are reported in Table 2.






Table 2






Cmpd#	Structure	EC ₅₀ (μ M)
1001		0.0454






Cmpd#	Structure	EC ₅₀ (uM)
2001		B
3001		A
3002		A
3003		0.0252
4001		C

Cmpd#	Structure	EC ₅₀ (uM)
4002		A
4003		A
4004		1.6920
4005		A
4006		B

Cmpd#	Structure	EC ₅₀ (uM)
4007		A
4008		0.1073
4009		A
4010		A

Cmpd#	Structure	EC ₅₀ (uM)
4011		A
4012		0.0271
4013		A
4014		A
4015		A

Cmpd#	Structure	EC ₅₀ (uM)
4016		0.0848
4017		C
4018		B
4019		C
4020		0.0450

Cmpd#	Structure	EC ₅₀ (uM)
4021		C
4022		A
4023		A
4024		0.0765
4025		A


Cmpd#	Structure	EC ₅₀ (uM)
4026		A
4027		C

It will be evident to one skilled in the art that the present disclosure is not limited to the foregoing illustrative examples, and that it can be embodied in other specific forms without departing from the essential attributes thereof. It is therefore desired that the examples be considered in all respects as illustrative and not restrictive, reference being made to the appended claims, rather than to the foregoing examples, and all changes which come within the meaning and range of equivalency of the claims are therefore intended to be embraced therein.

CLAIMS

We claim:

1. A compound of Formula I, including pharmaceutically acceptable salts thereof:



I

Wherein R^1 , R^2 are each independently selected from the group of hydrogen, halo, nitro, alkyl, cycloalkyl, haloalkyl, aminoalkyl, hydroxyalkyl, alkoxyalkyl, hydroxyl, alkoxy, OR^{17} , cycloalkoxy, amino, alkylamino, dialkylamino, alkylcarboxamido, alkoxy-carboxamido, alkoxyalkylcarboxamido, and Ar^1 ;

R^3 is selected from the group of cyano, alkoxy-carbonyl, (cycloalkyl)oxy-carbonyl, (alkylsulfonyl)aminocarbonyl, $CONR^{11}R^{12}$, $(R^{11})(R^{12})NCONH$, triazolyl, thiazolyl, and tetrazolyl;

R^4 is phenyl substituted with 0-2 halo substituents;

R^5 is selected from the group of hydrogen, alkyl, alkylsulfonyl, alkylcarbonyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, cyanoalkyl, cyanoalkenyl, cyanoalkynyl, cycloalkyl, halocycloalkyl, (haloalkyl)cycloalkyl, (halocycloalkyl)cycloalkyl, cyanocycloalkyl, (cyanoalkyl)cycloalkyl, (cyanocycloalkyl)cycloalkyl, hydroxycycloalkyl, (hydroxyalkyl)cycloalkyl, (hydroxycycloalkyl)cycloalkyl, alkoxy-cycloalkyl, (alkoxyalkyl)cycloalkyl, and (alkoxycycloalkyl)cycloalkyl;

R^6 is selected from the group of hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, alkylsulfonyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, cyanoalkyl, cyanoalkenyl, cyanoalkynyl, cycloalkyl, halocycloalkyl, (haloalkyl)cycloalkyl, (halocycloalkyl)cycloalkyl, cyanocycloalkyl, (cyanoalkyl)cycloalkyl, (cyanocycloalkyl)cycloalkyl, hydroxycycloalkyl, (hydroxyalkyl)cycloalkyl,

(hydroxycycloalkyl)cycloalkyl alkoxycycloalkyl, (alkoxyalkyl)cycloalkyl, and (alkoxycycloalkyl)cycloalkyl;

R^7R^8N taken together is azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, dihydroindolyl, or isoindoliny, and is substituted with 0-2 substituents selected from the group of alkyl, hydroxyalkyl, alkoxyalkyl, hydroxy, alkoxy, carboxy, alkoxy carbonyl, dialkylcarboxamido, alkylcarbonylamino, alkoxy carbonylamino, pyridiny and phenyl, and is further substituted with one or more additional substituent(s) selected from the group of halo, haloalkyl, cyano, cyanoalkyl, cyanoalkenyl, cyanocycloalkyl, cyanocycloalkenyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, urea, Ar^2 and phenyl groups, wherein said Ar^2 and phenyl groups are substituted with one or more substituents selected from the group of haloalkyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, and urea, and further wherein said pyridiny is substituted with one or more substituents selected from the group of halo, alkyl, haloalkyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, and urea;

Or

R^7R^8N taken together is selected from the group of azepany, 1,4-diazepany, 1,4-oxazepany, 1,4-thiazepany, 1,1-dioxo 1,4-thiazepany, azocany, 1,4-diazocany, 1,5-diazocany, 1,4-oxazocany, 1,5-oxazocany, 1,4-thiazocany, 1,5-thiazocany, 1,1-dioxo 1,4-thiazocany, and 1,1-dioxo 1,5-thiazocany, and is substituted with 0-2 substituents selected from the group of halo, haloalkyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, alkyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, aminocarbonyl, alkoxy carbonyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidiny, guanidiny, carbamide, urea, and Ar^3 ;

Or

R^7R^8N taken together is selected from the group of azetidiny, pyrrolidiny, piperidiny, piperaziny, morpholiny, dihydroindolyl, isoindoliny, azepany, 1,4-

diazepanyl, 1,4-oxazepanyl, 1,4-thiazepanyl, 1,1-dioxo 1,4-thiazepanyl, azocanyl, 1,4-diazocanyl, 1,5-diazocanyl, 1,4-oxazocanyl, 1,5-oxazocanyl, 1,4-thiazocanyl, 1,5-thiazocanyl, 1,1-dioxo 1,4-thiazocanyl, and 1,1-dioxo 1,5-thiazocanyl, and is substituted with a methylene group which is substituted with 0-2 substituents selected from the group of halo, haloalkyl, cyano, cyanoalkyl, cyanoalkenyl, cyanocycloalkyl, cyanocycloalkenyl, halocycloalkyl, haloalkenyl, halocycloalkenyl, cycloalkyl, alkenyl, alkynyl, cycloalkenyl, ester, amide, acid, amine, alcohol, ether, cycloether, cyclic amine, lactame, lactone, sulfonyl, aminosulfonyl, amidinyl, guanidinyl, carbamide, urea, and Ar³;

R¹¹ is hydrogen, alkyl, or cycloalkyl;

R¹² is hydrogen, alkyl, or cycloalkyl;

or R¹¹ and R¹² taken together with the nitrogen to which they are attached is azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, or morpholinyl;

R¹⁷ is selected from the group of haloalkyl, cyanoalkyl, (cycloalkyl)alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, (R¹⁸)alkyl, (Ar⁴)alkyl, alkynyl, and aminocycloalkyl;

R¹⁸ is selected from the group of CONH₂, H₂NCONH, dibenzylamino, phthalimido, amino, alkylamino, dialkylamino, azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl where azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, and morpholinyl, and is substituted with 0-3 alkyl or alkoxy carbonyl substituents;

R²⁰ is hydrogen, halo, alkyl, or alkoxy;

R²¹ is hydrogen, halo, alkyl, or alkoxy;

Ar¹ is selected from the group of phenyl, naphthalenyl, quinolinyl, isoquinolinyl, quinoxaliny, quinazolinyl, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, isoxazolyl, isothiazolyl, imidazolyl, oxazolyl, thiazolyl, oxadiazolyl, oxadiathiazolyl, triazolyl, tetrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazolyl, indolyl, azaindolyl, indazolyl, azaindazolyl, benzoxazolyl, benzoisoxazolyl, benzoisothiazolyl, benzimidazolyl, and benzothiazolyl, and is substituted with 0-2 substituents selected from the group of halo, alkyl, cycloalkyl, haloalkyl, alkoxyalkyl, hydroxy, alkoxy, amino, alkylamino, dialkylamino, aminocarbonyl, pyridinyl, phenyl, halophenyl, alkylphenyl, and alkoxyphenyl;

Ar² is selected from the group of naphthalenyl, quinolinyl, isoquinolinyl, quinoxaliny, quinazolinyl, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, isoxazolyl,


isothiazolyl, imidazolyl, oxazolyl, thiazolyl, oxadiazolyl, oxadiathiazolyl, triazolyl, tetrazolyl, pyrazinyl, pyridazinyl, triazolyl, indolyl, azaindolyl, indazolyl, azaindazolyl, benzoxazolyl, benzoisoaxazolyl, benzoisorhiazolyl, benzimidazolyl, and benzothiazolyl, and is substituted with 0-2 substituents selected from the group of halo, alkyl, cycloalkyl, haloalkyl, alkoxyalkyl, hydroxy, alkoxy, amino, alkylamino, dialkylamino, aminocarbonyl, halophenyl, alkylphenyl, alkoxyphenyl and Ar³;


Ar³ is selected from the group of phenyl, naphthalenyl, quinolinyl, isoquinolinyl, quinoxalyl, quinazolinyl, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, isoxazolyl, isothiazolyl, imidazolyl, oxazolyl, thiazolyl, oxadiazolyl, oxadiathiazolyl, triazolyl, tetrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazolyl, indolyl, azaindolyl, indazolyl, azaindazolyl, benzoxazolyl, benzoisoaxazolyl, benzoisorhiazolyl, benzimidazolyl, and benzothiazolyl, and is substituted with 0-2 substituents selected from the group of halo, alkyl, cycloalkyl, haloalkyl, alkoxyalkyl, hydroxy, alkoxy, amino, alkylamino, dialkylamino, aminocarbonyl, pyridinyl, phenyl, halophenyl, alkylphenyl, and alkoxyphenyl; and


Ar⁴ is selected from the group of furanyl, thienyl, pyrrolyl, pyrazolyl, isoxazolyl, isothiazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyridinyl, indolyl, and phenyl, and is substituted with 0-2 substituents selected from the group of halo, alkyl, haloalkyl, hydroxyl, and alkoxy.



2. A compound of claim 1 wherein R¹ and R² are each hydrogen.
3. A compound of claim 2 wherein R⁴ is phenyl substituted with one fluoro group.
4. A compound of claim 3 wherein R³ is -CONR¹¹R¹².
5. A compound of claim 4 wherein R¹¹ and R¹² are each independently hydrogen or alkyl.
6. A compound of claim 2 wherein R⁵ is alkyl or haloalkyl.

7. A compound of claim 6 wherein R⁶ is alkylsulfonyl.
8. A compound of claim 2 wherein R⁷R⁸N is selected from the group of azetidiny, pyrrolidiny, piperidiny, and piperaziny.
9. A compound of claim 8 wherein R⁷R⁸N is piperaziny.
10. A compound of claim 2 wherein R²⁰ is hydrogen.
11. A compound of claim 10 wherein R²¹ is hydrogen.
12. A compound, including pharmaceutically acceptable salts thereof, which is selected from the group of:









13. A composition comprising a compound of claim 1 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier, excipient and/or diluent.

14. A composition comprising a compound of claim 12 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier, excipient and/or diluent.

15. A method of treating hepatitis C infection comprising administering a therapeutically effective amount of a compound of claim 1 to a patient.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/018171

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D405/10 C07D405/12 C07D405/14 C07D307/84 C07D401/10
C07D401/14 C07D413/14 C07D417/14 A61K31/397 A61K31/496
A61P31/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BEILSTEIN Data, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2010/030592 A1 (SQUIBB BRISTOL MYERS CO [US]; YEUNG KAP-SUN [US]) 18 March 2010 (2010-03-18) schemes 1, 3; claims 1, 7, 10, 11; tables 1a-1c in particular table 1a, pages 42, 45, 47, 49, 52-56, 64-67, 70, 73 (-CONR7R8 residue), table 1b, pages 74, 79, 82-86, 90-92, ... (-NR5R6 residue) -----	1-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

22 March 2016

Date of mailing of the international search report

01/04/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Ladenburger, Claude

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2016/018171

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2010030592	A1	18-03-2010	
		AR 073740 A1	01-12-2010
		AU 2009291966 A1	18-03-2010
		CA 2736988 A1	18-03-2010
		CN 102209717 A	05-10-2011
		CO 6382110 A2	15-02-2012
		EA 201100390 A1	31-10-2011
		EP 2326633 A1	01-06-2011
		JP 5685191 B2	18-03-2015
		JP 2012502101 A	26-01-2012
		KR 20110059875 A	07-06-2011
		NZ 591172 A	21-12-2012
		PE 08062011 A1	31-10-2011
		TW 201014836 A	16-04-2010
		TW 201124403 A	16-07-2011
		US 2010093694 A1	15-04-2010
		WO 2010030592 A1	18-03-2010
