



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07F 5/02, C08F 10/00	A1	(11) International Publication Number: WO 99/06413 (43) International Publication Date: 11 February 1999 (11.02.99)
<p>(21) International Application Number: PCT/US98/14106</p> <p>(22) International Filing Date: 7 July 1998 (07.07.98)</p> <p>(30) Priority Data: 60/054,586 1 August 1997 (01.08.97) US</p> <p>(71) Applicants (for all designated States except US): THE DOW CHEMICAL COMPANY [US/US]; 2030 Dow Center, Midland, MI 48674 (US). NORTHWESTERN UNIVERSITY [US/US]; 633 Clark Street, Evanston, IL 60208 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): MCADON, Mark, H. [US/US]; 2810 Dawn Drive, Midland, MI 48642 (US). NICKIAS, Peter, N. [US/US]; 2705 Mt. Vernon Drive, Midland, MI 48642 (US). MARKS, Tobin, J. [US/US]; 2300 Central Park Avenue, Evanston, IL 60201 (US). SWARTZ, David, J. [US/US]; 510 That Way Boulevard, Lake Jackson, TX 77566 (US).</p> <p>(74) Agent: DELINE, Douglas, N.; Patent Dept., P.O. Box 1967, Midland, MI 48641-1967 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: CATALYST ACTIVATOR</p> <p>(57) Abstract</p> <p>A catalyst activator particularly adapted for use in the activation of metal complexes of metals of Group 3-10 for polymerization of ethylenically unsaturated polymerizable monomers, especially olefins, comprising two Group 13 metal or metalloid atoms and a ligand structure including at least one bridging group connecting ligands on the two Group 13 metal or metalloid atoms.</p>		

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CATALYST ACTIVATOR

The present invention relates to compounds that are useful as catalyst components. More particularly the present invention relates to such compounds that are particularly adapted for use in the coordination polymerization of unsaturated compounds comprising two Group 13 metal or metalloid atoms and a ligand structure including at least one bridging group connecting ligands on two Group 13 metal or metalloid atoms thereof. Such compounds are particularly advantageous for use in a polymerization process wherein catalyst, catalyst activator, and at least one polymerizable monomer are combined under polymerization conditions to form a polymeric product.

It is previously known in the art to activate Ziegler-Natta polymerization catalysts, particularly such catalysts comprising Group 3-10 metal complexes containing delocalized π -bonded ligand groups, by the use of Bronsted acid salts capable of transferring a proton to form a cationic derivative or other catalytically active derivative of such Group 3-10 metal complex. Preferred Bronsted acid salts are such compounds containing a cation/ anion pair that are capable of rendering the Group 3-10 metal complex catalytically active. Suitable activators comprise fluorinated arylborate anions, most preferably, the tetrakis(pentafluorophenyl)borate anion. Additional suitable anions include sterically shielded diboron anions of the formula:



wherein:


S is hydrogen, alkyl, fluoroalkyl, aryl, or fluoroaryl, Ar^{F} is fluoroaryl, and X^1 is either hydrogen or halide, disclosed in US-A-5,447,895. Additional examples include carborane compounds such as are disclosed and claimed in US-A-5, 407,884.

Examples of preferred charge separated (cation/ anion pair) activators are protonated ammonium, sulfonium, or phosphonium salts capable of transferring a hydrogen ion, disclosed in US-A-5,198,401, US-A-5,132,380, US-A-5,470,927, and US-A-5,153,157, as well as oxidizing salts such as carbonium, ferrocenium and silylium salts, disclosed in US-A-5,350,723, US-A-5,189,192 and US-A-5,626,087.

Further suitable activators for the above metal complexes include strong Lewis acids including (tris(perfluorophenyl)borane and tris(perfluorobiphenyl)borane. The former composition has been previously disclosed for the above stated end use in EP-A-520,732, whereas the latter composition is similarly disclosed by Marks, et al., in J. Am. Chem. Soc., 118, 12451-12452 (1996).

Despite the satisfactory performance of the foregoing catalyst activators under a variety of polymerization conditions, there is still a need for improved cocatalysts for use in the activation of various metal complexes under a variety of reaction conditions. Accordingly, it would be desirable if there were provided compounds that could be employed in solution, slurry, gas phase or high pressure polymerizations and under homogeneous or heterogeneous process conditions having improved activation properties.

According to the present invention there is now provided Group 13 containing compounds useful as catalyst activators in neutral (Lewis acid) or charge separated (cation/ anion pair) form, corresponding to the formula:



wherein:

B^1 and B^2 independently each occurrence are Group 13 metal or metalloid atoms, preferably boron;

Z^* is an optional divalent bridging group containing from 1 to 20 atoms, not counting hydrogen atoms;

R^1 and R^2 independently each occurrence are monovalent, anionic ligand groups containing from 1 to 40 atoms not counting hydrogen atoms, and, for cationic compounds, additionally comprising a dissociated cation moiety;

Ar^{f1} and Ar^{f2} independently each occurrence are monovalent, fluorinated organic groups containing from 6 to 100 carbon atoms, optionally, an Ar^{f1} and an R^2 group, or an Ar^{f2} and an R^1 group together form a divalent bridging group, and further optionally an Ar^{f1} group and an Ar^{f2} group together form a C₆₋₁₀₀ divalent bridging group,

z is 0 or 1,

r and s independently are 0, 1 or 2, and

m and n are 1, 2 or 3;

with the proviso that when z is 0, at least one of Ar^{f1} and Ar^{f2} are joined together, and the sum of r, z and m is 3 or 4, in the former event B^1 is neutral and in the latter event B^1 is negatively charged, said charge being balanced by a cation component of one R^1 ; and the sum of s, z and n is 3 or 4, in the former event B^2 is neutral and in the latter event B^2 is negatively charged, said charge being balanced by a cation component of one R^2 .

Additionally according to the present invention there is provided a catalyst composition for polymerization of an ethylenically unsaturated, polymerizable monomer comprising, in combination, the above described compound and a Group 3-10 metal complex, or the reaction product of such combination.

Additionally according to the present invention there is provided a process for polymerization of one or more ethylenically unsaturated, polymerizable monomers comprising contacting the same, optionally in the presence of an inert aliphatic, alicyclic or aromatic hydrocarbon, with the above catalyst composition.


The foregoing compounds are uniquely adapted for use in activation of a variety of metal complexes, especially Group 4 metal complexes, under standard and atypical olefin polymerization conditions. They are uniquely capable of forming monomeric and dimeric cationic metal complexes when combined with neutral metallocene complexes under such polymerization conditions. Because of this fact, the foregoing compounds are capable of forming highly desirable olefin polymers having enhanced levels of long chain branching, stereospecificity and comonomer distribution. In particular, bis-anions, due to the pairing of active catalyst sites in close proximity to one another, are capable of providing a higher local concentration of active catalyst site at the point of polymer formation. Moreover, such paired catalyst sites may be comprised of two disparate metals or metal ligand arrangements, or otherwise tailored to provide desirable polymer properties. For example, the use of symmetrical or unsymmetrical bis-anions results in two catalytically active sites that are held in close proximity during a polymerization reaction, thereby providing a large increase in local concentration of active catalyst sites. This increased local concentration of active catalyst sites leads to enhanced polymer stereostructure, molecular weight and microstructure. Certain catalyst sites in close proximity result in random comonomer incorporation, others affect the

stereospecificity of their close neighbor. By controlling the random versus clustered distribution of comonomer, blocky or non-blocky copolymers can be prepared. Additionally, the two catalysts associated with each bis-anion may themselves be nonspecific or stereospecific, such that the resulting combination catalyst is adapted to produce block copolymers via polymer interchange between such nonspecific and stereospecific catalysts. The degree of long-chain branching in polyolefins produced using multiple catalyst sites on bis-anions is enhanced due to the rate of reincorporation of *in situ* generated vinyl terminated macromonomer into the growing polymer chain due to the higher local concentration of catalyst.

10 DETAILED DESCRIPTION OF THE INVENTION


All references herein to elements belonging to a certain Group refer to the Periodic Table of the Elements published and copyrighted by CRC Press, Inc., 1995. Also any reference to the Group or Groups shall be to the Group or Groups as reflected in this Periodic Table of the Elements using the IUPAC system for numbering groups. When, in reference to a cation portion of any compound herein, it is stated that a ligand group comprises such cation, it is to be understood that the cation is not chemically or physically incorporated in said ligand, or necessarily chemically attached thereto, in as much as the cation may freely dissociate from the anion portion of the compound. Rather, such ligand group is said to "comprise" the cation in order to properly account for the correct number of cations as dictated by considerations of charge balance.

The catalyst activators of the invention are further characterized in the following manner. Preferred Group 13 metal or metalloids include aluminum and boron. Most highly preferably, both B¹ and B² are boron. The cocatalysts may be neutral Lewis acids or salts comprising one or more cation-anion pairs. Examples of suitable neutral Lewis acids according to the present invention correspond to the formula:



wherein all variables are as previously defined, and the sum of r, z and m and the sum of s, z and n are both 3.

More specific examples of the foregoing Lewis acid compounds correspond to the formula:




wherein:


- 5 R^1 and R^2 independently each occurrence are C_{1-20} hydrocarbyl, halohydrocarbyl, or halocarbyl, and

Ar^{f1} - Ar^{f2} in combination, independently each occurrence, is a divalent fluoro-substituted aromatic group of from 6 to 20 carbons.

Preferred examples of the foregoing Lewis acid compounds are the following:



Examples of suitable charge separated compounds according to the present invention correspond to the formula:




wherein:

- 15 R^1 and R^2 independently each occurrence are C_{1-20} hydrocarbyl, halohydrocarbyl, or halocarbyl groups, and (when the sum of r , z and m is 4) one R^1 additionally comprises a cation selected from the group consisting of protonated cations of Bronsted acids, ferrocenium cations, carbonium cations, silylium cations, and Ag^+ , and (when the sum of s , z and n is 4) one R^2 additionally comprises a

cation selected from the group consisting of protonated cations of Bronsted acids, ferrocenium cations, carbonium cations, silylium cations, and Ag^+ ;

- 5 r and s are 0, 1 or 2 with the proviso that at least one of r or s is not 0, and the sum of r, z and m is 3 or 4 and the sum of s, z and n is 3 or 4, with the proviso that at least one of the foregoing sums is 4.


More specific examples of the foregoing charge separated compounds correspond to the formula:



wherein:

- 10 R^1 and R^2 independently each occurrence are C_{1-20} hydrocarbyl, halohydrocarbyl, or halocarbyl groups, and (when connected to a negatively charged boron atom) one R^1 additionally comprises a cation selected from the group consisting of protonated cations of Bronsted acids, ferrocenium cations, carbonium cations, silylium cations, and Ag^+ , and (when connected to a negatively charged
- 15 boron atom) one R^2 additionally comprises a cation selected from the group consisting of protonated cations of Bronsted acids, ferrocenium, carbonium cations, silylium cations, and Ag^+ ; and
- an $\text{Ar}^{\text{f}1}$ group and an $\text{Ar}^{\text{f}2}$ group together form a C_{6-20} divalent fluoro-substituted aromatic group of from 6 to 20 carbons.

- 20 Specific examples of the foregoing salt compounds are the following:



wherein, R is a C₁₋₄₀ hydrocarbonyl ligand group, and L⁺ is a cation of a Bronsted acid, or a ferrocenium, carbonium, silylium, or Ag⁺ cation.


- 5 More preferably L⁺ is an ammonium cation of the formula HN⁺R₃, wherein R is C₁₋₅₀ hydrocarbonyl. Most preferably, one or two R groups are C₁₄₋₅₀ aliphatic groups, and the remaining R group(s) is (are) C₁₋₄ aliphatic.

The skilled artisan will appreciate that upon activation of a metal complex to a catalytically active state by the present compounds, to the extent a cationic derivative thereof is formed, the foregoing charge separated compounds may include therein the cationic derivative of such metal complex in place of the foregoing Bronsted acid, ferrocenium, carbonium, silylium, or Ag⁺ cations. For the preferred complexes the metal is selected from Groups 3-10 of the Periodic Table of the Elements, more preferably Group 4. Accordingly, such cationic derivative would be a Group 3-10 metal containing cation, more preferably a Group 4 metal containing cation.

Generally, solubility of the compounds of the invention in aliphatic compounds is increased by incorporation of one or more oleophilic groups such as long chain alkyl groups; long chain alkenyl groups; or halo-, alkoxy-, amino-, silyl-, or germlyl-substituted long chain alkyl groups or long chain alkenyl groups into the cation, L⁺. By the term "long chain" are meant groups having from 10 to 50 non-hydrogen atoms in such group, preferably in a non-branched form. It is understood that the

compound may comprise a mixture of oleophilic groups of differing lengths in the cation. For example, one suitable compound comprises the protonated ammonium salt derived from the commercially available long chain amine comprising a mixture of two C₁₄, C₁₆ or C₁₈ alkyl groups and one methyl group. Such amines are available
 5 from Witco Corp., under the trade name Kemamine™ T9701, and from Akzo-Nobel under the trade name Armeen™ M2HT.

Suitable catalysts for use in combination with the foregoing cocatalysts include any compound or complex of a metal of Groups 3-10 of the Periodic Table of the Elements capable of being activated to polymerize ethylenically unsaturated
 10 compounds by the present activators. Examples include Group 10 diimine derivatives corresponding to the formula:



M* is Ni(II) or Pd(II);

K is halo, hydrocarbyl, or hydrocarbyloxy;

15 Ar* is an aryl group, especially 2,6-diisopropylphenyl or aniline group;

CT-CT is 1,2-ethanediyl, 2,3-butanediyl, or form a fused ring system wherein the two T groups together are a 1,8-naphthanediyl group; and

A⁻ is the anionic component of the foregoing charge separated activators.

Similar catalysts to the foregoing are also disclosed by M. Brookhart, et al., in
 20 J. Am. Chem. Soc., 118, 267-268 (1996) and J. Am. Chem. Soc., 117, 6414-6415 (1995), as being active polymerization catalysts especially for polymerization of α -olefins, either alone or in combination with polar comonomers such as vinyl chloride, alkyl acrylates and alkyl methacrylates.


Additional catalysts include derivatives of Group 3, 4, or Lanthanide metals
 25 which are in the +2, +3, or +4 formal oxidation state. Preferred compounds include metal complexes containing from 1 to 3 π -bonded anionic or neutral ligand groups, which may be cyclic or non-cyclic delocalized π -bonded anionic ligand groups. Exemplary of such π -bonded anionic ligand groups are conjugated or nonconjugated, cyclic or non-cyclic dienyl groups, allyl groups, boratabenzene groups, and arene
 30 groups. By the term " π -bonded" is meant that the ligand group is bonded to the transition metal by a sharing of electrons from a partially delocalized π -bond.

Each atom in the delocalized π -bonded group may independently be substituted with a radical selected from the group consisting of hydrogen, halogen, hydrocarbyl, halohydrocarbyl, hydrocarbyl-substituted metalloid radicals wherein the

metalloid is selected from Group 14 of the Periodic Table of the Elements, and such hydrocarbyl- or hydrocarbyl-substituted metalloid radicals further substituted with a Group 15 or 16 hetero atom containing moiety. Included within the term "hydrocarbyl" are C₁₋₂₀ straight, branched and cyclic alkyl radicals, C₆₋₂₀ aromatic radicals, C₇₋₂₀ alkyl-substituted aromatic radicals, and C₇₋₂₀ aryl-substituted alkyl radicals. In addition two or more such radicals may together form a fused ring system, including partially or fully hydrogenated fused ring systems, or they may form a metallocycle with the metal. Suitable hydrocarbyl-substituted organometalloid radicals include mono-, di- and tri-substituted organometalloid radicals of Group 14 elements wherein each of the hydrocarbyl groups contains from 1 to 20 carbon atoms. Examples of suitable hydrocarbyl-substituted organometalloid radicals include trimethylsilyl, triethylsilyl, ethyldimethylsilyl, methyldiethylsilyl, triphenylgermyl, and trimethylgermyl groups. Examples of Group 15 or 16 hetero atom containing moieties include amine, phosphine, ether or thioether moieties or divalent derivatives thereof, e. g. amide, phosphide, ether or thioether groups bonded to the transition metal or Lanthanide metal, and bonded to the hydrocarbyl group or to the hydrocarbyl-substituted metalloid containing group.


Examples of suitable anionic, delocalized π -bonded groups include cyclopentadienyl, indenyl, fluorenyl, tetrahydroindenyl, tetrahydrofluorenyl, octahydrofluorenyl, pentadienyl, cyclohexadienyl, dihydroanthracenyl, hexahydroanthracenyl, decahydroanthracenyl groups, and boratabenzene groups, as well as C₁₋₁₀ hydrocarbyl-substituted or C₁₋₁₀ hydrocarbyl-substituted silyl substituted derivatives thereof. Preferred anionic delocalized π -bonded groups are cyclopentadienyl, pentamethylcyclopentadienyl, tetramethylcyclopentadienyl, tetramethylsilylcyclopentadienyl, indenyl, 2,3-dimethylindenyl, fluorenyl, 2-methylindenyl, 2-methyl-4-phenylindenyl, tetrahydrofluorenyl, octahydrofluorenyl, and tetrahydroindenyl.

The boratabenzenes are anionic ligands which are boron containing analogues to benzene. They are previously known in the art having been described by G. Herberich, et al., in Organometallics, 14,1, 471-480 (1995). Preferred boratabenzenes correspond to the formula:



wherein R" is selected from the group consisting of hydrocarbyl, silyl, or germyl, said R" having up to 20 non-hydrogen atoms. In complexes involving divalent derivatives of such delocalized π -bonded groups one atom thereof is bonded by means of a covalent bond or a covalently bonded divalent group to another atom of the complex thereby forming a bridged system.

A suitable class of catalysts are transition metal complexes corresponding to the formula:



wherein:

Lp is an anionic, delocalized, π -bonded group that is bound to M, containing up to 50 non-hydrogen atoms, optionally two Lp groups may be joined together forming a bridged structure, and further optionally one Lp may be bound to X;

M is a metal of Group 4 of the Periodic Table of the Elements in the +2, +3 or +4 formal oxidation state;

X is an optional, divalent substituent of up to 50 non-hydrogen atoms that together with Lp forms a metallocycle with M;

X' is an optional neutral ligand having up to 20 non-hydrogen atoms;

X" each occurrence is a monovalent, anionic moiety having up to 40 non-hydrogen atoms, optionally, two X" groups may be covalently bound together forming a divalent dianionic moiety having both valences bound to M, or, optionally 2 X" groups may be covalently bound together to form a neutral, conjugated or nonconjugated diene that is π -bonded to M (whereupon M is in the +2 oxidation state), or further optionally one or more X" and one or more X' groups may be bonded together thereby forming a moiety that is both covalently bound to M and coordinated thereto by means of Lewis base functionality;

l is 0, 1 or 2;

m is 0 or 1;


n is a number from 0 to 3;

p is an integer from 0 to 3; and

the sum, $l+m+p$, is equal to the formal oxidation state of M, except when 2 X" groups together form a neutral conjugated or non-conjugated diene that is π -bonded to M, in which case the sum $l+m$ is equal to the formal oxidation state of M.

- Preferred complexes include those containing either one or two Lp groups. The latter complexes include those containing a bridging group linking the two Lp groups. Preferred bridging groups are those corresponding to the formula $(ER^*_2)_x$ wherein E is silicon, germanium, tin, or carbon, R^* independently each occurrence is
- 5 hydrogen or a group selected from silyl, hydrocarbyl, hydrocarbyloxy and combinations thereof, said R^* having up to 30 carbon or silicon atoms, and x is 1 to 8. Preferably, R^* independently each occurrence is methyl, ethyl, propyl, benzyl, tert-butyl, phenyl, methoxy, ethoxy or phenoxy.

- Examples of the complexes containing two Lp groups are compounds
- 10 corresponding to the formula:



wherein:

- M is titanium, zirconium or hafnium, preferably zirconium or hafnium, in the +2 or +4 formal oxidation state;
- 15 R^3 in each occurrence independently is selected from the group consisting of hydrogen, hydrocarbyl, silyl, germyl, cyano, halo and combinations thereof, said R^3 having up to 20 non-hydrogen atoms, or adjacent R^3 groups together form a divalent derivative (that is, a hydrocarbadiyl, siladiyl or germadiyl group) thereby forming a fused ring system, and
- 20 X'' independently each occurrence is an anionic ligand group of up to 40 non-hydrogen atoms, or two X'' groups together form a divalent anionic ligand group of up to 40 non-hydrogen atoms or together are a conjugated diene having from 4 to 30 non-hydrogen atoms forming a π -complex with M, whereupon M is in the +2 formal oxidation state, and
- 25 R^* , E and x are as previously defined.

The foregoing metal complexes are especially suited for the preparation of polymers having stereoregular molecular structure. In such capacity it is preferred that the complex possesses C_s symmetry or possesses a chiral, stereorigid structure.

Examples of the first type are compounds possessing different delocalized π -bonded systems, such as one cyclopentadienyl group and one fluorenyl group. Similar systems based on Ti(IV) or Zr(IV) were disclosed for preparation of syndiotactic olefin polymers in Ewen, et al., J. Am. Chem. Soc. 110, 6255-6256 (1980). Examples of
 5 chiral structures include rac bis-indenyl complexes. Similar systems based on Ti(IV) or Zr(IV) were disclosed for preparation of isotactic olefin polymers in Wild et al., J. Organomet. Chem., 232, 233-47, (1982).


Exemplary bridged ligands containing two π -bonded groups are:
 dimethylbis(cyclopentadienyl)silane, dimethylbis(tetramethylcyclopentadienyl)silane,
 10 dimethylbis(2-ethylcyclopentadien-1-yl)silane, dimethylbis(2-t-butylcyclopentadien-1-yl)silane, 2,2-bis(tetramethylcyclopentadienyl)propane, dimethylbis(inden-1-yl)silane, dimethylbis(tetrahydroinden-1-yl)silane, dimethylbis(fluoren-1-yl)silane, dimethylbis(tetrahydrofluoren-1-yl)silane, dimethylbis(2-methyl-4-phenylinden-1-yl)-silane, dimethylbis(2-methylinden-1-yl)silane, dimethyl(cyclopentadienyl)(fluoren-1-yl)silane,
 15 dimethyl(cyclopentadienyl)(octahydrofluoren-1-yl)silane, dimethyl(cyclopentadienyl)(tetrahydrofluoren-1-yl)silane, (1, 1, 2, 2-tetramethyl)-1, 2-bis(cyclopentadienyl)disilane, (1, 2-bis(cyclopentadienyl)ethane, and dimethyl(cyclopentadienyl)-1-(fluoren-1-yl)methane.

Preferred X" groups are selected from hydride, hydrocarbyl, silyl, germyl,
 20 halohydrocarbyl, halosilyl, silylhydrocarbyl and aminohydrocarbyl groups, or two X" groups together form a divalent derivative of a conjugated diene or else together they form a neutral, π -bonded, conjugated diene. Most preferred X" groups are C₁₋₂₀ hydrocarbyl groups.

A further class of metal complexes utilized in the present invention
 25 corresponds to the preceding formula $L_pMX'_mX''_nX'''_p$, or a dimer thereof, wherein X is a divalent substituent of up to 50 non-hydrogen atoms that together with Lp forms a metallocycle with M.

Preferred divalent X substituents include groups containing up to 30 non-hydrogen atoms comprising at least one atom that is oxygen, sulfur, boron or a
 30 member of Group 14 of the Periodic Table of the Elements directly attached to the delocalized π -bonded group, and a different atom, selected from the group consisting of nitrogen, phosphorus, oxygen or sulfur that is covalently bonded to M.

A preferred class of such Group 4 metal coordination complexes used according to the present invention corresponds to the formula:



wherein:

M is titanium or zirconium, preferably titanium in the +2, +3, or +4 formal oxidation state;

5 R^3 in each occurrence independently is selected from the group consisting of hydrogen, hydrocarbyl, silyl, germyl, cyano, halo and combinations thereof, said R^3 having up to 20 non-hydrogen atoms, or adjacent R^3 groups together form a divalent derivative (that is, a hydrocarbadiyl, siladiyl or germadiyl group) thereby forming a fused ring system,

10 each X'' is a halo, hydrocarbyl, hydrocarbyloxy or silyl group, said group having up to 20 non-hydrogen atoms, or two X'' groups together form a neutral C_{5-30} conjugated diene or a divalent derivative thereof;

Y is -O-, -S-, -NR*, -PR*-; and

Z is SiR^*_2 , CR^*_2 , $SiR^*_2SiR^*_2$, $CR^*_2CR^*_2$, $CR^*=CR^*$, $CR^*_2SiR^*_2$, or GeR^*_2 ,

15 wherein R^* is as previously defined.

Illustrative Group 4 metal complexes that may be employed in the practice of the present invention include:

cyclopentadienyltitaniumtrimethyl,
 cyclopentadienyltitaniumtriethyl,
 20 cyclopentadienyltitaniumtriisopropyl,
 cyclopentadienyltitaniumtriphenyl,
 cyclopentadienyltitaniumtribenzyl,
 cyclopentadienyltitanium-2,4-dimethylpentadienyl,
 cyclopentadienyltitanium-2,4-dimethylpentadienyl•triethylphosphine,
 25 cyclopentadienyltitanium-2,4-dimethylpentadienyl•trimethylphosphine,
 cyclopentadienyltitaniumdimethylmethoxide,
 cyclopentadienyltitaniumdimethylchloride,
 pentamethylcyclopentadienyltitaniumtrimethyl,

- indenyltitaniumtrimethyl,
indenyltitaniumtriethyl,
indenyltitaniumtripropyl,
indenyltitaniumtriphenyl,
- 5 tetrahydroindenyltitaniumtribenzyl,
pentamethylcyclopentadienyltitaniumtriisopropyl,
pentamethylcyclopentadienyltitaniumtribenzyl,
pentamethylcyclopentadienyltitaniumdimethylmethoxide,
pentamethylcyclopentadienyltitaniumdimethylchloride,
- 10 bis(η^5 -2,4-dimethylpentadienyl)titanium,
bis(η^5 -2,4-dimethylpentadienyl)titanium•trimethylphosphine,
bis(η^5 -2,4-dimethylpentadienyl)titanium•triethylphosphine,
octahydrofluorenyltitaniumtrimethyl,
tetrahydroindenyltitaniumtrimethyl,
- 15 tetrahydrofluorenyltitaniumtrimethyl,
(tert-butylamido)(1,1-dimethyl-2,3,4,9,10- η -1,4,5,6,7,8-
hexahydronaphthalenyl)dimethylsilanetitaniumdimethyl,
(tert-butylamido)(1,1,2,3-tetramethyl-2,3,4,9,10- η -1,4,5,6,7,8-
hexahydronaphthalenyl)dimethylsilanetitaniumdimethyl,
- 20 (tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl) dimethylsilanetitanium dibenzyl,
(tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethylsilanetitanium dimethyl,
(tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)-1,2-ethanediyltitanium dimethyl,
(tert-butylamido)(tetramethyl- η^5 -indenyl)dimethylsilanetitanium dimethyl,
(tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethylsilane titanium (III)
- 25 2-(dimethylamino)benzyl;
(tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethylsilanetitanium (III) allyl,
(tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethylsilanetitanium (III)
2,4-dimethylpentadienyl,
(tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethylsilanetitanium (II)
- 30 1,4-diphenyl-1,3-butadiene,
(tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethylsilanetitanium (II)
1,3-pentadiene,
(tert-butylamido)(2-methylindenyl)dimethylsilanetitanium (II) 1,4-diphenyl-1,3-
butadiene,
- 35 (tert-butylamido)(2-methylindenyl)dimethylsilanetitanium (II) 2,4-hexadiene,
(tert-butylamido)(2-methylindenyl)dimethylsilanetitanium (IV) 2,3-dimethyl-1,3-
butadiene,

- (tert-butylamido)(2-methylindenyl)dimethylsilanetitanium (IV) isoprene,
 (tert-butylamido)(2-methylindenyl)dimethylsilanetitanium (IV) 1,3-butadiene,
 (tert-butylamido)(2,3-dimethylindenyl)dimethylsilanetitanium (IV)
 2,3-dimethyl-1,3-butadiene,
- 5 (tert-butylamido)(2,3-dimethylindenyl)dimethylsilanetitanium (IV)
 isoprene
 (tert-butylamido)(2,3-dimethylindenyl)dimethylsilanetitanium (IV) dimethyl
 (tert-butylamido)(2,3-dimethylindenyl)dimethylsilanetitanium (IV) dibenzyl
 (tert-butylamido)(2,3-dimethylindenyl)dimethylsilanetitanium (IV) 1,3-butadiene,
- 10 (tert-butylamido)(2,3-dimethylindenyl)dimethylsilanetitanium (II) 1,3-pentadiene,
 (tert-butylamido)(2,3-dimethylindenyl)dimethylsilanetitanium (II) 1,4-diphenyl-
 1,3-butadiene,
 (tert-butylamido)(2-methylindenyl)dimethylsilanetitanium (II) 1,3-pentadiene,
 (tert-butylamido)(2-methylindenyl)dimethylsilanetitanium (IV) dimethyl,
- 15 (tert-butylamido)(2-methylindenyl)dimethylsilanetitanium (IV) dibenzyl,
 (tert-butylamido)(2-methyl-4-phenylindenyl)dimethylsilanetitanium (II)
 1,4-diphenyl-1,3-butadiene,
 (tert-butylamido)(2-methyl-4-phenylindenyl)dimethylsilanetitanium (II) 1,3-pentadiene,
 (tert-butylamido)(2-methyl-4-phenylindenyl)dimethylsilanetitanium (II) 2,4-hexadiene,
- 20 (tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethyl- silanetitanium (IV)
 1,3-butadiene,
 (tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethylsilanetitanium (IV)
 2,3-dimethyl-1,3-butadiene,
 (tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethylsilanetitanium (IV)
- 25 isoprene,
 (tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethyl- silanetitanium (II)
 1,4-dibenzyl-1,3-butadiene,
 (tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethylsilanetitanium (II)
 2,4-hexadiene,
- 30 (tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl)dimethyl- silanetitanium (II)
 3-methyl-1,3-pentadiene,
 (tert-butylamido)(2,4-dimethylpentadien-3-yl)dimethylsilanetitaniumdimethyl,
 (tert-butylamido)(6,6-dimethylcyclohexadienyl)dimethylsilanetitaniumdimethyl,
 (tert-butylamido)(1,1-dimethyl-2,3,4,9,10- η -1,4,5,6,7,8-hexahydronaphthalen-4-
- 35 yl)dimethylsilanetitaniumdimethyl,
 (tert-butylamido)(1,1,2,3-tetramethyl-2,3,4,9,10- η -1,4,5,6,7,8-hexahydronaphthalen-
 4-yl)dimethylsilanetitaniumdimethyl

- (tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl methylphenylsilanetitanium (IV) dimethyl,
 (tert-butylamido)(tetramethyl- η^5 -cyclopentadienyl methylphenylsilanetitanium (II) 1,4-diphenyl-1,3-butadiene,
 5 1-(tert-butylamido)-2-(tetramethyl- η^5 -cyclopentadienyl)ethanediyltitanium (IV) dimethyl, and
 1-(tert-butylamido)-2-(tetramethyl- η^5 -cyclopentadienyl)ethanediyl- titanium (II) 1,4-diphenyl-1,3-butadiene.

Complexes containing two Lp groups including bridged complexes suitable for
 10 use in the present invention include:

- bis(cyclopentadienyl)zirconiumdimethyl,
 bis(cyclopentadienyl)zirconium dibenzyl,
 bis(cyclopentadienyl)zirconium methyl benzyl,
 bis(cyclopentadienyl)zirconium methyl phenyl,
 15 bis(cyclopentadienyl)zirconiumdiphenyl,
 bis(cyclopentadienyl)titanium-allyl,
 bis(cyclopentadienyl)zirconiummethoxymethoxide,
 bis(cyclopentadienyl)zirconiummethylchloride,
 bis(pentamethylcyclopentadienyl)zirconiumdimethyl,
 20 bis(pentamethylcyclopentadienyl)titaniumdimethyl,
 bis(indenyl)zirconiumdimethyl,
 indenylfluorenyl zirconiumdimethyl,
 bis(indenyl)zirconiummethyl(2-(dimethylamino)benzyl),
 bis(indenyl)zirconiummethyltrimethylsilyl,
 25 bis(tetrahydroindenyl)zirconiummethyltrimethylsilyl,
 bis(pentamethylcyclopentadienyl)zirconiummethylbenzyl,
 bis(pentamethylcyclopentadienyl)zirconiumdibenzyl,
 bis(pentamethylcyclopentadienyl)zirconiummethoxymethoxide,
 bis(pentamethylcyclopentadienyl)zirconiummethylchloride,
 30 bis(methylethylcyclopentadienyl)zirconiumdimethyl,
 bis(butylcyclopentadienyl)zirconiumdibenzyl,
 bis(t-butylcyclopentadienyl)zirconiumdimethyl,
 bis(ethyltetramethylcyclopentadienyl)zirconiumdimethyl,
 bis(methylpropylcyclopentadienyl)zirconiumdibenzyl,
 35 bis(trimethylsilylcyclopentadienyl)zirconiumdibenzyl,
 dimethylsilyl-bis(cyclopentadienyl)zirconiumdimethyl,
 dimethylsilyl-bis(tetramethylcyclopentadienyl)titanium (III) allyl

dimethylsilyl-bis(t-butylcyclopentadienyl)zirconiumdichloride,
 dimethylsilyl-bis(n-butylcyclopentadienyl)zirconiumdichloride,
 (methylene-bis(tetramethylcyclopentadienyl)titanium(III) 2-(dimethylamino)benzyl,
 (methylene-bis(n-butylcyclopentadienyl)titanium(III) 2-(dimethylamino)benzyl,
 5 dimethylsilyl-bis(indenyl)zirconiumbenzylchloride,
 dimethylsilyl-bis(2-methylindenyl)zirconiumdimethyl,
 dimethylsilyl-bis(2-methyl-4-phenylindenyl)zirconiumdimethyl,
 dimethylsilyl-bis(2-methylindenyl)zirconium-1,4-diphenyl-1,3-butadiene,
 dimethylsilyl-bis(2-methyl-4-phenylindenyl)zirconium (II) 1,4-diphenyl-1,3-butadiene,
 10 dimethylsilyl-bis(tetrahydroindenyl)zirconium(II) 1,4-diphenyl-1,3-butadiene,
 dimethylsilyl-bis(fluorenyl)zirconiummethylchloride,
 dimethylsilyl-bis(tetrahydrofluorenyl)zirconium bis(trimethylsilyl),
 (isopropylidene)(cyclopentadienyl)(fluorenyl)zirconiumdibenzyl, and
 dimethylsilyl(tetramethylcyclopentadienyl)(fluorenyl)zirconium dimethyl.

15 Other catalysts, especially catalysts containing other Group 4 metals, will, of course, be apparent to those skilled in the art.

The cocatalysts of the invention may also be used in combination with a an oligomeric or polymeric alumoxane compound, a tri(hydrocarbyl)aluminum compound, a di(hydrocarbyl)(hydrocarbyloxy)aluminum compound, a
 20 di(hydrocarbyl)(dihydrocarbyl-amido)aluminum compound, a bis(dihydrocarbyl-amido)(hydrocarbyl)aluminum compound, a di(hydrocarbyl)amido(disilyl)aluminum compound, a di(hydrocarbyl)-amido(hydrocarbyl)(silyl)aluminum compound, a bis(dihydrocarbylamido)(silyl)aluminum compound, or a mixture of the foregoing compounds, having from 1 to 20 non-hydrogen atoms in each hydrocarbyl,
 25 hydrocarbyloxy, or silyl group, if desired. These aluminum compounds are usefully employed for their beneficial ability to scavenge impurities such as oxygen, water, and aldehydes from the polymerization mixture.

Preferred aluminum compounds include C₂₋₆ trialkyl aluminum compounds, especially those wherein the alkyl groups are ethyl, propyl, isopropyl, n-butyl, isobutyl,
 30 pentyl, neopentyl, or isopentyl, dialkyl(aryloxy)aluminum compounds containing from 1-6 carbons in the alkyl group and from 6 to 18 carbons in the aryl group (especially (3,5-di(t-butyl)-4-methylphenoxy)diisobutylaluminum), methylalumoxane, modified methylalumoxane and diisobutylalumoxane. The molar ratio of aluminum compound to metal complex is preferably from 1:10,000 to 1000:1, more preferably from 1:5000
 35 to 100:1, most preferably from 1:100 to 100:1.

The molar ratio of catalyst/cocatalyst employed preferably ranges from 10:1 to 1:10, more preferably from 2.1:1 to 1:1.5, most preferably from 2.05:1 to 1:1. Mixtures of the activating cocatalysts of the present invention may also be employed if desired.

5 Suitable addition polymerizable monomers include ethylenically unsaturated monomers, acetylenic compounds, conjugated or non-conjugated dienes, and polyenes. Preferred monomers include olefins, for examples alpha-olefins having from 2 to 20,000, preferably from 2 to 20, more preferably from 2 to 8 carbon atoms and combinations of two or more of such alpha-olefins. Particularly suitable alpha-
10 olefins include, for example, ethylene, propylene, 1-butene, 1-pentene, 4-methylpentene-1, 1-hexene, 1-heptene, 1-octene, 1-nonene, 1-decene, 1-undecene, 1-dodecene, 1-tridecene, 1-tetradecene, 1-pentadecene, or combinations thereof, as well as long chain vinyl terminated oligomeric or polymeric reaction products formed during the polymerization, and C₁₀₋₃₀ α -olefins specifically added to the reaction
15 mixture in order to produce relatively long chain branches in the resulting polymers. Preferably, the alpha-olefins are ethylene, propene, 1-butene, 4-methyl-pentene-1, 1-hexene, 1-octene, and combinations of ethylene and/or propene with one or more of such other alpha-olefins. Other preferred monomers include styrene, halo- or alkyl substituted styrenes, tetrafluoroethylene, vinylcyclobutene, 1,4-hexadiene,
20 dicyclopentadiene, ethylidene norbornene, and 1,7-octadiene. Mixtures of the above-mentioned monomers may also be employed.

In general, the polymerization may be accomplished at conditions well known in the prior art for Ziegler-Natta or Kaminsky-Sinn type polymerization reactions. Suspension, solution, slurry, gas phase or high pressure, whether employed in batch
25 or continuous form or other process conditions, may be employed if desired. Examples of such well known polymerization processes are depicted in WO 88/02009, U.S. Patent Nos. 5,084,534, 5,405,922, 4,588,790, 5,032,652, 4,543,399, 4,564,647, 4,522,987, and elsewhere. Preferred polymerization temperatures are from 0-250°C. Preferred polymerization pressures are from atmospheric to 3000
30 atmospheres.

Preferred processing conditions include solution polymerization, more preferably continuous solution polymerization processes, conducted in the presence of an aliphatic or alicyclic liquid diluent. By the term "continuous polymerization" is meant that at least the products of the polymerization are continuously removed from
35 the reaction mixture, such as for example by devolatilization of a portion of the reaction mixture. Preferably one or more reactants are also continuously added to

the polymerization mixture during the polymerization. Examples of suitable aliphatic or alicyclic liquid diluents include straight and branched-chain hydrocarbons such as isobutane, butane, pentane, hexane, heptane, octane, and mixtures thereof; alicyclic hydrocarbons such as cyclohexane, cycloheptane, methylcyclohexane, methylcycloheptane, and mixtures thereof; and perfluorinated hydrocarbons such as perfluorinated C₄₋₁₀ alkanes. Suitable diluents also include aromatic hydrocarbons (particularly for use with aromatic α -olefins such as styrene or ring alkyl-substituted styrenes) including toluene, ethylbenzene or xylene, as well as liquid olefins (which may act as monomers or comonomers) including ethylene, propylene, butadiene, cyclopentene, 1-hexene, 3-methyl-1-pentene, 4-methyl-1-pentene, 1,4-hexadiene, 1-octene, 1-decene, styrene, divinylbenzene, allylbenzene, vinyltoluene (including all isomers alone or in admixture). Mixtures of the foregoing are also suitable.

In most polymerization reactions the molar ratio of catalyst:polymerizable compounds employed is from 10⁻¹²:1 to 10⁻¹:1, more preferably from 10⁻¹²:1 to 10⁻⁵:1.

The catalyst composition of the invention may also be utilized in combination with at least one additional homogeneous or heterogeneous polymerization catalyst in separate reactors connected in series or in parallel to prepare polymer blends having desirable properties. An example of such a process is disclosed in WO 94/00500, equivalent to U. S. Serial Number 07/904,770. A more specific process is disclosed in copending application U. S. Serial Number 08/10958, filed January 29, 1993.

Molecular weight control agents can be used in combination with the present cocatalysts. Examples of such molecular weight control agents include hydrogen, trialkyl aluminum compounds or other known chain transfer agents. A particular benefit of the use of the present cocatalysts is the ability (depending on reaction conditions) to produce narrow molecular weight distribution α -olefin homopolymers and copolymers in greatly improved catalyst efficiencies. Preferred polymers have Mw/Mn of less than 2.5, more preferably less than 2.3. Such narrow molecular weight distribution polymer products are highly desirable due to improved tensile strength properties.

The catalyst composition of the present invention can also be employed to advantage in the gas phase polymerization and copolymerization of olefins. Gas phase processes for the polymerization of olefins, especially the homopolymerization and copolymerization of ethylene and propylene, and the copolymerization of ethylene with higher α olefins such as, for example, 1-butene, 1-hexene, 4-

methyl-1-pentene are well known in the art. Such processes are used commercially on a large scale for the manufacture of high density polyethylene (HDPE), medium density polyethylene (MDPE), linear low density polyethylene (LLDPE) and polypropylene.

5 The gas phase process employed can be, for example, of the type which employs a mechanically stirred bed or a gas fluidized bed as the polymerization reaction zone. Preferred is the process wherein the polymerization reaction is carried out in a vertical cylindrical polymerization reactor containing a fluidized bed of polymer particles supported above a perforated plate, the fluidisation grid, by a flow
10 of fluidisation gas.

 The gas employed to fluidize the bed comprises the monomer or monomers to be polymerized, and also serves as a heat exchange medium to remove the heat of reaction from the bed. The hot gases emerge from the top of the reactor, normally via a tranquilization zone, also known as a velocity reduction zone, having a wider
15 diameter than the fluidized bed and wherein fine particles entrained in the gas stream have an opportunity to gravitate back into the bed. It can also be advantageous to use a cyclone to remove ultra-fine particles from the hot gas stream. The gas is then normally recycled to the bed by means of a blower or compressor and a one or more heat exchangers to strip the gas of the heat of polymerization.

20 A preferred method of cooling of the bed, in addition to the cooling provided by the cooled recycle gas, is to feed a volatile liquid to the bed to provide an evaporative cooling effect. The volatile liquid employed in this case can be, for example, a volatile inert liquid, for example, a saturated hydrocarbon having 3 to 8, preferably 4 to 6, carbon atoms. In the case that the monomer or comonomer itself
25 is a volatile liquid, or can be condensed to provide such a liquid this can be suitably be fed to the bed to provide an evaporative cooling effect. Examples of olefin monomers which can be employed in this manner are olefins containing from 3 to eight, preferably from 3 to six carbon atoms. The volatile liquid evaporates in the hot fluidized bed to form gas which mixes with the fluidizing gas. If the volatile liquid is a
30 monomer or comonomer, it will undergo some polymerization in the bed. The evaporated liquid then emerges from the reactor as part of the hot recycle gas, and enters the compression/heat exchange part of the recycle loop. The recycle gas is cooled in the heat exchanger and, if the temperature to which the gas is cooled is below the dew point, liquid will precipitate from the gas. This liquid is desirably
35 recycled continuously to the fluidized bed. It is possible to recycle the precipitated liquid to the bed as liquid droplets carried in the recycle gas stream, as described, for example, in EP-A-89691, US-A-4543399, WO 94/25495 and US-A-5352749. A

particularly preferred method of recycling the liquid to the bed is to separate the liquid from the recycle gas stream and to reinject this liquid directly into the bed, preferably using a method which generates fine droplets of the liquid within the bed. This type of process is described in WO 94/28032.

5 The polymerization reaction occurring in the gas fluidized bed is catalyzed by the continuous or semi-continuous addition of catalyst. [Such catalyst can be supported on an inorganic or organic support material if desired. The catalyst can also be subjected to a prepolymerization step, for example, by polymerizing a small quantity of olefin monomer in a liquid inert diluent, to provide a catalyst composite
10 comprising catalyst particles embedded in olefin polymer particles.

 The polymer is produced directly in the fluidized bed by catalyzed (co)polymerization of the monomer(s) on the fluidized particles of catalyst, supported catalyst or prepolymer within the bed. Start-up of the polymerization reaction is achieved using a bed of preformed polymer particles, which, preferably, is similar to
15 the target polyolefin, and conditioning the bed by drying with inert gas or nitrogen prior to introducing the catalyst, the monomer(s) and any other gases which it is desired to have in the recycle gas stream, such as a diluent gas, hydrogen chain transfer agent, or an inert condensable gas when operating in gas phase condensing mode. The produced polymer is discharged continuously or discontinuously from the
20 fluidized bed as desired, optionally exposed to a catalyst kill and optionally pelletized.

 It is understood that the present invention is operable in the absence of any component which has not been specifically disclosed. The following examples are provided in order to further illustrate the invention and are not to be construed as limiting. Unless stated to the contrary, all parts and percentages are expressed on a
25 weight basis.

Example 1

Preparation of (tetrafluoro-1,4-phenylene)-bis-(di(pentafluorophenyl)borane) ((C₆F₅)₂ B-C₆F₄-B (C₆F₅)₂)

 Into a thick-walled flask containing a J. Young valve, 1,4-C₆F₄(SnMe₃)₂ (0.60
30 g, 1.26 mmol) and (C₆F₅)₂BCl (2.87 g, 7.56 mmol) were added. Toluene (40 mL) was added, the flask was evacuated to 0.1 torr, and the J. Young valve was closed. The flask was heated at 140°C for 72 h. The solvent was removed *in vacuo*, and the residue washed with pentane (4 x 20 mL). The resulting light yellow solid was


exposed to dynamic vacuum (10^{-5} torr) for 12 h, giving the desired product as a microcrystalline pale yellow solid (0.75 g, 71 percent).

^{19}F NMR (CD_2Cl_2): δ -125.7 (br, 8F, ortho C_6F_5), -128.2 (br, 4F, C_6F_4), -141.1 (br, 4F, para C_6F_5), -159.1 (br, 8F, meta C_6F_5) ppm.

5 MS (EI, 6.3 V): m/e 838 (M^+ , 100 percent).

Activation of Metal Complex

Combination of the diborane compound of Example 1 with the zirconocene biscyclopentadienylzirconium dimethyl in CD_2Cl_2 at 25°C in a 1:1 and a 2:1 atomic ratio (Zr:B) gave two cationic reaction products corresponding to the monoanionic salt (1) and the dianionic salt (2) derivatives according to the following scheme.



NMR data for 1:

^1H NMR (CD_2Cl_2): δ 6.39 (s, 10H), 0.68 (s, 3H), 0.31 (br, 3H) ppm.

15 ^{19}F NMR (CD_2Cl_2): δ -127.5 (d, $^3J_{\text{FF}} = 21$ Hz, 3F), -129.3 (t, $^3J_{\text{FF}} = 18$ Hz, 1F), -131.8 (m, 1F), -132.4 (m, 6F), -136.7 (s, 1F), -144.8 (s, 1F), -159.0 (br, 1F), -160.0 (m, 4F), -163.8 (br, 2F), -164.2 (t, $^3J_{\text{FF}} = 19$ Hz, 3F)

NMR data for 2:

20 ^1H NMR (CD_2Cl_2): δ 6.29 (s, 20H), 0.45 (br, 6H), 0.29 (br, 6H) ppm.

^{19}F NMR (CD_2Cl_2): δ -131.5 (d, $^3J_{\text{FF}} = 18$ Hz, 8F, ortho C_6F_5), -142.8 (br, 4F, C_6F_4), -162.4 (br, 4F, para C_6F_5), -165.3 (br, 8F, meta C_6F_5) ppm.

Example 2

Preparation of (tetrafluoro-1,2-phenylene)-bis-(di(pentafluorophenyl)borane)

25 A) Preparation of $[1,2-\text{C}_6\text{F}_4\text{BCl}]_2$

Excess BCl_3 (ca. 5.0 g, 44 mmol) was condensed into a thick-walled flask containing a J. Young valve and $1,2\text{-C}_6\text{F}_4(\text{SnMe}_3)_2$ (5.3 g, 11.1 mmol), at -196°C . The flask was evacuated to 0.05 torr, and the J. Young valve was closed. The reaction mixture was heated at 180°C for 18 h. The excess BCl_3 was removed under
5 dynamic vacuum, giving a slightly moist, beige solid. The product was extracted with pentane (3 x 20 mL), leaving behind 65 percent of the Me_3SnCl byproduct. The remaining Me_3SnCl was sublimed away at $40^\circ\text{C}/10^{-5}$ torr. The product was then sublimed at $90^\circ\text{C}/10^{-5}$ torr, giving $[1,2\text{-C}_6\text{F}_4\text{BCl}]_2$ as a yellow solid (1.15 g, 53 percent).

10 ^{19}F NMR (C_6D_6): δ -122.7 (m, 4F), -143.9 (m, 4F) ppm.

^{13}C NMR (CDCl_3): δ 152.6 (d, $^1J_{\text{CF}} = 262$ Hz), 144.6 (d, $^1J_{\text{CF}} = 260$ Hz), 122.8 (br, B-C) ppm.

MS (EI, 8.7 V) (percent intensity): 392 (16), 391 (18), 390 (67), 389 (M^+ , 47), 388 (100), 387 (51), 342 (21), 318 (22), 304 (28), 250 (25), 201 (30).

15 Anal. Calcd for $\text{C}_{12}\text{F}_8\text{B}_2\text{Cl}_2$: C, 37.1; H, 0.0. Found: C, 38.2; H, 0.3.

B) Preparation of $[1,2\text{-C}_6\text{F}_4\text{B}(\text{C}_6\text{F}_5)]_2$,

Into a thick-walled flask containing a J. Young valve, $[1,2\text{-C}_6\text{F}_4\text{BCl}]_2$ (0.265 g, 0.68 mmol) and $(\text{C}_6\text{F}_5)_2\text{SnMe}_2$ (0.33 g, 0.68 mmol) were placed. Toluene (20 mL)
20 was added, the flask was evacuated to 0.1 torr, and the J. Young valve was closed. The reaction solution was heated to 140°C for 72 h, giving a bright yellow solution. The solution was concentrated to 10 mL, and then heated to dissolve all solids. Slow cooling of this solution to -78°C gave 2 as light yellow crystals. This solid was found to contain a small amount of Me_2SnCl_2 , which can be removed under dynamic
25 vacuum (10^{-5} torr/12 h), giving the desired product, $[1,2\text{-C}_6\text{F}_4\text{B}(\text{C}_6\text{F}_5)]_2$, as a light yellow crystalline solid (0.35 g, 68 percent). Alternatively, because of the sensitivity of the compound, the crude reaction solution can be exposed to dynamic vacuum (10^{-5} torr) for 12 h, giving the product in >95 percent purity, without the need for crystallization.

30 ^{19}F NMR (d_8 -toluene): δ -118.2 (br, 4F, ortho C_6F_4), -133.9 (dd, $^3J_{\text{FF}} = 25.1$ Hz; $^4J_{\text{FF}} = 7.9$ Hz, 4F, ortho C_6F_5), -138.9 (m, 4F, meta C_6F_4), -152.1 (t, $^3J_{\text{FF}} = 21$ Hz, 2F, para C_6F_5), -161.4 (ddd, $^3J_{\text{FF}} = 22$ Hz; $^3J_{\text{FF}} = 22$ Hz, $^5J_{\text{FF}} = 7$ Hz, 4F, meta C_6F_5) ppm.


^{13}C NMR (CDCl_3): δ 156.1 (d, $^1J_{\text{CF}} = 267$ Hz), 145.9 (d, $^1J_{\text{CF}} = 265$ Hz), 144.3 (d, $^1J_{\text{CF}} = 241$ Hz), 141.6 (d, $^1J_{\text{CF}} = 265$ Hz), 137.6 (d, $^1J_{\text{CF}} = 253$ Hz), 128.3 (br), 123.7 (br) ppm.

Crystal Data

- 5 $\text{C}_{24}\text{F}_{18}\text{B}_2 \cdot (2\text{C}_7\text{H}_8)$; monoclinic, space group $\text{P}2_1/\text{c}$; $a = 22.836(4)$, $b = 10.846(3)$, $c = 13.767(3)$ Å; $\beta = 99.66(2)^\circ$; $V = 3361(1)$ Å³; $Z = 4$; $d_{\text{calcd}} = 1.652$ g/cm³; at -120°C . The structure was solved by direct methods. Owing to the paucity of data, the fluorine atoms were refined anisotropically and the remaining non-hydrogen atoms were refined isotropically. Hydrogen atoms were included in "idealized"
- 10 positions and not refined. The final cycle of full-matrix least-squares refinement was based on 2042 observed reflections ($I > 3.00\sigma(I)$) and 324 variable parameters and converged (largest parameter shift was 0.12 times its esd) with unweighted and weighted agreement factors of $R = 0.051$ and $R_w = 0.039$. For clarity of the crystallographic discussion, it should be noted that there are two half molecules in the
- 15 asymmetric unit, and consequently there are two independent bond distances and angles for each bond distance/angle of 2.

Activation of Metal Complexes

- Combination of the diborane compound of Example 2 with the zirconocene biscyclopentadienylzirconium dimethyl in CD_2Cl_2 at 25°C in a 1:1 and a 2:1 atomic
- 20 ratio (Zr:B) gave two cationic reaction products corresponding to the monoanionic salt (3) and the dianionic salt (4) derivatives according to the following scheme.



NMR data for 3:

^{19}F (CD_2Cl_2 , 25°C): δ -123.1 (br, 2F), -132.4 (m, 3F), -134.0 (br, 2F), -134.8 (br, 1F), -145.0 (br, 2F), -155.4 (t, $^3J_{FF} = 21$ Hz, 1F), -158.9 (m, 2F), -160.0 (m, 1F),
 5 -161.8 (br, 1F), -162.5 (br, 1F), -164.1 (br, t, $^3J_{FF} = 21$ Hz, 2F) ppm.

1H (CD_2Cl_2 , 25°C): δ 6.34 (s, 10H), 0.66 (s, 3H), 0.17 (br, 3H) ppm.

NMR data for 4:

^{19}F (CD_2Cl_2 , 25°C): δ -132.7 (br, 2F, ortho C_6F_5), -134.7 (br, 4F, ortho C_6F_4),
 10 -136.4 (br, 2F, ortho C_6F_5), -161.6 (t, $^3J_{FF} = 20$ Hz, 2F, para C_6F_5), -162.6 (d, $^3J_{FF} = 19$
 Hz, 4F, meta C_6F_4), -164.4 (br, 2F, meta C_6F_5), -165.4 (br, 2F, meta C_6F_5) ppm.

1H (CD_2Cl_2 , 25°C): δ 6.22 (s, 10H), 0.69 (s, 3H), 0.19 (br, 3H) ppm; a slight excess of $(C_5H_5)_2ZrMe_2$ was present in this sample, and resonances attributable to this compound are also present.

Polymerizations

15 A two liter stirred reactor was charged with 640 mL Isopar™ E solvent, and 150 g of propylene. Hydrogen (25 ml at 35 Δ psi, 0.2 Δ MPa) was added as a molecular weight control agent. The reactor was heated to 70°C. The catalyst composition was prepared in a drybox by mixing together 0.005M toluene solutions of (t-butylamido)dimethyl(η^5 -tetramethylcyclopentadienyl)titanium dimethyl catalyst, and

- the compounds of examples 1 or 2 to give atomic ratios of B/Ti of 1:1 or 2:1. After a mixing time of 5 minutes the solutions were then transferred to an addition loop and injected into the reactor. The polymerization was allowed to proceed for 10 minutes while maintaining the reaction temperature at 70°C. The polymer solution was
- 5 transferred from the reactor into a glass kettle and dried in a vacuum oven for 16 hours at a maximum temperature of 120 °C. Results are contained in Table 1.

Table 1


run	μmol catalyst	cocatalyst (μmol)	reaction time min.	g polymer	efficiency (Kg polymer/ g Ti)
1*	6	FAB ¹ (6)	10	29.7	103
2	0.75	Ex. 2 (0.75)	10	70.9	1,974
3*	6	FAB ¹ (6)	10	37.0	129
4	1.5	Ex. 2 (0.75)	10	66.4	925
5	3	Ex. 1 (3)	18.5	29.7	207
6	6	Ex. 1 (3)	10	49.2	171

* not an example of the invention

¹. tris(pentafluorophenyl)borane

CLAIMS:

1. A compound corresponding to the formula:



wherein:

5 B^1 and B^2 independently each occurrence are Group 13 metal or metalloid atoms,

Z^* is an optional divalent bridging group containing from 1 to 20 atoms, not counting hydrogen atoms;

10 R^1 and R^2 independently each occurrence are monovalent, anionic ligand groups containing from 1 to 40 atoms, not counting hydrogen atoms, and, for cationic compounds, additionally comprising a dissociated cation moiety;

15 Ar^{f1} and Ar^{f2} independently each occurrence are monovalent, fluorinated organic groups containing from 6 to 100 carbon atoms, optionally, an Ar^{f1} and an R^2 group, or an Ar^{f2} and an R^1 group together form a divalent bridging group, and further optionally an Ar^{f1} group and an Ar^{f2} group together form a C_{6-100} divalent bridging group,


z is 0 or 1,

r and s independently are 0, 1 or 2, and

m and n are 1, 2 or 3;

20 with the proviso that when z is 0, at least one of Ar^{f1} and Ar^{f2} are joined together, and the sum of r , z and m is 3 or 4, in the former event B^1 is neutral and in the latter event B^1 is negatively charged, said charge being balanced by a cation component of one R^1 ; and the sum of s , z and n is 3 or 4, in the former event B^2 is neutral and in the latter event B^2 is negatively charged, said charge being balanced
25 by a cation component of one R^2 .

2. A compound according to claim 1 which is a neutral Lewis acid corresponding to the formula:



wherein:

B^1 and B^2 independently each occurrence are Group 13 metal or metalloid atoms,

5 Z^* is an optional divalent bridging group containing from 1 to 20 atoms, not counting hydrogen atoms;

R^1 and R^2 independently each occurrence are monovalent, anionic ligand groups containing from 1 to 40 atoms, not counting hydrogen atoms;

10 Ar^{f1} and Ar^{f2} independently each occurrence are monovalent, fluorinated organic groups containing from 6 to 100 carbon atoms, optionally, an Ar^{f1} and an R^2 group, or an Ar^{f2} and an R^1 group together form a divalent bridging group, and further optionally an Ar^{f1} group and an Ar^{f2} group together form a C_{6-100} divalent bridging group,

z is 0 or 1,


15 r and s independently are 0, 1 or 2, and

m and n are 1, 2 or 3; and

the sum of r , z and m is 3, and the sum of s , z and n is 3,

with the proviso that when z is 0, at least one of Ar^{f1} and Ar^{f2} are joined together.

20 3. The compound of Claim 2 corresponding to the formula:




wherein:


R^1 and R^2 independently each occurrence are C_{1-20} hydrocarbyl, halohydrocarbyl, or halocarbyl, and

Ar^{f1} - Ar^{f2} in combination, independently each occurrence, is a divalent fluoro-substituted aromatic group of from 6 to 20 carbons.

- 5 4. The compound of claim 3 corresponding to the formula::



- 5 5 A compound according to claim 1 which comprises a cation/ anion pair, corresponding to the formula:




- 10 wherein:

R^1 and R^2 independently each occurrence are C_{1-20} hydrocarbyl, halohydrocarbyl, or halocarbyl groups, and (when the sum of r , z and m is 4) one R^1 additionally comprises a cation selected from the group consisting of protonated cations of Bronsted acids, ferrocenium cations, carbonium cations, silylium cations, Ag^+ , and cationic derivatives of a Group 3-10 metal complex; and (when the sum of s , z and n is 4) one R^2 additionally comprises a cation selected from the group consisting of protonated cations of Bronsted acids, ferrocenium cations, carbonium cations, silylium cations, Ag^+ , and cationic derivatives of a Group 3-10 metal complex;

- 20 r and s are 0, 1 or 2 with the proviso that at least one of r or s is not 0, and the sum of r , z and m is 3 or 4 and the sum of s , z and n is 3 or 4, with the proviso that at least one of the foregoing sums is 4; and

Ar^{f1} and Ar^{f2} independently each occurrence are monovalent, fluorinated organic groups containing from 6 to 100 carbon atoms, and optionally an Ar^{f1} group and an Ar^{f2} group together form a C₆₋₁₀₀ divalent bridging group.

6. The compound of claim 5 corresponding to the formula:



5


wherein:

R¹ and R² independently each occurrence are C₁₋₂₀ hydrocarbyl, halohydrocarbyl, or halocarbyl groups, and (when connected to a negatively charged boron atom) one R¹ additionally comprises a cation selected from the group consisting of protonated cations of Bronsted acids, ferrocenium cations, carbonium cations, silylium cations, Ag⁺, and cationic derivatives of a Group 3-10 metal complex and (when connected to a negatively charged boron atom) one R² additionally comprises a cation selected from the group consisting of protonated cations of Bronsted acids, ferrocenium, carbonium cations, silylium cations, Ag⁺, and cationic derivatives of a Group 3-10 metal complex; and

15

an Ar^{f1} group and an Ar^{f2} group together form a C₆₋₂₀ divalent bridging group.

7. A compound according to claim 6 corresponding to the formula::



wherein, L^+ is a cation of a Bronsted acid, or a ferrocenium, carbonium, silylium, or Ag^+ cation.

- 5 8. A catalyst system for polymerization of α -olefins comprising, in combination, a Group 4 metal complex and a compound according to any one of claims 1-4 or 7, or the reaction product thereof.

9 A polymerization process comprising contacting one or more α -olefins under polymerization conditions with a catalyst system according to claim 8.

- 10 10. A process according to claim 9 which is a solution polymerization.

11. A polymerization process according to claim 10 which is a continuous solution polymerization.

12. A polymerization process according to claim 9 which is a gas phase polymerization.

INTERNATIONAL SEARCH REPORT

Intern al Application No

PCT/US 98/14106

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07F5/02 C08F10/00

According to International Patent Classification(IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07F C08F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 496 960 A (PIERS, W.E. ET AL.) 5 March 1996 see the whole document ---	1-12
X	US 5 447 895 A (MARKS, T.J. ET AL.) 5 September 1995 cited in the application see the whole document ---	1-12
P,X	DE 196 22 207 A (HOECHST) 4 December 1997 see the whole document ---	1-12
P,X	DE 196 32 557 A (HOECHST AG) 19 February 1998 see the whole document ---	1-12
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

10 November 1998

Date of mailing of the international search report

26/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Rinkel, L

INTERNATIONAL SEARCH REPORT

Intern al Application No

PCT/US 98/14106

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>PARKS, DANIEL J. ET AL: "Bis(pentafluorophenyl)borane: synthesis, properties, and hydroboration chemistry of a highly electrophilic borane reagent" ANGEW. CHEM., INT. ED. ENGL. (1995), 34(7), 809-11 CODEN: ACIEAY;ISSN: 0570-0833,1995, XP002083862 see Table 1</p> <p>---</p>	1-3
X	<p>CHEMICAL ABSTRACTS, vol. 128, no. 8, 23 February 1998 Columbus, Ohio, US; abstract no. 88939, GALSWORTHY, JANE R. ET AL: "Syntheses and characterization of amino(pentafluorophenyl)boranes. Crystal structure of '(Me3Si)2NB(C6F5)2!'" XP002083863 see abstract & POLYHEDRON (1997), VOLUME DATE 1998, 17(1), 119-124 CODEN: PLYHDE;ISSN: 0277-5387,1997,</p> <p>---</p>	1-3
X	<p>CHEMICAL ABSTRACTS, vol. 121, no. 20, 14 November 1994 Columbus, Ohio, US; abstract no. 231450, JIA, LI ET AL: "Cationic d0/f0 metallocene catalysts. properties of binuclear organoborane Lewis acid cocatalysts and weakly coordinating counteranions derived therefrom" XP002083864 see abstract & ORGANOMETALLICS (1994), 13(10), 3755-7 CODEN: ORGND7;ISSN: 0276-7333,1994,</p> <p>---</p>	1-3
X	<p>CHEMICAL ABSTRACTS, vol. 92, no. 13, 31 March 1980 Columbus, Ohio, US; abstract no. 111091, PHILLIPS, RODERIC J. ET AL: "Organothallium compounds. XVI. Formation of tris(polyfluorophenyl)thallium(III) compounds by reaction of bromobis(polyfluorophenyl)thallium(III) compounds with metal polyfluorobenzenesulfinates" XP002083865 see abstract & AUST. J. CHEM. (1979), 32(11), 2381-93 CODEN: AJCHAS;ISSN: 0004-9425,1979,</p> <p>---</p> <p>-/--</p>	1-3

INTERNATIONAL SEARCH REPORT

Intern al Application No

PCT/US 98/14106

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 69, no. 14, 30 September 1968 Columbus, Ohio, US; abstract no. 55894, DEACON, G. B. ET AL: "Vibrational spectra of organometallic compounds. III. Infrared spectra and assignments for bis(pentafluorophenyl)thallium(III) compounds" XP002083866 see abstract & SPECTROCHIM. ACTA, PART A (1968), 24(8), 1125-33 CODEN: SAMCAS, 1968, ---	1-3
P, X	CHEMICAL ABSTRACTS, vol. 129, Columbus, Ohio, US; abstract no. 161870, KOEHLER, KATRIN ET AL: "Bifunctional Lewis Acids. Synthesis and Olefin Polymerization Chemistry of the 1,1-Di'bis(perfluorophenyl)boryl!alkenes RCH:C'B(C6F5)2!2 (R = t-Bu, C6H5, C6F5)" XP002083867 see abstract & ORGANOMETALLICS (1998), 17(16), 3557-3566 CODEN: ORGND7; ISSN: 0276-7333, 1998, ---	1-3
P, X	CHEMICAL ABSTRACTS, vol. 129, Columbus, Ohio, US; abstract no. 41208, SPENCE, RUPERT E. V. H. ET AL: "Mechanistic Aspects of the Reactions of Bis(pentafluorophenyl)borane with the Dialkyl Zirconocenes Cp2ZrR2 (R = CH3, CH2SiMe3, and CH2C6H5)" XP002083868 see abstract & ORGANOMETALLICS (1998), 17(12), 2459-2469 CODEN: ORGND7; ISSN: 0276-7333, 1998, -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/14106

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5496960 A	05-03-1996	CA 2159651 A	24-05-1996
US 5447895 A	05-09-1995	CA 2185111 A	14-09-1995
		WO 9524269 A	14-09-1995
DE 19622207 A	04-12-1997	AU 2369597 A	11-12-1997
		CA 2206831 A	03-12-1997
		CN 1167767 A	17-12-1997
		EP 0811627 A	10-12-1997
		JP 10095786 A	14-04-1998
		NO 972507 A	04-12-1997
DE 19632557 A	19-02-1998	AU 3413397 A	19-02-1998
		CA 2212668 A	13-02-1998
		CN 1182093 A	20-05-1998
		CN 1174205 A	25-02-1998
		EP 0824112 A	18-02-1998
		EP 0824113 A	18-02-1998
		JP 10139806 A	26-05-1998
		JP 10158269 A	16-06-1998
		NO 973707 A	16-02-1998