



(86) Date de dépôt PCT/PCT Filing Date: 2007/10/18
(87) Date publication PCT/PCT Publication Date: 2008/04/24
(85) Entrée phase nationale/National Entry: 2009/04/17
(86) N° demande PCT/PCT Application No.: US 2007/022244
(87) N° publication PCT/PCT Publication No.: 2008/048668
(30) Priorité/Priority: 2006/10/18 (US60/852,604)

(51) Cl.Int./Int.Cl. *C07D 307/80* (2006.01),
A61K 31/343 (2006.01), *C07D 307/81* (2006.01)


(71) Demandeur/Applicant:
JANSSEN PHARMACEUTICA, N.V., BE

(72) Inventeurs/Inventors:
GRECO, MICHAEL N., US;
HAWKINS, MICHAEL J., US;
POWELL, EUGENE, US;
DE GARAVILLA, LAWRENCE, US;
MARYANOFF, BRUCE E., US

(74) Agent: OGILVY RENAULT LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : NOUVEAUX INHIBITEURS DE CHYMASE

(54) Title: NOVEL INHIBITORS OF CHYMASE



(57) Abrégé/Abstract:

The present invention is directed to a compound of Formula (I): or an enantiomer, diastereomer, polymorph or pharmaceutically acceptable salt thereof and methods for preparing said compounds and compositions, intermediates and derivatives thereof, and methods for treating inflammatory or serine protease mediated disorders.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 April 2008 (24.04.2008)

PCT

(10) International Publication Number
WO 2008/048668 A3

(51) International Patent Classification:

C07D 307/80 (2006.01) *A61K 31/343* (2006.01)
C07D 307/81 (2006.01)

(74) Agent: **STERCHO, Yuriy, P.**; Johnson & Johnson, 145
King of Prussia Road, Radnor, Pennsylvania 19087 (US).

(21) International Application Number:

PCT/US2007/022244

(22) International Filing Date: 18 October 2007 (18.10.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/852,604 18 October 2006 (18.10.2006) US

(71) Applicant (for all designated States except US):

JANSSEN PHARMACEUTICA, N.V. [BE/BE];
Turnhoutseweg 30, b-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GRECO, Michael, N.** [US/US]; 1634 Clearbrook Road, Lansdale, Pennsylvania 19446 (US). **HAWKINS, Michael, J.** [US/US]; 318 Heckler Street, Ambler, Pennsylvania 19002 (US). **POWELL, Eugene** [US/US]; 4801 Woodspring Drive, Pipersville, PA 18947-1064 (US). **DE GARAVILLA, Lawrence** [US/US]; 909 Noble Drive, Downingtown, Pennsylvania 19335 (US). **MARYANOFF, Bruce, E.** [US/US]; P.O. Box 239, New Hope, Pennsylvania 18938 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

(88) Date of publication of the international search report:

16 October 2008

(54) Title: NOVEL INHIBITORS OF CHYMASE



(I)

(57) Abstract: The present invention is directed to a compound of Formula (I): or an enantiomer, diastereomer, polymorph or pharmaceutically acceptable salt thereof and methods for preparing said compounds and compositions, intermediates and derivatives thereof, and methods for treating inflammatory or serine protease mediated disorders.

WO 2008/048668 A3

NOVEL INHIBITORS OF CHYMASE

CROSS REFERENCE TO RELATED APPLICATIONS

This present application claims benefit of U.S. Provisional Patent Application Serial No. 60/852604, filed October 18, 2006, which is incorporated
5 herein by reference in its entirety and for all purposes.

FIELD OF THE INVENTION

The present invention relates to certain novel compounds, methods for preparing compounds, compositions, intermediates and derivatives thereof and methods for treating inflammatory or serine protease mediated disorders. More
10 particularly, the compounds of the present invention are serine protease inhibitors useful for treating inflammatory or serine protease mediated disorders.

BACKGROUND OF THE INVENTION

Serine proteases represent a broad class of proteolytic enzymes that are
15 involved in physiological processes such as blood coagulation, complement activation, phagocytosis and turnover of damaged cell tissue. Human chymase (EC.3.4.21.39) is a glycosylated monomeric chymotrypsin-like serine protease (MW = 30 kDa) localized mainly in mast cell secretory granules. Chymase is thought to have a variety of functions, including degradation of extracellular
20 matrix proteins, cleavage of angiotensin I to angiotensin II (except in the rat), and activation of matrix proteases and cytokines. Endogenously, chymase is regulated by the serpins α 1-antichymotrypsin and α 1-protease.

Although the precise patho-physiological roles of chymase have yet to be determined, chymase has been implicated in microvascular leakage,
25 neutrophil accumulation, the stimulation of mucus secretion, and the modulation of cytokines. A potent, chymase-selective inhibitor may be indicated in inflammatory or serine protease (such as mast cell) mediated diseases such as asthma, pulmonary inflammation, and chronic obstructive pulmonary diseases (COPD). Because chymase can play a role in the

generation of cardiac and vascular wall angiotensin II, an inhibitor may have potential use as an antihypertensive treatment for vascular wall injury and inflammation (atherosclerosis/restenosis), as well as cardiac hypertrophy. Thus, small molecule inhibitors of chymase are likely to represent useful
5 therapeutic agents.

US Patent 5,508,273 to Beers, et al. and *Bioorganic & Med. Chem. Lett.*, **1995**, 5 (16), 1801-1806 describe phosphonic acid compounds useful in treating bone wasting diseases.


United States Patent Application 2005/0176769 describes aryl and
10 heteroaryl substituted phosphinic and phosphonic acid compounds useful in treating Inflammatory and serine protease mediated diseases.

Accordingly, it is an object of the present invention to provide phosphonic acid and phosphinic acid compounds that are serine protease inhibitors, in particular, inhibitors of chymase, useful for treating inflammatory or
15 serine protease mediated disorders. It is another object of the invention to provide a process for preparing phosphonic or phosphinic acid compounds, compositions, intermediates and derivatives thereof. It is a further object of the invention to provide methods for treating inflammatory or serine protease mediated disorders.

20

SUMMARY OF THE INVENTION

The present invention is directed to a compound of Formula (I):



25

and enantiomers, diastereomers, polymorphs or pharmaceutically acceptable salts thereof, wherein Ring A, R¹, R², R³, R⁴, n, W, X and Y are as defined herein.

The present invention provides phosphonic acid and phosphinic acid compounds that are serine protease inhibitors.

An example of a serine protease inhibited by a compound of the present invention is chymase.


5 Another example of the present invention includes chymase inhibitor compounds useful for treating inflammatory or serine protease mediated disorders.

The present invention further provides a process for preparing phosphonic or phosphinic acid compounds, compositions, intermediates and
10 derivatives thereof.

The present invention is also directed to a method for treating inflammatory or serine protease mediated disorders in a patient in need thereof comprising administering to the patient an effective amount of a compound of Formula (I).

15 DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to a compound of Formula (I):



wherein

R¹ is selected from the group consisting of hydrogen and C₁₋₄alkyl;

20 Ring A is selected from the group consisting of aryl, heteroaryl, benzo fused heterocyclyl, cycloalkyl and benzo fused cycloalkyl;

R² is one, two or three substituents independently selected from the group consisting of hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkoxy, -OCH₂-C₂₋₆alkenyl, C₁₋₆alkylthio, -OCF₃, -NH₂, -NH(C₁₋₆)alkyl,

25 -N(C₁₋₆)dialkyl, aryl, heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy

and nitro,

wherein, R^2 is optionally oxo when Ring A is heteroaryl or benzo fused heterocyclyl,

wherein any aryl-containing substituent of R^2 is optionally substituted with a

5 substituent independently selected from the group consisting of C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyl, C_{1-6} alkylthio, $-NH_2$, $-NH(C_{1-6})$ alkyl, $-N(C_{1-6})$ dialkyl, aryl, heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy and nitro,, and

wherein any of the foregoing C_{1-6} alkyl, C_{2-6} alkenyl or C_{1-6} alkoxy containing substituents of R^2 are optionally substituted with a substituent

10 independently selected from the group consisting of $-NR^{11}R^{12}$, aryl, heteroaryl, one, two or three halogen atoms and hydroxy;

R^{11} and R^{12} are independently hydrogen; C_{1-6} alkyl optionally substituted with hydroxy, aryl, $-C(=O)C_{1-4}$ alkoxy, or $-NR^{15}R^{16}$; or aryl;

15 R^{15} and R^{16} are substituents independently selected from the group consisting of hydrogen, C_{1-6} alkyl and aryl; optionally, R^{15} and R^{16} are each taken together with the atoms to which they are attached to form a ring of five to seven members;

R^3 is one, two or three substituents independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy,

20 C_{1-6} alkylthio, $-OCF_3$, $-OCH_2(C_{2-6})$ alkenyl, $-NH_2$, $-NH(C_{1-6})$ alkyl, $-N(C_{1-6})$ dialkyl, $-NHC(=O)Cy$, $-N(C_{1-6})alkylC(=O)Cy$, $-(NC(=O))_2NH_2$, $-C(=O)C_{1-4}$ alkoxy, $-C(=O)NR^{17}R^{18}$, $-C(=O)NH$ cycloalkyl,

$-C(=O)N(C_{1-6})alkyl$ cycloalkyl, $-C(=O)NHCy$, $-C(=O)N(C_{1-6})alkylCy$, $-C(=O)Cy$, $-OC(=O)C_{1-6}$ alkyl, $-OC(=O)NR^{19}R^{20}$, $-C(=O)O$ aryl,

25 $-C(=O)O$ heteroaryl, $-CO_2H$, ureido, halogen, hydroxy, nitro, cyano, aryl, heteroaryl, heteroaryloxy and aryloxy,

wherein any of the foregoing C_{1-6} alkyl or C_{1-6} alkoxy containing substituents of R^3 are optionally substituted with one, two or three substituents

30 independently selected from the group consisting of $-NR^{21}R^{22}$, $-NH$ (cycloalkyl), $-N(C_{1-6})alkyl$ (cycloalkyl), $-NHCy$, $-N(C_{1-6})alkylCy$, $-NHC(O)-C_{1-6}alkyl-C_{1-6}alkoxy$, aryl, heteroaryl, hydroxy, halogen, $-C(=O)NR^{23}R^{24}$, $-OC(=O)NR^{25}R^{26}$, $-C(=O)C_{1-4}$ alkoxy and $-C(=O)Cy$,

wherein any of the foregoing C₂₋₆alkenyl and C₂₋₆alkynyl containing substituents of R³ are optionally substituted with aryl or -C(=O)NR²⁷R²⁸, and

wherein the aryl, heteroaryl and cycloalkyl substituents of R³ are optionally substituted with one, two or three substituents independently selected from R¹⁴;

R¹⁴ is independently hydrogen, C₁₋₆alkyl, C₁₋₆alkoxy, C₂₋₆alkenyl, C₁₋₆alkylthio, -NH₂, -NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, aryl, heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy, or nitro;

wherein each of the C₁₋₆alkyl- or C₁₋₆alkoxy-containing substituents of R¹⁴ is optionally substituted on a terminal carbon atom with a substituent selected from -NR²⁹R³⁰, aryl, heteroaryl, one, two or three halogen atoms, or hydroxy;

R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are substituents independently selected from the group consisting of hydrogen, C₁₋₆alkyl and aryl,

wherein C₁₋₆alkyl and aryl are each optionally substituted with hydroxy, aryl, aryloxy, -C(=O)-aryl, -C(=O)C₁₋₄alkoxy, NH₂, -NH(C₁₋₆alkyl), or -N(C₁₋₆)dialkyl; optionally, R¹⁷ and R¹⁸, R¹⁹ and R²⁰, R²¹ and R²², R²³ and R²⁴, or R²⁵ and R²⁶ are each taken together with the atoms to which they are attached to form a ring of five to seven members;

R²⁷ and R²⁸ are independently hydrogen; C₁₋₆alkyl optionally substituted with hydroxy, aryl, -C(=O)C₁₋₄alkoxy, NH₂, -NH(C₁₋₆alkyl), or -N(C₁₋₆)dialkyl; or aryl; optionally, R²⁷ and R²⁸ are each taken together with the atoms to which they are attached to form a ring of five to seven members;

R²⁹ and R³⁰ are independently hydrogen, C₁₋₆alkyl or aryl, wherein C₁₋₆alkyl is optionally substituted with hydroxy, aryl, -C(=O)C₁₋₄alkoxy, NH₂, -NH(C₁₋₆alkyl), or -N(C₁₋₆)dialkyl, and, optionally, R²⁹ and R³⁰ are each taken together with the atoms to which they are attached to form a ring of five to seven members;

Cy is a heterocyclyl optionally substituted with a substituent selected from the group consisting of oxo, C₁₋₆alkyl, -C₁₋₆alkylC(=O)C₁₋₆alkyl, -C₁₋₆alkylC(=O)C₁₋₆alkoxy, -C₁₋₆alkyl-aryl, -C₁₋₆alkylC(=O)aryl,

-C(=O)(C₁₋₆)alkyl, -C(=O)(C₁₋₆)alkoxy, -C(=O)aryl, -SO₂aryl, aryl, heteroaryl and heterocyclyl,

wherein the aryl portion of any aryl-containing substituent of Cy is optionally substituted with one, two or three substituents independently selected from the group consisting of C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkylthio, halogen, hydroxy, NH₂, -NH(C₁₋₆alkyl) and -N(C₁₋₆)dialkyl, and wherein heterocyclyl is optionally substituted with aryl, one, two or three halogen atoms, or one, two or three oxo substituents; and heterocyclyl is optionally spiro-fused to said Cy;

n is 0 or 1;

W is O or S;

X is hydrogen or C₁₋₃alkyl;

Y is independently selected from the group consisting of C₁₋₆alkyl substituted with -OSO₂NH₂ or hydroxy; SO₃H, CO₂H, heteroaryl, -OC(=O)NH₂ and P(=O)OR⁵R⁶; provided that when Y is CO₂H, Ring A must be a bicyclic ring system;

R⁵ is selected from the group consisting of hydrogen, C₁₋₆alkyl and aryl, wherein C₁₋₆alkyl is optionally substituted with NH₂, -NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, 1,3-dioxolan-2-yl, C₁₋₆alkylcarbonyloxy-, C₁₋₆alkoxycarbonyloxy-, C₁₋₆alkylcarbonylthio-, (C₁₋₆)alkylaminocarbonyl-, di(C₁₋₆)alkylaminocarbonyl-, one, two or three halogen atoms, or hydroxy, and

wherein aryl is optionally substituted with C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkylthio, C₂₋₆alkenyl, -NH₂, -NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, aryl, heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy, or nitro, and, optionally, when R⁶ is C₁₋₈alkoxy, R⁵ and R⁶ are each taken together with the atoms to which they are attached to form a 5-8 membered monocyclic ring;

R⁶ is selected from the group consisting of C₁₋₈alkyl, C₁₋₈alkoxy, C₂₋₈alkenyl, heteroaryl, aryl and hydroxy,

wherein C₁₋₈alkyl, C₁₋₈alkoxy and C₂₋₈alkenyl are optionally substituted with a substituent selected from the group consisting of C₁₋₆alkoxy, aryl,

heterocyclyl, heteroaryl, NH_2 , $-\text{NH}(\text{C}_{1-6})\text{alkyl}$, $-\text{N}(\text{C}_{1-6})\text{dialkyl}$,
 $\text{C}_{1-6}\text{alkylcarbonyloxy-}$, $\text{C}_{1-6}\text{alkylcarbonylthio-}$, $\text{C}_{1-6}\text{alkoxycarbonyloxy-}$,
 $(\text{C}_{1-6})\text{alkylaminocarbonyl-}$, $\text{di}(\text{C}_{1-6})\text{alkylaminocarbonyl-}$, one, two or three
 halogen atoms and hydroxy,

- 5 wherein when R^6 is $\text{C}_{1-8}\text{alkyl}$, said $\text{C}_{1-8}\text{alkyl}$ is optionally substituted with
 halogen selected from up to three chlorine atoms or up to seven fluorine
 atoms, and

wherein the heteroaryl and aryl substituents of R^6 are optionally substituted
 with a substituent independently selected from the group consisting of
 10 $\text{C}_{1-6}\text{alkyl}$, $\text{C}_{1-6}\text{alkoxy}$, $\text{C}_{2-6}\text{alkenyl}$, $\text{C}_{1-6}\text{alkylthio}$, $-\text{NH}_2$, $-\text{NH}(\text{C}_{1-6})\text{alkyl}$,
 $-\text{N}(\text{C}_{1-6})\text{dialkyl}$, aryl, heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy
 and nitro;

R^4 is one, two or three substituents selected from the group consisting of
 hydrogen, $\text{C}_{1-6}\text{alkyl}$, $\text{C}_{2-6}\text{alkenyl}$, $\text{C}_{1-6}\text{alkoxy}$, $\text{C}_{1-6}\text{alkylthio}$, $\text{aryl}(\text{C}_{1-6})\text{alkyl}$,
 15 $\text{aryl}(\text{C}_{2-6})\text{alkenyl}$, halogen, hydroxy, $-\text{C}(=\text{O})\text{Cy}$, $-\text{C}(=\text{O})\text{NR}^{31}\text{R}^{32}$, aryl,
 $-\text{CO}_2\text{H}$, oxo and cyano,

wherein $\text{C}_{1-6}\text{alkyl}$, $\text{C}_{2-6}\text{alkenyl}$ and $\text{C}_{1-6}\text{alkoxy}$ are each optionally substituted
 with $-\text{NR}^{33}\text{R}^{34}$, aryl, heteroaryl, cycloalkyl, one, two or three halogen
 atoms, or hydroxy, and

- 20 wherein aryl and heteroaryl are each optionally substituted with a substituent
 independently selected from the group consisting of $\text{C}_{1-6}\text{alkyl}$, $\text{C}_{1-6}\text{alkoxy}$,
 $\text{C}_{2-6}\text{alkenyl}$, $\text{C}_{1-6}\text{alkylthio}$, $-\text{NH}_2$, $-\text{NH}(\text{C}_{1-6})\text{alkyl}$, $-\text{N}(\text{C}_{1-6})\text{dialkyl}$, aryl,
 heteroaryl, aryloxy, heteroaryloxy, one, two or three halogen atoms,
 hydroxy and nitro;

- 25 R^{31} , R^{32} , R^{33} and R^{34} are substituents independently selected from the group
 consisting of hydrogen, $\text{C}_{1-6}\text{alkyl}$ and aryl,

wherein $\text{C}_{1-6}\text{alkyl}$ is optionally substituted with hydroxy, aryl, $-\text{C}(=\text{O})\text{C}_{1-4}\text{alkoxy}$,
 NH_2 , $-\text{NH}(\text{C}_{1-6}\text{alkyl})$, or $-\text{N}(\text{C}_{1-6})\text{dialkyl}$, and, optionally, R^{31} and R^{32} or R^{33}
 and R^{34} are each taken together with the atoms to which they are
 30 attached to form a ring of five to seven members;

and enantiomers, diastereomers, polymorphs or pharmaceutically acceptable

salts thereof.

Illustrative of the invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and any of the compounds described above.

5 An illustration of the invention is a pharmaceutical composition made by mixing any of the compounds described above and a pharmaceutically acceptable carrier.

 Illustrating the invention is a process for making a pharmaceutical composition comprising mixing any of the compounds described above and a
10 pharmaceutically acceptable carrier.

 The present invention is also directed to methods for producing the instant compounds and pharmaceutical compositions and medicaments thereof.

 The present invention is further directed to methods for treating or
15 ameliorating an inflammatory or serine protease-mediated disorder. In particular, the method of the present invention is directed to treating or ameliorating an inflammatory or serine protease (such as chymase) mediated disorder such as, but not limited to, allergic rhinitis, viral rhinitis, asthma, chronic obstructive pulmonary diseases, bronchitis, pulmonary emphysema,
20 acute lung injury, psoriasis, arthritis, reperfusion injury, ischemia, hypertension, hypercardia myocardial infarction, heart failure damage associated with myocardial infarction and cardiac hypertrophy, arteriosclerosis, saroidosis, vascular stenosis or restenosis (e.g., associated with vascular injury, angioplasty, vascular stents or vascular grafts), pulmonary fibrosis, kidney
25 fibrosis (e.g., associated with glomerulonephritis), liver fibrosis, post surgical adhesion formation, systemic sclerosis, keloid scars, rheumatoid arthritis, bullous pemphigoid and atherosclerosis. Additionally, these compounds can be used for modulating wound healing and remodeling (e.g., cardiac hypertrophy) as well as immune modulation.

30 An embodiment of the present invention includes a method for treating or ameliorating an inflammatory or serine protease mediated disorder selected

from the group consisting of allergic rhinitis, asthma and heart failure damage associated with myocardial infarction and cardiac hypertrophy.


An embodiment of the present invention includes compounds of Formula (I) wherein R^1 is selected from the group consisting of hydrogen and C_{1-4} alkyl.

5 Examples of the present invention include compounds of Formula (I) wherein R^1 is hydrogen.

An embodiment of the present invention includes compounds of Formula (I) wherein Ring A is selected from the group consisting of aryl, heteroaryl, benzo fused heterocyclyl and benzo fused cycloalkyl.

10 An embodiment of the present invention includes compounds of Formula (I) wherein Ring A is selected from the group consisting of heteroaryl, benzo fused heterocyclyl, or aryl.

An embodiment of the present invention includes compounds of Formula (I) wherein Ring A is a bicyclic Ring a^1a^2 of the formula:



15

wherein the a^1 portion of said Ring a^1a^2 is optionally substituted with R^2 ; and the a^2 portion is optionally substituted with R^3 .

An embodiment of the present invention includes compounds of Formula (I) wherein the a^2 portion of Ring a^1a^2 is an aromatic ring.

20 An embodiment of the present invention includes compounds of Formula (I) wherein Ring A is selected from the group consisting of naphthyl, benzothiazolyl, benzothiophenyl, quinolinyl, isoquinolinyl, dihydronaphthyl, indanyl, tetrazolynyl and benzodioxolyl when n is equal to zero; and Ring A is phenyl, pyridin-2-yl, or pyridin-3-yl when n is equal to one.

25 In embodiments of the present invention wherein, for compounds of Formula (I), a bicyclic ring system is used for Ring A, the a^2 ring will be aromatic. For example, Ring A is selected from the group consisting of naphthyl, benzothiazolyl and benzothiophenyl, when n is equal to zero, and

Ring A is selected from phenyl, pyridin-2-yl and pyridin-3-yl when n is equal to one.

Embodiments of the present invention include compounds of Formula (I) wherein n is 0.

5 Examples of the present invention includes compounds of Formula (I) wherein Ring A is naphthyl and n is 0.

Embodiments of the present invention include compounds of Formula (I) wherein R^2 is one, two or three substituents independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{1-6} alkoxy, $-NH_2$, $-NH(C_{1-6}alkyl)$,
 10 $-N(C_{1-6})dialkyl$, aryl, heteroaryl, halogen, hydroxy and nitro, and wherein C_{1-6} alkyl and C_{1-6} alkoxy are optionally substituted with a substituent selected from $-NR^{11}R^{12}$, aryl, heteroaryl, one, two or three halogen atoms and hydroxy.

Embodiments of the present invention include compounds of Formula (I)
 15 wherein R^2 is a substituent independently selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{1-4} alkoxy, hydroxy, halogen and $-NH_2$.

Embodiments of the present invention include compounds of Formula (I) wherein R^2 is hydrogen, C_{1-4} alkyl, halogen, or $-NH_2$.

Examples of the present invention include compounds of Formula (I)
 20 wherein R^2 is hydrogen.

Embodiments of the present invention include compounds of Formula (I) wherein R^3 is one, two or three substituents independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} alkoxy,
 $-OCH_2(C_{2-6})alkenyl$, NH_2 , $-NH(C_{1-6}alkyl)$, $-N(C_{1-6})dialkyl$, $-NHC(=O)Cy$,
 25 $-N(C_{1-6}alkyl)C(=O)Cy$, $-C(=O)C_{1-4}alkoxy$, $-C(=O)NR^{17}R^{18}$,
 $-C(=O)NHcycloalkyl$, $-C(=O)N(C_{1-6}alkyl)cycloalkyl$, $-C(=O)NHCy$,
 $-C(=O)N(C_{1-6}alkyl)Cy$, $-C(=O)Cy$, $-OC(=O)NR^{19}R^{20}$, halogen, hydroxy, nitro, cyano, aryl and aryloxy,

wherein C_{1-6} alkyl and C_{1-6} alkoxy are optionally substituted with one, two or
 30 three substituents independently selected from the group consisting of $-NR^{21}R^{22}$, $-NHcycloalkyl$, $-N(C_{1-6}alkyl)cycloalkyl$, $-NHCy$, $-N(C_{1-6}alkyl)Cy$,

aryl, heteroaryl, halogen, $-C(=O)NR^{23}R^{24}$, $-OC(=O)NR^{25}R^{26}$,
 $-C(=O)(C_{1-4})alkoxy$ and $-C(=O)Cy$,

wherein $C_{2-6}alkenyl$ is optionally substituted on a terminal carbon with aryl or
 $-C(=O)NR^{27}R^{28}$, and

- 5 wherein aryl and cycloalkyl are optionally substituted with one, two or three
 substituents independently selected from R^{14} .

Embodiments of the present invention include compounds of Formula (I)

- wherein R^3 is one, two or three substituents independently selected from the
 group consisting of hydrogen, $C_{1-6}alkyl$, $C_{1-6}alkoxy$, $-NR^{19}R^{20}$,
 10 $-NHC(=O)Cy$, $-C(=O)NR^{17}R^{18}$, $-C(=O)NHcycloalkyl$,
 $-C(=O)N(C_{1-6}alkyl)cycloalkyl$, halogen and aryl,

wherein $C_{1-6}alkyl$ and $C_{1-6}alkoxy$ are optionally substituted on a terminal carbon
 atom with one, two or three fluorine atoms, $-NH_2$, $-NHCy$, or
 $-N(C_{1-4}alkyl)Cy$, and

- 15 wherein aryl and cycloalkyl are optionally substituted with a group
 independently selected from R^{14} .

Embodiments of the present invention include compounds of Formula (I)

- wherein R^3 is one, two or three substituents independently selected from
 hydrogen, trifluoromethyl, $C_{1-4}alkoxy$ optionally substituted with one, two or
 20 three fluorine atoms, $-NH_2$, $-NHC(=O)Cy$, or halogen.

Embodiments of the present invention include compounds of Formula (I)

- wherein when R^3 is $NHC(=O)Cy$, then Cy is preferably piperidinyl optionally
 substituted with a substituent selected from the group consisting of
 $C_{1-4}alkyl$, $C_{1-4}alkylC(=O)C_{1-4}alkyl$, $-C_{1-4}alkylC(=O)C_{1-4}alkoxy$,
 25 $C_{1-4}alkylC(=O)aryl$, $-C(=O)(C_{1-4})alkyl$, $-C(=O)(C_{1-4})alkoxy$, $-C(=O)aryl$,
 $-SO_2aryl$, aryl, heteroaryl and heterocyclyl,

- wherein aryl and the aryl portion of the $C_{1-4}alkylC(=O)aryl$, $-C(=O)aryl$ and
 $-SO_2aryl$ are each optionally substituted with one, two or three
 substituents independently selected from the group consisting of
 30 $C_{1-4}alkyl$, $C_{1-4}alkoxy$, halogen, hydroxy, NH_2 , $-NH(C_{1-6}alkyl)$ and
 $-N(C_{1-4})dialkyl$, and

wherein heterocyclyl is optionally substituted with aryl, one, two or three halogen atoms, or one oxo substituent.

Embodiments of the present invention include compounds of Formula (I) wherein R³ is hydrogen, trifluoromethyl, one or two fluorine atoms, chloro,
5 methoxy, trifluoromethoxy, or NH₂.

Examples of the present invention include compounds of Formula (I) wherein R³ is hydrogen.

Examples of the present invention include compounds of Formula (I) wherein W is O.

10 Embodiments of the present invention include compounds of Formula (I) wherein X is C₁₋₃alkyl.

Examples of the present invention include compounds of Formula (I) wherein X is hydrogen.

Examples of the present invention include compounds of Formula (I)
15 wherein R¹, R², R³ and X are each hydrogen and W is O.

Embodiments of the present invention include compounds of Formula (I) wherein Y is independently selected from a group consisting of C₁₋₃alkyl, SO₃H, CO₂H, heteroaryl, -OC(=O)NH₂ and P(=O)OR⁵R⁶, and
wherein C₁₋₃alkyl is substituted with a substituent selected from the group
20 consisting of -OSO₂NH₂ and hydroxy.

Examples of the present invention include compounds of Formula (I) wherein Y is independently SO₃H or P(=O)OR⁵R⁶.

Examples of the present invention include compounds of Formula (I) wherein Y is P(=O)OR⁵R⁶.

25 Embodiments of the present invention include compounds of Formula (I) wherein R⁵ is selected from the group consisting of hydrogen; C₁₋₆alkyl and aryl,

wherein C₁₋₆alkyl is optionally substituted with NH₂, -NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, C₁₋₆alkylcarbonyloxy-, C₁₋₆alkoxycarbonyloxy-,
30 C₁₋₆alkylcarbonylthio-, (C₁₋₆)alkylaminocarbonyl-,

di(C₁₋₆)alkylamino-carbonyl-, one, two or three halogen atoms, or hydroxy, and

wherein aryl is optionally substituted with C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkylthio, C₂₋₆alkenyl, -NH₂, -NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, aryl, heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy, or nitro; optionally, when R⁶ is C₁₋₈alkoxy, R⁵ and R⁶ are each taken together with the atoms to which they are attached to form a 5-8 membered monocyclic ring.

Embodiments of the present invention include compounds of Formula (I)

wherein R⁵ is selected from the group consisting of hydrogen, C₁₋₆alkyl and aryl,

wherein C₁₋₆alkyl is optionally substituted with C₁₋₆alkylcarbonyloxy-, C₁₋₆alkoxycarbonyloxy-, C₁₋₆alkyl-carbonylthio-, (C₁₋₆)alkylaminocarbonyl-, di(C₁₋₆)alkylaminocarbonyl-, one, two or three halogen atoms, or hydroxy and, optionally, when R⁶ is C₁₋₈alkoxy, R⁵ and R⁶ are each taken together with the atoms to which they are attached to form a 6-7 membered monocyclic ring.

Embodiments of the present invention include compounds of Formula (I)

wherein R⁵ is hydrogen or C₁₋₆alkyl optionally substituted with C₁₋₆alkylcarbonyloxy-, C₁₋₆alkoxycarbonyloxy-, C₁₋₆alkylcarbonylthio-, (C₁₋₆)alkylaminocarbonyl-, or di(C₁₋₆)alkylaminocarbonyl-, and, optionally, when R⁶ is C₁₋₈alkoxy, R⁵ and R⁶ are each taken together with the atoms to which they are attached to form a 6-membered monocyclic ring.

Examples of the present invention include compounds of Formula (I)

wherein R⁵ is hydrogen or C₁₋₆alkyl.

Examples of the present invention include compounds of Formula (I)

wherein R⁵ is hydrogen or methyl.

Embodiments of the present invention include compounds of Formula (I)

wherein R⁶ is selected from the group consisting of C₁₋₈alkyl, C₁₋₈alkoxy, C₂₋₈alkenyl, heteroaryl, aryl and hydroxy,

wherein C₁₋₈alkyl, C₁₋₈alkoxy and C₂₋₈alkenyl are each optionally substituted on a terminal carbon atom with a substituent independently selected from

the group consisting of C₁₋₄alkoxy, aryl, heteroaryl, heterocyclyl,
 C₁₋₆alkylcarbonyloxy-, C₁₋₆alkylcarbonylthio-, C₁₋₆alkoxycarbonyloxy-,
 (C₁₋₆)alkylaminocarbonyl-, di(C₁₋₆)alkyl-aminocarbonyl- and hydroxy, and
 wherein heteroaryl and aryl are optionally substituted with one, two or three
 5 substituents independently selected from the group consisting of aryl,
 hydroxy, C₁₋₆alkoxy and halogen.

Embodiments of the present invention include compounds of Formula (I)
 wherein R⁶ is selected from the group consisting of C₁₋₆alkyl, C₁₋₆alkoxy,
 heteroaryl, aryl and hydroxy,
 10 wherein C₁₋₆alkyl is optionally substituted on a terminal carbon atom with a
 substituent selected from C₁₋₃alkoxy, aryl, or hydroxy; and
 wherein C₁₋₆alkoxy is optionally substituted on a terminal carbon with a
 substituent independently selected from the group consisting of
 C₁₋₆alkylcarbonyloxy- and di(C₁₋₆)alkylaminocarbonyl-, and
 15 wherein heteroaryl and aryl are optionally substituted with one, two or three
 substituents independently selected from the group consisting of aryl,
 hydroxy, C₁₋₆alkoxy and halogen.

Examples of the present invention include compounds of Formula (I)
 wherein R⁶ is selected from the group consisting of C₁₋₆alkyl and hydroxy.
 20 Embodiments of the present invention include compounds of Formula (I)
 wherein R⁶ is selected from the group consisting of methyl, ethyl,
 methoxypropyl, phenethyl, benzo[1,3]dioxol-5-yl-propyl, hydroxy and C₁₋₃alkoxy
 optionally substituted on C₁₋₃alkoxy with C₁₋₆alkylcarbonyloxy- and
 di(C₁₋₆)alkylaminocarbonyl-.

25 Examples of the present invention include compounds of Formula (I)
 wherein R⁶ is selected from the group consisting of methyl and hydroxy.

Embodiments of the present invention include compounds of Formula (I)
 wherein R⁴ is one, two or three substituents selected from the group consisting
 of hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, C₁₋₆alkoxy, aryl(C₂₋₆)alkenyl, halogen,
 30 hydroxy, -C(=O)Cy, -C(=O)NR³¹R³², aryl, -CO₂H, oxo and cyano,
 wherein C₁₋₆alkyl and C₁₋₆alkoxy are optionally substituted on a terminal carbon

atom with a substituent selected from aryl, $-NR^{33}R^{34}$, one, two or three halogen atoms, or hydroxy, and

wherein aryl is optionally substituted with one, two or three substituents independently selected from the group consisting of hydrogen, C₁₋₆alkyl, C₁₋₆alkoxy, C₂₋₆alkenyl, -NH₂, -NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, aryl, heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy and nitro.

Embodiments of the present invention include compounds of Formula (I) wherein R⁴ is one, two or three substituents selected from the group consisting of hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, C₁₋₆alkoxy, aryl(C₂₋₆)alkenyl, halogen, hydroxy, -C(=O)Cy, -C(=O)NR³¹R³², aryl, -CO₂H, oxo and cyano,

wherein C₁₋₆alkyl and C₁₋₆alkoxy are optionally substituted with a substituent independently selected from $-NR^{33}R^{34}$, aryl, one, two or three halogen atoms, or hydroxy, and

wherein aryl is optionally substituted with a substituent independently selected from the group consisting of hydrogen, C₁₋₆alkyl, C₁₋₆alkoxy, aryl, halogen, hydroxy and nitro.

Embodiments of the present invention include compounds of Formula (I) wherein R⁴ is one, two or three substituents selected from the group consisting of hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, C₁₋₆alkoxy, aryl(C₂₋₆)alkenyl, halogen, hydroxy and -C(=O)Cy,


wherein aryl is optionally substituted with a substituent selected from halogen and C₁₋₄alkoxy.

Examples of the present invention include compounds of Formula (I) wherein R⁴ is one, two or three substituents selected from the group consisting of hydrogen, C₁₋₆alkyl, C₁₋₆alkoxy, halogen and hydroxy.

Embodiments of the present invention include compounds wherein R⁴ is one or two or three substituents selected from the group consisting of hydrogen, fluorine, chlorine, bromine, hydroxy, methyl and methoxy.

Examples of the present invention include compounds of Formula (I) wherein R⁴ is one or two or three substituents selected from the group consisting of hydrogen, chlorine, bromine, hydroxy, methyl and methoxy.

The present invention is further directed to a compound of Formula (Ia):



wherein

Ring A is aryl;

5 n is 0 or 1;

R⁵ is selected from the group consisting of hydrogen and C₁₋₆alkyl;

R⁶ is selected from the group consisting of C₁₋₈alkyl and hydroxy;

R⁴ is one, two or three substituents selected from the group consisting of hydrogen, C₁₋₆alkyl, C₁₋₆alkoxy, halogen and hydroxy;

10 and enantiomers, diastereomers, polymorphs or pharmaceutically acceptable salts thereof.

Examples of the present invention include compounds of Formula (Ia) wherein Ring A is naphthyl and n is 0.


15 Examples of the present invention include compounds of Formula (Ia) wherein R⁵ is hydrogen or methyl.

Examples of the present invention include compounds of Formula (Ia) wherein R⁶ is selected from the group consisting of methyl and hydroxy.


20 Examples of the present invention include compounds of Formula (Ia) wherein R⁴ is one or two or three substituents selected from the group consisting of hydrogen, chlorine, bromine, hydroxy, methyl and methoxy.

Embodiments of the phosphonic and phosphinic acids of the present invention include those compounds of Formula (Ia) wherein the R⁵ and R⁶ substituents are as previously defined for Formula (I) (which further include substitutions in any combination).


Embodiments of the present invention include a compound or an enantiomer, diastereomer, polymorph or pharmaceutically acceptable salt thereof selected from:




Cpd 1




Cpd 2




Cpd 3




Cpd 4




Cpd 5




Cpd 6




Cpd 7




Cpd 8




Cpd 9




Cpd 10




Cpd 11




Cpd 12




Cpd 13




Cpd 14




Cpd 15




Cpd 16




Cpd 17




Cpd 18




Cpd 19




Cpd 20




Cpd 21



Cpd 22



Cpd 23



Cpd 24

A representative compound of Formula (I) or an enantiomer, diastereomer, polymorph or pharmaceutically acceptable salt thereof includes a compound selected from the group consisting of:

Cpd	Name
1	[benzofuran-3-yl-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
2	[(5-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
3	[(5-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
4	[(6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
5	[(5,7-dibromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
6	[(5,7-dibromo-6-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
7	[(5,7-dibromo-6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,

Cpd	Name
8	[(5-chloro-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
9	[(4-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
10	[(5-bromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
11	[(7-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
12	[(5-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
13	[benzofuran-3-yl-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
14	methyl-[(5-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphinic acid,
15	[(5-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
16	[(6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
17	[(5,7-dibromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
18	[(5,7-dibromo-6-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
19	[(5,7-dibromo-6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
20	[(5-chloro-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
21	[(4-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
22	[(5-bromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
23	[(7-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid, and
24	[(5-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid.

Compound Forms

The compounds of the invention may be present in the form of pharmaceutically acceptable salts. For use in medicines, the “pharmaceutically acceptable salts” of the compounds of this invention refer to non-toxic

5 acidic/anionic or basic/cationic salt forms.

Suitable salt forms include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of an acid such as acetic acid, adipic acid, benzoic acid, carbonic acid, citric acid, fumaric acid, glycolic acid, hydrochloric acid, maleic acid, malonic acid, phosphoric acid, saccharinic acid, succinic acid, sulphuric acid, tartaric acid, trifluoroacetic acid and the like.

Furthermore when the compounds of the present invention carry an acidic moiety, suitable salts thereof may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with suitable organic ligands, e.g. quaternary ammonium salts.

Thus, representative salts include the following: acetate, adipate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium, camsylate (or camphosulphonate), carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, fumarate, gluconate, glutamate, glyconate, hydrabamine, hydrobromine, hydrochloride, iodide, isothionate, lactate, malate, maleate, malonate, mandelate, mesylate, nitrate, oleate, pamoate, palmitate, phosphate/diphosphate, saccharinate, salicylate, stearate, sulfate, succinate, tartrate, tosylate, trichloroacetate, trifluoroacetate and the like.

The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds, which are readily convertible *in vivo* into an active compound.

Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various disorders described with the compound specifically disclosed or a prodrug compound which would be obviously included within the scope of the invention although not specifically disclosed. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985. Phosphonic acid prodrugs (as described in De Lombaert S., et al, Non-Peptidic Inhibitors of Neutral Endopeptidase 24.11; Design and Pharmacology of Orally Active

Phosphonate Prodrugs, *Bioorganic and Medicinal Chemistry Letters*, **1995**, 5(2), 151-154; and, De Lombaert S., et al, *N*-Phosphonomethyl Dipeptides and Their Phosphonate Prodrugs, a New Generation Neutral Endopeptidase (NEP, EC 3.424.11) Inhibitors, *J. Med. Chem.*, **1994**, 37, 498-511) and phosphinic
5 acid prodrugs are intended to be included within the scope of the present invention.

The compounds according to this invention may have at least one chiral center and thus may exist as enantiomers. In addition, the compounds of the present invention may also possess two or more chiral centers and thus may
10 also exist as diastereomers. Where the processes for the preparation of the present compounds give rise to a mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. Accordingly, the compounds may be prepared as a racemic mixture or, by either enantiospecific synthesis or resolution, as individual enantiomers. The
15 compounds may, for example, be resolved from a racemic mixture into their component racemates by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active base, followed by fractional crystallization and regeneration of the compounds of this invention. The racemic mixture may also be resolved by formation of diastereomeric
20 esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention.

During any of the processes for preparation of the compounds of the
25 present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John
30 Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known in the art.

Furthermore, some of the crystalline forms for the compounds may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents and as such are also intended to be encompassed within the scope of this invention.

Chemical Definitions

As used herein, unless otherwise noted, "alkyl" whether used alone or as part of a substituent group refers to straight and branched carbon chains having 1 to 8 carbon atoms or any number within this range. Examples include methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, tertiary butyl, 1-pentyl, 2-pentyl, 3-pentyl, 1-hexyl, 2-hexyl, 3-hexyl and the like.

The term "alkoxy" refers to an -O-alkyl substituent group, wherein alkyl is as defined *supra*. Examples include methoxy, ethoxy, propoxy and the like.

The term "alkylthio" refers to an -S-alkyl substituent group, wherein the alkyl portion may be further substituted where allowed by available valences.

The terms "alkenyl" and "alkynyl" refer to straight and branched carbon chains having 2 to 8 carbon atoms or any number within this range, wherein an alkenyl chain has at least one double bond in the chain and an alkynyl chain has at least one triple bond in the chain. Examples for alkenyl include ethenyl (also referred to as vinyl), *iso*-propenyl, allyl (also referred to as propenyl), propylidene and the like. Examples for alkynyl include ethynyl, propynyl and the like. Alkenyl and alkynyl radicals may be attached to a core molecule and further substituted on any atom when allowed by available valences.

An alkyl and alkoxy chain may be substituted on a terminal carbon atom or, when acting as a linking group, within the carbon chain where allowed by available valences.

The term "alkylthio" refers to a -S-alkyl substituent group, wherein the alkyl portion may be further substituted where allowed by available valences.

The term "C₁₋₆alkylcarbonyloxy" refers to a -O-C(O)-C₁₋₆alkyl substituent group, wherein the alkyl portion may be further substituted where allowed by available valences.

The term "alkoxycarbonyl" refers to a $-C(O)-O-C_{1-6}$ alkyl substituent group, wherein the alkyl portion may be further substituted where allowed by available valences.

5 The term " C_{1-6} alkoxycarbonyloxy" refers to a $-O-C(O)-O-C_{1-6}$ alkyl substituent group, wherein the alkyl portion may be further substituted where allowed by available valences.

The term " C_{1-6} alkylcarbonylthio" refers to a $-S-C(O)-C_{1-6}$ alkyl substituent group, wherein the alkyl portion may be further substituted where allowed by available valences.

10 The term " (C_{1-6}) alkylaminocarbonyl" refers to a $-C(O)-NH-C_{1-6}$ alkyl or $-C(O)-NH(C_{1-6})$ alkyl substituent group, wherein the amino or alkyl portion may be further substituted where allowed by available valences.

The term " $di(C_{1-6})$ alkylaminocarbonyl" refers to a $-C(O)-N(C_{1-6}alkyl)_2$ or $-C(O)-N(C_{1-6})$ dialkyl substituent group, wherein the alkyl portion may be further substituted where allowed by available valences.

15

The term "cycloalkyl" refers to saturated or partially unsaturated, monocyclic or polycyclic hydrocarbon rings of from 3 to 20 carbon atom members (preferably from 3 to 14 carbon atom members). Further, a cycloalkyl ring may optionally be fused to one or more cycloalkyl rings. Examples of such rings include, and are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and adamantanyl.

20

The term "heterocyclyl" refers to a nonaromatic cyclic ring of 5 to 10 members in which 1 to 4 members are nitrogen or a nonaromatic cyclic ring of 5 to 10 members in which zero, one or two members are nitrogen and up to two members is oxygen or sulfur; wherein, optionally, the ring contains zero, one or two unsaturated bonds. Alternatively, the heterocyclyl ring may be fused to a benzene ring (benzo fused heterocyclyl), a 5 or 6 membered heteroaryl ring (containing one of O, S or N and, optionally, one additional nitrogen), a 5 to 7 membered cycloalkyl or cycloalkenyl ring, a 5 to 7 membered heterocyclyl ring (of the same definition as above but absent the option of a further fused ring) or fused with the carbon of attachment of a cycloalkyl, cycloalkenyl or heterocyclyl ring to form a spiro moiety (spiro-fused). For instant compounds of the

25

30

invention, the carbon atom ring members that form the heterocyclyl ring are fully saturated. Other compounds of the invention may have a partially saturated heterocyclyl ring. Additionally, the heterocyclyl can be bridged to form bicyclic rings. Preferred partially saturated heterocyclyl rings may have from one to two double bonds. Such compounds are not considered to be fully aromatic and are not referred to as heteroaryl compounds. Examples of heterocyclyl groups include, and are not limited to, pyrrolinyl (including 2*H*-pyrrole, 2-pyrrolinyl or 3-pyrrolinyl), pyrrolidinyl, 2-imidazolinyl, imidazolidinyl, 2-pyrazolinyl, pyrazolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, and piperazinyl.

The term "heterocyclylcarbonyl" refers to a -C(O)-heterocyclyl substituent group, wherein the heterocyclyl portion may be further substituted where allowed by available valences.

The term "aryl" refers to an unsaturated, aromatic monocyclic ring of 6 carbon members or to an unsaturated, aromatic polycyclic ring of from 10 to 20 carbon members. Examples of such aryl rings include, and are not limited to, phenyl, naphthalenyl and anthracenyl.

The term "benzo fused cycloalkyl" refers to a bicyclic or tricyclic ring structure wherein at least one of the ring substituents is phenyl or naphthalenyl and at least one of the other substituents is a cycloalkyl ring (as cycloalkyl was previously defined). For the purpose of these definitions, the cycloalkyl rings may be fused to an additional benzene ring (to provide fused multiple ring systems such as fluorene). Example of such benzo fused cycloalkyls include, but are not limited to, indanyl, 1,2,3,4-tetrahydronaphthalenyl and fluorenyl.

The term "heteroaryl" refers to an aromatic ring of 5 or 6 members wherein the ring consists of carbon atoms and has at least one heteroatom member. Suitable heteroatoms include nitrogen, oxygen or sulfur. In the case of 5 membered rings, the heteroaryl ring contains one member of nitrogen, oxygen or sulfur and, in addition, may contain up to three additional nitrogens. In the case of 6 membered rings, the heteroaryl ring may contain from one, two or three nitrogen atoms. For the case wherein the 6 membered ring has three nitrogens, at most two nitrogen atoms are adjacent. Optionally, the heteroaryl

ring is fused to a benzene ring (benzo fused heteroaryl), a 5 or 6 membered heteroaryl ring (containing one of O, S or N and, optionally, one additional nitrogen), a 5 to 7 membered cycloalkyl ring or a 5 to 7 membered heterocyclo ring (as defined supra but absent the option of a further fused ring). Examples of heteroaryl groups include, and are not limited to, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl or pyrazinyl; fused heteroaryl groups include indolyl, isoindolyl, indolinyl, benzofuryl, benzothienyl, indazolyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, benzisoxazolyl, benzothiadiazolyl, benzotriazolyl, quinoliziny, quinolinyl, isoquinolinyl, and quinazolinyl.

The term "aryl(C₁₋₆)alkyl" means an C₁₋₆alkyl group substituted with an aryl group (e.g., benzyl and phenethyl), as in the formula - C₁₋₆alkyl-aryl, wherein aryl may be substituted on any alkyl chain carbon atom where allowed by available valences.

The term "aryl(C₂₋₆)alkenyl" means an C₂₋₆alkenyl group substituted with an aryl group, as in the formula - C₂₋₆alkenyl-aryl, wherein aryl may be substituted on any alkenyl chain carbon atom where allowed by available valences.

The term "arylalkoxy" indicates an alkoxy group substituted with an aryl group (e.g., benzyloxy), as in the formula -O-alkyl-aryl, wherein aryl may be substituted on any alkyl chain carbon atom where allowed by available valences.


The term "aryloxy" indicates an oxy group substituted with an aryl group (e.g., benzoxy), as in the formula -O-aryl.

The term "heteroaryloxy" indicates an alkoxy group substituted with a heteroaryl group, as in the formula -O-alkyl-heteroaryl, wherein heteroaryl may be substituted on any alkyl chain carbon atom where allowed by available valences..

The term "halogen" refers to fluorine, chlorine, bromine, and iodine. Substituents that are substituted with multiple halogen atoms are substituted in a manner that provides compounds which are stable (e.g. -CF₃ or -OCF₃).

Whenever the term “alkyl” or “aryl” or either of their prefix roots appear in a name of a substituent (e.g., arylalkyl and alkylamino), it shall be interpreted as including those limitations given above for “alkyl” and “aryl.” Designated numbers of carbon atoms (e.g., C₁-C₆) shall refer independently to the number of carbon atoms in an alkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root. For alkyl, and alkoxy substituents the designated number of carbon atoms includes all of the independent member included in the range specified individually and all the combination of ranges within in the range specified.

In general, under standard nomenclature rules used throughout this disclosure, the terminal portion of the designated side chain is described first followed by the adjacent functionality toward the point of attachment. Thus, for example, a “phenylC₁-C₆ alkylamidoC₁-C₆alkyl” substituent refers to a group of the formula:



It is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of this invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth herein.

Pharmaceutical Compositions and Methods of Use

Illustrative of the invention is a composition comprising a pharmaceutically acceptable carrier and any of the compounds described above. Also illustrative of the invention is a composition made by mixing any of the compounds described above and a pharmaceutically acceptable carrier. A further illustration of the invention is a process for making a composition comprising mixing any of the compounds described above and a pharmaceutically acceptable carrier. The present invention also provides

compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier.

The compounds of the present invention are useful serine protease inhibitors (in particular, inhibitors of chymase) useful for treating inflammatory or serine protease mediated disorders. Serine proteases such as chymase produced by mast cells have been recognized to be involved in a variety of inflammatory and wound healing events (e.g., angiogenesis, collagen deposition and cell proliferation). Chymase plays these roles by activating a variety of pre-existing factors present in the microenvironment surrounding the mast cells. For example, chymase activates SCF, angiotensin I to angiotensin II, endothelin 1, type 1 procollagen, metalloproteinases, IL-1B, TGF- β , as well as, degrades the extracellular matrix (de Paulis *et al.* Int Arch Allerg Immunol 118 (1999) 422-425; Longley *et al.* Proc Natl Acad Sci USA 94 (1997) 9017-9021). Consequently, the release of chymase plays significant role in a variety of pathological conditions associated with vascular proliferation, fibrosis, tissue remodeling, inflammation, and the like.

Some of these, inflammatory or serine protease mediated disorders include, and are not limited to, allergic rhinitis, viral rhinitis, asthma, chronic obstructive pulmonary diseases, bronchitis, pulmonary emphysema, acute lung injury (e.g. adult (acute) respiratory distress syndrome), psoriasis, arthritis, reperfusion injury, ischemia, hypertension, hypercardia myocardial infarction, heart failure damage associated with myocardial infarction and cardiac hypertrophy, arteriosclerosis, saroidosis, vascular stenosis or restenosis (e.g., associated with vascular injury, angioplasty, vascular stents or vascular grafts), pulmonary fibrosis, kidney fibrosis (e.g., associated with glomerulonephritis), liver fibrosis, post surgical adhesion formation, systemic sclerosis, keloid scars rheumatoid arthritis, bullous pemphigoid and atherosclerosis.

Additionally, these compounds can be used for modulating wound healing and remodeling (e.g., cardiac hypertrophy) as well as immune modulation. The utility of the compounds to treat inflammatory or serine protease mediated disorders is illustrated by the following non-limiting discussions of the proposed mechanisms of actions of chymase. Other

disorders that can be treated with chymase inhibitors can be determined according to the procedures described herein and the use of animal knock-out models and the like.

As mentioned above, chymase converts angiotensin I into angiotensin II, and this activity has been associated with vascular proliferation. In human vascular extracts only about 8% of angiotensin II activity is inhibited with an angiotensin converting enzyme inhibitor (lisinopril) while 95% is inhibited by a chymase inhibitor. In vein grafts, vascular injury associated with catheter or balloon injury, chymase induces vascular hyperplasia and restenosis in dogs (Takai and Miyazaki, 21 (2003) 185-189). This same mechanism of action would also be expected to apply to restenosis associated with the use of vascular stents. Pathological serine protease mediated disorders associated with angiotensin II, including but not limited to hypertension, hypercardia myocardial infarction, arteriosclerosis, saroidosis, vascular stenosis or restenosis (e.g., associated with vascular injury, angioplasty, vascular stents or vascular grafts), and the like.

Pathological fibrosis can be associated with the degeneration of organs (e.g., skin, heart, kidneys or liver) or as an undesirable outcome of surgery. Preventing the formation of pathological fibrosis would be beneficial in a variety of diseases. For example, mast cell chymase has been implicated in pulmonary fibrosis, kidney fibrosis, liver fibrosis, post surgical adhesion formation, systemic sclerosis, keloid scars, and the like.

In the heart, mast cells have been implicated in cardiac hypertrophy, which involves both fibrosis and remodeling. Cardiac hypertrophy develops to preserve its function by normalizing chamber wall stress. Mast cells have been implicated as being involved in the development of myocardial fibrosis and systolic pressure over load induced hypertrophy (Hara *et al.*, J. Exp. Med. 195 (2002) 375-381). The remodeling of the heart associated under these conditions is believed to involve mast cell chymase, which activates endothelin 1, matrix metalloproteinases and TGF- β . Chymase inhibitors have been shown to exert favorable cardioprotective action in a dog model of hypertrophy (Matsumoto *et al.*, Circulation 107 (2003) 2555-2558).

In the kidneys, mast cell chymase has also been implicated in pathological fibrosis. For example, glomerulonephritis has also been reported to involve mast cells (Ehara and Shigematsu, *Kidney Inter.* 54 (1998) 1675-1683). The results of this found that mast cells were one of the constitutive cell types in the interstitium of IgA nephritis patients and contributed to interstitial fibrosis resulting in deterioration of renal function. Similarly, liver fibrosis has been associated with mast cells (Yamashiro *et al.*, *Virchows Arch.* 433 (1998) 471-479). Although, the mechanisms for fibrosis in the kidney and liver have not been as well defined as for coronary fibrosis, it is very likely that chymase is operating through similar signaling pathways to cause fibrosis (especially in liver fibrosis where fibrosis seem to be occurring more frequently where mast cells stained positive for chymase).

Chymase is also involved in the formation of fibrous adhesions associated with surgery. Chymase inhibitors have been tested in two different animals models and found to reduce the number of adhesions (Okamoto *et al.*, *J. Surg. Res.* 107 (2002) 219-222 and Lucas *et al.*, *J. Surg. Res.* 65 (1999) 135). It has been suggested that the prevention of adhesions is associated with blocking the activation of latent TGF- β by chymase (Yoa *et al.*, *J. Surg. Res.* 92 (2000) 40-44).

Collagen induced arthritic mice show increased numbers of mast cells and expression of chymase in fibroproliferative inflammation (Kakizoe *et al.*, *Inflamm. Res.* 48 (1999) 318-324). In human rheumatoid arthritis increased mast cell density in the superficial synovium is associated with the severity of the disease (Grotis-Graham and McNeil, *Arthritis & Rheumatism* 40 (1997) 479-489). It was theorized by these authors that chymase and its ability to activate metalloproteinases plays a significant role in the rapid functional deterioration observed in rheumatoid arthritis.

Mast cell chymase has been implicated in atherosclerosis via its ability to cleave apolipoprotein B-100 of LDL which facilitates lipoprotein aggregation and uptake by macrophages (Paananen *et al.*, *J. Biol. Chem.* 269 (1994) 2023-2031). Chymase also degrades apolipoprotein A of HDL, which would reduce cholesterol efflux and increases lipid deposition (Lindstedt *et al.*, *J. Clin. Invest.*

97 (1996) 2174-2182). Thus, chymase is involved in two different pathways to atherosclerosis.

An embodiment of the invention is a method for treating inflammatory or serine protease mediated disorders in a subject in need thereof which
5 comprises administering to the subject a therapeutically effective amount of any of the compounds or compositions described above.

An example of the invention is a method for treating an inflammatory or chymase mediated disorder selected from the group consisting of allergic rhinitis, viral rhinitis, asthma, chronic obstructive pulmonary diseases,
10 bronchitis, pulmonary emphysema, psoriasis, arthritis, reperfusion injury, ischemia, hypertension, hypercardia myocardial infarction, heart failure damage associated with myocardial infarction and cardiac hypertrophy, arteriosclerosis, sarcoidosis, vascular stenosis or restenosis, pulmonary fibrosis, kidney fibrosis, liver fibrosis, post surgical adhesion formation, systemic sclerosis, keloid scars,
15 rheumatoid arthritis, bullous pemphigoid and atherosclerosis in a subject in need thereof comprising administering to the subject a therapeutically effective amount of any of the compounds of Formula (I) or Formula (Ia) or compositions thereof as described above.

An example of the invention is a use of a compound of Formula (I) or
20 Formula (Ia) for the preparation of a medicament for treating an inflammatory or serine protease mediated disorder in a subject in need thereof.

An embodiment of a serine protease inhibited according to the use of a compound in a medicament is chymase.

An example of the use of a compound in a medicament for treating
25 inflammatory or chymase mediated disorders is a disorder selected from the group consisting of allergic rhinitis, viral rhinitis, asthma, chronic obstructive pulmonary diseases, bronchitis, pulmonary emphysema, psoriasis, arthritis, reperfusion injury, ischemia, hypertension, hypercardia myocardial infarction, heart failure damage associated with myocardial infarction and cardiac
30 hypertrophy, arteriosclerosis, sarcoidosis, vascular stenosis or restenosis, pulmonary fibrosis, kidney fibrosis, liver fibrosis, post surgical adhesion

formation, systemic sclerosis, keloid scars, rheumatoid arthritis, bullous pemphigoid and atherosclerosis.

An example of the use of a compound in a medicament for treating an inflammatory or serine protease mediated disorder is a disorder selected from the group consisting of allergic rhinitis, asthma, chronic obstructive pulmonary disease, bronchitis, pulmonary emphysema, acute lung injury, heart failure damage associated with myocardial infarction and cardiac hypertrophy.

An example of the use of a compound in a medicament for treating an inflammatory or serine protease mediated disorder is a disorder selected from the group consisting of allergic rhinitis, asthma and heart failure damage associated with myocardial infarction and cardiac hypertrophy.

The term "treating" as used herein refers to a method for improving, halting, retarding or palliating an inflammatory or serine protease mediated disorder in the subject in need thereof. All such methods of treatment are intended to be within the scope of the present invention.

In accordance with the methods of the present invention, the individual components of the compositions described herein can also be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted accordingly.

The term "subject" as used herein, refers to an animal (preferably, a mammal; most preferably, a human), having been the object of treatment, observation, or experiment.

The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human, that is being sought by a researcher, veterinarian, medical doctor, or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well

as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

To prepare the compositions of this invention, one or more compounds of Formula (I) or salt thereof as the active ingredient, is intimately admixed with
5 a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration (e.g. oral or parenteral). Suitable pharmaceutically acceptable carriers are well known in the art. Descriptions of some of these pharmaceutically acceptable carriers
10 may be found in The Handbook of Pharmaceutical Excipients, published by the American Pharmaceutical Association and the Pharmaceutical Society of Great Britain.

Methods of formulating compositions have been described in numerous publications such as Pharmaceutical Dosage Forms: Tablets, Second Edition,
15 Revised and Expanded, Volumes 1-3, edited by Lieberman et al; Pharmaceutical Dosage Forms: Parenteral Medications, Volumes 1-2, edited by Avis et al; and Pharmaceutical Dosage Forms: Disperse Systems, Volumes 1-2, edited by Lieberman et al; published by Marcel Dekker, Inc.

In preparing a composition of the present invention in liquid dosage form
20 for oral, topical, inhalation/insufflation and parenteral administration, any of the usual pharmaceutical media or excipients may be employed. Thus, for liquid dosage forms, such as suspensions (i.e., colloids, emulsions and dispersions) and solutions, suitable carriers and additives include but are not limited to pharmaceutically acceptable wetting agents, dispersants, flocculation agents,
25 thickeners, pH control agents (i.e., buffers), osmotic agents, coloring agents, flavors, fragrances, preservatives (i.e., to control microbial growth, etc.) and a liquid vehicle may be employed. Not all of the components listed above will be required for each liquid dosage form.

In solid oral preparations such as, for example, powders, granules,
30 capsules, caplets, gelcaps, pills and tablets (each including immediate release, timed release and sustained release formulations), suitable carriers and additives include but are not limited to diluents, granulating agents, lubricants,

binders, glidants, disintegrating agents, and the like. Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated, gelatin coated, film coated
5 or enteric coated by standard techniques.

Preferably these compositions are in unit dosage forms from such as tablets, pills, capsules, powders, granules, lozenges, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories for administration by oral, intranasal,
10 sublingual, intraocular, transdermal, parenteral, rectal, vaginal, inhalation or insufflation means. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection.

15 For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g., conventional tableting ingredients such as diluents, binders, adhesives, disintegrants, lubricants, antiadherents, and glidants.

Suitable diluents include, but are not limited to, starch (i.e., corn, wheat,
20 or potato starch, which may be hydrolized), lactose (granulated, spray dried or anhydrous), sucrose, sucrose-based diluents (confectioner's sugar; sucrose plus about 7 to 10 weight percent invert sugar; sucrose plus about 3 weight percent modified dextrans; sucrose plus invert sugar, about 4 weight percent invert sugar, about 0.1 to 0.2 weight percent cornstarch and magnesium
25 stearate), dextrose, inositol, mannitol, sorbitol, microcrystalline cellulose, dicalcium phosphate, calcium sulfate dihydrate, calcium lactate trihydrate, and the like.

Suitable binders and adhesives include, but are not limited to accacia gum, guar gum, tragacanth gum, sucrose, gelatin, glucose, starch, and
30 cellulosics (i.e. methylcellulose, sodium carboxymethy-cellulose, ethylcellulose, hydroxypropylmethylcellulose, hydroxypropylcellulose, and the like), water soluble or dispersible binders (i.e., alginic acid and salts thereof, magnesium

aluminum silicate, hydroxyethylcellulose, polyethylene glycol, polysaccharide acids, bentonites, polyvinylpyrrolidone, polymethacrylates and pregelatinized starch), and the like.

Suitable disintegrants include, but are not limited to, starches (corn, potato, etc.), sodium starch glycolates, pregelatinized starches, clays (magnesium aluminum silicate), celluloses (such as crosslinked sodium carboxymethylcellulose and microcrystalline cellulose), alginates, pregelatinized starches (i.e., corn starch, etc.), gums (i.e., agar, guar, locust bean, karaya, pectin, and tragacanth gum), cross-linked polyvinylpyrrolidone, and the like.

Suitable lubricants and antiadherents include, but are not limited to, stearates (magnesium, calcium and sodium), stearic acid, talc waxes, stearowet, boric acid, sodium chloride, DL-leucine, carbowax 4000, carbowax 6000, sodium oleate, sodium benzoate, sodium acetate, sodium lauryl sulfate, magnesium lauryl sulfate, and the like. Suitable gildants include, but are not limited to, talc, cornstarch, silica, and the like. Sweeteners and flavorants may be added to chewable solid dosage forms to improve the palatability of the oral dosage form. Additionally, colorants and coatings may be added or applied to the solid dosage form for ease of identification of the drug or for aesthetic purposes. These carriers are formulated with the pharmaceutical active to provide an accurate, appropriate dose of the pharmaceutical active with a therapeutic release profile.

Generally these carriers are mixed with the pharmaceutical active to form a solid preformulation composition containing a homogeneous mixture of the pharmaceutical active of the present invention, or a pharmaceutically acceptable salt thereof. Generally the preformulation will be formed by one of three common methods: (a) wet granulation, (b) dry granulation, and (c) dry blending. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of

the type described above containing from about 0.01 mg to about 500 mg of the active ingredient of the present invention.

The tablets or pills containing the novel compositions may also be formulated in multilayer tablets or pills to provide a sustained or provide dual-
5 release products. For example, a dual release tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer, which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in
10 release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric materials such as shellac, cellulose acetate, cellulose acetate phthalate, polyvinyl acetate phthalate, hydroxypropyl methylcellulose phthalate, hydroxy-propyl methylcellulose acetate succinate, methacrylate and ethylacrylate copolymers, and the like.
15 Sustained release tablets may also be made by film coating or wet granulation using slightly soluble or insoluble substances in solution (which for a wet granulation acts as the binding agents) or low melting solids a molten form (which in a wet granulation may incorporate the active ingredient). These materials include natural and synthetic polymers waxes, hydrogenated oils,
20 fatty acids and alcohols (i.e., beeswax, carnauba wax, cetyl alcohol, cetylstearyl alcohol, and the like), esters of fatty acids metallic soaps, and other acceptable materials that can be used to granulate, coat, entrap or otherwise limit the solubility of an active ingredient to achieve a prolonged or sustained release product.

25 The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, but are not limited to aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar
30 pharmaceutical vehicles.

Suitable suspending agents for aqueous suspensions, include synthetic and natural gums such as, acacia, agar, alginate (i.e., propylene alginate,

sodium alginate, and the like), guar, karaya, locust bean, pectin, tragacanth, and xanthan gum, cellulosics such as sodium carboxymethylcellulose, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose and hydroxypropyl methylcellulose, and combinations thereof,
5 synthetic polymers such as polyvinyl pyrrolidone, carbomer (i.e., carboxypolymethylene), and polyethylene glycol; clays such as bentonite, hectorite, attapulgite or sepiolite; and other pharmaceutically acceptable suspending agents such as lecithin, gelatin, or the like.

Suitable surfactants include but are not limited to sodium docusate,
10 sodium lauryl sulfate, polysorbate, octoxynol-9, nonoxynol-10, polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 80, polyoxamer 188, polyoxamer 235, and combinations thereof. Suitable deflocculating or dispersing agent include pharmaceutical grade lecithins. Suitable flocculating agent include but are not limited to simple neutral electrolytes (i.e., sodium chloride, potassium,
15 chloride, and the like), highly charged insoluble polymers and polyelectrolyte species, water soluble divalent or trivalent ions (i.e., calcium salts, alums or sulfates, citrates and phosphates (which can be used jointly in formulations as pH buffers and flocculating agents).

Suitable preservatives include but are not limited to parabens (i.e.,
20 methyl, ethyl, *n*-propyl and *n*-butyl), sorbic acid, thimerosal, quaternary ammonium salts, benzyl alcohol, benzoic acid, chlorhexidine gluconate, phenylethanol, and the like. There are many liquid vehicles that may be used in liquid pharmaceutical dosage forms, however, the liquid vehicle that is used in a particular dosage form must be compatible with the suspending agent(s).

25 For example, nonpolar liquid vehicles such as fatty esters and oils liquid vehicles are best used with suspending agents such as low HLB (Hydrophile-Lipophile Balance) surfactants, stearylalkonium hectorite, water insoluble resins, water insoluble film forming polymers, and the like. Conversely, polar liquids such as water, alcohols, polyols and glycols are best used with suspending
30 agents such as higher HLB surfactants, clays silicates, gums, water soluble cellulosics, water soluble polymers, and the like. For parenteral administration, sterile suspensions and solutions are desired. Liquid forms useful for parenteral

administration include sterile solutions, emulsions and suspensions. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

Furthermore, compounds of the present invention can be administered in an intranasal dosage form via topical use of suitable intranasal vehicles or via transdermal skin patches, the composition of which are well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the administration of a therapeutic dose will, of course, be continuous rather than intermittent throughout the dosage regimen.

Compounds of the present invention can also be administered in a form suitable for intranasal or inhalation therapy. For such therapy, compounds of the present invention are conveniently delivered in the form of a solution or suspension from a pump spray container that is squeezed or pumped or as an aerosol spray from a pressurized container or a nebulizer (such as, a metered dose inhaler, a dry powder inhaler or other conventional or non-conventional modes or devices for inhalation delivery) using a suitable propellant (such as, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas). In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurized container or nebulizer may contain a solution or suspension of the active compound. Capsules and cartridges (such as, those made from gelatin) for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

Compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, multilamellar vesicles and the like. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, phosphatidylcholines, and the like.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with

soluble polymers as targetable drug carriers. Such polymers can include, but are not limited to polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxy-ethylaspartamidephenol, and polyethyl eneoxidepolylysine substituted with palmitoyl residue. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, to homopolymers and copolymers (which means polymers containing two or more chemically distinguishable repeating units) of lactide (which includes lactic acid d-, l- and meso lactide), glycolide (including glycolic acid), ϵ -caprolactone, p-dioxanone (1,4-dioxan-2-one), trimethylene carbonate (1,3-dioxan-2-one), alkyl derivatives of trimethylene carbonate, δ -valerolactone, β -butyrolactone, γ -butyrolactone, ϵ -decalactone, hydroxybutyrate, hydroxyvalerate, 1,4-dioxepan-2-one (including its dimer 1,5,8,12-tetraoxacyclotetradecane-7,14-dione), 1,5-dioxepan-2-one, 6,6-dimethyl-1,4-dioxan-2-one, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates, and cross-linked or amphipathic block copolymers of hydrogels and blends thereof.

The therapeutically effective amount of a compound or composition thereof may be from about 0.001 mg/kg/dose to about 300 mg/kg/dose. Preferably, the therapeutically effective amount may be from about 0.001 mg/kg/dose to about 100 mg/kg/dose. More preferably, the therapeutically effective amount may be from about 0.001 mg/kg/dose to about 50 mg/kg/dose. Most preferably, the therapeutically effective amount may be from about 0.001 mg/kg/dose to about 30 mg/kg/dose. Therefore, the therapeutically effective amount of the active ingredient contained per dosage unit (e.g., tablet, capsule, powder, injection, suppository, teaspoonful, and the like) as described herein will be in the range of from about 1 mg/day to about 21,000 mg/day for a subject, for example, having an average weight of 70 kg. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the subject to be treated.

Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, and the advancement of the disease condition. In addition, factors associated with the particular subject being treated, including subject age, weight, diet and time of administration, will result in the need to adjust the dose to an appropriate therapeutic level. Advantageously, compounds of the present invention may be administered in a single daily dose or the total daily dosage may be administered in divided doses of two, three or four times daily.

Abbreviations used in the instant specification, particularly the Schemes and Examples, are as follows:

Abbreviation	Meaning
Boc	<i>tert</i> -butoxycarbonyl
BOC-ON	2-(<i>tert</i> -butoxycarbonyloxyimino)-2-phenylacetonitrile
BuLi	<i>n</i> -butyllithium
t-BuOH	<i>tert</i> -butanol
Cpd	compound
LiOH	lithium hydroxide
min/h/d	minutes/hour/hours/day/days
rt/RT	room temperature
TEA or Et ₃ N	triethylamine
THF	tetrahydrofuran
TFA	trifluoroacetic acid
TMSBr	bromotrimethylsilane

General Synthetic Methods

Representative compounds of the present invention can be synthesized in accordance with the general synthetic methods described below and are illustrated more particularly in the schemes that follow. Since the schemes are an illustration, the invention should not be construed as being limited by the chemical reactions and conditions expressed. The preparation of the various starting materials used in the schemes is well within the skill of persons versed in the art.

The following schemes describe general synthetic methods whereby intermediate and target compounds of the present invention may be prepared. Additional representative compounds and stereoisomers, racemic mixtures, diastereomers and enantiomers thereof can be synthesized using the


5 intermediates prepared in accordance to the general schemes and other materials, compounds and reagents known to those skilled in the art. All such compounds, stereoisomers, racemic mixtures, diastereomers and enantiomers thereof are intended to be encompassed within the scope of the present invention. Since the scheme is an illustration, the invention should not be

10 construed as being limited by the chemical reactions and conditions expressed. The preparation of the various starting materials used in the scheme is well within the skill of persons versed in the art.

SCHEME A

Scheme A illustrates the general method for the preparation of

15 compounds representative of the present invention.




A phosphonate or phosphinate anion (prepared from its corresponding phosphonate or phosphinate Compound **A2** and an organometallic base such as *n*-butyllithium) is reacted with an isocyanate Compound **A1** in a solvent such as THF to afford an amidophosphonate or amidophosphinate compound **A3**.

20 One versed in the art will recognize that conventional chemical transformations may be utilized to prepare certain R^2 and R^3 substituents of the present invention. For example, for the preparation of a compound wherein R^3 is amino, a nitro group may be reduced with hydrazine hydrate in the presence of

25 a palladium catalyst; or, for the preparation of a compound wherein R^3 is ureido, a compound in which R^3 is an amino group may be reacted with a cyanate salt or the like.

Compound **A2**, wherein R^5 and R^6 are as previously defined, may be made according to known methods (Katritsky *et. al. Org. Prep. Proced. Int.*, 1990, 22(2), 209-213; *J. Am. Chem. Soc.*, 2002, 124, 9386-9387; and *Chem. Ber.*, 1963, 96, 3184-3194), may be prepared from a commercially available or
 5 known haloalkyl substituted heteroaryl ring.


Fluorinated R^6 compounds can be made following methods known in the art such as the methods similar those set forth in Garabadzhia *et al.*, Journal General Chemistry USSR, English translation, 1981, pages 1905-1910.



10 Compound **A3** may be dealkylated with bromotrimethylsilane in a solvent such as pyridine, followed by treatment with dilute HCl to afford Compound **A4**.

SCHEME B

Scheme B shows a method for preparing Compound **B3** wherein R^6 is
 15 an alkyl or alkenyl substituent using methods described in the literature (*J. Organomet. Chem.* 2002, 643-644, 154-163; *J. Amer. Chem. Soc.* 2002, 124, 9386-9387).




Compound **B2** (wherein R^4 is other than hydrogen) is known; certain
 20 analogs of which may be commercially available and may be alternatively prepared according to *J. Med. Chem* 1997, 40 (17) 2706-2725. Another method for preparing such compounds (wherein R^4 is other than hydrogen) is


described in the literature (*Med. Chem.* **1995**, 38(17), 3297-3312; *Bioorg. Med. Chem.* **1999**, 7, 2697-2704).

Compound **B2** (wherein R^4 is hydrogen) is prepared according to a procedure described in *J. Med. Chem.*, **1997**, 40 (17) 2706-2725.


5




Compound **B4** is reacted with a solution of ethyl propiolate (in the presence of a solvent such as TEA and THF) to provide a Compound **B5**.




Compound **B5** is reacted in the presence of a catalyst (such as $Pd(OAc)_2$ and PPH_3) and a basic solution (such as $NaHCO_3$ in DMF) to provide a Compound **B6**.



Compound **B6** is reacted with a reagent solution (such as DIBAL-H in a solvent such as CH_2Cl_2) to provide a non-isolated intermediate which is then reacted in the presence of a catalyst solution (such as PPh_3Br_2 in a solvent such as CCl_4) to provide Compound **B2**.




Compound **B2** is reacted with a triethylphosphite solution (in a solvent such as dry toluene) to provide a Compound **B7** that is carried forward according to the procedure of Scheme A to provide compounds representative
5 of the present invention.




Compound **B2** is reacted with a methyl-diethylphosphite solution (in a solvent such as dry toluene) to provide a Compound **B8** that is carried forward according to the procedure of Scheme A to provide compounds representative
10 of the present invention.

SCHEME C


Scheme C illustrates a general method for the preparation of compounds representative of the present invention wherein Ring A of Formula (I) is an aryl substituent and n of Formula (I) is equal to 1.



Reaction of an α/β -unsaturated carboxylic acid, Compound **C1**, with phosphorazidic acid dialkyl ester Compound **C2** provides Compound **C3**. Compound **C3** may subsequently undergo a Curtius rearrangement to afford an isocyanate intermediate, Compound **C4**.



Compound **C4** is reacted with a phosphonate or phosphinate anion (using Compound **C2** from Scheme A or Compound **B3** from Scheme B) in a mixture of an aprotic solvent such as THF and an organometallic base (such as *n*-butyl-lithium) to yield amidophosphonate or amidophosphinate Compound **C5**.




Compound **C5** may be dealkylated with bromotrimethylsilane, followed by treatment with dilute HCl to afford Compound **C6**.

10


SCHEME D

Scheme D further illustrates the preparation of compounds representative of the present invention wherein Y of Formula (I) is a heteroaryl substituent.



Compound **D1** (where R^4 is hydrogen) is known and commercially available.


Compound **D1** may be dissolved in an aprotic solvent, treated with an organometallic base such as *n*-BuLi, and subsequently reacted with isocyanate Compound **A1** to afford Compound **D2**.



Compound **D2** may undergo a cycloaddition reaction with sodium azide to provide Compound **D3**.

SCHEME E


5 Scheme E shows the preparation of compounds representative of the present invention wherein Y of Formula (I) is a sulfonic acid.




Compound **B2** may be treated with sodium sulfite to afford Compound **E1**. Compound **E1** may then be treated with an organometallic base such as isopropylmagnesium bromide and reacted with isocyanate Compound **A1** to yield Compound **E2**.

SCHEME F


Scheme F illustrates the preparation of compounds representative of the present invention wherein Y of Formula (I) is a carboxylic acid.



Compound **F1** may be reacted with isobutylene under acidic conditions to provide ester Compound **F3**.



Compound **F3** may then be treated with a strong base such as lithium diethylamide and further reacted with isocyanate Compound **A1** to give Compound **F4**.




5

Compound **F4** is converted into its corresponding carboxylic acid Compound **F5** by treatment with TFA.


SCHEME G

Scheme G illustrates the preparation of compounds representative of the present invention wherein Y of Formula (I) is hydroxymethyl.

10




Nitrile Compound **D2** may be converted to an imidate in the presence of HCl gas, followed by hydrolysis to yield Compound **G1**.



Compound **G1** may be reduced to a primary alcohol in the presence of hydride source, such as sodium borohydride, to give methyl alcohol Compound **G2**.

SCHEME H

5 Scheme H illustrates the preparation of compounds representative of the present invention wherein Y of Formula (I) is a sulfamic acid methyl group.




Compound **G2** may be treated with a base such as sodium hydride, followed by the addition of sulfamoyl chloride to yield Compound **H1**.

10


SCHEME I

Scheme I illustrates the general method for the preparation of compounds representative of the present invention wherein R^3 is an amide substituent on ring A as defined by the invention.




15


Dinitro-substituted Compound **I1** may be reduced by hydrogenation in the presence of a palladium catalyst to give Compound **I2** which then may be acylated with BOC-ON to provide Compound **I3**.




Compound **I4** may be acylated with acid chloride Compound **I5** to yield Compound **I6**, followed by saponification of Compound **I6** (using a reagent such as LiOH) to provide carboxylic acid Compound **I7**.




- 5 Compound **I8** may be prepared by coupling Compound **I3** to Compound **I7** using an appropriate coupling agent, activating agent, and solvent. The Boc protecting group of Compound **I8** was removed under acidic conditions to afford the free amine, Compound **I9**.



- 10 Treatment of Compound **A2** with an organometallic base such as *n*-butyllithium, followed by reaction with carbon dioxide afforded the carboxylated phosphonic ester, Compound **I10**.



Compound **I10** was converted to its acid chloride by treatment with thionyl chloride followed by condensation with amine Compound **I9** to afford amide Compound **I11**.




5


Compound **I11** was dealkylated using bromotrimethylsilane and treated with HCl to provide Compound **I12**.

SCHEME J


Scheme J illustrates a method for the preparation of compounds
10 representative of the present invention.




A Compound **J1**, wherein R^3 is an alkoxycarbonyl substituent, may be reduced in the presence of a hydride source to the corresponding alcohol, Compound **J2**.




5 Compound **J2** may be oxidized to aldehyde Compound **J3**.




Reaction of Compound **J3** with a Wittig reagent affords alkene Compound **J4**.



10 Saponification of Compound **J4** provides carboxylic acid Compound **J5**.



Compound **J5** may be coupled with an amine, such as benzyl amine, in the presence of an appropriate coupling agent as described supra, to give amide Compound **J6**.




5 Compound **J6** may be dealkylated using the procedure previously described in Scheme A to yield Compound **J7**.

Alternatively, other compounds of the present invention wherein R^3 is alkoxy or $-C(=O)NR^{11}R^{12}$ may be derived from Compound **J2**. The hydroxy group of Compound **J2** may be alkylated using reagents and methods known to
10 one skilled in the art to afford compounds wherein R^3 is alkoxy.


Alternatively, the hydroxy group of Compound **J2** may be reacted with a variety of acylating agents known to one skilled in the art, such as isocyanates, to arrive at compounds of the present invention wherein R^3 is a carbamate.

SCHEME K

15 Scheme K illustrates a method for the preparation of compounds representative of the present invention.




As shown in Scheme K, Compound **J3** may be reacted with a variety of amines in the presence of a hydride source under acidic conditions to yield
20 Compound **K1**.




Dealkylation of Compound **K1** by the method described in Scheme A affords Compound **K2**.

SCHEME L


- 5 The preparation of compounds representative of the present invention wherein R^3 is $-C(=O)Cy$, as previously defined, and said Cy is attached through a nitrogen atom, is shown in Scheme L.




- Compound **J1** may be saponified under basic conditions to provide
10 Compound **L1** (wherein Ring A is phenyl or naphthyl).



Compound **L1** may be treated with thionyl chloride to give Compound **L2**.




Compound **L2** may be reacted with a substituted amine (wherein Cy is as previously defined) to provide Compound **L3**.



Dealkylation of Compound **L3** using methods previously described
5 affords Compound **L4**.

SCHEME M

Scheme M illustrates a method for the preparation of compounds representative of the present invention wherein R^5 and R^6 are appropriately substituted alkoxy substituents as defined herein.



10


A compound of formula **M1** wherein R^5 is hydrogen and R^6 is hydroxy may be coupled with an appropriately substituted alcohol in the presence of MSNT (1-(mesitylene-2-sulfonyl)-3-nitro-1,2,4-triazole) to afford a compound of formula **M2** wherein R^5 is a substituted alkyl and R^6 is a substituted alkoxy as
15 defined herein.

Alternatively, compounds of formula **M1** may be elaborated using an appropriately substituted alkylating agent to provide compounds of the present invention where either one or both hydroxy groups of the phosphonic acid are alkylated. An alkylating agent in this instance is an alkyl substituent that is
20 optionally substituted as defined for R^5 or R^6 , and said alkyl substituent is substituted with a leaving group. A leaving group is defined as a substituent


that is activated toward nucleophilic displacement, including halides, tosylates, and the like.

SCHEME N


Scheme N illustrates the preparation of compounds representative of the present invention wherein R^5 and R^6 (when R^6 is alkoxy) are taken together with the atoms to which they are both attached to form a monocyclic ring.



A diol of formula **N1** (wherein s is 0, 1 or 2) may be treated with a dichlorophosphite (wherein R_c is benzyl- or lower alkyl-) to form a cyclic phosphonate of formula **N2**.



A compound of formula **N2** may be condensed under refluxing conditions with a compound of formula **B2** to form a compound of formula **N3**.



The elaboration of a compound of formula **N3** to a compound of formula **N4** may be achieved using the methods described for Scheme A.

Specific Synthetic Examples


The following Examples are set forth to aid in the understanding of the invention, and are not intended and should not be construed to limit in any way the invention set forth in the claims which follow thereafter. The depicted intermediates may also be used in subsequent examples to produce additional compounds of the present invention. No attempt has been made to optimize the yields obtained in any of the reactions. One skilled in the art would know how to increase such yields through routine variations in reaction times, temperatures, solvents and/or reagents.

All chemicals were obtained from commercial suppliers and used without further purification. ^1H and ^{13}C NMR spectra were recorded on a Bruker AC[®] 300B (300 MHz proton) or a Bruker[®] AM-400 (400 MHz proton) spectrometer with Me_4Si as an internal standard (s = singlet, d = doublet, m = multiplet, t = triplet, br = broad). ES-MS were recorded on a Micromass[®] mass spectrometer or on an Agilent[®] HPLC mass spectrometer. TLC was performed with Whatman[®] 250- μm silica gel plates. Preparative TLC was performed with Analtech[®] tapered silica gel GF plates. Preparative HPLC separations were carried out on a Gilson[®] HPLC using a Phenomenex[®] Kromasil 100A C18 column (25 cm x 50 mm, or 10 cm x 21.2 mm) using gradients of CH_3CN /water/ 0.2% TFA; Analytical HPLC separations were carried out on a Supelco[®] ABZ+Plus column (5 cm x 2.1 mm) or a YMC[®] J'Sphere H80 S4 column (5 cm x 2 mm) with detection at 220 nm and 254 nm on a Hewlett Packard[®] 1100 UV detector. The gradient used was 10% to 90% CH_3CN /water/0.1% TFA in 6 min. Reported percent purity data is based on the 220 nm data. Microanalysis was performed by Robertson Microlit Laboratories, Inc.


EXAMPLE 1

[Benzofuran-3-yl-(naphthalen-2-ylcarbamoyl)-methyl]-
phosphonic acid (Cpd 1)


Compound **1d** was prepared according to a procedure described in *J. Med. Chem.*, **1997**, 40 (17) 2706-2725, as follows:




Compound **1a** was reacted with ethyl propiolate in the presence of TEA and THF at room temperature to provide Compound **1b**.




- 5 Compound **1b** was reacted in the presence of $\text{Pd}(\text{OAc})_2$ and PPh_3 and a base NaHCO_3 in DMF at 110°C for 16 h to provide Compound **1c**.




- 10 Compound **1c** was reacted with a solution of DIBAL-H in CH_2Cl_2 at -78°C and allowed to warm to RT to provide an intermediate which was not isolated. The intermediate was reacted in the presence of PPh_3Br_2 and CCl_4 at -0°C to provide Compound **1d**.



To a solution of Compound **1d** (0.29 g; 1.36 mmol) in 10 mL of dry toluene was added triethylphosphite (1.35 mL; 7.8 mmol) and the solution was heated at reflux for a total of 23 h. The reaction was concentrated under high vacuum (<0.5 mTorr) at 90°C. The crude product was purified by flash column
5 (EtOAc/heptane gradient) to yield 0.30 g of Compound **1e** as a clear viscous oil (HPLC: 100% 3.09 min.; MS (ES): $MH^+ = 269$).



To a solution of *n*-butyllithium (1.49 mmol) in 7 mL of dry THF under Ar at -78°C was added a solution of Compound **1e** (0.30 g; 1.10 mmol) in 7 mL of dry THF dropwise. The yellow solution was stirred at -78°C for 45 min, and a
10 mixture of 2-naphthyl isocyanate Compound **1f** (0.20 g; 1.21 mmol) in 7 mL of dry THF was added dropwise. The reaction was allowed to warm slowly in a dry ice/acetone bath for 2 h before being quenched with saturated aqueous NH_4Cl (2 mL). The mixture was warmed to rt, and water was added until two
15 clear layers appeared. The layers were separated, and the aqueous portion was extracted three times with 5 mL of EtOAc. The combined organics were washed once with 10 mL of brine and dried over Na_2SO_4 . The mixture was filtered, the organics were concentrated under reduced pressure, and the crude product was triturated with MeCN to yield Compound **1g** (0.29 g) as a white
20 solid. HPLC: 100% 3.93 min. retention time; MS (ES): $MH^+ = 438$.



To a solution of Compound **1g** (0.29 g; 0.65 mmol) in 5 mL of dry pyridine was added TMS-Br (0.69 mL; 5.24 mmol) in three portions, 15 min apart. After 90 min total reaction time, the reaction mixture was concentrated, and MeOH and 1N HCl were added to give a clear solution. The solution was concentrated, and the residue stirred for 1 h in 1N HCl. The solid was collected and dried in a filter funnel under N₂, then resuspended in MeCN, stirred for 45 min, then collected. A second crop was isolated from the filtrate in the same manner, and the combined product was suspended in a small volume of MeOH and tromethamine (1 equivalent; 63 mg) was added to yield a clear solution. The solution was filtered and concentrated under reduced pressure, and the crude salt was recrystallized from MeCN containing a small amount of EtOAc to yield Compound **1** as a white powder. HPLC: 3.75 min. retention time; MS (ES) MH⁺ = 382; ¹H NMR (DMSO-d₆) δ 4.30 (d, 1H, J = 21.8 Hz), 7.12-7.16 (m, 1H), 7.21-7.26 (m, 1H), 7.32-7.45 (m, 2H), 7.49-7.58 (m, 2H), 7.72-7.80 (m, 2H), 7.98 (d, 1H, J = 2.2 Hz), 8.26 (d, 1H, J = 1.1 Hz), 11.44 (s, 1H).

Using the procedure of Example 1 and appropriate starting materials, reagents and conditions known to those skilled in the art, other compounds or an enantiomer, diastereomer, polymorph or pharmaceutically acceptable salt thereof, which are representative of the present invention may be prepared:

Cpd	Name
2	[(5-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid
3	[(5-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid
4	[(6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid

Cpd	Name
5	[(5,7-dibromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid
6	[(5,7-dibromo-6-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid
7	[(5,7-dibromo-6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid
8	[(5-chloro-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid
9	[(4-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid
10	[(5-bromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid
11	[(7-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid
12	[(5-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid
13	[benzofuran-3-yl-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid
14	methyl-[(5-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphinic acid
15	[(5-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid
16	[(6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid
17	[(5,7-dibromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid
18	[(5,7-dibromo-6-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid
19	[(5,7-dibromo-6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid
20	[(5-chloro-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid
21	[(4-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid
22	[(5-bromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid
23	[(7-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid
24	[(5-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid

Biological Experimental Examples

The utility of the compounds of the present invention as a serine protease inhibitor and, particularly, as a chymase inhibitor useful for the treatment of inflammatory or serine protease mediated disorders can be
5 determined according to the procedures described herein.

EXAMPLE 1

Enzyme-Catalyzed Hydrolysis Assays


Enzyme-catalyzed hydrolysis rates were measured spectrophotometrically using human skin chymase (Cortex Biochem), a chromogenic
10 substrate (Suc-Ala-Ala-Pro-Phe-pNa) (Bachem) in aqueous buffer (450 mM Tris, 1800 mM NaCl, pH 8.0), and a microplate reader (Molecular Devices). IC₅₀ experiments were conducted by fixing the enzyme and substrate concentrations (10 nM enzyme, 0.7 mM substrate) and varying the inhibitor concentration. Changes in absorbance at 405 nM were monitored using the
15 software program Softmax (Molecular Devices), upon addition of enzyme, with and without inhibitor present at 37°C for 30 minutes. Percent inhibition was calculated by comparing the initial reaction slopes of the samples without inhibitor to those with inhibitor.

The IC₅₀ value ($2.54 \pm 0.56 \mu\text{M}$; N = 4) for Compound 1 as a chymase
20 inhibitor was determined using a four parameter fit logistics model.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the
25 following claims and their equivalents.

WHAT IS CLAIMED IS:

1. A compound of Formula (I)



wherein

- 5 R^1 is selected from the group consisting of hydrogen and C_{1-4} alkyl;

Ring A is selected from the group consisting of aryl, heteroaryl, benzo fused heterocyclyl, cycloalkyl and benzo fused cycloalkyl;

- 10 R^2 is one, two or three substituents independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, $-OCH_2-C_{2-6}$ alkenyl, C_{1-6} alkylthio, $-OCF_3$, $-NH_2$, $-NH(C_{1-6})$ alkyl, $-N(C_{1-6})$ dialkyl, aryl, heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy and nitro,

wherein, R^2 is optionally oxo when Ring A is heteroaryl or benzo fused heterocyclyl,

- 15 wherein any aryl-containing substituent of R^2 is optionally substituted with a substituent independently selected from the group consisting of C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyl, C_{1-6} alkylthio, $-NH_2$, $-NH(C_{1-6})$ alkyl, $-N(C_{1-6})$ dialkyl, aryl, heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy and nitro, and
- 20

wherein any of the foregoing C_{1-6} alkyl, C_{2-6} alkenyl or C_{1-6} alkoxy containing substituents of R^2 are optionally substituted with a substituent independently selected from the group consisting of $-NR^{11}R^{12}$, aryl, heteroaryl, one, two or three halogen atoms and hydroxy;

25

R^{11} and R^{12} are independently hydrogen; C_{1-6} alkyl optionally substituted with hydroxy, aryl, $-C(=O)C_{1-4}$ alkoxy, or $-NR^{15}R^{16}$; or aryl;

R^{15} and R^{16} are substituents independently selected from the group consisting of hydrogen, C_{1-6} alkyl and aryl; optionally, R^{15} and R^{16} are each taken together with the atoms to which they are attached to form a ring of five to seven members;

R^3 is one, two or three substituents independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkylthio, $-OCF_3$, $-OCH_2(C_{2-6})$ alkenyl, $-NH_2$, $-NH(C_{1-6})$ alkyl, $-N(C_{1-6})$ dialkyl, $-NHC(=O)Cy$, $-N(C_{1-6})C(=O)Cy$, $-(NC(=O))_2NH_2$, $-C(=O)C_{1-4}$ alkoxy, $-C(=O)NR^{17}R^{18}$, $-C(=O)NH$ cycloalkyl, $-C(=O)N(C_{1-6})$ cycloalkyl, $-C(=O)NHCy$, $-C(=O)N(C_{1-6})$ alkyl, $-C(=O)Cy$, $-OC(=O)C_{1-6}$ alkyl, $-OC(=O)NR^{19}R^{20}$, $-C(=O)O$ aryl, $-C(=O)O$ heteroaryl, $-CO_2H$, ureido, halogen, hydroxy, nitro, cyano, aryl, heteroaryl, heteroaryloxy and aryloxy,

wherein any of the foregoing C_{1-6} alkyl or C_{1-6} alkoxy containing substituents of R^3 are optionally substituted with one, two or three substituents independently selected from the group consisting of $-NR^{21}R^{22}$, $-NH$ (cycloalkyl), $-N(C_{1-6})$ alkyl(cycloalkyl), $-NHCy$, $-N(C_{1-6})$ alkyl, $-NHC(O)-C_{1-6}$ alkyl- C_{1-6} alkoxy, aryl, heteroaryl, hydroxy, halogen, $-C(=O)NR^{23}R^{24}$, $-OC(=O)NR^{25}R^{26}$, $-C(=O)C_{1-4}$ alkoxy, and $-C(=O)Cy$;

wherein any of the foregoing C_{2-6} alkenyl and C_{2-6} alkynyl containing substituents of R^3 are optionally substituted with aryl or $-C(=O)NR^{27}R^{28}$, and

wherein the aryl, heteroaryl and cycloalkyl substituents of R^3 are optionally substituted with one, two or to three substituents independently selected from R^{14} ;

R^{14} is independently hydrogen, C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyl, C_{1-6} alkylthio, $-NH_2$, $-NH(C_{1-6})$ alkyl, $-N(C_{1-6})$ dialkyl, aryl, heteroaryl,

aryloxy, heteroaryloxy, halogen, hydroxy, or nitro;

wherein each of the C₁₋₆alkyl- or C₁₋₆alkoxy-containing

substituents of R¹⁴ is optionally substituted on a terminal

carbon atom with a substituent selected from -NR²⁹R³⁰,

5 aryl, heteroaryl, one, two or three halogen atoms, or

hydroxy;

R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are substituents

independently selected from the group consisting of hydrogen,

C₁₋₆ alkyl, and aryl,

10 wherein C₁₋₆alkyl and aryl are each optionally substituted with hydroxy,

aryl, aryloxy, -C(=O)-aryl, -C(=O)C₁₋₄alkoxy, NH₂, -NH(C₁₋₆alkyl),

or -N(C₁₋₆)dialkyl; optionally, R¹⁷ and R¹⁸, R¹⁹ and R²⁰, R²¹ and

R²², R²³ and R²⁴, or R²⁵ and R²⁶ are each taken together with the

atoms to which they are attached to form a ring of five to seven

15 members;

R²⁷ and R²⁸ are independently hydrogen; C₁₋₆ alkyl optionally substituted

with hydroxy, aryl, -C(=O)C₁₋₄alkoxy, NH₂, -NH(C₁₋₆alkyl), or

-N(C₁₋₆)dialkyl; or aryl; optionally, R²⁷ and R²⁸ are each taken

together with the atoms to which they are attached to form a ring

20 of five to seven members;

R²⁹ and R³⁰ are independently hydrogen, C₁₋₆alkyl or aryl, wherein

C₁₋₆alkyl is optionally substituted with hydroxy, aryl,

-C(=O)C₁₋₄alkoxy, NH₂, -NH(C₁₋₆alkyl), or -N(C₁₋₆)dialkyl, and,

optionally, R²⁹ and R³⁰ are each taken together with the atoms to

25 which they are attached to form a ring of five to seven members;

Cy is a heterocyclyl optionally substituted with a substituent selected

from the group consisting of oxo, C₁₋₆alkyl,

-C₁₋₆alkylC(=O)C₁₋₆alkyl, -C₁₋₆alkylC(=O)C₁₋₆alkoxy, -C₁₋₆alkyl-aryl,

-C₁₋₆alkylC(=O)aryl, -C(=O)(C₁₋₆)alkyl, -C(=O)(C₁₋₆)alkoxy,

30 -C(=O)aryl, -SO₂aryl, aryl, heteroaryl, and heterocyclyl,

wherein the aryl portion of any aryl-containing substituent of Cy is

optionally substituted with one, two or three substituents

independently selected from the group consisting of C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkylthio, halogen, hydroxy, NH₂, -NH(C₁₋₆alkyl), and -N(C₁₋₆)dialkyl, and

wherein heterocyclyl is optionally substituted with aryl, one, two or three halogen atoms, or one, two or three oxo substituents; and heterocyclyl is optionally spiro-fused to said Cy;

n is 0 or 1;

W is O or S;

X is hydrogen or C₁₋₃alkyl;

Y is independently selected from the group consisting of C₁₋₆alkyl substituted with -OSO₂NH₂ or hydroxy; SO₃H, CO₂H, heteroaryl, -OC(=O)NH₂, and P(=O)OR⁵R⁶; provided that when Y is CO₂H, Ring A must be a bicyclic ring system;

R⁵ is selected from the group consisting of hydrogen; C₁₋₆alkyl and aryl,

wherein C₁₋₆alkyl is optionally substituted with NH₂, -NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, 1,3-dioxolan-2-yl, C₁₋₆alkylcarbonyloxy-, C₁₋₆alkoxycarbonyloxy-, C₁₋₆alkylcarbonylthio-, (C₁₋₆)alkylaminocarbonyl-, di(C₁₋₆)alkylaminocarbonyl-, one, two or three halogen atoms, or hydroxy, and

wherein aryl is optionally substituted with C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkylthio, C₂₋₆alkenyl, -NH₂, -NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, aryl, heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy, or nitro, and, optionally, when R⁶ is C₁₋₈alkoxy, R⁵ and R⁶ may each be taken together with the atoms to which they are attached to form a 5-8 membered monocyclic ring;

R⁶ is selected from the group consisting of C₁₋₈alkyl, C₁₋₈alkoxy, C₂₋₈alkenyl, heteroaryl, aryl, and hydroxy,

wherein C₁₋₈alkyl, C₁₋₈alkoxy, and C₂₋₈alkenyl are optionally substituted with a substituent selected from the group consisting of

C₁₋₆alkoxy, aryl, heterocyclyl, heteroaryl, NH₂, -NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, C₁₋₆alkylcarbonyloxy-, C₁₋₆alkylcarbonylthio-,

C₁₋₆alkoxycarbonyloxy-, (C₁₋₆)alkylaminocarbonyl-,
di(C₁₋₆)alkylaminocarbonyl-, one, two or three halogen atoms and
hydroxy,

wherein when R⁶ is C₁₋₈alkyl, said C₁₋₈alkyl is optionally

5 substituted with halogen selected from up to three chlorine
atoms or up to seven fluorine atoms, and

wherein the heteroaryl and aryl substituents of R⁶ are optionally
substituted with a substituent independently selected from
the group consisting of C₁₋₆alkyl, C₁₋₆alkoxy, C₂₋₆alkenyl,
10 C₁₋₆alkylthio, -NH₂, -NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, aryl,
heteroaryl, aryloxy, heteroaryloxy, halogen, hydroxy and
nitro;

R⁴ is one, two or three substituents selected from the group consisting of
hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, C₁₋₆alkoxy, C₁₋₆alkylthio,
15 aryl(C₁₋₆)alkyl, aryl(C₂₋₆)alkenyl, halogen, hydroxy, -C(=O)Cy,
-C(=O)NR³¹R³², aryl, -CO₂H, oxo and cyano,

wherein C₁₋₆alkyl, C₂₋₆alkenyl and C₁₋₆alkoxy are each optionally
substituted with -NR³³R³⁴, aryl, heteroaryl, cycloalkyl, one, two or
three halogen atoms, or hydroxy, and

20 wherein aryl and heteroaryl are each optionally substituted with a
substituent independently selected from the group consisting of
C₁₋₆alkyl, C₁₋₆alkoxy, C₂₋₆alkenyl, C₁₋₆alkylthio, -NH₂,
-NH(C₁₋₆)alkyl, -N(C₁₋₆)dialkyl, aryl, heteroaryl, aryloxy,
heteroaryloxy, one, two or three halogen atoms, hydroxy and
25 nitro;

R³¹, R³², R³³ and R³⁴ are substituents independently selected from the
group consisting of hydrogen, C₁₋₆alkyl and aryl,


wherein C₁₋₆alkyl is optionally substituted with hydroxy, aryl,

-C(=O)C₁₋₄alkoxy, NH₂, -NH(C₁₋₆alkyl), or -N(C₁₋₆)dialkyl,
30 and, optionally, R³¹ and R³² or R³³ and R³⁴ are each taken
together with the atoms to which they are attached to form a ring
of five to seven members;

and enantiomers, diastereomers, polymorphs or pharmaceutically acceptable salts thereof.

2. The compound of claim 1, wherein Ring A is naphthyl and n is 0.
3. The compound of claim 1, wherein R¹, R², R³ and X are each hydrogen
5 and W is O.
4. The compound of claim 1, wherein Y is independently SO₃H or P(=O)OR⁵R⁶.
5. The compound of claim 4, wherein Y is P(=O)OR⁵R⁶.
6. The compound of claim 1, wherein R⁵ is hydrogen or C₁₋₆alkyl.
- 10 7. The compound of claim 6, wherein R⁵ is hydrogen or methyl.
8. The compound of claim 1, wherein R⁶ is selected from the group consisting of C₁₋₆alkyl and hydroxy.
9. The compound of claim 8, wherein R⁶ is selected from the group consisting of methyl and hydroxy.
- 15 10. The compound of claim 1, wherein R⁴ is one, two or three substituents selected from the group consisting of hydrogen, C₁₋₆alkyl, C₁₋₆alkoxy, halogen and hydroxy.
11. The compound of claim 10, wherein R⁴ is one or two or three
20 substituents selected from the group consisting of hydrogen, chlorine, bromine, hydroxy, methyl and methoxy.

12. A compound of Formula (Ia):



wherein

Ring A is aryl;

5 n is 0 or 1;

R⁵ is selected from the group consisting of hydrogen and C₁₋₆alkyl;

R⁶ is selected from the group consisting of C₁₋₈alkyl and hydroxy;

R⁴ is one, two or three substituents selected from the group consisting of hydrogen, C₁₋₆alkyl, C₁₋₆alkoxy, halogen and hydroxy;

10 and enantiomers, diastereomers, polymorphs or pharmaceutically acceptable salts thereof.

13. The compound of claim 12, wherein Ring A is naphthyl and n is 0.

14. The compound of claim 12, wherein R⁵ is hydrogen or methyl.

15. The compound of claim 12, wherein R⁶ is selected from the group consisting of methyl and hydroxy.

16. The compound of claim 12, wherein R⁴ is one or two or three substituents selected from the group consisting of hydrogen, chlorine, bromine, hydroxy, methyl and methoxy.

17. A compound or an enantiomer, diastereomer, polymorph or pharmaceutically acceptable salt thereof selected from the group consisting of:

[benzofuran-3-yl-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,

[(5-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,

[(5-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
[(6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
[(5,7-dibromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
[(5,7-dibromo-6-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
[(5,7-dibromo-6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
[(5-chloro-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
[(4-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
[(5-bromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
[(7-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
[(5-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphonic acid,
[benzofuran-3-yl-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
methyl-[(5-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-phosphinic acid,
[(5-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
[(6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
[(5,7-dibromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
[(5,7-dibromo-6-methyl-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
[(5,7-dibromo-6-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
[(5-chloro-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
[(4-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
[(5-bromo-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid,
[(7-methoxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid, and

[(5-hydroxy-benzofuran-3-yl)-(naphthalen-2-ylcarbamoyl)-methyl]-methyl-phosphinic acid.

18. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier.
19. A composition comprising the compound of claim 12 and a pharmaceutically acceptable carrier.
- 5 20. Use of the compound of claim 1 for the manufacture of a medicament for treating inflammatory or chymase mediated disorders.
21. The use of claim 20 wherein the disorder is selected from the group consisting of allergic rhinitis, viral rhinitis, asthma, chronic obstructive pulmonary diseases, bronchitis, pulmonary emphysema, psoriasis,
10 arthritis, reperfusion injury, ischemia, hypertension, hypercardia myocardial infarction, heart failure damage associated with myocardial infarction and cardiac hypertrophy, arteriosclerosis, sarcoidosis, vascular stenosis or restenosis, pulmonary fibrosis, kidney fibrosis, liver fibrosis, post surgical adhesion formation, systemic sclerosis, keloid scars,
15 rheumatoid arthritis, bullous pemphigoid and atherosclerosis.
22. The use of claim 21 wherein the disorder is selected from the group consisting of allergic rhinitis, asthma, chronic obstructive pulmonary disease, bronchitis, pulmonary emphysema, acute lung injury, heart failure damage associated with myocardial infarction and cardiac
20 hypertrophy.
23. The use of claim 22 wherein the disorder is selected from the group consisting of allergic rhinitis, asthma and heart failure damage associated with myocardial infarction and cardiac hypertrophy.

