



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2610773 C 2014/02/11

(11)(21) **2 610 773**

(12) **BREVET CANADIEN**
CANADIAN PATENT

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2006/06/21
(87) Date publication PCT/PCT Publication Date: 2006/12/28
(45) Date de délivrance/Issue Date: 2014/02/11
(85) Entrée phase nationale/National Entry: 2007/12/05
(86) N° demande PCT/PCT Application No.: JP 2006/312385
(87) N° publication PCT/PCT Publication No.: 2006/137423
(30) Priorité/Priority: 2005/06/24 (US11/165,201)

(51) Cl.Int./Int.Cl. *A61K 31/7048* (2006.01),
A61P 11/00 (2006.01), *C07H 17/08* (2006.01)

(72) Inventeur/Inventor:
TAKAYAMA, KIYOSHI, JP

(73) Propriétaire/Owner:
TAISHO PHARMACEUTICAL CO., LTD., JP

(74) Agent: TORYS LLP

(54) Titre : CLARITHROMYCINE OU SEL DE CELLE-CI A UTILISER POUR LE TRAITEMENT OU LA PREVENTION DE
MALADIES PULMONAIRES INDUITES PAR LA DESTRUCTION D'ALVEOLES PULMONAIRES
(54) Title: CLARITHROMYCIN OR A SALT THEREOF FOR USE IN THE TREATMENT OR PREVENTION OF
PULMONARY DISEASE INDUCED BY DESTRUCTION OF PULMONARY ALVEOLI

(57) Abrégé/Abstract:

For the purpose of treatment and/or prevention of pulmonary disorders caused by the destruction of pulmonary alveoli resulting from smoking, air pollution, noxious gas, etc., there are provided, among others, a method of administering clarithromycin or a salt thereof to a mammal and a pharmaceutical composition comprising clarithromycin or a salt thereof.



ABSTRACT

For the purpose of treatment and/or prevention of pulmonary disorders caused by the destruction of pulmonary alveoli resulting from smoking, air pollution, noxious gas, 5 etc., there are provided, among others, a method of administering clarithromycin or a salt thereof to a mammal and a pharmaceutical composition comprising clarithromycin or a salt thereof.

DESCRIPTION

CLARITHROMYCIN OR A SALT THEREOF FOR USE IN THE TREATMENT OR PREVENTION OF PULMONARY DISEASE INDUCED BY DESTRUCTION OF PULMONARY ALVEOLI

5

Technical Field

[0001] This invention relates to the use of clarithromycin or a salt thereof for the treatment or prevention of pulmonary disorders caused by the destruction of pulmonary alveoli. More particularly, it relates to a therapeutic or prophylactic agent comprising as an effective ingredient, clarithromycin or a salt thereof, a pharmaceutical composition comprising clarithromycin or a salt thereof, and a method of administering clarithromycin or a salt thereof to a mammal, for the purpose of such treatment or prevention, as well as to the use of clarithromycin or a salt thereof in the manufacture of a medicament.

Background Art

[0002] The lung branches out from bronchi and is gradually subdivided finely to form the eventual baggy structures which are referred to as "alveoli (or alveolus)". The alveolus is composed of alveolar epithelial cells, vascular cells, and extracellular matrices (such as elastin or collagen). The alveolus is an important site where gas exchange in the body is conducted.

25

[0003] Pulmonary emphysema is a symptom where the alveolar walls are destroyed and the microstructures of the alveoli are hollowed out. Pulmonary emphysema reduces gas exchange efficiency and the elastic recoil of the whole lung, and it eventually lowers pulmonary functions.

[0004] It was known that inflammatory cells, such as alveolar macrophages or neutrophiles recruited into the lung, were strongly implicated in pulmonary emphysema which were activated by smoking, air pollution, noxious gas or the like (Barnes, P.J. et al., *Nature Reviews/Drug Discovery*, Vol. 1, 437-446 (2002)). It was reported that the macrophages and neutrophiles were activated by harmful substances in tobacco smoke or polluted air (although the detailed mechanism is unknown) and released substances (such as cytokines or proteases) capable of enhancing inflammation to take part in the destruction of the alveolar walls (Barnes, P.J. et al., *Pharmacol Rev.* 2004 Dec; 56(4):515-48).

[0005] At present, bronchodilators which improve the restriction of airflow, such as anticholinergic agents and β_2 receptor stimulants, are widely used as therapeutic agents for pulmonary emphysema. However, they have not been able to alleviate the destruction of pulmonary alveoli and to retard the progression of the pulmonary emphysema itself. The use of anti-inflammatory agents represented by steroids is recommended in the acute exacerbation stage of chronic obstructive pulmonary diseases (COPD), i.e., cases involving infection and deterioration of pulmonary functions. It has,

however, been reported that they are ineffective against the pulmonary emphysema itself. As pertinent art, azithromycins having a 15-membered ring were reported to display effectiveness against non-infectious inflammatory diseases (Japanese Published Application 2004-531539; JPA2004531539 or WO2002/087596). However, it was also pointed out that 14-membered macrolides differ from 15-membered macrolides with respect to pharmacological actions including anti-inflammatory action (*ibid.*).

10 Disclosure of Invention

Problems to Be Solved

[0006] It is an object of this invention to provide a therapeutic or prophylactic agent for a pulmonary disorder caused by the destruction of pulmonary alveoli.

15 Means for Solving the Problems

[0007] As a result of intensive and diligent research, the present inventors have accomplished this invention upon finding that clarithromycin reduces inflammatory reaction to improve a pulmonary emphysema condition in model mice with tobacco-smoke induced pulmonary emphysema mimicking human pulmonary emphysema and that clarithromycin is useful as a therapeutic or prophylactic agent for a pulmonary disease (particularly, pulmonary emphysema).

20 [0008] Specifically, this invention provides a method for treating or preventing a pulmonary disorder caused by the destruction of pulmonary alveoli in a mammal (principally, in human) in need of such treatment or prevention, the

method comprising administering to the mammal, an effective amount of clarithromycin or a salt thereof.

[0009] This invention also provides a pharmaceutical composition for the treatment or prevention of a pulmonary disorder caused by the destruction of pulmonary alveoli comprising an effective amount of clarithromycin or a salt thereof and a pharmaceutically acceptable carrier.

[0010] Further, this invention provides a therapeutic or prophylactic agent for a pulmonary disorder caused by the destruction of pulmonary alveoli comprising as an effective ingredient, clarithromycin or a salt thereof.

[0011] In addition, this invention provides the use of clarithromycin or a salt thereof in the manufacture of a medicament for the treatment or prevention of a pulmonary disorder caused by the destruction of pulmonary alveoli.

[0012] In the method of treatment/prevention and the therapeutic/prophylactic pharmaceutical composition mentioned above, the efficacy of clarithromycin or a salt thereof is excellent where the pulmonary disorder is pulmonary emphysema or a pulmonary emphysema condition.

[0013] In the method of treatment/prevention and the therapeutic/prophylactic pharmaceutical composition mentioned above, the efficacy of clarithromycin or a salt thereof is especially excellent where the pulmonary emphysema condition is chronic obstructive pulmonary disease.

[0014] This invention also provides an *in vivo* method for reducing the cell counts of macrophages and/or neutrophiles

in an mammal, the method comprising administering to the mammal, clarithromycin or a salt thereof in an amount sufficient to reduce the cell counts.

Effects of the Invention

5 [0015] According to this invention, it has been demonstrated that clarithromycin or a salt thereof is effective as a therapeutic or prophylactic agent for a pulmonary disorder caused by the destruction of pulmonary alveoli.

10 Best Mode for Carrying Out the Invention

15 [0016] As used in the present specification, the term "pulmonary disorder caused by the destruction of pulmonary alveoli" refers to a pulmonary disease triggered by the destruction of the cells of pulmonary alveoli due to smoking, air pollution, noxious gas, aging or the like. Especially, of note is "pulmonary emphysema (condition)" by which owing to the destruction of pulmonary alveoli, the pulmonary alveolar wall disintegrates to cause the adhesion of adjoining pulmonary alveoli and to form a cavity. In the pulmonary emphysema condition, the pulmonary alveoli of the peripheral respiratory tract are further destroyed, resulting in the obstructive disorder of airflow. This symptom is classified into "chronic obstructive pulmonary disease." The diseases displaying symptoms that are characterized by the aforementioned pulmonary alveoli destruction are 20 collectively referred to as "pulmonary disorder caused by 25

the destruction of pulmonary alveoli." These pulmonary disorders may often be accompanied by infections.

[0017] Clarithromycin used in this invention is a known compound; and its detailed description (production method, 5 antibacterial activity, etc.) is disclosed in, for example, Japanese Published Patent Application Sho 61-52839 (JP61052839B).

As used herein, the term "a salt of clarithromycin" refers to a 10 pharmaceutically acceptable salt of clarithromycin. Specifically, there are mentioned the salts with organic acids such as tartaric acid, citric acid, stearic acid, and succinic acid; the salt with methanesulfonic acid; the salt with aminoethanesulfonic acid; and the salts with amino 15 acids such as aspartic acid and glutamic acid. In place of clarithromycin a pharmaceutically acceptable derivative thereof may be used in this invention. This derivative refers to one that bears the basic skeleton of clarithromycin which has been derivatized and that displays 20 the same pharmacological function as does clarithromycin. The particularly useful derivative is an ester of clarithromycin that experiences hydrolysis *in vivo* to release clarithromycin. These derivatives are also referred 25 to as "prodrug(s)" and particular esters are well known to one skilled in the art.

[0018] When clarithromycin is used in this invention, it may be formulated into standard pharmaceutical preparations.

Specifically, clarithromycin or a salt thereof is compounded with a pharmaceutically acceptable carrier (e.g., an excipient, binder, disintegrating agent, flavoring agent, emulsifier, diluent, and solubilizer) to prepare a pharmaceutical composition. This pharmaceutical composition may be administered to a mammal in the form of a preparation that is suitable for an oral or parenteral route, including tablets, pills, powders, granules, capsules, solutions, emulsions, suspensions, injectables, suppositories, inhalants and transdermals.

[0019] In producing these preparations, there may be added a solvent, a solubilizer, an isotonicating agent, a preservative, an antioxidant, an excipient, a binder, a lubricant, and a stabilizer.

[0020] Specifically, the solvents include water, physiological saline, etc.; the solubilizers include ethanol, polysorbates, cremophor, etc.; the excipients include lactose, starch, crystalline cellulose, mannitol, maltose, calcium hydrogenphosphate, light anhydrous silicic anhydride, calcium carbonate, etc.; the binders include starch, polyvinylpyrrolidone, hydroxypropyl cellulose, ethyl cellulose, carboxymethylcellulose, gum arabic, etc.; the disintegrating agents include starch, calcium carboxymethylcellulose, etc.; the lubricants include magnesium stearate, talc, hydrogenated oils, etc.; and the stabilizers include lactose, mannitol, maltose, polysorbates, macrogols, polyoxyethylene hydrogenated castor oil, etc.

[0021] To the preparations, there may be added glycerol, dimethylacetamide, 70% sodium lactate, a surfactant, and a basic substance (e.g., sodium hydroxide, ethylenediamine, ethanolamine, sodium carbonate, arginine, meglumine, and Tris-aminomethane) where necessary.

[0022] As used herein, the term "effective amount" means an amount or an amount of composition sufficient to exert a desired pharmacological effect when clarithromycin or a salt thereof, preferably as a pharmaceutical composition, is administered to a mammal in need of treatment or prevention. When clarithromycin or a salt thereof is to be used for the treatment purpose, the desired pharmacological effect means that the cure or alleviation of the symptom can be achieved in the mammal showing the developed symptom. When clarithromycin or a salt thereof is to be used for the prevention purpose, the desired pharmacological effect means the inhibition of the crisis in a mammal. In this case, preventive administration is conducted to the mammal suspected of potential crisis or having high risk of crisis prior to such crisis.

[0023] The dosage of clarithromycin or a salt thereof will be determined in consideration of the animal test results and other various circumstances so that the full dose may not exceed a certain level when it is administered singly or repeatedly. Needless to indicate, the actual dosage will vary depending on the method of administration and the conditions of the patient or the animal to be

treated, specifically, within the range of from 10 to 1000 mg and on the age, the body weight, the gender, the susceptibility, the diet (feed), the administration time, any drug used together, or the degree of the symptom. The 5 suitable dosage and the frequency of administration need to be determined by the medical specialist after optimum dose determination according to the aforementioned guideline.

Examples

[0024] This invention will be concretely described further by referring to a preparation example and test 10 examples; however, the invention should not be limited to those examples.

Preparation Example 1

| | |
|-------------------------|----------|
| clarithromycin | 50 mg |
| lactose | 40 mg |
| corn starch | 49.75 mg |
| crystalline cellulose | 17 mg |
| carmellose calcium | 17 mg |
| hydroxypropyl cellulose | 5.25 mg |
| magnesium stearate | 1 mg |
| total | 180 mg |

[0025] Clarithromycin, lactose, corn starch, crystalline cellulose, and carmellose calcium were uniformly mixed. To this was added a 10% hydroxypropyl cellulose aqueous 20 solution. After blending the mixture, the granules were dried and sieved with a 30M-screen to make uniform granules.

Magnesium stearate was added to the granules and it was compressed to form tablets.

[0026] The pharmacological action of clarithromycin (improvement to a pulmonary emphysema condition) will be 5 next explained by way of test examples.

[0027] Test Example 1: Pharmacological effect in model mice with tobacco smoke-induced pulmonary emphysema

The model mice with tobacco smoke-induced pulmonary emphysema mimicking human pulmonary emphysema were 10 constructed according to the method as described in Hautamaki, R.D. et al., Science 1997, 277:2002-2004 or Shapiro, S.D. et al., Am. J. Pathol (2003) 163:2329-2335. Specifically, C57black/6 female mice of 12-weeks old were exposed to tobacco smoke at two cigarettes per day, 6 days 15 per week for 6 months.

[0028] Administration of clarithromycin was carried out orally at the rate of 25, 50 or 100 mg/body weight twice daily (morning and evening) during the tobacco smoke exposure period. The plasma $AUC_{0-\infty}$ of clarithromycin 20 obtained when 50 mg/kg was singly administered to the mice was $2.677 \mu\text{g} \cdot \text{hr}/\text{ml}$ while the plasma $AUC_{0-\infty}$ of clarithromycin obtained when 200 mg of clarithromycin was singly 25 administered to a normal human subject was $8.98 \mu\text{g} \cdot \text{hr}/\text{ml}$. The former concentration was substantially lower than the latter one and was an adequate concentration to be used clinically.

[0029] The index of the improvement to the pulmonary emphysema condition followed the method as described in Hautamaki, R.D. et al., *ibid.* or Shapiro, S.D. et al., *ibid.*

[0030] Hematoxylin eosin-stained pathological specimens were prepared from the model mice and ten visual fields under a microscope were randomly selected, where the average sizes of pulmonary alveoli were measured. An increase in the average size of pulmonary alveolus is considered as an index of the destruction of pulmonary alveoli, i.e., the pulmonary emphysema condition.

[0031] Table 1 shows the improvement factor of pulmonary emphysema in the model mice with tobacco smoke induced pulmonary emphysema. Six months' smoking (i.e., exposure to tobacco smoke) increased the pulmonary emphysema rate by 20% relative to the non-smoking group (control). The pulmonary emphysema was improved by 47% in the group administered with clarithromycin (25 mg/kg) and by a maximum of up to 87% in the group administered with clarithromycin (50 mg/kg). These results indicate that clarithromycin is effective against pulmonary emphysema (or pulmonary emphysema condition).

[0032]

Table 1

| | pulmonary alveolus size (μ m) | statistical significance | pulmonary emphysema rate (%) | improvement factor (%) |
|-----------------------------|--|-----------------------------|------------------------------------|---------------------------|
| non-smoking | 32.21 \pm 0.507 | | 0 | |
| smoking + solvent | 38.51 \pm 0.765 | 0.0004 (vs. non-smoking) | 20 | 0 |
| smoking + CAM(25 mg/kg) | 35.60 \pm 1.425 | 0.1001 (vs. smoking) | 10.6 | 47 |
| smoking + CAM(50 mg/kg) | 33.16 \pm 0.890 | 0.0046 (vs. smoking) | 3 | 85 |
| smoking + CAM(100 mg/kg) | 34.35 \pm 0.936 | 0.0092 (vs. smoking) | 6.7 | 66.5 |

(mean \pm SE)

[0033] The cell counts of inflammatory cells (macrophages and neutrophiles) in the bronchoalveolar lavage fluids after 5 six months' exposure were measured to provide an index of the inflammatory condition in the model mice with tobacco smoke-induced pulmonary emphysema. The results are shown in Fig. 1A and Fig. 1B.

[0034] The cell counts of macrophages and neutrophiles increased as the pulmonary emphysema progressed. These 10 cells produce the proteases that are involved in pulmonary emphysema. These cell increases exacerbate the pulmonary emphysema. Administration of clarithromycin reduced the cell counts of macrophages and neutrophiles in a dose-dependant manner. This result indicates that clarithromycin 15 inhibits inflammation reaction and suppresses the progression of pulmonary alveoli destruction.

[0035] It has been reported that as pulmonary emphysema progresses, pulmonary functions deteriorate. It has also 20 been known that the pulmonary compliance, which is an index of pulmonary functions, increases in model mice with tobacco

smoke-induced pulmonary emphysema. Thus, the pulmonary compliance was measured in the model mice with tobacco smoke-induced pulmonary emphysema after six months' exposure. The result is shown in Fig. 2.

5 [0036] As the pulmonary emphysema progressed, the pulmonary compliance (the index of pulmonary functions) was increasing in the model mice with tobacco smoke-induced pulmonary emphysema; however, the pulmonary compliance recovered upon the administration of clarithromycin. This 10 result was in accord with the improvement of the pulmonary emphysema ratio as well as with the decrease in the number of the inflammatory cells, and it indicates that the clarithromycin administration is useful for the treatment of pulmonary emphysema (or pulmonary emphysema condition).

15 Industrial Applicability

[0037] According to this invention, clarithromycin or a salt thereof is effective as a therapeutic or prophylactic agent for a pulmonary disorder caused by the destruction of pulmonary alveoli.

20 Brief Description of the Drawings

[0038] Fig. 1A and Fig. 1B show the cell counts of inflammatory cells (macrophages and neutrophiles) in bronchoalveolar lavage fluids from model mice with tobacco smoke-induced pulmonary emphysema after six months' exposure. Fig. 1A shows the cell count of macrophages. Fig. 1B shows 25 the cell count of neutrophiles.


[0039] Fig. 2 shows the results of pulmonary compliance measured in the model mice with tobacco smoke-induced pulmonary emphysema after six months' exposure.

CLAIMS

1. Use of an effective amount of clarithromycin or a salt thereof for suppressing the progression of the destruction of pulmonary alveoli in pulmonary emphysema in a mammal in need of such suppression.
2. The use according to claim 1, wherein the clarithromycin or a salt thereof is adapted for administration as a composition with a pharmaceutically acceptable carrier.
3. The use according to claim 2, wherein the composition further comprises one or more of a solvent, a solubilizer, an isotonicating agent, a preservative, an antioxidant, an excipient, a binder, a lubricant, and a stabilizer.

1/2

[Fig.1]



【Fig. 2】

