



(86) Date de dépôt PCT/PCT Filing Date: 2016/04/29
(87) Date publication PCT/PCT Publication Date: 2016/11/03
(45) Date de délivrance/Issue Date: 2024/04/16
(85) Entrée phase nationale/National Entry: 2017/09/25
(86) N° demande PCT/PCT Application No.: EP 2016/059669
(87) N° publication PCT/PCT Publication No.: 2016/174234
(30) Priorité/Priority: 2015/04/29 (EP15165808.5)

(51) Cl.Int./Int.Cl. *C11D 3/386* (2006.01),
C11D 1/83 (2006.01), *C11D 3/00* (2006.01),
C11D 3/395 (2006.01), *C11D 7/00* (2006.01),
C11D 7/42 (2006.01), *C11D 7/54* (2006.01)
(72) Inventeurs/Inventors:
BENIE, ASTRID, DK;
KNOTZEL, JURGEN C. F., DK;
BAUER, MIKAEL, DK;
CHRISTENSEN, LARS L. H., DK;
RANNES, JULIE B., DK;
SKAGERLIND, JAN PETER, SE;
GAO, NAN, CN
(73) Propriétaire/Owner:

(54) Titre : POLYPEPTIDES APPROPRIES POUR DETERGENT
(54) Title: POLYPEPTIDES SUITABLE FOR DETERGENT

(57) **Abrégé/Abstract:**

The present invention relates to novel subtilase polypeptides exhibiting increased wash performance and /or improved wash performance and to compositions comprising such polypeptides. The polypeptides of the invention are suitable for use in e.g. cleaning or detergent compositions, such as laundry detergent compositions and dish wash compositions, including automatic dish wash compositions. The present invention also relates to isolated DNA sequences encoding the polypeptides, expression vectors, host cells, and methods for producing and using the polypeptides of the invention.



(73) Propriétaires(suite)/Owners(continued):NOVOZYMES A/S, DK

(74) Agent: WILSON LUE LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2016/174234 A3

(43) International Publication Date
3 November 2016 (03.11.2016)

(51) International Patent Classification:
C11D 3/386 (2006.01)

(21) International Application Number:
PCT/EP2016/059669

(22) International Filing Date:
29 April 2016 (29.04.2016)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
15165808.5 29 April 2015 (29.04.2015) EP

(71) Applicant: NOVOZYMES A/S [DK/DK]; Krogshoejvej 36, 2880 Bagsvaerd (DK).

(72) Inventors: BENNE, Astrid; Krogshoejvej 36, 2880 Bagsvaerd (DK). KNÖTZEL, Jürgen, C, F.; Krogshoejvej 36, 2880 Bagsvaerd (DK). BAUER, Mikael; Krogshoejvej 36, 2880 Bagsvaerd (DK). CHRISTENSEN, Lars, L, H.; Krogshoejvej 36, 2880 Bagsvaerd (DK). RANNES, Julie, B.; Krogshoejvej 36, 2880 Bagsvaerd (DK). SKAGER-LIND, Jan, Peter; 25220 Helsingborg (SE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:
5 January 2017

(54) Title: POLYPEPTIDES SUITABLE FOR DETERGENT

(57) Abstract: The present invention relates to novel subtilase polypeptides exhibiting increased wash performance and /or improved wash performance and to compositions comprising such polypeptides. The polypeptides of the invention are suitable for use in e.g. cleaning or detergent compositions, such as laundry detergent compositions and dish wash compositions, including automatic dish wash compositions. The present invention also relates to isolated DNA sequences encoding the polypeptides, expression vectors, host cells, and methods for producing and using the polypeptides of the invention.



WO 2016/174234 A3

POLYPEPTIDES SUITABLE FOR DETERGENT

Reference to a Sequence Listing

This application contains a Sequence Listing in computer readable form.

Background of the Invention

5 Field of the Invention

The present invention relates to novel polypeptides having protease activity. The polypeptides of the invention are suitable for use in e.g. cleaning or detergent compositions, such as laundry detergent compositions, including liquid and powder laundry compositions, and dish wash compositions, including automatic- and hand- dish wash compositions. The present
10 invention also relates to detergent compositions, such as laundry and dish wash compositions comprising the polypeptides of the invention. The present invention also relates to isolated DNA sequences encoding the polypeptides, expression vectors, host cells, and methods for producing and using the polypeptides of the invention.

Description of the Related Art

15 In the detergent industry, enzymes have for many decades been implemented in washing formulations. Enzymes used in such formulations comprise proteases, lipases, amylases, cellulases, mannosidases as well as other enzymes or mixtures thereof. Commercially the most important enzymes are proteases.

An increasing number of commercially used proteases are protein engineered
20 polypeptides of naturally occurring wild type proteases Everlase®, Relase®, Ovozyme®, Polarzyme®, Liquease®, Liquease Ultra® and Kannase® (Novozymes A/S), Purafast®, Purafect OXP®, FN3®, FN4® and Excellase® (Genencor International, Inc.).

However, various factors make further improvement of the enzymes used in e.g. detergents such as the proteases advantageous. The washing conditions such as temperature
25 and pH changes over time and many stains are still difficult to completely remove under conventional washing conditions. Further, in wash conditions can result in inactivation of the enzymes (due to e.g. pH, temperature or chelation instability) resulting in loss of wash performance during the wash cycle. Thus, despite the intensive research in protease development there remains a need for new and improved proteases that have improved properties such as
30 increased stability and/or improved wash performance compared to the parent subtilase.

Summary of the Invention

The present invention relates to polypeptides having protease activity, said polypeptide comprising an amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID

NO 16 and SEQ ID NO 18.

The polypeptides of the invention have improved stability and/or improved wash performance compared to a reference protease. The present invention further relates to polynucleotides encoding the polypeptides and methods for obtaining the polypeptides of the invention.

The invention further relates to a composition comprising an active protease, wherein the protease is selected from a polypeptide comprising SEQ ID NO 4, SEQ ID NO 2, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

In some aspects the present invention further relates to a composition comprising an active protease selected from a polypeptide comprising SEQ ID NO 4, SEQ ID NO 2, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO

18, wherein the concentration of said active protease in said composition is at least 0.01 mg (e.g., at least 0.02 mg, at least 0.03 mg, at least 0.04 mg, at least 0.05 mg, at least 0.06 mg, at least 0.07 mg, at least 0.08 mg, at least 0.09 mg, at least 0.1 mg, at least 0.2 mg, at least 0.3 mg, at least 0.4 mg, at least 0.5 mg, at least 0.6 mg, at least 0.7 mg, at least 0.8 mg, at least 0.9 mg, at least 1 mg) of active protease per gram (or litre) of composition.

The invention further relates to detergent compositions such as:

A dish wash composition comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) at least one builder, and
- c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the

group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

5 In one aspect, the dish wash composition comprises a builder wherein the builder is selected among phosphates, sodium citrate builders, sodium carbonate, sodium silicate, sodium and zeolites, wherein the builder is added in an amount of about 0-65% by weight, preferably about 40-65% by weight, particularly about 20-65% by weight, particularly from 10% to 50% by weight.

The dish wash composition contains 0-50% by weight, such as about 5% to about 30%, of a detergent co-builder and/or the dish wash compositions is preferably phosphate free.

10 According to one aspect of the invention the builder is selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof. According to one aspect of the invention the bleach component is selected from bleaching catalysts, photobleaches, bleach activators, sources of hydrogen peroxide such as sodium percarbonate, sodium perborates and hydrogen peroxide, preformed peracids and mixtures

15 thereof. According to one aspect the bleach component is a peroxide such as percarbonate, persulfate, perphosphate, persilicate salts. In one aspect the bleaching component includes a percarbonate and bleach catalyst, preferably a manganese compound according to one aspect the bleach catalyst is 1,4,7-trimethyl-1,4,7-triazacyclononane or manganese (II) acetate tetrahydrate (MnTACN). In one aspect the dish wash composition comprising from 1-40 wt%,

20 preferably from 0.5-30 wt%, of bleaching components, wherein the bleach components are a peroxide, preferably percarbonate and a catalyst preferably a metal-containing bleach catalyst such as 1,4,7-trimethyl-1,4,7-triazacyclononane or manganese (II) acetate tetrahydrate (MnTACN).

One aspect of the invention relates to a granular detergent composition comprising:

- 25 a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) 5 wt % to 50 wt % anionic surfactant,
- 30 c) 1 wt % to 8 wt % nonionic surfactant,
- d) 5 wt % to 40 wt % builder such as carbonates, zeolites, phosphate builder, calcium sequestering builders or complexing agents.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16

35 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when

compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

5 In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100,
10 S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at
15 least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the
20 group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

In some aspects, the invention relates to a detergent composition (e.g., liquid or powder laundry detergent composition) comprising:

- 25 a) at least 0.01 mg of active protease per gram (or litre) of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) 5 wt % to 50 wt % anionic surfactant,
- c) 1 wt % to 8 wt % nonionic surfactant,
- 30 d) 5 wt % to 40 wt % builder such as carbonates, zeolites, phosphate builder, calcium sequestering builders or complexing agents.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or
35 deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at

least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

5 In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*,
10 Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence
15 identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10,
20 SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

In some aspects, the invention relates to a detergent composition (e.g., liquid or powder laundry detergent composition) comprising:

- a) at least 0.01 mg of active protease per litre detergent wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO
25 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) 2 wt% to 60 wt% of at least one surfactant,
- c) 5 wt% to 50 wt% of at least one builder.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO
30 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when
35 compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alteration (e.g., a substitution, insertion or deletion)s selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

In one aspect, the anion surfactant is one or more selected from linear alkylbenzenesulfonates (LAS) isomers of LAS, alcohol ether sulfate (AEO, AEOS) and sodium lauryl ether sulfate, sodium laureth sulfate (SLES). In one aspect the nonionic surfactant is selected from nonionic surfactants primary alcohol ethoxylates, such as Neodol 25-7 or Neodol 25-3, alkyl ester sulphates, especially C-9-15 alcohol ethersulfates (AES), C12-15 primary alcohol ethoxylate such as alcohol ethoxylates (AE or AEO), sodium toluene sulfonate (STS) and sodium dodecyl sulfate (SDS). In one aspect the builder is a carbonate, zeolites builder. In one aspect of the invention the builder is selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof. In one aspect the granular detergent composition comprises a bleach component selected from a peroxide such as percarbonate, persulfate, perphosphate, persilicate salts.

In one aspect of the invention, the polypeptides selected from the group consisting of SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 have improved wash performance on blood stains such as C-05 and CS-01 when compared to the polypeptide with SEQ ID NO 19.

In another aspect of the invention, the polypeptides selected from the group consisting of SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 have improved wash performance (e.g., in detergent applications, including liquid and powder laundry detergent applications) on blood stains such as C-05 and CS-01 when compared to the polypeptide with SEQ ID NO 19.

The invention further relates to the use of a dish wash, liquid, powder or granular laundry

composition as described above in a cleaning process, wherein the process is selected from a process of doing hard surface cleaning, such as dish wash, and a laundry process.

Definitions

5 Protease and polypeptides having protease activity

The term “protease” or “polypeptide having protease activity” is defined herein as an enzyme that hydrolyses peptide bonds. It includes any enzyme belonging to the EC 3.4 enzyme group (including each of the thirteen subclasses thereof). The EC number refers to Enzyme Nomenclature 1992 from NC-IUBMB, Academic Press, San Diego, California, including
10 supplements 1-5 published in Eur. J. Biochem. 1994, 223, 1-5; Eur. J. Biochem. 1995, 232, 1-6; Eur. J. Biochem. 1996, 237, 1-5; Eur. J. Biochem. 1997, 250, 1-6; and Eur. J. Biochem. 1999, 264, 610-650; respectively.

The term “protease activity” means a proteolytic activity (EC 3.4). Proteases of the invention are endopeptidases (EC 3.4.21). There are several protease activity types: The three
15 main activity types are: trypsin-like where there is cleavage of amide substrates following Arg or Lys at P1, chymotrypsin-like where cleavage occurs following one of the hydrophobic amino acids at P1, and elastase-like with cleavage following an Ala at P1. For purposes of the present invention, protease activity is determined according to the procedure described in “Materials and Methods” below. The polypeptides of the present invention have at least 20%, e.g., at least 40%,
20 at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, and at least 100% of the protease activity of the polypeptide of SEQ ID NO 19.

The term “reference protease” or “reference polypeptide” means a protease or polypeptide with protease activity these terms are used interchangeably in the application. The reference protease may be any prior art protease having at least one property which is improved by the
25 polypeptide of the invention. Such properties could be wash performance and/or stability in detergent. The reference protease may be a naturally occurring (wild-type) polypeptide or a variant thereof. In a particular embodiment the reference protease is a protease with at least 60 % identity, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 90%, at least 91%, at least 92%, at least
30 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% identity to a polypeptide with the polypeptide with SEQ ID NO 19. SEQ ID NO 19 is the subtilisin 309 protease also known as Savinase, which is well known for its wash performance in detergent and which is widely used in detergents today.

The term “wild-type protease” means a protease expressed by a naturally occurring
35 organism, such as a bacterium, archaea, yeast, fungus, plant or animal found in nature. An example of a wild-type subtilase is subtilisin 309 i.e. the polypeptide with the amino acid sequence SEQ ID NO 19.

The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "sequence identity". For purposes of the present invention, the sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, Trends Genet. 16: 276-277), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the *-nobrief* option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

For purposes of the present invention, the sequence identity between two deoxyribonucleotide sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *supra*) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, *supra*), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EDNAFULL (EMBOSS version of NCBI NUC4.4) substitution matrix. The output of Needle labeled "longest identity" (obtained using the *-nobrief* option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Deoxyribonucleotides} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

The term "variant" means a polypeptide having protease activity comprising an alteration, *i.e.*, a substitution, insertion, and/or deletion, at one or more (*e.g.*, several) positions compared to the precursor polypeptide or starting peptide. *E.g.* a variant of SEQ ID NO 19 is a polypeptide which has at least one an alteration, *i.e.*, a substitution, insertion, and/or deletion, at one or more (*e.g.*, several) positions compared SEQ ID NO 19. A substitution means replacement of the amino acid occupying a position with a different amino acid; a deletion means removal of the amino acid occupying a position; and an insertion means adding one or more (*e.g.* several) amino acids, *e.g.* 1, 2, 3, 4 or 5 amino acids adjacent to and immediately following the amino acid occupying a position.

The term "mature polypeptide" means a polypeptide in its final form following translation and any post-translational modifications, such as N-terminal processing, C-terminal truncation, glycosylation, phosphorylation, etc. It is known in the art that a host cell may produce a mixture of two or more different mature polypeptides (*i.e.*, with a different C-terminal and/or N-terminal amino acid) expressed by the same polynucleotide.

The term "mature polypeptide coding sequence" means a polynucleotide that encodes a mature polypeptide having protease activity.

The term "substantially pure polypeptide" means a preparation that contains at most 10%,

at most 8%, at most 6%, at most 5%, at most 4%, at most 3%, at most 2%, at most 1%, and at most 0.5% by weight of other polypeptide material with which it is natively or recombinantly associated. Preferably, the polypeptides of the invention are at least 92% pure, *e.g.*, at least 94% pure, at least 95% pure, at least 96% pure, at least 97% pure, at least 98% pure, at least 99%, at least 99.5% pure, and 100% pure by weight of the total polypeptide material present in the preparation. The polypeptides of the present invention are preferably in a substantially pure form. This can be accomplished, for example, by preparing the polypeptides by well-known recombinant methods or by classical purification methods.

The term "substantially pure polynucleotide" means a polynucleotide preparation free of other extraneous or unwanted nucleotides and in a form suitable for use within genetically engineered polypeptide production systems. Thus, a substantially pure polynucleotide contains at most 10%, at most 8%, at most 6%, at most 5%, at most 4%, at most 3%, at most 2%, at most 1%, and at most 0.5% by weight of other polynucleotide material with which it is natively or recombinantly associated. A substantially pure polynucleotide may, however, include naturally occurring 5'- and 3'- untranslated regions, such as promoters and terminators. It is preferred that the substantially pure polynucleotide is at least 90% pure, *e.g.*, at least 92% pure, at least 94% pure, at least 95% pure, at least 96% pure, at least 97% pure, at least 98% pure, at least 99% pure, and at least 99.5% pure by weight. The polynucleotides of the present invention are preferably in a substantially pure form.

The term "isolated" means a substance in a form or environment which does not occur in nature. Non-limiting examples of isolated substances include (1) any non-naturally occurring substance, (2) any substance including, but not limited to, any enzyme, variant, nucleic acid, protein, peptide or cofactor, that is at least partially removed from one or more or all of the naturally occurring constituents with which it is associated in nature; (3) any substance modified by the hand of man relative to that substance found in nature; or (4) any substance modified by increasing the amount of the substance relative to other components with which it is naturally associated (*e.g.*, multiple copies of a gene encoding the substance; use of a stronger promoter than the promoter naturally associated with the gene encoding the substance). An isolated substance may be present in a fermentation broth sample.

Detergents

The term "detergent composition" includes, unless otherwise indicated, all forms of detergent compositions such as gel, granulate, liquid, paste, powder, spray or tablet compositions including heavy-duty liquids (HDL), fine-fabric liquid detergents, liquid and/or solid laundry detergents and fine fabric detergents; hard surface cleaning formulations for *e.g.* glass, wood, ceramic and metal counter tops and windows; carpet cleaners; oven cleaners; fabric fresheners; fabric softeners; textile and laundry pre-spotters, as well as dish wash detergents such as hand dishwashing agents (HDW), light duty dishwashing agents, machine dishwashing agents such as automatic dish wash (ADW); all-purpose or heavy-duty washing agents, liquid, gel or paste-form

all-purpose washing agents, liquid cleaning and disinfecting agents, including antibacterial hand-wash types, cleaning bars, mouthwashes, denture cleaners, car or carpet shampoos, bathroom cleaners; hair shampoos and hair-rinses; shower gels, foam baths; metal cleaners; as well as cleaning auxiliaries such as bleach additives and "stain-stick" or pre-treat types.

5 In addition to containing a subtilase polypeptide of the invention, the detergent formulation may contain one or more additional enzymes (such as proteases, amylases, lipases, cutinases, cellulases, endoglucanases, xyloglucanases, pectinases, pectin lyases, xanthanases, peroxidaes, haloperoxygenases, catalases and mannanases, or any mixture thereof), and/or components such as surfactants, builders, chelators or chelating agents, bleach system or bleach
10 components, polymers, fabric conditioners, foam boosters, suds suppressors, dyes, perfume, tannish inhibitors, optical brighteners, bactericides, fungicides, soil suspending agents, anti-corrosion agents, enzyme inhibitors or stabilizers, enzyme activators, transferase(s), hydrolytic enzymes, oxido reductases, bluing agents and fluorescent dyes, antioxidants, and solubilizers.

The term "detergent component" is defined herein to mean the types of chemicals which
15 can be used in detergent compositions. Examples of detergent components are surfactants, hydrotropes, builders, co-builders, chelators or chelating agents, bleaching system or bleach components, polymers, fabric hueing agents, fabric conditioners, foam boosters, suds suppressors, dispersants, dye transfer inhibitors, fluorescent whitening agents, perfume, optical brighteners, bactericides, fungicides, soil suspending agents, soil release polymers, anti-
20 redeposition agents, enzyme inhibitors or stabilizers, enzyme activators, antioxidants, metal care agents and solubilizers. The detergent composition may comprise one or more of any type of detergent component.

The term "dish wash" refers to all forms of washing dishes, e.g. by hand (HDW) or automatic dish wash (ADW). Washing dishes includes, but is not limited to, the cleaning of all
25 forms of crockery such as plates, cups, glasses, bowls, all forms of cutlery such as spoons, knives, forks and serving utensils as well as ceramics, plastics such as melamine, metals, china, glass and acrylics.

The term "dish washing composition" refers to compositions intended for cleaning dishes, table ware, pots, pans, cutlery and all forms of compositions for cleaning hard surfaces
30 areas in kitchens. The terms encompass any materials/compounds selected for domestic or industrial washing applications and the form of the product can be liquid, powder or granulate. In addition to lipase, the automatic dishwashing composition may contain detergent components such as polymers, bleaching systems, bleach activators, bleach catalysts, silicates, dyestuff and metal care agents. The present invention is not restricted to any particular type of dish wash
35 composition or any particular detergent.

The term "automatic dishwashing composition" refers to compositions intended for cleaning dishware such as plates, cups, glasses, bowls, cutlery such as spoons, knives, forks, serving utensils, ceramics, plastics, metals, china, glass and acrylics in a dishwashing machine.

The terms encompass any materials/compounds selected for domestic or industrial washing applications and the form of the product can be liquid, powder or granulate. In addition to lipase, the automatic dishwashing composition may contain detergent components such as polymers, bleaching systems, bleach activators, bleach catalysts, silicates, dyestuff and metal care agents.

5 The term “Hard surface cleaning” is defined herein as cleaning of hard surfaces wherein hard surfaces may include floors, tables, walls, roofs etc. as well as surfaces of hard objects such as cars (car wash) and dishes (dish wash). Dish washing includes but are not limited to cleaning of plates, cups, glasses, bowls, cutlery such as spoons, knives, forks, serving utensils, ceramics, plastics, metals, china, glass and acrylics.

10 The term “improved property” means a characteristic associated with a polypeptide that is improved compared to a reference protease the reference protease is in the context of the present application the polypeptide with the amino acid sequence SEQ ID NO 19. Such improved properties include, but are not limited to, chelator stability, wash performance, protease activity, thermal activity profile, thermostability, pH activity profile, pH stability, substrate/cofactor
15 specificity, improved surface properties, substrate specificity, product specificity, increased stability or solubility in the presence of pretreated biomass, improved stability under storage conditions (storage stability), improved in wash stability and chemical stability. Preferred embodiments are improved wash performance and improved stability, preferably improved in wash stability.

20 The term “improved stability” covers all forms of improved stability, such as improved storage stability, improved pH stability, improved thermostability, improved chelator stability, improved chemical stability and improved in wash stability.

 The term “improved wash performance” is defined herein as a polypeptide displaying an alteration of the wash performance relative to a reference protease such as a protease with SEQ
25 ID NO 19, e.g. by increased stain removal. The term “wash performance” includes wash performance in dish wash but also in laundry. The wash performance may be determined by calculating the so-called intensity value (Int) as defined in the Automatic Mechanical Stress Assay (AMSA) for Automatic Dish Wash in the Materials and Methods section herein.

 The term “laundering” relates to both household laundering and industrial laundering and
30 means the process of treating textiles and/or fabrics with a solution containing a detergent composition of the present invention. The laundering process can for example be carried out using e.g. a household or an industrial washing machine or can be carried out by hand.

 The term “stability” includes storage stability and stability during use, e.g. during a wash process (in wash stability) and reflects the stability of the polypeptides according to the invention
35 as a function of time e.g. how much activity is retained when the polypeptide is kept in solution, in particular in a detergent solution. The stability is influenced by many factors e.g. pH, temperature, detergent composition e.g. amount of builder, surfactants etc.

 The term “textile” means any textile material including yarns, yarn intermediates, fibers,

non-woven materials, natural materials, synthetic materials, and any other textile material, fabrics made of these materials and products made from fabrics (e.g., garments and other articles). The textile or fabric may be in the form of knits, wovens, denims, non-wovens, felts, yarns, and towelling. The textile may be cellulose based such as natural cellulose, including cotton, flax/linen, jute, ramie, sisal or coir or manmade cellulose (e.g. originating from wood pulp) including viscose/rayon, cellulose acetate fibers (tricell), lyocell or blends thereof. The textile or fabric may also be non-cellulose based such as natural polyamides including wool, camel, cashmere, mohair, rabbit and silk or synthetic polymers such as nylon, aramid, polyester, acrylic, polypropylene and spandex/elastane, or blends thereof as well as blends of cellulose based and non-cellulose based fibers. Examples of blends are blends of cotton and/or rayon/viscose with one or more companion material such as wool, synthetic fiber (e.g. polyamide fiber, acrylic fiber, polyester fiber, polyvinyl chloride fiber, polyurethane fiber, polyurea fiber, aramid fiber), and/or cellulose-containing fiber (e.g. rayon/viscose, ramie, flax/linen, jute, cellulose acetate fiber, lyocell). Fabric may be conventional washable laundry, for example, stained household laundry. When the term fabric or garment is used, it is intended to include the broader term textiles as well.

The term "wash performance" is used as an enzyme's ability to remove stains present on the object to be cleaned during e.g. wash, such as laundry (e.g., liquid or powder laundry) or hard surface cleaning (e.g., ADW or HDW). The improvement in the wash performance may be quantified by calculating the so-called intensity value (Int) also defined in AMSA assay, as described in Materials and Methods section.

Delta remission value (ΔRem): The terms "Delta remission" or "Delta remission value" are defined herein as the result of a reflectance or remission measurement at a certain wavelength which typically is 460 nm. The swatch is measured with one swatch of similar colour as background, preferably a swatch from a repetition wash. A swatch representing each swatch type is measured before the wash. The Delta remission is the remission value of the washed swatch minus the remission value of the unwashed swatch.

One way of measuring the wash performance is the Delta enzyme performance value (ΔRem enzyme value): The term "Delta enzyme remission value" is defined herein as the result of a reflectance or remission measurement at 460 nm. The swatch is measured with one swatch of similar colour as background, preferably a swatch from a repetition wash. A swatch representing each swatch type is measured before wash. The Delta enzyme remission is the remission value of the swatch washed in detergent with an enzyme present minus the remission value of a similar swatch washed in a detergent without enzyme present.

Detailed Description of the Invention

The present invention relates to a polypeptide having protease activity, said polypeptide comprising an amino acid sequence having at least 60 % sequence identity to a polypeptide sequence selected from the group consisting of: SEQ ID NO 19.

One aspect of the invention relates to a polypeptide having protease activity, said polypeptide comprising an amino acid sequence having at least 60 %, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity to the amino acid sequence with SEQ ID NO 19.

The invention further relates to a polypeptide having protease activity, said polypeptide comprising an amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 2.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 4.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 6.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 8.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 10.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 12.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 14.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 16.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 18.

In a preferred embodiment, the polypeptides of the invention have improved wash performance compared to a reference protease e.g. compared to SEQ ID NO 19. In a preferred embodiment, the polypeptides of the invention have improved wash performance and/or improved detergent stability compared to a reference protease e.g. compared to SEQ ID NO 19.

The invention further relates to a polypeptide having protease activity, wherein the polypeptide comprise an amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18 wherein said polypeptide has improved wash performance compared to the polypeptide with SEQ ID NO 19.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 2,

and wherein the polypeptide has improved wash performance compared to the polypeptide with SEQ ID NO 19.

5 In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 4, and wherein the polypeptide has improved wash performance compared to the polypeptide with SEQ ID NO 19.

10 In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 6, and wherein the polypeptide has improved wash performance compared to the polypeptide with SEQ ID NO 19.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 8, and wherein the polypeptide has improved wash performance compared to the polypeptide with SEQ ID NO 19.

15 In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 10, and wherein the polypeptide has improved wash performance compared to the polypeptide with SEQ ID NO 19.

20 In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 12, and wherein the polypeptide has improved wash performance compared to the polypeptide with SEQ ID NO 19.

25 In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 14, and wherein the polypeptide has improved wash performance compared to the polypeptide with SEQ ID NO 19.

30 In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 16, and wherein the polypeptide has improved wash performance compared to the polypeptide with SEQ ID NO 19.

In one embodiment, the invention relates to a polypeptide having protease activity, wherein the polypeptide comprises or consists of the amino acid sequence with SEQ ID NO 18, and wherein the polypeptide has improved wash performance compared to the polypeptide with SEQ ID NO 19.

35 In one embodiment, the polypeptides comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 have improved performance, in particular improved in wash performance, compared to a reference polypeptide such as the polypeptide with SEQ ID NO 19. In a preferred embodiment,

the polypeptides comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 have improved stability, in particular improved in wash stability, and on par or improved wash performance compared to a reference polypeptide such as the polypeptide with SEQ ID NO 19.

5 In one embodiment, the invention relates to a polypeptide having protease activity, wherein said polypeptide comprises the amino acid sequence selected from the group consisting of SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18 and wherein the polypeptide sequence has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least
10 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% but less than 100% sequence identity to the amino acid sequence of SEQ ID NO 19.

In an embodiment, the polypeptide comprises the amino acid sequence selected from the group consisting of SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10,
15 SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18 has at least 65% but less than 100% sequence identity to the mature polypeptide of SEQ ID NO 19. In an embodiment, the polypeptide has at least 70% but less than 100% sequence identity to the mature polypeptide of the SEQ ID NO 19. In an embodiment, the polypeptide has at least 75% but less than 100% sequence identity to the mature polypeptide of the SEQ ID NO 19. In an embodiment, the
20 polypeptide has at least 80% but less than 100% sequence identity to the mature polypeptide of the SEQ ID NO 19. In an embodiment, the polypeptide has at least 85% but less than 100% sequence identity to the mature polypeptide of the SEQ ID NO 19. In an embodiment, the polypeptide has at least 90% but less than 100% sequence identity to the mature polypeptide of the SEQ ID NO 19. In an embodiment, the polypeptide has at least 93% but less than 100%
25 sequence identity to the mature polypeptide of the SEQ ID NO 19. In an embodiment, the polypeptide has at least 95% but less than 100% sequence identity to the mature polypeptide of the SEQ ID NO 19. In an embodiment, the polypeptide has at least 96% but less than 100% sequence identity to the mature polypeptide of the SEQ ID NO 19. In an embodiment, the polypeptide has at least 97% but less than 100% sequence identity to the mature polypeptide of the SEQ ID NO 19. In an embodiment, the polypeptide has at least 98% but less than 100%
30 sequence identity to the mature polypeptide of the SEQ ID NO 19.

In one aspect, the total number of alterations in the polypeptides of the invention compared to the SEQ ID NO 19 is between 5 and 30, preferably between 8 and 20, more preferably between 8 and 15, even more preferably between 8 and 10, most preferably between 3 and 8 alterations.
35 In another aspect, total number of alterations compared to the polypeptide with SEQ ID NO 19 is 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 alterations.

In one embodiment, the polypeptide of the invention has improved performance, in

particular improved wash performance, compared to the mature polypeptide of the SEQ ID NO 19. In a preferred embodiment, the polypeptides of the invention have improved performance, in particular improved wash performance, and on par or improved stability compared to the mature polypeptide of the SEQ ID NO 19.

5 In one embodiment, the polypeptides comprise the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18, wherein the polypeptides comprise an alteration at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182,
10 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein the polypeptide has at least 60% but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In one embodiment, the polypeptides comprising the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14,
15 SEQ ID NO 16, SEQ ID NO 18 or SEQ ID NO 19 wherein the polypeptides comprises one or more of the alterations selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S,
20 M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60% but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In one embodiment of the invention, the polypeptides consist of the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ
25 ID NO 14, SEQ ID NO 16 or SEQ ID NO 18. In one embodiment, the polypeptides have improved wash performance and/or improved stability, compared to a reference polypeptide or compared to SEQ ID NO 19.

The polypeptides of the invention may further comprise additional amino acid changes in addition to those listed above such amino acid changes may be of a minor nature, that is
30 conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of 1-30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding
35 domain.

Examples of conservative substitutions are within the groups of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine),

aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, *In, The Proteins*, Academic Press, New York. Common substitutions are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly.

Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

Essential amino acids in a polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for protease activity to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton *et al.*, 1996, *J. Biol. Chem.* 271: 4699-4708. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos *et al.*, 1992, *Science* 255: 306-312; Smith *et al.*, 1992, *J. Mol. Biol.* 224: 899-904; Wlodaver *et al.*, 1992, *FEBS Lett.* 309: 59-64. For Savinase (SEQ ID NO: 4) the catalytic triad comprising the amino acids S221, H64, and D32 (BPN' numbering) is essential for protease activity of the enzyme.

The polypeptides of the invention may consist of 150 to 350, e.g., 175 to 330, 200 to 310, 220 to 300, 240 to 290, 260 to 280 or 269 amino acids.

The polypeptides can be prepared using any mutagenesis procedure known in the art, such as site-directed mutagenesis, synthetic gene construction, semi-synthetic gene construction, random mutagenesis, shuffling, etc.

Site-directed mutagenesis is a technique in which one or more (e.g., several) mutations are introduced at one or more defined sites in a polynucleotide encoding the parent.

Site-directed mutagenesis can be accomplished in vitro by PCR involving the use of oligonucleotide primers containing the desired mutation. Site-directed mutagenesis can also be performed in vitro by cassette mutagenesis involving the cleavage by a restriction enzyme at a site in the plasmid comprising a polynucleotide encoding the parent and subsequent ligation of an oligonucleotide containing the mutation in the polynucleotide. Usually the restriction enzyme that digests the plasmid and the oligonucleotide is the same, permitting sticky ends of the plasmid and the insert to ligate to one another. See, e.g., Scherer and Davis, 1979, *Proc. Natl. Acad. Sci. USA* 76: 4949-4955; and Barton *et al.*, 1990, *Nucleic Acids Res.* 18: 7349-4966.

Site-directed mutagenesis can also be accomplished in vivo by methods known in the art. See, e.g., U.S. Patent Application Publication No. 2004/0171154; Storici et al., 2001, *Nature Biotechnol.* 19: 773-776; Kren et al., 1998, *Nat. Med.* 4: 285-290; and Calissano and Macino, 1996, *Fungal Genet. Newslett.* 43: 15-16.

5 Any site-directed mutagenesis procedure can be used in the present invention. There are many commercial kits available that can be used to prepare polypeptides.

Synthetic gene construction entails in vitro synthesis of a designed polynucleotide molecule to encode a polypeptide of interest. Gene synthesis can be performed utilizing a number of techniques, such as the multiplex microchip-based technology described by Tian et al. (2004, 10 *Nature* 432: 1050-1054) and similar technologies wherein oligonucleotides are synthesized and assembled upon photo-programmable microfluidic chips.

Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988, 15 *Science* 241: 53-57; Bowie and Sauer, 1989, *Proc. Natl. Acad. Sci. USA* 86: 2152-2156; WO 95/17413; or WO 95/22625. Other methods that can be used include error-prone PCR, phage display (e.g., Lowman et al., 1991, *Biochemistry* 30: 10832-10837; U.S. Patent No. 5,223,409; WO 92/06204) and region-directed mutagenesis (Derbyshire et al., 1986, *Gene* 46: 145; Ner et al., 1988, *DNA* 7: 127).

20 Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness et al., 1999, *Nature Biotechnology* 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual 25 amino acid residues in a polypeptide.

Semi-synthetic gene construction is accomplished by combining aspects of synthetic gene construction, and/or site-directed mutagenesis, and/or random mutagenesis, and/or shuffling. Semi-synthetic construction is typified by a process utilizing polynucleotide fragments that are synthesized, in combination with PCR techniques. Defined regions of genes may thus be 30 synthesized de novo, while other regions may be amplified using site-specific mutagenic primers, while yet other regions may be subjected to error-prone PCR or non-error prone PCR amplification. Polynucleotide sub-sequences may then be shuffled.

In an embodiment, the polypeptide selected from the group consisting of the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID 35 NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18 has improved stability, in particular improved in wash stability, compared to the reference polypeptide such as the polypeptide with SEQ ID NO 19 wherein in wash stability is measured using the 'in wash stability assay' as described in the Materials and Methods section herein.

In an embodiment, the polypeptide selected from the group consisting of the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18 has improved wash performance compared to a reference polypeptide such as the polypeptide with SEQ ID NO 19 wherein in wash performance is measured using the Automatic Mechanical Stress Assay (AMSA) for Automatic Dish Wash as described in the Materials and Methods section herein. In an embodiment, the polypeptide selected from the group consisting of the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18 has improved stability, in particular improved in wash stability and improved wash performance compared to the polypeptide of SEQ ID NO 19, wherein in wash stability is measured using the 'In Wash Stability Assay' and wash performance is measured using the Automatic Mechanical Stress Assay (AMSA) for Automatic Dish Wash as described in the Materials and Methods section herein.

The invention further relates to detergent compositions such as dish wash and laundry compositions (e.g., liquid or powder laundry compositions) and the use of such compositions in cleaning processes such as hard surface cleaning and laundry.

One embodiment of the invention relates to a dish wash composition (e.g., ADW or HDW) comprising:

- a) at least 0.01 mg of active protease per gram of composition, wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) at least one builder, and
- c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100,

S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The builder is preferably selected among phosphates, sodium citrate builders, sodium carbonate, sodium silicate, sodium aluminosilicate (zeolite). Suitable builders are alkali metal or ammonium phosphates, polyphosphates, phosphonates, polyphosphonates, carbonates, bicarbonates, borates, citrates, and polycarboxylates. Citrate builders, e.g., citric acid and soluble salts thereof (particularly sodium salt), are polycarboxylate builders. Citrates can be used in combination with zeolite, silicates like the BRITESIL types, and/or layered silicate builders. The builder is preferably added in an amount of about 0-65% by weight, such as about 5% to about 50% by weight. In a dish wash detergent, the level of builder is typically about 40-65% by weight, particularly about 50-65% by weight, particularly from 20% to 50% by weight. The builder and/or co-builder may particularly be a chelating agent that forms water-soluble complexes with Ca and Mg. Any builder and/or co-builder known in the art for use in cleaning detergents may be utilized. Non-limiting examples of builders include zeolites, diphosphates (pyrophosphates), triphosphates such as sodium triphosphate (STP or STPP), carbonates such as sodium carbonate, soluble silicates such as sodium metasilicate, layered silicates (e.g., SKS-6 from Hoechst), and (carboxymethyl)inulin (CMI), and combinations thereof. Further non-limiting examples of builders include citrate, chelators such as aminocarboxylates, aminopolycarboxylates and phosphonates, and alkyl- or alkenylsuccinic acid. Additional specific examples include 2,2',2"-nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTPA), iminodisuccinic acid (IDS), ethylenediamine-N,N'-disuccinic acid (EDDS), methylglycine-N,N-diacetic acid (MGDA), glutamic acid-N,N-diacetic acid (GLDA), 1-hydroxyethane-1,1-diphosphonic acid, N-(2-hydroxyethyl)iminodiacetic acid (EDG), aspartic acid-N-monoacetic acid (ASMA), aspartic acid-N,N-diacetic acid (ASDA), aspartic acid-N-monopropionic acid (ASMP), iminodisuccinic acid (IDA), N-(sulfomethyl)aspartic acid (SMAS), N-(2-sulfoethyl)-aspartic acid (SEAS), N-(sulfomethyl)glutamic acid (SMGL), N-(2-sulfoethyl)-glutamic acid (SEGL), N-methyliminodiacetic acid (MIDA), serine-N,N-diacetic acid (SEDA), isoserine-N,N-diacetic acid (ISDA), phenylalanine-N,N-diacetic acid (PHDA), anthranilic acid-N,N-diacetic acid (ANDA),

sulfanilic acid-N,N-diacetic acid (SLDA) , taurine-N,N-diacetic acid (TUDA) and N'-(2-hydroxyethyl)ethylenediamine-N,N,N'-triacetic acid (HEDTA), diethanolglycine (DEG), and combinations and salts thereof.

Phosphonates suitable for use herein include 1-hydroxyethane-1,1-diphosphonic acid (HEDP), ethylenediaminetetrakis(methylenephosphonic acid) (EDTMPA), diethylenetriaminepentakis(methylenephosphonic acid) (DTMPA or DTPMPA or DTPMP), nitrilotris(methylenephosphonic acid) (ATMP or NTMP), 2-phosphonobutane-1,2,4-tricarboxylic acid (PBTC), hexamethylenediaminetetrakis(methylenephosphonic acid) (HDTMP)

The dish wash composition may also contain 0-50% by weight, such as about 5% to about 30%, of a detergent co-builder.

The detergent composition may include a co-builder alone, or in combination with a builder, for example a zeolite builder. Non-limiting examples of co-builders include homopolymers of polyacrylates or copolymers thereof, such as poly(acrylic acid) (PAA) or copoly(acrylic acid/maleic acid) (PAA/PMA) or polyaspartic acid.

Further exemplary builders and/or co-builders are described in, e.g., WO 09/102854, US 5977053

In one preferred embodiment, the builder is a non-phosphorus based builder such as citric acid and/or methylglycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and / or salts thereof.

One embodiment relates a dish washing composition comprising at least one protease wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18, and a non-phosphate builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T,

N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18, and
- b) 10-50 wt % builder selected from citric acid, methylglycine-N,N-diacetic acid (MGDA) and/or glutamic acid-N,N-diacetic acid (GLDA) and mixtures thereof, and
- c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q,

N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, and
- b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof, and
- c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence

identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 4, and
- b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof, and
- c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the

group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

- 5 a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 6, and
- b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof, and
- c) at least one bleach component.

10 In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135,
15 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

20 In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100,
25 S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at
30 least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

 In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the
35 group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 8, and

b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof, and

5 c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42,
10 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%,
15 but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

35 a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 10, and

b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof, and

c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 12, and
- b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof, and
- c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or

deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 14, and
- b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof, and
- c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at

least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

5 In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*,
10 Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence
15 identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10,
20 SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 16, and
- 25 b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof, and
- c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at
30 least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO

4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 18, and
- b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof, and
- c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100,

S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18. One embodiment the dish wash composition is an automatic dish wash composition (ADW) comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and a second builder selected from carbonates, silicates and zeolites, and
- c) at least one bleach component.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q,

N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The bleach component is preferably selected from percarbonates, persulphates and peracids.

One embodiment concerns a dish wash composition comprising:

a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,

b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and a second builder selected from carbonates, silicates and zeolites, and

c) at least one bleach component, selected from percarbonates, persulphates and peracids.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q,

N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The detergent may contain 0-30% by weight, such as about 1% to about 20%, of a bleaching system. Any bleaching system comprising components known in the art for use in cleaning detergents may be utilized. Suitable bleaching system components include sources of hydrogen peroxide; sources of peracids; and bleach catalysts or boosters.

Sources of hydrogen peroxide:

Suitable sources of hydrogen peroxide are inorganic persalts, including alkali metal salts such as sodium percarbonate and sodium perborates (usually mono- or tetrahydrate), and hydrogen peroxide—urea (1/1).

Sources of peracids:

Peracids may be (a) incorporated directly as preformed peracids or (b) formed in situ in the wash liquor from hydrogen peroxide and a bleach activator (perhydrolysis) or (c) formed in situ in the wash liquor from hydrogen peroxide and a perhydrolase and a suitable substrate for the latter, e.g., an ester.

a) Suitable preformed peracids include, but are not limited to, peroxydicarboxylic acids such as peroxybenzoic acid and its ring-substituted derivatives, peroxy- α -naphthoic acid, peroxyphthalic acid, peroxyauric acid, peroxysearic acid, ϵ -phthalimidoperoxycaproic acid [phthalimidoperoxyhexanoic acid (PAP)], and o-carboxybenzamidoperoxycaproic acid; aliphatic and aromatic diperoxydicarboxylic acids such as diperoxydodecanedioic acid, diperoxyazelaic acid, diperoxysebacic acid, diperoxybrassylic acid, 2-decyldiperoxybutanedioic acid, and diperoxyphthalic, -isophthalic and -terephthalic acids; perimidic acids; peroxymonosulfuric acid; peroxydisulfuric acid; peroxyphosphoric acid; peroxyasilic acid; and mixtures of said compounds. It is understood that the peracids mentioned may in some cases be best added as suitable salts, such as alkali metal salts (e.g., Oxone®) or alkaline earth-metal salts.

b) Suitable bleach activators include those belonging to the class of esters, amides, imides, nitriles or anhydrides and, where applicable, salts thereof. Suitable examples are tetraacetylenediamine (TAED), sodium 4-[(3,5,5-trimethylhexanoyl)oxy]benzene-1-sulfonate (ISONOBS), sodium 4-(dodecanoyloxy)benzene-1-sulfonate (LOBS), sodium 4-(decanoyloxy)benzene-1-sulfonate, 4-(decanoyloxy)benzoic acid (DOBA), sodium 4-


(nonanoyloxy)benzene-1-sulfonate (NOBS), and/or those disclosed in WO98/17767. A particular family of bleach activators of interest was disclosed in EP624154 and particularly preferred in that family is acetyl triethyl citrate (ATC). ATC or a short chain triglyceride like triacetin has the advantage that they are environmentally friendly. Furthermore, acetyl triethyl citrate and triacetin have good hydrolytical stability in the product upon storage and are efficient bleach activators. Finally, ATC is multifunctional, as the citrate released in the perhydrolysis reaction may function as a builder.

Bleach catalysts and boosters

The bleaching system may also include a bleach catalyst or booster.

Some non-limiting examples of bleach catalysts that may be used in the compositions of the present invention include manganese oxalate, manganese acetate, manganese-collagen, cobalt-amine catalysts and manganese triazacyclononane (MnTACN) catalysts; particularly preferred are complexes of manganese with 1,4,7-trimethyl-1,4,7-triazacyclononane (Me3-TACN) or 1,2,4,7-tetramethyl-1,4,7-triazacyclononane (Me4-TACN), in particular Me3-TACN, such as the dinuclear manganese complex [(Me3-TACN)Mn(O)3Mn(Me3-TACN)](PF6)2, and [2,2',2''-nitritotris(ethane-1,2-diylazanylylidene-κN-methanylylidene)triphenolato-κ3O]manganese(III). The bleach catalysts may also be other metal compounds, such as iron or cobalt complexes.

In some embodiments, where a source of a peracid is included, an organic bleach catalyst or bleach booster may be used having one of the following formulae:



(iii) and mixtures thereof; wherein each R¹ is independently a branched alkyl group containing from 9 to 24 carbons or linear alkyl group containing from 11 to 24 carbons, preferably each R¹ is independently a branched alkyl group containing from 9 to 18 carbons or linear alkyl group containing from 11 to 18 carbons, more preferably each R¹ is independently selected from the group consisting of 2-propylheptyl, 2-butyloctyl, 2-pentylnonyl, 2-hexyldecyl, dodecyl, tetradecyl, hexadecyl, octadecyl, isononyl, isodecyl, isotridecyl and isopentadecyl.

Other exemplary bleaching systems are described, e.g. in WO2007/087258, WO2007/087244, WO2007/087259, EP1867708 (Vitamin K) and WO2007/087242. Suitable photobleaches may for example be sulfonated zinc or aluminium phthalocyanines.

Metal care agents

Metal care agents may prevent or reduce the tarnishing, corrosion or oxidation of metals, including aluminium, stainless steel and non-ferrous metals, such as silver and copper. Suitable examples include one or more of the following:

5 (a) benzotriazoles, including benzotriazole or bis-benzotriazole and substituted derivatives thereof. Benzotriazole derivatives are those compounds in which the available substitution sites on the aromatic ring are partially or completely substituted. Suitable substituents include linear or branch-chain C₁-C₂₀- alkyl groups (e.g., C₁-C₂₀- alkyl groups) and hydroxyl, thio, phenyl or halogen such as fluorine, chlorine, bromine and iodine.

10 (b) metal salts and complexes chosen from the group consisting of zinc, manganese, titanium, zirconium, hafnium, vanadium, cobalt, gallium and cerium salts and/or complexes, the metals being in one of the oxidation states II, III, IV, V or VI. In one aspect, suitable metal salts and/or metal complexes may be chosen from the group consisting of Mn(II) sulphate, Mn(II) citrate, Mn(II) stearate, Mn(II) acetylacetonate, K⁺TiF₆ (e.g., K₂TiF₆), K⁺ZrF₆ (e.g., K₂ZrF₆), CoSO₄, Co(NO₃)₂ and Ce(NO₃)₃, zinc salts, for example zinc sulphate, hydrozincite or zinc
15 acetate.;

(c) silicates, including sodium or potassium silicate, sodium disilicate, sodium metasilicate, crystalline phyllosilicate and mixtures thereof.

Further suitable organic and inorganic redox-active substances that act as silver/copper corrosion inhibitors are disclosed in WO 94/26860 and WO 94/26859. Preferably the composition
20 of the invention comprises from 0.1 to 5% by weight of the composition of a metal care agent, preferably the metal care agent is a zinc salt.

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

a) at least 0.01 mg of active protease per gram of composition wherein the protease
25 is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18, and

b) 10-50 wt % builder and

c) at least one bleach component, wherein the bleach is a peroxide and the bleach catalyst is a manganese compound.

30 In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135,
35 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%,

but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The oxygen bleach is preferably percarbonate and the manganese catalyst preferably 1,4,7-trimethyl-1,4,7-triazacyclononane or manganese (III) acetate tetrahydrate

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18, and
- b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof.
- c) at least one bleach component, wherein the bleach is an oxygen bleach and the bleach catalyst is a manganese compound.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%,

but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The oxygen bleach is preferably percarbonate and the manganese catalyst preferably 1,4,7-trimethyl-1,4,7-triazacyclo-nonane or manganese (II) acetate tetrahydrate

According to one embodiment, the dish wash composition is an automatic dish wash composition (ADW) comprising:

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) 10-50 wt % builder selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof, and
- c) 0.1-40 wt%, preferably from 0.5-30 wt%, of bleaching components, wherein the bleach components are a peroxide, preferably percarbonate and a metal-containing bleach catalyst preferably 1,4,7-trimethyl-1,4,7-triazacyclononane or manganese (II) acetate tetrahydrate (MnTACN).

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at

least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

5 In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*,
10 Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence
15 identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10,
20 SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The dish wash composition may also contain at least one surfactant, preferably anionic or nonionic surfactants but also amphoteric surfactants may be used. In general, bleach-stable surfactants are preferred. Preferred anionic surfactants are sulphate surfactants and in particular alkyl ether sulphates, especially C-9-15 alcohol ethersulfates, C12-15 primary alcohol ethoxylate,
25 C8-C16 ester sulphates and C10-C14 ester sulphates, such as mono dodecyl ester sulphates. Non-limiting examples of anionic surfactants include sulfates and sulfonates, in particular, linear alkylbenzenesulfonates (LAS), isomers of LAS, branched alkylbenzenesulfonates (BABS), phenylalkanesulfonates, alpha-olefinsulfonates (AOS), olefin sulfonates, alkene sulfonates, alkane-2,3-diylbis(sulfates), hydroxyalkanesulfonates and disulfonates, alkyl sulfates (AS) such
30 as sodium dodecyl sulfate (SDS), fatty alcohol sulfates (FAS), primary alcohol sulfates (PAS), alcohol ethersulfates (AES or AEOS or FES, also known as alcohol ethoxysulfates or fatty alcohol ether sulfates), secondary alkanesulfonates (SAS), paraffin sulfonates (PS), ester sulfonates, sulfonated fatty acid glycerol esters, alpha-sulfo fatty acid methyl esters (alpha-SFMe or SES) including methyl ester sulfonate (MES), alkyl- or alkenylsuccinic acid, dodeceny/tetradecenyl
35 succinic acid (DTSA), fatty acid derivatives of amino acids, diesters and monoesters of sulfo-succinic acid or salt of fatty acids (soap), and combinations thereof.

The anionic surfactants are preferably added to the detergent in the form of salts. Suitable cations in these salts are alkali metal ions, such as sodium, potassium and lithium and ammonium

salts, for example (2-hydroxyethyl) ammonium, bis(2-hydroxyethyl)ammonium and tris(2-hydroxyethyl)ammonium salts.

Non-limiting examples of nonionic surfactants include alcohol ethoxylates (AE or AEO), alcohol propoxylates, propoxylated fatty alcohols (PFA), alkoxylated fatty acid alkyl esters, such as ethoxylated and/or propoxylated fatty acid alkyl esters, alkylphenol ethoxylates (APE), nonylphenol ethoxylates (NPE), alkylpolyglycosides (APG), alkoxylated amines, fatty acid monoethanolamides (FAM), fatty acid diethanolamides (FADA), ethoxylated fatty acid monoethanolamides (EFAM), propoxylated fatty acid monoethanolamides (PFAM), polyhydroxyalkyl fatty acid amides, or *N*-acyl *N*-alkyl derivatives of glucosamine (glucamides, GA, or fatty acid glucamides, FAGA), as well as products available under the trade names SPAN and TWEEN, and combinations thereof.

Commercially available nonionic surfactants include Plurafac™, Lutensol™ and Pluronic™ range from BASF, Dehypon™ series from Cognis and Genapol™ series from Clariant.

The surfactants useful in the present invention are desirably included in the present detergent compositions at levels of from about 0.1% to about 15%, such as about 2 to about 8% of the composition. The total amount of surfactants typically included in the detergent compositions is typically at least 10% by weight, preferably of from 0.5% to 10% by weight and most preferably from 1% to 5% by weight.

The detergent composition may also be formulated into a granular, powder or liquid detergent for laundry. One embodiment of the invention concerns a granular (or powder or liquid) detergent composition comprising

- a) at least 0.01 mg of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) 5 wt % to 50 wt % anionic surfactant
- c) 1 wt % to 8 wt % nonionic surfactant
- d) 5 wt % to 40 wt % builder such as carbonates, zeolites, phosphate builder, calcium sequestering builders or complexing agents.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The surfactant may be selected among nonionic, anionic and/or amphoteric surfactants as described above, preferably anionic or nonionic surfactants but also amphoteric surfactants may be used. In general, bleach-stable surfactants are preferred. Preferred anionic surfactants are sulphate surfactants and in particular alkyl ether sulphates, especially C-9-15 alcohol ethersulfates, C12-15 primary alcohol ethoxylate, C8-C16 ester sulphates and C10-C14 ester sulphates, such as mono dodecyl ester sulphates. Non-limiting examples of anionic surfactants include sulfates and sulfonates, in particular, linear alkylbenzenesulfonates (LAS), isomers of LAS, branched alkylbenzenesulfonates (BABS), phenylalkanesulfonates, alpha-olefinsulfonates (AOS), olefin sulfonates, alkene sulfonates, alkane-2,3-diylbis(sulfates), hydroxyalkanesulfonates and disulfonates, alkyl sulfates (AS) such as sodium dodecyl sulfate (SDS), fatty alcohol sulfates (FAS), primary alcohol sulfates (PAS), alcohol ethersulfates (AES or AEOS or FES, also known as alcohol ethoxysulfates or fatty alcohol ether sulfates), secondary alkanesulfonates (SAS), paraffin sulfonates (PS), ester sulfonates, sulfonated fatty acid glycerol esters, alpha-sulfo fatty acid methyl esters (alpha-SFMe or SES) including methyl ester sulfonate (MES), alkyl- or alkenylsuccinic acid, dodecenyl/tetradecenyl succinic acid (DTSA), fatty acid derivatives of amino acids, diesters and monoesters of sulfo-succinic acid or salt of fatty acids (soap), and combinations thereof.

The anionic surfactants are preferably added to the detergent in the form of salts. Suitable cations in these salts are alkali metal ions, such as sodium, potassium and lithium and ammonium salts, for example (2-hydroxyethyl)ammonium, bis(2-hydroxyethyl)ammonium and tris(2-hydroxyethyl)ammonium salts.

Non-limiting examples of nonionic surfactants include alcohol ethoxylates (AE or AEO), alcohol propoxylates, propoxylated fatty alcohols (PFA), alkoxylated fatty acid alkyl esters, such as ethoxylated and/or propoxylated fatty acid alkyl esters, alkylphenol ethoxylates (APE), nonylphenol ethoxylates (NPE), alkylpolyglycosides (APG), alkoxylated amines, fatty acid
5 monoethanolamides (FAM), fatty acid diethanolamides (FADA), ethoxylated fatty acid monoethanolamides (EFAM), propoxylated fatty acid monoethanolamides (PFAM), polyhydroxyalkyl fatty acid amides, or N-acyl N-alkyl derivatives of glucosamine (glucamides, GA, or fatty acid glucamides, FAGA), as well as products available under the trade names SPAN and TWEEN, and combinations thereof.

10 Commercially available nonionic surfactants include Plurafac™, Lutensol™ and Pluronic™ range from BASF, Dehypon™ series from Cognis and Genapol™ series from Clariant. The builder is may be non-phosphate such as citrate preferably as a sodium salt and/or zeolites. Phosphonate builder may be any of those described above

The builder is preferably selected among phosphates and sodium citrate builders, sodium
15 carbonate, sodium silicate, sodium aluminosilicate (zeolite) as described above under the dish wash compositions. Suitable builders are described above and include alkali metal or ammonium phosphates, polyphosphates, phosphonates, polyphosphonates, carbonates, bicarbonates, borates, polyhydroxysulfonates, polyacetates, carboxylates, citrates, and polycarboxylates. Citrate builders, e.g., citric acid and soluble salts thereof (particularly sodium salt), are
20 polycarboxylate builders. The builder is preferably added in an amount of about 0-65% by weight, such as about 5% to about 50% by weight, such as 5 to 40 % by weight, such as 10 to 40 % by weight, such as 10 to 30 % by weight, such as 15 to 20 % by weight or such as 20 to 40 % by weight. The builder may be a phosphonate builder including 1-hydroxyethane-1,1-diphosphonic acid (HEDP), ethylenediaminetetra (methylenephosphonic acid) (EDTMPA),
25 diethylenetriaminepentakis (methylenephosphonic acid) (DTMPA or DTPMPA), diethylenetriamine penta (methylenephosphonic acid) (DTPMP), aminotris(methylenephosphonic acid) (ATMP), 2-phosphonobutane-1,2,4-tricarboxylic acid (PBTC) and hexamethylenediaminetetra (methylenephosphonic acid) (HDTMP).

Preferred phosphonates include 1-hydroxyethane-1,1-diphosphonic acid (HEDP) and/or
30 diethylenetriaminepentakis (methylenephosphonic acid) (DTMPA or DTPMPA). The phosphonate is preferably added in an amount of about in a level of from about 0.01 % to about 10 % by weight, preferably from 0.1 % to about 5 % by weight, more preferably from 0.5 % to 3 % by weight of the composition.

The laundry composition may also be phosphate free in the instance the preferred builders
35 include citrate, carbonates and/or sodium aluminosilicate (zeolite).

The proteases of the invention can also be formulated in liquid laundry compositions such as a liquid laundry compositions composition comprising:

a) at least 0.01 mg of active protease per litre detergent wherein the protease is

selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,

- b) 2 wt% to 60 wt% of at least one surfactant
- c) 5 wt% to 50 wt% of at least one builder

5 In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135,
10 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

15 In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100,
20 S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at
25 least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the
30 group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The proteases of the invention can also be formulated in powder laundry compositions such as a powder laundry compositions composition comprising:

- a) at least 0.01 mg of active protease per litre detergent wherein the protease is
35 selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) 2 wt% to 60 wt% of at least one surfactant,
- c) 5 wt% to 50 wt% of at least one builder.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alteration (e.g., a substitution, insertion or deletion)s selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The surfactant may be selected among nonionic, anionic and/or amphoteric surfactants as described above. Preferred surfactants include alkyl ester sulphates, especially C-9-15 alcohol ethersulfates (AES), C12-15 primary alcohol ethoxylate such as alcohol ethoxylates (AE or AEO), sodium toluene sulfonate (STS), sodium dodecyl sulfate (SDS). Commercially available nonionic surfactants include Plurafac™, Lutensol™ and Pluronic™ range from BASF, Dehypon™ series from Cognis and Genapol™ series from Clariant.

The surfactants useful in the present invention are desirably included in the present detergent compositions at levels of from about 0.1% to about 15%, such as about 2 to about 8% of the composition. The total amount of surfactants typically included in the detergent compositions is typically at least 15% by weight, preferably of from 0.5% to 15% by weight and most preferably from 1% to 10% by weight. The total amount of surfactants typically included in

the detergent compositions is typically at least 15% by weight, preferably of from 0.5% to 15% by weight and most preferably from 1% to 10% by weight.

The builder may be selected among nonionic, anionic and/or amphoteric surfactants as described above. One embodiment of the invention concerns a liquid laundry compositions composition comprising:

- a) at least 1 mg/L of active protease, wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18, and
- b) 1% to 15% by weight of at least one surfactant wherein the surfactant is LAS, AEOS, AEO and/or SLES, and
- c) at least one builder.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

One embodiment of the invention concerns a powder laundry compositions composition comprising:

- a) at least 1 mg/L of active protease, wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18, and
- b) 1% to 15% by weight of at least one surfactant wherein the surfactant is LAS, AEOS, AEO and/or SLES, and
- c) at least one builder.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alteration (e.g., a substitution, insertion or deletion)s selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The builder is preferably selected among phosphates and sodium citrate builders, sodium carbonate, sodium silicate, sodium aluminosilicate (zeolite). Suitable builders are described above and include alkali metal or ammonium phosphates, polyphosphates, phosphonates,

polyphosphonates, carbonates, bicarbonates, borates, polyhydroxysulfonates, polyacetates, carboxylates, citrates, and polycarboxylates. Citrate builders, e.g., citric acid and soluble salts thereof (particularly sodium salt), are polycarboxylate builders. The builder is preferably added in an amount of about 0-65% by weight, such as about 5% to about 50% by weight. In a dish wash detergent, the level of builder is typically about 40-65% by weight, particularly about 50-65% by weight, particularly from 20% to 50% by weight. The builder may be a phosphonate builder including 1-hydroxyethane-1,1-diphosphonic acid (HEDP), ethylenediaminetetrakis(methylenephosphonic acid) (EDTMPA), diethylenetriaminepentakis(methylenephosphonic acid) (DTMPA or DTPMPA), diethylenetriaminepentakis(methylenephosphonic acid) (DTPMP), aminotris(methylenephosphonic acid) (ATMP), 2-phosphonobutane-1,2,4-tricarboxylic acid (PBTC) and hexamethylenediaminetetrakis(methylenephosphonic acid) (HDTMP).

Preferred phosphonates include 1-hydroxyethane-1,1-diphosphonic acid (HEDP) and/or diethylenetriaminepentakis (methylenephosphonic acid) (DTMPA or DTPMPA). The phosphonate is preferably added in an amount of about in a level of from about 0.01 % to about 10 % by weight, preferably from 0.1 % to about 5 % by weight, more preferably from 0.5 % to 3 % by weight of the composition. Additional useful builders are described in the dish wash composition section.

The laundry composition (e.g., liquid or powder laundry composition) may also be phosphate free in the instance the preferred builders includes citrate and/or methylglycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and / or salts thereof.

One embodiment of the invention concerns a liquid laundry compositions composition comprising:

- a) at least 0.01 mg of active protease per litre of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18, and
- b) 1% to 15% by weight of at least one surfactant wherein the surfactant is LAS, AEOS and/or SLES, and
- c) 5% to 50% by weight of at least one builder selected from HEDP, DTMPA or DTPMPA.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at

least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

One embodiment of the invention concerns a powder laundry composition, composition comprising:

- a) at least 0.01 mg of active protease per litre of composition wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18, and
- b) 1% to 15% by weight of at least one surfactant wherein the surfactant is LAS, AEOS and/or SLES, and
- c) 5% to 50% by weight of at least one builder selected from HEDP, DTPMA or DTPMPA.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above comprises one or more of the alteration (e.g., a substitution, insertion or deletion)s selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

In some aspects, the polypeptide of the invention comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 as described above consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

The detergent compositions of the invention may contain 0-10% by weight, such as 0.5-5%, 2-5%, 0.5-2% or 0.2-1% of a polymer. Any polymer known in the art for use in detergents may be utilized. The polymer may function as a co-builder as mentioned above, or may provide antiredeposition, fiber protection, soil release, dye transfer inhibition, grease cleaning and/or anti-foaming properties. Some polymers may have more than one of the above-mentioned properties and/or more than one of the below-mentioned motifs. Exemplary polymers include (carboxymethyl)cellulose (CMC), poly(vinyl alcohol) (PVA), poly(vinylpyrrolidone) (PVP), poly(ethyleneglycol) or poly(ethylene oxide) (PEG), ethoxylated poly(ethyleneimine), (carboxymethyl)inulin (CMI), and polycarboxylates such as PAA, PAA/PMA, polyaspartic acid, and lauryl methacrylate/acrylic acid copolymers, hydrophobically modified CMC (HM-CMC) and silicones, copolymers of terephthalic acid and oligomeric glycols, copolymers of poly(ethylene terephthalate) and poly(oxyethylene terephthalate) (PET-POET), PVP, poly(vinylimidazole) (PVI), poly(vinylpyridine-N-oxide) (PVPO or PVPNO) and polyvinylpyrrolidone-vinylimidazole (PVPVI). Further exemplary polymers include sulfonated polycarboxylates, polyethylene oxide and polypropylene oxide (PEO-PPO) and diquaternium ethoxy sulfate. Other exemplary polymers are disclosed in, e.g., WO 2006/130575. Salts of the above-mentioned polymers are also contemplated.

The detergent compositions of the present invention (e.g., liquid or powder laundry compositions) may also include fabric hueing agents such as dyes or pigments, which when formulated in detergent compositions can deposit onto a fabric when said fabric is contacted with a wash liquor comprising said detergent compositions and thus altering the tint of said fabric

through absorption/reflection of visible light. Fluorescent whitening agents emit at least some visible light if subjected to ultraviolet light. In contrast, fabric hueing agents alter the tint of a surface as they absorb at least a portion of the visible light spectrum. Suitable fabric hueing agents include dyes and dye-clay conjugates, and may also include pigments. Suitable dyes include small molecule dyes and polymeric dyes. Suitable small molecule dyes include small molecule dyes selected from the group consisting of dyes falling into the Colour Index (C.I.) classifications of Direct Blue, Direct Red, Direct Violet, Acid Blue, Acid Red, Acid Violet, Basic Blue, Basic Violet and Basic Red, or mixtures thereof, for example as described in WO2005/03274, WO2005/03275, WO2005/03276 and EP1876226. The detergent composition preferably comprises from about 0.00003 wt% to about 0.2 wt%, from about 0.00008 wt% to about 0.05 wt%, or even from about 0.0001 wt% to about 0.04 wt% fabric hueing agent. The composition may comprise from 0.0001 wt% to 0.2 wt% fabric hueing agent, this may be especially preferred when the composition is in the form of a unit dose pouch. Suitable hueing agents are also disclosed in, e.g. WO 2007/087257 and WO2007/087243.

The detergent may contain 0-10% by weight, for example 0-5% by weight, such as about 0.5 to about 5%, or about 3% to about 5%, of a hydrotrope. Any hydrotrope known in the art for use in detergents may be utilized. Non-limiting examples of hydrotropes include sodium benzenesulfonate, sodium p-toluene sulfonate (STS), sodium xylene sulfonate (SXS), sodium cumene sulfonate (SCS), sodium cymene sulfonate, amine oxides, alcohols and polyglycolethers, sodium hydroxynaphthoate, sodium hydroxynaphthalene sulfonate, sodium ethylhexyl sulfate, and combinations thereof.

The detergent compositions of the present invention (e.g., liquid or powder laundry compositions) can also contain dispersants. In particular powdered detergents may comprise dispersants. Suitable water-soluble organic materials include the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals separated from each other by not more than two carbon atoms. Suitable dispersants are for example described in Powdered Detergents, Surfactant science series volume 71, Marcel Dekker, Inc.

The detergent compositions of the present invention (e.g., liquid or powder laundry compositions) may also include one or more dye transfer inhibiting agents. Suitable polymeric dye transfer inhibiting agents include, but are not limited to, polyvinylpyrrolidone polymers, polyamine-N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof. When present in a subject composition, the dye transfer inhibiting agents may be present at levels from about 0.0001 % to about 10%, from about 0.01% to about 5% or even from about 0.1% to about 3% by weight of the composition.

The detergent compositions of the present invention (e.g., liquid or powder laundry compositions) will preferably also contain additional components that may tint articles being cleaned, such as fluorescent whitening agent or optical brighteners. Where present the brightener

is preferably at a level of about 0.01% to about 0.5%. Any fluorescent whitening agent suitable for use in a laundry detergent composition may be used in the laundry composition of the present invention. The most commonly used fluorescent whitening agents are those belonging to the classes of diaminostilbene-sulfonic acid derivatives, diarylpyrazoline derivatives and biphenyl-distyryl derivatives. Examples of the diaminostilbene-sulfonic acid derivative type of fluorescent whitening agents include the sodium salts of: 4,4'-bis[(4-anilino-6-diethanolamino-s-triazin-2-yl)amino]stilbene-2,2'-disulfonate, 4,4'-bis[(4,6-dianilino-s-triazin-2-yl)amino]stilbene-2,2'-disulfonate, 4,4'-bis[4-anilino-6-[methyl(2-hydroxyethyl)amino]-s-triazin-2-ylamino]stilbene-2,2'-disulfonate, 4,4'-bis(4-phenyl-1,2,3-triazol-2-yl)stilbene-2,2'-disulfonate and sodium 5-(2H-naphtho[1,2-d][1,2,3]triazol-2-yl)-2-[(E)-2-phenylvinyl]benzenesulfonate. Preferred fluorescent whitening agents are Tinopal DMS and Tinopal CBS available from BASF. Tinopal DMS is the disodium salt of 4,4'-bis[(4-anilino-6-morpholino-s-triazin-2-yl)amino]stilbene-2,2'-disulfonate. Tinopal CBS is the disodium salt of 2,2'-[biphenyl-4,4'-di(2,1-ethenediyl)]dibenzene-1-sulfonate. Also preferred is the commercially available Parawhite KX, supplied by Paramount Minerals and Chemicals, Mumbai, India. Other fluorescers suitable for use in the invention include the 1-3-diarylpyrazolines and the 7-alkylaminocoumarins.

Suitable fluorescent brightener levels include lower levels of from about 0.01, from 0.05, from about 0.1 or even from about 0.2 wt % to upper levels of 0.5 or even 0.75 wt%.

The detergent compositions of the present invention (e.g., liquid or powder laundry compositions) may also include one or more soil-release polymers which aid the removal of soils from fabrics such as cotton and polyester-based fabrics, in particular the removal of hydrophobic soils from polyester-based fabrics. The soil release polymers may for example be nonionic or anionic terephthalate-based polymers, polyvinylcaprolactam and related copolymers, vinyl graft copolymers or polyester polyamides; see for example Chapter 7 in Powdered Detergents, Surfactant science series volume 71, Marcel Dekker, Inc. Another type of soil release polymers are amphiphilic alkoxyated grease cleaning polymers comprising a core structure and a plurality of alkoxyate groups attached to that core structure. The core structure may comprise a polyalkylenimine structure or a polyalkanolamine structure as described in detail in WO 2009/087523. Furthermore, random graft co-polymers are suitable soil-release polymers. Suitable graft co-polymers are described in more detail in WO 2007/138054, WO 2006/108856 and WO 2006/113314. Other soil-release polymers are substituted polysaccharide structures especially substituted cellulosic structures such as modified cellulose derivatives such as those described in EP 1867808 or WO 2003/040279. Suitable cellulosic polymers include cellulose, cellulose ethers, cellulose esters, cellulose amides and mixtures thereof. Suitable cellulosic polymers include anionically modified cellulose, nonionically modified cellulose, cationically modified cellulose, zwitterionically modified cellulose, and mixtures thereof.

The detergent compositions of the present invention (e.g., liquid or powder laundry compositions) may also include one or more anti-redeposition agents such as (carboxymethyl)

cellulose (CMC), poly(vinyl alcohol) (PVA), homopolymers of acrylic acid, copolymers of acrylic acid and maleic acid, and ethoxylated polyethyleneimines. The cellulose based polymers described under soil-release polymers above may also function as anti-redeposition agents.

5 The detergent composition of the invention (e.g., liquid or powder laundry compositions) may also contain one or more adjunct material. Suitable adjunct materials include, but are not limited to, anti-shrink agents, anti-wrinkling agents, bactericides, binders, carriers, dyes, enzyme stabilizers, fabric softeners, fillers, foam regulators, hydrotropes, perfumes, pigments, sod suppressors, solvents, and structurants for liquid detergents and/or structure elasticizing agents.

10 In addition to the proteases with the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 the detergents of the invention (e.g., liquid or powder laundry compositions) may further comprise cellulases. Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*,
15 e.g., the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum* disclosed in US 4,435,307, US 5,648,263, US 5,691,178, US 5,776,757 and WO 89/09259.

Especially suitable cellulases are the alkaline or neutral cellulases having color care benefits. Examples of such cellulases are cellulases described in EP 0 495 257, EP 0 531 372,
20 WO 96/11262, WO 96/29397, WO 98/08940. Other examples are cellulase polypeptides such as those described in WO 94/07998, EP 0 531 315, US 5,457,046, US 5,686,593, US 5,763,254, WO 95/24471, WO 98/12307 and PCT/DK98/00299.

Example of cellulases exhibiting endo-beta-1,4-glucanase activity (EC 3.2.1.4) are those having described in WO02/099091.

25 Other examples of cellulases include the family 45 cellulases described in WO96/29397, and especially polypeptides thereof having substitution, insertion and/or deletion at one or more of the positions corresponding to the following positions in SEQ ID NO: 8 of WO 02/099091: 2, 4, 7, 8, 10, 13, 15, 19, 20, 21, 25, 26, 29, 32, 33, 34, 35, 37, 40, 42, 42a, 43, 44, 48, 53, 54, 55, 58, 59, 63, 64, 65, 66, 67, 70, 72, 76, 79, 80, 82, 84, 86, 88, 90, 91, 93, 95, 95d, 95h, 95j, 97, 100,
30 101, 102, 103, 113, 114, 117, 119, 121, 133, 136, 137, 138, 139, 140a, 141, 143a, 145, 146, 147, 150e, 150j, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160c, 160e, 160k, 161, 162, 164, 165, 168, 170, 171, 172, 173, 175, 176, 178, 181, 183, 184, 185, 186, 188, 191, 192, 195, 196, 200, and/or 20, preferably selected among P19A, G20K, Q44K, N48E, Q119H or Q146 R.

Commercially available cellulases include Celluzyme™, Celluclean and Carezyme™
35 (Novozymes A/S), Clazinase™, and Puradax HA™ (Genencor International Inc.), and KAC-500(B)™ (Kao Corporation).

In addition to the proteases with the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or

SEQ ID NO 18 the detergents of the invention may further comprise additional proteases wherein the proteases include those of bacterial, fungal, plant, viral or animal origin e.g. vegetable or microbial origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. It may be an alkaline protease, such as a serine protease or a metalloprotease. A serine protease may for example be of the S1 family, such as trypsin, or the S8 family such as subtilisin. A metalloprotease protease may for example be a thermolysin from e.g. family M4 or other metalloprotease such as those from M5, M7 or M8 families.

Subtilases such as those derived from *Bacillus* such as *Bacillus lentus*, *B. alkalophilus*, *B. subtilis*, *B. amyloliquefaciens*, *Bacillus pumilus* and *Bacillus gibsonii* described in; US7262042 and WO09/021867, and *subtilisin lentus*, *subtilisin Novo*, *subtilisin Carlsberg*, *Bacillus licheniformis*, *subtilisin BPN'*, *subtilisin 309*, *subtilisin 147* and *subtilisin 168* described in WO89/06279 and protease PD138 described in (WO93/18140). Other useful proteases may be those described in WO92/175177, WO01/016285, WO02/026024 and WO02/016547. Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the *Fusarium* protease described in WO89/06270, WO94/25583 and WO05/040372, and the chymotrypsin proteases derived from *Cellulomonas* described in WO05/052161 and WO05/052146.

A further preferred protease is the alkaline protease from *Bacillus lentus* DSM 5483, as described for example in WO95/23221, and polypeptides thereof which are described in WO92/21760, WO95/23221, EP1921147 and EP1921148.

Examples of metalloproteases are the neutral metalloprotease as described in WO07/044993 (Genencor Int.) such as those derived from *Bacillus amyloliquefaciens*. Examples of useful proteases are the polypeptides described in: WO92/19729, WO96/034946, WO98/20115, WO98/20116, WO99/011768, WO01/44452, WO03/006602, WO04/03186, WO04/041979, WO07/006305, WO11/036263, WO11/036264.

Suitable commercially available protease enzymes include those sold under the trade names Alcalase®, Duralase™, Durazym™, Relase®, Relase® Ultra, Savinase®, Savinase® Ultra, Primase®, Polarzyme®, Kannase®, Liquanase®, Liquanase® Ultra, Ovozime®, Coronase®, Coronase® Ultra, Neutrase®, Everlase® and Esperase® (Novozymes A/S), those sold under the tradename Maxatase®, Maxacal®, Maxapem®, Purafect®, Purafect Prime®, Purafect MA®, Purafect Ox®, Purafect OxP®, Puramax®, Properase®, FN2®, FN3®, FN4®, Excellase®, Opticlean® and Optimase® (Danisco/DuPont), Axapem™ (Gist-Brocades N.V.), BLAP (sequence shown in Figure 29 of US5352604) and polypeptides hereof (Henkel AG) and KAP (*Bacillus alkalophilus* subtilisin) from Kao.

In addition to the proteases with the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 the detergents of the invention may further comprise lipases and cutinases which include those of bacterial or fungal origin. Chemically modified or protein engineered mutant enzymes are included. Examples include lipase from *Thermomyces*, e.g. from *T. lanuginosus*

(previously named *Humicola lanuginosa*) as described in EP258068 and EP305216, cutinase from *Humicola*, e.g. *H. insolens* (WO96/13580), lipase from strains of *Pseudomonas* (some of these now renamed to *Burkholderia*), e.g. *P. alcaligenes* or *P. pseudoalcaligenes* (EP218272), *P. cepacia* (EP331376), *P. sp.* strain SD705 (WO95/06720 & WO96/27002), *P. wisconsinensis* 5 (WO96/12012), GDSL-type *Streptomyces* lipases (WO10/065455), cutinase from *Magnaporthe grisea* (WO10/107560), cutinase from *Pseudomonas mendocina* (US5,389,536), lipase from *Thermobifida fusca* (WO11/084412), *Geobacillus stearothermophilus* lipase (WO11/084417), lipase from *Bacillus subtilis* (WO11/084599), and lipase from *Streptomyces griseus* (WO11/150157) and *S. pristinaespiralis* (WO12/137147).

10 Other examples are lipase polypeptides such as those described in EP407225, WO92/05249, WO94/01541, WO94/25578, WO95/14783, WO95/30744, WO95/35381, WO95/22615, WO96/00292, WO97/04079, WO97/07202, WO00/34450, WO00/60063, WO01/92502, WO07/87508 and WO09/109500.

Preferred commercial lipase products include Lipolase™, Lipex™, Lipolex™ and 15 Lipoclean™ (Novozymes A/S), Lumafast (originally from Genencor) and Lipomax (originally from Gist-Brocades).

Still other examples are lipases sometimes referred to as acyltransferases or perhydrolases, e.g. acyltransferases with homology to *Candida antarctica* lipase A (WO10/111143), acyltransferase from *Mycobacterium smegmatis* (WO05/56782), perhydrolases 20 from the CE 7 family (WO09/67279), and polypeptides of the *M. smegmatis* perhydrolase in particular the S54V variant used in the commercial product Gentle Power Bleach from Huntsman Textile Effects Pte Ltd (WO10/100028).

In addition to the proteases with the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or 25 SEQ ID NO 18 the detergents of the invention (e.g., liquid or powder laundry compositions) may further comprise amylases which can be used together with the proteases of the invention. The amylase may be an alpha-amylase or a glucoamylase and may be of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Amylases include, for example, alpha-amylases obtained from *Bacillus*, e.g., a special strain of *Bacillus licheniformis*, described 30 in more detail in GB 1,296,839.

Suitable amylases include amylases having SEQ ID NO: 3 in WO 95/10603 or polypeptides having 90% sequence identity to SEQ ID NO: 3 thereof. Preferred polypeptides are described in WO 94/02597, WO 94/18314, WO 97/43424 and SEQ ID NO: 4 of WO 99/019467, such as polypeptides with substitutions in one or more of the following positions: 15, 23, 105, 106, 35 124, 128, 133, 154, 156, 178, 179, 181, 188, 190, 197, 201, 202, 207, 208, 209, 211, 243, 264, 304, 305, 391, 408, and 444.

Different suitable amylases include amylases having SEQ ID NO: 6 in WO 02/010355 or polypeptides thereof having 90% sequence identity to SEQ ID NO: 6. Preferred polypeptides of

SEQ ID NO: 6 are those having a deletion in positions 181 and 182 and a substitution in position 193.

Other amylases which are suitable are hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO 2006/066594 and residues 36-483 of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4 of WO 2006/066594 or polypeptides having 90% sequence identity thereof. Preferred polypeptides of this hybrid alpha-amylase are those having a substitution, a deletion or an insertion in one or more of the following positions: G48, T49, G107, H156, A181, N190, M197, I201, A209 and Q264. Most preferred polypeptides of the hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO 2006/066594 and residues 36-483 of SEQ ID NO: 4 are those having the substitutions:

M197T;

H156Y+A181T+N190F+A209V+Q264S; or

G48A+T49I+G107A+H156Y+A181T+N190F+I201F+A209V+Q264S.

Further amylases which are suitable are amylases having SEQ ID NO: 6 in WO 99/019467 or polypeptides thereof having 90% sequence identity to SEQ ID NO: 6. Preferred polypeptides of SEQ ID NO: 6 are those having a substitution, a deletion or an insertion in one or more of the following positions: R181, G182, H183, G184, N195, I206, E212, E216 and K269. Particularly preferred amylases are those having deletion in positions R181 and G182, or positions H183 and G184.

Additional amylases which can be used are those having SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 2 or SEQ ID NO: 7 of WO 96/023873 or polypeptides thereof having 90% sequence identity to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7. Preferred polypeptides of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7 are those having a substitution, a deletion or an insertion in one or more of the following positions: 140, 181, 182, 183, 184, 195, 206, 212, 243, 260, 269, 304 and 476. More preferred polypeptides are those having a deletion in positions 181 and 182 or positions 183 and 184. Most preferred amylase polypeptides of SEQ ID NO: 1, SEQ ID NO: 2 or SEQ ID NO: 7 are those having a deletion in positions 183 and 184 and a substitution in one or more of positions 140, 195, 206, 243, 260, 304 and 476.

Other amylases which can be used are amylases having SEQ ID NO: 2 of WO 08/153815, SEQ ID NO: 10 in WO 01/66712 or polypeptides thereof having 90% sequence identity to SEQ ID NO: 2 of WO 08/153815 or 90% sequence identity to SEQ ID NO: 10 in WO 01/66712. Preferred polypeptides of SEQ ID NO: 10 in WO 01/66712 are those having a substitution, a deletion or an insertion in one or more of the following positions: 176, 177, 178, 179, 190, 201, 207, 211 and 264.

Further suitable amylases are amylases having SEQ ID NO: 2 of WO 09/061380 or polypeptides having 90% sequence identity to SEQ ID NO: 2 thereof. Preferred polypeptides of SEQ ID NO: 2 are those having a truncation of the C-terminus and/or a substitution, a deletion or

an insertion in one of more of the following positions: Q87, Q98, S125, N128, T131, T165, K178, R180, S181, T182, G183, M201, F202, N225, S243, N272, N282, Y305, R309, D319, Q320, Q359, K444 and G475. More preferred polypeptides of SEQ ID NO: 2 are those having the substitution in one of more of the following positions: Q87E,R, Q98R, S125A, N128C, T131I, T165I, K178L, T182G, M201L, F202Y, N225E,R, N272E,R, S243Q,A,E,D, Y305R, R309A, Q320R, Q359E, K444E and G475K and/or deletion in position R180 and/or S181 or of T182 and/or G183. Most preferred amylase polypeptides of SEQ ID NO: 2 are those having the substitutions:

N128C+K178L+T182G+Y305R+G475K;
10 N128C+K178L+T182G+F202Y+Y305R+D319T+G475K;
S125A+N128C+K178L+T182G+Y305R+G475K; or
S125A+N128C+T131I+T165I+K178L+T182G+Y305R+G475K wherein the polypeptides are C-terminally truncated and optionally further comprises a substitution at position 243 and/or a deletion at position 180 and/or position 181.

15 Other suitable amylases are the alpha-amylase having SEQ ID NO: 12 in WO01/66712 or a variant having at least 90% sequence identity to SEQ ID NO: 12. Preferred amylase polypeptides are those having a substitution, a deletion or an insertion in one of more of the following positions of SEQ ID NO: 12 in WO01/66712: R28, R118, N174; R181, G182, D183, G184, G186, W189, N195, M202, Y298, N299, K302, S303, N306, R310, N314; R320, H324, E345, Y396, R400, W439, R444, N445, K446, Q449, R458, N471, N484. Particular preferred
20 amylases include polypeptides having a deletion of D183 and G184 and having the substitutions R118K, N195F, R320K and R458K, and a variant additionally having substitutions in one or more position selected from the group: M9, G149, G182, G186, M202, T257, Y295, N299, M323, E345 and A339, most preferred a variant that additionally has substitutions in all these positions.

25 Other examples are amylase polypeptides such as those described in WO2011/098531, WO2013/001078 and WO2013/001087.

Commercially available amylases are DuramylTM, TermamylTM, FungamylTM, StainzymeTM, Stainzyme PlusTM, NatalaseTM, Liquozyme X and BANTM (from Novozymes A/S), and RapidaseTM, PurastarTM/EffectenzTM, Powerase and Preferenz S100 (from Genencor
30 International Inc./DuPont).

In addition to the proteases with the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 the detergents of the invention (e.g., liquid or powder laundry compositions) may further comprise peroxidases/oxidases including those of plant, bacterial or fungal origin.
35 Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinus*, e.g., from *C. cinereus*, and polypeptides thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257.

Commercially available peroxidases include GuardzymeTM (Novozymes A/S).

The detergent enzyme(s) may be included in a detergent composition by adding separate additives containing one or more enzymes, or by adding a combined additive comprising all of these enzymes. A detergent additive of the invention, *i.e.*, a separate additive or a combined additive, can be formulated, for example, as a granulate, liquid, slurry, etc. Preferred detergent
5 additive formulations are granulates, in particular non-dusting granulates, liquids, in particular stabilized liquids, or slurries.

Non-dusting granulates may be produced, *e.g.*, as disclosed in US 4,106,991 and 4,661,452 and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molar weights
10 of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591. Liquid enzyme preparations may, for instance, be
15 stabilized by adding a polyol such as propylene glycol, a sugar or sugar alcohol, lactic acid or boric acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

The detergent composition of the invention may be in any convenient form, *e.g.*, a bar, a homogenous tablet, a tablet having two or more layers, a pouch having one or more
20 compartments, a regular or compact powder, a granule, a paste, a gel, or a regular, compact or concentrated liquid.

Pouches can be configured as single or multicompartments. It can be of any form, shape and material which is suitable for hold the composition, *e.g.* without allowing the release of the composition to release of the composition from the pouch prior to water contact. The pouch is
25 made from water soluble film which encloses an inner volume. Said inner volume can be divided into compartments of the pouch. Preferred films are polymeric materials preferably polymers which are formed into a film or sheet. Preferred polymers, copolymers or derivatives thereof are selected polyacrylates, and water soluble acrylate copolymers, methyl cellulose, carboxy methyl cellulose, sodium dextrin, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose,
30 malto dextrin, poly methacrylates, most preferably polyvinyl alcohol copolymers and, hydroxypropyl methyl cellulose (HPMC). Preferably the level of polymer in the film for example PVA is at least about 60%. Preferred average molecular weight will typically be about 20,000 to about 150,000. Films can also be of blended compositions comprising hydrolytically degradable and water soluble polymer blends such as polylactide and polyvinyl alcohol (known under the
35 Trade reference M8630 as sold by MonoSol LLC, Indiana, USA) plus plasticisers like glycerol, ethylene glycerol, propylene glycol, sorbitol and mixtures thereof. The pouches can comprise a solid laundry cleaning composition or part components and/or a liquid cleaning composition or part components separated by the water soluble film. The compartment for liquid components

can be different in composition than compartments containing solids: US2009/0011970 A1.

Detergent ingredients can be separated physically from each other by compartments in water dissolvable pouches or in different layers of tablets. Thereby negative storage interaction between components can be avoided. Different dissolution profiles of each of the compartments
5 can also give rise to delayed dissolution of selected components in the wash solution.

A liquid or gel detergent, which is not unit dosed, may be aqueous, typically containing at least 20% by weight and up to 95% water, such as up to about 70% water, up to about 65% water, up to about 55% water, up to about 45% water, up to about 35% water. Other types of liquids, including without limitation, alkanols, amines, diols, ethers and polyols may be included in an
10 aqueous liquid or gel. An aqueous liquid or gel detergent may contain from 0-30% organic solvent.

The present invention is also directed to methods for using the polypeptides according to the invention or compositions thereof in laundering of textile and fabrics, such as house hold laundry washing and industrial laundry washing.

The invention is also directed to methods for using the polypeptides according to the
15 invention or compositions thereof in cleaning hard surfaces such as floors, tables, walls, roofs etc. as well as surfaces of hard objects such as cars (car wash) and dishes (dish wash).

The polypeptides of the present invention may be added to and thus become a component of a detergent composition. Thus, one aspect of the invention relates to the use of a polypeptide comprising the SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ
20 ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18 in a cleaning process such as laundering and/or hard surface cleaning.

Thus, one aspect of the invention relates to the use of the use of a polypeptide comprising the SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18 in a cleaning process such as laundering
25 and/or hard surface cleaning and wherein the polypeptide has improved wash performance, relative to a reference enzyme such as a protease with SEQ ID NO 19.

The cleaning process or the textile care process may for example be a laundry process, a dishwashing process or cleaning of hard surfaces such as bathroom tiles, floors, table tops, drains, sinks and washbasins. Laundry processes can for example be household laundering, but
30 it may also be industrial laundering. Furthermore, the invention relates to a process for laundering of fabrics and/or garments where the process comprises treating fabrics with a washing solution containing a detergent composition, and at least one protease variant of the invention. The cleaning process or a textile care process can for example be carried out in a machine washing process or in a manual washing process. The washing solution can for example be an aqueous
35 washing solution containing a detergent composition.

The last few years there has been an increasing interest in replacing components in detergents, which is derived from petrochemicals with renewable biological components such as enzymes and polypeptides without compromising the wash performance. When the components

of detergent compositions change new enzyme activities or new enzymes having alternative and/or improved properties compared to the commonly used detergent enzymes such as proteases, lipases and amylases are needed to achieve a similar or improved wash performance when compared to the traditional detergent compositions.

5 The invention further concerns the use of polypeptides of the invention in a proteinaceous stain removing processes. The proteinaceous stains may be stains such as food stains, e.g., baby food, sebum, cocoa, egg, blood, milk, ink, grass or a combination hereof, and more specific such as egg yolk, minced meat, Crème Brûlée, pasta, burnt-in milk. The protease comprising the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ
10 ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 and in particular the proteases comprising SEQ ID NO 2 and SEQ ID NO 4 are active on blood stains in particular aged blood and heat treated blood stains such as blood/milk/ink. Blood stains are particular resistant stain to remove and on the same time commercially very relevant e.g. in industrial cleaning. Often blood soiled textiles are not immediately washed and being exposed to elevated temperatures thus
15 wash performance or stain removal properties on such stains are advantageous in particular in textile cleaning. One aspect of the invention relates to a protease comprising the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 wherein the protease has improved wash performance on blood stains compared to the protease with SEQ ID NO 19. A particular
20 aspect of the invention relates to the use of a protease comprising or consisting of the amino acid sequences SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 for removal of blood stains from textiles.

 In a particular embodiment, the invention concerns the use of a composition comprising the polypeptides of the invention and one or more detergent components, such as surfactants,
25 hydrotropes, builders, co-builders, chelators or chelating agents, bleaching system or bleach components, polymers, fabric hueing agents, fabric conditioners, foam boosters, suds suppressors, dispersants, dye transfer inhibitors, fluorescent whitening agents, perfume, optical brighteners, bactericides, fungicides, soil suspending agents, soil release polymers, anti-redeposition agents, enzyme inhibitors or stabilizers, enzyme activators, antioxidants, and
30 solubilizers.

 In a particular embodiment, the invention concerns the use of a composition comprising a subtilase variant of the invention and one or more additional enzymes selected from the group consisting of proteases, amylases, lipases, cutinases, cellulases, endoglucanases, xyloglucanases, pectinases, pectin lyases, xanthanases, peroxidaes, haloperoxygenases,
35 catalases and mannanases, or any mixture thereof.

 In a particular embodiment, the invention concerns the use of a composition comprising a subtilase variant of the invention, one or more additional enzymes selected from the group consisting of proteases, amylases, lipases, cutinases, cellulases, endoglucanases,

xyloglucanases, pectinases, pectin lyases, xanthanases, peroxidaes, haloperoxygenases, catalases and mannanases, or any mixture thereof and one or more detergent components, such as surfactants, hydrotropes, builders, co-builders, chelators or chelating agents, bleaching system or bleach components, polymers, fabric hueing agents, fabric conditioners, foam boosters, suds suppressors, dispersants, dye transfer inhibitors, fluorescent whitening agents, perfume, optical brighteners, bactericides, fungicides, soil suspending agents, soil release polymers, anti-redeposition agents, enzyme inhibitors or stabilizers, enzyme activators, antioxidants, and solubilizers.

Washing Method

10 The present invention also relates to a method of cleaning a fabric, a dishware or hard surface with a detergent composition comprising a polypeptide of the invention.

 A preferred embodiment concerns a method of cleaning, said method comprising the steps of: contacting an object with a detergent composition comprising a polypeptide of the invention under conditions suitable for cleaning said object. In a preferred embodiment the
15 detergent composition is used in a laundry or a dish wash process.

 Still another embodiment relates to a method for removing stains from fabric or dishware which comprises contacting said fabric or dishware with a composition comprising a polypeptide of the invention under conditions suitable for cleaning said object.

 Also contemplated are compositions and methods of treating fabrics (e.g., to desize a textile) using one or more of the proteases of the invention. The protease can be used in any
20 fabric-treating method which is well known in the art (see, e.g., US6,077,316). For example, in one aspect, the feel and appearance of a fabric is improved by a method comprising contacting the fabric with a protease in a solution. In one aspect, the fabric is treated with the solution under pressure.

25 The detergent compositions of the present invention are suited for use in laundry and hard surface applications, including dish wash. Accordingly, the present invention includes a method for laundering a fabric or washing dishware. The method comprises the steps of contacting the fabric/dishware to be cleaned with a solution comprising the detergent composition according to the invention. The fabric may comprise any fabric capable of being laundered in normal consumer
30 use conditions. The dishware may comprise any dishware such as crockery, cutlery, ceramics, plastics such as melamine, metals, china, glass and acrylics. The solution preferably has a pH from about 5.5 to about 11.5. The compositions may be employed at concentrations from about 100 ppm, preferably 500 ppm to about 15,000 ppm in solution. The water temperatures typically range from about 5°C to about 95°C, including about 10°C, about 15°C, about 20°C, about 25°C,
35 about 30°C, about 35°C, about 40°C, about 45°C, about 50°C, about 55°C, about 60°C, about 65°C, about 70°C, about 75°C, about 80°C, about 85°C and about 90°C.

Variations in local and regional conditions, such as water hardness and wash temperature call for regional detergent compositions. Table 1 provides ranges for the composition of a typical European automatic dish wash (ADW) detergent.

Table 1: Typical European ADW detergent composition

P-Containing formulation	P-Free formulations
20-50% STPP	10-20% Na Citrate (or Chelating agent)
15-45% Soda (sodium carbonate)	25% Soda (sodium carbonate)
5-15% Sodium percarbonate	5-10% Sodium percarbonate
0-20% Sodium disilicate	5-25% Sodium disilicate
2-3% TAED	0-3% TAED
2-6% Polymers	2-6% Polymers
1-2% Phosphonate	1-20% Sodium sulfate
3-5% Surfactants	<5% Surfactants
<5% Enzymes	<5% Enzymes
To 100% Rest (perfume, dye, corrosion inhibitor. etc.) pH 9-11	To 100% Rest (perfume, dye, corrosion inhibitor. etc.) pH 9-11


5

The polypeptides of the detergent composition of the invention may be stabilized using conventional stabilizing agents and protease inhibitors, e.g. a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, different salts such as NaCl; KCl; lactic acid, formic acid, boric acid, or a boric acid derivative, e.g., an aromatic borate ester, or a phenyl boronic acid derivative such as 4-formylphenyl boronic acid, or a peptide aldehyde such as di-, tri- or tetrapeptide aldehydes or aldehyde analogues (either of the form B1-B0-R wherein, R is H, CH₃, CX₃, CHX₂, or CH₂X (X=halogen), B0 is a single amino acid residue (preferably with an optionally substituted aliphatic or aromatic side chain); and B1 consists of one or more amino acid residues (preferably one, two or three), optionally comprising an N-terminal protection group, or as described in WO09118375, WO98/13459) or a protease inhibitor of the protein type such as RASI, BASI, WASI (bifunctional alpha-amylase/subtilisin inhibitors of rice, barley and wheat) or CI2 or SSI. The composition may be formulated as described in e.g. WO 92/19709, WO 92/19708 and US6,472,364. In some embodiments, the enzymes employed herein are stabilized by the presence of water-soluble sources of zinc (II), calcium (II) and/or magnesium (II) ions in the finished compositions that provide such ions to the enzymes, as well as other metal ions (e.g., barium (II), scandium (II), iron (II), manganese (II), aluminum (III), Tin (II), cobalt (II), copper (II), Nickel (II), and oxovanadium (IV)).

In one embodiment, the polypeptides are stabilized using peptide aldehydes or ketones

Suitable peptide aldehydes are described in WO94/04651, WO95/25791, WO98/13458, WO98/13459, WO98/13460, WO98/13461, WO98/13462, WO07/141736, WO07/145963, WO09/118375, WO10/055052 and WO11/036153. A polypeptide of the present invention may also be incorporated in the detergent formulations disclosed in WO 97/07202.

- 5 In another embodiment, the polypeptides are stabilized using a phenyl boronic acid derivative is 4-formylphenylboronic acid (4-FPBA) with the following formula:



- 10 The detergent compositions may comprise two or more stabilizing agents e.g. such as those selected from the group consisting of propylene glycol, glycerol, 4-formylphenyl boronic acid and borate.

The stabilizing agent(s) is preferably present in the detergent composition in a quantity of from 0.001 to about 5.0 wt%, from 0.01 to about 2.0 wt%, from 0.1 to about 3 wt% or from 0.5% to about 1.5 wt%.

- 15 In some preferred embodiments, the detergent compositions provided herein are typically formulated such that, during use in aqueous cleaning operations, the wash water has a pH of from about 5.0 to about 11.5, or in alternative embodiments, even from about 6.0 to about 10.5. In some preferred embodiments, granular or liquid laundry products are formulated to have a pH from about 6 to about 8. Techniques for controlling pH at recommended usage levels include the
20 use of buffers, alkalis, acids, etc., and are well known to those skilled in the art.

The present invention is further described by the following examples that should not be construed as limiting the scope of the invention.

The invention is further defined in the following paragraphs:

1. A dish wash composition comprising:
- 25 a) at least 0.01 mg (e.g., at least 0.02 mg, at least 0.03 mg, at least 0.04 mg, at least 0.05 mg, at least 0.06 mg, at least 0.07 mg, at least 0.08 mg, at least 0.09 mg, at least 0.1 mg, at least 0.2 mg, at least 0.3 mg, at least 0.4 mg, at least 0.5 mg, at least 0.6 mg, at least 0.7 mg, at least 0.8 mg, at least 0.9 mg, at least 1 mg) of active protease per gram of composition, wherein the protease is selected from a polypeptide comprising SEQ ID
30 NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) at least one builder, and
- c) at least one bleach component.
- 35 2. The dish wash composition of paragraph 1, wherein said polypeptide comprises an

- alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein
- 5 said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
- 10 3. The dish wash composition of any of paragraphs 1-2, wherein said polypeptide comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A,
- 15 R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
- 20 4. The dish wash composition of any of paragraphs 1-3, wherein said polypeptide consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.
- 25 5. The dish wash composition of any of paragraphs 1-4, wherein the dish wash composition is an automatic dish wash (ADW) composition or a hand wash (HDW) composition.
- 30 6. The dish wash composition according to any of paragraphs 1-5, wherein the builder is selected among phosphates, sodium citrate builders, sodium carbonate, sodium silicate, sodium and zeolites.
- 35 7. The dish wash composition according to any of paragraphs 1-6, wherein the builder is added in an amount of about 0-65% by weight, preferably about 40-65% by weight, particularly about 20-65% by weight, particularly from 10% to 50% by weight.
8. The dish wash composition according to any of paragraphs 1-7, wherein the dish wash

composition contains 0-50% by weight, such as about 5% to about 30%, of a detergent co-builder.

- 5 9. The dish wash composition according to any of the preceding paragraphs, wherein the wash composition is phosphate free.
- 10 10. The dish wash composition according to any of the preceding paragraphs, wherein the builder is selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof.
- 15 11. The dish wash composition according to any of the preceding paragraphs, wherein the bleach component is selected from bleaching catalysts, photobleaches, bleach activators, sources of hydrogen peroxide such as sodium percarbonate, sodium perborates and hydrogen peroxide, preformed peracids and mixtures thereof.
- 20 12. The dish wash composition according to any of the preceding paragraphs, wherein the bleach component is a peroxide such as percarbonate, persulfate, perphosphate, persilicate salts.
- 25 13. The dish wash composition according to any of the preceding paragraphs, wherein the bleaching component includes a percarbonate and bleach catalyst, preferably a manganese compound.
- 30 14. The dish wash composition of paragraph 13, wherein the bleach catalyst is 1,4,7-trimethyl-1,4,7-triazacyclononane or manganese (II) acetate tetrahydrate (MnTACN).
- 35 15. The dish wash composition according to any of the preceding paragraphs comprising from 1-40 wt%, preferably from 0.5-30 wt%, of bleaching components, wherein the bleach components are a peroxide, preferably percarbonate and a catalyst preferably a metal-containing bleach catalyst such as 1,4,7-trimethyl-1,4,7-triazacyclononane or manganese (II) acetate tetrahydrate (MnTACN).
16. A granular detergent composition comprising:
 - a) at least 0.01 mg (e.g., at least 0.02 mg, at least 0.03 mg, at least 0.04 mg, at least 0.05 mg, at least 0.06 mg, at least 0.07 mg, at least 0.08 mg, at least 0.09 mg, at least 0.1 mg, at least 0.2 mg, at least 0.3 mg, at least 0.4 mg, at least 0.5 mg, at least 0.6 mg, at least 0.7 mg, at least 0.8 mg, at least 0.9 mg, at least 1 mg) of active protease per gram of composition wherein the protease is selected from a polypeptide comprising SEQ ID

- NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18,
- b) 5 wt % to 50 wt % (e.g., 5 wt % to 40 wt %, 5 wt % to 30 wt %, 5 wt % to 20 wt %, 5 wt % to 10 wt %) anionic surfactant,
- 5 c) 1 wt % to 8 wt % (e.g., 1 wt % to 7 wt %, 1 wt % to 6 wt %, 1 wt % to 5 wt %, 1 wt % to 4 wt %, 1 wt % to 3 wt %, 1 wt % to 2 wt %) nonionic surfactant,
- d) 5 wt % to 40 wt % (e.g., 5 wt % to 30 wt %, 5 wt % to 20 wt %, 5 wt % to 10 wt %) builder such as carbonates, zeolites, phosphate builder, calcium sequestering builders or complexing agents,
- 10 preferably said granular detergent composition further comprises a bleach component.
17. The granular detergent composition of paragraph 16, wherein said polypeptide comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97,
- 15 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less
- 20 than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
18. The granular detergent composition of any of paragraphs 16-17, wherein said polypeptide comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A,
- 25 N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least
- 30 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
19. The granular detergent composition of any of paragraphs 16-18, wherein said polypeptide consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.
- 35

20. The granular detergent composition of any of paragraphs 16-19, wherein the anion surfactant is one or more selected from linear alkylbenzenesulfonates (LAS) isomers of LAS, alcohol ether sulfate (AEO, AEOS) and sodium lauryl ether sulfate, sodium laureth sulfate (SLES).
- 5 21. The granular detergent composition of any of paragraphs 16-20, wherein the nonionic surfactant is selected from nonionic surfactants primary alcohol ethoxylates, such as Neodol 25-7 or Neodol 25-3, alkyl ester sulphates, especially C-9-15 alcohol ethersulfates (AES), C12-15 primary alcohol ethoxylate such as alcohol ethoxylates (AE or AEO), sodium
10 toluene sulfonate (STS) and sodium dodecyl sulfate (SDS).
22. The granular detergent composition according to and of paragraphs 16-21, wherein the builder is a carbonate, zeolites builder.
- 15 23. The granular detergent composition according to any of paragraphs 16-22, wherein the builder is selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof.
- 20 24. The granular detergent composition according to paragraphs 16-23, which additionally comprises a bleach component selected from a peroxide such as percarbonate, persulfate, perphosphate, persilicate salts.
- 25 25. A detergent composition comprising:
 - a) at least 0.01 mg (e.g., at least 0.02 mg, at least 0.03 mg, at least 0.04 mg, at least 0.05
25 mg, at least 0.06 mg, at least 0.07 mg, at least 0.08 mg, at least 0.09 mg, at least 0.1 mg, at least 0.2 mg, at least 0.3 mg, at least 0.4 mg, at least 0.5 mg, at least 0.6 mg, at least 0.7 mg, at least 0.8 mg, at least 0.9 mg, at least 1 mg) of active protease per litre detergent wherein the protease is selected from a polypeptide comprising SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO
30 14, SEQ ID NO 16 or SEQ ID NO 18,
 - b) 2 wt% to 60 wt% of at least one surfactant and
 - c) 5 wt% to 50 wt% of at least one builder,preferably said detergent composition further comprises a bleach component.
- 35 26. The detergent composition of paragraph 25, wherein said polypeptide comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193,

- 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
- 5
27. The detergent composition of any of paragraphs 25-26, wherein said polypeptide comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
- 10
- 15
28. The detergent composition of any of paragraphs 25-27, wherein said polypeptide consists of the amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.
- 20
29. The detergent composition of any of paragraphs 25-28, wherein the detergent composition is a liquid- or powder laundry detergent composition.
- 25
30. The detergent composition of any of paragraphs 25-29, wherein the surfactant is selected from the group consisting of: nonionic, anionic and/or amphoteric surfactants.
- 30
31. A composition according to any of the preceding paragraphs wherein the polypeptides selected from the group consisting of SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO 6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 or SEQ ID NO 18 have improved wash performance on blood stains such as C-05 and CS-01 when compared to the polypeptide with SEQ ID NO 19, preferably said composition further comprises a bleach component.
- 35
32. A polypeptide having protease activity, said polypeptide comprising an amino acid sequence selected from the group consisting of: SEQ ID NO 2, SEQ ID NO 4, SEQ ID NO

6, SEQ ID NO 8, SEQ ID NO 10, SEQ ID NO 12, SEQ ID NO 14, SEQ ID NO 16 and SEQ ID NO 18.

- 5
33. A composition comprising a polypeptide of paragraph 32.
34. The composition of paragraph 33 or the polypeptide of paragraph 32, wherein said polypeptide comprises an alteration (e.g., a substitution, insertion or deletion) at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
- 15
35. The composition of any of paragraphs 33-34 or the polypeptide of paragraph 32 or paragraph 34, wherein said polypeptide comprises one or more of the alterations (e.g., a substitution, insertion or deletion) selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, such as at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
- 20
- 25
- 30 36. The composition of any of paragraphs 33-35, wherein said composition is selected from the group consisting of: a dish wash composition, an automatic dish wash composition, a hand wash composition, a detergent composition, a granular detergent composition, a liquid detergent composition, a powder detergent composition.
- 35 37. Use of a composition or a polypeptide according to any of the preceding paragraphs in a cleaning process, wherein the process is selected from a process of doing hard surface cleaning, such as dish wash, and a laundry process.

38. The use of paragraph 37, for removal of blood stains, preferably said blood stains are aged and/or heated blood stains.

39. A method for removing a stain from a surface which comprises contacting the surface with a composition or a polypeptide according to any of the preceding paragraphs.

EXAMPLES

Materials and Methods

10 Protease Assay (Suc-AAPF-pNA assay)

pNA substrate : Suc-AAPF-pNA (Bachem L-1400).

Temperature : Room temperature (25°C)

Assay buffer : 100mM succinic acid, 100mM HEPES, 100mM CHES, 100mM CABS, 1mM CaCl₂, 150mM KCl, 0.01% Triton X-100, pH 9.0.

15

20µl protease (diluted in 0.01% Triton X-100) was mixed with 100µl assay buffer. The assay was started by adding 100µl pNA substrate (50mg dissolved in 1.0ml DMSO and further diluted 45x with 0.01% Triton X-100). The increase in OD₄₀₅ was monitored as a measure of the protease activity.

20 **Example 1: Preparation and Expression of polypeptides**

The following summarizes the mutation and introduction of an expression cassette into *Bacillus subtilis*. All DNA manipulations were done by PCR (e.g. Sambrook et al.; Molecular Cloning; Cold Spring Harbor Laboratory Press) and can be repeated by everybody skilled in the art.

25 Recombinant *B. subtilis* constructs encoding subtilase polypeptides were used to inoculate shakeflasks containing a rich media (e.g. PS-1: 100 g/L Sucrose (Danisco cat.no. 109-0429), 40 g/L crust soy (soy bean flour), 10g/L Na₂HPO₄.12H₂O (Merck cat.no. 6579), 0.1ml/L replace- Dowfax63N10 (Dow). Cultivation typically takes 4 days at 30°C shaking with 220 rpm.

Example 2: Fermentation of polypeptides

30 Fermentation may be performed by methods well known in the art or as follows. A *B. subtilis* strain harboring the relevant expression plasmid was streaked on a LB agar plate, and grown overnight at 37°C. The colonies were transferred to 100 ml PS-1 media in a 500 ml shaking flask. Cells and other undissolved material were removed from the fermentation broth by centrifugation at 4500 rpm for 20-25 minutes. Afterwards the supernatant was filtered to obtain a
35 clear solution.

Example 3: Purification of Polypeptides

The culture broth was centrifuged (26000 x g, 20 min) and the supernatant was carefully decanted from the precipitate. The supernatant was filtered through a Nalgene 0.2µm filtration unit in order to remove the rest of the *Bacillus* host cells. pH in the 0.2µm filtrate was adjusted to pH 8 with 3M Tris base and the pH adjusted filtrate was applied to a MEP Hypercel column (from Pall corporation) equilibrated in 20mM Tris/HCl, 1mM CaCl₂, pH 8.0. After washing the column with the equilibration buffer, the column was step-eluted with 20mM CH₃COOH/NaOH, 1mM CaCl₂, pH 4.5. Fractions from the column were analysed for protease activity (using the Suc-AAPF-pNA assay at pH 9) and peak-fractions were pooled. The pH of the pool from the MEP Hypercel column was adjusted to pH 6 with 20%(v/v) CH₃COOH or 3M Tris base and the pH adjusted pool was diluted with deionized water to the same conductivity as 20mM MES/NaOH, 2mM CaCl₂, pH 6.0. The diluted pool was applied to a SP-sepharose FF column (from GE Healthcare) equilibrated in 20mM MES/NaOH, 2mM CaCl₂, pH 6.0. After washing the column with the equilibration buffer, the protease was eluted with a linear NaCl gradient (0 → 0.5M) in the same buffer over five column volumes. Fractions from the column were analysed for protease activity (using the Suc-AAPF- pNA assay at pH 9) and active fractions were analysed by SDS-PAGE. The fractions, where only one band was seen on the coomassie stained SDS-PAGE gel, were pooled as the purified preparation and was used for further experiments.

Example 4: In Wash Stability

20 In Wash Stability Assay

In wash stability was measured according to the 'In wash stability assay' as described below using the MGDA and STPP model detergents as defined in table 2. The results are presented in table 2.

Table 2: Composition of the MGDA and STPP model detergents

Component	MGDA model detergent	STPP model detergent
MGDA (40% solution)	1.67 g/l	
STPP		1.65 g/l
Sodium carbonate	0.66 g/l	0.66 g/l
Sodium percarbonate (Dream)	0.33 g/l	0.33 g/l
Sodium disilicate	0.17 g/l	0.17 g/l
TAED (Dream)	0.10 g/l	0.10 g/l
Sokalan CP5 (39.5%)	0.42 g/l	0.42 g/l
Surfac 23-6.5 (100%)	0.17 g/l	0.17 g/l
Sodium sulphate	1.06 g/l	

Phosphonate (tetrasodium HEDP)		0.07 g/l
CaCl ₂	3 mM	3 mM
MgCl ₂	0.75 mM	0.75 mM
NaHCO ₃	7.5 mM	7.5 mM
pH	10.0	10.0

Both detergents are dissolved in 50 mM CHES buffer (N-Cyclohexyl-2-aminoethanesulfonic acid, Sigma C2885) to ensure that pH is maintained during the experiment at 10.0 also after addition of protease sample.

5 Protease culture supernatants are pre-diluted 2-4 times and purified protease samples are diluted to approximately 0.1 and 0.05 mg/ml using deionized water. 10 µl diluted protease sample is then mixed with 190 µl model detergent solution in a well of a 0.2 ml 96-well PCR plate. After mixing, 20 µl is transferred to a 96-well microtiter plate (Nunc F) and initial protease activity is measured by adding 100 µl Suc-AAPF-pNA substrate solution (0.72 mg/ml Suc-Ala-Ala-Pro-Phe-
10 pNA (Bachem L-1400) in 0.1 M Tris, pH 8.6) to each well, mixing and measuring absorbance at 405 nm every 20 s for 5 min on a SpectraMax Plus (Molecular Devices). Slope from linear regression on initial absorbance measurements is used for activity calculations.

The proteases in the PCR plate are then stressed by 30 min incubation at 58°C for STPP model detergent and 60 or 62°C for MGDA model detergent in a BioRad T100 Thermal Cycler.
15 After rapid cooling to room temperature, 20 µl is transferred to a 96 well microtiter plate and residual activity is measured as described for the initial protease activity. The temperatures in the stress step are chosen to give suitable residual activities of the Savinase reference (polypeptide of SEQ ID NO 19) and the polypeptides (preferably in the interval 10 to 80% of the initial activity).

The decrease in activity during the stress step is assumed to be exponential. Thus, the
20 half-life during the stress step is calculated using the formula:

$$T_{1/2} = T * \ln(2) / \ln(A(\text{Initial}) / A(\text{Residual}))$$

where $T_{1/2}$ is the half-life, T is the incubation time (30 min), A(Initial) is the initial protease
25 activity, and A(Residual) is the protease activity after the stress step. All protease samples are tested twice (using 2 times the same sample dilution for culture supernatants and 0.1 and 0.05 mg/ml for purified protease samples). Relative in wash stability improvement factor is then calculated by:

30 Relative In Wash Stability Improvement Factor = $\text{Avg}(T_{1/2}(\text{Sample})) / \text{Avg}(T_{1/2}(\text{Reference}))$

where $\text{Avg}(T_{1/2}(\text{Samples}))$ is the average of the half-lives for the given protease sample and

Avg($T_{1/2}$ (Reference)) is the average of the half-lives for the Savinase reference sample (SEQ ID NO 19).

The results show that the proteases with SEQ ID NO 4, 6, 8 and 10 are more stable than Savinase (SEQ ID NO 19).

5

Table 3: In Wash Stability Data using MGDA and STPP model detergents

Polypeptides	Relative Half Life ($T_{1/2}$)	
	STPP	MGDA
SEQ ID NO 10	3.08	2.93
SEQ ID NO 8	2.60	1.60
SEQ ID NO 6	1.92	1.13
SEQ ID NO 4	1.44	0.97

Example 5: Wash performance in Automatic Mechanical Stress Assay (AMSA) for Automatic Dish Wash

10 Washing experiments are performed in order to assess the wash performance of selected protease polypeptides in dish wash detergent compositions. The proteases of the present application are tested using the Automatic Mechanical Stress Assay (AMSA). With the AMSA, the wash performance of many small volume enzyme-detergent solutions can be examined. The AMSA plate has a number of slots for test solutions and a lid that firmly squeezes the melamine tile to be washed against the slot openings. During the wash, the plate, test solutions, melamine
15 tile and lid are vigorously shaken to bring the test solution in contact with the soiled melamine tile and apply mechanical stress in a regular, periodic oscillating manner. For further description see WO 02/42740 especially the paragraph "Special method embodiments" at page 23-24.

The experiment is conducted under the experimental conditions as specified in tables 4 and 5 below.

20

Table 4: AMSA Experimental Conditions using ADW model detergent with MGDA

ADW model detergent with MGDA	As defined in table 2
Detergent dosage	3.33 g/L
Test solution volume	160 micro L
pH	10,0
Wash time	20 minutes
Temperature	45 °C
Water hardness	21 °dH

Enzyme concentration in test solution	5.3, 10,7 mg enzyme protein/liter
Test material	Egg yolk melamine tile (DM-21)

Table 5: AMSA Experimental Conditions using ADW model detergent with STPP

ADW model detergent with STPP	As defined in table 2
Detergent dosage	3.33 g/L
Test solution volume	160 micro L
pH	10,0
Wash time	20 minutes
Temperature	45°C
Water hardness	21°dH
Enzyme concentration in test solution	5.3, 10,7 mg enzyme protein/liter
Test material	Egg yolk melamine tile (DM-21)

Water hardness is adjusted to 21°dH by addition of CaCl_2 , MgCl_2 , and NaHCO_3 ($\text{Ca}^{2+}:\text{Mg}^{2+}:\text{CO}_3^{2-} = 4:1:10$) to the test system. After washing the egg yolk melamine tiles are flushed in tap water and dried.

The performance of the enzyme variant is measured as the brightness of the colour of the melamine tile washed with that specific protease. Brightness can also be expressed as the intensity of the light reflected from the sample when illuminated with white light. When the sample is stained the intensity of the reflected light is lower, than that of a clean sample. Therefore, the intensity of the reflected light can be used to measure wash performance of a protease. Color measurements were made with a professional flatbed scanner (EPSON EXPRESSION 10000XL, Atea A/S, Lautrupvang 6, 2750 Ballerup, Denmark), which is used to capture an image of the washed melamine tiles.

To extract a value for the light intensity from the scanned images, 24-bit pixel values from the image are converted into values for red, green and blue (RGB). The intensity value (Int) is calculated by adding the RGB values together as vectors and then taking the length of the resulting vector:

$$Int = \sqrt{r^2 + g^2 + b^2}$$

Stains

Standard egg yolk melamine tiles (DM-21) were obtained from Center For Testmaterials BV, P.O. Box 120, 3133 KT Vlaardingen, the Netherlands.

The results are shown in table 6 below. The proteases with SEQ ID NO 4, 6 and 12 have

improved was performance in dish wash compared to the protease with SEQ ID NO 19.

Table 6

Polypeptides	Relative Performance AMSA	
	STPP	MGDA
SEQ ID NO 6	1.1	1.4
SEQ ID NO 4	1.2	1.1
SEQ ID NO 12	1.1	1.0

Example 6: Wash performance evaluation of SEQ ID NO 2 and SEQ ID NO 4 in powder laundry detergent

The proteases with SEQ ID NO 2 and 4 were investigated in full scale wash. Wash performance was tested in powder model detergent with percarbonate/TEAD and compared to the known laundry protease Savinase (SEQ ID NO 19). The results are shown in Table 9.

Wash performance was evaluated using the bleach containing laundry powder model detergent as defined in table 7 below and the experimental conditions defined in table 8.

Table 7. Composition of Model detergent T (powder) wt %	
LAS, sodium salt	10.0
AS, sodium salt	1.8
Soap, sodium salt	2.0
AEO	3.0
Soda ash	14.9
Hydrous sodium silicate	2.5
Zeolite A	15.0
HEDP-Na4	0.13
Sodium citrate	2.0
PCA, copoly(acrylic acid/maleic acid), sodium salt	1.5
SRP	0.5
Sodium sulfate	13.4
Sodium percarbonate	20.0

TAED	3.0
Foam regulator	1.0

CMC is added separately at 1.28 g/wash (around 13L water intake)

Table 8: Experimental condition for FSW

Machine	Miele Softtronic W2245 FLA
FSW Method	EU standard wash
Detergent	Model powder detergent with bleach (see table 5)
Detergent conc.	80 g Model T + 1.28 g CMC
Temperature	40 °C
Enzyme dosages	30 nM (0.10 wt% of SEQ ID NO 2 and 0.12 wt% of SEQ ID NO 4) 60 nM (0.20 wt% of SEQ ID NO 2 and 0.24 wt% of SEQ ID NO 4) 720 nM (2.38 wt% of SEQ ID NO 2 and 2.94 wt% of SEQ ID NO 4)
Water hardness	15°dH (Ca/Mg/HCO ₃ = 4:1:7.5)
Wash time	41 min main wash
Enzymes	Savinase (SEQ ID NO 19), SEQ ID NO 2 and SEQ ID NO 4
Stain/ swatch	C-05 Blood/milk/ink CS-01 Aged blood
Ballast	3kg mixed cotton/polyester (ratio 65:35)
Soil ballast	2 pieces SBL2004 and other enzyme sensitive stains like PC-03 (Chocolate milk/soot), C-H097 (Oatmeal/Chocolate) EMPA 112 (Cocoa), C-H010 (Cocoa cocked up with milk), CS-44 (Chocolate drink), WFK 10Z (Chocolate), CS-37 (Full egg/Pigment), WFK10N (Whole Egg), WFK10EG (Egg Yolk), 051KC (Egg), EMPA 164 (Grass), 062KC (Scrubbed Grass), WE5GMWKC (Grass/Mud), C-10 (Milk/Oil/Pigment) C-H165 (Minced meat) and KH-H172 (Meat Pate)

5

Test materials, were obtained from Center For Testmaterials BV, P.O. Box 120, 3133 KT Vlaardingen, the Netherlands, SBL2004 and the other enzyme sensitive stains were obtained either from Center For Testmaterials BV, P.O. Box 120, 3133 KT Vlaardingen, the Netherlands, or EMPA Testmaterials AG, Möwenstrasse 12, CH-9015 St. Gallen, Switzerland or WFK Testgewebe GmbH, Christenfeld 10, D-41379 Brüggen, Germany.

10

Wash performance was measured as delta remission on swatches, which after wash were

dried overnight on tray (12h-20h) at room (around 20-25C) temperature and 40% humidity, then measured at 460 nm in Coloreye equipment stacked 2 pieces in the second day.

Table 9 shows the performance as delta remission of the proteases with SEQ ID NO 2 and 4 on two different types of blood stains.

5

Table 9

	30 nM		60 nM		720 nM	
Polypeptides	SEQ ID NO 2	SEQ ID NO 4	SEQ ID NO 2	SEQ ID NO 4	SEQ ID NO 2	SEQ ID NO 4
Heat treated C-05 Blood/milk/ink	4.16	4.25	4.05	4.63	2.93	3.40
CS-01 Aged blood	3.46	5.39	5.30	7.29	8.98	12.94

The data clearly show an improved wash performance of protease with SEQ NO 2 and SEQ NO 4 relative to Savinase (SEQ ID NO 19) on the aged/heat treated blood stains C-05 and CS-01.

10 **Example 7 Performance of polypeptides on blood stains**

Detergent Model X pH profile (on EMPA117EH)

The assay was performed in a 96 well microtiter plate 96 with two EMPA117 EH (blood/milk/ink extra heated) swatches and the measurement performed in a 384 well microtiter plate.

15

Model detergent X	LAS	17.6 wt%
	AEO*	2.0 wt%
	Sodium carbonate	20.1 wt%
	Hydrous sodium silicate	12.4 wt%
	Zeolite A	15.0 wt%
	Sodium sulfate	31.8 wt%
	Polyacrylate	1.1 wt%
*AEO is added separately		

100 ml detergent:

0.2g Model X

233µl 0.713M CaCl₂
 233µl 0.357M MgCl₂
 700µl 0.535M NaHCO₃
 80ml Multibuffer 50mM pH 8-11

5

10µl of purified polypeptide normalized to a concentration of 10µM (µmol/Liter) in Buffer: 20mM MES; 0.01% Triton 100-X; 2mM CaCl₂ was diluted MilliQ/0.01% Triton 2000 times for a final concentration of 5 nM. The polypeptides were tested on the EMPA117 EH swatch. 96 well microtiter plates were shaken 1000 rpm for 30 min. Aspirate 50µl from the 96 well microtiter plates to clear 384 well microtiter plates and measure OD590 endpoint. Improvement factor is the measured endpoint value for the variant divided by the measured endpoint value for the reference enzyme (SEQ ID NO 19).

Table 10 shows the performance (improvement factor (IF) compared to the polypeptide with SEQ ID NO 19 with IF = 1) of the polypeptides in the assay described above at pH 8 and/or pH 9, pH 10 and pH 11 on the stain EMPA117EH (blood/milk/ink extra heated).

Table 10

Protease SEQ ID NO	pH 8	pH 9	pH 10	pH 11
SEQ ID NO 4	1.4	1.8	1.7	1.3
SEQ ID NO 10	1.6	1.4	1.4	1.3
SEQ ID NO 12	1.3	1.3	1.3	1.3
SEQ ID NO 14	1.3	1.4	1.4	1.2
SEQ ID NO 16	1.4	1.2	1.6	1.2
SEQ ID NO 18	1.7	2.3	2.0	2.3

The tested polypeptides with SEQ ID NO 4, 10, 12, 14, 16 and 18 show improved performance on the blood stain EMPA117 EH Blood/milk/ink on cotton/polyester at four different pH values.

AZCL-Hemoglobin assay:

Substrate: "AZCL-Hemoglobin" (Megazyme) substrate based on a Azurine-crosslinking to

the hemoglobin molecule.

The assay is performed in 96 well microtiter plates with AZCL-hemoglobin substrate and the measurement is performed in a 384 well microtiter plates. 10µl of purified polypeptide normalized to a concentration of 10µM (µmol/Liter) in Buffer: 20mM MES; 0.01% Triton 100-X; 2mM CaCl₂ was mixed with 90µl 0.01% Triton 100X and 5µl of this solution was mixed with 195µl Substrate mix (AZCL-Hemoglobin 10mg/ml in Multibuffer). The 96 well microtiter plates were shaken 1000rpm for 30 min. Aspirate 50µl from the 96 well microtiter plates to clear 384 well microtiter plates and measure OD590 endpoint. Improvement factor is the measured endpoint value for the variant divided by the measured endpoint value for the reference enzyme (SEQ ID NO 3). Table 11 shows the performance of the protease variants in the assay described above at pH 8 and/or pH 9, pH 10 and pH 11 on AZCL-Hemoglobin" (Megazyme).

Table 11

protease SEQ ID NO	pH 8	pH 9	pH 10	pH 11
SEQ ID NO 4	1.3	2.9	3.2	1.9
SEQ ID NO 12	1.9	2.3	2.5	1.6
SEQ ID NO 14	1.2	1.1	1.1	
SEQ ID NO 16	1.6	2.7	3.2	1.8

The tested polypeptides with SEQ ID NO 4, 12, 14 and 16 show improved performance on the blood stain AZCL-Hemoglobin" (Megazyme) at four different pH values.

Example 8. Full scale wash (FSW) with SEQ ID NO: 2 and SEQ ID NO: 4 in an EU powder model detergent T at 20°C and 40°C with and without bleach

In this example wash performance was evaluated using the EU powder model detergent T as defined in Tables 12-13 below and experimental conditions as defined below (e.g., Tables 14-15). Full scale wash method (FSW) was used for evaluation of stains (e.g., swatches). In the FSW wash performance was expressed as a delta remission value (ΔRem). After washing and rinsing, the swatches were spread out flat and allowed to air dry at room temperature overnight. All washes were evaluated the day after the wash. Light reflectance evaluations of the swatches were done using a Macbeth Color Eye 7000 reflectance spectrophotometer with very small aperture. The measurements were made without UV in the incident light and remission at 460 nm was extracted. Measurements were made on unwashed and washed swatches. The test swatch to be measured was placed on top of another swatch of same type and colour (twin swatch). With

only one swatch of each kind per beaker, a swatch from a replicate wash was used in this way. Remission values for individual swatches were calculated by subtracting the remission value of the unwashed swatch from the remission value of the washed swatch. The total wash performance for each stained swatch set was calculated as the sum of individual ΔRem .

- 5 Calculating the enzyme effect was done by taking the measurements from washed swatches with enzymes and subtract with the measurements from washed without enzyme for each stain. In this example laundry protease having SEQ ID NO: 19 (Savinase) was used as a benchmark (e.g., a corresponding reference).

Table 12. Composition of the EU powder model detergent T without bleach:	
Ingredient's name	Resulting content of active component in formulation, (% w/w)
LAS, sodium salt	10.0
AS, sodium salt	1.8
Soap, sodium salt	2.0
Nonionic (AEO)	3.0
Soda ash	15.0
Hydrous sodium silicate ("disilicate")	2.5
Zeolite 4A	15.1
PCA (copoly(acrylic acid/maleic acid), sodium salt)	1.6
HEDP-Na4	0.13
Sodium citrate	2.0
SRP (copolymer polyether/polyester)	0.5
Sodium sulfate	38.7
Foam regulator, antifoam (silicone)	1.0

Table 13. Composition of the EU powder model detergent T with bleach:	
Ingredient's name	Resulting content of active component in formulation, (% w/w)
LAS, sodium salt	10.0
AS, sodium salt	1.8
Soap, sodium salt	2.0
Nonionic (AEO)	3.0
Soda ash	15.0
Hydrous sodium silicate ("disilicate")	2.5
Zeolite 4A	15.1
PCA (copoly(acrylic acid/maleic acid), sodium salt)	1.6
HEDP-Na4	0.13
Sodium citrate	2.0
SRP (copolymer polyether/polyester)	0.5
Sodium sulfate	13.3
Sodium percarbonate	18.9
TAED (tetraacetylenediamine)	3.0
Foam regulator, antifoam (silicone)	1.0

Table 14. Experimental conditions used in the FSW:	
	Wash conditions
Washing machine	Miele W5841
Wash program	Cotton wash
Volume (water plus)	15.6 L
Standard EU conditions include activation of	

"Water Plus" (giving approx. 1 L water extra in the main wash).	
Temperature, °C (e.g. 20-30-40-60)	20°C and 40°C
Ballast	4 kg mixed cotton/polyester (ratio 65/35)

Table 15. Stains used in the FSW:		
	Name	Stain
1	WE5DASBWKC	Blood
2	051KC	Egg
3	062KC	Scrubbed grass
4	C-05	Blood/milk/ink
5	C-10	Milk/oil/pigment
6	C-H010	Cocoa cocked up milk
7	C-H097	Oatmeal/Chocolate
8	C-H165	Minced meat
9	CS-01	Aged blood
10	CS-37	Full egg/pigment
11	CS-44	Chocolate drink
12	EMPA112	Cocoa
13	EMPA117	Blood/milk/ink
14	EMPA164	Grass
15	KC-H172	Meat pate
16	PC-03	Chocolate milk/soot
17	WE5GMWKC	Grass/Mud

18	WFK 10Z	Chocolate
19	WFK10EG	Egg Yolk
20	WFK10N	Whole Egg

RESULTS

I. Wash performance in the EU powder model detergent T without bleach at 20°C:

- 5 It has been shown that in the EU powder model detergent T without bleach (Table 12) at 20°C the wash performance of SEQ ID NO: 2 and SEQ ID NO: 4 is superior to the wash performance of SEQ ID NO: 19, especially at low doses. In particular, the wash performance of SEQ ID NO: 4 has been found to be better than the wash performance of SEQ ID NO: 19 on certain stains, e.g., when analyzed for individual swatches, especially at low doses (e.g., at 30 and 60 nM) (Table 16).

10

Table 16. Delta remission-based comparison of the wash performances of SEQ ID NO: 2 and SEQ ID NO: 4 to the wash performance of SEQ ID NO: 19 (wash performance of SEQ ID NO: 19 has been used as a corresponding reference at the following concentrations 30 nM, 60 nM and 120 nM):						
20 °C Model T without bleach						
Stains	(SEQ ID NO: 2) 30 nM	(SEQ ID NO: 2) 60 nM	(SEQ ID NO: 2) 120 nM	(SEQ ID NO: 4) 30 nM	(SEQ ID NO: 4) 60 nM	(SEQ ID NO: 4) 120 nM
C-H010 Cocoa cocked up milk	5.18	2.85		8.42	3.00	1.73
C-H097 Oatmeal/Chocolate	3.28	1.28	4.24		4.10	3.15
CS-44 Chocolate drink		1.34			1.35	
EMPA112 Cocoa	2.73			1.21		
PC-03 Chocolate milk/soot	1.91			3.07		
WFK 10Z Chocolate	2.61			1.42	1.78	
WFK10EG Egg Yolk	1.77	1.38	1.49	1.85	1.59	1.08

CS-37 Full egg/pigment		1.85		1.40	3.95	1.38
WE5DASBWKC Blood	2.17				2.09	
C-05 Blood/milk/ink	2.46	2.87	1.58	3.84	3.30	2.20
CS-01 Aged blood	3.35	3.02	2.76	3.56	4.91	3.51
062KC Scrubbed grass					2.48	
EMPA164 Grass	1.83			2.70		
WE5GMWKC Grass/Mud	1.89	1.61			2.19	
C-10 Milk/oil/pigment	3.18	1.99		4.06	1.73	

II. Wash performance in the EU powder model detergent T with bleach at 20°C:

- 5 It has been shown that in the EU powder model detergent T with bleach (Table 13) at 20°C the wash performance of SEQ ID NO: 2 and SEQ ID NO: 4 is superior to the wash performance of SEQ ID NO: 19, especially at low doses. In particular, the wash performance of both SEQ ID NO: 2 and SEQ ID NO: 4 has been found to be better than the wash performance of SEQ ID NO: 19 on certain stains, e.g., when analyzed for individual swatches, especially on blood containing stains (Table 17).

Table 17. Delta remission-based comparison of the wash performances of SEQ ID NO: 2 and SEQ ID NO: 4 to the wash performance of SEQ ID NO: 19 (wash performance of SEQ ID NO: 19 has been used as a corresponding reference at the following concentrations 30 nM, 60 nM and 120 nM):						
20 °C Model T with bleach						
Stains	(SEQ ID NO: 2) 30 nM	(SEQ ID NO: 2) 60 nM	(SEQ ID NO: 2) 120 nM	(SEQ ID NO: 4) 30 nM	(SEQ ID NO: 4) 60 nM	(SEQ ID NO: 4) 120 nM
C-H010 Cocoa cocked up milk	3.74		1.31	8.91	3.80	1.64
C-H097 Oatmeal/Chocolate	1.94			3.29		
EMPA112 Cocoa			1.56		2.54	2.74

KC-H172 Meat pate				2.00		
WFK10EG Egg Yolk				1.53	2.07	2.17
CS-37 Full egg/pigment				5.50	4.24	
C-05 Blood/milk/ink		2.77	4.16	3.62	4.05	4.97
CS-01 Aged blood	1.33	3.05	3.54	3.36	4.42	3.82
EMPA117 Blood/milk/ink				11.13	4.55	3.27
062KC Scrubbed grass				3.39		
EMPA164 Grass		1.42			1.47	1.39
C-10 Milk/oil/pigment	1.75		1.40	2.45	1.47	

III. Wash performance in the EU powder model detergent T without bleach at 40 °C:

It has been shown that in the EU powder model detergent T without bleach (Table 12) at 40 °C the wash performance of both SEQ ID NO: 2 and SEQ ID NO: 4 is better than the wash performance of SEQ ID NO: 19, e.g., on certain stains when analyzed for individual swatches, especially on blood containing stains (Table 18).

Table 18. Delta remission-based comparison of the wash performances of SEQ ID NO: 2 and SEQ ID NO: 4 to the wash performance of SEQ ID NO: 19 (wash performance of SEQ ID NO: 19 has been used as a corresponding reference at the following concentrations 30 nM, 60 nM and 120 nM):

40 °C Model T without bleach						
Stains	(SEQ ID NO: 2) 30 nM	(SEQ ID NO: 2) 60 nM	(SEQ ID NO: 2) 120 nM	(SEQ ID NO: 4) 30 nM	(SEQ ID NO: 4) 60 nM	(SEQ ID NO: 4) 120 nM
C-H010 Cocoa cocked up milk	2.33			1.41		
EMPA112 Cocoa	2.16			3.72	2.56	1.13
WFK 10Z Chocolate					1.64	
C-H165 Minced meat		1.15			1.45	

051KC Egg						1.17
WFK10EG Egg Yolk		1.52	2.01	1.86	1.78	1.34
CS-37 Full egg/pigment		2.16			2.01	
WE5DASBWKC Blood					1.32	
C-05 Blood/milk/ink		2.31	1.63	1.21	3.17	
CS-01 Aged blood		4.65	3.13	2.07	4.97	4.55
EMPA117 Blood/milk/ink		1.17			1.66	
062KC Scrubbed grass		1.53		1.47		
WE5GMWKC Grass/Mud	1.42		4.90		1.94	
C-10 Milk/oil/pigment					1.12	

IV. Wash performance in the EU powder model detergent T with bleach at 40 °C:

- It has been shown that in the EU powder model detergent T with bleach (Table 13) at 40°C the wash performance of both SEQ ID NO: 2 and SEQ ID NO: 4 is better than the wash performance of SEQ ID NO: 19, e.g., on certain stains when analyzed for individual swatches, especially on blood containing stains (Table 19).

Table 19. Delta remission-based comparison of the wash performances of SEQ ID NO: 2 and SEQ ID NO: 4 to the wash performance of SEQ ID NO: 19 (wash performance of SEQ ID NO: 19 has been used as a corresponding reference at the following concentrations 30 nM, 60 nM and 120 nM):						
40 °C Model T with bleach						
Stains	(SEQ ID NO: 2) 30 nM	(SEQ ID NO: 2) 60 nM	(SEQ ID NO: 2) 120 nM	(SEQ ID NO: 4) 30 nM	(SEQ ID NO: 4) 60 nM	(SEQ ID NO: 4) 120 nM
C-H010 Cocoa cocked up milk		1.15				
CS-44 Chocolate drink	1.14		1.14	1.19		
EMPA112 Cocoa			2.01			
WFK10EG Egg Yolk				1.70	1.30	

CS-37 Full egg/pigment				1.61		
WE5DASBWKC Blood	1.29		1.89			
C-05 Blood/milk/ink	4.16	4.05	6.31	4.25	4.63	6.01
CS-01 Aged blood	3.46	5.30	6.20	5.39	7.29	6.38
EMPA117 Blood/milk/ink				2.21		

V. Conclusions:

Based on the results outlined above it has been shown that at 20°C SEQ ID NO: 4 and SEQ ID NO: 2 overall perform 1-2 remission units better than SEQ ID NO: 19, both in presence and absence of bleach. The superior wash performance of SEQ ID NO: 4 and SEQ ID NO: 2 is especially pronounced on blood stains. Furthermore, superior wash performance of SEQ ID NO: 2 and SEQ ID NO: 4 is also present at 40°C on blood stains.

Example 9. Full scale wash (FSW) with SEQ ID NO: 4 in a laundry application using a commercial detergent with bleach

In this example wash performance of SEQ ID NO: 4 has been evaluated using another (e.g., as compared to example 8 above) commercial detergent with bleach and experimental conditions as defined below (Table 20). Full scale wash was used as described in example 8 above:

Table 20. Experimental conditions used in the FSW:		
Wash Programs	Standard wash	Quick wash
Washing machine	Miele W5841	
Wash program	Cotton wash	Express20
Volume (water plus)	15 L	10.5 L
Ballast (mixed co/pe or pure cotton)	4 kg mixed cotton/polyester (ratio 65/35)	2 kg mixed cotton/polyester (ratio 65/35)
Main wash time	50 min	11 min
Rinse	3	2

In this example laundry protease having SEQ ID NO: 19 (Savinase) has been used as a benchmark (e.g., a corresponding reference) again. The wash performance of SEQ ID NO: 4 has been found to be superior to the wash performance of SEQ ID NO: 19 (Savinase) in the commercial detergent with bleach used herein at 40°C, especially on technical blood stains (Table 21). Furthermore, it has been shown that at low dosage and under quick wash conditions, besides blood stains, SEQ ID NO: 4 has better wash performance on e.g., milk, cocoa, some egg, grass and meat stains (Tables 21, 22). Therefore, SEQ ID NO: 4 can carry out its activity quicker than SEQ ID NO: 19 (Savinase), e.g., in the FSW. Furthermore, SEQ ID NO: 4 is much more resistant to bleach than SEQ ID NO: 19 (Savinase), e.g., on blood stains (Tables 21-22).

Table 21. Delta remission-based comparison of the wash performances of SEQ ID NO: 4 to the wash performance of SEQ ID NO: 19 in the commercial detergent with bleach at 40 °C (using SEQ ID NO: 19 as a benchmark):

Stain	15 nM	30 nM	60 nM	120 nM
C-05 Blood/milk/ink	4.0	4.1	3.8	2.4
PC05 Blood/milk/ink	5.1	3.6	3.3	3.7
CS-01 Aged blood	2.6	1.7	2.4	2.7
EMPA117 Blood/milk/ink	10.5	8.4	2.2	
E-117EH Extra heated Blood	8.5	9.3	8.8	5.3
CH010 Cocoa cocked up milk	6.7	2.3		
C03 Chocolate milk/soot	5.7	3.7	2.6	
C-10 Milk/oil/pigment	5.1			
PC-03 Chocolate milk/soot	6.9	1.5		
EMPA112 Cocoa	4.3	2.1		
WFK10EG Egg Yolk	1.1			
CS-37 Full egg/pigment	4.2			
EMPA164 Grass	1.5			
062KC Scrubbed grass	5.4			
KCH172 Meat pate	4.0			

Table 22. Delta remission-based comparison of the wash performances of SEQ ID NO: 4 to the wash performance of SEQ ID NO: 19 in the commercial detergent with bleach at 40°C under quick and normal wash conditions (using SEQ ID NO: 19 as a benchmark):

	Quick wash (11 min)			Standard wash (50 min)		
Stain	30 nM	60 nM	120 nM	30 nM	60 nM	120 nM
C-05 Blood/milk/ink	2.3	2.6	1.2	4.1	3.8	2.4
PC05 Blood/milk/ink	2.5	1.2		3.6	3.3	3.7
CS-01 Aged blood	1.6		2.0	1.7	2.4	2.7
EMPA117 Blood/milk/ink	4.6	4.9	1.9	8.4	2.2	
E-117EH Extra heated Blood	4.4	4.7	3.6	9.3	8.8	5.3
WE5DASBWKC (108KC) Blood	1.2					
CH010 Cocoa cocked up milk	4.2	3.9	2.9	2.3		
C03 Chocolate milk/soot	5.0	2.9		3.7	2.6	
C-10 Milk/oil/pigment	8.2	4.2	3.0			
PC-03 Chocolate milk/soot	5.7	3.6		1.5		
PC10 Milk/oil/pigment	10.6	3.6				
EMPA112 Cocoa	4.6	1.5		2.1		
WFK10N Whole Egg	1.8					
062KC Scrubbed grass	7.3	3.8	2.1			
KCH172 Meat pate	3.9					

CLAIMS

1. A detergent composition suitable for laundry or dishwashing comprising at least one detergent component and an active protease, wherein the protease is a polypeptide comprising the amino acid sequence of SEQ ID NO 4.
2. The composition of claim 1, wherein said composition is a dish wash composition comprising:
 - a) at least 0.01 mg of active protease per gram of composition, wherein the protease is a polypeptide comprising the amino acid sequence of SEQ ID NO 4,
 - b) at least one builder, and
 - c) at least one bleach component.
3. The dish wash composition of claim 2, wherein the dish wash composition is an automatic dish wash (ADW) composition or a hand wash (HDW) composition.
4. The dish wash composition according to claim 2 or 3, wherein the builder is a phosphate, a sodium citrate builder, sodium carbonate, sodium silicate, sodium or a zeolite.
5. The dish wash composition according to any one of claims 2-4, wherein the builder is added in an amount of about 10% to about 65% by weight.
6. The dish wash composition of claim 5, wherein the builder is added in an amount of about 40% to about 65% by weight.
7. The dish wash composition of claim 5, wherein the builder is added in an amount of about 20% to about 65% by weight.
8. The dish wash composition of claim 5, wherein the builder is added in an amount of about 10% to about 50% by weight.
9. The dish wash composition according to any one of claims 2-8, wherein the dish wash composition contains 0-50% by weight of a detergent co-builder.
10. The dish wash composition of claim 9, wherein the dish wash composition contains about 5% to about 30% by weight of the detergent co-builder.

11. The dish wash composition according to any one of claims 2-10, wherein the dish wash composition is phosphate free.
12. The dish wash composition according to any one of claims 2-11, wherein the builder is selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof.
13. The dish wash composition according to any one of claims 2-12, wherein the bleach component is selected from bleaching catalysts, photobleaches, bleach activators, sources of hydrogen peroxide, sodium perborates and hydrogen peroxide, preformed peracids and mixtures thereof.
14. The dish wash composition of claim 13, wherein the source of hydrogen peroxide is sodium percarbonate.
15. The dish wash composition according to any one of claims 2-14 wherein the bleach component is a peroxide.
16. The dish wash composition of claim 15, wherein the peroxide is a percarbonate, persulfate, perphosphate, or persilicate salt.
17. The dish wash composition according to any one of claims 2-16, wherein the bleaching component includes a percarbonate and bleach catalyst.
18. The dish wash composition of claim 17, wherein the bleach catalyst is a manganese compound.
19. The dish wash composition of claim 17, wherein the bleach catalyst is 1,4,7-trimethyl-1,4,7-triazacyclononane or manganese (II) acetate tetrahydrate (MnTACN).
20. The dish wash composition according to any one of claims 2-19, comprising from 1% to 40 wt% of bleach components, wherein the bleach components are a peroxide and a catalyst.
21. The dish wash composition of claim 20, comprising from 0.5% to 30 wt% of bleaching components.
22. The dish wash composition of claim 20 or 21, wherein the peroxide is a percarbonate.

23. The dish wash composition of any one of claims 20-22, wherein the catalyst is a metal-containing bleach catalyst.
24. The dish wash composition of claim 23, wherein the metal-containing bleach catalyst is 1,4,7-trimethyl-1,4,7-triazacyclononane or manganese (II) acetate tetrahydrate (MnTACN)
25. The composition of claim 1, wherein said composition is a granular detergent composition comprising:
- a) at least 0.01 mg of active protease per gram of composition, wherein the protease is a polypeptide comprising the amino acid sequence of SEQ ID NO 4,
 - b) 5 wt % to 50 wt % anionic surfactant,
 - c) 1 wt % to 8 wt % nonionic surfactant, and
 - d) 5 wt % to 40 wt % builder.
26. The granular detergent composition of claim 25, wherein the anionic surfactant is one or more selected from linear alkylbenzenesulfonates (LAS) isomers of LAS, alcohol ether sulfate (AEOS) and sodium lauryl ether sulfate, and sodium laureth sulfate (SLES).
27. The granular detergent composition of claim 25 or 26, wherein the nonionic surfactant is selected from primary alcohol ethoxylates, alkyl ester sulphates, sodium toluene sulfonate (STS) and sodium dodecyl sulfate (SDS).
28. The granular detergent composition of claim 27, wherein the primary alcohol ethoxylate is Neodol 25-7 or Neodol 25-3.
29. The granular detergent composition of claim 27, wherein the primary alcohol ethoxylate is a C12-15 primary alcohol ethoxylate.
30. The granular detergent composition of claim 27, wherein the alkyl ester sulphate is a C9-15 alkyl ester sulphate.
31. The granular detergent composition according to any one of claims 25-30, wherein the builder is a carbonate builder, a zeolite builder, a phosphate builder, a calcium sequestering builder or a complexing agent.
32. The granular detergent composition according to any one of claims 25-31, wherein the builder is a carbonate builder or a zeolite builder.

33. The granular detergent composition according to any one of claims 25-32, wherein the builder is selected from citric acid, methyl glycine-N,N-diacetic acid (MGDA) and/or glutamic-N,N-diacetic acid (GLDA) and mixtures thereof.
34. The granular detergent composition according to any one of claims 25-33, further comprising a bleach component.
35. The granular detergent composition of claim 34, wherein the bleach component is a peroxide.
36. The granular detergent composition of claim 35, wherein the peroxide is a percarbonate, persulfate, perphosphate, or persilicate salt.
37. The composition of claim 1, wherein said composition is a detergent composition comprising:
- a) at least 0.01 mg of active protease per litre or per gram detergent wherein the protease is selected from the polypeptide comprising SEQ ID NO 4,
 - b) 2 wt% to 60 wt% of at least one surfactant, and
 - c) 5 wt% to 50 wt% of at least one builder.
38. The detergent composition of claim 37, wherein said detergent composition is a liquid or powder laundry detergent composition.
39. The detergent composition of claim 37 or 38, wherein the surfactant is a nonionic surfactant, an anionic surfactant, an amphoteric surfactant or a combination thereof.
40. The detergent composition of any one of claims 37-39 further comprising a bleach component.
41. The composition of any one of claims 1-40, wherein said polypeptide comprises an alteration at one or more of the positions selected from the group consisting of 3, 4, 9, 15, 24, 42, 52, 59, 60, 66, 74, 76, 85, 96, 97, 99, 100, 101, 102, 103, 116, 118, 125, 126, 127, 128, 129, 135, 150, 160, 161, 164, 182, 188, 193, 199, 200, 203, 211, 212, 216, 239, 255 and 256 when compared to SEQ ID NO 19, wherein said polypeptide has at least 60%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
42. The composition of claim 41, wherein said polypeptide has at least 65%, but less than

- 100% sequence identity to the amino acid sequence SEQ ID NO 19.
43. The composition of claim 41, wherein said polypeptide has at least 70%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
44. The composition of claim 41, wherein said polypeptide has at least 75%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
45. The composition of claim 41, wherein said polypeptide has at least 80%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
46. The composition of claim 41, wherein said polypeptide has at least 85%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
47. The composition of claim 41, wherein said polypeptide has at least 90%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
48. The composition of claim 41, wherein said polypeptide has at least 91%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
49. The composition of claim 41, wherein said polypeptide has at least 92%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
50. The composition of claim 41, wherein said polypeptide has at least 93%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
51. The composition of claim 41, wherein said polypeptide has at least 94%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
52. The composition of claim 41, wherein said polypeptide has at least 95%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
53. The composition of claim 41, wherein said polypeptide has at least 96%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
54. The composition of claim 41, wherein said polypeptide has at least 97%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

55. The composition of claim 41, wherein said polypeptide has at least 98%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
56. The composition of claim 41, wherein said polypeptide has at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
57. The composition of any one of claims 1-56, wherein said polypeptide comprises one or more of the alterations selected from the group consisting of S3V/T, V4I, S9R, A15T, N42D/R/K, G59E/D, N60D/E, V66A, N74D, S76N, S85N/E, A96E/D, S97D/E, S99G/E/N/H, G100, S101A, V102I, S103, G116V, H118N/D, G125, S126L/F/V, P127Q/N/L, S128A, P129T, P129*, Q135H, S160G/K, Y161A, R164S/L, S182T, A188P, V193M, V199I, Q200E, Y203W, L211D/Q, N212D/S, M216A/S, Q239R, N255D/E and L256Y/E/D when compared to SEQ ID NO 19, wherein the polypeptide has at least 60%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
58. The composition of claim 57, wherein said polypeptide has at least 65%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19
59. The composition of claim 57, wherein said polypeptide has at least 70%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
60. The composition of claim 57, wherein said polypeptide has at least 75%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
61. The composition of claim 57, wherein said polypeptide has at least 80%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
62. The composition of claim 57, wherein said polypeptide has at least 85%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
63. The composition of claim 57, wherein said polypeptide has at least 90%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
64. The composition of claim 57, wherein said polypeptide has at least 91%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
65. The composition of claim 57, wherein said polypeptide has at least 92%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.

66. The composition of claim 57, wherein said polypeptide has at least 93%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
67. The composition of claim 57, wherein said polypeptide has at least 94%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
68. The composition of claim 57, wherein said polypeptide has at least 95%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
69. The composition of claim 57, wherein said polypeptide has at least 96%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
70. The composition of claim 57, wherein said polypeptide has at least 97%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
71. The composition of claim 57, wherein said polypeptide has at least 98%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
72. The composition of claim 57, wherein said polypeptide has at least 99%, but less than 100% sequence identity to the amino acid sequence SEQ ID NO 19.
73. The composition of any one of claims 1-72, wherein said polypeptide consists of the amino acid sequence of SEQ ID NO 4.
74. Use of the composition according to any one of claims 1-73 in a cleaning process, wherein the cleaning process is a process of doing hard surface cleaning or a laundry process.
75. The use of claim 74, wherein the hard surface cleaning process is dish wash.
76. The use of claim 74, for removal of blood stains.
77. The use of claim 76, wherein the blood stains are aged and/or heated blood stains.
78. A method for removing a stain from a surface which comprises contacting the surface with the composition according to any one of claims 1-73.