

PCT

WORLD INTELLECTUAL PROPER
International Bur



INTERNATIONAL APPLICATION PUBLISHED UNDER

WO 9605226A1

<p>(51) International Patent Classification ⁶ : C07K 14/705, 16/28, C12N 15/12</p>	<p>A1</p>	<p>(11) International Publication Number: WO 96/05226 (43) International Publication Date: 22 February 1996 (22.02.96)</p>
<p>(21) International Application Number: PCT/US94/09234 (22) International Filing Date: 16 August 1994 (16.08.94)</p> <p>(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; Suite 300, 9620 Medical Center Drive, Rockville, MD 20850-3338 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): LI, Yi [CN/US]; 16125 Howard Landing Drive, Gaithersburg, MD 20878 (US). ROSEN, Craig, A. [US/US]; 22400 Rollings Hills Road, Laytonsville, MD 20082 (US).</p> <p>(74) Agents: OLSTEIN, Elliot, M., et al.; Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart &, Olstein; 6 Becker Farm Road, Roseland, NJ 07068 (US).</p>	<p>(81) Designated States: AU, CA, CN, JP, KR, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report.</i></p>	
<p>(54) Title: POLYNUCLEOTIDE ENCODING C5a RECEPTOR</p> <p>(57) Abstract</p> <p>A human C5a receptor polypeptide and DNA (RNA) encoding such polypeptide and a procedure for producing such polypeptide by recombinant techniques is disclosed. Also disclosed are methods for utilizing such polypeptide for identifying antagonists and agonists to such polypeptide. Antagonists and agonists may be used therapeutically to inhibit or stimulate the C5a receptor.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

"POLYNUCLEOTIDE ENCODING C5A RECEPTOR".

This invention relates to newly identified polynucleotides, polypeptides encoded by such polynucleotides, the use of such polynucleotides and polypeptides, as well as the production of such polynucleotides and polypeptides. More particularly, the polypeptide of the present invention is a human 7-transmembrane receptor. The transmembrane receptor is a G-protein coupled receptor. More particularly, the 7-transmembrane receptor has been putatively identified as an anaphylatoxin C5a receptor, sometimes hereinafter referred to as "C5a". The invention also relates to inhibiting the action of such polypeptides.

It is well established that many medically significant biological processes are mediated by proteins participating in signal transduction pathways that involve G-proteins and/or second messengers, e.g., cAMP (Lefkowitz, Nature, 351:353-354 (1991)). Herein these proteins are referred to as proteins participating in pathways with G-proteins or PPG proteins. Some examples of these proteins include the GPC receptors, such as those for adrenergic agents and dopamine (Kobilka, B.K., et al., PNAS, 84:46-50 (1987); Kobilka, B.K., et al., Science, 238:650-656 (1987); Bunzow, J.R., et al., Nature, 336:783-787 (1988)), G-proteins themselves, effector

proteins, e.g., phospholipase C, adenylyl cyclase, and phosphodiesterase, and actuator proteins, e.g., protein kinase A and protein kinase C (Simon, M.I., et al., Science, 252:802-8 (1991)).

For example, in one form of signal transduction, the effect of hormone binding is activation of an enzyme, adenylyl cyclase, inside the cell. Enzyme activation by hormones is dependent on the presence of the nucleotide GTP, and GTP also influences hormone binding. A G-protein connects the hormone receptors to adenylyl cyclase. G-protein was shown to exchange GTP for bound GDP when activated by hormone receptors. The GTP-carrying form then binds to an activated adenylyl cyclase. Hydrolysis of GTP to GDP, catalyzed by the G-protein itself, returns the G-protein to its basal, inactive form. Thus, the G-protein serves a dual role, as an intermediate that relays the signal from receptor to effector, and as a clock that controls the duration of the signal.

A wide variety conditions, including infection by bacteria, viruses or fungi, infiltration by cancer cells, allergic or autoimmune disorders and physically or chemically-induced trauma causes an inflammatory response in humans. In all of these diseases and conditions in man and in most mammals, activation of the complement system (a set of proteins, regulatory factors and proteolytic enzymes) via either the classical or the alternative pathway results in the generation of biologically active peptides which serve to amplify and exacerbate the resulting inflammation.

The most active peptide, anaphylatoxin C5a, a 74-amino acid polypeptide, is generated by cleavage of the alpha-chain of native C5 at a specific site by convertase of the blood complement system, as well as by enzymes of the coagulation system. *In vivo*, C5a is thought to play a significant role in the inflammatory response and in a number of clinical disorders (Goldstein, I.M., Inflammation: Basic Principles

and Clinical Correlates, 309-323, Raven Press, New York (1988)). This peptide is a highly potent inflammatory agent, evoking dramatic responses in experimental animals (Bodammer, G. and Vogt, W., *Int. Arch. Allergy Appl. Immunol.*, 33:417-428 (1967)), and stimulating pulmonary, cardiac, vascular and gastrointestinal tissues *in vitro* (Stimler, N.P., et al., *Am. J. Pathol.*, 100:327-348 (1980)). C5a is a potent activator of polymorphonuclear neutrophils and macrophages, stimulating chemotaxis, hydrolytic enzyme release, and superoxide anion formation (Ward, P.A. and Newman, L.J., *J. Immunol.*, 102:93-99 (1969)). Several reports have additionally demonstrated actions of this peptide on eosinophils, including chemotaxis and increased hexose uptake, in addition to its actions on mast cells and basophils (Hugli, T.E., *Biological Response Mediators and Modulators*, 99-116, Academic Press, New York (1983)). In addition, the anaphylatoxin has been shown to have a spasmogenic effect on various tissues; it stimulates smooth muscle contraction (Stimler, N.P., et al., *J. Immunol.*, 126:2258-2261 (1981)); induces histamine release from mast cells, promotes serotonin release from platelets (Meuer, S., et al., *J. Immunol.*, 126:1506-1509 (1981)), and increases vascular permeability (Jose, P.J., et al., *J. Immunol.*, 127:2376-2380 (1981)).

The responses elicited by C5a in polymorphonuclear leukocytes result from the winding of the anaphylatoxin to a high-affinity receptor on the plasma membrane (Chenoweth, D.E. and Hugli, T.E., *Mol. Immunol.*, 17:151-161 (1980)). In these cells, it appears that the mechanism of signal transduction through the membrane involves one or more GTP-binding proteins (G proteins) as is the case with other chemotactic receptors. The receptor molecule for C5a on human neutrophils has been well characterized with respect to its kinetics and saturability and many of the structural requirements for its activity are known. Reports indicate that the neutrophil C5a receptor binds its ligand with a

nanomolar affinity constant, is expressed in approximately 100,000 copies per cell, and the binding sub-unit has an apparent mass of approximately 52 kDa.

The interaction of C5a with polymorphonuclear leukocytes and other target cells and tissues results in increased histamine release, vascular permeability, smooth muscle contraction, and an influx into tissues of inflammatory cells, including neutrophils, eosinophils and basophils (Hugli, T.E., Springer, *Semin. Immunopathol.*, 7:193-219 (1981)). C5a may also play an important role in mediating inflammatory effects of phagocytic mononuclear cells that accumulate at sites of chronic inflammation (Allison, A.C., et al., *H.U. Agents and Actions*, 8:27 (1978)). C5a can induce chemotaxis in monocytes and cause them to release lysosomal enzymes in a manner analogous to the neutrophil responses elicited by these agents. C5a may have an immunoregulatory role by enhancing antibody, particularly at sites of inflammation (Morgan, E.L., et al., *J. Exp. Med.*, 155:1412 (1982)).

In accordance with one aspect of the present invention, there are provided novel polypeptides which have been putatively identified as a C5a receptor, as well as fragments, analogs and derivatives thereof. The polypeptides of the present invention are of human origin.

In accordance with another aspect of the present invention, there are provided polynucleotides (DNA or RNA) which encode such polypeptides.

In accordance with a further aspect of the present invention, there is provided a process for producing such polypeptides by recombinant techniques.

In accordance with yet a further aspect of the present invention, there are provided antibodies against such polypeptides.

In accordance with another embodiment, there is provided a process for using the receptor to screen for receptor antagonists and/or agonists and/or receptor ligands.

In accordance with still another embodiment of the present invention there is provided a process of using such agonists for therapeutic purposes, for example, as a defense against bacterial infection, to stimulate the immunoregulatory effects of C5a, to treat cancers, immunodeficiency diseases and severe infections.

In accordance with another aspect of the present invention there is provided a process of using such antagonists for treating asthma, bronchial allergy, chronic inflammation, systemic lupus erythematosus, vasculitis, rheumatoid arthritis, osteoarthritis, gout, some auto-allergic diseases, transplant rejection, ulcerative colitis, in certain shock states, myocardial infarction, and post-viral encephalopathies.

These and other aspects of the present invention should be apparent to those skilled in the art from the teachings herein.

The following drawings are illustrative of embodiments of the invention and are not meant to limit the scope of the invention as encompassed by the claims.

Figure 1 shows the cDNA sequence and the corresponding deduced amino acid sequence of the putative mature G-protein coupled receptor of the present invention. The standard one-letter abbreviation for amino acids is used.

Figure 2 is an illustration of the secondary structural features of the G-protein coupled receptor. The first 7 illustrations set forth the regions of the amino acid sequence which are alpha helices, beta helices, turn regions or coiled regions. The boxed areas are the areas which correspond to the region indicated. The second set of figures illustrate areas of the amino acid sequence which are exposed to intracellular, cytoplasmic or are membrane-

spanning. The hydrophilicity plot illustrates areas of the protein sequence which are the lipid bilayer of the membrane and are, therefore, hydrophobic, and areas outside the lipid bilayer membrane which are hydrophilic. The antigenic index corresponds to the hydrophilicity plot, since antigenic areas are areas outside the lipid bilayer membrane and are capable of binding antigens. The surface probability plot further corresponds to the antigenic index and the hydrophilicity plot. The amphipathic plots show those regions of the protein sequences which are polar and non-polar. The flexible regions correspond to the second set of illustrations in the sense that flexible regions are those which are outside the membrane and inflexible regions are transmembrane regions.

Figure 3 illustrates an amino acid alignment of the G-protein coupled receptor of the present invention and C5a receptors from various species of animals. Faded areas are those areas which match with the other amino acid sequences in the figure.

It should be pointed out that sequencing inaccuracies are a common problem which occurs in polynucleotide sequences. Accordingly, the sequence of the drawing is based on several sequencing runs and the sequencing accuracy is considered to be at least 97%.

In accordance with an aspect of the present invention, there is provided an isolated nucleic acid (polynucleotide) which encodes for the mature polypeptide having the deduced amino acid sequence of Figure 1 or for the mature polypeptide encoded by the cDNA of the clone deposited as ATCC Deposit No. 75821 on June 24, 1994.

A polynucleotide encoding a polypeptide of the present invention is predominantly expressed in peripheral lymphocytes. The polynucleotide of this invention was discovered in a cDNA library derived from human osteoclastoma stromal cells. It is structurally related to the G protein-

coupled receptor family. It contains an open reading frame encoding a protein of 355 amino acid residues. The protein exhibits the highest degree of homology to a human C5a receptor with 27 % identity and 54 % similarity over the entire amino acid sequence.

The polynucleotide of the present invention may be in the form of RNA or in the form of DNA, which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. The coding sequence which encodes the mature polypeptide may be identical to the coding sequence shown in Figure 1 or that of the deposited clone or may be a different coding sequence which coding sequence, as a result of the redundancy or degeneracy of the genetic code, encodes the same mature polypeptide as the DNA of Figure 1 or the deposited cDNA.

The polynucleotide which encodes for the mature polypeptide of Figure 1 or for the mature polypeptide encoded by the deposited cDNA may include: only the coding sequence for the mature polypeptide; the coding sequence for the mature polypeptide and additional coding sequence such as a leader or secretory sequence or a proprotein sequence; the coding sequence for the mature polypeptide (and optionally additional coding sequence) and non-coding sequence, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the mature polypeptide.

Thus, the term "polynucleotide encoding a polypeptide" encompasses a polynucleotide which includes only coding sequence for the polypeptide as well as a polynucleotide which includes additional coding and/or non-coding sequence.

The present invention further relates to variants of the hereinabove described polynucleotides which encode for fragments, analogs and derivatives of the polypeptide having the deduced amino acid sequence of Figure 1 or the polypeptide encoded by the cDNA of the deposited clone. The

variant of the polynucleotide may be a naturally occurring allelic variant of the polynucleotide or a non-naturally occurring variant of the polynucleotide.

Thus, the present invention includes polynucleotides encoding the same mature polypeptide as shown in Figure 1 or the same mature polypeptide encoded by the cDNA of the deposited clone as well as variants of such polynucleotides which variants encode for a fragment, derivative or analog of the polypeptide of Figure 1 or the polypeptide encoded by the cDNA of the deposited clone. Such nucleotide variants include deletion variants, substitution variants and addition or insertion variants.

As hereinabove indicated, the polynucleotide may have a coding sequence which is a naturally occurring allelic variant of the coding sequence shown in Figure 1 or of the coding sequence of the deposited clone. As known in the art, an allelic variant is an alternate form of a polynucleotide sequence which may have a substitution, deletion or addition of one or more nucleotides, which does not substantially alter the function of the encoded polypeptide.

The present invention also includes polynucleotides, wherein the coding sequence for the mature polypeptide may be fused in the same reading frame to a polynucleotide sequence which aids in expression and secretion of a polypeptide from a host cell, for example, a leader sequence which functions as a secretory sequence for controlling transport of a polypeptide from the cell. The polypeptide having a leader sequence is a preprotein and may have the leader sequence cleaved by the host cell to form the mature form of the polypeptide. The polynucleotides may also encode for a proprotein which is the mature protein plus additional 5' amino acid residues. A mature protein having a prosequence is a proprotein and is an inactive form of the protein. Once the prosequence is cleaved an active mature protein remains.

Thus, for example, the polynucleotide of the present invention may encode for a mature protein, or for a protein having a prosequence or for a protein having both a prosequence and a presequence (leader sequence).

The polynucleotides of the present invention may also have the coding sequence fused in frame to a marker sequence which allows for purification of the polypeptide of the present invention. The marker sequence may be a hexahistidine tag supplied by a pQE-9 vector to provide for purification of the mature polypeptide fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a mammalian host, e.g. COS-7 cells, is used. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson, I., et al., Cell, 37:767 (1984)).

The present invention further relates to polynucleotides which hybridize to the hereinabove-described sequences if there is at least 50% and preferably 70% identity between the sequences. The present invention particularly relates to polynucleotides which hybridize under stringent conditions to the hereinabove-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. The polynucleotides which hybridize to the hereinabove described polynucleotides in a preferred embodiment encode polypeptides which either retain substantially the same biological function or activity as the mature polypeptide encoded by the cDNA of Figure 1 or the deposited cDNA, i.e. function as a G-protein coupled receptor or retain the ability to bind the ligand for the receptor even though the polypeptide does not function as a G-protein coupled receptor, for example, a soluble form of the receptor.

The deposit(s) referred to herein will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for purposes of Patent Procedure. These deposits are provided merely as convenience to those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112. The sequence of the polynucleotides contained in the deposited materials, as well as the amino acid sequence of the polypeptides encoded thereby, are incorporated herein by reference and are controlling in the event of any conflict with any description of sequences herein. A license may be required to make, use or sell the deposited materials, and no such license is hereby granted.

The present invention further relates to a G-protein coupled receptor polypeptide which has the deduced amino acid sequence of Figure 1 or which has the amino acid sequence encoded by the deposited cDNA, as well as fragments, analogs and derivatives of such polypeptide.

The terms "fragment," "derivative" and "analog" when referring to the polypeptide of Figure 1 or that encoded by the deposited cDNA, means a polypeptide which either retains substantially the same biological function or activity as such polypeptide, i.e. functions as a G-protein coupled receptor, or retains the ability to bind the ligand or the receptor even though the polypeptide does not function as a G-protein coupled receptor, for example, a soluble form of the receptor. An analog includes a proprotein which can be activated by cleavage of the proprotein portion to produce an active mature polypeptide.

The polypeptide of the present invention may be a recombinant polypeptide, a natural polypeptide or a synthetic polypeptide, preferably a recombinant polypeptide.

The fragment, derivative or analog of the polypeptide of Figure 1 or that encoded by the deposited cDNA may be (i) one in which one or more of the amino acid residues are

substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide, such as a leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

The polypeptides and polynucleotides of the present invention are preferably provided in an isolated form, and preferably are purified to homogeneity.

The term "isolated" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

The present invention also relates to vectors which include polynucleotides of the present invention, host cells which are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques.

Host cells are genetically engineered (transduced or transformed or transfected) with the vectors of this

invention which may be, for example, a cloning vector or an expression vector. The vector may be, for example, in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the C5a receptor genes. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

The polynucleotides of the present invention may be employed for producing polypeptides by recombinant techniques. Thus, for example, the polynucleotide may be included in any one of a variety of expression vectors for expressing a polypeptide. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other vector may be used as long as it is replicable and viable in the host.

The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and others are deemed to be within the scope of those skilled in the art.

The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct mRNA synthesis. As representative examples of such promoters, there may be mentioned: LTR or SV40 promoter, the *E. coli*. *lac* or *trp*, the phage lambda P_L promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses.

The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

In addition, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in E. coli.

The vector containing the appropriate DNA sequence as hereinabove described, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the protein.

As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as E. coli, Streptomyces, Salmonella typhimurium; fungal cells, such as yeast; insect cells such as Drosophila and Sf9; animal cells such as CHO, COS or Bowes melanoma; plant cells, etc. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein.

More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. The following vectors are provided by way of example. Bacterial: pQE70, pQE60, pQE-9 (Qiagen), pbs, pD10, phagescript, psiX174, pbluescript SK, pbsks, pNH8A, pNH16a, pNH18A, pNH46A (Stratagene); ptrc99a, pKK223-

3, pKK233-3, pDR540, pRIT5 (Pharmacia). Eukaryotic: pWLNEO, pSV2CAT, pOG44, pXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia). However, any other plasmid or vector may be used as long as they are replicable and viable in the host.

Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and PCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda P_R, P_L and trp. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

In a further embodiment, the present invention relates to host cells containing the above-described constructs. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation. (Davis, L., Dibner, M., Battey, I., Basic Methods in Molecular Biology, (1986)).

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, the polypeptides of the invention can be synthetically produced by conventional peptide synthesizers.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, N.Y., (1989), the disclosure of which is hereby incorporated by reference.

Transcription of the DNA encoding the polypeptides of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp that act on a promoter to increase its transcription. Examples including the SV40 enhancer on the late side of the replication origin bp 100 to 270, a cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), α -factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product.

Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation

initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include E. coli, Bacillus subtilis, Salmonella typhimurium and various species within the genera Pseudomonas, Streptomyces, and Staphylococcus, although others may also be employed as a matter of choice.

As a representative but nonlimiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, such methods are well known to those skilled in the art.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of

mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, Cell, 23:175 (1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

The G-protein coupled receptor polypeptides can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

The polypeptides of the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. Polypeptides of the invention may also include an initial methionine amino acid residue.

The G-protein coupled receptor of the present invention may be employed in a process for screening for antagonists and/or agonists for the receptor.

In general, such screening procedures involve providing appropriate cells which express the receptor on the surface thereof. In particular, a polynucleotide encoding the receptor of the present invention is employed to transfect cells to thereby express the G-protein coupled receptor. Such transfection may be accomplished by procedures as hereinabove described.

One such screening procedure involves the use of the melanophores which are transfected to express the G-protein coupled receptor of the present invention. Such a screening technique is described in PCT WO 92/01810 published February 6, 1992.

Thus, for example, such assay may be employed for screening for a receptor antagonist by contacting the melanophore cells which encode the G-protein coupled receptor with both the receptor ligand and a compound to be screened. Inhibition of the signal generated by the ligand indicates that a compound is a potential antagonist for the receptor, i.e., inhibits activation of the receptor.

The screen may be employed for determining an agonist by contacting such cells with compounds to be screened and determining whether such compound generates a signal, i.e., activates the receptor.

Other screening techniques include the use of cells which express the G-protein coupled receptor (for example, transfected CHO cells) in a system which measures extracellular pH changes caused by receptor activation, for example, as described in Science, volume 246, pages 181-296 (October 1989). For example, potential agonists or antagonists may be contacted with a cell which expresses the G-protein coupled receptor and a second messenger response, e.g. signal transduction or pH changes, may be measured to

determine whether the potential agonist or antagonist is effective.

Another such screening technique involves introducing RNA encoding the G-protein coupled receptor into xenopus oocytes to transiently express the receptor. The receptor oocytes may then be contacted in the case of antagonist screening with the receptor ligand and a compound to be screened, followed by detection of inhibition of a calcium signal.

Another screening technique involves expressing the G-protein coupled receptor in which the receptor is linked to a phospholipase C or D. As representative examples of such cells, there may be mentioned endothelial cells, smooth muscle cells, embryonic kidney cells, etc. The screening for an antagonist or agonist may be accomplished as hereinabove described by detecting activation of the receptor or inhibition of activation of the receptor from the phospholipase second signal.

Another method involves screening for antagonists by determining inhibition of binding of labeled ligand to cells which have the receptor on the surface thereof. Such a method involves transfecting a eukaryotic cell with DNA encoding the G-protein coupled receptor such that the cell expresses the receptor on its surface and contacting the cell with a potential antagonist in the presence of a labeled form of a known ligand. The ligand can be labeled, e.g., by radioactivity. The amount of labeled ligand bound to the receptors is measured, e.g., by measuring radioactivity of the receptors. If the potential antagonist binds to the receptor as determined by a reduction of labeled ligand which binds to the receptors, the binding of labeled ligand to the receptor is inhibited.

The present invention also provides a method for determining whether a ligand not known to be capable of binding to a G-protein coupled receptor can bind to such

receptor which comprises contacting a mammalian cell which expresses a G-protein coupled receptor with the ligand under conditions permitting binding of ligands to the G-protein coupled receptor, detecting the presence of a ligand which binds to the receptor and thereby determining whether the ligand binds to the G-protein coupled receptor. The systems hereinabove described for determining agonists and/or antagonists may also be employed for determining ligands which bind to the receptor.

In general, antagonists for G-protein coupled receptors which are determined by screening procedures may be employed for a variety of therapeutic purposes. For example, such antagonists have been employed for treatment of hypertension, angina pectoris, myocardial infarction, ulcers, asthma, allergies, psychoses, depression, migraine, vomiting, and benign prostatic hypertrophy.

Agonists for G-protein coupled receptors are also useful for therapeutic purposes, such as the treatment of asthma, Parkinson's disease, acute heart failure, hypotension, urinary retention, and osteoporosis.

A potential antagonist is an antibody, or in some cases an oligonucleotide, which binds to the G-protein coupled receptor but does not elicit a second messenger response such that the activity of the G-protein coupled receptor is prevented. Potential antagonists also include proteins which are closely related to the ligand of the G-protein coupled receptor, i.e. a fragment of the ligand, which have lost biological function and when binding to the G-protein coupled receptor, elicit no response.

A potential antagonist also includes an antisense construct prepared through the use of antisense technology. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the

polynucleotide sequence, which encodes for the mature polypeptides of the present invention, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix -see Lee et al., Nucl. Acids Res., 6:3073 (1979); Cooney et al, Science, 241:456 (1988); and Dervan et al., Science, 251: 1360 (1991)), thereby preventing transcription and the production of G-protein coupled receptor. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into the G-protein coupled receptors (antisense - Okano, J. Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of G-protein coupled receptors.

Another potential antagonist is a small molecule which binds to the G-protein coupled receptor, making it inaccessible to ligands such that normal biological activity is prevented. Examples of small molecules include but are not limited to small peptides or peptide-like molecules.

Potential antagonists also include a soluble form of a G-protein coupled receptor, e.g. a fragment of the receptor, which binds to the ligand and prevents the ligand from interacting with membrane bound G-protein coupled receptors.

The G-protein coupled receptor of the present invention has been putatively identified as a C5a receptor. This identification has been made as a result of amino acid sequence homology.

The antagonists may be used to treat all pathological conditions which result from anaphylaxis stimulated by the C5a polypeptide and mediated by the C5a receptor. These pathological conditions include asthma, bronchial allergy,

chronic inflammation, systemic lupus erythematosus, vasculitis, serum sickness, angioedema, rheumatoid arthritis, osteoarthritis, gout, bullous skin diseases, hypersensitivity, pneumonitis, idiopathic pulmonary fibrosis, immune complex-mediated glomerulonephritis, psoriasis, allergic rhinitis, adult respiratory distress syndrome, acute pulmonary disorders, endotoxin shock, hepatic cirrhosis, pancreatitis, inflammatory bowel diseases (including Crohn's disease and ulcerative colitis), thermal injury, gram-negative sepsis, necrosis in myocardial infarction, leukophoresis, exposure to medical devices (including, but not limited to, hemodialyzer membranes and extracorporeal blood circulation equipment), chronic hepatitis, transplant rejection, post-viral encephalopathies, and/or ischemia induced myocardial or brain injury. These antagonist may also be used as prophylactics for such conditions as shock accompanying Deng Urea fever. The antagonists may be employed in a composition with a pharmaceutically acceptable carrier, e.g., as hereinafter described.

The agonists identified by the screening method as described above, may be employed to enhance the C5a reactions mediated through the C5a receptor, which include defense against bacterial infection, stimulation of the immunoregulatory effects of C5a, treatment of cancers, immunodeficiency diseases and severe infections.

The C5a receptor and antagonists or agonists may be employed in combination with a suitable pharmaceutical carrier. Such compositions comprise a therapeutically effective amount of the polypeptide, and a pharmaceutically acceptable carrier or excipient. Such a carrier includes but is not limited to saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The formulation should suit the mode of administration.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of

the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the polypeptides of the present invention may be employed in conjunction with other therapeutic compounds.

The pharmaceutical compositions may be administered in a convenient manner such as by the topical, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes. The pharmaceutical compositions are administered in an amount which is effective for treating and/or prophylaxis of the specific indication. In general, the pharmaceutical compositions will be administered in an amount of at least about 10 $\mu\text{g}/\text{kg}$ body weight and in most cases they will be administered in an amount not in excess of about 8 mg/Kg body weight per day. In most cases, the dosage is from about 10 $\mu\text{g}/\text{kg}$ to about 1 mg/kg body weight daily, taking into account the routes of administration, symptoms, etc.

The C5a receptor polypeptides and antagonists or agonists which are polypeptides, may also be employed in accordance with the present invention by expression of such polypeptides *in vivo*, which is often referred to as "gene therapy."

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) encoding a polypeptide *ex vivo*, with the engineered cells then being provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For example, cells may be engineered by procedures known in the art by use of a retroviral particle containing RNA encoding a polypeptide of the present invention.

Similarly, cells may be engineered *in vivo* for expression of a polypeptide *in vivo* by, for example, procedures known in the art. As known in the art, a producer cell for producing a retroviral particle containing RNA encoding the polypeptide of the present invention may be administered to a patient for engineering cells *in vivo* and expression of the polypeptide *in vivo*. These and other methods for administering a polypeptide of the present invention by such method should be apparent to those skilled in the art from the teachings of the present invention. For example, the expression vehicle for engineering cells may be other than a retrovirus, for example, an adenovirus which may be used to engineer cells *in vivo* after combination with a suitable delivery vehicle.

The sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphisms) are presently available for marking chromosomal location. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis of the cDNA is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified fragment.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes or pools of large genomic clones in an analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include *in situ* hybridization, prescreening with labeled flow-sorted chromosomes and preselection by hybridization to construct chromosome specific-cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with cDNA as short as 500 or 600 bases; however, clones larger than 2,000 bp have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. FISH requires use of the clones from which the EST was derived, and the longer the better. For example, 2,000 bp is good, 4,000 is better, and more than 4,000 is probably not necessary to get good results a reasonable percentage of the time. For a review of this technique, see Verma et al., *Human Chromosomes: a Manual of Basic Techniques*, Pergamon Press, New York (1988).

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, *Mendelian Inheritance in Man* (available on line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the

affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

The polypeptides, their fragments or other derivatives, or analogs thereof, or cells expressing them can be used as an immunogen to produce antibodies thereto. These antibodies can be, for example, polyclonal or monoclonal antibodies. The present invention also includes chimeric, single chain, and humanized antibodies, as well as Fab fragments, or the product of an Fab expression library. Various procedures known in the art may be used for the production of such antibodies and fragments.

Antibodies generated against the polypeptides corresponding to a sequence of the present invention can be obtained by direct injection of the polypeptides into an animal or by administering the polypeptides to an animal, preferably a nonhuman. The antibody so obtained will then bind the polypeptides itself. In this manner, even a sequence encoding only a fragment of the polypeptides can be used to generate antibodies binding the whole native polypeptides. Such antibodies can then be used to isolate the polypeptide from tissue expressing that polypeptide.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, 1975, *Nature*, 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, *Immunology Today* 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies

(Cole, et al., 1985, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96).

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention.

The present invention will be further described with reference to the following examples; however, it is to be understood that the present invention is not limited to such examples. All parts or amounts, unless otherwise specified, are by weight.

In order to facilitate understanding of the following examples certain frequently occurring methods and/or terms will be described.

"Plasmids" are designated by a lower case p preceded and/or followed by capital letters and/or numbers. The starting plasmids herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the ordinarily skilled artisan.

"Digestion" of DNA refers to catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements were used as would be known to the ordinarily skilled artisan. For analytical purposes, typically 1 μ g of plasmid or DNA fragment is used with about 2 units of enzyme in about 20 μ l of buffer solution. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 μ g of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the

manufacturer. Incubation times of about 1 hour at 37°C are ordinarily used, but may vary in accordance with the supplier's instructions. After digestion the reaction is electrophoresed directly on a polyacrylamide gel to isolate the desired fragment.

Size separation of the cleaved fragments is performed using 8 percent polyacrylamide gel described by Goeddel, D. et al., *Nucleic Acids Res.*, 8:4057 (1980).

"Oligonucleotides" refers to either a single stranded polydeoxynucleotide or two complementary polydeoxynucleotide strands which may be chemically synthesized. Such synthetic oligonucleotides have no 5' phosphate and thus will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide will ligate to a fragment that has not been dephosphorylated.

"Ligation" refers to the process of forming phosphodiester bonds between two double stranded nucleic acid fragments (Maniatis, T., et al., *Id.*, p. 146). Unless otherwise provided, ligation may be accomplished using known buffers and conditions with 10 units to T4 DNA ligase ("ligase") per 0.5 µg of approximately equimolar amounts of the DNA fragments to be ligated.

Unless otherwise stated, transformation was performed as described in the method of Graham, F. and Van der Eb, A., *Virology*, 52:456-457 (1973).

Example 1

Bacterial Expression and Purification of C5a Receptor

The DNA sequence encoding the C5a receptor, ATCC # 75821, is initially amplified using PCR oligonucleotide primers corresponding to the 5' end sequences of the processed C5a receptor protein (minus the signal peptide sequence) and the vector sequences 3' to the gene. Additional nucleotides corresponding to the C5a receptor were added to the 5' and 3'

sequences respectively. The 5' oligonucleotide primer has the sequence 5' GACTAAAGCTTAATGGAAGATTTGGAGGAA 3' contains a HindIII restriction enzyme site followed by 19 nucleotides of C5a receptor coding sequence starting from the presumed terminal amino acid of the processed protein codon. The 3' sequence 5' GAACTTCTAGACCGTTATTGAGCTGTTTCCAGGAG 3' contains complementary sequences to an XbaI site and is followed by 18 nucleotides of the gene. The restriction enzyme sites correspond to the restriction enzyme sites on the bacterial expression vector pQE-9 (Qiagen, Inc. 9259 Eton Avenue, Chatsworth, CA, 91311). pQE-9 encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter operator (P/O), a ribosome binding site (RBS), a 6-His tag and restriction enzyme sites. pQE-9 was then digested with HindIII and XbaI. The amplified sequences were ligated into pQE-9 and were inserted in frame with the sequence encoding for the histidine tag and the RBS. The ligation mixture was then used to transform *E. coli* strain available from Qiagen under the trademark M15/rep 4 by the procedure described in Sambrook, J. et al., Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989). M15/rep4 contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies were selected. Plasmid DNA was isolated and confirmed by restriction analysis. Clones containing the desired constructs were grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells were grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG ("Isopropyl-B-D-thiogalacto pyranoside") was then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to

increased gene expression. Cells were grown an extra 3 to 4 hours. Cells were then harvested by centrifugation. The cell pellet was solubilized in the chaotropic agent 6 Molar Guanidine HCl. After clarification, solubilized C5a receptor was purified from this solution by chromatography on a Nickel-Chelate column under conditions that allow for tight binding by proteins containing the 6-His tag. Hochuli, E. et al., J. Chromatography 411:177-184 (1984). The C5a receptor was eluted from the column in 6 molar guanidine HCl pH 5.0 and for the purpose of renaturation adjusted to 3 molar guanidine HCl, 100mM sodium phosphate, 10 mmolar glutathione (reduced) and 2 mmolar glutathione (oxidized). After incubation in this solution for 12 hours the protein was dialyzed to 10 mmolar sodium phosphate.

Example 2

Expression of Recombinant C5a Receptor in COS cells

The expression of plasmid, pC5a HA is derived from a vector pcDNA1/Amp (Invitrogen) containing: 1) SV40 origin of replication, 2) ampicillin resistance gene, 3) E.coli replication origin, 4) CMV promoter followed by a polylinker region, a SV40 intron and polyadenylation site. A DNA fragment encoding the entire pC5a protein and a HA tag fused in frame to its 3' end was cloned into the polylinker region of the vector, therefore, the recombinant protein expression is directed under the CMV promoter. The HA tag correspond to an epitope derived from the influenza hemagglutinin protein as previously described (I. Wilson, H. Niman, R. Heighten, A. Cherenson, M. Connolly, and R. Lerner, 1984, Cell 37, 767). The infusion of HA tag to the target protein allows easy detection of the recombinant protein with an antibody that recognizes the HA epitope.

The plasmid construction strategy is described as follows:

The DNA sequence encoding for the C5a receptor, ATCC # 75821, was constructed by PCR on the full-length gene cloned using two primers: the 5' primer 5' GTCCGAAGCTTGCCACCATGGAA GATTTGGAGGAA 3' contains a HindIII site followed by 18 nucleotides of C5a receptor coding sequence starting from the initiation codon; the 3' sequence 5' CTAGCTCGAGTCAAGCGTAGTCTG GGACGTCGTATGGGTAGCATTGAGCTGTTTCCAGGAG 3' contains complementary sequences to an XhoI site, translation stop codon, HA tag and the last 18 nucleotides of the C5a receptor coding sequence (not including the stop codon). Therefore, the PCR product contains a HindIII site, C5a receptor coding sequence followed by HA tag fused in frame, a translation termination stop codon next to the HA tag, and an XhoI site. The PCR amplified DNA fragment and the vector, pcDNAI/Amp, were digested with HindIII and XhoI restriction enzyme and ligated. The ligation mixture was transformed into E. coli strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037) the transformed culture was plated on ampicillin media plates and resistant colonies were selected. Plasmid DNA was isolated from transformants and examined by restriction analysis for the presence of the correct fragment. For expression of the recombinant C5a receptor, COS cells were transfected with the expression vector by DEAE-DEXTRAN method. (J. Sambrook, E. Fritsch, T. Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989)). The expression of the C5a receptor HA protein was detected by radiolabelling and immunoprecipitation method. (E. Harlow, D. Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, (1988)). Cells were labelled for 8 hours with ³⁵S-cysteine two days post transfection. Culture media were then collected and cells were lysed with detergent (RIPA buffer (150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50mM Tris, pH 7.5). (Wilson, I. et al., Id. 37:767 (1984)). Both cell lysate and culture media were precipitated with a HA

specific monoclonal antibody. Proteins precipitated were analyzed on 15% SDS-PAGE gels.

Example 3

Cloning and expression of C5a receptor using the baculovirus expression system

The DNA sequence encoding the full length C5a receptor protein, ATCC # 75821, was amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene:

The 5' primer has the sequence 5' GCCGGATCCGCCA CCATGGAAGATTTGGAGGAA 3' and contains a BamHI restriction enzyme site (in bold) followed by 6 nucleotides resembling an efficient signal for the initiation of translation in eukaryotic cells (J. Mol. Biol. 1987, 196, 947-950, Kozak, M.), and is just behind the first 18 nucleotides of the gene (the initiation codon for translation "ATG" is underlined).

The 3' primer has the sequence 5' GCCGGATCCGT TATTGAGCTGTTCCAG 3' and contains the cleavage site for the restriction endonuclease BamHI and 18 nucleotides complementary to the 3' non-translated sequence of the C5a receptor gene. The amplified sequences were isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101 Inc., La Jolla, Ca.). The fragment was then digested with the endonucleases BamHI and then isolated again on a 1% agarose gel. This fragment is designated F2.

The vector pRG1 (modification of pVL941 vector, discussed below) is used for the expression of the C5a receptor protein using the baculovirus expression system (for review see: Summers, M.D. and Smith, G.E. 1987, A manual of methods for baculovirus vectors and insect cell culture procedures, Texas Agricultural Experimental Station Bulletin No. 1555). This expression vector contains the strong polyhedrin promoter of the Autographa californica nuclear polyhedrosis virus (AcMNPV) followed by the recognition sites

for the restriction endonuclease BamHI. The polyadenylation site of the simian virus (SV)40 is used for efficient polyadenylation. For an easy selection of recombinant viruses the beta-galactosidase gene from E.coli is inserted in the same orientation as the polyhedrin promoter followed by the polyadenylation signal of the polyhedrin gene. The polyhedrin sequences are flanked at both sides by viral sequences for the cell-mediated homologous recombination of co-transfected wild-type viral DNA. Many other baculovirus vectors could be used in place of pRG1 such as pAc373, pVL941 and pAcIM1 (Luckow, V.A. and Summers, M.D., Virology, 170:31-39).

The plasmid was digested with the restriction enzymes BamHI and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The DNA was then isolated from a 1% agarose gel as described above. This vector DNA is designated V2.

Fragment F2 and the dephosphorylated plasmid V2 were ligated with T4 DNA ligase. E.coli HB101 cells were then transformed and bacteria identified that contained the plasmid (pBacC5a) with the C5a receptor gene using the enzyme BamHI. The sequence of the cloned fragment was confirmed by DNA sequencing.

5 μ g of the plasmid pBacC5a was co-transfected with 1.0 μ g of a commercially available linearized baculovirus ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA.) using the lipofection method (Felgner et al. Proc. Natl. Acad. Sci. USA, 84:7413-7417 (1987)).

1 μ g of BaculoGold™ virus DNA and 5 μ g of the plasmid pBacC5a were mixed in a sterile well of a microtiter plate containing 50 μ l of serum free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards 10 μ l Lipofectin plus 90 μ l Grace's medium were added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture was added dropwise to the Sf9 insect

cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1ml Grace' medium without serum. The plate was rocked back and forth to mix the newly added solution. The plate was then incubated for 5 hours at 27°C. After 5 hours the transfection solution was removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum was added. The plate was put back into an incubator and cultivation continued at 27°C for four days.

After four days the supernatant was collected and a plaque assay performed similar as described by Summers and Smith (supra). As a modification an agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) was used which allows an easy isolation of blue stained plaques. (A detailed description of a "plaque assay" can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10).

Four days after the serial dilution of the viruses was added to the cells, blue stained plaques were picked with the tip of an Eppendorf pipette. The agar containing the recombinant viruses was then resuspended in an Eppendorf tube containing 200 μ l of Grace's medium. The agar was removed by a brief centrifugation and the supernatant containing the recombinant baculoviruses was used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes were harvested and then stored at 4°C.

Sf9 cells were grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells were infected with the recombinant baculovirus V-C5a at a multiplicity of infection (MOI) of 2. Six hours later the medium was removed and replaced with SF900 II medium minus methionine and cysteine (Life Technologies Inc., Gaithersburg). 42 hours later 5 μ Ci of 35 S-methionine and 5 μ Ci 35 S cysteine (Amersham) were added. The cells were further incubated for 16 hours before they

were harvested by centrifugation and the labelled proteins visualized by SDS-PAGE and autoradiography.

Example 4

Expression pattern of C5a receptor in human tissue

Northern blot analysis was carried out to examine the levels of expression of C5a receptor in human tissues. Total cellular RNA samples were isolated with RNazol™ B system (Biotechx Laboratories, Inc. 6023 South Loop East, Houston, TX 77033). About 10µg of total RNA isolated from each human tissue specified was separated on 1% agarose gel and blotted onto a nylon filter. (Sambrook, Fritsch, and Maniatis, Molecular Cloning, Cold Spring Harbor Press, (1989)). The labeling reaction was done according to the Stratagene Prime-It kit with 50ng DNA fragment. The labeled DNA was purified with a Select-G-50 column. (5 Prime - 3 Prime, Inc. 5603 Arapahoe Road, Boulder, CO 80303). The filter was then hybridized with radioactive labeled full length C5a receptor gene at 1,000,000 cpm/ml in 0.5 M NaPO₄, pH 7.4 and 7% SDS overnight at 65°C. After wash twice at room temperature and twice at 60°C with 0.5 x SSC, 0.1% SDS, the filter was then exposed at -70°C overnight with an intensifying screen. The message RNA for C5a receptor is abundant in peripheral lymphocytes.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, within the scope of the appended claims, the invention may be practiced otherwise than as particularly described.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: LI, ET AL.
- (ii) TITLE OF INVENTION: C5a Receptor
- (iii) NUMBER OF SEQUENCES: 2
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: CARELLA, BYRNE, BAIN, GILFILLAN,
CECCHI, STEWART & OLSTEIN
 - (B) STREET: 6 BECKER FARM ROAD
 - (C) CITY: ROSELAND
 - (D) STATE: NEW JERSEY
 - (E) COUNTRY: USA
 - (F) ZIP: 07068
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: 3.5 INCH DISKETTE
 - (B) COMPUTER: IBM PS/2
 - (C) OPERATING SYSTEM: MS-DOS
 - (D) SOFTWARE: WORD PERFECT 5.1
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: Concurrently
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:

(viii) ATTORNEY/AGENT INFORMATION:
 (A) NAME: FERRARO, GREGORY D.
 (B) REGISTRATION NUMBER: 36,134
 (C) REFERENCE/DOCKET NUMBER: 325800-195

(ix) TELECOMMUNICATION INFORMATION:
 (A) TELEPHONE: 201-994-1700
 (B) TELEFAX: 201-994-1744

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS
 (A) LENGTH: 2024 BASE PAIRS
 (B) TYPE: NUCLEIC ACID
 (C) STRANDEDNESS: SINGLE
 (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CGGCAAAGCA	GGCATGGACA	ATAGCTTCTC	TCCTCACAGA	AATTAACTG	ATTTCTTCAT	60
TCTCCATTTA	GCAAGGTCAT	GGAAGATTTG	GAGGAAACAT	TATTTGAAGA	ATTTGAAAAC	120
TATTCCTATG	ACCTAGACTA	TTACTCTCTG	GAGTCTGATT	TGGAGGAGAA	AGTCCAGCTG	180
GGAGTTGTTC	ACTGGGTCTC	CCTGGTGTTA	TATTGTTTGG	CTTTTGTTC	GGGAATCCCA	240
GGGAAATGCC	TCTATCATT	GTTTCACGGG	GTTCAAGTGG	AAGAAGACAG	TCACACTCTG	300
TGGTTCCTCA	ATCTAGCCAT	TGCGGATTC	ATTTTTCTTC	TCTTCTGCC	CCTGTACATC	360
TCCTATGTGG	CCATGAATTT	CCACTGGCCC	TTTGGCATCT	GGCTGTGCAA	AGCCAATTCC	420
TTCCTGCCC	AGTTGAACAT	GTTTGCCAGT	GTTTTTTTCC	TGACAGTGAT	CAGCCTGGAC	480
CACTATATCC	ACTTGATCCA	TCCTGTCTTA	TCTCATCGGC	ATCGAACCT	CAAGAACTCT	540
CTGATTGTCA	TTATATTCAT	CTGGCTGTG	GCTTCTCTAA	TTGGCGGTCC	TGCCCTGTAC	600
TTCCGGGATA	CTGTGGAGTT	CAATAATCAT	ACTCTTTGGT	ATAACAATTT	TCAGAAGCAT	660
GATCCTGACC	TCACTTGGAT	CAGGCACCAT	GTTCTGACTT	GGGTGAAATT	TATCATTGGT	720
TATCTCTTCC	CTTTGCTAAC	AATGAGTATT	CGGTACTTGT	GTCTCATCTT	CAAGGTGAAG	780
AAGCGAAGCA	TCCTGATCTC	CAGTAGGCAT	TTCTGGACAA	TTCTGGTTGT	GGTTGTGGCC	840
TTTGTGGTTT	GGTGGACTCC	TTATCACCTG	TTTAGCATTG	GGGAGCTCAC	CATTCACCAC	900
AATAGCTATT	CCCACCATGT	GATGCAGGCT	GGAATCCCCC	TCTCCACTGG	TTTGGCATT	960
CTCAATAGTT	GCTTGAACCC	CATCCTTTAT	GTCCTAGTTA	GTAAGAAGTT	CCAAGCTCGC	1020

```

TTCCGGTCCT CAGTTGCTGA GATACTCAAG TACACACTGT GGAAGTCAG CTGTTCTGGC      1080
ACAGTGAGTG AACAGCTCAG GAACTCAGAA ACCAAGAATC TGTGTCTCCT GGAAACAGCT      1140
CAATAAGTTA TTACTTTTCC ACAAATCAGT ATATGGCTTT TTATGTGGGT CCTCTGACTG      1200
ATGCTTTCAG ATTAATAATG TTTCCAAGAT AGAGAGCCGA CTCCACTTTC ATAGTTATTG      1260
TTTCTGGTCA CTATATAGGC ATCACATTTT TGTGTGGATA TGAAACTTAG GAAGGATCCT      1320
CTTGACTCCT TGTGATGTGG CAATAAATTT TTTTAAAAA ACTGAAAATA CTTAGGAAGG      1380
ATCCGCATAA TTTTTTCTG CAACTTAAAT GAAATGCATC ATTCTTGTTA ATCATACCAT      1440
GGTGAATTAA TCACTTTTGA AGCAATATCA GTTATTTTTT GAATAATAAC TTTTCTAAAG      1500
CCTTAAGTCT TAATATTAAA TATATGATTA GCCAGGCCCG GTGGCTGACA CCTGTAATCC      1560
CAGCACTTG  GGAGGCCAAG GTGGGGGGAT TACCCGAGGT CAGGAATTCG AGACCAGCCT      1620
GACCAACATG GAGAAACCCC GTCTCTACTA AAAATCCAAA ATTAGCCGGT CATGGTGGTG      1680
CATGTCTGCA AACCCAGCTA CTCGGGAGGC TGAAGCAGGA GAATCCACTT GAACCTGGGA      1740
GGCAGAGGTT GTGGTGAGCC AACATCACAC CATTGCACTC CAGCCTGGGC CACAAGAGTA      1800
AAACTCTGTC TCAAAAATAA ATAAATAAAA TAGATAAATA AATATATGAT TAACTAATTT      1860
TAAAAATGTT AAAATGTATT CTTAAATTCA TTTTAATTTT GTACAATAAC CTGCTAGACA      1920
CATTTTTTAAA ATGCAACATG TGTACTTAAT TTCTTTATGT AATCTATGTA TATACATTTA      1980
TGAATTAAG  TAATTGTTGG TTATCTTAAA AAAAAAAAAA AAAA      2024
    
```

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 355 AMINO ACIDS
- (B) TYPE: AMINO ACID
- (C) STRANDEDNESS:
- (D) TOPOLOGY: LINEAR

(ii) MOLECULE TYPE: PROTEIN

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Glu Asp Leu Glu Glu Thr Leu Phe Glu Glu Phe Glu Asn Tyr
      5                      10
Ser Tyr Asp Leu Asp Tyr Tyr Ser Leu Glu Ser Asp Leu Glu Glu
      20                      25
Lys Val Gln Leu Gly Val Val His Trp Val Ser Leu Val Leu Tyr
      35                      40
Cys Leu Ala Phe Val Leu Gly Ile Pro Gly Lys Cys Leu Asp His
      50                      55
Leu Val His Gly Val Gln Val Glu Glu Asp Ser His Thr Leu Trp
    
```

	65		70		75
Phe Leu Asn Leu Ala Ile Ala Asp Phe Ile Phe Leu Leu Phe Leu					
	80		85		90
Pro Leu Tyr Ile Ser Tyr Val Ala Met Asn Phe His Trp Pro Phe					
	95		100		105
Gly Ile Trp Leu Cys Lys Ala Asn Ser Phe Thr Ala Gln Leu Asn					
	110		115		120
Met Phe Ala Ser Val Phe Phe Leu Thr Val Ile Ser Leu Asp His					
	125		130		135
Tyr Ile His Leu Ile His Pro Val Leu Ser His Arg His Arg Thr					
	140		145		150
Leu Lys Asn Ser Leu Ile Val Ile Ile Phe Ile Trp Leu Val Ala					
	155		160		165
Ser Leu Ile Gly Gly Pro Ala Leu Tyr Phe Arg Asp Thr Val Glu					
	170		175		180
Phe Asn Asn His Thr Leu Trp Tyr Asn Asn Phe Gln Lys His Asp					
	185		190		195
Pro Asp Leu Thr Trp Ile Arg His His Val Leu Thr Trp Val Lys					
	200		205		210
Phe Ile Ile Gly Tyr Leu Phe Pro Leu Leu Thr Met Ser Ile Arg					
	215		220		225
Tyr Leu Cys Leu Ile Phe Lys Val Lys Lys Arg Ser Ile Leu Ile					
	230		235		240
Ser Ser Arg His Phe Trp Thr Ile Leu Val Val Val Val Ala Phe					
	245		250		255
Val Val Trp Trp Thy Pro Tyr His Leu Phe Ser Ile Gly Glu Leu					
	260		265		270
Thr Ile His His Asn Ser Tyr Ser His His Val Met Gln Ala Gly					
	275		280		285
Ile Pro Leu Ser Thr Gly Leu Ala Phe Leu Asn Ser Cys Leu Asn					
	290		295		300
Pro Ile Leu Tyr Val Leu Val Ser Lys Lys Phe Gln Ala Arg Phe					
	305		310		315
Arg Ser Ser Val Ala Glu Ile Leu Lys Tyr Thr Leu Trp Glu Val					
	320		325		330

Ser Cys Ser Gly Thr Val Ser Glu Gln Leu Arg Asn Ser Gly Thr
335 340 345
Lys Asn Leu Cys Leu Leu Glu Thr Ala Gln
350 355

WHAT IS CLAIMED IS:

1. An isolated polynucleotide selected from the group consisting of:
 - (a) a polynucleotide encoding the G-protein coupled receptor polypeptide having the deduced amino acid sequence of Figure 1 or a fragment, analog or derivative of said polypeptide;
 - (b) a polynucleotide encoding the G-protein coupled receptor polypeptide having the amino acid sequence encoded by the cDNA contained in ATCC Deposit No. 75821 or a fragment, analog or derivative of said polypeptide.
2. The polynucleotide of Claim 1 wherein the polynucleotide is DNA.
3. The polynucleotide of Claim 1 wherein the polynucleotide is RNA.
4. The polynucleotide of Claim 1 wherein the polynucleotide is genomic DNA.
5. The polynucleotide of Claim 2 wherein said polynucleotide encodes G-protein coupled receptor having the deduced amino acid sequence of Figure 1.
6. The polynucleotide of Claim 2 wherein said polynucleotide encodes the G-protein coupled receptor polypeptide encoded by the cDNA of ATCC Deposit No. 75821.
7. The polynucleotide of Claim 1 having the coding sequence of G-protein coupled receptor as shown in Figure 1.
8. The polynucleotide of Claim 2 having the coding sequence of G-protein coupled receptor deposited as ATCC Deposit No. 75821.
9. A vector containing the DNA of Claim 2.
10. A host cell genetically engineered with the vector of Claim 9.
11. A process for producing a polypeptide comprising: expressing from the host cell of Claim 10 the polypeptide encoded by said DNA.

12. A process for producing cells capable of expressing a polypeptide comprising genetically engineering cells with the vector of Claim 9.
13. An isolated DNA hybridizable to the DNA of Claim 2 and encoding a polypeptide having G-protein coupled receptor activity.
14. A polypeptide selected from the group consisting of (i) a G-protein coupled receptor polypeptide having the deduced amino acid sequence of Figure 1 and fragments, analogs and derivatives thereof and (ii) a G-protein coupled receptor polypeptide encoded by the cDNA of ATCC Deposit No. 75821 and fragments, analogs and derivatives of said polypeptide.
15. The polypeptide of Claim 14 wherein the polypeptide is a G-protein coupled receptor having the deduced amino acid sequence of Figure 1.
16. An antibody against the polypeptide of claim 14.
17. A compound which activates the G-protein coupled receptor polypeptide of claim 14.
18. A compound which inhibits activation of the G-protein coupled receptor polypeptide of claim 14.
19. A method for the treatment of a patient having need of activation of a G-protein coupled receptor polypeptide of claim 14 comprising: administering to the patient a therapeutically effective amount of a compound Claim 17.
20. A method for the treatment of a patient having need to inhibit activation of the G-protein coupled receptor of claim 14 comprising: administering to the patient a therapeutically effective amount of a compound of Claim 18.
21. The polypeptide of Claim 14 wherein the polypeptide is a soluble fragment of the G-protein coupled receptor and is capable of binding a ligand for the receptor.
22. A process for identifying antagonists and agonists to the G-protein coupled receptor comprising:
expressing the G-protein coupled receptor on the surface of a cell;

contacting the cell with a receptor ligand and compound to be screened;

determining whether a second signal is generated from the interaction of the ligand and the receptor; and

identifying if the compound to be screened is an agonist or antagonist.

23. A process for determining whether a ligand not known to be capable of binding to a G-protein coupled receptor can bind thereto comprising:

contacting a mammalian cell which expresses the G-protein coupled receptor with a potential ligand;

detecting the presence of the ligand which binds to the receptor; and

determining whether the ligand binds to the G-protein coupled receptor.

FIG. 1A

10
 30 50
 CGGCAAAGCAGGCATGGACAATAGCTTCTCTCCTCACAGAAATTTAACTGATTCTTTCAT
 70 90 110
 TCCTCCATTTAGCAAGGTCATGGAAGATTTGGAGGAAACATTTTGAAGAATTGAAAAC
 130 150 170
 M E D L E E T L F E E F E N
 TATTCCTATGACCTAGACTATTACTCTCTGGAGTCTGATTGGAGGAGAAAGTCCAGCTG
 190 210 230
 Y S Y D L D Y Y S L E S D L E E K V Q L
 GGAGTTGTTCACTGGGTCTCCCTGGTGTATATTTGTTGGCITTTGTTCTGGGAATTCCA
 250 270 290
 G V V H W V S L V L Y C L A F V L G I P
 GGGAAATGCCCTCGATCATTGGTTCAAGGTTCAAGTGGAAAGAAGACAGTCACACTCTG
 310 330 350
 G K C L D H L V H G V Q V E E D S H T L
 TGGTTCCTCAATCTAGCCATTGGGATTTTCATTTTCTCTCTTCTGCCCCTGTACATC
 370 390 410
 W F L N L A I A D F I F L L F L P L Y I
 TCCATGTGGCCATGAATTTCCACTGGCCCTTTGGCATCTGGCTGTGCAAGCCAATTC
 430 450 470
 S Y V A M N F H W P F G I W L C K A N S

MATCH WITH FIG. 1B

1 / 10

FIG. 1B

MATCH WITH FIG. 1A

TTCAC T GCCCAGTTGAACATGTTGCCAGTGTTTTCCAGTACAGTGATCAGCCCTGGAC
 F T A Q L N M F A S V F F L T V I S L D
 490 510 530

CACTATATCCACTTGATCCATCCGTGCTTATCTCATCGGCATCGAACCCCTCAAGAACTCT
 H Y I H L I H P V L S H R H R T L K N S
 550 570 590

CTGATTGTCATATATTCATCTGGCTTGTGGCTTCTCTAATTGGCGGTCCCTGCCCTGTAC
 L I V I I F I W L V A S L I G G P A L Y
 610 630 650

TTCCGGGATACTGTGGAGTTCAATAATCATACTTTGGIATAACAATTTTCAGAAGCAT
 F R D T V E F N N H T L W Y N N F Q K H
 670 690 710

GATCCCTGACCTTGGATCAGGCACCAATGTTCTGACTGGGTGAAATTTATCATTTGGT
 D P D L T W I R H H V L T W V K F I I G
 730 750 770

TATCTCTTCCCTTGGTAACAATGAGTATTCGGTACTTGTGTCATCTTCAAGGTGAAG
 Y L F P L L T M S I R Y L C L I F K V K
 790 810 830

AAGCGAAGCATCCATCCAGTAGGCATTTCTGGACAATTTCTGGTTGTGGTTGTGGCC
 K R S I L I S S R H F W T I L V V V A
 850 870 890

TTTGTGGTTGGTGGACTCCTTATCACCCTGTTTAGCATTTGGGGAGCTCACCATTCAACCAC
 F V V W W T P Y H L F S I G E L T I H H
 MATCH WITH FIG. 1C

FIG. 1D

MATCH WITH FIG. 1C

1390	1410	1430
A	T	C
1450	1470	1490
GG	T	G
1510	1530	1550
C	C	T
1570	1590	1610
C	A	G
1630	1650	1670
G	A	C
1690	1710	1730
C	A	T
1750	1770	1790
G	G	C
1810	1830	1850
A	A	A
1870	1890	1910
T	A	A
1930	1950	1970
C	A	T
1990	2010	
T	G	A

FIG. 2B

MATCH WITH FIG. 2A





FIG. 2A



7 / 10

FIG. 3A_a

MATCH WITH FIG. 3A_b →

M X X - - - T X - X X X N Y S - D X D X X X L X X D

10 20

1 M E D L E E T L F E E F E N Y S Y D L D Y Y S L E S D
 1 M N S F N Y T T - P D Y G H Y D - D K D T L D L N T P
 1 M D T - - - N M - S L L M N K S - A V N L M N V S G S
 1 M - - - - - - - - - I L N S S T E D G I K R I Q D D

V V G V L G N X L V V W V X G F X M K - X T V X T I W

60 70

50 V L G I P G K C L D H L V H G V Q V E - E D S H T L W
 49 L V G V L G N A L V V W V T A F E A K - R T I N A I W
 46 V L G V L G N G L V I W V A G F R M K - H T V T T I S
 40 V V G I F G N S L V V I V I Y F Y M K L K T V A S V F

X A M X X H W P F G X X L C K I X S X X I X L N L F A

110 120

97 V A M N F H W P F G I W L C K A N S F T A Q L N M F A
 96 I V Q H H H W P F G G A A C S I L P S L I L L N M Y A
 93 M V M G G H W P F G W F M C K F I Y T V I D I N L F G
 88 T A M E Y R W P F G N Y L C K I A S A S V S F N L Y A

N H R T X X L A K I V I I V I W L L A X L L T L P A I

160 170

147 R H R T L K N S L I V I I F I W L V A S L I G G P A -
 146 N F R G A G L A W I A C A V A W G L A L L L T I P S F
 143 N H R T V S L A K K V I I V P W I C A F L L T L P V I
 138 L R R T M L V A K V T C I I I W L L A G L A S L P A I

Y S X - - K - - - D K R X X R X T X L T - - G I V K F

210 220


188 Y N N F Q K H D P D L T W I R H H V L T - - - W V K F
 192 Y S H - - - - - D K R R E R A V - - - - - A I V R L
 191 F S P W T K D P V E K R K V A V T M L T V R G I I R F
 184 Y E S - - - - - - - - - Q N S T L P I G L G L T K N

FIG. 3Ab

← MATCH WITH FIG. 3Aa


X X K X S X X L X V X D X X S L V L F A V X F										Majority													
30						40				50													
L	E	K	V	Q	-	L	G	V	V	H	V	S	L	V	L	Y	C	L	A	F	34.PEP		
V	D	K	T	S	N	T	L	R	V	P	D	I	L	A	L	V	I	F	A	V	V	HuC5aR.PEP	
T	Q	S	V	S	A	G	Y	I	V	L	D	V	F	S	Y	L	I	F	A	V	T	MuFLMPR.PEP	
C	P	K	A	G	R	H	N	Y	I	F	V	M	I	P	-	T	L	Y	S	I	I	HuATIIR.PEP	
F L N L A I A D F - - C F L L T L P L Y I X S										Majority													
80										90											100		
F	L	N	L	A	I	A	D	F	I	-	-	F	L	L	F	L	P	L	Y	I	S	Y	34.PEP
F	L	N	L	A	V	A	D	F	L	S	C	L	A	-	-	L	P	I	L	F	T	S	HuC5aR.PEP
Y	L	N	L	A	I	A	D	F	-	-	C	F	T	S	T	L	P	F	Y	I	A	S	MuFLMPR.PEP
L	L	N	L	A	L	A	D	L	-	-	C	F	L	L	T	L	P	L	W	A	V	Y	HuATIIR.PEP
S V F L L A X I S L D R Y L X V X H P V W S Q										Majority													
130										140												150	
S	V	F	F	L	T	V	I	S	L	D	H	Y	I	H	L	I	H	P	V	L	S	H	34.PEP
S	I	L	L	L	A	T	I	S	A	D	R	F	L	L	V	F	K	R	I	W	C	Q	HuC5aR.PEP
S	V	F	L	I	A	L	I	A	L	D	R	C	I	C	V	L	H	P	V	W	A	O	MuFLMPR.PEP
S	V	F	L	L	T	C	L	S	I	D	R	Y	L	A	I	V	H	P	M	K	S	R	HuATIIR.PEP
I - - - L Y F V X N X E X X P X K T X C T F D										Majority													
180										190												200	
-	-	-	-	L	Y	F	R	D	T	V	E	F	N	N	H	-	-	-	-	T	L	W	34.PEP
-	-	-	-	L	Y	R	V	V	R	E	E	Y	F	P	P	K	V	L	C	G	V	D	HuC5aR.PEP
I	R	-	-	L	T	T	V	P	N	S	R	L	G	P	G	K	T	A	C	T	F	D	MuFLMPR.PEP
I	H	R	N	V	F	F	I	E	N	T	N	I	-	-	-	-	T	V	C	A	F	H	HuATIIR.PEP
I L G F L F P L L I X X I C Y T L I X X K X K										Majority													
230										240												250	
I	I	G	Y	L	F	P	L	L	T	M	S	I	R	Y	L	C	L	I	F	K	V	K	34.PEP
V	L	G	F	L	W	P	L	L	T	L	T	I	C	Y	T	F	I	L	L	R	T	W	HuC5aR.PEP
I	I	G	F	S	T	P	M	S	I	V	A	I	C	Y	G	L	I	T	T	K	I	H	MuFLMPR.PEP
I	L	G	F	L	F	P	F	L	I	I	L	T	S	Y	T	L	I	W	K	A	L	K	HuATIIR.PEP

FIG. 3Ba



Decoration 'Decoration #1': Shade with residues that match the Consensus exactly.

FIG. 3Bb



MATCH WITH FIG. 3B a

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/09234

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07K 14/705, 16/28; C12N 15/12
 US CL : 435/6, 69.1, 252.3, 320.1; 530/350, 388.22; 536/23.5
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 69.1, 252.3, 320.1; 530/350, 388.22; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, STN/MEDLINE
 search terms: C5a, anaphylatoxin#, receptor#, DNA, cDNA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Biochemistry Journal, Vol. 288, issued 1992, J. J. Perret, "Cloning and functional expression of the canine anaphylatoxin C5a receptor", pages 911-917, see entire document.	1-23
A	Biochemistry, Vol. 30, issued 1991, F. Boulay et al., "Expression Cloning of a Receptor for C5a Anaphylatoxin on Differentiated HL-60 Cells", pages 2993-2999, see entire document.	1-23

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

10 NOVEMBER 1994

Date of mailing of the international search report

07 DEC 1994

Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOHN D. ULM

Telephone No. (703) 308-0196