

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 March 2009 (19.03.2009)

PCT

(10) International Publication Number
WO 2009/033665 A2

(51) International Patent Classification:

A61K 38/08 (2006.01) A61P 31/00 (2006.01)
A61K 38/29 (2006.01) A61P 35/00 (2006.01)
A61P 3/00 (2006.01) A61P 37/00 (2006.01)
A61P 9/00 (2006.01) A61P 31/18 (2006.01)
A61P 11/00 (2006.01) A61P 31/20 (2006.01)
A61P 25/28 (2006.01)

(21) International Application Number:

PCT/EP2008/007452

(22) International Filing Date:

9 September 2008 (09.09.2008)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

07017746.4 11 September 2007 (11.09.2007) EP

(71) Applicant (for all designated States except US): **MONDO-BIOTECH LABORATORIES AG** [LI/LI]; Herrengasse 21, FL-9490 Vaduz (LI).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BEVEC, Dorian** [DE/DE]; Kriegerstrasse 62, 82110 Germering (DE). **CAVALLI, Fabio** [CH/CH]; Via Pasquée 23, CH-6925 Gentilino (CH). **CAVALLI, Vera** [CH/CH]; Via Pasquée, 23, CH-6925 Gentilino (CH). **BACHER, Gerald** [DE/DE]; Masurenweg 5, 82110 Germering (DE).

(74) Agent: **ARTH, Hans-Lothar**; ABK, Jasminweg 9, 14052 Berlin (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

— without international search report and to be republished upon receipt of that report
— with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(54) Title: USE OF A PEPTIDE AS A THERAPEUTIC AGENT

(57) Abstract: The present invention is directed to the use of the peptide compound Gly-Arg-Gly-Asp-Asn-Pro as a therapeutic agent for the prophylaxis and/or treatment of cancer, autoimmune diseases, fibrotic diseases, inflammatory diseases, neurodegenerative diseases, infectious diseases, lung diseases, heart and vascular diseases and metabolic diseases. Moreover the present invention relates to pharmaceutical compositions preferably in form of a lyophilisate or liquide buffer solution or artificial mother milk formulation or mother milk substitute containing the peptide Gly-Arg-Gly-Asp-Asn-Pro optionally together with at least one pharmaceutically acceptable carrier, cryoprotectant, lyoprotectant, excipient and/or diluent.



WO 2009/033665 A2

Use of a peptide as a therapeutic agent

Specification

5

The present invention is directed to the use of the peptide compound Gly-Arg-Gly-Asp-Asn-Pro-OH as a therapeutic agent for the prophylaxis and/or treatment of cancer, an infectious disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, an autoimmune disease, or a heart and vascular disease.

10

Background of the invention

15

The identification of a therapeutic compound effective for the prophylaxis and/or treatment of a disease can be based on the activity of the compound in a biological assay. A biological assay that mimics a disease causative mechanism can be used to test the therapeutic activity of a candidate peptide.

20

The causative mechanism of many diseases is the over activity of a biological pathway. A peptide that can reduce the activity of the biological pathway can be effective in the prophylaxis and/or treatment of the disease caused by the over activity of the biological pathway. Similarly the causative mechanism of many diseases is the over production of a biological molecule. A peptide that can reduce the production of the biological molecule or block the activity of the over produced biological molecule can be effective in the prophylaxis and/or treatment of the disease caused by the over production of the biological molecule.

25

30

Conversely, the causative mechanism of many diseases is the under activity of a biological pathway. A peptide that can increase the activity of the biological pathway can be effective in the prophylaxis and/or treatment of the disease caused by the under activity of the biological pathway. Also similarly the causative mechanism of many diseases is the under production of a biological molecule. A peptide that can increase the production of the biological molecule or mimic the biological activity of the under produced biological molecule can be effective in the prophylaxis and/or treatment of the disease caused by the under production of the biological molecule.

35

It is the object of the present invention to provide a peptide compound for the prophylaxis and/or treatment of cancer, autoimmune diseases, fibrotic diseases, inflammatory diseases, neurodegenerative diseases, infectious diseases, lung diseases, heart and vascular diseases and metabolic diseases.

The object of the present invention is solved by the teaching of the independent claims. Further advantageous features, aspects and details of the invention are evident from the dependent claims, the description, and the examples of the present application.

Description of the invention

The present invention relates to the use of the peptide Gly-Arg-Gly-Asp-Asn-Pro-OH, its use as a therapeutic in medicine and for the prophylaxis and/or treatment of cancer, autoimmune diseases, fibrotic diseases, inflammatory diseases, neurodegenerative diseases, infectious diseases, lung diseases, heart and vascular diseases and metabolic diseases. Also disclosed are pharmaceutical formulations preferably in form of a lyophilisate or liquid buffer solution or artificial mother milk formulation containing the inventive peptide. The peptide is especially useful for prophylaxis and/or treatment of AIDS, AIDS related diseases, HIV, Kaposi Sarcoma, lymphomas, anal dysplasia, cervical dysplasia, dementia complex, peripheral neuropathy, depression, anxiety, fatigue, anemia, thrombocytopenia, lipodystrophy, pulmonary hypertension, Wasting syndrome, aspergillosis, atypical mycobacteriosis, bacteraemia, candidiasis, HCMV infection, cryptococcosis, cryptosporidiosis, enteritis, histoplasmosis, nocardiosis, encephalitis, progressive multifocal leukoencephalopathy, pneumonia, toxoplasmosis, tuberculosis and other diseases which are described in the following.

Moreover the present invention relates to a peptide combination of the above mentioned peptide with the peptide compound Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe as well as to pharmaceutical compositions containing said peptide combination.

Cancer, tumors, proliferative diseases, malignancies and their metastases

The term "cancer" as used herein refers also to tumors, proliferative diseases, malignancies and their metastases. Examples for cancer diseases are adenocarcinoma, choroidal melanoma, acute leukemia, acoustic neurinoma, ampullary carcinoma, anal carcinoma, astrocytoma, basal cell carcinoma, pancreatic cancer, desmoid tumor, bladder cancer, bronchial carcinoma, non-small cell lung cancer (NSCLC), breast cancer, Burkitt's lymphoma, corpus cancer, CUP-syndrome (carcinoma of unknown primary), colorectal cancer, small intestine cancer, small intestinal tumors, ovarian cancer, endometrial carcinoma, ependymoma, epithelial cancer types, Ewing's tumors, gastrointestinal tumors, gastric cancer, gallbladder cancer, gall bladder carcinomas, uterine cancer, cervical cancer, cervix, glioblastomas,

gynecologic tumors, ear, nose and throat tumors, hematologic neoplasias, hairy cell leukemia, urethral cancer, skin cancer, skin testis cancer, brain tumors (gliomas), brain metastases, testicle cancer, hypophysis tumor, carcinoids, Kaposi's sarcoma, laryngeal cancer, germ cell tumor, bone cancer, colorectal carcinoma, head and neck tumors
5 (tumors of the ear, nose and throat area), colon carcinoma, craniopharyngiomas, oral cancer (cancer in the mouth area and on lips), cancer of the central nervous system, liver cancer, liver metastases, leukemia, eyelid tumor, lung cancer, lymph node cancer (Hodgkin's/Non-Hodgkin's), lymphomas, stomach cancer, malignant melanoma, malignant neoplasia, malignant tumors gastrointestinal tract, breast carcinoma, rectal
10 cancer, medulloblastomas, melanoma, meningiomas, Hodgkin's disease, mycosis fungoides, nasal cancer, neurinoma, neuroblastoma, kidney cancer, renal cell carcinomas, non-Hodgkin's lymphomas, oligodendroglioma, esophageal carcinoma, osteolytic carcinomas and osteoplastic carcinomas, osteosarcomas, ovarian carcinoma, pancreatic carcinoma, penile cancer, plasmocytoma, squamous cell carcinoma of the
15 head and neck (SCCHN), prostate cancer, pharyngeal cancer, rectal carcinoma, retinoblastoma, vaginal cancer, thyroid carcinoma, Schneeberger disease, esophageal cancer, spinalioma, T-cell lymphoma (mycosis fungoides), thymoma, tube carcinoma, eye tumors, urethral cancer, urologic tumors, urothelial carcinoma, vulva cancer, wart appearance, soft tissue tumors, soft tissue sarcoma, Wilm's tumor, cervical carcinoma
20 and tongue cancer.

The peptides and peptide combination of the present invention was tested using the assays described in Examples 1-7, 9-17 for their effect as active therapeutic agents in the prophylaxis and/or treatment of cancer, proliferative diseases, tumors and their
25 metastases.

Infectious disease

The immune system in higher vertebrates represents the first line of defense against various antigens that can enter the vertebrate body, including microorganisms such
30 as bacteria, fungi and viruses that are the causative agents of a variety of diseases.

Despite large immunization programs, viral infections, such as influenza virus, human immunodeficiency virus ("HIV"), herpes simplex virus ("HSV", type 1 or 2), human papilloma virus ("HPV", type 16 or 18), human cytomegalovirus ("HCMV") or human
35 hepatitis B or C virus ("HBV", Type B; "HCV", type C) infections, remain a serious source of morbidity and mortality throughout the world and a significant cause of illness and death among people with immune-deficiency associated with aging or different clinical conditions. Although antiviral chemotherapy with compounds such as amantadine and rimantadine have been shown to reduce the duration of

symptoms of clinical infections (i.e., influenza infection), major side effects and the emergence of drug-resistant variants have been described. New classes of antiviral agents designed to target particular viral proteins such as influenza neuraminidase are being developed. However, the ability of viruses to mutate the target proteins represents an obstacle for effective treatment with molecules which selectively inhibit the function of specific viral polypeptides. Thus, there is need for new therapeutic strategies to prevent and treat viral infections.

Additionally, there is a need for new therapies for the prevention and treatment of bacterial infections, especially bacterial infections caused by multiple drug resistant bacteria. Currently, bacterial infections are treated with various antibiotics. Although antibiotics have and can be effective in the treatment of various bacterial infections, there are a number of limitations to the effectiveness and safety of antibiotics. For example, some individuals have an allergic reaction to certain antibiotics and other individuals suffer from serious side effects. Moreover, continued use of antibiotics for the treatment of bacterial infections contributes to formation of antibiotic-resistant strains of bacteria.

Another aspect of the present invention is directed to the use of the peptide or the peptide combination for prophylaxis and/or treatment of infectious diseases including opportunistic infections.

Examples of infectious diseases are AIDS, alveolar hydatid disease (AHD, echinococcosis), amebiasis (*Entamoeba histolytica* infection), *Angiostrongylus* infection, anisakiasis, anthrax, babesiosis (*Babesia* infection), *Balantidium* infection (balantidiasis), *Baylisascaris* infection (raccoon roundworm), bilharzia (schistosomiasis), *Blastocystis hominis* infection (blastomycosis), boreliosis, botulism, Brainerd diarrhea, brucellosis, bovine spongiform encephalopathy (BSE), candidiasis, capillariasis (*Capillaria* infection), chronic fatigue syndrome (CFS), Chagas disease (American trypanosomiasis), chickenpox (*Varicella-Zoster* virus), *Chlamydia pneumoniae* infection, cholera, Creutzfeldt-Jakob disease (CJD), clonorchiasis (*Clonorchis* infection), cutaneous larva migrans (CLM) (hookworm infection), coccidioidomycosis, conjunctivitis, Coxsackievirus A16 (hand, foot and mouth disease), cryptococcosis, *Cryptosporidium* infection (cryptosporidiosis), *Culex* mosquito (West Nile virus vector), cyclosporiasis (*Cyclospora* infection), cysticercosis (neurocysticercosis), Cytomegalovirus infection, Dengue / Dengue fever, *Dipylidium* infection (dog and cat flea tapeworm), Ebola virus hemorrhagic fever, encephalitis, *Entamoeba coli* infection, *Entamoeba dispar* infection, *Entamoeba hartmanni* infection, *Entamoeba histolytica* infection (amebiasis),

Entamoeba polecki infection, enterobiasis (pinworm infection), enterovirus infection (non-polio), Epstein-Barr virus infection, Escherichia coli infection, foodborne infection, foot and mouth disease, fungal dermatitis, gastroenteritis, group A streptococcal disease, group B streptococcal disease, Hansen's disease (leprosy),
5 Hantavirus pulmonary syndrome, head lice infestation (pediculosis), Helicobacter pylori infection, hematologic disease, Hendra virus infection, hepatitis (HCV, HBV), herpes zoster (shingles), HIV Infection, human ehrlichiosis, human parainfluenza virus infection, influenza, isosporiasis (Isospora infection), Lassa fever, leishmaniasis, Kala-azar (Kala-azar, Leishmania Infection), lice (body lice, head lice,
10 pubic lice), Lyme disease, malaria, Marburg hemorrhagic fever, measles, meningitis, mosquito-borne diseases, Mycobacterium avium complex (MAC) infection, Naegleria infection, nosocomial infections, nonpathogenic intestinal ameobae infection, onchocerciasis (river blindness), opisthorciasis (Opisthorcis infection), parvovirus infection, plague, Pneumocystis carinii pneumonia (PCP),
15 polio, Q fever, rabies, respiratory syncytial virus (RSV) Infection, rheumatic fever, Rift Valley fever, river blindness (onchocerciasis), rotavirus infection, roundworm infection, salmonellosis, salmonella enteritidis, scabies, shigellosis, shingles, sleeping sickness, smallpox, streptococcal Infection, tapeworm infection (Taenia infection), tetanus, toxic shock syndrome, tuberculosis, ulcers (peptic ulcer
20 disease), valley fever, Vibrio parahaemolyticus infection, Vibrio vulnificus infection, viral hemorrhagic fever, warts, waterborne infectious diseases, West Nile virus infection (West Nile encephalitis), whooping cough, yellow fever.

Another aspect of the present invention is directed to the use of the peptide or the
25 peptide combination for prophylaxis and/or treatment of prion diseases.

Prions are infectious agents which do not have a nucleic acid genome. It seems that a protein alone is the infectious agent. A prion has been defined as "small proteinaceous infectious particle which resists inactivation by procedures that modify
30 nucleic acids". The discovery that proteins alone can transmit an infectious disease came as a considerable surprise to the scientific community. Prion diseases are often called "transmissible spongiform encephalopathies", because of the post mortem appearance of the brain with large vacuoles in the cortex and cerebellum. Probably most mammalian species develop these diseases. Prion diseases are a
35 group of neurodegenerative disorders of humans and animals and the prion diseases can manifest as sporadic, genetic or infectious disorders. Examples of prion diseases acquired by exogenous infection are bovine spongiform encephalitis (BSE) of cattle and the new variant of Creutzfeld-Jakob disease (vCJD) caused by BSE as well as scrapie of animals. Examples of human prion diseases include kuru,

sporadic Creutzfeldt-Jakob disease (sCJD), familial CJD (fCJD), iatrogenic CJD (iCJD), Gerstmann-Sträussler-Scheinker (GSS) disease, fatal familial insomnia (FFI), and especially the new variant CJD (nvCJD or vCJD).

- 5 The name "prion" is used to describe the causative agents which underlie the transmissible spongiform encephalopathies. A prion is proposed to be a novel infectious particle that differs from viruses and viroids. It is composed solely of one unique protein that resists most inactivation procedures such as heat, radiation, and proteases. The latter characteristic has led to the term protease-resistant isoform of the prion protein. The protease-resistant isoform has been proposed to slowly catalyze the conversion of the normal prion protein into the abnormal form.

The term "isoform" in the context of prions means two proteins with exactly the same amino acid sequence that can fold into molecules with dramatically different tertiary structures. The normal cellular isoform of the prion protein (PrP^C) has a high α -helix content, a low β -sheet content, and is sensitive to protease digestion. The abnormal, disease-causing isoform (PrP^{Sc}) has a lower α -helix content, a much higher β -sheet content, and is much more resistant to protease digestion.

- 20 As used herein the term "prion diseases" refers to transmissible spongiform encephalopathies. Examples for prion diseases comprise scrapie (sheep, goat), transmissible mink encephalopathy (TME; mink), chronic wasting disease (CWD; muledeer, deer, elk), bovine spongiform encephalopathy (BSE; cows, cattles), Creutzfeld-Jacob Disease (CJD), variant CJD (vCJD), sporadic Creutzfeldt-Jakob disease (sCJD), familial CJD (fCJD), iatrogenic CJD (iCJD, Gerstmann-Sträussler-Scheinker syndrome (GSS), fatal familial insomnia (FFI), and kuru. Preferred are BSE, vCJD, and CJD.

AIDS

- 30 AIDS stands for Acquired Immune Deficiency Syndrome. AIDS is the name historically chosen for a new medical syndrome which is essentially 100% fatal. Human Immunodeficiency Virus 1 (HIV-1) is responsible for the cause of AIDS. HIV-1 is a lentivirus. The HIV-1 preferably infects CD4+ cells which get lost in AIDS patients. In a person with AIDS, up to 13% of lymphocytes and monocytes in the blood were found infected. From 93% to 100% of the time, infectious virus and viral DNA could be recovered from the blood of asymptomatic people who were HIV-1 positive, and 100% of the time from people with AIDS.

Lentiviruses

The following table 1 gives an overview of the members of the lentivirus family.

Table 1: Lentiviruses

Virus	Host	Diseases
EIAV	Horse	anemia, wasting
VMV	Sheep	pneumonia, wasting, arthritis, mastitis, encephalitis
CAEV	Goat	arthritis, mastitis, encephalitis
BIV	Cattle	none
FIV	Cat	immunodeficiency, encephalitis, wasting
SIVs	various African monkey species	none
HIV-1, HIV-2	Humans	immunodeficiency, pneumonia, encephalopathy, wasting, gastroenteropathy, nephropathy

- 5 With respect to the numbers of HIV-1 infected individuals, each day about 16,000 people worldwide are infected with the virus, with 95% of new cases occurring in developing countries (mostly in Africa). More than 2.6 million people die of AIDS annually. According to the American Medical Association, about 45 million people worldwide are infected with HIV-1. Of the more than 2.6 million people died of AIDS-
 10 associated diseases annually, an estimated 550,000 were children under 15 years of age. Although life expectancy increased very significantly by introduction of new treatments, effective control of the AIDS pandemic remains elusive.

- 15 The origin of HIV-1 among nonhuman primates has been traced to a simian virus, SIVcpz, which infected several geographically isolated chimpanzee communities in southern Cameroon. This HIV-1 progenitor probably was passed from chimpanzees to human hunters through bloodborne transmission. Phylogenetic analysis of HIV-1 and related viruses from nonhuman primates suggest that three independent transmission events early in the 20th century spawned three HIV-1 groups: major (M,
 20 between 1915 and 1941), outlier (O), and nonmajor and nonoutlier (N). Although strains related to the M and N groups have been found in chimpanzees, recent evidence suggests that group O HIV-1 may have originated in gorillas, in which the closest relatives of this group have been identified. It is speculated that the virus then spread among humans along the Congo River into Kinshasa, Zaire, where the
 25 earliest documented case of HIV-1 infection (with group M strain) in humans has been traced to a blood sample from 1959.

HIV-1 has several intrinsic mechanisms that ensure rapid viral evolution. The reverse transcriptase of HIV lacks proofreading activity, the ability to confirm that the DNA transcript it makes is an accurate copy of the RNA code, and confers a mutation rate of approximately 3.4×10^{-5} mutations per base pair per replication cycle. Since the HIV genome is an estimated 10^4 base pairs in length and the baseline rate of viral production is approximately 10^{10} virions per day, millions of viral variants are produced within any infected person in a single day. HIV-1 recombination can lead to further viral diversity and occurs when one person is coinfectd with two separate strains of the virus that are multiplying in the same cell.

HIV-1 is a complex retrovirus encoding in addition to the structural proteins gag, pol, and env several regulatory genes. Specifically, the trans-acting regulator of viral gene expression, called Rev, is crucially important for HIV replication. The HIV-1 Rev protein is a nuclear phosphoprotein accumulating at steady state in the nucleoli and has the capacity to shuttle between the nuclear and cytoplasmic compartments. Within the Rev protein distinct biological domains exist that are essential for its biological activity. The cis-acting target for Rev is a highly structured sequence element termed the Rev Response Element (RRE) located on HIV-1 RNA. Upon binding to the HIV-1 RNA as a monomer, Rev has to multimerize and subsequently interact with cellularly encoded cofactors on its way to export the HIV-1 RNA from the nucleus to the cytoplasm where protein translation occurs. It was shown that the eukaryotic initiation factor 5A (eIF-5A) directly and specifically binds the HIV-1 Rev NES, and serves as a HIV genome carrier to the cellular ribosomes.

HIV-1 Classification

Group M is the predominant circulating HIV-1 group. It has been divided into subtypes, denoted with letters, and sub-subtypes, denoted with numerals. Subtypes A1, A2, A3, A4, B, C, D, F1, F2, G, H, J, and K are currently recognized. HIV-1 subtypes, also called clades, are phylogenetically linked strains of HIV-1 that are approximately the same genetic distance from one another; in some cases, subtypes are also linked geographically or epidemiologically. Genetic variation within a subtype can be 15 to 20%, whereas variation between subtypes is usually 25 to 35%. Over the past decade, advances in full-genome sequencing of HIV have led to the identification of circulating and unique recombinant forms (CRFs and URFs, respectively). These are the result of recombination between subtypes within a dually infected person, from whom the recombinant forms are then passed to other people. The recombinant progeny are classified as circulating recombinant forms if they are

identified in three or more people with no direct epidemiologic linkage; otherwise they are described as unique recombinant forms.

Classification	Definition	Examples
Subtypes or clades	Genetically related HIV-1 strains that are essentially phylogenetically equidistant, generating a starlike, rather than a treelike, phylogeny	Subtypes A, B, C, D, F, G, H, J, and K are currently known; A through D are highly prevalent, others have low prevalence and limited geographic distributions
Sub-subtypes	Distinct lineages within a subtype; genetic distance between sub-subtypes is smaller than that between subtypes	Subtypes A and F are subdivided into sub-subtypes A1 through A4 and F1 and F2, respectively; mostly these circulate in Central and West Africa
Intersubtype recombinant forms	Mosaic strains with segments from two or more subtypes alternating across the genome	Common in mixed-subtype epidemics; thought to result from infection of a person with more than one HIV-1 subtype
Circulating recombinant forms	Specific recombinant forms that are spreading in a population; new forms are defined when three people without direct epidemiologic linkage are found to be infected; the assigned name reflects sequence of discovery and subtype composition, with "cpx" indicating forms containing three or more subtypes	Currently, 43 forms are described; CRF01_AE and CRF02_AG are found principally in Southeast Asia and West Africa, respectively; others have more limited distributions
Unique recombinant forms	Intersubtype recombinant forms recovered from only a single person	Hundreds of forms have been described on the basis of partial or complete genome sequences; their potential for epidemic spread is unknown
Geographically distinct lineages	Lineages, often country-specific, that are distinguishable phylogenetically; unlike sub-subtypes, they are not phylogenetically equidistant within subtypes	Thai B, Indian C, West vs. East African D, and Former Soviet Union A (FSU-A)

5 Table 2: Phylogenetic Classifications of HIV-1

Receptor Use by HIV-1 Subtypes, and Disease Progression

Differential characteristics of viral subtypes and their interactions with the human host influence HIV transmission and disease progression. The HIV strains capable of using the chemokine coreceptor CCR5 (R5 viruses) are more frequently transmitted than strains that use the CXCR4 coreceptor (X4 viruses); X4 viruses emerge later in infected patients and are associated with more rapid disease progression. All HIV-1 subtypes can use both coreceptors, but subtype D may be dual-tropic (i.e., an R5X4 virus) most frequently. The percentage of X4 virus appears to be lower in subtype C than in subtype B, even when the viruses are obtained from patients with advanced AIDS. Early data on mother-to-child transmission implied that subtype C was transmitted more frequently than subtype B. Subtype differences result in variable rates of disease progression. HIV-1 disease progresses more rapidly, and the risk of death is greater among persons infected with subtype D, with recombinant forms, or with multiple subtypes than among those infected with subtype A. The propensity of subtype D to exhibit a greater degree of dual-tropism than other subtypes may help to

explain the observation that subtype D appears to be associated with a more rapid rate of disease progression than other HIV-1 subtypes.

Interactions between the host and HIV-1 that vary according to subtype are also important. The known differences in HIV-1 transmission and disease progression in hosts carrying specific HLA class I types may vary according to infecting HIV-1 subtype. In an infected person, T cells specific to HIV-1 can exhibit cross-subtype specificity and recognize viral epitopes within subtypes other than the one that generated the initial response. CD8+ T cells obtained from persons infected with subtype B recognize viral epitopes within conserved regions of the consensus sequences from genomes of subtypes A, B, and C. However, the immune response tends to be greatest against the infecting subtype, and CD8+ T-cell responses can wane over time.

Table 3: Response to Therapy

Currently there is a variety of medicaments available to treat HIV-infections.

FDA Antiretroviral Approval Dates				
Drug	Brand name	Sponsor	Approval date	Class
AZT (zidovudine)	Retrovir	Burroughs-Wellcome (now GlaxoSmithKline)	19 Mar 1987	NRTI
ddl (didanosine)	Videx	Bristol-Myers (now Bristol-Myers Squibb)	9 Oct 1991	NRTI
ddC (zalcitabine)*	HIVID*	Hoffmann-LaRoche	19 Jun 1992	NRTI
d4T (stavudine)	Zerit	Bristol-Myers Squibb (BMS)	24 Jun 1994	NRTI
3TC (lamivudine)	Epivir	GlaxoSmithKline (GSK)	17 Nov 1995	NRTI
saquinavir (SQV)	Invirase	Hoffmann-LaRoche	6 Dec 1995	PI
ritonavir (RTV)	Norvir	Abbott	1 Mar 1996	PI
indinavir (IDV)	Crixivan	Merck	13 Mar 1996	PI
nevirapine	Viramune	Boehringer Ingelheim	21 Jun 1996	NNRTI
nelfinavir (NFV)	Viracept	Agouron (now Pfizer)	14 Mar 1997	PI
delavirdine (DLV)	Rescriptor	Pharmacia & Upjohn (now Pfizer)	4 Apr 1997	NNRTI
AZT/3TC	Combivir	GSK	27 Sep 1997	NNRTI
saquinavir sgc*	Fortovase*	Hoffmann-LaRoche	7 Nov 1997	PI
efavirenz (EFV)	Sustiva	DuPont (now BMS)	17 Sep 1998	NNRTI
abacavir (ABC)	Ziagen	GSK	17 Dec 1998	NRTI
amprenavir	Agenerase	GSK	15 Apr 1999	PI
lopinavir (LPV)/ritonavir	Kaletra	Abbott	15 Sep 2000	PI
ddl enteric coated	Videx EC	BMS	31 Oct 2000	NRTI
AZT/3TC/ABC	Trizivir	GSK	14 Nov 2000	NRTI
tenofovir DF (TDF)	Viread	Gilead	26 Oct 2001	NRTI
enfuvirtide (T-20)	Fuzeon	Hoffmann-LaRoche / Trimeris	13 Mar 2003	EI
atazanavir	Reyataz	BMS	20 Jun 2003	PI
FTC (emtricitabine)	Emtriva	Gilead	2 Jul 2003	NRTI
ABC/3TC	Epzicom	GSK	2 Aug 2003	NRTI
FTC/tenofovir DF	Truvada	Gilead	2 Aug 2003	NRTI
fosamprenavir	Lexiva	GSK	20 Oct 2003	PI
tipranavir	Aptivus	Boehringer Ingelheim	22 Jun 2005	PI
darunavir	Prezista	Johnson & Johnson/Tibotec	23 Jun 2006	PI
TDF/3TC/efavirenz	Atripla	Gilead/BMS [Gilead/Merck in much of world]	12 Jul 2006	2 NRTI/1 NNRTI

* soft gel cap saquinavir (Fortovase) and zalcitabine (HIVID) were discontinued by Roche on 15 February 2006.

Nevertheless, the question rises whether HIV-1 subtypes influence the response to antiretroviral treatment, as only 12% of global infections are caused by the most studied subtype, B; and 50% of prevalent HIV infections and 47% of all new HIV-1 infections are with subtype C. This discrepancy in the availability of clinical data for non-B subtypes is exacerbated by the fact that, until the past few years, antiretroviral

treatment had been largely unavailable in many countries with non-B subtypes of HIV-1.

Initial data from treatment cohorts in Africa raised two concerns: first that certain subtypes of HIV-1 might spread or progress more rapidly than others, making treatment decisions more challenging, and second, that the data on baseline antiretroviral susceptibility derived from studies of subtype B may not be applicable to non-B subtypes. This concern is illustrated by HIV type 2 and group O strains of HIV-1, which possess intrinsic resistance to non-nucleoside reverse-transcriptase inhibitors.

Due to instantly high medical need, new compounds or vaccines are being developed to treat or prevent HIV-1 infections:

Table 4:

HIV Antiretroviral Drug Pipeline 2007

HIV Drugs in Clinical Trials*	
Phase II	Phase III
Maraviroc (Celsentri): CCR5 antagonist— <i>Pfizer</i>	
Raltegravir (Isentress, MK-0518): HIV integrase inhibitor— <i>Merck</i>	
Etravirine (TMC-125): NNRTI— <i>Tibotec</i>	
Rilpivirine (TMC-278): NNRTI— <i>Tibotec</i>	
Elvitegravir (GS-9137): HIV integrase inhibitor— <i>Gilead</i>	
Vicriviroc: CCR5 antagonist— <i>Schering</i>	
Bevirimat (PA-457): Maturation inhibitor— <i>Panacos</i>	
TNX-355: Anti-CD4 monoclonal antibody— <i>Genentech (Tanox)</i>	
BILR 355 BS: NNRTI— <i>Boehringer-Ingelheim</i>	
Racivir: NRTI— <i>Pharmasset</i>	
Dexelvucitabine (Reverset): NRTI— <i>Pharmasset</i>	
Amdoxovir: NRTI— <i>RFS Pharma</i>	
Apricitabine (AVX754): NRTI— <i>Avexa</i>	
Elvucitabine: NRTI— <i>Achillion Pharmaceuticals</i>	

DISCONTINUED: BrecaNavir (VX-385): Protease inhibitor—*GlaxoSmithKline*.

* This and subsequent tables only display compounds in phases II and beyond; compounds in

Table 5:

HIV Vaccines Pipeline 2007

HIV Vaccines in Clinical Trials	
Phase II	Phase III
ALVAC vCP1521: Canarypox— <i>Aventis Pasteur</i>	
AIDSVAX B/E: Protein— <i>VaxGen</i>	
MRK Ad5 trivalent: Adenovirus— <i>Merck</i>	
VRC HIV DNA 016/Ad5 boost: DNA plasmids— <i>NIH Vaccine Research Center</i>	
ALVAC vCP1452: Canarypox— <i>Aventis Pasteur</i>	
LIPO-5: Lipopeptides— <i>ANRS; Aventis</i>	
tgAAC09 AAV: Parvovirus— <i>Targeted Genetics; IAVI</i>	
ISS P-001: Tat protein— <i>ISS; ICAV; AIDS Vaccine Integrated Project</i>	

Table 6:

PrEP & Microbicides Pipeline 2007

Phase II	Phase III
Other Prevention Technologies in Clinical Trials	
Pre-Exposure Prophylaxis (PrEP)	
Tenofovir (Viread): Antiretroviral— <i>Gilead</i>	
Tenofovir/FTC (Truvada): Antiretroviral— <i>Gilead</i>	
Microbicides	
Carraguard: Barrier (adsorption inhibitor)— <i>Population Council</i>	
PRO 2000: Barrier (adsorption inhibitor)— <i>Indevus Pharmaceuticals</i>	
BufferGel: pH buffer/barrier— <i>Reprotect</i>	
Tenofovir DF: Antiretroviral— <i>CONRAD; Intl. Partnership for Microbicides</i>	
TMC 120 (daprivirine): Antiretroviral— <i>Intl. Partnership for Microbicides</i>	
VivaGel (SPL 7013): Fusion inhibitor— <i>Starpharma</i>	

Table 7:

HIV Therapeutic Vaccines Pipeline 2007

Therapeutic HIV Vaccines in Clinical Trials
Phase II
ALVAC vCP1452: Canarypox—Aventis Pasteur
Gag, Nef, Pol lipopeptides: Peptides—ANRS; Aventis
CD4-specific T-Cell Vaccine: Autologous T cells—Soroka Medical Center, Israel
DCV-2: Autologous dendritic cells—Hospital Clinic of Barcelona
AGS-004: Autologous dendritic cells—Argyros Therapeutics; ACTG
LC002 (DermaVir): DNA—Research Institute for Genetic & Human Therapy

Table 8:

Cytokine, Immunomodulator, and Gene Therapy Pipeline 2007

Cytokines, Immunomodulators, and Gene Therapies in Clinical Trials	
Phase II	Phase III
Interleukin-2 (Proleukin, IL-2): Cytokine—Novartis	
Human growth hormone (Serostim, HGHr): Growth hormone—Serono	
Cyclosporine A: Immunosuppressive—ACTG; AIEDRP	
VRX496: Ex vivo gene transfer—VIRxSYS	
OZ1: Gene therapy—J&J	
HGTV43: Ex vivo gene transfer—Enzo Biochem	
Palifermin (keratinocyte growth factor): Growth factor—Amgen	

5

Resistance to Antiretroviral Therapy

HIV resistance to antiretroviral therapy can be divided into two categories: primary resistance, which reflects acquisition of a drug-resistant strain of HIV by a newly infected person; and secondary, or acquired, resistance, which develops after a period of HIV treatment.

10

The Highly Active Antiretroviral Therapy (HAART), consisting of Nucleoside Analog Reverse Transcriptase Inhibitors (NRTIs), Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs) and Protease Inhibitors (PIs) is accompanied by quick emergence of resistant viral strains. In some patients, replication of resistant HIV-1 leads to emergence of more virulent variants of HIV-1, which are associated with an accelerated loss of CD4+ cells and confer a significantly increased risk of disease progression and death. Resistance emerges as a consequence of the selective pressure of incompletely suppressive therapy and is determined by mutations in the HIV-1 reverse transcriptase and protease genes. Primary mutations alter the binding of the drug to its target resulting in an increase in the amount of drug necessary to inhibit the enzyme. Secondary mutations increase resistance by improving the fitness of viruses carrying primary infections. In the case of NRTIs, the mutation M148V is an example of a key mutation leading to high-level resistance. Additional positions of mutations include M41L, K70R, T215Y (Zidovudine), L74V (Didanosine), T69D (Zalcitabine), Y115F (Abacavir). Multidrug resistance conferring cross-resistance to the entire NRTI class is well recognized (Q151M and insertion mutation T69SSS). Cross-resistance is also extensive among the three currently used NNRTIs Efavirenz, Nevirapine and Delavirdine which makes them inactive against the virus expressing the K103N mutation in the reverse transcriptase gene. The K103N mutation acts by inhibiting formation of the drug-binding pocket. For this class of drugs, the "first shot" is most frequently the "only shot". Also, cross-resistance between the PIs is rather a rule than an exception (Indinavir and Ritonavir have almost identical resistance patterns - K20M, V32I, M36I, M46I, I54V, L63P, A71V, V82A/T/F, I84V, L90M - and Saquinavir-resistant strains are also cross-resistant to other PIs - L10I, K20M, I84V, L90M). Even for anti HIV-1 drugs like T20, a 36 amino acid peptide derived from the HIVgp41 protein which inhibits fusion of the virus to the host cell, resistance has already been demonstrated. In addition, patients on HAART face long-term side effects, including pancreatitis, peripheral neuropathies, hepatotoxicity, diabetes, or metabolic abnormalities in body fat redistribution, as well as in the glucose metabolism and the cardiovascular field. For many patients, who experience the above mentioned drug toxicities, a structured treatment interruption is an unavoidable necessity ("Drug holidays"). However, during this time the suppressed virus may bounce back.

The inventive peptide or the peptide combination described in the present invention can be specifically combined with Hydroxyurea; with Integrase inhibitors like antraquinones, quinalizarin, L-chicoric acid, dicaffeoylquinic acid; with Zinc Finger inhibitors like dithiane compounds; with immunomodulators like Interleukin-2, Interferon-alpha, Interferon-beta, GM-CSF, G-CSF; with therapeutical vaccine strategies including live, attenuated and replication incompetent virus; killed, inactivated virus; envelope subunit protein; core subunit protein; peptides; nucleic

acids of the respective retroviruses, specifically of HIV-1 or HIV-2; with antiretroviral gene-therapy approaches like antisense or dominant-negative Rev mutants.

Listed below are diseases associated with HIV-1 infections:

Malignancies

- 5 • Anal Dysplasia/Cancer
- Cervical Dysplasia/Cancer
- Kaposi's Sarcoma (KS)
- Lymphomas

Neurological Conditions

- 10 • AIDS Dementia Complex
- Peripheral Neuropathy

Other Conditions and Complications

- 15 • Aphthous Ulcers
- Depression & Anxiety
- Fatigue and Anemia
- Nausea & Diarrhea
- Thrombocytopenia
- 20 • Wasting Syndrome & Lipodystrophy

Individuals with HIV-1 infection are at risk for both primary and secondary neurological and neurobehavioral disorders. Primary disorders occur as a result of the influence of HIV-1 on the central nervous system (CNS); whereas, secondary disorders usually occur as a result of immune system deficiencies or treatment effects. Neurobehavioral disorders associated with HIV-1 infection may be complicated by preexisting or new onset psychological and emotional disorders. Clinical research in neurological and particularly neurobehavioral disorders associated with HIV-1 infection has been complicated by methodological issues and controversies as well as ill-defined nosology.

Table 9: AIDS-related diseases are listed as follows:

Opportunistic disease or malignancy

Aspergillosis
Atypical mycobacteriosis
Bacteraemia
Candidiasis
CMV
Cryptococcosis
Cryptosporidiosis – Isosporiasis
Enteritis, non-specific ⁴
Herpes (systemic)
Histoplasmosis
Kaposi sarcoma
Lymphoma
Nocardiosis
Penicilliosis
Progressive multifocal leukoencephalopathy (PML) or HIV encephalitis
<i>Pneumocystis carinii</i> pneumonia
Pneumonia
Toxoplasmosis
Tuberculosis

In the following information from current research on HIV-1 related neurological disorders, diagnosis, and treatment with focus on the AIDS related neurobehavioral disorders and a brief review of secondary HIV-related neurological disorders are presented. The peptide or the peptide combination of the present invention are also useful for prophylaxis and/or treatment of the opportunistic diseases described below.

HIV-1 Related Neurological and Neurobehavioral Disorders

10 Primary Disorders

HIV-1 can directly invade the central nervous system (CNS). Viral infection in the CNS is most often seen in mono-nuclear microglial cells and multi-nucleated giant cells. Neuronal loss is usually secondary to the presence of HIV-1 in surrounding cells and not in the neurons themselves. The process by which neuronal death occurs is speculative, although proposed mechanisms include the production of cytokines that interfere with neuronal function, production of abnormal neurotransmitter metabolites that are neurotoxic, and the presence of certain viral fragments that interfere with neurotransmitter transmission.

20 HIV-1 associated CNS disorders include the neurobehavioral disorders, HIV associated minor cognitive disorder and HIV associated dementia, and the neurological disorder, HIV associated myelopathy.

HIV Associated Dementia

HIV-1 associated dementia (HAD) is a progressive disorder that initially presents as apathy, inertia, cognitive slowing, memory loss, and social withdrawal. As it progresses, multiple cognitive functions become increasingly impaired. The terminal phases are characterized by global cognitive impairment, mutism, and severe psychomotor retardation.

The peptide or the peptide combination of the present invention were tested using the assays described in Examples 1-7 for their effect as active therapeutic agents in the prophylaxis and/or treatment of infectious diseases and disorders.

Autoimmune disease

Autoimmune disease refers to any of a group of diseases or disorders in which tissue injury is associated with a humoral and/or cell-mediated immune response to body constituents or, in a broader sense, an immune response to self. The pathological immune response may be systemic or organ specific. That is, for example, the immune response directed to self may affect joints, skin, myelin sheath that protects neurons, kidney, liver, pancreas, thyroid, adrenals, and ovaries.

In fact, the list of autoimmune diseases is composed of more than eighty disorders. A few autoimmune diseases such as vitiligo, in which patches of skin lose pigmentation, are merely annoying. Most others are debilitating, often progressive with time and eventually fatal. Systemic lupus erythematosus (SLE), for example, is a chronic disease in which 10-15% of patients die within a decade of diagnosis, in all but a few autoimmune diseases, the sex ratio skews towards women. For example, in SLE the ratio of female to male patients is nine to one. In one particular case, Hashimoto's disease in which the immune system attacks the thyroid gland, the ratio is fifty to one.

It has long been known that immune complex formation plays a role in the etiology and progression of autoimmune disease. For example, inflammation in patients with arthritis has long been considered to involve phagocytosis by leukocytes of complexes of antigen, antibody and complement-immune complexes. However, only now it is being recognized that inflammation caused by immune complexes in the joints (arthritis), the kidneys (glomerulonephritis), and blood vessels (vasculitis) is a major cause of morbidity in autoimmune diseases. Increased immune complex formation correlates with the presence of antibodies directed to self or so-called

autoantibodies, and the presence of the latter can also contribute to tissue inflammation either as part of an immune complex or unbound to antigen (free antibody). In some autoimmune diseases, the presence of free autoantibody contributes significantly to disease pathology. This has been clearly demonstrated for example in SLE (anti-DNA antibodies), immune thrombocytopenia (antibody response directed to platelets), and to a lesser extent rheumatoid arthritis (IgG reactive rheumatoid factor). The important role of immune complexes and free autoantibodies is further demonstrated by the fact that successful treatment of certain autoimmune diseases has been achieved by the removal of immune complexes and free antibody by means of specific immunoadsorption procedures. For example, the use of an apheresis procedure in which immune complexes and antibodies are removed by passage of a patient's blood through an immunoaffinity column was approved by the U.S. FDA in 1987 for immune thrombocytopenia (ITP) and in 1999 for rheumatoid arthritis. However, currently there is no approved method for the treatment of autoimmune diseases which facilitates the elimination of immune complexes and autoantibodies by administration of a drug.

Another aspect of the etiology and progression of autoimmune disease is the role of proinflammatory cytokines. Under normal circumstances, proinflammatory cytokines such as tumor necrosis factor α (TNF α) and interleukin-1 (IL-1) play a protective role in the response to infection and cellular stress. However, the pathological consequences which result from chronic and/or excessive production of TNF α and IL-1 are believed to underlie the progression of many autoimmune diseases such as rheumatoid arthritis, Crohn's disease, inflammatory bowel disease, and psoriasis. Other proinflammatory cytokines include interleukin-6, interleukin-8, interleukin-17, and granulocyte-macrophage colony stimulating factor.

Naturally occurring CD4+CD25+ regulatory T cells (Tregs) play a critical role in the control of periphery tolerance to self-antigens. Interestingly, they also control immune responses to allergens and transplant antigens. Recent studies in animal models have shown that adoptive transfer of CD4+CD25+ Tregs can prevent or even cure allergic and autoimmune diseases, and appear to induce transplantation tolerance. Thus, adoptive cell therapy using patient-specific CD4+CD25+ Tregs has emerged as an individualized medicine for the treatment of inflammatory disease including allergy, autoimmune disease and transplant rejection. Furthermore, strategies to activate and expand antigen-specific CD4+CD25+ Tregs in vivo using pharmacological agents may represent a novel avenue for drug development.

The interaction of leukocytes with the vessel endothelium to facilitate the extravasation into the tissue represents a key process of the body's defense mechanisms. Excessive recruitment of leukocytes into the inflamed tissue in chronic diseases like autoimmune disorders could be prevented by interfering with the mechanisms of leukocyte extravasation. Significant progress in elucidating the molecular basis of the trafficking of leukocytes from the blood stream to the extravascular tissue has been achieved that enables new strategies for therapeutic approaches. The multistep process of leukocyte rolling, firm adhesion and transmigration through the endothelial wall is facilitated by a dynamic interplay of adhesion receptors on both leukocytes and on endothelial cells as well as chemokines. In preclinical studies using various animal models, promising results have been obtained demonstrating that blocking of adhesion receptors of the selectin and integrin families improved the inflammation process in models of ulcerative colitis, autoimmune encephalomyelitis or contact hypersensitivity. In addition to the targeting of adhesion receptors by antibodies, small molecules that mimic epitopes of adhesion receptor ligands have been developed and successfully applied in animal models. Clinical studies revealed a limited response using antibodies to selectins or leukocyte function-associated antigen 1 (LFA-1) integrins compared with animal models. However, using humanized antibodies to the alpha 4-integrin subunit significant efficacy has been demonstrated in autoimmune diseases like psoriasis, multiple sclerosis and inflammatory bowel disease.

Examples of autoimmune diseases of the eyes are idiopathic opticus-neuritis, ophthalmia sympathica, anterior uveitis and other uveitis forms, retina degeneration, and Mooren's ulcer.

Examples of autoimmune diseases of the skin are bullous pemphigoides, chronic urticaria (autoimmune subtype), dermatitis herpetiformis (morbus Duhring), epidermolysis bullosa acquisita (EBA), acquired angioedema, herpes gestationes, hypocomplementemic urticarial vasculitis syndrome (HUVS), linear IgA-dermatosis, and pemphigus.

Examples of hematological autoimmune diseases are autoimmune hemolytic anemia, autoimmune neutropenia, Evans syndrome, inhibitor hemophilia, idiopathic thrombocytopenial purpura (ITP) and pernicious anemia.

Examples of gynecological autoimmune diseases are habitual abortion and infertility.

Examples of autoimmune diseases of the heart are congenital heart block, idiopathic dilatative cardiomyopathy, peripartum-cardiomyopathy, postcardiotomy syndrome, and postinfarct syndrome (Dressler syndrome).

- 5 Examples of autoimmune diseases of the ear, nose and throat are chronic sensorineural hearing loss and morbus Menière.

Examples of autoimmune diseases of the colon are autoimmune enteropathy, colitis ulcerosa, indeterminant colitis, Crohn's disease and gluten-sensitive enteropathy.

10

Examples of autoimmune endocrinological autoimmune disorders are autoimmune polyglandular syndrome type 1, autoimmune polyglandular syndrome type 2, diabetes mellitus type 1 (IDDM), Hashimoto-thyroiditis, insulin-autoimmune-syndrome (IAS), idiopathic diabetes insipidus, idiopathic hypoparathyroidism, idiopathic Addison's disease and Graves-Basedow disease.

15

Examples of autoimmune diseases of the liver are autoimmune hepatitis (AIH type 1, 2 and 3), primary biliary cirrhosis (PBC), and primary sclerosing cholangitis.

- 20 Example of autoimmune diseases of the lung is Goodpasture's syndrome.

An example of an autoimmune disease of the stomach is chronic atrophic (type A) gastritis.

- 25 Examples of neurological autoimmune disorders are Guillain-Barré syndrome, IgM gammopathy-associated neuropathy, Lambert-Eaton syndrome, Miller-Fisher syndrome, multiple sclerosis, multifocal motoric neuropathy, myasthenia gravis, paraneoplastic neurological syndrome, Rasmussen's encephalitis, and stiff-man syndrome.

30

Examples of autoimmune diseases of the kidney are anti-TBM-nephritis, Goodpasture's syndrome/anti-GBM-nephritis, IgA-nephropathy, interstitial nephritis, and membrane proliferative glomerulonephritides.

- 35 Further diseases that may be caused by an autoimmune reaction are Behcet disease, chronic fatigue immune dysfunction syndrome (CFIDS), Cogan syndrome I, endometriosis, HELLP syndrome, Bechterew's disease, polymyalgia rheumatica, psoriasis, sarcoidosis and vitiligo.
- 40

During the last decade, new biotherapies have been developed for the treatment of systemic autoimmune diseases. The targets of these new treatments are all the steps of the immune response. These new therapies are: B lymphocyte (BL) inhibitors such as anti-CD20 monoclonal antibody, B lymphocyte stimulator (BLyS) antagonists and tolerogens of pathogenic-antibody secreting LB; inhibitors of the costimulation between antigen-presenting cells and T lymphocyte (TL) like monoclonal anti-CD40 ligand antibody or CTLA4-Ig (abatecept); TL antagonists which can inhibit the proliferation of autoreactive T cells; cytokine antagonists; chemokine and adhesion antagonists which inhibit trafficking of immunocompetent cells to target organs. These new approaches are based on a better understanding of the autoimmune response.

The peptide or the peptide combination of the present invention were tested using the assays described in Examples 14 - 15 for their effect as active therapeutic agents in the prophylaxis and/or treatment of autoimmune diseases and disorders.

Fibrotic disease

Fibrosis or fibrosis associated disorder affects the liver, epidermis, endodermis, muscle, tendon, cartilage, heart, pancreas, lung, uterus, nervous system, testis, ovary, adrenal gland, artery, vein, colon, small intestine, biliary tract, or stomach. In a further embodiment, the fibrosis or fibrosis associated disorder is interstitial lung fibrosis. In another embodiment the fibrosis or fibrosis associated disorder is the result of an infection with schistosoma. In another embodiment the fibrosis or fibrosis associated disorder is the result of wound healing.

Fibrosis is generally characterized by the pathologic or excessive accumulation of collagenous connective tissue. Fibrotic diseases and disorders include, but are not limited to, collagen disease, interstitial lung disease, human fibrotic lung disease (e.g., obliterative bronchiolitis, idiopathic pulmonary fibrosis,) pulmonary fibrosis from a known etiology, tumor stroma in lung disease, systemic sclerosis affecting the lungs, Hermansky-Pudlak syndrome, coal worker's pneumoconiosis, asbestosis, silicosis, chronic pulmonary hypertension, AIDS associated pulmonary hypertension, sarcoidosis, and the like), fibrotic vascular disease, tubulointerstitial and glomerular fibrosis, myocardial fibrosis, arterial sclerosis, atherosclerosis, varicose veins, coronary infarcts, cerebral infarcts, myocardial fibrosis, musculoskeletal fibrosis, post-surgical adhesions, human kidney disease (e.g., nephritic syndrome, Alport's syndrome, HIV associated nephropathy, polycystic kidney disease, Fabry's disease, diabetic nephropathy, chronic glomerulonephritis, nephritis associated with systemic lupus, and the like), cutis keloid formation, progressive systemic sclerosis (PSS),

primary sclerosing cholangitis (PSC), liver fibrosis, liver cirrhosis, renal fibrosis, pulmonary fibrosis, cystic fibrosis, chronic graft versus host disease, scleroderma (local and systemic), Grave's ophthalmopathy, diabetic retinopathy, glaucoma, Peyronie's disease, penis fibrosis, urethrostenosis after a test using a cystoscope, inner accretion after surgery, scarring, myelofibrosis, idiopathic retroperitoneal fibrosis, peritoneal fibrosis from a known etiology, drug induced ergotism, fibrosis incident to benign or malignant cancer, fibrosis incident to microbial infection (e.g., viral, bacterial, parasitic, fungal, etc.), Alzheimer's disease, fibrosis incident to inflammatory bowel disease (including stricture formation in Crohn's disease and microscopic colitis), fibrosis induced by chemical or environmental insult (e.g., cancer chemotherapy, pesticides, radiation/cancer radiotherapy), and the like.

Diseases associated with fibrosis include lupus, graft versus host disease, scleroderma, systemic sclerosis, scleroderma-like disorders, sine scleroderma, calcinosis, Raynaud's esophageal dysfunction, sclerodactyly, telangiectasiae, hypersensitivity pneumonitis, collagen vascular disease, asthma, pulmonary arterial hypertension, glomerulonephritis, chronic obstructive pulmonary disease, fibrosis following myocardial infarction, central nervous system fibrosis following a stroke or neuro-degenerative diseases (e.g. Alzheimer's disease), proliferative vitreoretinopathy (PVR) and arthritis, silicosis, asbestos induced pulmonary fibrosis, acute lung injury and acute respiratory distress syndrome (including bacterial pneumonia induced, trauma induced, viral pneumonia induced, tuberculosis, ventilator induced, non-pulmonary sepsis induced, and aspiration induced).

Increased number of activated myofibroblasts in fibrotic diseases

The emergence and disappearance of the myofibroblast appears to correlate with the initiation of active fibrosis and its resolution, respectively. In addition, the myofibroblast has many phenotypic features, which embody much of the pathologic alterations in fibrotic tissue, e.g. lung tissue. These features would seem to argue for an important role for the myofibroblast in the pathogenesis of fibrosis, e.g. lung fibrosis. Furthermore, the persistence of the myofibroblast may herald progressive disease, and, conversely, its disappearance may be an indicator of resolution. This in turn suggests that future therapeutic strategies targeting the myofibroblast would be productive.

Patients usually exhibit evidence of active fibrosis with increased numbers of activated fibroblasts, many of which have the phenotypic characteristics of myofibroblasts. At these sites, increased amounts of extracellular matrix deposition are evident with effacement of the normal alveolar architecture. Animal model

studies show the myofibroblast to be the primary source of type I collagen gene expression in active fibrotic sites. *In vitro* studies show differentiation of these cells from fibroblasts under the influence of certain cytokines but indicate their susceptibility to nitric oxide mediated apoptosis. In addition to promoting myofibroblast differentiation, transforming growth factor- β 1 (TGF- β 1) provides protection against apoptosis. Thus, this well-known fibrogenic cytokine is important both for the emergence of the myofibroblast and its survival against apoptotic stimuli. This is consistent with the critical importance of this cytokine in diverse models of fibrosis in various tissues. In view of these properties, the persistence or prolonged survival of the myofibroblast may be the key to understanding why certain forms of lung injury may result in progressive disease, terminating in end stage disease.

Although pulmonary fibrosis has diverse etiologies, there is a common feature characteristic of this process, namely, the abnormal deposition of extracellular matrix that effaces the normal lung tissue architecture. A key cellular source of this matrix is the mesenchymal cell population that occupies much of the fibrotic lesion during the active period of fibrosis. This population is heterogeneous with respect to a number of key phenotypes. One of these phenotypes is the myofibroblast, which is commonly identified by its expression in α -smooth muscle actin and by features that are intermediate between the bona fide smooth muscle cell and the fibroblast. The *de novo* appearance of myofibroblasts at sites of wound healing and tissue repair/fibrosis is associated with the period of active fibrosis and is considered to be involved in wound contraction. Furthermore, the localization of myofibroblasts at sites undergoing active extracellular matrix deposition suggests an important role for these cells in the genesis of the fibrotic lesion.

Increased TGF- β 1 family levels in fibrotic diseases

The transforming growth factor- β 1 (TGF- β 1) family of proteins has the most potent stimulatory effect on extracellular matrix deposition of any cytokines so far examined. In animal models of pulmonary fibrosis enhanced TGF- β 1 gene expression is temporally and spatially related to increased collagen gene expression and protein deposition. TGF- β 1 antibodies reduce collagen deposition in murine bleomycin-induced lung fibrosis and human fibrotic lung tissue shows enhanced TGF- β 1 gene and protein expression. Several lines of evidence suggest that TGF- β is a central regulator of pulmonary fibrosis. Several animal models over expressing TGF- β showed extensive progressive fibrosis but limited inflammation, indicating that TGF- β may play a predominant role in the progression of pulmonary fibrosis. Therapeutic efforts are therefore focusing on inhibition of TGF- β activity, for instance by anti-TGF- β 1-antibodies, or modulators of TGF- β 1 such as pirfenidone. Pirfenidone inhibits

TGF- β 1 gene expression *in vivo* resulting in inhibition of TGF- β 1-mediated collagen synthesis and appears to slow progression of IPF in patients. Other novel, promising antifibrotic agents include relaxin (inhibits TGF- β -mediated overexpression of collagen and increases collagenases), suramin (inhibits growth factors), prostaglandin E2 (inhibits collagen production) and lovastatin (blocks formation of granulation tissue by induction of fibroblast apoptosis).

Diseases involving the lung associated with increased levels of TGF- β include chronic lung disease of prematurity, idiopathic pulmonary fibrosis, rapid progressive pulmonary fibrosis, giant-cell interstitial pneumonia, acute rejection after lung transplantation, cytomegalovirus pneumonitis after lung transplantation, bronchiolitis obliterans, asbestosis, coal worker's pneumoconiosis, silicosis, histiocytosis, sarcoidosis, eosinophilic granuloma, scleroderma, systemic lupus erythematosus, lymphangioleiomyomatosis, central fibrosis in pulmonary adenocarcinoma, cystic fibrosis, chronic obstructive lung disease, and asthma.

Increased TNF- α levels in fibrotic diseases

An important role of tumor necrosis factor- α (TNF- α) in interstitial fibrosis has been established using transgenic mice, which either overexpress or display a deficiency of this cytokine. Mice transgenically modified to overexpress TNF- α develop lung fibrosis. In contrast, mice null for TNF- α show marked resistance to bleomycin induced fibrosis. TNF- α can stimulate fibroblast replication and collagen synthesis *in vitro*, and pulmonary TNF- α gene expression rises after administration of bleomycin in mice. Soluble TNF- α receptors reduce lung fibrosis in murine models and pulmonary overexpression of TNF- α in transgenic mice is characterized by lung fibrosis. In patients with CFA or asbestosis, bronchoalveolar lavage fluid-derived macrophages release increased amounts of TNF- α compared with controls.

Increased TNF- α may induce fibrosis or fibrosis-associated conditions affecting any tissue including, for example, fibrosis of an internal organ, a cutaneous or dermal fibrosing disorder, and fibrotic conditions of the eye. Fibrosis of internal organs (e.g., liver, lung, kidney, heart blood vessels, gastrointestinal tract) occurs in disorders such as pulmonary fibrosis, idiopathic fibrosis, autoimmune fibrosis, myelofibrosis, liver cirrhosis, veno-occlusive disease, mesangial proliferative glomerulonephritis, crescentic glomerulonephritis, diabetic nephropathy, renal interstitial fibrosis, renal fibrosis in subjects receiving cyclosporin, allograft rejection, HTV associated nephropathy. Other fibrosis-associated disorders include systemic sclerosis, eosinophilia-myalgia syndrome, and fibrosis-associated CNS disorders such as intraocular fibrosis. Dermal fibrosing disorders include, for example, scleroderma,

morphea, keloids, hypertrophic scars, familial cutaneous collagenoma, and connective tissue nevi of the collagen type. Fibrotic conditions of the eye include conditions such as diabetic retinopathy, post-surgical scarring (for example, after glaucoma filtering surgery and after crossed-eyes (strabismus) surgery), and proliferative vitreoretinopathy. Additional fibrotic conditions that may be treated by the methods of the present invention may result, for example, from rheumatoid arthritis, diseases associated with prolonged joint pain and deteriorated joints; progressive systemic sclerosis, polymyositis, dermatomyositis, eosinophilic fascitis, morphea, Raynaud's syndrome, and nasal polyposis.

Increased matrix metalloproteases levels in fibrotic diseases

The abnormal extracellular matrix (ECM) remodeling observed in the lungs of patients with interstitial pulmonary fibrosis (IPF) is due, at least in part, to an imbalance between matrix metalloproteases (MMPs) and tissue inhibitor of metalloproteinases (TIMPs). Normal lung fibroblasts do not make MMP-9 in vitro, whereas fibroblasts from IPF lungs strongly express MMP-9. In addition, fibroblasts from patients with IPF express increased levels of all TIMPs. In this setting, TIMPs may play a role in apoptosis in some cell populations. In vitro studies of alveolar macrophages obtained from untreated patients with idiopathic pulmonary fibrosis showed marked increase in MMP-9 secretion compared to macrophages collected from healthy individuals. In animals models of bleomycin-induced pulmonary fibrosis MMPs have been shown to be elevated in bronchoalveolar lavage (BAL) fluid. Indeed, a synthetic inhibitor of MMP, Batimastat, has been shown to significantly reduce bleomycin-induced lung fibrosis, again pointing to the importance of MMPs in the development of this fibrotic disease in the lung. A number of studies have shown that the actions of MMPs can result in the release of growth factors and cytokines. These profibrotic factors require proteolytic processing for their activation or release from extracellular matrix or carrier proteins before they can exert their activity. In fact, the proteolytic activity processing of several key factors involved in the pathogenesis of pulmonary fibrosis such as insulin-like growth factor (IGF), TGF- β_1 and TNF- α occur through the actions of MMPs, thereby activating or releasing them from inhibitory protein-protein interactions. For example, IGFs in vivo are sequestered by six high affinity IGF binding proteins (IGFBPs1-6), preventing their ability to interact with IGF receptors. Studies examining adults and children IPF and interstitial lung disease show that beside IPF, IGFBP-3 and IFPB-2 levels are increased in IPF BAL fluid. MMPs have recently been shown to regulate the cleavage of IGF binding proteins, thereby liberating the complexed ligand to affect IGF actions in target cells. Observations have also shown that the gelatinases, MMP-9 and MMP-2 may be involved in proteolytic activation of latent TGF- β

complexes. Furthermore, the MMP inhibitor Batimastat reduces MMP-9 activity in BAL fluid, which was associated with decreased amount of TGF- β and TNF- α .

Pulmonary fibrosis can be an all too common consequence of an acute inflammatory response of the lung to a host of inciting events. Chronic lung injury due to fibrotic changes can result from an identifiable inflammatory event or an insidious, unknown event. The inflammatory process can include infiltration of various inflammatory cell types, such as neutrophils and macrophages, the secretion of inflammatory cytokines and chemokines and the secretion of matrix remodeling proteinases.

Increased CCL18 levels in fibrotic diseases

The expression and regulation of cysteine-cysteine (CC) chemokine ligand 18 (CCL18), a marker of alternative activation, by human alveolar macrophages (AMs) is increased in patients with pulmonary fibrosis and correlates negatively with pulmonary function test parameters. Thus, CCL18 is an ideal diagnostic marker for pulmonary fibrosis.

The peptide or the peptide combination of the present invention were tested using the assays described in Examples 14 - 15 for their effect as active therapeutic agents in the prophylaxis and/or treatment of fibrotic diseases and disorders.

Inflammatory disease

Inflammation is the final common pathway of various insults, such as infection, trauma, and allergies to the human body. It is characterized by the activation of the immune system with recruitment of inflammatory cells, production of pro-inflammatory cells and production of pro-inflammatory cytokines. Most inflammatory diseases and disorders are characterized by abnormal accumulation of inflammatory cells including monocytes/macrophages, granulocytes, plasma cells, lymphocytes and platelets. Along with tissue endothelial cells and fibroblasts, these inflammatory cells release a complex array of lipids, growth factors, cytokines and destructive enzymes that cause local tissue damage.

One form of inflammatory response is neutrophilic inflammation which is characterized by infiltration of the inflamed tissue by neutrophil polymorphonuclear leukocytes (PMN), which are a major component of the host defense. Tissue infection by extracellular bacteria represents the prototype of this inflammatory response. On the other hand, various non-infectious diseases are characterized by extravascular recruitment of neutrophils. This group of inflammatory diseases includes chronic obstructive pulmonary disease, adult respiratory distress syndrome,

some types of immune-complex alveolitis, cystic fibrosis, bronchitis, bronchiectasis, emphysema, glomerulonephritis, rheumatoid arthritis, gouty arthritis, ulcerative colitis, certain dermatoses such as psoriasis and vasculitis. In these conditions neutrophils are thought to play a crucial role in the development of tissue injury which, when persistent, can lead to the irreversible destruction of the normal tissue architecture with consequent organ dysfunction. Tissue damage is primarily caused by the activation of neutrophils followed by their release of proteinases and increased production of oxygen species.

Chronic obstructive pulmonary disease (COPD) is described by the progressive development of airflow limitation that is not fully reversible. Most patients with COPD have three pathological conditions; bronchitis, emphysema and mucus plugging. This disease is characterized by a slowly progressive and irreversible decrease in forced expiratory volume in the first second of expiration (FEV_i), with relative preservation of forced vital capacity (FVC). In both asthma and COPD there is significant, but distinct, remodeling of airways. Most of the airflow obstruction is due to two major components, alveolar destruction (emphysema) and small airways obstruction (chronic obstructive bronchitis). COPD is mainly characterized by profound mucus cell hyperplasia. Neutrophil infiltration of the patient's lungs is a primary characteristic of COPD. Elevated levels of proinflammatory cytokines, like TNF- α , and especially chemokines like interleukin-8 (IL-8) and growth-regulated oncogene- α (GRO- α) play a very important role in pathogenesis of this disease. Platelet thromboxane synthesis is also enhanced in patients with COPD. Most of the tissue damage is caused by activation of neutrophils followed by their release of metalloproteinases, and increased production of oxygen species.

TNF- α has several biologic activities that are important in homeostasis as well as in pathophysiological conditions. The main sources of TNF- α are monocytes-macrophages, T-lymphocytes and mast cells. The finding that anti-TNF- α antibodies (cA2) are effective in the treatment of patients suffering from rheumatoid arthritis (RA) intensified the interest to find new TNF- α inhibitors as possible potent medicaments for RA. Rheumatoid arthritis is an autoimmune chronic inflammatory disease characterized by irreversible pathological changes of the joints. In addition to RA, TNF- α antagonists are also applicable to several other pathological conditions and diseases such as spondylitis, osteoarthritis, gout and other arthritic conditions, sepsis, septic shock, toxic shock syndrome, atopic dermatitis, contact dermatitis, psoriasis, glomerulonephritis, lupus erythematosus, scleroderma, asthma, cachexia, chronic obstructive lung disease, congestive heart failure, insulin resistance, lung

(pulmonary) fibrosis, multiple sclerosis, Crohn's disease, ulcerative colitis, viral infections and AIDS.

The term "immunoinflammatory disorder" encompasses a variety of conditions, including autoimmune diseases, proliferative skin diseases, and inflammatory dermatoses. Immunoinflammatory disorders result in the destruction of healthy tissue by an inflammatory process, dysregulation of the immune system, and unwanted proliferation of cells. Examples of immunoinflammatory disorders are acne vulgaris; acute respiratory distress syndrome; Addison's disease; allergic rhinitis; allergic intraocular inflammatory diseases, antineutrophil cytoplasmic antibody (ANCA)-associated small-vessel vasculitis; ankylosing spondylitis; arthritis, asthma; atherosclerosis; atopic dermatitis; autoimmune hepatitis; autoimmune hemolytic anemia; autoimmune hepatitis; Behcet's disease; Bell's palsy; bullous pemphigoid; cerebral ischemia; chronic obstructive pulmonary disease; cirrhosis; Cogan's syndrome; contact dermatitis; COPD; Crohn's disease; Cushing's syndrome; dermatomyositis; diabetes mellitus; discoid lupus erythematosus; eosinophilic fasciitis; erythema nodosum; exfoliative dermatitis; fibromyalgia; focal glomerulosclerosis; focal segmental glomerulosclerosis; giant cell arteritis; gout; gouty arthritis; graft versus host disease; hand eczema; Henoch-Schonlein purpura; herpes gestationis; hirsutism; idiopathic cerato-scleritis; idiopathic pulmonary fibrosis; idiopathic thrombocytopenic purpura; immune thrombocytopenic purpura; inflammatory bowel or gastrointestinal disorders, inflammatory dermatoses; lichen planus; lupus nephritis; lymphomatous tracheobronchitis; macular edema; multiple sclerosis; myasthenia gravis; myositis; nonspecific fibrosing lung disease; osteoarthritis; pancreatitis; pemphigoid gestationis; pemphigus vulgaris; periodontitis; polyarteritis nodosa; polymyalgia rheumatica; pruritus scroti; pruritis/inflammation, psoriasis; psoriatic arthritis; pulmonary histoplasmosis; rheumatoid arthritis; relapsing polychondritis; rosacea caused by sarcoidosis; rosacea caused by scleroderma; rosacea caused by Sweet's syndrome; rosacea caused by systemic lupus erythematosus; rosacea caused by urticaria; rosacea caused by zoster-associated pain; sarcoidosis; scleroderma; segmental glomerulosclerosis; septic shock syndrome; shoulder tendinitis or bursitis; Sjogren's syndrome; Still's disease; stroke-induced brain cell death; Sweet's disease; systemic lupus erythematosus; systemic sclerosis; Takayasu's arteritis; temporal arteritis; toxic epidermal necrolysis; transplant-rejection and transplant-rejection-related syndromes; tuberculosis; type-1 diabetes; ulcerative colitis; uveitis; vasculitis; and Wegener's granulomatosis.

As used herein, "non-dermal inflammatory disorders" include, for example, rheumatoid arthritis, inflammatory bowel disease, asthma, and chronic obstructive

pulmonary disease. By "dermal inflammatory disorders" or "inflammatory dermatoses" is meant an inflammatory disorder selected from psoriasis, guttate psoriasis, inverse psoriasis, pustular psoriasis, erythrodermic psoriasis, acute febrile neutrophilic dermatosis, eczema, asteatotic eczema, dyshidrotic eczema, vesicular
5 palmo-plantar eczema, acne vulgaris, atopic dermatitis, contact dermatitis, allergic contact dermatitis, dermatomyositis, exfoliative dermatitis, hand eczema, pompholyx, rosacea, rosacea caused by sarcoidosis, rosacea caused by scleroderma, rosacea caused by Sweet's syndrome, rosacea caused by systemic lupus erythematosus, rosacea caused by urticaria, rosacea caused by zoster-associated pain, Sweet's
10 disease, neutrophilic hidradenitis, sterile pustulosis, drug eruptions, seborrheic dermatitis, pityriasis rosea, cutaneous kikuchi disease, pruritic urticarial papules and plaques of pregnancy, Stevens-Johnson syndrome and toxic epidermal necrolysis, tattoo reactions, Wells syndrome (eosinophilic cellulitis), reactive arthritis (Reiter's syndrome), bowel-associated dermatosis-arthritis syndrome, rheumatoid neutrophilic
15 dermatosis, neutrophilic eccrine hidradenitis, neutrophilic dermatosis of the dorsal hands, balanitis circumscripta plasmacellularis, balanoposthitis, Behcet's disease, erythema annulare centrifugum, erythema dyschromicum perstans, erythema multiforme, granuloma annulare, hand dermatitis, lichen nitidus, lichen planus, lichen sclerosus et atrophicus, lichen simplex chronicus, lichen spinulosus, nummular
20 dermatitis, pyoderma gangrenosum, sarcoidosis, subcorneal pustular dermatosis, urticaria, and transient acantholytic dermatosis.

By "proliferative skin disease" is meant a benign or malignant disease that is characterized by accelerated cell division in the epidermis or dermis. Examples of
25 proliferative skin diseases are psoriasis, atopic dermatitis, nonspecific dermatitis, primary irritant contact dermatitis, allergic contact dermatitis, basal and squamous cell carcinomas of the skin, lamellar ichthyosis, epidermolytic hyperkeratosis, premalignant keratosis, acne, and seborrheic dermatitis. As will be appreciated by one skilled in the art, a particular disease, disorder, or condition may be
30 characterized as being both a proliferative skin disease and an inflammatory dermatosis. An example of such a disease is psoriasis.

Symptoms and signs of inflammation associated with specific conditions include:

- rheumatoid arthritis:- pain, swelling, warmth and tenderness of the involved joints;
35 generalized and morning stiffness;
- insulin-dependent diabetes mellitus-insulinitis; this condition can lead to a variety of complications with an inflammatory component, including:- retinopathy, neuropathy, nephropathy; coronary artery disease, peripheral vascular disease, and cerebrovascular disease;

- autoimmune thyroiditis:- weakness, constipation, shortness of breath, puffiness of the face, hands and feet, peripheral edema, bradycardia;
- multiple sclerosis:- spasticity, blurry vision, vertigo, limb weakness, paresthesias;
- uveoretinitis:- decreased night vision, loss of peripheral vision;
- 5 • lupus erythematosus:- joint pain, rash, photosensitivity, fever, muscle pain, puffiness of the hands and feet, abnormal urinalysis (hematuria, cylinduria, proteinuria), glomerulonephritis, cognitive dysfunction, vessel thrombosis, pericarditis;
- scleroderma:- Raynaud's disease; swelling of the hands, arms, legs and face; skin
- 10 thickening; pain, swelling and stiffness of the fingers and knees, gastrointestinal dysfunction, restrictive lung disease; pericarditis; renal failure;
- other arthritic conditions having an inflammatory component such as rheumatoid spondylitis, osteoarthritis, septic arthritis and polyarthritis:- fever, pain, swelling, tenderness;
- 15 • other inflammatory brain disorders, such as meningitis, Alzheimer's disease, AIDS dementia encephalitis:- photophobia, cognitive dysfunction, memory loss;
- other inflammatory eye inflammations, such as retinitis:- decreased visual acuity;
- inflammatory skin disorders, such as , eczema, other dermatites (e.g., atopic, contact), psoriasis, burns induced by UV radiation (sun rays and similar UV
- 20 sources):- erythema, pain, scaling, swelling, tenderness;
- inflammatory bowel disease, such as Crohn's disease, ulcerative colitis:- pain, diarrhea, constipation, rectal bleeding, fever, arthritis;
- asthma:- shortness of breath, wheezing;
- other allergy disorders, such as allergic rhinitis:- sneezing, itching, runny nose
- 25 • conditions associated with acute trauma such as cerebral injury following stroke-sensory loss, motor loss, cognitive loss;
- heart tissue injury due to myocardial ischemia:- pain, shortness of breath;
- lung injury such as that which occurs in adult respiratory distress syndrome:- shortness of breath, hyperventilation, decreased oxygenation, pulmonary infiltrates;
- 30 • inflammation accompanying infection, such as sepsis, septic shock, toxic shock syndrome:- fever, respiratory failure, tachycardia, hypotension, leukocytosis;
- other inflammatory conditions associated with particular organs or tissues, such as:
- (i) nephritis (e.g., glomerulonephritis):-oliguria, abnormal urinalysis;
- (ii) inflamed appendix:- fever, pain, tenderness, leukocytosis;
- 35 (iii) gout:- pain, tenderness, swelling and erythema of the involved joint, elevated serum and/or urinary uric acid;
- (iv) inflamed gall bladder:- abdominal pain and tenderness, fever, nausea, leukocytosis;
- (v) congestive heart failure:- shortness of breath, rales, peripheral edema;

(vi) Type II diabetes:- end organ complications including cardiovascular, ocular, renal, and peripheral vascular disease;

(vii) lung (pulmonary) fibrosis:- hyperventilation, shortness of breath, decreased oxygenation;

5 (viii) vascular disease, such as atherosclerosis and restenosis:- pain, loss of sensation, diminished pulses, loss of function; and

(ix) alloimmunity leading to transplant rejection:- pain, tenderness, fever.

10 A human peptide is "active" in an inflammatory disease if the inhibition is > 50% in one of the assays described below. Inhibition (as percentage) was calculated using the following formula: % inhibition = (1 - concentration of cytokines in sample/concentration of cytokines in positive control) x 100. The positive control refers to stimulated samples, not treated with substances.

15 The peptide or the peptide combination of the present invention were tested using the assays described in Examples 1-7, 9-17 for their effect as active therapeutic agents in the prophylaxis and/or treatment of inflammatory diseases and disorders.

Neurodegenerative disease

20 The present invention also relates generally to the fields of neurology and psychiatry and to methods of protecting the cells of a mammalian central nervous system from damage or injury.

25 Injuries or trauma of various kinds to the central nervous system (CNS) or the peripheral nervous system (PNS) can produce profound and long-lasting neurological and/or psychiatric symptoms and disorders. One form that this can take is the progressive death of neurons or other cells of the central nervous system (CNS), i.e., neurodegeneration or neuronal degeneration.

30 Neuronal degeneration as a result of, for example; Alzheimer's disease, multiple sclerosis, cerebral-vascular accidents (CVAs)/stroke, traumatic brain injury, spinal cord injuries, degeneration of the optic nerve, e.g., ischemic optic neuropathy or retinal degeneration and other central nervous system disorders is an enormous medical and public health problem by virtue of both its high incidence and the
35 frequency of long-term sequelae. Animal studies and clinical trials have shown that amino acid transmitters (especially glutamate), oxidative stress and inflammatory reactions contribute strongly to cell death in these conditions. Upon injury or upon ischemic insult, damaged neurons release massive amounts of the neurotransmitter glutamate, which is excitotoxic to the surrounding neurons. Glutamate is a negatively

charged amino acid that is an excitatory synaptic transmitter in the mammalian nervous system. Although the concentration of glutamate can reach the millimolar range in nerve terminals its extracellular concentration is maintained at a low level to prevent neurotoxicity. It has been noted that glutamate can be toxic to neurons if presented at a high concentration. The term "excitotoxicity" has been used to describe the cytotoxic effect that glutamate (and other such excitatory amino acids) can have on neurons when applied at high dosages.

Patients with injury or damage of any kind to the central (CNS) or peripheral (PNS) nervous system including the retina may benefit from neuroprotective methods. This nervous system injury may take the form of an abrupt insult or an acute injury to the nervous system as in, for example, acute neurodegenerative disorders including, but not limited to; acute injury, hypoxia-ischemia or the combination thereof resulting in neuronal cell death or compromise. Acute injury includes, but is not limited to, traumatic brain injury (TBI) including, closed, blunt or penetrating brain trauma, focal brain trauma, diffuse brain damage, spinal cord injury, intracranial or intravertebral lesions (including, but not limited to, contusion, penetration, shear, compression or laceration lesions of the spinal cord or whiplash shaken infant syndrome).

In addition, deprivation of oxygen or blood supply in general can cause acute injury as in hypoxia and/or ischemia including, but not limited to, cerebrovascular insufficiency, cerebral ischemia or cerebral infarction (including cerebral ischemia or infarctions originating from embolic occlusion and thrombosis, retinal ischemia (diabetic or otherwise), glaucoma, retinal degeneration, multiple sclerosis, toxic and ischemic optic neuropathy, reperfusion following acute ischemia, perinatal hypoxic-ischemic injury, cardiac arrest or intracranial hemorrhage of any type (including, but not limited to, epidural, subdural, subarachnoid or intracerebral hemorrhage).

Trauma or injury to tissues of the nervous system may also take the form of more chronic and progressive neurodegenerative disorders, such as those associated with progressive neuronal cell death or compromise over a period of time including, but not limited to, Alzheimer's disease, Pick's disease, diffuse Lewy body disease, progressive supranuclear palsy (Steel-Richardson syndrome), multisystem degeneration (Shy-Drager syndrome), chronic epileptic conditions associated with neurodegeneration, motor neuron diseases (amyotrophic lateral sclerosis), multiple sclerosis, degenerative ataxias, cortical basal degeneration, ALS-Parkinson's-dementia complex of Guam, subacute sclerosing panencephalitis, Huntington's disease, Parkinson's disease, synucleinopathies (including multiple system atrophy), primary progressive aphasia, striatonigral degeneration, Machado-Joseph disease or

spinocerebellar ataxia type 3 and olivopontocerebellar degenerations, bulbar and pseudobulbar palsy, spinal and spinobulbar muscular atrophy (Kennedy's disease), primary lateral sclerosis, familial spastic paraplegia, Werdnig-Hoffmann disease, Kugelberg-Welander disease, Tay-Sach's disease, Sandhoff disease, familial spastic
5 disease, Wohlfart-Kugelberg-Welander disease, spastic paraparesis, progressive multifocal leukoencephalopathy, familial dysautonomia (Riley-Day syndrome) or prion diseases (including, but not limited to Creutzfeld-Jakob disease, Gerstmann-Strussler-Scheinker disease, Kuru disease or fatal familial insomnia).

10 In addition, trauma and progressive injury to the nervous system can take place in various psychiatric disorders, including but not limited to, progressive, deteriorating forms of bipolar disorder or schizoaffective disorder or schizophrenia, impulse control disorders, obsessive compulsive disorder (OCD), behavioral changes in temporal lobe epilepsy and personality disorders.

15 In one preferred embodiment the compounds of the invention would be used to provide neuroprotection in disorders involving trauma and progressive injury to the nervous system in various psychiatric disorders. These disorders would be selected from the group consisting of; schizoaffective disorder, schizophrenia, impulse control
20 disorders, obsessive compulsive disorder (OCD) and personality disorders.

In addition, trauma and injury make take the form of disorders associated with overt and extensive memory loss including, but not limited to, neurodegenerative disorders associated with age-related dementia, vascular dementia, diffuse white matter
25 disease (Binswanger's disease), dementia of endocrine or metabolic origin, dementia of head trauma and diffuse brain damage, dementia pugilistica or frontal lobe dementia, including but not limited to Pick's Disease.

Other disorders associated with neuronal injury include, but are not limited to,
30 disorders associated with chemical, toxic, infectious and radiation injury of the nervous system including the retina, injury during fetal development, prematurity at time of birth, anoxic-ischemia, injury from hepatic, glycemic, uremic, electrolyte and endocrine origin, injury of psychiatric origin (including, but not limited to, psychopathology, depression or anxiety), injury from peripheral diseases and
35 plexopathies (including plexus palsies) or injury from neuropathy (including neuropathy selected from multifocal, sensory, motor, sensory-motor, autonomic, sensory-autonomic or demyelinating neuropathies (including, but not limited to Guillain-Barre syndrome or chronic inflammatory demyelinating polyradiculoneuropathy) or those neuropathies originating from infections,

inflammation, immune disorders, drug abuse, pharmacological treatments, toxins, trauma (including, but not limited to compression, crush, laceration or segmentation traumas), metabolic disorders (including, but not limited to, endocrine or paraneoplastic), Charcot-Marie-Tooth disease (including, but not limited to, type 1a, 1b, 2, 4a or 1-X linked), Friedreich's ataxia, metachromatic leukodystrophy, Refsum's disease, adrenomyeloneuropathy, ataxia-telangiectasia, Djerine-Sottas (including, but not limited to, types A or B), Lambert-Eaton syndrome or disorders of the cranial nerves).

Further indications are cognitive disorders. The term "cognitive disorder" shall refer to anxiety disorders, delirium, dementia, amnestic disorders, dissociative disorders, eating disorders, mood disorders, schizophrenia, psychotic disorders, sexual and gender identity disorders, sleep disorders, somatoform disorders, acute stress disorder, obsessive-compulsive disorder, panic disorder, posttraumatic stress disorder, specific phobia, social phobia, substance withdrawal delirium, Alzheimer's disease, Creutzfeldt-Jakob disease, head trauma, Huntington's disease, HIV disease, Parkinson's disease, Pick's disease, learning disorders, motor skills disorders, developmental coordination disorder, communication disorders, phonological disorder, pervasive developmental disorders, Asperger's disorder, autistic disorder, childhood disintegrative disorder, Rett's disorder, pervasive developmental disorder, attention-deficit/hyperactivity disorder (ADHD), conduct disorder, oppositional defiant disorder, pica, rumination disorder, tic disorders, chronic motor or vocal tic disorder, Tourette's disorder, elimination disorders, encopresis, enuresis, selective mutism, separation anxiety disorder, dissociative amnesia, depersonalization disorder, dissociative fugue, dissociative identity disorder, anorexia nervosa, bulimia nervosa, bipolar disorders, schizophreniform disorder, schizoaffective disorder, delusional disorder, psychotic disorder, shared psychotic disorder, delusions, hallucinations, substance-induced psychotic disorder, orgasmic disorders, sexual pain disorders, dyspareunia, vaginismus, sexual dysfunction, paraphilias, dyssomnias, breathing-related sleep disorder, circadian rhythm sleep disorder, hypersomnia, insomnia, narcolepsy, dyssomnia, parasomnias, nightmare disorder, sleep terror disorder, sleepwalking disorder, parasomnia, body dysmorphic disorder, conversion disorder, hypochondriasis, pain disorder, somatization disorder, alcohol related disorders, amphetamine related disorders, caffeine related disorders, cannabis related disorders, cocaine related disorders, hallucinogen related disorders, inhalant related disorders, nicotine related disorders, opioid related disorders, phencyclidine-related disorder, abuse, persisting amnestic disorder, intoxication, withdrawal.

The term "bipolar and clinical disorders" shall refer to adjustment disorders, anxiety disorders, delirium, dementia, amnesic and other cognitive disorders, disorders usually first diagnosed in infancy (e.g.), childhood, or adolescence, dissociative disorders (e.g. dissociative amnesia, depersonalization disorder, dissociative fugue and dissociative identity disorder), eating disorders, factitious disorders, impulse-control disorders, mental disorders due to a general medical condition, mood disorders, other conditions that may be a focus of clinical attention, personality disorders, schizophrenia and other psychotic disorders, sexual and gender identity disorders, sleep disorders, somatoform disorders, substance-related disorders, generalized anxiety disorder (e.g. acute stress disorder, posttraumatic stress disorder), panic disorder, phobia, agoraphobia, obsessive-compulsive disorder, stress, acute stress disorder, anxiety neurosis, nervousness, phobia, posttraumatic stress disorder, posttraumatic stress disorder (PTSD), abuse, obsessive-compulsive disorder (OCD), manic depressive psychosis, specific phobias, social phobia, adjustment disorder with anxious features.

Examples for disorders usually first diagnosed in infancy, childhood, or adolescence are: mental retardation, learning disorders, mathematics disorder, reading disorder, disorder of written expression, motor skills disorders, developmental coordination disorder, communication disorders, expressive language disorder, phonological disorder, mixed receptive-expressive language disorder, stuttering, pervasive developmental disorders, Asperger's disorder, autistic disorder, childhood disintegrative disorder, Rett's disorder, pervasive developmental disorder, attention-deficit/hyperactivity disorder (ADHD), conduct disorder, oppositional defiant disorder, feeding disorder of infancy or early childhood, pica, rumination disorder, tic disorders, chronic motor or vocal tic disorder, Tourette's syndrome, elimination disorders, encopresis, enuresis, selective mutism, separation anxiety disorder, reactive attachment disorder of infancy or early childhood, stereotypic movement disorder.

Examples for substance-related disorders are: alcohol related disorders, amphetamine related disorders, caffeine related disorders, cannabis related disorders, cocaine related disorders, hallucinogen related disorders, inhalant related disorders, nicotine related disorders, opioid related disorders, psychotic disorder, psychotic disorder, phencyclidine-related disorder, abuse, persisting amnesic disorder, anxiety disorder, persisting dementia, dependence, intoxication, intoxication delirium, mood disorder, psychotic disorder, withdrawal, withdrawal delirium, sexual dysfunction, sleep disorder.

The term "neuroprotection" as used herein shall mean; inhibiting, preventing, ameliorating or reducing the severity of the dysfunction, degeneration or death of nerve cells, axons or their supporting cells in the central or peripheral nervous system of a mammal, including a human. This includes the treatment or prophylaxis of a neurodegenerative disease; protection against excitotoxicity or ameliorating the cytotoxic effect of a compound (for example, a excitatory amino acid such as glutamate; a toxin; or a prophylactic or therapeutic compound that exerts an immediate or delayed cytotoxic side effect including but not limited to the immediate or delayed induction of apoptosis) in a patient in need thereof.

The term "a patient in need of treatment with a neuroprotective drug " as used herein will refer to any patient who currently has or may develop any of the above syndromes or disorders, or any disorder in which the patient's present clinical condition or prognosis could benefit from providing neuroprotection to prevent the development, extension, worsening or increased resistance to treatment of any neurological or psychiatric disorder.

The term "treating" or "treatment" as used herein, refers to any indicia of success in the prevention or amelioration of an injury, pathology or condition, including any objective or subjective parameter such as abatement; remission; diminishing of symptoms or making the injury, pathology, or condition more tolerable to the patient; slowing in the rate of degeneration or decline; making the final point of degeneration less debilitating; or improving a subject's physical or mental well-being. The treatment or amelioration of symptoms can be based on objective or subjective parameters; including the results of a physical examination, neurological examination, and/or psychiatric evaluations.

In some embodiments this invention provides methods of neuroprotection. In certain embodiments, these methods comprise administering a therapeutically effective amount of the peptide combination of the invention to a patient who has not yet developed overt, clinical signs or symptoms of injury or damage to the cells of the nervous system but who may be in a high risk group for the development of neuronal damage because of injury or trauma to the nervous system or because of some known predisposition either biochemical or genetic or the finding of a verified biomarker of one or more of these disorders.

Thus, in some embodiments, the methods and compositions of the present invention are directed toward neuroprotection in a subject who is at risk of developing neuronal damage but who has not yet developed clinical evidence. This patient may simply be

at "greater risk" as determined by the recognition of any factor in a subject's, or their families, medical history, physical exam or testing that is indicative of a greater than average risk for developing neuronal damage. Therefore, this determination that a patient may be at a "greater risk" by any available means can be used to determine whether the patient should be treated with the methods of the present invention.

Accordingly, in an exemplary embodiment, subjects who may benefit from treatment by the methods and peptide or the peptide combination of this invention can be identified using accepted screening methods to determine risk factors for neuronal damage. These screening methods include, for example, conventional work-ups to determine risk factors including but not limited to: for example, head trauma, either closed or penetrating, CNS infections, bacterial or viral, cerebrovascular disease including but not limited to stroke, brain tumors, brain edema, cysticercosis, porphyria, metabolic encephalopathy, drug withdrawal including but not limited to sedative-hypnotic or alcohol withdrawal, abnormal perinatal history including anoxia at birth or birth injury of any kind, cerebral palsy, learning disabilities, hyperactivity, history of febrile convulsions as a child, history of status epilepticus, family history of epilepsy or any seizure related disorder, inflammatory disease of the brain including lupis, drug intoxication either direct or by placental transfer, including but not limited to cocaine poisoning, parental consanguinity, and treatment with medications that are toxic to the nervous system including psychotropic medications.

The determination of which patients may benefit from treatment with a neuroprotective drug in patients who have no clinical signs or symptoms may be based on a variety of "surrogate markers" or "biomarkers".

As used herein, the terms "surrogate marker" and "biomarker" are used interchangeably and refer to any anatomical, biochemical, structural, electrical, genetic or chemical indicator or marker that can be reliably correlated with the present existence or future development of neuronal damage. In some instances, brain-imaging techniques, such as computer tomography (CT), magnetic resonance imaging (MRI) or positron emission tomography (PET), can be used to determine whether a subject is at risk for neuronal damage. Suitable biomarkers for the methods of this invention include, but are not limited to: the determination by MRI, CT or other imaging techniques, of sclerosis, atrophy or volume loss in the hippocampus or overt mesial temporal sclerosis (MTS) or similar relevant anatomical pathology; the detection in the patient's blood, serum or tissues of a molecular species such as a protein or other biochemical biomarker, e.g., elevated levels of ciliary neurotrophic factor (CNTF) or elevated serum levels of a neuronal degradation product; or other

evidence from surrogate markers or biomarkers that the patient is in need of treatment with a neuroprotective drug.

It is expected that many more such biomarkers utilizing a wide variety of detection techniques will be developed in the future. It is intended that any such marker or indicator of the existence or possible future development of neuronal damage, as the latter term is used herein, may be used in the methods of this invention for determining the need for treatment with the compounds and methods of this invention.

A determination that a subject has, or may be at risk for developing, neuronal damage would also include, for example, a medical evaluation that includes a thorough history, a physical examination, and a series of relevant bloods tests. It can also include an electroencephalogram (EEG), CT, MRI or PET scan. A determination of an increased risk of developing neuronal damage or injury may also be made by means of genetic testing, including gene expression profiling or proteomic techniques. For psychiatric disorders that may be stabilized or improved by a neuroprotective drug, e.g., bipolar disorder, schizoaffective disorder, schizophrenia, impulse control disorders, etc. the above tests may also include a present state exam and a detailed history of the course of the patients symptoms such as mood disorder symptoms and psychotic symptoms over time and in relation to other treatments the patient may have received over time, e.g., a life chart. These and other specialized and routine methods allow the clinician to select patients in need of therapy using the methods and formulations of this invention. In some embodiments of the present invention peptides suitable for use in the practice of this invention will be administered either singly or concomitantly with at least one or more other compounds or therapeutic agents, e.g., with other neuroprotective drugs or antiepileptic drugs, anticonvulsant drugs. In these embodiments, the present invention provides methods to treat or prevent neuronal injury in a patient. The method includes the step of; administering to a patient in need of treatment, an effective amount of one of the peptides disclosed herein in combination with an effective amount of one or more other compounds or therapeutic agents that have the ability to provide neuroprotection or to treat or prevent seizures or epileptogenesis or the ability to augment the neuroprotective effects of the compounds of the invention.

As used herein the term "combination administration" of a compound, therapeutic agent or known drug with the peptide combination of the present invention means administration of the drug and the one or more compounds at such time that both the

known drug and the peptide combination will have a therapeutic effect. In some cases this therapeutic effect will be synergistic. Such concomitant administration can involve concurrent (i.e. at the same time), prior, or subsequent administration of the drug with respect to the administration of the peptide combination of the present invention. A person of ordinary skill in the art would have no difficulty determining the appropriate timing, sequence and dosages of administration for particular drugs and peptides of the present invention.

The said one or more other compounds or therapeutic agents may be selected from compounds that have one or more of the following properties: antioxidant activity; NMDA receptor antagonist activity, augmentation of endogenous GABA inhibition; NO synthase inhibitor activity; iron binding ability, e.g., an iron chelator; calcium binding ability, e.g., a Ca (II) chelator; zinc binding ability, e.g., a Zn (II) chelator; the ability to effectively block sodium or calcium ion channels, or to open potassium or chloride ion channels in the CNS of a patient.

The peptide or the peptide combination of the present invention were tested using the assays described in Examples 1-7, 9-17 for their effect as active therapeutic agents in the prophylaxis and/or treatment of neurodegenerative diseases and disorders.

Heart and vascular disease

Heart disease is a general term used to describe many different heart conditions. For example, coronary artery disease, which is the most common heart disease, is characterized by constriction or narrowing of the arteries supplying the heart with oxygen-rich blood, and can lead to myocardial infarction, which is the death of a portion of the heart muscle. Heart failure is a condition resulting from the inability of the heart to pump an adequate amount of blood through the body. Heart failure is not a sudden, abrupt stop of heart activity but, rather, typically develops slowly over many years, as the heart gradually loses its ability to pump blood efficiently. Risk factors for heart failure include coronary artery disease, hypertension, valvular heart disease, cardiomyopathy, disease of the heart muscle, obesity, diabetes, and/or a family history of heart failure.

Examples of cardiovascular diseases and disorders are: aneurysm, stable angina, unstable angina, angina pectoris, angioneurotic edema, aortic valve stenosis, aortic aneurysm, arrhythmia, arrhythmogenic right ventricular dysplasia, arteriosclerosis, arteriovenous malformations, atrial fibrillation, Behcet syndrome, bradycardia, cardiac tamponade, cardiomegaly, congestive cardiomyopathy, hypertrophic

cardiomyopathy, restrictive cardiomyopathy, carotid stenosis, cerebral hemorrhage, Churg-Strauss syndrome, diabetes, Ebstein's Anomaly, Eisenmenger complex, cholesterol embolism, bacterial endocarditis, fibromuscular dysplasia, congenital heart defects, heart diseases, congestive heart failure, heart valve diseases, heart attack, epidural hematoma, hematoma, subdural, Hippel-Lindau disease, hyperemia, hypertension, pulmonary hypertension, cardiac hypertrophy, left ventricular hypertrophy, right ventricular hypertrophy, hypoplastic left heart syndrome, hypotension, intermittent claudication, ischemic heart disease, Klippel-Trenaunay-Weber syndrome, lateral medullary syndrome, long QT syndrome mitral valve prolapse, moyamoya disease, mucocutaneous lymph node syndrome, myocardial infarction, myocardial ischemia, myocarditis, pericarditis, peripheral vascular diseases, phlebitis, polyarteritis nodosa, pulmonary atresia, Raynaud disease, Sneddon syndrome, superior vena cava syndrome, syndrome X, tachycardia, Takayasu's arteritis, hereditary hemorrhagic telangiectasia, telangiectasis, temporal arteritis, tetralogy of Fallot, thromboangiitis obliterans, thrombosis, thromboembolism, tricuspid atresia, varicose veins, vascular diseases, vasculitis, vasospasm, ventricular fibrillation, Williams syndrome, peripheral vascular disease, varicose veins and leg ulcers, deep vein thrombosis, Wolff-Parkinson-White syndrome.

Vascular diseases are often the result of decreased perfusion in the vascular system or physical or biochemical injury to the blood vessel.

Peripheral vascular disease (PVD) is defined as a disease of blood vessels often encountered as narrowing of the vessels of the limbs. There are two main types of these disorders, functional disease which doesn't involve defects in the blood vessels but rather arises from stimuli such as cold, stress, or smoking, and organic disease which arises from structural defects in the vasculature such as atherosclerotic lesions, local inflammation, or traumatic injury. This can lead to occlusion of the vessel, aberrant blood flow, and ultimately to tissue ischemia.

One of the more clinically significant forms of PVD is peripheral artery disease (PAD). PAD is often treated by angioplasty and implantation of a stent or by artery bypass surgery. Clinical presentation depends on the location of the occluded vessel. For example, narrowing of the artery that supplies blood to the intestine can result in severe postprandial pain in the lower abdomen resulting from the inability of the occluded vessel to meet the increased oxygen demand arising from digestive and absorptive processes. In severe forms the ischemia can lead to intestinal necrosis. Similarly, PAD in the leg can lead to intermittent pain, usually in the calf, that comes and goes with activity. This disorder is known as intermittent claudication (IC) and

can progress to persistent pain while resting, ischemic ulceration, and even amputation.

5 Peripheral vascular disease is also manifested in atherosclerotic stenosis of the renal artery, which can lead to renal ischemia and kidney dysfunction.

10 One disease in which vascular diseases and their complications are very common is diabetes mellitus. Diabetes mellitus causes a variety of physiological and anatomical irregularities, the most prominent of which is the inability of the body to utilize glucose normally, which results in hyperglycemia. Chronic diabetes can lead to complications of the vascular system which include atherosclerosis, abnormalities involving large and medium size blood vessels (macroangiopathy) and abnormalities involving small blood vessels (microangiopathy) such as arterioles and capillaries.

15 Patients with diabetes mellitus are at increased risk of developing one or more foot ulcers as a result of established long-term complications of the disease, which include impaired nerve function (neuropathy) and/or ischemia. Local tissue ischemia is a key contributing factor to diabetic foot ulceration.

20 In addition to large vessel disease, patients with diabetes suffer further threat to their skin perfusion in at least two additional ways. First, by involvement of the non-conduit arteries, which are detrimentally affected by the process of atherosclerosis, and secondly, and perhaps more importantly, by impairment of the microcirculatory control mechanisms (small vessel disease). Normally, when a body part suffers
25 some form of trauma, the body part will, as part of the body's healing mechanism, experience an increased blood flow. When small vessel disease and ischemia are both present, as in the case of many diabetics, this natural increased blood flow response is significantly reduced. This fact, together with the tendency of diabetics to form blood clots (thrombosis) in the microcirculatory system during low levels of
30 blood flow, is believed to be an important factor in ulcer pathogenesis.

Neuropathy is a general term which describes a disease process which leads to the dysfunction of the nervous system, and is one of the major complications of diabetes mellitus, with no well-established therapies for either its symptomatic treatment or for
35 prevention of progressive decline in nerve function.

The thickening and leakage of capillaries caused by diabetes primarily affect the eyes (retinopathy) and kidneys (nephropathy). The thickening and leakage of

capillaries caused by diabetes are also associated with skin disorders and disorders of the nervous system (neuropathy).

The eye diseases associated with diabetes are nonproliferative diabetic retinopathy, proliferative diabetic retinopathy, diabetic maculopathy, glaucoma, cataracts and the like.

Other diseases, although not known to be related to diabetes are similar in their physiological effects on the peripheral vascular system. Such diseases include Raynaud syndrome, CREST syndrome, autoimmune diseases such as erythematosis, rheumatoid disease, and the like.

As used herein, the term "peripheral vascular diseases" comprises any peripheral vascular disease including peripheral and autonomic neuropathies. Examples of "peripheral vascular disease" include peripheral arterial disease, such as chronic arterial occlusion including arteriosclerosis, arteriosclerosis obliterans and thromboangiitis obliterans (Buerger's disease), macroangiopathy, microangiopathy, diabetes mellitus, thrombophlebitis, phlebemphraxis, Raynaud's disease, Raynaud's syndrome, CREST syndrome, health hazard due to vibration, Sudeck's syndrome, intermittent claudication, cold sense in extremities, abnormal sensation in extremities, sensitivity to the cold, Meniere's disease, Meniere's syndrome, numbness, lack of sensation, anesthesia, resting pain, causalgia (burning pain), disturbance of peripheral circulation function, disturbance of nerve function, disturbance of motor function, motor paralysis, diabetic peripheral circulation disorder, lumbar spinal canal stenosis, diabetic neuropathy, shock, autoimmune disease such as erythematosis, rheumatoid disease and rheumatoid arthritis, autonomic neuropathy, diabetic autonomic neuropathy, autonomic imbalance, orthostatic hypotension, erectile dysfunction, female sexual dysfunction, retrograde ejaculation, cystopathy, neurogenic bladder, defective vaginal lubrication, exercise intolerance, cardiac denervation, heat intolerance, gustatory sweating, diabetic complication, hyperglycemia, hypoglycemia unawareness, hypoglycemia unresponsiveness; glaucoma, neovascular glaucoma, cataract, retinopathy, diabetic retinopathy, diabetic maculopathy, occlusion of retinal artery, obstruction of central artery of retina, occlusion of retinal vein, macular edema, aged macular degeneration, aged disciform macular degeneration, cystoid macular edema, palpebral edema, retinal edema, chorioretinopathy, neovascular maculopathy, uveitis, iritis, retinal vasculitis,

endophthalmitis, panophthalmitis, metastatic ophthalmia, choroiditis, retinal pigment epithelitis, conjunctivitis, cyclitis, scleritis, episcleritis, optic neuritis, retrobulbar optic neuritis, keratitis, blepharitis, exudative retinal detachment, corneal ulcer, conjunctival ulcer, chronic nummular keratitis, Thygeson keratitis, progressive Mooren' s ulcer, 5 damage of skin, skin ulcer including foot ulcer, diabetic ulcer, burn ulcer, lower leg ulcer, postoperative ulcer, traumatic ulcer, ulcer after herpes zoster, radiation ulcer, drug induced ulcer, frostbite (cold injury) , chilblain, gangrene and sudden gangrene, angina pectoris/ variant angitis, coronary arteriosclerosis (chronic ischemic heart disease, asymptomatic ischemic heart disease, arteriosclerotic cardiovascular 10 disease), myocardial infarction, heart failure, congestive heart failure and painless ischemic heart disease, pulmonary edema, hypertension, pulmonary hypertension; portal hypertension, diabetic nephropathy, decubitus, renal failure.

15 The peptide or the peptide combination of the present invention were tested using the assays described in Examples 1-7, 9-17 for their effect as active therapeutic agents in the prophylaxis and/or treatment of heart and vascular diseases and disorders.

20 The peptides of the present invention were tested using the assays described in Examples 1-7, 9-17 for their effect as active therapeutic agents in the prophylaxis and/or treatment of heart and vascular diseases.

Angiogenesis

25 Angiogenesis is a physiological process involving the growth of new blood vessels from pre-existing vessels. Angiogenesis is a normal process in growth and development, as well as in wound healing. However, this is also a fundamental step in the transition of tumors from a dormant state to a malignant state. Angiogenesis occurs in several well-characterized stages. First, biological signals 30 known as angiogenic growth factors activate receptors present on endothelial cells present in pre-existing blood vessels. Second, the activated endothelial cells begin to release enzymes called proteases that degrade the basement membrane in order to allow endothelial cells to escape from the original (parent) vessel walls. The endothelial cells then proliferate into the surrounding matrix and form solid 35 sprouts connecting neighboring vessels. As sprouts extend toward the source of the angiogenic stimulus, endothelial cells migrate, using adhesion molecules, called integrins. These sprouts then form loops to become a full-fledged vessel

lumen as cells migrate to the site of angiogenesis. Sprouting occurs at a rate of several millimeters per day, and enables new vessels to grow across gaps in the vasculature.

Therapeutic angiogenesis is the application of specific compounds which may inhibit or induce the creation of new blood vessels in the body in order to combat disease. The presence of blood vessels where there should be none may affect the mechanical properties of a tissue, increasing the likelihood of failure. The absence of blood vessels in a repairing or otherwise metabolically active tissue may retard repair or some other function. Several diseases are the result of failure or insufficient blood vessel formation and may be treated by a local expansion of blood vessels, thus bringing new nutrients to the site, facilitating repair. Other diseases may be created by a local expansion of blood vessels, interfering with normal physiological processes.

Angiogenesis represents an excellent therapeutic target for the treatment of, for example, cardiovascular diseases. It is a potent, physiological process that underlies the natural manner in which the human body responds to a diminution of blood supply to vital organs, namely the production of new collateral vessels to overcome the ischemic insult.

The modern clinical application of the principle "angiogenesis" can be divided into two main areas:

1. Anti-angiogenic therapies
2. Pro-angiogenic therapies.

Whereas anti-angiogenic therapies are trying to fight:

- Any type of cancer and malignancies and their metastases in numerous organs, like hemangiomas (because tumors, in general, are nutrition- and oxygen-dependent, thus being in need of adequate blood supply),
- Infectious diseases,
- Vasculitis and excessive angiogenesis in autoimmune disorders such as systemic sclerosis (Scleroderma), multiple sclerosis, Sjögren's disease,
- Vascular malformations in blood and lymph vessels like DiGeorge syndrome, hereditary haemorrhagic telangiectasia, cavernous hemangioma, cutaneous hemangioma, lymphatic malformations, transplant arteriopathy, atherosclerosis, vascular anastomoses,
- Adipose tissue in obesity,
- Chronic allograft rejections,

- Skin diseases like psoriasis, warts, allergic dermatitis, scar keloids, pyogenic granulomas, blistering disease, Kaposi sarcoma in AIDS patients, systemic sclerosis (Scleroderma),
- Eye diseases like persistent hyperplastic vitreous syndrome, diabetic retinopathy, retinopathy of prematurity, choroidal neovascularization,
- Lung diseases like any type of pulmonary hypertension, asthma, nasal polyps, rhinitis, chronic airway inflammation and obstruction (COPD), cystic fibrosis, acute lung injury, bronchiolitis obliterans organizing pneumonia,
- Gastrointestinal tract diseases like inflammatory bowel disease, periodontal disease, ascites, peritoneal adhesions, liver cirrhoses,
- Reproductive system diseases like endometriosis, uterine bleeding, ovarian cysts, ovarian hyperstimulation,
- Bone and joint diseases like arthritis and synovitis, osteomyelitis, osteophyte formation, HIV-induced bone marrow angiogenesis,
- Kidney diseases like early diabetic nephropathy

The pro-angiogenic therapies are important in the search of new treatment options for diseases characterized or caused by insufficient angiogenesis or vessel regression:

- Nervous system diseases like Alzheimer's disease, amyotrophic lateral sclerosis, diabetic neuropathy, stroke,
- Blood and lymph vessels diseases like diabetic angiopathy, impaired reendothelialization in restenosis, lymphedema,
- Gastrointestinal ulcerations, oral ulcers, mucosal ischemia in Crohn's disease
- Skin diseases like lupus,
- Reproductive system diseases like preeclampsia, menorrhagia,
- Lung diseases like neonatal respiratory distress syndrome, pulmonary fibrosis, emphysema,
- Kidney diseases like nephropathy, glomerulosclerosis, tubulointerstitial fibrosis,
- Bone diseases like osteoporosis, impaired bone fracture healing,
- Heart diseases like ischaemic heart disease, cardiac failure,
- Any type of wound healing disorders.

Angiogenesis research is also a cutting edge field in cancer research, and traditional therapies, such as radiation therapy, may work in part by targeting the genomically stable endothelial cell compartment, rather than the genomically unstable tumor cell compartment. New blood vessel formation is a relatively

fragile process, subject to disruptive interference at several levels. In short, the therapy is the selection agent which is being used to kill a cell compartment. Tumor cells evolve resistance rapidly due to rapid generation time (days) and genomic instability (variation), whereas endothelial cells are a good target because of a long generation time (months) and genomic stability (low variation). Angiogenesis-based tumour therapy relies on natural and synthetic angiogenesis inhibitors like angiostatin, endostatin and tumstatin. These are proteins that mainly originate as specific fragments pre-existing structural proteins like collagen or plasminogen.

Recently, the 1st FDA-approved therapy targeted at angiogenesis in cancer came on the market in the US. This is a monoclonal antibody directed against an isoform of VEGF, and the therapy has been approved for use in colorectal cancer in combination with established chemotherapy. Therefore there is a wide medical need for additional medicaments in the field of angiogenesis.

In addition, in terms of tissue engineering, medicaments that influence angiogenesis in vascular grafts are needed. More than 450,000 vascular grafts were used in coronary bypass surgeries annually. Other uses for vascular grafts include treatments for blood vessel aneurysms and fistulas, as well as replacements for diseased arteries in other locations in the body. When possible, the best choice for a replacement vessel is an autograft, where sections of the patient's healthy blood vessels (usually veins) are harvested and implanted in the required location. Many patients, however, especially those with pre-existing vascular disease or patients that have already had autograft procedures, do not have blood vessels that are healthy enough to adequately serve as replacements. In these cases, the most common form of treatment has been the use of synthetic polymeric materials, like ePTFE (extended polytetrafluoroethylene) and Dacron (poly[ethylene terephthalate]), to form either permanent or resorbable replacements for the damaged vessels. In cases where the graft can be of a large diameter (greater than 5-6 mm), the synthetic material has been effective. However, in situations where a smaller vessel diameter is required, the synthetic materials cannot be used due to high rates of stenosis and thrombus formation. One possible solution is to use natural materials like collagen, either modified or combined with a synthetic material, to form a graft that more closely mimics the body's natural function and has low thrombogenicity and low incidence of stenosis.

Failure of the autograft is usually due to some form of occlusion that results from luminal narrowing. Damage of the vessel during removal and reimplantation may cause the recruitment of factors or cells that adhere to the autograft wall and decrease the diameter of the lumen. The restricted flow then increases the

thrombogenicity, making full occlusion even more likely. Other problems are preparation and preservation of the autograft, procedures that can result in vessel damage or diminished *in vivo* performance. Finally, due to increased and/or different mechanical forces, endothelial cells can shrink, diminishing barrier performance, and degrade, also resulting in increased thrombogenicity. In order to reduce thrombus formation anticoagulation drugs are necessary. The use of these drugs often results in undesirable systemic side effects and can be very problematic. Therefore synthetic materials are poor choices for materials for small diameter vascular grafts.

By incorporating biological materials into a synthetic vascular graft the host response can be modulated to help insure that the graft will not fail. The use of collagen as a material for a synthetic vascular graft is quite promising because it is biodegradable and has good mechanical properties. Since collagen is biodegradable, as the device degrades tissue can grow into the device. This is advantageous because ideally as the collagen implant degrades the newly formed tissue will replace it, which results in a gradual transfer of stress from the implanted device to the newly formed tissue.

If a collagen vascular implant material was seeded with endothelial cells so that they coat the lumen, the surface would theoretically be more biocompatible. Recently, endothelial cells have been cultured onto the collagen small diameter vascular grafts. Therefore by incorporating biodegradable peptides into the collagen vascular implant material, endothelial cells can be seeded onto the top of the material to create a luminal surface that is comprised of endothelial cells to more closely mimic the natural biological environment. Migration of endothelial cells on biomaterials is very important for the development of implantable devices. These cell property controls the rates of reendothelization and angiogenesis that are important for the success of the implant.

Angiogenesis is a complex, multi-stage process by which new blood vessels are formed from pre-existing vasculature. Two critical steps in this process are endothelial cell migration and assembly into new tubules. Over the last decade, diverse arrays of molecular regulators that participate in the process of angiogenesis have been identified. The receptor tyrosine kinases, for example, are one such family of angiogenesis regulators that play a prominent role in endothelial cell assembly and migration.

The peptides of the present invention were tested using the assays described in Examples 1-7, 9-17 for their effect as active therapeutic agents in the prophylaxis and/or treatment of heart and vascular diseases and disorders and of diseases and disorders dependent on increased or decreased angiogenesis.

Rare or Orphan diseases

Another aspect of the present invention is directed to the use of the peptide compound or the peptide combination as a therapeutic agent for the prophylaxis and/or treatment of the following orphan diseases as well as for the prophylaxis and/or treatment of an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, an infectious disease, or a heart and vascular disease in patients suffering from one or more of the following Rare or Orphan Diseases:

ABCD syndrome, AAE, ABSD, ACPS III, ACRP syndrome, ACS, ACTH deficiency, isolated ACTH resistance, ADANE, ADCA, ADCME, ADEM, ADLTE, ADULT syndrome, AEC syndrome, AGM2, AHDS, AIDS wasting syndrome, ALS, ALSG, AMME syndrome, ANOTHER syndrome, AOA1, AOS, APC, Autoimmune polyendocrinopathy - candidiasis - ectodermal dystrophy syndrome, APUDoma, ARCMT, ARC syndrome, ARCA, AREDYLD syndrome, ASD, ASPED, ASPWSCR duplication, ATLD, ATR16, ATRUS syndrome, ATS-MR, AVED Aagenaes syndrome, Aarskog like syndrome, Aarskog-Ose-Pande syndrome, Aarskog-Scott syndrome, Aase syndrome, Aase-Smith syndrome, Abdominal aortic aneurysm, Aberrant left pulmonary artery, Abetalipoproteinemia, Ablepharon macrostomia syndrome, Abruzzo-Erickson syndrome, Acalvaria, Acampomelic campomelic dysplasia, Acanthamoeba keratitis, Acanthocytic disorder, Acanthocytosis, Acanthosis nigricans, Acatalasemia, Aceruloplasminemia, Achalasia, Achard Thiers syndrome, Ad Acheiropodia, Achondroplasia, Achromatopsia, Acitretin embryofetopathy, Ackerman syndrome, Acoustic neurinoma, Acquired generalized lipodystrophy, Acquired hypoprothrombinemia, Acquired ichthyosis, Acquired idiopathic sideroblastic anaemia, Acquired lipoatrophic diabetes, Acquired prothrombin deficiency, Acrodermatitis enteropathica zinc deficiency type, Acrodysostosis, Acrodysplasia, Acrofacial dysostosis, Acrokeratoderma, Acrokeratoelastoidosis, Acromelanosis, Acromesomelic dwarfism, Acromicric dysplasia, Acroosteolysis dominant type, Acrorenal defect - ectodermal dysplasia - diabetes, Acrorenal syndrome, Actinic porokeratosis disseminated superficial, Actinic porokeratosis, Acute Respiratory Distress Syndrome, Acute basophilic leukaemia, Acute erythroblastic leukaemia, Acute febrile neutrophilic dermatosis, Acute inflammatory demyelinating polyradiculoneuropathy (aidp), Acute interstitial pneumonia, Acute leukaemia of ambiguous lineage, Acute leukaemia of indeterminate lineage, Acute liver failure, Acute lymphoblastic leukaemia, Acute medullary lesions, Acute megacaryoblastic leukaemia, Acute monoblastic leukaemia, Acute motor and sensory axonal neuropathy (AMSAN), Acute motor axonal neuropathy (AMAN), Acute myeloblastic leukaemia, Acute myelodysplasia with myelofibrosis, Acute myelofibrosis, Acute myeloid leukaemia in Down syndrome,

Acute myelomonocytic leukaemia, Acute myelosclerosis, Acute non lymphoblastic leukaemia, Acute panmyelosis with myelofibrosis, Acute peripheral arterial occlusion, Acute promyelocytic leukaemia, Acute tubulointerstitial nephritis and uveitis syndrome, Adactylia unilateral, Adamantinoma, Adams nance syndrome, Adams-Oliver syndrome, Addison's disease, Adenine phosphoribosyltransferase deficiency, Adenosine deaminase deficiency, Adenosylcobalamin deficiency, Adenovirus infection in immunocompromised patients, Adenylosuccinase deficiency Adhesive arachnoiditis, Adie syndrome, Adrenal adenoma, Adrenal hyperplasia, Adrenal incidentaloma, Adrenal insufficiency, Adrenocortical carcinoma, Adrenoleukodystrophy, Adrenomyeloneuropathy, Adrenomyodystrophy, Adult Onset Still's disease, Adult T-cell leukaemia/lymphoma, Adult idiopathic neutropenia, Adult neuronal ceroid lipofuscinosis (Kufs disease, CLN4), Adult spinal muscular atrophy, Afibrinogenemia, African tick typhus, African trypanosomiasis, Agammaglobulinemia, Age-related macular degeneration, Ahn-Lerman-Sagie syndrome, Ahumada-Del Castillo syndrome, Aicardi syndrome, Aicardi-Goutieres syndrome, AIDS, Akaba hayasaka syndrome, Akesson syndrome, Alagille syndrome, Alanine-glyoxylate aminotransferase deficiency (hyperoxaluria type 1), Albers-Schonberg disease, Albright hereditary osteodystrophy, Alcock syndrome, Aldolase A deficiency, Aldosterone synthase deficiency, Aldred syndrome, Alexander disease, Algodystrophy, Alkaptonuria, Alkylglycerone phosphate synthase deficiency, Allan-Herndon-Dudley syndrome, Allergic bronchopulmonary aspergillosis, Allgrove syndrome, Alopecia, Alpers syndrome, Alpers-Huttenlocher syndrome, Alpha-thalassemia, Alport syndrome, Alström syndrome, Alternating hemiplegia, Alveolar echinococcosis, Alves dos santos castello syndrome, Alzheimer disease, Amaurosis - hypertrichosis, Ambras syndrome, Amegacaryocytosis, Amelia, Aminoaciduria, Amoebiasis due to Entamoeba histolytica, Ampola syndrome, Amyloid cardiopathy, Amyloid nephropathy, Amyloid polyneuropathy, Amyloidosis, Amylopectinosis, Amyoplasia congenita, Amyotrophic lateral sclerosis, Amyotrophy fat tissue anomaly, Anemia, Anauxetic dysplasia, Ancylostomiasis, Andermann syndrome, Andersen disease, Aneurysmal subarachnoid haemorrhage, Angelman syndrome, Angio-osteohypertrophic syndrome, Angiodysgenetic necrotizing myelopathy, Angioedema, Angiofollicular ganglionic hyperplasia, Angiokeratoma, Angioma and vascular malformation, Angiomatosis systemic cystic seip syndrome, Angioneurotic oedema, Angiostrongyliasis, Anguillulosis, Aniridia, Anisakiasis, Ankylosing spondylarthritis, Ankylostomiasis, Annuloaortic ectasia, Anodontia, Anonychia, Anophthalmia - heart and pulmonary anomalies, Anorchidia, Anorexia nervosa, Anotia, Antenatal Epstein-Barr virus infection, Anterior horn cell disease, Anti-phospholipid syndrome, Antinolo nieto borrego syndrome, Antiplasmin deficiency, Antithrombin deficiency, Antley-Bixler syndrome, Anyane-Yeboa syndrome, Aorta coarctation, Aorta hypoplasia,

Aorta-pulmonary artery fistula, Aortic aneurysm syndrome, due to TGFbeta receptors anomalies, Aortic malformation, Aortic valve atresia, Aortic valve dysplasia, Aortic valve stenosis, APECED syndrome, Apert syndrome, Aphasia, Apical ballooning syndrome, Aplasia cutis, Aplastic anaemia, Apnea of infancy (AOI), Apnea of prematurity (AOP), Apo A-I deficiency, Apolipoprotein AI amyloidosis, Apple peel syndrome, Apraxia, Arbovirus fever, Arena syndrome, Areolar atrophy of the macula, Argyria, Argyrophilic grain disease, Arhinia choanal atresia microphthalmia, Arkless-Graham syndrome, Armfield syndrome, Arndt-Gottron disease, Arnold-Chiari malformation, Aromatase deficiency, Arrhinia, Arrhythmogenic right ventricular dysplasia, Arterial calcification, Arterial duct anomalies, Arterial occlusive disease, Arterial tortuosity, Arteriohepatic dysplasia, Arthritis juvenile, Arthrogryposis, Arthroophthalmopathy, Arthropathy, Arts syndrome, Asbestosis, Ascher syndrome, Aseptic abscesses syndrome, Aseptic osteitis, Asherman's syndrome, Aspartylglucosaminidase deficiency, Asperger syndrome, Aspergillosis, Asphyxiating thoracic dystrophy of the newborn, Astley-Kendall dysplasia, Astrocytoma, Ataxia, Atelencephaly, Atelosteogenesis, Atherosclerosis, Atkin-Flaitz syndrome, Atransferrinemia, Atresia, Atrial cardiomyopathy, Atrial myxoma, Atrial septal defect, Atrichia, Atrioventricular canal complete - fallot tetralogy, Atrophia aerata, Atrophoderma vermiculata, Atypical Mole syndrome, Atypical Werner syndrome, Aughton sloan milad syndrome, Aughton-Hufnagle syndrome, Ausems wittebol post hennekam syndrome, Autism, Autoimmune haemolytic anemia, Autoimmune lymphoproliferative syndrome, Autoimmune pancreatitis, Axenfeld-Rieger syndrome, Ayazi syndrome, B-cell chronic lymphocytic leukaemia, BAFME, BBB syndrome, X-linked, BCD, BEEC, BES, BIDS syndrome, BOD syndrome, BOFS, BOR syndrome, BOS syndrome, BPD, BRESEK syndrome, BRESHECK syndrome, BRIC, BS, BSCL, BTHS, BTK-deficiency, Babesiosis, Bacterial toxic-shock syndrome, Bahemuka brown syndrome, Baird syndrome, Balantidiasis, Ballard syndrome, Baller-Gerold syndrome, Ballooning cardiomyopathy, Balo diseases, Bamforth syndrome, Bangstad syndrome, Banti syndrome, Bannayan-Riley-Ruvalcaba syndrome, Barachydactyly type A4, Baraitser burn fixen syndrome, Baraitser-Brett-Piesowicz syndrome, Barakat syndrome, Barber-Say syndrome, Bardet-Biedl syndrome, Bare lymphocyte syndrome, Barnicoat baraitser syndrome, Barraquer-Simons syndrome, Barrett esophagus, Barth syndrome, Bartonellosis, Bartsocas-Papas syndrome, Bartter syndrome, Basan syndrome, Bassen-Kornzweig disease, Bassoe syndrome, Battaglia neri syndrome, Batten disease, Baughman syndrome, Bazex syndrome, Bazex-Dupre-Christol syndrome, Bazopoulou kyrkanidou syndrome, Bd syndrome, Beals syndrome, Beals-Hecht syndrome, Bean syndrome, Beare stevenson syndrome, Bechterew syndrome, Beckwith-Wiedemann, Beemer-Ertbruggen syndrome, Behcet disease, Behr syndrome, Behrens-Baumann-Vogel syndrome,

Bell's palsy, Bellini-Chiumello-Rimoldi syndrome, Benallegue Lacete syndrome, Bejel, Bencze syndrome, Bennion-Patterson syndrome, Benson's syndrome, Beradinelli-Seip syndrome, Berdon syndrome, Berger disease, Berk tabatznik syndrome, Berlin breakage syndrome, Bernard-soulier syndrome, Berylliosis, 5 Besnier-Boeck-Schaumann disease, Bessel-Hagen disease, Best disease, Beta thalassemia, Bethlehem myopathy, Bickel-Fanconi glycogenosis, Bickers-Adams syndrome, Bickerstaff's brainstem encephalitis, Bicuspid aortic valve, Biemond syndrome, Biermer disease, Bietti's crystalline dystrophy, Bile acid synthesis defect, Bile duct cancer, Biliary atresia, Biliary inflammatory disease, Bilineal acute 10 leukaemia, Billard-Toutain-Maheut syndrome, Binder syndrome, Bindewald-Ulmer-Muller syndrome, Binswanger disease, Birt-Hogg-Dube syndrome, Bixler christian gorlin syndrome, Bjornstad syndrome, Blackfan-Diamond anaemia, Blaichman syndrome, Blake's pouch cyst, Blau syndrome, Blepharophimosis, Blepharoptosis, Blepharospasm, Blethen wenick hawkins syndrome, Bloch-Sulzberger syndrome, 15 Bloom syndrome, Blount disease, Blue Diaper syndrome, Bohring syndrome, Bohring-Opitz syndrome, Boichis syndrome, Bone disease with defective bone mineralisation, Bone disease with increased bone density, Bone marrow failure, Bonneau-Beaumont syndrome, Bonnemann-Meinecke-Reich syndrome, Bonnet-Dechaume-Blanc syndrome, Book syndrome, Boomerang dysplasia, Booth haworth 20 dilling syndrome, Borjeson-Forssman-Lehmann syndrome, Bork syndrome, Bornholm eye disease, Bosley-Salih-Alorainy syndrome, Bosma henkin christiansen syndrome, Bothnia retinal dystrophy, Boucher-Neuhauser syndrome, Bourneville syndrome, Boutonneuse fever, Bouwes Bavinck syndrome, Bowen syndrome, Boyadjiev-Jabs syndrome, Boylan dew syndrome, Brachman-de Lange syndrome, 25 Brachydactyly - arterial hypertension, Brachymesophalangy II and V, Brachyolmia, Braddock carey syndrome, Bradyopsia, Brain inflammatory disease, Brain injury, Brain sclerosis, Brauer syndrome, Braun bayer syndrome, Braun-Tinschert, Breast cancer, Brill-Zinsser disease, Brittle bone disease, Brody myopathy, Bronchial carcinoid tumour, Bronchiectasis, Bronchiolitis obliterans organizing pneumonia, 30 Bronchiolitis obliterans with obstructive pulmonary disease, Bronchogenic cyst, Bronchopulmonary dysplasia, Bronsiegel-Zelnick syndrome, Brooke-Spiegler syndrome, Brown-Vialetto-van Laere syndrome, Bruce winship syndrome, Brucellosis, Bruck syndrome, Brugada syndrome, Brunner-Winter syndrome, Brunzell syndrome, Bruyn scheltens syndrome, Buckley syndrome, Budd-Chiari 35 syndrome, Buerger's disease, Bull-Nixon syndrome, Bulldog syndrome, Bulimia, Bullous systemic lupus erythematosus, Buntinx lormans martin syndrome, Burkitt lymphoma, Burn-McKeown syndrome, Burning Mouth syndrome, Buschke-Fischer-Brauer syndrome, Buschke-Ollendorff syndrome, Buttiens-Fryns syndrome, C syndrome, CACD, CACH syndrome, CADASIL, CAMAK syndrome, CAMFAK

syndrome, CAMOS syndrome, CANOMAD syndrome, CAP syndrome, CAPOS syndrome, CAPS (cryopyrin associated periodoc syndrome), CAR syndrome, CATCH 22, CATSHL syndrome, CAVC, CCFDN, CCGE syndrome, CDA type 1, CDG syndrome, CDGIIc, CDP, CDPD, CEDNIK syndrome, CFC syndrome, CHAND
5 syndrome, CREST syndrome, CRMO, CRV, CSD, CSID, CSWSS syndrome, CVID, Cacchi-Ricci disease, Cafe au lait spots syndrome, Caffey disease, Cahmr syndrome, Calcinosis, Calderon gonzalez cantu syndrome, Calpainopathy, Camera lituania cohen syndrome, Campomelia Cumming type, Camptodactyly, Camurati engelmann disease, Canale-Smith syndrome, Canavan disease, Candidiasis,
10 Cantalamessa baldini ambrosi syndrome, Canthus, Carbohydrate metabolism disorder, Cardiogenital syndrome, Cardiomyopathy, Cardioskeletal myopathy, Carey fineman ziter syndrome, Carnevale canun mendoza syndrome, Carnevale-Hernandez-del Castillo syndrome, Carnevale-Krajewska-Fischetto syndrome, Carney complex, Carney-Stratakis syndrome, Carnosinase deficiency, Carnosinemia,
15 Caroli's disease, Carpal Tunnel syndrome, Carpenter syndrome, Carpenter-Waziri syndrome, Carrington's disease, Carrion disease, Carvajal syndrome, Casamassima-Morton-Nance syndrome, Cassia Stocco dos Santos syndrome, Castleman disease, Castro gago pombo novo syndrome, Catalase deficiency, Cataract, Catel-Manzke syndrome, Cayler syndrome, Celiac disease, Celosomia, Cenani lenz syndactylism,
20 Central neurocytoma, Cephalopolysyndactyly, Ceramidase deficiency, Cerebellar hypoplasia, Cerebral arteriovenous shunt, Cerebral hemorrhage with amyloidosis, Cerebroretinal vasculopathy, Cfc syndrome, Chagas disease, Chanarin disease, Chandler syndrome, Chang-Davidson-Carlson syndrome, Chaotic atrial tachycardia, Char douglas dungan syndrome, Char syndrome, Charge syndrome, Charlevoix
25 disease, Charlie m syndrome, Chediak-Higashi like syndrome, Cheilitis glandularis, Chemke oliver mallek syndrome, Chemodectoma, Cherry-red-spot myoclonus syndrome, Cherubism, Chiari Frommel syndrome, Chitayat haj chahine syndrome, Chitayat moore del bigio syndrome, Chitayat-Meunier-Hodgkinson syndrome, Chitty hall webb syndrome, Chitty-Hall-Baraitser syndrome, Cholera, Cholestasis,
30 Cholesteryl ester storage disease, Choline acetyltransferase (ChAT) deficiency, Chondrocalcinosis, Chondrodysplasia, Chondrodystrophy, Chordoma, Choreoacanthocytosis, Chorioretinal atrophy, Choristoma, Choroidal dystrophy, Choroidal sclerosis, Choroideremia, Christ-Siemens-Touraine syndrome, Christian syndrome, Christian-Rosenberg syndrome, Christianson syndrome, Christianson-
35 Fourie syndrome, Christmas tree syndrome, Chromomycosis, Chronic eosinophilic pneumonia, Chronic fatigue syndrome, Chronic inflammatory demyelinating polyneuropathy, Chronic myeloproliferative disease, Chronic neutrophilic leukaemia, Chronic pain requiring intraspinal analgesia, Chronic pneumonitis of infancy, Chronic osteomyelitis, Chronic spinal muscular atrophy, Chudley rozdilsky syndrome,

Chudley-Lowry-Hoar syndrome, Churg-Strauss syndrome, Chylomicron retention disease, Ciliary dysentery, Ciliary dyskinesia-bronchiectasis, Cilliers-Beighton syndrome, Cirrhosis associated cardiac dysfunction, Cirrhotic cardiomyopathy, Clarkson disease, Classical Hodgkin disease, Classical homocystinuria, Claude-Bernard-Horner syndrome, Clayton-Smith-Donnai syndrome, Cleido rhizomelic syndrome, Cleidocranial dysostosis, Cleidocranial dysplasia, Clouston syndrome, Coagulation disorder, Coarctation of aorta, Coats disease, Cobb syndrome, Cocaine poisoning, Cockayne syndrome, Codas syndrome, Coeliac disease, Coenzyme Q cytochrome c reductase deficiency, Coffin syndrome, Coffin-Lowry syndrome, Coffin-Siris syndrome, Cogan syndrome, Cogan-reese syndrome, Cohen hayden syndrome, Cohen lockood wyborne syndrome, Cohen syndrome, Cole carpenter syndrome, Colitis, Collagen anomaly, Collins pope syndrome, Collins sakati syndrome, Coloboma, Colon cancer, Colonic atresia, Colorado tick encephalitis, Combined pituitary hormone deficiencies, Complement component deficiency, Congenital Lambert-Eaton-like syndrome, Congenital leptin deficiency, Congenital lobar emphysema, Conjunctival disease, Conjunctival vascular anomaly, Conn syndrome, Connective tissue disease, Conradi-Hünemann-Happle syndrome, Constrictive bronchiolitis, Cooks syndrome, Cooley anaemia, Cooper-Jabs syndrome, Cormier rustin munnich syndrome, Corneal dystrophy, Cornelia de Lange syndrome, Corneodermatoosseous syndrome, Corneogoniodysgenesis, Coronarocardiac fistula, Coronary arterial malformations, Coronary artery aneurysm, Coronary sinus type ASD, Cortada koussef matsumoto syndrome, Costeff optic atrophy syndrome, Costeff syndrome, Costello syndrome, Cote katsantoni syndrome, Cousin-Walbraum-Cegarra syndrome, Cowchock syndrome, Cowchock-Wapner-Kurtz syndrome, Cowden syndrome, Coxoauricular syndrome, Cramer-Niederellmann syndrome, Crandall syndrome, Crane heise syndrome, Cranial malformation, Craniopharyngioma, Craniorachischisis, Craniostenosis, Craniosynostosis, Craniotelencephalic dysplasia, Craniotubular syndrome, Creatine deficiency, Creeping disease, Creutzfeldt-Jakob disease, Cri du chat syndrome, Crigler-Najjar syndrome, Crimean-Congo haemorrhagic fever (CCHF), Crisponi syndrome, Criss-cross heart, Criswick-Schepens syndrome, Crohn disease, Crome syndrome, Cronkhite canada syndrome, Cross syndrome, Crouzon disease, Crow-Fukase syndrome, Cryoglobulinaemia mixed, Cryptococcosis, Cryptogenic organizing pneumonia, Cryptophthalmia, Cryptosporidiosis, Culler-Jones syndrome, Currarino triad, Curry-Hall syndrome, Curry-Jones syndrome, Cushing disease, Cutaneomeningospinal angiomas, Cutaneous lupus erythematosus, Cutaneous mastocytoma, Cutaneous mastocytosis, Cutaneous photosensitivity colitis, Cutaneous vasculitis, Cutaneous myiasis, Cutis laxa, Cutler bass romshe syndrome, Cyclosporosis, Cystathioninuria, Cystic fibrosis, Cystic hamartoma of lung

and kidney, Cystic lymphangioma, Cystic renal disease, Cystinosis, Cystinuria, Cytochrome c oxydase deficiency, Cytomegalovirus (CMV) disease in patients with impaired cell mediated immunity deemed at risk, Cytopenia, Czeizel brooser syndrome, Czeizel losonci syndrome, D ercole syndrome, D-2-
5 hydroxyglutaricaciduria, D-glycerate dehydrogenase deficiency (hyperoxaluria type 2), D-glycerate kinase deficiency, D-glycericacidemia, DCMA syndrome, DCMD, DEND syndrome, DI-CMT, DIDMOAD syndrome (Diabetes Insipidus - Diabetes Mellitus - Optic Atrophy - Deafness), DIS, DK phocomelia syndrome, DKC, DOOR syndrome, DORV, DTDP1, DYT6, Da silva syndrome, Dacryocystitis osteopoikilosis,
10 Daentl-Townsend-Siegel syndrome, Dahlberg-Borer-Newcomer syndrome, Daish hardman lamont syndrome, Dancing Eye syndrome, Dandy walker malformation, Daneman davy mancer syndrome, Danon disease, Darier disease, Darier-Gottron disease, Davenport donlan syndrome, David syndrome, Davies disease, Davis lafer syndrome, De Barsey syndrome, De Hauwere-Leroy-Adriaenssens syndrome, De
15 Santis-Cacchione syndrome, De Smet-Fabry-Fryns syndrome, De Vaal disease, De la Chapelle dysplasia, De morsier syndrome, Deafness - small bowel diverticulosis - neuropathy, Deal barratt dillon syndrome, Degos disease, Dejerine-Sottas syndrome, Dekaban-Arima syndrome, Delayed graft function after organ transplantation, Delleman-Oorthuys syndrome, Dementia associated with a metabolic disease,
20 Dementia associated with a neurodegenerative disease, Dementia associated with an infectious disease, Dementia associated with hepatic and renal failure, Demodicidosis, Dendritic cell sarcoma, Dendritic cell tumor, Dengue, Dennis cohen syndrome, Dennis fairhurst moore syndrome, Dense (delta) granule disease, Dent disease, Dentin dysplasia, Denys-Drash syndrome, Der Kaloustian-Jarudi-Khoury
25 syndrome, Der kaloustian mcintosh silver syndrome, Dercum's disease, Dermatofibrosarcoma protuberans, Dermatologic allergic disease, Dermatostomatitis Stevens Johnson type, Desbuquois syndrome, Desminopathy, Desmoid disease, Desmosterolosis, Devic's disease, Devriendt legius fryns syndrome, Devriendt vandenberghes fryns syndrome, DiGeorge syndrome, Diabetes, Dialysis-related
30 arthropathy, Diaphanospondylodysostosis, Diaphragmatic agenesis, Diaphragmatic spinal muscular atrophy, Diffuse alveolar haemorrhage, Diffuse large B cell lymphoma, Diffuse leiomyomatosis - Alport syndrome X-linked, Diffuse neonatal haemangiomas, Dihydropyrimidinuria, Dilated cardiomyopathy with ataxia, Dincsoy-Salih-Patel syndrome, Dinno shearer weisskopf syndrome, Diomedes bernardi
35 placidi syndrome, Dionisi-Vici-Sabetta-Gambarara syndrome, Diphtheria, Diprosopia, Discoid lupus erythematosus, Discrete fibromuscular subaortic stenosis, Distichiasis - congenital heart defects - peripheral vascular anomalies, Distomatosis, Dobrow syndrome, Donath-Landsteiner syndrome, Donnai-Barrow syndrome, Donohue syndrome, Doose syndrome, Dorfman-cheranin disease, Dowling-Degos disease,

Dowling-Degos-Kitamura disease, Down syndrome, Doyne honeycomb retinal dystrophy (DHRD), Drachtman weinblatt sitarz syndrome, Drash syndrome, Dravet syndrome, Drummond syndrome, Du Pan syndrome, Duane syndrome, Dubin-Johnson syndrome, Dubowitz syndrome, Duhring brocq disease, Duker-Weiss-Siber syndrome, Dunnigan syndrome, Dupont sellier chochillon syndrome, Dyggve-Melchior-Clausen disease, Dykes-Markes-Harper syndrome, Dyschondrosteosis, Dyschromatosis universalis, Dysferlinopathy, Dysfibrinogenemia, Dyskeratosis, Dysmorphic syndrome with connective tissue involvement, Dysosteosclerosis, Dysostosis, Dysphagia lusoria, Dysplasia, Dysprothrombinemia, Dyssegmental dysplasia glaucoma, Dysspondyloenchondromatosis, Dystoni-like syndrome with paroxysmal disease, Dystonia, EBD, EBJ, EBS, ECP syndrome, EDS III, EEC syndrome, EEM syndrome, EGE, ENT, ERA, ESS1, Eagle-Barret syndrome, Eales disease, Ebola virus disease, Echinocytic disorder, Ectodermal dysplasia, Ectromelia, Ectropion, Eczema-thrombocytopenia-immunodeficiency syndrome, Edinburgh malformation syndrome, Edward syndrome, Edwards-Patton-Dilly syndrome, Ehlers-Danlos syndrome, Ehrlichiosis, Eiken syndrome, Eisenmenger syndrome, Elastosis perforans serpiginosa, Elejalde syndrome, Elliott ludman teebi syndrome, Elliptocytosis, Ellis Van Creveld syndrome, Ellis yale winter syndrome, Elsching syndrome, Emanuel syndrome, Emery-Dreifuss muscular dystrophy, Emery-Nelson syndrome, Empty Sella syndrome, Encephalitis, Encephalomyelitis, Encephalopathy, Enchondromatosis, Endometriosis, Endotheliitis, Eng strom syndrome, Engel congenital myasthenia, Engelhard yatziv syndrome, Enolase deficiency, Enteric anendocrinosis, Enteropathy, Enterovirus antenatal infection, Entropion, Envenomization, Eosinophilic endocarditis, Eosinophilic pneumonia, Ependymoma, Epidermolysis bullosa, Epilepsy, Epiphyseal dysplasia, Episodic ataxia, Epispadias, Epithelial ovarian cancer, Epithelioma, Epstein-Barr virus infection, Erdheim disease, Erdheim-Chester disease, Eronen-Somer-Gustafsson syndrome, Erythema, Erythermalgia, Erythroblastopenia, Erythrocytosis, Erythroderma, Erythrokeratoderma, Erythromelalgia, Escher hirt syndrome, Escobar syndrome, Esophageal adenocarcinoma, Esophageal atresia, Essential cryoglobulinaemia, Essential iris atrophy, Essential osteolysis, Esthesioneuroblastoma, Estrogen receptor deficiency, Estrogen resistance syndrome, Evans syndrome, Ewing sarcoma, Exner syndrome, Exostoses, Exsudative retinopathy, Extracutaneous mastocytoma, Extrinsic allergic alveolitis, Eye disease, F syndrome, FAP, FAS deficiency, FCS syndrome, FCU, FENIB, FEOM, FFDD type I, FG syndrome, FLOTCH syndrome, FOP, FOSMN syndrome, FPS/AML syndrome, FRAXA syndrome, FRAXE syndrome, FRAXF syndrome, FSH resistance, Fabry disease, Factor VII deficiency, Factor VIII deficiency, Factor X deficiency, Factor XI deficiency, Factor XII deficiency, Factor XIII deficiency, Factors

II,VII,IX and X, combined deficiency, Fahr syndrome, Fallot complex, Familial LCAT deficiency, Fanconi anaemia, Fanconi ichthyosis dysmorphism, Fanconi syndrome, Fanconi-Bickel disease, Fara-Chlupackova syndrome, Farber lipogranulomatosis, Farmer's lung disease, Fatal infantile COX deficiency, Faulk-Epstein-Jones syndrome, Favism, Fazio-Londe disease, Fechtner syndrome, Feigenbaum-Bergeron-Richardson syndrome, Feingold syndrome, Felty syndrome, Fenton wilkinson toselano syndrome, Ferlini-Ragno-Calzolari syndrome, Fernhoff-Blackston-Oakley syndrome, Fetal cytomegalovirus syndrome, Fetal edema, Fetal left ventricular aneurysm, Fibrinogen disorder, Fibrochondrogenesis, Fibrodysplasia ossificans progressiva, Fibromatosis, Fibromuscular dysplasia of arteries, Fibromyalgia, Fibronectin glomerulopathy, Fibrosarcoma, Fibrosing mediastinitis, Fibrosis of extraocular muscles, Fiessinger-Leroy-Reiter's syndrome, Figuera syndrome, Filamin anomaly, Filariasis, Filippi syndrome, Fine-Lubinsky syndrome, Finlay-Markes syndrome, Finucane kurtz scott syndrome, Fitz Hugh Curtis syndrome, Fitzsimmons-Guilbert syndrome, Fitzsimmons-McLachlan-Gilbert syndrome, Fitzsimmons-Walson-Mellor syndrome, Fixed subaortic stenosis, Flegel disease, Floating-Harbor syndrome, Florid cemento-osseous dysplasia, Flynn aird syndrome, Foix chavany marie syndrome, Foix-Alajouanine syndrome, Follicular atrophoderma-basal cell carcinoma, Follicular dendritic cell sarcoma, Follicular dyskeratoma, Follicular ichthyosis, Follicular lymphoma, Fontaine-Farriaux-Blanckaert syndrome, Forbes disease, Forney-Robinson-Pascoe syndrome, Forunculoid myiasis, Fountain syndrome, Fowler-Christmas-Chapple syndrome, Fox Fordyce disease, Fra-X syndrome, Fragile X syndrome, Fragoso cid garcia hernandez syndrome, Franceschetti-Klein syndrome, Francois dyscephalic syndrome, Francois syndrome, Franek bocker kahlen syndrome, Frank-Ter Haar syndrome, Franklin disease, Fraser like syndrome, Fraser syndrome, Frasier syndrome, Freeman-Sheldon syndrome, Freiberg's disease, Freire maia pinheiro opitz syndrome, Frey's syndrome, Frias syndrome, Fried syndrome, Fried-Goldberg-Mundel syndrome, Friedman goodman syndrome, Friedreich ataxia, Froelich's syndrome, Froster-Huch syndrome, Froster-Iskenius-Watson syndrome, Fructosuria, Frydman-Cohen-Karmon syndrome, Fryns macrocephaly, Fryns-Aftimos syndrome, Fryns-Hofkens-Fabry syndrome, Fuhrmann-Rieger-de Sousa syndrome, Fukuda miyanomae nakata syndrome, Fukuhara syndrome, Fuqua-Berkovitz syndrome, Furlong syndrome, Furukawa takagi nakao syndrome, G syndrome, G6PD deficiency, GABA metabolism disease, GAMT deficiency, GAPO syndrome, GIST, GM1 gangliosidosis, GOSHS, GRACILE syndrome, GRF Tumour, GSD, GTN, GVH, Gaisbock syndrome, Galactokinase deficiency, Galactosemia, Galactosialidosis, Galloway syndrome, Galloway-Mowat syndrome, Gamborg nielsen syndrome, Game-Friedman-Paradise syndrome, Gamstorp episodic adynamy, Ganglioglioma, Garcia torres guarner syndrome,

Garcia-Lurie syndrome, Gardner silengo wachtel syndrome, Gardner-Morrison-Abbott syndrome, Garret tripp syndrome, Gastric cancer, Gastroschisis, Gaucher disease, Gaucher-like disease, Geen sandford davison syndrome, Gelineau disease, Gemignani syndrome, Gemss syndrome, Genes syndrome, Genochondromatosis, 5 Gerbode defect, Gerhardt syndrome, German syndrome, Gershonibaruch-Leibo syndrome, Gerstmann-Straussler-Scheinker syndrome, Ghosal syndrome, Gianotti Crosti syndrome, Giant cell arteritis, Giant platelet syndrome, Gilbert syndrome, Gilles de la Tourette syndrome, Gillespie syndrome, Gitelman syndrome, Glanzmann thrombasthenia, Glass bone disease, Glass-Chapman-Hockley syndrome, 10 Glaucoma, Glioblastoma, Glomerular disease, Glomerulonephritis, Glomerulopathy with fibronectin deposits (GFND), Gloomy syndrome, Glucagonoma, Glucocorticoid resistance, Glycogen storage disease, Gms syndrome, Goiter-deafness syndrome, Golabi-Rosen syndrome, Goldberg syndrome, Goldberg-Maxwell syndrome, Goldberg-Shprintzen megacolon syndrome, Goldblatt viljoen syndrome, Goldblatt 15 wallis syndrome, Goldenhar syndrome, Goldmann-Favre syndrome, Goldstein hutt syndrome, Goldston syndrome, Gollop syndrome, Gollop wolfgang complex, Goltz syndrome, Goltz-Gorlin syndrome, Gombo syndrome, Gonzales del angel syndrome, Goodman syndrome, Goodpasture syndrome, Goossens-Devriendt syndrome, Gordon syndrom, Gorham syndrome, Gorham-Stout disease, Gorlin syndrome, 20 Gorlin-Chaudhry-Moss syndrome, Graft rejection after lung transplantation, Graft versus host disease, Graham boyle troxell syndrome, Graham-Cox syndrome, Grand-Kaine-Fulling syndrome, Grange occlusive arterial syndrome, Grant syndrome, Granulocytic sarcoma, Granulomatous allergic angiitis, Granulomatous inflammatory arthritis, dermatitis, and uveitis, Granulomatous mastitis, Graves' 25 disease, Gray platelet syndrome, Greenberg dysplasia, Greig syndrome, Greither's disease, Griscelli disease, Grix blankenship peterson syndrome, Groll hirschowitz syndrome, Gronblad-Strandberg-Touraine syndrome, Grosse syndrome, Grover's disease, Growth hormone deficiency, Grubben de cock borghgraef syndrome, Gräsbeck-Imerslund disease, Guam disease, Guanidinoacetate methyltransferase 30 deficiency, Guibaud-Vainsel syndrome, Guillain-Barré syndrome, Guizar-Vasquez-Luengas syndrome, GuizarVazquez-Sanchez-Manzano syndrome, Gunal seber basaran syndrome, Gurrieri-Sammito-Bellussi syndrome, Gusher syndrome, Gynandroblastoma, Günther disease, HAD deficiency, HAE, HAIRAN syndrome, HANAC syndrome, HARD syndrome (Hydrocephalus - agyria - retinal dysplasia), 35 HCDD, HCL, HDL metabolism disorder, HEM, HEP, HERNS syndrome, HHE syndrome, HHT, HHV-8, HID syndrome, HIGM1, HIT, HMSN 5, HMSNP, HNPCC, HNSCC, HPA-1 deficiency, HPE, HSAN 1, HSD deficiency, HSV encephalitis, HSV keratitis, HUS, HVR, Haas-Robinson syndrome, Haddad syndrome, Haematologic cancers, Haemochromatosis, Haemoglobin disorders, Haemolysis, Haemolytic

anaemia, Haemolytic uremic syndrome, Haemorrhagic fever, Haemorrhagiparous thrombocytic dystrophy, Hageman factor deficiency, Hagemoser weinstein bresnick syndrome, Hailey-Hailey disease, Haim-Munk syndrome, Hairy cell leukaemia, Hajdu-Cheney syndrome, Hal-Berg-Rudolph syndrome, Halal syndrome, Halal-Setton-Wang syndrome, Hallermam streiff like syndrome, Hallermann-Streiff-Francois syndrome, Hallervorden-Spatz disease, Hamanishi ueba tsuji syndrome, Hamano tsukamoto syndrome, Hamman-Rich syndrome, Hanhart syndrome, Hand Foot Mouth syndrome, Hand-Shuller-Christian disease, Hanot syndrome, Hantavirus pulmonary syndrome, Hapnes boman skeie syndrome, Happy puppet syndrome, Harboyan syndrome, Hardcastle syndrome, Harding ataxia, Harrod syndrome, Harrod-Keele syndrome, Hartnup disorder, Hartsfield bixler demyer syndrome, Hashimoto struma, Hashimoto-Pritzker syndrome, Haspeslagh-Fryns-Muelenaere syndrome, Hawkinsinuria, Hay wells syndrome, Heart block progressive, Heart-hand syndrome, Heavy chain deposition disease, Hec syndrome, Hecht scott syndrome, Heckenlively syndrome, Heide syndrome, Heimler syndrome, Heiner syndrome (cow's milk hypersensitivity), Helmerhorst heaton crossen syndrome, Hemangioma-thrombocytopenia syndrome, Hemangiopericytoma, Hematopoietic hypoplasia, Hemeralopia, Hemi 3 syndrome, Hemiconvulsion-Hemiplegia-Epilepsy syndrome, Hemifacial hyperplasia strabismus, Hemihypertrophy intestinal web corneal opacity, Hemimelia, Hemitruncus, Hemochromatosis, Hemoglobin C disease, Hemoglobin E disease, Hemoglobin H disease, Hemolytic anaemia, Hemophilia, Hemorrhagiparous thrombocytic dystrophy, Hennekam koss de geest syndrome, Hennekam syndrome, Hennekam-Beemer syndrome, Henoch-Schoenlein purpura, Hepatic cystic hamartoma, Hepatic fibrosis, Hepatic cancer, Hepatic venoocclusive disease, Hepatitis B re-infection following liver transplantation, Hepatitis, Hepatoblastoma, Hepatocellular adenoma, Hepatocellular carcinoma, Hepatoerythropoeitic porphyria, Hepatoportal sclerosis, Hereditary coproporphyria, Hereditary endotheliopathy - retinopathy - nephropathy - stroke, Hereditary lymphoedema type I, Hereditary motor and sensory neuropathy, Hereditary vascular retinopathie - Raynaud phenomenon - migraine, Hermansky-Pudlak syndrome, Hernandez fragoso syndrome, Hernandez-Aguirre Negrete syndrome, Herpes virus infection, Herrmann opitz arthrogryposis syndrome, Hers disease, Hersh-Podruch-Weisskopf syndrome, Herva disease, Heterotaxia, Heterozygous OSMED, Hillig syndrome, Hinman syndrome, Hinson-Pepys disease, Hipo syndrome, Hirayama disease, Hirschsprung disease, Hirsutism, His bundle tachycardia, Histidine metabolism disorder, Histidinuria renal tubular defect, Histiocytic and dendritic cell tumour, Histiocytic sarcoma, Histiocytoid cardiomyopathy, Histiocytosis X, Histoplasmosis, Hittner hirsch kreh syndrome, Hmc syndrome, Hodgkin lymphoma, Hoepffner dreyer reimers syndrome, Hoffman's syndrome, Holmes benacerraf syndrome, Holmes collins syndrome, Holmes-Gang

syndrome, Holoacardius amorphus, Holoprosencephaly, Holt-Oram syndrome,
 Holzgreve wagner rehder syndrome, Homocarnosinosis, Homocystinuria,
 Homogentisic acid oxydase deficiency, Hoon hall syndrome, Horner syndrome,
 Horton disease, Houlston iron-ton temple syndrome, House allergic alveolitis, Howard
 5 young syndrome, Howell-Evans syndrome, Hoyeraal-Hreidarsson syndrome,
 Humeroradial synostosis, Humeroradioulnar synostosis, Humerospinal dysostosis,
 Hunter carpenter mc donald syndrome, Hunter jurenka thompson syndrome, Hunter
 syndrome, Hunter-Rudd-Hoffmann syndrome, Hunter-Thompson-Reed syndrome,
 Huntington disease, Huriez syndrome, Hurler syndrome, Hurler-Scheie syndrome,
 10 Hutchinson-Gilford syndrome, Hutteroth spranger syndrome, Hyaline membrane
 disease, Hyaluronidase deficiency, Hydatidosis, Hyde-Forster-Mccarthy-Berry
 syndrome, Hygroma cysticum, Hyperaldosteronism, Hyperargininemia,
 Hyperbilirubinemia, Hypercalciuria idiopathic, Hypercholesterolemia,
 Hyperchylomicronemia, Hypercortisolism, Hyperexplexia, Hyperglycinemia,
 15 Hyperimidodipeptiduria, Hyperinsulinism, Hyperkeratosis, Hyperlipidaemia,
 Hyperlipoproteinemia, Hyperlysinemia, Hypermethioninemia, Hyperornithinemia,
 Hyperostosis, Hyperoxaluria, Hyperparathyroidism, Hyperphalangism dysmorphism
 bronchomalacia, Hyperphenylalaninemic embryopathy, Hyperpipecolatemia,
 Hypersensitivity pneumonitis, Hypertelorism, Hyperthermia, Hyperthyroidism,
 20 Hypertrichosis, Hypertrophic neuropathy, Hypertrophic or verrucous lupus
 erythematosus, Hypertrophic subaortic stenosis, Hypobetalipoproteinemia,
 Hypobetalipoproteinemia, Hypochondroplasia, Hypocomplementaemic
 leucocytoclastic vasculitis, Hypodontia, Hypofibrinogenemia, Hypokalemic alkalosis,
 Hypokeratosis, Hypomyelination, Hypoparathyroidism, Hypopituitarism, Hypoplastic
 25 left heart syndrome, Hypoplastic right heart syndrome, Hypospadias, Hypothalamic
 hamartoblastoma syndrome, Hypothyroidism, Hypotrichosis, Hypoxanthine guanine
 phosphoribosyltransferase (HPRT) complete deficiency, I-cell disease, IBIDS
 syndrome, ICCA syndrome, ICE syndrome, ICF syndrome, ICOS deficiency, IDI, IED,
 IFAP syndrome, IGDA, IGF-1 deficiency, IGHD, IMAGE syndrome, INAD, INCL,
 30 IOMID syndrome, IOSCA, IPEX, IPSID, IRAK4 deficiency, ISOD, ITP, IVC stenosis,
 Ichthyosis, Idaho syndrome, Idiopathic dystonia DYT1, Idiopathic granulomatous
 mastitis, Idiopathic hypereosinophilic syndrome, Idiopathic infantile arterial
 calcification, Idiopathic infection caused by BCG or atypical mycobacteria, Idiopathic
 interstitial pneumonia, Idiopathic juvenile osteoporosis, Idiopathic myelofibrosis,
 35 Idiopathic obliterative arteriopathy, Idiopathic orthostatic hypotension, Idiopathic
 pulmonary fibrosis, Idiopathic thrombocytopenic purpura, Ieshima-Koeda-Inagaki
 syndrome, Illum syndrome, Ilyina amoashy grygory syndrome, Imaizumi kuroki
 syndrome, Immune thrombocytopaenia, Immunodeficiency, Immunoproliferative
 small intestinal disease, Infant respiratory distress syndrome, Insulin-resistance

syndrome, Insulinoma, Interdigitating dendritic cell sarcoma, Intermediate DEND
 syndrome, Intermediate spinal muscular atrophy, Internal carotid agenesis,
 Interstitial cystitis, Interstitial granulomatous dermatitis with arthritis, Interstitial
 pneumonia, Interventricular septum aneurysm, Intestinal atresia multiple, Intestinal
 5 epithelial dysplasia, Intestinal hypomagnesemia with secondary hypocalcemia,
 Intestinal lipodystrophy, Intestinal lipophagic granulomatosis, Intestinal
 lymphangiectasia, Intestinal pseudoobstruction, Intracerebral haemorrhage,
 Intracranial aneurysms, Intracranial arteriovenous malformation, Inverse Marcus-
 Gunn phenomenon, Iridocorneal endothelial syndrome, Iridogoniodysgenesis, Irons-
 10 Bhan syndrome, Irritable bowel syndrome, Isaac's syndrome, Isaacs mertens
 syndrome, Ischaemic brain injury, Ischemia/perfusion injury associated with solid
 organ transplantation procedure, Ischio-vertebral dysplasia, Iso-Kikuchi syndrome,
 Isosporiasis, Isotretinoin syndrome, Isotretinoin-like syndrome, Isovaleric acidemia,
 Itin syndrome, Ito hypomelanosis, Ivemark syndrome, JAE, JWS, Jackson-Barr
 15 syndrome, Jackson-Weiss syndrome, Jacobs syndrome, Jacobsen syndrome, Jaffe
 campanacci syndrome, Jaffe-Lichtenstein disease, Jagell holmgren hofer syndrome,
 Jalili syndrome, Jancar syndrome, Japanese encephalitis, Jarcho-Levin syndrome,
 Jaw-Winking syndrome, Jensen syndrome, Jequier-Kozlowski syndrome, Jervell and
 Lange-Nielsen syndrome, Jeune syndrome, Job syndrome, Johanson-Blizzard
 20 syndrome, Johnson syndrome, Johnson-McMillin syndrome, Johnson-Munson
 syndrome, Johnston-Aarons-Schellely syndrome, Jones syndrome, Jorgenson lenz
 syndrome, Joubert syndrome, Joubert-Boltshauser syndrome, Juberg hayward
 syndrome, Juberg-Marsidi syndrome, Judge misch wright syndrome, Jumping
 Frenchman of Maine, Jung wolff back stahl syndrome, Juvenile chronic
 25 myelomonocytic leukaemia, Juvenile gastrointestinal polyposis, Juvenile glaucoma,
 Juvenile hemochromatosis, Juvenile hyaline fibromatosis, Juvenile idiopathic arthritis,
 Juvenile macular degeneration, Juvenile myelomonocytic leukaemia, Juvenile
 polyposis syndrome (JPS), Juvenile temporal arteritis, KBG syndrome, KBG-like
 syndrome, KID syndrome, Kabuki syndrome, Kaeser syndrome, Kahler's disease,
 30 Kaler garrity stern syndrome, Kallin syndrome, Kallmann syndrome, Kalyanaraman
 syndrome, Kanzaki disease, Kaplan-Plauchu-Fitch syndrome, Kaplowitz-Bodurtha
 syndrome, Kaposi's sarcoma, Kaposiform hemangioendothelioma, Kapur-Toriello
 syndrome, Karandikar-Maria-Kamble syndrome, Karsch neugebauer syndrome,
 Kartagener syndrome, Kasabach-Merritt syndrome, Kashani-Strom-Utley syndrome,
 35 Kasznica carlson coppedge syndrome, Katsantoni papadakou lagoyanni syndrome,
 Kaufman-Mckusick syndrome, Kawasaki disease, Kawashima syndrome,
 Kawashima-Tsuji syndrome, Kearns-Sayre syndrome, Kelley-Seegmiller syndrome,
 Kelly-Kirson-Wyatt syndrome, Kennedy disease, Kennedy-Teebi syndrome,
 Kennerknecht syndrome, Kenny syndrome, Kenny-Caffey syndrome, Kenya tick-bite

fever, Keratinisation disorder associated with genetic eye disease, Keratitis, Keratoacanthoma, Keratoconus, Keratoderma, Keratosis, Kerion celsi, Kersey syndrome, Ketoacidosis, Ketoaciduria, Ketolysis disorder, Keutel syndrome, KGB syndrome, Khalifa-Graham syndrome, Kienbock disease, Kikuchi disease, Kikuchi-Fujimoto disease, Kimura disease, King-Denborough syndrome, Kinsbourne syndrome, Klatskin tumour, Klein-Waardenburg syndrome, Kleine-Levin syndrome, Kleiner holmes syndrome, Klinefelter syndrome, Klippel-Feil malformation, Klippel-Trenaunay syndrome, Kluver-Bucy syndrome, Kniest dysplasia, Knobloch layer syndrome, Kocher-Debre-Semelaigne syndrome, Kohler's disease, Kohlschutter-Tonz syndrome, Kok disease, Komar syndrome, Konigsmark knox hussels syndrome, Kopysc barczyk kroi syndrome, Kosenow syndrome, Kostmann syndrome, Kosztolanyi syndrome, Koussef nichols syndrome, Kousseff syndrome, Kowarski syndrome, Kozlowski brown hardwick syndrome, Kozlowski massen syndrome, Kozlowski ouvrier syndrome, Kozlowski tsuruta syndrome, Kozlowski-Krajewska syndrome, Krabbe disease, Krasnow-Qazi syndrome, Krauss herman holmes syndrome, Kudo tamura fuse syndrome, Kugelberg-Welander disease, Kumar-Levick syndrome, Kunze riehms syndrome, Kurczynski-Casperson syndrome, Kuskokwim disease, Kuzniecky syndrome, Kynureninase deficiency, Kyphomelic dysplasia, Kyphosis brachyphalangy optic atrophy, Küssmaul-Maier disease, L1 syndrome, L-2-hydroxyglutaricaciduria, LADD syndrome, LBSL, LBWC syndrome, LCAD, LCAT deficiency, LCCS, LCDD, LCH, LCHAD deficiency, LDD, LEOPARD syndrome, LGMD, LHCDD, LIG4 syndrome, LMS, LORD, LPI, Laband syndrome, Lachiewicz sibley syndrome, Lactate dehydrogenase deficiency, Lactic acidosis, Lactotroph adenoma, Ladda zonana ramer syndrome, Lafora disease, Laing distal myopathy, Lambdoid synostosis, Lambert syndrome, Lambert-Eaton myasthenic syndrome, Lamellar ichthyosis, Laminopathy, Landau-Kleffner syndrome (LKS), Landing disease, Landouzy-Dejerine myopathy, Langer-Giedion syndrome, Langerhans cell granulomatosis, Langerhans cell histiocytosis, Langerhans cell sarcoma, Laparoschisis, Laplane fontaine lagardere syndrome, Laron syndrome, Larsen syndrome, Larsen-like syndrome, Laryngeal abductor paralysis, Laryngo onycho cutaneous syndrome, Laryngo-tracheo-esophageal cleft pulmonary hypoplasia, Lassa fever, Lassueur-Graham-Little syndrome, Late infantile neuronal ceroid lipofuscinosis, Late onset sepsis in premature infants, Lathosterolosis, Laubry-pezzi syndrome, Launois-Bensaude adenolipomatosis, Laurence-Moon syndrome, Laurin-Sandrow syndrome, Lawrence syndrome, Lawrence-Seip syndrome, Laxova-Opitz syndrome, Le Merrer syndrome, Le marec bracq picaud syndrome, Leao-da Silva syndrome, Learman syndrome, Leber 'plus' disease, Leber congenital amaurosis, Leber miliary aneurysm, Left renal vein entrapment syndrome, Left ventricular hypertrabeculation, Left ventricular noncompaction, Legg-Calve-Perthes

disease, Legionellosis, Legionnaires' disease, Leichtman-Wood-Rohn syndrome, Leifer lai buyse syndrome, Leigh disease, Leiner disease, Leiomyomatosis of esophagus cataract hematuria, Leiomyomatosis, Leiomyosarcoma, Leipala kaitila syndrome, Leishmaniasis, Leisti-Hollister-Rimoin syndrome, Lemierre syndrome, 5 Lenegre disease, Lennox-Gastaut syndrome, Leprechaunism, Leprosy, Leptospirosis, Leri pleonosteosis, Leri-Weill syndrome, Lesch-Nyhan syndrome, Lethal arthrogryposis with anterior horn cell disease (LAAHD), Lethal chondrodysplasia moerman type, Lethal congenital contracture syndrome, Lethal osteosclerotic bone dysplasia, Letterer-Siwe disease, Leucinosi, Leukaemia, 10 Leukocyte adhesion deficiency (LAD), Leukodystrophy, Leukoencephalopathy, Leukonychia totalis, Leukotriene C4 (LTC4) synthase deficiency, Levic stefanovic nikolic syndrome, Levine-Critchley syndrome, Levocardia, Levy-Hollister syndrome, Levy-Yeboah syndrome, Lewis-Pashayan syndrome, Lewis-Sumner syndrome, Lewy body dementia, Leydig cell hypoplasia, Lhermitte-Duclos disease, Li-Fraumeni 15 syndrome, Lichen, Lichstenstein syndrome, Liddle syndrome, Lindsay-Burn syndrome, Linear hamartoma syndrome, Linitis plastica, Lip-pit syndrome, Lipid storage disease, Lipodystrophy, Lipodystrophy-HIV related, Lipoedema, Lipoid proteinosis, Lipomatosis, Lipoprotein metabolism disease, Liposarcoma, Lisker-Garcia-Ramos syndrome, Lissencephaly, Listeriosis, Little syndrome, Lobar atrophy 20 of brain, Lobstein disease, Lobster-claw deformity, Localized Castleman disease, Localized scleroderma, Locked-in syndrome, Loeffler's endocarditis, Loeys-Dietz syndrome, Loffredo cennamo cecio syndrome, Logic syndrome, Loiasis, Long QT syndrome, Longman-Tolmie syndrome, Loose anagen syndrome, Lopes gorlin syndrome, Lopes marques de faria syndrome, Lopez-Hernandez syndrome, Lou- 25 Gehrig disease, Louis-Bar syndrome, Lowe kohn cohen syndrome, Lowe oculocerebrorenal syndrome, Lowe syndrome, Lower mesodermal defects, Lown-Ganong-Levine syndrome, Lowry syndrome, Lowry-MacLean syndrome, Lowry-Yong syndrome, Lubani-Al Saleh-Teebi syndrome, Lubinsky syndrome, Lubs-Arena Syndrome, Lucey driscoll syndrome, Lucky gelehrter syndrome, Lujan-Fryns 30 syndrome, Lunatomalacia, Lundberg syndrome, Lung agenesis heart defect thumb anomalies, Lung cancer small cell, Lung fibrosis, Lupus erythematosus, Lurie kletsky syndrome, Luteinizing hormone releasing hormone deficiency with ataxia, Lutz-Richner-Landolt syndrome, Lyell syndrome, Lyme borreliosis, Lyme disease, Lymphangioliomyomatosis, Lymphangioma, Lymphatic filariasis, Lymphatic 35 malformation, Lymphedema, Lymphocyte apoptosis anomaly, Lymphocyte-depleted classical hodgkin lymphoma, Lymphocyte-rich classical hodgkin lymphoma, Lymphocytic colitis, Lymphoid interstitial pneumonia, Lymphomatoid granulomatosis, Lymphoproliferative disease associated with primary immune disease, Lynch lee murday syndrome, Lynch syndrome, Lyngstadaas syndrome, Lysosomal disease,

Lytico-bodig disease, M-CMTC, M/SCHAD, MAD, MADSAM, MAE, MALT lymphoma, MASA syndrome, MCA, MCAD deficiency, MCOPS1, MDC1A, MEB (Muscle-Eye-Brain) syndrome, MEHMO syndrome, MELAS, MEN 1, MEN 2, MERRF syndrome, MGA type I, MHBD deficiency, MIDD, MIRAS, MMEP syndrome, MMND, MNGIE syndrome, MOBA syndrome, MOCOD, MODY syndrome, MORM syndrome, MPPH syndrome, MPS, MRGH, MRKH syndrome, MRXS7, MSA, MTHFR deficiency, MVA syndrome, MYH9, Mac Duffie's syndrome, Mac dermat winter syndrome, Maccario mena syndrome, Macdermot-Patton-Williams syndrome, Machado-Joseph disease, Macias flores garcia cruz rivera syndrome, Mackay shek carr syndrome, Macroglossia, Macrophage or histiocytic tumour, Macrophagic activation syndrome, Macrophagic myofasciitis, Macrothrombocytopenia with leukocyte inclusions, Macular amyloidosis, Macular dystrophy, Macular edema, Madelung's disease, Madras motor neuron disease, Maffucci syndrome, Majeed syndrome, Majewski ozturk syndrome, Major airway collapse, Meleda disease, Malakoplakia, Malakoplasia, Malaria, Malignant fibrous histiocytoma, Malignant germ cell tumor, Malignant hyperpyrexia, Malignant hyperthermia, Malignant mesenchymal tumor, Malignant paroxysmal ventricular tachycardia, Mallory Weiss syndrome, Malouf syndrome, Maltase-glucoamylase deficiency, Maniac-depressive disorders, Manouvrier syndrome, Mansonellosis, Mantle cell lymphoma, Maple syrup urine disease, Marashi gorlin syndrome, Marble brain disease, Marburg disease, Marchiafava-Micheli disease, Marcus-Gunn syndrome, Marden walker like syndrome, Marfan syndrome, Margarita island ectodermal dysplasia, Marin-Amat syndrome, Marinesco-Sjogren syndrome, Marion mayers syndrome, Markel-Vikkula -Mulliken syndrome, Marles greenberg persaud syndrome, Maroteaux cohen solal bonaventure syndrome, Maroteaux le merrer bensahel syndrome, Maroteaux stanescu cousin syndrome, Maroteaux-Lamy syndrome, Maroteaux-Malamut syndrome, Marsden nyhan sakati syndrome, Marshall syndrome, Marshall-Smith syndrome, Martinez monasterio pinheiro syndrome, Martinez-Frias syndrome, Martsof syndrome, Massa casaer ceulemans syndrome, Mast cell leukaemia, Mast cell sarcoma, Mastocytosis, Mastroiacovo de rosa satta syndrome, Mathieu de broca bony syndrome, Matsoukas liarikos giannika syndrome, Matthew-Wood syndrome, Mature B-cell tumour, Mature T-cell and NK-cell tumour, May-Hegglin thrombocytopenia, Mayer-Rokitansky-Küster-Hauser syndrome, Mazabraud syndrome, McArdle disease, McCabe's disease, McCune-Albright syndrome, McDonough syndrome, McDowall syndrome, McGrath syndrome, McKusick-Kaufman syndrome, McLeod syndrome, McPherson-Hall syndrome, Mcalister crane syndrome, McCallum macadam johnston syndrome, McGillivray syndrome, McLain - Dekaban syndrome, Mcpherson clemens syndrome, Meacham winn culler syndrome, Meadows' syndrome, Meckel like syndrome, Meckel syndrome, Meckel-Gruber

syndrome, Meconium aspiration syndrome, Medeira dennis donnai syndrome, Mediastinal (thymic) large b-cell lymphoma, Mediastinal diffuse large-cell lymphoma with sclerosis, Mediastinal fibrosis, Medrano roldan syndrome, Medullar disease, Medullary cystic kidney disease, Medulloblastoma, Megacalycosis, Megaduodenum
5 and/or megacystis, Megaloblastic anaemia, Megarbane-Loiselet syndrome, Mehes syndrome, Mehta-Lewis-Patton syndrome, Meier blumberg imahorn syndrome, Meier-Gorlin syndrome, Meige disease, Meinecke pepper syndrome, Meinecke syndrome, Melanoma, Meleda disease, Melhem fahl syndrome, Melioidosis, Melkersson rosenthal syndrome, Melnick-Needles syndrome, Melorheostosis,
10 Membranoproliferative glomerulonephritis, Membranous glomerulopathy, Menetrier's disease, Mengel konigsmark syndrome, Meniere's disease, Meningioma, Meningitis, Menkes syndrome, Mental retardation, Meretoja syndrome, Merkel cell carcinoma (MCC), Merlob grunebaum reisner syndrome, Mesangial sclerosis, Mesodermic dysplasia, Mesothelioma, Mesulam syndrome, Metabolic intoxication disease,
15 Metabolic liver disease, Metaphyseal dysplasia, Michels syndrome, Mickleson syndrome, Micro syndrome, Microcephaly, Microcoria, Microcystic infiltrating lymphatic malformation, Microcytic anaemia, Microphthalmia, Microscopic colitis Microtia, Microvillous inclusion disease, Mid-aortic dysplastic syndrome, Midas syndrome, Middle aortic syndrome, Midline heart, Mietens syndrome, Mievis verellen
20 dumoulin syndrome, Mikati najjar sahli syndrome, Mikulicz disease, Mild campomelic dysplasia, Miller syndrome, Miller-Dieker syndrome, Miller-Fisher syndrome (MFS), Mills syndrome, Milroy disease, Minimal change nephrotic syndrome (MCNS), Minkowski-Chauffard disease, Mirhosseini-Holmes-Walton syndrome, Mitral valve prolapse disease, Miura syndrome, Mixed connective tissue disease, Mixed
25 phenotype acute leukaemia, Mixed sclerosing bone dystrophy, Miyoshi myopathy, Mls syndrome, Moderate and severe traumatic brain injury, Moebius syndrome, Moerman vandenbergh frans syndrome, Moersch-Woltman syndrome, Moeschler clarren syndrome, Mohr syndrome, Mohr-Tranebjaerg syndrome, Mollica pavone antener syndrome, Moloney syndrome, Momo syndrome, Monilethrix, Mononen-
30 Karnes-Senac syndrome, Monostotic fibrous dysplasia, Montefiore syndrome, Moore-Federman syndrome, Morava-Mehes syndrome, Morgagni-Stewart-Morel syndrome, Morillo cucci passarge syndrome, Morning glory syndrome, Morquio disease, Morris syndrome, Morse rawnsley sargent syndrome, Morvan syndrome, Moschcowitz disease, Mounier-Kuhn syndrome, Mousa-Al Din-Al Nassar syndrome,
35 Movement disease, Mowat-Wilson syndrome, Moya-moya disease, Moynahan syndrome, Mpo deficiency, Msbd syndrome, Mseleni joint disease (MJD), Mucha Habermann Disease, Muckle-Wells syndrome, Mucoepithelial dysplasia, Mucopolysaccharidosis, Mucormycosis, Mucosal pemphigoid, Mucosulfatidosis, Muenke syndrome, Muir-Torre syndrome, Mullerian aplasia,

Multicentric Castleman disease (MCD), Multicentric giant lymph node hyperplasia, Multicentric osteolysis, Multifocal acquired demyelinating sensory and motor neuropathy, Multifocal pattern dystrophy simulating fundus flavimaculatus, Multiglandular hyperplasia, Multiminicore disease (MmD), Multinodular goiter cystic
5 kidney polydactyly, Multiple carboxylase deficiency, Multiple contracture syndrome, Multiple cutaneous and uterine leiomyomas, Multiple endocrine neoplasia, Multiple epiphyseal dysplasia, Multiple fibrofolliculoma, Multiple hamartoma syndrome, Multiple keratoacanthoma, Multiple pterygium syndrome, Multiple sclerosis, Multiple sulfatase deficiency, Multiple system atrophy, Multiple ventricular septal defects,
10 Mulvihill-Smith syndrome, MURCS association, Murray-Puretic-Drescher syndrome, Muscular channelopathy, Muscular dystrophy, Muscular fibrosis multifocal obstructed vessels, Mutchinick syndrome, Myalgia eosinophilia associated with tryptophan, Myasthenia gravis, Myasthenic syndromes, Mycetoma, Mycoplasma encephalitis, Mycosis fungoides, Myelinoclastic diffuse sclerosis, Myelinosis centralis diffusa,
15 Myelocerebellar disorder, Myelodysplastic or myeloproliferative disease, Myelofibrosis with myeloid metaplasia, Myeloid sarcoma, Myeloma, Myhre syndrome, Myiasis, Myoclonic dystonia, Myoclonic epilepsy, Myodysplasia, Myofibrillar myopathy, Myoglobinuria, Myopathy and diabetes mellitus, Myopathy, Myopia, Myositis ossificans progressiva, Myotilinopathy, Myotonia congenita,
20 Myotonic disease, Myotubular myopathy, Myxofibrosarcoma, Myxoid liposarcoma, Myxoid malignant fibrous histiocytoma, Myxoma with fibrous dysplasia, Möbius syndrome, N syndrome, NACG, NAGS deficiency, NAME syndrome, NAO syndrome, NARP syndrome, NASH syndrome, NBS, NCL, NCMD, NF 1, NFJ syndrome, NHL, NHPP, NISCH syndrome, NOMID syndrome, NPLCA, NSIP, NTD,
25 Naegeli syndrome, Naegeli-Franceschetti-Jadassohn syndrome, Nager syndrome, Naguib syndrome, Nail anomaly, Nail dysplasia, Naito-Oyanagi disease, Nakagawa's angioblastoma, Nakajo nishimura syndrome, Nakajo syndrome, Nakamura osame syndrome, Nance-Horan syndrome, Narcolepsy without cataplexy, Narcolepsy-Cataplexy, Nasodigitoacoustic syndrome, Nasopharyngeal cancer, Nasu-Hakola
30 disease, Nathalie syndrome, Navajo brainstem syndrome, Naxos disease, Necrotising hypophysitis, Necrotizing myelitis, Nemaline myopathy, Neonatal Onset Multisystem Inflammatory Disease, Neonatal death immune deficiency, Neonatal hemochromatosis, Neonatal neutropenia, Neonatal respiratory distress syndrome, Nephroblastoma, Nephrogenic fibrosing dermopathy, Nephrogenic systemic fibrosis,
35 Nephrolithiasis, Nephronophthisis - hepatic fibrosis, Nephropathy, Nephrosis, Nephrotic syndrome with diffuse mesangial sclerosis, Nephrotic syndrome, Nervous system tumour, Netherton disease, Neu-Laxova syndrome, Neuhauser daly magnelli syndrome, Neuhauser eichner opitz syndrome, Neuhauser's anomaly, Neural crest tumour, Neuroacanthocytosis, Neuroaxonal dystrophy, Neuroblastoma,

Neurocutaneous melanosis, Neurodegeneration due to 3-hydroxyisobutyryl-CoA hydrolase deficiency, Neurodegeneration with brain iron accumulation (NBIA), Neurodegenerative disease, Neuroectodermal syndrome, Neuroepithelioma, Neurofibromatosis, Neurolipomatosis, Neuromuscular junction disease, Neuromyelitis optica, Neuromyotonia, Neuropathy, Neutral Lipid Storage Disease, Neutropaenia, Nevo syndrome, Nevoid hypermelanosis, Nezelof syndrome, Nicolaides baraitser syndrome, Niemann-Pick disease, Nievergelt syndrome, Niikawa-Kuroki syndrome, Nijmegen breakage syndrome, Nivelon-Nivelon-Mabille syndrome, Noack syndrome, Noble bass sherman syndrome, Nocardiosis, Nodular lymphocyte predominant Hodgkin lymphoma, Nodulosis-arthropathy-osteolysis syndrome, Noma, Non-24-Hour Sleep-Wake syndrome, Non-DYT1 idiopathic torsion dystonia, Non-Hodgkin malignant lymphoma, Non-alcoholic steatohepatitis, Non-amyloid monoclonal immunoglobulin deposition disease, Non-giant cell granulomatous temporal arteritis with eosinophilia, Non-infectious uveitis affecting the posterior segment of the eye, Nonaka myopathy, Nondysgerminomatous germ cell tumor, Noonan like contracture myopathy hyperpyrexia, Noonan like syndrome, Noonan syndrome, Normomorphich sialidosis, Norrie disease, Norum disease, Nova syndrome, Novak syndrome, Nuclear cell envelopathy, O donnell pappas syndrome, O'Doherty syndrome, O'Sullivan-McLeod syndrome, OA-1, OCA, OCRL1, OFC syndrome, OFCD syndrome, OHSS, OLEDAID, ONMR syndrome, OPPG, ORW 2, OSLAM syndrome, OSMED, OTUDP syndrome, Obliterative portal venopathy, Occlusive infantile arteriopathy, Occupational allergic alveolitis, Ochoa syndrome, Ochronosis, Oculo skeletal renal syndrome, Oculo-osteo-cutaneous syndrome, Oculoectodermal syndrome, Oculogastrointestinal muscular dystrophy, Oculomotor palsy, Oculomotor paralysis, Oculopharyngodistal myopathy, Odontologic disease, Odontomatosis, Oerter-Friedman-Anderson syndrome, Oesophageal atresia, Oguchi disease, Ohaha syndrome, Ohdo madokoro sonoda syndrome, Ohtahara syndrome, Okamoto syndrome, Okihiro syndrome, Oligocone syndrome, Oligomeganephronia, Oliver mcfarlane syndrome, Oliver syndrome, Ollier disease, Olmsted syndrome, Omenn syndrome, Omodysplasia, Onat syndrome, Onchocerciasis, Ondine syndrome, Ondine-Hirschsprung disease, Onychodystrophy, Oochs syndrome, Ophtalmic ichthyosis, Ophtalmoplegia, Opitz BBB/G syndrome, Opitz reynolds fitzgerald syndrome, Opitz-Caltabiano syndrome, Opitz-Frias syndrome, Oppenheim's dystonia, Opsismodysplasia, Opsoclonus-myoclonus syndrome, Optic atrophy, Optic nerve hypoplasia, Optic neuropathy, Optic pathway glioma, Orbital leiomyoma, Ormond's disease, Ornithine aminotransferase deficiency, Orofaciodigital syndrome, Oromandibular dystonia, Oroticaciduria, Oroya fever, Osebold-Remondini syndrome, Osgood-Schlatter disease, Osler-Vaquez disease, Osteoarthropathy, Osteoblastoma, Osteochondritis, Osteochondromas, Osteochondrosis,

Osteocraniostenosis, Osteodysplasia, Osteoectasia, Osteogenic sarcoma, Osteolysis, Osteomesopyknosis, Osteonecrosis, Osteopaenia, Osteopathia striata - cranial sclerosis, Osteopetrosis, Osteopoikilosis, Osteoporosis, Osteosarcoma, Osteosclerosis, Ostravik lindemann solberg syndrome, Otosclerosis, Ouvrier billson syndrome, Ovarian Sertoli-Leydig cell tumor, Ovarian cancer, Ovarian germ cell malignant tumor, Ovarioleukodystrophy, Oxalosis, PAF, PAGOD syndrome, PAN, PANDAS, PAP, PAPA syndrome, PARC syndrome, PCA, PCARP, PCH with optic atrophy, PCT, PDALS, PEHO syndrome, PEL, PELVIS syndrome, PFAPA syndrome, PFIC, PHACE syndrome, PIBIDS syndrome, PJS, PLOSL, PMD, PNDM, POADS, POEMS syndrome, POF, POMC deficiency, PPA, PPHS, PPM-X, PPoma, PSEK, PSP, PTC-RCC, PTLAH, PTLD, Pachygyria, Pachyonychia, Pacman dysplasia, Paediatric Autoimmune Disorders Associated with Streptococcus infections, Paediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcus infections, Paediatric granulomatous arthritis, Paget disease, Pagon stephan syndrome, Pai syndrome, Pallister W syndrome, Pallister-Hall syndrome, Pallister-Killian syndrome, Palmer pagon syndrome, Palpebral disease, Panayiotopoulos syndrome, Pancreatic carcinoma, Pancreatitis, Panner disease, Panniculitis, Panostotic fibrous dysplasia, Papillo-renal syndrome, Papillomatosis, recurrent respiratory, Papillon-Leage-Psaume syndrome, Papillon-Lefevre syndrome, Papular and sclerodermoid lichen myxedematosus, Papular atrichia, Papular mucinosis of infancy, Paramyotonia, Paraneoplastic pemphigus, Paraneoplastic retinopathy, Paraplegia, Parathyroid carcinoma, Parenchymatous liver disease, Paris-Trousseau thrombocytopenia, Parkes-Weber syndrome, Parkinson disease, Parkinsonism-dementia-ALS complex, Paroxysmal cold haemoglobinuria, Paroxysmal exertion-induced dyskinesia, Paroxysmal ventricular fibrillation, Parry-Romberg syndrome, Parsonage-Turner syndrome, Partial deep dermal and full thickness burns, Partington amyloidosis, Partington disease, Partington-Anderson syndrome, Partington-Mulley syndrome, Parvovirus antenatal infection, Pascual castroviejo syndrome, Pashayan syndrome, Passwell-Goodman-Siprkowski syndrome, Patau syndrome, Patterned dystrophy of the retinal pigment epithelium, Patterson pseudoleprechaunism syndrome, Patterson stevenson syndrome, Patterson-Lowry rhizomelic dysplasia, Pauciarticular chronic arthritis, Pavone fiumara rizzo syndrome, Pearson syndrome, Peeling skin syndrome, Pelget-Huer anomaly, Pelizaeus-Merzbacher brain sclerosis, Pellagra, Pemphigus, Pena-Shokeir syndrome, Pendred syndrome, Penta X syndrome, Pentosuria, Peptide metabolism disease, Peptidic growth factors deficiency, Perheentupa syndrome, Periarthritis nodosa, Pericardial defect diaphragmatic hernia, Pericarditis, Perineurioma, Peripartum cardiomyopathy, Peripheral T-cell lymphoma, Peripheral neuropathy and optic atrophy, Peritoneal leiomyomatosis, Peritumoral oedema derived from brain tumours,

Periventricular nodular heterotopia, Perlman syndrome, Pernicious anaemia, Perniola krajewska carnevale syndrome, Peroxisomal beta-oxidation disease, Perrault syndrome, Persistent Mullerian duct syndrome, Peters anomaly, Peters-plus syndrome, Petges-Clejat syndrome, Petit-Fryns syndrome, Petty laxova wiedemann syndrome, Peutz-Jeghers syndrome, Peyronie syndrome, Pfeiffer mayer syndrome, Pfeiffer palm teller syndrome, Pfeiffer rockelein syndrome, Pfeiffer syndrome, Pfeiffer-Kapferer syndrome, Pfeiffer-Singer-Zschiesche syndrome, Pfeiffer-Weber-Christian syndrome, Phacomatosis, Phaeochromocytoma, Phagocyte function anomaly, Phaver syndrome, Phelan-McDermid syndrome, Phenotypic diarrhoea, Phenylketonuria, Phocomelia, Phytosterolemia, Picardi-Lassueur-Little syndrome, Pick disease of brain, Piebaldism, Pierre Robin sequence associated with branchial archs anomalies, Pierre Robin sequence associated with collagen diseases, Pigeon-breeder's lung disease, Pillay syndrome, Pilomatrixoma, Pilotto syndrome, Pinheiro freire maia miranda syndrome, Pinsky-Di George-Harley syndrome, Pitt-Hopkins syndrome, Pitt-Williams brachydactyly, Pitt-rogers-danks syndrome, Pituitary adenoma, Pituitary agenesis, Pituitary hormone deficiency, Pituitary lactotrophic adenoma, Pityriasis rubra pilaris, Piussan-Lenaerts-Mathieu syndrome, Plasma cell tumour, Platelet function disease, Platyspondylic dysplasia, Plectin deficiency, Pleomorphic liposarcoma, Pleuro-pulmonary blastoma, Pleuro-pulmonary endometriosis, Plott syndrome, Plum syndrome, Plummer-Vinson syndrome, Pneumoblastoma, Pneumocystosis, Pneumonia caused by Pseudomonas Aeruginosa, Poikilo-dermatomyositis, Pollitt syndrome, Polyarteritis nodosa, Polyarthrititis, Polycystic kidney disease, Polycystic liver disease, Polycystic ovarian disease, Polycythaemia, Polydactyly, Polyepiphyseal dysplasia, Polymicrogyria, Polymorphic catecholergic ventricular tachycardia, Polymyositis, Polyostotic fibrous dysplasia, Polyposis, Polysyndactyly - cardiac malformation, Pompe disease, Popliteal web syndrome, Porokeratosis, Porphyria, Portal hypertension, Portal vein thrombosis, Post polio syndrome, Post transplantation graft dysfunction, Post-poliomyelitic syndrome, Post-transplant lymphoproliferative disease, Post-traumatic syringomyelia, Postanginal sepsis secondary to orophyngal infection, Posterior cortical atrophy, Postpartum cardiomyopathy, Postviral Fatigue Syndrome, Potocki-Shaffer syndrome, Potter sequence, Powell chandra saal syndrome, Powell venencie gordon syndrome, Prader-Willi syndrome, Prata liberal goncalves syndrome, Preauricular pits renal disease, Precursor B-cell acute lymphoblastic leukaemia, Precursor T-cell acute lymphoblastic leukaemia, Preeyasombat-Varavithya syndrome, Pregnancy-related cholestasis, Premature aging, Pressure-induced localized lipotrophy, Prieto-Badia-Mulas syndrome, Prieur-Griscelli syndrome, Primary biliary cirrhosis, Primary ciliary dyskinesia, Primary cutaneous CD30-positive T-cell lymphoproliferative disorders, Primary effusion lymphoma, Primary effusion

lymphoma associated with the human immunodeficiency virus (HIV) infection, Primary intestinal lymphangiectasia, Primary lateral sclerosis, Primary lipodystrophy, Primary lymphoedema, Primary pulmonary lymphoma, Primary sclerosing cholangitis, Primerose syndrome, Progeria, Progressive bulbar paralysis of childhood, Progressive cone dystrophy, Progressive diaphyseal dysplasia, Progressive massive osteolysis, Progressive nephropathy with hypertension, Progressive neuronal degeneration of childhood with liver disease, Prolactinoma, Popping Zerres syndrome, Prostate cancer, Proteus syndrome, Proud-Levine-Carpenter syndrome, Prune belly syndrome, Psoriatic arthritis, PTEN Hamartoma syndrome, Pterygia, Pudendal neuralgia, Pudendal neuropathy, Pulmonar arterioveinous aneurysm, Pulmonary Langerhans' cell histiocytosis, Pulmonary alveolar microlithiasis, Pulmonary alveolar proteinosis, Pulmonary aortic stenosis, Pulmonary arterial hypertension, Pulmonary arterio-veinous fistula, Pulmonary artery hypoplasia, Pulmonary atresia, Pulmonary blastoma, Pulmonary branch stenosis, Pulmonary endometriosis, Pulmonary haemosiderosis, Pulmonary insufficiency, Pulmonary lymphangiectasia, Pulmonary lymphangiomatosis, Pulmonary nodular lymphoid hyperplasia, Pulmonary supra-valvular stenosis, Pulmonary surfactant protein anomalies, Pulmonary valve agenesis (PVA), Pulmonary venoocclusive disease, Pulp stones, Pulpal dysplasia, Pueric syndrome, Purtilo syndrome, Pycnodysostosis, Pyknoachondrogenesis, Pyknolepsy, Pyle disease, Pyoderma gangrenosum, Pyomyositis, Pyropoikilocytosis, Q fever, Qazi-Markouizos syndrome, Quattrin mcpherson syndrome, RAEB-1, RAPADILINO syndrome, RB-ILD, RECQ2, RECQL3, RHS, Rabson-Mendenhall syndrome, Radiation syndromes, Radio renal syndrome, Raine syndrome, Rajab-Spranger syndrome, Rambam-Hasharon syndrome, Rambaud galian syndrome, Ramon syndrome, Ramos arroyo clark syndrome, Ramsay hunt syndrome, Randall disease, Rapp-Hodgkin ectodermal dysplasia syndrome, Rapp-Hodgkin syndrome, Rasmussen johnsen thomsen syndrome, Rasmussen syndrome, Rathburn disease, Ray peterson scott syndrome, Raynaud phenomenon, Reardon-Baraitser syndrome, Reardon-Hall-Slaney syndrome, Recurrent hepatitis C virus induced liver disease in liver transplant recipients, Red cell aplasia, Refetoff syndrome, Reflex sympathetic dystrophy syndrome, Refsum disease, Reginato-Schiapachasse syndrome, Reifenstein syndrome, Reinhardt pfeiffer syndrome, Reiter's syndrome, Renal adysplasia, Renal cell carcinoma, Renal dysplasia, Renal glucosuria, Renal hypertension, Renal hypoplasia, Renal nutcracker syndrome, Renal tubular acidosis, Renal tubular disorder, Renal-coboloma syndrome, Rendu-Osler-Weber disease, Renier-Gabreels-Jasper syndrome, Renpenning syndrome, Resistance to activated protein C, Resistance to thyroid stimulating hormone, Respiratory bronchiolitis, Restless legs syndrome, Restrictive cardiomyopathy, Reticular perineurioma, Retinal

arteriolar tortuosity, Retinal degeneration, Retinal dystrophy, Retinal hemorrhage, Retinoblastoma, Retinohepatoendocrinologic syndrome, Retinopathy of prematurity, Retinoschisis with early hemeralopia, Retinoschisis, Retraction syndrome, Retroperitoneal fibrosis, Rett like syndrome, Rett syndrome, Revesz-Debuse
 5 syndrome, Reye's syndrome, Reynolds syndrome, Rh deficiency syndrome, Rhabdomyosarcoma, Rheumatic fever, Rhizomelic dysplasia, Rhnull syndrome, Richards-Rundle syndrome, Richardson's syndrome, Richieri Costa-Guion Almeida-Cohen syndrome, Richieri costa da silva syndrome, Richieri costa gorlin syndrome, Richieri-Costa-Colletto syndrome, RichieriCosta-Pereira syndrome, Richner-Hanhart
 10 syndrome, Ricker syndrome, Rickettesiae disease, Riedel Thyroiditis, Rieger syndrome, Right atrium familial dilatation, Right ventricle hypoplasia, Rigid spine syndrome, Riley-Day syndrome, Riley-Smith syndrome, Rippberger aase syndrome, Rippling muscle disease, Ritscher schinzel syndrome, Rivera-Perez-Salas syndrome, Roberts syndrome, Robinow syndrome, Robinow-Sorauf syndrome, Robinow-Unger
 15 syndrome, Robinow-like syndrome, Roch-Leri mesosomatous lipomatosis, Rocky Mountain spotted fever, Rodini richieri costa syndrome, Roger disease, Roifman-Melamed syndrome, Rokitansky syndrome, Romano-Ward long QT syndrome, Rombo syndrome, Rommen mueller sybert syndrome, Rosai-Dorfman disease, Rosenberg lohr syndrome, Rosenberg Chutorian syndrome, Rothmund-Thomson
 20 syndrome, Rotor syndrome, Roy maroteaux kremp syndrome, Rozin-Hertz-Goodman syndrome, Rubinstein-Taybi syndrome, Rudd-Klimek syndrome, Rudiger syndrome, Russell Silver syndrome, Russell weaver bull syndrome, Rutherford syndrome, Ruvalcaba syndrome, Ruvalcaba-Myhre-Smith syndrome, SADDAN, SANDO, SAPHO syndrome, SC phocomelia, SCA, SCAN 2, SCAR1, SCARF syndrome,
 25 SCASI, SCD, SCID, SCLC, SE(M)D, SGBS, SGS, SHORT syndrome, SIADH, SIBIDS syndrome, SJS, SLK, SMD, SMEI, SMMCI, SOD, SOLAMEN syndrome, SPG, SPONASTRIME dysplasia, SPS, SRP, SUNCT syndrome, Saal-Greenstein syndrome, Saccharopinuria, Sack-Barabas syndrome, Saethre-Chotzen syndrome, Saito kuba tsuruta syndrome, Sakati syndrome, Sakati-Nyhan syndrome, Sakati-
 30 Nyhan-Tisdale syndrome, Salcedo syndrome, Salla disease, Salmonellosis, Salti salem syndrome, Sammartino decreccio syndrome, San Luis Valley syndrome, Sandhoff disease, Sandifer syndrome, Sandrow syndrome, Sanfilippo disease, Sanjad-Sakati syndrome, Santavuori disease, Santos-Mateus-Leal syndrome, Sarcocystosis, Sarcoidosis, Sarcosinemia, Sarcosporidiosis, Satoyoshi syndrome,
 35 Say barber hobbs syndrome, Say barber miller syndrome, Say field coldwell syndrome, Say meyer syndrome, Scarring in glaucoma filtration surgical procedures, Schaap taylor baraitser syndrome, Scheie syndrome, Scheuermann disease, Schilbach-Rott syndrome, Schilder disease, Schimke syndrome, Schimmelpenning syndrome, Schindler disease, Schinzel syndrome, Schisis association,

Schistosomiasis, Schmidt syndrome, Schmitt gillenwater kelly syndrome,
 Schneckenbecken dysplasia, Schnitzler syndrome, Schofer-Beetz-Bohl syndrome,
 Scholte begeer van essen syndrome, Schopf-Schulz-Passarge syndrome,
 Schwannomatosis, Schwartz-Jampel syndrome, Scimitar syndrome, Scleroatrophic
 5 syndrome, Scleroderma, Scleromyxedema, Sclerosing mediastinitis, Sclerosteosis,
 Scott syndrome, Scott-Bryant-Graham syndrome, Scott-Taor syndrome, Seaver
 cassidy syndrome, Sebastian syndrome, Seckel like syndrome, Seckel syndrome,
 Sedlackova syndrome, Seemanova lesny syndrome, Segawa syndrome, Seghers
 syndrome, Seitelberger disease, Selig-Benacerraf-Greene syndrome, Sellars-
 10 Beighton syndrome, Sengers syndrome, Sengers-Hamel-Otten syndrome, Senior
 syndrome, Senior-Boichis syndrome, Senior-Loken syndrome, Sensenbrenner
 syndrome, Senter syndrome, Sepsis, Septic phlebitis of the internal jugular vein,
 Sequeiros sack syndrome, Servelle-Martorell syndrome, Setleis syndrome, Severe
 closed traumatic brain injury, Severe combined immunodeficiency T- B-, Severe
 15 combined immunodeficiency with hypereosinophilia, Severe combined
 immunodeficiency with leukopenia, Severe pneumococemia, Sezary's lymphoma,
 Shapiro syndrome, Sharma kapoor ramji syndrome, Sharp syndrome, Sheehan
 syndrome, Shigellosis, Shokeir syndrome, Shone syndrome, Short QT syndrome,
 Short bowel syndrome due to necrotizing enterocolitis, Short bowel syndrome due to
 20 thrombosis, Short bowel syndrome, Shprintzen omphalocele syndrome, Shprintzen-
 Goldberg syndrome, Shulman syndrome, Shwachman-diamond syndrome, Shy-
 drager syndrome, Sialidosis, Sick cell anaemia, Sideroblastic anaemia, Sidransky-
 Feinstein-Goodman syndrome, Siegler brewer carey syndrome, Silengo lerone
 pelizzo syndrome, Sillence syndrome, Simosa penchaszadeh bustos syndrome,
 25 Simpson dysmorphia syndrome (SDYS), Simpson-Golabi-Behmel syndrome,
 Sinding-Larsen-Johansson disease, Singh chhaparwal dhanda syndrome, Singh-
 Williams-McAlister syndrome, Single ventricular septal defect, Singleton-Merten
 dysplasia, Singleton-Merten syndrome, Sino-auricular heart block, Sinus node
 disease and myopia, Sipple syndrome, Sirenomelia, Sitosterolemia, Situs inversus
 30 viscerum-cardiopathy, Sjögren syndrome, Sjögren-Larsson syndrome, Skeletal
 dysplasia, Skeletal muscle disease, Skin collagen disease, Skin vascular disease,
 Sleep disorder, Sleeping seekness, Sly disease, Small bowel adenocarcinoma, Small
 bowel leiomyosarcoma, Small non-cleaved cell lymphoma, Smith martin dodd
 syndrome, Smith-Fineman-Myers syndrome, Smith-Lemli-Opitz syndrome, Smith-
 35 Magenis syndrome, Sneddon syndrome, Sneddon-Wilkinson disease, Snyder-
 Robinson syndrome, Soft tissue perineurioma, Soft tissue sarcomas, Sohval suffer
 syndrome, Solitary plasmacytoma, Solomon syndrome, Somatotroph adenoma,
 Sommer hines syndrome, Sommer rathbun battles syndrome, Sommer-Young-Wee-
 Frye syndrome, Sondheimer syndrome, Sonoda syndrome, Sorsby syndrome,

Sorsby's fundus dystrophy, Sotos syndrome, Spastic paraplegia, Spellacy gibbs
 watts syndrome, Spherophakia-brachymorphia, Sphingolipidosis, Spina bifida,
 Spinal atrophy, Spirillosis, Splenic marginal zone lymphoma, Spondylarthropathy,
 Spondylo camptodactyly syndrome, Spondylocostal dysostosis,
 5 Spondyloenchondrodysplasia, Spondyloepiphyseal dysplasia, Spongy degeneration
 of central nervous system, Spongy myocardium, Spontaneous pneumothorax familial
 type, Sporotrichosis, Squamous cell carcinoma of head and neck, St Louis
 encephalitis, Stalker chitayat syndrome, Stampe sorensen syndrome, Stapedo-
 vestibular ankylosis, Staphylococcal necrotizing pneumonia, Staphylococcal scarlet
 10 fever, Staphylococcal toxic shock syndrome, Stargardt disease, Stark-Kaesler
 syndrome, Startle disease, Steatocystoma, Steele-Richardson-Olszewski disease,
 Stein-Leventhal syndrome, Steinert myotonic dystrophy, Steinfeld syndrome, Stern-
 Lubinsky-Durrie syndrome, Stevens-Johnson syndrome, Stickler syndrome, Stiff
 person syndrome, Still disease, Stimmler syndrome, Stoelinga de koomen davis
 15 syndrome, Stoll alembik finck syndrome, Stoll geraudel chauvin syndrome, Stoll
 kieny dott syndrome, Stoll-Levy-Francfort syndrome, Stomach cancer, Stormorken-
 Sjaastad-Langslet syndrome, Stratton garcia young syndrome, Stratton parker
 syndrome, Streptobacillosis, Streptococcal toxic-shock syndrome, Stress
 cardiomyopathie, Strumpell-Lorrain disease, Sturge-Weber syndrome, Stuve-
 20 Wiedemann dysplasia, Subcutaneous panniculitis-like T-cell lymphoma,
 Subpulmonary stenosis, Sucking/swallowing disorder, Sudden infant death
 syndrome, Sugarman syndrome, Sujansky-Leonard syndrome, Sulfocysteinuria,
 Summerskill-Walsh-Tygstrup syndrome, Summitt syndrome, Supravalvar aortic
 stenosis, Susac syndrome, Sutton disease II, Sweet syndrome, Swyer syndrome,
 25 Symphalangism, Syncopal paroxysmal tachycardia, Syncopal tachyarrhythmia,
 Syndromic diarrhea, Synoviosarcoma, Synovitis, Synspondylism,
 Syntelencephaly, Syringocystadenoma papilliferum, Syringomyelia, Systemic
 capillary leak syndrome, Systemic lupus erythematosus, Systemic mastocytosis,
 Systemic scleroderma (systemic sclerosis), Systemic vasculitis, T cell
 30 immunodeficiency, T-cell leukaemia, T-cell chronic lymphocytic leukaemia, TAC, TAR
 syndrome, TCP, TDO syndrome, TEMF, TGA, TINU syndrome, TNF receptor 1
 associated periodic syndrome, TOS, TRAPS syndrome, TTP, TTR amyloid
 cardiopathy, TTR amyloid neuropathy, Tabatznik syndrome, Takatsuki syndrome,
 Takayasu arteritis, Takotsubo cardiomyopathy, Tang hsi ryu syndrome, Tangier
 35 disease, Tardive dyskinesia, Tarsal Tunnel syndrome, Tarui disease, Tauopathy,
 Taussig-Bing syndrome, Tay syndrome, Tay-Sachs disease, Taybi syndrome, Taybi-
 Linder syndrome, Teebi al saleh hassoon syndrome, Teebi kaurah syndrome, Teebi
 naguib alawadi syndrome, Teebi shaltout syndrome, Telangiectasia, Telecanthus,
 Telfer sugar jaeger syndrome, Temtamy-Shalash syndrome, Ter Haar syndrome,

Teratoma, Tetraamelia, Tetralogy of Fallot, Thakker donnai syndrome, Thalassaemia syndrome, Thanatophoric dysplasia, Theodore's syndrome, Thiele syndrome, Thiemann disease, Thies-Reis syndrome, Thomas jewett raines syndrome, Thomas syndrome, Thompson baraitser syndrome, Thomsen and Becker disease, Thong douglas ferrante syndrome, Thoracic aortic aneurysm and/or dissection, Thoracic outlet syndrome, Three M disease, Thromboangiitis obliterans, Thrombocytopaenia, Thrombocytopenic purpura autoimmune, Thrombocytopenic purpura idiopathic, Thrombocytosis, Thromboembolic pulmonary hypertension, Thrombotic disease of haematologic origin, Thymic aplasia, Thymic carcinoma, Thyroid tumor, Tick-borne encephalitis, Tietze syndrome, Timothy syndrome, Tollner horst manzke syndrome, Tolosa-Hunt syndrome, Tomaculous neuropathy, Tome brune fardeau syndrome, Toni-Debré-Fanconi disease, Tonoki-Ohura-Niikawa syndrome, TORCH syndrome, Toriello syndrome, Toriello-Carey syndrome, Toriello-Higgins-Miller syndrome, Toriello-Lacassie-Droste syndrome, Torres ayber syndrome, Tourette syndrome, Townes-Brocks syndrome, Toxocariasis, Toxoplasma embryopathy, Toxoplasmosis, Tracheopathia osteoplastica, Tranebjaerg-Svejgaard syndrome, Transmissible spongiform encephalopathies, Transposition of the great arteries with pulmonary stenosis, Transthyretin amyloid polyneuropathy, Treacher-Collins syndrome, Aspiration pneumonitis requiring intubation and mechanical ventilation, Cardiogenic shock, Treft-Sanborn-Carey syndrome, Trench fever, Trevor disease, Triatrial heart, Trichinosis, Tricho onychic dysplasia, Tricho-dento-osseous syndrome, Tricho-hepato-enteric syndrome, Trichorhinophalangeal, Trichorrhexis nodosa syndrome, Trichothiodystrophy, Tricuspid atresia, Triopia, Triple A syndrome, Triple H (HHH) syndrome, Triplo-X syndrome, Trisomy, Tritanopia, Trochlear dysplasia, Tropical calcific chronic pancreatitis, Tropical endomyocardial fibrosis, Trueb burg bottani syndrome, Tsao-Ellingson syndrome, Tsukahara-Kajii syndrome, Tsukahara syndrome, Tsutsugamushi disease, Tsutsugamushi fever, Tuberculosis, Tuberous sclerosis, Tubular duplication of the oesophagus, Tubular dysplasia, Tubular renal disease - cardiomyopathy, Tubulointerstitial nephritis and uveitis syndrome, Tucker syndrome, Tuffli-Laxova syndrome, Tularaemia, Tungiasis, Tungland-Bellman syndrome, Tunnel subaortic stenosis, Turcot syndrome, Turner syndrome, Turner-Kieser syndrome, Twin.twin transfusion syndrome, Tylosis, ULD, UPDM, UPDP, USH, Uhl anomaly, Ulbright hodes syndrome, Ulcerative colitis, Ulerythema ophryogenesis, Ulick syndrome, Ullrich disease, Umbilical cord ulceration, Univentricular cardiopathy, Unverricht-Lundborg disease, Upington disease, Upshaw-Schulman syndrome, Urbach-Wiethe disease, Urban-Rogers-Meyer syndrome, Urban-Schosser-Spohn syndrome, Uremic pruritus, Urrets-Zavalía syndrome, Usher syndrome, Usual interstitial pneumonia (UIP), Uveitis, VIPoma, VMCM, VODI syndrome, VSD, VWS, Vagueur triolle ripert syndrome, Van Allen-

Myhre syndrome, Van Benthem-Driessen-Hanveld syndrome, Van Bogaert disease, Van Der Woude syndrome, Van biervliet hendrickx van ertbruggen syndrome, Van de berghe-Dequeker syndrome, Van den Bosch syndrome, Van den ende brunner syndrome, Van der Knapp syndrome, Van goethem syndrome, Van maldergem
5 wetzburger verloes syndrome, Van regemorter pierquin vamos syndrome, Varadi-Papp syndrome, Vascular leukoencephalopathy, Vasculitis, Vasquez-Hurst-Sotos syndrome, Vasterbotten dystrophy, Vein of Galen aneurysm, Venencie powell winkelmann syndrome, Ventricular septal defect, Ventricular septal defect with aortic insufficiency, Verloes - Gillerot - Fryns syndrome, Verloes bourguignon syndrome,
10 Verloes david syndrome, Verloes van maldergem marneffe syndrome, Verloes-Deprez syndrome, Verloove vanhorick brubakk syndrome, Verneuil disease, Viljoen winship syndrome, Viljoen-Kallis-Voges syndrome, Viljoen-Smart syndrome, Viral hemorrhagic fever, Viral hepatitis, Viral vasculitis, Visceral neuropathy, Vitiligo, Vitreoretinal degeneration, Vogt-Koyanagi-Harada disease, Vohwinkel syndrome,
15 Volcke-Soekarman syndrome, Von Gierke disease, Von Hippel-Lindau disease, Von Recklinghausen disease, Von Voss-Cherstvoy syndrome, Von Willebrand disease, Von hippel anomaly, Vsr syndrome, Vuopala disease, W syndrome, WAGR syndrome, WARBM1, WHIM syndrome, WL syndrome, WT limb-blood syndrome, Waaler-Aarskog syndrome, Waardenburg syndrome, Waardenburg-Shah syndrome,
20 Wagner disease, Waisman syndrome, Waldenström macroglobulinemia, Waldmann disease, Walker-Dyson syndrome, Walker-Warburg syndrome, Wallis cremin beighton syndrome, Wallis zieff goldblatt syndrome, Warburg Micro syndrome, Warburg thomsen syndrome, Warburton-Anyane-Yeboa syndrome, Warman-Mulliken-Hayward syndrome, Water-West syndrome, Waterhouse-Friedrickson
25 syndrome, Watson syndrome, Weaver like syndrome, Weaver syndrome, Weaver-Williams syndrome, Weber-Christian disease (WCD), Weber-Christian panniculitis, Webster deming syndrome, Wegener granulomatosis, Weil syndrome, Weill-Marchesani syndrome, Weismann Netter Stuhl syndrome, Weissenbacher-Zweymuller syndrome, Wellesley-Carman-French syndrome, Wells syndrome, Wells-Jankovic syndrome, Werdnig-Hoffmann disease, Wermer syndrome, Werner syndrome, Wernicke's encephalopathy, Wernicke-Korsakoff syndrome, West syndrome, West-Nile encephalitis, Westerhof-Beemer-Cormane syndrome, Western equine encephalomyelitis, Westphall disease, Whelan syndrome, Whipple disease, Whistling face syndrome, Whooping cough, Whyte-Murphy syndrome, Wieacker-
35 Wolff syndrome, Wiedemann grosse dibbern syndrome, Wiedemann oldigs oppermann syndrome, Wiedemann-Beckwith syndrome, Wiedemann-Rautenstrauch syndrome, Wildervanck syndrome, Wilkes stevenson syndrome, Wilkie-Taylor-Scambler syndrome, Willebrand disease, Willi-Prader syndrome, Williams syndrome, Williams-Beuren syndrome, Wilms tumor, Wilson disease, Wilson-Turner syndrome,

Winchester disease, Winkelman bethge pfeiffer syndrome, Winkelmann's cytophagic
 panniculitis, Winship viljoen leary syndrome, Winter harding hyde syndrome, Winter-
 Shortland-Temple syndrome, Wiskott-Aldrich syndrome, Wissler-Fanconi syndrome,
 Witkop syndrome, Wittwer syndrome, Wolcott-Rallison syndrome, Wolf-Hirschhorn
 5 syndrome, Wolff zimmermann syndrome, Wolff-Parkinson-White syndrome, Wolfram
 syndrome, Wolman disease, Woodhouse sakati syndrome, Woods black norbury
 syndrome, Woods leversha rogers syndrome, Woods-Crouchman-Huson syndrome,
 Worster drought syndrome, Worth syndrome, Wrinkly skin syndrome, Wyburn-Mason
 syndrome, XHIGM, XLAG syndrome, XMEA, XP, Xanthic urolithiasis, Xanthinuria,
 10 Xanthogranulomatous hypophysitis, Xanthomatosis cerebrotendinous, Xerocytosis,
 Xeroderma pigmentosum, Yellow fever, Yellow nail syndrome, Yersiniosis, Yorifuji-
 Okuno syndrome, Yoshimura-takeshita syndrom, Young maders syndrome, Young
 syndrome, Young-Hugues syndrome, Young-Simpson syndrome, Yunis-Varon
 syndrome, ZASP-related myofibrillar myopathy, Zadik-Barak-Levin syndrome,
 15 Zellweger syndrome, Zellweger-like syndrome, Zimmer phocomelia, Zimmerman
 laband syndrome, Zinsser-Cole-Engman syndrome, Zlotogora-Ogur syndrome,
 Zlotogura-Martinez syndrome, Zollinger-Ellison syndrome, Zori stalker williams
 syndrome, Zurich-Kaye syndrome, Zygomycosis, 2,8 dihydroxy-adenine urolithiasis,
 2-aminoadipic aciduria, 2-hydroxyglutaricaciduria, 2-methylbutyric aciduria, 3
 20 hydroxyisobutyric aciduria, 3-hydroxy-3-methylglutaric aciduria, 3-
 methylcrotonylglycinuria, 3-methylglutaconic aciduria, 3C syndrome, 3M syndrome,
 4-hydroxybutyricaciduria, Visceral leishmaniasis, Vernal keratoconjunctivitis, UV-A
 and visible light-induced photosensitivity disorders (chronic actinic dermatitis,
 cutaneous porphyrias, actinic prurigo and solar urticaria), Uremic pruritus, Tricyclic
 25 antidepressants poisoning, Traumatic spinal cord injury, Renal cell carcinoma,
 Superficial bladder cancer, *Staphylococcus aureus* bacteraemia, Spinal cord injury,
 Spina bifida, Soft tissue sarcoma, Small cell lung cancer, Sickle cell disease, Severe
 myoclonic epilepsy in infancy, Severe combined immunodeficiency (SCID), Severe
 closed traumatic brain injury, Retinopathy of prematurity, Retinitis pigmentosa,
 30 Respiratory distress syndrome in premature neonates of less than 32 weeks of
 gestational age, Recurrent hepatitis C virus induced liver disease in liver transplant
 recipients, Radiation proctitis, *Pseudomonas aeruginosa* lung infection in cystic
 fibrosis, Progressive myoclonic epilepsies, Primary malignant bone tumors, Primary
 apnoea of premature newborns, Post-transplant lymphoproliferative disorders, Post-
 35 neonatal intracerebral haemorrhage, Post transplantation graft dysfunction,
 Polycythemia vera, Peritumoral oedema derived from brain tumors, Peripheral T-cell
 lymphoma (nodal, other extranodal and leukaemic/ disseminated), Ductus arteriosus
 in premature neonates of less than 34 weeks of gestational age, Partial deep dermal
 and full thickness burns, Paroxysmal nocturnal haemoglobinuria, Pancreatic cancer,

Painful HIV-associated neuropathy, Ovarian cancer, Osteosarcoma, Orthostatic hypotension in patients with pure autonomic failure, Orthostatic hypotension in patients with multiple system atrophy, Ornithine-transcarbamylase deficiency, Oral mucositis in head and neck cancer patients undergoing radiation therapy,

5 Oesophageal cancer, Non-traumatic osteonecrosis, Non-ketotic hyperglycinaemia, Non-infectious uveitis affecting the posterior segment of the eye, Non-24-hour sleep-wake disorders in blind people with no light perception, Neuroblastoma, Neovascular glaucoma, Nephritic syndrome, Myelodysplastic syndromes, Myasthenia gravis, Moderate and severe traumatic brain injury, Metachromatic leukodystrophy,

10 Medullary thyroid carcinoma, Mastocytosis, Mantle cell lymphoma, Malignant melanoma, Malignant gastrointestinal stromal tumors, Malabsorption due to exocrine pancreatic enzyme insufficiency, Low flow priapism, Lipoprotein lipase deficiency, Ligneous conjunctivitis, Leber's hereditary optic neuropathy, Leber's congenital amaurosis, Late onset sepsis in premature infants of less than or equal to 32 weeks

15 gestational age, Juvenile myelomonocytic leukaemia, Japanese encephalitis, Intestinal graft-versus-host disease, Indolent non-Hodgkin's lymphoma, Inborn errors in primary bile acid synthesis, Hyperphenylalaninemia, Hypereosinophilic syndrome, Glioma, High-grade dysplasia in Barrett's oesophagus, *Herpes simplex* virus stromal keratitis, Hereditary factor XIII deficiency, Hepatocellular carcinoma, Hepatitis B re-

20 infection following liver transplantation, Hepatic veno-occlusive disease, Gram negative bacterial lung infection in cystic fibrosis, Gastric cancer, Gamma sarcoglycanopathy, Follicular lymphoma, Familial adenomatous polyposis, Emphysema secondary to congenital alpha-1 antitrypsin deficiency, Duchenne muscular dystrophy, Diffuse large B cell lymphoma, Diffuse alveolar haemorrhage,

25 Diarrhoea associated with intestinal microsporidial infection, Cutaneous T-cell lymphoma, Cutaneous forms of lupus erythematosus, Cushing's syndrome secondary to ectopic ACTH secretion, Corneal graft rejection, Congenital venous malformations, Congenital lymphatic malformations, Congenital alpha-1 antitrypsin deficiency, Congenital adrenal hyperplasia, Chronic pain, Cocaine poisoning, Chronic

30 myeloid leukaemia, Chronic lymphocytic leukaemia, Chronic iron overload requiring chelation therapy, Chronic idiopathic myelofibrosis, Chronic eosinophilic leukaemia and the hypereosinophilic syndrome, Cholangiocarcinoma, Charcot-Marie-Tooth disease type 1A, Cardiogenic shock, Bronchopulmonary dysplasia in premature neonates of less than 30 weeks of gestational age, B-cell chronic lymphocytic

35 leukemia, Autoimmune uveitis, Atypical Haemolytic Uraemic Syndrome (aHUS) associated with an inherited abnormality of the complement system, Aspiration pneumonitis requiring intubation and mechanical ventilation, Aneurysmal subarachnoid haemorrhage, Anaplastic thyroid cancer, Anal fistula, Acute sensorineural hearing loss (acute acoustic trauma, sudden deafness and surgery

induced acoustic trauma), Acute peripheral arterial occlusion, Acute intermittent porphyria, Active phase of Peyronie's disease, Acanthamoeba keratitis, A-mannosidosis, 5q spinal muscular atrophy, Cavopulmonary Anastomosis, Atrial Septal Defects (ASD), Partial Anomalous Pulmonary Venous Return, Persistent
5 Common Atrio Ventricular Canal Endocardial Cushion Defect. Ostium Primum, Single Atrium, Patent Ductus Arteriosus (PDA), Total Anomalous Pulmonary Venous Return, Ventricular Septal Defects (VSD), Pulmonary Valve Stenosis, Pulmonary Artery Stenosis and Stenosis of Pulmonary Artery Branches, Pulmonary Atresia with Intact Ventricular Septum, Congenital Mitral Valve Disease, Aortic Valvular Stenosis
10 and Congenital Aortic Valvular Regurgitation, Supravalvular Aortic Stenosis, Transposition of the Great Arteries, Double Outlet Right Ventricle, Corrected Transposition of the Great Arteries, Truncus Arteriosus, Aorto Pulmonary Window, Tricuspid Atresia, Ebstein Anomaly, Malformations of the Vena Cava, Coarctation of the Aorta, Atresia of Aortic Valve, Anomalies of the Aortic Arch, Anomalous Origin of
15 the Right Subclavian Artery with Coarctation of the Aorta, Idiopathic Dilatation of the Pulmonary Artery, Left Pulmonary Artery Arising from Right Pulmonary Artery, Dextrocardia - Situs Inversus Totalis, Association of Heart Malformations with Asplenia, Malformations of the Vena Cava, Congenital Coronary Artery Arterio-Venous Fistula, Abnormal Origin of the Coronary Arteries, Aneurysm of the Sinus of
20 Valsalva (Aortic Sinus Aneurysm), Endocardial Fibroelastosis, Idiopathic Hypertrophic Subaortic Stenosis (IHSS), Mitral Valve Prolapse - Barlow's Syndrome, Hypoplastic Left Heart.

Pharmaceutical compositions

25 Still another aspect of the present invention relates to the use of the peptide of the invention and the inventive peptide combination as an active ingredient, together with at least one pharmaceutically acceptable carrier, excipient and/or diluents for the manufacture of a pharmaceutical composition for the treatment and/or prophylaxis of cancer, an autoimmune disease, a fibrotic disease, an inflammatory disease, a
30 neurodegenerative disease, an infectious disease, a lung disease, a heart and vascular disease or a metabolic disease or any other disease disclosed herein.

Such pharmaceutical compositions comprise the peptide or the peptide combination as an active ingredient, together with at least one pharmaceutically acceptable
35 carrier, excipient, binders, disintegrates, glidants, diluents, lubricants, coloring agents, sweetening agents, flavoring agents, preservatives or the like. The pharmaceutical compositions of the present invention can be prepared in a conventional solid or liquid carrier or diluents and a conventional pharmaceutically-made adjuvant at suitable dosage level in a known way. Preferably, the two

peptides are contained in the combination in an amount from 20% by weight of peptide 1 to 80% by weight of peptide 2 to 80% by weight of peptide 1 to 20% by weight of peptide 2. More preferably, the two peptides are contained in the combination in an amount from 30% by weight of peptide 1 to 70% by weight of peptide 2 to 70% by weight of peptide 1 to 30% by weight of peptide 2. Still more preferably the two peptides are contained in the combination in an amount from 40% by weight of peptide 1 to 60% by weight of peptide 2 to 60% by weight of peptide 1 to 40% by weight of peptide 2.

- 10 Preferably the peptide or the peptide combination is suitable for intravenous administration or suitable for oral administration or suitable for administration by inhalation.

Administration forms include, for example, pills, tablets, film tablets, coated tablets, capsules, liposomal formulations, micro- and nano-formulations, powders and deposits. Furthermore, the present invention also includes pharmaceutical preparations for parenteral application, including dermal, intradermal, intragastral, intracutan, intravasal, intravenous, intramuscular, intraperitoneal, intranasal, intravaginal, intrabuccal, percutan, rectal, subcutaneous, sublingual, topical, or transdermal application, which preparations in addition to typical vehicles and/or diluents contain the peptide or the peptide combination according to the present invention.

25 The present invention also includes mammalian milk, artificial mammalian milk as well as mammalian milk substitutes as a formulation for oral administration of the peptide combination to newborns, toddlers, and infants, either as pharmaceutical preparations, and/or as dietary food supplements.

30 The peptide or the peptide combination of the invention can also be administered in form of its pharmaceutically active salts. Suitable pharmaceutically active salts comprise acid addition salts and alkali or earth alkali salts. For instance, sodium, potassium, lithium, magnesium or calcium salts can be obtained.

35 The peptide or the peptide combination of the invention forms pharmaceutically acceptable salts with organic and inorganic acids. Examples of suitable acids for such acid addition salt formation are hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, acetic acid, citric acid, oxalic acid, malonic acid, salicylic acid, p-aminosalicylic acid, malic acid, fumaric acid, succinic acid, ascorbic acid, maleic acid, sulfonic acid, phosphonic acid, perchloric acid, nitric acid, formic acid, propionic

acid, gluconic acid, lactic acid, tartaric acid, hydroxymaleic acid, pyruvic acid, phenylacetic acid, benzoic acid, p-aminobenzoic acid, p-hydroxybenzoic acid, methanesulfonic acid, ethanesulfonic acid, nitrous acid, hydroxyethanesulfonic acid, ethylenesulfonic acid, p-toluenesulfonic acid, naphthylsulfonic acid, sulfanilic acid, camphersulfonic acid, china acid, mandelic acid, o-methylmandelic acid, hydrogen-benzenesulfonic acid, picric acid, adipic acid, D-o-tolytartaric acid, tartronic acid, α -toluic acid, (o, m, p)-toluic acid, naphthylamine sulfonic acid, and other mineral or carboxylic acids well known to those skilled in the art. The salts are prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt in the conventional manner.

The pharmaceutical compositions according to the present invention will typically be administered together with suitable carrier materials selected with respect to the intended form of administration, i.e. for oral administration in the form of tablets, capsules (either solid filled, semi-solid filled or liquid filled), powders for constitution, aerosol preparations consistent with conventional pharmaceutical practices. Other suitable formulations are gels, elixirs, dispersible granules, syrups, suspensions, creams, lotions, solutions, emulsions, suspensions, dispersions, and the like. Suitable dosage forms for sustained release include tablets having layers of varying disintegration rates or controlled release polymeric matrices impregnated with the active components and shaped in tablet form or capsules containing such impregnated or encapsulated porous polymeric matrices. The pharmaceutical compositions may be comprised of 5 to 95% by weight of the peptide or the peptide combination, while also up to 100% of the pharmaceutical composition can consist of the peptide combination.

As pharmaceutically acceptable carrier, excipient and/or diluents can be used lactose, starch, sucrose, cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, talc, mannitol, ethyl alcohol (liquid filled capsules).

Suitable binders include starch, gelatin, natural sugars, corn sweeteners, natural and synthetic gums such as acacia, sodium alginate, carboxymethyl-cellulose, polyethylene glycol and waxes. Among the lubricants that may be mentioned for use in these dosage forms, boric acid, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrants include starch, methylcellulose, guar gum and the like. Sweetening and flavoring agents and preservatives may also be included where appropriate. Some of the terms noted above, namely disintegrants, diluents, lubricants, binders and the like, are discussed in more detail below.

Additionally, the compositions of the present invention may be formulated in sustained release form to provide the rate controlled release of any one or more of the components or active ingredients to optimize the therapeutic effects. Suitable dosage forms for sustained release include layered tablets containing layers of varying disintegration rates or controlled release polymeric matrices impregnated with the active components and shaped in tablet form or capsules containing such impregnated or encapsulated porous polymeric matrices.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier such as inert compressed gas, e.g. nitrogen.

For preparing suppositories, a low melting wax such as a mixture of fatty acid glycerides such as cocoa butter is first melted, and the active ingredient is dispersed homogeneously therein by stirring or similar mixing. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool and thereby solidify.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The peptide or the peptide combination of the present invention may also be deliverable transdermally. The transdermal compositions may take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

The transdermal formulation of the peptide or the peptide combination of the invention is understood to increase the bioavailability of said peptide into the circulating blood. One problem in the administration of peptide(s) is the loss of bioactivity due to the formation of insolubles in aqueous environments or due to degradation. Therefore stabilization of peptide(s) for maintaining their fluidity and maintaining their biological activity upon administration to the patients in need thereof needs to be achieved.

Prior efforts to provide active agents for medication include incorporating the medication in a polymeric matrix whereby the active ingredient is released into the systemic circulation. Known sustained-release delivery means of active agents are disclosed, for example, in US4235988, US4188373, US4100271, US447471, US4474752, US4474753, or US4478822 relating to polymeric pharmaceutical

vehicles for delivery of pharmaceutically active chemical materials to mucous membranes. The pharmaceutical carriers are aqueous solutions of certain polyoxyethylene-polyoxypropylene condensates. These polymeric pharmaceutical vehicles are described as providing for increased drug absorption by the mucous membrane and prolonged drug action by a factor of two or more. The substituents are block copolymers of polyoxypropylene and polyoxyethylene used for stabilization of drugs such as insulin.

Aqueous solutions of polyoxyethylene-polyoxypropylene block copolymers (poloxamers) are useful as stabilizers for peptide(s). Aside from serving as a stabilizer for the peptide(s), poloxamers provide excellent vehicles for the delivery of the peptide(s), and they are physiologically acceptable. Poloxamers, also known by the trade name Pluronic (e.g. Pluronic F127, Pluronic P85, Pluronic F68) have surfactant properties that make them useful in industrial applications. Among other things, they can be used to increase the water solubility of hydrophobic, oily substances or otherwise increase the miscibility of two substances with different hydrophobicities. For this reason, these polymers are commonly used in industrial applications, cosmetics, and pharmaceuticals. They have also been used as model systems for drug delivery applications. In situ gelation of pharmaceutical compositions based on poloxamer that are biologically triggered are known in the art (e.g. US5256396), describing compositions containing poloxamer 407 and water at specified concentrations.

The term capsule refers to a special container or enclosure made of methyl cellulose, polyvinyl alcohols, or denatured gelatins or starch for holding or containing compositions comprising the active ingredients. Hard shell capsules are typically made of blends of relatively high gel strength bone and pork skin gelatins. The capsule itself may contain small amounts of dyes, opaquing agents, plasticizers and preservatives.

Tablet means compressed or molded solid dosage form containing the active ingredients with suitable diluents. The tablet can be prepared by compression of mixtures or granulations obtained by wet granulation, dry granulation or by compaction well known to a person skilled in the art.

Oral gels refers to the active ingredients dispersed or solubilized in a hydrophilic semi-solid matrix.

Powders for constitution refer to powder blends containing the active ingredients and suitable diluents which can be suspended in water or juices. One example for such an oral administration form for newborns, toddlers and/or infants is a human breast milk substitute which is produced from milk powder and milk whey powder, optionally and partially substituted with lactose.

Human breast milk is a complex fluid, rich in nutrients and in non-nutritional bioactive components. It contains all of the nutrients needed by the newborn baby. These include the metabolic components (fat, protein, and carbohydrates), water, and the raw materials for tissue growth and development, such as fatty acids, amino acids, minerals, vitamins, and trace elements.

More than 98% of the fat in is in the form of triglycerides. Oleic acid and palmitic acid are the most abundant fatty acids in breastmilk triglycerides, with comparatively high proportions of the essential fatty acids, and linolenic acid, followed by long-chain polyunsaturated fatty acids, such as arachidonic acid and docosahexaenoic acid. These long-chain fatty acids are constituents of brain and neural tissue and are needed in early life for mental and visual development. The lipid component of breast milk is the transport vehicle for fat-soluble micronutrients such as prostaglandins and vitamins A, D, E, and K.

Proteins account for approximately 75 % of the nitrogen-containing compounds in breast milk. Non-protein nitrogen substances include urea, nucleotides, peptides, free amino acids, and DNA. The proteins of breast milk can be divided into two categories: micellar caseins and aqueous whey proteins, present in the ratio of about 40:60. Casein forms micelles of relatively small volume and produces a soft, flocculent curd in the infant's stomach. The major whey proteins are lactalbumin, lactoferrin, secretory IgA, and serum albumin, with a large number of other proteins and peptides present in smaller amounts.

The principal carbohydrate is lactose, a disaccharide produced in the mammary epithelial cell from glucose by a reaction involving lactalbumin.

In addition to the nutritional components, breast milk contains a wealth of bioactive components that have beneficial non-nutritional functions. These include a wide range of specific and non-specific antimicrobial factors; cytokines and anti-inflammatory substances; and hormones, growth modulators, and digestive enzymes (Table 1), many of which have multiple activities. These components may be of particular importance for young infants because of the immaturity of the host defense and digestive systems early in life.

TABLE 10: Examples of the non-nutritional components of breast milk

Antimicrobial factors	Growth factors
secretory IgA, IgM, IgG lactoferrin lysozyme complement C3 leucocytes bifidus factor lipids and fatty acids antiviral mucins, GAGs oligosaccharides	epidermal (EGF) nerve (NGF) insulin-like (IGF) transforming (TGF) taurine polyamines
Cytokines and anti-inflammatory factors	Digestive enzymes
tumor necrosis factor interleukins interferons prostaglandins antichymotrypsin antitrypsin platelet-activating factor	amylase bile acid-stimulating esterase bile acid-stimulating lipases lipoprotein lipase
Hormones	Transporters
feedback inhibitor of lactation (FIL) insulin prolactin thyroid hormones corticosteroids ACTH oxytocin calcitonin parathyroid hormone erythropoietin	lactoferrin (Fe) folate binder cobalamin binder IgF binder thyroxine binder corticosteroid binder

Besides breast milk, infant formula is the only other infant milk which the medical community considers nutritionally acceptable for infants under the age of one year. Cow's milk is not recommended because of its high protein and electrolyte (salt) content which may harm infant's immature kidneys. The nutrient content of infant

formula should comprise: Protein, Fat, Linoleic acid, Vitamins: A, C, D, E, K, thiamin (B1), riboflavin (B2), B6, B12, Niacin, Folic acid, Pantothenic acid, Calcium, Metals: magnesium, iron, zinc, manganese, copper; Phosphorus, Iodine, Sodium chloride, Potassium chloride. In addition, formulas not made with cow's milk must include
5 biotin, choline, and inositol. Hypoallergenic formulas reduce the likelihood of certain medical complications in babies with specific health problems. Baby formula can be synthesized from raw amino acids. This kind of formula is sometimes referred to as elemental infant formula or as medical food because of its specialized nature.

Powder blends containing the active ingredients and suitable diluents which can be
10 suspended in water or juices can be produced by spray drying.

Spray drying has been found the most suitable process for removing the last part of the water, since spray drying can convert milk concentrate into a powder while still keeping the valuable properties of the milk. The principle of all spray dryers is to
15 transform the concentrate into many small droplets which are then exposed to a fast current of hot air. Because of the very large surface area of the droplets, the water evaporates almost instantaneously and the droplets are transformed into powder particles.

20 Powdered milk is a powder made from dried milk solids. Powdered milk has a far longer shelf life than liquid milk and does not need to be refrigerated due to its low moisture content.

Instant milk powder is produced by partially rehydrating the dried milk powder
25 particles causing them to become sticky and agglomerate. The water is then removed by drying resulting in an increased amount of air incorporated between the powder particles.

Milk powder manufacture is a process carried out on a large scale. It involves the
30 gentle removal of water, while retaining all the desirable natural properties of the milk like colour, flavour, solubility, nutritional value.

Milk powder process includes spray drying, fluid bed processing, extraction, evaporation and freeze drying. Other processes are freeze concentration, filtration, and homogenisation.

35 The artificial mother milk formulations or mother milk substitutes of the present invention are preferably prepared by adding to a mother milk formulation including commercially available mother milk formulations especially in powder form the peptide or inventive peptide combination. The peptide or peptide combination is

preferably added in an amount of 3 – 100 µg peptide or peptide combination per 100 ml (commercially available) mother milk formulation, more preferably in an amount of 5 – 70 µg / 100 ml and most preferably in an amount of 10 – 40 µg / 100 ml mother milk formulation.

5

Suitable diluents are substances that usually make up the major portion of the composition or dosage form. Suitable diluents include sugars such as lactose, sucrose, mannitol and sorbitol, starches derived from wheat, corn rice and potato, and celluloses such as microcrystalline cellulose. The amount of diluents in the composition can range from about 5 to about 95% by weight of the total composition, preferably from about 25 to about 75%, more preferably from about 30 to about 60% by weight, and most preferably from about 40 to 50% by weight.

10

15

The term disintegrants refers to materials added to the composition to help it break apart (disintegrate) and release the medicaments. Suitable disintegrants include starches, "cold water soluble" modified starches such as sodium carboxymethyl starch, natural and synthetic gums such as locust bean, karaya, guar, tragacanth and agar, cellulose derivatives such as methylcellulose and sodium carboxymethylcellulose, microcrystalline celluloses and cross-linked microcrystalline celluloses such as sodium croscarmellose, alginates such as alginic acid and sodium alginate, clays such as bentonites, and effervescent mixtures. The amount of disintegrant in the composition can range from about 1 to about 40% by weight of the composition, preferably 2 to about 30% by weight of the composition, more preferably from about 3 to 20% by weight of the composition, and most preferably from about 5 to about 10% by weight.

20

25

Binders characterize substances that bind or "glue" powders together and make them cohesive by forming granules, thus serving as the "adhesive" in the formulation. Binders add cohesive strength already available in the diluents or bulking agent.

30

Suitable binders include sugars such as sucrose, starches derived from wheat, corn rice and potato; natural gums such as acacia, gelatin and tragacanth; derivatives of seaweed such as alginic acid, sodium alginate and ammonium calcium alginate; cellulosic materials such as methylcellulose and sodium carboxymethylcellulose and hydroxypropyl-methylcellulose; polyvinylpyrrolidone; and inorganics such as magnesium aluminum silicate. The amount of binder in the composition can range from about 1 to 30% by weight of the composition, preferably from about 2 to about 20% by weight of the composition, more preferably from about 3 to about 10% by weight, even more preferably from about 3 to about 6% by weight.

35

Lubricant refers to a substance added to the dosage form to enable the tablet, granules, etc. after it has been compressed, to release from the mold or die by reducing friction or wear. Suitable lubricants include metallic stearates such as magnesium stearate, calcium stearate or potassium stearate; stearic acid; high melting point waxes; and water soluble lubricants such as sodium chloride, sodium benzoate, sodium acetate, sodium oleate, polyethylene glycols and d'l-leucine. Lubricants are usually added at the very last step before compression, since they must be present on the surfaces of the granules and in between them and the parts of the tablet press. The amount of lubricant in the composition can range from about 0.05 to about 15% by weight of the composition, preferably 0.2 to about 5% by weight of the composition, more preferably from about 0.3 to about 3%, and most preferably from about 0.3 to about 1.5% by weight of the composition.

Glidants are materials that prevent caking and improve the flow characteristics of granulations, so that flow is smooth and uniform. Suitable glidants include silicon dioxide and talc. The amount of glident in the composition can range from about 0.01 to 10% by weight of the composition, preferably 0.1% to about 7% by weight of the total composition, more preferably from about 0.2 to 5% by weight, and most preferably from about 0.5 to about 2% by weight.

Coloring agents are excipients that provide coloration to the composition or the dosage form. Such excipients can include food grade dyes and food grade dyes adsorbed onto a suitable adsorbent such as clay or aluminum oxide. The amount of the coloring agent can vary from about 0.01 to 10% by weight of the composition, preferably from about 0.05 to 6% by weight, more preferably from about 0.1 to about 4% by weight of the composition, and most preferably from about 0.1 to about 1%.

Peptide(s) of the invention can be used to form multiparticulates, discrete particles, well known dosage forms, whose totality represents the intended therapeutically useful dose of a drug. When taken orally, multiparticulates generally disperse freely in the gastrointestinal tract, and maximize absorption. A specific example is described in US 6068859, disclosing multiparticulates that provide controlled release of azithromycin. Another advantage of the multiparticulates is the improved stability of the drug. The poloxamer component of the multiparticulate is very inert, thus minimizing degradation of the drug.

However, formulation problems result from the melt-congeal process often used to form multiparticulates. The multiparticulates are preferably formed into round beads or spheres. Some carriers, when melted and then solidified, do not form round beads but may solidify into rods, strings, or other non-spherical shapes. The result is very

irregularly shaped multiparticulates that are difficult to process into dosage forms. This problem is solved by e.g. WO 2007104173 where the particles consist of a poloxamer, a resin, and/or a tocopherol, creating together with the medicament (e.g. insulin) micelles. Micelle formation is essential for the absorption of many nutrients within the human body. Bile salts formed in the liver and secreted by the gall bladder allow micelles of fatty acids to form. This allows the absorption of complicated lipids and lipid soluble vitamins within the micelle by the small intestine. Micelles are approximately spherical in shape. Preferably, the peptide or the peptide combination of the invention are formulated with a poloxamer and a resin to form micelles suitable for oral administration to patients in need of the medicament.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injections or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

Other preferred pharmaceutical compositions are buffered solutions. The term buffer, buffer system, buffer solution and buffered solution, when used with reference to hydrogen-ion concentration or pH, refers to the ability of a system, particularly an aqueous solution, to resist a change of pH on adding acid or alkali, or on dilution with a solvent. Preferred buffer systems can be selected from the group consisting of formate (pKa=3.75), lactate (pKa=3.86), benzoic acid (pKa=4.2) oxalate (pKa=4.29), fumarate (pKa=4.38), aniline (pKa=4.63), acetate buffer (pKa=4.76), citrate buffer (pKa2=4.76, pKa3=6.4), glutamate buffer (pKa=4.3), phosphate buffer (pKa=7.20), succinate (pKa1=4.93; pKa2=5.62), pyridine (pKa=5.23), phthalate (pKa=5.41); histidine (pKa=6.04), MES (2-(N-morpholino)ethanesulphonic acid; pKa=6.15); maleic acid (pKa=6.26); cacodylate (dimethylarsinate, pKa=6.27), carbonic acid (pKa=6.35), ADA (N-(2-acetamido)imino-diacetic acid (pKa=6.62); PIPES (4-piperazinebis-(ethanesulfonic acid; BIS-TRIS-propane (1,3-bis[tris(hydroxymethyl)methylamino]-propane), pKa=6.80), ethylenediamine (pKa=6.85), ACES 2-[(2-amino-2-oxoethyl)amino]ethanesulphonic acid; pKa=6.9), imidazole (pKa=6.95), MOPS (3-(N-morphin)-propane-sulfonic acid; pKa=7.20), diethylmalonic acid (pKa=7.2), TES (2-[tris(hydroxymethyl)methyl]aminoethanesulphonic acid; pKa=7.50) and HEPES (N-2-hydroxyethylpiperazin-N'-2-ethanesulfonic acid; pKa=7.55) buffers or other buffers having a pKa between 3.8 to 7.7.

Preferred is the group of carboxylic acid buffers such as acetate and carboxylic diacid buffers such as fumarate, tartrate and phthalate and carboxylic triacid buffers such as citrate. Another group of preferred buffers is represented by inorganic buffers such as sulfate, borate, carbonate, oxalate, calcium hydroxyde and phosphate buffers. Another group of preferred buffers are nitrogen containing buffers such as imidazole, diethylenediamine, and piperazine.

Also preferred are sulfonic acid buffers such as TES, HEPES, ACES, PIPES, [(2-hydroxy-1,1-bis(hydroxymethyl)ethyl)amino]-1-propanesulfonic acid (TAPS), 4-(2-hydroxyethyl)piperazine-1-propanesulfonic acid (EPPS), 4-Morpholinepropanesulfonic acid (MOPS) and N,N-bis(2-hydroxyethyl)-2-aminoethanesulfonic acid (BES).

Another group of preferred buffers are glycine buffers such as glycine, glycyl-glycine, glycyl-glycyl-glycine, N,N-bis(2-hydroxyethyl)glycine and N-[2-hydroxy-1,1-bis(hydroxy-methyl)ethyl]glycine (Tricine).

Preferred are also amino acid buffers such as glycine, alanine, valine, leucine, isoleucine, serine, threonine, phenylalanine, tyrosine, tryptophane, lysine, arginine, histidine, aspartate, glutamate, asparagine, glutamine, cysteine, methionine, proline, 4-hydroxyproline, N,N,N-trimethyllysine, 3-methylhistidine, 5-hydroxylysine, O-phosphoserine, γ -carboxyglutamate, ϵ -N-acetyllysine, ω -N-methylarginine, citrulline, ornithine and derivatives thereof.

Table 11: Also preferred are the following buffers:

effective pH range	pKa 25°C	buffer
2.7-4.2	3.40	malate (pK1)
3.0-4.5	3.75	formate
3.0-6.2	4.76	citrate (pK2)
3.2-5.2	4.21	succinate (pK1)
3.6-5.6	4.76	acetate
3.8-5.6	4.87	propionate
4.0-6.0	5.13	malate (pK2)
4.9-5.9	5.23	pyridine
5.0-6.0	5.33	piperazine (pK1)
5.0-7.4	6.27	cacodylate
5.5-6.5	5.64	succinate (pK2)

5.5-6.7	6.10	MES
5.5-7.2	6.40	citrate (pK3)
5.5-7.2	6.24	maleate (pK2)
5.5-7.4	1.70, 6.04, 9.09	histidine
5.8-7.2	6.46	bis-tris
5.8-8.0	7.20	phosphate (pK2)
6.0-12.0	9.50	ethanolamine
6.0-7.2	6.59	ADA
6.0-8.0	6.35	carbonate (pK1)
6.1-7.5	6.78	ACES
6.1-7.5	6.76	PIPES
6.2-7.6	6.87	MOPSO
6.2-7.8	6.95	imidazole
6.3-9.5	6.80, 9.00	BIS-TRIS propane
6.4-7.8	7.09	BES
6.5-7.9	7.14	MOPS
6.8-8.2	7.48	HEPES
6.8-8.2	7.40	TES
6.9-8.3	7.60	MOBS
7.0-8.2	7.52	DIPSO
7.0-8.2	7.61	TAPSO
7.0-8.3	7.76	triethanolamine (TEA)
7.0-9.0	0.91, 2.10, 6.70, 9.32	pyrophosphate
7.1-8.5	7.85	HEPPSO
7.2-8.5	7.78	POPSO

Preferred are the buffers having an effective pH range of from 2.7 to 8.5, and more preferred of from 3.8 to 7.7. The effective pH range for each buffer can be defined as $pK_a - 1$ to $pK_a + 1$, where K_a is the ionization constant for the weak acid in the buffer and $pK_a = -\log K$.

Most preferred are buffers suitable for pharmaceutical use e.g. buffers suitable for administration to a patient such as acetate, carbonate, citrate, fumarate, glutamate, lactate, phosphate, phthalate, and succinate buffers. Particularly preferred examples of commonly used pharmaceutical buffers are acetate buffer, citrate buffer, glutamate buffer and phosphate buffer. Also most preferred is the group of carboxylic acid

buffers. The term "carboxylic acid buffers" as used herein shall refer to carboxylic mono acid buffers and carboxylic diacid buffers as well as carboxylic triacid buffers. Of course also combinations of buffers, especially of the buffers mentioned herein are useful for the present invention.

5

Some suitable pharmaceutical buffers are a citrate buffer (preferably at a final formulation concentration of from about 20 to 200 mM, more preferably at a final concentration of from about 30 to 120 mM) or an acetate buffer (preferably at a final formulation concentration of about 20 to 200 mM) or a phosphate buffer (preferably

10

Techniques for the formulation and administration of the peptide or the peptide combination of the present invention may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton PA. A suitable composition comprising at least one peptide mentioned herein may be a solution of the peptide or the peptide combination in a suitable liquid pharmaceutical carrier or any other formulation such as tablets, pills, film tablets, coated tablets, dragees, capsules, powders and deposits, gels, syrups, slurries, suspensions, emulsions, and the like.

15

A particularly preferred pharmaceutical composition is a lyophilised (freeze-dried) preparation (lyophilisate) suitable for administration by inhalation or for intravenous administration. To prepare the preferred lyophilised preparation the peptide or the peptide combination of the invention are solubilised in a 4 to 5% (w/v) mannitol solution and the solution is then lyophilised. The mannitol solution can also be prepared in a suitable buffer solution as described above.

25

Further examples of suitable cryo- / lyoprotectants (otherwise referred to as bulking agents or stabilizers) include thiol-free albumin, immunoglobulins, polyalkyleneoxides (e.g. PEG, polypropylene glycols), trehalose, glucose, sucrose, sorbitol, dextran, maltose, raffinose, stachyose and other saccharides (cf. for instance WO 97/29782), while mannitol is used preferably. These can be used in conventional amounts in conventional lyophilization techniques. Methods of lyophilisation are well known in the art of preparing pharmaceutical formulations.

30

For administration by inhalation the particle diameter of the lyophilised preparation is preferably between 2 to 5 μm , more preferably between 3 to 4 μm . The lyophilised preparation is particularly suitable for administration using an inhalator, for example the OPTINEB[®] or VENTA-NEB[®] inhalator (NEBU-TEC, Elsenfeld, Germany). The

35

lyophilised product can be rehydrated in sterile distilled water or any other suitable liquid for inhalation administration.

Alternatively for intravenous administration the lyophilised product can be rehydrated
5 in sterile distilled water or any other suitable liquid for intravenous administration.

After rehydration for administration in sterile distilled water or another suitable liquid the lyophilised preparation should have the approximate physiological osmolality of the target tissue for the rehydrated peptide preparation i.e. blood for intravenous
10 administration or lung tissue for inhalation administration. Thus it is preferred that the rehydrated formulation is substantially isotonic.

The preferred dosage concentration for either intravenous, oral, or inhalation administration is between 100 to 2000 $\mu\text{mole/ml}$, and more preferably is between 200
15 to 800 $\mu\text{mole/ml}$. These are also the preferred ranges of the peptide combination in the mother milk substitute or artificial mother milk formulation or the pharmaceutical compositions disclosed herein.

Dietary supplement

20 Still another aspect of the present invention relates to the use of disclosed peptide and peptide combination as a dietary supplement. That dietary supplement is preferably for oral administration and especially but not limited to administration to newborns, toddlers, and/or infants. A dietary supplement is intended to supplement the diet. The "dietary ingredients" in these products may in addition include: vitamins,
25 minerals, herbs or other botanicals, amino acids, and substances such as enzymes, organ tissues, glandulars, and metabolites. Dietary supplements may be manufactured in forms such as tablets, capsules, softgels, gelcaps, liquids, or powders.

Method of treatment

30 Another aspect of the present invention relates to a method of prophylaxis and/or treatment of cancer, an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, an infectious disease, a lung disease, a heart and vascular disease or a metabolic disease or any other disease disclosed herein
35 comprising administering to a patient in need thereof a pharmaceutical composition comprising the peptide or the peptide combination according to the present invention in a therapeutically effective amount effective to treat the afore-mentioned disease.

Accordingly, the terms "prophylaxis" or "treatment" includes the administration of the peptide or the peptide combination of the present invention to prevent, inhibit, or arrest the symptoms of an infectious disease, an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, or a heart and vascular disease. In some instances, treatment with the peptide or the peptide combination of the present invention will be done in combination with other protective compounds to prevent, inhibit, or arrest the symptoms of an infectious disease, an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, or a heart and vascular disease.

The term "active agent" or "therapeutic agent" as used herein refers to an agent that can prevent, inhibit, or arrest the symptoms and/or progression of an infectious, an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, a heart and vascular disease or any other disease disclosed herein.

The term "therapeutic effect" as used herein, refers to the effective provision of protection effects to prevent, inhibit, or arrest the symptoms and/or progression of an infectious, an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, or a heart and vascular disease.

The term "a therapeutically effective amount" as used herein means a sufficient amount of the peptide or the peptide combination of the invention to produce a therapeutic effect, as defined above, in a subject or patient in need of treatment.

The terms "subject" or "patient" are used herein mean any mammal, including but not limited to human beings, including a human patient or subject to which the compositions of the invention can be administered. The term mammals include human patients and non-human primates, as well as experimental animals such as rabbits, rats, and mice, and other animals.

The peptide or the peptide combination of the present invention can be used for the prophylaxis and/or treatment of cancer, an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, an infectious disease, a lung disease, a heart and vascular disease or a metabolic disease or any other disease mentioned herein in combination administration with another therapeutic compound. As used herein the term "combination administration" of a compound, therapeutic agent or known drug with the peptide or the peptide combination of the present invention means administration of the drug and the peptide or the peptide

combination at such time that both the known drug and the peptide or the peptide combination will have a therapeutic effect. In some cases this therapeutic effect will be synergistic. Such concomitant administration can involve concurrent (i.e. at the same time), prior, or subsequent administration of the drug with respect to the administration of the peptide or the peptide combination of the present invention. A person of ordinary skill in the art would have no difficulty determining the appropriate timing, sequence and dosages of administration for particular drugs and peptide(s) of the present invention.

Definition of peptide activity

A peptide or peptide combination is deemed to have therapeutic activity if it demonstrated any one of the following activities listed in a) to g).

a) The peptide could inhibit the activity of an over active biological pathway.

b) The peptide could inhibit the production of an over produced biological molecule.

c) The peptide could inhibit the activity of an over produced biological molecule.

d) The peptide could increase the activity of an under active biological pathway.

e) The peptide could increase the production of an under produced biological molecule.

f) The peptide could mimic the activity of an under produced biological molecule.

g) The peptide could prevent, inhibit, or arrest the symptoms and/or progression of cancer, an infectious disease, an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, or a heart and vascular disease or any other disease disclosed herein.

As used herein "inhibition" is defined as a reduction of the activity or production of a biological pathway or molecule activity of between 10 to 100%. More preferably the reduction of the activity or production of a biological pathway or molecule activity is between 25 to 100%. Even more preferably the reduction of the activity or production of a biological pathway or molecule activity is between 50 to 100%.

As used herein "increase" is defined as an increase of the activity or production of a biological pathway or molecule of between 10 to 100%. More preferably the increase

of the activity or production of a biological pathway or molecule activity is between 25 to 100%. Even more preferably the increase of the activity or production of a biological pathway or molecule activity is between 50 to 100%.

- 5 As used herein "mimic" is defined as an increase in the activity of a biological pathway dependent on the under produced biological molecule of between 10 to 100%. More preferably the increase of the activity of the biological pathway is between 25 to 100%. Even more preferably the increase of the activity of the biological pathway is between 50 to 100%.

10

Peptides and peptide combination

The following peptides were tested alone and in combination for their activity as a therapeutic agent for the prophylaxis and/or treatment of cancer, an infectious disease, an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, or a heart and vascular disease.

15

Peptide 1: having the amino acid sequence:

Gly-Arg-Gly-Asp-Asn-Pro-OH

- 20 Peptide 2: Parathyroid Hormone (pTH)(1-34) having the amino acid sequence:
Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe

(from the Calcium metabolism peptide family also comprising:

- 25 Parathyroid Hormone (PTH)(1-38) having the amino acid sequence:
Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe-Val-Ala-Leu-Gly,

Parathyroid Hormone (PTH)(1-44) having the amino acid sequence:

- 30 Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe-Val-Ala-Leu-Gly-Ala-Pro-Leu-Ala-Pro-Arg,

Parathyroid Hormone (PTH)(13-34) having the amino acid sequence:

- 35 Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe,

Parathyroid Hormone (PTH)(39-68) having the amino acid sequence:

Ala-Pro-Leu-Ala-Pro-Arg-Asp-Ala-Gly-Ser-Gln-Arg-Pro-Arg-Lys-Lys-Glu-Asp-Asn-Val-Leu-Val-Glu-Ser-His-Glu-Lys-Ser-Leu-Gly,

Parathyroid Hormone (PTH)(44-68) having the amino acid sequence:

- 5 Arg-Asp-Ala-Gly-Ser-Gln-Arg-Pro-Arg-Lys-Lys-Glu-Asp-Asn-Val-Leu-Val-Glu-Ser-His-Glu-Lys-Ser-Leu-Gly,

Parathyroid Hormone (PTH)(1-34)-Amide [Nle⁸ 18 Tyr³⁴] having the amino acid sequence:

- 10 Ser-Val-Ser-Glu-Ile-Gln-Leu-Nle-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Nle-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Tyr-NH₂,

pTH-Related Protein having the amino acid sequence:

- Ala-Val-Ser-Glu-His-Gln-Leu-Leu-His-Asp-Lys-Gly-Lys-Ser-Ile-Gln-Asp-Leu-Arg-Arg-15 Arg-Phe-Phe-Leu-His-His-Leu-Ile-Ala-Glu-Ile-His-Thr-Ala,

pTH-Related Protein (1-16) having the amino acid sequence:

Ala-Val-Ser-Glu-His-Gln-Leu-Leu-His-Asp-Lys-Gly-Lys-Ser-Ile-Gln,

- 20 Teriparatide having the amino acid sequence:

Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe,

Parathyrin having the amino acid sequence:

- 25 Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe-Val-Ala-Leu,

pTH (18-48) having the amino acid sequence:

- Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe-Val-Ala-30 Leu-Gly-Ala-Pro-Leu-Ala-Pro-Arg-Asp-Ala-Gly-Ser,

Calcitonin having the amino acid sequence:

Cys-Gly-Asn-Leu-Ser-Thr-Cys-Met-Leu-Gly-Thr-Tyr-Thr-Gln-Asp-Phe-Asn-Lys-Phe-His-Thr-Phe-Pro-Gln-Thr-Ala-Ile-Gly-Val-Gly-Ala-Pro-NH₂,

- 35 Calcitonin Gene Related Peptide (CGRP) having the amino acid sequence:

Ala-Cys-Asp-Thr-Ala-Thr-Cys-Val-Thr-His-Arg-Leu-Ala-Gly-Leu-Leu-Ser-Arg-Ser-Gly-Gly-Val-Val-Lys-Asn-Asn-Phe-Val-Pro-Thr-Asn-Val-Gly-Ser-Lys-Ala-Phe-NH₂,

Calcitonin Gene Related Peptide (CGRP)[Tyr⁰] having the amino acid sequence:

Tyr-Ala-Cys-Asp-Thr-Ala-Thr-Cys-Val-Thr-His-Arg-Leu-Ala-Gly-Leu-Leu-Ser-Arg-Ser-Gly-Gly-Val-Val-Lys-Asn-Asn-Phe-Val-Pro-Thr- Asn-Val-Gly-Ser-Lys-Ala-Phe-NH₂,

Calcitonin Gene Related Peptide (CGRP)(8-37) having the amino acid sequence:

- 5 Val-Thr-His-Arg-Leu-Ala-Gly-Leu-Leu-Ser-Arg-Ser-Gly-Gly-Val-Val-Lys-Asn-Asn-Phe-Val-Pro-Thr-Asn-Val-Gly-Ser-Lys-Ala-Phe-NH₂,

Calcitonin Gene Related Peptide (CGRP)(17-37) having the amino acid sequence:

- 10 Ser-Arg-Ser-Gly-Gly-Val-Val-Lys-Asn-Asn-Phe-Val-Pro-Thr-Asn-Val-Gly-Ser-Lys-Ala-Phe-NH₂,

Calcitonin Gene Related Peptide (CGRP) Alpha [Tyr27 Ala31 Pro34 Phe35](27-37) having the amino acid sequence:

- 15 Tyr-Val-Pro-Thr-Ala-Val-Gly-Pro-Phe-Ala-Phe-NH₂,

Calcitonin Gene Related Peptide II (CGRP II) having the amino acid sequence:

Ala-Cys-Asn-Thr-Ala-Thr-Cys-Val-Thr-His-Arg-Leu-Ala-Gly-Leu-Leu-Ser-Arg-Ser-Gly-Gly-Met-Val-Lys-Ser-Asn-Phe-Val-Pro-Thr- Asn-Val-Gly-Ser-Lys-Ala-Phe-NH₂,

- 20 Intermedin-53 having the amino acid sequence:

His-Ser-Gly-Pro-Arg-Arg-Thr-Gln-Ala-Gln-Leu-Leu-Arg-Val-Gly-Cys-Val-Leu-Gly-Thr-Cys-Gln-Val-Gln-Asn-Leu-Ser-His-Arg-Leu-Trp-Gln-Leu-Met-Gly-Pro-Ala-Gly-Arg-Gln-Asp-Ser-Ala-Pro-Val-Asp-Pro-Ser-Ser-Pro-His-Ser-Tyr-NH₂,

- 25 Intermedin having the amino acid sequence:

Thr-Gln-Ala-Gln-Leu-Leu-Arg-Val-Gly-Cys-Val-Leu-Gly-Thr-Cys-Gln-Val-Gln-Asn-Leu-Ser-His-Arg-Leu-Trp-Gln-Leu-Met-Gly-Pro-Ala-Gly-Arg-Gln-Asp-Ser-Ala-Pro-Val-Asp-Pro-Ser-Ser-Pro-His-Ser-Tyr-NH₂).

- 30 Both peptides are preferably contained in the inventive combination in a molar ratio of 1 mole peptide 1 to 5 mole peptide 2 to 5 mole peptide 1 to 1 mole peptide 2, more preferred in a molar ratio of 1 mole peptide 1 to 4 mole peptide 2 to 4 mole peptide 1 to 1 mole peptide 2, still more preferred in a molar ratio of 1 mole peptide 1 to 3 mole peptide 2 to 3 mole peptide 1 to 1 mole peptide 2, still more preferred in a molar ratio of 1 mole peptide 1 to 2 mole peptide 2 to 2 mole peptide 1 to 1 mole peptide 2, and
35 of 1 mole peptide 1 to 1.5 mole peptide 2 to 1.5 mole peptide 1 to 1 mole peptide 2. Preferred ratios of the peptides in % by weight are disclosed above which can be used instead of the ratios mentioned as molar rates.

Furthermore the present invention relates to the use of the above-mentioned peptide combination as pharmaceutically active agents in medicine, i.e. as medicament. Advantage of the inventive peptide combination is that the peptides are less toxic in comparison to the commonly used drugs for the certain indications mentioned herein and that the peptide combination has less side effects, can be used for a long term treatment of certain diseases and can be easily administered. Moreover the peptide combination is selective for certain targets and under physiological conditions no toxic or noxious degradation products are formed.

As used herein, the term "peptide(s)" or "peptide combination" shall also refer to salts, deprotected or deacetylated forms, enantiomers, diastereomers, racemates, prodrugs and hydrates of the above-mentioned peptides. Diastereomers of a peptide are obtained when the stereochemical or chiral center of one or more amino acids is changed. The enantiomer has the opposite stereochemistry at all chiral centers.

The term "prodrug" refers to any precursor compound which is able to generate or to release the above-mentioned peptide under physiological conditions. Such prodrugs, i.e. such precursor molecules are for instance larger peptides which are selectively cleaved in order to form one of the above-mentioned peptides. Further prodrugs are protected amino acids having especially protecting groups at the carboxylic acid and/or amino group.

Suitable protecting groups for amino groups are the benzyloxycarbonyl, t-butyloxycarbonyl (BOC), formyl, and acetyl or acyl group. Suitable protecting groups for the carboxylic acid group are esters such as benzyl esters or t-butyl esters.

The present invention also includes the above peptides having amino acid substitutions, deletions, additions, the substitutions and additions including the standard D and L amino acids and modified amino acids such as for example amidated and acetylated amino acids, wherein the therapeutic activity of the base peptide sequence as shown above is maintained.

In the listed peptide sequences "Ac" indicates an acetylated residue and "NH₂" indicates an amidated residue, "cyclo" indicates a cyclic peptide, and "D" indicates a D optical isomer. Deacetyled amino or NH-group refers to the free amino (–HH₂) group.

The following abbreviations are used for the common amino acids referred to herein.

Table 12:

Abbreviation	Amino acid
Ala	Alanine
Arg	Arginine
Asn	Asparagine
Asp	Aspartic acid (Aspartate)
Cys	Cysteine
Gln	Glutamine
Glu	Glutamic acid (Glutamate)
Gly	Glycine
His	Histidine
Ile	Isoleucine
Leu	Leucine
Lys	Lysine
Met	Methionine
Phe	Phenylalanine
Pro	Proline
Pyl	Pyrrolysine
Ser	Serine
Sec	Selenocysteine
Thr	Threonine
Trp	Tryptophan
Tyr	Tyrosine
Val	Valine
Asx	Aspartic acid or Asparagine
Glx	Glutamine or Glutamic acid
Xaa	Any amino acid
Xle	Leucine or Isoleucine

5

Some modified amino acids are indicated as follows:

"D-2-Nal" is 2-naphthyl-D-alanine,

"SertBu" is t-butyl serine,

10 "Azagly" is aza glycine,

"Me" is methyl,

Met(O) is methionine sulfoxide,

"Pyr" and "pGlu" are pyroglutamic acid,

"Tyr(SO₃H)" is sulphated tyrosine,

15 "Tyr(Me)" is methyltyrosine,

"NHEt" is ethylamide.

EXAMPLES

The peptides as listed above and the inventive peptide combination with approximately equimolar amounts of the two peptides (deviation $\pm 10\%$) were tested for activity using the assays described in Examples 1 to 17. The tested peptides are all commercially available and are all known peptides and well described and characterized in the state of the art literature. The inventive peptide combination was prepared by simply mixing the two commercially available peptides in a molar ratio, for instance, between 0.9 to 1.1 and 1.1 to 0.9 (referred to as "approximately equimolar amounts") or other ratios such as from 0.5 – 1.5 to 1.5 – 0.5. It has to be mentioned that the biological data obtained with the peptide combinations in molar ratios between 0.4 to 1.6 mole (peptide 1 to peptide 2) and 1.6 to 0.4 mol (peptide 1 to peptide 2) were in general very similar to the biological data obtained with the equimolar peptide combinations so that in most cases the data for the equimolar (from 0.9 : 1.1 to 1.1 : 0.9) combinations are cited herein. Further data are presented only in these cases where the biological data of the peptide combinations with non equimolar amounts of the peptides differ more than 20% from the biological data of the equimolar combination.

The term "peptides" as used in the following examples refers to peptide 1, peptide 2 and the peptide combination and the concentration of "10 micrograms per ml" refers to 10 μg peptide 1 per ml or 10 μg peptide 2 per ml or 10 μg peptide combination per ml.

Thus the term "peptides" in the following examples indicates that the test disclosed in the corresponding example was conducted with peptide 1 alone and peptide 2 alone and with the peptide combination generally in equimolar ratios (molar ratio about 1 : 1 for peptide 1 : peptide 2) if no other molar ratio is mentioned in the corresponding example.

EXAMPLE 1:

HIV-1 Experiments

CEM-SS cells were passaged in T-75 flasks prior to use in the antiviral assay. On the day preceding the assay, the cells were split 1:2 to assure they were in an exponential growth phase at the time of infection. Total cell viability quantification was performed using a hemacytometer and trypan blue exclusion. Cell viability was greater than 95% for the cells to be utilized in the assay. The cells were resuspended at 5×10^4 cells/ml in tissue culture medium and added to the peptides-containing microtiter plates in a volume of 50 microliters.

The virus used was the lymphocytotropic strain HIV-1_{IIIB}. Virus was obtained from NIH AIDS Research and Reference Reagent Program and was grown in CEM-SS cells for the production of stock virus pools. For each assay, a pre-titered aliquot of virus was removed from the freezer (-80°C) and allowed to thaw slowly to room temperature in a biological safety cabinet. The virus was resuspended and diluted into tissue culture medium such that the amount of virus added to each well in a volume of 50 microliters was the amount determined to give between 85% to 95% cell killing after 6 days post-infection. TCID₅₀ calculations by endpoint titration in CEM-SS cells indicated that the multiplicity of infection was approximately 0.01. AZT (nucleoside reverse transcriptase inhibitor; NRTI) and indinavir (protease inhibitor; PI) were used as positive control antiviral compounds.

Plate format

Each plate contained cell control wells (cells only), virus control wells (cells plus virus), drug cytotoxicity wells (cells plus peptides only), peptide colorimetric control wells (peptide only) as well as experimental wells (peptides - 10 micrograms per ml - plus cells plus virus). Samples were evaluated for antiviral efficacy with triplicate measurements and with duplicate measurements to determine cellular cytotoxicity, if detectable.

	1	2	3	4	5	6
A	Reagent Background Control					
B	Drug 1 Tox	Cell Control	Drug 1 Antiviral Efficacy			Drug 1 Tox
C	Drug 2 Tox		Drug 2 Antiviral Efficacy			Drug 2 Tox
D	Drug 3 Tox		Drug 3 Antiviral Efficacy			Drug 3 Tox
E	Drug 4 Tox	Virus Control	Drug 4 Antiviral Efficacy			Drug 4 Tox
F	Drug 5 Tox		Drug 5 Antiviral Efficacy			Drug 5 Tox
G	Drug 6 Tox		Drug 6 Antiviral Efficacy			Drug 6 Tox
H	Drug 6 Color	Drug 5 Color	Drug 4 Color	Drug 3 Color	Drug 2 Color	Drug 1 Color

	7	8	9	10	11	12
A	Plastic Control					
B	Drug 7 Tox	Drug 7 Antiviral Efficacy			Cell Control	Drug 7 Tox
C	Drug 8 Tox	Drug 8 Antiviral Efficacy				Drug 8 Tox
D	Drug 9 Tox	Drug 9 Antiviral Efficacy				Drug 9 Tox
E	Drug 10 Tox	Drug 10 Antiviral Efficacy			Virus Control	Drug 10 Tox
F	Drug 11 Tox	Drug 11 Antiviral Efficacy				Drug 11 Tox
G	Drug 12 Tox	Drug 12 Antiviral Efficacy				Drug 12 Tox
H	Drug 12 Color	Drug 11Color	Drug 10 Color	Drug 9 Color	Drug 8 Color	Drug 7 Color

At assay termination, the plates were stained with the soluble tetrazolium-based dye MTS (CellTiter 96 Reagent, Promega) to determine cell viability and quantify peptides' toxicity. MTS is metabolized by the mitochondrial enzymes of metabolically active cells to yield a soluble formazan product, allowing the rapid quantitative analysis of cell viability and peptide cytotoxicity. This reagent is a stable, single solution that does not require preparation before use. At assay termination, 20-25 microliters of MTS reagent was added per well and the microtiter plates were then incubated for 5 hours at 37°C, and 5% CO₂ to assess cell viability. Adhesive plate sealers were used in place of lids, the sealed plates were inverted several times to mix the soluble formazan product and the plate was read spectrophotometrically at 490/560 nm with a Molecular Devices Vmax plate reader.

The overall assay performance was valid based upon judgement of the positive control compounds AZT and indinavir exhibiting the expected levels of antiviral activity. Macroscopic observation of the cells in each well of the microtiter plate confirmed the cytotoxicity results obtained following staining of the cells with the MTS metabolic dye.

Results from HIV experiments: Peptide 1 reduced the cytopathic effect (CPE) in HIV-1 infected CEM-SS cells by 100%. Peptide 2 reduced CPE in HIV-1 infected CEM-SS cells by 18.1%. The peptide combination (0.95 mole peptide 1 and 1.05 mole peptide 2) reduced CPE in HIV-1 infected CEM-SS cells by 60% and the peptide combination (1.50 mole peptide 1 and 0.50 mole peptide 2) reduced CPE in HIV-1 infected CEM-SS cells by 80%. In addition, the peptides of the invention and the peptide combination did not show any significant inhibitory effects on cell viability in these human liver cells.

The selectivity and specificity of the obtained data was examined in the following examples

EXAMPLE 2:

HBV Experimental Assay System

HepG2-2.2.15 is a stable cell line containing the hepatitis B virus (HBV) ayw strain genome (ATCC Cat. No. CRL-11997). Antiviral compounds blocking any late step of viral replication such as transcription, translation, pregenome encapsidation, reverse transcription, particle assembly and release can be identified and characterized using this cell line. In this assay, an active compound will reduce the production of secreted HBV from cells, measured by utilizing real

time quantitative PCR (TaqMan) assay to directly and accurately measure HBV DNA copies. The analysis of this data allows to calculate:

- * Antiviral activity
- * Compound Cytotoxicity

5 HepG2-2.2.15 cells were plated in 96-well microtiter plates. After 16-24 hours the confluent monolayer of HepG2-2.2.15 cells was washed and the medium was replaced with complete medium containing test peptides - 10 micrograms per ml - in duplicate. Lamivudine (3TC) was used as the positive control, while media alone was added to the cells as a negative control (virus control). Three days later
10 the culture medium was replaced with fresh medium containing the peptides. Six days following the initial administration of the peptides, the cell culture supernatants was collected, treated with pronase and DNase and then used in a real-time quantitative TaqMan PCR assay. The PCR-amplified HBV DNA was detected in real-time by monitoring increases in fluorescence signals that result
15 from the exonucleolytic degradation of a quenched fluorescence probe molecule that hybridizes to the amplified HBV DNA. For each PCR amplification, a standard curve was simultaneously generated using dilutions of purified HBV DNA. Antiviral activity was calculated from the reduction in HBV DNA levels (% virus control). A novel dye uptake assay was then employed to measure cell viability,
20 which is used to calculate toxicity (% cell control).
Results from HBV experiments: Peptide 1 inhibited by 36.4% HBV replication as compared to the virus control infection. Peptide 2 inhibited by 0% HBV replication as compared to the virus control infection. The peptide combination (0.95 mole peptide 1 and 1.05 mole peptide 2) inhibited by 18% HBV replication as compared to
25 the virus control infection. In addition, the peptides of the invention did not show any significant inhibitory effects on cell viability in these human liver cells.

EXAMPLE 3:

HCMV experimental assay system

30 MRC-5 cells (human embryonal lung fibroblasts) were obtained from the American Type Culture Collection (ATCC CCL-171; Rockville, Maryland) and grown in Eagle's Minimum Essential Medium with Earle's BSS (EMEM) supplemented with 10% fetal bovine serum (FBS), 0.1 mM non-essential amino acids, 1.0 mM sodium pyruvate, 2.0 mM L-Glutamine, 100 units/ml Pencillin and 100 micrograms/ml Streptomycin. Cells
35 were split twice a week 1:2.

HCMV strain AD169 was obtained from ATCC (ATCC VR-538). Virus stocks were prepared by infecting 80% confluent MRC-5 cells at a minimal multiplicity of infection in MRC-5 growth medium containing 2% FBS. Monolayers were incubated at 37°C, 5% CO₂ until 90%-95% viral cytopathic effect (CPE) was observed (10-13 days). Culture

medium was then collected from the cells, centrifuged at low speed to remove cellular debris, aliquoted in 1ml volumes and stored at -80°C as stock virus.

MRC-5 cells were seeded at 75,000 cells/well in 24 well plates using MRC-5 growth medium. The plates were incubated overnight at 37°C, 5% CO₂. The following day, media was removed and 100 plaque forming units (pfu) of HCMV was added to the wells. Virus was allowed to adsorb onto the cells for 1 hour at 37°C, 5% CO₂. Peptides were diluted - 10 micrograms per ml - in assay medium containing 0.5% Methylcellulose. After the incubation period, 1ml of each peptide solution was added to the wells without aspirating the virus inoculums. The plates were incubated for 7-10 days to allow for plaque formation. Ganciclovir was used as positive control. Cultures were examined microscopically and toxicities were noted. The media was aspirated from the wells and the cells were fixed and stained using 20% methanol containing Crystal Violet followed by enumeration of plaques by microscopic inspection.

For cytotoxicity testing, MRC-5 cells were seeded at 2,500 cells/well in 96 well plates using growth medium. The plates were incubated overnight at 37°C, 5% CO₂. The following day, peptides were added and tested in duplicates. After a 6 days incubation period, cell viability was measured using CellTiter 96 Solution (Promega). Plates were incubated for additional 4 hours at 37°C. Adhesive plate sealers were used in place of lids, the sealed plates were inverted several times to mix the soluble formazan product and the plate was read spectrophotometrically at 490/560 nm with a Molecular Devices Vmax plate reader.

The overall assay performance was valid based upon judgement of the positive control compound Ganciclovir exhibiting the expected levels of antiviral activity.

Macroscopic observation of the cells in each well of the microtiter plate confirmed the cytotoxicity results obtained following staining of the cells with the MTS metabolic dye.

Results from HCMV assay: Peptides 1 and 2 did not inhibit HCMV plaque formation as compared to the virus control experiment. The peptide combination did not provide synergistic effects. In addition, the peptides of the invention did not show any significant inhibitory effects on cell viability in these human lung cells.

EXAMPLE 4:

Methicillin Resistant *Staphylococcus Aureus* (MRSA) assay

The antibacterial assay was conducted using clear, U-bottom 96-well microtiter plates. Cation-adjusted Mueller-Hinton Broth (MHB) was used for testing MRSA. The peptides of the invention (0.1 ml of each - 10 micrograms per ml) were dispensed into wells in duplicate. Then the wells were inoculated with 5 x 10⁵ CFU/mL MRSA in 0.1 ml volume. For control purposes, each plate included 4 wells containing media

without bacterial inoculum and 4 wells containing medium with inoculum but without peptides. The plates were incubated for 12 h at 37 °C, and read visually 18-24 hours post-incubation. Growth control of MRSA was examined first to determine adequacy of media preparations and growth conditions. Acceptable growth is defined as ≥ 2 mm wide button of cells at the bottom of each sample well, or obvious turbidity in the culture supernatant. Test wells were examined and scored as positive/negative for activity. A positive score for activity is based on complete inhibition of macroscopic growth of the test MRSA.

Results from MRSA assay: Peptides of the invention and the peptide combination showed no inhibitory effects on the growth of MRSA.

EXAMPLE 5:

***Pseudomonas aeruginosa* assay**

The antibacterial assay was conducted using clear, U-bottom 96-well microtiter plates. Cation-adjusted Mueller-Hinton Broth (MHB) was used for testing *Pseudomonas aeruginosa*. The peptides of the invention (0.1 ml of each - 10 micrograms per ml) were dispensed into wells in duplicate. Then the wells were inoculated with 5×10^5 CFU/mL *Pseudomonas aeruginosa* in 0.1 ml volume. For control purposes, each plate included 4 wells containing media without bacterial inoculum and 4 wells containing medium with inoculum but without peptides. The plates were incubated for 12 h at 37 °C, and read visually 18-24 hours post-incubation. Growth control of *Pseudomonas aeruginosa* was examined first to determine adequacy of media preparations and growth conditions. Acceptable growth is defined as ≥ 2 mm wide button of cells at the bottom of each sample well, or obvious turbidity in the culture supernatant. Test wells were examined and scored as positive/negative for activity. A positive score for activity is based on complete inhibition of macroscopic growth of the test *Pseudomonas aeruginosa*.

Results from *Pseudomonas aeruginosa* assay: Peptides of the invention and the peptide combination showed no inhibitory effects on the growth of *Pseudomonas aeruginosa*.

EXAMPLE 6:

***Streptococcus pneumoniae* assay**

The antibacterial assay was conducted using clear, U-bottom 96-well microtiter plates. Cation-adjusted Mueller-Hinton Broth (MHB) was used for testing *Streptococcus pneumoniae*. The peptides of the invention (0.1 ml of each - 10 micrograms per ml) were dispensed into wells in duplicate. Then the wells were inoculated with 5×10^5 CFU/mL *Streptococcus pneumoniae* in 0.1 ml volume. For

control purposes, each plate included 4 wells containing media without bacterial inoculum and 4 wells containing medium with inoculum but without peptides. The plates were incubated for 12 h at 37 °C, and read visually 18-24 hours post-incubation. Growth control of *Streptococcus pneumoniae* was examined first to determine adequacy of media preparations and growth conditions. Acceptable growth is defined as ≥ 2 mm wide button of cells at the bottom of each sample well, or obvious turbidity in the culture supernatant. Test wells were examined and scored as positive/negative for activity. A positive score for activity is based on complete inhibition of macroscopic growth of the test *Streptococcus pneumoniae*.

Results from *Streptococcus pneumoniae* assay: Peptide 1 inhibited by 100% the growth of *Streptococcus pneumoniae*. Peptide 2 showed no effect. The peptide combination showed no significant inhibitory effects on the growth of *Streptococcus pneumoniae*.

EXAMPLE 7:

***Mycobacterium tuberculosis* assay**

The antibacterial assay was conducted using clear, U-bottom 96-well microtiter plates. Middlebrook 7H12 assay medium was used for testing drug-resistant *Mycobacterium tuberculosis*. The peptides of the invention (0.1 ml of each - 10 micrograms per ml) were dispensed into wells in duplicate. Then the wells were inoculated with 5×10^5 CFU/mL *Mycobacterium tuberculosis* in 0.1 ml volume. For control purposes, each plate included 4 wells containing media without bacterial inoculum and 4 wells containing medium with inoculum but without peptides. The plates were incubated for seven days at 37 °C, and read visually thereafter. Growth control of *Mycobacterium tuberculosis* was examined first to determine adequacy of media preparations and growth conditions. Acceptable growth is defined as ≥ 2 mm wide button of cells at the bottom of each sample well, or obvious turbidity in the culture supernatant. Test wells were examined and scored as positive/negative for activity. A positive score for activity is based on complete inhibition of macroscopic growth of the test *Mycobacterium tuberculosis*. The drug-resistant *Mycobacterium tuberculosis* that was used in the assay is resistant against following medicaments: para-aminosalicylic acid (PAS), streptomycin and isoniazid (INH).

Results from *Mycobacterium tuberculosis* assay: Peptides of the invention and the peptide combination showed no inhibitory effects on the growth of *Mycobacterium tuberculosis*.

EXAMPLE 8:**Cell cycle assay**

Human A549 cells (carcinomic human alveolar basal epithelial cells) were utilized in the experiments employing the Propidium iodide cell cycle assay. The eukaryotic cell cycle is a series of events that take place in a cell leading to its replication.

The regulation of the cell cycle involves steps crucial to the cell, including detecting and repairing genetic damage, and provision of various checks to prevent uncontrolled cell division. The molecular events that control the cell cycle are ordered and directional; that is, each process occurs in a sequential fashion.

The cell cycle consists of four distinct phases: G₁ phase, S phase, G₂ phase (collectively known as interphase) and M phase. M phase is itself composed of two tightly coupled processes: mitosis, in which the cell's chromosomes are divided between the two daughter cells, and cytokinesis, in which the cell's cytoplasm divides forming distinct cells. Activation of each phase is dependent on the proper progression and completion of the previous one. Cells that have temporarily or reversibly stopped dividing are said to have entered a state of quiescence called G₀ phase. The relatively brief M phase consists of nuclear division and cytoplasmic division. The first phase within interphase, from the end of the previous M phase till the beginning of DNA synthesis is called G₁ (G indicating gap or growth). During this phase the biosynthetic activities of the cell resume at a high rate. This phase is marked by synthesis of various enzymes that are required in S phase, mainly those needed for DNA replication. The ensuing S phase starts when DNA synthesis commences; when it is complete, all of the chromosomes have been replicated. The cell then enters the G₂ phase, which lasts until the cell enters mitosis. Significant protein synthesis occurs during this phase, mainly involving the production of microtubules, which are required during the process of mitosis. Inhibition of protein synthesis during G₂ phase prevents the cell from undergoing mitosis.

Disregulation of the cell cycle components may lead to tumor formation.

Propidium iodide is an intercalating agent and a fluorescent molecule that can be used to stain DNA. Cells were incubated for 24 hours with test peptides - 10 micrograms per ml - or left untreated. After that cells were trypsinized, suspended in medium + 10% FCS, centrifuged (1000 rpm, 5 min), and the cell pellet resuspended in PBS (1 ml). The cells were pipetted into 2.5 ml absolute EtOH (final concentration approx. 70%) and incubated on ice for 15 min. Thereafter, cells were pelleted at 1500 rpm for 5 min and resuspended in Propidium iodide solution in PBS. After incubation for 40 min at 37°C, cells were analyzed in the FACS.

Results from cell cycle assay: Peptides of the invention and the peptide combination showed no inhibitory or irregular effects on the cell cycle of the tested human lung cells.

EXAMPLE 9:**T cell proliferation assay**

Human Peripheral Blood Mononuclear Cells (PBMC) were obtained from normal human donors. The T cell proliferation was induced by stimulation of the cells with the T cell mitogen phytohemagglutinin (PHA), either in the absence (positive proliferation control), or in the presence of test peptides - 10 micrograms per ml - to examine their effects on the T cell proliferating response. 10^5 /well PBMC were plated in 96-well microtiter plates and assayed in duplicate with the peptides. Cell cultures were incubated at 37°C for 3 days in a 5% CO₂ incubator and were thereafter pulsed with 1microCi/well ³H-thymidine for additional 12 hours of culture. At the end of incubation time, the plates were harvested and the cells counted by liquid scintillation for the incorporation of ³H-thymidine as a measure of T cell proliferation.

Results from T cell proliferation assay: Peptides of the invention and the peptide combination showed no significant inhibitory effects on the proliferation of specifically stimulated human T-cells.

EXAMPLE 10:**B cell proliferation assay**

Human Peripheral Blood Mononuclear Cells (PBMC) were obtained from normal human donors. The B cell proliferation was induced by stimulation of the cells with the B cell mitogen Staphylococcus aureus Cowans I (SAC) plus Interleukin-2, either in the absence (positive proliferation control), or in the presence of test peptides - 10 micrograms per ml - to examine their effects on the B cell response. 10^5 /well PBMC were plated in 96-well microtiter plates and assayed in duplicate with the peptides. Cell cultures were incubated at 37°C for 3 days in a 5% CO₂ incubator and were thereafter pulsed with 1microCi/well ³H-thymidine for additional 12 hours of culture. At the end of incubation time, the plates were harvested and the cells counted by liquid scintillation for the incorporation of ³H-thymidine as a measure of B cell proliferation.

Results from B cell proliferation assay: Peptides of the invention and the peptide combination showed no significant inhibitory effects on the proliferation of specifically stimulated human B-cells.

EXAMPLE 11:**Phagocytosis assay**

RAW 264.7 (Mouse leukaemic monocyte macrophage cell line) cells were obtained from ATCC and grown in RPMI 1640 medium containing 10% FBS. Cells were incubated in 12x75 mm tubes at 37°C with test peptides - 10 micrograms per ml - for

30 min prior to adding Fluorescein-labeled *Escherichia coli* bacteria as the agent to be ingested. After the cells were incubated for additional 60 min at 37°C and allowed to ingest the Fluorescein-labeled *Escherichia coli* bacteria, cells were fixed with 1% paraformaldehyde. The samples were then analyzed by flow cytometry to determine the amount of phagocytosis as a function of brightness (the greater the phagocytic activity, the more fluorescence in the macrophage population). Data are reported as % positive and the mean fluorescence intensity (MFI) of positively stained cells.

- 10 Results from phagocytosis assay: Peptides of the invention and the peptide combination showed no inhibitory effects on the phagocytic activity of murine macrophages.

EXAMPLE 12:

15 Apoptosis induction assay

Human A549 cells (carcinomic human alveolar basal epithelial cells) were utilized in the experiments employing the Annexin-5 apoptosis assay. Annexin-5 is a member of a highly conserved protein family that binds acidic phospholipids in a calcium-dependent manner. Annexin-5 possesses a high affinity for phosphatidylserine. Phosphatidylserine is translocated from the inner side of the plasma membrane to the outer layer when cells undergo death by apoptosis or cell necrosis and serves as a signal by which cell destined for death are recognized by phagocytes. Test peptides - 10 micrograms per ml - were exposed for 24 hours to the A549 cells before they were analyzed for signs of apoptosis.

- 25 Results from apoptosis induction assay: Peptides of the invention and the peptide combination showed no significant induction of apoptosis on human lung cells.

EXAMPLE 13:

Apoptosis prevention assay

- 30 Human A549 cells (carcinomic human alveolar basal epithelial cells) were utilized in the experiments employing the Annexin-5 apoptosis assay. Annexin-5 is a member of a highly conserved protein family that binds acidic phospholipids in a calcium-dependent manner. Annexin-5 possesses a high affinity for phosphatidylserine. Phosphatidylserine is translocated from the inner side of the plasma membrane to the outer layer when cells undergo death by apoptosis or cell necrosis and serves as a signal by which cell destined for death are recognized by phagocytes. A549 cells were pretreated for 30 min with test peptides - 10 micrograms per ml - followed by the exposure to C2 ceramide. Ceramide mediates cell apoptosis through the activation of the mitogen activating protein kinase (MAPK) and the stress activated

kinase (JNK/SAPK). C2 ceramide is a synthetic, membrane soluble analog of ceramide.

Results from apoptosis prevention assay: Peptide 2 reduced by 23.9% and peptide 1 did not reduce the ceramide-induced apoptosis on human lung cells. The peptide combination (peptide 1 : peptide 2 = 0.95 : 1.05 mole) did not show any synergistic effect on the ceramide-induced apoptosis on human lung cells.

EXAMPLE 14:

Th1/Th2 cytokine profiling assay

The Balb/c mice (originated in 1923, it is a popular strain and is used in many different research disciplines. Also classified as an inbred from the production of 20 or more successive brother-sister matings, the Balb/c mouse is albino and small in size) were immunized on Days 1, 15, and 29 with Ovalbumin (Ovalbumin is the main protein found in egg white, commonly used to stimulate an immunological reaction in test animals) in PBS (5 micrograms/injection). On day 50, spleens of the mice were harvested (3 weeks after last boost with Ovalbumin). Cells were cultured (2×10^5 /well in triplicate) and incubated with culture medium or test peptides - 10 micrograms per ml - for 30 min. Thereafter, additional Ovalbumin was added to the cells at 10 micrograms/ml for in vitro restimulation of the cells. 72 hours later, cell supernatants were harvested and assayed using the Becton Dickinson Mouse Th1/Th2 Cytokine CBA Kit. This kit can be used to measure Interleukin-2 (IL-2), Interleukin-4 (IL-4), Interleukin-5 (IL-5), Interferon- γ (IFN- γ), and Tumor Necrosis Factor- α (TNF- α) protein levels in a single sample. The kit performance has been optimized for analysis of physiologically relevant concentrations (pg/ml levels) of specific cytokine proteins in tissue culture supernatants and serum samples.

Results from Th1/Th2 Cytokine assay: Peptide 1 induced by 23.7% the production of TNF- α , by 58.8% the production of IL-2, by 55.6% the production of IL-4 and did not significantly change the production of IFN- γ and IL-5 in murine spleen cells. Peptide 2 decreased by 31.4% the production of IFN- γ , increased by 80.4% the production of IL-2, increased by 35.2% the production of IL-4 and did not significantly change the production of TNF- α and IL-5 in murine spleen cells. The peptide combination (0.95 mole peptide 1 and 1.05 mole peptide 2) increased by 82% the production of IL-2, increased by 46% the production of IL-4 and did not significantly change the production of TNF- α , IFN- γ and IL-5 in murine spleen cells.

EXAMPLE 15:

TNF alpha production assay

Human Peripheral Blood Mononuclear Cells (PBMC) were obtained from normal human donors. The macrophages were prepared by adherence of PBMC to the

plastic wells of the plates. After 8 days in culture in the presence of recombinant human macrophage-colony stimulating factor at 2ng/ml, differentiated macrophages were preincubated with test peptides - 10 micrograms per ml - for 30 min, followed by in-well stimulation by the addition of lipopolysaccharide at a final concentration of 200ng/ml. Not stimulated macrophages served as negative background control.

After overnight incubation, supernatants from the control and LPS-stimulated cultures were harvested and assayed for TNF alpha production employing a TNF alpha specific ELISA.

Results from TNF alpha assay: Peptides of the invention and the peptide combination did not significantly change the LPS-induced TNF-alpha production in human macrophages.

EXAMPLE 16:

Endothelial cell migration assay

Endothelial cell migration is a prerequisite for the process of neo-vascularization or angiogenesis which is crucial for on-site recruitment of blood vessel formation. Primary Human endothelial cells (HUVEC) were seeded in insert chambers with 3 micrometer pore size of multi-transwell plate for 6 hours at 37°C in Endothelial Cell Basal Medium (EBM) supplemented with 0.1% bovine serum albumin. Thereafter, designated concentration of testing peptides - 10 micrograms per ml - was added in duplicate wells. The endothelia were allowed to migrate for 22 hours at 37°C, then, migrated cells were fixed and stained with Hoechst 33342 dye. Images of 3 fields per insert were taken and the number of migrated cells per field were quantified using the ImageProPlus software. Data were analyzed for the average number of the migrated cells and standard deviation of six data points for each treatment condition. Active test peptides against HUVEC migration was determined based on 50% inhibition of migrated cells as compared with the control. Statistic p values were computed using the Student's t-test.

Results from endothelial cell migration assay: Peptide 1 inhibited by 31% the migration of human endothelial cells, peptide 2 had no effect on the migration of endothelial cells, and the peptide combination (peptide 1 : peptide 2 (1.10 mole : 0.90 mole) did not show any synergistic effect on the inhibition of the migration of human endothelial cells.

EXAMPLE 17:

Endothelial tube formation assay

The endothelial tube formation assay is based on the ability of endothelial cells to form three-dimensional capillary-like tubular structures when cultured on a gel of

basement membrane extract. The endothelial tube formation assay represents a powerful model for studying inhibition and induction of angiogenesis. Pre-labeled HUVEC with Calcein AM were seeded in a 96-well culture plate coated with extracellular matrix (Chemicon international Cat. ECM625) and treated with test peptides - 10 micrograms per ml - in full growth medium. Positive control was vehicle only. The endothelial cells were allowed to form tubes for 20 hours and were then examined under an inverted fluorescent microscope. Duplicate wells for each treatment were photographed and quantitatively analyzed for an average tubule length using image analysis software ImageProPlus. Raw data were expressed as average tubule lengths in pixels $\pm$ standard deviation. Statistic p values were computed using the Student's t-test.

Results from endothelial tube formation assay: Peptide 1 had no effect on the tube formation of human endothelial cells, peptide 2 inhibited by 28% the tube formation of endothelial cells, and the peptide combination (peptide 1 : peptide 2 (1.10 mole : 0.90 mole) did not show any synergistic effect on the inhibition of the tube formation of human endothelial cells.

EXAMPLE 18

20 Mother milk formulation

Methods to prepare mother milk or artificial mother milk formulations or mother milk substitutes are described in WO03043429, US5962062, WO0030461, EP0527283, EP0832565

One example of an artificial mother milk or mother milk substitute formulation is provided in the following while also the other formulations disclosed in the above mentioned references can be used and are included herewith by reference.

The milk substitute contains, by weight, approximately 15% skimmed milk solids, approximately 75% demineralized water, approximately 9% soya oil, approximately 0.02% of carrageenates, 0.2% lecithin, and approximately 0.2% of disodium hydrogenphosphate.

In a first step, the solubilizing aqueous medium is produced, comprises, by weight, approximately 75% of water, approximately 0.02% of carrageenate and approximately 0.2% of disodium hydrogenphosphate.

The skimmed milk powder is then added to the solution for 10 min at 60°C and dissolved in the liquid.

Then soya oil and lecithin are added to the milk substitute composition at 60°C. The milk composition is allowed to stand 30 min at 55°C. After pasteurization, the

peptide 1 of the invention is added in liquid or powder form in such a quantity that the milk composition obtained comprises an amount of 5-50 micrograms, preferably 10-40 micrograms per 100 ml of milk composition. Optionally peptide 2 could be added in similar or smaller amounts to the obtained composition.

5

EXAMPLE 19**Gel formulation**

0.5 g of peptide 1

1.6 g of isopropanol

10 1.0 g of glycerol

1.6 g of polyoxyethylene-polyoxypropylene copolymer 12500 (Pluronic F127)

5.3 g of water

are mixed for 10 minutes and then heated to 85°C under continuous stirring for 15 minutes. The solution is cooled to room temperature under stirring. During the cooling phase the solution begins to gel at a temperature of about 45°C to form a clear gel. The gel contains 5% of the peptide combination for medical use. Optionally peptide 2 could be added in an amount from 0.01 to 0.5 g.

15

20 **EXAMPLE 20****Lotion formulation**

0.5 g of peptide 1

1.9 g of isopropanol

1.0 g of dimethylisosorbide

25 1.0 g of polyoxyethylene-polyoxypropylene copolymer 12500 (Pluronic F127)

5.6 g of water

are stirred and heated at 50°C, until a clear solution has been formed. Then the composition is cooled to room temperature under stirring. The lotion contains 5% of peptide combination for medical use. Optionally peptide 2 could be added in an amount from 0.01 to 0.5 g.

30

Claims

1. Peptide having the formula Gly-Arg-Gly-Asp-Asn-Pro for use in medicine.
- 5 2. Peptide according to claim 1 in combination with peptide having the formula Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe for use in medicine.
- 10 3. Combination of the peptides Gly-Arg-Gly-Asp-Asn-Pro and Ser-Val-Ser-Glu-Ile-Gln-Leu-Met-His-Asn-Leu-Gly-Lys-His-Leu-Asn-Ser-Met-Glu-Arg-Val-Glu-Trp-Leu-Arg-Lys-Lys-Leu-Gln-Asp-Val-His-Asn-Phe and salts and hydrates thereof.
- 15 4. Combination according to claim 3, wherein the peptides are contained in the combination in an amount from 30% by weight to 70% by weight to 70% by weight to 30% by weight.
5. Combination according to claim 3 or 4 for use in medicine.
- 20 6. Use of the peptide according to claim 1 or 2 or the combination according to claim 3 or 4 or 5 for the manufacture of a pharmaceutical composition for treatment and/or prophylaxis of cancer, autoimmune diseases, fibrotic diseases, inflammatory diseases, neurodegenerative diseases, infectious diseases, lung diseases, heart and vascular diseases and metabolic diseases.
- 25 7. Use of the peptide according to claim 6, wherein cancer, the autoimmune disease, fibrotic disease, inflammatory disease, neurodegenerative disease, infectious disease, lung disease, heart and vascular disease or metabolic disease is selected from AIDS, AIDS related diseases, HIV, Kaposi Sarcoma, lymphomas, anal dysplasia, cervical dysplasia, dementia complex, peripheral neuropathy, depression, anxiety, fatigue, anemia, thrombocytopenia, lipodystrophy, pulmonary hypertension, Wasting syndrome, aspergillosis, atypical mycobacteriosis, bacteraemia, candidiasis, HCMV infection, cryptococcosis, cryptosporidiosis, enteritis, histoplasmosis, nocardiosis, 30 encephalitis, progressive multifocal leukoencephalopathy, pneumonia, toxoplasmosis, tuberculosis.
- 35

8. Use of the peptide according to claim 1 or 2 or the combination according to claim 3 or 4 or 5 for the preparation of a formulation for oral administration to newborns, toddlers, and/or infants.
- 5 9. Use of the peptide according to claim 1 or 2 or the combination according to claim 3 or 4 or 5 for the preparation of a lyophilized formulation or a buffered liquide formulation.
- 10 10. Pharmaceutical composition containing the peptide according to claim 1 or the combination according to claim 3 together with at least one pharmaceutically acceptable carrier, cryoprotectant, lyoprotectant, excipient and/or diluent.
- 15 11. Pharmaceutical composition according to claim 9 in the form of a lyophilisate or liquide buffer solution.
- 20 12. Pharmaceutical composition according to claim 10 or 11 suitable for intravenous administration, oral administration, or for administration by inhalation.
- 25 13. Pharmaceutical composition according to claim 10, 11 or 12 in the form of an artificial mother milk formulation or mother milk substitute suitable for oral delivery to newborns, toddlers and infants.
- 30 14. Pharmaceutical composition according to claim 10, 11, 12 or 13 suitable for the treatment and/or prophylaxis of cancer, an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, an infectious disease, a lung disease, a heart and vascular disease or a metabolic disease.
- 35 15. Method for treating cancer, an autoimmune disease, a fibrotic disease, an inflammatory disease, a neurodegenerative disease, an infectious disease, a lung disease, a heart and vascular disease or a metabolic disease in a mammal, including a human, which comprises administering to the mammal a pharmaceutically effective amount of the peptides Gly-Arg-Gly-Asp-Asn-Pro or salts and hydrates thereof effective to treat the cancer, the autoimmune disease, the fibrotic disease, the inflammatory disease, the neurodegenerative disease, the infectious disease, the lung disease, the heart and vascular disease or the metabolic disease.