

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/11, C07H 21/04, 21/02, A61K 31/70	A2	(11) International Publication Number: WO 94/23028 (43) International Publication Date: 13 October 1994 (13.10.94)
(21) International Application Number: PCT/US94/03454 (22) International Filing Date: 30 March 1994 (30.03.94) (30) Priority Data: 08/040,752 31 March 1993 (31.03.93) US (60) Parent Application or Grant (63) Related by Continuation US 08/040,752 (CIP) Filed on 31 March 1993 (31.03.93) (71) Applicant (for all designated States except US): HYBRIDON, INC. [US/US]; One Innovation Drive, Worcester, MA 01605 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): AGRAWAL, Sudhir [IN/US]; 61 Lamplighter Drive, Shrewsbury, MA 01545 (US). TANG, Jin-Yan [CN/US]; 19 Sheridan Drive, #10, Shrewsbury, MA 01545 (US). PADMAPRIYA, Abeyasinghe [LK/US]; 23 Pal Drive, Shrewsbury, MA 01545 (US). (74) Agent: GREENFIELD, Michael, S.; Allegretti & Witcoff, Ltd., Ten South Wacker Drive, Chicago, IL 60606 (US).		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: MODIFIED OLIGONUCLEOTIDES HAVING IMPROVED ANTI-INFLUENZA ACTIVITY		
(57) Abstract The invention provides anti-influenza modified oligonucleotides that have greater efficacy in inhibiting influenza replication or propagation than previously described oligonucleotides. The greater efficacy arises from structural features such as chimeric or hybrid backbones, nuclease resistance-conferring terminal capping structures and/or self-complementary regions.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

MODIFIED OLIGONUCLEOTIDES HAVING IMPROVED
ANTI-INFLUENZA ACTIVITY

This is a continuation-in-part of Ser. No. 07/909,069,
5 filed July 2, 1992. This is also a continuation-in-part of
Ser. No. 07/918,239, filed July 23, 1992. This is also a
continuation-in-part of Ser. No. 07/698,568, filed May 10,
1991.

BACKGROUND OF THE INVENTION

Field Of The Invention

The invention relates to anti-sense oligonucleotides.
More particularly, the invention relates to modified
oligonucleotides that are capable of inhibiting replication or
propagation of influenza virus.

Summary Of The Related Art

Influenza A virus is a membrane-enclosed virus whose
genome is a segmented minus strand of RNA. The ten influenza
virus genes are present on eight segments of the single-
stranded RNA of strains A and B, and on seven segments of
20 strain C. The segments are of varying sizes (ranging from 890
to 2341 nucleotides in length) and each is a template for
synthesis for a different mRNA. The influenza virus virion
contains virus-specific RNA polymerases necessary for mRNA
synthesis from these templates and, in the absence of such

- 2 -

specific polymerases, the minus strand of influenza virus RNA is not infectious. Initiation of transcription of the mRNAs occurs when the influenza virus mRNA polymerase takes 12 to 15 nucleotides from the 5' end of a cellular mRNA or mRNA precursor and uses the borrowed oligonucleotide as a primer. Generally, the mRNAs made through this process encode only one protein. The M RNA and the NS RNA also encode spliced mRNAs, which results in production of two different proteins for each of these two segments.

Influenza viruses infect humans and animals (e.g., pigs, birds, horses) and may cause acute respiratory disease. There have been numerous attempts to produce vaccines effective against influenza virus. None, however, have been completely successful, particularly on a long-term basis. This may be due, at least in part, to the segmented characteristic of the influenza virus genome, which makes it possible, through reassortment of the segments, for numerous forms to exist. For example, it has been suggested that there could be an interchange of RNA segment between animal and human influenza viruses, which would result in the introduction of new antigenic subtypes into both populations. Thus, a long-term vaccination approach has failed, due to the emergence of new subtypes (antigenic "shift"). In addition, the surface proteins of the virus, hemagglutinin and neuraminidase, constantly undergo minor antigenic changes (antigenic

- 3 -

"drift"). This high degree of variation explains why specific immunity developed against a particular influenza virus does not establish protection against new variants. Hence, alternative antiviral strategies are needed. Although influenza B and C viruses cause less clinical disease than the A types, chemical antivirals should also be helpful in curbing infections caused by these agents.

Consequently, there is considerable interest in the development of anti-sense oligonucleotides that are capable of inhibiting the replication or propagation of influenza virus. Such anti-sense oligonucleotides hold the promise of providing broader protection against different strains of influenza, because they can be designed to hybridize to conserved regions of the influenza genome that are present in multiple strains of influenza virus.

Agrawal et al., U.S. Patent No. 5,194,428 the teachings of which are hereby incorporated by reference, discloses oligonucleotides having certain modified internucleotide linkages that are capable of inhibiting influenza replication by hybridizing to the influenza PB1 polymerase gene. Cowser et al., W092/03454 (1992) discloses antisense oligonucleotides that inhibit influenza virus propagation by hybridizing to the influenza polymerase 1, 2 or 3, or the hemagglutinin, nucleoprotein, neuraminidase, matrix protein, or nonstructural

- 4 -

protein gene, or to various splice junction sites or packaging sequences.

Pederson et al., U.S. Patent Nos. 5,149,797 and 5,XXX,XXX (Serial No. 07/839,472, allowed December 24, 1992), the teachings of which are hereby incorporated by reference, discloses chimeric mixed phosphate backbone oligonucleotides having RNase H activating segments adjacent to RNase H inactivating segments.

Thus, anti-sense oligonucleotides show promise as anti-influenza therapeutic agents. As with any such agents, however, there remains a need for improved agents that have even greater efficacy in inhibiting influenza virus replication or propagation. Such improved anti-sense oligonucleotides would be useful for studying which regions of the influenza virus genome are the best candidates for broad cross-strain inhibition as well as for use, and as anti-influenza therapeutic agents.

- 5 -

BRIEF SUMMARY OF THE INVENTION

The invention provides modified oligonucleotides having greater efficacy in inhibiting the replication or propagation of influenza virus than previously known oligonucleotides. These modified oligonucleotides are characterized by having a nucleotide sequence sufficiently complementary to an essential nucleic acid of influenza virus origin to hybridize to such influenza virus nucleic acid in a cell. In preferred embodiments, the essential nucleic acid is a portion of the PB1, 2 or 3 polymerase gene or to the hemagglutinin, nucleoprotein, neuraminidase, matrix protein, or nonstructural protein gene of influenza or to the various influenza splice junction sites or packaging sequences.

In various embodiments, modified oligonucleotides according to the invention have one or more type of modified internucleotide linkage. In a preferred embodiment, at least some of the modified internucleotide linkages are phosphorothioate or phosphorodithioate linkages. In certain preferred embodiments, phosphorothioate internucleotide linkages are present in a modified oligonucleotide that also contains other modified, i.e. nonphosphodiester, linkages. Preferably these other modified internucleotide linkages are alkylphosphonate or alkylphosphonothioate linkages. Most preferably, these other modified internucleotide linkages are present at or near one or both ends of the oligonucleotide.

- 6 -

In additional preferred embodiments of modified oligonucleotides according to the invention, phosphorothioate or phosphorodithioate internucleotide linkages are present in an oligonucleotide having one or more modified nucleoside. Preferably, such modified nucleoside is a 2'-O alkyl nucleoside. Most preferably, at least some modified nucleosides are present at or near one or both ends of the oligonucleotide.

In yet additional preferred embodiments of modified oligonucleotides according to the invention, phosphorothioate or phosphorodithioate internucleotide linkages are present in an oligonucleotide having an exonuclease resistance-conferring cap structure at one or both ends. Preferably, such a cap structure is present at least at the 3' end of the molecule. Such cap structures may also be present at one or both ends of all embodiments of modified oligonucleotides according to the invention and preferably are present at least at the 3' end of the oligonucleotide.

In further preferred embodiments of modified oligonucleotides according to the invention, phosphorothioate or phosphorodithioate internucleotide linkages are present in an oligonucleotide that is self-stabilized by having a self-complementary region involving nucleotides at or near one or both ends, and preferably at least at or near the 3' end, whereby the oligonucleotide forms a hairpin-like structure.

- 7 -

Each of these modified oligonucleotides according to the invention provides greater efficacy in inhibiting replication or propagation of influenza than any known oligonucleotide or modified oligonucleotide that hybridizes to the same gene.

5 Each of these modified oligonucleotides according to the invention may optionally also contain other modifications to the sugars or bases of the oligonucleotide and may also optionally have additional internucleotide linkages other than phosphorothioate, phosphorodithioate, alkylphosphonate or
10 alkylphosphonothioate linkages.

- 8 -

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide sequence of an influenza virus PB1 (polymerase 1) gene, against which complementary antisense oligonucleotides can be prepared.

5 Figure 2 shows a self-stabilized anti-influenza modified oligonucleotide according to the invention.

Figure 3 shows an alternative form of a self-stabilized anti-influenza modified oligonucleotide according to the invention.

10 Figure 4 shows certain preferred cap structures that confer exonuclease resistance upon oligonucleotides.

Figure 5 shows in vivo nucleolytic degradation patterns for 5'-capped, 3'-capped and uncapped oligonucleotides.

15 Figure 6 shows DNA Polymerase I 3'-exonuclease degradation patterns for self-stabilized and non-self-stabilized oligodeoxynucleotide phosphodiesteres.

Figure 7 shows DNA Polymerase I 3'-exonuclease degradation patterns for self-stabilized and non-self-stabilized oligodeoxynucleotide phosphorothioates.

- 9 -

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The invention relates to anti-sense oligonucleotides having anti-influenza activity. Modified oligonucleotides having in vivo activity against influenza are referred to herein as anti-influenza modified oligonucleotides. The invention provides anti-influenza modified oligonucleotides that have greater efficacy in inhibiting replication or propagation of influenza virus than known oligonucleotides or modified oligonucleotides that hybridize to the same gene. Modified oligonucleotides according to the invention have specific preferred characteristics that are discussed in greater detail for each preferred embodiment below. In addition to these characteristics, modified oligonucleotides according to the invention may optionally have additional ribonucleotide, 2'-substituted ribonucleotide, and/or deoxyribonucleotide monomers, any of which are connected together via 5' to 3' linkages which may include any of the internucleotide linkages known in the art. Preferably, such modified oligonucleotides may optionally contain phosphodiester, phosphotriester, phosphoramidate, siloxane, carbonate, carboxymethylester, acetamidate, carbamate, thioether, bridged phosphoramidate, bridged methylene phosphonate, bridged phosphoramidate, bridged methylene phosphonate, bridged phosphorothioate and/or sulfone internucleotide linkages. Those skilled in the art will

- 10 -

recognize that the synthesis of oligonucleotides containing any of these internucleotide linkages is well known to those skilled in the art, as is illustrated by articles by Uhlmann and Peyman, Chemical Reviews 90:543-584 (1990) and Schneider and Banner, Tetrahedron Lett. 31:335 (1990). Preferably, modified oligonucleotides according to the invention should contain from about 6 to about 100 monomers in total. Such modified oligonucleotides may also optionally contain modified nucleic acid bases and/or sugars, as well as added substituents, such as diamines, cholesteryl or other lipophilic groups.

Various preferred embodiments of modified oligonucleotides according to the invention are illustrated in Table I, below. Although these embodiments all have a nucleotide sequence from the same region of the influenza PBI polymerase gene, those skilled in the art will recognize that the anti-influenza efficacy of oligonucleotides having nucleotide sequences complementary to other essential nucleic acid sequences of influenza virus can also be enhanced by incorporating into such oligonucleotides the structural features of preferred embodiments of modified oligonucleotides according to the invention. For purposes of the invention, complementary means having a sequence that hybridizes to the essential nucleic acid sequence under physiological conditions. An essential nucleic acid sequence of influenza

- 11 -

virus means a nucleic acid sequence that is required for replication or propagation of influenza virus. Such essential nucleic acid sequences can be from any known strain of influenza virus. For example such oligonucleotides can have other sequences from the influenza PB1 polymerase gene (polymerase 1), as shown in Table I of U.S. Patent No. 5,194,428, the teachings of which are hereby incorporated by reference. Indeed, any sequence from the influenza PB1 (polymerase 1) gene [SEQ. ID. No. 1] (see Figure 1) should serve as the basis for modified oligonucleotides according to the invention. Alternatively, sequences from other influenza sequences or genes can be used (see Table II). As a practical matter, the structural features of preferred embodiments of modified oligonucleotides according to the invention should enhance the anti-influenza activity of any antisense oligonucleotide having a nucleotide sequence that hybridizes in a cell with any essential nucleic acid sequence of influenza virus.

Each preferred embodiment of modified oligonucleotides according to the invention is separately discussed in greater detail below.

- 12 -

TABLE I

Modified Oligonucleotides Having Superior
Anti-Influenza Activity

<u>Compound Identification NO.]</u>	<u>Sequence and Structure</u>	<u>[SEQ. ID</u>
CMPD A 2]	<u>CAGAGCAAAATCATCAGAAGA</u> ¹	[SEQ. ID NO.
CMPD B 3]	<u>CAGAGCAAAATCATCAGAAGA</u> ²	[SEQ. ID NO.
CMPD C 4]	<u>CAGAGCAAAATCATCAGAAGA</u> ³	[SEQ. ID NO.
CMPD D 5]	<u>CAGAGCAAAATCATCAGAAGA</u> ³	[SEQ. ID NO.
CMPD E 6]	<u>CAGAGCAAAATCATCAGAAGA</u> ²	[SEQ. ID NO.
CMPD F 7]	<u>CAGAGCAAAATCATCAGAAGA</u> ²	[SEQ. ID NO.
CMPD G 8]	<u>CAGAGCAAAATCATCAGAAGA</u> ³	[SEQ. ID NO.
CMPD H 9]	<u>CAGAGCAAAATCATCAGAAGA</u> ³	[SEQ. ID NO.
CMPD I 10]	<u>CAGAGCAAAATCATCAGAAGA</u> -C ₁₂ ⁴	[SEQ. ID NO.
CMPD J 12]	<u>AGAGCAAAATCATCAGAAG</u> ³	[SEQ. ID NO.
CMPD K 14]	<u>GAGCAAAATCATCAGAA</u> ³	[SEQ. ID NO.
CMPD L 15]	<u>CAGAGCAAAATCATCAGAAGA</u> ³	[SEQ. ID NO.
CMPD M 16]	<u>CAGAGCAAAATCATCAGAAGATTCTGATGA</u> ⁵	[SEQ. ID NO.

- 13 -

- 0 All sequences shown 5' to 3'; nonunderlined regions are oligonucleotide phosphorothioates.
- 1 Underscoring represents nucleotides connected by methylphosphonate internucleotide linkage.
- 2 Underscoring represents nucleotides connected by methylphosphonothioate linkage.
- 3 Underscoring represents 2'-OMe nucleotides.
- 4 C₁₂ represents cap structure.
- 5 Self-stabilized.

- 14 -

TABLE II
Additional Anti-Influenza Oligonucleotide Potential Sequences

<u>Sequence</u>	<u>Target</u>	<u>[SEQ. ID. No.]</u>
CTT TCC ATA TTG AAT ATA AT	AUG segment 1 polymerase 3	[SEQ. ID. No. 17]
ACA TCC ATT CAA ATG GTT TG	AUG segment 2 polymerase 1	[SEQ. ID. No. 18]
TCT TCC ATT TTG GAT CAG TA	AUG segment 3 polymerase 2	[SEQ. ID. No. 19]
GCC TTC ATT TTG GTT GTT TT	AUG segment 4 hemagglutinin	[SEQ. ID. No. 20]
GAC GCC ATG ATT TTG ATG TC	AUG segment 5 nucleoprotein	[SEQ. ID. No. 21]
GGA TTC ATT TTA AAC CCC TG	AUG segment 6 neuraminidase	[SEQ. ID. No. 22]
AGA CTC ATC TTT CAA TAT CT	AUG segment 7 matrix protein	[SEQ. ID. No. 23]
GAT AGA GAG AAC GTA CGT TT	left splice junction segment 7	[SEQ. ID. No. 24]
TCT GAT AGG CCT GCA AAT TT	right splice junction segment 7	[SEQ. ID. No. 25]
GGA TCC ATT ATG TCT TTG TC	AUG segment 8 nonstructural protein	[SEQ. ID. No. 26]
CAT GTC GGT TAG GTA ACG CG	splice branch segment 8	[SEQ. ID. No. 27]
GCA ATC TAC CTG AAA GCT TG	right splice junction segment 8	[SEQ. ID. No. 28]
AGC AGT ATG TCC TGG AAG AG	left splice junction segment 8	[SEQ. ID. No. 29]
AAA ACG ACC TTG TTT CTA CT	packaging sequence segment 1	[SEQ. ID. No. 30]
AAA AAT GCC TTG TTC CTA CT	packaging sequence segment 2	[SEQ. ID. No. 31]

- 15 -

AAA AGT ACC TTG TTT CTA CT	packaging [SEQ. ID. No. 32] sequence segment 3
AAA ACA CCC TTG TTT CTA CT	packaging [SEQ. ID. No. 33] sequence segment 4
AAA ATA CCC TTG TTT CTA CT	packaging [SEQ. ID. No. 34] sequenc segment 5
AAA AAC TCC TTG TTT CTA CT	packaging [SEQ. ID. No. 35] sequence segment 6
AAA ACT ACC TTG TTT CTA CT	packaging [SEQ. ID. No. 36] sequence segment 7
AAA ACA CCC TTG TTT CTA CT	packaging [SEQ. ID. No. 37] sequence segment 8

In a first preferred embodiment, anti-influenza modified oligonucleotides according to the invention are in the form of a mixed backbone chimeric oligonucleotide having one or more regions of nucleotides connected by phosphorothioate or phosphorodithioate internucleotide linkages ("phosphorothioate or phosphorodithioate region") as well as one or more regions of nucleotides connected by alkylphosphonate internucleotide linkages ("alkylphosphonate region"). In this embodiment, at least one alkylphosphonate region preferably includes nucleotides at or near the 5' end and/or the 3' end of the oligonucleotide. For purposes of the invention, "at or near the 5' or the 3' end of the oligonucleotide" means involving at least one nucleotide within about 5 nucleotides from the 5' or 3' end of the oligonucleotide. Preferably, the alkylphosphonate region comprises from about 2 to about 10

- 16 -

contiguous nucleotides connected by alkylphosphonate linkages. Preferably, the phosphorothioate or phosphorodithioate region comprises at least 3, and up to about 100 contiguous nucleotides connected by phosphorothioate or phosphorodithioate linkages. An example of an anti-influenza modified oligonucleotide according to this embodiment of the invention is shown in Table I as CMPD A.

Anti-influenza modified oligonucleotides according to this embodiment of the invention are synthesized by solid phase methods, alternating H-phosphonate chemistry and sulfur oxidation for phosphorothioate regions, and alkylphosphonamidate chemistry for alkylphosphonate regions. A preferred H-phosphonate approach is taught by Agrawal et al., U.S. Patent No. 5,149,798, the teachings of which are hereby incorporated by reference. Alkylphosphonamidite chemistry is well known in the art, as illustrated by Agrawal and Goodchild, Tetrahedron Lett. 28:3539-3542 (1987). Synthesis of phosphorodithioate-containing oligonucleotides is also well known in the art, as illustrated by U.S. Patent No. 5,151,510, the teachings of which are hereby incorporated by reference (See also, e.g., Marshall and Caruthers, Science 259: 1564-1570 (1993) and references cited therein).

In a second preferred embodiment, anti-influenza modified oligonucleotides according to the invention are in the form of a mixed backbone chimeric oligonucleotide having one or more

- 17 -

region of nucleotides connected by phosphorothioate or phosphorodithioate internucleotide linkages ("phosphorothioate or phosphorodithioate region"), as well as one or more region of nucleotides connected by alkylphosphonothioate or arylphosphonothioate internucleotide linkages ("alkylphosphonothioate region"). In this embodiment, at least one alkylphosphonothioate region preferably includes nucleotides at or near the 5' end and/or the 3' end of the oligonucleotide. Preferably, the alkylphosphonothioate region comprises from about 2 to about 10 contiguous nucleotides connected by alkylphosphonothioate linkages. Preferably, the phosphorothioate or phosphorodithioate region comprises at least 3, and up to about 100 contiguous nucleotides connected by phosphorothioate or phosphorodithioate linkages. Examples of anti-influenza modified oligonucleotides according to this embodiment of the invention are shown in Table I as CMPD B, CMPD E and CMPD F.

Anti-influenza modified oligonucleotides according to this embodiment of the invention are synthesized by solid phase methods, alternating chemistries for each region to be synthesized. Phosphorothioate or phosphorodithioate regions are synthesized as described for the first embodiment. Alkylphosphonothioate regions are synthesized by coupling together two or more nucleosides via alkylphosphite linkages, then oxidatively thiolating the alkylphosphite linkages to

- 18 -

produce alkylphosphonothioate linkages. This synthesis procedure is set forth in detail in Example 1.

In a third preferred embodiment, anti-influenza modified oligonucleotides according to the invention are in the form of a hybrid oligonucleotide having regions of deoxyribonucleotides ("deoxyribonucleotide regions") and regions of ribonucleotides or 2'-substituted ribonucleotides ("ribonucleotide regions"). Preferably, from about one to about all of the internucleotide linkages are phosphorothioate or phosphorodithioate linkages. Preferred 2'-substituted ribonucleotides are halo, amino, alkyl, aryl or lower alkyl (1-6 carbon atoms) substituted ribonucleotides, especially 2'-OMe-ribonucleotides. Preferably, at least some of the ribonucleotide regions include nucleotides present at or near the 5' end and/or the 3' end of the oligonucleotide. Most preferably, the ribonucleotide regions each comprise from about 2 and preferably from about 4 to about 100 contiguous ribonucleotides and/or 2'-substitute oligonucleotides. The deoxyribonucleotide regions are optional, and when present may contain from about 1 to about 100 contiguous deoxyribonucleotides. Examples of anti-influenza modified oligonucleotides according to this embodiment of the invention are shown in Table I as CMPD C, CMPD D, CMPD G, CMPD H, CMPD K, CMPD M and CMPD N.

- 19 -

Anti-influenza modified oligonucleotides according to this embodiment of the invention are typically synthesized by solid phase methods, preferably by the H-phosphonate approach, using deoxynucleotide H-phosphonates for deoxyribonucleotide regions, and ribonucleotide or 2'-substituted ribonucleotide H-phosphonates for ribonucleotide regions.

In a fourth preferred embodiment, anti-influenza modified oligonucleotides according to the invention are in the form of an oligonucleotide having at its 5' and/or 3' end a cap structure that confers exonuclease resistance to the oligonucleotide. Such modified oligonucleotides preferably also have from 1 to about all modified (non-phosphodiester) internucleotide linkage. Preferred cap structures include those shown in Figure 4, as well as lower alkyl (C1-C12) or alcohol groups. Preferred modified internucleotide linkages include phosphotriester, phosphoramidate, siloxane, carbonate, carboxymethylester, acetamidate, carbamate, thioether, bridged phosphoramidate, bridged methylene phosphonate, bridged phosphorothioate, sulfone, phosphorothioate and phosphorodithioate linkages.

Anti-influenza modified oligonucleotides according to this embodiment of the invention are synthesized according to procedures well known in the art (see e.g., Uhlmann and Peyman, Chemical Reviews 90:543-584 (1990); Schneider and Banner, Tetrahedron Lett. 31:335 (1990)). For

- 20 -

oligonucleotides having cap structures at the 3' end, the cap structure is reversibly attached to the solid support and is then coupled to the first nucleotide monomer in the synthesis scheme. For oligonucleotides having cap structures at the 5' end, the cap structure is coupled to the end of the oligonucleotide after addition of the last nucleotide monomer in the synthesis scheme.

In a fifth embodiment, anti-influenza modified oligonucleotides are self-stabilized by having a self-complementary region that hybridizes intramolecularly with the oligonucleotide to form an exonuclease resistant hairpin-like structure. Anti-influenza modified oligonucleotides according to this embodiment of the invention are generally characterized by having two regions: an influenza hybridizing region and a self-complementary region. The influenza hybridizing region has a nucleotide sequence that is complementary to an essential nucleic acid sequence of influenza virus. Preferably, this region has from about 6 to about 100 nucleotides. One form of this embodiment of the invention is shown in Figure 2. In this form, the influenza hybridizing region is shown as connected rectangular squares, and the self-complementary region is shown as connected circles. The complementary nucleic acid sequence in a target influenza messenger RNA molecule is represented by connected diamonds. Hydrogen bonding between nucleotides is indicated

- 21 -

by dots. The oligonucleotide is stabilized, i.e., rendered resistant to exonucleolytic degradation by base-pairing between the target hybridizing region and the self-complementary region and/or by base-pairing between complementary sequences within the self-complementary region. When the oligonucleotide encounters an influenza nucleic acid molecule having a complementary nucleic acid sequence, base-pairing between the influenza hybridizing region and the self-complementary region of the oligonucleotide is disrupted and replaced by base-pairing between the influenza hybridizing region of the oligonucleotide and the complementary nucleic acid sequence of the nucleic acid molecule. This disruption and replacement of base-pairing takes place because the intermolecular base-paired structure formed by the hybrid between the target nucleic acid sequence and the target hybridizing region is more thermodynamically stable than the intramolecular base-paired structure formed by the self-complementary oligonucleotide.

A second form of an oligonucleotide according to this embodiment of the invention operates in a similar way as the first form, but forms a different structure upon self-complementary base-pairing. This alternative form forms a hammer-like structure as shown in Figure 3. In this form, the self-complementary region contains oligonucleotide sequences that can base pair with other oligonucleotide sequences within

- 22 -

the self-complementary region. The self-complementary region may also contain oligonucleotide sequences that are complementary to the influenza hybridizing region.

5 The second significant region of self-stabilized oligonucleotides according to the invention is the self-complementary region. The self-complementary region contains oligonucleotide sequences that are complementary to other oligonucleotide sequences within the oligonucleotide. These other oligonucleotide sequences may be within the influenza
10 hybridizing region or within the self-complementary region, or they may span both regions. The complementary sequences form base pairs, resulting in the formation of a hairpin structure, as shown in Figure 2, or a hammer-like structure, as shown in Figure 3. Either the hairpin structure or the hammer-like
15 structure can have loops resulting from non-base-paired nucleotides, as shown in Figure 2 for the hairpin structure, or can be devoid of such loops, as shown in Figure 3 for the hammer-like structure. The number of base-pairs to be formed by intra-molecular hybridization involving the self-
20 complementary region may vary, but should be adequate to maintain a double-stranded structure so that the 3' end is not accessible to endonucleases. Generally, about 4 or more base-pairs will be necessary to maintain such a double-stranded structure. In a preferred embodiment, there are about 10
25 intramolecular base-pairs formed in the self-stabilized

- 23 -

oligonucleotide, with the 10 base pairs being consecutive and involving the 3'-most nucleotides. Of course, the intramolecular base-pairing can be so extensive as to involve every nucleotide of the oligonucleotide. Preferably, this will involve a self-complementary region of about 50 nucleotides or less.

Oligonucleotides according to this embodiment may have from 1 to about all modified internucleotide linkages, as described for the fourth embodiment. Preferably, at least either the influenza hybridizing region or the self-complementary region, and most preferably both, will contain from about 2 to about all nucleotides being coupled by phosphorothioate and/or phosphorodithioate linkages.

An example of an anti-influenza modified oligonucleotide according to this embodiment of the invention is shown in Table I as CMPD O.

Those skilled in the art will recognize that the features of the various preferred embodiments described above can be combined to produce additional embodiments that may have even greater anti-influenza activity. Thus, the invention contemplates anti-influenza modified oligonucleotides having every possible combination of chimeric features, hybrid features, cap structures and self-stabilizing character, all as described herein.

- 24 -

Each of the preferred embodiments of the present invention has greater ability to inhibit influenza replication or propagation than anti-sense oligonucleotides of the prior art that hybridize to the same influenza mRNA. For example, U.S. Patent No. 5,194,428, teaches modified oligonucleotides that inhibit influenza virus replication. Among those modified oligonucleotides are oligonucleotide phosphorothioates that hybridize to the influenza PB1 (polymerase 1) gene. In the present study, an oligonucleotide phosphorothioate that hybridizes to the influenza PB1 polymerase gene was tested for its ability to inhibit influenza virus replication in comparison with various preferred embodiments of the present invention. Each embodiment of the present invention tested demonstrated surprisingly improved efficacy in anti-influenza activity, relative to the oligonucleotide phosphorothioate that binds to the same site on the same gene. As shown in Table III, below, anti-influenza modified oligonucleotides according to the invention gave a reduction in the 50 per cent inhibitory concentration (IC_{50}) ranging from 2 to nearly fifteen fold.

- 25 -

TABLE III

Anti-Influenza Activity Of Oligonucleotide

<u>Oligonucleotide Improvement</u>	<u>IC₅₀ (μg/ml)</u>	<u>X - F o l d</u>
CAGAGCAAAATCATCAGAAGA ¹	317 ²	-
CMPD A	N.T.	-
CMPD B	34	9.3
CMPD C	55	5.8
CMPD D	58 ²	5.5
CMPD E	N.T.	-
CMPD F	178	1.8
CMPD G	147	2.2
CMPD H	59	5.4
CMPD I	45	7.0
CMPD J	27 ²	11.7
CMPD K	22 ²	14.4
CMPD L	61	5.2
CMPD M	175	1.8
CMPD N	104 ²	3.0
CMPD O	23 ²	13.8

¹ oligodeoxynucleotide phosphorothioate control² average of two experiments

Such oligonucleotides are useful for a variety of purposes. First, they can be used in studies to determine which influenza genes and sites within such genes provide the best basis for anti-sense oligonucleotides having broad effectiveness against multiple strains of influenza virus. Second, they can be used to determine what structural features

- 26 -

or combination of structural features provide the greatest effectiveness against influenza in vitro and in vivo. Finally, such oligonucleotides are useful as therapeutic agents for treating influenza infections. For such treatment, oligonucleotides may be administered intraperitoneally, intranasally, orally or anally. Preferably, such oligonucleotides will be administered at a concentration of from about 1 to about 50 mg/kg body weight.

The following examples are intended to further illustrate certain preferred embodiments of the invention and are not intended to be limiting in nature.

EXAMPLE 1

Synthesis Of A Chimeric Oligonucleotide Having Methylphosphonate And Phosphorothioate Regions

Chimeric oligonucleotides having methylphosphonate and phosphorothioate regions were synthesized using methylphosphonamidites for methylphosphonate regions and nucleoside H-phosphonates for phosphorothioate regions.

Methylphosphonamidite synthesis was as follows:

Methylchloro-N, N-diisopropylaminophosphine was prepared by reaction of methyldichlorophosphine (51 mmol) with diisopropylamine (102 mmol) in ether at 15° under nitrogen. After removal of salt by filtration and evaporation of solvent, methylchloro-N,N-diisopropylaminophosphine was obtained as an oil (48 mmol, 95% of theory) that was

- 27 -

characterized by ^1H and ^{31}P NMR and was stable at -20° for at least eight weeks. The product was reacted with the usual protected nucleosides in dichloromethane containing N, N-diisopropylethylamine at room temperature for 10-20 min.

5 Aqueous work up and precipitation from ethyl acetate using pentane at -30° to -40° gave product as white solids in 80-90% yield. Products were pure by tlc on silica in $\text{CH}_2\text{Cl}_2:\text{EtOAc}:\text{Et}^3\text{N}(9:9:2)$ and were characterized by ^1H and ^{31}P NMR.

10 These products were used in automated DNA synthesizer using the same conditions and program used for standard phosphoramidite reagents. Nucleotides were dissolved in acetonitrile at a concentration of 33 mg/ml and activated with tetrazole. Synthesis on prepacked CPG support was performed
15 using a coupling time of 1 minute.

Coupling efficiency was followed by the dimethoxytrityl assay and was found to be the same as for control syntheses run in parallel using phosphoramidites.

20 After coupling, the product was detritylated then cleaved from the support with NH_4OH at room temperature for 2 hrs and deblocked using ethylenediamine: ethanol (1:1) at room temperature for 4 hrs. This basic treatment caused about 1% degradation of the internucleoside phosphonate group in a model study assayed by HPLC.

- 28 -

H-phosphonate synthesis was as exemplified in U.S. Patent No. 5,149,798, the teachings of which are hereby incorporated by reference. Then, the H-phosphonates were converted to phosphorothioates by oxidation with 0.2 M sulfur in carbon disulfide/pyridine/triethylamine (9:9:1 vol/vol).

EXAMPLE 2

Synthesis Of A Chimeric Oligonucleotide Having Methylphosphonothioate And Phosphorothioate Regions

For preparing a modified oligonucleotide having regions of four contiguous nucleotide methylphosphonothioates at either end and a region of 12 contiguous oligonucleotide phosphorothioates in the middle, the following procedure on 8 micromole scale was used. A first monomer was pre-bound to a control pore glass (CPG) solid support. A second monomer, which was a deoxynucleotide methylphosphonamidite, was coupled to the first monomer using a standard amidite coupling cycle (see e.g., Agrawal and Goodchild, Tetrahedron Lett. 28:3539-3542 (1987)). In separate cycles, three more deoxynucleotide methylphosphonamidites were sequentially coupled to the growing chain. Then oxidative thiolation was carried out, using 1% Beaucage reagent (3H-1,2-benzodithiole-2-one) in acetonitrile for 5 minutes at ambient temperature to generate a CPG-bound pentanucleotide methylphosphonothioate. The next 12 monomers were added sequentially by the H-phosphonate approach of Agrawal and Zamecnik, U.S. Patent No. 5,149,798, the teachings

- 29 -

of which are hereby incorporated by reference. Then, the H-phosphonates were converted to phosphorothioates by oxidation with 0.2 M sulfur in carbon disulfide/pyridine/triethylamine (9:9:1 vol/vol). The final four monomers, which were deoxynucleotide methylphosphonothioates, were added, then oxidatively thiolated as described above. The resulting oligonucleotide was deprotected at room temperature for 30 minutes in 0.5 ml ethylene diamine, then kept at room temperature for 6 hours with occasional stirring. Finally, the mixture was filtered and evaporated in vacuo to obtain a solid mass and the mass was dissolved in water and desalted on Sep Pak C₁₈.

EXAMPLE 3

Resistance Of Oligonucleotides Having Methylphosphonothioate Linkages To Nucleolytic Degradation

Oligonucleotides having 2-4 contiguous deoxynucleotide methylphosphonates at 3' ends and otherwise having all deoxynucleotide phosphodiester were tested for their relative resistance to 3' exonucleolytic degradation compared with an oligonucleotide phosphodiester. For each oligonucleotide, 0.4 A₂₆₀ units of oligonucleotide was lyophilized, dissolved in 0.5 ml buffer (10 mM Tris, 10 mM MgCl₂, pH 8.5) and mixed with 5 µl (1.5 milliunits) of snake venom phosphodiesterase. The mixture was incubated at 37°C in a thermally regulated cell

- 30 -

and A_{260} was plotted against time. Increase in hyperchromicity was used as the indicator for oligonucleotide degradation. The results are shown in Table IV, below.

5 These results demonstrate that oligonucleotides having methylphosphonothioate linkages near the 3' end were far more stable than the oligonucleotide lacking such linkages. In addition, oligonucleotide stability increased with increasing numbers of methylphosphonothioate linkages (4 linkages >> 3 linkages > > 2 linkages).

TABLE IV
Resistance Of Oligonucleotides To Nucleolytic Degradation

<u>in</u> <u>Oligonucleotide</u> <u>hyperchromicity</u>	<u>t 1/2 (seconds)</u>	% increase
Oligonucleotide phosphodiester	44	22.56
Oligonucleotide with 2 3' terminal deoxynucleotide methylphosphonothioates	210	24.58
Oligonucleotide with 3 3' terminal deoxynucleotide methylphosphonothioates	264	18
Oligonucleotide with 4 3' terminal deoxynucleotide methylphosphonothioates	401	15.54

- 31 -

EXAMPLE 4Synthesis Of A Hybrid Oligonucleotide Phosphorothioate Having Deoxyribonucleotide And 2'-OMe-Ribonucleotide Regions

Hybrid oligonucleotide phosphorothioates were synthesized on CPG on a 5-6 μ mole scale on an automated synthesizer (model 8700, Millipore, Milford, MA) using the H-phosphonate approach described in U.S. Patent No. 5,149,798, the teachings of which are hereby incorporated by reference.

Deoxyribonucleoside H-phosphonates were obtained from Millipore. 2'-OMe ribonucleotide H-phosphonates were synthesized by standard procedures. Segments of oligonucleotides containing 2'-OMe nucleoside were assembled by using 2'-OMe ribonucleoside H-phosphonates for the desired cycles. Similarly, segments of oligonucleotides containing deoxyribonucleosides were assembled by using deoxynucleoside H-phosphonates for the desired cycles. After assembly, CPG bound oligonucleotide H-phosphonate was oxidized with sulfur as described in Example 2 to generate the phosphorothioate linkage. Oligonucleotides were then deprotected in concentrated NH_4OH at 40°C for 48 hours.

Crude oligonucleotide (about 500 A_{260} units) was analyzed on reverse low pressure chromatography on a C_{18} reversed phase medium. The DMT group was removed by treatment with 80% aqueous acetic acid, then the oligonucleotides were dialyzed against distilled water and lyophilized.

- 32 -

EXAMPLE 5Relative Nuclease Resistance Of
Hybrid Oligonucleotide Phosphorothioates

To test the relative nuclease resistance of various hybrid oligonucleotide phosphorothioates, the oligonucleotides were treated with snake venom phosphodiesterase (SVPD). About 0.2 A₂₆₀ units of oligos having no 2'-OMe-RNA region, or having 3 and 4 contiguous 2'-OMe ribonucleotides at 5' and 3' ends, respectively, or having all 2'-OMe nucleotides, were dissolved in 500 µl buffer (40 mM NH₄CO₃, pH 4.0 + 20 mM MgCl₂), and mixed with 0.1 units SVPD. The mixture was incubated at 37°C for 420 minutes. After 0, 200 and 420 minutes, 165 µl aliquots were removed and analyzed using ion exchange HPLC. The oligonucleotide having all 2'-OMe-ribonucleotides was very resistant to SVPD, whereas the oligonucleotide having no 2'-OMe-ribonucleotides was digested almost to completion and the oligonucleotide having 5' and 3' terminal 2'-OMe-RNA regions was digested to 50%. An oligonucleotide phosphodiester was digested to about 80% in one minute using one tenth of the concentration of SVPD.

These results indicate that the presence of 2'-OMe ribonucleotides in an oligonucleotide phosphorothioate enhances resistance to exonucleolytic digestion and that this enhanced resistance increases when a larger proportion of 2'-OMe ribonucleotides are used. Due to the similar character

- 33 -

and behavior of ribonucleotides, other 2'-substituted ribonucleotides and 2'-OMe ribonucleotides, these results also suggest that similar enhancement of nuclease resistance would be obtained for hybrid oligonucleotide phosphorothioates and/or phosphorodithioates having ribonucleotides, 2'-substituted ribonucleotides, or a mixture of ribonucleotides and 2'-substituted ribonucleotides.

EXAMPLE 6

Synthesis Of An Oligonucleotide Phosphorothioate Having A 3' Cap Structure

Oligonucleoside phosphorothioates were synthesized on a Model 8700 automated synthesizer (Milligen-Bioscience, Burlington, MA) using H-phosphonate chemistry on controlled pore glass (CPG), followed by oxidation with 0.2 M sulfur in carbon disulfide/pyridine/triethylamine (9:9:1 vol/vol). Synthesis was carried out on a 5-10 micromolar scale. Oligonucleoside phosphorothioates were purified by low pressure ion exchange chromatography (DEAE-cellulose, DE-50 Whatman), followed by reverse phase chromatography (C₁₈) and dialysis. 5'-capped oligonucleoside phosphorothioates were prepared by carrying out the last coupling, after the assembly of the required sequence, with N-Fmoc-O'-DMTr-3-amino-1,2-propanediol-H-phosphonate. The 5'-capped oligonucleoside H-phosphonate was then oxidized with sulfur. 3'-capped oligonucleoside phosphorothioates were assembled on N-Fmoc-O'-

- 34 -

DMTr-3-amino-1,2-propanediol-CPG, followed by sulfur oxidation. Combination of these procedures was used to produce 3',5'-capped oligonucleoside phosphorothioates.

Alternatively, oligonucleoside phosphorothioates having other 3' or 5' cap structures, (see e.g., Figure 4), are prepared by substituting the phosphonate or CPG-derivatized cap structures for the N-Fmoc-O'-DMTr-3-amino-1,2-propanediol-H phosphonate or CPG in the capping procedure. Similarly, capped modified oligonucleotides other than oligonucleotide phosphorothioates are prepared in an analogous manner by appending the capping procedure to the appropriate synthetic procedure.

EXAMPLE 7

In Vivo Stability Of Oligonucleotide Phosphorothioates Having Terminal Cap Structures

Male CDC2F1 mice (average weight 20 grams) were treated by intravenous or intraperitoneal injection with a 30 mg/kg dose of radiolabelled oligonucleotides dissolved in 200 microlitres physiological saline. Each capped or uncapped oligonucleotide was administered to three mice. Urine was collected separately from each animal up to 24 hours post-dosing, then extracted with proteinase K. (2 mg/ml, final concentration) in 0.5% SDS, 10 mM 20 mM Tris Cl (pH 7.6), 10 mM EDTA for one hour at 37°C, followed by phenol-chloroform extraction and ethanol precipitation. Recovered

- 35 -

oligonucleotides were then analyzed by PAGE (20% polyacrylamide/7 M urea) followed by autoradiography. Radioactivity was also measure from cage rinse to account for urine spill.

5 Twenty-four hours after dosing, about 30% of oligonucleoside phosphorothioates were excreted, whether capped or uncapped. Excreted uncapped and 5'-capped oligonucleoside phosphorothioates were extensively degraded, as shown in Figure 5. Excreted 3'-capped and 3',5'-capped
10 oligonucleoside phosphorothioates, in contrast, demonstrated virtually no degradation. This indicates that in vivo degradation of oligonucleoside phosphorothioates excreted in urine is mediated by 3'-exonuclease activity which can be inhibited by adding a cap to the 3' hydroxyl group of the
15 oligonucleotide.

EXAMPLE 8

Nuclease Resistance Of Self-Stabilized Oligonucleotide Phosphodiester

20 The control oligonucleotide used for this study was an oligodeoxynucleotide phosphodiester without a self-complementary region. The test compound was identical, except that it had a 3' self complementary region of 10 nucleotides. To control for any size effects, another control oligodeoxynucleotide phosphodiester was used that was

- 36 -

identical to the first control oligonucleotide except for having at its 3' end 10 mismatched nucleotides (T_{10}).

The oligonucleotides were tested for their relative resistance to 3' exonucleolytic degradation. For each oligonucleotide, 0.4 A_{260} units of oligonucleotide was lyophilized, dissolved in 0.5 ml buffer (10 mM Tris, 10 mM $MgCl_2$, pH 8.5) and mixed with 5 μ l (1.5 milliunits) of snake venom phosphodiesterase (SVPD). The mixture was incubated at 37° C in a thermally regulated cell and A_{260} was plotted against time. Oligonucleotide degradation was measured as function of increase in hyperchromicity.

The results of these experiments are shown in Table V, below. These results demonstrate that self-stabilized oligonucleotide phosphodiesterases according to the invention are far more resistant to 3' exonucleolytic degradation than either oligonucleotide phosphodiesterases or oligonucleotide phosphodiesterases having a non-complementary tail.

In addition to the testing described above, the oligonucleotides were also subjected to DNA Polymerase I 3'-exonuclease digestion. As shown in Figure 6 the non-self-stabilized oligonucleotides were digested to completion in 30 minutes, whereas the self-stabilized oligonucleotide with a 10 nucleotide self-complementary region was only partly digested over 30 minutes.

- 37 -

TABLE V
SVPD Half-Life Of Oligonucleotides

<u>Supplementary Region</u> <u>Digestion</u>	<u>Half-Life</u> <u>For</u> <u>SVPD</u>
Absent	75 seconds
10 nucleotides	950 seconds
mismatched	75 seconds

EXAMPLE 9

Nuclease Resistance of Self-Stabilized
Oligonucleotide Phosphorothioates

To test the relative nuclease resistance of self-stabilized and non-self-stabilized oligonucleotide phosphorothioates, a DNA Polymerase I 3'-exonuclease activity assay was used, because of the slow degradation of oligonucleotide phosphorothioates by SVPD.

All oligonucleotides were labelled at the 5-end with gamma-³²P-ATP and kinase. To a solution of 40 pmole 5'-labelled oligonucleotide in 20 µl buffer (40 mM Tris HCl, pH 8.0,

10 mM MgCl₂, 5 mM DTT, 50 mM KCl, 50 µg/ml BSA), 5 units DNA polymerase I was added and incubated at 37° C. Aliquots of 4 µl were taken at 0, 30, 60, 120 minutes and were mixed with 6 µl stop solution (98% formamide, 10 mM EDTA, 0.1% xylene cyanol, 0.1% bromophenol blue). The samples were analyzed by 15% acrylamide gel (urea) and autoradiography.

- 38 -

The results are shown in Figure 7. Oligonucleotide phosphorothioates having no self-complementary region or only mismatched nucleotides (T_{10}) at the 3' end (as described in Example 8 for oligonucleotide phosphodiester) were digested to almost 50% within 4 hours. The oligonucleotide phosphorothioate having a 10 nucleotide self-complementary region was undegraded after 4 hours. Oligonucleotide phosphorothioates having 6 or 4 nucleotide self-complementary regions were also found to be stable. These results demonstrate that self-stabilized oligonucleotide phosphorothioates are far more resistant to nucleolytic degradation than are non-self-stabilized oligonucleotide phosphorothioates.

EXAMPLE 10

Anti-Influenza Activity Of Modified Oligonucleotides

MDCK canine kidney cells were seeded in Minimum Essential Medium (MEM) with non-essential amino acids (GIBCO BRL, Grand Island, New York) with 5% fetal bovine serum (Hyclone Laboratories, Logan, Utah), 0.1% NaHCO_3 , at a concentration of 4×10^5 cells/ml in 96 well tissue culture plates (Corning, Corning, New York) at 0.2 ml/well. The cells were incubated overnight to establish monolayers of cells. Growth medium was then removed and 0.1 ml of oligonucleotide at pre-selected concentrations in serum-free MEM containing 0.18% NaHCO_3 and

- 39 -

50 µg/ml gentamycin. This was done for each compound for each of 4 wells: 1 well as toxicity control and 3 wells as anti-viral tests. Three cell control wells and six virus control wells received 0.1 ml of serum-free MEM containing 0.18% NaHCO₃ and 50 µg/ml gentamycin. Within 10 minutes of addition of the oligonucleotide compounds, influenza A/NWS/33(H1N1) virus was added to each test well and virus control well in 0.1 ml MEM containing 0.18% NaHCO₃, 20 µg/ml trypsin, 2 µg/ml EDTA and

50 µg/ml gentamycin. Cell control and toxicity control wells received 0.1 ml of this same medium without virus.

Plates were incubated at 37° C in a humidified incubator with 5% CO₂, 95% air atmosphere. The cells were examined by microscopic observation for evidence of virus-specific cytopathic effect (CPE) and for morphological changes due to compound effect in non-infected toxicity controls. Virus CPE was graded on a scale of 0-4, with 4 being 95-100% CPE. The effective dose, 50% endpoint (ED₅₀) was calculated by regression analysis of the mean CPE grade at each concentration of the compound where activity was seen to bracket the CPE grade that was 50% of that seen in the virus controls. Visible changes in the morphology of cells in toxicity control wells were graded by microscopic observation, using a scale from no toxicity (0%) to complete destruction of the cells (100%) in 20% increments. The cytotoxic dose, 50%

- 40 -

endpoint (CD_{50}) was calculated by regression analysis of those toxicity grades bracketing the 50% endpoint, compared to the concentrations of compound used for those toxicity grades. A therapeutic index (TI) was calculated for each compound using the formula $TI = CD_{50}/ED_{50}$. These results are shown in Table VI, below.

TABLE VI
Anti-Influenza Activity Of Modified Oligonucleotides

<u>CMPD</u>	<u>ED₅₀ (μg/ml)</u>	<u>CD₅₀ (μg/ml)</u>	<u>TI</u>
A	Not Tested		
B	34.3	147	4.3
C	55	562	10
D	61	649	11
E	Not Tested		
F	178	261	1.5
G	147	649	4.4
H	59	562	9.5
I	45	750	17
J	175	750	4.3
K	22	825	38
L	32	422	13
M	23.7	383	16
Control ¹	440	422	<1.0

¹Control is oligodeoxynucleotide phosphorothioate having the sequence 5' CAGAGCAAATCATCAGAAGA 3'.

These results demonstrate that all of the preferred structural features of anti-influenza modified

- 41 -

oligonucleotides according to the invention, i.e., chimeric, hybrid, capped and self-stabilized features, are capable of improving the efficacy of antisense oligonucleotides in inhibiting influenza virus replication or propagation. These results further suggest that combinations of such structural features within an antisense oligonucleotide should have even greater efficacy.

EXAMPLE 11

Inhibition Of Various Strains Of Influenza Virus By Anti-influenza Modified Oligonucleotides

To test whether anti-fluenza modified oligonucleotides according to the invention can inhibit other strains of influenza virus, the experiment described in Example 10 was repeated, using compound M against various influenza strains. The influenza strains chosen for this study were the H1N1 strains A/NWS/33 and A/PR/8/34, the H3N2 strains A/Washington/897/80, A/Victoria/3/75 and A/Port Chalmers/1/73, and the H2N2 strain A2/Japan/305/57. The results were calculated as described in Example 10, and are shown in Table VII below.

- 42 -

TABLE VII
Effect Of Compound M On Various Strains of Influenza

<u>Influenza Strain</u>	<u>ED₅₀ (μg/ml)</u>	<u>CD₅₀ (μg/ml)</u>	<u>TI</u>
A/NWS/33	7.6	>100	>13
A/PR/8/34	>100	>100	?
A2/Japan/305/57	61	>100	>1.6
A/Washington/897/80	6	>100	17
A/Victoria/3/75	6.2	>100	>16
A/Port Chalmers/1/73	>100	>100	?

According to these results, compound M showed great efficacy in inhibiting three of the six strains of influenza tested and had some efficacy against a fourth strain. These results demonstrate that anti-influenza modified oligonucleotides according to the invention can be effective against multiple strains of influenza virus. To maximize the breadth of cross-strain efficacy, nucleotide sequences of various genes from several different influenza viruses can be compared and the most conserved nucleotide sequences can be used to prepare inhibitory oligonucleotides.

- 43 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: HYBRIDON, INC.
- (ii) TITLE OF INVENTION: Novel Anti-Influenza Oligonucleotides
- (iii) NUMBER OF SEQUENCES: 37
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Alleghetti & Witcoff, Ltd.
 - (B) STREET: Ten South Wacker Drive
 - (C) CITY: Chicago
 - (D) STATE: Illinois
 - (E) COUNTRY: U.S.A.
 - (F) ZIP: 60606
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Michael S. Greenfield
 - (B) REGISTRATION NUMBER: 37,142
 - (C) REFERENCE/DOCKET NUMBER: 93,161-A
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 312/715-1000
 - (B) TELEFAX: 617/715-1234

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 2149 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AGCGAAAGCA GGCAAACCAT TTGAATGGAT GTCAATCCGA CTTTACTTTT CTAAAAAGTG	60
CCAGCACAAA ATGCTATAAG CACAACTTTC CCTTATACTG GAGACCCTCC TTACAGCCAT	120
GGGACAGGAA CAGGATACAC CATGGATACT GTCAACAGGA CACATCAGTA CTCAGAAAGG	180

- 44 -

GGAAGATGGA	CAACAAACAC	CGAAACTGGA	GCACCGCAAC	TCAACCCGAT	TGATGGGCCA	240
CTGCCAGAAG	ACAATGAACC	AAGTGGTTAT	GCCCAAACAG	ATTGTGTATT	GGAAGCAATG	300
GCCTTCCTTG	AGGAATCCCA	TCCTGGTATC	TTTGAGACCT	CGTGTCTTGA	AACGATGGAG	360
GTTGTTTCAGC	AAACACGAGT	GGACAAGCTG	ACACAAGGCC	GACAGACCTA	TGACTGGACT	420
CTAAATAGGA	ACCAGCCTGC	TGCAACAGCA	TTGGCCAAAC	CAATAGAAGT	GTTTCAGATCA	480
AATGGCCTCA	CGGCCAATGA	ATCCGGAAGG	CTCATAGACT	TCCTTAAGGA	TGTAATGGAG	540
TCAATGAACA	AAGAAGAAAT	GGAGATCACA	ACTCATTTTC	AGAGAAAGAG	ACGAGTGAGA	600
GACAATATGA	CTAAGAAAAT	GGTGACACAG	AGAACAATAG	GTAAAAGGAA	GCAGAGATTG	660
AACAAAAGGA	GTTATCTAAT	TAGGGCATTG	ACCCTGAACA	CAATGACCAA	AGATGCTGAG	720
AGAGGGAAGC	TAAAACGGAG	AGCAATTGCA	ACCCAGGGA	TGCAAATAAG	GGGGTTTGTA	780
TACTTTGTTG	AGACACTAGC	AAGGAGTATA	TGTGAGAAAC	TTGAACAATC	AGGATTGCCA	840
GTTGGAGGCA	ATGAGAAGAA	AGCAAAGTTG	GCAAATGTTG	TAAGGAAGAT	GATGACCAAT	900
TCTCAGGACA	CTGAAATTTT	TTTCACATCA	CTGGAGATAA	CACCAAATGG	AACGAAAATC	960
AGAACCCTCG	GATGTTTTTT	GCCATGATCA	CATATATAAC	CAGAAATCAG	CCCGAATGGT	1020
TCAGAAATGT	TCTAAGTATT	GCTCCAATAA	TGTTCTCAAA	CAAATGGCG	AGACTGGGAA	1080
AGGGGTACAT	GTTTGAGAGC	AAGAGTATTA	AAATTAGAAC	TCAAATACCT	GCAGAAATGC	1140
TAGCAAGCAT	CGATTTGAAA	TACTTCAATG	ATTCAACTAG	AAAGAAGATT	GAAAAAATCC	1200
GGCCGCTCTT	AATAGATGGG	ACTGCATCAT	TGAGCCCTGG	AATGATGATG	GGCATGTTCA	1260
ATATGTTAAG	TACTGTATTA	GGCGTCTCCA	TCCTGAATCT	TGGACAAAAG	AGACACACCA	1320
AGACTACTTA	CTGGTGGGAT	GGTCTTCAAT	CTTCTGATGA	TTTTGCTCTG	ATTGTCAATG	1380
CACCCAATCA	TGAAGGGATT	CAAGCCGGAG	TCAACAGGTT	TTATCGAACC	TGTAAGCTAC	1440
TTGGAATTAA	TATGAGCAAG	AAAAAGTCTT	ACATAAACAG	AACAGGTACA	TTTGAATTCA	1500
CAAGTTTTTT	CTATCGTTAT	GGGTTTGTG	CCAATTTTCA	CATGGAGCTT	CCCAGCTTTG	1560
GGGTGTCTGG	GATCAACGAG	TCTGCGGACA	TGAGTATTGG	AGTTACTGTC	ATCAAAAACA	1620
ATATGATAAA	CAATGATCTT	GGTCCAGCAA	CCGCTCAAAT	GGCCCTTCAG	CTGTTCATCA	1680
AAGATTACAG	GTACACGTAC	CGCTGCCATA	GAGGTGACAC	ACAAATACAA	ACCCGAAGAT	1740
CATTTGAAAT	AAAGAAACTG	TGGGAGCAAA	CCCATTCCAA	AGCTGGACTG	CTGGTCTCCG	1800
ACGGAGGCCC	AAATTTATAC	AACATTAGAA	ATCTCCACAT	TCCTGAAGTC	TGCTTGAAAT	1860
GGGAATTAAT	GGATGAGGAT	TACCAGGGGC	GTTTATGCAA	CCCACTGAAC	CCATTTGTCA	1920
ACCATAAAGA	CATTGAATCA	GTGAACAATG	CAGTGATAAT	GCCAGCACAT	GGTCCAGCCA	1980
AAAACATGGA	GTATGATGCT	GTTGCAACAA	CACACTCCTG	GATCCCCAAA	AGAAATCGAT	2040
CCATCTTGAA	TACAAGCCAA	AGAGGAATAC	TTGAAGATGA	ACAAATGTAC	CAAAAGTGCT	2100

SUBSTITUTE SHEET (RULE 26)

- 45 -

GCAACTTATT TGAAAAATTC TTCCCCAGCA GTTCATACAG AAGACCAGT

2149

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CAGAGCAAAA TCATCAGAAG A

21

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CAGAGCAAAA TCATCAGAAG A

21

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CAGAGCAAAA TCATCAGAAG A

21

- 46 -

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CAGAGCAAAA TCATCAGAAG A

21

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CAGAGCAAAA TCATCAGAAG A

21

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 47 -

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CAGAGCAAAA TCATCAGAAG A

21

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CAGAGCAAAA TCATCAGAAG A

21

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CAGAGCAAAA TCATCAGAAG A

21

- 48 -

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CAGAGCAAAA TCATCAGAAG A

21

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CAGAGCAAAA TCATCAGAAG A

21

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 49 -

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CAGAGCAAAA TCATCAGAAG A

21

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

AGAGCAAAAT CATCAGAAG

19

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

AGAGCAAAAT CATCAGAAG

19

- 50 -

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GAGCAAAATC ATCAGAA

17

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 17 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GAGCAAAATC ATCAGAA

17

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 51 -

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CAGAGCAAAA TCATCAGAAG A

21

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CAGAGCAAAA TCATCAGAAG ATTCTGATGA

30

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CTTCCATAT TGAATATAAT

20

- 52 -

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

ACATCCATTC AAATGGTTTG

20

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

TCTTCCATTT TGGATCAGTA

20

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- 53 -

- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: YES
- (iv) ANTI-SENSE: NO
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GCCTTCATTT TGGTTGTTTT

20

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: YES
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GACGCCATGA TTTTGATGTC

20

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: YES
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GGATTCATTT TAAACCCCTG

20

- 54 -

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

AGACTCATCT TTCAATATCT

20

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GATAGAGAGA ACGTACGTTT

20

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 55 -

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

TCTGATAGGC CTGCAAATTT

20

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GGATCCATTA TGTCTTTGTC

20

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

CATGTCGGTT AGGTAACGCG

20

- 56 -

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: YES
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

GCAATCTACC TGAAAGCTTG

20

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: YES
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

AGCAGTATGT CCTGGAAGAG

20

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)

- 57 -

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

AAAACGACCT TGTTTCTACT

20

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

AAAAATGCCT TGTTCTACT

20

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

AAAAGTACCT TGTTTCTACT

20

- 58 -

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

AAAACACCCT TGTTTCTACT

20

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

AAAATACCCT TGTTTCTACT

20

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- 59 -

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

AAAAACTCCT TGTTTCTACT

20

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

AAACTACCT TGTTTCTACT

20

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

AAAACACCCT TGTTTCTACT

20

- 60 -

We claim:

1. An anti-influenza modified oligonucleotide comprising a nucleotide sequence that is complementary to an essential nucleic acid sequence of influenza virus, wherein the oligonucleotide is a mixed backbone chimeric oligonucleotide comprising a phosphorothioate or phosphorodithioate region and an alkylphosphonate region.

2. The oligonucleotide according to claim 1, wherein an alkylphosphonate region is at or near the 5' or 3' end of the oligonucleotide.

3. The oligonucleotide according to claim 1, wherein an alkylphosphonate region comprises from about 2 to about 10 contiguous nucleotides connected by alkylphosphonate linkages.

4. The oligonucleotide according to claim 2, wherein an alkylphosphonate region comprises from about 2 to about 10 contiguous nucleotides connected by alkylphosphonate linkages.

5. The oligonucleotide according to claim 1, wherein a phosphorothioate or phosphorodithioate region comprises from about 3 to about 100 contiguous nucleotides connected by phosphorothioate or phosphorodithioate linkages.

- 61 -

6. The oligonucleotide according to claim 2, wherein a phosphorothioate or phosphorodithioate region comprises from about 3 to about 100 contiguous nucleotides connected by phosphorothioate or phosphorodithioate linkages.

7. The oligonucleotide according to claim 3, wherein a phosphorothioate or phosphorodithioate region comprises from about 3 to about 100 contiguous nucleotides connected by phosphorothioate or phosphorodithioate linkages.

8. The oligonucleotide according to claim 4, wherein a phosphorothioate or phosphorodithioate region comprises from about 3 to about 100 contiguous nucleotides connected by phosphorothioate or phosphorodithioate linkages.

9. An anti-influenza modified oligonucleotide comprising a nucleotide sequence that is complementary to an essential nucleic acid sequence of influenza virus, wherein the oligonucleotide is a mixed backbone chimeric oligonucleotide comprising a phosphorothioate or phosphorodithioate region and an alkylphosphonothioate region.

10. The oligonucleotide according to claim 10, wherein an alkylphosphonothioate region is at or near the 5' or 3' end of the oligonucleotide.

- 62 -

11. The oligonucleotide according to claim 9, wherein an alkylphosphonothioate region comprises from about 2 to about 10 contiguous nucleotides connected by alkylphosphonothioate linkages.

12. The oligonucleotide according to claim 10, wherein an alkylphosphonothioate region comprises from about 2 to about 10 contiguous nucleotides connected by alkylphosphonothioate linkages.

13. The oligonucleotide according to claim 9, wherein a phosphorothioate or phosphorodithioate region comprises from about 3 to about 100 contiguous nucleotides connected by phosphorothioate or phosphorodithioate linkages.

14. The oligonucleotide according to claim 10, wherein a phosphorothioate or phosphorodithioate region comprises from about 3 to about 100 contiguous nucleotides connected by phosphorothioate or phosphorodithioate linkages.

15. The oligonucleotide according to claim 11, wherein a phosphorothioate or phosphorodithioate region comprises from about 3 to about 100 contiguous nucleotides connected by phosphorothioate or phosphorodithioate linkages.

- 63 -

16. The oligonucleotide according to claim 12, wherein a phosphorothioate or phosphorodithioate region comprises from about 3 to about 100 contiguous nucleotides connected by phosphorothioate or phosphorodithioate linkages.

17. An anti-influenza modified oligonucleotide comprising a nucleotide sequence that is complementary to an essential nucleic acid sequence of influenza virus, wherein the oligonucleotide is a hybrid oligonucleotide comprising a deoxyribonucleotide region and a ribonucleotide region.

18. The oligonucleotide according to claim 17, further comprising from about 1 to about all phosphorothioate or phosphorodithioate internucleotide linkages.

19. The oligonucleotide according to claim 17, wherein a ribonucleotide region is at or near the 5' or 3' end of the oligonucleotide.

20. The oligonucleotide according to claim 17, wherein a ribonucleotide region comprises from about 2 to about 100 contiguous ribonucleotides.

- 64 -

21. The oligonucleotide according to claim 18, wherein a ribonucleotide region comprises from about 2 to about 100 contiguous ribonucleotides.

22. The oligonucleotide according to claim 19, wherein a ribonucleotide region comprises from about 2 to about 100 contiguous ribonucleotides.

23. The oligonucleotide according to claim 17, wherein a deoxyribonucleotide region comprises from about 0 to about 100 deoxyribonucleotides.

24. The oligonucleotide according to claim 18, wherein a deoxyribonucleotide region comprises from about 0 to about 100 deoxyribonucleotides.

25. The oligonucleotide according to claim 19, wherein a deoxyribonucleotide region comprises from about 0 to about 100 deoxyribonucleotides.

26. The oligonucleotide according to claim 20, wherein a deoxyribonucleotide region comprises from about 0 to about 100 deoxyribonucleotides.

- 65 -

27. The oligonucleotide according to claim 21, wherein a deoxyribonucleotide region comprises from about 0 to about 100 deoxyribonucleotides.

28. The oligonucleotide according to claim 22, wherein a deoxyribonucleotide region comprises from about 0 to about 100 deoxyribonucleotides.

29. An anti-influenza modified oligonucleotide comprising a nucleotide sequence that is complementary to an essential nucleic acid sequence of influenza virus, wherein the oligonucleotide has at its 5' and/or 3' end a nuclease resistance conferring cap structure selected from the group consisting of the cap structures shown in Figure 4 and lower alkyl (C1-C12) or alcohol groups, and wherein the oligonucleotide has from 1 to about all modified internucleotide linkages selected from the group consisting of phosphotriester, phosphoramidate, siloxane, carbonate, carboxymethylester, acetamidate, carbamate, thioether, bridged phosphoramidate, bridged methylene phosphonate, bridged phosphorothioate, sulfone, phosphorothioate and phosphorodithioate linkages.

- 66 -

30. An anti-influenza modified oligonucleotide comprising an anti-influenza hybridizing region and a self-complementary region.

31. The oligonucleotide according to claim 30, wherein the influenza hybridizing region comprises from about 6 to about 100 nucleotides that are complementary to an essential nucleic acid sequence of influenza virus.

32. The oligonucleotide according to claim 30, wherein the self-complementary region comprises from about 4 to about 50 nucleotides that form intramolecular base pairs.

33. The oligonucleotide according to claim 30, wherein a self-complementary region is at or near the 5' or 3' end of the oligonucleotide.

34. The oligonucleotide according to claim 31, wherein a self-complementary region is at or near the 5' or 3' end of the oligonucleotide.

35. The oligonucleotide according to claim 32, wherein a self-complementary region is at or near the 5' or 3' end of the oligonucleotide.

- 67 -

36. The oligonucleotide according to claim 30, wherein from about 2 to about all nucleotides are coupled by phosphorothioate and/or phosphorodithioate internucleotide linkages.

37. The oligonucleotide according to claim 31, wherein from about 2 to about all nucleotides are coupled by phosphorothioate and/or phosphorodithioate internucleotide linkages.

38. The oligonucleotide according to claim 32, wherein from about 2 to about all nucleotides are coupled by phosphorothioate and/or phosphorodithioate internucleotide linkages.

39. The oligonucleotide according to claim 33, wherein from about 2 to about all nucleotides are coupled by phosphorothioate and/or phosphorodithioate internucleotide linkages.

40. The oligonucleotide according to claim 34, wherein from about 2 to about all nucleotides are coupled by phosphorothioate and/or phosphorodithioate internucleotide linkages.

- 68 -

41. The oligonucleotide according to claim 35, wherein from about 2 to about all nucleotides are coupled by phosphorothioate and/or phosphorodithioate internucleotide linkages.

42. The oligonucleotide according to claim 1, wherein the essential nucleic acid sequence is selected from the group consisting of influenza polymerase 3 gene, influenza polymerase 1 gene, influenza polymerase 2 gene, influenza hemagglutinin gene, influenza nucleoprotein gene, influenza neuraminidase gene, influenza matrix protein gene, influenza left or right splice junctions of segments 7 or 8, influenza splice branch of segment 8 and influenza packaging sequences of segment 1, 2, 3, 4, 5, 6, 7 or 8.

43. The oligonucleotide according to claim 9, wherein the essential nucleic acid sequence is selected from the group consisting of influenza polymerase 3 gene, influenza polymerase 1 gene, influenza polymerase 2 gene, influenza hemagglutinin gene, influenza nucleoprotein gene, influenza neuraminidase gene, influenza matrix protein gene, influenza left or right splice junctions of segments 7 or 8, influenza splice branch of segment 8 and influenza packaging sequences of segment 1, 2, 3, 4, 5, 6, 7 or 8.

- 69 -

44. The oligonucleotide according to claim 17, wherein the essential nucleic acid sequence is selected from the group consisting of influenza polymerase 3 gene, influenza polymerase 1 gene, influenza polymerase 2 gene, influenza hemagglutinin gene, influenza nucleoprotein gene, influenza neuraminidase gene, influenza matrix protein gene, influenza left or right splice junctions of segments 7 or 8, influenza splice branch of segment 8 and influenza packaging sequences of segment 1, 2, 3, 4, 5, 6, 7 or 8.

45. The oligonucleotide according to claim 29, wherein the essential nucleic acid sequence is selected from the group consisting of influenza polymerase 3 gene, influenza polymerase 1 gene, influenza polymerase 2 gene, influenza hemagglutinin gene, influenza nucleoprotein gene, influenza neuraminidase gene, influenza matrix protein gene, influenza left or right splice junctions of segments 7 or 8, influenza splice branch of segment 8 and influenza packaging sequences of segment 1, 2, 3, 4, 5, 6, 7 or 8.

46. The oligonucleotide according to claim 31, wherein the essential nucleic acid sequence is selected from the group consisting of influenza polymerase 3 gene, influenza polymerase 1 gene, influenza polymerase 2 gene, influenza hemagglutinin gene, influenza nucleoprotein gene, influenza

- 70 -

neuraminidase gene, influenza matrix protein gene, influenza left or right splice junctions of segments 7 or 8, influenza splice branch of segment 8 and influenza packaging sequences of segment 1, 2, 3, 4, 5, 6, 7 or 8.

1/8

1 AGCGAAAGCATGGCAUACCTTATTGATGGAT GTCAATCCGA CTTTACTTTT
51 CTTAAAAGTG CCAGCACAAA ATGCTATAAG CACAACTTTC CCTTATACTG
101 GAGACCTCC TTACAGCCAT GGGACAGGA CAGGATACAC CATGGATACT
151 GTCAACAGGA CACATCAGTA CTCAGAAAGG GGAAGATGGA CAACAAACAC
201 CGAAACTGGA GCACCGCAAC TENACCCGAT TGATGGGCCA CTGCCAGAAG
251 ACAATGAACC AAGTGGTTAT GCCCAAACAG ATTGTGTATT GGAAGCAATG
301 GCCTTCCTTG AGGAATCCCA TCCTGGTATC TTTGAGACCT CGTGTCTTGA
351 AACGATGGAG GTTGTTCAGC AACACGAGT GGACAAGCTG ACACAAGGCC
401 GACAGACCTA TGACTGGACT CTAATAGGA ACCAGCCTGC TGCAACAGCA
451 TTGGCCAACA CAATAGAAGT GTTCAGATCA AATGGCCTCA CGGCCAATGA
501 ATCCGGAAGG CTCATAGACT TCCTTAAGGA TGTAAATGGAG TCAATGAACA
551 AAGAAGAAAT GGAGATCACA ACTCATTTTC AGAGAAAGAG ACGAGTGAGA
601 GACAAATATGA CTAAGAAAAT GGTGACACAG AGAACAAATAG GTAAAAGGAA
651 GCAGAGATTG AACAAAAGGA GTTATCTAAT TAGGGCATTG ACCCTGAACA
701 CAATGACCAA AGATGCTGAG AGAGGGAAGC TAAACGGAG AGCAATTGCA
751 ACCCCAGGGA TGCAATAAG GGGGTTTGTA TACTTTGTTG AGACACTAGC
801 AAGGAGTATA TGTGAGAAAC TTGAACAATC AGGATTGCCA GTTGGAGGCA
851 ATGAGAAGAA AGCAAAGTTG GCAAATGTTG TAAGGAAGAT GATGACCAAT
901 TCTCAGGACA CTGAAATTC TTTCACCATC ACTGGAGATA ACACCAAATG
951 GAACGAAAAT CAGAACCTC GGATGTTTTT GGCCATGATC ACATATATAA
1001 CCAGAAATCA GCCCGAATGG TTCAGAAATG TTCTAAGTAT TGCTCCAATA
1051 ATGTTCTCAA ACAAATGGC GAGACTGGGA AAGGGGTACA TGTTTGAGAG


Fig. 1

2 / 8

1101 CAAGAGTATT AAAATTAGAA CTCAAATACC TGCAGAAATC CTAGCAAGCA
1151 TCGATTTGAA ATACTTCAAT GATTCAACTA GAAAGAAGAT TGA AAAAATC
1201 CGGCCGCTCT TAATAGATGG GACTGCATCA TTGAGCCCTG GAATGATGAT
1251 GGGCATGTTT ~~AAATATGTTAA~~ GTACTGTATT AGGCGTCTCC ATCCTGAATC
1301 TTGGACAAAA GAGACACACC AAGACTACTT ACTGGTGGGA TGGTCTTCAA
1351 ~~TCTTCTGATGATTTTGCTCTGATTGTGAT~~ GCACCCATC ATGAAGGGAT
1401 TCAAGCCGGA GTCAACAGGT TTTATCGAAC CTGTAGCTA CTTGGAATTA
1451 ATATGAGCAA GAAAAGTCT TACATAACA GAACAGGTAC ATTTGAATTC
1501 ACAAGTTTTT TCTATCGTTA TGGGTTTGT GCCAATTCA GCATGGAGCT
1551 TCCCAGCTTT GGGGTGTCG GGATCAACA GTCTGCGGAC ATGAGTATTG
1601 GAGTTACTGT CATCAAAAC AATATGATA ACAATGATCT TGGTCCAGCA
1651 ACCGCTCAA TGGCCCTTCA GCTGTTCATC AAGATTACA GGTACACGTA
1701 CCGGTGCCAT AGAGGTGACA CACAAATACA AACCAGAGA TCATTTGAAA
1751 TAAAGAACT GTGGGAGCAA ACCCATTCOA AAGCTGGACT GCTGGTCTCC
1801 GACGGAGGCC CAAATTTATA CAACATTAGA AATCTCCACA TTCCTGAAGT
1851 CTGCTTGAAA TGGGAATTAA TGGATGAGGA TTACCAGGGG CGTTTATGCA
1901 ACCCACTGAA CCCATTTGTC AACCATAAAG ACATTGAATC AGTGAACAAT
1951 GCAGTGATAA TGCCAGCACA TGGTCCAGCC AAAAACATGG AGTATGATGC
2001 TGTTCGAACA ACACACTECT GGATCCCCAA AAGAAATCGA TCCATCTTGA
2051 ATACAAGCCA AAGAGGAATA CTTGAAGATG AACAATGTA CCAAAGTGC
2101 TGCAACTTAT TTGAAAATT CTTCCCCAGC AGTTCATACA GAAGACCAGT

Fig. 1
(cont'd)

3/8

**Fig. 2**




Fig. 3

5 / 8





Fig. 4

6 / 8

*Fig. 5*

7/8




Fig. 6

8 / 8




Fig. 7