

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 2094872 T4**

(12) Oversættelse af ændret
europæisk patentskrift

-
- (51) Int.Cl.: **A 61 K 39/12 (2006.01)** **A 61 K 39/00 (2006.01)** **A 61 K 49/00 (2006.01)**
C 12 N 7/00 (2006.01) **C 12 Q 1/70 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2020-05-18**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om opretholdelse af patentet i ændret form: **2020-02-19**
- (86) Europæisk ansøgning nr.: **07869309.0**
- (86) Europæisk indleveringsdag: **2007-12-14**
- (87) Den europæiske ansøgnings publiceringsdag: **2009-09-02**
- (86) International ansøgning nr.: **US2007087628**
- (87) Internationalt publikationsnr.: **WO2008076915**
- (30) Prioritet: **2006-12-15 US 870311 P**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR**
- (73) Patenthaver: **Boehringer Ingelheim Animal Health USA Inc., 3239 Satellite Blvd, Duluth, GA 30096, USA**
- (72) Opfinder: **FACHINGER, Vicky, Bismarckstrasse 13, 65812 Bad Soden, Tyskland**
PIONTKOWSKI, Michael, 4114 Pawnee Road, Perry, KS 66073, USA
ELBERS, Knut, Caprino-Veronese-Str. 4a, 55435 Gau Algesheim, Tyskland
LISCHEWSKI, Axel, Boehringer Ingelheim GmbH, CD Patents, Binger Strasse 173, 55216 Ingelheim an Rhein, Tyskland
KIXMOELLER, Marion, Mariahilfstr. 1, 81541 München, Tyskland
ORVEILLON, Francois-Xavier, Neutorstrasse 17, 55116 Mainz, Tyskland
FREIIN VON RICHTHOFEN, Isabelle, Ordrupvej 171, 2th, 2920 Charlottenlund, Tyskland
- (74) Fuldmægtig i Danmark: **AWA Denmark A/S, Strandgade 56, 1401 København K, Danmark**
- (54) Benævnelse: **Behandling af anti-PCV2-antistofseropositive svin med PCV2-antigen**
- (56) Fremdragne publikationer:
BLANCHARD ET AL.: 'Protection of swine against post-weanling multisystemic wasting syndrome (PMWS) by porcine circovirus type 2 (PCV2) proteins' VACCINE vol. 21, 2003, pages 4565 - 4575, XP004467337
FENAUX ET AL.: 'A Chimeric Porcine Circovirus (PCV) with the Immunogenic Capsid Gene of the Pathogenic PCV Type 2 (PCV2) Cloned into the Genomic Backbone of the Nonpathogenic PCV I Induces Protective Immunity against PCV2 Infection in Pigs' JOURNAL OF VIROLOGY vol. 78, no. 12, June 2004, pages 6297 - 6303, XP008039321
EP-A1- 1 926 496
WO-A2-2005/009462
WO-A2-2006/072065
WO-A2-2008/076915
US-B1- 6 703 023

Fortsættes ...

Assignment of inventors to Boehringer Ingelheim Vetmedica Inc.
 IFA_tech_sheet
 IFA_ihc_guidebook_5th_edition_chapter_10
 Robert Desrosiers; Et Al: "Preliminary results with IngelvacÂ® CircoFLEX to protect multiple ages of Quebec pigs against PCVAD", American Association of swine Veterinarians, 1 January 2007 (2007-01-01), pages 143-146, XP055446462,
 Proof of date AASV 2007
 CircoFLEX Symposium 2007
 Urniza et al: "DOI of a PCV2 vaccine", IPVS 2006,
 Proof of date IPVS 2006
 FENAUX M; ET AL: "A CHIMERIC PORCINE CIRCOVIRUS (PCV) WITH THE IMMUNOGENIC CAPSID GENE FOR THE PATHOGENIC PCV TYPE 2 (PCV2) CLONED INTO THE GENOMIC BACKBONE OF THE NONPATHOGENIC PCV1 INDUCES PROTECTIVE IMMUNITY AGAINST PCV2 INFECTION IN PIGS", JOURNAL OF VIROLOGY, vol. 78, no. 12, 1 June 2004 (2004-06-01), pages 6297-6303, XP008039321,
 Thomas et al: "Effect of MDA on a PCV2 vaccine", AASV 2005,
 Segales et al: "Epidemiology of PCV2", IPVS 2004,
 Allen et al. in J Vet Diagn Invest 2000
 EMA document regarding Ingelvac CircoFLEX, 2008
 minutes of oral proceedings held 29.09.2015
 Minutes oral proceedings T1415/16
 Blanchard et al: Vaccine, vol. 21, 2003, pages 4565-4575,
 Opriessnig et al: "Results of Suvaxyn PCV2 One shot in MDA positive piglets", AASV 2007,
 Pages of opposition filed against EP1926496 by Boehringer Ingelheim on 26 march 2014
 Vaccine 21 (2003) 3406-3412, review article
 Curriculum Vitae Dr. Urs Bruderer
 Vaccine 2008, 26, 1488-1499
 proof of publication date D41
 Veterinary Microbiology 168 (2014) 272-280
 front page and page 10 of EP 2371385 A1
 front page and page 28 of EP 2275132 A2
 declaration prof. Savelkoul
 Curriculum Vitae prof. Savelkoul
 Report Dr. Urs Bruderer
 Applicant's letter 25.08.2015
 CircoFLEX USDA license
 CircoFLEX label
 MSDS CircoFLEX
 Boehringer Ingelheim press release
 Boehringer Ingelheim Health Seminar
 Charreyre, IPVS congress 2000
 Rodriguez, AJVR, 2002
 Meerts, Int symposium, 2003
 Fort Dodge News Release
 Interlocutory decision EP 1104244
 Preliminary opinion on EP application 05777317
 Preliminary opinion on EP application 15165133.8
 Preliminary opinion on EP application 08756672.5
 Preliminary opinion on EP application 06720702.7
 Preliminary opinion by the Board in T0239/16
 Decision on EP 1951304
 Declaration Prof. Saalmuller
 Kixmoller et al: Vaccine, vol. 26, 2008, pages 3443-3451,
 Declaration Dr. Gassel
 PCV2 study on various adjuvants
 Overdose study
 Excerpts of submission by the opponent on case EP1926496
 WO-A-99/29871
 WO-A-03/049703
 WO-A-2005/092069
 WO-A-2006/072065
 WO-A-2007/094893

WO-A-2008/073464

WO-A1-2007/028823

WO-A2-2005/092069

US-B1- 6 368 601

OSTANELLO F ET AL: "Experimental infection of 3-week-old conventional colostrum-fed pigs with porcine circovirus type 2 and porcine parvovirus" VETERINARY MICROBIOLOGY, ELSEVIER BV, NL, vol. 108, no. 3-4, 1 July 2005 (2005-07-01), pages 179-186, XP004933318 ISSN: 0378-1135

MCKEOWN N E ET AL: "Effects of porcine circovirus type 2 (PCV2) maternal antibodies on experimental infection of piglets with PCV2." CLINICAL AND DIAGNOSTIC LABORATORY IMMUNOLOGY NOV 2005, vol. 12, no. 11, November 2005 (2005-11), pages 1347-1351, XP002557104 ISSN: 1071-412X

BOEHRINGER INGELHEIM VETMEDICA ET AL: "IngelvacRCircoFLEXTM. Material safety data sheet" INTERNET CITATION, [Online] 23 October 2006 (2006-10-23), XP002433251 Retrieved from the Internet:

URL:http://www.bimsds.us/msds/IngelvacCirc oFlex_msds.pdf [retrieved on 2007-05-10]

CHAE C: "A review of porcine circovirus 2-associated syndromes and diseases" VETERINARY JOURNAL, BAILLIERE TINDALL, LONDON, GB, vol. 169, no. 3, 1 May 2005 (2005-05-01), pages 326-336, XP004858799 ISSN: 1090-0233

SEGALÉS JOAQUIM ET AL: "Pathological findings associated with naturally acquired porcine circovirus type 2 associated disease." VETERINARY MICROBIOLOGY 4 FEB 2004, vol. 98, no. 2, 4 February 2004 (2004-02-04), pages 137-149, XP002557105 ISSN: 0378-1135

KAMSTRUP S ET AL: "Immunisation against PCV2 structural protein by DNA vaccination of mice" VACCINE, BUTTERWORTH SCIENTIFIC. GUILDFORD, GB, vol. 22, no. 11-12, 29 March 2004 (2004-03-29), pages 1358-1361, XP004500378 ISSN: 0264-410X

BLANCHARD ET AL.: 'Protection of swine against post-weanling multisystemic wasting syndrome (PMWS) by porcine circovirus type 2 (PCV2) proteins' VACCINE vol. 21, 2003, pages 4565 - 4575, XP004467337

FENAUX ET AL.: 'A Chimeric Porcine Circovirus (PCV) with the Immunogenic Capsid Gene of the Pathogenic PCV Type 2 (PVC2) Cloned into the Genomic Backbone of the Nonpathogenic PCV1 Induces Protective Immunity against PCV2 Infection in Pigs' JOURNAL OF VIROLOGY vol. 78, no. 12, June 2004, pages 6297 - 6303, XP008039321

Description

SEQUENCE LISTING

5 **[0001]** This application contains a sequence listing in paper format and in computer readable format the teachings and content of which are hereby incorporated by reference. The sequence listing is also identical with that incorporated in WO06/072065.

BACKGROUND OF THE INVENTION

Field of the Invention

10 **[0002]** The present invention relates to the use of an immunogenic composition comprising a porcine circovirus type 2 (PCV2) antigen for the prevention, reduction in severity of clinical signs, reduction in the incidence of infection and/or clinical signs, and treatment of several clinical manifestations (diseases) in animals having anti-PCV2 specific antibodies. Preferably, those anti-PCV-2 specific antibodies are maternal antibodies.

Description of the Prior Art

20 **[0003]** Porcine circovirus type 2 (PCV2) is a small (17 - 22 nm in diameter), icosahedral, non-enveloped DNA virus, which contains a single-stranded circular genome. PCV2 shares approximately 80% sequence identity with porcine circovirus type 1 (PCV1). However, in contrast with PCV1, which is generally non-virulent, infection of swine with PCV2 has recently associated with a number of disease syndromes which have been collectively named Porcine Circovirus-Associated Diseases (PCVAD) (also known as Porcine Circovirus Diseases (PCVD)) (Allan et al. 2006, IPVS Congress).
 25 Postweaning Multisystemic Wasting Syndrome (PMWS) is generally regarded to be the major clinical manifestation of PCVAD. (Harding et al., 1997, Swine Health Prod; 5: 201-203; Kennedy et al., 2000, J Comp Pathol; 122: 9-24). PMWS affects pigs between 5-18 weeks of age. PMWS is clinically characterized by wasting, paleness of the skin, unthriftiness, respiratory distress, diarrhea, icterus, and jaundice. In some affected swine, a combination of all symptoms will be apparent while other affected swine will only have one or two of these symptoms. (Muirhead, 2002, Vet. Rec.; 150: 456)
 30 During necropsy, microscopic and macroscopic lesions also appear on multiple tissues and organs, with lymphoid organs being the most common site for lesions. (Allan and Ellis, 2000; J Vet. Diagn. Invest. 12: 3-14) A strong correlation has been observed between the amount of PCV2 nucleic acid or antigen and the severity of microscopic lymphoid lesions. Mortality rates for swine infected with PCV2 can approach 80%. In addition to PMWS, PCV2 has been associated with several other infections including pseudorabies, porcine reproductive and respiratory syndrome (PRRS), Glasser's disease, streptococcal meningitis, salmonellosis, postweaning colibacillosis, dietetic hepatitis, and suppurative bronchopneumonia. However, research thus far has not confirmed whether any of these clinical symptoms are in fact, the direct result of a PCV2 infection. Moreover, it is not yet known whether any of these clinical symptoms can be effectively reduced or cured by an active agent directed against PCV2.

35 **[0004]** Approaches to treat PCV2 infections based on a DNA vaccine are described in U.S. Patent No. 6,703,023. In WO03/049703 production of a live chimeric vaccine is described, comprising a PCV-1 backbone in which an immunogenic gene of a pathogenic PCV2 strains replaces a gene of the PCV-1 backbone. WO99/18214 has provided several PCV2 strains and procedures for the preparation of a killed PCV2 vaccine. However, no efficacy data have been reported. An effective ORF-2 based subunit vaccine has been reported in WO06/072065 and in WO2007/028823. Any of such vaccines are intended to be used for the vaccination/treatment of swine or pigs older than 3 weeks of age. None of these vaccines have been described for use in young piglets, younger than 3 or 2 weeks of age.

40 **[0005]** Maternally derived immunity has been shown to confer a certain degree of protection against PCV2 infection and clinical diseases associated with PCV2 infections. This protection has been shown to be titer dependent: higher titers are generally protective whereas lower titers are not (McKeown et al., 2005; Clin. Diagn. Lab. Immunol.; 12: 1347-1351). The mean antibody half-life in weanlings has been estimated to be 19.0 days and the window for PCV2-passive antibody decay within a population is relatively wide (Opriessnig et al. 2004, J. Swine Health Prod. 12:186-191). Low titers of PCV2 passively acquired antibodies present at 10-12 days of age were found to decay by approximately 4.9 $\pm$ 1.2 weeks of age, moderate levels of antibodies were found to decay by approximately 8.1 $\pm$ 1.9 weeks of age and high levels of antibodies were found to decay by approximately 11.1 $\pm$ 2.5 weeks of age (Opriessnig et al., 2006, 37th Annual Meeting of the American Association of Swine Veterinarians). In a timely close correlation with the waning antibody titer stands the occurrence of first clinical signs of PCVAD which occur when piglets are approximately 5 and 12 weeks old (Allan et al, 2000, Vet. Diagn. Investigation, 12: 3-14). Furthermore, PCV2 has also been isolated out of lymphnodes of neonatal piglets (Hirai et al, 2001, Vet. Record; 148:482-484) indicating that even younger piglets may be affected from PCVAD in the absence of protective maternal antibody titers. The obvious correlation between the

antibody titer and protection has been proven in a Spanish Field study: Pigs with low antibody titers at 7 weeks of age (mean antibody titer 1:100, range 0 to 1:320) had a significantly higher mortality rate over the following 5 weeks than animals with higher antibody titers (Rodriguez-Arrioja et al., 2002, Am. J Vet. Res. 63:354-357).

[0006] The presence of maternally-derived antibody not only may confer a certain degree of protection against viral infections, which however is not predictable, but also be known to impair the efficacy of immunization. For example higher titers of maternally-derived antibodies to classical swine fever virus (CSFV) inhibit both cell-mediated and humoral immune response to a CSFV vaccine, but lower titers have no significant influence (Suradhat and Damrongwatanapokin, 2003, Vet. Microbiol; 92: 187-194). Also, for live PCV2 vaccines, it has been predicted that they will work most efficiently when given to piglets older than 7 or 8 weeks of age, because the maternal antibodies have mostly waned at that time. Maternal antibody interference is influenced by the type of elicited immune response (Th1 versus Th2) which is dependent (beyond others) on the type of vaccine, type of antigen, type of adjuvant as well as on the amount of administered antigen. Consequently, possible maternal antibody interference may differ for vaccines even if they protect against the same pathogen. Altogether, maternally-derived anti-PCV2 antibodies may confer a certain degree of protection against PCV2, but on the other hand those antibodies may impair the efficacy of any PCV2 vaccine.

[0007] The protection of animals by active immunization is further complicated by the fact that a) the time for the decay of maternally derived antibodies (MDA) varies from animal to animal and b) many diseases occur shortly after the decay of antibodies. To face this problem several vaccination strategies foresee a two shot vaccination regime for young animals: The first vaccination is given early in life in order to protect those animals with low MDA. It is accepted that this first vaccination may not be effective in animals with high MDA titers due to an interference with the vaccine antigen. In order to also protect these animals, a second vaccination is required, when high MDA levels are expected to have declined. This kind of vaccination schedule is used for many small animal vaccines (against e.g. canine parvovirus, canine hepatitis, etc.), equine vaccines (against e.g. equine influenza vaccines) and porcine vaccines (against e.g. *Actinobacillus pleuropneumoniae*, *Haemophilus parasuis*). As the onset of PCVAD in animals 5 weeks of age or older seems to be linked to the decay of PCV2 antibodies, which is reported to occur in animals aged 4-11 weeks, several vaccine approaches against PCVAD have been described using a two shot vaccination regime in order to circumvent a possible maternal antibody interference. In WO 2007/028823 vaccination of piglets having maternally-derived anti-PCV2 antibodies with more than 20 µg/dose antigen using a two shot vaccination regime is described. Initial vaccination was administered between 1 and 4 weeks of age. All animals were re-vaccinated three weeks after the initial vaccination, when the maternally-derived antibodies in animals with high MDA levels at the time of first vaccination had declined or ceased. Thus, yet no information exist which describes the exact influence of maternally-derived anti-PCV2 antibodies on degree of protection or interference. For that reason, it is recommended not to vaccinate piglets prior to three (3) weeks of age at least with a single shot vaccine regime. Vaccination prior to weeks 3 of age is connected with a certain degree of uncertainty with respect to immunization efficacy. On the other hand, piglets with lower levels of maternally-derived anti-PCV2 antibodies, whereas yet nobody knows what lower levels exactly means, are not sufficiently protected against PCV2 infection prior to week 3 of age. In other words, herds with low MDA titers which are not vaccinated before 3 weeks of age have an immanent risk of PCV2 infections due to lack of a sufficient immune status.

[0008] Moreover, such vaccines have not been described to confer protective immunity against PCV2 infection or reducing, lessening the severity of, lessening the incidence of, or curing any clinical symptoms associated therewith in pigs already having anti-PCV2 antibodies, preferably having maternal anti-PC2 antibodies.

BRIEF DESCRIPTION OF THE DRAWINGS

[0009]

Figure 1 is a graph of anti-PCV2 antibody titer classes at the time of vaccination;

Fig. 2 is a graph comparing the live body weight in vaccinated animals with low (< 1:100) and high (> 1:1000) anti-PCV2 antibodies; and

Fig. 3 is a graph illustrating body weight difference in vaccinated (IVP) as compared to placebo-treated control animals (CP).

DISCLOSURE OF THE INVENTION

[0010] The claims define the matter for which protection is sought. The present invention overcomes the problems inherent in the prior art and provides a distinct advance in the state of the art. According to general aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in animals having anti-PCV2 antibodies, comprising the step of administering an effective amount of PCV2 antigen or an immunogenic composition comprising an PCV2 antigen to that animal in need of such treatment. It was an unpredictable and surprising finding, that the presence of anti-PCV2 antibodies,

and in particular of maternal origin, does not impair the efficacy of vaccine comprising PCV2 antigen.

[0011] The terms "vaccine" or "immunogenic composition" (both terms are used synonymously) as used herein refer to any pharmaceutical composition containing a PCV2 antigen, which composition can be used to prevent or treat a PCV2 infection-associated disease or condition in a subject. A preferred immunogenic composition can induce, stimulate or enhance the immune response against PCV2. The term thus encompasses subunit immunogenic compositions, as described below.

[0012] Thus according to another aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in animals having anti-PCV2 antibodies, in particular maternally-derived anti-PCV2 antibodies, comprising the step of administering an effective amount of PCV2 antigen or an immunogenic composition comprising an PCV2 antigen to that animal in need of such treatment, wherein the immunogenic composition is a subunit immunogenic composition, a compositions containing whole killed, or attenuated and/or inactivated PCV2.

[0013] The term "subunit immunogenic composition" as used herein refers to a composition containing at least one immunogenic polypeptide or antigen, but not all antigens, derived from or homologous to an antigen from PCV2. Such a composition is substantially free of intact PCV2. Thus, a "subunit immunogenic composition" is prepared from at least partially purified or fractionated (preferably substantially purified) immunogenic polypeptides from PCV2, or recombinant analogs thereof. A subunit immunogenic composition can comprise the subunit antigen or antigens of interest substantially free of other antigens or polypeptides from PCV2, or in fractionated from. A preferred immunogenic subunit composition comprises the PCV2 ORF-2 protein as described below. Most preferred are immunogenic subunit compositions, which comprise any of the PCV2 antigens provided in WO06/072065. An "immune response" means but is not limited to the development in a host of a cellular and/or antibody-mediated immune response to the composition or vaccine of interest. Usually, an "immune response" includes but is not limited to one or more of the following effects: the production or activation of antibodies, B cells, helper T cells, suppressor T cells, and/or cytotoxic T cells, directed specifically to an antigen or antigens included in the composition or vaccine of interest. Preferably, the host will display either a therapeutic or a protective immunological (memory) response such that resistance to new infection will be enhanced and/or the clinical severity of the disease reduced. Such protection will be demonstrated by either a reduction in number or severity of, or lack of one or more of the symptoms associated with PCV2 infections, in delay of onset of viremia, in a reduced viral persistence, in a reduction of the overall viral load and/or a reduction of viral excretion.

[0014] The terms "antigen" as used herein refers to an amino acid sequence which elicits an immunological response as described above. An antigen, as used herein, includes the full-length sequence of any PCV2 proteins, analogs thereof, or immunogenic fragments thereof. The term "immunogenic fragment" refers to a fragment of a protein which includes one or more epitopes and thus elicits the immunological response described above. Such fragments can be identified using any number of epitope mapping techniques, well known in the art. See, e.g., *Epitope Mapping Protocols in Methods in Molecular Biology*, Vol. 66 (Glenn E. Morris, Ed., 1996) Humana Press, Totowa, New Jersey. For example, linear epitopes may be determined by e.g., concurrently synthesizing large numbers of peptides on solid supports, the peptides corresponding to portions of the protein molecule, and reacting the peptides with antibodies while the peptides are still attached to the supports. Such techniques are known in the art and described in, e.g., U.S. Patent No. 4,708,871; Geysen et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:3998-4002; Geysen et al. (1986) *Molec. Immunol.* 23:709-715. Similarly, conformational epitopes are readily identified by determining spatial conformation of amino acids such as by, e.g., x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, e.g., *Epitope Mapping Protocols*, *supra*.

[0015] Synthetic antigens are also included within the definition, for example, polyepitopes, flanking epitopes, and other recombinant or synthetically derived antigens. See, e.g., Bergmann et al. (1993) *Eur. J. Immunol.* 23:2777-2781; Bergmann et al. (1996), *J. Immunol.* 157:3242-3249; Suhrbier, A. (1997), *Immunol. and Cell Biol.* 75:402-408; Gardner et al., (1998) 12th World AIDS Conference, Geneva, Switzerland, June 28-July 3, 1998.

[0016] According to further aspect, the immunogenic composition as used herein most preferably comprises the polypeptide, or a fragment thereof, expressed by ORF-2 of PCV2. PCV2 ORF-2 DNA and protein, used herein for the preparation of the compositions and within the processes provided herein is a highly conserved domain within PCV2 isolates and thereby, any PCV2 ORF-2 would be effective as the source of the PCV ORF-2 DNA and/or polypeptide as used herein. A preferred PCV2 ORF-2 protein is that of SEQ ID NO: 11 herein and of WO06/072065. A further preferred PCV ORF-2 polypeptide is provided as SEQ ID NO: 5 herein and in WO06/072065. However, it is understood by those of skill in the art that this sequence could vary by as much as 6-10% in sequence homology and still retain the antigenic characteristics that render it useful in immunogenic compositions. The antigenic characteristics of an immunological composition can be, for example, estimated by the challenge experiment as provided by Example 4 of WO06/072065. Moreover, the antigenic characteristic of a modified antigen is still retained, when the modified antigen confers at least 70%, preferably 80%, more preferably 90% of the protective immunity as compared to the PCV2 ORF-2 protein, encoded by the polynucleotide sequence of SEQ ID NO:3 or SEQ ID NO:4 as provided herein and in WO06/072065.

[0017] Thus according to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in animals having

anti-PCV2 antibodies, in particular maternally-derived anti-PCV2 antibodies, comprising the step of administering an effective amount of PCV2 antigen or an immunogenic composition comprising a PCV2 antigen to that animal in need of such treatment, wherein the PCV2 antigen is an antigen of PCV2 ORF-2 protein that has at least 70%, preferably, 80% even more preferably 90% of the protective immunity as compared to the PCV2 ORF-2 protein, encoded by the polynucleotide sequence of SEQ ID NO:3 or SEQ ID NO:4 as provided herein and in WO06/072065. Preferably said PCV2 ORF-2 sequences have the sequence of SEQ ID NO: 11 or SEQ ID NO: 5 as provided herein and in WO06/072065.

[0018] In some forms, immunogenic portions of PCV2 ORF-2 protein are used as the antigenic component in the immunogenic composition, comprising PCV2 antigen. The term "immunogenic portion" as used herein refers to truncated and/or substituted forms, or fragments of PCV2 ORF-2 protein and/or polynucleotide, respectively. Preferably, such truncated and/or substituted forms, or fragments will comprise at least 6 contiguous amino acids from the full-length ORF-2 polypeptide. More preferably, the truncated or substituted forms, or fragments will have at least 10, more preferably at least 15, and still more preferably at least 19 contiguous amino acids from the full-length PCV ORF-2 polypeptide. Two preferred sequences in this respect are provided as SEQ ID NO: 9 and SEQ ID NO:10 herein and in WO06/072065. It is further understood that such sequences may be a part of larger fragments or truncated forms.

[0019] As mentioned above, a further preferred PCV2 ORF-2 polypeptide is any one encoded by the nucleotide sequences of SEQ ID NO: 3 or SEQ ID NO: 4. However, it is understood by those of skill in the art that this sequence could vary by as much as 6-20% in sequence homology and still retain the antigenic characteristics that render it useful in immunogenic compositions. In some forms, a truncated or substituted form, or fragment of this PCV2 ORF-2 polypeptide is used as the antigenic component in the composition. Preferably, such truncated or substituted forms, or fragments will comprise at least 18 contiguous nucleotides from the full-length PCV2 ORF-2 nucleotide sequence, e.g. of SEQ ID NO: 3 or SEQ ID NO: 4. More preferably, the truncated or substituted forms, or fragments, will have at least 30, more preferably at least 45, and still more preferably at least 57 contiguous nucleotides of the full-length PCV2 ORF-2 nucleotide sequence, e.g. SEQ ID NO: 3 or SEQ ID NO: 4.

[0020] "Sequence identity" as it is known in the art refers to a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, namely a reference sequence and a given sequence to be compared with the reference sequence. Sequence identity is determined by comparing the given sequence to the reference sequence after the sequences have been optimally aligned to produce the highest degree of sequence similarity, as determined by the match between strings of such sequences. Upon such alignment, sequence identity is ascertained on a position-by-position basis, e.g., the sequences are "identical" at a particular position if at that position, the nucleotides or amino acid residues are identical. The total number of such position identities is then divided by the total number of nucleotides or residues in the reference sequence to give % sequence identity. Sequence identity can be readily calculated by known methods, including but not limited to, those described in Computational Molecular Biology, Lesk, A. N., ed., Oxford University Press, New York (1988), Biocomputing: Informatics and Genome Projects, Smith, D.W., ed., Academic Press, New York (1993); Computer Analysis of Sequence Data, Part I, Griffin, A.M., and Griffin, H. G., eds., Humana Press, New Jersey (1994); Sequence Analysis in Molecular Biology, von Heinge, G., Academic Press (1987); Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York (1991); and Carillo, H., and Lipman, D., SIAM J. Applied Math., 48: 1073 (1988). Preferred methods to determine the sequence identity are designed to give the largest match between the sequences tested. Methods to determine sequence identity are codified in publicly available computer programs which determine sequence identity between given sequences. Examples of such programs include, but are not limited to, the GCG program package (Devereux, J., et al., Nucleic Acids Research, 12(1):387 (1984)), BLASTP, BLASTN and FASTA (Altschul, S. F. et al., J. Molec. Biol., 215:403-410 (1990)). The BLASTX program is publicly available from NCBI and other sources (BLAST Manual, Altschul, S. et al., NCVI NLM NIH Bethesda, MD 20894, Altschul, S. F. et al., J. Molec. Biol., 215:403-410 (1990)). These programs optimally align sequences using default gap weights in order to produce the highest level of sequence identity between the given and reference sequences. As an illustration by a polynucleotide having a nucleotide sequence having at least, for example, 85%, preferably 90%, even more preferably 95% "sequence identity" to a reference nucleotide sequence, it is intended that the nucleotide sequence of the given polynucleotide is identical to the reference sequence except that the given polynucleotide sequence may include up to 15, preferably up to 10, even more preferably up to 5 point mutations per each 100 nucleotides of the reference nucleotide sequence. In other words, in a polynucleotide having a nucleotide sequence having at least 85%, preferably 90%, even more preferably 95% identity relative to the reference nucleotide sequence, up to 15%, preferably 10%, even more preferably 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 15%, preferably 10%, even more preferably 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. Analogously, by a polypeptide having a given amino acid sequence having at least, for example, 85%, preferably 90%, even more preferably 95% sequence identity to a reference amino acid sequence, it is intended

that the given amino acid sequence of the polypeptide is identical to the reference sequence except that the given polypeptide sequence may include up to 15, preferably up to 10, even more preferably up to 5 amino acid alterations per each 100 amino acids of the reference amino acid sequence. In other words, to obtain a given polypeptide sequence having at least 85%, preferably 90%, even more preferably 95% sequence identity with a reference amino acid sequence, up to 15%, preferably up to 10%, even more preferably up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 15%, preferably up to 10%, even more preferably up to 5% of the total number of amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or the carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in the one or more contiguous groups within the reference sequence. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. However, conservative substitutions are not included as a match when determining sequence identity.

[0021] "Sequence homology", as used herein, refers to a method of determining the relatedness of two sequences. To determine sequence homology, two or more sequences are optimally aligned, and gaps are introduced if necessary. However, in contrast to "sequence identity", conservative amino acid substitutions are counted as a match when determining sequence homology. In other words, to obtain a polypeptide or polynucleotide having 95% sequence homology with a reference sequence, 85%, preferably 90%, even more preferably 95% of the amino acid residues or nucleotides in the reference sequence must match or comprise a conservative substitution with another amino acid or nucleotide, or a number of amino acids or nucleotides up to 15%, preferably up to 10%, even more preferably up to 5% of the total amino acid residues or nucleotides, not including conservative substitutions, in the reference sequence may be inserted into the reference sequence. Preferably the homolog sequence comprises at least a stretch of 50, even more preferably at least 100, even more preferably at least 250, and even more preferably at least 500 nucleotides.

[0022] A "conservative substitution" refers to the substitution of an amino acid residue or nucleotide with another amino acid residue or nucleotide having similar characteristics or properties including size, hydrophobicity, etc., such that the overall functionality does not change significantly.

[0023] "Isolated" means altered "by the hand of man" from its natural state, i.e., if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

[0024] Thus, according to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in animals having anti-PCV2 antibodies, in particular maternally-derived anti-PCV2 antibodies, comprising the step of administering an effective amount of PCV2 antigen or an immunogenic composition comprising a PCV2 antigen to that animal in need of such treatment, wherein said PCV2 ORF-2 protein is any one of those, described above. Preferably, said PCV2 ORF-2 protein is

i) a polypeptide comprising the sequence of SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 9, SEQ ID NO: 10 or SEQ ID NO: 11 herein or of WO06/07065;

ii) any polypeptide that is at least 80% homologous to the polypeptide of i),

iii) any immunogenic portion of the polypeptides of i) and/or ii)

iv) the immunogenic portion of iii), comprising at least 10 contiguous amino acids included in the sequences of SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 9, SEQ ID NO: 10 or SEQ ID NO: 11 herein or of WO06/072065,

v) a polypeptide that is encoded by a DNA comprising the sequence of SEQ ID NO: 3 or SEQ ID NO: 4 herein or of WO06/072065.

vi) any polypeptide that is encoded by a polynucleotide that is at least 80% homologous to the polynucleotide of v),

vii) any immunogenic portion of the polypeptides encoded by the polynucleotide of v) and/or vi)

viii) the immunogenic portion of vii), wherein polynucleotide coding for said immunogenic portion comprises at least 30 contiguous nucleotides included in the sequences of SEQ ID NO: 3, or SEQ ID NO: 4 herein or of WO06/072065.

[0025] Preferably any of those immunogenic portions have the immunogenic characteristics of PCV2 ORF-2 protein that is encoded by the sequence of SEQ ID NO: 3 or SEQ ID NO: 4 herein or of WO06/07065.

[0026] According to a further aspect, PCV2 ORF-2 protein is provided in the immunogenic composition at an antigen inclusion level effective for inducing the desired immune response, namely reducing the incidence of lessening the severity of, or preventing or reducing one or more clinical symptoms resulting from or associated with a PCV2 infection. Preferably, the PCV2 ORF-2 protein inclusion level is at least 0.2 μg antigen / ml of the final immunogenic composition ($\mu\text{g}/\text{ml}$), more preferably from about 0.2 to about 400 $\mu\text{g}/\text{ml}$, still more preferably from about 0.3 to about 200 $\mu\text{g}/\text{ml}$, even more preferably from about 0.35 to about 100 $\mu\text{g}/\text{ml}$, still more preferably from about 0.4 to about 50 $\mu\text{g}/\text{ml}$, still more preferably from about 0.45 to about 30 $\mu\text{g}/\text{mL}$ still more preferably from about 0.5 to about 18 $\mu\text{g}/\text{ml}$, even more

preferably from about 0.6 to about 15 $\mu\text{g/ml}$ even more preferably from about 0.75 to about 8 $\mu\text{g/ml}$, even more preferably from about 1.0 to about 6 $\mu\text{g/ml}$, still more preferably from about 1.3 to about 3.0 $\mu\text{g/ml}$, even more preferably from about 1.4 to about 2.5 $\mu\text{g/ml}$, even more preferably from about 1.5 to about 2.0 $\mu\text{g/ml}$, and most preferably about 1.6 $\mu\text{g/ml}$.

[0027] According to a further aspect, the PCV2 ORF-2 antigen inclusion level is at least 0.2 $\mu\text{g/PCV2 ORF-2 protein}$ as described above per dose of the final antigenic composition ($\mu\text{g/dose}$), more preferably from about 0.2 to about 400 $\mu\text{g/dose}$, still more preferably from about 0.3 to about 200 $\mu\text{g/dose}$, even more preferably from about 0.35 to about 100 $\mu\text{g/dose}$, still more preferably from about 0.4 to about 50 $\mu\text{g/dose}$, still more preferably from about 0.45 to about 30 $\mu\text{g/dose}$, still more preferably from about 0.5 to about 18 $\mu\text{g/dose}$, even more preferably from about 0.6 to about 15 $\mu\text{g/ml}$, even more preferably from about 0.75 to about 8 $\mu\text{g/dose}$, even more preferably from about 1.0 to about 6 $\mu\text{g/dose}$, still more preferably from about 1.3 to about 3.0 $\mu\text{g/dose}$, even more preferably from about 1.4 to about 2.5 $\mu\text{g/dose}$, even more preferably from about 1.5 to about 2.0 $\mu\text{g/dose}$, and most preferably about 1.6 $\mu\text{g/dose}$. It has been surprisingly found, that a PCV2 ORF-2 protein inclusion level (antigen content) of less than 20 $\mu\text{g/dose}$, preferably of about 0.5 to 18 $\mu\text{g/dose}$ is suitable to confer immunity in young animals and/or in animals which are positive for PCV2 antibodies, in particular which are positive for anti-PCV2 maternally-derived antibodies. Thus, according to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in animals having anti-PCV2 antibodies, in particular maternally-derived anti-PCV2 antibodies, comprising the step of administering less than 20 $\mu\text{g/dose}$, preferably of about 0.5 to 18 $\mu\text{g/dose}$ of PCV2 antigen or an immunogenic composition comprising an PCV2 antigen to that animal in need of such treatment. Said PCV2 antigen is any one described in this patent application. Preferably, said PCV2 antigen is any PCV2 ORF-2 protein, more preferably, any PCV2 ORF-2 protein described herein.

[0028] The PCV2 ORF-2 polypeptide used in the immunogenic composition in accordance with the present invention can be derived in any fashion including isolation and purification of PCV2 ORF2, standard protein synthesis, and recombinant methodology. Preferred methods for obtaining PCV2 ORF-2 polypeptide are provided in WO06/072065. Briefly, susceptible cells are infected with a recombinant viral vector containing PCV2 ORF-2 DNA coding sequences, PCV2 ORF-2 polypeptide is expressed by the recombinant virus, and the expressed PCV2 ORF-2 polypeptide is recovered from the supernatant by filtration and inactivated by any conventional method, preferably using binary ethylenimine, which is then neutralized to stop the inactivation process.

[0029] The immunogenic composition as used herein also refers to a composition that comprises i) any of the PCV2 ORF-2 protein described above, preferably in concentrations described above, and ii) at least a portion of the viral vector expressing said PCV2 ORF-2 protein, preferably of a recombinant baculovirus. Moreover, the immunogenic composition can comprise i) any of the PCV2 ORF-2 proteins described above, preferably in concentrations described above, ii) at least a portion of the viral vector expressing said PCV2 ORF-2 protein, preferably of a recombinant baculovirus, and iii) a portion of the cell culture supernatant.

[0030] Thus, according to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in animals having anti-PCV2 antibodies, in particular maternally-derived anti-PCV2 antibodies, comprising the step of administering an effective amount of PCV2 antigen or an immunogenic composition comprising an PCV2 antigen to that animal in need of such treatment, wherein the PCV2 antigen is recombinant PCV2 ORF-2, preferably a baculovirus expressed PCV2 ORF-2. Preferably those recombinant or baculovirus expressed PCV2 ORF-2 having the sequence as described above.

[0031] The immunogenic composition as used herein also refers to a composition that comprises i) any of the PCV2 ORF-2 proteins described above, preferably in concentrations described above, ii) at least a portion of the viral vector expressing said PCV2 ORF-2 protein, preferably of a recombinant baculovirus, and iii) a portion of the cell culture; wherein about 90% of the components have a size smaller than 1 μm .

[0032] The immunogenic composition as used herein also refers to a composition that comprises i) any of the PCV2 ORF-2 proteins described above, preferably in concentrations described above, ii) at least a portion of the viral vector expressing said PCV2 ORF-2 protein, iii) a portion of the cell culture, iv) and inactivating agent to inactivate the recombinant viral vector preferably BEI, wherein about 90% of the components i) to iii) have a size smaller than 1 μm . Preferably, BEI is present in concentrations effective to inactivate the baculovirus, preferably in an amount of 2 to about 8 mM BEI, preferably of about 5 mM BEI.

[0033] The immunogenic composition as used herein also refers to a composition that comprises i) any of the PCV2 ORF-2 proteins described above, preferably in concentrations described above, ii) at least a portion of the viral vector expressing said PCV2 ORF-2 protein, iii) a portion of the cell culture, iv) an inactivating agent to inactivate the recombinant viral vector preferably BEI, and v) a neutralization agent to stop the inactivation mediated by the inactivating agent, wherein about 90% of the components i) to iii) have a size smaller than 1 μm . Preferably, if the inactivating agent is BEI, said composition comprises sodium thiosulfate in equivalent amounts to BEI.

[0034] The polypeptide is incorporated into a composition that can be administered to an animal susceptible to PCV2 infection. In preferred forms, the composition may also include additional components known to those of skill in the art (see also Remington's Pharmaceutical Sciences. (1990). 18th ed. Mack Publ., Easton). Additionally, the composition

may include one or more veterinary-acceptable carriers. As used herein, "a veterinary-acceptable carrier" includes any and all solvents, dispersion media, coatings, adjuvants, stabilizing agents, diluents, preservatives, antibacterial and antifungal agents, isotonic agents, adsorption delaying agents, and the like. In a preferred embodiment, the immunogenic composition comprises PCV2 ORF-2 protein as provided herewith, preferably in concentrations described above, which is mixed with an adjuvant, preferably Carbopol, and physiological saline.

[0035] Those of skill in the art will understand that the composition used herein may incorporate known injectable, physiologically acceptable sterile solutions. For preparing a ready-to-use solution for parenteral injection or infusion, aqueous isotonic solutions, such as e.g. saline or corresponding plasma protein solutions, are readily available. In addition, the immunogenic and vaccine compositions used in the present invention can include diluents, isotonic agents, stabilizers, or adjuvants. Diluents can include water, saline, dextrose, ethanol, glycerol, and the like. Isotonic agents can include sodium chloride, dextrose, mannitol, sorbitol, and lactose, among others. Stabilizers include albumin and alkali salts of ethylenediaminetetracetic acid, among others.

[0036] "adjuvants" as used herein, can include aluminium hydroxide and aluminium phosphate, saponins e.g., Quil A, QS-21 (Cambridge Biotech Inc., Cambridge MA), GPI-0100 (Galenica Pharmaceuticals, Inc., Birmingham, AL), water-in-oil emulsion, oil-in-water emulsion, water-in-oil-in-water emulsion. The emulsion can be based in particular on light liquid paraffin oil (European Pharmacopoeia type); isoprenoid oil such as squalane or squalene oil resulting from the oligomerization of alkenes, in particular of isobutene or decene; esters of acids or of alcohols containing a linear alkyl group, more particularly plant oils, ethyl oleate, propylene glycol di-(caprylate/caprate), glyceryl tri-(caprylate/caprate) or propylene glycol dioleate; esters of branched fatty acids or alcohols, in particular isostearic acid esters. The oil is used in combination with emulsifiers to form the emulsion. The emulsifiers are preferably nonionic surfactants, in particular esters of sorbitan, of mannide (e.g. anhydromannitol oleate), of glycol, of polyglycerol, of propylene glycol and of oleic, isostearic, ricinoleic or hydroxystearic acid, which are optionally ethoxylated, and polyoxypropylene-polyoxyethylene copolymer blocks, in particular the Pluronic products, especially L121. See Hunter et al., *The Theory and Practical Application of Adjuvants* (Ed. Stewart-Tull, D. E. S.). John Wiley and Sons, NY, pp51-94 (1995) and Todd et al., *Vaccine* 15:564-570 (1997).

[0037] For example, it is possible to use the SPT emulsion described on page 147 of "Vaccine Design, The Subunit and Adjuvant Approach" edited by M. Powell and M. Newman, Plenum Press, 1995, and the emulsion MF59 described on page 183 of this same book.

[0038] A further instance of an adjuvant is a compound chosen from the polymers of acrylic or methacrylic acid and the copolymers of maleic anhydride and alkenyl derivative. Advantageous adjuvant compounds are the polymers of acrylic or methacrylic acid which are cross-linked, especially with polyalkenyl ethers of sugars or polyalcohols. These compounds are known by the term carbomer (Pharmeuropa Vol. 8, No. 2, June 1996). Persons skilled in the art can also refer to U. S. Patent No. 2,909,462 which describes such acrylic polymers cross-linked with a polyhydroxylated compound having at least 3 hydroxyl groups, preferably not more than 8, the hydrogen atoms of at least three hydroxyls being replaced by unsaturated aliphatic radicals having at least 2 carbon atoms. The preferred radicals are those containing from 2 to 4 carbon atoms, e.g. vinyls, allyls and other ethylenically unsaturated groups. The unsaturated radicals may themselves contain other substituents, such as methyl. The products sold under the name Carbopol; (BF Goodrich, Ohio, USA) are particularly appropriate. Carbopol™ as used throughout this specification is a trademark designation for polyacrylic acid polymers also known as Carbomer. They are cross-linked with an allyl sucrose or with allyl pentaerythritol. Among them, there may be mentioned Carbopol 974P, 934P and 971P. Most preferred is the use of Carbopol, in particular the use of Carbopol 971P, preferably in amounts of about 500 µg to about 5 mg per dose, even more preferred in an amount of about 750 µg to about 2.5 mg per dose and most preferred in an amount of about 1 mg per dose.

[0039] Further suitable adjuvants include, but are not limited to, the RIBI adjuvant system (Ribi Inc.), Block co-polymer (CytRx, Atlanta GA), SAF-M (Chiron, Emeryville CA), monophosphoryl lipid A, Avridine lipid-amine adjuvant, heat-labile enterotoxin from *E. coli* (recombinant or otherwise), cholera toxin, IMS 1314, or muramyl dipeptide among many others.

[0040] Preferably, the adjuvant is added in an amount of about 100 µg to about 10 mg per dose. Even more preferably, the adjuvant is added in an amount of about 100 µg to about 10 mg per dose. Even more preferably, the adjuvant is added in an amount of about 500 µg to about 5 mg per dose. Even more preferably, the adjuvant is added in an amount of about 750 µg to about 2.5 mg per dose. Most preferably, the adjuvant is added in an amount of about 1 mg per dose.

[0041] Additionally, the composition can include one or more pharmaceutical-acceptable carriers. As used herein, "a pharmaceutical-acceptable carrier" includes any and all solvents, dispersion media, coatings, stabilizing agents, diluents, preservatives, antibacterial and antifungal agents, isotonic agents, adsorption delaying agents, and the like. Most preferably, the composition provided herewith, contains PCV2 ORF-2 protein recovered from the supernatant of in vitro cultured cells, wherein said cells were infected with a recombinant viral vector containing PCV2 ORF-2 DNA and expressing PCV2 ORF-2 protein, and wherein said cell culture was treated with about 2 to about 8 mM BEI, preferably with about 5 mM BEI to inactivate the viral vector, and an equivalent concentration of a neutralization agent, preferably sodium thiosulfate solution to a final concentration of about 2 to about 8 mM, preferably of about 5 mM.

[0042] The present disclosure also relates to an immunogenic composition that comprises i) any of the PCV2 ORF-

2 proteins described above, preferably in concentrations described above, ii) at least a portion of the viral vector expressing said PCV2 ORF-2 protein, iii) a portion of the cell culture, iv) an inactivating agent to inactivate the recombinant viral vector (preferably BEI), and v) a neutralization agent to stop the inactivation mediated by the inactivating agent, preferably sodium thiosulfate in equivalent amounts to BEI; and vi) a suitable adjuvant, preferably Carbopol 971 in amounts described above; wherein about 90% of the components i) to iii) have a size smaller than 1 μm . According to a further aspect, this immunogenic composition further comprises a pharmaceutical acceptable salt, preferably a phosphate salt in physiologically acceptable concentrations. Preferably, the pH of said immunogenic composition is adjusted to a physiological pH, meaning between about 6.5 and 7.5.

[0043] The immunogenic composition as used herein also refers to a composition that comprises per one ml i) at least 1.6 μg of PCV2 ORF-2 protein described above, preferably less than 20 μg ii) at least a portion of baculovirus expressing said PCV2 ORF-2 protein iii) a portion of the cell culture, iv) about 2 to 8 mM BEI, v) sodium thiosulfate in equivalent amounts to BEI; and vi) about 1 mg Carbopol 971, and vii) phosphate salt in a physiologically acceptable concentration; wherein about 90% of the components i) to iii) have a size smaller than 1 μm and the pH of said immunogenic composition is adjusted to about 6.5 to 7.5.

[0044] The immunogenic compositions can further include one or more other immuno-modulatory agents such as, e.g., interleukins, interferons, or other cytokines. The immunogenic compositions can also include Gentamicin and Merthiolate. While the amounts and concentrations of adjuvants and additives useful in the context of the present invention can readily be determined by the skilled artisan, the present invention contemplates compositions comprising from about 50 μg to about 2000 μg of adjuvant and preferably about 250 μg /ml dose of the vaccine composition. Thus, the immunogenic composition as used herein also refers to a composition that comprises from about 1 μg /ml to about 60 μg /ml of antibiotics, and more preferably less than about 30 μg /ml of antibiotics.

[0045] The immunogenic composition as used herein also refers to a composition that comprises i) any of the PCV2 ORF-2 proteins described above, preferably in concentrations described above, ii) at least a portion of the viral vector expressing said PCV2 ORF-2 protein, iii) a portion of the cell culture, iv) an inactivating agent to inactivate the recombinant viral vector preferably BEI, and v) an neutralization agent to stop the inactivation mediated by the inactivating agent, preferably sodium thiosulfate in equivalent amounts to BEI; vi) a suitable adjuvant, preferably Carbopol 971 in amounts described above; vii) a pharmaceutical acceptable concentration of a saline buffer, preferably of a phosphate salt, and viii) an anti-microbiological active agent; wherein about 90% of the components i) to iii) have a size smaller than 1 μm .

[0046] The immunogenic composition as used herein also refers to Ingelvac® CircoFLEX™, (Boehringer Ingelheim Vetmedica Inc, St Joseph, MO, USA).

[0047] Thus according to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in animals in animals having anti-PCV2 maternal antibodies, comprising the step of administering an effective amount of a PCV2 antigen to that animal in need of such treatment, wherein the immunogenic composition is CircoFLEX®, CircoVac®, CircoVent or Suvaxyn PCV-2 One Dose®. Most preferably, the immunogenic composition is Ingelvac® CircoFLEX™, and/or the PCV2 antigen is PCV2 ORF-2, preferably, baculovirus expressed PCV2 ORF-2, most preferably as included in Ingelvac® CircoFLEX™.

[0048] For investigation of a possible interference of PCV2 antigen with the maternal antibody a study was conducted in which the antibody titers of study animals were determined at the time of vaccination which were then grouped into a low, moderate and high antibody class: Geometric mean titers of < 1:100 were considered as low antibody titers, titers of 1:100 to 1:1000 were considered as moderate antibody titers and titers of > 1:1000 were considered as high antibody titers. This grouping pattern is comparable to that done in a Canadian field study where antibody titers of 1:80 were considered as low, antibody titers of 1:640 as moderate and antibody titers of > 1:1280 as high (Laroche et al., 2003, Can. J. Vet. Res.; 67: 114-120). In order to analyze the impact of low, medium and high antibody titers at the time of vaccination on viremia, vaccinated and placebo-treated animals were compared with regard to the onset, end, duration of viremia, the number of positive sampling days and the virus load. It was surprisingly found, that the presence of anti-PCV2 antibodies, in particular of maternally-derived antibodies, had no significant impact of any of those parameters. In other words, it was surprisingly found that the efficacy of the PCV2 antigen in prevention and treatment of a PCV2 infection or in reduction of clinical symptoms caused by or associated with a PCV2 infection in animals was not affected at the day of vaccination by the presence of anti-PCV2 antibodies, by anti-PCV2 antibody titers of more than 1:1000. This effect could be shown in a one shot vaccination experiment, which means that the PCV2 antigen was administered only once and without any subsequent administration of PCV2 antigen.

[0049] Methods for detection and quantification of anti-PCV2 antibodies are well known in the art. For example detection and quantification of PCV2 antibodies can be performed by indirect immunofluorescence as described in Magar et al., 2000, Can. J. Vet. Res.; 64: 184-186 or Magar et al., 2000, J. Comp. Pathol.; 123: 258-269. Further assays for quantification of anti-PCV2 antibodies are described in Opriessnig et al., 2006, 37th Annual Meeting of the American Association of Swine Veterinarians. Moreover, example 2 also describes an indirect immunofluorescence assay that can be used by a person skilled in the art. In cases of controversial results and in any question of doubt, anti-PCV2 titers as mentioned

herein, refer to those which are /can be estimated by the assay as described in Example 2.

[0050] Thus according to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in animals having anti-PCV2 antibodies, in particular maternal antibodies, comprising the step of administering an effective amount of a PCV2 antigen to that animal in need of such treatment, preferably of less than 20 µg/dose wherein said animal has a detectable anti-PCV2 antibody titer of more than 1:1000. Preferably, such an anti-PCV2 antibody titer is detectable and quantifiable in a specific anti-PCV2 immune assay, preferably in the assay as described in Example 2. More preferably, those anti-PCV-2 antibodies are maternally-derived antibodies. The PCV2 antigen is only administered once, preferably with a dose of less than 20 µg/dose.

[0051] Piglets with only low titers (< 1:100) or moderate titers (< 1:1000) of maternally-derived anti-PCV2 antibodies are not sufficiently protected against PCV2 infections which occur prior to week 3 of age. Therefore, vaccination at a very early stage of life is desirable. Due to the unpredictable and unexpected results provided herein and demonstrating the lack of interference of anti-PCV2 antibodies with PCV2 antigen, vaccination/treatment of animals before 3 weeks of age becomes realistic. Moreover, it has been surprisingly found that anti-PCV2 antibody titers of more than 1:1000 had no influence on the efficacy of the PCV2 vaccine regardless of the level of the existing initial antibody titer. For example, vaccination of high-titer animals (anti-PCV2 antibody titer > 1:1000) resulted in a 9.5 day shorter duration of viremia, 11.9 days earlier end of viremia, 1.9 days less viremic sampling days and an approximately 2-fold reduction of the sum of genomic equivalents/ml as compared to non vaccinated control animals. Upon comparison of vaccinated "high" "moderate" and "low titer animals" no significant differences were observed with regard to the different parameters of PCV2 viraemia. These results indicate that also in the presence of high anti-PCV2 antibody titers, the PCV2 antigen used for vaccination can still significantly reduce viremia in blood (end of viremia, duration of viremia, virus load). In line with this finding, no differences could be found with regard to the live body weight when comparing low and high titer animals of the vaccinated group. Furthermore vaccinated animals with a high anti-PCV2 antibody titer at the time of vaccination/treatment (> 1:1000) also showed a significantly higher body weight after the onset of viremia compared to placebo-treated animals with initial high antibody titers (see Figure 3). Consequently, vaccination/treatment of animals of 1 day of age or older with PCV2 antigen is possible. However, vaccination should be done within the first 8, preferably within the first 7 weeks of age. Thus according to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in animals, comprising the step of administering to that animal in need of such treatment at day 1 of age or later, preferably but not later than at week 8 of age an effective amount of a PCV2 antigen. According to a preferred embodiment, less than 20 µg/dose PCV2 antigen are required to confer immunity in such animal. The PCV2 antigen, preferably less than 20 µg/dose thereof, is only administered once to the animal in need of such treatment.

[0052] According to a further, more general aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in young animals, comprising the step of administering an effective amount of a PCV2 antigen to that animal in need of such treatment.

[0053] The term "young animal" as used herein refers to an animal of 1 to 22 days of age. Preferably, by the term young animal, an animal of 1 to 20 days of age is meant. More preferably, the term young animal refers to an animal of 1 to 15 days of age, even more preferably of 1 day of age to 14 days of age, even more preferably of 1 to 12 days of age, even more preferably of 1 to 10 days of age, even more preferably of 1 to 8 days of age, even more preferably of 1 to 7 days of age, even more preferably of 1 to 6 days of age, even more preferably of 1 to 5 days of age, even more preferably of 1 to 4 days of age, even more preferably of 1 to 3 days of age, even more preferably of 1 or 2 day(s) of age, and most preferably to an animal of 1 day of age. Thus according to a further aspect, the present disclosure a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in young animals, comprising the step of administering an effective amount of a PCV2 antigen to an animal of 1 to 22 days of age, preferably of 1 to 20 days of age, more preferably of 1 to 15 days of age, even more preferably of 1 to 14 days of age, even more preferably of 1 to 12 days of age, even more preferably of 1 to 10 days of age, even more preferably of 1 to 8 days of age, even more preferably of 1 to 7 days of age, even more preferably of 1 to 6 days of age, even more preferably of 1 to 5 days of age, even more preferably of 1 to 4 days of age, even more preferably of 1 to 3 days of age, even more preferably of 1 or 2 day(s) of age, most preferably at 1 day of age in need of such treatment. For example, evidence is given that vaccination/treatment on 19 to 22 days of age shows high efficacy of vaccination. Moreover, vaccination/treatment at 12 to 18, preferably 12 to 14 days of age has also be shown to be very effective in the reduction of clinical symptoms associated with PCV2 infections, reduction of overall viral load, reduction of duration of viremia, delay in onset of viremia, and weight gain. Moreover, vaccination at 1 week of age has also been shown to be very effective in reduction of clinical symptoms associated with PCV2 infections, reduction of overall viral load, reduction of duration of viremia, delay in onset of viremia, weight gain. Preferably less than 20 µg/dose PCV2 antigen are required to confer immunity in those young animals. The PCV2 antigen, preferably less than 20 µg, is only administered once to that young animal in need of such treatment.

[0054] Due to the ubiquity of PCV2 in the field, most of the young piglets are seropositive in respect to PCV2. Thus according to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in young animals having anti-PCV2 antibodies at the day of vaccination, comprising the step of administering an effective amount of a PCV2 antigen to an animal of 1 to 22 days of age, preferably of 1 to 20 days of age, more preferably of 1 to 15 days of age, even more preferably of 1 to 14 days of age, even more preferably of 1 to 12 days of age, even more preferably of 1 to 10 days of age, even more preferably of 1 to 8 days of age, even more preferably of 1 to 7 days of age, even more preferably of 1 to 6 days of age, even more preferably of 1 to 5 days of age, even more preferably of 1 to 4 days of age, even more preferably of 1 to 3 days of age, even more preferably at 1 or 2 day(s) of age, and most preferably at 1 day of age in need of such treatment. Preferably, said young animals, at the day of vaccination/treatment, have a detectable anti-PCV2 antibody titer of up to 1:100, preferably of more than 1:100, even more preferably of more than 1:250, even more preferably of more than 1:500, even more preferably of 1:640, even more preferably of more than 1:750, most preferably of more than 1:1000 at the day of vaccination/treatment. Preferably less than 20 µg/dose PCV2 antigen are required to confer a sufficient immunity in those young animals. The PCV2 antigen, preferably less than 20 µg, is only administered once to that young animal in need of such treatment.

[0055] As described above, vaccination/treatment of young animals with PCV2 antigen resulted in a shortening of viremic phase as compared to non vaccinated control animals. The average shortening time was 9.5 days as compared to non vaccinated control animals of the same species. Therefore, according to a further aspect, the present disclosure also provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in young animals, comprising the step of administering an effective amount of a PCV2 antigen to that animal in need of such treatment, wherein the treatment or prophylaxis results in shortening of the viremia phase of 5 or more days, preferably 6 or more day, even more preferably of 7 or more days, even more preferably of 8 or more days, even more preferably of 9, even more preferably of 10, even more preferably of 12, even more preferably of 14, and most preferably of more than 16 days as compared to animals of a non-treated control group of the same species. In some cases viremic phase is shortened by more than 20 days. In general, the vaccination of young piglets resulted in a reduction in the loss of weight gain, a shorter duration of viremia, an earlier end to viremia, and a lower virus load. Therefore, according to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in young animals, comprising the step of administering an effective amount of a PCV2 antigen to that animal in need of such treatment, wherein said treatment or prophylaxis of PCV2 infection results in an improvement in comparison to animals of a non-treated control group of the same species in a vaccine efficacy parameter selected from the group consisting of a reduction in the loss of weight gain, a shorter duration of viremia, an earlier end to viremia, a lower virus load, or combinations thereof. Preferably less than 20 µg/dose PCV2 antigen are required to cause any of the improved vaccine efficacy parameters mentioned above.

[0056] Moreover such improved vaccine efficacy parameter(s) are achieved by a single administration of only one dose.

[0057] The term "an effective amount" as used herein means, but is not limited to, an amount of antigen, that elicits or is able to elicit an immune response in an animal, to which said effective dose of PCV2 antigen is administered. Preferably, an effective amount is defined as an amount of antigen that confers a duration of immunity (DOI) of at least 10 weeks, preferably at least 12 weeks, more preferably at least 15 weeks, and most preferably at least 20 weeks.

[0058] The amount that is effective depends on the ingredients of the vaccine and the schedule of administration. Typically, when an inactivated virus or a modified live virus preparation is used in the combination vaccine, an amount of the vaccine containing about $10^{2.0}$ to about $10^{9.0}$ TCID₅₀ per dose, preferably about $10^{3.0}$ to about $10^{8.0}$ TCID₅₀ per dose, and more preferably, about $10^{4.0}$ to about $10^{8.0}$ TCID₅₀ per dose is used. In particular, when modified live PCV2 is used in the vaccines, the recommended dose to be administered to the susceptible animal is preferably about $10^{3.0}$ TCID₅₀ (tissue culture infective dose 50% end point)/dose to about $10^{6.0}$ TCID₅₀/dose and more preferably about $10^{4.0}$ TCID₅₀/dose to about $10^{5.0}$ TCID₅₀/dose. In general, the quantity of antigen will be between 0.2 and 5000 micrograms, and between $10^{2.0}$ and $10^{9.0}$ TCID₅₀, preferably between $10^{3.0}$ and $10^{6.0}$ TCID₅₀, more preferably between $10^{4.0}$ and $10^{5.0}$ TCID₅₀, when purified antigen is used.

[0059] Sub-unit vaccines are normally administered with an antigen inclusion level of at least 0.2 µg antigen per dose, preferably with about 0.2 to about 400 µg/dose, still more preferably with about 0.3 to about 200 µg/dose, even more preferably with about 0.35 to about 100 µg/dose, still more preferably with about 0.4 to about 50 µg/dose, still more preferably with about 0.45 to about 30 µg/dose, still more preferably with about 0.5 to about 18 g/dose, still more preferably with about 0.6 to about 16 µg/dose, even more preferably with about 0.75 to about 8 µg/dose, even more preferably with about 1.0 to about 6 µg/dose, and still more preferably with about 1.3 to about 3.0 µg/dose.

[0060] Unexpectedly, it was found that the prophylactic use of the immunogenic compositions described *supra*, is effective for the reduction of clinical symptoms caused by or associated with PCV2 infections, preferably in young animals and/or in animals having passive immunity against PCV2 at the day of treatment. In particular, it was discovered that the prophylactic use of the immunogenic compositions as described herein, and specifically of compositions comprising

PCV2 ORF-2 antigen, is effective for reducing lymphadenopathy, lymphoid depletion and/or multinucleated/giant histiocytes in animals infected with PCV2 and having maternal anti-PCV-2 antibodies at the day of treatment/vaccination. Furthermore, it was discovered that the prophylactic use of the immunogenic compositions as described herein, and specifically of compositions comprising PCV2 ORF-2 antigen, is effective for reducing (1) interstitial pneumonia with interlobular edema, (2) cutaneous pallor or icterus, (3) mottled atrophic livers, (4) gastric ulcers, (5) nephritis (6) reproductive disorders, e.g. abortion, stillbirths, mummies, etc, (7) Pia like lesions, normally known to be associated with *Lawsonia intracellularis* infections (Ileitis), (8) lymphadenopathy, (9) lymphoid depletion and/or (10) multinucleated/giant histiocytes (11) Porcine Dermatitis and Nephropathy Syndrome (PDNS), (12) PCVAD associated mortality, (13) PCVAD associated weight loss, (14), reduced growth variability (15), reduced frequency of 'runts' (16), reduced co-infections with Porcine Reproductive and Respiratory Disease Complex (PRRSV). Such an immunogenic composition is also effective in improving economical important growth parameters such as time to slaughter, carcass weight, and/or lean meat ratio. Thus the term "clinical symptoms" as used herein, means, but is not limited to (1) interstitial pneumonia with interlobular edema, (2) cutaneous pallor or icterus, (3) mottled atrophic livers, (4) gastric ulcers, (5) nephritis (6) reproductive disorders, e.g. abortion, stillbirths, mummies, etc, (7) Pia like lesions, normally known to be associated with *Lawsonia intracellularis* infections (Ileitis), (8) lymphadenopathy, (9) lymphoid depletion and/or (10) multinucleated/giant histiocytes (11) Porcine Dermatitis and Nephropathy Syndrome (PDNS), (12) PCVAD associated mortality, (13) PCVAD associated weight loss, (14) reduced growth variability (15) reduced frequency of 'runts' and (16) reduced co-infections with Porcine Reproductive and Respiratory Disease Complex (PRRSV). Moreover, the antigenic composition described herein reduces the overall circovirus load including a later onset, a shorter duration, an earlier end of viremia, and a reduced viral load and its immunosuppressive impact in young animals, in particular in those having anti-PCV2 antibodies at the day of vaccination, thereby resulting in a higher level of general disease resistance and a reduced incidence of PCV2 associated diseases and symptoms.

[0061] Thus, according to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in young animals and/or in animals having anti-PCV2 antibodies, comprising the step of administering an effective amount of PCV2 antigen or an immunogenic composition comprising an PCV2 antigen to that animal in need of such treatment, wherein those clinical symptoms are selected from the group consisting of: (1) interstitial pneumonia with interlobular edema, (2) cutaneous pallor or icterus, (3) mottled atrophic livers, (4) gastric ulcers, (5) nephritis (6) reproductive disorders, e.g. abortion, stillbirths, mummies, etc, (7) Pia like lesions, normally known to be associated with *Lawsonia intracellularis* infections (Ileitis), (8) lymphadenopathy, (9) lymphoid depletion and/or (10) multinucleated/giant histiocytes (11) Porcine Dermatitis and Nephropathy Syndrome (PDNS), (12) PCVAD associated mortality, (13) PCVAD associated weight loss, (14) reduced growth variability (15) reduced frequency of 'runts' and (16) reduced co-infections with Porcine Reproductive and Respiratory Disease Complex (PRRSV). According to a further aspect, the present disclosure provides a method for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in young animals, comprising the step of administering an effective amount of a PCV2 antigen to that animal in need of such treatment, wherein those clinical symptoms are selected from the group consisting of: (1) interstitial pneumonia with interlobular edema, (2) cutaneous pallor or icterus, (3) mottled atrophic livers, (4) gastric ulcers, (5) nephritis (6) reproductive disorders, e.g. abortion, stillbirths, mummies, etc, (7) Pia like lesions, normally known to be associated with *Lawsonia intracellularis* infections (Ileitis), (8) lymphadenopathy, (9) lymphoid depletion and/or (10) multinucleated/giant histiocytes (11) Porcine Dermatitis and Nephropathy Syndrome (PDNS), (12) PCVAD associated mortality, (13) PCVAD associated weight loss, (14) reduced growth variability (15) reduced frequency of 'runts' and (16) reduced co-infections with Porcine Reproductive and Respiratory Disease Complex (PRRSV).

[0062] The composition according to the disclosure may be administered or applied, orally, intradermally, intratracheally, or intravaginally. The composition preferably may be administered or applied intramuscularly or intranasally, most preferably intramuscularly. In an animal body, it can prove advantageous to apply the pharmaceutical compositions as described above via an intravenous or by direct injection into target tissues. For systemic application, the intravenous, intravascular, intramuscular, intranasal, intraarterial, intraperitoneal, oral, or intrathecal routes are preferred. A more local application can be effected subcutaneously, intradermally, intracutaneously, intracardially, intralobally, intramedullarily, intrapulmonarily or directly in or near the tissue to be treated (connective-, bone-, muscle-, nerve-, epithelial tissue). Depending on the desired duration and effectiveness of the treatment, the compositions according to the disclosure may be administered once or several times, also intermittently, for instance on a daily basis for several days, weeks or months and in different dosages.

[0063] Preferably, one dose of the immunogenic composition as described above is intramuscularly administered to the subject in need thereof. According to a further aspect, the PCV2 antigen or the immunogenic composition comprising any such PCV2 antigen as described herein is bottled in and administered at one (1) mL per dose. Thus, according to a further aspect, the present disclosure also provides a 1 ml immunogenic composition, comprising PCV-2 antigen as described herein, for the treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in young animals, comprising the step of administering an effective amount of a

PCV2 antigen to that animal in need of such treatment. According to a further aspect, the present disclosure also provides a 1 ml immunogenic composition, comprising PCV-2 antigen as described herein, treatment or prophylaxis of a PCV2 infection or for reduction of clinical symptoms caused by or associated with a PCV2 infection in animals having anti-PCV2 antibodies, comprising the step of administering an effective amount of PCV2 antigen or an immunogenic composition comprising an PCV2 antigen to that animal in need of such treatment.

[0064] Preferably, the immunogenic composition is administered with an immune stimulant. Preferably, said immune stimulant is given at least twice. Preferably, at least 3 days, more preferably at least 5 days, even more preferably at least 7 days are in between the first and the second or any further administration of the immune stimulant. Preferably, the immune stimulant is given at least 10 days, preferably 15 days, even more preferably 20, and even more preferably at least 22 days beyond the initial administration of the immunogenic composition provided herein. A preferred immune stimulant is, for example, keyhole limpet hemocyanin (KLH), preferably emulsified with incomplete Freund's adjuvant (KLH/ICFA). However, it is herewith understood, that any other immune stimulant known to a person skilled in the art can also be used. The term "immune stimulant" as used herein, means any agent or composition that can trigger the immune response, preferably without initiating or increasing a specific immune response, for example the immune response against a specific pathogen. It is further instructed to administer the immune stimulant in a suitable dose.

[0065] The "animal" as used herein means swine, pig or piglet.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0066] The following examples set forth preferred materials and procedures in accordance with the present invention. It is to be understood, however, that these examples are provided by way of illustration only, and nothing therein should be deemed a limitation upon the overall scope of the invention.

EXAMPLE 1

Preparation of PCV2 ORF-2 antigen

[0067] Initial SF+ cell cultures from liquid nitrogen storage were grown in Excell 420 media (JRH Biosciences, Inc., Lenexa, KS) in suspension in sterile spinner flasks with constant agitation. The cultures were grown in 100 mL to 250mL spinner flasks with 25 to 150 mL of Excell 420 serum-free media. When the cells had multiplied to a cell density of $1.0 - 8.0 \times 10^6$ cells/mL, they were split to new vessels with a planting density of $0.5 - 1.5 \times 10^6$ cells/mL. Subsequent expansion cultures were grown in spinner flasks up to 36 liters in size or in stainless steel bioreactors of up to 300 liters for a period of 2-7 days at 25 - 29°C.

[0068] After seeding, the flasks were incubated at 27°C for four hours. Subsequently, each flask was seeded with a recombinant baculovirus containing the PCV2 ORF-2 gene (SEQ ID NO: 4). The recombinant baculovirus containing the PCV2 ORF-2 gene was generated as described in WO06/072065. After being seeded with the baculovirus, the flasks were then incubated at $27 \pm 2^\circ\text{C}$ for 7 days and were also agitated at 100 rpm during that time. The flasks used ventilated caps to allow for air flow.

[0069] After incubation, the resulting supernatant were harvested, filtered in order to remove cell debris and inactivated. The supernatant was inactivated by bringing its temperature to $37 \pm 2^\circ\text{C}$ and binary ethylenimine (BEI) is added to the supernatant to a final concentration of 5mM. The samples were then stirred continuously for 72 to 96 hrs. A 1.0 M sodium thiosulfate solution to give a final minimum concentration of 5 mM was added to neutralize any residual BEI. After inactivation, PCV2 ORF-2 buffered with phosphate buffer and Carbopol was added to about 0.5 to 2.5 mg/dose. The final dose comprises about 16 µg PCV2 ORF-2 antigen.

EXAMPLE 2

Anti PCV-2 Immuno assay

[0070] PK15 (e.g. ATCC CCL-33) or VIDO R1 cells described in WO 02/077210, are seeded onto a 96 well plate (about 20.000 to 60.000 cells per wells). Cells are infected with a PCV2 isolate, when monolayers are approximately 65 to 85% confluent. Infected cells are incubated for 48 hours. Medium is removed and wells are washed 2 times with PBS. The wash buffer is discarded and cells are treated with cold 50/50 methanol/acetone fixative ($\sim 100 \mu\text{l/well}$) for about 15 min at about -20°C . The fixative is discarded and the plates are air dried. Serial dilutions of porcine serum samples are prepared in PBS, added to the plates and incubated to allow antibodies to bind if present in the serum samples for about 1 hr at $36,5 \pm 1^\circ\text{C}$. In addition, serial dilutions of an anti-PCV2 positive and negative control sample (Positive Control and Negative Control Samples) are run in parallel. The plates are then washed three times with PBS. The PBS is discarded. Plates are then stained with an commercial Goat anti-Swine FITC conjugate diluted 1:100 in PBS and incubated for about

1 hr at $36.5 \pm 1^\circ\text{C}$, which allows detection of antibodies bound to infected cells. After incubation is complete, the microplates are removed from incubator, the conjugate is discarded and the plates are washed 2 times with PBS. The plates were read using UV microscopy and individual wells reported as positive or negative. The Positive Control and Negative Control samples are used to monitor the test system. If the controls are within expected ranges the test results are acceptable in regard to test method parameters. The serum antibody titers were calculated using the highest dilution showing specific IFA reactivity and the number of wells positive per dilution, or a 50% endpoint is calculated using the appropriate Reed-Muench formula.

EXAMPLE 3

Efficacy of PCV2 ORF-2 (Ingelvac® CircoFLEX™) in young animals having low or high anti-PCV2 antibodies

[0071] For investigation of a possible interference of the vaccine with the maternal antibody a study was conducted in which the antibody titers of all study animals were determined at the time of vaccination which were then grouped into a low, moderate and high antibody class: Geometric mean titers of $< 1:100$ were considered as low antibody titers, titers of $1:100$ to $1:1000$ were considered as moderate antibody titers and titers of $> 1:1000$ were considered as high antibody titers.

STUDY PERFORMANCE

[0072] Approximately 500 animals were included into the study. The study animals were balanced and equally distributed among both treatment groups with regard to initial body weight and litter assignment. At 20 days of age all study animals received a single dose (1 ml) of the PCV2 vaccine (Investigational Veterinary Product, IVP) or a single (1 ml) dose of a placebo containing adjuvanted cell culture supernatant (Control Product, CP) by intramuscular injection in the right side of the neck. Study animals were followed until the end of fattening. Blood samples from all study animals were collected and subsequently analyzed by IFAT in order to determine the antibody titers at the time of vaccination. Following this, the initial antibody titers were correlated with the weight gain. In addition, dependent on the initial antibody titer, animals were grouped into three classes (low, moderate and high initial antibody titers) and 'high titer animals' of both treatment groups were then compared for possible differences with regard to weight gain and viremia.

RESULTS

Initial Antibody Titers

[0073] At the time of vaccination the majority of animals had either moderate antibody titers (defined as $1:100$ to $1:1000$) or high antibody titers (defined as $> 1:1000$). Only approximately 13% percent of animals had low antibody titers (defined as $< 1:100$). Due to the absence of PCV2 infection at the time of study initiation it can be concluded that the antibody titers on study day 0 were possibly maternally derived. No significant differences in the antibody titers of study day 0 were observed between the two treatment groups. An overview about the percentage of animals per titer class is given in Figure 1.

Correlation of Antibody Titers at the Time of Vaccination with Viremia in Blood

[0074] In order to determine whether a high antibody titer at the time of vaccination ($> 1:1000$) had an impact on viremia, vaccinated and placebo-treated animals with high initial antibody titers were compared with regard to the onset, end, duration of viremia, the number of positive sampling days and the virus load. Table 1 summarizes the comparison of viremia parameters of the 'high-titer animals' from both treatment groups.

Table 1: Comparison of viremia in 'high titer animals' from both treatment groups

Investigated Parameter	Treatment Group	Number of pigs	Mean	Median	P
Onset of	CP	38	111.90 days	113.00 days	0.7843
Viremia	IVP CP-IVP	36	109.50 days 2.4 days	113.00 days	ns

(continued)

Investigated Parameter	Treatment Group	Number of pigs	Mean	Median	P
Duration of Viremia	CP	38	27.00 days	27.50 days	<0.0001 ***
	IVP	36	17.50 days	6.50 days	
	CP-IVP		9.50 days		
End of Viremia	CP	38	138.90 days	141.00 days	0.0033 **
	IVP	36	127.00 days	122.50 days	
	CP-IVP		11.9 days		
Positive Sampling days	CP	39	3.70 days	3.00 days	0.0082 **
	IVP	47	1.80 days	1.00 days	
	CP-IVP		1.9 days		
Mean Sum gE (log10)	CP	39	18.79 gE	17.21 gE	< 0.0001 ***
	IVP	47	9.12 gE	5.38 gE	
	CP-IVP		9.67 gE		

gE: sum of genomic equivalents per ml
P: p-value of the Wilcoxon Mann-Whitney test for comparisons between groups;
ns: not significant, p>0.05;
** significant, p ≤ 0.01;
*** significant, p < 0.001

[0075] Compared to the placebo-treated high-titer animals, vaccinated high-titer animals had a 9.5 day shorter duration of viremia, a 11.9 days earlier end of viremia, 1.9 days less viremic sampling days and an approximately 2-fold reduction of the sum of genomic equivalents/ml over the course of the study. These results indicate that also in the presence of high maternal antibody titers the IVP can still significantly reduce viremia in blood (end of viremia, duration of viremia, virus load).

Correlation of Antibody Titers at the Time of Vaccination with Weight Gain

[0076] It was next investigated, whether the initial antibody titer had any effect on the weight gain over the course of the study. Table 2 presents the correlation of the initial antibody titer with the weight gain at different time intervals as determined by the calculation of the Spearman rank coefficient and the p-value.

[0077] A statistically significant negative correlation between the antibody titer and the weight gain was found for both treatment groups at study weeks 0 to 7 indicating that a high maternal antibody titer negatively influences the weight gain development in the rearing phase. No other statistically significant correlations between the initial antibody titer and the weight gain during different time intervals were observed. It can therefore be concluded that the level of maternal antibody titer did not have an influence on the weight gain from 10 weeks of age (study week 7) onwards for neither the vaccinated or for the placebo-treated animals.

Table 2: Correlation of the PCV2 antibody titer at the time of vaccination with body weight gain over the course of the study

		Correlation of antibody titer at the time of vaccination with weight gain			
		Study week 0-7	Study week 7-12	Study week 12-17	Study week 17-22
CP	r	-0.09623	0.03501	-0.00521	-0.02774
	P	0.0086 **	0.3425 ns	0.8884 ns	0.4617 ns
	n	744	737	728	706
IVP	r	-0.09748	0.04309	-0.00954	0.02694
	P	0.0077 **	0.2440 ns	0.7974 ns	0.4710 ns
	n	746	733	727	718
r: Spearman rank correlation coefficient					
P: p-value of test on $r=0$: ns: not significant, $p > 0.05$,					

(continued)

	Correlation of antibody titer at the time of vaccination with weight gain			
	Study week 0-7	Study week 7-12	Study week 12-17	Study week 17-22
** significant, $p \leq 0.01$ n: Number of animals				

[0078] In line with this finding, no differences could be found with regard to the live body weight when comparing low and high titer animals of the vaccinated group. Figure 2 shows that the body weight after the onset of viremia (study week 17 and 22) was comparable irrespective of the level of initial antibody titer (Figure 2).

[0079] Furthermore vaccinated animals with a high antibody titer at the time of vaccination ($> 1:1000$) also showed a significantly higher body weight after the onset of viremia compared to placebo-treated animals with initial high antibody titers. As can be seen in Figure 3 the body weight (LSMean) at study week 17 and at study week 22 was indeed significantly higher in vaccinated 'high titer animals' (study week 17: 1.55 kg, $p = 0.0328$; study week 22: 3.06 kg, $p = 0.0007$) than in placebo-treated 'high titer animals'. Together these findings demonstrate that there is no interference of the IVP with the antibody titer at the time of vaccination.

Conclusion

[0080] For analysis of a possible maternal antibody interference the initial antibody titer was correlated with the two efficacy parameters viremia in blood and live body weight. Compared to the placebo-treated 'high titer animals' the following statistical significant findings were noted for the vaccinated 'high titer animals':

- reduction in loss of weight gain
- shorter duration of viremia and earlier end of viremia
- lower virus load

EXAMPLE 4

Efficacy of PCV2 ORF-2 (Ingelvac® CircoFLEX™) in young animals having anti-PCV2 antibodies with respect to lymphoid depletion, lymphoid inflammation, and lymphoid immunohistochemistry (IHC)

[0081] The objective of this blinded vaccination-challenge study was to evaluate at what age pigs vaccinated with Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector established immunity in the presence of Porcine Circovirus Type 2 (PCV2) maternally-derived antibodies. Three primary parameters were analyzed following challenge. These three parameters included lymphoid depletion, lymphoid inflammation, and lymphoid immunohistochemistry (IHC). To demonstrate immunity in the presence of PCV2 maternally-derived antibodies, conventionally raised pigs vaccinated with PCV2 vaccine at 3 weeks of age or at 8 weeks of age, must demonstrate statistically significant differences ($p < 0.05$) for lymphoid depletion, lymphoid inflammation, and lymphoid IHC, compared with challenge control pigs treated with Control Product at 3 weeks of age.

STUDY PERFORMANCE

[0082] One hundred twenty (120) conventionally raised pigs, 21 days of age on Day 0 (DO), were assigned completely at random to one of five treatment groups. On D0, blood samples were collected from all pigs,

- Group 1a was treated with Investigational Veterinary Product (IVP; PCV2 reference vaccine) at 3 weeks of age.
- Group 1b was treated with Investigational Veterinary Product (IVP; PCV2 reference vaccine) at 8 weeks of age.
- Group 2 was treated with Control Product (CP) at 3 weeks of age.

[0083] Pigs were observed for clinical assessments post-vaccination from D-1 to D59. Additional pre-challenge blood samples were collected on D14, D28, D42, D56 and D63. A summary of Group PCV2 serological Geometric Mean Titers (GMT) pre-challenge are shown below in Table 3.

Table 3: Group PCV2 Serological Geometric Mean Titers Pre-challenge

Group-Treatment	PCV2 Serology - GMT					
	D0	D14	D28	D42	D56	D63
Group 1a IVP administered 3 weeks of age	556.5	252.8	142.0	56.2	32.0	51.3
Group 1b IVP administered at 8 weeks of age	476.2	308.2	151.6	36.2	29.3	48.3
Group 2 CP administered at 3 weeks of age	513.8	310.7	134.3	36.9	16.9	24.5

[0084] All remaining pigs received 2.0 mL of keyhole limpet hemocyanin (KLH) emulsified in incomplete Freund's adjuvant (ICFA) IM on D60 (Day Post-Challenge (DPC) -3) and D66 (DPC 3). On D63 (DPC 0), remaining pigs received 1.0 mL of PCV2 Iowa State University Veterinary Diagnostic Laboratory (ISUVDL) challenge material (4.75 log₁₀ TCID₅₀/mL) IM and 1.0 mL of the same material IN. Body weights, rectal temperatures, clinical observations, blood samples and nasal swabs were collected on the day of challenge and periodically post-challenge. At necropsy for each pig, gross lesions were noted and lung and lymphoid tissue samples were collected. Lung and lymphoid tissues were examined microscopically by ISUVDL for lesions and for the presence of PCV2 antigen by IHC testing. A general description of the challenge phase of the study is shown below in table 4.

Table 4: Challenge Phase of Study

Group - Treatment	Number	KLH/ICFA On D60 (DPC -3)	PCV2 Challenge on D63 (DPC 0)	KLH/ ICFA On D66 (DPC 3)	Day of Necropsy
Group 1a IVP administered 3 weeks of age	20	Yes	Yes	Yes	D87 (DPC 24) or D88 (DPC 25)
Group 1b IVP administered at 8 weeks of age	21	Yes	Yes	Yes	D87 (DPC 24) or D88 (DPC 25)
Group 2 CP administered at 3 weeks of age	20	Yes	Yes	Yes	D87 (DPC 24) or D88 (DPC 25)

[0085] On D86, the geometric mean titers were 906.6, 2447.1, 2014.9, respectively.

RESULTS

[0086] Following PCV2 challenge exposure on D63 and subsequent necropsy, Group 1a had a statistically significant lower proportion of pigs positive for lymphoid depletion (**p=0.0084**), a lower proportion of pigs positive for lymphoid inflammation (**p=0.0079**), and a lower proportion of pigs with IHC lymphoid-positive tissues (**p=0.0031**), all in comparison to Group 2. Following PCV2 challenge, Group 1b had a statistically significant lower proportion of pigs positive for lymphoid depletion (**p=0.0148**), a lower proportion of pigs positive for lymphoid inflammation (**p=0.0036**), and a lower proportion of pigs with IHC lymphoid-positive tissues (**p=0.0013**), all in comparison to Group 2. A summary of primary efficacy parameter results for Groups 1a, 1b and 2 are shown below in table 5.

Table 5: Summary of Primary Efficacy Parameter Results for Groups 1a and 1b compared with Group 2

Group-Treatment	PCV2 Serological status on Day 0	Lymphoid Depletion (+/total)	Lymphoid Inflammation (+/total)	Lymphoid IHC (+/total)
Group 1a IVP at 3 weeks of age	Seropositive	1/20 (5%) *p=0.0084	3/20(15%) *p=0.0079	3/20 (15%) *p=0.0031
Group 1b IVP at 8 weeks of age	Seropositive	2/21 (9.5%) *p=0.0148	3/21 (14.3%) *p=0.0036	3/21 (14.3%) *p=0.0013

(continued)

Group-Treatment	PCV2 Serological status on Day 0	Lymphoid Depletion (+/total)	Lymphoid Inflammation (+/total)	Lymphoid IHC (+/total)
Group 2 CP at 3 weeks of age	Seropositive	9/20 (45%)	12/20 (60%)	13/20 (65%)
* p value compared with Group 2 - Fisher's Exact Test				

[0087] There were significant differences between Groups 1a and 1b compared with Group 2 for microscopic lung inflammation ($p < 0.0407$), but no significant differences between these groups for lung tissue testing positive for PCV2 antigen by IHC testing ($p \geq 0.2317$). There were no significant differences between Groups 1a and 1b compared with Group 2 for clinical assessments post-vaccination, ADG, clinical signs post-challenge, pyrexia, nasal shedding of PCV2, % total lung scores and lymphadenopathy.

[0088] In conclusion, Group 1a, vaccinated at 3 weeks of age and having a GMT of 556.6 at the time of vaccination, was significantly protected from lymphoid depletion, lymphoid inflammation, and lymphoid tissues testing positive for PCV2 antigen by IHC testing, compared with Group 2. Group 1b, vaccinated at 8 weeks of age and having a GMT of 151.6 one week prior to vaccination, was significantly protected from lymphoid depletion, lymphoid inflammation and lymphoid tissues testing positive for PCV2 antigen by IHC testing, compared with Group 2. Pigs with PCV2 maternally-derived antibodies were protected from Porcine Circovirus Associated Disease (PCVAD) when vaccinated as early as 3 weeks of age.

SEQUENCE LISTING

[0089]

<110> BOEHRINGER INGELHEIM VETMEDICA GMBH

<120> TREATMENT OF PIGS WITH PCV2 ANTIGEN

<130> P01-2169

<160> 11

<170> PatentIn version 3.3

<210> 1

<211> 8

<212> DNA

<213> Artificial

<220>

<223> This is a modified Kozak's sequence.

<400> 1

ccgccatg 8

<210> 2

<211> 6

<212> DNA

<213> Artificial

<220>

<223> This is a recombinant Eco R1 sequence.

<400> 2

gaattc 6

<210> 3

<211> 713

5 <212> DNA

<213> Porcine circovirus

<400> 3

```

10      cagctatgac gtatccaagg aggcgttacc gcagaagaag acaccgcccc cgcagccatc      60
      ttggccagat cctccgccgc cgcccctggc tcgtccaccc ccgccaccgc taccgttgga      120
      gaaggaaaaa tggcatcttc aacacccgcc tctcccgcac ctccggatat actgtggaga      180
      aggaaaaatg gcatcttcaa caccgcctc tcccgcacct tcggatatac tgtgacgact      240
      ttgttcccc  gggagggggg accaacaaaa tctctatacc ctttgaatac tacagaataa      300
15      gaaagggttaa gggtgaattc tggccctgct ccccatcac ccagggtgat aggggagtgg      360
      gctccactgc tgttattcta gatgataact ttgtaacaaa ggccacagcc ctaacctatg      420
      acccatatgt aaactactcc tcccgccata caatcccca acccttctcc taccactccc      480
      gttacttcac acccaaacct gttcttgact ccactattga ttacttccaa ccaaataaca      540
      aaaggaatca gctttggctg aggtacaaa cctctagaaa tgtggaccac gtaggcctcg      600
20      gcaactgcgtt cgaaaacagt aaatacgacc aggactacaa tatccgtgta accatgtatg      660
      tacaattcag agaatttaat cttaaagacc cccacttaa accctaaatg aat      713

```

<210> 4

<211> 713

25 <212> DNA

<213> Porcine circovirus

<400> 4

```

30      ccgccatgac gtatccaagg aggcgttacc gcagaagaag acaccgcccc cgcagccatc      60
      ttggccagat cctccgccgc cgcccctggc tcgtccaccc ccgccaccgc taccgttgga      120
      gaaggaaaaa tggcatcttc aacacccgcc tctcccgcac ctccggatat actgtcaagg      180
      ctaccacagt cacaacgccc tcctgggcgg tggacatgat gagatttaat attgacgact      240
      ttgttcccc  gggagggggg accaacaaaa tctctatacc ctttgaatac tacagaataa      300
35      gaaagggttaa gggtgaattc tggccctgct ccccatcac ccagggtgat aggggagtgg      360
      gctccactgc tgttattcta gatgataact ttgtaacaaa ggccacagcc ctaacctatg      420
      acccatatgt aaactactcc tcccgccata caatcccca acccttctcc taccactccc      480
      gttacttcac acccaaacct gttcttgact ccactattga ttacttccaa ccaaataaca      540
      aaaggaatca gctttggctg aggtacaaa cctctagaaa tgtggaccac gtaggcctcg      600
40      gcaactgcgtt cgaaaacagt aaatacgacc aggactacaa tatccgtgta accatgtatg      660
      tacaattcag agaatttaat cttaaagacc cccacttga accctaagaa ttc      713

```

<210> 5

<211> 233

45 <212> PRT

<213> Porcine circovirus

<400> 5

50

55

Met Thr Tyr Pro Arg Arg Arg Tyr Arg Arg Arg Arg His Arg Pro Arg
 1 5 10 15
 Ser His Leu Gly Gln Ile Leu Arg Arg Arg Pro Trp Leu Val His Pro
 5 20 25 30
 Arg His Arg Tyr Arg Trp Arg Arg Lys Asn Gly Ile Phe Asn Thr Arg
 35 40 45
 Leu Ser Arg Thr Phe Gly Tyr Thr Val Lys Ala Thr Thr Val Thr Thr
 10 50 55 60
 Pro Ser Trp Ala Val Asp Met Met Arg Phe Asn Ile Asp Asp Phe Val
 65 70 75 80
 Pro Pro Gly Gly Gly Thr Asn Lys Ile Ser Ile Pro Phe Glu Tyr Tyr
 15 85 90 95
 Arg Ile Arg Lys Val Lys Val Glu Phe Trp Pro Cys Ser Pro Ile Thr
 100 105 110
 Gln Gly Asp Arg Gly Val Gly Ser Thr Ala Val Ile Leu Asp Asp Asn
 115 120 125
 Phe Val Thr Lys Ala Thr Ala Leu Thr Tyr Asp Pro Tyr Val Asn Tyr
 130 135 140
 Ser Ser Arg His Thr Ile Pro Gln Pro Phe Ser Tyr His Ser Arg Tyr
 145 150 155 160
 Phe Thr Pro Lys Pro Val Leu Asp Ser Thr Ile Asp Tyr Phe Gln Pro
 165 170 175
 Asn Asn Lys Arg Asn Gln Leu Trp Leu Arg Leu Gln Thr Ser Arg Asn
 180 185 190
 Val Asp His Val Gly Leu Gly Thr Ala Phe Glu Asn Ser Lys Tyr Asp
 195 200 205
 Gln Asp Tyr Asn Ile Arg Val Thr Met Tyr Val Gln Phe Arg Glu Phe
 210 215 220
 Asn Leu Lys Asp Pro Pro Leu Lys Pro
 225 230

<210> 6

<211> 233

<212> PRT

<213> Porcine circovirus

<400> 6

Met Thr Tyr Pro Arg Arg Arg Tyr Arg Arg Arg Arg His Arg Pro Arg
 1 5 10 15
 5 Ser His Leu Gly Gln Ile Leu Arg Arg Arg Pro Trp Leu Val His Pro
 20 25 30
 Arg His Arg Tyr Arg Trp Arg Arg Lys Asn Gly Ile Phe Asn Thr Arg
 35 40 45
 10 Leu Ser Arg Thr Phe Gly Tyr Thr Val Lys Ala Thr Thr Val Thr Thr
 50 55 60
 Pro Ser Trp Ala Val Asp Met Met Arg Phe Asn Ile Asp Asp Phe Val
 65 70 75 80
 15 Pro Pro Gly Gly Gly Thr Asn Lys Ile Ser Ile Pro Phe Glu Tyr Tyr
 85 90 95
 Arg Ile Arg Lys Val Lys Val Glu Phe Trp Pro Cys Ser Pro Ile Thr
 100 105 110
 20 Gln Gly Asp Arg Gly Val Gly Ser Thr Ala Val Ile Leu Asp Asp Asn
 115 120 125
 Phe Val Thr Lys Ala Thr Ala Leu Thr Tyr Asp Pro Tyr Val Asn Tyr
 130 135 140
 25 Ser Ser Arg His Thr Ile Pro Gln Pro Phe Ser Tyr His Ser Arg Tyr
 145 150 155 160
 30 Phe Thr Pro Lys Pro Val Leu Asp Ser Thr Ile Asp Tyr Phe Gln Pro
 165 170 175
 Asn Asn Lys Arg Asn Gln Leu Trp Leu Arg Leu Gln Thr Ser Arg Asn
 180 185 190
 35 Val Asp His Val Gly Leu Gly Thr Ala Phe Glu Asn Ser Lys Tyr Asp
 195 200 205
 Gln Asp Tyr Asn Ile Arg Val Thr Met Tyr Val Gln Phe Arg Glu Phe
 210 215 220
 40 Asn Leu Lys Asp Pro Pro Leu Glu Pro
 225 230

<210> 7

45 <211> 756

<212> DNA

<213> Artificial

<220>

50 <223> This sequence is from porcine circovirus type 2, open reading frame 2, together with a portion from the pGEM T-easy vector.

<400> 7

55

	gcggccgcgg	gaattcgatc	cgccatgacg	tatccaagga	ggcgttaccg	cagaagaaga	60
	caccgcccc	gcagccatct	tggccagatc	ctccgcccgc	gcccctggct	cgtdccacccc	120
	cgccaccgct	accgttgagg	aaggaaaaat	ggcatcttca	acaccgcgct	ctcccgcacc	180
5	ttcggatata	ctgtcaaggc	taccacagtc	acaacgcctt	cctggggcgg	ggacatgatg	240
	agattttaata	ttgacgactt	tgttcccccg	ggagggggga	ccaacaaaat	ctctataccc	300
	tttgaatact	acagaataag	aaagggttaag	gttgaattct	ggccctgctc	ccccatcacc	360
	caggggtgata	ggggagtggg	ctccactgct	gttattctag	atgataactt	tgtaacaaaag	420
	gccacagccc	taacctatga	cccatatgta	aactactcct	cccgccatac	aatcccccaa	480
	cccttctcct	accactcccg	ttacttcaca	cccaaacctg	ttcttgactc	cactattgat	540
10	tacttccaac	caaataacaa	aaggaatcag	ctttggctga	ggctacaaac	ctctagaaat	600
	gtggaccacg	taggcctcgg	cactgcgttc	gaaaacagta	aatacgacca	ggactacaat	660
	atccgtgtaa	ccatgtatgt	acaattcaga	gaatttaatc	ttaaagacc	cccacttgaa	720
	ccctaagaat	tctatcacta	gtgaattcgc	ggccgc			756
15	<210> 8						
	<211> 10387						
	<212> DNA						
	<213> Artificial						
20	<220>						
	<223> This is the porcine circovirus type 2, ORF-2construct, which includes baculovirus and pGEM T-easy coding sequences.						
25	<400> 8						
	aagctttact	cgtaaagcga	gttgaaggat	catatttagt	tgcgtttatg	agataagatt	60
	gaaagcacgt	gtaaaatggt	tcccgcgcgt	tggcacaact	atttacaatg	cggccaagtt	120
	ataaaagatt	ctaactctgat	atgtttttaa	acacctttgc	ggcccgagtt	gtttgcgtac	180
	gtgactagcg	aagaagatgt	gtggaccgca	gaacagatag	taaaacaaaa	ccctagtatt	240
30	ggagcaataa	tcgatttaac	caacacgtct	aaatattatg	atggtgtgca	ttttttgcgg	300
	gcgggcctgt	tatacaaaaa	aattcaagta	cctggccaga	ctttgccgcc	tgaaagcata	360
	gttcaagaat	ttattgacac	ggtaaaagaa	tttacagaaa	agtgtcccgg	catgttgggtg	420
	ggcgtgcact	gcacacacgg	tattaatcgc	accggttaca	tgggtgtgcag	atattttaatg	480
	cacaccctgg	gtattgcgcc	gcaggaagcc	atagatagat	tcgaaaaagc	cagaggtcac	540
35	aaaattgaaa	gacaaaatta	cgttcaagat	ttattaattt	aattaatatt	atltgcattc	600
	tttaacaaat	actttatcct	atlttcaa	tgttgcgctt	cttcacagca	acaaaaacta	660
	tgttcgctt	gctccgttta	gcttgtagcc	gatcagtggc	gttgttccaa	tcgacggtag	720
	gattaggccg	gatattctcc	accacaatgt	tggcaacggt	gatgttacgt	ttatgctttt	780
	ggttttccac	gtacgtcttt	tggccggtaa	tagccgtaaa	cgtagtgccg	tcgcgcgtca	840
	cgcacaacac	cggatgtttg	cgttgtccg	cggggtattg	aaccgcgcga	tccgacaaat	900
40	ccaccacttt	ggcaactaaa	tcggtgacct	gcgcgtcttt	tttctgcatt	atltcgtctt	960
	tcttttgcat	ggtttccctg	aagccggtgt	acatgcggtt	tagatcagtc	atgacgcgcg	1020
	tgacctgcaa	atctttggcc	tcgatctgct	tgtccttgat	ggcaacgatg	cgttcaataa	1080
	actcttgttt	tttaacaagt	tcctcggttt	tttgcgccac	caccgcttgc	agcgcgtttg	1140
	tgtgctcgg	gaatgtcgca	atcagcttag	tcaccaactg	tttgcctctc	tcctcccgtt	1200
45	gtttgatcgc	gggatcgtag	ttgccgggtgc	agagcacttg	aggaattact	tcttctaaaa	1260
	gccattcttg	taattctatg	gcgtaaggca	atlttgactt	cataatcagc	tgaatcacgc	1320
	cggatttagt	aatgagcact	gtatgcggct	gcaaatacag	cgggtcgccc	cttttcacga	1380
	cgctgttaga	ggtagggccc	ccatttttga	tggctctgctc	aaataacgat	ttgtatttat	1440
	tgtctacatg	aacacgtata	gctttatcac	aaactgtata	ttttaaactg	ttagcgacgt	1500
	ccttggccac	gaaccggacc	tgttggctgc	gctctagcac	gtaccgcagg	ttgaacgtat	1560
50	cttctccaaa	tttaaatctt	ccaattttta	cgcgagccat	tttgatacac	gtgtgtcgat	1620

55

	tttgcaacaa	ctattgtttt	ttaacgcaaa	ctaaacttat	tgtggtaagc	aataattaaa	1680
	tatgggggaa	catgcgcgc	tacaacactc	gtcgttatga	acgcagacgg	cgccggtctc	1740
	ggcgcaagcg	gctaaaaagt	gttgcgcggt	caacgcggca	aacatcgcaa	aagccaatag	1800
	tacagttttg	atthgcatat	taacggcgat	tttttaaat	atcttattta	ataaatagtt	1860
5	atgacgccta	caactccccg	cccgcggtga	ctcgctgcac	ctcgagcagt	tcgttgacgc	1920
	cttctctcgt	gtggccgaac	acgtcgagcg	ggtaggtcgat	gaccagcggc	gtcccgcaacg	1980
	cgacgcacaa	gtatctgtac	accgaatgat	cgtcgggcga	aggcacgtcg	gcctccaagt	2040
	ggcaatattg	gcaaattcga	aaatatatac	agttgggttg	tttgcgcata	tctatcgtgg	2100
	cgttgggcat	gtacgtccga	acgttgattt	gcatgcaagc	cgaaattaaa	tcattgcgat	2160
10	tagtgcgatt	aaaacgttgt	acatcctcgc	ttttaatcat	gccgtcgatt	aaatcgcgca	2220
	atcgagtcaa	gtgatcaaa	gtgggaataa	tgttttcttt	gtattcccga	gtcaagcgca	2280
	gcgcgtatth	taacaaacta	gccatcttgt	aagttagttt	catttaaatgc	aactttatcc	2340
	aataatatat	tatgtatcgc	acgtcaagaa	ttacaatgc	gccggttgtc	gcatctcaac	2400
	acgactatga	tagagatcaa	ataaagcgcg	aattaaatag	cttgcgacgc	aacgtgcacg	2460
	atctgtgcac	gcgttccggc	acgagctttg	attgtaataa	gtttttacga	agcgatgaca	2520
15	tgacccccgt	agtgacaacg	atcacgcca	aaagaactgc	cgactacaaa	attaccgagt	2580
	atgtcgggtga	cgttaaaact	attaagccat	ccaatcgacc	gttagtcgaa	tcaggaccgc	2640
	tggtagcgaga	agccgcgaag	tatggcgaat	gcattcgtata	acgtgtggag	tcgcgtcatt	2700
	agagcgtcat	gtttagacaa	gaaagctaca	tatttaattg	atcccgatga	ttttattgat	2760
	aaattgaccc	taactccata	cacggtattc	tacaatggcg	gggttttggt	caaaatttcc	2820
	ggactgcgat	tgtacatgct	gttaacggct	ccgccacta	ttaatgaaat	taaaaattcc	2880
20	aattttaaaa	aacgcagcaa	gagaaacatt	tgtatgaaag	aatgcgtaga	aggaaagaaa	2940
	aatgtcgtcg	acatgctgaa	caacaagatt	aatatgcctc	cgtgtataaa	aaaaatattg	3000
	aacgattttga	aagaaaacaa	tgtaccgcgc	ggcggatgtg	acagggaagag	gtttataacta	3060
	aactgtttaca	ttgcaaacgt	ggtttcgtgt	gccaaagtgtg	aaaaccgatg	tttaatacaag	3120
	gctctgacgc	atttctacaa	ccacgactcc	aagtgtgtgg	gtgaagtcat	gcattctttta	3180
25	atcaaatccc	aagatgtgta	taaaccacca	aactgccaaa	aatgaaaac	tgtcgacaag	3240
	ctctgtccgt	ttgctggcaa	ctgcaagggt	ctcaatccta	tttgtaatta	ttgaataata	3300
	aaacaattat	aaatgctaaa	tttgtttttt	attaacgata	caaaccacaa	gcaacaagaa	3360
	catttgtagt	attatctata	attgaaaacg	cgtagttata	atcgctgagg	taataatttaa	3420
	aatcattttc	aaatgattca	cagtttaattt	gcgacaatat	aattttattt	tcacataaac	3480
	tagacgcctt	gtcgtcttct	tcttcgtatt	ccttctcttt	ttcatatttt	tcctcataaa	3540
30	aattaacata	gttattatcg	tatccatata	tgtatctatc	gtatagagta	aattttttgt	3600
	tgtcataaat	atatatgtct	tttttaattg	ggtgtatagt	acgcgtgcgc	atagtttttc	3660
	tgtaatattac	aacagtgtct	ttttctggta	gttcttcgga	gtgtgttgct	tttaattatta	3720
	aattttatata	atcaatgaat	ttgggacgt	cggttttgta	caatatgttg	ccggcatagt	3780
	acgcagcttc	ttctagtcca	attacaccat	tttttagcag	caccggatta	acataacttt	3840
	ccaaaatggt	gtacgaaccg	ttaaacaaaa	acagttcacc	tcctttttct	atactattgt	3900
35	ctgcgagcag	ttgtttgttg	ttaaaaataa	cagccattgt	aatgagacgc	acaaactaat	3960
	atcacaaact	ggaaatgtct	atcaatatat	agttgtctgat	atcatggaga	taattaaaaa	4020
	gataaccatc	tcgcaaataa	ataagtattt	tactgttttc	gtaacagttt	tgtaataaaa	4080
	aaacctataa	atattccgga	ttattcatac	cgtcccacca	tcggggcgcg	atcagatctg	4140
	cagcgccgcg	gggaattcga	tcgcgccatga	cgtatccaag	gaggcggttac	cgcagaagaa	4200
40	gacaccgccc	ccgcagccat	cttgcccaga	tcctccgcgc	ccgcccctgg	ctcgtccacc	4260
	cccgccaccg	ctaccgttgg	agaaggaaaa	atggcatctt	caacaccgcg	ctctcccgca	4320
	ccttcggata	tactgtcaag	gctaccacag	tcacaacgcc	ctcctgggcg	gtggacatga	4380
	tgagattttaa	tattgacgac	tttgttcccc	cgggaggggg	gaccaacaaa	atctctatag	4440
	cctttgaata	ctacagaata	agaaagggtta	aggttgaatt	ctggccctgc	tcctccatca	4500
	cccagggtga	taggggagtg	ggctccactg	ctgttattct	agatgataac	tttgtaacaa	4560
45	aggccacagc	cctaacctat	gacccatattg	ttaaactactc	ctcccgccat	acaatcccc	4620
	aacccttctc	ctaccactcc	cgttacttca	cacccaaacc	tgttcttgac	tcactatttg	4680
	attacttcca	accaaataac	aaaaggaatc	agctttggct	gaggctacaa	acctctagaa	4740
	atgtggacca	cgtaggcctc	ggcactgcgt	tcgaaaacag	taaatacgac	caggactaca	4800
	atatccgtgt	aacctatgat	gtacaattca	gagaatttaa	tcttaaagac	ccccacttg	4860
	aaccctaaga	attctatcac	tagtgaattc	gcggccgcgc	gccgctccag	aattctagaa	4920
50	ggtacccggg	atcctttcct	gggacccggc	aagaacaaaa	aactcactct	cttcaaggaa	4980
	atccgtaatg	ttaaacccga	cacgatgaag	cttgctgttg	gatggaaagg	aaaagagttc	5040
	tacagggaaa	cttggaaccg	cttcatggaa	gacagcttcc	ccattgttaa	cgaccaagaa	5100
	gtgatggatg	ttttccttgt	tgtcaacatg	cgtcccacta	gacccaaccg	ttgttacaaa	5160
	ttcctggccc	aacacgctct	gcgttgcgac	cccagactatg	tacctcatga	cgtgattagg	5220
55	atcgctcgagc	cttcatgggt	gggcagcaac	aacgagtacc	gcacagcct	ggctaagaag	5280

	ggcggcggct	gcccataat	gaaccttcac	tctgagtaca	ccaactcgtt	cgaacagttc	5340
	atcgatcgtg	tcatctggga	gaacttctac	aagcccatcg	tttacatcgg	taccgactct	5400
	gctgaagagg	aggaaattct	ccttgaagtt	tccctgggtg	tcaaagtaaa	ggagtttgca	5460
	ccagacgcac	ctctgttcac	tgggtccggcg	tattaaaaca	cgatacattg	ttatttagtac	5520
5	atltattaag	cgctagattc	tgtgcgttgt	tgattttacag	acaattggtg	tacgtatttt	5580
	aataattcat	taaaattata	atcttttaggg	tgggtatgta	gagcgaaaat	caaatgattt	5640
	tcagcgtctt	tatatctgaa	tttaaatatt	aaatcctcaa	tagatttgta	aaataggttt	5700
	cgattagttt	caaacaaggg	ttgtttttcc	gaaccgatgg	ctggactatc	taatggattt	5760
	tcgctcaacg	ccacaaaact	tgccaaatct	tgtagcagca	atctagcttt	gtcgatattc	5820
10	gtttgtgttt	tgttttgtaa	taaagggttcg	acgtcgttca	aaatattatg	cgcttttgta	5880
	tttctttcat	cactgtcgtt	agtgtacaat	tgactcgacg	taaacacggt	aaataaagct	5940
	tggacatatt	taacatcggg	cgtgttagct	ttattaggcc	gattatcgct	gtcgtcccaa	6000
	ccctgctcgt	tagaagttgc	ttccgaagac	gatttttgcca	tagccacacg	acgcctatta	6060
	attgtgtcgg	ctaacacgtc	cgcgatcaaa	ttttagattg	agctttttgg	aattattttct	6120
	gattgcgggc	gtttttgggc	gggtttcaat	otaactgtgc	ccgattttta	ttcagacaa	6180
15	acgttagaaa	gcgatggtgc	aggcgggtgg	aacatttcag	acggcaaatc	tactaatggc	6240
	ggcgggtggtg	gagctgatga	taaatctacc	atcgggtggag	gcgcaggcgg	ggctggcggc	6300
	ggaggcggag	gcggaggtgg	tggcgggtgat	gcagacggcg	gttttaggctc	aaatgtctct	6360
	ttaggcaaca	cagtcggcac	ctcaactatt	gtactggttt	cgggcgcgct	ttttggtttg	6420
	accggtctga	gacgagtgcg	atttttttcgt	tttctaatag	cttccaacaa	gttggtctctg	6480
	tcgtctaaag	gtgcagcggg	ttgaggttcc	gtcggcattg	gtggagcggg	cggcaattca	6540
20	gacatcgatg	gtggtggtgg	tgggtggaggg	gctggaatgt	taggcacggg	agaaggtggt	6600
	ggcggcgggtg	ccgccgtat	aatttgttct	ggtttagttt	gttcgcgcac	gattgtgggc	6660
	accggcgcag	gcgccgctgg	ctgcacaacg	gaaggctcgtc	tgcttcgagg	cagcgtctgg	6720
	ggtggtggca	attcaatatt	ataattggaa	tacaaatcgt	aaaaatctgc	tataagcatt	6780
	gtaatttcgc	tatcgtttac	cgtgccgata	tttaacaacc	gctcaatgta	agcaattgta	6840
	ttgtaaagag	attgtctcaa	gctcgcgcga	atctcgataac	aagccttttc	atttttacta	6900
25	cagcattgta	gtggcgagac	acttcgctgt	cgtcgacgta	catgtatgct	ttgttgtcaa	6960
	aaacgtcgtt	ggcaagcttt	aaaatattta	aaagaacatc	tctgttcagc	accactgtgt	7020
	tgtcgtaaat	gttgtttttg	ataatttgcg	cttcgcgagt	atcgacacgt	tcaaaaaatt	7080
	gatgcgcac	aattttgttg	ttcctattat	tgaataaata	agattgtaca	gattcatatc	7140
	tacgattcgt	catggccacc	acaaatgcta	cgtcgcaaac	gctggtacaa	ttttacgaaa	7200
30	actgcaaaaa	cgtcaaaaact	cgggtataaaa	taattcaacg	gcgctttggc	aaaaatatca	7260
	ttttatcgca	caagccact	agcaaatgtg	atltgcagaa	aacaatttcg	gcgcacaatt	7320
	ttaacgctga	cgaataaaaa	gttcaccagt	taatgagcga	ccacccaaat	tttataaaaa	7380
	tctattttta	tcacggttcc	atcaacaacc	aagtgatcgt	gatggactac	attgactgtc	7440
	ccgattttatt	tgaacacta	caaattaaag	gcgagctttc	gtaccaactt	gttagcaata	7500
	ttattagaca	gctgtgtgaa	gcgctcaacg	atltgcacaa	gcacaatttc	atacacaacg	7560
35	acataaaaact	cgaatatgtc	ttatatlttcg	aagcacttga	tcgcgtgtat	gtttgcgatt	7620
	acggattgtg	caaacacgaa	aactcactta	gcgtgcacga	cggcacgttg	gagtatttta	7680
	gtccggaaaa	aattcgacac	acaactatgc	acgttttcgtt	tgactggtac	gcggcgtggt	7740
	aacatacaag	ttgctaacgt	aatcatggtc	atagctgttt	cctgtgtgaa	attgttatoc	7800
	gctcacaaat	ccacacaaca	tacgagccgg	aagcataaag	tgtaaagcct	ggggtgccta	7860
40	atgagtgagc	taactcacat	taattgcgtt	gcgctcactg	cccgttttcc	agtcgggaaa	7920
	cctgtcgtgc	cagctgcatt	aatgaatcgg	ccaacgcgcg	gggagaggcg	gtttgcggtat	7980
	tgggcgtctc	tccgcttcc	cgtcactga	ctcgtcgcgc	tcggtcgttc	ggctgcggcg	8040
	agcggtatca	gctcactcaa	aggcggtaat	acggttatcc	acagaatcag	gggataacgc	8100
	aggaaagaac	atgtgagcaa	aaggccagca	aaaggccagg	aaccgtaaaa	aggccgcgtt	8160
	gctggcgttt	ttccataggc	tccgcccccc	tgacgagcat	cacaaaaatc	gacgtcaag	8220
	tcagaggtgg	cgaataccga	caggactata	aagataccag	gcgtttcccc	ctggaagctc	8280
45	cctcgtgcgc	tctcctgttc	cgaccctgcc	gcttaccgga	tacctgtccg	cctttctccc	8340
	ttcgggaagc	gtggcgcttt	ctcatagctc	acgctgtagg	tatctcagtt	cgggtgtaggt	8400
	cgttcgctcc	aagctgggct	gtgtgcacga	acccccggtt	cagcccgacc	gctgcgcctt	8460
	atccggtaac	tatcgtcttg	agtccaaccc	ggtaagacac	gacttatcgc	cactggcagc	8520
	agccactggg	aacaggatta	gcagagcgag	gtatgtaggc	ggtgctacag	agttccttgaa	8580
50	gtgggtggcct	aactacggct	acactagaag	gacagtattt	ggtatctcgc	ctctgctgaa	8640
	gccagttacc	ttcggaaaaa	gagttggtag	ctcttgatcc	ggcaaacaaa	ccaccgctgg	8700
	tagcgggtgg	ttttttgttt	gcaagcagca	gattacgcgc	agaaaaaaag	gatctcaaga	8760
	agatccctttg	atcttttcta	cggggctctga	cgtcagtggt	aacgaaaact	caggttaagg	8820
	gatttttggtc	atgagattat	caaaaaggat	cttcacctag	atccttttaa	attaaaaatg	8880
55	aagttttaaa	tcaatctaaa	gtatatatga	gtaaaacttg	tctgacagtt	accaatgctt	8940

```

    aatcagtgag gcacctatct cagcgatctg tctatttcgt tcatccatag ttgcctgact 9000
    ccccgctcgtg tagataacta cgatacggga gggcttacca tctggcccca gtgctgcaat 9060
    gataccgcga gacccacgct caccggctcc agatttatca gcaataaacc agccagccgg 9120
    aagggccgag cgcagaagtg gtcctgcaac tttatccgcc tccatccagt ctattaattg 9180
5    ttgccgggaa gctagagtaa gtagttcgcc agttaatagt ttgcgcaacg ttgttgccat 9240
    tgctacaggc atcgtgggtg caccgctcgtc gtttggtagt gcttcattca gctccggttc 9300
    ccaacgatca aggcgagtta catgatcccc catgttgtgc aaaaaagcgg ttagctcctt 9360
    cggctctccg atcgttgtca gaagtaagtt ggccgcagtg ttatcactca tggttatggc 9420
    agcactgcat aattctctta ctgtcatgcc atccgtaaga tgcttttctg tgactggtga 9480
10   gtactcaacc aagtcattct gagaatagtg tatgcggcga ccgagttgct cttgcccggc 9540
    gtcaatacgg gataataccg cgccacatag cagaacttta aaagtgtca tcattggaaa 9600
    acgttcttcg gggcgaaaac tctcaaggat cttaccgctg ttgagatcca gttcgatgta 9660
    acccactcgt gcacccaact gatcttcagc atcttttact ttcaccagcg tttctgggtg 9720
    agcaaaaaaca ggaaggcaaa atgccgcaaa aaagggaata agggcgacac ggaaatgttg 9780
15   aatactcata ctcttccttt ttcaatatta ttgaagcatt tatcagggtt attgtctcat 9840
    gagcggatag atatttgaat gtatttagaa aaataaacia atagggggtc cgcgcacatt 9900
    tccccgaaaa gtgccacctg acgtctaaga aaccattatt atcatgacat taacctataa 9960
    aaataggcgt atcacgaggc cctttcgtct cgcgcgtttc ggtgatgacg gtgaaaacct 10020
    ctgacacatg cagctcccgg agacggtcac agcttgtctg taagcggatg ccgggagcag 10080
    acaagcccgt cagggcgcggt cagcgggtgt tggcgggtgt cggggctggc ttaactatgc 10140
20   ggcatcagag cagattgtac tgagagtgca ccatatgcgg tgtgaaatac cgcacagatg 10200
    cgtaaggaga aaataccgca tcaggcgcca ttcgccattc aggctgcgca actgttgggg 10260
    agggcgatcg gtgcgggcct cttcgctatt acgccagctg gcgaaagggg gatgtgctgc 10320
    aaggcgatta agttgggtaa cgccagggtt ttcccagtca cgacgttgta aaacgacggc 10380
    cagtgcc

```

10387

```

25
    <210> 9
    <211> 20
    <212> PRT
    <213> Porcine circovirus

```

```

30
    <400> 9

```

```

35   Ser Tyr Pro Arg Arg Arg Tyr Arg Arg Arg Arg His His Pro Pro Ser
      1           5           10           15

```

```

      His Leu Gly Gln
           20

```

```

40
    <210> 10
    <211> 19
    <212> PRT
    <213> Porcine circovirus

```

```

45
    <400> 10

```

```

50   Pro Arg His His Tyr Arg Pro Arg Arg Lys Asn Gly Ile Phe Asn Thr
      1           5           10           15

```

```

      Thr Leu Ser

```

```

55   <210> 11
    <211> 233
    <212> PRT
    <213> Artificial

```

<220>

<223> This is an amino acid sequence for porcine circovirus type 2, open reading frame 2.

<400> 11

```

5
Met Thr Tyr Pro Arg Arg Arg Tyr Arg Arg Arg Arg His Arg Pro Arg
1      5      10      15

10
Ser His Leu Gly Gln Ile Leu Arg Arg Arg Pro Trp Leu Val His Pro
20      25      30

Arg His Arg Tyr Arg Trp Arg Arg Lys Asn Gly Ile Phe Asn Thr Arg
35      40      45

15
Leu Ser Arg Thr Phe Gly Tyr Thr Val Lys Ala Thr Thr Val Arg Thr
50      55      60

Pro Ser Trp Ala Val Asp Met Met Arg Phe Asn Ile Asp Asp Phe Val
65      70      75      80

20
Pro Pro Gly Gly Gly Thr Asn Lys Ile Ser Ile Pro Phe Glu Tyr Tyr
85      90      95

Arg Ile Lys Lys Val Lys Val Glu Phe Trp Pro Cys Ser Pro Ile Thr
100     105     110

25
Gln Gly Asp Arg Gly Val Gly Ser Thr Ala Val Ile Leu Asp Asp Asn
115     120     125

Phe Val Thr Lys Ala Thr Ala Leu Thr Tyr Asp Pro Tyr Val Asn Tyr
130     135     140

30
Ser Ser Arg His Thr Ile Pro Gln Pro Phe Ser Tyr His Ser Arg Tyr
145     150     155     160

Phe Thr Pro Lys Pro Val Leu Asp Ser Thr Ile Asp Tyr Phe Gln Pro
165     170     175

35
Asn Asn Lys Arg Asn Gln Leu Trp Leu Arg Leu Gln Thr Ser Arg Asn
180     185     190

Val Asp His Val Gly Leu Gly Thr Ala Phe Glu Asn Ser Ile Tyr Asp
195     200     205

Gln Asp Tyr Asn Ile Arg Val Thr Met Tyr Val Gln Phe Arg Glu Phe
210     215     220

45
Asn Leu Lys Asp Pro Pro Leu Lys Pro
225     230

```

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 06072065 A [0001] [0004] [0013] [0016] [0017] [0018] [0024] [0028] [0068]
- US 6703023 B [0004]
- WO 03049703 A [0004]
- WO 9918214 A [0004]
- WO 2007028823 A [0004] [0007]
- US 4708871 A [0014]
- WO 0607065 A [0024] [0025]
- US 2909462 A [0038]
- WO 02077210 A [0070]

Non-patent literature cited in the description

- ALLAN et al. *IPVS Congress*, 2006 [0003]
- HARDING et al. *Swine Health Prod*, 1997, vol. 5, 201-203 [0003]
- KENNEDY et al. *J Comp Pathol*, 2000, vol. 122, 9-24 [0003]
- MUIRHEAD. *Vet. Rec.*, 2002, vol. 150, 456 [0003]
- ALLAN ; ELLIS. *J Vet. Diagn. Invest.*, 2000, vol. 12, 3-14 [0003]
- MCKEOWN et al. *Clin. Diagn. Lab. Immunol.*, 2005, vol. 12, 1347-1351 [0005]
- OPRIESSNIG et al. *J. Swine Health Prod.*, 2004, vol. 12, 186-191 [0005]
- OPRIESSNIG et al. *37th Annual Meeting of the American Association of Swine Veterinarians*, 2006 [0005] [0049]
- ALLAN et al. *Vet. Diagn. Investigation*, 2000, vol. 12, 3-14 [0005]
- HIRAI et al. *Vet. Record*, 2001, vol. 148, 482-484 [0005]
- RODRIGUEZ-ARRIOJA et al. *Am. J Vet. Res.*, 2002, vol. 63, 354-357 [0005]
- SURADHAT ; DAMRONGWATANAPOKIN. *Vet. Microbiol*, 2003, vol. 92, 187-194 [0006]
- Epitope Mapping Protocols in *Methods in Molecular Biology*. Humana Press, 1996, vol. 66 [0014]
- GEYSEN et al. *Proc. Natl. Acad. Sci. USA*, 1984, vol. 81, 3998-4002 [0014]
- GEYSEN et al. *Molec. Immunol.*, 1986, vol. 23, 709-715 [0014]
- BERGMANN et al. *Eur. J. Immunol.*, 1993, vol. 23, 2777-2781 [0015]
- BERGMANN et al. *J. Immunol.*, 1996, vol. 157, 3242-3249 [0015]
- SUHRBIER, A. *Immunol. and Cell Biol.*, 1997, vol. 75, 402-408 [0015]
- GARDNER et al. *12th World AIDS Conference*, 1998 [0015]
- Computational Molecular Biology. Oxford University Press, 1988 [0020]
- Biocomputing: Informatics and Genome Projects. Academic Press, 1993 [0020]
- Computer Analysis of Sequence Data. Humana Press, 1994 [0020]
- Sequence Analysis in Molecular Biology. Academic Press, 1987 [0020]
- Sequence Analysis Primer. M. Stockton Press, 1991 [0020]
- CARILLO, H. ; LIPMAN, D. *SIAM J. Applied Math.*, 1988, vol. 48, 1073 [0020]
- DEVEREUX, J. et al. *Nucleic Acids Research*, 1984, vol. 12 (1), 387 [0020]
- ALTSCHUL, S. F. et al. *J. Molec. Biol.*, 1990, vol. 215, 403-410 [0020]
- BLAST Manual. NCBI NLM NIH [0020]
- Remington's Pharmaceutical Sciences. Mack Publ, 1990 [0034]
- HUNTER et al. *The Theory and Practical Application of Adjuvants*. JohnWiley and Sons, 1995, 51-94 [0036]
- TODD et al. *Vaccine*, 1997, vol. 15, 564-570 [0036]
- Vaccine Design, The Subunit and Adjuvant Approach. Plenum Press, 1995, 147 [0037]
- *Pharmeuropa*, June 1996, vol. 8 (2 [0038]
- LAROCHELLE et al. *Can. J. Vet. Res.*, 2003, vol. 67, 114-120 [0048]
- MAGAR et al. *Can. J. Vet Res.*, 2000, vol. 64, 184-186 [0049]
- MAGAR et al. *J. Comp. Pathol.*, 2000, vol. 123, 258-269 [0049]

P A T E N T K R A V

1. Anvendelse af en virksom mængde af et type 2-svinecircovirus(PCV2)-antigen til fremstilling af en immunogen sammensætning til behandling eller forebyggelse af
 - a) PCV2-infektion hos dyr; eller
 - b) til reduktion af kliniske symptomer forvoldt af eller forbundet med en PCV2-infektion hos dyr,
hvor dyrene har anti-PCV2-antistoffer med en anti-PCV2-antistoftiter på mere end 1:1000 som bestemt ved en PCV-specifik indirekte immunfluorescensanalyse,
hvor antigenet er et rekombinant baculovirusudtrykt PCV2-ORF-2-protein, og den immunogene sammensætning indgives kun én gang.
2. Immunogen sammensætning omfattende et type 2-svinecircovirus(PCV2)-antigen til anvendelse i en fremgangsmåde til behandling eller forebyggelse af
 - a) PCV2-infektion hos et dyr; eller
 - b) til reduktion af kliniske symptomer forvoldt af eller forbundet med en PCV2-infektion hos dyr,
hvor dyrene har anti-PCV2-antistoffer med en anti-PCV2-antistoftiter på mere end 1:1000 som bestemt ved en PCV-specifik indirekte immunfluorescensanalyse,
hvor antigenet er et rekombinant baculovirusudtrykt PCV2-ORF-2-protein, og den immunogene sammensætning indgives kun én gang.
3. Anvendelse eller immunogen sammensætning ifølge krav 1 eller 2, hvor anti-PCV2-antistofferne er maternelt afledte antistoffer.
4. Anvendelse eller immunogen sammensætning ifølge et hvilket som helst af kravene 1 til 3, hvor PCV2-antigenet indgives ved alderen 7 dage eller senere.
5. Anvendelse eller immunogen sammensætning ifølge et hvilket som helst af kravene 1 til 4, hvor PCV2-antigenet indgives ved alderen 14 dage eller senere.
6. Anvendelse eller immunogen sammensætning ifølge et hvilket som helst af kravene 1 til 5, hvor PCV2-antigenet indgives senest ved alderen 7 uger.
7. Anvendelse eller immunogen sammensætning ifølge et hvilket som helst af kravene 1 til 6 til forkortelse af viræmifasen på 5 eller flere dage i sammenligning med dyr fra en ikke-behandlet kontrolgruppe af samme art.
8. Anvendelse eller immunogen sammensætning ifølge et hvilket som helst af kravene 1 til 7, hvor PCV2-ORF-2-antigeninklusionsniveauet i den immunogene sammensætning er 15 til 400 µg.
9. Anvendelse eller immunogen sammensætning ifølge et hvilket som helst af kravene 1 til 8, hvor behandlingen eller forebyggelsen af PCV2-infektion resulterer i en forbedring i sammenligning med dyr fra en ikke-behandlet kontrolgruppe af samme art med hensyn til en vaccineeffektivitetsparameter udvalgt fra gruppen bestående af en reduktion af tilvæksttabet, kortere viræmivarighed, tidligere viræmiafslutning, lavere virusbelastning eller kombinationer deraf.
10. Anvendelse eller immunogen sammensætning ifølge et hvilket som helst af kra-

2

vene 1 til 9, hvor dyret er et svin.

5

DRAWINGS

Figure 1:




Figure 2:




Figure 3:

