

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199748748 B2
(10) Patent No. 731929

(54) Title
Heterocycle derivatives which inhibit factor Xa

(51)⁷ International Patent Classification(s)
C07D 239/26 C07D 213/73
A61K 031/44 C07D 213/84
A61K 031/505 C07D 213/89
C07D 213/56 C07D 403/10
C07D 213/61

(21) Application No: **199748748**

(22) Application Date: **1997.11.04**

(87) WIPO No: **WO98/21188**

(30) Priority Data

(31) Number	(32) Date	(33) Country
9623283	1996.11.08	GB
9715893	1997.07.29	GB

(43) Publication Date : **1998.06.03**

(43) Publication Journal Date : **1998.07.30**

(44) Accepted Journal Date : **2001.04.05**

(71) Applicant(s)
AstraZeneca AB

(72) Inventor(s)
John Preston; Andrew Stocker; Paul Turner; Michael James Smithers; John Wall Rayner

(74) Agent/Attorney
PHILLIPS ORMONDE and FITZPATRICK, 367 Collins Street, MELBOURNE VIC 3000



(51) International Patent Classification ⁶ : C07D 239/26, A61K 31/505, C07D 213/56, A61K 31/44, C07D 213/61, 213/84, 213/89, 213/73, 403/10		A1	(11) International Publication Number: WO 98/21188 (43) International Publication Date: 22 May 1998 (22.05.98)
(21) International Application Number: PCT/GB97/03033 (22) International Filing Date: 4 November 1997 (04.11.97) (30) Priority Data: 9623283.0 8 November 1996 (08.11.96) GB 9715893.5 29 July 1997 (29.07.97) GB (71) Applicant (for all designated States except US): ZENECA LIMITED (GB/GB); 15 Stanhope Gate, London W1Y 6LN (GB) AstraZeneca AB, 151 85 Sodertalje, Sweden. (72) Inventors; and (75) Inventors/Applicants (for US only): PRESTON, John [GB/GB]; Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). STOCKER, Andrew [GB/GB]; Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). TURNER, Paul [GB/GB]; Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). SMITHERS, Michael, James [GB/GB]; Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). RAYNER, John, Wall [GB/GB]; Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).		(74) Agent: BROWN, Andrew, Stephen; Zeneca Pharmaceuticals, Intellectual Property Dept., Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: HETEROCYCLE DERIVATIVES WHICH INHIBIT FACTOR Xa			
(57) Abstract The invention relates to heterocyclic derivatives of the formula (I): A-B-X ¹ -T ¹ (R ²)-L ¹ -T ² (R ³)-X ² -Q or pharmaceutically acceptable salts thereof, which possess antithrombotic and anticoagulant properties due to their inhibition of Factor Xa and are accordingly useful in methods of treatment of humans or animals. The invention also relates to processes for the preparation of the heterocyclic derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments for use in the production of an antithrombotic or anticoagulant effect.			



HETEROCYCLIC DERIVATIVES

The invention relates to heterocyclic derivatives, or pharmaceutically-acceptable salts thereof, which possess antithrombotic and anticoagulant properties and are accordingly useful
5 in methods of treatment of humans or animals. The invention also relates to processes for the preparation of the heterocyclic derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments for use in the production of an antithrombotic or anticoagulant effect.

The antithrombotic and anticoagulant effect produced by the compounds of the
10 invention is believed to be attributable to their strong inhibitory effect against the activated coagulation protease known as Factor Xa. Factor Xa is one of a cascade of proteases involved in the complex process of blood coagulation. The protease known as thrombin is the final protease in the cascade and Factor Xa is the preceding protease which cleaves prothrombin to generate thrombin.

15 Certain compounds are known to possess Factor Xa inhibitory properties and the field has been reviewed by R.B. Wallis, Current Opinion in Therapeutic Patents, 1993, 1173-1179. Thus it is known that two proteins, one known as antistatin and the other known as tick anticoagulant protein (TAP), are specific Factor Xa inhibitors which possess antithrombotic properties in various animal models of thrombotic disease.

20 It is also known that certain non-peptidic compounds possess Factor Xa inhibitory properties. Of the low molecular weight inhibitors mentioned in the review by R.B. Wallis, all possessed a strongly basic group such as an amidinophenyl or amidinonaphthyl group.

We have now found that certain heterocyclic derivatives possess Factor Xa inhibitory activity. Many of the compounds of the present invention also possess the advantage of being
25 selective Factor Xa inhibitors, that is the enzyme Factor Xa is inhibited strongly at concentrations of test compound which do not inhibit or which inhibit to a lesser extent the enzyme thrombin which is also a member of the blood coagulation enzymatic cascade.

PCT International Publication Number WO9610022 describes certain heterocyclic piperazine compounds as Factor Xa inhibitors. PCT International Publication Number
30 WO9728129, published after the priority date of this application, describes also certain heterocyclic piperazine compounds as Factor Xa inhibitors. PCT International Publication



AMENDED SHEET

Number WO9730971, published after the priority date of this application, discloses benzamidine type compounds as Factor Xa inhibitors.

The compounds of the present invention possess activity in the treatment or prevention of a variety of medical disorders where anticoagulant therapy is indicated, for example in the treatment or prevention of thrombotic conditions such as coronary artery and


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2

cerebro-vascular disease. Further examples of such medical disorders include various cardiovascular and cerebrovascular conditions such as myocardial infarction, the formation of atherosclerotic plaques, venous or arterial thrombosis, coagulation syndromes, vascular injury including reocclusion and restenosis following angioplasty and coronary artery bypass

5 surgery, thrombus formation after the application of blood vessel operative techniques or after general surgery such as hip replacement surgery, the introduction of artificial heart valves or on the recirculation of blood, cerebral infarction, cerebral thrombosis, stroke, cerebral embolism, pulmonary embolism, ischaemia and angina (including unstable angina).

The compounds of the invention are also useful as inhibitors of blood coagulation in
10 an ex-vivo situation such as, for example, the storage of whole blood or other biological samples suspected to contain Factor Xa and in which coagulation is detrimental.

Accordingly in one aspect the present invention provides compounds of formula (I)



15

wherein:

A is an optionally substituted 5- or 6-membered monocyclic aromatic ring containing 1, 2 or 3 ring heteroatoms selected from nitrogen, oxygen and sulphur atoms;

B is an optionally substituted phenylene ring;

20 T^1 is CH or N;

T^2 is CH or N;

with the proviso that at least one of T^1 and T^2 is N;

X^1 is SO, SO₂, C(R⁴)₂ or CO when T^1 is CH or N; or in addition X^1 is O or S when T^1 is CH;

and wherein each R⁴ is independently hydrogen or C₁₋₄alkyl;

25 L^1 is C₁₋₄alkylene or C₁₋₃alkylenecarbonyl;

R² is hydrogen or C₁₋₄alkyl;

R³ is hydrogen or C₁₋₄alkyl;

or R² and R³ are joined to form a C₁₋₄alkylene or -CH₂CO- group; wherein the ring formed by

T^1 , R², R³, T^2 and L^1 is optionally substituted; with the proviso that when T^1 and T^2 are both N,

30 L^1 is not methylene and R² and R³ together are not methylene;

X^2 is S(O)_y, wherein y is one or two, C(R⁵)₂ or CO; and each R⁵ is hydrogen or C₁₋₄alkyl;

Q is phenyl, naphthyl, phenylC₁₋₄alkyl, phenylC₂₋₄alkenyl, phenylC₂₋₄alkynyl or a heterocyclic moiety containing up to 4 heteroatoms selected from nitrogen, oxygen and sulphur and Q is optionally substituted;

and pharmaceutically acceptable salts thereof.

5 In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. An analogous convention applies to other generic terms.

It is to be understood that certain heterocyclic derivatives of the present invention can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to
10 be understood that the invention encompasses all such solvated forms which possess Factor Xa inhibitory activity.

It is further to be understood that, insofar as certain of the compounds of the formula defined above may exist in optically active or racemic forms by virtue of one or more asymmetric carbon atoms, the invention encompasses any such optically active or racemic
15 form which possesses Factor Xa inhibitory activity. The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form.

A is an optionally substituted 5- or 6-membered monocyclic aromatic ring
20 containing 1, 2 or 3 ring nitrogen atoms. Preferably A is a pyridyl, pyrimidinyl or pyridazinyl ring for example 4-pyridyl, 2-pyridyl, 4-pyridazinyl, 5-pyrimidinyl, 4-pyrimidinyl or 3-pyridyl. Of these 4-pyrimidinyl, 4-pyridazinyl and 4-pyridyl are preferred of which 4-pyrimidinyl and 4-pyridyl are most preferred.

In one aspect A is unsubstituted. In another aspect A is substituted by one, two or
25 three atoms or groups selected from halo (for example fluoro, chloro or bromo), oxo, carboxy, trifluoromethyl, cyano, amino, hydroxy, nitro, C₁₋₄alkyl (for example methyl or ethyl), C₁₋₄alkoxy (for example methoxy or ethoxy), C₁₋₄alkoxycarbonyl, C₁₋₄alkylamino (for example methylamino or ethylamino) or di-C₁₋₄alkylamino (for example dimethylamino or diethylamino). For the avoidance of doubt substituents on A may also be present, where
30 possible, on the heteroatom of the ring, such as, for example, N-oxides. Preferred substituents are C₁₋₄alkyl and halo. Preferably A is unsubstituted

B is an optionally substituted phenylene ring wherein the bonds to A and X¹ may suitably be in the meta or para disposition. Preferably the bonds to A and X¹ are in para disposition, that is B is a para-phenylene group.

In one aspect B is unsubstituted. In another aspect B is substituted by one or two
 5 substituents selected from halo, trifluoromethyl, trifluoromethoxy, cyano, nitro, C₁₋₄alkyl, C₂₋₄alkenyl and C₂₋₄alkynyl, from the substituent -(CH₂)_n Y¹ wherein n is 0-4 and Y¹ is selected from hydroxy, amino, carboxy, C₁₋₄alkoxy, C₂₋₄alkenyloxy, C₂₋₄alkynyloxy, C₁₋₄alkylamino, di-C₁₋₄alkylamino, pyrrolidin-1-yl, piperidino, morpholino, thiomorpholino, 1-oxothiomorpholino, 1,1-dioxothiomorpholino, piperazin-1-yl, 4-C₁₋₄alkylpiperazin-1-yl, C₁₋₄alkylthio, C₁₋₄alkylsulphinyl, C₁₋₄alkylsulphonyl, C₂₋₄alkanoylamino, benzamido, C₁₋₄alkylsulphonamido and phenylsulphonamido, from the substituent -(CH₂)_n Y² wherein n is 0-4 and Y² is selected from carboxy, carbamoyl, C₁₋₄alkoxycarbonyl, N-C₁₋₄alkylcarbamoyl, N,N-di-C₁₋₄alkylcarbamoyl, pyrrolidin-1-ylcarbonyl, piperidinocarbonyl, morpholinocarbonyl, thiomorpholinocarbonyl, 1-oxothiomorpholinocarbonyl, 1,1-dioxothiomorpholinocarbonyl, piperazin-1-ylcarbonyl, 4-C₁₋₄alkylpiperazin-1-ylcarbonyl, C₁₋₄alkylsulphonamidocarbonyl, phenylsulphonamidocarbonyl and benzylsulphonamidocarbonyl, from a substituent of the formula -X³-L²-Y² wherein X³ is a group of the formula CON(R⁵), CON(L²-Y²), C(R⁵)₂O, O, N(R⁵) or N(L²-Y²), L² is C₁₋₄alkylene, Y² has any of the meanings
 20 defined immediately hereinbefore and each R⁵ is independently hydrogen or C₁₋₄alkyl, and from a substituent of the formula -X³-L³-Y¹ wherein X³ is a group of the formula CON(R⁵), CON(L³-Y¹), C(R⁵)₂O, O, N(R⁵) or N(L³-Y¹), L³ is C₂₋₄alkylene, Y¹ has any of the meanings defined immediately hereinbefore and each R⁵ is independently hydrogen or C₁₋₄alkyl, and wherein any heterocyclic group in a substituent of B optionally bears 1 or 2 substituents
 25 selected from carboxy, carbamoyl, C₁₋₄alkyl, C₁₋₄alkoxycarbonyl, N-C₁₋₄alkylcarbamoyl and N,N-di-C₁₋₄alkylcarbamoyl, and wherein any phenyl group in a substituent of B optionally bears 1 or 2 substituents selected from halo, trifluoromethyl, cyano, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy, C₂₋₄alkenyloxy and C₂₋₄alkynyloxy. Preferably B is substituted by carboxy, C₁₋₄alkoxy or C₁₋₄alkoxycarbonyl. Preferably B is unsubstituted.

30 In a particular aspect, when T¹ is CH or N, X¹ is CO, SO₂, or CH₂ or, when T¹ is CH, X¹ in addition is O or S. Preferably X¹ is CO.

For the avoidance of doubt T^1 is directly attached to the groups X^1 and L^1 and T^2 is directly attached to the groups L^1 and X^2 .

L^1 is C_{1-4} alkylene for example methylene, ethylene or propylene (preferably ethylene) or is C_{1-3} alkylenecarbonyl for example methylenecarbonyl ($-\text{CH}_2\text{CO}-$).

5 R^2 is hydrogen or C_{1-4} alkyl for example methyl or ethyl. R^3 is hydrogen or C_{1-4} alkyl for example methyl or ethyl.

In a preferred aspect R^2 and R^3 are joined to form a C_{1-4} alkylene group, for example a methylene, ethylene or propylene group (preferably ethylene), or a methylenecarbonyl ($-\text{CH}_2\text{CO}-$) group.

10 In a particular aspect R^2 and R^3 are joined to form, together with T^1 , T^2 and L^1 , a heterocyclic ring wherein at least one of T^1 and T^2 is N. Examples of such heterocyclic rings are piperazine (wherein T^1 and T^2 are both N), piperidine (wherein either T^1 or T^2 is N and the other is CH) and pyrrolidine (wherein either T^1 or T^2 is N and other is CH). Preferably the heterocyclic ring formed by R and R^1 is piperazine.

15 In one aspect the heterocyclic ring formed by T^1 , T^2 , L^1 , R^2 and R^3 is unsubstituted. In another aspect this ring is substituted by one or two substituents selected from hydroxy, oxo, carboxy and C_{1-4} alkoxycarbonyl; or one of the following:

$-(\text{CH}_2)_n-\text{R}$, $-(\text{CH}_2)_n-\text{NRR}^1$, $-\text{CO}-\text{R}$, $-\text{CO}-\text{NRR}^1$, $-(\text{CH}_2)_n-\text{CO}-\text{R}$ and $-(\text{CH}_2)_n-\text{CO}-\text{NRR}^1$;

20

wherein n is 0, 1 or 2, preferably n is 1 or 2;

R and R^1 are independently selected from hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, hydroxy C_{1-4} alkyl, carboxy C_{1-4} alkyl and C_{1-4} alkoxycarbonyl C_{1-4} alkyl or where possible R and R^1 may together form a 5- or 6-membered optionally substituted saturated or partially

25 unsaturated (preferably saturated) heterocyclic ring which may include in addition to the nitrogen to which R and R^1 are attached 1 or 2 additional heteroatoms selected from nitrogen, oxygen and sulphur.

In a particular aspect the heterocyclic ring formed by R and R^1 are preferably selected from 1-pyrrolidinyl, 1-imidazolyl, 1-piperidino, 1-piperazinyl, 4-morpholino and
30 4-thiomorpholino. In a particular aspect the heterocyclic ring formed by R and R^1 may be unsubstituted. In an alternative aspect the ring formed by R and R^1 is substituted by 1 or 2

substituents selected from oxo, hydroxy and carboxy. Preferably the heterocyclic ring formed by T¹, T², L¹, R² and R³ is substituted by oxo, carboxy, C₁₋₄alkoxy or C₁₋₄alkoxycarbonyl. Preferably the heterocyclic ring formed by T¹, T², L¹, R² and R³ is unsubstituted.

In a particular aspect X² is SO₂, CH₂ or CO. Preferably X² is SO₂.

- 5 In one aspect Q is unsubstituted. In another aspect Q is substituted by one, two or three substituents selected from halo, trifluoromethyl, trifluoromethoxy, cyano, hydroxy, amino, nitro, trifluoromethylsulphonyl, carboxy, carbamoyl, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy, C₂₋₄alkenyloxy, C₂₋₄alkynyloxy, C₁₋₄alkylthio, C₁₋₄alkylsulphinyl, C₁₋₄alkylsulphonyl, C₁₋₄alkylamino, di-C₁₋₄alkylamino, C₁₋₄alkoxycarbonyl, N-C₁₋₄alkylcarbamoyl, N,N-di-C₁₋₄alkylcarbamoyl, C₂₋₄alkanoyl, C₂₋₄alkanoylamino, hydroxyC₁₋₄alkyl, C₁₋₄alkoxyC₁₋₄alkyl, carboxyC₁₋₄alkyl, C₁₋₄alkoxycarbonylC₁₋₄alkyl, carbamoylC₁₋₄alkyl, N-C₁₋₄alkylcarbamoylC₁₋₄alkyl, N,N-di-C₁₋₄alkylcarbamoylC₁₋₄alkyl, phenyl, heteroaryl, phenoxy, phenylthio, phenylsulphinyl, phenylsulphonyl, benzyl, benzoyl, heteroaryloxy, heteroarylthio, heteroarylsulphinyl and heteroarylsulphonyl, and wherein said
- 10 heteroaryl substituent or the heteroaryl group in a heteroaryl-containing substituent is a 5- or 6-membered monocyclic heteroaryl ring containing up to 3 heteroatoms selected from nitrogen, oxygen and sulphur, and wherein said phenyl, heteroaryl, phenoxy, phenylthio, phenylsulphinyl, phenylsulphonyl, heteroaryloxy, heteroarylthio, heteroarylsulphinyl, heteroarylsulphonyl, benzyl or benzoyl substituent optionally bears 1, 2 or 3 substituents
- 20 selected from halo, trifluoromethyl, cyano, hydroxy, amino, nitro, carboxy, carbamoyl, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkylamino, di-C₁₋₄alkylamino, C₁₋₄alkoxycarbonyl, N-C₁₋₄alkylcarbamoyl, N,N-di-C₁₋₄alkylcarbamoyl and C₂₋₄alkanoylamino.

A suitable value for Q when it is naphthyl is, for example, 1-naphthyl or 2-naphthyl; when it is phenyl-C₁₋₄alkyl is, for example, benzyl, phenethyl and 3-phenylpropyl, when it is

25 phenyl-C₂₋₄alkenyl is, for example, styryl, cinnamyl or 3-phenylprop-2-enyl; and when it is phenyl-C₂₋₄alkynyl is, for example, 2-phenylethynyl, 3-phenylprop-2-ynyl and 3-phenylprop-1-ynyl. Preferably Q is naphthyl in particular 2-naphthyl.

A suitable value for Q when it is a heterocyclic moiety containing up to 4 heteroatoms selected from nitrogen, oxygen and sulphur is, for example, a 5- or 6-membered

30 heterocyclic moiety which is a single ring or is fused to one or two benzo rings such as furyl, benzofuranyl, tetrahydrofuryl, chromanyl, thienyl, benzothienyl, pyridyl, piperidino, quinolyl,

- 1,2,3,4-tetrahydroquinoliny, isoquinolyl, 1,2,3,4-tetrahydroisoquinoliny, pyrazinyl, piperazinyl, pyrimidinyl, pyridazinyl, quinoxaliny, quinazoliny, cinnoliny, pyrrolyl, pyrrolidinyl, indolyl, indoliny, imidazolyl, benzimidazolyl, pyrazolyl, indazolyl, oxazolyl, benzoxazolyl, isoxazolyl, thiazolyl, benzothiazolyl, isothiazolyl, morpholino, 4H-1,4-benzoxazinyl, 4H-1,4-benzothiazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, oxadiazolyl, furazanyl, thiadiazolyl, tetrazolyl, dibenzofuranyl and dibenzothienyl, which may be attached through any available position including, for an appropriate X² group such as, for example, SO₂, C(R⁵)₂ or CO, through any available nitrogen atom, and which may bear up to three substituents as defined hereinabove including a substituent on any available nitrogen atom.
- 10 A suitable value for the heteroaryl substituent on Q or the heteroaryl group in a heteroaryl-containing substituent on Q which comprises a 5- or 6-membered monocyclic heteroaryl ring containing up to 3 heteroatoms selected from oxygen, nitrogen and sulphur is, for example, furyl, thienyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 15 oxadiazolyl, furazanyl and thiadiazolyl which may be attached through any available position including, for an appropriate X² group such as, for example, SO₂, C(R⁵)₂ or CO, through any available nitrogen atom, and which may be up to three substituents as defined hereinabove including a substituent on any available nitrogen atom.

Suitable values for optional substituents for B and Q are:

- | | |
|--|---|
| 20 for C ₁₋₄ alkyl: | methyl, ethyl and propyl; |
| for C ₁₋₄ alkoxycarbonyl: | methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl and <u>tert</u> -butoxycarbonyl; |
| for <u>N</u> -C ₁₋₄ alkylcarbamoyl: | <u>N</u> -methylcarbamoyl, <u>N</u> -ethylcarbamoyl and <u>N</u> -propylcarbamoyl; |
| 25 for <u>N,N</u> -di-C ₁₋₄ alkylcarbamoyl: | <u>N,N</u> -dimethylcarbamoyl, <u>N</u> -ethyl- <u>N</u> -methylcarbamoyl and <u>N,N</u> -diethylcarbamoyl; |
| for hydroxyC ₁₋₄ alkyl: | hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl and 3-hydroxypropyl; |
| 30 for C ₁₋₄ alkoxyC ₁₋₄ alkyl: | methoxymethyl, ethoxymethyl, 1-methoxymethyl, 2-methoxyethyl, |

for carboxyC₁₋₄alkyl:

for C₁₋₄alkoxycarbonylC₁₋₄alkyl:

5

10

for carbamoylC₁₋₄alkyl:

15 for N-C₁₋₄alkylcarbamoylC₁₋₄alkyl:

20

for N,N-di-C₁₋₄alkylcarbamoyl-
C₁₋₄alkyl:

25

30

2-ethoxyethyl and 3-methoxypropyl;
carboxymethyl, 1-carboxyethyl,
2-carboxyethyl and 3-carboxypropyl;
methoxycarbonylmethyl,
ethoxycarbonylmethyl, tert-butoxy-
carbonylmethyl, 1-methoxycarbonylethyl,
1-ethoxycarbonylethyl,
2-methoxycarbonylethyl,
2-ethoxycarbonylethyl,
3-methoxycarbonylpropyl and
3-ethoxycarbonylpropyl;
carbamoylmethyl, 1-carbamoylethyl,
2-carbamoylethyl and
3-carbamoylpropyl;
N-methylcarbamoylmethyl,
N-ethylcarbamoylmethyl,
N-propylcarbamoylmethyl,
1-(N-methylcarbamoyl)ethyl,
1-(N-ethylcarbamoyl)ethyl,
2-(N-methylcarbamoyl)ethyl,
2-(N-ethylcarbamoyl)ethyl and
3-(N-methylcarbamoyl)propyl;
N,N-dimethylcarbamoylmethyl,
N-ethyl-N-methylcarbamoylmethyl,
N,N-diethylcarbamoylmethyl,
1-(N,N-dimethylcarbamoyl)ethyl,
1-(N,N-diethylcarbamoyl)ethyl,
2-(N,N-dimethylcarbamoyl)ethyl,
2-(N,N-diethylcarbamoyl)ethyl and
3-(N,N-dimethylcarbamoyl)propyl;

- for halo: fluoro, chloro, bromo;
- for C₁₋₄alkoxy: methoxy, ethoxy;
- for C₁₋₄alkylamino: methylamino, ethylamino;
- for di-C₁₋₄alkylamino: dimethylamino, diethylamino;
- 5 for C₁₋₄alkenyl: vinyl and allyl;
- for C₂₋₄alkynyl: ethynyl and prop-2-ynyl;
- for C₂₋₄alkenyloxy: vinyloxy and allyloxy;
- for C₂₋₄alkynyloxy: ethynyloxy and prop-2-ynyloxy;
- for C₁₋₄alkylthio: methylthio, ethylthio and propylthio;
- 10 for C₁₋₄alkylsulphinyl: methylsulphinyl, ethylsulphinyl and propylsulphinyl;
- for C₁₋₄alkylsulphonyl: methylsulphonyl, ethylsulphonyl and propylsulphonyl;
- for C₂₋₄alkanoylamino: acetamido, propionamido and butyramido;
- 15 A preferred class of compounds of the present invention is that wherein:
- A is pyridyl, pyrimidinyl or pyridazinyl;
- B is para-phenylene;
- X¹ is CO, SO₂ or CH₂, preferably CO;
- T¹ and T² are both N;
- 20 L¹ is ethylene or propylene;
- R² and R³ are joined to form an ethylene or propylene or methylenecarbonyl group;
- wherein the heterocyclic ring formed by T¹, T², L¹, R² and R³ is unsubstituted or is substituted;
- X² is SO₂;
- Q is styryl optionally substituted (preferably 4-substituted), naphthyl optionally substituted
- 25 (preferably 6-substituted) or is phenyl optionally substituted (preferably 4-substituted) by fluoro, chloro or bromo;
- and pharmaceutically-acceptable salts thereof.

Particular compounds of the invention include:

- 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyridyl)benzoyl]piperazine;
- 30 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(2-pyridyl)benzoyl]piperazine;
- 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyrimidinyl)benzoyl]piperazine;

- 1-(6-chloronaphth-2-ylsulphonyl)-4-[4-(4-pyridinyl)benzoyl]piperazine;
 1-(6-chloronaphth-2-ylsulphonyl)-3-methoxycarbonyl-4-[4-(4-pyrimidinyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(2-methylpyrimidin-4-yl)benzoyl]piperazine;
 5 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(2,6-dimethylpyrimidin-4-yl)benzoyl]piperazine;
 1-(6-chloronaphth-2-ylsulphonyl)-4-[4-(4-pyrimidinyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(3-fluoro-4-pyridyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-3-hydroxymethyl-4-[4-(4-pyridyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-3-ethoxycarbonyl-4-[4-(4-pyridyl)benzoyl]piperazine;
 10 1-(6-bromonaphth-2-ylsulphonyl)-4-[2-methoxycarbonyl-4-(4-pyridyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyridazinyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(2-methyl-4-pyridyl)benzoyl]piperazine;
 1-(4-chloro-E-styrylsulphonyl)-4-[4-(4-pyridyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(3-pyridazinyl)benzoyl]piperazine;
 15 1-(6-bromonaphth-2-ylsulphonyl)-3-oxo-4-[4-(4-pyrimidinyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(2-cyano-4-pyridyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[2-methoxy-4-(4-pyridazinyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(1,2,3-thiadiazol-4-yl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyridyl)benzoyl]homopiperazine;
 20 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(3,5-diamino-1,2,4-triazol-1-yl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-3-(4-thiomorpholinocarbonyl)-4-[4-(4-pyrimidinyl)benzoyl]piperazine;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(3-furanyl)benzoyl]piperazine; and
 1-(6-methoxynaphth-2-ylsulphonyl)-4-[4-(4-pyridyl)benzoyl]piperazine.
- 25 Particularly preferred compounds of the invention are;
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyrimidinyl)benzoyl]piperazine;
 1-(6-chloronaphthyl-2-ylsulphonyl)-4-[4-(4-pyridyl)benzoyl]piperazine; and
 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyridazinyl)benzoyl]piperazine.


A heterocyclic derivative of the formula I, or pharmaceutically-acceptable salt
 30 thereof, may be prepared by any process known to be applicable to the preparation of related compounds. Such procedures are provided as a further feature of the invention and are

illustrated by the following representative processes in which, unless otherwise stated A, B, X¹, T¹, T², L¹, R², R³, X² and Q have any of the meanings defined hereinbefore wherein any functional group, for example amino, alkylamino, carboxy or hydroxy, is optionally protected by a protecting group which may be removed when necessary.

5 Necessary starting materials may be obtained by standard procedures of organic chemistry.

According to another aspect, the present invention provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt thereof, which comprises:

10 (a) For the production of those compounds of the formula (I) wherein T¹ is N and X¹ is CO, the reaction, conveniently in the presence of a suitable base, of an amine of formula (II)



15 with an acid of the formula (III)



20 or a reactive derivative thereof.

A suitable reactive derivative of an acid of the formula (III) is, for example, an acyl halide, for example an acyl chloride formed by the reaction of the acid and an inorganic acid chloride, for example thionyl chloride; a mixed anhydride, for example an anhydride formed by the reaction of the acid with a chloroformate such as isobutyl chloroformate or with an
25 activated amide such as 1,1'-carbonyldiimidazole; an active ester, for example an ester formed by the reaction of the acid and a phenol such as pentafluorophenol, an ester such as pentafluorophenyl trifluoroacetate or an alcohol such as N-hydroxybenzotriazole or N-hydroxysuccinimide; an acyl azide, for example an azide formed by the reaction of the acid and an azide such as diphenylphosphoryl azide; an acyl cyanide, for example a cyanide
30 formed by the reaction of an acid and a cyanide such as diethylphosphoryl cyanide; or the

product of the reaction of the acid and a carbodiimide such as

N,N'-dicyclohexylcarbodiimide or N-(3-dimethylaminopropyl)-N'-ethyl-carbodiimide.

The reaction is conveniently carried out in the presence of a suitable base such as, for example, an alkali or alkaline earth metal carbonate, alkoxide, hydroxide or hydride, for example sodium carbonate, potassium carbonate, sodium ethoxide, potassium butoxide, sodium hydroxide, potassium hydroxide, sodium hydride or potassium hydride, or an organometallic base such as an alkyl-lithium, for example n-butyl-lithium, or a dialkylamino-lithium, for example lithium di-isopropylamide, or, for example, an organic amine base such as, for example, pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, triethylamine, morpholine or diazabicyclo[5.4.0]undec-7-ene. The reaction is also preferably carried out in a suitable inert solvent or diluent, for example methylene chloride, chloroform, carbon tetrachloride, tetrahydrofuran, 1,2-dimethoxyethane, N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidin-2-one, dimethylsulphoxide or acetone, and at a temperature in the range, for example, -78° to 150°C, conveniently at or near ambient temperature.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or tert-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a tert-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid such as hydrochloric, sulphuric, phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.


A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl
 5 group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. An arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying
 10 group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a tert-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

15

(b) For the production of those compounds of the formula (I) wherein T^1 is CH and X^1 is O by the reaction, conveniently in the presence of a suitable coupling agent, of a compound of the formula (IV):

20



wherein Z is a displaceable group, with a phenolic compound of the formula (V):

25



A suitable value for the displaceable group Z is, for example, a halo or sulphonyloxy group, for example a fluoro, chloro, bromo, mesyloxy or 4-tolylsulphonyloxy group.

A suitable reagent for the coupling reaction when Z is a halo or sulphonyloxy
 30 group is, for example, a suitable base, for example, an alkali or alkaline earth metal carbonate, hydroxide or hydride, for example sodium carbonate, hydroxide or hydride, for

example sodium carbonate, potassium carbonate, sodium hydroxide, potassium hydroxide, sodium hydride or potassium hydride. The alkylation reaction is preferably performed in a suitable inert solvent or diluent, for example N,N-dimethylformamide, N,N-dimethylacetamide, dimethylsulphoxide, acetone, 1,2-dimethoxyethane or tetrahydrofuran, and at a temperature in the range, for example, -10° to 150°C, conveniently at or near ambient temperature. An analogous procedure may be employed for the preparation of those compounds of the formula (I) wherein T¹ is CH and X¹ is S.

A suitable reagent for the coupling reaction of the alcohol of the formula (IV) wherein Z is a hydroxy group, where the hydroxy group is converted in situ to a displaceable group as defined above, is, for example, the reagent obtained when said alcohol is reacted with a di-C₁₋₄alkyl azodicarboxylate in the presence of a triarylphosphine or tri-C₁₋₄alkylphosphine, for example with diethyl azodicarboxylate in the presence of triphenylphosphine or tributylphosphine. The reaction is preferably performed in a suitable inert solvent or diluent, for example acetone, 1,2-dimethoxyethane or tetrahydrofuran, and at a temperature in the range, for example, 10° to 80°C, conveniently at or near ambient temperature.

(c) For the production of those compounds of the formula (I) wherein T¹ is N and X¹ is CH(R⁴), the reductive amination of a keto compound of the formula (VI):




wherein R⁴ is hydrogen or C₁₋₄ alkyl, with an amine of the formula (II) as defined above.

Any reducing agent known in the art for promoting a reductive amination reaction may be employed. A suitable reducing agent is, for example, a hydride reducing agent, for example an alkali metal aluminium hydride such as lithium aluminium hydride or, preferably, an alkali metal borohydride such as sodium borohydride, sodium cyanoborohydride, sodium triethylborohydride, sodium trimethoxyborohydride and sodium triacetoxyborohydride. The reaction is conveniently performed in a suitable inert solvent or diluent, for example tetrahydrofuran and diethyl ether for the more powerful reducing agents such as lithium aluminium hydride, and, for example, methylene chloride or a protic solvent

such as methanol and ethanol for the less powerful reducing agents such as sodium triacetoxyborohydride. The reaction is performed at a temperature in the range, for example, 10° to 80°C, conveniently at or near ambient temperature.

- 5 (d) The reaction of a compound of the formula (VII):



wherein Z is a displaceable group such as halo, with an activated derivative of ring A.

- 10 Suitable activated derivatives include metalised derivatives, such as with zinc or tin, and borane derivatives. The activated derivative of ring A is reacted with a compound of the formula (VII) to effect cross coupling where Z is triflate or a halo group, such as iodo, bromo or chloro. Suitably the reaction is catalysed by use of a transition state metal catalyst, such as palladium, for example tetrakis (triphenylphosphine) palladium (0).

- 15 Alternatively it is possible that ring A contains the displaceable group Z and ring B is activated, and the reaction performed as described above.

Compounds of the formula (VII) not suitable for this method are those which contain a halo substituent on B, Q or L¹.

- 20 (e) By forming A ring on compounds of formula (VII), wherein Z is a functional group capable of cyclisation. Suitable reagents and conditions are described below in preparing compounds of formula (III) by cyclisation.

- (f) For the production of compounds wherein T² is N, the reaction of a compound of the
25 formula (VIII):




with a compound of the formula (IX):

30



wherein Z is a displaceable group for example chloro, under conditions similar to those of process (a) above.

- 5 (g) For the production of compounds wherein T^1 is N and X^1 is SO or SO_2 , the reaction of a compound of the formula (II) as defined above with a compound of the formula (X):



- 10 wherein x is one or two and Z is a displaceable group; under appropriate conventional coupling conditions, similar to those of process variant (a) above.

(h) For production of compounds of the formula (I) by coupling T^2 to Q and thus preparing the $-T^2-X^2-Q$ moiety, methods analogous to those described in process variants (a),

- 15 (c) and (g) for preparing the $B-X^1-T^1$ moiety may be employed.

- (i) For the production of compounds of the formula (I) wherein X^1 is a group of the formula SO, SO_2 , wherein B bears a C_{1-4} alkylsulphinyl, C_{1-4} alkylsulphonyl, 1-oxothiomorpholino or 1,1-dioxothiomorpholino group, wherein X^2 is a group of the formula
20 SO or SO_2 , wherein Q bears a C_{1-4} alkylsulphinyl, C_{1-4} alkylsulphonyl, phenylsulphinyl, phenylsulphonyl, heteroarylsulphinyl or heteroarylsulphonyl group, the oxidation of the corresponding compound of the formula (I) which contains X^1 as a thio group.

A suitable oxidising agent is, for example, any agent known in the art for the oxidation of thio to sulphinyl and/or sulphonyl, for example, hydrogen peroxide, a peracid
25 (such as 3-chloroperoxybenzoic or peroxyacetic acid), an alkali metal peroxysulphate (such as potassium peroxymonosulphate), chromium trioxide or gaseous oxygen in the presence of platinum. The oxidation is generally carried out under as mild conditions as possible and with the required stoichiometric amount of oxidising agent in order to reduce the risk of over oxidation and damage to other functional groups. In general the reaction is carried out in a
30 suitable solvent or diluent such as methylene chloride, chloroform, acetone, tetrahydrofuran or tert-butyl methyl ether and at a temperature, for example, at or near ambient temperature,

that is in the range 15 to 35°C. Suitable reagents and conditions are described in, for example, Page G. O.; Synth. Commun. 23, (1993) 6, 765-769. When a compound carrying a sulphinyl group is required a milder oxidising agent may also be used, for example sodium or potassium metaperiodate, conveniently in a polar solvent such as acetic acid or ethanol. It will be appreciated that when a compound of the formula I containing a sulphonyl group is required, it may be obtained by oxidation of the corresponding sulphinyl compound as well as of the corresponding thio compound. Those compounds of the formula (I) which contain oxygen labile groups (such as A ring is pyridyl) are probably not suitable intermediates for this process step, unless oxidation of such groups is desired.

- 10 Compounds of formula (II) wherein T² is N may be prepared by the reaction of a compound of the formula (XI)



- 15 , wherein P is a protecting group, with a compound of formula (IX), as defined above, in an analogous manner as described above in method (f) above, and subsequently removing the protecting group. In addition compounds of formula (II) may be prepared in an analogous manner as described above in methods (h) and (i).

- 20 Compounds of formula (IV) may be prepared in an analogous manner as described for the preparation of compounds of formula (II).

 Compounds of formula (III) may be prepared by the coupling of a compound of formula (XII), wherein Z is a displaceable group, preferably halo,



- 25 with an activated derivative of ring A as described, for example, in method (d) above. Ideally the reaction is catalysed with a palladium catalyst as described in Example 1 (c) and Example 3 (a) below. Suitable reagents and conditions are described in Martin A.R.; Acta.Chem.Scand., 47, 221-230, (1993); Mitchell T.N.; Synthesis, 803, (1992) and Stille, 30 J.K., Angew. Chem. Int. Ed. Engl. 25, 508-524, (1986).

Suitable non-catalysed coupling reactions include those described in Shiao, M-J. et. al., Synlett., 655, (1992).

Synthesis of stannane intermediates which may be required for palladium catalysed reactions are described in Hylarides, M.D. et. al., Journal of Organometallic Chemistry, 367, 5 259-265, (1989).

Alternatively compounds of formula (III) may be prepared by forming A rings on compounds of formula (XII), wherein Z is a functional group capable of cyclisation, by cyclisation reaction. Suitable reagents and conditions are described in Bredereck H. Chem.Ber.; 96, 1505, (1963); Fuchigami, T., Bull. Chem. Soc. Jpn., 49, p3607, (1976); 10 Huffman, K.R., J. Org. Chem., 28, p1812, (1963); Palusso, G., Gazz. Chim. Ital., 90, p1290, (1960) and Ainsworth C.J., Heterocycl. Chem., 3, p470, (1966). Such reactions are particularly suited to the formation of 5-membered A rings. Processes suitable for synthesis of starting materials in such cyclisation reactions are described, for example, in Zhang M.Q. et.al; J.Heterocyclic. Chem.; 28, 673, (1991) and Kosugi, M. et al., Bull. Chem. Soc. Jpn., 15 60, 767-768 (1987).

Compounds of formula (V), (VI) and (X) may be prepared in an analogous manner as described for preparing compounds of formula (III).

Compounds of formula (VII) wherein T² is N may be prepared by the reaction of a compound of the formula (XIII)


20



with a compound of formula (IX), as defined above, in an analogous manner as described above in method (f).


25

Compounds of formula (XIII) may be prepared by the reaction of a compound of the formula (XIV)



wherein Z is a displaceable group or hydrogen and P is a protecting group in an analogous manner is described in method (a), (b) and (c) above and subsequently effecting removal of the protecting group.

Compounds of formula (VIII) may be prepared by the reaction of a compound of
5 formula (XV)



wherein Z is a displaceable group or hydrogen and P is a protecting group, in an analogous
10 method as described in method (a), (b) and (c) above and subsequently effecting removal of the protecting group.

Compounds of formula (IX), where X^2 is SO or SO₂, may be prepared by oxidation of compound of formula (IX), where X^2 is S, in an analogous method as described in method (h) above. Suitable reagents and conditions are described in Newman, M.S., et. al., Organic
15 Synthesis. Vol. 51, p139. Methods for preparation of the thio analogues of Q are described in Kharasch, N. et. al., J. Am. Chem. Soc., 73, p3240, 1951.

When a pharmaceutically-acceptable salt of a compound of the formula (I) is required, it may be obtained, for example, by reaction of said compound with a suitable acid or base using a conventional procedure.

20 When an optically active form of a compound of the formula (I) is required, it may be obtained, for example, by carrying out one of the aforesaid procedures using an optically active starting material or by resolution of a racemic form of said compound using a conventional procedure, for example by the formation of diastereomeric salts, use of chromatographic techniques, conversion using chirally specific enzymatic processes, or by
25 addition of temporary extra chiral group to aid separation.

As stated previously, the compounds of the formula (I) are inhibitors of the enzyme Factor Xa. The effects of this inhibition may be demonstrated using one or more of the standard procedures set out hereinafter:-

30 a) Measurement of Factor Xa Inhibition

An *in vitro* assay system based on the method of Kettner *et al.*, *J. Biol. Chem.*, 1990, 265, 18289-18297, whereby various concentrations of a test compound are dissolved in a pH7.5 buffer containing 0.5% of a polyethylene glycol (PEG 6000) and incubated at 37°C with human Factor Xa (0.001 Units/ml, 0.3 ml) for 15 minutes. The chromogenic substrate
5 S-2765 (KabiVitrum AB, 20 µM) is added and the mixture is incubated at 37°C for 20 minutes whilst the absorbance at 405 nm is measured. The maximum reaction velocity (V_{max}) is determined and compared with that of a control sample containing no test compound. Inhibitor potency is expressed as an IC₅₀ value.

b) Measurement of Thrombin Inhibition

10 The procedure of method a) is repeated except that human thrombin (0.005 Units/ml) and the chromogenic substrate S-2238 (KabiVitrum AB, 7 µM) are employed.

c) Measurement of Anticoagulant Activity

An *in vitro* assay whereby human, rat or rabbit venous blood is collected and added directly to a sodium citrate solution (3.2 g/100 ml, 9 parts blood to 1 part citrate solution). Blood
15 plasma is prepared by centrifugation (1000 g, 15 minutes) and stored at 2-4°C. Conventional prothrombin time (PT) tests are carried out in the presence of various concentrations of a test compound and the concentration of test compound required to double the clotting time, hereinafter referred to as CT₂, is determined. In the PT test, the test compound and blood plasma are incubated at 37°C for 10 minutes. Tissue thromboplastin with calcium (Sigma
20 Limited, Poole, England) is added and fibrin formation and the time required for a clot to form are determined.

d) Rat Disseminated Intravascular Coagulation in vivo activity test:

Fasted male Alderley Park rats (300-450 g) are pre-dosed by oral gavage (5 mls/kg) with
25 compound or vehicle (5% DMSO/PEG200) at various times before being anaesthetised with Intraval® (120 mg/kg i.p.). The left jugular vein and the right carotid artery are exposed and cannulated. A 1 mL blood sample is taken from the carotid canular into 3.2% trisodium citrate. 0.5 mL of the whole blood is then treated with EDTA and used for platelet count determination whilst the remainder is centrifuged (5 mins, 20000g) and the resultant plasma
30 frozen for subsequent drug level, fibrinogen or thrombin antithrombin (TAT) complex determinations. Recombinant human tissue factor (Dade Innovin Cat.B4212-50), reconstituted to the manufacturers specification, is infused (2 mL/kg/hr) into the venous canular for 60

minutes. Immediately after the infusion is stopped a 2 mL blood sample is taken and platelet count, drug level, plasma fibrinogen concentration and TAT complex are determined as before. Platelet counting is performed using at Coulter T540 blood analyser. Plasma fibrinogen and TAT levels are dertermining using a clotting assay (Sigma Cat.880-B) and
5 TAT ELISA (Behring) respectively. The plasma concentration of the compound is bioassayed using human Factor Xa and a chromogenic substrate S2765 (Kabi), extrapolated from a standard curve (Fragmin) and expressed in Anti-Factor Xa units. The data is analysed as follows; tissue factor-induced reductions in platelet count are normalised with respect to pre-dose platelet count and drug activity expressed as a percent inhibition of tissue factor-
10 induced thrombocytopenia when compared to vehicle treated animals. Compounds are active if there is statistically significant ($p < 0.05$) inhibition of TF-induced thrombocytopenia.

e) An ex vivo Assay of Anticoagulant Activity

The test compound is administered intravenously or orally to a group of Alderley Park
15 Wistar rats. At various times thereafter animals are anaesthetised, blood is collected and PT coagulation assays analogous to those described hereinbefore are conducted.

f) An in vivo Measurement of Antithrombotic Activity

Thrombus formation is induced using an analogous method to that described by Vogel
20 et al., Thromb. Research, 1989, 54, 399-410. A group of Alderley Park Wistar rats is anaesthetised and surgery is performed to expose the vena cava. Collateral veins are ligated and two loose sutures are located, 0.7 cm apart, round the inferior vena cava. Test compound is administered intravenously or orally. At an appropriate time thereafter tissue thromboplastin (30 $\mu\text{l/kg}$) is administered via the jugular vein and, after 10 seconds, the two
25 sutures are tightened to induce stasis within the ligated portion of vena cava. After 10 minutes the ligated tissue is excised and the thrombus therein is isolated, blotted and weighed.

In general compounds of the formula I possess activity at the following concentrations or doses in at least one of the above tests a) to c):-

- 30 test a): IC_{50} (Factor Xa) in the range, for example, 0.001-25 μM ;
test b): IC_{50} (thrombin), for example, greater than 40 μM ;

test c): CT2 (PT) in the range, for example, 0.1-50 μ M.

A feature of the invention is a compound of formula (I), or a pharmaceutically acceptable salt thereof, for use in medical therapy.

According to a further feature of the invention there is provided a pharmaceutical
5 composition which comprises a heterocyclic derivative of the formula (I), or a pharmaceutically-acceptable salt thereof, in association with a pharmaceutically-acceptable diluent or carrier.

The composition may be in a form suitable for oral use, for example a tablet, capsule, aqueous or oily solution, suspension or emulsion; for topical use, for example a
10 cream, ointment, gel or aqueous or oily solution or suspension; for nasal use, for example a snuff, nasal spray or nasal drops; for vaginal or rectal use, for example a suppository; for administration by inhalation, for example as a finely divided powder such as a dry powder, a microcrystalline form or a liquid aerosol; for sub-lingual or buccal use, for example a tablet or capsule; or for parenteral use (including intravenous, subcutaneous, intramuscular,
15 intravascular or infusion), for example a sterile aqueous or oily solution or suspension. In general the above compositions may be prepared in a conventional manner using conventional excipients.

The amount of active ingredient (that is a heterocyclic derivative of the formula (I), or a pharmaceutically-acceptable salt thereof) that is combined with one or more excipients
20 to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will
25 generally contain about 1 mg to about 500 mg of an active ingredient.

According to a further feature of the invention there is provided a heterocyclic derivative of formula (I), or a pharmaceutically-acceptable salt thereof, for use in a method of treatment of the human or animal body by therapy.

The invention also includes the use of such an active ingredient in the production of
30 a medicament for use in:-

- (i) producing a Factor Xa inhibitory effect;

- (ii) producing an anticoagulant effect;
- (iii) producing an antithrombotic effect;
- (iv) treating a Factor Xa mediated disease or medical condition;
- (v) treating a thrombosis mediated disease or medical condition;
- 5 (vi) treating coagulation disorders; and/or
- (vii) treating thrombosis or embolism involving Factor Xa mediated coagulation.

The invention also includes a method of producing an effect as defined hereinbefore or treating a disease or disorder as defined hereinbefore which comprises administering to a warm-blooded animal requiring such treatment an effective amount of an
10 active ingredient as defined hereinbefore.

The size of the dose for therapeutic or prophylactic purposes of a compound of the formula (I) will naturally vary according to the nature and severity of the medical condition, the age and sex of the animal or patient being treated and the route of administration, according to well known principles of medicine. As mentioned above, compounds of the
15 formula (I) are useful in the treatment or prevention of a variety of medical disorders where anticoagulant therapy is indicated. In using a compound of the formula (I) for such a purpose, it will generally be administered so that a daily oral dose in the range, for example, 0.5 to 100 mg/kg body weight/day is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed, for example a dose
20 for intravenous administration in the range, for example, 0.01 to 10 mg/kg body weight/day will generally be used. For preferred and especially preferred compounds of the invention, in general, lower doses will be employed, for example a daily dose in the range, for example, 0.1 to 10 mg/kg body weight/day. In general a preferred dose range for either oral or parenteral administration would be 0.01 to 10 mg/kg body weight/day.

25 Although the compounds of formula (I) are primarily of value as therapeutic or prophylactic agents for use in warm-blooded animals including man, they are also useful whenever it is required to produce an anticoagulant effect, for example during the ex-vivo storage of whole blood or in the development of biological tests for compounds having anticoagulant properties.

30 The compounds of the invention may be administered as a sole therapy or they may be administered in conjunction with other pharmacologically active agents such as a

thrombolytic agent, for example tissue plasminogen activator or derivatives thereof or streptokinase. The compounds of the invention may also be administered with, for example, a known platelet aggregation inhibitor (for example aspirin, a thromboxane antagonist or a thromboxane synthase inhibitor), a known hypolipidaemic agent or a known

5 anti-hypertensive agent.

The invention will now be illustrated in the following Examples in which, unless otherwise stated:-

(i) evaporations were carried out by rotary evaporation in vacuo and work-up procedures were carried out after removal of residual solids by filtration;

10 (ii) operations were carried out at room temperature, that is in the range 18-25°C and under an atmosphere of an inert gas such as argon;

(iii) column chromatography (by the flash procedure) and medium pressure liquid chromatography (MPLC) were generally performed on Merck Kieselgel silica (Art. 9385) or Merck Lichroprep RP-18 (Art. 9303) reversed-phase silica obtained from E. Merck,
15 Darmstadt, Germany; alternatively high pressure liquid chromatography (HPLC) was performed on a Dynamax C-18 60Å preparative reversed-phase column;

(iv) yields are given for illustration only and are not necessarily the maximum attainable;

(v) the end-products of the formula (I) have satisfactory microanalyses and their
20 structures were confirmed by nuclear magnetic resonance (NMR) and mass spectral techniques; unless otherwise stated, CD₃SOCD₃ solutions of the end-products of the formula I were used for the determination of NMR spectral data, chemical shift values were measured on the delta scale; the following abbreviations have been used: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet;

25 (vi) intermediates were not generally fully characterised and purity was assessed by thin layer chromatographic, infra-red (IR) or NMR analysis;

(vii) melting points were determined using a Mettler SP62 automatic melting point apparatus or an oil-bath apparatus; melting points for the end-products of the formula I were generally determined after crystallisation from a conventional organic solvent such as
30 ethanol, methanol, acetone, ether or hexane, alone or in admixture; and

(viii) the following abbreviations have been used:-

DMF	<u>N,N</u> -dimethylformamide;
EtOAc	ethyl acetate;
DMSO	dimethylsulphoxide.

5 Example 1

Methyl 4-(4-pyrimidinyl)benzoate (0.41g, 1.9mmol) was stirred at room temperature in ethanol (20mL) and 2*N* NaOH_(aq) (20mL) for 1 hour. 2*N* HCl_(aq) was added until a precipitate formed. The resulting suspension was concentrated *in vacuo* and azeotroped with toluene. Thionyl chloride (100mL) and DMF (1 drop) were added and the reaction mixture refluxed for 1 hour. The reaction mixture was concentrated *in vacuo* and azeotroped twice with dichloromethane to yield 4-(4-pyrimidinyl)benzoyl chloride. The acid chloride was suspended in dichloromethane (100mL) and 1-(6-bromonaphth-2-ylsulphonyl)piperazine hydrochloride (0.545g, 1.5mmol) added as a solid in two portions followed by triethylamine (2.2mL, 15mmol). The reaction mixture was stirred overnight at room temperature then concentrated *in vacuo*. The resulting solid was separated between ethyl acetate (100mL) and water (2x100mL). The organic layer was dried over magnesium sulphate, filtered and concentrated *in vacuo* to yield a black oil which was subjected to chromatography (SiO₂: 40%, 50%, 60% Ethyl acetate / Hexane) to yield 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyrimidinyl)benzoyl]piperazine as a white solid;

¹H NMR (250MHz, DMSO-d₆) δ = 2.94 to 3.18ppm (m,4H), δ = 3.40 to 3.83ppm (m,4H), δ = 7.49ppm (d,2H), δ = 7.83ppm (m,2H), δ = 8.10ppm (dd,1H), δ = 8.14 to 8.23ppm (m,4H), δ = 8.43ppm (d,1H), δ = 8.49ppm (s,1H), δ = 8.89ppm (d,1H), δ = 9.26ppm (s,1H); MS (M+H)⁺ 536.

Methyl 4-(4-pyrimidinyl)benzoate was prepared as follows:

(a) 4(3H)-Pyrimidone (7.00g, 72.8mmol) was refluxed in thionyl chloride (50mL) and DMF (3mL) for 1 hour. A clear solution was not obtained so a further portion of thionyl chloride (50ml) was added and the reaction mixture refluxed for a further hour. The reaction mixture was concentrated *in vacuo*. Ether (500ml) was added and the solid "scratched". The resulting yellow solid was filtered off and washed with ether. A precipitate formed in the filtrate. This was filtered off to yield 4-chloropyrimidine as a pale brown solid (4.97g); ¹H

NMR (250MHz, DMSO- d_6) δ = 6.62ppm (d,1H), δ = 8.07ppm (dd,1H), δ = 9.11ppm (s,1H); MS (M+H)⁺ 115.

(b) 4-Carboxybenzene boronic acid (10.22g, 61.2mmol) was refluxed overnight in 5 methanol (70mL) with concentrated sulphuric acid (0.5mL). The reaction mixture was allowed to cool to room temperature then concentrated *in vacuo*. The resulting oil was separated between ethyl acetate (200mL) and water (200mL). The organic layer was dried over magnesium sulphate, filtered and concentrated *in vacuo* to yield the corresponding methyl ester as a white solid (9.85g); ¹H NMR (250MHz, DMSO- d_6) δ = 3.85ppm (s,3H), 10 δ = 7.89ppm (s,4H), δ = 8.19ppm (s,2H); MS (M-H)⁻ 179.

(c) 1,1'-Bis(diphenylphosphino)ferrocene (2.48g, 4.5mmol) and palladium (II) acetate (1.0g, 4.5mmol) were stirred at 50°C in toluene (25mL) under an atmosphere of nitrogen for 30 minutes, then allowed to cool to room temperature. The boronic acid methyl ester from b) 15 above (2.2g, 11.2mmol), 4-chloropyrimidine hydrochloride from a) above (1.69g, 11.2mmol) and potassium fluoride (3.9g, 67mmol) were added followed by water (25mL). The reaction mixture was refluxed overnight under an atmosphere of nitrogen. The reaction mixture was separated between ethyl acetate (100ml) and water (100ml). The organic layer was dried over magnesium sulphate, filtered and concentrated *in vacuo* to yield a black oil which was 20 subjected to chromatography (SiO₂: 100% Ethyl acetate) to yield methyl 4-(4-pyrimidinyl)benzoate as a brown solid (1.17g); ¹H NMR (250MHz, DMSO- d_6) δ = 3.91ppm (s,3H), δ = 8.13 & 8.36ppm (dd,4H), δ = 8.19ppm (dd,1H), δ = 8.94ppm (d,1H), δ = 9.32ppm (d,1H); MS (M+H)⁺ 215.

25 **Example 2**

Methyl 4-(4-pyrimidinyl)benzoate (0.72g, 3.4mmol) was stirred at room temperature in ethanol (100mL) and 2N NaOH_(aq) (20mL) for 1 hour. 2N HCl_(aq) was added until a precipitate formed. The resulting suspension was concentrated *in vacuo* and azeotroped with toluene. Thionyl chloride (100mL) and DMF (1 drop) were added and the reaction mixture 30 refluxed for 1 hour. The reaction mixture was concentrated *in vacuo* and azeotroped with toluene to yield 4-(4-pyrimidinyl)benzoate. The acid chloride was suspended in

dichloromethane (100mL) and 4-(6-chloronaphth-2-ylsulphonyl)-2-methoxycarbonylpiperazine hydrochloride (1.0g, 2.7mmol) added as a solid in one portion followed by triethylamine (3.8mL, 27mmol). The reaction mixture was stirred overnight at room temperature then concentrated *in vacuo*. The resulting solid was separated between
5 ethyl acetate (200mL) and water(2x200mL). The organic layer was dried over magnesium sulphate, filtered and concentrated *in vacuo* to yield a brown solid which was subjected to chromatography (SiO₂: 100% Ethyl acetate) to yield 1-(6-chloronaphth-2-ylsulphonyl)-3-methoxycarbonyl-4-[4-(4-pyrimidinyl)benzoyl]piperazine as an off white solid. This solid was again subjected to chromatography (SiO₂: 30%, 40%, 60%, 65% Ethyl acetate / Hexane)
10 to yield a white crystalline solid (1.051g); ¹H NMR (250MHz, DMSO-d₆)
δ = 2.75ppm (m,1H), δ = 2.92 to 4.64ppm & 5.35ppm (m,9H), δ = 7.52ppm (d,2H),
δ = 7.73ppm (dd,1H), δ = 7.80ppm (dd,1H), δ = 8.10ppm (d,1H), δ = 8.14 to 8.31ppm
(m,5H), δ = 8.51ppm (d,1H), δ = 8.89ppm (d,1H), δ = 8.89ppm (d,1H), δ = 9.26ppm (s,1H);
MS (M+H)⁺ 551.

15

Example 3

4-(4-Pyridyl)benzoic acid (238mg, 1.2mmol) was suspended in DMF (5mL) with triethylamine (0.17mL, 1.2mmol). The reaction mixture was stirred at room temperature for 15 minutes then cooled to 5°C. Carbonyldiimidazole (194mg, 1.2mmol) was added and the
20 reaction mixture allowed to warm slowly to room temperature over one hour. 4-(6-Bromonaphth-2-ylsulphonyl)piperazine hydrochloride (470mg, 1.2mmol) was added as a solid in one portion and the reaction mixture stirred for 2 days at room temperature. The reaction mixture was diluted with ethyl acetate (50mL) and washed twice with saturated aqueous sodium hydrogen carbonate solution (2x50mL). The organic layer was dried over
25 MgSO₄, filtered and concentrated *in vacuo*. The resulting solid was subjected to chromatography (SiO₂: 2%, 4%, 6%, 8%, 10% methanol / ethyl acetate) to yield 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyridyl)benzoyl]piperazine as a white solid (90mg), ¹H NMR (DMSO-d₆) δ = 3.07ppm (s,4H), δ = 3.59ppm (s,4H), δ = 7.27 & 7.46ppm (dd,4H), δ = 7.18 & 8.64ppm (dd,4H), δ = 7.34ppm (m,2H), δ = 8.17ppm (dd,2H),
30 δ = 8.39ppm (d,1H), δ = 8.47ppm (s,1H); MS (M+H)⁺ 536.

4-(4-Pyridyl)benzoic acid was prepared as follows:

- a) 4-Cyanophenylboronic acid (1.49g, 10mmol), 4-bromopyridine hydrochloride (1.97g, 10mmol), 10% palladium-on-carbon (322mg) and anhydrous sodium carbonate (2.15g, 20mmol) were refluxed in a mixture of ethanol (12mL) and water (3mL), overnight under an atmosphere of argon. The reaction mixture was filtered through diatomaceous earth, then concentrated *in vacuo*. The resulting white solid was separated between ethyl acetate (3x100mL) and water (100mL). The combined organic layers were dried over MgSO_4 , filtered and concentrated *in vacuo*. The resulting solid was subjected to chromatography (SiO₂; 50%, 60%, 70%, 80% ethyl acetate / *iso*-Hexane) to yield 4-(4-pyridyl)benzonitrile as a crystalline white solid (1.60g), ¹H NMR (DMSO-*d*₆) δ = 7.77ppm (dd, 2H), δ = 7.99ppm (m, 4H), δ = 8.69ppm (dd, 2H); MS (M+H)⁺ 181.
- b) 4-(4-Pyridyl)benzonitrile (0.5g, 2.8mmol) was dissolved in concentrated sulphuric acid (10mL) and heated at 110°C overnight. The reaction was allowed to cool to room temperature then poured into ice / water. Sodium hydroxide pellets were added until a precipitate formed. This was filtered off to yield 4-(4-pyridyl)benzamide as a white solid (0.40g), ¹H NMR (DMSO-*d*₆) δ = 7.77ppm (d, 2H), δ = 7.89ppm (d, 2H), δ = 8.02ppm (d, 2H), δ = 8.65ppm (br s, 2H); MS (M+H)⁺ 199.
- c) 4-(4-Pyridyl)benzamide (0.35g, 1.8mmol) was suspended in ethanol (5mL). 10% w/w aqueous sodium hydroxide solution was added and the reaction mixture refluxed for two hours then allowed to cool to room temperature. The reaction mixture was adjusted to pH 7 with concentrated sulphuric acid. A white precipitate formed which was isolated by filtration to give 4-(4-pyridyl)benzoic acid (238mg).

Example 4

4-(2-Pyridyl)benzoic acid (199mg, 1mmol) was suspended in DMF (5 mL) and triethylamine (0.14mL, 1mmol) was added. The reaction mixture was stirred at room temperature for 15 minutes then cooled to 5°C. Carbonyl diimidazole (162mg, 1mmol) was added and the reaction mixture allowed to warm slowly to room temperature over one hour.

1-(6-Bromonaphth-2-ylsulphonyl)piperazine hydrochloride (393mg, 1mmol) was added as a solid in one portion and the reaction mixture was stirred overnight at room temperature then concentrated *in vacuo*. The crude product was dissolved in ethyl acetate (50mL) and washed with aqueous sodium bicarbonate solution (2x50mL). The organic layer was dried over
5 MgSO₄, filtered and concentrated *in vacuo*. The resulting off white solid was subjected to chromatography (SiO₂; 100% EtOAc) to yield 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(2-pyridyl)benzoyl]piperazine as a white solid (92mg); ¹H NMR (DMSO-*d*₆) δ = 3.09ppm (s,4H), δ = 3.62ppm (s,4H), δ = 7.37ppm (t,1H), δ = 7.51 to 7.99ppm (m,4H), δ = 7.42 and 8.08ppm (dd,4H), δ = 8.17ppm (dd,2H), δ = 8.40ppm (d,1H), δ = 8.47ppm (s,1H), δ =
10 8.66ppm (dd,1H); MS (M+H)⁺ 536.

Example 5

1-(6-Chloronaphth-2-ylsulphonyl)-4-(4-iodobenzoyl)piperazine (920mg, 1.7mmol), diethyl 3-pyridylborane (250mg, 1.7mmol), tetrabutyl ammonium bromide (110mg, 0.34mmol), tetrakis(triphenylphosphine) palladium (0) (69mg,0.06mmol) and potassium
15 hydroxide (286mg, 5.1mmol) were refluxed in dry tetrahydrofuran (100mL) for two hours under an atmosphere of nitrogen. The reaction mixture was allowed to cool to room temperature then concentrated *in vacuo*. The resulting solid was subjected to flash chromatography (SiO₂: 100% CH₂Cl₂, 50% EtOAc / Hexane, 100% EtOAc) to yield an oil. The oil was dissolved in EtOAc (100mL) and washed with saturated sodium hydrogen
20 carbonate solution (100mL). The EtOAc layer was separated, dried over MgSO₄, filtered and concentrated *in vacuo* to yield an off white foam which was triturated with hexane. This solid was subjected to chromatography (SiO₂: 40%, 50%, 60%, 70% EtOAc / Hexane) to yield 1-(6-chloronaphth-2-ylsulphonyl)-4-[4-(3-pyridyl)benzoyl]piperazine as a white solid (322.4mg); ¹H NMR (DMSO-*d*₆) δ = 3.1ppm (s,4H), δ = 3.6ppm (s,4H),
25 δ = 7.46ppm (d,2H), δ = 7.5 to 7.67ppm (m,1H), δ = 7.70 to 7.79ppm (m,3H), δ = 7.83ppm (dd,1H), δ = 8.09ppm (dt,1H), δ = 8.15ppm to 8.31ppm (m,3H), δ = 8.50ppm (s,1H), δ = 8.60ppm (dd,1H), δ = 8.90ppm (d,1H); MS (M+H)⁺ 492.

1-(6-Chloronaphth-2-ylsulphonyl)-4-(4-iodobenzoyl)piperazine was prepared as
30 follows:

1-(6-Chloronaphth-2-ylsulphonyl)piperazine (0.65g, 2.1mmol) was dissolved in dichloromethane (50mL) and triethylamine (2.9mL, 21mmol) was added at room temperature. 4-Iodobenzoyl chloride (0.56g, 2.1mmol) was added as a solid in one portion and the reaction mixture stirred for one hour at room temperature. The reaction mixture was concentrated *in vacuo*, then separated between EtOAc (100mL) and saturated sodium hydrogen carbonate solution (100mL). The EtOAc layer was separated, dried over MgSO₄, filtered and concentrated *in vacuo*. The resulting white solid was subjected to chromatography (SiO₂: 50% EtOAc / Hexane) to yield 1-(6-chloronaphth-2-ylsulphonyl)-4-(4-iodobenzoyl)piperazine as a white solid (0.97g); ¹H NMR (DMSO-*d*₆) δ = 3.05ppm (s,4H), δ = 3.53ppm (s,4H), δ = 7.11ppm & 7.74ppm (dd,4H), δ = 7.65 to 7.84ppm (m,2H), δ = 8.10ppm to 8.28ppm (m,3H), δ = 8.47ppm (s,1H); MS (M+H)⁺ 540.

Example 6

4-(2-Methyl-4-(3-pyridyl)benzoyl)piperazine (0.41g, 1.5mmol) was dissolved in dichloromethane (50mL) at room temperature. 6-Chloronaphthalenesulphonyl chloride (0.38g, 1.5mmol) and triethylamine (1ml, 7.5mmol) were added and the reaction mixture stirred overnight. The reaction mixture was concentrated *in vacuo*. The crude product was dissolved in EtOAc (100mL) and washed with saturated sodium hydrogen carbonate solution (100mL). The EtOAc layer was separated, dried over MgSO₄, filtered and concentrated *in vacuo*. The resulting solid was subjected to chromatography (SiO₂: 50%, 60% EtOAc / Hexane) to yield 1-(6-chloronaphth-2-ylsulphonyl)-4-[2-methyl-4-(3-pyridyl)benzoyl]piperazine as a white solid (508.4mg); ¹H NMR (DMSO-*d*₆) δ = 2.15ppm (s,3H), δ = 2.94 to 3.25ppm (m,6H), δ = 3.78ppm (m,2H), δ = 7.23ppm (d,1H), δ = 7.44 to 7.62ppm (m,3H), δ = 7.70ppm (dd,1H), δ = 7.84ppm (dd, 1H), δ = 8.08ppm (m,1H), pyridyl 5-H), δ = 8.12 to 8.28ppm (m,3H), δ = 8.48ppm (s,1H), δ = 8.59ppm (dd,1H), δ = 8.88ppm (d,1H); MS (M+H)⁺ 506.

4-(2-Methyl-4-(3-pyridyl)benzoyl)piperazine was prepared as follows:

- 30 a) 4-Bromo-2-methylbenzoic acid (11.55g, 53.7mmol) was suspended in thionyl chloride (40mL). One drop of DMF was added and the resulting mixture was stirred at 69°C

until gas evolution ceased. The reaction mixture was concentrated *in vacuo* and azeotroped twice with toluene. The resulting oil was dissolved in dichloromethane (100mL) and cooled to 5°C. *tert*-Butyl-1-piperazine carboxylate (10.0g, 53.7mmol) was added portionwise over one hour followed by triethylamine (37.5mL, 260mmol). The reaction mixture was stirred 5 overnight at room temperature then concentrated *in vacuo*. The crude product was dissolved in ethyl acetate (750mL) and washed with aqueous 2*N* sodium hydroxide solution and then with brine. The organic layer was dried over MgSO₄, filtered and concentrated *in vacuo* to yield a brown oil. The product crystallised from EtOAc / Hexane to yield beige crystals (9.01g); ¹H NMR (DMSO-*d*₆) δ = 1.40ppm (s,9H), δ = 2.20ppm (s,3H), δ = 3.10ppm (m,2H), 10 δ = 3.26ppm (m,2H), δ = 3.41ppm (m,2H), δ = 3.62ppm (s,2H), δ = 7.15ppm (d,1H), δ = 7.44ppm (dd,1H), δ = 7.52ppm (d,1H); MS (M+H)⁺ 383.

b) The product from a) above (3.83g, 10mmol), diethyl 3-pyridylborane (1.47g, 15 10mmol), tetrabutyl ammonium bromide (0.65g, 2mmol), tetrakis(triphenylphosphine) palladium (0) (0.40g, 0.35mmol) and potassium hydroxide (1.68g, 30mmol) were refluxed in dry tetrahydrofuran (75mL) for two hours under an atmosphere of nitrogen. The reaction mixture was allowed to cool to room temperature then separated between EtOAc (250mL) and water (2x500mL). The EtOAc layer was separated, dried over MgSO₄, filtered and 20 concentrated *in vacuo*. The resulting oil was subjected to flash chromatography (SiO₂: 100% EtOAc) to yield *t*-butyl 4-[2-methyl-4-(3-pyridyl)benzoyl]piperazine carboxylate as a brown solid (1.78g); ¹H NMR (DMSO-*d*₆) δ = 1.41ppm (s,9H), δ = 2.60ppm (s,3H), δ = 3.17ppm (m,2H), δ = 3.27ppm (m,2H), δ = 3.44ppm (m,2H), δ = 3.66ppm (s,2H), δ = 7.61ppm (d,1H), δ = 7.49ppm (dd,1H), 25 δ = 7.59ppm (dd,1H), δ = 7.61ppm (s,1H), δ = 8.08ppm (m,1H), δ = 8.58ppm (dd,1H), δ = 8.90ppm (d,1H); MS (M+H)⁺ 382.

c) The product from b) above (1.66g, 4.35mmol) was dissolved in dichloromethane (50mL) at room temperature. Trifluoroacetic acid (10mL) and the reaction mixture stirred for 30 one hour at room temperature. The reaction mixture was concentrated *in vacuo*, then

separated between EtOAc and 2*N* sodium hydroxide solution. The EtOAc layer was separated, dried over MgSO₄, filtered and concentrated *in vacuo* to yield 4-[2-methyl-4-(3-pyridyl)benzoyl]piperazine as a colourless oil (410g).

5 Example 7

A solution of 4-(4-pyridyl)benzoic acid (398 mg), hydroxybenztriazole (338 mg) and dicyclohexylcarbodiimide (453 mg) were stirred at ambient temperature for 1 hour then 1-(6-chloronaphth-2-ylsulphonyl)piperazine (621 mg) was added to the resultant white suspension and stirring continued at ambient temperature for a further 16 hours. The reaction mixture was filtered and the filtrate concentrated *in vacuo* to a foam which was partially purified by flash chromatography at 3psi on silica (Merck ART 9385) eluting with 2.5% v/v methanol in dichloromethane plus 0.1% SG 0.88 ammonia solution. This gave a white foam (473 mg). A portion of this foam (110 mg) was purified by preparative HPLC on a Dynamax® C-18 60Å column eluting with 0.1% trifluoroacetic acid in aqueous acetonitrile over a gradient of 30%-70% acetonitrile. The solution was freeze-dried to produce a white foam. There was thus obtained 1-(6-chloronaphth-2-ylsulphonyl)-4-[4-(4-pyridyl)benzoyl]piperazine the title compound as a trifluoroacetate salt (83.5mg); mp 175-176°C; ¹H NMR (250MHz, DMSO-d₆ at 373 °K) δ = 3.17 (t,4H), 3.47 (t,4H), 7.47 (d,2H), 7.62-7.86 (m,6H), 8.08-8.48 (m,3H), 8.43 (s,1H), 8.66 (d,2H) ppm; MS: m/z 492/494 (M+H)⁺ (1 Cl pattern).

Example 8

4-(2-Methyl-4-pyridyl)benzoic acid (62mg), 1-(6-bromonaphth-2-ylsulphonyl)piperazine (94mg) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (61mg) were dissolved in DMF (2.5ml) and the resultant solution stirred at ambient temperature for 16 hours. Excess DMF was removed *in vacuo*, water (10ml) was added and the precipitate that formed was filtered, washed thoroughly with cold water and dried over P₂O₅. The solid thus obtained was purified by flash chromatography at 3psi on silica (Merck ART 9385) eluting with 2.5% v/v methanol in dichloromethane. There was thus obtained the 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(2-methyl-4-pyridyl)benzoyl]piperazine (99mg); mp 204-205°C; ¹H NMR (300 MHz, DMSO-d₆) δ = 2.48 (s,3H), 3.03 (s,4H), 3.57 (broad s, 4H), 7.46

(t,3H), 7.56 (s,1H), 7.77(d,2H), 7.83(d,2H), 8.17(q,2H), 8.42 (s,1H), 8.48 (d,2H) ppm; MS: m/z 550/552 (M+H)⁺ (1 Br pattern).

4-(2-Methyl-4-pyridyl) benzoic acid was prepared as follows :-

5

(a) iso-Amyl nitrite (7.9g) was slowly added to a solution of ethyl 4-aminobenzoate (4.95g) in 2-picoline (100ml) at ambient temperature. The resultant mixture was heated at 100°C for 2 hours then excess 2-picoline was removed *in vacuo* to give a black oil. The mixture of isomers thus obtained was purified by flash chromatography at 3psi on silica (Merck ART 10 9385) eluting with 25% v/v ethyl acetate in iso-hexane. There was thus obtained ethyl 4-(2-methyl-4-pyridyl)benzoate (0.2g) as brown gum of sufficient purity to continue; ¹H NMR (300MHz, DMSO-d₆) δ = 1.32 (t,3H), 2.53 (s,3H), 4.33 (q,2H), 7.52 (d,1H), 7.61 (s,1H), 7.92 (d,2H), 8.05 (d,2H), 8.52 (d,1H) ppm; MS: m/z 242 (M+H)⁺.

15 (b) Ethyl 4-(2-methyl-4-pyridyl)benzoate (185mg) was dissolved in methanol (7.5ml) and 1.0M NaOH (3.75ml) and heated at 60°C for 3 hours. The resulting mixture was reduced to low volume then water (10ml) added, the solution neutralised to pH 7 with 1.0M HCl, the resulting precipitate filtered and dried over P₂O₅ to give 4-(2-methyl-4-pyridyl)benzoic acid as a light brown solid (73mg); mp 293-294°C; ¹H NMR (300MHz, DMSO-d₆) δ = 2.52 (s,3H), 20 7.53 (d,1H), 7.62 (s,1H), 7.89 (d,2H), 8.04 (d,2H), 8.52 (d,1H) ppm. MS: m/z 214 (M+H)⁺.

Example 9

4-(4-Pyridazinyl)benzoic acid (300mg), 1-(6-bromonaphth-2-ylsulphonyl) piperazine 25 (484mg) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (317mg) were dissolved in DMF (7.5ml) and the resultant solution stirred at ambient temperature for 16 hours. Water (50ml) was added and the precipitate that formed was filtered, washed thoroughly with cold water and dried over P₂O₅. There was thus obtained 1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyridazinyl)benzoyl]piperazine (535mg); mp 128-130°C; ¹H NMR 30 (300MHz, DMSO-d₆ at 373°K) δ = 3.12 (s,4H), 3.57 (s,4H), 7.48 (d,2H), 7.80 (m,2H), 7.89 (d,2H), 7.94 (m,1H), 8.14 (d,2H), 8.39 (s,1H), 8.46 (s,1H), 9.26 (d,1H), 9.58 (s,1H) ppm; MS: m/z 537/539 (M+H)⁺ (1 Br pattern).

4-(4-Pyridazinyl)benzoic acid was prepared as follows :-

(a) A solution of sodium nitrite (1.44g) in water (3.0ml) was added slowly to a stirred solution of ethyl 4-aminobenzoate (3.3g) in 48% fluoroboric acid (9.4ml) at 0°C. After the final addition the mixture was stirred at 0°C for a further 0.5 hours, then filtered and washed with cold fluoboric acid (5.0ml), then with ethanol and finally with diethylether. The ethyl (4-diazonium tetrafluoroborate)benzoate (4.60g) thus obtained was mixed dry with potassium acetate (3.40g) and 18-crown-6 (0.23g) then treated with pyridazine (25ml) at ambient temperature. The mixture rapidly turned black with evolution of nitrogen gas. After stirring at ambient temperature for 16 hours the excess pyridazine was removed *in vacuo* and the black residue thus obtained dissolved in ethyl acetate (50ml) and washed with water (50ml). The organic layer was dried (MgSO₄) and reduced to a black residue. The mixture of isomers thus obtained was then purified by flash chromatography on Merck ART 9385 silica eluting with ethyl acetate to give ethyl 4-(4-pyridazinyl) benzoate (1.04g); mp 110-112°C; ¹H NMR (300MHz, CDCl₃) δ = 1.42 (t,3H), 4.43 (q,2H), 7.68 (dxd,1H), 7.75 (d,2H), 8.22 (d,2H), 9.28, (d,1H), 9.50 (d,1H) ppm; MS: m/z 229 (M+H)⁺.

(b) Ethyl 4-(4-pyridazinyl)benzoate (580mg) was dissolved in methanol (12.5ml) and 1.0M NaOH (12.7ml) and heated at 60°C for 4 hours. The resulting mixture was reduced to low volume then water (25ml) added, the solution neutralised to pH 7 with 1.0M HCl, the resulting precipitate filtered and dried over P₂O₅ to give 4-(4-pyridazinyl)benzoic acid as a light brown solid (503mg); mp >330°C; ¹H NMR (300 MHz, DMSO-d₆) δ = 8.05 (m,5H), 9.32 (d,1H), 9.67 (s,1H) ppm; MS: m/z 201 (M+H)⁺.





Example 10

6-Bromonaphth-2-ylsulphonyl chloride (470mg) was added in one portion to a mixture of 1-[2-methoxycarbonyl-4-(4-pyridyl)benzoyl]piperazine (500mg) and triethylamine (311mg) in dichloromethane (5ml) at ambient temperature. After 10 minutes the mixture was reduced *in vacuo* and the residue thus obtained purified by flash chromatography at 3psi on silica (Merck ART 9385) eluting first with dichloromethane, then 1% and 2% v/v

methanol in dichloromethane. There was thus obtained 1-(6-bromonaphth-2-ylsulphonyl)-4-[2-methoxycarbonyl-4-(4-pyridyl)benzoyl]piperazine (866mg) as a foam; ^1H NMR (250MHz, CDCl_3) δ = 3.03 (t,2H), 3.27 (t,2H), 3.34 (t,2H), 3.56 (s,3H), 3.94 (broad s,2H), 7.32 (d,1H) 7.48 (dxd,2H), 7.70-7.82 (m,3H), 7.86 (d,1H), 7.92(d,1H), 8.13(d,1H), 8.24(d,1H), 8.32(s,1H), 8.71(dxd,2H) ppm; MS: m/z 594/596 ($\text{M}+\text{H}$) $^+$ (1 Br pattern).

1-[2-methoxycarbonyl-4-(4-pyridyl)benzoyl]piperazine was prepared as follows :-

10



15

A solution of t-butyloxycarbonylpiperazine (14.4g) in chloroform (20ml) was added dropwise to a solution of 4-bromophthalic anhydride (17.5g) in chloroform (50ml) at ambient temperature. The reaction mixture was stirred at 60°C for 1 hour then reduced *in vacuo* to an oil. A mixture of isomers (31.9g) was obtained (structures 'C' and 'D').

20

Potassium carbonate (10.7g) and dimethyl sulphate (9.71g) were added to a solution of the isomeric mixture of acids (C+D) in acetone (60ml) and the mixture stirred for 2 hours at ambient temperature. The solution containing the esters (E+F) was filtered and the filtrate reduced *in vacuo* to a gum (33.0g) which was purified by preparative HPLC using PhaseSep NP Silica, 15-35 μm , 60Å and eluting with 25% - 50% v/v ethyl acetate in iso-hexane. There

was thus obtained structure 'E' (12.86g), the slower running isomer; mp 131-132°C; ¹H NMR (400MHz, CDCl₃) δ = 1.46 (s,9H), 3.14 (t,2H), 3.36 (broad s, 2H), 3.56 (broad s, 2H), 3.77 (broad s,2H), 3.89 (s,3H), 7.16 (d,1H), 7.71 (dxd,1H), 8.17 (d,1H) ppm; MS: m/z 427/429 (M+H)⁺ (1 Br pattern).

5 A solution of structure 'E' (4.27g) in dry DMF (40ml) was deoxygenated by bubbling argon for 5 minutes then tetrakis-triphenylphosphine palladium(0) (1.15g), bis-triphenylphosphine palladium dichloride (0.70g) and silver(1)oxide (2.32g) were added and the mixture stirred for 5 minutes at 100°C then trimethyl-(4-pyridyl)-stannane (3.63g) was added and heating at 100°C continued for 15 minutes. The mixture was allowed to cool and
10 stirred at ambient temperature for 20 hours then filtered through diatomaceous earth and reduced *in vacuo* to a black residue which was stirred with 1.0M potassium fluoride (20ml) for 1 hour then extracted with ethyl acetate (3x25ml). The extracts were dried (MgSO₄), filtered and reduced to a black oil which was partially purified by flash chromatography at 3psi on silica (Merck ART 9385) eluting with 25% and 50% v/v ethyl acetate in iso-hexane,
15 then 2% and 4% v/v methanol in dichloromethane. A final purification by BIOTAGE® P45 MPLC eluting with ethyl acetate at 10psi gave compound 1-tert-butoxycarbonyl-4-[2-methoxycarbonyl-4-(4-pyridyl)benzoyl]piperazine (1.94g); mp 144-146°C; ¹H NMR (300MHz, CDCl₃) δ = 1.46 (s,9H), 3.20 (t,2H), 3.39 (m,2H), 3.60 (m,2H), 3.81 (m,2H), 3.93 (s,3H), 7.42 (d,1H), 7.52 (d,2H), 7.84 (dxd,1H), 8.3 (d,1H), 8.73 (d,2H) ppm; MS: m/z 370
20 (M+H)⁺.

A 2.2 molar solution of hydrogen chloride in diethyl ether (2.9ml) was added to a solution of structure 'B' (1.05g) in dichloromethane (10ml) and the mixture stirred for 4 hours at ambient temperature. The supernatant liquors were decanted and the remaining gum was triturated with diethyl ether to give a white solid which was treated with saturated aqueous
25 sodium bicarbonate then extracted with ethyl acetate (3 x 10ml). The combined organic extracts were dried (MgSO₄), filtered and reduced to give structure 1-[2-methoxycarbonyl-4-(4-pyridyl)benzoyl]piperazine (500mg) as a white foam; ¹H NMR (250MHz, DMSO-d₆) δ = 2.45 (m,2H), 2.62 (m,2H), 2.90 (m,2H), 3.39 (m,2H), 3.68 (s,3H), 7.33 (d,1H), 7.62 (dxd,2H), 7.93 (dxd,1H), 8.10 (d,1H), 8.53 (dxd,2H) ppm; MS: m/z 326 (M+H)⁺.

Example 11

A stirred suspension of 4-(4-pyridyl)benzoic acid (sodium salt) (190 mg, 0.86 mmol) in dichloromethane (10 ml) was treated with oxalyl chloride (0.2 ml, 2.3 mmol) and DMF (catalytic amount). After stirring for 2 hours, further oxalyl chloride 0.2 ml, 2.3 mmol) and DMF (catalytic amount) was added and the suspension stirred a further 4 hours. The solvent was removed *in vacuo* and the residue, after drying, was suspended in dichloromethane (20 ml) and treated with 2-(hydroxymethyl)-4-(6-bromonaphth-2-ylsulphonyl)piperazine (300 mg, 0.78 mmol) and triethylamine (0.36 ml, 2.5 mmol). After stirring at room temperature overnight the reaction mixture was diluted with dichloromethane (20 ml) and water (20 ml). A copious precipitate appeared which was filtered off, dried and recrystallised from ethyl acetate (10 ml) to yield 1-(6-bromonaphth-2-ylsulphonyl)-3-(hydroxymethyl)-4-[4-(4-pyridyl)benzoyl] piperazine as a colourless solid (250 mg); ^1H NMR (300MHz, DMSO- d_6) δ = 3-4ppm (broad, 9H), δ = 7.2ppm (d, 2H), δ = 7.7ppm (d, 2H), δ = 7.8ppm (m, 4H), δ = 8.2ppm (t, 2H), δ = 8.4ppm (s, 1H), δ = 8.45ppm (s, 1H), δ = 8.6ppm (d, 2H); signals due to ethyl acetate (1 mol eq) were also present; MS: (M+H) $^+$ 566/568 (1 Br pattern); analysis; found: C, 56.8; H, 4.9; N, 6.3 %; $\text{C}_{27}\text{H}_{24}\text{BrN}_3\text{SO}_4 \cdot \text{C}_4\text{H}_8\text{O}_2$ requires: C, 56.9; H, 4.9; N, 6.4 %.

3-(Hydroxymethyl)-4-(6-bromonaphth-2-ylsulphonyl) piperazine was prepared as follows:

20

3-(Hydroxymethyl)piperazine monoacetate (1.1 g, 6.25 mmol) and triethylamine (2.2 ml, 2.5 eq) were stirred at room temperature in dichloromethane (50 ml) and the suspension treated with 6-bromonaphth-2-ylsulphonyl chloride (2.0 g, 6.5 mmol). The mixture was stirred overnight and then diluted with further dichloromethane (50 ml); the solution was washed sequentially with water, saturated sodium hydrogen carbonate solution and brine. Drying (PS paper) and evaporation yielded a colourless foam (1 g). This was subjected to chromatography (SiO_2 ; dichloromethane : methanol 19 : 1 v/v) to yield 2-(hydroxymethyl)-4-(6-bromonaphth-2-ylsulphonyl)piperazine as a colourless foam (670 mg) which was used without further purification, ^1H NMR (300 MHz, CDCl_3) δ = 2.3ppm (t, 1H), δ = 2.5ppm (dt, 1H), δ = 2.9 - 3.1ppm (m, 3H), δ = 3.5ppm (dd, 1H), δ = 3.6ppm (m, 3H), δ = 7.6 - 8.0ppm (m, 4H), δ = 8.1ppm (s, 1H), δ = 8.3ppm (s, 1H); MS (M+H) $^+$ 385/387 (1 Br pattern).

30

Example 12

A stirred suspension of 4-(4-pyridyl)benzoic acid (133 mg, 0.67 mmol) in dimethylformamide (5 ml) was treated sequentially with 1-hydroxybenzotriazole hydrate (108 mg, 0.8 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (153 mg, 0.8 mmol) and 1-(5-chlorobenzofuran-2-ylsulphonyl) piperazine (201 mg, 0.67 mmol). After stirring overnight the solvent was removed *in vacuo* and the residue chromatographed (Merck Art 9385 silica, eluting with dichloromethane containing 2% v/v of methanol) to yield 1-(5-chlorobenzofuran-2-ylsulphonyl)-4-[4-(4-pyridyl)benzoyl] piperazine as a colourless solid (40 mg), ¹H NMR (CDCl₃) δ = 3.2 - 3.4 ppm (broad s, 4H), δ = 3.6 - 4.0 ppm (broad s, 4H), δ = 7.35 ppm (s, 1H), δ = 7.5 ppm (m, 6H), δ = 7.7 ppm (m, 3H), δ = 8.7 ppm (d, 2H), MS (M+H)⁺ 482/484.

1-(5-Chlorobenzofuran-2-ylsulfonyl) piperazine was prepared as follows:




15 A stirred solution of piperazine (1.15g, 13.4 mmol) and triethylamine (4.7 ml, 46.5 mmol) in dichloromethane (30 ml) was cooled to ~5 °C, and a solution of 5-chlorobenzofuran-2-sulphonyl chloride (1.69g, 7.8 mmol) in dichloromethane (10 ml) was added. Stirring was continued for 15 mins, and the reaction mixture then allowed to warm to ambient temperature over 2 hrs with stirring. Water was added to the reaction mixture, and the organic layer separated; this was washed with water (twice), brine (once), then dried (MgSO₄), filtered and evaporated to give a yellow gum. This was chromatographed (Merck Art 9385 silica, eluting with dichloromethane containing increasing amounts of methanol, up to 10% v/v) to give a yellow solid; trituration with diethyl ether gave 5-chlorobenzofuran-2-ylsulphonyl piperazine 25 as a colourless solid (1.11g) which was used without further purification. ¹H NMR (CDCl₃) δ = 2.8 - 3.0 ppm (t, 4H), δ = 3.2 - 3.4 ppm (t, 4H), δ = 7.3 ppm (s, 1H), δ = 7.45 ppm (dd, 2H), δ = 7.7 ppm (s, 1H); MS (M+H)⁺ 301/303.

The requisite 5-chlorobenzofuran-2-ylsulphonyl chloride starting material was prepared as 30 described in European Patent Application 0 355 827 (Mochida, Hydantoin derivatives).

Example 13

Further examples are described in Table 1




TABLE 1

No.	Compound	Method	MS: m/z	¹ H NMR (MHz, solvent)
1		4-(2-methylpyrimidin-4-yl)benzoyl chloride ¹ + 1-(6-bromonaphth-2-ylsulphonyl)piperazine	551/553 (M+H) ⁺ 1 Br pattern	(300MHz, DMSO-d ₆) δ = 2.66 (s,3H), 3.08 (brs,4H), 3.53 (brm,4H), 7.46 (d,2H), 7.84 (m,3H), 8.18 (m,4H), 8.40 (s,1H), 8.48 (s, 1H), 8.76 (d,1H) ppm.
2		4-(2,6-dimethylpyrimidin-4-yl)benzoic acid ² + 1-(6-bromonaphth-2-ylsulphonyl)piperazine: reaction using 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide	565/567 (M+H) ⁺ 1 Br pattern	(300MHz, DMSO-d ₆) δ = 2.46 (s, 3H), 2.61 (s,3H), 3.08 (brm,4H), 3.55 (brm,4H), 7.46 (d,2H), 7.80 (m,3H), 8.16 (m,4H), 8.41 (s,1H), 8.46 (s,1H) ppm
3		4-(4-pyrimidinyl)benzoic acid + 1-(6-chloronaphth-2-ylsulphonyl)piperazine: reaction using 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.	493/495 (M+H) ⁺ 1 Cl pattern	(300MHz, DMSO-d ₆) δ = 3.0-3.2 (m,4H), 3.4-3.8 (m,4H), 7.25 (d,2H), 7.7 (d,1H), 7.8 (d,1H), 8.05 (d,1H), 8.1-8.3 (m,5H), 8.5 (s,1H), 8.9 (d,1H), 9.25 (s,1H) ppm

5 ¹ Prepared according to the method described in Example 1 & 1 (c) starting from 2-methyl-4-chloropyrimidine

² Prepared according to the method described in Example 1 (c) starting from 2,6-dimethyl-4-chloropyrimidine and subsequent ester hydrolysis




TABLE 1 (cont.)

No.	Compound	Method	MS: m/z	¹ H NMR (MHz, solvent)
4		4-(4-pyridyl)benzoyl chloride + 4-(6-bromonaphth-2-ylsulphonyl)-2-ethoxycarbonyl piperazine.	608/610 (M+H) ⁺ 1 Br pattern	(300MHz, CDCl ₃) δ = 2.6-3.2 (t,3H), 2.3-2.56 (b,1H), 2.56-2.74 (b,1H), 3.60-3.82 (m,2H), 4.16-4.40 (m,3H), 4.40-4.56 (m,1H), 5.44-5.56 (b,1H), 7.38-7.56 (m,4H), 7.56-7.70 (d,2H), 7.70-7.82 (td,2H), 7.82-7.94 (m,2H), 8.14 (s,1H), 8.32 (s,1H), 8.64-8.72 (d,2H) ppm.
5		4-(3-fluoro-4-pyridyl)benzoic acid ³ + 1-(6-bromonaphth-2-ylsulphonyl)piperazine: reaction using 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.	554/556 (M+H) ⁺ 1 Br pattern	(300MHz, DMSO-d ₆) δ = 3.15 (s,4H), 3.48-3.88 (bm,4H), 7.57 (d,2H), 7.68-7.80 (m,3H), 7.92 (t,2H), 8.25 (t,2H), 8.50 (s,1H), 8.58 (m,2H), 8.74 (s,1H) ppm.
6		2-methoxy-4-(4-pyridazinyl)benzoic acid ⁴ + 1-(6-bromonaphth-2-ylsulphonyl)piperazine: reaction using 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.	567/569 (M+H) ⁺ 1 Br pattern	(300MHz, DMSO-d ₆) δ = 2.94 (m,4H), 3.22 (m,2H), 3.63 (s,3H), 3.72 (s,2H), 7.26 (d,1H), 7.45 (d,1H), 7.47 (s,1H), 7.82 (t,2H), 8.02 (dxd,1H), 8.26 (d,1H), 8.02 (d,1H), 8.40 (s,1H), 8.48 (s,1H), 9.27 (d,1H), 9.64 (m,1H) ppm.

³ Prepared according to the method described in Example 1(c) starting from 3-fluoro-4-iodopyridine and subsequent ester hydrolysis

5 ⁴ Prepared according to methods described in Example 9 (a) and (b) starting from methyl 4-amino-2-methoxybenzoate

TABLE 1 (cont.)




No.	Compound	Method	MS: m/z	¹ H NMR (MHz, solvent)
7		4-(1-oxo-4-pyridyl)benzoic acid ⁵ + 1-(6-bromonaphth-2-ylsulphonyl)piperazine: reaction using 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide + 1-hydroxybenztriazole.	552/554 (M+H) ⁺ 1 Br pattern	(300MHz, CDCl ₃) δ = 3.15 (bs,4H), 3.75 (b,4H), 7.45 (m, 4H), 7.58 (d,2H), 7.75 (t,2H), 7.85 (d,1H), 7.92 (d,1H), 8.12 (s,1H), 8.25 (d,2H), 8.30 (s,1H) ppm.
8		4-(2-cyano-4-pyridyl)benzoic acid ⁶ + 1-(6-bromonaphth-2-ylsulphonyl)piperazine: reaction using 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.	561/563 (M+H) ⁺ 1 Br pattern	(300MHz, CDCl ₃) δ = 3.15 (bs,4H), 3.72 (b,4H), 7.46 (2h,d), 7.60 - 7.80 (m,5H), 7.86 (d,2H), 7.93 (d,1H), 8.13 (s,1H), 8.30 (s,1H), 8.79 (s,1H) ppm.
9		4-(2-amino-4-pyridyl)benzoic acid ⁷ + 1-(6-bromonaphth-2-ylsulphonyl)piperazine: reaction using 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.	551/553 (M+H) ⁺ 1 Br pattern	(300MHz, DMSO-d ₆) δ = 3.06 (bs,4H), 3.20 - 3.80 (b, 4H + H ₂ O), 7.15 (s,1H), 7.18 (s,1H), 7.49 (d,2H), 7.76 (d,2H), 7.81 (t,2H), 7.90 - 8.04 (m,3H), 8.18 (t,2H), 8.40 (s,1H), 8.47 (s,1H) ppm.

⁵ Prepared from 4-(4-pyridyl)benzoic acid described in Example 3 (c) by esterification, then oxidation with 3-chloroperoxybenzoic acid, and subsequent ester hydrolysis.

⁶ Prepared by a palladium catalysed coupling of 4-chloro-2-cyanopyridine and 4-carboxyphenylboronic acid using bis-(tri-σ-tolylphosphine)palladiumchloride.

⁷ Prepared by a palladium catalysed coupling of 2-(N-tert-butoxycarbonylamino)-4-bromopyridine and 4-carboxyethylphenylboronic acid using [1,1'-bis(diphenylphosphino)ferrocene]diacetyl palladium.

TABLE 1 (cont.)

No.	Compound	Method	MS: m/z	¹ H NMR (MHz, solvent)
10		1-(6-bromonaphth-2-ylsulphonyl)-4-[2-carboxy-4-(4-pyridyl)benzoyl]piperazine ⁸ + diethanolamine: reaction using 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide + 1-hydroxybenztriazole.	667/669 (M+H) ⁺ 1 Br pattern	(300MHz, DMSO-d ₆ at 373°K) δ = 3.18 (t,4H), 3.31 (t,4H), 3.47 (m,8H), 7.38 (d,1H), 7.63 (dd,2H), 7.74 (m,2H), 7.79 (m,1H), 7.82 (dd,1H), 8.08 (d,1H), 8.13 (d,1H), 8.31 (d,1H), 8.42 (d,1H), 8.63 (m,2H) ppm.
11		4-(4-pyridyl)benzoic acid + 1-[(E)-4-chlorostyrylsulphonyl]piperazine ⁹ : reaction using 1,3-dicyclohexylcarbodiimide + 1-hydroxybenztriazole.	468/470 (M+H) ⁺ 1 Cl pattern	(300MHz, DMSO-d ₆) δ = 3.20 (m, 4H), 3.65 (bs,4H), 7.33 (d,1H), 7.46 (d,1H), 7.49 (d,2H), 7.56 (d,2H), 7.72 (d, 2H), 7.81 (d,2H), 7.86 (d,2H), 8.65 (d,2H) ppm.
12		6-bromonaphth-2-ylsulphonyl chloride + 2-oxo-1-[4-(4-pyrimidinyl)benzoyl]piperazine ¹⁰	552 (M+H) ⁺	(300MHz, CDCl ₃) δ = 3.64 (dd,2H), 4.00 (m,4H), 7.45 (d,2H), 7.70 - 7.95 (m,4H), 7.97 - 8.05 (m,3H), 8.18 (s,1H), 8.42 (s,1H), 8.82 (d,1H), 9.30 (s,1H) ppm.

⁸ Prepared from the ester described in Example 10 by base hydrolysis.


⁹ Described in WO96/10022 Ex57 page109 last paragraph

¹⁰ Prepared by reacting 4-(4-pyrimidinyl)benzoyl chloride with 1-tert-butoxycarbonyl-3-oxopiperazine and subsequent treatment with trifluoroacetic acid.

Claims

1. A compound of formula (I)

5



wherein:

A is 5- or 6-membered monocyclic aromatic ring containing 1, 2 or 3 ring heteroatoms

10 selected from nitrogen, oxygen and sulphur atoms optionally substituted by one, two or three atoms or groups selected from halo, oxo, carboxy, trifluoromethyl, cyano, amino, hydroxy, nitro, C₁₋₄alkyl (for example methyl or ethyl), C₁₋₄alkoxy (for example methoxy or ethoxy), C₁₋₄alkoxycarbonyl, C₁₋₄alkylamino (for example methylamino or ethylamino) or di-C₁₋₄alkylamino (for example dimethylamino or diethylamino);

15

B is a phenylene ring optionally substituted by one or two substituents selected from halo, trifluoromethyl, trifluoromethoxy, cyano, nitro, C₁₋₄alkyl, C₂₋₄alkenyl and C₂₋₄alkynyl, from the substituent -(CH₂)_nY¹ wherein n is 0-4 and Y¹ is selected from hydroxy, amino, carboxy, C₁₋₄alkoxy, C₂₋₄alkenyloxy, C₂₋₄alkynyloxy, C₁₋₄alkylamino, di-C₁₋₄alkylamino, pyrrolidin-

20 1-yl, piperidino, morpholino, thiomorpholino, 1-oxothiomorpholino, 1,1-

dioxothiomorpholino, piperazin-1-yl, 4-C₁₋₄alkylpiperazin-1-yl, C₁₋₄alkylthio, C₁₋₄alkylsulphanyl, C₁₋₄alkylsulphonyl, C₂₋₄alkanoylamino, benzamido, C₁₋₄alkylsulphonamido and phenylsulphonamido, from the substituent -(CH₂)_nY² wherein n is 0-4 and Y² is selected from carboxy, carbamoyl, C₁₋₄alkoxycarbonyl, N-C₁₋₄alkylcarbamoyl, N,N-di-C₁₋

25 4alkylcarbamoyl, pyrrolidin-1-ylcarbonyl, piperidinocarbonyl, morpholinocarbonyl, thiomorpholinocarbonyl,

1-oxothiomorpholinocarbonyl, 1,1-dioxothiomorpholinocarbonyl, piperazin-1-ylcarbonyl, 4-C₁₋₄alkylpiperazin-1-ylcarbonyl, C₁₋₄alkylsulphonamidocarbonyl,

phenylsulphonamidocarbonyl and benzylsulphonamidocarbonyl, from a substituent of the

30 formula -X³-L²-Y² wherein X³ is a group of the formula CON(R³), CON(L²-Y²), C(R³)₂O, O, N(R³) or N(L²-Y²), L² is C₁₋₄alkylene, Y² has any of the meanings defined immediately hereinbefore and each R³ is independently hydrogen or C₁₋₄alkyl, and from a substituent of the formula -X³-L³-Y¹ wherein X³ is a group of the formula CON(R³), CON(L³-Y¹), C(R³)₂O,



AMENDED SHEET

O, N(R⁵) or N(L³-Y¹), L³ is C₂₋₄alkylene, Y¹ has any of the meanings defined immediately hereinbefore and each R⁵ is independently hydrogen or C₁₋₄alkyl, and wherein any heterocyclic group in a substituent of B optionally bears 1 or 2 substituents selected from carboxy, carbamoyl, C₁₋₄alkyl, C₁₋₄alkoxycarbonyl, N-C₁₋₄alkylcarbamoyl and

5 N,N-di-C₁₋₄alkylcarbamoyl, and wherein any phenyl group in a substituent of B optionally bears 1 or 2 substituents selected from halo, trifluoromethyl, cyano, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy, C₂₋₄alkenyloxy and C₂₋₄alkynyloxy;

T¹ is CH or N;

10 T¹ is CH or N;

wherein the groups T¹(R²)-L¹-T²(R³) form a piperazine ring with the proviso that at least one of T¹ and T² is N and wherein the piperazine ring formed by T¹, T², L¹, R² and R³ is optionally substituted by one or two substituents selected from hydroxy, oxo, carboxy and C₁₋₄alkoxycarbonyl; or one of the following:

-(CH₂)_n-R, -(CH₂)_n-NRR¹, -CO-R, -CO-NRR¹, -(CH₂)_n-CO-R and -(CH₂)_n-CO-NRR¹;

wherein n is 0, 1 or 2, preferably n is 1 or 2;

R and R¹ are independently selected from hydrogen, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, hydroxyC₁₋₄alkyl, carboxyC₁₋₄alkyl and C₁₋₄alkoxycarbonylC₁₋₄alkyl or where possible R and R¹ may together form a 5- or 6-membered optionally substituted saturated or partially unsaturated (preferably saturated) heterocyclic ring which may include in addition to the nitrogen to which R and R¹ are attached 1 or 2 additional heteroatoms selected from nitrogen, oxygen and sulphur

X¹ is SO, SO₂, C(R⁴)₂ or CO when T¹ is CH or N; or in addition X¹ is O or S when T¹ is CH; and wherein each R⁴ is independently hydrogen or C₁₋₄alkyl;

L¹ is C₁₋₄alkylene or C₁₋₄alkylenecarbonyl;

R² is hydrogen or C₁₋₄alkyl;

R³ is hydrogen or C₁₋₄alkyl;

or R² and R³ are joined to form a C₁₋₄alkylene or -CH₂CO- group; wherein the ring formed by

T¹, R², R³, T² and L¹ is optionally substituted; with the proviso that when T¹ and T² are both N,

L¹ is not methylene and R² and R³ together are not methylene;



AMENDED SHEET

- X^2 is $S(O)_y$, wherein y is one or two, $C(R^3)_2$ or CO ; and each R^3 is hydrogen or C_{1-4} alkyl;
 Q is phenyl, naphthyl, phenyl C_{1-4} alkyl, phenyl C_{2-4} alkenyl, phenyl C_{2-4} alkynyl or a heterocyclic moiety containing up to 4 heteroatoms selected from nitrogen, oxygen and sulphur and Q is optionally substituted by one, two or three substituents selected from halo, trifluoromethyl, trifluoromethoxy, cyano, hydroxy, amino, nitro, trifluoromethylsulphonyl, carboxy, carbamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{2-4} alkenyloxy, C_{2-4} alkynyloxy, C_{1-4} alkylthio, C_{1-4} alkylsulphinyl, C_{1-4} alkylsulphonyl, C_{1-4} alkylamino, di- C_{1-4} alkylamino, C_{1-4} alkoxycarbonyl, N - C_{1-4} alkylcarbamoyl, N,N -di- C_{1-4} alkylcarbamoyl, C_{2-4} alkanoyl, C_{2-4} alkanoylamino, hydroxy C_{1-4} alkyl, C_{1-4} alkoxy C_{1-4} alkyl, carboxy C_{1-4} alkyl, C_{1-4} alkoxycarbonyl C_{1-4} alkyl, carbamoyl C_{1-4} alkyl, N - C_{1-4} alkylcarbamoyl C_{1-4} alkyl, N,N -di- C_{1-4} alkylcarbamoyl C_{1-4} alkyl, phenyl, heteroaryl, phenoxy, phenylthio, phenylsulphinyl, phenylsulphonyl, benzyl, benzoyl, heteroaryloxy, heteroarylthio, heteroarylsulphinyl and heteroarylsulphonyl, and wherein said heteroaryl substituent or the heteroaryl group in a heteroaryl-containing substituent is a 5- or 6-membered monocyclic heteroaryl ring containing up to 3 heteroatoms selected from nitrogen, oxygen and sulphur, and wherein said phenyl, heteroaryl, phenoxy, phenylthio, phenylsulphinyl, phenylsulphonyl, heteroaryloxy, heteroarylthio, heteroarylsulphinyl, heteroarylsulphonyl, benzyl or benzoyl substituent optionally bears 1, 2 or 3 substituents selected from halo, trifluoromethyl, cyano, hydroxy, amino, nitro, carboxy, carbamoyl, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkylamino, di- C_{1-4} alkylamino, C_{1-4} alkoxycarbonyl, N - C_{1-4} alkylcarbamoyl, N,N -di- C_{1-4} alkylcarbamoyl and C_{2-4} alkanoylamino; and pharmaceutically acceptable salts thereof.

2. A compound of formula (I) according to claim 1 wherein A is a pyridyl, pyrimidinyl or pyridazinyl ring.

3. A compound of formula (I) according to claim 2 wherein A is 4-pyrimidinyl or 4-pyridyl.

4. A compound of formula (I) according to any one of claims 1 to 3 wherein B is paraphenylene.



5. A compound of formula (I) according to any one of claims 1 to 4 wherein the ring formed by T¹, R², R³, T² and L is 1,4-piperazinediyl.
- 5 6. A compound of formula (I) according to any one of claims 1 to 5 wherein X¹ is CO.
7. A compound of formula (I) according to any one of claims 1 to 6 wherein X² is SO₂.
8. A compound of formula (I), as defined in claim 1, wherein
- 10 A is pyridyl, pyrimidinyl, or pyridazinyl;
B is para-phenylene;
X¹ is CO, SO₂ or CH₂;
T¹ and T² are both N;
L¹ is ethylene or propylene;
- 15 R² and R³ are joined to form an ethylene or propylene or methylenecarbonyl group;
X² is SO₂;
Q is styryl or naphthyl optionally substituted by fluoro, chloro or bromo or is phenyl optionally substituted by fluorophenyl, chlorophenyl, or bromophenyl;
and pharmaceutically-acceptable salts thereof.
- 20 9. A compound of formula (I) selected from:
1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyrimidinyl)benzoyl]piperazine;
1-(6-chloronaphth-2-ylsulphonyl)-4-[4-(4-pyridyl)benzoyl]piperazine;
1-(6-bromonaphth-2-ylsulphonyl)-4-[4-(4-pyridazinyl)benzoyl]piperazine;
- 25 and pharmaceutically-acceptable salts thereof.
10. A compound of formula (I) according to any one of claims 1 to 9 for use in medical therapy.
- 30 11. A pharmaceutical formulation comprising a compound of formula (I) according to any one of claims 1 to 9 and a pharmaceutically-acceptable diluent or carrier.



AMENDED SHEET

12. Use of a compound of formula (I) according to any one of claims 1 to 9 in the preparation of a medicament for use in producing a Factor Xa inhibiting effect.

5 13. A method of preventing or treating a Factor Xa mediated disease or medical condition comprising administering to a patient a pharmaceutically effective amount of a compound of formula (I), as defined in any one of claims 1 to 9.

14. A process for preparing a compound of formula (I), are defined in claim 1,
10 comprising:

(a) for the production of those compounds of the formula (I) wherein T' is N and X' is CO, the reaction, conveniently in the presence of a suitable base, of an amine of formula (II)

$$15 \quad \text{HN}(\mathbf{R}^2) \cdot \mathbf{L}^1 \cdot \mathbf{T}^2(\mathbf{R}^3) \cdot \mathbf{X}^3 \cdot \mathbf{Q} \quad (\text{II})$$

with an acid of the formula (III)

A-B-COOH (III)

20 or a reactive derivative thereof;

(b) for the production of those compounds of the formula (I) wherein T' is CH and X' is O by the reaction, conveniently in the presence of a suitable coupling agent, of a compound of the formula (IV):

$$Z-CH(R^1)-L^1-T^1(R^1)-X^1-O \quad (IV)$$


wherein Z is a displaceable group, with a phenolic compound of the formula (V):

30 $A \cdot B \cdot OH$ (V);




AMENDED SHEET


5



10



(f) for the production of compounds wherein T² is N, the reaction of a compound of the formula (VIII):



(g) for the production of compounds wherein T¹ is N and X¹ is SO or SO₂, the reaction of a compound of the formula (II) as defined above with a compound of the formula (X):



AMENDED SHEET

wherein x is one or two and Z is a displaceable group;

- (h) for production of compounds of formula (I) by coupling T² to Q and thus preparing
5 the -T²-X²-Q moiety, methods analogous to those described in process variants (a), (c) and (g)
for preparing the B-X¹-T¹ moiety may be employed;
- (i) for the production of compounds of formula (I) wherein X¹ is a group of the
formula SO, SO₂, wherein B bears a C₁₋₄alkylsulphinyl, C₁₋₄alkylsulphonyl,
10 1-oxothiomorpholino or 1,1-dioxothiomorpholino group, wherein X² is a group of the formula
SO or SO₂ wherein Q bears a C₁₋₄alkylsulphinyl, C₁₋₄alkylsulphonyl, phenylsulphinyl,
phenylsulphonyl, heteroarylsulphinyl or heteroarylsulphonyl group, the oxidation of the
corresponding compound of the formula (I) which contains X¹ as a thio group.

DATED: 22 January, 2001

PHILLIPS ORMONDE & FITZPATRICK

Attorneys for:

ZENECA LIMITED



AMENDED SHEET