

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Vaccine composition

(51)⁶ International Patent Classification(s)
C12N 015/74 A61K 039/118
C12N 001/21 A61K 039/116
C07K 014/195 A61K 039/095
A61K 039/295 C12N 015/67
A61K 039/102 C12N 015/31
A61K 039/104 A61K 039/02
A61K 039/09

(21) Application No: **2003271337**

(22) Application Date: **2003.12.17**

(43) Publication Date : **2004.01.29**

(43) Publication Journal Date : **2004.01.29**

(62) Divisional of:
68336/00

(71) Applicant(s)
SmithKline Beecham Biologicals S.A.

(72) Inventor(s)
Voet, Pierre; Thonnard, Joelle; Thiry, Georges; Poolman,
Jan ; Lobet, Yves; Feron, Christiane; Dequesne, Guy; Denoel,
Philippe; Dalemans, Wilfried L. J.; Berthet, Francois-Xavier Jacques7

(74) Agent/Attorney
Phillips Ormonde & Fitzpatrick, 367 Collins Street, Melbourne, VIC, 3000

(56) Related Art
WO 9957280
WO 0022430
INFECT IMMUN (2003) 71(2):647-55

ABSTRACT

The present invention relates to an immuno-protective and non-toxic Gram-negative bleb vaccine suitable for paediatric use. Examples of the Gram-negative strains from which the blebs are made are *N. meningitidis*, *M. catarrhalis* and *H. influenzae*. The blebs of the invention are improved by one or more genetic changes to the chromosome of the bacterium, including up-regulation of protective antigens, down-regulation of immunodominant non-protective antigens, and detoxification of the Lipid A moiety of LPS.

10

15

20

25

30

AUSTRALIA

Patents Act

**COMPLETE SPECIFICATION
(ORIGINAL)**

Class Int. Class

Application Number:
Lodged:

Complete Specification Lodged:
Accepted:
Published:

Priority

Related Art:

Name of Applicant:

SmithKline Beecham Biologicals S.A.

Actual Inventor(s):

Francois-Xavier Jacques Berthet, Wilfried L. J. Dalemans, Philippe Denoel, Guy Dequesne, Christiane Feron, Yves Lobet, Jan Poolman, Georges Thiry, Joelle Thonnard, Pierre Voet

Address for Service and Correspondence:

PHILLIPS ORMONDE & FITZPATRICK
Patent and Trade Mark Attorneys
367 Collins Street
Melbourne 3000 AUSTRALIA

Invention Title:

VACCINE COMPOSITION

Our Ref : 709569
POF Code: 456838/257598

The following statement is a full description of this invention, including the best method of performing it known to applicant(s):

- 1 -

00000

VACCINE COMPOSITION

The present application is a divisional application of Australian patent application number 68336/00 the entire content of which is incorporated herein by
5 reference.

FIELD OF THE INVENTION

The present invention relates to the field of Gram-negative bacterial vaccine compositions, their manufacture, and the use of such compositions in medicine. More
10 particularly it relates to the field of novel outer-membrane vesicle (or bleb) vaccines, and advantageous methods of rendering these vaccines more effective and safer.

BACKGROUND OF THE INVENTION

Gram-negative bacteria are separated from the external medium by two
15 successive layers of membrane structures. These structures, referred to as the cytoplasmic membrane and the outer membrane (OM), differ both structurally and functionally. The outer membrane plays an important role in the interaction of pathogenic bacteria with their respective hosts. Consequently, the surface exposed bacterial molecules represent important targets for the host immune response, making
20 outer-membrane components attractive candidates in providing vaccine, diagnostic and therapeutics reagents.

Whole cell bacterial vaccines (killed or attenuated) have the advantage of supplying multiple antigens in their natural micro-environment. Drawbacks around this approach are the side effects induced by bacterial components such as endotoxin
25 and peptidoglycan fragments. On the other hand, acellular subunit vaccines containing purified components from the outer membrane may supply only limited protection and may not present the antigens properly to the immune system of the host.

Proteins, phospholipids and lipopolysaccharides are the three major constituents found in the outer-membrane of all Gram-negative bacteria. These
30 molecules are distributed asymmetrically: membrane phospholipids (mostly in the inner leaflet), lipooligosaccharides (exclusively in the outer leaflet) and proteins (inner and outer leaflet lipoproteins, integral or polytopic membrane proteins). For many bacterial pathogens which impact on human health, lipopolysaccharide and outer-

membrane proteins have been shown to be immunogenic and amenable to confer protection against the corresponding disease by way of immunization.

The OM of Gram-negative bacteria is dynamic and, depending on the environmental conditions, can undergo drastic morphological transformations. Among these manifestations, the formation of outer-membrane vesicles or “blebs” has been studied and documented in many Gram-negative bacteria (Zhou, L *et al.* 1998. *FEMS Microbiol. Lett.* 163: 223-228). Among these, a non-exhaustive list of bacterial pathogens reported to produce blebs include: *Bordetella pertussis*, *Borrelia burgdorferi*, *Brucella melitensis*, *Brucella ovis*, *Chlamydia psittaci*, *Chlamydia trachomatis*, *Esherichia coli*, *Haemophilus influenzae*, *Legionella pneumophila*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Pseudomonas aeruginosa* and *Yersinia enterocolitica*. Although the biochemical mechanism responsible for the production of OM blebs is not fully understood, these outer membrane vesicles have been extensively studied as they represent a powerful methodology in order to isolate outer-membrane protein preparations in their native conformation. In that context, the use of outer-membrane preparations is of particular interest to develop vaccines against *Neisseria*, *Moraxella catarrhalis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa* and *Chlamydia*. Moreover, outer membrane blebs combine multiple proteinaceous and non-proteinaceous antigens that are likely to confer extended protection against intra-species variants.

In comparison with the other, more widely used, types of bacterial vaccine (whole cell bacterial and purified subunit vaccines), the inventors will show that outer membrane bleb vaccines (if modified in certain ways) may represent the ideal compromise.

The wide-spread use of bacterial subunit vaccines has been due to the intensive study of bacterial surface proteins that have been found to be useful in vaccine applications [for instance *B. pertussis* pertactin]. These proteins are loosely associated with the bacterial outer membrane and can be purified from culture supernatant or easily extracted from the bacterial cells. However it has also been shown that structural, integral outer membrane proteins are also protective antigens. Examples are PorA for *N. meningitidis* serogroup B; D15 for *H. influenzae*; OMP CD for *M. catarrhalis*; OMP F for *P. Aeruginosa*. Such proteins however have rather

specific structural features, particularly multiple amphipathic β -sheets, which complicates their straightforward use as purified (recombinant) subunit vaccines.

In addition, it has become clear that multiple component vaccines are needed (in terms of bacterial surface proteins and integral membrane proteins) to supply a reasonable level of protection. For instance, in the case of *B. pertussis* subunit vaccines multicomponent vaccines are superior to mono or bicomponent products.

In order to incorporate integral outer-membrane proteins into such a subunit product, native (or near-native) conformational folding of the proteins must be present in the product in order to have a useful immunological effect. The use of excreted outer membrane vesicles or blebs may be an elegant solution to the problem of including protective integral membrane proteins into a subunit vaccine whilst still ensuring that they fold properly.

N. meningitidis serogroup B (menB) excretes outer membrane blebs in sufficient quantities to allow their manufacture on an industrial scale. Such multicomponent outer-membrane protein vaccines from naturally-occurring menB strains have been found to be efficacious in protecting teenagers from menB disease and have become registered in Latin America. An alternative method of preparing outer-membrane vesicles is via the process of detergent extraction of the bacterial cells (EP 11243).

Examples of bacterial species from which bleb vaccines can be made are the following.

Neisseria meningitidis:

Neisseria meningitidis (meningococcus) is a Gram-negative bacterium frequently isolated from the human upper respiratory tract. It occasionally causes invasive bacterial diseases such as bacteremia and meningitis. The incidence of meningococcal disease shows geographical seasonal and annual differences (Schwartz, B., Moore, P.S., Broome, C.V.; Clin. Microbiol. Rev. 2 (Supplement), S18-S24, 1989). Most disease in temperate countries is due to strains of serogroup B and varies in incidence from 1-10/100,000/year total population sometimes reaching higher values (Kaczmarek, E.B. (1997), Commun. Dis. Rep. Rev. 7: R55-9, 1995; Scholten, R.J.P.M., Bijlmer, H.A., Poolman, J.T. et al. Clin. Infect. Dis. 16: 237-246, 1993; Cruz, C., Pavez, G., Aguilar, E., et al. Epidemiol. Infect. 105: 119-126, 1990).

Age-specific incidences in the two high risk-groups, infants and teenagers, reach higher levels.

Epidemics dominated by serogroup A meningococci occur, mostly in central Africa, sometimes reaching levels up to 1000/100,000/year (Schwartz, B., Moore, P.S., Broome, C.V. Clin. Microbiol. Rev. 2 (Supplement), S18-S24, 1989). Nearly all cases of meningococcal disease as a whole are caused by serogroup A, B, C, W-135 and Y meningococci. A tetravalent A, C, W-135, Y capsular polysaccharide vaccine is available (Armand, J., Arminjon, F., Mynard, M.C., Lafaix, C., J. Biol. Stand. 10: 335-339, 1982).

The polysaccharide vaccines are currently being improved by way of chemically conjugating them to carrier proteins (Lieberman, J.M., Chiu, S.S., Wong, V.K., et al. JAMA 275 : 1499-1503, 1996). A serogroup B vaccine is not available, since the B capsular polysaccharide is non-immunogenic, most likely because it shares structural similarity to host components (Wyle, F.A., Artenstein, M.S., Brandt, M.L. et al. J. Infect. Dis. 126: 514-522, 1972; Finne, J.M., Leinonen, M., Mäkelä, P.M. Lancet ii.: 355-357, 1983).

For many years efforts have been focused on developing meningococcal outer membrane based vaccines (de Moraes, J.C., Perkins, B., Camargo, M.C. et al. Lancet 340: 1074-1078, 1992; Bjune, G., Hoiby, E.A. Gronnesby, J.K. et al. 338: 1093-1096, 1991). Such vaccines have demonstrated efficacies from 57% - 85% in older children (>4 years) and adolescents. Most of these efficacy trials were performed with OMV (outer membrane vesicles, derived by LPS depletion from blebs) vaccines derived from wild-type *N. meningitidis* B strains.

Many bacterial outer membrane components are present in these vaccines, such as PorA, PorB, Rmp, Opc, Opa, FrpB and the contribution of these components to the observed protection still needs further definition. Other bacterial outer membrane components have been defined (using animal or human antibodies) as potentially being relevant to the induction of protective immunity, such as TbpB, NspA (Martin, D., Cadieux, N., Hamel, J., Brodeux, B.R., J. Exp. Med. 185: 1173-1183, 1997; Lissolo, L., Maitre-Wilmotte, C., Dumas, p. et al., Inf. Immun. 63: 884-890, 1995). The mechanism of protective immunity will involve antibody mediated bactericidal activity and opsonophagocytosis.

The frequency of *Neisseria meningitidis* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiple antibiotic resistant strains, and increased exposure due to an increase in social activities (for instance swimming pools or theatres). It is no longer uncommon to isolate *Neisseria meningitidis* strains that are resistant to some or all of the standard antibiotics. This phenomenon has created an unmet medical need and demand for new anti-microbial agents, vaccines, drug screening methods, and diagnostic tests for this organism.

Moraxella catarrhalis

Moraxella catarrhalis (also named *Branhamella catarrhalis*) is a Gram-negative bacterium frequently isolated from the human upper respiratory tract. It is responsible for several pathologies, the main ones being otitis media in infants and children, and pneumonia the elderly. It is also responsible for sinusitis, nosocomial infections and, less frequently, for invasive diseases.

Bactericidal antibodies have been identified in most adults tested (Chapman, AJ et al. (1985) J. Infect.Dis. 151:878). Strains of *M. catarrhalis* present variations in their capacity to resist serum bactericidal activity: in general, isolates from diseased individuals are more resistant than those who are simply colonized (Hol, C et al. (1993) Lancet 341:1281, Jordan, KL et al. (1990) Am.J.Med. 88 (suppl. 5A):28S). Serum resistance could therefore be considered as a virulence factor of the bacteria. An opsonizing activity has been observed in the sera of children recovering from otitis media.

The antigens targeted by these different immune responses in humans have not been identified, with the exception of OMP B1, a 84 kDa protein, the expression of which is regulated by iron, and that is recognized by the sera of patients with pneumonia (Sethi, S, et al. (1995) Infect.Immun. 63:1516), and of UspA1 and UspA2 (Chen D. et al.(1999), Infect.Immun. 67:1310).

A few other membrane proteins present on the surface of *M. catarrhalis* have been characterized using biochemical methods for their potential implication in the induction of a protective immunity (for review, see Murphy, TF (1996) Microbiol.Rev. 60:267). In a mouse pneumonia model, the presence of antibodies raised against some of them (UspA, CopB) favors a faster clearance of the pulmonary infection. Another polypeptide (OMP CD) is highly conserved among *M. catarrhalis* strains, and presents homologies with a

porin of *Pseudomonas aeruginosa*, which has been demonstrated to be efficacious against this bacterium in animal models.

M. catarrhalis produces outer membrane vesicles (Blebs). These Blebs have been isolated or extracted by using different methods (Murphy T.F., Loeb M.R. 1989. Microb. Pathog. 6: 159-174; Unhanand M., Maciver, I., Ramillo O., Arencibia-Mireles O., Argyle J.C., McCracken G.H. Jr., Hansen E.J. 1992. J. Infect. Dis. 165:644-650). The protective capacity of such Bleb preparations has been tested in a murine model for pulmonary clearance of *M. catarrhalis*. It has been shown that active immunization with Bleb vaccine or passive transfer of anti-Blebs antibody induces significant protection in this model (Maciver I., Unhanand M., McCracken G.H. Jr., Hansen, E.J. 1993. J. Infect. Dis. 168: 469-472).

Haemophilus influenzae

Haemophilus influenzae is a non-motile Gram-negative bacterium. Man is its only natural host. *H. influenzae* isolates are usually classified according to their polysaccharide capsule. Six different capsular types designated 'a' through 'f' have been identified. Isolates that fail to agglutinate with antisera raised against one of these six serotypes are classified as nontypeable, and do not express a capsule.

H. influenzae type b (Hib) is clearly different from the other types in that it is a major cause of bacterial meningitis and systemic diseases. Nontypeable strains of *H. influenzae* (NTHi) are only occasionally isolated from the blood of patients with systemic disease. NTHi is a common cause of pneumonia, exacerbation of chronic bronchitis, sinusitis and otitis media. NTHi strains demonstrate a large variability as identified by clonal analysis, whilst Hib strains as a whole are more homogeneous.

Various proteins of *H. influenzae* have been shown to be involved in pathogenesis or have been shown to confer protection upon vaccination in animal models.

Adherence of NTHi to human nasopharyngeal epithelial cells has been reported (Read RC. et al. 1991. J. Infect. Dis. 163:549). Apart from fimbriae and pili (Brinton CC. et al. 1989. Pediatr. Infect. Dis. J. 8:S54; Kar S. et al. 1990. Infect. Immun. 58:903; Gildorf JR. et al. 1992. Infect. Immun. 60:374; St. Geme JW et al. 1991. Infect. Immun. 59:3366; St. Geme JW et al. 1993. Infect. Immun. 61: 2233), many adhesins have been identified in NTHi. Among them, two surface exposed high-molecular-weight proteins

designated HMW1 and HMW2 have been shown to mediate adhesion of NTHi to epithelial cells (St. Geme JW. et al. 1993. Proc. Natl. Acad. Sci. USA 90:2875). Another family of high-molecular-weight proteins has been identified in NTHi strains that lack proteins belonging to HMW1/HMW2 family. The NTHi 115-kDa Hia protein (Barenkamp SJ., St Geme S.W. 1996. Mol. Microbiol. In press) is highly similar to the Hsf adhesin expressed by *H. influenzae* type b strains (St. Geme JW. et al. 1996. J. Bact. 178:6281). Another protein, the Hap protein shows similarity to IgA1 serine proteases and has been shown to be involved in both adhesion and cell entry (St. Geme JW. et al. 1994. Mol. Microbiol. 14:217).

10 Five major outer membrane proteins (OMP) have been identified and numerically numbered. Original studies using *H. influenzae* type b strains showed that antibodies specific for P1 and P2 OMPs protected infant rats from subsequent challenge (Loeb MR. et al. 1987. Infect. Immun. 55:2612; Musson RS. Jr. et al. 1983. J. Clin. Invest. 72:677). P2 was found to be able to induce bactericidal and opsonic antibodies, which are directed against the variable regions present within surface exposed loop structures of this integral OMP (Haase EM. et al. 1994 Infect. Immun. 62:3712; Troelstra A. et al. 1994 Infect. Immun. 62:779). The lipoprotein P4 also may induce bactericidal antibodies (Green BA. et al. 1991. Infect.Immun.59:3191).

20 OMP P6 is a conserved peptidoglycan associated lipoprotein making up 1-5 % of the outer membrane (Nelson MB. et al. 1991. Infect. Immun. 59:2658). Later a lipoprotein of about the same molecular weight was recognized called PCP (P6 cross-reactive protein) (Deich RM. et al. 1990. Infect. Immun. 58:3388). A mixture of the conserved lipoproteins P4, P6 and PCP did not reveal protection as measured in a chinchilla otitis-media model (Green BA. et al. 1993. Infect.immun. 61:1950). P6 alone appears to induce protection in the chinchilla model (Demaria TF. et al. 1996. Infect. Immun. 64:5187).

30 A fimbriin protein (Miyamoto N., Bakaletz, LO. 1996. Microb. Pathog. 21:343) has also been described with homology to OMP P5, which itself has sequence homology to the integral *Escherichia coli* OmpA (Miyamoto N., Bakaletz, LO. 1996. Microb. Pathog. 21:343; Munson RS. Jr. et al. 1993. Infect. Immun. 61:1017). NTHi seem to adhere to mucus by way of fimbriae.

In line with the observations made with gonococci and meningococci, NTHi expresses a dual human transferrin receptor composed of TbpA and TbpB when grown under iron limitation. Anti-TbpB antibody protected infant rats (Loosmore SM. et al. 1996. Mol. Microbiol. 19:575). Hemoglobin / haptoglobin receptor also have been described for NTHi (Maciver I. et al. 1996. Infect. Immun. 64:3703). A receptor for Haem:Hemopexin has also been identified (Cope LD. et al. 1994. Mol.Microbiol. 13:868). A lactoferrin receptor is also present amongst NTHi, but is not yet characterized (Schryvers AB. et al. 1989. J. Med. Microbiol. 29:121). A protein similar to neisserial FrpB-protein has not been described amongst NTHi.

10 An 80kDa OMP, the D15 surface antigen, provides protection against NTHi in a mouse challenge model. (Flack FS. et al. 1995. Gene 156:97). A 42kDa outer membrane lipoprotein, LPD is conserved amongst *Haemophilus influenzae* and induces bactericidal antibodies (Akkoyunlu M. et al. 1996. Infect. Immun. 64:4586). A minor 98kDa OMP (Kimura A. et al. 1985. Infect. Immun. 47:253), was found to be a protective antigen, this OMP may very well be one of the Fe-limitation inducible OMPs or high molecular weight adhesins that have been characterized thereafter. *H. Influenzae* produces IgA1-protease activity (Mulks MH., Shoberg RJ. 1994. Meth. Enzymol. 235:543). IgA1-proteases of NTHi have a high degree of antigenic variability (Lomholt H., van Alphen L., Kilian, M. 1993. Infect. Immun. 61:4575).

20 Another OMP of NTHi, OMP26, a 26-kDa protein has been shown to enhance pulmonary clearance in a rat model (Kyd, J.M., Cripps, A.W. 1998. Infect. Immun. 66:2272). The NTHi HtrA protein has also been shown to be a protective antigen. Indeed, this protein protected Chinchilla against otitis media and protected infant rats against *H. influenzae* type b bacteremia (Loosmore S.M. et al. 1998. Infect. Immun. 66:899).

25 Outer membrane vesicles (or blebs) have been isolated from *H. influenzae* (Loeb M.R., Zachary A.L., Smith D.H. 1981. J. Bacteriol. 145:569-604; Stull T.L., Mack K., Haas J.E., Smit J., Smith A.L. 1985. Anal. Biochem. 150: 471-480). The vesicles have been associated with the induction of blood-brain barrier permeability (Wiwelwey B., Hansen E.J., Scheld W.M. 1989 Infect. Immun. 57: 2559-2560), the induction of meningeal inflammation (Mustafa M.M., Ramilo O., Syrogiannopoulos G.A., Olsen K.D., McCracken G.H. Jr., Hansen, E.J. 1989. J. Infect. Dis. 159: 917-922) and to DNA uptake (Concino M.F., Goodgal S.H. 1982 J. Bacteriol. 152: 441-450).

These vesicles are able to bind and be absorbed by the nasal mucosal epithelium (Harada T., Shimuzu T., Nishimoto K., Sakakura Y. 1989. *Acta Otorhinolarygol.* 246: 218-221) showing that adhesins and/or colonization factors could be present in blebs. Immune response to proteins present in outer membrane vesicles has been observed in patients with various *H. influenzae* diseases (Sakakura Y., Harada T., Hamaguchi Y., Jin C.S. 1988. *Acta Otorhinolarygol. Suppl. (Stockh.)* 454: 222-226; Harada T., Sakakura Y., Miyoshi Y. 1986. *Rhinology* 24: 61-66).

Pseudomonas aeruginosa:

10 The genus *Pseudomonas* consists of Gram-negative, polarly flagellated, straight and slightly curved rods that grow aerobically and do not form spores. Because of their limited metabolic requirements, *Pseudomonas spp.* are ubiquitous and are widely distributed in the soil, the air, sewage water and in plants. Numerous species of *Pseudomonas* such as *P. aeruginosa*, *P. pseudomallei*, *P. mallei*, *P.*
15 *maltophilia* and *P. cepacia* have also been shown to be pathogenic for humans. Among this list, *P. aeruginosa* is considered as an important human pathogen since it is associated with opportunistic infection of immuno-compromised host and is responsible for high morbidity in hospitalized patients. Nosocomial infection with *P. aeruginosa* afflicts primarily patients submitted for prolonged treatment and receiving
20 immuno-suppressive agents, corticosteroids, antimetabolites, antibiotics or radiation.

The *Pseudomonas*, and particularly *P. aeruginosa*, produces a variety of toxins (such as hemolysins, fibrinolysins, esterases, coagulases, phospholipases, endo- and exo-toxins) that contribute to the pathogenicity of these bacteria. Moreover, these organisms have high intrinsic resistance to antibiotics due to the presence of multiple
25 drug efflux pumps. This latter characteristic often complicates the outcome of the disease.

Due to the uncontrolled use of antibacterial chemotherapeutics the frequency of nosocomial infection caused by *P. aeruginosa* has increased considerably over the last 30 years. In the US, for example, the economic burden of *P. aeruginosa*
30 nosocomial infection is estimated to 4.5 billion US\$ annually. Therefore, the development of a vaccine for active or passive immunization against *P. aeruginosa* is actively needed (for review see Stanislavsky et al. 1997. *FEMS Microbiol. Lett.* 21: 243-277).

Various cell-associated and secreted antigens of *P. aeruginosa* have been the subject of vaccine development. Among *Pseudomonas* antigens, the mucoid substance, which is an extracellular slime consisting predominantly of alginate, was found to be heterogenous in terms of size and immunogenicity. High molecular mass
5 alginate components (30-300 kDa) appear to contain conserved epitopes while lower molecular mass alginate components (10-30 kDa) possess conserved epitopes in addition to unique epitopes. Among surface-associated proteins, PcrV, which is part of the type III secretion-translocation apparatus, has also been shown to be an interesting target for vaccination (Sawa et al. 1999. Nature Medicine 5:392-398).

10 Surface-exposed antigens including O-antigens (O-specific polysaccharide of LPS) or H-antigens (flagellar antigens) have been used for serotyping due to their highly immunogenic nature. Chemical structures of repeating units of O-specific polysaccharides have been elucidated and these data allowed the identification of 31 chemotypes of *P. aeruginosa*. Conserved epitopes among all serotypes of *P.*
15 *aeruginosa* are located in the core oligosaccharide and the lipid A region of LPS and immunogens containing these epitopes induce cross-protective immunity in mice against different *P. aeruginosa* immunotypes. The outer core of LPS was implicated to be a ligand for binding of *P. aeruginosa* to airway and ocular epithelial cells of animals. However, heterogeneity exists in this outer core region among different
20 serotypes. Epitopes in the inner core are highly conserved and have been demonstrated to be surface-accessible, and not masked by O-specific polysaccharide.

To examine the protective properties of OM proteins, a vaccine containing *P. aeruginosa* OM proteins of molecular masses ranging from 20 to 100 kDa has been used in pre-clinical and clinical trials. This vaccine was efficacious in animal models
25 against *P. aeruginosa* challenge and induced high levels of specific antibodies in human volunteers. Plasma from human volunteers containing anti-*P. aeruginosa* antibodies provided passive protection and helped the recovery of 87% of patients with severe forms of *P. aeruginosa* infection. More recently, a hybrid protein containing parts of the outer membrane proteins OprF (amino acids 190-342) and OprI
30 (amino acids 21-83) from *Pseudomonas aeruginosa* fused to the glutathione-S-transferase was shown to protect mice against a 975-fold 50% lethal dose of *P. aeruginosa* (Knapp et al. 1999. Vaccine. 17:1663-1669).

The present inventors have realised a number of drawbacks associated with the above wild-type bleb vaccines (either naturally occurring or chemically made).

Examples of such problems are the following:

- 5 - the presence of immunodominant but variable proteins on the bleb (PorA; TbpB, Opa [*N. meningitidis* B]; P2, P5 [non-typeable *H. influenzae*]) – such blebs being effective only against a restricted selection of bacterial species. Type-specificity of the bactericidal antibody response may preclude the use of such vaccines in infancy.
- 10 - the presence of unprotective (non relevant) antigens (Rmp, H8, ...) on the bleb – antigens that are decoys for the immune system
- the lack of presence of important molecules which are produced conditionally (for instance iron-regulated outer membrane proteins, IROMP's, in vivo regulated expression mechanisms) – such conditions are hard to control in bleb production in order to optimise the amount of antigen on the surface
- 15 - the low level of expression of protective, (particularly conserved) antigens (NspA, P6)
- the toxicity of the LPS remaining on the surface of the bleb
- 20 - the potential induction of an autoimmune response because of host-identical structures (for example the capsular polysaccharide in *Neisseria meningitidis* serogroup B, the lacto-N-neotetraose in *Neisseria* LPS, saccharide structure within nHi LPS, saccharide structures within Pili).

Such problems may prevent the use of bleb vaccines as human vaccine reagents. This is particularly so for paediatric use (<4 years) where reactogenicity against bleb vaccines is particularly important, and where bleb vaccines (for instance the previously mentioned marketed MenB bleb vaccine) have been shown to be ineffective at immuno-protecting. Accordingly, the present invention provides methods of alleviating the above problems using genetically engineered bacterial strains, which result in improved bleb vaccines. Such methods will be especially useful in the generation of new vaccines against bacterial pathogens such as *Neisseria meningitidis*, *Moraxella catarrhalis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa*, and others.

The bleb vaccines of the invention are designed to focus the immune response on a few protective (preferably conserved) antigens or epitopes - formulated in a multiple component vaccine. Where such antigens are integral OMPs, the outer membrane vesicles of bleb vaccines will ensure their proper folding. This invention provides methods to optimize the OMP and LPS composition of OMV (bleb) vaccines by deleting immunodominant variable as well as non protective OMPs, by creating conserved OMPs by deletion of variable regions, by upregulating expression of protective OMPs, and by eliminating control mechanisms for expression (such as iron restriction) of protective OMPs. In addition the invention provides for the reduction in toxicity of lipid A by modification of the lipid portion or by changing the phosphoryl composition whilst retaining its adjuvant activity or by masking it. Each of these new methods of improvement individually improve the bleb vaccine, however a combination of one or more of these methods work in conjunction so as to produce an optimised engineered bleb vaccine which is immuno-protective and non-toxic - particularly suitable for paediatric use.

The discussion of the background to the invention herein is included to explain the context of the invention. This is not to be taken as an admission that any of the material referred to was published, known or part of the common general knowledge in Australia as at the priority date of any of the claims.

Throughout the description and claims of the specification the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.

SUMMARY OF THE INVENTION

The present invention provides a genetically-engineered bleb preparation from a Gram-negative bacterial strain characterized in that said preparation is obtainable by employing one or more processes selected from the following group:

- a) a process of reducing immunodominant variable or non-protective antigens within the bleb preparation comprising the steps of determining the identity of such antigen, engineering a bacterial strain to produce less or none of said antigen, and making blebs from said strain;
- b) a process of upregulating expression of protective, endogenous (and preferably conserved) OMP antigens within the bleb preparation

-
- comprising the steps of identifying such antigen, engineering a bacterial strain so as to introduce a stronger promoter sequence upstream of a gene encoding said antigen such that said gene is expressed at a level higher than in the non-modified bleb, and making blebs from said strain;
- 5 c) a process of upregulating expression of conditionally-expressed, protective (and preferably conserved) OMP antigens within the bleb preparation comprising the steps of identifying such an antigen, engineering a bacterial strain so as to remove the repressive control mechanisms of its expression (such as iron restriction), and making blebs from said strain;
- 10 d) a process of modifying lipid A portion of bacterial LPS within the bleb preparation, comprising the steps of identifying a gene involved in rendering the lipid A portion of LPS toxic, engineering a bacterial strain so as to reduce or switch off expression of said gene, and making blebs from said strain;
- 15 e) a process of modifying lipid A portion of bacterial LPS within the bleb preparation, comprising the steps of identifying a gene involved in rendering the lipid A portion of LPS less toxic, engineering a bacterial strain so as to introduce a stronger promoter sequence upstream of said gene such that said gene is expressed at a level higher than in the non-
- 20 modified bleb, and making blebs from said strain;
- f) a process of reducing lipid A toxicity within the bleb preparation and increasing the levels of protective antigens, comprising the steps of engineering the chromosome of a bacterial strain to incorporate a gene encoding a Polymyxin A peptide, or a derivative or analogue thereof, fused
- 25 to a protective antigen, and making blebs from said strain;
- g) a process of creating conserved OMP antigens on the bleb preparation comprising the steps of identifying such antigen, engineering a bacterial strain so as to delete variable regions of a gene encoding said antigen, and making blebs from said strain;
- 30 h) a process of reducing expression within the bleb preparation of an antigen which shares a structural similarity with a human structure and may be capable of inducing an auto-immune response in humans (such as the capsular polysaccharide of *N. meningitidis* B), comprising the steps of

identifying a gene involved in the biosynthesis of the antigen, engineering a bacterial strain so as to reduce or switch off expression of said gene, and making blebs from said strain; or

5 i) a process of upregulating expression of protective, endogenous (and preferably conserved) OMP antigens within the bleb preparation comprising the steps of identifying such antigen, engineering a bacterial strain so as to introduce into the chromosome one or more further copies of a gene encoding said antigen controlled by a heterologous, stronger promoter sequence, and making blebs from said strain.

10 According to one aspect of the present invention, there is provided a modified *Neisseria meningitidis* strain characterized in that said strain is obtainable by employing the following processes:

15 d) a process of detoxifying the lipid A portion of LPS within the strain, comprising the steps of identifying an *msbB* gene involved in rendering the lipid A portion of LPS toxic, and engineering the strain so as to reduce or switch off expression of said gene; and

h) a process of engineering the strain such that it is free of capsular polysaccharide.

20 Further aspects of the invention include, preferential processes for obtaining the above bleb preparation, including optimal positioning of strong promoters for the upregulation of expression of antigens within blebs, preferential antigens for upregulation and downregulation for various bacterial strains in order to obtain bleb preparations particularly suitable for vaccine use. Preferential formulations comprising the blebs of the invention are also provided which are particularly suitable
25 for global vaccines against certain disease states. Vectors for producing the blebs of the invention, and modified bacterial strains from which the blebs of the invention are produced are still further aspects of the invention.

The present invention provides for the first time a bleb vaccine which is immuno-protective and non-toxic when used with children under 4 years of age.

30 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: Reactivity of the 735 mAb on different colonies.

Figure 2: Reactivities of specific monoclonal antibodies by whole cell Elisa.

Figure 3: Schematic representation of the pCMK vectors used to deliver genes,
5 operons and/or expression cassettes in the genome of *Neisseria meningitidis*.

Figure 4: Analysis of PorA expression in total protein extracts of recombinant *N. meningitidis* serogroupB (H44/76 derivatives). Total proteins were recovered from cps- (lanes 3 and 4), cps- porA::pCMK+ (lanes 2 and 5) and cps- porA::nspA (lanes 1 and 6) recombinant *N. meningitidis* serogroupB strains and were analyzed under SDS-PAGE conditions in a 12% polyacrylamide gel. Gels were stained with Coomassie blue (lanes 1 to 3) or transferred to a nitrocellulose membrane and immuno-stained with an anti-PorA monoclonal antibody.
10

Figure 5: Analysis of NspA expression in protein extracts of recombinant *N. meningitidis* serogroupB strains (H44/76 derivatives). Proteins were extracted from whole bacteria (lanes 1 to 3) or outer-membrane blebs preparations (lanes 4 to 6) separated by SDS-PAGE on a 12% acrylamide gel and analyzed by immuno-blotting using an anti-NspA polyclonal serum. Samples corresponding to cps- (lanes 1 and 6),
20 cps- porA::pCMK+ (lanes 3 and 4) and cps- porA::nspA (lanes 2 and 5) were analyzed. Two forms of NspA were detected: a mature form (18kDa) co-migrating with the recombinant purified NspA, and a shorter form (15kDa).

Figure 6: Analysis of D15/omp85 expression in protein extracts of recombinant *N. meningitidis* serogroupB strains (H44/76 derivatives). Proteins were extracted from outer-membrane blebs preparations and were separated by SDS-PAGE on a 12% acrylamide gel and analyzed by immuno-blotting using an anti-omp85 polyclonal serum. Samples corresponding to cps- (lane 2), and cps-, PorA+, pCMK+Omp85/D15 (lane 1) recombinant *N. meningitidis* serogroupB strains were analyzed.
25

30

Figure 7: General strategy for modulating gene expression by promoter delivery (RS stands for restriction site).

Figure 8: Analysis of outer-membrane blebs produced by recombinant *N. meningitidis* serogroup B *cps*- strains (H44/76 derivatives). Proteins were extracted from outer-membrane bleb preparations and were separated by SDS-PAGE under reducing conditions on a 4-20% gradient polyacrylamide gel. The gel was stained with Coomassie brilliant blue R250. Lanes 2, 4, 6 corresponded to 5µg of total proteins whereas lanes 3, 5 and 7 were loaded with 10µg proteins.

Figure 9: Construction of a promoter replacement plasmid used to up-regulate the expression/production of Omp85/D15 in *Neisseria meningitidis* H44/76.

Figure 10: Analysis of OMP85 expression in total protein extracts of recombinant NmB (H44/76 derivatives). Gels were stained with Coomassie blue (A) or transferred to nitrocellulose membrane and immuno-stained with rabbit anti-OMP85 (*N.gono*) monoclonal antibody (B).

Figure 11: Analysis of OMP85 expression in OMV preparations from recombinant NmB (H44/76 derivatives). Gels were stained with Coomassie blue (A) or transferred to nitrocellulose membrane and immuno-stained with rabbit anti-OMP85 polyclonal antibody (B).

Figure 12: Schematic representation of the recombinant PCR strategy used to delete the *lacO* in the chimeric *porA/lacO* promoter.

Figure 13: Analysis of Hsf expression in total protein extracts of recombinant *N. meningitidis* serogroup B (H44/76 derivatives). Total proteins were recovered from Cps-PorA+(lanes 1), and Cps-PorA+/Hsf (lanes 2) recombinant *N. meningitidis* serogroup B strains and were analyzed under SDS-PAGE conditions in a 12% polyacrylamide gel. Gels were stained with Coomassie blue .

Figure 14: Analysis of GFP expression in total protein extracts of recombinant *N. meningitidis* (H44/76 derivative). Total protein were recovered from Cps-, PorA+ (lane1), Cps-, PorA- GFP+ (lane2 & 3) recombinant strains. Proteins were separated by PAGE-SDS in a 12% polyacrylamide gel and then stained with Coomassie blue.

Figure 15: Illustration of the pattern of major proteins on the surface of various recombinant bleb preparations as analysed by SDS-PAGE (Coomassie staining).

5 **Figure 16:** Specific anti-Hsf response for various bleb and recombinant bleb preparations using purified recombinant Hsf protein.

Figure 17: Analysis of NspA expression in total protein extracts of recombinant *NmB* (serogroup B derivatives). Gels were stained with Coomassie blue (A) or transferred to nitrocellulose membrane and immuno-stained with mouse anti-PorA monoclonal antibody (B) or mouse anti-NspA polyclonal antibody (C).

10

DESCRIPTION OF THE INVENTION

The present invention relates to a general set of tools and methods capable of being used for manufacturing improved, genetically engineered blebs from Gram-negative bacterial strains. The invention includes methods used to make recombinant blebs more immunogenic, less toxic and safer for their use in a human and/or animal vaccine. Moreover, the present invention also describes specific methods necessary for constructing, producing, obtaining and using recombinant, engineered blebs from various Gram-negative bacteria, for vaccine, therapeutic and/or diagnostic purposes. By the methods of the invention, the biochemical composition of bacterial blebs can be manipulated by acting upon/altering the expression of bacterial genes encoding products present in or associated with bacterial outer-membrane blebs (outer membrane proteins or OMPs). The production of blebs using a method of genetic modification to increase, decrease or render conditional the expression of one or more genes encoding outer-membrane components are also included in the scope of this invention.

15

20

25

For clarity, the term “expression cassette” will refer herein to all the genetic elements necessary to express a gene or an operon and to produce and target the corresponding protein(s) of interest to outer-membrane blebs, derived from a given bacterial host. A non-exhaustive list of these features includes control elements (transcriptional and/or translational), protein coding regions and targeting signals, with appropriate spacing between them. Reference to the insertion of promoter

30

sequences means, for the purposes of this invention, the insertion of a sequence with at least a promoter function, and preferably one or more other genetic regulatory elements comprised within an expression cassette. Moreover, the term “integrative cassette” will refer herein to all the genetic elements required to integrate a DNA segment in given bacterial host. A non-exhaustive list of these features includes a delivery vehicle (or vector), with recombinogenic regions, and selectable and counter selectable markers.

Again for the purpose of clarity, the terms ‘engineering a bacterial strain to produce less of said antigen’ refers to any means to reduce the expression of an antigen of interest, relative to that of the non-modified (i.e., naturally occurring) bleb such that expression is at least 10% lower than that of the non-modified bleb. Preferably it is at least 50% lower. “Stronger promoter sequence” refers to a regulatory control element that increases transcription for a gene encoding antigen of interest. “Upregulating expression” refers to any means to enhance the expression of an antigen of interest, relative to that of the non-modified (i.e., naturally occurring) bleb. It is understood that the amount of ‘upregulation’ will vary depending on the particular antigen of interest but will not exceed an amount that will disrupt the membrane integrity of the bleb. Upregulation of an antigen refers to expression that is at least 10% higher than that of the non-modified bleb. Preferably it is at least 50% higher. More preferably it is at least 100% (2 fold) higher.

Aspects of the invention relate to individual methods for making improved engineered blebs, to a combination of such methods, and to the bleb compositions made as a result of these methods. Another aspect of the invention relates to the genetic tools used in order to genetically modify a chosen bacterial strain in order to extract improved engineered blebs from said strain.

The engineering steps of the processes of the invention can be carried out in a variety of ways known to the skilled person. For instance, sequences (e.g. promoters or open reading frames) can be inserted, and promoters/genes can be disrupted by the technique of transposon insertion. For instance, for upregulating a gene’s expression, a strong promoter could be inserted via a transposon up to 2 kb upstream of the gene’s initiation codon (more preferably 200-600 bp upstream, most preferably approximately 400 bp upstream). Point mutation or deletion may also be used (particularly for down-regulating expression of a gene).

Such methods, however, may be quite unstable or uncertain, and therefore it is preferred that the engineering step [particularly for processes a), b), c), d), e), h) and i) as described below] is performed via a homologous recombination event. Preferably, the event takes place between a sequence (a recombinogenic region) of at least 30 nucleotides on the bacterial chromosome, and a sequence (a second recombinogenic region) of at least 30 nucleotides on a vector transformed within the strain. Preferably the regions are 40-1000 nucleotides, more preferably 100-800 nucleotides, most preferably 500 nucleotides). These recombinogenic regions should be sufficiently similar that they are capable of hybridising to one another under highly stringent conditions (as defined later).

Recombination events may take place using a single recombinogenic region on chromosome and vector, or via a double cross-over event (with 2 regions on chromosome and vector). In order to perform a single recombination event, the vector should be a circular DNA molecule. In order to perform a double recombination event, the vector could be a circular or linear DNA molecule (see Figure 7). It is preferable that a double recombination event is employed and that the vector used is linear, as the modified bacterium so produced will be more stable in terms of reversion events. Preferably the two recombinogenic regions on the chromosome (and on the vector) are of similar (most preferably the same) length so as to promote double cross-overs. The double cross-over functions such that the two recombinogenic regions on the chromosome (separated by nucleotide sequence 'X') and the two recombinogenic regions on the vector (separated by nucleotide sequence 'Y') recombine to leave a chromosome unaltered except that X and Y have interchanged. The position of the recombinogenic regions can both be positioned upstream or downstream of, or may flank, an open reading frame of interest. These regions can consist of coding, non-coding, or a mixture of coding and non-coding sequence. The identity of X and Y will depend on the effect desired. X may be all or part of an open reading frame, and Y no nucleotides at all, which would result in sequence X being deleted from the chromosome. Alternatively Y may be a strong promoter region for insertion upstream of an open reading frame, and therefore X may be no nucleotides at all.

Suitable vectors will vary in composition depending what type of recombination event is to be performed, and what the ultimate purpose of the recombination event is. Integrative vectors used to deliver region Y can be

conditionally replicative or suicide plasmids, bacteriophages, transposons or linear DNA fragments obtained by restriction hydrolysis or PCR amplification. Selection of the recombination event is selected by means of selectable genetic marker such as genes conferring resistance to antibiotics (for instance kanamycin, erythromycin, chloramphenicol, or gentamycin), genes conferring resistance to heavy metals and/or toxic compounds or genes complementing auxotrophic mutations (for instance *pur*, *leu*, *met*, *aro*).

Process a) and f) – Down regulation/Removal of Variable and non-protective immunodominant antigens

Many surface antigens are variable among bacterial strains and as a consequence are protective only against a limited set of closely related strains. An aspect of this invention covers the reduction in expression, or, preferably, the deletion of the gene(s) encoding variable surface protein(s) which results in a bacterial strain producing blebs which, when administered in a vaccine, have a stronger potential for cross-reactivity against various strains due to a higher influence exerted by conserved proteins (retained on the outer membranes) on the vaccinee's immune system. Examples of such variable antigens include: for *Neisseria* - pili (PilC) which undergoes antigenic variations, PorA, Opa, TbpB, FrpB; for *H. influenzae* - P2, P5, pilin, IgA1-protease; and for *Moraxella* - CopB, OMP106.

Other types of gene that could be down-regulated or switched off are genes which, *in vivo*, can easily be switched on (expressed) or off by the bacterium. As outer membrane proteins encoded by such genes are not always present on the bacteria, the presence of such proteins in the bleb preparations can also be detrimental to the effectiveness of the vaccine for the reasons stated above. A preferred example to down-regulate or delete is *Neisseria* Opc protein. Anti-Opc immunity induced by an Opc containing bleb vaccine would only have limited protective capacity as the infecting organism could easily become Opc⁻. *H. influenzae* HgpA and HgpB are other examples of such proteins.

In process a), these variable or non-protective genes are down-regulated in expression, or terminally switched off. This has the above-mentioned surprising advantage of concentrating the immune system on better antigens that are present in low amounts on the outer surface of blebs.

The strain can be engineered in this way by a number of strategies including transposon insertion to disrupt the coding region or promoter region of the gene, or point mutations or deletions to achieve a similar result. Homologous recombination may also be used to delete a gene from a chromosome (where sequence X comprises part (preferably all) of the coding sequence of the gene of interest). It may additionally be used to change its strong promoter for a weaker (or no) promoter (where nucleotide sequence X comprises part (preferably all) of the promoter region of the gene, and nucleotide sequence Y comprises either a weaker promoter region [resulting in a decreased expression of the gene(s)/operon(s) of interest], or no promoter region). In this case it is preferable for the recombination event to occur within the region of the chromosome 1000 bp upstream of the gene of interest.

Alternatively, Y may confer a conditional transcriptional activity, resulting in a conditional expression of the gene(s)/operon(s) of interest (down-regulation). This is useful in the expression of molecules that are toxic to or not well supported by the bacterial host.

Most of the above-exemplified proteins are integral OMPs and their variability may be limited only to one or few of their surface exposed loops. Another aspect of this invention [process g]) covers the deletion of DNA regions coding for these surface exposed loops which leads to the expression of an integral OMP containing conserved surface exposed loops. Surface exposed loops of *H. influenzae* P2 and P5 are preferred examples of proteins that could be transformed into cross-reactive antigens by using such a method. Again, homologous recombination is a preferred method of performing this engineering process.

Process b) - Promoter delivery and modulation:

A further aspect of the invention relates to modifying the composition of blebs by altering *in situ* the regulatory region controlling the expression of gene(s) and/or operon(s) of interest. This alteration may include partial or total replacement of the endogenous promoter controlling the expression of a gene of interest, with one conferring a distinct transcriptional activity. This distinct transcriptional activity may be conferred by variants (point mutations, deletions and/or insertions) of the endogenous control regions, by naturally occurring or modified heterologous promoters or by a combination of both. Such alterations will preferably confer a

transcriptional activity stronger than the endogenous one (introduction of a strong promoter), resulting in an enhanced expression of the gene(s)/operon(s) of interest (up-regulation). Such a method is particularly useful for enhancing the production of immunologically relevant Bleb components such as outer-membrane proteins and lipoproteins (preferably conserved OMPs, usually present in blebs at low concentrations).

Typical strong promoters that may be integrated in *Neisseria* are *porA* [SEQ ID NO: 24], *porB* [SEQ ID NO:26], *igtF*, *Opa*, *p110*, *lst*, and *hpuAB*. *PorA* and *PorB* are preferred as constitutive, strong promoters. It has been established (Example 9) that the *PorB* promoter activity is contained in a fragment corresponding to nucleotides -1 to -250 upstream of the initiation codon of *porB*. In *Moraxella*, it is preferred to use the *ompH*, *ompG*, *ompE*, *OmpB1*, *ompB2*, *ompA*, *OMP*_{CD} and *Omp106* promoters, and in *H. influenzae*, it is preferred to integrate the P2, P4, P1, P5 and P6 promoters.

Using the preferred double cross-over homologous recombination technology to introduce the promoter in the 1000 bp upstream region, promoters can be placed anywhere from 30-970 bp upstream of the initiation codon of the gene to be up-regulated. Although conventionally it is thought the promoter region should be relatively close to the open reading frame in order to obtain optimal expression of the gene, the present inventors have surprisingly found that placement of the promoter further away from the initiation codon results in large increases in expression levels. Thus it is preferred if the promoter is inserted 200-600 bp from the initiation codon of the gene, more preferably 300-500 bp, and most preferably approximately 400 bp from the initiation ATG.

Process c) - Bleb components produced conditionally

The expression of some genes coding for certain bleb components is carefully regulated. The production of the components is conditionally modulated and depends upon various metabolic and/or environmental signals. Such signals include, for example, iron-limitation, modulation of the redox potential, pH and temperature variations, nutritional changes. Some examples of bleb components known to be produced conditionally include iron-regulated outer-membrane proteins from *Neisseria* and *Moraxella* (for instance *TbpB*, *LbpB*), and substrate-inducible outer-

membrane porins. The present invention covers the use of the genetic methods described previously (process a) or b)) to render constitutive the expression of such molecules. In this way, the influence of environmental signal upon the expression of gene(s) of interest can be overcome by modifying/replacing the gene's corresponding control region so that it becomes constitutively active (for instance by deleting part [preferably all] or the repressive control sequence – e.g. the operator region), or inserting a constitutive strong promoter. For iron regulated genes the *fur* operator may be removed. Alternatively, process i) may be used to deliver an additional copy of the gene/operon of interest in the chromosome which is placed artificially under the control of a constitutive promoter.

Processes d), and e) - Detoxification of LPS

The toxicity of bleb vaccines presents one of the largest problems in the use of blebs in vaccines. A further aspect of the invention relates to methods of genetically detoxifying the LPS present in blebs. Lipid A is the primary component of LPS responsible for cell activation. Many mutations in genes involved in this pathway lead to essential phenotypes. However, mutations in the genes responsible for the terminal modifications steps lead to temperature-sensitive (*htrB*) or permissive (*msbB*) phenotypes. Mutations resulting in a decreased (or no) expression of these genes result in altered toxic activity of lipid A. Indeed, the non-lauroylated (*htrB* mutant) or non-myristoylated (*msbB* mutant) lipid A are less toxic than the wild-type lipid A. Mutations in the lipid A 4'-kinase encoding gene (*lpxK*) also decreases the toxic activity of lipid A.

Process d) thus involves either the deletion of part (or preferably all) of one or more of the above open reading frames or promoters. Alternatively, the promoters could be replaced with weaker promoters. Preferably the homologous recombination techniques described above are used to carry out the process.

The sequences of the *htrB* and *msbB* genes from *Neisseria meningitidis* B, *Moraxella catarrhalis*, and *Haemophilus influenzae* are additionally provided for this purpose.

LPS toxic activity could also be altered by introducing mutations in genes/loci involved in polymyxin B resistance (such resistance has been correlated with addition of aminoarabinose on the 4' phosphate of lipid A). These genes/loci could be *pmrE*

that encodes a UDP-glucose dehydrogenase, or a region of antimicrobial peptide-resistance genes common to many enterobacteriaceae which could be involved in aminoarabinose synthesis and transfer. The gene *pmrF* that is present in this region encodes a dolicol-phosphate manosyl transferase (Gunn J.S., Kheng, B.L., Krueger J., Kim K., Guo L., Hackett M., Miller S.I. 1998. *Mol. Microbiol.* 27: 1171-1182).

5 Mutations in the PhoP-PhoQ regulatory system, which is a phospho-relay two component regulatory system (i. e. PhoP constitutive phenotype, PhoP^c), or low Mg⁺⁺ environmental or culture conditions (that activate the PhoP-PhoQ regulatory system) lead to the addition of aminoarabinose on the 4'-phosphate and 2-hydroxymyristate replacing myristate (hydroxylation of myristate). This modified lipid A displays reduced ability to stimulate E-selectin expression by human endothelial cells and TNF- α secretion from human monocytes.

Process e) involves the upregulation of these genes using a strategy as described above (strong promoters being incorporated, preferably using homologous recombination techniques to carry out the process).

Alternatively, rather than performing any such mutation, a polymyxin B resistant strain could be used as a vaccine production strain (in conjunction with one or more of the other processes of the invention), as blebs from such strains also have reduced LPS toxicity (for instance as shown for meningococcus - van der Ley, P, Hamstra, HJ, Kramer, M, Steeghs, L, Petrov, A and Poolman, JT. 1994. *In: Proceedings of the ninth international pathogenic Neisseria conference. The Guildhall, Winchester, England.*)

As a further alternative (and further aspect of the invention) the inventors provide a method of detoxifying a Gram-negative bacterial strain comprising the step of culturing the strain in a growth medium containing 0.1mg-100g of aminoarabinose per litre medium.

As a further still alternative, synthetic peptides that mimic the binding activity of polymyxin B (described below) may be added to the Bleb preparation in order to reduce LPS toxic activity (Rustici, A, Velucchi, M, Faggioni, R, Sironi, M, Ghezzi, P, Quataert, S, Green, B and Porro M 1993. *Science* 259: 361-365; Velucchi, M, Rustici, A, Meazza, C, Villa, P, Ghezzi, P and Porro, M. 1997. *J. Endotox. Res.* 4:).

Process f) - Anchoring homologous or heterologous proteins to outer-membrane blebs whilst reducing the toxicity of LPS

A further aspect of this invention covers the use of genetic sequences encoding polymyxin B peptides (or analogues thereof) as a means to target fusion proteins to the outer-membrane. Polymyxin B is a cyclic peptide composed of non tRNA-encoded amino acids (produced by Gram-positive actinomycetal organisms) that binds very strongly to the Lipid A part of LPS present in the outer-membrane. This binding decreases the intrinsic toxicity of LPS (endotoxin activity). Peptides mimicking the structure of Polymyxin B and composed of canonical (tRNA encoded) amino acids have been developed and also bind lipid A with a strong affinity. These peptides have been used for detoxifying LPS. One of these peptides known as SAEP-2 (Nterminus-Lys-Thr-Lys-Cys-Lys-Phe-Leu-Lys-Lys-Cys-Cterminus) was shown to be very promising in that respect (Molecular Mapping and detoxifying of the Lipid A binding site by synthetic peptides (1993). Rustici, A., Velucchi, M., Faggioni, R., Sironi, M., Ghezzi, P., Quataert, S., Green, B. and M. Porro. *Science* 259, 361-365).

The present process f) of the invention provides an improvement of this use. It has been found that the use of DNA sequences coding for the SEAP-2 peptide (or derivatives thereof), fused genetically to a gene of interest (encoding for instance a T cell antigen or a protective antigen that is usually secreted such as a toxin, or a cytosolic or periplasmic protein) is a means for targeting the corresponding recombinant protein to the outer-membrane of a preferred bacterial host (whilst at the same time reducing the toxicity of the LPS).

This system is suitable for labile proteins which would not be directly exposed to the outside of the bleb. The bleb would therefore act as a delivery vehicle which would expose the protein to the immune system once the blebs had been engulfed by T-cells. Alternatively, the genetic fusion should also comprise a signal peptide or transmembrane domain such that the recombinant protein may cross the outer membrane for exposure to the host's immune system.

This targeting strategy might be of particular interest in the case of genes encoding proteins that are not normally targeted to the outer-membrane. This methodology also allows the isolation of recombinant blebs enriched in the protein of interest. Preferably, such a peptide targeting signal allows the enrichment of outer membrane blebs in one or several proteins of interest, which are naturally not found in

that given subcellular localization. A non exhaustive list of bacteria that can be used as a recipient host for such a production of recombinant blebs includes *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Moraxella catarrhalis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa*, *Chlamydia trachomatis*, and *Chlamydia pneumoniae*.

5 Although it is preferred that the gene for the construct is engineered into the chromosome of the bacterium [using process i)], an alternative preferred embodiment is for SAEP-2-tagged recombinant proteins to be made independently, and attached at a later stage to a bleb preparation.

10 A further embodiment is the use of such constructs in a method of protein purification. The system could be used as part of an expression system for producing recombinant proteins in general. The SAEP-2 peptide tag can be used for affinity purification of the protein to which it is attached using a column containing immobilised lipid A molecules.

15 Process h) - Cross-reactive polysaccharides

 The isolation of bacterial outer-membrane blebs from encapsulated Gram-negative bacteria often results in the co-purification of capsular polysaccharide. In some cases, this “contaminant” material may prove useful since polysaccharide may enhance the immune response conferred by other bleb components. In other cases
20 however, the presence of contaminating polysaccharide material in bacterial bleb preparations may prove detrimental to the use of the blebs in a vaccine. For instance, it has been shown at least in the case of *N. meningitidis* that the serogroup B capsular polysaccharide does not confer protective immunity and is susceptible to induce an adverse auto-immune response in humans. Consequently, process h) of the invention
25 is the engineering of the bacterial strain for bleb production such that it is free of capsular polysaccharide. The blebs will then be suitable for use in humans. A particularly preferred example of such a bleb preparation is one from *N. meningitidis* serogroup B devoid of capsular polysaccharide.

 This may be achieved by using modified bleb production strains in which the
30 genes necessary for capsular biosynthesis and/or export have been impaired. Inactivation of the gene coding for capsular polysaccharide biosynthesis or export can be achieved by mutating (point mutation, deletion or insertion) either the control region, the coding region or both (preferably using the homologous recombination

techniques described above). Moreover, inactivation of capsular biosynthesis genes may also be achieved by antisense over-expression or transposon mutagenesis. A preferred method is the deletion of some or all of the *Neisseria meningitidis* *cps* genes required for polysaccharide biosynthesis and export. For this purpose, the replacement
5 plasmid pMF121 (described in Frosh et al.1990, *Mol. Microbiol.* 4:1215-1218) can be used to deliver a mutation deleting the *cpsCAD* (+ *galE*) gene cluster. Alternatively the *siaD* gene could be deleted, or down-regulated in expression (the meningococcal *siaD* gene encodes alpha-2,3-sialyltransferase, an enzyme required for capsular polysaccharide and LOS synthesis). Such mutations may also remove host-similar
10 structures on the saccharide portion of the LPS of the bacteria.

Process i) - Delivery of one or more further copies of a Gene and/or operon in a host chromosome, or delivery of a heterologous gene and/or operon in a host chromosome.

An efficient strategy to modulate the composition of a Bleb preparation is to
15 deliver one or more copies of a DNA segment containing an expression cassette into the genome of a Gram-negative bacterium. A non exhaustive list of preferred bacterial species that could be used as a recipient for such a cassette includes *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Moraxella catarrhalis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa*, *Chlamydia trachomatis*, *Chlamydia*
20 *pneumoniae*. The gene(s) contained in the expression cassette may be homologous (or endogenous) (i.e. exist naturally in the genome of the manipulated bacterium) or heterologous (i.e. do not exist naturally in the genome of the manipulated bacterium). The reintroduced expression cassette may consist of unmodified, "natural" promoter/gene/operon sequences or engineered expression cassettes in which the
25 promoter region and/or the coding region or both have been altered. A non-exhaustive list of preferred promoters that could be used for expression includes the promoters *porA*, *porB*, *lbpB*, *tbpB*, *p110*, *lst*, *hpuAB* from *N. meningitidis* or *N. gonorrhoeae*, the promoters *p2*, *p5*, *p4*, *ompF*, *p1*, *ompH*, *p6*, *hin47* from *H. influenzae*, the promoters *ompH*, *ompG*, *ompCD*, *ompE*, *ompB1*, *ompB2*, *ompA* of *M. catarrhalis*, the
30 promoter λ pL, *lac*, *tac*, *araB* of *Escherichia coli* or promoters recognized specifically by bacteriophage RNA polymerase such as the *E. coli* bacteriophage T7. A non-exhaustive list of preferred genes that could be expressed in such a system includes *Neisseria* *NspA*, *Omp85*, *PilQ*, *TbpA/B* complex, *Hsf*, *PldA*, *HasR*; *Chlamydia*

MOMP, HMWP; *Moraxella* OMP106, HasR, PilQ, OMP85, PldA; *Bordetella pertussis* FHA, PRN, PT.

In a preferred embodiment of the invention the expression cassette is delivered and integrated in the bacterial chromosome by means of homologous and/or site specific recombination. Integrative vectors used to deliver such genes and/or operons can be conditionally replicative or suicide plasmids, bacteriophages, transposons or linear DNA fragments obtained by restriction hydrolysis or PCR amplification. Integration is preferably targeted to chromosomal regions dispensable for growth *in vitro*. A non exhaustive list of preferred loci that can be used to target DNA integration includes the *porA*, *porB*, *opa*, *opc*, *rmp*, *omp26*, *lecA*, *cps*, *lgtB* genes of *Neisseria meningitidis* and *Neisseria gonorrhoeae*, the *P1*, *P5*, *hmw1/2*, *IgA-protease*, *fimE* genes of NTHi; the *lecA1*, *lecA2*, *omp106*, *uspA1*, *uspA2* genes of *Moraxella catarrhalis*. Alternatively, the expression cassette used to modulate the expression of bleb component(s) can be delivered into a bacterium of choice by means of episomal vectors such as circular/linear replicative plasmids, cosmids, phasmids, lysogenic bacteriophages or bacterial artificial chromosomes. Selection of the recombination event can be selected by means of selectable genetic marker such as genes conferring resistance to antibiotics (for instance kanamycin, erythromycin, chloramphenicol, or gentamycin), genes conferring resistance to heavy metals and/or toxic compounds or genes complementing auxotrophic mutations (for instance *pur*, *leu*, *met*, *aro*).

Heterologous Genes - Expression of foreign proteins in outer-membrane blebs

Outer-membrane bacterial blebs represent a very attractive system to produce, isolate and deliver recombinant proteins for vaccine, therapeutic and/or diagnostic uses. A further aspect of this invention is in respect of the expression, production and targeting of foreign, heterologous proteins to the outer-membrane, and the use of the bacteria to produce recombinant blebs.

A preferred method of achieving this is via a process comprising the steps of: introducing a heterologous gene, optionally controlled by a strong promoter sequence, into the chromosome of a Gram-negative strain by homologous recombination. Blebs may be made from the resulting modified strain.

A non-exhaustive list of bacteria that can be used as a recipient host for production of recombinant blebs includes *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Moraxella catarrhalis*, *Haemophilus influenzae*, *Pseudomonas aeruginosa*, *Chlamydia trachomatis*, *Chlamydia pneumoniae*. The gene expressed in such a system can be of viral, bacterial, fungal, parasitic or higher eukaryotic origin.

A preferred application of the invention includes a process for the expression of *Moraxella*, *Haemophilus* and/or *Pseudomonas* outer-membrane proteins (integral, polytopic and/or lipoproteins) in *Neisseria meningitidis* recombinant blebs. The preferable integration loci are stated above, and genes that are preferably introduced are those that provide protection against the bacterium from which they were isolated. Preferred protective genes for each bacterium are described below.

Further preferred applications are: blebs produced from a modified *Haemophilus influenzae* strain where the heterologous gene is a protective OMP from *Moraxella catarrhalis*; and blebs produced from a modified *Moraxella catarrhalis* strain where the heterologous gene is a protective OMP from *Haemophilus influenzae* (preferred loci for gene insertion are given above, and preferred protective antigens are described below).

A particularly preferred application of this aspect is in the field of the prophylaxis or treatment of sexually-transmitted diseases (STDs). It is often difficult for practitioners to determine whether the principal cause of a STD is due to gonococcus or *Chlamydia trachomatis* infection. These two organisms are the main causes of salpingitis – a disease which can lead to sterility in the host. It would therefore be useful if a STD could be vaccinated against or treated with a combined vaccine effective against disease caused by both organisms. The Major Outer Membrane Protein (MOMP) of *C. trachomatis* has been shown to be the target of protective antibodies. However, the structural integrity of this integral membrane protein is important for inducing such antibodies. In addition, the epitopes recognised by these antibodies are variable and define more than 10 serovars. The previously described aspect of this invention allows the proper folding of one or more membrane proteins within a bleb outer membrane preparation. The engineering of a gonococcal strain expressing multiple *C. trachomatis* MOMP serovars in the outer membrane, and the production of blebs therefrom, produces a single solution to the multiple problems of correctly folded membrane proteins, the presentation of sufficient MOMP serovars

to protect against a wide spectrum of serovars, and the simultaneous prophylaxis/treatment of gonococcal infection (and consequently the non-requirement of practitioners to initially decide which organism is causing particular clinical symptoms – both organisms can be vaccinated against simultaneously thus allowing the treatment of the STD at a very early stage). Preferred loci for gene insertion in the gonococcal chromosome are give above. Other preferred, protective *C. trachomatis* genes that could be incorporated are HMWP, PmpG and those OMPs disclosed in WO 99/28475.

10 *Targeting of heterologous proteins to outer-membrane blebs:*

The expression of some heterologous proteins in bacterial blebs may require the addition of outer-membrane targeting signal(s). The preferred method to solve this problem is by creating a genetic fusion between a heterologous gene and a gene coding for a resident OMP as a specific approach to target recombinant proteins to blebs. Most preferably, the heterologous gene is fused to the signal peptides sequences of such an OMP.

Neisserial bleb preparations

One or more of the following genes (encoding protective antigens) are preferred for upregulation via processes b) and/or i) when carried out on a Neisserial strain, including gonococcus, and meningococcus (particularly *N. meningitidis* B): NspA (WO 96/29412), Hsf-like (WO 99/31132), Hap (PCT/EP99/02766), PorA, PorB, OMP85 (WO 00/23595), PilQ (PCT/EP99/03603), PldA (PCT/EP99/06718), FrpB (WO 96/31618), TbpA (US 5,912,336), TbpB, FrpA/FrpC (WO 92/01460), LbpA/LbpB (PCT/EP98/05117), FhaB (WO 98/02547), HasR (PCT/EP99/05989), lipo02 (PCT/EP99/08315), Tbp2 (WO 99/57280), MltA (WO 99/57280), and ctrA (PCT/EP00/00135). They are also preferred as genes which may be heterologously introduced into other Gram-negative bacteria.

One or more of the following genes are preferred for downregulation via process a): PorA, PorB, PilC, TbpA, TbpB, LbpA, LbpB, Opa, and Opc.

One or more of the following genes are preferred for downregulation via process d): htrB, msbB and lpxK.

One or more of the following genes are preferred for upregulation via process e): pmrA, pmrB, pmrE, and pmrF.

Preferred repressive control sequences for process c) are: the *fur* operator region (particularly for either or both of the TbpB or LbpB genes); and the DtxR operator region.

One or more of the following genes are preferred for downregulation via process h): galE, siaA, siaB, siaC, siaD, ctrA, ctrB, ctrC, and ctrD.

Pseudomonas aeruginosa bleb preparations

One or more of the following genes (encoding protective antigens) are preferred for upregulation via processes b) and/or i): PcrV, OprF, OprI. They are also preferred as genes which may be heterologously introduced into other Gram-negative bacteria.

Moraxella catarrhalis bleb preparations

One or more of the following genes (encoding protective antigens) are preferred for upregulation via processes b) and/or i): OMP106 (WO 97/41731 & WO 96/34960), HasR (PCT/EP99/03824), PilQ (PCT/EP99/03823), OMP85 (PCT/EP00/01468), lipo06 (GB 9917977.2), lipo10 (GB 9918208.1), lipo11 (GB 9918302.2), lipo18 (GB 9918038.2), P6 (PCT/EP99/03038), ompCD, CopB (Helminen ME, et al (1993) Infect. Immun. 61:2003-2010), D15 (PCT/EP99/03822), OmpIA1 (PCT/EP99/06781), Hly3 (PCT/EP99/03257), LbpA and LbpB (WO 98/55606), TbpA and TbpB (WO 97/13785 & WO 97/32980), OmpE, UspA1 and UspA2 (WO 93/03761), and Omp21. They are also preferred as genes which may be heterologously introduced into other Gram-negative bacteria.

One or more of the following genes are preferred for downregulation via process a): CopB, OMP106, OmpB1, TbpA, TbpB, LbpA, and LbpB.

One or more of the following genes are preferred for downregulation via process d): htrB, msbB and lpxK.

One or more of the following genes are preferred for upregulation via process e): pmrA, pmrB, pmrE, and pmrF.

Haemophilus influenzae bleb preparations

One or more of the following genes (encoding protective antigens) are preferred for upregulation via processes b) and/or i): D15 (WO 94/12641), P6 (EP 281673), TbpA, TbpB, P2, P5 (WO 94/26304), OMP26 (WO 97/01638), HMW1, HMW2, HMW3, HMW4, Hia, Hsf, Hap, Hin47, and Hif (all genes in this operon should be upregulated in order to upregulate pilin). They are also preferred as genes which may be heterologously introduced into other Gram-negative bacteria.

One or more of the following genes are preferred for downregulation via process a): P2, P5, Hif, IgA1-protease, HgpA, HgpB, HMW1, HMW2, Hxu, TbpA, and TbpB.

One or more of the following genes are preferred for downregulation via process d): htrB, msbB and lpxK.

One or more of the following genes are preferred for upregulation via process e): pmrA, pmrB, pmrE, and pmrF.

Vaccine Formulations

A preferred embodiment of the invention is the formulation of the bleb preparations of the invention in a vaccine which may also comprise a pharmaceutically acceptable excipient.

The manufacture of bleb preparations from any of the aforementioned modified strains may be achieved by any of the methods well known to a skilled person. Preferably the methods disclosed in EP 301992, US 5,597,572, EP 11243 or US 4,271,147 are used. Most preferably, the method described in Example 8 is used.

Vaccine preparation is generally described in Vaccine Design ("The subunit and adjuvant approach" (eds Powell M.F. & Newman M.J.) (1995) Plenum Press New York).

The bleb preparations of the present invention may be adjuvanted in the vaccine formulation of the invention. Suitable adjuvants include an aluminium salt such as aluminum hydroxide gel (alum) or aluminium phosphate, but may also be a salt of calcium (particularly calcium carbonate), iron or zinc, or may be an insoluble suspension of acylated tyrosine, or acylated sugars, cationically or anionically derivatised polysaccharides, or polyphosphazenes.

Suitable Th1 adjuvant systems that may be used include, Monophosphoryl lipid A, particularly 3-de-O-acylated monophosphoryl lipid A, and a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL) together with an aluminium salt. An enhanced system involves the combination of a monophosphoryl lipid A and a saponin derivative particularly the combination of
5 QS21 and 3D-MPL as disclosed in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol as disclosed in WO96/33739. A particularly potent adjuvant formulation involving QS21 3D-MPL and tocopherol in an oil in water emulsion is described in WO95/17210 and is a preferred formulation.

10 The vaccine may comprise a saponin, more preferably QS21. It may also comprise an oil in water emulsion and tocopherol. Unmethylated CpG containing oligo nucleotides (WO 96/02555) are also preferential inducers of a TH1 response and are suitable for use in the present invention.

The vaccine preparation of the present invention may be used to protect or
15 treat a mammal susceptible to infection, by means of administering said vaccine via systemic or mucosal route. These administrations may include injection *via* the intramuscular, intraperitoneal, intradermal or subcutaneous routes; or *via* mucosal administration to the oral/alimentary, respiratory, genitourinary tracts. Thus one aspect of the present invention is a method of immunizing a human host against a
20 disease caused by infection of a gram-negative bacteria, which method comprises administering to the host an immunoprotective dose of the bleb preparation of the present invention.

The amount of antigen in each vaccine dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in
25 typical vaccinees. Such amount will vary depending upon which specific immunogen is employed and how it is presented. Generally, it is expected that each dose will comprise 1-100 μ g of protein antigen, preferably 5-50 μ g, and most typically in the range 5 - 25 μ g.

An optimal amount for a particular vaccine can be ascertained by standard
30 studies involving observation of appropriate immune responses in subjects. Following an initial vaccination, subjects may receive one or several booster immunisations adequately spaced.

Ghost or Killed Whole cell vaccines

The inventors envisage that the above improvements to bleb preparations and vaccines can be easily extended to ghost or killed whole cell preparations and vaccines (with identical advantages). The modified Gram-negative strains of the invention from which the bleb preparations are made can also be used to made ghost and killed whole cell preparations. Methods of making ghost preparations (empty cells with intact envelopes) from Gram-negative strains are well known in the art (see for example WO 92/01791). Methods of killing whole cells to make inactivated cell preparations for use in vaccines are also well known. The terms 'bleb preparations' and 'bleb vaccines' as well as the processes described throughout this document are therefore applicable to the terms 'ghost preparation' and 'ghost vaccine', and 'killed whole cell preparation' and 'killed whole cell vaccine', respectively, for the purposes of this invention.

15 Combinations of methods a) – i)

It may be appreciated that one or more of the above processes may be used to produce a modified strain from which to make improved bleb preparations of the invention. Preferably one such process is used, more preferably two or more (2, 3, 4, 5, 6, 7, 8 or 9) of the processes are used in order to manufacture the bleb vaccine. As each additional method is used in the manufacture of the bleb vaccine, each improvement works in conjunction with the other methods used in order to make an optimised engineered bleb preparation.

A preferred meningococcal (particularly *N. meningitidis* B) bleb preparation comprises the use of processes a), b), d) and/or e), and h). Such bleb preparations are safe (no structures similar to host structures), non-toxic, and structured such that the host immune response will be focused on high levels of protective (and preferably conserved) antigens. All the above elements work together in order to provide an optimised bleb vaccine.

Similarly for *M. catarrhalis* and non-typeable *H. influenzae*, preferred bleb preparations comprise the use of processes a), b), and d) and/or e).

A further aspect of the invention is thus an immuno-protective and non-toxic Gram-negative bleb, ghost, or killed whole cell vaccine suitable for paediatric use.

By paediatric use it is meant use in infants less than 4 years old.

By immunoprotective it is meant that at least 40% (and preferably 50, 60, 70, 80, 90 and 100%) of infants seroconvert (4-fold increase in bactericidal activity [the dilution of antisera at which 50% of bacteria die – see for example PCT/EP98/05117]) against a set of heterologous strains to be selected from the major clonal groups known. For meningococcus B these stains should have a different PorA type from the bleb production strain, and should preferably be 2, 3, 4 or, most preferably, all 5 of strains H44/76, M97/252078, BZ10, NGP165 and CU385. For non-typeable *H. influenzae*, the strains should preferably be 2, 3, 4 or, most preferably, all 5 of strains 3224A, 3219C, 3241A, 640645, and A840177. For *M. catarrhalis*, the strains should preferably be 2, 3, 4 or, most preferably, all 5 of strains ATCC 43617, 14, 358, 216 and 2926.

By non-toxic it is meant that there is a significant (2-4 fold, preferably 10 fold) decrease of endotoxin activity as measured by the well-known LAL and pyrogenicity assays.

Vaccine Combinations

A further aspect of the invention are vaccine combinations comprising the bleb preparations of the invention with other antigens which are advantageously used against certain disease states. It has been found that blebs are particularly suitable for formulating with other antigens, as they advantageously have an adjuvant effect on the antigens they are mixed with.

In one preferred combination, the meningococcus B bleb preparations of the invention are formulated with 1, 2, 3 or preferably all 4 of the following meningococcal capsular polysaccharides which may be plain or conjugated to a protein carrier: A, C, Y or W. Such a vaccine may be advantageously used as a global meningococcus vaccine. Rather than use the meningococcus B bleb preparations of the invention, it is also envisaged that the formulation could alternatively contain wild-type meningococcus B bleb preparations from 2 or more (preferably several) strains belonging to several subtype/serotypes (for instance chosen from P1.15, P1.7,16, P1.4, and P1.2).

In a further preferred embodiment, the meningococcus B bleb preparations of the invention [or the aforementioned mix of 2 or more wild-type meningococcus B bleb preparations], preferably formulated with 1, 2, 3 or all 4 of the plain or

conjugated meningococcal capsular polysaccharides A, C, Y or W, are formulated with a conjugated *H. influenzae* b capsular polysaccharide, and one or more plain or conjugated pneumococcal capsular polysaccharides. Optionally, the vaccine may also comprises one or more protein antigens that can protect a host against *Streptococcus*
5 *pneumoniae* infection. Such a vaccine may be advantageously used as a global meningitis vaccine.

The pneumococcal capsular polysaccharide antigens are preferably selected from serotypes 1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F and 33F (most preferably from serotypes 1, 3, 4, 5, 6B, 7F,
10 9V, 14, 18C, 19F and 23F).

Preferred pneumococcal proteins antigens are those pneumococcal proteins which are exposed on the outer surface of the pneumococcus (capable of being recognised by a host's immune system during at least part of the life cycle of the pneumococcus), or are proteins which are secreted or released by the pneumococcus.

15 Most preferably, the protein is a toxin, adhesin, 2-component signal transducer, or lipoprotein of *Streptococcus pneumoniae*, or fragments thereof. Particularly preferred proteins include, but are not limited to: pneumolysin (preferably detoxified by chemical treatment or mutation) [Mitchell *et al.* Nucleic Acids Res. 1990 Jul 11; 18(13): 4010 "Comparison of pneumolysin genes and proteins from *Streptococcus*
20 *pneumoniae* types 1 and 2.", Mitchell *et al.* Biochim Biophys Acta 1989 Jan 23; 1007(1): 67-72 "Expression of the pneumolysin gene in *Escherichia coli*: rapid purification and biological properties.", WO 96/05859 (A. Cyanamid), WO 90/06951 (Paton *et al.*), WO 99/03884 (NAVA)]; PspA and transmembrane deletion variants thereof (US 5804193 - Briles *et al.*); PspC and transmembrane deletion variants
25 thereof (WO 97/09994 - Briles *et al.*); PsaA and transmembrane deletion variants thereof (Berry & Paton, Infect Immun 1996 Dec;64(12):5255-62 "Sequence heterogeneity of PsaA, a 37-kilodalton putative adhesin essential for virulence of *Streptococcus pneumoniae*"); pneumococcal choline binding proteins and transmembrane deletion variants thereof; CbpA and transmembrane deletion variants
30 thereof (WO 97/41151; WO 99/51266); Glyceraldehyde-3-phosphate - dehydrogenase (Infect. Immun. 1996 64:3544); HSP70 (WO 96/40928); PcpA (Sanchez-Beato *et al.* *FEMS Microbiol Lett* 1998, 164:207-14); M like protein, SB patent application No. EP 0837130; and adhesin 18627, SB Patent application No. EP

0834568. Further preferred pneumococcal protein antigens are those disclosed in WO 98/18931, particularly those selected in WO 98/18930 and PCT/US99/30390.

In a further preferred embodiment, the *Moraxella catarrhalis* bleb preparations of the invention are formulated with one or more plain or conjugated pneumococcal capsular polysaccharides, and one or more antigens that can protect a host against non-typeable *H. influenzae* infection. Optionally, the vaccine may also comprise one or more protein antigens that can protect a host against *Streptococcus pneumoniae* infection. The vaccine may also optionally comprise one or more antigens that can protect a host against RSV and/or one or more antigens that can protect a host against influenza virus. Such a vaccine may be advantageously used as a global otitis media vaccine.

Preferred non-typeable *H. influenzae* protein antigens include Fimbrin protein (US 5766608) and fusions comprising peptides therefrom (eg LB1 Fusion) (US 5843464 - Ohio State Research Foundation), OMP26, P6, protein D, TbpA, TbpB, Hia, Hmw1, Hmw2, Hap, and D15.

Preferred influenza virus antigens include whole, live or inactivated virus, split influenza virus, grown in eggs or MDCK cells, or Vero cells or whole flu virosomes (as described by R. Gluck, Vaccine, 1992, 10, 915-920) or purified or recombinant proteins thereof, such as HA, NP, NA, or M proteins, or combinations thereof.

Preferred RSV (Respiratory Syncytial Virus) antigens include the F glycoprotein, the G glycoprotein, the HN protein, or derivatives thereof.

In a still further preferred embodiment, the non-typeable *H. influenzae* bleb preparations of the invention are formulated with one or more plain or conjugated pneumococcal capsular polysaccharides, and one or more antigens that can protect a host against *M. catarrhalis* infection. Optionally, the vaccine may also comprise one or more protein antigens that can protect a host against *Streptococcus pneumoniae* infection. The vaccine may also optionally comprise one or more antigens that can protect a host against RSV and/or one or more antigens that can protect a host against influenza virus. Such a vaccine may be advantageously used as a global otitis media vaccine.

Nucleotide sequences of the invention

A further aspect of the invention relates to the provision of new nucleotide sequences which may be used in the processes of the invention. Specific upstream regions from various genes from various strains are provided which can be used in, for instance, processes a), b), d) and h). In addition, coding regions are provided for performing process d).

General method for the analysis of the non-coding flanking region of a bacterial gene, and its exploitation for modulated expression of the gene in blebs

The non-coding flanking regions of a specific gene contain regulatory elements important in the expression of the gene. This regulation takes place both at the transcriptional and translational level. The sequence of these regions, either upstream or downstream of the open reading frame of the gene, can be obtained by DNA sequencing. This sequence information allows the determination of potential regulatory motifs such as the different promoter elements, terminator sequences, inducible sequence elements, repressors, elements responsible for phase variation, the Shine-Dalgarno sequence, regions with potential secondary structure involved in regulation, as well as other types of regulatory motifs or sequences.

This sequence information allows the modulation of the natural expression of the gene in question. The upregulation of the gene expression may be accomplished by altering the promoter, the Shine-Dalgarno sequence, potential repressor or operator elements, or any other elements involved. Likewise, downregulation of expression can be achieved by similar types of modifications. Alternatively, by changing phase variation sequences, the expression of the gene can be put under phase variation control, or may be uncoupled from this regulation. In another approach, the expression of the gene can be put under the control of one or more inducible elements allowing regulated expression. Examples of such regulation includes, but is not limited to, induction by temperature shift, addition of inductor substrates like selected carbohydrates or their derivatives, trace elements, vitamins, co-factors, metal ions, etc.

Such modifications as described above can be introduced by several different means. The modification of sequences involved in gene expression can be done *in vivo* by random mutagenesis followed by selection for the desired phenotype. Another approach consists in isolating the region of interest and modifying it by random

mutagenesis, or site-directed replacement, insertion or deletion mutagenesis. The modified region can then be reintroduced into the bacterial genome by homologous recombination, and the effect on gene expression can be assessed. In another approach, the sequence knowledge of the region of interest can be used to replace or delete all or part of the natural regulatory sequences. In this case, the regulatory region targeted is isolated and modified so as to contain the regulatory elements from another gene, a combination of regulatory elements from different genes, a synthetic regulatory region, or any other regulatory region, or to delete selected parts of the wild-type regulatory sequences. These modified sequences can then be reintroduced into the bacterium via homologous recombination into the genome.

In process b), for example, the expression of a gene can be modulated by exchanging its promoter with a stronger promoter (through isolating the upstream sequence of the gene, in vitro modification of this sequence, and reintroduction into the genome by homologous recombination). Upregulated expression can be obtained in both the bacterium as well as in the outer membrane vesicles shed (or made) from the bacterium.

In other preferred examples, the described approaches can be used to generate recombinant bacterial strains with improved characteristics for vaccine applications, as described above. These can be, but are not limited to, attenuated strains, strains with increased expression of selected antigens, strains with knock-outs (or decreased expression) of genes interfering with the immune response, and strains with modulated expression of immunodominant proteins.

SEQ ID NO:2-23, 25, 27-38 are all Neisserial upstream sequences (upstream of the initiation codon of various preferred genes) comprising approximately 1000 bp each. SEQ ID NO: 39-62 are all *M. catarrhalis* upstream sequences (upstream of the initiation codon of various preferred genes) comprising approximately 1000 bp each. SEQ ID NO: 63-75 are all *H. influenzae* upstream sequences (upstream of the initiation codon of various preferred genes) comprising approximately 1000 bp each. All of these can be used in genetic methods (particularly homologous recombination) for up-regulating, or down-regulating the open reading frames to which they are associated (as described before). SEQ ID NO: 76-81 are the coding regions for the HtrB and MsbB genes from Neisseria, *M. catarrhalis*, and *Haemophilus influenzae*. These can be used in genetic methods (particularly homologous recombination) for

down-regulating (in particular deleting) part (preferably all) of these genes [process d)].

Another aspect of the invention is thus an isolated polynucleotide sequence which hybridises under highly stringent conditions to at least a 30 nucleotide portion of the nucleotides in SEQ ID NO: 2-23, 25, 27-81 or a complementary strand thereof. Preferably the isolated sequence should be long enough to perform homologous recombination with the chromosomal sequence if it is part of a suitable vector – namely at least 30 nucleotides (preferably at least 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, or 500 nucleotides). More preferably the isolated polynucleotide should comprise at least 30 nucleotides (preferably at least 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, or 500 nucleotides) of SEQ ID NO: 2-23, 25, 27-81 or a complementary strand thereof.

As used herein, highly stringent hybridization conditions include, for example, 6X SSC, 5X Denhardt, 0.5% SDS, and 100 µg/mL fragmented and denatured salmon sperm DNA hybridized overnight at 65 °C and washed in 2X SSC, 0.1% SDS one time at room temperature for about 10 minutes followed by one time at 65 °C for about 15 minutes followed by at least one wash in 0.2X SSC, 0.1% SDS at room temperature for at least 3-5 minutes.

A further aspect is the use of the isolated polynucleotide sequences of the invention in performing a genetic engineering event (such as transposon insertion, or site specific mutation or deletion, but preferably a homologous recombination event) within 1000 bp upstream of a Gram-negative bacterial chromosomal gene in order to either increase or decrease expression of the gene. Preferably the strain in which the recombination event is to take place is the same as the strain from which the upstream sequences of the invention were obtained. However, the meningococcus A, B, C, Y and W and gonococcus genomes are sufficiently similar that upstream sequence from any of these strains may be suitable for designing vectors for performing such events in the other strains. This is may also be the case for *Haemophilus influenzae* and non-typeable *Haemophilus influenzae*.

30

EXAMPLES

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The

5 examples are illustrative, but do not limit the invention.

Example 1: Construction of a *Neisseria meningitidis* serogroup B strain lacking capsular polysaccharides.

10 The plasmid pMF121 (Frosch et al., 1990) has been used to construct a *Neisseria meningitidis* B strain lacking the capsular polysaccharide. This plasmid contains the flanking regions of the gene locus coding for the biosynthesis pathway of the group B polysaccharide (B PS), and the erythromycin resistance gene. Deletion of the B PS resulted in loss of expression of the group B capsular polysaccharide as well as a
15 deletion in the active copy of *galE* leading to the synthesis of galactose deficient LPS.

Strain transformation:

Neisseria meningitidis B H44/76 strain (B:15:P1.7, 16;Los 3,7,9) was selected for transformation. After an overnight CO₂ incubation on MH plate (without
20 erythromycin), cells were collected in liquid MH containing 10 mM MgCl₂ (2 ml were used per MH plate) and diluted up to an OD of 0.1 (550 nm). To this 2 ml solution, 4 µl of the plasmid pMF121 stock solution (0.5 µg/ml) were added for a 6 hours incubation period at 37°C (with shaking). A control group was done with the same amount of *Neisseria meningitidis* B bacteria, but without addition of plasmid. After
25 the incubation period, 100 µl of culture, as such, at 1/10, 1/100 and 1/1000 dilutions, were put in MH plates containing 5, 10, 20, 40 or 80 µg erythromycin /ml before incubation for 48 hours at 37°C.

Colony blotting:

30 After plate incubation, 20 colonies were grown and selected from the 10 and 20 µg erythromycin/ml MH plates, while there was no colony growth in the control group without plasmid transformation. The H44/76 wild type strain was unable to grow in the selected erythromycin plates (10 to 80 µg erythromycin/ml). The day after, all the

visible colonies were placed on new MH plates without erythromycin in order to let them grow. Afterwards, they were transferred onto nitrocellulose sheets (colony blotting) for presence of B polysaccharide. Briefly, colonies were blotted onto a nitrocellulose sheet and rinsed directly in PBS-0.05 % Tween 20 before cell
5 inactivation for 1 hour at 56°C in PBS-0.05% Tween 20 (diluant buffer). Afterwards, the membrane was overlaid for one hour in the diluant buffer at room temperature (RT). Then, sheets were washed again for three times 5 minutes in the diluant buffer before incubation with the anti-B PS 735 Mab (Boehringer) diluted at 1/3000 in the diluant buffer for 2 hours at RT. After a new washing step (3 times 5 minutes), the
10 monoclonal antibody was detected with a biotinylated anti-mouse Ig from Amersham (RPN 1001) diluted 500 times in the diluant buffer (one hour at RT) before the next washing step (as described above). Afterwards, sheets were incubated for one hour at RT with a solution of streptavidin-peroxidase complex diluted 1/1000 in the diluant buffer. After the last three washing steps using the same method, nitrocellulose sheets
15 were incubated for 15 min in the dark using the revelation solution (30 mg of 4-chloro-1-naphtol solution in 10 ml methanol plus 40 ml PBS and 30 mcl of H₂O₂ 37% from Merck). The reaction was stopped with a distilled water-washing step.

Whole cell Elisas:

20 Whole cell Elisas were also done using the two transformed colonies (“D” and “R”) and the wild type strain (H44/76) as coated bacteria (20 µg protein/ml), and a set of different monoclonal antibodies used to characterize *Neisseria meningitidis* strains. The following Mabs were tested: anti-B PS (735 from Dr Frosch), and the other Mabs from NIBSC: anti-B PS (Ref 95/750) anti-P1.7 (A-PorA, Ref 4025), anti-P1.16 (A-PorA, Ref 95/720), anti-Los 3,7,9 (A-LPS, Ref 4047), anti-Los 8 (A-LPS, Ref 4048),
25 and anti-P1.2 (A-PorA Ref 95/696).

Microtiter plates (Maxisorp, Nunc) were coated with 100 µl of the recombinant meningococcal B cells solution overnight (ON) at 37°C at around 20 µg/ml in PBS. Afterwards, plates are washed three times with 300 µl of 150 mM
30 NaCl - 0.05 % Tween 20, and were overlaid with 100 µl of PBS-0.3 % Casein and incubated for 30 min at room temperature with shaking. Plates were washed again using the same procedure before incubation with antibodies. Monoclonal antibodies (100 µl) were used at different dilutions (as shown in Figure 2) in PBS-0.3 % Casein

0.05 % Tween 20 and put onto the microplates before incubation at room temperature for 30 min with shaking, before the next identical washing step. 100 µl of the anti-mouse Ig (from rabbit, Dakopatts E0413) conjugated to biotin and diluted at 1/2000 in PBS-0.3 % Casein - 0.05 % Tween 20 were added to the wells to detect bound
5 monoclonal antibodies. After the washing step (as before), plates were incubated with a streptavidin-peroxidase complex solution (100 µl of the Amersham RPN 1051) diluted at 1/4000 in the same working solution for 30 min at room temperature under shaking conditions. After this incubation and the last washing step, plates are incubated with 100 µl of the chromogen solution (4 mg orthophenylenediamine
10 (OPD) in 10 ml 0.1 M citrate buffer pH4.5 with 5 µl H₂O₂) for 15 min in the dark. Plates are then read at 490/620 nm using a spectrophotometer.

Results:

Figure 1 shows that from the 20 isolated colonies, which were able to growth
15 on the selected medium with erythromycin, only two (the "D" and the "R") colonies were shown negative for presence of B polysaccharide. Among the others, 16 were clearly positive for B PS and still resistant to erythromycin. This indicated that they integrated the plasmid into their genome, but in the wrong orientation, and keeping intact the B PS and LPS gene (no double crossing-over). Positive and negative
20 controls were also tested on the plates, and showed that the H44/76 wild type NmB strain was clearly positive for the B polysaccharide, while meningococcus A (A1) and meningococcus C (C11) strains were clearly negative with this anti-B PS 735 Mab. These results indicate that around 10 % of the selected colonies correctly integrated the plasmid in their genome by making a double crossing-over, while the other
25 strains/colonies were obtained after a simple crossing-over, keeping the B PS and LPS genes intact and expressed.

Using whole cell Elisa, results (Figure 2 and the Table below) clearly indicate that the two "D" and "R" transformants (derived from D and R colonies) can not be recognized anymore by the anti-B PS Mabs (735 and 95/750), nor by the anti-Los
30 3,7,9 and anti-Los 8 Mabs. However, when using specific anti-PorA Mabs, there is a clear reaction with the anti-P1.7 and anti-P1.16 Mabs on the cells, as also observed in the wild-type strain. No reaction was observed with a non-specific anti-PorA Mab (anti-P1.2 mab). These results confirm that the PorA protein, and specifically P1.7

and P1.16 epitopes are still present after transformation, while B polysaccharide and Los 3,7,9 and Los 8 epitopes (LPS) were not.

Table : Specificities of the monoclonal antibodies tested

Mabs Tested	Directed against	Result
Anti-B PS 735	B polysaccharide	++ on the wild type strain (-) on the "D" and "R" mutants
Anti-B PS 95/750 from NIBSC	B PS	++ on the wild type strain (-) on the "D" and "R" mutants
Anti-P1.7 (NIBSC)	Loop 1 of Porin A	++ on all wild type and mutants strains
Anti-P1.16 (NIBSC)	Loop 4 of Porin A	++ on all wild type and mutants strains
Anti-Los 3,7,9	LPS	++ on the wild type strain (-) on the "D" and "R" mutants
Anti-Los 8 (NIBSC)	LPS	+/- on the wild type strain (-) on the "D" and "R" mutants
Anti-P1.2 (NIBSC)	Anti-Porin A Sero-subtype 1.2	(-) on all wild type and mutants strains

5

Example 2: Construction of versatile gene delivery vectors (the pCMK series) targeting integration in the *porA* locus of *Neisseria meningitidis*.

10 A plasmid allowing homologous recombination and stable integration of foreign DNA in the *porA* locus of *Neisseria meningitidis* was constructed. This delivery vector (genes, operons and/or expression cassettes) is useful for constructing *Neisseria meningitidis* strains producing recombinant, improved blebs. Typically, such a vector contains at least: (1) a plasmid backbone replicative in *E. coli* but not in
15 *Neisseria meningitidis* (a suicide plasmid), (2) at least one, but preferably two regions of homology for targeting the integration in a chromosomal locus such as *porA*, (3) Efficient transcriptional (promoter, regulatory region and terminator) and translational (optimised ribosome binding site and initiation codon) signals functional in *Neisseria meningitidis*, (4) a multiple cloning site and (5) selectable gene(s) allowing the

maintenance of the plasmid in *E. coli* and the selection of integrants in *Neisseria meningitidis*. Additional elements include, for example, uptake sequences to facilitate the entry of foreign DNA in *Neisseria meningitidis*, and counter selectable markers such as *sacB*, *rpsL*, *gltS* to enhance the frequency of double cross-over events.

5 A schematic drawing of the vector constructed in this example and designated pCMK is represented in Figure 3. Its corresponding complete nucleotide sequence is shown in SEQ. ID NO:1. pCMK derives from a pSL1180 backbone (PharmaciaBiotech, Sweden), a high copy-number plasmid replicative in *E. coli*, harbouring the *bla* gene (and thereby conferring resistance to ampicillin). In addition
10 to this, pCMK functionally contains two *porA* flanking regions (*porA5'* and *porA3'* containing a transcription terminator) necessary for homologous recombination, a selectable marker conferring resistance to kanamycin, two uptake sequences, a *porA/lacO* chimeric promoter repressed in the *E. coli* host expressing *lacI^f* but transcriptionally active in *Neisseria meningitidis*, and a multiple cloning site (5 sites
15 present: *NdeI*, *KpnI*, *NheI*, *PinA1* and *SphI*) necessary for the insertion of foreign DNA in pCMK.

pCMK was constructed as follows. The *porA5'* and *porA3'* recombinogenic regions, the *porA/lacO* promoter were PCR amplified using the oligonucleotides listed in the table below, cloned in pTOPO and sequenced. These DNA fragments were
20 successively excised from pTOPO and recloned in pSL1180. The kanamycin resistance cassette was excised from pUC4K (PharmaciaBiotech, Sweden) and was introduced between the *porA5'* flanking region and the *porA/lacO* promoter region.

Table: Oligonucleotides used in this work

Oligonucleotides	Sequence	Remark(s)
PorA5' Fwd	5'-CCC AAG CTT GCC GTC TGA ATA CAT CCC GTC ATT CCT CA-3'	<i>Hind</i> III cloning site Uptake sequence ()
PorA5' Rev	5'-CGA TGC TCG CGA CTC CAG AGA CCT CGT GCG GGC C-3'	<i>Nru</i> I cloning site
PorA3' Fwd	5'-GGA AGA TCT GAT TAA ATA GGC GAA AAT ACC AGC TAC GA-3'	<i>Bgl</i> II cloning site Stop codons ()
PorA3' Rev	5'-GCC GAA TTC TTC AGA CCG C GC AGC AGG AAT TTA TCG G-3'	<i>Eco</i> RI cloning site Uptake sequence ()
PoLa Rev1	5'- GAA TTG TTA TCC GCT CAC AAT TCC GGG CAA ACA CCC GAT AC-3'	
PoLa Rev2	5'-GAA TTC CAT ATG ATC GGC TTC CTT TTG TAA ATT TGA TAA AAA CCT AAA AAC ATC GAA TTG TTA TCC GCT C-3'	<i>Nde</i> I cloning site
PorAlacO Fwd	5'-AAG CTC TGC AGG AGG TCT GCG CTT GAA TTG-3'	<i>Pst</i> I cloning site
PorAlacO Rev	5'-CTT AAG GCA TAT GGG CTT CCT TTT GTA A-3'	<i>Nde</i> I cloning site
PPA1	5'-CGG GCC GTT GCC GAT GTC AGC C-3'	
PPA2	5'-GGC ATA GCT GAT GCG TGG AAC TGC-3'	
N-full-01:	5'-GGG AAT TCC ATA TGA AAA AAG CAC TTG CCA CAC-3'	<i>Nde</i> I cloning site
Nde-NspA-3:	5'- GGA ATT CCA TAT GTC AGA ATT TGA CGC GCA C -3'	<i>Nde</i> I cloning site
PNS1	5'- CCG CGA ATT CGG AAC CGA ACA CGC CGT TCG-3'	<i>Eco</i> RI cloning site
PNS1	5'-CGT CTA GAC GTA GCG GTA TCC GGC TGC -3'	<i>Xba</i> I cloning site
PromD15-S1X	5'- GGG CGA ATT CGC GGC CGC CGT CAA CGG CAC ACC CGT TG-3'	<i>Eco</i> RI and <i>Nru</i> I cloning sites
PromD15-S2	5'- GCT CTA GAG CGG AAT GCG GTT TCA GAC G- 3'	<i>Xba</i> I cloning site
PNS4	5'- AGC TTT ATT TAA ATC CTT AAT TAA CGC GTC CGG AAA ATA TGC TTA TC_34	<i>Swa</i> I and <i>Pac</i> I cloning sites
PNS5	5'- AGC TTT GTT TAA ACC CTG TTC CGC TGC TTC GGC-3'	<i>Pme</i> I cloning site
D15-S4	5'- GTC CGC ATT TAA ATC CTT AAT TAA GCA GCC GGA CAG GGC GTG G-3'	<i>Swa</i> I and <i>Pac</i> I cloning sites
D15-S5	5'- AGC TTT GTT TAA AGG ATC AGG GTG TGG TCG GGC-3'	<i>Pme</i> I cloning site

Example 3: Construction of a *Neisseria meningitidis* serogroup B strain lacking both capsular polysaccharides and the major immunodominant antigen PorA.

Modulating the antigenic content of outer membrane blebs may be advantageous in improving their safety and efficacy in their use in vaccines, or diagnostic or therapeutic uses. Components such as the *Neisseria meningitidis* serogroup B capsular polysaccharides should be removed to exclude the risk of inducing autoimmunity (see example 1). Similarly, it is beneficial to suppress the immunodominance of major outer-membrane antigens such as PorA, which induce strain-specific bactericidal antibodies but fail to confer cross-protection. To illustrate such an approach, we used the pCMK(+) vector to construct a *Neisseria meningitidis* serogroup B strain lacking both capsular polysaccharides and the immunodominant PorA outer membrane protein antigen. For this purpose, a deletion of the *porA* gene was introduced in the H44/76 *cps*- strain, described in example 1 by means of homologous recombination.

The H44/76 *cps*- strain was prepared competent and transformed with two 2µg of supercoiled pCMK(+) plasmid DNA as described previously. Aliquot fractions of the transformation mixture (100µl) were plated on Mueller-Hinton plates supplemented with Kanamycin (200µg/ml) and incubated at 37°C for 24 to 48 hours. Kanamycin-resistant colonies were selected, restreaked on MH-Kn and grown for an additional 24 hours at 37°C. At that stage half of the bacterial culture was used to prepare glycerol stocks (15 % vol./vol.) and was kept frozen at -70°C. Another fraction (estimated to be 10⁸ bacteria) was resuspended in 15 µl of distilled water, boiled ten minutes and used as a template for PCR screening. Two *porA* internal primers named , PPA1 and PPA2, were synthesized and used to perform PCR amplification on boiled bacterial lysates in the conditions described by the supplier (HiFi DNA polymerase, Boehringer Mannheim, GmbH). The thermal cycling used was the following: 25 times (94°C 1min., 52°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). Since a double crossing-over between pCMK DNA and the chromosomal *porA* locus deletes the region required for #1 and #2 annealing, clones lacking a 1170bp PCR amplification fragment were selected as *porA* deletion mutants. These PCR results were further confirmed by analyzing in parallel, the presence of PorA in the corresponding bacterial protein extracts. For that purpose,

another aliquot of bacteria (estimated to be 5.10^8 bacteria) was re-suspended in 50 μ l of PAGE-SDS buffer (SDS 5%, Glycerol 30%, Beta-mercaptoethanol 15%, Bromophenol blue 0.3mg/ml, Tris-HCl 250 mM pH6.8), boiled (100°C) frozen (-20°C) / boiled (100°C) three times and was separated by PAGE-SDS electrophoresis on a 12.5 % gel. Gels were then stained by Coomassie Brilliant blue R250 or transferred to a nitrocellulose membrane and probed with an anti-PorA monoclonal antibody as described in Maniatis *et al.* As represented in Figure 4, both Coomassie and immunoblot staining confirmed that *porA* PCR negative clones do not produce detectable levels of PorA. This result confirm that the pCMK vector is functional and can be used successfully to target DNA insertion in the *porA* gene, abolishing concomitantly the production of the PorA outer membrane protein antigen.

Example 4: Up-regulation of the NspA outer membrane protein production in blebs derived from a recombinant *Neisseria meningitidis* serogroup B strain lacking functional *porA* and *cps* genes.

Enriching bleb vesicles with protective antigens is advantageous for improving the efficiency and the coverage of outer membrane protein-based vaccines. In that context, recombinant *Neisseria meningitidis* strains lacking functional *cps* and *porA* genes were engineered so that the expressions level of the outer-membrane protein NspA was up-regulated. For that purpose, the gene coding for NspA was PCR amplified using the N01-full-*NdeI* and *NdeI*-3' oligonucleotide primers (see table in example 2). The conditions used for PCR amplification were those described by the supplier (HiFi DNA polymerase, Boehringer Mannheim, GmbH). Thermal cycling done was the following: 25 times (94°C 1min., 52°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). The corresponding amplicon was digested with *NdeI* and inserted in the *NdeI* restriction site of the pCMK(+) delivery vector. Insert orientation was checked and recombinant plasmids, designed pCMK(+)-NspA, were purified at a large scale using the QIAGEN maxiprep kit and 2 μ g of this material was used to transform a *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes (strain described in example 1). Integration resulting from a double crossing-over between the pCMK(+)-NspA vector and the chromosomal *porA* locus were

selected using a combination of PCR and Western blot screening procedures presented in example 3.

Bacteria (corresponding to about $5 \cdot 10^8$ bacteria) were re-suspended in 50 μ l of PAGE-SDS buffer, frozen(-20°C) / boiled (100°C) three times and then were separated by PAGE-SDS electrophoresis on a 12.5 % gel. Gels were then stained by Coomassie Brilliant blue R250 or transferred to a nitrocellulose membrane and probed with an anti-NspA polyclonal serum. Both Coomassie (data not shown) and immunoblot staining (see figure 4) confirmed that *porA* PCR negative clones do not produce detectable levels of PorA. The expression of NspA was examined in Whole-cell bacterial lysates (WCBL) or outer-membrane bleb preparations derived from NmB [*cps*-, *porA*-] or NmB [*cps*-, *porA*-, *Nspa*+]. Although no difference was observable by Coomassie staining, immunoblotting with the anti-NspA polyclonal serum detected a 3-5 fold increased in the expression of NspA (with respect to the endogenous NspA level), both in WCBL and outer-membrane bleb preparations (see figure 5). This result confirm that the pCMK(+)-NspA vector is functional and can be used successfully to up-regulate the expression of outer membrane proteins such as NspA, abolishing concomitantly the production of the PorA outer membrane protein antigen.

Example 5: Up-regulation of the D15/Omp85 outer membrane protein antigen in blebs derived from a recombinant *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes but expressing PorA.

Certain geographically isolated human populations (such as Cuba) are infected by a limited number of *Neisseria meningitidis* isolates belonging largely to one or few outer membrane protein serotypes. Since PorA is a major outer-membrane protein antigen inducing protective and strain-specific bactericidal antibodies, it is then possible to confer vaccine protection using a limited number of *porA* serotypes in a vaccine. In such a context, the presence of PorA in outer membrane vesicles may be advantageous, strengthening the vaccine efficacy of such recombinant improved blebs. Such PorA containing vaccines, however, can be improved still further by increasing the level of other cross-reactive OMPs such as omp85/D15.

In the following example, the pCMK(+) vector was used to up-regulate the expression of the Omp85/D15 outer membrane protein antigen in a strain lacking functional *cps* genes but expressing *porA*. For that purpose, the gene coding for Omp85/D15 was PCR amplified using the D15-*NdeI* and D15-*NotI* oligonucleotide primers. The conditions used for PCR amplification were those described by the supplier (HiFi DNA polymerase, Boehringer Mannheim, GmbH). Thermal cycling done was the following: 25 times (94°C 1min., 52°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). The corresponding amplicon was inserted in the pTOPO cloning vector according to the manufacturer's specifications and confirmatory sequencing was performed. This Omp85/D15 DNA fragment was excised from pTOPO by restriction hydrolysis using *NdeI/NsiI* and subsequently cloned in the corresponding restriction sites of the pCMK(+) delivery vector. Recombinant plasmids, designed pCMK(+)-D15 were purified on a large scale using the QIAGEN maxiprep kit and 2 µg of this material was used to transform a *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes (strain described in example 1). In order to preserve the expression of *porA*, integration resulting from a single crossing-over (either in Omp85/D15 or in *porA*) were selected by a combination of PCR and Western blot screening procedures. Kanamycin resistant clones testing positive by *porA*-specific PCR and western blot were stored at -70°C as glycerol stocks and used for further studies.

Bacteria (corresponding to about $5 \cdot 10^8$ bacteria) were re-suspended in 50 µl of PAGE-SDS buffer, frozen(-20°C) / boiled (100°C) three times and then were separated by PAGE-SDS electrophoresis on a 12.5 % gel. Gels were then stained by Coomassie Brilliant blue R250 or transferred to a nitrocellulose membrane and probed with an anti-*porA* monoclonal antibody. As represented in Figure 6, both Coomassie and immunoblot staining confirmed that *porA* PCR positive clones produce PorA.

The expression of D15 was examined using outer-membrane bleb preparations derived from NmB [*cps*-, *porA*-] or NmB [*cps*-, *porA*+, D15+]. Coomassie detected a significant increase in the expression of D15 (with respect to the endogenous D15 level), preparations (see Figure 6). This result confirmed that the pCMK(+)-D15 vector is functional and can be used successfully to up-regulate the expression of outer membrane proteins such as D15, without abolishing the production of the major PorA outer membrane protein antigen.

Example 6: Construction of versatile promoter delivery vectors

Rational: The rational of this approach is represented in Figure 7 and can be summarized in 7 essential steps. Some of these steps are illustrated below with the construction of Vector for up-regulating the expression of NspA and D15/Omp85.

Vector for up-regulating the expression of the NspA gene.

Step 1. A DNA region (997bp) located upstream from the NspA coding gene was discovered (SEQ. ID NO:2) in the private Incyte PathoSeq data base containing unfinished genomic DNA sequences of the *Neisseria meningitidis* strain ATCC 13090. Two oligonucleotide primers referred to as PNS1 and PNS2 (see table in example 2) were designed using this sequence and synthesized. These primers were used for PCR amplification using genomic DNA extracted from the H44/76 strain.

Step 2. The corresponding amplicons were cleaned-up using the Wizard PCR kit (Promega, USA) and submitted to digestion with the *EcoRI/XbaI* restriction enzymes for 24 hours using the conditions described by the supplier (Boehringer Mannheim, Germany). The corresponding DNA fragments were gel purified and inserted in the corresponding sites of the pUC18 cloning vector. **Step 3.** Recombinant plasmids were prepared on a large scale and an aliquot fraction was used as a template for inverse PCR amplification. Inverse PCR was performed using the PNS4 and PNS5 oligonucleotides using the following thermal cycling conditions: 25 times (94°C 1min., 50°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). Linearized pUC 18 vectors harbouring a deletion in the NspA upstream region insert were obtained.

Vector for up-regulating the expression of the D15/omp85 gene.

Step 1. A DNA region (1000 bp) located upstream from the D15/omp85 coding gene was discovered (SEQ. ID NO:3) in the private Incyte PathoSeq database containing unfinished genomic DNA sequences of the *Neisseria meningitidis* strain ATCC 13090. Two oligonucleotide primers referred to as PromD15-51X and PromD15-S2 (see table in example 2) were designed using this sequence and synthesized. These primers were used for PCR amplification using genomic DNA extracted from the H44/76 strain. **Step 2.** The corresponding amplicons were cleaned-up using the

Wizard PCR kit (Promega, USA) and submitted to digestion with the *EcoRI/XbaI* restriction enzymes for 24 hours in the conditions described by the supplier (Boehringer Mannheim, Germany). The corresponding DNA fragments were gel purified and inserted in the corresponding sites of the pUC18 cloning vector. **Step 3.**

5 Recombinant plasmids were prepared on a large scale and an aliquot fraction was used as a template for inverse PCR amplification. Inverse PCR was performed using the D15-S4 and D15-S5 oligonucleotides using the following thermal cycling conditions: 25 times (94°C 1min., 50°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery). Linearized pUC 18 vectors harbouring a deletion in the D15/omp85
10 upstream region insert were obtained.

Example 7: Fermentation processes for producing recombinant blebs.

The examples listed below describe methods for producing recombinant blebs lacking
15 either capsular polysaccharides or capsular polysaccharides and PorA. Such a procedure may be used for a wide range of *Neisseria meningitidis* recombinant strains and may be adapted over an extended scale range.

Culture media: *Neisseria meningitidis* serogroup B strains were propagated in solid
20 (FNE 004 AA, FNE 010 AA) or liquid (FNE 008 AA) culture media. These new media for growing meningococcus are advantageously free of animal products, and are considered a further aspect of the invention.

Components	FNE 004 AA	FNE 008 AA	FNE 010 AA
Agar	18 g/L	-	18 g/L
NaCl	6 g/L	6 g/L	6 g/L
Na-Glutamate	-	1.52 g/L	-
NaH ₂ PO ₄ .2H ₂ O	2.2 g/L	2.2 g/L	2.2 g/L
KCl	0.09 g/L	0.09 g/L	0.09 g/L
NH ₄ Cl	1.25 g/L	1.25 g/L	1.25 g/L
Glucose	5 g/L	20 g/L	5 g/L
Yeast Extract UF	-	2.5 g/L	-
Soy Pepton	5 g/L	30 g/L	5 g/L
CaCl ₂ .2H ₂ O	0.015 g/L	-	0.015 g/L
MgSO ₄ .7H ₂ O	0.6 g/L	0.6 g/L	0.6 g/L
Erythromycine :	0.015 g/L	-	-
Kanamycine	-	-	0.2 g/L

5 Flask cultivation of *Neisseria meningitidis* serogroup B cps- recombinant blebs: This was performed in two steps comprising preculture on solid medium followed by liquid cultivation. **Solid pre-culture** A vial of seed was removed from freezer (-80°C), thawed to room temperature and 0.1 mL was streaked into a Petri dish containing 15 mL of FNE004AA (see above). The Petri dish was incubated at 37°C for 18 ± 2 hours. The surface growth was resuspended in 8 mL of FNE008AA (see above) supplemented with 15 mg/L of erythromycin. **Flask culture.** 2 mL of resuspended solid pre-culture were added to a 2 litre flask containing 400 mL of FNE008AA supplemented with 15 mg/L of erythromycin. The flask was placed on a shaking table (200 rpm) and incubated at 37°C for 16 ± 2 hours. The cells were separated from the culture broth by centrifugation at 5000g at 4°C for 15 minutes.

15 Batch mode cultivation of *Neisseria meningitidis* serogroup B cps- recombinant blebs: This was performed in three steps comprising preculture on solid medium, liquid cultivation and batch mode cultivation. **Solid pre-culture.** A vial of seed was removed from freezer (-80°C), thawed to room temperature and 0.1 mL was streaked into a Petri dish containing 15 mL of FNE004AA (see above). The Petri dish was incubated at 37°C for 18 ± 2 hours. The surface growth was resuspended in 8 mL of

FNE008AA (see above) supplemented with 15 mg/L of erythromycin. **Liquid pre-culture.** 2 mL of resuspended solid pre-culture were added to one 2 liters flask containing 400 mL of FNE008AA supplemented with 15 mg/L of erythromycin. The flask was placed on a shaking table (200 rpm) and incubated at 37°C for 16 ± 2 hours.

5 The content of the flask was used to inoculate the 20 liters fermenter. **Batch mode culture in fermenter.** The inoculum (400 mL) was added to a pre-sterilized 20 liters (total volume) fermenter containing 10 L of FNE008AA supplemented with 15 mg/L of erythromycin. The pH was adjusted to and maintained at 7.0 by the automated addition of NaOH (25% w/v) and H₃PO₄ (25% v/v). The temperature was regulated
10 at 37°C. The aeration rate was maintained at 20 L of air / min and the dissolved oxygen concentration was maintained at 20% of saturation by the agitation speed control. The overpressure in the fermenter was maintained at 300 g/cm². After 9 ± 1 hours, the culture was in stationary phase. The cells were separated from the culture broth by centrifugation at 5000g at 4°C for 15 minutes.

15

Flask cultivation of *Neisseria meningitidis* serogroup B cps-, PorA- recombinant blebs: This was performed in two steps comprising preculture on solid medium followed by liquid cultivation. **Solid pre-culture.** A vial of seed was removed from freezer (-80°C), thawed to room temperature and 0.1 mL was streaked into a Petri dish containing 15 mL of FNE010AA (see above). The Petri dish was incubated at 37°C
20 for 18 ± 2 hours. The surface growth was resuspended in 8 mL of FNE008AA (see above) supplemented with 200 mg/L of kanamycin. **Flask culture.** 2 mL of resuspended solid pre-culture were added to a 2 litre flask containing 400 mL of FNE008AA supplemented with 200 mg/L of kanamycin. The flask was placed on a
25 shaking table (200 rpm) and incubated at 37°C for 16 ± 2 hours. The cells were separated from the culture broth by centrifugation at 5000g at 4°C for 15 minutes.

Example 8: Isolation and purification of blebs from meningococci devoid of capsular polysaccharide

30

Recombinant blebs were purified as described below. The cell paste (42gr) was suspended in 211 ml of 0.1M Tris-Cl buffer pH 8.6 containing 10 mM EDTA and 0.5% Sodium Deoxycholate (DOC). The ratio of buffer to biomass was 5/1 (V/W).

The biomass was extracted by magnetic stirring for 30 minutes at room temperature. Total extract was then centrifuged at 20,000g for 30 minutes at 4°C (13,000 rpm in a JA-20 rotor, Beckman J2-HS centrifuge). The pellet was discarded. The supernatant was ultracentrifuged at 125,000g for 2 hours at 4°C (40,000 rpm in a 50.2Ti rotor, 5 Beckman L8-70M ultracentrifuge) in order to concentrate vesicles. The supernatant was discarded. The pellet was gently suspended in 25 ml of 50 mM Tris-Cl buffer pH 8.6 containing 2 mM EDTA, 1.2% DOC and 20% sucrose. After a second ultracentrifugation step at 125,000g for 2 hours at 4°C, vesicles were gently suspended in 44 ml of 3% sucrose and stored at 4°C. All solutions used for bleb extraction and 10 purification contained 0.01% thiomersalate. As illustrated in Figure 8, this procedure yields protein preparations highly enriched in outer-membrane proteins such as PorA and PorB.

Example 9: Identification of bacterial promoters suitable for up-regulation antigens-
15 **coding genes**

The use of strong bacterial promoter elements is essential to obtain up-regulation of genes coding for outer membrane proteins. In that context, we have shown previously that up-regulating the *Neisseria meningitidis nspA*, *hsf*, and *omp85* genes using the *porA* promoter has allowed us to isolate recombinant blebs enriched in 20 the corresponding NspA, Hsf and Omp85 proteins. Alternatives to the *porA* promoter may be useful to obtain different levels of up-regulation, to overcome potential *porA* phase variation and/or to achieve conditional gene expression (iron-regulated promoters). Here we describe a method allowing the identification of a precise transcriptional start site of strong promoter elements likely to confer high level of 25 expression in bacteria. Since promoter regulatory elements are classically encompassed within 200 bp upstream and 50bp downstream from the +1 site (Collado-Vides J, Magasanik B, Gralla JD, 1991, *Microbiol Rev* 55(3):371-94), the result of such an experiment allows us to identify DNA fragments of about 250 bp carrying strong promoter activities. Major outer membrane proteins such as *Neisseria* 30 *meningitidis* PorA, PorB & Rmp, *Haemophilus influenzae* P1, P2, P5 & P6, *Moraxella catarrhalis* OmpCD, OmpE, as well as some cytoplasmic and/or iron regulated proteins of these bacteria possess strong promoter elements. As a validation

of this general methodology, we mapped the transcriptional start site of the strong *Neisseria meningitidis* *porA* and *porB* promoters using rapid amplification of cDNA elements (5' RACE).

The principles of 5' RACE are the following: 1) Total RNA extraction using
5 QIAGEN "RNeasy" Kit. Genomic DNA removing by DNase treatment followed by
QIAGEN purification; 2) mRNA reverse transcription with a *porA* specific 3' end
primer (named porA3). Expected cDNA size: 307 nt. RNA removing by alkaline
hydrolysis; 3) Ligation of a single-stranded DNA oligo anchor (named DT88) to the 3'
10 end of the cDNA using T4 RNA ligase. Expected product size: 335 nt. Amplification
of the anchor-ligated cDNA using a combination of hemi-nested PCR; 4) PCR
amplification of the anchor-ligated cDNA using a complementary-sequence anchor
primer as the 5' end primer (named DT89) and a 3'end primer (named p1-2) which is
internal to the 3'end RT primer porA3. Expected product size: 292 bp; 5) PCR
15 amplification of previous PCR products using DT89 as 5'end primer and p1-1 as 3'end
primer (internal to p1-2). Expected product size: 211bp; and 6) Sequencing with p1-1
primer (expected products size can be calculated because *porA* transcription start site
is known: 59 nt before the "ATG" translation start site).

Experimental procedure

20 Total RNA was extracted from approximately 10^9 cells of *Neisseria meningitidis*
serogroup B *cps- porA+* strain. Extraction of 1 ml of a liquid culture at appropriate optical
density ($OD_{600} = 1$) was performed by the QIAGEN "RNAeasy" kit according to the
manufacturer's instructions. Chromosomal DNA was removed by addition of 10U of RNase-
free DNase (Roche Diagnostics, Mannheim, Germany) to the 30 μ l of eluted RNA and was
25 incubated at 37°C for 15 min. The DNA-free RNA was purified with the same QIAGEN kit
according to instructions.

Reverse transcription reactions were performed using primer porA3 and 200U of
SUPERScript II reverse transcriptase (Life Technologies). The RT reactions were performed in
a 50 μ l volume containing: 5 μ l of 2mM dNTP, 20 pmol of porA3 pimer, 5 μ l of 10X
30 SUPERScript II buffer, 9 μ l of 25mM MgCl₂, 4 μ l of 0.1M DTT, 40U of recombinant
ribonuclease inhibitor and 1 μ g of total RNA. The porA3 primer was annealed stepwise
(70°C for 2 min, 65°C for 1 min, 60°C for 1 min, 55°C for 1 min, 50°C for 1 min, and 45°C
for 1 min) before the SUPERScript II was added. The RT reaction was performed at 42°C for

30 min, followed by 5 cycles (50°C for 1 min, 53°C for 1 min and 56°C for 1 min) to destabilize RNA secondary structure. Two parallel reactions were performed with the reverse transcriptase omitted from one reaction as negative control.

5 The RNA was removed by alkaline hydrolysis cleavage with the addition of 1 µl of 0.5M EDTA followed by 12.5 µl of 0.2 M NaOH before incubation at 68°C for 5 min. The reactions were neutralized by adding 12.5 µl of 1 M Tris-HCl (pH7.4) and precipitated by the addition of 20 µg of glycogen (Roche Molecular Biochemicals, Mannheim, Germany), 5 µl of 3 M sodium acetate and 60 µl of isopropanol. Both samples were resuspended in 20 µl of 10:1 TE (10 mM Tris-HCl, pH 7.4; 1 mM EDTA, pH8).

10 T4 RNA ligase was used to anchor a 5'-phosphorylated, 3'end ddCTP-blocked anchor oligonucleotide DT88 (see table below). Two parallel ligations were performed overnight at room temperature with each containing: 1.3 µl of 10X RNA ligase buffer (Roche Molecular Biochemicals), 0.4 µM DT88, 10 µl of either cDNA or RT control sample and 3 U of T4 RNA ligase. As negative controls, a second set of ligations reactions was performed, omitting
15 the T4 RNA ligase. The resulting ligation-reaction mixtures were used directly without purification in the subsequent PCR.

The anchor-ligated cDNA was amplified using a combination of hemi-nested and hot-started PCR approaches to increase specificity and product yield. Four separate first-round PCR were performed on the RT/ligase reaction and controls in a 30 µl volume, each
20 containing: 3 µl of 10X Taq Platinum buffer, 3µl of 25 mM MgCl₂, 1 µl of 10mM dNTP, 10 pmol of each primers and 1 µl of corresponding RNA ligation reaction. The PCR were hot started by the use of Taq Platinum (Life Technologies) DNA polymerase (2U added). The first ligation-anchored PCR (LA-PCR) was performed using 10 pmol of both the anchor-specific primer DT89 and the transcript-specific primer p1-2 (see table below) which is
25 internal to the 3' end RT primer porA3. The PCR was performed using an initial 95°C for a 5 min step (for DNA polymerase activation) followed by 10 cycles at 95°C for 10 s and 70°C for 1 min (reducing one degree per cycle), 15 cycles at 95°C for 10 s and 60°C for 1 min. The second hemi-nested LA-PCR was performed under the same conditions using primer DT89 and the p1-2 internal primer, together with 10 pmol of p1-1 (see table below) and 1 µl of
30 first-round PCR. Amplification products were purified using the QIAGEN "QIAquick PCR purification" kit according to manufacturer instructions before submitted to sequencing.

The CEQ™ Dye Terminator Cycle Sequencing kit (Beckman, France) was used to sequence the RACE PCR products using 10 pmol of primer p1-1. Sequencing reactions were performed according to the provided instructions and sequencing products were analyzed by
35 the Ceq2000 DNA Analysis System (Beckman-Coulter).

DT88	5' GAAGAGAAGGTGGAAATGGCGTTTGGC 3'
DT89	5' CAAAACGCCATTTCCACCTTCTCTC 3'
porA3	5' CCAAATCCTCGCTCCCCTTAAAGCC 3'
p1-2	5' CGCTGATTTTCGTCCTGATGCGGC 3'
p1-1	5' GGTC AATTGCGCCTGGATGTTCTG 3'

Results for the *Neisseria meningitidis* porA promoter

The start of transcription for *Neisseria meningitidis* serogroup B (strain H44/76) *porA*-mRNA was mapped 59 bp upstream of the ATG start codon using the described 5'-RACE procedure. This result confirms the mapping performed by primer extension and published by van der Ende *et al* (1995). This result supports that a DNA fragment containing nucleotides -9 to -259 with regard to the *porA* ATG is suitable for driving strong gene expression in *Neisseria meningitidis* and possibly in other bacterial species such as *Haemophilus*, *Moraxella*, *Pseudomonas*.

Results for the *Neisseria meningitidis* porB promoter

The same experimental strategy has been applied for *Neisseria meningitidis* serogroup B (strain H44/76) *porB* transcription start site mapping. Primers listed in the table below correspond to 3' end RT primer (porB3), transcript-specific primer that is internal to the porB3 (porB2) and internal to the porB2 (porB1). porB3, porB2 and porB1 are respectively located 265 bp, 195 bp and 150 bp downstream the ATG start codon.

porB1	5' GGTAGCGGTGTAACTCAGTAACTT 3'
porB2	5' GTCTTCTTGGCCTTGAAGCCGATT 3'
porB3	5' GGAGTCAGTACCGGCGATAGATGCT 3'

Using porB1 and DT89 primers a ~200 bp PCR amplicon was obtained by performing 5' - RACE mapping. Since porB1 is located 150 bp from the *porB* ATG start codon, this result supports that the *porB* transcriptional start site is located about 50 bp (+/- 30 bp) upstream of the *porB* ATG.

The exact nucleotide corresponding to transcription initiation is presently being determined by DNA sequencing. The above PCR result supports that a DNA fragment containing nucleotides -1 to -250 with regard to the *porB* ATG start codon is suitable for driving strong gene expression in *Neisseria meningitidis* and possibly in other bacterial species such as *Haemophilus*, *Moraxella*, *Pseudomonas*.

Example 10: Up-regulation of the *N. meningitidis* serogroup B *Omp85* gene by promoter replacement.

The aim of the experiment was to replace the endogenous promoter region of the D15/*Omp85* gene by the strong *porA* promoter in order to up-regulate the production of the D15/*Omp85* antigen. For that purpose, a promoter replacement plasmid was constructed using *E. coli* cloning methodologies. A DNA region (1000 bp) located upstream from the *D15/omp85* coding gene was discovered (SEQ ID NO:3) in the private Incyte PathoSeq data base containing unfinished genomic DNA sequences of the *Neisseria meningitidis* strain ATCC 13090. The main steps of this procedure are represented in Figure 9. Briefly, a DNA fragment (1000bp) covering nucleotides -48 to -983 with respect to the *D15/Omp85* gene start codon (ATG) was PCR amplified using oligonucleotides **ProD15-51X** (5'-GGG CGA ATT CGC GGC CGC CGT CAA CGG CAC ACC GTT G-3') and **ProD15-52** (5'-GCT CTA GAG CGG AAT GCG GTT TCA GAC G-3') containing *EcoRI* and *XbaI* restriction sites (underlined) respectively. This fragment was submitted to restriction and inserted in pUC18 plasmid restricted with the same enzymes. The construct obtained was submitted to *in vitro* mutagenesis using the Genome Priming system (using the pGPS2 donor plasmid) commercialized by New England Biolabs (MA, USA). Clones having inserted a mini-transposon (derived from Tn7 and harboring a chloramphenicol resistance gene) were selected. One clone containing a mini-transposon insertion located in the *D15/Omp85* 5' flanking region, 401 bp downstream from the *EcoRI* site was isolated and used for further studies. This plasmid was submitted to circle PCR mutagenesis (Jones & Winistofer (1992), Biotechniques12: 528-534) in order to (i) delete a repeated DNA sequence (Tn7R) generated by the transposition process, (ii) insert meningococcal uptake sequences required for transformation, and (iii) insert suitable restriction sites allowing cloning of foreign DNA material such as promoters. The circle PCR was performed using the **TnRD15-KpnI/XbaI + US** (5'-CGC CGG TAC CTC TAG AGC CGT CTG AAC CAC TCG TGG ACA ACC C-3') & **TnR03Cam(KpnI)** (5'-CGC CGG TAC CGC CGC TAA CTA TAA CGG TC-3') oligonucleotides containing uptake sequences and suitable restriction sites (*KpnI* and *XbaI*) underlined. The resulting PCR fragment was gel-purified, digested with *Asp718* (isoschizomer of *KpnI*) and ligated to a 184bp DNA fragment containing the *porA* promoter and generated by PCR using the

PorA-01 (5'-CGC CGG TAC CGA GGT CTG CGC TTG AAT TGT G-3') and **PorA02** (5'-CGC CGG TAC CTC TAG ACA TCG GGC AAA CAC CCG-3') oligonucleotides containing *KpnI* restriction sites. Recombinant clones carrying a *porA* promoter inserted in the correct orientation (transcription proceeding in the *EcoRI* to *XbaI* direction) were selected and used to transform a strain of *Neisseria meningitidis* serogroup B lacking capsular polysaccharide (*cps*-) and one of the major outer membrane proteins - PorA (*porA*-). Recombinant *Neisseria meningitidis* clones resulting from a double crossing over event (PCR screening using oligonucleotides **Cam-05** (5'-GTA CTG CGA TGA GTG GCA GG-3') & **proD15-52**) were selected on GC medium containing 5µg/ml chloramphenicol and analyzed for D15/Omp85 expression. As represented in Figure 10, the production of D15/Omp85 was significantly increased in the total protein extracts of *Nm* strains resulting from promoter replacement, when compared to parental strain (*cps*-). This result was also observed when analyzing outer-membrane blebs prepared from the same strains (see Figure 17). These results are attributable to the replacement of the endogenous *D15* promoter by the strong *porA* promoter. In addition, it was surprisingly found that expression, where the *porA* promoter was introduced approximately 400 bp upstream of the initiator codon, was approximately 50 times greater than when the promoter was placed approximately 100 bp upstream. Altogether, these experiments support that the promoter replacement strategy works and allows the up-regulation of the synthesis of integral outer-membrane proteins in outer-membrane blebs.

Certain geographically isolated human populations (such as Cuba) are infected by a limited number of *Neisseria meningitidis* isolates belonging largely to one or few outer membrane protein serotypes. Since PorA is a major outer-membrane protein antigen which can induce protective and strain-specific bactericidal antibodies, it may be possible to confer vaccine protection in such a population using a limited number of *porA* serotypes. Moreover, PorA may interact with or stabilize some other outer membrane proteins. In this context, the presence of PorA in outer membrane vesicles may be advantageous, strengthening the vaccine efficacy of such recombinant improved blebs.

For such a reason, it may be desirable to up-regulate the expression of D15/Omp85 outer membrane protein in a *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes but expressing PorA. Genomic DNA was extracted from

the recombinant *Neisseria meningitidis* serogroup B *cps*-, *porA*-, D15/Omp85+ strain using the QIAGEN Genomic Tips 100-G kit. 10µgr of this material was linearized and used to transform *Neisseria meningitidis* serogroup B *cps*- following a classical transformation protocol. Recombinant *Neisseria* were obtained on GC agar plates containing 5µgr/ml chloramphenicol.

Integrations resulting from a double crossing-over upstream of the *D15* gene were screened by PCR as described previously. As homologous recombinations can occur everywhere in the chromosome, a second PCR screening was performed to control the integrity of the *porA* locus in the recombinant strain. For this purpose, internal *porA* primers PPA1 (5- GCG GCC GTT GCC GAT GTC AGC C -3') and PPA2 (5- GGC ATA GCT GAT GCG TGG AAC TGC -3') were used in a PCR screening experiment. The amplification of an 1170bp fragment confirms the presence of the *porA* gene in the recombinant bacteria.

Recombinant bacteria (corresponding to about 5.10^8 bacteria) can be resuspended in 50 µl of PAGE-SDS buffer, frozen(-20°C) / boiled (100°C) three times and then separated by PAGE-SDS electrophoresis on a 12.5 % gel. Gels can then be stained by Coomassie Brilliant blue R250 or transferred to a nitrocellulose membrane and probed either with an anti-*porA* monoclonal antibody or with an anti-D15/Omp85 rabbit polyclonal antibody. Analysis of outer-membrane blebs prepared from the same strains can also be performed.

Example 11: Up-regulation of the Hsf protein antigen in a recombinant *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes but expressing *PorA*.

As described above, in certain countries, the presence of *PorA* in outer membrane vesicles may be advantageous, and can strengthen the vaccine efficacy of recombinant improved blebs. In the following example, we have used a modified pCMK(+) vector to up-regulate the expression of the Hsf protein antigen in a strain lacking functional *cps* genes but expressing *PorA*. The original pCMK(+) vector contains a chimeric *porA/lacO* promoter repressed in *E. coli* host expressing *lacI^f* but transcriptionally active in *Neisseria meningitidis*. In the modified pCMK(+), the native *porA* promoter was used to drive the transcription of the *hsf* gene. The gene coding for Hsf was PCR amplified using the HSF 01-*NdeI* and HSF 02-*NheI* oligonucleotide primers, presented in the table below. Because of the sequence of the

HSF 01-*NdeI* primer the Hsf protein expressed will contain two methionine residues at the 5' end. The conditions used for PCR amplification were those described by the supplier (HiFi DNA polymerase, Boehringer Mannheim, GmbH). Thermal cycling was the following: 25 times (94°C 1min., 48°C 1min., 72°C 3min.) and 1 time (72°C

5 10min., 4°C up to recovery). The corresponding amplicon was subsequently cloned in the corresponding restriction sites of pCMK(+) delivery vector. In this recombinant plasmid, designed pCMK(+)-Hsf, we deleted the *lacO* present in the chimeric *porA/lacO* promoter by a recombinant PCR strategy (See Figure 12). The pCMK(+)-Hsf plasmid was used as a template to PCR amplify 2 separate DNA fragments:

10 -fragment 1 contains the *porA* 5' recombinogenic region, the Kanamycin resistance gene and the *porA* promoter. Oligonucleotide primers used, RP1(*SacII*) and RP2, are presented in the table below. RP1 primer is homologous to the sequence just upstream of the *lac* operator.

15 -fragment 2 contains the Shine-Dalgarno sequence from the *porA* gene, the *hsf* gene and the *porA* 3' recombinogenic region. Oligonucleotide primers used, RP3 and RP4(*ApaI*), are presented in the table below. RP3 primer is homologous to the sequence just downstream of the *lac* operator. The 3' end of fragment 1 and the 5' end of fragment 2 have 48 bases overlapping. 500ng of each PCR (1 and 2) were used for a final PCR reaction using primers RP1 and RP4. The final amplicon obtained was

20 subcloned in pSL1180 vector restricted with *SacII* and *ApaI*. The modified plasmid pCMK(+)-Hsf was purified at a large scale using the QIAGEN maxiprep kit and 2 µg of this material was used to transform a *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes (the strain described in example 1). In order to preserve the expression of *porA*, integration resulting from a single crossing-over was selected

25 by a combination of PCR and Western blot screening procedures. Kanamycin resistant clones testing positive by *porA*-specific PCR and western blot were stored at -70°C as glycerol stocks and used for further studies. Bacteria (corresponding to about 5.10^8 bacteria) were re-suspended in 50 µl of PAGE-SDS buffer, frozen (-20°C) / boiled (100°C) three times and then were separated by PAGE-SDS electrophoresis on a 12.5

30 % gel. The expression of Hsf was examined in Whole-cell bacterial lysates (WCBL) derived from *NmB* [Cps-, PorA+] or *NmB* [Cps-, PorA+, Hsf+]. Coomassie staining detected a significant increase in the expression of Hsf (with respect to the endogenous Hsf level) (See in Figure 13). This result confirms that the modified

pCMK(+)-Hsf vector is functional and can be used successfully to up-regulate the expression of outer membrane proteins, without abolishing the production of the major PorA outer membrane protein antigen.

5 Oligonucleotides used in this work

Oligonucleotides	Sequence	Remark(s)
Hsf 01-Nde	5'- GGA ATT CCA TAT GAT GAA CAA AAT ATA CCG C-3'	<i>Nde</i> I cloning site
Hsf 02-Nhe	5'-GTA GCT AGC TAG CTT ACC ACT GAT AAC CGA C-3'	<i>Nhe</i> I cloning site
GFP-mut-Asn	5'-AAC TGC AGA ATT AAT ATG AAA GGA GAA GAA CTT TTC-3'	<i>Asn</i> I cloning site Compatible with <i>Nde</i> I
GFP-Spe	5'-GAC ATA CTA GTT TAT TTG TAG AGC TCA TCC ATG -3'	<i>Spe</i> I cloning site Compatible with <i>Nhe</i> I
RP1 (SacII)	5'- TCC CCG CGG GCC GTC TGA ATA CAT CCC GTC-3'	<i>Sac</i> II cloning site
RP2	5'-CAT ATG GGC TTC CTT TTG TAA ATT TGA GGG CAA ACA CCC GAT ACG TCT TCA-3'	
RP3	5'-AGA CGT ATC GGG TGT TTG CCC TCA AAT TTA CAA AAG GAA GCC CAT ATG -3'	
RP4(ApaI)	5'-GGG TAT TCC GGG CCC TTC AGA CGG CGC AGC AGG -3'	<i>Apa</i> I cloning site

Example 12: Expression of the Green Fluorescent Protein in a recombinant *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes but expressing PorA.

10 In the following example, the pCMK vector was used to test the expression of a cytoplasmic heterologous protein in *Neisseria meningitidis*. The Green Fluorescent Protein was amplified from the pKen-Gfpmut2 plasmid with the primers GFP-Asn-mut2 and GFP-Spe (see table in Example 11). *Asn*I gives cohesive ends compatible with *Nde*I, *Spe*I gives cohesive ends compatible with *Nhe*I. The conditions used for
15 PCR amplification were those described by the supplier (HiFi DNA polymerase, Boehringer Mannheim, GmbH). Thermal cycling was the following: 25 times (94°C 1min., 48°C 1min., 72°C 3min.) and 1 time (72°C 10min., 4°C up to recovery).The

corresponding amplicon was subsequently cloned in the pCMK(+) delivery vector digested with *NdeI* and *NheI* restriction enzymes. In this recombinant plasmid, designed pCMK(+)-GFP, we deleted the *lacO* present in the chimeric *porA/lacO* promoter by a recombinant PCR strategy. The pCMK(+)-GFP plasmid was used as

5 template to PCR amplify 2 separate DNA fragments:

-fragment 1 contained the *porA* 5' recombinogenic region, the Kanamycin resistance gene and the *porA* promoter. Oligonucleotide primers used, RP1(*SacII*) and RP2 (see table in example 11). RP1 primer is homologous to the sequence just upstream of the *lac* operator.

10 -fragment 2 contains the PorA Shine-Dalgarno sequence, the *gfp* gene and the *porA* 3' recombinogenic region. Oligonucleotide primers used, RP3 and RP4(*ApaI*), are presented in the table in example 11. RP3 primer is homologous to the sequence just downstream of the *lac* operator.

The 3'end of fragment 1 and the 5'end of fragment 2 have 48 bases

15 overlapping. 500ng of each PCR (1 and 2) were used for a final PCR reaction using primers RP1 and RP4. Twenty µg of this PCR fragment were used to transform a *Neisseria meningitidis* serogroup B strain lacking functional *cps* genes.

Transformation with linear DNA is less efficient than with circular plasmid DNA but all the recombinants obtained performed a double crossing-over (confirmed

20 by a combination of PCR and Western blot screening procedures). Kanamycin resistant clones were stored at -70°C as glycerol stocks and used for further studies. Bacteria (corresponding to about 5.10⁸ bacteria) were re-suspended in 50 µl of PAGE-SDS buffer, frozen (-20°C) / boiled (100°C) three times and then were separated by PAGE-SDS electrophoresis on a 12.5 % gel.

25 The expression of GFP was examined in Whole-cell bacterial lysates (WCBL) derived from *NmB* [Cps-, PorA+] or *NmB* [Cps-, PorA-, GFP+]. Coomassie staining detected an expression of GFP absent in the recipient *Neisseria meningitidis* strain (see figure 14).

30 **Example 13: Up-regulation of the *N. meningitidis* serogroup B *NspA* gene by promoter replacement**

The aim of the experiment was to replace the endogenous promoter region of the *NspA* gene by the strong *porA* promoter, in order to up-regulate the production of

the NspA antigen. For that purpose, a promoter replacement plasmid was constructed using *E. coli* cloning methodologies. A DNA region (924bp) located upstream from the *NspA* coding gene was discovered (SEQ ID NO: 7) in the private Incyte PathoSeq data base containing unfinished genomic DNA sequences of the *Neisseria meningitidis* strain ATCC 13090. A DNA fragment (675bp) covering nucleotides -115 to -790 with respect to the *NspA* gene start codon (ATG) was PCR amplified using oligonucleotides **PNS1**' (5'-CCG CGA ATT CGA CGA AGC CGC CCT CGA C-3') and **PNS2** (5'-CGT CTA GAC GTA GCG GTA TCC GGC TGC -3') containing *EcoRI* and *XbaI* restriction sites (underlined) respectively. The PCR fragment was submitted to restriction with *EcoRI* and *XbaI* and inserted in pUC18. This plasmid was submitted to circle PCR mutagenesis (Jones & Winistofer (1992), Biotechniques12: 528-534) in order to insert meningococcal uptake sequences required for transformation, and suitable restriction sites allowing cloning of a *CmR/PorA* promoter cassette. The circle PCR was performed using the **BAD01-2** (5'- GGC GCC CGG GCT CGA GCT TAT CGA TGG AAA ACG CAG C-3') & **BAD02-2** (5'-GGC GCC CGG GCT CGA GTT CAG ACG GCG CGC TTA TAT AGT GGA TTA AC -3') oligonucleotides containing uptake sequences and suitable restriction sites (*XmaI* and *XhoI*) underlined. The resulting PCR fragment was gel-purified and digested with *XhoI*. The *CmR/PorA* promoter cassette was amplified from the pUC D15/Omp85 plasmid previously described, using primers **BAD 15-2** (5'-GGC GCC CGG GCT CGA GTC TAG ACA TCG GGC AAA CAC CCG-3') & **BAD 03-2** (5'- GGC GCC CGG GCT CGA GCA CTA GTA TTA CCC TGT TAT CCC-3') oligonucleotides containing suitable restriction sites (*XmaI*, *XbaI*, *SpeI* and *XhoI*) underlined. The PCR fragment obtained was submitted to digestion and inserted in the circle PCR plasmid restricted with the corresponding enzymes. 10 µg of the recombinant plasmid were linearized and used to transform a strain of *Neisseria meningitidis* serogroup B lacking capsular polysaccharide (*cps-*) and one of the major outer membrane proteins - PorA (*porA-*). Recombinant *Neisseria meningitidis* clones resulting from a double crossing over event (PCR screening using oligonucleotides **BAD 25** (5'-GAG CGA AGC CGT CGA ACG C -3') & **BAD08** (5'-CTT AAG CGT CGG ACA TTT CC-3')) were selected on GC agar plates containing 5µg/ml chloramphenicol and analyzed for NspA expression. Recombinant bacteria (corresponding to about 5.10⁸ bacteria) were re-suspended in 50 µl of PAGE-SDS

buffer, frozen (-20°C) / boiled (100°C) three times and then were separated by PAGE-SDS electrophoresis on a 12.5 % gel. Gels were then stained by Coomassie Brilliant blue R250 or transferred to a nitrocellulose membrane and probed either with an anti-PorA monoclonal antibody or with anti-NspA polyclonal antibody (figure 17). As for
5 Omp85, there is a surprising indication that insertion of the promoter approximately 400 bp upstream of the NspA initiation codon expresses more protein than if placed approximately 100 bp upstream.

The same recombinant pUC plasmid can be used to up-regulate the expression of NspA in a *Neisseria meningitidis* serogroup B strain lacking functional *cps* gene
10 but still expressing PorA.

Example 14: Up-regulation of the *N. meningitidis* serogroup B *pldA* (*omplA*) gene by promoter replacement.

The aim of the experiment was to replace the endogenous promoter region of
15 the *pldA* (*omplA*) gene by the strong *porA* promoter in order to up-regulate the production of the PldA (OmplA1) antigen. For that purpose, a promoter replacement plasmid was constructed using *E. coli* cloning methodologies. A DNA region (373bp) located upstream from the *pldA* coding sequence was discovered (SEQ ID NO: 18) in the private Incyte PathoSeq data base of the *Neisseria meningitidis* strain ATCC
20 13090. This DNA contains the sequence coding for a putative *rpsT* gene. The stop codon of *rpsT* is located 169bp upstream the *pldA* ATG. To avoid the disruption of this potentially important gene, we decided to insert the *CmR*/*PorA* promoter cassette just upstream of the ATG of *pldA*. For that purpose, a DNA fragment of 992 bp corresponding to the the *rpsT* gene, the 169 bp intergenic sequence and the 499 first
25 nucleotides of *pldA* gene was PCR amplified from *Neisseria meningitidis* serogroup B genomic DNA using oligonucleotides **PLA1 Amo5** (5'-GCC GTC TGA ATT TAA AAT TGC GCG TTT ACA G-3') and **PLA1 Amo3** (5'-GTA GTC TAG ATT CAG ACG GCG CAA TTT GGT TTC CGC AC -3') containing uptake sequences (underlined). PLA1 Amo3 contains also a *Xba*I restriction site. This PCR fragment
30 was cleaned with a High Pure Kit (Roche, Mannheim, Germany) and directly cloned in a pGemT vector (Promega, USA). This plasmid was submitted to circle PCR mutagenesis (Jones & Winistofer (1992)) in order to insert suitable restriction sites allowing cloning of a *CmR*/*PorA* promoter cassette. The circle PCR was performed

using the **CIRC1-Bgl** (5'CCT AGA TCT CTC CGC CCC CCA TTG TCG -3') & either **CIRC1-XH-RBS/2** (5'-CCG CTC GAG TAC AAA AGG AAG CCG ATA TGA ATA TAC GGA ATA TGC G-3') or **CIRC2-XHO/2** (5'-CCG CTC GAG ATG AAT ATA CGG AAT -3') oligonucleotides containing suitable restriction sites (*Bgl*II and *Xho*I) underlined. The *CmR*/PorA promoter cassette was amplified from the pUC D15/Omp85 plasmid previously described, using primers **BAD20** (5'- TCC CCC GGG AGA TCT CAC TAG TAT TAC CCT GTT ATC CC-3') and **CM-PORA-3** (5'- CCG CTC GAG ATA AAA ACC TAA AAA CAT CGG GC-3') containing suitable restriction sites (*Bgl*II and *Xho*I) underlined. This PCR fragment was cloned in the circle PCR plasmid obtained with primers CIRC1-Bgl and CIRC1-XH-RBS/2. This plasmid can be used to transform *Neisseria meningitidis* serogroup B (*cps*-) and (*cps*- *porA*-) strains. Integration by double crossing-over in the upstream region of *pldA* will direct the insertion of the *porA* promoter directly upstream of the *pldA* ATG. Another cassette was amplified from the genomic DNA of the recombinant *Neisseria meningitidis* serogroup B (*cps*-, *porA*-, D15/Omp85+) over-expressing D15/Omp85 by promoter replacement. This cassette contains the *cmR* gene, the *porA* promoter and 400bp corresponding to the 5' flanking sequence of the *D15/Omp85* gene. This sequence has been proven to be efficacious for up-regulation of the expression of D15/Omp85 in *Neisseria* and will be tested for the up-regulation of the expression of other *Neisseria* antigens. Primers used for the amplification were **BAD 20** and **CM-PORA-D15/3** (5'- CGG CTC GAG TGT CAG TTC CTT GTG GTG C-3') containing *Xho*I restriction sites (underlined). This PCR fragment was cloned in the circle PCR plasmid obtained with primers CIRC1-Bgl and CIRC2-XHO/2. This plasmid will be used to transform *Neisseria meningitidis* serogroup B (*cps*-) and (*cps*-, *porA*-) strains. Integration by double crossing-over in the upstream region of *pldA* will direct the insertion of the *porA* promoter 400bp upstream the *pldA* ATG.

Example 15: Up-regulation of the *N. meningitidis* serogroup B *tbpA* gene by promoter replacement.

The aim of the experiment was to replace the endogenous promoter region of the *tbpA* gene by the strong *porA* promoter, in order to up-regulate the production of the TbpA antigen. For that purpose, a promoter replacement plasmid was constructed using *E. coli* cloning methodologies. A DNA region (731bp) located upstream from

the *tbpA* coding sequence was discovered (SEQ ID NO: 17) in the private Incyte PathoSeq data base of the *Neisseria meningitidis* strain ATCC 13090. This DNA contains the sequence coding for TbpB antigen. The genes are organized in an operon. The *tbpB* gene will be deleted and replaced by the *CmR/porA* promoter cassette. For that purpose, a DNA fragment of 3218bp corresponding to the 509bp 5' flanking region of *tbpB* gene, the 2139bp *tbpB* coding sequence, the 87bp intergenic sequence and the 483 first nucleotides of *tbpA* coding sequence was PCR amplified from *Neisseria meningitidis* serogroup B genomic DNA using oligonucleotides **BAD16** (5'-GGC CTA GCT AGC CGT CTG AAG CGA TTA GAG TTT CAA AAT TTA TTC-3') and **BAD17** (5'-GGC CAA GCT TCA GAC GGC GTT CGA CCG AGT TTG AGC CTT TGC-3') containing uptake sequences and *NheI* and *HindIII* restriction sites (underlined). This PCR fragment was cleaned with a High Pure Kit (Boehringer Mannheim, Germany) and directly cloned in a pGemT vector (Promega, USA). This plasmid was submitted to circle PCR mutagenesis (Jones & Winistofer (1992)) in order to (i) insert suitable restriction sites allowing cloning of a *CmR/porA* promoter cassette and (ii) to delete 209bp of the 5' flanking sequence of *tbpB* and the *tbpB* coding sequence. The circle PCR was performed using the **BAD 18** (5'-TCC CCC GGG AAG ATC TGG ACG AAA AAT CTC AAG AAA CCG-3') & the **BAD 19** (5'-GGA AGA TCT CCG CTC GAG CAA ATT TAC AAA AGG AAG CCG ATA TGC AAC AGC AAC ATT TGT TCC G -3') oligonucleotides containing suitable restriction sites *XmaI*, *BglIII* and *XhoI* (underlined). The *CmR/porA* promoter cassette was amplified from the pUC D15/Omp85 plasmid previously described, using primers **BAD21** (5'- GGA AGA TCT CCG CTC GAG ACA TCG GGC AAA CAC CCG-3') & **BAD20** (5'- TCC CCC GGG AGA TCT CAC TAG TAT TAC CCT GTT ATC CC-3') containing suitable restriction sites *XmaI*, *SpeI*, *BglIII* and *XhoI* (underlined). This PCR fragment was cloned in the circle PCR plasmid. This plasmid will be used to transform *Neisseria meningitidis* serogroup B (*cps-*) and (*cps- porA-*) strains. Integration by double crossing-over in the upstream region of *tbpA* will direct the insertion of the *porA* promoter directly upstream of the *tbpA* ATG.

30

Example 16: Up-regulation of the *N. meningitidis* serogroup B *pilQ* gene by promoter replacement.

The aim of the experiment was to replace the endogenous promoter region of the *pilQ* gene by the strong *porA* promoter, in order to up-regulate the production of the PilQ antigen. For that purpose, a promoter replacement plasmid was constructed using *E. coli* cloning methodologies. A DNA region (772bp) located upstream from the *pilQ* coding gene was discovered (SEQ ID NO: 12) in the private Incyte PathoSeq data base of the *Neisseria meningitidis* strain ATCC 13090. This DNA contains the sequence coding for PilP antigen. The *pilQ* gene is part of an operon we do not want to disturb, pilins being essential elements of the bacteria. The *CmR/porA* promoter cassette was introduced upstream the *pilQ* gene following the same strategy described for the up-regulation of the expression of the *pldA* gene. For that purpose, a DNA fragment of 866 bp corresponding to the 3' part of the *pilP* coding sequence, the 18bp intergenic sequence and the 392 first nucleotides of *pilQ* gene was PCR amplified from *Neisseria* serogroup B genomic DNA using **PQ-rec5-Nhe** (5'-CTA GCT AGC GCC GTC TGA ACG ACG CGA AGC CAA AGC-3') and **PQ-rec3-Hin** (GCC AAG CTT TTC AGA CGG CAC GGT ATC GTC CGA TTC G-3') oligonucleotides containing uptake sequences and *NheI* and *HindIII* restriction sites (underlined). This PCR fragment was directly cloned in a pGemT vector (Promega, USA). This plasmid was submitted to circle PCR mutagenesis (Jones & Winistofer (1992)) in order to insert suitable restriction sites allowing cloning of a *CmR/porA* promoter cassette. The circle PCR was performed using the **CIRC1-PQ-Bgl** (5'-GGA AGA TCT AAT GGA GTA ATC CTC TTC TTA-3') & either **CIRC1-PQ-XHO** (5'-CCG CTC GAG TAC AAA AGG AAG CCG ATA TGA TTA CCA AAC TGA CAA AAA TC-3') or **CIRC2-PQ-X** (5'-CCG CTC GAG ATG AAT ACC AAA CTG ACA AAA ATC -3') oligonucleotides containing suitable restriction sites *BglII* and *XhoI* (underlined). The *CmR/porA* promoter cassette was amplified from the pUC D15/Omp85 plasmid previously described, using primers **BAD20** (5'- TCC CCC GGG AGA TCT CAC TAG TAT TAC CCT GTT ATC CC-3') and **CM-PORA-3** (5'- CCG CTC GAG ATA AAA ACC TAA AAA CAT CGG GCA AAC ACC C-3') containing suitable restriction sites *BglII* and *XhoI* (underlined). This PCR fragment was cloned in the circle PCR plasmid obtained with primers CIRC1-PQ-Bgl and CIRC1-PQ-XHO. This plasmid can be used to transform *Neisseria meningitidis* serogroup B (*cps-*) and (*cps-*),

porA-) strains. Integration by double crossing-over in the upstream region of *pilQ* will direct the insertion of the *porA* promoter directly upstream of the *pilQ* ATG.

Another cassette was amplified from the genomic DNA of the recombinant *Neisseria meningitidis* serogroup B (*cps-*, *porA-*, *D15/Omp85+*) over-expressing D15/Omp85 by promoter replacement. This cassette contains the *cmR* gene, the *porA* promoter and 400bp corresponding to the 5' flanking sequence of the *D15/Omp85* gene. This sequence has been proven to be efficacious for up-regulation of the expression of D15/Omp85 in *Neisseria meningitidis* and will be tested for the up-regulation of the expression of other *Neisseria* antigens. Primers used for the amplification were **BAD 20** and **CM-PORA-D153** (5'- GGG CTC GAG TGT CAG TTC CTT GTG GTG C-3') containing *XhoI* restriction sites (underlined). This PCR fragment was cloned in the circle PCR plasmid obtained with primers CIRC1-PQ-Bgl and CIRC2-PQ-X. This plasmid can be used to transform *Neisseria meningitidis* serogroup B (*cps-*) and (*cps-*, *porA-*) strains. Integration by double crossing-over in the upstream region of *pilQ* will direct the insertion of the *porA* promoter 400bp upstream the *pilQ* ATG.

Example 17: Construction of a *kanR/sacB* cassette for introducing “clean”, unmarked mutations in the *N. meningitidis* chromosome.

The aim of the experiment is to construct a versatile DNA cassette containing a selectable marker for the positive screening of recombination in the chromosome of *Neisseria meningitidis* (ie: *kanR* gene), and a counter selectable marker to delete the cassette from the chromosome after recombination (ie: *sacB* gene). By this method, any heterologous DNA introduced during homologous recombination will be removed from the *Neisseria* chromosome.

A DNA fragment containing the *neoR* gene and the *sacB* gene expressed under the control of its own promoter was obtained by restriction of the pIB 279 plasmid (Blomfield IC, Vaughn V, Rest RF, Eisenstein BI (1991), Mol Microbiol 5:1447-57) with *BamHI* restriction enzyme. The recipient vector was derived from plasmid pCMK, previously described. The *kanR* gene of the pCMK was deleted by restriction with enzymes *NruI* and *EcoRV*. This plasmid was named pCMKs. The *neoR/sacB* cassette was inserted in the pCMKs at a *BglIII* restriction site compatible with *BamHI* ends.

E. coli harboring the plasmid is unable to grow in the presence of 2% sucrose in the culture medium, confirming the functionality of the *sacB* promoter.

This plasmid contains recombinogenic sequences allowing the insertion of the cassette at the *porA* locus in the chromosome of *Neisseria meningitidis* serogroup B.

5 Recombinant *Neisseria* were obtained on GC agar plates containing 200µg/ml of kanamycin. Unfortunately, the *sacB* promoter was not functional in *Neisseria meningitidis*: no growth difference was observed on GC agar plates containing 2% sucrose.

10 A new cassette was constructed containing the *sacB* gene under the control of the *kanR* promoter. A circle PCR was performed using the plasmid pUC4K ((Amersham Pharmacia Biotech, USA)) as a template with **CIRC-Kan-Nco** (5'-CAT GCC ATG GTT AGA AAA ACT CAT CGA GCA TC-3') & **CIRC-Kan-Xba** (5'-CTA GTC TAG ATC AGA ATT GGT TAA TTG GTT G-3') oligonucleotides containing *NcoI* and *XbaI* restriction sites (underlined). The resulting PCR fragment
15 was gel-purified, digested with *NcoI* and ligated to the *sacB* gene generated by PCR from the pIB279 plasmid with **SAC/NCO/NEWS**

(5'-CAT GCC ATG GGA GGA TGA ACG ATG AAC ATC AAA AAG TTT GCA A-3') oligonucleotide containing a *NcoI* restriction site (underlined) and a RBS (bold) & **SAC/NCO/NEW3** (5'-GAT CCC ATG GTT ATT TGT TAA CTG TTA ATT
20 GTC-3') oligonucleotide containing a *NcoI* restriction site (underlined). The recombinant *E. coli* clones can be tested for their sensitivity on agar plates containing 2% sucrose. The new *kanR/sacB* cassette can be subcloned in the pCMKs and used to transform a *Neisseria meningitidis* serogroup B *cps*- strain. The acquired sucrose sensitivity will be confirmed in *Neisseria*. The pCMKs plasmid will be used to
25 transform the recombinant *kanR/SacB Neisseria* to delete the entire cassette inserted in the chromosome at the *porA* locus. Clean recombinant *Neisseria* will be obtained on GC agar plates containing 2% sucrose.

30 **Example 18: Use of small recombinogenic sequences (43bp) to allow homologous recombination in the chromosome of *Neisseria meningitidis*.**

The aim of the experiment is to use small recombinogenic sequences (43bp) to drive insertions, modifications or deletions in the chromosome of *Neisseria*. The achievement of this experiment will greatly facilitate future work, in terms of avoiding

subcloning steps of homologous sequences in *E. coli* (recombinogenic sequences of 43bp can easily be added in the PCR amplification primer). The *kanR* gene was PCR amplified from plasmid pUC4K with oligonucleotides **Kan-PorA-5** (5'-GCC GTC TGA ACC CGT CAT TCC CGC GCA GGC GGG AAT CCA GTC CGT TCA **GTT TCG GGA AAG CCA CGT TGT GTC**-3') containing 43bp homologous to the 5' flanking sequence of *NmB porA* gene (bold) and an uptake sequence (underlined) & **Kan-PorA-3** (5'-TTC AGA CGG CGC AGC AGG AAT TTA TCG GAA ATA ACT GAA ACC GAA CAG ACT AGG CTG AGG TCT GCC TCG-3') containing 43bp homologous to the 3' flanking sequence of *NmB porA* gene (bold) and an uptake sequence (underlined). The 1300bp DNA fragment obtained was cloned in pGemT vector (Promega, USA). This plasmid can be used to transform a *Neisseria meningitidis* serogroupB *cps*- strain. Recombinant *Neisseria* will be obtained on GC plates containing 200µg/ml kanamycin. Integrations resulting from a double crossing-over at the *porA* locus will be screened by PCR with primers PPA1 & PPA2 as described previously.

Example 19: Active protection of mice immunized with WT and recombinant *Neisseria meningitidis* blebs

Animals were immunised three times (IP route) with 5 µg of the different OMVs adsorbed on Al(OH)₃ on days 0, 14 and 28. Bleedings were done on days 28 (day 14 Post II) and 35 (day 7 post III), and they were challenged on day 35 (IP route). The challenge dose was 20 x LD50 (~10⁷ CFU/mouse). Mortality rate was monitored for 7 days after challenge.

OMVs injected were:

- Group1: Cps-, PorA+ blebs
- Group2: Cps-, PorA- blebs
- Group3: Cps-, PorA-, NspA+ blebs
- Group4: Cps-, PorA-, Omp85+ blebs
- Group5: Cps-, PorA-, Hsf+ blebs

Figure 15 illustrates the pattern of these OMVs by analyzed SDS Page (Coomassie staining).

24 hours after the challenge, there was 100% mortality (8/8) in the negative control group (immunised with Al(OH)₃ alone) while mice immunised with the 5

different OMVs preparations were still alive (7 to 8/8 mice survived). Sickness was also monitored during the 7 days and the mice immunised with the NSPA over-expressed blebs appeared to be less sick than the other groups. PorA present in PorA+ blebs is likely to confer extensive protection against infection by the homologous strain. However, protection induced by PorA- up-regulated blebs is likely to be due at least to some extent, to the presence of increased amount of NspA, Omp85 or Hsf.

Example 20: Immunogenicity of recombinant blebs measured by whole cell & specific ELISA methods

To measure the ability of the antibodies to recognize the antigens present on the MenB cell surface, the pooled mice sera (from Example 19) were tested by whole cell ELISA (using tetracyclin inactivated cells), and titers were expressed as mid-point titers. All types of bleb antibodies induce a high whole cell Ab titer while the negative control group was clearly negative.

15

Bleb	WCE(H4476) mid-point titer	
	14P2	14P3
CPS(-)		
PorA(+)	23849	65539
CPS(-)		
PorA(-)	20130	40150
CPS(-)		
PorA(-)		
NSPA(+)	8435	23846
CPS(-)		
PorA(-)		
OMP85(+)	4747	16116
CPS(-)		
PorA(-)		
HSF(+)	6964	22504
(-)	51	82

The specific Ab response to available recombinant HSF protein was carried out. Microplates were coated with 1 µg/ml full length HSF molecule.

The results illustrated in Figure 16 show that there was a good specific HSF response when HSF over-expressed OMVs were used to immunize mice (using purified recombinant HSF on the plates). The HSF over-expressed blebs induce a good level of specific antibodies.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A modified *Neisseria meningitidis* strain wherein said strain is obtained by employing the following processes:
 - 5 e) a process of detoxifying the lipid A portion of LPS within the strain, comprising the steps of identifying an *msbB* gene involved in rendering the lipid A portion of LPS toxic, and engineering the strain so as to reduce or switch off expression of said gene; and
 - i) a process of engineering the strain such that it is free of capsular polysaccharide.
- 10 2. The modified *Neisseria meningitidis* strain of claim 1, wherein the *msbB* gene has a nucleotide sequence of SEQ ID NO: 81.
- 15 3. The modified *Neisseria meningitidis* strain of claim 1 or 2, wherein the *msbB* gene is downregulated via point mutation or deletion.
4. The modified *Neisseria meningitidis* strain of claim 1 or 2, wherein part or all of the *msbB* open reading frame or promoter is deleted.
- 20 5. The modified *Neisseria meningitidis* strain of claims 1-4, wherein a gene coding for capsular polysaccharide biosynthesis or export is inactivated by mutating the control region or the coding region of said gene.
- 25 6. The modified *Neisseria meningitidis* strain of claims 1-4, wherein a gene coding for capsular polysaccharide biosynthesis or export is downregulated via point mutation or deletion.
7. The modified *Neisseria meningitidis* strain of claims 1-6, wherein in process
30 h) one or more genes are downregulated from a list consisting of: *galE*, *siaA*, *siaB*, *siaC*, *siaD*, *ctrA*, *ctrB*, *ctrC*, and *ctrD*.

-
8. The modified *Neisseria meningitidis* strain of claims 1-7 which is a serogroup B strain.
9. A genetically-engineered bleb preparation made from the modified *Neisseria meningitidis* strain of claims 1-8.
10. A vaccine comprising the genetically-engineered bleb preparation of claim 9, and a pharmaceutically acceptable excipient.
11. A vaccine against *Neisseria meningitidis* generated from the modified *Neisseria meningitidis* strain of claims 1-8.
12. The vaccine of claim 10 or 11 further comprising one or more plain or conjugated meningococcal capsular polysaccharides selected from the serotypes A, C, Y or W.
13. The vaccine of claim 10 or 11 further comprising a conjugated *H. influenzae* b capsular polysaccharide, and one or more plain or conjugated pneumococcal capsular polysaccharides.
14. A method of making a vaccine against *Neisseria meningitidis* comprising the step of generating said vaccine from the modified *Neisseria meningitidis* strain of claims 1-8.
15. Use of the modified *Neisseria meningitidis* strain of claims 1-8 in the manufacture of a medicament for immunizing a human host against a disease caused by infection of *Neisseria meningitidis*.
16. Use of the genetically-engineered bleb preparation of claim 9 in the manufacture of a medicament for immunizing a human host against a disease caused by infection of *Neisseria meningitidis*.

17. A method of immunising a human host against a disease caused by infection of *Neisseria meningitidis* comprising administering an effective amount of a modified *Neisseria meningitidis* strain according to any one of claims 1-8.

5 18. A method of immunising a human host against a disease caused by infection of *Neisseria meningitidis* comprising administering an effective amount of a bleb preparation according to claim 9.

DATED: 15 December 2003


10

PHILLIPS ORMONDE & FITZPATRICK

Patent Attorneys for:

SMITHKLINE BEECHAM BIOLOGICALS S.A.

15



20

25

30

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11

11/17




Figure 12

12/17

Schematic representation of the recombinant PCR strategy used to delete the *lacO* in the chimeric *porA/lacO* promoter.




Figure 13




Figure 14

14/17




Figure 15




Figure 16





Figure 17



SEQ. ID NO:1

Nucleotide sequence of the pCMK(+) vector

TCCTCCGCTTCCCTCGCTCACTGACTCGCTCGCTCGGTCTTCCGCTCGCGGAGCGGTATCAGCTCACTCAAAGCGGT
AATACGGTTATCCACAGAATCAGGGGATAACGCGAGAAAGAACATGTGAGCAAAGGCCAGAAAAGGCCAGGAACCGTA
5 AAAAGCCGCGTTCCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCAAAAATCGACGCTCAAGTCAGAGG
TGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTTCCCTCGGAAGCTCCCTCGTCCGCTCTCCTGTTCCGACCT
CGCGCTTACCGGATACCTGTCCGCCITTCCTCCTTCGGGAAGGTGGCGCTTTCATAGCTCAGCTGTAGGTATCTCA
GTTCCGGTGTAGTCTGTCCGCTCCAAGCTGGGCTGTGTGACGAAACCCCGCTCAGCCGACCGCTGCGCCTTATCCGGT
AAGTATCGTCTTAGTCCAACCCGTAAGCAGCACTTATCGCCACTGGCAGCAGCCACTGGTACAGGATTTAGCAGAGC
10 GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCCTAATCAGGCTACACTAGAAGAACAGTATTTGGTATCT
CGCTCTGCTGAAGCCAGTTACCTTCGGAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGGT
GGTTTTTTTGTGTGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGTCTTTGATCTTTTACGCGGTC
TGACGCTCAGTGGAAACGAAAACCTCACGTTAAGGGATTTTGGTCAATGAGTATCAAAAAGGATCTCACCTAGATCCTTT
TAAATTAATAAGTAACTTTAAATCAATCTAAAGTATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGT
15 GAGCCACTATCTCAGCGATCTGTCTATTTTCGTTCACTCATAGTTGCTGACTCCCGCTCGTGTAGTAACTACGATAACG
GGAGGCTTACCATCTGGCCCCAGTGTCAATGATACCGCAGAACCCAGCTCACCGGCTCCAGATTTATCAGCAATAA
ACCAGCCAGCCGGAAGGCGCAGAGTGGTCTTGCACACTTATCCGCTCCATCCAGTCTATTAATTTGTCGCGG
GAGCTAGATAGTAGTTCGCGAGTAAATAGTTTGGCAGCTTGTGCGCATGCTACAGGCATCGTGGTGTCAAGCTC
GTGCTTGGTATGGCTTCACTCAGTCCGGTTCACACGATCAAGGCGAGTTACATGATCCCCATGTTGTGCAAAAAG
20 CGGTAGCTCCTCGTCCGATCGTTGTGAGAATAGTTGGCAGGTGTATCACTCATGGTTATGGCAGCACTG
CATAAATCTTACTGTATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGTACTCAACCAAGTCTTCTGAGAATA
GTGATATCGCGGACCCAGGTTGCTCTTGCCCGCGTCAATACGGGATAAATCCGCGCACATAGCAGAACTTTAAAGTGC
TCATCATTTGAAAACGTTCTTCGGGGCAAAAACCTCAAGGATCTTACCCTGTTGAGATCCAGTTCGATGTAACCCACT
CGTGCACCCAACTGATCTTTCAGCATCTTTACTTTCACCGCGCTTCTGGTGAGCAAAAACGAGGAAGCAAAATGCCG
25 AAAAAAGGGAATAGGGCGACAGGAAATGTTGAATCTCATACTCTCCCTTTTCAATATATTGAAGCATTATCAGG
GTATATGTCTCATAGCGGGATACATATTTGAATGATTTGAGAAAAAATAACAAATAGGGGTTCCGCGCACATTTCCCGA
AAAGTGCACCTGACGCTTAAGAAACCATATTATTCATGACATTAACCTATAAAAATAGGCGTATCACGAGCCCTTTCC
TTCGCGCGTTCGCTGATACGGGTGAAAACCTTGACACATGACGACTCCCGGAGACGGTCAACGCTTGTGTGAAGCG
ATGCGGGAGCAGACAAAGCCGTCAGGCGCGTCAGCGGTTGGCGGTTGCGGGTGGCTTAACTATGGCGCATCA
30 GACCATGCTACTGAGAGTGACACATAAAATGTAACCTTAATATTTGTTAAATTCGCGTAAATTTTGTAAAT
CAGCTCATTTTTAAACCAATAGCCGAAATCGSCAAAATCCCTTATAAATCAAAGAATAGCCCGAGATAGGGTTGAGTG
TTGTTCCAGTTTGGAAACAGAGTCCACTATTAAGAACGTTGACTCCAACTCAAAGGGCAAAAACCGTCTATCAGGGC
GATGGCCACTACGTGAACCATCACCAATCAAGTTTTTTGGGGTGGAGTGCCTAAAGCACTAAATCGGAACCTAA
35 AGGGACCCCGATTAGAGCTTGACGGGAAAGCCGCAAGCGTGGAGAAAGGAAGGAAGAAAGCAAGGAGCGG
GCGTACGCGGCTGGCAAGTGTAGCGGTACGCTGCGCGTAAACACACCCCGCGCTTAAATGCGCGCTACAGGGC
GCGTACTATGTTGCTTTGACGTATGCGGTGAAATACCGCACAGATGCGTAAGGAGAAAATCCGCTACAGGCGCCAT
TCGCCATTCAGGCTGCGCAACTGTTGGGAAGGCGATCGGTGCGGGCTCTTCGCTATACGCCAGCTGGCGAAAGGGGG
ATGTGCTGAAGCGATTAAGTTGGGTAACCCAGGTTTTCCAGTCAAGAGTTGTAACGACGCGCCAGTCCCAAGC
40 TTGCGCTGAAATACATCCCGTCACTTCCATAAAACAGAAAACAAATCAGAAAACCTAAAATCCCGTCACTCCGCGCA
GGCGGAATCCAGTCCGTTCACTTCGGTCAITTCGATAAATTCCTGCTGCTTTTCAATTTCTAGATTTCCCACTTTGCTG
GGAATGACGCGGAAAGGTTTTGGTTTTTCCGATAAATCTTGAGGCATTGAAATTCAGATTTCCCGCTGCGCGGAA
TGACCGCTGTAGATGCCGATGTTTATAGCGGATTAACAAAATCAGGACAGGCGCAGAAAGCCGACAGTACAG
ATAGTACGGAACCGATTCACTGGTGTTCAGCACTTAGAGAAATCGTTCTTTGAGCTAAGGCGAGGCAACCGCTTAC
45 TTGTTTTTTTAAATCACTATAAAGTGCAGGCTGTGTTTTTTTATGGCGTTTTAAAAGCGAGACTGATCCGGGCAGC
AGCGCATCGGCCCGCAGAGGCTCTGGAGTGCAGGACTCAAGGGGAAATTCGAGGGGGGGGGAAAGCCAGTT
GTGTCTCAAAATCTCTGATTTACATTTGCACAGATAAAAATATATCATGATGAACAATAAACTGTCTGCTTACATAAA
CAGTAATCAAGGGGTGTTATGAGCCATATTCAACGGAAAACGCTTGTCTCGAGCCCGGATTAATTTCCAACTGGATG
CTGATTTATAGGGTATAAATGGGCTCGGATAATGTGGGCAATCAGGTGGCACTATCTATCGATTTGATGGGAAGCC
GATGCGCCAGAGTGTCTGAAACATGGCAAAGTAGGTTGCCAATGATGTTACAGATGAGTGGTCAACTAACTG
50 GCTGACGGAATTTATGCTCTTCGACCATCAAGCATTATCCGTAATCTCCTGATGATGATGATGATGATGATGATGATG
TCCCGGGAAAAACAGCATTCCAGGTTATGAGAAATATCCGATTCAGGTGAAAATTTGTTGATGGCTGGCAGTGTTC
CTGCGCGGTTGCAATTCGATTCCTGTTGTAATTTGCTTTTAAACAGCATCCGATTTCTGCTCCTCGTCAGGCGCAATC
ACGAATGAATAACGGTTTTGGTTGATGCGAGTATTTGATGACGAGCTAATGGCTGGCTGTGAACAAGTCTGGAAAG

AAATGCATAAGCTTTTGCCATTCTCACCGGATTGAGTCTGCTACTCATGGTATTTCTCACTTGATAACCTTATTTTTGAC
GAGGGGAAATTAATAGGTTGTATGTGATGTTGGACGAGTCGGAATCGCAGACCGATACCGAGATCTTGCCATCCTATGGAA
CTGCGTCGGTGAGTTTCTCCTTCATTACAGAAACGGCTTTTCAAAAATATGGTATTGATAATCTGATATGAATAAAT
TGCAATTTTCAATGATGCTCGATGAGTTTTTCTAATCAGAATTGGTTAATGGTTGTAACACTGGCAGAGCATTACGCTG
5 ACTTGACGGGACGGCGGCTTTGTTGAATAAATCGAACTTTGCTGAGTTGAAGGATCAGATCAGCATCTTCCGACAAAC
GCAGACCGTTCGCTGGCAAGCAAAAGTTCAAAAACCAACTGGTCCACCTACAAACAAAGCTCTCATCAACCGTGGCTC
CCTCACTTTCTGGCTGATGATGGGCGATTACGGCTGATGAGTACGCAACACCTTTTACAGGCGAGACCTCAGC
GCCCCCCCCCTGACGAGGCTCTGCGCTGAAATGTTGTTGTAGAAACACAACGTTTTTGAAAAATAAGCTATTGTTT
TATATCAAAATATAATCAATTTTAAAAAAGGTTGCGGCAATTTATCAGATATTGTTCTGAAAAATGGTTTTTGCGGG
10 GGGGGGGTATAATTGAAGCGTATCGGGTGTTCGCCGGAATGTTGAGCGGATAACAAATCGATGTTTTAGGTTTTTA
TCAAAATTTCAAAAGCAAGCCCATATGCATCCTAGGCCATTAATATTTCCGGAGTATACGTAGCCGGCTAACGTTAAACA
CCGGTACCTTAGAATATAGCTAGCATGCGCAAAATTAAGCGCTGATATCGATCGCGCAGATCTGATTAATAGGC
GAAATACACAGTACGATCAAAATCATCGCGGCTGATTATGATTTTCCAAACGCACTTCGCGCATCGTGTCTGGCGC
TTGGCTGAAACGCAATACCGGCATCGGCAACTACACTCAAAATTAATGCCCTCGTGGTTTTGCGCCAAAAATCTAAA
15 TATCGGGCGGTTGAGCGGATAGCTTTGTTTTGACGGCTTGCCTTCACTTTGATTGCAATCTGACTGCCAATCTGC
TTCAGCCCAAAACAAAACCGGATACGGAAAGAAAAACGGCAATAAAGACAGCAAAATACCGTCTGAAAGATTTTCAGAG
GTATTTGCGATTTTGGCTTGGTTGCGCATATAGTGAGACCTTGGCAAAAATAGTCTGTTAACGAAATTTGACGCATAA
AAATGCGCCAAAAATTTCAATGGCTAAAACCTTCTAATATTGAGCAAAAATAGGAAAAATCAGAAAAGTTTTGCA
TTTTGAAAAATGAGATTGAGCATAAAAATTTAGTAACCTATGTTATTGCAAAAGGTCTGAAATTTTCATCCACGAGGG
20 GGAATCTAGTCTGTTGCGTTTCAGTTATTTCCGATAAATTCCTGCTGCGCGCTGGAAGAATCGTAATCATGTCATAG
CTGTTCTCTGTGAAATGTTATCCGCTCACAAATCCACACAACATACGAGCCGGAAGCATAAAGTGAAGCTGGGG
TGCTAATGAGTGAAGTAACTACATTAATTCGTTGCGCTCACTGCGCGCTTCCAGTGGGAAACCTGCTGCTGCCAGC
TGCAATTAATGAATCGCCCAACGCGGGGAGGCGGTTGCGTATTGGGCGC

25 SEQ. ID NO:2

Nucleotide sequence of DNA region (997 bp) up-stream from the NspA gene in the *Neisseria meningitidis* serogroup A strain Z2491.

GGAAACCGAAACCGCGTTCCGTCATACGCGCCGAAAGGTTTGGCGAAGAAGAAAGCCCGCTCGACATCGAAGACGGG
TACACGGCCGGCTGAAAGCGGGTTTTGTCCACTACGAAACATCGGCTTTTGGAAACAGCCATGCAATGCGCCAC
30 AAATTTGAATCTGCGAGTTTCGGCGATTTATGAGCATAGGCGGGCGCGCACGCAAAATTTCTATCCCGACCGCAT
CGAGCGCAACCGTCCGCGCCGCAACCGCAACGACTACCTCGCTTAATGCAAAACCGACGAGCGAAGCCGTCGAACGA
AAACCGTCCGCGCAAGATTTCCGTTGCAATTCATGATGAACGCCCTGCGCCTGACCGAGCGGTAACCCACCGGATG
TTGCAGGAGCGCACGGCGTACCGAGTCCAAAATCATGGCGCAATCGAAACGGCAAGGCAAAAAGCCGCTGGAAC
CGACCCCGCGTATTCGCGCGAAGAAAGGACGCTTGTTTTTAAACGATTTGCTGCAAGTTTTTATAGTGGATTA
35 ACAAAAACCGATCGCGGTTGCTGCGCTTAGCTCAAGAGAACGATTCTTAAGGTGCTGAAGCACCAAGTGAATCGGT
TCCGTAATATCTGATCTGCTGCGGCTTCTGCTGCTGCTGATTTTGTAACTCACTATATAAGCGCAAAACAAATCG
GCGCCGCGCGGAAACCCCGCAACCGGTCGGGAAATATGCTTATCGATGGAACCGACGCGCATCCCCCGCGG
GCGTTTTAGAGCGGACAGCCGCGCGGAAATGTCGACGCTTAAGGCAACAGACGACACAAAAACCGTATGCTGAC
40 CTGCAACATCGACAGATACCGCTGTTTTTCCAAACGTTTGAAGTTTACCCATCCGCGCGTATGCGCGCACCA
CCATTTAAAGGCAACGCGGGTTAACGGCTTTGCGG

SEQ. ID NO:3

Nucleotide sequence of DNA region (1000 bp) up-stream from the D15/Omp85 gene in the *Neisseria meningitidis* serogroup B strain ATCC13090.

ACCATTTGCGCCCGCGCGCTTCCAAAGCGGCGCAAAAATACAATCCGTCAACGGCACACCGGTTGAGATTGGGGCAG
CGCGCAAAACGAAATCGTCTCAACCTCGAAGCGGCAAAAGTCCGCTCGGTTTACAGCGGCATCAGGCGGCAAAACCGT
CCGCACCATCGATGCGCGGACGCGGAAAGCGGTAATAATCGCAAAAAACCAAGGCTACATCGGACTGATGCCCCTTA
AAATCAACACCGTTGCGGTTGCGTGGAAAAAGGACGCGCCCGCAAAAAGCAGGCTGAAACCGGCGACAGGCTGACT
GCCGCGGACGGCAAAACCACTTACCTCATGCGAAGAAATGGGCAAACTGACCCGCAAAAGCCCGCAAAAATACCTT
50 GAACCTAGAAAGCGGCGGCAAAACCACTACCGCGGACATCGCGCGGATACTGTGAAACAGCCGACCAACCCCTGATCG
GGCGGCTCGGCTCCGTCGCGAGCGGACAGGCGTGGAGCGGCAAAATCCGCGCAGCTACCGTCCGCTGTTATCCGCG

5 GCATTCGGCATGGGCTGGAAAAACCGTTTCCCACTCGTGGACAAACCTCAAATTTTTCCGCAAACTAATCAGCGGCAA
CGCCTCCGTCAGCCATATTTCCGGGCGCTGACCATTCGCGACATTGCGGACAGTCCGCGGAACTGGCTTGCAAAGTT
ATTTGGAAATTTTGGCACTGGTCAGCATCAGCCTCGGCGTGTGAACCTGCTCCCGTCCCGTGGTGGACGGGCGCAC
CTCGTGTATTAATCTCCGAATGGATACGCGGCAACCTTTGGGCGAACCGCTCAAAAACATCGGTTTGGCTTCGGGT
TGCCCTCATGATGCTGATGGCGGTCCGCTTCTTCAACGACGTTACCGGCTGCTCGGTTAGATTTACGTTTCGGAA
TGCCGCTG6AAACCGCATTCCGACCAACAAGGAAGTACA

SEQ. ID NO:4

Nucleotide sequence of DNA region (1000 bp) up-stream from the Hsf-like gene from *Neisseria meningitidis*

10 ATTCCTCCGCGAGCGGGAATCCAGAAAACGCAACGCAACAGGAATTTATCGGAAAAACAGAAACCTCACCGCCGTCATT
CCGCAAAAGCGGAAATCTAGAAAACAACCGCGCAGGACTTTATCAGAAAAACAGAAACCCACCGCGTCAATCCCGC
AAAAGCGGAAATCCAGACCGCTCGGCACGGAAACTTACCGGATAAAAACAGTTTCTTAGATTCCAGGTCCTAGATTCCCG
CTTTCCGGGAAATGACGAGATTTAGATTTATGGGAATTTATCAGGAATGATGAATCCATAGAAAAACCAAGGAATCTA
15 TCAGAAAAACAGAAAACCCCAACCGCTCATTCCCGCGCAGGCGGAAATCCAGAAAACAACCGCGCAGGACTTTATCGG
AAAAACCGAAACCCACCGACCGTCAATCCCGCAAAAGTTGGAATCCAAAAACGCAACGCAACAGGAATTTATCGGAAA
AAACAGAAAACCCCAACCGCTCATTCCCGCGCAGGCGGGAATCCAGAAAACAACCGCGCAGGACTTTATCGGAAAAAC
AGAAAACCCCAACGACGCTCATTCCCGCAAAAGCGGAAATCCAGAAAACAACCGGAAATCTATCAGCAAAAACAG
AAACCCCAACCGACCGTCAATCCCGCGCAGGCGGGAATCCAGAAAACAACCGCGCAGGACTTTATCGGAAAAACAGAA
20 ACCCAACCGACGCTCATTCCCGCAAAAGCTGGAATCCAAAAACGCAACGCAACAGGAATTTATCGGAAAAACAGAAAC
CCACCGCGCTCATTCCCGCAAAAGCGGAAATCCAGACCGCTCGGCACGGAATTTACCGGATAAAAACAGTTTCTTAGAT
TCCAGCTCCCGAATCCCGCTTCCCGGAAATGACGAGATTTAAGTTGGGGAATTTATCAGAAAAACCCCAACCCCA
AAAACCGGCGGATGCCACCATCCGCCCCAAACCCGATTTAACATTCAAACAAACCAAAAGAAAAACAA

25 SEQ. ID NO:5

Nucleotide sequence of DNA region (772 bp) up-stream from the PilQ gene from *Neisseria meningitidis*

30 GCGATGTCGGGAAGCCTTCTCCCGAATCATTACCCCTTGAGTCGCTGAAATCGCCAAATCTCCGAAAAACGGGCGCAAT
CATGACGGCAAGAGCAGCATCTGAACTCAGTGCATTGCCACCACTACCAAGCAAAATCCGTAGAGAGCTTGCCGC
AGAAGCGGCACAAAATGCCAGCAAAAATAACTTACGTTAGGGAAACCAAGAAACACTATGCCTTACTCATCAGCTTTCT
GGCTCTCTCCGCGTGTCCCAAGGTTCTGAGGACCTAAACGAATGGATGGCAACAACCGCGACGCGAAGCCAAAGCAGAAA
TCATACTTTCCCAAGCACCTACCCCTGCGGTTGCGCGGTATACAGCCCGCCGAGCTTACAGGGCCGAAACGATTCGAC
TTCCGCGCATGGAACCGCAAAAAAGGGAAAAATGCCCCGACACCAAGCGTATTAAAGAAACGCTGAAAAAATCAG
35 TTTGAAAAATATGCGTTATGTCGGCATTTTGAAGTCTGGACAGAAAGTCTCCGCTTCATCGAGGCTGAAGTTATGCT
ACACTGTCGCTGTCGGCACTATTGGGACAAAACACTACGGTAGAATCGAAAGCATTACCGACGACAGCATCGCTCTGAAC
GAGCTGATAGAGACAGCAGCGGCAACTGGGTTTCCCGTAAAGCAGAACTGCTGTTGAATTTCCGACAAAAACACCGA
ACAAGCGGACGACCTGCCGAGAACAAAATTAAGAAGAGGATTACTCCATT

SEQ. ID NO:6

Nucleotide sequence of DNA region (1000 bp) up-stream from the Hap gene from *Neisseria meningitidis*

40 GTGCGGCAAAAAACAGCAAAAGCCCGCTGCGATTGCTGACCGTCCGCGTCCGTAATAACAGCATAGGTTGCCACGCGC
GGCTTGGCGTTTTCCCAACAAGCCTCTGCCATCGGCGCAGGTTTTTCCCGATATGCGTATCAGCCCAACGCGCC
GCGCCGCGTGGCGTAGCGACTGCCCAATCGTTGGAACGTTATCCGACATAAAACCCCGAAAAATCAAACAGCCGCG
45 ATTATAGCAAAATCCGCTGAGTCCGACGGTTTGGCTTTCAGACGGCATAAAACCGAAAAATGCTTGATAAATCCGTC
CGCCTGACCTAATAACCATATGAAAAACGAAACAATACGCTTCCCTGCTCGGTATAGGCTCGCTGCTGGGTCTGTT
CCATCCGCAAAAACCGCATCCGCCCAATCCCGCGACGATCTCAAAAACATCGGCGCGATTTTCAACGCGCCATAG
AGAAAAGCGGAAAATGACGAAAAACGCAACGACAGGCAAGGCGCGCAGGCTGCGAAAACCGTCTCAAAATCCCGGAGCTTG
TCGAGCAAAATCCTGTCGACGAGTACGTGCAATAATGATAGCCCGCGTTTCCATTGCGGATCGTTGCCGCGCGCTCC
50 GACTTGGCGCAATACACGACATTAACAGCAACGCGGACAGCCGATTAAGCAATGGCGGAAAAAGAACAGCCGCTCCG
GCACGAAACCATACGCAAGCAAAACCTTCAACAGGCGCGGCAACTGTACGCGCTCATCAGCGTCATCTGATACTGC
TTTTTCCGCTCTCTCGTATGGAGGGTACCCCGCAACCGCGCTCCCTTCCGCGCGCACAGTGGTTGCTTGGCG
GGTCTTTCGATTTGGAAGAGCGAGACCAAGGCAAAAATTAATGCAAAATCCTAGGCGTGTCTCATATCCGCGCGA
ACGCGCAACCGCACATATAGGCACATCCCGCGCGCCGCGAAGCGAAGCCGCGCTCCCAACAAACCCGAAATCCCG
55 TCAGATAAGGAAAAATA

SEQ. ID NO:7

Nucleotide sequence of DNA region (924 bp) up-stream from the NspA gene from *Neisseria meningitidis* (serogroup B) (ATCC13090)

5
10
15
GGAAACGAAACACGCGTTCGGTTCATACGCGCGAAAGTTGCGCGAAGACGAAACCGCCCTCGACATCGAAGACGCGG
TACACGGCGCGTGGAAAGCGCGGGTTTTGTCACACTACGAAACATCGGCTTTTGC3AAACAGCCATGCAGTCCCGCAC
AATTTGAACACTAGCGAGTTCGGCGATTATTAGGCATAGGCGGGGCGCTCACGG3CAAAAATTTCTATCCCGACCGCAT
CGAGCGACCGTCCGCGCGCCACCCCAAGACTACCTCGCCTTAATGCAAAGCCAAACAGGATGAAGCGTCGAACGCA
AAACCGTTGCCCGAAGATTGCGGTTTGAATTCATGATGAACGCCCTGCGCCT3ACCGACGCGTACCGCCGCGATGT
20 TGCAAGGAGCGCAGGGCGTACCGAGTCCAAAATCATGGCGAAATCGAAACCGCAAGGCAAAAAGGCGCTGTGGAAACC
GACCCCGCGTATTCGCGCGACCGAAAAGGACGCTTGTTTTAAACGATTTGCTGCGAGTGTTTTTATAGTGGATTAA
CAAAAACAGTACGCGTTCGCTCGCCTTAGCTCAAAGAGAACGATTTCTTAAGGTCGAAAGCACCAGTGAATCGGTT
CGGTACTATTTTGACTGCTGCGGCTTCGTCGCTTGTCTGATTTTGTATTAACCTATATAAGCGCAAAACAAATCGG
CGGCGCGCGGAAAACCGCGCGAAACGCGTCCGAAAATATGCTTATCGATGAAAACGACGCGCATCCCGCGCGG
GCGTTTCAGACGGCACAGCGCGCGCGAAAATGTCGACGCTTAAGGCACAGCGCACAAAACCGTATGCTCGACCT
30 GCAACAATCCGACAGATACCGCTGTTTTTCCAAACCGTTTGA

SEQ. ID NO:8

Nucleotide sequence of DNA region (1000 bp) up-stream from the FrpB gene from *Neisseria meningitidis* (serogroup B)

20
25
30
AAGTGGGAATCTAAAATGAAAACCAACAGGAATTTATCGGAAATGACCGAACT3AACGGACTGGATTCCCGCTTTCCG
GGAAATGACGGCGCAGGGTTGCTGTTATAGTGGATGAACAAAACAGTACGTC3TTGCTCGCCTTAGCTCAAAGAGA
ACGATTCCTAAGGTGCTGAAGCACAAGTGAATCGGTTCCGCTTATTTGACTGCTGCGGGCTTCGTCGCTTTGCTCT
GATTTCTGTTGTTTTCGGTTATTCGGATAAATACCGCGCTTCTCGTCAATTTCTTAAACCTTCGTCATTCCCGCGC
25 AGGCGGGAATCTAGTTTTTTGAGTCCAGTGTCTGATAAATCTTCGAGCTTTGAGTTCCTAGATTCACCTTTCC
TGGGAATGAGGTGGAAAAGTTGCCGTGATTTCCGATAAATTTGTAACGCAATTTCCGTTTTACCGGATAAATGCC
CGCAATCTCAAAATCCGCTATTCGCAAAAACAAAATAAAAACAGAAATATGCTCAATCCCGCGCAGGCGGGAATCT
AGAAGCTTAGAAAACAGCAATATTCAAAAGATTATCTGAAAAGTCGAGATTTAGATTTCCCACTTTGCGGAAATGACGA
30 TTTAGTTTTCTGTTTTGTTTTGTTCTGCTTCCGCGGAATGATGAATTTAAGTTTAGGAATTTAAGGAAAACAG
AAACCGCTTCGCGCTCATTCCGCAACAGGCTTCGTCATTCCCGCGCAGGCTTCGTCATTCCCGCATTTGTTAATCCACTA
TATTTCCCGCGTTTTTACATTTCCGCAAAAACCTGTCACAAAACAAACACTTCGCAATAAAAACGATAATCAGCTT
TGCAAAAATCCCGCGCGGTTTAAATAAAAATAAAAATAAATAAATTTTCTTATCTCGCAAACTTAAACGGT
TTGGATTTACTTCCCTCATACTCAAGAGGACGATTGA

SEQ. ID NO:9

Nucleotide sequence of DNA region (1000 bp) up-stream from the FrpA gene from *Neisseria meningitidis* (serogroup B)

35
40
45
50
CTATAAAGATGTAATAAAAATCTCGGTAACSGTAACTTTGGCTCAGCAAGCAGCTACACAAAACAGACGGTACAA
CCGCAAAAATGGGGATTTACTTTTAGCAGCCGCAATCTGCACAGCCGCTCAC3PACAAAATGCTATCCATTAGCCAT
GTTCCGGAAAACAGATTTCCCGTTTTTTAGCGCTGTCTAAACAATAACGTAAGGTATCATTTAATAATA
ATAAAGATTTAATCTTTATGACGAAATTTAGAGAAAGGCTAGACTGTCGATTAATGACAAACAAATCTGAGAAAG
GAANTATTTACTATCCGAGCACAGACATTTTAGTAGCCTGTAAGTTCCTCGCGGAAAGGATGAAGTGGGA
CTTACCAGGAATAAATGCTCGTGTGTGATATGATGCCATCCGCGAAGCAATGATGCAATCAGCGAGTCTACT
TGAATGAAACCTGCGTGTGAGANTTTGAAAACGCTATTTAAGAAAGGATAAGGGGAAAGAAATTTGGTTTTTA
45 GCTGATGAAACCGTTGGAAATAAGTACACCTACGATAAATAAATTTTGGTTTTTATTCTACAGACTATTATA
TAGATGCTTAAAGTTTTAATTTTAGATGCCAAAATAATTTTATATCTCATATGTTTTATATGCTTTATTTTGA
ATATCTTACGATGGGAAATTTATATATTTTAAATAAATTTACTCATTTGCTAATATGCTATGGAATATTACTT
GATTTTGTAGAAATTTTCCATATGAAAATATTCATTACTATTTTCTGAACCTTATAGTTTTATTTAATTTTT
ACCTCTATATTTACCATAAGAGAGCTAATGATTCATATATTTAGTCGATTAATTTATTTCTTAATTTAATTT
50 CTCAGTTATTTTTAATTTACTTGAAGGAAACGAGAT

SEQ. ID NO:10

Nucleotide sequence of DNA region (1000 bp) up-stream from the FrpC gene from *Neisseria meningitidis* (serogroup B)

55
60
65
GGAAACAGAAAAAGTTTCTCTCTATCTGGATAAATATATTTACCCCTCAGTTTGTAGTAAAGTATGGAATTTATACC
TAAGTAGTAAAAGTTAGTAATATTTTTAACTAAAGAGTTAGTATCTACCATAATAITA'CTTTAACTAATTTCTAGGC
TTGAAATTTAGAGACCATATGCTACTACATTTA'CAAGCTTTTATTTGTTTATTTGGAGTGTTTTTACTATGACCTCA
TGTAACCTGTGAATGAAAAGACAGATCAAAAAGCAGTAAAGTGGCAACAGGCTAAAGAAACAAACAGTTTCAACATCC
60 CGAGCCATGACAGGATTTGAACATACGGTTACATTTGATTTTACAGGCAACAAAAGGTTATCCCTATGGCTATCTTG
CACGGTATGCGAAGCAATGCCAAAATGGCTTTCCGACACGCGCGGCGAGATGCTTACTCCATTAATTTGATAGAG
ATTAGCTTATTTACAAAACCCGACCAAGGCTGGTCTTGGCCATACAAACGCAAAAACAAAGCACATTTATCCA
ATTTCTACGCGCGGTTGGATAGCGTGGACGATTTGTTATCCGAAAAGATGCGTGTAGTTTAAAGTACGACTATGGGAG
AAAGATTGCTTACTACGGGTTAAAAAATGCCATCTGCGCTATCCTGAATACGAGCTTATGAAGATAAAGACATATT
CCTGAAAATCCATTTTTCATGAATTTTACTATATTAATAAAGGAGAAAATCCGGGATTTA'CTCATCGGAATAATCG
65 AATAAACCAACTGAAAGATAGTTATAGCACTAGCGTAGGTTCTGTATTAACGGTTTCCAGGTCAGATTAACCGGT
TTTTCGAAAAGCAGCAGCTCACACGACAGGATTTGGTAGGTTACCAACAAGTAGAGCAATTTGGTACAGAGTTTT
GTAACAATTCAAATAAAAATAAATTTAAAGGATCTTATT

SEQ. ID NO:11

Nucleotide sequence of DNA region (1000 bp) up-stream from the Omp85 gene from *Neisseria meningitidis* (serogroup B)

5 ACSTCCGAAACCGTGAATCCGCAAGCCGCGCCCAAAACCAAGCCCAAGCCAAATGCCGATATAGTTGGCATTGGCAAT
CGCTTTAATCGGGTTGGCGACCAAGTTTCATCGCAGCGATTCAACACTTCCACAAATGCCGAAAGCCGCGCGGACA
CATCGCCCGCGCCGCAAAACAATGTSOGTCCGGGAAACCATACCGCGATGACACCGCGTCCAGGGTGGCGAAACCGTA
20 CCAATGAGGTAAAGGATGATAATCGGCCTGATATGCGCCTTGTTCGCTTTTGGTCTGCGCBAATTTGGCCGCAACAA
AATAAATACCAAACCGGCGACCCGCTTGGCGCCCGCAACAGGCTGCCGAAACGAGCTGCCCGCCAGCCAGTT
GCGGGAAACCGCAACCGATTACGATGCCCCAACCCGCGGCAATCTGCCCTACCCAGSCTGACCGGCCGATCGCA
10 GAAATAAGGTTTGGCGAACGCAATTTCTTCTATGTTGTGATATGTTTAAATAAGTTTGTATTTTAAAGAAACT
CACTCTCGTGTTTTCTTATTTTTCGGCTGCTTTTAAAGTTGCGTTGATTTGCCCTATGCACTGCCGCGCAGCTTTG
CTTTATCAATTCGGCGCAAGCTTTAATTTATTGAACGAAATAAATTTAATTAATCTGCTTATTTCCGCACTATTC
GAAACGACCTGTTTCCATATCGGATTTGAAACAAAATACCTTAAACAGCAGATACATTTCCGGCGGCGCAAC
CTCCGAAATACCGCGGAGTATGCGCTCTGAAGTCTCCGCGCGTCCGAAACACAAACCAAGCCCTTCCAAACCT
15 GTCCGAAACAGTTTGAATCGAATCTGCCACAGATGACACACCGCTACCATGATGATCAAAACCGACCGCCCTGCT
CCTCGCGCTTATTTCTTTCCGACGCACTACCGCT

SEQ. ID NO:12

Nucleotide sequence of DNA region (772 bp) up-stream from the PilQ gene from *Neisseria meningitidis* (serogroup B) (ATCC13090)

20 GCGATGTCGGGAAGCTTCTCCGCAATCATACCCCTTGGTTCGCTGAAATCGCCCAATCTCCGGAACCGGCGCAAT
CATGACGCGAAGAGCAGCATCTGAACCTCAGTGCCATGCGCACCACTACCAAGCAAAATCGTAGAAGAGCTTCCGCG
AGAAGCGCAGAAATGCCGACAAAATAACTTACGTTAGGGAACCAATGAAACACTATGCTTACTCATCAGCTTCT
25 GCTCTCTCCGAGTTCCTCAAGCTTCTGAGGACTAACGAAATGGATGGCAAAACGCGACCGCAAGCCAAAGCAGAAA
TCACTCTTCCAAAGCCTACCCCTGCGGTTGCGCGGATACAGCCCGCGCAGCTTACAGGGCCGAAACGCAATCGAC
TTCGCGCCGATGAAACCGCAAAAAGGGGAAAATGCCCCGACCAACAGCTTAAAGAAACCTGGGAAAATTCAG
20 TTTGAAAATATGCGTATGTCGCGCATTTTGAAGTCTGGAACAGAAAGTCTCCGCTTCACTGAGGCTGAAAGTTATGCT
ACA CTGCGGTGCGCAACTATTTGGGACAAAATACGCTGAAATCGAAAGCATTACCGACGACAGCATGCTGCTGAC
GAGCTGATAGAAGACGACCGGCACTGGGTTCCCGTAAAGCAGAACTGCTGTTGAAATCTCCGCAAAAACCGCA
30 ACAAGCGCAGCACTCCGCGAGAACAAAATTAAGAAGAGGATTACTCCATT

SEQ. ID NO:13

Nucleotide sequence of DNA region (1000 bp) up-stream from the Hsf-like gene from *Neisseria meningitidis* (serogroup B)

35 TTGTGTTTTCTTTGGTTGTTGAAATGGTTAAATCGGGTTTGGGGCGGATGTCGCGCAICCGCCCGGTTTTTGGG
GGTTGGGGGTTTTCTGATAAATCCCCCACTTAAATCTGCTCATTTCCGGAAGCGGGAAITCGGACCTGGAATCT
AAGGAAACGTTTTATCCGGTAAGTTTCCGTCGCGGCTGCGATTCCCGCTTTTGGCGGAATGACCGCGGTGGGGTT
TCGTTTTTCCGATAAATCTCTGTGCGTTGCGTTTTGATGCTTTTGGCGAATGACGCTCGGTGGGGTTTTCT
40 GTTTTTTCCGATAAATCTCCGCGCTGCTTTCTGATCCCGCGGAAATGACCGGTCGCGGTGGGGTTTTCTGT
TTTTGCTGATAGTTCCGTTGTTTTTCCGTTGCTGATTTCCGCTTTTGGCGGAAATGACGCTCGGTGGGGTTTTCTGT
40 TTTCCGATAAATCCGTTGCTGTTGTTTTGATTCGCGCTCCGCGGAATGACGCGGTTGGGGTTTTCTGTTTTTTC
CGATAAATCCGTTGCTGCTGTTTTGATTCGCACTTTCCGCGAATGACGCTCGGTGGGGTTTTCTGTTTTTCCGA
TAAAGTTCGCGCGGTTTTGTTTTCTGATTTCCGCTGCGCGGAAATGACGCGGTTGGGGTTTTCTGTTTTTCTGATGA
TCTCGTGGTTTTCTATGATTCATCATCTGATAAATCCCAATATCTAAATCTGCTCATTTCCGCGAAGCGGG
45 AATCTAGGACGTTGAATCTAAGGAACTGTTTTATCCGTAAGTTTCCGTCGCGGCTGATGATTTCCGCGTTTTGCGG
GAATGACGCGGTTTTCTGTTTTTCTGATAAAGTCTCCGCGTGTGTTTTCTAGATTTCCGCTTTTGGCGGAAAT
GACGGCGGTGAGGTTTTCTGTTTTTCCGATAAATTCCTGT

SEQ. ID NO:14

Nucleotide sequence of DNA region (1000 bp) up-stream from the Hap gene from *Neisseria meningitidis* (serogroup B)

50 AATCAGCATAGTTGCCACGCGCGCTTGGGOSTTTCCACACAAAAGCTCTGCCATCGGCAGCAGGTTTTTCCCGGAT
ATGCGTATCACGCCACGCGCGCGCCCGGTTGCGGTAGCGACTGCCGCAATCGTTGGAACGTTATCCGACATAAAACC
CCGAAAATTAACAAACAGCGCGGATATAGCAATGCGCTCTGAAGTCCGCGGTTTTGGCTTTTCCAGCGGATAAAACCG
55 CAAAAATCTGATAAATCCGTCGCGCTGACTAATATAACCATATGGAAAAACGAAACACATACGCTTCCGCTCGGT
ATAGGCTCGCTGCTGTTTCCATCCCGCAAAAACCGCCATCCGCCCCAATCCGCGCAGCATCAAAAAACATCGG
CGCGGATTTTCAAAGCGCATAGAGAAAGCGGAAAATGACCGAAAACGCAAGGACGCGCGGCGGCTGCGAAAC
CGTCGTAATCCCGGAGCTTGTCCAGCAATCTCTCCGACGACTGCTGCAAAATGATAGCCGCGGTTTTCCATT
CGGATCGTTGCGCGCGCTCGACTTGGCGCAATACAAACGACTATACGCAACGGGCGAGCCGCTTATGGCAATG
60 GCGGAAAAGCAACGCGCTCGGACGAAACCATACGCAAGCAACACTTCAACAGGCGCGGCGCAATGTACGCGTT
CATGAGCTATCCCTGATCTGTTTTTTCGCTTCTCTGATGGAGCGGTAACCCGCAACCGCGCTCCCTTGGCG
CGGCGACAGTGTGCTTGGCGGTTCTTCTGATTTGAAAGAACCGGACCAAGGCAAAAATTAATGTCAAATCTTA
GGCGTGTCTCATATCCGCGGAAACCGCAACATATAGGCAATCCGCGCGCGCGGAAAGCGGAAAGCGCGCC
CTCCAAACAAACCGAATCCGTCAGATAAGGAAAAATA

SEQ. ID NO:15

Nucleotide sequence of DNA region (1000 bp) up-stream from the LbpA gene from *Neisseria meningitidis* (serogroup B)

70 GATTTTGGTCACTCCGCAAGCTTCTTCTGAAAGGCGTGAATTCCTTTGGTTAGCAAGAGAAACCATCAAGCTTGC
CGATGGCAGGAAATGACCGTCCGTGTTTTCGCACTTTTGAACCTATGTAAACTCGGACGATATAAAACCGAACCC
CGCAAGTAAACCAAGGCGAAGATAAAGGAGGATGAAGAGTGCAGGCGTTGGTACGTCGAAAGAGCGAAGGC
GAAGTTTCCGAAGATGAAGGCAAGAGCCGAAAGAAATCGTCAAGAGAAACCGAAGAAAGAGCTGAAGAGGAAAGGC
TGAACCAAGAAAGTTGAAGAAACCGAAGAAAAATCGCGACAGAAAGGCGGCGGTTCAACCGCATCTGCTGCTG

CCTCGGAAGCCTCTAAAGGCAGGACATCGACCTTTCTCTGAAAGSTATCGCACGGCGGAAGCCGACATTTCCAAGAACC
GGAAAAGCACACTATACCGGCACTTTGGGAGGCGGTATCGGCACACCCATTCATGGGCAATCAGGCGGATAAAGAAGC
GGCAAAGCAGAATTTACCGTTAATTTCCGGGAGAAATCGAATTTCCGGAAACGCTGACGGGAGAAAAAGCGTGTACAACCTG
5 CTTTCTATATTGAAAAAGCGCAAGATTGAGGGCAGCGTTTCCACGCAACAGCACGCACTCTGTGAGAACGGCATCAATCTT
TCGGGAAATGGTTCGACCAACCCAGAACCCTTCCAGCTAGTGTATCTTCTGTGAGAAGGAGGATTTTACGGCCCGCAGCG
GAGGAATTTGGCGGTATTTTCAATAAGGATGGGAAATCTCTGTGATAAAGTGAAGTACTGAAAATAAAGTTGAAGT
TGAAGCTGAAGTTGAAGTTGAAGCTGAAACTGGTGTGTGTAACAGTTAGAACCTGATGAAGTTAAACCCCAATTCGGCG
TGTATTCTGGTGGAAAGATATAAAGAGTTGGAAAA

10 SEQ. ID NO:16
Nucleotide sequence of DNA region (1000 bp) up-stream from the LbpB gene
from *Neisseria meningitidis* (serogroup A)

CGGCTTAGAGTTTGGGCGAGTAAGGGCGGTCGCGCTTAGATCTGTAAGTTACGATTCGTTAAATACTTTTACTGA
CTTTGAGTTTTTGGACTAAGGGTGAAGCACCTTACTGCTTAAAGTCCAAAGCAAAAACAAAGACAAAAACACTT
15 TTATTACCTTAAATCGAACCCATAAATGACCTTTTGTCTTTGGCGAGCGGAGTAAGGGCGGCTCCGCGCTTAG
ATCTGTAAGTTATGATTCGTTAAATAGCCTTTACTGACTTTGAGTTTTTGGACTAAGGGTGAAGCACCTTACTGCT
TACCTTCAATGGGCTTTGAATTTTGTCTGCTTTGGCTTGTCTGACTAAGGGTGAAGCACCTTACTGCGCTCGCC
AAGACGAAAGGGTTTATTCGGGGTGGATTTTGGCAGTAAGGGCGGTCGCGCTTAGATCTGTAAGTTATGAT
20 CCGTTAAATAGCTTTACTGACTTTGAGTTTTTGGACTAAGGGTGAAGCACCTTACTGCTTCACTTCAATGGGCTT
TGAATTTTGTCTGCTTTGGCTTGTCTGATCTAAGGGTGAAGCACCTTACTGCGCTCGCGAAGCAACGAGGGCTA
TTTACGGCTTAGAGTTTGGGCGAGTAAGGGCGGTCGCGCTTAGATCTGTAAGTTATGATTTGAAATAGTCAATTT
GCCAAGAAGCTTAAACCGCAGGACCTAAGGGTGAAGCACCTTACTGCGCTTACATCCAAGCACCTTACTGCGCTA
TCCAGCACCTTACTGCGCTACGTCACGACCTTACTGCGCTTACATCCAAGCACCTTACTGCGCTTACATGACATG
25 ACAGACCGGAGCAGCGGAACAGGACTAAAAACAATTAAGTATATTTTGGCCCACTATAATAGACATGTAATATA
TTACTATTAATAATAATAGTTTACTCTCTTTCATCCC

SEQ. ID NO:17
Nucleotide sequence of DNA region (731 bp) up-stream from the TbpA gene
from *Neisseria meningitidis* (serogroup B) (ATCC13090)

TATGAAGTCGAAGTCTGCTGTTCCACCTTCAATTAATGAAATACGGAATGTGACCGC
30 AAAACAGCAAGTCCGCGATGACGAGGAGGAAAGCAGTAGTCAAGCTGATGTAAAAAG
GAACAAAGTTGGACAAAGTATGTTCTTCAAGGGCAGCGCACCGATGAAAAAGAGATTCCA
AACGACCAAAACGCTGTTTATCGGGGCTTTGGTACGGGCATATGCCAACGCGCACAAAG
TGGAGCGCAATGCTTCCGATAAAGAGGGCGCAACAGGGCGGACTTTACTGTGAATTT
35 GGTACGAAAAAATTAACGCGACCTTAAACGCTGACCAACGCGCAGGCGCGCAACCTTTAGC
ATTGCGGGCATTTTGGGGCAACGGTTTTTCCGGTACGGCAAAACCTGCTGACTCAGT
TTTGTCTCGATCAAGCAATACCGCGCAGCTTAAAGCAATATATCAAAAGCGCAAG
CTCCAGGGCGGTTTTTACGGCCCAAGCGCAAGTTGGCGGATGTTTGGCTTTTCG
GAGATAAACAAACGAAATGCAACAGATGCTTCCGCAATGGAATTCAGCAAGCAAT
40 GCAATGCTGATTTGGTTCGAAACCGCAAGCGCTGCAATTAAGCAGGTTGCCGAAC
ATCAAGAATAAGGCTCAGACGCGCACCGCTCTCCGATACCGCTGAAAGCGAAGAT
AGGAAACACT

SEQ. ID NO:18
Nucleotide sequence of DNA region (373 bp) up-stream from the OmpA gene
from *Neisseria meningitidis* (serogroup B) (ATCC13090)

CGTACCGCATTCGCACTGCAAGTAAAAAGTATTGAAAGCAGTCAAGCAGGCGATAAAGCTGCCGACAAAGCGGTTTA
45 CCAAGAGTCCGCAAAAGTCAATCGACCGCATCGCCGCAAGGGCGTGTCCATAAAAAACAAGCGGCTCGCCACAAAACCC
GTTTTGCTCAAAAAGTAAAAACCTTGGCTTGAATTTTGCAAAACCTGCAATCCGGTTTTTCACTGCTGATTCGAAAAACCC
50 TGAAGCCGACGGTTTTGGGGTTTTCTGTATTGGGGGCAAAAATCCGAAAATGGCGGAAAGGGTGGGTTTTTATCCG
AATCCGCTATAAAATCCGCTGAAAAACAATATGCGCAATGGGGTGGAG

SEQ. ID NO:19
Nucleotide sequence of DNA region (1000 bp) up-stream from the Plal gene
from *Neisseria meningitidis* (serogroup B)

TTTTGGCTTCCAGCGTTTCATTGTTTTCTGACAAAGTCTAAGTCAAGTTCAGATTCAGATTTG
CTTTTTAAAGTCTTCGACCGTACTCTCATCAACATAGTTTCGACCAAGTTGATGATGTTCA
55 GAGTATCGGTGGCAGCGGCTTCGGCATTTGGCAGCAGACGCGAGCGTTCTGCTTGAAGTTGCA
CGCGTTTTTTTTGCTTCCGCGCAGGCTTCCAGACAGCGCGCGCCAAACCGCTAATA
60 CGAATTTTTCAAGCGGCAAGTTCTGATGAAAGGTTGCAAAAAAGAAATCCCGC
GCCCCATGTTACCCCGGCAAGGTTGGCATTGTAAGTAAATTTGCGAACTCAA
AGCATATTGACTGATTTCTAAAAAATATCTGTTTTCAAAAAGGGGAGAAAAAGT
CGCCCGATTTTGGCTTTTTTGGCTGTCAAGGTTGTCGACGGGCGGATAGAGAGAAA
65 AGCTTGCATATAAGTAAACCCCTTAAATTTGCGGTTTACAGAAATTTATTTTCTT
CCAGGAGATTTCAATAGGCAACAGCGCACAGCGCAACGTTGCCGCGCCGCTCCGTC
AAACAAAGCGCCCAATGCTAGCTGCTGCTACCGCATTCGCAACCGCAGTGA AAAAAGTA
TTGAAAGCAGTGAAGCAGGCGGATAAAGGTTCCGCAACAGCGGTTTCAACAGATTCGCT
AAAGTCAACGCGCATCGCGCAAGGGCGGTTCCAAAAAACAAGCGGCAAGCGCCAC
70 AAAAGCGCTGCTGCAAAAGTAAAGGCTTGGCTGATTTTCAAAAAGCGGCAAGG
GGTTGATACGCGATAAGCGGAAACCTGAAAGCGGCGGTTTTGGGGTTTTCTGATTC
CGGGGCAAAATCCGAAATGGCGGAAAGGGTGCATTTTTATCCGAATCCGCTATAAA
ATCGGTTTTGAAAAACAATAGCGCAATGGGGCGGAG

SEQ. ID NO:20

Nucleotide sequence of DNA region (1000 bp) up-stream from the FhaB gene from *Neisseria meningitidis* (serogroup B)

5 TACGGAACTGCAAGCGGATCCGAAAGTTAGAGCGTGCATTTTCGGTCCCGTAAAAAATGGCTGTTTCTTTAATC
ACAATGGACATCGTACCACGAAGCAACCGAAGGCTATCCGTCATATACGATTACACGGCAACAATATCAGAGCAGC
CTGGCCGCGAGCGCATGCTTTGGCCTAACAGACTTCATAAACTCAGTCCGAATGAAATATGGACAGCCAAACCTA
10 TAAATACATGAGATGCCGAATCCAAAGTCAACCGCCGCTCAGAGCTGGGAAGCCGAATTCGCCACCGTGCTT
ACCTCAACCGTTGGCAGCTTACGGCAAGTTGCTTACAAACGCGGACCGCATGCGCCAAAGTATGCTGCACCGGAA
GAAACGCGGCGATATCTTCCAGGTACATCTCGTATGAAAATCATTACTGCGGTTTGGAGCAGCGCCCATTTAT
15 TTTAGCAAAACAGCAGTCTTTACGCAACCGCAATCAAGCTCAATGGAACAAAACGCGCTTGGTGGCCAGATAAAT
TGTCATCGGACCGCTACACCGTTCCGGATTTGATGGGAGCAGAGTCTTTCCGGAGAGCGAGTCTTACTCGGCAG
AATCTTTAACTTGGTATTTTCAATCGAACCATCAGTCTATCTCGGTGCGACTATGGCCGCTATTTGGCGAAAGTGC
ACAATATGATCGGGCAAGCAGTGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGT
ATGATCTGTTTGGCCGCAAGCGCTTCAAAACCCAAAGGCTTTCAGACGACCAACCGCTTACCGCTTCAACTGAAAT
20 TACAGTTTCTAACCTCGAATTTTACTGATATTTAGACGGTCTTTCCTTATCTCAGACCGTCAAACTTACTAGCT
ACTTGGCGCAGTACGTTTCAATTTCAAAATGGAATAGAC

SEQ. ID NO:21

Nucleotide sequence of DNA region (1000 bp) up-stream from the Lipo02 gene from *Neisseria meningitidis* (serogroup B)

20 TTATCTTGGTCAAAACTTTGTCCGGGTCGGACTGCTACGGCTTTGGGTTTGGACCCGCTCATCGGTCTGATTACCGGT
TCGGTGTGCGTACGCGCGGACACGCTACGTCAGGTGCGTGGGACCTAAATTTGAAACGCAATACGGCTTGGTCCGCGC
AACCGGTTTGGGATTTGCTACCGCTACTTCCGGCTGGTGTTCGGCGGCTGATCGGCGGCGGTTGGCGCGCGCTGA
25 TCAAACAAATGGCGCAACCGGTTGAAAACAAAAACAGGATCAGGACGCAACCGGACGAGTGTTCGAGCAGGCA
AAACGACCGCGCTGATTAACGGGAAATTCGCCGTTGAAAACGTTGCCATGTTGGCGGCTGTTGGCGTTTGGCGAGAT
TATGAGCGGCTTGCACAAAGAAATATCTGTTCCAGCTGCGCAAAATTCGTTGGTGTCTGTTTGGCGGCGTGGTCAATCGCA
ACATCTCAGCTCGCATCAAGGTCAATATGTTCCGACCGCGCATCGATGGTTCGGCAATGCTTCCGCTTTCGCTTTTC
20 TGGCAATGGCGGTTGCTGAATTTGAAACTGTGGAGCTGACCGGTTTGGCGGCGCTGTAACCGTATCTTCCGCTACA
AACCGTGGTATGTTTGTACGCGACTTTGTACTCTATGCTTATGGGGCGGACTATGATGGCGCAGTATGGCTG
30 CCGGCTATGGCGCTTCCGCTTGGGTTGGGTTGCAACCGCACGCGCGTGGCAAAATATGCAATCCGTCACGATCTTCCGCGCG
TCGCAATAAGCGTTTTGATGTTGCTATGGTCCGCGGCTTCTCGTGGATTTGATTAATGCGCGATCTCACCGGTT
TGTGAATTTCTTAAAGGCTGATTTTCGCTTCCGCAAAAGCACCTGCAAGGTTTACCGCTGAGGTCCTTTTGTCTA
TGAATCCGCTACGCTGCAACCGTTTGGAGGAACATC

SEQ. ID NO:22

Nucleotide sequence of DNA region (1000 bp) up-stream from the Tbp2 gene from *Neisseria meningitidis* (serogroup B)

40 CCTACTCCACGATTCAAATGCTGGCGGACCCACGAAGCCAAAGACTTGGAAATTTTGAACCTGGGCAATCAAAATC
GTCAAACCATATATGGGCGTTGCTTTTGGGACGAAACGTTGAGTACAGCCCGAAGAGTCAAGCTGGCGTTGAAAGA
AGCGTCCGCTGCTACATGAAAGCGCAAAAGTACCGGACCCCGTGAATCTTCTTCCGAAAGCCAAACCGCATCGGCGCC
GCCACGGCTTGGGATGAGCGACCAAAATCGAAACCGCATCATCGAAGCCAAATCGGCGGATCTACGAAAGCCCGGTT
ATGGCGTTGTTCACATCGCTACGAAACGCTTGGTACCGGATCCACAAAGGACCAATCGAAACATACCGCATCAA
45 CGGCTCCGCTCGGCGGTTGCTTACCAAGCGCTGGTTCGACAGCCAGCCCTGATGTTGGCGAAACCGCCAAAC
GCTGGTCCCAAGCGTTACCGGCAAGTACCTCGAATCGCGGCGGCAACGACTACTGATTTCTGAAACCGAA
TCCGCAACCTGACTACCAACCGAAACCGCTGAGTATGAAAGTGAAGTGGCGGCTTACCGCGCTCGACCGCAT
CGACAGCTCAAGTGCAGCAACCTCGACATCCGACACCGCGCCAAACTGGGCTACTCTCGAAAGCGGTTTGTGT
50 CGTGGCGGAAAGCTCGGATATCCGCAAGTTGGCAATAAGAAATAAAGTTGCTGTTTGCATCATTAGCAACTTAAG
GGTCTCTGAAAGATGATCCCTTATGTTAAAGGAATCCTATGAAAGAAATCAAAGTGTCTTATCAGGAAAGCCAG
TGTCCAGCTGTTTTTCGGCGCGCAAGGTCACCCCGTCAATTTACGCGGCTTCTCAACAAACAAACCCCGGAAAG
GCTGGCGGTCGAGACTTTCGAATAACATAGGTTACTAA

SEQ. ID NO:23

Nucleotide sequence of DNA region (1000 bp) up-stream from the PorA gene from *Neisseria meningitidis* (serogroup B)

55 GAAATGACATTCATAGTTTTCCGAAATCCCAACATAACCGAAACCTGACAAATACCGTAGCAACTGAACCGTCATTCCC
GCAAAAGCGGAAATCCAGTCCGTTCAAGTTTCGGTCAATTCGGATAAATGCTGCTTTCATTTCTAGATCCACCTT
TCTGGAAATGACGCGGAAAGGTTTTGGTTTTTCCGATAAATCTTGGGCAATTTGAAATTCGAAATTCGCGCTGGCG
60 GGAATCAAGCTCCAGATCCCGACCGCTTTATAGTGAATTAACAAATCAGGACAGGCGACGAGCTGCAGAGT
ACAGATAGTACGGAACCGATTCACTTAGTCTTCAATCTTAGAATGCTTCTTTCAGCTACGGGACGCAACCTC
GTACTGTTTTGTTTCATCCACTATATATGACACGGAAGCGCGCTCCAAACATGCGCTGAAAGAACTACACA
GATACCGCGCTTATATTCAATCCGCGCCCGTGGTTCGAAACCTCCACACTAABAACTAAGAAACCCCTATGTTCC
65 CCGCAACGAAAGAGCTGCAAGTATCTCGTTTTGGTAATCAAAAACCAATATCCGCGGAAATACGCGCCGAAAT
TTTGAAGCGTTGCAACAAACATCCGCAACGACTATTCGTAATTCGTTCCGCAAGTTCACCGGCTTCGCGCTGCG
CCATGACCGGCGAGCCGACTTCGCCACCATCGTCAATCCGCTACATTCGCGACATCAAAATGGTGAAGCAATCCGTC
AACTTACTCTTCAAGTTCGCGCAACCGGCGATTTTATGAAAGCTGCGTCAACATCATCATGAAAGACTCATTTG
CCTGATGGATCCGAAATACATCGAAGTATTCGGGAGTTCACACCGCGGCGGCTGCGCATTCATCTTTCCGCAAT
70 ACGGCAAGCAGGACCGAGTTTGAAGCATTGGCGGCTAA

SEQ. ID NO:24

***Neisseria meningitidis* (serogroup B) PorA Promoter Region**

GATATCGAGGCTCGCCCTGAATTGTGTGTAGAAAACACACCGTTTTGAAAAAATAAGCTATTGTTTTATATCAAAATA
TAATCATTTTTAAATAAAGGTTGCGGCATTTATCAGATATTTGTTCTGAAAAATGGTTTTTTCGCGGGGGGGGGTATA
ATTGAAGACGTATCGGGTGTTCGCCGATGTTTTTAGGTTTTTATCAAATTTACAAAAGGAAGCCCAT

5 SEQ. ID NO:25

Nucleotide sequence of DNA region (1000 bp) up-stream from the PorB gene from *Neisseria meningitidis* (serogroup A)

gttttctgtttttgaggggaatgacgggatgtaggttcgtaagaatgacgggataggtttccgtgaggatggatctcgc
atccccgcgaggggggaatctagaacgtggaatctaaagaacggtttatccgataagtttccgtgagcaagtttgg
atccccgcctgaggggaatgacgggatcttaggtttctaaatttgggtttctgttttgaggggaatgacgggatgagg
ttcgtaggaatgacgggataggtttccgtgaggtggttcgtcattccccgcgaggggaatctagacctagaac
aacagcaatattcaagattatctgaaagtcgagatctagattccccgcctgaggggaatgacgaaaagtggcgggaa
tgacggttagcgtgctcgccttagctcaagagaaacgattctcaaggtgctgaagcacaagtgaaatcggttccgta
ctattgtactgtctgaggggttcgtcgcctgctgctgatttttggtaactcaactatctctgcccaggggggggttttg
catccgcgcttcgaaagaaacccgctgctggtttttgcccctttataacccccggttgcaatgccctccaataacc
ctcccgatgaagttgtaaaaatgcaaatctaaaaaattaaataaccatagttataaacaataataaccataat
atctctatccgtcctcaaaatgcacatgcaatccacacaaaacagcagaagttgttttcagacaggaacatct
atagttcaagacatgtaacgagccctcggcgttaaatgcaagctaacggcctggaaaagccggcctgcttaaa
ttcttaacaaaaaggaatcagcaatgaaaaatccctgattgcccctgacttggcagccctcctgttgacgacat
20 ggcgagcttaccctgacggcaaccatacaaacggcgtg

SEQ. ID NO:26

***Neisseria meningitidis* (serogroup B) PorB Promoter Region**

25 GTTTCTGTTTTGAGGGAATGACGGGATAGGTTCTGTAAGAATGACGGGATAGGTTTCCGTGCGGATGGATTCTGTC
ATCCCGCGCAGGGGGGAATCTAGAAGCTGGAATCTAAGAAACCGTTTTATCCGATAAGTTTTCCGTGCGGACAAAGTTG
GATTCCCGCCTCGCGGGAATGACGGGATTTAGGTTCTAATTTTGGTTTTCTGTTTTGAGGGAATGACGGGATGATG
GTTCTAGGAATGACGGGATAGGTTTCCGTGCGGATGGATTCTGATTCCCGCGCAGGGGGGAATCCAGACTTAGAA
CAACAGCATATCTAAAGATTATCTGAAAGTCCGAGATTCTAGATTCCCGCCTGAGCGGGAATGACGAAAAGTGGCGGGA
ATGACGGTTAGCGTTGCTCGCCTTAGCTCAAAGAGAAAGGATTTCTAAGGTGCTGAAGCACTAAGTGAATCGGTTCCGT
30 ACTATTTGACTGCTCGCGCTTCGTCGCTTGTCTGATTTTTGTTAATCCACTAT

SEQ. ID NO:27

Nucleotide sequence of DNA region (1000 bp) up-stream from the siaABC gene from *Neisseria meningitidis* (serogroup B)

35 ATACGGCAATGGCTTCAGAAAGCGATAGCCTCTGSGTGA AAAACCGATTTCCTGTGTTCTCCCAACCGCACCCATAGA
CGTAAAGGATAGGGATGGTAATCATGTAACACATCACCGCGACGCGAGCAAAATATTTTTCGCGGATTTGCAACTA
AATCTTCCAAAGCAACAGTTCGTACTACATTGCCAGGTGTCAGCTGCACATTCTGATCTCCACATTTTCGCTTGAACCA
CCTACCGCAGCCAGCATCCAAACACACGCTCACCGGCTGCGCTCAGCGGATACGCACACTATTTCCAGCAGCAATCAC
CGACACTTCCGCGATTATTTGTCACCAAAAGCACCACTCACTTGGCTGATTTGGCCATTTTTTCAGGCGGCTTTAA
40 TAATTTCTGAACTGACCAAGCGTTTTACCGACCCGAAATATGCGCAACAAACGGCACAGAAACCGTACCACTGCTCC
GTGACCAACTGCTCGCAACTTAGTTTATGCGCACTCCCGAGCCCATCGAAAGAAAGGCCACACCAACCAATCTGSC
CGGCGGCGCTTCCCAATCTAATACTCAATACATCACCAATTTTTCGCTGATCCAGCGAAGCACTGCGCAACT
GAGTGAATGATGATTTATCTCGAGCTTATATATTAATCTGAGCACTCCGCTGATTTCTCATCATCACTCCACTCCAGTA
45 ATTGAACTTCAGATTGTTCCCTAAGAGACAATTTTTTTGCGCTGCGGCTGATGAAAGAAATCGCAGCATCTTAC
ATTTAATCTCCACCAATTAATCTGCTGAGCAATATAAATTTTCACTTTAAACAAAGCAATCTTATATTAAT
TATAATGGCTAATTAAGCACTTAATCGAAATAATTTATGACTACGTAGAGTAAATTAGTATTCTTCTTCCAACT
TCCTTACTTATATATATATCTTATAGATTCTAAATC

SEQ. ID NO:28

Nucleotide sequence of DNA region (1000 bp) up-stream from the lgt gene from *Neisseria meningitidis* (serogroup B)

50 GCCAAAGCATTGGCGCGGATGCGCGCTGCGGAAACGCGCGCGCTTTGCCAAAGCCGACTTGGTAAACCGAAATGGT
CGCGGAGTCCCAGAACTGCAAGGACGATGGCAATACTATGCTGTTTGGACGGGAAACCGAAGAAATGCGCAAG
CGCTCGAGCAGCACTATCAGCGGTTTTGCGCGCAACAGCTGCGCAAGCAAAATGCGCGCGCTGGCACTGGCC
55 GCACAACTAGAAACCTTGGTGGCATTGGGGCATGGTCTGATTCGCAACGGCGCAAAAGCCCTTACGCTTGGCGCG
CGCTGCTGGGTATTTGCGTATGCTGATGATGATGTTGGACGTGAACGAACTGATTAGACGGCAATCGACAGCT
TCCCAAAAGTTTGTCAACGAAAAAGCCGCTGTAACCGCGCACTTATGACGGCGCGCTTGGCGCTTGGCTGCA
AAGGATTATCGCAAGACATCGTTGCGCGCTACTCGCCAAACAGCGCGCGCTTTGGACGATTGACCGCCAACTGCA
60 GGCCTTGGCGGTTCAAACAACCTGCCGCGCGCTGCGGAAAGGCAATTTCCAAACCGCTTGTCCGAATGCTTCCGTA
AAGCCGATGCGGAGTTGGCGCGGTTAAGCAAGCCTGTGCAACAGGACGAAAGAAAGCCCTTTTGGCGCGCGCA
GGCTTGCAGCGGAAATGCGCGCGCGCTGCGGAAAGGCAATTTCCAAACCGCTTGTCCGAATGCTTCCGTA
65 CAAAGTGCATGCTTCTTGAAGCGGTGATGTAATGGCGGAGATGCGCGCTTAAACAAACCGCTGAACTGCTGA
ACCGTTGCGAGCAATGAAAGCGGTGAGCGCATCGCGCTTTGGCGGAGTAAACCGTTGACAGTCAATGCGCGT
TGAAGCCTTCAGACGCACTCGTCCATCGGAGATAAA

SEQ. ID NO:29

Nucleotide sequence of DNA region (1000 bp) up-stream from the TbpB gene from *Neisseria meningitidis* (strain MC58)

70 CAACGACCGGATTTCCACTTTCGTTGGAAATGACGAATTCAGGTTACTGTTTTGGTTTTCTGTTTTTGGAAAAATAA
GGATTTCCAGCTTGGGATTTTACCGGAAAAACAGAAACCGCTCGCGCTCATTTCCGCGCAGCGGGGAATCTAGGT
TTCGCTGCGGAAACTTATCGGATAAAACGTTTCTGAGATTTTCGTTCTGGATTCCCACTTTCGTGGAAATGACGCG
ACAGAAACCGCTCCGCGCTATCCGCGCAGCGGGAACTAGACATTCATGCTAAGCAATTTATCGGAAATGACT

5 GAAACTCAAAAACTGGATTCCCACCTTCGTGGGAATGACGTGGTGCAGGTTCCGATATGATGGATTTCGTCAATCCCGC
GCAGGCGGGAACTAGACCTTCRACTAAGGCAATTTATCGGAATGACTGAACTCGAAAACTCGGATCCCACCTTTT
GTGGGAATGACCGGATAGATTTCALAAATTTTCTAAATAGCTGAACTCAACACACTGGATTCCCGCTCGCGGA
ATGACGAAATGCGAGTTACCGAATCTTAAACCAAGCGAAACCGAAGCAACTGATCCCGCTTCGTGGAAATGACGA
ATGATGATTCGTGGAAATGACGCGGAGGGTTCTCGCTTTTCCAAATAATGACCCCAACTTAAATCCCGTCAATCCC
GCGCAGCGGGAACTAGGCTCTCGGTGGGAATCTATCGGTAATAAGGTTCTTTAGATTTTGGTCTCGATPCC
CACTTTCCGTGGAAATGACGAAATGATGCTTCGTGGAAATGACGGGATATAGGTTTCCGTGGACCGCTTCGGATTCATG
ACTGCGGGAAATGACGGGATTTGGTGTATTCCTAAAAAATAAAAAATGATTTCGAAATTTGTAATAATAATAA
ATAAATACTTCAATTCCTTAATGAAATGGATTATT

10 SEQ. ID NO:30
Nucleotide sequence of DNA region (1000 bp) up-stream from the *opc* gene from
Neisseria meningitidis (serogroup A)

15 CAAAGGCTACGACAGTGGGAAACCGGCAACCTCTGGAAGAACATCAGTTGTTGSAAGCATTATGCGCAAGCCCTGCC
GCAACCGCTCCGCTGTGGAAACCGCAAAACCAACGCAACCGGATTTGTCGAAGACCGCTTATAGTGGATTAATTTAAAT
CAGGCAAGCGGACGAAAGCCGACAGACAGTACAAATAGTACGGCAAGCGGAGGCAACCGCGTACTGTTAAATTTAATCC
ACTATATGGTTCGAAACGAGCTTCGGTACCGTGCACCCATAATTCGCTATGCGGGGCGCCTATTCGGACTGATTA
AAGTACGTCGCAAAACCACTGAAAGCGATGTTGTTGAACTGTTGAAGCGCCTAACACTAAGTGGCGCCGCTCGC
GCTTAAAGAGGAGCCGATGCTGATTCGGTATCCGGGAGGCTTAAAGCGCTATTTGGTAAATTTGGAGGATTT
20 TGGCGGAAATGACGAAACCTCTCTGCTTTCGGCTTCGGCTCGGAGGGAAGSAAATTTGCAAGATCTCATCTGT
TATTTTCAAAAACGAAACCAAAAACGCAACCTGAAATTCCTCATTCGCGCAGGCGGGAAATCCAGACCCCAAC
GCGGAGGAATCTATCGAAATTAACGAAACCGGACGAACTAGATTCCCGCTTCGCGGAAATGACGCGAGTGGTTT
CAGTGTCTCCGATTAATCCGCTCTCAAGTCTCGTCTTCCCTTAAACGAAACCGAAATCGAAACCTAAATTT
25 TCGTCAATCCCAATAAAAACGAAACCAAGTGAAGATAACAATTCGTTAAACAAATAACTATTTGTAATTTTAT
AATATATGTAATAATCCCGCCCGCCCGCGAAGCTTAAGATAATAATGTAAGCTAAGCATTATTTACGTTATGTT
ACCATATCCGACTCAATCCAAATTTGGAGTTTAACT

30 SEQ. ID NO:31
Nucleotide sequence of DNA region (1000 bp) up-stream from the *siaD* gene
from *Neisseria meningitidis* (serogroup B)

35 ATAAATGACAGCGCTGAAGTGTAAACATCAACACACATCGTTGAAGCAAAATGCTGATGAGGCCAAACAAAGTCAAT
CCAGGCAATGACAGATGCTCTATTTATGAAATTAAGAACGTTGCGCCCTGAATGAAGAAGATGAGATTAATTAAGA
ATACTAGAGAGTAAAGGATGATTTTATCAGTACTCTTCTCTGTCAGCTGCTTACGATTAACAAGTATGGATA
TTCGAGCATATAAATCGGCTCTGGCGAATGTAATACTAACCATAATTAACCTGGTGGCCTCTTTGGTAAGCCTATT
40 ATTCTCTACCGGCAATGATTTATGAAAGCATCAAAAAGTCGGTAGAAATTAATCGAAGACAGGGGTACCTTATGC
TTTGCTTCACTGACCAACATCTACCCAAACCCCTTACGAAGATGTTGATGGTGGTATGAAAGATTATGAAAGCTT
TTCCAGACCAATCATTTGGCTGTCTGACCAATACCTAGATAACTATGCTTGTAGGAGCAGTAGCTTTAGCGGTTG
ATTTTAGAGGCTCACTTACTGACCGCATGGATCGCCAGTCCGGATATGTAATGCTCTATGAAATCCGATACTTTAA
45 AGAGCTCAAGCAAGCGCTCATGCTTTAAATTTGGCAGCGCGGCAAAAAGACAGATTAATCGCGGAGAAAGCCAA
CTAAAGATTTCGCTTTGCTATGCTGATGAGTAAAGACATTAAAAAGGAGAACTGTTGCGGAGATAAACCCTATGG
GTTAAAGCCCAAGGCAATGAGACTTCAGCTCAACGAATGAAACATTTATTTGTAAGTTCGCTGCTGCAATTTG
CAAAGGCTCAAAATCAAAAACGATATGAAATGCTTATTAACCTAGTACTTTATTAACAGAGGATGGCTATT
ACATATAGCTAATTCATTAATTTTAAAGATACAATA

50 SEQ. ID NO:32
Nucleotide sequence of DNA region (1000 bp) up-stream from the *ctrA* gene
from *Neisseria meningitidis* (serogroup B)

55 ATACTGCACTGATGTCGCGCAATAAATTAGCATGTTTCAATAAGCTAAAAATATTCAAATGGAATGGAAGGAAA
TGCATAAATTTATCAGATGATATTTAATAATCTTGCAGAACTTTCAGTCCAGTGTCAATTATAGGTAGATGC
TATGATATTTTGGCCACTTATCTATGCTTGGAAATTTGGCCGCAATTTGGGCAATTAATGTTGCTCTGATGTC
60 ACGGGGTGAAACATGAAATACCATAATTTGATGTTAAACCGTAAATTTCTTGAATCTTCTAAGATGGGAGGT
GGAAGGCGCAATACATTAATCGGGGAGCGATGATGTAATATGCTTCTTTTCTCCCAATTTGCACTAGGCGAG
TGACAGCTTGTTCATTTGCTACCAAGTGGATGAGAAAGTTACTAATAGAATGACGAATGGAGTCATCTACTGATCCA
GATAGTCCACACTTCGATATGGCAAACTAAACGGCTGTTAATGCACTACAGCTGCGCTGCTAGTGTCTTAAACG
65 GTCGCGTGAATCATGACCATATCAGTTCATTTTCAATCAGATAGACGAGATAAAGCTAATGTTATGCTAAAAGG
CACCCTTGGTTCACCTTGGATTTGATTTGAAAACGATATGATGTTGATGATTTTCTCGAGTACTTCTTGTAGGTT
CTGGCATATGTTTTCATCATATGCAATACCAGTTCAATCAATGCAATTCAGGCTGSGGTGATTTCAATATAGGCTAA
TAAAGGTTTATGCTTCCGAAAGTCCGCTCTGCTACCTGTAATGCAAGAATTTCTTTCATGATTTAGAATCTAAGTA
70 TATATATAGATATAGGAAGTTGGAAGAGAACTAATATACTACTACTCATAAATTTATTTGATTAAGT
CTATAATAGGCCATTTATAATTTATATAGGATTTGGCTT

75 SEQ. ID NO:33
Nucleotide sequence of DNA region (1000 bp) up-stream from the *lgtF* gene
from *Neisseria meningitidis* (serogroup A)

80 TCTTTTTCGGAATGAAAGGACGATCATCCGACATCGAGCGGCTGTTGTCGCGGAGCCAAAGCATAGTTATGCTTAC
GAGGCCATCAAAATACGCTGACCGATATGATGCTGCGGGGCGAGGGAAGAAATTTTCCGCTCCGAAAGTATGTTTT
CGACTCGCTTTATCGCCAGCCGCGCAACCGGCAACCGGAAAAAACCAGCCCATACGACCGGAAACCGGACGGGC
TGGTCACTCGGCAAGGCGCGGGATTTTCGTGCTGGAAGAAATGGAACACGCAACCGCGCGTGGCAATTTACGCC
85 GAACCTGTCGCTACGGAGCAACAGCGATGCTACCATATTTCCAGCCCGCCCGGCGCAAGGCGCAATCCTTGC
CTTTTCAGACGCTTTCGAACGCGAGACTTTGCGCCGAGACGCTCGGCTGATTAATCTGACCGGACCGGAGCGCAC
ACACGACGATATGAAAGCCGCGCTTTCGCGAACTATGCGCTGCACTGCAAGCAAGCCGAAACCGGAAACCG
GGACACGCTGCGCGCGGCGCAATCGAAGCGCGTTTCGCTGGGCAATGCTACCGGAAAGCAATCCGAAAGG
90 GAAACTTCGCGCCAGCTTTGGGACGGGAGAACGATCCGACCTTCCGCTACTCACTGACCGGACGCGGACCGCT

GGGAAACCGAAAAACGCATTGCCGCCAGCTCGTGGTTGCCCTTCGAGGAAAGCAACTGCGTTTTACTCATCGATGAAAT
AAGTTTGTCAATCCACCGCTATGCTATACAATACCGCCCTACTCTTATGGGTCTGTAGCTCAGGGGTAGAGCAGGGG
ACTCATATACCTTTGGTGGTGGTTCGAGCCACCGGACCCCAATTTCCCAAGCCCGGACGTATGTTTGGGCTTTT
CGCCCTGTGAAACCAAAATGCTTTGAGAACTTGGATA

5

SEQ. ID NO:34

Nucleotide sequence of DNA region (1000 bp) up-stream from the lgtB gene from *Neisseria meningitidis* (serogroup B)

TAGAAAAATATTTGCCCAATCATTAGCCGCGCTGTTGAATCAGACTTGGCGCAACTTGGAGATTTTGATTGTCGATGAC
GGCTCGACGACAGCTACGCTTGCATTGCCAAGGATTTCAAAGCGGGACAGCCGATCAAATCCTTGACAAAGCTCA
AAATTCGCGCTGATTCCTCTTTAAACATCGCGCTGGACGAAATGGCAAGTCAGGAATGGGGAAATATATTGACAGCA
CCGATCGCGAGATATTGCCGCCCGACTGGATTGAGAAAAATCGTGGGGAGATGAAAAAGACCGGACATCATCGCG
ATGGCGCGTGGCTGGAAGTTTGTGGAGAAAAAGGACGGCAACCGGCTGGCGGCATCACAGGACCGCAAAATTTG
GAAAAAGCCGACCCGACCAAGATATTGCCGACTTTTCCCTTTTCGGCAACCCATACACAACAACACGATGATATGA
GGCGCAGCGTATGACGGCGTTTGGCTTACAAACCGAGCGGGATTGGCGGAAAGATTACCAATTTGGTACGATGTC
AGCAATTTGGCGAGCTGGCTTATTATCCGAAAGCTTGTGCAAAATACCGCTTACCGCAATCAGGTTTACATCAAATA
15 CAGCATCCGCAACAGCAAAATCGCGAAGCATCAAATAACCGCAGAAACGATTTTTCAGTCTATGGTTTTAAAA
CCGGTTTCGACAGCTTGAATACCGCAAAATAAAGCAGTAGCGTATGAAATGCTGGAGAAACATTGCGGAAAGAGAT
TTTGAACGCGCCGCGTTTTGTACCAATGCTTCAAACGGACGGACCGCTGCCCGCGCGTGGCTGGATTTGCG
20 GGCAGACGCGAGGATCGCGCGTGTTCCTTGAAGCAATCTTCGGCATTTCGACCGATTGCGAAAAACCGTTGAA
AAACCGCGCTTATCCAACAGACAAAAACAGGATAAAT

SEQ. ID NO:35

Nucleotide sequence of DNA region (1000 bp) up-stream from the lst gene from *Neisseria meningitidis* (serogroup B)

GGCCACGGCTTTTCTTCATCGGTTGAGGGTCGGCAGGATAATCGGGACGGCAAGCCCTTAGACTGCAATCTTTAA
TCCGCGCGCTGATGAGGTACGGATGCGCTGATGTTTCGGCAGTTGATTACGTTTGCATCGGCTGTTTACACAGTTCG
CCCAATTCGGCAAGCGCTCGGTAAGCGCTGCGCTTCGGTCAGATATTCGGGAAATGCCGCAAAATACGCGCAGCAG
GGAAATGTCGCGCAGTTTGCATCAATATCGCGTGGCGGGCAAAACCGCTCGACAAATCGGCAGCAGGATTTGGTGGCC
30 TCCGCGGGGCTTCGTCGATGGGATAAAACAATGGTGGATTTTGGAGTATAGGATTTCTCTTGGAGTGGTGGT
TCTTTGGAAACACTTCCCGGGGAATGTCGCGCTATTATGGCATAATTCGGCGCTTGTTCGGCTTTGTTGGATC
TTGGCGTTTGAACCGCGCGCTGAAAGGAAAGGGGAAATGGTTCCCGCTTTGGCGCGCTGCTGGAGGCTGCTG
CGCTGATGTCGGCGCAATTTTCGGTGAATGATTTTATAGTGGTAAATTTAAACCACTACAGCGTTCGCTCGCC
35 TTGCTACTATCTGTACTCTGCGCTTCTGCGCTTCTGCTGATTTAAATTTAAACCACTATATATTCGCTAATCTG
TCGAAATATCTGTAATAATTCGCAATTTTCCGCTCGGCACTCGGGCGTATGTTAAATTTGTCGAAATGGAGTTT
AGGGAT

SEQ. ID NO:36

Nucleotide sequence of DNA region (1000 bp) up-stream from the msbB gene from *Neisseria meningitidis* (serogroup B)

CGCCGACGGGAAACAGACAGTCTGGAATCAACCGCTTGGACAGTACGGCGGCAATACGACATGCTTGCAGGTTATC
TTGAAAGACTTCCGCAAAAAACCGACCGTTGGCGTGGCGCTACCGCAAAATGCCGCTGAACACCGGATATCTCTTT
GAAAGCGCATATGCCCCATACCTTCCCGATATTTCCCAATGATACAGCAAAATTTGGAACAATATTTCAAAGACT
45 GAACGGTACGAACTTGGCGCGTGTACGATATGCTTGCATCAGGTGGAATAACCGCTGCTGTTGTTGTTGATGGAAC
AATGGCGCGCAACAGTCCAAAGCTCCGTATGTTGGACTGAAACCGCAATACTTGCCTAAAAAATGATTTCAAAC
GGTTTGTCTGAAATATGTCGGCAACCGTCCGTATCTTGGGTAATGACCCGGGCACTGCGTAAACGGGTTTCGGTTCATC
GATGTCAGGGGGCGGATCATTTTACGTGCTTCCGGCTGCAATAAAACCGCTGCGGATGCGCTCTTGGCAGACAGGAT
TGCCGTGATTTGTCGGCATATCGGGAAGTCTTACCGTTTAAAGCTCAACAGGCGGCGAGTGAACAGGTTTCTGTC
50 ACCTCAATCCGCAATCAACGCTGATGCTCGGTACGCTAGGGCGCGGCAATGGCGGCAATGCTCAGCCATAAGCTGCC
GTTTTCGGAATACACCGCTTGCAGGTCAACAGCGCTAGTTCGGCAAGGGCAAGCGGCAAAAGAACAGGTGACGATAT
GGTGTGCGAGTGTCCGGGCTTTCGGGAACCGCGAGGANTGGCGGCGGACGCTTGCCTGCGCTGACCCACCGCTT
ACGCAACCGCGGCTTCCGCGCAACTCAATCTTCCGGGATGACAGTCAAGCGCGGAGGTTTCAATAGTTTCAGACGG
CATTGATTTTTCGGCTCGAAAGAAAAATGATATCGAG

55

SEQ. ID NO:37

Nucleotide sequence of DNA region (1000 bp) up-stream from the htrB gene from *Neisseria meningitidis* (serogroup B)

CGCCCAAGCTTTCCCTTTCTCGGCTTAACATTTGCTTGTACGGCAGACTTTTCCCTTCAATACGGCGCTTCC
GAAAGACGATGAGTGGCGGACGATATCTCAACCTTAAAGTACGGTTGAGCAAAAGTTTCTCTTTCAATCAACCT
60 GCAACTTTTCGGCTACCGAGTGTCTCGTATGTTTGGCGCACTACGCCCTTAAAAAAGCGGACTTTTTCGATG
CCGCTCTAGGTTTCAAGTAAATTTACCTTATAAAGAGTGTACTTACCATACTTCCCTTAAAAAGATATAAAG
CGGACAGCTGTAATAACAAGTATGTTGTCGGGAGACTTCTTCAACAAACAAAAAGTTCTTTTATAGATTTCTCGCTTA
TAGAACAATGAAGGCTTAGCATAGGCTCCGGTAGGCTATTTCAACGGCTGTTTCAACGGCTACGCTAAAACTACGG
TAGAACCGGCTTCTGGGTTTCCGCAACGCGCGCTTTGGAACAGTTGTTGTCGGAACGCAATAACCGCCGCTTTA
65 ATGGTGTGGCGGTTTCACTGATGATGTTTCAAGCTGCGCTTTCGTTAGTTCGTTGAGCCGATGTTGAGGAGCTC
CGAACTACGGTTCGCGACGCGCAGCGCACATGATGCTGTTGTTAGAGGCTGTAGCGGTTCCGCACTTGTCTCCG
CTTCCGTAAGTAAATGTTCCGACCGCTGTTTCAAACTCAACAGCGGATACGAGCTGCTTTGGGGTGGGACG
CGGCAACGGTAGATAATGTCGGTGGCGGACTACGTCGAGTTTTCGCTTAAATGCGTTTCCGCGGAGGACGGAACCGACG
70 CAGGGTGGCTTTTCGGGTTGACTGGCAACAAATGCTATAGGCGTTCATTTTTCGTAACATGCGCAGCAGGAG
AGATACGTTGCTGGGCTTTAGCCAACTTCTCAACT

SEQ. ID NO:38

Nucleotide sequence of DNA region (1000 bp) up-stream from the MltA gene from *Neisseria meningitidis* (serogroup B)

5 CACAAAAACCAAGTTATGACGGGAATAAGGTACAGCAGCCAAACCAAGGCCCTCGCCCTGC
GTCGGATGGTCCGATAGCCGAAAAATCCGCCGAGCAGCAGCCCAACGGGCTGCTCTCG
TGCAGATATTTTGTAGTGAAGCAACAATGTCCTGAAGCGCGTCCAAATGCTGCTCG
TGCAGCGCAGCAGCGAAACCGGCAAGCAGACAGCGGCAACGATAATCAGAAAACGGCCCT
6 TCCCAACGGAAAAACCTTCCAGATTTCAGGCGCATCCACCTGTAAATCAACCGGCCA
ATCACGGCGCAGCCAAAAACCCCGCTACCGCACCGGCCGCTCTGCCAGTCCGGGCTC
TGTGTAATACGGCAGCAGGAAAAAAGCTCTCCAAACCTTCGCGGCCACGGCAAGA
10 AACGCCATACCGAACAAAGCCATCCTTCAACGCTGCCAGCTTCAAAACCGCCGCA
GAATCCTGAAGCTCCGCTTTCATCGAACGGGCGCTTTTTCATCCATAAAATCATATA
GTCCAGATCGCAGCAACAAACCGATAATGCCAGCAGCAACTCTGCTGCTTCTGG
GGAATCTCGCCGTTCCGAAATGATCCGTAACCCAGCCCAACACATCAAAAGAGCA
15 AGAACAAACCCGACACAGACTTAGGCATCAGTTGGAATGTCGGGACTGTTTCAGAAAA
CCGGCAACGATGCCGACGATGAGCGGCTTCGATACCTCCGCGCAACATAATAAAAAA
CGCACAGCATAAACCGCAACGAAACAGGATGATGAATAATATATATTCGGAATATTTTC
ATTGCTGTAATAACAAATGCAAGTATTTTATCTGCAGTACCGCGCGGCGAAAGTTC
CGCAGCTCAGCTCGCCCTGTGTAATAATCCCTCTCCACGGCTCGCGCAACCGCCGCC
20 GAAACCATCTTCTTATTAATCGCGGCAACATTGTCCATT

SEQ. ID NO:39

Nucleotide sequence of DNA region (1000 bp) up-stream from the ompCD gene from *Moraxella catarrhalis*

25 GCTGATTTGTGAGCAAGCGGGCGCATCAGGGATTACCTTGCATTTGCGAGAAGATCGTCG
ACATATTCAGATGAAGATGTTTATGAATTGATTGGCAATTGACAAACCGCATGAATCT
TGAGATGGCAGTCACTGATGAGATGCTAAATATTCGCCAAAGGTACGACACGATGGGT
GTGTTTAGTACAGAAAAACGCAAGAGCTGACTACAGAAAGTGGGCTGTATATCGCCAA
TTTATCAATATTCAGGATTTATACACAGTCTTCAGCAGGCGGATATTAAGGTTCTTT
30 ATTCATCGATCCAGATCCGATCAAAATGATGCTGCAATGCTTGGTGCTGATCGGAT
TGAGCTGCATACGGGAGCTTATGCTCAAGCGACTTTCACAAAATAATCAAAAGCTGTGTA
TAAAGAGCTTGACCGTATTCAAAAAGCCGTGCAATGGCAGAAAAAATCATCATTATT
GATTAATGCAGGTCATGGTTGACGCGTGATAATGTCGACGCGATTCGCCAAATGATGG
TATTCATGAGCTGAATATCGGGCATGCTTATTCAGATGCGATATTTATGGGCGTGA
35 TAATGCAGTCAAGGCAATGAAAAAGGCTTTTATTCAGATAAAAACGACCAATCATTTGATG
CGTTAGAAAAGAAATCGTAAATAATGATGACTATTTGTAATATTAATGTAATTTGTTCA
AAAAAGGTTGTAAAAAAATTCATTTACCATTAAAGTAAAGCCACAGCCACAAATGAATA
CCTATTGGTTGACTCATTTAGTCACTAAGAACTCGCAAAATTTTAAACAGATTAATGGC
40 AGGCTTGGATCGCTATGCTAAATAGGTTGGGTAATCTTGAAAAACCAACCATTCCTTC
GCAATTTTATGAAAAGGATATAAACCTCTTCGCGTCACTCGCAGCGGTTGCGCTCC
AGTTGCAGCTCCAGTTGCTCAGCTGCTGTCAGCT

SEQ. ID NO:40

Nucleotide sequence of DNA region (1000 bp) up-stream from the copB gene from *Moraxella catarrhalis*

45 GATGCTGTTAAAGTGGTATTTGGTCTGTTCTATTGTAACAACCGTATTTGTTGACGGC
ATTGGGCTCCCGCAGATAAGTGCCATTGATAGTGGCAAGTGGTAAAGATCGCAT
CCTTTGATTCCGATGCGGTATTCGTTTTCCGGGTGATATCGCCAAAGCCATCGCAGCA
GGCGCTTCATGATTTAGTGGGTAGCTTGTGGCAGGTACCGAAGAACACCTGTGTGAG
50 GTGGAATATTCCAAGTCTGTTATTAAGGCTTATCGTGGTATGGCAGCTTGGGCGCA
ATGCTGGTCAAAATGGCTCATCGGATCGTTATTTCAAGATGCCAAAGATGGTGTGAA
AAACTGGTTCAGAGGATCGAAGGCGGTGTTCCCTATAAAGGCCGTGGCAGGCATC
ATCGTCAATTTGGCAGGTTGCTAAGATCATCCATGGGTTATACAGGTTGCCAGACCATC
GAACGATCGGTAAAGATACAGCTTTGTCAAAGTGAATTCAGGATGAGGAAATCG
55 CATGTACAGGATGACAGATACCAAGAGACCCCAATATCGCCAAATTAACCTAT
TATAGCAAAATACAGCACTATAGATAGGTTGGGTTGCTTTTAGAGCATAAAAAATA
ACTGACATGACTTATGCTATTTTTAAATGCTTTAATTTAGATTTTAAATTTAG
ATATGCTAAAAATACAGATATTAATTTAAAGTTTTCAAAATCAAGGATTCATGTA
AATTTGAAAAATAAACAATAATCTGATTTATTTAAACCAATATTCATATTCAT
60 TTACAAAGAAAATTTTTTGTAAATTTCTACTTACCTTGTCTATTTTTCTTATTT
ATCATTTTGGCGTATTTCTGTTGATTTTAAAGTAAAGTAAAGTAAAGTAAAGTAAAG
AAAAACAAATTTGATTTCAAGCTCATTAATCGAGTTATT

SEQ. ID NO:41

Nucleotide sequence of DNA region (1000 bp) up-stream from the D15 gene from *Moraxella catarrhalis*

65 AAAACTGGTGTGCTTCACTGCTATTCATGGTGAACCAATCAATGATTGGCTAAGTGCC
ACCAAGATTATTACGGCAAAATCCAGAAAACCATGCTGATGACAGTCACTCGCTCAAGGT
AAGCAGGTTGATTTAAATTAATGCCCCGGTGGTAAAGACACAAAACGGCTAGTCGGT
70 CAATCGGATTCGCCCCAGATTGATTCGATACGCTCATTCCTGATGAATATGATG
ACGATTCAAATGATGTCGGTGAAGCATTACTCAAGCCATCCGACGAACCTATGATTTA
TCAATAATGACCTTAGATGCGATGGTAAAGATGATACAGGATGATGGCATGAAAT
CTATCAGGTCCTTGGCATTGCGGATGTTCTAAGACCACTTTGAGTTGGGATTTCAA
GAAGTGTATCCGACAGCCCAATCATCAGTTTAAAGTGGCAGTACTGAATCTTTTACCA
ATTCCAGTGTAGATGGGGCATTGGTATTTATACTTATGAATGATATGGGCAAA

TCTATGAATGAAGCGGTGCAGATGCCAGCATTAAAGCGGGTGGCTTATGCTTTTTTGT
TTCATGTTACTTGCATTCAGTAACGATATCAATGCCGATTTTGGCTAAGTTCGATTTAT
CGTACCCTTAAACAAATTTTTGGCTTTTTAAGCTGAATACCTTGCACAAATTAACCTTT
TGGCTTACCTTTACACATATAAATTTGGGTGTAGAAAATTTGSGATACATTTTATACC
5 TATTTTTAGAAAATTTAAAAAATAAGTTGGATGACCTTATGCCGTAATCATATTTAA
AGTTTTTCAGGTCAGTGCATGACAAATGGCTCATGATGGTAATCTCAACTCATGCACA
AGCGCGGATTTTTATGSCAAATGACATGCCATCACAGGACTACAGCGAGTGACCAATTA
AAGCTTACAAAGCGTGCCTCCGTTTCGCTTGGGTCAAGT

10 SEQ. ID NO:42
Nucleotide sequence of DNA region (1000 bp) up-stream from the ompA gene
from *Moraxella catarrhalis*

ACTTGGCGAAAATACCATTATATCGATTGTGATGTTATACAGGCAGATGGCGGTACACG
CACAGCCAGTATCAGTGGTGTGGGTGGCACTTATGATGCTTTAGAACACTTGCAGCG
15 TCGTAAAAAGCTTACCCAAAGATCGCTTTTGGGCTTGGTGGCAGCGTTTCGTGGGTGT
TAATCAAGGCGGTGATTTGCTTGTATTGGATTGCTGAAGATTCAACTTGTGATACCGA
TTTAAATGTGGTCATGACGCAAGCAGGTGGGTTATTGAGATTCAAGGCACAGCAGAAGA
AAAGCCATTACTCGTGGCTGAAGCTAATGCGATGCTTGTATTGGCAGAGCTGGGAATGG
GCAGATTATCGAAGCCAAAAGCAAGTATTAGGCTGGTGTATGCTAATCGTGAAGATA
20 ATGCGCTGATCATACATTAATGSCAAAGTAAAGACCCATTATTTGGTGGTGCATGTA
TATGCTCCTGCTGGGTGCTTGGTGGCAATCATTTGTTGATTGACCCGTTTTTTATG
CAATCGTGGTGGCTTTATTGCAATTTGGTATTTGGTATTTGCTTTTAAATTCAAAGCCAA
AAGCCAAACTTGTCAATGTTTTCCAAAGTCCGTTGAAGATTACHTCCAAAGCCCTTG
AGATTCNTAACAAATCACTAACCTTATCAGCATCGGCAACATATCTGCTAAGATPAAC
25 AAATGACAAATTTGATCGGGCATTGAATATCATTTTACGGTTTTGCTGATGACCGTG
AAATTAATATAGCCAAACAGTACTTTGGGAAAGTCAATCAAAACCAATGCGGTGGCG
TAACATTGGCTAAGTATGTTGGTATGACAGCAGGTTGGATGGCTTTAATCTCACCCA
CCTAACTTTTTCTTTGTTGATTTAAGATATGTTATGATGGCAGGATTTATTTTAA
GTCATCATTTAATGCAATCAGTTGTCAGAGTAGCCGTTTC

30 SEQ. ID NO:43
Nucleotide sequence of DNA region (1000 bp) up-stream from the hly3 gene
from *Moraxella catarrhalis*

GTGATCGGCAACACCCACCATTTCAGGAGCAACAAAATGCCCGTGCCTTGCCTTCTT
35 GGTGATACATTTGGCAGGCAATGTGGCTAAGTAGTGGTGTGCCATCAGGTGCGTGGT
GGTGGTGTACGATTCGTTATGTCATATAAATATATCCTTTTGGGTTGGATGATATCAATG
AAATACCTTACGGTTGATGGAAATTTATCCATTGTACCAGGATTTGGTCTTTTTAAAT
TAACACGCGCTTCTAGCAAGTCAAAGTTTTTATGCTATTTTTTCAGATTTTAAGGTAC
40 AATTAAGCCAAATGTTAATAATATGGTATTGTCATGATTTATGATGAATGGCACAAJA
TTTTGGGAAAATATCCCTTAGATGCTTTAACAGATGCTGAATGGGAAGCATTAATGAGC
CGATGTGGCCGCTTGTGTTGTGAATTTCTGATGATGACAAATGTTAAATGACCGAA
TATACCGATTTGGCTGCCAGCTATTGGATTTGCTCAACAGGATTTTCCCAAACTATGCC
AAGCGTCAACAGTTTGGCAGATTTGATTCGCTTAACACTGATATGCTGCTGATATG
45 CTGTGGTTGCCACGCCATTGTGCTTATAAGCGTTGTATCTTGGCAAAATCTGCCAGCA
TGGCAGCGCTCATTAACATAGCCAAACCTGGTGCAGGATTTGCCAAGTTTCAACT
GCTGGCGATGTGTGAGTGGCTTGTATGATGATGAAGACATAGAAGGCGAGTGGTG
AAATGGTTAAACCTTGCATGATTTGTGATGATGACAGCAATAAAAATTTGCCAAA
TTGATAAAATTTGGTATGTTGATTTTATCAAAAGCACTTGAATAAAAACCGAGTGA
50 TAGCTAAAATTTAGCAAAACCAATCAATCATATAAATTTAATGAACAGGTTAAAT
TATACTGCTATGCTGATGACAAATCAAGCACTTGGTGC

SEQ. ID NO:44
Nucleotide sequence of DNA region (1000 bp) up-stream from the lbpA gene
from *Moraxella catarrhalis*

55 TAACAAAGCAACCCACAGCGAGTTATTTGTGCAAGCGGTCAAGCGGATGTCAGTAC
TCAGCTGCCAGTGCAGTAAATTCACCTATAATGGTCTTTGGCAGGCTACCTGACCCA
GAAAAGACAAAGGTTATAGCAAGTGGGATACCCTCAAGCAAAGGCTTAAAGA
60 TTATATTTGCAAAAGACTTTATCCCAAGATGACGATGACGATGACGATGACGATG
TTTGACCGCATCTGATGATTCACAGATGATAATACACATGCGATGATGATTTGCTTC
ATCTGATGATTCAAGATGATGACGAGATGGCGATGACGATTCAGATGATTTGGGTGA
TGGTGCAGATGATGACCGCCAGGCAAGTGTATCATGAGGTAATTTGCCCTGATTT
70 TGAAAACAAATACTTGGCCATTATGAGCTACTCATGAAAAACCTTTGCCCTGATGG
TAAAATAAGGCTAAGTTGATGTAACCTTTGACACCAACAGCCCTACTGTAATTA
CGATGAGAGAGGTGATATGCTCTTTGATATCAAAAATGGCAAAATGATGGCAGGAT
TACCGCAAAGCCGATGTCGCAACTATCGTGAAGAGTGGCTAACAAACAGGTTGGCGG
65 TTTCTTATACAACATCAAAGATATTGATGTTAAGGGGCAATTTTTGGCACAATGGCGA
AGAGTTGGCAGGACGGTTACATCATGACAAAGCGATGGCATCACTGACACCGCGAAA
AGCAGGGGCTGTCTTTGGGCTGTTAAAGATTAATAAAGCCCCCTCATCATCGTTTGT
CGCTTGACCGACAGTTGATGACGCCCTTGGCAATGCTTAAACAGCAGCTTTGAAACAGT
70 GCCTTGGGCGAATCTTGGATAAATGCACAGATTTGCTCGGGCTAATATCTTGTATAAA
ACATCGCCATAAAATAGAAAATAAAGTTTAGGATTTTTT

SEQ. ID NO:45

Nucleotide sequence of DNA region (1000 bp) up-stream from the lbpB gene from *Moraxella catarrhalis*

5 CAGCTGTACCATTTGGTGAATATATACCAATTTGGTGGTTTGGATATTTTACAGGG
 CTGAGGGTGTGGTAGCCTAAGCCGTGGCGATGATRAGCAACCCSCTCAAATTTGGC
 GCGCGGTGGCGATACGATTTGGTGGGCAATTTCTATGAGGTGGCATCTCTGAGACG
 ACSCGTAAAATGCACTTGGCAGTAAATTTTATTAACCGTCTCAAACGATGCTTGGTTT
 GGTACAAACGAGTCCCTTTGCAGCATTACAATGGTCAAAATGGAGCTTGGAGACG
 10 GGGCGATGGTTTGGCGTGCACAAACCAACGGAGTACGCAATTAATTGACCATCAAGGA
 CGGATATCAAGCAGATACCGCAGTTTACGGAGATATTTGCGAGGTGATGTACCCAGT
 TATGTTGGACACAGCCCTTATATGTTTGGGGGATATCCCATGTTGGGGTTTCTTTG
 GTGCTGATTTTCTTAGTATCATGGCAAGAAAATGAAAATACCAACCGCAACGAGAA
 AAAATTTATACCGTGTATGGTGTGGTAGACCGCTGAATTTGCCACTTTGGGGGTAGAG
 CATGAGCAAGATTAGCCGTTGGGTGAGCTTTGGTTGATTTACTCATCAGCCTACCCGAAA
 15 CCTGCCAAACATCACCGCCAAACCTAAACATACAATGGTAAAAATACAGAAAATAA
 CTGTGTATTGTAATCTTATGTTTATCATGTGATAAATAATCATTAGTACCAAGAT
 ATCCATTACTAACTTCAATCCCCCATCTTAACAGTTACCAAGCGGTGAGCGGATTAATCCG
 ATTGACAGCAAGCTTAGCATGATGGCATCGGCTGATGCTTTTTGCCCTTGTGTGTGT
 20 TGTGGGAGTTGATGTACTTACCTTAGTGGTGGATGCTTGGGCTGATTAATAAATTTG
 ATCAAAGCGGCTTTCACAAACACCAAAACGAGATATCACC

SEQ. ID NO:46

Nucleotide sequence of DNA region (1000 bp) up-stream from the tbpB gene from *Moraxella catarrhalis*

25 AGTTTGCCTTGATTTGGAGGCCACTGCCATCATGAATTTGTGGCGTAAACCCACTCG
 TATTTCTTCGGTTTCCCTTTCCATGCAAAACACAGGATACACGCGCGCCATFGGCA
 GCGCGCGGTGCTTGGGTGCTAAAAATTTGCATGATGTCAGCGAATCTTGCACCC
 AAGCAACCAAGTCTAATCAGCACCGCTGTTGAATGGTCATGTGGATACAGCTTAGG
 30 ATTTAGCACCCCTAAGTGGTGTGGTCTTGATAGCGTTTCTTAACCCCATCAGCGCT
 GGCATCTCAGCTTTCGCAAGGCAATCTCAGCGGACCATTAATCGGCTAAACGGATATCA
 GGCATTTATAATCGTGAAGTTTGGTGGTACTGGATGATGAGGTAGGCAATCT
 TTATTCCTAAGCTATTTAAAGTATTTTAAACATAATTTGATGAATTTGAGATAATTG
 ATGTAAGAGGTTGAATGACCAACCATCGTAAACATAAGAAAAGACATTTTAAAGCAT
 35 AAAAAGCAAAATGTCTTGGTGGCTTATATAACAGTATATGATAAATTTGGGTAAGAA
 AGTAAATGGATCGTTGGTAACTTGTGGCTATCTTAATTAATATAATTTTAAAT
 AATGCTTTTACTTTAATTTAAAAATAGAGTAAAAAATGGTGGCTTTGGGTTTATCTC
 ACTATGTAGATAAAATGATACAAAATGGTTGTATTACTTGTATTGATATATAA
 TTTTACTTATTTTACAACTATACACTAAAAACAAAATTAATCACTTTGGTTGGTGG
 40 TTTTAGCAAGCAAAATGGTATTTTGGTAAACAAATTAAGTTCTTAAAAACGATACACGCT
 ATAAAACAGATGGTTTTGGCATCTGCAATTTGATGCTGCTTGTGATGTTGGGGTGT
 ATCGGTGATCAAAGTCAAAAAGCCAAACAGGTGGTCAATG

SEQ. ID NO:47

Nucleotide sequence of DNA region (1000 bp) up-stream from the tbpA gene from *Moraxella catarrhalis*

45 TTGGGGCGGATAAAAAGTGGTCTTTGCCAAAGGGGCATATGTGGGAGCGAAACCCCAA
 ATCTATGGCAAAACATCATCAAAAATCAAAAATAACAACGACCATTTGGGCGAGCTGGGG
 GCAAAATTTGGGCTTTGCTGATGCCAAAAAGACCTTAGCATGAGACCTATGGTGAATAA
 50 AGATTTTATGGGATGAGCGTTTATCCGACACCATCGGCATGCGATGTCGGTTGATFAT
 AGAATCAACCCAAAATTTCAAAGCCTAAACCGCATAGACATATCAGCCTAACCAACCAT
 CGGACGCCAGGGCTGACAGTAAACACTTTATACAGCACATCATGATTTATATCCCA
 AATGCCACAGCTATATCTTTTGGGGCAGACTTTTATGATGAAAAAGTCCCAACAGC
 CCATCTGACAGCTATGAGCTCGTGGCATACGACAGCGTGGGGCAAGATGGGCGGTT
 55 GGTCTTCAAGCGTGGCCAAATCAGCATCAACAAACGCAATTAACCAAGGGCAACCTA
 ACCAGTGGCGGCAAAATCGCCATGATAAACAGATCAAGCGCTTTTATCGCTTTGGCAC
 AGAGACATTCAAAATGGGCGATCAGCCACGGCTGACCATCAGTACAAAATCAATAA
 AGCAATGACATCAAGTCAAAATATCAAAAATCAAAATGTTGTTGAGTTTAGTCGATT
 60 TTTTGAAGGATAAGCAGCCCTACTTTGTTTTGTAAAAAAATGTGCCATCATAGACA
 ATATCAAGAAAAAATCAAGAAAAAAGATTAACAATTAATGATAATTTGTTATGTTTAT
 GTTATTTATCAATGTAATTTGCCGTATTTGTCCATCAAAAACGCAATTTATCATCA
 ATGCCAGACAAAATCGCCAAATGACATGTCACATGCCAAAATAGGCATTAACAGAC
 TTTTTAGATAATACCATCAACCCATCAGAGGATTTTTT

SEQ. ID NO:48

Nucleotide sequence of DNA region (1000 bp) up-stream from the ompE gene from *Moraxella catarrhalis*

65 AAAGACATTACACATCATCAATCAAAACGCCCAACCATGTACCTCTGCCCCGGTGGCAC
 GCCAACGCTTTTGTATGCGGTGGTGGTTGAGATGGCTTGTCAATCATTGGTTTTAT
 TAAAATTCATACCTTTGGTAGTTGGCTTACCTGATATGCTATTGATTAACGAAACAA
 70 TACGCAAGTTGACCAAAACATCAATTTTAGCCATTTGCCAAGCACTCAATATTAACGCTCA
 TACGACCATGCTTGGTATTAATCATCACATAAAGATACCTTACATCCATTTGAATTGAC

ATTACCCAAATACGGCCATGCCCAAATATGATGATGAATGGTGCAAAAACATCCATT
GGCTATTTTCATCAACTGCTGCCGCTGCGGATATTTTATACCCAAACGGTTTGTAT
TGTTGGCGGTGAAAGCTCAGGGAAAACACTACCTTGGTGCAAAAACCTTGCCAAATATTTATGG
TGCCAGCATCGCACCTGAAATGGGTGATATACACACATCCCATCTCGGGGTAGCGA
5 ACTTGGCCCTTCAATACAGCGACTACGCAATCCATTCGCATCAATACGCCAACGGCTATCGA
AACCSCCTGTAACACTGCGAGCTCTGCTGTATCACTGATGATCTGATTTTGGCACACAC
GCAAGCATTTTGGAAATTTAGAGGGGGAAACCATCCGCTTGTCCAGATTTTCTAA
ACAAATCGGATTCGATTTTACGATTTATTTAGATAAATATCTTGGCTGGCTCGCTATGG
CATGGTAGGCTTGGTATGATCATCAACGAGTTTGTCCCAATAAATTTGCTTSGAT
10 TTTGGCAGSATATGATATAGTTATCATATCATTAATGACACCGACTACCAAAACCGTA
TCTACAAGCATTAAGCTTATAGACATCATATTTTATCATTTTACAAAATTTATGA
CAATTAATTAGGAAAATCGATGAAAATGATATTTTAG

SEQ. ID NO:49

15 **Nucleotide sequence of DNA region (1000 bp) up-stream from the *uspa1* gene from *Moraxella catarrhalis***

GGATGTGGCATATCTGCCATCGACCCAAATACACATCGGTGAGGCTATCAAGATGTGT
ACGAATTAATACGAGCTCAGGTAAAGGCGGGTCTGCGTATATCTTGACGGCCATTTTGG
20 TTTTAATTAACACGCTGGACACAGATTGATTTTGGCTGGTACAGGCTTATGSCAGA
AAGTATGGCGGCTGAACTAAAACATGATGAGCTGCTTGAATTTTACCACAGCGTATCT
TAAGCAAGATAAATCCGCTAAGTGAATATACCATCAGCAATAAAGGCGATGCTGTCCAG
CTTCAAAGGCAAGTAGCGACACCCAAAGCGGTGTTGAGGTGATTTGGTCAAGGCAATGG
TGCGTTATCTGCGTTCAATGATGGCTTGGTGAATCCACGGGACAGAGATTCAATGTAC
25 CAATTAACCCGAACACGCGCATGATAACAAAACCCATCAAAAACCGGATCGGATAACCA
AACCAGTCCCGCGTGGCTTATATCCAGCTGTGCTGAGGGGCGATTATTAACAGGC
ATCCGACCTTCCGATGACCGGTATCCGCGATGCTAAAAGCTCATTTACCGCTTGGCA
CAGGCGTGGTATCTGACCAATCAAAATCCGATGATGGCAGGATTTTATTTTATGAT
GGCTGCCAAACATGATGATCATCAGCATGTGAGCAATGACTGGCTAANTGACTGAT
GAGTGTCTATTTAATGAAAGATATCAATATATAAAGTTGACTATAGCGATGCAATACAG
30 TAAAATTTGTACCGCTAARCATAAACGCGGTCCAAGATGGCGATATCGCCATTTACCA
ACCTGATAATCAGTTGATAGCCATTAAGCATGGCATCAAGTTGTTGTTGATATGTCAT
TATAAACGGTAAAATTTGGTTGGTGGATGCCCATCTGATTTACCGTCCCCCTAATAAGT
GAGGGGGGGAGACCCAGTCAATTTATAGGAGACTAAG

35 SEQ. ID NO:50

Nucleotide sequence of DNA region (1000 bp) up-stream from the *uspa2* gene from *Moraxella catarrhalis*

CCCCAAGCTTCCGTTTGTGTGCTGCTGSGTGTGSGGCGTCAATACCATGCTGGTGGTGT
CCGATGGTGGCGGTGTATCAGATATTTGGGCGAGTCTATCGATGATGGGTGGGTGAAT
40 GCTTTGATAAAAACGGCAAAAATGCTCAAACTGCCCTATCCTGGTGGCCAAAATTCGAAA
AATTAGCCAAAACGGCAAAACCCACAGCGCTATGAGCTGCCAAGACCCATGACAGATAAAG
GGCTGATTTTTCGTTACGTGGCATGAAACCGCCATTCATATCTCATCAAGACACAC
CAAACCGCCAAAGCCACCCGCTACAGCGAGCATCCCGCTGAGCTTTGAGTATGGG
TGSTGATACTTTGSTCAAAAATGACCAAGCACTACAGATGACAGGCTATGSCACAG
45 TGCTGTCCAGGSSGCTCTCTGCCAATCAGATGCTACGCGCCACCTGACCGAGAGCC
TCCGCCAAATCGATGCGTCCGTTACTATGCCCCGCGAGCTATGCAAGGATATGGTG
CGATGATCGCTATGCTGGCTTTTGTCCGCTCAGCTGTGACAGTCCGATGACTTGGCGG
TTCCGCTATTTCCCGATGGGATATGACGAGCTTGGCGTATCGGCTCATAGATAGCCAC
50 ATCAATCATACCAACAAATCGTACAAAACGGTGTATACATGCCAAAATACCATTTGAA
AGTAGGGTTTGGGTATTTATGTAACCTTATATCTAATTTGGTGTGATACCTTTGATAA
AGCCCTGCTATAGTAACTAAATGGATATGATAGAGATTTTCCATTTATGCCAGCAA
AAGAGATAGATAGATAGATAGATAGATAGAACTCTGCTTTTATCTGTCGCTGATGCTT
TCTGCTGCCACCGATGATCATTTATCTGCTTTTGGGATCAGTTATTTACCCGTGA
TGACTGATGATGACTTAACCAAAAAGAGAGTCTAA

55 SEQ. ID NO:51

Nucleotide sequence of DNA region (1000 bp) up-stream from the *omp21* gene from *Moraxella catarrhalis*

GAGTGAACCTTATGTAATATGATTCATTAAGATACAAATCATCAACCGCAGCATC
AGGTTTGTAAAACAAATTTTACCATAATATAGCCATAACGACAGCAAGCTGAT
60 TATGCGAGCGGCTGCCCCTGATAAATATCATTTTGGAAATCAACCAACCTAATGSCATC
AGTCGGTGGATGCCAGTGTATGACACAGCTATAAATAAGCGTGGTGGGCTTTT
GACGCTGAGCGTATCACGCCAATCACTGGTCAACAGTGTGCGCATCCAAAATGTA
TAAAATTTAGGCAAAATAACGCTCAGGCTTATGTTACAAATTTGCCAAATAAAAACCCCG
65 TGCTTTAATCGTTCAAGCCCTTGTATAAACCCTGCATAGCTTGGGATTTTCAATPTGT
TTTATGGGCATATCTGCCAAAATAACTCATGGGCATGTTGAATCATAGTCGATCATA
GATATGATGCTGCTGCTCAACCAATTTTAGCGAACCATTTGCCACCCAGCT
TTTATGATGATCAATTTGGCATAGGCGTAAAGTTAAGCTTGGCATACATGCCATTTAGCCG
CGCCGCAAAATCAGGGGCACTATCGATAAGCGTACCATCCAATCAATATAATCAGTTT
70 TTTGCCAGTCAATGACAGTGTGGATGCTTTTCTTATTTAAAATTTGGCGGTGTT
TGGTATTTTAAAATCAGTCAATTTTACCATTGCTCATATAATGACAAAATGACAAAAT
AGCAATATTTTAGTGCAATTTTGGCGAAGTTTATGAAAACCTGCTCATTTGGTTGCAAAA
CTTTACACAGTACCTATAAACTTGACAGTTAATAAGAAATATTTGTTACTATAGGGG

CGTCAATGGGAAACAGACAGTTATTTGTAATAGTTATTTGCAAAAGACGGCTAAAAGAC
AGAACAGCGTTTGGTTTCAGTGATTAAC TAGGAGAAAAACA

SEQ. ID NO:52

5 **Nucleotide sequence of DNA region (1000 bp) up-stream from the omp106 gene from *Moraxella catarrhalis***

TTGATCGGTTTTGCCCACTGTTTCATGATTTACTCAAAACAGCGGCTTGATCGTGTG
GCAGGCTGACCCAAAACCAAACCCAAAGCGGTATCGATGCTACTGCGCTTATGTTACG
CTTGATACGCCAATTTGTTATGAGATGCCAAGACTGCCAATGGCAAGCGCTAAGCGCG
20 ATCAAACTCAACCCATTAAGCGATAGCCATGAGCCCAAACTTAAGCCAAACCGCGT
ATATCAACAAGTTGAGCAGACCGCAAGCGTTATTTGAGACATTGGGCGATGCTCACT
TCATGATGCTATGCCACTTTTGGCCGAATTTGAAAAACCGTGTCTATCGCGCACT
CAATCACACGCACGGCAATCAGTCAAAAACCGCCAAATCCTGGTATCAATCGTGGCAC
15 ATACGCACCAAATGAAAACCCATCACTTACTTTAGACCGCCAGTTATCGCCATGGATA
TGGCAGGTGTGCTCGCTGCGGTATGATGGCGATGACCCCAATTTGCCCATATCTGC
ACGATTTGACATGATTAACATGTGATGATTTAAACATGACATGATTTAAACATGTT
TAATACTGTTGCCATCATTACCAATAATTTAGTAAAGCATTGTAATAATCAATGCCCTT
TTTTATGTTGATCAITGAATAGAAATATATGATGTTATGATTTATGATCAAGAAAT
GGTATGCTACGAGTTGATTTGGGTTAATCACTCTATTTGATATGTTTGAACAATA
20 ATCTATTGACTTAAATCACCATATGGTTAATAATTTAGCAATAATGGTAGGCTTTTGTAAA
AATCACATCGCAATATGTTCTACTGTTACCAACATGCTTGAATGACGATCCAAATCACC
AGATTCATTCAGTGTGTTGTTATACGCAACCTTACCCATAATTTCAATCAAT
GCCTATGTCAGCATGATCAITTTTTTAAAGTAAACACC

SEQ. ID NO:53

25 **Nucleotide sequence of DNA region (1000 bp) up-stream from the HtrB gene from *Moraxella catarrhalis***

ACTATTCGCTTTTTGTTTTTCAAGAAATGCGAATGCCCAACTCAGCAACTGCGGATTAT
CAACTTCAGCAGGTGCTTCGGTCAATGGGCAATCTGCCGCTTGGTTTTGGGAAAGCGCA
30 TCACATCAGGATGAGCTGGCAACCAACCTCAGCATAATCAGCGATCTAGACCAAAATG
CCAAACCAACCGTGGCGGTCACCAAAAGCAATGATCCATCAAAAACCTTAAACTTAA
GCTCTGCTTCTTTTAGAAATACCAAGGCATCAAAATACCGCTCTGATGTCACCG
TATTAATACGCGAAGCCGCAACCAATTTCTGTCGCAATTTAGTACCATGTCATAGCGAA
TGGATAGGGCGTTTCGGACTTTGTTGAGTTTCCTCAACGAGCCTTTGGCGGTFGA
35 AAGGATGATGAATGATGCTCACTTACCAATCATCAGTTTCCTCAAACTTGGAAATCAA
CGCCCAAGCGGTGCCCAATCAGAGTAAATAAATTTAAATCAGTACCGATTTTAAAC
GCAATGCAACCATAGCATATGACGATTTTGGCTTTTTCGGCAACAAAGAAATGATAT
CGCCAGTTTGGGCAATCGGTACGCTCAATCAGTCAATCAAAAACCTCATCGGTCAATTT
40 TAATGATGGTGATGTTATCTGATTTCTTTTCAAGCGCAATTTGATATGCTTGGCTG
CATTCACCTTAATATGCCAATCCACGAGCCCTAAATGCAACAAATTTGGTACT
CATCAATCTGTTGGACTCATGTTACCGCATTGGAATGCTAAGCAACACACCGC
CTTTAGGATCTTGGCGGGCCCTGAAATACTTAAATCAACATGTTGATGATGTCAG
CAACATCAATAGTTTAAAGGAATGCTAAATCAGGCTTATCTGAGGCATAATCAGCA
45 TGGCATCTGCTAAGTATGTCGGGGAAGTATCAAACTCA

SEQ. ID NO:54

50 **Nucleotide sequence of DNA region (1000 bp) up-stream from the MsbB gene from *Moraxella catarrhalis***

TGGATCATATCTTTAATTAATGGTACTGTTTAAACCTGATTTTAAAGTTTATGGGTCA
TATTTTCAAGCTCATCCATCGCTCAAGCTTTCATCAAAAAGCTCATCAATCTCTACCA
ATCGCTCACAGCCTTCGTTGCTGCGCCAAATCGGTATTAACCATGAACCATCTTCAA
55 TCTTTTGGCAAGCTGTGCTGCTTTGTTCAAGTGCAGCAATTTCAATAGGCAAACTT
CAAGTTCAAGCTGCTCTTTATAGCTGAGTTGCGTTTTTGGCAACGCTGATGAGGTG
GTTTGAATTTGGATGGGTTAGCGGTTTTGTCGCTTAGGTTTATGCTGTGGCGTAT
GAGCAAGCCATCTTTCATGCTGTTGATCATAGTCTTCAAAACCGCAACATTTCCAAAA
CGATACCGCTGCGCTACTTATCAGTATCAAAATCCCAAGTTTGGGTAACACATTTATCCA
60 TAAAGCAGGTCATGGCTGATGAGTAAATCCGTCCTTTAAATGCAACCAAAATCTT
CTAAAAGCTCAAGTGTGCAATATCCAATCATGGTAGGCTCACTCAAGCACCAAAACAT
TGGCAGGTTTTAGCAATAATTTGGCAATAAAACCGCGTCTTTTCAAGCCTGATAGT
CTTTAACAGGTGTGCGAGCAGATTTGGCGTGAATAAAAATCTTGCAATAGCTTAAAA
TGTGCGTAGTTTTTCCACCAACATCGACATGGTCAGAGCCTTCTGAAACATTTCTGCGA
TAGATTTTTCAGGCTAGGTCGCTTTGAGTTGTTCAAAAAGCAATTTAGATTTGG
65 TGGCAAGCTTAAGTAACTGACTGACTGAATCGCTGAATCATCCAAACCAAAATGCTTTAA
TTAAGGTTGTTTTTACCAAGCCTTTTTTGGCAATGATACCAACTTTATCACCAGAACAA
GCAGGCTGAAAAATCTTAACTAAGGTTTTATGTCGTAT

SEQ. ID NO:55

70 **Nucleotide sequence of DNA region (1000 bp) up-stream from the PilQ gene from *Moraxella catarrhalis***

CAACTTGAAAATCAGCTCAATGCTCTGCCACGCACGACCGATGAGCGAGATTATCGGA

ATGATAAATACCAAAGCACAAGCGGTTAATGTGCAGGTGGTGGAGTGCATCAGTTCAAGCA
GGTCGTGAACAGGATTTATATACCGAACCGCCCTATCGCAGTGGAGTGCACAGGGGATTT
CATGCTTTGGGTCGATGGTTACTTGAAGTTGTACAGAGGTAACCAATTTGCTGACAGTGCAT
GATTTGATCTGAAGGCTGGTTTGAACCATCAGCTGATGATGATTTGTCAGATGAAAACT
5 TATCAAGCGAAACAAACGCCAAAACCAAGTTGCTCAGCAGGTGCTGATGTTCAATGAATA
TTATCGGTGGGGCATTTGGGTGCTTGGATTTGGGTGGGATTTGGATGCTGATAGCAG
CAGTCAAGTTGTTGATGATAAGCTTGCACATATTACCCATGAAGAGCGTATGGCGATCAG
TGAGCCTGTCCGATACCCCTTATCTGTGCCGATGATATACAGCAAGGCAAAAGATCCTTT
10 TATCAATCCTTATAGAAAATGTTGAGGTTCTTGATACCAATCATGCCGTGATCAGCAAGA
TGACCCAAAAACCAATCTACCAAAAGCTTGGCTATGGCAGACATATGCCATCTCAGCC
ATCTGATCTCATCAGTCTGCCAAGGCTCAGGCACAAGTCTTCAAAAGCGATCCGATAGT
CATTSATACCAACCGTGTTCGAGAGCCTTGAAGAGCTATGACTTATCAAGCTACGCTA
TCATGCTGATTTTTGATGATGTTAGACTGTGGCACTCATATGAGTCTGATGGCAT
15 CGTTATCGTGTGAGTACTGGCAATATCTTGTAAAATCAAGGAAAAATACCCATAT
TGACAGTCTGATACATCTGATTAAGCGGTGCTGATACACAAGGTGGCTATTATCG
CCGTGATGTAACATTCATTTTATCATAGCAATGACAC

SEQ. ID NO:56

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipol8 gene
from *Moraxella catarrhalis*

20 TTCATGCAACAGCGACCATCTTGGCCGATGATACCATCCTGCTCACTAAGAAAAATCAG
TTTATCAGCTTGCAGGCAATGGCTGTGGTCACTCATCTTCTGCCAATAGATTA
AAATTCGCCGTAACCGAAAAACCTGTGGTCTAGTAGGACAATATGGTCATTTACCAA
25 ATTTAGGCGAATGGCATCGACATCAATTTGAGCGTACCTCAGCTGATCTGATATCCAT
ACCATCTCGATGCGTAAGGGCGAGCGGTGACAAAAATACCGAAATGGCATCAATACG
AGATCCGTACATTTGGGAGTTAGCAAGCCCAACGACAGCGAGCTTCGATTTGTAGAGC
AATGAGCCGACTGCTTCAAGATGGCAGGCAATAGATTCATACGGTTTACACGCCAATT
CTCATGTAGTTGATATCAGCTTGGGATTTTGTAAATTTTTTCCACTTGTGGCGTAC
30 ACCATGCAACAGCAAAATTTGATGCCAAGCTGTGTAGCAGTGCAAAATCATGAATCAG
CGTACTAAAAATGTCAAGAGCGACCGCTCATACCAAAACATAACCAAAAGGTTTGGCC
ACGATGGGTGTTAATGTACGGGGCAGAAATACGAAAAATGACAGGTGTGAGTTCAGG
AGTGTCTGATAGGTGCTGACAGAAATTCATGAATGCTCCAAAGAGTCAATGGCTGTAA
ATAAGAAATGGCGAACAATATAGCGAGAGCGTCTGATGTTGGTCAAATGTCCTAATA
35 AACTATCAAGTATACCATCATACCATAGCAAAAGTTTGGGAGATGCCAAGCGAATTTATC
AGCTGTGATAGGTGGCATATGATAAAAATCAACATCATCGTCCCGAGTTTGGAGCATG
GTAAGTGTACCATAAATTAACAGTCAAGAAATTCACACCGTCAATCAGCTGTGCTATG
CTTATGGGCACATAAACTTGACCAACACAGGATAAATTTA

SEQ. ID NO:57

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipol1 gene
from *Moraxella catarrhalis*

40 GGCATACITTTGCGAATGCTTTATTTGGCATAACTGCTATAGCCCATGCTACTTTTTA
TCATTTATCCATATGTCATAAATGCTTTATGTAATTTAGGCACACTTAACTCGTG
CCACTGTAAACATTCAGCATAAAAATCTTAACAATGAATCAAGCATCGTATTTGGTGT
45 AAATGATAAGCTTATATTTATTTAAATTCAGCATAAATGATTTAATATGGACATATCA
GGTGAATCAAAAATTTGGGAGGTTATGATGATATGAATAAAAATGGCCACATC
GTAGGGTGTGATACGGTGTCTTATATGGCATAATGTTGGTCACTGGCTATATTTA
TGGGGCTGGATTTGGTGTGACAGGATTTACTCGTGGCAGTACTTTGGCTGATCGCTTGGAT
50 ATTGTGGGTACGATTTGCTGATAGAAATCTGATACCGATATTTTGGACCGTGGTGGT
TTATTTTCTATCTTTTGA AAAAAGGGGATAATTTGGTATTTTTTCAAAAAAATCAT
GATTTTTTTGTAACATCTAAAAATCAATATGTTATATATGATGATAAAAAGTGGG
CATGCTTAAGTTTGGATGCAAAAAATCAATATCATCACTGACCAAGCTGTGATGAT
ATCAAACTTTATCAAGTCTTAGGGTATATCAAGATATCATACCAATGAATACTTA
55 CCCAATCTACTATAAAAATCAATGATATGACTGTGATTTATATCATAGATCAAAAA
TCAAAAACGATGAGCAGGATGATGATGAAATCAAAAATTTGGCAGCATTAATGACA
ATCTAAATGTCGCGAGAACCGTGAACATTTGGGTGATTTGGTGGGATAGGGTCAAGCCA
GTGCGATTAAGCTAAATTTTTATGTTGGCAATCGCTGACTTTATTTTTATTTGGCCAGTT
60 GGAACAATTCGTGCTAATGATTTATTTTAAAGAGATAA

SEQ. ID NO:58

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipol0 gene
from *Moraxella catarrhalis*

65 TCTGGTCTACATCCAACTATTTACACAAGAAAACATAAGCAGTGGAGCAGATGACG
CTCAAAAAGGCATCTTATAGTAAATTTGACAGTTAATTTTCGTCAAGTGCCTGTACAAAA
TACACCATCGTGCAGAAAGTTTGTACCAATTTAAGCACAATCATTTTGGCACACACTGTC
AAGCAATGCTTCAGGCAAAATAGCTGTGGTAAAGATCTTTGGGTCAATGCAATGSCA
70 TCAACCTTTCTGCTCGTTGAAAGGTAAGTTTCCACTTCCAAAATTTGACCAATGSI
TTAGACCTTGGCGTGGCTTGTGATATGCTCCCAATGTCGATGAGTATGATTTGAGTCA
TATCAAAAGGCGATGAAATGATGGAATGTCATATAGCCATTTGTAGACTTGTG
75 TCATCAGGCTTGGCCATGACTATTAACCTGTGTCACCAATAATATCAGACACAA
ACGCAAGTGTGCTGATGAAATGGCAATGCTGCTCCCAATATCAGCATACGCCACC
AATTTGGCTTGGCGTGGAGCTCAGCTTGGGTGTCAAGGGTCTCGTGGTGTGACGSGGC
CGATTTGACATGACGCTGACGCAATAATCTGCTTTGAGTGGTTCGGAAATTTATGTA
AATAATCCGCTTTGAGTCTTGTGCGTTTTTAGGCTTTTCAAGGGGTGATAATCAGSCA
TGGTTCTTGGTAAATCAAGCCGCTTCCATGGGTGAAAATGAGGCAATCATGAAAAAA

TGACCTGTTTCATTGGTGTATGATTACCGTTTTTGTGGTGGTGGCCACATATTGCACCG
CAATGACTTCTCGAGCTGATAAACTGCGTCCATCAGTAAGCGGCGTACTGTGATGATGA
CTGGTAGACGAATATCGCCACCTCGTAAAAAATAACCTGTAGGCTATGACACAGGTTTAT
CAATCGTTAATGTGTTAGCACCCAGCAGCGCTTGGGCA

5

SEQ. ID NO:59

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipo2 gene from *Moraxella catarrhalis*

10 TAAAAAATGACCTTACAAAATAAAAATATATGTTCAAAAATCGCTTAAAGTATTGAAAAAAGC
TATAAAAACTTATCTATTAAGCATAAAAGATATTAAGCATAAAAGACGAGAAAAGAGC
AAGCGTCAATGATGATATTTTCATATAAAAACTTATGAAATTTTCAATTTTTATCGATT
GATTCAGCTTGGCTATCGGTGGTCAACTTGGCTGCCAAGCATCGCCGGCTTTTGA
AATCATCACAATGGCAACAATGATGATGGTGAATCCACTTGACATATACCATGTTGGG
ATGCTCACCATAGTTAATCGCAAGGCTTCCCAAGCCACACCGCAACACACCTGCCAT
15 TGCAGAATAACCAATCAAAGACACCAAGGTCATGACCGCATTAATCAAATGGGCAG
GCTTTCAGCAAAATAGTATTTGCTGCAACCTGCCAATGGCTGACCATAGATTGGC
AGCTTCGGTCAGTCTGTGGTACTTCTAATAAAGCATGGCACTCAAGCGTGCAAAAA
TGGAAATGCTGCCACACTCAAAGGACGATGGCGCTGTGTGCCAAGGTTGTCCAC
CAAATCGTGGTGCATGAGAAATAAGAGCAAAATAAAAAGGAACGGAGCGAC
20 AATATTAATAAATAACATCAAATAAATAAATACTGCGATTTCAGGATACGCCCCCTT
ATCGGTTAAAAATGCCAAAAACCTTATCGGTAGCCCAACAAAAAGGATGSCAATGGC
AGCAAGCCCATATAGATGGTTTCCCAAGTGGATTGGGCAACCATCTCCACATCTTGG
GTGCAATTCACAGCAAAATTTGTGACGATTTCAATCCACATAGCCGATTAATCTCATAT
TGACCCGATGGTGGTTAAAAATCTATTTGCTGATGACCGAGGTGCTTCAACCATAA
25 GCTCAGCAATGGTAAAGCCAAATTTTATATCACCTGCATAA

SEQ. ID NO:60

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipo7 gene from *Moraxella catarrhalis*

30 AGTAAACAATGGTAAACAATACAGCAGTGTCCACAGTCTCAGTACGATGATTCTGAAT
TTGAATATGCGAGATTTTGGATAGSATTGTGGCATGCTTGTGATAATTTAATTTGTA
TGATTAATAATGACCGTATTTGGTTTTATAATATACGCAATGATGGCCATGCTGCTG
ACCAATACCGTTTTATAGTGTGGGATGCAATCTTTATAGTGTGGGATGCTATCC
35 TAACTTATGATGGCGGGCGTGTGGTTTTGGTAAAAGGTTGCAACCCAG
GTAAATGCTCTTTGGGCTGCAAGTCCGTGATGCCAAGCAGGCAATTTATCAGTGTG
CAAGGGCATTTAGGATATTTAGTTATCTGATTTCACTCGTATTTTGTGGGAC
TTATTGGTGGTTTTGATAGAAAACAGGCTGGCATGATAAATGCAAACTG
TTGTGTAACCACTTCGCTGATGGGTGCCAGTAAACAAATAAACCACTCAAAGCA
40 CGAGGGCATGTTTGGCAGTGGCGGTAGATAAGCTAAACAAAGCAGGCTATSAAT
TATTTAAACCACTTACACCCCAATTTGTGTGAGATGATATATCGCCAGACAGCC
TTGGAGCAATCGCACCGTTTGGTGCAGCAAGATTTGTACGGTATTTGTGAAGTGG
TAGCGAACAAGTCTGTGATGTTACAGCGCTTGAGAGGTTTCACTCAAAGGACAGCC
AAAAATCTACCAAGCAGCAGAACGATTTTAACTCAATTAATCCAAATATATGATGATG
45 ATACCGTTTGTGATCATGTTTGTGATGGTGTGATGGATGATGAACATGAATGGAT
AAAAAATCGTTTGTGATGGTCAATGGTGTGAAATTAATCAATCAAGCAATCGGTAG
CTTACTATAAGATATATCCAGTAAATGGAAAATAGCA

SEQ. ID NO:61

Nucleotide sequence of DNA region (1000 bp) up-stream from the lipo6 gene from *Moraxella catarrhalis*

50 CGTTTAGCTTCATACGACACCTTGTGCACCTTCGGGCAACCGAAGCATCACGCCAGCAT
CACGCATCCGCACAAAACCCATCATGCCATCAATTCGCTGCTGATATGATATACCCCA
CCAAAGTAAACCGCTTAAATCGTGAATAACCGCTGCTGAGGGTGGGCTCAGGCA
55 AAACCAAGGTAACCTTATCCCAACTTAAAGTCCATGTCAGAGCAATGGACTCACCTA
ATATAATACCAACTCGCCGATATGTAATATCCAAATTCCTGCGGTATATGCTCAT
CAATGATAGAAAATGCTTTTCTGATCAGGCTCAATGCCAGAAACCGATTCAGTCA
CCTGACCTTCAGCGTTAACATACCTTGTAGTTGAATATAAGGGCAACTGCTTGCACCT
CTGGATTTGCAATTTGATTTTTTGGCAAGTTCTTGCCAAATTTGTCAAAATTTCTGTG
60 AGGTAAGTGAAGCTTGGGCAACATGCCAAGAAATGCGTGAATTAATTTCAAGGTCAAAAG
CATTCATGACCGCAAAACCGTATAGCACTGCAACCCCAAGCGTAAAGCCAAATGGTGTG
AGATCAAGAAATAAGGAATAAGCAATTTTACCTTAGCTTTGGTATATCTAAGCC
CAATAAATAACCGCAAGGACCAAACTAAGCTGTGTCCAAACGACCCACGCTGCTAG
TTTAGCACTTTTGGCAAAATCCAAACATCACTAAACAAATGAATCATCAGTGTGTT
65 TGTGCGCTTGTGATCTGATGATAGTTCTTGTCAAAACGCTTTTTATCTCAGA
ATACGAAAGGATATATCTATATTTTAACTTTAAATAGATCTGCTTTTATATCCGA
TGATTTGGCATGAAGTTATCGGTGATATGCTGATATAAGTTTATCGGCTGATATA
AATTTAATTAATCACTCAAAATTTTAAGGAATTTATCATTA

SEQ. ID NO:62

70 **Nucleotide sequence of DNA region (1000 bp) up-stream from the P6 gene from *Moraxella catarrhalis***

TAAGGATACCAAGATTTGGCTGTCAATCGTGTGTTAATCATTGTAACGGTTTATAGTG
ATTTGCAATTAATAAGGGTAAAAAGTATTTATCAAGTAATAATCTTCTTATATGTGAA
TATAATGACAAATTTATCACATTTTACAAGGATTTTTATCAAGATTAGGATATGTTC

AGCTTAATTATTAGTGATGAGCGGTGATTATTTGGCATCGTTAAATTTATGAGTCTAA
AATTGGCAAATGATTAATAATTTTGGCTAACATGATAGCCCTTTGGTAGGCTTTATTGGT
ATTGATGAGCAATAATAATATACCGAGTTAAATGGATTAACTTAAACATACGCCAAAAT
5 TAACACGAAAAGTAGATGATTATGACAGATACAGTACAAAAGATACAGCCAGTCCC
CAAAAAGTTTATCTAAAAGACTACAGCCGCGAGTATGCGATTAAATAACTGGT
GGATATCCGCTTTTGGATGATCATCTGTCTGGTGGTCCAACTTAAATGACAGGAGC
ACAGCGAGGAGCTTCCGGCTTTGGCGAGATTAAAGCTTAAAGCMTTCCCTAAA
TGCTCAGGAATPAGACTCCAGCGGTATCATCTTGATAANGAAGCTTAACAATTTAGA
10 TGACAGATGTCAGATGATGAGACATGGTGGAGATTCCACCACAAACCAACACAC
ACTTGAAGGGCTATATCAAGCAGGAACAGGTGATGATAAGATCTTTGTGACAACTGGA
ACCTGAGGGCTTTCCGAAAATCACCTTTTCCCTGACCGCCCTGATTTTGCAGGAATA
CACACAGCGCTTGAAGCACCAGCAATTTAAACCTTGCCTGCAATGGTAATTTGGT
TGAGTCAGGAGATGGATGAAAATCGCCATTAACCAATTTGGCATGATCTTCCAAAA
ACCCAGCTATCTATTCGCCGCTGCATTTGCCAATCTAGAAG

15 SEQ. ID NO:63
Nucleotide sequence of DNA region (1000 bp) up-stream from the MsbB gene
from *Haemophilus influenzae* (HiRd)

20 AATCAAGCGCCTGTGCTGCTGGTATGGTGTGGAGACGAATATATTTGGTTGA
ACCGCAAACACAGGCCTTCACTGAGCAAACTAAAGTTACACCGCCTGAGCCGTTTTT
GTGCCAAGGATTTTGAACCTACCGAATCGGAGAGAATGGTAGAATAGCATTGAGGTAA
ATCAATATGAGATATCGGCATTGATCTTTAGCAATATGTTTTGGTGGTTTTGCGCA
TCATTTATCGATGCAATGCTGGCGGTGGTGAATTAATCACCATTCCAGCGTTACCATG
ACAGGTATGCCACAGCAATGGCGTTAGGCACCAACAAATGCAAGCTATGGCGGTGCA
25 TATCCGCAAGCCTTTATTTCTGGAAAAGAGCGGTCAATTAACGGATATTTGGTTT
ATTTGATTTGGGTTTTCTTAGGTTCTGCCCTAGTACATTAATTAATCAATCAATGAC
GTGGCGATTTCAAATAATGCTTCTTTTTGATTTTAGCCATTGGTCAATATTTTATA
TTTACTCTAAATTAGGTGATGAAGATCGAAAACAGGATTAAGTTATCTGTTATTTGGT
CTTTTAGTTAGCCATTTTAGGTTTTTATGATGGCTCTTTGGCCAGGGACTGGCTCA
30 ATCATGAGTTAGCCTGTGTTACTTTGCTAGGATTTAACTCCGAAAGCGGCAGCAGAT
GCAAAAGTGAATGAACTTCACTTCGAACCTTGCTCTTTTGCACCTTTCTTATTTGGCGGA
CAAAATCTTTGGAAAAGTGGTTTTGATGATGGCTGGGAGCATTTAGGTGCAAAATTA
GGTGCACAAATGGTATGACGAAAGTAAACCTTGAATCGACCGATGGTGTGTTATCATG
35 TCTTTATGATGACGGCTAAAATGGTTACGATCAGGGTGGTTTCATTTTAAATTCGGA
AAGCGCGAAAAGTGGGTTAAAATTAATACATTTATTA

SEQ. ID NO:64
Nucleotide sequence of DNA region (1000 bp) up-stream from the HtrB gene
from *Haemophilus influenzae* (HiRd)

40 TTGAAGTCCCAATTTACCACCAATCTCCTGGGCAACATTTGGTAGGTAACAGATT
CTTCGAAAGAGCTCCATCTGCTAATGGTGGTCTAATACACTAATGACNCTGTCCCGG
CTCCGTCACATCAAGACTCTTTTGCACAGTGGCAAAATGAAAGGCTTTGATTTG
GGCTAATTAATGTCATGCTTTTTCAGAACGCTCACCAAAAGTGGGTTAATCAATAT
45 CAGAAATTAATTTAAACCTTCTTAATATCTCTCTCTGATATACATTTACCTACAA
CGGCTTCAAAATCAGACATATGGGTGTCATATATGAGCCCAAGATAACGTTCAAAAT
CAGTCCCTTTGGATCGATCAACAGGCACATTCGCTTGGCGCAATTTGAATCATTT
TCTGAAATCTTTAAGCGTGCCTTTGCCGTAATCAGAAAGATCAAGCACCCTAATTT
TCACCGCACTTTCAACTTCGCTAATAAATCTTGCATCTACATATTGAATCTTCTT
CAAAATCAAGCGGAGCAGCTGTGATGACGAGATAAAATACGTAATTTAGTAATGGTTG
50 GATGGGTTCTAATGCAACAAAATTAACATCAATCTTTGTTTTCTAATAGTGGGAAA
GTGCAAGACTGTCTCATCTTGTCCAAATCAATCCCAATTAAGTGAACGGGTACATGAGTG
AAGCAATATTCAGCCACATTTGACGACCGCCCGCGCTTCTCATTTCTTGTACGC
GAACTACTGGCACTGGTCTTCTGTTGAAAATACGGTTGGTTCACCGAACCAATAACGAT
CAAGCATCACATCGCTAATACAGTACTTTGCTTGCCTTAAATCTGCTGAATTTGAG
55 CCAATTTAAAATCTCTATTTGAATAACCAAAATTTGGCGATTTTACCACAACTCAA
TTTACGATAAATACGCCCTTAACTTACGTGAAAAGAACAA

SEQ. ID NO:65
Nucleotide sequence of DNA region (1000 bp) up-stream from the protein D
gene from *Haemophilus influenzae* (HiRd)

60 AGCAATAATTATAGCTGGAATATCTTTAAAGATGAAAGAGATCGTATAAGCAAAAA
ATTTTATATGGAGAATATTAGCAATATTGGTTTCGCTAATATCGTAATAAATAGTTC
AAATAATGATGGAAAACAGACTTTTTCTGGGGCAATTTCTTTTACAGCTATTTT
TATTCATCTGTACAGAAATTTAATTTGAAAAAAGTAGCCAAAAGATAAATGCTGTTGT
65 AATAAGTGCATCGACAGCAACAAATTCAGGAGTATTTTATGTTAGCTTTTAAATAC
TAAACAAATATATTTATCAAGATGTTGGCAATGGAAATGATAGGTTAGTTTGGC
TGCTTTTATGGAGATCAAGAGGATGTTGATGCTTTTATATCTCAAACAGGAG
AATCACCTTTTAAACCTTTGCAATATTAATCTCTCTTCAACTCCGATAATAGGTTA
CTTAAATAGATGAAAGATAAATATCTATCAGGAAATTAGCGGATATTTGATATAT
70 TGCTTCTATTTGGCAATTAAGAAAGAAAACAGGAGTGTGATATAAAGTATGATGAT
GTGGTGGATAGTATAGTAAATATCTGTCAATTCCTTTAATGAGCGGTTAGCA
ATCTCATTTAAGTTATGTTGAATAGATATTTGGAAAAGATGGAAGATAAAGC
TGTTAAATATGCTGAACATATGAATATACCAATACTCAAAATAAATAATCTGCTG
GAAATGATTAATAATCTCAATCTAATCTGATGATAGGATAATGACTATCAATCAGAAA
75 GAATGATTAAGCCATTAATAATATTTCAATTTGATGACTAATAATCACTCAACCTATT

CAAGTGCTGAACATCTTTAGAACTATTAATAATTATTCACGGAAATAATTAATTCGTGTA
AAATAGGTAATGATCCTAACCAATATATAAGGAGAATAAGT

SEQ. ID NO:66

5 Nucleotide sequence of DNA region (1000 bp) up-stream from the Hin47 gene
from *Haemophilus influenzae* (HiRd)

10 TAAATCTCCAAATAAATTTCCAGTAACTGGTCTCTAAGCAAAAAAATAAAAAAAT
GTTCAATAGAGGGAGCAAAATTAATCTGTTTAAAGGAAATCGGAGCAGTACAAAAACG
GCTTACAGTAGCAAAATTTCTATAAATTTATGTTTCTAATACGGCAATTTTCTAGTCAAT
15 AAAAAAGTCAAAAAATGAGCTGGATTAACCGAATTTTATGAAAAGTCTTCTCTTCCA
CTCGAAAAGCCAAATGTCAGAGGCGTATGGACAAAATGTAAGTCTGTTGAAACAGTAC
TTTATAGTGAAGAAGTCAACAGTAATCTGTATGTTGCCGAAATGGTGCATCATATGC
GTATGATGCTCGTGGAGCTTTATTAATTTTATGGACGAAGTTCAAGCCAGAAATGG
CGCCAGATTTAGAACCAAAAGATATTTAAAAATCAAGATTTAAAGAAATATAAGATC
20 GTATCAATCGCGCAAAAAGAAACGGGCGAGAAAGATGCGCTAATTAATCATGACAGGTA
CACTTTATAATATGCCAATCGTTGGCTGCAATCGAATTTGCTTTTATGGGCGGTTCAA
TGGGTTCTGTAGTTGGTCAAAAATTTGTTAAAGCGGCTGAAAAGCGATGGAATGAAT
GTCCATTTGTGTTTCTCTGCGAGTGGTGGTCTGTATGAGGAAAGCATTATTTCTCT
25 TAATGCAATGGCAAAAATAGTGGCGTACTTGTCTCAATGCGTGAAGGGTGTGCCAT
TTATTTCAATTAACGATCCGACTTTAGGCGGCTATCAGCCAGTTTTCGATGTTAG
GGGATTTAAATATGCGGAGCCAAAAGCCTTAATGGTTTTCAGGGCCACGCGTTATG
AACAACCTGCGTGA AAAATGCGCAAGGTTCCACAGTGTAGTGTCTTACTTGAGA
AAGGGCAATTTGATGATCGTGAACGTTTCAAGATGCGT

SEQ. ID NO:67

25 Nucleotide sequence of DNA region (1000 bp) up-stream from the P5 gene from
Haemophilus influenzae (HiRd)

30 TCACCTAATCAAGCGCATCAATGTTTCTAAAAACCAACAGAATTGACCGCACTTGTA
TCTAAAAATTCGCCATTTATTAAGACTGCGCGTAATGCCAAAACATGATTAGAGGTTTA
CCATATTTGCAATGAGCCTTCCCGAGGCGATCGGTTTAAATCATTCCACCTAAAGTCGT
CGATGCTGGTGGACGTTCTGGGGCAAGAAACAAACATGTTGTTTAAAAATTTGATTA
AGTTGATCTTTACTACGCTGCTTGTACTCGAAACCAACGTTCTTTACATGAGTTCT
35 AAGATGGCTGCATATGACGAGAAAGATCCACTATTAATTTGTTTATGATGATTTGCCA
TTTGTGCCAGTGCCTCCACCGCGAGGCGTAAAGCTGATGATGATATTCAGGTAATTT
GCCAATTTGTTATCCGCACTATATCAGCAACCGTTTTCGAAAAGAAATGCTTTGGGA
AGTTGTTGGTAAACGCTGTTATCCGTAGCCAGACTTAATCTATCTGATAGTTTGGCGCA
ATATCCCTCCAAAATGTTGGCAATGAGATCATCAGATAATCAAGTACATTTTCTCA
40 ACTGAGGAATCGCATTTAGATTTGCCAACAATGTAATTCACCTTTAAACATATCAGA
TGGAGGCTTGTATAATCTAAGGCTAGAATAATGTCGATTTAGGAAGAGAGGAGCAAA
AGTAAAAGCTGTTTAAAGAGGTTTATTTGGATAAAACATAAACA AAAATTTCAAAA
GAATTTGATCTTTCAATTTTATAGGATAATAGCGCACTTTGAACTTCTTTGGGG
45 TAAACATTAAGCAAGGAATGGAATTTGTCAAAAGGTAATAAGTGGCCAAATTTCAAAAC
CCTAGTTAAGTGAAGTGTATAATGATGCTTTAATTAAGGTTCAAGTATAAACAAGGACA
CTTTTATTAATTCGATCACTAATAAGAGGACATCAAAA

SEQ. ID NO:68

45 Nucleotide sequence of DNA region (1000 bp) up-stream from the D15 gene
from *Haemophilus influenzae* (HiRd)

50 TCGATGTAATCCTATATAAATTTATAGACGTAAAAAATCATAAATAATGCAAAACCGTT
AAGCTTAATAACAGTCTGCGCCAAATTCGATAACAGATGCTTTGACCCCGCTCAGAAACA
GGTTTTCCTTTAACAGCTTCCATTTGTTAAAAAACTAAATGACCGCCATTAATACTGGT
AATGGAATAAATTCATAATCCCTAAATTTACACTAATCAATGCCATAAACTTAAAAAA
55 TACACCAATCCRAATTTGCTGATGCGCCAGCAGCCTTTGCAATAGAAATGGCCCACTT
AAATTAATTAATGCAAAATCGCCAGTAAGTAATTTCCCTAATATTTCAAGGTTAAAAAG
GAAAGCTGCTGTTTTTCAATGCTTTTTGTAAGATTCAAGAAATCAATAATTTAAT
TCAGTACGCTATTCATCCGTAATTTGTTAAGGCTGGGCTAACCCCAAAAACCAATTTG
60 CCATTTGATTAAGCACTGGAGTTAGGACTTTGTCAAATGTTTCCATTACGTTCAACT
TTAATAGAAAAGATTGCGCTTGTTCGACCTGTTTATAAATCTTGGCAAGAAATGGG
GTTAAATTTCTTTAAAAATTTTATCACGATTTGTAACCGACTTCTCAGCGGAGAA
TTTTGAACAACCTTTAGAAAGCACAATTTCAATTTAGGACGATAGGCATAATCCCTAAT
65 GCCTCAAAGCCTTTCTTTTTCAGGATCGAATGTCCAATTTGTAAGATTTAAAGTCCGT
TGTTGTTCAATTTAGAATTTGAAAGGAGAAAGGCTAATCTCAACATTAGGCTCCCCCAAT
TTTGGCCAGTAGCATTTGATGTTTTCCCAATCTGAGTTCTTCCCATCAATTTGTA
AGAATTTGCGTATTTGGGTTCAATGTTGGGCTTGTGCTGCGATTTGATTTGGTGTATTGAT
TCAATCACTGGTTTAAACCGTTGGCAATTCATAAAGGTAAT

SEQ. ID NO:69

70 Nucleotide sequence of DNA region (1000 bp) up-stream from the Omp26 gene
from *Haemophilus influenzae* (HiRd)

TTTGATAAATATCCTTAATTAATGATGGGTTTAAATTTTCTCTGCCAAATTAATTTAG
CGAGAGAACGTTGTTTTGAGTTCTGATGAAGAAAAAGTTCAATTTATAGAAAGAAC
TCCAATCAATAATGGAACGTTTCGACATCATCAATTTCAATTTTTTAATTTGGTTGGC
ATAAGAGAATACCAATGGCCCAATAGGAGATTGCCATTGGAATCCGACACTGTAGAGGC

5 GCGAATACGGCTTGATTGGCCATAATCGGGTAAGCTTTTTAAATACATTGTTATCTAACCC
ACTCTTATCCGATTCCACTTAGTATTCACAAACACTTGGCCGATCAACAAATAGGGAGGT
TCGGACTGTATTTGGCTTTATCACTCACAACCGTGGTGGTACAATAAGTCTGCACCT
CGCAGTTGGATTGCATTACCAACCAATCACATCAGAACTTATCTCTTAAAGTACCATT
ACCAATTACCATGTTCTGCATAAAATGGGTTAGTCCAATCTACCATAGCAAAACCCAG
TAATGAACCGATGCCACCGCTGTAAAGTTGATAGAACCGTAAACCGCTTGTTCCTAAA
ACCAATTGCATATCCTGCAGATGCTTTGACAGATACACCCAGAGGTGATCTCTGTCTAA
TGGGTAGAAACCCCTGACGCTGCACCTAGTTTTGAGTATTTGTTATCAGAACTGSAAT
10 AGTAACTCGTCCCAACGCTTGTCTTAAACCCCTTAAAGTTGGGAAATAGCCTCTATTAAG
GCTGTTATACTTCCAAACCAAGAAAATCAAAAGTCAATTCCTTTAAATGCCATTAACCTT
AAATTTCAATGATGAATATAAAATACGGCTTATATCTAGAGCAAACTTACTAATTTT
ATATAGGTATGGCTAATCTACATAATAGGATTTATTTTCAATTTACAGGAAACCTAA
AGTAACTACTTCCATAAGTCTGACGCTTATAGTTAGAGG

15 SEQ. ID NO:70
Nucleotide sequence of DNA region (1000 bp) up-stream from the P6 gene from
Haemophilus influenzae (HiRd)

20 TTGATTTCTCCTAAATGAGTTTTTATTTAGTAAAGTATGGAGACCAAGCTGGAATTT
AACTTGACCATCACTTCTGGAAGGCTCGCTTAAAGCGACCATCTGGGAAACCAATTTG
TAGCACTTTCCTAAGCCCTGTGAGAACTATAAATAATCAATAATCCATTTGGAGAGAG
GCTTGGGCTTTCGCTAGAAAAGATGACTAAGTACCTCGAAAACCGCCCTTGTGAGATC
TTGTTAACTACATTATGTTACCATTAATCAACAGTGTTTTTCCATCTGCACTAAT
TTGTGCGTACCGGACCAACCACTGCTGTGCACTACCAACCGCTTGCATCCATTCGATA
25 AACTTGGGCGAACCACTTCTATCGGATGTAATAAATGAAATTCGCTTGGGACCA
CGCTGGTTCAGTATTAACCCGACCACTCGTCAATGAGTAGGTGACCGCCATTTGC
TCCATAACGTAATATTCAGAACCACTACCGAGAAGAAAGCAAAAGTAAACGAGAAC
ATCTGGGAAAAGGCTGGTGGCCATATGCGCTTGAAGAAGTGCACACTTAACTGTC
GCCAGAAATTTAAATCCTGTACAAAGTTGTGATTTTTTTCAAAACGATACATAAGC
30 CAAACGCTGGCCGCTGGAGACCAAGCTGGAGACATAATGGTGGGCACTACGATGAC
GATAAATGATTATAGCCATCAATCTGCTACACGAACTTCAAGGTTGGGAAACCGCC
ATTTTTTGCACAACATAAGCGATACGAGTCTTAAAGGCAACCGATGCGAGTTAATTT
TTCAAAAATCTCATCGCTACAGTATGCGCCGATAGCGTAAACCATTTATTTGTAAGT
TAGCTATTTGCATTAATACAGTCCCTGGCGTACTGATGACCAACCGTATCAATTTAA
TTGATAAGTAATACTATAACCATACCCGATGGAACCACTT

35 SEQ. ID NO:71
Nucleotide sequence of DNA region (1000 bp) up-stream from the TbpA gene
from *Haemophilus influenzae* (non-typeable)

40 GCGATTAACCGAGTTTTGGGATTTAGTGCCTAAGAAAGCCCAAAAACCAAAATTA
TCCAGAGAAACCTTAATGATGGCAAGCTACTACTTTTAAAGAACTGATGCAAAAAC
AATCAACAGCCGATACAAACCAATAAACAACCAATGCAATAACCGATGAAAAAACC
TTTAAAGCCGAGATATCAAGTTTGGTGAAGCTGATTAATCTTAAATGCAATCAG
45 CCGTTCGCTTTTACCTGAAAAAATACTGATGATTTTCAATAGTAGTAGGCATCACT
GTAGGAATAAAGCTATAAAGTGAAGCATGTGCAAGAATCTAAGTATGTAATAATTT
GGTATGTTATGAAGACCCCTTAAAGAAAGAAAGAAAGAAAGAAAGAAAGAAAGAAAG
CARGAAAAAAGAAAGAAAGAAAGAAAGAAAGCAACGACGCAACATCTAGAGACTTATATCAA
TTCTTATAGGTACCCGCTACTGCAAGCCGACATACCTGCAACGGGAAACGTAAGATAT
50 CGCGTAAATGGTTTGGTTATATGTTGATGATGACAGCATCTTACTCACTACTGAGAT
AAAAATGCTCTCGCCGAGTTGATGTAATTTTGGCGATAAAAAGCTAACAGGCGAATTA
AAACGACAGCAATAAGAAATACCGTATTTAAAAATACGACAGCTTCAAGTGGTAAAG
AATGACTTCACTGGTACAGCAACCGCAAAATTTGTAATAGATGGTAAACATAGTCAA
ACTGGAATAACCAAAATTAATTAATAAGTAAATGGGGCAATTTATGGACCTAAG
GCTACAGAAATAGGCGGTTATTTCACTATAACGGAAATCTACAGCTAAAAATTCCTCA
55 ACCGTACCTTACCACCAATTCACCAATGCAAGAGCTGCAATTTGGTTGGAGCTAAA
AAACCAACAGTAGAAACCAACCAAGTAAATGGAATACTAAAA

SEQ. ID NO:72
Nucleotide sequence of DNA region (1000 bp) up-stream from the TbpB gene
from *Haemophilus influenzae* (HiRd)

60 TAGAATATATCTTATACAAAATGATAATGTTGCGATTATCATTTTTTTTTGTAAT
AATGCAACTTATAATTTTTAAAGTTCATGGATAAAAATGAAAAATGGCGTAAAAACAC
TTTTCTCTTATCAATTAATAGGCTTATCATTAACGAAATGAGCTTGGGCGAAGTTGCAC
GTCTAAAAATGATACNTTGCAAAATCAGATTCAAAGTGGGAAATAAAAACCTCTCTTT
65 TTCTCTATGCTTAAAGAAAGAAATACCAATAGGCAATATTTCTCTTTCCAAAAGCC
AATTAGCCACCACTCCAGGCTTGTGTTGGTGGGTTAATCTCTCTTATTCAAAATA
ACACTAGGCACTTCAACTCTTATTAACCACTATATAAACAATTTCTCAACAGATAAT
TCTTACTACTTGGGCAAGGCTATTGAAGCTGTAACAGGTGATTTACTCAATCTA
TTGCTTATATCGGAAATTTCCGCTGAGACGCACTTTACTACTTTTACGTTATTAAT
70 TAGCTCAAGCTCTATTTTTAACTATGAAAATGAAAGTGCACAAATCAATTTGAAAAT
TAGTACAGAGGTAGATGATAAAAATTTTAGGTGTTATGATCAGTATCTTTAACAC
TAAATCAGCGAATCAATGGATATGGCAAGTAGGATTAATTTTTTAAATGATGATAAT
TGAATAACGCTCCAAAAGTGCACAAAATTTGATGTTGACCGCTTGGGAAAAGAAA
GTGGCAGGGGTAGGCTATTTCTTATCAGTAGAAAAAATGGCCATGGGAGATCAAT
75 TTTTATGTAATAACTATGTTTAAATGGAATGAAAAATATTTGGGATAATAAAAAATACA
ATGAGGCTACTGTGCTATAGTGGTGGTTAGGCTATCAAACTGCCTCAGTTGAAGTCT

CGTTGTTCCTTTTCAAGAAAAACGCTGGTATGCAGCGGT

SEQ. ID NO:73

Nucleotide sequence of DNA region (1000 bp) up-stream from the HifA (pilin)

5 gene from *Haemophilus influenzae* (LKP serotype 1 genome)

TTATAAATGCTCCATNAAGAGGTTTGGCCCTATAAATNAGGCAATAAGATTATATA
AACCGTTTATAAATGCCAAAGCCCTTAATAACAGCAAACCTTTCTTTCCGAAAAAAG
TAAAAAATCTTCCATTATATATATATATATATATAANTTAAAGCCCTTTTGAAAAATTT
CATATTTTTTTGAATTAATTCGCTGTAGGTTGGGTTTTGCCACACTGGAGACATAAAA
10 AAAGATTTGTAGGTTGGCCCTAAGCCCAAGCGAACACTCAAAACACTGTANTGTGT
ATTAGGCACGGTGGGCTTATGCCCTCGCTACGGGAAATGAATAGGTAATATGGGCT
TAGCCCACTTATGGATTAATTAATGTGAAATGGGAAAAACATGTTAAAAAACACT
TTATTTTTTACCCACTATTTTTTGGCCGACTTTGTGCATTTTCCGCAATGCAGATGT
GATFATCACTGGCAACAGAGTGATTTATCCCGCTGGGCAAAAAATGTTATCGTGAAGTT
15 AGAAAAACAATGATGATTCGGCAGCATTTGGTGCAGCCCTGGATTGATATGGCAATCCAAA
TGCCGATCCAAAATACACCAAAAACCCCTTTGTGATTACCCCGCCTTTGTGCGAGTGA
AGCGAAATCAGGGCAAAGTTTGGGATACGTTTACAGCAGGACGAGCCTTTACCTGATGA
TCGCGAAAGCCTCTTTTAAATTTGTTAGATATTCGCGGAAACCTGATGCGGCATT
20 TCGGCAAAACAGCGAGCTTTATGCAAATGGCATTGCTCACGTTTGAAGTTGTTTTA
TCGCCCTGCAAACTCTCGATGGATTCTCGTATGCAATGAAAAAAGTAGTGTAAAGC
CACACCTGAAGGGGTTGGTGGATAATCAAAACCCCTTATATATGAACACTTGGTTT
GTTACATCAAAAATAACCTGGAAAAATGTCAAAATGGTTG

SEQ. ID NO:73

25 Nucleotide sequence of DNA region (1000 bp) up-stream from the HifE (tip pilin) gene from *Haemophilus influenzae* (LKP serotype 1 genome)

TAGTAGATTTCCGCAAGGGCAAAAAACAATGGTGTATTTAACTCACTTGGCAAATG
GCGAGCCAGTCCAAATGGCATCCACCGCACAAAGTAGCGAAGGGGCTTTGTGGCGATG
30 TGCTGCAAGGTTGGTGTCTTTCCGCTAATAAACTTCCCAAGCCAAAGGGGAGTTAATCG
TCAAAATGGGGTGGCGAGAAAGCGAAACAATCGGTTTCCCAATCAAGTTGATTGGATA
ACGCACAAAATCAAAAGTACAGATTTCAATGCAAAAACCGCAAAAATAAATAATGAAGAGG
ATTTATGCAAAAAACCCAAAAAATTAACCGCGCTTTTCCATCAAAAAATCCACTGTAC
TTGTAGTGGAGCAAAATATAGTGGAGCAAAATATAGTGGCTCAAAAATGCTTAGGTTTCA
35 TCGCTGCGCTTGTCTTGTCTGGGCGCTCTGCTTGTGATTGCAATGGCACTGCGCTCTTA
TGCTTACGATGGCAGAGTGACCTTTCAAGGGGAGATTTAAGTGAATGGCACTTGTAAAAT
TGAACACAGCAGCCCAAAATCGCACGGTTACCTTGCACACTGGAAAGCTAATTTAG
ACACGTAAGGCAAAAGCGCCCTGCTGCTTTTCCATCACTTAAGAATGCAATGCAATGC
AGATGATGCTAAGAAGCTAATCTGCTATTTAAAGGGGGAGCAACACACAGGGCAATC
40 TTATCTTTCCAAATAGGCGGCAAGCGCAAGCCCAAGCTGGCATTCAAATTTGCAA
AGCCGATGGCATAGGCAAGCCCTTCAAGGTGGAGCGCACCGAAGCCAAACCGGAAAAAG
CCCGACACAGGTAAGCGCAAAAGCGCACAGTTTATCAACCCCTTTTGGTACTTTGG
CTCGTATTACGCCACAGGTAAGCGCACCGGAGGACGTTGAAGCCACTGCAACTTTTG
AAGTGCAGTATACTAAAATATTTTATTCAGTGAAAAA

45 SEQ. ID NO:75

Nucleotide sequence of DNA region (1000 bp) up-stream from the P2 gene from

Haemophilus influenzae (HiRd)

1 TTATCCGCTA ACATTTCATC AGTAATTTCCA TGAACCTTAA TCGCATCAGG
51 ATCANCAGGG CGATCTGGCT TAATATAAAT ATGAYAATTA TIACCTGTGT
101 AAGCACGATT TATTAATTCA ACTGCACCAA TTTCAAATAT GCAGTGTCTCT
151 TCATAATGGC CGCCAAGCTG ATTCATACCT GTAGTTTCAG TATCTAATAC
201 AATTTGGCGA TTGGGATTAA TCATTITGTC AACCTATCTC TTTCCATTAA
251 AATATCTGCC ATTCACACA ACAACCTTTC TGTATGCK AAACAGATTG
301 AATATTTTAC TGATGGATCT TGCTTAGGTA ATCCAGGGC GGGCGAATT
55 351 GGTGCCGTAT TCCGTATTA ACAACATGAA AAAACCTCT CCAAGGCTA
401 TTTCCAAAC ACCAATAATC GAATGGAAAT ACCCGCTGT ATTGAAGCAT
451 TAAATACNTT AAAAGAACCT TGCTTGATCA CGCTTATAG TGATAGCCAA
501 TATATGAAA ATGGCATAAC CAAATGGATC TTTACTGGA AAAAAATAA
551 TTGGAAGCA AGTCTCGAA AGCCTGTAAA AAACCAAGAT TTTGGGATAG
60 601 CCTTAGATGA ATCCATCCAA CGTCATAAAA TTAATTTGGA ATGGGTAAAA
651 GGCCATGCTG GACACAGAGA AAATGAAAT TGCGATGAAT TAGCAAAAA
701 AGGGGACGAA AATCCGACAT TGGAAAGTAT GGGGTACATA GAAGAATAAT
751 ACAACTGATA TAACGTCATA TTTTTCGATA CCTAAAAATA TTTAATACTT
801 AAACCTAAAA CAGAATAAAA AATAATCAAA TTCATTAAA AATGTGATC
851 TCGATCAGAT TTCAGAAAA TTAATAATTT GGAGTATTGA CATCAAAAAT
901 TTTTTTTGTA AAGATGCAGC TCGTCCGTTT TGGCGATTGG ACAATTTCTAT
951 TGGAGAAAAG TTCATCATA GATAGTAAAC AACCATAAGG AATACAAAT
1001 A

70 SEQ. ID NO:76

Nucleotide sequence of DNA coding region (partial) of the *Moraxella Catarrhalis* HtrB gene

1 TCAGTGGCTTG GTTTTTAAAG ATATGTACCG CTGTCAGTCC TGCAIGGATT
 51 GCGGGCTGT GCGTCTTATA TTCTCTATCA TTGCAGGCTT AGTATTTATC
 101 GCRGATCCA AGCCAAATTA ATCTTGCTTC ACCCGAAGAT GCCAGACGCA
 151 CAGCGGCAAA AACTCGGCAA ACAARTCCTA AAAAATCAGC TCATCAGTGC
 201 AGTCGACAGT CTTAAACTT GGGCAATGCC ACCAAAATGG TCTATCGCAC
 251 AAATTTAAAC GGTTCATCAT GAGATATCC TAATCAAGC ACTTGCCAAAT
 301 CCAAGTGCTA TGCTTGGCAT TGTGCTCAT ATCGGCACTT GGGAGATGAT
 351 GAATGCTTGG CTCATACCT TTGGCTCCC TACTATCATG TATAAGCCCA
 401 TCAAAAATGC GCGGTAGAT CGCTTGTGT TACAGGGGCG TGAAGACTA
 451 AATGCCAGCC TTGTACCCAC AGATGCTAGT GGTGTTAAGG CAATTTTTAA
 501 AACACTCAA GCAGGTGGAT TTAGTATCAT ACTGCCCGAC CATGTACCTG
 551 ATCCATCAGG TGGTGAGATT GCTCCTTTT TTGGTATTA AACCCTAACC
 601 AGTACGCTGG CGTCAAAGCT TGCTGCAAAA ACTGGTTTGG CTCTTGTGG
 651 CTTAAGTGT ATTCCGGGTG AAGATGGCGA TGGTTTGA AATTTTTGTT
 701 ATGAATTTAA TGATGAACAA CTTTATTCAA AAAATACCAA AATTGCAACC
 751 ACTGCTTTAA ATGGTGGAT GGAACAAATG ATTTATCCAC ATTTTTGCA
 801 TTATATGTGG AGCTATGCT GGTTCAGCA TACACCCTA TTAATAATC
 851 CTTATTTACT TAATGAAAT GAGCTAAAA AAATAGCCAT AAAGCTCAA
 901 GCCATGTCAA AGGATAGTTA TGAG

Protein Seq: 25% identity and 35% similarity with HtrB from *E. coli*

1 SVLGFRLVVP LSVLHGLAAC ASYISYHCLR SIYRSIQANL ILVHPKMPDA
 51 QRKLAQKIL KNQLISAVDS LKTWAMPKWK SIAQIKTVHH EDLILKALAN
 101 PSGMLAIVPH IGTWEMNMAN LNTFGSPTIM YKPIKNAVD RFLVQGRERL
 151 NASLVPTDAS GVKAIFKTLK AGGFSIILPD HVPDPSGGEI APFFGKILT
 201 STLASKLAAK TGCAVLVGLSC IRREDGDGFE IFCYELNDEQ LYSKNTKIAT
 251 TALNGAMEQM IYPHFLHYWV SYRRFKHTPL LNNPYLLNEN ELKKIAIKLQ
 301 AMSKDSYE

SEQ. ID NO:77

Nucleotide sequence of DNA coding region of the *Neisseria* (meningococcus B)

HtrB gene

1 ATGTTTCGTT TACAATCCG GCGTTCGCC CCTTGGCAA CCGCATSCA
 51 CATCCTGTTG ACCGCCCTGC TCAAAATGCT CTCCCTGCTG GCACCTTCTT
 101 GTCTGCACAC GCTGGGAAAC CGGCTCGGAC ATCTGGCCTT TTACCTTTTA
 151 AAGGAAGACC GCGCGCGCAT CGTCCCAAT ATGCTCAGG CAGCATGAA
 201 TCCCGACCCC AAAACAGTCA AAGCCGTTT TCGGAAACG GCAAAAGSCG
 251 GTTTGGAACT TGCCCCCGCG TTTTTCAGAA AACCGAAGA CATGAAACA
 301 ATGTTCAAAG CGGTACACGG CTGGAAACAT GTGCAGCAGG CTTTGGACAA
 351 ACAOAGAGGG CTGTATTTCA TCACGCCGCA CATCGCAGC TACGATTTGG
 401 GCGGACGCTA CATCAGCCAG CAGCTTCCGT TCCCGCTGAC CGCCATGTAC
 451 AAACCGCCGA AAATCAAAGC GATAGACAAA ATCATGCAAG CCGGCAAGST
 501 TCGGGGCAAA GGAACAAACG CGCCTACCA CATACAAGG GTCAACACAA
 551 TCATCAAAGC CCTGCGTTGG GGGGAAGCAA CCATCGTCTT GCCCGACCC
 601 GTCCCTCCCT CTAAGAAGG CCGGGAAGCG GTATGGGTGG ATTTCTTCGG
 651 CAACCTGCC TATACCATGA CGCTGCGGCG AAAATTGGCA CACGTCAAAG
 701 GCGTGAACAC CCGTTTTTTC TGCTGCGAAC GCCTGCTGG CCGACAAGGT
 751 TTGATTTTGC ACATCCGCCG CCGTCAAGGG GAATTGAACG CCGACAAGC
 801 CCAATGATGC CCGGTGTTC ACCGCAATCG CGAATATTGG ATACCCGCTT
 851 TTCGACGCA GTATCTGTTT ATGTACAACC GCTACAAAAT GCCG

Protein Sequence - 30% identity and 38% similarity with HtrB *E. coli*

1 MFRLQGLFPP PLRTAMHILL TALLKCLSLP PLSCLHTLGN RLGHIAFYLL
 51 KEDRARIVAN MRQAGMNPDP KTVKAVFAET AKGGLLELAPA FFRKPEDLET
 101 MFKAVHGWNEH VQQALDKHEG LLFITPHIGS YDLGGYISQ QLPPPLTAMY
 151 KPPKIKAIKDK IMQAGRVKRG GKTAPTSISQ VKQIILKALRS GEATIVLPEDH
 201 VPSBQGGEG VVVDFFGKPA YMTLAAKLA HVKGVKTLFF CCERLPGGQG
 251 FDLHIRPVQG ELNGDKAHD AVFNRAEYV IRRFPPTYLF MYNRYKM?

SEQ. ID NO:78

Nucleotide sequence of DNA coding region of the *Haemophilus influenzae* (non-

typeable) HtrB gene

1 ATGAAAACG AAAAATCCCT TCAATTTCAA CCGCACTTT TAGCCCCAAA
 51 ATACTGGTCT TTTTGGCTAG GCGTGGCAAT TTGGCGAAGT ATTTTATGTC
 101 TTCCCTATCC TATTTTGGCG CATATTGGTC ATGTTTCCG TTGGCTGTTT
 151 TCACATTTAA AAGTGGGTAA ACGTCGAGCT GCCATTGCAC GCCGTAACT
 201 TGAACCTTGT TTCCCTGATA TGCTGAAAA CGAACGTGAG ACGATTTTGC
 251 AAGAAAATCT TCGTTCAGTA GGCATGGCAA TTATCGAAAC TGGCATGGCT
 301 TGGTTTTGGT CCGATTACAG TATCAAAAAA TGGTCGAAAG TTGAAGGCTT
 351 ACRTTATCTA AAAGAAAATC AAAAAGATGG AATTGTTCTC GTCCGTGTTT
 401 ATTTCCTAAC CCFAGACTT GCGGACGSCA TCATGCGTTT ACATCATCTT
 451 GGCATTTGGT TTATCGTCC AAATGATAAT CCTTGCCTG ATTGCCTA
 501 AACACAAGCG CGTTTACGCT CCAATTAAGA TATGCTTGT CCAATGAT
 551 TACGCGAAT GATCAAGCT TTACGCCACG AAGAACCAT TTGGTATGG
 601 CCTGATCACG ATTAGGCGAG AAAAATGCCC GTTTTTGTTT CTTTTTTTGC

651 AGTACCTGAC ACTTGCACTA CTACTGGTAG TTATTATTTA TTGAATCCT
 701 CGCAAAACAG CAAAGTGATT CCATTGGCGC CATTACGCAA TAAAGATGGT
 751 TCGGGCTATA CGGTGAGTAT TTCAGCGCCT GTTGTATTTA CGGATTTACA
 801 AGATGAAACG GCGATTGCTG CCGCAATGAA TCAATTCGTA GAAAGGAAA
 851 TCATGAAGGG CATATCACA TATATGTGGC TACATCGCCG TTTTAAAAACA
 901 CGTCCAGATG AAAATACGCC TAGTTTATAC GATTAA

Protein Sequence - 57% identity and 66% similarity with HtrB *E. coli*
 1 MKNEKLPQFQ PHFLAPKYWL FWLGVAIWRS ILCLPYPILR HIGHGFGWLF
 51 SHLVKGRRA AIARRNLELC FDPMPENERE TILQENLRV GMALIEISMA
 101 WFWSDSRIKK WSKVEGLHYL KENQKDGIVL VGVHFLTLEL GARIIGLHHP
 151 GIGVYRPNND PLLDWLQIQG RLRSNKDMLD RKDLRGMKA LRHEETIWA
 201 PDHDIYGRKNA VFVFFFAVED TCTTTGYYL LKSSQNSKVI PFAFLRNKDG
 251 SGYTVSISAP VDFDLQDET AIAARMNQIV EKEIMKGISQ YMWLHRRFKT
 301 RPDENTPSLY D*

SEQ. ID NO:79

Nucleotide sequence of DNA coding region of the Haemophilus influenzae (non-typeable) MsbB gene

20 1 ATGTCGGATA ATCAACAAAA TTTACGTTG ACGGCGAGAG TGGGCTAIGA
 51 AGCGCACTTT TCATGGTCTG ATTTAAAGCC TCAATATTGG GGGATTTSGC
 101 TTGGTATTTT CTTTTTATTG TTGTAGACAT TTGTGCCTTT TCGTCTGCGC
 151 GATAAATTGA CGGAAAAATT AGGTATTGG ATTGGGCATA AAGCAAAGAA
 201 ACAGCGTACG CGTGACACAA CTAACCTGCA ATATTGTTTC CCTCATTSGA
 251 CTGAACAACA ACGTGAGCAA GTGATTGATA AAATGTTTC GGTGTGCTCT
 301 CAGGTATGT TTGGTATGG TGAGATTGCC ATCCGTTCAA AGAAACATTT
 351 GCAAAAAAGC AGCGAATTTA TCGGCTTGA ACATATCGAA CAGCAAAAAG
 401 CTGAAGGAAA GAATATTATT CTTATGGTGC CACATGGCTG GCGGATTSAT
 451 GCGTCTGGCA TTTTGGTGA CACTCAAGGC ATGCCAATGA CTTCATSTTA
 501 TAATCCACAC CGTAATCCAT TGGTGGATTG GCTTTGGACG ATTACAC3CC
 551 AACGTTTCGG CGGAAAAAGT CATGCACGCC AAAATGGTAT TAAACCTTTT
 601 TTAAGTCATG TTCGTAAGG CGAAATGGT TATTACTTAC CCGATGAAGA
 651 TTTTGGGGCG GAACAAAGCG TATTTGTTGA TTTCTTTGGG ACTTATPAAG
 701 CGACATTACC AGGGTTAAAT AAAATGGCAA AACTTCTAA AGCCGTTSTT
 751 ATTCCAATGT TTCTCGTTA TAACGCTGAA ACGGGCAAA ATGAAATSGA
 801 AATTCATCCT CCAATGATTT TAAGTAGTGA TCCGAAACA TCAGCCCSAG
 851 CAATGAACGA AGAATAGAA TCTTTGTTA CCGCACGCC AGAGCATAT
 901 GTTSGATTT TGCAATTATT CGCTACAAG AAAGATGCGC AAGATCTTTA
 951 TGATTTAA

Protein Sequence - 45% identity and 56% similarity with MsbB *E. coli*
 1 MSDNQQLRL TARVYEAHF SMSYLKQVW GSWLGFLL LLAFVDFRLR
 51 DKLTLKLIW TGHAKKQRT RAGTLLQYCF PWFTEQREQ VIDKMFNVA
 101 QVMPGLGEIA IRSKHLQKR SFVIGLEHIE QAKAEGXNLI LMPVHGMAD
 151 ASGIIILHTQG MPMTSMYNPH RNPLVDLWLT ITRQRPGKM HARONGIKPF
 201 LSHVRKGMG VYLPDEDFGA EQSVFVDFRG TYKATLPLN KMAKLSKAVV
 251 IPMFRYNAE TGKYEIEHP AMNLSDDEPQ SARAMEIEE SFVTPAPEQV
 301 VVILQLLRTR KDGEDLYD*

SEQ. ID NO:80

Nucleotide sequence of DNA coding region of the Moraxella catarrhalis MsbB gene

1 ATGAGTTGCC ATCATCAGCA TAAGCAGACA CCCAAACAGC CCATATCCAT
 51 TAAGCATATG CCAAGCTTGA CAGATACTCA TAAACAAGT AGCCAAGCTG
 101 AGCCAAAATC GTTTGAATGG GCGTTTTTAC ATCCCAAATA TTGGGGAGTT
 151 TGGCTGGCTT TTGCGTTGAT TTIACCGCTG ATTTTCTAC CGGTGCGSTG
 201 GCAGTTTTGG ATCGGCAAGC GTCTGGGATC TTTGGTACAT TACTTAGCTA
 251 AAAGCCGAGT TCAAGCAGCT CTAACCAACC TGCAGCTTAC CTTCACAAAT
 301 CAACCAAAT CAAAAACACAA GGCCACCGCA CGCAAGTAT TTATTAATCA
 351 AGTATGGT ATTTTGAAGA GTTTATGTCG ATGGTTTGG CCTAATGTCT
 401 TTAACCGCAC TTTTACGATT TCGGTTTAC AGCAATTTGAT TGAAGCCAA
 451 AANCAAAAATA AAGCGGTGAT TTTACTTGGT GGACATCGCA CACCGCTTGA
 501 TTTGGGCGGT CGTATATGTA CACAGTTTTT TCGCGCGGAC TCGGTGTATC
 551 CCCCACAAA CAACCCTTTC CTTGAATGGT TTATCTATRA TGCAGCGCCG
 601 TGTATCTTTG ATGAGCAJAT CTCAAATCGT GATATGAAA AACTCATCAC
 651 TCGGCTCAAA CAAGGTCGSA TAATTTGGTA TTCACCTGAT CAAGATTTTG
 701 GTCTTGAGCA TGGCGTGATG GCGACCTTTT TTGGTGTGCG TGACGCAAG
 751 ATTACCGCTC AGCGTGTCT TATTRAGCTG GGTGATAAAG CCAATCTCC
 801 TGTATCATC ATGATGATA TGCTCAGACA AACGCCGCTG TATATCGCAA
 851 AAGGTCACCG TCCACATTAT CACATCAGCC TAAGCGCTGT GTTAAAAAT
 901 TATCCACGCG ATGACGAAAC CGCGATGCT GAACGATCA ATCGACTGAT
 951 TGAGCAAAAT ATTCAAAAG ATTTAACCCA GTGGATGTGG TTTCACTGCG
 1001 GCTTTAAAA TCAAGCCGAT GACACCAATT ACTATCAACA TTAATG

Protein Sequence - 28% identity and 37 similarity with MsbB of *E. coli*
 1 MSCHHQHKQT PKHAISIKHM PSLTDTHKQS SQABPKSFEW AFLHPKYGVV

51 WLAFAILILPL IFLPLRWQFW IGKRLGILVH YLAKS RVQDT LTNLQLTFPN
 101 QPKSKHKATA RQVFINQIG IFESLCAWFR PNVFKRTFSI SGLQHLIDAQ
 151 KQNKAVTLLG GHRITLDLGG RLCTQFFAD CVYRFPQNPPL LEWFLYNARR
 201 CTFDEQISNR DMKLLITRLK QGRITWYSPD QDFLEHGVN ATEFQVPAAT
 251 ITAQRRLLIKL GDKANPPVII MMDMLRQTFD YIAKGRPHY HILSLAVLKN
 301 YPSDDETADA ERINRLIEQN IQKDLTQWNV FRRRFKQAD DTNYIQH*

SEQ. ID NO:81

Nucleotide sequence of DNA coding region of the *Neisseria* (meningococcus B)

10 MsbB gene

1 ATGAAATTTA TATTTTTTGT ACTGTATGTT TTGCAGTTTC TGCCGTTTGC
 51 GCTGCTGCAC AAACCTGCGC ACCTGACGGG TTGCTCGCC TACCTTTTGG
 101 TCAAACCCCG CCGCCGTATC GCGGAATCA ATTTGGCAA ATGCTTTCC
 151 GAGTGGGACG GAAAAAGCG CGAAACCGTA TTGAAGCAGC ATTTCAAACA
 201 TATGGCGAAA CTGATGCTTG AATACGGCTT ATATTGGTAC GCGCCTGCC
 251 GCGCTTTGAA ATCGCTGGTG CGTTACCGCA ATAAGCATT TTTGACGAC
 301 GCGCTGGCGG CCGGGGAAAA AGTCATCATT CTGTACCCGC ACTTCACCGC
 351 GTTCGAGATG GCGGTGTACG CGCTTAATCA GGATGTACCG CTGATCAGTA
 401 TGTAATCCCA CCAAAAAAC AAGATAATGG ACGCACAGAT TTTGAAAGGC
 451 CGCAACCGCT ACGACAATGT CTTCCTTATC GGGCGCACCG AAGCGGTGCG
 501 CGCCCTCGTC AAACAGTTCG GCAAAAGCAG CCGCCCGTTT CTGTATCTGC
 551 CCGATCAGGA TTTCGGACGC AACGATTGCG TTTTGTGGA TTTTTCGGT
 601 ATTCAGACGG CAACGATTAC CGGCTTGAGC CGCATTGCCG CGCTTGCAA
 651 TGCAAAAAGT ATACCCGCCA TCCCGTCCG CGAGGCGGAC AATACGGTTA
 701 CATTGCATTT CTACCCGGCT TGGGAATCCT TTCCGAGTGA AGATGCGCAG
 751 CCGGACGCGC AGCGCATGAA CCGTTTTATC GAGGAACCGT GCGCGAACAT
 801 CCGGACGAGT ATTTTGGCT GCACAAGCGT TTCAAAAACC GTCGGAAGG
 851 CAGCCCCGAT TTTTACTGAT ACGTAA

30 Protein Sequence - 25% identity and 36% identity with MsbB *E. coli*

1 MKFIFPVLYV LQFLPFALLH KLADLTGLLA YLLVKPRRRI GEINLAKCFP
 51 EWDGKKRETV LKQHFHMAK LMLEYGLYNY APAGRLKSLV RYRNKHYLDD
 101 ALAAGEKVII LYPHPTAFEM AVYALNQDVP LISMSYSHQN KILDAQILKG
 151 RNRYDNVFLI GRTEGVRLV KQFRKSSAFP LYLDPDQFGR NDSVPVDFPG
 201 IQTATITGLS RIAAALANAKV IPALPVREAD NTVLHFYPA WESFPSEDAQ
 251 ADAQRMNRFI EEPKANIPSS IPGCTSVSKP VRKAAPIFTD T*