

**(12) PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 199663312 B2**  
**(10) Patent No. 707023**

(54) Title  
**Monkey monoclonal antibodies specific to human B7.1 and/or B7.2 primatized forms, pharmaceutical compositions**

(51)<sup>6</sup> International Patent Classification(s)  
**C12N 005/12 C07K 016/00**  
**A61K 039/395**

(21) Application No: **199663312**

(22) Application Date: **1996.06.06**

(87) WIPO No: **WO96/40878**

(30) Priority Data

(31) Number	(32) Date	(33) Country
<b>08/487550</b>	<b>1995.06.07</b>	<b>US</b>

(43) Publication Date : **1996.12.30**

(43) Publication Journal Date : **1997.02.20**

(44) Accepted Journal Date : **1999.07.01**

(71) Applicant(s)  
**IDEC Pharmaceuticals Corporation**

(72) Inventor(s)  
**Darrell R. Anderson; Peter Brams; Nabil Hanna; William S. Shestowsky**

(74) Agent/Attorney  
**WATERMARK PATENT and TRADEMARK ATTORNEYS, Locked Bag 5, HAWTHORN VIC 3122**



<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 5/12, A61K 39/395, C07K 16/00</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/40878</b>
			<b>(43) International Publication Date:</b> 19 December 1996 (19.12.96)
<b>(21) International Application Number:</b> PCT/US96/10053 <b>(22) International Filing Date:</b> 6 June 1996 (06.06.96) <b>(30) Priority Data:</b> 08/487,550 7 June 1995 (07.06.95) US <b>(71) Applicant:</b> IDEC PHARMACEUTICALS CORPORATION [US/US]; 11011 Torreyana Road, San Diego, CA 92121 (US). <b>(72) Inventors:</b> ANDERSON, Darrell, R.; 1851 Navajo Place, Escondido, CA 92029 (US). BRAMS, Peter; 4303 Proctor Place, San Diego, CA 92116 (US). HANNA, Nabil; 3255 Fortuna Ranch Road, Olivenhain, CA 92024 (US). SHESTOWSKY, William, S.; 1155 Thomas Avenue, San Diego, CA 92109 (US). <b>(74) Agents:</b> GESS, E., Joseph et al.; Burns, Doane, Swecker & Mathis L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).			<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> MONKEY MONOCLONAL ANTIBODIES SPECIFIC TO HUMAN B7.1 AND/OR B7.2 PRIMATIZED FORMS, PHARMACEUTICAL COMPOSITIONS			
<b>(57) Abstract</b> <p>The present invention relates to the identification of macaque antibodies to human B7.1 and B7.2 by screening of phase display libraries or monkey heterohybridomas obtained using B lymphocytes from B7.1 and/or B7.2 immunized monkeys. More specifically, the invention provides four monkey monoclonal antibodies 7B6, 16C10, 7C10 and 20C9 which inhibit the B7:CD28 pathway and thereby function as effective immunosuppressants. The invention further provides the complete DNA and amino acid sequences of the light and heavy chain of three primatized antibodies derived from those monkey monoclonal antibodies which bind B7.1 and possibly B7.2, primatized 7C10, primatized 7B6 and primatized 16C10. These primatized and monkey antibodies may be used as specific immunosuppressants, e.g., for the treatment of autoimmune diseases and to prevent organ transplant rejection.</p>			

-1-

**MONKEY MONOCLONAL ANTIBODIES SPECIFIC TO HUMAN B7.1 AND/OR B7.2 PRIMATIZED FORMS. PHARMACEUTICAL COMPOSITIONS**

5

**FIELD OF THE INVENTION**

The present invention relates to the manufacture and identification of novel monoclonal antibodies to human B7, i.e., human B7.1 and human B7.2 and primatized forms thereof. More specifically, the present invention relates to the production and identification of macaque antibodies to human B7, i.e., human B7.1 and human B7.2 produced by screening of phage display libraries and monkey heterohybridomas using B lymphocytes obtained from B7 immunized monkeys.

15

The invention further relates to specific primatized antibodies which bind to human B7, i.e., human B7.1 and B7.2 as well as their corresponding amino acid and nucleic acid sequences.

20

Also, the present invention relates to pharmaceutical compositions containing monkey monoclonal or primatized antibodies specific to human B7.1 and/or human B7.2 and their use as immunosuppressants by modulating the B7:CD28 pathway, e.g., for the treatment of autoimmune disorders, and the prevention of organ rejection.

25

**BACKGROUND OF THE INVENTION**

30

The clinical interface between immunology, hematology, and oncology has long been appreciated. Many conditions treated by the hematologist or oncologist have either an autoimmune or immunodeficient component to their pathophysiology that has led to the widespread adoption of immunosuppressive medications by hematologists, whereas oncologists have sought

-2-

immunologic adjuvants that might enhance endogenous immunity to tumors. To date, these interventions have generally consisted of nonspecific modes of immunosuppression and immune stimulation. In addition to the limited efficacy of these interventions, toxicities secondary to their nonspecificity have also limited their overall success. Therefore, alternative strategies have been sought.

Elucidation of the functional role of a rapidly increasing number of cell surface molecules has contributed greatly to the integration of immunology with clinical hematology and oncology. Nearly 200 cell surface antigens have been identified on cells of the immune and hematopoietic systems (Schlossman SF, Boumsell L, Gilks JM, Harlan T, Kishimoto, C, Morimoto C, Ritz J, Shaw S, Silverstein RL, Springer TA, Tedder TF, Todd RF: CD antigens (1993), Blood 83:879, 1994). These antigens represent both lineage-restricted and more widely distributed molecules involved in a variety of processes, including cellular recognition, adhesion, induction and maintenance of proliferation, cytokine secretion, effector function, and even cell death. Recognition of the functional attributes of these molecules has fostered novel attempts to manipulate the immune response. Although molecules involved in cellular adhesion and antigen-specific recognition have previously been evaluated as targets of therapeutic immunologic intervention, recent attention has focused on a subgroup of cell surface molecules termed co-stimulatory molecules (Bretscher P: "The two-signal model of lymphocyte activation twenty-one years later." Immunol. Today 13:73, (1992); Jenkins MK, Johnson JG: "Molecules involved in T-cell co-stimulation." Curr Opin Immunol 5:351, 1993; Geppert T, Davis L, Gur H, Wacholtz

-3-

M. Lipsky P: "Accessory cell signals involved in T-cell activation." Immunol Rev 117:5, (1990); Weaver CT, Unanue ER: "The co-stimulatory function of antigen-presenting cells." Immunol Today 11:49, (1990); Stennam RM, Young JW: "Signals arising from antigen-presenting cells." Curr Opin Immunol 3:361, (1991)).

Co-stimulatory molecules do not initiate but rather enable the generation and amplification of antigen-specific T-cell responses and effector function

(Bretscher P: "The two-signal model of lymphocyte activation twenty-one years later." Immunol. Today 13:73, (1992); Jenkins MK, Johnson JG: "Molecules involved in T-cell co-stimulation." Curr Opin Immunol 5:351, (1993); Geppert T, Davis L. Gur H. Wacholtz M.

Lipsky P: "Accessory cell signals involved in T-cell activation." Immunol Rev 117:5, (1990); Weaver CT, Unanue ER: "The co-stimulatory function of antigen-presenting cells." Immunol Today 11:49, (1990); Stennam RM, Young JW: "Signals arising from antigen-presenting cells." Curr Opin Immunol 3:361, (1991); June CH, Bluestone JA, Linsley PS, Thompson CD: "Role of the CD28 receptor in T-cell activation." Immunol Today 15:321, (1994).

Recently, one specific co-stimulatory pathway termed B7:CD28 has been studied by different research groups because of its significant role in B and T cell activation (June CH, Bluestone JA, Linsley PS, Thompson CD: "Role of the CD28 receptor in T-cell activation." Immunol Today 15:321, (1994); June CH, Ledbetter JA: "The role of the CD28 receptor during T-cell responses to antigen." Annu Rev Immunol 11:191, (1993); Schwartz RH: "Co-stimulation of T lymphocytes: The role of CD28, CTLA-4, and B7/BB1 in interleukin-2 production and immunotherapy." Cell 71:1065, (1992)). Since this

-4-

ligand:receptor pathway was discovered four years ago, a large body of evidence has accumulated suggesting that B7:CD28 interactions represent one of the critical junctures in determining immune reactivity versus anergy (June CH, Bluestone JA, Linsley PS, Thompson CD: "Role of the CD28 receptor in T-cell activation." Immunol Today 15:321, (1994); June CH, Ledbetter JA: "The role of the CD28 receptor during T-cell responses to antigen." Annu Rev Immunol 11:191, (1993); Schwartz RH: "Co-stimulation of T lymphocytes: The role of CD28, CTLA-4, and B7/BB1 in interleukin-2 production and immunotherapy." Cell 71:1065, (1992); Cohen J: "Mounting a targeted strike on unwanted immune responses" (news; comment). Science 257:751, (1992); Cohen J: "New protein steals the show as 'co-stimulator' of T cells" (news; comment). Science 262:844, (1993)).

In particular, the role of the human B7 antigens, i.e., human B7.1 and B7.2, has been reported to play a co-stimulatory role in T-cell activation.

#### 1. B7.1 and B7.2 Co-stimulatory Role in T Cell Activation

The elaboration of a successful immune response depends on a series of specific interactions between a T cell and an antigen presenting cell. Although the essential first step in this process depends upon the binding of antigen to the T cell receptor, in the context of the MHC class II molecule (Lane, P.J.L., F.M. McConnell, G.L. Schieven, E.A. Clark, and J.A. Ledbetter, (1990), "The Role of Class II Molecules in Human B Cell Activation." The Journal of Immunology, 144:3684-3692), this interaction alone is not sufficient to induce all the events necessary for a sustained

-5-

- response to a given antigen (Schwartz, R.H. (1990), "A Cell Culture Model for T Lymphocyte Clonal Anergy." Science, 248:1349; Jenkins, M.K. (1992). "The Role of Cell Division in the Induction of Clonal Anergy."
- 5 Immunology Today, 13:69; Azuma, M., M. Catabyab, D. Buck, J.H. Phillips, and L.L. Lanier, (1992). "Involvement of CD28 in MHC-unrestricted Cytotoxicity Mediated by a Human Natural Killer Leukemia Cell Line." The Journal of Immunology, 149:1556-1561; Azuma, M., M.
- 10 Catabyab, D. Buck, J.H. Phillips, and L.L. Lanier, (1992). "CD28 Interaction with B7 Costimulates Primary Allogeneic Proliferative Responses and Cytotoxicity Mediated by Small Resting T Lymphocytes." J. Exp. Med., 175:353-360).
- 15       The involvement of certain other co-stimulatory molecules is necessary (Norton, S.D., L. Zuckerman, K.B. Urdahl, R. Shefner, J. Miller, and M.K. Jenkins. (1992), "The CD28 Ligand, B7, Enhances IL-2 Production by Providing A Costimulatory Signal to T Cells." The
- 20 Journal of Immunology, 149:1556-1561). "The homodimers CD28 and CTLA-4 expressed on T cells" (June, C.H., J.A. Ledbetter, P.S. Linsley, and C.B. Thompson, (1990), "Role of the CD28 Receptor in T-Cell Activation." Immunology Today, 11:211-216; Linsley, P.S., W. Brady,
- 25 M. Urnes, L.S. Grosmaire, N.K. Damle, and J.A. Ledbetter, (1991), "CTLA-4 is a Second Receptor for the B Cell Activation Antigen B7." J. Exp. Med., 174:561), together with B7.1 (CD80) and B7.2 (CD86) expressed on antigen presenting cells, are major pairs of co-
- 30 stimulatory molecules necessary for a sustained immune response (Azuma, M., H. Yssel, J.H. Phillips, H. Spits, and L.L. Lanier, (1993), "Functional Expression of B7/BB1 on Activated T Lymphocytes." J. Exp. Med., 177:845-850; Freeman, G.J., A.S. Freedman, J.M. Segil,

-6-

- G. Lee, J.F. Whitman, and L.M. Nadler, (1989), "B7, A New Member of the Ig Superfamily with Unique Expression on Activated and Neoplastic B Cells." The Journal of Immunology, 143:2714-2722; Hathcock, K.S., G. Laslo, H.B. Dickler, J. Bradshaw, P. Linsley, and R.J. Hodes, (1993), "Identification of an Alternative CTLA-4 Ligand Costimulatory for T Cell Activation." Science, 262:905-911; Hart, D.N.J., G.C. Starling, V.L. Calder, and N.S. Fernando, (1993). "B7/BB-1 is a Leucocyte Differentiation Antigen on Human Dendritic Cells Induced by Activation." Immunology, 79:616-620). It can be shown *in vitro* that the absence of these co-stimulatory signals leads to an aborted T cell activation pathway and the development of unresponsiveness to the specific antigen, or anergy. (See, e.g., Harding, F.A., J.G. McArthur, J.A. Gross, D.M. Raulet, and J.P. Allison, (1992). "CD28 Mediated Signalling Co-stimulates Murine T Cells and Prevents Induction of Anergy in T Cell Clones." Nature, 356:607-609; Gimmi, C.D., G.J. Freeman, J.G. Gribben, G. Gray, and L.M. Nadler, (1993). "Human T-Cell Clonal Anergy is Induced by Antigen Presentation in the Absence of B7 Costimulation." Proc. Natl. Acad. Sci., 90:6586-6590; Tan, P., C. Anasetti, J.A. Hansen, J. Melrose, M. Brunvand, J. Bradshaw, J.A. Ledbetter, and P.S. Linsley, (1993), "Induction of Alloantigen-specific Hyporesponsiveness in Human T Lymphocytes by Blocking Interaction of CD28 with Its Natural Ligand B7/BB1." J. Exp. Med., 177:165-173). Achievement of *in vivo* tolerance constitutes a mechanism for immunosuppression and a viable therapy for organ transplant rejection and for the treatment of autoimmune diseases. This has been achieved in experimental models following the administration of CTLA4-Ig (Lenschow, D.J., Y. Zeng, R.J. Thistlethwaite, A. Montag, W. Brady,



-7-

M.G. Gibson, P.S. Linsley, and J.A. Bluestone, (1992), "Long-Term Survival of Xenogeneic Pancreatic Islet Grafts Induced by CTLA-4Ig." Science, 257:789-795).

The molecules B7.1 and B7.2 can bind to either CD28  
5 or CTLA-4, although B7.1 binds to CD28 with a Kd of 200  
Nm and to CTLA-4 with a 20-fold higher affinity  
(Linsley, P.S., E.A. Clark, and J.A. Ledbetter, (1990),  
"T-Cell Antigen CD28 Mediates Adhesion with B Cells by  
Interacting with Activation Antigen B7/BB-1." Proc.  
10 Natl. Acad. Sci., 87:5031-5035; Linsley et al, (1993),  
"The Role of the CD28 receptor during T cell responses  
to antigen," Annu. Rev. Immunol., 11:191-192; Linesley  
et al, (1993), "CD28 Engagement by B7/BB-1 Induces  
Transient Down-Regulation of CD28 Synthesis and  
15 Prolonged Unresponsiveness to CD28 Signaling," The  
Journal of Immunology, 150:3151-3169). B7.2 is  
expressed on activated B cells and interferon induced  
monocytes, but not resting B cells (Freeman, G.J., G.S.  
Gray, C.D. Gimmi, D.B. Lomarrd, L-J. Zhou, M. White,  
20 J.D. Fingerioth, J.G. Gribben, and LM. Nadler, (1991).  
"Structure, Expression and T Cell Costimulatory Activity  
of the Murine Homologue of the Human B Lymphocyte  
Activation Antigen B7," J. Exp. Med., 174:625-631).  
B7.2, on the other hand, is constitutively expressed at  
25 very low levels on resting monocytes, dendritic cells  
and B cells, and its expression is enhanced on activated  
T cells, NK cells and B lymphocytes (Azuma, M. D. Ito,  
H. Yagita, K. Okumura, J.H. Phillips, L.L. Lanier, and  
C. Somoza, "1993", "B70 Antigen is a Second Ligand for  
30 CTLA-4 and CD28," Nature, 366:76-79). Although B7.1  
and B7.2 can be expressed on the same cell type, their  
expression on B cells occurs with different kinetics  
(Lenschow, D.J., G.H. Su, L.A. Zuckerman, N. Nabavi,  
C.L. Jellis, G.S. Gray, J. Miller, and J.A. Bluestone,

-8-

(1993), "Expression and Functional Significance of an Additional Ligand for CTLA-4," Proc. Natl. Acad. Sci., USA, 90:11054-11058; Boussiotis, V.A., G.J. Freeman, J.G. Gribben, J. Daley, G. Gray, and L.M. Nadler, (1993), "Activated Human B Lymphocytes Express Three CTLA-4 Counter-receptors that Co-stimulate T-Cell Activation." Proc. Natl. Acad. Sci., USA, 90:11059-11063). Further analysis at the RNA level has demonstrated that B7.2 mRNA is constitutively expressed, whereas B7.1 MRNA is detected 4 hours after activation and initial low levels of B7.1 protein are not detectable until 24 hours after stimulation (Boussiotis, V.A., G.J. Freeman, J.G. Gribben, J. Daley, G. Gray, and L.M. Nadler, (1993), "Activated Human B Lymphocytes Express Three CTLA-4 Counter-receptors that Co-stimulate T-Cell Activation," Proc. Natl. Acad. Sci., USA, 90:11059-11063). CTLA-4/CD28 counter receptors, therefore, may be expressed at various times after B Cell activation.

The differential temporal expression of B7.1 and B7.2 suggests that the interaction of these two molecules with CTLA-4 and/or CD28 deliver distinct but related signals to the T cell (LaSalle, J.M., P.J. Tolentino, G.J. Freeman, L.M. Nadler, and D.A. Hafler, (1992), "CD28 and T Cell Antigen Receptor Signal Transduction Coordinately Regulate Intedeukin 2 Gene Expression In Response to Superantigen Stimulation," J. Exp. Med., 176:177-186; Vandenberghe, P., G.J. Freeman, L.M. Nadler, M.C. Fletcher, M. Kamoun, L.A. Turka, J.A. Ledbetter, C.B. Thompson, and C.H. June, (1992), "Antibody and B7/BB1-mediated Ligation of the CD28 Receptor Induces Tyrosine Phosphorylation in Human T Cells," The Journal of Experimental Medicine, 175:951-960). The exact signaling functions of CTLA-4

-9-

and CD28 on the T cell are currently unknown (Janeway, C.A., Jr. and K. Bottomly, (1994), "Signals and Signs for Lymphocyte Responses," Cell, 76:275-285). However, it is possible that one set of receptors could provide the initial stimulus for T cell activation and the second, a sustained signal to allow further elaboration of the pathway and clonal expansion to take place (Linsley, P.S., J.L. Greene, P. Tan, J. Bradshaw, J.A. Ledbetter, C. Anasetti, and N.K. Damle, (1992), "Coexpression and Functional Cooperation of CTLA-4 and CD28 on Activated T Lymphocytes," J. Exp. Med., 176:1595-1604). The current data supports the two-signal hypothesis proposed by Jenkins and Schwartz (Schwartz, R.H., (1990), "A Cell Culture Model for T Lymphocyte Clonal Anergy," Science, 248:1349; Jenkins, M.K., (1992), "The Role of Cell Division in the Induction of Clonal Anergy," Immunology Today, 13:69) that both a TCR and co-stimulatory signal are necessary for T cell expansion, lymphokine secretion and the full development of effector function (Greenan, V. and G. Kroemer, (1993), "Multiple Ways to Cellular Immune Tolerance," Immunology Today, 14:573). The failure to deliver the second signal results in the inability of T cells to secrete IL-2 and renders the cell unresponsive to antigen.

Structurally, both B7.1 and B7.2 contain extracellular immunoglobulin superfamily V and C-like domains, a hydrophobic transmembrane region and a cytoplasmic tail (Freeman, G.J., J.G. Gribben, V.A. Boussiotis, J.W. Ng, V. Restivo, Jr., L.A. Lombard, G.S. Gray, and L.M. Nadler, (1993), "Cloning of B7-2: A CTLA-4 Counter-receptor that Co-stimulates Human T Cell Proliferation," Science, 262:909). Both B7.1 and B7.2 are heavily glycosylated. B7.1 is a 44-54kD

-10-

glycoprotein comprised of a 223 amino acid extracellular domain, a 23 amino acid transmembrane domain, and a 61 amino acid cytoplasmic tail. B7.1 contains 3 potential protein kinase phosphorylation sites. (Azuma, M., H. Yssel, J.H. Phillips, H. Spits, and L.L. Lanier, (1993), "Functional Expression of B7/BB1 on Activated T Lymphocytes," J. Exp. Med., 177:845-850). B7.2 is a 306 amino acid membrane glycoprotein. It consists of a 220 amino acid extracellular region, a 23 amino acid hydrophobic transmembrane domain and a 60 amino acid cytoplasmic tail (Freeman, G.J., A.S. Freedman, J.M. Segil, G. Lee, J.F. Whitman, and L.M. Nadler, (1989), "B7, A New Member of the Ig Superfamily with Unique Expression on Activated and Neoplastic B Cells," The Journal of Immunology, 143:2714-2722). Although both B7.1 and B7.2 genes are localized in the same chromosomal region (Freeman, G.J., D.B. Lombard, C.D. Gimmi, S.A. Brod, L. Lee, J.C. Laning, D.A. Hafler, M.E. Dorf, G.S. Gray, H. Reiser, C.H. June, C.B. Thompson, and L.M. Nadler, (1992), "CTLA-4 and CD28 mRNA are Coexpressed in Most T Cells After Activation," The Journal of Immunology, 149:3795-3801; Schwartz, R.H., (1992), "Costimulation of T Lymphocytes: The Role of CD28, CTLA-4, and B7/BB1" in Selvakumar, A., B.K. Mohanraj, R.L. Eddy, T.B. Shows, P.C. White, C. Perrin, and B. Dupont, (1992), "Genomic Organization and Chromosomal Location of the Human Gene Encoding the B-Lymphocyte Activation Antigen B7," Immunogenetics, 36:175-181), these antigens do not share a high level of homology. The overall homology between B7.1 and B7.2 is 26% and between murine B7.1 and human S7 is 27% (Azuma, M., H. Yssel, J.H. Phillips, H. Spits, and L.L. Lanier, (1993), "Functional Expression of B7/BB1 on Activated T Lymphocytes," J. Exp. Med., 177:845-850; Freeman, G.J.,

-11-

A.S. Freedman, J.M. Segil, G. Lee, J.F. Whitman, and L.M. Nadler, (1989), "B7, A New Member of the Ig Superfamily with Unique Expression on Activated and Neoplastic B Cells," The Journal of Immunology, 143:2714-2722).

5 Although alignment of human B7.1 human B7.2 and murine B.1 sequences shows few stretches of lengthy homology, it is known that all three molecules bind to human CTLA-4 and CD28. Thus, there is most likely a common, or  
10 closely homologous region shared by the three molecules that may be either contiguous or conformational. This region may constitute the binding site of the B7.1 and B7.2 molecules to their counter-receptors. Antibodies raised against these epitopes could potentially inhibit the interaction of B7 with its counter-receptor on the T  
15 cell. Furthermore, antibodies that cross-reacted with this region on both B7.1 and B7.2 molecules would potentially have practical advantages over antibodies directed against B7.1 or B7.2 separately.

## 2. Blockade of the B7/CD28 Interaction

20 Blocking of the B7/CD28 interaction offers the possibility of inducing specific immunosuppression, with potential for generating long lasting antigen-specific therapeutic effects. Antibodies to either B7.1 or B7.2 have been shown to block T cell activation, as measured  
25 by the inhibition of IL-2 production *in vitro* (DeBoer, M., P. Parren, J. Dove, F. Ossendorp, G. van der Horst, and J. Reeder, (1992), "Functional Characterization of a Novel Anti-B7 Monoclonal Antibody," Eur. Journal of Immunology, 22:3071-3075; Azuma, M., H. Yssel, J.H.  
30 Phillips, H. Spits, and L.L. Lanier, (1993), "Functional Expression of B7/BB1 on Activated T Lymphocytes," J. Exp. Med., 177:845-850). However, different antibodies have been shown to vary in their immunosuppressive

-12-

potency, which may reflect either their affinity or epitope specificity. CTLA-4/Ig fusion protein and anti-CD28 Fabs were shown to have similar effects on the down regulation of IL-2 production.

5        *In vivo* administration of a soluble CTLA-4/Ig fusion protein has been shown to suppress T cell - dependent antibody responses in mice (Linsley, P.S., J.L. Greene, P. Tan, J. Bradshaw, J.A. Ledbetter, C. Anasetti, and N.K. Damle, (1992), "Coexpression and  
10    Functional Cooperation of CTLA-4 and CD28 on Activated T Lymphocytes," J. Exp. Med., 176:1595-1604; Lin, H., S.F. Builing, P.S. Linsley, R.O. Wei, C.D. Thompson, and L.A. Turka, (1993), "Long-term Acceptance of Major  
15    Histocompatibility Complex Mismatched Cardiac Allografts Induced by CTLA-4-Ig Plus Donor Specific Transfusion," J. Exp. Med., 178:1801) and, furthermore, larger doses were also able to suppress responses to a second  
20    immunization, demonstrating the feasibility of this approach for the treatment of antibody mediated  
25    autoimmune disease. In addition, CTLA-4/Ig was able to prevent pancreatic islet cell rejection in mice by directly inhibiting the interaction of T cells and B7.1/B7.2 antigen presenting cells (Lenschow, D.J., G.H. Su, L.A. Zuckerman, N. Nabavi, C.L. Jellis, G.S. Gray, J. Miller, and J.A. Bluestone, (1993), "Expression and Functional Significance of an Additional Ligand for CTLA-4," Proc. Natl. Acad. Sci., USA, 90:11054-11058). In this case, long term donor specific tolerance was achieved.

-13-

### 3. Recombinant Phage Display Technology for Antibody Selection

To date, no monoclonal antibodies which crossreact with both B7.1 and B7.2 have been reported. As noted, such antibodies would potentially be highly desirable as immunosuppressants. Phage display technology is beginning to replace traditional methods for isolating antibodies generated during the immune response, because a much greater percentage of the immune repertoire can be assessed than is possible using traditional methods. This is in part due to PEG fusion inefficiency, chromosomal instability, and the large amount of tissue culture and screening associated with heterohybridoma production. Phage display technology, by contrast, relies on molecular techniques for potentially capturing the entire repertoire of immunoglobulin genes associated with the response to a given antigen.

This technique is described by Barber et al, Proc. Natl. Acad. Sci., USA, 88, 7978-7982, (1991). Essentially, immunoglobulin heavy chain genes are PCR amplified and cloned into a vector containing the gene encoding the minor coat protein of the filamentous phage M13 in such a way that a heavy chain fusion protein is created. The heavy chain fusion protein is incorporated into the M13 phage particle together with the light chain genes as it assembles. Each recombinant phage contains, within its genome, the genes for a different antibody Fab molecule which it displays on its surface. Within these libraries, in excess of  $10^6$  different antibodies can be cloned and displayed. The phage library is panned on antigen coated microliter wells, non-specific phage are washed off, and antigen binding phage are eluted. The genome from the antigen-specific

-14-

clones is isolated and the gene III is excised, so that antibody can be expressed in soluble Fab form for further characterization. Once a single Fab is selected as a potential therapeutic candidate, it may easily be converted to a whole antibody. A previously described expression system for converting Fab sequences to whole antibodies is IDEC's mammalian expression vector NEOSPLA. This vector contains either human gamma 1 or gamma 4 constant region genes. CHO cells are transfected with the NEOSPLA vectors and after amplification this vector system has been reported to provide very high expression levels (> 30 pg/cell/day) can be achieved.

#### 4. Primatized Antibodies

Another highly efficient means for generating recombinant antibodies is disclosed by Newman, (1992), Biotechnology, 10, 1455-1460. More particularly, this technique results in the generation of primatized antibodies which contain monkey variable domains and human constant sequences. This reference is incorporated by reference in its entirety herein. Moreover, this technique is also described in commonly assigned U.S. Application No. 08/379,072, filed on January 25, 1995, which is a continuation of U.S. Serial No. 07/912,292, filed July 10, 1992, which is a continuation-in-part of U.S. Serial No. 07/856,281, filed March 23, 1992, which is finally a continuation-in-part of U.S. Serial No. 07/735,064, filed July 25, 1991. 08/379,072 and the parent application thereof are incorporated by reference in their entirety herein.

This technique modifies antibodies such that they are not antigenically rejected upon administration in humans. This technique relies on immunization of



-15-

cynomolgus monkeys with human antigens or receptors. This technique was developed to create high affinity monoclonal antibodies directed to human cell surface antigens.

5       Antibodies generated in this manner have previously been reported to display human effector function, have reduced immunogenicity, and long serum half-life. The technology relies on the fact that despite the fact that  
10       cynomolgus monkeys are phylogenetically similar to humans, they still recognize many human proteins as foreign and therefore mount an immune response. Moreover, because the cynomolgus monkeys are phylogenetically close to humans, the antibodies generated in these monkeys have been discovered to have  
15       a high degree of amino acid homology to those produced in humans. Indeed, after sequencing macaque immunoglobulin light and heavy chain variable region genes, it was found that the sequence of each gene family was 85-98% homologous to its human counterpart  
20       (Newman et al, (1992), Id.). The first antibody generated in this way, an anti-CD4 antibody, was 91-92% homologous to the consensus sequence of human immunoglobulin framework regions. Newman et al, Biotechnology, 10:1458-1460, (1992).

25       Monoclonal antibodies specific to the human B7 antigen have been previously described in the literature. For example, Weyl et al, Hum. Immunol., 31(4), 271-276, (1991) describe epitope mapping of human monoclonal antibodies against HLA-B-27 using natural and  
30       mutated antigenic variants. Also, Toubert et al, Clin. Exp. Immunol., 82(1), 16-20, (1990) describe epitope mapping of an HLA-B27 monoclonal antibody that also reacts with a 35-KD bacterial outer membrane protein. Also, Valle et al, Immunol., 69(4), 531-535, (1990)

-16-

describe a monoclonal antibody of the IgG1 subclass which recognizes the B7 antigen expressed in activated B cells and HTLV-1-transformed T cells. Further, Toubert et al, J. Immunol., 141(7), 2503-9, (1988) describe epitope mapping of HLA-B27 and HLA-B7 antigens using intradomain recombinants constructed by making hybrid genes between these two alleles in *E. coli*.

High expression of B7 antigen has been correlated to autoimmune diseases by some researchers. For example, Ionesco-Tirgoviste et al, Med. Interre, 24(1), 11-17, (1986) report increased B7 antigen expression in type 1 insulin-dependent diabetes. Also, the involvement of B7 antigen expression on dermal dendritic cells obtained from psoriasis patients has been reported. (Nestle et al, J. Clin. Invest., 94(1), 202-209, (1994)).

Further, the inhibition of anti-HLA-B7 alloreactive CTL using affinity-purified soluble HLA-B7 has been reported in the literature. (Zavazava et al, Transplantation, 51(4), 838-42, (1991)). Further, the use of B7 receptor soluble ligand, CTLA-4-Ig to block B7 activity (See, e.g., Lenschow et al, Science, 257, 789, 7955 (1992)) in animal models and a B7-1-Ig fusion protein capable of inhibiting B7 has been reported.

#### SUMMARY AND OBJECTS OF THE INVENTION

An object of the invention is to produce and identify novel macaque antibodies to human B7 antigen, more specifically to human B7.1 antigen and/or human B7.2 antigen.

More specifically, it is an object of the present invention to produce and identify novel macaque antibodies to human B7 antigen, i.e., human B7.1 and human B7.2 antigen by screening of phage display

-17-

libraries and/or monkey heterohybridomas using B lymphocytes obtained from human B7 antigen, i.e., human B7.1 or B7.2 antigen immunized monkeys.

5 It is another specific object of the invention to provide anti-B7 monkey monoclonal antibodies and primatized forms thereof which specifically bind human B7.1 and/or B7.2 antigen which inhibit the B7/CD86 pathway and B7 stimulation of activated T cells, thereby inhibiting IL-2 production and T cell proliferation and  
10 functioning as effective immunosuppressants.

It is another object of the invention to provide anti-human B7.1 and anti-human B7.2 monkey monoclonal antibodies and primatized forms thereof which inhibit antigen driven responses in donor spleen cell cultures,  
15 e.g., antigen specific IgG responses, IL-2 production and cell proliferation.

It is another specific object of the invention to identify particular monkey monoclonal antibodies specific to human B7.1 and human B7.2 antigen and  
20 primatized forms thereof having advantageous properties, i.e., affinity, immunosuppressive activity, which are useful as therapeutics. More specifically, these monkey antibodies and primatized forms thereof are to be used, e.g., as immunosuppressants, i.e., to block antigen  
25 driven immune responses, to treat autoimmune diseases such as psoriasis, rheumatoid arthritis, systemic erythematosus (SLE), type 1 diabetes mellitus, idiopathic thrombocytopenia purpura (ITP), and to prevent organ rejection.

30 It is another object of the invention to provide pharmaceutical compositions containing one or more monkey monoclonal antibodies specific to human B7 antigen, i.e., human B7.1 and/or human B7.2 antigen, or primatized forms thereof, and a pharmaceutically

-18-

acceptable carrier or excipient. These compositions will be used, e.g., as immunosuppressants to treat autoimmune diseases, e.g., idiopathic thrombocytopenia purpura (ITP) and systemic lupus erythematosus (SLE), to  
5 block antigen driven immune responses, and to prevent organ rejection in transplant recipients.

It is another object of the invention to provide novel methods of therapy by administration of therapeutically effective amounts of one or more monkey  
10 or primatized monoclonal antibodies which specifically bind to B7 antigen, i.e., human B7.1 and/or B7.2 antigens. Such therapeutic methods are useful for treatment of diseases treatable by inhibition of the B7:CD28 pathway e.g., autoimmune diseases such as  
15 idiopathic thrombocytopenia purpura (ITP), systemic lupus erythematosus (SLE), type 1 diabetes mellitus, psoriasis, rheumatoid arthritis, multiple sclerosis, aplastic anemia, as well as for preventing rejection in transplantation subjects.

20 It is still another object of the invention to provide transfectants, e.g., CHO cells, which express at least the variable heavy and light domains of monkey monoclonal antibodies specific to the human B7.1 and/or B7.2 antigen.

25 It is another object of the invention to provide nucleic acid sequences which encode the variable heavy and/or light domains of monkey monoclonal antibodies specific to human B7.1 and/or human B7.2 antigen, and expression vectors which provide for the expression of  
30 primatized antibodies containing these nucleic acid sequences.

-19-

### Definitions

The following terms are defined so that the invention may be more clearly understood.

5 Depleting antibody - an antibody which kills activated B cells or other antigen presenting cells.

Non-depleting antibody - an antibody which blocks the co-stimulatory action of B7 and T cell activating ligands CD28 and CTLA-4. Thus, it anergizes but does not eliminate the antigen presenting cell.

10 Primatized antibody - a recombinant antibody which has been engineered to contain the variable heavy and light domains of a monkey antibody, in particular, a cynomolgus monkey antibody, and which contains human constant domain sequences, preferably the human  
15 immunoglobulin gamma 1 or gamma 4 constant domain (or PE variant). The preparation of such antibodies is described in Newman et al, (1992), "Primatization of Recombinant Antibodies for Immunotherapy of Human Diseases: A Macaque/Human Chimeric Antibody Against  
20 Human CDH, Biotechnology, 10:1458-1460; also in commonly assigned 08/379,072 both of which are incorporated by reference in their entirety herein. These antibodies have been reported to exhibit a high degree of homology to human antibodies, i.e., 85-98%, display human  
25 effector functions, have reduced immunogenicity, and may exhibit high affinity to human antigens.

B7 antigens - B7 antigens in this application include, e.g., human B7, B7.1 and B7.2 antigens. These antigens bind to CD28 and/or CTLA-4. These antigens have a co-  
30 stimulatory role in T cell activation. Also, these B7 antigens all contain extracellular immunoglobulin superfamily V and C-like domains, a hydrophobic transmembrane region and a cytoplasmic tail. (See,

-20-

Freeman et al, Science, 262:909, (1993)), and are heavily glycosylated.

Anti-B7 antibodies - Antibodies, preferably monkey monoclonal antibodies or primatized forms thereof, which specifically bind human B7 antigens, e.g., human B7.1 and/or B7.2 antigen with a sufficient affinity to block the B7:CD28 interaction and thereby induce immunosuppression.

#### BRIEF DESCRIPTION OF THE FIGURES

10        Figure 1 depicts the pMS vector used to screen recombinant immunoglobulin libraries produced against B7 displayed on the surface of filamentous phage which contains primers based on macaque immunoglobulin sequences.

15        Figure 2 depicts the NEOSPLA expression vector used to express the subject primatized antibodies specific to human B7.1 antigen.

20        Figure 3 depicts monkey serum anti-B7.1 titers directed against cell surface B7.1 on transfected CHO cells.

      Figure 4 depicts inhibition of radiolabeled sB7.1 binding by SB7.1 affinity-purified monkey antibodies in the presence of unlabeled SB7 and Mab L307.4 murine anti-B7.1.

25        Figure 5 depicts inhibition of binding of radiolabeled monkey 135 and L3707.4 anti-B7.1 antibodies to B7 positive human SB cells by competition with affinity-purified SB7.1.

30        Figure 6 depicts inhibition of radiolabeled B7-Ig binding to activated human peripheral blood T cells by competing with unlabeled SB7.1 murine anti-B7.1 (L307.4) and monkey 1127 affinity purified serum antibodies.

-21-

Figure 7 depicts inhibition of IL-2 protein in mixed lymphocyte cultures by anti-B7.1 affinity-purified monkey serum antibodies.

5 Figure 8a depicts the amino acid and nucleic acid sequence of a primatized form of the light chain of 7C10.

Figure 8b depicts the amino acid and nucleic acid sequence of a primatized form of the heavy chain of 7C10.

10 Figure 9a depicts the amino acid and nucleic acid sequence of a primatized form of the light chain of 7B6.

Figure 9b depicts the amino acid and nucleic acid sequence of a primatized form of the heavy chain of 7B6.

15 Figure 10a depicts the amino acid and nucleic acid sequence of a primatized light chain 16C10.

Figure 10b depicts the amino acid and nucleic acid sequence of a primatized heavy chain 16C10.

#### DETAILED DESCRIPTION OF THE INVENTION

20 As described above, the present invention relates to the manufacture of novel monkey monoclonal antibodies which specifically bind human B7.1 and/or human B7.2 antigen, as well as primatized antibodies derived therefrom. These antibodies possess high affinity to human B7.1 and/or B7.2 and therefore may be used as  
25 immunosuppressants which inhibit the B7:CD86 pathway.

Preparation of monkey monoclonal antibodies will preferably be effected by screening of phage display libraries or by preparation of monkey heterohybridomas using B lymphocytes obtained from B7 (e.g., human B7.1  
30 and/or B7.2) immunized monkeys.

As noted, the first method for generating anti-B7 antibodies involves recombinant phage display

-22-

technology. This technique is generally described *supra*.

Essentially, this will comprise synthesis of recombinant immunoglobulin libraries against B7 antigen displayed on the surface of filamentous phage and selection of phage which secrete antibodies having high affinity to B7.1 and/or B7.2 antigen. As noted *supra*, preferably antibodies will be selected which bind to both human B7.1 and B7.2. To effect such methodology, the present inventors have created a unique library for monkey libraries which reduces the possibility of recombination and improves stability. This vector, pMS, is described in detail *infra*, and is shown in Figure 1.

Essentially, to adopt phage display for use with macaque libraries, this vector contains specific primers for PCR amplifying monkey immunoglobulin genes. These primers are based on macaque sequences obtained while developing the primatized technology and databases containing human sequences.

Suitable primers are disclosed in commonly assigned 08/379,072 incorporated by reference herein.

The second method involves the immunization of monkeys, i.e., macaques, against human B7 antigen, preferably against human B7.1 and B7.2 antigen. The inherent advantage of macaques for generation of monoclonal antibodies is discussed *supra*. In particular, such monkeys, i.e., cynomolgus monkeys, may be immunized against human antigens or receptors. Moreover, the resultant antibodies may be used to make primatized antibodies according to the methodology of Newman et al, Biotechnology, 10, 1455-1460, (1992), and Newman et al, commonly assigned U.S. Serial No. 08/379,072, filed January 25, 1995, which are incorporated by reference in their entirety.



-23-

The significant advantage of antibodies obtained from cynomolgus monkeys is that these monkeys recognize many human proteins as foreign and thereby provide for the formation of antibodies, some with high affinity to  
5 desired human antigens, e.g., human surface proteins and cell receptors. Moreover, because they are phylogenetically close to humans, the resultant antibodies exhibit a high degree of amino acid homology to those produced in humans. As noted above, after sequencing  
10 macaque immunoglobulin light and heavy variable region genes, it was found that the sequence of each gene family was 85-88% homologous to its human counterpart (Newman et al, (1992), *Id.*).

Essentially, cynomolgus macaque monkeys are  
15 administered human B7 antigen, e.g., human B7.1 and/or human B7.2 antigen, B cells are isolated therefrom, e.g., lymph node biopsies are taken from the animals, and B lymphocytes are then fused with KH6/B5 (mouse x human) heteromyeloma cells using polyethylene glycol  
20 (PEG). Heterohybridomas secreting antibodies which bind human B7 antigen, e.g., human B7.1 and/or human B7.2 antigen, are then identified.

Antibodies which bind to both B7.1 and B7.2 are desirable because such antibodies potentially may be  
25 used to inhibit the interaction of B7.1 and B7.2, as well as B7 with their counter-receptors, i.e., human CTLA-4 and CD28. Antibodies against these epitopes may inhibit the interaction of both human B7.1 and human B7.2 with their counter receptors on the T cell. This  
30 may potentially provide synergistic effects.

However, antibodies which bind to only one of human B7 antigen, B7.1 antigen or B7.2 antigen, are also highly desirable because of the co-involvement of these molecules in T cell activation, clonal expansion

-24-

lymphokine (IL-2) secretion, and responsiveness to antigen. Given that both human B7.1 and B7.2 bind to human CTLA-4 and CD28, it is probable that there is at least one common or homologous region (perhaps a shared  
5 conformational epitope or epitopes) to which macaque antibodies may potentially be raised.

The present inventors elected to immunize macaques against human B7.1 antigen using recombinant soluble B7.1 antigen produced in CHO cells and purified by  
10 affinity chromatography using a L307.4-sepharose affinity column. However, the particular source of human B7 antigen, human B7.1 antigen or human B7.2 antigen is not critical, provided that it is of sufficient purity to result in a specific antibody  
15 response to the particular administered B7 antigen and potentially to other B7 antigens.

The human B7 antigen, human B7.1 antigen (also called CD80) and human B7.2 antigen (also called CD86) genes have been cloned, and sequenced, and therefore may  
20 readily be manufactured by recombinant methods.

Preferably, the administered human B7 antigen, human B7.1 antigen and/or human B7.2 antigen will be administered in soluble form, e.g., by expression of a B7, B7.1 or B7.2 gene which has its transmembrane and  
25 cytoplasmic domains removed, thereby leaving only the extracellular portion, i.e., the extracellular superfamily V and C-like domains. (See, e.g., Grumet et al, Hum. Immunol., 40(3), p. 228-234, 1994, which teaches expression of a soluble form of human B7, which  
30 is incorporated by reference in its entirety herein).

The macaques will be immunized with the B7, B7.1 and/or B7.2 antigen, preferably a soluble form thereof, under conditions which result in the production of antibodies specific thereto. Preferably, the soluble

-25-

human B7, B7.1 or B7.2 antigen will be administered in combination with an adjuvant, e.g., Complete Freund's Adjuvant (CFA), Alum, Saponin, or other known adjuvants, as well as combinations thereof. In general, this will  
5 require repeated immunization, e.g., by repeated injection, over several months. For example, administration of soluble B7.1 antigen was effected in adjuvant, with booster immunizations, over a 3 to 4 month period, with resultant production of serum  
10 containing antibodies which bound human B7.1 antigen.

After immunization B cells are collected, e.g., by lymph node biopsies taken from the immunized animals and B lymphocytes fused with KH6/B5 (mouse x human) heteromyeloma cells using polyethylene glycol. Methods  
15 for preparation of such heteromyelomas are known and may be found in U.S. Serial No. 08/379,072 by Newman et al, filed on January 25, 1995 and incorporated by reference herein.

Heterohybridomas which secrete antibodies which  
20 bind human B7, B7.1 and/or B7.2 are then identified. This may be effected by known techniques. For example, this may be determined by ELISA or radioimmunoassay using enzyme or radionuclide labelled human B7, B7.1 and/or B7.2 antigen.

25 Cell lines which secrete antibodies having the desired specificity to human B7, B7.1 and/or B7.2 antigen are then subcloned to monoclonality.

In the present invention, the inventors screened purified antibodies for their ability to bind to soluble  
30 B7.1 antigen coated plates in an ELISA assay, antigen positive B cells, and CHO transfectomas which express human B7.1 antigen on their cell surface. In addition, the antibodies were screened for their ability to block B cell/T cell interactions as measured by IL-2

-26-

production and tritiated thymidine uptake in a mixed lymphocyte reaction (MLR), with B7 binding being detected using  $^{125}\text{I}$ -radiolabeled soluble B7.1 (SB7.1).

Also, affinity purified antibodies from macaques  
5 were tested for their reactivity against CHO  
transfectants which expressed B7.1/Ig fusion proteins,  
and against CHO cells which produced human B7.2 antigen.  
These results indicated that the B7.1 immune sera bound  
to the B7.2 transfectomas. Binding of antibodies to  
10 B7.2 antigen may be confirmed using soluble B7.2-Ig  
reagents. As discussed in the examples, this may be  
effected by producing and purifying B7.2-Ig from CHO  
transfectomas in sufficient quantities to prepare a  
B7.2-Ig-sepharose affinity column. Those antibodies  
15 which cross-react with B7.2 will bind the B7.2-Ig-  
sepharose column.

Cell lines which express antibodies which  
specifically bind to human B7 antigen, B7.1 antigen  
and/or B7.2 antigen are then used to clone variable  
20 domain sequences for the manufacture of primatized  
antibodies essentially as described in Newman et al,  
(1992), Id. and Newman et al, U.S. Serial No. 379,072,  
filed January 25, 1995, both of which are incorporated  
by reference herein. Essentially, this entails  
25 extraction of RNA therefrom, conversion to cDNA, and  
amplification thereof by PCR using Ig specific primers.  
Suitable primers are described in Newman et al, 1992,  
Id. and in U.S. Serial No. 379,072. (See, in  
particular, Figure 1 of U.S. Serial No. 379,072).

30 The cloned monkey variable genes are then inserted  
into an expression vector which contains human heavy and  
light chain constant region genes. Preferably, this is  
effected using a proprietary expression vector of IDEC,  
Inc., referred to as NEOSPLA. This vector is shown in

-27-

Figure 2 and contains the cytomegalovirus promoter/enhancer, the mouse beta globin major promoter, the SV40 origin of replication, the bovine growth hormone polyadenylation sequence, neomycin phosphotransferase exon 1 and exon 2, human immunoglobulin kappa or lambda constant region, the dihydrofolate reductase gene, the human immunoglobulin gamma 1 or gamma 4 PE constant region and leader sequence. This vector has been found to result in very high level expression of primatized antibodies upon incorporation of monkey variable region genes, transfection in CHO cells, followed by selection in G418 containing medium and methotrexate amplification.

For example, this expression system has been previously disclosed to result in primatized antibodies having high avidity ( $K_d \leq 10^{-10}$  M) against CD4 and other human cell surface receptors. Moreover, the antibodies have been found to exhibit the same affinity, specificity and functional activity as the original monkey antibody. This vector system is substantially disclosed in commonly assigned U.S. Serial No. 379,072, incorporated by reference herein as well as U.S. Serial No. 08/149,099, filed on November 3, 1993, also incorporated by reference in its entirety herein. This system provides for high expression levels, i.e., > 30 pg/cell/day.

As discussed *infra*, the subject inventors have selected four lead candidate monkey monoclonal antibodies which specifically bind the B7.1 antigen, and which may also bind the B7.2 antigen. These monkey monoclonal antibodies are referred to herein as 7B6, 16C10, 7C10 and 20C9.

As discussed in greater detail *infra*, these antibodies were evaluated for their ability to block B

-28-

cell/T cell interactions as measured by IL-2 production and tritiated thymidine uptake in a mixed lymphocyte reaction for T cell binding experiments for T cell binding, human body coat peripheral blood lymphocytes were cultured for 3-6 days in the presence of PHA stimulator. B7 binding was radioassayed using <sup>125</sup>I-radiolabeled soluble B7.1. The observed results indicate that all of these antibodies bind B7.1 antigen with high affinity and effectively block B cell/T cell interactions as evidenced by reduced IL-2 production and reduced proliferation of mixed lymphocyte cultures.

The properties of these particular monkey monoclonal antibodies are summarized below:

1. To demonstrate the monkey antibodies' ability to block the physical interaction between CTLA4-Ig, varying concentrations of the monkey anti-B7.1 antibodies and unlabeled CTLA4-IG were incubated with radiolabeled CTLA4-Ig<sup>125</sup>. The results of the inhibition assay showed that the IC50 (the concentration of inhibitor which results in 50% inhibition) for the monkey antibodies are:

a:	7C10:	0.39 µg/Ml
b:	16C10:	1.60 µg/Ml
c:	20C9:	3.90 µg/Ml
d:	7B6:	39.0 µg/Ml

2. Scatchard analysis showed that the apparent affinity constants (Kd) for the monkey antibodies binding to B7-Ig coated plates were approximated to be:

a:	7C10:	6.2 x 10 <sup>-9</sup> M
b:	16C10:	8.1 x 10 <sup>-9</sup> M
c:	7B6:	10.7 x 10 <sup>-9</sup> M

-29-

d: 20C9:  $16.8 \times 10^{-9}M$

3. The antibodies were tested in vitro in a mixed lymphocyte reaction assay (MLR). The MLR showed that all 4 anti-B7.1 antibodies inhibit IL-2 production to different extents as shown by the following Ibgo values:

a: 7B6: 5.0  $\mu g/M$   
b: 16C10:  $<0.1 \mu g/M$   
c: 20C9: 2.0  $\mu g/M$   
d: 7C10: 5.0  $\mu g/M$

4. The monkey anti-B7.1 antibodies were tested for their ability to bind B7 on human peripheral blood lymphocytes (PBL). FACS analysis showed that all 4 monkey antibodies tested positive.

5. Monkey antibodies 16C10, 7B6, 7C10 and 20C9 were tested for C1q binding by FACS analysis. Results showed 7C10 monkey Ig had strong human C1q binding after incubating with B7.1 CHO-transfected cells. 16C10 was positive, while 20C9 and 7B6 monkey antibodies were negative.

6. To select an animal model for path-tox studies, the monkey antibodies were tested with animal blood from different species. It was determined that the monkey anti-B7.1 antibodies cross-reacted with human, chimpanzee, and possibly baboon.

Based on these properties, it would appear that three monkey monoclonal antibodies possess the most advantageous properties, 16C10, 7C10 and 20C9, with 16C10 and 7C10 being somewhat better than 20C9.

-30-

Using the techniques described *supra*, and in commonly assigned U.S. Serial No. 08/379,072, the present inventors have cloned the variable domains of 7C10, 7B6 and 16C10, and provide the amino acid and  
5 nucleic acid sequences of primatized forms of the 7C10 light chain, 7C10 heavy chain, 7B6 light chain, 7B6 heavy chain, 16C10 light chain and 16C10 heavy chain. These amino acid and nucleic acid sequences may be found in Figures 8a and 8b, 9a and 9b, and 10a and 10b. The  
10 DNA and amino acid sequence for the human gamma 1 constant domain may be found in 08/379,072.

As discussed *supra*, these primatized antibodies are preferably expressed using the NEOSPLA expression vector shown in Figure 2 which is substantially described in  
15 commonly assigned 08/379,072 and 08/149,099, both of which applications are incorporated by reference herein.

As previously noted, the subject primatized antibodies will preferably contain either the human immunoglobulin gamma 1 or gamma 4 constant region, with  
20 gamma 4 preferably mutated at two positions to create gamma 4 PE. The gamma 4 PE mutant contains two mutations, a glutamic acid in the CH2 region introduced to eliminate residual FCR binding, and a proline substitution in the hinge region, intended to enhance  
25 the stability of the heavy chain disulfide bond interaction. (See, Alegre et al, J. Immunol., 148, 3461-3468, (1992); and Angel et al, Mol. Immunol., 30, 105-158, (1993), both of which are incorporated by reference herein).

30 Whether the subject primatized antibodies contain the gamma 1, gamma 4 or gamma 4 PE constant region largely depends on the particular disease target. Preferably, depleting and non-depleting primatized IgG1



-31-

and IgG4 antibodies are created and tested against specific disease targets.

Given the described binding and functional properties of the subject monkey monoclonal antibodies, these anti-B7.1 monoclonal antibodies and primatized forms thereof should be well suited as therapeutic agents for blocking the B7:CD28 interaction thereby providing for immunosuppression. In particular, given their high affinity to B7.1 antigen and ability to block B cell/T cell interactions as measured by IL-2 production and tritiated thymidine uptake in mixed lymphocyte culture as well as their ability to effectively inhibit antigen driven responses in donor spleen cell cultures as shown by reduced antigen specific IgG responses, IL-2 production and cell proliferation, these monkey monoclonal antibodies and primatized forms thereof should function as effective immunosuppressants which modulate the B7:CD28 pathway. This is significant for the treatment of many diseases wherein immunosuppression is therapeutically desirable, e.g., autoimmune diseases, to inhibit undesirable antigen specific IgG responses, and also for prevention of organ rejection and graft-versus-host disease. Essentially, the subject antibodies will be useful in treating any disease wherein suppression of the B7:CD28 pathway is therapeutically desirable.

Key therapeutic indications for the subject anti-B7.1 antibodies include, by way of example, autoimmune diseases such as idiopathic thrombocytopenia purpura (ITP), systemic lupus erythematosus (SLE), type 1 diabetes mellitus, multiple sclerosis, aplastic anemia, psoriasis and rheumatoid arthritis.

Another significant therapeutic indication of the subject anti-B7.1 antibodies is for prevention of graft-

-32-

versus-host-disease (GVHD) during organ transplant and bone marrow transplant (BMT). The subject antibodies may be used to induce host tolerance to donor-specific alloantigens and thereby facilitate engraftment and  
5 reduce the incidence of graft rejection. It has been shown in a murine model of allogeneic cardiac transplantation that intravenous administration of CTLA4-Ig can result in immunosuppression or even induction of tolerance to alloantigen. (Lin et al, J. Exp. Med. 178:1801, 1993; Torka et al, Proc. Natl. Acad. Sci., USA, 89:11102, 1992). It is expected that the  
10 subject primatized anti-B7.1 antibodies will exhibit similar or greater activity.

Antibodies produced in the manner described above,  
15 or by equivalent techniques, can be purified by a combination of affinity and size exclusion chromatography for characterization in functional biological assays. These assays include determination of specificity and binding affinity as well as effector  
20 function associated with the expressed isotype, e.g., ADCC, or complement fixation. Such antibodies may be used as passive or active therapeutic agents against a number of human diseases, including B cell lymphoma, infectious diseases including AIDS, autoimmune and  
25 inflammatory diseases, and transplantation. The antibodies can be used either in their native form, or as part of an antibody/chelate, antibody/drug or antibody/toxin complex. Additionally, whole antibodies or antibody fragments (Fab<sub>2</sub>, Fab, Fv) may be used as  
30 imaging reagents or as potential vaccines or immunogens in active immunotherapy for the generation of anti-idiotypic responses.

The amount of antibody useful to produce a therapeutic effect can be determined by standard

-33-

techniques well known to those of ordinary skill in the art. The antibodies will generally be provided by standard technique within a pharmaceutically acceptable buffer, and may be administered by any desired route.

5 Because of the efficacy of the presently claimed antibodies and their tolerance by humans it is possible to administer these antibodies repetitively in order to combat various diseases or disease states within a human.

10 The anti-B7.1 antibodies (or fragments thereof) of this invention are useful for inducing immunosuppression, i.e., inducing a suppression of a human's or animal's immune system. This invention therefore relates to a method of prophylactically or  
15 therapeutically inducing immunosuppression in a human or other animal in need thereof by administering an effective, non-toxic amount of such an antibody of this invention to such human or other animal.

The ability of the compounds of this invention to  
20 induce immunosuppression has been demonstrated in standard tests used for this purpose, for example, a mixed lymphocyte reaction test or a test measuring inhibition of T-cell proliferation measured by thymidine uptake.

25 The fact that the antibodies of this invention have utility in inducing immunosuppression indicates that they should be useful in the treatment or prevention of resistance to or rejection of transplanted organs or tissues (e.g., kidney, heart, lung, bone marrow, skin,  
30 cornea, etc.); the treatment or prevention of autoimmune, inflammatory, proliferative and hyperproliferative diseases, and of cutaneous manifestations of immunologically mediated diseases (e.g., rheumatoid arthritis, lupus erythematosus,

-34-

systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, type 1 diabetes, uveitis, nephrotic syndrome, psoriasis, atopic dermatitis, contact dermatitis and further eczematous dermatitides, seborrheic dermatitis, Lichen planus, Pemphigus, bullous pemphigus, Epidermolysis bullosa, urticaria, angioedemas, vasculitides, erythema, cutaneous eosinophilias, Alopecia areata, etc.); the treatment of reversible obstructive airways disease, intestinal inflammations and allergies (e.g., Coeliac disease, proctitis, eosinophilia gastroenteritis, mastocytosis, Crohn's disease and ulcerative colitis) and food-related allergies (e.g., migraine, rhinitis and eczema).

One skilled in the art would be able, by routine experimentation, to determine what an effective, non-toxic amount of antibody would be for the purpose of inducing immunosuppression. Generally, however, an effective dosage will be in the range of about 0.05 to 100 milligrams per kilogram body weight per day.

The antibodies (or fragments thereof) of this invention should also be useful for treating tumors in a mammal. More specifically, they should be useful for reducing tumor size, inhibiting tumor growth and/or prolonging the survival time of tumor-bearing animals. Accordingly, this invention also relates to a method of treating tumors in a human or other animal by administering to such human or animal an effective, non-toxic amount of an antibody. One skilled in the art would be able, by routine experimentation, to determine what an effective, non-toxic amount of anti-B7 antibody would be for the purpose of treating carcinogenic tumors. Generally, however, an effective dosage is

-35-

expected to be in the range of about 0.05 to 100 milligrams per kilogram body weight per day.

The antibodies of the invention may be administered to a human or other animal in accordance with the  
5   aforementioned methods of treatment in an amount sufficient to produce such effect to a therapeutic or prophylactic degree. Such antibodies of the invention can be administered to such human or other animal in a conventional dosage form prepared by combining the  
10   antibody of the invention with a conventional pharmaceutically acceptable carrier or diluent according to known techniques. It will be recognized by one of skill in the art that the form and character of the pharmaceutically acceptable carrier or diluent is  
15   dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables.

The route of administration of the antibody (or fragment thereof) of the invention may be oral,  
20   parenteral, by inhalation or topical. The term parenteral as used herein includes intravenous, intraperitoneal, intramuscular, subcutaneous, rectal or vaginal administration. The subcutaneous and intramuscular forms of parenteral administration are  
25   generally preferred.

The daily parenteral and oral dosage regimens for employing compounds of the invention to prophylactically or therapeutically induce immunosuppression, or to therapeutically treat carcinogenic tumors will generally  
30   be in the range of about 0.05 to 100, but preferably about 0.5 to 10, milligrams per kilogram body weight per day.

The antibodies of the invention may also be administered by inhalation. By "inhalation" is meant

-36-

intranasal and oral inhalation administration.

Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional techniques. The preferred dosage amount of a compound of the invention to be employed is generally within the range of about 10 to 100 milligrams.

The antibodies of the invention may also be administered topically. By topical administration is meant non-systemic administration and includes the application of an antibody (or fragment thereof) compound of the invention externally to the epidermis, to the buccal cavity and instillation of such an antibody into the ear, eye and nose, and where it does not significantly enter the blood stream. By systemic administration is meant oral, intravenous, intraperitoneal and intramuscular administration. The amount of an antibody required for therapeutic or prophylactic effect will, of course, vary with the antibody chosen, the nature and severity of the condition being treated and the animal undergoing treatment, and is ultimately at the discretion of the physician. A suitable topical dose of an antibody of the invention will generally be within the range of about 1 to 100 milligrams per kilogram body weight daily.

#### Formulations

While it is possible for an antibody or fragment thereof to be administered alone, it is preferable to present it as a pharmaceutical formulation. The active ingredient may include for topical administration, from 0.001% to 10% w/w, e.g., from 1% to 2% by weight of the formulation, although it may include as much as 10%



-37-

w/w but preferably not in excess of 5% w/w and more preferably from 0.1% to 1% w/w of the formulation.

The topical formulations of the present invention, include an active ingredient together with one or more acceptable carrier(s) therefor and optionally any other therapeutic ingredients(s). The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of where treatment is required, such as liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose.

Drops according to the present invention may include sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active ingredient in a suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining at 90°-100°C for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol and propylene glycol.



-38-

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared  
5 by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor  
10 oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely-divided  
15 or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy basis. The basis may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap;  
20 a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives, or a fatty acid such as stearic or oleic acid together with an alcohol such as propylene glycol or macrogols. The formulation may incorporate any  
25 suitable surface active agent such as an anionic, cationic or non-ionic surface active such as sorbitan esters or polyoxyethylene derivatives thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous  
30 silicas, and other ingredients such as lanolin, may also be included.

The subject anti-B7.1 antibodies or fragments thereof may also be administered in combination with other moieties which modulate the B7:CD28 pathway. Such



-39-

moieties include, by way of example, cytokines such as IL-7 and IL-10, CTLA4-Ig, soluble CTLA-4 and anti-CD28 antibodies and fragments thereof.

It will be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of an antibody or fragment thereof of the invention will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular animal being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of an antibody or fragment thereof of the invention given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following formulations are, therefore, to be construed as merely illustrative embodiments and not a limitation of the scope of the present invention in any way.

#### 25 Capsule Composition

A pharmaceutical composition of this invention in the form of a capsule is prepared by filling a standard two-piece hard gelatin capsule with 50 mg. of an antibody or fragment thereof of the invention, in powdered form, 100 mg. of lactose, 32 mg. of talc and 8 mg. of magnesium stearate.

-40-

Injectable Parenteral Composition

A pharmaceutical composition of this invention in a form suitable for administration by injection is prepared by stirring 1.5% by weight of an antibody or  
5 fragment thereof of the invention in 10% by volume propylene glycol and water. The solution is sterilized by filtration.

Ointment Composition

Antibody or fragment thereof of the invention  
10 1.0 g.

White soft paraffin to 100.0 g.

The antibody or fragment thereof of the invention is dispersed in a small volume of the vehicle to produce a smooth, homogeneous product. Collapsible metal tubes  
15 are then filled with the dispersion.

Topical Cream Composition

Antibody or fragment thereof of the invention  
1.0 g.

Polawax GP 200 20.0 g.

20 Lanolin Anhydrous 2.0 g.

White Beeswax 2.5 g.

Methyl hydroxybenzoate 0.1 g.

Distilled Water to 100.0 g.

The polawax, beeswax and lanolin are heated  
25 together at 60°C. A solution of methyl hydroxybenzoate is added and homogenization is achieved using high speed stirring. The temperature is then allowed to fall to 50°C. The antibody or fragment thereof of the invention is then added and dispersed throughout, and the  
30 composition is allowed to cool with slow speed stirring.

-41-

Topical Lotion Composition

Antibody or fragment thereof of the invention  
1.0 g.  
Sorbitan Monolaurate 0.6 g.  
5 Polysorbate 20 0.6 g.  
Cetostearyl Alcohol 1.2 g.  
Glycerin 6.0 g.  
Methyl Hydroxybenzoate 0.2 g.  
Purified Water B.P. to 100-00 ml. (B.P. = British  
10 Pharmacopeia)

The methyl hydroxybenzoate and glycerin are dissolved in 70 ml. of the water at 75°C. The sorbitan monolaurate, polysorbate 20 and cetostearyl alcohol are melted together at 75°C and added to the aqueous  
15 solution. The resulting emulsion is homogenized, allowed to cool with continuous stirring and the antibody or fragment thereof of the invention is added as a suspension in the remaining water. The whole suspension is stirred until homogenized.

20 Eye Drop Composition

Antibody or fragment thereof of the invention  
0.5 g.

Methyl Hydroxybenzoate 0.01 g.  
Propyl Hydroxybenzoate 0.04 g.  
25 Purified Water B.P. to 100-00 ml.

The methyl and propyl hydroxybenzoates are dissolved in 70 ml. purified water at 75°C and the resulting solution is allowed to cool. The antibody or fragment thereof of the invention is then added, and the  
30 solution is sterilized by filtration through a membrane filter (0.022  $\mu$ m pore size), and packed aseptically into suitable sterile containers.

-42-

Composition for Administration by Inhalation

For an aerosol container with a capacity of 15-20 ml: mix 10 mg. of an antibody or fragment thereof of the invention with 0.2-0.5% of a lubricating agent, such as polysorbate 85 or oleic acid, and disperse such mixture in a propellant, such as freon, preferably in a combination of (1,2 dichlorotetrafluoroethane) and difluorochloro-methane and put into an appropriate aerosol container adapted for either intranasal or oral inhalation administration.

Composition for Administration by Inhalation

For an aerosol container with a capacity of 15-20 ml: dissolve 10 mg. of an antibody or fragment thereof of the invention in ethanol (6-8 ml.), add 0.1-0.2% of a lubricating agent, such as polysorbate 85 or oleic acid; and disperse such in a propellant, such as freon, preferably in combination of (1,2 dichlorotetrafluoroethane) and difluorochloromethane, and put into an appropriate aerosol container adapted for either intranasal or oral inhalation administration.

The antibodies and pharmaceutical compositions of the invention are particularly useful for parenteral administration, i.e., subcutaneously, intramuscularly or intravenously. The compositions for parenteral administration will commonly comprise a solution of an antibody or fragment thereof of the invention or a cocktail thereof dissolved in an acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers may be employed, e.g., water, buffered water, 0.4% saline, 0.3% glycine, and the like. These solutions are sterile and generally free of particulate matter. These solutions may be sterilized by conventional, well-known sterilization techniques. The

-43-

compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as Ph adjusting and buffering agents, etc. The concentration of the  
5 antibody or fragment thereof of the invention in such pharmaceutical formulation can vary widely, i.e., from less than about 0.5%, usually at or at least about 1% to as much as 15 or 20% by weight, and will be selected primarily based on fluid volumes, viscosities, etc.,  
10 according to the particular mode of administration selected.

Thus, a pharmaceutical composition of the invention for intramuscular injection could be prepared to contain 1 Ml sterile buffered water, and 50 mg. of an antibody  
15 or fragment thereof of the invention. Similarly, a pharmaceutical composition of the invention for intravenous infusion could be made up to contain 250 ml. of sterile Ringer's solution, and 150 mg. of an antibody or fragment thereof of the invention. Actual methods  
20 for preparing parenterally administrable compositions are well known or will be apparent to those skilled in the art, and are described in more detail in, for example, Remington's Pharmaceutical Science, 15th ed., Mack Publishing Company, Easton, Pennsylvania, hereby  
25 incorporated by reference herein.

The antibodies (or fragments thereof) of the invention can be lyophilized for storage and reconstituted in a suitable carrier prior to use. This technique has been shown to be effective with  
30 conventional immune globulins and art-known lyophilization and reconstitution techniques can be employed.

Depending on the intended result, the pharmaceutical composition of the invention can be

-44-

administered for prophylactic and/or therapeutic treatments. In therapeutic application, compositions are administered to a patient already suffering from a disease, in an amount sufficient to cure or at least partially arrest the disease and its complications. In prophylactic applications, compositions containing the present antibodies or a cocktail thereof are administered to a patient not already in a disease state to enhance the patient's resistance.

Single or multiple administrations of the pharmaceutical compositions can be carried out with dose levels and pattern being selected by the treating physician. In any event, the pharmaceutical composition of the invention should provide a quantity of the altered antibodies (or fragments thereof) of the invention sufficient to effectively treat the patient.

It should also be noted that the antibodies of this invention may be used for the design and synthesis of either peptide or non-peptide compounds (mimetics) which would be useful in the same therapy as the antibody. See, e.g., Saragovi et al., Science, 253, 792-795 (1991).

To further illustrate the invention, the following examples are provided. These examples are not intended, nor are they to be construed, as further limiting the invention.

#### Example 1

Recombinant immunoglobulin libraries displayed on the surface of filamentous phage were first described by McCafferty et al, Nature, 348:552-554, 1990 and Barbas et al, Proc. Natl. Acad. Sci., USA 88:7978-7982, 1991. Using this technology, high affinity antibodies have been isolated from immune human recombinant libraries

-45-

(Barbas et al, Proc. Natl. Acad. Sci., USA 589:10164-10168, 1992). Although the phage display concept used is substantially similar to that described by Barbas, 1991, Id. the technique has been modified by the

5 substitution of a unique vector for monkey libraries to reduce the possibility of recombination and improve stability. This vector, pMS, Figure 1 contains a single lac promoter/operator for efficient transcription and translation of polycistronic heavy and light chain

10 monkey DNA. This vector contains two different leader sequences, the omp A (Movva et al, J. Biol. Chem., 255: 27-29, (1980), for the light chain and the pel B (Lei, J. Bact., 4379-109:4383 (1987) for the heavy chain Fd. Both leader sequences are translated into hydrophobic

15 signal peptides that direct the secretion of the heavy and light chain cloned products into the periplasmic space. In the oxidative environment of the periplasm, the two chains fold and disulfide bonds form to create stable Fab fragments. We derived the backbone of the

20 vector from the phagemid bluescript. (Stratagene, La Jolla, CA). It contains the gene for the enzyme beta-lactamase that confers ampicillin (carbenicillin) resistance to bacteria that harbor pMS DNA. We also derived, from bluescript, the origin of replication of

25 the multicopy plasmid ColE1 and the origin of replication of the filamentous bacteriophage f1. The origin of replication of phage f1 (the so-called intragenic region), signals the initiation of synthesis of single stranded pMS DNA, the initiation of capsid

30 formation and the termination of RNA synthesis by viral enzymes. The replication and assembly of pMS DNA strands into phage particles requires viral proteins that must be provided by a helper phage. We have used helper phage VCSM13 which is particularly suited for

-46-

this, since it also contains a gene coding for kanamycin resistance. Bacteria infected with VCSM13 and pMS can be selected by adding both kanamycin and carbenicillin to the growth medium. The bacteria will ultimately  
5 produce filamentous phage particles containing either pMS or VCSM13 genomes. Packaging of the helper phage is less efficient than that of pMS, resulting in a mixed phage population that contains predominately recombinant  
10 pMS phages. The ends of the phage pick up minor coat proteins specific to each end. Of particular interest here is the gene III product which is present in three to five copies at one end of the phage. The gene III product is 406 amino acid residues and is required for phage infection of *E. coli* via the F pili. The first  
15 two domains of the heavy chain, the variable and the CH1 domain, are fused to the carboxy-terminal half of the gene III protein. This recombinant pili protein, directed by the pel B leader, is secreted to the periplasm where it accumulates and forms disulfide bonds  
20 with the light chain before it is incorporated in the coat of the phage. Also, another vector contains a FLAG sequence engineered downstream of the gene III. The FLAG is an 8 amino acid peptide expressed at the carboxy terminal of the Fd protein. We are using commercially  
25 available monoclonal anti-FLAG M2 for both purification and detection of phage Fab by ELISA (Brizzard, Bio Techniques, 16(4):730-731, (1994)).

After constructing the vector pMS, we tested its ability to produce phage bound Fab using control  
30 antibody genes. We cloned an anti-tetanus toxoid antibody, (obtained from Dr. Carlos Barbas), into pMS and transformed XLI-blue. We co-infected our cells with VCSM13 and generated phage displaying the anti-tetanus toxoid antibody. We performed efficiency experiments



-47-

where anti-tetanus toxoid phage were combined with phage beading an irrelevant antibody at 1:100,000. We performed three rounds of panning by applying 50  $\mu$ l of the mixed phage to antigen (tetanus toxoid) coated polystyrene wells. Non-adherent phage were washed off and the adherent phage were eluted with acid. The eluted phage were used to infect a fresh aliquot of XL1-Blue bacteria and helper phage was added. After overnight amplification, phage were prepared and again panned on antigen coated plates. After three rounds of panning, we were able to show that we had successfully enriched for the anti-tetanus toxoid phage. The success of this technology also depends on the ability to prepare soluble Fabs for characterization of the final panned product. This was achieved by excising gene III from the pMS DNA using the restriction enzyme Nhe I followed by re-ligation. After the gene III was excised, the Fab was no longer displayed on the phage surface but accumulated in the periplasmic space. Lysates were prepared from bacteria expressing soluble Fab and tested for antigen specificity using an ELISA. High levels of soluble Fab were detected.

In order to adapt phage display technology for use with macaque libraries, we developed specific primers for PCR amplifying monkey immunoglobulin genes. These were based on macaque sequences we obtained while developing the PRIMATIZED<sup>TM</sup> antibody technology (See, 08/379,072, incorporated by reference herein) and databases containing human sequences. (Kabat et al, (1991), "Sequences of Proteins of Immunological Interest," U.S. Dept. of Health and Human Services, National Institute of Health).

We developed three sets of primers to cover amplification of the macaque repertoire. Our first set

-48-

of primers was designed for amplification of the heavy chain VH and CH1 (Fd) domains. It consisted of a 3' CH1 domain primer and six 5' VH family specific primers that bind in the framework 1 region. Our second set of  
5 primers, for amplifying the whole lambda chain, covers the many lambda chain subgroups. It consists of a 3' primer and three 5' degenerate primers that bind in the VL framework 1 region. Our third set of primers was designed for amplification of the kappa chain subgroups.  
10 It consists of one 3' primer and five VK framework 1 primers. Using each of these sets, PCR parameters were optimized to obtain strong enough signals from each primer pair so that ample material was available for cloning of the library. We recently created macaque  
15 combinatorial libraries in our pMS vector using these optimized PCR conditions. Bone marrow biopsies were taken from CD4 immune monkeys as the source of immunoglobulin RNA. The libraries contained approximately  $10^6$  members and are currently being panned  
20 for specific binders on antigen coated wells.

### Example 2

#### **Development of B7/CTLA-4 Reagents**

We have generated a number of reagents for the purpose of immunizing monkeys, developing binding and  
25 functional assays in vitro, screening heterohybridomas and panning phage libraries. Table 1 lists each reagent and its intended purpose. In the case of B7.1, RNA was extracted from SB cells and converted to cDNA using reverse transcriptase. The first strand cDNA was PCR  
30 amplified using B7.1 specific primers and cloned into IDEC's NEOSPLA mammalian expression vectors. CHO cells were transfected with B7.1 NEOSPLA DNA and clones expressing membrane associated B7.1 were identified.

-49-

The B7.1 fusion protein was generated similarly, except that the PCR amplified B7.1 gene was cloned into a NEOSPLA cassette vector containing the human CH2 and CH3 immunoglobulin genes. CHO cells were transformed with the B7.1/Ig NEOSPLA DNA and stable clones secreting B7.1/Ig fusion protein were amplified. In general, the B7.2 and CTLA4 reagents were generated in the same manner, except that for B7.2 the RNA was isolated from human spleen cells that had been stimulated 24 hours with anti-Ig and IL-4, and for the CTLA4 constructs the gene source was PHA activated human T cells.

Table 1

Reagent	Purpose	CHO Expression
Soluble B7.1	Immunization, immunoassays	Yes
B7.1 Transfectant	Screening, ELISA	Yes
B7.1/Ig Fusion Protein	Inhibition studies, panning	Yes
B7.2 Transfectant	Screening, ELISA	Yes
B7.2/Ig Fusion Protein	Inhibition studies, panning	To be completed
CTLA4 Transfectant	Inhibition studies	To be completed
CTLA4/Ig	Inhibition studies	To be completed

The availability of these reagents, together with monoclonal antibodies to B7.1 (L3074) (Becton Dickinson, 1994) and B7.2 (Fun-1 (Engel et al, Blood, 84, 1402-1407, (1994) and purified goat and rabbit antisera, specifically developed to detect monkey Fab fragments, facilitates identification of antibodies having the desired properties.

-50-

Example 3**Investigation of the Immune Response in Cynomolgus Monkeys to Soluble and Cell Associated Human B7.1**

To evaluate the feasibility of producing monkey  
5 antibodies to human B7.1 antigen, we first purified  
recombinant SB7.1 from CHO cell media by affinity  
chromatography using a L307.4-sepharose affinity column.  
SB7.1 was then injected, with adjuvant, into five mature  
cynomolgus macaques. After a 3 to 4 month period of  
10 booster immunizations, sera from the monkeys immunized  
with SB7.1 or human SB cells were tested for antigen  
binding.

Serum samples from the five monkeys immunized with  
SB7.1. and three additional animals immunized with B7.1  
15 positive human SB cells, were tested for antibody titers  
against membrane associated B7.1 expressed in  
transfected CHO cells. The results summarized in Figure  
3 showed that four out of five monkeys immunized with  
affinity-purified SB7.1 produced antibody titers in  
20 excess of 1:5000. The three animals immunized with SB  
cells containing cell associated B7.1 expressed lower  
titers of antibodies ranging from 1:1400 to 1:2800.

Example 4

We purified antibodies from sera of all eight  
25 immunized monkeys using SB7.1-sepharose and then tested  
their ability to bind to 1) SB7.1 coated plates in  
ELISA; 2) antigen positive B cells and 3) B7.1 CHO  
transfectomas. In addition, they were evaluated for  
their ability to block B cell interactions as measured  
30 by IL-2 production and tritiated thymidine uptake in a  
mixed lymphocyte reaction (MLR). For T cell binding  
experiments, human buffy coat peripheral blood

-51-

lymphocytes were cultured for 3-6 days in the presence of PHA stimulator. B7 binding was detected by radio assay using  $^{125}\text{I}$ -radiolabeled soluble B7.1 (SB7.1).

#### Example 5

5 **Direct binding of monkey antibodies to radiolabeled SB7.**  
 $^{125}\text{I}$  radiolabeled SB7.1 was tested for binding to anti-B7.1 antibodies at 4, 1 and 0.25  $\mu\text{g/ml}$  in solution. The results shown in Table 2 suggest that most of the antibodies produced by monkeys immunized with SB7.1 were  
10 capable of binding the affinity-purified  $^{125}\text{I}$ -SB7.1 in a concentration dependent manner. To evaluate the specificity of binding to labeled SB7.1, unlabelled SB7.1 competition experiments were done with antibodies from two animals. Affinity-purified antibodies from  
15 monkeys 1133 and 1144 were coated onto microwell plates at 400 ng/well. Affinity-purified unlabeled SB7.1 (500 and 100 ng/well) was used as competitor. The results shown in Figure 4 demonstrated that SB7.1 preparations are effective in inhibiting the  $^{125}\text{I}$ -SB7.1 from binding to  
20 the antibodies.

Table 2

**Binding of SB7- $^{125}\text{I}$  to Monkey Antibodies Affinity  
Purified on a SB7-Sepharose Affinity Column**

Antibody ( $\mu\text{g/ml}$ )	<u>Monkey Numbers</u>							
	769	908	1133	1135	1137	1139	1144	1146
4	175	213	9,056	12,771	4,318	226	5,781	108
1	106	142	6,569	7,940	3,401	110	3,901	80
0.25	95	104	1,803	2,673	1,219	100	1,186	94

-52-

Data are mean values of duplicate assays and represent cpm SB7-I<sup>125</sup> bound.

#### Example 6

5     **Direct binding of radiolabeled affinity-purified monkey antibodies to B7<sup>+</sup> cells and inhibition by SB7.1.**

       Affinity-purified radiolabeled monkey anti-B7.1 antibodies from monkey PRI135 were compared with radiolabeled L307.4 MAbs for direct binding to B7 positive human SB cells. As a specificity control,  
10     unlabeled SB7.1 (0.002 - 20 µg/ml) was added to compete with both radiolabeled antibodies. We demonstrated that monkey antibodies can bind cell associated B7.1 and are inhibited with SB7.1, as shown in Figure 5. Inhibition as high as 90% was observed with SB7.1.

15                     Example 7

**Direct binding of radiolabeled B7-Ig fusion protein to activated T cells and Inhibition by affinity-purified monkey antibodies.**

       Human peripheral blood T lymphocytes were activated  
20     for 3-6 days and tested for direct binding of <sup>125</sup>I-B7.1-Ig. Because of Fc receptor upregulation on activated human T cells, it was necessary to pre-incubate the cells with heat-aggregated pre-immune immunoglobulin to block Fc binding sites prior to  
25     addition of B7.1-Ig to the cells. A background control using SP2/0 murine myeloma cells was included to allow correction of the background binding. Figure 6 shows that inhibition of <sup>125</sup>I-B7.1-Ig fusion protein binding to

-53-

activated T cells was achieved with affinity-purified monkey antibodies at concentrations from 200 to 8  $\mu\text{g/ml}$ . Unlabeled SB7.1 and L307.4 MAb used as controls were also effective in inhibiting B7.1-Ig fusion protein cell binding.

#### Example 8

#### **Inhibition of IL-2 production in mixed lymphocyte reactions by monkey anti-B7 antibodies.**

The blocking of CD28/B7 interaction leads to inhibition of IL-2 production by T lymphocytes. In the experiment shown in Figure 7, affinity-purified monkey antibodies from two monkeys immunized with SB7.1 (monkeys 1137 and 1135) and one immunized with B7 positive SB cells (monkey 1146) were evaluated for their abilities to inhibit human T cell activation in mixed lymphocyte reaction (MLR), as measured by inhibition of IL-2 production. The results of this experiment show that affinity-purified anti-B7.1 antibodies from monkeys 1146 and 1137 inhibited IL-2 production when added at concentrations of 50  $\mu\text{g/ml}$ . Monkey 1135 antibodies could not be evaluated at the two highest concentrations due to lack of material, yet gave significant inhibition at lower concentrations. The murine MAb L307.4 was inhibitory at concentrations of 10  $\mu\text{g/ml}$ . Other monkey sera tested at these concentrations were negative (data not shown). These results demonstrate that at least three of the monkeys immunized with both soluble and membrane associated forms of the B7 antigen are producing B7-blocking antibodies with immunosuppressive potential.

-54-

Example 9**Investigation of Cross-reactivity in B7.1 Immunized Monkey Serum to B7.2 Antigen.**

Antibodies raised against B7.1 are to be tested for cross-reactivity to B7.2. Preliminary results using B7.1 affinity-purified antibodies from B7.1 immune sera provided suggestive evidence of binding to B7.2 transfected CHO cells (not shown). These data should be confirmed by using soluble B7.2Ig reagents. We will first purify additional monkey antibodies from B7.1 immunized animals by affinity chromatography on B7.1Ig--sepharose. We will then produce and purify B7.2Ig from CHO cells in sufficient quantities to prepare a B7.2Ig-sepharose affinity column. We will select from the B7.1 specific antibody population those antibodies which cross-react with B7.2 by binding to the B7.2Ig-sepharose column. Any cross-reactive antibodies identified will be further characterized by direct binding to both B7.1 and B7.2 transfected CHO cells and inhibition of binding to B7.2 transfected cells by B7.1Ig.

Example 10**Generation of a Phage Display Library**

Recombinant phage display libraries are generated from B7.1 and B7.2 immune monkeys. Lymph node and bone marrow biopsies are performed 7-12 days after immunization to harvest RNA rich B cells and plasma cells. RNA is isolated from the lymphocytes using the method described by Chomczynski Anal. Biochem., 162(1), 156-159, (1987). RNA is converted to cDNA using an oligo dT primer and reverse transcriptase. The first



-55-

strand cDNA is divided into aliquots and PCR amplified using the sets of kappa, lambda, and heavy chain Fd region primers described earlier and either Pfu polymerase (Stratagene, San Diego) or Taq polymerase (Promega, Madison). The heavy chain PCR amplified products are pooled, cut with Xho VSpe I restriction enzymes and cloned into the vector pMS. Subsequently, the light chain PCR products are pooled, cut with Sac I/Xba I restriction enzymes, and cloned to create the recombinant library. XLI-Blue *E. coli* is transformed with the library DNA and super-infected with VCSM13 to produce the phage displaying antibodies. The library is panned four rounds on polystyrene wells coated with B7.1 or B7.2 antigen. Individual phage clones from each round of panning are analyzed. The pMS vector DNA is isolated and the gene III excised. Soluble Fab fragments are generated and tested in ELISA for binding to B7.1 and B7.2.

#### Example 11

##### 20 Characterization of Phage Fab Fragments

The monkey phage Fab fragments are characterized for their specificity and the ability to block B7.1-Ig and B7.2-Ig binding to CTLA-4-Ig or CTLA-4 transfected cells. Phage fragments are also characterized for cross-reactivity after first panning for 4 rounds on the B7 species used for immunization in order to select for high affinity fragments. Fab fragments identified from four rounds of panning either on B7.1 or B7.2 antigen coated surfaces are scaled up by infection and grown in 24 hour fermentation cultures of *E. coli*. Fragments are purified by Kodak FLAG binding to a anti-FLAG affinity column. Purified phage Fabs are tested for affinity by an ELISA based direct binding modified Scatchard

-56-

analysis (Kato et al, J. Chem. BioEng., 76:451-454, (1993)) using Goat anti-monkey Fab antibodies or anti-FLAG MAb conjugated with horseradish peroxidase. The anti-monkey Fab reagents will be absorbed against human heavy chain constant region Ig to remove any cross-reactivity to B7-Ig. Kd values are calculated for each fragment after measurements of direct binding to B7.1-Ig or B7.2-Ig coated plates.

#### Example 12

##### 10 Phage Fab Fragment Blocking of CTLA-4/B7 Binding

Fab fragments most effectively blocking the binding of B7-Ig at the lowest concentrations are selected as lead candidates. Selections are made by competing off <sup>125</sup>I-B7-Ig binding to CTLA-4-Ig or CTLA-4 transfected cells. Additional selection criteria include, blocking of mixed lymphocyte reaction (MLR), as measured by inhibiting 3H-thymidine uptake in responder cells (Azuma et al, J. Exp. Med., 177:845-850,; Azuma et al, Nature, 301:76-79, (1993)) and direct analysis of IL-2 production using IL-2 assay kits. The three or four candidates which are most effective in inhibiting of MLR and CTLA-4 binding assays are chosen for cloning into the above-described mammalian expression vector for transfection into CHO cells and expression of chimeric monkey/human antibodies.

#### Example 13

##### Generation of Monkey Heterohybridomas

Monkey heterohybridomas secreting monoclonal antibodies are generated from existing immunized animals whose sera tested positive for B7.1 and/or B7.2. Lymph node biopsies are taken from animals positive to either, or both, antigens. The method of hybridoma production

-57-

is similar to the established method used for the generation of monkey anti-CD4 antibodies (Newman, 1992(Id.)). Monkeys with high serum titers will have sections of inguinal lymph nodes removed under  
5 anesthesia. Lymphocytes are washed from the tissue and fused with KH6/B5 heteromyeloma cells (Carrol et al, J. Immunol. Meth., 89:61-72, (1986)) using polyethylene glycol (PEG). Hybridomas are selected on H.A.T. media and stabilized by repeated subcloning in 96 well plates.

10 Monkey monoclonal antibodies specific for B7.1 antigen are screened for cross-reactivity to B7.2. Monkey anti-B7 antibodies will be characterized for blocking of B7/CTLA-4 binding using the <sup>125</sup>I-B7-Ig binding  
15 assay. Inhibition of MLR by 3H-Thymidine uptake and direct measurement of IL-2 production is used to select three candidates. Two candidates will be brought forward in Phase II studies and expressed in CHO cells while repeating all functional studies. For the  
20 purposes of developing an animal model for *in vivo* pharmacology, anti-B7 antibodies will be tested on cells of several animal species. The establishment of an animal model will allow preclinical studies to be carried out for the selected clinical indication.

#### Example 14

25 As discussed *supra*, using the above heterohybridoma methods, 4 lead monkey anti-B7.1 antibodies have been identified: 16C10, 7B6, 7C10 and 20C9. These antibodies were characterized as follows:

30 To demonstrate the monkey antibodies' ability to block the physical interaction between CTLA4-Ig, varying concentrations of the monkey anti-B7.1 antibodies and unlabeled CTLA4-Ig were incubated

-58-

with radiolabeled CTLA4-Ig<sup>1125</sup>. The results of the inhibition assay showed that the IC<sub>50</sub> (the concentration of inhibitor which results in 50% inhibition) for the monkey antibodies are:

5	a:	7C10:	0.39 µg/Ml
	b:	16C10:	1.60 µg/Ml
	c:	20C9:	3.90 µg/Ml
	d:	7B6:	39.0 µg/Ml

Scatchard analysis showed that the apparent affinity constants (K<sub>d</sub>) for the monkey antibodies binding to B7-Ig coated plates were approximated to be:

	a:	7C10:	6.2 x 10 <sup>-9</sup> M
	b:	16C10:	8.1 x 10 <sup>-9</sup> M
15	c:	7B6:	10.7 x 10 <sup>-9</sup> M
	d:	20C9:	16.8 x 10 <sup>-9</sup> M

The antibodies were tested *in vitro* in a mixed lymphocyte reaction assay (MLR). The MLR showed that all 4 anti-B7.1 antibodies inhibit IL-2 production to different extents:

20	a:	7B6:	5.0 µg/Ml
	b:	16C10:	0.1 µg/Ml
	c:	20C9:	2.0 µg/Ml
	d:	7C10:	5.0 µg/Ml

25 The monkey anti-B7.1 antibodies were tested for their ability to bind B7 on human peripheral blood lymphocytes (PBL). FACS analysis showed that all 4 monkey antibodies tested positive.

30 Monkey antibodies 16C10, 7B6, 7C10 and 20C9 were tested for Clq binding by FACS analysis. Results

-59-

showed 7C10 monkey Ig had strong human C1q binding after incubating with B7.1 CHO-transfected cells. 16C10 was negative, as were the 20C9 and 7B6 monkey antibodies.

5

Example 15

Using the primatized antibody methodology incorporated by reference to commonly assigned U.S. Serial No. 08/379,072, and using the NEOSPLA vector system shown in Figure 2, the heavy and light variable  
10 domains of 7C10, 7B6 and 16C10 were cloned and primatized forms thereof have been synthesized in CHO cells using the NEOSPLA vector system. The amino acid and nucleic acid sequences for the primatized 7C10 light and heavy chain, 7B6 light and heavy chain, and 16C10  
15 light and heavy chain are respectively shown in Figures 8a, 8b, 9a, 9b, 10a and 10b.

It is expected that these primatized antibodies, given their probable low antigenicity and human effector function, will be well suited as therapeutics. In fact,  
20 it has recently been shown that primatized 16C10 exhibits human C1<sub>q</sub> binding, whereas 16C10 does not.

Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation many equivalents to the specific embodiments of the  
25 invention described herein. Such equivalents are intended to be embraced by the following claims.

-60-

## Claims:

1. A biologically pure monkey monoclonal antibody or a primatized form thereof which specifically binds human B7.1 antigen and/or human B7.2 antigen.

5        2. The antibody of claim 1 which is selected from the group consisting of 16C10, 7C10, 20C9 and 7B6.

3. The antibody of claim 1 which is a depleting antibody.

10       4. The antibody of claim 1 which is a non-depleting antibody.

5. A primatized antibody which specifically binds to human B7.1 antigen which contains the variable heavy and light domains of an antibody selected from the group consisting of 16C10, 7C10, 20C9 and 7B6.

15       6. The primatized antibody of claim 5 wherein said antibody is derived from 7C10 and has the amino acid sequence set forth in Figures 8a and 8b.

20       7. The primatized antibody of claim 6 which is encoded by the nucleic acid sequence set forth in Figures 8a and 8b.

25       8. The primatized antibody of claim 5 wherein said antibody is derived from 7B6 and has the amino acid sequence set forth in Figures 9a and 9b.

9. The primatized antibody of claim 8 wherein said antibody is encoded by the nucleic acid sequence set forth in Figures 9a and 9b.



-61-

10. The primatized antibody of claim 5 wherein said antibody is derived from 16C10 and has the amino acid sequence set forth in Figures 4a and 4b.

11. The primatized antibody of claim 5 wherein said antibody is encoded by the nucleic acid sequence set forth in Figures 10a and 10b.

12. A transfectoma which expresses a primatized antibody which specifically binds to human B7.1 and/or human B7.2 antigen.


13. The transfectoma of claim 12 which is a CHO cell.

14. The transfectoma of claim 13 wherein said cell expresses a primatized antibody having the amino acid sequence set forth in any one of Figures 8a, 8b, 9a, 9b, 10a and 10b.

15. A pharmaceutical composition suitable for treatment of a disease treatable by inhibition of B7-CD28 binding which includes an antibody according to any one of claims 1 to 11 together with a pharmaceutically acceptable additive, adjuvant or carrier.

16. A method of treating a disease by inhibition of the B7:CD28 pathway which includes administering to a subject in need of such treatment a therapeutically effective amount of at least one antibody according to any one of claims 1 to 11.

17. The method of claim 16 wherein said antibody is 16C10, 7C10, 20C9, 7B6 or a primatized form thereof.



-62-

18. The method of claim 16 wherein said disease is an autoimmune disorder.

19. The method of claim 16 wherein said disease is selected from idiopathic thrombocytopenia purpura,  
5 systemic lupus erythematosus, type 1 diabetes mellitus, rheumatoid arthritis, psoriasis and multiple sclerosis.

20. The method of claim 16 wherein said disease is graft-versus-host disease.



1 / 15





FIG. 1

FIG. 2



3 / 15

FIG. 3






FIG. 4




4 / 15

FIG. 5




5 / 15



6 / 15

FIG. 7



7 / 15

Frame 1 M R V P A Q L L G L L L L W L P G A R  
 ATG AGG GTC CCC GCT CAG CTC CTG GGG CTC CTG CTC TGG CTC CCA GGT GCA CGA  
 9 18 27 36 45 54  
 C A Y E L T Q P P S V S V S P G Q T A R I  
 TGT GCC TAT GAA CTG ACT CAG CCA CCC TCG GTG TCA GTG TCC CCA GGA CAG ACG GCC AGG ATC  
 63 72 81 90 99 108 117  
 T C G G D N S R N E Y V H W Y Q Q K P A R  
 ACC TGT GGG GGA GAC AAC AGT AGA AAT GAA TAT GTG CAC TGG TAC CAG CAG AAG CCA ACG CGG  
 126 135 144 153 162 171 180  
 A P I L V I Y D D S D R P S G I P E R F S  
 GCC CCT ATA CTG GTC ATC TAT GAT GAT AGT GAC CGG CCC TCA GGG ATC CCT GAG CGA TTC TCT  
 189 198 207 216 225 234 243  
 G S K S G N T A T L T I N G V E A G D E A  
 GGC TCC AAA TCA GGG AAC ACC GCC ACC CTG ACC ATC AAC GGG GTC GAG GCC GGG GAT GAG GCT  
 252 261 270 279 288 297 306  
 D Y Y C Q V W D R A S D H P V F G G G T R  
 GAC TAT TAC TGT CAG GTG TGG GAC AGG GCT AGT GAT CAT CCG GTC TTC GGA GGA GGG ACC CGG  
 315 324 333 342 351 360 369  
 V T V L G Q P K A A P S V T L F P P S S E  
 GTG ACC GTC CTA GGT CAG CCC AAG GCT GCC CCC TCG GTC ACT CTG TTC CCG CCC TCC TCT GAG  
 378 387 396 405 414 423 432  
 E L Q A N K A T L V C L I S D F Y P G A V  
 GAG CTT CAA GCC AAC AAG GCC ACA CTG GTG TGT CTC ATA AGT GAC TTC TAC CCG GGA GCC GTG  
 441 450 459 468 477 486 495  
 T V A W K A D S S P V K A G V E T T T P S  
 ACA GTG GCC TGG AAG GCA GAT AGC AGC CCC GTG AAG GCG GGA GTG GAG ACC ACC ACA CCC TCC  
 504 513 522 531 540 549 558  
 K Q S N N K Y A A S S Y L S L T P E Q W K  
 AAA CAA AGC AAC AAC AAG TAC GCG GCC AGC AGC TAC CTG AGC CTG ACG CCT GAG CAG TGG AAG  
 567 576 585 594 603 612 621  
 S H R S Y S C Q V T H E G S T V E K T V A  
 TCC CAC AGA AGC TAC AGC TGC CAG GTC ACG CAT GAA GGG AGC ACC GTG GAG AAG ACA GTG GCC  
 630 639 648 657 666 675 684  
 P T E C S  
 CCT ACA GAA TGT TCA TGA  
 693 702

FIG. 8a

8 / 15

Frame 1 M K H L W F F L L L V A A P R W V L S  
 ATG AAA CAC CTG TGG TTC TTC CTC CTC GTG GCA GCT CCC AGA TGG GTC CTG TCC  
 9 18 27 36 45 54  
 Q V K L Q Q W G E G L L Q P S E T L S R T  
 CAG GTG AAG CTG CAG CAG TGG GGC GAA GGA CTT CTG CAG CCT TCG GAG ACC CTG TCC CGC ACC  
 63 72 81 90 99 108 117  
 C V V S G G S I S G Y Y Y W T W I R Q T P  
 TGC GTT GTC TCT GGT GGC TCC ATC AGC GGT TAC TAC TAC TGG ACC TGG ATC CGC CAG ACC CCA  
 126 135 144 153 162 171 180  
 G R G L E W I G H I Y G N G A T T N Y N P  
 GGG AGG GGA CTG GAG TGG ATT GGC CAT ATT TAT GGT AAT GGT GCG ACC ACC AAC TAC AAT CCC  
 189 198 207 216 225 234 243  
 S L K S R V T I S K D T S K N Q F F L N L  
 TCC CTC AAG AGT CGA GTC ACC ATT TCA AAA GAC ACG TCC AAG AAC CAG TTC TTC CTG AAC TTG  
 252 261 270 279 288 297 306  
 N S V T D A D T A V Y Y C A R G P R P D C  
 AAT TCT GTG ACC GAC GCG GAC ACG GCC GTC TAT TAC TGT GCG AGA GGC CCT CGC CCT GAT TGC  
 315 324 333 342 351 360 369  
 T T I C Y G G W V D V W G P G D L V T V S  
 ACA ACC ATT TGT TAT GGC GGC TGG GTC GAT GTC TGG GGC CCG GGA GAC CTG GTC ACC GTC TCC  
 378 387 396 405 414 423 432  
 S A S T K G P S V F P L A P S S K S T S G  
 TCA GCT AGC ACC AAG GGC CCA TCG GTC TTC CCC CTG GCA CCC TCC TCC AAG AGC ACC TCT GGG  
 441 450 459 468 477 486 495  
 G T A A L G C L V K D Y F P E P V T V S W  
 GGC ACA GCG GCC CTG GGC TGC CTG GTC AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG TCG TGG  
 504 513 522 531 540 549 558  
 N S G A L T S G V H T F P A V L Q S S G L  
 AAC TCA GGC GCC CTG ACC AGC GGC GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC TCA GGA CTC  
 567 576 585 594 603 612 621  
 Y S L S S V V T V P S S S L G T Q T Y I C  
 TAC TCC CTC AGC AGC GTG GTG ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG ACC TAC ATC TGC  
 630 639 648 657 666 675 684  
 N V N H K P S N T K V D K K A E P K S C D  
 AAC GTG AAT CAC AAG CCC AGC AAC ACC AAG GTG GAC AAG AAA GCA GAG CCC AAA TCT TGT GAC  
 693 702 711 720 729 738 747  
 K T H T C P P C P A P E L L G G P S V F L  
 AAA ACT CAC ACA TGC CCA CCG TGC CCA GCA CCT GAA CTC CTG GGG GGA CCG TCA GTC TTC CTC  
 756 765 774 783 792 801 810  
 F P P K P K D T L M I S R T P E V T C V V  
 TTC CCC CCA AAA CCC AAG GAC ACC CTC ATG ATC TCC CGG ACC CCT GAG GTC ACA TGC GTG GTG  
 819 828 837 846 855 864 873  
 V D V S H E D P E V K F N W Y V D G V E V  
 GTG GAC GTG AGC CAC GAA GAC CCT GAG GTC AAG TTC AAC TGG TAC GTG GAC GGC GTG GAG GTG  
 882 891 900 909 918 927 936

FIG. 8b-1

SUBSTITUTE SHEET (RULE 26)



9 / 15

H	N	A	K	T	K	P	R	E	E	Q	Y	N	S	T	Y	R	V	V	S	V
CAT	AAT	GCC	AAG	ACA	AAG	CCG	CGG	GAG	GAG	CAG	TAC	AAC	AGC	ACG	TAC	CGT	GTG	GTC	AGC	GTC
945				954			963			972			981			990			999	
L	T	V	L	H	Q	D	W	L	N	G	K	E	Y	K	C	K	V	S	N	K
CTC	ACC	GTC	CTG	CAC	CAG	GAC	TGG	CTG	AAT	GGC	AAG	GAG	TAC	AAG	TGC	AAG	GTC	TCC	AAC	AAA
1008			1017				1026			1035			1044			1053			1062	
A	L	P	A	P	I	E	K	T	I	S	K	A	K	G	Q	P	R	E	P	Q
GCC	CTC	CCA	GCC	CCC	ATC	GAG	AAA	ACC	ATC	TCC	AAA	GCC	AAA	GGG	CAG	CCC	CGA	GAA	CCA	CAG
1071			1080				1089			1098			1107			1116			1125	
V	Y	T	L	P	P	S	R	D	E	L	T	K	N	Q	V	S	L	T	C	L
GTG	TAC	ACC	CTG	CCC	CCA	TCC	CGG	GAT	GAG	CTG	ACC	AAG	AAC	CAG	GTC	AGC	CTG	ACC	TGC	CTG
1134			1143				1152			1161			1170			1179			1188	
V	K	G	F	Y	P	S	D	I	A	V	E	W	E	S	N	G	Q	P	E	N
GTC	AAA	GGC	TTC	TAT	CCC	AGC	GAC	ATC	GCC	GTG	GAG	TGG	GAG	AGC	AAT	GGG	CAG	CCG	GAG	AAC
1197			1206				1215			1224			1233			1242			1251	
N	Y	K	T	T	P	P	V	L	D	S	D	G	S	F	F	L	Y	S	K	L
AAC	TAC	AAG	ACC	ACG	CCT	CCC	GTG	CTG	GAC	TCC	GAC	GGC	TCC	TTC	TTC	CTC	TAC	AGC	AAG	CTC
1260			1269				1278			1287			1296			1305			1314	
T	V	D	K	S	R	W	Q	Q	G	N	V	F	S	C	S	V	M	H	E	A
ACC	GTG	GAC	AAG	AGC	AGG	TGG	CAG	CAG	GGG	AAC	GTC	TTC	TCA	TGC	TCC	GTG	ATG	CAT	GAG	GCT
1323			1332				1341			1350			1359			1368			1377	
L	H	N	H	Y	T	Q	K	S	L	S	L	S	P	G	K					
CTG	CAC	AAC	CAC	TAC	ACG	CAG	AAG	AGC	CTC	TCC	CTG	TCT	CCG	GGT	AAA	TGA				
1386			1395				1404			1413			1422			1431				

FIG. 8b-2

10 / 15

Frame 1 M S L P A Q L L G L L L L C V P G S S  
 ATG AGC CTC CCT GCT CAG CTC CTC GGG CTG CTA TTG CTC TGC GTC CCC GGG TCC AGT  
 9 18 27 36 45 54  
 G E V V M T Q S P L S L P I T P G E P A S  
 GGG GAA GTT GTG ATG ACT CAG TCT CCA CTG TCC CTT CCC ATC ACA CCT GGA GAG CCG GCC TCC  
 63 72 81 90 99 108 117  
 I S C R S S Q S L K H S N G D T F L S W Y  
 ATC TCC TGT AGG TCT AGT CAA AGC CTT AAA CAC AGT AAT GGA GAC ACC TTC CTG AGT TGG TAT  
 126 135 144 153 162 171 180  
 Q Q K P G Q P P R L L I Y K V S N R D S G  
 CAG CAG AAG CCA GGC CAA CCT CCA AGG CTC CTG ATT TAT AAG GTT TCT AAC CCG GAC TCT GGG  
 189 198 207 216 225 234 243  
 V P D R F S G S G A G T D F T L K I S A V  
 GTC CCA GAC AGA TTC AGC GGC AGT GGG GCA GGG ACA GAT TTC ACA CTG AAA ATC AGC GCA GTG  
 252 261 270 279 288 297 306  
 E A E D V G V Y F C G Q G T R T P P T F G  
 GAG GCT GAA GAT GTT GGG GTT TAT TTC TGC GGG CAA GGT TCA AGG ACT CCT CCC ACT TTC GGC  
 315 324 333 342 351 360 369  
 G G T K V E I K R T V A A P S V F I F P P  
 GGA GGG ACC AAG GTG GAA ATC AAA CGT ACG GTG GCT GCA CCA TCT GTC TTC ATC TTC CCG CCA  
 378 387 396 405 414 423 432  
 S D E Q L K S G T A S V V C L L N N F Y P  
 TCT GAT GAG CAG TTG AAA TCT GGA ACT GCC TCT GTT GTG TGC CTG CTG AAT AAC TTC TAT CCC  
 441 450 459 468 477 486 495  
 R E A K V Q W K V D N A L Q S G N S Q E S  
 AGA GAG GCC AAA GTA CAG TGG AAG GTG GAT AAC GCC CTC CAA TCG GGT AAC TCC CAG GAG AGT  
 504 513 522 531 540 549 558  
 V T E Q D S K D S T Y S L S S T L T L S K  
 GTC ACA GAG CAG GAC AGC AAG GAC AGC ACC TAC AGC CTC AGC AGC ACC CTG ACG CTG AGC AAA  
 567 576 585 594 603 612 621  
 A D Y E K H K V Y A C E V T H Q G L S S P  
 GCA GAC TAC GAG AAA CAC AAA GTC TAC GCC TGC GAA GTC ACC CAT CAG GGC CTG AGC TCG CCC  
 630 639 648 657 666 675 684  
 V T K S F N R G E C  
 GTC ACA AAG AGC TTC AAC AGG GGA GAG TGT TGA  
 693 702 711 720

FIG. 9a

11 / 15

Frame 1 M G W S L I L L F L V A V A T R V Q C  
 ATG GGT TGG AGC CTC ATC TTG CTC TTC CTT GTC GCT GTT GCT ACG CGT GTC CAG TGT  
 9 18 27 36 45 54  
 E V Q L V E S G G G L V Q P G G S L R V S  
 GAG GTG CAA CTG GTG GAG TCT GGG GGA GGC TTG GTC CAG CCT GGC GGG TCC CTG AGA GTC TCC  
 63 72 81 90 99 108 117  
 C A V S G F T F S D H Y M Y W F R Q A P G  
 TGT GCA GTC TCT GGA TTC ACC TTC AGT GAC CAC TAC ATG TAT TGG TTC CGC CAG GCT CCA GGG  
 126 135 144 153 162 171 180  
 K G P E W V G F I R N K P N G G T T E Y A  
 AAG GGG CCG GAA TGG GGA GGT TTC ATT AGA AAC AAA CCG AAC GGT GGC ACA ACA GAA TAC GCC  
 189 198 207 216 225 234 243  
 A S V K D R F T I S R D D S K S I A Y L Q  
 GCG TCT GTG AAA GAC AGA TTC ACC ATC TCC AGA GAT GAT TCC AAA AGC ATC GCC TAT CTG CAA  
 252 261 270 279 288 297 306  
 M S S L K I E D T A V Y Y C T T S Y I S H  
 ATG AGC AGC CTG AAA ATC GAG GAC ACG GCC GTC TAT TAC TGT ACT ACA TCC TAC ATT TCA CAT  
 315 324 333 342 351 360 369  
 C R G G V C Y G G Y F E F W G Q G A L V T  
 TGT CCG GGT GGT GTC TGC TAT GGA GGT TAC TTC GAA TTC TGG GGC CAG GGC GCC CTG GTC ACC  
 378 387 396 405 414 423 432  
 V S S A S T K G P S V F P L A P S S K S T  
 GTC TCC TCA GCT AGC ACC AAG GGC CCA TCG GTC TTC CCC CTG GCA CCC TCC TCC AAG AGC ACC  
 441 450 459 468 477 486 495  
 S G G T A A L G C L V K D Y F P E P V T V  
 TCT GGG GGC ACA GCG GCC CTG GGC TGC CTG GTC AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG  
 504 513 522 531 540 549 558  
 S W N S G A L T S G V H T F P A V L Q S S  
 TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC TCA  
 567 576 585 594 603 612 621  
 G L Y S L S S V V T V P S S S L G T Q T Y  
 GGA CTC TAC TCC CTC AGC AGC GTG GTG ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG ACC TAC  
 630 639 648 657 666 675 684  
 I C N V N H K P S N T K V D K K A E P K S  
 ATC TGC AAC GTG AAT CAC AAG CCC AGC AAC ACC AAG GTG GAC AAG AAA GCA GAG CCC AAA TCT  
 693 702 711 720 729 738 747  
 C D K T H T C P P C P A P E L L G G P S V  
 TGT GAC AAA ACT CAC ACA TGC CCA CCG TGC CCA GCA CCT GAA CTC CTG GGG GGA CCG TCA GTC  
 756 765 774 783 792 801 810  
 F L F P P K P K D T L M I S R T P E V T C  
 TTC CTC TTC CCC CCA AAA CCC AAG GAC ACC CTC ATG ATC TCC CGG ACC CCT GAG GTC ACA TGC  
 819 828 837 846 855 864 873  
 V V V D V S H E D P E V K F N W Y V D G V  
 GTG GTG GTG GAC GTG AGC CAC GAA GAC CCT GAG GTC AAG TTC AAC TGG TAC GTG GAC GGC GTG  
 882 891 900 909 918 927 936

FIG. 9b-1

SUBSTITUTE SHEET (RULE 26)

12 / 15

E	V	H	N	A	K	T	K	P	R	E	E	Q	Y	N	S	T	Y	R	V	V
GAG	GTG	CAT	AAT	GCC	AAG	ACA	AAG	CCG	CGG	GAG	GAG	CAG	TAC	AAC	AGC	ACG	TAC	CGT	GTG	GTC
945				954			963			972			981		990			999		
S	V	L	T	V	L	H	Q	D	W	L	N	G	K	E	Y	K	C	K	V	S
AGC	GTC	CTC	ACC	GTC	CTG	CAC	CAG	GAC	TGG	CTG	AAT	GGC	AAG	GAG	TAC	AAG	TGC	AAG	GTC	TCC
1008				1017			1026			1035			1044		1053			1062		
N	K	A	L	P	A	P	I	E	K	T	I	S	K	A	K	G	Q	P	R	E
AAC	AAA	GCC	CTC	CCA	GCC	CCC	ATC	GAG	AAA	ACC	ATC	TCC	AAA	GCC	AAA	GGG	CAG	CCC	CGA	GAA
1071				1080			1089			1098			1107		1116			1125		
P	Q	V	Y	T	L	P	P	S	R	D	E	L	T	K	N	Q	V	S	L	T
CCA	CAG	GTG	TAC	ACC	CTG	CCC	CCA	TCC	CGG	GAT	GAG	CTG	ACC	AAG	AAC	CAG	GTC	AGC	CTG	ACC
1134				1143			1152			1161			1170		1179			1188		
C	L	V	K	G	F	Y	P	S	D	I	A	V	E	W	E	S	N	G	Q	P
TGC	CTG	GTC	AAA	GGC	TTC	TAT	CCC	AGC	GAC	ATC	GCC	GTG	GAG	TGG	GAG	AGC	AAT	GGG	CAG	CCG
1197				1206			1215			1224			1233		1242			1251		
E	N	N	Y	K	T	T	P	P	V	L	D	S	D	G	S	F	F	L	Y	S
GAG	AAC	AAC	TAC	AAG	ACC	ACG	CCT	CCC	GTG	CTG	GAC	TCC	GAC	GGC	TCC	TTC	TTC	CTC	TAC	ACC
1260				1269			1278			1287			1296		1305			1314		
K	L	T	V	D	K	S	R	W	Q	Q	G	N	V	F	S	C	S	V	M	H
AAG	CTC	ACC	GTG	GAC	AAG	AGC	AGG	TGG	CAG	CAG	GGG	AAC	GTC	TTC	TCA	TGC	TCC	GTG	ATG	CAT
1323				1332			1341			1350			1359		1368			1377		
E	A	L	H	N	H	Y	T	Q	K	S	L	S	L	S	P	G	K			
GAG	GCT	CTG	CAC	AAC	CAC	TAC	ACG	CAG	AAG	AGC	CTC	TCC	CTG	TCT	CCG	GGT	AAA	TGA		
1386				1395			1404			1413			1422		1431					

FIG. 9b-2

13 / 15

Frame 1 M R V P A Q L L G L L L L W L P G A R  
 ATG AGG GTC CCC GCT CAG CTC CTG GGG CTC CTG CTG CTC TGG CTC CCA GGT GCA CGA  
 9 18 27 36 45 54  
 C E S V L T Q P P S V S G A P G Q K V T I  
 TGT GAG TCT GTC CTG ACA CAG CCG CCC TCA GTG TCT GGG GCC CCA GGG CAG AAG GTC ACC ATC  
 63 72 81 90 99 108 117  
 S C T G S T S N I G G Y D L H W Y Q Q L P  
 TCG TGC ACT GGG AGC ACC TCC AAC ATT GGA GGT TAT GAT CTA CAT TGG TAC CAG CAG CTC CCA  
 126 135 144 153 162 171 180  
 G T A P K L L I Y D I N K R P S G I S D R  
 GGA ACG GCC CCC AAA CTC CTC ATC TAT GAC ATT AAC AAG CGA CCC TCA GGA ATT TCT GAC CGA  
 189 198 207 216 225 234 243  
 F S G S K S G T A A S L A I T G L Q T E D  
 TTC TCT GGC TCC AAG TCT GGT ACC GCG GCC TCC CTG GCC ATC ACT GGG CTC CAG ACT GAG GAT  
 252 261 270 279 288 297 306  
 E A D Y Y C Q S Y D S S L N A Q V F G G G  
 GAG GCT GAT TAT TAC TGC CAG TCC TAT GAC AGC AGC CTG AAT GCT CAG GTA TTC GGA GGA GGG  
 315 324 333 342 351 360 369  
 T R L T V L G Q P K A A P S V T L F P P S  
 ACC CGG CTG ACC GTC CTA GGT CAG CCC AAG GCT GCC CCC TCG GTC ACT CTG TTC CCG CCC TCC  
 378 387 396 405 414 423 432  
 S E E L Q A N K A T L V C L I S D F Y P G  
 TCT GAG GAG CTT CAA GCC AAC AAG GCC ACA CTG GTG TGT CTC ATA AGT GAC TTC TAC CCG GGA  
 441 450 459 468 477 486 495  
 A V T V A W K A D S S P V K A G V E T T T  
 GCC GTG ACA GTG GCC TGG AAG GCA GAT AGC AGC CCC GTC AAG GCG GGA GTG GAG ACC ACC ACA  
 504 513 522 531 540 549 558  
 P S K Q S N N K Y A A S S Y L S L T P E Q  
 CCC TCC AAA CAA AGC AAC AAC AAG TAC GCG GCC AGC AGC TAC CTG AGC CTG ACG CCT GAG CAG  
 567 576 585 594 603 612 621  
 W K S H R S Y S C Q V T H E G S T V E K T  
 TGG AAG TCC CAC AGA AGC TAC AGC TGC CAG GTC ACG CAT GAA GGG AGC ACC GTG GAG AAG ACA  
 630 639 648 657 666 675 684  
 V A P T E C S  
 GTG GCC CCT ACA GAA TGT TCA TGA  
 693 702 711

FIG. 10a

14 / 15

Frame 1 M K H L W F F L L L V A A P R W V L S  
 ATG AAA CAC CTG TGG TTC TTC CTC CTC CTG GTG GCA GCT CCC AGA TGG GTC CTG TCC  
                   9                  18                  27                  36                  45                  54

Q V Q L Q E S G P G L V K P S E T L S L T  
 CAG GTG CAG CTG CAG GAG TCG GGC CCA GGA CTG GTG AAG CCT TCG GAG ACC CTG TCC CTC ACC  
           63                  72                  81                  90                  99                  108                  117

C A V S G G S I S G G Y G W G W I R Q P P  
 TGC GCT GTC TCT GGT GGC TCC ATC AGC GGT GGT TAT GGC TGG GGC TGG ATC CGC CAG CCC CCA  
           126                  135                  144                  153                  162                  171                  180

G K G L E W I G S F Y S S S G N T Y Y N P  
 GGG AAG GGG CTG GAG TGG ATT GGG AGT TTC TAT AGT AGT AGT GGG AAC ACC TAC TAC AAC CCC  
           189                  198                  207                  216                  225                  234                  243

S L K S Q V T I S T D T S K N Q F S L K L  
 TCC CTC AAG AGT CAA GTC ACC ATT TCA ACA GAC ACG TCC KAG AAC CAG TTC TCC CTG AAG CTG  
           252                  261                  270                  279                  288                  297                  306

N S M T A A D T A V Y Y C V R D R L F S V  
 AAC TCT ATG ACC GCC GCG GAC ACG GCC GTG TAT TAC TGT GTG AGA GAT CGT CTT TTT TCA GTT  
           315                  324                  333                  342                  351                  360                  369

V G M V Y N N W F D V W G P G V L V T V S  
 GTT GGA ATG GTT TAC AAC AAC TGG TTC GAT GTC TGG GGC CCG GGA GTC CTG GTC ACC GTC TCC  
           378                  387                  396                  405                  414                  423                  432

S A S T K G P S V F P L A P S S K S T S G  
 TCA GCT AGC ACC AAG GGC CCA TCG GTC TTC CCC CTG GCA CCC TCC TCC AAG AGC ACC TCT GGG  
           441                  450                  459                  468                  477                  486                  495

G T A A L G C L V K D Y F P E P V T V S W  
 GGC ACA GCG GCC CTG GGC TGC CTG GTC AAG GAC TAC TTC CCC GAA CCG GTG ACG GTG TCG TGG  
           504                  513                  522                  531                  540                  549                  558

N S G A L T S G V H T F P A V L Q S S G L  
 AAC TCA GGC GCC CTG ACC AGC GGC GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC TCA GGA CTC  
           567                  576                  585                  594                  603                  612                  621

Y S L S S V V T V P S S S L G T Q T Y I C  
 TAC TCC CTC AGC AGC GTG GTG ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG ACC TAC ATC TGC  
           630                  639                  648                  657                  666                  675                  684

N V N H K P S N T K V D K K A E P K S C D  
 AAC GTG AAT CAC AAG CCC AGC AAC ACC AAG GTG GAC AAG AAA GCA GAG CCC AAA TCT TGT GAC  
           693                  702                  711                  720                  729                  738                  747

K T H T C P P C P A P E L L G G P S V F L  
 AAA ACT CAC ACA TGC CCA CCG TGC CCA GCA CCT GAA CTC CTG GGG GGA CCG TCA GTC TTC CTC  
           756                  765                  774                  783                  792                  801                  810

F P P K P K D T L M I S R T P E V T C V V  
 TTC CCC CCA AAA CCC AAG GAC ACC CTC ATG ATC TCC CGG ACC CCT GAG GTC ACA TGC GTG GTG  
           819                  828                  837                  846                  855                  864                  873

V D V S H E D P E V K F N W Y V D G V E V  
 GTG GAC GTG AGC CAC GAA GAC CCT GAG GTC AAG TTC AAC TGG TAC GTG GAC GGC GTG GAG GTG

FIG. 10b-1

15 / 15

882	891	900	909	918	927	936
H N A K T K P R E E Q Y N S T Y R V V S V						
CAT AAT GCC AAG ACA AAG CCG CGG GAG GAG CAG TAC AAC AGC ACG TAC CGT GTG GTC AGC GTC						
945 954 963 972 981 990 999						
L T V L H Q D W L N G K E Y K C K V S N K						
CTC ACC GTC CTG CAC CAG GAC TGG CTG AAT GGC AAG GAG TAC AAG TGC AAG GTC TCC AAC AAA						
1008 1017 1026 1035 1044 1053 1062						
A L P A P I E K T I S K A K G Q P R E P Q						
GCC CTC CCA GCC CCC ATC GAG AAA ACC ATC TCC AAA GGC AAA GGG CAG CCC CGA GAA CCA CAG						
1071 1080 1089 1098 1107 1116 1125						
V Y T L P P S R D E L T K N Q V S L T C L						
GTG TAC ACC CTG CCC CCA TCC CGG GAT GAG CTG ACC AAG AAC CAG GTC AGC CTG ACC TGC CTG						
1134 1143 1152 1161 1170 1179 1188						
V K G F Y P S D I A V E W E S N G Q P E N						
GTG AAA GGC TTC TAT CCC AGC GAC ATC GCC GTG GAG TGG GAG AGC AAT GGG CAG CCG GAG AAC						
1197 1206 1215 1224 1233 1242 1251						
N Y K T T P P V L D S D G S F F L Y S K L						
AAC TAC AAG ACC ACG CCT CCC GTG CTG GAC TCC GAC GGC TCC TTC TTC CTC TAC AGC AAG CTC						
1260 1269 1278 1287 1296 1305 1314						
T V D K S R W Q Q G N V F S C S V M H E A						
ACC GTG GAC AAG AGC AGG TGG CAG CAG GGG AAC GTC TTC TCA TGC TCC GTG ATG CAT GAG GCT						
1323 1332 1341 1350 1359 1368 1377						
L H N H Y T Q K S L S L S P G K .						
CTG CAC AAC CAC TAC ACG CAG AAG AGC CTC TCC CTG TCT CCG GGT AAA TGA						
1386 1395 1404 1413 1422 1431						

FIG. 10b-2