

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 April 2002 (25.04.2002)

PCT

(10) International Publication Number
WO 02/32874 A2

(51) International Patent Classification⁷: **C07D 215/20**,
A61K 31/435, A61P 35/00, C07D 215/22, 295/20, 295/14,
401/12

(21) International Application Number: PCT/US01/32422

(22) International Filing Date: 16 October 2001 (16.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/241,127 17 October 2000 (17.10.2000) US
09/740,391 19 December 2000 (19.12.2000) US

(71) Applicant: **THE PROCTER & GAMBLE COMPANY**
[US/US]; One Procter & Gamble Plaza, Cincinnati, OH
45202 (US).

(72) Inventors: **DEGENHARDT, Charles, Raymond**; 10555
Wellington Court, Cincinnati, OH 45240 (US). **EICK-
HOFF, David, Joseph**; 505 Garden Way, Edgewood, KY
41017 (US).

(74) Agents: **REED, T., David** et al.; The Procter & Gam-
ble Company, 5299 Spring Grove Avenue, Cincinnati, OH
45217-1087 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AT
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE
(utility model), DK, DK (utility model), DM, DZ, EC, EE,
EE (utility model), ES, FI, FI (utility model), GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA,
UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: SUBSTITUTED HETEROCYCLIC COMPOUNDS FOR TREATING MULTIDRUG RESISTANCE

(57) Abstract: Substituted heterocyclic compounds for treating multidrug resistance are disclosed. Compositions and methods of use for the substituted heterocyclic compounds are disclosed. Suitable substituted heterocyclic compounds include:



WO 02/32874 A2

SUBSTITUTED HETEROCYCLIC COMPOUNDS FOR TREATING MULTIDRUG RESISTANCE

FIELD OF THE INVENTION

5 This invention relates to compounds for treating multidrug resistance and methods for their preparation and use. More particularly, this invention relates to compounds that regulate the cellular transport proteins P-glycoprotein or MRP1, or both, which are the proteins believed to be largely responsible for causing multidrug resistance in cancer patients.

BACKGROUND OF THE INVENTION

10 “Drug resistance” means a circumstance when a disease (e.g., cancer) does not respond to a therapeutic agent. Drug resistance can be intrinsic, which means that the disease has never been responsive to the therapeutic agent, or acquired, which means that the disease ceases responding to the agent or agents to which the disease had previously been responsive. “Multidrug resistance” is a type of drug resistance wherein a disease is resistant to a variety of drugs that can be functionally unrelated, structurally unrelated, or both. Multidrug resistance is a problem associated with cancer and other conditions, such as bacterial, viral, protozoal, and fungal diseases.

20 One cause of multidrug resistance in cancer patients is that many cancer cells express high levels of the transmembrane transport proteins, such as Pleiotropic-glycoprotein (also known as Pgp, P-glycoprotein, gp-170, or MDR1) and MRP1 (see Borst, P., “Multidrug resistance: A solvable problem?” *Annals of Oncology*, 10, suppl. 4, pp. S162-S164 (1999)). In adenosine-triphosphate driven processes, these transport proteins export hydrophobic compounds (such as vinblastine, daunorubicin, doxorubicin, etoposide, vincristine, and TAXOL®, which are cytotoxic drugs useful for treating cancer) from the cell in an effort to protect the cell from harm. The transport proteins remove the compounds from the cell prior to their having a lethal effect on the cell (see Legrand, et. al, “Simultaneous Activity of MRP1 and Pgp Is Correlated With In Vitro
25 Resistance to Daunorubicin and With In Vivo Resistance in Adult Acute Myeloid
30

Leukemia", *Blood*, Vol. 94, No. 3, pp. 1046-1056 (1999); and Zhu, B.T.; "A Novel Hypothesis for the Mechanism of Action of P-glycoprotein as a Multidrug Transporter," *Molecular Carcinogenesis* 25, pp.1-14 (1999)). Although it is not currently known which of these two classes of proteins is more important for multidrug resistance, and indeed it may be that the class (or classes) of protein which is important depends on the type of cancer and the particular drug or drugs used to treat the cancer, Pgp is known to be highly expressed in approximately 50% of human cancers which require drug therapy. Consequently, Pgp is believed to be a major cause of multidrug resistance.

Other types of multidrug resistance, such as antibacterial, antiviral, and antifungal multidrug resistance may also be caused by the action of transport proteins that are similar to Pgp, and others (see "Annual Reports on Medicinal Chemistry - 33; Section III Cancer and Infectious Diseases" ed. Plattner, J., Academic Press, Ch. 12, pp. 121 - 130 (1998)).

Furthermore, Pgp is also expressed at high levels in the gastrointestinal tract, liver, kidneys, and brain, and therefore Pgp represents a major pharmacological barrier to the bioavailability of many drugs (see Amudkar, et. al in "Biochemical, Cellular, and Pharmacological Aspects of the Multidrug Transporter," *Annu. Rev. Pharmacol. Toxicol.*, 39, pp. 361-398 (1999)). For example, the oral bioavailability of many nutrients and drugs is negatively affected by the action of Pgp present in the gastrointestinal tract. "Oral bioavailability" means the ability of a drug or nutrient that is administered orally to be transported across the gastrointestinal tract and enter into the bloodstream. In addition, Pgp adversely affects penetration of many drugs through the blood-brain barrier.

SUMMARY OF THE INVENTION

This invention relates to novel compounds useful in treating or preventing multidrug resistance ("MDR"). More specifically, these compounds are useful in treating or preventing P-glycoprotein-mediated MDR and MRP1-mediated MDR. This invention further relates to compositions comprising these compounds. This invention further relates to methods for the preparation and use of the compounds and compositions. The compounds and compositions of this invention are well suited for treatment of multidrug

resistant cells, for prevention of the development of multidrug resistance, and for use in multidrug resistant chemotherapies.

DETAILED DESCRIPTION OF THE INVENTION

5 Publications and patents are referred to throughout this disclosure. All U.S. Patents cited herein are hereby incorporated by reference.

All percentages, ratios, and proportions used herein are by weight unless otherwise specified.

Definitions and Usage of Terms

10 The following is a list of definitions, as used herein.

“Aromatic group” means a group having a monocyclic or polycyclic ring structure. Monocyclic aromatic groups contain 4 to 10 carbon atoms, preferably 4 to 7 carbon atoms, and more preferably 4 to 6 carbon atoms in the ring. Preferred polycyclic ring structures have two or three rings. Polycyclic structures having two rings typically have 8
15 to 12 carbon atoms, preferably 8 to 10 carbon atoms in the rings. Polycyclic aromatic groups include groups wherein at least one, but not all, of the rings are aromatic.

“Carbocyclic group” means a saturated or unsaturated hydrocarbon ring. Carbocyclic groups are not aromatic. Carbocyclic groups are monocyclic or polycyclic. Polycyclic carbocyclic groups can be fused, spiro, or bridged ring systems. Monocyclic
20 carbocyclic groups contain 4 to 10 carbon atoms, preferably 4 to 7 carbon atoms, and more preferably 5 to 6 carbon atoms in the ring. Bicyclic carbocyclic groups contain 8 to 12 carbon atoms, preferably 9 to 10 carbon atoms in the rings.

“Carrier” means one or more substances that are suitable for administration to a subject (i.e., mammal) and that can be combined with the active compound according to
25 this invention. Carrier includes solid and liquid diluents, hydrotropes, surface-active agents, and encapsulating substances.

“Chemosensitizing agent” means a noncytotoxic compound that sensitizes drug resistant cells to the action of cytotoxic drugs. As used in this application, the term “chemosensitizing agent”, excludes the active compounds of this invention.

30 “Halogen atom” means F, Cl, Br, or I.

“Heteroaromatic group” means an aromatic group containing carbon and 1 to 4 heteroatoms in the ring. Monocyclic heteroaromatic groups contain 4 to 10 member atoms, preferably 4 to 7 member atoms, and more preferably 4 to 6 member atoms in the ring. Preferred polycyclic ring structures have two or three rings. Polycyclic structures having two rings typically have 8 to 12 member atoms, preferably 8 to 10 member atoms in the rings. Polycyclic heteroaromatic groups include groups wherein at least one, but not all, of the rings are heteroaromatic.

“Heteroatom” means an atom other than carbon e.g., in the ring of a heterocyclic group or the chain of a heterogeneous group. Preferably, heteroatoms are selected from the group consisting of sulfur, phosphorous, nitrogen and oxygen atoms. Groups containing more than one heteroatom may contain different heteroatoms.

“Heterocyclic group” means a saturated or unsaturated ring structure containing carbon atoms and 1 or more heteroatoms in the ring. Heterocyclic groups are not aromatic. Heterocyclic groups are monocyclic or polycyclic. Polycyclic heteroaromatic groups can be fused, spiro, or bridged ring systems. Monocyclic heterocyclic groups contain 4 to 10 member atoms (i.e., including both carbon atoms and at least 1 heteroatom), preferably 4 to 7, and more preferably 5 to 6 in the ring. Bicyclic heterocyclic groups contain 8 to 18 member atoms, preferably 9 or 10 in the rings.

“Heterogeneous group” means a saturated or unsaturated chain of non-hydrogen member atoms comprising carbon atoms and at least one heteroatom. Heterogeneous groups typically have 1 to 25 member atoms. Preferably, the chain contains 1 to 12 member atoms, more preferably 1 to 10, and most preferably 1 to 6. The chain may be linear or branched. Preferred branched heterogeneous groups have one or two branches, preferably one branch. Preferred heterogeneous groups are saturated. Unsaturated heterogeneous groups have one or more double bonds, one or more triple bonds, or both. Preferred unsaturated heterogeneous groups have one or two double bonds or one triple bond. More preferably, the unsaturated heterogeneous group has one double bond.

“Hydrocarbon group” means a chain of 1 to 25 carbon atoms, preferably 1 to 12 carbon atoms, more preferably 1 to 10 carbon atoms, and most preferably 1 to 8 carbon atoms. Hydrocarbon groups may have a linear or branched chain structure. Preferred

hydrocarbon groups have one or two branches, preferably 1 branch. Preferred hydrocarbon groups are saturated. Unsaturated hydrocarbon groups have one or more double bonds, one or more triple bonds, or combinations thereof. Preferred unsaturated hydrocarbon groups have one or two double bonds or one triple bond; more preferred unsaturated hydrocarbon groups have one double bond.

“IC₅₀” means concentration of drug required to produce a 50% inhibition of growth of cancer cells or 50% inhibition of activity.

“MDR” means multidrug resistance.

“Parenteral” as used herein includes subcutaneous, intravenous, intramuscular, intraarticular, intrasynovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques.

“Pgp” means P-glycoprotein.

“Pharmaceutically acceptable” means suitable for use in a human or other mammal.

“Protecting group” is a group that replaces the active hydrogen of a -OH, -COOH, or -NH₂ moiety thus preventing undesired side reaction at the moiety. Use of protecting groups in organic synthesis is well known in the art. Examples of protecting groups are found in Protecting Groups in Organic Synthesis by Greene, T. W. and Wuts, P. G. M., 2nd ed., Wiley & Sons, Inc., 1991. Preferred protecting groups for hydroxyl moieties include silyl ethers, alkoxymethyl ethers, tetrahydropyranyl, tetrahydrofuranyl, esters, and substituted or unsubstituted benzyl ethers. Other preferred protecting groups include carbamates.

“Subject” means a living vertebrate animal such as a mammal (preferably human).

“Substituted aromatic group” means an aromatic group wherein 1 or more of the hydrogen atoms bonded to carbon atoms in the ring have been replaced with other substituents. Preferred substituents include hydrocarbon groups such as methyl groups and heterogeneous groups including alkoxy groups such as methoxy groups. The substituents may be substituted at the ortho, meta, or para position on the ring, or any combination thereof.

“Substituted carbocyclic group” means a carbocyclic group wherein 1 or more hydrogen atoms bonded to carbon atoms in the ring have been replaced with other substituents. Preferred substituents include hydrocarbon groups such as alkyl groups (e.g., methyl groups) and heterogeneous groups such as alkoxy groups (e.g., methoxy groups).

5 “Substituted heteroaromatic group” means a heteroaromatic group wherein 1 or more hydrogen atoms bonded to carbon atoms in the ring have been replaced with other substituents. Preferred substituents include monovalent hydrocarbon groups including alkyl groups such as methyl groups and monovalent heterogeneous groups including alkoxy groups such as methoxy groups.

10 “Substituted heterocyclic group” means a heterocyclic group wherein 1 or more hydrogen atoms bonded to carbon atoms in the ring have been replaced with other substituents. Preferred substituents include monovalent hydrocarbon groups including alkyl groups such as methyl groups and monovalent heterogeneous groups including alkoxy groups such as methoxy groups. Substituted heterocyclic groups are not aromatic.

15 “Substituted heterogeneous group” means a heterogeneous group, wherein 1 or more of the hydrogen atoms bonded to carbon atoms in the chain have been replaced with other substituents. Preferred substituents include monovalent hydrocarbon groups including alkyl groups such as methyl groups and monovalent heterogeneous groups including alkoxy groups such as methoxy groups.

20 “Substituted hydrocarbon group” means a hydrocarbon group wherein 1 or more of the hydrogen atoms bonded to carbon atoms in the chain have been replaced with other substituents. Preferred substituents include monovalent aromatic groups, monovalent substituted aromatic groups, monovalent hydrocarbon groups including alkyl groups such as methyl groups, monovalent substituted hydrocarbon groups such as benzyl, and
25 monovalent heterogeneous groups including alkoxy groups such as methoxy groups. Substituted hydrocarbon groups include groups wherein a $-\text{CH}_2-$ is changed to $-\text{C}(\text{O})-$.

“Substrate potential” means the likelihood that a compound for use in treating multidrug resistance will be transported out of a cell by cellular transport proteins before effectively preventing or reversing multidrug resistance.

“Transport protein” means a protein that acts to remove cytotoxic substances from cells through the cell membrane. Transport protein includes P-glycoprotein, MRP1, and others.

“Treating multidrug resistance” means preventing multidrug resistance from developing in nonresistant cells, increasing or restoring sensitivity of multidrug resistant cells to therapeutic or prophylactic agents, or both.


“Treating” means 1) preventing a disease (i.e., causing the clinical symptoms of the disease not to develop), 2) inhibiting the disease (i.e., arresting the development of clinical symptoms of the disease), 3) relieving the disease (i.e., causing regression of the clinical symptoms), and combinations thereof.

“Wax” means a lower-melting organic mixture or compound of high molecular weight, solid at room temperature and generally similar in formulation to fats and oils except that they contain no glycerides.

15

Active Compounds Used in this Invention

The active compounds of this invention can have the structure:




wherein s is an integer from about 1 to about 3. In a preferred embodiment of the invention, s is 1.

20

A is selected from the group consisting of A¹ and A³. Preferably, A is A¹.

A¹ is a group of the formula:




wherein --- denotes a point of attachment.

Each R^1 is independently selected from the group consisting of a hydrogen atom, a hydroxyl group, a hydrocarbon group, a substituted hydrocarbon group, a heterogeneous group, a substituted heterogeneous group, a carbocyclic group, a substituted carbocyclic group, a heterocyclic group, a substituted heterocyclic group, an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group. Preferably, R^1 is a hydrogen atom or a hydroxyl group.


R^2 is selected from the group consisting of a hydrocarbon group, a substituted hydrocarbon group, a heterogeneous group, a substituted heterogeneous group, a carbocyclic group, a substituted carbocyclic group, a heterocyclic group, a substituted heterocyclic group, an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group.

In a preferred embodiment of the invention, R^2 is selected from the group consisting of an aromatic group; a substituted aromatic group; a heteroaromatic group; a substituted heteroaromatic group; a substituted hydrocarbon group, wherein the substituted hydrocarbon group is substituted with a group selected from the group consisting of an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group; and a substituted heterogeneous group, wherein the substituted heterogeneous group is substituted with a group selected from the group consisting of an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group. More preferably, R^2 is a substituted hydrocarbon group or a substituted heterogeneous group, wherein said group is substituted with a group selected from the group consisting of an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group.

In a more preferred embodiment of the invention, R^2 is selected from the group consisting of:



and



, wherein

wherein a is at least about 2, b is at least about 2, c is about 1 to about 3, and d is about 1 to about 3. Preferably, a and b are each about 3 to about 10. More preferably, a and b are each about 3.

R^{12} and R^{13} are each independently selected from the group consisting of hydrocarbon groups and substituted hydrocarbon groups. Preferably, R^{12} and R^{13} are substituted hydrocarbon groups such as alkoxy groups. Preferred alkoxy groups include methoxy, ethoxy, propoxy, and butoxy.

Each R^{14} is independently selected from the group consisting of CH and a heteroatom. Preferably, the heteroatom is nitrogen. More preferably, each R^{14} is CH.

D^1 and D^2 are each independently selected from the group consisting of $-C(O)-$ and $-NR^3-$, wherein R^3 is selected from the group consisting of a hydrogen atom and R^2 , and with the proviso that optionally, R^2 and R^3 may be bonded together thereby forming a ring structure selected from the group consisting of heterocyclic groups and substituted heterocyclic groups when D^2 is $-NR^3-$.

The ring structure typically has about 4 to about 9 members, preferably 4 to 7 members, more preferably 5 to 6 members. The ring structure is preferably a substituted heterocyclic group, such as substituted piperidyl or substituted piperazinyl. The substituted heterocyclic group is preferably substituted with a group selected from the group consisting of an aromatic group; a substituted aromatic group; a heteroaromatic group; a substituted heteroaromatic group; a substituted hydrocarbon group, wherein the substituted hydrocarbon group is substituted with a group selected from the group consisting of an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group; and a substituted heterogenous group, wherein the substituted heterogenous group is substituted with a group selected from the group consisting of an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group.


In a preferred embodiment of the invention, D^1 is $-C(O)-$ and D^2 is $-NR^3-$. In this embodiment, preferably, R^3 is hydrogen or a hydrocarbon group.

In an alternative embodiment of the invention, D^1 is $-NR^3-$ and D^2 is $-C(O)-$. In this embodiment, preferably, R^3 is hydrogen or a hydrocarbon group.

In the formula above, x is 0 to about 10, y is 0 or 1, and z is 0 or 1. However, when y is 0; then z is 1. When y is 1; then z is 0. When y is 0 and D^1 is $-NR^3-$; then D^2 is $-C(O)-$. When y is 0 and D^2 is $-NR^3-$; then D^1 is $-C(O)-$. When x is 0, D^1 is $-C(O)-$, y is 0, D^2 is $-NR^3-$, and D^3 is $-C(O)-$ or $-S(O)_2-$; then R^2 is selected from the group consisting of a hydrocarbon group and a substituted hydrocarbon group, wherein the substituted hydrocarbon group is substituted with an aromatic group.

In an alternative embodiment of the invention, D^1 is $-C(O)-$, y is 1, and z is 0.

A^2 is selected from the group consisting of a hydrogen atom and groups of the formula




wherein u is 0 to about 10, p is 0 to about 10, and v is 0 or 1. Preferably, u is about 1 to about 3. Preferably, p is about 1 to about 3. Preferably, v is 1. More preferably, u is about 1 and p is about 1.

D^3 is selected from the group consisting of $-S(O)_2-$, $-C(O)-$, and $-CR^1(OH)-$. D^3 is preferably $-CR^1(OH)-$. However, when D^3 is $-S(O)_2-$, then D^1 is $-C(O)-$, D^2 is $-NR^3-$, and R^2 and R^3 are bonded together to form the ring structure.


R^5 is selected from the group consisting of a hydrocarbon group, a substituted hydrocarbon group, a heterogeneous group, a substituted heterogeneous group, a carbocyclic group, a substituted carbocyclic group, a heterocyclic group, a substituted heterocyclic group, an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group. However, when D^3 is $-C(O)-$, v is 0, and R^5 contains a $-C(O)-$ group, then p is not 0. Preferably, R^5 is selected from the group consisting of substituted hydrocarbon groups of at least 2 carbon atoms and substituted heterogeneous groups of at least 2 member atoms, wherein R^5 is substituted with at least one group selected from the group consisting of aromatic groups, heteroaromatic groups, substituted aromatic groups, and substituted heteroaromatic groups;

In a preferred embodiment of the invention, u is 0, and R⁵ is selected from the group consisting of an aromatic group; a substituted aromatic group; a heteroaromatic group; a substituted heteroaromatic group; a hydrocarbon group; a substituted hydrocarbon group, wherein the substituted hydrocarbon group is substituted with a group selected from the group consisting of an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group; and a substituted heterogenous group, wherein the substituted heterogenous group is substituted with a group selected from the group consisting of an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group. In one embodiment, preferably p is 0, and D³ is -SO₂-. In a more preferred embodiment, D³ is -CR¹(OH)- and R⁵ is a heteroaromatic group of the formula:



wherein each X is independently selected from the group consisting of CH and a heteroatom, with the proviso that at least one X is a heteroatom. The heteroatom is preferably nitrogen. Preferably, one X is a heteroatom. Examples of heteroaromatic groups for R⁵ include quinolyl and isoquinolyl groups. Preferred quinolyl groups for R⁵ include 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, and 8-quinolyl. More preferably, R⁵ is 5-quinolyl.

A³ has the formula



wherein t is about 1 to about 6. Preferably in A³, at least one R¹ is a hydroxyl group.

D⁴ is selected from the group consisting of -C(O)- and -CHR¹-.

D⁵ is selected from the group consisting of -NR⁶(R⁷), -O_rR⁶, and -C(O)R⁶

wherein r is 0 or 1.

R^6 is selected from the group consisting of a hydrocarbon group, a substituted hydrocarbon group, a heterogeneous group, a substituted heterogeneous group, a carbocyclic group, a substituted carbocyclic group, a heterocyclic group, a substituted heterocyclic group, an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group.

R^7 is selected from the group consisting of a hydrogen atom and R^6 , with the proviso that when u is 0, D^3 and D^4 are not both $-C(O)-$.

In a preferred embodiment of the invention, D^4 is $-C(O)-$, t is 0, and D^5 is $-C(O)R^6$.

10 In an alternative preferred embodiment of the invention, D^4 is $-C(O)-$ and D^5 is $-O_rR^6$.

In an alternative preferred embodiment of the invention, D^4 is $-CH(R^1)-$ and D^5 is $-O_rR^6$.







15 In an alternative preferred embodiment of the invention, D^4 is $-CH(R^1)-$ and D^5 is $-NR^6(R^7)-$.

In an alternative preferred embodiment of the invention, D^4 is $-C(O)-$ and D^5 is $-NR^6(R^7)$.

20 In an alternative embodiment of the invention, the active compound can be an optical isomer, a diastereomer, an enantiomer, a pharmaceutically-acceptable salt, a biohydrolyzable amide, a biohydrolyzable ester, and a biohydrolyzable imide of the structure.







25 In one embodiment of the invention, the active compound has the structure above wherein A is A^1 , R^2 and R^3 are bonded together and the ring structure has 4 to 6 members in the ring. Examples of compounds according to this embodiment of the invention are shown below in Table 1.

Table 1


	
	
	

In a preferred embodiment of the invention, the active compound has the structure above wherein A is A¹, D¹ is -C(O)-, and D² is -NR³- where R² and R³ are not bonded together in a ring structure. Examples of compounds according to this embodiment of the invention are shown below in Table 2.

Table 2

In an alternative embodiment of the invention, the active compound has the structure above wherein A is A³. An example compound according to this embodiment is shown below in Table 3.



The active compound of this invention inhibits at least one transport protein. The active compound preferably inhibits Pgp or MRP1. More preferably, the active compound inhibits both Pgp and MRP1. In a preferred embodiment of this invention, the active compound inhibits Pgp and has low substrate potential for Pgp. In an alternative preferred embodiment, the active compound inhibits MRP1 and has low substrate potential for MRP1. In the most preferred embodiment of this invention, the active compound inhibits both Pgp and MRP1 and the active compound has low substrate potential for both Pgp and MRP1.

The degree to which a compound inhibits a transport protein can be measured by quantitating the effectiveness of the compound toward restoring drug sensitivity to multidrug resistant cells. Methods for quantitating the effectiveness of the active compounds toward restoring drug sensitivity are readily available to one skilled in the art without undue experimentation (see U.S. Patent Nos. 5,935,954 and 5,272,159, which are hereby incorporated by reference for the purpose of disclosing these methods). Any assay known to measure the restoration of the anti-proliferative activity of a drug may be employed to test the compounds of this invention. These assays use cell lines resistant to particular drugs, and characterized by the presence of one or both of Pgp and MRP1. These cell lines include L1210, HL60, P388, CHO, and MCF7. Alternatively, resistant cell lines can be developed by methods readily available to one of ordinary skill in the art without undue experimentation (see Chaudhary, et al., "Induction of Multidrug Resistance

in Human Cells by Transient Exposure to Different Chemotherapeutic Agents,” *Journal of the National Cancer Institute*, Vol. 85, No. 8, pp. 632-639 (1993)). The cell line is then exposed to compounds of this invention in the presence or absence of the drug to which it is resistant, such as TAXOL®. The viability of the cells treated with both the active
5 compound and the drug can then be compared to the viability of the cells treated only with the drug.

The active compound preferably also has low substrate potential for Pgp or MRP1. More preferably, the active compound has low substrate potential for both Pgp and MRP1. Substrate potential for a transport protein can be determined by using an assay for
10 measuring ATPase activity of the Pgp or MRP1 pumps (see, for example, Reference Example 4, below).

Methods for quantitating accumulation of the active compounds are readily available to one skilled in the art without undue experimentation (see U.S. 5,272,159 which is hereby incorporated by reference for the purpose of disclosing assays for
15 quantitating accumulation). These assays use cell lines resistant to particular chemotherapeutic agents, and characterized by the presence of one or both of Pgp and MRP1. The cell line is exposed to a labeled form of the active compound (e.g., radioactivity or fluorescence labeling) and the accumulation of the active compound is monitored over time. The amount of active compound accumulated in the cell can be
20 compared with a compound which is readily transported by these proteins, e.g. labeled TAXOL®.

Compositions of this Invention

This invention further relates to a composition. The composition can be used for
25 treating various conditions or disease states. The composition is preferably a pharmaceutical composition administered for treatment or prevention of multidrug resistance. Standard pharmaceutical formulation techniques are used, such as those disclosed in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA. (1990) and U.S. Patent No. 5,091,187, which is hereby incorporated by reference.

The composition comprises component (A) the active compound described above and component (B) a carrier. The composition may further comprise component (C) an optional ingredient, such as a therapeutic agent.

Component (B) is a carrier. A carrier is one or more compatible substances that are suitable for administration to a mammal. "Compatible" means that the components of the composition are capable of being commingled with component (A), and with each other, in a manner such that there is no interaction which would substantially reduce the efficacy of the composition under ordinary use situations. Carriers must be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the mammal being treated. The carrier can be inert, or it can possess pharmaceutical benefits, cosmetic benefits, or both, depending on the intended use as described herein.

The choice of carrier for component (B) depends on the route by which component (A) will be administered and the form of the composition. The composition may be in a variety of forms, suitable, for example, for systemic administration (e.g., oral, rectal, nasal, sublingual, buccal, or parenteral) or topical administration (e.g., local application on the skin, ocular, liposome delivery systems, or iontophoresis).

Systemic Compositions

Carriers for systemic administration typically comprise one or more ingredients selected from the group consisting of a) diluents, b) lubricants, c) binders, d) disintegrants, e) colorants, f) flavors, g) sweeteners, h) antioxidants, j) preservatives, k) glidants, m) solvents, n) suspending agents, o) surfactants, combinations thereof, and others.

Ingredient a) is a diluent. Suitable diluents include sugars such as glucose, lactose, dextrose, and sucrose; polyols such as propylene glycol; calcium carbonate; sodium carbonate; glycerin; mannitol; sorbitol; and maltodextrin. The amount of ingredient a) in the composition is typically about 1 to about 99 %.

Ingredient b) is a lubricant. Suitable lubricants are exemplified by solid lubricants including silica, talc, stearic acid and its magnesium salts and calcium salts, calcium sulfate; and liquid lubricants such as polyethylene glycol and vegetable oils such as

peanut oil, cottonseed oil, sesame oil, olive oil, corn oil, and oil of theobroma. The amount of ingredient b) in the composition is typically about 1 to about 99 %.

Ingredient c) is a binder. Suitable binders include polyvinylpyrrolidone; magnesium aluminum silicate; starches such as corn starch and potato starch; gelatin; 5 tragacanth; and cellulose and its derivatives, such as sodium carboxymethylcellulose, ethylcellulose, methylcellulose, microcrystalline cellulose, and hydroxypropylmethylcellulose; carbomer; providone; acacia; guar gum; and xanthan gum. The amount of ingredient c) in the composition is typically about 1 to about 99 %.

Ingredient d) is a disintegrant. Suitable disintegrants include agar, alginic acid and 10 the sodium salt thereof, effervescent mixtures, croscarmellose, crospovidone, sodium carboxymethyl starch, sodium starch glycolate, clays, and ion exchange resins. The amount of ingredient d) in the composition is typically about 1 to about 99 %.

Ingredient e) is a colorant such as an FD&C dye. The amount of ingredient e) in the composition is typically about 1 to about 99 %.

15 Ingredient f) is a flavor such as menthol, peppermint, and fruit flavors. The amount of ingredient f) in the composition is typically about 1 to about 99 %.

Ingredient g) is a sweetener such as saccharin and aspartame. The amount of ingredient g) in the composition is typically about 1 to about 99 %.

Ingredient h) is an antioxidant such as butylated hydroxyanisole, butylated 20 hydroxytoluene, and vitamin E. The amount of ingredient h) in the composition is typically about 1 to about 99 %.

Ingredient j) is a preservative such as phenol, alkyl esters of parahydroxybenzoic acid, benzoic acid and the salts thereof, boric acid and the salts thereof, sorbic acid and the salts thereof, chlorbutanol, benzyl alcohol, thimerosal, phenylmercuric acetate and 25 nitrate, nitromersol, benzalkonium chloride, cetylpyridinium chloride, methyl paraben, ethyl paraben, and propyl paraben. Particularly preferred are the salts of benzoic acid, cetylpyridinium chloride, methyl paraben and propyl paraben, and sodium benzoate. The amount of ingredient j) in the composition is typically about 1 to about 99 %.

Ingredient k) is a glidant such as silicon dioxide. The amount of ingredient k) in 30 the composition is typically about 1 to about 99 %.

Ingredient m) is a solvent, such as water, isotonic saline, ethyl oleate, alcohols such as ethanol, glycerin, cremaphor, glycols (e.g., polypropylene glycol and polyethylene glycol), and buffer solutions (e.g., phosphate, potassium acetate, boric carbonic, phosphoric, succinic, malic, tartaric, citric, acetic, benzoic, lactic, glyceric, gluconic, glutaric, and glutamic). The amount of ingredient m) in the composition is typically about 1 to about 99 %.

Ingredient n) is a suspending agent. Suitable suspending agents include AVICEL® RC-591 from FMC Corporation of Philadelphia, Pennsylvania and sodium alginate. The amount of ingredient n) in the composition is typically about 1 to about 99 %.

Ingredient o) is a surfactant such as lecithin, polysorbate 80, sodium lauryl sulfate, polyoxyethylene sorbitan fatty acid esters, polyoxyethylene monoalkyl ethers, sucrose monoesters, lanolin esters, and lanolin ethers. Suitable surfactants are known in the art and commercially available, e.g., the TWEENS® from Atlas Powder Company of Wilmington, Delaware. Suitable surfactants are disclosed in the C.T.F.A. Cosmetic Ingredient Handbook, pp. 587-592 (1992); Remington's Pharmaceutical Sciences, 15th Ed., pp. 335-337 (1975); and McCutcheon's Volume 1, Emulsifiers & Detergents, North American Edition, pp. 236-239 (1994). The amount of ingredient o) in the composition is typically about 1 to about 99%.

The carrier ingredients discussed above are exemplary and not limiting. One skilled in the art would recognize that different carrier ingredients may be added to or substituted for the carrier ingredients above. One skilled in the art would be able to select appropriate carrier ingredients for systemic compositions without undue experimentation.

Compositions for parenteral administration typically comprise (A) about 0.1 to about 10% of an active compound and (B) about 90 to about 99.9% of a carrier comprising a) a diluent and m) a solvent. Preferably, component a) is propylene glycol and m) is selected from the group consisting of ethanol, ethyl oleate, water, isotonic saline, and combinations thereof.

Compositions for oral administration can have various dosage forms. For example, solid forms include tablets, capsules, granules, and bulk powders. These oral dosage

forms comprise a safe and effective amount, usually at least about 1%, and preferably from about 5% to about 50%, of component (A). The oral dosage compositions further comprise (B) about 50 to about 99% of a carrier, preferably about 50 to about 95%.

Tablets can be compressed, tablet triturates, enteric-coated, sugar-coated, film-coated, or multiple-compressed. Tablets typically comprise (A) the active compound, and
5 (B) a carrier comprising ingredients selected from the group consisting of a) diluents, b) lubricants, c) binders, d) disintegrants, e) colorants, f) flavors, g) sweeteners, k) glidants, and combinations thereof. Preferred diluents include calcium carbonate, sodium carbonate, mannitol, lactose, and sucrose. Preferred binders include starch, and gelatin.
10 Preferred disintegrants include alginic acid, and croscarmellose. Preferred lubricants include magnesium stearate, stearic acid, and talc. Preferred colorants are the FD&C dyes, which can be added for appearance. Chewable tablets preferably contain g) sweeteners such as aspartame and saccharin or f) flavors such as menthol, peppermint, and fruit flavors, or both.

15 Capsules (including time release and sustained release compositions) typically comprise (A) the active compound and (B) the carrier comprising one or more a) diluents disclosed above in a capsule comprising gelatin. Granules typically comprise (A) the active compound, and preferably further comprise k) glidants such as silicon dioxide to improve flow characteristics.

20 The selection of ingredients in the carrier for oral compositions depends on secondary considerations like taste, cost, and shelf stability, which are not critical for the purposes of this invention. One skilled in the art can optimize appropriate ingredients without undue experimentation.

The solid compositions may also be coated by conventional methods, typically
25 with pH or time-dependent coatings, such that component (A) is released in the gastrointestinal tract at various times to extend the desired action. The coatings typically comprise one or more components selected from the group consisting of cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropyl methyl cellulose phthalate, ethyl cellulose, acrylic resins such as EUDRAGIT® coatings (available from Rohm & Haas
30 G.M.B.H. of Darmstadt, Germany), waxes, shellac, polyvinylpyrrolidone, and other

commercially available film-coating preparations such as Dri-Klear, manufactured by Crompton & Knowles Corp., Mahwah, NJ or OPADRY® manufactured by Colorcon, Inc., of West Point, Pennsylvania.

Compositions for oral administration can also have liquid forms. For example, suitable liquid forms include aqueous solutions, emulsions, suspensions, solutions reconstituted from non-effervescent granules, suspensions reconstituted from non-effervescent granules, effervescent preparations reconstituted from effervescent granules, elixirs, tinctures, syrups, and the like. Liquid orally administered compositions typically comprise (A) the active compound and (B) a carrier comprising ingredients selected from the group consisting of a) diluents, e) colorants, and f) flavors, g) sweeteners, j) preservatives, m) solvents, n) suspending agents, and o) surfactants. *Peroral* liquid compositions preferably comprise one or more ingredients selected from the group consisting of e) colorants, f) flavors, and g) sweeteners.

Other compositions useful for attaining systemic delivery of the active compounds include sublingual, buccal and nasal dosage forms. Such compositions typically comprise one or more of soluble filler substances such as a) diluents including sucrose, sorbitol and mannitol; and c) binders such as acacia, microcrystalline cellulose, carboxymethylcellulose, and hydroxypropylmethylcellulose. Such compositions may further comprise b) lubricants, e) colorants, f) flavors, g) sweeteners, h) antioxidants, and k) glidants.

The composition may further comprise component (C) one or more optional ingredients. Component (C) can be a therapeutic agent used to treat the underlying disease from which the subject suffers. For example, component (C) can be (i) a cancer therapeutic agent, such as a chemotherapeutic agent or a chemosensitizing agent, or a combination thereof; (ii) an antibacterial agent, (iii) an antiviral agent, (iv) an antifungal agent, and combinations thereof. Component (C) can be coadministered with component (A) to increase the susceptibility of the multidrug resistant cells within the subject to the therapeutic agent.

Suitable (i) cancer therapeutic agents are known in the art. Cancer therapeutic agents include chemotherapeutic agents, chemosensitizing agents, and combinations

thereof. Suitable chemotherapeutic agents are disclosed in U.S. Patent No. 5,416,091, which is hereby incorporated by reference for the purpose of disclosing chemotherapeutic agents. Suitable chemotherapeutic agents include actinomycin D, adriamycin, amsacrine, colchicine, daunorubicin, docetaxel (which is commercially available as TAXOTERE®
5 from Aventis Pharmaceuticals Products, Inc.), doxorubicin, etoposide, mitoxantrone, mytomycin C, paclitaxel (which is commercially available as TAXOL® from Bristol-Myers Squibb Company of New York, NY), tenipaside, vinblastine, vincristine, and combinations thereof.

Suitable chemosensitizing agents include calcium channel blockers, calmodulin
10 antagonists, cyclic peptides, cyclosporins and their analogs, phenothiazines, quinidine, reserpine, steroids, thioxantheres, transflupentixol, trifluoperazine, and combinations thereof. Suitable chemosensitizing agents are disclosed by Amudkar, et. al in "Biochemical, Cellular, and Pharmacological Aspects of the Multidrug Transporter," *Annu. Rev. Pharmacol. Toxicol.*, 39, pp. 361-398 (1999).

15 Suitable (ii) antibacterial agents, (iii) antiviral agents, and (iv) antifungal agents are known in the art (see "Annual Reports on Medicinal Chemistry - 33; Section III Cancer and Infectious Diseases" ed. Plattner, J., Academic Press, Ch. 12, pp. 121 - 130 (1998)). Suitable antibacterial agents include quinolones, fluoroquinolones, β -lactam antibiotics, aminoglycosides, macrolides, glycopeptides, tetracyclines, and combinations
20 thereof.

Suitable (iii) antiviral agents include protease inhibitors, DNA synthase inhibitors, reverse transcription inhibitors, and combinations thereof.

Suitable (iv) antifungal agents include azoles, such as ketoconazole, fluconazole, itraconazole, and combinations thereof.

25 One skilled in the art will recognize that these therapeutic agents are exemplary and not limiting, and that some may be used in the treatment of various multidrug resistant conditions and diseases. One skilled in the art would be able to select therapeutic agents without undue experimentation.

The amount of component (C) used in combination with component (A), whether

included in the same composition or separately coadministered, will be less than or equal to that used in a monotherapy. Preferably, the amount of component (C) is less than 80% of the dosage used in a monotherapy. Monotherapeutic dosages of such agents are known in the art.

5 Component (C) may be part of a single pharmaceutical composition or may be separately administered at a time before, during, or after administration of component (A), or combinations thereof.

 In a preferred embodiment, the composition of this invention comprises component (A), component (B), and (C) a chemotherapeutic agent. In an alternative
10 preferred embodiment, the composition comprises component (A), component (B), and (C) a chemosensitizing agent. In another preferred alternative embodiment, the composition comprises component (A), component (B), and (C) both a chemotherapeutic agent and a chemosensitizing agent.

 The exact amounts of each component in the systemic compositions depend on
15 various factors. These factors include the specific compound selected as component (A), and the mode by which the composition will be administered. The amount of component (A) in the systemic composition is typically about 1 to about 99 %.

 The systemic composition preferably further comprises 0 to 99 % component (C), and a sufficient amount of component (B) such that the amounts of components (A), (B),
20 and (C), combined equal 100%. The amount of (B) the carrier employed in conjunction with component (A) is sufficient to provide a practical quantity of composition for administration per unit dose of the compound. Techniques and compositions for making dosage forms useful in the methods of this invention are described in the following references: Modern Pharmaceutics, Chapters 9 and 10, Banker & Rhodes, eds. (1979);
25 Lieberman et al., Pharmaceutical Dosage Forms: Tablets (1981); and Ansel, Introduction to Pharmaceutical Dosage Forms, 2nd Ed., (1976).

Topical Compositions

 Topical compositions comprise: component (A), described above, and component
30 (B) a carrier. The carrier of the topical composition preferably aids penetration of

component (A) into the skin. Topical compositions preferably further comprise (C) the optional ingredient described above.

Component (B) the carrier may comprise a single ingredient or a combination of two or more ingredients. In the topical compositions, component (B) is a topical carrier.

5 Preferred topical carriers comprise one or more ingredients selected from the group consisting of water, alcohols, aloe vera gel, allantoin, glycerin, vitamin A and E oils, mineral oil, propylene glycol, polypropylene glycol-2 myristyl propionate, dimethyl isosorbide, combinations thereof, and the like. More preferred carriers include propylene glycol, dimethyl isosorbide, and water.

10 The topical carrier may comprise one or more ingredients selected from the group consisting of q) emollients, r) propellants, s) solvents, t) humectants, u) thickeners, v) powders, and w) fragrances in addition to, or instead of, the preferred topical carrier ingredients listed above. One skilled in the art would be able to optimize carrier ingredients for the topical compositions without undue experimentation.

15 Ingredient q) is an emollient. The amount of ingredient q) in the topical composition is typically about 5 to about 95%. Suitable emollients include stearyl alcohol, glyceryl monoricinoleate, glyceryl monostearate, propane-1,2-diol, butane-1,3-diol, mink oil, cetyl alcohol, isopropyl isostearate, stearic acid, isobutyl palmitate, isocetyl stearate, oleyl alcohol, isopropyl laurate, hexyl laurate, decyl oleate, octadecan-2-ol, isocetyl
20 alcohol, cetyl palmitate, di-n-butyl sebacate, isopropyl myristate, isopropyl palmitate, isopropyl stearate, butyl stearate, polyethylene glycol, triethylene glycol, lanolin, sesame oil, coconut oil, arachis oil, castor oil, acetylated lanolin alcohols, petrolatum, mineral oil, butyl myristate, isostearic acid, palmitic acid, isopropyl linoleate, lauryl lactate, myristyl lactate, decyl oleate, myristyl myristate, polydimethylsiloxane, and combinations thereof.
25 Preferred emollients include stearyl alcohol and polydimethylsiloxane.

Ingredient r) is a propellant. The amount of ingredient r) in the topical composition is typically about 5 to about 95%. Suitable propellants include propane, butane, isobutane, dimethyl ether, carbon dioxide, nitrous oxide, nitrogen, and combinations thereof.

Ingredient s) is a solvent. The amount of ingredient s) in the topical composition is typically about 5 to about 95 %. Suitable solvents include water, ethyl alcohol, methylene chloride, isopropanol, castor oil, ethylene glycol monoethyl ether, diethylene glycol monobutyl ether, diethylene glycol monoethyl ether, dimethylsulfoxide, dimethyl
5 formamide, tetrahydrofuran, and combinations thereof. Preferred solvents include ethyl alcohol.

Ingredient t) is a humectant. The amount of ingredient t) in the topical composition is typically about 5 to about 95 %. Suitable humectants include glycerin, sorbitol, sodium 2-pyrrolidone-5-carboxylate, soluble collagen, dibutyl phthalate, gelatin,
10 and combinations thereof. Preferred humectants include glycerin.

Ingredient u) is a thickener. The amount of ingredient u) in the topical composition is typically 0 to about 95%.

Ingredient v) is a powder. The amount of ingredient v) in the topical composition is typically 0 to about 95 %. Suitable powders include chalk, talc, fullers earth, kaolin,
15 starch, gums, colloidal silicon dioxide, sodium polyacrylate, tetraalkyl ammonium smectites, trialkyl aryl ammonium smectites, chemically modified magnesium aluminum silicate, organically modified montmorillonite clay, hydrated aluminum silicate, fumed silica, carboxyvinyl polymer, sodium carboxymethyl cellulose, ethylene glycol monostearate, and combinations thereof.

20 Ingredient w) is a fragrance. The amount of ingredient w) in the topical composition is typically about 0.001 to about 0.5%, preferably about 0.001 to about 0.1%.

Ingredient x) is a wax. Waxes useful in this invention are selected from the group consisting of animal waxes, vegetable waxes, mineral waxes, various fractions of natural waxes, synthetic waxes, petroleum waxes, ethylenic polymers, hydrocarbon types such as
25 Fischer-Tropsch waxes, silicone waxes, and mixtures thereof wherein the waxes have a melting point between 40 and 100°C. The amount of ingredient x) in the topical composition is typically about 1 to about 99%.

In an alternative embodiment of the invention, the active compounds may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles,
30 large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a

variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines. A preferred composition for topical delivery of the present compounds uses liposomes as described in Dowton et al., "Influence of Liposomal Composition on Topical Delivery of Encapsulated Cyclosporin A: I. An *in vitro* Study Using Hairless Mouse Skin", *S.T.P.*

5 *Pharma Sciences*, Vol. 3, pp. 404 - 407 (1993); Wallach and Philippot, "New Type of Lipid Vesicle: Novasome®", *Liposome Technology*, Vol. 1, pp. 141 - 156 (1993); U.S. Patent No. 4,911,928, and U.S. Patent No. 5,834,014.

The exact amounts of each component in the topical composition depend on various factors. Including the specific compound selected for component (A) and the
10 mode by which the composition will be administered. However, the amount of component (A) typically added to the topical composition is about 0.1 to about 99%, preferably about 1 to about 10%.

The topical composition preferably further comprises 0 to about 99% component (C), more preferably 0 to about 10%, and a sufficient amount of component (B) such that
15 the amounts of components (A), (B), and (C), combined equal 100%. The amount of (B) the carrier employed in conjunction with component (A) is sufficient to provide a practical quantity of composition for administration per unit dose of the compound. Techniques and compositions for making dosage forms useful in the methods of this invention are described in the following references: Modern Pharmaceutics, Chapters 9
20 and 10, Banker & Rhodes, eds. (1979); Lieberman et al., Pharmaceutical Dosage Forms: Tablets (1981); and Ansel, Introduction to Pharmaceutical Dosage Forms, 2nd Ed., (1976).

Topical compositions that can be applied locally to the skin may be in any form including solutions, oils, creams, ointments, gels, lotions, shampoos, leave-on and rinse-
25 out hair conditioners, milks, cleansers, moisturizers, sprays, skin patches, and the like. Component (A) may be included in kits comprising component (A), a systemic or topical composition described above, or both; and information, instructions, or both that use of the kit will provide treatment for multidrug resistance (particularly in humans). The information and instructions may be in the form of words, pictures, or both, and the like.
30 In addition or in the alternative, the kit may comprise component (A), a composition, or

both; and information, instructions, or both, regarding methods of administration of component (A) or the composition, preferably with the benefit of treating multidrug resistance in mammals.

5 In an alternative embodiment of the invention, components (A) and (C) may be included in kits comprising components (A) and (C), systemic or topical compositions described above, or both; and information, instructions, or both that use of the kit will provide treatment for multidrug resistance (particularly humans). The information and instructions may be in the form of words, pictures, or both, and the like. In addition or in the alternative, the kit may comprise components (A) and (C), compositions, or both; and 10 information, instructions, or both, regarding methods of administration of components (A) and (C) or the compositions, preferably with the benefit of treating multidrug resistance in mammals.

Methods of Use of the Invention

15 This invention relates to a method for inhibiting transport protein activity. The method comprises administering to a subject (A) an active compound described above.

This invention further relates to a method for treating multidrug resistance. The method comprises administering to a subject (preferably a human) suffering from multidrug resistance, (A) an active compound described above. For example, a subject 20 diagnosed with multidrug resistant cancer can be treated by the methods of this invention. Preferably, a systemic or topical composition comprising (A) the active compound and (B) the carrier is administered to the subject. More preferably, the composition is a systemic composition comprising (A) the active compound, (B) the carrier, and (C) an optional ingredient such as a therapeutic agent. Component (A) may be administered 25 before, during, or after administration of component (C). A preferred administration schedule is a continuous infusion over the 24 hour period during which component (C) is also administered.

The dosage of component (A) administered depends on various factors, including the method of administration, the physical attributes of the subject (e.g., age, weight, and 30 gender), and the condition from which the subject suffers. Effective dosage levels for

treating or preventing MDR range from about 0.01 to about 100 mg/kg body weight per day, preferably about 0.5 to about 50 mg/kg body weight per day of (A) a compound of this invention. These dosage ranges are merely exemplary, and daily administration can be adjusted depending on various factors. The specific dosage of the active compound to be administered, as well as the duration of treatment, and whether the treatment is topical or systemic are interdependent. The dosage and treatment regimen will also depend upon such factors as the specific active compound used, the treatment indication, the efficacy of the active compound, the personal attributes of the subject (such as, for example, weight, age, sex, and medical condition of the subject), compliance with the treatment regimen, and the presence and severity of any side effects of the treatment.

In addition to the benefits in treating multidrug resistance in subjects suffering from cancer, the active compounds in the compositions and methods of this invention can also be used to treat other conditions. These other conditions include other types of multidrug resistance (i.e., in addition to cancer multidrug resistance) such as bacterial, viral, and fungal multidrug resistance. For example, many of the FDA approved HIV protease inhibitors used to treat AIDS patients suffering from the HIV virus are substrates for Pgp. Therefore, in an alternative embodiment of this invention, an active compound of this invention is coadministered with a therapeutic agent such as an HIV protease inhibitor.

The active compounds and compositions of this invention can also be administered with other therapeutic agents such as oral drugs. The active compounds and compositions can be used to enhance oral drug absorption and increase bioavailability of various drugs.

The active compounds and compositions can also be used to aid drug delivery through the blood-brain barrier for, e.g., enhancing the effectiveness of drugs to treat Alzheimer's disease, treating memory disorders, enhancing memory performance, or treating any other central nervous system disorder where drug delivery is compromised via this transport pump mechanism.

The active compounds and compositions can also be administered to treat subjects suffering from neurological disorders such as spinal injuries, diabetic neuropathy, and macular degeneration.

5 The active compounds and compositions can also be administered to treat subjects suffering from vision disorders and to improve vision.

The active compounds and compositions can also be administered to treat hair loss. "Treating hair loss" includes arresting hair loss, reversing hair loss, and promoting hair growth.

10 The active compounds and compositions can also be administered to treat inflammatory diseases. Inflammatory diseases include irritable bowel disease, arthritis, and asthma.

EXAMPLES

These examples are intended to illustrate the invention to those skilled in the art and should not be interpreted as limiting the scope of the invention set forth in the claims.

15 The active compounds of this invention can be made using conventional organic syntheses, which are readily available to one skilled in the art without undue experimentation. Such syntheses can be found in standard texts such as J. March, Advanced Organic Chemistry, John Wiley & Sons, 1992. One of ordinary skill in the art will appreciate that certain reactions are best carried out when other functionalities are
20 masked or protected in the compound, thus increasing the yield of the reaction or avoiding any undesirable side reactions. The skilled artisan may use protecting groups to accomplish the increased yields or to avoid the undesired reactions. These reactions can be found in the literature, see for example, Greene, T.W. and Wuts, P.G.M., Protecting Groups in Organic Synthesis, 2nd ed., John Wiley & Sons, 1991.

25 The starting materials for preparing the compounds of the invention are known, made by known methods, or commercially available. The starting materials for preparing the compounds of the invention may include the following.

The following reagents are available from Aldrich Chemical Company, Milwaukee, WI: 1-bromo-3-phenylpropane, 5-hydroxyquinoline, (R)-(-)-glycidyl tosylate,
30 3,4-pyridinedicarboxylic acid, 4-phenylbutylamine, 3-pyridinepropionic acid, *tert*-butyl[S-

(R*, R*)]-(-)-(1-oxiranyl)-2-phenylethyl)carbamate, epichlorohydrin, 3,4,5-trimethoxybenzoyl chloride, N,N-diisopropylethylamine, 4-dimethylaminopyridine, 1-hydroxybenzotriazole, 4-*trans*-aminomethylcyclohexanecarboxylic acid, 3,4,5-trimethoxybenzylamine, and 2,2,4-trimethyl-2-oxazoline.

5 The following reagents are available from Lancaster Synthesis Inc., Windham, NH: 4-phenylbutyronitrile, 1-*tert*-butoxycarbonyl-piperidine-3-carboxylic acid, 1-benzyl-4-aminopiperidine, 3,4-dimethoxybenzenesulfonyl chloride, and 1-benzyl-4-homopiperazine.

The following reagents are available from Fluka Chemie AG, Milwaukee, WI:
10 1-*tert*-butoxycarbonyl-piperidine-4-carboxylic, and (benzotriazol-1-yl-*oxy*)tripyrrolidinophosphonium hexafluorophosphate ("PyBOP"), N-(*tert*-butoxycarbonyl)-iminodiacetic acid, and 1-(diphenylmethyl)piperazine.

The following reagents are available from Acros Organics, Pittsburgh, PA: quinoline-6-carboxylic acid and quinoline-5-carboxylic acid.

15 The following reagent is available from Bachem Bioscience, King of Prussia, PA: *tert*-butoxycarbonyl- β -(3-pyridyl)-alanine.

The following reagent is available from Sigma Chemical Company, Milwaukee, Wisconsin: N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride.


Various abbreviations are used herein. Abbreviations that can be used and their
20 definitions are shown below in Table 4.

Table 4 - Abbreviations

Abbreviation	Definition
"AM"	acetoxymethyl ester
"Boc"	<i>tert</i> -butoxycarbonyl
"CIMS"	chemical ionization mass spectrometry
"DMF"	dimethylformamide
"ESMS"	electrospray mass spectrometry
"Et"	an ethyl group

"Me"	a methyl group
"MH ⁺ "	parent ion in ESMS
"MS"	mass spectrometry
"MTT"	3-[4,5-dimethyl-thiazoyl-2-yl]2,5-diphenyl-tetrazolium bromide
"NIH"	National Institute of Health
"PBS"	Phosphate-buffered saline
"THF"	tetrahydrofuran

Example 1 - 1,7-Diphenyl-4-aminoheptane hydrochloride (1)




- 5 Magnesium (40.2 g, 1.65 mol) and anhydrous ether (3.2 L) are combined in a reaction vessel with stirring. A solution of 1-bromo-3-phenyl propane in 1.6 L of anhydrous ether is added to an addition funnel. The bromide solution is added dropwise to the stirring reaction vessel over a 1 hour period. Upon completion of addition, the mixture stirs for 1 - 2 hours. A solution of 4-phenylbutyronitrile (160 g, 1.1 mol) in anhydrous ether (2.4 L)
- 10 is placed in the addition funnel. The solution is added to the reaction vessel over a 1 hour time period. Upon complete addition the solution is heated to reflux for 10 hours, and then stirs at room temperature for six hours. The reaction mixture is diluted with methanol (3.2 L) using an addition funnel. Sodium borohydride (83.4 g, 2.2 mol) is added in portions. Upon complete addition the reaction is stirred at room temperature for
- 15 six hours. The reaction mixture is quenched by a slow addition of water (3.2 L). The mixture is diluted with ether (3.2 L) and water (1.6 L). The ether layer is separated and the aqueous layer is extracted twice with ether (3.2 L x 2). The combined ether extracts are washed once with sodium chloride solution, dried, filtered, and concentrated *in vacuo* to give the crude product. This product is diluted in ether (1.2 L) and acidified by slow
- 20 addition of 1M HCl (1.2 L). The mixture stirs for one hour and is concentrated *in vacuo*.

The resulting precipitate is diluted with acetonitrile and is stirred for 16 hours. The desired 1,7-diphenyl-4-aminoheptane hydrochloride is collected by filtration.


Example 2 - (R)-5-Oxiranylmethoxy-quinoline (2)

5



Sodium hydride (60 weight %; 1.79 g; 44.8 mmol) is washed with hexanes (3 x 10 mL) under an argon blanket. DMF (17 mL) is then added at ambient temperature and the stirred slurry is cooled to 5°C. A solution of 5-hydroxyquinoline (5.00 g; 34.4 mmol) in DMF (65 mL) is added dropwise over 30 minutes. The resulting mixture is allowed to warm to ambient temperature over 1 hour affording a clear, reddish-brown solution. A solution of (R)-(-)-glycidyl tosylate (10.22 g; 44.8 mmol) in DMF (50 mL) is added dropwise over 20 minutes. The resulting mixture is stirred at ambient temperature for 4 hours, quenched by the addition of saturated aqueous ammonium chloride (25 mL), poured onto water (750 mL), and extracted with ether (3 x 375 mL). The combined ether layers are washed with saturated aqueous sodium bicarbonate (2 x 375 mL), then dried over MgSO₄, filtered, and concentrated *in vacuo*. The residue is purified via silica gel chromatography with gradient elution (33% → 50% ethyl acetate in hexanes) affording the desired product (4.95 g) as a tan solid. ESMS: MH⁺ 202.2 (base).


Example 3 - 4-(*tert*-Butoxycarbonyl)-1-(ethoxycarbonylmethyl)piperazine (3):



1-(Ethoxycarbonylmethyl)-piperazine (5.00 g, 29.0 mmol) is dissolved in 35 mL of 50% aqueous dioxane. Triethylamine (4.41 g, 43.5 mmol) is added followed by 2-(*tert*-butoxycarbonyloxyimino)-2-phenylacetonitrile (7.86 g, 31.9 mmol). The reaction is stirred for 14.5 hours at ambient temperature, then poured onto water and extracted with ethyl acetate (2 x 100 mL). The combined organic extracts are washed with water (3 x 100 mL), brine, dried over MgSO_4 , filtered, and concentrated in vacuo. The residue is purified via silica gel chromatography with gradient elution (20% $\rightarrow$ 50% ethyl acetate in hexanes) affording the desired product as an oil. CIMS: MH^+ 273.


10

Example 4 - Lithium 4-(*tert*-butoxycarbonyl)-1-(carboxymethyl)piperazine (4):




4-(*tert*-Butoxycarbonyl)-1-(ethoxycarbonylmethyl)piperazine (3) (3.45 g, 12.7 mmol) is dissolved in 85 mL of 2:2:1 tetrahydrofuran:water:methanol. Lithium hydroxide (0.32 g, 13.3 mmol) is added and the solution stirred at ambient temperature for 3.5 hours. The reaction mixture is then concentrated in vacuo using absolute ethanol to azeotropically remove the residual water. The residue is dried in vacuo to afford the desired product as a white solid. ESMS: MH^+ for free acid 245.2.

Example 5 - 4-(*tert*-Butoxycarbonyl)-1-(carboxymethyl)piperazine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (5):




- 5 Lithium 4-(*tert*-butoxycarbonyl)-1-(carboxymethyl)piperazine (4) (1.00 g; 4.00 mmol) is dissolved in N,N-dimethylformamide (20 mL) at ambient temperature. 1-Hydroxybenzotriazole (1.62 g, 12.0 mmol) and N,N-diisopropylethylamine (2.07 g, 16.0 mmol) are added. The solution is cooled in an ice-bath and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (0.92 g, 4.8 mmol) is added. After 30 minutes 1,7-diphenyl-4-heptylamine hydrochloride (1) (1.04 g, 4.40) is added. The mixture is allowed to warm to ambient temperature and stirred for 30 hours. The reaction mixture is poured onto water (50 mL) and extracted with methylene chloride (3 x 75 mL). The combined organic extracts are washed successively with saturated sodium bicarbonate solution, water and brine. The organic solution is dried over MgSO₄, filtered and concentrated in vacuo. The residue is purified via silica gel chromatography (40%→60% ethyl acetate in hexanes) affording the desired product (41) as an oil. CIMS: MH⁺ 494

Example 6 - 1-(Carboxymethyl)piperazine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (6):



4-(*tert*-Butoxycarbonyl)-1-(carboxymethyl)piperazine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (5) (0.61 g; 1.24 mmol) is dissolved in methylene chloride (15 mL) at ambient temperature. Trifluoroacetic acid (5 mL) is added in a slow stream, and the solution is stirred for 3 hours at ambient temperature. The solution is concentrated *in vacuo* at 40°C. The residue is dissolved in methylene chloride (100 mL) and poured onto saturated sodium bicarbonate solution. The pH is adjusted to 9 with saturated potassium carbonate solution. The mixture is shaken and the layers separated. The water layer is extracted with methylene chloride (3 x 50 mL). The combined organic extracts are washed with water, dried over MgSO₄, filtered, and concentrated in *vacuo* affording the desired product (0.39 g) as a solid. ESMS: MH⁺ 394.2


Example 7 - 1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]]-4-(carboxymethyl)piperazine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (7):



1-(Carboxymethyl)piperazine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (6) (182.3 mg; 0.463 mmol) is dissolved in ethanol (12 mL) at ambient temperature. (R)-5-


Oxiranylmethoxy-quinoline (2) (93.2 mg; 0.463 mmol) is added, then the mixture is refluxed for 20 hours. After cooling to ambient temperature, the solution is concentrated in vacuo at 40°C. The residue is purified via silica gel chromatography with gradient elution (80% → 90% ethyl acetate in hexanes, 50%→100% acetone in hexanes, then 5% ethanol in acetone) affording the desired product as an oil. ESMS: MH^+ 595.4

Example 8 - (R)-4-Oxiranylmethoxy-quinoline (8):




Sodium hydride (60 weight %; 1.79 g; 44.8 mmol) is washed with hexanes (3 x 10 mL) under an argon blanket. DMF (17 mL) is then added at ambient temperature and the stirred slurry is cooled to 5°C. A solution of 4-hydroxyquinoline (5.00 g; 34.4 mmol) in DMF (65 mL) is added dropwise over 10 minutes. The resulting mixture is allowed to warm to ambient temperature over 1 hour affording a clear, reddish-brown solution. A solution of (R)-(-)-glycidyl tosylate (10.22 g; 44.8 mmol) in DMF (50 mL) is added dropwise over 10 minutes. The resulting mixture is stirred at ambient temperature for 20.5 hours, quenched by the addition of saturated aqueous ammonium chloride (25 mL), poured onto water (750 mL), and extracted with ether (3 x 375 mL). The combined ether layers are washed with saturated aqueous sodium bicarbonate (2 x 375 mL), then dried over $MgSO_4$, filtered, and concentrated *in vacuo*. The residue is purified via silica gel chromatography with gradient elution (50% → 60% acetone in hexanes) affording the desired product (1.11 g) as a tan solid. ESMS: MH^+ 202.2.

Example 9 - 1-[(R)-1-[2-Hydroxy-3-(quinolin-4-yloxy)-propyl]]-4-(carboxymethyl)piperazine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (9):



1-(Carboxymethyl)piperazine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (6) (125.4 mg; 0.319 mmol) is dissolved in ethanol (10 mL) at ambient temperature. (R)-4-Oxiranylmethoxy-quinoline (8) (64.1 mg; 0.319 mmol) is added, then the mixture is
 5 refluxed for 21.5 hours. After cooling to ambient temperature, the solution is concentrated in vacuo at 40°C. The residue is purified via silica gel chromatography with gradient elution (50% → 100% ethyl acetone in hexanes, 5% → 20% ethanol in acetone) affording the desired product as a solid. ESMS: MH^+ 595.4


10 Example 10 - 1-{(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]}-4-(carboxymethyl)piperazine ethyl ester (10):



1-(Carboxymethyl)piperazine ethyl ester (1.71 g; 9.94 mmol) is dissolved in ethanol (200 mL) at ambient temperature. (R)-5-Oxiranylmethoxy-quinoline (2) (2.00 g; 9.94 mmol) is
 15 added, then the mixture is refluxed for 18 hours. After cooling to ambient temperature,

the solution is concentrated in vacuo at 40°C. The residue is purified via silica gel chromatography with gradient elution (70%→100% acetone in hexanes, then 5% ethanol in acetone) affording the desired product as an oil. ESMS: MH^+ 374.2

5 Example 11 - 1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]]-4-(carboxymethyl)piperazine lithium salt (11):




1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]]-4-(carboxymethyl)piperazine ethyl ester (10) (1.34 g; 3.59 mmol) is dissolved in 40 mL of 2:2:1 tetrahydrofuran: water:

10 methanol. Lithium hydroxide (90.2 mg; 3.77 mmol) is added and the solution stirred for 17 hours at ambient temperature. The solution is concentrated at reduced pressure to afford the desired product as a solid.

Example 12 - 1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]]-4-


15 (carboxymethyl)piperazine (4-benzhydrylpiperazine-1-yl) amide (12):



1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]]-4-(carboxymethyl)piperazine lithium salt (11) (100 mg; 0.285 mmol) is dissolved in N,N-dimethylformamide (3 mL) at ambient temperature. 1-(Diphenylmethyl)piperazine (86.2 mg; 0.342 mmol), N,N-diisopropylethylamine (80.9 mg; 0.626 mmol) and PyBOP (177.7 mg; 0.342 mmol) are added sequentially. The reaction is stirred for 22 hours at ambient temperature. Water (3 mL) is added and the mixture shaken. The layers are separated and the water layer extracted with methylene chloride (2 x 2mL). The combined methylene chloride extracts are dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue is purified via silica gel chromatography (90% ethyl acetate in hexanes, then 50%→100% acetone in hexanes, then 5%→20% ethanol in acetone) affording the desired product as a solid. ESMS: MH^+ 580.4


Example 13 - 1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]]-4-

(carboxymethyl)piperazine N,N-dibenzylamine amide (13):




1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]]-4-(carboxymethyl)piperazine lithium salt (11) (100 mg; 0.285 mmol) is dissolved in N,N-dimethylformamide (3 mL) at ambient temperature. N,N-Dibenzylamine (67.4 mg; 0.342 mmol), N,N-diisopropylethylamine (80.9 mg; 0.626 mmol) and PyBOP (177.7 mg; 0.342 mmol) are added sequentially. The reaction is stirred for 22 hours at ambient temperature. Water (3 mL) is added and the mixture shaken. The layers are separated and the water layer extracted with methylene chloride (2 x 2 mL). The combined methylene chloride extracts are dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue is purified via silica gel chromatography (90% ethyl acetate in hexanes, then 50%→100% acetone in hexanes, then 5%→20% ethanol in acetone) affording the desired product as a solid. ESMS: MH^+ 525.0

Example 14 - 1-(tert-Butoxycarbonyl)-piperidine-2-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (14):




15 1-tert-Butoxycarbonyl-piperidine-2-carboxylic acid (3 g; 13.1 mmol) is dissolved in methylene chloride (100 mL) at ambient temperature. 1,7-Diphenyl-4-aminoheptane hydrochloride (1) (4.77 g; 15.7 mmol), diisopropylethylamine (7.3 mL; 41.9 mmol), and PyBOP (8.17 g; 15.7 mmol) are added sequentially. The mixture is stirred for 17 hours at ambient temperature then concentrated *in vacuo* at 40°C. The residue is purified via silica gel chromatography with gradient elution (10% → 30% ethyl acetate in hexanes) affording the desired product as an oil. ESMS: MH^+ 479.4

25 Example 15 - Piperidine-2-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (15):



1-(*tert*-Butoxycarbonyl)-piperidine-2-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (14) (6.77 g; 14.1 mmol) is dissolved in methylene chloride (60 mL) at ambient temperature. Trifluoroacetic acid (40 mL) is added in a slow stream, and the solution is stirred for 1.25 hours at ambient temperature. The solution is concentrated in vacuo at 40°C. The residue is dissolved in methylene chloride (300 mL) and poured onto saturated sodium bicarbonate solution. The pH is adjusted to 9 with saturated potassium carbonate solution. The mixture is shaken and the layers separated. The water layer is extracted with methylene chloride (3 x 100 mL). The combined organic extracts are washed with water, dried over MgSO₄, filtered, and concentrated in vacuo affording the desired product (5.34 g) as a white solid. ESMS: MH⁺ 379.2


Example 16 - 1-{1-[2-(R)-Hydroxy-3-(quinolin-5-yloxy)-propyl]-piperazine-4-acetyl}-piperidine-2-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (16):



1-{(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]}-4-(carboxymethyl)piperazine lithium salt (11) (100 mg; 0.285 mmol) is dissolved in N,N-dimethylformamide (3 mL) at ambient temperature. Piperidine-2-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-


amide (15) (129.3 mg; 0.342 mmol), N,N-diisopropylethylamine (80.9 mg; 0.626 mmol) and PyBOP (177.7 mg; 0.342 mmol) are added sequentially. The reaction is stirred for 46 hours at ambient temperature. Water (3 mL) is added and the mixture shaken. The layers are separated and the water layer extracted with methylene chloride (2 x 2 mL). The combined methylene chloride extracts are dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue is purified via silica gel chromatography (50%→100% acetone in hexanes, then 5%→20% ethanol in acetone) affording the desired product. ESMS: MH^+ 706.6

10 Example 17 - 1-(*tert*-Butoxycarbonyl)-piperidine-3-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (17):




15 1-*tert*-Butoxycarbonyl-piperidine-3-carboxylic acid (3.00 g; 13.1 mmol) is dissolved in methylene chloride (100 mL) at ambient temperature. 1,7-Diphenyl-4-aminoheptane hydrochloride (1) (4.77 g; 15.7 mmol), diisopropylethylamine (5.41 g; 41.9 mmol), and PyBOP (8.17 g; 15.7 mmol) are added sequentially. The mixture is stirred for 17 hours at ambient temperature then concentrated in vacuo at 40°C. The residue is purified via silica gel chromatography with gradient elution (20% → 40% ethyl acetate in hexanes) affording the desired product as an oil. ESMS: MH^+ 479.4

20 Example 18 - Piperidine-3-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (18):



1-(*tert*-Butoxycarbonyl)-piperidine-3-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (17) (6.30 g; 13.2 mmol) is dissolved in methylene chloride (60 mL) at ambient temperature. Trifluoroacetic acid (40 mL) is added in a slow stream, and the solution is stirred for 1 hour at ambient temperature. The solution is concentrated in vacuo at 40°C. The residue is dissolved in methylene chloride (300 mL) and poured onto saturated sodium bicarbonate solution. The pH is adjusted to 9 with saturated potassium carbonate solution. The mixture is shaken and the layers separated. The water layer is extracted with methylene chloride (3 x 100 mL). The combined organic extracts are washed with water, dried over MgSO₄, filtered, and concentrated in vacuo affording the desired product (5.34 g) as a white solid. ESMS: MH⁺ 379.0


Example 19 - 1-{1-[2-(R)-Hydroxy-3-(quinolin-5-yloxy)-propyl]-piperazine-4-acetyl}-piperidine-3-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (19):



1-{(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]}-4-(carboxymethyl)piperazine lithium salt (11) (100 mg; 0.285 mmol) is dissolved in N,N-dimethylformamide (3 mL) at ambient temperature. Piperidine-3-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (18) (129.3 mg; 0.342 mmol), N,N-diisopropylethylamine (80.9 mg; 0.626 mmol)


and PyBOP (177.7 mg; 0.342 mmol) are added sequentially. The reaction is stirred for 46 hours at ambient temperature. Water (3 mL) is added and the mixture shaken. The layers are separated and the water layer extracted with methylene chloride (2 x 2 mL). The combined methylene chloride extracts are dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue is purified via silica gel chromatography (50%→100% acetone in hexanes, then 5%→20% ethanol in acetone) affording the desired product. ESMS: MH^+ 706.6

Example 20 - 1-(*tert*-Butoxycarbonyl)-piperidine-4-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (20):




1-(*tert*-Butoxycarbonyl)-piperidine-4-carboxylic acid (4.00 g; 17.4 mmol) is dissolved in methylene chloride (125 mL) at ambient temperature. 1,7-Diphenyl-4-aminoheptane hydrochloride (1) (6.36 g; 20.9 mmol), diisopropylethylamine (7.27 g; 55.8 mmol), and PyBOP (10.89 g; 20.9 mmol) are added sequentially. The mixture is stirred for 14 hours at ambient temperature then concentrated in vacuo at 40°C. The residue is purified via silica gel chromatography with gradient elution (20% → 40% ethyl acetate in hexanes) affording the desired product as an oil.

Example 21 - Piperidine-4-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (21):



1-(*tert*-Butoxycarbonyl)-piperidine-4-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (20) (8.17 g; 17.1 mmol) is dissolved in methylene chloride (60 mL) at ambient temperature. Trifluoroacetic acid (40 mL) is added in a slow stream, and the solution is stirred for 2 hours at ambient temperature. The solution is concentrated in vacuo at 40°C. The residue is dissolved in methylene chloride (400 mL) and poured onto saturated sodium bicarbonate solution. The pH is adjusted to 9 with saturated potassium carbonate solution. The mixture is shaken and the layers separated. The water layer is extracted with methylene chloride (3 x 100 mL). The combined organic extracts are washed with water, dried over MgSO₄, filtered, and concentrated in vacuo affording the desired product (5.91 g) as a white solid. ESMS: MH⁺ 379.0


Example 22 - 1-{1-[2-(R)-Hydroxy-3-(quinolin-5-yloxy)-propyl]-piperazine-4-acetyl}-piperidine-4-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (22):



1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]}-4-(carboxymethyl)piperazine lithium salt (11) (100 mg; 0.285 mmol) is dissolved in N,N-dimethylformamide (3 mL) at ambient temperature. Piperidine-4-carboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (21) (129.3 mg; 0.342 mmol), N,N-diisopropylethylamine (80.9 mg; 0.626 mmol) and PyBOP (177.7 mg; 0.342 mmol) are added sequentially. The reaction is stirred for 46 hours at ambient temperature. Water (3 mL) is added and the mixture shaken. The layers are separated and the water layer extracted with methylene chloride (2 x 2 mL). The combined methylene chloride extracts are dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue is purified via silica gel

chromatography (50%→100% acetone in hexanes, then 5%→20% ethanol in acetone) affording the desired product. ESMS: MH^+ 706.6

Example 23 - N-(*tert*-Butoxycarbonyl)-*cis*-4-amino-1-cyclohexanecarboxylic acid (23):



5


23

cis-4-Amino-1-cyclohexanecarboxylic acid (2.00 g, 14.0 mmol) is dissolved in 1N sodium hydroxide (15.2 mL, 15.2 mmol) and *tert*-butanol (18 mL). Di-*tert*-butyl dicarbonate (3.06 g, 14.0 mmol) is added and the reaction stirred for 2 hours at ambient temperature. The mixture is washed with hexane (3 x 20 mL). The aqueous phase is treated with 1N HCl (20 mL) and extracted with ethyl acetate (3 x 20 mL). The combined organic extracts are dried over magnesium sulfate, filtered and concentrated in vacuo to afford the desired product as a white solid. CIMS: MH^+ 244

10

Example 24 - N-(*tert*-Butoxycarbonyl)-*cis*-4-amino-1-cyclohexanecarboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (24):

15



23

1


24

N-(*tert*-Butoxycarbonyl)-*cis*-4-amino-1-cyclohexanecarboxylic acid (23) (1.00 g; 4.11 mmol) is dissolved in methylene chloride (30 mL) at ambient temperature. 1,7-Diphenyl-4-aminoheptane hydrochloride (1) (1.50 g; 4.93 mmol), N,N-diisopropylethylamine (1.70

g; 13.15 mmol) and PyBOP (2.57 g; 4.93 mmol) are added sequentially. The reaction is stirred 4 hours at room temperature, then concentrated under reduced pressure. The residue is purified via silica gel chromatography (20%→40% ethyl acetate in hexanes) affording the desired product as a solid. CIMS: MH^+ 493

5


Example 25 - *cis*-4-Amino-1-cyclohexanecarboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (25):



- 10 N-(*tert*-Butoxycarbonyl)-*cis*-4-amino-1-cyclohexanecarboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (24) (1.80 g; 3.65 mmol) is dissolved in methylene chloride (30 mL) at ambient temperature. Trifluoroacetic acid (15 mL) is added in a slow stream, and the solution is stirred for 6 hours at ambient temperature. The solution is concentrated *in vacuo* at 40°C. The residue is dissolved in methylene chloride (200 mL)
- 15 and poured onto saturated sodium bicarbonate solution. The pH is adjusted to 9 with saturated potassium carbonate solution. The mixture is shaken and the layers separated. The water layer is extracted with methylene chloride (3 x 50 mL). The combined organic extracts are washed with water, dried over $MgSO_4$, filtered, and concentrated *in vacuo* affording the desired product. CIMS: MH^+ 393


20

Example 26 - N-{1-[2-(*R*)-Hydroxy-3-(quinolin-5-yloxy)-propyl]-piperazine-4-acetyl}-*cis*-4-amino-1-cyclohexanecarboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (26):



1-{(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]}-4-(carboxymethyl)piperazine lithium salt (11) (100 mg; 0.285 mmol) is dissolved in N,N-dimethylformamide (3 mL) at ambient temperature. *cis*-4-Amino-1-cyclohexanecarboxylic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (25) (134.1 mg; 0.342 mmol), N,N-diisopropylethylamine (80.9 mg; 0.626 mmol) and PyBOP (177.7 mg; 0.342 mmol) are added sequentially. The reaction is stirred for 46 hours at ambient temperature. Water (3 mL) is added and the mixture shaken. The layers are separated and the water layer extracted with methylene chloride (2 x 2 mL). The combined methylene chloride extracts are dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue is purified via silica gel chromatography (50%→100% acetone in hexanes, then 5%→20% ethanol in acetone) affording the desired product. ESMS: MH^+ 720.6

Example 27 - N-*tert*-Butoxycarbonyl-N-methyl-2-aminoacetic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (27):




(*N*-*tert*-Butoxycarbonyl)-(N-methyl)-2-aminoacetic acid (Sigma Chemical Company) (1.00 g; 5.29 mmol) is dissolved in methylene chloride (40 mL) at ambient temperature. 1,7-Diphenyl-4-aminoheptane hydrochloride (1) (1.93 g; 6.34 mmol), N,N-diisopropylethylamine (2.19 g; 16.9 mmol) and PyBOP (3.30 g; 3.30 mmol) are added

sequentially. The reaction is stirred for 1 hour at room temperature, then concentrated under reduced pressure. The residue is purified via silica gel chromatography (20%→40% ethyl acetate in hexanes) affording the desired product as a solid. CIMS: MH^+ 439


5

Example 28 - N-Methyl-2-aminoacetic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (28):




N-tert-Butoxycarbonyl-N-methyl-2-aminoacetic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (27) (2.19 g; 4.99 mmol) is dissolved in methylene chloride (30 mL) at ambient temperature. Trifluoroacetic acid (20 mL) is added in a slow stream, and the solution is stirred for 2.5 hours at ambient temperature. The solution is concentrated *in vacuo* at 40°C. The residue is dissolved in methylene chloride (200 mL) and poured onto saturated sodium bicarbonate solution. The pH is adjusted to 9 with saturated potassium carbonate solution. The mixture is shaken and the layers separated. The water layer is extracted with methylene chloride (3 x 50 mL). The combined organic extracts are washed with water, dried over $MgSO_4$, filtered, and concentrated *in vacuo* affording the desired product (1.65 g) as a white solid. CIMS: MH^+ 339

20 Example 29 - N-{1-[2-(R)-Hydroxy-3-(quinolin-5-yloxy)-propyl]-piperazine-4-acetyl}-N-methyl-2-aminoacetic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (29):




1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]-4-(carboxymethyl)piperazine lithium salt (11) (100 mg; 0.285 mmol) is dissolved in N,N-dimethylformamide (3 mL) at ambient temperature. N-Methyl-2-aminoacetic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (28) (115.6 mg; 0.342 mmol), N,N-diisopropylethylamine (80.9 mg; 0.626 mmol) and PyBOP (177.7 mg; 0.342 mmol) are added sequentially. The reaction is stirred for 46 hours at ambient temperature. Water (3 mL) is added and the mixture shaken. The layers are separated and the water layer extracted with methylene chloride (2 x 2 mL). The combined methylene chloride extracts are dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue is purified via silica gel chromatography (50%→100% acetone in hexanes, then 5%→20% ethanol in acetone) affording the desired product. ESMS: MH^+ 666.4

Example 30 - 1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]-piperazine-4-acetyl]-piperazine-4-acetic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (31):




1-[(R)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]]-4-(carboxymethyl)piperazine lithium salt (11) (100 mg; 0.285 mmol) is dissolved in N,N-dimethylformamide (3 mL) at ambient temperature. Piperazine-4-acetic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (6) (80.0 mg; 0.342 mmol), N,N-diisopropylethylamine (80.9 mg; 0.626 mmol) and
 5 PyBOP (177.7 mg; 0.342 mmol) are added sequentially. The reaction is stirred for 46 hours at ambient temperature. Water (3 mL) is added and the mixture shaken. The layers are separated and the water layer extracted with methylene chloride (2 x 2 mL). The combined methylene chloride extracts are dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue is purified via silica gel
 10 chromatography (50%→100% acetone in hexanes, then 5%→20% ethanol in acetone) affording the desired product. ESMS: MH^+ 721.6

Example 31 - N-tert-Butoxycarbonyl-3-(3-pyridyl)alanine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (32):




15 (N-tert-Butoxycarbonyl)-3-(3-pyridyl)alanine (1.00 g; 3.76 mmol) is dissolved in methylene chloride (25 mL) at ambient temperature. 1,7-Diphenyl-4-aminoheptane hydrochloride (1) (1.37 g; 4.51 mmol), N,N-diisopropylethylamine (1.55 g; 12.0 mmol) and PyBOP (2.34 g; 4.51 mmol) are added sequentially. The reaction is stirred for 2.5
 20 hours at room temperature, then concentrated under reduced pressure. The residue is purified via silica gel chromatography (60%→80% ethyl acetate in hexanes) affording the desired product as a solid. ESMS: MH^+ 516.2

Example 32 - 3-(3-Pyridyl)alanine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (33):



N-*tert*-Butoxycarbonyl-3-(3-pyridyl)alanine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (32) (2.08 g; 4.03 mmol) is dissolved in methylene chloride (40 mL) at ambient temperature. Trifluoroacetic acid (20 mL) is added in a slow stream, and the solution is stirred for 4 hours at ambient temperature. The solution is concentrated in vacuo at 40°C. The residue is dissolved in methylene chloride (200 mL) and poured onto saturated sodium bicarbonate solution. The pH is adjusted to 9 with saturated potassium carbonate solution. The mixture is shaken and the layers separated. The water layer is extracted with methylene chloride (3 x 50 mL). The combined organic extracts are washed with water, dried over MgSO₄, filtered, and concentrated in vacuo affording the desired product (1.65 g) as an oil. ESMS: MH⁺ 416.2


Example 33 - N-{1-[2-(*R*)-Hydroxy-3-(quinolin-5-yloxy)-propyl]-piperazine-4-acetyl}-3-(3-pyridyl)alanine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (34):



1-[(*R*)-1-[2-Hydroxy-3-(quinolin-5-yloxy)-propyl]]-4-(carboxymethyl)piperazine lithium salt (11) (100 mg; 0.285 mmol) is dissolved in *N,N*-dimethylformamide (3 mL) at ambient temperature. 3-(3-Pyridyl)alanine [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (33) (141.9 mg; 0.342 mmol), *N,N*-diisopropylethylamine (80.9 mg; 0.626 mmol) and


PyBOP (177.7 mg; 0.342 mmol) are added sequentially. The reaction is stirred for 42 hours at ambient temperature. Water (3 mL) is added and the mixture shaken. The layers are separated and the water layer extracted with methylene chloride (2 x 2 mL). The combined methylene chloride extracts are dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue is purified via silica gel chromatography (70%→100% acetone in hexanes, then 5%→50% ethanol in acetone) affording the desired product. ESMS: MH^+ 743.4

Example 34 - 3-[4-(*tert*-Butoxycarbonyl)piperazin-1-yl]-3-oxopropionic acid methyl ester (35):




N-(*tert*-Butoxycarbonyl)piperazine (2.50 g; 13.4 mmol) and triethylamine (1.77 g; 2.43 mmol) are dissolved in methylene chloride (40 mL). The solution is cooled in an ice-bath and a solution of methyl 3-chloro-3-oxopropionate (2.02 g; 14.8 mmol) in 10 mL methylene chloride is added dropwise over a 10-minute period. The ice-bath is removed and the reaction mixture is stirred for 1 hour at room temperature. The reaction mixture is then poured onto ice-cold 0.1N HCl (100 mL) and extracted with methylene chloride (200 mL). The organic extract is washed with saturated sodium bicarbonate and water, dried over magnesium sulfate, filtered and concentrated in vacuo. The residue is purified by chromatography on silica gel (60%→80% ethyl acetate in hexanes) to afford the desired product as a white solid. ESMS: MH^+ 287.2

Example 35 - 3-[4-(*tert*-Butoxycarbonyl)piperazin-1-yl]-3-oxopropionic acid (36):



3-[4-(*tert*-Butoxycarbonyl)piperazin-1-yl]-3-oxopropionic acid methyl ester (35) (3.30 g; 11.5 mmol) is dissolved in 2:2:2 tetrahydrofuran: water: methanol. Lithium chloride (2.76 g; 115.3 mmol) is added and the mixture is stirred for 1 hour at ambient temperature. The solution is concentrated in vacuo to remove the tetrahydrofuran and methanol. The concentrate is poured onto citric acid solution (115 mL) and extracted with ethyl acetate (230 mL). The organic layer is washed with water (3 x 115 mL) and brine (1 x 115 mL), then dried over magnesium sulfate, filtered and concentrated in vacuo to afford the desired product as a solid. ESMS: MH^+ 273.0


Example 36 - 3-[4-(*tert*-Butoxycarbonyl)piperazin-1-yl]-3-oxopropionic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (37):



3-[4-(*tert*-Butoxycarbonyl)piperazin-1-yl]-3-oxopropionic acid (36) (1.20 g; 4.41 mmol) is dissolved in methylene chloride (40 mL) at ambient temperature. 1,7-Diphenyl-4-aminoheptane hydrochloride (1) (1.61 g; 5.29 mmol), *N,N*-diisopropylethylamine (1.82 g; 14.1 mmol) and PyBOP (2.75 g; 5.29 mmol) are added sequentially. The reaction is


stirred for 14 hours at room temperature, then concentrated under reduced pressure. The residue is purified via silica gel chromatography (40%→70% ethyl acetate in hexanes) affording the desired product as a solid. ESMS: MH^+ 522.4

5 Example 37 - 3-(Piperazin-1-yl)-3-oxopropionic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (38):



3-[4-(*tert*-Butoxycarbonyl)piperazin-1-yl]-3-oxopropionic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (37) (2.05 g; 3.93 mmol) is dissolved in methylene chloride (30 mL) at ambient temperature. Trifluoroacetic acid (15 mL) is added in a slow stream, and the solution is stirred for 1 hour at ambient temperature. The solution is concentrated in vacuo at 40°C. The residue is dissolved in methylene chloride (200 mL) and poured onto saturated sodium bicarbonate solution. The pH is adjusted to 9 with saturated potassium carbonate solution. The mixture is shaken and the layers separated. The water layer is extracted with methylene chloride (3 x 50 mL). The combined organic extracts are washed with water, dried over $MgSO_4$, filtered, and concentrated in vacuo affording the desired product as an oil. ESMS: MH^+ 422.2

20 Example 38 - 3-{4-[2-(R)-Hydroxy-3-(quinolin-5-yloxy)-propyl]-piperazin-1-yl}-3-oxopropionic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (39):



3-(Piperazin-1-yl)-3-oxopropionic acid [4-phenyl-1-(3-phenyl-propyl)-butyl]-amide (38) (209.5mg; 0.497 mmol) is dissolved in ethanol (12 mL) at ambient temperature. (R)-5-Oxiranylmethoxy-quinoline (2) (100.0 mg; 0.497 mmol) is added, then the mixture is refluxed for 15 hours. After cooling to ambient temperature, the solution is concentrated in vacuo at 40°C. The residue is purified via silica gel chromatography with gradient elution (neat ethyl acetate, then 50%→100% acetone in hexanes, then 5% ethanol in acetone) affording the desired product as a solid. ESMS: MH^+ 623.4

10 Reference Example 1 - Method for Measuring Activity to Inhibit Pgp (Reversal Assay)

NIH-MDR1-G185 cells (obtained from M. Gottesman, NIH) were harvested and resuspended at 6×10^4 cells/ml in RPMI 1640 containing L-glutamine, 10% Cosmic calf serum, and penicillin-streptomycin. Cell suspension aliquots of 100 microliters were added to individual wells of a 96 well microtiter plate and incubated overnight at 37°C to allow cells to adhere. Cell viability in the presence of an anticancer drug was determined in the presence and absence of an MDR modifying agent using an MTT assay (P. A. Nelson, et. al, *J. Immunol*, 150:2139-2147 (1993)).

Briefly, cells were preincubated with an MDR modulating agent (final concentration 5 micromolar) for 15 min at 37° C, then treated with varying concentrations of an anticancer agent for 72 hr at 37° C. MTT dye (20 microliters of 5 mg/ml PBS solution) was added to each well and incubated for 4 hr at 37° C. Media was carefully removed and dye was solubilized with 100 microliters of acidified isopropyl alcohol. Absorption was measured on a spectrophotometric plate reader at 570 nm and corrected

for background by subtraction at 630 nm. Reversal index was calculated for each MDR modulator and normalized to the reversal index of a benchmark modulator, VX-710 as below:

- 5 Reversal index = IC_{50} in the absence of modulator / IC_{50} in the presence of modulator
Normalized reversal index = Reversal index of modulator / Reversal index of VX-710

VX-710 is (S)-N-[2-Oxo-2-(3,4,5-trimethoxyphenyl)acetyl]piperidine-2-carboxylic acid 1,7-bis(3-pyridyl)-4-heptyl ester.

10

Reference Example 2 - Method for Measuring Activity to Inhibit Pgp and MRP1 (Calcein AM Extrusion Assay)

- Pgp-dependent calcein AM extrusion was measured in NIH-MDR1-G185 cells or HL60-MDR1 cells. MRP1-dependent calcein AM extrusion was measured in HL60/ADR
15 cells. Dye uptake was measured by incubating $0.5 - 1 \times 10^6$ cells/ml in cell culture medium containing 0.25 mM calcein AM at 37° C at an excitation wavelength = 493 nm and an emission wavelength = 515 nm. Inhibition of calcein AM transport by varying concentrations of MDR modulators was determined by measuring the rate of increase in fluorescence of free calcein for 5 min periods. The IC_{50} values were obtained by
20 determining the concentration of modulator resulting in 50% of the maximum transport inhibition. Maximum transport inhibition was the % inhibition produced in the presence of 50 - 60 micromolar verapamil.

Reference Example 3 - Fluorescent Substrate Accumulation Assay

- 25 NIH-MDR1-G185 cells (obtained from M. Gottesman, NIH) were harvested and resuspended in RPMI-1640 containing L-glutamine, 10% Cosmic Calf Serum and penicillin-streptomycin. Cell suspension aliquots of 175 microliters (1×10^5 cells) were added to individual wells of a 96 well microtiter plate and preincubated for 15 min at 37° C with 20 microliters MDR modulator diluted in cell culture media to give a final
30 concentration of 10 micromolar. Control wells received no modulating agent. BODIPY-

FL Taxol (Molecular Probes, Eugene, Ore.) was added to each well in 10 microliter aliquots to give a final concentration of 500 nM and cells were incubated for 40 min at 37° C. Cells were centrifuged at 100 x g for 5 min at 4° C and the cell pellet washed with 200 microliters cold PBS to remove fluorescent medium from wells. Cells were

5 centrifuged once more, media removed, and cells resuspended in 200 microliters cold PBS. Fluorescence accumulation was measured in a fluorescence plate reader fitted with an excitation filter of 485 nm and an emission filter of 538 nm. BODIPY-FL taxol accumulation in the cells was calculated as follows:

- 10 Accumulation Index = (fluorescence in NIH-MDR1-G185 cells in the presence of modulator) / (fluorescence in NIH-MDR1-G185 cells in absence of modulator)

Reference Example 4 - Method for Measuring Substrate Potential for MDR1 (MDR1 ATPase assay)

- 15 Recombinant baculovirus carrying the human MDR1 gene was generated and Sf9 cells infected with virus. The virus-infected cells were harvested and their membranes isolated. MDR1-ATPase activity of the isolated Sf9 cell membranes was estimated by measuring inorganic phosphate liberation as previously described (B. Sarkadi, *J. Biol. Chem.*, 1992, 267:4854 - 4858). The differences between the ATPase activities measured
- 20 in the absence and presence of 100 micromolar vanadate were determined as activity specific to MDR1. MDR modulator concentrations causing half-maximum activation (K_a) or half-maximum inhibition of the MDR1-ATPase stimulated by 30 - 40 micromolar verapamil (K_i) were determined.

25 Example A - Activity of the Compounds

Accumulation Index of various compounds prepared above was tested according to the method in Reference Example 3. The results are in Table 5.

Table 5 - Accumulation Index of the Active Compounds

Example	Compound	Accumulation Index
---------	----------	--------------------

7	7	10
9	9	8
12	12	7
13	13	6
16	16	10
19	19	10
22	22	9
26	26	9
29	29	8
30	31	8
33	34	7
38	39	9

Example B - Oral Composition for the Active Compound of this Invention

A composition for oral administration is prepared by reducing an active compound according to this invention to a No. 60 powder. Starch and magnesium stearate are passed
5 through a No. 60 bolting cloth onto the powder. The combined ingredients are mixed for 10 minutes and filled into a hard shell capsule of a suitable size at a fill weight of 100 mg per capsule. The capsule contains the following composition:

Active Compound	5 mg
Starch	88 mg
10 Magnesium Stearate	7 mg

Example C - Oral Composition for the Active Compound of this Invention with a Chemotherapeutic Agent

A mixture of vinblastine and an active compound of this invention is reduced to a
15 No. 60 powder. Lactose and magnesium stearate are passed through a No. 60 bolting cloth onto the powder. The combined ingredients are mixed for 10 minutes, and then filled into a No. 1 dry gelatin capsule. Each capsule contains the following composition:

Active Compound	5 mg
-----------------	------

Vinblastine	5 mg
Lactose	580 mg
Magnesium Stearate	10 mg

5 Example D - Parenteral Composition for the Active Compound of this Invention

An active compound according to this invention (1 mg) is dissolved in 1 mL of a solution of 10% cremaphor, 10% ethanol, and 80% water. The solution is sterilized by filtration.


10 Example E - Parenteral Composition for the Active Compound of this Invention

A sufficient amount of an active compound according to this invention and TAXOL® are dissolved in a 0.9% sodium chloride solution such that the resulting mixture contains 0.9 mg/mL of the active compound of this invention and 1.2 mg/mL TAXOL®.

15 A sufficient amount of the solution to deliver 135 mg/sq m TAXOL® is administered intravenously over 24 hours to a patient suffering from ovarian cancer.


WHAT IS CLAIMED IS:

1. A compound having the structure:



or an optical isomer, diastereomer, enantiomer, pharmaceutically-acceptable salt, biohydrolyzable amide, biohydrolyzable ester, or biohydrolyzable imide thereof, wherein:


- (a) s is an integer from about 1 to about 3;
- (b) A is selected from the group consisting of A^1 and A^3 ;
- (c) A^1 is a group having the structure:




- (d) each R^1 is independently selected from the group consisting of a hydrogen atom, a hydroxyl group, a hydrocarbon group, a substituted hydrocarbon group, a heterogeneous group, a substituted heterogeneous group, a carbocyclic group, a substituted carbocyclic group, a heterocyclic group, a substituted heterocyclic group, an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group;
- (e) x is 0 to about 10;
- (f) R^2 is selected from the group consisting of a hydrocarbon group, a substituted hydrocarbon group, a heterogeneous group, a substituted heterogeneous group, a carbocyclic group, a substituted carbocyclic group, a heterocyclic group, a substituted heterocyclic group, an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group;
- (g) D^1 and D^2 are each independently selected from the group consisting of ---C(O)--- and $\text{---NR}^3\text{---}$;
- (h) R^3 is selected from the group consisting of a hydrogen atom and R^2 , and with the proviso that optionally, R^2 and R^3 may be bonded together thereby forming a ring

selected from the group consisting of heterocyclic groups and substituted heterocyclic groups;

- (i) y is 0 or 1 and z is 0 or 1, with the proviso that when y is 0, z is 1 and when y is 1, z is 0, when y is 0 and D^1 is $-NR^3-$, then D^2 is $-C(O)-$, when y is 0 and D^2 is $-NR^3-$, then D^1 is $-C(O)-$, and when x is 0, D^1 is $-C(O)-$, y is 0, D^2 is $-NR^3-$, and D^3 is $-C(O)-$ or $-S(O)_2-$, then R^2 is selected from the group consisting of a hydrocarbon group and a substituted hydrocarbon group, wherein the substituted hydrocarbon group is substituted with an aromatic group; and
- (j) A^2 is a group having the structure:



- (k) u is 0 to about 10; p is 0 to about 10; v is 0 or 1;
- (l) D^3 is selected from the group consisting of $-S(O)_2-$, $-C(O)-$, and $-CR^1(OH)-$, with the proviso that when D^3 is $-S(O)_2-$, then D^1 is $-C(O)-$, D^2 is $-NR^3-$, and R^2 and R^3 are bonded together to form the ring structure, and when D^3 is $-C(O)-$, v is 0, and R^5 contains a $-C(O)-$ group, then p is not 0;
- (m) R^5 is selected from the group consisting of a substituted hydrocarbon groups of at least 2 carbon atoms and substituted heterogeneous groups of at least 2 member atoms, wherein R^5 is substituted with at least one group selected from the group consisting of aromatic groups, heteroaromatic groups, substituted aromatic groups, and substituted heteroaromatic groups;
- (n) A^3 has the structure:




- (o) t is about 1 to about 6;
- (p) D^4 is selected from the group consisting of $-C(O)-$ and $-CHR^1-$;
- (q) D^5 is selected from the group consisting of $-NR^6(R^7)$, $-O_rR^6$, and $-C(O)R^6$;
- (r) r is 0 or 1;
- (s) R^6 is selected from the group consisting of a hydrocarbon group, a substituted hydrocarbon group, a heterogeneous group, a substituted heterogeneous group, a

carbocyclic group, a substituted carbocyclic group, a heterocyclic group, a substituted heterocyclic group, an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group; and


- (t) R^7 is selected from the group consisting of a hydrogen atom and R^6 , with the proviso that when u is 0, D^3 and D^4 are not both $-C(O)-$.


2. The compound according to Claim 1, wherein u is about 1 to about 3, p is about 1 to about 3, and v is 1.
3. The compound according to any of the preceding claims, wherein D^3 is $-CR^1(OH)-$.
4. The compound according to any of the preceding claims, wherein R^5 is a heteroaromatic group of the formula:




wherein each X is independently selected from the group consisting of CH and a heteroatom, with the proviso that at least one X is a heteroatom.

5. The compound according to any of the preceding claims, wherein A is A^1 , D^1 is $-C(O)-$, and D^2 is $-NR^3-$.
6. The compound according to any of the preceding claims, wherein the compound has a structure selected from the group consisting of:





7. The compound according to any of the preceding claims, wherein the compound has the structure:



8. A composition characterized by the compound according to any of the preceding claims and a pharmaceutically acceptable carrier.

9. A composition according to Claim 8, further characterized by a therapeutic agent selected from the group consisting of cancer therapeutic agents, antibacterial agents, antiviral agents, antifungal agents, and combinations thereof.
10. Use of the compound according to any of the preceding claims for the manufacture of a medicament suitable for treatment of multidrug resistance.