

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2017202079 B2**

(54) Title
Prevention of Adverse Effects Caused by CD3 Specific Binding Domains

(51) International Patent Classification(s)
C07K 16/28 (2006.01) **A61K 39/395** (2006.01)
A61K 31/573 (2006.01)

(21) Application No: **2017202079** (22) Date of Filing: **2017.03.29**

(43) Publication Date: **2017.04.20**

(43) Publication Journal Date: **2017.04.20**

(44) Accepted Journal Date: **2017.11.30**

(62) Divisional of:
2011328393

(71) Applicant(s)
Amgen Research (Munich) GmbH

(72) Inventor(s)
ZUGMAIER, Gerhard;NAGORSEN, Dirk;SCHEELE, Juergen

(74) Agent / Attorney
WRAYS PTY LTD, 56 Ord St, West Perth, WA, 6005, AU

(56) Related Art
WO 1999/054440 A1
WO 2007/068354 A1

ABSTRACT

5 The present invention relates to a glucocorticoid (GC) for use in the amelioration, treatment or prophylaxis of neurological/psychiatric adverse events caused by a CD3 binding domain. Kits comprising a GC, a CD3 binding domain and instructions for use which indicate that the GC is to be employed for the treatment amelioration and/or prophylaxis of neurological adverse events caused by said CD3 binding domain, are also disclosed.

2017202079 29 Mar 2017

5

Prevention of adverse effects caused by CD3 specific binding domains

10 [0001] The present invention relates to a glucocorticoid (GC) for use in the amelioration, treatment or prophylaxis of neurological/psychiatric adverse events caused by a CD3 binding domain. Kits comprising a GC, a CD3 binding domain and instructions for use which indicate that the GC is to be employed for the treatment amelioration and/or prophylaxis of neurological adverse events caused by said CD3 binding domain, are also disclosed.

15

[0002] Antibody-based cancer therapies require a target antigen firmly bound to the surface of cancer cells in order to be active. By binding to the surface target, the antibody can directly deliver a deadly signal to the cancer cell or indirectly by, for example, recruiting a cytotoxic T cell, if it is a bispecific antibody. In an ideal treatment scenario, a target antigen is abundantly present and accessible on every cancer cell and is absent, shielded or much less abundant on normal cells. This situation provides the basis for a therapeutic window in which a defined amount of the antibody-based therapeutic effectively hits cancer cells but spares normal cells.

25 [0003] Though binding domains like antibodies are an effective means in treating many disorders, in particular cancer, their administration is not necessarily devoid of side effects. Adverse effects may cause a reversible or irreversible change in the health status of a patient. As adverse effects could be harmful and undesired, it is highly desirable to avoid them. However, though it is known that a medicament can cause adverse effects, its prescription and administration could not be avoided or is accepted, since the medicament has an outstanding beneficial therapeutic effect or may even be life-saving.

35 [0004] In clinical trials, a general distinction can be made between adverse effects (AEs) and serious adverse effects (SAEs). Specifically, adverse effects can be classified in 5 grades in accordance with the Common Terminology Criteria for Adverse Events

2017202079 29 Mar 2017

(CTCAE). Grade 1 relates to mild AE, Grade 2 to moderate AE, Grade 3 to severe AE, Grade 4 to life-threatening or disabling AE, while Grade 5 means death related to AE.

5 [0005] An adverse effect observed in antibody therapy is the occurrence of infusion-related side effects, such as the cytokine release syndrome ("CRS"). Other adverse side effects described to be associated with CRS are fatigue, vomiting, tachycardia, hypertension, back pain, but also central nervous system reactions (CNS reactions, neurological and/ or psychiatric), such as seizures, encephalopathy, cerebral edema, aseptic meningitis, and headache.

10 [0006] Adverse events such as cytokine release and neurological/psychiatric reactions have not only been observed with antibodies binding to the T cell receptor but also with a CD19xCD3 bispecific single chain antibody binding to the CD3 part of the T cell receptor (called Blinatumomab (MT103)). Blinatumomab (MT103) is a lymphoma-
15 directed, recombinant bispecific single-chain CD19xCD3 antibody that binds to CD19 on the surface of almost all B cells and B tumor cells and concomitantly can engage a T cell, thereby triggering the T-cell to kill the target B cell or B tumor cell. Blinatumomab consists of four immunoglobulin variable domains assembled into a single polypeptide chain. Two of the variable domains form the binding site for CD19, a cell surface antigen
20 expressed on most B cells and B tumor cells. The other two variable domains form the binding site for the CD3 complex on T cells. Blinatumomab is designed to direct the body's cytotoxic, or cell-destroying, T cells against tumor cells, and represent a new therapeutic approach to cancer therapy. Blinatumomab is presently in clinical trials.

25 [0007] As described, for instance, in WO 99/54440, adverse effects have been observed in a previous study performed with Blinatumomab applied in repeated bolus infusions to a patient with B-cell derived chronic lymphatic leukaemia (B-CLL). As shown in Figures 19 und 20 of WO 99/54440, release of TNF, IL-6 and IL-8 has been found in response to each of the administered 20 minute-infusions of 3 microgram and 10 microgram of the
30 mentioned bispecific single chain antibody, respectively, with cytokine release after each administration. Maximal cytokine release was observed after administration of 10 microgram of bispecific single chain antibody. In a following clinical trial study, in which escalating doses of the CD19xCD3 bispecific single chain antibody have been administered to patients with B cell malignancies as bolus infusions, adverse effects

2017202079 29 Mar 2017

have also been observed. According to a retrospective analysis, 7 out of 22 patients showed an early neurological reaction, including, for example, confusion, ataxia, speech disorder, or disorientation.

5 [0008] As shown in Bargou et al. (Science 321 (2008): 974-7), doses as low as 0.005 milligrams per square meter per day continuously administered to non-Hodgkin's lymphoma patients over four weeks led to an elimination of lymphoma target cells in blood. Partial and complete tumor regressions were first observed at a dose level of 0.015 milligrams/m²/d, and all seven patients treated at a dose level of 0.06
10 milligrams/m²/d experienced a tumor regression (Bargou et al., cited above). The CD19xCD3 bispecific single chain antibody also led to clearance of tumor cells from bone marrow and liver. However, though this (still ongoing) study established clinical proof of concept for the therapeutic potency of the CD19xCD3 bispecific single chain antibody format in the treatment of blood-cell derived cancer, neurological reactions
15 have been found in the course of the aforementioned clinical trial. In order to get these undesired side effects under control, the mode of administration of the CD19xCD3 bispecific single chain antibody has been changed in that it has been switched over from bolus infusion to a continuous intravenous administration of said antibody for a longer period of time. Accordingly, since Blinatumomab is a very promising candidate
20 medicament for treating non-Hodgkin's lymphoma (NHL), such as, diffuse large B-cell lymphoma (DLBCL), follicular lymphoma and mantle cell lymphoma (MCL), acute lymphoblastic leukemia (ALL), and/or chronic lymphocytic leukemia (CLL), it is highly desirable to reduce or even completely avoid undesired side-effects in the treatment of patients in need thereof with the CD19xCD3 bispecific single chain antibody.

25 [0009] It is however difficult to design a CD19xCD3 antibody-based therapy, which does not cause CNS (neurological and/or psychiatric) reactions or, to put it differently, it is desired to provide a CD19xCD13 antibody-based medical therapies with increased patient tolerability, i.e., reduced or even no undesired adverse effects such as CNS
30 reactions.

[0010] Though pharmaceutical means and methods which allow a more gradual activation of T cell populations (see WO 2007/068354) already helped to avoid significant adverse side effects in patients treated with the CD19xCD3 bispecific single

2017202079 29 Mar 2017

chain antibody, neurological reactions could unfortunately not be prevented by these measures, in particular in cases in which doses of more than 5 to 10 microgram per square meter per day (i.e. 24h) of the antibody have been administered.

5 [0011] Thus, the technical problem underlying the present invention was to provide means and methods to overcome the above problems.

[0012] The present invention addresses this need and thus provides embodiments concerning means and methods for use in the amelioration, treatment or prophylaxis of
10 neurological adverse effects caused by a CD3 binding domain, such as a CD19xCD3 bispecific antibody.

[0013] These embodiments are characterized and described herein and reflected in the claims.

15 ***

[0014] It must be noted that as used herein, the singular forms "a", "an", and "the", include plural references unless the context clearly indicates otherwise. Thus, for example, reference to "a reagent" includes one or more of such different reagents and reference to "the method" includes reference to equivalent steps and methods known to
20 those of ordinary skill in the art that could be modified or substituted for the methods described herein.

[0015] All publications and patents cited in this disclosure are incorporated by reference in their entirety. To the extent the material incorporated by reference contradicts or is
25 inconsistent with this specification, the specification will supersede any such material.

Unless otherwise indicated, the term "at least" preceding a series of elements is to be understood to refer to every element in the series. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are
30 intended to be encompassed by the present invention.

[0016] Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and

29 Mar 2017

2017202079

"comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integer or step.

In each instance herein any of the terms "comprising", "consisting essentially of" and
5 "consisting of" may be replaced with either of the other two terms.

[0017] Several documents are cited throughout the text of this specification. Each of the documents cited herein (including all patents, patent applications, scientific publications, manufacturer's specifications, instructions, etc.), whether supra or infra, are hereby
10 incorporated by reference in their entirety. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

[0018] In view of the adverse events described in the sections above, particularly the
15 adverse CNS effects including neurological reactions observed with CD3 specific binding domains, the finding that these adverse effects can be mitigated or even prevented if the administration of the CD3-specific binding domain is accompanied or preceded by the administration of a glucocorticoid, is definitely remarkable.

[0019] Specifically, the present inventors observed that those patients, to whom a
20 CD19xCD3 bispecific antibody was administered, encountered neurological side effects, and, further, that these neurological side effects could be prevented or alleviated by means of a glucocorticoid (pre) and/or (co)therapy.

[0020] Accordingly, the present invention establishes for the first time that
25 glucocorticoids such as dexamethasone mitigate or even prevent neurological/psychiatric adverse effects which might occur in the course of a treatment with CD3 specific binding domains (see also the Example section).

[0021] Glucocorticoids (GCs) are still the most widely used immunosuppressive agents
30 for the treatment of inflammatory disorders and autoimmune diseases. Glucocorticoids (GC) are a class of steroid hormones that bind to the glucocorticoid receptor (GR), which is present in almost every vertebrate animal cell, including humans. These compounds are potent anti-inflammatory agents, regardless of the inflammation's cause.

2017202079 29 Mar 2017

Glucocorticoids suppress, inter alia, the cell-mediated immunity by inhibiting genes that code for the cytokines IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-8 and IFN- γ .

[0022] Cortisone which belongs to the group of GCs is an important therapeutic drug
5 which is used to fight many ailments ranging from Addison's disease to rheumatoid
arthritis. Ever since the discovery of its anti-rheumatic properties, which led to its
acclaim as a wonder drug, many derivatives of cortisone with enhanced properties to
better fight a specific ailment have been produced. Cortisone belongs to a group of
steroids known as corticosteroids. These steroids are produced by the adrenal cortex,
10 which is the outer part of the adrenal glands, near the kidneys. The corticosteroids are
divided into two main groups: the glucocorticoids (GCs), which control fat, protein,
calcium and carbohydrate metabolism, and the mineralocorticoids controlling sodium
and potassium levels. Cortisone belongs to the former group, i.e. to the GCs. Cortisone
and its many derivatives are used for a variety of diseases. Cortisone also helped to
15 make organ transplants a reality due to its ability to minimize the defence reaction of the
body towards foreign proteins present in the implanted organ and thus damage the
functionality of the implanted organ. However, despite clinical use during more than 50
years, the specific anti-inflammatory effects of GC on different cellular compartments of
the immune system are not yet clear. GCs affect nearly every cell of the immune
20 system, and there is growing evidence for cell type-specific mechanisms.

[0023] In a first embodiment, the present invention relates to a glucocorticoid (GC) for
use in the amelioration, treatment or prophylaxis of neurological adverse effects caused
by a CD3 binding domain. As outlined above, these unwanted adverse effects frequently
25 accompany a therapy with a CD3 binding domain. The present invention remedies these
disadvantages and provides glucocorticoid(s) for use in the amelioration, treatment or
prophylaxis of neurological adverse effects in a patient wherein said patient is subject to
therapy with a CD3 binding domain. Accordingly, the present invention relates to a
glucocorticoid (GC) for use in a method in the amelioration, treatment or prophylaxis of
30 neurological adverse effects caused by a CD3 binding domain.

[0024] The present invention thus relates to a GC for use in the amelioration, treatment
or prophylaxis of neurological adverse effects caused by a CD3 binding domain in a

human patient, wherein said GC is to be administered prior to, concurrently with and/or subsequently to the administration of said binding domain.

5 **[0024A]** Also, the present invention relates to glucocorticoid (GC) when used in the amelioration, treatment or prophylaxis of neurological adverse events caused by the administration of a CD3 binding domain, wherein the GC is administered prior to the administration of the CD3 binding domain.

10 **[0025]** Also, the present invention relates to a method of amelioration, treatment or prophylaxis of neurological adverse effects caused by a CD3 binding domain, said method comprising administering to a patient in need thereof a glucocorticoid (GC). The GC is preferably administered in an amount which is sufficient to ameliorate, treat or prevent said neurological adverse effects caused by a CD3 binding domain.

15 **[0025A]** Also, the present invention relates to a kit when used for the treatment, amelioration and/or prophylaxis of neurological adverse events caused by CD3 binding domain, comprising a dexamethasone and a CD3 binding domain and instructions for use and/or an imprint, wherein the dexamethasone is administered prior to the administration of a first dose of the CD3 binding domain and prior to the
20 administration of a second dose and/or a third dose of the CD3 binding domain.

[0025B] Also, the present invention relates to a kit when used for the treatment, amelioration and/or prophylaxis of neurological adverse events caused by the CD3 binding domain, comprising a dexamethasone and one or more doses of
25 CD3 binding domain wherein the dexamethasone is administered prior to administration of a first dose of the CD3 binding domain and prior to the administration of a second dose and/or a third dose of the CD3 binding domain and instructions for use and/or an imprint.

30 **[0026]** The neurological side effects are “caused by” the administration of a CD3 binding domain to a patient. The term “caused by” means that the CD3 binding domain is causative for the neurological side effects. The skilled person can easily evaluate whether the administration a CD3 binding domain is causative for a neurological effect or not. To this end, it is just required to closely monitor the patient

5 during the course of the administration and to detect, thereby, that the administration of the CD3 binding domain was causative for the neurological side effects. Likewise, it is envisaged to discontinue the administration of the CD3 binding domain and to evaluate whether the neurological side effects are thereby ameliorated or even fade away, which also indicates that the neurological side effects were caused by said CD3 binding domain.

10 **[0027]** The term "glucocorticoid" means compounds that bind, preferably specifically, to the glucocorticoid receptor. Said term includes compound(s) selected from the group consisting of cortisone, cortisol (hydrocortisone), cloprednol, prednisone, prednisolone, methylprednisolone, deflazacort, flucortolone, triamcinolone, (including triamcinolonacetone) , dexamethasone, betamethasone, cortivazol, paramethasone, fluticasonepropionate, triamcinolonacetone. and/or fluticasone
15 thereof. In the context of the embodiments of the present invention, the mentioned compounds may be used alone or in combination. Dexamethasone is preferred. The present invention is however not limited to the above mentioned specific GCs. It is envisaged that all substances which already are or will be classified as a "glucocorticoid", may be employed in the context of the present invention as well.
20 Such future glucocorticoids include compounds which specifically bind to and activate the glucocorticoid receptor. The term "specifically binds to the GC receptor" means in accordance with the present

25

30

35

[Text continued on page 8]

2017202079 29 Mar 2017

invention that the GC (or a compound which is assumed to act like a GC) associates with (e.g., interacts with) the GC receptor (also known as NR3C1) to a statistically significant degree as compared to association with proteins/receptors generally (i.e., non-specific binding). When the GC receptor binds to glucocorticoids, its primary mechanism of action is the regulation of gene transcription. In the absence of GC, the glucocorticoid receptor (GR) resides in the cytosol complexed with a variety of proteins including Heat shock protein 90 (hsp90), the heat shock protein 70 (hsp70) and the protein FKBP52 (FK506-binding protein 52). The binding of the GC to the glucocorticoid receptor (GR) results in release of the heat shock proteins. It is thus envisaged that a future GC, or a pharmaceutically acceptable derivative or salt of a GC is preferably able to bind to the GC receptor and to release the above mentioned heat shock proteins. The activated GR complex up-regulates the expression of anti-inflammatory proteins in the nucleus or represses the expression of pro-inflammatory proteins in the cytosol (by preventing the translocation of other transcription factors from the cytosol into the nucleus).

[0028] In a preferred embodiment, said GC is selected from the most clinical used and relevant GCs like dexamethasone, fluticasonepropionate, prednisolone, methylprednisolone, betamethasone, triamcinolonacetonide or combinations thereof.

20

[0029] In an even more preferred embodiment, said GC is dexamethasone.

[0030] Dexamethasone has the highest glucocorticoid potency of the most commonly used steroids and also has the longest half-life (see Table below). But a person skilled in the field can select one of the other known glucocorticoids, some of which are disclosed herein, and select an appropriate effective dose to ameliorate or prevent neurological adverse events that may result from the treatment of a patient in need thereof, such as a DLBCL patient with a bispecific antibody molecule that contains a CD3 binding domain, such as CD19xCD3 bispecific single chain antibody.

30

2017202079 29 Mar 2017

<i>Agent</i>	<i>Approx. equiv. dose (mg)</i>	<i>Relative anti-inflammatory (glucocorticoid) potency</i>	<i>Relative mineralocorticoid (Na⁺ retaining) potency</i>	<i>Biologic half-life (hrs)</i>
Cortisone	25	0.8	0.8	8-12
Hydrocortisone	20	1	1	8-12
Prednisone	5	4	0.8	18-36
Prednisolone	5	4	0.8	18-36
Methylprednisolone	5	5	0.5	18-36
Dexamethasone	0.75	25	0	36-54

[0031] Dexamethasone also possesses a beneficial effect in malignant central nervous system (CNS) disease (e.g. CNS lymphoma or brain metastases) – possibly due to specific penetration to the CNS. It is also preferentially (over other steroids) used to treat brain edema. Although corticosteroids decrease capillary permeability in the tumor itself, it has been found in animal models that dexamethasone may act differently and decrease edema by effects on bulk flow away from the tumor (Molnar, Lapin, & Gothuis, 1995, Neurooncol. 1995; 25(1):19-28.

[0032] For the clinical trials in connection with the application of a CD19xCD3 bispecific single chain antibody for the treatment of tumorous mass of lymph node tissue and/or extranodal lymphoma caused by DLBCL, the present inventors had to develop a treatment regime which was efficient and would be well tolerated by most of the patients. To this end, the present inventors applied a step-wise application of a CD19xCD3 bispecific single chain antibody in that 5/15/60 µg/m²/24h was administered to patients. Thereby, adverse effects, in particular neurological/psychiatric events could be reduced in number, ameliorated and even prevented. Also contemplated in the step-wise administration of a CD19xCD3 bispecific single chain antibody is a treatment regime using two of the dosages, such as 5/15 µg/m²/24h, 5/60 µg/m²/24h, or 15/60 µg/m²/24h for the duration of the patient's treatment. The appropriate dosage can be selected by the clinician on the basis of efficacy, tolerability and safety with a minimum of adverse effects in the patient.

But the inventors also contemplate the treatment of tumorous mass of lymph node tissue and/or extranodal lymphoma caused by DLBCL to include the continuous administration of a flat dose without escalation to a subsequent higher dose. For

2017202079 29 Mar 2017

example, the present treatment regime includes the administration of $5\mu\text{g}/\text{m}^2/24\text{h}$, $15\mu\text{g}/\text{m}^2/24\text{h}$, or $60\mu\text{g}/\text{m}^2/24\text{h}$ of a CD19xCD3 bispecific single chain antibody until the conclusion of a course of the treatment up to 8 weeks [56 days] with good tolerability and no adverse effects, and even longer if determined to be safe and effective.

5

[0033] It is generally preferred that each of the doses disclosed herein can be converted from amount (in μg)/ m^2/d into $\mu\text{g}/\text{d}$ by multiplying the respective dose with the factor 1.9. Accordingly, each of the doses disclosed herein can be applied in the methods and uses by multiplying it with the factor 1.9. For example, a dose of $5\mu\text{g}/\text{m}^2/\text{d}$ is converted into $9.5\mu\text{g}/\text{d}$, a dose of $15\mu\text{g}/\text{m}^2$ is converted into $28.5\mu\text{g}/\text{m}^2/$ and a dose of $60\mu\text{g}/\text{m}^2/$ is converted into $114\mu\text{g}/\text{m}^2$. It is preferred that a decimal digit that results from the multiplication is either rounded up or rounded down, respectively, to a whole number. For example, a dose of $9.5\mu\text{g}/\text{d}$ can be rounded down to $9\mu\text{g}/\text{d}$ and a dose of $28.5\mu\text{g}/\text{m}^2$ can be rounded down to $28\mu\text{g}/\text{d}$. Likewise, a dose of $9.5\mu\text{g}/\text{d}$ can be rounded up to $10\mu\text{g}/\text{d}$ and a dose of $28.5\mu\text{g}/\text{m}^2$ can be rounded up to $29\mu\text{g}/\text{d}$.

10
15

[0034] The term "pharmaceutically acceptable derivatives" includes salts, esters, enol ethers, enol esters, acetals, ketals, orthoesters, hemiacetals, hemiketals, acids, bases, solvates, hydrates or prodrugs thereof. Such derivatives may be readily prepared by those of skill in this art using known methods for such derivatisation.

20

[0035] The dose of the GC that is to be used in accordance with the embodiments of the present invention is not limited, i.e. it will depend on the circumstances of the individual patient. GC can be administered intravenously or orally. Preferred dosages of the GC include, however, between 1 to 6 mg (dexamethasone equivalent) at the lower end of dosing to 40 mg/d (dexamethasone equivalent). Said dosage can be administered all at once or subdivided into smaller dosages (see the appended examples). Particularly preferred is a dosage of 4 to 24 mg/d. Daily dosages of 4, 8, 12, 16, 20 or 24 mg/d are even more preferred. A dosage of 1 x 4mg per day, 2 x 4mg per day, 1 x 8 mg per day; 1x4 mg plus 1x8 mg per day, 2 x 8 mg per day, 2 x 10mg per day or 5 x 4mg per day and 3 x 8 mg per day of dexamethasone is particularly preferred. "d" denotes one day. Further dosage regimens are derivable from the appended examples. All dosages given in this paragraph refer to dexamethasone equivalents.

25
30

2017202079 29 Mar 2017

[0036] An "adverse effect", which is sometimes also denoted as "side effect" or "adverse event (in clinical studies)" is a harmful and undesired effect resulting from medication in the treatment of a patient with a CD3 binding domain. A "neurological/psychiatric adverse effect" which is sometimes also denoted as neurological symptom or CNS adverse effect, includes conditions of a human patient such as all forms of pain, including headache and back pain, muscle weakness or incoordination, abnormal sensations in the skin, and disturbances of the senses, seizures, encephalopathy, cerebral edema, confusion, ataxia, speech disorder, hallucinations, apraxia, paresis, tremor, headache, or disorientation. A neurological effect as used herein preferably includes psychiatric adverse effects. Sometimes, however, the terms "neurological adverse effect" and "psychiatric adverse effects" can be used interchangeable.

[0037] Specifically, neurological/psychiatric reactions observed during treatment with a CD3 binding domain include for example confusion and disorientation. "Confusion" as used herein refers to loss of orientation which is the ability to place oneself correctly in the world by time, location, and personal identity, and often memory which is the ability to correctly recall previous events or learn new material. The patients usually have difficulties to concentrate and thinking is not only blurred and unclear but often significantly slowed down. Patients with neurological/psychiatric reactions also suffer from loss of memory. Frequently, the confusion leads to the loss of ability to recognize people and/or places, or to tell time and the date. Feelings of disorientation are common in confusion, and the decision-making ability is impaired. Neurological reactions further comprise blurred speech and/or word finding difficulties. This disorder may impair both, the expression and understanding of language as well as reading and writing. Besides urinary incontinence, also vertigo and dizziness may accompany neurological reactions in some patients.

[0038] The herein mentioned "patient" is a mammal, preferably a human, who will be or (already) is treated with a CD3 binding domain.

[0039] It is also envisaged that the patient is characterized by a B/T-cell ratio of less than 1:5 (see PCT/EP2010/066207).

29 Mar 2017

2017202079

5 [0040] In a preferred embodiment, the patient is suspected/assumed to comprise or already comprises malignant CD19 positive lymphocytes (in particular B cells). In the latter case, said patient has already been diagnosed to comprise such cells. These malignant CD19 positive lymphocytes (in particular B cells) are present in a patient developing and/or suffering from leukemia and/or lymphoma.

10 [0041] A "CD3 binding domain" characterizes in connection with the present invention a binding domain which comprises a framework region and an "antigen-binding-site" or "antigen-interaction site" which is able to specifically interact with a CD3 antigen. Said binding/interaction is also understood to define a "specific recognition". The term "specifically interact/interacting" means in accordance with this invention that the binding domain is capable of binding to at least two, preferably at least three, more preferably at least four amino acids of the CD3 antigen, preferably the CD3epsilon antigen, and more preferably the human CD3epsilon antigen. Such CD3 binding domains as well as
15 specific CD3epsilon epitopes are well-known to the skilled person and exemplified in great detail for example in WO2008119567 or in WO2008119566, both of which are included herein by way of reference thereto.

20 [0042] As used herein, "CD3" denotes a molecule expressed as part of the T cell receptor and has the meaning as typically ascribed to it in the prior art. In human, it encompasses in individual or independently combined form all known CD3 subunits, for example CD3 epsilon, CD3 delta, CD3 gamma, CD3 zeta, CD3 alpha and CD3 beta. The human CD3 epsilon is indicated in GenBank Accession No.NM_000733.

25 [0043] A CD3 binding molecule which binds to the human CD3 epsilon is preferred. The CD3epsilon epitope disclosed in great detail in WO2008119567 or in WO2008119566 is even more preferred.

30 [0044] The term "framework" includes a scaffold for antigen-binding sites. For example, such a scaffold could be provided by protein A, in particular, the Z-domain thereof (affibodies), ImmE7 (immunity proteins), BPTI/APPI (Kunitz domains), Ras-binding protein AF-6 (PDZ-domains), charybdotoxin (Scorpion toxin), CTLA-4, Min-23 (knottins), lipocalins (anticalins), neokarzinostatin, a fibronectin domain, an ankyrin consensus repeat domain or thioredoxin (Skerra, Curr. Opin. Biotechnol. 18, 295-304 (2005); Hosse et al., Protein Sci. 15, 14-27 (2006); Nicaise et al., Protein Sci. 13, 1882-1891 (2004); Nygren and Uhlen, Curr. Opin. Struc. Biol. 7, 463-469 (1997)).

2017202079 29 Mar 2017

[0045] A preferred “framework” is, in the context of the present invention, the art-recognized portions of an antibody variable region that exist between the more divergent (i.e., hypervariable) complementarity determining regions (CDRs) within the variable region of an antibody. Such framework regions are typically referred to as frameworks 1 through 4 (FR1, FR2, FR3, and FR4) and provide a scaffold for the presentation of the six CDRs (three from the heavy chain and three from the light chain) in three dimensional space, to form an antigen-binding surface.

[0046] A preferred example of a CD3 binding domain in line with the present invention is an antibody. The binding domain may be a monoclonal or polyclonal antibody or derived from a monoclonal or polyclonal antibody. The term “antibody” comprises derivatives or functional fragments thereof which still retain the binding specificity. Techniques for the production of antibodies are well known in the art and described, e.g. in Harlow and Lane “Antibodies, A Laboratory Manual”, Cold Spring Harbor Laboratory Press, 1988 and Harlow and Lane “Using Antibodies: A Laboratory Manual” Cold Spring Harbor Laboratory Press, 1999. The term “antibody” also comprises immunoglobulins (Ig’s) of different classes (i.e. IgA, IgG, IgM, IgD and IgE) and subclasses (such as IgG1, IgG2 etc.).

[0047] The definition of the term “antibody” also includes embodiments such as chimeric, single chain, de-immunized and humanized antibodies, as well as antibody fragments, like, inter alia, Fab fragments. Antibody fragments or derivatives further comprise F(ab’)₂, Fv, scFv fragments or single domain antibodies, single variable domain antibodies or immunoglobulin single variable domain comprising merely one variable domain, which might be VH or VL, that specifically bind to an antigen or epitope independently of other V regions or domains; see, for example, Harlow and Lane (1988) and (1999), cited above. Such immunoglobulin single variable domain encompasses not only an isolated antibody single variable domain polypeptide, but also larger polypeptides that comprise one or more monomers of an antibody single variable domain polypeptide sequence.

[0048] Bispecific antibody formats are preferred; however other multispecific antibody formats (trispesific, tetrabodies etc.) are not excluded.

2017202079 29 Mar 2017

[0049] In a further preferred embodiment, the present invention relates to methods of treatment/dosage regimen which employ CD19xCD3 bispecific antibodies, comprising a first binding domain capable of binding to an epitope of human CD3 epsilon chain and a second binding domain capable of binding to human CD19. Examples for bispecific molecules according to the methods of the invention are described in great detail in 5 WO 99/54440 and WO 2004/106381 and WO 2008/119565. All the specific CD19xCD3 bispecific antibodies disclosed therein, including their variants, fragments, equivalents etc. are particularly preferred CD19xCD3 bispecific antibodies of the present invention.

10 [0050] As used herein, a "CD19xCD3 bispecific antibody" (including a CD19xCD3 bispecific single chain antibody) denotes a single polypeptide chain comprising two binding domains. Such single chain antibodies are preferred in the context of the methods/dosage regimen of the present invention. Each binding domain comprises at least one variable region from an antibody heavy chain ("VH or H region"), wherein the 15 VH region of the first binding domain specifically binds to the CD3 epsilon molecule, and the VH region of the second binding domain specifically binds to CD19. The two binding domains are optionally linked to one another by a short polypeptide spacer. A non-limiting example for a polypeptide spacer is Gly-Gly-Gly-Gly-Ser (G-G-G-G-S) and repeats thereof. Each binding domain may additionally comprise one variable region 20 from an antibody light chain ("VL or L region"), the VH region and VL region within each of the first and second binding domains being linked to one another via a polypeptide linker, for example of the type disclosed and claimed in EP 623679 B1, but in any case long enough to allow the VH region and VL region of the first binding domain and the VH region and VL region of the second binding domain to pair with one another such that, 25 together, they are able to specifically bind to the respective first and second binding domains. Such CD19CD3 bispecific single chain antibodies are described in great detail in WO 99/54440 and WO 2004/106381.

30 [0051] The human CD19 protein is indicated in GenBank Accession No. AAA69966.

[0052] Preferably, the bispecific antibody applied in the methods/dosage regimens of the present invention has the domain arrangement VL(CD19)-VH(CD19)-VH(CD3)-VL(CD3).

2017202079 29 Mar 2017

[0053] It is, however, also envisaged that the methods of the invention can be carried out with CD19xCD3 bispecific single chain antibodies of other domain arrangements, such as

- VH(CD19)-VL(CD19)-VH(CD3)-VL(CD3),
 5 VL(CD19)-VH(CD19)-VL(CD3)-VH(CD3),
 VH(CD19)-VL(CD19)-VL(CD3)-VH(CD3),
 VL(CD3)-VH(CD3)-VH(CD19)-VL(CD19),
 VH(CD3)-VL(CD3)-VH(CD19)-VL(CD19),
 VL(CD3)-VH(CD3)-VL(CD19)-VH(CD19), or
 10 VH(CD3)-VL(CD3)-VL(CD19)-VH(CD19).

[0054] A preferred CD19xCD3 bispecific antibody applied in the methods of the present invention comprises the

- (a) anti-CD3 CDRs of the heavy chain shown as CD3 CDR-H1 in SEQ ID NO: 11
 15 (RYTMH), more preferably in SEQ ID NO: 11 (GYTFTRYTMH), CD3 CDR-H2 in
 SEQ ID NO: 12 (YINPSRGYTNYNQKFKD) and CD3 CDR-H3 in SEQ ID NO: 13
 (YYDDHYCLDY); and/or
- (b) anti-CD3 CDRs of the light chain shown as CD3 CDR-L1 in SEQ ID NO: 14
 (RASSSVSYMN), CD3 CDR-L2 in SEQ ID NO: 15 (DTSKVAS) and CD3 CDR-L3
 20 in SEQ ID NO: 16 (QQWSSNPLT); and/or
- (c) anti-CD19 CDRs of the heavy chain shown as CD19 CDR-H1 in SEQ ID NO: 17
 (SYWMN), more preferably in SEQ ID NO: 17 (GYAFSSYWMN), CD19 CDR-H2
 in SEQ ID NO: 18 (QIWPGDGDTNYNGKFKG) and CD19 CDR-H3 in SEQ ID
 NO: 19 (RETTTVGRYYYAMDY); and/or
- 25 (d) anti-CD19 CDRs of the light chain shown as CD19 CDR-L1 in SEQ ID NO: 20
 (KASQSVDYDGDSYLN), CD19 CDR-L2 in SEQ ID NO: 21 (DASNLVS) and
 CD19 CDR-L3 in SEQ ID NO: 22 (QQSTEDPWT).

[0055] It is more preferred that the CD19xCD3 bispecific antibody applied in the
 30 methods of the present invention comprises the CD3 CDRs of the heavy and light chain.
 Even more preferably, the CD19xCD3 bispecific antibody applied in the methods of the
 present invention comprises the CD3 CDRs of the heavy and light chain as well as the
 CD19 CDRs of the heavy and light chain.

2017202079 29 Mar 2017

The CDRs referred to herein are in accordance with the Kabat numbering system. The Kabat numbering scheme is a widely adopted standard for numbering the residues in an antibody in a consistent manner (Kabat et al., Sequences of Proteins of Immunological Interest, 1991).

5

[0056] Alternatively, it is preferred that the CD19xCD3 bispecific antibody applied in the methods of the present invention comprises the

- (a) CD19 variable heavy chain shown in SEQ ID NO: 3 (nucleotide sequence is shown in SEQ ID NO: 4); and/or
- 10 (b) CD19 variable light chain shown in SEQ ID NO: 5 (nucleotide sequence is shown in SEQ ID NO: 6); and/or
- (c) CD3 variable heavy chain shown in SEQ ID NO: 7 (nucleotide sequence is shown in SEQ ID NO: 8); and/or
- (d) CD3 variable light chain shown in SEQ ID NO: 9 (nucleotide sequence is shown in SEQ ID NO: 10).

15

More preferably, the CD19xCD3 bispecific antibody applied in the methods of the present invention comprises the CD19 variable heavy and light chain and/or the CD3 variable heavy and light chain. Even more preferably, the CD19xCD3 bispecific antibody applied in the methods of the present invention comprises the CD19 variable heavy and light chain as well as the CD3 variable heavy and light chain.

20

[0057] In another alternative, it is also preferred that said bispecific single chain antibody comprises an amino acid sequence selected from the group consisting of

- (a) an amino acid sequence as depicted in SEQ ID NO: 1;
- 25 (b) an amino acid sequence encoded by a nucleic acid sequence as shown in SEQ ID NO: 2;
- (c) an amino acid sequence encoded by a nucleic acid sequence having at least 70%, 80%, 90%, 95% or 99% identity to a nucleic acid sequence of (b), wherein said amino acid sequence is capable of specifically binding to CD3 and CD19;
- 30 and
- (d) an amino acid sequence encoded by a nucleic acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence of (b), wherein said amino acid sequence is capable of specifically binding to CD3 and CD19.

2017202079 29 Mar 2017

[0058] It is to be understood that the sequence identity is determined over the entire amino acid sequence. For sequence alignments, for example, the programs Gap or BestFit can be used (Needleman and Wunsch J. Mol. Biol. 48 (1970), 443-453; Smith and Waterman, Adv. Appl. Math 2 (1981), 482-489), which is contained in the GCG software package (Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711 (1991)). It is a routine method for those skilled in the art to determine and identify an amino acid sequence having e.g. 70%, 80%, 90%, 95%, 96%, 97%, 98% or 99% sequence identity to the amino acid sequences of the CD19xCD3 bispecific antibody described herein (preferably MT103). For example, according to Crick's Wobble hypothesis, the 5' base on the anti-codon is not as spatially confined as the other two bases, and could thus have non-standard base pairing. Put in other words: the third position in a codon triplet may vary so that two triplets which differ in this third position may encode the same amino acid residue. Said hypothesis is well known to the person skilled in the art (see e.g. http://en.wikipedia.org/wiki/Wobble_Hypothesis; Crick, J Mol Biol 19 (1966): 548-55). It is furthermore a routine procedure for those skilled in the art to determine cytotoxic activity of such an amino acid sequence having e.g. 70%, 80%, 90%, 95%, 96%, 97%, 98% or 99% sequence identity to the nucleotide or amino acid sequences of the CD19xCD3 bispecific single chain antibody described herein. Cytotoxic activity of the CD19xCD3 bispecific single chain antibody or an antibody construct having e.g. 70%, 80%, 90%, 95%, 96%, 97%, 98% or 99% sequence identity to the amino acid sequences of the CD19xCD3 bispecific single chain antibody can be determined by methods as illustrated e.g. in WO 99/54440.

[0059] Particularly preferred, said CD19xCD3 bispecific single chain antibody has the amino acid sequence shown in SEQ ID NO: 1.

[0060] Also particularly preferred is the CD19xCD3 bispecific antibody MT103 described in WO 99/54440 as well as those CD19xCD3 bispecific antibodies described in WO 2004/106381 and WO 2008/119565.

[0061] The present invention also relates to a CD19xCD3 bispecific antibody for use in the treatment of malignant CD19 positive lymphocytes in a human patient, wherein said antibody is to be administered prior to, concurrently with or subsequently to the administration of a GC.

2017202079 29 Mar 2017

[0062] The present invention further relates to a method for

- (i) administering a CD3 binding domain, such as a CD19xCD3 bispecific antibody to a human patient, or
- (ii) treating malignant CD19 positive lymphocytes in a human patient by administering the CD3 binding domain;

wherein said antibody is to be administered prior to, subsequently to and/or in combination with a GC.

[0063] The administration of the CD3 binding domain or of a pharmaceutical composition comprising said CD3 binding domain is preferably an intravenous administration. The administration of the GC or a pharmaceutical composition comprising said GC is preferably intravenous or per os (p.o.). They may be administered as a bolus injection or continually (continuously), with continually being preferred. A continual administration refers to an administration which is essentially without interruption. "Essentially without interruption" includes a continual administration usually without an uninterrupted flow or spatial extension. By way of example, WO 2007/068354 discloses a treatment regimen which is specifically included herein by way of reference thereto. Other treatment regimens which are envisaged in the context of the present invention are disclosed in PCT/EP2010/066207.

[0064] In a preferred embodiment, a first dose of the CD3 binding domain is administered for a first period of time; and optionally consecutively a second dose of the CD3 binding domain is administered for a second period of time, wherein the second dose exceeds the first dose.

[0065] The term "exceeds" means that the second period of time is at least one day longer than the first period of time.

[0066] In another preferred embodiment of the present application, a third dose of the CD3 binding domain is administered for a third period of time after administering a first and second dose for a first and second period of time. Accordingly, the present invention provides a three-stage (three-step) administration scheme (dosage regimen) to be applied in the uses and methods described herein.

2017202079 29 Mar 2017

[0067] In a further preferred embodiment, the present invention encompasses the continuous administration of a flat dose of the CD3 binding domain without escalation to a subsequent higher dose. For example, the present administration includes the administration of $60\mu\text{g}/\text{m}^2/24\text{h}$, $15\mu\text{g}/\text{m}^2/24\text{h}$ or $5\mu\text{g}/\text{m}^2/24\text{h}$ of a CD3 binding domain, in particular a CD19xCD3 bispecific single chain antibody until the conclusion of a course of the treatment up to 8 weeks [56 days] and even longer if determined to be safe and effective.

[0068] More specifically, in a three-stage administration scheme, dexamethasone is administered in the range of between 6 and 48 hours before the administration of the first dose of the CD3 binding domain, more preferably between 6 and 12 hours, and more preferably 12 hours, before the first dose administration. Then approximately 1 hour before (range 15min - 2h including 30 min, 45 min, 60 min, 75 min, 90 min) the first dose of the binding domain is administered, a dose of dexamethasone is again administered to the patient. Then dexamethasone is administered 1 or more days, preferably 2 to 3 days, after the first dose of the antibody, preferably on the two days after the first binding domain administration and administered 1 or more days, preferably 2 or more days after each dose increase, preferably on the two days after the administration of the dose increase of the binding domain. Each of the dexamethasone doses is preferably between 6 and 40 mg, and preferably at approximately 20 or 24 mg per dose.

[0069] The time range between 6 and 48 hours includes the administration of the dexamethasone dose and means that the times prior to the first administration of the binding domain are 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47 and 48 hours. Similarly, the preferred time range between 6 and 12 hours includes the administration of the dexamethasone dose prior to the first administration of the binding domain and includes 6, 7, 8, 9, 10, 11 and 12 hours,

[0070] In a further embodiment, dexamethasone is administered for a period of two, three, four or five days during which the dose of the binding domain is increased. For example, dexamethasone is administered at a first point of time in a dose of 6 to 40 mg or 6 to 48 mg, preferably 20, 24, 28, 32, 36, 40, 44 or 48 mg, with 20 or 24 mg being

2017202079 29 Mar 2017

preferred, at a second point of time in a dose of preferably 8, 12, 16, 20, or 24 mg, with 16 mg being preferred and/or at a third point of time in a dose of preferably 2, 4, 6, 8, 10 or 12 mg, with 8 mg being preferred. It may also be administered at a fourth point of time or fourth and fifth point of time in a dose of preferably 2, 4, 6, 8, 10 or 12 mg, with 8 mg being preferred.

[0071] It is also envisaged that the human patient of the present invention is characterized by a B/T-cell ratio of less than 1:5 (see PCT/EP2010/066207). As disclosed in great detail in PCT/EP2010/066207, neurological side effects frequently accompany the administration of a CD19xCD3 bispecific antibody in patients which are characterized by a B/T-cell ratio of less than 1:5. The prevention, amelioration or treatment of neurological side effects caused by a CD3 binding domain by way of a GC therapy disclosed herein, is however also applicable to patients which are characterized by a B/T-cell ratio of more than 1:5 (see PCT/EP2010/066207).

[0072] The present invention also relates to a (pharmaceutical) kit or package comprising a GC and/or a CD3 binding domain, and instructions and/or an imprint indicating that the GC is to be employed for the treatment amelioration and/or prophylaxis of neurological adverse events caused by said CD3 binding domain. Said GC and CD3 binding domain are preferably packaged together in one sealed package or kit. It is also envisaged that the package or kit of the present invention, further comprises means to administer the GC and/or Cd3 binding domain to a patient and/or buffers, vials, teflon bags or infusion bags which are normally used for the infusion of therapeutic agents. "Means" thereby includes one or more article(s) selected from the group consisting of a syringe, a hypodermic needle, a cannula, a catheter, an infusion bag for intravenous administration, intravenous vehicles, vials, buffers, stabilizers, written instructions which aid the skilled person in the preparation of the respective doses and infusions of the invention etc.

[0073] By "complete response" is intended an absence of clinically detectable disease with normalization of any previously abnormal imaging studies such as radiographic studies. Such a response preferably persists for at least 4 to 8 weeks, sometimes 6 to 8 weeks or more than 8, 10, 12, 14, 16, 18 or 20 weeks or longer, following treatment according to the invention. Alternatively, an improvement in the disease may be categorized as being a partial response.

Preferably, complete remission in ALL is defined as a blast count below 5% in the bone marrow and recovery of bone marrow function. Detailed remission and response definitions for NHL patients are used according to Cheson et al., 1999, J Clin Oncol. Apr;17(4):1244

5

[0074A] By "partial response" is intended at least about a 50% decrease in all measurable tumor burden (i.e., the number of malignant cells present in the patient, or the measured bulk of tumor masses) in the absence of new lesions and persisting for 4 to 8 weeks or more than 8, 10, 12, 14, 16, 18 or 20 weeks or longer. A "complete response" does, however, not necessarily indicate that a disease has been cured, since a patient may relapse. However, if so, the patient can again be treated with a composition comprising a CD19xCD3 bispecific antibody as described herein.

10

[0074B] Throughout the specification and claims, unless the context requires otherwise, the word "comprise" or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.

15

29 Mar 2017

2017202079

Examples

5 [0075] The following examples illustrate the invention. These examples should not be construed as to limit the scope of this invention. The examples are included for purposes of illustration and the present invention is limited only by the claims.

10 [0076] Patients have been treated with blinatumomab and the co-medication of dexamethasone worked as amelioration, treatment or prophylaxis of neurological and/or psychiatric adverse events.

[0077] The administration of therapeutic dexamethasone was beneficial in that neurological and/or psychiatric symptoms disappeared without necessity to stop treatment by Blinatumomab.

15 [0078] Patient 109-015 with follicular lymphoma received blinatumomab at a dose of $60\mu\text{g}/\text{m}^2/\text{d}$ right from treatment start (i.e. without dose escalation step) for 4 weeks, the patient additionally received dexamethasone on days 4-5: 3x 8mg p.o. and on days 6-7: 1x8mg p.o. in order to treat headache. This treatment cycle had not to be discontinued due to neurological/psychiatric adverse events.

20 [0079] Patient 109-017 with follicular lymphoma received blinatumomab at a dose of $60\mu\text{g}/\text{m}^2/\text{d}$ right from treatment start (i.e. without dose escalation step) for 5 weeks; in order to treat tremor dexamethasone was given p.o.: day 7: 8mg-4mg-4mg; day 8+9: 8mg-4mg-0mg; day 10-11: 4mg-4mg-0mg; day 12-15: 4mg- 0mg – 0mg)). This treatment cycle had not to be discontinued due to neurological/psychiatric adverse events.

30 [0080] Patient 109-026 with MCL received blinatumomab at a dose of $5\mu\text{g}/\text{m}^2/\text{d}$ for 1 week followed by $60\mu\text{g}/\text{m}^2/\text{d}$ for the remaining 6 weeks. The patient received dexamethasone p.o. in order to treat apraxia on day d11: 3x8mg and on day 12: 2x8mg. This treatment cycle had not to be discontinued due to neurological/psychiatric adverse events.

2017202079 29 Mar 2017

- 5 [0081] Patient 135-001 with DLBCL received blinatumomab at 5 $\mu\text{g}/\text{m}^2/\text{d}$ for one week, at 15 $\mu\text{g}/\text{m}^2/\text{d}$ for the 2nd week, and 60 $\mu\text{g}/\text{m}^2/\text{d}$ for 4 more days. The patient received dexamethasone in order to treat tremor at 5 $\mu\text{g}/\text{m}^2/\text{d}$ on d3 (3x8mg) continued in decreasing dose over less than a week. There were no neurological events after escalation to 15ug, which could be explained by a prophylactic effect of dexamethasone. However, on day 4 after step to 60ug/m²/d the patient had to stop due to neurological/psychiatric adverse events which occurred too fast to intervene with dexamethasone.
- 10 [0082] Patient 109-036 with follicular lymphoma was treated with 5 $\mu\text{g}/\text{m}^2/\text{d}$ for the first week followed by treatment at 60 $\mu\text{g}/\text{m}^2/\text{d}$ blinatumomab for the remaining 6 weeks of treatment. The patient experienced neurological adverse events (tremor and apraxia) starting on day 11, i.e. four days after the step to 60 $\mu\text{g}/\text{m}^2/\text{d}$. Dexamethasone was administered i.v. (3x8 mg) and the neurological adverse events resolved under
15 continued treatment with blinatumomab.
- [0083] Patient 109-038 with DLBCL received blinatumomab at 5 $\mu\text{g}/\text{m}^2/\text{d}$ for one week, at 15 $\mu\text{g}/\text{m}^2/\text{d}$ for the 2nd week, and 60 $\mu\text{g}/\text{m}^2/\text{d}$ for the remaining 6 weeks of treatment. On day 15, the patient developed intention tremor which resolved after dexamethasone
20 (3x8mg i.v.) was given (continued blinatumomab treatment).
- [0084] Patient 109-039 with MCL received blinatumomab at 5 $\mu\text{g}/\text{m}^2/\text{d}$ for one week, at 15 $\mu\text{g}/\text{m}^2/\text{d}$ for the 2nd week, and 60 $\mu\text{g}/\text{m}^2/\text{d}$ for the remaining 12 days of treatment. On day 17, the patient developed tremor and slight speech disturbance which resolved after
25 dexamethasone; 3x8mg of dexamethasone i.v. was given (continued blinatumomab treatment).
- [0085] Patient 153-002 with MCL received blinatumomab at 5 $\mu\text{g}/\text{m}^2/\text{d}$ for one week, at 15 $\mu\text{g}/\text{m}^2/\text{d}$ for the 2nd week, and 60 $\mu\text{g}/\text{m}^2/\text{d}$ for one week. The patient developed slight
30 speech problems on day 4 which completely resolved after dexamethasone was given (3x8mg p.o.), in decreasing doses over a few days. On day 22 the patient had to stop treatment due to paresis which disappeared without intervention.

2017202079 29 Mar 2017

Besides the therapeutic use of dexamethasone to mitigate neurological adverse events, dexamethasone was also used as prophylaxis in order to prevent neurological adverse events.

5 [0086] Patient 135-002 with DLBCL received blinatumomab at 5 $\mu\text{g}/\text{m}^2/\text{d}$ for one week, at 15 $\mu\text{g}/\text{m}^2/\text{d}$ for the 2nd week, and 60 $\mu\text{g}/\text{m}^2/\text{d}$ for the remaining 2 weeks and in addition received dexamethasone prophylaxis (dose: 3x8 mg on day of start of blinatumomab treatment and on days of dose escalation steps). The patient did not have to discontinue treatment with blinatumomab due to neurological adverse events.

10

[0087] In patient 155-001 in trial 103-206 with relapsed ALL, was treated with a continuous infusion of blinatumomab at a dosage of 15 $\mu\text{g}/\text{m}^2/\text{d}$ for 16 days. The patient was administered dexamethasone before the administration of blinatumomab in accordance with the teaching of the present application Specifically, dexamethasone prophylaxis was given (8mg before treatment start): The patient did not have to discontinue treatment due to adverse events. The patient achieved a complete remission.

15

Each of the following 5 patients in trial 103-104 (all with DLBCL) were treated with blinatumomab continuous infusion at 5 $\mu\text{g}/\text{m}^2/\text{d}$ for the 1st week, then 15 $\mu\text{g}/\text{m}^2/\text{d}$ for the 2nd week, and then 60 $\mu\text{g}/\text{m}^2/\text{d}$ for the remaining treatment period that could an additional 2–6 weeks. Also each of the following patients were treated prophylactically with 20 mg of dexamethasone at twelve hours and 1 hour before the start of blinatumomab treatment and before each dose increase from the 5 to 15 μg and from 15 to 60 μg .

20

25

[0088] Patients 135-003 did not have to discontinue due to neurological/psychiatric adverse events.

30 [0089] Patient 108-007 did not have to discontinue due to neurological/psychiatric adverse events. This patient achieved a complete remission of the lymphoma.

[0090] Patient 108-008 did not have to discontinue due to neurological/psychiatric adverse events.

2017202079 29 Mar 2017

[0091] Patient 108-009 did not have to discontinue due to neurological/psychiatric adverse events. This patient achieved an objective response of the lymphoma.

5 [0092] Patient 108-010 additionally received 100 mg prednisolone one hour before start of infusion. The patient did not have to discontinue due to neurological/psychiatric adverse events.

10 [0093] Dexamethasone was used as a prophylaxis for neurological adverse events in a 14-y female patient with ALL who was treated with 15 μ g/m²/d blinatumomab on a named patient compassionate use base. 4 weeks in first cycle The patient received dexamethasone on day 1: 3x6 mg p.o., on day 2: 2x 6mg p.o. and on day 3: 1x 6mg). The patient did not have to discontinue treatment due to adverse events. The patient achieved a complete remission.

15 [0094] A Phase I clinical trial was performed in patients with various B-NHL including DLBCL to evaluate the CD19xCD3 bispecific antibody construct in DLBCL patients. Patients were treated for 4-8-weeks by continuous i.v. administration of the antibody with the following step-wise dosing regimen: first week at 5 μ g/m²/d, second week at 15 μ g/m²/d and for the remaining treatment period at 60 μ g/m²/d.

20 Two cohorts each with 6 DLBCL patients were enrolled. The two cohorts solely differed by the dose and schedule of the glucocorticoid medication administered at the beginning of the antibody infusion for mitigation of adverse events.

25 Out of the twelve patients, 5 were male and 7 female. The median age was 57 years (range from 24 to 78 years). Patients had received a median of 4 prior regimens (range from 2-6). All patients had been exposed to rituximab. Eight of the 12 patients had undergone ASCT. International prognostic index (IPI) at screening ranged from 1 to 3 with a median of 2. In the first cohort 100 mg prednisolone was administered 1 hour prior to start; and in the second cohort patients received dexamethasone (3x 8 mg) on days 1, 2, and 3. Before treatment start in the second cohort 20 mg dexamethasone was administered at 12 hours and 1 hour prior to the administration of a CD19xCD3 bispecific antibody construct

30

Although just one DLT (reversible CNS event grade 3) occurred in the prednisolone DLBCL cohort and, thus, the cohort is considered safe, a further DLBCL cohort applying prophylactic dexamethasone (3x8 mg at start of infusion or dose increase and reduction

2017202079 29 Mar 2017

to 3x6 mg or 3x4 mg on the following 2 days, respectively) was opened to optimize management of CNS events. In light of one of the first two patients having a DLT due to a reversible CNS adverse event, a modified "early dexamethasone" schedule (20 mg at -12 to -6 hours and -1 hour, at start of infusion or dose increase, and 3x8 mg during the following 2 days) was introduced to test if earlier and more intensive administration of dexamethasone may ameliorate CNS adverse events. No further DLTs were observed after this adjustment of the dexamethasone schedule. Thus, both the dexamethasone cohort as well as the "early dexamethasone administration" are considered safe. Among a total of 5 patients with DLBCL treated with the "early dexamethasone schedule" no DLT was observed. Therefore, it was concluded that additional administration of "early dexamethasone" is the safest way to administer blinatumomab to patients with DLBCL. Also for patients with "early dexamethasone" objective responses have been observed.

15

Claims

1. Glucocorticoid (GC) when used in the amelioration, treatment or prophylaxis of neurological adverse events caused by the administration of a CD3 binding domain, wherein the GC is administered prior to the administration of the CD3 binding domain.
2. The use of claim 1, wherein said GC is dexamethasone.
3. The use of claim 1 or 2, wherein a first dose of $5\mu\text{g}/\text{m}^2/\text{d}$ of the CD3 binding domain is administered for a first period of time and consecutively a second dose of $15\mu\text{g}/\text{m}^2/\text{d}$ of the CD3 binding domain is administered for a second period of time, wherein the GC is administered prior to the administration of the first and second dose of the CD3 binding domain.
4. The use of claim 3, wherein after a first and second dose of the CD3 binding domain for a first and second period of time, a third dose of $60\mu\text{g}/\text{m}^2/\text{d}$ of the CD3 binding domain is administered, wherein the GC is administered prior to the administration of the first and second and third dose of the CD3 binding domain.
5. The use of claim 1 or 2, wherein a first dose of $5\mu\text{g}/\text{m}^2/\text{d}$ of the CD3 binding domain is administered for a first period of time and consecutively a second dose of $60\mu\text{g}/\text{m}^2/\text{d}$ of the CD3 binding domain is administered for a second period of time, wherein the GC is administered prior to the administration of the first and second dose of the CD3 binding domain.
6. The use of claim 1 or 2, wherein a first dose of $15\mu\text{g}/\text{m}^2/\text{d}$ of the CD3 binding domain is administered for a first period of time and consecutively a second dose of $60\mu\text{g}/\text{m}^2/\text{d}$ of the CD3 binding domain is administered for a second period of time, wherein the GC is administered prior to the administration of the first and second dose of the CD3 binding domain.
7. The use of claim 1 or 2, wherein a flat dose of $5\mu\text{g}/\text{m}^2/\text{d}$ of the CD3 binding domain is administered for up to 8 weeks.
8. The use of claim 1 or 2, wherein a flat dose of $15\mu\text{g}/\text{m}^2/\text{d}$ of the CD3 binding domain is administered for up to 8 weeks.

- 5
- 0
- 5
- 0
- 25
- 30
- 35
9. The use of claim 1 or 2, wherein a flat dose of $60\mu\text{g}/\text{m}^2/\text{d}$ of the CD3 binding domain is administered for up to 8 weeks.
 10. The use of claim 1 or 2, wherein the CD3 binding domain is administered continuously.
 11. The use of any one of claims 1-10, wherein said neurological adverse event is one or more of disturbances of the senses, seizures, encephalopathy, cerebral edema, confusion, ataxia, apraxia, speech disorder, hallucinations, paresis, tremor, headache or disorientation.
 12. The use any one of claims 1-11, wherein the GC is administered subsequent to the appearance of the neurological adverse event.
 13. The use of any one of claims 1-12, wherein said CD3 binding domain is a bispecific single chain antibody.
 14. The use of claim 13, wherein said bispecific single chain antibody is a CD19 x CD3 bispecific single chain antibody.
 15. The use of claim 14, wherein said CD19 x CD3 bispecific single chain antibody is MT103.
 16. The use of any one of claims 1-15, wherein the administration of the CD3 binding domain is to a human patient.
 17. The use of claim 16, wherein said patient is characterized by a B/T-cell ratio of less than 1:5.
 18. The use of any one of claims 1-17, wherein said CD3 binding domain binds to human CD3 epsilon.
 19. A kit when used for the treatment amelioration and/or prophylaxis of neurological adverse events caused by CD3 binding domain: comprising a dexamethasone and a CD3 binding domain and instructions for use and/or an imprint, wherein the dexamethasone is administered prior to the administration of a first dose of the CD3

binding domain and prior to the administration of a second dose and/or a third dose of the CD3 binding domain.

20. A kit when used for the treatment, amelioration and/or prophylaxis of neurological adverse events caused by the CD3 binding domain, comprising a dexamethasone and one or more doses of CD3 binding domain wherein the dexamethasone is administered prior to administration of a first dose of the CD3 binding domain and prior to the administration of a second dose and/or a third dose of the CD3 binding domain, and instructions for use and/or an imprint.

2017202079 29 Mar 2017

eol f - seq l . t x t
SEQUENCE LI STI NG

<110> M cromet GrbH
<120> Prevention of adverse effects caused by CD3 specific binding domains
<130> M M13896PCT
<150> US 61/ 412, 229
<151> 2010-11-10
<160> 22
<170> Patent In version 3.5
<210> 1
<211> 498
<212> PRT
<213> Artificial Sequence
<220>
<223> CD19xCD3 bispecific single chain antibody
<400> 1

Asp Ile G n Leu Thr G n Ser Pro Ala Ser Leu Ala Val Ser Leu G y
1 5 10 15

G n Arg Ala Thr Ile Ser Cys Lys Ala Ser G n Ser Val Asp Tyr Asp
20 25 30

G y Asp Ser Tyr Leu Asn Trp Tyr G n G n Ile Pro G y G n Pro Pro
35 40 45

Lys Leu Leu Ile Tyr Asp Ala Ser Asn Leu Val Ser G y Ile Pro Pro
50 55 60

Arg Phe Ser G y Ser G y Ser G y Thr Asp Phe Thr Leu Asn Ile His
65 70 75 80

Pro Val G u Lys Val Asp Ala Ala Thr Tyr His Cys G n G n Ser Thr
85 90 95

G u Asp Pro Trp Thr Phe G y G y G y Thr Lys Leu G u Ile Lys G y
100 105 110

G y G y G y Ser G y G y G y G y Ser G y G y G y G y Ser G n Val
115 120 125

G n Leu G n G n Ser G y Ala G u Leu Val Arg Pro G y Ser Ser Val
130 135 140

Lys Ile Ser Cys Lys Ala Ser G y Tyr Ala Phe Ser Ser Tyr Trp Met
145 150 155 160

Asn Trp Val Lys G n Arg Pro G y G n G y Leu G u Trp Ile G y G n
165 170 175

2017202079 29 Mar 2017

eol f - seq1 . t x t

I l e T r p P r o G y 180 A s p G y A s p T h r A s n 185 T y r A s n G y L y s P h e 190 L y s G y
L y s A l a T h r 195 L e u T h r A l a A s p G u 200 S e r S e r S e r T h r A l a 205 T y r M e t G n
L e u S e r 210 S e r L e u A l a S e r G u 215 A s p S e r A l a V a l T y r 220 P h e C y s A l a A r g
A r g G u 225 T h r T h r T h r V a l 230 G y A r g T y r T y r T y r 235 A l a M e t A s p T y r T r p 240
G y G n G y T h r T h r 245 V a l T h r V a l S e r S e r 250 G y G y G y G y S e r 255 A s p
I l e L y s L e u G n 260 G n S e r G y A l a G u 265 L e u A l a A r g P r o G y 270 A l a S e r
V a l L y s M e t 275 S e r C y s L y s T h r S e r 280 G y T y r T h r P h e T h r 285 A r g T y r T h r
M e t H i s 290 T r p V a l L y s G n A r g 295 P r o G y G n G y L e u 300 G u T r p I l e G y
T y r 305 I l e A s n P r o S e r A r g 310 G y T y r T h r A s n T y r 315 A s n G n L y s P h e L y s 320
A s p L y s A l a T h r L e u 325 T h r T h r A s p L y s S e r 330 S e r S e r T h r A l a T y r M e t 335
G n L e u S e r S e r 340 L e u T h r S e r G u 345 A s p S e r A l a V a l T y r T y r 350 C y s A l a
A r g T y r T y r 355 A s p A s p H i s T y r C y s 360 L e u A s p T y r T r p G y 365 G n G y T h r
T h r L e u 370 T h r V a l S e r S e r V a l 375 G u G y G y S e r G y 380 G y S e r G y G y
S e r 385 G y G y S e r G y G y 390 V a l A s p A s p I l e G n 395 L e u T h r G n S e r P r o 400
A l a I l e M e t S e r A l a 405 S e r P r o G y G u L y s 410 V a l T h r M e t T h r C y s 415 A r g
A l a S e r S e r S e r 420 V a l S e r T y r M e t A s n 425 T r p T y r G n G n L y s S e r G y 430
T h r S e r P r o 435 L y s A r g T r p I l e T y r 440 A s p T h r S e r L y s V a l 445 A l a S e r G y

eol f - seq1 . t x t

Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu
 450 455 460

Thr Ile Ser Ser Met Gu Ala Gu Asp Ala Ala Thr Tyr Tyr Cys Gn
 465 470 475 480

Gn Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Gu
 485 490 495

Leu Lys

<210> 2
 <211> 1494
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> CD19xCD3 bispecific single chain antibody

<400> 2
 gat at ccagc t gaccagc t ccagct t ct t t ggct gt gt ct ct agggca gagggccacc 60
 at ct cct gca aggcagcca aagt gt t gat t at gat ggt g at agt t at t t gaact ggt ac 120
 caacagat t c caggacagcc acccaaact c ct cat ct at g at gcat ccaa t ct agt t t ct 180
 gggat cccac ccaggt t t ag t ggcagt ggg t ct gggacag act t caccct caacat ccat 240
 cct gt ggaga aggt ggat gc t gcaacct at cact gt cagc aaagt act ga ggat ccgt gg 300
 acgt t cggt g gagggaccaa gct cgagat c aaaggt ggt g gt ggt t ct gg cggcggcggc 360
 t ccggt ggt g gt ggt t ct ca ggt gcagct g cagcagt ct g gggct gagct ggt gaggcct 420
 gggc cct cag t gaagat t t c ct gcaaggct t ct ggct at g cat t cagt ag ct act ggat g 480
 aact ggt ga agcagaggcc t ggacagggt ct t gagt gga t t ggacagat t t ggcct gga 540
 gat ggt gat a ct aact acaa t ggaaagt t c aagggt aaag ccact ct gac t gcagacgaa 600
 t cct ccagca cagcct acat gcaact cagc agcct agcat ct gaggact c t gcggt ct at 660
 t t ct gt gcaa gacgggagac t acgacggt a ggccgt t at t act at gct at ggact act gg 720
 ggccaaggga ccacggt cac cgt ct cct cc ggaggt ggt g gat ccgat at caaact gcag 780
 cagt cagggg ct gaact ggc aagacct ggg gcct cagt ga agat gt cct g caagact t ct 840
 ggct acacct t t act aggt a cacgat gcac t ggggt aaaac agaggcct gg acagggt ct g 900
 gaat ggat t g gat acat t aa t cct agccgt ggt t at act a at t acaat ca gaagt t caag 960
 gacaaggcca cat t gact ac agacaaat cc t ccagcacag cct acat gca act gagcagc 1020
 ct gacat ct g aggact ct gc agt ct at t ac t gt gcaagat at t at gat ga t cat t act gc 1080
 ct t gact act ggggccaagg caccact ct c acagt ct cct cagt cgaagg t ggaagt gga 1140
 ggt t ct ggt g gaagt ggagg t t caggt gga gt cgacgaca t t cagct gac ccagt ct cca 1200
 gcaat cat gt ct gcat ct cc aggggagaag gt caccat ga cct gcagagc cagt t caagt 1260

eol f - seq| . t x t

```

gt aagt t aca t gaact ggt a ccagcagaag t caggcacct cccccaaaag at ggat t t at      1320
gacacat cca aagt ggct t c t ggagt ccct t at cgct t ca gt ggcagt gg gt ct gggacc      1380
t cat act ct c t cacaat cag cagcat ggag gct gaagat g ct gccact t a t t act gccaa      1440
cagt ggagt a gt aacccgct cacgt t cggt gct gggacca agct ggagct gaaa      1494

```

```

<210> 3
<211> 124
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> VH anti CD19

```

```

<400> 3

```

```

G n Val G n Leu G n G n Ser G y Al a G u Leu Val Arg Pro G y Ser
1          5          10          15

```

```

Ser Val Lys Ile Ser Cys Lys Al a Ser G y Tyr Al a Phe Ser Ser Tyr
          20          25          30

```

```

Trp Met Asn Trp Val Lys G n Arg Pro G y G n G y Leu G u Trp Ile
          35          40          45

```

```

G y G n Ile Trp Pro G y Asp G y Asp Thr Asn Tyr Asn G y Lys Phe
          50          55          60

```

```

Lys G y Lys Al a Thr Leu Thr Al a Asp G u Ser Ser Ser Thr Al a Tyr
65          70          75          80

```

```

Met G n Leu Ser Ser Leu Al a Ser G u Asp Ser Al a Val Tyr Phe Cys
          85          90          95

```

```

Al a Arg Arg G u Thr Thr Thr Val G y Arg Tyr Tyr Tyr Al a Met Asp
          100          105          110

```

```

Tyr Trp G y G n G y Thr Thr Val Thr Val Ser Ser
          115          120

```

```

<210> 4
<211> 372
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> VH anti CD19

```

```

<400> 4

```

```

caggt gcagc t gcagcagt c t ggggct gag ct ggt gaggc ct gggc cct c agt gaagat t      60
t cct gcaagg ct t ct ggct a t gcat t cagt agct act gga t gaact gggc gaagcagagg      120
cct ggacagg gt ct t gagt g gat t ggacag at t t ggcct g gagat ggt ga t act aact ac      180
aat ggaaagt t caagggt aa agccact ct g act gcagacg aat cct ccag cacagcct ac      240

```

eol f - seq| . t x t

at gcaact ca gcagcct agc at ct gaggac t ct gcggt ct at t t ct gt gc aagacgggag 300
 act acgacgg t aggccgt t a t t act at gct at ggact act ggggccaagg gaccacggt c 360
 accgt ct cct cc 372

<210> 5
 <211> 111
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> VL anti CD19

<400> 5

Asp Ile G n Leu Thr G n Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1 5 10 15

G n Arg Ala Thr Ile Ser Cys Lys Ala Ser G n Ser Val Asp Tyr Asp
 20 25 30

Gly Asp Ser Tyr Leu Asn Trp Tyr G n G n Ile Pro Gly G n Pro Pro
 35 40 45

Lys Leu Leu Ile Tyr Asp Ala Ser Asn Leu Val Ser Gly Ile Pro Pro
 50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His
 65 70 75 80

Pro Val Gu Lys Val Asp Ala Ala Thr Tyr His Cys G n G n Ser Thr
 85 90 95

Gu Asp Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Gu Ile Lys
 100 105 110

<210> 6
 <211> 333
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> VL anti CD19

<400> 6

gat at ccagc t gaccagc t ccagct t ct t t ggct gt gt ct ct agggca gagggccacc 60
 at ct cct gca aggccagcca aagt gt t gat t at gat ggt g at agt t at t t gaact ggt ac 120
 caacagat t c caggacagcc acccaaact c ct cat ct at g at gcat ccaa t ct agt t t ct 180
 gggat cccac ccaggt t t ag t ggca ggt ggg t ct gggacag act t caccct caacat ccat 240
 cct gt ggaga aggt ggat gc t gcaacct at cact gt cagc aaagt act ga ggat ccgt gg 300
 acgt t cggg g gagggaccaa gct cgagat c aaa 333

<210> 7

29 Mar 2017 2017202079

eol f - seq| . t x t

<211> 119
<212> PRT
<213> Artificial Sequence

<220>
<223> VH anti CD3

<400> 7

Asp Ile Lys Leu G n G n Ser G y Ala G u Leu Ala Arg Pro G y Ala
1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser G y Tyr Thr Phe Thr Arg Tyr
20 25 30

Thr Met His Trp Val Lys G n Arg Pro G y G n G y Leu G u Trp Ile
35 40 45

G y Tyr Ile Asn Pro Ser Arg G y Tyr Thr Asn Tyr Asn G n Lys Phe
50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met G n Leu Ser Ser Leu Thr Ser G u Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp G y G n G y
100 105 110

Thr Thr Leu Thr Val Ser Ser
115

<210> 8
<211> 357
<212> DNA
<213> Artificial Sequence

<220>
<223> VH anti CD3

<400> 8

gat at caaac tgcagcagt c aggggct gaa ct ggcaagac ct ggggcct c agt gaagat g 60
t cct gcaaga ct t ct ggct a cacct t t act aggt acacga t gcact gggg t aaaacagagg 120
cct ggacagg gt ct ggaat g gat t ggat ac at t aat cct a gccgt ggt t a t act aat t ac 180
aat cagaagt t caaggacaa ggccacat t g act acagaca aat cct ccag cacagcct ac 240
at gcaact ga gcagcct gac at ct gaggac t ct gcagt ct at t act gt gc aagat at t at 300
gat gat cat t act gcct t ga ct act ggggc caaggcacca ct ct cacagt ct cct ca 357

<210> 9
<211> 106
<212> PRT
<213> Artificial Sequence

29 Mar 2017 2017202079

eol f - seq| . t x t

<220>
<223> VL anti CD3

<400> 9

Asp Ile G n Leu Thr G n Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
1 5 10 15

G u Lys Val Thr Met Thr Cys Arg Ala Ser Ser Ser Val Ser Tyr Met
20 25 30

Asn Trp Tyr G n G n Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
35 40 45

Asp Thr Ser Lys Val Ala Ser Gly Val Pro Tyr Arg Phe Ser Gly Ser
50 55 60

G y Ser G y Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met G u Ala G u
65 70 75 80

Asp Ala Ala Thr Tyr Tyr Cys G n G n Trp Ser Ser Asn Pro Leu Thr
85 90 95

Phe G y Ala G y Thr Lys Leu G u Leu Lys
100 105

<210> 10
<211> 318
<212> DNA
<213> Artificial Sequence

<220>
<223> VL anti CD3

<400> 10
gacattcagc t gaccagc t ccagcaat c atgtctgcat ctccagggga gaaggtcacc 60
atgacctgca gagccagtc aagtgt aagt t acatgaact ggtaccagca gaagtcaggc 120
acctccccca aaagatggat ttatgacaca tccaaagtg gcttctggagt cccttatcgc 180
ttcagtgga gtgggtctgg gacctcat ac tctctcaca t cagcagcat ggaggctgaa 240
gatgctgcca cttat t act g ccaacagtg g agt agt aacc cgctcacgt t cggtgctggg 300
accaagctgg agctgaaa 318

<210> 11
<211> 10
<212> PRT
<213> artificial

<220>
<223> CD3 CDR- H1

<400> 11

G y Tyr Thr Phe Thr Arg Tyr Thr Met His
1 5 10

<210> 12
 <211> 17
 <212> PRT
 <213> a r t i f i c i a l

<220>
 <223> CD3 CDR- H2

<400> 12

Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn G n Lys Phe Lys
 1 5 10 15

Asp

<210> 13
 <211> 10
 <212> PRT
 <213> a r t i f i c i a l

<220>
 <223> CD3 CDR- H3

<400> 13

Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr
 1 5 10

<210> 14
 <211> 10
 <212> PRT
 <213> a r t i f i c i a l

<220>
 <223> CD3 CDR- L1

<400> 14

Arg Ala Ser Ser Ser Val Ser Tyr Met Asn
 1 5 10

<210> 15
 <211> 7
 <212> PRT
 <213> a r t i f i c i a l

<220>
 <223> CD3 CDR- L2

<400> 15

Asp Thr Ser Lys Val Ala Ser
 1 5

<210> 16
 <211> 9
 <212> PRT
 <213> a r t i f i c i a l

<220>
 <223> CD3 CDR- L3

<400> 16

G n G n Trp Ser Ser Asn Pro Leu Thr
1 5

<210> 17

<211> 10

<212> PRT

<213> a r t i f i c i a l

<220>

<223> CD19 CDR- H1

<400> 17

G y Tyr Ala Phe Ser Ser Tyr Trp Met Asn
1 5 10

<210> 18

<211> 17

<212> PRT

<213> a r t i f i c i a l

<220>

<223> CD19 CDR- H2

<400> 18

G n Ile Trp Pro Gly Asp Gly Asp Thr Asn Tyr Asn Gly Lys Phe Lys
1 5 10 15

G y

<210> 19

<211> 15

<212> PRT

<213> a r t i f i c i a l

<220>

<223> CD19 CDR- H3

<400> 19

Arg G u Thr Thr Thr Val Gly Arg Tyr Tyr Tyr Ala Met Asp Tyr
1 5 10 15

<210> 20

<211> 15

<212> PRT

<213> a r t i f i c i a l

<220>

<223> CD19 CDR- L1

<400> 20

Lys Ala Ser G n Ser Val Asp Tyr Asp Gly Asp Ser Tyr Leu Asn
1 5 10 15

<210> 21

29 Mar 2017

2017202079

eol f - seq| . t x t

<211> 7
<212> PRT
<213> artificial

<220>
<223> CD19 CDR- L2

<400> 21

Asp Al a Ser Asn Leu Val Ser
1 5

<210> 22
<211> 9
<212> PRT
<213> artificial

<220>
<223> CD19 CDR- L3

<400> 22

G n G n Ser Thr G u Asp Pro Trp Thr
1 5