

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2004299286 B2**

- (54) Title
1-N-phenyl-amino-1H-imidazole derivatives and pharmaceutical compositions containing them
- (51) International Patent Classification(s)
C07D 233/54 (2006.01) **C07D 419/12** (2006.01)
C07D 409/12 (2006.01)
- (21) Application No: **2004299286** (22) Date of Filing: **2004.12.15**
- (87) WIPO No: **WO05/058842**
- (30) Priority Data
- | (31) Number | (32) Date | (33) Country |
|-------------------|-------------------|--------------|
| 03293152.9 | 2003.12.15 | EP |
| 04292681.6 | 2004.11.12 | EP |
- (43) Publication Date: **2005.06.30**
(44) Accepted Journal Date: **2011.06.30**
- (71) Applicant(s)
Laboratoire Theramex
- (72) Inventor(s)
Puccio, Francois;Clerc, Thierry;Duc, Igor;Blot, Christian;Rondot, Benoit;Lafay, Jean;Shields, Jacqueline;Duranti, Eric;Maillos, Philippe;Bonnet, Paule
- (74) Agent / Attorney
Spruson & Ferguson, Level 35 St Martins Tower 31 Market Street, Sydney, NSW, 2000
- (56) Related Art
WO2002068395
AU2003298259 (family member equivalent of EP1431292)
US5071861
EP0641785

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
30 June 2005 (30.06.2005)

PCT

(10) International Publication Number
WO 2005/058842 A1

(51) International Patent Classification⁷: **C07D 233/56**,
409/12, 419/12, A61K 31/4164, 31/5415, A61P 5/00,
35/00

François [FR/FR]; 74 Avenue du Mont Alban, F-06300
Nice (FR). **BLÖT, Christian** [FR/FR]; 1582 Route de
Cagnes, F-06480 La Colle Sur Loup (FR). **MAILLOS,**
Philippe [FR/FR]; EN CASSE, F-81500 Lavaur (FR).

(21) International Application Number:
PCT/EP2004/014847

(74) Agents: **GILLARD, Marie Louise** et al.; Cabinet BEAU
DE LOMENIE, 158 Rue de l'Université, F-75340 Paris
cedex 07 (FR).

(22) International Filing Date:
15 December 2004 (15.12.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03293152.9 15 December 2003 (15.12.2003) EP
04292681.6 12 November 2004 (12.11.2004) EP

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AF, AG, AI, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(71) Applicant (for all designated States except US): **LAB-
ORATOIRE THERAMEX** [MC/MC]; 6 Avenue Prince
Héréditaire Albert, MC-98000 MONACO (MC).

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (for US only): **LAFAY, Jean**
[FR/FR]; 1 Rue Clément Ader, F-06100 Nice (FR). **RON-
DOT, Benoît** [FR/FR]; Parc de Montfort, lot 8, Chemin
du Caminon, F-06480 La Colle Sur Loup (FR). **BON-
NET, Paule** [FR/FR]; Villa les Rocailles, Chemin Glena,
37 Route de Sospel, F-06500 Menton (FR). **CLERC,**
Thierry [FR/FR]; Les Balcons d'Eze, 601 Avenue de Ver-
dun, F-06360 Eze (FR). **SHIELDS, Jacqueline** [FR/FR];
6 Avenue George Sand, F-06100 Nice (FR). **DUC, Igor**
[FR/FR]; Villa Casimir, 26 Rue Borniol, F-06400 Cannes
(FR). **DURANTI, Eric** [FR/FR]; 258 Avenue des Fi-
lagnes, F-06700 Saint Laurent Du Var (FR). **PUCCIO,**

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 1-N-PHENYL-AMINO-1H-IMIDAZOLE DERIVATIVES AND PHARMACEUTICAL COMPOSITIONS CONTAIN-
ING THEM



(57) Abstract: The invention relates to the compounds of formula (I), in
which R₁, R₂, R₃, R₄, Q and Z are as defined in the specification. The inven-
tion also relates to the pharmaceutical compositions containing these com-
pounds. The invention generally relates to the field of hormone- and non
hormone-dependent cancer and endocrine disorders.

1-N-phenyl-amino-1H-imidazole derivatives and pharmaceutical compositions containing them.

The present invention relates to 1-N-phenylamino-1H-imidazole derivatives and to pharmaceutical compositions containing them.

- 5 The invention generally relates to the field of hormone- and non hormone-dependent cancer and endocrine disorders.

Aromatase is the physiological enzyme responsible for the specific conversion of androgens such as androstenedione or testosterone, into estrogens such as estrone and estradiol, respectively (Simpson ER et al., Endocrine Reviews, 1994, 10 15 : 342-355). Inhibition of aromatase is, therefore, a strategy of choice to interfere with normal or pathological estrogen-induced or estrogen-dependent biological processes such as female sexual differentiation, ovulation, implantation, pregnancy, breast and endometrial cell proliferation as well as regulations of spermatogenesis or prostate cell proliferation in male or of non-reproductive functions such as bone formation or immune T cell and cytokine 15 balance (see Simpson ER et al., Recent Progress in Hormone Research, 1997, 52 : 185-213 and the whole issues of Endocrine Related Cancer (1999, volume 6, n°2) and Breast Cancer Research Treatment (1998, volume 49, supplement n°1)).

- 20 The enzyme steroid sulfatase (E.C. 3.1.6.2., STS) catalyses the hydrolysis of estrone sulfate to estrone and the DHEA sulfate to DHEA (Dibbelt L, Biol. Chem, Hoppe-Seyler, 1991, 372, 173-185 and Stein C, J. Biol. Chem., 1989, 264, 13865-13872).


The steroid sulfatase pathway has been the focus of recent interest in the 25 context of breast cancer, with regard to the local intra-tissue formation of estrogens from the abundant circulating pool of estrone sulfate (E_1S) (Pasqualini JR, J. Steroid Biochem. Mol. Biol., 1999, 69, 287-292 and Purohit A, Mol. Cell. Endocrinol., 2001, 171, 129-135).

- Inhibition of this enzyme would prevent E_1S from yielding free estrone (E_1), 30 which can be transformed into estradiol (E_2) by enzymatic reduction. In addition to the estrone sulfatase pathway, it is now believed that another potent estrogen, androstenediol (adiol) obtained from DHEA after hydrolysis of DHEA-S,


could be another important contributor, in the support of growth and development of hormone-dependent breast tumors.

The formation of estrogens in women is schematically represented in figure 1.

In patients with hormone-dependent cancers, aromatase inhibitors are currently
5 used to prevent estrogen synthesis. However, clinical trials showed a relative lack of efficacy for patients with estrogen receptor-positive tumors (Castiglione-Gertsch M, Eur. J. Cancer, 1996, 32A, 393-395 and Jonat W, Eur. J. Cancer, 1996, 32A, 404-412). As an explanation, steroid sulfatase pathway could be another important route for estrogen formation in breast tumors.
10 EMATE (Ahmed S, Curr. Med. Chem., 2002, 9, 2, 263-273), estrone-3-sulfamate, is the historical standard steroidal sulfatase inhibitor but has the major drawback of being estrogenic because of its mechanism of inhibition: the sulfamate moiety is cleaved during the process of enzyme inactivation, which releases E_1 , not from E_1S but from EMATE itself (Ahmed S, J. Steroid Biochem. Mol. Biol., 2002, 80,
15 429-440).




Estradiol



EMATE


Other non-steroid sulfamate compounds which release derivatives without estrogenic properties have been presented as acceptable drug candidates such
20 as 6,6,7-COUMATE, a standard non-estrogenic sulfatase inhibitor from the literature (Purohit A, Cancer Res, 2000, 60, 3394-3396).

**6,6,7 COUMATE**

Human carbonic anhydrases catalyse the conversion between carbon dioxide (CO₂) and the bicarbonate ion (HCO₃⁻), and are involved in physiological and pathological processes. They include hormone-dependent and non-hormone-dependent cancerogenesis, metastasis invasive process and hypoxic tumors that express these enzymes which are less responsive to classical chemo/radiotherapy inhibitors. In particular, EMATE was found to have a human carbonic anhydrase inhibitory potency similar to that of acetazolamide, a well-known sulfonamide human carbonic anhydrase inhibitor (Winum J and al., J. Med. Chem. 2003, 46, 2197-2204).

It is therefore of particular interest to have compounds with at least one, preferably at least two of the following activities: aromatase inhibition, steroid sulfatase inhibition and carbonic anhydrase inhibition.

Recently, B. Potter et al. (J. Med. Chem., 46, 2003, 3193-3196) reported that sulfamoylated-derivatives of the aromatase inhibitor YM 511 inhibited sulfatase and aromatase activity in JEG-3 cells.

**YM 511**

Compounds presented as useful for the treatment of estrogen-dependent diseases, are described in US 2003/0008862A. Imidazole derivatives with anti-aromatase properties are described in WO 2004/054983.


It has now been found that imidazole derivatives which contain a 1-N-phenylamino group, demonstrate an unexpectedly high potency to inhibit aromatase and/or steroid sulfatase and/or carbonic anhydrase.

Accordingly, one object of this invention is to provide 1-N-phenylamino-1H-imidazole derivatives, which are potent aromatase and/or steroid sulfatase and/or carbonic anhydrase inhibitors.

Another object of this invention is to provide a pharmaceutical composition containing, as active ingredient, a 1-N-aryl-amino-1H-imidazole derivative as depicted below.


10 A further object of this invention is to provide the use of 1-N-phenylamino-1H-imidazole derivatives in the manufacture of a medicament for treating or preventing various diseases and for managing reproductive functions in women, in men as well as in female and male wild or domestic animals.

15 The 1-N-phenylamino-1H-imidazole derivatives of this invention are represented by the following general formula (I):




and acid addition salts and stereoisomeric forms thereof, wherein :


- R_1 and R_2 are each independently hydrogen, a (C_1-C_6) alkyl or a (C_3-C_8) cycloalkyl ; or R_1 and R_2 together form a saturated or unsaturated 5-, 6- or 7- membered carbocyclic ring;
- Q is $(CH_2)_m-X-(CH_2)_n-A$;
- A is a direct link, O, S, SO, SO_2 , NR_5 ;
- X is a direct link, CF_2 , O, S, SO, SO_2 , C(O), NR_5 or CR_6R_7 ;
- Z is a group selected from:



- m and n are each independently 0, 1, 2, 3 or 4;
- p is 1, 2, 3 or 4;
- q is 0, 1 or 2;
- the dotted line means that R_8 and/or R_9 can be on any position of the benzothiophene ring;
- R_3 and R_8 are each independently hydrogen or a hydroxy, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, (C_1-C_6) alkylthio, (C_1-C_6) alkylsulfonyl, acyl, (C_1-C_6) alkoxycarbonyl, carboxamido, $OPO(OR_{10})_2$, $NR_{10}R_{11}$, $SO_2NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, OSO_2OR_{10} , SO_2OR_{10} , $SSO_2NR_{10}R_{11}$, $CF_2SO_2OR_{10}$, $CF_2SO_2NR_{10}R_{11}$, CF_2 -tetrazolyl or $NR_{12}SO_2NR_{10}R_{11}$, $OSO_2NR_{12}SO_2NR_{10}R_{11}$, CO_2R_{10} , $CONR_{10}R_{11}$, $OCHO$, $OCONR_{10}R_{11}$, $OCSNR_{10}R_{11}$, $SCONR_{10}R_{11}$, $SCSNR_{10}R_{11}$, tetrazolyl, $NR_{12}CONR_{10}R_{11}$, NR_{10} -CHO group;
- when Q-Z is




- n is 0, 1 or 2 and p is 1, one of R_3 and R_8 is a hydroxy, nitro, $OPO(OR_{10})_2$, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, OSO_2OR_{10} , SO_2OR_{10} , $SSO_2NR_{10}R_{11}$, $CF_2SO_2OR_{10}$, $CF_2SO_2NR_{10}R_{11}$, CF_2 -tetrazolyl, $NR_{12}SO_2NR_{10}R_{11}$, $OSO_2NR_{10}SO_2NR_{11}R_{12}$, CO_2R_{10} , $CONR_{10}R_{11}$, $OCHO$, $CONR_{10}R_{11}$, $OCSNR_{10}R_{11}$, $SCONR_{10}R_{11}$, $SCSNR_{10}R_{11}$, tetrazolyl, $NR_{12}CONR_{10}R_{11}$, NR_{10} -CHO group and the other is hydrogen or a hydroxy, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, (C_1-C_6) alkylthio, (C_1-C_6) alkylsulfonyl, acyl, (C_1-C_6) alkoxycarbonyl, carboxamido, $NR_{10}R_{11}$, $SO_2NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, OSO_2OR_{10} , SO_2OR_{10} , $SSO_2NR_{10}R_{11}$, $CF_2SO_2OR_{10}$, $CF_2SO_2NR_{10}R_{11}$, CF_2 -tetrazolyl, $NR_{12}SO_2NR_{10}R_{11}$, $OSO_2NR_{12}SO_2NR_{10}R_{11}$, CO_2R_{10} , $CONR_{10}R_{11}$, $OCHO$, $CONR_{10}R_{11}$, $OCSNR_{10}R_{11}$, $SCONR_{10}R_{11}$, $SCSNR_{10}R_{11}$, tetrazolyl, $NR_{12}CONR_{10}R_{11}$, NR_{10} -CHO group;
- R_4 and R_9 are each independently hydrogen or a hydroxy, cyano, halogen, nitro, $OPO(OR_{10})_2$, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, (C_1-C_6) alkylthio, (C_1-C_6) alkylsulfonyl, acyl, (C_1-C_6) alkoxycarbonyl, carboxamido, $NR_{10}R_{11}$, $SO_2NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, OSO_2OR_{10} , SO_2OR_{10} , $SSO_2NR_{10}R_{11}$, $CF_2SO_2OR_{10}$, $CF_2SO_2NR_{10}R_{11}$, CF_2 -tetrazolyl, $NR_{12}SO_2NR_{10}R_{11}$, $OSO_2NR_{12}SO_2NR_{10}R_{11}$, CO_2R_{10} , CHO , $CONR_{10}R_{11}$, $OCHO$, $CONR_{10}R_{11}$, $OCSNR_{10}R_{11}$, $SCONR_{10}R_{11}$, $SCSNR_{10}R_{11}$, tetrazolyl, $NR_{12}CONR_{10}R_{11}$, NR_{10} -CHO group;
 - when p is 2, 3 or 4 the R_9 s can be the same or different;
 - R_6 and R_7 are independently hydrogen, halogen, a (C_1-C_6) alkyl or a (C_3-C_8) cycloalkyl;
 - R_5 , R_{10} , R_{11} and R_{12} are each independently hydrogen, hydroxy, a (C_1-C_6) alkyl, or a (C_3-C_8) cycloalkyl; R_{10} can also be a salt; R_{10} and R_{11} can also form, together with the nitrogen atom to which they are bound, a 5- to 7-membered heterocycle containing one or two heteroatoms selected from O, S and N;
 - when Z is



and p is 1,


then R_8 and R_9 can also form together with the phenyl ring a benzoxathiazine dioxide, a dihydrobenzoxathiazine dioxide, a benzoxathiazinone dioxide, a benzoxathiazole dioxide, a benzoxadithiadiazine tetraoxide, a benzodithiazine tetraoxide or a benzodioxadithiine tetraoxide;

- when Z is






R_3 and R_4 together with the phenyl ring bearing them can also form a benzofurane or a N-methylbenzotriazole, provided that when p is 1 and Q is $(CH_2)_n$, then R_8 and R_9 are independently a hydroxy, nitro, $OPO(OR_{10})_2$, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, OSO_2OR_{10} , SO_2OR_{10} , $SSO_2NR_{10}R_{11}$, $CF_2SO_2OR_{10}$, $CF_2SO_2NR_{10}R_{11}$, CF_2 -tetrazolyl, $NR_{12}SO_2NR_{10}R_{11}$, $OSO_2NR_{12}SO_2NR_{10}R_{11}$, CO_2R_{10} , $CONR_{10}R_{11}$, $OCHO$, $OCONR_{10}R_{11}$, $OCSNR_{10}R_{11}$, $SCONR_{10}R_{11}$, $SCSNR_{10}R_{11}$, tetrazolyl, $NR_{12}CONR_{10}R_{11}$, or $NR_{10}-CHO$ group.

According to an aspect of the present invention there is provided an imidazole derivative of formula (I) :




or an acid addition salt or stereoisomeric form thereof, wherein :

- R_1 and R_2 are each independently hydrogen or a (C_1-C_6) alkyl;
- Q is selected from a direct link, $C(O)$, SO_2 , $CONH$, $C(O)(CH_2)_n$, $(CH_2)_n(O)$ or $(CH_2)_n$ in which n is 0, 1 or 2.
- Z is a group selected from:




- p is 1, 2, 3 or 4;
- q is 0, 1 or 2;
- the dotted line means that R₈ and/or R₉ can be on any position of the benzothiophene ring;

- 5
- one of R₃ and R₈ is hydroxy, cyano, (C₁-C₆)alkoxy or OSO₂NR₁₀R₁₁; and
 - the other is hydrogen or a hydroxy, halogen, nitro, cyano, (C₁-C₆)alkoxy, NR₁₀R₁₁, SO₂NR₁₀R₁₁, OSO₂NR₁₀R₁₁, NR₁₂SO₂NR₁₀R₁₁, OSO₂NR₁₀SO₂NR₁₁R₁₂ group;
 - R₄ is hydrogen and R₉ is hydrogen, hydroxy, cyano, halogen, nitro, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, acyl, NR₁₀R₁₁, OSO₂NR₁₀R₁₁, NR₁₂SO₂NR₁₀R₁₁, CO₂R₁₀ group;
- 10
- when Q-Z is



n is 0, 1 or 2 and p is 1, one of R₃ and R₈ is a hydroxy or OSO₂NR₁₀R₁₁;

- 15
- when p is 2, 3 or 4 the R₉s are other than hydrogen and can be the same or different;
 - R₁₀, R₁₁ and R₁₂ are each independently hydrogen or a (C₁-C₆)alkyl;
 - when Z is



and p is 1,

- 20
- then R₈ and R₉ can also form together with the phenyl ring a benzoxathiazine dioxide, a dihydrobenzoxathiazine dioxide or a benzoxathiazole dioxide.

The present invention also concerns the addition salts of the invention compounds or the stereoisomeric forms thereof, when they exist.

- 25
- In the description and claims, the term "(C₁-C₆)alkyl" is understood as meaning a linear or branched hydrocarbon chain having 1 to 6 carbon atoms. A (C₁-C₆)alkyl radical is for example a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl,

isopentyl or hexyl radical. Preferred alkyl radicals are those having 1, 2 or 3 carbon atoms.

The term "halogen" is understood as meaning a chlorine, bromine, iodine or fluorine atom.

5 The term "(C₃-C₈)cycloalkyl" is understood as meaning a saturated monocyclic hydrocarbon having 3 to 8 carbon atoms. A (C₃-C₈)cycloalkyl radical is for example a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl radical.

The term "(C₁-C₆)alkoxy" is understood as meaning a group OR in which R is a (C₁-C₆)alkyl as defined above. A (C₁-C₆)alkoxy radical is for example a methoxy, ethoxy,
10 propoxy, isopropoxy, butoxy, isobutoxy, tert-butoxy, n-pentyloxy or

isopentyloxy radical. Preferred alkoxy radicals are those having 1, 2 or 3 carbon atoms.

The term "acyl" is understood as meaning a group $\text{R}'-\text{C}(=\text{O})$ in which R' is hydrogen or a

(C₁-C₄)alkyl wherein the term "alkyl" is as defined above. An acyl radical is for example a formyl, an acetyl, a propionyl, a butyryl or a valeryl radical. Preferred acyl radicals are formyl and acetyl.

In the definition of R₁₀, a "salt" is understood as meaning an alkali metal salt or alkaline earth metal salt, such as a sodium, potassium, magnesium or calcium salt, or a salt with an ammonium or with an organic amine such as triethylamine, ethanolamine or tris-(2-hydroxyethyl)amine. In the context of the invention, this applies to groups having an OR₁₀ moiety.

The 5- to 7-membered heterocycle can be saturated or unsaturated, and includes for example tetrazole, triazole, pyrazole, pyrazolidine, imidazole, imidazolidine, piperidine, piperazine, morpholine, pyrrolidine.

Compounds of formula (I) form acid addition salts, for example with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like or with organic carboxylic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid and the like. Especially preferred are those salts which are pharmaceutically acceptable.

Among the compounds of formula (I), those which fulfil at least one of the following conditions are preferred:

- 25 • R₃ and R₈ are each independently hydrogen or a hydroxy, cyano, halogen, nitro, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, (C₁-C₆)alkylthio, (C₁-C₆)alkylsulfonyl, acyl, (C₁-C₆)alkoxycarbonyl, carboxamido, NR₁₀R₁₁, SO₂NR₁₀R₁₁, OSO₂NR₁₀R₁₁, OSO₂NR₁₂SO₂NR₁₀R₁₁, OCHO, NR₁₂SO₂NR₁₀R₁₁ group;
- 30 • R₄ and R₉ are each independently hydrogen or a hydroxy, cyano, halogen, nitro, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, (C₁-C₆)alkylthio,

(C₁-C₆)alkylsulfonyl, acyl, (C₁-C₆)alkoxycarbonyl, carboxamido, NR₁₀R₁₁, OSO₂NR₁₀R₁₁, CO₂R₁₀, CHO, NR₁₂SO₂NR₁₀R₁₁ group;

- R₁ and R₂ are each independently hydrogen or (C₁-C₆)alkyl;
- R₁₀, R₁₁ and R₁₂ are each independently hydrogen or (C₁-C₆)alkyl.

5.

Particularly preferred compounds of a formula (I) are those in which:

- one of R₃ and R₈ is a hydroxy, nitro, NR₁₀R₁₁, OSO₂NR₁₀R₁₁ or NR₁₂SO₂NR₁₀R₁₁ group; and
- the other is hydrogen or a hydroxy, cyano, halogen, nitro, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, (C₁-C₆)alkylthio, (C₁-C₆)alkylsulfonyl, acyl, (C₁-C₆)alkoxycarbonyl, carboxamido, NR₁₀R₁₁, OSO₂NR₁₀R₁₁, NR₁₂SO₂NR₁₀R₁₁ group.

A particularly preferred group of these compounds are those in which:

- one of R₃ and R₈ is hydroxy, cyano, (C₁-C₆)alkoxy or OSO₂NR₁₀R₁₁; and
- the other is hydrogen or a hydroxy, halogen, nitro, cyano, (C₁-C₆)alkoxy, NR₁₀R₁₁, SO₂NR₁₀R₁₁, OSO₂NR₁₀R₁₁, NR₁₂SO₂NR₁₀R₁₁, OSO₂NR₁₀SO₂NR₁₁R₁₂ group.

20 Preferably, the compounds of formula (I) are those in which:

- one of R₃ and R₈ is cyano; and
- the other is hydrogen or a hydroxy, halogen, nitro, (C₁-C₆)alkoxy, trifluoromethyl, NR₁₀R₁₁, SO₂NR₁₀R₁₁, OSO₂NR₁₀R₁₁, NR₁₂SO₂NR₁₀R₁₁ group.

25 Further preferred compounds are those in which:

- one of R₄ and R₉ is hydrogen or a hydroxy, cyano, OSO₂NR₁₀R₁₁ group; and
- the other is hydrogen or a hydroxy, cyano, halogen, nitro, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, trifluoromethyl, NR₁₀R₁₁, OSO₂NR₁₀R₁₁, CO₂R₁₀, CHO, NR₁₂SO₂NR₁₀R₁₁ group.

30


An other group of preferred compounds are those in which:

- R₄ is hydrogen, hydroxy, cyano, or OSO₂NR₁₀R₁₁;

- R_9 is a hydrogen or a hydroxy, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, CO_2R_{10} , CHO group.

Especially preferred compounds of formula (I) are those where:

- 5
- R_4 is hydrogen; and
 - R_9 is a hydroxyl, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, CO_2R_{10} , CHO or $NR_{12}SO_2NR_{10}R_{11}$ group.
- 10
- Particularly preferred compounds of formula (I) are those in which Z is:




in which R_8 and R_9 are as hereinabove defined.

- 15
- In the above defined compounds, R_8 and R_9 are preferably as follows:
- R_8 is hydrogen, hydroxy, halogen, nitro, cyano, (C_1-C_6) alkoxy, $NR_{10}R_{11}$, $SO_2NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, $NR_{12}SO_2NR_{10}R_{11}$ or $OSO_2NR_{10}SO_2NR_{11}R_{12}$ group;
 - R_9 hydrogen or a hydroxy, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, CO_2R_{10} , CHO, $NR_{12}SO_2NR_{10}R_{11}$ group.
- 20
- p and q are as defined above.

Among the compounds of formula (I), those in which Q is selected from a direct link, C(O), SO_2 , CONH, $C(O)(CH_2)_n$, $(CH_2)_n(O)$ or $(CH_2)_n$ in which n is 0,1 or 2, are also particularly preferred.

Particularly preferred are the compounds of formula (II)



in which:

- Q is $(CH_2)_n$ in which n is 0, 1 or 2;
- 5 • one of R_3 and R_8 is a hydroxy, nitro, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$ or $NR_{12}SO_2NR_{10}R_{11}$ group and the other is hydrogen or a hydroxy, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, (C_1-C_6) alkylthio, (C_1-C_6) alkylsulfonyl, acyl, (C_1-C_6) alkoxycarbonyl, carboxamido, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$ or $NR_{12}SO_2NR_{10}R_{11}$ group;
- 10 • R_4 and R_9 are each independently hydrogen, hydroxy, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, (C_1-C_6) alkylthio, (C_1-C_6) alkylsulfonyl, acyl, (C_1-C_6) alkoxycarbonyl, carboxamido, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$ or $NR_{12}SO_2NR_{10}R_{11}$ group .
- R_{10} and R_{11} are each independently hydrogen, a (C_1-C_6) alkyl or a
- 15 (C_3-C_8) cycloalkyl;
- p is 1, 2, 3 or 4;
- R_8 and R_9 together with the phenyl ring bearing them can also form a benzoxathiazine dioxide or a dihydrobenzoxathiazine dioxide;
- R_3 and R_4 together with the phenyl ring bearing them can also form a
- 20 benzofurane or a N-methylbenzotriazole.

Among these compounds of formula (II), those which fulfil at least one of the following conditions are preferred:

- Q is $(CH_2)_n$ in which n 0, 1 or 2;
- 25 • R_8 is hydroxy, halogen, nitro, cyano or a (C_1-C_6) alkoxy, $NR_{10}R_{11}$, $SO_2NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$ or $NR_{12}SO_2NR_{10}R_{11}$ group;

- R_9 is hydrogen, hydroxy, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$;
- p is 1, 2, 3 or 4.

5 Most preferred compounds of formula (II) are those in which:

- n is 0 or 1;
- R_1 and R_2 are each independently hydrogen or (C_1-C_6) alkyl;
- R_4 and R_9 are each independently hydrogen, halogen, (C_1-C_6) alkoxy, acyl, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$ or $NR_{12}SO_2NR_{10}R_{11}$.

10

Especially preferred compounds of formula (II) are those in which:

- n is 0 or 1;
- R_1 , R_2 and R_4 are each hydrogen;
- R_9 is hydrogen, halogen, (C_1-C_6) alkoxy or $OSO_2NR_{10}R_{11}$.

15

Particularly preferred compounds of formula (II) are those in which:


- n and p are 1;
- R_8 is a hydroxy, halogen, nitro, cyano, (C_1-C_6) alkoxy, $NR_{10}R_{11}$, $SO_2NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, $NR_{12}SO_2NR_{10}R_{11}$, or $OSO_2NR_{10}SO_2NR_{11}R_{12}$ group;
- 20 • R_9 is a hydroxy, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, CO_2R_{10} or CHO group;
- R_3 is cyano, hydroxy, $OSO_2NR_{10}R_{11}$ or $NR_{12}SO_2NR_{10}R_{11}$;
- R_4 is hydrogen, hydroxy, halogen, cyano or $OSO_2NR_{10}R_{11}$.

25 Among these compounds, those which fulfil at least one of the following conditions are also preferred:

- one of R_3 and R_8 is hydroxy, cyano or $OSO_2NR_{10}R_{11}$, preferably cyano or $OSO_2NR_{10}R_{11}$; and
- the other is hydroxy, nitro, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$ or $NR_{12}SO_2NR_{10}R_{11}$,
- 30 preferably hydroxy or $OSO_2NR_{10}R_{11}$.

Among these compounds of formula (II), those wherein R_{10} and R_{11} are hydrogen are mostly preferred.

Particularly preferred are also the compounds of formula (III):



in which:

- Q is $(CH_2)_m-X-(CH_2)_n-A-$;
- A is a direct bond or O, S, SO, SO₂, NR₅;
- X is a direct bond, CF₂, O, S, SO, SO₂, C(O), NR₅ or CR₆R₇;
- m and n are each independently 0, 1, 2, 3 or 4;
- R₃, R₄, R₈ and R₉ are each independently hydrogen or a hydroxy, cyano, halogen, nitro, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, benzyloxy, trifluoromethyl, (C₁-C₆)alkylthio, (C₁-C₆)alkylsulfonyl, acyl, (C₁-C₆)alkoxycarbonyl, NR₁₀R₁₁, OPO(OR₁₀)₂, OCHO, COOR₁₀, SO₂NR₁₀R₁₁, OSO₂NR₁₀R₁₁, SO₂OR₁₀, OSO₂OR₁₀, SSO₂NR₁₀R₁₁, CONR₁₀R₁₁, OCONR₁₀R₁₁, OCSNR₁₀R₁₁, SCONR₁₀R₁₁, SCSNR₁₀R₁₁, NR₁₂SO₂NR₁₀R₁₁, tetrazolyl, NR₁₀CONR₁₁OH, NR₁₀SO₂NR₁₁OH, NOH-CHO, NOHSO₂NR₁₀R₁₁, or OSO₂NR₁₀OH group;
- p is 1, 2, 3 or 4;
- q is 0, 1 or 2.
- R₅, R₆, R₇, R₁₀, R₁₁ and R₁₂ are each independently hydrogen, a (C₁-C₆)alkyl or a (C₃-C₈)cycloalkyl; R₁₀ can also be a salt; R₁₀ and R₁₁ can also form, together with the nitrogen atom to which they are bound, a 5- to 7-membered heterocycle containing one or two heteroatoms selected from O, S and N;
- The dotted line means that Q and/or R₈ and/or R₉ can be on any position of the benzothiophene ring.


Among the compounds of formula (III) those which fulfil at least one of the following conditions are also preferred:

- R₃ is hydrogen, halogen or cyano;
- R₈ is OSO₂NR₁₀R₁₁ or NR₁₂SO₂NR₁₀R₁₁;
- R₉ is hydrogen, halogen, nitro, COOR₁₀ or cyano;
- R₄ is hydrogen, halogen, cyano, (C₁-C₆)alkoxy, NR₁₀R₁₁, OSO₂NR₁₀R₁₁ or
- 5 NR₁₂SO₂NR₁₀R₁₁;
- R₁₀, R₁₁ and R₁₂ are each independently hydrogen or (C₁-C₆)alkyl.

The compounds of formula (III), wherein Q is (CH₂)_m-X-(CH₂)_n-A where m is 0, 1 or 2 and X is a direct bond, SO₂ or CO, n is 0 and A is a direct bond are also

10 preferred.

Particularly preferred are the compounds of formula (IV):



in which R₁, R₂, R₃, R₄, R₈, R₉ and p are as defined for compounds of formula (I).

15

Particularly preferred compounds of formula (IV) are those in which:

- R₃ is cyano or OSO₂NR₁₀R₁₁;
- R₄ is hydrogen, hydroxyl, halogen, cyano, OSO₂NR₁₀R₁₁;
- R₈ is hydroxy, cyano, OSO₂NR₁₀R₁₁, NR₁₀R₁₁, NR₁₂SO₂NR₁₀R₁₁, OCHO,
- 20 tetrazolyl;
- R₉ is hydrogen, halogen, nitro, cyano or CO₂R₁₀; and
- Q is as defined above for compounds of formula (I).

By virtue of their capability to inhibit the enzymes aromatase and/or steroid

25 sulfatase and/or carbonic anhydrase, the compounds of the present invention can be used alone or in combination with other active ingredients for the

treatment or the prevention of any hormone or non hormone-dependent cancer, in humans as well as in wild or domestic animals. Because of their inhibition activity of aromatase and/or steroid sulfatase, the compounds of the invention are suitable for the management of estrogen-regulated reproductive functions, in humans as well as in wild or domestic animals.

In the treatment or prevention of the above conditions, the compounds of the invention can be used alone or in combination with an antiestrogen, a SERM (selective estrogen receptor modulator), an aromatase inhibitor, a carbonic anhydrase inhibitor, an antiandrogen, a steroid sulfatase inhibitor, a lyase inhibitor, a progestin, or a LH-RH agonist or antagonist. The compounds of the invention can also be used in combination with a radiotherapeutic agent; a chemiotherapeutic agent such as a nitrogenated mustard analogue like cyclophosphamide, melphalan, iphosphamide, or trophosphamide; an ethylenimine like thiotepa; a nitrosourea like carmustine; a lysed agent like temozolomide or dacarbazine; an antimetabolite of folic acid like methotrexate or raltitrexed; a purine analogue like thioguanine, cladribine or fludarabine; a pyrimidine analogue like fluorouracil, tegafur or gemcitabine; an alkaloid of vinca or analogue like vinblastine, vincristine or vinorelbine; a podophyllotoxin derivative like etoposide, taxanes, docetaxel or paclitaxel; an anthracycline or analogue like doxorubicin, epirubicin, idarubicin or mitoxantrone; a cytotoxic antibiotic like bleomycin or mitomycin; a platinum compound like cisplatin, carboplatin or oxaliplatin; a monoclonal antibody like rituximab; an antineoplastic agent like pentostatin, miltefosine, estramustine, topotecan, irinotecan or bicalutamide; or with a prostaglandin inhibitor (COX 2/COX 1 inhibitor).

The compounds of the invention can also be used for the control or management of estrogen-regulated reproductive functions such as male or female fertility, pregnancy, abortion or delivery, in humans as well as in wild or domestic animal species, alone or in combination with one or several other therapeutic agents such as a LH-RH agonist or antagonist, an estroprogestative contraceptive, a progestin, an antiprogesterone or a prostaglandin inhibitor.

Breast tissue being a sensitive target of estrogen-stimulated proliferation and/or differentiation, inhibitors of aromatase and/or steroid sulfatase and/or carbonic anhydrase can be used in the treatment or prevention of benign breast diseases

in women, gynecomastia in men and in benign or malignant breast tumors with or without metastasis both in men and women or in male or female domestic animals. The compounds of the invention can also be used in the treatment or prevention of benign or malignant disease of the uterus or the ovary. In each
5 case, the compounds of the invention can be used alone or in combination with one or several other sexual endocrine therapeutic agents such as an antiandrogen, an anti-estrogen, a progestin or a LH-RH agonist or antagonist.

As the enzyme steroid sulfatase transforms DHEA sulfate into DHEA, a precursor of active androgens (testosterone and dihydrotestosterone), the compounds of
10 the invention can be used in the treatment or prevention of androgen-dependent diseases such as androgenic alopecia (male pattern loss) (Hoffman R et al., J. Invest. Dermatol., 2001, 117, 1342-1348), hirsutism, acne (Billich A et al., WO 9952890), benign or malignant diseases of the prostate or the testis (Reed MJ, Rev. Endocr. Relat. Cancer, 1993, 45, 51-62), alone or in combination with one
15 or several other sexual endocrine therapeutic agents such as an antiandrogen, an antiestrogen, a SERM, an antiaromatase, a progestin, a lyase inhibitor or a LH-RH agonist or antagonist.

Inhibitors of steroid sulfatase are also potentially involved in the treatment of cognitive dysfunction, because they are able to enhance learning and spatial
20 memory in the rat (Johnson DA, Brain Res, 2000, 865, 286-290). DHEA sulfate as a neurosteroid affects a number of neurotransmitter systems including those involving acetylcholine, glutamate, and GABA, resulting in increased neuronal excitability (Wolf OT, Brain Res. Rev, 1999, 30, 264-288). The compounds of the present invention are thus also useful for enhancing the cognitive function,
25 especially for the treatment of senile dementia, including Alzheimer's diseases, by increasing the DHEA levels in the central nervous system.

In addition, estrogens are involved in the regulation of the balance between Th₁ and Th₂ predominant immune functions and may therefore be useful in the treatment or prevention of gender-dependent auto-immune diseases such as
30 lupus, multiple sclerosis, rheumatoid arthritis and the like (Daynes RA, J. Exp. Med, 1990, 171, 979-996). Steroid sulfatase inhibition was further shown to be protective in models of contact allergy and collagen-induced arthritis in rodents (Sutters AJ, Immunology, 1997, 91, 314-321).

Studies using 2-MeOEMATE have shown that steroid sulfatase inhibitors have potent estradiol-independent growth-inhibitory effect (MacCarthy-Moorogh L, Cancer Research, 2000, 60, 5441-5450). A decrease in tumor volume was surprisingly observed with the compounds of the invention, with low tumor
5 steroid sulfatase inhibition. In view of this, the compounds of the invention could lead to a decrease in cellular division because of the large interaction between such new chemical entities and the microtubular network within the cancerous cell, whatever the tissue, including breast, endometrium, uterus, prostate, testis or metastasis generated therefrom. The compounds of the invention could
10 therefore be useful in the treatment of non-estrogen dependent cancer. The compounds of the invention are of particular value for the treatment or prevention of estrogen-dependent diseases or disorders, i.e. estrogen-induced or estrogen-stimulated diseases or disorders (Golob T, Bioorg. Med. Chem., 2002, 10, 3941-3953).

15 In addition, the compounds of the present invention are inhibitors of carbonic anhydrase (CA). This property could explain the interest of such compounds in non-hormone-dependent cancer. Immunohistochemical studies of CA II have shown that it is expressed in malignant brain tumors (Parkkila A-K. et al., Histochem. J., 1995, 27: 974-982) and gastric and pancreatic carcinomas
20 (Parkkila S et al., Histochem. J., 1995, 27: 133-138), and recent evidence has shown that CA IX and XII are also expressed in some tumors and may be functionally related to oncogenesis. Ivanov et al. (Proc. Natl. Acad. Sci. USA, 1998, 95: 12596-12601) recently hypothesized that tumor-associated CA IX and XII may be implicated in acidification of the extracellular medium surrounding
25 cancer cells, which would create a microenvironment conducive to tumor growth and spreading. It has been shown that acetazolamide markedly inhibited invasion capacity in four renal cancer cell lines (Parkkila S et al., Proc. Natl. Acad. Sci. USA, 2000, 97: 2220-2224), an effect attributable to CA II, IX, and XII, which were expressed in these cells. Leukemia cells can easily spread from bone
30 marrow to other organs via circulation, but various leukemias differ in their ability to form extramedullary tumors i.e., metastases. If CA activities were essential for invasion by other cancer cells, one could analogously predict that active CA(s) could also function in leukemia cells.

As used herein, the term "combined" or "combination" refers to any protocol for the co-administration of a compound of the invention and one or more other pharmaceutical substances, irrespective of the nature of the time of administration and the variation of dose over time of any of the substances. The co-administration can for example be parallel, sequential or over a period of time.

For the treatment/prevention of any of the above-mentioned diseases or disorders, the compounds of the invention may be administered, for example, orally, topically, parenterally, in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. These dosage forms are given as examples, but other dosage forms may be developed by those skilled in the art of formulation, for the administration of the compounds of the invention. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition to the treatment of humans, the compounds of the invention are effective in the treatment of warm-blooded animals such as mice, rats, horses, cattle sheep, dogs, cats, etc.

The pharmaceutical compositions containing the active ingredient(s) may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavouring agents, colouring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient(s) in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by

known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

- 5 They may also be coated by the technique described in the U.S. Patent 4,256,108; 4,166,452 and 4,265,874 to form osmotic therapeutic tablets for controlled release.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient(s) is (are) mixed with an inert solid diluent, for example,
10 calcium carbonate, calcium phosphate, or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active ingredient(s) in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients
15 are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example
20 polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol
25 anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more colouring agents, one or more flavouring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

- 30 Oily suspensions may be formulated by suspending the active ingredient(s) in a vegetable oil, for example peanut oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or acetyl alcohol. Sweetening agents

such as those set forth above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

5 Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient(s) in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavouring and colouring agents, may also be present. The
10 pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsion. The oily phase may be a vegetable oil, for example olive oil or peanut oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally occurring phosphatides, for example soybean, lecithin, and esters or partial esters derived from fatty acids
15 and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavouring agents.

The pharmaceutical compositions may be in the form of a sterile injectable
20 aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-
25 butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find
30 use in the preparation of injectables.

Dosage levels of the order of from about 0.0001 mg to about 20 mg/kg of body weight per day are useful in the treatment of the above-indicated conditions, or alternatively about 0.1 mg to about 2000 mg per patient per day.


The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 0.1 mg to about 400 mg of compound of the invention, typically 0.1 mg, 1 mg, 2 mg, 5 mg, 10 mg, 20 mg, 40 mg, 80 mg, 100 mg, 200 mg or 400 mg.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

According to another object, the invention relates to a method for the treatment or prevention of the above-mentioned diseases, disorders or conditions. The method comprises administering to a subject (human or animal) in need thereof a therapeutically effective amount of a compound of the invention or a pharmaceutically acceptable acid addition salt thereof.

The 1-N-aryl-amino-1H-imidazole derivatives of formula (I) and their acid addition salts can be prepared following general schemes I a, I b, IIa, IIb, III, IV and V in which $(\text{O})_m$ represents $(\text{CH}_2)_m$ and $(\text{O})_n$ represents $(\text{CH}_2)_n$.

Scheme Ia



According to scheme I a, N,N-disubstituted aniline (3) can be obtained by
 5 condensation of the aniline derivative (1) with halogeno derivatives, alkoxy
 derivatives, sulfonyl derivatives or sulfinyl derivatives (2) using standard
 conditions (March J., *Advanced Organic Chemistry*, Fourth edition, Wiley
 Interscience, New-York). Most of the halogeno derivatives, alkoxy derivatives,
 sulfonyl derivatives or sulfinyl derivatives (2) are commercially available or are
 10 synthesized by usual chemistry methods (see experimental part).

Compound (3) is converted to its nitroso derivative using standard conditions, then reduced to afford the 1,1-disubstituted hydrazine of formula (4).

Alternatively, the 1,1-disubstituted hydrazine (4) can be prepared by selective N-alkylation of a hydrazine of formula (5) with a compound of formula (2) using the

5 conditions described by U. Lerch and J. König (*Synthesis*, 1983, 2, 157-8) or the conditions described by J. Chung et al. (*Tetrahedron Letters*, 1992, 33, 4717-20).

Then, condensation of (4) with a dialkyloxy-alkyl-isothiocyanate derivative or an ethylenedioxy-alkyl-isocyanate derivative, affords the thiosemicarbazide (6) which is transformed to the 1-amino-imidazole-2-thione (7) by treatment with an

10 acid like acetic acid or sulphuric acid.

Desulfurization of (7) in acetic acid, following the conditions described by S.Grivas and E.Ronne in *Acta Chemica Scandinavia*, 1995, 49, 225-229, gives the final 1-N-phenyl-amino-1H-imidazole (8), which is optionally converted to one of its acid addition salts.

15

Alternatively said compounds (8) where R₃ or R₄ is an electron-withdrawing group can be obtained by condensation of the N-imidazoloaniline (9) with halogeno derivatives, alcoyl derivatives, sulfonyl derivatives or sulfinyl derivatives (2) using standard conditions (March J., *Advanced Organic Chemistry*, Fourth

20 edition, Wiley Interscience, New-York).


If R₈ is ester, saponification of compound (8) gives carboxylic derivative by usual chemistry methods.

If R₈ is sulfamide, it can be directly obtained with correct substituted alcoyl derivative or halogeno derivative (2).

25 if R₈ is cyano group, reaction with sodium azide gives tetrazolyl group (Kiyoto K., *Synthesis*, 1998, 910-14).

The compound (9) is obtained by a process similar to the process disclosed herein above for obtaining the compound (8) starting from the compound (4).

Scheme Ib



- 5 According to scheme Ib deprotection of methoxy or benzyloxy derivatives (8) with tribromoborane (McOmie. J.F.W, *Tetrahedron*, 1968, 24, 2289-92) or piperidine (Nishioka H. *Synthesis*, 200, 2, 243-46) or by hydrogenation (Felix A., *J Org Chem*, 1978, 43, 4194-97) gives the hydroxyl compound (10). Reduction of nitro compound (8) by stannous chloride or with ruthenium and hydrazine (WO 10 02051821) gives amino-compound (12) (Matassa V., *J Med Chem*, 1990, 33, 2621-29).

These compounds are transformed into the corresponding sulfamates (11) or aminosulfonylamines (13) by treatment with sodium hydride and sulfamoyl chloride (Nussbaumer. P, *J Med Chem*, 2002, 45, 4310-20) or by reaction with 15 sulfamoyl chloride in dimethylacetamide (DMAc) (Makoto O, *Tetrahedron letters*, 2000, 41, 7047-51).


These hydroxyl compounds can be transformed into formate derivatives by treatment with formyl acetate (Schreiner E., *Bioorg Med Chem Lett*, 2004, 14,

4999-5002) or into 1H-imidazole-1-carboxylate derivatives by treatment with N,N'-carbonyldiimidazole (Fischer, W., *Synthesis*, 2002, 1, 29-30)

Compounds (10) or (12) having both an amino and an hydroxyl in an ortho position, can be transformed to the corresponding benzoxathiazoles following the conditions proposed by K. K. Andersen (*J Org Chem*, 1991, 56, 23, 6508-6516).

When X is CO or CS, reduction of ketone, thioketone, amide, thioamide is performed following standard conditions (March J., *Advanced Organic Chemistry*, Fourth edition, Wiley Interscience, New-York). These reduction steps can be effective at any steps of synthesis process.

10

Scheme IIa

According to scheme IIa, compounds (15) can be obtained by the same procedure as compound (3) in scheme Ia starting from compounds (1) and (14) or by condensation of the N-imidazoloaniline (9) with halogeno derivatives, alcoyl derivatives, sulfonyl derivatives or sulfinyl derivatives (14) using standard conditions (March J., *Advanced Organic Chemistry*, Fourth edition, Wiley Interscience, New-York). Most of the halogeno derivatives, alcoyl derivatives, sulfonyl derivatives or sulfinyl derivatives (14) are commercially available or are synthesized by usual chemistry methods (see experimental part).

5 If R_8 is ester, saponification of compound (15) gives carboxylic derivative by usual chemistry methods.

If R_8 is sulfamide, it can be directly obtained with correct substituted alcoyl derivative or halogeno derivative (14).

if R_8 is cyano group, reaction with sodium azide gives tetrazolyl group (Kiyoto K., *Synthesis*, 1998, 910-14).

15 Deprotection of methoxy or benzyloxy benzothiophene (15) with tribromoborane (McOmie. J.F.W, *Tetrahedron*, 1968, 24, 2289-92) or by hydrogenation (Felix A., *J Org Chem*, 1978, 43, 4194-97) gives the hydroxy benzothiophene (16). Reduction of nitro benzothiophene compound (15) by stannous chloride gives amino-benzothiophene (18) (Matassa V., *J Med Chem*, 1990, 33, 2621-29).

20

These compounds (16) and (18) are transformed into the corresponding sulfamates (17) or aminosulfonylamines (19) using the same conditions as for the synthesis of compounds (11) or (13).


Oxidation of sulphur on benzothiophene by hydrogen peroxide in trifluoroacetic acid, following the conditions described by Grivas S. and Ronne E. (*Acta Chemica Scandinavia*, 1995, 49, 225-229) or by meta-chloro-per-benzoic acid in methylene chloride gives the oxydised benzothiophenes.(Ong H. H., *J Med Chem*, 1987, 30, 12, 2295-2303).

25 Reduction of amide, thioamide, ketone, thioketone when X is CO or CS or reduction of oxydised sulphur function is performed following the conditions described by Ellefson C. (*J Med Chem*, 1981, 24, 1107-10), Hajos J. (*Complex Hydrides*, Elsevier, New York, 1979) or Drabowicz S. (*Org Prep Proced Int*, 1977, 9, 63-83) and Bordwell J. (*J Am Chem Soc*, 1951, 73, 2251-53) or following

30

standard conditions (March J., *Advanced Organic Chemistry*, Fourth edition, Wiley Interscience, New-York). These oxidation and reduction steps can be effective at any steps of synthesis process.

5

Scheme IIb

According to scheme IIb, compounds (21) are synthesized following the same synthetic methods as for compounds (3) starting from compounds (1) and (20) or (9) and (20).

10

3-Halogeno benzothienopyridine derivative (21) treated by aqueous metal hydroxide (Svoboda J., *Collect Czech Chem comm*, 2000, 65, 7, 1082-92 or Sall D., *J Med Chem*, 2000, 43, 4, 649-63) gives 3-hydroxy benzothienopyridine derivative (22) or

treated by aqueous ammonia in acetone or ethanol (Bordwell F., *J. A. C. S.*, 1948, 70, 1955-58) gives 3-amino benzothiophene derivative (24).

2-Hydroxy benzothiophene derivative (22) or 2-amino benzothiophene derivative (24) are performed respectively by deprotection of 2-methoxy benzothiophene derivative (21) with pyridine hydrochloride (Cannizzo S., *J Heterocyclic Chem*, 5 1990, 27, 2175-79) and by reduction of 2-nitro benzothiophene compound (21) by stannous chloride (Matassa V, *J Med Chem*, 1990, 33, 2621-29).

These compounds (22) and (24) can be sulfamoylated to give (23) and (25) using the same conditions as for the synthesis of compounds (11) or (13).

10


Deprotonation of 2-H-benzothiophene derivative (21) with lithium amides or alkyls leads to lithiation at C-2 position. Addition of sulfonyl chloride yields chlorosulfonyl compound that is treated by aqueous ammonia in acetone (Graham S., *J Med Chem*, 1989, 32, 2548-54) to give 2-sulfonamide 15 benzothiophene derivative (21) or addition of dry ice followed by hydrolysis gives 2-carboxylic acid benzothiophene derivative (21) (Matecka D., *J Med Chem*, 1997, 40, 705-16).

3-Carboxylic or carboxamide benzothiophene derivative (21) is prepared by 20 treatment of 3-H-benzothiophene derivative with trichloroacetyl chloride/aluminium chloride followed by hydrolysis with water (Bonjouklian R., *Synth Comm*, 1985, 15, 8, 711-13) or aqueous ammonia (Turnbull K., *J Heterocycl Chem*, 2000, 37, 2, 383-88).

3-Sulfonamide derivative (21) is prepared following the conditions described by 25 Chapman N. (*J. Chem. Soc.*, 1970, 18, 2431-35) or Hageman W. (Ger. Offen., 3435173, 11 April 1985).

Oxidation of sulfur on benzothiophene and reduction of carboxamide, thioamide, ketone, thioketone, oxydised sulphur function can be performed at any step of 30 synthesis following the same conditions already presented in this description.

Scheme III




- 5 According to scheme III, compounds (27) are obtained by condensation of the N-imidazoloaniline (9) with isocyanate derivatives (26) using standard conditions (March J., *Advanced Organic Chemistry*, Fourth edition, Wiley Interscience, New-York). Most of the isocyanate derivatives (26) are commercially available or are synthesized by usual chemistry methods.
- 10 Compounds (28) and (30) are synthesized following the same synthetic methods as for compounds (10) and (12) in scheme Ib.

Compounds (29) and (31) are synthesized following the same synthetic methods as for compounds (11) and (13) in scheme Ib.

Scheme IV


5



According to scheme IV, bi-aryl compounds (33) can be synthesized following the same synthetic methods as for compounds (8) with derivatives (32). Derivatives (32) are commercially available or synthesized by usual chemistry methods (as ex Buraway S, J Chem Soc, 1955, 2557; Tilley J W, J Med Chem, 1989, 32, 8, 1814).

15

Scheme V



According to scheme V, compounds (35) are synthesized following the same synthetic methods as for compounds (8) with heterocycles (34) (one or two nitrogen atoms in any positions). Heterocycles (34) are commercially available or synthesized by usual chemistry methods (see following examples). Halogeno pyridine derivatives (34) can be obtained following publications (Biorg Med Chem

- Lett, 1996, 6, 21, 2613; Myers A G, J Org Chem, 1996, 61, 813; Tetrahedron, 1993, 49, 19, 4085) or from carboxylic acid (WO 0177078). Halogeno pyrimidine derivatives (34) can be obtained from alkyl pyrimidines (Budesinsky, Collect, Czech, Chem Commun, 1968, 33, 7, 2266; Kunieda T, J Am Chem Soc, 1971, 93,3487) by usual halogenation (Isoda S, Chem pharm Bull, 1980, 28, 5, 1408; March J., *Advanced Organic Chemistry*, Fourth edition, Wiley Interscience, New-York), or from carbaldehydes (Bredereck, Chem Ber, 1967, 100, 11, 3664; Adams J L, Bioorg Med Chem Lett, 1998, 8, 22, 3111), and carboxylic acids (Huffman K R, J Org Chem, 27, 1962, 551; Daves J Org Chem, 1961, 26, 2755) can be transformed to acid chlorides (34). Halogeno pyrazine derivatives (34) can be obtained from alkyl pyrazines (Lutz W B, J Org Chem, 1964, 29, 415) by bromination, or from carbaldehydes (US 3558625), and carboxylic acids (Sato N, J Heterocycl Chem, 19, 1982, 407-408; Felder P, Helv Chim Acta, 1964, 47, 873) can be transformed to acid chlorides (34). Halogeno pyridazine derivatives (34) can be obtained following Piras S (Farmaco, 1993, 48, 9, 1249) Yanai, (Heterocycles, 1976, 4, 1331), or by halogenation of alkyl pyridazines (Becker, J Prakt Chem, 1970, 312, 591; DE 1950491) and carboxylic acids (Boger D L, J Am Chem Soc 1987, 109, 9, 2717) can be transformed to acid chlorides (34).
- For all schemes IIa, IIb, III, IV, V, carboxylate derivative, sulfamide derivative and tetrazole derivative are synthesized by the methods already described for scheme Ia and R8 and (R9)p are performed following the same conditions already presented in this description.
- The groups described for R₃, R₄, R₈ and R₉ can be obtained by usual chemistry methods (for references see review on sulfatase (Nussbaumer P, *Medecinal Research*, 2004, 24, 4, 529-76), on carbonic anhydrase (Supuran C T, *Carbonic anhydrase*, 2004, C R C press) and articles from Park J D (*J Heterocycl Chem*, 2000, 37, 2, 383-88), Schreiner E P (*Bioorg Med Chem Lett*, 2004, 14, 4999-5002) and Taylor S D (*Bioorg Med Chem Lett*, 2004, 14, 151-155).

The following examples are intended to illustrate and not to limit the scope of the invention.

PREPARATION OF N, N-DISUBSTITUTED HYDRAZINES (4)**EXAMPLE 1****N¹-(4-cyanophenylmethyl)-N¹-(4-methoxyphenyl)hydrazine**

- Chloromethylbenzonitrile (25 g, 164.90 mmol) was introduced with stirring into a flask containing toluene (200 ml) and triethylamine (46.40 ml, 329.80 mmol). 4-methoxy-phenylhydrazine hydrochloride (28.80 g, 164.90 mmol) was added portionwise and the reaction mixture was stirred 3 h at reflux. After cooling, the mixture was filtered, washed with toluene (50 ml) and with water (200 ml) to give a white solid (27.20 g, 65%), mp: 115°C.
- ¹H-NMR (DMSO d₆) : 3.65 (s, 3H), 4.30 (s, 2H), 4.57 (s, 2H), 6.77 (d, 2H), 6.94 (d, 2H), 7.48 (d, 2H), 7.76 (d, 2H).

PREPARATION OF IMIDAZOLES (9)**EXAMPLE 2**

- 4-[N-(1H-imidazol-1-yl)amino]benzonitrile**
- a) 4-[N-(2,3-dihydro-1H-imidazol-1-yl-2-thione)amino]benzonitrile
- To a suspension of 4-cyanophenylhydrazine hydrochloride (6.00 g, 35.40 mmol) in ethanol (60 ml) was added dropwise 2,2-dimethoxyethylisothiocyanate (6.25 g, 42.4 mmol) and the reaction mixture was heated to reflux for 2 h. After cooling the solvent was evaporated under vacuum, the resulting oil was diluted with acetic acid/ water (9/1, 32 ml) and the suspension was heated to reflux for 1.5 h and at room temperature overnight. The resulting residue was poured into water (300 ml) and a brown precipitate was collected. After trituration from ethanol, the brown solid afforded a white solid (4.60 g, 58 %).
- ¹H-NMR (DMSO d₆) : 6.54 (d, 2H), 7.00 (t, 1H), 7.23 (t, 1H), 7.62 (d, 2H), 9.83 (s, 1H), 12.40 (s, 1H).
- b) 4-[N-(1H-imidazol-1-yl)amino]benzonitrile
- 35 % hydrogen peroxide (4.90 ml, 55.5 mmol) was added dropwise to an ice-cooled suspension of 4-[N-(2,3-dihydro-1H-imidazol-1-yl-2-thione)amino]benzonitrile (4.00 g, 18.50 mmol) in acetic acid (20 ml). When TLC showed complete reaction, the reaction mixture was diluted with water, adjusted to pH 11 with sodium hydroxide, treated with sodium hydrogen sulfite and extracted with ethyl acetate. The organic layer was dried over sodium sulfate and

concentrated under vacuum. Flash chromatography on silica gel (toluene / dioxane : 6/4) yielded a pure oil and crystallization from ethanol afforded white crystals (4.40 g, 58 %), mp: 162°C.

¹H-NMR (DMSO-d₆) : 6.50 (d, 2H), 7.08 (s, 1H), 7.30 (s, 1H), 7.66 (d, 2H), 7.83 (s, 1H).

PREPARATION OF BENZOTHIOPHENE DERIVATIVES (14) AND (20)

EXAMPLE 3

1-Chloro-3-(3-methoxyphenyl)sulfanyl-propan-2-one

To a stirred solution of 1,3-dichloro-2-propanone (12.70 g, 0.1 mol) in methanol/water (100 ml, 1:3) at 0°C is added a suspension of 3-methoxybenzenethiol (14.02 g, 0.1 mol) and sodium hydroxide (4.00 g, 0.10 g) in water (100 ml). The mixture is stirred at 0°C for 7 h and at room temperature for 10 h. The precipitated product is extracted with dichloromethane (100 ml), washed with water (80 ml), and dried with sodium sulfate. After removal of the solvent, we obtained the good product (oil, 18.70 g).

¹H-NMR (CDCl₃): 3.80 (s, 3H), 3.83 (s, 2H), 4.29 (s, 2H), 6.78 (dd, 1H), 6.98 (d, 1H), 6.90 (dd, 1H), 7.21 (t, 1H).

EXAMPLE 4

3-Chloromethyl-6-methoxy-benzothiophene

A solution of the above thio compound (17.50 g, 75.85 mmol) in CH₂Cl₂ (1700 ml) was added dropwise to a solution of BF₃.Et₂O (10.60 ml, 83.44 mmol) in CH₂Cl₂ (100 ml) at room temperature under nitrogen atmosphere. The mixture was stirred overnight and after hydrolysis with aqueous NaHCO₃ solution, the reaction mixture was stirred until both phases became clear. The CH₂Cl₂ layer was separated, and the aqueous layer was extracted with CH₂Cl₂. The combined organic phases were dried over Na₂SO₄, filtered, and concentrated under vacuum to give an oil (18.00 g). Flash chromatography on silica gel (toluene / petroleum spirit 40-60°C : 5/5) yielded a 1:10 mixture of 3-chloromethyl-4-methoxy-benzothiophene: 3-chloromethyl-6-methoxy-benzothiophene as an oil (12.35 g, 58 %).

Major isomer ¹H-NMR (CDCl₃): 3.89 (s, 3H), 4.82 (s, 2H), 7.08 (dd, 1H), 7.30 (s, 1H), 7.35 (d, 1H), 7.78 (d, 1H).

EXAMPLE 5**3-Bromo-6-benzyloxy-benzothiophene**

N-bromosuccinimide (15.70 g, 83.92 mmol) and p-toluenesulfonic acid (2.70 g, 15.68 mmol) were added to a solution of 6-benzyloxy-benzothiophene (Zhengying C., CN 1370533 A, 21.2 g, 88.33 mmol) in 1,2-dichloroethane (120 ml). The mixture was maintained at 80°C for 35 min, cooled in an ice bath, and the succinimide was removed by filtration. The solution was extracted with saturated sodium bicarbonate solution, dried over Na₂SO₄, filtered, and concentrated under vacuum to give an oil. Crystallisation from pentane afforded a white solid (21.60 g, 92 %, mp: 68°C).

¹H-NMR (DMSO-d₆): 5.14 (s, 2H), 7.08 (dd, 1H), 7.25-7.55 (m, 6H), 7.65 (d, 1H), 7.76 (d, 1H).

EXAMPLE 6**3-Bromo-6-benzyloxy-benzothiophene-1,1-dioxyde**

To a solution of 3-bromo-6-benzyloxy-benzothiophene (2.00 g, 6.27 mmol) in dichloromethane (50 ml) and trifluoroacetic acid (1.5 ml) was added 35% aqueous hydrogen peroxide (2.00 ml, 19.54 mmol). After 8 h at 50°C the mixture was hydrolysed with saturated aqueous NaHCO₃, extracted with dichloromethane, dried over Na₂SO₄, filtered, and concentrated under vacuum to give the crude product. Flash chromatography on silica gel (toluene / ethyl acetate: 9/1) yielded a limpid oil (1.10 g, 55%).

¹H-NMR (DMSO-d₆): 5.20 (s, 2H), 7.20-7.60 (m, 7H), 7.72 (d, 1H), 7.83 (s, 1H).

EXAMPLE 7**(6-Benzyloxy-benzothien-2-yl)methanol**

To a solution of 6-benzyloxy-benzothiophene-2-carbaldehyde (described by Nomura Y., WO 9635688 A1, 6.50 g, 24.20 mmol) in THF (50 ml) was added dropwise to a -30°C cooled suspension of LiAlH₄ (0.85 g, 22.26 mmol). After warming up to room temperature the mixture was stirred overnight, cooled to -10°C, hydrolyzed with ice water, extracted with dichloromethane, dried over Na₂SO₄, filtered and concentrated under vacuum to give the crude product. Flash chromatography on silica gel (toluene / ethyl acetate: 7/3) yielded a limpid oil (4.50 g, 69%).

¹H-NMR (DMSO-d₆): 4.68 (s, 2H), 5.13 (s, 2H), 5.60 (s, 1H), 7.00 (dd, 1H), 7.14 (s, 1H), 7.25-7.80 (m, 7H).

EXAMPLE 8

6-Benzyloxy-2-(chloromethyl)-benzothiophene

- 5 Sulfonyl chloride (20 ml) is added to a solution of (6-benzyloxy-benzothien-2-yl) methanol (4.20 g, 15.50 mmol) in dichloromethane (40 ml). The mixture was maintained at reflux for 2 h, cooled at room temperature then concentrated under vacuum to give 4.20 g as an oil.

¹H-NMR (CDCl₃): 4.75 (s, 2H), 5.04 (s, 2H), 6.95 (dd, 1H), 7.10 (s, 1H), 7.20-
10 7.60 (m, 7H).

PREPARATION OF IMIDAZOLES OF FORMULA (8, 33, 35)

Using the same procedure as described in example 2, but replacing the 4-cyanophenylhydrazine, hydrochloride by:

- 15 - N¹-(4-cyanophenylmethyl)-N¹-(4-methoxyphenyl)hydrazine,
the following compound was obtained:

EXAMPLE 9

4-[N-(1H-imidazol-1-yl)-N-(4-methoxyphenyl)amino]methylbenzonitrile

- 20 ¹H-NMR (DMSO d₆) : 3.70 (s, 3H), 4.90 (s, 2H), 6.60-7.00 (m, 5H), 7.40 (s, 1H), 7.55 (d, 2H), 7.70 (s, 1H), 7.78 (d, 2H).

Crystallization from hydrochloric ethanol yielded white crystals (5.70 g, 66 %).

mp: 207°C

- ¹H-NMR (DMSO d₆) : 3.70 (s, 3H), 4.97 (s, 2H), 6.93 (d, 2H), 7.13 (d, 2H), 7.45
25 (d, 2H), 7.70 (s, 1H), 7.84 (d, 2H), 8.04 (s, 1H), 8.18 (s, 1H), 9.55 (s, 1H).

EXAMPLE 10

4-[N-(4-hydroxyphenyl)-N-(1H-imidazol-1-yl)amino]methylbenzonitrile

- A solution of boron tribromide (60 ml, 60.00 mmol) in 20 ml of dichloromethane
30 is added to a cold (0-5°C) solution of 4-[N-(1H-imidazol-1-yl)-N-(4-methoxyphenyl)amino]methylbenzonitrile (4.60 g, 15.11 mmol). After 1 h at room temperature the mixture was hydrolysed with saturated aqueous NaHCO₃, filtered, washed with water (50 ml) and with dichloromethane (20 ml) to give a

brown solid (4.00 g). Crystallization from acetone yielded a brown solid (3.00 g, 68%) mp: 150°C.

¹H-NMR (DMSO d₆) : 4.84 (s, 2H), 6.70 (s, 4H), 6.90 (s, 1H), 7.45-7.62 (m, 3H), 7.62-7.90 (m, 3H), 9.25 (s, 1H).

5 **EXAMPLE 11**

4-[N-(4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile

4-hydroxybenzylbromide (15.6 g, 84.3 mmol, prepared following Wissner A. et al., J.Med.Chem. 1992, 35, 1650) was added to a mixture of 4-[N-(1H-imidazol-1-yl)amino]benzonitrile (10.00 g, 54.30 mmol) and K₂CO₃ (8.20 g, 59.70 mmol)
10 in dry THF (150 ml) at room temperature. The mixture was then stirred at room temperature for 2 h and after poured into water and extracted with ethyl acetate, dried over Na₂SO₄, filtered, and concentrated under vacuum to give the crude product (16.00 g as solid). Crystallization from ethyl acetate with ethanol yielded
15 the expected product (6.50 g, 41%, mp: 180°C)

¹H-NMR (DMSO d₆) : 4.80 (s, 2H), 6.65 (d, 2H), 6.91 (s, 1H), 7.04 (d, 1H), 7.20 (s, 1H), 7.56 (s, 1H), 7.63 (d, 2H).

Using the same procedure but replacing the 4-hydroxybenzylbromide by:

- 20 - 3-chloro-4-hydroxybenzylbromide
- 3-bromo-4-hydroxybenzylbromide
- 4-hydroxy-3-methoxybenzylbromide
- 2,3,5,6-tetrafluoro-4-hydroxybenzylchloride (prepared following Angyal S.J. et al., J.Chem. Soc. 1950, 2141)
- 25 - 3-formyl-4-hydroxybenzylchloride (prepared following Angyal S.J. et al., J.Chem. Soc. 1950, 2141)
- 1-benzyloxy-4-(2-bromo-ethoxy)-benzene (prepared following Brinkman J. and al. Bioorg. Med. Chem. Lett., 1996, 6, 21, 2491-94)
- 2-chloro-5-chloromethyl-pyridine
- 30 - 4-(bromomethyl)benzene sulfonamide (prepared following Colescott R. and all. J. Am. Chem. Soc., 1957, 79, 4232-35)
- 4-(chloromethyl)-2-nitro-phenol (prepared following Bayer Patent: DE 132475)

- 5-chloromethyl-2-methoxy-benzoic acid (prepared following Leonard F. and all. J. Med. Chem., 1965, 8, 812-15)

the following compounds were respectively obtained:

EXAMPLE 12

5 **4-[N-(3-chloro-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile**

mp 195°C

¹H-NMR (DMSO_d₆) : 4.88 (s, 2H), 6.67 (d, 2H), 6.88 (d, 2H), 6.98 (s, 1H), 7.05 (dd, 1H), 7.24 (d, 1H), 7.33 (s, 1H), 7.70 (s, 1H), 7.72 (d, 2H), 10.28 (s, 1H).

10 **EXAMPLE 13**

4-[N-(3-bromo-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile

mp 198°C

15 ¹H-NMR (DMSO_d₆) : 4.90 (s, 2H), 6.65 (d, 2H), 6.85 (d, 1H), 6.99 (s, 1H), 7.07 (d, 1H), 7.30 (s, 1H), 7.40 (s, 1H), 7.65 (s, 1H), 7.67 (d, 2H), 10.40 (s, 1H).

EXAMPLE 14

4-[N-(4-hydroxy-3-methoxyphenylmethyl)-N-(1H-imidazol-1-yl)amino] benzonitrile

mp 215°C

20 ¹H-NMR (DMSO_d₆) : 3.70 (s, 3H), 4.89 (s, 2H), 6.68 (s, 2H), 6.70 (d, 2H), 6.80 (s, 1H), 6.99 (s, 1H), 7.30 (s, 1H), 7.63 (s, 1H), 7.72 (d, 2H), 9.20 (s, 1H).

EXAMPLE 15

4-[N-(2,3,5,6-tetrafluoro-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino] benzonitrile

25 mp 243°C

¹H-NMR (DMSO_d₆): 5.09 (s, 2H), 6.72 (d, 2H), 7.00 (s, 1H), 7.32 (s, 1H), 7.69 (d, 2H), 7.77 (s, 1H), 11.80 (s, 1H).

EXAMPLE 16

30 **4-[N-(3-formyl-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile**

mp 160°C

¹H-NMR (DMSO-d₆) : 4.95 (s, 2H), 6.70 (d, 2H), 6.90 (s, 1H), 6.96 (d, 1H), 7.35 (s, 1H), 7.44 (dd, 1H), 7.58 (d, 1H), 7.67 (s, 1H), 7.71 (d, 2H), 10.20 (s, 1H), 10.75 (s, 1H).

EXAMPLE 17

- 5 **4-{N-[2-(4-benzyloxy-phenoxy)ethyl]-N-(1H-imidazol-1-yl)amino}benzonitrile**

¹H-NMR (DMSO d₆) : 3.95-4.10 (m, 2H), 4.11-4.28 (m, 2H), 5.01 (s, 2H), 6.60 (d, 2H), 6.82 (d, 2H), 7.95 (d, 2H), 7.03-7.50 (m, 7H), 7.69 (d, 2H), 7.88 (s, 1H).

EXAMPLE 18

- 10 **4-{N-[(6-chloropyridin-3-yl)methyl]-N-(1H-imidazol-1-yl)amino}benzonitrile**

mp 156°C

¹H-NMR (DMSO d₆) : 5.10 (s, 2H), 6.71 (d, 2H), 7.00 (s, 1H), 7.42 (s, 1H), 7.49 (d, 1H), 7.55-7.90 (m, 4H), 8.34 (d, 1H).

- 15 **EXAMPLE 19**

4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}benzene sulfonamide

mp 150°C

- 20 ¹H-NMR (DMSO d₆) : 5.15 (s, 2H), 6.62 (d, 2H), 7.00 (s, 1H), 7.36 (s, 2H), 7.45 (s, 1H), 7.55 (d, 2H), 7.65-7.90 (m, 5H).

EXAMPLE 20

4-[N-(4-hydroxy-3-nitrophenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile

mp 205°C

- 25 ¹H-NMR (DMSO d₆) : 4.98 (s, 2H), 6.72 (d, 2H), 7.00 (s, 1H), 7.06 (d, 1H), 7.39 (s, 1H), 7.48 (d, 1H), 7.60-7.78 (m, 3H), 7.82 (s, 1H).

EXAMPLE 21

5-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-2-methoxybenzoic acid

- 30 mp 187°C

¹H-NMR (DMSO d₆) : 3.79 (s, 3H), 5.00 (s, 2H), 6.69 (d, 2H), 6.98 (s, 1H), 7.05 (d, 1H), 7.31 (s, 1H), 7.39 (dd, 1H), 7.55 (d, 1H), 7.62-7.85 (m, 3H).

EXAMPLE 22**4-[N-(1H-imidazol-1-yl)-N-(4-nitrophenyl)amino]benzonitrile**

4-[N-(1H-imidazol-1-yl)amino]benzonitrile (10.00 g, 54.30 mmol) was added portionwise to a suspension of potassium tert-butoxide (6.69 g, 59.73 mM) in DMSO (100 ml) at (10-15°C) with stirring. The mixture was stirred for 30 mn at room temperature, and then 4-nitro-fluorobenzene (7.60 g, 54.00 mM) in DMSO (15 ml) was added dropwise while keeping the temperature below 30°C. After 2h, the mixture was poured into water (800 ml) and the resulting precipitate was collected by filtration and purified by crystallization from ethanol (1.00 g, 48%, mp: 188°C).

¹H-NMR (DMSO-d₆) : 7.00 (d, 2H), 7.17 (s, 1H), 7.26 (d, 2H), 7.65 (s, 1H), 7.90 (d, 2H), 8.20 (s,1H), 7.22 (d,2H).

Using the same procedure but replacing the 4-nitro-fluorobenzene by:

- 6-chloro-nicotinoyl chloride
- 4-fluorophenylacetyl chloride
- 4-hydroxyphenylacetyl chloride
- 4-hydroxyphenylpropanoyl chloride (prepared following Elias H. and all. Macromol. Chem. Phys., 1981, 182, 681-86)
- 4-phenylmethoxybenzene sulfonyl chloride (prepared following Toja E. and all. Eur. J.Med.Chem. 1991, 26, 403-13)

the following compounds were respectively obtained:

EXAMPLE 23**6-chloro-N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)nicotinamide**

mp 132°C

¹H-NMR (DMSO d₆) : 6.98 (s, 1H), 7.40-7.62 (m, 3H), 7.70 (s, 1H), 7.95 (d, 2H), 8.00 (d, 1H), 8.19 (s,1H), 8.57 (d, 1H).

EXAMPLE 24**N-(1H-imidazol-1-yl)-N-(4-cyanophenyl)-2-(4-fluorophenyl)acetamide**

mp 131°C

¹H-NMR (DMSO d₆) : 3.57 (s, 2H), 7.00-7.35 (m, 5H), 7.55 (d, 2H), 7.70 (s, 1H), 7.93 (d, 2H), 8.19 (s,1H).

EXAMPLE 25**N-(1H-imidazol-1-yl)-N-(4-cyanophenyl)-2-(4-hydroxyphenyl)acetamide**

¹H-NMR (DMSO d₆) : 3.32 (s, 2H), 6.65 (d, 2H), 6.87 (d, 2H), 7.08 (s, 1H), 7.50 (d, 2H), 7.70 (s, 1H), 7.90 (d, 2H), 8.10 (s, 1H), 9.30 (s, 1H).

EXAMPLE 26**N-(4-cyanophenyl)-3-(4-hydroxyphenyl)-N-(1H-imidazol-1-yl)propanamide**

mp 172°C

¹H-NMR (DMSO d₆) : 2.25-2.60 (m, 2H), 2.65-2.90 (m, 2H), 6.63 (d, 2H), 6.90 (d, 2H), 7.08 (s, 1H), 7.51 (d, 2H), 7.61 (s, 1H), 7.90 (d, 2H), 8.10 (s, 1H), 9.20 (s, 1H).

EXAMPLE 27**N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-4-(phenylmethoxy)-benzensulfonamide**

¹H-NMR (CDCl₃) : 5.14 (s, 2H), 6.93 (t, 1H), 7.00-7.15 (m, 3H), 7.30-7.45 (m, 7H), 7.50 (s, 1H), 7.66 (d, 2H), 7.68 (d, 2H).

PREPARATION OF IMIDAZOLES OF FORMULA (15), (21)**EXAMPLE 28****5-Nitro-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)]-benzothiophene-2-carboxamide**

5-Nitro-benzothiophene-2-carbonyl chloride (commercial compound, 10.00 g, 41.00 mmol) was added to a mixture of 4-[N-(1H-imidazol-1-yl)amino]benzonitrile (7.55 g, 41.00 mmol), TEA (20 ml, 143.00 mmol) in dry THF (150 ml) at room temperature. The mixture was then stirred at room temperature overnight and the precipitate was filtrated, washed with THF, water to give the crude product as a solid (9.26g). Crystallization with ethanol yielded white crystals (3.50 g, mp: 221°C).

¹H-NMR (DMSO d₆): 7.10 (s, 1H), 7.54 (s, 1H), 7.70 (d, 2H), 7.82 (s, 1H), 7.98 (d, 2H), 8.15-8.40 (m, 3H), 8.89 (s, 1H).

Using the same procedure but replacing the 5-nitro-benzothiophene-2-carbonyl chloride by:

6-methoxy-benzothiophene-3-acetyl chloride (described by Sauter F., Monatshefte fuer Chemie, 1968, 99, 2, 610-15) the following compound was

5 obtained:

EXAMPLE 29

N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-2-(6-methoxy-benzothien-3-yl) acetamide

mp: 104°C

10 ¹H-NMR (DMSO d₆): 3.75-3.85 (m, 5H), 7.02 (dd, 1H), 7.13 (s, 1H), 7.30 (s, 1H), 7.50-7.75 (m, 4H), 7.80 (s, 1H), 7.90 (d, 2H), 8.25 (s, 1H).

EXAMPLE 30

4-{N-[1H-imidazol-1-yl]-N-[(6-methoxy-benzothien-3-yl)methyl]amino} benzonitrile

15 3-Chloromethyl-6-methoxy-benzothiophene (12.35 g, 58.06 mmol) was added to a mixture of 4-[N-(1H-imidazol-1-yl)amino]benzonitrile (9.72 g, 52.78 mmol), K₂CO₃ (14.60 g, 105.56 mmol) and potassium iodide (0.10g, 0.60 mmol) in dry DMF (70 ml) at room temperature. The mixture was then stirred at room temperature overnight and after poured into water and extracted with ethyl

20 acetate, dried over Na₂SO₄, filtered, and concentrated under vacuum to give the crude product as solid (14.30 g). Flash chromatography on silica gel (toluene / dioxan: 6/4) yielded the expected product (10.50 g, 55 %, powder). Crystallization with ethanol yielded white crystals (7.30 g, mp: 164°C).

¹H-NMR (DMSO d₆): 3.80 (s, 3H), 5.25 (s, 2H), 6.74 (d, 2H), 6.93 (s, 1H), 7.02 (dd, 1H), 7.28 (s, 1H), 7.40 (s, 1H), 7.53 (s, 1H), 7.55 (s, 1H), 7.68 (d, 1H), 7.75 (d, 2H).

25

EXAMPLE 31

4-[N-(6-benzyloxy-1,1-dioxido-benzothien-3-yl)-N-(1H-imidazol-1-yl)amino] benzonitrile

30 4-[N-(1H-imidazol-1-yl)amino]benzonitrile (0.50 g, 27.14 mmol) was added portionwise to a suspension of potassium tert-butoxide (0.35 g, 31.00 mmol) in THF (20 ml) at (10-15°C) with stirring. The mixture was stirred for 30 mn at room temperature, and then 3-bromo-6-benzyloxy-benzothiophene-1,1-dioxyde

(1.10 g, 31.33 mM) in THF (5 ml) was added dropwise while keeping the temperature below 30°C. After one night, the mixture was poured into water (200 ml) and extracted with ethyl acetate, dried over Na₂SO₄, filtered, and concentrated under vacuum to give the crude product as an oil (2.50 g). Flash chromatography on silica gel (toluene / 1,4-dioxan: 7/3) and crystallisation in ethanol yielded light brown crystals (1.20 g, 95%, mp: 146°C).

¹H-NMR (DMSO d₆): 5.22 (s, 2H), 6.48 (s, 1H), 6.49 (d, 1H), 7.05-7.20 (m, 2H), 7.25-7.50 (m, 8H), 7.60 (d, 1H), 7.73 (s, 1H), 7.94 (d, 2H).

10 Using the same procedure but replacing the 3-bromo-6-benzyloxy-benzothiophene-1,1-dioxide by:

6-benzyloxy-2-(chloromethyl)-benzothiophene

the following compound was obtained:

EXAMPLE 32

15 **4-[N-[(6-benzyloxy-benzothien-2-yl)methyl]-N-(1H-imidazol-1-yl)amino] benzonitrile**

¹H-NMR (DMSO d₆): 5.12 (s, 2H), 5.30 (s, 2H), 6.72 (d, 2H), 7.00 (s, 1H), 7.04 (dd, 1H), 7.23 (s, 1H), 7.27-7.90 (m, 11H).

20 PREPARATION OF IMIDAZOLES OF FORMULA (10), (16)

EXAMPLE 33

4-{N-[(6-hydroxy-benzothien-3-yl)methyl]-N-[1H-imidazol-1-yl]amino} benzonitrile

A solution of 4-{N-[1H-imidazol-1-yl]-N-[(6-methoxy-benzothien-3-yl)methyl]amino} benzonitrile (0.50 g, 1.39 mmol) in 10 ml of methylene chloride is added at room temperature to a solution 1M of boron tribromide in methylene chloride (1.50 ml, 1.52 mmol). After 2h at room temperature the mixture was hydrolysed with saturated aqueous NaHCO₃, extracted with dichloromethane, dried over Na₂SO₄, filtered, and concentrated in vacuum. The crude product was purified by flash chromatography on silica gel (toluene / dioxan: 6/4) to give the expected product (0.30 g, 62 %, powder). Crystallization with ethanol yielded white crystals (0.10 g, mp: 169°C).

¹H-NMR (DMSO d₆): 5.24 (s, 2H), 6.72 (d, 2H), 6.87 (dd, 1H), 6.94 (s, 1H), 7.27 (d, 2H), 7.29 (s, 1H), 7.55 (s, 1H), 7.56 (d, 1H), 7.75 (d, 2H), 9.67 (s, 1H).

Using the same procedure but replacing the 4-{N-[1H-imidazol-1-yl]-N-[(6-methoxy-benzothiophen-3-yl)methyl]amino}benzonitrile by:

N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-2-(6-methoxy-benzothien-3-yl)-acetamide, the following compound was obtained:

EXAMPLE 34

N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-2-(6-hydroxy-benzothien-3-yl) acetamide

mp 179°C

¹H-NMR (DMSO d₆): 3.72 (s, 2H), 6.85 (dd, 1H), 7.10 (d, 2H), 7.25 (d, 1H), 7.40-7.70 (m, 3H), 7.80 (s, 1H), 7.91 (d, 2H), 8.24 (s, 1H), 9.60 (s, 1H).

EXAMPLE 35

4-[N-(3-amino-4-hydroxy-phenylmethyl)-N-(1H-imidazol-1-yl)amino] benzonitrile

4-[N-(4-hydroxy-3-nitro-phenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile (2 g, 5.97 mmol) was hydrogenated over a suspension of Pd/C (10 % Wt on carbon, 0.2 g) in ethanol (30 ml). When TLC showed complete reaction, the mixture was filtered on celatum, rinsed with EtOH. The solvent was concentrated under vacuum. Crystallisation from EtOH/petroleum ether yielded orange crystals (1.1 g, 60.5 %, mp: 208°C).

¹H-NMR (DMSO d₆) : 4.60 (s, 2H), 4.79 (s, 2H), 6.80 (d, 1H), 6.40-6.80 (m, 4H), 6.99 (s, 1H), 7.25 (s, 1H), 7.50-7.80 (m, 3H), 9.07 (s, 1H).

Using the same procedure but replacing the 4-[N-(4-hydroxy-3-nitro-benzyl)-N-(1H-imidazol-1-yl)amino]benzonitrile by:

- 4-{N-[2-(4-benzyloxy-phenoxy)ethyl]-N-(1H-imidazol-1-yl)amino]}
- benzonitrile
- N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-4-(phenylmethoxy)-benzensulfonamide

the following compounds were respectively obtained:

EXAMPLE 36

4-{N-[2-(4-hydroxyphenoxy)ethyl]-N-(1H-imidazol-1-yl)amino}benzonitrile

mp 188°C

- 5 ¹H-NMR (DMSO d₆) : 4.99 (t, 2H), 4.17 (t, 2H), 6.40-6.85 (m, 6H), 7.10 (s, 1H), 7.40 (s, 1H), 7.68 (d, 2H), 7.86 (s, 1H), 8.95 (s, 1H).

EXAMPLE 37

N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-4-hydroxybenzensulfonamide

mp 248°C

- 10 ¹H-NMR (DMSO d₆) : 6.95 (d, 2H), 7.00 (s, 1H), 7.24 (s, 1H), 7.49 (d, 2H), 7.52 (d, 2H), 7.88 (s, 1H), 7.92 (d, 2H), 10.95 (s, 1H).

EXAMPLE 38

4-[N-[(6-hydroxy-1,1-dioxido-benzothien-3-yl)]-N-(1H-imidazol-1-yl)amino] benzonitrile

- 15 A mixture of 4-[N-[(6-benzyloxy-1,1-dioxido-benzothien-3-yl)]-N-(1H-imidazol-1-yl) amino]benzonitrile (3.00 g, 6.78 mmol), 10% Pd/C (0.50 g), THF (30 ml) and solution of ammonium formate (25% in H₂O, 30 ml) was stirred at ambient temperature for 6 h and filtered. The mixture was poured into water and extracted with ethyl acetate, dried over Na₂SO₄, filtered, and concentrated under vacuum to give the crude product (2.50 g as solid). Crystallisation from ethanol afforded white crystals (0.80 g, 26 %, mp: 260°C).
- 20

¹H-NMR (DMSO d₆): 6.25 (s, 1H), 6.29 (d, 1H), 6.82 (dd, 1H), 7.10 (s, 1H), 7.15 (s, 1H), 7.30 (d, 2H), 7.70 (s, 1H), 7.91 (d, 2H), 8.25 (s, 1H).

- 25 Using the same procedure but replacing the 4-[N-[(6-benzyloxy-1,1-dioxido-benzothien-3-yl)]-N-(1H-imidazol-1-yl)amino]benzonitrile by:

4-[N-[(6-benzyloxy-benzothien-2-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile

the following compound was obtained:

- 30 **EXAMPLE 39**

4-[N-[(6-hydroxy-benzothien-2-yl)methyl]-N-(1H-imidazol-1-yl)amino] benzonitrile

mp 230°C

¹H-NMR (DMSO d₆): 5.28 (s, 2H), 6.70 (d, 2H), 6.82 (dd, 1H), 7.00 (s, 1H), 7.15-7.21 (m, 2H), 7.31 (s, 1H), 7.55 (d, 1H), 7.69-7.80 (m, 3H), 9.63 (s, 1H).

PREPARATION OF IMIDAZOLES OF FORMULA (12), (18)

5 **EXAMPLE 40**

4-[N-(4-aminophenyl)-N-(1H-imidazol-1-yl)amino]benzonitrile

Hydrazine (1.52 ml, 49.00 mmol) was added portionwise to a suspension of 4-[N-(1H-imidazol-1-yl)-N-(4-nitrophenyl)amino]benzonitrile (3.00 g, 9.80 mM) and ruthenium, 5 wt. % on carbon (0.30 g, 0.15 mM) in ethanol (35 ml) at reflux
10 with stirring. When TLC showed complete reaction, the mixture was cooled and the catalyst was filtered. The solvent was concentrated under vacuum. The residue was poured into water and extracted with dichloromethane, dried over Na₂SO₄, filtered, and concentrated under vacuum to give the crude product (2.50g as solid). Crystallization from ethyl acetate with ethanol yielded the
15 expected product (1.30 g, 50%, mp: 147°C).

¹H-NMR (DMSO d₆) : 5.50 (s, 2H), 6.30 (d, 2H), 6.69 (d, 2H), 7.09 (s, 1H), 7.29 (d, 2H), 7.63 (s, 1H), 7.65 (d, 2H), 8.14 (s, 1H).

EXAMPLE 41

20 **5-Amino-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)]-benzothiophene-2-carboxamide**

Stannous chloride dihydrate (13.10 g, 58.00 mmol) was added portionwise to a stirred solution of 5-nitro-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)]-benzothiophene-2-carboxamide (4.50 g, 11.6 mM) in ethanol (100 ml). The mixture was heated under reflux. When TLC showed complete reaction, the
25 mixture was cooled and basified with saturated bicarbonate solution. The mixture was extracted with ethyl acetate, dried over Na₂SO₄, filtered, and concentrated under vacuum to give the crude product as a solid (3.90g). Crystallization from methanol yielded the expected product (2.60 g, 63%, mp: 214°C).

¹H-NMR (DMSO d₆): 6.70-6.98 (m, 3H), 7.07 (s, 1H), 7.40-7.25 (m, 4H), 7.26-
30 7.96 (m, 3H).

GENERAL PROCEDURE OF SULFAMOYLATION**PREPARATION OF SULFAMATES (11, 17, 23) AND AMINOSULFONYLAMINES (13, 19, 25)****EXAMPLE 42****5 Sulfamic acid 4-[N-(4-cyanophenylmethyl)-N-(1H-imidazol-1-yl)amino]phenyl ester**

Sulfamoyl chloride (2.39 g, 20.69 mmol) was added to a solution of 4-[N-(4-hydroxyphenyl)-N-(1H-imidazol-1-yl)amino]methylbenzonitrile (1.00 g, 3.45 mmol) in dry DMAc (36 ml) with ice cooling. The mixture was then stirred at
 10 room temperature for 6 h. After addition of TEA (3.40 ml, 24.73 ml), the mixture was poured into cold brine and extracted with ethyl acetate, dried over Na₂SO₄, filtered, and concentrated under vacuum to give the crude product (0.70 g as solid). Crystallization from ethyl acetate yielded the expected product (0.40 g, 31%, mp: 60°C).

15 ¹H-NMR (DMSO d₆) : 5.00 (s, 2H), 6.70 (d, 2H), 6.92 (s, 1H), 7.19 (d, 2H), 7.40 (s, 1H), 7.55 (d, 2H), 7.74 (s, 1H), 7.77 (d, 2H), 7.91 (s, 2H).

Using the same procedure but replacing the 4-[N-(4-hydroxyphenyl)-N-(1H-imidazol-1-yl)amino]methylbenzonitrile by:

- 20 • 4-[N-(4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile
- 4-[N-(3-chloro-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile
- 4-[N-(3-bromo-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile
- 25 • 4-[N-(3-methoxy-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino] benzonitrile
- 4-[N-(2,3,5,6-tetrafluoro-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino] benzonitrile
- 30 • 4-[N-(3-formyl-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino] benzonitrile
- 4-[N-(4-aminophenyl)-N-(1H-imidazol-1-yl)amino]benzonitrile
- N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-4-hydroxybenzensulfonamide

- 4-{N-[2-(4-hydroxyphenoxy)ethyl]-N-(1H-imidazol-1-yl)amino}benzonitrile
- N-(1H-imidazol-1-yl)-N-(4-cyanophenyl)-2-(4-hydroxyphenyl)acetamide
- 5 • N-(4-cyanophenyl)-3-(4-hydroxyphenyl)-N-(1H-imidazol-1-yl)propanamide
- 4-[N-(3-amino-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile
- 5-Amino-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)]-benzothiophene-2-carboxamide
- 10 • 4-[N-[(6-hydroxy-1,1-dioxido-benzothien-3-yl)]-N-(1H-imidazol-1-yl)amino]benzonitrile
- N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-2-[(6-hydroxy-benzothien-2-yl)] acetamide
- 15 • 4-{N-[(6-hydroxy-benzothien-3-yl)methyl]-N-[1H-imidazol-1-yl]amino}benzonitrile
- 4-{N-[(6-hydroxy-benzothien-2-yl)methyl]-N-[1H-imidazol-1-yl]amino}benzonitrile

the following compounds were respectively obtained:

20 **EXAMPLE 43**

Sulfamic acid-4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1yl)amino]methyl}phenyl ester

mp 172°C

¹H-NMR (DMSO-d₆) : 5.00 (s, 2H), 6.65 (d, 2H), 7.00 (s, 1H), 7.22 (d, 2H), 7.40 (s, 1H), 7.42 (d, 2H), 7.70 (s, 1H), 7.75 (d, 2H), 8.00 (s, 2H).

EXAMPLE 44

Sulfamic acid 2-chloro-4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1yl)amino]methyl}phenyl ester

¹H-NMR (DMSO-d₆) : 5.05 (s, 2H), 6.63 (d, 2H), 7.00 (s, 1H), 7.35-7.45 (m, 3H), 7.54 (s, 1H), 7.70 (d, 2H), 7.80 (s, 1H), 8.29 (s, 2H).

EXAMPLE 45

Sulfamic acid 2-bromo-4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1yl)amino] methyl}phenyl ester , hydrochloride

Crystallisation was performed in EtOH/HCl (mp 145°C)

¹H-NMR (DMSO-d₆) : 5.15 (s, 2H), 6.90 (d, 2H), 7.48 (s, 2H), 7.60-7.85 (s, 5H), 8.08 (s, 1H), 8.32 (s, 2H), 9.51 (s, 1H).

EXAMPLE 46

Sulfamic acid 2-methoxy-4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino] methyl}phenyl ester

mp 211°C

¹H-NMR (DMSO-d₆) : 3.77 (s, 3H), 5.02 (s, 2H), 6.75 (d, 2H), 6.92 (d, 1H), 7.05 (s, 2H), 7.25 (d, 1H), 7.45 (s, 1H), 7.71 (d, 2H), 7.80 (s, 1H), 7.93 (s, 2H).

EXAMPLE 47

Sulfamic acid 2,3,5,6-tetrafluoro-4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino] methyl}phenyl ester

¹H-NMR (DMSO-d₆) : 5.37 (s, 2H), 5.83 (s, 2H), 6.86 (d, 2H), 7.08 (s, 1H), 7.41 (s, 1H), 7.79 (s, 1H), 7.81 (d, 2H).

EXAMPLE 48

4-[N-[(2,2-dioxido-1,2,3-benzoxathiazin-6-yl)methyl]-N-(1H-imidazol-1-yl)amino] benzonitrile

mp 180°C

¹H-NMR (DMSO-d₆) : 5.17 (s, 2H), 6.67 (d, 2H), 7.02 (s, 1H), 7.46 (s, 1H), 7.50 (d, 2H), 7.74 (d, 2H), 7.82 (dd, 1H), 7.87 (s, 1H), 8.02 (d, 1H), 9.19 (s, 1H).

EXAMPLE 49

N-{4-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]phenyl}sulfamide

mp 121°C

¹H-NMR (DMSO-d₆) : 6.38 (d, 2H), 7.09 (s, 1H), 7.20 (s, 1H), 7.22 (d, 2H), 7.51 (d, 2H), 7.63 (s, 1H), 7.69 (d, 2H), 8.19 (s, 1H)

EXAMPLE 50

Sulfamic acid 4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]sulfonyl}phenyl ester, hydrochloride

Crystallisation was performed in EtOH/HCl

mp 200°C

¹H-NMR (DMSO-d₆) : 7.57 (d, 2H), 7.60 (s, 1H), 7.70 (d, 2H), 7.80-8.10 (m, 5H), 8.40 (s, 2H), 9.30 (s, 1H).

EXAMPLE 51

Sulfamic acid 4-{2-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]ethoxy} phenyl ester

mp 174°C

- 5 ¹H-NMR (DMSO-d₆) : 4.08 (t, 2H), 4.23 (t, 2H), 6.60 (d, 2H), 6.95 (d, 2H), 7.10 (s, 1H), 7.18 (d, 2H), 7.42 (s, 1H), 7.70 (d, 2H), 7.88 (s, 3H).

EXAMPLE 52

Sulfamic acid 4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-carbamoyl]-methyl} phenyl ester

- 10 ¹H-NMR (DMSO-d₆) : 3.32 (s, 2H), 7.12 (s, 1H), 7.18-7.32 (m, 4H), 7.50-7.60 (d, 2H), 7.71 (s, 1H), 7.85-8.05 (m, 4H), 8.20 (s, 1H).

EXAMPLE 53

Sulfamic acid 4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]-3-oxopropyl} phenyl ester

- 15 mp 100°C
¹H-NMR (DMSO-d₆) : 2.45 (t, 2H), 3.85 (t, 2H), 7.10 (s, 1H); 7.12-7.30 (m, 4H), 7.55 (d, 2H), 7.65 (s, 1H), 7.90 (m, 4H), 8.12 (s, 1H).

EXAMPLE 54

- 20 **Sulfamic acid 3-(aminosulfonyl)amino-4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}phenyl ester**

mp 197°C

¹H-NMR (DMSO-d₆) : 5.00 (s, 2H), 6.71 (d, 2H), 6.93 (d, 1H), 7.02 (s, 2H), 7.21 (d, 1H), 7.48 (s, 1H), 7.61 (d, 2H), 7.74 (s, 1H), 7.92 (s, 2H), 8.10 (s, 2H).

EXAMPLE 55

- 25 **5-(Aminosulfonyl)amino-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)]-benzothiophene-2-carboxamide**

mp: 169°C

¹H-NMR (DMSO d₆): 6.87 (d, 1H), 7.05-7.30 (m, 4H), 7.56 (s, 1H), 7.67 (d, 2H), 7.80-8.10 (m, 4H), 7.86 (s, 1H), 9.65 (s, 1H).

- 30 **EXAMPLE 56**

Sulfamic acid 3-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]-1,1-dioxido-benzothien-6-yl ester

mp 172°C

¹H-NMR (DMSO d₆): 6.24 (s, 1H), 6.29 (d, 1H), 6.85 (dd, 1H), 7.15 (s, 1H), 7.20 (s, 1H), 7.30 (d, 2H), 7.65 (s, 1H), 7.95 (d, 2H), 8.05 (s, 2H), 8.25 (s, 1H).

EXAMPLE 57

Sulfamic acid 3-{2-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]-2-oxoethyl}-benzothien-6-yl ester

¹H-NMR (DMSO d₆): 3.84 (s, 2H), 7.15 (s, 1H), 7.28-7.37 (dd, 1H), 7.51-7.65 (m, 3H), 7.80-7.98 (m, 5H), 8.01 (s, 2H), 8.29 (s, 1H).

EXAMPLE 58

Sulfamic acid 3-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-benzothien-6-yl ester

mp 193°C

¹H-NMR (DMSO d₆): 5.30 (s, 2H), 6.80 (d, 2H), 6.97 (s, 1H), 7.30-7.40 (m, 2H), 7.56 (s, 1H), 7.68 (s, 1H), 7.75 (d, 2H), 7.80-7.95 (m, 2H), 8.05 (s, 2H).

EXAMPLE 59

Sulfamic acid 2-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-benzothien-6-yl ester

mp 178 °C

¹H-NMR (DMSO d₆): 5.17 (s, 2H), 6.65 (d, 2H), 6.89 (dd, 1H), 6.94 (s, 1H), 7.20 (d, 2H), 7.27 (s, 1H), 7.45 (s, 1H), 7.58 (d, 1H), 7.72 (d, 2H), 8.00 (s, 2H).

20

Using the same procedure as example 13 but replacing the 4-[N-(4-hydroxyphenyl)-N-(1H-imidazol-1-yl)amino]methylbenzonitrile by:

- 4-[N-(3-bromo-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile

25 The following compounds was obtained as a by-product:

EXAMPLE 60

2-Bromo-4-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}phenyl amidimidodisulfate acid

mp 220°C

30 ¹H-NMR (DMSO d₆): 5.00 (s, 2H), 5.70 (s, 2H), 6.65 (d, 2H), 7.03 (s, 1H), 7.42 (dd, 1H), 7.50 (s, 1H), 7.55 (d, 1H), 7.65 (d, 1H), 7.72 (d, 2H), 7.81 (s, 1H).

PREPARATION OF OTHER COMPOUNDS (8)**EXAMPLE 61****4-[N-[(2,2-dioxido-3,4-dihydro-1,2,3-benzoxathiazin-6-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile**

- 5 To a suspension of 4-[N-[(2,2-dioxido-1,2,3-benzoxathiazin-6-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile (0.40 g, 1.05 mmol) in methanol (8 ml) was added portionwise NaBH₄ (0.08 g, 2.11 mmol) and the reaction mixture was stirred at room temperature for 3 h. After addition of a saturated solution of NH₄Cl (3 ml) and water (40 ml), the precipitate obtained was filtered, washed
10 with water and dried to give a white solid. Crystallization from ethyl acetate with ethanol yielded the expected product (0.40 g, 82%, mp: 190°C).

¹H-NMR (DMSO-d₆) : 4.55 (d, 2H), 5.04 (s, 2H), 6.64 (d, 2H), 7.02 (s, 1H), 7.03 (d, 1H), 7.27 (d, 1H), 7.30 (s, 1H), 7.41 (s, 1H), 7.70 (d, 2H), 7.85 (s, 1H), 8.56 (t, 1H).

15 EXAMPLE 62**5-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-2-hydroxybenzoic acid**

- A mixture of 5-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-2-methoxybenzoic acid (1.00g, 2.85 mmol) and piperidine (0.42ml, 8.57 mmol) in
20 dimethylacetamide (DMA) (2 ml) was heated at 150°C. When starting material had disappeared (TLC monitoring), the solvent was removed in vacuo. Flash chromatography on silica gel (MeOH / dichloromethane : 5/95) and crystallization from ethanol afforded white crystals (53 mg, 6 %, mp: 260°C).

- ¹H-NMR (DMSO d₆) : 4.85 (s, 2H), 6.54 (d, 1H), 6.66 (d, 2H), 6.95 (s, 1H), 7.05
25 (dd, 1H), 7.24 (s, 1H), 7.50-7.60 (m, 2H), 7.68 (d, 2H).

EXAMPLE 63**4-[N-(1H-imidazol-1-yl)-N-(phenyl)amino]benzonitrile**

- 4-[N-(1H-imidazol-1-yl)-N-(4-nitrophenyl)amino]benzonitrile (3.91 g, 12.80 mmol) was placed in a hydrogenation flask and dissolved in acetic anhydride (60 ml) and acetic acid (60 ml). Palladium 10 % on C, 0.20 g) was added, and the
30 bottle was attached to a Parr hydrogenation apparatus. Hydrogenation was carried out with shaking for 3 h at 25 psi H₂. The catalyst was removed by filtration and the solution was cooled in a bath for 30 mn. Sodium nitrite (0.97 g,

14.00 mmol) was added to the mixture, and the vessels are lightly capped. The flask is kept in the ice bath for 2 h and allowed to warm to room temperature overnight. The mixture was poured into ice/water and the solid was filtered, washed with water (50 ml) to give crude product. Flash chromatography
5 (toluen/dioxan: 6/4) and crystallization from ethyl acetate with ethanol yielded the expected product (0.40 g, 8 %, mp: 162°C).

¹H-NMR (DMSO d₆) : 6.60 (d, 2H), 7.00-7.50 (m, 9H), 7.69 (s, 1H).

PREPARATION OF BENZOXATHIAZOLE (8)

10 **EXAMPLE 64**

4-[N-(3-tosylamino-4-hydroxy-benzyl)-N-(1H-imidazol-1-yl)amino]benzonitrile

Tosyl chloride (3.5 g, 18.59 mmol) in CH₂Cl₂ (20 ml) was added dropwise to a solution of (TX 1840) (5.4 g, 17.7 mmol) and pyridine (19.47 mmol, 1.6 ml) in
15 CH₂Cl₂ (50 ml) at 0°C.

The mixture was then stirred at room temperature for 4h then poured into water and extracted with EtOAc, dried over Na₂SO₄, filtered and concentrated under vacuum to give the crude product (8.1 g as orange solid).

Flash chromatography on silica gel (toluene dioxane 7/3) yielded yellow solid.

20 ¹H-NMR (DMSO d₆) : 2.32 (s, 3H), 4.85 (s, 2H), 6.60 (d, 2H), 6.62 (s, 1H), 6.80 (dd, 1H), 6.97 (s, 1H), 7.10-7.30 (m, 5H), 7.00 (d, 2H), 7.03 (s, 1H), 7.52 (d, 2H), 9.45 (s, 2H).

EXAMPLE 65

25 **4-[N-[(2,2-dioxido-3-tosyl-3H-1,2,3-benzoxathiazol-5-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile**

Sulfuryl chloride (0.60 ml, 7.40 mmol) in dichloromethane (50 ml) was added dropwise with stirring over a 30 mn period to 4-[N-(3-tosylamino-4-hydroxy-benzyl)-N-(1H-imidazol-1-yl)amino]benzonitrile (3.40 g, 7.40 mmol) and triethylamine (2.10 ml, 14.81 mmol) in dichloromethane (6 ml) at -78°C. After
30 an additional 15 mn, the mixture was allowed to room temperature for 4 h. The reaction mixture was diluted with water, extracted with dichloromethane. The organic layer was dried over sodium sulfate and concentrated under vacuum.

Flash chromatography on silica gel (toluene / dioxane : 5/5) yielded a pure oil (1.60 g, 41 %).

¹H-NMR (DMSO d₆) : 2.39 (s, 3H), 5.18 (s, 2H), 6.80 (d, 2H), 7.00-7.50 (m, 6H), 7.60-7.85 (m, 5H), 8.02 (s, 1H).

5 **EXAMPLE 66**

4-[N-[(2,2-dioxido-3H-1,2,3-benzoxathiazol-5-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile

Potassium fluoride (0.36 g, 6.14 mmol) in water (5 ml) was added to 4-[N-[(2,2-dioxido-3-tosyl-3H-1,2,3-benzoxathiazol-5-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile (1.60 g, 3.07 mmol) in acetonitrile (15 ml) at room temperature. The solution was stirred overnight, concentrated, extracted with dichloromethane. The organic layer was dried over sodium sulfate and concentrated under vacuum. Flash chromatography on silica gel (toluene / dioxane : 5/5) yielded an oil and crystallization from ethanol yielded the expected product (0.17 g, 15 %, mp: 230°C).

¹H-NMR (DMSO d₆) : 4.84 (s, 2H), 6.38 (dd, 1H), 6.42 (d, 1H), 6.63 (d, 1H), 6.69 (d, 2H), 7.15 (s, 1H), 7.43 (s, 1H), 7.70 (d, 2H), 8.02 (s, 1H).

PREPARATION OF PHENYL UREAS (27)

20 **EXAMPLE 67**

N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-N'-phenylurea

Phenyl isocyanate (3.6 ml, 32.6 mmol) was added to a solution of 4-[N-(1H-imidazol-1-yl)amino] benzonitrile (5 g, 27.17 mmol) in THF (50 ml). The mixture was stirred at 50°C overnight and after evaporation was crystallised in acetone/EtOH to give white crystals (2.5 g, 30.3 %, 178°C).

¹H-NMR (DMSO d₆) : 6.59 (s, 1H), 7.05-7.40 (m, 9H), 7.52 (d, 2H), 7.83 (s, 1H).

BIOLOGICAL TEST RESULTS

INHIBITION OF STEROID SULFATASE, AROMATASE AND CARBONIC ANHYDRASE II *IN VITRO*

Estrone sulfate (E_1S) is a major circulating plasma estrogen that is converted by
5 the steroid sulfatase enzyme into estrone (E_1), which in turn can be transformed
into estradiol (E_2) by enzymatic reduction.

The inhibition of E_2 synthesis by aromatase inhibitors has been clinically proved
to be a good way to halt the progress of hormone-dependent breast tumors.
More recently, the inhibition of the steroid sulfatase pathway emerged as an
10 alternative. Therefore, the development of compounds able to inhibit both
aromatase and steroid sulfatase inhibition appears as a new and suitable
approach to inhibit tumor growth.

Human carbonic anhydrases catalyses the conversion between carbon dioxide
(CO_2) and the bicarbonate ion (HCO_3^-), and are involved in numerous
15 physiological and pathological processes. They include hormone-dependent and
non-hormone-dependent cancerogenesis, metastasis process and hypoxic tumors
that are less responsive to classical chemo/radio-therapy. The inhibition of
human carbonic anhydrases could be therefore a valuable additional activity for
dual aromatase and sulfatase inhibitor (DASI) compounds. In particular, EMATE
20 was found to have a human carbonic anhydrases inhibitory potency similar to
that of acetazolamide, a well-known sulfonamide human carbonic anhydrases
inhibitor. The aim of these experiments was to evaluate *in vitro* the inhibitory
potential of new compounds on aromatase and/or steroid sulfatase activity. In
addition, inhibition on human carbonic anhydrase II activity (as an exemple of
25 human carbonic anhydrase) for some of them were evaluated in comparison with
6,6,7 COUMATE and acetazolamide.

Materials & Methods

a) Aromatase activity

30 The JEG-3 cell line, derived from a human placental choriocarcinoma,
constitutively overexpresses human aromatase and represents a relevant model
for assaying putative aromatase inhibitors *in vitro*. Aromatase activity was
determined by the tritiated water method. In brief, cells were first seeded into

96-well microplates in decompemented fetal calf serum (dFCS) supplemented medium. 24 hours later, cells were rinsed and fresh medium containing 1β - 3 H-androstenedione as aromatase substrate was added together with test compounds at concentrations ranging from 10^{-12} M to 10^{-5} M. After 2 hours of incubation, a fraction of the medium was transferred to homologous new 96-well microplates and a dextran-coated charcoal solution was added to each well. Following standing on ice for 10 minutes, microplates were centrifuged (1500 g; 4°C). All steroids, including the radioactive substrate and the newly biosynthesized estrogens, were adsorbed on charcoal; only 3 H-water specifically formed during aromatisation of 1β - 3 H-androstenedione, remained in the supernatant. The radioactivity in the supernatant was measured by liquid scintillation counting. In parallel, cell solubilization was performed in ethylenediamine tetraacetate solution. DNA content was measured by a standard fluorimetric method using the Hoechst 33258 fluorochrome. Finally, aromatase activity was expressed in fmoles of 3 H-water formed/2 hours/ μ g DNA and aromatase inhibition as a percentage of control activity without inhibitor. A non-linear fit analysis (GraphPad Prism Software) of % of inhibition vs. inhibitor concentration allowed the determination of the 50 % inhibitory concentration (IC_{50}) : the lowest IC_{50} s correspond to the most potent inhibitors (**Table 1**).

20

b) Steroid sulfatase activity

The JEG-3 cell line is intrinsically very rich in human estrone sulfatase and therefore is a useful biological system to evaluate new steroid sulfatase inhibitors *in vitro*. Assays were carried out with cells in logarithmic growth phase on 96-well microplates. 24h before studies, cells were seeded in decompemented fetal calf serum (dFCS) supplemented medium. The medium was removed 24h later and cells were rinsed with PBS to eliminate any trace of dFCS. Then, 3 H- E_1 S was added in dFCS free medium, followed by test compounds at concentrations ranging from 10^{-12} M to 10^{-5} M. After 4 h of treatment, the medium was transferred into 96-deep-well microplates and centrifuged at 200 x g for 10 min to pellet cells before toluene extraction. A fraction of the medium was used for toluene extraction in order to separate the conjugated substrate from non-conjugated products. The radioactivity in the toluene phase was measured by

30

liquid scintillation counting. In parallel, after cell solubilization in a ethylenediamine tetraacetate solution, DNA content was measured by a standard fluorimetric method using the Hoechst 33258 fluorochrome. Finally, estrone sulfatase activity was expressed in pmoles of $^3\text{H-E}_1 + ^3\text{H-E}_2$ formed/4 hours/mg DNA and estrone sulfatase inhibition as a percentage of control activity without inhibitor. A non linear fit analysis (GraphPad Prism Software) of % inhibition vs. inhibitor concentration allowed the determination of the 50 % inhibitory concentration (IC_{50}): the lowest IC_{50} s correspond to the most potent inhibitors (Table 1).

c) Dual aromatase and steroid sulfatase activities

In order to evaluate the inhibitory potency of DASI compounds in parallel on both human aromatase and steroid sulfatase activities, a new *in vitro* model was set up using JEG-3 cells. The experimental conditions of the steroid sulfatase model previously described was adopted with slight modifications: the presence of the two substrates $^3\text{-HE}_1\text{S}$ and $1\beta\text{-}^3\text{H}$ -androstenedione into the medium and an incubation period of 2 hours. Results were expressed in pmoles of product formed ($^3\text{H-E}_1 + ^3\text{H-E}_2$ or $^3\text{H}_2\text{O}$)/2 hours/mg DNA (Table 1).

d) Alkaline phosphatase activity

Ishikawa cells were plated into 96-well microplates 48 hours before studies. The next day, the medium was replaced by phenol red free medium supplemented with 5 % charcoal-stripped dFCS. 24 hours later, the medium was renewed and compounds were added to the plated cells and incubated for an additional four-day period. For each compound, the tested concentrations ranged from 10^{-12} M to 10^{-5} M, and the final vehicle concentration did not exceed 0.1 %. At the end of the incubation period, alkaline phosphatase activity (APase) was assayed by a method involving the hydrolysis of p-nitrophenyl phosphate to p-nitrophenol and spectrophotometric determination of the product at 405 nm.

In brief, the microplates were first rinsed twice with cold phosphate buffered solution and then placed at -80°C for at least 15 minutes. After thawing at room temperature, 50 μl ice-cold solution containing 5 mM p-nitrophenyl phosphate was added to each well. After a 15 to 60 minute incubation period at room

temperature, the intensity of the yellow color generated by the production of *p*-nitrophenol was measured into each well at 405 nm.

For each tested concentration, APase activity, reflected by absorbance, was first expressed as fold increase over control (FI) and then as percentage of E₂ activity (10⁻⁸ M) chosen equal to 100 %. Sigmoidal dose-response curves were plotted (GraphPad Prism Software) and 50 % effective concentrations (EC₅₀) were calculated for each compound (**Table 1**).

e) Human carbonic anhydrase II activity

This assay was performed as described in the literature (Armstrong J. et al. Purification and properties of human erythrocyte carbonic anhydrases, J Biol Chem, 1966, 241: 5137-5149). Briefly, in this assay, human carbonic anhydrase II catalyses the conversion of *p*-nitrophenyl acetate into *p*-nitrophenol. The potential inhibitory effect of test compounds was evaluated by colorimetric determination of *p*-nitrophenol produced during the enzymatic reaction. The optical density levels obtained without inhibitor will be referred to as "total activity". The levels obtained without inhibitor and without the enzyme will be referred to as "blank" in order to assess any interference with the substrate during the assay (**Table 2**).

20

Table 1: Inhibition of aromatase, estrone sulfatase and estrogenic potency

Compounds	Aromatase activity		Sulfatase activity		Estrogenic potency	
	IC ₅₀ (nM) ± S.E.M.	n	IC ₅₀ (nM) ± S.E.M.	n	EC ₅₀ (nM) ± S.E.M.	n
letrozole	0.47 ± 0.08 ^a	8	nd ^b	6	nd	1
anastrozole	6.93 ± 1.07 ^a	6	nd ^b	6	nd	1
	2.5 ± 0.6 ^c	4	nd ^c			
A*	nd ^a	3	2.5 ± 0.5 ^b	4	nd	4
	nd ^c	4	3.6 ± 0.8 ^c	4		
Ex 11	0.17 ± 0.01 ^a	8	nd ^b	4	nd	4

Compounds	Aromatase activity		Sulfatase activity		Estrogenic potency	
	IC ₅₀ (nM) ± S.E.M.	n	IC ₅₀ (nM) ± S.E.M.	n	EC ₅₀ (nM) ± S.E.M.	n
Ex 13	0.17 ± 0.03 ^a	8	--	--	nd	4
	0.05 ± 0.02 ^c	4	nd ^c	4		
Ex 14	0.21 ± 0.05 ^a	4	nd ^b	1	nd	4
Ex 20	0.3 ^a	1	nd ^b	4	nd	4
Ex 22	0.32 ± 0.40 ^a	4	--	--	nd	1
Ex 26	50.6 ^a	1	nd ^b	1	--	--
Ex 30	0.06 ^c	1	--	--	--	--
Ex 35	2.2 ± 0.7 ^c	4	nd ^c	4	--	--
Ex 36	0.61 ± 0.21 ^c	4	nd ^c	4	--	--
Ex 37	52.0 ± 9.6 ^c	4	nd ^c	4	--	--
Ex 40	0.67 ± 0.10 ^a	8	--	--	nd	3
Ex 41	4.3 ^c	1	--	--	--	--
Ex 43	0.57 ± 0.16 ^a	4	5.5 ± 1.1 ^b	4	nd	4
Ex 45	0.13 ± 0.07 ^c	4	3.4 ± 0.4 ^c	4	nd	4
Ex 46	0.79 ± 0.26 ^a	4	13.8 ± 2.5 ^b	4	nd	4
Ex 48	0.82 ± 0.12 ^a	4	2657 ± 257 ^b	4	nd	1
Ex 51	0.34 ± 0.13 ^c	4	597.2 ± 67.4 ^c	4	--	--
Ex 61	1.06 ± 0.21 ^a	4	nd ^b	4	nd	1
Ex 62	81.8 ^c	1	nd ^c	1	--	--

A*:sulfamic acid, 3-cycloheptylmethylbenzothiophen-6-yl-1,1-dioxide-ester
(described in WO2004/101545)

nd: not detected

Materials & methods: ^a aromatase activity alone; ^b sulfatase activity alone; ^c dual
5 aromatase and sulfatase activity, ^d estrogenic potency.

Among the tested compounds, Ex 43, Ex 45 and Ex 46 showed a strong inhibition (IC_{50} about 10 nM) of human estrone sulfatase activity. In addition, the same compounds were shown to be strong inhibitors of aromatase activity ($IC_{50} < 1$ nM). Despite this dual activity, They were not estrogenic *in vitro*.

5

Table 2: Inhibition of human carbonic anhydrase II

Human carbonic anhydrase II activity		
Compounds	IC_{50} (nM) \pm S.E.M.	n
acetazolamide	11.3 ± 1.9	4
6,6,7 COUMATE	16.1 ± 1.5	4
EX 45	16.2 ± 3.3	4
EX 51	6.5 ± 0.9	4

Ex 45 and Ex 51 inhibited human carbonic anhydrase II *in vitro*.

10 **ANTI UTEROTROPHIC/ANTI STEROID SULFATASE ACTIVITY *IN VIVO***

Wistar female rats were ovariectomized and left to rest for 4 weeks. Prior to treatment, the absence of cyclicity was checked by vaginal smears. Animals were supplemented with estrone sulfate (E_1S) at 50 μ g/kg/day s.c., alone or combined with oral administration of potential sulfatase inhibitors, at 1 mg/kg/day for 4 days. The uteri were removed, freed of adjacent tissue and wet weighed (**Table 3**).

Estrone sulfatase activity is measured according to the method described by Purohit et al., with slight modifications. Briefly, uteri were thawed, weighed and homogenized. Aliquots of the supernatant were treated with dextran-coated charcoal and assayed for sulfatase. E_1S activity is assessed after 30 min of incubation with 5 nM of 3H - E_1S and 20 μ M of unlabelled E_1S as substrate. Estrone sulfatase activity is expressed as pmol /h/mg protein and reported as percentage of inhibition vs E_1S ; for uterus weights, results are expressed as % of inhibition of the E_1S induced stimulation

25

Table 3: Inhibition of uterus weight

Compound	% inhibition (uterus weight)	n
6,6,7 COUMATE	81	8
Ex 45	70	8

Since there is a direct linear correlation between sulfatase inhibition and uterus weight inhibition (WO 2004/101545), it can be concluded that this uterus activity results from significant inhibition of E_1S (> 90%). Ex 45 was chosen as a potential inhibitor of estrone sulfatase activity because of its lack of estrogenicity, and significant steroid sulfatase inhibition. These *in vivo* results were in agreement with *in vitro* results obtained in JEG-3 cells.

10 INHIBITION OF ESTRADIOL PEAK, AS *IN VIVO* AROMATASE INHIBITION MODEL

The aim of this experimentation was to determine a dose-related activity of test compounds in comparison with anastrozole on 17β -estradiol level 24 hours after one oral administration in female rats. Anastrozole, is a potent, non steroidal inhibitor of aromatase, which significantly inhibits estradiol levels at 3 $\mu\text{g/kg}$, 24 hours after one single oral administration in female rats. One hundred IOPS Wistar female rats, weighing 180 to 200 g, were accommodated by four in stainless steel mesh cages. Animals were allowed free access to a standard diet from Harlan Teklad 2016 pellets. Vaginal smears were performed for each animal, each morning, in order to establish the different phases of cycle. The rats which did not exhibit a regular estrous cycle, were excluded from the experimentation. Starting at 4.00 pm, animals received an oral administration of formulations. 24 hours later, samples of serum were collected and estradiol level were determined as previously described (Table 4).

Table 4: Inhibition of estradiol peak

Compound	% inhibition (estradiol peak)
Anastrozole	56
Ex 45	28

Ex 45 inhibited estradiol peak *in vivo*.

5 INHIBITION OF TUMOR GROWTH

MCF-7 xenograft into nude mice

MCF-7 cells, derived from human breast adenocarcinoma, are injected subcutaneously in ovariectomized athymic nude mice supplemented with a daily administration of subcutaneous estrone sulfate. Xenograft volumes are determined weekly. When tumor volumes reach significant increase, tested compound is orally administered at 1mg/kg/day for 8 weeks. Xenografts are measured, removed, weighed, and deep frozen for the determination of sulfatase activity, according to methods used for rat uteri

15 ER+ Human breast tumor tissue xenografted into nude rats

MCF-7 cells (ATCC), derived from human breast adenocarcinoma, were injected subcutaneously in ovariectomized athymic nude mice supplemented with a daily administration of subcutaneous estradiol. Xenograft volumes were determined weekly. After 8 weeks, tumor volumes reached 1000 mm³, providing 500 mg samples of estrogen responsive tissues for each animals. At this time each tumor was sliced and regrafted into naïve ovariectomized nude rat (Rnu/Rnu). Then animals were treated with estrone sulfate as precursor of estradiol by the steroid sulfatase pathway. When tumor volumes reached significant increase, distinct from the placebo, animals were randomized and separated in two groups. The first group was treated with Ex 45 and supplemented with daily administration of estrone sulfate. The second one was sham administered with the vehicle and supplemented with the same dosage of estrone sulfate. Ex 45 was orally administered at 5mg/kg/day for 5 weeks.

Xenografts were measured weekly by electronic Caliper and compared to the estrone sulfate group. (Table 5).

Table 5: Inhibition of human breast tumor

Parameters		Volume variations (expressed as % of E1S positive control group)			
Treatment	Time	Day 7	Day 28	Day 35	Day 42
E ₁ S + Ex 45		0	-26	-44	-39

5

When animals were treated with Ex 45, tumor volumes did not increased. On the other hand, simultaneously, groups only treated with estrone sulfate (E1S), tumor volumes were higher. At the end of the experiments, after five weeks of daily treatment, Ex 45 led to a 40% decrease in tumor volume, showing a clear and well established anticancerous potency.

10

JEG-3 xenograft into nude mice

The JEG-3 cell line over-expresses both human aromatase and estrone sulfatase enzymes. It is injected subcutaneously into ovariectomized athymic nude mice supplemented with a daily administration of subcutaneous estrone sulfate with or without test compounds (administered at 5mg/kg/day). In that particular case, and according to their semi-liquid status (choriocarcinoma origin), tumor measures are irrelevant. Nevertheless, an indirect effect of estradiol which is mainly synthesised by the tumour tissue, is obtained on uteri weights. On the other hand, because of the aromatase and sulfatase overexpression in such tumours, enzyme levels are measurable. Enzyme activities are determined according to the above-mentioned methods

15

20

JEG-3 xenograft within nude rat supplemented with estrone sulfate and Δ^4 -androstenedione

25


15 days before JEG-3 cell injection into rats, blood samples are taken to measure basal levels of estradiol plasma. Then, JEG-3 cells are injected subcutaneously in

ovariectomized athymic (Rnu/Rnu) nude rat. Animals are supplemented with a daily administration of subcutaneous estrone sulfate and $\Delta 4$ -androstenedione with or without tested compound (administered at 1mg/kg/day). After a 21-day period, blood sampling are performed one day after cancerous xenograft and at

5 the end of the experiment. In this experiment estrone sulfate and $\Delta 4$ -androstenedione are the precursors of estradiol. The effect of estradiol is reflected on the uterus weight after sacrifice. Plasma hormone levels are assayed at the end of the experiments according to the supplier's standard method (DSL, Webster, TX, USA).


The claims defining the invention are as follows:

1. An imidazole derivative of formula (I) :




or an acid addition salt or stereoisomeric form thereof, wherein :

- R_1 and R_2 are each independently hydrogen or a (C_1-C_6) alkyl;
- Q is selected from a direct link, $C(O)$, SO_2 , $CONH$, $C(O)(CH_2)_n$, $(CH_2)_n(O)$ or $(CH_2)_n$ in which n is 0, 1 or 2.
- Z is a group selected from:




- p is 1, 2, 3 or 4;
- q is 0, 1 or 2;
- the dotted line means that R_8 and/or R_9 can be on any position of the benzothiophene ring;
- one of R_3 and R_8 is hydroxy, cyano, (C_1-C_6) alkoxy or $OSO_2NR_{10}R_{11}$; and
- the other is hydrogen or a hydroxy, halogen, nitro, cyano, (C_1-C_6) alkoxy, $NR_{10}R_{11}$, $SO_2NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, $NR_{12}SO_2NR_{10}R_{11}$, $OSO_2NR_{10}SO_2NR_{11}R_{12}$ group;
- R_4 is hydrogen and R_9 is hydrogen, hydroxy, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, acyl, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, $NR_{12}SO_2NR_{10}R_{11}$, CO_2R_{10} group;
- when $Q-Z$ is



n is 0, 1 or 2 and p is 1, one of R_3 and R_8 is a hydroxy or $\text{OSO}_2\text{NR}_{10}\text{R}_{11}$;

- when p is 2, 3 or 4 the R_9 s are other than hydrogen and can be the same or different;

- R_{10} , R_{11} and R_{12} are each independently hydrogen or a $(\text{C}_1\text{-C}_6)$ alkyl;
- when Z is



and p is 1,

then R_8 and R_9 can also form together with the phenyl ring a benzoxathiazine dioxide, a dihydrobenzoxathiazine dioxide or a benzoxathiazole dioxide.

2. The derivative according to claim 1, or an acid addition salt or stereoisomeric form thereof, wherein:

- one of R_3 and R_8 is cyano; and
- the other is hydrogen or a hydroxy, halogen, nitro, $(\text{C}_1\text{-C}_6)$ alkoxy, $\text{NR}_{10}\text{R}_{11}$, $\text{SO}_2\text{NR}_{10}\text{R}_{11}$, $\text{OSO}_2\text{NR}_{10}\text{R}_{11}$, $\text{NR}_{12}\text{SO}_2\text{NR}_{10}\text{R}_{11}$ group.

3. The derivative according to claim 1, or an acid addition salt or stereoisomeric form thereof, wherein R_9 is hydrogen or a hydroxy, cyano, halogen, nitro, $(\text{C}_1\text{-C}_6)$ alkyl, $(\text{C}_1\text{-C}_6)$ alkoxy, trifluoromethyl, $\text{NR}_{10}\text{R}_{11}$, $\text{OSO}_2\text{NR}_{10}\text{R}_{11}$, CO_2R_{10} , CHO , $\text{NR}_{12}\text{SO}_2\text{NR}_{10}\text{R}_{11}$ group.

4. The derivative according to claim 3, or an acid addition salt or stereoisomeric form thereof, wherein R_9 is hydroxy, cyano, halogen, nitro, $(\text{C}_1\text{-C}_6)$ alkyl, $(\text{C}_1\text{-C}_6)$ alkoxy, trifluoromethyl, $\text{NR}_{10}\text{R}_{11}$, $\text{OSO}_2\text{NR}_{10}\text{R}_{11}$, CO_2R_{10} , CHO or $\text{NR}_{12}\text{SO}_2\text{NR}_{10}\text{R}_{11}$.


5. The derivative according to any one of claims 1 to 4, or an acid addition salt or stereoisomeric form thereof, in which:

- R_8 is hydrogen, hydroxy, halogen, nitro, cyano, $(\text{C}_1\text{-C}_6)$ alkoxy, $\text{NR}_{10}\text{R}_{11}$, $\text{SO}_2\text{NR}_{10}\text{R}_{11}$, $\text{OSO}_2\text{NR}_{10}\text{R}_{11}$, $\text{NR}_{12}\text{SO}_2\text{NR}_{10}\text{R}_{11}$ or $\text{OSO}_2\text{NR}_{10}\text{SO}_2\text{NR}_{11}\text{R}_{12}$ group;
- R_9 hydrogen or a hydroxy, cyano, halogen, nitro, $(\text{C}_1\text{-C}_6)$ alkyl, $(\text{C}_1\text{-C}_6)$ alkoxy, trifluoromethyl, $\text{NR}_{10}\text{R}_{11}$, $\text{OSO}_2\text{NR}_{10}\text{R}_{11}$, CO_2R_{10} , CHO , $\text{NR}_{12}\text{SO}_2\text{NR}_{10}\text{R}_{11}$ group;

- p and q are as defined in claim 1.

6. The derivative according to claim 1, or an acid addition salt or stereoisomeric form thereof, wherein:


- Z is



in which R_8 , R_9 and p are as defined in claim 1.

7. The derivative according to claim 6, or an acid addition salt or stereoisomeric form thereof, wherein:

- Z is



- Q is $(CH_2)_n$ in which n is 0, 1 or 2;
- R_8 is hydroxy, halogen, nitro, cyano or a (C_1-C_6) alkoxy, $NR_{10}R_{11}$, $SO_2NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$, or $NR_{12}SO_2NR_{10}R_{11}$ group;
- R_9 is hydrogen, hydroxy, cyano, halogen, nitro, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, trifluoromethyl, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$;
- p is as defined in claim 1.

8. The derivative according to any one of claims 6 to 7, or an acid addition salt or stereoisomeric form thereof, wherein:


- n is 0 or 1;
- R_1 , R_2 and R_4 are each hydrogen;
- R_9 is hydrogen, halogen, (C_1-C_6) alkyl or $OSO_2NR_{10}R_{11}$.

9. The derivative according to any one of claims 6 to 8, or an acid addition salt or stereoisomeric form thereof, wherein one of R_3 and R_8 is hydroxy, cyano or $OSO_2NR_{10}R_{11}$ and the other is hydroxy, nitro, $NR_{10}R_{11}$, $OSO_2NR_{10}R_{11}$ or $NR_{12}SO_2NR_{10}R_{11}$.

10. The derivative according to claim 9, or an acid addition salt or stereoisomeric form thereof, wherein one of R_3 and R_8 is cyano or $\text{OSO}_2\text{NR}_{10}\text{R}_{11}$ and the other is hydroxy or $\text{OSO}_2\text{NR}_{10}\text{R}_{11}$.

11. The derivative according to claim 1, or an acid addition salt or stereoisomeric form thereof, wherein:

- Z is



in which:

- R_8 , R_9 , p and q are as defined in claim 1;
- The dotted line means that Q and/or R_8 and/or R_9 can be on any position of the benzothiophene ring.

12. The derivative according to claim 11, or an acid addition salt or stereoisomeric form thereof, wherein R_8 is $\text{OSO}_2\text{NR}_{10}\text{R}_{11}$ or $\text{NR}_{12}\text{SO}_2\text{NR}_{10}\text{R}_{11}$.


13. The derivative according to claim 11 or 12, wherein R_9 is hydrogen, halogen, nitro, COOR_{10} or cyano.

14. The derivative according to any one of claims 11 to 13, wherein Q is SO_2 or CO.

15. The derivative according to any one of claims 11 to 14, wherein R_3 is hydrogen, halogen or cyano.

16. The derivative according to claim 1, or an acid addition salt or stereoisomeric form thereof, wherein:

- Z is a group:



in which R_8 , R_9 and p are as defined in claim 1.

17. The derivative according to any one of claims 1 to 16, or an acid addition salt or stereoisomeric form thereof, wherein R_{10} and R_{11} are hydrogen.

18. The derivative according to any one of claims 1 to 17 or a pharmaceutically acceptable salt thereof for use as an active therapeutic substance.

19. The pharmaceutical composition comprising a derivative according to any one of claims 1 to 17, or a pharmaceutically acceptable acid addition salt thereof, and a pharmaceutically acceptable carrier.

20. The pharmaceutical composition according to claim 19, comprising from 0.1 to 400 mg of said derivative.

21. Use of a derivative according to any one of claims 1 to 17 or a pharmaceutically acceptable acid addition salt thereof in the manufacture of a medicament for the treatment or prevention of hormone- or non hormone-dependent tumors, wherein said derivative is optionally combined with a sexual endocrine therapeutic agent.

22. Use of a derivative according to any one of claims 1 to 17 or a pharmaceutically acceptable acid addition salt thereof in the manufacture of a medicament for the control or management of reproductive functions, wherein said derivative is optionally combined with a LH-RH agonist or antagonist, an estroprogestative contraceptive, a progestin, an anti-progestin or a prostaglandin.

23. Use of a derivative according to any one of claims 1 to 17 or a pharmaceutically acceptable acid addition salt thereof in the manufacture of a medicament for the treatment or prevention of benign or malignant diseases of the breast, the uterus or the ovary, wherein said derivative is optionally combined with an antiandrogen, an anti-estrogen, a progestin or a LH-RH agonist or antagonist.

24. Use of a derivative according to any one of claims 1 to 17 or a pharmaceutically acceptable acid addition salt thereof in the manufacture of a medicament for the treatment or prevention of androgen-dependent diseases or benign or malignant diseases of the prostate or the testis, wherein said derivative is optionally combined with an antiandrogen, a progestin, a lyase inhibitor or a LH-RH agonist or antagonist.

25. Use of a derivative according to any one of claims 1 to 17 or a pharmaceutically acceptable acid addition salt thereof in the manufacture of a medicament for the treatment or prevention of cognitive function disorders, especially senile dementia, in particular Alzheimer's disease.

26. Use of a derivative according to any one of claims 1 to 17 or a pharmaceutically acceptable acid addition salt thereof in the manufacture of a medicament for the treatment or prevention of immunodisorders.

27. Use of a derivative according to any one of claims 1 to 17 or a pharmaceutically acceptable acid addition salt thereof in the manufacture of a medicament for the treatment or prevention of pathologies in which inhibition of aromatase and/or steroid sulfatase and/or carbonic anhydrase is required.

28. A method of treating a disease in which aromatase and/or steroid sulfatase and/or carbonic anhydrase is involved, which comprises administering to a subject in need thereof a therapeutically effective amount of a compound according to any one of claims 1 to 17.

29. An imidazole derivative according to any one of claims 1 to 10 and 17, which is selected among the following compounds:

4-[N-(1H-imidazol-1-yl)-N-(4-methoxyphenyl)amino]methylbenzonitrile;

4-[N-(4-hydroxyphenyl)-N-(1H-imidazol-1-yl)amino]methylbenzonitrile;

4-[N-(4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile;

4-[N-(3-chloro-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile;

4-[N-(3-bromo-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile;

4-[N-(4-hydroxy-3-methoxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]

benzonitrile;

4-[N-(2,3,5,6-tetrafluoro-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino] benzonitrile;

4-[N-(3-formyl-4-hydroxyphenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile;

4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino}methyl}benzene sulfonamide;

4-[N-(4-hydroxy-3-nitrophenylmethyl)-N-(1H-imidazol-1-yl)amino]benzonitrile;

4-[N-(1H-imidazol-1-yl)-N-(4-nitrophenyl)amino]benzonitrile;

N-(1H-imidazol-1-yl)-N-(4-cyanophenyl)-2-(4-fluorophenyl)acetamide;

N-(1H-imidazol-1-yl)-N-(4-cyanophenyl)-2-(4-hydroxyphenyl)acetamide;

N-(4-cyanophenyl)-3-(4-hydroxyphenyl)-N-(1H-imidazol-1-yl)propanamide;

N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-4-(phenylmethoxy)-benzensulfonamide;

4-[N-(3-amino-4-hydroxy-phenylmethyl)-N-(1H-imidazol-1-yl)amino]

benzonitrile;

4-{{N-[2-(4-hydroxyphenoxy)ethyl]-N-(1H-imidazol-1-yl)amino}benzonitrile;

N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-4-hydroxybenzensulfonamide;

- 4-[N-(4-aminophenyl)-N-(1H-imidazol-1-yl)amino]benzonitrile;
 Sulfamic acid 4-[N-(4-cyanophenylmethyl)-N-(1H-imidazol-1-yl)amino]phenyl
 ester;
 Sulfamic acid-4- {[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}phenyl
 5 ester;
 Sulfamic acid 2-chloro-4- {[N-(4-cyanophenyl)-N-(1H-imidazol-
 1-yl)amino]methyl}phenyl ester;
 Sulfamic acid 2-bromo-4- {[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]
 methyl}phenyl ester, hydrochloride;
 10 Sulfamic acid 2-methoxy-4- {[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]
 methyl}phenyl ester;
 Sulfamic acid 2,3,5,6-tetrafluoro-4- {[N-(4-cyanophenyl)-N-(1H-imidazol-
 1-yl)amino] methyl}phenyl ester;
 4-[N-[(2,2-dioxido-1,2,3-benzoxathiazin-6-yl)methyl]-N-(1H-imidazol-1-yl)amino]
 15 benzonitrile;
 N- {4-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]phenyl}sulfamide;
 Sulfamic acid 4- {[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]sulfonyl}phenyl
 ester, hydrochloride;
 Sulfamic acid 4- {2-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]ethoxy}phenyl
 20 ester;
 Sulfamic acid 4- {[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-carbamoyl]-methyl}
 phenyl ester;
 Sulfamic acid 4- {[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]-3-oxopropyl}
 phenyl ester;
 25 Sulfamic acid 3-(aminosulfonyl)amino-4- {[N-(4-cyanophenyl)-N-(1H-imidazol-
 1-yl)amino]methyl}phenyl ester;
 2-Bromo-4- {[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}phenyl
 amidimidodisulfate acid;
 4-[N-[(2,2-dioxido-3,4-dihydro-1,2,3-benzoxathiazin-6-yl)methyl]-N-(1H-
 30 imidazol-1-yl)amino]benzonitrile;
 5- {[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-2-hydroxybenzoic
 acid;
 4-[N-(1H-imidazol-1-yl)-N-(phenyl)amino]benzonitrile
 4-[N-(3-tosylamino-4-hydroxy-benzyl)-N-(1H-imidazol-1-yl)amino]benzonitrile;

4-[N-[(2,2-dioxido-3-tosyl-3H-1,2,3-benzoxathiazol-5-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile;

4-[N-[(2,2-dioxido-3H-1,2,3-benzoxathiazol-5-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile;

5 N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-N'-phenylurea.

30. An imidazole derivative according to claim 29, which is selected among the following compounds:

Sulfamic acid 4-[N-(4-cyanophenylmethyl)-N-(1H-imidazol-1-yl)amino]phenyl ester;

10 Sulfamic acid-4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino}methyl}phenyl ester;

Sulfamic acid 2-chloro-4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino}methyl}phenyl ester;

15 Sulfamic acid 2-bromo-4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino}methyl}phenyl ester, hydrochloride;

Sulfamic acid 2-methoxy-4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino}methyl}phenyl ester;

Sulfamic acid 2,3,5,6-tetrafluoro-4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino}methyl}phenyl ester;

20 4-[N-[(2,2-dioxido-1,2,3-benzoxathiazin-6-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile;

Sulfamic acid 4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino}sulfonyl}phenyl ester, hydrochloride;

25 Sulfamic acid 4-{2-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]ethoxy}phenyl ester;

Sulfamic acid 4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-carbamoyl}-methyl}phenyl ester;

Sulfamic acid 4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino}-3-oxopropyl}phenyl ester;

30 Sulfamic acid 3-(aminosulfonyl)amino-4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino}methyl}phenyl ester.

31. An imidazole derivative according to claim 30, which is sulfamic acid 2-bromo-4-{{N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino}methyl}phenyl ester, hydrochloride.

32. An imidazole derivative according to any one of claims 1 to 5, 11 to 15 and 17, which is selected among the following compounds:

5-Nitro-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)]-benzothiophene-2-carboxamide;

N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-2-(6-methoxy-benzothien-3-yl)acetamide;

4-{N-[1H-imidazol-1-yl]-N-[(6-methoxy-benzothien-3-yl)methyl]amino}benzonitrile;

4-[N-(6-benzyloxy-1,1-dioxido-benzothien-3-yl)-N-(1H-imidazol-1-yl)amino]benzonitrile;

4-[N-[(6-benzyloxy-benzothien-2-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile;

4-{N-[(6-hydroxy-benzothien-3-yl)methyl]-N-[1H-imidazol-1-yl]amino}benzonitrile;

N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)-2-(6-hydroxy-benzothien-3-yl)acetamide;

4-[N-[(6-hydroxy-1,1-dioxido-benzothien-3-yl)]-N-(1H-imidazol-1-yl)amino]benzonitrile;

4-[N-[(6-hydroxy-benzothien-2-yl)methyl]-N-(1H-imidazol-1-yl)amino]benzonitrile;

5-Amino-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)]-benzothiophene-2-carboxamide;

5-(Aminosulfonyl)amino-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)]-benzothiophene-2-carboxamide;

Sulfamic acid 3-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]-1,1-dioxido-benzothien-6-yl ester;

Sulfamic acid 3-{2-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]-2-oxoethyl}-benzothien-6-yl ester;

Sulfamic acid 3-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-benzothien-6-yl ester;

Sulfamic acid 2-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-benzothien-6-yl ester.

33. An imidazole derivative according to claim 32, which is selected among the following compounds:

Sulfamic acid 3-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]-1,1-dioxido-benzothien-6-yl ester;

5 Sulfamic acid 3-{2-[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]-2-oxoethyl}-benzothien-6-yl ester;

Sulfamic acid 3-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-benzothien-6-yl ester;

10 Sulfamic acid 2-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-benzothien-6-yl ester;

34. An imidazole derivative according to any one of claims 1 to 5, 16 and 17, which is selected among the following compounds:

4-{N-[(6-chloropyridin-3-yl)methyl]-N-(1H-imidazol-1-yl)amino}benzonitrile;

6-chloro-N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)nicotinamide.

15 35. An imidazole derivative which is 5-{[N-(4-cyanophenyl)-N-(1H-imidazol-1-yl)amino]methyl}-2-methoxybenzoic acid.

Dated 16 May, 2011

Laboratoire Theramex

Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON

