



(51) International Patent Classification:

A61K 39/245 (2006.01) *C07K 14/05* (2006.01)
A61K 39/395 (2006.01) *C07K 16/08* (2006.01)

(21) International Application Number:

PCT/US2018/030030

(22) International Filing Date:

27 April 2018 (27.04.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/491,945 28 April 2017 (28.04.2017) US

(71) Applicants: **CITY OF HOPE** [US/US]; 1500 East Duarte Road, Duarte, California 91010-3000 (US). **CALIFORNIA INSTITUTE OF TECHNOLOGY** [US/US]; 1200 E. California Blvd., Pasadena, California 91125 (US).


(72) Inventors: **OGEMBO, Javier Gordon**; 1767 La Mesa Oaks Drive, San Dimas, California 91773 (US). **MUTSVUNGUMA, Lorraine Zvichapera**; 1260 Duarte Road, Duarte, California 91010 (US). **ONDRUS, Alison Evelyn**; 1318 Cordova Street, Pasadena, California 91106 (US).

(74) Agent: **WANG, Yingli**; Perkins Coie LLP, P.O. Box 1247, Seattle, Washington 98111-1247 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,

(54) Title: EPSTEIN-BARR VIRUS ANTIBODIES AND USES THEREOF

Figure 4



(57) **Abstract:** Disclosed herein are antibodies or immunogenic fragments thereof that specifically bind to Epstein-Barr virus (EBV) glycoprotein 350 (gp350) or 220 or one or more epitopes of EBV gp350 and neutralize EBV infection. Also disclosed are immunogenic peptides comprising one or more gp350 epitopes, EBV antibody-small molecule conjugates and pharmaceutical compositions comprising the antibody or an immunogenic fragment thereof, one or more epitopes of EBV gp350, one or more immunogenic peptides, or the EBV antibody-small molecule conjugate. The antibodies, epitopes, immunogenic peptides, conjugates, and pharmaceutical compositions can be used to treat or prevent EBV infections and EBV-associated conditions and diseases.



SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*

EPSTEIN-BARR VIRUS ANTIBODIES AND USES THEREOF

PRIORITY CLAIM

[0001] The present invention claims the benefit of U.S. Provisional Patent Application No. 62/491,945, entitled "EPSTEIN-BARR VIRUS ANTIBODIES AND USES THEREOF," filed April 28, 2017, the content of which is incorporated herein by reference in its entirety.

STATEMENT OF GOVERNMENT INTEREST

[0002] The present invention was made with government support under Grant No. R21CA205106, awarded by U.S. Public Health Service. The Government has certain rights in the invention.

BACKGROUND

[0003] Epstein-Barr virus (EBV) infection is the causal agent of acute infectious mononucleosis (62, 63). Persistent EBV infection in immunodeficient individuals is associated with numerous epithelial and lymphoid malignancies, such as nasopharyngeal carcinoma, gastric carcinoma, Burkitt lymphoma, Hodgkin lymphoma, and post-transplant lymphoproliferative diseases (PTLD) (1). Transplantation is the treatment of choice for a variety of patients with end-stage organ failure or hematologic malignancies, or in need of reconstructive transplantation (1). Transplantation success depends entirely on potent immunosuppressive drugs to prevent stem cell/organ rejection. However, these drugs impose several serious side effects, including an increased risk of infection with or reactivation of Epstein-Barr virus (EBV), and the resultant development of PTLDs, which are aggressive, life-threatening complications (2, 3). Through the early 2000s, PTLD patients who had been EBV-naïve prior to transplantation showed mortality rates of 50–90% for stem cell and solid organ transplants; while recent data suggest outcomes have improved, challenges remain. PTLDs usually develop in EBV-naïve patients, particularly pediatric patients, who receive organs from EBV+ donors. A variety of non-standardized, non-specific treatments are used to treat EBV+ PTLD cases (4-9). Initial clinical management typically involves reduction of immunosuppression; however, this can lead to graft-versus-host disease. Other treatments including radiation/chemotherapy and excision of PTLD lesions all have undesirable side effects. Second-line treatment often includes antibodies (Abs)

against the B cell antigen, CD20; however, this also targets healthy B cells, further weakening the immune system and exposing patients to other opportunistic infections.

[0004] In over 50 years of EBV vaccine research, few candidates have demonstrated partial clinical efficacy, and none have been efficacious enough to elicit sterilizing immunity and be licensed (24). Antibodies, whether elicited in the host naturally or via passive immunization, provide an effective first-line of defense against viral infection.

[0005] Thus, there is an urgent need for a novel EBV-specific therapy that targets EBV+ cells to neutralize EBV infection and prevent subsequent PTLD development in EBV-naïve patients.

SUMMARY

[0006] In one aspect, this disclosure relates to an Epstein-Barr virus (EBV) antibody or an immunogenic fragment thereof. In some embodiments, the EBV antibody or an immunogenic fragment thereof specifically binds to EBV glycoprotein 350/220. In some embodiments, the EBV antibody or an immunogenic fragment thereof specifically binds to EBV glycoprotein 350 or one or more epitopes represented by SEQ ID NOs: 1-3. In some embodiments, the EBV antibody comprises a VH region comprising CDR-1, CDR-2, and CDR-3 represented by SEQ ID NOs: 4-19, 20-35, and 36-51, respectively. In some embodiments, the EBV antibody comprises a VL region comprising CDR-1, CDR-2, and CDR-3 represented by SEQ ID NOs: 52-67, 68-83, and 84-99, respectively. In some embodiments, the antibody is not antibody clone 72A1 or an antibody comprising the CDRs of antibody clone 72A1. In some embodiments, the EBV antibody is a monoclonal antibody. In some embodiments, the EBV antibody is a chimeric antibody, a human antibody, or a humanized antibody.

[0007] In another aspect, this disclosure relates to an immunogenic peptide comprising one or more gp350 epitopes having an amino acid sequence identical to or sharing at least 60% similarity to SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3. In some embodiments, the immunogenic peptide comprises a first domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO:1, a second domain comprising an amino acid sequence having at least 60% similarity

to SEQ ID NO: 2, and a third domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO: 3. In some embodiments, the immunogenic peptide comprises a first domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3, and a second domain comprising a known immunogenic peptide such as keyhole limpet hemocyanin (KLH) peptide.

[0008] In another aspect, this disclosure relates to an EBV antibody-small molecule conjugate. The EBV antibodies disclosed herein can be conjugated to small molecules having activities against EBV-transformed cells. For example, the small molecules have anti-proliferative activities against EBV-transformed B lymphoma cells. In some embodiments, the small molecules are growth inhibitors of EBV infected B cells. In some embodiments, the small molecule is L₂P₄, 2-butyne diamide, or a derivative thereof. In some embodiments, the small molecule is conjugated to the antibody via a linker or an adaptor. In some embodiments, the small molecule is conjugated to the constant region of the heavy chain or the light chain of the antibody.

[0009] In a related aspect, this disclosure relates to a pharmaceutical composition comprising the EBV antibody disclosed herein or an immunogenic fragment thereof, one or more epitopes represented by SEQ ID NOs: 1-3, one or more immunogenic peptides disclosed herein, or the EBV antibody-small molecule conjugate disclosed herein. The pharmaceutical composition can further comprise one or more pharmaceutically acceptable excipients. The pharmaceutical composition can be formulated into any suitable formulation depending on the administration route.

[0010] In another aspect, this disclosure relates to a method of neutralizing EBV infection. The method includes administering to a subject infected with EBV a therapeutically effective amount of the EBV antibody disclosed herein or an immunogenic fragment thereof, one or more epitopes represented by SEQ ID NOs: 1-3, one or more immunogenic peptides disclosed herein, the EBV antibody-small molecule conjugate, or the pharmaceutical composition described above. In some embodiments, the subject is human.

[0011] In another aspect, this disclosure relates to a method of preventing EBV infection. The method includes administering to a subject at an elevated risk of EBV infection a therapeutically effective amount of the EBV antibody disclosed herein or an immunogenic fragment thereof, one or more epitopes represented by SEQ ID NOs: 1-3, one or more immunogenic peptides disclosed herein, the EBV antibody-small molecule conjugate, or the pharmaceutical composition described above. In some embodiments, the subject is human.

[0012] In another aspect, this disclosure relates to a method of preventing a post-transplant lymphoproliferative disease (PTLD). PTLD is associated with EBV infection of B cells, either as a consequence of reactivation of the virus post transplantation or from primary EBV infection. The method includes administering to a subject who is a transplant recipient a therapeutically effective amount of the EBV antibody disclosed herein or an immunogenic fragment thereof, one or more epitopes represented by SEQ ID NOs: 1-3, one or more immunogenic peptides disclosed herein, the EBV antibody-small molecule conjugate, or the pharmaceutical composition described above. The administration can be before, during, and/or after the transplant. In some embodiments, the subject is a pediatric transplant recipient who is EBV naïve. In some embodiments, the subject is an adult transplant recipient. In some embodiments, the subject is human.

[0013] In another aspect, this disclosure relates to a method of treating an EBV-associated cancer. The method includes administering to a subject suffering from an EBV-associated cancer a therapeutically effective amount of the EBV antibody disclosed herein or an immunogenic fragment thereof, one or more epitopes represented by SEQ ID NOs: 1-3, one or more immunogenic peptides disclosed herein, the EBV antibody-small molecule conjugate, or the pharmaceutical composition described above. In some embodiments, the examples of EBV-associated cancer include but are not limited to Hodgkin lymphoma, Burkitt lymphoma, gastric cancer, and nasopharyngeal carcinoma. In some embodiments, the subject is human.

[0014] In another aspect, this disclosure relates to a method of immunizing or vaccinating a subject against an EBV infection. The method includes administering to a subject suffering from an EBV infection a therapeutically effective amount of the EBV antibody disclosed herein or an immunogenic fragment thereof, one or more

epitopes represented by SEQ ID NOs: 1-3, one or more immunogenic peptides disclosed herein, or the pharmaceutical composition thereof as described above. In some embodiments, the subject is human.

[0015] In another aspect, this disclosure relates to a method of inducing the production of neutralizing antibodies against a EBV in a subject. The method includes administering to a subject an effective amount of a gp350 epitope represented by SEQ ID NO: 3 or an immunogenic peptide comprising the gp350 epitope represented by SEQ ID NO: 3. In some embodiments, the subject is human.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] Figures 1A-1D show characterization of novel anti-gp350 mAbs. Figure 1A: ELISA screening of hybridoma (HB) supernatants for anti-gp350-specific antibodies. Soluble EBV gp350 protein was used as the target antigen at 0.5 µg/ml. nAb-72A1 at 10 µg/ml and 1x phosphate buffered saline (PBS) were used as positive and negative (not shown) controls, respectively. Bound antibodies were detected using HRP-conjugated anti-mouse IgG (1:2,000). Twenty-three HB clones with ELISA signals two times greater than those of PBS control were considered as positive hybridomas. Figure 1B: Determining specificity of anti-gp350 producing hybridoma supernatant by immunoblotting with gp350-transfected stable CHO lysate. Western blot analysis was conducted on untransfected and pCAGGS-gp350 transfected CHO lysate. Anti-gp350 mAb 72A1 was used as a positive control (1:100) and anti-gp350 hybridoma clone supernatants were used at 1:50. Anti-mouse secondary antibody was used at 1:2000. Figure 1C: Flow cytometric analysis of surface expression of gp350 protein on gp350 expressing CHO cells for all 23 HB clones. 10⁶ CHO cells were transfected with 1 µg of pCAGGS-gp350. gp350 expressing cells were resuspended in PBS, stained with anti-gp350 mAb (1:250), which detects the gp350 ED, followed by secondary goat anti-mouse conjugated to AF488. Figure 1D: Flow cytometric analysis of HB5, HB17 and HB19.

[0017] Figures 2A and 2B show PROMALS3D multiple sequence alignment of VH (Figure 2A) and VL (Figure 2B) regions of 15 mAbs and nAb-72A1. The highly variable complementarity determining regions (CDR) 1-3, indicated by black boxes, define the antigen binding specificity. The conserved framework regions (FR) 1-4 flank the CDRs. Consensus amino acid (aa) are in bold and upper case.

Consensus predicted secondary structure (ss) symbols: alpha-helix: h and beta-strand: e.

[0018] Figures 3A and 3B show neutralization activity of novel anti-gp350 mAbs against EBV-eGFP in Raji cells. EBV-eGFP was pre-incubated with 15 anti-gp350 mAbs at 10 µg/ml (Figure 3A) and 50 µg/ml (Figure 3B), followed by incubation with Raji cells for 48 h. EBV-eGFP+ cells were enumerated using flow cytometry. Anti-gp350 (nAb-72A1) and anti-gH/gL (E1D1) mAbs served as positive controls and non-neutralizing anti-gp350 (2L10) mAb served as negative control.

[0019] Figures 4A-4C show construction of chimeric gp350 nAbs. Figure 4A: Construction of chimeric Ab. Figure 4B: Cloning strategy of heavy chain and light chain variable regions into expression vectors. Figure 4C: PCR amplification of heavy chain and light chain variable regions. 72A1 and clone 19 were used as examples of PCR amplification.

[0020] Figure 5 illustrates an antibody-L2P4 conjugate.

DETAILED DESCRIPTION

[0021] Disclosed herein are EBV antibodies or immunogenic fragments thereof that specifically bind to gp350/gp220 or an epitope of gp350, immunogenic peptides comprising one or more gp350 epitopes, or one or more amino acid sequences having at least 60% similarity to one or more gp350 epitopes, EBV antibody-small molecule conjugates for treating or preventing EBV infection, in particular, in subjects receiving a transplant. In some embodiments, chimeric (human/mouse) anti-gp350 nAbs are conjugated to L₂P₄ to obtain an EBV-specific ADC that improves the therapeutic efficacy of treating EBV-associated PTLDs. L₂P₄ described by Jiang et al., *Nature Biomedical Engineering* 1: 0042 (2017), is an example of the small molecules encompassed by this disclosure.

[0022] The term “antibody” as used herein refers to an immunoglobulin molecule or an immunologically active portion thereof that specifically binds to, or is immunologically reactive with a particular antigen, for example, EBV gp350/gp220, or a particular domain or fragment of gp350/gp220. In some embodiments, the antibody disclosed herein specifically binds to one or more epitopes on EBV gp350. The epitopes can be neutralizing epitopes or immunodominant epitopes. In some embodiments, the epitope has an amino acid sequence selected from

²⁵³TIPIGTGYAYSLRLTPRPVSRFL²⁷⁵ (SEQ ID NO: 1),
⁸⁷⁵LLLLVMADCAFRRLSTSHTYTTTPPY⁸⁹⁹ (SEQ ID NO: 2), and
³⁸¹GAFASNRTFDIT³⁹² (SEQ ID NO: 3).

[0023] In a related aspect, this disclosure relates to a method of producing an EBV antibody using the epitopes disclosed herein according to any known technology. The method entails immunizing an animal such as a mouse or a rabbit with an immunogenic peptide disclosed herein or the epitope represented by SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3 alone, and screening for and isolating a hybridoma producing an EBV antibody. The EBV antibody produced from such hybridomas can be used for treating or preventing EBV infections.

[0024] In certain embodiments an antibody for use in the present methods or compositions is a full-length immunoglobulin molecule, which comprises two heavy chains and two light chains, with each heavy and light chain containing three complementary determining regions (CDRs). The CDRs of various antibodies are identified and listed in Table 1 below.

Table 1: CDR Sequences						
Antibodies	VH			VL		
	CDR-1	CDR-2	CDR-3	CDR-1	CDR-2	CDR-3
72A1	GSSFTDY (SEQ ID NO: 4)	INPYNGG (SEQ ID NO: 20)	GGLRRVNWFAIW (SEQ ID NO: 36)	TGAVTTSNY (SEQ ID NO: 52)	GTN (SEQ ID NO: 68)	VLWHSNHWV (SEQ ID NO: 84)
HB-1	GFLTTY (SEQ ID NO: 5)	IWAGGS (SEQ ID NO: 21)	RDRGYGYLYAMDYW (SEQ ID NO: 37)	QNVGTN (SEQ ID NO: 53)	STD (SEQ ID NO: 69)	QQYNTYPYT (SEQ ID NO: 85)
HB-2	GYTFTAY (SEQ ID NO: 6)	INYKTGE (SEQ ID NO: 22)	PYGYALDYW (SEQ ID NO: 38)	SSVNY* (SEQ ID NO: 54)	ATS* (SEQ ID NO: 70)	QQWSSNPPT* (SEQ ID NO: 86)
HB-3	GYTFASY (SEQ ID NO: 7)	INPNNGH* (SEQ ID NO: 23)	RNLYYYGRPDYW* (SEQ ID NO: 39)	QDIGNY* (SEQ ID NO: 55)	YTS* (SEQ ID NO: 71)	QQGNTLPPT* (SEQ ID NO: 87)
HB-5	GYTFTNH (SEQ ID NO: 8)	INPYNDY (SEQ ID NO: 24)	RSEGWLRRGAWFAY (SEQ ID NO: 40)	QSIGTS (SEQ ID NO: 56)	YAS (SEQ ID NO: 72)	QQSNSWPMLT (SEQ ID NO: 88)
HB-6	GYTFTDY* (SEQ ID NO: 9)	INTRTGE (SEQ ID NO: 25)	PYGYALDYW (SEQ ID NO: 41)	SSVNY* (SEQ ID NO: 57)	ATS* (SEQ ID NO: 73)	QQWSSNPPT* (SEQ ID NO: 89)
HB-7	GYTFTDY* (SEQ ID NO: 10)	ISPGRSG (SEQ ID NO: 26)	RYGHPSYLDVW (SEQ ID NO: 42)	QSVGNA (SEQ ID NO: 58)	SAS (SEQ ID NO: 74)	QQYSSYPLT (SEQ ID NO: 90)
HB-8	GYSFTNY* (SEQ ID NO: 11)	INTYTGE (SEQ ID NO: 27)	RYYYGSVYSAWFAYW (SEQ ID NO: 43)	QSIVHSNGNTY* (SEQ ID NO: 59)	KVS* (SEQ ID NO: 75)	FQGSHVPYT* (SEQ ID NO: 91)
HB-9	GFTFSSY (SEQ ID NO: 12)	ISSGGSY (SEQ ID NO: 28)	REDFYGGSSYGFFDVW (SEQ ID NO: 44)	QSIVHSNGNTY* (SEQ ID NO: 60)	KVS* (SEQ ID NO: 76)	FQGSHVPYT* (SEQ ID NO: 92)
HB-10	GYTFTSY* (SEQ ID NO: 13)	INPSNGH (SEQ ID NO: 29)	RNLYYYGRPDYW (SEQ ID NO: 45)	QDIGNY* (SEQ ID NO: 61)	YTS* (SEQ ID NO: 77)	QQNTLPPT (SEQ ID NO: 93)
HB-11	GDSITSG (SEQ ID NO: 14)	ISYSGS (SEQ ID NO: 30)	RGNGGNYDWYFDVW (SEQ ID NO: 46)	SSVNF (SEQ ID NO: 62)	YIS (SEQ ID NO: 78)	QQFTSSPSWT (SEQ ID NO: 94)
HB-12	GYTFTNY* (SEQ ID NO: 15)	INPNNGH* (SEQ ID NO: 31)	RNLYYYGRPDYW* (SEQ ID NO: 47)	QSLVHSNGNTY (SEQ ID NO: 63)	KVS* (SEQ ID NO: 79)	SQSTHVPLT (SEQ ID NO: 95)
HB-14	GYTFTDY* (SEQ ID NO: 16)	IHPRRGG (SEQ ID NO: 32)	RYGYPWYFDVW (SEQ ID NO: 48)	QSIVHDNGNTY (SEQ ID NO: 64)	KVS* (SEQ ID NO: 80)	FQGSHVPPT (SEQ ID NO: 96)
HB-17	GYTFTSY* (SEQ ID NO: 17)	INPNNGH* (SEQ ID NO: 33)	RNLFYYSRPDYW (SEQ ID NO: 49)	QDIGNY* (SEQ ID NO: 65)	YTS* (SEQ ID NO: 81)	QQGNTLPPT* (SEQ ID NO: 97)
HB-20	GYTFTSY* (SEQ ID NO: 18)	INPTNGH (SEQ ID NO: 34)	RNLYYYGRPDYW* (SEQ ID NO: 50)	QDIGNY* (SEQ ID NO: 66)	YTS* (SEQ ID NO: 82)	QQGNALPPT (SEQ ID NO: 98)
HB-22	GFSLTNY (SEQ ID NO: 19)	IWSDGS (SEQ ID NO: 35)	RNYGNSYPAWFAYW (SEQ ID NO: 51)	QSIVHSNGNTY (SEQ ID NO: 67)	KVS* (SEQ ID NO: 83)	FQGSHVPWT (SEQ ID NO: 99)

[0025] The term “antibody,” in addition to natural antibodies, also includes genetically engineered or otherwise modified forms of immunoglobulins, such as synthetic antibodies, intrabodies, chimeric antibodies, fully human antibodies, humanized antibodies, peptibodies and heteroconjugate antibodies (e.g., bispecific antibodies, multispecific antibodies, dual-specific antibodies, anti-idiotypic antibodies, diabodies, triabodies, and tetrabodies). The antibodies disclosed herein can be monoclonal antibodies or polyclonal antibodies. In those embodiments wherein an antibody is an immunologically active portion of an immunoglobulin molecule, the antibody may be, for example, a Fab, Fab', Fv, Fab' F(ab')₂, disulfide-linked Fv, single chain Fv antibody (scFv), single domain antibody (dAb), or diabody. The antibodies disclosed herein, including those that are immunologically active portion

of an immunoglobulin molecule, retain the ability to bind a specific antigen, for example, EBV gp350 or one or more epitopes thereof, EBV gp220 or one or more epitopes thereof, or to bind a specific fragment of gp350/gp220.

[0026] In some embodiments, the EBV antibodies disclosed herein have undergone post-translational modifications such as phosphorylation, methylation, acetylation, ubiquitination, nitrosylation, glycosylation, or lipidation associated with expression in a mammalian cell line, including a human or a non-human host cell. Techniques for producing recombinant antibodies and for *in vitro* and *in vivo* modifications of recombinant antibodies are known in the art.

[0027] Provided in certain embodiments herein are chimeric, and/or humanized EBV antibodies. Various techniques are known in the art for humanizing antibodies from non-human species such that the antibodies are modified to increase their similarity to antibodies naturally occurring in humans. Six CDRs are present in each antigen binding domain of a natural antibody. These CDRs are short, non-contiguous sequences of amino acids that are specifically positioned to form the antigen binding domain as the antibody assumes its three dimensional configuration. CDR sequences of certain antibodies identified herein are shown in Table 1. The remainder of the amino acids in the antigen binding domains, referred to as “framework” regions, show less inter-molecular variability and form a scaffold to allow correct positioning of the CDRs. This disclosure also relates to antibodies comprising VH and VL regions comprising the CDRs shown in Table 1.

[0028] “Treating” or “treatment” of a disease or a condition may refer to preventing the disease or condition, slowing the onset or rate of development of the disease or condition, reducing the risk of developing the disease or condition, preventing or delaying the development of symptoms associated with the disease or condition, reducing or ending symptoms associated with the disease or condition, generating a complete or partial regression of the disease or condition, or some combinations thereof.

[0029] As used herein, the term “subject” refers to mammalian subject, preferably a human. The phrases “subject” and “patient” are used interchangeably herein.

[0030] The method for treating a condition or a viral infection includes administering a therapeutically effective amount of a therapeutic agent or a pharmaceutical composition. An “effective amount,” “therapeutically effective amount” or “effective dose” is an amount of a composition (e.g., a therapeutic agent or a pharmaceutical composition) that produces a desired therapeutic effect in a subject, such as preventing or treating a target disease or condition or alleviating symptoms associated with the disease or condition. The precise therapeutically effective amount is an amount of the composition that will yield the most effective results in terms of efficacy of treatment in a given subject. This amount will vary depending upon a variety of factors, including but not limited to the characteristics of the therapeutic agent (including activity, pharmacokinetics, pharmacodynamics, and bioavailability), the physiological condition of the subject (including age, sex, disease type and stage, general physical condition, responsiveness to a given dosage, and type of medication), the nature of the pharmaceutically acceptable carrier or carriers in the formulation, and the route of administration. One skilled in the clinical and pharmacological arts will be able to determine a therapeutically effective amount through routine experimentation, namely by monitoring a subject’s response to administration of a compound and adjusting the dosage accordingly. For additional guidance, see Remington: The Science and Practice of Pharmacy 21st Edition, Univ. of Sciences in Philadelphia (USIP), Lippincott Williams & Wilkins, Philadelphia, PA, 2005.

[0031] The pharmaceutical composition may include, among other things, an EBV antibody disclosed herein or an immunogenic fragment thereof, one or more gp350 epitopes represented by SEQ ID NOs: 1-3, one or more immunogenic peptides disclosed herein, or an EBV antibody-small molecule conjugate disclosed herein.

[0032] The pharmaceutical composition may also include one or more pharmaceutically acceptable carriers. A “pharmaceutically acceptable carrier” refers to a pharmaceutically acceptable material, composition, or vehicle that is involved in carrying or transporting a compound of interest from one tissue, organ, or portion of the body to another tissue, organ, or portion of the body. For example, the carrier may be a liquid or solid filler, diluent, excipient, solvent, or encapsulating material, or some combination thereof. Each component of the carrier must be

“pharmaceutically acceptable” in that it must be compatible with the other ingredients of the formulation. It also must be suitable for contact with any tissue, organ, or portion of the body that it may encounter, meaning that it must not carry a risk of toxicity, irritation, allergic response, immunogenicity, or any other complication that excessively outweighs its therapeutic benefits.

[0033] The pharmaceutical compositions described herein may be administered by any suitable route of administration. A route of administration may refer to any administration pathway known in the art, including but not limited to aerosol, enteral, nasal, ophthalmic, oral, parenteral, rectal, transdermal (e.g., topical cream or ointment, patch), or vaginal. “Transdermal” administration may be accomplished using a topical cream or ointment or by means of a transdermal patch. “Parenteral” refers to a route of administration that is generally associated with injection, including infraorbital, infusion, intraarterial, intracapsular, intracardiac, intradermal, intramuscular, intraperitoneal, intrapulmonary, intraspinal, intrasternal, intrathecal, intrauterine, intravenous, subarachnoid, subcapsular, subcutaneous, transmucosal, or transtracheal. In some embodiments, the therapeutic compositions described herein are administered by intravenous injection or intraperitoneal injection.

[0034] The epitopes disclosed herein have various uses. For example, the epitopes can be used to produce immunogenic peptides. Such an immunogenic peptide comprises one or more gp350 epitopes having an amino acid sequence identical to or sharing at least 60% similarity to SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3. In some embodiments, the immunogenic peptide comprises a first domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO:1, a second domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO: 2, and a third domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO: 3. In some embodiments, the immunogenic peptide comprises a first domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3, and a second domain comprising a known immunogenic peptide such as keyhole limpet hemocyanin (KLH) peptide.

[0035] Additionally, the epitopes or immunogenic peptides disclosed herein can be used for producing anti-gp350 antibodies using any existing technology.

[0036] In certain embodiments, disclosed herein is a method of treating or preventing EBV infection in a subject, comprising administering a therapeutically effective amount of an anti-gp350 antibody disclosed herein or an immunogenic fragment thereof, one or more epitopes represented by SEQ ID NOs: 1-3, an immunogenic peptide described herein, an EBV antibody-drug conjugate described herein, or a pharmaceutical composition comprising the anti-gp350 antibody or an immunogenic fragment thereof, one or more epitopes, one or more immunogenic peptides, or the EBV antibody-drug conjugate.

[0037] In certain embodiments, disclosed herein is a method of treating or preventing EBV infection in a subject, comprising administering a therapeutically effective amount of an anti-gp350 antibody disclosed herein or an immunogenic fragment thereof, one or more epitopes represented by SEQ ID NOs: 1-3, an immunogenic peptide described herein, an EBV antibody-drug conjugate described herein, or a pharmaceutical composition comprising the anti-gp350 antibody or an immunogenic fragment thereof, one or more epitopes, one or more immunogenic peptides, or the EBV antibody-drug conjugate.

[0038] In certain embodiments, disclosed herein is a method of treating or preventing EBV-associated PTLD in a subject, comprising administering a therapeutically effective amount of an anti-gp350 antibody disclosed herein or an immunogenic fragment thereof, one or more epitopes described herein, one or more immunogenic peptides described herein, an EBV antibody-drug conjugate described herein, or a pharmaceutical composition comprising the anti-gp350 antibody or an immunogenic fragment thereof, one or more epitopes, one or more immunogenic peptides, or the EBV antibody-drug conjugate, before or after the transplant in the subject.

[0039] As shown in the working examples, 15 novel EBV gp350-specific mAbs were generated, their binding to gp350 was characterized, their neutralization activity against EBV infection *in vitro* was determined, their cognate epitopes were mapped, and the exact aa residues they recognize on gp350 were defined. Six of the eight previously described epitopes responsible for generating neutralizing and non-nAbs were confirmed and the exact aa residues that they bind were defined. This study also confirmed that the binding epitopes on gp350 that elicit nAbs are between aa 4–443 (59). An additional neutralizing epitope and two new non-neutralizing epitopes,

with one located downstream of the gp350 ectodomain (aa 1-841) were identified. Importantly, the newly developed mAbs have many uses in vaccine development, diagnosis of viral infection, and therapeutic/prophylactic management of post-transplant lymphoproliferative diseases, either individually, in combination with nAb-72A1, or with other mAbs such as anti-gH/gL (E1D1).

[0040] To overcome the existing challenges facing PTLD treatment, novel EBV antibodies and EBV antibody-drug conjugates (ADCs) are developed. The EBV neutralizing antibodies (nAbs) that specifically block new or reactivated EBV infection are conjugated with small molecules that specifically target latent viral protein, EBV nuclear antigen 1 expressed in all EBV+ malignancies. The recent identification and isolation of nAbs against the highly variable viruses HIV-1 (10, 11), influenza (12-14), and respiratory syncytial virus (15) has direct implications for successful EBV protection. Indeed, in 2012, an international, multidisciplinary expert panel recommended use of intravenous (IV) anti-viral nAbs for preventing or treating EBV+ PTLD (16). EBV uses multiple surface glycoproteins (gps), including the major gp350, to infect host cells (17, 18). These gps are expressed on EBV virions and in EBV+ cells (19, 20), and stimulate immune responses in humans and in animal models (21-23), making them attractive targets for an EBV vaccine (24). Multiple lines of evidence suggest that use of anti-gp350 nAbs to protect against EBV-PTLDs is feasible (16): (A) Maternal Abs protect against EBV infection in neonates (25, 26); (B) gp350-expressing EBV+ cells activate complement (27) and mediate Ab-dependent cellular cytotoxicity (28); (C) gp350 vaccines reduce EBV load and protect against EBV+ lymphomas in marmosets (29-32) and protect EBV-naïve adults from EBV-induced mononucleosis (32-34); (D) Compared to control mice, SCID mice injected with peripheral blood mononuclear cells from EBV-naïve donors and immunized with anti-gp350 (72A1) mouse nAb are completely protected against EBV and development of EBV+ tumors or PTLD-like lesions (35); and (E) 72A1 also conferred short-term protection against acquiring EBV after transplantation in 3 out of 4 pediatric patients in a small phase 1 clinical trial (35). However, there was a major drawback: all 4 patients who received 72A1 developed human anti-mouse Abs (HAMA), which can cause side effects and limit treatment efficacy, with one developing a hypersensitivity reaction. This suggests that 72A1 in its native form is not a safe treatment for humans (35). Thus, chimeric (human/mouse), humanized,

or human nAbs, which are safe and effective in the treatment of various cancers, are needed (7, 36-38).

[0041] Pre-existing antibodies provide the primary defense against viral infection. Prophylactic prevention of EBV primary infection has mainly focused on blocking the first step of viral entry by generating neutralizing antibodies (nAbs) that target EBV envelope glycoproteins. Five glycoproteins, in particular, gp350/220 (gp350), gp42, gH, gL, and gB, are required for efficient infection of permissible host cells and have emerged as potential prophylactic targets (23, 24, 61, 64).

[0042] Several studies have indicated that the EBV gp350 as the major immunodominant glycoprotein is an ideal target for EBV nAbs production. Although the ectodomain of EBV gp350 (aa 1-841) has been shown to contain at least eight unique CD21 binding epitopes, only one of these epitopes (aa 142-161) is capable of eliciting nAbs (57-58). The aa residues that constitute the other epitopes and their role in generating nAb has not been elucidated, as this information would be valuable in the precise design of effective EBV peptide vaccine. To date, nAb-72A1 remains the only EBV antibody with proven clinical prophylactic efficacy, as it has been shown to confer short-term protection by reducing and delaying EBV infection onset in immunized pediatric transplant patients (35).

[0043] EBV predominantly infects epithelial cells and B cells, reflecting the viral tropism and the cellular ontogeny for EBV-associated malignancies (17). There are two schools of thought on how the initial EBV transmission into the human host occurs. In the first infection model, the incoming virus engages with ephrin receptor A2 via heterodimeric gH/gL, which triggers gB fusion with the epithelial cell membrane and entry of the virus into the cytoplasm (17). This interaction is thought to occur in the oral mucosa, where the virus undergoes lytic replication to release virions that subsequently infect B cells. In the alternative model, the incoming virus binds to the host cell via complement receptor type 1 (CR1)/CD35 (45) and/or CR2/CD21 through its major immunodominant glycoprotein, gp350 (65). The interaction between gp350 and CD35 and/or CD21 triggers viral adsorption, capping, and endocytosis into the B cell (66), which subsequently leads to interaction between heterotrimeric viral glycoproteins complex, gp42/gH/gL, binding to HLA class II molecules to activate gB membrane fusion and entry. Because these two models are not necessarily mutually exclusive, and given that both gp350 and gH/gL

complex are important in initiating the first viral contact with host cells, use of nAbs that target either gp350 or gH/gL complex or both may potentially block incoming virus at the oral mucosa.

[0044] Nearly all EBV-infected individuals develop nAbs directed to the ectodomains of these glycoproteins (52, 67). These antibodies can prevent neonatal infection, and can protect against acute infectious mononucleosis in adolescents and several human lymphoid and epithelial malignancies associated with EBV infection (30, 32-34). Although numerous monoclonal antibodies (mAbs) have been generated against EBV gp350 (53, 68, 69), only two murine mAbs, the non-neutralizing 2L10 and the neutralizing 72A1, have been extensively characterized and made commercially available (68, 69). Importantly, nAb-72A1 conferred short-term clinical protection against EBV transmission after transplantation in pediatric patients in a small phase I clinical trial (35).

[0045] EBV gp350 is the most immunogenic envelope glycoproteins on the virions. It is a type 1 membrane protein that encodes for 907 amino acid (aa) residues. A single splice of the primary transcript deletes 197 codons and joins gp350 codons 501 and 699, in frame, to generate the gp220 messenger RNA. Both gp350 and gp220 are comprised of the same 18-aa residue at the C terminus that is located within the viral membrane, a 25-aa residue at the transmembrane-spanning domain, and a large highly glycosylated N-terminal ectodomain, aa 1-841 (57). The first 470 aa of gp350 are sufficient for binding CD21 in B cells, as demonstrated by a truncated gp350 (aa 1-470) blocking the binding of EBV to B cells and reducing viral infectivity (17). The gp350-binding domain on CD21 is mapped to N-terminal short consensus repeats (SCRs) 1 and 2, which also bind to a bioactive fragment of complement protein 3 (C3d) (34, 68). A soluble truncated EBV gp350 fragment (aa 1-470) and soluble CD21 SCR1 and SCR2 can block EBV infection and immortalization of primary B cells (57). However, gp350 binding to CD35 is not restricted to N-terminal SCRs; it binds long homologous repeat regions as well as SCRs 29-30 (57).

[0046] The gp350 ectodomain is heavily glycosylated, with both N- and O-linked sugars, which accounts for over half of the molecular mass of the protein. Currently there is only one crystal structure available for gp350, comprised of a truncated structure between 4-443 aa, with at least 14 glycosylated arginines coating

the protein with sugars, with the exception of a single glycan-free patch (59). Mutational studies of several residues in the glycan-free patch resulted in the loss of CD21 binding (59), suggesting that binding of CD35 and CD21 by gp350 is mediated within this region.

[0047] There are at least eight unique CD21 binding epitopes located at the N-terminus of the gp350 ectodomain (58); at least one of these epitopes (aa 142-161) is capable of eliciting nAbs (57, 58). The aa residues 142-161 are also the binding site for nAb-72A1 (59, 68). Using gp350 synthetic peptides binding to CD21 on the surface of a B cell line, additional gp350 epitope was identified in the C-terminal region of gp350 (aa 822-841), suggesting it is involved in EBV invasion of B cells (58). The role of other epitopes in eliciting nAbs has not been fully investigated. Furthermore, the exact aa residues that comprise the core binding sites for epitope capable of eliciting neutralization and non-neutralization antibodies have not been determined. Mapping the EBV gp350 protein residues defining immunodominant epitopes, identifying the critical aa residues of the eight epitopes, and defining their roles in generating neutralizing and non-nAbs will guide rational design and construction of an efficacious EBV gp350-based vaccine that would focus the B-cell responses to the protective epitopes.

[0048] As demonstrated in the working examples, 23 hybridomas producing antibodies against EBV gp350 were generated. To assess their clinical potential and utility in informing future prophylactic and therapeutic vaccine design: (1) the ability of the antibodies produced by the new hybridomas to detect gp350 protein was tested by enzyme-linked immunosorbent assay (ELISA), flow cytometry, and immunoblot; (2) the unique CDRs of the heavy and light chains of all 23 hybridomas were sequenced to identify novel mAbs; (3) the efficacy of each mAb to neutralize EBV infection *in vitro* was measured; and (4) RepliTope peptide microarrays were used to identify gp350 core aa residues recognized by neutralizing and non-neutralizing mAbs. Using the newly generated antibodies, a new epitope bound preferentially by nAbs was identified, distinct from the canonical neutralizing epitope bound by nAb-72A1, as well as two immunodominant epitopes bound by both neutralizing and non-nAbs.

[0049] Figure 1 demonstrates characterization of the 23 gp350-specific antibodies produced by hybridomas and their ability to bind gp350 protein. Figure 2

demonstrates that out of 23 hybridomas, 15 hybridomas were determined to be monoclonal and novel based on the VH and VL CDR sequences compared to those of the reported nAb-72A1 (57).

[0050] Following confirmation that the new mAbs recognized EBV gp350 antigen and had unique VH-VL sequences, further characterization revealed that mAbs HB9, HB10, HB11, HB17, and HB20 inhibited EBV infection in a dose-dependent manner, with HB17 and HB20 being the best neutralizers (Figure 3). Thus, provided herein are new nAbs and non-nAbs against EBV infection for therapeutic and prophylactic uses.

[0051] Various methods, including lectin/ricin immune-affinity assay (69), purified mAbs (53, 70), purified soluble gp350 mutants (57), synthetic peptides (51, 55, 54), cell binding assays (58), and crystal structure of partial gp350 protein (4-443) (59), have been used to identify the critical gp350 epitopes responsible for its interaction with the CD35 and CD21 cellular receptors (for a detailed summary review see Table 2).

Table 2: Summary of gp350 Epitope Mapping over Time Using Different Methodologies			
Method	mAbs/ protein/ peptides	Number of epitopes	Reference
Competitive binding assay Tagged mAb vs untagged mAbs	Newly generated mAbs from purified virus or whole cell prep after induction	8 epitopes—Sequence not defined	Qualtiere et al., 1987 (53)
Binding studies—determine the effects of anti-gp350 mAbs on gp350 binding to CR2	Newly generated mAbs (61)	2 possible regions identified by sequence alignment to C3d sequence. 1. aa 21-28 2. aa 372-378	Nemerow et al 1987 (54)
Binding studies	Peptide and protein	1. aa 21-28 2. N-terminus of gp350	Nemerow et al 1989 (55)
Dot Blot immunoassay Purified truncated protein incubated with mAbs	Protein – 8 clones overlapping N and C terminus portions of protein mAbs from Qualtiere et al., 1987	3 sequences defined 1. aa 310-325 2. aa 326-439 3. aa 733-841	Zhang et al., 1991 (56)

Peptide digest and immunoprecipitation	Truncated and mutant protein mAbs (72A1 and BOS-1)	Narrowed down to the first 470 residues	Tanner et al., 1998 (57)
Peptide cell binding assay- to 2 CR2 positive (Raji and Ramos) and 1 negative (P3HR-1) cell line	Synthesized peptides covering gp350 (907aa)	7 regions 3 identified 1. aa 142-161 2. aa 282-301 3. aa 822-841	Urquiza et al 2005 (58)
Crystal structure and binding studies	Mutant proteins, mAbs 72a1	3 epitopes (based on 72A1 binding and gp350 4-443) 1. aa 16-29 2. aa 142-161 3. aa 282-301	Szakonyi et al., 2006 (59)
Structural docking studied and antigenicity mapping	Gp350 and CR2 crystal structure alignment/docking	Single epitope (based on gp350 (1-470)) 1. aa 147-165	Sitompul et al., 2012 (60)
Structural alignment (computer modeling of gp350 and 72A1) / docking studies	Peptides (used in immunization) and mAb (72A1)	3 epitopes 1. aa 14-20 2. aa 144-161 3. aa 194-211	Tanner et al., 2015 (51)

[0052] The epitope mapping assay identified a total of nine epitopes including two new epitopes, one bound by all mAbs, including nAb-72A1 and 2L10, regardless of the mAb neutralizing capability suggesting it is an immunodominant epitope on the gp350 protein. Portions of six of the previously described epitopes, including the only currently recognized neutralizing epitope were also identified and their exact aa residues were defined. However, two previously reported epitopes located at aa 282-301 and aa 194-211, which have been reported to be involved in the binding of nAb-72A1 or CD21, respectively (51, 58, 59), were not identified.

[0053] As demonstrated in the working examples, the comparative epitope mapping analysis results identified a novel neutralizing epitope, ₃₈₁GAFASNRTFDIT₃₉₂ (SEQ ID NO; 3), which was bound preferentially by the nAbs HB20 and nAb-72A1, but not the non-neutralizing mAb HB5. The new neutralizing epitope is distinct from the reported canonical nAb-72A1 binding epitope aa 142-161, on gp350, suggesting that epitope 381-392 is a novel epitope on gp350 capable of eliciting nAbs using existing technology and protocols.

[0054] Virus-specific treatments are less likely to target basic metabolic

mechanisms of healthy cells, making them more likely to efficiently kill virus-infected cells with fewer side effects. Until recently, few drug regimens have specifically targeted EBV+ lymphomas. However, in 2015, a few small molecules showed activity against EBV-transformed cells (39). Furthermore, in 2017, Jiang et al. described a novel small molecule (L₂P₄) that shows discriminating anti-proliferative activities against EBV-transformed B lymphoma cells (40).

[0055] Having described the invention with reference to the embodiments and illustrative examples, those in the art may appreciate modifications to the invention as described and illustrated that do not depart from the spirit and scope of the invention as disclosed in the specification. The examples are set forth to aid in understanding the invention but are not intended to, and should not be construed to limit its scope in any way. The examples do not include detailed descriptions of conventional methods. Such methods are well known to those of ordinary skill in the art and are described in numerous publications. Further, all references cited above and in the examples below are hereby incorporated by reference in their entirety, as if fully set forth herein.

EXAMPLES

Example 1: Materials and Methods

[0056] Cells and viruses. EBV-AGS, a human gastric carcinoma cell line infected with a recombinant Akata virus expressing enhanced fluorescent green protein (eGFP) was a kind gift of Dr. Lisa Selin (University of Massachusetts Medical School). Anti-EBV gH/gL (E1D1) hybridoma cell line was a kind gift of Dr. Lindsey Hutt-Fletcher (Louisiana State University Health Sciences Center). Chinese hamster ovary cells (CHO); human embryonic kidney cells expressing SV-40 T antigen (HEK-293T); HEK-293 6E suspension cells; EBV-positive Burkitt lymphoma cells (Raji); myeloma cells (P3X63Ag8.653); and anti-EBV gp350 nAb-72A1 hybridoma cells (HB168) were purchased from American Type Culture Collection (ATCC). EBV-AGS cells were maintained in Ham's F-12 media supplemented with 500 µg/ml neomycin (G418, Gibco). Raji, P3X63Ag8.653, and HB168 hybridoma cells were maintained in RPMI 1640. CHO and HEK-293T cells were maintained in DMEM. HEK-293 6E cells were maintained in FreeStyle F17 Expression Medium supplemented with 0.1% Pluronic F-68. All culture media were supplemented with 10% fetal bovine serum

(FBS), 2% penicillin-streptomycin, and 1% l-glutamine, with the exception of Freestyle F17 expression medium.

[0057] Antibodies and plasmids. Primary antibodies: EBV anti-gp350 nAb (72A1) and anti-gH/gL (E1D1) were purified from the supernatant of HB168 and E1D1 hybridoma cell lines, respectively, using Capturem™ Protein A Maxiprep spin columns (Takara). Anti-gp350/220 mAb (2L10) was purchased from Millipore Sigma.

[0058] Secondary antibodies: Horseradish peroxidase (HRP)-conjugated goat anti-mouse IgG for immunoblot or ELISA were purchased from Bio-Rad. Alexa Fluor (AF) 488-conjugated goat anti-mouse IgG (H+L) for flow cytometry was purchased from Life Sciences Tech. Goat anti-mouse IgG (H+L) secondary antibody and DyLight 650 for epitope mapping were purchased from Thermo Fisher Scientific.

[0059] The construction of the pCI-puro vector and pCAGGS-gp350/220-F has been described (23, 45).

[0060] Virus production and purification. eGFP-tagged EBV was produced from the EBV-infected AGS cell line as described (46). Briefly, EBV-AGS cells were seeded to 90% confluency in T-75 flasks in Ham's F-12 medium containing G418 antibiotic. After the cells reached confluency, G418 media was replaced with Ham's F-12 medium containing 33 ng/ml 12-O-tetradecanoylphorbol-13-acetate (TPA) and 3 mM sodium butyrate (NaB) to induce lytic replication of the virus. Twenty-four h post-induction, the media was replaced with complete Ham's F-12 media without G418, TPA, or NaB and cells were incubated for 4 days at 37°C. The cell supernatant was collected, centrifuged, and filtered using 0.8 µm filter to remove cell debris. The filtered supernatant was ultra-centrifuged using a Beckman-Coulter type 19 rotor for 70 min at 10,000 rpm to pellet the virus. EBV-eGFP virus was titrated in both HEK-293T cells and Raji cells, and stocks were stored at -80°C for subsequent experiments.

[0061] Generation and purification of gp350 virus-like particles. To generate gp350 VLPs, equal amounts (8 µg/plasmid) of the relevant plasmids (pCAGGS-Newcastle disease virus (NDV) M, -NP, and gp350 ectodomain fused to fusion protein cytoplasmic and transmembrane domains) were co-transfected into 80% confluent CHO cells seeded in T-175cm² flasks; supernatant from transfected

cells was collected and VLPs were purified and composition characterized as previously described (47).

[0062] Production of hybridoma cell lines. Seven days prior to immunization, two eight-week-old BALB/c mice were bled for collection of pre-immune serum. The mice were immunized with purified UV-inactivated EBV three times (Day 0, 21, and 35), and boosted every 7 days three times (Day 42, 49, and 56) with VLPs incorporating gp350 on the surface after Day 35. The mice were sacrificed, and their splenocytes were isolated, purified, and used to fuse with P3X63Ag8.653 myeloma cells at a ratio of 3:1 in the presence of polyethylene glycol (PEG, Sigma). Hybridoma cells were seeded in flat-bottom 96-well plates and selected in specialized hybridoma growth media with HAT (Sigma) and 10% FBS.

[0063] Indirect ELISA. Hybridoma cell culture supernatant from wells that had colony-forming cells were tested for antibody production by indirect ELISA. Briefly, immunoplates (Costar 3590; Corning Incorporated) were coated with 50 μ l of 0.5 μ g/ml recombinant EBV gp350/220 (Millipore) protein diluted in phosphate buffered saline (PBS, pH 7.4) and incubated overnight at 4°C. After washing three times with PBS containing 0.05% (v/v) Tween 20 (washing buffer), plates were blocked with 100 μ l washing buffer containing 2% (w/v) bovine serum albumin (BSA) then incubated for 1 h at room temperature and washed as above. 100 μ l of hybridoma supernatant was added to each well (in triplicate) and incubated for 2 h at room temperature. PBS and nAb-72A1 were added as negative and positive controls, respectively. The plates were washed as described above, followed by incubation with goat anti-mouse IgG horseradish peroxidase-conjugated secondary antibody (1:2,000 diluted in PBS) at room temperature for 1 h. The plates were washed again and the chromogenic substrate 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS, Life Science Technologies) was added. The reaction was stopped using ABTS peroxidase stop solution containing 5% sodium dodecyl sulfate (SDS) in water. The absorbance was read at an optical density of 405 nm using an ELISA reader (Molecular Devices). Three independent were performed.

[0064] Antibody purification, quantification, and isotyping. Hybridoma cells from selected individual positive clones were expanded stepwise from 96-well plates to T-75 flasks. At confluence in T-75 flasks, supernatant from individual clones was collected, clarified by centrifugation (4,000 g, 10 min, 4°C), and filtered

through a 0.22- μ m-membrane filter (Millipore). Antibodies were further purified by Capturem™ Protein A Maxiprep (Takara) and stored in PBS (pH 7.4) at 4°C. Antibodies were analyzed by SDS-PAGE to determine purity. Bicinchoninic acid assay (BCA assay; Thermo Fisher Scientific) was conducted to determine the concentration of purified antibodies. Isotype identification was performed with the Rapid ELISA mouse mAb isotyping kit (Thermo Fisher Scientific).

[0065] RNA extraction, cDNA synthesis, and sequencing of the variable region of the mAbs. Total RNA was extracted from 1×10^6 hybridoma cells using the RNeasy Mini Kit (Qiagen). Each hybridoma clone cDNA was synthesized in a total volume of 20 μ l using Tetro Reverse Transcriptase (200 u), RiboSafe RNase Inhibitor, Oligo(dT)18 primer, dNTP mix (10 mM each nucleotide), and 100-200 ng RNA. Reverse transcription was performed at 45°C for 30 min, and terminated at 85°C for 5 min. The cDNA was stored at -20°C. Immunoglobulin (Ig) VH and VL were amplified using the mouse Ig specific primer set purchased from Novagen (48). The VH and VL genes were amplified in separate reactions and PCR products were visualized on 1% agarose gel.

[0066] The VH and VL amplicons were sequenced using an Illumina MiSeq platform: duplicate 50 μ l PCR reactions were performed, each containing 50 ng of purified cDNA, 0.2 mM dNTPs, 1.5 mM MgCl₂, 1.25 U Platinum Taq DNA polymerase, 2.5 μ l of 10x PCR buffer, and 0.5 μ M of each primer designed to amplify the VH and VL. The amplicons were purified using AxyPrep Mag PCR Clean-up kit (Thermo Fisher Scientific). The Illumina primer PCR PE1.0 and index primers were used to allow multiplexing of samples. The library was quantified using ViiA™ 7 Real-Time PCR System (Life Technologies) and visualized for size validation on an Agilent 2100 Bioanalyzer (Agilent Technologies) using a high-sensitivity cDNA assay. The sequencing library pool was diluted to 4 nM and run on a MiSeq desktop sequencer (Illumina). The 600-cycle MiSeq Reagent Kit (Illumina) was used to run the 6 pM library with 20% PhiX (Illumina), and FASTQ files were used for data analysis.

[0067] Chimeric mAb construct generation. To generate chimeric mAbs, the VH and VL sequences were cloned into the dual-vector system pFUSE CHlg/pFUSE CLlg (InvivoGen), which express the constant region of the heavy and light chains of human immunoglobulins, respectively (Genewiz). The constructs

were transiently transfected into HEK-293 6E cells. The supernatants were collected at 72 h post-transfection and IgG was purified using protein A/G affinity chromatography.

[0068] Immunoblot analysis. CHO cells were cultured and stably co-transfected with pCAGGS-gp350/220 F and pCI-puro vector containing a puromycin resistance gene. Forty-eight h post-transfection, DMEM media containing 10 µg/ml of puromycin was added to enrich for cells expressing gp350 protein. Puromycin-resistant clones were expanded, followed by flow cytometry sorting using nAb-72A1 to a purity >90%. EBV gp350-positive CHO cells were harvested and lysed in radioimmunoprecipitation assay buffer (RIPA) followed by centrifugation at 15,000 g for 15 min on a benchtop centrifuge. The supernatants were collected and heated at 95°C for 10 min in SDS sample buffer containing β-mercaptoethanol, then separated using SDS-PAGE. Proteins were transferred onto a nitrocellulose membrane using an iBlot™ Transfer System (Thermo Fisher Scientific) followed by incubation in blocking buffer (1% BSA; 20 mM Tris-HCl, pH 7.5; 137 mM NaCl; and 0.1% Tween-20 [TBST]) for 1 h. The blots were incubated in TBST containing purified anti-gp350 antibodies (1:50) overnight at 4°C. After three washes with TBST, the blots were incubated with a goat anti-mouse conjugated to horseradish peroxidase (1:2000) in TBST for 1 h. After three washes, the antibody-protein complexes were detected using the Amersham ECL Prime Western Blotting Detection Reagent (GE Healthcare). All experiments were independently repeated three times.

[0069] Flow cytometry. To assess the ability of purified anti-gp350 mAb to detect surface expression of EBV gp350 protein by flow cytometry, CHO cells that stably express EBV gp350 were harvested and stained with purified anti-gp350 (10 µg/ml), followed by Alexa Fluor® 488 goat anti-mouse IgG secondary antibody. Flow cytometric analysis was performed on a C-6 FC (BD Biosciences) and data was analyzed using FlowJo Cytometry Analysis software (FlowJo, LLC) as described (47). All the experiments were independently repeated three times.

[0070] EBV neutralization assay. Purified individual anti-gp350 mAbs were incubated with purified AGS-EBV-eGFP (titer calculated to infect at least 20% of HEK293 cells seeded in 100 µl of serum-free DMEM) for 2 h at 37°C. To represent EBV infection of B cells, the pre-incubated anti-gp350 mAbs/AGS-EBV-eGFP were used to infect 5x10⁵ Raji cells seeded in a 96-well plate. Anti-gp350 neutralizing

72A1 and non-neutralizing 2L10 mAbs served as positive and negative controls, respectively. Plates were incubated at 37°C and the number of eGFP+ cells was determined using flow cytometry 48 h post-infection. All dilutions were performed in quintuplicate and the assays repeated three times for Raji cells. Antibody EBV neutralization activity was calculated as: % neutralization = (EBValone–EBVmAb) / (EBValone) x 100.

[0071] Epitope mapping. Anti-gp350 mAbs were incubated with a multi-well EBV GP350/GP340 RepliTope (JPT) peptide microarray displaying 224 peptides (15mers with 11 aa overlap) in 3x7 subarrays. Briefly, anti-gp350 mAbs were diluted in blocking solution (TBS-T and 2% BSA) to a final concentration of 10 µg/ml and incubated with the microarray slide for 1 h at 30°C with shaking. Slides was washed 5 times with wash buffer (TBS-T), followed by incubation with 1 µg/ml secondary antibody Dylight 650 (Thermo Fisher Scientific) for 1 h at 30°C. After washing 5 times with wash buffer and 2 times with distilled water, microarray slides were dried by centrifugation. Detection was performed using the Agilent DNA microarray scanner.

Example 2: Development and characterization of monoclonal antibodies against EBV gp350

[0072] New EBV gp350-specific mAbs were generated and biochemically characterized, and their ability to neutralize EBV infection was evaluated. In addition, the antibodies were used to map immunodominant epitopes on the EBV gp350 protein. 23 novel monoclonal antibodies specific against EBV gp350 were developed. To generate hybridomas, BALB/c mice were immunized with purified UV-inactivated EBV and boosted with virus-like particles (VLPs) that incorporate EBV gp350 ectodomain on the surface to enrich for production of anti-gp350 antibodies, then splenocytes were isolated from the immunized mice and fused with myeloma cells. Specifically, five eight-week-old BALB/c mice were immunized with virus-like particles incorporating gp350/220 on the surface, four times (day 0, 14, 28, and 56) via intraperitoneal injection without adjuvants. At day 64, immunized mice were boosted once intravenously. Animals were sacrificed at Day 70 to harvest splenocytes for fusion with the mouse myeloma P3X63Ag8 cell line.

[0073] Indirect ELISA was used to screen supernatants from the hybridomas for specificity against purified EBV gp350 ectodomain protein (aa 4-863) and 23 hybridomas producing gp350 specific antibodies were identified. These novel antibodies were analyzed by flow cytometric analysis for surface expression of gp350 protein on 10^6 CHO cells transfected with 1 μ g of pCAGGS-gp350. gp350 expressing cells were resuspended in PBS, stained with anti-gp350 mAb, which detects the gp350 ED, followed by secondary Ab goat anti-mouse conjugated to AF488. Additionally, western blot analysis was conducted on untransfected and pCAGGS-gp350 transfected CHO lysate. Anti-gp350 mAb 72A1 was used as a positive control (1:100) and anti-gp350 hybridoma clone supernatants were used at 1:50, and anti-mouse secondary antibody was used at 1:2000.

[0074] The isotypes of the new antibodies were determined to be IgG1 (n=14), IgG2a (n=5), IgG2b (n=1), a mixture of IgG1 and IgG2b (n=1), and a mixture of IgG1 and IgM (n=2). It was found that all 23 hybridoma producing antibodies, designated HB1–23, recognized the gp350 antigen in an initial ELISA screening using unfractionated and unpurified hybridoma supernatants (data not shown). Affinity purification with protein A followed by SDS-PAGE was used to confirm the purity of the antibodies. When quantified amount of the purified antibodies (10 μ g/ml) was re-evaluated using indirect ELISA, all of the 23 antibodies had ELISA signals two times greater than those of phosphate buffered saline (negative control), and were considered as positive or specific to gp350. Of these, five (HB4, HB5, HB7, HB13, and HB14) demonstrated binding affinity equal to or greater than that of the positive control, nAb-72A1 (Figure 1A). This difference in binding of the 23 antibodies could be due to differential exposure of cognate epitopes on gp350 in the assay performed.

[0075] Determining the nature of the binding between an antibody and its target antigen is an important consideration for the performance and specificity of an antibody, as it can involve the recognition of a linear or conformational epitope (49). The antibodies were characterized using immunoblot analysis of denatured gp350 antigen expressed from Chinese hamster ovary (CHO) cells, and 16 of the antibodies reacted to both the 350 kDa and the 220 kDa splice variant. In contrast, HB2, HB3, HB6, HB7, HB13, HB20, and HB21 failed to recognize either of the denatured isoforms of gp350 (Figure 1B). The antibodies were further characterized by flow cytometric analysis of CHO cells stably expressing gp350 on the cell surface,

and HB1, HB2, HB3, HB5, HB6, HB9, HB11, HB12, HB15, HB17, HB19, HB20, and HB21 antibodies readily recognized gp350 (Figures 1C, 1D). Given that HB2, HB3, HB20, and HB21 detected gp350 by flow cytometry, but not by immunoblot, suggests that these four antibodies recognized conformational epitopes on gp350, whereas HB5, HB9, HB11, HB15, HB17, and HB19 recognized both linear and conformational epitopes (Figures 1B-1D). The observation that all 23 anti-gp350 antibodies recognized the gp350 antigen either by indirect ELISA, flow cytometry, or immunoblot assay suggests that antibodies that are specific to EBV gp350 protein were successfully produced.

Example 3: Analysis of the variable heavy and variable light chain sequences

[0076] The complementarity determining regions (CDRs) of the anti-gp350 mAbs were sequenced by amplifying mRNA representing unique heavy/light variable chain sequences using RT-PCR. RT-PCR products were sequenced and analyzed. The sequences of the heavy and light chain variable region genes (VH and VL, respectively) of the 23 new anti-gp350 antibodies, as well as the nAb-72A1, were determined and compared the sequences to published nAb-72A1 sequences (50, 51). The sequence of the CDR of this antibody was recently determined and published, revealing two unique IgG1 heavy chains and two unique light chains, one kappa and one lambda (50, 51). PCR was used to amplify the genes encoding the VH and VL chain regions in cDNA generated from the 23 hybridoma cells as well as from HB168 (nAb-72A1). The PCR products presented distinct bands at approximately 350–400 bp and 450–500 bp for VH and VL, respectively (data not shown). The purified fragments were sequenced using Illumina MiSeq, followed by in silico analysis and CDRs for both VH and VL were identified (Figures 2A, 2B). Two VH and VL sequences of nAb-72A1 were identified as >94% identical to the previously published sequences (50), suggesting that nAb-72A1 exists as a mixed antibody, instead of the reported mAb (51). Similar to nAb-72A1, HB4, HB13, HB15, and HB23 hybridomas each produced a mixture of two antibodies, with unique sequences of the VH chain showing at >5% frequencies, suggesting that they are not mAbs (Table 3). Coding sequences for VL chains for HB7, HB9, and HB17 were unable to be identified, unless the frequencies were lowered to >1% (Table 3); in this case, the identified coding VL chain sequences were identical.

Table 3. Summary of Illumina Dual Demultiplex of V_H and V_L Regions

Sample	Chain	Starting Pairs	PEAR Merged Reads	Length Filtered Reads	Primer Matched Reads	3x Reads	>5% Unique Coding	>5% Unique Non-Coding	
HB1	HEAVY	51,641	51,210	46,655	32,482	22,725	1	0	
	LIGHT	280,048	279,012	100,725	68,041	58,570	1	1	
HB2	HEAVY	22,793	22,621	16,475	11,415	7,429	1	0	
	LIGHT	167,230	166,496	161,764	132,752	115,886	1	1	
HB3	HEAVY	26,382	26,162	25,542	16,910	11,709	1	0	
	LIGHT	12,681	12,609	11,753	9,809	8,023	1	1	
HB4	HEAVY	38,811	38,238	17,151	11,957	7,217	2	0	§
	LIGHT	179,249	129,752	111,996	78,392	66,419	1	1	
HB.5	HEAVY	42,951	42,173	35,793	25,842	17,173	1	0	
	LIGHT	176,073	175,267	168,806	141,712	127,045	1	1	
HB6	HEAVY	26,142	25,981	22,245	15,658	10,453	1	0	
	LIGHT	171,996	171,370	167,730	138,348	122,397	1	1	
HB7	HEAVY	32,443	32,094	25,449	17,615	11,836	1	0	
	LIGHT	67,031	63,924	37,344	26,271	22,378	1	1	*
HB8	HEAVY	140,091	103,349	92,744	58,292	31,583	1	0	
	LIGHT	151,244	115,527	102,439	82,154	70,803	1	1	
HB9	HEAVY	37,057	36,473	19,544	11,585	7,358	1	0	
	LIGHT	409,432	310,529	136,074	106,820	90,063	1	2	*
HB10	HEAVY	38,181	37,981	26,043	17,104	11,391	1	0	
	LIGHT	114,255	112,498	106,914	84,368	75,370	1	1	
HB11	HEAVY	22,225	21,841	6,956	4,408	2,465	1	0	
	LIGHT	106,465	102,278	65,332	50,232	44,527	1	0	
HB12	HEAVY	83,044	82,355	46,350	30,276	20,886	1	0	
	LIGHT	53,098	47,336	15,823	7,560	5,845	1	1	
HB13	HEAVY	81,451	80,372	47,995	32,216	20,139	2	0	§
	LIGHT	27,314	24,774	8,987	5,457	4,104	2	1	
HB14	HEAVY	76,299	75,357	28,309	19,104	12,939	1	0	
	LIGHT	153,011	149,264	48,474	29,710	25,133	1	1	
HB.15	HEAVY	26,551	26,410	16,387	11,434	7,002	2	0	§
	LIGHT	78,525	77,778	43,509	29,504	24,731	1	1	
HB16	HEAVY	54,249	53,943	9,517	7,128	4,179	1	0	
	LIGHT	42,048	40,351	30,602	22,758	18,251	2	1	§
HB17	HEAVY	111,614	110,882	81,428	50,844	35,949	1	0	
	LIGHT	102,490	100,488	83,925	65,925	57,727	1	1	*
HB18	HEAVY	211,215	155,410	146,256	91,009	50,308	1	0	
	LIGHT	212,261	161,879	155,235	123,096	105,959	1	1	
HB19	HEAVY	109,692	82,221	20,546	12,587	7,274	1	1	
	LIGHT	70,828	69,744	62,572	48,354	42,051	1	1	
HB20	HEAVY	15,781	15,632	12,789	7,757	4,852	1	0	
	LIGHT	135,527	133,208	118,513	90,717	78,701	1	1	
HB21	HEAVY	15,312	15,202	8,577	5,645	3,420	1	0	
	LIGHT	102,450	100,171	89,059	68,552	60,500	1	1	
HB22	HEAVY	217,959	156,488	154,008	95,755	50,245	1	0	
	LIGHT	205,334	156,986	143,386	108,728	85,136	1	0	
HB23	HEAVY	196,390	143,929	123,028	71,076	39,358	2	0	§
	LIGHT	158,594	120,140	115,476	90,004	78,787	2	0	
72A1	HEAVY	213,480	158,199	156,215	107,395	68,486	2	0	§
	LIGHT	187,216	140,964	132,945	105,783	91,208	1	1	

* Hybridoma with V_L chain sequences identified with >1% frequency, § Hybridoma with more than one unique, plausible-coding V_H chain sequence with > 5% frequency.

The term “unique” refers to unique sequence counts (so, identical sequences found in a substantial frequency of merged reads, not necessarily unique compared to other samples).

[0077] The analysis and comparison of the VH and VL chain gene sequences of the 23 hybridomas compared to HB168 (nAb-72A1) showed unique sequences within the CDR 1-3 region. Only HB8 and HB18 had identical VH and VL chain gene sequences, suggesting that the two are the same clone isolated separately; therefore, HB18 was excluded from subsequent experiments. One of the two HB15 antibodies had identical VH and VL gene sequences to that of HB10, however based on the previous characterization, the presence of the additional antibody in HB15 was sufficient to confer subtle differences in biochemical characterizations for gp350 between the two antibodies. Thus, based on the sequence analysis (Figure 2), 15 unique anti-gp350 mAbs were generated, with distinct biochemical and sequence identity from commercially available nAb-72A1.

Example 4: Neutralization Assay

[0078] The ability of the 15 mAbs (10 µg/ml or 50 µg/ml) to neutralize purified eGFP-tagged AGS-EBV infection of the Raji B cell line *in vitro* was evaluated following standardized procedures (52) and determined the percentage of eGFP+ cells using flow cytometry as described (45, 47). The nAbs 72A1 and E1D1 were used as positive controls, whereas the non-neutralizing mAb 2L10 was used as a negative control. Because HB4, HB7, HB13, HB15, HB16, HB19, HB21, and HB23 were confirmed to be mixtures based on isotyping or sequence data, they were eliminated from further consideration in the neutralization assay. An antibody was considered a neutralizer if it inhibited EBV infection >20% at 10 µg/ml and >60% at 50 µg/ml. Several mAbs inhibited EBV infection in a dose-dependent manner. HB20 and HB17 were the most effective at preventing EBV infection of Raji cells *in vitro*, whereby they reduced infection by 40% and >60%, and >30% and 80% at 10 µg/ml and 50 µg/ml, respectively (Figures 3A and 3B). The HB9 and HB10 antibodies prevented EBV infection of Raji cells by ~25% at 10 µg/ml and ~60% at 50 µg/ml. The HB11 antibody neutralized <20% at 10 µg/ml, but showed a dose-dependent increase at 50 µg/ml by neutralizing EBV infection by 60%. By the set neutralization parameters, HB1–3, HB5–8, HB12, HB14, and HB22 did not neutralize EBV infection of Raji cells. In comparison, both nAb-72A1 and nAb-E1D1 neutralized EBV

infection by >70% and 40%, respectively, at 10 µg/ml. The nAb-72A1 neutralized EBV infection by >95% at 50 µg/ml, whereas nAb- E1D1 neutralized infection by >60% at 50 µg/ml. As expected, the negative control, mAb-2L10, did not neutralize viral infection at either concentration. Based on the neutralization assay results, the remaining 15 mAbs were organized into two distinct groups, neutralizers (+) and non-neutralizers (-) as summarized in Table 4.

Table 4: Summarized Biochemical and Functional Characterizations of Anti-gp350 Antibodies						
Antibody	IgG sub-class	Light chain	ELISA binding to purified EBV gp350/220	FACS (CHO Cells)	Western blot	Neutralization activity
HB1	IgG1	κ	+	+	+	-
HB2	IgG2a	κ	+	+	-	-
HB3	IgG2a	κ	-	+	-	-
HB4	IgG1	κ	+	-	+	ND
HB5	IgG2a	κ	+	+	+	-
HB6	IgG1	κ	-	+	-	-
HB7	IgG1	κ	+	-	-	-
HB8	IgG1	κ	+	-	+	-
HB9	IgG2a	κ	-	+	+	+
HB10	IgG1	κ	-	-	+	+
HB11	IgG1	κ	+	+	+	+
HB12	IgG1	κ	-	+	+	-
HB13	IgG1	κ	+	-	-	ND
HB14	IgG1	κ	+	-	+	-
HB15	IgG1	κ	+	+	+	ND
HB16	IgG1/ IgM	κ	+	-	+	ND
HB17	IgG2b	κ	+	+	+	+
HB19	IgG1/ IgM	κ	+	+	+	ND
HB20	IgG2a	κ	-	+	-	+
HB21	IgG1/IgG2b	κ	-	+	-	ND
HB22	IgG1	κ	-	-	+	-
HB23	IgG1	κ	-	+	+	ND
72A1	IgG1	κ	+	+	+	+
E1D1 (anti gH/gL)	IgG1	κ	-	-	-	+

ND=Not determined, + = positive, - = negative, ELISA = enzyme-linked immunosorbent assay, κ = Kappa.

Example 5: Epitope Mapping

[0079] The 15 new anti-gp350 mAbs (neutralizers vs. non-neutralizers) were used to identify most, if not all, of the relevant immunodominant aa residues targeted by both the nAbs and non-nAbs. RepliTope approach was used, in which overlapping peptides [15-mer with 11 aa overlap] that cover the complete sequence of gp350 (aa 1-907) were immobilized on microarray slides and probed with the purified anti-gp350 antibodies in an ELISA format. nAb-72A1 was used as positive control, because the cognate epitopes bound by the antibody have previously been reported.

[0080] Two epitopes, ²⁵³TPIPGTGYAYSLRLTPRPVSRFL₂₇₅ and ³⁹³VSGLGTAPKTLIITRTATNATTT₄₁₅, were both bound by all 15 mAbs, as well as nAb-72A1 and 2L10, regardless of their neutralizing or non-neutralizing capabilities. This consensus suggests that these epitopes are immunodominant. Several of the 15 mAbs (HB2, HB3, HB8, HB11, HB12, HB14, HB17, and HB22), as well as nAb-72A1, bound to ³⁴¹ANSPNVTVTAFWAWPNNTE₃₅₉. Two epitopes, aa 341-359 and aa 393-415, were found within the previously identified single epitope II, which is encoded by nucleotides between 3,186 bp and 3,528 bp, corresponding to aa 326-439 of gp350 (56). Two mAbs, HB1 and HB10, bound ⁶⁰⁵TTPTPNATGPTVGETSPQA₆₂₃, an epitope located within the gp350 (501-699) splice region that is involved in generation of the 220 kDa splice variant. A total of eight mAbs (HB1-3, HB8, HB10-12, and HB22) also bound to the region between ¹MEAALLVCQYTIQSLIHLTGEDPG₂₄, which includes a region homologous to C3d, another molecule known to interact with CD21 (51, 55). Two epitopes common between most nAbs and non-nAbs, ⁸²¹PPSTSSKLRPRWTFSTPPV₈₃₉ and ⁸⁷⁵LLLLVMADCAFRRNLSTSHTYTPPY₈₉₉, were located upstream and downstream, respectively, of the transmembrane domain on the C-terminus of gp350. Epitope aa 821-839 is located within the previously identified epitope I, which is located between aa 733-841 (56). Furthermore, epitope aa 821-839 is potentially involved in EBV infection of B cells (58). However, the study could not identify two epitopes located at aa 282-301 and aa 194-211, which were previously shown to be involved in the binding of nAb-72A1 and CD21, respectively (51, 58, 59). It was shown that nAb-72A1 bound ¹⁴⁵EMQNPVYLIPETVPYIKWDN₁₆₄, one of the neutralizing epitopes on gp350 (51, 58, 59). Nine gp350 epitopes identified by 15 neutralizing and non-

neutralizing mAbs are shown in the sequence below and summarized in Table 5). Residues in bold represent the gp220 splice variant region. Residues in boxes represent RepliTope-identified epitopes and exact residues. Italic residues represent canonical neutralizing epitope, underlined residues represent epitope bound by all assayed mAb.

```

1  MEAALLVCQY  TIQSLIHLTG  EDPGFFNVEI  PEFPPFYPTCN  VCTADVNVTI
NFDVGGKKHQ

61  LDLDFGQLTP  HTKAVYQPRG  AFGGSENATN  LFLLELLGAG  ELALTMRSKK
LPINVTTGEE

121  QQVSLESVDV  YFQDVEGTMW  CHHAEMQNPV  YLIPETVPYI  KWDNCNSTNI
TAVVRAQGLD

181  VTLPLSLPTS  AQDSNFSVKT  EMLGNEIDIE  CIMEDGEISQ  VLPGDNKFNI
TCSGYESHVP

241  SGGILTSTSP  VATPIPGTGY  AYSLRLTPRP  VSRFLGNNSI  LYVFYSGNGP
KASGGDYCIQ

301  SNIVFSDEIP  ASQDMPTNTT  DITYVGDNAT  YSVPMVTSED  ANSPNVTVTA
FWAWPNNTET

361  DFKCKWTLTS  GTPSGCENIS  GAFASNRTFD  IT/VSGLGATAP  KTLIITRTAT
NATTTTHKVI

421  FSKAPESTTT  SPTLNTTGFA  DPNTTGLPS  STHVPTNLTA  PASTGPTVST
ADVTSPTPAG

481  TTSGASPVTP  SPSPWDNGTE  SKAPDMTSST  SPVTTPTPNA  TSPTPAVTPP
TPNATSPTPA

541  VTTPTPNATS  PTLGKTSPTS  AVTTPTPNAT  SPTLGKTSPT  SAVTTPTPNA
TSPTLGKTSPT

601  TSAVTTPTPN  ATGPTVGETS  PQANATNHTL  GGTSPTPVVT  SQPKNATSAV
TTGQHNITSS

661  STSSMSLRPS  SNPETLSPST  SDNSTSHMPL  L TSAHPTGGE  NITQVTPASI
STHHVSTSSP

721  APRPGTTSQA  SGPGNSSTST  KPGEVNVTKG  TPPQNATSPQ  APSGQKTAVP
TVTSTGGKAN

```

781 STTGKHTTG HGARTSTEPT TDYGGDSTTP RPRYNATTYL PPSTSSKLRP
 RWTFTSPPVT

841 TAQATVPVPP TSQPRFSNLS MLVLQWASIA VLTLLLLLVM ADCAFRRLS
 TSHTYTTTPPY

901 DDAETTV

Table 5: Antibody Binding gp350 Epitopes

Epitope	Amino acid sequence	Epitope Properties
1	¹ MEAALLVCQYTIQSLIHLTGEDPG ₂₄	
2	¹⁴⁵ EMQNPVYLIPETVPYIKWDN ₁₆₄	Neutralizing
3	²⁵³ TPIPGTGYAYSLRLTPRPVSRFL ₂₇₅	Immunodominant/ novel
4	³⁴¹ ANSPNVTVTAFWAWPNNTE ₃₅₉	
5	³⁸¹ GAFASNRTFDIT ₃₉₂	Neutralizing/ novel
6	³⁹³ VSGLGTAPKTLIITRTATNATTT ₄₁₅	Immunodominant
7	⁶⁰⁵ TTPTPNATGPTVGETSPQA ₆₂₃	
8	⁸²¹ PPSTSSKLRPRWTFTSPPV ₈₃₉	
9	⁸⁷⁵ LLLLVMADCAFRRLSTSHTYTTTPPY ₈₉₉	novel

Example 6: Construction of chimeric gp350 nAbs

[0081] Chimeric gp350 nAbs were constructed according to the diagrams of Figure 4. First, mouse antibodies against human gp350 were developed, and then the mouse antibody variable region is fused to human constant region, for example, to human IgM. The heavy chain and light chain variable regions of the mouse antibody were cloned into pFUSE-CHlg and pFUSE2-CLlg vectors, respectively, followed by co-transfection of mammalian cells with recombinant pFUSE-CHlg and pFUSE2-CLlg vectors. The expression vectors were obtained from InvivoGen, and the expression was conducted in CHO cells or HEK293 cells, available from ATCC. The construction schemes and expression of heavy and light chains of clone 19 are shown in Figure 4.

[0082] Analysis of VH-VL sequence from the HB168 (nAb-72A1) hybridoma revealed that the hybridoma produced two antibodies: one that is gp350-specific and another that recognizes mineral oil-induced plasmacytoma (MOPC) (57). To

further investigate gp350 for additional neutralizing epitopes, the gp350-specific nAb-72A1 VH-VL sequence was used to generate chimeric (mouse/human) recombinant antibodies. Similarly, the VH-VL sequence for the HB20 antibody, which the neutralization analysis above showed to be one of the best nAb, was used to generate chimeric antibody. A negative control chimeric recombinant antibody was generated using VH-VL sequences from the gp350-specific but non-neutralizing HB5 antibody mentioned above (Figure 3).

[0083] Comparative analysis of the epitope binding pattern was performed using the chimeric recombinant antibodies and a novel epitope was revealed, ³⁸¹GAFASNRTFDIT₃₉₂, which was bound by HB20 and nAb-72A1, but not by HB5, and is distinct from the ¹⁴⁵EMQNPVYLIPETVPYIKWDN₁₆₄ epitope, which is bound only by the nAb-72A1. The sequence below shows the novel gp350-neutralizing epitope identified by epitope mapping of neutralizing (nAb-72A1 and HB20) vs non-neutralizing (HB5) anti-gp350 mAbs. Epitopes bound by nAb-72A1 (double underlined), HB20 (wavy underlined), and HB5 (boxed) are indicated. Residues 501-700 represent splice variant region (bold), single underlined residues represent epitopes bound by nAb-72A1, HB20 and HB5, and italic residues represent epitopes bound by nAbs.

```

1  MEAALLVCQY TIQSLIHLTG EDPGFFNVEI PEPFFYPTCN VCTADVNVTI NFDVGGKKHQ
61  LDLDFGQLTP HTKAVYQPRG AFGGSENATN LFLLELLGAG ELALTMRSKK LPINVTTGEE
121 QQVSLESVDV YFQDVFGTMW CHHAEMQNPV YLIPETVPYI KWDNCNSTNI TAVVRAQGLD
181 VTLPLSLPTS AQDSNFSVKT EMLGNEIDIE CIMEDGEISQ VLPGDNKFNI TCSGYESHVP
241 SGGILTSTSP VATPIPGTGY AYSLRLTPRP VSRFLGNNSI LYVFYSGNGP KASGGDYCIQ
301 SNIVFSDEIP ASQDMPTNTT DITYVGDNAT YSVPMVTSSE ANSPNVTVTA FWAHPNNTET
361 DFKCKWTLTS GTPSGCENIS GAFASNRTFD ITVSGLGATP KTLIITRTAT NATTTTHKVI
421 FSKAPESTTT SPTLNNTGFA DPNTTGLPS STHVPTNLTA PASTGPTVST ADVTSPTPAG
481 TTSGASPVTP SPSPWDNGTE SKAPDMTSST SPVTTPTPNA TSPTPAVTTT TPNATSPTPA
541 VTTPTPNATS PTLGKTSPTS AVTTPTPNAT SPTLGKTSPT SAVTTPTPNA TSPTLGKTSPT
601 TSAVTTPTPN ATGPTVGETS PQANATNHTL GGTSPTPVVT SQPKNATSAV TTGQHNTSS
661 STSSMSLRPS SNPETLSPST SDNSTSHMPL LSAHPTGGE NITQVTPASI STHHVSTSSP
721 APRPGTTSQA SGPGNSSTST KPGEVNVTKG TPPQNATSPQ APSGQKTAVP TVTSTGGKAN
781 STTGKHTTG HGARTSTEPT TDYGGDSTTP RPRYNATTYL PPSTSSKLRP RWTFTSPPVIT

```

841 TAQATVPVPP TSQPRFSNLS MLVLQWASLA VLTLLLLLVLM ADCAFRRNLS TSHTYTTPPY
901 DDAETTV

Example 7: Development of antibody-small molecule conjugates (ADCs)

[0084] Using small molecule L₂P₄ as an example, antibody-small molecule conjugates can be developed as illustrated in Figure 5. One or more small molecules can be conjugated to the antibody heavy chain or light chain via a reactive donor group and a C-terminal acceptor group. Small molecule L₂P₄ was disclosed in the publication by Jiang et al.⁴⁰ After validating the function of the purified chimeric gp350 nAbs using ELISA, flow cytometry (FC), and surface plasmon resonance, the nAbs can be conjugated to L₂P₄ in a site-specific manner via a Val-Cit dipeptide linker, which releases the active agent upon internalization of the ADC.^{41, 42}

Example 8: Efficacy test of ADCs

[0085] To screen for optimal dose and identify the best nAb clone for pre-clinical studies, the ability of the ADC to neutralize EBV *in vitro* and to protect against PTLD *in vivo* is tested using a humanized mouse, as described^{43,44}. In brief, purified chimeric ADC (or controls: PBS, or isotype-matched non-nAbs) is injected by I.V. into humanized mice, followed by EBV-B95-8-eGFP challenge. Mice are monitored regularly and euthanized upon signs of illness or after a preset limit of 100 days. Routine histology and necropsy are conducted to assess the efficacy of the ADC to protect against EBV infection and PTLD development.

REFERENCES

The references listed below, and all references cited in the specification are hereby incorporated by reference in their entireties, as if fully set forth herein.

1. Rickinson AB, Kieff E. Epstein-Barr Virus. In: Knipe DM, Howley PM, eds. *Fields Virology*. Philadelphia: Lippincott, Williams & Wilkins; 2007:2680-2700.
2. Goedert JJ, Cote TR, Virgo P, et al. Spectrum of AIDS-associated malignant disorders. *Lancet* 1998;351:1833-9.
3. Coté TR, Biggar RJ, Rosenberg PS, et al. Non-Hodgkin's lymphoma among people with AIDS: Incidence, presentation and public health burden. *International Journal of Cancer* 1998;73:645-50.
4. Benkerrou M, Jais JP, Leblond V, et al. Anti-B-cell monoclonal antibody treatment of severe posttransplant B-lymphoproliferative disorder: prognostic factors and long-term outcome. *Blood* 1998;92:3137-47.
5. Cruz RJ, Jr., Ramachandra S, Sasatomi E, et al. Surgical management of gastrointestinal posttransplant lymphoproliferative disorders in liver transplant recipients. *Transplantation* 2012;94:417-23.
6. Faro A, Kurland G, Michaels MG, et al. Interferon-alpha affects the immune response in post-transplant lymphoproliferative disorder. *Am J Respir Crit Care Med* 1996;153:1442-7.
7. Milpied N, Vasseur B, Parquet N, et al. Humanized anti-CD20 monoclonal antibody (Rituximab) in post transplant B-lymphoproliferative disorder: a retrospective analysis on 32 patients. *Ann Oncol* 2000;11 Suppl 1:113-6.
8. Papadopoulos EB, Ladanyi M, Emanuel D, et al. Infusions of donor leukocytes to treat Epstein-Barr virus-associated lymphoproliferative disorders after allogeneic bone marrow transplantation. *N Engl J Med* 1994;330:1185-91.
9. Starzl TE, Holmes JH. Experience in renal transplantation. 1964.
10. Stamatatos L, Morris L, Burton DR, Mascola JR. Neutralizing antibodies generated during natural HIV-1 infection: good news for an HIV-1 vaccine? *Nature medicine* 2009;15:866-70.
11. Walker LM, Phogat SK, Chan-Hui P-Y, et al. Broad and potent neutralizing antibodies from an African donor reveal a new HIV-1 vaccine target. *Science* 2009;326:285-9.

12. Sui J, Hwang WC, Perez S, et al. Structural and functional bases for broad-spectrum neutralization of avian and human influenza A viruses. *Nature structural & molecular biology* 2009;16:265-73.
13. Han T, Marasco WA. Structural basis of influenza virus neutralization. *Annals of the New York Academy of Sciences* 2011;1217:178-90.
14. Wrammert J, Smith K, Miller J, et al. Rapid cloning of high-affinity human monoclonal antibodies against influenza virus. *Nature* 2008;453:667-71.
15. Piedimonte G, King KA, Holmgren NL, Bertrand PJ, Rodriguez MM, Hirsch RL. A humanized monoclonal antibody against respiratory syncytial virus (palivizumab) inhibits RSV-induced neurogenic-mediated inflammation in rat airways. *Pediatr Res* 2000;47:351-6.
16. Glotz D, Chapman JR, Dharnidharka VR, et al. The seville expert workshop for progress in posttransplant lymphoproliferative disorders. *Transplantation* 2012;94:784-93.
17. Connolly SA, Jackson JO, Jardetzky TS, Longnecker R. Fusing structure and function: a structural view of the herpesvirus entry machinery: A structural view of herpesvirus entry machinery. *Nat Rev Microbiol* 2011;9:369-81.
18. Eisenberg RJ, Atanasiu D, Cairns TM, Gallagher JR, Krummenacher C, Cohen GH. Herpes virus fusion and entry: a story with many characters. *Viruses* 2012;4:800-32.
19. Henle G, Henle W, Diehl V. Relation of Burkitt's tumor-associated herpes-type virus to infectious mononucleosis. *Proceedings of the National Academy of Sciences of the United States of America* 1968;59:94.
20. Cohen JI, Fauci AS, Varmus H, Nabel GJ. Epstein-Barr Virus: An Important Vaccine Target for Cancer Prevention. *Science Translational Medicine* 2011;3:107fs7-fs7.
21. Khanna R, Sherritt M, Burrows SR. EBV structural antigens, gp350 and gp85, as targets for ex vivo virus-specific CTL during acute infectious mononucleosis: potential use of gp350/gp85 CTL epitopes for vaccine design. *The Journal of Immunology* 1999;162:3063-9.
22. Thorley-Lawson DA, Geilinger K. Monoclonal antibodies against the major glycoprotein (gp350/220) of Epstein-Barr virus neutralize infectivity. *Proceedings of the National Academy of Sciences of the United States of America* 1980;77:5307-11.

23. Perez EM, Foley J, Tison T, Silva R, Ogembo JG. Novel Epstein-Barr virus-like particles incorporating gH/gL-EBNA1 or gB-LMP2 induce high neutralizing antibody titers and EBV-specific T-cell responses in immunized mice. *Oncotarget* 2016.
24. Cohen JL. Epstein-barr virus vaccines. *Clin Transl Immunology* 2015;4:e32.
25. Biggar RJ, Henle W, Fleisher G, Böcker J, Lennette ET, Henle G. Primary Epstein-Barr virus infections in african infants. I. Decline of maternal antibodies and time of infection. *International Journal of Cancer* 1978;22:239-43.
26. Biggar RJ, Henle G, Böcker J, Lennette ET, Fleisher G, Henle W. Primary Epstein-Barr virus infections in African infants. II. Clinical and serological observations during seroconversion. *International Journal of Cancer* 1978;22:244-50.
27. Mold C, Bradt B, Nemerow G, Cooper N. Activation of the alternative complement pathway by EBV and the viral envelope glycoprotein, gp350. *The Journal of Immunology* 1988;140:3867-74.
28. Khyatti M, Patel PC, Stefanescu I, Menezes J. Epstein-Barr virus (EBV) glycoprotein gp350 expressed on transfected cells resistant to natural killer cell activity serves as a target antigen for EBV-specific antibody-dependent cellular cytotoxicity. *Journal of virology* 1991;65:996-1001.
29. Finerty S, Mackett M, Arrand JR, Watkins PE, Tarlton J, Morgan AJ. Immunization of cottontop tamarins and rabbits with a candidate vaccine against the Epstein-Barr virus based on the major viral envelope glycoprotein gp340 and alum. *Vaccine* 1994;12:1180-4.
30. Gu SY, Huang TM, Ruan L, et al. First EBV vaccine trial in humans using recombinant vaccinia virus expressing the major membrane antigen. *Dev Biol Stand* 1995;84:171-7.
31. Mok H, Cheng X, Xu Q, et al. Evaluation of Measles Vaccine Virus as a Vector to Deliver Respiratory Syncytial Virus Fusion Protein or Epstein-Barr Virus Glycoprotein gp350. *Open Virol J* 2012;6:12-22.
32. Moutschen M, Leonard P, Sokal EM, et al. Phase I/II studies to evaluate safety and immunogenicity of a recombinant gp350 Epstein-Barr virus vaccine in healthy adults. *Vaccine* 2007;25:4697-705.
33. Sokal EM, Hoppenbrouwers K, Vandermeulen C, et al. Recombinant gp350 vaccine for infectious mononucleosis: a phase 2, randomized, double-blind, placebo-controlled trial to evaluate the safety, immunogenicity, and efficacy of an Epstein-

Barr virus vaccine in healthy young adults. *Journal of Infectious Diseases* 2007;196:1749-53.

34. Rees L, Tizard EJ, Morgan AJ, et al. A phase I trial of epstein-barr virus gp350 vaccine for children with chronic kidney disease awaiting transplantation. *Transplantation* 2009;88:1025-9.

35. Haque T, Johannessen I, Dombagoda D, et al. A mouse monoclonal antibody against Epstein-Barr virus envelope glycoprotein 350 prevents infection both in vitro and in vivo. *J Infect Dis* 2006;194:584-7.

36. Ferrara N, Hillan KJ, Gerber H-P, Novotny W. Discovery and development of bevacizumab, an anti-VEGF antibody for treating cancer. *Nat Rev Drug Discov* 2004;3:391-400.

37. Hudis CA. Trastuzumab — Mechanism of Action and Use in Clinical Practice. *New England Journal of Medicine* 2007;357:39-51.

38. Jonker DJ, O'Callaghan CJ, Karapetis CS, et al. Cetuximab for the Treatment of Colorectal Cancer. *New England Journal of Medicine* 2007;357:2040-8.

39. Dzung RK, Jha HC, Lu J, Saha A, Banerjee S, Robertson ES. Small molecule growth inhibitors of human oncogenic gammaherpesvirus infected B-cells. *Mol Oncol* 2015;9:365-76.

40. Jiang L, Lan R, Huang T, et al. EBNA1-targeted probe for the imaging and growth inhibition of tumours associated with the Epstein–Barr virus. *Nature Biomedical Engineering* 2017;1:0042.

41. Dubowchik GM, Firestone RA. Cathepsin B-sensitive dipeptide prodrugs. A model study of structural requirements for efficient release of doxorubicin. *Bioorganic & medicinal chemistry letters* 1998;8:3341-6.

42. Senter PD, Sievers EL. The discovery and development of brentuximab vedotin for use in relapsed Hodgkin lymphoma and systemic anaplastic large cell lymphoma. *Nature biotechnology* 2012;30:631-7.

43. Haque T, Johannessen I, Dombagoda D, et al. A mouse monoclonal antibody against Epstein-Barr virus envelope glycoprotein 350 prevents infection both in vitro and in vivo. *J Infect Dis* 2006;194:584-7.

44. Jangalwe S, Shultz LD, Mathew A, Brehm MA. Improved B cell development in humanized NOD-scid IL2R γ manu mice transgenically expressing human stem cell factor, granulocyte-macrophage colony-stimulating factor and interleukin-3. *Immun Inflamm Dis* 2016;4:427-40.

45. Ogembo JG, Kannan L, Ghiran I, Nicholson-Weller A, Finberg RW, Tsokos GC, Fingerroth JD. 2013. Human complement receptor type 1/CD35 is an Epstein-Barr Virus receptor. *Cell Rep* 3:371-385.
46. Speck P, Longnecker R. 1999. Epstein-Barr virus (EBV) infection visualized by EGFP expression demonstrates dependence on known mediators of EBV entry. *Arch Virol* 144:1123-1137.
47. Ogembo JG, Muraswki MR, McGinnes LW, Parcharidou A, Sutiwisesak R, Tison T, Avendano J, Agnani D, Finberg RW, Morrison TG. 2015. A chimeric EBV gp350/220-based VLP replicates the virion B-cell attachment mechanism and elicits long-lasting neutralizing antibodies in mice. *Journal of Translational Medicine* 13:50.
48. Jones ST, Bendig MM. 1991. Rapid PCR-cloning of full-length mouse immunoglobulin variable regions. *Biotechnology (N Y)* 9:579.
49. Sela M, Schechter B, Schechter I, Borek F. 1967. Antibodies to Sequential and Conformational Determinants. *Cold Spring Harbor Symposia on Quantitative Biology* 32:537-545.
50. Herrman M, Muhe J, Quink C, Wang F. 2015. Epstein-Barr Virus gp350 Can Functionally Replace the Rhesus Lymphocryptovirus Major Membrane Glycoprotein and Does Not Restrict Infection of Rhesus Macaques. *J Virol* 90:1222-1230.
51. Tanner JE, Coincon M, Leblond V, Hu J, Fang JM, Sygusch J, Alfieri C. 2015. Peptides designed to spatially depict the Epstein-Barr virus major virion glycoprotein gp350 neutralization epitope elicit antibodies that block virus-neutralizing antibody 72A1 interaction with the native gp350 molecule. *J Virol* 89:4932-4941.
52. Sashihara J, Burbelo PD, Savoldo B, Pierson TC, Cohen JI. 2009. Human antibody titers to Epstein-Barr Virus (EBV) gp350 correlate with neutralization of infectivity better than antibody titers to EBV gp42 using a rapid flow cytometry-based EBV neutralization assay. *Virology* 391:249-256.
53. Qualtiere LF, Decoteau JF, Hassan Nasr-el-Din M. 1987. Epitope mapping of the major Epstein-Barr virus outer envelope glycoprotein gp350/220. *J Gen Virol* 68 (Pt 2):535-543.
54. Nemerow GR, Mold C, Schwend VK, Tollefson V, Cooper NR. 1987. Identification of Gp350 as the Viral Glycoprotein Mediating Attachment of Epstein-Barr-Virus (Ebv) to the Ebv/C3d Receptor of B-Cells - Sequence Homology of Gp350 and C3-Complement Fragment C3d. *Journal of Virology* 61:1416-1420.

55. Nemerow GR, Houghten RA, Moore MD, Cooper NR. 1989. Identification of an epitope in the major envelope protein of Epstein-Barr virus that mediates viral binding to the B lymphocyte EBV receptor (CR2). *Cell* 56:369-377.
56. Zhang PF, Klutch M, Armstrong G, Qualtiere L, Pearson G, Marcus-Sekura CJ. 1991. Mapping of the epitopes of Epstein-Barr virus gp350 using monoclonal antibodies and recombinant proteins expressed in *Escherichia coli* defines three antigenic determinants. *J Gen Virol* 72 (Pt 11):2747-2755.
57. Tanner J, Whang Y, Sample J, Sears A, Kieff E. 1988. Soluble gp350/220 and deletion mutant glycoproteins block Epstein-Barr virus adsorption to lymphocytes. *J Virol* 62:4452-4464.
58. Urquiza M, Lopez R, Patino H, Rosas JE, Patarroyo ME. 2005. Identification of three gp350/220 regions involved in Epstein-Barr virus invasion of host cells. *J Biol Chem* 280:35598-35605.
59. Szakonyi G, Klein MG, Hannan JP, Young KA, Ma RZ, Asokan R, Holers VM, Chen XS. 2006. Structure of the Epstein-Barr virus major envelope glycoprotein. *Nat Struct Mol Biol* 13:996-1001.
60. Sitompul LS, Widodo N, Djati MS, Utomo DH. 2012. Epitope mapping of gp350/220 conserved domain of Epstein Barr virus to develop nasopharyngeal carcinoma (npc) vaccine. *Bioinformation* 8:479-482.
61. Balfour HH, Jr. 2014. Progress, prospects, and problems in Epstein-Barr virus vaccine development. *Curr Opin Virol* 6C:1-5.
62. Henle G, Henle W. 1979. The virus as the etiologic agent of infectious mononucleosis, p 297-320, *The Epstein-Barr Virus*. Springer.
63. Luzuriaga K, Sullivan JL. 2010. Infectious mononucleosis. *N Engl J Med* 362:1993-2000.
64. Cui X, Cao Z, Chen Q, Arjunaraja S, Snow AL, Snapper CM. 2016. Rabbits immunized with Epstein-Barr virus gH/gL or gB recombinant proteins elicit higher serum virus neutralizing activity than gp350. *Vaccine*.
65. Fingerroth JD, Weis JJ, Tedder TF, Strominger JL, Biro PA, Fearon DT. 1984. Epstein-Barr virus receptor of human B lymphocytes is the C3d receptor CR2. *Proc Natl Acad Sci U S A* 81:4510-4514.
66. Tanner J, Weis J, Fearon D, Whang Y, Kieff E. 1987. Epstein-Barr virus gp350/220 binding to the B lymphocyte C3d receptor mediates adsorption, capping, and endocytosis. *Cell* 50:203-213.

67. Weiss ER, Alter G, Ogembo JG, Henderson JL, Tabak B, Bakis Y, Somasundaran M, Garber M, Selin L, Luzuriaga K. 2017. High Epstein-Barr Virus Load and Genomic Diversity Are Associated with Generation of gp350-Specific Neutralizing Antibodies following Acute Infectious Mononucleosis. *J Virol* 91.
68. Hoffman GJ, Lazarowitz SG, Hayward SD. 1980. Monoclonal antibody against a 250,000-dalton glycoprotein of Epstein-Barr virus identifies a membrane antigen and a neutralizing antigen. *Proc Natl Acad Sci U S A* 77:2979-2983.
69. Thorley-Lawson DA, Geilinger K. 1980. Monoclonal antibodies against the major glycoprotein (gp350/220) of Epstein-Barr virus neutralize infectivity. *Proc Natl Acad Sci U S A* 77:5307-5311.
70. Alfarano C, Andrade CE, Anthony K, Bahroos N, Bajec M, Bantoft K, Betel D, Bobechko B, Boutilier K, Burgess E, Buzadzija K, Caverio R, D'Abreo C, Donaldson I, Dorairajoo D, Dumontier MJ, Dumontier MR, Earles V, Farrall R, Feldman H, Garderman E, Gong Y, Gonzaga R, Grytsan V, Gryz E, Gu V, Haldorsen E, Halupa A, Haw R, Hrvojic A, Hurrell L, Isserlin R, Jack F, Juma F, Khan A, Kon T, Konopinsky S, Le V, Lee E, Ling S, Magidin M, Moniakakis J, Montojo J, Moore S, Muskat B, Ng I, Paraiso JP, Parker B, Pintilie G, Pirone R, et al. 2005. The Biomolecular Interaction Network Database and related tools 2005 update. *Nucleic Acids Res* 33:D418-424.

CLAIMS

1. An antibody or an immunogenic fragment thereof that specifically binds to Epstein-Barr virus (EBV) glycoprotein (gp) 350 or glycoprotein 220.

2. The antibody or the immunogenic fragment thereof of claim 1, wherein the antibody or the immunogenic fragment thereof specifically binds to a gp350 epitope comprising the amino acid sequence of SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3.

3. The antibody or the immunogenic fragment thereof of claim 1 or claim 2, comprising a VH region comprising CDR1, CDR2 and CDR3 having an amino acid sequence of SEQ ID NOs: 4-51.

4. The antibody or the immunogenic fragment thereof of claim 1 or claim 2, comprising a VL region comprising CDR1, CDR2 and CDR3 having an amino acid sequence of SEQ ID NOs: 52-99.

5. The antibody of any one of claims 1-4, wherein the antibody is a monoclonal antibody.

6. The antibody of any one of claims 1-4, wherein the antibody is a chimeric antibody, a humanized antibody or a human antibody.

7. An immunogenic peptide comprising one or more gp350 epitopes, wherein each epitope has an amino acid sequence identical to or sharing at least 60% similarity to SEQ ID NO: 1, SEQ ID NO: 2, or SEQ ID NO: 3.

8. The immunogenic peptide of claim 7, comprising a first domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO:1, a second domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO: 2, and a third domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO: 3.

9. The immunogenic peptide of claim 7, comprising a first domain comprising an amino acid sequence having at least 60% similarity to SEQ ID NO: 1,

SEQ ID NO: 2, or SEQ ID NO: 3, and a second domain comprising a known immunogenic peptide such as keyhole limpet hemocyanin (KLH) peptide.

10. An antibody-small molecule conjugate comprising:
an antibody or an immunogenic fragment thereof that specifically binds to EBV gp350 or EBV gp220; and
a small molecule having an anti-proliferative activity against EBV-transformed cells,

wherein the small molecule is conjugated to the antibody.

11. The conjugate of claim 10, wherein the antibody is a monoclonal antibody.

12. The conjugate of claim 10, wherein the antibody is a chimeric antibody, a humanized antibody or a human antibody.

13. The conjugate of any one of claims 10-12, wherein the small molecule is a growth inhibitor of EBV infected B cells.

14. The conjugate of any one of claims 10-13, wherein the small molecule is L₂P₄, 2-butyne diamide, or a derivative thereof.

15. The conjugate of any one of claims 10-14, wherein the small molecule is conjugated to the antibody via a linker or an adaptor.

16. The conjugate of any one of claims 10-15, wherein the small molecule is conjugated to the constant region of the heavy chain or the light chain of the antibody.

17. A pharmaceutical composition comprising the antibody or the immunogenic fragment thereof of any one of claims 1-6, the immunogenic peptide of any one of claims 7-9, or the conjugate of any one of claim 10-16.

18. A method of neutralizing EBV infection comprising administering to a subject infected with EBV a therapeutically effective amount of the antibody or the immunogenic fragment thereof of any one of claims 1-6, the immunogenic peptide of

any one of claims 7-9, the conjugate of any one of claim 10-16, or the pharmaceutical composition of claim 17.

19. A method of preventing EBV infection comprising administering to a subject at an elevated risk of EBV infection a therapeutically effective amount of the antibody or the immunogenic fragment thereof of any one of claims 1-6, the immunogenic peptide of any one of claims 7-9, the conjugate of any one of claim 10-16, or the pharmaceutical composition of claim 17.

20. A method of preventing a post-transplant lymphoproliferative disease (PTLD) comprising administering to a subject who is a transplant recipient a therapeutically effective amount of the antibody or the immunogenic fragment thereof of any one of claims 1-6, the immunogenic peptide of any one of claims 7-9, the conjugate of any one of claim 10-16, or the pharmaceutical composition of claim 17.

21. The method of claim 20, wherein the subject is a pediatric transplant recipient who is EBV naïve or an adult transplant recipient.

22. The method of claim 20 or claim 21, wherein the administration is before, during, and/or after the transplant.

23. A method of treating an EBV-associated cancer comprising administering to a subject suffering from an EBV-associated cancer a therapeutically effective amount of the antibody or the immunogenic fragment thereof of any one of claims 1-6, the immunogenic peptide of any one of claims 7-9, the conjugate of any one of claim 10-16, or the pharmaceutical composition of claim 17.

24. The method of claim 23, wherein the EBV-associated cancer is selected from the group consisting of Hodgkin lymphoma, Burkitt lymphoma, gastric cancer, and nasopharyngeal carcinoma.

25. A method of immunizing or vaccinating a subject against EBV infection comprising administering to the subject a therapeutically effective amount of the antibody or the immunogenic fragment thereof of any one of claims 1-6, the immunogenic peptide of any one of claims 7-9, or a pharmaceutical composition thereof.

26. A method of inducing the production of neutralizing antibodies against a EBV in a subject, comprising administering an effective amount of a gp350 epitope represented by SEQ ID NO: 3 or an immunogenic peptide comprising SEQ ID NO: 3.

27. The method of any one of claims 18-26, wherein the subject is human.

28. Use of the antibody or the immunogenic fragment thereof of any one of claims 1-6, the immunogenic peptide of any one of claims 7-9 for inducing the production of a neutralizing antibody against EBV in a subject.

29. Use of the antibody or the immunogenic fragment thereof of any one of claims 1-6, the immunogenic peptide of any one of claims 7-9 for the manufacture of a medicament for inducing the production of a neutralizing antibody against EBV in a subject.

Figure 1A






Figure 1B





Anti-EBV gp350/220- Alexa Fluor 488

Figure 1D




Figure 2A

FR-1		95	96	6	69	6	9	55	5	9565	75	757	5	9	5	95	7	6	7	7	6	66	77
Conservation:		1	--	FELVPGTSMKISCKAS	ESSTEDTMMWQKSHGKLEWIGL	INPNNGH	ETYNQKFKGKATILDKSSSTAYMEVLSL	78															
72A1		1	--	PGLVAFPSQSLSITCTVS	SHLLTYGVHVVWQPPGKGLKLV	WAGCG	--SYNIGKMSQVILKMSL	77															
HB1		1	--	PELAKPGETVKLSCKAS	SYFTTASGMHWKLTPEKGLKMMW	INTKCTGE	PTYADFKGRFAFLESASTAVLIQINN	78															
HB2		1	--	PELAKPGASVLSCKAS	SYTFASVMQVWKQWPGQGLEWIS	IMPNNHGH	INYNDEFKFKASITVDKSSSTAYMQLSSL	78															
HB3		1	--	AEVLVPGASVLSCKAS	SYFTFTNHNINWTKQRPQGLDWIG	IMPNNYDI	TSYNQKFKGKATILVDKSSSTAYMELSSL	78															
HB5		1	--	PELRPGETVKLSCKAS	SYFTDTEMHWKQTFPGKGLKMMW	INTRICE	FRYADDFKGRFAFLESASTAVLIQINN	78															
HB6		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB7		1	--	PELRPGETVKLSCKAS	SYFTYGMWVKQCAPGKGLKMMW	INTYCTGE	PTYADDFKGRFAFLESASTAVLIQINN	78															
HB8		1	--	CGLVKPGGSLKLSAAS	HTFSSYTNMWKQTFPEARLEWVA	ISSGGSY	TYYPDSVYKGRFTISROKANTILXIQMSSL	78															
HB9		1	--	AEVLVPGASVLSCKAS	SYFTTSYGMHWKQWPGSQSPEN	IRFIMPNSGH	INYNDEFKFKATILVDKSSSTAYMQLSSL	78															
HB10		1	--	PELVKPGQSLKLSAAS	HTFSSYTNMWKQWPGKLEWIG	IMPNNHGH	ETYNQKFKGKATILDKSSSTAYMELSSL	78															
HB11		1	--	AEVLVPGASVLSCKAS	SYFTFTNHNINWTKQRPQGLDWIG	IMPNNYDI	TSYNQKFKGKATILVDKSSSTAYMELSSL	78															
HB12		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB13		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB14		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB15		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB16		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB17		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB18		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB19		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB20		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB21		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	78															
HB22		1	--	AEVLVPGASVLSCKAS	SYFTDTEMHWKQTFVGLGLEWIG	ISPGKSGE	TAYNCKFKGKATILADKSSSTAYMELSSL	77															
Consensus_aa1				..s.IV.Pt.o.floCpkoC64oTay..hVY+Q.Psp.L-W.G.Lps.st.f.YappKt+hisio.DpSsp.pdtphsSL																			
Consensus_aa1				..s.IV.Pt.o.floCpkoC64oTay..hVY+Q.Psp.L-W.G.Lps.st.f.YappKt+hisio.DpSsp.pdtphsSL																			

CDR-1

CDR-2

FR-4		7869	95955	6699699	97998999
------	--	------	-------	---------	--

Figure 3A




Figure 3B




Figure 4





Figure 5



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US18/30030

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
 - a. ☐ forming part of the international application as filed:
 - ☐ in the form of an Annex C/ST.25 text file.
 - ☐ on paper or in the form of an image file.
 - b. ☐ furnished together with the international application under PCT Rule 13ter.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
 - c. ☐ furnished subsequent to the international filing date for the purposes of international search only:
 - ☐ in the form of an Annex C/ST.25 text file (Rule 13ter.1(a)).
 - ☐ on paper or in the form of an image file (Rule 13ter.1(b) and Administrative Instructions, Section 713).
2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

An invitation to Furnish Nucleotide and/or Amino Acid Sequence Listing and to Pay, Where Applicable, Late Furnishing Fee ("ISA/225") was mailed on 11 May 2018 (11.05.2017). No approved electronic sequence listing was received in response to the ISA/225. Therefore, the international search was carried out only to the extent possible.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US18/30030

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 5-6, 14-25, 27-29
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US18/30030

A. CLASSIFICATION OF SUBJECT MATTER

IPC - A61K 39/245, 39/395; C07K 14/05, 16/08 (2018.01)

CPC - A61K 39/245, 39/395; C07K 14/05, 16/085

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

See Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	(TANNER, JE et al.) Peptides designed to spatially depict the Epstein-Barr virus major virion glycoprotein gp350 neutralization epitope elicit antibodies that block virus neutralizing antibody 72A1 interaction with the native gp350 molecule. Journal of Virology. May 2015, Vol. 80, No. 9, pages 4932-4941, ePub 18 February 2015; abstract; page 4933, 1st column, 1st paragraph; page 4933, 1st column, 4th-5th paragraphs; page 4933, 2nd column, 1st paragraph; page 4934, 1st column, 2nd-3rd paragraphs; page 4939, 1st column, 1st paragraph – page 4939, 2nd column, 3rd paragraph; Figure 1; Table 2; DOI: 10.1128/JVI.03269-14	1-2, 3/1-2, 4/1-2, 7-9, 26 -- 10-12, 13/10-12
Y	(JIANG, L et al.) EBNA 1-targeted probe for the imaging and growth inhibition of tumours associated with the Epstein-Barr virus. Nature: Biomedical Engineering. 13 March 2017, Vol. 1, No. 0042, pages 1-10; page 8, 1st column, 3rd paragraph; Figure 4J; DOI: 10.1038/s41551-017-0042	10-12, 13/10-12
A	(BAGHIAN, A et al.) Glycoprotein B of Human Herpesvirus 8 Is a Component of the Virion in a Cleaved Form Composed of Amino- and Carboxyl-Terminal Fragments. Virology. 30 March 2000, Vol. 269, No. 1, pages 18-25; DOI: 10.1006/viro.2000.0198	1-2, 3/1-2, 4/1-2, 7-13, 26

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

26 July 2018 (26.07.2018)

Date of mailing of the international search report

07 AUG 2018

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Shane Thomas

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774