

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
25 May 2001 (25.05.2001)

PCT


(10) International Publication Number
WO 01/35949 A2

- (51) International Patent Classification⁷: **A61K 31/00**
- (21) International Application Number: PCT/US00/30130
- (22) International Filing Date:
13 November 2000 (13.11.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/165,108 12 November 1999 (12.11.1999) US
09/710,754 10 November 2000 (10.11.2000) US
- (71) Applicant (for all designated States except US):
GENELABS TECHNOLOGIES, INC. [US/US];
505 Penobscot Drive, Redwood City, CA 94063 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **SCHWARTZ, Kenneth, E.** [US/US]; 511 Virginia Avenue, San Mateo, CA 94402 (US). **GURWITH, Marc, J.** [US/US]; 13157 La Paloma Road, Los Altos, CA 94022 (US).
- (74) Agents: **HALIDAY, Emily, M.** et al.; Skjerven, Morrill, MacPherson, Suite 700, 25 Metro Drive, San Jose, CA 95110 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TREATMENT OF SLE WITH DEHYDROEPIANDROSTERONE



(57) Abstract: This invention provides a method of treating systemic lupus erythematosus (SLE) with (DHEA) and a related pharmaceutical product.



WO 01/35949 A2

TREATMENT OF SLE WITH DEHYDROEPIANDROSTERONE**CROSS-REFERENCE TO RELATED APPLICATION**

This application claims the benefit of U.S. Provisional Application No. 60/165,108 filed November 12, 1999, entitled "Treatment of SLE with Dehydroepiandrosterone" and naming Kenneth Schwartz as the inventor. This prior application is hereby incorporated by reference in its entirety.

FIELD OF THE INVENTION

The field of this invention concerns improvements in the treatment of systemic lupus erythematosus (SLE).

BACKGROUND OF THE INVENTION**References**

- Alcocer-Varela, et al., J. Clin. Invest. 69:1388, (1982).
Barrett-Connor, et al. New Engl. J. Med. 315:1519, (1986).
Gutierrez-Ramos, et al., Nature 346:27, (1990).
15 Heinz, D., et al., Steroids Lip Res. 5(4):216 (1974).
Jungers, et al., Arthritis Rheum. 25: 454, (1982).
Krupp, LP, et al., Arch Neurol, 46:1121 (1989).
Lahita, et al., Arthritis Rheum 26:1517, (1983).
Liang, et al., Arthritis Rheum 32:1107 (1989).
20 Linker-Israeli, et.al., J. Immunol. 130:2651, (1983).
Lucas, et al. J. Clin. Invest. 75:2091, (1985).
Murakawy, et al., J. Immunol 134:187, (1985).
Roubinian, et al., Arthritis Rheum. 22:1399, (1979).
Steinberg, et al., Arthritis Rheum. 22:1170, (1979).
25 Vande Wiele, et al. Recent Prog. Horm. Res. 19:75, (1963).

SUMMARY OF THE INVENTION

The present invention provides a method for treating systemic lupus erythematosus (SLE). The method entails administering an effective amount of a

pharmaceutically active form of DHEA to an individual with SLE, and at least about 4 weeks after initiating DHEA administration, determining the following disease-activity and constitutional-symptom variables characterizing the individual's SLE condition: the SLE Disease activity index (SLEDAI), the Systemic Lupus Activity Measurement (SLAM), the Patient Visual Analog Scale (Patient VAS), and the Krupp Fatigue Severity Score (KFSS). The differences between the values for SLEDAI, KFSS, VAS, and SLAM after initiating DHEA administration and baseline values for SLEDAI, KFSS, VAS, and SLAM before initiating DHEA administration are then determined. A decrease in three of these four variables and either a decrease, no change, or an increase of no more than about 5% of a baseline value in the fourth variable indicates that the individual is responding to said DHEA administration. Preferably, the individual is a human SLE patient, and more preferably a human SLE patient with a SLEDAI value greater than 2.

The DHEA polymorphs known as forms I and II are preferred for use in the treatment method of the invention. Accordingly, in preferred embodiments, at least about 85%, and more preferably at least about 95%, of the DHEA administered is present as the form I polymorph, the form II polymorph, or a combination thereof.

In one embodiment, DHEA is administered at a dose and for a period effective to produce a decrease in three of the four disease-activity and constitutional-symptom variables characterizing an individual's SLE condition and either a decrease, no change, or an increase of no more than about 3% of a pretreatment baseline value in the fourth variable. In another embodiment, DHEA is administered at a dose effective to reduce the risk of an SLE flare at about 200 days of DHEA administration by at least about 5%. Preferably, the method entails administering a daily oral dose of at least about 100 mg, and more preferably about 200 mg, of a pharmaceutically active DHEA. In one embodiment, DHEA administration is continued for a period of at least about 40 weeks.

One advantage of the treatment method of the invention is that it can be combined with other therapies. For example, the treatment method can be carried out with an SLE patient receiving an orally administered drug, such as a glucocorticoid, a non-steroidal anti-inflammatory agent, an immunosuppressant, or an anti-malarial drug. In this case, the treatment method includes continuing administration of the drug during the period of DHEA administration.

The invention also provides a pharmaceutical product for use in treating systemic lupus erythematosus (SLE) in an individual. The pharmaceutical product includes a plurality of doses of a pharmaceutically active form of DHEA, and instructions for performing the treatment method of the invention.

5 The invention includes the use of the pharmaceutical product of the invention, for treating systemic lupus erythematosus (SLE) in a human patient.

 Another aspect of the invention is the use of a pharmaceutically active acid, salt, or ester form dehydroepiandrosterone (DHEA) in the preparation of a tablet-form medicament for use in treating systemic lupus erythematosus (SLE) in a human patient, with
10 a greater than 50% expectation of achieving improvement in the measured values of at least three of the disease-activity and constitutional-symptom variables characterizing a patient's SLE condition consisting of SLEDAI, KFSS, VAS, and SLAM, and an increase of no greater than 5% of a pretreatment baseline value in the fourth variable, where the change in each variable is determined from the difference between a pretreatment baseline value and
15 values during treatment, when the patient has a pre-treatment SLEDAI value greater than 2, and is treated with a dose of at least about 100 mg per day DHEA, administered orally. In a preferred variation of this embodiment, the patient is treated with a dose of at least about 200 mg DHEA per day for a period of at least about 40 weeks.

BRIEF DESCRIPTION OF THE DRAWING

20 Fig. 1 shows the results of a clinical trial conducted to determine the effect of DHEA treatment on SLE flares, which is described below in connection with Tables 6-8. Patients received 200 mg DHEA per day, administered orally in capsule form, or with a non-drug capsule (placebo). Fig. 1 shows a graph of the percentage of patients surviving without a flare over the indicated duration of the study.

25 DETAILED DESCRIPTION

 The present invention relates to the treatment of systemic lupus erythematosus (SLE) with dehydroepiandrosterone (DHEA). In preferred embodiments, the method employs four SLE evaluation criteria: the SLE Disease activity index (SLEDAI), the Systemic Lupus Activity Measurement (SLAM), the Patient Visual Analog Scale (Patient
30 VAS), and the Krupp Fatigue Severity Score (KFSS). Preferably, individuals with a

SLEDAI value greater than 2.0 are selected for DHEA treatment. In addition, the treatment method entails comparing the values for these disease-activity and constitutional-symptom variables before and after initiating DHEA administration as an indication of response to DHEA administration.

5 **I. SLE Evaluation Criteria**

A. The SLE Disease Activity Index

The SLEDAI (SLE Disease Activity Index) was developed as a clinical index for the measurement of disease activity (Bombardier). It consists of a weighted index of 24 questions covering 9 organ systems for disease activity in SLE. Table 1 shows the SLEDAI
10 questionnaire employed in the present study. As seen, the SLEDAI score is weighted, with 8 points being assigned for central nervous system and vascular, 4 points for renal and musculoskeletal, 2 points for serosal, dermal, and immunologic, and 1 point for constitutional and hematologic findings. Each system is rated as present or absent over the ten day period before and including the day of evaluation. Thus, SLEDAI, as a quantitative measure of
15 SLE disease activity, assesses only recent disease activity, which must be present within the previous 10 days as an indication of current disease state.

B. The Systemic Lupus Activity Measurement

The SLAM (Systemic Lupus Activity Measurement) covers symptoms that occurred during the previous month and includes 24 clinical manifestations and 8 laboratory
20 parameters (Liang). Clinical and laboratory parameters are scored for both activity and severity. A manifestation or symptom is determined to be either active or not active; severity is then used to expand a scale's graduations and is "judged by the need to treat with immunosuppressive agents, the need to follow the patient more closely, or the functional or prognostic consequences of the manifestations" (Liang). Table 2 below shows the SLAM
25 questionnaire used in the present study. The theoretical range for SLAM is 0 to 86, with scores in SLE patients typically falling in the range of 5 to 20.

C. The Patient Visual Analog Scale

In determining Patient VAS (Patient Visual Analog Scale), patient overall (or “global”) assessment is based on a visual analog scale, with a range from "no problems at all" (0) to "the worst I have ever felt" (100). The scale is shown at the bottom (question 2) in Table 3. The patient is asked to mark on the scale how they have felt the past week. The distance in mm's from "0" is measured to arrive at the score.

D. The Krupp Fatigue Severity Score

The KFSS (Krupp Fatigue Severity Score) is a composite score based on the mean of 9 questions, with a possible range from 1-7, where higher scores indicate greater fatigue (Krupp), as shown at the top (question 1) in Table 3. The theoretical range of KFSS is 0 to 7, with scores ranging typically in SLE patients between 4 and 7.

E. Changes in Evaluation Criteria Values

For all four variables, an increase in the measured or determined value represents a more severe condition or assessment, and a decrease in the value represents an improvement in the condition or assessment.

For each of the disease activity variables (SLEDAI and SLAM) and constitutional symptom variables (Patient VAS and KFSS), the value of interest for each patient is determined as the difference between a pretreatment baseline value, calculated as the mean of two baseline values, taken prior to any treatment with DHEA), and values taken during treatment, calculated as the mean of all values obtained during on-treatment scheduled visits, e.g., 13, 26, 39, and 52 weeks of treatment.

For each of the variables, an improvement in the variable is defined as either (i) no change in the variable or (ii) a decrease in the variable. Thus, for example, an improvement in a baseline SLEDAI score of 5.00 would be any value 5.00 or less. A change in a variable that represents a worsening of the state or condition is any increase in the measured value.

A successful responder in the clinical trial studies that were carried out, in accordance with the invention, is defined as a patient that shows an improvement in three of the four variables (that is, SLEDAI, SLAM, Patient VAS, and KFSS) and an increase in the

fourth variable that is no greater than 5% of the pretreatment baseline value. Thus, for a baseline SLEDAI value of 5.0, and a 3% cap, the maximum allowed increase in SLEDAI value, for classification as a 3% responder is 0.15

Table 1

SLEDAI SCORE

VISIT: Screening Qualifying 1 2 3 Completion/Early Termination Other
 WEEK: (0) (13) (26) (39) (52)

Check box: If descriptor is present at the time of visit or in the preceding 10 days.

Wt.	Present	Descriptor	Definition
8	<input type="checkbox"/>	Seizure	- Recent onset. Exclude Metabolic, infectious or drug cause.
8	<input type="checkbox"/>	Psychosis	- Altered ability to function in normal activity due to severe disturbance in the perception of reality. Include hallucinations, incoherence, marked loose associations, impoverished thought content, marked illogical thinking, bizarre, disorganized, or catatonic behavior. Exclude uremia and drug causes.
8	<input type="checkbox"/>	Organic Brain Syndrome	- Altered mental function with impaired orientation, memory or other intellectual function, with rapid onset and fluctuating clinical features. Include clouding of consciousness with reduced capacity to focus, and inability to sustain attention to environment, plus at least two of the following: perceptual disturbance, incoherent speech, insomnia or daytime drowsiness, or increased or decreased psychomotor activity. Exclude metabolic, infectious or drug causes.
8	<input type="checkbox"/>	Visual Disturbance	- Retinal changes of SLE. Include cytoid bodies, retinal hemorrhages, serous exudate or hemorrhages in the choroid, or optic neuritis. Exclude hypertension, infection, or drug causes.
8	<input type="checkbox"/>	Cranial Nerve Disorder	- New onset of sensory or motor neuropathy involving cranial nerves.
8	<input type="checkbox"/>	Lupus Headache	- Severe persistent headache; may be migrainous, but must be non-responsive to narcotic analgesia.
8	<input type="checkbox"/>	CVA	- New onset of cerebrovascular accident(s). Exclude arteriosclerosis.
8	<input type="checkbox"/>	Vasculitis	- Ulceration, gangrene, tender finger nodules, periungual infarction, splinter hemorrhages, or biopsy or angiogram proof of vasculitis.
4	<input type="checkbox"/>	Arthritis	- More than 2 joints with pain and signs of inflammation (i.e. tenderness, swelling, or effusion).
4	<input type="checkbox"/>	Myositis	- Proximal muscle aching/weakness, associated with elevated creatine phosphokinase/adolase or electromyogram changes or a biopsy showing myositis.
4	<input type="checkbox"/>	Urinary Casts	- Heme-granular or red blood cell casts.
4	<input type="checkbox"/>	Hematuria	- >5 red blood cells/high power field. Exclude stone, infection or other cause.
4	<input type="checkbox"/>	Proteinuria	- >0.5 gm/24 hours. New onset or recent increase of more than 0.5 gm/24 hours.
4	<input type="checkbox"/>	Pyuria	- >5 white blood cells/high power field. Exclude infection.
2	<input type="checkbox"/>	New Rash	- New onset or recurrence of inflammatory type rash.
2	<input type="checkbox"/>	Alopecia	- New onset or recurrence of abnormal, patchy or diffuse loss of hair.
2	<input type="checkbox"/>	Mucosal ulcers	- New onset or recurrence of oral or nasal ulcerations.
2	<input type="checkbox"/>	Pleurisy	- Pleuritic chest pain with pleural rub or effusion, or pleural thickening.
2	<input type="checkbox"/>	Pericarditis	- Pericardial pain with at least 1 of the following: rub, effusion, or electrocardiogram confirmation.
2	<input type="checkbox"/>	Low complement	- Decrease in CH50, C3, or C4 below the lower limit of normal for testing laboratory
2	<input type="checkbox"/>	Increased DNA binding	- >25% binding by Farr assay or above normal range for testing laboratory.
1	<input type="checkbox"/>	Fever	- >38° C. Exclude infectious cause.
1	<input type="checkbox"/>	Thrombocytopenia	- <100,000 platelets/mm ³ .
1	<input type="checkbox"/>	Leukopenia	- <3,000 White blood cells/mm ³ . Exclude drug causes.
<p>— TOTAL SCORE (Sum of weights next to descriptors marked present)</p>			

Table 2

SLAM ASSESSMENT PAGE 1

VISIT: <input type="checkbox"/> Screening <input type="checkbox"/> Qualifying <input type="checkbox"/> 1 <input type="checkbox"/> 2 <input type="checkbox"/> 3 <input type="checkbox"/> Completion/Early Termination <input type="checkbox"/> Other WEEK: (0) (13) (26) (39) (52)					
Constitutional					
	ABSENT or NORMAL	MILD/MODERATE		SEVERE	NOT RECORDED
1. Weight Loss	<input type="checkbox"/> 0	<input type="checkbox"/> 1 < 10% body weight		<input type="checkbox"/> 3 > 10%	<input type="checkbox"/>
2. Fatigue	<input type="checkbox"/> 0	<input type="checkbox"/> 1 No limits on activity		<input type="checkbox"/> 3 Functional limitation	<input type="checkbox"/>
3. Fever	<input type="checkbox"/> 0	<input type="checkbox"/> 1 37.5 - 38.5 °C		<input type="checkbox"/> 3 > 38.5 °C	<input type="checkbox"/>
Integument					
	ABSENT	MILD	MODERATE	SEVERE	NOT RECORDED
4. Oral/nasal ulcers, or periungual erythema, malar rash, photosensitive rash, or nail fold infarct	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Present			<input type="checkbox"/>
5. Alopecia	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Hair loss with trauma	<input type="checkbox"/> 2 Spontaneous hair loss		<input type="checkbox"/>
6. Erythematous, maculopapular rash, discoid lupus, lupus profundus, or bullous lesions	<input type="checkbox"/> 0	<input type="checkbox"/> 1 < 20% total body surface (TBA)	<input type="checkbox"/> 2 20 - 50% TBA	<input type="checkbox"/> 3 > 50% TBA	<input type="checkbox"/>
7. Vasculitis (leucocytoclastic vasculitis, urticaria, palpable purpura, livedo reticularis, ulcer or panniculitis)	<input type="checkbox"/> 0	<input type="checkbox"/> 1 < 20% (TBA)	<input type="checkbox"/> 2 20 - 50% TBA	<input type="checkbox"/> 3 > 50% TBA or necrosis	<input type="checkbox"/>
Eye					
	ABSENT	MILD	MODERATE	SEVERE	NOT RECORDED
8. Cytoid bodies	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Present		<input type="checkbox"/> 3 Visual acuity < 20/200	<input type="checkbox"/>
9. Hemorrhage (retinal or choroidal) or episcleritis	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Present		<input type="checkbox"/> 3 Visual acuity < 20/200	<input type="checkbox"/>
10. Papillitis or pseudotumor cerebri	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Present		<input type="checkbox"/> 3 Visual acuity < 20/200 or field cut	<input type="checkbox"/>

SLAM ASSESSMENT PAGE 2

VISIT: <input type="checkbox"/> Screening <input type="checkbox"/> Qualifying <input type="checkbox"/> 1 <input type="checkbox"/> 2 <input type="checkbox"/> 3 <input type="checkbox"/> Completion/Early Termination <input type="checkbox"/> Other WEEK: (0) (13) (26) (39) (52)					
Reticuloendothelial					
	ABSENT or NORMAL	MILD	MODERATE	SEVERE	NOT RECORDED
11. Diffuse lymphadenopathy (cervical, axillary, epitrochlear)	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Shotty	<input type="checkbox"/> 2 > 1cm X 1.5cm		<input type="checkbox"/>
12. Hepato - or splenomegaly	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Palpable only with inspiration	<input type="checkbox"/> 2 Palpable without inspiration		<input type="checkbox"/>
Pulmonary					
	ABSENT or NORMAL	MILD	MODERATE	SEVERE	NOT RECORDED
13. Pleural effusion/pleurisy	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Shortness of breath or pain only with prompting. exam normal or near normal	<input type="checkbox"/> 2 Shortness of breath or pain with exercise, decreased breath sounds and dull lower lobe(s)	<input type="checkbox"/> 3 Shortness of breath or pain at rest, decreased breath sounds and dull middle and lower lobe(s)	<input type="checkbox"/>
14. Pneumonitis	<input type="checkbox"/> 0	<input type="checkbox"/> 1 X-ray infiltrates only	<input type="checkbox"/> 2 Shortness of breath with exercise	<input type="checkbox"/> 3 Shortness of breath at rest	<input type="checkbox"/>
Cardiovascular					
	ABSENT or NORMAL	MILD	MODERATE	SEVERE	NOT RECORDED
15. Raynaud's	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Present			<input type="checkbox"/>
16. Hypertension	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Diast. 90-105	<input type="checkbox"/> 2 Diast. 105-115	<input type="checkbox"/> 3 Diast. > 115	<input type="checkbox"/>
17. Carditis	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Pericarditis by EKG &/or RUB &/or effusion by echo; no sx	<input type="checkbox"/> 2 Chest pain or arrhythmia	<input type="checkbox"/> 3 Myocarditis with hemodynamic compromise &/or arrhythmia	<input type="checkbox"/>
Gastrointestinal					
	ABSENT or NORMAL	MILD	MODERATE	SEVERE	NOT RECORDED
18. Abdominal pain (Serositis, pancreatitis, ischemic bowel, etc.)	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Complaint	<input type="checkbox"/> 2 Limiting pain	<input type="checkbox"/> 3 Peritoneal signs/ascites	<input type="checkbox"/>
Neuromotor					
	ABSENT or NORMAL	MILD	MODERATE	SEVERE	NOT RECORDED
19. Stroke syndrome (includes mononeuritis multiplex, transient ischemic attack (TIA), reversible ischemic neurologic deficit (RIND) cerebrovascular accident (CVA) retinal vascular thrombosis)	<input type="checkbox"/> 0	<input type="checkbox"/> 1 Single TIA	<input type="checkbox"/> 2 Multiple TIA/RIND or mononeuritis multiplex or cranial neuropathy or chorea	<input type="checkbox"/> 3 CVA/myelitis, retinal vascular occlusion	<input type="checkbox"/>
20. Seizure	<input type="checkbox"/> 0	<input type="checkbox"/> 1 1-2/month	<input type="checkbox"/> 2 > 2/month	<input type="checkbox"/> 3 Status epilepticus	<input type="checkbox"/>

SLAM ASSESSMENT PAGE 3

VISIT: <input type="checkbox"/> Screening <input type="checkbox"/> Qualifying <input type="checkbox"/> 1 <input type="checkbox"/> 2 <input type="checkbox"/> 3 <input type="checkbox"/> Completion/Early Termination <input type="checkbox"/> Other WEEK: (0) (13) (26) (39) (52)					
Neuromotor (continued)					
	ABSENT or NORMAL	MILD	MODERATE	SEVERE	NOT RECORDED
21. Cortical dysfunction	<input checked="" type="checkbox"/> 0	<input checked="" type="checkbox"/> 1 Mild depression/ personality disorder or cognitive deficit	<input checked="" type="checkbox"/> 2 Δ in sensorium, severe depression, or limiting cognitive impairment	<input checked="" type="checkbox"/> 3 Psychosis, dementia, or coma	<input type="checkbox"/>
22. Headache (including migraine equivalents)	<input checked="" type="checkbox"/> 0	<input checked="" type="checkbox"/> 1 Symptoms or transient neuro deficit	<input checked="" type="checkbox"/> 2 Interferes somewhat with normal activities	<input checked="" type="checkbox"/> 3 Incapacitating/ aseptic meningitis	<input type="checkbox"/>
23. Myalgia/myositis	<input checked="" type="checkbox"/> 0	<input checked="" type="checkbox"/> 1 Complaint	<input checked="" type="checkbox"/> 2 Limits some activity	<input checked="" type="checkbox"/> 3 Incapacitating	<input type="checkbox"/>
Joints					
	ABSENT or NORMAL	MILD	MODERATE	SEVERE	NOT RECORDED
24. Joint pain from synovitis and/or tenosynovitis	<input checked="" type="checkbox"/> 0	<input checked="" type="checkbox"/> 1 Arthralgia only	<input checked="" type="checkbox"/> 2 Objective inflammation	<input checked="" type="checkbox"/> 3 Limited function	<input type="checkbox"/>
Laboratory					
	NORMAL	MILD	MODERATE	SEVERE	UNKNOWN NOT RECORDED
25. Hematocrit	<input checked="" type="checkbox"/> 0 > 35	<input checked="" type="checkbox"/> 1 30 - 35	<input checked="" type="checkbox"/> 2 25 - 29.9	<input checked="" type="checkbox"/> 3 < 25	<input type="checkbox"/>
26. WBC	<input checked="" type="checkbox"/> 0 > 3500	<input checked="" type="checkbox"/> 1 3500 - 2000	<input checked="" type="checkbox"/> 2 2000 - 1000	<input checked="" type="checkbox"/> 3 < 1000	<input type="checkbox"/>
27. Lymphocyte count	<input checked="" type="checkbox"/> 0 1500 - 4000	<input checked="" type="checkbox"/> 1 1499 - 1000	<input checked="" type="checkbox"/> 2 999 - 500	<input checked="" type="checkbox"/> 3 < 499	<input type="checkbox"/>
28. Platelet count	<input checked="" type="checkbox"/> 0 > 150T	<input checked="" type="checkbox"/> 1 100 - 150T	<input checked="" type="checkbox"/> 2 99 - 50T	<input checked="" type="checkbox"/> 3 < 50T	<input type="checkbox"/>
29. ESR (westergren)	<input checked="" type="checkbox"/> 0 < 25	<input checked="" type="checkbox"/> 1 25 - 50	<input checked="" type="checkbox"/> 2 51 - 75	<input checked="" type="checkbox"/> 3 > 75	<input type="checkbox"/>
30. Serum creatine or creatinine clearance	<input checked="" type="checkbox"/> 0 0.5 - 1.3mg/dl or 80 - 100% CrCl	<input checked="" type="checkbox"/> 1 1.4 - 2mg/dl or 79 - 60% CrCl	<input checked="" type="checkbox"/> 2 2.1 - 4mg/dl or 30 - 60% CrCl	<input checked="" type="checkbox"/> 3 > 4mg/dl or < 30% CrCl	<input type="checkbox"/>
31. Urine sediment	<input checked="" type="checkbox"/> 0 Urine protein < 150 mg/24 hours	<input checked="" type="checkbox"/> 1 > 5 RBC &/or WBC/hpf &/or 0 to 1-3 granular &/or cellular casts /hpf &/or 1-2+ proteinuria &/or < 500 mg/L 24* urine protein	<input checked="" type="checkbox"/> 2 > 10 RBC &/or WBC/hpf &/or > 3 granular &/or cellular casts/hpf &/or 3 or 4+ &/or 500 mg/L-3.5 g/L 24* urine protein	<input checked="" type="checkbox"/> 3 > 25 RBC or WBC/hpf &/or Red cell cast &/or > 4+ proteinuria &/or > 3.5 g/L 24* urine protein	<input type="checkbox"/>

Table 3

PATIENT SELF-ASSESSMENT QUESTIONNAIRE


VISIT: Screening Qualifying 1 2 3 Completion/Early Termination Other
 WEEK: (0) (13) (26) (39) (52)

SELF-ADMINISTERED BY THE PATIENT: We are interested in learning whether or not you are affected by fatigue because of your illness, as well as the overall effects of your illness on your general well-being.

1) Circle a number between 1 and 7 that indicates your degree of agreement with each of the statements below for the past week, where 1 indicates that you strongly disagree and 7 means that you strongly agree...

	DISAGREE					AGREE	
a) My motivation is lower when I am fatigued.	1	2	3	4	5	6	7
b) Exercise brings on my fatigue.	1	2	3	4	5	6	7
c) I am easily fatigued.	1	2	3	4	5	6	7
d) Fatigue interferes with my physical functioning.	1	2	3	4	5	6	7
e) Fatigue causes frequent problems for me.	1	2	3	4	5	6	7
f) My fatigue prevents sustained physical functioning.	1	2	3	4	5	6	7
g) Fatigue interferes with carrying out certain duties and responsibilities.	1	2	3	4	5	6	7
h) Fatigue is among my three most disabling symptoms.	1	2	3	4	5	6	7
i) Fatigue interferes with my work, family or social life.	1	2	3	4	5	6	7

2) Please indicate on the scale below, using a vertical line, how you have felt in the past week (including psychological and physical factors).



Score:

(for Genetec use only)

II. Clinical-Trial Data

A. Effect of DHEA Treatment on SLEDAI, SLAM, Patient VAS, and KFSS

5 Table 4 shows baseline characteristics of two patient groups employed in a clinical trial study, a 176-patient placebo group and a 170-patient treatment group. As seen, the two groups are comparable for all characteristics that were determined, including mean baseline SLEDAI, SLAM, Patient and Physician VAS, and fatigue score. The mean baseline scores represent the mean of two value determinations made prior to treatment.

10

Table 4
Baseline Comparability
(Per-protocol population)

Variable	Placebo (N=176)	GL701 (N=170)
Menopausal	86(49%)	74(44%)
Prednisone use	98(56%)	91(54%)
SLEDAI total>2	133(76%)	132(78%)
Race (white)	125(71%)	132(78%)
Smoking now	25(14%)	32(18%)
Age at screening (sd)	43.8(10.5)	44.1(11.1)
SLEDAI total (sd)	5.9(4.4)	6.5(4.3)
SLAM (sd)	12.0(2.9)	12.3(2.8)
Patient VAS (sd)	55.1(18.8)	55.2(18.6)
Physician VAS (sd)	30.6(13.5)	30.4(13.2)
Fatigue score (sd)	5.6(1.2)	5.5(1.2)
SLICC Score(sd)	1.3(1.5)	1.2(1.5)
SF36-Mental (sd)*	42.1(11.8)	43.3(10.4)
SF36-Physical (sd)*	31.3(8.4)	31.5(8.4)

N=174 and N=169 for placebo and GL170 groups, respectively

Each group, in turn, can be broken down into three subgroups: (I) total patient population (II) all patients with SLEDAI values greater than 2, and (III) patients having SLEDAI values greater than 2 and currently being treated with prednisone (at dose greater than 2 mg/prednisone /day), where the number of patients in each subgroup is indicated by N in the Table 5.

Table 5 shows the results of a clinical-trial in which the three placebo and three treatment subgroups were treated with DHEA over an extended period. Treatment was with 200 mg DHEA per day, administered orally in capsule form, or with a non-drug capsule (placebo). Mean duration of the treatment for each group was somewhat higher for the placebo group (308 days for placebo vs 288.4 days for DHEA, but median durations were almost the identical (362 vs. 359 days, respectively).

The three columns in the table represent total of the three subgroups for both the placebo and treatment group. The three rows represent responders in which an improvement (no change or decrease in value) in three the three of the four variables was observed and an increase in the fourth variable of no more than 3% (first row), 5% (second row), and 10% (third row) of the pretreatment baseline value.

Table 5
Window Analyses (3%, 5%, and 10%)

Variable	Group I: All patients			Group II: All patients with SLEDAI>2			Group III: All patients with SLEDAI>2, Pred>0		
	Placebo (N=176)	GL701 (N=170)	P value	Placebo (N=133)	GL701 (N=132)	P value	Placebo (N=80)	GL701 (N=73)	P value
			Improvement			Improvement			Improvement
Responders (3% window)*	57 32.4%	77 45.3%	**P=0.014	47 35.3%	72 54.5%	P=0.002	26 32.5%	39 53.4%	P=0.010
			39.8%			54.4%			64.3%
Responders (5% window)*	62 35.2%	79 46.5%	**P=0.034	50 37.6%	73 55.3%	P=0.004	26 32.5%	40 54.8%	P=0.006
			32.1%			47.1%			68.6%
Responders (10% window)*	69 39.2%	92 54.1%	**P=0.006	56 42.1%	84 63.6%	P=0.001	27 33.8%	45 61.6%	P=0.001
			38.0%			51.1%			82.2%

5 * baseline mean + 3%, 5%, or 10% of patients baseline mean

**P value not valid due to statistically significant treatment by SLEDAI interaction

As seen, patients in the treatment subgroups II (SLEDAI values greater than 2) and III (SLEDAI values greater than 2 and daily prednisone treatment) showed a greater than 50% responder rate, as defined by either a 3%, 5%, or 10% "increase" in the fourth variable, compared with a responder rates substantially less than 50% for the same two placebo subgroups.

Thus, in accordance with one aspect of the invention, the responder rate in DHEA treatment can be substantially improved, in both treatment by DHEA alone, or in combination with a second anti-SLE drug, such as prednisone, by (i) prescreening SLE patients for SLEDAI value, (ii) selecting for DHEA treatment, those patients with a SLEDAI value greater than 2.0, and (iii) treating the selected patient with a daily oral dose of DHEA.

In particular, this method can be practiced with a greater than 50% expectation of achieving improvement in the measured values of at least three of the disease-activity and constitutional-symptom variables characterizing a patient's SLE condition consisting of SLEDAI, KFSS, VAS, and SLAM, with an increase of no more than a 5% of a pretreatment baseline value in the fourth variable, where the changes in each variable are determined from the difference between a pretreatment baseline value and the mean of all values obtained at regularly scheduled intervals during treatment.

B. Effect of DHEA Treatment on SLE Flares

A clinical trial was conducted to determine the effect of DHEA treatment on SLE flares. An SLE flare is a significant new clinical manifestation of the disease (i.e., one that was not previously present in the patient or not previously as severe) and/or a clinical intervention. Table 6 identifies the clinical findings or interventions that were scored as flares in the present study. Any patient having at least one of these finding or interventions was scored as having a flare.

TABLE 6. SLE FLARES	
Type of Clinical Finding/Intervention	Definition of a Flare
New/worse CNS Lupus	Scored on SLEDAI & not present on previous visit.
Vasculitis	Scored on SLEDAI & not present on previous visit.
Myositis	Scored on SLEDAI & not present on previous visit.
Hematologic	Platelets < 60,000 or hemoglobin < 7mg/dL or decrease of at least 3 mg/dL.
Nephritis	Proteinuria with pyuria and/or hematuria treated with new/increased dose of corticosteroids or immunosuppressives.
Steroids	An increase of ≥ 2.5 mg for at least 7 days for SLE related reasons.
Immunosuppressives or anti-malarials	New use of or increase in dose for at least 7 days for SLE related reasons.
Hospitalization	Hospitalization for new SLE manifestation

Table 7 shows baseline characteristics of two patient groups employed in a clinical trial study including a 109-patient placebo group and a 189-patient treatment group. As seen, the two groups are comparable for all characteristics that were determined, including mean baseline SLEDAI, SLAM, Patient VAS (indicated as “Patient Global Assessment Score”), and KFSS. (This table shows the group mean baseline scores
5 determined prior to treatment.)

**TABLE 7. BASELINE CHARACTERISTICS OF FEMALE SLE PATIENTS
BY TREATMENT GROUP**

	PLACEBO (N=192)*	DHEA (N=189)*
Mean Age (yrs)	43.8	44.4
Caucasian (yes)	71.4%	77.2%
Post-Menopausal (yes)	47.9%	43.9%
Mean (Median) Prednisone Dose	3.7 (2.5) mg/d	3.5 (3.8) mg/d
Prednisone Use at Baseline (yes)	53.7%	54.5%
Immunosuppressive Use at Baseline (yes)	14.6%	16.9%
Anti-Malarial Use at Baseline (yes)	25.0%	23.3%
Mean (Median) SLEDAI† Score	5.8 (5.0)	6.5 (6.0)
Mean (Median) SLAM‡ Score	12.0 (12.0)	12.2 (12.0)
Mean (Median) Patient global assessment§ Score	55.4 (57.0)	55.2 (57.0)
Mean (Median) KFSS § Score	5.6 (5.7)	5.5 (5.9)
Mean (Median) DHEA-S	103(50) µg/dl	107(61) µg/dl
Mean (Median) C3 Complement	103.0 (102.0) mg/dl	102.8 (100.0) mg/dl
Mean (Median) C4 Complement	18.0 (16.0) mg/dl	17.9 (17.0) mg/dl
Mean (Median) Double-Stranded DNA	24.4 (1.9) IU/dl	34.8 (2.6) IU/dl
Antibody		

* Baseline values were not obtained on all patients for some clinical laboratory tests. For DHEA-S, N=163, placebo, and N=165, DHEA. For C3, C4 and double-stranded DNA antibody, N=178, placebo, and N=169, DHEA.

5 † Systemic Lupus Erythematosus Disease Activity Index

‡ Systemic Lupus Activity Measure

§ Krupp Fatigue Severity Scale

|| Dehydroepiandrosterone sulfate

10 To convert DHEA-S to µmol/liter, multiply by 0.027. To convert C3 and C4 complement to g/liter, multiply by 0.01.

Clinical trial results were analyzed for each group (placebo and DHEA-treated) as a whole (“per-protocol”), as well as broken down into three subgroups: all patients with SLEDAI values greater than 2 (“active SLE”), and patients having SLEDAI values greater than 2 and receiving corticosteroids and/or immunosuppressives (“more severe SLE”).

Table 8 shows the results of a clinical-trial in which the patients were treated with DHEA or placebo over an extended period. Patients received 200 mg DHEA per day, administered orally in capsule form, or with a non-drug capsule (placebo). Mean duration of the treatment for each group was somewhat higher for the placebo group (308.4 days for placebo vs 288.4 days for DHEA, but median durations were almost the identical (362 vs. 359 days, respectively.).

The first column of the table shows the patient population or sub-population analyzed, the next three columns show the results of the study for patients identified as responders according to the following criteria: (1) Weighted average change from baseline for Systemic Lupus Activity Measure (SLAM) <1; for Systemic Lupus Erythematosus Disease Activity Index (SLEDAI) <0.5; for Krupp Fatigue Severity Scale (KFSS) <0.5; for patient global assessment <10; and (2) no clinical deterioration. The last three columns of the table show the results of the study for patients experiencing at least one flare during the study.

The results indicate that the magnitude of the response to DHEA treatment increased with the severity of disease (i.e., from the less severe “active SLE” category to the “more severe” category to the “SLE” category in which patients were receiving corticosteroids and/or immunosuppressives). In addition, DHEA treatment reduced the occurrence of flares, and the magnitude of this effect also increased with the severity of disease.

The results of this trial are also shown in Fig. 1, which is a graph of the percentage of patients who did not experience a flare over the indicated duration of the study. Fig. 1 demonstrates that DHEA treatment reduces the risk of flare in “active SLE” patients, as compared to the placebo group, beginning at about 85 days of treatment. At about 200 days of treatment, incidence of flare is at least about 5% lower in the treated patients, and from this point on, the difference in flare incidence between the treated and placebo patients grows to at least about 10%.

TABLE 8. PERCENT RESPONDERS* AND PATIENTS WITH AT LEAST ONE DEFINITE FLARE

VARIABLE	RESPONDERS*			PATIENTS WITH AT LEAST ONE FLARE		
	Placebo	DHEA	P Value†	Placebo	DHEA	P value†
Population‡	45%	58%	0.018§	27%	22%	0.335
	(80/176)	(99/170)		(47/176)	(37/170)	
Active SLE	49%	66%	0.005	31%	24%	0.201
	(65/133)	(87/132)		(41/133)	(31/132)	
More Severe SLE¶	44%	64%	0.010	39%	26%	0.056
	(37/85)	(51/80)		(33/85)	(21/80)	

* A responder is defined as a patient who satisfies the following conditions: (1) Weighted average change from baseline for Systemic Lupus Activity Measure (SLAM) <1; for Systemic Lupus Erythematosus Disease Activity Index (SLEDAI) <0.5; for Krupp Fatigue Severity Scale (KFSS) <0.5; for patient global assessment <10; and (2) no clinical deterioration.

† P-value for responder is from a logistic regression analysis with treatment as a factor; P-value for flare is from a log-rank test for time to first definite flare.

‡ On study drug for > 60 days, had measurements of SLE scores or other data (pertinent to flare determination and clinical deterioration) beyond 60 days), and had no major protocol violations.

§ A significant treatment interaction with baseline SLEDAI (baseline SLEDAI >2 and ≤2) was noted, p=0.0003. Therefore, patients with baseline SLEDAI 0-2 should not be pooled with those with baseline SLEDAI >2 to determine treatment effect.

|| Per-protocol patients with SLEDAI >2 at baseline.

¶ Per-protocol patients with SLEDAI >2 and receiving corticosteroids and/or immunosuppressives at baseline

15

Thus, in accordance with one aspect of the invention, DHEA treatment can substantially reduce the risk of flare, especially in patients with a SLEDAI value greater than 2.0.

III. Treatment Method

The invention provides a method for treating SLE in an individual. The individual can be any animal that has SLE or an SLE-like condition. Generally, the individual is a mammal, and preferably a human SLE patient. Preferably, the individual has a SLEDAI value of greater than 2.0. According to the method, DHEA is administered to the individual and then the four disease-activity and constitutional symptom variables discussed above (SLEDAI, KFSS, VAS, AND SLAM) are determined and compared to baseline values determined before initiating DHEA administration. A decrease in three of these four variables and either a decrease, no change, or an increase of no more than about 5% of a baseline value in the fourth variable indicates that the individual is responding to said DHEA administration.

A. DHEA

In the SLE treatment method of the invention, an effective amount of a pharmaceutically active form of dehydroepiandrosterone (“DHEA”) is administered to an individual with SLE. As used herein, the term “pharmaceutically active form of DHEA” includes pharmaceutically active acid, salt, and ester forms of DHEA, such as DHEA sulfate (7 α -3H-DHEA sulfate, e.g., Heinz).

DHEA can be isolated in at least 6 different polymorphic forms, as described in detail in co-owned PCT Application No. PCT/US/00/06987 (International Publication No. WO 00/54763). DHEA was previously known, via analytical techniques such as x-ray diffraction, infrared (IR) spectroscopy, and differential scanning calorimetry (DSC), to occur in several different hydrate and anhydrate crystal forms. The anhydrate forms include forms I, II, III, IV and V, although the latter two forms have been observed only transiently by DSC. The hydrates (solvates) include forms S1 (1/4 hydrate), S2 (monohydrate), S3 (monohydrate), and S4 (1/2 methanolate). PCT Application No. PCT/US/00/06987 describes an additional form, form VI, which is detectable only by solid state NMR.

In preferred embodiments of the present invention, the DHEA employed has defined bioavailabilities and pharmacokinetic properties, which can be achieved by using preparations containing polymorphs that provide the desired properties.

In one aspect, the treatment method employs a DHEA preparation that is at least about 85%, preferably at least about 90%, more preferably at least about 95%, and most

preferably at least about 99% form I. The form I polymorph has the following characteristics:

(1) X-Ray Powder Diffraction unique peaks at 15.0 (s), 16.8 (w), 18.0 (m), 18.7 (m), 19.1 (w), 19.3 (w), 20.2 (w), 24.8 (w) 25.0 (w), 25.2 (w) (peak positions are given in degrees 2θ ; s = strong, m = medium, w = weak); and

(2) Solid State ^{13}C -NMR peaks: 14.8, 14.1 ppm carbon no. 18, 120.4, 118.9 ppm carbon no. 6, where these characteristics are measured as described in PCT Application No. PCT/US/00/06987.

DHEA form I-containing preparations exhibit good uptake by the GI tract upon oral administration, show good therapeutic activity, and are highly stable under ambient conditions.

In another aspect, the DHEA preparation is at least about 85%, preferably at least about 90%, more preferably at least about 95%, and most preferably at least about 99% form II. The form II polymorph has the following characteristics:

(1) X-Ray Powder Diffraction unique peaks at 8.6 (w), 17.3 (w), 20.9 (m), 22.0 (w), 22.2 (w), 27.1 (w) (peak positions are given in degrees 2θ ; s = strong, m = medium, w = weak); and

(2) Solid State ^{13}C -NMR peaks: 13.1 ppm carbon no. 18, 119.9 ppm carbon no. 6, where these characteristics are measured as described in PCT Application No. PCT/US/00/06987.

DHEA form II-containing preparations exhibit good uptake by the GI tract upon oral administration, a rapid rate of absorption (greater than the form I polymorph) and good therapeutic activity, and are also stable under ambient conditions.

Additionally, DHEA preparations useful in the treatment method can contain mixtures of the form I and II polymorphs. Generally, the combined form I and II polymorphs account for at least about 85%, preferably at least about 90%, more preferably at least about 95%, and most preferably at least about 99% of the DHEA in such preparations. Preparations enriched in form I and/or form II, as described herein, provide more predictable pharmacokinetic profiles than are provided by compositions having random polymorphic compositions.

Such compositions, including DHEA, and precursors such as DHEA acetate, are commercially available from various sources (e.g., Sigma Chemical Co., St. Louis, MO;

Aldrich Chemical Company, Inc.; Diosynth, Inc.; Pfaltz & Bauer, Inc.; Schering AG). DHEA compositions enriched for selected polymorphs can be prepared by crystallization of commercial DHEA in selected solvents under appropriate cooling or evaporation conditions.

In one preferred method, pure form I is prepared by (a) crystallizing DHEA from anhydrous 2-propanol (or, alternatively, acetone or acetonitrile) under a nitrogen stream at room temperature over about 2 days, producing a crystalline precipitate that contains predominantly form I and some amount of form VI, followed by (b) suspending the precipitate in ethyl acetate (about 100 mL/30 g of DHEA) and stirring the resulting slurry at room temperature for about one week, followed by filtration. The filter cake is allowed to dry at room temperature overnight. ¹³C-SSNMR analysis (discussed below) showed that product prepared by this method consisted of pure or nearly pure (>99%) form I; no other forms were detected by ¹³C-SSNMR.

DHEA highly enriched for form II can be obtained by rapid crystallization from tetrahydrofuran (THF), dioxane, chloroform or mixtures of chloroform and THF. Example 1 of PCT Application No. PCT/US/00/06987 provides a specific procedure for crystallization from THF, which produced a product shown by X-ray powder diffraction to be pure form II.

B. Formulation and Administration of DHEA

DHEA may be administered in a variety of ways, orally, including orally, parenterally, transcutaneously, transmucosally, or by inhalation, although oral administration is generally preferred.

Depending upon the manner of introduction, the DHEA may be formulated in a variety of ways. DHEA formulations can be prepared in various pharmaceutical forms, such as granules, tablets, capsules, suppositories, powders, controlled release formulations, suspensions, emulsions, creams, ointments, salves, lotions, or aerosols and the like.

Preferably, DHEA formulations are employed in solid dosage forms suitable for simple, and preferably oral, administration of precise dosages. Solid dosage forms for oral administration are preferably tablets, capsules, or the like.

DHEA formulations useful in the invention can include one or more pharmaceutical grade organic or inorganic carriers, excipients, and/or diluents, especially those suitable for oral or topical use. Such carriers include tocopherol, dimethyl sulfoxide,

and the like. For oral administration, suitable excipients include lactose, mannitol, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, gelatin, sucrose, magnesium carbonate, and the like.

To prepare orally deliverable tablets, DHEA is mixed with at least one
5 pharmaceutical excipient, and the solid formulation is compressed to form a tablet according to known methods, for delivery to the gastrointestinal tract. The tablet composition is typically formulated with additives, e.g. a saccharide or cellulose carrier, a binder such as starch paste or methyl cellulose, a filler, a disintegrator, or other additives typically usually used in the manufacture of medical preparations. To prepare orally deliverable capsules, DHEA is mixed
10 with at least one pharmaceutical excipient, and the solid formulation is placed in a capsular container suitable for delivery to the gastrointestinal tract.

Diluents known in the art include, for example, vegetable and animal oils and fats. Stabilizing agents, wetting and emulsifying agents, salts for varying the osmotic pressure, buffers for securing an adequate pH value, and/or skin penetration enhancers can
15 be used as auxiliary agents in the DHEA formulations. Methods for preparing various conventional dosage forms are known or will be apparent to those skilled in the art; for example, see *Remington's Pharmaceutical Sciences* (19th Ed., Williams & Wilkins, 1995).

The proportion of pharmaceutically active DHEA to carrier and/or other substances may vary from about 0.5 to about 100 wt.% (weight percent). For oral use, the
20 pharmaceutical formulation will generally contain from about 5 to about 100% by weight of the active material. For other uses, the formulation will generally have from about 0.5 to about 50 wt.% of the active material.

DHEA formulations employed in the invention provide an effective amount of DHEA upon administration to an individual. As used in this context, an "effective
25 amount" of DHEA is an amount that is effective to ameliorate a symptom of SLE. Such a therapeutic effect is generally observed within about 4 to about 6 weeks of initiating administration of an effective amount of DHEA.

The subject formulations are preferably, though not necessarily administered daily, in an amount to provide at least about a 10%, and more usually at least about 25%,
30 increase in the blood level of DHEA. Generally, the total daily dosage will be at least about 50 mg, preferably at least about 100 mg, and more preferably at least about 200 mg, and preferably not more than 500 mg per day, administered orally, e.g., in 4 capsules or tablets,

each containing 50 mg DHEA. Although capsules or tablets for oral delivery can conveniently contain up to a full daily oral dose, e.g., 200 mg or more. Where administration by other than an oral route, the DHEA may be delivered over an extended period, e.g., 3-10 days, in an amount effective to produce at least an average daily dose of, e.g., 50 mg.

DHEA treatment is carried out for an extended period, typically at least about 20, at least about 40, or at least about 60 weeks, and preferably as long as the patient is receiving noticeable benefit from the treatment method.

In preferred embodiments, DHEA is administered at a dose and for a period effective to produce a decrease in three of the four disease-activity and constitutional-symptom variables characterizing an individual's SLE condition and either a decrease, no change, or an increase of no more than about 3% of a pretreatment baseline value in the fourth variable. Preferably, the DHEA is administered at a dose effective to reduce the risk that an individual will experience an SLE flare at about 200 days of DHEA administration by at least about 5%. The risk of an SLE flare is said to be reduced by at least about 5% if at 200 days of DHEA treatment, the incidence of flare is at least about 5% lower in a DHEA-treated population, as compared to a placebo-treated population. The relevant populations are those in which the severity of disease is matched to the subject individual, i.e., the reduction in risk that an individual will experience a flare is defined in terms of DHEA- and placebo-treated populations that have the same severity of disease as the individual. For this purpose, the severity of disease is the same if the individual and two populations fall into one of the population groups identified in Table 8 above.

DHEA treatment can be combined with administration of one or more other drugs that are used in accordance with conventional SLE treatments, which include corticosteroids, such as glucocorticoids; non-steroidal anti-inflammatory agents; immunosuppressants; and anti-malarials. Examples of such drugs include hydroxychloroquine, prednisone, quinacrine, azathioprine, and immunosuppressants, such as anticytokines, including anti-TNF- α , TNF- α receptor antagonists, anti-IL-1, anti-IL-6, and anti-CD40 ligand. Dosages for the glucocorticoid prednisone, for example, are generally from about 1-15, more usually from about 1-12 mg/day, and typically more than 2 mg per day. The additional drugs may be administered separately or in conjunction with DHEA and may, if desired, be formulated in the same formulation with DHEA.

IV. Pharmaceutical Product

The invention also provides a pharmaceutical product for use in treating SLE in an individual including a plurality of doses of a pharmaceutically active form of DHEA, and instructions for performing the treatment method of the invention.

5 Specifically, the instructions direct that:

- 1) an effective amount of a pharmaceutically active form of DHEA be administered to an individual with SLE;
- 2) the following disease-activity and constitutional-symptom variables characterizing the individual's SLE condition be determined: SLEDAI, KFSS, VAS, and
10 SLAM at least about after initiating DHEA administration; and
- 3) the differences between the values for SLEDAI, KFSS, VAS, and SLAM after initiating DHEA administration and baseline values for SLEDAI, KFSS, VAS, and SLAM before initiating DHEA administration be determined, wherein a decrease in
15 three of these four variables and either a decrease, no change, or an increase of no more than about 5% of a baseline value in the fourth variable indicates that the individual is responding to said DHEA administration.

The pharmaceutically active DHEA can be formulated as described above with reference to the treatment method of the invention and can be packaged in any convenient manner.

20 Generally, the instructions direct the administration of DHEA as described above with reference to the treatment method. Oral administration is preferred. In a preferred embodiment, the instructions specify selecting those individuals having a SLEDAI value greater than 2.0 for DHEA administration.

The instructions can be affixed to the packaging material or can be included
25 as a package insert. While the instructions typically comprise written or printed materials they are not limited to such. Any medium capable of storing such instructions and communicating them to an end user is contemplated by this invention. Such media include, but are not limited to, electronic storage media (e.g., magnetic discs, tapes, cartridges, chips), optical media (e.g., CD ROM), and the like. As used herein, the term "instructions" can
30 include the address of an internet site that provides the instructions.

The invention also includes the use of the above-described pharmaceutical product for the treatment of SLE in a human patient.

* * *

All publications and patents cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

5 Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the scope of the appended claims.

CLAIMS

WHAT IS CLAIMED IS:

1. A method for treating systemic lupus erythematosus (SLE),
comprising:
 - 5 a) administering an effective amount of a pharmaceutically active form of DHEA to an individual with SLE;
 - b) at least about 4 weeks after initiating DHEA administration, determining the following disease-activity and constitutional-symptom variables characterizing the individual's SLE condition: SLEDAI, KFSS, VAS, and SLAM; and
 - 10 c) determining the differences between the values for SLEDAI, KFSS, VAS, and SLAM after initiating DHEA administration and baseline values for SLEDAI, KFSS, VAS, and SLAM before initiating DHEA administration, wherein a decrease in three of these four variables and either a decrease, no change, or an increase of no more than about 5% of a baseline value in the fourth variable indicates that the individual
15 is responding to said DHEA administration.
2. The method of claim 1, wherein the individual is a human SLE patient.
3. The method of claim 2, wherein the SLE patient has a SLEDAI value greater than 2.
- 20 4. The method of claim 3, wherein at least about 85% of the DHEA administered is present as the form I polymorph, the form II polymorph, or a combination thereof.
5. The method of claim 4, wherein at least about 95% of the DHEA administered is present as the form I polymorph, the form II polymorph, or a combination
25 thereof.
6. The method of claim 5, wherein at least about 95% of the DHEA administered is present as the form I polymorph.

7. The method of claim 5, wherein as least about 95% of the DHEA administered is present as the form II polymorph.

8. The method of claim 3, wherein the DHEA is administered at a dose and for a period effective to produce a decrease in three of the four disease-activity and constitutional-symptom variables characterizing an individual's SLE condition and either a
5 decrease, no change, or an increase of no more than about 3% of a pretreatment baseline value in the fourth variable.

9. The method of claim 3, wherein the DHEA is administered at a dose effective to reduce the risk of an SLE flare at about 200 days of DHEA administration by at
10 least about 5%.

10. The method of claim 3, wherein said administering comprises administering a daily oral dose of at least about 100 mg of a pharmaceutically active DHEA to the SLE patient.

11. The method of claim 10, wherein the dose is at least about 200 mg
15 DHEA/day, for a period of at least about 40 weeks.

12. The method of claim 10, wherein the SLE patient is receiving an orally administered drug selected from the group consisting of: a glucocorticoid, a non-steroidal anti-inflammatory agent, an immunosuppressant, and an anti-malarial drug prior to administration of DHEA, and said method includes continuing administration of said drug
20 during the period of DHEA administration.

13. The method of claim 12, wherein said drug is prednisone, at a daily dose of at least 2 mg.

14. A pharmaceutical product for use in treating systemic lupus erythematosus (SLE) in an individual, comprising:

- 25
- a) a plurality of doses of a pharmaceutically active form of DHEA, and
 - b) instructions directing that:

i) an effective amount of a pharmaceutically active form of DHEA be administered to an individual with SLE

ii) the following disease-activity and constitutional-symptom variables characterizing the individual's SLE condition be determined:
5 SLEDAI, KFSS, VAS, and SLAM at least about after initiating DHEA administration; and

ii) the differences between the values for SLEDAI, KFSS, VAS, and SLAM after initiating DHEA administration and baseline values for SLEDAI, KFSS, VAS, and SLAM before initiating DHEA administration be
10 determined, wherein a decrease in three of these four variables and either a decrease, no change, or an increase of no more than about 5% of a baseline value in the fourth variable indicates that the individual is responding to said DHEA administration.

15 15. The pharmaceutical product of claim 14, wherein the individual is a human SLE patient.

15 16. The pharmaceutical product of claim 15, wherein the instructions specify selecting those SLE patients having a SLEDAI value is greater than 2 for DHEA administration.

20 17. The pharmaceutical product of claim 16, wherein at least about 85% of the DHEA is present as the form I polymorph, the form II polymorph, or a combination thereof.

18. The pharmaceutical product of claim 17, wherein at least about 95% of the DHEA is present as the form I polymorph, the form II polymorph, or a combination thereof.

25 19. The pharmaceutical product of claim 18, wherein at least about 95% of the DHEA is present as the form I polymorph.

20. The pharmaceutical product of claim 18, wherein at least about 95% of the DHEA is present as the form II polymorph.

21. The pharmaceutical product of claim 14, wherein the doses of DHEA are capsules or tablets comprising 50 mgs. DHEA per capsule or tablet, respectively.

22. The pharmaceutical product of claim 14, wherein the instructions specify administering DHEA orally at a dose of at least about 100 mg DHEA per day.

5 23. The pharmaceutical product of claim 24, wherein the instructions specify administering DHEA orally at a dose of at least about 200 mg DHEA per day.

24. Use of the pharmaceutical product of claim 14, for treating systemic lupus erythematosus (SLE) in a human patient.




FIG. 1