

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 December 2010 (23.12.2010)

(10) International Publication Number
WO 2010/145796 A2

PCT

(51) International Patent Classification:
C12Q 1/68 (2006.01) *G01N 33/574* (2006.01)

(21) International Application Number:
PCT/EP2010/003563

(22) International Filing Date:
15 June 2010 (15.06.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09008042.5 19 June 2009 (19.06.2009) EP
09012197.1 25 September 2009 (25.09.2009) EP

(71) Applicant (for all designated States except US): **MERCK PATENT GMBH** [DE/DE]; Frankfurter Strasse 250, 64293 Darmstadt (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **STROH, Christopher** [DE/DE]; Waldstrasse 63D, 64807 Dieburg (DE).
VON HEYDEBRECK, Anja [DE/DE]; Frankfurter Strasse 66, 64293 Darmstadt (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: BIOMARKERS AND METHODS FOR DETERMINING EFFICACY OF ANTI-EGFR ANTIBODIES IN CANCER THERAPY

(57) Abstract: The invention relates to biomarkers based on gene expression products and methods for determining the efficacy of anti-EGFR antibodies in the treatment of EGFR expressing cancer. The invention is further related to the prediction of sensitivity or resistance of a patient suffering from EGFR expressing cancer to the treatment of said patient with a specific anti-EGFR antibody. The invention is preferably related to the identification of respective biomarkers that allow a better prediction of the clinical outcome of the treatment with anti-EGFR antibodies in patients with KRAS wild-type tumors. In this context, the invention especially relates to anti-EGFR antibody c225/cetuximab (Erbitux®) and its use in patients suffering from colorectal Cancer (CRC).



WO 2010/145796 A2

BIOMARKERS AND METHODS FOR DETERMINING EFFICACY OF ANTI-EGFR ANTIBODIES IN CANCER THERAPY

FIELD OF THE INVENTION:

5 The invention relates to biomarkers based on genes or gene expression products and methods for determining the efficacy of anti-EGFR antibodies in the treatment of EGFR expressing cancer. The invention is further related to the prediction of sensitivity or resistance of a patient suffering from EGFR expressing cancer to the treatment of said patient with a specific anti-EGFR antibody. The invention is preferably related to the identification of
10 respective biomarkers that allow a better prediction of the clinical outcome of the treatment with anti-EGFR antibodies in patients with KRAS wild-type tumors. In this context, the invention especially relates to anti-EGFR antibody c225/cetuximab (Erbix®) and its use in patients suffering especially from colorectal cancer (CRC).

15 BACKGROUND OF THE INVENTION

Monoclonal antibodies are commonly used in the treatment of cancer. Since antibodies are expensive drugs charging the national health care authorities, and often cause undesired side-effects, which provoke an additional burden and stress for the patient suffering from a severe and often terminal disease, it would be very desirable to know in advance whether
20 treating a specific patient with a specific antibody drug could really improve or even cure the patient's condition. There are already a couple of clinical and histological parameters which are used to obtain a prognosis whether a specific drug and / or treatment regimen may be successful for treating a disease. However, it has been shown that tumors very often elicit a diverse genetic pattern, which may change from one individual to another one. Thus, a
25 specific drug or a specific treatment, which may improve the clinical condition of a first patient, is less or not effective in a second patient suffering from the same cancer. For example, patients suffering from colorectal cancer may respond to a certain specific antibody drug differently. In the worst case a specific patient group does not respond at all to the drug, whereas another group of patients elicit a satisfying therapeutic response thereto dependent
30 on a different genetic tumor pattern and disposition.

Although modern molecular biology and biochemistry have revealed hundreds of genes whose activities influence the behavior of tumor cells, the state of their differentiation, and

their sensitivity or resistance to certain therapeutic drugs, such as antibodies, the status of these genes usually has not been exploited for the purpose of routinely making clinical decisions about drug treatments. Exception are the use of ErbB2 (Her2) protein expression in breast carcinomas to select patients with the Her2 antagonist drug Herceptin® (Genentech)

5 Another exception is the finding that mutations in the KRAS gene of EGFR expressing tumors are associated with the lack of sensitivity or response to anti-EGFR antibodies (Allegra et al. 2009, J Clin Oncol [Epub ahead of print])

New prognostic and predictive markers, which would facilitate a selection of patients for therapy, are needed to better predict patient response to treatments, such as small molecule

10 or biological molecule drugs, in the clinic. The classification of patient samples is a crucial aspect of cancer diagnosis and treatment. The association of a patient's response to a treatment with molecular and genetic markers can open up new opportunities for treatment development in non-responding patients, or distinguish a treatment's indication among other treatment choices because of higher confidence in the efficacy. Further, the pre-selection of

15 patients who are likely to respond well to a drug, or a combination of drugs or a specific regimen may reduce the number of patients needed in a clinical study or accelerate the time needed to complete a clinical development program.

The epidermal growth factor receptor (EGFR) and its downstream signaling effectors, notably members of the Ras/Raf/MAP kinase pathway, play an important role in both normal and

20 malignant epithelial cell biology (Normanno et al., Gene 366, 2-16 (2006)) and have therefore become established targets for therapeutic development.

Several monoclonal chimeric, humanized or fully human monoclonal antibodies have been developed which recognize and inhibit the EGFR. Two examples of antibodies, which have been marketed in meantime are cetuximab (ERBITUX®) and panitumumab (VECTIBIX®)

25 The anti-EGFR antibody c225 (cetuximab), a chimeric IgG1, which was demonstrated to inhibit EGF-mediated tumor cell growth in vitro and to inhibit human colorectal tumor growth in vivo, received marked approval in 2003. Its sequence was first disclosed in WO 96/40210.

The antibody as well as in general all anti-EGFR antibodies, appear to act, above all, in synergy with certain chemotherapeutic agents (i.e., doxorubicin, adriamycin, taxol, and

30 cisplatin) to eradicate human tumors in vivo in xenograft mouse models (e.g.

EP 0667165). Furthermore, it could be shown that the combination of the anti-EGFR antibody c225 with a second humanized anti-EGFR antibody matuzumab (Mab h425) shows a

synergistic effect in vitro models, indicating that these two antibodies although directed to the same receptor bind to different epitopes on the receptor (WO 2004/032960).

It was shown in the past that about 75% of patients treated with anti-EGFR antibodies, including cetuximab, or respective small molecule inhibitors develop more or less severe skin rash very fast after starting treatment. Although frequently tolerable and manageable, approximately 10% of patients require dose interruption and/or dose reduction due to the severe symptoms, and a few patients discontinue therapy. The increasingly apparent association of cutaneous toxicity with favorable clinical outcomes to EGFR inhibitors makes "achievement" of rash a desirable but potentially troublesome toxicity in patients experiencing therapeutic benefit, sometimes limiting the utility of these agents. Nonetheless, the occurrence of skin rash during anti-EGFR antibody treatment can be taken as reliable surrogate marker for therapeutic response to the treatment with said antibody, for example, cetuximab.

However, selecting skin rash occurrence as surrogate marker is not optimal because identification of cancer patients that generally do not respond to the treatment with an anti-EGFR antibodies is possible not before having administered the drug to the patient for a longer time.

Therefore, the finding that mutation of the KRAS gene (codon 12/13) and gene product in EGFR expressing tumor is responsible for insensitivity to the treatment of metastatic colorectal cancer with EGFR inhibitors can be regarded as improvement in the prediction whether a treatment is successful or not. WO 2008/112269 reports that panitumumab, a human Anti-EGFR antibody is effective only in KRAS wild-type tumor patients. Khambata-Ford et al. (2007, J. Clin. Oncol. 25, 3230) describe that metastatic colorectal cancer patients with tumors that have high gene expression levels of epiregulin and amphiregulin, as well as patients with wild-type KRAS tumors are more likely to have disease control on cetuximab treatment.

Despite above-said recent advances, a major challenge in cancer treatment remains to select patients for specific treatment regimens based on pathogenetic and / or genetic markers in order to optimize outcome. In context of treating EGFR expressing tumors with anti-EGFR antibodies inhibiting growth of these tumors, it would be helpful to know and better understand which patients are able to respond to an intended treatment with these antibodies, especially in view of newer findings that even in the KRAS-wild type tumor group

not all patients (ca. 40%) do respond well to the treatment of anti-EGFR antibodies and other EGFR inhibitors.

Therefore, a need exists for diagnostic tests, methods and tools using biomarkers that simultaneously can provide predictive information about patient responses to the variety of treatment options including personally different tumor genotypes.

SUMMARY OF THE INVENTION

The invention discloses predictive biomarkers for determining the efficacy of an anti-EGFR antibody in the treatment of cancer.

10 In one embodiment of the invention specific biomarkers are described which can be used to predict before administration in a patient the efficacy of an anti-EGFR antibody in the treatment of tumors in KRAS wild-type patients or patients having a mutation in the KRAS gene.

15 In a further embodiment of the invention specific biomarkers are disclosed which can be used to predict more exactly the efficacy or the degree of efficacy of an anti-EGFR antibody in the treatment of tumors in patients having no mutation in the KRAS gene (KRAS wild type), which usually respond statistically but not necessarily individually positive to an anti-EGFR antibody therapy.

20 Another embodiment of the invention is related to biomarkers which indicate a high likelihood of a good (positive biomarkers) or a bad (negative biomarkers) response to an anti-EGFR antibody treatment in patients suffering from colorectal cancer (CRC), preferably metastatic colorectal cancer (mCRC), squamous cell head and neck cancer (SCCHN) or non-small cell lung cancer (NSCLC).

25 In a specific embodiment of the invention biomarkers are disclosed which are predictive for the efficacy or non-efficacy of a tumor related (solid or metastatic tumors) therapy with the anti-EGFR antibody cetuximab (Erbix®).

In a preferred embodiment of the invention are disclosed which are predictive for the efficacy or non-efficacy of a tumor related (solid or metastatic tumors) therapy with the anti-EGFR antibody cetuximab (Erbix®), wherein the tumor is CRC, mCRC, SCCHN or NSCLC.

In a preferred embodiment of the invention are disclosed which are predictive for the efficacy or non-efficacy of a tumor related (solid or metastatic tumors) therapy with the anti-EGFR antibody cetuximab (Erbix®), wherein the tumor is CRC, mCRC, SCCHN or NSCLC, and wherein the patients suffering from said cancers preferably show an individual KRAS wild-type gene pattern.

In another embodiment the invention relates to an *in vitro* method for predicting by diagnostic means and / or diagnostic apparatus the likelihood that a patient suffering from KRAS wild type EGFR expressing tumor, who is a candidate for treatment with an EGFR antibody, will respond or not respond to the treatment with said anti-EGFR antibody.

- 10 According to the invention the method comprises determining the expression level of one or more prognostic genes or gene expression products thereof in a tissue sample obtained from said patient, wherein high or low expression of the gene/gene product compared to a clinical relevant reference value indicates that the patient is likely to respond to said treatment or is likely not to respond to the treatment.
- 15 Genes or gene products causing high expression in a sample of a tumor patient that has a high likelihood to respond or do respond to the treatment with an anti-EGFR antibody, preferably cetuximab are according to the invention selected from the group of genes consisting of: ADAMDEC1, BSDC1, C1orf144, CAPZB, CDC42, DHCR7, DNAJC8, ECSIT, EXOSC10, FADS1, GBA, GLT25D1, GOSR2, IMPDH1, KLHL21, KPNA6, KPNB1, LSM12,
20 MAN1B1, MIDN, PPAN, SH3BP2, SQLE, SSH3, TNFRSF1B, URM1, ZFYVE26, RGMB, SPIRE2, ABCC5, ACSL5, AOA1, AXIN2, CD24, CEACAM5, CEACAM6, ETS2, FMNL2, GPSM2, HDAC2, JUN, ME3, MED17, MYB, MYC, NEBL, NOSIP, PITX2, POF1B, PPARG, PPP1R14C, PRR15, PSMG1, RAB15, RAB40B, RNF43, RPS23, SLC44A3, SOX4, THEM2, VAV3, ZNF337, EPDR1, KCNK5, KHDRBS3, PGM2L1, STK38, and SHROOM2.
- 25 Genes or gene products causing high expression in a sample of a tumor patient that has a low likelihood to respond or do not respond to the treatment with an anti-EGFR antibody, preferably cetuximab, are according to the invention selected from the group of genes consisting of: C7orf46, CAST, DCP2, DIP2B, ERAP1, INSIG2, KIF21A, KLK6, NGRN, NRIP1, PHGDH, PPP1R9A, QPCT, RABEP1, RPA1, RPL22L1, SKP1, SLC25A27,
30 SLC25A46, SOCS6, TPD52, ZDHHC2, ZNF654, ASB6, ATM, BMI1, CDC42EP2, EDEM3, PLLP, RALBP1, SLC4A11, TNFSF15, TPK1, C11orf9, C1QC, CABLES1, CDK6, EHBP1, EXOC6, EXT1, FLRT3, GCNT2, MTHFS, PIK3AP1, ST3GAL1, TK2, ZDHHC14, ARFGAP3, AXUD1, CAPZB, CHSY1, DNAJB9, GOLT1B, HSPA5, LEPROTL1, LIMS1, MAPK6, MYO6,

PROSC, RAB8B, RAP2B, RWDD2B, SERTAD2, SOCS5, TERF2IP, TIAL1, TIPARP, TRIM8, TSC22D2, TGFA, VAPA, and UBE2K, indicates that the patient is likely not to respond to said treatment compared to a reference value.

5 In another embodiment of the invention preferred biomarkers which are above-average expressed and indicate that the patient will likely respond or will likely respond superior to the treatment with the respective anti-EGFR antibody (e.g. cetuximab), are selected from the group consisting of EPDR1, KCNK5, KHDRBS3, PGM2L1, SHROOM2, STK38, RGMB, SPIRE2 and VAV3 or the respective gene expression products of said group.

10 In another embodiment of the invention preferred biomarkers which are above-average expressed and indicate that the patient will not likely respond or will not likely respond superior to the treatment with the respective anti-EGFR antibody (e.g. cetuximab), are selected from the group consisting of ASB6, ATM, BMI1, CDC42EP2, EDEM3, PLLP, RALBP1, SLC4A11, TNFSF15, TPK1, ARFGAP3, AXUD1, CAPZB, CHSY1, DNAJB9, GOLT1B, HSPA5, LEPROTL1, LIMS1, MAPK6, MYO6, PROSC, RAB8B, RAP2B, RWDD2B, 15 SERTAD2, SOCS5, TERF2IP, TIAL1, TIPARP, TRIM8, TSC22D2, TGFA, VAPA, and UBE2K or the respective gene expression products of said group.

In a further embodiment the preferred biomarker according to the invention predictive for a positive response of the patient to an anti-EGFR antibody is VAV3 and the preferred biomarker according to the invention predictive for a negative or negligible response of the 20 patient to an anti-EGFR antibody is TGFA.

In another embodiment a respective method is applied, wherein at least one considerably or highly expressed first biomarker as specified above and in the claims is used which indicates that the patient will probably respond well or extraordinary or superior to the treatment with the anti-EGFR antibody preferably cetuximab (compared to a clinical average or standard 25 response and / or expression value calculated from a respective average patient cohort), and at least one considerably or highly expressed second biomarker as specified above and in the claims is used which indicates that the patient probably will not respond well or extraordinary or superior to the treatment with the anti-EGFR antibody preferably cetuximab (compared to a clinical average or standard response and / or expression value calculated from a respective 30 average patient cohort).

In another embodiment a respective method is applied for an *in vitro* method for predicting the likelihood that a patient suffering from KRAS wild type EGFR expressing tumor and is a candidate for treatment with an EGFR antibody, will respond to the treatment with said anti-

EGFR antibody, wherein expression levels of one or more of the above and below specified biomarkers are determined in combination with a AREG and / or EREG in context with the treatment of a tumor patient with an anti-EGFR antibody, preferably cetuximab.

Preferably, a respective method is applied, wherein the gene or gene product expression
5 levels of VAV3 and ARAG or EREG are determined in context with the treatment of a tumor patient, preferably a CRC or mCRC tumor patient, with an anti-EGFR antibody, preferably cetuximab.

In a further specific embodiment, a respective method is applied, wherein gene or gene
product expression levels of VAV3 and ARAG or EREG are determined in context with the
10 treatment of a tumor patient, preferably a CRC or mCRC tumor patient, with an anti-EGFR antibody, preferably cetuximab.

In another preferred embodiment according to the invention, a respective method is applied,
wherein gene or gene product expression levels TGFa and ARAG or EREG are determined in
context with the treatment of a tumor patient, preferably a CRC or mCRC tumor patient, with
15 an anti-EGFR antibody, preferably cetuximab.

In another preferred embodiment according to the invention, a respective method is applied,
wherein gene or gene product expression levels VAV3 and TGFa determined in context with
the treatment of a tumor patient, preferably a CRC or mCRC tumor patient, with an anti-
EGFR antibody, preferably cetuximab.

20 In another preferred embodiment according to the invention, a respective method is applied,
wherein gene or gene product expression levels VAV3, TGFa and ARAG or EREG are
determined in context with the treatment of a tumor patient, preferably a CRC or mCRC tumor
patient, with an anti-EGFR antibody, preferably cetuximab.

In a further aspect, the invention relates to an *in vitro* method for predicting the likelihood that
25 a patient suffering from KRAS wild type EGFR expressing cancer will respond therapeutically
to the treatment with an anti-EGFR antibody, preferably cetuximab, the method comprises:
(a) measuring by diagnostic means and / or diagnostic apparatus in a biopsy tissue sample
from tumor tissue or plasma of said patient the expression level of one or more biomarkers
selected from the group (i) consisting of ADAMDEC1, BSDC1, C1orf144, CAPZB, CDC42,
30 DHCR7, DNAJC8, ECSIT, EXOSC10, FADS1, GBA, GLT25D1, GOSR2, IMPDH1, KLHL21,
KPNA6, KPNB1, LSM12, MAN1B1, MIDN, PPAN, SH3BP2, SQLE, SSH3, TNFRSF1B,
URM1, ZFYVE26, RGMB, SPIRE2, ABCC5, ACSL5, AOA, AXIN2, CD24, CEACAM5,

CEACAM6, ETS2, FMNL2, GPSM2, HDAC2, JUN, ME3, MED17, MYB, MYC, NEBL, NOSIP, PITX2, POF1B, PPARG, PPP1R14C, PRR15, PSMG1, RAB15, RAB40B, RNF43, RPS23, SLC44A3, SOX4, THEM2, VAV3, ZNF337, EPDR1, KCNK5, KHDRBS3, PGM2L1, STK38, SHROOM2, and / or from group (ii) consisting of C7orf46, CAST, DCP2, DIP2B, ERAP1,
 5 INSIG2, KIF21A, KLK6, NGRN, NRIP1, PHGDH, PPP1R9A, QPCT, RABEP1, RPA1, RPL22L1, SKP1, SLC25A27, SLC25A46, SOCS6, TPD52, ZDHHC2, ZNF654, ASB6, ATM, BMI1, CDC42EP2, EDEM3, PLLP, RALBP1, SLC4A11, TNFSF15, TPK1, C11orf9, C1QC, CABLES1, CDK6, EHBP1, EXOC6, EXT1, FLRT3, GCNT2, MTHFS, PIK3AP1, ST3GAL1, TK2, ZDHHC14, ARFGAP3, AXUD1, CAPZB, CHSY1, DNAJB9, GOLT1B,
 10 HSPA5, LEPROTL1, LIMS1, MAPK6, MYO6, PROSC, RAB8B, RAP2B, RWDD2B, SERTAD2, SOCS5, TERF2IP, TIAL1, TIPARP, TRIM8, TSC22D2, TGFA, VAPA, and UBE2K,

(b) exposing ex-vivo a tissue sample from tumor or plasma of said patient to said anti-EGFR antibody, (c) measuring again in said exposed tissue sample of step (b) the expression level
 15 of one or more biomarkers specified in step (a), and (d) calculating the differences in expression levels measured in steps (b) and (c), wherein an increase in the expression level of the biomarkers of group (i) obtained in step (c) compared to step (a) indicates an increased likelihood that said patient responds therapeutically to the treatment with said anti-EGFR antibody, and wherein an increase in the
 20 expression level of the biomarkers of group (ii) obtained in step (c) compared to step (a) indicates a decreased likelihood that said patient responds therapeutically to the treatment with said anti-EGFR antibody.

In another aspect, the invention relates to a respective *in vitro* method as disclosed above and below, wherein the patient does not only suffer from KRAS wild type EGFR expressing
 25 tumor but in addition shows a mutation in the EGFR gene of the tumor tissue. In a specific embodiment this mutation is responsible for skin rash associated with the administration of the anti-EGFR antibody, preferably cetuximab. This mutation causes preferably a R521K polymorphism in EGFR.

In another aspect, the invention relates to a respective *in vitro* method as disclosed above
 30 and below, wherein the patient does not only suffer from KRAS wild type EGFR expressing tumor, especially CRC / mCRC, but in addition shows a mutation in the BRAF gene of the tumor tissue.

In a further aspect, the invention relates to DNA or RNA array comprising an arrangement of polynucleotides presented or hybridizing to one or more of the following genes:

- ADAMDEC1, BSDC1, C1orf144, CAPZB, CDC42, DHCR7, DNAJC8, ECSIT, EXOSC10, FADS1, GBA, GLT25D1, GOSR2, IMPDH1, KLHL21, KPNA6, KPNB1, LSM12, MAN1B1, MIDN, PPAN, SH3BP2, SQLE, SSH3, TNFRSF1B, URM1, ZFYVE26, RGMB, SPIRE2, ABCC5, ACSL5, AOA1, AXIN2, CD24, CEACAM5, CEACAM6, ETS2, FMNL2, GPSM2, HDAC2, JUN, ME3, MED17, MYB, MYC, NEBL, NOSIP, PITX2, POF1B, PPARG, PPP1R14C, PRR15, PSMG1, RAB15, RAB40B, RNF43, RPS23, SLC44A3, SOX4, THEM2, VAV3, ZNF337, EPDR1, KCNK5, KHDRBS3, PGM2L1, STK38, SHROOM2, C7orf46, CAST, DCP2, DIP2B, ERAP1, INSIG2, KIF21A, KLK6, NGRN, NRIP1, PHGDH, PPP1R9A, QPCT, RABEP1, RPA1, RPL22L1, SKP1, SLC25A27, SLC25A46, SOCS6, TPD52, ZDHHC2, ZNF654, ASB6, ATM, BMI1, CDC42EP2, EDEM3, PLLP, RALBP1, SLC4A11, TNFSF15, TPK1, C11orf9, C1QC, CABLES1, CDK6, EHBP1, EXOC6, EXT1, FLRT3, GCNT2, MTHFS, PIK3AP1, ST3GAL1, TK2, ZDHHC14, ARFGAP3, AXUD1, CAPZB, CHSY1, DNAJB9, GOLT1B, HSPA5, LEPROTL1, LIMS1, MAPK6, MYO6, PROSC, RAB8B, RAP2B, RWDD2B, SERTAD2, SOCS5, TERF2IP, TIAL1, TIPARP, TRIM8, TSC22D2, TGFA, VAPA, and UBE2K, wherein the gene or gene products are immobilized on a solid surface.

- In a preferred embodiment the DNA or RNA array comprises one or more of the following genes or hybridizes to said genes: TNFRSF1B, DNAJC8, ECSIT, GOSR2, PPP1R9A, KLK6, C7orf46, SSH3, SERTAD2, AXIN2, C10orf99, ETS2, PITX2, PRR15, VAV3, gene coding for IKK interacting protein, EDEM3, LY6G6D, and optionally in addition AREG and / or EREG and / or TGFA.

- In another preferred embodiment the DNA or RNA array, according to the invention, is consisting of the following arrangement of polynucleotides presented by or hybridizing to the following genes

- (a) TNFRSF1B, DNAJC8, ECSIT, GOSR2, PPP1R9A, KLK6, C7orf46, SSH3, SERTAD2, AXIN2, C10orf99, ETS2, PITX2, PRR15, VAV3, gene coding for IKK interacting protein, EDEM3, LY6G6D, AREG, EREG and TGFA; or
- (b) TNFRSF1B, DNAJC8, VAV3, AREG or EREG, TNFa; or
- (c) VAV3; AREG or EREG, TNFa; or
- (d) VAV3, TNFa; or
- (e) AREG or EREG, TNFa.

SHORT DESCRIPTION OF THE FIGURES:

Fig. 1: Genes whose expression in baseline samples is significantly associated with disease control after six weeks of cetuximab monotherapy in patients with wildtype KRAS gene ($p < 0.002$, moderated t-test). Based on study EMR 62202-045 (first-line treatment of metastatic CRC)

Fig. 2: Genes whose expression in baseline samples is significantly associated with disease control after six weeks of cetuximab monotherapy ($p < 0.002$, moderated t-test). Based on study EMR 62202-045 (first-line treatment of metastatic CRC).

Fig. 3: Study EMR 62202-502 (cetuximab plus irinotecan treatment of irinotecan-refractory metastatic CRC patients), analysis of patients with wildtype KRAS gene: genes whose expression in baseline samples is significantly associated with best overall response ($p < 0.002$, Welch t-test)

Fig. 4: Study EMR 62202-502 (cetuximab plus irinotecan treatment of irinotecan-refractory metastatic CRC patients): genes whose expression in baseline samples is significantly associated with best overall response ($p < 0.002$, Welch t-test).

Fig. 5: Study EMR 62202-502 (cetuximab plus irinotecan treatment in irinotecan-refractory metastatic CRC patients), analysis of patients with wildtype KRAS gene: genes whose expression in baseline samples is significantly associated with overall survival time ($p < 0.002$, Cox proportional hazards regression)

Fig. 6: Study EMR 62202-502 (cetuximab plus irinotecan treatment in irinotecan-refractory metastatic CRC patients): genes whose expression in baseline samples is significantly associated with overall survival time ($p < 0.002$, Cox proportional hazards regression)

Fig. 7: Study EMR 62202-502 (cetuximab plus irinotecan treatment in irinotecan-refractory metastatic CRC patients): genes whose expression in baseline samples is significantly associated with progression-free survival time ($p < 0.002$, Cox proportional hazards regression).

Fig. 8: Study EMR 62202-502 (cetuximab plus irinotecan treatment in irinotecan-refractory metastatic CRC patients), analysis of patients with wildtype KRAS gene: genes whose expression in baseline samples is significantly associated with progression-free survival time ($p < 0.002$, Cox proportional hazards regression)

Fig. 9: Affymetrix probe sets used to evaluate the degree of liver tissue contamination of tumor biopsies.

Fig. 10: Association of baseline expression data with disease control at week 6 in patients with *KRAS* wild-type tumors: 57 probe sets with $P < .002$. Log-ratio values are mean $\log_2$ expression levels of patients with disease control minus those of patients with progressive disease, adjusted for the degree of liver contamination of the samples.

Fig. 11: Forty-seven probe sets showing an association of on-treatment changes in expression from baseline to week 4 with best overall response. Log-ratio values represent means of on-treatment changes of patients with partial response minus those of patients with stable or progressive disease, adjusted for the degree of liver contamination of the samples.

Fig. 12: Association of on-treatment changes in candidate gene expression from baseline to week 4 with best overall response. Log-ratio values represent means of on-treatment changes of patients with partial response minus those of patients with stable or progressive disease, adjusted for the degree of liver contamination of the samples.

Fig. 13: Association of baseline expression of candidate genes with disease control at week 6 in tumors with *KRAS* wild-type status. Log-ratio values are mean $\log_2$ expression levels of patients with disease control minus those of patients with progressive disease, adjusted for the degree of liver contamination of the samples.

Fig. 14: Results of the statistical analysis of Luminex plasma proteomics data.

Analyses refer to general changes between baseline and week 4 samples, associations of on-treatment changes with response at week 6 among all patients, as well as among the patients with *KRAS* wild-type tumors. For each of these analyses, $\log_2$ -ratio values, p-values and q-values are given for each measured protein. Log2-ratio values refer to the mean difference of $\log_2$ concentrations between week 4 and baseline samples (general change), or to the difference between these mean differences between responders and non-responders (association with response).

Fig. 15: Antibody reagents and immunohistochemistry assay conditions.

Fig. 16: Identification of RNA samples with high (green), medium (red) and low (black) liver contamination based on expression of genes predominantly expressed in colorectal cancer (blue box) and normal liver (purple box). The color scale reflects the absolute element signal intensity after normalization.

Fig. 17: Association of on-treatment changes in expression from baseline to week 4 with best overall response (partial response vs stable disease plus progressive disease). Forty-seven probe sets with $P < 0.002$ are shown. Gene names, followed by the element ID are given on the right of the image. The element intensity represents the $\log_2$ ratio of gene expression at week 4 over gene expression at baseline.

Abbreviations; PD, progressive disease; SD, stable disease; PR, partial response.

Fig. 18: Association of on-treatment changes in candidate gene expression from baseline to week 4 with best overall response (partial response vs stable disease plus progressive disease). Gene names, followed by the element ID are given on the right of the image. The element intensity represents the $\log_2$ ratio of gene expression at week 4 over gene expression at baseline.

Abbreviations; PD, progressive disease; SD, stable disease; PR, partial response.

Fig. 19: Association of baseline expression of candidate genes with disease control (partial response plus stable disease vs progressive disease) in tumors with *KRAS* wild-type status. Gene names, followed by the element ID are given on the right of the image. The intensity scale reflects the $\log_2$ ratio for each element, relative to the mean for each probe set across all samples. Abbreviations; PD, progressive disease; SD, stable disease; PR, partial response.

Fig. 20: Correlation of *KRAS* status with response rate and progression-free survival in mCRC patients treated with cetuximab (Erbixux)

Fig 21. Immunohistochemical analysis of the expression of selected EGFR signaling pathway associated markers in skin (A) and tumor (B) samples: changes between paired week4/baseline samples.

Fig. 22. Proportion of patients free of disease progression versus progression-free survival time in months according to *KRAS* tumor mutation status.

Fig. 23. Association of baseline gene expression data with disease control (partial response plus stable disease vs progressive disease) at week 6 in patients with *KRAS* wild-type tumors. 57 probe sets with $P < .002$ are shown. Gene names, followed by the element ID are given on the right of the image. The color scale reflects the $\log_2$ ratio for each element, relative to the mean for each probe set across all samples.

Abbreviations; PD, progressive disease; SD, stable disease; PR, partial response.

Fig.24 . *AREG* (element 205239_at), *EREG* (element 205767_at) and *TGFA* (element 205016_at) expression levels in baseline samples according to response at week 6 in all patients (Panel A, C, and E, respectively) and in patients whose tumors were wild-type for *KRAS* (Panels B, D, and F, respectively). *P*-values refer to the association with disease control (partial response, PR and stable disease, SD versus progressive disease, PD).

Fig. 25. Association of on-treatment changes in plasma protein concentrations from baseline to week 4 with response at week 6 (partial response, PR vs stable disease, SD plus progressive disease, PD) among 45 patients in the intention to treat (ITT) population (Panel A) and among 24 ITT patients with *KRAS* wild-type tumors (Panel B). All proteins with $P < .01$ are shown. The element intensity represents the $\log_2$ ratio of protein concentration at week 4 over protein concentration at baseline.

Fig. 26: Boxplots showing the association of VAV3 with response. PD: progressive disease, PR: partial response, SD: stable disease. Green dots: Patients with *KRAS* and *BRAF* wild-type tumors, red dots: patients with *KRAS* mutations, black dots: patients with *BRAF* mutations, blue dots: mutation status unknown. *P*-values are based on Welch t-tests.

Fig. 27: Kaplan-Meier plot showing estimated progression-free survival distribution functions stratified by VAV3 expression. Patients have been classified as high or low VAV3 expressors, depending on whether their baseline VAV3 expression levels were above or below the median level across patients, respectively. The *p*-value is derived from a Cox proportional hazards model.

Fig. 28: Kaplan-Meier plot showing estimated overall survival distribution functions stratified by VAV3 expression. Patients have been classified as high or low VAV3 expressors, depending on whether their baseline VAV3 expression levels were above or below the median level across patients, respectively. The *p*-value is derived from a Cox proportional hazards model.

Fig. 29: Kaplan-Meier plot showing estimated progression-free survival distribution functions stratified by VAV3 expression and *KRAS* mutation status. Patients have been grouped into four strata, representing all possible combinations of *KRAS* mutation status and baseline VAV3 expression (above or below the median).

Fig. 30: Vav3 interacts with activated EGFR. After transfection of HEK 293 cells with VAV3 and EGFR alone or in combination, cells were lysed and subjected to immunoprecipitation (IP) and Western blotting (WB).

DETAILED DESCRIPTION OF THE INVENTION

The EGFR-targeting immunoglobulin (Ig)G1 monoclonal antibody, cetuximab, was the first monoclonal antibody to be approved for the treatment of a solid tumor.

Intensive research on usable biomarkers to predict cetuximab response has been conducted
5 to identify those patients who will benefit most significantly from cetuximab treatment. Tumor
EGFR expression as assessed by immunohistochemistry has proved to be a disappointing
biomarker for the efficacy of EGFR-targeted treatment in CRC. More promising data have
been reported for mutations of the KRAS gene, which encodes a GDP/GTP-binding protein
linking ligand-dependent receptor activation to intracellular pathways of the EGFR signaling
10 cascade. Retrospective analyses of the KRAS mutation status in a multitude of clinical
studies including two randomized studies of first-line treatment in metastatic CRC (mCRC),
EMR 62202-047 and EMR 62202-013, as well as the randomized C0.17 study (investigating
cetuximab monotherapy in patients with mCRC who had failed prior chemotherapy) have
demonstrated that KRAS codon 12/13 mutation status is predictive for cetuximab activity in
15 CRC. Tumor responses are predominantly seen in subgroups of patients whose tumors were
wild-type for KRAS and patients carrying a KRAS codon 12/13 mutation do not benefit from
cetuximab therapy. The mutation status of KRAS therefore appears to be a powerful
predictive biomarker for cetuximab activity in CRC, allowing the exclusion from treatment of a
subpopulation unlikely to derive a significant benefit.

20 Yet, not all of the about 60% of CRC patients with KRAS wild-type tumors do benefit from
cetuximab treatment. About 40% of patients with KRAS wild-type tumors do not respond to
cetuximab treatment and a substantial fraction of these patients progresses early and has a
short overall survival.

Therefore, there is a need for the identification and use of further biomarkers that can be used
25 in addition to the KRAS mutation status to better predict the clinical outcome of cetuximab
treatment in CRC patients. It is further a need to identify of biomarkers that allow a better
prediction of cetuximab efficacy in the treatment of CRC in addition to KRAS mutation status.

Microarray analysis of fresh frozen liver metastasis biopsies was performed in the two
cetuximab CRC studies EMR 62202-502 and EMR 62202-045 to identify genes whose
30 expression is associated with response, progression-free survival or overall survival in the
general patient population or in patients with KRAS wild-type tumors. The expression of these
genes can be used as a predictive biomarker for efficacy of cetuximab treatment in CRC and

to better identify those patients who will derive most benefit from cetuximab treatment in CRC.

The expression of the above described genes are used as biomarkers for predicting efficacy of cetuximab and other anti-EGFR directed therapeutic antibodies in patients with CRC and facilitate treatment decisions in the clinic, i.e. if a patient will receive cetuximab or other anti-EGFR directed therapeutic antibodies or not. The application in clinical practice is:

1. Analysis of mRNA expression of these genes from formalin-fixed paraffin embedded (FFPE) or fresh tumor biopsies (the latter have to be either directly frozen in liquid nitrogen or treated with RNA-later to conserve RNA integrity). The biopsies can be obtained from primary tumor or metatasis. Analysis of mRNA expression is performed by PCR-based methods (e.g. real-time PCR, qPCR) by using gene specific primers to amplify the gene of interest or by hybridization of the mRNA of the gene of interest to gene specific, immobilized hybridization probes on gene arrays.
2. Analysis of protein expression of these genes from FFPE or fresh tumor biopsies (the latter have to be either directly frozen in liquid nitrogen or treated with RNA-later to conserve RNA integrity). The biopsies can be obtained from primary tumor or metatasis. Analysis of protein expression includes methods such as immunohistochemistry, enzyme-linked immunosorbent assay (ELISA), Luminex, blotting and detection of proteins on membranes, mass spectrometry.
- For soluble proteins: Analysis of protein expression from plasma or serum comprising methodologies such as ELISA, Luminex, mass spectrometry.

For establishing a diagnostic assay for clinical practice, the expression levels of the candidate gene(s) or protein(s) need to be normalized against the expression of another gene or protein (or a combination of genes and proteins) that is (are) assessed from the same biopsy with the same assay method. These "normalization" genes or proteins can comprise cellular house-keeping genes that are known to display very low variation from patient to patient.

Alternatively, the ratio of the expression level of a gene or protein (or a combination of genes or proteins) from the "good-prognosis" (or whatever terminology we will use in the end (e.g. "sensitivity") group and the expression level of a gene or protein (or a combination of genes or proteins) from the "bad-prognosis" (or e.g. "resistance") group can be determined. The latter approach offers the advantage of using only expression levels of genes or proteins that are directly linked to the efficacy of the anti-EGFR therapy. This approach results in a high dynamic range and is independent of suitable house-keeping genes or proteins.

Prior to implementation of the diagnostic assays a threshold has to be defined which is the ratio of expression levels of the applied markers (as described above) that has to be achieved in order to trigger a positive decision for treating a patient with the respective anti-EGFR therapy. This threshold should discriminate between patients who benefit and patients who do not benefit from the anti-EGFR treatment in the best possible way. Such a threshold has to be derived from a "training-set" of tumor samples from patients treated with the anti-EGFR therapy. Then the threshold has to be prospectively validated in a different set of tumor samples from a sufficient number of patients to prove its ability to select patients who will derive most benefit or to exclude patients who will not benefit from treatment.

10 In two independent clinical studies (EMR 62202-502 and EMR 62202-045) for treatment of CRC patients with cetuximab the expression of the genes described above and in the following was found to be associated with response and/or progression-free survival and/or overall survival.

- EGFR gene amplification may predict favorable outcome to anti-EGFR therapy
- 15 • Subsequent studies found a lower incidence of EGFR gene amplification
- Methodology/comparability issues
- K-Ras mutations "override" benefit in patients with EGFR gene amplification
- Skin-rash is up to now probably the best predictable biomarker for Erbitux activity (mCRC, NSCLC)
- 20 • PhIII study FLEX of Erbitux: early onset of skin rash (first 21 days) associated with prolonged over-all survival (OS)
- BRAF mutations found in 5% of patients
- BRAF and KRAS mutations were mutually exclusive
- BRAF mutation appears to be rather a bad prognostic marker than a predictive marker for Erbitux efficacy.
- 25 • AREG and EREG expression in mCRC is independent of the KRAS mutation status
- Additional predictive power by combining AREG and EREG expression status with KRAS mutation status

Cetuximab treatment according to the experiments of the invention was associated with substantial downregulation of p-EGFR, p-MAPK and proliferation and substantial upregulation of p27^{Kip1} and p-STAT3 levels in basal keratinocytes. No marked difference in

30

these effects was noted for the different schedules of administration and dose levels. In the cetuximab monotherapy phase, responses were achieved only in patients whose tumors were wild-type for *KRAS* (8/29 vs 0/19 for *KRAS* mutant tumors; $P=.015$). Progression-free survival was longer for patients with *KRAS* wild-type compared with *KRAS* mutant tumors (logrank, $P=.048$). Genomics/proteomics analyses identified candidate markers associated with response.

This translational study conducted in a phase I dose-escalation trial of cetuximab monotherapy constitutes to our knowledge the first attempt to use pharmacogenomic and pharmacoproteomic analyses to identify predictive pretreatment biomarkers for cetuximab-responsive mCRC in the first-line setting.

The clinical study (to be reported elsewhere) demonstrated that cetuximab can be safely administered as first-line therapy to patients with mCRC every second week at doses of 400-700 mg/m². The MTD was not reached at the highest dose level, and there were no marked differences in the incidence or severity of adverse events or the activity of cetuximab at different dose levels. Using the skin to measure target impact, the IHC data in the pharmacodynamic biomarker evaluation showed consistent inhibition of signaling proteins within the EGFR pathway across the dose-escalation groups. These data provide a biological rationale supporting the functional equivalence of weekly and every second week dosing regimens.

Retrospective analyses of single arm and randomized mCRC studies have confirmed that the mutation status in tumors of *KRAS* at codons 12 and 13 is a strong predictor of cetuximab activity, with treatment benefit tightly linked to wild-type status.^{4,7-13} The current study addressed for the first time the influence of *KRAS* mutation status on cetuximab monotherapy in the first-line treatment of mCRC patients. Consistent with the data from previous series of chemorefractory patients treated with cetuximab as a single agent or in combination with chemotherapy,^{8,11,13} objective responses in the monotherapy part of the study were only observed in those patients with *KRAS* wild-type tumors (8 out of 29 patients, 28%), with no responses (0 out of 19) seen in patients with tumors that carried mutations in the gene ($P=.015$). The best overall response rates (including the responses after adding FOLFIRI) in patients with *KRAS* wild-type tumors (55%) and *KRAS* mutated tumors (32%) were comparable to results obtained in the CRYSTAL and OPUS studies, where cetuximab was combined as first-line treatment respectively with FOLFIRI and FOLFOX.^{4,12} The observed responses in mCRC patients with *KRAS* mutated tumors receiving cetuximab in combination with chemotherapy as first-line treatment in the current study are most likely attributable to the

effect of chemotherapy. Overall PFS was significantly longer for patients whose tumors were wild-type for *KRAS*, demonstrating the clinical significance of *KRAS* tumor mutation status as a predictive biomarker in relation to cetuximab treatment.

5 A subset of patients with *KRAS* wild-type tumors do not appear to benefit from cetuximab treatment. It may therefore be possible to identify further predictive biomarkers which will facilitate the more accurate tailoring of treatment to those patients who will respond to cetuximab. Recently, the negative predictive value of *BRAF*¹⁸ and *PI3K*^{19,20} mutations as well as PTEN deregulation¹⁹⁻²¹ has been preliminarily described. Other molecular markers which have been putatively associated with the clinical activity of cetuximab include tumor
10 expression levels of *VEGF*, *IL8*, *EGFR*, and *PTGS2* (*COX2*);²² circulating levels of VEGF during treatment;²³ constitutional polymorphisms of *PTGS2* and *EGFR*²⁴ and *TP53* tumor mutation status.²⁵

For a range of anticancer agents high-throughput genomics technologies are increasingly being utilized in the search for predictive biomarkers.²⁶⁻²⁹ In the case of cetuximab, high-level
15 expression of genes encoding the EGFR ligands AREG (amphiregulin) and EREG (epiregulin) in tumors has been shown to be associated with clinical activity in mCRC patients, both by microarray^{10,30} (unselected population and a population with *KRAS* wild-type tumors) and quantitative reverse transcriptase-PCR³¹ (patients receiving cetuximab plus irinotecan) approaches. Similarly, in the current first-line study, AREG and EREG expression
20 appeared to be elevated in tumors of patients without disease progression, in both the total population and the *KRAS* wild-type tumor subgroup. In contrast, *TGF α* (encoding TGF- α), showed lower levels of expression in patients without disease progression. The global gene expression analysis of *KRAS* wild-type tumors identified 57 genes putatively associated with disease control at week 6 ($P < .002$). Among these candidates, six genes (*TNFRSF1B*,
25 *DNAJC8*, *ECSIT*, *GOSR2*, *PPP1R9A*, and *KLK6*) were found to have a False Discovery Rate < 0.1 . The value of these putative biomarkers for improving prediction of cetuximab efficacy in *KRAS* wild-type mCRC needs further exploration.

Luminex analysis of plasma proteins revealed a strong increase in the levels of amphiregulin and TGF- α during cetuximab monotherapy treatment, a trend that was also seen for EGF.
30 The upregulation of these EGFR ligands might be a compensating reaction to EGFR inhibition. Interestingly, the increase in amphiregulin levels was significantly lower in patients who responded to cetuximab treatment. A significant decrease of carcinoembryonic antigen and the cancer antigens 125 and 19-9 was observed under cetuximab monotherapy in responders. Remarkably also, the decrease in IL-8 levels was significantly associated with

response in all tumors as well as in *KRAS* wild-type tumors. IL-8 is a pro-inflammatory cytokine that promotes proliferation and survival of tumor cells and has profound effects on the tumor microenvironment.³² IL-8 seems to be a predictive biomarker for cetuximab efficacy.

- 5 Furthermore according to the invention a direct interaction of EGFR and VAV3 could be detected when EGFR and VAV3 were expressed in HEK 293 cells. This indicates a direct and outstanding role for VAV3 in EGFR signaling and a direct link between observed high VAV3 expression levels and modulation of the activity of anti-EGFR therapy with cetuximab.

The invention shows for the first time that treatment with anti-EGFR antibodies, preferably
10 cetuximab (every second week and weekly administration) as a single agent in a first-line setting benefits mCRC patients who have *KRAS* wild-type tumors. In addition, the global gene expression analyses of this early-phase study have generated a number of interesting results regarding the expression of certain genes and the clinical activity of cetuximab. These observations enable validation on larger patient series using different methodologies. The
15 results of these studies provide a rational foundation for optimizing treatment in patients suffering from different cancers, especially CRC or mCRC with cetuximab or anti-EGFR antibodies being similarly active.

Immunohistochemical Analysis of EGFR Pathway Components

Evaluable paired baseline/week 4 skin biopsies to analyze pharmacodynamic changes of the
20 assessed markers were available from up to 35 patients. Substantial downregulation of p-EGFR, p-MAPK and proliferation (as assessed by Ki67 staining) was observed in the 4-week compared with baseline samples. In parallel, a substantial upregulation of p27^{Kip1} and p-STAT3 was observed. In the analysis of different schedules of administration and dose levels, no relevant differences in relation to changes in the levels of these markers between
25 groups of patients were present for baseline to on-treatment timepoints (Fig. 21A).

Evaluable paired baseline/week 4 tumor biopsies were available from up to 17 patients. Reduction in proliferation and a profound downregulation of p-EGFR and p-MAPK were observed in tumor cells after therapy (Fig. 21B). However, p27^{Kip1}, p-STAT3 and p-AKT levels were not markedly modified by cetuximab treatment (data not shown). The small
30 number of available paired tumor biopsies did not allow a comparison of changes in biomarker levels with dose groups and response variables.

KRAS Mutation Analysis

KRAS codon 12 or 13 mutations were detected in 19/48 (40%) patient samples (G12V, 9 patients; G13D, 5 patients; G12D, 4 patients; G12A, 1 patient). In the cetuximab monotherapy phase, there were eight partial responses (PRs) among the 48 patients. All were in patients
5 whose tumors were wild-type for *KRAS* (8/29; 28%). No responses were reported in the 19 patients whose tumors carried *KRAS* mutations ($P=.015$) (Table 1). In the study overall (monotherapy and combination therapy phases), responses were seen in 16/29 (55%) of patients with *KRAS* wild-type and 6/19 (32%) with *KRAS* mutant tumors ($P=.144$). PFS was significantly longer in patients whose tumors were wild-type for *KRAS*, compared to those
10 whose tumors carried mutations (Fig. 22; median 9.4 vs 5.6 months, hazard ratio 0.47; logrank $P=.048$).

Microarray Analysis of Gene Expression

A total of 106 tumor-derived samples from baseline and 4-week timepoints were hybridized to Affymetrix GeneChip HG-U133 Plus 2.0 arrays. Four arrays were excluded from further
15 analysis on the basis of general quality control parameters and 24 samples were excluded due to presence of normal liver tissue contamination (Fig. 16; supplemental material) and were not further analyzed. After the exclusion of duplicates, 62 array data sets from 42 ITT patients (36 baseline, 26 week 4: 20 pairs) remained for analysis.

For the analyzed tumor samples, the data from the 54,675 probe sets was pre-filtered on the
20 basis of variance, signal intensity and probe set annotation (see Supplementary Methods). This process restricted the tumor expression analysis to 15,230 probe sets, representing 10,538 genes. In global comparisons of baseline pre-filtered data according to response: progressive disease (PD; $n=12$) versus disease control at 6 weeks ($n=23$; 1 patient not evaluable) and for best overall response: PD and stable disease (SD) ($n=19$) versus PR
25 ($n=14$; 3 patients not evaluable), the distribution of P values (data not shown) was essentially as expected by chance, suggesting that a gene expression profile predictive of response had not been identified for the global population. However, restricting the analysis to *KRAS* wild-type tumors (8 patients with PD versus 11 patients with disease control), 57 probe sets with expression patterns putatively associated with disease control at week 6 were identified
30 ($P<.002$; Fig 23). Imposing a False Discovery Rate (FDR) threshold of 0.1 (for FDR definition see Supplementary Methods section), six genes were found to be significantly associated with disease control (*TNFRSF1B*, $P=6.90E-07$; *DNAJC8*, $P=1.60E-06$; *ECSIT*, $P=6.80E-06$;

GOSR2, $P=3.90E-05$ with higher expression in patients showing disease control and PPP1R9A, $P=8.90E-07$ and KLK6, $P=3.00E-05$ with higher expression in patients with PD).

On treatment changes associated with response were examined using data from patients with available samples from the baseline and the week 4 timepoints. No expression changes were identified that appeared to be tightly associated with disease control at week 6 ($n=12$) compared with PD ($n=8$). Considering the combination with chemotherapy, the comparison of profiles for PR ($n=7$) versus SD/PD ($n=13$), revealed 47 probe sets showing differences in on-treatment changes ($P<.002$, moderated t-test, see Fig. 17).

In patients with *KRAS* wild-type tumors, as well as in the complete set of analyzed patients, baseline *EREG* (epiregulin) and *AREG* (amphiregulin) expression levels were higher in those tumors responding to cetuximab (Fig. 24; Fig. 18). These findings are in line with results reported by Khambata-Ford et al.¹⁰ Interestingly, *TGFA* (TGF- α) demonstrated a reciprocal expression pattern (Fig. 24; Fig. 18). Among other genes known to be directly or indirectly involved in EGFR signaling such as ERBB receptors and ligands *ERBB3* (*HER3*) and *ERBB2* (*HER2*) showed a trend for stronger downregulation in tumors with a PR as best overall response (Fig. 19).

Plasma Proteomics

The concentrations of 97 different proteins were analyzed in plasma samples using Luminex technology. The protein panel included EGFR ligands, other growth factors, interleukins and a variety of other candidate proteins. During the cetuximab monotherapy phase, the decrease in plasma levels of interleukin (IL)-8, macrophage inflammatory protein (MIP)-1 α as well as the tumor markers carcinoembryonic antigen, cancer antigens 125 and 19-9 between baseline and week 4 was significantly associated ($P<.01$) with response at week 6 (Fig. 25A). The general strong increase in plasma concentrations of amphiregulin was significantly ($P<.01$) weaker in patients with partial response to cetuximab monotherapy (Fig. 25). The association with response at week 6 was also found for carcinoembryonic antigen, cancer antigen 19-9, IL-8 and amphiregulin when the analysis was restricted to patients with *KRAS* wild-type tumors (data from 24 patients, Fig. 25B). Furthermore a general increase (independent of response) of TGF- α and EGF levels and a decrease of soluble EGFR were observed in plasma during the first 4 weeks of cetuximab monotherapy treatment.

Among the investigated genes and candidates showing an association between expression levels and success of therapy in patients with metastatic colorectal cancer (mCRC) receiving cetuximab, VAV3 is of particular interest. In study EMR 62202-502 high tumoral VAV3 mRNA

expression levels were not only strongly associated with better response to cetuximab in combination with irinotecan (Figure 26) but also with progression free survival (PFS) (Figure 27) and overall survival (OS) (Figure 28). Furthermore, high tumoral VAV3 expression levels were found to be particularly associated with response and prolonged PFS in patients with KRAS wild-type tumors (see Figure 1 and Figure 29). Therefore, VAV3 expression appears to be a good biomarker candidate for predicting clinical outcome of cetuximab therapy in CRC in patients with KRAS wild-type tumors which would help to further optimize selection of patients deriving most benefit from cetuximab therapy.

Interestingly, a direct interaction of EGFR and VAV3 could be detected when EGFR and VAV3 were expressed in HEK 293 cells (Figure 30). This suggests a direct role for VAV3 in EGFR signaling and a direct link between VAV3 expression levels and modulation of the activity of anti-EGFR therapy.

Examples

Clinical studies:

EMR 62202-502: This randomized study investigated cetuximab dose-escalation in patients (pts) with EGFR-expressing mCRC failing irinotecan-including therapy. Pts were randomized 22 days after starting cetuximab (400 mg/m² initial dose then 250 mg/m²/week [w]) with I (180 mg/m² q 2 w) if they had not experienced >grade (G) 1 skin reaction, any other >G 2 cetuximab-related adverse event and were tolerant to I. Randomization was to standard cetuximab dose (Arm A; 250 mg/m²/w) or dose-escalation (Arm B; cetuximab dose increased by 50 mg/m² q 2 w, until >G 2 toxicity, tumor response or dose = 500 mg/m²). Pts not randomized (Arm C) continued on standard cetuximab dose. Primary endpoint was to compare in skin and tumor biopsies, taken before and during treatment, the effects of dose-escalation on EGFR and downstream signalling markers with those of the standard cetuximab regimen. Secondary endpoints were PK, efficacy, safety, tolerability, biomarker analyses on tumor biopsies and plasma samples. The KRAS mutation status was analyzed from tumor biopsies.

EMR 62202-045: This study examined the safety and pharmacokinetics of every second week administration of cetuximab in patients with metastatic colorectal cancer. Secondary objectives included a pharmacodynamic biomarker analysis. Patients received cetuximab monotherapy for 6 weeks, followed by cetuximab plus FOLFIRI until disease progression.

Patients in the control arm received cetuximab as a 400 mg/m² initial dose then 250 mg/m²/week and in the dose-escalation arms, at 400-700 mg/m², every second week. The KRAS mutation status was analyzed from tumor biopsies.

Tumor material for Gene expression (Microarray) analysis:

5 EMR 62202-502: Tumor material was taken by open surgery, endoscopy or core/fine needle biopsy at baseline (pre-treatment), at day 22 and if possible, at disease progression of patients in the dose-escalation arm (Arm B). The samples were snap-frozen in liquid nitrogen.

EMR 62202-045: Tumor material was taken by open surgery, endoscopy or core/fine needle biopsy at baseline, at week 4 and if possible, at disease progression. The samples were
10 snap-frozen in liquid nitrogen.

RNA Expression Profiling

Experimental procedures related to the microarray analysis are detailed in the Supplementary Methods section. Briefly, snap-frozen tumor biopsies were homogenized and total RNA was extracted using an RNeasy Micro Kit® (Qiagen, Hilden). Biotinylated target cRNAs for the
15 array hybridization experiments were prepared from all samples according to the Affymetrix Two-Cycle Eukaryotic Target labeling protocol. For each tumor analyzed, an initial 50 ng of total RNA was included in the first cDNA synthesis reaction of this cRNA amplification/labeling process. Labeled cRNA was subsequently hybridized to Affymetrix GeneChip HG-U133 Plus 2.0 gene expression arrays for 16 hours at 45 °C at 60 rpm.
20 Following hybridization, arrays were stained on an Affymetrix Fluidics Station 450 and signal quantified using a GeneChip Scanner. Quality control and preprocessing of the raw expression data were carried out using the proprietary Affymetrix GCOS software and the Bioconductor package, affyPLM.

EMR 62202-502: After all quality control checks and pre-processing steps had been
25 performed a total of 68 array data sets from 47 subjects of the intention to treat (ITT) population were eligible for further analysis. Baseline samples were available from 35 subjects.

EMR 62202-045: After all quality control checks and pre-processing steps had been performed a total of 62 array data sets from 42 ITT patients were eligible for further analysis.
30 Baseline samples were available from 36 subjects.

Statistical analyses were conducted for all Affymetrix probe sets with reliable gene annotation that passed initial filters based on variability and signal intensity in at least one of the two studies (16414 Affymetrix probe sets representing 10785 genes).

Genes whose expression is associated with clinical response were identified with Welch t-test
5 comparisons between responders and non-responders (EMR 62202-502), or between patients with disease control after six weeks of cetuximab monotherapy versus those with progressive disease (EMR 62202-045), respectively. Genes whose expression is associated with progression-free survival or with overall survival were identified using Cox proportional hazards regression (EMR 62202-502). These analyses were conducted for the complete sets
10 of patients, as well as only for the patients with KRAS wildtype tumors.

A meta-analysis to identify response-associated genes across both studies was conducted using the products of single-study one-sided p-values as test statistic and deriving p-values from the null distribution of these products.

P-values in a range below 0.01 and 0.0001, specifically below 0.01, preferably 0.005, more
15 preferably 0.002, most preferably 0.0005 or 0.0001 (from the meta-analysis for association with clinical response, and from the analysis of EMR 62202-502 for the association with progression-free and overall survival), were considered as statistically significant. This criterion was fulfilled for 200 Affymetrix probe sets representing 179 known genes in at least one of the comparisons.

20 ***Patient Eligibility and Study Design***

Eligibility criteria and study design have been reported in full in a separate manuscript. Briefly, the study was divided into two parts; a cetuximab monotherapy phase lasting 6 weeks and a combination therapy phase, during which patients received cetuximab, at the same
25 dose/schedule as during the monotherapy phase, and the irinotecan-based schedule FOLFIRI. Patients were assigned sequentially to either the standard weekly schedule and dose of cetuximab (400 mg/m², followed by weekly doses of 250 mg/m²) or a cetuximab dose-escalation treatment on a bi-weekly schedule with different cohorts from 400 to 700 mg/m². Clinical response was reported after 6 weeks of cetuximab monotherapy and as best overall response (monotherapy and combination therapy phases).

30 ***Collection and Storage of Patient Material***

Skin biopsies were obtained at baseline and on day 26-28 (week 4) . If skin rash was present, samples were taken from a rash-free area. The biopsy was immediately immersed into ≥20

times its volume of neutral-buffered formaldehyde solution at 4 °C, and held for 8-16 hours at room temperature. The fixed specimen was dehydrated to xylene using a graded ethanol series and embedded longitudinally in paraffin wax under vacuum at 60 °C. Tumor material was taken by open surgery, endoscopy or core/fine needle biopsy at baseline, at week 4 and
5 if possible, at disease progression. One sample per timepoint was formalin fixed and paraffin embedded, as previously described^{13a} and three samples were snap-frozen in liquid nitrogen. To provide normal DNA, 10 ml of whole blood was obtained from each patient at baseline and stored at -20 °C or lower until use. Plasma (2.5 ml) was collected for Luminex analysis at baseline and week 4, and stored at -80 °C.

10 ***Immunohistochemistry***

Immunohistochemical (IHC) analysis of formalin fixed paraffin embedded (FFPE) tissue was used to investigate the expression of the following proteins: EGFR, phospho(p)-EGFR, p-MAPK, Ki67 (MIB1), p27^{Kip1} (CDKN1B) and p-STAT3 (skin and tumor biopsies); HER2, p-HER2 and p-AKT (tumor biopsies). Immunohistochemistry analysis was performed as
15 previously described.^{13a} Details of the antibodies and methods used are provided in the Supplementary Methods section.

KRAS Mutation Analysis

FFPE patient-derived archival tumor tissue was available from 48 patients from the intention to treat (ITT) population. DNA was extracted and screened for the presence of *KRAS* codon
20 12 and 13 mutations using a polymerase chain reaction (PCR) clamping and melting curve technique adapted from Chen et al, 2004¹⁴ (LightMix, k-ras Gly12, TIB MOLBIOL, Berlin, Germany), as previously described.¹²

RNA Expression Profiling

Experimental procedures related to the microarray analysis are detailed in the Supplementary
25 Methods section. Briefly, snap-frozen tumor biopsies were homogenized and total RNA was extracted using an RNeasy Micro Kit® (Qiagen, Hilden). Biotinylated target cRNAs for the array hybridization experiments were prepared from all samples according to the Affymetrix Two-Cycle Eukaryotic Target labeling protocol. For each tumor analyzed, an initial 50 ng of total RNA was included in the first cDNA synthesis reaction of this cRNA
30 amplification/labeling process. Labeled cRNA was subsequently hybridized to Affymetrix GeneChip HG-U133 Plus 2.0 gene expression arrays for 16 hours at 45 °C at 60 rpm. Following hybridization, arrays were stained on an Affymetrix Fluidics Station 450 and signal

quantified using a GeneChip Scanner. Quality control of the raw expression data was carried out using the proprietary Affymetrix GCOS software and the Bioconductor package affyPLM.¹⁵ If replicate arrays were available from individual samples, the data set with the best quality control assessment was selected for analysis. Preprocessing of the raw probe-level intensity data was performed using the GCRMA algorithm.¹⁶

Proteomic Analysis

A multiplex analysis of 97 proteins (HumanMAP version 1.6 plus amphiregulin, betacellulin, EGFR, heparin-binding (HB)-EGF, epiregulin, interleukin-18, transforming growth factor (TGF)- α , and thrombospondin-1) from plasma using the Luminex xMAP[®] technology platform (as described in the Supplementary Methods section) was performed at Rules-Based Medicine (Austin, Texas, US). Betacellulin, EGFR and HB-EGF were only assessed in 23 samples from patients who were enrolled later in the trial.

Statistical analysis

Response rates and progression-free survival (PFS), defined as the duration from the first infusion of cetuximab until the first radiologically confirmed disease progression under the combination of cetuximab and FOLFIRI, for patients whose tumors were wild-type or mutant with respect to *KRAS* were compared using Fisher's exact and logrank tests, respectively.

All statistical analyses of the IHC, microarray and proteomics data (see Supplementary Methods) were performed using Bioconductor software¹⁵ and SAS version 9.1. These exploratory analyses were viewed as hypothesis-generating.

REFERENCES (relevant and/or used in this description)

1. Cunningham D, Humblet Y, Siena S, et al: Cetuximab monotherapy and cetuximab plus irinotecan in irinotecan-refractory metastatic colorectal cancer. *N Engl J Med* 351:337-345, 2004
- 25 2. Jonker DJ, O'Callaghan CJ, Karapetis CS, et al: Cetuximab for the treatment of colorectal cancer. *N Engl J Med* 357:2040-2048, 2007
3. Sobrero AF, Maurel J, Fehrenbacher L, et al: EPIC: phase III trial of cetuximab plus irinotecan after fluoropyrimidine and oxaliplatin failure in patients with metastatic colorectal cancer. *J Clin Oncol* 26:2311-2319, 2008
- 30 4. Van Cutsem E, Kohne CH, Hitre E, et al: Cetuximab and chemotherapy as initial treatment for metastatic colorectal cancer. *N Engl J Med* 360:1408-1417, 2009

5. Cappuzzo F, Finocchiaro G, Rossi E, et al: EGFR FISH assay predicts for response to cetuximab in chemotherapy refractory colorectal cancer patients. *Ann Oncol* 19:717-723, 2008
6. Chung KY, Shia J, Kemeny NE, et al: Cetuximab Shows Activity in Colorectal Cancer Patients With Tumors That Do Not Express the Epidermal Growth Factor Receptor by Immunohistochemistry. *J Clin Oncol* 23:1803-1810, 2005
7. De Roock W, Piessevaux H, De Schutter J, et al: KRAS wild-type state predicts survival and is associated to early radiological response in metastatic colorectal cancer treated with cetuximab. *Ann Oncol* 19:508-515, 2008
- 10 8. Di Fiore F, Blanchard F, Charbonnier F, et al: Clinical relevance of KRAS mutation detection in metastatic colorectal cancer treated by cetuximab plus chemotherapy. *Br J Cancer* 96:1166-1169, 2007
9. Finocchiaro G, Cappuzzo F, Jänne PA, et al: EGFR, HER2 and Kras as predictive factors for cetuximab sensitivity in colorectal cancer. *J Clin Oncol* 25, 2007:(suppl; abstr 15 4021)
10. Khambata-Ford S, Garrett CR, Meropol NJ, et al: Expression of epiregulin and amphiregulin and K-ras mutation status predict disease control in metastatic colorectal cancer patients treated with cetuximab. *J Clin Oncol* 25:3230-3237, 2007
11. Lievre A, Bachet JB, Boige V, et al: KRAS mutations as an independent prognostic factor in patients with advanced colorectal cancer treated with cetuximab. *J Clin Oncol* 26:374-379, 2008
- 20 12. Bokemeyer C, Bondarenko I, Makhson A, et al: Fluorouracil, leucovorin, and oxaliplatin with and without cetuximab in the first-line treatment of metastatic colorectal cancer. *J Clin Oncol* 27:663-671, 2009
- 25 13. Karapetis CS, Khambata-Ford S, Jonker DJ, et al: K-ras mutations and benefit from cetuximab in advanced colorectal cancer. *N Engl J Med* 359:1757-1765, 2008
14. Rojo F, Tabernero J, Albanell J, et al: Pharmacodynamic studies of gefitinib in tumor biopsy specimens from patients with advanced gastric carcinoma. *J Clin Oncol* 24:4309-4316, 2006
- 30 15. Chen CY, Shiesh SC, Wu SJ: Rapid detection of K-ras mutations in bile by peptide nucleic acid-mediated PCR clamping and melting curve analysis: comparison with restriction fragment length polymorphism analysis. *Clin Chem* 50:481-489, 2004

16. Gentleman RC, Carey VJ, Bates DM, et al: Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol* 5:R80, 2004
17. Wu Z, Irizarry RA, Gentleman R, et al: A model-based background adjustment for oligonucleotide expression arrays. *JASA* 99:909-917, 2004
- 5 18. Di Nicolantonio F, Martini M, Molinari F, et al: Wild-type BRAF is required for response to panitumumab or cetuximab in metastatic colorectal cancer. *J Clin Oncol* 26:5705-5712, 2008
19. Perrone F, Lampis A, Orsenigo M, et al: PI3KCA/PTEN deregulation contributes to impaired responses to cetuximab in metastatic colorectal cancer patients. *Ann Oncol* 20:84-
10 90, 2009
20. Jhawer M, Goel S, Wilson AJ, et al: PIK3CA mutation/PTEN expression status predicts response of colon cancer cells to the epidermal growth factor receptor inhibitor cetuximab. *Cancer Res* 68:1953-1961, 2008
21. Loupakis F, Pollina L, Stasi I, et al: PTEN Expression and KRAS Mutations on Primary
15 Tumors and Metastases in the Prediction of Benefit From Cetuximab Plus Irinotecan for Patients With Metastatic Colorectal Cancer. *J Clin Oncol* 27:2622-2629, 2009
22. Vallbohmer D, Zhang W, Gordon M, et al: *Molecular determinants of cetuximab efficacy.* *J Clin Oncol* 23:3536-3544, 2005
23. Vincenzi B, Santini D, Russo A, et al: Circulating VEGF reduction, response and
20 outcome in advanced colorectal cancer patients treated with cetuximab plus irinotecan. *Pharmacogenomics* 8:319-327, 2007
24. Lurje G, Nagashima F, Zhang W, et al: Polymorphisms in cyclooxygenase-2 and epidermal growth factor receptor are associated with progression-free survival independent of K-ras in metastatic colorectal cancer patients treated with single-agent cetuximab. *Clin*
25 *Cancer Res* 14:7884-7895, 2008
25. Oden-Gangloff A, Di Fiore F, Bibeau F, et al: TP53 mutations predict disease control in metastatic colorectal cancer treated with cetuximab-based chemotherapy. *Br J Cancer* 100:1330-1335, 2009
26. Hsu DS, Balakumaran BS, Acharya CR, et al: Pharmacogenomic strategies provide a
30 rational approach to the treatment of cisplatin-resistant patients with advanced cancer. *J Clin Oncol* 25:4350-4357, 2007

27. Bonnefoi H, Potti A, Delorenzi M, et al: Validation of gene signatures that predict the response of breast cancer to neoadjuvant chemotherapy: a substudy of the EORTC 10994/BIG 00-01 clinical trial. *Lancet Oncol* 8:1071-1078, 2007
28. Burington B, Barlogie B, Zhan F, et al: Tumor cell gene expression changes following
5 short-term in vivo exposure to single agent chemotherapeutics are related to survival in multiple myeloma. *Clin Cancer Res* 14:4821-4829, 2008
29. Kunz M: Genomic signatures for individualized treatment of malignant tumors. *Curr Drug Discov Technol* 5:9-14, 2008
30. de Reynies A, Boige V, Milano G, et al: KRAS mutation signature in colorectal tumors
10 significantly overlaps with the cetuximab response signature. *J Clin Oncol* 26:2228-2230; author reply 2230-2221, 2008
31. Tejpar S, De Roock W, Biesmans B, et al: High amphiregulin and epiregulin expression in KRAS wild type colorectal primaries predicts response and survival benefit after treatment with cetuximab and irinotecan for metastatic disease. *ASCO Gastrointestinal
15 Cancers Symposium*, January 25-27, 2008, Orlando, FL (abstr 411)
32. Waugh DJ, Wilson C: The interleukin-8 pathway in cancer. *Clin Cancer Res* 14:6735-6741, 2008

PATENT CLAIMS:

1. An *in vitro* method for predicting the likelihood that a patient suffering from KRAS wild type EGFR expressing tumor, who is a candidate for treatment with an EGFR antibody, will respond to the treatment with said anti-EGFR antibody, comprising determining the expression level of one or more prognostic genes or gene expression products thereof in a tissue sample obtained from said patient by subjecting a nucleic acid sample from the tumor sample from the patient to PCR or an RNA or DNA array or a comparable diagnostic tool or apparatus, wherein
 - (i) high expression of the gene or the gene product selected from the group of genes consisting of: ADAMDEC1, BSDC1, C1orf144, CAPZB, CDC42, DHCR7, DNAJC8, ECSIT, EXOSC10, FADS1, GBA, GLT25D1, GOSR2, IMPDH1, KLHL21, KPNA6, KPNB1, LSM12, MAN1B1, MIDN, PPAN, SH3BP2, SQLE, SSH3, TNFRSF1B, URM1, ZFYVE26, RGMB, SPIRE2, ABCC5, ACSL5, AOA1, AXIN2, CD24, CEACAM5, CEACAM6, ETS2, FMNL2, GPSM2, HDAC2, JUN, ME3, MED17, MYB, MYC, NEBL, NOSIP, PITX2, POF1B, PPARG, PPP1R14C, PRR15, PSMG1, RAB15, RAB40B, RNF43, RPS23, SLC44A3, SOX4, THEM2, VAV3, ZNF337, EPDR1, KCNK5, KHDRBS3, PGM2L1, STK38, and SHROOM2, indicates that the patient is likely to respond to said treatment compared to a reference value, and
 - (ii) high expression of the gene or the gene product selected from the group of genes consisting of: C7orf46, CAST, DCP2, DIP2B, ERAP1, INSIG2, KIF21A, KLK6, NGRN, NRIP1, PHGDH, PPP1R9A, QPCT, RABEP1, RPA1, RPL22L1, SKP1, SLC25A27, SLC25A46, SOCS6, TPD52, ZDHHC2, ZNF654, ASB6, ATM, BMI1, CDC42EP2, EDEM3, PLLP, RALBP1, SLC4A11, TNFSF15, TPK1, C11orf9, C1QC, CABLES1, CDK6, EHBP1, EXOC6, EXT1, FLRT3, GCNT2, MTHFS, PIK3AP1, ST3GAL1, TK2, ZDHHC14, ARFGAP3, AXUD1, CAPZB, CHSY1, DNAJB9, GOLT1B, HSPA5, LEPROTL1, LIMS1, MAPK6, MYO6, PROSC, RAB8B, RAP2B, RWDD2B, SERTAD2, SOCS5, TERF2IP, TIAL1, TIPARP, TRIM8, TSC22D2, TGFA, VAPA, and UBE2K, indicates that the patient is likely not to respond to said treatment compared to a reference value.
2. A method of claim 1, wherein the treatment with said anti-EGFR antibody as a first-line therapy and the selected genes or gene expression products from genes of group (i) are one or more of ADAMDEC1, BSDC1, C1orf144, CAPZB, CDC42, DHCR7, DNAJC8, ECSIT, EXOSC10, FADS1, GBA, GLT25D1, GOSR2, IMPDH1, KLHL21, KPNA6, KPNB1, LSM12, MAN1B1, MIDN, PPAN, SH3BP2, SQLE, SSH3, TNFRSF1B, URM1,

VAV3 and ZFYVE26, and from group (ii) are one or more of C7orf46, CAST, DCP2, DIP2B, ERAP1, INSIG2, KIF21A, KLK6, NGRN, NRIP1, PHGDH, PPP1R9A, QPCT, RABEP1, RPA1, RPL22L1, SKP1, SLC25A27, SLC25A46, SOCS6, TPD52, TGFA, ZDHHC2, and ZNF654.

- 5 3. A method of claim 1, wherein the treatment with said anti-EGFR antibody is a combination therapy with a chemotherapeutic agent after the patient has developed a chemo-refractory tumor, and the selected genes or gene expression products from group (i) are one or more of RGMB, SPIRE2, ABCC5, ACSL5, AOA, AXIN2, CD24, CEACAM5, CEACAM6, ETS2, FMNL2, GPSM2, HDAC2, JUN, ME3, MED17, MYB,

10 MYC, NEBL, NOSIP, PITX2, POF1B, PPARG, PPP1R14C, PRR15, PSMG1, RAB15, RAB40B, RNF43, RPS23, SLC44A3, SOX4, THEM2, VAV3, ZNF337, EPDR1, KCNK5, KHDRBS3, PGM2L1, STK38, and SHROOM2, and from group (ii) are one or more of ASB6, ATM, BMI1, CDC42EP2, EDEM3, PLLP, RALBP1, SLC4A11, TNFSF15, TPK1, C11orf9, C1QC, CABLES1, CDK6, EHBP1, EXOC6, EXT1, FLRT3, GCNT2, MTHFS,

15 PIK3AP1, ST3GAL1, TK2, ZDHHC14, ARFGAP3, AXUD1, CAPZB, CHSY1, DNAJB9, GOLT1B, HSPA5, LEPROTL1, LIMS1, MAPK6, MYO6, PROSC, RAB8B, RAP2B, RWDD2B, SERTAD2, SOCS5, TERF2IP, TIAL1, TIPARP, TRIM8, TSC22D2, TGFA, VAPA, and UBE2K.
4. A method of claim 3, wherein the clinical response is overall survival time (OS), and the

20 selected genes from group (i) are one or more of EPDR1, KCNK5, KHDRBS3, PGM2L1, SHROOM2, STK38, and VAV3, and from group (ii) are one or more of ASB6, ATM, BMI1, CDC42EP2, EDEM3, PLLP, RALBP1, SLC4A11, TNFSF15, and TPK1, or the respective gene expression products of each of said groups.
5. A method of claim 3, wherein the clinical response is progression free survival time

25 (PFS) and the selected genes from group (i) are one or more of ACSL5, AOA, AXIN2, CD24, CEACAM5, CEACAM6, ETS2, FMNL2, GPSM2, HDAC2, JUN, ME3, MED17, MYB, MYC, NEBL, NOSIP, PITX2, POF1B, PPARG, PPP1R14C, PRR15, PSMG1, RAB15, RAB40B, RNF43, RPS23, SLC44A3, SOX4, THEM2, VAV3, and ZNF337, and from group (ii) are one or more of C11orf9, C1QC, CABLES1, CDK6, EHBP1, EXOC6,

30 EXT1, FLRT3, GCNT2, MTHFS1, PIK3AP1, ST3GAL1, TK2, and ZDHHC14, or the respective gene expression products of each of said groups.
6. A method of claim 3, wherein the clinical overall response (OR) is measured as partial response versus stable or progressive disease, and the selected genes from group (i)

are one or more of VAV3, RGMB, and SPIRE2, and from group (ii) are one or more of ARFGAP3, AXUD1, CAPZB, CHSY1, DNAJB9, GOLT1B, HSPA5, LEPROTL1, LIMS1, MAPK6, MYO6, PROSC, RAB8B, RAP2B, RWDD2B, SERTAD2, SOCS5, TERF2IP, TIAL1, TIPARP, TRIM8, TSC22D2, TGFA, VAPA, and UBE2K, or the respective gene expression products of each of said groups.

7. A method of any of the claims 1 – 6, wherein the reference value is defined by one or more of a specific functional or clinical property, and / or a specific expression profile obtained from a reference patient or reference patient group.

8. A method of claim 7, wherein said reference value is obtained from a reference patient or patient group that does not express or express little said gene or gene product.

9. A method of any of the claims 1 – 8, wherein the reference value is an expression threshold value of a control gene or the ratio of gene expression of selected genes from group (i) in comparison to gene expression of selected genes from group (ii) or the reference value is an expression threshold value defined by specific clinical response parameters to be determined or by specific pre-treatment or treatment conditions.

10. A method of claim 9, wherein the clinical response parameter is progression free survival time (PFS), overall survival time (OS), partial response (PR), stable disease (SD), progressive disease (PD) or combinations thereof.

11. A method of any of the claims 1 – 10, wherein the tissue samples are taken from the patient before treatment with said anti-EGFR antibody.

12. A method of claim 11, wherein additionally tissue samples are taken from the patient on treatment with said anti-EGFR antibody.

13. A method of claim 12, wherein the expression levels of the genes or gene expression products obtained on treatment are compared with the values obtained before starting treatment of said patient.

14. A method of any of the claims 1 – 13, wherein the patient sample derives from tumor tissue.

15. A method of any of the claims 1 – 13, wherein the patient sample derives from plasma.

16. A method of any of the claims 1 – 15, wherein the level of the expressed proteins encoded by said genes is determined.

17. An *in vitro* method for predicting the likelihood that a patient suffering from KRAS wild type EGFR expressing cancer will respond therapeutically to the treatment with an anti-EGFR antibody, the method comprises:

(a) measuring by diagnostic means and / or diagnostic apparatus in a biopsy tissue

sample from tumor tissue or plasma of said patient the expression level of one or more

biomarkers selected from the group (i) consisting of ADAMDEC1, BSDC1, C1orf144,

CAPZB, CDC42, DHCR7, DNAJC8, ECSIT, EXOSC10, FADS1, GBA, GLT25D1,

GOSR2, IMPDH1, KLHL21, KPNA6, KPNB1, LSM12, MAN1B1, MIDN, PPAN, SH3BP2,

SQLE, SSH3, TNFRSF1B, URM1, ZFYVE26, RGMB, SPIRE2, ABCC5, ACSL5, AOA, H,

AXIN2, CD24, CEACAM5, CEACAM6, ETS2, FMNL2, GPSM2, HDAC2, JUN, ME3,

MED17, MYB, MYC, NEBL, NOSIP, PITX2, POF1B, PPARG, PPP1R14C, PRR15,

PSMG1, RAB15, RAB40B, RNF43, RPS23, SLC44A3, SOX4, THEM2, VAV3, ZNF337,

EPDR1, KCNK5, KHDRBS3, PGM2L1, STK38, SHROOM2,

and / or from group (ii) consisting of C7orf46, CAST, DCP2, DIP2B, ERAP1, INSIG2,

KIF21A, KLK6, NGRN, NRIP1, PHGDH, PPP1R9A, QPCT, RABEP1, RPA1, RPL22L1,

SKP1, SLC25A27, SLC25A46, SOCS6, TPD52, ZDHHC2, ZNF654, ASB6, ATM, BMI1,

CDC42EP2, EDEM3, PLLP, RALBP1, SLC4A11, TNFSF15, TPK1, C11orf9, C1QC,

CABLES1, CDK6, EHBP1, EXOC6, EXT1, FLRT3, GCNT2, MTHFS, PIK3AP1,

ST3GAL1, TK2, ZDHHC14, ARFGAP3, AXUD1, CAPZB, CHSY1, DNAJB9, GOLT1B,

HSPA5, LEPROTL1, LIMS1, MAPK6, MYO6, PROSC, RAB8B, RAP2B, RWDD2B,

SERTAD2, SOCS5, TERF2IP, TIAL1, TIPARP, TRIM8, TSC22D2, TGFA, VAPA, and

UBE2K,

(b) exposing ex-vivo a tissue sample from tumor or plasma of said patient to said anti-

EGFR antibody, (c) measuring again in said exposed tissue sample of step (b) the

expression level of one or more biomarkers specified in step (a), and (d) calculating the

differences in expression levels measured in steps (b) and (c),

wherein an increase in the expression level of the biomarkers of group (i) obtained in

step (c) compared to step (a) indicates an increased likelihood that said patient responds

therapeutically to the treatment with said anti-EGFR antibody, and wherein an increase

in the expression level of the biomarkers of group (ii) obtained in step (c) compared to

step (a) indicates a decreased likelihood that said patient responds therapeutically to the

treatment with said anti-EGFR antibody.

18. A method of any of the claims 1 – 17, wherein the genes or gene expression products

are selected from the group consisting of TNFRSF1B, DNAJC8, ECSIT, GOSR2,

PPP1R9A, VAV3 and KLK6.

19. A method of any of the claims 1 – 17, wherein one of the genes or gene expression products from group (i) and TGFa.
20. A method of claim 19, wherein the expression levels of VAV3 and TGFa are determined.
21. A method of any of the claims 1 – 17, wherein one of the genes or gene expression products from group (ii) and AREG or EREG.
22. A method of any of the claims 19 – 21, the expression levels of TGFa, and AREG or EREG and optionally of VAV3 and / or EGF are determined.
23. A method of any of the claims 1 – 22, wherein the patient suffering from a KRAS wild type EGFR expressing tumor additionally has a EGFR mutation in tumor tissue.
24. A method of claim 23, wherein the EGFR mutation is a R521K polymorphism.
25. A method of any of the claims 1 – 24, wherein the anti-EGFR antibody is c225 (cetuximab).
26. A method of claim 25, wherein the tumor from which the patient suffers is colorectal cancer (CRC) or metastatic colorectal cancer (mCRC).
27. A method of claim 26, wherein additionally the expression level of AREG and / or EREG or their expression products is determined and compared to one or more of the genes or gene expression products of group (i) and / or group (ii).
28. A method of claim 27, wherein the expression level of AREG and / or EREG and at least of TGFA and / or VAV3, and / or EGF or of their gene expression products is determined.
29. A DNA or RNA array comprising an arrangement of polynucleotides presented by or hybridizing to the following genes TNFRSF1B, DNAJC8, ECSIT, GOSR2, PPP1R9A, KLK6, C7orf46, SSH3 immobilized on a solid surface.
30. A gene array of claim 29 further comprising an arrangement of polynucleotides presented by or hybridizing to the following genes SERTAD2, AXIN2, C10orf99, ETS2, PITX2, PRR15, VAV3, gene coding for IKK interacting protein, EDEM3, LY6G6D.
31. A gene array of claim 29 or 30 comprising additionally polynucleotides hybridizing to the genes AREG and / or EREG and / or TGFA.

32. A DNA or RNA array consisting of an arrangement of polynucleotides presented by or hybridizing to the following genes TNFRSF1B, DNAJC8, ECSIT, GOSR2, PPP1R9A, KLK6, C7orf46, SSH3, SERTAD2, AXIN2, C10orf99, ETS2, PITX2, PRR15, VAV3, gene coding for IKK interacting protein, EDEM3, LY6G6D, AREG, EREG and TGFA.
- 5 33. A DNA or RNA array comprising an arrangement of polynucleotides immobilized on a solid surface, said polynucleotides being presented by or hybridizing to the genes arranged on the array, wherein the arrangement of polynucleotides is selected from the group consisting of:
- (a) ADAMEC1, C7orf46, CAST, DHCR7, ERAP1, FADS1, INSIG2, KLK6, MIDN,
 10 PHGDN, PPP1R9A, QPCT, RABEP1, RPL22L1, SLC25A27, SOCS6, SQLE, TNFRSF1B, TPD52, ZDHHC2,
- (b) GOLT1B, HSPA5, LEPROTL1, MAPK6, PROSC, RGMB, RWDD2B, SERTAD2, SPIRE2,
- (c) ASB6, ATM, BMI1, CDC42EP2, EDEM3, KCNK5, PGM2L1, RALBP1, SHROOM2,
 15 TNFSF15;
- (d) ACSL5, C1QC, CBALES1, CDK6, EHBP1, ETS2, EXOC6, EXT1, FMNL2, HDAC2, JUN, MED17, MTHFS, PITX2, POF1B, PRR15, PSMG1, RAB15, RAB40B, RNF43, SLC44A3, SOX4, TK2, TNFSF15, ZDHHC14,
- (e) ZDHHC14, TK2, PRR15, MTHFS, CBALES1, EHBP1, MED17, BMI1, CDC42EP2,
 20 EDEM3, PGM2L1, RGMB, ADAMEC1, ERAP1, FADS1, KLK6, PHGDH, PPP1R9A, QPCT, SLC25A27, TNFRSF1B, ZDHHC2.
34. A gene array of claim 33 comprising additionally polynucleotides presented by or hybridizing to the genes AREG and / or EREG and / or TGFA.
35. A kit for real-time PCR amplification of genetic anti-EGFR antibody biomarkers
 25 comprising a first package comprising the DNA or RNA of one or more of the genes of group (i) and / or group (ii) as specified in claim 1, a second package comprising PCR primers which specifically hybridize with said DNA/RNA molecules of said first package, a third package comprising a well-plate, and a fourth package comprising diagnostic means and solvents by means of which real-time PCR amplification can be carried out.
- 30 36. A kit of claim 35, wherein the first package comprises DNA/RNA of the following genes: TNFRSF1B, DNAJC8, ECSIT, GOSR2, PPP1R9A, KLK6, C7orf46, SSH3, SERTAD2, AXIN2, C10orf99, ETS2, PITX2, PRR15, VAV3, gene coding for IKK interacting protein, EDEM3, LY6G6D, AREG, EREG and TGFA, optionally in addition with AREG or EREG.

37. Use of one or more of the genetic biomarkers or a gene array or a kit for real-time PCR amplification of genetic anti-EGFR antibody biomarkers comprising one or more of said biomarkers selected from group consisting of

ADAMDEC1, BSDC1, C1orf144, CAPZB, CDC42, DHCR7, DNAJC8, ECSIT, EXOSC10,
 5 FADS1, GBA, GLT25D1, GOSR2, IMPDH1, KLHL21, KPNA6, KPNB1, LSM12,
 MAN1B1, MIDN, PPAN, SH3BP2, SQLE, SSH3, TNFRSF1B, URM1, ZFYVE26, RGMB,
 SPIRE2, ABCC5, ACSL5, AOA, AXIN2, CD24, CEACAM5, CEACAM6, ETS2, FMNL2,
 GPSM2, HDAC2, JUN, ME3, MED17, MYB, MYC, NEBL, NOSIP, PITX2, POF1B,
 PPARG, PPP1R14C, PRR15, PSMG1, RAB15, RAB40B, RNF43, RPS23, SLC44A3,
 10 SOX4, THEM2, VAV3, ZNF337, EPDR1, KCNK5, KHDRBS3, PGM2L1, STK38,
 SHROOM2;

C7orf46, CAST, DCP2, DIP2B, ERAP1, INSIG2, KIF21A, KLK6, NGRN, NRIP1,
 PHGDH, PPP1R9A, QPCT, RABEP1, RPA1, RPL22L1, SKP1, SLC25A27, SLC25A46,
 SOCS6, TPD52, ZDHHC2, ZNF654, ASB6, ATM, BMI1, CDC42EP2, EDEM3, PLLP,
 15 RALBP1, SLC4A11, TNFSF15, TPK1, C11orf9, C1QC, CABLES1, CDK6, EHBP1,
 EXOC6, EXT1, FLRT3, GCNT2, MTHFS, PIK3AP1, ST3GAL1, TK2, ZDHHC14,
 ARFGAP3, AXUD1, CAPZB, CHSY1, DNAJB9, GOLT1B, HSPA5, LEPROTL1, LIMS1,
 MAPK6, MYO6, PROSC, RAB8B, RAP2B, RWDD2B, SERTAD2, SOCS5, TERF2IP,
 TIAL1, TIPARP, TRIM8, TSC22D2, TGFA, VAPA, and UBE2K, optionally in combination
 20 with AREG and / or EREG,

or a respective protein expression product thereof, for predicting the pharmaceutical efficacy and / or clinical response of a patient suffering from KRAS wild type EGFR expressing cancer to an anti-EGFR antibody intended to be used for treatment, wherein said prediction results from calculating the differences in expression levels towards a
 25 threshold value to be determined from the underlying clinical determination parameters.

38. Use of claim 37, wherein at least one of the following biomarkers is used:

TNFRSF1B, DNAJC8, ECSIT, GOSR2, PPP1R9A, KLK6, C7orf46, SSH3, SERTAD2,
 AXIN2, C10orf99, ETS2, PITX2, PRR15, VAV3, gene coding for IKK interacting protein,
 EDEM3, LY6G6D, TGFA or combinations thereof including AREG and / or EREG .

39. Use of one or more of the genetic biomarkers or a gene array or a kit for real-time PCR amplification of genetic anti-EGFR antibody biomarkers comprising one or more of said biomarkers selected from group consisting of TNFRSF1B, DNAJC8, ECSIT, GOSR2,
 30 PPP1R9A, KLK6, C7orf46, SSH3, SERTAD2, AXIN2, C10orf99, ETS2, PITX2, PRR15,
 VAV3, gene coding for IKK interacting protein, EDEM3, LY6G6D, TGFA or combinations
 35 thereof including additionally AREG and / or EREG, for the manufacture of a

medicament which is an anti-EGFR antibody for the treatment of KRAS wild type EGFR expressing CRC or mCRC in a patient when one or more of said genes or gene products are expressed or overexpressed in a tumor sample of said patient.

- 5 40. Use of any of the claims 37 – 39, wherein said protein expression product is determined from a body fluid of the patient including plasma.
41. Use of any of the claims 37 – 40, wherein the intended underlying treatment is a first-line treatment.
- 10 42. Use of claim 37 – 41, wherein the intended underlying treatment is a combination treatment of said anti-EGFR antibody with a chemotherapeutic agent, and said patient has developed chemo-refractory cancer.
43. Use according to any of the claims 37 – 42, wherein the anti-EGFR antibody is c225 (cetuximab).
44. Use according to any of the claims 37 – 43, wherein the and the cancer is colorectal cancer (CRC) or metastatic colorectal cancer (mCRC).
- 15 45. Use of a monoclonal or polyclonal antibody which binds specifically to a protein expression product of a gene or a gene product as specified in any of the claims 37 to 39 for determining in vitro the pharmaceutical efficacy and / or clinical response of a patient suffering from KRAS wild type EGFR expressing cancer to an anti-EGFR antibody intended to be used for treatment.

Fig. 1: Genes whose expression in baseline samples is significantly associated with disease control after six weeks of cetuximab monotherapy in patients with wildtype KRAS gene. Based on study EMR 62202-045 (first-line treatment of metastatic CRC).

	p-value	log2-ratio	HUGO.GeneSymbol	EntrezGeneID	Description
206134_at	0,00031	1,76	ADAMDEC1	27299	ADAM-like, decysin 1
216008_s_at	2,00E-04	0,57	ARIH2	10425	ariadne homolog 2 (Drosophila)
218004_at	0,00068	0,66	BSDC1	55108	BSD domain containing 1
222200_s_at	0,0017	0,65	BSDC1	55108	BSD domain containing 1
212002_at	0,00065	1,08	C1orf144	26099	chromosome 1 open reading frame 144
228600_x_at	8,20E-05	-1,26	C7orf46	340277	chromosome 7 open reading frame 46
37012_at	0,00075	0,82	CAPZB	832	capping protein (actin filament) muscle Z-line, beta
212586_at	0,00057	-1,08	CAST	831	calpastatin
208908_s_at	0,0016	-1,12	CAST	831	calpastatin
208728_s_at	0,0012	0,63	CDC42	998	cell division cycle 42 (GTP binding protein, 25kDa)
204258_at	0,0019	-0,57	CHDI	1105	chromodomain helicase DNA binding protein 1
235258_at	0,00097	-0,62	DGP2	167227	DGP2 decapping enzyme homolog (S. cerevisiae)
201791_s_at	0,00094	1,19	DHCR7	1717	7-dehydrocholesterol reductase
224872_at	0,00035	-0,94	DIP2B	57609	DIP2 disco-interacting protein 2 homolog B (Drosophila)
205545_x_at	1,60E-06	0,79	DNAJC8	22826	DnaJ (Hsp40) homolog, subfamily C, member 8
218225_at	6,80E-06	0,86	ECST	51295	ECST homolog (Drosophila)
209788_s_at	0,00045	-1,46	ERAP1	51752	endoplasmic reticulum aminopeptidase 1
207541_s_at	0,0017	0,8	EXOSC10	5394	exosome component 10
208963_x_at	0,0014	1,61	FADS1	3992	fatty acid desaturase 1
210589_s_at	0,00098	0,96	GBA	2629	glucosidase, beta; acid (includes glucosylceramidase)
218473_s_at	0,0012	0,67	GLT25D1	79709	glycosyltransferase 25 domain containing 1
213144_at	3,90E-05	0,83	GOSR2	9570	golgi SNAP receptor complex member 2
204169_at	7,00E-04	0,86	IMPDH1	3614	IMP (inosine monophosphate) dehydrogenase 1
209566_at	0,00031	-1,11	INSIG2	51141	insulin induced gene 2
226003_at	0,0015	-0,9	KIF21A	55605	kinesin family member 21A
203068_at	0,0014	0,86	KLHL21	9903	kelch-like 21 (Drosophila)
204733_at	3,00E-05	-4,18	KLK6	5653	kalikrein-related peptidase 6
212101_at	9,00E-04	0,69	KPNA6	23633	karyopherin alpha 6 (importin alpha 7)
208974_x_at	0,0013	0,69	KPNB1	3837	karyopherin (importin) beta 1
212529_at	0,00035	0,82	LSM12	124801	LSM12 homolog (S. cerevisiae)
218636_s_at	0,00084	0,78	MAN1B1	11253	mannosidase, alpha, class 1B, member 1
225954_s_at	0,0012	1,08	MIDN	90007	midnolin

Fig 1 contd.

217722_s_at	0,00087	-0,66	NGRN	51335	neurin, neurite outgrowth associated
202599_s_at	0,0019	-0,61	NRIP1	8204	nuclear receptor interacting protein 1
201397_at	0,00077	-2,26	PHGDH	26227	phosphoglycerate dehydrogenase
221649_s_at	0,0013	0,81	PPAN	56342	peter pan homolog (Drosophila)
228494_at	8,90E-07	-4,71	PPP1R9A	55607	protein phosphatase 1, regulatory (inhibitor) subunit 9A
221088_s_at	0,00038	-3,28	PPP1R9A	55607	protein phosphatase 1, regulatory (inhibitor) subunit 9A
205174_s_at	0,0018	-2,03	QPCT	25797	glutaminyl-peptide cyclotransferase (glutaminyl cyclase)
225092_at	0,0018	-1,05	RABEP1	9135	rabaptin, RAB GTPase binding effector protein 1
201529_s_at	0,002	-0,74	RPA1	6117	replication protein A1, 70kDa
225541_at	0,0018	-1,37	RPL22L1	200916	ribosomal protein L22-like 1
209370_s_at	0,00069	0,62	SH3BP2	6452	SH3-domain binding protein 2
200719_at	0,00017	-0,94	SKP1	6500	S-phase kinase-associated protein 1
230624_at	0,00049	-1,47	SLC25A27	9481	solute carrier family 25, member 27
212833_at	2,00E-04	-0,79	SLC25A46	91137	solute carrier family 25, member 46
227542_at	0,00042	-1,31	SOC56	9306	suppressor of cytokine signaling 6
213577_at	0,00039	1,8	SQLE	6713	squalene epoxidase
219919_s_at	6,00E-05	0,46	SSH3	54961	slingshot homolog 3 (Drosophila)
203508_at	6,90E-07	2,08	TNFRSF1B	7133	tumor necrosis factor receptor superfamily, member 1B
201691_s_at	0,0019	-1,38	TPD52	7163	tumor protein D52
208101_s_at	3,00E-04	0,62	URM1	81605	ubiquitin related modifier 1 homolog (S. cerevisiae)
202682_s_at	0,0016	0,47	USP4	7375	ubiquitin specific peptidase 4 (proto-oncogene)
222731_at	0,00046	-1,91	ZDHHC2	51201	zinc finger, DHHC-type containing 2
37943_at	0,00084	0,59	ZFYVE26	23503	zinc finger, FYVE domain containing 26
241348_at	0,00099	-0,58	ZNF654	55279	zinc finger protein 654
224890_s_at	0,0013	0,63	NA	389541	similar to CG14977-PA

Fig. 2: Genes whose expression in baseline samples is significantly associated with disease control after six weeks of cetuximab monotherapy Based on study EMR 62202-045 (first-line treatment of metastatic CRC).

	p-value	log2-ratio	HUGO.GeneSymbol	EntrezGeneID	Description
221815_at	0,0013	-0,73	ABHD2	11057	abhydrolase domain containing 2
212738_at	0,00048	0,58	ARHGAP19	84986	Rho GTPase activating protein 19
227598_at	0,00047	-0,68	C7orf29	113763	chromosome 7 open reading frame 29
228600_x_at	0,0014	-0,88	C7orf46	340277	chromosome 7 open reading frame 46
226803_at	0,0011	-1,46	CHMP4C	92421	chromatin modifying protein 4C
205545_x_at	1,90E-05	0,53	DNAJC8	22826	DnaJ (Hsp40) homolog, subfamily C, member 8
236073_at	0,00065	-0,69	EPHA10	284656	EPH receptor A10
225023_at	0,0018	0,56	GOPC	57120	golgi associated PDZ and coiled-coil motif containing
212439_at	0,00088	0,43	HPK1	9807	inositol hexaphosphate kinase 1
209566_at	0,00047	-0,66	INSIG2	51141	insulin induced gene 2
205470_s_at	0,00024	-1,45	KLK11	11012	kalikrein-related peptidase 11
208974_x_at	0,00039	0,55	KPNB1	3837	karyopherin (importin) beta 1
212473_s_at	0,0011	0,77	MICAL2	9645	microtubule associated monooxygenase, calponin and LIM domain containing 2
225954_s_at	0,00025	0,71	MIDN	90007	midnolin
230518_at	0,0013	-0,76	MPZL2	10205	myelin protein zero-like 2
201710_at	0,002	1,01	MYBL2	4605	v-myb myeloblastosis viral oncogene homolog (avian)-like 2
213328_at	0,0011	0,89	NEK1	4750	NIMA (never in mitosis gene a)-related kinase 1
201158_at	0,00011	0,58	NMT1	4836	N-myristoyltransferase 1
225005_at	0,00017	0,42	PHF13	148479	PHD finger protein 13
201251_at	0,0013	-0,68	PKM2	5315	pyruvate kinase, muscle
228494_at	0,00036	-2,92	PPP1R9A	55607	protein phosphatase 1, regulatory (inhibitor) subunit 9A
235638_at	0,00024	-1,38	RASSF6	166824	Ras association (RalGDS/AF-6) domain family member 6
225541_at	0,00049	-0,95	RPL22L1	200916	ribosomal protein L22-like 1
209257_s_at	0,0016	1,01	SMC3	9126	structural maintenance of chromosomes 3
227542_at	0,0011	-0,86	SOC56	9306	suppressor of cytokine signaling 6
213577_at	0,0017	1,02	SQLE	6713	squalene epoxidase
222634_s_at	0,00078	0,94	TBL1XR1	79718	transducin (beta)-like 1 X-linked receptor 1
227112_at	0,0016	-0,74	TMCCI	23023	transmembrane and coiled-coil domain family 1
203508_at	0,00012	1,37	TNFRSF1B	7133	tumor necrosis factor receptor superfamily, member 1B
217975_at	0,0019	-0,93	WBP5	51186	WW domain binding protein 5
226124_at	0,0016	0,62	ZFP90	146198	zinc finger protein 90 homolog (mouse)
205182_s_at	0,0015	0,55	ZNF324	25799	zinc finger protein 324
227045_at	0,00022	0,87	ZNF614	80110	zinc finger protein 614

Fig. 3: Study EMR 62202-502 (cetuximab plus irinotecan treatment of irinotecan-refractory metastatic CRC patients), analysis of patients with wildtype KRAS gene: genes whose expression in baseline samples is significantly associated with best overall response.

	p-value	log2-ratio	HUGO.GeneSymbol	EntrezGeneID	Description
202211_at	0,00052	-0,76	ARFGAP3	26286	ADP-ribosylation factor GTPase activating protein 3
225557_at	0,00053	-0,74	AXUD1	64651	AXIN1 up-regulated 1
201949_x_at	0,0018	-0,65	CAPZB	832	capping protein (actin filament) muscle Z-line, beta
203044_at	0,0019	-0,87	CHSY1	22856	chondroitin sulfate synthase 1
202842_s_at	0,0018	-1,1	DNAJB9	4189	DnaJ (Hsp40) homolog, subfamily B, member 9
218193_s_at	0,0011	-1,08	GOLT1B	51026	golgi transport 1 homolog B (S. cerevisiae)
230031_at	0,002	-1,02	HSPA5	3309	heat shock 70kDa protein 5 (glucose-regulated protein, 78kDa)
202595_s_at	0,00071	-1,08	LEPROTL1	23484	leptin receptor overlapping transcript-like 1
207198_s_at	0,0017	-0,65	LIMS1	3987	LIM and senescent cell antigen-like domains 1
207121_s_at	0,0013	-1,08	MAPK6	5597	mitogen-activated protein kinase 6
201152_s_at	0,00097	-0,58	MBNL1	4154	muscleblind-like (Drosophila)
210480_s_at	0,0013	-0,92	MYO6	4646	myosin VI
202215_s_at	0,0011	-0,52	NFYC	4802	nuclear transcription factor Y, gamma
209384_at	0,00046	-1,12	PROSC	11212	proline synthetase co-transcribed homolog (bacterial)
209385_s_at	0,0011	-1,06	PROSC	11212	proline synthetase co-transcribed homolog (bacterial)
226633_at	0,0018	-0,94	RAB8B	51762	RAB8B, member RAS oncogene family
202362_at	0,00051	-0,56	RAP1A	5906	RAP1A, member of RAS oncogene family
213923_at	0,0017	-0,89	RAP2B	5912	RAP2B, member of RAS oncogene family
227339_at	0,0015	2,35	RGMB	285704	RGM domain family, member B
218377_s_at	0,0015	-1,4	RWDD2B	10069	RWD domain containing 2B
202656_s_at	0,00029	-1,16	SERTAD2	9792	SERTA domain containing 2
202657_s_at	8,00E-04	-0,98	SERTAD2	9792	SERTA domain containing 2
209648_x_at	0,00081	-0,73	SOC5	9655	suppressor of cytokine signaling 5
243594_x_at	0,00037	1,32	SPIRE2	84501	spire homolog 2 (Drosophila)
227706_at	0,00058	0,86	SPIRE2	84501	spire homolog 2 (Drosophila)
217790_s_at	0,0013	-0,56	SSR3	6747	signal sequence receptor, gamma (translocon-associated protein gamma)
201174_s_at	0,00063	-0,82	TERF2IP	54386	telomeric repeat binding factor 2, interacting protein
202405_at	0,0012	-0,78	TIAL1	7073	TIA1 cytotoxic granule-associated RNA binding protein-like 1
212665_at	0,0019	-0,99	TIPARP	25976	TCDD-inducible poly(ADP-ribose) polymerase
223132_s_at	0,0016	-0,62	TRIM8	81603	tripartite motif-containing 8
204094_s_at	0,00041	-0,88	TSC22D2	9819	TSC22 domain family, member 2
202347_s_at	0,00059	-0,85	UBE2K	3093	ubiquitin-conjugating enzyme E2K (UBC1 homolog, yeast)
208780_x_at	0,0018	-1,02	VAPA	9218	VAMP (vesicle-associated membrane protein)-associated protein A, 33kDa

Fig. 4: Study EMR 62202-502 (cetuximab plus irinotecan treatment of irinotecan-refractory metastatic CRC patients): genes whose expression in baseline samples is significantly associated with best overall response.

	p-value	log2-ratio	HUGO_GeneSymbol	EntrezGeneID	Description
203180_at	0,00065	-1,24	ALDH1A3	220	aldehyde dehydrogenase 1 family, member A3
202211_at	0,0016	-0,73	ARFGAP3	26286	ADP-ribosylation factor GTPase activating protein 3
201658_at	0,00028	-0,66	ARL1	400	ADP-ribosylation factor-like 1
225557_at	0,00065	-0,59	AXUD1	64651	AXIN1 up-regulated 1
223916_s_at	0,0017	-0,72	BCOR	54880	BCL6 co-repressor
211518_s_at	0,0016	-2,04	BMP4	652	bone morphogenetic protein 4
201848_s_at	0,0018	-1,13	BNIP3	664	BCL2/adenovirus E1B 19kDa interacting protein 3
229399_at	0,00098	-1,1	C10orf118	55088	chromosome 10 open reading frame 118
217893_s_at	0,0018	-0,76	C1orf108	79647	chromosome 1 open reading frame 108
203024_s_at	0,002	-0,57	C5orf15	56951	chromosome 5 open reading frame 15
223433_at	0,00053	0,94	C7orf36	57002	chromosome 7 open reading frame 36
208022_s_at	0,0015	0,92	CDC14B	8555	CDC14 cell division cycle 14 homolog B (S. cerevisiae)
202284_s_at	0,0015	-1,17	CDKN1A	1026	cyclin-dependent kinase inhibitor 1A (p21, Cip1)
213957_s_at	0,00089	-0,88	CEP350	9857	centrosomal protein 350kDa
212653_s_at	0,00096	-0,66	EHBP1	23301	EH domain binding protein 1
201143_s_at	0,0011	-0,69	EIF2S1	1965	eukaryotic translation initiation factor 2, subunit 1 alpha, 35kDa
202815_s_at	0,00028	-0,65	HEXIM1	10614	hexamethylene bis-acetamide inducible 1
218611_at	0,00044	-1,11	IER5	51278	immediate early response 5
209912_s_at	0,00046	0,49	KIAA0415	9907	KIAA0415
202595_s_at	0,00065	-1,04	LEPROTL1	23484	leptin receptor overlapping transcript-like 1
207121_s_at	0,00067	-0,9	MAPK6	5597	mitogen-activated protein kinase 6
200797_s_at	0,0012	-0,62	MCL1	4170	myeloid cell leukemia sequence 1 (BCL2-related)
213696_s_at	0,0017	-0,92	MED8	112950	mediator complex subunit 8
208919_s_at	0,0015	-0,38	NADK	65220	NAD kinase
203675_at	2,00E-04	-0,85	NUCB2	4925	nucleobindin 2
205552_s_at	0,0015	-1,37	OAS1	4938	2",5"-oligoadenylate synthetase 1, 40/46kDa
224701_at	0,00064	-1,15	PARP14	54625	poly (ADP-ribose) polymerase family, member 14
201939_at	0,0013	-1,33	PLK2	10769	polo-like kinase 2 (Drosophila)
209384_at	0,0015	-0,9	PROSC	11212	proline synthetase co-transcribed homolog (bacterial)
225074_at	0,00029	-0,79	RAB2B	84932	RAB2B, member RAS oncogene family
226633_at	0,0014	-0,8	RAB8B	51762	RAB8B, member RAS oncogene family
227339_at	0,0013	2,31	RGMB	285704	RGM domain family, member B
202129_s_at	0,0014	-0,92	RIOK3	8780	RIO kinase 3 (yeast)

Fig. 4 contd.

219684_at	0,0011	-1,1	RTP4	64108	receptor (chemosensory) transporter protein 4
228531_at	0,00066	-1,54	SAMD9	54809	sterile alpha motif domain containing 9
226603_at	0,00083	-1,53	SAMD9L	219285	sterile alpha motif domain containing 9-like
230036_at	0,0016	-1,02	SAMD9L	219285	sterile alpha motif domain containing 9-like
202657_s_at	0,001	-0,67	SERTAD2	9792	SERTA domain containing 2
202656_s_at	0,0016	-0,77	SERTAD2	9792	SERTA domain containing 2
201250_s_at	0,00023	-1,97	SLC2A1	6513	solute carrier family 2 (facilitated glucose transporter), member 1
217789_at	0,0014	-0,77	SNX6	58533	sorting nexin 6
243594_x_at	5,10E-05	1,31	SPIRE2	84501	spire homolog 2 (Drosophila)
227706_at	0,00014	0,91	SPIRE2	84501	spire homolog 2 (Drosophila)
209436_at	0,0016	-2,14	SPON1	10418	spondin 1, extracellular matrix protein
221489_s_at	0,0016	-1,61	SPRY4	81848	sprouty homolog 4 (Drosophila)
201174_s_at	0,00059	-0,71	TERF2IP	54386	telomeric repeat binding factor 2, interacting protein
203167_at	0,00052	-1,28	TIMP2	7077	TIMP metalloproteinase inhibitor 2
212665_at	0,001	-0,85	TIPARP	25976	TCDD-inducible poly(ADP-ribose) polymerase
214168_s_at	0,0015	-0,9	TJP1	7082	tight junction protein 1 (zona occludens 1)
210260_s_at	0,00022	-1,67	TNFAIP8	25816	tumor necrosis factor, alpha-induced protein 8
208296_x_at	0,0015	-1,28	TNFAIP8	25816	tumor necrosis factor, alpha-induced protein 8
223132_s_at	5,90E-05	-0,63	TRIM8	81603	tripartite motif-containing 8
202347_s_at	0,00082	-0,71	UBE2K	3093	ubiquitin-conjugating enzyme E2K (UBC1 homolog, yeast)
225925_s_at	0,0016	-0,55	USP48	84196	ubiquitin specific peptidase 48
208780_x_at	0,00049	-0,81	VAPA	9218	VAMP (vesicle-associated membrane protein)-associated protein A, 33kDa
225198_at	8,00E-04	-1,19	VAPA	9218	VAMP (vesicle-associated membrane protein)-associated protein A, 33kDa

Fig. 5: Study EMR 62202-502 (cetuximab plus irinotecan treatment in irinotecan-refractory metastatic CRC patients), analysis of patients with wildtype KRAS gene: genes whose expression in baseline samples is significantly associated with overall survival time.

	p-value	log2-ratio	HUGO.GeneSymbol	EntrezGeneID	Description
221657_s_at	0,0015	-1,83	ASB6	140459	ankyrin repeat and SOCS box-containing 6
208442_s_at	0,00089	-1,88	ATM	472	ataxia telangiectasia mutated
222696_at	0,0016	0,48	AXIN2	8313	axin 2 (conductin, axil)
202265_at	0,0017	-2,29	BMI1	648	BMI1 polycomb ring finger oncogene
227736_at	0,001	0,48	C10orf99	387695	chromosome 10 open reading frame 99
209850_s_at	0,00035	-2,92	CDC42EP2	10435	CDC42 effector protein (Rho GTPase binding) 2
223243_s_at	9,00E-04	-1,51	EDEM3	80267	ER degradation enhancer, mannosidase alpha-like 3
223253_at	0,00053	0,66	EPDR1	54749	ependymin related protein 1 (zebrafish)
219615_s_at	0,001	1,21	KCNK5	8645	potassium channel, subfamily K, member 5
209781_s_at	0,0013	0,88	KHDRBS3	10656	KH domain containing, RNA binding, signal transduction associated 3
207457_s_at	6,80E-05	0,49	LY6G6D	58530	lymphocyte antigen 6 complex, locus G6D
238417_at	0,00052	3,38	PGM2L1	283209	phosphoglucomutase 2-like 1
204519_s_at	0,0018	-0,56	PLLP	51090	plasma membrane proteolipid (plasmolipin)
226957_x_at	0,0016	-1,86	RALBP1	10928	ralA binding protein 1
204967_at	0,00037	1,03	SHROOM2	357	shroom family member 2
223748_at	0,0014	-0,78	SLC4A11	83959	solute carrier family 4, sodium borate transporter, member 11
1553117_a_at	0,0017	0,92	STK38	11329	serine/threonine kinase 38
221085_at	0,00047	-1,18	TNFSF15	9966	tumor necrosis factor (ligand) superfamily, member 15
221218_s_at	0,00059	-0,82	TPK1	27010	thiamin pyrophosphokinase 1
218806_s_at	6,70E-05	0,67	VAV3	10451	vav 3 guanine nucleotide exchange factor
218807_at	8,00E-04	0,39	VAV3	10451	vav 3 guanine nucleotide exchange factor
207705_s_at	1,50E-05	0,86	NA	22981	ninein-like
1554452_a_at	0,00048	1,38	NA	29923	hypoxia-inducible protein 2
218692_at	0,002	0,63	NA	55638	Golgi-localized protein

Fig. 6: Study EMR 62202-502 (cetuximab plus irinotecan treatment in irinotecan-refractory metastatic CRC patients): genes whose expression in baseline samples is significantly associated with overall survival time.

	p-value	log2-ratio	HUGO.GeneSymbol	EntrezGeneID	Description
213017_at	0,0019	-1,07	ABHD3	171586	abhydrolase domain containing 3
223253_at	0,00049	0,55	EPDR1	54749	ependymin related protein 1 (zebrafish)
226259_at	0,0016	-1,3	EXOC6	54536	exocyst complex component 6 formin-like
226184_at	0,0016	0,95	FMNL2	114793	2
219020_at	0,002	1,43	HS1BP3	64342	HCLS1 binding protein 3
226944_at	0,00062	1,3	HTRA3	94031	HtrA serine peptidase 3
219615_s_at	0,00031	1,02	KCNK5	8645	potassium channel, subfamily K, member 5
205968_at	0,00081	0,69	KCNS3	3790	potassium voltage-gated channel, delayed-rectifier, subfamily S, member 3
206006_s_at	0,00056	1,45	KIAA1009	22832	KIAA1009
223098_s_at	0,0011	-1,58	LONP2	83752	lon peptidase 2, peroxisomal nucleotide-binding oligomerization domain containing
221073_s_at	0,0018	1,4	NOD1	10392	1
243623_at	0,0017	0,81	SLC1A7	6512	solute carrier family 1 (glutamate transporter), member 7
223748_at	0,0011	-0,58	SLC4A11	83959	solute carrier family 4, sodium borate transporter, member 11
203343_at	0,00034	-1,25	UGDH	7358	UDP-glucose dehydrogenase
218806_s_at	0,00061	0,47	VAV3	10451	vav 3 guanine nucleotide exchange factor
207705_s_at	1,20E-05	0,67	NA	22981	ninein-like hypoxia-inducible protein
1554452_a_at	0,00019	1,21	NA	29923	2
205414_s_at	0,001	0,93	NA	9912	Rho-type GTPase-activating protein RICH2

Fig. 7: Study EMR 62202-502 (cetuximab plus irinotecan treatment in irinotecan-refractory metastatic CRC patients): genes whose expression in baseline samples is significantly associated with progression-free survival time.

	p-value	log2-ratio	HUGO.GeneSymbol	EntrezGeneID	Description
219623_at	0,00089	1,05	ACTR5	79913	ARP5 actin-related protein 5 homolog (yeast)
39248_at	0,0013	-0,4	AQP3	360	aquaporin 3 (Gill blood group)
204073_s_at	0,0015	-0,45	C11orf9	745	chromosome 11 open reading frame 9
206387_at	0,0014	0,77	CDX2	1045	caudal type homeobox 2
201328_at	0,0014	0,6	ETS2	2114	v-ets erythroblastosis virus E26 oncogene homolog 2 (avian)
226259_at	0,0017	-1,11	EXOC6	54536	exocyst complex component 6
219250_s_at	0,0013	-0,5	FLRT3	23767	fibronectin leucine rich transmembrane protein 3
230788_at	0,00045	-0,56	GCNT2	2651	glucosaminyl (N-acetyl) transferase 2, l-branching enzyme (l blood group)
219615_s_at	6,40E-05	0,97	KCNK5	8645	potassium channel, subfamily K, member 5
206006_s_at	0,00085	1,34	KIAA1009	22832	KIAA1009
203713_s_at	0,0017	0,96	LLGL2	3993	lethal giant larvae homolog 2 (Drosophila)
223098_s_at	0,0015	-1,39	LONP2	83752	lon peptidase 2, peroxisomal
202180_s_at	0,002	1,13	MVP	9961	major vault protein
207558_s_at	2,90E-05	1,34	PITX2	5308	paired-like homeodomain 2
204554_at	8,00E-04	1,21	PPP1R3D	5509	protein phosphatase 1, regulatory (inhibitor) subunit 3D
226961_at	0,00038	0,64	PRR15	222171	proline rich 15
203405_at	0,00022	0,82	PSMG1	8624	proteasome (prosome, macropain) assembly chaperone 1
204547_at	0,0018	0,84	RAB40B	10966	RAB40B, member RAS oncogene family
243623_at	0,00031	0,8	SLC1A7	6512	solute carrier family 1 (glutamate transporter), member 7
224959_at	0,00088	0,39	SLC26A2	1836	solute carrier family 26 (sulfate transporter), member 2
222513_s_at	0,001	0,66	SORBS1	10580	sorbin and SH3 domain containing 1
228038_at	0,0013	-0,36	SOX2	6657	SRY (sex determining region Y)-box 2
228867_at	0,0019	-1,15	TATDN3	128387	TatD DNase domain containing 3
218806_s_at	0,00013	0,5	VAV3	10451	vav 3 guanine nucleotide exchange factor
218807_at	0,00083	0,31	VAV3	10451	vav 3 guanine nucleotide exchange factor
209048_s_at	0,0019	0,8	ZMYND8	23613	zinc finger, MYND-type containing 8
1554452_a_at	0,00039	0,86	NA	29923	hypoxia-inducible protein 2
207705_s_at	0,00095	0,42	NA	22981	ninein-like
218507_at	0,0013	0,54	NA	29923	hypoxia-inducible protein 2
236249_at	0,0019	-1,12	NA	121457	IKK interacting protein

Fig. 8: Study EMR 62202-502 (cetuximab plus irinotecan treatment in irinotecan-refractory metastatic CRC patients), analysis of patients with wildtype KRAS gene: genes whose expression in baseline samples is significantly associated with progression-free survival time.

	p-value	regression coefficient	HUGO_GeneSymbol	EntrezGeneID	Description
226363_at	0,002		1 ABCC5	10057	ATP-binding cassette, sub-family C (CFTR/MRP), member 5
218322_s_at	0,0011		1,02 ACSL5	51703	acyl-CoA synthetase long-chain family member 5
205639_at	0,00062		0,95 AOA3	313	acyloxyacyl hydrolase (neutrophil)
39248_at	0,0017		-0,51 AQP3	360	aquaporin 3 (Gill blood group)
222696_at	5,10E-05		0,69 AXIN2	8313	axin 2 (conductin, axil)
227736_at	8,00E-04		0,47 C10orf99	387695	chromosome 10 open reading frame 99
204073_s_at	0,00013		-0,72 C11orf9	745	chromosome 11 open reading frame 9
225353_s_at	0,001		-1,27 C1QC	714	complement component 1, q subcomponent, C chain
225532_at	0,0013		-1,42 CABLES1	91768	Cdk5 and Abl enzyme substrate 1
208651_x_at	0,00088		0,73 CD24	934	CD24 molecule
209771_x_at	0,0018		0,82 CD24	934	CD24 molecule
216379_x_at	0,0019		0,78 CD24	934	CD24 molecule
224851_at	0,0015		-1,22 CDK6	1021	cyclin-dependent kinase 6
201884_at	0,00017		0,76 CEACAM5	1048	carcinoembryonic antigen-related cell adhesion molecule 5
203757_s_at	0,00042		0,55 CEACAM6	4680	carcinoembryonic antigen-related cell adhesion molecule 6 (non-specific cross reacting antigen)
211657_at	0,00064		0,69 CEACAM6	4680	carcinoembryonic antigen-related cell adhesion molecule 6 (non-specific cross reacting antigen)
228915_at	0,00038		0,45 DACH1	1602	dachshund homolog 1 (Drosophila)
212653_s_at	0,00043		-1,49 EHBP1	23301	EH domain binding protein 1
201328_at	7,20E-05		1,06 ETS2	2114	v-ets erythroblastosis virus E26 oncogene homolog 2 (avian)
226259_at	0,0018		-1,46 EXOC6	54536	exocyst complex component 6
201995_at	0,00031		-1,45 EXT1	2131	exostoses (multiple) 1
219250_s_at	0,00023		-0,75 FLRT3	23767	fibronectin leucine rich transmembrane protein 3
222853_at	0,0015		-0,55 FLRT3	23767	fibronectin leucine rich transmembrane protein 3
226184_at	0,0019		1,29 FMNL2	114793	formin-like 2
230788_at	0,00071		-0,75 GCNT2	2651	glucosaminyl (N-acetyl) transferase 2, l-branching enzyme (l blood group)
221922_at	0,00091		0,79 GPM2	29899	G-protein signaling modulator 2
201833_at	0,00099		1,81 HDAC2	3066	(AGS3-like, C. elegans)
201466_s_at	0,00081		1,03 JUN	3725	histone deacetylase 2
207457_s_at	0,0011		0,35 LY6G6D	58530	jun oncogene
204663_at	0,00097		0,73 ME3	10873	lymphocyte antigen 6 complex, locus G6D
					malic enzyme 3, NADP(+)-dependent, mitochondrial

Fig. 8 contd.

223115_at	0,00093	2,19	MED17	9440	mediator complex subunit 17
207480_s_at	0,00074	-0,53	MEIS2	4212	Meis homeobox 2
203433_at	0,0011	-1,47	MTHFS	10588	5,10-methylenetetrahydrofolate synthetase (5-
204798_at	0,00086	0,7	MYB	4602	formyltetrahydrofolate cyclo-ligase)
					v-myb myeloblastosis viral oncogene homolog (avian)
					v-myc myelocytomatosis viral
202431_s_at	0,0019	0,86	MYC	4609	oncogene homolog (avian)
203962_s_at	0,00021	0,77	NEBL	10529	nebullette
217950_at	0,00051	1,7	NOSIP	51070	nitric oxide synthase interacting protein
226459_at	0,001	-0,63	PIK3AP1	118788	phosphoinositide-3-kinase adaptor protein 1
207558_s_at	4,30E-05	1,85	PITX2	5308	paired-like homeodomain 2
219756_s_at	0,0015	1,18	POF1B	79983	premature ovarian failure, 1B
208510_s_at	0,001	0,68	PPARG	5468	peroxisome proliferator-activated receptor gamma
226907_at	0,00033	0,67	PPP1R14C	81706	protein phosphatase 1, regulatory (inhibitor) subunit 14C
226961_at	1,00E-06	1,49	PRR15	222171	proline rich 15
203405_at	0,00034	1,05	PSMG1	8624	proteasome (prosome, macropain) assembly chaperone 1
221810_at	0,00056	1,29	RAB15	376267	RAB15, member RAS oncogene family
204547_at	0,0011	1,2	RAB40B	10966	RAB40B, member RAS oncogene family
218704_at	0,00015	1,07	RNF43	54894	ring finger protein 43
227722_at	0,0017	0,8	RPS23	6228	ribosomal protein S23
228221_at	0,0016	1,03	SLC44A3	126969	solute carrier family 44, member 3
201416_at	0,0014	1,17	SOX4	6659	SRY (sex determining region Y)-box 4
213994_s_at	0,002	-0,46	SPON1	10418	spondin 1, extracellular matrix protein
208322_s_at	0,00087	-0,7	ST3GAL1	6482	ST3 beta-galactoside alpha-2,3-sialyltransferase 1
204565_at	0,0013	0,99	THEM2	55856	thioesterase superfamily member 2
204227_s_at	0,0015	-1,68	TK2	7084	thymidine kinase 2, mitochondrial
221085_at	0,00093	-0,88	TNFSF15	9966	tumor necrosis factor (ligand) superfamily, member 15
221218_s_at	0,00082	-0,68	TPK1	27010	thiamin pyrophosphokinase 1
218806_s_at	4,20E-05	0,72	VAV3	10451	vav 3 guanine nucleotide exchange factor
218807_at	2,00E-04	0,46	VAV3	10451	vav 3 guanine nucleotide exchange factor
219247_s_at	0,0017	-1,4	ZDHHC14	79683	zinc finger, DHHC-type containing 14
214760_at	0,0011	0,92	ZNF337	26152	zinc finger protein 337
236249_at	5,70E-05	-2,42	NA	121457	IKK interacting protein
218692_at	0,00016	0,95	NA	55638	Golgi-localized protein
227862_at	0,00057	-0,85	NA	388610	hypothetical LOC388610

Fig. 9: Affymetrix probe sets used to evaluate the degree of liver tissue contamination of tumor biopsies

Affymetrix element ID	Gene name	Predominantly expressed
60474_at	<i>FERMT1</i>	Colorectal cancer
218687_s_at	<i>MUC13</i>	Colorectal cancer
203397_s_at	<i>GALNT3</i>	Colorectal cancer
201650_at	<i>KRT19</i>	Colorectal cancer
217867_x_at	<i>BACE2</i>	Colorectal cancer
204771_s_at	<i>TTF1</i>	Colorectal cancer
219520_s_at	<i>WWC3</i>	Colorectal cancer
202869_at	<i>OAS1</i>	Colorectal cancer
206606_at	<i>LIPC</i>	Liver
207097_s_at	<i>SLC17A2</i>	Liver
210558_at	<i>AKR1C4</i>	Liver
210324_at	<i>C8G</i>	Liver
206753_at	<i>RDH16</i>	Liver
206505_at	<i>UGT2B4</i>	Liver
206535_at	<i>SLC2A2</i>	Liver
209976_s_at	<i>CYP2E1</i>	Liver

Fig. 10: Association of baseline expression data with disease control at week 6 in patients with *KRAS* wild-type tumors.

Affymetix probe set ID	Entrez GeneID	Gene Symbol	P value	q-value	Log ₂ ratio
200719_at	6500	<i>SKP1</i>	0.00017	0.27	-0.94
201397_at	26227	<i>PHGDH</i>	0.00077	0.4	-2.26
201529_s_at	6117	<i>RPA1</i>	0.002	0.52	-0.74
201691_s_at	7163	<i>TPD52</i>	0.0019	0.52	-1.38
201791_s_at	1717	<i>DHCR7</i>	0.00094	0.41	1.19
202599_s_at	8204	<i>NRIP1</i>	0.0019	0.52	-0.61
202682_s_at	7375	<i>USP4</i>	0.0016	0.5	0.47
203068_at	9903	<i>KLHL21</i>	0.0014	0.47	0.86
203508_at	7133	<i>TNFRSF1B</i>	6.90E-07	0.0068	2.08
204169_at	3614	<i>IMPDH1</i>	7.00E-04	0.4	0.86
204258_at	1105	<i>CHD1</i>	0.0019	0.52	-0.57
204733_at	5653	<i>KLK6</i>	3.00E-05	0.091	-4.18
205174_s_at	25797	<i>QPCT</i>	0.0018	0.52	-2.03
205545_x_at	22826	<i>DNAJC8</i>	1.60E-06	0.008	0.79
206134_at	27299	<i>ADAMDEC1</i>	0.00031	0.33	1.76
207541_s_at	5394	<i>EXOSC10</i>	0.0017	0.52	0.8
208101_s_at	81605	<i>URM1</i>	3.00E-04	0.33	0.62
208728_s_at	998	<i>CDC42</i>	0.0012	0.47	0.63
208908_s_at	831	<i>CAST</i>	0.0016	0.5	-1.12
208963_x_at	3992	<i>FADS1</i>	0.0014	0.47	1.61
208974_x_at	3837	<i>KPNB1</i>	0.0013	0.47	0.69
209370_s_at	6452	<i>SH3BP2</i>	0.00069	0.4	0.62
209566_at	51141	<i>INSIG2</i>	0.00031	0.33	-1.11
209788_s_at	51752	<i>ERAP1</i>	0.00045	0.34	-1.46
210589_s_at	2629	<i>GBA</i>	0.00098	0.41	0.96
212002_at	26099	<i>C1orf144</i>	0.00065	0.4	1.08
212101_at	23633	<i>KPNA6</i>	9.00E-04	0.41	0.69
212529_at	124801	<i>LSM12</i>	0.00035	0.33	0.82
212586_at	831	<i>CAST</i>	0.00057	0.38	-1.08
212833_at	91137	<i>SLC25A46</i>	2.00E-04	0.27	-0.79
213144_at	9570	<i>GOSR2</i>	3.90E-05	0.098	0.83
213577_at	6713	<i>SQLE</i>	0.00039	0.33	1.8
216008_s_at	10425	<i>ARIH2</i>	2.00E-04	0.27	0.57
217722_s_at	51335	<i>NGRN</i>	0.00087	0.41	-0.66
218004_at	55108	<i>BSDC1</i>	0.00068	0.4	0.66
218225_at	51295	<i>ECSIT</i>	6.80E-06	0.026	0.86
218473_s_at	79709	<i>GLT25D1</i>	0.0012	0.46	0.67
218636_s_at	11253	<i>MAN1B1</i>	0.00084	0.41	0.78
219919_s_at	54961	<i>SSH3</i>	6.00E-05	0.13	0.46
221088_s_at	55607	<i>PPP1R9A</i>	0.00038	0.33	-3.28
221649_s_at	56342	<i>PPAN</i>	0.0013	0.47	0.81

Fig. 10: contd.

222200_s_at	55108	<i>BSDC1</i>	0.0017	0.52	0.65
222731_at	51201	<i>ZDHHC2</i>	0.00046	0.34	-1.91
224872_at	57609	<i>DIP2B</i>	0.00035	0.33	-0.94
224890_s_at	389541	<i>NA</i>	0.0013	0.47	0.63
225092_at	9135	<i>RABEP1</i>	0.0018	0.52	-1.05
225541_at	200916	<i>RPL22L1</i>	0.0018	0.52	-1.37
225954_s_at	90007	<i>MIDN</i>	0.0012	0.46	1.08
226003_at	55605	<i>KIF21A</i>	0.0015	0.5	-0.9
227542_at	9306	<i>SOCS6</i>	0.00042	0.33	-1.31
228494_at	55607	<i>PPP1R9A</i>	8.90E-07	0.0068	-4.71
228600_x_at	340277	<i>C7orf46</i>	8.20E-05	0.16	-1.26
230624_at	9481	<i>SLC25A27</i>	0.00049	0.34	-1.47
235258_at	167227	<i>DCP2</i>	0.00097	0.41	-0.62
241348_at	55279	<i>ZNF654</i>	0.00099	0.41	-0.58
37012_at	832	<i>CAPZB</i>	0.00075	0.4	0.82
37943_at	23503	<i>ZFYVE26</i>	0.00084	0.41	0.59

Fig. 11. Forty-seven probe sets showing an association of on-treatment changes in expression from baseline to week 4 with best overall response.

Affymetix probe set ID	Entrez GeneID	Gene Symbol	P value	q-value	Log ₂ ratio
1554078_s_at	9093	DNAJA3	0.00097	0.55	-0.86
1559633_a_at	1131	CHRM3	0.0015	0.56	1.44
1563321_s_at	8028	MLLT10	0.0012	0.56	1.2
200653_s_at	801	CALM1	0.00059	0.55	0.88
201502_s_at	4792	NFKBIA	0.0014	0.56	0.95
201870_at	10953	TOMM34	0.002	0.56	-1.02
202388_at	5997	RGS2	0.0011	0.56	2.08
203542_s_at	687	KLF9	0.00077	0.55	2.03
204040_at	9781	RNF144A	0.0013	0.56	1.67
204676_at	25880	TMEM186	0.0013	0.56	-0.91
204985_s_at	79090	TRAPPC6A	0.00052	0.55	-1.1
205092_x_at	22890	ZBTB1	0.0017	0.56	1.16
206383_s_at	9908	G3BP2	0.0018	0.56	1
208994_s_at	9360	PPIG	0.0016	0.56	0.76
210933_s_at	6624	FSCN1	0.00063	0.55	0.86
211948_x_at	23215	BAT2D1	0.00038	0.55	0.71
212451_at	9728	NA	0.00022	0.55	1.15
212650_at	23301	EHBP1	0.00041	0.55	1.78
213956_at	9857	CEP350	0.00023	0.55	0.94
213998_s_at	10521	DDX17	0.0017	0.56	2.02
214055_x_at	23215	BAT2D1	0.00057	0.55	0.7
214081_at	57125	PLXDC1	0.0014	0.56	2.4
214374_s_at	8496	PPFIBP1	0.00098	0.55	1.53
215280_s_at	8541	PPFIA3	0.00048	0.55	-1.25
216392_s_at	11196	SEC23IP	0.00084	0.55	0.83
217813_s_at	10927	SPIN1	0.00057	0.55	0.89
218324_s_at	65244	SPATS2	0.001	0.56	0.6
218916_at	79724	ZNF768	0.002	0.56	-1.02
219001_s_at	79269	WDR32	0.0014	0.56	-1.23
219384_s_at	23536	ADAT1	0.00054	0.55	-1.27
219596_at	56906	THAP10	7.00E-04	0.55	-1.35
222457_s_at	51474	LIMA1	0.0019	0.56	1.17
222527_s_at	55696	RBM22	0.00053	0.55	1.13
222586_s_at	114885	OSBPL11	0.00032	0.55	0.9
222810_s_at	9462	RASAL2	0.00025	0.55	1.24
222890_at	29070	CCDC113	0.0016	0.56	-1.75
223259_at	94103	ORMDL3	0.00088	0.55	-0.81
223672_at	84251	SGIP1	0.0018	0.56	2.16
226987_at	29890	RBM15B	0.00091	0.55	-1.04
227148_at	130271	PLEKHH2	0.0018	0.56	1.83
229586_at	80205	CHD9	0.00061	0.55	1.34
230629_s_at	57634	EP400	0.00073	0.55	1.06

- 16/40 -

Fig. 11: contd.

235192_at	112858	TP53RK	0.00081	0.55	-1.28
235435_at	132949	AASDH	0.00065	0.55	0.92
237411_at	11174	ADAMTS6	0.00091	0.55	1.1
239147_at	153642	ARSK	0.00011	0.55	0.88
35436_at	2801	GOLGA2	0.0014	0.56	0.91

Fig. 12: . Association of on-treatment changes in candidate gene expression from baseline to week 4 with best overall response.

Affymetix probe set ID	Entrez GeneID	Gene Symbol	P value	Log ₂ ratio
1565483_at	1956	EGFR	0.37	0.45
201983_s_at	1956	EGFR	0.71	-0.13
201984_s_at	1956	EGFR	0.18	-0.58
216836_s_at	2064	ERBB2	0.06	-0.7
226213_at	2065	ERBB3	0.046	-0.68
202454_s_at	2065	ERBB3	0.011	-0.93
203821_at	1839	HBEGF	0.78	-0.1
205016_at	7039	TGFA	0.59	0.27
205239_at	374	AREG	0.39	-0.59
205767_at	2069	EREG	0.2	-1.17
38037_at	1839	HBEGF	0.59	0.19

Fig. 13: Association of baseline expression of candidate genes with disease control.

Affymetix probe set ID	Entrez GeneID	Gene Symbol	P value	Log ₂ ratio
1565483_at	1956	EGFR	0.81	0.1
201983_s_at	1956	EGFR	0.23	0.38
201984_s_at	1956	EGFR	0.42	0.34
216836_s_at	2064	ERBB2	0.27	-0.86
226213_at	2065	ERBB3	0.16	-0.81
202454_s_at	2065	ERBB3	0.14	-0.96
203821_at	1839	HBEGF	0.76	-0.13
205016_at	7039	TGFA	0.0073	-1.63
205239_at	374	AREG	0.13	0.99
205767_at	2069	EREG	0.099	1.45
38037_at	1839	HBEGF	0.97	-0.01

Fig. 14: contd.

EGF-R	-5.16	5.50E-06	0.00018	0.14	0.92	0.94	-0.18	0.93	0.99
ENA-78	0.26	0.24	0.41	-0.71	0.36	0.87	-0.89	0.34	0.93
Endothelin-1	-0.26	0.16	0.31	-0.69	0.1	0.68	-0.63	0.26	0.88
EN-RAGE	-0.1	0.65	0.78	-0.76	0.16	0.8	-0.63	0.33	0.93
Eotaxin	-0.04	0.87	0.89	-0.57	0.5	0.87	-1.11	0.26	0.88
Epregrulin	-0.07	0.43	0.62	0.32	0.15	0.8	0.4	0.19	0.81
Erythropoietin	0.12	0.49	0.68	0.06	0.8	0.93	0.25	0.5	0.98
Fatty Acid Binding Protein	0.3	0.032	0.1	0.18	0.53	0.87	0.02	0.95	0.99
Factor VII	-0.12	0.072	0.19	0.08	0.71	0.89	0.08	0.76	0.98
Ferritin	-0.37	0.0057	0.029	-0.34	0.39	0.87	-0.3	0.51	0.98
FGF basic	0.26	0.097	0.23	-0.48	0.032	0.34	-0.56	0.099	0.73
Fibrinogen	-0.14	0.029	0.098	-0.13	0.47	0.87	0.03	0.88	0.99
G-CSF	0.61	0.00013	0.0018	0.09	0.72	0.89	-0.24	0.52	0.98
Growth Hormone	-0.1	0.73	0.82	0.31	0.64	0.87	0.66	0.43	0.98
GM-CSF	0.33	0.11	0.24	0.05	0.91	0.94	-0.11	0.84	0.99
Glutathione S-Transferase	-0.12	0.27	0.45	-0.2	0.6	0.87	0.06	0.85	0.99
Haptoglobin	-0.35	0.1	0.24	-0.83	0.37	0.87	0.13	0.7	0.98
HB-EGF	-0.21	0.36	0.56	-0.02	0.97	0.97	0.31	0.78	0.98
ICAM-1	-0.07	0.16	0.32	-0.18	0.32	0.87	-0.11	0.49	0.98
IFN-gamma	0.1	0.08	0.21	0.15	0.26	0.87	0.08	0.66	0.98
IgA	-0.03	0.54	0.69	-0.06	0.64	0.87	0	0.98	0.99
IgE	0.02	0.82	0.87	-0.09	0.71	0.89	-0.18	0.59	0.98
IGF-1	0.98	0.036	0.11	3.15	0.014	0.19	2.32	0.068	0.69
IgM	-0.07	0.11	0.25	-0.04	0.78	0.92	-0.02	0.9	0.99
IL-10	0.38	0.0036	0.021	0.13	0.69	0.89	-0.27	0.49	0.98
IL-12p40	0.06	0.58	0.72	0.15	0.59	0.87	0.18	0.6	0.98
IL-12p70	-0.33	0.0074	0.034	0.14	0.55	0.87	-0.01	0.98	0.99
IL-13	0.4	0.0054	0.029	0.1	0.73	0.89	-0.03	0.94	0.99

Fig. 14: contd.

IL-15	0.13	0.099	0.23	0.27	0.18	0.8	0.12	0.54	0.98
IL-16	0.02	0.76	0.85	-0.13	0.6	0.87	-0.38	0.2	0.81
IL-18	0.24	0.00098	0.0097	-0.15	0.4	0.87	-0.18	0.34	0.93
IL-1alpha	0.2	0.0057	0.029	0.12	0.5	0.87	0.02	0.93	0.99
IL-1beta	0.11	0.48	0.67	-0.08	0.73	0.89	0.24	0.46	0.98
IL-1ra	0.52	0.0024	0.017	0.06	0.92	0.94	-0.35	0.6	0.98
IL-2	NA	NA	NA	NA	NA	NA	NA	NA	NA
IL-3	0.47	0.025	0.09	0.3	0.51	0.87	-0.14	0.79	0.99
IL-4	0.07	0.59	0.72	0.15	0.43	0.87	0.18	0.47	0.98
IL-5	0.34	0.17	0.33	-1.18	0.072	0.58	-1.46	0.079	0.69
IL-6	-0.21	0.36	0.56	-0.55	0.49	0.87	-0.42	0.65	0.98
IL-7	0.42	0.062	0.17	-0.12	0.72	0.89	-0.5	0.28	0.88
IL-8	-0.33	0.041	0.12	-1.33	0.0092	0.15	-1.66	0.0072	0.17
Insulin	1.07	0.0014	0.011	0.38	0.56	0.87	0.86	0.28	0.88
Leptin	-0.02	0.83	0.88	0.19	0.51	0.87	-0.04	0.89	0.99
Lipoprotein (a)	-0.28	0.00084	0.0097	0.16	0.46	0.87	0.15	0.55	0.98
Lymphotoctin	0.15	0.39	0.59	1.03	0.11	0.71	0.77	0.22	0.83
MCP-1	-0.02	0.81	0.87	-0.18	0.45	0.87	-0.41	0.17	0.81
MDC	-0.12	0.21	0.38	-0.3	0.38	0.87	-0.11	0.72	0.98
MIP-1alpha	0.02	0.89	0.89	-0.65	0.0061	0.15	-0.69	0.016	0.3
MIP-1beta	0.24	0.0013	0.011	-0.23	0.19	0.8	-0.4	0.075	0.69
MMP-2	0.39	0.086	0.21	-1.09	0.022	0.27	-0.94	0.044	0.6
MMP-3	0.19	0.15	0.31	0.52	0.18	0.8	0.12	0.69	0.98
MMP-9	0.47	0.026	0.09	0.18	0.66	0.88	-0.03	0.95	0.99
Myeloperoxidase	0.31	0.18	0.34	-0.32	0.63	0.87	-0.29	0.71	0.98
Myoglobin	0.08	0.55	0.69	0.38	0.2	0.8	-0.14	0.7	0.98
PAI-1	0.05	0.37	0.57	-0.1	0.29	0.87	-0.16	0.18	0.81
Prostatic Acid Phosphatase	0.33	0.00012	0.0018	-0.22	0.42	0.87	-0.48	0.15	0.81
PAPP-A	0.02	0.87	0.89	0.59	0.071	0.58	0.52	0.19	0.81


Fig. 14: contd.

Prostate Specific Antigen Free	0.4	0.012	0.052	0.18	0.49	0.87	0.1	0.75	0.98
RANTES	0.21	0.19	0.35	0.05	0.91	0.94	-0.09	0.86	0.99
Serum Amyloid P	-0.21	0.00011	0.0018	0.03	0.84	0.94	0.03	0.85	0.99
Stem Cell Factor	0.08	0.51	0.69	-0.18	0.33	0.87	-0.58	0.029	0.47
SGOT	-0.15	0.52	0.69	-0.09	0.91	0.94	0.03	0.97	0.99
SHBG	-0.07	0.28	0.46	0.15	0.42	0.87	0.2	0.28	0.88
Thyroxine Binding Globulin	-0.1	0.019	0.074	0.08	0.54	0.87	0.08	0.59	0.98
Tissue Factor	0.01	0.95	0.95	0.08	0.87	0.94	-0.26	0.45	0.98
		4.00E-06							
TGF-alpha	1.63	0.00018	0.00018	-0.7	0.35	0.87	-0.55	0.51	0.98
TIMP-1	-0.28	0.001	0.0097	-0.4	0.073	0.58	-0.49	0.064	0.69
TNF RII	-0.03	0.57	0.71	-0.08	0.61	0.87	-0.08	0.62	0.98
TNF-alpha	0.09	0.34	0.54	-0.22	0.31	0.87	-0.31	0.17	0.81
TNF-beta	0.05	0.72	0.82	0.21	0.37	0.87	0.39	0.088	0.7
Thrombopoietin	0.17	0.25	0.42	0.21	0.66	0.88	0.23	0.66	0.98
Thyroid Stimulating Hormone	-0.45	0.00013	0.0018	-0.5	0.19	0.8	-0.44	0.2	0.81
Thrombospondin-1	0.28	0.2	0.35	-0.07	0.89	0.94	-0.29	0.65	0.98
VCAM-1	-0.03	0.55	0.69	-0.03	0.8	0.93	-0.02	0.86	0.99
VEGF	-0.05	0.42	0.61	-0.03	0.87	0.94	-0.09	0.62	0.98
von Willebrand Factor	0.03	0.78	0.86	-0.38	0.27	0.87	-0.13	0.74	0.98

Fig. 15 Antibody reagents and immunohistochemistry assay conditions

Probe	Source	Antigen Retrieval	Dilution	Incubation time	Supplier	Code
EGFR	Mouse	Proteinase K	Ready to use	30 min	Dako	K1494
phospho-EGFR	Mouse	Water bath pH10	1:50	30 min	Chemicon	MAB3052
HER2	Rabbit	Water bath pH6	Ready to use	30 min	Dako	5207
phospho-HER2 (Tyr1248)	Rabbit	Microwave pH10	1:20	120 min	Cell Signaling Technology	2244
phospho-AKT (Ser473)	Rabbit	Microwave pH10	1:20	120 min	Cell Signaling Technology	9277L
phospho-MAPK (Thr202/Tyr204)	Rabbit	Microwave pH6	1:50	120 min	Cell Signaling Technology	9101L
p27 ^{Kip1}	Mouse	Microwave pH6	1:80	60 min	Dako	M7203
phospho-STAT3 (Tyr705)	Rabbit	Microwave pH6	1:100	120 min	Cell Signaling Technology	9131L
Ki67	Mouse	Microwave pH6	1:100	60 min	Dako	M7240

Fig. 16: Identification of RNA samples with high (grey), medium (dark grey) and low (black) liver contamination based on expression of genes predominantly expressed in colorectal cancer (upper box) and normal liver (lower box). The color scale reflects the absolute element signal intensity after normalization.



- 23/40 -

Fig. 17: Association of on-treatment changes in expression from baseline to week 4 with best overall response (partial response vs stable disease plus progressive disease).





Fig. 18: . Association of on-treatment changes in candidate gene expression from baseline to week 4 with best overall response (partial response vs stable disease plus progressive disease). Abbreviations; PD, progressive disease; SD, stable disease; PR, partial response.



- 25/40 -

Fig. 19: Association of baseline expression of candidate genes with disease control (partial response plus stable disease vs progressive disease) in tumors with *KRAS* wild-type status. Abbreviations; PD, progressive disease; SD, stable disease; PR, partial response.




Fig. 20:Correlation of *KRAS* status with response rate and progression-free survival

	Cetuximab monotherapy		Cetuximab plus FOLFIRI	
	Tumor <i>KRAS</i> status		Tumor <i>KRAS</i> status	
	Wild-type N=29	Mutation N=19	Wild-type N=29	Mutation N=19
Response, N (%)				
Complete response	0	0	0	0
Partial response	8 (28)	0	16 (55)	6 (32)
Stable disease	12 (41)	11 (58)	11 (38)	10 (53)
Progressive disease	9 (31)	8 (42)	0	3 (16)
Not evaluable	0	0	2 (7)	0
Overall response rate, N (%)	8 (28)	0 (-)	16 (55)	6 (32)
[95% CI]	[13 to 47]	[0 to 18]	[36 to 74]	[13 to 57]
<i>P</i> -value*	0.015		0.144	
Median PFS, months	-	-	9.4	5.6
[95% CI]			[7.0 to 11.3]	[3.3 to 12.2]
Hazard ratio			0.47	
<i>P</i> -value**			0.0475	

*Fisher's exact test; **logrank test

Abbreviation: PFS, progression-free survival


Fig. 21.: **A: Skin**

Fig. 21 (ctd.)




Fig. 21 (ctd.)



**B: Tumor:**

Fig. 21 (ctd.)




Fig. 22.






Fig. 23.

Fig. 24.

- 34/40 -

Fig. 24 (ctd.)








Fig. 24 (ctd.)




- 36/40 -

Fig. 25.**A**

- 37/40 -

Fig. 25 (ctd.)

B

- 38/40 -





Fig. 26:

Fig. 27:**Fig. 28**

- 40/40 -

Fig. 29:**Fig. 30:**