



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A01N 63/00	A2	(11) International Publication Number: WO 94/10845 (43) International Publication Date: 26 May 1994 (26.05.94)
(21) International Application Number: PCT/US93/11150 (22) International Filing Date: 16 November 1993 (16.11.93) (30) Priority data: 07/977,318 17 November 1992 (17.11.92) US (60) Parent Application or Grant (63) Related by Continuation US 07/977,318 (CIP) Filed on 17 November 1992 (17.11.92) (71) Applicant (for all designated States except US): CIBA-GEIGY AG [CH/CH]; Klybeckstrasse 141, CH-4002 Basle (CH).		(72) Inventors; and (75) Inventors/Applicants (for US only) : BECKER, J., Ole [DE/US]; 6164 Oswego Drive, Riverside, CA 92506 (US). TORKEWITZ, Nancy, R. [US/US]; 7301 Gates Road, Hurdle Mills, NC 27514 (US). MORTON, H., Vince [US/US]; 151 Olive Road, Reidsville, NC 27320 (US). (74) Agents: GLYNN, Michael, W. et al.; Ciba-Geigy Corporation, Patent Department, 7 Skyline Drive, Hawthorne, NY 10532 (US). (81) Designated States: AU, BR, CA, JP, KR, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: SYNERGISTIC MICROBICIDAL COMPOSITIONS (57) Abstract Microbicidal compositions which comprise, as active ingredients, an effective amount of at least one bacterial biocontrol strain of the genus <i>Pseudomonas</i> in combination with at least one synthetic microbicidal compound selected from the group consisting of halogenated nitrobenzenes, 3-phenylpyrrole derivatives, carboxin and thiram. The combination demonstrates synergistic effects which allow effective control using lower concentrations of microbicidal compounds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

Synergistic microbicidal compositions

The present invention relates to compositions comprising bacterial strains in combination with synthetic pesticidal compounds which are effective for the control of plant pests and pathogens.

More specifically, the invention relates to compositions which comprise, as active ingredients, an effective amount of at least one bacterial biocontrol strain of the genus Pseudomonas in combination with at least one synthetic microbicidal compound, together with a suitable carrier material, with the proviso that said synthetic microbicidal compound is not metalaxyl, furalaxyl, oxadixyl, benalaxyl or ofurace.

Such compositions exhibit synergistic effects and are useful in inhibiting plant pathogens, preferably phytopathogenic fungi, especially Rhizoctonia and Pythium species such as Rhizoctonia solani and Pythium ultimum.

In order to achieve effective control of fungal pathogens via biological control, it is important that a certain level of inhibitory bacteria be achieved. However, it may be difficult to obtain a level of bacteria sufficient to achieve effective control of plant pathogens through biocontrol alone. Furthermore, it may not be economically feasible to apply a large enough population of bacteria to achieve effective control. Thus, it is desirable to obtain combinations of bacterial biocontrol strains with other means for control of plant pathogens, particularly fungi, such as Rhizoctonia solani and Pythium ultimum, in which lower levels of Pseudomonas bacteria are able to effectively control the plant pathogens.

It has become known from EP-A-472,494 that purified bacterial strains of the species Pseudomonas fluorescens are effective for the inhibition of plant pathogens and that compositions comprising a microbicidal Pseudomonas fluorescens strain in combination with a synthetic microbicide of the acylalanine type, in particular metalaxyl, furalaxyl, oxadixyl, benalaxyl or ofurace, are useful for controlling fungi.

- 2 -

Surprisingly it has been found that compositions which comprise, as active ingredients, an effective amount of at least one bacterial biocontrol strain of the genus Pseudomonas in combination with at least one synthetic microbicidal compound, particularly selected from the group consisting of halogenated nitrobenzenes, 3-phenylpyrrole derivatives, carboxin and thiram display synergistic effects which allow for effective control using lower concentrations of microbicidal compounds, be that the Ps. organism or the microbicidal partner from chemical synthesis.

The combinations of the present invention are important for several reasons. First, Rhizoctonia solani is a particularly pernicious plant pathogen. The affected plants include ornamentals, vegetables, beans, wheat, tomato, potato and cotton. Secondly, presently available fungicide treatments are expensive, and may not provide efficiently for the protection of crops from Rhizoctonia solani. Therefore, the use of combinations of biocontrol agents and chemical microbicides to control or prevent pest infections in crop plants and other plants may provide an environmentally safe, economical and efficient method of control of plant pathogens such as Rhizoctonia solani. In addition, the compositions of the present invention can be used in mixtures in conjunction with bacterial strains that otherwise are not effective biocontrol agents for the particular plant pest. For example, the compositions can be used with strains that otherwise are not effective biocontrol agents for Rhizoctonia solani and Pythium ultimum and thereby increase the effective range of these biocontrol strains. The use of the biocontrol agents of the present invention in mixtures in order to improve the biocontrol capabilities of other strains of rhizosphere biocontrol agents is also a part of the present invention.

For example, US Patent No. 4,456,684, (Weller et al.) discloses that take-all, a disease of wheat caused by the fungus Gaeumannomyces graminis, can be controlled in some cases by the application of bacteria inhibitory to this pathogen to wheat seeds prior to planting. However, where the growth of G. graminis is effectively under control, R. solani may become a growing problem pathogen of wheat. Thus, the biocontrol agents of the present invention can be used together with biocontrol agents intended to protect wheat from take-all and extend their range of effectiveness to include R. solani and P. ultimum.

Methods are available in the art for selecting and isolating bacterial strains which can suppress and control plant pathogens; for Example EP-A- 472,494 (U.S. application Serial No. 705,424).

- 3 -

A number of such organisms are reported in the literature. See, for example, U.S. patent Nos. 4,900,348; 4,642,131; 4,975,277; 4,751,081; 4,714,614; 4,456,684; 4,647,533; 4,569,841; and 4,479,936; as well as EPA 0 376,775; EPA 0 353,689; WO91/05475; and EPA 0 200,344.

The structure of one of several antibiotic compounds produced by the biocontrol agent has been identified as that of the compound pyrrolnitrin. The structure of pyrrolnitrin and a proposed pathway for its biosynthesis from L-tryptophan is disclosed by Chang et al., J. Antibiot., 5: (1981). Pyrrolnitrin has previously been identified as an effective antifungal agent. See, for example, US Patent No.4,636,520.

The term 'biocontrol agent', as used throughout the present specification and the claims, encompasses purified biological disease controlling bacterial strains alone, referred to as biocontrol strains, or in combination with one or more synthetic chemical pesticides. The term biocontrol agent further includes the active compound produced by, or extracted from, the biocontrol strains, including antifungal, antibacterial, and other pesticidal metabolites, such as antibiotic compounds, whether used alone or in combination with one or more synthetic chemical compounds.

There is some uncertainty as to whether the Pseudomonas strains described herein belong to the species Pseudomonas fluorescens or Pseudomonas aurantiaca. In the present specification and the claims the term Pseudomonas fluorescens encompasses as well the species Pseudomonas aurantiaca.

Quintozone is the common name for pentachloronitrobenzene.

Carboxin is the common name for 5,6-dihydro-2-methyl-N-phenyl-1,4-oxanthiin-3-carboxamide.

Thiram is the common name for Bis(dimethylthio-carbamoyl)disulfide.

Metalaxyl is the common name for N-(2-methoxyacetyl)-N-(2,6-xylyl)-DL-alaninate.

Furalaxyl is the common name for N-(2-furoyl)-N-(2,6-xylyl)-DL-alaninate.

Oxadixyl is the common name for 2-methoxy-N-(2-oxo-1,3-oxazolidin-3-yl)acetat-2',6'-xylylide.

Benalaxyl is the common name for Methyl-N-phenylacetyl-N-2,6-xylyl-DL-alaninate.

Ofurace is the common name for α -2-chloroN-2,6-xylyl- γ -butyrolactone.

Preferred synthetic microbicidal compounds are selected from the group consisting of halogenated nitrobenzenes, 3-phenylpyrrole derivatives, carboxin and thiram.

- 4 -


The following preferred bacterial strains of Pseudomonas fluorescens have been deposited with the American Type Culture Collection in Rockville, Maryland on April 24, 1991:

CGA 266446 (ATCC Accession No. 55171);
 CGA 266447 (ATCC Accession No. 55170);
 CGA 267356 (ATCC Accession No. 55169);
 CGA 270293 (ATCC Accession No. 55175);
 CGA 270294 (ATCC Accession No. 55174); and
 CGA 281836 (ATCC Accession No. 55168).

Further preferred embodiments of this invention are:

(1) A composition, wherein the synthetic microbicidal compound is a halogenated nitrobenzene, in particular pentachloronitrobenzene (quintozene).

(2) A composition wherein the 3-phenylpyrrole derivative is of the formula



wherein

R₁ represents hydrogen, halogen, methyl, methoxy, trifluoromethyl, trifluoromethoxy,

R₂ represents halogen, trifluoromethyl or trifluoromethoxy, or

R₁ and R₂ together form a methylenedioxy, an ethylenedioxy or an ethylenoxy bridge, each of them being unsubstituted or substituted by methyl, chlorine or fluorine,

R₃ represents hydrogen or halogen,

X represents cyano, trifluoromethyl or COOCH₃, and

R represents hydrogen, C₁-C₄-acyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkoxy-C₁-C₄-acyl, C₂-C₄-alkenyloxycarbonyl or carbamoyl;

in particular a composition wherein X is cyano; or wherein

R₁ represents hydrogen, trifluoromethyl or halogen,

- 5 -

R₂ represents halogen or trifluoromethyl, or

R₁ and R₂ together form a difluoromethylendioxy bridge,

R₃ represents hydrogen,

X represents cyano and

R represents hydrogen, C₁-C₄-acyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkoxy-C₁-C₄-acyl, or C₂-C₄-alkenyloxycarbonyl.

Especially preferred thereof are the following 3-phenylpyrrole derivatives:

- a) 3-(2,3-dichlorophenyl)-4-cyanopyrrole (fenpiclonil).
- b) 3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole (fludioxonil);
- c) 1-acetyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole;
- d) 1-methoxyacetyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole;
- e) 1-methoxycarbonyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole;
- f) 1-allyloxycarbonyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole; and
- g) 1-n-propoxyacetyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole.

Particularly preferred is

3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole (fludioxonil).

(3) A composition wherein the synthetic microbicidal compounds are carboxin and/or thiram.

When used in combination with either of the Ps. strains synthetic microbicides can usually be applied in lower rates of application thereby still being sufficiently active.

The active ingredients of the present invention are normally applied in the form of compositions and can be applied to the crop area or plant to be treated, simultaneously or in succession, with other compounds. These compounds can be both fertilizers or micronutrient donors or other preparations that influence plant growth. They can also be selective herbicides, insecticides, fungicides, bactericides, nematocides, molluscides or mixtures of several of these preparations, if desired, together with further agriculturally acceptable carriers, surfactants or application-promoting adjuvants customarily employed in the art of formulation. Suitable carriers and adjuvants can be solid or liquid and correspond to the substances ordinarily employed in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, binders or fertilizers. Examples for microbicides are:

(E)-methyl 2-(2-(6-(2-cyanophenoxy)pyrimidin-4-yloxy)-phenyl)-3-methoxypropenoate, (+)-cis-1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)-oxiran-2-ylmethyl-1H-1,2,4-triazole, (RS)-1-aminopropyl- phosphonic acid, (RS)-4-(4-chloro-phenyl)-2-phenyl-2-(1H-1,2,4,-triazol-1-ylmethyl)butyronitrile, (Z)-N-but-2-enyloxymethyl-2-chloro-2',6'-diethylacetanilide, 1-(2-cyano-2-methoxy-iminoacetyl)-3-ethyl urea, 3-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-yl)quinazolin-4(3H)-one, 4-bromo-2-cyano-N,N-dimethyl-6-trifluoromethyl-benzimidazole-1-sulphonamide, 5-ethyl-5,8-dihydro-8-oxo-(1,3)-dioxol-(4,5-g) guinoline-7-carboxylic acid, a-[N-(3-chloro-2,6-xylyl)-2-methoxyacetamido]-butyrolactone, alanycarb, aldimorph, ampropylfos, anilazine, azaconazole, benomyl, biloxazol, binapacryl, bitertanol, blasticidin S, bromuconazole, bupirimate, butenachlor, buthiobate, captafol, captan, carbendazim, carbendazim chlorhydrate, chinomethionate, chlorbenzthiazole, chloroneb, chlorothalonil, chlorozolate, clozylacon, copper containing compounds such as copper oxychloride, copper oxyquinolate, copper sulphate and Bordeaux mixture, cycloheximide, cymoxanil, cyproconazole, di-2-pyridyl disulphide 1,1'-dioxide, dichlofluanid, dichlone, diclobutrazol, diclomezine, dicloran, didecyl dimethyl ammonium chloride, diethodencarb, difenoconazole, O,O-di-iso-propyl-S-benzyl thiophosphate, dimefluaazole, dimetconazole, dimethomorph, dimethirimol, diniconazole, dinocap, dipyrithione, ditalimfos, dithianon, dodemorph, dodine, doguadien, edifenphos, epoxyconazole, etaconazole, ethirimol, ethoxyguin, ethyl (2)-N-benzyle-N-([methyl(methyl-thioethylideneamino-oxy-carbon y l)amino]thio)-b-alaninate, etridiazole, fenaminosulph, fenapanil, fenarimol, fenbuconazole, fenfuram, fenpropidin, fenpropimorph, fentin acetate, fentin hydroxide, ferbam, ferimzone, fluazinam, fluoroimide, fluotrimazole, dilutolanil, flutriafol, flusilazole, folpet, fuberidazole, furconazole-cis, guazatine, hexaconazole, hydroxyisoxazole, hymexazole, imazalil, imibenconazole, ipconazole, iprobenfos, iprodione, isopropanyl butyl carbamate, isoprothiolane, kasugamycin, mancozeb, maneb, mepanipyrim, mepronil, methfuroxam, metiram, metiram-zinc, metsulfovax, myclobutanil, neoasozin, nickel dimethyldithiocarbamate, nitrothal-isopropyl, nuarimol, organomercury compounds, oxolinic acid, oxycarboxin, pefurazoate, penconazole, pencycuron, phenzin oxide, phosetyl-Al, phosphorus acids, phthalide, polyoxin D, polyram, probenazole, prochloraz, procymidone, propamocarb, propamocart hydrochloride, propiconazole, propineb, propionic acid, prothiocarb, pyracarbolid, pyrazophos, pyrifenox, pyroquilon, pyoxyfur, pyrrolnitrin, quaternary ammonium, compounds, quinconazole, quinomethionate, rabenazole, sodium pentachlorophenolate, streptomycin, sulphur, tebuconazole, techlofthalam, tecnazene,

- 7 -

tetraconazole, thiabendazole, thicarbamil, thicyofen,
 2-(thio-dyanomethylthio)benzothiazole thiophanate-methyl, thiram, thimibenconazole,
 tolclofos-methyl, tolylfluanid, triacetate salt of 1,1'-iminodi-(octamethylene)diguanidine,
 triadimefon, triadimenol, triazbutyl, triazoxide, tricyclazole, tridemorph, triforine,
 triflumizole, triticonazole, validamycin A, vapam, vinclozolin, zineb, ziram, terbufos,
 carbofuran, chloropicrin, ethyl 3-methyl-4-(methylthio)phenyl (1-methylethyl)-
 phosphoramidate, ethoprop, aldicarb [2-methyl-2-(methylthio)-propionaldehyde
 O-(methylcarbamoyl)oxime], and oxamyl [methyl-N'-N'-dimethyl-N-[methylcarbamoyl]-
 oxy]-1-thiooxamimidate].

Favorable mixing ratios of the two active principles bacterial biocontrol strain (= I) and the chemical compound (= II) are: I:II = 100:1 to 1:100, preferably 20:1 to 1:20. Other preferred ratios are 10:1 to 1:10; 5:1 to 1:5; 7:2 to 2:3.

The compositions of the present invention are effective, for example, against the phytopathogenic fungi belonging to the following classes: Ascomycetes, e.g., *Fusarium*; Basidiomycetes, e.g., *Rhizoctonia*; Oomycetes belonging to the class of Phycomycetes, e.g., *Phytophthora*, *Pythium* and *Plasmopara*. As plant protective agents, the compositions of the present invention can be used against important noxious fungi of the Fungi imperfecti family, e.g., *Cercospora* and *Botrytis*. *Botrytis* and the gray mould on vines, strawberries, apples, onions and other varieties of fruit and vegetables are a source of considerable economic damage. Thus, the compositions of the present invention may be particularly useful because they demonstrate excellent microbicidal activity against a wide spectrum of fungi. They control mold fungi such as *Penicillium*, *Aspergillus*, *Rhizopus*, *Fusarium*, *Helminthosporium*, *Nigrospora* and *Alternaria*, as well as bacteria such as butyric acid bacteria and yeast fungi such as *Candida*. Furthermore, the combinations of the present invention have excellent activity against fungi which occur in seeds or in the soil. As plant protective agents, the combinations are advantageous for practical application in agriculture for protecting cultivated plants, without damaging said plants by harmful side-effects.

Target crops to be protected within the scope of the present invention comprise e.g., the following species of plants: cereals (wheat, barley, maize, rye, oats, rice, sorghum and related crops), beet (sugar beet and fodder beet), pomes, drupes and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries), leguminous plants (beans, lentils, peas, soybeans), oil plants (rape, mustard, poppy, olives,

- 8 -

sunflowers, coconut, castor oil plants, cocoa beans, groundnuts), cucumber plants (cucumber, marrows, melons), fibre plants (cotton, flax, hemp, jute), citrus fruit (oranges, lemons, grapefruit, mandarins), vegetables (spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, paprika), lauraceae (avocados, cinnamon, camphor), or plants such as maize, tobacco, nuts, coffee, sugar cane, tea, vines, hops, bananas and natural rubber plants, as well as ornamentals (composites). The compositions may also be useful for storage protection of natural substances which are in freshly harvested or further processed form.

The biocontrol agents may be applied in any method known for treatment of seed or soil with bacterial strains. For example, see US Patent No. 4,863,866. The strains are effective for biocontrol even if the bacterium is not living. Preferred is, however, the application of the living bacterium.

The active ingredients may be used in unmodified form or together with any suitable agriculturally acceptable carrier. Such carriers are adjuvants conventionally employed in the art of agricultural formulation, and are therefore formulated in known manner to emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granulates, and also encapsulations, for example, in polymer substances. Like the nature of the compositions, the methods of application, such as spraying, atomizing, dusting, scattering or pouring, are chosen in accordance with the intended object and the prevailing circumstances. Advantageous rates of application are normally from about 50 g to about 5 kg of active ingredient (a.i.) per hectare ("ha", approximately 2.471 acres), preferably from about 100 g to about 2 kg a.i./ha. Important rates of application are about 200 g to about 1 kg a.i./ha and 200 g to 500 g a.i./ha.

For seed dressing advantageous application rates are 0.5 g to 1000 g a.i. per 100 kg seed, preferably 3 g to 100 g a.i. per 100 kg seed or 10 g to 50 g a.i. per 100 kg seed.

Preferred methods of applying an active ingredient of the present invention or an agrochemical composition of the present invention are leaf application, seed coating and soil application. The number of applications and the rate of application depend on the intensity of infestation by the corresponding pathogen (type of fungus). However, the active ingredients can also penetrate the plant through their roots via the soil (systemic action) by impregnating the locus of the plant with a liquid composition, or by applying the compounds in solid form to the soil, e.g. in granular form (soil application). The

- 9 -

active ingredients may also be applied to seeds (coating) by impregnating the seeds either with a liquid formulation containing active ingredients, or coating them with a solid formulation. In special cases, further types of application are also possible, for example, selective treatment of the plant stems or buds.

The formulations, compositions or preparations containing the active ingredients and, where appropriate, a solid or liquid adjuvant, are prepared in known manner, for example by homogeneously mixing and/or grinding the active ingredients with extenders, for example solvents, solid carriers and, where appropriate, surface-active compounds (surfactants).

Suitable solvents for compositions including those that contain the pesticidal metabolites produced by the biocontrol bacterial strains of the present invention include aromatic hydrocarbons, preferably the fractions having 8 to 12 carbon atoms, for example, xylene mixtures or substituted naphthalenes, phthalates such as dibutyl phthalate or dioctyl phthalate, aliphatic hydrocarbons such as cyclohexane or paraffins, alcohols and glycols and their ethers and esters, such as ethanol, ethylene glycol monomethyl or monethyl ether, ketones such as cyclohexanone, strongly polar solvents such as N-methyl-2-pyrrolidone, dimethyl sulfoxide or dimethyl formamide, as well as epoxidized vegetable oils such as epoxidized coconut oil or soybean oil; or water.

The solid carriers used e.g. for dusts and dispersible powders, are normally natural mineral fillers such as calcite, talcum, kaolin, montmorillonite or attapulgite. In order to improve the physical properties it is also possible to add highly dispersed silicic acid or highly dispersed absorbent polymers. Suitable granulated adsorptive carriers are porous types, for example pumice, broken brick, sepiolite or bentonite; and suitable nonsorbent carriers are materials such as calcite or sand. In addition, a great number of pregranulated materials of inorganic or organic nature can be used, e.g. especially dolomite or pulverized plant residues.

Depending on the nature of the active ingredient to be used in the formulation, suitable surface-active compounds are nonionic, cationic and/or anionic surfactants having good emulsifying, dispersing and wetting properties. The term "surfactants" will also be understood as comprising mixtures of surfactants.

Suitable anionic surfactants can be both water-soluble soaps and water-soluble synthetic

- 10 -

surface-active compounds.

Suitable soaps are the alkali metal salts, alkaline earth metal salts or unsubstituted or substituted ammonium salts of higher fatty acids (chains of 10 to 22 carbon atoms), for example the sodium or potassium salts of oleic or stearic acid, or of natural fatty acid mixtures which can be obtained for example from coconut oil or tallow oil. The fatty acid methyltaurin salts may be used.

More frequently, however, so-called synthetic surfactants are used, especially fatty sulfonates, fatty sulfates, sulfonated benzimidazole derivatives or alkylarylsulfonates.

The fatty sulfonates or sulfates are usually in the form of alkali metal salts, alkaline earth metal salts or unsubstituted or substituted ammonium salts and have a 8 to 22 carbon alkyl radical which also includes the alkyl moiety of alkyl radicals, for example, the sodium or calcium salt of lignonsulfonic acid, of dodecylsulfate or of a mixture of fatty alcohol sulfates obtained from natural fatty acids. These compounds also comprise the salts of sulfuric acid esters and sulfonic acids of fatty alcohol/ethylene oxide adducts. The sulfonated benzimidazole derivatives preferably contain 2 sulfonic acid groups and one fatty acid radical containing 8 to 22 carbon atoms. Examples of alkylarylsulfonates are the sodium, calcium or triethanolamine salts of dodecylbenzenesulfonic acid, dibutyl-naphthalene-sulfonic acid, or of anaphthalenesulfonic acid/formaldehyde condensation product. Also suitable are corresponding phosphates, e.g. salts are preferably in the form of halides, methylsulfates or ethylsulfates, e.g. stearyltrimethylammonium chloride or benzyldi(2-chloroethyl) ethylammonium bromide or salts of the phosphoric acid ester of an adduct of p-nonyl-phenol with 4 to 14 moles of ethylene oxide.

Non-ionic surfactants are preferably polyglycol ether derivatives of aliphatic or cycloaliphatic alcohols, or saturated or unsaturated fatty acids and alkylphenols, said derivatives containing 3 to 30 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon moiety and 6 to 18 carbon atoms in the alkyl moiety of the alkylphenols.

Further suitable non-ionic surfactants are the water-soluble adducts of polyethylene oxide with polypropylene glycol, ethylenediamine propylene glycol and alkylpolypropylene glycol containing 1 to 10 carbon atoms in the alkyl chain, which adducts contain 20 to 250 ethylene glycol ether groups and 10 to 100 propylene glycol ether groups. These

- 11 -

compounds usually contain 1 to 5 ethylene glycol units per propylene glycol unit.

Representative examples of non-ionic surfactants are nonylphenolpolyethoxyethanols, castor oil polyglycol ethers, polypropylene/polyethylene oxide adducts, tributylphenoxy-polyethoxyethanol, polyethylene glycol and octylphenoxyethoxyethanol. Fatty acid esters of polyoxyethylene sorbitan and polyoxyethylene sorbitan trioleate are also suitable non-ionic surfactants.

Cationic surfactants are preferably quaternary ammonium salts which have, as N-substituent, at least one C₈-C₂₂alkyl radical and, as further substituents, lower unsubstituted or halogenated alkyl, benzyl or lower hydroxyalkyl radicals. The salts are preferably in the form of halides, methylsulfates or ethylsulfates, e.g. stearyltrimethylammonium chloride or benzyldi(2-chloroethyl)ethylammonium bromide.

The surfactants customarily employed in the art of formulation are described, for example, in "McCutcheon's Detergents and Emulsifiers Annual, "MC Publishing Corp. Ringwood, New Jersey, 1979, and Sisely and Wood, "Encyclopedia of Surface Active Agents, "Chemical Publishing Co., Inc. New York, 1980.

The agrochemical compositions usually contain from about 0.1 to about 99%, preferably about 0.1 to about 95%, and most preferably from about 3 to about 90% of the active ingredient, from about 1 to about 99.9%, preferably from about 1 to about 99%, and most preferably from about 5 to about 95% of a solid or liquid adjuvant, and from about 0 to about 25%, preferably about 0.1 to about 25%, and most preferably from about 0.1 to about 20% of a surfactant.

Whereas commercial products are preferably formulated as concentrates, the end user will normally employ dilute formulations.

Formulation Examples

In the following examples, percentages of composition are given by weight. The active ingredient is a combination of a Pseudomonas fluorescens strain CGA 266446, CGA 266447, CGA 267356, CGA 270293, CGA 270294 or CGA 281836 with a synthetic chemical fungicide such as a halogenated nitrobenzene, a 3-phenylpyrrole derivative, carboxin or thiram. [cfu = colony-forming units]

- 12 -

<u>Dusts</u>	a)	b)
Active ingredient (incl. 10^{13} cfu/kg)	5 %	6 %
Talcum	85 %	-
Kaolin	-	84%
Residual moisture	10 %	10 %

Ready-to-use dusts are obtained by mixing the active ingredient with the carrier. Such powders may be used as a composition for dry-dressing seeds.

Extruder granules

Active ingredient	10 % (10^{13} cfu/kg)
Sodium ligninsulfonate	2 %
Carboxymethylcellulose	1 %
Kaolin	77 %
Moisture	10 %

The active ingredient (organism + chemical) is mixed with the additives, and the mixture is ground and moistened with water. This mixture is extruded and subsequently dried in a stream of air.

Coated granules

Active ingredient	3 % (10^{13} cfu/kg)
Polyethylene glycol (MW 200)	3 %
Kaolin	84 %
Water	10 %
(MW = molecular weight)	

In a mixer, the active ingredient is applied uniformly to the kaolin which has been moistened with polyethylene glycol. In this manner, dust-free coated granules are obtained.

Suspension concentrate

Active ingredient	40 % (10^{10} cfu/kg)
Propylene glycol	10 %
Nonylphenol polyethylene glycol ether	6 %

- 13 -

(15 moles of ethylene oxide)	
Sodium ligninsulfonate	10 %
Carboxymethyl cellulose	1 %
Silicone oil	1 %
(in the form of a 75 % aqueous emulsion)	
Water	32 %

The active ingredient (organism + chemical) is mixed intimately with the additives. This gives a suspension concentrate from which suspensions of any desired lower dilution can be prepared by dilution with water. Live plants and plant propagation material can be treated and protected from infestation with microorganisms by spraying them with such dilutions, pouring such dilutions or immersing the live plants or plant propagation material in such dilutions.

Biological Examples

If the activity achieved by said combinations is decisively greater than the activity to be expected by adding together the activities of the components individually, there exists a synergistic effect of the combination.

In the following examples,

* = indicates the expected value of the additive effects of two treatments in the absence of synergy, as computed by the method described in Richer, Pestic. Sci. 19:309-315 (1987).

** = indicates that the result exceeds the expected value of the additive effects of the individual treatments, and therefore is considered to be synergistic.

Example 1: Synergistic Effects of Combinations of Bacterial Biocontrol Strain CGA 267356 with Fludioxonil.

Soil is prepared by mixing potting soil (Metro-mix 360, WR Grace) with vermiculite and sand. 15 cm x 15 cm pots are filled with the soil mix, and 3/4" deep furrows are made near the perimeter of each pot. Thus, each pot contains a circular 12" furrow.

Stoneville 506 cotton seeds are obtained from the CIBA-GEIGY Delta Research Station. Fludioxonil is prepared as a 50 WP(50% wettable powder). 60 mg of powder is

- 14 -

resuspended as a 1 ml slurry which is used to coat 60 g of seeds. This is equal to an application rate of 50 g ai/100 kg seeds, which is defined as the 1x treatment for Fludioxonil. Dilutions of this treatment are made by resuspending lesser amounts of the powder in the slurry used to coat the seed (e.g., 30 mg powder/60 g seed = 25 g ai/100 kg seed = 1/2x treatment). The treated seeds are allowed to air dry for about one hour.

Strain CGA267356 is grown overnight in Luria broth at 28°C. Cells are centrifuged, then resuspended in sterile H₂O, and diluted in various proportions to give a range of cfu/ml dilutions. The 1x rate is defined as 20 ml of cell suspension at a concentration of 2×10^9 colony-forming units (cfu)/ml/ft.

Combinations of bacterial strains and Fludioxonil are tested as follows. In one set of experiments, the chemical is applied as a seed treatment, then bacteria are applied as an in-furrow drench. Ten seeds, either treated or untreated, are placed in each furrow. Dried ground *Rhizoctonia*-infested millet powder is sprinkled evenly over the seeds in the furrows. 20 ml of Strain CGA267356 suspensions or sterile H₂O are pipetted over the appropriate treatments. Each treatment consists of three or four replicate pots for a total of 30 or 40 seeds per treatment. Additional experiments are conducted in which both chemical and bacteria are applied as seed treatments. First, the seeds are treated with fenpiclonil, as described above. 4 g of seeds are then coated either with 1 ml of sterile H₂O or with a 1 ml suspension of Strain CGA267356 at a concentration of about 5×10^9 cfu/ml, or approximately 10^8 cfu/seed. The seeds are planted as described above, except that no additional water is pipetted over the seeds in the furrows.

The experiments are conducted in a phytotron set to a day/night temperature of 26/21°C. All pots are watered on day 7 and day 13 after planting.

Approximately two weeks (16 days) after planting, all plants are gently removed from the pots and the number of symptom-free plants is recorded. For each experiment, the percent control is calculated by comparison to the healthy check (no pathogen added, defined as 100% symptom-free) and the disease check (pathogen added, H₂O treated only, defined as 0% symptom-free). In these experiments, both pre-and post-emergence damping-off occurs.

- 15 -

Table 1

Control of Damping off of Cotton caused by *Rhizoctonia solani*: Synergism between Fludioxonil (Seed treatment) and Strain CGA267356 (Bacterial drench).

Experiment 1	% Disease Control	
<u>Treatment</u>	<u>Expected Value*</u>	<u>Actual Value</u>
Disease check		0
CGA 267356 1x		55
Fludioxonil 1/2x		34
CGA267356 1x + Fludioxonil 1/2x	70	86**

Experiment 2: Effect of Reduced Concentrations of Bacterial Drench

<u>Treatment</u>	<u>Expected Value*</u>	<u>Actual Value</u>
Disease check		0
Fludioxonil 1/2x		49
CGA267356 1x		98
CGA267356 1/10x		91
CGA267356 1/100x		56
Fludioxonil 1/2x + CGA267356 1/10x	95	98**
Fludioxonil 1/2x + CGA267356 1/100x	77	95**

Experiment 3: Effect of Reduced Concentrations of Chemical Seed Treatment

<u>Treatment</u>	<u>Expected Value*</u>	<u>Actual Value</u>
Disease check		0
CGA267356 1/10x		61
Fludioxonil 1x		33
Fludioxonil 1/3x		22
Fludioxonil 1/4x		6
CGA267356 1/10x + Fludioxonil 1x	74	96**
CGA267356 1/10x + Fludioxonil 1/3x	70	88**
CGA267356 1/10x + Fludioxonil 1/4x	65	93**

- 16 -

Table 2

Control of Damping off of Cotton caused by Rhizoctonia solani: Synergism between Fludioxonil (Seed treatment) and CGA 267356 (Seed treatment).

<u>Treatment</u>	<u>E Value*</u>	Trial 1	<u>E Value*</u>	Trial 2
		<u>Actual Value</u>		<u>Actual Value*</u>
Healthy check		100		100
Disease check		0		0
Fludioxonil 1/2x		37		45
CGA267356#		0		18
Fludioxonil 1/2x + CGA267356	37	52**	55	72**

= application rate for Strain CGA267356 seed treatment is approximately 10^8 cfu/seed.

Example 2: Synergistic Effects of Combinations of Bacterial Biocontrol Strain CGA 267356 with Fludioxonil under Field Conditions.

Combinations of CGA 267356 and Fludioxonil are tested under field conditions in Sanger, California, Vero Beach, Florida and Dewey, Illinois as follows. Cotton seeds are treated at the rates of 25 or 15 g ai/100 kg seed with Fludioxonil. Green bean seeds are treated with 25g, 7.5g, or 3.75g ai/100kg seed. The treated seeds are allowed to air dry for about one hour.

Strain CGA267356 is grown overnight in Luria broth at 28°C. Cells are centrifuged, then resuspended in sterile H₂O, and diluted in various proportions to give a range of cfu/ml dilutions. The 1x rate is defined as 100 ml of cell suspension at a concentration of 4×10^9 colony-forming units (cfu)/ml/10 ft.

Combinations of bacterial strains and a 3-phenyl-4-cyanopyrrole compound are tested as follows. The chemical (e.g. Fludioxonil or fenpiclonil) is applied as a seed treatment, then bacteria are applied as an in-furrow drench. 100 seeds, either treated or untreated, are placed in each 10 foot furrow. Dried ground Rhizoctonia-infested millet powder is

- 17 -

sprinkled evenly over the seeds in the furrows. 100 ml of Strain CGA267356 suspensions or sterile H₂O are sprayed over the appropriate treatments using a backpack CO₂ sprayer. Each treatment consist of four replicate plots for a total of 400 seeds per treatment.

Approximately three weeks after planting, the number of healthy standing plants is recorded. For each experiment, the percent control is calculated by comparison to the number of healthy plants in the check (no pathogen added, defined as 100% stand) and the disease check (pathogen added, H₂O treated only, defined as 0% stand). In these experiments, both pre-and post-emergence damping-off occurs.

Table 3

Control of Damping off of Cotton (Field Trials 1 and 2) and green bean (Feild Trial 3) caused by *Rhizoctonia solani*: Synergism between Fludioxonil (chemical seed treatment) and Strain CGA267356 (bacterial drench) under field conditions.

Field Trial 1

% Disease Control

<u>Treatment</u>	<u>Expected Value*</u>	<u>Actual Value</u>
Disease check		0
CGA267356 1x		86.5
CGA267356 1/10x		20.2
Fludioxonil 1/2x		97.7
Fludioxonil 1/3x		77.5
CGA2673561x + Fludioxonil 1/2x	99.7	114**
CGA2673561x + Fludioxonil 1/3x	96.9	102**
CGA2673561/10x + Fludioxonil 1/3x	82	84.4**

Field Trial 2

% Disease Control

<u>Treatment</u>	<u>Expected Value*</u>	<u>Actual Value</u>
Disease check		0
CGA267356 1x		5.3
CGA267356 1/10x		13.3
Fludioxonil 1/2x		71.6
CGA2673561x + Fludioxonil 1/2x	73.1	89**

- 18 -

CGA2673561 1/10x + Fludioxonil 1/2x 75.4 89**

Field Trial 3 % Disease Control

<u>Treatment</u>	<u>Expected Value*</u>	<u>Actual Value</u>
Disease check		0
CGA 267356 1x		9
CGA 267356 1/10x		6
Fludioxonil 1x		58 "
Fludioxonil 1/3x		9
Fludioxonil 1/7x		0
CGA 267356 1x + Fludioxonil 1/3x	17	64**
CGA 267356 1x + Fludioxonil 1/7x	9	55**
CGA 267356 1/10x + Fludioxonil 1/7x	6	24**

Example 3: Synergistic Effects of Combinations of Bacterial Biocontrol Strain CGA 267356 with carboxin.

Soil is prepared by mixing potting soil (Metro-mix 360, WR Grace) with vermiculite and sand. 15 cm x 15 cm pots are filled with the soil mix, and 3/4" deep furrows are made near the perimeter of each pot. Thus, each pot contains a circular 12" furrow.

Stoneville 506 cotton seeds are obtained from the CIBA-GEIGY Delta Research Station. Seeds are treated with carboxin at the rate of 85g/100kg seed.

Strain CGA 267356 is grown overnight in Luria broth at 28°C. Cells are centrifuged, then resuspended in sterile H₂O, and diluted in various proportions to give a range of cfu/ml dilutions. The 1x rate is defined as 20 ml of cell suspension at a concentration of 2×10^9 colony-forming units (cfu)/ml/ft.

Combination of bacterial strains and carboxin are tested as follows. Ten treated or untreated seeds are placed in each furrow. Dried ground Rhizoctonia-infested millet powder is sprinkled evenly over the seeds in the furrows. 20 ml of H₂O or CGA 267356 suspensions are pipetted over the appropriate treatments. Each treatment consists of three replicate pots for a total of 30 seeds per treatment.

- 19 -

The experiments are conducted in a phytotron set to a day/night temperature of 26/21°C. All pots are watered on day 7 and day 13 after planting.

Approximately two weeks (16 days) after planting, all plants are gently removed from the pots and the number of symptom-free plants is recorded. For each experiment, the percent control is calculated by comparison to the healthy check (no pathogen added, defined as 100% symptom-free) and the disease check (pathogen added, H₂O treated only, defined as 0% symptom-free).

In these experiments, both pre-and post-emergence damping-off occurs.

Table 4

Control of Damping off of Cotton caused by Rhizoctonia solani: Synergism between carboxin (chemical seed treatment) and CGA 267356 (drench treatment).

Treatment

	<u>E Value*</u>	<u>Actual Value</u>
Disease check		0
CGA267356 1x		100
CGA267356 1/100x		50
carboxin		72
carboxin + CGA267356 1/100x	86	89**

Example 4: Synergistic Effects of Combinations of Bacterial Biocontrol Strain CGA 267356 with quintozone.

Soil is prepared by mixing potting soil (Metro-mix 360, WR Grace) with vermiculite and sand. 15 cm x 15 cm pots are filled with the soil mix, and 3/4" deep furrows are made near the perimeter of each pot. Thus, each pot contains a circular 12" furrow.

Stoneville 506 cotton seeds are obtained from the CIBA-GEIGY Delta Research Station. Quintozone suspensions are prepared at the following concentrations: 0.002 oz/10 ml (= 1x field use application rate recommended for cotton); 0.001 oz/10 ml (1/2x) and 0.0002 oz/10 ml (1/10x).

- 20 -

Strain CGA267356 is grown overnight in Luria broth at 28°C. Cells are centrifuged, then resuspended in sterile H₂O, and diluted in various proportions to give a range of cfu/ml dilutions. The 1x rate is defined as 20 ml of cell suspension at a concentration of 2×10^9 colony-forming unites (cfu) /ml/ft.

Combination of bacterial strains and quintocene are tested as follows. The untreated seeds are placed in each furrow. Dried ground Rhizoctonia-infested millet powder is sprinkled evenly over the seeds in the furrows. 10 ml of quintocene or H₂O are pipetted over the appropriate treatments, followed by 10 ml of CGA267356 or H₂O. Each treatment consists of three replicated pots for a total of 30 seeds per treatment.

The experiments are conducted in a phytotron set to a day/night temperature of 26/21°C. All pots are watered on day 7 and day 13 after planting.

Approximately two weeks (16 days) after planting, all plants are gently removed from the pots and the number of symptom-free plants is recorded. For each experiment, the percent control is calculated by comparison to the healthy check (no pathogen added, defined as 100% symptom-free) and the disease check (pathogen added, H₂O treated only, defined as 0% symptom-free).

In these experiments, both pre- and post-emergence damping-off occurs.

Table 5

Control of Damping off of Cotton caused by Rhizoctonia solani: Synergism between quintocene (chemical seed treatment) and CGA 267356 (bacterial drench)

<u>Treatment</u>	<u>E Value*</u>	<u>Actual Value</u>
Disease check		0
CGA267356 1x		84
CGA267356 1/100x		23
quintocene 1x		64
quintocene 1/2x		60
quintocene 1/10x		8

- 21 -


quintozene 1x + CGA267356 1/100x	72	86**
quintozene 1/2x + CGA267356 1/100x	69	78**
quintozene 1/10x + CGA 267356 1/100x	29	49**

- 22 -

What is claimed is:

1. A microbicidal composition which comprises, as active ingredients, an effective amount of at least one bacterial biocontrol strain of the genus *Pseudomonas* in combination with at least one synthetic microbicidal compound, together with a suitable carrier material, with the proviso that said synthetic microbicidal compound is not metalaxyl, furalaxyl, oxadixyl, benalaxyl or ofurace.
2. A composition according to claim 1 wherein the weight ratio of *Pseudomonas* strain(I):synthetic microbicide(II) is 100:1 to 1:100, preferably 20:1 to 1:20.
3. The composition of claim 1, wherein said synthetic microbicidal compound is selected from the group consisting of halogenated nitrobenzenes, 3-phenylpyrrole derivatives, carboxin and thiram.
4. The composition of claim 3, wherein the microbicidal biocontrol strain is of the species *Pseudomonas fluorescens*.
5. The composition of claim 4, wherein said bacterial strain is selected from the group consisting of the following strains of *Pseudomonas fluorescens*
 - a) CGA 266446 (ATCC Accession No. 55171);
 - b) CGA 266447 (ATCC Accession No. 55170);
 - c) CGA 267356 (ATCC Accession No. 55169);
 - d) CGA 270293 (ATCC Accession No. 55175);
 - e) CGA 270294 (ATCC Accession No. 55174); and
 - f) CGA 281836 (ATCC Accession No. 55168).
6. The composition of claim 3, wherein the synthetic microbicidal compound is a halogenated nitrobenzene.
7. The composition of claim 6, wherein the halogenated nitrobenzene is pentachloro-nitrobenzene.
8. The composition of claim 4, wherein the 3-phenylpyrrole derivative is of the formula

- 23 -



wherein

R_1 represents hydrogen, halogen, methyl, methoxy, trifluoromethyl, trifluoromethoxy,

R_2 represents halogen, trifluoromethyl or trifluoromethoxy, or

R_1 and R_2 together form a methylenedioxy, an ethylenedioxy or an ethylenoxy bridge, each of them being unsubstituted or substituted by methyl, chlorine or fluorine,

R_3 represents hydrogen or halogen,

X represents cyano, trifluoromethyl or COOCH_3 , and

R represents hydrogen, $\text{C}_1\text{-C}_4$ -acyl, $\text{C}_1\text{-C}_4$ -alkoxycarbonyl, $\text{C}_1\text{-C}_4$ -alkoxy- $\text{C}_1\text{-C}_4$ -acyl, $\text{C}_2\text{-C}_4$ -alkenyloxycarbonyl or carbamoyl.

9. The composition of claim 8, wherein X is cyano.

10. The composition of claim 9, wherein

R_1 represents hydrogen, trifluoromethyl or halogen,

R_2 represents halogen or trifluoromethyl, or

R_1 and R_2 together form a difluoromethylenedioxy bridge,

R_3 represents hydrogen,

X represents cyano and

R represents hydrogen, $\text{C}_1\text{-C}_4$ -acyl, $\text{C}_1\text{-C}_4$ -alkoxycarbonyl, $\text{C}_1\text{-C}_4$ -alkoxy- $\text{C}_1\text{-C}_4$ -acyl, or $\text{C}_2\text{-C}_4$ -alkenyloxycarbonyl.

11. The composition of claim 10, wherein the 3-phenyl-4-cyanopyrrole is 3-(2,3-dichlorophenyl)-4-cyanopyrrole (fenciclonil).

12. The composition of claim 10, wherein the 3-phenyl-4-cyanopyrrole is selected from the group consisting of:

- a) 3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole (=fludioxonil);
- b) 1-acetyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole;
- c) 1-methoxyacetyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole;

- 24 -

- d) 1-methoxycarbonyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole;
- e) 1-allyloxycarbonyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole; and
- f) 1-n-propoxyacetyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole.

13. The composition of claim 10, wherein the 3-phenyl-4-cyanopyrrole is 3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole (=fludioxonil).

14. The composition of claim 3, wherein the synthetic microbicial compound is carboxin and/or thiram.

15. A method of controlling or preventing infestation of cultivated plants by plant pathogens, which method comprises applying to said plants, parts of plants, or to a locus thereof, in either sequence or simultaneously at least one microbicial biocontrol strain of the genus *Pseudomonas* and at least one synthetic microbicial compound.

16. The method of claim 15, wherein said synthetic microbicial compound is selected from the group consisting of halogenated nitrobenzenes, 3-phenylpyrrole derivatives, carboxin and thiram.

17. The method of claim 15, wherein the microbicial biocontrol strain is of the species *Pseudomonas fluorescens*.


18. The method of claim 17, wherein said bacterial strain is selected from the group consisting of the following strains of *Pseudomonas fluorescens*:

- a) CGA 266446 (ATCC Accession No. 55171);
- b) CGA 266447 (ATCC Accession No. 55170);
- c) CGA 267356 (ATCC Accession No. 55169);
- d) CGA 270293 (ATCC Accession No. 55175);
- e) CGA 270294 (ATCC Accession No. 55174); and
- f) CGA 281836 (ATCC Accession No. 55168).

19. The method of claim 16, wherein the halogenated nitrobenzene is pentachloro-nitrobenzene.

20. The method of claim 16, wherein the 3-phenylpyrrole derivative is of the formula

- 25 -



wherein

R_1 represents hydrogen, halogen, methyl, methoxy, trifluoromethyl, trifluoromethoxy,

R_2 represents halogen, trifluoromethyl or trifluoromethoxy, or

R_1 and R_2 together form a methylenedioxy, an ethylenedioxy or an ethylenoxy bridge, each of them being unsubstituted or substituted by methyl, chlorine or fluorine,

R_3 represents hydrogen or halogen,

X represents cyano, trifluoromethyl or COOCH_3 , and

R represents hydrogen, $\text{C}_1\text{-C}_4\text{-acyl}$, $\text{C}_1\text{-C}_4\text{-alkoxycarbonyl}$, $\text{C}_1\text{-C}_4\text{-alkoxy-C}_1\text{-C}_4\text{-acyl}$, $\text{C}_2\text{-C}_4\text{-alkenyloxycarbonyl}$ or carbamoyl.

21. The method of claim 20, wherein X is cyano.

22. The method of claim 21, wherein

R_1 represents hydrogen, trifluoromethyl or halogen,

R_2 represents halogen or trifluoromethyl, or

R_1 and R_2 together form a difluoromethylenedioxy bridge,

R_3 represents hydrogen,

X represents cyano and

R represents hydrogen, $\text{C}_1\text{-C}_4\text{-acyl}$, $\text{C}_1\text{-C}_4\text{-alkoxycarbonyl}$, $\text{C}_1\text{-C}_4\text{-alkoxy-C}_1\text{-C}_4\text{-acyl}$, or $\text{C}_2\text{-C}_4\text{-alkenyloxycarbonyl}$.

23. The method of claim 22, wherein the 3-phenyl-4-cyanopyrrole is 3-(2,3-dichlorophenyl)-4-cyanopyrrole (=fenpiclonil).

24. The method of claim 22, wherein the 3-phenyl-4-cyanopyrrole is selected from the group consisting of:

- a) 3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole (fludioxonil);
- b) 1-acetyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole;
- c) 1-methoxyacetyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole;

- 26 -

- d) 1-methoxycarbonyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole;
- e) 1-allyloxycarbonyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole; and
- f) 1-n-propoxyacetyl-3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole.

25. The method of claim 24, wherein the 3-phenyl-4-cyanopyrrole is 3-(2,2-difluorobenzodioxol-4-yl)-4-cyanopyrrole (=fludioxonil).

26. The method of claim 16, wherein the synthetic microbicidal compound is carboxin and/or thiram.

27. The method of claim 15, wherein the parts of plants are seeds.

28. A dressed seed having reduced susceptibility to phytopathogenic fungi which is dressed according to the method of claim 15.