



US 20210308273A1

(19) **United States**

(12) **Patent Application Publication**
Subramanian et al.

(10) **Pub. No.: US 2021/0308273 A1**

(43) **Pub. Date: Oct. 7, 2021**

(54) **MUSCLE TARGETING COMPLEXES AND USES THEREOF FOR TREATING DYSTROPHINOPATHIES**

(71) Applicant: **Dyne Therapeutics, Inc.**, Waltham, MA (US)

(72) Inventors: **Romesh R. Subramanian**, Framingham, MA (US); **Mohammed T. Qatanani**, Waltham, MA (US); **Timothy Weeden**, Waltham, MA (US)

(73) Assignee: **Dyne Therapeutics, Inc.**, Waltham, MA (US)

(21) Appl. No.: **17/265,024**

(22) PCT Filed: **Aug. 2, 2019**

(86) PCT No.: **PCT/US2019/044949**

§ 371 (c)(1),

(2) Date: **Feb. 1, 2021**

Related U.S. Application Data

(60) Provisional application No. 62/855,766, filed on May 31, 2019, provisional application No. 62/714,031, filed on Aug. 2, 2018.

Publication Classification

(51) **Int. Cl.**

A61K 47/68 (2006.01)

C07K 16/28 (2006.01)

A61P 21/00 (2006.01)

C12N 15/113 (2006.01)

(52) **U.S. Cl.**

CPC *A61K 47/6807* (2017.08); *A61K 47/6849*

(2017.08); *C07K 16/2881* (2013.01); *A61P*

21/00 (2018.01); *C12N 15/1137* (2013.01);

A61K 2039/505 (2013.01); *C07K 2317/33*

(2013.01); *C07K 2317/55* (2013.01); *C07K*

2317/92 (2013.01); *C07K 2317/24* (2013.01);

C07K 2317/77 (2013.01); *C12N 15/113*

(2013.01)

(57)

ABSTRACT

Aspects of the disclosure relate to complexes comprising a muscle-targeting agent covalently linked to a molecular payload. In some embodiments, the muscle-targeting agent specifically binds to an internalizing cell surface receptor on muscle cells. In some embodiments, the molecular payload promotes the expression or activity of a functional dystrophin protein. In some embodiments, the molecular payload is an oligonucleotide, such as an antisense oligonucleotide, e.g., an oligonucleotide that causes exon skipping in a mRNA expressed from a mutant DMD allele.

Specification includes a Sequence Listing.




FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D




FIG. 4E

MUSCLE TARGETING COMPLEXES AND USES THEREOF FOR TREATING DYSTROPHINOPATHIES

RELATED APPLICATIONS

[0001] This application claims the benefit of the filing date of U.S. Provisional Application No. 62/714,031, entitled “MUSCLE TARGETING COMPLEXES AND USES THEREOF FOR TREATING DYSTROPHINOPATHIES”, filed Aug. 2, 2018; and U.S. Provisional Application No. 62/855,766, entitled “MUSCLE TARGETING COMPLEXES AND USES THEREOF FOR TREATING DYSTROPHINOPATHIES”, filed May 31, 2019; the contents of each of which are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

[0002] The present application relates to targeting complexes for delivering molecular payloads (e.g., oligonucleotides) to cells and uses thereof, particularly uses relating to treatment of disease.

REFERENCE TO THE SEQUENCE LISTING

[0003] The present application is being filed along with a Sequence Listing in electronic format. The Sequence Listing is provided as a file entitled D082470008WO00-SEQ.txt created on Jul. 31, 2019 which is 102 kilobytes in size. The information in electronic format of the sequence listing is incorporated herein by reference in its entirety.

BACKGROUND OF INVENTION

[0004] Dystrophinopathies are a group of distinct neuromuscular diseases that result from mutations in dystrophin gene. Dystrophinopathies include Duchenne muscular dystrophy, Becker muscular dystrophy, and X-linked dilated cardiomyopathy. Dystrophin (DMD) is a large gene, containing 79 exons and ~2.6 million total base pairs. Numerous mutations in DMD, including exonic frameshift, deletion, substitution, and duplicative mutations, are able to diminish the expression of functional dystrophin, leading to dystrophinopathies. One agent that targets exon 51 of human DMD, eteplirsen, has been preliminarily approved by the U.S. Food and Drug Administration (FDA) however its efficacy is still being evaluated.

SUMMARY OF INVENTION

[0005] According to some aspects, the disclosure provides complexes that target muscle cells for purposes of delivering molecular payloads to those cells. In some embodiments, complexes provided herein are particularly useful for delivering molecular payloads that increase or restore expression or activity of functional DMD. In some embodiments, complexes comprise oligonucleotide based molecular payloads that promote normal expression of functional DMD through an in-frame exon skipping mechanism or suppression of stop codons. In other embodiments, complexes are configured for delivering a mini-dystrophin gene or synthetic mRNA that increases or restores functional dystrophin activity. Accordingly, in some embodiments, complexes provided herein comprise muscle-targeting agents (e.g., muscle targeting antibodies) that specifically bind to receptors on the surface of muscle cells for purposes of delivering

molecular payloads to the muscle cells. In some embodiments, the complexes are taken up into the cells via a receptor mediated internalization, following which the molecular payload may be released to perform a function inside the cells. For example, complexes engineered to deliver oligonucleotides may release the oligonucleotides such that the oligonucleotides can promote expression of functional DMD (e.g., through an exon skipping mechanism) in the muscle cells. In some embodiments, the oligonucleotides are released by endosomal cleavage of covalent linkers connecting oligonucleotides and muscle-targeting agents of the complexes.

[0006] Some aspects of the disclosure comprise a complex comprising a muscle-targeting agent covalently linked to a molecular payload configured for promoting the expression or activity of a DMD, wherein the muscle-targeting agent specifically binds to an internalizing cell surface receptor on muscle cells.

[0007] In some embodiments, the muscle-targeting agent is a muscle-targeting antibody. In some embodiments, a muscle-targeting antibody is an antibody that specifically binds to an extracellular epitope of a transferrin receptor (e.g., an epitope of the apical domain of the transferrin receptor). A muscle-targeting antibody may specifically binds to an epitope of a sequence in the range of C89 to F760 of SEQ ID NO: 1-3. In some embodiments, the equilibrium dissociation constant (Kd) of binding of a muscle-targeting antibody to a transferrin receptor is in a range from 10^{-11} M to 10^{-6} M.

[0008] In some embodiments, a muscle-targeting antibody of a complex competes for specific binding to an epitope of a transferrin receptor with an antibody listed in Table 1 (e.g., competes for specific binding to an epitope of a transferrin receptor with an Kd of less than or equal to 10^{-6} M, e.g., in a range of 10^{-11} M to 10^{-6} M).

[0009] In some embodiments, a muscle-targeting antibody of a complex does not specifically bind to the transferrin binding site of a transferrin receptor and/or does not inhibit binding of transferrin to a transferrin receptor. In some embodiments, a muscle-targeting antibody of a complex is cross-reactive with extracellular epitopes of two or more of a human, non-human primate and rodent transferrin receptor. In some embodiments, a muscle-targeting antibody of a complex is configured to promote transferrin receptor mediated internalization of the molecular payload into a muscle cell.

[0010] A muscle-targeting antibody (e.g., muscle-targeting antibody is an antibody that specifically binds to an extracellular epitope of a transferrin receptor) is a chimeric antibody, wherein optionally the chimeric antibody is a humanized monoclonal antibody. A muscle-targeting antibody may be in the form of a ScFv, Fab fragment, Fab' fragment, F(ab')₂ fragment, or Fv fragment.

[0011] In some embodiments, a molecular payload of a complex is an oligonucleotide. In some embodiments, an oligonucleotide comprises a sequence listed in Table 2. In some embodiments, an oligonucleotide comprises a sequence as provided in any one of SEQ ID NO: 15-266. In some embodiments, an oligonucleotide comprises a region of complementarity to a mutated DMD allele.

[0012] In some embodiments, an oligonucleotide is configured to suppress a truncating mutation in a DMD allele by mono- or multi-exon skipping. In some embodiments, an oligonucleotide promotes antisense-mediated exon skipping

to produce in-frame dystrophin mRNA. In some embodiments, an oligonucleotide promotes skipping of an exon of DMD in the range of exon 8 to exon 55 (e.g., exon 23 to exon 53). In some embodiments, an oligonucleotide promotes skipping of exon 8, exon 23, exon 44, exon 45, exon 50, exon 51, exon 52, exon 53, and/or exon 55. In some embodiments, an oligonucleotide promotes skipping of multiple exons in the range of exon 44 to exon 53.

[0013] An oligonucleotide of the disclosure may comprise at least one modified internucleotide linkage (e.g., a phosphorothioate linkage). In some embodiments, an oligonucleotide comprises phosphorothioate linkages in the Rp stereochemical conformation and in the Sp stereochemical conformation. In some embodiments, an oligonucleotide comprises phosphorothioate linkages that are all in the Rp stereochemical conformation. In other embodiments, an oligonucleotide comprises phosphorothioate linkages that are all in the Sp stereochemical conformation.

[0014] An oligonucleotide of the disclosure may comprise one or more modified nucleotides (e.g., 2'-modified nucleotides). In some embodiments, a modified nucleotide is a 2'-O-methyl, 2'-fluoro (2'-F), 2'-O-methoxyethyl (2'-MOE), or 2', 4'-bridged nucleotide. In some embodiments, a modified nucleotide is a bridged nucleotide (e.g., selected from: 2',4'-constrained 2'-O-ethyl (cEt) and locked nucleic acid (LNA) nucleotides).

[0015] In some embodiments, an oligonucleotide is a gapmer oligonucleotide that directs RNase H-mediated cleavage of an miRNA that negatively regulates DMD expression in a cell, optionally wherein the miRNA is miR-31. A gapmer oligonucleotide may comprise a central portion of 5 to 15 deoxyribonucleotides flanked by wings of 2 to 8 modified nucleotides (e.g., 2'-modified nucleotides).

[0016] In some embodiments, an oligonucleotide is a mixmer oligonucleotide. In some embodiments, a mixmer oligonucleotide promotes exon skipping. A mixmer oligonucleotide may comprise two or more different 2' modified nucleotides.

[0017] In some embodiments, an oligonucleotide is an RNAi oligonucleotide that promotes RNAi-mediated cleavage of an miRNA that negatively regulates DMD expression in a cell, optionally wherein the miRNA is miR-31. An RNAi oligonucleotide may be a double-stranded oligonucleotide of 19 to 25 nucleotides in length. In some embodiments, an RNAi oligonucleotide comprises at least one 2' modified nucleotide.

[0018] In some embodiments, an oligonucleotide comprises a guide sequence for a genome editing nuclease.

[0019] In some embodiments, an oligonucleotide is a phosphorodiamidite morpholino oligomer (PMO).

[0020] In other embodiments, a molecular payload is a polypeptide. In some embodiments, a molecular payload is a polypeptide that is a functional fragment of dystrophin protein.

[0021] In some embodiments, a muscle-targeting agent is covalently linked to a molecular payload via a cleavable linker (e.g., a protease-sensitive linker, pH-sensitive linker, or glutathione-sensitive linker). A protease-sensitive linker may comprise a sequence cleavable by a lysosomal protease and/or an endosomal protease. In some embodiments, a protease-sensitive linker comprises a valine-citrulline dipeptide sequence. A pH-sensitive linker may be cleaved at a pH in a range of 4 to 6.

[0022] In some embodiments, a muscle-targeting agent is covalently linked to a molecular payload via a non-cleavable linker (e.g., an alkane linker).

[0023] In some embodiments, a muscle-targeting antibody comprises a non-natural amino acid to which an oligonucleotide can be covalently linked. In some embodiments, a muscle-targeting antibody is covalently linked to an oligonucleotide via conjugation to a lysine residue or a cysteine residue of the antibody. In some embodiments, an oligonucleotide is conjugated to a cysteine residue of the antibody via a maleimide-containing linker, optionally wherein the maleimide-containing linker comprises a maleimidocaproyl or maleimidomethyl cyclohexane-1-carboxylate group.

[0024] In some embodiments, a muscle-targeting antibody is a glycosylated antibody that comprises at least one sugar moiety to which an oligonucleotide is covalently linked. In some embodiments, a glycosylated antibody that comprises at least one sugar moiety that is a branched mannose. In some embodiments, a muscle-targeting antibody is a glycosylated antibody that comprises one to four sugar moieties each of which is covalently linked to a separate oligonucleotide. In some embodiments, a muscle-targeting antibody is a fully-glycosylated antibody or a partially-glycosylated antibody. A partially-glycosylated antibody may be produced via chemical or enzymatic means. In some embodiments, a partially-glycosylated antibody is produced in a cell that is deficient for an enzyme in the N- or O-glycosylation pathway.

[0025] Some aspects of the disclosure comprise a method of delivering an molecular payload to a cell expressing transferrin receptor. In some embodiments, methods of delivering an molecular payload to a cell expressing transferrin receptor comprise contacting the cell with a complex comprising a muscle-targeting agent covalently linked to a molecular payload configured for promoting the expression or activity of a DMD.

[0026] Some aspects of the disclosure comprise a method of promoting the expression or activity of a DMD protein in a cell. In some embodiments, methods of promoting the expression or activity of a DMD protein in a cell comprise contacting the cell with a complex comprising a muscle-targeting agent covalently linked to a molecular payload configured for promoting the expression or activity of a DMD in an amount effective for promoting internalization of the molecular payload to the cell. In some embodiments, the cell is in vitro. In some embodiments, the cell is in a subject. In some embodiments, the subject is a human.

[0027] Further aspects of the disclosure comprise a method of treating a subject having a mutated DMD allele that is associated with a dystrophinopathy. In some embodiments, methods of treating a subject having a mutated DMD allele that is associated with a dystrophinopathy comprise administering to the subject an effective amount of a complex comprising a muscle-targeting agent covalently linked to a molecular payload configured for promoting the expression or activity of a DMD.

[0028] Yet further aspects of the disclosure comprise a method of promoting skipping of an exon of a DMD mRNA transcript in a cell. In some embodiments, methods of promoting skipping of an exon of a DMD mRNA transcript comprise administering to the cell an effective amount of the complex comprising a muscle-targeting agent covalently linked to a molecular payload configured for promoting the expression or activity of a DMD. In some embodiments,

methods promote skipping of exon 8, exon 23, exon 44, exon 45, exon 50, exon 51, exon 52, exon 53, and/or exon 55 of the DMD mRNA transcript.

BRIEF DESCRIPTION OF THE DRAWINGS

[0029] FIG. 1 depicts a non-limiting schematic showing the effect of transfecting cells with an siRNA.

[0030] FIG. 2 depicts a non-limiting schematic showing the activity of a muscle targeting complex comprising an siRNA.

[0031] FIGS. 3A-3B depict non-limiting schematics showing the activity of a muscle targeting complex comprising an siRNA in mouse muscle tissues (gastrocnemius and heart) in vivo, relative to control experiments. (N=4 C57BL/6 WT mice)

[0032] FIGS. 4A-4E depict non-limiting schematics showing the tissue selectivity of a muscle targeting complex comprising an siRNA.

DETAILED DESCRIPTION OF INVENTION

[0033] Aspects of the disclosure relate to a recognition that while certain molecular payloads (e.g., oligonucleotides, peptides, small molecules) can have beneficial effects in muscle cells, it has proven challenging to effectively target such cells. As described herein, the present disclosure provides complexes comprising muscle-targeting agents covalently linked to molecular payloads in order to overcome such challenges. In some embodiments, the complexes are particularly useful for delivering molecular payloads that modulate (e.g., promote) the expression or activity of target genes in muscle cells, e.g., in a subject having or suspected of having a rare muscle disease. For example, in some embodiments, complexes are provided for targeting DMD, e.g., a mutated DMD allele. In some embodiments, complexes provided herein may comprise oligonucleotides that promote normal expression and activity of DMD. As another example, complexes may comprise oligonucleotides that induce skipping of exon of DMD mRNA. In some embodiments, synthetic nucleic acid payloads (e.g., DNA or RNA payloads) may be used that express one or more proteins that promote normal expression and activity of DMD.

[0034] In some embodiments, complexes may comprise molecular payloads of synthetic cDNAs and/or synthetic mRNAs, e.g., that express dystrophin or fragments thereof (e.g., a dystrophin mini gene). In some embodiments, complexes may comprise molecular payloads such as guide molecules (e.g., guide RNAs) that are capable of targeting nucleic acid programmable nucleases (e.g., Cas9) to a sequence at or near a disease-associated mutation of DMD, e.g., a mutated DMD exon. In some embodiments, such nucleic programmable nucleases could be used to cleave part or all of a disease-associated mutation of DMD, e.g., a mutated DMD exon, to promote expression of functional DMD. In some embodiments, complexes may comprise molecular payloads that upregulate the expression and/or activity of genes that can replace the function of dystrophin, such as utrophin.

[0035] Further aspects of the disclosure, including a description of defined terms, are provided below.

I. Definitions

[0036] Administering: As used herein, the terms “administering” or “administration” means to provide a complex to

a subject in a manner that is physiologically and/or pharmacologically useful (e.g., to treat a condition in the subject).

[0037] Approximately: As used herein, the term “approximately” or “about,” as applied to one or more values of interest, refers to a value that is similar to a stated reference value. In certain embodiments, the term “approximately” or “about” refers to a range of values that fall within 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[0038] Antibody: As used herein, the term “antibody” refers to a polypeptide that includes at least one immunoglobulin variable domain or at least one antigenic determinant, e.g., paratope that specifically binds to an antigen. In some embodiments, an antibody is a full-length antibody. In some embodiments, an antibody is a chimeric antibody. In some embodiments, an antibody is a humanized antibody. However, in some embodiments, an antibody is a Fab fragment, a F(ab')₂ fragment, a Fv fragment or a scFv fragment. In some embodiments, an antibody is a nanobody derived from a camelid antibody or a nanobody derived from shark antibody. In some embodiments, an antibody is a diabody. In some embodiments, an antibody comprises a framework having a human germline sequence. In another embodiment, an antibody comprises a heavy chain constant domain selected from the group consisting of IgG, IgG1, IgG2, IgG2A, IgG2B, IgG2C, IgG3, IgG4, IgA1, IgA2, IgD, IgM, and IgE constant domains. In some embodiments, an antibody comprises a heavy (H) chain variable region (abbreviated herein as VH), and/or a light (L) chain variable region (abbreviated herein as VL). In some embodiments, an antibody comprises a constant domain, e.g., an Fc region. An immunoglobulin constant domain refers to a heavy or light chain constant domain. Human IgG heavy chain and light chain constant domain amino acid sequences and their functional variations are known. With respect to the heavy chain, in some embodiments, the heavy chain of an antibody described herein can be an alpha (α), delta (Δ), epsilon (ε), gamma (γ) or mu (μ) heavy chain. In some embodiments, the heavy chain of an antibody described herein can comprise a human alpha (α), delta (Δ), epsilon (ε), gamma (γ) or mu (μ) heavy chain. In a particular embodiment, an antibody described herein comprises a human gamma 1 CH1, CH2, and/or CH3 domain. In some embodiments, the amino acid sequence of the VH domain comprises the amino acid sequence of a human gamma (γ) heavy chain constant region, such as any known in the art. Non-limiting examples of human constant region sequences have been described in the art, e.g., see U.S. Pat. No. 5,693,780 and Kabat E A et al., (1991) supra. In some embodiments, the VH domain comprises an amino acid sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or at least 99% identical to any of the variable chain constant regions provided herein. In some embodiments, an antibody is modified, e.g., modified via glycosylation, phosphorylation, sumoylation, and/or methylation. In some embodiments, an antibody is a glycosylated antibody, which is conjugated to one or more sugar or carbohydrate molecules. In some embodiments, the one or more sugar or carbohydrate molecule are conjugated to the antibody via N-glycosylation, O-glycosylation, C-glycosylation, glypiation (GPI anchor attachment), and/or

phosphoglycosylation. In some embodiments, the one or more sugar or carbohydrate molecule are monosaccharides, disaccharides, oligosaccharides, or glycans. In some embodiments, the one or more sugar or carbohydrate molecule is a branched oligosaccharide or a branched glycan. In some embodiments, the one or more sugar or carbohydrate molecule includes a mannose unit, a glucose unit, an N-acetylglucosamine unit, an N-acetylgalactosamine unit, a galactose unit, a fucose unit, or a phospholipid unit. In some embodiments, an antibody is a construct that comprises a polypeptide comprising one or more antigen binding fragments of the disclosure linked to a linker polypeptide or an immunoglobulin constant domain. Linker polypeptides comprise two or more amino acid residues joined by peptide bonds and are used to link one or more antigen binding portions. Examples of linker polypeptides have been reported (see e.g., Holliger, P., et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R. J., et al. (1994) *Structure* 2:1121-1123). Still further, an antibody may be part of a larger immunoadhesion molecule, formed by covalent or noncovalent association of the antibody or antibody portion with one or more other proteins or peptides. Examples of such immunoadhesion molecules include use of the streptavidin core region to make a tetrameric scFv molecule (Kipriyanov, S. M., et al. (1995) *Human Antibodies and Hybridomas* 6:93-101) and use of a cysteine residue, a marker peptide and a C-terminal polyhistidine tag to make bivalent and biotinylated scFv molecules (Kipriyanov, S. M., et al. (1994) *Mol. Immunol.* 31:1047-1058).

[0039] CDR: As used herein, the term “CDR” refers to the complementarity determining region within antibody variable sequences. There are three CDRs in each of the variable regions of the heavy chain and the light chain, which are designated CDR1, CDR2 and CDR3, for each of the variable regions. The term “CDR set” as used herein refers to a group of three CDRs that occur in a single variable region capable of binding the antigen. The exact boundaries of these CDRs have been defined differently according to different systems. The system described by Kabat (Kabat et al., *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, Md. (1987) and (1991)) not only provides an unambiguous residue numbering system applicable to any variable region of an antibody, but also provides precise residue boundaries defining the three CDRs. These CDRs may be referred to as Kabat CDRs. Sub-portions of CDRs may be designated as L1, L2 and L3 or H1, H2 and H3 where the “L” and the “H” designates the light chain and the heavy chains regions, respectively. These regions may be referred to as Chothia CDRs, which have boundaries that overlap with Kabat CDRs. Other boundaries defining CDRs overlapping with the Kabat CDRs have been described by Padlan (FASEB J. 9:133-139 (1995)) and MacCallum (*J Mol Biol* 262(5):732-45 (1996)). Still other CDR boundary definitions may not strictly follow one of the above systems, but will nonetheless overlap with the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. The methods used herein may utilize CDRs defined according to any of these systems, although preferred embodiments use Kabat or Chothia defined CDRs.

[0040] CDR-grafted antibody: The term “CDR-grafted antibody” refers to antibodies which comprise heavy and light chain variable region sequences from one species but

in which the sequences of one or more of the CDR regions of VH and/or VL are replaced with CDR sequences of another species, such as antibodies having murine heavy and light chain variable regions in which one or more of the murine CDRs (e.g., CDR3) has been replaced with human CDR sequences.

[0041] Chimeric antibody: The term “chimeric antibody” refers to antibodies which comprise heavy and light chain variable region sequences from one species and constant region sequences from another species, such as antibodies having murine heavy and light chain variable regions linked to human constant regions.

[0042] Complementary: As used herein, the term “complementary” refers to the capacity for precise pairing between two nucleotides or two sets of nucleotides. In particular, complementary is a term that characterizes an extent of hydrogen bond pairing that brings about binding between two nucleotides or two sets of nucleotides. For example, if a base at one position of an oligonucleotide is capable of hydrogen bonding with a base at the corresponding position of a target nucleic acid (e.g., an mRNA), then the bases are considered to be complementary to each other at that position. Base pairings may include both canonical Watson-Crick base pairing and non-Watson-Crick base pairing (e.g., Wobble base pairing and Hoogsteen base pairing). For example, in some embodiments, for complementary base pairings, adenosine-type bases (A) are complementary to thymidine-type bases (T) or uracil-type bases (U), that cytosine-type bases (C) are complementary to guanosine-type bases (G), and that universal bases such as 3-nitropyrrole or 5-nitroindole can hybridize to and are considered complementary to any A, C, U, or T. Inosine (I) has also been considered in the art to be a universal base and is considered complementary to any A, C, U or T.

[0043] Conservative amino acid substitution: As used herein, a “conservative amino acid substitution” refers to an amino acid substitution that does not alter the relative charge or size characteristics of the protein in which the amino acid substitution is made. Variants can be prepared according to methods for altering polypeptide sequence known to one of ordinary skill in the art such as are found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Fourth Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2012, or *Current Protocols in Molecular Biology*, F. M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. Conservative substitutions of amino acids include substitutions made amongst amino acids within the following groups: (a) M, I, L, V; (b) F, Y, W; (c) K, R, H; (d) A, G; (e) S, T; (f) Q, N; and (g) E, D.

[0044] Covalently linked: As used herein, the term “covalently linked” refers to a characteristic of two or more molecules being linked together via at least one covalent bond. In some embodiments, two molecules can be covalently linked together by a single bond, e.g., a disulfide bond or disulfide bridge, that serves as a linker between the molecules. However, in some embodiments, two or more molecules can be covalently linked together via a molecule that serves as a linker that joins the two or more molecules together through multiple covalent bonds. In some embodiments, a linker may be a cleavable linker. However, in some embodiments, a linker may be a non-cleavable linker.

[0045] Cross-reactive: As used herein and in the context of a targeting agent (e.g., antibody), the term “cross-reactive,”

refers to a property of the agent being capable of specifically binding to more than one antigen of a similar type or class (e.g., antigens of multiple homologs, paralogs, or orthologs) with similar affinity or avidity. For example, in some embodiments, an antibody that is cross-reactive against human and non-human primate antigens of a similar type or class (e.g., a human transferrin receptor and non-human primate transferrin receptor) is capable of binding to the human antigen and non-human primate antigens with a similar affinity or avidity. In some embodiments, an antibody is cross-reactive against a human antigen and a rodent antigen of a similar type or class. In some embodiments, an antibody is cross-reactive against a rodent antigen and a non-human primate antigen of a similar type or class. In some embodiments, an antibody is cross-reactive against a human antigen, a non-human primate antigen, and a rodent antigen of a similar type or class.

[0046] DMD: As used herein, the term “DMD” refers to a gene that encodes dystrophin protein, a key component of the dystrophin-glycoprotein complex, which bridges the inner cytoskeleton and the extracellular matrix in muscle cells, particularly muscle fibers. Deletions, duplications, and point mutations in DMD may cause dystrophinopathies, such as Duchenne muscular dystrophy, Becker muscular dystrophy, or cardiomyopathy. Alternative promoter usage and alternative splicing result in numerous distinct transcript variants and protein isoforms for this gene. In some embodiments, a dystrophin gene may be a human (Gene ID: 1756), non-human primate (e.g., Gene ID: 465559), or rodent gene (e.g., Gene ID: 13405; Gene ID: 24907). In addition, multiple human transcript variants (e.g., as annotated under GenBank RefSeq Accession Numbers: NM_000109.3, NM_004006.2, NM_004009.3, NM_004010.3 and NM_004011.3) have been characterized that encode different protein isoforms.

[0047] DMD allele: As used herein, the term “DMD allele” refers to any one of alternative forms (e.g., wild-type or mutant forms) of a DMD gene. In some embodiments, a DMD allele may encode for dystrophin that retains its normal and typical functions. In some embodiments, a DMD allele may comprise one or more mutations that results in muscular dystrophy. Common mutations that lead to Duchenne muscular dystrophy involve frameshift, deletion, substitution, and duplicative mutations of one or more of 79 exons present in a dystrophin allele, e.g., exon 8, exon 23, exon 41, exon 44, exon 50, exon 51, exon 52, exon 53, or exon 55. Further examples of DMD mutations are disclosed, for example, in Flanigan K M, et al., *Mutational spectrum of DMD mutations in dystrophinopathy patients: application of modern diagnostic techniques to a large cohort*. Hum Mutat. 2009 December; 30 (12):1657-66, the contents of which are incorporated herein by reference in its entirety.

[0048] Dystrophinopathy: As used herein, the term “dystrophinopathy” refers to a muscle disease results from one or more mutated DMD alleles. Dystrophinopathies include a spectrum of conditions (ranging from mild to severe) that includes Duchenne muscular dystrophy, Becker muscular dystrophy, and DMD-associated dilated cardiomyopathy (DCM). In some embodiments, at one end of the spectrum, dystrophinopathy is phenotypically associated with an asymptomatic increase in serum concentration of creatine phosphokinase (CK) and/or muscle cramps with myoglobinuria. In some embodiments, at the other end of the spectrum, dystrophinopathy is phenotypically associated with progres-

sive muscle diseases that are generally classified as Duchenne or Becker muscular dystrophy when skeletal muscle is primarily affected and as DMD-associated dilated cardiomyopathy (DCM) when the heart is primarily affected. Symptoms of Duchenne muscular dystrophy include muscle loss or degeneration, diminished muscle function, pseudo-hypertrophy of the tongue and calf muscles, higher risk of neurological abnormalities, and a shortened lifespan. Duchenne muscular dystrophy is associated with Online Mendelian Inheritance in Man (OMIM) Entry #310200. Becker muscular dystrophy is associated with OMIM Entry #300376. Dilated cardiomyopathy is associated with OMIM Entry X #302045.

[0049] Framework: As used herein, the term “framework” or “framework sequence” refers to the remaining sequences of a variable region minus the CDRs. Because the exact definition of a CDR sequence can be determined by different systems, the meaning of a framework sequence is subject to correspondingly different interpretations. The six CDRs (CDR-L1, CDR-L2, and CDR-L3 of light chain and CDR-H1, CDR-H2, and CDR-H3 of heavy chain) also divide the framework regions on the light chain and the heavy chain into four sub-regions (FR1, FR2, FR3 and FR4) on each chain, in which CDR1 is positioned between FR1 and FR2, CDR2 between FR2 and FR3, and CDR3 between FR3 and FR4. Without specifying the particular sub-regions as FR1, FR2, FR3 or FR4, a framework region, as referred by others, represents the combined FRs within the variable region of a single, naturally occurring immunoglobulin chain. As used herein, a FR represents one of the four sub-regions, and FRs represents two or more of the four sub-regions constituting a framework region. Human heavy chain and light chain acceptor sequences are known in the art. In one embodiment, the acceptor sequences known in the art may be used in the antibodies disclosed herein.

[0050] Human antibody: The term “human antibody”, as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the disclosure may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo), for example in the CDRs and in particular CDR3. However, the term “human antibody”, as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

[0051] Humanized antibody: The term “humanized antibody” refers to antibodies which comprise heavy and light chain variable region sequences from a non-human species (e.g., a mouse) but in which at least a portion of the VH and/or VL sequence has been altered to be more “human-like”, i.e., more similar to human germline variable sequences. One type of humanized antibody is a CDR-grafted antibody, in which human CDR sequences are introduced into non-human VH and VL sequences to replace the corresponding nonhuman CDR sequences. In one embodiment, humanized anti-transferrin receptor antibodies and antigen binding portions are provided. Such antibodies may be generated by obtaining murine anti-transferrin receptor monoclonal antibodies using traditional hybridoma technology followed by humanization using in vitro genetic engi-

neering, such as those disclosed in Kasaian et al PCT publication No. WO 2005/123126 A2.

[0052] Internalizing cell surface receptor: As used herein, the term, “internalizing cell surface receptor” refers to a cell surface receptor that is internalized by cells, e.g., upon external stimulation, e.g., ligand binding to the receptor. In some embodiments, an internalizing cell surface receptor is internalized by endocytosis. In some embodiments, an internalizing cell surface receptor is internalized by clathrin-mediated endocytosis. However, in some embodiments, an internalizing cell surface receptor is internalized by a clathrin-independent pathway, such as, for example, phagocytosis, macropinocytosis, caveolae- and raft-mediated uptake or constitutive clathrin-independent endocytosis. In some embodiments, the internalizing cell surface receptor comprises an intracellular domain, a transmembrane domain, and/or an extracellular domain, which may optionally further comprise a ligand-binding domain. In some embodiments, a cell surface receptor becomes internalized by a cell after ligand binding. In some embodiments, a ligand may be a muscle-targeting agent or a muscle-targeting antibody. In some embodiments, an internalizing cell surface receptor is a transferrin receptor.

[0053] Isolated antibody: An “isolated antibody”, as used herein, is intended to refer to an antibody that is substantially free of other antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds transferrin receptor is substantially free of antibodies that specifically bind antigens other than transferrin receptor). An isolated antibody that specifically binds transferrin receptor complex may, however, have cross-reactivity to other antigens, such as transferrin receptor molecules from other species. Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals.

[0054] Kabat numbering: The terms “Kabat numbering”, “Kabat definitions” and “Kabat labeling” are used interchangeably herein. These terms, which are recognized in the art, refer to a system of numbering amino acid residues which are more variable (i.e. hypervariable) than other amino acid residues in the heavy and light chain variable regions of an antibody, or an antigen binding portion thereof (Kabat et al. (1971) *Ann. NY Acad. Sci.* 190:382-391 and, Kabat, E. A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). For the heavy chain variable region, the hypervariable region ranges from amino acid positions 31 to 35 for CDR1, amino acid positions 50 to 65 for CDR2, and amino acid positions 95 to 102 for CDR3. For the light chain variable region, the hypervariable region ranges from amino acid positions 24 to 34 for CDR1, amino acid positions 50 to 56 for CDR2, and amino acid positions 89 to 97 for CDR3.

[0055] Molecular payload: As used herein, the term “molecular payload” refers to a molecule or species that functions to modulate a biological outcome. In some embodiments, a molecular payload is linked to, or otherwise associated with a muscle-targeting agent. In some embodiments, the molecular payload is a small molecule, a protein, a peptide, a nucleic acid, or an oligonucleotide. In some embodiments, the molecular payload functions to modulate the transcription of a DNA sequence, to modulate the expression of a protein, or to modulate the activity of a protein. In some embodiments, the molecular payload is an

oligonucleotide that comprises a strand having a region of complementarity to a target gene.

[0056] Muscle-targeting agent: As used herein, the term, “muscle-targeting agent,” refers to a molecule that specifically binds to an antigen expressed on muscle cells. The antigen in or on muscle cells may be a membrane protein, for example an integral membrane protein or a peripheral membrane protein. Typically, a muscle-targeting agent specifically binds to an antigen on muscle cells that facilitates internalization of the muscle-targeting agent (and any associated molecular payload) into the muscle cells. In some embodiments, a muscle-targeting agent specifically binds to an internalizing, cell surface receptor on muscles and is capable of being internalized into muscle cells through receptor mediated internalization. In some embodiments, the muscle-targeting agent is a small molecule, a protein, a peptide, a nucleic acid (e.g., an aptamer), or an antibody. In some embodiments, the muscle-targeting agent is linked to a molecular payload.

[0057] Muscle-targeting antibody: As used herein, the term, “muscle-targeting antibody,” refers to a muscle-targeting agent that is an antibody that specifically binds to an antigen found in or on muscle cells. In some embodiments, a muscle-targeting antibody specifically binds to an antigen on muscle cells that facilitates internalization of the muscle-targeting antibody (and any associated molecular payload) into the muscle cells. In some embodiments, the muscle-targeting antibody specifically binds to an internalizing, cell surface receptor present on muscle cells. In some embodiments, the muscle-targeting antibody is an antibody that specifically binds to a transferrin receptor.

[0058] Oligonucleotide: As used herein, the term “oligonucleotide” refers to an oligomeric nucleic acid compound of up to 200 nucleotides in length. Examples of oligonucleotides include, but are not limited to, RNAi oligonucleotides (e.g., siRNAs, shRNAs), microRNAs, gapmers, mixmers, phosphorodiamidite morpholinos, peptide nucleic acids, aptamers, guide nucleic acids (e.g., Cas9 guide RNAs), etc. Oligonucleotides may be single-stranded or double-stranded. In some embodiments, an oligonucleotide may comprise one or more modified nucleotides (e.g. 2'-O-methyl sugar modifications, purine or pyrimidine modifications). In some embodiments, an oligonucleotide may comprise one or more modified internucleotide linkage. In some embodiments, an oligonucleotide may comprise one or more phosphorothioate linkages, which may be in the Rp or Sp stereochemical conformation.

[0059] Recombinant antibody: The term “recombinant human antibody”, as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell (described in more details in this disclosure), antibodies isolated from a recombinant, combinatorial human antibody library (Hoogenboom H. R., (1997) *TIB Tech.* 15:62-70; Azzazy H., and Highsmith W. E., (2002) *Clin. Biochem.* 35:425-445; Gavilondo J. V., and Larrick J. W. (2002) *BioTechniques* 29:128-145; Hoogenboom H., and Chames P. (2000) *Immunology Today* 21:371-378), antibodies isolated from an animal (e.g., a mouse) that is transgenic for human immunoglobulin genes (see e.g., Taylor, L. D., et al. (1992) *Nucl. Acids Res.* 20:6287-6295; Kellermann S-A., and Green L. L. (2002) *Current Opinion in Biotechnology* 13:593-597; Little M. et al (2000) *Immu-*

nology Today 21:364-370) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies are subjected to in vitro mutagenesis (or, when an animal transgenic for human Ig sequences is used, in vivo somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire in vivo. One embodiment of the disclosure provides fully human antibodies capable of binding human transferrin receptor which can be generated using techniques well known in the art, such as, but not limited to, using human Ig phage libraries such as those disclosed in Jermutus et al., PCT publication No. WO 2005/007699 A2.

[0060] Region of complementarity: As used herein, the term “region of complementarity” refers to a nucleotide sequence, e.g., of a oligonucleotide, that is sufficiently complementary to a cognate nucleotide sequence, e.g., of a target nucleic acid, such that the two nucleotide sequences are capable of annealing to one another under physiological conditions (e.g., in a cell). In some embodiments, a region of complementarity is fully complementary to a cognate nucleotide sequence of target nucleic acid. However, in some embodiments, a region of complementarity is partially complementary to a cognate nucleotide sequence of target nucleic acid (e.g., at least 80%, 90%, 95% or 99% complementarity). In some embodiments, a region of complementarity contains 1, 2, 3, or 4 mismatches compared with a cognate nucleotide sequence of a target nucleic acid.

[0061] Specifically binds: As used herein, the term “specifically binds” refers to the ability of a molecule to bind to a binding partner with a degree of affinity or avidity that enables the molecule to be used to distinguish the binding partner from an appropriate control in a binding assay or other binding context. With respect to an antibody, the term, “specifically binds”, refers to the ability of the antibody to bind to a specific antigen with a degree of affinity or avidity, compared with an appropriate reference antigen or antigens, that enables the antibody to be used to distinguish the specific antigen from others, e.g., to an extent that permits preferential targeting to certain cells, e.g., muscle cells, through binding to the antigen, as described herein. In some embodiments, an antibody specifically binds to a target if the antibody has a K_D for binding the target of at least about 10^{-4} M, 10^{-5} M, 10^{-6} M, 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M, 10^{-12} M, 10^{-13} M, or less. In some embodiments, an antibody specifically binds to the transferrin receptor, e.g., an epitope of the apical domain of transferrin receptor.

[0062] Subject: As used herein, the term “subject” refers to a mammal. In some embodiments, a subject is non-human primate, or rodent. In some embodiments, a subject is a human. In some embodiments, a subject is a patient, e.g., a human patient that has or is suspected of having a disease. In some embodiments, the subject is a human patient who has or is suspected of having a disease resulting from a mutated DMD gene sequence, e.g., a mutation in an exon of a DMD gene sequence. In some embodiments, a subject has a dystrophinopathy, e.g., Duchenne muscular dystrophy.

[0063] Transferrin receptor: As used herein, the term, “transferrin receptor” (also known as TFRC, CD71, p90, or TFR1) refers to an internalizing cell surface receptor that binds transferrin to facilitate iron uptake by endocytosis. In some embodiments, a transferrin receptor may be of human (NCBI Gene ID 7037), non-human primate (e.g., NCBI Gene ID 711568 or NCBI Gene ID 102136007), or rodent (e.g., NCBI Gene ID 22042) origin. In addition, multiple human transcript variants have been characterized that encoded different isoforms of the receptor (e.g., as annotated under GenBank RefSeq Accession Numbers: NP_001121620.1, NP_003225.2, NP_001300894.1, and NP_001300895.1).

II. Complexes

[0064] Provided herein are complexes that comprise a targeting agent, e.g. an antibody, covalently linked to a molecular payload. In some embodiments, a complex comprises a muscle-targeting antibody covalently linked to a oligonucleotide. A complex may comprise an antibody that specifically binds a single antigenic site or that binds to at least two antigenic sites that may exist on the same or different antigens.

[0065] A complex may be used to modulate the activity or function of at least one gene, protein, and/or nucleic acid. In some embodiments, the molecular payload present with a complex is responsible for the modulation of a gene, protein, and/or nucleic acids. A molecular payload may be a small molecule, protein, nucleic acid, oligonucleotide, or any molecular entity capable of modulating the activity or function of a gene, protein, and/or nucleic acid in a cell. In some embodiments, a molecular payload is an oligonucleotide that targets a disease-associated repeat in muscle cells.

[0066] In some embodiments, a complex comprises a muscle-targeting agent, e.g. an anti-transferrin receptor antibody, covalently linked to a molecular payload, e.g. a mixmer antisense oligonucleotide that targets a mutated DMD allele to promote exon skipping.

[0067] A. Muscle-Targeting Agents

[0068] Some aspects of the disclosure provide muscle-targeting agents, e.g., for delivering a molecular payload to a muscle cell. In some embodiments, such muscle-targeting agents are capable of binding to a muscle cell, e.g., via specifically binding to an antigen on the muscle cell, and delivering an associated molecular payload to the muscle cell. In some embodiments, the molecular payload is bound (e.g., covalently bound) to the muscle targeting agent and is internalized into the muscle cell upon binding of the muscle targeting agent to an antigen on the muscle cell, e.g., via endocytosis. It should be appreciated that various types of muscle-targeting agents may be used in accordance with the disclosure. For example, the muscle-targeting agent may comprise, or consist of, a nucleic acid (e.g., DNA or RNA), a peptide (e.g., an antibody), a lipid (e.g., a microvesicle), or a sugar moiety (e.g., a polysaccharide). Exemplary muscle-targeting agents are described in further detail herein, however, it should be appreciated that the exemplary muscle-targeting agents provided herein are not meant to be limiting.

[0069] Some aspects of the disclosure provide muscle-targeting agents that specifically bind to an antigen on muscle, such as skeletal muscle, smooth muscle, or cardiac muscle. In some embodiments, any of the muscle-targeting

agents provided herein bind to (e.g., specifically bind to) an antigen on a skeletal muscle cell, a smooth muscle cell, and/or a cardiac muscle cell.

[0070] By interacting with muscle-specific cell surface recognition elements (e.g., cell membrane proteins), both tissue localization and selective uptake into muscle cells can be achieved. In some embodiments, molecules that are substrates for muscle uptake transporters are useful for delivering a molecular payload into muscle tissue. Binding to muscle surface recognition elements followed by endocytosis can allow even large molecules such as antibodies to enter muscle cells. As another example molecular payloads conjugated to transferrin or anti-transferrin receptor antibodies can be taken up by muscle cells via binding to transferrin receptor, which may then be endocytosed, e.g., via clathrin-mediated endocytosis.

[0071] The use of muscle-targeting agents may be useful for concentrating a molecular payload (e.g., oligonucleotide) in muscle while reducing toxicity associated with effects in other tissues. In some embodiments, the muscle-targeting agent concentrates a bound molecular payload in muscle cells as compared to another cell type within a subject. In some embodiments, the muscle-targeting agent concentrates a bound molecular payload in muscle cells (e.g., skeletal, smooth, or cardiac muscle cells) in an amount that is at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, or 100 times greater than an amount in non-muscle cells (e.g., liver, neuronal, blood, or fat cells). In some embodiments, a toxicity of the molecular payload in a subject is reduced by at least 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, or 95% when it is delivered to the subject when bound to the muscle-targeting agent.

[0072] In some embodiments, to achieve muscle selectivity, a muscle recognition element (e.g., a muscle cell antigen) may be required. As one example, a muscle-targeting agent may be a small molecule that is a substrate for a muscle-specific uptake transporter. As another example, a muscle-targeting agent may be an antibody that enters a muscle cell via transporter-mediated endocytosis. As another example, a muscle targeting agent may be a ligand that binds to cell surface receptor on a muscle cell. It should be appreciated that while transporter-based approaches provide a direct path for cellular entry, receptor-based targeting may involve stimulated endocytosis to reach the desired site of action.

[0073] i. Muscle-Targeting Antibodies

[0074] In some embodiments, the muscle-targeting agent is an antibody. Generally, the high specificity of antibodies for their target antigen provides the potential for selectively targeting muscle cells (e.g., skeletal, smooth, and/or cardiac muscle cells). This specificity may also limit off-target toxicity. Examples of antibodies that are capable of targeting a surface antigen of muscle cells have been reported and are within the scope of the disclosure. For example, antibodies that target the surface of muscle cells are described in Arahata K., et al. "Immunostaining of skeletal and cardiac muscle surface membrane with antibody against Duchenne muscular dystrophy peptide" *Nature* 1988; 333: 861-3; Song K. S., et al. "Expression of caveolin-3 in skeletal, cardiac, and smooth muscle cells. Caveolin-3 is a component of the sarcolemma and co-fractionates with dystrophin and dystrophin-associated glycoproteins" *J Biol Chem* 1996; 271: 15160-5; and Weisbart R. H. et al., "Cell type specific

targeted intracellular delivery into muscle of a monoclonal antibody that binds myosin IIb" *Mol Immunol.* 2003 March, 39(13):78309; the entire contents of each of which are incorporated herein by reference.

[0075] a. Anti-Transferrin Receptor Antibodies

[0076] Some aspects of the disclosure are based on the recognition that agents binding to transferrin receptor, e.g., anti-transferrin-receptor antibodies, are capable of targeting muscle cell. Transferrin receptors are internalizing cell surface receptors that transport transferrin across the cellular membrane and participate in the regulation and homeostasis of intracellular iron levels. Some aspects of the disclosure provide transferrin receptor binding proteins, which are capable of binding to transferrin receptor. Accordingly, aspects of the disclosure provide binding proteins (e.g., antibodies) that bind to transferrin receptor. In some embodiments, binding proteins that bind to transferrin receptor are internalized, along with any bound molecular payload, into a muscle cell. As used herein, an antibody that binds to a transferrin receptor may be referred to as an anti-transferrin receptor antibody. Antibodies that bind, e.g. specifically bind, to a transferrin receptor may be internalized into the cell, e.g. through receptor-mediated endocytosis, upon binding to a transferrin receptor.

[0077] It should be appreciated that anti-transferrin receptor antibodies may be produced, synthesized, and/or derivatized using several known methodologies, e.g. library design using phage display. Exemplary methodologies have been characterized in the art and are incorporated by reference (Díez, P. et al. "High-throughput phage-display screening in array format", *Enzyme and microbial technology*, 2015, 79, 34-41.; Christoph M. H. and Stanley, J. R. "Antibody Phage Display: Technique and Applications" *J Invest Dermatol.* 2014, 134:2.; Engleman, Edgar (Ed.) "Human Hybridomas and Monoclonal Antibodies." 1985, Springer.). In other embodiments, an anti-transferrin antibody has been previously characterized or disclosed. Antibodies that specifically bind to transferrin receptor are known in the art (see, e.g. U.S. Pat. No. 4,364,934, filed Dec. 4, 1979, "Monoclonal antibody to a human early thymocyte antigen and methods for preparing same"; U.S. Pat. No. 8,409,573, filed Jun. 14, 2006, "Anti-CD71 monoclonal antibodies and uses thereof for treating malignant tumor cells"; U.S. Pat. No. 9,708,406, filed May 20, 2014, "Anti-transferrin receptor antibodies and methods of use"; U.S. Pat. No. 9,611,323, filed Dec. 19, 2014, "Low affinity blood brain barrier receptor antibodies and uses therefor"; WO 2015/098989, filed Dec. 24, 2014, "Novel anti-Transferrin receptor antibody that passes through blood-brain barrier"; Schneider C. et al. "Structural features of the cell surface receptor for transferrin that is recognized by the monoclonal antibody OKT9." *J Biol Chem.* 1982, 257:14, 8516-8522.; Lee et al. "Targeting Rat Anti-Mouse Transferrin Receptor Monoclonal Antibodies through Blood-Brain Barrier in Mouse" 2000, *J Pharmacol. Exp. Ther.*, 292: 1048-1052.).

[0078] Any appropriate anti-transferrin receptor antibodies may be used in the complexes disclosed herein. Examples of anti-transferrin receptor antibodies, including associated references and binding epitopes are listed in Table 1. In some embodiments, the anti-transferrin receptor antibody comprises the complementarity determining regions (CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2, and CDR-L3) of any of the anti-transferrin receptor antibodies provided herein, e.g., anti-transferrin receptor antibodies listed in Table 1.

TABLE 1

List of anti-transferrin receptor antibody clones, including associated references and binding epitope information.		
Antibody Clone Name	Reference(s)	Epitope/Notes
OKT9	U.S. Pat. No. 4,364,934, filed Dec. 4, 1979, entitled "MONOCLONAL ANTIBODY TO A HUMAN EARLY THYMOCYTE ANTIGEN AND METHODS FOR PREPARING SAME" Schneider C. et al. "Structural features of the cell surface receptor for transferrin that is recognized by the monoclonal antibody OKT9." J Biol Chem. 1982, 257: 14, 8516-8522.	Apical domain of TfR (residues 305-366 of human TfR sequence XM_052730.3, available in GenBank)
(From JCR) Clone M11 Clone M23 Clone M27 Clone B84	WO 2015/098989, filed Dec. 24, 2014, "Novel anti-Transferrin receptor antibody that passes through blood-brain barrier" U.S. Pat. No. 9,994,641, filed Dec. 24, 2014, "Novel anti-Transferrin receptor antibody that passes through blood-brain barrier"	Apical domain (residues 230-244 and 326-347 of TfR) and protease-like domain (residues 461-473)
(From Genentech) 7A4, 8A2, 15D2, 10D11, 7B10, 15G11, 16G5, 13C3, 16G4, 16F6, 7G7, 4C2, 1B12, and 13D4 (From Armagen) 8D3	WO 2016/081643, filed May 26, 2016, entitled "ANTI-TRANSFERRIN RECEPTOR ANTIBODIES AND METHODS OF USE" U.S. Pat. No. 9,708,406, filed May 20, 2014, "Anti-transferrin receptor antibodies and methods of use" Lee et al. "Targeting Rat Anti-Mouse Transferrin Receptor Monoclonal Antibodies through Blood-Brain Barrier in Mouse" 2000, J Pharmacol. Exp. Ther., 292: 1048-1052. US Patent App. 2010/077498, filed Sep. 9, 2008, entitled "COMPOSITIONS AND METHODS FOR BLOOD-BRAIN BARRIER DELIVERY IN THE MOUSE"	Apical domain and non-apical regions
OX26	Haobam, B. et al. 2014. Rab17-mediated recycling endosomes contribute to autophagosome formation in response to Group A Streptococcus invasion. Cellular microbiology. 16: 1806-21.	
DF1513	Ortiz-Zapater E et al. Trafficking of the human transferrin receptor in plant cells: effects of tyrphostin A23 and brefeldin A. Plant J 48: 757-70 (2006).	
1A1B2, 66IG10, MEM-189, JF0956, 29806, 1A1B2, TFRC/1818, 1E6, 66Igl0, TFRC/1059, Q1/71, 23D10, 13E4, TFRC/1149, ER-MP21, YTA74.4, BU54, 2B6, R17 217 (From INSERM) BA120g	Commercially available anti-transferrin receptor antibodies. US Patent App. 2011/0311544A1, filed Jun. 15, 2005, entitled "ANTI-CD71 MONOCLONAL ANTIBODIES AND USES THEREOF FOR TREATING MALIGNANT TUMOR CELLS"	Novus Biologicals 8100 Southpark Way, A-8 Littleton CO 80120 Does not compete with OKT9
LUCA31	U.S. Pat. No. 7,572,895, filed Jun. 7, 2004, entitled "TRANSFERRIN RECEPTOR ANTIBODIES"	"LUCA31 epitope"

TABLE 1-continued

List of anti-transferrin receptor antibody clones, including associated references and binding epitope information.		
Antibody Clone Name	Reference(s)	Epitope/Notes
(Salk Institute) B3/25 T58/30	Trowbridge, I. S. et al. "Anti-transferrin receptor monoclonal antibody and toxin-antibody conjugates affect growth of human tumour cells." <i>Nature</i> , 1981, volume 294, pages 171-173	
R17 217.1.3, 5E9C11, OKT9 (BE0023 clone)	Commercially available anti-transferrin receptor antibodies.	BioXcell 10 Technology Dr., Suite 2B West Lebanon, NH 03784-1671 USA
BK19.9, B3/25, T56/14 and T58/1	Gatter, K. C. et al. "Transferrin receptors in human tissues: their distribution and possible clinical relevance." <i>J Clin Pathol.</i> 1983 May; 36(5): 539-45.	

[0079] In some embodiments, the muscle-targeting agent is an anti-transferrin receptor antibody. In some embodiment, an anti-transferrin receptor antibody specifically binds to a transferrin protein having an amino acid sequence as disclosed herein. In some embodiments, an anti-transferrin receptor antibody may specifically bind to any extracellular epitope of a transferrin receptor or an epitope that becomes exposed to an antibody, including the apical domain, the transferrin binding domain, and the protease-like domain. In some embodiments, an anti-transferrin receptor antibody binds to an amino acid segment of a human or non-human primate transferrin receptor, as provided in SEQ ID Nos. 1-3 in the range of amino acids C89 to F760. In some embodiments, an anti-transferrin receptor antibody specifically binds with binding affinity of at least about 10^{-4} M, 10^{-5} M, 10^{-6} M, 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M, 10^{-12} M, 10^{-13} M, or less. Anti-transferrin receptor antibodies used herein may be capable of competing for binding with other anti-transferrin receptor antibodies, e.g. OKT9, 8D3, that bind to transferrin receptor with 10^{-3} M, 10^{-4} M, 10^{-5} M, 10^{-6} M, 10^{-7} M, or less.

[0080] An example human transferrin receptor amino acid sequence, corresponding to NCBI sequence NP_003225.2 (transferrin receptor protein 1 isoform 1, *Homo sapiens*) is as follows:

(SEQ ID NO: 1)

MMDQARSAFSNLFGGPELSYTRFSLARQVDGDNHSHVEMKLVDEEENADN
 NTKANVTKPKRCSGSI CYGTIAIVIFFLIGFMIGLYGCKGVEPKTECER
 LAGTESPVREEPEGDFPAARRLYWDDLKRKLSSEKLDSTDFGTIKLLNEN
 SYVPREAGSQKDENLALYVENQFREFKLSKVWRDQHFVKIQVKDSAQNSV
 IIVDKNGRLVYLVENPGGYVAYSKAATVTGKLVHANFGTKKDFEDLYTPV
 NGSIVIVRAGKITFAEKVANAESLNAIGVLIYMDQTKFPIVNAELSFPGH
 AHLGTGDPYTPGFPSFNHTQFPSPSSGLPNI PVQTI SRAAAEKLFGNME
 GDCPSDWKTDSTCKMVTSESKNVKLTVSNVLKEIKILNIFGVIKGFVEPD
 HYVVVGAQRDAWGPGAAGKSGVTALLLKLQMFSDMVLKDFGQPSRSIIF

-continued

ASWSAGDFGSGATEWLEGYLSLHLKAPTYYINLDKAVLGTSNPKVVSASP
 LLYTLIEKTMQNVKHPVTGQFLYQDSNWASKVEKLTLDNAAFPPLAYSGI
 PAVSFCPCEDTDYPYLGTTMDTYKELIERIPELNKVARAAAEVAGQFVIK
 LTHDVELNLDYERYNSQLLSFVRDLNQYRADIKEMGLSLQWLYSARGDFF
 RATSRLTTDFGNAEKTRFVVMKLNDRVMRVEYHFLSPYVSPKESPPFRHV
 FWGSGSHTLPALLENLKLKQNGAFNETLFRNQALALATWTIQGAANALS
 GDVWDIDNEF.

[0081] An example non-human primate transferrin receptor amino acid sequence, corresponding to NCBI sequence NP_001244232.1 (transferrin receptor protein 1, *Macaca mulatta*) is as follows:

(SEQ ID NO: 2)

MMDQARSAFSNLFGGPELSYTRFSLARQVDGDNHSHVEMKLVDEEENADN
 NTKPNGTKPKRCGGNICYGTIAVIIFFLIGFMIGLYGCKGVEPKTECER
 LAGTESPAREEPEEDFPAAPRLYWDDLKRKLSSEKLDSTDFSTIKLLNEN
 LYVPREAGSQKDENLALYIENQFREFKLSKVWRDQHFVKIQVKDSAQNSV
 IIVDKNGGLVYLVENPGGYVAYSKAATVTGKLVHANFGTKKDFEDLDSPV
 NGSIVIVRAGKITFAEKVANAESLNAIGVLIYMDQTKFPIVNAELSFPGH
 AHLGTGDPYTPGFPSFNHTQFPSPSSGLPNI PVQTI SRAAAEKLFGNME
 GDCPSDWKTDSTCKMVTSESKNVKLTVSNVLKETKILNIFGVIKGFVEPD
 HYVVVGAQRDAWGPGAAGKSGVTALLLKLQMFSDMVLKDFGQPSRSIIF
 ASWSAGDFGSGATEWLEGYLSLHLKAPTYYINLDKAVLGTSNPKVVSASP
 LLYTLIEKTMQDVKHPVTGRSLYQDSNWASKVEKLTLDNAAFPPLAYSGI
 PAVSFCPCEDTDYPYLGTTMDTYKELVERIPELNKVARAAAEVAGQFVIK
 LTHDTELNLDYERYNSQLLFLRDLNQYRADVKEMGLSLQWLYSARGDFF
 RATSRLTTDFRANAEKRDKFVVMKLNDRVMRVEYHFLSPYVSPKESPPFRHV
 FWGSGSHTLSALLESKLRQNSAFNETLFRNQALALATWTIQGAANALS
 GDVWDIDNEF

[0082] An example non-human primate transferrin receptor amino acid sequence, corresponding to NCBI sequence XP_005545315.1 (transferrin receptor protein 1, *Macaca fascicularis*) is as follows:

(SEQ ID NO: 3)

```
MMDQARSAFSNLFGGEP LSYTRFSLARQVDGDN SHVEMKLG VDEEENTDN
NTKANGTKPKRCGGNICYGTIAV IIFFLIGFMIGYLG YCKGVPEKTECER
LAGTESPAREEPEEDFPAAPRL YWDDLKRKLS EKLDTTDFSTIKLLNEN
LYVPREAGSQKDENLALYIENQ FREFKLSKVWRD QHFVKIQVKDS AQNSV
IIVDKNGGLVYLVENPGGYVAYS KAATVTGKLVHAN FGTKKDFEDLSPV
NGSIVIVRAGKITFAEKVANAES LNAIGVLIYMDQT KFPPIVKADLSFFGH
AHLGTGDPYTPGPPS FNHTQFP PPSQSSGLPNI PVQTI SRAAA EKLFGNME
GDCPSDWKTDSTCKMVTSENK SVKLTVSNVLK ETKILNIFGVIKGFVEPD
HYVVVGAQRDAWGPAAKSSVGT ALLLLKLAQMFS DMVLKDGFPQRS IIF
ASWSAGDFGSVGATEWLEGYL SSSLHLKAF TYINLDKAVLGT SNFKV SASP
LLYTLIEKTMQDVKHVPTGRS LYQDSNWASKVE KLTLDNAAFPPLAYSGI
PAVSFCFCEDTDYPYLGTMTD YKELVERIPEL NKVARAAA EVAGQFVIK
LTHDTELNDLYERYNSQLLFL RDLNQRADVKEM GLSLQWLYSARGDFF
RATSRLTTDFRNAEK RDKFVMMKLNDR VMRVEY YFLSPYVSPKES PFRHV
FWGSGSHTLSALLESLKLR RQNNSAFNETL FRNQALATWTTI QGAANALS
GDVWDIDNEF.
```

[0083] An example mouse transferrin receptor amino acid sequence, corresponding to NCBI sequence NP_001344227.1 (transferrin receptor protein 1, *Mus musculus*) is as follows:

(SEQ ID NO: 4)

```
MMDQARSAFSNLFGGEP LSYTRFSLARQVDGDN SHVEMKLA DEEENADN
NMKASVRKPKRFNGRLCFAAIALV IIFFLIGFMSGY LGYCKRVEQKEECVK
LAETEETDKSETMETEDVPTSS RLYWADLKTLLS EKLNSIEFADTIKQLS
QNTYTPREAGSQKDESLAYYIEN QFHEPKFSKVWR DEHYVKIQVKSSIGQ
NMVTIVQSNGLNDPVESPEGYVAF SKPTEVSGKLV HANFGTKKDFEELSY
SVNGSLVIVRAGEITFAEKVANAQ SFNAIGVLIYMD KNKFPVVEADLALF
GHAHLGTGDPYTPGPPS FNHTQFP PPSQSSGLPNI PVQTI SRAAA EKLPKGK
MEGSCPARWNIDSSCKLELSQN QNVKLI VKNVLKERRI LNIFGVIKGYEE
PDRYVVVGAQRDALGAGVAAKSSV GTGLLLKLAQV FSDMISKDGFPRPSR
IIFASWTAGDFGAVGATEWLEGYL SSSLHLKAF TYINLDKAVLGT SNFKV SASP
ASPLLYTLMGKIMQDVKHVPTGRS LYRDSNWI SKVEKLSFDNAAYPPLAY
SGIPAVSFCFCEDADYPYLGT RLDTYEALTQKVP QLNQMVRTAAEVAGQL
I I K L T H D V E L N D Y E M Y N S K L L S F M K D L N Q F K T D I R D M G L S L Q W L Y S A R G
D Y F R A T S R L T T D F H N A E K T N R F V M R E I N D R I M K V E Y H F L S P Y V S P R E S P F
R H I F W G S G S H T L S A L V E N L K L R Q K N I T A F N E T L F R N Q L A L A T W T T I Q G V A N
A L S G D I W N I D N E F
```

[0084] In some embodiments, an anti-transferrin receptor antibody binds to an amino acid segment of the receptor as follows:

(SEQ ID NO: 5)

```
FVKIQVKDS AQNSV IIVDKNGRLVYLVENPGGYVAYS KAATVTGKLVHAN
FGTKKDFEDLTPVNGSIVIVRAGKITFAEKVANAES LNAIGVLIYMDQT
KFPPIVNAELSPFGHAHLGTGDPYTPGPPS FNHTQFP PPSRSGLPNIPVQT
ISRAAAEKLFGNMEGDCPSDWKTDSTCRMVTS ESKNVKLTVSNVLKE
```

and does not inhibit the binding interactions between transferrin receptors and transferrin and/or human hemochromatosis protein (also known as HFE).

[0085] Appropriate methodologies may be used to obtain and/or produce antibodies, antibody fragments, or antigen-binding agents, e.g., through the use of recombinant DNA protocols. In some embodiments, an antibody may also be produced through the generation of hybridomas (see, e.g., Kohler, G and Milstein, C. “Continuous cultures of fused cells secreting antibody of predefined specificity” *Nature*, 1975, 256: 495-497). The antigen-of-interest may be used as the immunogen in any form or entity, e.g., recombinant or a naturally occurring form or entity. Hybridomas are screened using standard methods, e.g. ELISA screening, to find at least one hybridoma that produces an antibody that targets a particular antigen. Antibodies may also be produced through screening of protein expression libraries that express antibodies, e.g., phage display libraries. Phage display library design may also be used, in some embodiments, (see, e.g. U.S. Pat. No. 5,223,409, filed Mar. 1, 1991, “Directed evolution of novel binding proteins”; WO 1992/18619, filed Apr. 10, 1992, “Heterodimeric receptor libraries using phagemids”; WO 1991/17271, filed May 1, 1991, “Recombinant library screening methods”; WO 1992/20791, filed May 15, 1992, “Methods for producing members of specific binding pairs”; WO 1992/15679, filed Feb. 28, 1992, and “Improved epitope displaying phage”). In some embodiments, an antigen-of-interest may be used to immunize a non-human animal, e.g., a rodent or a goat. In some embodiments, an antibody is then obtained from the non-human animal, and may be optionally modified using a number of methodologies, e.g., using recombinant DNA techniques. Additional examples of antibody production and methodologies are known in the art (see, e.g. Harlow et al. “Antibodies: A Laboratory Manual”, Cold Spring Harbor Laboratory, 1988.).

[0086] In some embodiments, an antibody is modified, e.g., modified via glycosylation, phosphorylation, sumoylation, and/or methylation. In some embodiments, an antibody is a glycosylated antibody, which is conjugated to one or more sugar or carbohydrate molecules. In some embodiments, the one or more sugar or carbohydrate molecule are conjugated to the antibody via N-glycosylation, O-glycosylation, C-glycosylation, glypiation (GPI anchor attachment), and/or phosphoglycosylation. In some embodiments, the one or more sugar or carbohydrate molecules are monosaccharides, disaccharides, oligosaccharides, or glycans. In some embodiments, the one or more sugar or carbohydrate molecule is a branched oligosaccharide or a branched glycan. In some embodiments, the one or more sugar or carbohydrate molecule includes a mannose unit, a glucose unit, an N-acetylglucosamine unit, an N-acetylglucalac-

tosamine unit, a galactose unit, a fucose unit, or a phospholipid unit. In some embodiments, there are about 1-10, about 1-5, about 5-10, about 1-4, about 1-3, or about 2 sugar molecules. In some embodiments, a glycosylated antibody is fully or partially glycosylated. In some embodiments, an antibody is glycosylated by chemical reactions or by enzymatic means. In some embodiments, an antibody is glycosylated in vitro or inside a cell, which may optionally be deficient in an enzyme in the N- or O-glycosylation pathway, e.g. a glycosyltransferase. In some embodiments, an antibody is functionalized with sugar or carbohydrate molecules as described in International Patent Application Publication WO2014065661, published on May 1, 2014, entitled, "*Modified antibody, antibody-conjugate and process for the preparation thereof*".

[0087] Some aspects of the disclosure provide proteins that bind to transferrin receptor (e.g., an extracellular portion of the transferrin receptor). In some embodiments, transferrin receptor antibodies provided herein bind specifically to transferrin receptor (e.g., human transferrin receptor). Transferrin receptors are internalizing cell surface receptors that transport transferrin across the cellular membrane and participate in the regulation and homeostasis of intracellular iron levels. In some embodiments, transferrin receptor antibodies provided herein bind specifically to transferrin receptor from human, non-human primates, mouse, rat, etc. In some embodiments, transferrin receptor antibodies provided herein bind to human transferrin receptor. In some embodiments, transferrin receptor antibodies provided herein specifically bind to human transferrin receptor. In some embodiments, transferrin receptor antibodies provided herein bind to an apical domain of human transferrin receptor. In some embodiments, transferrin receptor antibodies provided herein specifically bind to an apical domain of human transferrin receptor.

[0088] In some embodiments, transferrin receptor antibodies of the present disclosure include one or more of the CDR-H (e.g., CDR-H1, CDR-H2, and CDR-H3) amino acid sequences from any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, transferrin receptor antibodies include the CDR-H1, CDR-H2, and CDR-H3 as provided for any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, anti-transferrin receptor antibodies include the CDR-L1, CDR-L2, and CDR-L3 as provided for any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, anti-transferrin receptor antibodies include the CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2, and CDR-L3 as provided for any one of the anti-transferrin receptor antibodies selected from Table 1. The disclosure also includes any nucleic acid sequence that encodes a molecule comprising a CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2, or CDR-L3 as provided for any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, antibody heavy and light chain CDR3 domains may play a particularly important role in the binding specificity/affinity of an antibody for an antigen. Accordingly, anti-transferrin receptor antibodies of the disclosure may include at least the heavy and/or light chain CDR3s of any one of the anti-transferrin receptor antibodies selected from Table 1.

[0089] In some examples, any of the anti-transferrin receptor antibodies of the disclosure have one or more CDR (e.g., CDR-H or CDR-L) sequences substantially similar to

any of the CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2, and/or CDR-L3 sequences from one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, the position of one or more CDRs along the VH (e.g., CDR-H1, CDR-H2, or CDR-H3) and/or VL (e.g., CDR-L1, CDR-L2, or CDR-L3) region of an antibody described herein can vary by one, two, three, four, five, or six amino acid positions so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% of the binding of the original antibody from which it is derived). For example, in some embodiments, the position defining a CDR of any antibody described herein can vary by shifting the N-terminal and/or C-terminal boundary of the CDR by one, two, three, four, five, or six amino acids, relative to the CDR position of any one of the antibodies described herein, so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% of the binding of the original antibody from which it is derived). In another embodiment, the length of one or more CDRs along the VH (e.g., CDR-H1, CDR-H2, or CDR-H3) and/or VL (e.g., CDR-L1, CDR-L2, or CDR-L3) region of an antibody described herein can vary (e.g., be shorter or longer) by one, two, three, four, five, or more amino acids, so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% of the binding of the original antibody from which it is derived).

[0090] Accordingly, in some embodiments, a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or CDR-H3 described herein may be one, two, three, four, five or more amino acids shorter than one or more of the CDRs described herein (e.g., CDRs from any of the anti-transferrin receptor antibodies selected from Table 1) so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or CDR-H3 described herein may be one, two, three, four, five or more amino acids longer than one or more of the CDRs described herein (e.g., CDRs from any of the anti-transferrin receptor antibodies selected from Table 1) so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, the amino portion of a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or CDR-H3 described herein can be extended by one, two, three, four, five or more amino acids compared to one or more of the CDRs described herein (e.g., CDRs from any of the anti-transferrin receptor antibodies selected from Table 1) so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the

binding of the original antibody from which it is derived). In some embodiments, the carboxy portion of a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or CDR-H3 described herein can be extended by one, two, three, four, five or more amino acids compared to one or more of the CDRs described herein (e.g., CDRS from any of the anti-transferrin receptor antibodies selected from Table 1) so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, the amino portion of a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or CDR-H3 described herein can be shortened by one, two, three, four, five or more amino acids compared to one or more of the CDRs described herein (e.g., CDRS from any of the anti-transferrin receptor antibodies selected from Table 1) so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, the carboxy portion of a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or CDR-H3 described herein can be shortened by one, two, three, four, five or more amino acids compared to one or more of the CDRs described herein (e.g., CDRS from any of the anti-transferrin receptor antibodies selected from Table 1) so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). Any method can be used to ascertain whether immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained, for example, using binding assays and conditions described in the art.

[0091] In some examples, any of the anti-transferrin receptor antibodies of the disclosure have one or more CDR (e.g., CDR-H or CDR-L) sequences substantially similar to any one of the anti-transferrin receptor antibodies selected from Table 1. For example, the antibodies may include one or more CDR sequence(s) from any of the anti-transferrin receptor antibodies selected from Table 1 containing up to 5, 4, 3, 2, or 1 amino acid residue variations as compared to the corresponding CDR region in any one of the CDRs provided herein (e.g., CDRs from any of the anti-transferrin receptor antibodies selected from Table 1) so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, any of the amino acid variations in any of the CDRs provided herein may be conservative variations. Conservative variations can be introduced into the CDRs at positions where the residues are not likely to be involved in interacting with a transferrin receptor protein (e.g., a human transferrin receptor protein), for example, as determined based on a crystal structure. Some aspects of the disclosure provide transferrin receptor antibodies that comprise one or more of the heavy chain variable (VH) and/or light chain variable (VL) domains provided herein. In some embodiments, any

of the VH domains provided herein include one or more of the CDR-H sequences (e.g., CDR-H1, CDR-H2, and CDR-H3) provided herein, for example, any of the CDR-H sequences provided in any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, any of the VL domains provided herein include one or more of the CDR-L sequences (e.g., CDR-L1, CDR-L2, and CDR-L3) provided herein, for example, any of the CDR-L sequences provided in any one of the anti-transferrin receptor antibodies selected from Table 1.

[0092] In some embodiments, anti-transferrin receptor antibodies of the disclosure include any antibody that includes a heavy chain variable domain and/or a light chain variable domain of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, anti-transferrin receptor antibodies of the disclosure include any antibody that includes the heavy chain variable and light chain variable pairs of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1.

[0093] Aspects of the disclosure provide anti-transferrin receptor antibodies having a heavy chain variable (VH) and/or a light chain variable (VL) domain amino acid sequence homologous to any of those described herein. In some embodiments, the anti-transferrin receptor antibody comprises a heavy chain variable sequence or a light chain variable sequence that is at least 75% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to the heavy chain variable sequence and/or any light chain variable sequence of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, the homologous heavy chain variable and/or a light chain variable amino acid sequences do not vary within any of the CDR sequences provided herein. For example, in some embodiments, the degree of sequence variation (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) may occur within a heavy chain variable and/or a light chain variable sequence excluding any of the CDR sequences provided herein. In some embodiments, any of the anti-transferrin receptor antibodies provided herein comprise a heavy chain variable sequence and a light chain variable sequence that comprises a framework sequence that is at least 75%, 80%, 85%, 90%, 95%, 98%, or 99% identical to the framework sequence of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1.

[0094] In some embodiments, an anti-transferrin receptor antibody, which specifically binds to transferrin receptor (e.g., human transferrin receptor), comprises a light chain variable VL domain comprising any of the CDR-L domains (CDR-L1, CDR-L2, and CDR-L3), or CDR-L domain variants provided herein, of any of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, an anti-transferrin receptor antibody, which specifically binds to transferrin receptor (e.g., human transferrin receptor), comprises a light chain variable VL domain comprising the CDR-L1, the CDR-L2, and the CDR-L3 of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, the anti-transferrin receptor antibody comprises a light chain variable (VL) region sequence comprising one, two, three or four of the framework regions of the light chain variable region sequence of any anti-

transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, the anti-transferrin receptor antibody comprises one, two, three or four of the framework regions of a light chain variable region sequence which is at least 75%, 80%, 85%, 90%, 95%, or 100% identical to one, two, three or four of the framework regions of the light chain variable region sequence of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, the light chain variable framework region that is derived from said amino acid sequence consists of said amino acid sequence but for the presence of up to 10 amino acid substitutions, deletions, and/or insertions, preferably up to 10 amino acid substitutions. In some embodiments, the light chain variable framework region that is derived from said amino acid sequence consists of said amino acid sequence with 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 amino acid residues being substituted for an amino acid found in an analogous position in a corresponding non-human, primate, or human light chain variable framework region.

[0095] In some embodiments, an anti-transferrin receptor antibody that specifically binds to transferrin receptor comprises the CDR-L1, the CDR-L2, and the CDR-L3 of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, the antibody further comprises one, two, three or all four VL framework regions derived from the VL of a human or primate antibody. The primate or human light chain framework region of the antibody selected for use with the light chain CDR sequences described herein, can have, for example, at least 70% (e.g., at least 75%, 80%, 85%, 90%, 95%, 98%, or at least 99%) identity with a light chain framework region of a non-human parent antibody. The primate or human antibody selected can have the same or substantially the same number of amino acids in its light chain complementarity determining regions to that of the light chain complementarity determining regions of any of the antibodies provided herein, e.g., any of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, the primate or human light chain framework region amino acid residues are from a natural primate or human antibody light chain framework region having at least 75% identity, at least 80% identity, at least 85% identity, at least 90% identity, at least 95% identity, at least 98% identity, at least 99% (or more) identity with the light chain framework regions of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1. In some embodiments, an anti-transferrin receptor antibody further comprises one, two, three or all four VL framework regions derived from a human light chain variable kappa subfamily. In some embodiments, an anti-transferrin receptor antibody further comprises one, two, three or all four VL framework regions derived from a human light chain variable lambda subfamily.

[0096] In some embodiments, any of the anti-transferrin receptor antibodies provided herein comprise a light chain variable domain that further comprises a light chain constant region. In some embodiments, the light chain constant region is a kappa, or a lambda light chain constant region. In some embodiments, the kappa or lambda light chain constant region is from a mammal, e.g., from a human, monkey, rat, or mouse. In some embodiments, the light chain constant

region is a human kappa light chain constant region. In some embodiments, the light chain constant region is a human lambda light chain constant region. It should be appreciated that any of the light chain constant regions provided herein may be variants of any of the light chain constant regions provided herein. In some embodiments, the light chain constant region comprises an amino acid sequence that is at least 75%, 80%, 85%, 90%, 95%, 98%, or 99% identical to any of the light chain constant regions of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1.

[0097] In some embodiments, the anti-transferrin receptor antibody is any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1.

[0098] In some embodiments, an anti-transferrin receptor antibody comprises a VL domain comprising the amino acid sequence of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 1, and wherein the constant regions comprise the amino acid sequences of the constant regions of an IgG, IgE, IgM, IgD, IgA or IgY immunoglobulin molecule, or a human IgG, IgE, IgM, IgD, IgA or IgY immunoglobulin molecule. In some embodiments, an anti-transferrin receptor antibody comprises any of the VL domains, or VL domain variants, and any of the VH domains, or VH domain variants, wherein the VL and VH domains, or variants thereof, are from the same antibody clone, and wherein the constant regions comprise the amino acid sequences of the constant regions of an IgG, IgE, IgM, IgD, IgA or IgY immunoglobulin molecule, any class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2), or any subclass (e.g., IgG2a and IgG2b) of immunoglobulin molecule. Non-limiting examples of human constant regions are described in the art, e.g., see Kabat E A et al., (1991) supra.

[0099] In some embodiments, an antibody of the disclosure can bind to a target antigen (e.g., transferrin receptor) with relatively high affinity, e.g., with a K_D less than 10^{-6} M, 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M or lower. For example, anti-transferrin receptor antibodies can bind to a transferrin receptor protein (e.g., human transferrin receptor) with an affinity between 5 pM and 500 nM, e.g., between 50 pM and 100 nM, e.g., between 500 pM and 50 nM. The disclosure also includes antibodies that compete with any of the antibodies described herein for binding to a transferrin receptor protein (e.g., human transferrin receptor) and that have an affinity of 50 nM or lower (e.g., 20 nM or lower, 10 nM or lower, 500 pM or lower, 50 pM or lower, or 5 pM or lower). The affinity and binding kinetics of the anti-transferrin receptor antibody can be tested using any suitable method including but not limited to biosensor technology (e.g., OCTET or BIACORE).

[0100] In some embodiments, an antibody of the disclosure can bind to a target antigen (e.g., transferrin receptor) with relatively high affinity, e.g., with a K_D less than 10^{-6} M, 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M or lower. For example, anti-transferrin receptor antibodies can bind to a transferrin receptor protein (e.g., human transferrin receptor) with an affinity between 5 pM and 500 nM, e.g., between 50 pM and 100 nM, e.g., between 500 pM and 50 nM. The disclosure also includes antibodies that compete with any of the antibodies described herein for binding to a transferrin receptor protein (e.g., human transferrin receptor) and that have an affinity of 50 nM or lower (e.g., 20 nM or lower, 10

nM or lower, 500 pM or lower, 50 pM or lower, or 5 pM or lower). The affinity and binding kinetics of the anti-transferrin receptor antibody can be tested using any suitable method including but not limited to biosensor technology (e.g., OCTET or BIACORE).

[0101] In some embodiments, the muscle-targeting agent is a transferrin receptor antibody (e.g., antibody and variants thereof as described in International Application Publication WO 2016/081643, incorporated herein by reference).

[0102] In some embodiments, the heavy chain and light chain CDRs of an example antibody according to different definition systems are provided in Table 1.1. The different definition systems, e.g., the Kabat definition, the Chothia definition, and/or the contact definition have been described. See, e.g., (e.g., Kabat, E. A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242, Chothia et al., (1989) Nature 342:877; Chothia, C. et al. (1987) J. Mol. Biol. 196:901-917, Al-lazikani et al (1997) J. Molec. Biol. 273:927-948; and Almagro, J. Mol. Recognit. 17:132-143 (2004). See also hgmp.mrc.ac.uk and bioinf.org.uk/abs).

TABLE 1.1

Heavy chain and light chain CDRs of the mouse transferrin receptor antibody			
CDRs	Kabat	Chothia	Contact
CDR-H1	SYWMH (SEQ ID NO: 267)	GYTFTSY (SEQ ID NO: 273)	TSYWMH (SEQ ID NO: 275)
CDR-H2	EINPTN- GRTNYIEKFKS (SEQ ID NO: 268)	NPTNGR (SEQ ID NO: 274)	WIGEINPTNGRTN (SEQ ID NO: 276)
CDR-H3	GTRAYHY (SEQ ID NO: 269)	GTRAYHY (SEQ ID NO: 269)	ARGTRA (SEQ ID NO: 277)
CDR-L1	RASDNLYSNLA (SEQ ID NO: 270)	RASDNLYSNLYSNLAWY A (SEQ ID NO: 270)	(SEQ ID NO: 278)
CDR-L2	DATNLAD (SEQ ID NO: 271)	DATNLAD (SEQ ID NO: 271)	LLVYDATNLA (SEQ ID NO: 279)
CDR-L3	QHFAGTPLT (SEQ ID NO: 272)	QHFAGTPLT (SEQ ID NO: 272)	QHFAGTPL (SEQ ID NO: 280)

[0103] Example heavy chain variable domain (VH) and light chain variable domain sequences are also provided:

VH
(SEQ ID NO: 283)
QVQLQQPGAELVKPGASVKLSCKASGYFTSYWMHWVKRPGQGLEWIGE
INPTNGRTNYIEKFKSKATLTVDKSSSTAYMQLSSLTSEDSAVVYCARGT
RAYHYWGQGTSVTVSS

-continued

VL
(SEQ ID NO: 284)
DIQMTQSPASLSVSVGETVTTITCRASDNLYSNLAWYQQKQKSPQLLVYD
ATNLADGVPSRFPSSGSGSTQYSLKINSLQSEDFGTYCYQHPFGTPLTPGA
GTKLELEK

[0104] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-H1, a CDR-H2, and a CDR-H3 that are the same as the CDR-H1, CDR-H2, and CDR-H3 shown in Table 1.1. Alternatively or in addition, the transferrin receptor antibody of the present disclosure comprises a CDR-L1, a CDR-L2, and a CDR-L3 that are the same as the CDR-L1, CDR-L2, and CDR-L3 shown in Table 1.1.

[0105] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-H1, a CDR-H2, and a CDR-H3, which collectively contains no more than 5 amino acid variations (e.g., no more than 5, 4, 3, 2, or 1 amino acid variation) as compared with the CDR-H1, CDR-H2, and CDR-H3 as shown in Table 1.1. “Collectively” means that the total number of amino acid variations in all of the three heavy chain CDRs is within the defined range. Alternatively or in addition, the transferrin receptor antibody of the present disclosure may comprise a CDR-L1, a CDR-L2, and a CDR-L3, which collectively contains no more than 5 amino acid variations (e.g., no more than 5, 4, 3, 2 or 1 amino acid variation) as compared with the CDR-L1, CDR-L2, and CDR-L3 as shown in Table 1.1.

[0106] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-H1, a CDR-H2, and a CDR-H3, at least one of which contains no more than 3 amino acid variations (e.g., no more than 3, 2, or 1 amino acid variation) as compared with the counterpart heavy chain CDR as shown in Table 1.1. Alternatively or in addition, the transferrin receptor antibody of the present disclosure may comprise CDR-L1, a CDR-L2, and a CDR-L3, at least one of which contains no more than 3 amino acid variations (e.g., no more than 3, 2, or 1 amino acid variation) as compared with the counterpart light chain CDR as shown in Table 1.1.

[0107] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-L3, which contains no more than 3 amino acid variations (e.g., no more than 3, 2, or 1 amino acid variation) as compared with the CDR-L3 as shown in Table 1.1. In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-L3 containing one amino acid variation as compared with the CDR-L3 as shown in Table 1.1. In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-L3 of QHFAGTPLT (SEQ ID NO: 281 according to the Kabat and Chothia definition system) or QHFAGTPL (SEQ ID NO: 282 according to the Contact definition system). In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-H1, a CDR-H2, a CDR-H3, a CDR-L1 and a CDR-L2 that are the same as the CDR-H1, CDR-H2, and CDR-H3 shown in Table 1.1, and comprises a CDR-L3 of QHFAGTPLT (SEQ ID NO: 281 according to the Kabat and Chothia definition system) or QHFAGTPL (SEQ ID NO: 282 according to the Contact definition system).

[0108] In some embodiments, the transferrin receptor antibody of the present disclosure comprises heavy chain CDRs

that collectively are at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the heavy chain CDRs as shown in Table 1.1. Alternatively or in addition, the transferrin receptor antibody of the present disclosure comprises light chain CDRs that collectively are at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the light chain CDRs as shown in Table 1.1.

[0109] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH comprising the amino acid sequence of SEQ ID NO: 283. Alternatively or in addition, the transferrin receptor antibody of the present disclosure comprises a VL comprising the amino acid sequence of SEQ ID NO: 284.

[0110] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH containing no more than 20 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the VH as set forth in SEQ ID NO: 283. Alternatively or in addition, the transferrin receptor antibody of the present disclosure comprises a VL containing no more than 15 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the VL as set forth in SEQ ID NO: 284.

[0111] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the VH as set forth in SEQ ID NO: 283. Alternatively or in addition, the transferrin receptor antibody of the present disclosure comprises a VL comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the VL as set forth in SEQ ID NO: 284.

[0112] In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized antibody (e.g., a humanized variant). In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-H1, a CDR-H2, a CDR-H3, a CDR-L1, a CDR-L2, and a CDR-L3 that are the same as the CDR-H1, CDR-H2, and CDR-H3 shown in Table 1.1, and comprises a humanized heavy chain variable region and/or a humanized light chain variable region.

[0113] Humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some embodiments, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, the humanized antibody may comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences, but are included to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region or domain (Fc), typically that of a human immunoglobulin. Antibodies may have Fc regions modified as described in WO 99/58572. Other forms

of humanized antibodies have one or more CDRs (one, two, three, four, five, six) which are altered with respect to the original antibody, which are also termed one or more CDRs derived from one or more CDRs from the original antibody. Humanized antibodies may also involve affinity maturation.

[0114] In some embodiments, humanization is achieved by grafting the CDRs (e.g., as shown in Table 1.1) into the IGKV1-NL1*01 and IGHV1-3*01 human variable domains. In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized variant comprising one or more amino acid substitutions at positions 9, 13, 17, 18, 40, 45, and 70 as compared with the VL as set forth in SEQ ID NO: 284, and/or one or more amino acid substitutions at positions 1, 5, 7, 11, 12, 20, 38, 40, 44, 66, 75, 81, 83, 87, and 108 as compared with the VH as set forth in SEQ ID NO: 283. In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized variant comprising amino acid substitutions at all of positions 9, 13, 17, 18, 40, 45, and 70 as compared with the VL as set forth in SEQ ID NO: 284, and/or amino acid substitutions at all of positions 1, 5, 7, 11, 12, 20, 38, 40, 44, 66, 75, 81, 83, 87, and 108 as compared with the VH as set forth in SEQ ID NO: 283.

[0115] In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized antibody and contains residues at positions 43 and 48 of the VL as set forth in SEQ ID NO: 284. Alternatively or in addition, the transferrin receptor antibody of the present disclosure is a humanized antibody and contains the residues at positions 48, 67, 69, 71, and 73 of the VH as set forth in SEQ ID NO: 283.

[0116] The VH and VL amino acid sequences of an example humanized antibody that may be used in accordance with the present disclosure are provided:

Humanized VH
(SEQ ID NO: 285)
EVQLVQSGAEVKKPGASVKVCKASGYTFTSYWMHWVRQAPGQRLIEWIGE
INPTNGRTNYIEKFKSRATLTVDKASATYMELSLRSEDTAVYYCARGT
RAYHYWGQGTMTVSS

Humanized VL
(SEQ ID NO: 286)
DIQMTQSPSSLSASVGDRTVITCRASDNLYSNLAWYQQKPKGSKPLLVDYD
ATNLADGVPSRFRFSGSGSDYSLKINSLSQSEDFGTYQCQHFVGTPLTFGA
GTKLELK

[0117] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH comprising the amino acid sequence of SEQ ID NO: 285. Alternatively or in addition, the transferrin receptor antibody of the present disclosure comprises a VL comprising the amino acid sequence of SEQ ID NO: 286.

[0118] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH containing no more than 20 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the VH as set forth in SEQ ID NO: 285. Alternatively or in addition, the transferrin receptor antibody of the present disclosure comprises a VL containing no more than 15 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12,

11, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the VL as set forth in SEQ ID NO: 286.

[0119] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the VH as set forth in SEQ ID NO: 285. Alternatively or in addition, the transferrin receptor antibody of the present disclosure comprises a VL comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the VL as set forth in SEQ ID NO: 286.

[0120] In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized variant comprising amino acid substitutions at one or more of positions 43 and 48 as compared with the VL as set forth in SEQ ID NO: 284, and/or amino acid substitutions at one or more of positions 48, 67, 69, 71, and 73 as compared with the VH as set forth in SEQ ID NO: 283. In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized variant comprising a S43A and/or a V48L mutation as compared with the VL as set forth in SEQ ID NO: 284, and/or one or more of A67V, L69I, V71R, and K73T mutations as compared with the VH as set forth in SEQ ID NO: 283.

[0121] In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized variant comprising amino acid substitutions at one or more of positions 9, 13, 17, 18, 40, 43, 48, 45, and 70 as compared with the VL as set forth in SEQ ID NO: 284, and/or amino acid substitutions at one or more of positions 1, 5, 7, 11, 12, 20, 38, 40, 44, 48, 66, 67, 69, 71, 73, 75, 81, 83, 87, and 108 as compared with the VH as set forth in SEQ ID NO: 283.

[0122] In some embodiments, the transferrin receptor antibody of the present disclosure is a chimeric antibody, which can include a heavy constant region and a light constant region from a human antibody. Chimeric antibodies refer to antibodies having a variable region or part of variable region from a first species and a constant region from a second species. Typically, in these chimeric antibodies, the variable region of both light and heavy chains mimics the variable regions of antibodies derived from one species of mammals (e.g., a non-human mammal such as mouse, rabbit, and rat), while the constant portions are homologous to the sequences in antibodies derived from another mammal such as human. In some embodiments, amino acid modifications can be made in the variable region and/or the constant region.

[0123] In some embodiments, the transferrin receptor antibody described herein is a chimeric antibody, which can include a heavy constant region and a light constant region from a human antibody. Chimeric antibodies refer to antibodies having a variable region or part of variable region from a first species and a constant region from a second species. Typically, in these chimeric antibodies, the variable region of both light and heavy chains mimics the variable regions of antibodies derived from one species of mammals (e.g., a non-human mammal such as mouse, rabbit, and rat), while the constant portions are homologous to the sequences in antibodies derived from another mammal such as human. In some embodiments, amino acid modifications can be made in the variable region and/or the constant region.

[0124] In some embodiments, the heavy chain of any of the transferrin receptor antibodies as described herein may comprise a heavy chain constant region (CH) or a portion thereof (e.g., CH1, CH2, CH3, or a combination thereof).

The heavy chain constant region can of any suitable origin, e.g., human, mouse, rat, or rabbit. In one specific example, the heavy chain constant region is from a human IgG (a gamma heavy chain), e.g., IgG1, IgG2, or IgG4. An exemplary human IgG1 constant region is given below:

(SEQ ID NO: 287)

```

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGV
HTFPQAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKEVP
KSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVDS
HEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK
EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCL
LVKGFYPSDIAVEWESNGQPENNYKTPPPVLDSDGSFFLYSKLTVDKSRW
QQGNVFSCSVMHEALHNHYTQKSLSLSPGK
    
```

[0125] In some embodiments, the light chain of any of the transferrin receptor antibodies described herein may further comprise a light chain constant region (CL), which can be any CL known in the art. In some examples, the CL is a kappa light chain. In other examples, the CL is a lambda light chain. In some embodiments, the CL is a kappa light chain, the sequence of which is provided below:

(SEQ ID NO: 288)

```

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGV
HTFPQAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKEVP
KSCDKTHTCP
    
```

[0126] Other antibody heavy and light chain constant regions are well known in the art, e.g., those provided in the IMGT database (www.imgt.org) or at www.vbase2.org/vb-stat.php, both of which are incorporated by reference herein.

[0127] Exemplary heavy chain and light chain amino acid sequences of the transferrin receptor antibodies described are provided below:

Heavy Chain (VH + human IgG1 constant region)

(SEQ ID NO: 289)

```

QVQLQQPQGAELVKPGASVKLSCKASGYTFTSYWMHWVKRPGQGLEWIGE
INPTNGRNTNYIEKFKSKATLTVDKSSSTAYMQLSSLTSEDSAVYYCARGT
RAYHWYWGQGTSTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP
EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICN
VNHKPSNTKVDKKEPKKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTL
MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR
VVSIVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL
PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPPVLDSD
GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
    
```

-continued

Light Chain (VL + kappa light chain)
 (SEQ ID NO: 290)
 QVQLQQPGAELVKPGASVKLSCKASGYTFTSYWMHWKQRPQGLEWIGE
 INPTNGRNTNYIEKFKSKATLTVDKSSSTAYMQLSSLTSEDSAVYYCARGT
 RAYHYWGQGTSTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP
 EPVTVSWNSGALTSKVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYICN
 VNHKPSNTKVDKKVEPKSCDKTHTCP
 Heavy Chain (humanized VH + human IgG1
 constant region)
 (SEQ ID NO: 291)
 EVQLVQSGAEVKKPGASVKVCKASGYTFTSYWMHWVRQAPQRLWEWIGE
 INPTNGRNTNYIEKFKSRATLTVDKASASTAYMELSSLRSEDTAVYYCARGT
 RAYHYWGQGTMTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP
 EPVTVSWNSGALTSKVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYICN
 VNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELGGPSVFLFPPKPKDTL
 MISRTPVEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR
 VVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL
 PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD
 GSFPLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK
 Light Chain (humanized VL + kappa light
 chain)
 (SEQ ID NO: 292)
 DIQMTQSPSSLSASVGDRVTITCRASDNLAYSINLAWYQQKPKGKPKLLVYD
 ATNLADGVPSRFRSGSGSDYSLKINSLSQSEDFGTYICQHFNGTPLTFGA
 GTKLELKASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNS
 GALTSGVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYICNVNHKPSNTK
 VDKKVEPKSCDKTHTCP

[0128] In some embodiments, the transferrin receptor antibody described herein comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to SEQ ID NO: 289. Alternatively or in addition, the transferrin receptor antibody described herein comprises a light chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to SEQ ID NO: 290. In some embodiments, the transferrin receptor antibody described herein comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 289. Alternatively or in addition, the transferrin receptor antibody described herein comprises a light chain comprising the amino acid sequence of SEQ ID NO: 290.

[0129] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a heavy chain containing no more than 20 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the heavy chain as set forth in SEQ ID NO: 289. Alternatively or in addition, the transferrin receptor antibody of the present disclosure comprises a light chain containing no more than 15 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the light chain as set forth in SEQ ID NO: 290.

[0130] In some embodiments, the transferrin receptor antibody described herein comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to SEQ ID NO: 291. Alternatively or in addition, the transferrin receptor antibody described herein comprises a light chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to SEQ ID NO: 292. In some embodiments, the transferrin receptor antibody described herein comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 291. Alternatively or in addition, the transferrin receptor antibody described herein comprises a light chain comprising the amino acid sequence of SEQ ID NO: 292.

[0131] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a heavy chain containing no more than 20 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the heavy chain of a humanized antibody as set forth in SEQ ID NO: 289. Alternatively or in addition, the transferrin receptor antibody of the present disclosure comprises a light chain containing no more than 15 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the light chain of humanized antibody as set forth in SEQ ID NO: 290.

[0132] In some embodiments, the transferrin receptor antibody is an antigen binding fragment (FAB) of an intact antibody (full-length antibody). Antigen binding fragment of an intact antibody (full-length antibody) can be prepared via routine methods. For example, F(ab')₂ fragments can be produced by pepsin digestion of an antibody molecule, and Fab fragments that can be generated by reducing the disulfide bridges of F(ab')₂ fragments. Exemplary FAB amino acid sequences of the transferrin receptor antibodies described herein are provided below:

Heavy Chain FAB (VH + a portion of
 human IgG1 constant region)
 (SEQ ID NO: 293)
 QVQLQQPGAELVKPGASVKLSCKASGYTFTSYWMHWKQRPQGLEWIGE
 INPTNGRNTNYIEKFKSKATLTVDKSSSTAYMQLSSLTSEDSAVYYCARGT
 RAYHYWGQGTSTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP
 EPVTVSWNSGALTSKVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYICN
 VNHKPSNTKVDKKVEPKSCDKTHTCP
 Heavy Chain FAB (humanized VH + a
 portion of human IgG1 constant region)
 (SEQ ID NO: 294)
 EVQLVQSGAEVKKPGASVKVCKASGYTFTSYWMHWVRQAPQRLWEWIGE
 INPTNGRNTNYIEKFKSRATLTVDKASASTAYMELSSLRSEDTAVYYCARGT
 RAYHYWGQGTMTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP
 EPVTVSWNSGALTSKVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYICN
 VNHKPSNTKVDKKVEPKSCDKTHTCP

[0133] The transferrin receptor antibodies described herein can be in any antibody form, including, but not limited to, intact (i.e., full-length) antibodies, antigen-binding fragments thereof (such as Fab, Fab', F(ab')₂, Fv), single chain antibodies, bi-specific antibodies, or nanobodies. In

some embodiments, the transferrin receptor antibody described herein is a scFv. In some embodiments, the transferrin receptor antibody described herein is a scFv-Fab (e.g., scFv fused to a portion of a constant region). In some embodiments, the transferrin receptor antibody described herein is a scFv fused to a constant region (e.g., human IgG1 constant region as set forth in SEQ ID NO: 289).

[0134] b. Other Muscle-Targeting Antibodies

[0135] In some embodiments, the muscle-targeting antibody is an antibody that specifically binds hemojuvelin, caveolin-3, Duchenne muscular dystrophy peptide, myosin lib, or CD63. In some embodiments, the muscle-targeting antibody is an antibody that specifically binds a myogenic precursor protein. Exemplary myogenic precursor proteins include, without limitation, ABCG2, M-Cadherin/Cadherin-15, Caveolin-1, CD34, FoxK1, Integrin alpha 7, Integrin alpha 7 beta 1, MYF-5, MyoD, Myogenin, NCAM-1/CD56, Pax3, Pax7, and Pax9. In some embodiments, the muscle-targeting antibody is an antibody that specifically binds a skeletal muscle protein. Exemplary skeletal muscle proteins include, without limitation, alpha-Sarcoglycan, beta-Sarcoglycan, Calpain Inhibitors, Creatine Kinase MM/CKMM, eIF5A, Enolase 2/Neuron-specific Enolase, epsilon-Sarcoglycan, FABP3/H-FABP, GDF-8/Myostatin, GDF-11/GDF-8, Integrin alpha 7, Integrin alpha 7 beta 1, Integrin beta 1/CD29, MCAM/CD146, MyoD, Myogenin, Myosin Light Chain Kinase Inhibitors, NCAM-1/CD56, and Troponin I. In some embodiments, the muscle-targeting antibody is an antibody that specifically binds a smooth muscle protein. Exemplary smooth muscle proteins include, without limitation, alpha-Smooth Muscle Actin, VE-Cadherin, Caldesmon/CALDI, Calponin 1, Desmin, Histamine H2 R, Motilin R/GPR38, Transgelin/TAGLN, and Vimentin. However, it should be appreciated that antibodies to additional targets are within the scope of this disclosure and the exemplary lists of targets provided herein are not meant to be limiting.

[0136] c. Antibody Features/Alterations

[0137] In some embodiments, conservative mutations can be introduced into antibody sequences (e.g., CDRs or framework sequences) at positions where the residues are not likely to be involved in interacting with a target antigen (e.g., transferrin receptor), for example, as determined based on a crystal structure. In some embodiments, one, two or more mutations (e.g., amino acid substitutions) are introduced into the Fc region of a muscle-targeting antibody described herein (e.g., in a CH2 domain (residues 231-340 of human IgG1) and/or CH3 domain (residues 341-447 of human IgG1) and/or the hinge region, with numbering according to the Kabat numbering system (e.g., the EU index in Kabat)) to alter one or more functional properties of the antibody, such as serum half-life, complement fixation, Fc receptor binding and/or antigen-dependent cellular cytotoxicity.

[0138] In some embodiments, one, two or more mutations (e.g., amino acid substitutions) are introduced into the hinge region of the Fc region (CH1 domain) such that the number of cysteine residues in the hinge region are altered (e.g., increased or decreased) as described in, e.g., U.S. Pat. No. 5,677,425. The number of cysteine residues in the hinge region of the CH1 domain can be altered to, e.g., facilitate assembly of the light and heavy chains, or to alter (e.g., increase or decrease) the stability of the antibody or to facilitate linker conjugation.

[0139] In some embodiments, one, two or more mutations (e.g., amino acid substitutions) are introduced into the Fc

region of a muscle-targeting antibody described herein (e.g., in a CH2 domain (residues 231-340 of human IgG1) and/or CH3 domain (residues 341-447 of human IgG1) and/or the hinge region, with numbering according to the Kabat numbering system (e.g., the EU index in Kabat)) to increase or decrease the affinity of the antibody for an Fc receptor (e.g., an activated Fc receptor) on the surface of an effector cell. Mutations in the Fc region of an antibody that decrease or increase the affinity of an antibody for an Fc receptor and techniques for introducing such mutations into the Fc receptor or fragment thereof are known to one of skill in the art. Examples of mutations in the Fc receptor of an antibody that can be made to alter the affinity of the antibody for an Fc receptor are described in, e.g., Smith P et al., (2012) PNAS 109: 6181-6186, U.S. Pat. No. 6,737,056, and International Publication Nos. WO 02/060919; WO 98/23289; and WO 97/34631, which are incorporated herein by reference.

[0140] In some embodiments, one, two or more amino acid mutations (i.e., substitutions, insertions or deletions) are introduced into an IgG constant domain, or FcRn-binding fragment thereof (preferably an Fc or hinge-Fc domain fragment) to alter (e.g., decrease or increase) half-life of the antibody in vivo. See, e.g., International Publication Nos. WO 02/060919; WO 98/23289; and WO 97/34631; and U.S. Pat. Nos. 5,869,046, 6,121,022, 6,277,375 and 6,165,745 for examples of mutations that will alter (e.g., decrease or increase) the half-life of an antibody in vivo.

[0141] In some embodiments, one, two or more amino acid mutations (i.e., substitutions, insertions or deletions) are introduced into an IgG constant domain, or FcRn-binding fragment thereof (preferably an Fc or hinge-Fc domain fragment) to decrease the half-life of the anti-transferrin receptor antibody in vivo. In some embodiments, one, two or more amino acid mutations (i.e., substitutions, insertions or deletions) are introduced into an IgG constant domain, or FcRn-binding fragment thereof (preferably an Fc or hinge-Fc domain fragment) to increase the half-life of the antibody in vivo. In some embodiments, the antibodies can have one or more amino acid mutations (e.g., substitutions) in the second constant (CH2) domain (residues 231-340 of human IgG1) and/or the third constant (CH3) domain (residues 341-447 of human IgG1), with numbering according to the EU index in Kabat (Kabat E A et al., (1991) supra). In some embodiments, the constant region of the IgG1 of an antibody described herein comprises a methionine (M) to tyrosine (Y) substitution in position 252, a serine (S) to threonine (T) substitution in position 254, and a threonine (T) to glutamic acid (E) substitution in position 256, numbered according to the EU index as in Kabat. See U.S. Pat. No. 7,658,921, which is incorporated herein by reference. This type of mutant IgG, referred to as "YTE mutant" has been shown to display fourfold increased half-life as compared to wild-type versions of the same antibody (see Dall'Acqua W F et al., (2006) J Biol Chem 281: 23514-24). In some embodiments, an antibody comprises an IgG constant domain comprising one, two, three or more amino acid substitutions of amino acid residues at positions 251-257, 285-290, 308-314, 385-389, and 428-436, numbered according to the EU index as in Kabat.

[0142] In some embodiments, one, two or more amino acid substitutions are introduced into an IgG constant domain Fc region to alter the effector function(s) of the anti-transferrin receptor antibody. The effector ligand to which affinity is altered can be, for example, an Fc receptor

or the C1 component of complement. This approach is described in further detail in U.S. Pat. Nos. 5,624,821 and 5,648,260. In some embodiments, the deletion or inactivation (through point mutations or other means) of a constant region domain can reduce Fc receptor binding of the circulating antibody thereby increasing tumor localization. See, e.g., U.S. Pat. Nos. 5,585,097 and 8,591,886 for a description of mutations that delete or inactivate the constant domain and thereby increase tumor localization. In some embodiments, one or more amino acid substitutions may be introduced into the Fc region of an antibody described herein to remove potential glycosylation sites on Fc region, which may reduce Fc receptor binding (see, e.g., Shields R L et al., (2001) *J Biol Chem* 276: 6591-604).

[0143] In some embodiments, one or more amino in the constant region of a muscle-targeting antibody described herein can be replaced with a different amino acid residue such that the antibody has altered C1q binding and/or reduced or abolished complement dependent cytotoxicity (CDC). This approach is described in further detail in U.S. Pat. No. 6,194,551 (Idusogie et al). In some embodiments, one or more amino acid residues in the N-terminal region of the CH2 domain of an antibody described herein are altered to thereby alter the ability of the antibody to fix complement. This approach is described further in International Publication No. WO 94/29351. In some embodiments, the Fc region of an antibody described herein is modified to increase the ability of the antibody to mediate antibody dependent cellular cytotoxicity (ADCC) and/or to increase the affinity of the antibody for an Fcγ receptor. This approach is described further in International Publication No. WO 00/42072.

[0144] In some embodiments, the heavy and/or light chain variable domain(s) sequence(s) of the antibodies provided herein can be used to generate, for example, CDR-grafted, chimeric, humanized, or composite human antibodies or antigen-binding fragments, as described elsewhere herein. As understood by one of ordinary skill in the art, any variant, CDR-grafted, chimeric, humanized, or composite antibodies derived from any of the antibodies provided herein may be useful in the compositions and methods described herein and will maintain the ability to specifically bind transferrin receptor, such that the variant, CDR-grafted, chimeric, humanized, or composite antibody has at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or more binding to transferrin receptor relative to the original antibody from which it is derived.

[0145] In some embodiments, the antibodies provided herein comprise mutations that confer desirable properties to the antibodies. For example, to avoid potential complications due to Fab-arm exchange, which is known to occur with native IgG4 mAbs, the antibodies provided herein may comprise a stabilizing 'Adair' mutation (Angal S., et al., "A single amino acid substitution abolishes the heterogeneity of chimeric mouse/human (IgG4) antibody," *Mol Immunol* 30, 105-108; 1993), where serine 228 (EU numbering; residue 241 Kabat numbering) is converted to proline resulting in an IgG1-like hinge sequence. Accordingly, any of the antibodies may include a stabilizing 'Adair' mutation.

[0146] As provided herein, antibodies of this disclosure may optionally comprise constant regions or parts thereof. For example, a VL domain may be attached at its C-terminal end to a light chain constant domain like Cκ or Cλ. Similarly, a VH domain or portion thereof may be attached to all or part of a heavy chain like IgA, IgD, IgE, IgG, and IgM,

and any isotype subclass. Antibodies may include suitable constant regions (see, for example, Kabat et al., *Sequences of Proteins of Immunological Interest*, No. 91-3242, National Institutes of Health Publications, Bethesda, Md. (1991)). Therefore, antibodies within the scope of this may disclosure include VH and VL domains, or an antigen binding portion thereof, combined with any suitable constant regions.

[0147] ii. Muscle-Targeting Peptides

[0148] Some aspects of the disclosure provide muscle-targeting peptides as muscle-targeting agents. Short peptide sequences (e.g., peptide sequences of 5-20 amino acids in length) that bind to specific cell types have been described. For example, cell-targeting peptides have been described in Vines et al., A. "Cell-penetrating and cell-targeting peptides in drug delivery" *Biochim Biophys Acta* 2008, 1786: 126-38; Jarver P., et al., "In vivo biodistribution and efficacy of peptide mediated delivery" *Trends Pharmacol Sci* 2010; 31: 528-35; Samoylova T. I., et al., "Elucidation of muscle-binding peptides by phage display screening" *Muscle Nerve* 1999; 22: 460-6; U.S. Pat. No. 6,329,501, issued on Dec. 11, 2001, entitled "METHODS AND COMPOSITIONS FOR TARGETING COMPOUNDS TO MUSCLE"; and Samoylov A. M., et al., "Recognition of cell-specific binding of phage display derived peptides using an acoustic wave sensor." *Biomol Eng* 2002; 18: 269-72; the entire contents of each of which are incorporated herein by reference. By designing peptides to interact with specific cell surface antigens (e.g., receptors), selectivity for a desired tissue, e.g., muscle, can be achieved. Skeletal muscle-targeting has been investigated and a range of molecular payloads are able to be delivered. These approaches may have high selectivity for muscle tissue without many of the practical disadvantages of a large antibody or viral particle. Accordingly, in some embodiments, the muscle-targeting agent is a muscle-targeting peptide that is from 4 to 50 amino acids in length. In some embodiments, the muscle-targeting peptide is 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 amino acids in length. Muscle-targeting peptides can be generated using any of several methods, such as phage display.

[0149] In some embodiments, a muscle-targeting peptide may bind to an internalizing cell surface receptor that is overexpressed or relatively highly expressed in muscle cells, e.g. a transferrin receptor, compared with certain other cells. In some embodiments, a muscle-targeting peptide may target, e.g., bind to, a transferrin receptor. In some embodiments, a peptide that targets a transferrin receptor may comprise a segment of a naturally occurring ligand, e.g., transferrin. In some embodiments, a peptide that targets a transferrin receptor is as described in U.S. Pat. No. 6,743, 893, filed Nov. 30, 2000, "RECEPTOR-MEDIATED UPTAKE OF PEPTIDES THAT BIND THE HUMAN TRANSFERRIN RECEPTOR". In some embodiments, a peptide that targets a transferrin receptor is as described in Kawamoto, M. et al, "A novel transferrin receptor-targeted hybrid peptide disintegrates cancer cell membrane to induce rapid killing of cancer cells." *BMC Cancer*. 2011 Aug. 18; 11:359. In some embodiments, a peptide that targets a transferrin receptor is as described in U.S. Pat. No. 8,399, 653, filed May 20, 2011, "TRANSFERRIN/TRANSFERIN RECEPTOR-MEDIATED SIRNA DELIVERY".

[0150] As discussed above, examples of muscle targeting peptides have been reported. For example, muscle-specific peptides were identified using phage display library presenting surface heptapeptides. As one example a peptide having the amino acid sequence ASSLNIA (SEQ ID NO: 6) bound to C2C12 murine myotubes in vitro, and bound to mouse muscle tissue in vivo. Accordingly, in some embodiments, the muscle-targeting agent comprises the amino acid sequence ASSLNIA (SEQ ID NO: 6). This peptide displayed improved specificity for binding to heart and skeletal muscle tissue after intravenous injection in mice with reduced binding to liver, kidney, and brain. Additional muscle-specific peptides have been identified using phage display. For example, a 12 amino acid peptide was identified by phage display library for muscle targeting in the context of treatment for DMD. See, Yoshida D., et al., "Targeting of salicylate to skin and muscle following topical injections in rats." *Int J Pharm* 2002; 231: 177-84; the entire contents of which are hereby incorporated by reference. Here, a 12 amino acid peptide having the sequence SKTFNTHPQSTP (SEQ ID NO: 7) was identified and this muscle-targeting peptide showed improved binding to C2C12 cells relative to the ASSLNIA (SEQ ID NO: 6) peptide.

[0151] An additional method for identifying peptides selective for muscle (e.g., skeletal muscle) over other cell types includes in vitro selection, which has been described in Ghosh D., et al., "Selection of muscle-binding peptides from context-specific peptide-presenting phage libraries for adenoviral vector targeting" *J Virol* 2005; 79: 13667-72; the entire contents of which are incorporated herein by reference. By pre-incubating a random 12-mer peptide phage display library with a mixture of non-muscle cell types, non-specific cell binders were selected out. Following rounds of selection the 12 amino acid peptide TARGEHKEEELI (SEQ ID NO: 8) appeared most frequently. Accordingly, in some embodiments, the muscle-targeting agent comprises the amino acid sequence TARGEHKEEELI (SEQ ID NO: 8).

[0152] A muscle-targeting agent may an amino acid-containing molecule or peptide. A muscle-targeting peptide may correspond to a sequence of a protein that preferentially binds to a protein receptor found in muscle cells. In some embodiments, a muscle-targeting peptide contains a high propensity of hydrophobic amino acids, e.g. valine, such that the peptide preferentially targets muscle cells. In some embodiments, a muscle-targeting peptide has not been previously characterized or disclosed. These peptides may be conceived of, produced, synthesized, and/or derivatized using any of several methodologies, e.g. phage displayed peptide libraries, one-bead one-compound peptide libraries, or positional scanning synthetic peptide combinatorial libraries. Exemplary methodologies have been characterized in the art and are incorporated by reference (Gray, B. P. and Brown, K. C. "Combinatorial Peptide Libraries: Mining for Cell-Binding Peptides" *Chem Rev.* 2014, 114:2, 1020-1081.; Samoylova, T. I. and Smith, B. F. "Elucidation of muscle-binding peptides by phage display screening." *Muscle Nerve*, 1999, 22:4. 460-6.). In some embodiments, a muscle-targeting peptide has been previously disclosed (see, e.g. Writer M. J. et al. "Targeted gene delivery to human airway epithelial cells with synthetic vectors incorporating novel targeting peptides selected by phage display." *J. Drug Targeting*, 2004; 12:185; Cal, D. "BDNF-mediated enhancement of inflammation and injury in the aging heart." *Physiol*

Genomics, 2006, 24:3, 191-7.; Zhang, L. "Molecular profiling of heart endothelial cells." *Circulation*, 2005, 112:11, 1601-11.; McGuire, M. J. et al. "In vitro selection of a peptide with high selectivity for cardiomyocytes in vivo." *J Mol Biol.* 2004, 342:1, 171-82.). Exemplary muscle-targeting peptides comprise an amino acid sequence of the following group: CQAQGQLVC (SEQ ID NO: 9), CSERSMNFC (SEQ ID NO: 10), CPKTRRVPC (SEQ ID NO: 11), WLSEAGPVVTVRALRGTGSW (SEQ ID NO: 12), ASSLNIA (SEQ ID NO: 6), CMQHSMRVC (SEQ ID NO: 13), and DDTRHWG (SEQ ID NO: 14). In some embodiments, a muscle-targeting peptide may comprise about 2-25 amino acids, about 2-20 amino acids, about 2-15 amino acids, about 2-10 amino acids, or about 2-5 amino acids. Muscle-targeting peptides may comprise naturally-occurring amino acids, e.g. cysteine, alanine, or non-naturally-occurring or modified amino acids. Non-naturally occurring amino acids include β -amino acids, homo-amino acids, proline derivatives, 3-substituted alanine derivatives, linear core amino acids, N-methyl amino acids, and others known in the art. In some embodiments, a muscle-targeting peptide may be linear; in other embodiments, a muscle-targeting peptide may be cyclic, e.g. bicyclic (see, e.g. Silvana, M. G. et al. *Mol. Therapy*, 2018, 26:1, 132-147.).

[0153] iii. Muscle-Targeting Receptor Ligands

[0154] A muscle-targeting agent may be a ligand, e.g. a ligand that binds to a receptor protein. A muscle-targeting ligand may be a protein, e.g. transferrin, which binds to an internalizing cell surface receptor expressed by a muscle cell. Accordingly, in some embodiments, the muscle-targeting agent is transferrin, or a derivative thereof that binds to a transferrin receptor. A muscle-targeting ligand may alternatively be a small molecule, e.g. a lipophilic small molecule that preferentially targets muscle cells relative to other cell types. Exemplary lipophilic small molecules that may target muscle cells include compounds comprising cholesterol, cholesteryl, stearic acid, palmitic acid, oleic acid, oleyl, linolen, linoleic acid, myristic acid, sterols, dihydrotestosterone, testosterone derivatives, glycerine, alkyl chains, trityl groups, and alkoxy acids.

[0155] iv. Muscle-Targeting Aptamers

[0156] A muscle-targeting agent may be an aptamer, e.g. an RNA aptamer, which preferentially targets muscle cells relative to other cell types. In some embodiments, a muscle-targeting aptamer has not been previously characterized or disclosed. These aptamers may be conceived of, produced, synthesized, and/or derivatized using any of several methodologies, e.g. Systematic Evolution of Ligands by Exponential Enrichment. Exemplary methodologies have been characterized in the art and are incorporated by reference (Yan, A. C. and Levy, M. "Aptamers and aptamer targeted delivery" *RNA biology*, 2009, 6:3, 316-20.; Germer, K. et al. "RNA aptamers and their therapeutic and diagnostic applications." *Int. J. Biochem. Mol. Biol.* 2013; 4: 27-40.). In some embodiments, a muscle-targeting aptamer has been previously disclosed (see, e.g. Phillippou, S. et al. "Selection and Identification of Skeletal-Muscle-Targeted RNA Aptamers." *Mol Ther Nucleic Acids*. 2018, 10:199-214.; Thiel, W. H. et al. "Smooth Muscle Cell-targeted RNA Aptamer Inhibits Neointimal Formation." *Mol Ther.* 2016, 24:4, 779-87.). Exemplary muscle-targeting aptamers include the A01B RNA aptamer and RNA Apt 14. In some embodiments, an aptamer is a nucleic acid-based aptamer, an oligonucleotide aptamer or a peptide aptamer. In some

embodiments, an aptamer may be about 5-15 kDa, about 5-10 kDa, about 10-15 kDa, about 1-5 Da, about 1-3 kDa, or smaller.

[0157] v. Other Muscle-Targeting Agents

[0158] One strategy for targeting a muscle cell (e.g., a skeletal muscle cell) is to use a substrate of a muscle transporter protein, such as a transporter protein expressed on the sarcolemma. In some embodiments, the muscle-targeting agent is a substrate of an influx transporter that is specific to muscle tissue. In some embodiments, the influx transporter is specific to skeletal muscle tissue. Two main classes of transporters are expressed on the skeletal muscle sarcolemma, (1) the adenosine triphosphate (ATP) binding cassette (ABC) superfamily, which facilitate efflux from skeletal muscle tissue and (2) the solute carrier (SLC) superfamily, which can facilitate the influx of substrates into skeletal muscle. In some embodiments, the muscle-targeting agent is a substrate that binds to an ABC superfamily or an SLC superfamily of transporters. In some embodiments, the substrate that binds to the ABC or SLC superfamily of transporters is a naturally-occurring substrate. In some embodiments, the substrate that binds to the ABC or SLC superfamily of transporters is a non-naturally occurring substrate, for example, a synthetic derivative thereof that binds to the ABC or SLC superfamily of transporters.

[0159] In some embodiments, the muscle-targeting agent is a substrate of an SLC superfamily of transporters. SLC transporters are either equilibrative or use proton or sodium ion gradients created across the membrane to drive transport of substrates. Exemplary SLC transporters that have high skeletal muscle expression include, without limitation, the SATT transporter (ASCT1; SLC1A4), GLUT4 transporter (SLC2A4), GLUT7 transporter (GLUT7; SLC2A7), ATRC2 transporter (CAT-2; SLC7A2), LAT3 transporter (KIAA0245; SLC7A6), PHT1 transporter (PTR4; SLC15A4), OATP-J transporter (OATP5A1; SLC21A15), OCT3 transporter (EMT; SLC22A3), OCTN2 transporter (FLJ46769; SLC22A5), ENT transporters (ENT1; SLC29A1 and ENT2; SLC29A2), PAT2 transporter (SLC36A2), and SAT2 transporter (KIAA1382; SLC38A2). These transporters can facilitate the influx of substrates into skeletal muscle, providing opportunities for muscle targeting.

[0160] In some embodiments, the muscle-targeting agent is a substrate of an equilibrative nucleoside transporter 2 (ENT2) transporter. Relative to other transporters, ENT2 has one of the highest mRNA expressions in skeletal muscle. While human ENT2 (hENT2) is expressed in most body organs such as brain, heart, placenta, thymus, pancreas, prostate, and kidney, it is especially abundant in skeletal muscle. Human ENT2 facilitates the uptake of its substrates depending on their concentration gradient. ENT2 plays a role in maintaining nucleoside homeostasis by transporting a wide range of purine and pyrimidine nucleobases. The hENT2 transporter has a low affinity for all nucleosides (adenosine, guanosine, uridine, thymidine, and cytidine) except for inosine. Accordingly, in some embodiments, the muscle-targeting agent is an ENT2 substrate. Exemplary ENT2 substrates include, without limitation, inosine, 2',3'-dideoxyinosine, and calofarabine. In some embodiments, any of the muscle-targeting agents provided herein are associated with a molecular payload (e.g., oligonucleotide payload). In some embodiments, the muscle-targeting agent is covalently linked to the molecular payload. In some

embodiments, the muscle-targeting agent is non-covalently linked to the molecular payload.

[0161] In some embodiments, the muscle-targeting agent is a substrate of an organic cation/carnitine transporter (OCTN2), which is a sodium ion-dependent, high affinity carnitine transporter. In some embodiments, the muscle-targeting agent is carnitine, mildronate, acetylcarnitine, or any derivative thereof that binds to OCTN2. In some embodiments, the carnitine, mildronate, acetylcarnitine, or derivative thereof is covalently linked to the molecular payload (e.g., oligonucleotide payload).

[0162] A muscle-targeting agent may be a protein that is protein that exists in at least one soluble form that targets muscle cells. In some embodiments, a muscle-targeting protein may be hemojuvelin (also known as repulsive guidance molecule C or hemochromatosis type 2 protein), a protein involved in iron overload and homeostasis. In some embodiments, hemojuvelin may be full length or a fragment, or a mutant with at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% sequence identity to a functional hemojuvelin protein. In some embodiments, a hemojuvelin mutant may be a soluble fragment, may lack a N-terminal signaling, and/or lack a C-terminal anchoring domain. In some embodiments, hemojuvelin may be annotated under GenBank RefSeq Accession Numbers NM_001316767.1, NM_145277.4, NM_202004.3, NM_213652.3, or NM_213653.3. It should be appreciated that a hemojuvelin may be of human, non-human primate, or rodent origin.

[0163] B. Molecular Payloads

[0164] Some aspects of the disclosure provide molecular payloads, e.g., for modulating a biological outcome, e.g., the transcription of a DNA sequence, the splicing and processing of a RNA sequence, the expression of a protein, or the activity of a protein. In some embodiments, a molecular payload is linked to, or otherwise associated with a muscle-targeting agent. In some embodiments, such molecular payloads are capable of targeting to a muscle cell, e.g., via specifically binding to a nucleic acid or protein in the muscle cell following delivery to the muscle cell by an associated muscle-targeting agent. It should be appreciated that various types of muscle-targeting agents may be used in accordance with the disclosure. For example, the molecular payload may comprise, or consist of, an oligonucleotide (e.g., anti-sense oligonucleotide), a peptide (e.g., a peptide that binds a nucleic acid or protein associated with disease in a muscle cell), a protein (e.g., a protein that binds a nucleic acid or protein associated with disease in a muscle cell), or a small molecule (e.g., a small molecule that modulates the function of a nucleic acid or protein associated with disease in a muscle cell). In some embodiments, the molecular payload is an oligonucleotide that comprises a strand having a region of complementarity to a mutated DMD allele. Exemplary molecular payloads are described in further detail herein, however, it should be appreciated that the exemplary molecular payloads provided herein are not meant to be limiting.

[0165] i. Oligonucleotides

[0166] Any suitable oligonucleotide may be used as a molecular payload, as described herein. In some embodiments, the oligonucleotide may be designed to induce exon skipping, e.g., EXONDYS 51 oligonucleotide (Sarepta Therapeutics, Inc.), which comprises SEQ ID NO: 195 (CUCCAACAUCAAGGAAGAUGGCAUUUCUAG);

WVE-210201 (Wave Life Sciences), which comprises SEQ ID NO: 186 (UCAAGGAAGAUGGCAUUUCU); Casimersen (Sarepta Therapeutics, Inc.), which comprises SEQ ID NO: 159 (CAAUGCCAUGGAGUUCUG); or Golodirsen (Sarepta Therapeutics, Inc.), which comprises SEQ ID NO: 231 (GUUGCCUCCGGUUCUGAAGGU-GUUC).

[0167] In some embodiments, the oligonucleotide may be designed to cause degradation of an mRNA (e.g., the oligonucleotide may be a gapmer, an siRNA, a ribozyme or an aptamer that causes degradation). In some embodiments, the oligonucleotide may be designed to block translation of an mRNA (e.g., the oligonucleotide may be a mixmer, an siRNA or an aptamer that blocks translation). In some embodiments, an oligonucleotide may be designed to cause degradation and block translation of an mRNA. In some embodiments, the oligonucleotide may be designed to promote stability of an mRNA. In some embodiments, the oligonucleotide may be designed to promote translation of an mRNA. In some embodiments, an oligonucleotide may be designed to promote stability and promote translation of an mRNA. In some embodiments, an oligonucleotide may be a guide nucleic acid (e.g., guide RNA) for directing activity of an enzyme (e.g., a gene editing enzyme). In some embodiments, a guide nucleic acid may direct an enzyme to delete the entirety or a part of a mutated DMD allele (e.g., to facilitate in-frame exon skipping). In some embodiments, the oligonucleotide may be designed to target repressive regulators of DMD expression, e.g., miR-31. Other examples of oligonucleotides are provided herein. It should be appreciated that, in some embodiments, oligonucleotides in one format (e.g., antisense oligonucleotides) may be suitably adapted to another format (e.g., siRNA oligonucleotides) by incorporating functional sequences (e.g., antisense strand sequences) from one format to the other format.

[0168] Examples of oligonucleotides useful for targeting DMD are provided in U.S. Patent Application Publication US20100130591A1, published on May 27, 2010, entitled “MULTIPLE EXON SKIPPING COMPOSITIONS FOR DMD”; U.S. Pat. No. 8,361,979, issued Jan. 29, 2013, entitled “MEANS AND METHOD FOR INDUCING EXON-SKIPPING”; U.S. Patent Application Publication 20120059042, published Mar. 8, 2012, entitled “METHOD FOR EFFICIENT EXON (44) SKIPPING IN DUCHENNE MUSCULAR DYSTROPHY AND ASSOCIATED MEANS”; U.S. Patent Application Publication 20140329881, published Nov. 6, 2014, entitled “EXON SKIPPING COMPOSITIONS FOR TREATING MUSCULAR DYSTROPHY”; U.S. Pat. No. 8,232,384, issued Jul. 31, 2012, entitled “ANTISENSE OLIGONUCLEOTIDES FOR INDUCING EXON SKIPPING AND METHODS OF USE THEREOF”; U.S. Patent Application Publication 20120022134A1, published Jan. 26, 2012, entitled “METHODS AND MEANS FOR EFFICIENT SKIPPING OF EXON 45 IN DUCHENNE MUSCULAR DYSTROPHY PRE-MRNA”; U.S. Patent Application Publication 20120077860, published Mar. 29, 2012, entitled “ADENO-ASSOCIATED VIRAL VECTOR FOR EXON SKIPPING IN A GENE ENCODING A DISPENSABLE DOMAIN PROTEIN”; U.S. Pat. No. 8,324,371, issued Dec. 4, 2012, entitled “OLIGOMERS”; U.S. Pat. No. 9,078,911, issued Jul. 14, 2015, entitled “ANTISENSE OLIGONUCLEOTIDES”; U.S. Pat. No. 9,079,934, issued Jul. 14, 2015, entitled “ANTISENSE NUCLEIC ACIDS”; U.S. Pat. No.

9,034,838, issued May 19, 2015, entitled “MIR-31 IN DUCHENNE MUSCULAR DYSTROPHY THERAPY”; and International Patent Publication WO2017062862A3, published Apr. 13, 2017, entitled “OLIGONUCLEOTIDE COMPOSITIONS AND METHODS THEREOF”; the contents of each of which are incorporated herein in their entireties.

[0169] Table 2 provides non-limiting examples of sequences of oligonucleotide that are useful for targeting DMD, e.g., for exon skipping. In some embodiments, an oligonucleotide may comprise any sequence provided in Table 2.

TABLE 2

Oligonucleotide sequences for targeting DMD.		
EXON	SEQ ID NO:	SEQUENCE
8	15	CUUCCUGGAUGGCUUCAAU
8	16	GUACAUUAGAUGGACUUC
8	17	UAUCUGGAUAGGUGGUAUCAAGAUUCUGUAA
8	18	AUGUAAACUGAAAUGUUCUUCUUUA
8	19	UGGAUAGGUGGUAUCAACAUCUGUAAAGCAC
8	20	GAUAGGUGGUAUCAACAUCUGU
8	21	UAUCUGGAUAGGUGGUAUCAACAUCUGUAA
8	22	AAACUUGGAAGAGUGAUGUGAUGUA
8	23	GCUCACUUGUUGAGGCAAAACUUGGAA
8	24	GCCUUGGCAACAUUUCCACUUCUG
8	25	UACACACUUUACCUUGUUGAGAAUAG
8	26	GAUAGGUGGUAUCAACAUCUGUAA
8	27	GAUAGGUGGUAUCAACAUCUG
8	28	GAUAGGUGGUAUCAACAUCUGUAA
8	29	GGUGGUAUCAACAUCUGUAA
8	30	GUAUCAACAUCUGUAAAGCAC
23	31	CGGCUAAUUUCAGAGGGCGCUUUCUUNGAC
23	32	ACAGUGGUGCUGAGAUAGUAUAGGCC
23	33	UAGGCCACUUUUGUUCUUCUUC
23	34	UUCAGAGGGCGCUUUCUUC
23	253	GGCCAAACCUCGGCUUACCGUAAAU
23	254	GGCCAAACCUCGGCUUACCU
35	35	UCUUCAGGUGCACCUUCUGUUUCUCAAUCU
35	36	UCUGUGAUACUCUUCAGGUGCACCUUCUGU
35	37	UCUUCUGCUCGGGAGGUGACA
35	38	CCAGUUACUAUUCAGAAGAC
35	39	UCUUCAGGUGCACCUUCUGU
43	40	UGCUGCUGUCUUCUUGCU

TABLE 2-continued

Oligonucleotide sequences for targeting DMD.		
EXON	SEQ ID NO:	SEQUENCE
43	41	UUGUUAACUUUUUCCCAUU
43	42	UGUUAACUUUUUCCCAUUGG
43	43	CAUUUUGUUAAACUUUUUCC
43	44	CUGUAGCUUCACCCUUUC
43	45	GAGAGCUUCCUGUAGCUUCACCCUUU
43	46	UCCUGUAGCUUCACCCUUUCACAGGCG
43	47	UGUGUUACCUACCCUUGUCG
43	48	UAGACUAUCUUUAUAUUCUGUAAUAU
43	49	GAGAGCUUCCUGUAGCUUCACCCUUUCCA
43	50	UUCUGUAGCUUCACCCUUUCACAGGCGUU
43	51	AGCUUCCUGUAGCUUCACCCUUU
43	52	GGAGAGAGCUUCCUGUAGCUUCACCCUUU
43	53	GAGAGCUUCCUGUAGCUUCACCC
43	54	UAUGUGUUACCUACCCUUGUCGGUC
43	55	GGAGAGAGCUUCCUGUAGCU
43	56	UCACCCUUUCCACAGGCGUUGCA
43	57	GCUGGGAGAGAGCUUCCUGUAGCUUCAC
43	58	UGUUACCUACCCUUGUCGGUCUUGUAC
43	59	CUGCUGUCUUUUGCUAUGAAUAUUGUC
43	60	GGCGUUGCACUUUGCAAUGCUGCUGUCU
43	61	UUGGAAAUCAAGCUGGGAGAGAGCUUCC
43	62	CUACCCUUGUCGGUCCUUGUACAUUUUG
43	63	GUCAAUCCGACCUGAGCUUUGUUGUAGA
43	64	CUUGCUAUGAAUAUUGUCAAUCCGACC
43	65	UAUAUGUGUUACCUACCCUUGUCGGUCC
43	255	AAUCAGCUGGGAGAGAGCUUCCUGUAGCU
43	256	UCGUUCUUCUGUCGUCGUAACGUUUC
44	66	UUUGUGUCUUUCUGAGAAAC
44	67	AAAGACUUACCUUAAGAUAAC
44	68	AUCUGUCAAAUCGCCUGCAG
44	69	CGCCGCCAUUUCUCAACAG
44	70	UUUGUAUUUAGCAUGUUCCC
44	71	CCGCCAUUUCUCAACAG
44	72	UUCUCAGGAAUUUGUGUCUUU
44	73	GACAACUCUUU
44	74	UCAGCUUCUGUUAGCCACUG

TABLE 2-continued

Oligonucleotide sequences for targeting DMD.		
EXON	SEQ ID NO:	SEQUENCE
44	75	UGUUCAGCUUCUGUUAGCCACUGA
44	76	CUGUUCAGCUUCUGUUAGCCACUGAUU
44	77	UUCUCAACAGAUUCUGUCAAAUCGCCUGCAG
44	78	GCCACUGAUUAAAUAUCUUUAUAUC
44	79	UCUGUUAGCCACUGAUUAAAUAUCUUUAUA
44	80	GAGAAACUGUUCAGCUUCUGUUAGCCACUGA
44	81	UCUUUCUGAGAAACUGUUCAGCUUCUGUUAG
44	82	CAGAUCUGUCAAAUCGCCUGCAGGUA
44	83	CAACAGAUUCUGUCAAAUCGCCUGCAG
44	84	AAACUGUUCAGCUUCUGUUAGCCACUGAUUAAA
44	85	GAAACUGUUCAGCUUCUGUUAGCCACUGAUU
44	86	AAACUGUUCAGCUUCUGUUAGCCACUGA
44	87	UGAGAAACUGUUCAGCUUCUGUUAGCCA
44	88	UUCUGAGAAACUGUUCAGCUUCUGUUAGCCAC
44	89	UUCUGAGAAACUGUUCAGCUUCUGUU
44	90	GAUCUGUCAAAUCGCCUGCAGGUA
44	91	AUAAUGAAAACGCCCAUUUCUCA
44	92	AAACUGUUCAGCUUCUGUUAGCCAC
44	93	UUGUGUCUUUCUGAGAAACUGUUCA
44	94	CCAAUUCUCAGGAAUUUGUGUCUUU
44	95	AUCGCCUGCAGGUAAGCAUAUUGG
44	96	UGAAAACGCCGCCAUUUCUCAACAGAUUCUG
44	97	CAUAAUGAAAACGCCCAUUUCUCAACAG
44	98	UGUUCAGCUUCUGUUAGCCACUGAUUAAA
44	99	CAGAUCUGUCAAAUCGCCUGCAGG
44	100	CAACAGAUUCUGUCAAAUCGCCUGCAGG
44	101	CUCAACAGAUUCUGUCAAAUCGCCUGCAGG
44	102	GAUCUGUCAAAUCGCCUGCAGGU
44	103	GAUCUGUCAAAUCGCCUGCAGG
44	104	GAUCUGUCAAAUCGCCUGCAG
44	105	CAGAUCUGUCAAAUCGCCUGCAGGU
44	106	CAGAUCUGUCAAAUCGCCUGCAG
44	107	GUGUCUUUCUGAGAAACUGUUCAGC
44	108	GAGAAACUGUUCAGCUUCUGUUAGCCAC
44	109	GAAACUGUUCAGCUUCUGUUAGCCACUG
44	110	CUGUUCAGCUUCUGUUAGCCACUG

TABLE 2-continued

Oligonucleotide sequences for targeting DMD.		
EXON	SEQ ID NO:	SEQUENCE
44	111	AUCUGUCAAAUCCGCGCAGGUA AAAAG
44	112	GAUCUGUCAAAUCCGCGCAGGUA AAAAGC
44	257	CACCGAUUGUCUUCGA
44	258	CCCUUGUACGAUUUAUG
44	259	UCUGUGUUUAAGGACUCU
45	113	GCUGAAUUUUUCUCCCC
45	114	UUUUUCUGUCUGACAGCUG
45	115	UCUGUUUUUGAGGAUUGC
45	116	CCACCGCAGAUUCAGGC
45	117	GCCCAAUGCCAUC CUGG
45	118	UUUGCAGACCUC CUGCC
45	119	CAGUUUGCCG CUGCCCA
45	120	GUUGCAUCAAUGUUCUGAC
45	121	AUUUUUCUGUAGAAUACUGG
45	122	GCUGCCCAAUGCGAUCCUGGAGUUC CUGUAAGAU
45	123	GCUGCCCAAUGCCAUC CUGGAGUUC CUG
45	124	GCUGCCCAAUGCCAUC CUGGAGUUC CUGUAA
45	125	CAAUGCCAUC CUGGAGUUC CUGUAAGAUACC
45	126	GCUGCCCAAUGCCAUC CUGGAGUUC CUGUAAG
45	127	CCAAUGCCAUC CUGGAGUUC CUGUAAGAU
45	128	UUGCCG CUGCCCAAUGCCAUC CUGGAGUUC CUGU AAGAU
45	129	GCUGCCCAAUGCCAUC CUGGAGUUC CUGUAAGAU
45	130	CAAUGCCAUC CUGGAGUUC CUGUAAGA
45	131	CAGUUUGCCG CUGCCCAAUGCCAUC C
45	132	CUUCCCCAGUUGCAUUC A AUGUUC
45	133	CUGGCAUCUGUUUUUGAGGAUUG
45	134	UUAGAUCUGUCGCCCCUACCU
45	135	GCUGCCCAAUGCCAUC CUGGAGUUC CUGUAAGAU ACCAA
45	136	GCCCAAUGCCAUC CUGGAGUUC CUGUAAGAUACC
45	137	CAUC CUGGAGUUC CUGUAAGAUACC
45	138	UGCAUC CUGGAGUUC CUGUAAGAUACC
45	139	UGCAUC CUGGAGUUC CUGUAAGAU
45	140	CAAUGCCAUC CUGGAGUUC CUGUAAGAU
45	141	GCCCAAUGCCAUC CUGGAGUUC CUGUAAGAU
45	142	GCCCAAUGCCAUC CUGGAGUUC CUGUAA

TABLE 2-continued

Oligonucleotide sequences for targeting DMD.		
EXON	SEQ ID NO:	SEQUENCE
45	143	GCCG CUGCCCAAUGGACAUC CUGGAGUUC CUGUAA
45	144	GCCAUC CUGGAGUUC CUGUAAGAU
45	145	CCAAUGCCAUC CUGGAGUUC CUGUA
45	146	CUGACAACAGUUUGCCG CUGCCCAA
45	147	UUUGAGGAUUG CUGAAUUUUUCU
45	148	CAGUUUGCCG CUGCCCAAUGCCAUC CUGGA
45	149	UUGCCG CUGCCCAAUGCCAUC CUGGAGUUC
45	150	UUUGCCG CUGCCCAAUGCCAUC CUG
45	151	CCAAUGCCAUC CUGGAGUUC CUG
45	152	CCCAAUGCCAUC CUGGAGUUC CUGUAAGA
45	153	CCG CUGCCCAAUGCCAUC CUGGAGUUC C
45	154	CCCAAUGCCAUC CUGGAGUUC CUGUAAGAU
45	155	CCG CUGCCCAAUGCCAUC CUGGAGUUC CUG
45	156	UGCCCAAUGCCAUC CUGGAGUUC CUGUAAG
45	157	CCCAAUGCCAUC CUGGAGUUC CUGUAAG
45	158	UGCCCAAUGCCAUC CUGGAGUUC CUGUA
45	159	CAAUGCCAUC CUGGAGUUC CUG
45	260	GCCG CUGCCCAAUGCCAUC CUGGAGUUC CUG
45	261	AUUAGAUCUGUCGCCCCUACCUUUUUUC
45	262	UGUCGCCCCUACCUUUUUUCUGUCUG
45	263	GCCCAAUGCCAUC CUGGAGUUC CUG
55	160	AGCCUCUGCUCACUCACC CUGCAAAGGA
50	161	CCACUCAGAGCUCAGAUUCUUAACUUC C
50	162	CUUCCACUCAGAGCUCAGAUUCUUAAC
50	163	GGGAUCCAGUAUAUCUACAGGCUC C
50	164	CUCAGAGCUCAGAUUCU
50	165	GGCUGCUUUGCCUC
50	166	CUCAGAUUCUUAACUUC CUCUUUAAC
50	167	CUCAGAGCUCAGAUUCUUAACUUC CUCU
50	168	CGCCUUCCACUCAGAGCUCAGAUUC C
50	169	UCAGCUCUUGAAGUAAACGGUUUACCG
50	170	UUUGCCUCAGCUCUUGAAGUAAACGG
50	171	GGCUGCUUUGCCUCAGCUCUUGAAGU
50	172	CAGGAGCUAGGUCAGGCUCUUUGCC
50	173	UCCAAUAGUGGUCAGUCCAGGAGCU
50	174	AAAGAGAAUGGGAUCCAGUAUACUUAAC

TABLE 2-continued

Oligonucleotide sequences for targeting DMD.		
EXON	SEQ ID NO:	SEQUENCE
50	175	AAAUAGCUAGAGCCAAAGAGAAUGGGA
50	176	GGCUGCUUUGCCCUCAGCUCUUGAAGUAAACGG
50	177	AGGCUGCUUUGCCCUCAGCUCUUGAAGUAA
50	178	GUCAGGCUGCUUUGCCCUCAGCUCUUGAAG
50	179	AGGUCAGGCUGCUUUGCCCUCAGCUCUUGA
50	180	CAGAGCUCAGAUUCUUAACUUCU
50	181	CUUACAGGCUCCAUAUAGUGGUCAGU
50	182	AUGGGAUCCAGUAUACUUAACAGGCU
50	183	AGAGAAUGGGAUCCAGUAUACUUAAC
50	184	AACUUCUCUUUAAACAGAAAAGCAUAC
50	264	GAGCCUCUCGCUCACUCACCCUGCAAAGGA
51	185	CUCAUACCUUCUGCUUGAUGAUC
51	186	UCAAGGAAGAUGGCAUUUCU
51	187	GAAAGCCAGUCGGUAAGUUC
51	188	CACCCACCAUCACCC
51	189	CCUCUGAUUUUAUAACUUGAU
51	190	UGAUUCCUCAAGGUCACCC
51	191	GGUACCUCCAACAUCAAGGAAGAUGGCAUU
51	192	AUUUCUAGUUUGGAGAUGGCAGUUUC
51	193	CAUCAAGGAAGAUGGCAUUUCUAGUU
51	194	GAGCAGGUACCUCCAACAUCAAGGAA
51	195	CUCCAACAUCAAGGAAGAUGGCAUUUCUAG
51	196	ACCAGAGUAAACAGUCUGAGUAGGAG
51	197	CACCAGAGUAAACAGUCUGAGUAGGA
51	198	UCACCAGAGUAAACAGUCUGAGUAGG
51	199	GUCACCAGAGUAAACAGUCUGAGUAG
51	200	ACCAGAGUAAACAGUCUGAGUAGGAGC
51	201	UUCUGUCCAAGCCCGGUUGAAAUC
51	202	ACAUCAAGGAAGAUGGCAUUUCUAGUUUGG
51	203	ACAUCAAGGAAGAUGGCAUUUCUAG
51	204	AUCAUUUUUCUCAUACCUUCUGCU
51	205	CACCCACCAUCACCCUCUGUG
51	206	AUCAUCUCGUUGAUUCCUCA
51	207	CUCCAACAUCAAGGAAGAUGGCAUUUCU
51	208	CAUCAAGGAAGAUGGCAUUUCUAGU
51	265	AUCAUUUUUCUCAUACCUUCUGCUAGGAGCUAA AA

TABLE 2-continued

Oligonucleotide sequences for targeting DMD.		
EXON	SEQ ID NO:	SEQUENCE
52	209	UUGCUGGUCUUGUUUUUC
52	210	CCGUAAUGAUUGUUCU
52	211	GCUGGUCUUGUUUUUCA
52	212	UGGUCUUGUUUUUCAAAUUU
52	213	GUCUUGUUUUUCAAAUUUUG
52	214	CUUGUUUUUCAAAUUUUGG
52	215	UGUUUUUCAAAUUUUGGGC
52	216	UCCAACUGGGGACGCCUCUGUCCAAAUCUGC
52	217	UCCUGCAUUGUUGCCUGUAAG
52	218	UCCAACUGGGGACGCCUCUGUCCAAAUC
52	219	ACUGGGGACGCCUCUGUCCA
52	220	CCGUAAUGAUUGUUCUAGCC
52	221	UGUUAAAAACUUAUCUUGA
53	222	CUGUUGCCUCCGGUUCUG
53	223	UUGGCUCUGGCCUGUCCU
53	224	UUCAACUGUUGCCUCCGGUUCUGAAGGUGUUC
53	225	UACUUCAUCCACUGAUUCUGAAUU
53	226	CUGAAGGUGUUCUGUACUUCAUCC
53	227	CUGUUGCCUCCGGUUCUGAAGGUGU
53	228	CUGUUGCCUCCGGUUCUGAAGGUGUUCUUG
53	229	CAACUGUUGCCUCCGGUUCUGAAGGUGUUC
53	230	UUGCCUCCGGUUCUGAAGGUGUUCUUGAUC
53	231	GUUGCCUCCGGUUCUGAAGGUGUUC
53	232	CUCCGGUUCUGAAGGUGUUCUUG
53	233	CUCCGGUUCUGAAGGUGUUCU
53	234	CUCCGGUUCUGAAGGUGUUCU
53	235	CUCCGGUUCUGAAGGUGUUC
53	236	CUCCGGUUCUGAAGGUGUUC
53	237	CAUUAACUGUUGCCUCCGGUUCUG
53	238	CUGUUGCCUCCGGUUCUGAAGGUGU
53	239	CAUUAACUGUUGCCUCCGGUUCUGAAGGUGU
53	240	UACUAACCUUGGUUUCUGUGA
53	241	UGUAUAGGGACCCUCCUUCUGAUCUC
53	242	CUAACCUGGUUUCUGUGAUUUUCU
53	243	GGUAUCUUUGAUACUAACCUUGGUUUC
53	244	AUUCUUUCAACUAGAAUAAAAG

TABLE 2-continued

Oligonucleotide sequences for targeting DMD.		
EXON	SEQ ID NO:	SEQUENCE
53	245	GAUUCUGAAUUCUUUCAACUAGAAU
53	246	AUCCACUGAAUUCUGAAUUC
53	247	AACCGAGACCCGACAGGAUUCU
53	266	GGAAGCUAAGGAAGAAGCUGAGCAGG
55	248	CUGUUGCAGUAAUCUAUGAG
55	249	UGCCAUUGUUUCAUCAGCUCUUU
55	250	UGCAGUAAUCUAUGAGUUUC
55	251	UCCUGUAGGACAUUGGCAGU
55	252	GAGUCUUCUAGGAGCCUU

[0170] Examples of oligonucleotides for promoting DMD gene editing include International Patent Publication WO2018053632A1, published Mar. 29, 2018, entitled “METHODS OF MODIFYING THE DYSTROPHIN GENE AND RESTORING DYSTROPHIN EXPRESSION AND USES THEREOF”; International Patent Publication WO2017049407A1, published Mar. 30, 2017, entitled “MODIFICATION OF THE DYSTROPHIN GENE AND USES THEREOF”; International Patent Publication WO2016161380A1, published Oct. 6, 2016, entitled “CRISPR/CAS-RELATED METHODS AND COMPOSITIONS FOR TREATING DUCHENNE MUSCULAR DYSTROPHY AND BECKER MUSCULAR DYSTROPHY”; International Patent Publication WO2017095967, published Jun. 8, 2017, entitled “THERAPEUTIC TARGETS FOR THE CORRECTION OF THE HUMAN DYSTROPHIN GENE BY GENE EDITING AND METHODS OF USE”; International Patent Publication WO2017072590A1, published May 4, 2017, entitled “MATERIALS AND METHODS FOR TREATMENT OF DUCHENNE MUSCULAR DYSTROPHY”; International Patent Publication WO2018098480A1, published May 31, 2018, entitled “PREVENTION OF MUSCULAR DYSTROPHY BY CRISPR/CPFI-MEDIATED GENE EDITING”; US Patent Application Publication US20170266320A1, published Sep. 21, 2017, entitled “RNA-Guided Systems for In Vivo Gene Editing”; International Patent Publication WO2016025469A1, published Feb. 18, 2016, entitled “PREVENTION OF MUSCULAR DYSTROPHY BY CRISPR/CAS9-MEDIATED GENE EDITING”; U.S. Patent Application Publication 2016/0201089, published Jul. 14, 2016, entitled “RNA-GUIDED GENE EDITING AND GENE REGULATION”; and U.S. Patent Application Publication 2013/0145487, published Jun. 6, 2013, entitled “MEGANUCLEASE VARIANTS CLEAVING A DNA TARGET SEQUENCE FROM THE DYSTROPHIN GENE AND USES THEREOF”, the contents of each of which are incorporated herein in their entireties. In some embodiments, an oligonucleotide may have a region of complementarity to DMD gene sequences of multiple species, e.g., selected from human, mouse and non-human species.

[0171] In some embodiments, the oligonucleotide may have region of complementarity to a mutant DMD allele, for example, a DMD allele with at least one mutation in any of exons 1-79 of DMD in humans that leads to a frameshift and improper RNA splicing/processing.

[0172] In some embodiments, the oligonucleotide may target lncRNA or mRNA, e.g., for degradation. In some embodiments, the oligonucleotide may target, e.g., for degradation, a nucleic acid encoding a protein involved in a mismatch repair pathway, e.g., MSH2, MutLalpha, MutS-beta, MutLalpha. Non-limiting examples of proteins involved in mismatch repair pathways, for which mRNAs encoding such proteins may be targeted by oligonucleotides described herein, are described in Iyer, R. R. et al., “DNA triplet repeat expansion and mismatch repair” *Annu Rev Biochem.* 2015; 84:199-226.; and Schmidt M. H. and Pearson C. E., “Disease-associated repeat instability and mismatch repair” *DNA Repair (Amst).* 2016 February; 38:117-26.

[0173] a. Oligonucleotide Size/Sequence

[0174] Oligonucleotides may be of a variety of different lengths, e.g., depending on the format. In some embodiments, an oligonucleotide is 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 75, or more nucleotides in length. In some embodiments, the oligonucleotide is 8 to 50 nucleotides in length, 8 to 40 nucleotides in length, 8 to 30 nucleotides in length, 10 to 15 nucleotides in length, 10 to 20 nucleotides in length, 15 to 25 nucleotides in length, 21 to 23 nucleotides in lengths, etc.

[0175] In some embodiments, a complementary nucleic acid sequence of an oligonucleotide for purposes of the present disclosure is specifically hybridizable or specific for the target nucleic acid when binding of the sequence to the target molecule (e.g., mRNA) interferes with the function of the target (e.g., mRNA) to cause a change of activity (e.g., inhibiting translation, altering splicing, exon skipping) or expression (e.g., degrading a target mRNA) and there is a sufficient degree of complementarity to avoid non-specific binding of the sequence to non-target sequences under conditions in which avoidance of non-specific binding is desired, e.g., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed under suitable conditions of stringency. Thus, in some embodiments, an oligonucleotide may be at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% complementary to the consecutive nucleotides of a target nucleic acid. In some embodiments a complementary nucleotide sequence need not be 100% complementary to that of its target to be specifically hybridizable or specific for a target nucleic acid.

[0176] In some embodiments, an oligonucleotide comprises region of complementarity to a target nucleic acid that is in the range of 8 to 15, 8 to 30, 8 to 40, or 10 to 50, or 5 to 50, or 5 to 40 nucleotides in length. In some embodiments, a region of complementarity of an oligonucleotide to a target nucleic acid is 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides in length. In some embodiments, the region of complementarity is complementary with at least 8 consecutive nucleotides of a target nucleic acid. In some

embodiments, an oligonucleotide may contain 1, 2 or 3 base mismatches compared to the portion of the consecutive nucleotides of target nucleic acid. In some embodiments the oligonucleotide may have up to 3 mismatches over 15 bases, or up to 2 mismatches over 10 bases.

[0177] b. Oligonucleotide Modifications:

[0178] The oligonucleotides described herein may be modified, e.g., comprise a modified sugar moiety, a modified internucleoside linkage, a modified nucleotide and/or combinations thereof. In addition, in some embodiments, oligonucleotides may exhibit one or more of the following properties: do not mediate alternative splicing; are not immune stimulatory; are nuclease resistant; have improved cell uptake compared to unmodified oligonucleotides; are not toxic to cells or mammals; have improved endosomal exit internally in a cell; minimizes TLR stimulation; or avoid pattern recognition receptors. Any of the modified chemistries or formats of oligonucleotides described herein can be combined with each other. For example, one, two, three, four, five, or more different types of modifications can be included within the same oligonucleotide.

[0179] In some embodiments, certain nucleotide modifications may be used that make an oligonucleotide into which they are incorporated more resistant to nuclease digestion than the native oligodeoxynucleotide or oligoribonucleotide molecules; these modified oligonucleotides survive intact for a longer time than unmodified oligonucleotides. Specific examples of modified oligonucleotides include those comprising modified backbones, for example, modified internucleoside linkages such as phosphorothioates, phosphotriesters, methyl phosphonates, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages. Accordingly, oligonucleotides of the disclosure can be stabilized against nucleolytic degradation such as by the incorporation of a modification, e.g., a nucleotide modification.

[0180] In some embodiments, an oligonucleotide may be of up to 50 or up to 100 nucleotides in length in which 2 to 10, 2 to 15, 2 to 16, 2 to 17, 2 to 18, 2 to 19, 2 to 20, 2 to 25, 2 to 30, 2 to 40, 2 to 45, or more nucleotides of the oligonucleotide are modified nucleotides. The oligonucleotide may be of 8 to 30 nucleotides in length in which 2 to 10, 2 to 15, 2 to 16, 2 to 17, 2 to 18, 2 to 19, 2 to 20, 2 to 25, 2 to 30 nucleotides of the oligonucleotide are modified nucleotides. The oligonucleotide may be of 8 to 15 nucleotides in length in which 2 to 4, 2 to 5, 2 to 6, 2 to 7, 2 to 8, 2 to 9, 2 to 10, 2 to 11, 2 to 12, 2 to 13, 2 to 14 nucleotides of the oligonucleotide are modified nucleotides. Optionally, the oligonucleotides may have every nucleotide except 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides modified. Oligonucleotide modifications are described further herein.

[0181] c. Modified Nucleotides

[0182] In some embodiments, an oligonucleotide include a 2'-modified nucleotide, e.g., a 2'-deoxy, 2'-deoxy-2'-fluoro, 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-amino-propyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA).

[0183] In some embodiments, an oligonucleotide can include at least one 2'-O-methyl-modified nucleotide, and in some embodiments, all of the nucleotides include a 2'-O-methyl modification. In some embodiments, an oligonucleotide comprises modified nucleotides in which the ribose

ring comprises a bridge moiety connecting two atoms in the ring, e.g., connecting the 2'-O atom to the 4'-C atom. In some embodiments, the oligonucleotides are "locked," e.g., comprise modified nucleotides in which the ribose ring is "locked" by a methylene bridge connecting the 2'-O atom and the 4'-C atom. Examples of LNAs are described in International Patent Application Publication WO/2008/043753, published on Apr. 17, 2008, and entitled "RNA Antagonist Compounds For The Modulation Of PCSK9", the contents of which are incorporated herein by reference in its entirety.

[0184] Other modifications that may be used in the oligonucleotides disclosed herein include ethylene-bridged nucleic acids (ENAs). ENAs include, but are not limited to, 2'-O,4'-C-ethylene-bridged nucleic acids. Examples of ENAs are provided in International Patent Publication No. WO 2005/042777, published on May 12, 2005, and entitled "APP/ENA Antisense"; Morita et al., *Nucleic Acid Res., Suppl* 1:241-242, 2001; Suroño et al., *Hum. Gene Ther.*, 15:749-757, 2004; Koizumi, *Curr. Opin. Mol. Ther.*, 8:144-149, 2006 and Horie et al., *Nucleic Acids Symp. Ser (Oxf)*, 49:171-172, 2005; the disclosures of which are incorporated herein by reference in their entireties.

[0185] In some embodiments, the oligonucleotide may comprise a bridged nucleotide, such as a locked nucleic acid (LNA) nucleotide, a constrained ethyl (cEt) nucleotide, or an ethylene bridged nucleic acid (ENA) nucleotide. In some embodiments, the oligonucleotide comprises a modified nucleotide disclosed in one of the following United States Patent or Patent Application Publications: U.S. Pat. No. 7,399,845, issued on Jul. 15, 2008, and entitled "6-Modified Bicyclic Nucleic Acid Analogs"; U.S. Pat. No. 7,741,457, issued on Jun. 22, 2010, and entitled "6-Modified Bicyclic Nucleic Acid Analogs"; U.S. Pat. No. 8,022,193, issued on Sep. 20, 2011, and entitled "6-Modified Bicyclic Nucleic Acid Analogs"; U.S. Pat. No. 7,569,686, issued on Aug. 4, 2009, and entitled "Compounds And Methods For Synthesis Of Bicyclic Nucleic Acid Analogs"; U.S. Pat. No. 7,335,765, issued on Feb. 26, 2008, and entitled "Novel Nucleoside And Oligonucleotide Analogues"; U.S. Pat. No. 7,314,923, issued on Jan. 1, 2008, and entitled "Novel Nucleoside And Oligonucleotide Analogues"; U.S. Pat. No. 7,816,333, issued on Oct. 19, 2010, and entitled "Oligonucleotide Analogues And Methods Utilizing The Same" and US Publication Number 2011/0009471 now U.S. Pat. No. 8,957,201, issued on Feb. 17, 2015, and entitled "Oligonucleotide Analogues And Methods Utilizing The Same", the entire contents of each of which are incorporated herein by reference for all purposes.

[0186] In some embodiments, the oligonucleotide comprises at least one nucleotide modified at the 2' position of the sugar, preferably a 2'-O-alkyl, 2'-O-alkyl-O-alkyl or 2'-fluoro-modified nucleotide. In other preferred embodiments, RNA modifications include 2'-fluoro, 2'-amino and 2'-O-methyl modifications on the ribose of pyrimidines, abasic residues or an inverted base at the 3' end of the RNA.

[0187] In some embodiments, the oligonucleotide may have at least one modified nucleotide that results in an increase in T_m of the oligonucleotide in a range of 1° C., 2° C., 3° C., 4° C., or 5° C. compared with an oligonucleotide that does not have the at least one modified nucleotide. The oligonucleotide may have a plurality of modified nucleotides that result in a total increase in T_m of the oligonucleotide in a range of 2° C., 3° C., 4° C., 5° C., 6° C., 7° C., 8° C., 9°

C., 10° C., 15° C., 20° C., 25° C., 30° C., 35° C., 40° C., 45° C. or more compared with an oligonucleotide that does not have the modified nucleotide.

[0188] The oligonucleotide may comprise alternating nucleotides of different kinds. For example, an oligonucleotide may comprise alternating deoxyribonucleotides or ribonucleotides and 2'-fluoro-deoxyribonucleotides. An oligonucleotide may comprise alternating deoxyribonucleotides or ribonucleotides and 2'-O-methyl nucleotides. An oligonucleotide may comprise alternating 2'-fluoro nucleotides and 2'-O-methyl nucleotides. An oligonucleotide may comprise alternating bridged nucleotides and 2'-fluoro or 2'-O-methyl nucleotides.

[0189] d. Internucleotide Linkages/Backbones

[0190] In some embodiments, oligonucleotide may contain a phosphorothioate or other modified internucleotide linkage. In some embodiments, the oligonucleotide comprises phosphorothioate internucleoside linkages. In some embodiments, the oligonucleotide comprises phosphorothioate internucleoside linkages between at least two nucleotides. In some embodiments, the oligonucleotide comprises phosphorothioate internucleoside linkages between all nucleotides. For example, in some embodiments, oligonucleotides comprise modified internucleotide linkages at the first, second, and/or third internucleoside linkage at the 5' or 3' end of the nucleotide sequence.

[0191] Phosphorus-containing linkages that may be used include, but are not limited to, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates comprising 3' alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates comprising 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'; see U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050.

[0192] In some embodiments, oligonucleotides may have heteroatom backbones, such as methylene(methylimino) or MMI backbones; amide backbones (see De Mesmaeker et al. *Ace. Chem. Res.* 1995, 28:366-374); morpholino backbones (see Summerton and Weller, U.S. Pat. No. 5,034,506); or peptide nucleic acid (PNA) backbones (wherein the phosphodiester backbone of the oligonucleotide is replaced with a polyamide backbone, the nucleotides being bound directly or indirectly to the aza nitrogen atoms of the polyamide backbone, see Nielsen et al., *Science* 1991, 254, 1497).

[0193] e. Stereospecific Oligonucleotides

[0194] In some embodiments, internucleotidic phosphorus atoms of oligonucleotides are chiral, and the properties of the oligonucleotides by adjusted based on the configuration of the chiral phosphorus atoms. In some embodiments, appropriate methods may be used to synthesize P-chiral oligonucleotide analogs in a stereocontrolled manner (e.g., as described in Oka N, Wada T, Stereocontrolled synthesis of oligonucleotide analogs containing chiral internucleotidic phosphorus atoms. *Chem Soc Rev.* 2011 December; 40(12):

5829-43.) In some embodiments, phosphorothioate containing oligonucleotides comprise nucleoside units that are joined together by either substantially all Sp or substantially all Rp phosphorothioate intersugar linkages are provided. In some embodiments, such phosphorothioate oligonucleotides having substantially chirally pure intersugar linkages are prepared by enzymatic or chemical synthesis, as described, for example, in U.S. Pat. No. 5,587,261, issued on Dec. 12, 1996, the contents of which are incorporated herein by reference in their entirety. In some embodiments, chirally controlled oligonucleotides provide selective cleavage patterns of a target nucleic acid. For example, in some embodiments, a chirally controlled oligonucleotide provides single site cleavage within a complementary sequence of a nucleic acid, as described, for example, in US Patent Application Publication 20170037399 A1, published on Feb. 2, 2017, entitled "CHIRAL DESIGN", the contents of which are incorporated herein by reference in their entirety.

[0195] f. Morpholinos

[0196] In some embodiments, the oligonucleotide may be a morpholino-based compounds. Morpholino-based oligomeric compounds are described in Dwayne A. Braasch and David R. Corey, *Biochemistry*, 2002, 41(14), 4503-4510; *Genesis*, volume 30, issue 3, 2001; Heasman, J., *Dev. Biol.*, 2002, 243, 209-214; Nasevicius et al., *Nat. Genet.*, 2000, 26, 216-220; Lacerra et al., *Proc. Natl. Acad. Sci.*, 2000, 97, 9591-9596; and U.S. Pat. No. 5,034,506, issued Jul. 23, 1991. In some embodiments, the morpholino-based oligomeric compound is a phosphorodiamidate morpholino oligomer (PMO) (e.g., as described in Iverson, *Curr. Opin. Mol. Ther.*, 3:235-238, 2001; and Wang et al., *J. Gene Med.*, 12:354-364, 2010; the disclosures of which are incorporated herein by reference in their entireties).

[0197] g. Peptide Nucleic Acids (PNAs)

[0198] In some embodiments, both a sugar and an internucleoside linkage (the backbone) of the nucleotide units of an oligonucleotide are replaced with novel groups. In some embodiments, the base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, for example, an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative publication that report the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

[0199] h. Gappers

[0200] In some embodiments, the oligonucleotide is a gapper. A gapper oligonucleotide generally has the formula 5'-X—Y—Z-3', with X and Z as flanking regions around a gap region Y. In some embodiments, the Y region is a contiguous stretch of nucleotides, e.g., a region of at least 6 DNA nucleotides, which are capable of recruiting an RNase, such as RNase H. In some embodiments, the gapper binds to the target nucleic acid, at which point an RNase is recruited and can then cleave the target nucleic acid. In some embodiments, the Y region is flanked both 5' and 3' by regions X and Z comprising high-affinity modified

nucleotides, e.g., one to six modified nucleotides. Examples of modified nucleotides include, but are not limited to, 2' MOE or 2'OMe or Locked Nucleic Acid bases (LNA). The flanking sequences X and Z may be of one to twenty nucleotides, one to eight nucleotides or one to five nucleotides in length, in some embodiments. The flanking sequences X and Z may be of similar length or of dissimilar lengths. The gap-segment Y may be a nucleotide sequence of five to twenty nucleotides, size to twelve nucleotides or six to ten nucleotides in length, in some embodiments.

[0201] In some embodiments, the gap region of the gapmer oligonucleotides may contain modified nucleotides known to be acceptable for efficient RNase H action in addition to DNA nucleotides, such as C4'-substituted nucleotides, acyclic nucleotides, and arabino-configured nucleotides. In some embodiments, the gap region comprises one or more unmodified internucleosides. In some embodiments, one or both flanking regions each independently comprise one or more phosphorothioate internucleoside linkages (e.g., phosphorothioate internucleoside linkages or other linkages) between at least two, at least three, at least four, at least five or more nucleotides. In some embodiments, the gap region and two flanking regions each independently comprise modified internucleoside linkages (e.g., phosphorothioate internucleoside linkages or other linkages) between at least two, at least three, at least four, at least five or more nucleotides.

[0202] A gapmer may be produced using appropriate methods. Representative U.S. patents, U.S. patent publications, and PCT publications that teach the preparation of gapmers include, but are not limited to, U.S. Pat. Nos. 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; 5,700,922; 5,898,031; 7,432,250; and 7,683,036; U.S. patent publication Nos. US20090286969, US20100197762, and US20110112170; and PCT publication Nos. WO2008049085 and WO2009090182, each of which is herein incorporated by reference in its entirety.

[0203] i. Mixmers

[0204] In some embodiments, an oligonucleotide described herein may be a mixmer or comprise a mixmer sequence pattern. In general, mixmers are oligonucleotides that comprise both naturally and non-naturally occurring nucleotides or comprise two different types of non-naturally occurring nucleotides typically in an alternating pattern. Mixmers generally have higher binding affinity than unmodified oligonucleotides and may be used to specifically bind a target molecule, e.g., to block a binding site on the target molecule. Generally, mixmers do not recruit an RNase to the target molecule and thus do not promote cleavage of the target molecule. Such oligonucleotides that are incapable of recruiting RNase H have been described, for example, see WO2007/112754 or WO2007/112753.

[0205] In some embodiments, the mixmer comprises or consists of a repeating pattern of nucleotide analogues and naturally occurring nucleotides, or one type of nucleotide analogue and a second type of nucleotide analogue. However, a mixmer need not comprise a repeating pattern and may instead comprise any arrangement of modified nucleotides and naturally occurring nucleotides or any arrangement of one type of modified nucleotide and a second type of modified nucleotide. The repeating pattern, may, for instance be every second or every third nucleotide is a modified nucleotide, such as LNA, and the remaining nucleotides are

naturally occurring nucleotides, such as DNA, or are a 2' substituted nucleotide analogue such as 2'MOE or 2' fluoro analogues, or any other modified nucleotide described herein. It is recognized that the repeating pattern of modified nucleotide, such as LNA units, may be combined with modified nucleotide at fixed positions—e.g. at the 5' or 3' termini.

[0206] In some embodiments, a mixmer does not comprise a region of more than 5, more than 4, more than 3, or more than 2 consecutive naturally occurring nucleotides, such as DNA nucleotides. In some embodiments, the mixmer comprises at least a region consisting of at least two consecutive modified nucleotide, such as at least two consecutive LNAs. In some embodiments, the mixmer comprises at least a region consisting of at least three consecutive modified nucleotide units, such as at least three consecutive LNAs.

[0207] In some embodiments, the mixmer does not comprise a region of more than 7, more than 6, more than 5, more than 4, more than 3, or more than 2 consecutive nucleotide analogues, such as LNAs. In some embodiments, LNA units may be replaced with other nucleotide analogues, such as those referred to herein.

[0208] Mixmers may be designed to comprise a mixture of affinity enhancing modified nucleotides, such as in non-limiting example LNA nucleotides and 2'-O-methyl nucleotides. In some embodiments, a mixmer comprises modified internucleoside linkages (e.g., phosphorothioate internucleoside linkages or other linkages) between at least two, at least three, at least four, at least five or more nucleotides.

[0209] A mixmer may be produced using any suitable method. Representative U.S. patents, U.S. patent publications, and PCT publications that teach the preparation of mixmers include U.S. patent publication Nos. US20060128646, US20090209748, US20090298916, US20110077288, and US20120322851, and U.S. Pat. No. 7,687,617.

[0210] In some embodiments, a mixmer comprises one or more morpholino nucleotides. For example, in some embodiments, a mixmer may comprise morpholino nucleotides mixed (e.g., in an alternating manner) with one or more other nucleotides (e.g., DNA, RNA nucleotides) or modified nucleotides (e.g., LNA, 2'-O-Methyl nucleotides).

[0211] In some embodiments, mixmers are useful for splice correcting or exon skipping, for example, as reported in Touznik A., et al., *LNA/DNA mixmer-based antisense oligonucleotides correct alternative splicing of the SMN2 gene and restore SMN protein expression in type 1 SMA fibroblasts* *Scientific Reports*, volume 7, Article number: 3672 (2017), Chen S. et al., *Synthesis of a Morpholino Nucleic Acid (MNA)-Uridine Phosphoramidite, and Exon Skipping Using MNA/2'-O-Methyl Mixmer Antisense Oligonucleotide*, *Molecules* 2016, 21, 1582, the contents of each which are incorporated herein by reference.

[0212] j. RNA Interference (RNAi)

[0213] In some embodiments, oligonucleotides provided herein may be in the form of small interfering RNAs (siRNA), also known as short interfering RNA or silencing RNA. siRNA, is a class of double-stranded RNA molecules, typically about 20-25 base pairs in length that target nucleic acids (e.g., mRNAs) for degradation via the RNA interference (RNAi) pathway in cells. Specificity of siRNA molecules may be determined by the binding of the antisense strand of the molecule to its target RNA. Effective siRNA molecules are generally less than 30 to 35 base pairs in

length to prevent the triggering of non-specific RNA interference pathways in the cell via the interferon response, although longer siRNA can also be effective.

[0214] Following selection of an appropriate target RNA sequence, siRNA molecules that comprise a nucleotide sequence complementary to all or a portion of the target sequence, i.e. an antisense sequence, can be designed and prepared using appropriate methods (see, e.g., PCT Publication Number WO 2004/016735; and U.S. Patent Publication Nos. 2004/0077574 and 2008/0081791).

[0215] The siRNA molecule can be double stranded (i.e. a dsRNA molecule comprising an antisense strand and a complementary sense strand) or single-stranded (i.e. a ssRNA molecule comprising just an antisense strand). The siRNA molecules can comprise a duplex, asymmetric duplex, hairpin or asymmetric hairpin secondary structure, having self-complementary sense and antisense strands.

[0216] Double-stranded siRNA may comprise RNA strands that are the same length or different lengths. Double-stranded siRNA molecules can also be assembled from a single oligonucleotide in a stem-loop structure, wherein self-complementary sense and antisense regions of the siRNA molecule are linked by means of a nucleic acid based or non-nucleic acid-based linker(s), as well as circular single-stranded RNA having two or more loop structures and a stem comprising self-complementary sense and antisense strands, wherein the circular RNA can be processed either in vivo or in vitro to generate an active siRNA molecule capable of mediating RNAi. Small hairpin RNA (shRNA) molecules thus are also contemplated herein. These molecules comprise a specific antisense sequence in addition to the reverse complement (sense) sequence, typically separated by a spacer or loop sequence. Cleavage of the spacer or loop provides a single-stranded RNA molecule and its reverse complement, such that they may anneal to form a dsRNA molecule (optionally with additional processing steps that may result in addition or removal of one, two, three or more nucleotides from the 3' end and/or the 5' end of either or both strands). A spacer can be of a sufficient length to permit the antisense and sense sequences to anneal and form a double-stranded structure (or stem) prior to cleavage of the spacer (and, optionally, subsequent processing steps that may result in addition or removal of one, two, three, four, or more nucleotides from the 3' end and/or the 5' end of either or both strands). A spacer sequence may be an unrelated nucleotide sequence that is situated between two complementary nucleotide sequence regions which, when annealed into a double-stranded nucleic acid, comprise a shRNA.

[0217] The overall length of the siRNA molecules can vary from about 14 to about 100 nucleotides depending on the type of siRNA molecule being designed. Generally between about 14 and about 50 of these nucleotides are complementary to the RNA target sequence, i.e. constitute the specific antisense sequence of the siRNA molecule. For example, when the siRNA is a double- or single-stranded siRNA, the length can vary from about 14 to about 50 nucleotides, whereas when the siRNA is a shRNA or circular molecule, the length can vary from about 40 nucleotides to about 100 nucleotides.

[0218] An siRNA molecule may comprise a 3' overhang at one end of the molecule, The other end may be blunt-ended or have also an overhang (5' or 3'). When the siRNA molecule comprises an overhang at both ends of the mol-

ecule, the length of the overhangs may be the same or different. In one embodiment, the siRNA molecule of the present disclosure comprises 3' overhangs of about 1 to about 3 nucleotides on both ends of the molecule.

[0219] k. microRNA (miRNAs)

[0220] In some embodiments, an oligonucleotide may be a microRNA (miRNA). MicroRNAs (referred to as "miRNAs") are small non-coding RNAs, belonging to a class of regulatory molecules that control gene expression by binding to complementary sites on a target RNA transcript. Typically, miRNAs are generated from large RNA precursors (termed pri-miRNAs) that are processed in the nucleus into approximately 70 nucleotide pre-miRNAs, which fold into imperfect stem-loop structures. These pre-miRNAs typically undergo an additional processing step within the cytoplasm where mature miRNAs of 18-25 nucleotides in length are excised from one side of the pre-miRNA hairpin by an RNase III enzyme, Dicer.

[0221] As used herein, miRNAs including pri-miRNA, pre-miRNA, mature miRNA or fragments of variants thereof that retain the biological activity of mature miRNA. In one embodiment, the size range of the miRNA can be from 21 nucleotides to 170 nucleotides. In one embodiment the size range of the miRNA is from 70 to 170 nucleotides in length. In another embodiment, mature miRNAs of from 21 to 25 nucleotides in length can be used.

[0222] l. Aptamers

[0223] In some embodiments, oligonucleotides provided herein may be in the form of aptamers. Generally, in the context of molecular payloads, aptamer is any nucleic acid that binds specifically to a target, such as a small molecule, protein, nucleic acid in a cell. In some embodiments, the aptamer is a DNA aptamer or an RNA aptamer. In some embodiments, a nucleic acid aptamer is a single-stranded DNA or RNA (ssDNA or ssRNA). It is to be understood that a single-stranded nucleic acid aptamer may form helices and/or loop structures. The nucleic acid that forms the nucleic acid aptamer may comprise naturally occurring nucleotides, modified nucleotides, naturally occurring nucleotides with hydrocarbon linkers (e.g., an alkylene) or a polyether linker (e.g., a PEG linker) inserted between one or more nucleotides, modified nucleotides with hydrocarbon or PEG linkers inserted between one or more nucleotides, or a combination of thereof. Exemplary publications and patents describing aptamers and method of producing aptamers include, e.g., Lorsch and Szostak, 1996; Jayasena, 1999; U.S. Pat. Nos. 5,270,163; 5,567,588; 5,650,275; 5,670,637; 5,683,867; 5,696,249; 5,789,157; 5,843,653; 5,864,026; 5,989,823; 6,569,630; 8,318,438 and PCT application WO 99/31275, each incorporated herein by reference.

[0224] m. Ribozymes

[0225] In some embodiments, oligonucleotides provided herein may be in the form of a ribozyme. A ribozyme (ribonucleic acid enzyme) is a molecule, typically an RNA molecule, that is capable of performing specific biochemical reactions, similar to the action of protein enzymes. Ribozymes are molecules with catalytic activities including the ability to cleave at specific phosphodiester linkages in RNA molecules to which they have hybridized, such as mRNAs, RNA-containing substrates, lncRNAs, and ribozymes, themselves.

[0226] Ribozymes may assume one of several physical structures, one of which is called a "hammerhead." A hammerhead ribozyme is composed of a catalytic core

containing nine conserved bases, a double-stranded stem and loop structure (stem-loop II), and two regions complementary to the target RNA flanking regions the catalytic core. The flanking regions enable the ribozyme to bind to the target RNA specifically by forming double-stranded stems I and III. Cleavage occurs in cis (i.e., cleavage of the same RNA molecule that contains the hammerhead motif) or in trans (cleavage of an RNA substrate other than that containing the ribozyme) next to a specific ribonucleotide triplet by a transesterification reaction from a 3', 5'-phosphate diester to a 2', 3'-cyclic phosphate diester. Without wishing to be bound by theory, it is believed that this catalytic activity requires the presence of specific, highly conserved sequences in the catalytic region of the ribozyme.

[0227] Modifications in ribozyme structure have also included the substitution or replacement of various non-core portions of the molecule with non-nucleotidic molecules. For example, Benseler et al. (J. Am. Chem. Soc. (1993) 115:8483-8484) disclosed hammerhead-like molecules in which two of the base pairs of stem II, and all four of the nucleotides of loop II were replaced with non-nucleoside linkers based on hexaethylene glycol, propanediol, bis(triethylene glycol) phosphate, tris(propanediol)bisphosphate, or bis(propanediol) phosphate. Ma et al. (Biochem. (1993) 32:1751-1758; Nucleic Acids Res. (1993) 21:2585-2589) replaced the six nucleotide loop of the TAR ribozyme hairpin with non-nucleotidic, ethylene glycol-related linkers. Thomson et al. (Nucleic Acids Res. (1993) 21:5600-5603) replaced loop II with linear, non-nucleotidic linkers of 13, 17, and 19 atoms in length.

[0228] Ribozyme oligonucleotides can be prepared using well known methods (see, e.g., PCT Publications WO9118624; WO9413688; WO9201806; and WO 92/07065; and U.S. Pat. Nos. 5,436,143 and 5,650,502) or can be purchased from commercial sources (e.g., US Biochemicals) and, if desired, can incorporate nucleotide analogs to increase the resistance of the oligonucleotide to degradation by nucleases in a cell. The ribozyme may be synthesized in any known manner, e.g., by use of a commercially available synthesizer produced, e.g., by Applied Biosystems, Inc. or Milligen. The ribozyme may also be produced in recombinant vectors by conventional means. See, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory (Current edition). The ribozyme RNA sequences may be synthesized conventionally, for example, by using RNA polymerases such as T7 or SP6.

[0229] n. Guide Nucleic Acids

[0230] In some embodiments, oligonucleotides are guide nucleic acid, e.g., guide RNA (gRNA) molecules. Generally, a guide RNA is a short synthetic RNA composed of (1) a scaffold sequence that binds to a nucleic acid programmable DNA binding protein (napDNAbp), such as Cas9, and (2) a nucleotide spacer portion that defines the DNA target sequence (e.g., genomic DNA target) to which the gRNA binds in order to bring the nucleic acid programmable DNA binding protein in proximity to the DNA target sequence. In some embodiments, the napDNAbp is a nucleic acid-programmable protein that forms a complex with (e.g., binds or associates with) one or more RNA(s) that targets the nucleic acid-programmable protein to a target DNA sequence (e.g., a target genomic DNA sequence). In some embodiments, a nucleic acid-programmable nuclease, when in a complex with an RNA, may be referred to as a nuclease:RNA

complex. Guide RNAs can exist as a complex of two or more RNAs, or as a single RNA molecule.

[0231] Guide RNAs (gRNAs) that exist as a single RNA molecule may be referred to as single-guide RNAs (sgRNAs), though gRNA is also used to refer to guide RNAs that exist as either single molecules or as a complex of two or more molecules. Typically, gRNAs that exist as a single RNA species comprise two domains: (1) a domain that shares homology to a target nucleic acid (i.e., directs binding of a Cas9 complex to the target); and (2) a domain that binds a Cas9 protein. In some embodiments, domain (2) corresponds to a sequence known as a tracrRNA and comprises a stem-loop structure. In some embodiments, domain (2) is identical or homologous to a tracrRNA as provided in Jinek et al., Science 337:816-821 (2012), the entire contents of which is incorporated herein by reference.

[0232] In some embodiments, a gRNA comprises two or more of domains (1) and (2), and may be referred to as an extended gRNA. For example, an extended gRNA will bind two or more Cas9 proteins and bind a target nucleic acid at two or more distinct regions, as described herein. The gRNA comprises a nucleotide sequence that complements a target site, which mediates binding of the nuclease:RNA complex to said target site, providing the sequence specificity of the nuclease:RNA complex. In some embodiments, the RNA-programmable nuclease is the (CRISPR-associated system) Cas9 endonuclease, for example, Cas9 (CsnI) from *Streptococcus pyogenes* (see, e.g., "Complete genome sequence of an M1 strain of *Streptococcus pyogenes*." Ferretti J. J., McShan W. M., Ajdic D. J., Savic D. J., Savic G., Lyon K., Primeaux C., Sezate S., Suvorov A. N., Kenton S., Lai H. S., Lin S. P., Qian Y., Jia H. G., Najjar F. Z., Ren Q., Zhu H., Song L., White J., Yuan X., Clifton S. W., Roe B. A., McLaughlin R. E., Proc. Natl. Acad. Sci. U.S.A. 98:4658-4663 (2001); "CRISPR RNA maturation by trans-encoded small RNA and host factor RNase III." Deltcheva E., Chylinski K., Sharma C. M., Gonzales K., Chao Y., Pirzada Z. A., Eckert M. R., Vogel J., Charpentier E., Nature 471:602-607 (2011); and "A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity." Jinek M., Chylinski K., Fonfara I., Hauer M., Doudna J. A., Charpentier E. Science 337:816-821 (2012), the entire contents of each of which are incorporated herein by reference.

[0233] o. Multimers

[0234] In some embodiments, molecular payloads may comprise multimers (e.g., concatemers) of 2 or more oligonucleotides connected by a linker. In this way, in some embodiments, the oligonucleotide loading of a complex can be increased beyond the available linking sites on a targeting agent (e.g., available thiol sites on an antibody) or otherwise tuned to achieve a particular payload loading content. Oligonucleotides in a multimer can be the same or different (e.g., targeting different genes or different sites on the same gene or products thereof).

[0235] In some embodiments, multimers comprise 2 or more oligonucleotides linked together by a cleavable linker. However, in some embodiments, multimers comprise 2 or more oligonucleotides linked together by a non-cleavable linker. In some embodiments, a multimer comprises 2, 3, 4, 5, 6, 7, 8, 9, 10 or more oligonucleotides linked together. In some embodiments, a multimer comprises 2 to 5, 2 to 10 or 4 to 20 oligonucleotides linked together.

[0236] In some embodiments, a multimer comprises 2 or more oligonucleotides linked end-to-end (in a linear

arrangement). In some embodiments, a multimer comprises 2 or more oligonucleotides linked end-to-end via a oligonucleotide based linker (e.g., poly-dT linker, an abasic linker). In some embodiments, a multimer comprises a 5' end of one oligonucleotide linked to a 3' end of another oligonucleotide. In some embodiments, a multimer comprises a 3' end of one oligonucleotide linked to a 3' end of another oligonucleotide. In some embodiments, a multimer comprises a 5' end of one oligonucleotide linked to a 5' end of another oligonucleotide. Still, in some embodiments, multimers can comprise a branched structure comprising multiple oligonucleotides linked together by a branching linker.

[0237] Further examples of multimers that may be used in the complexes provided herein are disclosed, for example, in US Patent Application Number 2015/0315588 A1, entitled *Methods of delivering multiple targeting oligonucleotides to a cell using cleavable linkers*, which was published on Nov. 5, 2015; US Patent Application Number 2015/0247141 A1, entitled *Multimeric Oligonucleotide Compounds*, which was published on Sep. 3, 2015, US Patent Application Number US 2011/0158937 A1, entitled *Immunostimulatory Oligonucleotide Multimers*, which was published on Jun. 30, 2011; and U.S. Pat. No. 5,693,773, entitled *Triplex-Forming Antisense Oligonucleotides Having Abasic Linkers Targeting Nucleic Acids Comprising Mixed Sequences Of Purines And Pyrimidines*, which issued on Dec. 2, 1997, the contents of each of which are incorporated herein by reference in their entireties.

[0238] ii. Small Molecules:

[0239] Any suitable small molecule may be used as a molecular payload, as described herein. In some embodiments, the small molecule enhances exon skipping of DMD mutant sequences. In some embodiments, the small molecule is as described in US Patent Application Publication US20140080896A1, published Mar. 20, 2014, entitled "IDENTIFICATION OF SMALL MOLECULES THAT FACILITATE THERAPEUTIC EXON SKIPPING". Further examples of small molecule payloads are provided in U.S. Pat. No. 9,982,260, issued May 29, 2018, entitled "Identification of structurally similar small molecules that enhance therapeutic exon skipping". For example, in some embodiments, the small molecule is an enhancer of exon skipping such as perphenazine, flupentixol, zuclopenthixol or corynanthine. In some embodiments, a small molecule enhancer of exon skipping inhibits the ryanodine receptor or calmodulin. In some embodiments, the small molecule is an H-Ras pathway inhibitor such as manumycin A. In some embodiments, the small molecule is a suppressor of stop codons and desensitizes ribosomes to premature stop codons. In some embodiments, the small molecule is ataluren, as described in McElroy S. P. et al. "A Lack of Premature Termination Codon Read Through Efficacy of PTC124 (Ataluren) in a Diverse Array of Reporter Assays." *PLOS Biology*, published Jun. 25, 2013. In some embodiments, the small molecule is a corticosteroid, e.g., as described in Manzur, A. Y. et al. "Glucocorticoid corticosteroids for Duchenne muscular dystrophy". *Cochrane Database Syst Rev*. 2004; (2):CD003725. In some embodiments, the small molecule upregulates the expression and/or activity of genes that can replace the function of dystrophin, such as utrophin. In some embodiments, a utrophin modulator is as described in International Publication No. WO2007091106, published Aug. 16, 2007, entitled

"TREATMENT OF DUCHENNE MUSCULAR DYSTROPHY" and/or International Publication No. WO/2017/168151, published Oct. 5, 2017, entitled "COMPOSITION FOR THE TREATMENT OF DUCHENNE MUSCULAR DYSTROPHY".

[0240] iii. Peptides/Proteins

[0241] Any suitable peptide or protein may be used as a molecular payload, as described herein. In some embodiments, a protein is an enzyme. In some embodiments, peptides or proteins may be produced, synthesized, and/or derivatized using several methodologies, e.g. phage displayed peptide libraries, one-bead one-compound peptide libraries, or positional scanning synthetic peptide combinatorial libraries. Exemplary methodologies have been characterized in the art and are incorporated by reference (Gray, B. P. and Brown, K. C. "Combinatorial Peptide Libraries: Mining for Cell-Binding Peptides" *Chem Rev*. 2014, 114:2, 1020-1081.; Samoylova, T. I. and Smith, B. F. "Elucidation of muscle-binding peptides by phage display screening." *Muscle Nerve*, 1999, 22:4. 460-6.).

[0242] In some embodiments, a peptide may facilitate exon skipping in an mRNA expressed from a mutated DMD allele. In some embodiments, a peptide may promote the expression of functional dystrophin and/or the expression of a protein capable of functioning in place of dystrophin. In some embodiments, payload is a protein that is a functional fragment of dystrophin, e.g. an amino acid segment of a functional dystrophin protein.

[0243] In some embodiments, the peptide or protein comprises at least one zinc finger.

[0244] In some embodiments, the peptide or protein may comprise about 2-25 amino acids, about 2-20 amino acids, about 2-15 amino acids, about 2-10 amino acids, or about 2-5 amino acids. The peptide or protein may comprise naturally-occurring amino acids, e.g. cysteine, alanine, or non-naturally-occurring or modified amino acids. Non-naturally occurring amino acids include β -amino acids, homo-amino acids, proline derivatives, 3-substituted alanine derivatives, linear core amino acids, N-methyl amino acids, and others known in the art. In some embodiments, the peptide may be linear; in other embodiments, the peptide may be cyclic, e.g. bicyclic.

[0245] iv. Nucleic Acid Constructs

[0246] Any suitable gene expression construct may be used as a molecular payload, as described herein. In some embodiments, a gene expression construct may be a vector or a cDNA fragment. In some embodiments, a gene expression construct may be messenger RNA (mRNA). In some embodiments, a mRNA used herein may be a modified mRNA, e.g., as described in U.S. Pat. No. 8,710,200, issued on Apr. 24, 2014, entitled "*Engineered nucleic acids encoding a modified erythropoietin and their expression*". In some embodiments, a mRNA may comprise a 5' methyl cap. In some embodiments, a mRNA may comprise a polyA tail, optionally of up to 160 nucleotides in length. A gene expression construct may encode a sequence of a dystrophin protein, a dystrophin fragment, a mini-dystrophin, a utrophin protein, or any protein that shares a common function with dystrophin. In some embodiments, the gene expression construct may be expressed, e.g., overexpressed, within the nucleus of a muscle cell. In some embodiments, the gene expression constructs encodes a protein that comprises at least one zinc finger. In some embodiments, the gene expression construct encodes a protein that promotes the expres-

sion of dystrophin or a protein that shares function with dystrophin, e.g., utrophin. In some embodiments, the gene expression construct encodes a gene editing enzyme. In some embodiments, the gene expression construct is as described in U.S. Patent Application Publication US20170368198A1, published Dec. 28, 2017, entitled "Optimized mini-dystrophin genes and expression cassettes and their use"; Duan D. "Myodys, a full-length dystrophin plasmid vector for Duchenne and Becker muscular dystrophy gene therapy." *Curr Opin Mol Ther* 2008; 10:86-94; and expression cassettes disclosed in Tang, Y. et al., "AAV-directed muscular dystrophy gene therapy" *Expert Opin Biol Ther.* 2010 March; 10(3):395-408; the contents of each of which are incorporated herein by reference in their entireties.

[0247] C. Linkers

[0248] Complexes described herein generally comprise a linker that connects a muscle-targeting agent to a molecular payload. A linker comprises at least one covalent bond. In some embodiments, a linker may be a single bond, e.g., a disulfide bond or disulfide bridge, that connects a muscle-targeting agent to a molecular payload. However, in some embodiments, a linker may connect a muscle-targeting agent to a molecular through multiple covalent bonds. In some embodiments, a linker may be a cleavable linker. However, in some embodiments, a linker may be a non-cleavable linker. A linker is generally stable in vitro and in vivo, and may be stable in certain cellular environments. Additionally, generally a linker does not negatively impact the functional properties of either the muscle-targeting agent or the molecular payload. Examples and methods of synthesis of linkers are known in the art (see, e.g. Kline, T. et al. "Methods to Make Homogenous Antibody Drug Conjugates." *Pharmaceutical Research*, 2015, 32:11, 3480-3493.; Jain, N. et al. "Current ADC Linker Chemistry" *Pharm Res.* 2015, 32:11, 3526-3540.; McCombs, J. R. and Owen, S. C. "Antibody Drug Conjugates: Design and Selection of Linker, Payload and Conjugation Chemistry" *AAPS J.* 2015, 17:2, 339-351.).

[0249] A precursor to a linker typically will contain two different reactive species that allow for attachment to both the muscle-targeting agent and a molecular payload. In some embodiments, the two different reactive species may be a nucleophile and/or an electrophile. In some embodiments, a linker is connected to a muscle-targeting agent via conjugation to a lysine residue or a cysteine residue of the muscle-targeting agent. In some embodiments, a linker is connected to a cysteine residue of a muscle-targeting agent via a maleimide-containing linker, wherein optionally the

maleimide-containing linker comprises a maleimidocaproyl or maleimidomethyl cyclohexane-1-carboxylate group. In some embodiments, a linker is connected to a cysteine residue of a muscle-targeting agent or thiol functionalized molecular payload via a 3-arylpropionitrile functional group. In some embodiments, a linker is connected to a muscle-targeting agent and/or a molecular payload via an amide bond, a hydrazide, a trizaole, a thioether, or a disulfide bond.

[0250] i. Cleavable Linkers


[0251] A cleavable linker may be a protease-sensitive linker, a pH-sensitive linker, or a glutathione-sensitive linker. These linkers are generally cleavable only intracellularly and are preferably stable in extracellular environments, e.g. extracellular to a muscle cell.

[0252] Protease-sensitive linkers are cleavable by protease enzymatic activity. These linkers typically comprise peptide sequences and may be 2-10 amino acids, about 2-5 amino acids, about 5-10 amino acids, about 10 amino acids, about 5 amino acids, about 3 amino acids, or about 2 amino acids in length. In some embodiments, a peptide sequence may comprise naturally-occurring amino acids, e.g. cysteine, alanine, or non-naturally-occurring or modified amino acids. Non-naturally occurring amino acids include β -amino acids, homo-amino acids, proline derivatives, 3-substituted alanine derivatives, linear core amino acids, N-methyl amino acids, and others known in the art. In some embodiments, a protease-sensitive linker comprises a valine-citrulline or alanine-citrulline dipeptide sequence. In some embodiments, a protease-sensitive linker can be cleaved by a lysosomal protease, e.g. cathepsin B, and/or an endosomal protease.


[0253] A pH-sensitive linker is a covalent linkage that readily degrades in high or low pH environments. In some embodiments, a pH-sensitive linker may be cleaved at a pH in a range of 4 to 6. In some embodiments, a pH-sensitive linker comprises a hydrazone or cyclic acetal. In some embodiments, a pH-sensitive linker is cleaved within an endosome or a lysosome.

[0254] In some embodiments, a glutathione-sensitive linker comprises a disulfide moiety. In some embodiments, a glutathione-sensitive linker is cleaved by a disulfide exchange reaction with a glutathione species inside a cell. In some embodiments, the disulfide moiety further comprises at least one amino acid, e.g. a cysteine residue.

[0255] In some embodiments, the linker is a Val-cit linker (e.g., as described in U.S. Pat. No. 6,214,345, incorporated herein by reference). In some embodiments, before conjugation, the val-cit linker has a structure of:



[0256] In some embodiments, after conjugation, the val-cit linker has a structure of:



agent, e.g. an antibody, through a lysine or cysteine residue present on the muscle-targeting agent

[0257] ii. Non-Cleavable Linkers

[0258] In some embodiments, non-cleavable linkers may be used. Generally, a non-cleavable linker cannot be readily degraded in a cellular or physiological environment. In some embodiments, a non-cleavable linker comprises an optionally substituted alkyl group, wherein the substitutions may include halogens, hydroxyl groups, oxygen species, and other common substitutions. In some embodiments, a linker may comprise an optionally substituted alkyl, an optionally substituted alkylene, an optionally substituted arylene, a heteroarylene, a peptide sequence comprising at least one non-natural amino acid, a truncated glycan, a sugar or sugars that cannot be enzymatically degraded, an azide, an alkyne-azide, a peptide sequence comprising a LPXT sequence, a thioether, a biotin, a biphenyl, repeating units of polyethylene glycol or equivalent compounds, acid esters, acid amides, sulfamides, and/or an alkoxy-amine linker. In some embodiments, sortase-mediated ligation will be utilized to covalently link a muscle-targeting agent comprising a LPXT sequence to a molecular payload comprising a (G)_n sequence (see, e.g. Proft T. Sortase-mediated protein ligation: an emerging biotechnology tool for protein modification and immobilization. *Biotechnol Lett.* 2010, 32(1):1-10.).

[0259] In some embodiments, a linker may comprise a substituted alkylene, an optionally substituted alkenylene, an optionally substituted alkynylene, an optionally substituted cycloalkylene, an optionally substituted cycloalkenylene, an optionally substituted arylene, an optionally substituted heteroarylene further comprising at least one heteroatom selected from N, O, and S; an optionally substituted heterocyclylene further comprising at least one heteroatom selected from N, O, and S; an imino, an optionally substituted nitrogen species, an optionally substituted oxygen species O, an optionally substituted sulfur species, or a poly(alkylene oxide), e.g. polyethylene oxide or polypropylene oxide.

[0260] iii. Linker Conjugation

[0261] In some embodiments, a linker is connected to a muscle-targeting agent and/or molecular payload via a phosphate, thioether, ether, carbon-carbon, or amide bond. In some embodiments, a linker is connected to an oligonucleotide through a phosphate or phosphorothioate group, e.g. a terminal phosphate of an oligonucleotide backbone. In some embodiments, a linker is connected to an muscle-targeting

[0262] In some embodiments, a linker is connected to a muscle-targeting agent and/or molecular payload by a cycloaddition reaction between an azide and an alkyne to form a triazole, wherein the azide and the alkyne may be located on the muscle-targeting agent, molecular payload, or the linker. In some embodiments, an alkyne may be a cyclic alkyne, e.g., a cyclooctyne. In some embodiments, an alkyne may be bicyclononyne (also known as bicyclo[6.1.0]nonyne or BCN) or substituted bicyclononyne. In some embodiments, a cyclooctane is as described in International Patent Application Publication WO2011136645, published on Nov. 3, 2011, entitled, “*Fused Cyclooctyne Compounds And Their Use In Metal free Click Reactions*”. In some embodiments, an azide may be a sugar or carbohydrate molecule that comprises an azide. In some embodiments, an azide may be 6-azido-6-deoxygalactose or 6-azido-N-acetylgalactosamine. In some embodiments, a sugar or carbohydrate molecule that comprises an azide is as described in International Patent Application Publication WO2016170186, published on Oct. 27, 2016, entitled, “*Process For The Modification Of A Glycoprotein Using A Glycosyltransferase That Is Or Is Derived From A β (1,4)-N-Acetylgalactosaminyltransferase*”. In some embodiments, a cycloaddition reaction between an azide and an alkyne to form a triazole, wherein the azide and the alkyne may be located on the muscle-targeting agent, molecular payload, or the linker is as described in International Patent Application Publication WO2014065661, published on May 1, 2014, entitled, “*Modified antibody, antibody-conjugate and process for the preparation thereof*”; or International Patent Application Publication WO2016170186, published on Oct. 27, 2016, entitled, “*Process For The Modification Of A Glycoprotein Using A Glycosyltransferase That Is Or Is Derived From A β (1,4)-N-Acetylgalactosaminyltransferase*”.


[0263] In some embodiments, a linker further comprises a spacer, e.g., a polyethylene glycol spacer or an acyl/carbomoyl sulfamide spacer, e.g., a HydraSpace™ spacer. In some embodiments, a spacer is as described in Verkade, J. M. M. et al., “*A Polar Sulfamide Spacer Significantly Enhances the Manufacturability, Stability, and Therapeutic Index of Antibody-Drug Conjugates*”, *Antibodies*, 2018, 7, 12.

[0264] In some embodiments, a linker is connected to a muscle-targeting agent and/or molecular payload by the Diels-Alder reaction between a dienophile and a diene/

hetero-diene, wherein the dienophile and the diene/hetero-diene may be located on the muscle-targeting agent, molecular payload, or the linker. In some embodiments a linker is connected to a muscle-targeting agent and/or molecular payload by other pericyclic reactions, e.g. ene reaction. In some embodiments, a linker is connected to a muscle-targeting agent and/or molecular payload by an amide, thioamide, or sulfonamide bond reaction. In some embodiments, a linker is connected to a muscle-targeting agent

herein may be used. In some embodiments, the linker is linked to the 5' end, the 3' end, or internally of the oligonucleotide. In some embodiments, the linker is linked to the antibody via a thiol-reactive linkage (e.g., via a cysteine in the antibody).

[0268] An exemplary structure of a complex comprising a transferrin receptor antibody covalently linked to an oligonucleotide via a Val-cit linker is provided below:



and/or molecular payload by a condensation reaction to form an oxime, hydrazone, or semicarbazide group existing between the linker and the muscle-targeting agent and/or molecular payload.

[0265] In some embodiments, a linker is connected to a muscle-targeting agent and/or molecular payload by a conjugate addition reactions between a nucleophile, e.g. an amine or a hydroxyl group, and an electrophile, e.g. a carboxylic acid or an aldehyde. In some embodiments, a nucleophile may exist on a linker and an electrophile may exist on a muscle-targeting agent or molecular payload prior to a reaction between a linker and a muscle-targeting agent or molecular payload. In some embodiments, an electrophile may exist on a linker and a nucleophile may exist on a muscle-targeting agent or molecular payload prior to a reaction between a linker and a muscle-targeting agent or molecular payload. In some embodiments, an electrophile may be an azide, a silicon centers, a carbonyl, a carboxylic acid, an anhydride, an isocyanate, a thioisocyanate, a succinimidyl ester, a sulfosuccinimidyl ester, a maleimide, an alkyl halide, an alkyl pseudohalide, an epoxide, an episulfide, an aziridine, an aryl, an activated phosphorus center, and/or an activated sulfur center. In some embodiments, a nucleophile may be an optionally substituted alkene, an optionally substituted alkyne, an optionally substituted aryl, an optionally substituted heterocyclyl, a hydroxyl group, an amino group, an alkylamino group, an anilido group, or a thiol group.

[0266] D. Examples of Antibody-Molecular Payload Complexes

[0267] Other aspects of the present disclosure provide complexes comprising any one the muscle targeting agent (e.g., a transferrin receptor antibodies) described herein covalently linked to any of the molecular payloads (e.g., an oligonucleotide) described herein. In some embodiments, the muscle targeting agent (e.g., a transferrin receptor antibody) is covalently linked to a molecular payload (e.g., an oligonucleotide) via a linker. Any of the linkers described

wherein the linker is linked to the 5' end, the 3' end, or internally of the oligonucleotide, and wherein the linker is linked to the antibody via a thiol-reactive linkage (e.g., via a cysteine in the antibody).

[0269] It should be appreciated that antibodies can be linked to oligonucleotides with different stoichiometries, a property that may be referred to as a drug to antibody ratios (DAR) with the “drug” being the oligonucleotide. In some embodiments, one oligonucleotide is linked to an antibody (DAR=1). In some embodiments, two oligonucleotides are linked to an antibody (DAR=2). In some embodiments, three oligonucleotides are linked to an antibody (DAR=3). In some embodiments, four oligonucleotides are linked to an antibody (DAR=4). In some embodiments, a mixture of different complexes, each having a different DAR, is provided. In some embodiments, an average DAR of complexes in such a mixture may be in a range of 1 to 3, 1 to 4, 1 to 5 or more. DAR may be increased by conjugating oligonucleotides to different sites on an antibody and/or by conjugating multimers to one or more sites on antibody. For example, a DAR of 2 may be achieved by conjugating a single oligonucleotide to two different sites on an antibody or by conjugating a dimer oligonucleotide to a single site of an antibody.

[0270] In some embodiments, the complex described herein comprises a transferrin receptor antibody (e.g., an antibody or any variant thereof as described herein) covalently linked to an oligonucleotide (e.g., an oligonucleotide that is capable of inducing exon skipping of DMD). In some embodiments, the complex described herein comprises a transferrin receptor antibody (e.g., an antibody or any variant thereof as described herein) covalently linked to an oligonucleotide (e.g., an oligonucleotide that is capable of inducing exon skipping of DMD) via a linker (e.g., a Val-cit linker). In some embodiments, the linker (e.g., a Val-cit linker) is linked to the 5' end, the 3' end, or internally of the nucleotide (e.g., an oligonucleotide that is capable of inducing exon skipping of DMD). In some embodiments, the linker (e.g., a Val-cit linker) is linked to the antibody (e.g.,

an antibody or any variant thereof as described herein) via a thiol-reactive linkage (e.g., via a cysteine in the antibody).

[0271] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide, wherein the transferrin receptor antibody comprises a CDR-H1, a CDR-H2, and a CDR-H3 that are the same as the CDR-H1, CDR-H2, and CDR-H3 shown in Table 1.1; and a CDR-L1, a CDR-L2, and a CDR-L3 that are the same as the CDR-L1, CDR-L2, and CDR-L3 shown in Table 1.1.


[0272] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide, wherein the transferrin receptor antibody comprises a VH having the amino acid sequence of SEQ ID NO: 283 and a VL having the amino acid sequence of SEQ ID NO: 284. In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide, wherein the transferrin receptor antibody comprises a VH having the amino acid sequence of SEQ ID NO: 285 and a VL having the amino acid sequence of SEQ ID NO: 286.

[0273] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide, wherein the transferrin receptor antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO: 289 and a light chain having the amino acid sequence of SEQ ID NO: 290. In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide, wherein the transferrin receptor antibody comprises a heavy chain having the amino

acid sequence of SEQ ID NO: 283 and a VL having the amino acid sequence of SEQ ID NO: 284. In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide via a linker (e.g., a Val-cit linker), wherein the transferrin receptor antibody comprises a VH having the amino acid sequence of SEQ ID NO: 285 and a VL having the amino acid sequence of SEQ ID NO: 286.

[0276] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide via a linker (e.g., a Val-cit linker), wherein the transferrin receptor antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO: 289 and a light chain having the amino acid sequence of SEQ ID NO: 290. In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide via a linker (e.g., a Val-cit linker), wherein the transferrin receptor antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO: 291 and a light chain having the amino acid sequence of SEQ ID NO: 292.

[0277] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide via a Val-cit linker, wherein the transferrin receptor antibody comprises a CDR-H1, a CDR-H2, and a CDR-H3 that are the same as the CDR-H1, CDR-H2, and CDR-H3 shown in Table 1.1; and a CDR-L1, a CDR-L2, and a CDR-L3 that are the same as the CDR-L1, CDR-L2, and CDR-L3 shown in Table 1.1, and wherein the complex comprises the structure of:




acid sequence of SEQ ID NO: 291 and a light chain having the amino acid sequence of SEQ ID NO: 292.

[0274] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide via a linker (e.g., a Val-cit linker), wherein the transferrin receptor antibody comprises a CDR-H1, a CDR-H2, and a CDR-H3 that are the same as the CDR-H1, CDR-H2, and CDR-H3 shown in Table 1.1; and a CDR-L1, a CDR-L2, and a CDR-L3 that are the same as the CDR-L1, CDR-L2, and CDR-L3 shown in Table 1.1.

[0275] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide via a linker (e.g.,

[0278] wherein the linker Val-cit linker is linked to the 5' end, the 3' end, or internally of the DMD targeting oligonucleotide, and wherein the Val-cit linker is linked to the antibody (e.g., an antibody or any variant thereof as described herein) via a thiol-reactive linkage (e.g., via a cysteine in the antibody).

[0279] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide via a Val-cit linker, wherein the transferrin receptor antibody comprises a VH having the amino acid sequence of SEQ ID NO: 283 and a VL having the amino acid sequence of SEQ ID NO: 284, and wherein the complex comprises the structure of:




wherein the linker Val-cit linker is linked to the 5' end, the 3' end, or internally of DMD targeting oligonucleotide, and wherein the Val-cit linker is linked to the antibody (e.g., an antibody or any variant thereof as described herein) via a thiol-reactive linkage (e.g., via a cysteine in the antibody).

[0280] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide via a Val-cit linker, wherein the transferrin receptor antibody comprises a VH having the amino acid sequence of SEQ ID NO: 285 and a VL having the amino acid sequence of SEQ ID NO: 286, and wherein the complex comprises the structure of:


and wherein the Val-cit linker is linked to the antibody (e.g., an antibody or any variant thereof as described herein) via a thiol-reactive linkage (e.g., via a cysteine in the antibody).

[0281] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide via a Val-cit linker, wherein the transferrin receptor antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO: 289 and a light chain having the amino acid sequence of




wherein the linker Val-cit linker is linked to the 5' end, the 3' end, or internally of the DMD targeting oligonucleotide,

SEQ ID NO: 290, and wherein the complex comprises the structure of:



[0282] wherein the linker Val-cit linker is linked to the 5' end, the 3' end, or internally of DMD targeting oligonucleotide, and wherein the Val-cit linker is linked to the antibody (e.g., an antibody or any variant thereof as described herein) via a thiol-reactive linkage (e.g., via a cysteine in the antibody).

[0283] In some embodiments, the complex described herein comprises a transferrin receptor antibody covalently linked to a DMD targeting oligonucleotide via a Val-cit linker, wherein the transferrin receptor antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO: 291 and a light chain having the amino acid sequence of SEQ ID NO: 292, and wherein the complex comprises the structure of:



wherein the linker Val-cit linker is linked to the 5' end, the 3' end, or internally of DMD targeting oligonucleotide, and wherein the Val-cit linker is linked to the antibody (e.g., an antibody or any variant thereof as described herein) via a thiol-reactive linkage (e.g., via a cysteine in the antibody).

III. Formulations

[0284] Complexes provided herein may be formulated in any suitable manner. Generally, complexes provided herein are formulated in a manner suitable for pharmaceutical use. For example, complexes can be delivered to a subject using a formulation that minimizes degradation, facilitates delivery and/or uptake, or provides another beneficial property to the complexes in the formulation. In some embodiments, provided herein are compositions comprising complexes and pharmaceutically acceptable carriers. Such compositions can be suitably formulated such that when administered to a subject, either into the immediate environment of a target cell or systemically, a sufficient amount of the complexes enter target muscle cells. In some embodiments, complexes are formulated in buffer solutions such as phosphate-buffered saline solutions, liposomes, micellar structures, and capsids.

[0285] It should be appreciated that, in some embodiments, compositions may include separately one or more components of complexes provided herein (e.g., muscle-targeting agents, linkers, molecular payloads, or precursor molecules of any one of them).

[0286] In some embodiments, complexes are formulated in water or in an aqueous solution (e.g., water with pH adjustments). In some embodiments, complexes are formulated in basic buffered aqueous solutions (e.g., PBS). In some embodiments, formulations as disclosed herein com-

prise an excipient. In some embodiments, an excipient confers to a composition improved stability, improved absorption, improved solubility and/or therapeutic enhancement of the active ingredient. In some embodiments, an excipient is a buffering agent (e.g., sodium citrate, sodium phosphate, a tris base, or sodium hydroxide) or a vehicle (e.g., a buffered solution, petrolatum, dimethyl sulfoxide, or mineral oil).

[0287] In some embodiments, a complex or component thereof (e.g., oligonucleotide or antibody) is lyophilized for extending its shelf-life and then made into a solution before use (e.g., administration to a subject). Accordingly, an excipient in a composition comprising a complex, or component thereof, described herein may be a lyoprotectant

(e.g., mannitol, lactose, polyethylene glycol, or polyvinyl pyrrolidone), or a collapse temperature modifier (e.g., dextran, ficoll, or gelatin).

[0288] In some embodiments, a pharmaceutical composition is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, administration. Typically, the route of administration is intravenous or subcutaneous.

[0289] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. In some embodiments, formulations include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, and sodium chloride in the composition. Sterile injectable solutions can be prepared by incorporating the a complexes in a required amount in a selected solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization.

[0290] In some embodiments, a composition may contain at least about 0.1% of the a complex, or component thereof, or more, although the percentage of the active ingredient(s) may be between about 1% and about 80% or more of the weight or volume of the total composition. Factors such as solubility, bioavailability, biological half-life, route of administration, product shelf life, as well as other pharmacological considerations will be contemplated by one skilled

in the art of preparing such pharmaceutical formulations, and as such, a variety of dosages and treatment regimens may be desirable.

IV. Methods of Use/Treatment

[0291] Complexes comprising a muscle-targeting agent covalently to a molecular payload as described herein are effective in treating a subject having a dystrophinopathy, e.g., duchenne muscular dystrophy. In some embodiments, complexes comprise a molecular payload that is an oligonucleotide, e.g., an antisense oligonucleotide that facilitates exon skipping of an mRNA expressed from a mutated DMD allele.

[0292] In some embodiments, a subject may be a human subject, a non-human primate subject, a rodent subject, or any suitable mammalian subject. In some embodiments, a subject may have duchenne muscular dystrophy or other dystrophinopathy. In some embodiments, a subject has a mutated DMD allele, which may optionally comprise at least one mutation in a DMD exon that causes a frameshift mutation and leads to improper RNA splicing/processing. In some embodiments, a subject is suffering from symptoms of a severe dystrophinopathy, e.g. muscle atrophy or muscle loss. In some embodiments, a subject has an asymptomatic increase in serum concentration of creatine phosphokinase (CK) and/or muscle cramps with myoglobinuria. In some embodiments, a subject has a progressive muscle disease, such as Duchenne or Becker muscular dystrophy or DMD-associated dilated cardiomyopathy (DCM). In some embodiments, a subject is not suffering from symptoms of a dystrophinopathy.

[0293] An aspect of the disclosure includes a methods involving administering to a subject an effective amount of a complex as described herein. In some embodiments, an effective amount of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently to a molecular payload can be administered to a subject in need of treatment. In some embodiments, a pharmaceutical composition comprising a complex as described herein may be administered by a suitable route, which may include intravenous administration, e.g., as a bolus or by continuous infusion over a period of time. In some embodiments, intravenous administration may be performed by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, or intrathecal routes. In some embodiments, a pharmaceutical composition may be in solid form, aqueous form, or a liquid form. In some embodiments, an aqueous or liquid form may be nebulized or lyophilized. In some embodiments, a nebulized or lyophilized form may be reconstituted with an aqueous or liquid solution.

[0294] Compositions for intravenous administration may contain various carriers such as vegetable oils, dimethylacetamide, dimethylformamide, ethyl lactate, ethyl carbonate, isopropyl myristate, ethanol, and polyols (glycerol, propylene glycol, liquid polyethylene glycol, and the like). For intravenous injection, water soluble antibodies can be administered by the drip method, whereby a pharmaceutical formulation containing the antibody and a physiologically acceptable excipients is infused. Physiologically acceptable excipients may include, for example, 5% dextrose, 0.9% saline, Ringer's solution or other suitable excipients. Intramuscular preparations, e.g., a sterile formulation of a suitable soluble salt form of the antibody, can be dissolved and

administered in a pharmaceutical excipient such as Water-for-Injection, 0.9% saline, or 5% glucose solution.

[0295] In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently to a molecular payload is administered via site-specific or local delivery techniques. Examples of these techniques include implantable depot sources of the complex, local delivery catheters, site specific carriers, direct injection, or direct application.

[0296] In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently to a molecular payload is administered at an effective concentration that confers therapeutic effect on a subject. Effective amounts vary, as recognized by those skilled in the art, depending on the severity of the disease, unique characteristics of the subject being treated, e.g. age, physical conditions, health, or weight, the duration of the treatment, the nature of any concurrent therapies, the route of administration and related factors. These related factors are known to those in the art and may be addressed with no more than routine experimentation. In some embodiments, an effective concentration is the maximum dose that is considered to be safe for the patient. In some embodiments, an effective concentration will be the lowest possible concentration that provides maximum efficacy.

[0297] Empirical considerations, e.g. the half-life of the complex in a subject, generally will contribute to determination of the concentration of pharmaceutical composition that is used for treatment. The frequency of administration may be empirically determined and adjusted to maximize the efficacy of the treatment.

[0298] Generally, for administration of any of the complexes described herein, an initial candidate dosage may be about 1 to 100 mg/kg, or more, depending on the factors described above, e.g. safety or efficacy. In some embodiments, a treatment will be administered once. In some embodiments, a treatment will be administered daily, biweekly, weekly, bimonthly, monthly, or at any time interval that provide maximum efficacy while minimizing safety risks to the subject. Generally, the efficacy and the treatment and safety risks may be monitored throughout the course of treatment

[0299] The efficacy of treatment may be assessed using any suitable methods. In some embodiments, the efficacy of treatment may be assessed by evaluation of observation of symptoms associated with a dystrophinopathy, e.g. muscle atrophy or muscle weakness, through measures of a subject's self-reported outcomes, e.g. mobility, self-care, usual activities, pain/discomfort, and anxiety/depression, or by quality-of-life indicators, e.g. lifespan.

[0300] In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently to a molecular payload described herein is administered to a subject at an effective concentration sufficient to modulate activity or expression of a target gene by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 95% relative to a control, e.g. baseline level of gene expression prior to treatment.

[0301] In some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently to a molecular payload described herein to a subject is sufficient to inhibit activity or expression of a target gene for at

least 1-5, 1-10, 5-15, 10-20, 15-30, 20-40, 25-50, or more days. In some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently to a molecular payload described herein to a subject is sufficient to inhibit activity or expression of a target gene for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 weeks. In some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently to a molecular payload described herein to a subject is sufficient to inhibit activity or expression of a target gene for at least 1, 2, 3, 4, 5, or 6 months.

[0302] In some embodiments, a pharmaceutical composition may comprise more than one complex comprising a muscle-targeting agent covalently to a molecular payload. In some embodiments, a pharmaceutical composition may further comprise any other suitable therapeutic agent for treatment of a subject, e.g. a human subject having a dystrophinopathy. In some embodiments, the other therapeutic agents may enhance or supplement the effectiveness of the complexes described herein. In some embodiments, the other therapeutic agents may function to treat a different symptom or disease than the complexes described herein.

EXAMPLES

Example 1: Targeting HPRT with Transfected Antisense Oligonucleotides

[0303] A siRNA that targets hypoxanthine phosphoribosyltransferase (HPRT) was tested in vitro for its ability to reduce expression levels of HPRT in an immortalized cell line. Briefly, Hepa 1-6 cells were transfected with either a control siRNA (siCTRL; 100 nM) or the siRNA that targets HPRT (siHPRT; 100 nM), formulated with lipofectamine 2000. HPRT expression levels were evaluated 48 hours following transfection. A control experiment was also performed in which vehicle (phosphate-buffered saline) was delivered to Hepa 1-6 cells in culture and the cells were maintained for 48 hours. As shown in FIG. 1, it was found that the HPRT siRNA reduced HPRT expression levels by ~90% compared with controls.

TABLE 3

Sequences of siHPRT and siCTRL	
	Sequence
siHPRT sense strand	5' - UcCuAuGaCuGuAgAuUdUaU - (CH ₂) ₆ NH ₂ -3'
siHPRT antisense strand	5' - paUaAaAuCuAcAgUcAuAg GasAsu-3'
siCTRL sense strand	5' - UgUaAuAaCcAuAuCuAcCuU - (CH ₂) ₆ NH ₂ -3'
siCTRL antisense strand	5' - aAgGuAgAuAuGgUuAuUaCas Asa-3'

*Lower case - 2'Ome ribose;

Capital letter - 2'Fluoro ribose;

p - phosphate linkage;

s - phosphorothioate linkage

Example 2: Targeting HPRT with a Muscle-Targeting Complex

[0304] A muscle-targeting complex was generated comprising the HPRT siRNA used in Example 1 (siHPRT) covalently linked, via a non-cleavable N-gamma-maleimido-butyl-oxysuccinimide ester (GMBS) linker, to DTX-A-002, an anti-transferrin receptor antibody.

[0305] Briefly, the GMBS linker was dissolved in dry DMSO and coupled to the 3' end of the sense strand of siHPRT through amide bond formation under aqueous conditions. Completion of the reaction was verified by Kaiser test. Excess linker and organic solvents were removed by gel permeation chromatography. The purified, maleimide functionalized sense strand of siHPRT was then coupled to DTX-A-002 antibody using a Michael addition reaction.

[0306] The product of the antibody coupling reaction was then subjected to hydrophobic interaction chromatography (HIC-HPLC). antiTfR-siHPRT complexes comprising one or two siHPRT molecules covalently attached to DTX-A-002 antibody were purified. Densitometry confirmed that the purified sample of complexes had an average siHPRT to antibody ratio of 1.46. SDS-PAGE analysis demonstrated that >90% of the purified sample of complexes comprised DTX-A-002 linked to either one or two siHPRT molecules.

[0307] Using the same methods as described above, a control IgG2a-siHPRT complex was generated comprising the HPRT siRNA used in Example 1 (siHPRT) covalently linked via the GMBS linker to an IgG2a (Fab) antibody (DTX-A-003). Densitometry confirmed that DTX-C-001 had an average siHPRT to antibody ratio of 1.46 and SDS-PAGE demonstrated that >90% of the purified sample of control complexes comprised DTX-A-003 linked to either one or two siHPRT molecules.

[0308] The antiTfR-siHPRT complex was then tested for cellular internalization and inhibition of HPRT in cellulo. Hepa 1-6 cells, which have relatively high expression levels of transferrin receptor, were incubated in the presence of vehicle (phosphate-buffered saline), IgG2a-siHPRT (100 nM), antiTfR-siCTRL (100 nM), or antiTfR-siHPRT (100 nM), for 72 hours. After the 72 hour incubation, the cells were isolated and assayed for expression levels of HPRT (FIG. 2). Cells treated with the antiTfR-siHPRT demonstrated a reduction in HPRT expression by ~50% relative to the cells treated with the vehicle control. Meanwhile, cells treated with either of the IgG2a-siHPRT or antiTfR-siCTRL had HPRT expression levels comparable to the vehicle control (no reduction in HPRT expression). These data indicate that the anti-transferrin receptor antibody of the antiTfR-siHPRT enabled cellular internalization of the complex, thereby allowing the siHPRT to inhibit expression of HPRT.

Example 3: Targeting HPRT in Mouse Muscle Tissues with a Muscle-Targeting Complex

[0309] The muscle-targeting complex described in Example 2, antiTfR-siHPRT, was tested for inhibition of HPRT in mouse tissues. C57BL/6 wild-type mice were intravenously injected with a single dose of a vehicle control (phosphate-buffered saline); siHPRT (2 mg/kg of RNA); IgG2a-siHPRT (2 mg/kg of RNA, corresponding to 9 mg/kg antibody complex); or antiTfR-siHPRT (2 mg/kg of RNA, corresponding to 9 mg/kg antibody complex). Each experimental condition was replicated in four individual C57BL/6

wild-type mice. Following a three-day period after injection, the mice were euthanized and segmented into isolated tissue types. Individual tissue samples were subsequently assayed for expression levels of HPRT (FIGS. 3A-3B and 4A-4E).

[0310] Mice treated with the antiTfR-siHPRT complex demonstrated a reduction in HPRT expression in gastrocnemius (31% reduction; $p < 0.05$) and heart (30% reduction; $p < 0.05$), relative to the mice treated with the siHPRT control (FIGS. 3A-3B). Meanwhile, mice treated with the IgG2a-siHPRT complex had HPRT expression levels comparable to the siHPRT control (little or no reduction in HPRT expression) for all assayed muscle tissue types.

[0311] Mice treated with the antiTfR-siHPRT complex demonstrated no change in HPRT expression in non-muscle tissues such as brain, liver, lung, kidney, and spleen tissues (FIGS. 4A-4E). These data indicate that the anti-transferrin receptor antibody of the antiTfR-siHPRT complex enabled cellular internalization of the complex into muscle-specific tissues in an in vivo mouse model, thereby allowing the siHPRT to inhibit expression of HPRT. These data further demonstrate that the antiTfR-oligonucleotide complexes of the current disclosure are capable of specifically targeting muscle tissues.

Example 4: Targeting DMD with a Muscle-Targeting Complex

[0312] A muscle-targeting complex is generated comprising an antisense oligonucleotide that targets a mutant allele of DMD (DMD ASO), for exon skipping, e.g., an oligonucleotide having a sequence as disclosed in Table 2, covalently linked, via a cathepsin cleavable linker, to DTX-A-002 (R17 217 (Fab)), an anti-transferrin receptor antibody.

[0313] Briefly, a maleimidocaproyl-L-valine-L-citrulline-p-aminobenzyl alcohol p-nitrophenyl carbonate (MC-Val-Cit-PABC-PNP) linker molecule is coupled to $\text{NH}_2\text{-C}_6\text{-DMD ASO}$ using an amide coupling reaction. Excess linker and organic solvents are removed by gel permeation chromatography. The purified Val-Cit-linker-DMD ASO is then coupled to a thiol-reactive anti-transferrin receptor antibody (DTX-A-002).

[0314] The product of the antibody coupling reaction is then subjected to hydrophobic interaction chromatography (HIC-HPLC) to purify the muscle-targeting complex. Densitometry and SDS-PAGE analysis of the purified complex allow for determination of the average ratio of ASO-to-antibody and total purity, respectively.

[0315] Using the same methods as described above, a control complex is generated comprising DMD ASO covalently linked via a Val-Cit linker to an IgG2a (Fab) antibody. The purified muscle-targeting complex comprising DTX-A-002 covalently linked to DMD ASO is then tested for cellular internalization and modulation of DMD exon skipping. Disease-relevant muscle cells that have relatively high expression levels of transferrin receptor, are incubated in the presence of vehicle control (saline), muscle-targeting complex (100 nM), or control complex (100 nM) for 72 hours. After the 72 hour incubation, the cells are isolated and assayed for expression levels of DMD.

EQUIVALENTS AND TERMINOLOGY

[0316] The disclosure illustratively described herein suitably can be practiced in the absence of any element or elements, limitation or limitations that are not specifically

disclosed herein. Thus, for example, in each instance herein any of the terms “comprising”, “consisting essentially of”, and “consisting of” may be replaced with either of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the disclosure. Thus, it should be understood that although the present disclosure has been specifically disclosed by preferred embodiments, optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this disclosure.

[0317] In addition, where features or aspects of the disclosure are described in terms of Markush groups or other grouping of alternatives, those skilled in the art will recognize that the disclosure is also thereby described in terms of any individual member or subgroup of members of the Markush group or other group.

[0318] It should be appreciated that, in some embodiments, sequences presented in the sequence listing may be referred to in describing the structure of an oligonucleotide or other nucleic acid. In such embodiments, the actual oligonucleotide or other nucleic acid may have one or more alternative nucleotides (e.g., an RNA counterpart of a DNA nucleotide or a DNA counterpart of an RNA nucleotide) and/or one or more modified nucleotides and/or one or more modified internucleotide linkages and/or one or more other modification compared with the specified sequence while retaining essentially same or similar complementary properties as the specified sequence.

[0319] The use of the terms “a” and “an” and “the” and similar referents in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The terms “comprising,” “having,” “including,” and “containing” are to be construed as open-ended terms (i.e., meaning “including, but not limited to,”) unless otherwise noted. Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein, is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

[0320] Embodiments of this invention are described herein. Variations of those embodiments may become apparent to those of ordinary skill in the art upon reading the foregoing description.

[0321] The inventors expect skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than as specifically described herein. Accordingly, this invention includes all

modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise

clearly contradicted by context. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 294

<210> SEQ ID NO 1

<211> LENGTH: 760

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Met Met Asp Gln Ala Arg Ser Ala Phe Ser Asn Leu Phe Gly Gly Glu
1          5          10          15
Pro Leu Ser Tyr Thr Arg Phe Ser Leu Ala Arg Gln Val Asp Gly Asp
20        25        30
Asn Ser His Val Glu Met Lys Leu Ala Val Asp Glu Glu Glu Asn Ala
35        40        45
Asp Asn Asn Thr Lys Ala Asn Val Thr Lys Pro Lys Arg Cys Ser Gly
50        55        60
Ser Ile Cys Tyr Gly Thr Ile Ala Val Ile Val Phe Phe Leu Ile Gly
65        70        75        80
Phe Met Ile Gly Tyr Leu Gly Tyr Cys Lys Gly Val Glu Pro Lys Thr
85        90        95
Glu Cys Glu Arg Leu Ala Gly Thr Glu Ser Pro Val Arg Glu Glu Pro
100       105       110
Gly Glu Asp Phe Pro Ala Ala Arg Arg Leu Tyr Trp Asp Asp Leu Lys
115       120       125
Arg Lys Leu Ser Glu Lys Leu Asp Ser Thr Asp Phe Thr Gly Thr Ile
130       135       140
Lys Leu Leu Asn Glu Asn Ser Tyr Val Pro Arg Glu Ala Gly Ser Gln
145       150       155       160
Lys Asp Glu Asn Leu Ala Leu Tyr Val Glu Asn Gln Phe Arg Glu Phe
165       170       175
Lys Leu Ser Lys Val Trp Arg Asp Gln His Phe Val Lys Ile Gln Val
180       185       190
Lys Asp Ser Ala Gln Asn Ser Val Ile Ile Val Asp Lys Asn Gly Arg
195       200       205
Leu Val Tyr Leu Val Glu Asn Pro Gly Gly Tyr Val Ala Tyr Ser Lys
210       215       220
Ala Ala Thr Val Thr Gly Lys Leu Val His Ala Asn Phe Gly Thr Lys
225       230       235       240
Lys Asp Phe Glu Asp Leu Tyr Thr Pro Val Asn Gly Ser Ile Val Ile
245       250       255
Val Arg Ala Gly Lys Ile Thr Phe Ala Glu Lys Val Ala Asn Ala Glu
260       265       270
Ser Leu Asn Ala Ile Gly Val Leu Ile Tyr Met Asp Gln Thr Lys Phe
275       280       285
Pro Ile Val Asn Ala Glu Leu Ser Phe Phe Gly His Ala His Leu Gly
290       295       300
Thr Gly Asp Pro Tyr Thr Pro Gly Phe Pro Ser Phe Asn His Thr Gln
305       310       315       320

```


-continued

Gln Asn Asn Gly Ala Phe Asn Glu Thr Leu Phe Arg Asn Gln Leu Ala
725 730 735

Leu Ala Thr Trp Thr Ile Gln Gly Ala Ala Asn Ala Leu Ser Gly Asp
740 745 750

Val Trp Asp Ile Asp Asn Glu Phe
755 760

<210> SEQ ID NO 2
<211> LENGTH: 760
<212> TYPE: PRT
<213> ORGANISM: Macaca mulatta

<400> SEQUENCE: 2

Met Met Asp Gln Ala Arg Ser Ala Phe Ser Asn Leu Phe Gly Gly Glu
1 5 10 15

Pro Leu Ser Tyr Thr Arg Phe Ser Leu Ala Arg Gln Val Asp Gly Asp
20 25 30

Asn Ser His Val Glu Met Lys Leu Gly Val Asp Glu Glu Glu Asn Thr
35 40 45

Asp Asn Asn Thr Lys Pro Asn Gly Thr Lys Pro Lys Arg Cys Gly Gly
50 55 60

Asn Ile Cys Tyr Gly Thr Ile Ala Val Ile Ile Phe Phe Leu Ile Gly
65 70 75 80

Phe Met Ile Gly Tyr Leu Gly Tyr Cys Lys Gly Val Glu Pro Lys Thr
85 90 95

Glu Cys Glu Arg Leu Ala Gly Thr Glu Ser Pro Ala Arg Glu Glu Pro
100 105 110

Glu Glu Asp Phe Pro Ala Ala Pro Arg Leu Tyr Trp Asp Asp Leu Lys
115 120 125

Arg Lys Leu Ser Glu Lys Leu Asp Thr Thr Asp Phe Thr Ser Thr Ile
130 135 140

Lys Leu Leu Asn Glu Asn Leu Tyr Val Pro Arg Glu Ala Gly Ser Gln
145 150 155 160

Lys Asp Glu Asn Leu Ala Leu Tyr Ile Glu Asn Gln Phe Arg Glu Phe
165 170 175

Lys Leu Ser Lys Val Trp Arg Asp Gln His Phe Val Lys Ile Gln Val
180 185 190

Lys Asp Ser Ala Gln Asn Ser Val Ile Ile Val Asp Lys Asn Gly Gly
195 200 205

Leu Val Tyr Leu Val Glu Asn Pro Gly Gly Tyr Val Ala Tyr Ser Lys
210 215 220

Ala Ala Thr Val Thr Gly Lys Leu Val His Ala Asn Phe Gly Thr Lys
225 230 235 240

Lys Asp Phe Glu Asp Leu Asp Ser Pro Val Asn Gly Ser Ile Val Ile
245 250 255

Val Arg Ala Gly Lys Ile Thr Phe Ala Glu Lys Val Ala Asn Ala Glu
260 265 270

Ser Leu Asn Ala Ile Gly Val Leu Ile Tyr Met Asp Gln Thr Lys Phe
275 280 285

Pro Ile Val Lys Ala Asp Leu Ser Phe Phe Gly His Ala His Leu Gly
290 295 300

Thr Gly Asp Pro Tyr Thr Pro Gly Phe Pro Ser Phe Asn His Thr Gln
305 310 315 320

-continued

Phe Pro Pro Ser Gln Ser Ser Gly Leu Pro Asn Ile Pro Val Gln Thr
325 330 335
Ile Ser Arg Ala Ala Ala Glu Lys Leu Phe Gly Asn Met Glu Gly Asp
340 345 350
Cys Pro Ser Asp Trp Lys Thr Asp Ser Thr Cys Lys Met Val Thr Ser
355 360 365
Glu Asn Lys Ser Val Lys Leu Thr Val Ser Asn Val Leu Lys Glu Thr
370 375 380
Lys Ile Leu Asn Ile Phe Gly Val Ile Lys Gly Phe Val Glu Pro Asp
385 390 395 400
His Tyr Val Val Val Gly Ala Gln Arg Asp Ala Trp Gly Pro Gly Ala
405 410 415
Ala Lys Ser Ser Val Gly Thr Ala Leu Leu Leu Lys Leu Ala Gln Met
420 425 430
Phe Ser Asp Met Val Leu Lys Asp Gly Phe Gln Pro Ser Arg Ser Ile
435 440 445
Ile Phe Ala Ser Trp Ser Ala Gly Asp Phe Gly Ser Val Gly Ala Thr
450 455 460
Glu Trp Leu Glu Gly Tyr Leu Ser Ser Leu His Leu Lys Ala Phe Thr
465 470 475 480
Tyr Ile Asn Leu Asp Lys Ala Val Leu Gly Thr Ser Asn Phe Lys Val
485 490 495
Ser Ala Ser Pro Leu Leu Tyr Thr Leu Ile Glu Lys Thr Met Gln Asp
500 505 510
Val Lys His Pro Val Thr Gly Arg Ser Leu Tyr Gln Asp Ser Asn Trp
515 520 525
Ala Ser Lys Val Glu Lys Leu Thr Leu Asp Asn Ala Ala Phe Pro Phe
530 535 540
Leu Ala Tyr Ser Gly Ile Pro Ala Val Ser Phe Cys Phe Cys Glu Asp
545 550 555 560
Thr Asp Tyr Pro Tyr Leu Gly Thr Thr Met Asp Thr Tyr Lys Glu Leu
565 570 575
Val Glu Arg Ile Pro Glu Leu Asn Lys Val Ala Arg Ala Ala Ala Glu
580 585 590
Val Ala Gly Gln Phe Val Ile Lys Leu Thr His Asp Thr Glu Leu Asn
595 600 605
Leu Asp Tyr Glu Arg Tyr Asn Ser Gln Leu Leu Leu Phe Leu Arg Asp
610 615 620
Leu Asn Gln Tyr Arg Ala Asp Val Lys Glu Met Gly Leu Ser Leu Gln
625 630 635 640
Trp Leu Tyr Ser Ala Arg Gly Asp Phe Phe Arg Ala Thr Ser Arg Leu
645 650 655
Thr Thr Asp Phe Arg Asn Ala Glu Lys Arg Asp Lys Phe Val Met Lys
660 665 670
Lys Leu Asn Asp Arg Val Met Arg Val Glu Tyr Tyr Phe Leu Ser Pro
675 680 685
Tyr Val Ser Pro Lys Glu Ser Pro Phe Arg His Val Phe Trp Gly Ser
690 695 700
Gly Ser His Thr Leu Ser Ala Leu Leu Glu Ser Leu Lys Leu Arg Arg
705 710 715 720

-continued

Gln Asn Asn Ser Ala Phe Asn Glu Thr Leu Phe Arg Asn Gln Leu Ala
 725 730 735

Leu Ala Thr Trp Thr Ile Gln Gly Ala Ala Asn Ala Leu Ser Gly Asp
 740 745 750

Val Trp Asp Ile Asp Asn Glu Phe
 755 760

<210> SEQ ID NO 3
 <211> LENGTH: 760
 <212> TYPE: PRT
 <213> ORGANISM: Macaca fascicularis

<400> SEQUENCE: 3

Met Met Asp Gln Ala Arg Ser Ala Phe Ser Asn Leu Phe Gly Gly Glu
 1 5 10 15

Pro Leu Ser Tyr Thr Arg Phe Ser Leu Ala Arg Gln Val Asp Gly Asp
 20 25 30

Asn Ser His Val Glu Met Lys Leu Gly Val Asp Glu Glu Glu Asn Thr
 35 40 45

Asp Asn Asn Thr Lys Ala Asn Gly Thr Lys Pro Lys Arg Cys Gly Gly
 50 55 60

Asn Ile Cys Tyr Gly Thr Ile Ala Val Ile Ile Phe Phe Leu Ile Gly
 65 70 75 80

Phe Met Ile Gly Tyr Leu Gly Tyr Cys Lys Gly Val Glu Pro Lys Thr
 85 90 95

Glu Cys Glu Arg Leu Ala Gly Thr Glu Ser Pro Ala Arg Glu Glu Pro
 100 105 110

Glu Glu Asp Phe Pro Ala Ala Pro Arg Leu Tyr Trp Asp Asp Leu Lys
 115 120 125

Arg Lys Leu Ser Glu Lys Leu Asp Thr Thr Asp Phe Thr Ser Thr Ile
 130 135 140

Lys Leu Leu Asn Glu Asn Leu Tyr Val Pro Arg Glu Ala Gly Ser Gln
 145 150 155 160

Lys Asp Glu Asn Leu Ala Leu Tyr Ile Glu Asn Gln Phe Arg Glu Phe
 165 170 175

Lys Leu Ser Lys Val Trp Arg Asp Gln His Phe Val Lys Ile Gln Val
 180 185 190

Lys Asp Ser Ala Gln Asn Ser Val Ile Ile Val Asp Lys Asn Gly Gly
 195 200 205

Leu Val Tyr Leu Val Glu Asn Pro Gly Gly Tyr Val Ala Tyr Ser Lys
 210 215 220

Ala Ala Thr Val Thr Gly Lys Leu Val His Ala Asn Phe Gly Thr Lys
 225 230 235 240

Lys Asp Phe Glu Asp Leu Asp Ser Pro Val Asn Gly Ser Ile Val Ile
 245 250 255

Val Arg Ala Gly Lys Ile Thr Phe Ala Glu Lys Val Ala Asn Ala Glu
 260 265 270

Ser Leu Asn Ala Ile Gly Val Leu Ile Tyr Met Asp Gln Thr Lys Phe
 275 280 285

Pro Ile Val Lys Ala Asp Leu Ser Phe Phe Gly His Ala His Leu Gly
 290 295 300

Thr Gly Asp Pro Tyr Thr Pro Gly Phe Pro Ser Phe Asn His Thr Gln
 305 310 315 320

-continued

Phe Pro Pro Ser Gln Ser Ser Gly Leu Pro Asn Ile Pro Val Gln Thr
325 330 335
Ile Ser Arg Ala Ala Ala Glu Lys Leu Phe Gly Asn Met Glu Gly Asp
340 345 350
Cys Pro Ser Asp Trp Lys Thr Asp Ser Thr Cys Lys Met Val Thr Ser
355 360 365
Glu Asn Lys Ser Val Lys Leu Thr Val Ser Asn Val Leu Lys Glu Thr
370 375 380
Lys Ile Leu Asn Ile Phe Gly Val Ile Lys Gly Phe Val Glu Pro Asp
385 390 395 400
His Tyr Val Val Val Gly Ala Gln Arg Asp Ala Trp Gly Pro Gly Ala
405 410 415
Ala Lys Ser Ser Val Gly Thr Ala Leu Leu Leu Lys Leu Ala Gln Met
420 425 430
Phe Ser Asp Met Val Leu Lys Asp Gly Phe Gln Pro Ser Arg Ser Ile
435 440 445
Ile Phe Ala Ser Trp Ser Ala Gly Asp Phe Gly Ser Val Gly Ala Thr
450 455 460
Glu Trp Leu Glu Gly Tyr Leu Ser Ser Leu His Leu Lys Ala Phe Thr
465 470 475 480
Tyr Ile Asn Leu Asp Lys Ala Val Leu Gly Thr Ser Asn Phe Lys Val
485 490 495
Ser Ala Ser Pro Leu Leu Tyr Thr Leu Ile Glu Lys Thr Met Gln Asp
500 505 510
Val Lys His Pro Val Thr Gly Arg Ser Leu Tyr Gln Asp Ser Asn Trp
515 520 525
Ala Ser Lys Val Glu Lys Leu Thr Leu Asp Asn Ala Ala Phe Pro Phe
530 535 540
Leu Ala Tyr Ser Gly Ile Pro Ala Val Ser Phe Cys Phe Cys Glu Asp
545 550 555 560
Thr Asp Tyr Pro Tyr Leu Gly Thr Thr Met Asp Thr Tyr Lys Glu Leu
565 570 575
Val Glu Arg Ile Pro Glu Leu Asn Lys Val Ala Arg Ala Ala Ala Glu
580 585 590
Val Ala Gly Gln Phe Val Ile Lys Leu Thr His Asp Thr Glu Leu Asn
595 600 605
Leu Asp Tyr Glu Arg Tyr Asn Ser Gln Leu Leu Leu Phe Leu Arg Asp
610 615 620
Leu Asn Gln Tyr Arg Ala Asp Val Lys Glu Met Gly Leu Ser Leu Gln
625 630 635 640
Trp Leu Tyr Ser Ala Arg Gly Asp Phe Phe Arg Ala Thr Ser Arg Leu
645 650 655
Thr Thr Asp Phe Arg Asn Ala Glu Lys Arg Asp Lys Phe Val Met Lys
660 665 670
Lys Leu Asn Asp Arg Val Met Arg Val Glu Tyr Tyr Phe Leu Ser Pro
675 680 685
Tyr Val Ser Pro Lys Glu Ser Pro Phe Arg His Val Phe Trp Gly Ser
690 695 700
Gly Ser His Thr Leu Ser Ala Leu Leu Glu Ser Leu Lys Leu Arg Arg
705 710 715 720

-continued

Gln Asn Asn Ser Ala Phe Asn Glu Thr Leu Phe Arg Asn Gln Leu Ala
725 730 735

Leu Ala Thr Trp Thr Ile Gln Gly Ala Ala Asn Ala Leu Ser Gly Asp
740 745 750

Val Trp Asp Ile Asp Asn Glu Phe
755 760

<210> SEQ ID NO 4
<211> LENGTH: 763
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 4

Met Met Asp Gln Ala Arg Ser Ala Phe Ser Asn Leu Phe Gly Gly Glu
1 5 10 15

Pro Leu Ser Tyr Thr Arg Phe Ser Leu Ala Arg Gln Val Asp Gly Asp
20 25 30

Asn Ser His Val Glu Met Lys Leu Ala Ala Asp Glu Glu Glu Asn Ala
35 40 45

Asp Asn Asn Met Lys Ala Ser Val Arg Lys Pro Lys Arg Phe Asn Gly
50 55 60

Arg Leu Cys Phe Ala Ala Ile Ala Leu Val Ile Phe Phe Leu Ile Gly
65 70 75 80

Phe Met Ser Gly Tyr Leu Gly Tyr Cys Lys Arg Val Glu Gln Lys Glu
85 90 95

Glu Cys Val Lys Leu Ala Glu Thr Glu Glu Thr Asp Lys Ser Glu Thr
100 105 110

Met Glu Thr Glu Asp Val Pro Thr Ser Ser Arg Leu Tyr Trp Ala Asp
115 120 125

Leu Lys Thr Leu Leu Ser Glu Lys Leu Asn Ser Ile Glu Phe Ala Asp
130 135 140

Thr Ile Lys Gln Leu Ser Gln Asn Thr Tyr Thr Pro Arg Glu Ala Gly
145 150 155 160

Ser Gln Lys Asp Glu Ser Leu Ala Tyr Tyr Ile Glu Asn Gln Phe His
165 170 175

Glu Phe Lys Phe Ser Lys Val Trp Arg Asp Glu His Tyr Val Lys Ile
180 185 190

Gln Val Lys Ser Ser Ile Gly Gln Asn Met Val Thr Ile Val Gln Ser
195 200 205

Asn Gly Asn Leu Asp Pro Val Glu Ser Pro Glu Gly Tyr Val Ala Phe
210 215 220

Ser Lys Pro Thr Glu Val Ser Gly Lys Leu Val His Ala Asn Phe Gly
225 230 235 240

Thr Lys Lys Asp Phe Glu Glu Leu Ser Tyr Ser Val Asn Gly Ser Leu
245 250 255

Val Ile Val Arg Ala Gly Glu Ile Thr Phe Ala Glu Lys Val Ala Asn
260 265 270

Ala Gln Ser Phe Asn Ala Ile Gly Val Leu Ile Tyr Met Asp Lys Asn
275 280 285

Lys Phe Pro Val Val Glu Ala Asp Leu Ala Leu Phe Gly His Ala His
290 295 300

Leu Gly Thr Gly Asp Pro Tyr Thr Pro Gly Phe Pro Ser Phe Asn His
305 310 315 320

-continued

Leu Arg Gln Lys Asn Ile Thr Ala Phe Asn Glu Thr Leu Phe Arg Asn
725 730 735

Gln Leu Ala Leu Ala Thr Trp Thr Ile Gln Gly Val Ala Asn Ala Leu
740 745 750

Ser Gly Asp Ile Trp Asn Ile Asp Asn Glu Phe
755 760

<210> SEQ ID NO 5
<211> LENGTH: 197
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 5

Phe Val Lys Ile Gln Val Lys Asp Ser Ala Gln Asn Ser Val Ile Ile
1 5 10 15

Val Asp Lys Asn Gly Arg Leu Val Tyr Leu Val Glu Asn Pro Gly Gly
20 25 30

Tyr Val Ala Tyr Ser Lys Ala Ala Thr Val Thr Gly Lys Leu Val His
35 40 45

Ala Asn Phe Gly Thr Lys Lys Asp Phe Glu Asp Leu Tyr Thr Pro Val
50 55 60

Asn Gly Ser Ile Val Ile Val Arg Ala Gly Lys Ile Thr Phe Ala Glu
65 70 75 80

Lys Val Ala Asn Ala Glu Ser Leu Asn Ala Ile Gly Val Leu Ile Tyr
85 90 95

Met Asp Gln Thr Lys Phe Pro Ile Val Asn Ala Glu Leu Ser Phe Phe
100 105 110

Gly His Ala His Leu Gly Thr Gly Asp Pro Tyr Thr Pro Gly Phe Pro
115 120 125

Ser Phe Asn His Thr Gln Phe Pro Pro Ser Arg Ser Ser Gly Leu Pro
130 135 140

Asn Ile Pro Val Gln Thr Ile Ser Arg Ala Ala Ala Glu Lys Leu Phe
145 150 155 160

Gly Asn Met Glu Gly Asp Cys Pro Ser Asp Trp Lys Thr Asp Ser Thr
165 170 175

Cys Arg Met Val Thr Ser Glu Ser Lys Asn Val Lys Leu Thr Val Ser
180 185 190

Asn Val Leu Lys Glu
195

<210> SEQ ID NO 6
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 6

Ala Ser Ser Leu Asn Ile Ala
1 5

<210> SEQ ID NO 7
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 7

Ser Lys Thr Phe Asn Thr His Pro Gln Ser Thr Pro
1 5 10

<210> SEQ ID NO 8

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 8

Thr Ala Arg Gly Glu His Lys Glu Glu Glu Leu Ile
1 5 10

<210> SEQ ID NO 9

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 9

Cys Gln Ala Gln Gly Gln Leu Val Cys
1 5

<210> SEQ ID NO 10

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 10

Cys Ser Glu Arg Ser Met Asn Phe Cys
1 5

<210> SEQ ID NO 11

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 11

Cys Pro Lys Thr Arg Arg Val Pro Cys
1 5

<210> SEQ ID NO 12

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 12

Trp Leu Ser Glu Ala Gly Pro Val Val Thr Val Arg Ala Leu Arg Gly
1 5 10 15

Thr Gly Ser Trp
20

-continued

<210> SEQ ID NO 13
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 13

Cys Met Gln His Ser Met Arg Val Cys
1 5

<210> SEQ ID NO 14
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 14

Asp Asp Thr Arg His Trp Gly
1 5

<210> SEQ ID NO 15
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 15

cuuccuggau ggcuucaau 19

<210> SEQ ID NO 16
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 16

guacauaaag auggacuuc 19

<210> SEQ ID NO 17
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 17

uaucuggaua ggugguauca agaucuguaa 30

<210> SEQ ID NO 18
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 18

auguaacuga aaaguucuu cuuua 25

<210> SEQ ID NO 19

-continued

<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 19

uggauaggug guaucaacau cuguagcac 30

<210> SEQ ID NO 20
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 20

gauagguggu aucaacau cu 22

<210> SEQ ID NO 21
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 21

uaucuggaua ggugguauca acaucuguaa 30

<210> SEQ ID NO 22
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 22

aaacuuggaa gagugaugug augua 25

<210> SEQ ID NO 23
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 23

gcucacuugu ugaggcaaaa cuuggaa 27

<210> SEQ ID NO 24
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 24

gccuuggcaa cauuuccacu uccug 25

<210> SEQ ID NO 25
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 25

uacacacuuu accuguugag aauag 25

<210> SEQ ID NO 26

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 26

gauagguggu aucaacaucu guaa 24

<210> SEQ ID NO 27

<211> LENGTH: 21

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 27

gauagguggu aucaacaucu g 21

<210> SEQ ID NO 28

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 28

gauagguggu aucaacaucu gaaag 25

<210> SEQ ID NO 29

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 29

ggugguauca acaucuguaa 20

<210> SEQ ID NO 30

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 30

guaucaacau cuguaagcac 20

<210> SEQ ID NO 31

<211> LENGTH: 30

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (27)..(27)

-continued

<223> OTHER INFORMATION: n can be any nucleouide

<400> SEQUENCE: 31

cggcuaauuu cagagggcgc uuucuungac 30

<210> SEQ ID NO 32

<211> LENGTH: 26

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 32

acaguggcgc ugagauagua uaggcc 26

<210> SEQ ID NO 33

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 33

uaggccacuu uguugcucu gc 22

<210> SEQ ID NO 34

<211> LENGTH: 19

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 34

uucagagggc gcuuucuuc 19

<210> SEQ ID NO 35

<211> LENGTH: 30

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 35

ucuucaggug caccuucugu uucucaauuc 30

<210> SEQ ID NO 36

<211> LENGTH: 30

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 36

ucugugauac ucuucaggug caccuucugu 30

<210> SEQ ID NO 37

<211> LENGTH: 21

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 37

-continued

ucuucugcuc gggaggugac a 21

<210> SEQ ID NO 38
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 38

ccaguuacua uucagaagac 20

<210> SEQ ID NO 39
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 39

ucuucaggug caccuucugu 20

<210> SEQ ID NO 40
<211> LENGTH: 18
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 40

ugcugcuguc uucuugcu 18

<210> SEQ ID NO 41
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 41

uuguuaacuu uuucccauu 19

<210> SEQ ID NO 42
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 42

uguaaacuuu uucccauugg 20

<210> SEQ ID NO 43
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 43

cauuuuguua acuuuuuccc 20

<210> SEQ ID NO 44

-continued

<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 44

cuguagcuuc acccuuucc 19

<210> SEQ ID NO 45
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 45

gagagcuucc uguagcuuca cccuuu 26

<210> SEQ ID NO 46
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 46

uccuguagcu ucacccuuuc cacaggcg 28

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 47

uguguuaccu acccuugucg 20

<210> SEQ ID NO 48
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 48

uagacuaucu uuuaauuucu guaaau 27

<210> SEQ ID NO 49
<211> LENGTH: 29
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 49

gagagcuucc uguagcuuca cccuuucca 29

<210> SEQ ID NO 50
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 50

uuccuguagc uacacccuuu ccacagcgcu u 31

<210> SEQ ID NO 51
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 51

agcuuccugu agcuacccc uuu 23

<210> SEQ ID NO 52
<211> LENGTH: 29
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 52

ggagagagcu uccuguagcu ucacccuuu 29

<210> SEQ ID NO 53
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 53

gagagcuucc uguagcuca ccc 23

<210> SEQ ID NO 54
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 54

uauguguuac cuacccuugu cgguc 25

<210> SEQ ID NO 55
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 55

ggagagagcu uccuguagcu 20

<210> SEQ ID NO 56
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 56

-continued

ucacccuuuc cacaggcggu gca 23

<210> SEQ ID NO 57
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 57

gcugggagag agcuuccugu agcuucac 28

<210> SEQ ID NO 58
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 58

uguuaccuac ccuugucggu ccuuguac 28

<210> SEQ ID NO 59
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 59

cugcugucuu cuugcauga auaauguc 28

<210> SEQ ID NO 60
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 60

ggcguugcac uuugcaaugc ugcugucu 28

<210> SEQ ID NO 61
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 61

uuggaaauca agcugggaga gagcuucc 28

<210> SEQ ID NO 62
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 62

cuacccuugu cgguccuugu acauuuug 28

<210> SEQ ID NO 63

-continued

<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 63

gucaauccga ccugagcuuu guuguaga 28

<210> SEQ ID NO 64
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 64

cuugcuaua auaaugucaa uccgacc 27

<210> SEQ ID NO 65
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 65

uauauguguu accuacccuu gucggucc 28

<210> SEQ ID NO 66
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 66

uuugugucuu ucugagaaac 20

<210> SEQ ID NO 67
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 67

aaagacuuac cuuaagauac 20

<210> SEQ ID NO 68
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 68

aucugucaaaa ucgccugcag 20

<210> SEQ ID NO 69
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 69

cgccgccaau ucucaacag 19

<210> SEQ ID NO 70
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 70

uuuguuuua gcauguucc 20

<210> SEQ ID NO 71
<211> LENGTH: 17
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 71

ccgccauuuc ucaacag 17

<210> SEQ ID NO 72
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 72

uucucaggaa uuugugucu u 21

<210> SEQ ID NO 73
<211> LENGTH: 11
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 73

gacaacucuu u 11

<210> SEQ ID NO 74
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 74

ucagcuucug uuagccacug 20

<210> SEQ ID NO 75
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 75

-continued

uguucagcuu cuguuagcca cuga 24

<210> SEQ ID NO 76
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 76

cuguucagcu ucuguuagcc acugauu 27

<210> SEQ ID NO 77
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 77

uucucaacag aucugucaaaa ucgccugcag 30

<210> SEQ ID NO 78
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 78

gccacugauu aaauaucuuu auauc 25

<210> SEQ ID NO 79
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 79

ucuguuagcc acugauuaaa uaucuuuua 30

<210> SEQ ID NO 80
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 80

gagaaacugu ucagcuucug uuagccacug a 31

<210> SEQ ID NO 81
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 81

ucuuucugag aaacuguuca gcuucuguua g 31

<210> SEQ ID NO 82

-continued

<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 82

cagaucuguc aaaucgccug caggua 26

<210> SEQ ID NO 83
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 83

caacagaucu gucaaaucgc cugcag 26

<210> SEQ ID NO 84
<211> LENGTH: 33
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 84

aaacuguuca gcuucuguaa gccacugauu aaa 33

<210> SEQ ID NO 85
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 85

gaaacuguuc agcuucuguu agccacugau u 31

<210> SEQ ID NO 86
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 86

aaacuguuca gcuucuguaa gccacuga 28

<210> SEQ ID NO 87
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 87

ugagaaacug uucagcuucu guuagcca 28

<210> SEQ ID NO 88
<211> LENGTH: 32
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid
<400> SEQUENCE: 88
uucugagaaa cuguucagcu ucuguuagcc ac 32

<210> SEQ ID NO 89
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid
<400> SEQUENCE: 89
uucugagaaa cuguucagcu ucuguu 26

<210> SEQ ID NO 90
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid
<400> SEQUENCE: 90
gaucugucuaa aucgccugca gguaa 25

<210> SEQ ID NO 91
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid
<400> SEQUENCE: 91
auaaugaaaa cgccgcauu ucua 25

<210> SEQ ID NO 92
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid
<400> SEQUENCE: 92
aaacuguaa gcuucuguaa gccac 25

<210> SEQ ID NO 93
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid
<400> SEQUENCE: 93
uugugucuuu cugagaaacu guca 25

<210> SEQ ID NO 94
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid
<400> SEQUENCE: 94

-continued

ccaauucuca ggaauuugug ucuuu 25

<210> SEQ ID NO 95
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 95

aucgccugca gguaaaagca uaugg 25

<210> SEQ ID NO 96
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 96

ugaaaacgcc gccauuucuc aacagaucug 30

<210> SEQ ID NO 97
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 97

cauaaugaaa acgccgccau uucucaacag 30

<210> SEQ ID NO 98
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 98

uguucagcuu cuguuagcca cugauuaau 30

<210> SEQ ID NO 99
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 99

cagaucuguc aaaucgccug cagg 24

<210> SEQ ID NO 100
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 100

caacagaucu gucaaaucgc cugcagg 27

<210> SEQ ID NO 101

-continued

<211> LENGTH: 29
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 101

cucaacagau cugucaaauc gccugcagg 29

<210> SEQ ID NO 102
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 102

gaucugucuaa aucgccugca ggu 23

<210> SEQ ID NO 103
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 103

gaucugucuaa aucgccugca gg 22

<210> SEQ ID NO 104
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 104

gaucugucuaa aucgccugca g 21

<210> SEQ ID NO 105
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 105

cagaucuguc aaaucgccug caggu 25

<210> SEQ ID NO 106
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 106

cagaucuguc aaaucgccug cag 23

<210> SEQ ID NO 107
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 107

gugucuuucu gagaacugu ucagc

25

<210> SEQ ID NO 108

<211> LENGTH: 28

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 108

gagaacugu ucagcuucug uuagccac

28

<210> SEQ ID NO 109

<211> LENGTH: 28

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 109

gaaacuguuc agcuucuguu agccacug

28

<210> SEQ ID NO 110

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 110

cuguucagcu ucuguuagcc acug

24

<210> SEQ ID NO 111

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 111

aucugucaaa ucgccugcag guaaaag

27

<210> SEQ ID NO 112

<211> LENGTH: 29

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 112

gaucugucaa aucgccugca gguaaaagc

29

<210> SEQ ID NO 113

<211> LENGTH: 19

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 113

-continued

gcugaauuau uucuucucc 19

<210> SEQ ID NO 114
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 114

uuuuucuguc ugacagcug 19

<210> SEQ ID NO 115
<211> LENGTH: 18
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 115

ucuguuuuug aggauugc 18

<210> SEQ ID NO 116
<211> LENGTH: 17
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 116

ccaccgcaga uucaggc 17

<210> SEQ ID NO 117
<211> LENGTH: 17
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 117

gcccraagcc auccugg 17

<210> SEQ ID NO 118
<211> LENGTH: 17
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 118

uuugcagacc uccugcc 17

<210> SEQ ID NO 119
<211> LENGTH: 17
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 119

caguugccg cugccca 17

<210> SEQ ID NO 120

-continued

<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 120

guugcauca auguucugac 20

<210> SEQ ID NO 121
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 121

auuuuuccug uagaauacug g 21

<210> SEQ ID NO 122
<211> LENGTH: 34
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 122

gcugcccaau gcgauccugg aguuccugua agau 34

<210> SEQ ID NO 123
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 123

gcugcccaau gccauccugg aguuccug 28

<210> SEQ ID NO 124
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 124

gcugcccaau gccauccugg aguuccugua a 31

<210> SEQ ID NO 125
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 125

caaugccauc cuggaguucc uguaagauac c 31

<210> SEQ ID NO 126
<211> LENGTH: 32
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 126

gcugcccaau gccauccugg aguuccugua ag 32

<210> SEQ ID NO 127

<211> LENGTH: 30

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 127

ccaugccau ccuggaguuc cuguaagaua 30

<210> SEQ ID NO 128

<211> LENGTH: 39

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 128

uugccgcugc ccaaugccau ccuggaguuc cuguaagau 39

<210> SEQ ID NO 129

<211> LENGTH: 34

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 129

gcugcccaau gccauccugg aguuccugua agau 34

<210> SEQ ID NO 130

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 130

caaugccauc cuggaguucc uguaaga 27

<210> SEQ ID NO 131

<211> LENGTH: 26

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 131

caguugccg cugcccaaug ccaucc 26

<210> SEQ ID NO 132

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 132

-continued

cuuccccagu ugcauuccaau guuc 24

<210> SEQ ID NO 133
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 133

cuggcaucug uuuuugagga uug 23

<210> SEQ ID NO 134
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 134

uuagaucugu cgcccuaccu 20

<210> SEQ ID NO 135
<211> LENGTH: 39
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 135

gcugcccaau gccauccugg aguuccugua agauaccaa 39

<210> SEQ ID NO 136
<211> LENGTH: 34
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 136

gcccaaugcc auccuggagu uccuguaaga uacc 34

<210> SEQ ID NO 137
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 137

cauccuggag uuccuguaag auacc 25

<210> SEQ ID NO 138
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 138

ugccauccug gaguuccugu aagauacc 28

<210> SEQ ID NO 139

-continued

<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 139

ugccauccug gaguuccugu aagau 25

<210> SEQ ID NO 140
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 140

caaugccauc cuggaguucc uguagau 28

<210> SEQ ID NO 141
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 141

gcccaaugcc auccuggagu uccuguaaga u 31

<210> SEQ ID NO 142
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 142

gcccaaugcc auccuggagu uccuguaa 28

<210> SEQ ID NO 143
<211> LENGTH: 34
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 143

gccgcugccc aaugacauc uggaguuccu guaa 34

<210> SEQ ID NO 144
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 144

gcccauccugg aguuccugua agaua 25

<210> SEQ ID NO 145
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 145

ccaaugccau ccuggaguuc cugua

25

<210> SEQ ID NO 146

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 146

cugacaacag uuugccgcug cccaa

25

<210> SEQ ID NO 147

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 147

uuugaggauu gcugaauuuu uucuu

25

<210> SEQ ID NO 148

<211> LENGTH: 30

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 148

caguuugccg cugcccaaug ccauccugga

30

<210> SEQ ID NO 149

<211> LENGTH: 30

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 149

uuugccgcug ccaaugccau ccuggaguuc

30

<210> SEQ ID NO 150

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 150

uuugccgcug cccaugcca uccug

25

<210> SEQ ID NO 151

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 151

-continued

ccaauGCCAU ccuggaguuc cu	22
<210> SEQ ID NO 152	
<211> LENGTH: 29	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 152	
cccaaugCCA uccuggaguu ccguaaga	29
<210> SEQ ID NO 153	
<211> LENGTH: 28	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 153	
ccgcugCCA augccauccu ggaguucc	28
<210> SEQ ID NO 154	
<211> LENGTH: 30	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 154	
cccaaugCCA uccuggaguu ccguaagau	30
<210> SEQ ID NO 155	
<211> LENGTH: 30	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 155	
ccgcugCCA augccauccu ggaguuccug	30
<210> SEQ ID NO 156	
<211> LENGTH: 30	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 156	
ugcccaaugc cauccuggag uuccguaag	30
<210> SEQ ID NO 157	
<211> LENGTH: 28	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 157	
cccaaugCCA uccuggaguu ccguaag	28
<210> SEQ ID NO 158	

-continued

<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 158

ugcccaaugc cauccuggag uuccugua 28

<210> SEQ ID NO 159
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 159

caaugccauc cuggaguucc ug 22

<210> SEQ ID NO 160
<211> LENGTH: 29
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 160

agccucucgc ucacucaccc ugcaaagga 29

<210> SEQ ID NO 161
<211> LENGTH: 29
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 161

ccacucagag cucagauuu cuaacuucc 29

<210> SEQ ID NO 162
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 162

cuuccacuca gagcucagau cuucuaa 27

<210> SEQ ID NO 163
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 163

gggauccagu auacuuacag gcucc 25

<210> SEQ ID NO 164
<211> LENGTH: 17
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 164

cucagagcuc agaucuu 17

<210> SEQ ID NO 165

<211> LENGTH: 15

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 165

ggcugcuuug cccuc 15

<210> SEQ ID NO 166

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 166

cucagaucuu cuaacuuccu cuuaaac 27

<210> SEQ ID NO 167

<211> LENGTH: 29

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 167

cucagagcuc agaucuucua acuuccucu 29

<210> SEQ ID NO 168

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 168

cgccuuccac ucagagcuca gauucuc 27

<210> SEQ ID NO 169

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 169

ucagcucuug aaguaaacgg uuuaccg 27

<210> SEQ ID NO 170

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 170

-continued

uuugcccuca gcucuugaag uaaacgg 27

<210> SEQ ID NO 171
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 171

ggcugcuuug cccucagcuc uugaagu 27

<210> SEQ ID NO 172
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 172

caggagcuag gucaggcugc uuugcc 26

<210> SEQ ID NO 173
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 173

uccaaauagug gucaguccag gagcu 25

<210> SEQ ID NO 174
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 174

aaagagaug ggauccagua uacuuac 27

<210> SEQ ID NO 175
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 175

aaaauagcuag agccaaagag aauggga 27

<210> SEQ ID NO 176
<211> LENGTH: 33
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 176

ggcugcuuug cccucagcuc uugaaguaaa cgg 33

<210> SEQ ID NO 177

-continued

<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 177

aggcugcuuu gccucagcu cuugaagua 30

<210> SEQ ID NO 178
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 178

gucaggcugc uuugccuca gcucuugaag 30

<210> SEQ ID NO 179
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 179

aggucaggcu gcuuugccu cagcucuuga 30

<210> SEQ ID NO 180
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 180

cagagcucag aucuucuaac uuccu 25

<210> SEQ ID NO 181
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 181

cuuacaggcu ccaauagugg ucagu 25

<210> SEQ ID NO 182
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 182

augggaucca guauacuuac aggcu 25

<210> SEQ ID NO 183
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 183

agagaauuggg auccaguaua cuuac 25

<210> SEQ ID NO 184
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 184

aacuuccucu uuaacagaaa agcauac 27

<210> SEQ ID NO 185
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 185

cucauaccuu cugcuugaug auc 23

<210> SEQ ID NO 186
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 186

ucaaggaaga uggcauuucu 20

<210> SEQ ID NO 187
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 187

gaaagccagu cgguaaguuc 20

<210> SEQ ID NO 188
<211> LENGTH: 15
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 188

caccaccau cacc 15

<210> SEQ ID NO 189
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 189

-continued

ccucugugau uuuauaacuu gau	23
<210> SEQ ID NO 190	
<211> LENGTH: 20	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 190	
ugauauccuc aaggucaccc	20
<210> SEQ ID NO 191	
<211> LENGTH: 30	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 191	
gguaccucca acaucaagga agauggcauu	30
<210> SEQ ID NO 192	
<211> LENGTH: 26	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 192	
auuucuaguu uggagauggc aguuuc	26
<210> SEQ ID NO 193	
<211> LENGTH: 26	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 193	
caucaaggaa gauggcauuu cuaguu	26
<210> SEQ ID NO 194	
<211> LENGTH: 26	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 194	
gagcagguac cuccaacauc aaggaa	26
<210> SEQ ID NO 195	
<211> LENGTH: 30	
<212> TYPE: RNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Nucleic acid	
<400> SEQUENCE: 195	
cuccaacauc aaggaagaug gcauuucuag	30
<210> SEQ ID NO 196	

-continued

<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 196

accagaguaa cagucugagu aggag 25

<210> SEQ ID NO 197
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 197

caccagagua acagucugag uagga 25

<210> SEQ ID NO 198
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 198

ucaccagagu aacagucuga guagg 25

<210> SEQ ID NO 199
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 199

gucaccagag uaacagucug aguag 25

<210> SEQ ID NO 200
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 200

accagaguaa cagucugagu aggagc 26

<210> SEQ ID NO 201
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 201

uucuguccaa gcccgguuga aauc 24

<210> SEQ ID NO 202
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 202

acaUCAAGGA agaUGGCAUu ucuAGUuUGG 30

<210> SEQ ID NO 203
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 203

acaUCAAGGA agaUGGCAUu ucuAG 25

<210> SEQ ID NO 204
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 204

aucaUUUUUU cucaUACCUu cUGcu 25

<210> SEQ ID NO 205
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 205

caccCACCAU caccCUCUGu g 21

<210> SEQ ID NO 206
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 206

aucaUCUCGU uGauAUCCUc aa 22

<210> SEQ ID NO 207
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 207

cuccAACAUc aaggaAGAUg gcaUUUcu 28

<210> SEQ ID NO 208
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 208

-continued

caucaaggaa gauggcauuu cuagu 25

<210> SEQ ID NO 209
<211> LENGTH: 18
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 209

uugcuggucu uguuuuuc 18

<210> SEQ ID NO 210
<211> LENGTH: 16
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 210

ccgaaugau uguucu 16

<210> SEQ ID NO 211
<211> LENGTH: 18
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 211

gcuggucuug uuuuucaa 18

<210> SEQ ID NO 212
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 212

uggucuuguu uucaaaaa 20

<210> SEQ ID NO 213
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 213

gucuuguuuu ucaaaaauug 20

<210> SEQ ID NO 214
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 214

cuuguuuuuc aaauuuggg 20

<210> SEQ ID NO 215

-continued

<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 215

uguuuuucaa auuuugggc 19

<210> SEQ ID NO 216
<211> LENGTH: 33
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 216

uccaacuggg gacgccucug uucaaaaucc ugc 33

<210> SEQ ID NO 217
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 217

uccugcauug uugccguaa g 21

<210> SEQ ID NO 218
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 218

uccaacuggg gacgccucug uucaaaaucc 30

<210> SEQ ID NO 219
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 219

acuggggacg ccucuguucc a 21

<210> SEQ ID NO 220
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 220

ccgaaugau uguucuagcc 20

<210> SEQ ID NO 221
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 221

uguuaaaaaa cuuacuucga

20

<210> SEQ ID NO 222

<211> LENGTH: 18

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 222

cuguugccuc egguucug

18

<210> SEQ ID NO 223

<211> LENGTH: 18

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 223

uuggcucugg ccuguccu

18

<210> SEQ ID NO 224

<211> LENGTH: 33

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 224

uucaacuguu gccuccgguu cugaaggugu ucu

33

<210> SEQ ID NO 225

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 225

uacuucaucc cacugauucu gaauu

25

<210> SEQ ID NO 226

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 226

cugaaggugu ucuuguacuu caucc

25

<210> SEQ ID NO 227

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 227

-continued

cuguugccuc cgguucugaa ggugu 25

<210> SEQ ID NO 228
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 228

cuguugccuc cgguucugaa gguguucuu 30

<210> SEQ ID NO 229
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 229

caacuguugc cuccgguucu gaagguguuc 30

<210> SEQ ID NO 230
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 230

uugccuccgg uucugaaggu guucuuguac 30

<210> SEQ ID NO 231
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 231

guugccuccg guucugaagg uguuc 25

<210> SEQ ID NO 232
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 232

cuccgguucu gaagguguuc uug 23

<210> SEQ ID NO 233
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 233

cuccgguucu gaagguguuc uu 22

<210> SEQ ID NO 234

-continued

<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 234

cuccgguucu gaagguguuc u 21

<210> SEQ ID NO 235
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 235

cuccgguucu gaagguguuc 20

<210> SEQ ID NO 236
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 236

cuccgguucu gaagguguu 19

<210> SEQ ID NO 237
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 237

caucaacug uugccuccgg uucug 25

<210> SEQ ID NO 238
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 238

cuguugccuc cgguucugaa ggug 24

<210> SEQ ID NO 239
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 239

caucaacug uugccuccgg uucugaaggu g 31

<210> SEQ ID NO 240
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 240

uacuaaccuu gguuucugug a 21

<210> SEQ ID NO 241

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 241

uguauaggga ccuccuucc augacuc 27

<210> SEQ ID NO 242

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 242

cuaaccuugg uuucugugau uuucu 25

<210> SEQ ID NO 243

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 243

gguaucuuug auacuaaccu ugguuuc 27

<210> SEQ ID NO 244

<211> LENGTH: 22

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 244

auucuuuca cuagaauaaa ag 22

<210> SEQ ID NO 245

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 245

gauucugaau ucuucaacu agaau 25

<210> SEQ ID NO 246

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 246

-continued

aucccacuga uucugaauc 20

<210> SEQ ID NO 247
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 247

aaccgagacc ggacaggauu cu 22

<210> SEQ ID NO 248
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 248

cuguugcagu aaucuaugag 20

<210> SEQ ID NO 249
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 249

ugccauuguu ucaucagcuc uuu 23

<210> SEQ ID NO 250
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 250

ugcaguaauc uaugaguuuuc 20

<210> SEQ ID NO 251
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 251

uccuguagga cauuggcagu 20

<210> SEQ ID NO 252
<211> LENGTH: 18
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 252

gagucuucua ggagccuu 18

<210> SEQ ID NO 253

-continued

<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 253

ggccaaaccu cggcuuaccu gaaau 25

<210> SEQ ID NO 254
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 254

ggccaaaccu cggcuuaccu 20

<210> SEQ ID NO 255
<211> LENGTH: 29
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 255

aaucagcugg gagagagcuu ccuguagcu 29

<210> SEQ ID NO 256
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 256

ucguucuucu gucgucguaa cguuuc 26

<210> SEQ ID NO 257
<211> LENGTH: 16
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 257

caccgauugu cuucga 16

<210> SEQ ID NO 258
<211> LENGTH: 17
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 258

cccuuguacg auuuuug 17

<210> SEQ ID NO 259
<211> LENGTH: 18
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 259

ucuguguuuu aggacucu 18

<210> SEQ ID NO 260

<211> LENGTH: 31

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 260

gccgcugccc aaugccauc uggaguuccu g 31

<210> SEQ ID NO 261

<211> LENGTH: 29

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 261

auuagaucug ucgccuacc ucuuuuuuc 29

<210> SEQ ID NO 262

<211> LENGTH: 27

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 262

ugucgccua ccucuuuuu cugucug 27

<210> SEQ ID NO 263

<211> LENGTH: 25

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 263

gcccraagcc auccuggagu uccug 25

<210> SEQ ID NO 264

<211> LENGTH: 30

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 264

gagccucug cucacucacc cugcaaagga 30

<210> SEQ ID NO 265

<211> LENGTH: 36

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 265

-continued

aucauuuuuuu cucauaccuu cugcuaggag cuaaaa 36

<210> SEQ ID NO 266
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleic acid

<400> SEQUENCE: 266

ggaagcuaag gaagaagcug agcagg 26

<210> SEQ ID NO 267
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 267

Ser Tyr Trp Met His
1 5

<210> SEQ ID NO 268
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 268

Glu Ile Asn Pro Thr Asn Gly Arg Thr Asn Tyr Ile Glu Lys Phe Lys
1 5 10 15

Ser

<210> SEQ ID NO 269
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 269

Gly Thr Arg Ala Tyr His Tyr
1 5

<210> SEQ ID NO 270
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 270

Arg Ala Ser Asp Asn Leu Tyr Ser Asn Leu Ala
1 5 10

<210> SEQ ID NO 271
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

-continued

<400> SEQUENCE: 271

Asp Ala Thr Asn Leu Ala Asp
1 5

<210> SEQ ID NO 272

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 272

Gln His Phe Trp Gly Thr Pro Leu Thr
1 5

<210> SEQ ID NO 273

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 273

Gly Tyr Thr Phe Thr Ser Tyr
1 5

<210> SEQ ID NO 274

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 274

Asn Pro Thr Asn Gly Arg
1 5

<210> SEQ ID NO 275

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 275

Thr Ser Tyr Trp Met His
1 5

<210> SEQ ID NO 276

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 276

Trp Ile Gly Glu Ile Asn Pro Thr Asn Gly Arg Thr Asn
1 5 10

<210> SEQ ID NO 277

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 277

Ala Arg Gly Thr Arg Ala
1 5

<210> SEQ ID NO 278

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 278

Tyr Ser Asn Leu Ala Trp Tyr
1 5

<210> SEQ ID NO 279

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 279

Leu Leu Val Tyr Asp Ala Thr Asn Leu Ala
1 5 10

<210> SEQ ID NO 280

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 280

Gln His Phe Trp Gly Thr Pro Leu
1 5

<210> SEQ ID NO 281

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 281

Gln His Phe Ala Gly Thr Pro Leu Thr
1 5

<210> SEQ ID NO 282

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 282

Gln His Phe Ala Gly Thr Pro Leu
1 5

<210> SEQ ID NO 283

<211> LENGTH: 116

<212> TYPE: PRT

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 283

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30
 Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Glu Ile Asn Pro Thr Asn Gly Arg Thr Asn Tyr Ile Glu Lys Phe
 50 55 60
 Lys Ser Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Gly Thr Arg Ala Tyr His Tyr Trp Gly Gln Gly Thr Ser Val
 100 105 110
 Thr Val Ser Ser
 115

<210> SEQ ID NO 284
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 284

Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Val Ser Val Gly
 1 5 10 15
 Glu Thr Val Thr Ile Thr Cys Arg Ala Ser Asp Asn Leu Tyr Ser Asn
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val
 35 40 45
 Tyr Asp Ala Thr Asn Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Ser
 65 70 75 80
 Glu Asp Phe Gly Thr Tyr Tyr Cys Gln His Phe Trp Gly Thr Pro Leu
 85 90 95
 Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 100 105

<210> SEQ ID NO 285
 <211> LENGTH: 116
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 285

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30

-continued

Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
 35 40 45
 Gly Glu Ile Asn Pro Thr Asn Gly Arg Thr Asn Tyr Ile Glu Lys Phe
 50 55 60
 Lys Ser Arg Ala Thr Leu Thr Val Asp Lys Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Gly Thr Arg Ala Tyr His Tyr Trp Gly Gln Gly Thr Met Val
 100 105 110
 Thr Val Ser Ser
 115

<210> SEQ ID NO 286
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 286

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Asp Asn Leu Tyr Ser Asn
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ser Pro Lys Leu Leu Val
 35 40 45
 Tyr Asp Ala Thr Asn Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Tyr Ser Leu Lys Ile Asn Ser Leu Gln Ser
 65 70 75 80
 Glu Asp Phe Gly Thr Tyr Tyr Cys Gln His Phe Trp Gly Thr Pro Leu
 85 90 95
 Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 100 105

<210> SEQ ID NO 287
 <211> LENGTH: 330
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 287

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80
 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys

-continued

85	90	95
Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys 100 105 110		
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro 115 120 125		
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys 130 135 140		
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp 145 150 155 160		
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu 165 170 175		
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu 180 185 190		
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn 195 200 205		
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly 210 215 220		
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu 225 230 235 240		
Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr 245 250 255		
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn 260 265 270		
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe 275 280 285		
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn 290 295 300		
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr 305 310 315 320		
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys 325 330		

<210> SEQ ID NO 288
 <211> LENGTH: 110
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 288

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys 1 5 10 15		
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr 20 25 30		
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser 35 40 45		
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser 50 55 60		
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr 65 70 75 80		
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys 85 90 95		
Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro		

-continued

100 105 110

<210> SEQ ID NO 289
 <211> LENGTH: 446
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 289

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30

Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Glu Ile Asn Pro Thr Asn Gly Arg Thr Asn Tyr Ile Glu Lys Phe
 50 55 60

Lys Ser Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Gly Thr Arg Ala Tyr His Tyr Trp Gly Gln Gly Thr Ser Val
 100 105 110

Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala
 115 120 125

Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu
 130 135 140

Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly
 145 150 155 160

Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser
 165 170 175

Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu
 180 185 190

Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr
 195 200 205

Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr
 210 215 220

Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe
 225 230 235 240

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro
 245 250 255

Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val
 260 265 270

Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
 275 280 285

Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val
 290 295 300

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys
 305 310 315 320

Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser
 325 330 335

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro

-continued

```

<210> SEQ ID NO 291
<211> LENGTH: 446
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 291

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1          5          10          15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20          25          30
Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
35          40          45
Gly Glu Ile Asn Pro Thr Asn Gly Arg Thr Asn Tyr Ile Glu Lys Phe
50          55          60
Lys Ser Arg Ala Thr Leu Thr Val Asp Lys Ser Ala Ser Thr Ala Tyr
65          70          75          80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85          90          95
Ala Arg Gly Thr Arg Ala Tyr His Tyr Trp Gly Gln Gly Thr Met Val
100         105         110
Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala
115         120         125
Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu
130         135         140
Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly
145         150         155         160
Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser
165         170         175
Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu
180         185         190
Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr
195         200         205
Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr
210         215         220
Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe
225         230         235         240
Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro
245         250         255
Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val
260         265         270
Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
275         280         285
Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val
290         295         300
Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys
305         310         315         320
Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser
325         330         335
Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro
340         345         350
Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val

```

-continued

```

      355              360              365
Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
  370              375              380

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp
  385              390              395              400

Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp
              405              410              415

Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His
              420              425              430

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
              435              440              445

```

```

<210> SEQ ID NO 292
<211> LENGTH: 217
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

```

```

<400> SEQUENCE: 292

```

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
  1              5              10              15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Asp Asn Leu Tyr Ser Asn
              20              25              30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ser Pro Lys Leu Leu Val
              35              40              45

Tyr Asp Ala Thr Asn Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
  50              55              60

Ser Gly Ser Gly Thr Asp Tyr Ser Leu Lys Ile Asn Ser Leu Gln Ser
  65              70              75              80

Glu Asp Phe Gly Thr Tyr Tyr Cys Gln His Phe Trp Gly Thr Pro Leu
              85              90              95

Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys Ala Ser Thr Lys Gly
              100              105              110

Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly
              115              120              125

Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val
  130              135              140

Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe
  145              150              155              160

Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val
              165              170              175

Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val
              180              185              190

Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys
  195              200              205

Ser Cys Asp Lys Thr His Thr Cys Pro
  210              215

```

```

<210> SEQ ID NO 293
<211> LENGTH: 226
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

```

-continued

<400> SEQUENCE: 293

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30
 Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Glu Ile Asn Pro Thr Asn Gly Arg Thr Asn Tyr Ile Glu Lys Phe
 50 55 60
 Lys Ser Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Gly Thr Arg Ala Tyr His Tyr Trp Gly Gln Gly Thr Ser Val
 100 105 110
 Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala
 115 120 125
 Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu
 130 135 140
 Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly
 145 150 155 160
 Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser
 165 170 175
 Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu
 180 185 190
 Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr
 195 200 205
 Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr
 210 215 220
 Cys Pro
 225

<210> SEQ ID NO 294

<211> LENGTH: 226

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 294

Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30
 Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
 35 40 45
 Gly Glu Ile Asn Pro Thr Asn Gly Arg Thr Asn Tyr Ile Glu Lys Phe
 50 55 60
 Lys Ser Arg Ala Thr Leu Thr Val Asp Lys Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

-continued

Ala	Arg	Gly	Thr	Arg	Ala	Tyr	His	Tyr	Trp	Gly	Gln	Gly	Thr	Met	Val
			100					105					110		
Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala
		115					120					125			
Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu
	130				135						140				
Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly
145					150					155					160
Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser
				165					170					175	
Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu
		180						185					190		
Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr
		195					200					205			
Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr
	210					215					220				
Cys	Pro														
225															

1.-72. (canceled)

73. A complex comprising an anti-transferrin receptor antibody covalently linked to an oligonucleotide that targets a dystrophin (DMD) pre-mRNA in a muscle cell,

wherein the oligonucleotide is 15 to 35 nucleotides in length and comprises a region of complementarity to the DMD pre-mRNA, wherein the region of complementarity is at least 12 nucleotides in length; and

wherein the anti-transferrin receptor antibody binds in the range of C89 to F760 of human transferrin receptor protein 1 (TfR1) having an amino acid sequence as set forth in SEQ ID NO: 1, permits transferrin binding to TfR1, and cross-reacts with human TfR1 and cynomolgus TfR1.

74. The complex of claim **73**, wherein the anti-transferrin receptor antibody is in the form of a ScFv, Fab fragment, Fab' fragment, F(ab')₂ fragment, or Fv fragment.

75. The complex of claim **73**, wherein the anti-transferrin receptor antibody is in the form of a Fab fragment.

76. The complex of claim **73**, wherein the anti-transferrin receptor antibody binds human TfR1 with a K_D of 10^{-11} M to 10^{-6} M.

77. The complex of claim **73**, wherein the anti-transferrin receptor antibody is a humanized antibody.

78. The complex of claim **73**, wherein the oligonucleotide comprises one or more phosphorodiamidate morpholinos.

79. The complex of claim **73**, wherein the oligonucleotide is a phosphorodiamidate morpholino oligomer (PMO).

80. The complex of claim **73**, wherein the oligonucleotide is 16-30 nucleotides in length.

81. The complex of claim **73**, wherein the region of complementarity is at least 16 nucleotides in length.

82. The complex of claim **73**, wherein the oligonucleotide comprises a region of complementarity that is at least 12 nucleotides in length and is complementary to the target sequence of an oligonucleotide listed in Table 2.

83. The complex of claim **73**, wherein the oligonucleotide comprises at least 16 consecutive nucleotides of an oligonucleotide listed in Table 2.

84. The complex of claim **73**, wherein the anti-transferrin receptor antibody is covalently linked to the oligonucleotide via a cleavable linker comprising a valine-citrulline sequence.

85. The complex of claim **73**, wherein the anti-transferrin receptor antibody is covalently linked to the oligonucleotide via conjugation to a lysine residue or a cysteine residue of the anti-transferrin receptor antibody.

86. The complex of claim **73**, wherein the complex is configured to promote transferrin receptor mediated internalization of the oligonucleotide into a muscle cell.

87. The complex of claim **73**, wherein the DMD pre-mRNA comprises one or more frameshift mutations.

88. The complex of claim **87**, wherein the frameshift mutation is in exon 8, exon 23, exon 41, exon 44, exon 50, exon 51, exon 52, exon 53, or exon 55.

89. The complex of claim **86**, wherein internalization of the complex in a muscle cell induces skipping of exon 44, exon 45, exon 51, exon 8, exon 23, exon 50, exon 52, exon 53, or exon 55 of DMD pre-mRNA in the muscle cell.

90. The complex of claim **89**, wherein internalization of the complex in a muscle cell promotes the expression or activity of a functional dystrophin protein.

91. A method of inducing skipping of an exon in a dystrophin (DMD) pre-mRNA in a muscle cell, the method comprising contacting the muscle cell with a complex comprising an anti-transferrin receptor antibody covalently linked to an oligonucleotide that targets the DMD pre-mRNA,

wherein the oligonucleotide is 15 to 35 nucleotides in length and comprises a region of complementarity to the DMD pre-mRNA, wherein the region of complementarity is at least 12 nucleotides in length; and

wherein the anti-transferrin receptor antibody binds in the range of C89 to F760 of human transferrin receptor protein 1 (TfR1) having an amino acid sequence as set forth in SEQ ID NO: 1, permits transferrin binding to TfR1, and cross-reacts with human TfR1 and cynomolgus TfR1.

92. A method of treating Duchenne Muscular Dystrophy (DMD) in a subject expressing a DMD pre-mRNA comprising one or more frameshift mutations, the method comprising administering to the subject a complex comprising an anti-transferrin receptor antibody covalently linked to an oligonucleotide that targets the DMD pre-mRNA in a muscle cell,

wherein the oligonucleotide is 15 to 35 nucleotides in length and comprises a region of complementarity to the DMD pre-mRNA, wherein the region of complementarity is at least 12 nucleotides in length; and

wherein the anti-transferrin receptor antibody binds in the range of C89 to F760 of human transferrin receptor protein 1 (TfR1) having an amino acid sequence as set forth in SEQ ID NO: 1, permits transferrin binding to TfR1, and cross-reacts with human TfR1 and cynomolgus TfR1.

* * * * *