

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
3 July 2008 (03.07.2008)

PCT

(10) International Publication Number
WO 2008/077404 A1

(51) International Patent Classification:

C07D 213/61 (2006.01) *A61P 25/00* (2006.01)
C07D 413/12 (2006.01) *A61P 29/00* (2006.01)
C07D 417/12 (2006.01) *A61P 31/00* (2006.01)
A61K 31/44 (2006.01) *A61P 35/00* (2006.01)
A61K 31/4427 (2006.01)

(DK). **NIELSEN, Simon, Feldbæk** [DK/DK]; Åfaldet 72, DK-2730 Herlev (DK).

(21) International Application Number:

PCT/2007/000564

(74) **Agent:** **DEUSSEN, Heinz-Josef**; Principal Patent Counsel, Leo Pharma A/S, Industriparken 55, DK-2750 Ballerup (DK).

(22) International Filing Date:

21 December 2007 (21.12.2007)

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language:

English

(26) Publication Language:

English

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(30) Priority Data:

60/871,689 22 December 2006 (22.12.2006) US
 60/945,470 21 June 2007 (21.06.2007) US

(71) **Applicant** (for all designated States except US): **LEO PHARMA A/S** [DK/DK]; Industriparken 55, DK-2750 Ballerup (DK).


(72) Inventors; and

(75) **Inventors/Applicants** (for US only): **FELDING, Jakob** [DK/DK]; Ordruphøje 24, 1., DK-2920 Charlottenlund

Published:

— with international search report

(54) Title: SUBSTITUTED ACETOPHENONES USEFUL AS PDE4 INHIBITORS



(57) **Abstract:** The present invention relates to a compound according to formula: (I); wherein X_1 , X_2 , X_3 , X_4 and X_5 independently of each other represent -CH- or N; or X_3 , X_4 and X_5 independently of each other represent -CH- or N, and X_1 and X_2 independently of each other represent C and form part of an additional 6- membered aromatic ring; R_1 represents hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, hydroxylalkyl, or alkylcarbonyl, all of which are optionally substituted; R_2 and R_3 independently represent hydrogen, $-CH_2-C(O)NR'-R'$, alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, hydroxylalkyl, heterocycloalkenyl, alkylaryl, alkylalkoxycarbonyl, alkylcarbonyloxy, or alkoxyalkyl, all of which are optionally substituted; R_{11} represents hydrogen, halogen, cyano, amino, alkoxy or alkylamino, X_1 - X_5 represent -CH- or N, including N-oxides, enantiomers and diastereomers; and pharmaceutically acceptable salts, hyd rates, or solvates thereof. The invention relates further to processes for the preparation of said compounds, to said compounds for use in therapy, to pharmaceutical compositions comprising said compounds, to methods of treating diseases, e.g. dermal diseases, with said compounds, and to the use of said compounds in the manufacture of medicaments.

WO 2008/077404 A1

SUBSTITUTED ACETOPHENONES USEFUL AS PDE4 INHIBITORS

FIELD OF THE INVENTION

This invention relates to novel substituted acetophenones and derivatives thereof, processes for the preparation thereof, to said compounds for use in therapy, to pharmaceutical compositions comprising said compounds, to methods of treating diseases with said compounds, and to the use of said compounds in the manufacture of medicaments.

10 BACKGROUND OF THE INVENTION

Phosphodiesterases are enzymes that catalyse the hydrolysis of cyclic AMP and/or cyclic GMP in cells to 5-AMP and 5-GMP, respectively, and as such they are critical to cellular regulation of cAMP or cGMP levels. Of the 11 phosphodiesterases identified so far, phosphodiesterase (PDE) 4, PDE7 and 15 PDE8 are selective for cAMP. PDE4 is the most important modulator of cAMP expressed in immune and inflammatory cells such as neutrophils, macrophages and T-lymphocytes (Z. Huang and J.A. Mancini, *Current Med. Chem.* 13, 2006, pp. 3253-3262). As cAMP is a key second messenger in the modulation of inflammatory responses, PDE4 has been found to regulate inflammatory 20 responses of inflammatory cells by modulating proinflammatory cytokines such as TNF α , IL-2, IFN- γ , GM-CSF and LTB4. Inhibition of PDE4 has therefore become an attractive target for the therapy of inflammatory diseases such as asthma, chronic obstructive pulmonary disease (COPD), rheumatoid arthritis, atopic dermatitis, Crohn's disease etc. (M.D. Houslay et al., *Drug Discovery 25 Today* 10 (22), 2005, pp. 1503-1519). As atopic dermatitis (AD) patients have increased PDE-activity, PDE4-inhibition would also appear to be a viable treatment of AD (*Journal of Investigative Dermatology* (1986), 87(3), 372-6). The PDE4 gene family consists at least of four genes, A, B, C and D, which have a high degree of homology (V. Boswell Smith and D. Spina, *Curr. Opinion 30 Investig. Drugs* 6(11), 2006, pp. 1136-1141). The four PDE4 isoforms are differentially expressed in different tissues and cell types. Thus, PDE4B is predominantly expressed in monocytes and neutrophils, but not in cortex and epithelial cells, while PDE4D is expressed in lung, cortex, cerebellum and T-cells (C. Kroegel and M. Foerster, *Exp. Opinion Investig. Drugs* 16(1), 2007, pp. 109-

124). It has been speculated that inhibition of PDE4D in the brain is associated with the adverse effects found when administering PDE4 inhibitors clinically, primarily nausea and emesis, whereas inhibition of PDE4B is associated with anti-inflammatory effects (B. Lipworth, *Lancet* 365, 2005, pp. 167-175).

5 However, the PDE inhibitors developed so far are not believed to be specific for any of the four PDE4 isoforms. Numerous PDE4 inhibitors have been studied for their therapeutic effect on inflammatory diseases, primarily asthma and COPD.

The first of these, theophylline, is a weak, non-selective phosphodiesterase 10 inhibitor used in the treatment of respiratory diseases such as asthma and COPD. Treatment with theophylline may, however, give rise to both mild and severe adverse effects, e.g. arrhythmia and convulsions, restricting the clinical utility of theophylline (Kroegel and Foerster, *supra*). As phosphodiesterase has remained an attractive target for anti-inflammatory therapy, several other, more 15 selective PDE4 inhibitors have been developed and investigated in a clinical setting. The clinical development of many of the first-generation PDE4 inhibitors such as rolipram was discontinued due to dose-limiting side effects, primarily nausea and emesis. Second-generation PDE4 inhibitors with apparently less pronounced adverse effects are currently in clinical trials (Houslay, *supra*).

20 PDE-4 inhibitors are for example disclosed in EP 0771794 and EP 0943613. WO95/20578 and WO 96/31476 disclose structurally different 4-substituted-3,5-dichloropyridines which are inhibitors of cyclic AMP phosphodiesterase. There is a continued need for developing novel PDE4 inhibitors which have a 25 more favourable therapeutic window, i.e. fewer adverse effects, while retaining their therapeutic anti-inflammatory effect. An overview of preclinical and clinical trials with selective PDE4 inhibitors, including such inhibitors aimed for the treatment of for psoriasis, was recently given in *Inflammation & Allergy: Drug Targets*, 2007, 6(1), 17-26.

30 SUMMARY OF THE INVENTION


The inventors have surprisingly found that novel compounds of the present invention exhibit selective PDE4 inhibitory activity (having a >10 times reduced activity against PDE 1, 2, 3, 5, 7, 8, 9, 10, or 11), which may be useful as therapeutic agents for inflammatory allergic diseases such as bronchial asthma,

allergic rhinitis, and nephritis; autoimmune diseases such as rheumatoid arthritis, multiple sclerosis, inflammatory Bowel disease, Crohn's disease, and systemic lupus erythematosus; diseases of the central nervous system such as depression, amnesia, and dementia; organopathy associated with ischemic 5 reflux caused by cardiac failure, shock, and cerebrovascular diseases, and the like; insulin-resistant diabetes; wounds; AIDS, and the like.

Compounds of the present invention may also be beneficial in preventing, 10 treating or ameliorating a variety of diseases, such as dermal diseases or conditions, such as proliferative and inflammatory skin disorders and in particular psoriasis, epidermal inflammation, alopecia, skin atrophy, steroid induced skin atrophy, skin ageing, photo skin ageing, acne, dermatitis, atopic dermatitis, seborrheic dermatitis, contact dermatitis, urticaria, pruritis, and eczema.

15 Compounds of the present invention may also have advantageous properties such as low cytotoxicity, reduced HERG inhibition, low genotoxicity, reduced skin irritation, improved metabolic stability properties and metabolic elimination properties, dermal delivery properties, systemic exposure characteristics after dermal delivery, all of which may make them especially suitable to be used as 20 active pharmaceutical ingredients in drug formulations for the treatment of dermal diseases.

Accordingly, the present invention relates to a compound according to formula I



I

wherein X_1 , X_2 , X_3 , X_4 and X_5 independently of each other represent $-\text{CH-}$ or N ;

5 or X_3 , X_4 and X_5 independently of each other represent $-\text{CH-}$ or N , and X_1 and X_2 independently of each other represent C and form part of an additional 6-membered aromatic ring;

10 wherein R_1 represents alkyl, alkenyl, alkynyl, haloalkyl, hydroxyalkyl, or alkylcarbonyl, all of which are optionally substituted by one or more, same or different substituents selected from R_4 , or R_1 represents hydrogen;

15 R_2 represents alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, hydroxyalkyl, heterocycloalkenyl, alkylaryl, arylalkyl, alkylalkoxycarbonyl, alkylcarbonyloxy, or alkoxyalkyl, all of which are optionally substituted by one or more, same or different substituents selected from R_5 ; or R_2 represents hydrogen or $-\text{CH}_2\text{-C(O)NR}_9\text{-R}_{12}$;

20 R_3 represents alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, hydroxyalkyl, heterocycloalkenyl, alkylaryl, arylalkyl, alkylalkoxycarbonyl, alkylcarbonyloxy, or alkoxyalkyl, all of which are optionally substituted by one or more, same or different substituents selected from R_6 ; or R_3 represents hydrogen, $-\text{CH}_2\text{-C(O)-heterocycloalkyl}$ or $-\text{CH}_2\text{-C(O)NR}_9\text{-R}_{12}$;

25 R_4 represents hydrogen, alkyl, alkenyl, alkynyl, halogen, oxo, alkoxy, hydroxy, or haloalkyl;

30 R_5 represents alkylaryl, carboxy, alkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, carbamoyl, hydroxyalkyl, aryloxy, alkoxy carbonyloxy, alkoxy carbonyl, alkoxy, alkoxyalkyl, aryl, a heterocyclic ring, aminocarbonyl, alkylthio, alkylcarbonylamino, hydroxy, alkylcarbonyl, arylcarbonyl, alkylcarbonyloxy, or amino, all of which are optionally substituted by one or more, same or different substituents selected from R_7 ;

or R_5 represents hydrogen, oxo, halogen, cyano, or nitro;

R₆ represents alkylaryl, carboxy, alkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, carbamoyl, hydroxyalkyl, aryloxy, alkoxy carbonyloxy, alkoxy carbonyl, alkoxy, alkoxyalkyl, aryl, a heterocyclic ring, aminocarbonyl, 5 alkylthio, alkylcarbonylamino, arylcarbonyl, hydroxy, alkylcarbonyl, alkylcarbonyloxy, or amino, all of which are optionally substituted by one or more, same or different substituents selected from R₈;
or R₆ represents hydrogen, oxo, halogen, cyano, or nitro;

10 R₇ represents alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkynyl, alkoxy, haloalkyl, alkylthio, heterocycloalkenyl, heterocycloalkyl, aryl, alkylcarbonyl, heteroaryl, aryloxy, alkoxy carbonyl, hydroxyalkyl, amino, hydroxy, or carboxy; all of which are optionally substituted by one or more, same or different substituents selected from R₁₀;
15 or R₇ represents hydrogen, halogen, or oxo;

R₈ represents alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkynyl, alkoxy, haloalkyl, alkylthio, alkylsulfonyl, alkylsulfinyl, heterocycloalkenyl, heterocycloalkyl, aryl, alkylcarbonyl, heteroaryl, aryloxy, alkoxy carbonyl, hydroxyalkyl, amino, 20 hydroxy, or carboxy; all of which are optionally substituted by one or more, same or different substituents selected from R₁₀;
or R₈ represents hydrogen, halogen, or oxo;

R₉ represents hydrogen, alkyl, haloalkyl, or hydroxyalkyl;

25 R₁₀ represents hydrogen, alkyl, oxo, hydroxy, halogen, carboxy, amino, alkoxy, haloalkyl, or hydroxyalkyl;

R₁₁ represents one or more, same or different substituents selected from 30 hydrogen, halogen, cyano, amino, alkyl, methylsulfinyl, methylsulfonyl, amino, cyano, or alkoxy;

R₁₂ represents alkylaryl, arylalkyl, carboxy, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, carbamoyl, hydroxyalkyl, aryloxy, alkoxy carbonyloxy,

alkoxycarbonyl, alkoxy, alkoxyalkyl, aryl, a heterocyclic ring, aminocarbonyl, alkylthio, alkylcarbonylamino, hydroxy, alkylcarbonyl, arylcarbonyl, alkylcarbonyloxy, or amino, all of which are optionally substituted by one or more, same or different substituents selected from R₈,

5 or R₁₂ represents hydrogen;

with the proviso R₁, R₂ and R₃ cannot all be methyl;

with the proviso that when R₂ and R₃ are both hydrogen, R₁ cannot be methyl or hydrogen;

10 with the proviso that when R₁ is methyl or hydrogen, R₂ is methyl and R₃ is hydrogen, then ring B cannot be phenyl;

and pharmaceutically acceptable and physiologically cleavable esters, pharmaceutically acceptable salts, hydrates, N-oxides or solvates thereof.

15

In another aspect, the invention relates to a pharmaceutical composition comprising a compound of general formula I as defined herein together with a pharmaceutically acceptable vehicle or excipient or pharmaceutically acceptable carrier(s), optionally together with one or more other therapeutically active 20 compound(s).

In yet another aspect, the invention relates to the use of a compound according to formula I as defined herein, and pharmaceutically acceptable and physiologically cleavable esters, pharmaceutically acceptable salts, hydrates, N-oxides or solvates thereof, in the manufacture of a medicament for the 25 prophylaxis, treatment or amelioration of dermal diseases or conditions, or acute or chronic cutaneous wound disorders.

In yet another aspect, the invention relates to a method of preventing, treating 30 or ameliorating dermal diseases or conditions, or acute or chronic cutaneous wound disorders, the method comprising administering to a person suffering from at least one of said diseases an effective amount of one or more compounds of formula I as defined herein and pharmaceutically acceptable and

physiologically cleavable esters, pharmaceutically acceptable salts, hydrates, N-oxides or solvates thereof;
optionally together with a pharmaceutically acceptable carrier or one or more excipients, optionally in combination with other therapeutically active
5 compounds.

In yet another aspect, the present invention relates to a method for the preparation or manufacture of a compound of general formula I comprising a method as anywhere described herein, such as method a), b), c) d), or such as
10 any one of the general methods or procedures described in the examples or preparations herein, and optionally further processing of a compound obtained, to give a compound of general formula I as anywhere defined herein.

DETAILED DESCRIPTION OF THE INVENTION

15 The term "hydrocarbon radical" is intended to indicate a radical containing only hydrogen and carbon atoms, it may contain one or more double and/or triple carbon-carbon bonds, and it may comprise cyclic moieties in combination with branched or linear moieties. Said hydrocarbon comprises 1-20 carbon atoms, and preferably comprises 1-12, e.g. 1-6, e.g. 1-4, e.g. 1-3, e.g. 1-2 carbon
20 atoms. The term includes alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkynyl and aryl, as indicated below.

The term "aryl" is intended to indicate a radical of aromatic carbocyclic rings comprising 6-20 carbon atoms, such as 6-14 carbon atoms, preferably 6-10
25 carbon atoms, in particular 5- or 6-membered rings, including fused carbocyclic rings with at least one aromatic ring, such as phenyl, naphthyl, indenyl and indanyl.

The term "heteroaryl" is intended to indicate radicals of heterocyclic aromatic rings comprising 1-6 heteroatoms (selected from O, S and N) and 1-20 carbon atoms, such as 1-5 heteroatoms and 1-10 carbon atoms, such as 1-5 heteroatoms and 1-6 carbon atoms, such as 1-5 heteroatoms and 1-3 carbon atoms, in particular 5- or 6-membered rings with 1-4 heteroatoms selected from O, S and N, including fused bicyclic rings with 1-4 heteroatoms, and wherein at

least one ring is aromatic, e.g. pyridyl, quinolyl, isoquinolyl, indolyl, thiadiazolyl, oxodiazolyl, tetrazolyl, furanyl, pyridyl, thiazolyl, benzoazolyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thienyl, pyrazinyl, isothiazolyl, benzimidazolyl, and benzofuranyl.

5

In the present context, the term "alkyl" is intended to indicate a radical obtained when one hydrogen atom is removed from a hydrocarbon. Said alkyl comprises 1-20, preferably 1-12, such as 1-6, such as 1-4 or such as 1-3 carbon atoms.

10 The term includes the subclasses normal alkyl (*n*-alkyl), secondary and tertiary alkyl, such as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, sec.-butyl, *tert*.-butyl, pentyl, isopentyl, hexyl and isohexyl.

15 The term "cycloalkyl" is intended to indicate a saturated cycloalkane radical comprising 3-20 carbon atoms, preferably 3-10 carbon atoms, in particular 3-8 carbon atoms, such as 3-6 carbon atoms, including fused bicyclic rings, e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, or cycloheptyl.

20 The term "heterocycloalkyl" is intended to indicate a cycloalkane radical as described above, wherein one or more carbon atoms are replaced by heteroatoms, comprising 1-20 carbon atoms, e.g. 2-5 or 2-4 carbon atoms, further comprising 1-6 heteroatoms, preferably 1, 2, or 3 heteroatoms, selected from O, N, or S, e.g. piperidinyl, pyrrolidinyl, morpholinyl, [1,3]dioxolanyl and [1,3]dioxolyl, or including fused bicyclic rings with 1-4 heteroatoms, wherein at 25 least one ring comprises a heteroatom, and wherein the other ring may for example be a carbocyclic ring, e.g. isoindolyl.

30 The term "alkenyl" is intended to indicate a mono-, di-, tri-, tetra- or pentaunsaturated hydrocarbon radical comprising 2-10 carbon atoms, in particular 2-6 carbon atoms, such as 2-4 carbon atoms, e.g. ethenyl, propenyl (allyl), methylbutenyl, butenyl, pentenyl or hexenyl.

The term "cycloalkenyl" is intended to indicate mono-, di- tri- or tetraunsaturated non-aromatic cyclic hydrocarbon radicals, comprising 3-20 carbon atoms, including fused bicyclic rings, typically comprising 3-10 carbon

atoms, such as 3, 4, or 6 carbon atoms, e.g. cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl.

The term "heterocycloalkenyl" is intended to indicate a cycloalkene radical as

5 described above, wherein one or more carbon atoms are replaced by heteroatoms, comprising 1-20 carbon atoms, e.g. 2-4 carbon atoms, further comprising 1-6 heteroatoms, preferably 1, 2, or 3 heteroatoms, selected from O, N, or S, including fused bicyclic rings with 1-4 heteroatoms, wherein at least one ring comprises a heteroatom and wherein the other ring may for example be a
10 carbocyclic ring, e.g. dihydrofuranyl, or 2,5-dihydro-1H-pyrrolyl.

The term "alkynyl" is intended to indicate an hydrocarbon radical comprising 1-5 triple C-C bonds and 2-20 carbon atoms, typically comprising 2-10 carbon atoms, in particular 2-6 carbon atoms, such as 2-4 carbon atoms, e.g. ethynyl, propynyl, butynyl, pentynyl or hexynyl.

The term "halogen" is intended to indicate a substituent from the 7th main group of the periodic table, such as fluoro, chloro and bromo.

20 The term "haloalkyl" is intended to indicate an alkyl group as defined above substituted with one or more halogen atoms as defined above, e.g. fluoro or chloro, such as difluoromethyl, or trifluoromethyl.

The term "alkoxy" is intended to indicate a radical of the formula -OR', wherein

25 R' is alkyl as indicated above, e.g. methoxy, ethoxy, n-propoxy, isopropoxy, butoxy, etc.

The term "hydroxyalkyl" is intended to indicate an alkyl group as defined above substituted with one or more hydroxy, e.g. hydroxymethyl, hydroxyethyl,

30 hydroxypropyl.

The term "amino" is intended to indicate a radical of the formula -NR''₂, wherein each R'' independently represents hydrogen, or a hydrocarbon radical as indicated above, e.g. -NH₂, dimethylamino, -NHMe, -NHEt, *tert*.-butylamino.

The term "alkylthio" is intended to indicate a radical of the formula $-S-R'$, wherein R' is alkyl as indicated above, e.g. $-SMe$.

- 5 The term "alkoxycarbonyl" is intended to indicate a radical of the formula $-C(O)-O-R'$, wherein R' is alkyl as indicated above, e.g. methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, isopropoxycarbonyl, tert-butoxycarbonyl etc.
- 10 The term "aminocarbonyl" is intended to indicate a radical of the formula $-C(O)-NR''_2$, wherein each R'' is as indicated above.

The term "alkylcarbonylamino" is intended to indicate a radical of the formula $-NR''-C(O)-R'$, wherein R' is alkyl as indicated above, and each R'' is as indicated above.

The term "alkylcarbonyl" is intended to indicate a radical of the formula $-C(O)-R'$, wherein R' is alkyl as indicated above, e.g. acetyl.

- 20 The term "alkylcarbonyloxy" is intended to indicate a radical of the formula $-O-C(O)-R'$, wherein R' is alkyl as indicated above.

The term "alkoxycarbonyloxy" is intended to indicate a radical of the formula $-O-C(O)-O-R'$, wherein R' is alkyl as indicated above.

- 25 The term "heterocyclic ring" is intended to include the definitions heteroaryl, heterocycloalkyl and heterocycloalkenyl as defined above, including annelated ring systems with each other or with cyclic hydrocarbons, e.g. 2,5-dihydrobenzo(b)dioxocine, 2,3,5,8-tetrahydro-[1,4]dioxocine, 5,8-dihydro-[1,4]dioxocine, 2,3-dihydro-1H-isoindole.
- 30

The term "alkylaryl" is intended to indicate an aryl radical as defined above, which is substituted with an alkyl radical as defined above, e.g. toyl (=toloyl), ethylbenzene, etc.

The term "arylalkyl" is intended to indicate an alkyl radical as defined above, which is substituted with an aryl radical as defined above, e.g. benzyl, phenylethyl, naphthylmethyl, etc.

5

The term "alkoxyalkyl" is intended to indicate an alkyl radical as defined above, which is substituted with an alkoxy radical as defined above, i.e. $-R'-O-R'$, wherein each R' is alkyl, same or different, as indicated above, e.g. methoxymethyl, ethoxymethyl.

10

The term "aryloxy" is intended to indicate $-O-R''$, wherein R'' is aryl as indicated above, e.g. phenoxy.

15 The term "arylcarbonyl" is intended to indicate $-C(O)-R'''$, wherein R''' is an aryl radical as defined above, e.g. benzoyl, naphthylcarbonyl.

The term "alkylalkoxycarbonyl" is intended to indicate $-R'-C(O)-O-R'$, wherein each R' is alkyl, same or different, as indicated above, e.g. tert-butoxycarbonylmethyl, methoxycarbonylmethyl.

20

The term "pharmaceutically acceptable salt" is intended to indicate salts prepared by reacting a compound of formula I with a suitable inorganic or organic acid, such as hydrochloric, hydrobromic, hydroiodic, sulfuric, nitric, phosphoric, formic, acetic, 2,2-dichloroacetic, adipic, ascorbic, L-aspartic, L-glutamic, galactaric, lactic, maleic, L-malic, phthalic, citric, propionic, benzoic, glutaric, gluconic, D-glucuronic, methanesulfonic, salicylic, succinic, malonic, tartaric, benzenesulfonic, ethane-1,2-disulfonic, 2-hydroxy ethanesulfonic acid, toluenesulfonic, sulfamic or fumaric acid. Pharmaceutically acceptable salts of compounds of formula I may also be prepared by reaction with a suitable base such as sodium hydroxide, potassium hydroxide, magnesium hydroxide, calcium hydroxide, silver hydroxide, ammonia or the like, or suitable non-toxic amines, such as lower alkylamines, for example triethylamine, hydroxy-lower alkylamines, for example 2-hydroxyethylamine, bis-(2-hydroxyethyl)-amine, cycloalkylamines, for example dicyclohexylamine, or benzylamines, for example

N,N'-dibenzylethylenediamine, and dibenzylamine, or L-arginine or L-lysine. Salts obtained by reaction with a suitable base include, but are not limited to sodium salts, choline salts, 2-(dimethylamino)-ethanol salts, 4-(2-hydroxyethyl)-morpholin salts, L-lysine salts, N-(2-hydroxyethyl)-pyrrolidine salts, 5 ethanolamine salts, potassium salts, tetrabutylammonium salts, benzyltrimethylammonium salts, cetyltrimethylammonium salts, tetramethylammonium salts, tetrapropylammonium salts, tris(hydroxymethyl)aminomethane salts, N-methyl-D-glucamine salts, silver salts, benzethonium salts, and triethanolamine salts.

10

The term "solvate" is intended to indicate a species formed by interaction between a compound, e.g. a compound of formula I, and a solvent, e.g. alcohol, glycerol or water, wherein said species are in a solid form. When water is the solvent, said species is referred to as a hydrate.

15

Compounds of the invention which comprise free hydroxyl groups or free carboxylic acid groups may also exist in the form of pharmaceutically acceptable, physiologically cleavable esters, and as such are included within the scope of the present invention. Such pharmaceutically acceptable esters are 20 preferably prodrug ester derivatives, such being convertible by solvolysis or cleavage under physiologically conditions to the corresponding compounds of the invention which comprise free hydroxyl groups or free carboxylic acid groups respectively, e.g. *in-vivo* hydrolysable.

25 Embodiments of the present invention


In one or more embodiments of the present invention ring B represents pyridyl, pyrazinyl, quinolyl, pyrimidinyl or pyridazinyl optionally substituted with one or more same or different substituents selected from fluoro, chloro, bromo, cyano, methoxy, -NH₂ or C₁₋₄amino.

30

In one or more embodiments of the present invention ring B, optionally substituted with R₁₁, represents 2-(6-chloro-pyrazinyl), 2-pyrazinyl, 4-(3-bromo-pyridyl), 4-(3,5-dibromo-pyridyl), 4-(6-chloro-pyrimidinyl), 2-(4-chloro-pyridyl), 3-(2-chloro-pyridyl), 4-(2-methoxy-pyridyl), 4-(2-cyano-pyridyl), 3-pyridazinyl,

4-(2-*tert*-butylamino-3,5-dichloro-pyridyl), 4-(2-amino-3,5-dichloro-pyridyl), 4-(3,5-dichloro-pyridyl), 2-(3-bromo-pyrazinyl), 4-pyridyl, 4-quinolyl or 4-(3,5-dichloro-1-oxy-pyridyl).

5 In one or more embodiments of the present invention formula I represents general formula Iz,



Iz

10 wherein X_3 represents $-\text{CH-}$ or N , and wherein R_1 , R_2 , and R_3 are as anywhere defined herein.

In one or more preferred embodiments of the present invention R_1 represents alkyl, alkenyl, alkynyl, haloalkyl, hydroxyalkyl, or alkylcarbonyl, all of which are 15 optionally substituted by one or more, same or different substituents selected from R_4 , and/or R_2 represents alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, hydroxyalkyl, heterocycloalkenyl, alkylaryl, alkylalkoxycarbonyl, arylalkyl, alkylcarbonyloxy, or alkoxyalkyl, all of which are optionally substituted by one or 20 more, same or different substituents selected from R_5 ; or R_2 represents $-\text{CH}_2-\text{C}(\text{O})\text{NR}_9\text{R}_{12}$.

In one or more preferred embodiments of the present invention R_1 represents methyl or ethyl.

In one or more embodiments of the present invention R_2 represents C_1 - C_6 alkyl, C_1 - C_6 alkenyl, C_1 - C_6 alkoxy C_1 - C_6 alkyl, hydroxy C_1 - C_6 alkyl, halo C_1 - C_6 alkyl, C_1 - C_6 alkynyl, C_1 - C_6 cycloalkyl, C_1 - C_6 alkyl C_6 - C_{10} aryl, C_1 - C_6 alkyl C_{1-6} alkoxycarbonyl or C_1 - C_6 alkylcarbonyloxy, all of which are optionally substituted by one or more, 5 same or different substituents selected from R_5 .

In one or more embodiments of the present invention R_2 represents methyl, ethyl, propyl, tert-butoxycarbonylmethyl, allyl, difluoromethyl, ethylbenzene, methylbenzene, butenyl, hydroxyethyl, ethylphenyl, tolyl, pentenyl, 10 methoxyethyl, butinyl, propinyl, cyclopentyl, all of which are optionally substituted by one or more, same or different substituents selected from R_5 .

In one or more embodiments of the present invention R_2 represents cyclopropyl, methylcyclopropyl, cyclopentyl, methyl, ethyl, propyl, or allyl, all of which are 15 optionally substituted by one or more, same or different substituents selected from hydroxy, fluoro, or alkoxy.

In one or more embodiments of the present invention R_3 represents methyl, ethyl, propyl, butyl, pentyl, hydroxyethyl, hexyl, butenyl, pentenyl, allyl, butinyl, 20 methylbenzene, ethylbenzene, tolyl, toluoyl, propylbenzene, methylnaphthyl, ethylnaphthyl, methylcarbonylmethoxy, methylcarbonylethoxy, methoxyethyl, methoxypropyl, all of which are optionally substituted by one or more, same or different substituents selected from R_6 , all of which are optionally substituted by 25 one or more, same or different substituents selected from R_8 , or R_3 represents hydrogen, $-CH_2-C(O)$ -heterocycloalkyl or $-CH_2-C(O)NR_9-R_{12}$.

In one or more embodiments of the present invention R_3 represents methyl, ethyl, propyl, or allyl, all of which are optionally substituted by one or more, same or different substituents selected from hydroxy, fluoro, or alkoxy.

30 In one or more embodiments of the present invention R_5 represents methyl, tert-butoxy, ethenyl, cyclopropyl, propenyl, phenyl, butenyl, propinyl, ethylhydroxy, ethinyl, allyl, ethyl, or methoxy,

all of which are optionally substituted by one or more, same or different substituents selected from R₇, or R₅ represents hydrogen, oxo, chloro, fluoro, or hydroxy.

- 5 In one or more embodiments of the present invention R₆ represents ethenyl, methyl, tert-butoxy, isoxazolyl, methoxy, propinyl, butenyl, phenyl, pyridyl, benzoxazolyl, thiazolyl, [1,3,4]thiadiazolyl, [1,2,4]oxadiazolyl, 2,3-dihydro-1H-isoindolyl, ethoxy, thiophenyl, propyl, ethyl, butyl, pentyl, allyl, isopropoxy, isopropyl, naphthyl, cyclohexyl, hydroxy, cyclopentyl, phenoxy, tolyl, toluoyl, 10 benzoyl, carbonylnaphthalene, ethylbenzene, quinolinyl, -NH₂, ethoxycarbonyl, methoxycarbonyl, carbamoyl, isoindole, methylamine, pyrrolidyl, morpholinyl, methylsulfonyl, methylsulfinyl, butylamine, propylamine, ethylamine, cycloheptyl, hydroxyethyl, hydroxypropyl, indanyl, or ethoxyethyl, all of which are optionally substituted by one or more, same or different substituents 15 selected from R₈, or R₆ represents hydrogen, oxo, fluoro, chloro, or cyano.

In one or more embodiments of the present invention R₈ represents represents, methyl, ethyl, propyl, butyl, phenyl, cyclopropyl, ethoxy, methoxy, allyl, ethenyl, ethoxycarbonyl, hydroxy, naphthyl, cyclohexyl, methoxycarbonyl, phenoxy, 20 isopropoxy, -NH₂, methylamine, pyrrolidinyl, morpholinyl, methylsulfonyl, methylsulfinyl, cycloheptyl, cyclopentyl, hydroxymethyl, hydroxyethyl, dimethylamino, furanyl, pyridyl, tolyl, piperidinyl, acetyl, thiophenyl, cycloheptyl, all of which are optionally substituted by one or more, same or different substituents selected from R₁₀, or R₈ represents hydrogen, oxo, chloro, bromo, 25 fluoro, cyano, or trifluoromethyl.

In one or more embodiments of the present invention R₉ represents hydrogen, methyl, or ethyl.

- 30 In one or more embodiments of the present invention R₁₀ represents hydrogen, oxo, methyl, hydroxy, fluoro, cyano, chloro, or methoxy.

In one or more embodiments of the present invention R₂ represents methyl.

In one or more embodiments of the present invention R₁ and R₂ both represent methyl.

5 In one or more embodiments of the present invention R₁ and R₃ represent methyl and/or difluoromethyl.

In one or more embodiments of the present invention R₃ represents -CH₂-C(O)NH-R₁₂, -CH₂-C(O)NH-heterocycloalkyl, -CH₂CH₂-phenyl-R₆, or -CH₂-phenyl-R₆.

10 In one or more embodiments of the present invention R₂ and/or R₃ represents -CH₂COOH, methyl, hydrogen, allyl, ethyl, tert-butoxycarbonylmethyl, difluoromethyl, 3-methyl-5-methylisoxazole, 2-methoxy-ethane, 2-butyne, 2-methyl-2-butene, 2-phenylethane, benzyl, 2-methyl-1,3-benzoxazole, 4-methyl-15 2-methylthiazole, 2-methyl-5-cyclopropyl-[1,3,4]thiadiazole, 3-methyl-[1,2,4]oxadiazole, ethyl acetate, 4-chlorobenzyl, 5-chloro-2-methyl-thiophene, phenoxyethane, (4-methylphenyl)ethane, 3-phenylpropane, (3-methoxyphenyl)ethane, (4-methoxyphenyl)ethane, (3-bromophenyl)ethane, (2-methoxyphenyl)ethane, (4-fluorophenyl)ethane, (2-fluorophenyl)ethane, (3,4-20 dimethoxyphenyl)ethane, benzyl acetate, isopropyl acetate, 3-methyl benzoic acid methyl ester, 3-methyl-butane, 1-hexyl, but-1-ene, pent-1-ene, 1-propyl, 1-butyl, 2-methyl-propane, butyric acid ethyl ester, 4-methyl-benzyl, 3-chlorobenzyl, propoxybenzene, 1-(4-methoxy-phenyl)-ethanone, 4-methyl-benzonitrile, 2-methyl-naphthalene, 1-pentyl, methyl-cyclohexane, 3-methyl-benzonitrile, 1-ethoxy-4-chloro-benzene, 2-ethyl-butane, 2-hydroxy-ethane, 4-methyl benzoic acid methyl ester, 1-naphthalen-2-yl-ethanone, 2,5-dimethoxy-phenyl-ethanone, 1-p-tolyl-ethanone, 4-fluoro-benzyl, 2-fluoro-benzyl, 5-trifluoromethyl-benzyl, 5-trifluoromethoxy-benzyl, 3-fluoro-5-trifluoromethyl-benzyl, 1-(2-methoxy-phenyl)-ethanone, 1-(2,4-dimethyl-phenyl)-ethanone, 4-chloro-benzyl, 2-difluoromethoxy-benzyl, 4-isopropyl-benzyl, 2-fluoro-6-trifluoromethyl-benzyl, 2,3-difluoro-4-methyl-benzyl, 2-methyl-benzyl, 3-methyl-benzyl, pent-2-ene, 6-methyl-2-methyl-quinoline, 2-chloro-benzyl, 3-methoxy-benzyl, 4-methoxy-benzyl, (3-Chloro-phenyl)-ethane, 5-methyl-hexane, ethyl-cyclohexane, pentanoic acid ethyl ester, (propoxymethyl)-

benzene, acetamide, 2-ethyl-isoindole-1,3-dione, 2-propyl-isoindole-1,3-dione,
N-methyl-acetamide, methyl-cyclopropane, but-1-ene, 4-yl-but-1-ene, 2-
methyl-pent-2-ene, ethanol, benzyl, pent-2-ene, 2-methoxy-ethane, but-2-yne,
propyne, acetate, 1-pyrrolidin-1-yl-ethanone, N-benzylacetamide, 1-morpholin-
5 4-yl-ethanone, N-phenyl-acetamide, N-methyl-N-phenyl-acetamide, N-(3-
hydroxy-3-methyl-butyl)-acetamide, N-n-propyl-acetamide, N-ethyl-acetamide,
N-isopropyl-acetamide, N-butyl-acetamide, N-cyclopentyl-acetamide, N-(3-
methyl-butyl)-acetamide, N-(4-methoxy-benzyl)-acetamide, N-(2,2-dimethyl-
propyl)-acetamide, N-cyclohexyl-acetamide, N-(3-methoxy-benzyl)-acetamide,
10 N-cycloheptyl- acetamide, N-(2-methoxy-benzyl)-acetamide, N-
cyclohexylmethyl-acetamide, N-(2-hydroxy-ethyl)-acetamide, N-(1-phenyl-
ethyl)-acetamide, N-(3-hydroxy-propyl)-acetamide, N-(2-methoxy-ethyl)-
acetamide, N-(2-dimethylamino-ethyl)-acetamide, N-(3-dimethylamino-propyl)-
acetamide, N-(1-phenyl-ethyl)-acetamide, N-(3-isopropoxy-propyl)-acetamide,
15 N-furan-2-ylmethyl-acetamide, N-pyridin-2-ylmethyl-acetamide, N-pyridin-3-
ylmethyl-acetamide, N-(2-phenoxy-ethyl)-acetamide, N-pyridin-4-ylmethyl-
acetamide, N-(4-ethyl-benzyl)-acetamide, N-(3,5-difluoro-benzyl)-acetamide, N-
(2,3-difluoro-benzyl)-acetamide, N-(2-pyridin-2-yl-ethyl)-acetamide, N-(2-
methyl-benzyl)-acetamide, N-(3-fluoro-benzyl)-acetamide, N-(3-methyl-benzyl)-
20 acetamide, N-(4-methyl-benzyl)-acetamide, N-phenethyl-acetamide, N-(2-
pyridin-4-yl-ethyl)-acetamide, N-(3-phenyl-propyl)-acetamide, N-(2-chloro-
benzyl)-acetamide, N-(2-piperidin-1-yl-ethyl)-acetamide, N-(3-chloro-benzyl)-
acetamide, N-(2-morpholin-4-yl-ethyl)-acetamide, N-(4-chloro-benzyl)-
acetamide, N-(2-pyridin-3-yl-ethyl)-acetamide, N-(2-pyrrolidin-1-yl-ethyl)-
25 acetamide, N-(2-acetylamino-ethyl)-acetamide, (R)-N-(2-hydroxy-2-phenyl-
ethyl)-acetamide, (S)-N-(2-hydroxy-2-phenyl-ethyl)-acetamide, N-thiophen-2-
ylmethyl-acetamide, N-[3-(2-oxo-pyrrolidin-1-yl)-propyl]-acetamide, N-(2-
hydroxy-indan-1-yl)-acetamide, N-cycloheptylmethyl-acetamide, N-[2-(2-
hydroxy-ethoxy)-ethyl]-acetamide, N-(4-dimethylamino-butyl)-acetamide,
30 cyclopentan, cyclopropylmethyl, ethyl, phenyl-ethane, acetic acid benzyl ester,
2-methyl-benzonitrile, (1-oxy-pyridin-4-yl)ethane, (4-pyridyl)ethane, (3-
pyridyl)ethane, (2-pyridyl)ethane, (4-benzonitrile)ethane, (4-methylsulfinyl-
phenyl)ethane, (4-methylsulfonyl-phenyl)ethane, 1-phenyl-propane, 2-phenyl-
propane or 1-methyl-2-phenyl-ethane.

In one or more embodiments of the present invention R_{12} represents alkyl, cycloalkyl, hydroxyalkyl, aryl, arylalkyl, alkylcarbonylamino, all of which are optionally substituted with one or more same or different substituents selected

5 from alkyl, cycloalkyl, alkoxy, heterocycloalkyl, heteroaryl, aryloxy, amino, hydroxy, halogen, oxy, all of which are optionally substituted with oxo or hydroxyl, or R_{12} represents hydrogen;

In one or more embodiments of the present invention, R_{12} represents methyl,

10 ethyl, 1-propyl, or 2-propyl, optionally substituted with one or more, same or different substituents selected from fluoro, chloro, bromo, or methyl; or R_{12} represents hydrogen.

In yet another embodiment, the present invention relates to a compound of

15 general formula Iz , wherein X_3 represents CH , R_1 and R_2 are both methyl, and R_3 is as anywhere described herein.

In one or more embodiments of the present invention, R_1 and R_2 cannot both be hydrogen.

20

In one or more embodiments of the present invention, R_2 and/or R_3 are as described anywhere in the following exemplified compounds.

The present invention includes all embodiments wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_6 ,

25 R_7 , R_8 , R_9 , R_{10} , R_{11} and R_{12} are combined in any combination as anywhere described herein.

In one or more embodiments of the present invention, the compounds of

general structure I have a molecular weight below 800 Dalton, such as below

30 750 Dalton, e.g. below 700 Dalton, or below 650, 600, 550, or 500 Dalton.

In particular compounds of formula I may be selected from

2-(3,5-Dichloro-pyridin-4-yl)-1-(2-hydroxy-3,4-dimethoxy-phenyl)-ethanone
(compound 101),

2-(3,5-Dichloro-pyridin-4-yl)-1-(3-hydroxy-2,4-dimethoxy-phenyl)-ethanone (compound 102),
1-(2-Allyloxy-3-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 103),
5 2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3-diethoxy-4-methoxy-phenyl)-ethanone (compound 104),
{2-tert-Butoxycarbonylmethoxy-6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-3-methoxy-phenoxy}-acetic acid tert-butyl ester (compound 105),
1-(2,3-Bis-allyloxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone
10 (compound 106),
1-(2,3-Bis-difluoromethoxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 107),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(5-methyl-isoxazol-3-ylmethoxy)-phenyl]-ethanone (compound 108),
15 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methoxy-ethoxy)-phenyl]-ethanone (compound 109),
1-(2-But-2-ynylmethoxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 110),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methyl-but-2-enyloxy)-phenyl]-ethanone (compound 111),
20 2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-phenethoxy-phenyl)-ethanone (compound 112),
1-(2-Benzylmethoxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 113),
25 1-(2-Allyloxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 114),
1-[2-(Benzoxazol-2-ylmethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 115),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methyl-thiazol-4-ylmethoxy)-phenyl]-ethanone (compound 116),
30 1-[2-(5-Cyclopropyl-[1,3,4]thiadiazol-2-ylmethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 117),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-([1,2,4]oxadiazol-3-ylmethoxy)-phenyl]-ethanone (compound 118),

{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid ethyl ester (compound 119),
1-{2-[2-(4-Chloro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 120),
5 1-[2-(5-Chloro-thiophen-2-ylmethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 121),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-phenoxy-ethoxy)-phenyl]-ethanone (compound 122),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-p-tolyl-ethoxy)-phenyl]-ethanone (compound 123),
10 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-phenyl-propoxy)-phenyl]-ethanone (compound 124),
2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(3-methoxy-phenyl)-ethoxy]-phenyl}-ethanone (compound 125),
15 2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(4-methoxy-phenyl)-ethoxy]-phenyl}-ethanone (compound 126),
1-{2-[2-(3-Bromo-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 127),
2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(2-methoxy-phenyl)-ethoxy]-phenyl}-ethanone (compound 128),
20 2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 129),
2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(2-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 130),
25 2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(3,4-dimethoxy-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 131),
{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid benzyl ester (compound 132),
30 {6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid isopropyl ester (compound 133),
3-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy-methyl}-benzoic acid methyl ester (compound 134),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methyl-butoxy)-phenyl]-ethanone (compound 135),

2-(3,5-Dichloro-pyridin-4-yl)-1-(2-hexyloxy-3,4-dimethoxy-phenyl)-ethanone
(compound 136),

1-(2-But-3-enyloxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-
ethanone (compound 137),

5 2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-pent-4-enyloxy-phenyl)-
ethanone (compound 138),

2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-propoxy-phenyl)-ethanone
(compound 139),

1-(2-Butoxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone
10 ethanone (compound 140),

2-(3,5-Dichloro-pyridin-4-yl)-1-(2-isobutoxy-3,4-dimethoxy-phenyl)-ethanone
(compound 141),

4-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-butyric
acid ethyl ester (compound 142),

15 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(4-methyl-benzyloxy)-
phenyl]-ethanone (compound 143),

1-[2-(3-Chloro-benzyloxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-
ethanone (compound 144),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-phenoxy-propoxy)-phenyl]-
20 ethanone (compound 145),

2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(4-methoxy-phenyl)-2-
oxo-ethoxy]-phenyl}-ethanone (compound 146),

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy-methyl}-
benzonitrile (compound 147),

25 4-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy-methyl}-
benzonitrile (compound 148),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(naphthalen-2-ylmethoxy)-
phenyl]-ethanone (compound 149),

2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-pentyloxy-phenyl)-ethanone
30 (compound 150),

1-(2-Cyclohexylmethoxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-
ethanone (compound 151),

3-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy-methyl}-
benzonitrile (compound 152),

1-{2-[2-(4-Chloro-phenoxy)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 153),
2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-ethyl-butoxy)-3,4-dimethoxy-phenyl]-ethanone (compound 154),
5 2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-hydroxy-ethoxy)-3,4-dimethoxy-phenyl]-ethanone (compound 155),
4-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxyethyl}-benzoic acid methyl ester (compound 156),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-naphthalen-2-yl-2-oxo-ethoxy)-phenyl]-ethanone (compound 157),
10 2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(2,5-dimethoxy-phenyl)-2-oxo-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 158),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-oxo-2-p-tolyl-ethoxy)-phenyl]-ethanone (compound 159),
15 2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(4-fluoro-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone (compound 160),
2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-fluoro-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone (compound 161),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-trifluoromethyl-benzyloxy)-phenyl]-ethanone (compound 162),
20 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-trifluoromethoxy-benzyloxy)-phenyl]-ethanone (compound 163),
2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(3-fluoro-5-trifluoromethyl-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone (compound 164),
25 2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(2-methoxy-phenyl)-2-oxo-ethoxy]-phenyl}-ethanone (compound 165),
2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(2,4-dimethyl-phenyl)-2-oxo-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 166),
1-[2-(4-Chloro-benzyloxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 167),
30 2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-difluoromethoxy-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone (compound 168),
2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(4-isopropyl-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone (compound 169),

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-fluoro-6-trifluoromethyl-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone (compound 170),
2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2,3-difluoro-4-methyl-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone (compound 171),
5 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methyl-benzyloxy)-phenyl]-ethanone (compound 172),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methyl-benzyloxy)-phenyl]-ethanone (compound 173),
2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-pent-2-enyloxy-phenyl)-ethanone (compound 174),
10 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methyl-quinolin-6-ylmethoxy)-phenyl]-ethanone (compound 175),
1-[2-(2-Chloro-benzyloxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 176),
15 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methoxy-benzyloxy)-phenyl]-ethanone (compound 177),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(4-methoxy-benzyloxy)-phenyl]-ethanone (compound 178),
1-{2-[2-(3-Chloro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 179),
20 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(5-methyl-hexyloxy)-phenyl]-ethanone (compound 180),
1-[2-(2-Cyclohexyl-ethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 181),
25 5-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-pentanoic acid ethyl ester (compound 182),
1-[2-(3-Benzyl-ethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 183),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide
30 (compound 184),
2-(2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-ethyl)-isoindole-1,3-dione (compound 185),
2-(3-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-propyl)-isoindole-1,3-dione (compound 186),

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-methyl-acetamide (compound 187),
2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-2,4-dimethoxy-phenyl)-ethanone (compound 188),
5 1-(3-Cyclopropylmethoxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 189),
1-(2-Allyloxy-3-but-3-enyloxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 190),
1-(3-But-3-enyloxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 191),
10 2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-propoxy-phenyl)-ethanone (compound 192),
1-(3-Allyloxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 193),
15 2-(3,5-Dichloro-pyridin-4-yl)-1-[2,4-dimethoxy-3-(4-methyl-pent-3-enyloxy)-phenyl]-ethanone (compound 194),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3-(2-hydroxy-ethoxy)-2,4-dimethoxy-phenyl]-ethanone (compound 195),
2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-phenethyloxy-phenyl)-ethanone (compound 196),
20 1-(3-Benzylxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 197),
2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-pent-2-enyloxy-phenyl)-ethanone (compound 198),
25 2-(3,5-Dichloro-pyridin-4-yl)-1-[2,4-dimethoxy-3-(2-methoxy-ethoxy)-phenyl]-ethanone (compound 199),
1-(3-But-2-nyloxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 200),
2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-prop-2-nyloxy-phenyl)-ethanone (compound 201),
30 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-oxo-2-pyrrolidin-1-yl-ethoxy)-phenyl]-ethanone (compound 202),
N-Benzyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 203),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-morpholin-4-yl-2-oxo-ethoxy)-phenyl]-ethanone (compound 204),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-phenyl-acetamide (compound 205),
5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-methyl-N-phenyl-acetamide (compound 206),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-hydroxy-3-methyl-butyl)-acetamide (compound 207),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-
10 acetamide (compound 208),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-isopropyl-acetamide (compound 209),
N-Butyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 210),
15 N-Cyclopentyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 211),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-methyl-butyl)-acetamide (compound 212),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-
20 methoxy-benzyl)-acetamide (compound 213),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2,2-dimethyl-propyl)-acetamide (compound 214),
N-Cyclohexyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 215),
25 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-methoxy-benzyl)-acetamide (compound 216),
N-Cycloheptyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 217),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-
30 methoxy-benzyl)-acetamide (compound 218),
N-Cyclohexylmethyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 219),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-ethyl)-acetamide (compound 220),

(R)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(1-phenyl-ethyl)-acetamide (compound 221),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-hydroxy-propyl)-acetamide (compound 222),
5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-methoxy-ethyl)-acetamide (compound 223),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-dimethylamino-ethyl)-acetamide (compound 224),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-dimethylamino-propyl)-acetamide (compound 225),
10 (S)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(1-phenyl-ethyl)-acetamide (compound 226),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-isopropoxy-propyl)-acetamide (compound 227),
15 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-furan-2-ylmethyl-acetamide (compound 228),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-pyridin-2-ylmethyl-acetamide (compound 229),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-pyridin-
20 3-ylmethyl-acetamide (compound 230),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-phenoxy-ethyl)-acetamide (compound 231),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-pyridin-4-ylmethyl-acetamide (compound 232),
25 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-ethyl-benzyl)-acetamide (compound 233),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3,5-difluoro-benzyl)-acetamide (compound 234),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2,3-
30 difluoro-benzyl)-acetamide (compound 235),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyridin-2-yl-ethyl)-acetamide (compound 236),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-methyl-benzyl)-acetamide (compound 237),

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-fluoro-benzyl)-acetamide (compound 238),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-methyl-benzyl)-acetamide (compound 239),
5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-methyl-benzyl)-acetamide (compound 240),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-phenethyl-acetamide (compound 241),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyridin-4-yl-ethyl)-acetamide (compound 242),
10 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-phenyl-propyl)-acetamide (compound 243),
N-(2-Chloro-benzyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 244),
15 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-piperidin-1-yl-ethyl)-acetamide (compound 245),
N-(3-Chloro-benzyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 246),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-morpholin-4-yl-ethyl)-acetamide (compound 247),
20 N-(4-Chloro-benzyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 248),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyridin-3-yl-ethyl)-acetamide (compound 249),
25 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyrrolidin-1-yl-ethyl)-acetamide (compound 250),
N-(2-Acetylamino-ethyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 251),
(R)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-2-phenyl-ethyl)-acetamide (compound 252),
30 (S)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-2-phenyl-ethyl)-acetamide (compound 253),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-thiophen-2-ylmethyl-acetamide (compound 254),

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-[3-(2-oxo-pyrrolidin-1-yl)-propyl]-acetamide (compound 255),
(2R)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-indan-1-yl)-acetamide (compound 256),
5 N-Cycloheptylmethyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 257),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide (compound 258),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-
10 dimethylamino-butyl)-acetamide (compound 259),
1-(3-Cyclopentyloxy-2-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 260),
1-(3-Cyclopropylmethoxy-2-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-
pyridin-4-yl)-ethanone (compound 261),
15 2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-2-hydroxy-4-methoxy-phenyl)-
ethanone (compound 262),
2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-4-methoxy-2-phenethyloxy-phenyl)-
ethanone (compound 263),
1-[2-(5-Cyclopropyl-[1,3,4]thiadiazol-2-ylmethoxy)-3-ethoxy-4-methoxy-
20 phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 264),
{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2-ethoxy-3-methoxy-phenoxy}-acetic
acid benzyl ester (compound 265),
1-(3-Allyloxy-2-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-
ethanone (compound 266),
25 2-{2-Allyloxy-6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-3-methoxy-
phenoxyethyl}-benzonitrile (compound 267),
1-(3-Allyloxy-4-methoxy-2-phenethyloxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-
ethanone (compound 268),
1-{3-Allyloxy-2-[2-(4-fluoro-phenyl)-ethoxy]-4-methoxy-phenyl}-2-(3,5-
30 dichloro-pyridin-4-yl)-ethanone (compound 269),
N-Benzyl-2-{6-[2-(3,5-dichloro-1-oxy-pyridin-4-yl)-acetyl]-2,3-dimethoxy-
phenoxy}-acetamide (compound 270),
2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-(3,4-dimethoxy-2-phenethyloxy-phenyl)-
ethanone (compound 271),

2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 272),
2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(1-oxy-pyridin-4-yl)-ethoxy]-phenyl}-ethanone (compound 274),
5 2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-(2-hydroxy-3,4-dimethoxy-phenyl)-ethanone (compound 275),
4-(2-{6-[2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-ethyl)-benzonitrile (compound 276),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-pyridin-4-yl-ethoxy)-phenyl]-ethanone (compound 277),
10 4-(2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-ethyl)-benzonitrile (compound 278),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-pyridin-2-yl-ethoxy)-phenyl]-ethanone (compound 279),
15 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-pyridin-3-yl-ethoxy)-phenyl]-ethanone (compound 280),
2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(4-methanesulfinyl-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 281),
2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(4-methanesulfonyl-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 282),
20 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(1-phenyl-propoxy)-phenyl]-ethanone (compound 283),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-phenyl-propoxy)-phenyl]-ethanone (compound 284),
25 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(1-methyl-2-phenyl-ethoxy)-phenyl]-ethanone (compound 285),
2-{6-[2-(6-Chloro-pyrazin-2-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 286),
2-{6-[2-(3-Bromo-pyrazin-2-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 287),
30 2-{6-[2-(2,6-Dichloro-phenyl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 288),
2-[2,3-Dimethoxy-6-(2-pyridin-4-yl-acetyl)-phenoxy]-N-propyl-acetamide (compound 289),

2-[2,3-Dimethoxy-6-(2-quinolin-4-yl-acetyl)-phenoxy]-N-propyl-acetamide (compound 290),

2-[2,3-Dimethoxy-6-(2-pyrazin-2-yl-acetyl)-phenoxy]-N-propyl-acetamide (compound 291),

5 2-{6-[2-(3-Bromo-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 292),

2-{6-[2-(3,5-Dibromo-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 293),

2-{6-[2-(6-Chloro-pyrimidin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 294),

10 2-{6-[2-(4-Chloro-pyridin-2-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 295),

2-{6-[2-(2-Chloro-pyridin-3-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 296),

15 2-{2,3-Dimethoxy-6-[2-(2-methoxy-pyridin-4-yl)-acetyl]-phenoxy}-N-propyl-acetamide (compound 297),

2-{6-[2-(2-Cyano-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 298),

2-[2,3-Dimethoxy-6-(2-pyridazin-3-yl-acetyl)-phenoxy]-N-propyl-acetamide (compound 299),

20 2-(2-tert-Butylamino-3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 300),

2-(2-amino-3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 301),

25 2-(3,5-Dichloro-pyridin-4-yl)-1-(4-ethoxy-3-methoxy-2-phenethoxy-phenyl)-ethanone (compound 302),

{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid (compound 504),

2-tert-butoxycarbonylmethoxy-3,4-dimethoxy-benzoic acid methyl ester (compound 506a),

30 2-carboxymethoxy-3,4-dimethoxy-benzoic acid methyl ester (compound 506b),

3,4-Dimethoxy-2-propylcarbamoylmethoxy-benzoic acid methyl ester (compound 506c), or

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-ethyl-acetamide (compound 305),
and pharmaceutically acceptable and physiologically cleavable esters,
pharmaceutically acceptable salts, hydrates, N-oxides or solvates thereof.

5

In one preferred embodiment of the present invention X_3 represents N, and X_1 , X_2 , X_4 and X_5 represent -CH-. In another embodiment X_1 represents N, and X_2 , X_3 , X_4 and X_5 represent -CH-. In another embodiment X_2 represents N, and X_1 , X_3 , X_4 and X_5 represent -CH-. In another embodiment X_4 represents N, and X_1 , X_2 , X_3 and X_5 represent -CH-, or X_5 represents N, and X_1 , X_2 , X_3 and X_4

10 represent -CH-. In a further preferred embodiment X_1 , X_2 , X_3 , X_4 and X_5 represent -CH-. In yet another embodiment X_1 and X_4 represent N, and X_2 , X_3 and X_5 represent -CH-, or X_2 and X_5 represent N and X_1 , X_3 and X_4 represent -CH-, or X_3 and X_5 represent N and X_1 , X_2 and X_4 represent -CH-. In a further
15 embodiment X_1 and X_2 represent N, and X_3 , X_4 and X_5 represent -CH-, or X_4 and X_5 represent N and X_1 , X_2 and X_3 represent -CH-.

Suitable substituents R_{11} of ring B comprise halogens such as chloro, bromo or
20 fluoro, cyano, methoxy, ethoxy, propoxy or alkylamines such as *tert*-butylamine.

Substituents R_{11} of ring B may preferably be attached in the ortho (2 and 6 position) and/or meta (3 and 5 position) position from where ring B is attached to the acetophenone.

25

In another embodiment when R_3 is hydrogen, then one of X_1 , X_2 , X_3 , X_4 or X_5 is nitrogen.

30 In one or more embodiments of the present invention, the compounds of formula I as defined above are useful in therapy and in particular useful for treatment of dermal diseases.

In one or more embodiments of the present invention, the dermal disease or condition is selected from proliferative and inflammatory skin disorders,

psoriasis, cancer, epidermal inflammation, alopecia, skin atrophy, steroid induced skin atrophy, skin ageing, photo skin ageing, acne, dermatitis, atopic dermatitis, seborrheic dermatitis, contact dermatitis, urticaria, pruritis, and eczema.

5

The compounds of formula I may be obtained in crystalline form either directly by concentration from an organic solvent or by crystallisation or recrystallisation from an organic solvent or mixture of said solvent and a cosolvent that may be organic or inorganic, such as water. The crystals may be isolated in essentially solvent-free form or as a solvate, such as a hydrate. The invention covers all crystalline forms, such as polymorphs and pseudopolymorphs, and also mixtures thereof.

Compounds of formula I may or may not comprise asymmetrically substituted (chiral) carbon atoms which give rise to the existence of isomeric forms, e.g. enantiomers and possibly diastereomers. The present invention relates to all such isomers, either in pure form or as mixtures thereof (e.g. racemates). Pure stereoisomeric forms of the compounds and the intermediates of this invention may be obtained by the application of procedures known in the art. The various isomeric forms may be separated by physical separation methods such as selective crystallization and chromatographic techniques, e.g. liquid chromatography using chiral stationary phases. Enantiomers may be separated from each other by the selective crystallization of their diastereomeric salts with optically active amines, such as L-ephedrine. Alternatively, enantiomers may be separated by chromatographic techniques using chiral stationary phases. Said pure stereoisomeric forms may also be derived from the corresponding pure stereoisomeric forms of the appropriate starting materials, provided that the reaction occurs stereoselectively or stereospecifically. Preferably, if a specific stereoisomer is desired, said compound will be synthesized by stereoselective or stereospecific methods of preparation. These methods will advantageously employ chiral pure starting materials.

Compounds of the present invention, optionally in combination with other active compounds, would be useful for the treatment of dermal diseases or conditions,

or acute or chronic cutaneous wound disorders, in particular for the treatment of proliferative and inflammatory skin disorders, psoriasis, cancer, epidermal inflammation, alopecia, skin atrophy, steroid induced skin atrophy, skin ageing, photo skin ageing, acne, dermatitis, atopic dermatitis, seborrheic dermatitis, 5 contact dermatitis, urticaria, pruritis, and eczema.

Besides being useful for human treatment, the compounds of the present invention may also be useful for veterinary treatment of animals including mammals such as horses, cattle, sheep, pigs, dogs, and cats.

10 For use in therapy, compounds of the present invention are typically in the form of a pharmaceutical composition. The invention therefore relates to a pharmaceutical composition comprising a compound of formula I, optionally together with one or more other therapeutically active compound(s), together 15 with a pharmaceutically acceptable excipient, vehicle or carrier(s). The excipient must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipient thereof.

20 Conveniently, the active ingredient comprises from 0.05-99.9% by weight of the formulation.

In the form of a dosage unit, the compound may be administered one or more times a day at appropriate intervals, always depending, however, on the condition of the patient, and in accordance with the prescription made by the 25 medical practitioner. Conveniently, a dosage unit of a formulation contain between 0.1 mg and 1000 mg, preferably between 1 mg and 100 mg, such as 5-50 mg of a compound of formula I.

A suitable dosage of the compound of the invention will depend, *inter alia*, on the age and condition of the patient, the severity of the disease to be treated 30 and other factors well known to the practising physician. The compound may be administered either orally, parenterally or topically according to different dosing schedules, e.g. daily or with weekly intervals. In general a single dose will be in the range from 0.01 to 400 mg/kg body weight. The compound may be

administered as a bolus (i.e. the entire daily dose is administered at once) or in divided doses two or more times a day.

In the context of topical treatment it may be more appropriate to refer to a

5 "usage unit", which denotes a single dose which is capable of being administered to a patient, and which may be readily handled and packed, remaining as a physically and chemically stable unit dose comprising either the active material as such or a mixture of it with solid or liquid pharmaceutical diluents or carriers.

10 The term "usage unit" in connection with topical use means a unitary, i.e. a single dose, capable of being administered topically to a patient in an application per square centimetre of the infected area of from 0.1 mg to 10 mg and preferably from 0.2 mg to 1 mg of the active ingredient in question.

15 It is also envisaged that in certain treatment regimes, administration with longer intervals, e.g. every other day, every week, or even with longer intervals may be beneficial.

If the treatment involves administration of another therapeutically active

20 compound it is recommended to consult *Goodman & Gilman's The Pharmacological Basis of Therapeutics*, 9th Ed., J.G. Hardman and L.E. Limbird (Eds.), McGraw-Hill 1995, for useful dosages of said compounds.

25 The administration of a compound of the present invention with one or more other active compounds may be either concomitantly or sequentially.

The formulations include e.g. those in a form suitable for oral (including sustained or timed release), rectal, parenteral (including subcutaneous, intraperitoneal, intramuscular, intraarticular and intravenous), transdermal, 30 ophthalmic, topical, nasal or buccal administration.

The formulations may conveniently be presented in dosage unit form and may be prepared by any of the methods well known in the art of pharmacy, e.g. as disclosed in Remington, *The Science and Practice of Pharmacy*, 21ed ed., 2005.

All methods include the step of bringing the active ingredient into association with the carrier, which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing the active ingredient into association with a liquid carrier or a finely divided solid carrier or both, and then, if necessary, shaping the product into the desired formulation.

Formulations of the present invention suitable for oral administration may be in the form of discrete units as capsules, sachets, tablets or lozenges, each 10 containing a predetermined amount of the active ingredient; in the form of a powder or granules; in the form of a solution or a suspension in an aqueous liquid or non-aqueous liquid, such as ethanol or glycerol; or in the form of an oil-in-water emulsion or a water-in-oil emulsion. Such oils may be edible oils, such as e.g. cottonseed oil, sesame oil, coconut oil or peanut oil. Suitable 15 dispersing or suspending agents for aqueous suspensions include synthetic or natural gums such as tragacanth, alginate, acacia, dextran, sodium carboxymethylcellulose, gelatin, methylcellulose, hydroxypropylmethylcellulose, hydroxypropylcellulose, carbomers and polyvinylpyrrolidone. The active ingredients may also be administered in the form of a bolus, electuary or paste.

20 A tablet may be made by compressing or moulding the active ingredient optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient(s) in a free-flowing form such as a powder or granules, optionally mixed by a binder, 25 such as e.g. lactose, glucose, starch, gelatine, acacia gum, tragacanth gum, sodium alginate, carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, polyethylene glycol, waxes or the like; a lubricant such as e.g. sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride or the like; a disintegrating agent 30 such as e.g. starch, methylcellulose, agar, bentonite, croscarmellose sodium, sodium starch glycollate, crospovidone or the like or a dispersing agent, such as polysorbate 80. Moulded tablets may be made by moulding, in a suitable machine, a mixture of the powdered active ingredient and suitable carrier moistened with an inert liquid diluent.

Formulations for rectal administration may be in the form of suppositories in which the compound of the present invention is admixed with low melting water soluble or insoluble solids such as cocoa butter, hydrogenated vegetable oils, 5 polyethylene glycol or fatty acids esters of polyethylene glycols, while elixirs may be prepared using myristyl palmitate.

Formulations suitable for parenteral administration conveniently comprise a sterile oily or aqueous preparation of the active ingredients, which is preferably 10 isotonic with the blood of the recipient, e.g. isotonic saline, isotonic glucose solution or buffer solution. The formulation may be conveniently sterilised by for instance filtration through a bacteria retaining filter, addition of sterilising agent to the formulation, irradiation of the formulation or heating of the formulation. Liposomal formulations as disclosed in e.g. *Encyclopedia of Pharmaceutical 15 Technology*, vol.9, 1994, are also suitable for parenteral administration.

Alternatively, the compounds of formula I may be presented as a sterile, solid preparation, e.g. a freeze-dried powder, which is readily dissolved in a sterile solvent immediately prior to use.

20

Transdermal formulations may be in the form of a plaster or a patch.

Formulations suitable ophthalmic administration may be in the form of a sterile aqueous preparation of the active ingredients, which may be in microcrystalline 25 form, for example, in the form of an aqueous microcrystalline suspension. Liposomal formulations or biodegradable polymer systems e.g. as disclosed in *Encyclopedia of Pharmaceutical Technology*, vol.2, 1989, may also be used to present the active ingredient for ophthalmic administration.

30 Formulations suitable for topical or ophthalmic administration include liquid or semi-liquid preparations such as liniments, lotions, gels, applicants, oil-in-water or water-in-oil emulsions such as creams, ointments or pastes; or solutions or suspensions such as drops. Compositions for ophthalmic treatment may preferably additionally contain a cyclodextrin.

For topical administration, the compound of formula I may typically be present in an amount of from 0.01 to 20% by weight of the composition, such as 0.1% to about 10 %, but may also be present in an amount of up to about 50% of the 5 composition.

Formulations suitable for nasal or buccal administration include powder, self-propelling and spray formulations, such as aerosols and atomisers. Such formulations are disclosed in greater detail in e.g. Modern Pharmaceutics, 2nd 10 ed., G.S. Banker and C.T. Rhodes (Eds.), page 427-432, Marcel Dekker, New York; Modern Pharmaceutics, 3rd ed., G.S. Banker and C.T. Rhodes (Eds.), page 618-619 and 718-721, Marcel Dekker, New York and Encyclopedia of Pharmaceutical Technology, vol. 10, J. Swarbrick and J.C. Boylan (Eds), page 191-221, Marcel Dekker, New York.

15 In addition to the aforementioned ingredients, the formulations of a compound of formula I may include one or more additional ingredients such as diluents, buffers, flavouring agents, colourant, surface active agents, thickeners, preservatives, e.g. methyl hydroxybenzoate (including anti-oxidants), 20 emulsifying agents and the like.

When the active ingredient is administered in the form of salts with pharmaceutically acceptable non-toxic acids or bases, preferred salts are for instance easily water-soluble or slightly soluble in water, in order to obtain a particular and 25 appropriate rate of absorption.

The pharmaceutical composition may additionally comprise one or more other active components conventionally used in the treatment of dermal disease or conditions, e.g. selected from the group consisting of proliferative and 30 inflammatory skin disorders, psoriasis, cancer, epidermal inflammation, alopecia, skin atrophy, steroid induced skin atrophy, skin ageing, photo skin ageing, acne, dermatitis, atopic dermatitis, seborrheic dermatitis, contact dermatitis, urticaria, pruritis, and eczema.

Examples of such additional active components may for example be selected from the group consisting of glucocorticoids, vitamin D and vitamin D analogues, antihistamines, platelet activating factor (PAF) antagonists, anticholinergic agents, methylxanthines, β -adrenergic agents, COX-2 inhibitors, salicylates, 5 indomethacin, flufenamate, naproxen, timogadine, gold salts, penicillamine, serum cholesterol lowering agents, retinoids, zinc salts and salicylazosulfapyridine.

METHODS OF PREPARATION

10 The compounds of the present invention can be prepared in a number of ways well known to those skilled in the art of synthesis. The compounds of formula I may for example be prepared using the reactions and techniques outlined below together with methods known in the art of synthetic organic chemistry, or variations thereof as appreciated by those skilled in the art. Preferred methods 15 include, but are not limited to, those described below. The reactions are carried out in solvents appropriate to the reagents and materials employed and suitable for the transformations being effected. Also, in the synthetic methods described below, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of 20 experiment and work-up procedures, are chosen to be conditions of standard for that reaction, which should be readily recognized by one skilled in the art. Not all compounds falling into a given class may be compatible with some of the reaction conditions required in some of the methods described. Such restrictions to the substituents which are compatible with the reaction conditions will be 25 readily apparent to one skilled in the art and alternative methods can be used. The compounds of the present invention or any intermediate may be purified if required using standard methods well known to a synthetic organic chemist, e.g. methods described in "Purification of Laboratory Chemicals", 5th ed. 2003. Starting materials are either known compounds, commercially available, or they 30 may be prepared by routine synthetic methods well known to a person skilled in the art.

GENERAL PROCEDURES, PREPARATIONS AND EXAMPLES

¹H nuclear magnetic resonance (NMR) spectra were usually recorded at 300 MHz. Chemical shift values (δ , in ppm) are quoted in the specified solvent relative to internal tetramethylsilane (δ = 0.00) or chloroform (δ = 7.25) standards. The value of a multiplet, either defined (doublet (d), triplet (t), quartet (q)) or not (m) at the approximate mid point is given unless a range is quoted. (bs) indicates a broad singlet. The organic solvents used were usually anhydrous. Chromatography was performed on Merck silica gel 60 (0.040 - 0-063 mm). The solvent ratios indicated refer to v:v unless otherwise noted.

10 The following abbreviations have been used throughout:

ADDP	1,1'--(Azodicarbonyl)dipiperidine
AIBN	Azobisisobutyronitrile
Ar-Me	aryl-methyl
DCE	dichloroethane
15 DCM	dichloromethane
DMF	N,N'-Dimethylformamide
DMSO	dimethyl sulfoxide
Et	ethyl
h	hour(s)
20 HATU	O-(7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HetAr-Me	heteroaryl-methyl
L	litre
LDA	lithium diisopropylamide
25 LiHMDS	lithium hexamethyldisilazide
m	milli
MCPBA	m-chloroperoxybenzoic acid
Me	methyl
NBS	N-bromosuccinimide
30 NMP	1-methyl-2-pyrrolidinone
NMR	nuclear magnetic resonance
rt	room temperature
RT	retention Time
TBAD	di-tert-butyl-azodicarboxylate

TFA	trifluoroacetic acid
THF	tetrahydrofuran
TIS	triethylsilane
v	volume

5

Preparative HPLC/MS

Preparative HPLC/MS was performed on a Dionex APS-system with two Shimadzu PP150 prep. pumps and a Thermo MSQ Plus mass spectrometer. Column: Waters XTerra C-18, 150 mm x 19 mm, 5 µm; solventsystem: A = water (0.1 % formic acid) and B = acetonitrile (0.1 % formic acid); flow rate = 18 mL/min; method (10 min): Linear gradient method going from 10 % B to 100 % B in 6 minutes and staying at 100 % B for another 2 minutes. The fractions were collected based on ion traces of relevant ions and PDA signal (240-400 nm).

15

Analytical HPLC/MS

Method A:

Analytical HPLC/MS was performed on a Dionex APS-system with a P680A analytical pump and a Thermo MSQ Plus mass spectrometer. Column: Waters 20 XTerra C-18, 150 mm x 4.6 mm, 5 µm; solventsystem: A = water (0.1 % formic acid) and B = acetonitrile (0.1 % formic acid); flow rate = 1.0 mL/min; method (10 min): Linear gradient method going from 10 % B to 100 % B in 6.6 minutes and staying at 100 % B for another 1.5 minutes.

Method B:

25 Analytical HPLC/MS was performed on a system consisting of a Waters 2795 HPLC, Micromass ZQ mass spectrometer, Waters 996 PDA. Column: Waters XTerra C-18, 50 mm x 3.0 mm, 5 µm; solventsystem: A = water:acetonitrile 95:5 (0.05 % formic acid) and B = acetonitrile (0.05 % formic acid); flow rate = 1.0 mL/min; method (8 min): Linear gradient method going from 10 % B to 100 30 % B in 6.0 minutes and staying at 100 % B for 1 minute.


General Methods and Examples:

The compounds of the invention may for example be prepared according to the following non-limiting general methods and examples:

Method a)

Mono or dialkylation of compounds with formula Ia

5




wherein R1 is as described herein, with suitable alkyl or alkenyl chlorides, bromides, iodides, mesylates or tosylates under basic conditions (Protective

10 Groups in Organic Chemistry, John Wiley & sons, Ed: T. Greene and P. G. Wuts, 3rd edition (1999), p 249-72), or by suitable alcohols, e.g. alkyl alcohols, using Mitsunobu conditions (see *Synthesis* (1981), 1; *Tet. Lett.* (1993), **34**, 1639-42 and *Eur. J. Org. Chem.* (2004), 2763-72), in a suitable solvent such as THF, benzene or DMF. The reaction may either be one-pot or in two subsequent

15 steps.

Method b)

Alkylation of compounds with formula Ic,




wherein R1 and R3 (R3 ≠ H) are as described herein, with alkyl, alkenyl or alkynyl chlorides, bromides, iodides, mesylates or tosylates under basic conditions or with alkyl alcohols using Mitsunobu conditions in a suitable solvent such as THF, acetone, benzene, toluene or DMF.

5

Method c)

Alkylation of compounds with formula Id (I where R3 = hydrogen),




wherein R1 and R2 (R2 ≠ H) are as described herein, with alkyl, alkenyl or alkynyl chlorides, bromides, iodides, mesylates or tosylates under basic conditions or by alkyl alcohols using Mitsunobu conditions in a suitable solvent such as THF, acetone, benzene, toluene or DMF.

15

Method d)

Amide formation of compounds with formula Ie,



wherein R1 and R2 are as described herein, with primary or secondary amines

5 using for example a coupling reagent such as HATU (Carpino, L. A. *J. Am. Chem. Soc.* (1993), **115**, 4397), or any other standard peptide coupling reagents (Larry Yet, Albany Molecular Research, Technical Reports Vol 4, number 1, p 1-7).

10 Starting materials of formula Ia-Ie may be prepared according to standard procedures well known to a person skilled in the art. E.g. commercially available 2,3,4-Methoxy benzoic acid (Aldrich) is esterified with MeI in the presence of a suitable base, such as K_2CO_3 or Et_3N in a suitable solvent such as DMF, THF or DCM at temperatures from rt to 100°C. The resulting ester is then condensed 15 with 3,5-dichloro-4-methylpyridine (e.g. prepared according to either *J. Org. Chem.* (1961), **26**, 789-92, *Heterocycles* (2001), **55**, 2075-84 or PCT9414742) in the presence of a suitable base, such as LDA or LiHMDS, in a suitable solvent such as THF, at temperatures from minus 78°C to rt. The resulting ketone is deprotected either:

20 A) In the presence of a suitable acid, such as HI or HBr in a suitable solvent such as AcOH at temperatures from 50°C to 120°C to provide compounds of formula Ia.

B) In the presence of a suitable Lewis acid, such as BCl_3 , BBr_3 or $AlCl_3$ in a suitable solvent, such as dichloromethane at temperatures from 0° to rt 25 to provide compounds of formula Id.

Dialkylation of compounds with formula Ia with alkenyl chlorides, bromides, iodides, mesylates or tosylates in the presence of a suitable base, such as K_2CO_3 or Et_3N in a suitable solvent such as DMF, NMP, THF or DCM at temperatures from rt to 100°C provide compounds of formula Ib. Subsequent deprotection of

5 compounds with formula Ib in the presence of a suitable Lewis acid, such as BCl_3 , BBr_3 or $AlCl_3$ in a suitable solvent, such as dichloromethane at temperatures from 0° to rt, followed by re-alkylation with alkenyl chlorides, bromides, iodides, mesylates or tosylates in the presence of a suitable base, such as K_2CO_3 or Et_3N in a suitable solvent such as DMF, NMP, THF or DCM at

10 temperatures from rt to 100°C provide unsymmetrical compounds of formula Ib.


Selective monoalkylation of compounds with formula Ia with alkyl, alkenyl or alkynyl bromides or iodides in the presence of a suitable base, such as K_2CO_3 in a suitable solvent such as DMF or NMP at temperatures from rt to 100°C provide compounds of formula Ic or Id.

15 Alkylation of compounds with formula Id with alkyl chloro or bromoacetate in the presence of suitable base, such as K_2CO_3 in a suitable solvent such as DMF or NMP at temperatures from rt to 100°C followed by hydrolysis of the carboxylic ester under standard conditions (Protective Groups in Organic Chemistry, John Wiley & sons, Ed: T. Greene and P. G. Wuts, 3rd edition (1999), p 384-86)

20 provide compounds of formula Ie.

Preparation 1 (compound 501)

Methyl 2,3,4-trimethoxybenzoate




2,3,4-Trimethoxybenzoic acid (Aldrich) (20g, 94 mmol) was dissolved in dry DMF (250 mL). K_2CO_3 (13g, 94 mmol) and MeI (6.5 mL, 103.4 mmol) were added and the reaction mixture was heated to 50°C for 5h. The reaction mixture

30 was cooled to rt and water (250 mL) was added. EtOAc (1L) was added and the

organic phase was washed with water (3x 500 mL) and finally with NaCl (sat.). The organic phase washed dried over MgSO₄, filtered and concentrated *in vacuo*. Redissolved in toluene and evaporated. Methyl 2,3,4-trimethoxybenzoate was obtained as a slightly yellow oil. ¹H NMR (CDCl₃) δ = 7.60 (1H, d), 6.70 (1H, d), 3.99 (3H, s), 3.91 (3H, s), 3.89 (3H, s), 3.88 (3H, s). Yield 21g (99%).

Preparation 2 (compound 502)

2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3,4-trimethoxy-phenyl)-ethanone




10

Methyl 2,3,4-trimethoxybenzoate from preparation 1 (2.7g, 10 mmol) and 3,5-dichloro-4-methylpyridine (prepared according to literature procedures) was dissolved in dry THF (40 mL). The reaction was cooled on ice and treated dropwise (during 5 minutes) with 1M lithium bis(trimethylsilyl)amide (12 mL, 12 mmol). After 1.5 hours at 0°C the reaction was quenched with NH₄Cl (sat, 100 mL). The organic products were extracted with EtOAc (2x100 mL), and the combined organic phases washed with NaCl (sat, 100 mL). The organic phase was dried over Na₂SO₄, evaporated *in vacuo* and purified by flash chromatography using a gradient of EtOAc in heptane as eluent. 2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3,4-trimethoxy-phenyl)-ethanone was obtained as a white solid. ¹H NMR (CDCl₃) δ = 8.43 (2H, s), 7.55 (1H, d), 6.69 (1H, d), 4.61 (2H, s), 4.02 (3H, s), 3.87 (3H, s), 3.84 (3H, s). Yield 2.3g (65%).

Example 1 (compound 101)


25 2-(3,5-Dichloro-pyridin-4-yl)-1-(2-hydroxy-3,4-dimethoxy-phenyl)-ethanone



2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3,4-trimethoxy-phenyl)-ethanone from preparation 2 (20g, 56 mmol) was dissolved in DCM (75 mL) under argon. BCl_3 (95 mL of a 1M solution in DCM) was added slowly at room temperature for 2 hours. Water (50 mL) was added slowly followed by EtOH (200 mL). The white precipitate was filtered and re-crystallised from EtOH. 2-(3,5-Dichloro-pyridin-4-yl)-1-(2-hydroxy-3,4-dimethoxy-phenyl)-ethanone was obtained as a white solid. ^1H NMR (CDCl_3) δ = 11.92 (1H, s), 8.54 (2H, s), 7.68 (1H, d), 6.60 (1H, d), 4.67 (2H, s), 3.98 (3H, s), 3.90 (3H, s). Yield 14.2g (74%)

Preparation 3 (compound 503)

2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3-dihydroxy-4-methoxy-phenyl)-ethanone




2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3,4-trimethoxy-phenyl)-ethanone from preparation 2 (1.25g, 3.5 mmol) was dissolved in AcOH (100%, 9.25 mL) under argon. HI (55-58%, 4.55 mL) was added and the reaction mixture was heated at

80°C for 18 hours. The reaction mixture was cooled to rt and evaporated *in vacuo*. NaHCO₃ (sat., 150 mL) was added slowly followed by NaCl (sat., 50 mL). The organic products were extracted with EtOAc (4x100 mL). The combined organic phases were washed twice with NaCl (sat, 100 mL), Na₂S₂O₃ (10%, 100 mL), dried over Na₂SO₄ and evaporated *in vacuo*. 2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3-dihydroxy-4-methoxy-phenyl)-ethanone was obtained as a brown solid.

5 ¹H NMR (CDCl₃) δ = 11.88 (1H, s), 8.54 (2H, s), 7.51 (1H, d), 6.60 (1H, d), 5.62 (1H, bs), 4.67 (2H, s), 4.00 (3H, s). Yield 0.935g (81%)

10 General procedure for preparation of compounds with formula If, wherein R3 is as defined herein (R3 ≠ hydrogen):



2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3-dihydroxy-4-methoxy-phenyl)-ethanone
 15 (0.1 mmol) from preparation 3 was dissolved in dry DMF (0.75 mL). K₂CO₃ (0.1 mmol) was added followed by 0.1 mmol of an alkyl bromide or iodide. The reaction mixture was heated to 70°C for 18 hours. After cooling to rt, water (1 mL) was added and the organic products were extracted with EtOAc (3x1mL). The combined organic phases were washed with NaCl (sat, 1mL) and then dried over Na₂SO₄. The pure compounds were obtained by redissolving the reaction mixture in DMSO followed by standard preparative HPLC purification.

20

Using this procedure the following compounds were obtained:

25 Example 2 (compound 102)

2-(3,5-Dichloro-pyridin-4-yl)-1-(3-hydroxy-2,4-dimethoxy-phenyl)-ethanone

LC/MS (METHOD B): (m/z) 342.1 (MH⁺); RT = 3.34 min; purity (UV) = 100%


Alkyl halide: Methyl iodide

Example 3 (compound 103)

5 1-(2-Allyloxy-3-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone LC/MS (METHOD B): (m/z) 368.2 (MH⁺); RT = 3.81 min ; purity (UV) = 100%

Alkyl halide: Allyl bromide

10 General procedure for preparation of compounds with formula Ig, wherein R2 = R3 and are as defined herein (R3 ≠ hydrogen):



15 2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3-dihydroxy-4-methoxy-phenyl)-ethanone (0.1 mmol) from preparation 3 was dissolved in dry DMF (0.75 mL). K₂CO₃ (0.2 mmol) was added followed by 0.2 mmol of an alkyl chloride, bromide or iodide. In the case of the alkyl chlorides additional KI (0.05 mmol) was added. The reaction mixture was heated to 70°C for 18 hours. After cooling to rt, water (1 mL) was added and the organic products were extracted with EtOAc (3x1 mL). The combined organic phases were washed with NaCl (sat, 1mL) and then dried over Na₂SO₄. The pure compounds were obtained by redissolving the reaction mixture in DMSO followed by standard preparative HPLC purification.

20

25 Using this procedure the following compounds were obtained:

Example 4 (compound 104)

2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3-diethoxy-4-methoxy-phenyl)-ethanone

LC/MS (METHOD B): (m/z) 384.1 (MH⁺); RT = 4.69 min; purity (UV) = 100%

Alkyl halide: Ethyl iodide

5

Example 5 (compound 105)

{2-tert-Butoxycarbonylmethoxy-6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-3-methoxy-phenoxy}-acetic acid tert-butyl ester

LC/MS (METHOD B): (m/z) 554.3 (MH⁺); RT = 5.39 min; purity (UV) = 100%

10 Alkyl halide: Tert-butyl bromoacetate

Example 6 (compound 106)

1-(2,3-Bis-allyloxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone

LC/MS (METHOD B): (m/z) 408.2 (MH⁺); RT = 4.94 min; purity (UV) = 100%

15 Alkyl halide: Allyl bromide

Example 107 (compound 107)


1-(2,3-Bis-difluoromethoxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone LC/MS (METHOD B): (m/z) 428.1 (MH⁺); RT = 4.23 min; purity (UV)

20 = 100%

Alkyl halide: Sodium chlorodifluoroacetate. Heated to 100°C for 30 minutes.

General procedure for preparation of compounds with formula Ih, wherein R3 (R3 ≠ hydrogen) is as defined above:

25



2-(3,5-Dichloro-pyridin-4-yl)-1-(2-hydroxy-3,4-methoxy-phenyl)-ethanone (0.035 mmol) from example 1 was dissolved in dry DMSO (0.25 mL). Aqueous K_2CO_3 (0.025 mL of a 2M) was added followed by 0.053 mmol of an alkyl chloride, bromide or iodide dissolved in 0.025 mL DMSO. The reaction mixture was left at rt for 48 hours. The pure compounds were obtained by standard preparative HPLC purification.

Using this procedure the following compounds were obtained:

10

Example 108 (compound 108)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(5-methyl-isoxazol-3-ylmethoxy)-phenyl]-ethanone

LC/MS (METHOD B): (m/z) 437.1 (MH⁺); RT = 4.01 min; purity (UV) = 100%

15 Alkyl halide: 3-Chloromethyl-5-methylisoxazole

Example 109 (compound 109)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methoxy-ethoxy)-phenyl]-ethanone

20 LC/MS (METHOD B): (m/z) 400.1 (MH⁺); RT = 3.90 min; purity (UV) = 100%

Alkyl halide: 1-Bromo-2-methoxy-ethane

Example 110 (compound 110)

1-(2-But-2-ynloxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-

25 ethanone LC/MS (METHOD B): (m/z) 394.1 (MH⁺); RT = 4.30 min; purity (UV) = 100%

Alkyl halide: 1-Bromo-2-butyne

Example 111 (compound 111)

30 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methyl-but-2-enyloxy)-phenyl]-ethanone

LC/MS (METHOD B): (m/z) 410.1 (MH⁺); RT = 4.83 min; purity (UV) = 100%

Alkyl halide: 1-Bromo-2-methyl-2-butene

Example 112 (compound 112)

2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-phenethyloxy-phenyl)-ethanone

¹H NMR (CDCl₃) δ = 8.48 (2H, s), 7.61 (1H, d), 7.31-7.08 (5H, m), 6.75 (1H, d),

5 4.50-4.46 (4H, m), 3.92 (3H, s), 3.77 (3H, s), 3.18 (2H, t).

Alkyl halide: 1-Bromo-2-phenyl-ethane

Example 113 (compound 113)

1-(2-Benzyl-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone

10 LC/MS (METHOD B): (m/z) 432.1 (MH⁺); RT = 4.70 min; purity (UV) = 100%

Alkyl halide: Benzyl bromide

Example 114 (compound 114)

1-(2-Allyloxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone

15 LC/MS (METHOD B): (m/z) 382 (MH⁺); RT = 4.34 min; purity (UV) = 100%

Alkyl halide: Allyl bromide

Example 115 (compound 115)

1-[2-(Benzoxazol-2-ylmethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-

20 pyridin-4-yl)-ethanone

LC/MS (METHOD B): (m/z) 473 (MH⁺); RT = 4.36 min; purity (UV) = 100%

Alkyl halide: 2-(chloromethyl)-1,3-benzoxazole

Example 116 (compound 116)

25 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methyl-thiazol-4-ylmethoxy)-phenyl]-ethanone

LC/MS (METHOD B): (m/z) 453 (MH⁺); RT = 4.02 min; purity (UV) = 100%

Alkyl halide: 4-(chloromethyl)-2-methylthiazole

30 Example 117 (compound 117)

1-[2-(5-Cyclopropyl-[1,3,4]thiadiazol-2-ylmethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone

LC/MS (METHOD B): (m/z) 480.1 (MH⁺); RT = 3.86 min; purity (UV) = 100%

Alkyl halide: 2-(chloromethyl)-5-cyclopropyl-[1,3,4]thiadiazole

Example 118 (compound 118)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-([1,2,4]oxadiazol-3-ylmethoxy)-phenyl]-ethanone

5 LC/MS (METHOD B): (m/z) 424 (MH+); RT = 3.65 min; purity (UV) = 100%
Alkyl halide: 3-(Chloromethyl)-[1,2,4]oxadiazole

Example 119 (compound 119)

{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid

10 ethyl ester
¹H NMR (CDCl₃) δ = 8.48 (2H, s), 7.61 (1H, d), 6.76 (1H, d), 4.92 (1H, s), 4.79 (2H, s), 4.25 (2H, q), 3.93 (3H, s), 3.88 (3H, s), 1.27 (3H, t).
Alkyl halide: Ethyl bromoacetate

15 Example 120 (compound 120)

1-{2-[2-(4-Chloro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone

¹H NMR (CDCl₃) δ = 8.49 (2H, s), 7.61 (1H, d), 7.23-7.18 (4H, m), 6.75 (1H, d), 4.45 (2H, t), 4.37 (2H, s), 3.93 (3H, s), 3.77 (3H, s), 3.14 (2H, t).

20 Alkyl halide: 4-Chlorobenzyl chloride

Example 121 (compound 121)

1-[2-(5-Chloro-thiophen-2-ylmethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone

25 LC/MS (METHOD B): (m/z) 474 (MH+); RT = 5.01 min; purity (UV) = 100%
Alkyl halide: 5-Chloro-2-(chloromethyl)-thiophene

Example 122 (compound 122)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-phenoxy-ethoxy)-phenyl]-

30 ethanone ¹H NMR (CDCl₃) δ = 8.48 (2H, s), 7.65 (1H, d), 7.23 (2H, t), 6.92 (1H, t), 6.85 (2H, d), 6.76 (1H, d), 4.78 (2H, s), 4.69-4.63 (2H, m), 4.39-4.33 (2H, m), 3.94 (3H, s), 3.92 (3H, s).

Alkyl halide: 1-Bromo-2-phenoxy-ethane

Example 123 (compound 123)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-p-tolyl-ethoxy)-phenyl]-ethanone LC/MS (METHOD B): (m/z) 460 (MH⁺); RT = 5.29 min; purity (UV) = 100%

5 Alkyl halide: 1-Bromo-2-(4-methylphenyl)-ethane

Example 124 (compound 124)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-phenyl-propoxy)-phenyl]-ethanone ¹H NMR (CDCl₃) δ = 8.49 (2H, s), 7.59 (1H, d), 7.22 (5H, m), 6.75 (1H, d), 4.69 (2H, s), 4.28 (2H, t), 3.93 (3H, s), 3.88 (3H, s), 2.85 (2H, t), 2.21 (2H, m).

Alkyl halide: 1-Bromo-3-phenyl-propane

Example 125 (compound 125)

15 2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(3-methoxy-phenyl)-ethoxy]-phenyl}-ethanone
LC/MS (METHOD B): (m/z) 476.2 (MH⁺); RT = 4.86 min; purity (UV) = 100%
Alkyl halide: 1-Bromo-2-(3-methoxyphenyl)-ethane

20 Example 126 (compound 126)

2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(4-methoxy-phenyl)-ethoxy]-phenyl}-ethanone
LC/MS (METHOD B): (m/z) 476.2 (MH⁺); RT = 4.84 min; purity (UV) = 100%
Alkyl halide: 1-Bromo-2-(4-methoxyphenyl)-ethane

25

Example 127 (compound 127)

1-{2-[2-(3-Bromo-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone
¹H NMR (DMSO-d₆) δ = 8.62 (2H, s), 7.54 (2H, m), 7.26 (3H, m), 6.97 (1H, d), 4.43 (2H, t), 4.36 (2H, s), 3.88 (3H, s), 3.70 (3H, s), 3.14 (2H, t).
Alkyl halide: 1-Bromo-2-(3-bromophenyl)-ethane

Example 128 (compound 128)

2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(2-methoxy-phenyl)-ethoxy]-phenyl}-ethanone

LC/MS (METHOD B): (m/z) 476.2 (MH⁺); RT = 5.08 min; purity (UV) = 100%

Alkyl halide: 1-Bromo-2-(2-methoxyphenyl)-ethane

5

Example 129 (compound 129)

2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone

LC/MS (METHOD B): (m/z) 464.3 (MH⁺); RT = 4.83 min; purity (UV) = 100%

10 Alkyl halide: 1-Bromo-2-(4-fluorophenyl)-ethane

Example 130 (compound 130)

2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(2-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone

15 LC/MS (METHOD B): (m/z) 464.2 (MH⁺); RT = 4.99 min; purity (UV) = 100%

Alkyl halide: 1-Bromo-2-(2-fluorophenyl)-ethane

Example 131 (compound 131)

2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(3,4-dimethoxy-phenyl)-ethoxy]-3,4-

20 dimethoxy-phenyl}-ethanone

¹H NMR (DMSO-d₆) δ = 8.60 (2H, s), 7.52 (1H, d), 6.95 (1H, d), 6.89 (1H, d),
6.81 (1H, d), 6.73 (1H, d), 4.43 (2H, t), 4.26 (2H, s), 3.88 (3H, s), 3.76 (3H, s),
3.68 (3H, s), 3.55 (3H, s), 3.06 (2H, t).

Alkyl halide: 1-Bromo-2-(3,4-dimethoxyphenyl)-ethane

25

Example 132 (compound 132)

{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid

benzyl ester

LC/MS (METHOD B): (m/z) 490.2 (MH⁺); RT = 4.66 min; purity (UV) = 100%

30 Alkyl halide: Benzyl bromoacetate

Example 133 (compound 133)

{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid
isopropyl ester

LC/MS (METHOD B): (m/z) 442.2 (MH+); RT = 4.44 min; purity (UV) = 100%
Alkyl halide: Isopropyl bromoacetate

Example 134 (compound 134)

5 3-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy-methyl}-
benzoic acid methyl ester
LC/MS (METHOD A): (m/z) 489.9 (MH+); RT = 8.14 min; purity (UV) = 100%
Alkyl halide: 3-Chloromethyl benzoic acid methyl ester

10 Example 135 (compound 135)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methyl-butoxy)-phenyl]-
ethanone LC/MS (METHOD A): (m/z) 412 (MH+); RT = 8.99 min; purity (UV) =
100%
Alkyl halide: 1-Bromo-3-methyl-butan

15

Example 136 (compound 136)

2-(3,5-Dichloro-pyridin-4-yl)-1-(2-hexyloxy-3,4-dimethoxy-phenyl)-ethanone
LC/MS (METHOD A): (m/z) 395.9 (MH+); RT = 9.42 min; purity (UV) = 100%
Alkyl halide: 1-Bromo-hexan

20

Example 137 (compound 137)

1-(2-But-3-enyloxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-
ethanone LC/MS (METHOD A): (m/z) 395.9 (MH+); RT = 8.27 min; purity (UV)
= 100%

25 Alkyl halide: 4-Bromo-but-1-ene

Example 138 (compound 138)

2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-pent-4-enyloxy-phenyl)-
ethanone LC/MS (METHOD A): (m/z) 410 (MH+); RT = 8.62 min; purity (UV) =
30 100%

Alkyl halide: 5-Bromo-pent-1-ene

Example 139 (compound 139)

2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-propoxy-phenyl)-ethanone

LC/MS (METHOD A): (m/z) 383.9 (MH+); RT = 8.34 min; purity (UV) = 100%
Alkyl halide: 1-Iodo-propan

Example 140 (compound 140)

5 1-(2-Butoxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone
LC/MS (METHOD A): (m/z) 397.9 (MH+); RT = 8.72 min; purity (UV) = 100%
Alkyl halide: 1-Iodo-butan

Example 141 (compound 141)

10 2-(3,5-Dichloro-pyridin-4-yl)-1-(2-isobutoxy-3,4-dimethoxy-phenyl)-ethanone
LC/MS (METHOD A): (m/z) 397.9 (MH+); RT = 8.72 min; purity (UV) = 100%
Alkyl halide: 1-Iodo-2-methyl-propane

Example 142 (compound 142)

15 4-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-butyric
acid ethyl ester
LC/MS (METHOD A): (m/z) 456 (MH+); RT = 7.89 min; purity (UV) = 100%
Alkyl halide: 4-Bromo-butyric acid ethyl ester

20 Example 143 (compound 143)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(4-methyl-benzyloxy)-
phenyl]-ethanone
LC/MS (METHOD A): (m/z) 445.9 (MH+); RT = 8.69 min; purity (UV) = 97.1%
Alkyl halide: 4-Methyl-benzylchloride

25

Example 144 (compound 144)

1-[2-(3-Chloro-benzyloxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-
ethanone
LC/MS (METHOD A): (m/z) 465.9 (MH+); RT = 8.69 min; purity (UV) = 100%
30 Alkyl halide: 3-Chloro-benzylchloride

Example 145 (compound 145)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-phenoxy-propoxy)-phenyl]-
ethanone

LC/MS (METHOD A): (m/z) 476 (MH+); RT = 8.59 min; purity (UV) = 98.2%
Alkyl halide: (3-Bromo-propoxy)-benzene

Example 146 (compound 146)

5 2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(4-methoxy-phenyl)-2-oxo-ethoxy]-phenyl}-ethanone

LC/MS (METHOD A): (m/z) 490 (MH+); RT = 7.74 min; purity (UV) = 97.1%
Alkyl halide: 2-Bromo-1-(4-methoxy-phenyl)-ethanone

10 Example 147 (compound 147)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxyethyl}-benzonitrile LC/MS (METHOD A): (m/z) 456.9 (MH+); RT = 7.81 min; purity (UV) = 97.1%. ^1H NMR (DMSO-d₆) δ = 8.60 (2H, s), 7.91 (1H, d), 7.81 (1H, d), 7.78 (1H, t), 7.59 (1H, t), 7.55 (1H, d), 7.02 (1H, d), 5.43 (2H, s), 4.50 (2H, s), 3.92 (3H, s), 3.81 (3H, s).

15 Alkyl halide: 2-Chloromethyl-benzonitrile

Example 148 (compound 148)

20 4-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxyethyl}-benzonitrile LC/MS (METHOD A): (m/z) 456.9 (MH+); RT = 7.82 min; purity (UV) = 98.5%

Alkyl halide: 4-Chloromethyl-benzonitrile

Example 149 (compound 149)

25 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(naphthalen-2-ylmethoxy)-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 481.9 (MH+); RT = 8.86 min; purity (UV) = 77.3%

Alkyl halide: 2-Bromomethyl-naphthalene

30 Example 150 (compound 150)

2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-pentyloxy-phenyl)-ethanone

LC/MS (METHOD A): (m/z) 411.9 (MH+); RT = 9.06 min; purity (UV) = 100%

Alkyl halide: 1-Iodo-pentane

Example 151 (compound 151)

1-(2-Cyclohexylmethoxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone LC/MS (METHOD A): (m/z) 438 (MH⁺); RT = 9.59 min ; purity (UV) = 100%

5 Alkyl halide: Bromomethyl-cyclohexane

Example 152 (compound 152)

3-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy-methyl}-benzonitrile LC/MS (METHOD A): (m/z) 456.9 (MH⁺); RT = 7.81 min; purity (UV) = 96.5%

10 Alkyl halide: 3-Chloromethyl-benzonitrile

Example 153 (compound 153)

1-{2-[2-(4-Chloro-phenoxy)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone

15 LC/MS (METHOD A): (m/z) 495.9 (MH⁺); RT = 8.56 min; purity (UV) = 100%

Alkyl halide: 1-(2-Bromo-ethoxy)-4-chloro-benzene

Example 154 (compound 154)

20 2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-ethyl-butoxy)-3,4-dimethoxy-phenyl]-ethanone LC/MS (METHOD A): (m/z) 426 (MH⁺); RT = 9.39 min; purity (UV) = 92.5%

Alkyl halide: 1-Iodo-2-ethyl-butane

Example 155 (compound 155)

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-hydroxy-ethoxy)-3,4-dimethoxy-phenyl]-ethanone LC/MS (METHOD A): (m/z) 385.9 (MH⁺); RT = 6.22 min; purity (UV) = 88.0%

Alkyl halide: 1-Iodo-2-hydroxy-ethane

30

Example 156 (compound 156)

4-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy-methyl}-benzoic acid methyl ester

LC/MS (METHOD A): (m/z) 489.9 (MH⁺); RT = 8.17 min; purity (UV) = 89.6%

Alkyl halide: 4-Chloromethyl benzoic acid methyl ester

Example 157 (compound 157)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-naphthalen-2-yl-2-oxo-

5 ethoxy)-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 509.9 (MH⁺); RT = 8.41 min; purity (UV) = 89.9%

Alkyl halide: 2-Bromo-1-naphthalen-2-yl-ethanone

Example 158 (compound 158)

10 2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(2,5-dimethoxy-phenyl)-2-oxo-ethoxy]-

3,4-dimethoxy-phenyl}-ethanone

LC/MS (METHOD A): (m/z) 519.9 (MH⁺); RT = 7.99 min; purity (UV) = 95.7%

Alkyl halide: 2-Bromo-2,5-dimethoxy-phenyl-ethanone

15 Example 159 (compound 159)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-oxo-2-p-tolyl-ethoxy)-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 473.9 (MH⁺); RT = 8.11 min; purity (UV) = 88.1%

Alkyl halide: 2-Bromo-1-p-tolyl-ethanone

20

Example 160 (compound 160)

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(4-fluoro-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 449.9 (MH⁺); RT = 8.32 min; purity (UV) = 100%

25 Alkyl halide: 4-Fluoro-benzylchloride

Example 161 (compound 161)

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-fluoro-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone

30 LC/MS (METHOD A): (m/z) 449.9 (MH⁺); RT = 8.34 min; purity (UV) = 95.7%

Alkyl halide: 2-Fluoro-benzylchloride

Example 162 (compound 162)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-trifluoromethyl-benzyloxy)-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 499.9 (MH+); RT = 8.61 min; purity (UV) = 98.9%

Alkyl halide: 5-Trifluoromethyl-benzylchloride

5

Example 163 (compound 163)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-trifluoromethoxy-benzyloxy)-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 515.9 (MH+); RT = 8.72 min; purity (UV) = 98.9%

10 Alkyl halide: 5-Trifluoromethoxy-benzylchloride

Example 164 (compound 164)

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(3-fluoro-5-trifluoromethyl-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone

15 LC/MS (METHOD A): (m/z) 558.9 (MH+); RT = 8.71 min; purity (UV) = 98.9%

Alkyl halide: 3-Fluoro-5-trifluoromethyl-benzylchloride

Example 165 (compound 165)

2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(2-methoxy-phenyl)-2-oxo-ethoxy]-phenyl}-ethanone

20 LC/MS (METHOD A): (m/z) 490 (MH+); RT = 7.96 min; purity (UV) = 95.8%

Alkyl halide: 2-Bromo-1-(2-methoxy-phenyl)-ethanone

Example 166 (compound 166)

25 2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(2,4-dimethyl-phenyl)-2-oxo-ethoxy]-3,4-dimethoxy-phenyl}-ethanone

LC/MS (METHOD A): (m/z) 487.9 (MH+); RT = 8.49 min; purity (UV) = 99.1%

Alkyl halide: 2-Bromo-1-(2,4-dimethyl-phenyl)-ethanone

30 Example 167 (compound 167)

1-[2-(4-Chloro-benzyloxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone

LC/MS (METHOD A): (m/z) 465.9 (MH+); RT = 8.72 min; purity (UV) = 98.5%

Alkyl halide: 4-Chloro-benzylchloride

Example 168 (compound 168)

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-difluoromethoxy-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone

5 LC/MS (METHOD A): (m/z) 497.8 (MH+); RT = 8.19 min; purity (UV) = 98.5%
Alkyl halide: 2-Difluoromethoxy-benzylchloride

Example 169 (compound 169)

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(4-isopropyl-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone

10 LC/MS (METHOD A): (m/z) 474 (MH+); RT = 9.21 min; purity (UV) = 100%
Alkyl halide: 4-Isopropyl-benzylchloride

Example 170 (compound 170)

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-fluoro-6-trifluoromethyl-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 517.9 (MH+); RT = 8.52 min; purity (UV) = 98.9%
Alkyl halide: 2-Fluoro-6-trifluoromethyl-benzylchloride

20 Example 171 (compound 171)

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2,3-difluoro-4-methyl-benzyloxy)-3,4-dimethoxy-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 481.9 (MH+); RT = 8.71 min; purity (UV) = 98.6%
Alkyl halide: 2,3-Difluoro-4-methyl-benzylchloride

25

Example 172 (compound 172)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methyl-benzyloxy)-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 445.9 (MH+); RT = 8.67 min; purity (UV) = 98.5%
30 Alkyl halide: 2-Methyl-benzylchloride

Example 173 (compound 173)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methyl-benzyloxy)-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 446 (MH+); RT = 8.69 min; purity (UV) = 98.4%

Alkyl halide: 3-Methyl-benzylchloride

Example 174 (compound 174)

5 2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-pent-2-enyloxy-phenyl)-ethanone

LC/MS (METHOD A): (m/z) 409.9 (MH+); RT = 8.67 min; purity (UV) = 98.3%

Alkyl halide: 1-Bromo-pent-2-ene

10 Example 175 (compound 175)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methyl-quinolin-6-ylmethoxy)-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 497 (MH+); RT = 5.27 min; purity (UV) = 98.1%

Alkyl halide: 6-Bromomethyl-2-methyl-quinoline

15

Example 176 (compound 176)

1-[2-(2-Chloro-benzylxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone

LC/MS (METHOD A): (m/z) 465.9 (MH+); RT = 8.71 min; purity (UV) = 96.3%

20 Alkyl halide: 2-Chloro-benzylchloride

Example 177 (compound 177)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methoxy-benzylxy)-phenyl]-ethanone

25 LC/MS (METHOD A): (m/z) 461.9 (MH+); RT = 8.26 min; purity (UV) = 96.3%

Alkyl halide: 3-Methoxy-benzylchloride

Example 178 (compound 178)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(4-methoxy-benzylxy)-phenyl]-ethanone

30 LC/MS (METHOD A): (m/z) 461.9 (MH+); RT = 8.22 min; purity (UV) = 77.9%

Alkyl halide: 4-Methoxy-benzylchloride

Example 179 (compound 179)

1-{2-[2-(3-Chloro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone

LC/MS (METHOD A): (m/z) 479.9 (MH⁺); RT = 8.81 min; purity (UV) = 98.6%

Alkyl halide: 1-Bromo-2-(3-Chloro-phenyl)-ethane

5

Example 180 (compound 180)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(5-methyl-hexyloxy)-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 440 (MH⁺); RT = 9.69 min; purity (UV) = 100%

10 Alkyl halide: 1-Bromo-5-methyl-hexane

Example 181 (compound 181)

1-[2-(2-Cyclohexyl-ethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone

15 LC/MS (METHOD A): (m/z) 452 (MH⁺); RT = 9.92 min; purity (UV) = 97.7%

Alkyl halide: (2-Bromo-ethyl)-cyclohexane

Example 182 (compound 182)

5-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-pentanoic

20 acid ethyl ester

LC/MS (METHOD A): (m/z) 470 (MH⁺); RT = 8.11 min; purity (UV) = 100%

Alkyl halide: 4-Bromo-pentanoic acid ethyl ester

Example 183 (compound 183)

25 1-[2-(3-Benzyl-ethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone

LC/MS (METHOD A): (m/z) 490 (MH⁺); RT = 8.64 min; purity (UV) = 97.5%

Alkyl halide: (3-Bromo-propoxymethyl)-benzene

30 Example 184 (compound 184)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

¹H NMR (DMSO-d₆) δ = 8.64 (2H, s), 7.67-7.58 (2H, m), 7.37 (1H, bs), 6.98 (1H, d), 4.79 (2H, s), 4.62 (2H, s), 3.90 (3H, s), 3.80 (3H, s).

Alkyl halide: 2-Chloro-acetamide

Example 185 (compound 185)

2-(2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-ethyl)-isoindole-1,3-dione

5 ¹H NMR (CDCl₃) δ = 8.36 (2H, s), 7.74 (2H, m), 7.60 (3H, m), 6.75 (1H, d), 4.52 (2H, t), 4.48 (2H, s), 4.21 (2H, t), 3.91 (3H, s), 3.83 (3H, s).

Alkyl halide: 2-(2-Bromo-ethyl)-isoindole-1,3-dione

Example 186 (compound 186)

10 2-(3-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-propyl)-isoindole-1,3-dione

1¹H NMR (CDCl₃) δ = 8.48 (2H, s), 7.81 (2H, m), 7.70 (2H, m), 7.59 (1H, d), 6.75 (1H, d), 4.67 (2H, s), 4.31 (2H, t), 3.96 (2H, t), 3.93 (3H, s), 3.89 (3H, s), 2.30 (2H, m).

15 Alkyl halide: 2-(3-Bromo-propyl)-isoindole-1,3-dione

Example 187 (compound 187)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-methyl-acetamide

20 ¹H NMR (CDCl₃) δ = 8.52 (2H, s), 7.75-7.64 (2H, m), 6.79 (1H, d), 4.67 (2H, s), 4.60 (2H, s), 3.96 (3H, s), 3.87 (3H, s), 2.86 (3H, d).

Alkyl halide: 2-Chloro-N-methyl-acetamide


Example 305 (compound 305)

25 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-ethyl-acetamide

1¹H NMR (DMSO-d₆) δ = 8.64 (2H, s), 8.18 (1H, bs), 7.61 (1H, d), 6.99 (1H, d), 4.77 (2H, s), 4.62 (2H, s), 3.91 (3H, s), 3.79 (3H, s), 3.15 (2H, m), 1.01 (3H, t).

30 Alkyl halide: 2-Chloro-N-ethyl-acetamide

General procedure for preparation of compounds with formula II or Ij, wherein R₂ is as defined above:



5 2-(3,5-Dichloro-pyridin-4-yl)-1-(3-hydroxy-2,4-methoxy-phenyl)-ethanone (0.035 mmol) from example 2 (Ii) or 1-(2-allyloxy-3-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (0.035 mmol) from example 3 (Ij) was dissolved in dry 3-pentanone (0.4 mL). Solid K_2CO_3 (0.052 mmol) was added followed by 0.052 mmol of an alkyl bromide or iodide dissolved in 3-pentanone (0.1 mL). In the case of the alkyl bromides additional KI (0.5 eq) was added. The reaction mixture was heated to 70°C for 18 hours. The samples were filtered in the solvent removed *in vacuo*. The pure compounds were obtained by redissolving the reaction mixture in DMSO followed by standard preparative HPLC purification.

10

Using this procedure the following compounds were obtained:

Example 188 (compound 188)

2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-2,4-dimethoxy-phenyl)-ethanone

LC/MS (METHOD B): (m/z) 370 (MH+); RT = 4.45 min; purity (UV) = 100%

Alkyl halide: Iodoethane

Example 189 (compound 189)

5 1-(3-Cyclopropylmethoxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone LC/MS (METHOD B): (m/z) 396.1 (MH+); RT = 4.78 min; purity (UV) = 100%

Alkyl halide: Bromomethyl-cyclopropane

10 Example 190 (compound 190)

1-(2-Allyloxy-3-but-3-enyloxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone

LC/MS (METHOD B): (m/z) 422.1 (MH+); RT = 5.27 min; purity (UV) = 100%

Alkyl halide: 4-Bromo-but-1-ene

15

Example 191 (compound 191)

1-(3-But-3-enyloxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone LC/MS (METHOD A): (m/z) 396.3 (MH+); RT = 8.39 min; purity (UV) = 94.0%

20 Alkyl halide: 4-Bromo-but-1-ene

Example 192 (compound 192)

2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-propoxy-phenyl)-ethanone

LC/MS (METHOD A): (m/z) 384.3 (MH+); RT = 8.42 min; purity (UV) = 100%

25 Alkyl halide: 1-Iodoethane

Example 193 (compound 193)

1-(3-Allyloxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone

LC/MS (METHOD A): (m/z) 382.3 (MH+); RT = 8.01 min; purity (UV) = 100%

30 Alkyl halide: Allyl bromide

Example 194 (compound 194)

2-(3,5-Dichloro-pyridin-4-yl)-1-[2,4-dimethoxy-3-(4-methyl-pent-3-enyloxy)-phenyl]-ethanone

LC/MS (METHOD A): (m/z) 424.3 (MH+); RT = 8.99 min; purity (UV) = 100%
Alkyl halide: 5-Bromo-2-methyl-pent-2-ene

Example 195 (compound 195)

5 2-(3,5-Dichloro-pyridin-4-yl)-1-[3-(2-hydroxy-ethoxy)-2,4-dimethoxy-phenyl]-ethanone LC/MS (METHOD A): (m/z) 386.3 (MH+); RT = 6.21 min; purity (UV) = 70.5%

Alkyl halide: 2-Iodo-ethanol

10 Example 196 (compound 196)

2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-phenethyloxy-phenyl)-ethanone LC/MS (METHOD A): (m/z) 446.2 (MH+); RT = 8.71 min ; purity (UV) = 82.0%

Alkyl halide: 2-Phenyl-1-bromo-ethane

15

Example 197 (compound 197)

1-(3-Benzyl-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone

LC/MS (METHOD A): (m/z) 432.2 (MH+); RT = 8.49 min; purity (UV) = 96.3%

Alkyl halide: Benzyl bromide

20

Example 198 (compound 198)

2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-pent-2-enyloxy-phenyl)-ethanone LC/MS (METHOD A): (m/z) 410.3 (MH+); RT = 8.67 min ; purity (UV) = 91.9%

25 Alkyl halide: 1-Bromo-pent-2-ene

Example 199 (compound 199)

2-(3,5-Dichloro-pyridin-4-yl)-1-[2,4-dimethoxy-3-(2-methoxy-ethoxy)-phenyl]-ethanone

30 LC/MS (METHOD A): (m/z) 400.3 (MH+); RT = 7.34 min; purity (UV) = 91.6%

Alkyl halide: 1-Bromo-2-methoxy-ethane

Example 200 (compound 200)

1-(3-But-2-ynyloxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone LC/MS (METHOD A): (m/z) 394.3 (MH⁺); RT = 7.81 min; purity (UV) = 100%

Alkyl halide: 1-Bromo-but-2-yne

5

Example 201 (compound 201)


2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-prop-2-ynyloxy-phenyl)-ethanone LC/MS (METHOD A): (m/z) 380.2 (MH⁺); RT = 7.44 min; purity (UV) = 100%

10 Alkyl halide: 3-Bromo-propyne

Preparation 4 / Example 273 (compound 504)

{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid


15



2-(3,5-Dichloro-pyridin-4-yl)-1-(2-hydroxy-3,4-methoxy-phenyl)-ethanone (1.37 g, 4.1 mmol) from example 1 was dissolved in dry NMP (50 mL). Ethyl bromoacetate (660 μ L, 6.1 mmol) was added followed by K_2CO_3 (830 mg, 6.1 mmol) and the reaction mixture was stirred at rt overnight. Water (500 mL) was added and the product extracted with EtOAc (2x250 mL). The combined organic phases were washed with water (250 mL), brine (250 mL) and dried over $MgSO_4$. The solvent were removed *in vacuo* and the pure product obtained be standard silica gel chromatography to provide {6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid ethyl ester as a white solid. Yield 990 mg (78%). The compound was re-dissolved in MeOH-water (1:1, 200 mL)

by heating. LiOH (490 mg, 12 mmol, 5 eq.) was added and the reaction mixture was stirred at rt for 1 hour. The reaction mixture was acidified to pH = 1 by addition of 1N HCl (15 mL) and extracted with EtOAc (2x250 mL). The combined organic phases were washed with brine (250 mL) and dried over MgSO₄. The solvent was removed *in vacuo* to provide {6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid as a white solid. ¹H NMR (CDCl₃) δ = 8.52 (2H, s), 7.76 (1H, d), 6.81 (1H, d), 4.85 (2H, s), 4.66 (2H, s), 3.99 (3H, s), 3.88 (3H, s). Yield 0.85g (92%).

10 General procedure for preparation of compounds with formula I_k, wherein R₉ and R₁₂ are as defined herein:



15 {6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid (0.037 mmol) from preparation 4 was dissolved in dry DMF (0.25 mL). A primary or secondary amine (0.045 mmol) dissolved in DMF (0.05 mL) was added followed by HATU (0.045 mmol) dissolved in DMF (0.050 mL). The reaction mixture was left at rt for 48 hours. The pure compounds were obtained by standard preperative HPLC purification.

20

Using this procedure the following compounds were obtained:

Example 202 (compound 202)

25 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-oxo-2-pyrrolidin-1-yl-ethoxy)-phenyl]-ethanone

¹H NMR (CDCl₃) δ = 8.47 (2H, s), 7.64 (1H, d), 6.76 (1H, d), 4.92 (2H, s), 4.88 (2H, s), 3.94 (3H, s), 3.89 (3H, s), 3.53 (2H, t), 3.39 (2H, t), 1.98 (2H, m), 1.86 (2H, m).

Amine: Pyrrolidine

5

Example 203 (compound 203)

N-Benzyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

¹H NMR (CDCl₃) δ = 8.49 (2H, s), 8.12 (1H, bs), 7.66 (1H, d), 7.29-7.20 (5H, m), 6.77 (1H, d), 4.72 (2H, s), 4.55 (2H, s), 4.50 (2H, d), 3.95 (3H, s), 3.81 (3H, s).

Amine: Benzylamine

Example 204 (compound 204)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-morpholin-4-yl-2-oxo-ethoxy)-phenyl]-ethanone

¹H NMR (CDCl₃) δ = 8.48 (2H, s), 7.63 (1H, d), 6.78 (1H, d), 4.96 (2H, s), 4.80 (2H, s), 3.94 (3H, s), 3.89 (3H, s), 3.72-3.61 (6H, m), 3.54-3.45 (2H, m).

Amine: Morpholine

20

Example 205 (compound 205)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-phenyl-acetamide

¹H NMR (CDCl₃) δ = 9.97 (1H, bs), 8.50 (2H, s), 7.75 (1H, d), 7.59 (2H, d), 7.25 (2H, t), 7.07 (1H, t), 6.81 (1H, d), 4.80 (2H, s), 4.65 (2H, s), 3.99 (3H, s), 3.89 (3H, s).

Amine: Aniline

Example 206 (compound 206)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-methyl-N-phenyl-acetamide

¹H NMR (CDCl₃) δ = 8.48 (2H, s), 7.59 (1H, d), 7.45-7.32 (3H, m), 7.25-7.19 (2H, m), 6.70 (1H, d), 4.83 (2H, s), 4.68 (2H, s), 3.90 (3H, s), 3.77 (3H, s), 3.32 (3H, s).

Amine: N-Methyl aniline

Example 207 (compound 207)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-

5 hydroxy-3-methyl-butyl)-acetamide

LC/MS (METHOD A): (m/z) 485 (MH⁺); RT = 5.89 min; purity (UV) = 100%

Amine: 3-Hydroxy-3-methyl-butylamine

Example 208 (compound 208)

10 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-
acetamide

LC/MS (METHOD A): (m/z) 441 (MH⁺); RT = 6.71 min; purity (UV) = 89.3%

¹H NMR (CDCl₃) δ = 8.51 (2H, s), 7.79 (1H, bs), 7.67 (1H, d), 6.78 (1H, d), 4.67
(2H, s), 4.60 (2H, s), 3.97 (3H, s), 3.87 (3H, s), 3.27 (2H, q), 1.50 (2H, m),

15 0.87 (3H, t).

Amine: n-Propylamine

Example 209 (compound 209)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-

20 isopropyl-acetamide

LC/MS (METHOD A): (m/z) 441 (MH⁺); RT = 6.72 min; purity (UV) = 87.7%

¹H NMR (DMSO-d₆) δ = 8.64 (2H, s), 8.01 (1H, d), 7.65 (1H, d), 6.99 (1H, d),
4.77 (2H, s), 4.61 (2H, s), 3.91 (4H, m), 3.80 (3H, s), 1.04 (6H, d)

Amine: Isopropylamine

25

Example 210 (compound 210)

N-Butyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-
acetamide LC/MS (METHOD A): (m/z) 455 (MH⁺); RT = 7.09 min ; purity (UV)
= 100%

30 Amine: Butylamine

Example 211 (compound 211)

N-Cyclopentyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-
phenoxy}-acetamide

LC/MS (METHOD A): (m/z) 466.9 (MH+); RT = 7.19 min; purity (UV) = 100%
Amine: Cyclopentylamine

Example 212 (compound 212)

5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-methyl-butyl)-acetamide

LC/MS (METHOD A): (m/z) 469 (MH+); RT = 7.42 min; purity (UV) = 100%
Amine: 3-Methyl-butylamine

10 Example 213 (compound 213)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-methoxy-benzyl)-acetamide

¹H NMR (DMSO-d₆) δ = 8.67 (1H, t), 8.64 (2H, s), 7.61 (1H, d), 7.15 (2H, d), 6.99 (1H, d), 6.75 (2H, d), 4.75 (2H, s), 4.70 (2H, s), 4.26 (2H, d), 3.90 (3H, s), 3.76 (3H, s), 3.70 (3H, s)

15 Amine: 4-Methoxy-benzylamine

Example 214 (compound 214)

20 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2,2-dimethyl-propyl)-acetamide

LC/MS (METHOD A): (m/z) 469 (MH+); RT = 7.44 min; purity (UV) = 100%
Amine: 2,2-Dimethyl-propylamine

Example 215 (compound 215)

25 N-Cyclohexyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

LC/MS (METHOD A): (m/z) 481 (MH+); RT = 7.52 min; purity (UV) = 100%
Amine: Cyclohexylamine

30 Example 216 (compound 216)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-methoxy-benzyl)-acetamide

¹H NMR (DMSO-d₆) δ = 8.72 (1H, t), 8.63 (2H, s), 7.60 (1H, d), 7.12 (1H, t), 6.99 (1H, d), 6.83-6.74 (3H, m), 4.75 (2H, s), 4.73 (2H, s), 4.30 (2H, d), 3.90 (3H, s), 3.78 (3H, s), 3.69 (3H, s).

Amine: 3-Methoxy-benzylamine

5

Example 217 (compound 217)

N-Cycloheptyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

LC/MS (METHOD A): (m/z) 495 (MH⁺); RT = 7.84 min; purity (UV) = 92.2%

10 Amine: Cycloheptylamine

Example 218 (compound 218)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-methoxy-benzyl)-acetamide

15 ¹H NMR (DMSO-d₆) δ = 8.64 (2H, s), 8.50 (1H, t), 7.62 (1H, d), 7.19 (1H, t), 7.12 (1H, d), 6.99 (1H, d), 6.93 (1H, d), 6.76 (1H, t), 4.78 (2H, s), 4.74 (2H, s), 4.31 (2H, d), 3.91 (3H, s), 3.79 (3H, s), 3.76 (3H, s).

Amine: 2-Methoxy-benzylamine

20 Example 219 (compound 219)

N-Cyclohexylmethyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

LC/MS (METHOD A): (m/z) 495 (MH⁺); RT = 7.86 min; purity (UV) = 100%

Amine: Cyclohexylmethylamine

25

Example 220 (compound 220)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-ethyl)-acetamide

LC/MS (METHOD A): (m/z) 442.9 (MH⁺); RT = 5.41 min; purity (UV) = 100%

30 Amine: 2-Hydroxy-ethylamine

Example 221 (compound 221)

(R)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(1-phenyl-ethyl)-acetamide

LC/MS (METHOD A): (m/z) 502.9 (MH+); RT = 7.37 min; purity (UV) = 100%
Amine: (R)-1-Phenyl-ethylamine

Example 222 (compound 222)

5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-hydroxy-propyl)-acetamide
LC/MS (METHOD A): (m/z) 457 (MH+); RT = 5.51 min; purity (UV) = 100%
Amine: 3-Hydroxy-propylamine

10 Example 223 (compound 223)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-methoxy-ethyl)-acetamide
LC/MS (METHOD A): (m/z) 457 (MH+); RT = 6.12 min; purity (UV) = 100%
Amine: 2-Methoxy-ethylamine

15

Example 224 (compound 224)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-dimethylamino-ethyl)-acetamide
LC/MS (METHOD A): (m/z) 470 (MH+); RT = 4.26 min; purity (UV) = 100%
20 Amine: 2-Dimethylamino-ethylamine

Example 225 (compound 225)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-dimethylamino-propyl)-acetamide
25 LC/MS (METHOD A): (m/z) 484 (MH+); RT = 4.29 min; purity (UV) = 100%
Amine: 3-Dimethylamino-propylamine

Example 226 (compound 226)

(S)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(1-phenyl-ethyl)-acetamide
30 LC/MS (METHOD A): (m/z) 503 (MH+); RT = 7.39 min; purity (UV) = 100%
Amine: (S)-1-Phenyl-ethylamine

Example 227 (compound 227)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-isopropoxy-propyl)-acetamide

LC/MS (METHOD A): (m/z) 498.9 (MH⁺); RT = 6.89 min; purity (UV) = 100%

Amine: 3-Isopropoxy-propylamine

5

Example 228 (compound 228)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-furan-2-ylmethyl-acetamide

LC/MS (METHOD A): (m/z) 478.9 (MH⁺); RT = 6.74 min; purity (UV) = 100%

10 Amine: Furan-2-yl-methylamine

Example 229 (compound 229)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-pyridin-2-ylmethyl-acetamide

15 LC/MS (METHOD A): (m/z) 490 (MH⁺); RT = 5.17 min; purity (UV) = 100%

Amine: Pyridine-2-yl-methylamine

Example 230 (compound 230)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-pyridin-3-ylmethyl-acetamide

20 LC/MS (METHOD A): (m/z) 490 (MH⁺); RT = 4.62 min; purity (UV) = 100%

Amine: Pyridine-3-yl-methylamine

Example 231 (compound 231)

25 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-phenoxy-ethyl)-acetamide

LC/MS (METHOD A): (m/z) 518.9 (MH⁺); RT = 7.19 min; purity (UV) = 98.3%

Amine: 2-Phenoxy-ethylamine

30 Example 232 (compound 232)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-pyridin-4-ylmethyl-acetamide

LC/MS (METHOD A): (m/z) 490 (MH⁺); RT = 4.46 min; purity (UV) = 100%

Amine: Pyridine-4-yl-methylamine

Example 233 (compound 233)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-ethyl-benzyl)-acetamide

5 LC/MS (METHOD A): (m/z) 517 (MH⁺); RT = 7.71 min; purity (UV) = 96.8%
Amine: 4-Ethyl-benzylamine

Example 234 (compound 234)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3,5-difluoro-benzyl)-acetamide

10 LC/MS (METHOD A): (m/z) 524.9 (MH⁺); RT = 7.27 min; purity (UV) = 95.3%
Amine: 3,5-Difluoro-benzylamine

Example 235 (compound 235)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2,3-difluoro-benzyl)-acetamide

LC/MS (METHOD A): (m/z) 524.9 (MH⁺); RT = 7.22 min; purity (UV) = 100%
Amine: 2,3-Difluoro-benzylamine

20 Example 236 (compound 236)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyridin-2-yl-ethyl)-acetamide

LC/MS (METHOD A): (m/z) 503.9 (MH⁺); RT = 4.54 min; purity (UV) = 100%
Amine: 2-Pyridine-2-yl-ethylamine

25

Example 237 (compound 237)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-methyl-benzyl)-acetamide

LC/MS (METHOD A): (m/z) 503 (MH⁺); RT = 7.37 min; purity (UV) = 100%

30 Amine: 2-Methyl-benzylamine

Example 238 (compound 238)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-fluoro-benzyl)-acetamide

LC/MS (METHOD A): (m/z) 506.9 (MH⁺); RT = 7.14 min; purity (UV) = 100%

Amine: 3-Fluoro-benzylamine

Example 239 (compound 239)

5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-methyl-benzyl)-acetamide

LC/MS (METHOD A): (m/z) 503 (MH⁺); RT = 7.39 min; purity (UV) = 100%

Amine: 3-Methyl-benzylamine

10 Example 240 (compound 240)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-methyl-benzyl)-acetamide

LC/MS (METHOD A): (m/z) 502.9 (MH⁺); RT = 7.39 min; purity (UV) = 100%

Amine: 4-Methyl-benzylamine

15

Example 241 (compound 241)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-phenethyl-acetamide

LC/MS (METHOD A): (m/z) 503 (MH⁺); RT = 7.29 min; purity (UV) = 100%

20 Amine: 2-Phenyl-ethylamine

Example 242 (compound 242)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyridin-4-yl-ethyl)-acetamide

25 ¹H NMR (DMSO-d₆) δ = 8.65 (2H, s), 8.44 (2H, d), 8.33 (1H, t), 7.60 (1H, d), 7.30 (2H, d), 6.99 (1H, d), 4.74 (2H, s), 4.62 (2H, s), 3.90 (3H, s), 3.75 (3H, s), 3.44 (2H, q), 2.81 (2H, t).

Amine: 2-Pyridine-4-yl-ethylamine

30 Example 243 (compound 243)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-phenyl-propyl)-acetamide

LC/MS (METHOD A): (m/z) 517 (MH⁺); RT = 7.56 min; purity (UV) = 100%

Amine: 3-Phenyl-propylamine

Example 244 (compound 244)

N-(2-Chloro-benzyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

5 LC/MS (METHOD A): (m/z) 522.9 (MH+); RT = 7.46 min; purity (UV) = 100%
Amine: 2-Chloro-benzylamine

Example 245 (compound 245)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-

10 piperidin-1-yl-ethyl)-acetamide
LC/MS (METHOD A): (m/z) 510 (MH+); RT = 4.46 min; purity (UV) = 92.5%
Amine: 2-Piperidin-1-yl-ethylamine

Example 246 (compound 246)

15 N-(3-Chloro-benzyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

LC/MS (METHOD A): (m/z) 522.8 (MH+); RT = 7.46 min; purity (UV) = 100%
Amine: 3-Chloro-benzylamine

20 Example 247 (compound 247)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-morpholin-4-yl-ethyl)-acetamide

LC/MS (METHOD A): (m/z) 512 (MH+); RT = 4.31 min; purity (UV) = 98.1%
Amine: 2-Morpholin-4-yl-ethylamine

25

Example 248 (compound 248)

N-(4-Chloro-benzyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

LC/MS (METHOD A): (m/z) 522.9 (MH+); RT = 7.49 min; purity (UV) = 98.7%

30 Amine: 4-Chloro-benzylamine

Example 249 (compound 249)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyridin-3-yl-ethyl)-acetamide

LC/MS (METHOD A): (m/z) 504 (MH+); RT = 4.47 min; purity (UV) = 100%
Amine: 2-Pyridine-3-yl-ethylamine

Example 250 (compound 250)

5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyrrolidin-1-yl-ethyl)-acetamide

LC/MS (METHOD A): (m/z) 496 (MH+); RT = 4.37 min; purity (UV) = 100%
Amine: 2-Pyrrolidin-1-yl-ethylamine

10 Example 251 (compound 251)

N-(2-Acetylaminio-ethyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

LC/MS (METHOD A): (m/z) 484 (MH+); RT = 5.31 min; purity (UV) = 100%
Amine: 2-Acetylaminio-ethylamine

15

Example 252 (compound 252)

(R)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-2-phenyl-ethyl)-acetamide

LC/MS (METHOD A): (m/z) 518.9 (MH+); RT = 6.47 min; purity (UV) = 73.4%

20 Amine: (R)-2-Hydroxy-2-phenyl-ethylamine

Example 253 (compound 253)

(S)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-2-phenyl-ethyl)-acetamide

25 LC/MS (METHOD A): (m/z) 519 (MH+); RT = 6.47 min; purity (UV) = 73.8%

Amine: (S)-2-Hydroxy-2-phenyl-ethylamine

Example 254 (compound 254)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-30 thiophen-2-ylmethyl-acetamide

LC/MS (METHOD A): (m/z) 494.9 (MH+); RT = 6.97 min; purity (UV) = 100%

Amine: Thiophen-2-yl-methylamine

Example 255 (compound 255)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-[3-(2-oxo-pyrrolidin-1-yl)-propyl]-acetamide

LC/MS (METHOD A): (m/z) 524 (MH⁺); RT = 5.66 min; purity (UV) = 95.7%

Amine: 3-(2-Oxo-pyrrolidin-1-yl)-propylamine

5

Example 256 (compound 256)

(2R)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-indan-1-yl)-acetamide

LC/MS (METHOD A): (m/z) 530.8 (MH⁺); RT = 6.62 min; purity (UV) = 90.4%

10 Amine: (2R)-2-Hydroxy-indan-1-yl-amine

Example 257 (compound 257)

N-Cycloheptylmethyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

15 LC/MS (METHOD A): (m/z) 509 (MH⁺); RT = 8.19 min; purity (UV) = 100%

Amine: Cycloheptylmethylamine

Example 258 (compound 258)

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-[2-(2-

20 hydroxy-ethoxy)-ethyl]-acetamide

LC/MS (METHOD A): (m/z) 487 (MH⁺); RT = 5.41 min; purity (UV) = 100%

Amine: 2-(2-Hydroxy-ethoxy)-ethylamine

Example 259 (compound 259)


25 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-dimethylamino-butyl)-acetamide

LC/MS (METHOD A): (m/z) 497.9 (MH⁺); RT = 4.36 min; purity (UV) = 100%

Amine: 4-Dimethylamino-butylamine


30 Preparation 5 (compound 505)

Resin bound 2-(3,5-dichloro-pyridin-4-yl)-1-(2,3-dihydroxy-4-methoxy-phenyl)-ethanone



2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3-dihydroxy-4-methoxy-phenyl)-ethanone (530 mg, 1.6 mmol) from preparation 3 was dissolved in dry NMP (5 mL), K_2CO_3 (210 mg, 1.5 mmol) was added followed by bromomethyl-Wang resin (560 mg, L = 1.45 mmol/g, 0.81 mmol). The reaction mixture was vortexed at rt overnight. The resin was filtered and washed with MeOH:water (4:1, 3x10 mL), NMP (3x25 mL), MeOH (3x25mL), dry THF (5x25 mL) and dried *in vacuo* to provide 760 mg of resin bound 2-(3,5-dichloro-pyridin-4-yl)-1-(2,3-dihydroxy-4-methoxy-phenyl)-ethanone.

General procedure for preparation of compounds with formula Im, wherein R2 is as defined above:



Resin bound 2-(3,5-dichloro-pyridin-4-yl)-1-(2,3-dihydroxy-4-methoxy-phenyl)-ethanone (90 mg, L = 1.07 mmol/g, 0.1 mmol) from preparation 5 was treated with a 1 mL of a stock solution of R2-OH (0.5 mmol) and PBU_3 (0.5 mmol.)

cooled to 0°C and then treated with 0.5 mL of a stock solution of TBAD (0.5 mmol). After 3 hours at rt the reaction mixture was filtered and 1 mL of the stock solution of R2-OH (0.5 mmol) and PBu_3 (0.5 mmol) was added. After cooling to 0°C 0.5 mL of the stock solution of TBAD (0.5 mmol) was added. The 5 reaction mixture was vortexed at rt overnight. The resin was washed with THF (5x3 mL) and DCE (5x3 mL). Cleavage of the products was performed by addition of a stock solution of DCE-TFA-TIS (90:10:1, 1 mL). After 30 min the cleavage solution was replaced by 1 mL of a fresh cleavage solution. After 10 additional 30 min the combined cleavage solutions was evaporated *in vacuo*. The pure compounds were obtained by redissolving the reaction mixture in DMSO followed by standard preparative HPLC purification.

Using this procedure the following compounds were obtained:

15 Example 260 (compound 260)

1-(3-Cyclopentyloxy-2-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone

LC/MS (METHOD B): (m/z) 396 (MH⁺); RT = 4.84 min; purity (UV) = 100%

Alcohol: Cyclopentanol

20

Example 261 (compound 261)

1-(3-Cyclopropylmethoxy-2-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone


LC/MS (METHOD B): (m/z) 382.1 (MH⁺); RT = 4.36 min; purity (UV) = 100%

25 Alcohol: Cyclopropylmethylalcohol

Example 262 (compound 262)


2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-2-hydroxy-4-methoxy-phenyl)-ethanone

30



2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3-dihydroxy-4-methoxy-phenyl)-ethanone
 5 (450 mg, 1.4 mmol) from preparation 3 was dissolved in dry DMF (10 mL). K_2CO_3 (566 mg, 4.1 mmol) was added followed by Ethyl iodide (442 μL , 5.5 mmol). The reaction mixture was stirred at rt overnight. Water (10 mL) was added and the organic products were extracted with EtOAc (2x25 mL). The combined organic phases were washed with water (10 mL) and brine (10 mL)
 10 and then dried over MgSO_4 . The solvent was removed *in vacuo* and the pure product obtained be standard silica gel chromatography to provide 2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3-diethoxy-4-methoxy-phenyl)-ethanone as a white solid. Yield 220 mg (42%). The compound was redissolved in DCM (1 mL) under argon. BCl_3 (0.97 mL of a 1M solution in DCM) was added slowly at room
 15 temperature. EtOH (2 mL) was added slowly after 2 hours. The solvent were removed *in vacuo* and the pure product obtained be standard silica gel chromatography to provide 2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-2-hydroxy-4-methoxyphenyl)-ethanone was obtained as a white solid. ^1H NMR (DMSO- d_6) δ = 11.36 (1H, s), 8.67 (2H, s), 7.88 (1H, d), 6.77 (1H, d), 4.80 (2H, s), 3.96 (2H, q), 3.91 (3H, s), 1.26 (3H, t). Yield 100 mg (49%)
 20

General procedure for preparation of compounds with formula In, wherein R3 (R3 \neq hydrogen) is as defined above:



2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-2-hydroxy-4-methoxy-phenyl)-
 5 ethanone from example 263 (12 mg, 0.034 mmol) was dissolved in dry DMSO
 (0.25 mL). Aqueous K_2CO_3 (0.025 mL of a 2M) was added followed by 1.5 eq. of
 an alkyl bromide or iodide dissolved in 0.025 mL DMSO. The reaction mixture
 was left at rt for 48 hours. The pure compounds were obtained by standard
 preperative HPLC purification.

10

Using this procedure the following compounds were obtained:

Example 263 (compound 263)

2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-4-methoxy-2-phenethyloxy-phenyl)-
 15 ethanone 1H NMR ($CDCl_3$) δ = 8.47 (2H, s), 7.61 (1H, d), 7.26 (4H, m), 7.10
 (1H, t), 6.75 (1H, d), 4.49 (2H, t), 4.43 (2H, s), 4.01 (2H, q), 3.91 (3H, s), 3.17
 (2H, t), 1.35 (3H, t).

Alkyl halide: 2-Bromo-1-phenyl-ethane

20 Example 264 (compound 264)

1-[2-(5-Cyclopropyl-[1,3,4]thiadiazol-2-ylmethoxy)-3-ethoxy-4-methoxy-
 phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone

1H NMR ($CDCl_3$) δ = 8.47 (2H, s), 7.61 (1H, d), 6.81 (1H, d), 5.64 (2H, s), 4.56
 (2H, s), 4.15 (2H, q), 3.95 (3H, s), 2.39 (1H, m), 1.41 (3H, t), 1.56-1.11 (4H,
 m).

Alkyl halide: 2-Chloromethyl-5-cyclopropyl-[1,3,4]thiadiazol


Example 265 (compound 265)

{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2-ethoxy-3-methoxy-phenoxy}-acetic acid benzyl ester

5 ^1H NMR (DMSO-d₆) δ = 8.62 (2H, s), 7.51 (1H, d), 7.32 (5H, m), 6.96 (1H, d),
5.19 (2H, s), 4.74 (2H, s), 3.98 (2H, q), 3.88 (3H, s), 1.27 (3H, t).
Alkyl halide: 2-Bromo-acetic acid benzyl ester


Example 266 (compound 266)

10 1-(3-Allyloxy-2-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone



15 1-(2,3-Bis-allyloxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone
from example 6 (3.9g, 9.55 mmol) was dissolved in MeOH (96 mL) and cooled
to 0°C under argon. Pd(PPh₃)₄ (110mg, 0.0955 mmol) was added followed by
K₂CO₃ (1.3g, 9.55 mmol). The cooling bath was removed and the reaction
mixture was stirred for 3 hours and then evaporated *in vacuo*. The crude was
20 redissolved in EtOAc (100 mL) and washed twice with NH₄Cl (sat, 50 mL). The
organic phase was dried over Na₂SO₄ and evaporated *in vacuo* and purified by
flash chromatography using a gradient of toluene and 2% EtOAc as eluent. 1-(3-
Allyloxy-2-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone
was obtained as a white solid. ^1H NMR (CDCl₃) δ = 11.90 (1H, s), 8.53 (2H, s),
7.68 (1H, d), 6.58 (1H, d), 6.10 (1H, m), 5.26 (2H, m), 4.66 (2H, s), 4.59 (2H,
25 d), 3.96 (3H, s). Yield 2.2g (64%)

General procedure for preparation of compounds with formula Io, wherein R3 (R3 ≠ hydrogen) is as defined above:



2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-2-hydroxy-4-methoxy-phenyl)-ethanone from example 263 (12 mg, 0.034 mmol) was dissolved in dry DMSO (0.25 mL). Aqueous K_2CO_3 (0.025 mL of a 2M) was added followed by 1.5 eq. of an alkyl bromide or iodide dissolved in 0.025 mL DMSO. The reaction mixture was left at rt for 48 hours. The pure compounds were obtained by standard preoperative HPLC purification.

Using this procedure the following compounds were obtained:

15

Example 267 (compound 267)

2-{2-Allyloxy-6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-3-methoxy-phenoxy-methyl}-benzonitrile

LC/MS (METHOD B): (m/z) 383.2 (MH⁺); RT = 4.72 min; purity (UV) = 100%

20 Alkyl halide: 2-Chloromethyl-benzonitrile

Example 268 (compound 268)

1-(3-Allyloxy-4-methoxy-2-phenethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone

25 LC/MS (METHOD B): (m/z) 472.3 (MH⁺); RT = 5.32 min; purity (UV) = 100%

Alkyl halide: 1-Bromo-2-phenyl-ethane


Example 269 (compound 269)

1-{3-Allyloxy-2-[2-(4-fluoro-phenyl)-ethoxy]-4-methoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone

5 LC/MS (METHOD B): (m/z) 490.2 (MH⁺); RT = 5.26 min; purity (UV) = 100%
 Alkyl halide: 1-Bromo-2-(4-fluoro-phenyl)-ethane

General procedure for preparation of compounds with formula Ip, wherein R3 is as defined above:

10



1-(2-Alkoxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone from the previous examples (0.021 mmol) was dissolved in dry DCM (0.25 mL).

15 Methyltrioxorhenium (0.105 mmol) was added followed by hydrogenperoxide (0.042 mmol) and the reaction mixture was stirred at rt overnight. MnO₂ (0.063 mmol) was added. After 2 minutes the reaction mixture was after filtered and the reaction mixture evaporated *in vacuo*. The crude was re-dissolved in DMSO (0.4 mL). The pure compounds were obtained by standard preperative HPLC
 20 purification.

Using this procedure the following compounds were obtained:

Example 270 (compound 270)

N-Benzyl-2-{6-[2-(3,5-dichloro-1-oxy-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide

¹H NMR (CDCl₃) δ = 8.15 (2H, s), 8.10 (1H, bs), 7.62 (1H, d), 7.30 (5H, m),

6.76 (1H, d), 4.72 (2H, s), 4.50 (2H, d), 4.46 (2H, s), 3.95 (3H, s), 3.80 (3H,

5 s). Using N-benzyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide from example 203 as starting material.

Example 271 (compound 271)

2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-(3,4-dimethoxy-2-phenethyloxy-phenyl)-

10 ethanone

¹H NMR (CDCl₃) δ = 8.13 (2H, s), 7.61 (1H, d), 7.26 (5H, m), 6.75 (1H, d), 4.40 (2H, t), 4.30 (2H, s), 3.93 (3H, s), 3.79 (3H, s), 3.17 (2H, t).

Using 2-(3,5-dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-phenethyloxy-phenyl)-ethanone from example 112 as starting material.

15

Example 272 (compound 272)

2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone

¹H NMR (CDCl₃) δ = 8.19 (2H, s), 7.61 (1H, d), 7.26 (2H, t), 6.95 (2H, t), 6.75

20 (1H, d), 4.44 (2H, t), 4.34 (2H, s), 3.93 (3H, s), 3.77 (3H, s), 3.14 (2H, t).

Using 2-(3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone from example 129 as starting material.

Example 274 (compound 274)

2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(1-oxy-pyridin-4-yl)-ethoxy]-phenyl}-ethanone

¹H NMR (DMSO-d₆) δ = 8.62 (2H, s), 8.11 (2H, d), 7.54 (1H, d), 7.39 (2H, d), 6.97 (1H, d), 4.41 (4H, m), 3.88 (3H, s), 3.67 (3H, s), 3.11 (2H, t).

Using example 100 as starting material.

30

Example 275 (compound 275)

2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-(2-hydroxy-3,4-dimethoxy-phenyl)-ethanone

¹H NMR (DMSO-d₆) δ = 11.20 (1H, bs), 8.64 (2H, s), 7.84 (1H, d), 6.77 (1H, d), 4.70 (2H, bs), 3.91 (3H, s), 3.72 (3H, s).

Using 2-(3,5-dichloro-pyridin-4-yl)-1-(2-hydroxy-3,4-dimethoxy-phenyl)-ethanone from example 1 as starting material.

5

Example 276 (compound 276)

4-(2-{6-[2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-ethyl)-benzonitrile


¹H NMR (DMSO-d₆) δ = 8.61 (2H, s), 7.73 (2H, d), 7.55 (3H, m), 6.96 (1H, d),

10 4.45 (2H, t), 4.20 (2H, s), 3.88 (3H, s), 3.66 (3H, s), 3.22 (2H, t).

Using example 100 as starting material.

General procedure for preparation of compounds with formula Iq, wherein R3 (R3 ≠ hydrogen) is as defined above:

15



2-(3,5-Dichloro-pyridin-4-yl)-1-(2-hydroxy-3,4-dimethoxy-phenyl)-ethanone

from example 1 (1.5 mmol) or 2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-(2-

20 hydroxy-3,4-dimethoxy-phenyl)-ethanone from example 274 (1.5 mmol), PBu₃ (1.5 mmol) and R3-OH (1.0 mmol) was dissolved in dry benzene (10 mL) under argon. The reaction mixture was cooled to 0°C and then treated with ADDP (1.5 mmol). After 0.5 hours the reaction mixture was brought to rt and stirred overnight. Decalite was added and the reaction mixture was evaporated *in vacuo* 25 and purified by flash chromatography using a gradient of petroleum ether and EtOAc as eluent.

Using this procedure the following compounds were obtained:

Example 277 (compound 277)

5 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-pyridin-4-yl-ethoxy)-phenyl]-ethanone
¹H NMR (DMSO-d₆) δ = 8.63 (2H, s), 8.44 (2H, d), 7.53 (1H, d), 7.37 (2H, d),
6.97 (1H, d), 4.46 (4H, m), 3.88 (3H, s), 3.66 (3H, s), 3.15 (2H, t).
R3-OH: 2-Pyridin-4-yl-ethanol

10

Example 278 (compound 278)

4-(2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-ethyl)-benzonitrile
¹H NMR (DMSO-d₆) δ = 8.63 (2H, s), 7.70 (2H, d), 7.55 (3H, m), 6.97 (1H, d),
4.47 (2H, t), 4.30 (2H, s), 3.89 (3H, s), 3.68 (3H, s), 3.23 (2H, t).
R3-OH: 4-(2-Hydroxy-ethyl)-benzonitrile prepared from 2-(4-bromo-phenyl)-ethanol according to

Example 279 (compound 279)

20 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-pyridin-2-yl-ethoxy)-phenyl]-ethanone
¹H NMR (DMSO-d₆) δ = 8.62 (2H, s), 8.41 (1H, t), 7.65 (1H, t), 7.50 (1H, d),
7.38 (1H, d), 7.10 (1H, m), 6.95 (1H, d), 4.61 (2H, t), 4.38 (2H, s), 3.88 (3H, s),
3.72 (3H, s), 3.28 (2H, t).
25 R3-OH: 2-Pyridin-2-yl-ethanol

Example 280 (compound 280)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-pyridin-3-yl-ethoxy)-phenyl]-ethanone
30 ¹H NMR (DMSO-d₆) δ = 8.64 (2H, s), 8.55 (1H, s), 8.34 (1H, d), 7.77 (1H, d),
7.53 (1H, d), 7.28 (1H, m), 6.97 (1H, d), 4.47 (2H, s), 4.43 (2H, t), 3.88 (3H, s),
3.64 (3H, s), 3.15 (2H, t).
R3-OH: 2-Pyridin-3-yl-ethanol

Example 281 (compound 281)

2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(4-methanesulfinyl-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone

(CDCl₃) δ = 8.50 (2H, s), 7.59 (3H, m), 7.49 (2H, d), 6.78 (1H, d), 4.48 (4H,

5 m), 3.93 (3H, s), 3.74 (3H, s), 3.24 (2H, t), 2.63 (3H, s).

R3-OH: 2-(4-Methanesulfinyl-phenyl)-ethanol, prepared by oxidising 2-(4-methylsulfanyl-phenyl)-ethanol with 1.5 eq MCPBA in DCM rt overnight. ¹H NMR

(CDCl₃) δ = 7.88 (2H, d), 7.45 (2H, d), 3.92 (2H, t), 3.04 (3H, s), 2.97 (2H, t),

1.70 (1H, bs). LC/MS (METHOD B): (m/z) 185 (MH⁺); RT = 1.27 min

10

Example 282 (compound 282)

2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(4-methanesulfonyl-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone

¹H NMR (CDCl₃) δ = 8.50 (2H, s), 7.88 (2H, d), 7.59 (1H, d), 7.53 (2H, d), 6.77

15 (1H, d), 4.51 (2H, s), 4.49 (2H, t), 3.93 (3H, s), 3.72 (3H, s), 3.27 (2H, t), 2.98 (3H, s).

R3-OH: 2-(4-Methanesulfonyl-phenyl)-ethanol, prepared by oxidising 2-(4-methylsulfanyl-phenyl)-ethanol with 1.5 eq MCPBA in DCM rt overnight. ¹H NMR

(CDCl₃) δ = 7.59 (2H, d), 7.40 (2H, d), 3.90 (2H, t), 2.94 (2H, t), 2.72 (3H, s),

20 1.85 (1H, bs). LC/MS (METHOD B): (m/z) 201 (MH⁺); RT = 1.48 min

Example 283 (compound 283)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(1-phenyl-propoxy)-phenyl]-ethanone

25 ¹H NMR (DMSO-d₆) δ = 8.65 (2H, s), 7.34 (6H, m), 6.85 (1H, d), 5.48 (1H, t),

4.63 (2H, s), 3.85 (3H, s), 3.72 (3H, s), 2.14 (1H, m), 1.95 (1H, m), 0.88 (3H,

t).

R3-OH: 1-Phenyl-propan-1-ol

30 Example 284 (compound 284)

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-phenyl-propoxy)-phenyl]-ethanone

¹H NMR (DMSO-d₆) δ = 8.62 (2H, s), 7.51 (1H, d), 7.35 (2H, d), 7.23 (2H, t), 7.06 (1H, t), 6.95 (1H, d), 4.33 (4H, m), 3.88 (3H, s), 3.69 (3H, s), 3.33 (1H, m), 1.37 (3H, d).

R3-OH: 2-Phenyl-propan-1-ol

5

Example 285 (compound 285)


2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(1-methyl-2-phenyl-ethoxy)-phenyl]-ethanone

10 ¹H NMR (DMSO-d₆) δ = 8.66 (2H, s), 7.46 (1H, d), 7.28 (4H, d), 7.17 (1H, m), 6.93 (1H, d), 4.94 (1H, m), 4.60 (2H, m), 3.88 (3H, s), 3.58 (3H, s), 3.02 (2H, m), 1.18 (3H, d).

R3-OH: 1-Phenyl-propan-2-ol

Preparation 6 (compound 506):


15



Methyl 2,3,4-trimethoxybenzoic acid (25.7g, 114 mmol) was dissolved in DCM (25 mL) under argon. BCl₃ (133 mL of an 1M solution in DCM, 133 mmol) was 20 added dropwise and the reaction mixture was left at rt for 2 hours. EtOH (200 mL) was added and the reaction mixture was stirred for 2 hours. Filter the precipitate and recryst. from EtOAc-petrol ether to provide 6g (25%) of 2-hydroxy-3,4-dimethoxy-benzoic acid methyl ester as a white solid. ¹H NMR (CDCl₃) δ = 10.90 (1H, s), 7.59 (1H, d), 6.48 (1H, d), 3.93 (3H, s), 3.92 (3H, s), 3.89 (3H, s). 2-Hydroxy-3,4-dimethoxy-benzoic acid methyl ester (6g, 28.2 mmol) was redissolved in NMP (35 mL) and treated with *tert*-butyl bromoacetate (42.4 mmol) and K₂CO₃ (42.4 mmol). The reaction mixture was heated to 50°C overnight. The reaction mixture was poured into water (300 mL). The organic products were extracted with EtOAc (2x100 mL), and the combined organic

phases washed with NaCl (sat, 100 mL). The organic phase was dried over Na₂SO₄, evaporated *in vacuo* and purified by flash chromatography using a gradient of EtOAc in heptane as eluent to provide 2-tert-butoxycarbonylmethoxy-3,4-dimethoxy-benzoic acid methyl ester (compound 506a) as a colourless oil. ¹H NMR (CDCl₃) δ = 7.58 (1H, d), 6.71 (1H, d), 4.56 (2H, s), 3.90 (3H, s), 3.87 (3H, s), 3.86 (3H, s), 1.50 (9H, s). Yield 8.19g (89%). 2-tert-Butoxycarbonylmethoxy-3,4-dimethoxy-benzoic acid methyl ester (8.19g, 25.1 mmol) was dissolved in DCM (50 mL) and treated with triethylsilane (4mL, 25.1 mmol) and TFA (9.66 mL, 125.5 mmol). The reaction mixture was left overnight. The reaction mixture was evaporated *in vacuo*, redissolved in toluene (200 mL) and evaporated to dryness. This procedure was repeated three times to afford 2-carboxymethoxy-3,4-dimethoxy-benzoic acid methyl ester (compound 506b) as a white solid. ¹H NMR (CDCl₃) δ = 12.6 (1H, bs), 7.75 (1H, d), 6.73 (1H, d), 4.84 (2H, s), 3.95 (3H, s), 3.92 (3H, s), 3.85 (3H, s). Yield 6.71g (99%). 2-Carboxymethoxy-3,4-dimethoxy-benzoic acid methyl ester (6.71g, 24.8 mmol) was dissolved in dry DMF (130 mL) under argon. Propylamine (4.1 mL, 49.8 mmol) was added followed by HATU (11.32g, 29.8 mmol) and the reaction mixture was left overnight. The solvent was evaporated *in vacuo* and the crude mixture was redissolved in EtOAc (150 mL). Wash the organic phase with CaCl₂ aq. (2x50 mL), water (2x50 mL) and brine (50 mL). The organic phase was dried over Na₂SO₄, evaporated *in vacuo* and purified by flash chromatography using a gradient of EtOAc in heptane as eluent. 3,4-Dimethoxy-2-propylcarbamoylmethoxy-benzoic acid methyl ester (compound 506c) was obtained as a white solid. ¹H NMR (CDCl₃) δ = 8.47 (1H, bs), 7.71 (1H, d), 6.70 (1H, d), 4.69 (2H, s), 3.93 (3H, s), 3.87 (3H, s), 3.83 (3H, s), 3.33 (2H, q), 1.64 (2H, m), 0.99 (3H, t). Yield 6.67g (86%).

General procedure for preparation of compounds with formula I_r, wherein Ar and HetAr is as defined above:



3,4-Dimethoxy-2-propylcarbamoylmethoxy-benzoic acid methyl ester (28mg, 0.09 mmol) obtained from Preparation 6 and a Ar-Me or HetAr-Me compound 5 (1.2 eq, see below) was dissolved in dry THF (1 mL) under argon. The mixture was cooled to 0° and treated dropwise with LiHMDS (0.273 mL of a 1M solution). The reaction mixture was brought to room temperature and left overnight. The reaction was quenched with NH₄Cl (sat., 2 mL) and the organic products were extracted with EtOAc (2x2 mL). The combined organic phases was washed with 10 NaCl (sat, 2 mL). The organic phase was dried over Na₂SO₄, evaporated *in vacuo* and redissolved in MeOH (0.350 mL) followed by standard HPLC purification.

Using this procedure the following compounds were obtained:

15 Example 286 (compound 286)

2-{6-[2-(6-Chloro-pyrazin-2-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide

LC/MS (METHOD B): (m/z) 408.4 (MH⁺); RT = 3.33 min; purity (UV) = 91%
HetAr-Me: 2-Chloro-6-methyl-pyrazine

20

Example 287 (compound 287)

2-{6-[2-(3-Bromo-pyrazin-2-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide

LC/MS (METHOD B): (m/z) 454.3 (MH⁺); RT = 3.26 min; purity (UV) = 86%

25 HetAr-Me: 2-Bromo-3-methyl-pyrazine

Example 288 (compound 288)

2-{6-[2-(2,6-Dichloro-phenyl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide

¹H NMR (CDCl₃) δ = 7.84 (1H, bs), 7.68 (1H, d), 7.34 (2H, d), 7.19 (1H, t), 6.77 (1H, d), 4.66 (2H, s), 4.60 (2H, s), 3.95 (3H, s), 3.86 (3H, s), 3.25 (2H, q), 1.48 (2H, m), 0.86 (3H, t).

Ar-Me: 1,3-Dichloro-2-methyl-benzene

5

Example 289 (compound 289)

2-[2,3-Dimethoxy-6-(2-pyridin-4-yl-acetyl)-phenoxy]-N-propyl-acetamide

¹H NMR (DMSO-d₆) δ = 8.49 (2H, bs), 8.15 (1H, t), 7.55 (1H, d), 7.24 (2H, d), 6.95 (1H, d), 4.56 (2H, s), 4.44 (2H, s), 3.88 (3H, s), 3.77 (3H, s), 3.09 (2H, q), 1.42 (2H, m), 0.81 (3H, t).

HetAr-Me: 4-Methyl-pyridine

Example 290 (compound 290)

2-[2,3-Dimethoxy-6-(2-quinolin-4-yl-acetyl)-phenoxy]-N-propyl-acetamide

15 LC/MS (METHOD B): (m/z) 423.4 (MH⁺); RT = 2.45 min; purity (UV) = 100%

HetAr-Me: 4-Methyl-quinoline

Example 291 (compound 291)

2-[2,3-Dimethoxy-6-(2-pyrazin-2-yl-acetyl)-phenoxy]-N-propyl-acetamide

20 LC/MS (METHOD B): (m/z) 374.1 (MH⁺); RT = 2.58 min; purity (UV) = 81%

HetAr-Me: 2-Methyl-pyrazine

Example 292 (compound 292)

2-{6-[2-(3-Bromo-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide

LC/MS (METHOD B): (m/z) 453.07 (MH⁺); RT = 3.13 min; purity (UV) = 100%

HetAr-Me: 3-Bromo-4-methyl-pyridine

Example 293 (compound 293)

30 2-{6-[2-(3,5-Dibromo-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide

LC/MS (METHOD B): (m/z) 531.0 (MH⁺); RT = 3.68 min; purity (UV) = 100%

HetAr-Me: 3,5-Dibromo-4-methyl-pyridine

Example 294 (compound 294)

2-{6-[2-(6-Chloro-pyrimidin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide

LC/MS (METHOD B): (m/z) 408.3 (MH⁺); RT = 3.19 and 3.79 min (probably

5 keton and enol isomer); purity (UV) = 100%

HetAr-Me: 4-Chloro-6-methyl-pyrimidine

Example 295 (compound 295)

2-{6-[2-(4-Chloro-pyridin-2-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide

10 LC/MS (METHOD B): (m/z) 407.4 (MH⁺); RT = 3.50 min; purity (UV) = 100%

HetAr-Me: 4-Chloro-2-methyl-pyridine

Example 296 (compound 296)

15 2-{6-[2-(2-Chloro-pyridin-3-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide

LC/MS (METHOD B): (m/z) 407.4 (MH⁺); RT = 3.26 min; purity (UV) = 100%

HetAr-Me: 2-Chloro-3-methyl-pyridine

20 Example 297 (compound 297)

2-{2,3-Dimethoxy-6-[2-(2-methoxy-pyridin-4-yl)-acetyl]-phenoxy}-N-propyl-acetamide

LC/MS (METHOD B): (m/z) 403.3 (MH⁺); RT = 3.24 min; purity (UV) = 100%

HetAr-Me: 2-Methoxy-4-methyl-pyridine

25

Example 298 (compound 298)

2-{6-[2-(2-Cyano-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide

LC/MS (METHOD B): (m/z) 398.4 (MH⁺); RT = 3.23 min; purity (UV) = 94%

30 HetAr-Me: 4-Methyl-pyridine-2-carbonitrile

Example 299 (compound 299)


2-[2,3-Dimethoxy-6-(2-pyridazin-3-yl-acetyl)-phenoxy]-N-propyl-acetamide

LC/MS (METHOD B): (m/z) 374.2 (MH⁺); RT = 2.39 min; purity (UV) = 90%

HetAr-Me: 3-Methyl-pyridazine

Example 300 (compound 300)

2-(2-tert-Butylamino-3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone




2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-

10 dimethoxy-phenyl}-ethanone obtained from example 272 was converted into 2-(2-tert-Butylamino-3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone according to the procedure described in *J. Org. Chem.* (2007), **72**, 4554-57. Pure 2-(2-tert-butylamino-3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone was obtained by standard HPLC purification.

LC/MS (METHOD B): (m/z) 535.2 (MH⁺); RT = 6.21 min; purity (UV) = 100%

Example 301 (compound 301)

2-(2-amino-3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone




2-(2-tert-Butylamino-3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone obtained from example 300 (16.5 mg, 0.0308 mmol), triethylsilane (2 eq.) and TFA (0.1 mL) in dichloroethane (0.2 mL) was heated to 50°C for 24 hours. The crude mixture was evaporated *in vacuo*, redissolved in dichloroethane (1 mL) and washed with NaHCO₃ (sat, 2x0.5 mL). The organic phase was dried over Na₂SO₄ and the solvent evaporated *in vacuo*. Pure 2-(2-amino-3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone was obtained by flash chromatography using a gradient of MeOH in dichloromethane as eluent.

LC/MS (METHOD B): (m/z) 479.3 (MH⁺); RT = 4.53 min; purity (UV) = 100%

15 Example 302 (compound 302)

2-(3,5-Dichloro-pyridin-4-yl)-1-(4-ethoxy-3-methoxy-2-phenethyloxy-phenyl)-ethanone



2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-phenethoxy-phenyl)-ethanone obtained from example 112 (89.3 mg, 0.1 mmol) was treated with 5 piperidine (0.8 mL) and water (0.32 mL). The yellow suspension was heated to 90°C for 54 hours. The solvent was evaporated *in vacuo*. The crude mixture was treated with NH₄Cl (sat., 1 mL) and the organic products were extracted with EtOAc (3x2 mL). The combined organic phases was dried over Na₂SO₄ and evaporated *in vacuo*. Purification by flash chromatography using a gradient of 10 EtOAc in toluene as eluent provided 2-(3,5-dichloro-pyridin-4-yl)-1-(4-hydroxy-3-methoxy-2-phenethoxy-phenyl)-ethanone as a white solid. Yield 8.6 mg (20%). 2-(3,5-Dichloro-pyridin-4-yl)-1-(4-hydroxy-3-methoxy-2-phenethoxy-phenyl)-ethanone (5.5 mg, 0.013 mmol) was dissolved in DMSO (0.45 mL) and treated with K₂CO₃ (0.019 mL of a 1M aq. solution) followed by ethyliodide 15 (0.019 mL of a 1M solution in DMSO). The reaction mixture was left at rt for 48 hours and purified by standard HPLC purification. 2-(3,5-Dichloro-pyridin-4-yl)-1-(4-ethoxy-3-methoxy-2-phenethoxy-phenyl)-ethanone was obtained as a colourless solid. Yield 3.2 mg (36%).

20 ¹H NMR (CDCl₃) δ = 8.48 (2H, s), 7.59 (1H, d), 7.27 (4H, m), 7.11 (1H, t), 6.73 (1H, d), 4.48 (4H, m), 4.14 (2H, q), 3.78 (3H, s), 3.18 (2H, t), 1.49 (3H, t).

Example 303

PDE4 assay

25 Human recombinant PDE4 (Gene bank accession no NM_006203) are incubated for 1 hour, respectively, with the test compound at concentrations up to 10 uM,

with cAMP (1×10^{-5} M), and with a low amount (0.021 MBq) of radioactively labelled cAMP. At the end of the incubation, the cleavage of the substrate is evaluated by the binding of the AMP product to SPA beads, which generate chemo luminescence when bound to the radioactive tracer. The AMP product 5 inhibits the binding of the radioactive tracer to the beads, and the luminescent signal is competed.

The results are calculated as the molar concentrations resulting in 50% inhibition of the substrate cleavage compared to controls samples, and are expressed as $-\log IC_{50}$ (M).

10 Compounds 101, 103, 104, 106-148, 150-155, 157-206, 208-234, 236-241, 243, 244, 246, 248, 249, 251,-258, 260-272, 274-5, 277-285, 287-290, 292-293, 296, 301 and 302 were shown to be efficient inhibitors with $-\log IC_{50}$ (M) above 6.

15 Example 304

TNFalpha- release

Human peripheral blood mononuclear cells (PBMC) are isolated from buffy coats.

The blood is mixed with saline at a ratio of 1:1, and the PBMC is isolated by the use of Lymphoprep tubesTM (Nycomed, Norway). The PBMC is suspended in


20 RPMI1640 with 2 % foetal calf serum (FCS), pen/strep and 2 mM L-glutamine at a concentration of 5×10^5 c/ml. The cells are pre-incubated for 30 minutes with the test compounds in 96 well tissue culture plates and stimulated for 18 hours with lipopolysaccharide 1 mg/ml (Sigma). The level of TNF-a is measured in the culture supernatant by enzyme immuno assays using primary and secondary 25 biotinylated antibodies from R&D systems. Results are expressed as pIC₅₀ values calculated from inhibition curves using as positive controls the secretion in LPS stimulated wells and as negative controls the secretion in unstimulated cells.

Compounds 101, 104, 106-138, 140-148, 150-153, 155, 157,161-164, 168,

30 170, 171, 176, 177, 182-186, 188-190, 193, 202-206, 208-241, 243, 244, 246, 248, 249, 251-258, 260-275, 277-282, 284, 285, 288-290, 292 and 293 were shown to be efficient inhibitors with $-\log IC_{50}$ (M) above 6.

CLAIMS

1. A compound according to formula I



wherein X_1 , X_2 , X_3 , X_4 and X_5 independently of each other represent $-CH-$ or N ;

10 or X_3 , X_4 and X_5 independently of each other represent $-CH-$ or N , and X_1 and X_2 independently of each other represent C and form part of an additional 6-membered aromatic ring;

15 wherein R_1 represents alkyl, alkenyl, alkynyl, haloalkyl, hydroxyalkyl, or alkylcarbonyl, all of which are optionally substituted by one or more, same or different substituents selected from R_4 , or R_1 represents hydrogen;

20 R_2 represents alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, hydroxyalkyl, heterocycloalkenyl, alkylaryl, arylalkyl, alkylalkoxycarbonyl, alkylcarbonyloxy, or alkoxyalkyl, all of which are optionally substituted by one or more, same or different substituents selected from R_5 ; or R_2 represents hydrogen or $-CH_2-C(O)NR_9R_{12}$;

25 R_3 represents alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, hydroxyalkyl, heterocycloalkenyl, alkylaryl, arylalkyl, alkylalkoxycarbonyl,

alkylcarbonyloxy, or alkoxyalkyl, all of which are optionally substituted by one or more, same or different substituents selected from R₆; or R₃ represents hydrogen, -CH₂-C(O)-heterocycloalkyl or -CH₂-C(O)NR₉-R₁₂;

5 R₄ represents hydrogen, alkyl, alkenyl, alkynyl, halogen, oxo, alkoxy, hydroxy, or haloalkyl;

R₅ represents alkylaryl, carboxy, alkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, carbamoyl, hydroxyalkyl, aryloxy, alkoxy carbonyloxy, 10 alkoxy carbonyl, alkoxy, alkoxyalkyl, aryl, a heterocyclic ring, aminocarbonyl, alkylthio, alkylcarbonylamino, hydroxy, alkylcarbonyl, arylcarbonyl, alkylcarbonyloxy, or amino, all of which are optionally substituted by one or more, same or different substituents selected from R₇;

or R₅ represents hydrogen, oxo, halogen, cyano, or nitro;

15 R₆ represents alkylaryl, carboxy, alkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, carbamoyl, hydroxyalkyl, aryloxy, alkoxy carbonyloxy, alkoxy carbonyl, alkoxy, alkoxyalkyl, aryl, a heterocyclic ring, aminocarbonyl, alkylthio, alkylcarbonylamino, arylcarbonyl, hydroxy, alkylcarbonyl, 20 alkylcarbonyloxy, or amino, all of which are optionally substituted by one or more, same or different substituents selected from R₈;

or R₆ represents hydrogen, oxo, halogen, cyano, or nitro;

R₇ represents alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkynyl, alkoxy, haloalkyl, 25 alkylthio, heterocycloalkenyl, heterocycloalkyl, aryl, alkylcarbonyl, heteroaryl, aryloxy, alkoxy carbonyl, hydroxyalkyl, amino, hydroxy, or carboxy; all of which are optionally substituted by one or more, same or different substituents selected from R₁₀;

or R₇ represents hydrogen, halogen, or oxo;

30 R₈ represents alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkynyl, alkoxy, haloalkyl, alkylthio, alkylsulfonyl, alkylsulfinyl, heterocycloalkenyl, heterocycloalkyl, aryl, alkylcarbonyl, heteroaryl, aryloxy, alkoxy carbonyl, hydroxyalkyl, amino, hydroxy, or carboxy; all of which are optionally substituted by one or more,

same or different substituents selected from R₁₀;
or R₈ represents hydrogen, halogen, or oxo;

R₉ represents hydrogen, alkyl, haloalkyl, or hydroxyalkyl;

5

R₁₀ represents hydrogen, alkyl, oxo, hydroxy, halogen, carboxy, amino, alkoxy, haloalkyl, or hydroxyalkyl;

R₁₁ represents one or more, same or different substituents selected from

10 hydrogen, halogen, cyano, amino, alkyl, methylsulfinyl, methylsulfonyl, amino, cyano, or alkoxy;

R₁₂ represents alkylaryl, arylalkyl, carboxy, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, carbamoyl, hydroxyalkyl, aryloxy, alkoxy carbonyloxy, alkoxy carbonyl, alkoxy, alkoxyalkyl, aryl, a heterocyclic ring, aminocarbonyl, alkylthio, alkylcarbonylamino, hydroxy, alkylcarbonyl, arylcarbonyl, alkylcarbonyloxy, or amino, all of which are optionally substituted by one or more, same or different substituents selected from R₈, or R₁₂ represents hydrogen;

20

with the proviso R₁, R₂ and R₃ cannot all be methyl;

with the proviso that when R₂ and R₃ are both hydrogen, R₁ cannot be methyl or hydrogen;

with the proviso that when R₁ is methyl or hydrogen, R₂ is methyl and R₃ is


25 hydrogen, then ring B cannot be phenyl;

and pharmaceutically acceptable and physiologically cleavable esters, pharmaceutically acceptable salts, hydrates, N-oxides or solvates thereof.

30 2. A compound according to claim 1, wherein ring B represents pyridyl, pyrazinyl, quinolyl, pyrimidinyl or pyridazinyl optionally substituted with one or more same or different substituents selected from fluoro, chloro, bromo, cyano, methoxy, -NH₂ or C₁₋₄amino.

3. A compound according to any one of claims 1-2, wherein ring B, optionally substituted with R₁₁, represents 2-(6-chloro-pyrazinyl), 2-pyrazinyl, 4-(3-bromo-pyridyl), 4-(3,5-dibromo-pyridyl), 4-(6-chloro-pyrimidinyl), 2-(4-chloro-pyridyl), 3-(2-chloro-pyridyl), 4-(2-methoxy-pyridyl), 4-(2-cyano-pyridyl), 3-pyridazinyl,
 5 4-(2-*tert*-butylamino-3,5-dichloro-pyridyl), 4-(2-amino-3,5-dichloro-pyridyl), 4-(3,5-dichloro-pyridyl), 2-(3-bromo-pyrazinyl), 4-pyridyl, 4-quinolyl or 4-(3,5-dichloro-1-oxy-pyridyl).

10 4. A compound according to claim 1, wherein formula I represents general formula I_z,

I_z

15 wherein X₃ represents -CH- or N.

5. A compound according to any one of claims 1-4 wherein R₁ represents methyl or ethyl.

20 6. A compound according to any one of the preceding claims wherein R₂ represents C₁-C₆alkyl, C₁-C₆alkenyl, C₁-C₆alkoxyC₁-C₆alkyl, hydroxyC₁-C₆alkyl, haloC₁-C₆alkyl, C₁-C₆alkynyl, C₁-C₆cycloalkyl, C₁-C₆alkylC₆-C₁₀aryl, C₁-C₆alkylC₁-C₆alkoxycarbonyl or C₁-C₆alkylcarbonyloxy, all of which are optionally substituted by one or more, same or different substituents selected from R₅.

7. A compound according to any one of the preceding claims wherein R₂ represents methyl, ethyl, propyl, tert-butoxycarbonylmethyl, allyl, difluoromethyl, ethylbenzene, methylbenzene, butenyl, hydroxyethyl, tolyl, pentenyl, methoxyethyl, butinyl, propinyl, cyclopentyl, all of which are optionally substituted by one or more, same or different substituents selected from R₅.

8. A compound according to any one of the preceding claims wherein R₃ represents methyl, ethyl, propyl, butyl, pentyl, hexyl, hydroxyethyl, butenyl, pentenyl, allyl, butinyl, benzyl, methylbenzene, ethylbenzene, ethylpyridine, tolyl, toluoyl, propylbenzene, methylnaphthyl, ethylnaphthyl, methylcarbonylmethoxy, methylcarbonylethoxy, methoxyethyl, methoxypropyl, all of which are optionally substituted by one or more, same or different substituents selected from R₆, said R₆ substituent being optionally substituted with one or more, same or different substituents selected from R₈, or R₃ represents hydrogen, -CH₂-C(O)-heterocycloalkyl or -CH₂-C(O)NR₉-R₁₂.

9. A compound according to any one of the preceding claims wherein R₅ represents methyl, tert-butoxy, ethenyl, cyclopropyl, propenyl, phenyl, butenyl, propinyl, methylhydroxy, ethinyl, allyl, ethyl, or methoxy, all of which are optionally substituted by one or more, same or different substituents selected from R₇, or R₅ represents hydrogen, oxo, chloro, fluoro, or hydroxy.

10. A compound according to any one of the preceding claims wherein R₆ represents ethenyl, methyl, tert-butoxy, isoxazolyl, methoxy, propinyl, butenyl, phenyl, pyridyl, benzoxazolyl, thiazolyl, [1,3,4]thiadiazolyl, [1,2,4]oxadiazolyl, 2,3-dihydro-1H-isoindolyl, ethoxy, thiophenyl, propyl, ethyl, butyl, pentyl, allyl, isopropoxy, isopropyl, naphthyl, cyclohexyl, hydroxy, cyclopentyl, phenoxy, tolyl, toluoyl, benzoyl, carbonylnaphthalene, ethylbenzene, quinolinyl, -NH₂, ethoxycarbonyl, methoxycarbonyl, carbamoyl, isoindole, methylamine, pyrrolidyl, morpholinyl, methylsulfonyl, methylsulfinyl, butylamine, propylamine, ethylamine, cycloheptyl, hydroxyethyl, hydroxypropyl, indanyl, or ethoxyethyl, all of which are optionally substituted by one or more, same or different

substituents selected from R₈, or R₆ represents hydrogen, oxo, fluoro, chloro, or cyano.

11. A compound according to any one of the preceding claims wherein R₈ represents, methyl, ethyl, propyl, butyl, phenyl, cyclopropyl, ethoxy, methoxy, allyl, ethenyl, ethoxycarbonyl, hydroxy, naphthyl, cyclohexyl, methoxycarbonyl, phenoxo, isopropoxy, -NH₂, methylamine, pyrrolidinyl, morpholinyl, methylsulfonyl, methylsulfinyl, cycloheptyl, cyclopentyl, hydroxymethyl, hydroxyethyl, dimethylamino, furanyl, pyridyl, tolyl, piperidinyl, acetyl, thiophenyl, cycloheptyl, all of which are optionally substituted by one or more, same or different substituents selected from R₁₀, or R₈ represents hydrogen, oxo, chloro, bromo, fluoro, cyano, or trifluoromethyl.
12. A compound according to any one of the preceding claims wherein R₉ represents hydrogen, methyl or ethyl.
13. A compound according to any one of the preceding claims wherein R₁₀ represents hydrogen, oxo, methyl, hydroxy, fluoro, cyano, chloro, or methoxy.
14. A compound according to any one of the preceding claims wherein R₃ represents -CH₂-C(O)NH-R₁₂, -CH₂-C(O)-heterocycloalkyl, -CH₂CH₂-phenyl-R₆, or -CH₂-phenyl-R₆.
15. A compound according to any one of the preceding claims wherein R₂ and/or R₃ represents -CH₂COOH, methyl, hydrogen, allyl, ethyl, tert-butoxycarbonylmethyl, difluoromethyl, 3-methyl-5-methylisoxazole, 2-methoxyethane, 2-butyne, 2-methyl-2-butene, 2-phenylethane, benzyl, 2-methyl-1,3-benzoxazole, 4-methyl-2-methylthiazole, 2-methyl-5-cyclopropyl-[1,3,4]thiadiazole, 3-methyl-[1,2,4]oxadiazole, ethyl acetate, (4-chlorophenyl)ethane, 5-chloro-2-methyl-thiophene, phenoxyethane, (4-methylphenyl)ethane, 3-phenylpropane, (3-methoxyphenyl)ethane, (4-methoxyphenyl)ethane, (3-bromophenyl)ethane, (2-methoxyphenyl)ethane, (4-fluorophenyl)ethane, (2-fluorophenyl)ethane, (3,4-dimethoxyphenyl)ethane, benzyl acetate, isopropyl acetate, 3-methyl benzoic acid methyl ester, 3-methyl-

butane, 1-hexyl, but-1-ene, pent-1-ene, 1-propyl, 1-butyl, 2-methyl-propane, butyric acid ethyl ester, 4-methyl-benzyl, 3-chloro-benzyl, propoxybenzene, 1-(4-methoxy-phenyl)-ethanone, 4-methyl-benzonitrile, 2-methyl-naphthalene, 1-pentyl, methyl-cyclohexane, 3-methyl-benzonitrile, 1-ethoxy-4-chloro-benzene, 5 2-ethyl-butane, 2-hydroxy-ethane, 4-methyl benzoic acid methyl ester, 1-naphthalen-2-yl-ethanone, 2,5-dimethoxy-phenyl-ethanone, 1-p-tolyl-ethanone, 4-fluoro-benzyl, 2-fluoro-benzyl, 5-trifluoromethyl-benzyl, 5-trifluoromethoxy-benzyl, 3-fluoro-5-trifluoromethyl-benzyl, 1-(2-methoxy-phenyl)-ethanone, 1-(2,4-dimethyl-phenyl)-ethanone, 4-chloro-benzyl, 2-difluoromethoxy-benzyl, 4-10 isopropyl-benzyl, 2-fluoro-6-trifluoromethyl-benzyl, 2,3-difluoro-4-methyl-benzyl, 2-methyl-benzyl, 3-methyl-benzyl, pent-2-ene, 6-methyl-2-methyl-quinoline, 2-chloro-benzyl, 3-methoxy-benzyl, 4-methoxy-benzyl, (3-Chloro-phenyl)-ethane, 5-methyl-hexane, ethyl-cyclohexane, pentanoic acid ethyl ester, (propoxymethyl)-benzene, acetamide, 2-ethyl-isoindole-1,3-dione, 2-propyl-15 isoindole-1,3-dione, N-methyl-acetamide, methyl-cyclopropane, but-1-ene, 4-yl-but-1-ene, 2-methyl-pent-2-ene, ethanol, 2-methoxy-ethane, but-2-yne, propyne, acetate, 1-pyrrolidin-1-yl-ethanone, N-benzylacetamide, 1-morpholin-4-yl-ethanone, N-phenyl-acetamide, N-methyl-N-phenyl-acetamide, N-(3-hydroxy-3-methyl-butyl)-acetamide, N-n-propyl-acetamide, N-ethyl-acetamide, 20 N-isopropyl-acetamide, N-butyl-acetamide, N-cyclopentyl-acetamide, N-(3-methyl-butyl)-acetamide, N-(4-methoxy-benzyl)-acetamide, N-(2,2-dimethyl-propyl)-acetamide, N-cyclohexyl-acetamide, N-(3-methoxy-benzyl)-acetamide, N-cycloheptyl-acetamide, N-(2-methoxy-benzyl)-acetamide, N-cyclohexylmethyl-acetamide, N-(2-hydroxy-ethyl)-acetamide, N-(1-phenyl-ethyl)-acetamide, N-(3-hydroxy-propyl)-acetamide, N-(2-methoxy-ethyl)-acetamide, N-(2-dimethylamino-ethyl)-acetamide, N-(3-dimethylamino-propyl)-acetamide, N-(1-phenyl-ethyl)-acetamide, N-(3-isopropoxy-propyl)-acetamide, N-furan-2-ylmethyl-acetamide, N-pyridin-2-ylmethyl-acetamide, N-pyridin-3-ylmethyl-acetamide, N-(2-phenoxy-ethyl)-acetamide, N-pyridin-4-ylmethyl-acetamide, 25 N-(4-ethyl-benzyl)-acetamide, N-(3,5-difluoro-benzyl)-acetamide, N-(2,3-difluoro-benzyl)-acetamide, N-(2-pyridin-2-yl-ethyl)-acetamide, N-(2-methyl-benzyl)-acetamide, N-(3-fluoro-benzyl)-acetamide, N-(3-methyl-benzyl)-acetamide, N-(4-methyl-benzyl)-acetamide, N-phenethyl-acetamide, N-(2-pyridin-4-yl-ethyl)-acetamide, N-(3-phenyl-propyl)-acetamide, N-(2-chloro-30

benzyl)-acetamide, N-(2-piperidin-1-yl-ethyl)-acetamide, N-(3-chloro-benzyl)-acetamide, N-(2-morpholin-4-yl-ethyl)-acetamide, N-(4-chloro-benzyl)-acetamide, N-(2-pyridin-3-yl-ethyl)-acetamide, N-(2-pyrrolidin-1-yl-ethyl)-acetamide, N-(2-acetyl-amino-ethyl)-acetamide, (R)-N-(2-hydroxy-2-phenyl-ethyl)-acetamide, (S)-N-(2-hydroxy-2-phenyl-ethyl)-acetamide, N-thiophen-2-ylmethyl-acetamide, N-[3-(2-oxo-pyrrolidin-1-yl)-propyl]-acetamide, N-(2-hydroxy-indan-1-yl)-acetamide, N-cycloheptylmethyl-acetamide, N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide, N-(4-dimethylamino-butyl)-acetamide, cyclopentan, cyclopropylmethyl, phenyl-ethane, acetic acid benzyl ester, 2-methyl-benzonitrile, 2-(1-oxy-pyridin-4-yl)ethane, (4-pyridyl)ethane, (3-pyridyl)ethane, (2-pyridyl)ethane, (4-benzonitrile)ethane, (4-methylsulfinyl-phenyl)ethane, (4-methylsulfonyl-phenyl)ethane, 1-phenyl-propane, 2-phenyl-propane or 1-methyl-2-phenyl-ethane.

15 16. A compound according to any one of the preceding claims wherein R₂ represents methyl.

17. A compound according to claim 16, wherein R₁₂ represents alkyl, cycloalkyl, hydroxyalkyl, aryl, arylalkyl, alkylcarbonylamino, all of which is optionally substituted with one or more same or different substituents selected from alkyl, cycloalkyl, alkoxy, heterocycloalkyl, heteroaryl, aryloxy, amino, hydroxy, halogen, oxy, all of which is optionally substituted with oxo or hydroxyl, or R₁₂ represents hydrogen.

20 25 18. A compound according to any one of the preceding claims with a molecular weight below 800 Dalton.

19. A compound according to claim 1 selected from 2-(3,5-Dichloro-pyridin-4-yl)-1-(2-hydroxy-3,4-dimethoxy-phenyl)-ethanone (compound 101), 2-(3,5-Dichloro-pyridin-4-yl)-1-(3-hydroxy-2,4-dimethoxy-phenyl)-ethanone (compound 102), 1-(2-Allyloxy-3-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 103),

2-(3,5-Dichloro-pyridin-4-yl)-1-(2,3-diethoxy-4-methoxy-phenyl)-ethanone (compound 104),
{2-tert-Butoxycarbonylmethoxy-6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-3-methoxy-phenoxy}-acetic acid tert-butyl ester (compound 105),
5 1-(2,3-Bis-allyloxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 106),
1-(2,3-Bis-difluoromethoxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 107),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(5-methyl-isoxazol-3-ylmethoxy)-phenyl]-ethanone (compound 108),
10 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methoxy-ethoxy)-phenyl]-ethanone (compound 109),
1-(2-But-2-ynyloxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 110),
15 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methyl-but-2-enyloxy)-phenyl]-ethanone (compound 111),
2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-phenethyloxy-phenyl)-ethanone (compound 112),
1-(2-Benzylxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone
20 (compound 113),
1-(2-Allyloxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 114),
1-[2-(Benzooxazol-2-ylmethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 115),
25 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methyl-thiazol-4-ylmethoxy)-phenyl]-ethanone (compound 116),
1-[2-(5-Cyclopropyl-[1,3,4]thiadiazol-2-ylmethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 117),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-([1,2,4]oxadiazol-3-ylmethoxy)-phenyl]-ethanone (compound 118),
30 {6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid ethyl ester (compound 119),
1-{2-[2-(4-Chloro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 120),

1-[2-(5-Chloro-thiophen-2-ylmethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 121),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-phenoxy-ethoxy)-phenyl]-ethanone (compound 122),
5 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-p-tolyl-ethoxy)-phenyl]-ethanone (compound 123),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-phenyl-propoxy)-phenyl]-ethanone (compound 124),
2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(3-methoxy-phenyl)-ethoxy]-phenyl}-ethanone (compound 125),
10 2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(4-methoxy-phenyl)-ethoxy]-phenyl}-ethanone (compound 126),
1-{2-[2-(3-Bromo-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 127),
15 2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(2-methoxy-phenyl)-ethoxy]-phenyl}-ethanone (compound 128),
2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 129),
2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(2-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 130),
20 2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(3,4-dimethoxy-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 131),
{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid benzyl ester (compound 132),
25 {6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid isopropyl ester (compound 133),
3-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy-methyl}-benzoic acid methyl ester (compound 134),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methyl-butoxy)-phenyl]-ethanone (compound 135),
30 2-(3,5-Dichloro-pyridin-4-yl)-1-(2-hexyloxy-3,4-dimethoxy-phenyl)-ethanone (compound 136),
1-(2-But-3-enyloxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 137),

2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-pent-4-enyloxy-phenyl)-ethanone (compound 138),

2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-propoxy-phenyl)-ethanone (compound 139),

5 1-(2-Butoxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone ethanone (compound 140),

2-(3,5-Dichloro-pyridin-4-yl)-1-(2-isobutoxy-3,4-dimethoxy-phenyl)-ethanone (compound 141),

10 4-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-butyric acid ethyl ester (compound 142),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(4-methyl-benzylxy)-phenyl]-ethanone (compound 143),

1-[2-(3-Chloro-benzylxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 144),

15 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-phenoxy-propoxy)-phenyl]-ethanone (compound 145),

2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(4-methoxy-phenyl)-2-oxo-ethoxy]-phenyl}-ethanone (compound 146),

20 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-methyl-benzonitrile (compound 147),

4-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-methyl-benzonitrile (compound 148),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(naphthalen-2-ylmethoxy)-phenyl]-ethanone (compound 149),

25 2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-pentyloxy-phenyl)-ethanone (compound 150),

1-(2-Cyclohexylmethoxy-3,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 151),

30 3-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-methyl-benzonitrile (compound 152),

1-{2-[2-(4-Chloro-phenoxy)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 153),

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-ethyl-butoxy)-3,4-dimethoxy-phenyl]-ethanone (compound 154),

2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-hydroxy-ethoxy)-3,4-dimethoxy-phenyl]-ethanone (compound 155),
4-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy-methyl}-benzoic acid methyl ester (compound 156),
5 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-naphthalen-2-yl-2-oxo-ethoxy)-phenyl]-ethanone (compound 157),
2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(2,5-dimethoxy-phenyl)-2-oxo-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 158),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-oxo-2-p-tolyl-ethoxy)-phenyl]-ethanone (compound 159),
10 2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(4-fluoro-benzyl-oxo)-3,4-dimethoxy-phenyl]-ethanone (compound 160),
2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-fluoro-benzyl-oxo)-3,4-dimethoxy-phenyl]-ethanone (compound 161),
15 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-trifluoromethyl-benzyl-oxo)-phenyl]-ethanone (compound 162),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-trifluoromethoxy-benzyl-oxo)-phenyl]-ethanone (compound 163),
2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(3-fluoro-5-trifluoromethyl-benzyl-oxo)-3,4-dimethoxy-phenyl]-ethanone (compound 164),
20 2-(3,5-Dichloro-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(2-methoxy-phenyl)-2-oxo-ethoxy]-phenyl}-ethanone (compound 165),
2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(2,4-dimethyl-phenyl)-2-oxo-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 166),
25 1-[2-(4-Chloro-benzyl-oxo)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 167),
2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-difluoromethoxy-benzyl-oxo)-3,4-dimethoxy-phenyl]-ethanone (compound 168),
30 2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(4-isopropyl-benzyl-oxo)-3,4-dimethoxy-phenyl]-ethanone (compound 169),
2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2-fluoro-6-trifluoromethyl-benzyl-oxo)-3,4-dimethoxy-phenyl]-ethanone (compound 170),
2-(3,5-Dichloro-pyridin-4-yl)-1-[2-(2,3-difluoro-4-methyl-benzyl-oxo)-3,4-dimethoxy-phenyl]-ethanone (compound 171),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methyl-benzyloxy)-phenyl]-ethanone (compound 172),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methyl-benzyloxy)-phenyl]-ethanone (compound 173),
5 2-(3,5-Dichloro-pyridin-4-yl)-1-(3,4-dimethoxy-2-pent-2-enyloxy-phenyl)-ethanone (compound 174),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-methyl-quinolin-6-ylmethoxy)-phenyl]-ethanone (compound 175),
1-[2-(2-Chloro-benzyloxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 176),
10 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(3-methoxy-benzyloxy)-phenyl]-ethanone (compound 177),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(4-methoxy-benzyloxy)-phenyl]-ethanone (compound 178),
15 1-{2-[2-(3-Chloro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 179),
2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(5-methyl-hexyloxy)-phenyl]-ethanone (compound 180),
1-[2-(2-Cyclohexyl-ethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 181),
20 5-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-pentanoic acid ethyl ester (compound 182),
1-[2-(3-Benzyl-ethoxy)-3,4-dimethoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 183),
25 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 184),
2-(2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-ethyl)-isoindole-1,3-dione (compound 185),
2-(3-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-propyl)-isoindole-1,3-dione (compound 186),
30 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-methyl-acetamide (compound 187),
2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-2,4-dimethoxy-phenyl)-ethanone (compound 188),

1-(3-Cyclopropylmethoxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 189),

1-(2-Allyloxy-3-but-3-enyloxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 190),

5 1-(3-But-3-enyloxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 191),

2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-propoxy-phenyl)-ethanone (compound 192),

1-(3-Allyloxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone

10 (compound 193),

2-(3,5-Dichloro-pyridin-4-yl)-1-[2,4-dimethoxy-3-(4-methyl-pent-3-enyloxy)-phenyl]-ethanone (compound 194),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3-(2-hydroxy-ethoxy)-2,4-dimethoxy-phenyl]-ethanone (compound 195),

15 2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-phenethyloxy-phenyl)-ethanone (compound 196),

1-(3-Benzylxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 197),

2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-pent-2-enyloxy-phenyl)-

20 ethanone (compound 198),

2-(3,5-Dichloro-pyridin-4-yl)-1-[2,4-dimethoxy-3-(2-methoxy-ethoxy)-phenyl]-ethanone (compound 199),

1-(3-But-2-ynyloxy-2,4-dimethoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 200),

25 2-(3,5-Dichloro-pyridin-4-yl)-1-(2,4-dimethoxy-3-prop-2-ynyloxy-phenyl)-ethanone (compound 201),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-oxo-2-pyrrolidin-1-yl-ethoxy)-phenyl]-ethanone (compound 202),

N-Benzyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-

30 acetamide (compound 203),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-morpholin-4-yl-2-oxo-ethoxy)-phenyl]-ethanone (compound 204),

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-phenyl-acetamide (compound 205),

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-methyl-N-phenyl-acetamide (compound 206),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-hydroxy-3-methyl-butyl)-acetamide (compound 207),
5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 208),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-isopropyl-acetamide (compound 209),
N-Butyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 210),
N-Cyclopentyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 211),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-methyl-butyl)-acetamide (compound 212),
15 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-methoxy-benzyl)-acetamide (compound 213),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2,2-dimethyl-propyl)-acetamide (compound 214),
N-Cyclohexyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 215),
20 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-methoxy-benzyl)-acetamide (compound 216),
N-Cycloheptyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 217),
25 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-methoxy-benzyl)-acetamide (compound 218),
N-Cyclohexylmethyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 219),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-30 hydroxy-ethyl)-acetamide (compound 220),
(R)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(1-phenyl-ethyl)-acetamide (compound 221),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-hydroxy-propyl)-acetamide (compound 222),

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-methoxy-ethyl)-acetamide (compound 223),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-dimethylamino-ethyl)-acetamide (compound 224),
5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-dimethylamino-propyl)-acetamide (compound 225),
(S)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(1-phenyl-ethyl)-acetamide (compound 226),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-isopropoxy-propyl)-acetamide (compound 227),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-furan-2-ylmethyl-acetamide (compound 228),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-pyridin-2-ylmethyl-acetamide (compound 229),
15 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-pyridin-3-ylmethyl-acetamide (compound 230),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-phenoxy-ethyl)-acetamide (compound 231),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-pyridin-
20 4-ylmethyl-acetamide (compound 232),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-ethyl-benzyl)-acetamide (compound 233),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3,5-difluoro-benzyl)-acetamide (compound 234),
25 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2,3-difluoro-benzyl)-acetamide (compound 235),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyridin-2-yl-ethyl)-acetamide (compound 236),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-
30 methyl-benzyl)-acetamide (compound 237),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-fluoro-benzyl)-acetamide (compound 238),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-methyl-benzyl)-acetamide (compound 239),

2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-methyl-benzyl)-acetamide (compound 240),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-phenethyl-acetamide (compound 241),
5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyridin-4-yl-ethyl)-acetamide (compound 242),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(3-phenyl-propyl)-acetamide (compound 243),
N-(2-Chloro-benzyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 244),
10 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-piperidin-1-yl-ethyl)-acetamide (compound 245),
N-(3-Chloro-benzyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 246),
15 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-morpholin-4-yl-ethyl)-acetamide (compound 247),
N-(4-Chloro-benzyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 248),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyridin-3-yl-ethyl)-acetamide (compound 249),
20 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-pyrrolidin-1-yl-ethyl)-acetamide (compound 250),
N-(2-Acetylamino-ethyl)-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 251),
25 (R)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-2-phenyl-ethyl)-acetamide (compound 252),
(S)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-2-phenyl-ethyl)-acetamide (compound 253),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-
30 thiophen-2-ylmethyl-acetamide (compound 254),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-[3-(2-oxo-pyrrolidin-1-yl)-propyl]-acetamide (compound 255),
(2R)-2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(2-hydroxy-indan-1-yl)-acetamide (compound 256),

N-Cycloheptylmethyl-2-{6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 257),
2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-[2-(2-hydroxy-ethoxy)-ethyl]-acetamide (compound 258),
5 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-(4-dimethylamino-butyl)-acetamide (compound 259),
1-(3-Cyclopentyloxy-2-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 260),
1-(3-Cyclopropylmethoxy-2-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 261),
10 2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-2-hydroxy-4-methoxy-phenyl)-ethanone (compound 262),
2-(3,5-Dichloro-pyridin-4-yl)-1-(3-ethoxy-4-methoxy-2-phenethyloxy-phenyl)-ethanone (compound 263),
15 1-[2-(5-Cyclopropyl-[1,3,4]thiadiazol-2-ylmethoxy)-3-ethoxy-4-methoxy-phenyl]-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 264),
{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2-ethoxy-3-methoxy-phenoxy}-acetic acid benzyl ester (compound 265),
1-(3-Allyloxy-2-hydroxy-4-methoxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 266),
20 2-{2-Allyloxy-6-[2-(3,5-dichloro-pyridin-4-yl)-acetyl]-3-methoxy-phenoxy}-benzonitrile (compound 267),
1-(3-Allyloxy-4-methoxy-2-phenethyloxy-phenyl)-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 268),
25 1-{3-Allyloxy-2-[2-(4-fluoro-phenyl)-ethoxy]-4-methoxy-phenyl}-2-(3,5-dichloro-pyridin-4-yl)-ethanone (compound 269),
N-Benzyl-2-{6-[2-(3,5-dichloro-1-oxy-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetamide (compound 270),
2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-(3,4-dimethoxy-2-phenethyloxy-phenyl)-ethanone (compound 271),
30 2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 272),
2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-{3,4-dimethoxy-2-[2-(1-oxy-pyridin-4-yl)-ethoxy]-phenyl}-ethanone (compound 274),

2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-1-(2-hydroxy-3,4-dimethoxy-phenyl)-ethanone (compound 275),

4-(2-{6-[2-(3,5-Dichloro-1-oxy-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-ethyl)-benzonitrile (compound 276),

5 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-pyridin-4-yl-ethoxy)-phenyl]-ethanone (compound 277),

4-(2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-ethyl)-benzonitrile (compound 278),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-pyridin-2-yl-ethoxy)-phenyl]-ethanone (compound 279),

10 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-pyridin-3-yl-ethoxy)-phenyl]-ethanone (compound 280),

2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(4-methanesulfinyl-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 281),

15 2-(3,5-Dichloro-pyridin-4-yl)-1-{2-[2-(4-methanesulfonyl-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 282),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(1-phenyl-propoxy)-phenyl]-ethanone (compound 283),

2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(2-phenyl-propoxy)-phenyl]-ethanone (compound 284),

20 2-(3,5-Dichloro-pyridin-4-yl)-1-[3,4-dimethoxy-2-(1-methyl-2-phenyl-ethoxy)-phenyl]-ethanone (compound 285),

2-{6-[2-(6-Chloro-pyrazin-2-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 286),

25 2-{6-[2-(3-Bromo-pyrazin-2-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 287),

2-{6-[2-(2,6-Dichloro-phenyl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 288),

2-[2,3-Dimethoxy-6-(2-pyridin-4-yl-acetyl)-phenoxy]-N-propyl-acetamide

30 (compound 289),

2-[2,3-Dimethoxy-6-(2-quinolin-4-yl-acetyl)-phenoxy]-N-propyl-acetamide

(compound 290),

2-[2,3-Dimethoxy-6-(2-pyrazin-2-yl-acetyl)-phenoxy]-N-propyl-acetamide

(compound 291),

2-{6-[2-(3-Bromo-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 292),
2-{6-[2-(3,5-Dibromo-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 293),
5 2-{6-[2-(6-Chloro-pyrimidin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 294),
2-{6-[2-(4-Chloro-pyridin-2-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 295),
2-{6-[2-(2-Chloro-pyridin-3-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 296),
10 2-{2,3-Dimethoxy-6-[2-(2-methoxy-pyridin-4-yl)-acetyl]-phenoxy}-N-propyl-acetamide (compound 297),
2-{6-[2-(2-Cyano-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-propyl-acetamide (compound 298),
15 2-[2,3-Dimethoxy-6-(2-pyridazin-3-yl-acetyl)-phenoxy]-N-propyl-acetamide (compound 299),
2-(2-tert-Butylamino-3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-dimethoxy-phenyl}-ethanone (compound 300),
2-(2-amino-3,5-dichloro-pyridin-4-yl)-1-{2-[2-(4-fluoro-phenyl)-ethoxy]-3,4-
20 dimethoxy-phenyl}-ethanone (compound 301),
2-(3,5-Dichloro-pyridin-4-yl)-1-(4-ethoxy-3-methoxy-2-phenethoxy-phenyl)-ethanone (compound 302),
{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-acetic acid (compound 504),
25 2-tert-butoxycarbonylmethoxy-3,4-dimethoxy-benzoic acid methyl ester (compound 506a),
2-carboxymethoxy-3,4-dimethoxy-benzoic acid methyl ester (compound 506b),
3,4-Dimethoxy-2-propylcarbamoylmethoxy-benzoic acid methyl ester (compound 506c), or
30 2-{6-[2-(3,5-Dichloro-pyridin-4-yl)-acetyl]-2,3-dimethoxy-phenoxy}-N-ethyl-acetamide (compound 305),
and pharmaceutically acceptable and physiologically cleavable esters,
pharmaceutically acceptable salts, hydrates, N-oxides or solvates thereof.

20. A compound according to any one of claims 1-19 for use in therapy.


21. A compound according to any one of claims 1-19 for use in treatment of dermal diseases.

5 22. A pharmaceutical composition comprising a compound according to any one of claims 1-19 together with a pharmaceutically acceptable vehicle or excipient or pharmaceutically acceptable carrier(s).

10 23. A pharmaceutical composition according to claim 22 together with one or more other therapeutically active compound(s).

24. The use of a compound according to formula I

15



I

wherein X_1 , X_2 , X_3 , X_4 and X_5 independently of each other represent $-CH-$ or N ;

20 or X_3 , X_4 and X_5 independently of each other represent $-CH-$ or N , and X_1 and X_2 independently of each other represent C and form part of an additional 6-membered aromatic ring;

wherein R₁ represents alkyl, alkenyl, alkynyl, haloalkyl, hydroxyalkyl, or alkylcarbonyl, all of which are optionally substituted by one or more, same or different substituents selected from R₄, or R₁ represents hydrogen;

5 R₂ represents alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, hydroxyalkyl, heterocycloalkenyl, alkylaryl, arylalkyl, alkylalkoxycarbonyl, alkylcarbonyloxy, or alkoxyalkyl, all of which are optionally substituted by one or more, same or different substituents selected from R₅; or R₂ represents hydrogen or -CH₂-C(O)NR₉-R₁₂;

10 R₃ represents alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, hydroxyalkyl, heterocycloalkenyl, alkylaryl, arylalkyl, alkylalkoxycarbonyl, alkylcarbonyloxy, or alkoxyalkyl, all of which are optionally substituted by one or more, same or different substituents selected from R₆; or R₃ represents hydrogen, -CH₂-C(O)-heterocycloalkyl or -CH₂-C(O)NR₉-R₁₂;

R₄ represents hydrogen, alkyl, alkenyl, alkynyl, halogen, oxo, alkoxy, hydroxy, or haloalkyl;

20 R₅ represents alkylaryl, carboxy, alkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, carbamoyl, hydroxyalkyl, aryloxy, alkoxy carbonyloxy, alkoxy carbonyl, alkoxy, alkoxyalkyl, aryl, a heterocyclic ring, aminocarbonyl, alkylthio, alkylcarbonylamino, hydroxy, alkylcarbonyl, arylcarbonyl, alkylcarbonyloxy, or amino, all of which are optionally substituted by one or more, same or different substituents selected from R₇;

25 or R₅ represents hydrogen, oxo, halogen, cyano, or nitro;

30 R₆ represents alkylaryl, carboxy, alkyl, alkenyl, cycloalkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, carbamoyl, hydroxyalkyl, aryloxy, alkoxy carbonyloxy, alkoxy carbonyl, alkoxy, alkoxyalkyl, aryl, a heterocyclic ring, aminocarbonyl, alkylthio, alkylcarbonylamino, arylcarbonyl, hydroxy, alkylcarbonyl, alkylcarbonyloxy, or amino, all of which are optionally substituted by one or more, same or different substituents selected from R₈;

or R₆ represents hydrogen, oxo, halogen, cyano, or nitro;

R₇ represents alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkynyl, alkoxy, haloalkyl, alkylthio, heterocycloalkenyl, heterocycloalkyl, aryl, alkylcarbonyl, heteroaryl, aryloxy, alkoxycarbonyl, hydroxyalkyl, amino, hydroxy, or carboxy; all of which
5 are optionally substituted by one or more, same or different substituents selected from R₁₀;
or R₇ represents hydrogen, halogen, or oxo;

R₈ represents alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkynyl, alkoxy, haloalkyl, alkylthio, alkylsulfonyl, alkylsulfinyl, heterocycloalkenyl, heterocycloalkyl, aryl, alkylcarbonyl, heteroaryl, aryloxy, alkoxycarbonyl, hydroxyalkyl, amino, hydroxy, or carboxy; all of which are optionally substituted by one or more, same or different substituents selected from R₁₀;
10 or R₈ represents hydrogen, halogen, or oxo;

R₉ represents hydrogen, alkyl, haloalkyl, or hydroxyalkyl;

R₁₀ represents hydrogen, alkyl, oxo, hydroxy, halogen, carboxy, amino, alkoxy, haloalkyl, or hydroxyalkyl;

R₁₁ represents one or more, same or different substituents selected from hydrogen, halogen, cyano, amino, alkyl, methylsulfinyl, methylsulfonyl, amino, cyano, or alkoxy;

R₁₂ represents alkylaryl, arylalkyl, carboxy, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, carbamoyl, hydroxyalkyl, aryloxy, alkoxycarbonyloxy, alkoxycarbonyl, alkoxy, alkoxyalkyl, aryl, a heterocyclic ring, aminocarbonyl, alkylthio, alkylcarbonylamino, hydroxy, alkylcarbonyl, arylcarbonyl, alkylcarbonyloxy, or amino, all of which are optionally substituted by one or
25 more, same or different substituents selected from R₈, or R₁₂ represents hydrogen;

30 with the proviso R₁, R₂ and R₃ cannot all be methyl;

with the proviso that when R_2 and R_3 are both hydrogen, R_1 cannot be methyl or hydrogen;

and pharmaceutically acceptable and physiologically cleavable esters,

5 pharmaceutically acceptable salts, hydrates, N-oxides or solvates thereof

in the manufacture of a medicament for the prophylaxis, treatment or amelioration of dermal diseases or conditions, or acute or chronic cutaneous wound disorders.

10

25. The use according to claim 24, wherein the dermal disease or condition is selected from the group consisting of proliferative and inflammatory skin disorders, psoriasis, cancer, epidermal inflammation, alopecia, skin atrophy, steroid induced skin atrophy, skin ageing, photo skin ageing, acne, dermatitis, 15 atopic dermatitis, seborrheic dermatitis, contact dermatitis, urticaria, pruritis, and eczema.

26. A method of preventing, treating or ameliorating dermal diseases or conditions, or acute or chronic cutaneous wound disorders, the method

20 comprising administering to a person suffering from at least one of said diseases an effective amount of one or more compounds according to according to any one of claims 1-19, optionally together with a pharmaceutically acceptable carrier or one or more excipients, optionally in combination with other therapeutically active compounds.

25

27. A method according to claim 26, wherein the dermal disease or condition is selected from the group consisting of proliferative and inflammatory skin disorders, psoriasis, cancer, epidermal inflammation, alopecia, skin atrophy, steroid induced skin atrophy, skin ageing, photo skin ageing, acne, dermatitis, 30 atopic dermatitis, seborrheic dermatitis, contact dermatitis, urticaria, pruritis, and eczema.

INTERNATIONAL SEARCH REPORT

International application No
PCT/DK2007/000564

A. CLASSIFICATION OF SUBJECT MATTER				
INV.	C07D213/61	C07D413/12	C07D417/12	A61K31/44
	A61P25/00	A61P29/00	A61P31/00	A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 943 613 A (KYOWA HAKKO KOGYO KK [JP]) 22 September 1999 (1999-09-22) cited in the application claims; examples	1-27
A	WO 2006/135828 A (MEMORY PHARMACEUTICALS CORP [US]; HOFFMANN LA ROCHE [CH]; TALAMAS FRAN) 21 December 2006 (2006-12-21) claims; examples	1-27
A	WO 95/20578 A (RHONE POULENC RORER LTD [GB]; FENTON GARRY [GB]; PALFREYMAN MALCOLM NO) 3 August 1995 (1995-08-03) cited in the application the whole document	1-27

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
17 March 2008	26/03/2008
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer Bosma, Peter

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK2007/000564

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 26 and 27 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers allsearchable claims.
2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.

The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/DK2007/000564

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
EP 0943613	A 22-09-1999	AU	4967697 A	10-06-1998
		CA	2272327 A1	28-05-1998
		WO	9822455 A1	28-05-1998
WO 2006135828	A 21-12-2006	AU	2006257863 A1	21-12-2006
		CA	2611562 A1	21-12-2006
		EP	1888528 A2	20-02-2008
WO 9520578	A 03-08-1995	AT	164575 T	15-04-1998
		AU	1463195 A	15-08-1995
		DE	69501947 D1	07-05-1998
		DE	69501947 T2	27-08-1998
		EP	0741707 A1	13-11-1996
		JP	3953096 B2	01-08-2007
		JP	9509654 T	30-09-1997
		ZA	9500639 A	26-07-1996