




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07H 19/01, A61K 31/71		A1	(11) International Publication Number: WO 93/18049
			(43) International Publication Date: 16 September 1993 (16.09.93)
(21) International Application Number: PCT/US93/00428 (22) International Filing Date: 27 January 1993 (27.01.93) (30) Priority data: 844,802 2 March 1992 (02.03.92) US 947,239 18 September 1992 (18.09.92) US (60) Parent Applications or Grants (63) Related by Continuation US 844,802 (CIP) Filed on 2 March 1992 (02.03.92) US 947,239 (CIP) Filed on 18 September 1992 (18.09.92) (71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).		(72) Inventor; and (75) Inventor/Applicant (for US only): KOCH, Kevin [US/US]; 34 Linda Avenue, Mystic, CT 06340 (US). (74) Agents: RICHARDSON, Peter, C. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US). (81) Designated States: CA, FI, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report.	

(54) Title: FLUOROSUGAR DERIVATIVES OF MACROLIDES



(57) Abstract

Macrolides of formula (I) and methods of treatment of resistance to transplantation, fungal infections and autoimmune diseases such as rheumatoid arthritis and psoriasis using said macrolides of formula (I), wherein n is 1 or 2; A and B are taken together and form =O or A and B are taken separately and are each OH or A is OH and B is H; R¹ is a fluoroglycosyl group; R² is OH or a fluoroglycosyloxy group; and R³ is an alkyl or allyl group.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

5

FLUOROSUGAR DERIVATIVES OF MACROLIDESBackground of the Invention

This invention relates to new chemical compounds which have value in the field of medical science. More particularly, it relates to new chemical compounds which are of value for administration to a mammalian subject, particularly man, as immunosuppressive agents. These new immunosuppressive agents can be compared to the macrolides known as FK-506 and FK-520, which are described in further detail in United States Patent No. 4,894,366. The new compounds of this invention will find especial utility in preventing or treating graft rejection following skin or organ transplant surgery and in preventing or treating autoimmune diseases such as rheumatoid arthritis and psoriasis. Additionally, these macrolide derivatives will find use in preventing or treating infectious diseases caused by fungi.

Graft or organ transplant rejection following transplant surgery is a common occurrence which arises when foreign antigens are recognized by the host's immune response system. The host's immune response system, in an effort to "protect" itself from the foreign tissue, then releases its cellular and humoral arsenal. Both activated lymphocytes and antibodies attack the foreign tissue, resulting in complications which often end in rejection of said tissue.

Similarly, the occurrence of immunoregulatory irregularities in autoimmune and chronic inflammatory diseases is well known. Irrespective of the underlying etiology of the condition, a variety of autoantibodies and self-reactive lymphocytes often arise to complicate the condition.

Treatments which target the immune response system often result in a complete shutdown of the system, leading to a lowering of the body's ability to combat infection. This can be as dangerous as the original condition which led to the shutdown.

Currently the leading medicinal agent for the prevention or treatment of graft rejection is cyclosporin A, approved by the United States Food and Drug Administration in 1983. The drug acts by inhibiting the body's immune response system from mobilizing its arsenal of natural protecting agents to reject the transplant's foreign protein. Although cyclosporin is effective in fighting graft rejection, it suffers drawbacks in that it can cause kidney failure, liver damage and ulcers; which in many cases can be very severe. Safer drugs which are more selective in their ability to affect the

-2-

immune response system and which have fewer side effects are constantly being pursued.

United States Patent No. 4,894,366 discloses the macrolides FK-506 and FK-520, *inter alia*, as immunosuppressants, including the treatment of "resistance to transplantation," autoimmune diseases and infectious diseases. International Patent Publication No. WO 91/02736 discloses derivatives of FK-506, FK-520 and related macrolides. European Patent Publication No. 428,365 A1 discloses various other derivatives of FK-506, FK-520 and related macrolides.

10


Summary of the Invention

The present invention is directed to compounds of the formula

15

20

25



(I)

or a pharmaceutically acceptable salt thereof;

wherein n is 1 or 2;

30 the dotted line represents an optional double bond in the case where R² is H;

A and B are taken separately and A is H and B is H or OH, or A and B are taken together and form =O;


R² is H, (C₂-C₅)alkanoyloxy or -OR⁰;

-3-


R^3 is (C_1 to C_3)alkyl or allyl;

R^1 and R^0 are each H,

5



or



10

R^4 is, for each occurrence, independently $-CO_2R^8$, $-CO_2H$,

$-CH_2OH$, H, $-CH_3$, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CONH_2$, $-CONHR^8$, $-CON(R^8)_2$,

$-CH_2OCOR^8$, $-CH_2OCO_2R^8$, $-CH_2OCONR_2^8$ or $-CH_2OR^8$;

15 R^5 is, for each occurrence, independently (C_1 to C_4)alkoxy, benzyloxy, $-OH$, $-OCOR^8$, $-OCOCH_2R^8$, $-OCO_2R^8$ or $-OSi(R^8)_3$;

t is 1, 2 or 3;

m is 0 or 1; and

20 R^8 is (C_1 - C_6)alkyl, (C_3 - C_6)cycloalkyl, allyl, pyridyl, thienyl, benzyl, benzyl variously substituted with one to five halogen atoms, $-OH$ groups or (C_1 - C_4)alkoxy groups, phenyl or phenyl variously substituted with one to five halogen atoms, $-OH$ groups or (C_1 - C_4)alkoxy groups;

provided that R^1 and R^0 are not both H; when m is 0, R^4 is


25 $-CH_2F$, $-CHF_2$ or $-CF_3$; and in formulae (II) and (III), each ring carbon must bear at least one hydrogen atom.

A preferred group of compounds of this invention is the group of compounds of formula (I) wherein n is 2; the dotted line represents no bond; A and B are taken together and form $=O$;


R^1 is

-4-

5



10




R^2 is -OH; R^3 is ethyl; R^4 is H, $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{F}$, $-\text{CH}_2\text{OCOCH}_3$ or $-\text{CH}_2\text{OCH}_2\text{C}_6\text{H}_5$; and R^5 is -OH, $-\text{OCOCH}_2\text{C}_6\text{H}_5$ or $-\text{OCOCH}_3$.

Especially preferred within this group are the compounds of the formula (I)

15 where R^1 is

-5-



A second preferred group of compounds of this invention is the group of
 20 compounds of formula (I) wherein n is 2; A and B are taken separately and A is H and
 B is H; the dotted line represents no bond; R² is -OH; and R³ is ethyl.

The compounds of formula I are active as immunosuppressants. This activity
 makes these compounds useful in treating and preventing graft and transplant
 rejection. Further, this activity makes these compounds useful in preventing and
 25 treating autoimmune diseases such as rheumatoid arthritis and psoriasis in a mammal,
 especially man.

Accordingly this invention also embraces a method of treating resistance to
 transplantation in a mammal in need of such treatment comprising administering to said
 mammal a compound of formula (I) or a pharmaceutically acceptable salt thereof.

30 The term "transplantation," when used above and hereinafter, refers to the
 implantation in one part of an individual of a tissue or organ taken from another part
 of that individual or from another individual. Typical transplantations include, but are

-6-

The term "graft" when used above and hereinafter, refers to any unattached tissue or organ which is used for transplantations. Typical grafts include, but are not limited to, skin, bone, fat and nerve grafts.

5 Additionally this invention embraces a method of treating autoimmune disease (such as rheumatoid arthritis or psoriasis) in a mammal in need of such treatment comprising administering to said mammal a compound of formula (I) or a pharmaceutically acceptable salt thereof.

10 Further, this invention embraces a pharmaceutical composition comprising a resistance to transplantation treating effective amount of a compound of formula (I) and a pharmaceutically acceptable carrier.

Still further this invention embraces a pharmaceutical composition comprising an autoimmune disease (such as rheumatoid arthritis or psoriasis) treating effective amount of a compound of formula (I) and a pharmaceutically acceptable carrier.

15 Yet further the compounds of this invention of formula (I) have antifungal activity. Hence these compounds can be used to treat or prevent infections in mammals caused by fungi.

Accordingly, this invention embraces a method of treating diseases caused by fungi in a mammal in need of such treatment comprising administering to said mammal an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

20 Additionally, this invention embraces a pharmaceutical composition comprising a fungal infectious disease treating effective amount of a compound of formula (I) and a pharmaceutically acceptable carrier.


25

Detailed Description


The compounds of formula (I) of the present invention are readily prepared. Most generally, a macrolide of formula (IV) or (V) below is coupled with an appropriate sugar halide derivative of the formula

30

-7-




- 10 wherein X is halo,
e.g., chloro or bromo. The coupled (or glycosylated) macrolide is then further modified
as described hereinbelow.



30

-8-



15

The production of macrolides of formulae (IV) and (V) is well-known in the literature. The generally preferred route to these macrolides is *via* biological fermentation of microorganisms belonging to the genus *Streptomyces*. The compounds of formulae (IV) and (V) wherein R³ is allyl are obtained by fermentation of *Streptomyces tsukubaensis* No. 9993 (Ferm BP-927). The compound of formula (IV) wherein R³ is ethyl and the compound for formula (IV) wherein R³ is methyl are obtained by fermentation of *Streptomyces hygroscopicus* subsp. *ascomyceticus* ATCC 14891.

A lyophilized sample of *Streptomyces hygroscopicus* subsp. *ascomyceticus* ATCC 14891 has been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland, 20852, U.S.A., under the terms of the Budapest Treaty on January 13, 1992. This newly deposited culture was given the new deposit number of ATCC 55276.

Streptomyces tsukubaensis No. 9993 (Ferm BP-927) is currently on deposit with the Fermentation Research Institute, Agency of Industrial Science and Technology (No. 1-3, Higashi-1-chome, Yatabemachi, Tsukuba-gun, Ibaraki Prefecture, Japan), under the provisions of the Budapest Treaty. A fresh sample of the microorganism will

be deposited with the American Type Culture Collection in accordance with the terms of the Budapest Treaty.

The above-mentioned microorganisms, when placed separately in aqueous nutrient media, will produce the aforementioned compounds of formulae IV and V. The fermentation of said microorganisms to produce these macrolides is accomplished substantially as disclosed in U.S. Patent 4,894,366, which is hereby incorporated by reference. Any changes made to the disclosed procedure are made in order to accommodate existing equipment at the facility and are described in Preparations 1 and 2 hereinbelow.

10 To prepare the compound of formula (I) wherein R^0 is H and R^1 is a sugar substituent of formula (II) or (III), a macrolide of formula (IV) or (V) is coupled with a sugar halide of formula (VI) or (VII).

The coupling (or glycosylation) reaction of the sugar halides of formula (VI) or (VII) and the macrolide of formula (IV) or (V) is accomplished in a straightforward manner, using chemistry well known to one of ordinary skill in the art. The coupling reaction is generally carried out using the peracetylated form of the fluorosugar. About 2-4 molar equivalents of the appropriate sugar halide of formula (VI) or (VII) is mixed with the macrolide of formula (IV) or (V) in a reaction inert solvent. Reaction inert solvents useful for this type of reaction include chlorinated solvents such as chloroform, 20 methylene chloride and ethylene dichloride; ether solvents such as diethyl ether, tetrahydrofuran, dioxane and dimethoxyethane; aromatic solvents such as benzene, toluene and xylene; and dipolar aprotic solvents such as N,N-dimethylformamide, acetonitrile and N-methylpyrrolidone. Preferred solvents are chlorinated solvents and a particularly preferred solvent is methylene chloride. Generally it is desirable to employ 25 enough solvent such that the reactants are dissolved or suspended by the solvent. Typically the amount of solvent used is varied to give a 10^{-1} to 10^{-3} Molar solution of macrolide with 10^{-1} Molar being preferred. Dry conditions are maintained during the course of the reaction by the utilization of anhydrous solvents and by the addition of a drying agent to the reaction mixture. Drying agents typically used for this purpose 30 are molecular sieves, calcium sulfate and magnesium sulfate. A preferred drying agent is 4Å molecular sieves.

Initial mixing of the reagents is performed at a temperature of from about -78°C to about 70°C . Preferred are temperatures ranging from about -78°C to about 0°C .

-10-

Especially preferred for ease of preparation is a cooling bath which maintains the reaction temperature at -78°C .

After the above-mentioned reactants have been mixed and the temperature has equilibrated to -78°C , the reaction mixture is treated with a suitable base such as
5 mercuric carbonate, silver carbonate, mercuric nitrate or silver nitrate. The preferred base for this reaction is silver carbonate. Following addition of said base, the reaction mixture is treated with a catalyst. Typical catalysts for this reaction include triflate, perchlorate and tetrafluoroborate salts of the cation associated with the particular base used. The preferred catalyst is silver triflate.

10 After all reactants and reagents have been added, the reaction mixture is warmed to 0°C , stirred for 0.5-24 hours at 0°C and then warmed slowly to room temperature. The reaction mixture is stirred for an additional 0.5-24 hours at room temperature. Generally, the reaction mixture is stirred at 0°C for 5 hours and allowed to warm to room temperature over 3 hours followed by stirring at room temperature for
15 16 hours. The product is then isolated from the reaction mixture using techniques familiar to one of ordinary skill in the art. Thus, simple filtration through a filter aid such as Celite[®] followed by evaporation affords a residue which is purified by column chromatography. One of ordinary skill in the art will recognize that column chromatography entails the use of a solid phase component such as silica gel and a
20 liquid phase component comprised of an advantageous mixture of solvents for the separation and purification of compounds from a mixture. Removal of solvents after chromatography affords the fluoroglycosylmacrolide.

Generally, the coupling reaction only takes place at one of the three alcohol sites of the macrolide, this site being the C-4' alcoholic functionality (see Formula IV).
25 This selectivity is possibly due to the greater availability of the hydroxyl group of this position in the macrolide's preferred conformation. On occasion, however, with particularly reactive sugar halides, small amounts of diglycosylated material (wherein $\text{R}^2 = \text{OR}^0$) are formed. This material is detected during the monitoring of the progress of the reaction, which is generally accomplished *via* thin layer chromatography, according to
30 standard practice. The diglycosylated material is isolated and purified as for the monoglycosylated material with the notable exception that the diglycosylated material is the first material isolated from the chromatography, with the monoglycosylated material being isolated in later fractions. The use of added equivalents of sugar

chloride or the altering of other parameters such as solvent, base or catalyst can affect the yield of diglycosylated material.

To prepare the compound of formula (I) wherein R^1 is H and R^0 is a sugar substituent of the formula (II) or (III), a macrolide of formula (IV) or (V) is first protected with a hydroxyl protecting group at the C-4" position. Hydroxyl protecting groups suitable for such purposes include but are not limited to such groups as silyl ethers, carboxylic esters and carbonic esters of the alcohol. The protecting groups are appended to the alcohol utilizing the well known methods of organic chemistry. Bulky silyl ethers are preferred for their selectivity, ease of attachment and ease of removal.

Conveniently, a macrolide of formula (IV) or (V) is dissolved in a reaction inert solvent at a temperature of about 0°C to about 30°C. Reaction inert solvents for this type of reaction include dipolar aprotic solvents such as dimethylformamide, acetonitrile and N-methyl pyrrolidone; chlorinated solvents such as chloroform, dichloromethane and 1,2-dichloroethane; and ether solvents such as diethyl ether, dioxane and tetrahydrofuran. The solvent of choice is often dimethylformamide. A silylating agent, usually a silyl chloride such as dimethyl-t-butylsilyl chloride, trimethylchlorosilane or triphenylchlorosilane, is added along with an organic amine such as triethylamine, trimethylamine or imidazole. Ordinarily imidazole is the preferred base. The reaction mixture is stirred for about one hour to about 24 hours, typically at room temperature, after which time the product is isolated from the reaction broth in a manner well known to one of ordinary skill in the art.

The macrolide, now protected at the C-4" position, can be coupled with a sugar halide of formula (VI) or (VII) as described hereinabove. The product of such a coupling reaction is a derivative of a compound of formula (I) with a sugar derivative attached by way of oxygen to the C-14 position and with a protected C-4" position. The C-4" position can be deprotected to afford the free hydroxy compound by employing standard methods of organic chemistry well known to one of ordinary skill in the art. Typically, to remove a preferred silyl ether protecting group, the C-4"-silyl protected compound of formula (I) is dissolved in an ether solvent such as tetrahydrofuran or diethyl ether at a temperature of about 0°C to 30°C and is treated with a fluoride source such as tetra-N-butylammonium fluoride. The reaction is stirred for about one hour to about 24 hours and the product is then isolated by employing standard methods of organic chemistry well known to one of ordinary skill in the art.

-12-

To prepare the compounds of the invention of formula (I) wherein A and B are taken separately and are each H (hereinafter referred to as the C-2 desoxo macrolide), a compound of formula (I) wherein A and B are taken together and are =O is reduced using standard conditions for the reduction of α -ketoamides. This reduction procedure selectively reduces the carbonyl adjacent to the amide without affecting other carbonyls in the molecule. Generally the macrolide is dissolved in a reaction inert solvent or mixture of solvents and hydrogen sulfide gas is bubbled through the mixture for 6-24 hours at room temperature. For convenience, the gas is generally bubbled through the reaction mixture overnight. Suitable reaction inert solvents for this reaction include, but are not limited to, organic bases such as diethylamine, triethylamine, dimethylamine, trimethylamine and aniline; dipolar aprotic solvents such as N,N-dimethylformamide, dimethylsulfoxide and N-methylpyrrolidone; and alcoholic solvents such as methanol, ethanol and propanol. A combination of two or more of these solvents is sometimes used to achieve optimum yield or to affect the course of reduction. For example, the macrolide wherein A is H and B is OH is prepared by using methanol as solvent. A particularly preferred solvent system for providing the C-2 desoxo macrolide is pyridine and N,N-dimethylformamide in equal amounts. When the reaction is completed, the product is isolated using the standard techniques of organic chemistry as would be understood by one skilled in the art.


Alternatively, the macrolide of formula (IV) or (V) can be reduced prior to glycosylation, using the foregoing procedure. Following reduction, the macrolide can be glycosylated as recited hereinabove.

To prepare compounds of the invention of formula (I) wherein the dotted line represents a bond and R² is hydrogen, the compound of formula (I) wherein R² is -OH and the dotted line represents no bond (hereinafter referred to as the β -hydroxy ketone) is dehydrated as disclosed in European Patent Application No. 323042. Generally the β -hydroxy ketone is dissolved in a reaction inert solvent containing a catalytic amount of an organic acid. Suitable reaction inert solvents are aromatic solvents such as benzene, toluene, xylene and the like, with toluene being preferred. The organic acid is generally selected from such acids as toluenesulfonic acid, camphorsulfonic acid and the like with toluenesulfonic acid being preferred. The reaction mixture is heated at about 50°C to about 120°C for about five minutes to about one hour. Generally steam bath temperatures (about 100°C) are preferred and


-13-

five minutes is generally sufficient for complete reaction. The reaction product is isolated according to methods well understood by one of ordinary skill in the art. The reaction is generally carried out on compounds which have already been glycosylated.

- To prepare compounds of the invention of formula (I) wherein R^5 is hydroxy and R^4 is hydroxymethyl, a compound of formula (I) wherein R^5 is acetoxy and R^4 is acetoxymethyl is deacetylated using standard conditions known to one of ordinary skill in the art as recited hereinbelow. This selective deacetylation does not affect any amides which are present and is readily accomplished by the addition of an alkoxide base to a solution of the material to be deacetylated in an alcoholic solvent at 0°C .
- Generally a catalytic amount, such as 0.01 equivalents, of base is used. Usually the alkoxide base of the particular alcoholic solvent in use is preferred. Most preferred, for its ease of use and reactivity, is the system wherein methanol is the solvent and sodium methoxide is the base. Isolation of the product is achieved *via* standard methods well known to those of ordinary skill in the art.
- The fluorosugar halide derivatives of formulae (VI) and (VII) are conveniently prepared from the fluorosugar derivatives of formulae (VIII) and (IX) wherein R^6 is H, (C_1-C_4) alkyl or (C_2-C_4) alkanoyl by employing standard methods of halogenation well known to one of ordinary skill in the art.



VIII



IX

- Bromination is the method of choice; chlorination may also be employed in certain cases. Bromination is effected by dissolving a 1-hydroxy, alkoxy or alkanoyloxy fluorosugar derivative in an organic acid solvent such as acetic acid. When the sugar is a 1-hydroxy or 1-alkoxy sugar, one to ten equivalents of acetic anhydride are generally added. The preferred substrates are 1-acetoxy sugar derivatives. The reaction mixture is cooled so that the temperature falls within the range of about -20°C

to about 0°C. The generally preferred temperature is about 0°C. The cooled reaction mixture is treated with a solution of hydrobromic acid in the acidic solvent. Generally a large excess, such as 10-40 molar equivalents, of hydrobromic acid is employed. The reaction mixture is warmed to room temperature and stirred until the reaction is
5 complete. Generally, for convenience, the reaction mixture is left stirring overnight. The isolation of the brominated product is achieved in a straightforward manner well known to one of ordinary skill in the art. Often this merely involves removing the solvent *in vacuo*. Occasionally, to more fully effect solvent removal, a cosolvent such as toluene, which azeotropes the reaction solvent, is utilized. Further purification is sometimes
10 achieved by the use of column chromatography.

To prepare compounds of formulae (VIII) and (IX) wherein R⁴ is -CH₂OCOR⁸, R⁵ is -OCOR⁸ and R⁶ is -COR⁸, a compound of formulae (VIII) or (IX) wherein R⁴ is -CH₂OH, R⁵ is -OH and R⁶ is H is used as a substrate in a standard acylation reaction. Typically, said substrate is reacted with a suitable acylating agent such as, but not
15 limited to, acetic anhydride, in a suitable solvent such as acetic acid at about 0°C to about 25°C. The product is isolated utilizing the standard techniques of organic chemistry well known to one of ordinary skill in the art.

The fluorosugar compounds of formulae (VIII) and (IX) can be prepared as taught in the literature. (See, for example, Kovac, P.; Yeh, H.J.C. and
20 Glaudemans, C.P.J., Carbohydrate Research, 140, 277 (1985) and Sharma, M. and Korytnyk, W., Tetrahedron Letters, 1977, 573.)

Sugar derivatives of formula (VIII) wherein m is 1 and the fluorine atom is attached to the 4-position of the sugar molecule can be prepared as described hereinbelow.

25 In order to prepare a 4-fluorosugar derivative (a compound of formula (VIII) wherein m is 1 and the fluorine atom is attached to the 4-position) a methyl pyranoside (such as methyl glucopyranoside or methyl galactopyranoside) is regioselectively acetylated at the 2,3 and 6 hydroxyl groups. The remaining hydroxyl group is replaced with fluoride *via* S_N² substitution and the methoxy group at the
30 1-position is replaced with an acetoxy group to afford the tetraacetylfluoropyranoside.


Thus, an alkyl pyranoside (such as methyl- α -D-glucopyranoside or methyl- α -D-galactopyranoside) is activated with a bulky organometallic agent such as bis(tributyltin)oxide. The reaction is refluxed in a water-azeotroping solvent such as toluene,

benzene or xylene. Toluene is the most preferred solvent. After about 1 hour to about 24 hours, and generally after about 3 hours, all of the non-hindered hydroxyl groups have been activated. The reaction mixture is treated with exactly one equivalent of acetyl chloride for every activated hydroxyl group. Generally, for a six-carbon sugar this
5 requires the addition of 3 equivalents of acetyl chloride where the 1-hydroxyl group is already methylated. The reaction is complete after about 1 hour to about 24 hours. Generally the reaction mixture is stirred for about 16 hours (overnight). Concentration of the reaction mixture followed by purification using standard chromatographic techniques affords the desired triacetylated pyranoside derivative.

10 The introduction of fluorine to said pyranoside derivative is accomplished as described hereinbelow. The pyranoside is dissolved in a reaction inert solvent such as chloroform, methylene chloride or ethylene dichloride. Particularly preferred is methylene chloride. The reaction mixture is cooled to about -78°C to about 0°C. A particularly preferred temperature is -40°C. The reaction mixture is treated with 4-DMAP
15 followed by dimethylaminosulfurtrifluoride (DAST). A person of ordinary skill in the art would recognize that about one equivalent each of 4-DMAP and DAST would generally be sufficient for the reaction to proceed to completion. However, depending upon the particular reaction conditions or the nature of the substrate, it may be required to add additional equivalents of either or both of these reagents. The reaction mixture is
20 stirred at room temperature for a period of about 1 hour to about 24 hours. Conveniently the reaction mixture is stirred overnight at room temperature. The reaction mixture is quenched with a proton source such as a (C₁-C₄)alkanol with methanol being preferred, and the product is isolated in a way familiar to one of ordinary skill in the art. The S_N² nature of the substitution reaction converts a glucopyranoside starting material
25 into a galactopyranoside and a galactopyranoside starting material into a glucopyranoside. The alkyl pyranoside thus obtained may be directly halogenated at the 1-position or, in order to provide a better leaving group, the methoxy group may be converted into an acetoxyl group prior to bromination as illustrated in the Preparation Section for methyl 2,3,6-tri-O-acetyl-4-deoxy-4-fluoro- α -D-galactopyranoside.

30 To prepare the precursors of the compounds of formulae (VIII) and (IX) wherein R⁴ is -COOH or a derivative of -COOH such as -CO₂R⁸, -CONH₂, -CONHR⁸ or -CONR₂⁸, the corresponding uronic acid of formula (X) or (XI)

-16-



10

is converted to the above-mentioned derivatives utilizing methods well known to one of ordinary skill in the art. For example, to prepare compounds wherein R^4 is $-\text{CO}_2R^8$, a compound of formulae (X) or (XI) is reacted under the standard esterification procedures of organic chemistry with a suitable $R^8\text{-OH}$ derivative. To prepare compounds wherein R^8 is $-\text{CONH}_2$, $-\text{CONHR}^8$ or $-\text{CONR}_2^8$, a compound of formulae (X) or (XI) is reacted under the standard amidation conditions of organic chemistry with a suitable amine. The sugar derivatives so prepared can be fluorinated employing the methods described above, and coupled to the macrolides of formulae (IV) and (V) as described hereinabove. To obtain the compound wherein R^4 is $-\text{COOH}$, an appropriate compound wherein R^4 is $-\text{CO}_2R^8$ can be selectively deesterified, after glycosylation, utilizing the methods of deesterification well known to one of ordinary skill in the art.

When the compounds of formula (I) of the present invention are acidic, as when R^4 is $-\text{CO}_2\text{H}$, the invention also embraces pharmaceutically acceptable salts of said compounds of formula (I).

Typical pharmaceutically acceptable cationic salts for such use include alkali metal salts (e.g., sodium and potassium), alkaline earth metal salts (e.g., magnesium and calcium), aluminum salts, ammonium salts and salts with organic amines such as benzathine (N,N'-dibenzylethylenediamine), choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine), benethamine (N-benzylphenethylamine), diethylamine, piperazine, tromethamine (2-amino-2-hydroxymethyl-1,3-propanediol) and procaine. An especially preferred such salt is the sodium salt.

The pharmaceutically acceptable cationic salts of the present invention are readily prepared by reacting the acid forms with an appropriate base, usually one equivalent, in a co-solvent. Typical bases are sodium hydroxide, sodium methoxide, magnesium hydroxide, calcium hydroxide, benzathine, choline, diethanolamine, 5 piperazine and tromethamine. The salt is isolated by concentration to dryness or by addition of a non-solvent. In many cases, salts are preferably prepared by mixing a solution of the acid with a solution of a different salt of the cation (sodium or potassium ethylhexanoate, magnesium oleate), employing a solvent (e.g., ethyl acetate) from which the desired cationic salt precipitates, or can be otherwise isolated by 10 concentration and/or addition of a non-solvent.

With respect to the macrolides of formula (I) of this invention, it is to be understood that there are conformer(s) or stereoisomeric forms such as optical and geometrical isomers due to asymmetric carbon atom(s) and double bond(s), and such isomers are also included within the scope of this invention.

15 The compounds of formula (I) thus prepared are useful in the treatment of resistance to transplantation and autoimmune diseases such as rheumatoid arthritis or psoriasis. In the treatment of resistance to transplantation, a compound of formula (I) may be used either prophylactically or in response to an adverse reaction by the human subject to a transplanted organ or tissue. When used prophylactically, a 20 compound of formula (I) is administered to the patient or to the tissue or organ to be transplanted in advance of the transplantation operation. Prophylactic treatment may also include administration of the medication after the transplantation operation but before any signs of adverse reaction to transplantation are observed. When administered in response to an adverse reaction, a compound of formula (I) is admin- 25 istered directly to the patient in order to treat said resistance to transplantation after outward signs of the resistance have been manifested.

For use in the treatment of resistance to transplantation and autoimmune diseases such as rheumatoid arthritis or psoriasis in a mammal, including man, a compound of formula (I) is formulated into a suitable pharmaceutical composition 30 containing a disease treating effective amount. Depending upon the potency of the particular compound of formula (I) being administered, about 0.05 mg/kg of body weight per day to about 30 mg/kg of body weight per day, in single or multiple daily doses, is administered to the mammal being treated. A more preferred range is

0.1 mg/kg of body weight per day to about 20 mg/kg of body weight per day, although in particular cases, at the discretion of the attending physician, doses outside the broader range may be required. The preferred route of administration is generally oral, but parenteral administration (e.g., intravenous, intramuscular, subcutaneous or intramedullary) will be preferred in special cases such as where oral administration is inappropriate to the instant target or where the patient is unable for various reasons to ingest the drug. Topical administration may also be indicated, as where the patient is suffering from a skin disease such as psoriasis or whenever the medication is best applied to the surface of a tissue or organ as determined by the attending physician.

The compounds of formula (I) thus prepared are also useful in the treatment of infections caused by fungi. For use in the treatment of said fungal infections in a mammal, including man, a compound of formula (I) is formulated into a pharmaceutical composition containing a disease treating effective amount. Depending upon the potency of the particular compound of formula (I) being administered, about 0.05 mg/kg of body weight per day to about 30 mg/kg of body weight per day, in single or multiple daily doses, is administered to the mammal being treated. A more preferred range is 0.1 mg/kg of body weight per day to about 20 mg/kg of body weight per day, although in particular cases, at the discretion of the attending physician, doses outside the broader range may be required. The preferred route of administration is generally oral, but parenteral administration (e.g., intravenous, intramuscular, subcutaneous or intramedullary) will be preferred in special cases such as where oral administration is inappropriate to the instant target or where the patient is unable for various reasons to ingest the drug. Topical administration may also be indicated whenever the medication is best applied to the surface of a tissue or organ as determined by the attending physician.

The compounds of the present invention are generally administered in the form of a pharmaceutical composition comprising at least one of the compounds of formula (I) together with a pharmaceutically acceptable vehicle or diluent. Such compositions are generally formulated in a conventional manner utilizing solid or liquid vehicles or diluents as appropriate to the mode of administration: for oral administration, in the form of tablets, hard or soft gelatin capsules, suspensions, granules, powders and the like; for parenteral administration, in the form of injectable

solutions or suspensions and the like; and for topical administration, in the form of solutions, lotions, ointments, salves and the like.

The utility of the compounds of the present invention as medical agents in the treatment of resistance to transplantation and autoimmune diseases such as
5 rheumatoid arthritis or psoriasis is demonstrated by the activity of said compounds in the biological screen described hereinbelow. Said biological screen also provides a means whereby the activities of the compounds of formula (I) can be compared with the activities of other known compounds. The results of these comparisons are useful for determining dosage levels in mammals, including man, for the treatment of
10 resistance to transplantation and autoimmune diseases such as rheumatoid arthritis and psoriasis.

The human mixed lymphocyte reaction (MLR) is used to generate an immune response *in vitro* which is measured *via* ^3H -thymidine uptake. This screen uses peripheral blood mononuclear cells in a modified two-way MLR. To ensure disparity of
15 HLA type D antigens and therefore maximize stimulation, a pool of frozen donor cells is used as the stimulator population; freshly isolated cells are used as the responder population.

Freshly drawn mononuclear cells are suspended in RPMI-1640 enriched with: 0.5% MEM non-essential amino acids (100x) solution, 1% L-glutamine (200 mM), 1%
20 MEM vitamins (100x), 1% penicillin streptomycin solution (10,000 units/mL) and 15% heat-inactivated human AB serum (NABI). The cells are counted and the concentration is adjusted to 5×10^5 cells/mL. The solution is then transferred to round bottom 96 well plates in 100 μL /well quantities. These plates now contain the responder cells.

The stimulator cells are prepared by pooling the mononuclear cells collected
25 from several different individuals. The cells are suspended in 90% human AB serum and 10% DMSO such that the cell count is 2×10^7 cells/mL. The cells are stored in liquid nitrogen. For an MLR, the viable cells are diluted to 5×10^5 cells/mL, and 100 μL /well is added to the plates containing the responder cells. To each well, containing a mixture of responder cells and stimulator cells, is added 50 μL of compound
30 solution. Triplicate wells are run for each dose. The plates are incubated at 37°C under an atmosphere of 5% CO_2 and are humidified for five days. To each well is added 1 μCi of ^3H -thymidine and incubation is continued for another eighteen hours. The cells are harvested using the LKB Beta Plate system.

The percent inhibition of stimulated control is obtained using the following equation:

$$5 \quad \% \text{ Inhibition} = \left[100 - \left(\frac{\text{avg. cpm of drug}}{\text{avg. cpm of stimulated control}} \right) \right] \times 100$$

The abbreviation cpm is defined as counts per minute. RPMI-1640 is a tissue culture medium which is available from Sigma.

10 Activity in the MLR screen recited above is indicative of usefulness of the active compound in the treatment of resistance to transplantation and autoimmune diseases such as rheumatoid arthritis and psoriasis.

Antimicrobial activities of the macrolides of the present invention against various fungi are determined by a serial agar dilution method in a Sabouraud agar. Minimum inhibitory concentrations (MIC) are obtained after incubation for 24 hours at 30°C.

The present invention is illustrated by the following examples. However, it should be understood that the invention is not limited to the specific details of these examples. All reactions are conducted under an inert atmosphere, such as nitrogen, unless otherwise specified. The abbreviations THF, DMSO, DAST, DMAP and Ac, where used, refer to tetrahydrofuran, dimethyl sulfoxide, dimethylamino sulfurtrifluoride, 4-dimethylaminopyridine and acetyl, respectively. The sugar halides were purchased from a reliable vendor such as Sigma or Aldrich, as were the sugars, unless specifically mentioned. Anhydrous solvents were used, anhydrous being defined as substantially free from water.

25 The expression "reaction inert solvent," where used hereinabove, refers to any solvent which does not interact with starting materials, reagents, intermediates or products in a manner which adversely affects the yield of the desired product.

Terms or acronyms which appear in Preparations 1 and 2 are described in further detail hereinbelow.

30 PYEA agar is prepared by dissolving Difco maltose (10 g), Difco yeast extract (4 g), dextrose (4 g), Difco agar (15 g) and fresh coconut milk (50 mL) in enough deionized water to yield a one liter solution; and the solution is adjusted to pH 7.3 with 1N NaOH.

-21-

ATCC 172 medium is prepared by dissolving glucose (10 g), soluble starch (20 g), yeast extract (5 g), NZ-amine A (Difco, 5 g) and calcium carbonate (1 g) in enough deionized water to yield a one liter solution; and the solution is adjusted to pH 7.0 with 1N KOH.

- 5 JDYTT medium is prepared by dissolving cerelese (10 g), corn starch (5 g), corn steep liquor (5 g), NZ-amine YTT (5 g), cobalt chloride (0.002 g) and calcium carbonate (3 g) in enough deionized water to yield a one liter solution; and the solution is adjusted to pH 7.2 with 1N NaOH.

- 10 NZ-amine A and NA-amine YTT are available from Difco, as are most of the ingredients of the above media.

In the MLR protocol provided hereinabove, RPMI-1640 is a standard medium for MLR studies; MEM is defined as "minimum essential media"; and NABI is a supplier.

-22-

EXAMPLE 1

17-Ethyl-1,14-dihydroxy-12-[2'-(4''-(2''',3''',6'''-tri-O-acetyl-4'''-deoxy-4'''-fluoro- α -D-galactopyranosyloxy)-3''-methoxycyclohexyl)-1'-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatricyclo[22.3.1.O^{4,9}]-octacos-18-ene-2,3,10,16-tetraone

To a stirred slurry of FK520 (7.1 g, 8.9 mmol), the title compound of Preparation Four (3.3 g, 8.9 mmol) and 4Å Molecular Sieves (crushed, 3.3 g) in methylene chloride (90 mL) at -78°C was added silver carbonate (4.9 g, 17.8 mmol) followed by silver triflate (0.46 g, 1.8 mmol). The reaction mixture was allowed to warm to room temperature over 10 hours and was then stirred for an additional 8 hours. The resultant tan slurry was filtered through Celite and the filtrate was evaporated *in vacuo*. The residue was purified on silica gel, eluting with ethyl acetate/hexane (1/1), to afford the product as a single anomer (2.94 g, 30%). FAB Ms (M⁺ + Na⁺) 1104.

EXAMPLES 2-4

Using substantially the same procedure as recited in Example 1, but substituting one molar equivalent of the appropriate sugar halide for the title compound of Preparation 4, the following compounds were prepared.

2. 17-Ethyl-1,14-dihydroxy-12-[2'-(4''-(2''',3''',4'''-tri-O-acetyl-6'''-deoxy-6'''-fluoro- α -D-galactopyranosyloxy)-3''-methoxycyclohexyl)-1'-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatricyclo[22.3.1.O^{4,9}]-octacos-18-ene-2,3,10,16-tetraone

Mass spectrum (FAB): 1104 (molecular ion + Na⁺).

3. 17-Ethyl-1,14-dihydroxy-12-[2'-(4''-(2''',4''',6'''-tri-O-acetyl-3'''-deoxy-3'''-fluoro- α -D-galactopyranosyloxy)-3''-methoxycyclohexyl)-1'-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatricyclo[22.3.1.O^{4,9}]-octacos-18-ene-2,3,10,16-tetraone

Mass spectrum (FAB): 1104 (molecular ion + Na⁺).

4. 17-Ethyl-1,14-dihydroxy-12-[2'-(4''-(3''',5'''-di-O-benzoyl-2'''-deoxy-2'''-fluoro- α -D-arabinofuranosyl-oxy)-3''-methoxycyclohexyl)-1'-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatri-cyclo[22.3.1.O^{4,9}]-octacos-18-ene-2,3,10,16-tetraone

Mass spectrum (FAB): 1156 (molecular ion + Na⁺).

EXAMPLE 5

17-Ethyl-1,14-dihydroxy-12-[2'-(4''-(4'''-deoxy-4'''-fluoro-galactopyranosyloxy)-3''-methoxycyclohexyl)-1'-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatricyclo[22.3.1.O^{4,9}]-octacos-18-ene-2,3,10,16-tetraone

5

The title compound of Example 1 (1.1 g) is dissolved in methanol (10 mL) and treated with sodium methoxide (10 mg). The reaction mixture is stirred for 48 hours at 0°C, then is treated with one drop of acetic acid from a disposable pipet. The solvent is removed *in vacuo* and the residue is purified on silica gel eluting with methylene chloride/methanol (15/1) to afford the title compound of this Example.

10

EXAMPLES 6 AND 7

Using substantially the same procedure as recited in Example 5, but substituting the appropriate title compound of either Example 2 or Example 3 for the title compound of Example 1, the following compounds can be prepared.

15

6. 17-Ethyl-1,14-dihydroxy-12-[2'-(4''-(6'''-deoxy-6'''-fluoro- α -D-galactopyranosyloxy)-3''-methoxycyclohexyl)-1'-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatricyclo[22.3.1.O^{4,9}]-octacos-18-ene-2,3,10,16-tetraone

20

7. 17-Ethyl-1,14-dihydroxy-12-[2'-(4''-(3'''-deoxy-3'''-fluoro- α -D-galactopyranosyloxy)-3''-methoxycyclohexyl)-1'-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatricyclo[22.3.1.O^{4,9}]-octacos-18-ene-2,3,10,16-tetraone

EXAMPLE 8

25 17-Ethyl-1,14-dihydroxy-12-[2'-(4''-(3'''',4'''',6'''-tri-O-acetyl-2'''-deoxy-2'''-fluoro- α -D-galactopyranosyloxy)-3''-methoxycyclohexyl)-1'-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatricyclo[22.3.1.O^{4,9}]-octacos-18-ene-2,3,10,16-tetraone

30

Using substantially the same procedure as recited in Example 1, but substituting the title compound of Preparation 11 for the title compound of Preparation 4, the title compound of the Example was prepared.

Mass spec (LSIMS;FAB) - M⁺_{N₂} = 1104.5, base = 291.1

PREPARATION 1

17-Ethyl-1,14-dihydroxy-12-[2'-(4"-hydroxy-3"-methoxy-cyclohexyl)-1'-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatricyclo[22.3.1.O^{4,9}]-octacos-18-ene-2,3,10,16-tetraone

5

Streptomyces hygroscopicus subsp. *ascomyceticus* culture ATCC 14891 was carried on PYEA agar slants (10 g/L of Difco maltose, 4 g/L of Difco yeast extract, 4 g/L of dextrose, 15 g/L of Difco agar and 50 mL of fresh coconut milk which was diluted up to one liter with deionized water, then adjusted to pH 7.3 with 1N NaOH). The preparation was incubated for 10 to 12 days at 28°C, and the spores were then transferred to sterile 1x6 shake tubes containing 10 mL of ATCC 172 medium (10 g/L of glucose, 20 g/L of soluble starch, 5 g/L of yeast extract, 5 g/L of NZ-amine A and 1 g/L of calcium carbonate. The pH was adjusted to 7.0 with 1N KOH). The tubes were incubated at 28°C and shaken on a rotary shaker at about 150 to 200 cycles/minute. After 4 to 5 days the broth was diluted to 40% with glycerol and ATCC 172 medium and then transferred aseptically to cryotubes. Each tube was charged with 1/2 mL of broth. The tubes were frozen at -80°C during storage.

The tubes from the ultra cold stock were used as seed inoculum for the preparation of inoculum flasks, one tube per 50 mL of sterile JDYTT medium in 300 mL shake flasks. The composition of the JDYTT medium was 10 g/L of cerelose, 5 g/L of NZ-amine YTT, 0.002 g/L of cobalt chloride and 3 g/L of calcium carbonate. The pH of the JDYTT medium was adjusted to pH 7.2 with 1N NaOH. The shake flasks were shaken and incubated on a rotary shaker at about 150-200 cycles/minute and 28°C.

Two mL of an about 3 to 5 day-old shake flask inoculum was used to inoculate the second stage flask inoculum containing 80 mL of JDYTT medium in a 3L jar fermenter. The fermenter medium was 45 g/L of corn starch, 10 g/L of corn steep liquor, 10 g/L of amber or Baker dried yeast, 3 g/L of calcium carbonate and 0.005 g/L of cobalt chloride. The pH was adjusted to about 6.4 to 6.8 with 1N NaOH. One mL of antifoam P-2000 was added to the jar fermenters together with 100 mL of soya bean oil. The pH was adjusted to about 6.4 to 6.8 with 1N NaOH and the material was agitated at 1700 rpm. The temperature was maintained at 28°C and sterile air was sparged through the medium at the rate of one volume per volume per minute.

After inoculation, sterile soya bean oil was used to control foaming. In longer fermentations, and, depending on media used, the sugar content can be monitored and sugar feeds used at 40, 60 and 90 hours to maintain the reducing sugar level at or above 0.05%. The fermentation was run for 46 hours.

5 Using standard methods of thin-layer chromatography and HPLC, the fermentation broth was monitored and relative potency was calculated.

The product was found primarily in the mycelium, but workup of the whole broth is preferred. Thus, after the fermentation has run its course, the whole broth was extracted twice with one-third to one-half of its volume of methylisobutylketone (MIBK).
10 The layers were separated by means of a DeLaval separator or a Podbielnack extractor. The solvent layer was clarified and concentrated first in a vacuum pan and then in a rotary evaporator. The concentrate was subjected to four tube counter current distribution in 20 liter carbuoys using 10 liter top layer and 1 liter bottom layer per carbuoy of a heptane/acetonitrile 10/1 system. The active bottom layers were collected,
15 combined and concentrated. The material was further purified *via* filtration through Florisil (washing with hexane, hexane/methylene chloride and methylene chloride, successively, with a gradual increase in methylene chloride). Most of the activity was found in the methylene chloride fractions. These were combined and concentrated. A second filtration step was performed, this time through silica gel (washing with
20 heptane, methylene chloride, methylene chloride/ethyl acetate and ethyl acetate). The activity was mostly found in the fractions containing a methylene chloride/ethyl acetate mixture and the fractions containing only ethyl acetate. These were combined and concentrated, redissolved in methylene chloride and treated with DARCO G60. The sample was then divided into 12 to 15 g portions and each sample was further
25 chromatographed on a Prep 500 liquid chromatograph using silica gel columns and eluting using a linear gradient beginning with 100% methylene chloride and ending with 100% ethyl acetate. The active cuts were combined, concentrated and chromatographed on a Prep 500, using reversed phase (^{18}C) silica gel and eluting with a linear gradient beginning with acetone and ending with 100% water. Clean product
30 was obtained as the last component isolated off the column.

The active fractions in the foregoing fermentation procedure were determined using the following bioassay.

-26-

A 12.5 mm disc was applied directly to the agar surface. *Candida albicans* ATCC 14053, *Saccharomyces pastorianus* FD3737 and a sensitive strain of *Byssoschlamys fulva* FM 10,300(S) and FM 10,464(R) were used. The *Candida* and *Saccharomyces* plates were incubated at 37°C for 18 hours, then the plates were
5 examined for activity. The *Byssoschlamys* plates were incubated at 28°C and read after 18 hours. Plates containing only FK506 and FK520 (CP-105051) were active against the *Byssoschlamys* strain. Impure fractions (containing nigericin) were active against the other strains as well.

An HPLC method for determining the purity of the fractions was also used. The
10 method entailed using a Dupont Zorbax CN column (4.6 mm x 25 cm) and an isocratic system composed of 55/45 water/acetonitrile and a flow rate of one mL/min. Detection was accomplished at 214 nm. The broth sample (20 mL) was mixed with MIBK (20 mL) and shaken for about 4 to 5 minutes. The layers were separated and the solvent was concentrated to near dryness. The residue was taken up in 1 mL of neat acetonitrile
15 and a 5 μ L sample was injected into the HPLC. The retention time for FK520 is approximately 12.7 minutes under these conditions.

PREPARATION 2

20 17-Allyl-1,14-dihydroxy-12-[2'-(4"-hydroxy-3"-methoxy-cyclohexyl)-1'-methylvinyl]-23,25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxa-4-azatricyclo[22.3.1.O^{4,9}]-octacos-18-ene-2,3,10,16-tetraone

Streptomyces tsukubaensis No. 9993 FERM BP-927 was carried on PYEA agar slants (10 g/L of Difco maltose, 4 g/L of Difco yeast extract, 4 g/L of dextrose, 15 g/L
25 of Difco agar and 50 mL of fresh coconut milk which was diluted up to one liter with deionized water, then adjusted to pH 7.3 with 1N NaOH). The preparation was incubated for 10 to 12 days at 28°C, and the spores were then transferred to sterile 1x6 shake tubes containing 10 mL of ATCC 172 medium (10 g/L of glucose, 20 g/L of soluble starch, 5 g/L of yeast extract, 5 g/L of NZ-amine A and 1 g/L of calcium carbonate. The pH was adjusted to 7.0 with 1N KOH). The tubes were incubated at
30 28°C and shaken on a rotary shaker at about 150 to 200 cycles/minute. After 4 to 5 days the broth was diluted to 40% with glycerol and ATCC 172 medium and then transferred aseptically to cryotubes. Each tube was charged with 1/2 mL of broth. The tubes were frozen at -80°C during storage.

-27-

The tubes from the ultra cold stock were used as seed inoculum for the preparation of inoculum flasks, one tube per 50 mL of sterile JDYTT medium in 300 mL shake flasks. The composition of the JDYTT medium was 10 g/L of cerelose, 5 g/L of NZ-amine YTT, 0.002 g/L of cobalt chloride and 3 g/L of calcium carbonate.

- 5 The pH of the JDYTT medium was adjusted to pH 7.2 with 1N NaOH. The shake flasks were shaken and incubated on a rotary shaker at about 150-200 cycles/minute and 28°C.

- Two mL of an about 3 to 5 day-old shake flask inoculum was used to inoculate the second stage flask inoculum containing 80 mL of JDYTT medium in a
10 3L jar fermenter. The fermenter medium was 45 g/L of corn starch, 10 g/L of corn steep liquor, 10 g/L of amber or Baker dried yeast, 3 g/L of calcium carbonate and 0.005 g/L of cobalt chloride. The pH was adjusted to about 6.4 to 6.8 with 1N NaOH. One mL of antifoam P-2000 was added to the jar fermenters together with 100 mL of soya bean oil. The pH was adjusted to about 6.4 to 6.8 with 1N NaOH and the material
15 was agitated at 1700 rpm. The temperature was maintained at 28°C and sterile air was sparged through the medium at the rate of one volume per volume per minute.

- After inoculation, sterile soya bean oil was used to control foaming. In longer fermentations, and, depending on media used, the sugar content can be monitored and sugar feeds used at 40, 60 and 90 hours to maintain the reducing sugar level at or
20 above 0.05%. The fermentation was run for 46 hours.

Using standard methods of thin-layer chromatography and HPLC, the fermentation broth was monitored and relative potency was calculated.

- The fermenters were stopped and extracted twice with ½ its volume of methylisobutylketone (MIBK). The solvent layer was separated by aspiration and
25 concentration *in vacuo* to a viscous oil. The oil was triturated with hexane, diethyl ether and methylene chloride and the active cuts (the diethyl ether cuts) were chromatographed on florisil. The florisil was eluted with, successively, diethyl ether methylene chloride, ethyl acetate and acetone. The eluate was concentrated and treated with activated charcoal. The concentrate was filtered and dissolved in ethyl
30 acetate. Hexane was added to crystallize the product.

The bioactivity of the broth and subsequent recovery streams was followed by using a strain of *Byssochlamys fulva*. The components in the broth and recovery streams were visualized by chromatography on Analtech silica gel GF (Trademark)

-28-

plates using neat ethyl acetate as the eluant. The developed plates were sprayed with vanillin reagent (3 g of vanillin in 75 mL of ethanol and 25 mL of 85% phosphoric acid) and heated to 80°C. The product appeared as a violet spot.

PREPARATION 3

5 Methyl-2,3,6-tri-O-acetyl- α -D-glucopyranoside

Methyl- α -D-glucopyranoside (Aldrich, 15.0 g, 77 mmol) in toluene (300 mL) was treated with bis(tributyltin)oxide (Aldrich, 78 mL, 154 mmol) and the resultant mixture was refluxed under a Dean-Stark trap for three hours. The mixture was cooled to 0°C and then acetyl chloride (17 mL, 231 mmol, 3 equivalents) was added. The mixture
10 was stirred overnight, then concentrated *in vacuo* to yield a syrup. Purification on silica gel [eluting with ethyl acetate/hexane (3/2)] afforded an oil (10.4 g, 42%).

PREPARATION 4

Methyl-2,3,6-tri-O-acetyl-4-deoxy- 4-fluoro- α -D-galactopyranoside

15 The product of Preparation 3 (10.4 g, 32 mmol) was mixed with 4-dimethylaminopyridine (8.2 g, 67 mmol) in methylene chloride (100 mL) and cooled to -40°C. The reaction mixture was treated dropwise with DAST (8.6 mL, 65 mmol) and then allowed to warm slowly to room temperature. After stirring at room temperature overnight (16 hours), the reaction mixture was cooled to -40°C and quenched with
20 methanol (20 mL). The quenched mixture was diluted with ethyl acetate (800 mL) and washed with 1N HCl (2 x 80 mL), water (1 x 80 mL), saturated aqueous sodium bicarbonate (1 x 80 mL) and brine (1 x 80 mL). The solvent was dried with MgSO₄ and the solvent was removed to yield an oily residue which was purified on silica gel [eluting with ethyl acetate/hexane (2/3)] to afford 3.7 g (36%) of a solid, m.p. 87-89°C.

25 PREPARATION 5

1,2,3,6-Tetra-O-acetyl-4- deoxy-4-fluoro- α -D-galactose

The title compound of Preparation 4 (100 mg, 0.31 mmol) was mixed with acetic acid (1 mL), acetic anhydride (1 mL) and sulfuric acid (5 drops) and stirred for 18 hours.
30 The reaction mixture was diluted with ethyl acetate (200 mL) and washed with saturated aqueous sodium bicarbonate (2 x 20 mL), water (1 x 20 mL) and brine (1 x 20 mL) and then treated with MgSO₄. The solvent was removed *in vacuo* to yield an oily residue which was purified on silica gel [eluting with ethyl acetate/hexane(2/3)] to afford 90 mg (82%) of an oil.

PREPARATION 62,3,6-Tri-O-acetyl-4-deoxy-4-fluoro- α -D-galactosylbromide

The title compound of Preparation 5 (3.4 g, 9.7 mmol) was mixed with a 4.1M solution of hydrobromic acid in acetic acid (50 mL) and stirred at room temperature for five hours. The mixture was concentrated *in vacuo* to give 5 g of an oil which was purified on silica gel [eluting with ethyl acetate/hexane (2/3)] to afford 3.3 g (92%) of an oil.

Analysis calculated for $C_{12}H_{16}O_7BrF$:

10 C, 38.83; H, 4.31.

Found: C, 38.85, H, 4.48.

PREPARATION 72,4,6-Tri-O-acetyl-3-deoxy-3-fluoro- α -D-galactopyranosylbromide

15 Prepared as described by Kovac, P.; Yeh, H.J.C. and Glaudemans, C.P.J., Carbohydrate Research, 140, 277 (1985).

PREPARATION 83,5-Di-O-benzoyl-2-deoxy-2-fluoro- α -D-arabinofuranosyl Bromide

20 1,3,5-Tri-O-benzoyl-2-deoxy-2-fluoro- α -D-arabinofuranose (Pfanstiehl, 1000 mg, 2.2 mmol) was dissolved/suspended in acetic acid (10 mL) and treated with a solution of HBr/acetic acid (10 mL, 30%) at 0°C for 2 hours. The solvent was removed *in vacuo* and residual acid was removed by azeotroping with toluene. The bromide was used without further purification. 1H NMR (partial): δ 4.6 (3H), 5.18 (m, 1H), 5.34 (d, 1H) and
25 6.45 (d, 1H).

-30-

PREPARATION 92,3,4-Tri-O-acetyl-6-deoxy-6-fluoro- α -D-galactosylbromide

A. 1,2,3,4-Tetraacetyl-6-deoxy-6-fluoro-galactopyranoside was prepared as
5 described by Sharma, M. and Korytnyk, W. (Tetrahedron Letters, 1977, 573).

B. 1,2,3,4-Tetraacetyl-6-deoxy-6-fluoro-galactopyranoside (100 mg) was
dissolved in a 30% solution of HBr in acetic acid (10 mL) and was stirred at room
temperature for two hours. The solvent was removed *in vacuo* and the residual acid
was removed *via* repeated azeotroping with toluene. The residue was
10 chromatographed on silica gel (eluted with ethyl acetate:hexane::30:70) to afford a
white solid; m.p. 131-133°C. Mass spectrum: $m/z = 371$ (M^+), 373.

PREPARATION 103,4,6-Tri-O-acetyl-2-deoxy-2-fluoro-galactopyranosyl fluoride

15 The title compound of this preparation was prepared as described by Korytnyk,
W. et al., Tetrahedron, 38 (16), 2547 (1982).

PREPARATION 113,4,6-Tri-O-acetyl-2-deoxy-2-fluoro-galactopyranosyl bromide

20 The title compound of this preparation was prepared substantially as recited in
Preparation 6, but substituting the title compound of Preparation 10 for the title
compound of Preparation 5. (Mass spectrum - $M^+ = 370.1$).

-31-

CLAIMS


1. A compound of the formula

5

10

15

20



or a pharmaceutically acceptable salt thereof;

wherein n is 1 or 2;

the dotted line represents an optional double bond in the case where R² is H;

- 25 A and B are taken separately and A is H and B is H or -OH, or A and B are taken together and form =O;


R² is H, (C₂-C₅)alkanoyloxy or -OR⁰;

R³ is (C₁ to C₃)alkyl or allyl;

R⁰ and R¹ are each H,

30

-32-



10

R^4 is, for each occurrence, independently $-\text{CO}_2R^8$, $-\text{CO}_2\text{H}$,
 $-\text{CH}_2\text{OH}$, H , $-\text{CH}_3$, $-\text{CH}_2\text{F}$, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{CONH}_2$, $-\text{CONHR}^8$, $-\text{CON}(\text{R}^8)_2$,
 $-\text{CH}_2\text{OCOR}^8$, $-\text{CH}_2\text{OCO}_2\text{R}^8$, $-\text{CH}_2\text{OCONR}_2^8$ or $-\text{CH}_2\text{OR}^8$;

R^5 is, for each occurrence, independently $(\text{C}_1 \text{ to } \text{C}_4)\text{alkoxy}$, benzyloxy, $-\text{OH}$, $-\text{OCOR}^8$,
 15 OCOCH_2R^8 , $-\text{OCO}_2\text{R}^8$ or $-\text{OSi}(\text{R}^8)_3$;

t is 1, 2 or 3;

m is 0 or 1; and

R^8 is $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_3\text{-C}_6)\text{cycloalkyl}$, allyl, pyridyl, thienyl, benzyl, benzyl variously
 20 substituted with one to five halogen atoms, $-\text{OH}$ groups or $(\text{C}_1\text{-C}_4)\text{alkoxy}$ groups, phenyl
 or phenyl variously substituted with one to five halogen atoms, $-\text{OH}$ groups or $(\text{C}_1\text{-C}_4)\text{alkoxy}$ groups;

provided that R^1 and R^0 are not both H ; when m is 0, R^4 is

$-\text{CH}_2\text{F}$, $-\text{CHF}_2$ or $-\text{CF}_3$; and in formulae (II) and (III), each ring carbon atom must bear
 at least one hydrogen atom.


25 2. A compound according to claim 1 wherein n is 2; A and B are taken
 together and form $=\text{O}$; the dotted line represents no bond.

3. A compound according to claim 2 wherein R^3 is methyl, ethyl or allyl.

4. A compound according to claim 3 wherein R^3 is ethyl and R^2 is $-\text{OH}$.


-33-

5. A compound according to claim 4 wherein R^1 is



R^4 is H, $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{F}$, $-\text{CH}_2\text{OCOCH}_3$ or $-\text{CH}_2\text{OCH}_2\text{C}_6\text{H}_5$; and
 R^5 is $-\text{OH}$, $-\text{OCOCH}_2\text{C}_6\text{H}_5$ or $-\text{OCOCH}_3$.

6. A compound according to claim 5 wherein R^1 is




R^4 is H, $-\text{CH}_2\text{OH}$ or $-\text{CH}_2\text{OCOCH}_3$; and R^5 is $-\text{OH}$ or $-\text{OCOCH}_3$.

7. The compound according to claim 6 wherein R^4 is
 $-\text{CH}_2\text{OH}$ and R^5 is $-\text{OH}$.

- 30 8. The compound according to claim 6 wherein R^4 is
 $-\text{CH}_2\text{OCOCH}_3$ and R^5 is $-\text{OCOCH}_3$.

9. A compound according to claim 5 wherein R^1 is

-34-



5


R^4 is H, $-\text{CH}_2\text{OH}$ or $-\text{CH}_2\text{OCOCH}_3$; and R^5 is $-\text{OH}$ or $-\text{OCOCH}_3$.

10 10. The compound according to claim 9 wherein R^4 is $-\text{CH}_2\text{OCOCH}_3$ and R^5 is $-\text{OCOCH}_3$.

11. The compound according to claim 9 wherein R^4 is $-\text{CH}_2\text{OH}$ and R^5 is $-\text{OH}$.

12. A compound according to claim 5 wherein R^1 is

15



20


R^4 is $-\text{CH}_2\text{F}$ and R^5 is $-\text{OH}$ or $-\text{OCOCH}_3$.

13. The compound according to claim 12 wherein R^5 is $-\text{OCOCH}_3$.

14. The compound according to claim 12 wherein R^5 is $-\text{OH}$.

25

15. A compound according to claim 5 wherein R^1 is



30

R^4 is H, $-\text{CH}_2\text{OH}$ or $-\text{CH}_2\text{OCOR}^8$; R^5 is $-\text{OH}$ or $-\text{OCOR}^8$; and R^8 is benzyl.


16. The compound according to claim 15 wherein R^4 is

-35-

$-\text{CH}_2\text{OCOR}^8$ and R^5 is $-\text{OCOR}^8$.

17. A compound according to claim 1 wherein n is 2; A and B taken separately and are each H; the dotted line represents no bond; R^2 is OH; and R^3 is ethyl.

5 18. A compound according to claim 17 wherein R^1 is



19. A compound according to claim 18 wherein R^4 is $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OCOCH}_3$, $-\text{CH}_2\text{OCOCH}_2\text{C}_6\text{H}_5$ or $-\text{CH}_2\text{F}$; and R^5 is $-\text{OH}$, $-\text{OCOCH}_3$ or $-\text{OCOCH}_2\text{C}_6\text{H}_5$.

20. A method for treating resistance to transplantation in a mammal in need of such treatment comprising administering to said mammal a resistance to transplantation treating effective amount of a compound according to claim 1 or a pharmaceutically-acceptable salt thereof.

21. A method for treating autoimmune disease in a mammal in need of such treatment comprising administering to said mammal an autoimmune disease treating effective amount of a compound according to claim 1 or a pharmaceutically-acceptable salt thereof.

22. A method for treating fungal diseases in a mammal in need of such treatment comprising administering to said mammal a fungal disease treating effective

amount of a compound according to claim 1 or a pharmaceutically-acceptable salt thereof.


23. A pharmaceutical composition comprising a resistance to transplantation treating effective amount of a compound of claim 1 and a pharmaceutically-acceptable carrier.

24. A pharmaceutical composition comprising an autoimmune disease treating effective amount of a compound of claim 1 and a pharmaceutically-acceptable carrier.

25. A pharmaceutical composition comprising a fungal disease treating effective amount of a compound of claim 1 and a pharmaceutically-acceptable carrier.

26. The compound according to claim 1 wherein said compound is 17-ethyl-1,14-dihydroxy-12-[2'-(4''-(3''',4''',6'''-tri-O-acetyl-2'''-deoxy-2'''-fluoro- α -D-galactopyranosyloxy)-3''-methoxycyclohexyl)-1'-methylvinyl]-23-25-dimethoxy-13,19,21,27-tetramethyl-11,28-dioxo-4-azatricyclo[22.3.1.0.^{4,9}]-octacos-18-ene-2,3,10,16-tetraone.

27. A process for preparing a compound of the formula




30 wherein n is 1 or 2;

R² is (C₂-C₅)alkanoyloxy or -OR⁰;

R³ is (C₁ to C₃)alkyl or allyl;

R⁰ and R¹ are each H,

-37-



10

R^4 is, for each occurrence, independently $-\text{CO}_2R^8$, $-\text{CO}_2\text{H}$,
 H , $-\text{CH}_3$, $-\text{CH}_2\text{F}$, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{CONH}_2$, $-\text{CONHR}^8$, $-\text{CON}(\text{R}^8)_2$,
 $-\text{CH}_2\text{OCOR}^8$, $-\text{CH}_2\text{OCO}_2\text{R}^8$, $-\text{CH}_2\text{OCONR}_2^8$ or $-\text{CH}_2\text{OR}^8$;

15 R^5 is, for each occurrence, independently $(\text{C}_1 \text{ to } \text{C}_4)\text{alkoxy}$, benzyloxy, $-\text{OCOR}^8$, $-\text{OCOCH}_2\text{R}^8$, $-\text{OCO}_2\text{R}^8$ or $-\text{OSi}(\text{R}^8)_3$;

t is 1, 2 or 3;

m is 0 or 1; and

R^8 is $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_3\text{-C}_6)\text{cycloalkyl}$, allyl, pyridyl, thienyl, benzyl, benzyl variously
 20 substituted with one to five halogen atoms, $-\text{OH}$ groups or $(\text{C}_1\text{-C}_4)\text{alkoxy}$ groups, phenyl
 or phenyl variously substituted with one to five halogen atoms, $-\text{OH}$ groups or $(\text{C}_1\text{-C}_4)\text{alkoxy}$ groups;


provided that R^1 and R^0 are not both H ; when m is 0, R^4 is

$-\text{CH}_2\text{F}$, $-\text{CHF}_2$ or $-\text{CF}_3$; and in formulae (II) and (III), each ring carbon atom must bear

25 at least one hydrogen atom, comprising reacting a compound of the formula


30

-38-



wherein R^3 is (C_1 to C_3)alkyl or allyl and n is 1 or 2 with 2 to 4 molar equivalents of a compound selected from the group consisting of

15



wherein X is halo;

R^4 is, for each occurrence, independently $-\text{CO}_2R^8$, $-\text{CO}_2H$, H ,

$-\text{CH}_3$, $-\text{CH}_2F$, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{CONH}_2$, $-\text{CONHR}^8$, $-\text{CON}(R^8)_2$,

25 $-\text{CH}_2\text{OCOR}^8$, $-\text{CH}_2\text{OCO}_2R^8$, $-\text{CH}_2\text{OCONR}_2^8$ or $-\text{CH}_2\text{OR}^8$;

R^5 is, for each occurrence, independently (C_1 to C_4)alkoxy, benzyloxy, $-\text{OCOR}^8$, $-\text{OCOCH}_2R^8$, $-\text{OCO}_2R^8$ or $-\text{OSi}(R^8)_3$;

t is 1, 2 or 3;


m is 0 or 1; and

30 R^8 is (C_1 - C_6)alkyl, (C_3 - C_6)cycloalkyl, allyl, pyridyl, thienyl, benzyl, benzyl variously substituted with one to five halogen atoms, $-\text{OH}$ groups or (C_1 - C_4)alkoxy groups, phenyl or phenyl variously substituted with one to five halogen atoms, $-\text{OH}$ groups or (C_1 - C_4)alkoxy groups in the presence of a drying agent selected from the group consisting

-39-

of molecular sieves, calcium sulfate and magnesium sulfate; a base selected from the group consisting of mercuric carbonate, silver carbonate, mercuric nitrate and silver nitrate; and a catalyst selected from the group consisting of silver triflate, silver perchlorate, silver tetrafluoroborate, mercuric triflate, mercuric perchlorate and mercuric tetrafluoroborate in a reaction inert solvent at about -78°C to about -70°C with warming to about 0°C for about 0.5 to about 24 hours and subsequent stirring at room temperature for about 0.5 to 24 hours.

28. A process for preparing a compound of the formula



wherein n is 1 or 2;


R^2 is (C_2-C_5) alkanoyloxy or $-OR^0$;

R^3 is $(C_1 \text{ to } C_3)$ alkyl or allyl;

25 R^0 and R^1 are each H,

30

-40-




- 10 R^4 is, for each occurrence, independently $-\text{CO}_2R^8$, $-\text{CO}_2\text{H}$, $-\text{CH}_2\text{OH}$, H , $-\text{CH}_3$, $-\text{CH}_2\text{F}$, $-\text{CHF}_2$, $-\text{CF}_3$, $-\text{CONH}_2$, $-\text{CONHR}^8$, $-\text{CON}(\text{R}^8)_2$, $-\text{CH}_2\text{OCO}_2\text{R}^8$, $-\text{CH}_2\text{OCONR}_2^8$ or $-\text{CH}_2\text{OR}^8$;
 R^5 is, for each occurrence, independently $(\text{C}_1 \text{ to } \text{C}_4)\text{alkoxy}$, benzyloxy, $-\text{OH}$, $-\text{OCOCH}_2\text{R}^8$, $-\text{OCO}_2\text{R}^8$ or $-\text{OSi}(\text{R}^8)_3$;
 15 t is 1, 2 or 3;
 m is 0 or 1; and
 R^8 is $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_3\text{-C}_6)\text{cycloalkyl}$, allyl, pyridyl, thienyl, benzyl, benzyl variously substituted with one to five halogen atoms, $-\text{OH}$ groups or $(\text{C}_1\text{-C}_4)\text{alkoxy}$ groups, phenyl or phenyl variously substituted with one to five halogen atoms, $-\text{OH}$ groups or $(\text{C}_1\text{-C}_4)\text{alkoxy}$ groups;
 20 provided that R^1 and R^9 are not both H ; when m is 0, R^4 is $-\text{CH}_2\text{F}$, $-\text{CHF}_2$ or $-\text{CF}_3$; and in formulae (II) and (III), each ring carbon atom must bear at least one hydrogen atom comprising reacting a compound of the formula

25

30

-41-



wherein n is 1 or 2;


the dotted line represents an optional double bond in the case where R² is H;

15 R² is H, (C₂-C₅)alkanoyloxy or -OR⁰;

R³ is (C₁ to C₃)alkyl or allyl;

R⁰ and R¹ are each H,

20



25

R⁴ is, for each occurrence, independently -CO₂R⁸, -CO₂H,

30 H, -CH₃, -CH₂F, -CHF₂, -CF₃, -CONH₂, -CONHR⁸, -CON(R⁸)₂,
-CH₂OCOR⁸, -CH₂OCO₂R⁸, -CH₂OCONR₂⁸ or -CH₂OR⁸;

R⁵ is, for each occurrence, independently (C₁ to C₄)alkoxy, benzyloxy, -OCOR⁸, -
OCOCH₂R⁸, -OCO₂R⁸ or -OSi(R⁸)₃;

-42-

t is 1, 2 or 3;

m is 0 or 1; and

R⁸ is (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, allyl, pyridyl, thienyl, benzyl, benzyl variously substituted with one to five halogen atoms, -OH groups or (C₁-C₄)alkoxy groups, phenyl
5 or phenyl variously substituted with one to five halogen atoms, -OH groups or (C₁-C₄)alkoxy groups;

provided that R¹ and R⁰ are not both H; when m is 0, R⁴ is -CH₂F, -CHF₂ or -CF₃; and
in formulae (II) and (III), each ring carbon atom must bear at least one hydrogen atom
with a catalytic amount of an alkoxide base in an alcohol solvent at 0°C.

10

INTERNATIONAL SEARCH REPORT

PCT/US 93/00428

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC Int.Cl. 5 C07H19/01; A61K31/71		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C07H ; A61K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	EP,A,0 466 365 (MERCK & CO. INC.) 15 January 1992 see page 8, line 47 - line 52; claims 1-9 ---	1-19, 23-26
Y	EP,A,0 184 162 (FUJISAWA) 11 June 1986 see page 91, line 14 - page 92, line 11; claims 1,5,16-18 -----	1-19, 23-26
¹⁰ Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search 21 MAY 1993		Date of Mailing of this International Search Report 02.06.93
International Searching Authority EUROPEAN PATENT OFFICE		Signature of Authorized Officer BRENNAN J.

INTERNATIONAL SEARCH REPORT

I. national application No.

PCT/US 93/00428

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 20-22 are directed to a method of treatment of the human/animal body, the search has been carried out based on the alleged effects of the compounds.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9300428
SA 69663

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

21/05/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0466365	15-01-92	CA-A- 2046040	04-01-92
		JP-A- 4230399	19-08-92

EP-A-0184162	11-06-86	AU-B- 592067	04-01-90
		AU-A- 5059685	12-06-86
		CN-A- 1056103	13-11-91
		JP-C- 1686568	11-08-92
		JP-B- 3046445	16-07-91
		JP-A- 3072483	27-03-91
		JP-A- 3072484	27-03-91
		JP-C- 1670486	12-06-92
		JP-B- 3038276	10-06-91
		JP-A- 61148181	05-07-86
		US-A- 4956352	11-09-90
		US-A- 5110811	05-05-92
		US-A- 4894366	16-01-90
		US-A- 4929611	29-05-90
