

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12) Oversættelse af
europæisk patentskrift

(10) **DK/EP 3331876 T3**

-
- (51) Int.Cl.: **C 07 D 405/14 (2006.01)** **A 61 K 31/437 (2006.01)** **A 61 P 3/08 (2006.01)**
A 61 P 11/06 (2006.01) **A 61 P 17/06 (2006.01)** **A 61 P 25/08 (2006.01)**
A 61 P 25/18 (2006.01) **A 61 P 25/24 (2006.01)** **A 61 P 27/02 (2006.01)**
A 61 P 35/00 (2006.01) **A 61 P 37/00 (2006.01)** **C 07 D 235/10 (2006.01)**
C 07 D 235/12 (2006.01) **C 07 D 401/06 (2006.01)** **C 07 D 401/12 (2006.01)**
C 07 D 405/06 (2006.01) **C 07 D 471/04 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2021-01-11**
- (80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2020-10-07**
- (86) Europæisk ansøgning nr.: **16750587.4**
- (86) Europæisk indleveringsdag: **2016-08-03**
- (87) Den europæiske ansøgnings publiceringsdag: **2018-06-13**
- (86) International ansøgning nr.: **US2016045318**
- (87) Internationalt publikationsnr.: **WO2017024018**
- (30) Prioritet: **2015-08-05 US 201562201348 P** **2016-04-11 US 201662320890 P**
2016-05-26 US 201662341999 P
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV
MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Vitae Pharmaceuticals, LLC, 5 Giralda Farms, Madison, NJ 07940, USA**
- (72) Opfinder: **CLAREMON, David, A., 1508 Aidenn Lair Road, Maple Glen, PA 19002, USA**
DILLARD, Lawrence Wayne, 496 Kings Road, Yardley, PA 19067, USA
FAN, Yi, 40 Bittersweet Drive, Doylestown, PA 18901, USA
LOTESTA, Stephen, D., 99 Jennifer Lane, Burlington, NJ 08016, USA
SINGH, Suresh, B., 4 Adams Road, Kendall Park, NJ 08824, USA
TICE, Colin, M., 1325 Pinebrook Court, Maple Glen, PA 19002, USA
ZHAO, Wei, 11306 Royal Manor Way, North Potomac, MD 20878, USA
ZHUANG, Linghang, 3135 Fox Drive, Chalfont, PA 18914, USA
- (74) Fuldmægtig i Danmark: **Potter Clarkson A/S, Havnegade 39. kl, 1058 København K, Danmark**
- (54) Benævnelse: **MODULATORS OF ROR-GAMMA**
- (56) Fremdragne publikationer:
WO-A1-2013/019682
WO-A1-2014/044738
WO-A1-2015/159233

DESCRIPTION

TECHNICAL FIELD

[0001] The present disclosure is directed to novel retinoic acid receptor-related orphan receptor gamma ("ROR γ " or "ROR-gamma") modulators, processes for their preparation, pharmaceutical compositions containing these modulators, and their use in the treatment of inflammatory, metabolic, autoimmune and other diseases mediated by ROR γ .

BACKGROUND

[0002] Retinoic acid receptor-related orphan receptors (RORs) are a subfamily of transcription factors in the steroid hormone nuclear receptor superfamily (Jetten & Joo (2006) Adv. Dev. Biol. 2006, 16, 313-355). The ROR family consists of ROR alpha (ROR α), ROR beta (ROR β) and ROR gamma (ROR γ), each encoded by a separate gene (in human: *RORA*, *RORB* and *RORC*, respectively; in mouse: *rora*, *rorb* and *rorc*, respectively). RORs contain four principal domains shared by the majority of nuclear receptors: an N-terminal domain, a highly conserved DNA-binding domain (DBD) consisting of two zinc finger motifs, a hinge domain, and a ligand binding domain (LBD). Each ROR gene generates several isoforms, differing only in their N-terminal domains. ROR γ has two isoforms: ROR γ 1 and ROR γ 2 (also known as ROR γ t). ROR γ refers to ROR γ 1 and/or ROR γ t. ROR γ 1 is expressed in a variety of tissues including thymus, muscle, kidney and liver, but ROR γ t is exclusively expressed in the cells of the immune system, has a critical role in thymopoiesis and the development of several secondary lymphoid tissues, and is a key regulator of Th17 cell differentiation (Jetten, 2009, Nucl. Recept. Signal., 7:e003, doi:10.1621/nrs.07003, Epub 2009 Apr 3).

[0003] Th17 cells are a subset of T helper cells which preferentially produce the pro-inflammatory cytokines IL-17A, IL-17F, IL-21 and IL-22. Th17 cells and their effector molecules, such as IL-17, IL-21, IL-22, GM-CSF and CCL20, are associated with the pathogenesis of several autoimmune and inflammatory diseases, such as rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, psoriasis, inflammatory bowel disease, allergy and asthma (Maddur et al., 2012, Am. J. Pathol., 181:8-18). Recent findings support a role for IL17 and Th17 cells in the pathogenesis of acne (Thiboutot et al., 2014, J. Invest. Dermatol., 134(2):307-10, doi: 10.1038/jid.2013.400; Agak et al., 2014, J. Invest. Dermatol., 134(2):366-73, doi: 10.1038/jid.2013.334, Epub 2013 Aug 7). Th17 cells are also potent inducers of inflammation associated with endometriosis, a chronic inflammatory disease (Hirata et al., 2010, Endocrinol., 151:5468-5476; Hirata et al., 2011, Fertil Steril., Jul;96(1):113-7, doi: 10.1016/j.fertnstert.2011.04.060, Epub 2011 May 20). Additionally, Th17 cells have a key role in the mouse autoimmune models of experimental autoimmune encephalomyelitis (EAE), collagen-induced arthritis (CIA) and adjuvant-induced arthritis (AIA) (Bedoya et al., 2013, Clin. Dev. Immunol., 2013:986789. Epub 2013 Dec 26. Th17 cells are activated during inflammatory

and autoimmune disease processes and are responsible for recruiting other inflammatory cell types, particularly neutrophils, to mediate pathology in target tissues (Miossec & Kolls, 2012, *Nature Rev.*, 11:763-776; Korn et al., 2009, *Annu. Rev. Immunol.*, 27:485-517). Aberrant Th17 cell function has been implicated in a variety of autoimmune diseases, including multiple sclerosis and rheumatoid arthritis. Autoimmune disease is believed to arise from the disruption of the equilibrium between effector and regulatory T cells (Solt et al., 2012, *ACS Chem. Biol.*, 7:1515-1519, Epub 2012 July 9). The importance of ROR γ t to Th17 cell differentiation and the pathogenic role of Th17 cells is evidenced by the fact that ROR γ t-deficient mice have very few Th17 cells and have a reduction in severity of EAE (Ivanov et al., 2006, *Cell*, 126:1121-1133).

[0004] Recently, IL-17-producing neutrophils have been identified as promoting inflammation leading to both microbial clearance and IL-17-associated tissue damage in the cornea and other tissues (Taylor et al., 2014, *J. Immunol*, 192:3319-3327; Taylor et al., 2014, *Nat. Immunol.*, 15:143-151), supporting a role for compounds that inhibit ROR γ activity in the treatment of corneal ulcers and other diseases and disorders associated with IL-17 expressing neutrophils.

[0005] Circadian rhythms are daily cycles of behavioral and physiological changes that are regulated by endogenous circadian clocks. A number of studies have established links between nuclear receptor (including ROR γ) function and expression, the circadian regulatory circuitry, and the regulation of various physiological processes (Jetten (2009) op. cit.).

[0006] Obstructive sleep apnea syndrome (OSAS) is a chronic inflammatory disease regulated by T lymphocytes. OSAS patients have a significant increase in peripheral Th17 cell frequency, IL-17 and ROR γ t levels (Ye et al., 2012, *Mediators Inflamm.*, 815308, doi: 10.1155/2012/815308, Epub 2012 Dec 31).

[0007] A number of studies have provided evidence of a role of RORs in cancer. Mice deficient in the expression of ROR γ exhibit a high incidence of thymic lymphomas that metastasize frequently to liver and spleen. High expression of Th17-associated genes (including ROR γ) and high levels of Th17 cells in the tumor microenvironment has been shown to correlate with a poor prognosis in various cancers, including lung, gastric, breast and colon cancer (Tosolini et al., 2011, *Cancer Res.*, 71:1263-1271, doi: 10.1158/0008-5472.CAN-10-2907, Epub 2011 Feb 8; Su et al., 2014, *Immunol. Res.*, 58:118-124, doi: 10.1007/s12026-013-8483-y, Epub 2014 Jan 9; Carmi et al., 2011, *J. Immunol.*, 186:3462-3471, doi: 10.4049/jimmunol.1002901, Epub 2011 Feb 7; Chen et al., 2013, *Histopathology*, 63:225-233, doi: 10.1111/his.12156, Epub 2013 Jun 6). Recent evidence also shows that ROR γ is overexpressed and amplified in metastatic castration-resistant prostate cancer tumors, and that ROR γ antagonists suppressed tumor growth in multiple androgen receptor-expressing xenograft prostate cancer models. See e.g., *Nature Medicine*, March 28, 2016, advance online publication, doi: 10.1038/nm.4070.


[0008] ROR γ has also been identified to have a regulatory role in lipid/glucose homeostasis, and has been implicated in metabolic syndrome, obesity (Meissburger et al., 2011, *EMBO Mol. Med.*, 3:637-651), hepatosteatosis, insulin resistance and diabetes.

[0009] Further support for the role of ROR γ in the pathogenesis of inflammatory, metabolic, circadian effect, cancer, and autoimmune diseases and disorders can be found in the following references: Chang et al., 2012, J. Exp. Pharmacol., 4:141-148; Jetten et al., 2013, Frontiers Endocrinol., 4:1-8; Huh & Littman, 2012, Eur. J. Immunol., 42:2232-2237; Martinez et al., 2008, Ann. N.Y. Acad. Sci., 1143:188-211; Pantelyushin et al., 2012, J. Clin. Invest., 122:2252-2256; Jetten & Ueda, 2002, Cell Death Differen., 9:1167-1171; Solt et al., 2010, Curr. Opin. Lipidol., 21:204-211.

[0010] In light of the role that ROR γ plays in disease pathogenesis, inhibition of ROR γ activity and Th17 cell differentiation and activity, including IL17 production, will be of significant therapeutic benefit. It is therefore desirable to prepare compounds that inhibit ROR γ activity and hence have utility in the treatment of inflammatory, autoimmune, metabolic, circadian effect, cancer, and other diseases mediated by ROR γ , such as *e.g.*, asthma, atopic dermatitis, acne, Crohn's disease, regional enteritis, ulcerative colitis, Sjögren's syndrome, uveitis, Behçet's disease, dermatomyositis, multiple sclerosis, ankylosing spondylitis, systemic lupus erythematosus, scleroderma, psoriasis, psoriatic arthritis, steroid resistant asthma and rheumatoid arthritis.

SUMMARY

[0011] It has now been found that compounds disclosed herein, and pharmaceutically acceptable compositions thereof, are effective modulators of ROR γ (see *e.g.*, Table 1). Also described herein are compounds of Formula I.



or a pharmaceutically acceptable salt thereof, wherein each of Cy², L², R², A, Y, R⁹, p, X, n, L¹, and Cy¹ are as defined and described herein.

According to the present invention, there is provided a compound having the Formula IV or V in accordance with Claim 1.


[0012] The provided compounds, and pharmaceutically acceptable compositions thereof, are modulators of ROR γ and are useful for treating a variety of diseases, disorders or conditions. Such diseases, disorders, or conditions include those described herein.

[0013] The provided compounds can be used alone (*i.e.*, as a monotherapy) or in combination with one or more other therapeutic agent effective for treating any of the indications described herein.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

1. General Description of Compounds

[0014] Described herein is a compound of Formula I:



or a pharmaceutically acceptable salt thereof, wherein

X is -C(O)NH- or -NHC(O)-;

A is CH or N;

Y is CR¹⁰ or N;

n is 0, 1, 2, or 3;

p is 0, 1, or 2;

L² is CH₂, CH₂CH₂, CHMe, O, C(=O), CH(OH), or CH₂O, where the oxygen atom in CH₂O can either be attached to Cy² or to the carbon atom on the center ring bridging NR² and A;

L¹ is absent or is CR⁷R⁸;

Cy¹ is selected from aryl, heteroaryl, heterocyclyl, and cycloalkyl, wherein the aryl, heteroaryl, heterocyclyl, and cycloalkyl are each substituted with 1 to 3 groups independently selected from R⁵;

Cy² is selected from aryl, heteroaryl, monocyclic cycloalkyl, and monocyclic heterocyclyl, wherein the aryl, heteroaryl, monocyclic cycloalkyl, and monocyclic heterocyclyl are each optionally substituted with 1 to 3 groups independently selected from R⁶;

R² is (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₁-C₄)haloalkyl, or monocyclic cycloalkyl;

R⁵ and R⁶ are each independently selected from halogen, -CN, -OR^c, -NR^dR^e, -S(O)_kR^c, -NR^cS(O)₂R^c, -S(O)₂NR^dR^e, -C(=O)OR^c, -OC(=O)OR^c, -OC(=O)R^c, -OC(=S)OR^c, -C(=S)OR^c, -OC(=S)R^c, -C(=O)NR^dR^e, -NR^cC(=O)R^e, -C(=S)NR^dR^e, -NR^cC(=S)R^c, -NR^cC(=O)OR^c, -OC(=O)NR^dR^e, -NR^c(C=S)OR^c, -OC(=S)NR^dR^e, -NR^cC(=O)NR^dR^e, -NR^c(C=S)NR^dR^e, -C(=S)R^c, -C(=O)R^c, (C₁-C₆)alkyl, cycloalkyl, -(CH₂)₁₋₄-cycloalkyl, heterocyclyl, -(CH₂)₁₋₄-heterocyclyl, aryl, -NHC(=O)-heterocyclyl, -NHC(=O)-cycloalkyl, -(CH₂)₁₋₄-aryl, heteroaryl and -

(CH₂)₁₋₄-heteroaryl,

wherein the alkyl, cycloalkyl, heterocyclyl, aryl or heteroaryl portion present in each of said (C₁-C₆)alkyl, cycloalkyl, -(CH₂)₁₋₄-cycloalkyl, heterocyclyl, -(CH₂)₁₋₄-heterocyclyl, aryl, -(CH₂)₁₋₄-aryl, heteroaryl and -(CH₂)₁₋₄-heteroaryl substituent for R⁶ are further optionally substituted with halogen, OR^c, -NO₂, -CN, -NR^cC(=O)R^c, -NR^dR^e, -S(O)_kR^c, -C(=O)OR^c, -C(=O)NR^dR^e, -C(=O)R^c, (C₁-C₃)alkyl, halo(C₁-C₃)alkyl, (C₁-C₃)alkoxy(C₁-C₃)alkyl, (C₁-C₃)alkoxy, or halo(C₁-C₃)alkoxy;

each R^c is independently selected from hydrogen and (C₁-C₆)alkyl optionally substituted with 1 to 3 halogen;

each R^d and R^e is independently selected from hydrogen and (C₁-C₆)alkyl;

k is 0, 1 or 2;

any heterocyclyl or heteroaryl portion of Cy¹ or Cy² is further optionally substituted with =O;

R⁷ and R⁸ are each independently hydrogen, OH, (C₁-C₃)alkyl, hydroxy(C₁-C₃)alkyl, -O(C₁-C₃)alkyl, -(C₁-C₃)alkyl-O-(C₁-C₃)alkyl, -(C₁-C₃)alkyl-NH₂, -(C₁-C₃)alkyl-N-di(C₁-C₃)alkyl, CO₂H, (CH₂)₁₋₃COOH, monocyclic heterocyclyl, -CO-(C₁-C₃)alkoxy, -(C₁-C₃)alkoxy-C(O)-NH₂, -(C₁-C₃)alkyl-O-(C₁-C₃)alkyl-C(O)-NH₂, -(C₁-C₃)alkyl-C(O)-(C₁-C₃)alkoxy, -(C₁-C₃)alkyl-C(O)-NH₂, halophenyl, -(C₁-C₃)alkyl-halophenyl, or quinolin-2(1H)one-4yl-methyl;

R⁹ is independently (C₁-C₄)alkyl, (C₁-C₄)haloalkyl, (C₁-C₄)alkoxy, or halogen; and

R¹⁰ is hydrogen, (C₁-C₄)alkyl, (C₁-C₄)haloalkyl, (C₁-C₄)alkoxy, or halogen;

provided that:

when A is N, Y is CR¹⁰, R¹⁰ is hydrogen, R² is cyclohexyl, L² is CH₂, Cy² is phenyl, X is -C(O)NH, n is 0, p is 0, L¹ is CR⁷R⁸, and one of R⁷ or R⁸ is COOH and the other is hydrogen, then Cy¹ is not 1H-indol-5-yl-3yl;

when A is N, Y is CR¹⁰, R¹⁰ is hydrogen, R² is (C₁-C₄)alkyl, X is -NHC(O), L² is CH₂, p is 0, n is 0, L¹ is CR⁷R⁸, and both of R⁷ and R⁸ are hydrogen, then Cy¹ is not piperidinyl; and

when A is N, Y is CR¹⁰, R¹⁰ is hydrogen, R² is methyl, Cy¹ is aryl, L² is CH₂, p is 0, X is -C(O)NH, n is 0, L¹ is CR⁷R⁸, and both of R⁷ or R⁸ are hydrogen, then Cy² is not phenyl substituted with CN.

2. Compounds and Definitions

[0015] The terms "halo" and "halogen" as used herein refer to an atom selected from fluorine (fluoro, -F), chlorine (chloro, -Cl), bromine (bromo, -Br), and iodine (iodo, -I).

[0016] The term "alkyl", used alone or as a part of a larger moiety such as e.g., "haloalkyl", means a saturated monovalent straight or branched hydrocarbon radical having, unless otherwise specified, 1-10 carbon atoms and includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, n-decyl and the like. "Monovalent" means attached to the rest of the molecule at one point.

[0017] The term "haloalkyl" or "halocycloalkyl" include mono, poly, and perhaloalkyl groups where the halogens are independently selected from fluorine, chlorine, and bromine.

[0018] The terms "cycloalkyl" and "cycloaliphatic", used alone or as part of a larger moiety, refer to a saturated cyclic aliphatic monocyclic or bicyclic ring system, as described herein, having from, unless otherwise specified, 3 to 10 carbon ring atoms. Monocyclic cycloalkyl groups include, without limitation, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl, cycloheptenyl, and cyclooctyl. It will be understood that when specified, optional substituents on a cycloalkyl or cycloaliphatic group may be present on any substitutable position and, include, e.g., the position at which the cycloalkyl or cycloaliphatic group is attached.

[0019] The term "carbocycle", "carbocyclyl", "carbocyclo", or "carbocyclic" used alone or as part of a larger moiety refer to saturated, partially saturated, or aromatic ring systems comprising all carbon atoms having, unless otherwise specified, a total of 3 to 10 ring members. It will be understood that when specified, optional substituents on a carbocycle, carbocyclyl, carbocyclo, or carbocyclic may be present on any substitutable position and, include, e.g., the position at which the cycloalkyl is attached.

[0020] The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy", or "aryloxyalkyl", refers to an aromatic carbocyclic ring system having, unless otherwise specified, a total of 6 to 10 ring members. The term "aryl" may be used interchangeably with the term "aryl ring", "aryl group", "aryl moiety," or "aryl radical". In certain embodiments of the present disclosure, "aryl" refers to an aromatic ring system which includes, but is not limited to, phenyl (abbreviated as "Ph"), naphthyl and the like. It will be understood that when specified, optional substituents on an aryl group may be present on any substitutable position and, include, e.g., the position at which the aryl is attached.

[0021] The term "heteroaryl" used alone or as part of a larger moiety as in "heteroarylalkyl", "heteroarylalkoxy", or "heteroarylaminoalkyl", refers to a 5-10 -membered aromatic radical containing 1-4 heteroatoms selected from N, O, and S and includes, for example, thienyl,

furanyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolizinyl, purinyl, naphthyridinyl, and pteridinyl. The term "heteroaryl" may be used interchangeably with the terms "heteroaryl ring", "heteroaryl group", or "heteroaromatic". The terms "heteroaryl" and "heteroar-", as used herein, also include groups in which a heteroaromatic ring is fused to one or more aryl rings, where the radical or point of attachment is on the heteroaromatic ring. Nonlimiting examples include indolyl, indazolyl, benzimidazolyl, benzthiazolyl, quinolyl, quinazolinyl, and quinoxalinyl. A heteroaryl group may be mono- or bicyclic. It will be understood that when specified, optional substituents on a heteroaryl group may be present on any substitutable position and, include, e.g., the position at which the heteroaryl is attached.

[0022] The term "heterocyclyl" means a 4-, 5-, 6- and 7-membered saturated or partially unsaturated heterocyclic ring containing 1 to 4 heteroatoms independently selected from N, O, and S. The terms "heterocycle", "heterocyclyl", "heterocyclyl ring", "heterocyclic group", "heterocyclic moiety", and "heterocyclic radical", are used interchangeably herein. A heterocyclyl ring can be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure. Examples of such saturated or partially unsaturated heterocyclic radicals include, without limitation, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, pyrrolidinyl, pyrrolidonyl, piperidinyl, oxazolidinyl, piperazinyl, dioxanyl, dioxolanyl, morpholinyl, dihydrofuranyl, dihydropyranyl, dihydropyridinyl, tetrahydropyridinyl, dihydropyrimidinyl, and tetrahydropyrimidinyl. A heterocyclyl group may be mono- or bicyclic. Unless otherwise specified, bicyclic heterocyclyl groups include, e.g., unsaturated heterocyclic radicals fused to another unsaturated heterocyclic radical or aromatic or heteroaryl ring, such as for example, tetrahydronaphthyridine, indolinone, dihydropyrrolotriazole, imidazopyrimidine, quinolinone, dioxaspirodecane. It will also be understood that when specified, optional substituents on a heterocyclyl group may be present on any substitutable position and, include, e.g., the position at which the heterocyclyl is attached.

[0023] As used herein the terms "subject" and "patient" may be used interchangeably, and means a mammal in need of treatment, e.g., companion animals (e.g., dogs, cats, and the like), farm animals (e.g., cows, pigs, horses, sheep, goats and the like) and laboratory animals (e.g., rats, mice, guinea pigs and the like). Typically, the subject is a human in need of treatment.

[0024] Certain of the disclosed compounds may exist in various stereoisomeric forms. Stereoisomers are compounds that differ only in their spatial arrangement. Enantiomers are pairs of stereoisomers whose mirror images are not superimposable, most commonly because they contain an asymmetrically substituted carbon atom that acts as a chiral center. "Enantiomer" means one of a pair of molecules that are mirror images of each other and are not superimposable. Diastereomers are stereoisomers that contain two or more asymmetrically substituted carbon atoms. The symbol "*" in a structural formula represents the presence of a chiral carbon center. "R" and "S" represent the configuration of substituents around one or more chiral carbon atoms. Thus, "R*" and "S*" denote the relative configurations of substituents around one or more chiral carbon atoms.

[0025] As used herein, a hyphen ("-") at the beginning or end of a recited group designates the point at which a recited group is attached to a defined group. For example, -SO₂-(C₁-C₃)alkyl-(C₂-C₆)cycloalkyl means that the group is attached via the sulfonyl.

[0026] "Racemate" or "racemic mixture" means a compound of equimolar quantities of two enantiomers, wherein such mixtures exhibit no optical activity, *i.e.*, they do not rotate the plane of polarized light.

[0027] "Geometric isomer" means isomers that differ in the orientation of substituent atoms in relationship to a carbon-carbon double bond, to a cycloalkyl ring, or to a bridged bicyclic system. Atoms (other than H) on each side of a carbon-carbon double bond may be in an E (substituents are on opposite sides of the carbon-carbon double bond) or Z (substituents are oriented on the same side) configuration. "R," "S," "S*," "R*," "E," "Z," "cis," and "trans," indicate configurations relative to the core molecule. When a disclosed compound is named or depicted by structure without indicating a particular geometric isomer form, it is to be understood that the name or structure encompasses one geometric isomer free of other geometric isomers, mixtures of geometric isomers, or all geometric isomers.

[0028] The compounds of the herein may be prepared as individual enantiomers by either enantio-specific synthesis or resolved from an enantiomerically enriched mixture. Conventional resolution techniques include forming the salt of a free base of each isomer of an enantiomeric pair using an optically active acid (followed by fractional crystallization and regeneration of the free base), forming the salt of the acid form of each enantiomer of an enantiomeric pair using an optically active amine (followed by fractional crystallization and regeneration of the free acid), forming an ester or amide of each of the enantiomers of an enantiomeric pair using an optically pure acid, amine or alcohol (followed by chromatographic separation and removal of the chiral auxiliary), or resolving an enantiomeric mixture of either a starting material or a final product using various well known chromatographic methods.

[0029] When the stereochemistry of a disclosed compound is named or depicted by structure, the named or depicted stereoisomer is at least 60%, 70%, 80%, 90%, 99% or 99.9% by weight pure relative to all of the other stereoisomers. Percent by weight pure relative to all of the other stereoisomers is the ratio of the weight of one stereoisomer over the weight of the the other stereoisomers. When a single enantiomer is named or depicted by structure, the depicted or named enantiomer is at least 60%, 70%, 80%, 90%, 99% or 99.9% by weight optically pure. Percent optical purity by weight is the ratio of the weight of the enantiomer over the weight of the enantiomer plus the weight of its optical isomer.

[0030] When the stereochemistry of a disclosed compound is named or depicted by structure, and the named or depicted structure encompasses more than one stereoisomer (*e.g.*, as in a diastereomeric pair), it is to be understood that one of the encompassed stereoisomers or any mixture of the encompassed stereoisomers are included. It is to be further understood that the stereoisomeric purity of the named or depicted stereoisomers at least 60%, 70%, 80%, 90%,

99% or 99.9% by weight pure relative to all of the other stereoisomers. The stereoisomeric purity in this case is determined by dividing the total weight in the mixture of the stereoisomers encompassed by the name or structure by the total weight in the mixture of all of the stereoisomers.

[0031] When a disclosed compound is named or depicted by structure without indicating the stereochemistry, and the compound has one chiral center, it is to be understood that the name or structure encompasses one enantiomer of compound free from the corresponding optical isomer, a racemic mixture of the compound and mixtures enriched in one enantiomer relative to its corresponding optical isomer.

[0032] When a disclosed compound is named or depicted by structure without indicating the stereochemistry and *e.g.*, the compound has at least two chiral centers, it is to be understood that the name or structure encompasses one stereoisomer free of other stereoisomers, mixtures of stereoisomers, and mixtures of stereoisomers in which one or more stereoisomers is enriched relative to the other stereoisomer(s). For example, the name or structure may encompass one stereoisomer free of other diastereomers, mixtures of stereoisomers, and mixtures of stereoisomers in which one or more diastereomers is enriched relative to the other diastereomer(s).


[0033] The compounds of the herein may be present in the form of pharmaceutically acceptable salts. For use in medicines, the salts of the compounds of the invention refer to nontoxic "pharmaceutically acceptable salts." Pharmaceutically acceptable salt forms include pharmaceutically acceptable acidic/anionic or basic/cationic salts.

[0034] Pharmaceutically acceptable basic/cationic salts include, the sodium, potassium, calcium, magnesium, diethanolamine, n-methyl-D-glucamine, L-lysine, L-arginine, ammonium, ethanolamine, piperazine and triethanolamine salts.

[0035] Pharmaceutically acceptable acidic/anionic salts include, *e.g.*, the acetate, benzenesulfonate, benzoate, bicarbonate, bitartrate, carbonate, citrate, dihydrochloride, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrobromide, hydrochloride, malate, maleate, malonate, mesylate, nitrate, salicylate, stearate, succinate, sulfate, tartrate, and tosylate.


3. Description of Exemplary Compounds

[0036] Described herein is a compound of Formula I,




or a pharmaceutically acceptable salt thereof, wherein the variables are as described above.

[0037] Also described herein is a compound of Formula I,




or a pharmaceutically acceptable salt thereof, wherein the variables are as described above, provided that when A is N, Y is CR¹⁰, R¹⁰ is hydrogen, and R² is monocyclic cycloalkyl, neither R⁷ nor R⁸ are (C₁-C₃)alkyl; and when A is N, Y is CR¹⁰, R¹⁰ is hydrogen, and R² is (C₄)alkyl, neither R⁷ nor R⁸ are (C₁-C₃)alkyl.


[0038] Also described herein is a compound of Formula II:




or a pharmaceutically acceptable salt thereof, wherein R⁷ and R⁸ are each independently hydrogen, OH, (C₁-C₂)alkyl, hydroxy(C₁-C₃)alkyl, -O(C₁-C₃)alkyl, -(C₁-C₃)alkyl-O-(C₁-C₃)alkyl, -(C₁-C₃)alkyl-NH₂, -(C₁-C₃)alkyl-N-di(C₁-C₃)alkyl, (CH₂)₁₋₃COOH, monocyclic heterocyclyl, -CO-(C₁-C₃)alkoxy, -(C₁-C₃)alkoxy-C(O)-NH₂, -(C₁-C₃)alkoxy-O-(C₁-C₃)alkyl-C(O)-NH₂, -(C₁-C₃)alkyl-C(O)-(C₁-C₃)alkoxy, -(C₁-C₃)alkyl-C(O)-NH₂, halophenyl, -(C₁-C₃)alkyl-halophenyl, or quinolin-2(1H)one-4yl-methyl, and wherein the variables in structural Formula II are as described for Formula I. Alternatively, as herein described, the compound of Formula I is of Formula II:



or a pharmaceutically acceptable salt thereof, wherein R⁷ and R⁸ are each independently hydrogen, OH, (C₁-C₂)alkyl, hydroxy(C₁-C₃)alkyl, -O(C₁-C₃)alkyl, -(C₁-C₃)alkyl-O-(C₁-C₃)alkyl, -(C₁-C₃)alkyl-NH₂, -(C₁-C₃)alkyl-N-di(C₁-C₃)alkyl, (CH₂)₁₋₃COOH, monocyclic heterocyclyl, -CO-(C₁-C₃)alkoxy, -(C₁-C₃)alkoxy-C(O)-NH₂, -(C₁-C₃)alkoxy-O-(C₁-C₃)alkyl-C(O)-NH₂, -(C₁-C₃)alkyl-C(O)-(C₁-C₃)alkoxy, -(C₁-C₃)alkyl-C(O)-NH₂, halophenyl, -(C₁-C₃)alkyl-halophenyl, or quinolin-2(1H)one-4yl-methyl; and p is 1, and wherein the variables in structural Formula II are as described for Formula I. In another alternative described herein, the compound of Formula I is of Formula II:






(II);

or a pharmaceutically acceptable salt thereof, wherein R^7 and R^8 are each independently hydrogen, OH, (C_1-C_2) alkyl, hydroxy (C_1-C_3) alkyl, $-O(C_1-C_3)$ alkyl, $-(C_1-C_3)$ alkyl- $O-(C_1-C_3)$ alkyl, $-(C_1-C_3)$ alkyl- NH_2 , $-(C_1-C_3)$ alkyl- $N-di(C_1-C_3)$ alkyl, $(CH_2)_{1-3}COOH$, monocyclic heterocyclyl, $-CO-(C_1-C_3)$ alkoxy, $-(C_1-C_3)$ alkoxy- $C(O)-NH_2$, $-(C_1-C_3)$ alkoxy- $O-(C_1-C_3)$ alkyl- $C(O)-NH_2$, $-(C_1-C_3)$ alkyl- $C(O)-(C_1-C_3)$ alkoxy, $-(C_1-C_3)$ alkyl- $C(O)-NH_2$, halophenyl, $-(C_1-C_3)$ alkyl-halophenyl, or quinolin-2(1H)one-4yl-methyl; p is 1; and R^9 is (C_1-C_4) alkoxy or halogen, and wherein the variables in structural Formula II are as described for Formula I.


[0039] Also described herein is a compound of Formula IIa:



(IIa);

or a pharmaceutically acceptable salt thereof, wherein the variables in structural Formula III are as specified for Formulae I or II as herein described.


[0040] Also described herein is a compound of Formula III:



(III);

or a pharmaceutically acceptable salt thereof, wherein the variables in structural Formula III are as specified for Formulae I or II as herein described.

[0041] Also described herein is a compound of Formula IIIa:






(IIIa);


or a pharmaceutically acceptable salt thereof, wherein the variables in structural Formula IIIa are as described for Formula I or the third embodiment.

[0042] L^2 in Formulas I to III described herein is CH_2 , $CHMe$, O , or $C(=O)$, wherein the remaining variables are as described for Formulae I or II.

[0043] In an embodiment of the present invention, there is provided a compound of Formula IV, IVa, V, or Va:



or



or a pharmaceutically acceptable salt thereof, wherein

A is CH or N;

Y is CH

Cy¹ is phenyl or pyridyl, each substituted with 1 to 3 groups independently selected from R⁵, wherein at least one R⁵ is -SO₂-(C₁-C₃)alkyl

Cy² is selected from phenyl, pyridinyl, piperidinyl, cyclohexyl, tetrahydropyranyl, cyclopropyl, and cyclobutyl, each optionally substituted with 1 to 3 groups independently selected from R⁶

R² is (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₁-C₄)haloalkyl, or monocyclic cycloalkyl;

R⁵ and R⁶ are each independently selected from halogen, -CN, -ORᶜ, -NRᵈRᵉ, -S(O)ₖRᶜ, -NRᶜS(O)₂Rᶜ, -S(O)₂NRᵈRᵉ, -C(=O)ORᶜ, -OC(=O)ORᶜ, -OC(=O)Rᶜ, -OC(=S)ORᶜ, -C(=S)ORᶜ,-

$\text{OC}(=\text{S})\text{R}^c$, $-\text{C}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c\text{C}(=\text{O})\text{R}^c$, $-\text{C}(=\text{S})\text{NR}^d\text{R}^e$, $-\text{NR}^c\text{C}(=\text{S})\text{R}^c$, $-\text{NR}^c\text{C}(=\text{O})\text{OR}^c$,
 $\text{OC}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c(\text{C}=\text{S})\text{OR}^c$, $-\text{OC}(=\text{S})\text{NR}^d\text{R}^e$, $-\text{NR}^c\text{C}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{NR}^c(\text{C}=\text{S})\text{NR}^d\text{R}^e$,
 $\text{C}(=\text{S})\text{R}^c$, $-\text{C}(=\text{O})\text{R}^c$, $(\text{C}_1\text{-C}_6)\text{alkyl}$, cycloalkyl , $-(\text{CH}_2)_{1-4}\text{-cycloalkyl}$, heterocyclyl , $-(\text{CH}_2)_{1-4}\text{-}$
 heterocyclyl , aryl , $-\text{NHC}(=\text{O})\text{-heterocyclyl}$, $-\text{NHC}(=\text{O})\text{-cycloalkyl}$, $-(\text{CH}_2)_{1-4}\text{-aryl}$, heteroaryl and $-(\text{CH}_2)_{1-4}\text{-heteroaryl}$,

wherein the alkyl , cycloalkyl , heterocyclyl , aryl or heteroaryl portion present in each of said $(\text{C}_1\text{-C}_6)\text{alkyl}$, cycloalkyl , $-(\text{CH}_2)_{1-4}\text{-cycloalkyl}$, heterocyclyl , $-(\text{CH}_2)_{1-4}\text{-heterocyclyl}$, aryl , $-(\text{CH}_2)_{1-4}\text{-aryl}$,
 heteroaryl and $-(\text{CH}_2)_{1-4}\text{-heteroaryl}$ substituent for R^6 are further optionally substituted with
 halogen , OR^c , $-\text{NO}_2$, $-\text{CN}$, $-\text{NR}^c\text{C}(=\text{O})\text{R}^c$, $-\text{NR}^d\text{R}^e$, $-\text{S}(\text{O})_k\text{R}^c$, $-\text{C}(=\text{O})\text{OR}^c$, $-\text{C}(=\text{O})\text{NR}^d\text{R}^e$, $-\text{C}(=\text{O})\text{R}^c$, $(\text{C}_1\text{-C}_3)\text{alkyl}$, $\text{halo}(\text{C}_1\text{-C}_3)\text{alkyl}$, $(\text{C}_1\text{-C}_3)\text{alkoxy}(\text{C}_1\text{-C}_3)\text{alkyl}$, $(\text{C}_1\text{-C}_3)\text{alkoxy}$, or $\text{halo}(\text{C}_1\text{-C}_3)\text{alkoxy}$;

each R^c is independently selected from hydrogen and $(\text{C}_1\text{-C}_6)\text{alkyl}$ optionally substituted with 1 to 3 halogen;

each R^d and R^e is independently selected from hydrogen and $(\text{C}_1\text{-C}_6)\text{alkyl}$;

k is 0, 1 or 2;



any heterocyclyl or heteroaryl portion of Cy^1 or Cy^2 is further optionally substituted with $=\text{O}$; and



R^7 is hydrogen, $(\text{C}_1\text{-C}_2)\text{alkyl}$, $\text{hydroxy}(\text{C}_1\text{-C}_3)\text{alkyl}$, $-(\text{C}_1\text{-C}_3)\text{alkylNH}_2$, $-(\text{C}_1\text{-C}_3)\text{alkyl-N-di}(\text{C}_1\text{-C}_3)\text{alkyl}$, or $-(\text{C}_1\text{-C}_3)\text{alkoxy-C}(\text{O})\text{-NH}_2$,

provided that:


when A is N , Y is CH , R^2 is methyl, Cy^1 is phenyl and the compound is of formula IV, then Cy^2 is not phenyl substituted with CN

[0044] In another embodiment of the present invention, there is provided a compound of Formula VI, VIa, VII, or VIIa:






or




or a pharmaceutically acceptable salt thereof, wherein the variables in structural Formula VI and structural Formula VII are as described for Formulae IV, IVa, V and Va.

[0045] . In one embodiment of the present invention, R⁷ is hydroxy(C₁-C₃)alkyl.

[0046] According to another embodiment of the present invention, there is provided a compound of Formula VIII or Formula IX:



or



or a pharmaceutically acceptable salt thereof, wherein the variables in structural Formula (VIII) or structural Formula (IX) are as described for Formulae IV, IVa, V and Va.

[0047] In another embodiment, R² in the compounds according to the present invention is (C₁-C₄)alkyl or cyclopropyl.

[0048] As described herein, Cy¹ in Formulae I to IX is aryl or heteroaryl, each substituted with

1 to 3 groups independently selected from R^5 ,...

[0049] . In one embodiment of the present invention, Cy^1 in Formulae IV to IX is pyridinyl substituted with 1 to 3 groups independently selected from R^5 , wherein at least one R^5 is $-SO_2-(C_1-C_3)alkyl$.

[0050] In an embodiment of the present invention, Cy^2 is selected from phenyl, pyridinyl, cyclohexyl, tetrahydropyranyl, cyclopropyl, and cyclobutyl, each optionally substituted with 1 to 3 groups independently selected from R^6 .

[0051] In another embodiment of the present invention, Cy^2 is selected from phenyl, pyridinyl, piperidinyl, cyclohexyl, or tetrahydropyranyl, each optionally substituted with 1 to 3 groups independently selected from R^6 .

[0052] In an embodiment of the present invention, Cy^2 is selected from phenyl and cyclohexyl, each optionally substituted with 1 to 3 groups independently selected from R^6 .

[0053] In another embodiment of the present invention, Cy^1 is phenyl substituted with 1 to 3 groups independently selected from R^5 , wherein at least one R^5 is $-SO_2-(C_1-C_3)alkyl$; and Cy^2 is cyclohexyl substituted with 1 to 3 groups independently selected from R^6 .

[0054] In another embodiment of the present invention, R^5 is selected from halogen, $-CN$, $-OR^c$, $-NR^dR^e$, $-NR^cS(O)_2R^c$, $-S(O)_2NR^dR^e$, $-C(=O)OR^c$, $-C(=O)NR^dR^e$, $-NR^cC(=O)R^c$, $-NR^cC(=O)OR^c$, $-OC(=S)NR^dR^e$, $-C(=O)R^c$, $-SO_2-(C_1-C_3)alkyl$, and $(C_1-C_4)alkyl$ optionally substituted with halogen; and R^6 is selected from halogen, $-CN$, $-OR^c$, $-NR^dR^e$, $-NR^cS(O)_2R^c$, $-S(O)_2NR^dR^e$, $-C(=O)OR^c$, $-OC(=O)OR^c$, $-OC(=O)R^c$, $-C(=O)NR^dR^e$, $-NR^cC(=O)R^c$, $-C(=S)NR^dR^e$, $-NR^cC(=S)R^c$, $-NR^cC(=O)OR^c$, $-OC(=O)NR^dR^e$, $-NR^c(C=S)OR^c$, $-OC(=S)NR^dR^e$, $-NR^cC(=O)NR^dR^e$, $-NR^c(C=S)NR^dR^e$, $-C(=S)R^c$, $-C(=O)R^c$, $-SO_2-(C_1-C_3)alkyl$, and $(C_1-C_4)alkyl$ optionally substituted with halogen.

[0055] In another embodiment of the present invention, R^5 is $-SO_2-(C_1-C_3)alkyl$; R^6 is selected from halo, $-OR^c$, and $(C_1-C_4)alkyl$ optionally substituted with halogen and R^c is $(C_1-C_4)alkyl$ optionally substituted with halogen.

[0056] In a further embodiment of the present invention, R^6 is CF_3 .

[0057] Specific examples of compounds are provided in the EXEMPLIFICATION.

Pharmaceutically acceptable salts as well as the neutral forms of these compounds are included herein.

[0058] In certain embodiments, the present disclosure provides a method of treating a patient (e.g., a human) with a disorder mediated by ROR γ comprising the step of administering to the patient an effective amount of the compound with any compound described herein, or a pharmaceutically acceptable salt or composition thereof.

4. Uses, Formulation and Administration

Pharmaceutically acceptable compositions

[0059] According to another embodiment, the present disclosure provides a method of treating a subject (e.g., a human) with a disorder mediated by ROR γ using a composition comprising a compound of the present invention and a pharmaceutically acceptable carrier, adjuvant, or vehicle. In certain embodiments, the amount of compound of the present invention in a provided composition is such that it is effective as an inverse agonist or antagonist to ROR γ in a biological sample or in a subject. In certain embodiments, a provided composition is formulated for administration to a subject in need of such composition. In some embodiments, a provided composition is formulated for oral administration to a subject.

[0060] The term "pharmaceutically acceptable carrier, adjuvant, or vehicle" refers to a nontoxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the compositions of this disclosure include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[0061] Compositions described herein may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques.

[0062] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and

elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[0063] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[0064] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[0065] In order to prolong the effect of a provided compound, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[0066] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic

acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar--agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[0067] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[0068] Provided compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[0069] Dosage forms for topical or transdermal administration of a compound herein include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, and eye drops are also contemplated as being within the scope herein. Additionally, the present disclosure contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption

enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[0070] Pharmaceutically acceptable compositions provided herein may be formulated for oral administration. Such formulations may be administered with or without food. In some embodiments, pharmaceutically acceptable compositions of this disclosure are administered without food. In other embodiments, pharmaceutically acceptable compositions of this disclosure are administered with food.

[0071] The amount of provided compounds that may be combined with carrier materials to produce a composition in a single dosage form will vary depending upon the patient to be treated and the particular mode of administration.

[0072] It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, the judgment of the treating physician, and the severity of the particular disease being treated. The amount of a provided compound in the composition will also depend upon the particular compound in the composition.

Uses of Compounds and Pharmaceutically Acceptable Compositions

[0073] Compounds and compositions disclosed herein are generally useful for modulating ROR γ . Thus, in some embodiments, the present disclosure provides a method of treating inflammatory, metabolic and autoimmune diseases or disorders mediated by ROR γ , comprising administering a provided compound or composition. More particularly, the compounds and compositions described herein act as inverse agonists or antagonists of ROR γ .

[0074] As used herein, the terms "treatment," "treat," and "treating" refer to reversing, alleviating, delaying the onset of, or inhibiting the progress of a disease or disorder, or one or more symptoms thereof, as described herein. In some embodiments, treatment may be administered after one or more symptoms have developed, *i.e.*, therapeutic treatment. In other embodiments, treatment may be administered in the absence of symptoms. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms (*e.g.*, in light of a history of symptoms and/or in light of genetic or other susceptibility factors), *i.e.*, prophylactic treatment. Treatment may also be continued after symptoms have resolved, for example to prevent or delay their recurrence.

[0075] Modulation of ROR γ (or to modulate ROR γ), means that a change or alternation in the activity of ROR γ has occurred from the administration of one or more of the compounds disclosed herein. Modulation may be an upregulation (increase) or a downregulation

(decrease) in the magnitude of the activity or function of ROR γ . Exemplary activities and functions include e.g., binding characteristics, enzymatic activity, cell receptor activation, transcriptional activity, and signal transduction. In one aspect, the compounds disclosed herein inhibit ROR γ . In further aspects, the compounds disclosed herein act as agonists, antagonists, or inverse agonists of ROR γ .

[0076] In another aspect, compounds and compositions disclosed herein are useful for reducing the amount of IL-17 in a subject. Thus, in some embodiments, provided herein are methods of reducing the amount of IL-17 in a subject comprising administering an effective amount of a provided compound or composition. ROR γ modulators disclosed in WO 2014/179564, WO 2015/116904, WO 2016/061160 and U.S. patent application Nos. 14/933,468 and 14/933,524 can also be used in such methods.

[0077] In another aspect, compounds and compositions disclosed herein are useful for inhibiting the synthesis IL-17 in a subject. Thus, in some embodiments, provided herein are methods of inhibiting the synthesis of IL-17 in a subject comprising administering an effective amount of a provided compound or composition. ROR γ modulators disclosed in WO 2014/179564, WO 2015/116904, WO 2016/061160 and U.S. patent application Nos. 14/933,468 and 14/933,524 can also be used in such methods.

[0078] Diseases and conditions treatable according to the methods disclosed herein include, but are not limited to, inflammatory, metabolic and autoimmune diseases or disorders mediated by ROR γ . These diseases and conditions include, for example, asthma, chronic obstructive pulmonary disease (COPD), bronchitis, allergic rhinitis, atopic dermatitis, contact dermatitis, acne, cystic fibrosis, allograft rejection, multiple sclerosis, scleroderma, arthritis, rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, systemic lupus erythematosus (SLE), psoriasis, Hashimoto's disease, pancreatitis, autoimmune diabetes, type I diabetes, autoimmune ocular disease, ulcerative colitis, Crohn's disease, regional enteritis, inflammatory bowel disease (IBD), inflammatory bowel syndrome (IBS), Sjögren's syndrome, optic neuritis, obesity, hepatosteatosis, adipose tissue-associated inflammation, insulin resistance, type II diabetes, neuromyelitis optica, myasthenia gravis, age related macular degeneration, dry eye, uveitis, Guillain-Barré syndrome, psoriasis, psoriatic arthritis (PsA), steroid resistant asthma, Graves' disease, scleritis, endometriosis, obstructive sleep apnea syndrome (OSAS), Behçet's disease, dermatomyositis, polymyositis, graft versus host disease, primary biliary cirrhosis, liver fibrosis, non-alcoholic fatty liver disease (NAFLD), sarcoidosis, primary sclerosing cholangitis, autoimmune thyroid disease, autoimmune polyendocrine syndrome type I, autoimmune polyendocrine syndrome type II, celiac disease, neuromyelitis, juvenile idiopathic arthritis, systemic sclerosis, myocardial infarction, pulmonary hypertension, osteoarthritis, cutaneous leishmaniasis, sinonasal polyposis, and cancer, including but not limited to lung cancer, gastric cancer, breast cancer and colon cancer. In one aspect, an exemplified form of cancer treatable according to the methods herein also includes prostate cancer e.g., (metastatic castration-resistant prostate cancer tumors). In another aspect, an exemplified form of cancer treatable according to the methods herein includes e.g., malignant tumor, angiogenesis glaucoma, infantile hemangioma, multiple myeloma, acute

myeloblastic leukemia, chronic sarcoma, chronic myelogenous leukemia, metastasis melanoma, Kaposi's sarcoma, vascular proliferation, cachexia, colorectal cancer (e.g., familial colorectal cancer, hereditary nonpolyposis colorectal cancer, and gastrointestinal stromal tumor), lung cancer (e.g., non-small cell lung cancer, small cell lung cancer and malignant mesothelioma), mesothelioma, pancreatic cancer (e.g., pancreatic duct cancer), gastric cancer (e.g., papillary adenocarcinoma, mucinous adenocarcinoma and adenosquamous carcinoma), breast cancer (e.g., invasive ductal carcinoma, ductal carcinoma *in situ*, inflammatory breast cancer and metastatic breast cancer), ovarian cancer (e.g., ovarian epithelial carcinoma, extragonadal germ cell tumor, ovarian germ cell tumor, and ovarian low malignant potential tumor), hormone-dependent prostate cancer, non-hormone dependent prostate cancer, liver cancer (e.g., primary liver cancer and extrahepatic bile duct cancer), thyroid cancer (e.g., medullary thyroid carcinoma), kidney cancer (e.g., renal cell carcinoma, and transitional cell carcinoma in kidney and urinary duct), uterine cancer, endometrial cancer, brain tumor (e.g., pineal astrocytoma, pilocytic astrocytoma, diffuse astrocytoma and anaplastic astrocytoma), melanoma, sarcoma, urinary bladder cancer, hematologic cancer, hypophyseal adenoma, glioma, acoustic neurinoma, retinoblastoma, head and neck cancer, head and neck squamous cell carcinoma, pharyngeal cancer, laryngeal cancer, cancer of the tongue, thymoma, esophagus cancer, duodenal cancer, colorectal cancer, rectal cancer, hepatoma, pancreatic endocrine tumor, cancer of the bile duct, gallbladder cancer, penile cancer, urinary duct cancer, testis tumor, vulvar cancer, cervical cancer, endometrial cancer, uterus sarcoma, vaginal cancer, skin cancer, fungoid mycosis, basal cell tumor, soft tissue sarcoma, malignant lymphoma, Hodgkin's disease, myelodysplastic syndrome, acute lymphocytic leukemia, chronic lymphocytic leukemia, chronic myelocytic leukemia, malignant myeloma, adult T cell leukemia, chronic bone marrow proliferative disease, pancreatic endocrine tumor, fibrous histiocytoma, leiomyosarcoma, rhabdomyosarcoma, cancer of unknown primary, cancer-driven myelopoiesis, tumor growth, and metastasis.

[0079] Also included are diseases or disorders which are implicated by the regulation of the circadian rhythm of individuals and include, e.g., major depression, seasonal affective disorder, post-traumatic stress disorder (PTSD), bipolar disorder, autism, epilepsy, Alzheimer's disease and other central nervous system (CNS) disorders associated with altered sleep and/or circadian rhythms.

[0080] Also included in the diseases and disorders treatable by the compounds disclosed herein are diseases and disorders mediated by IL-17 expression, including STAT3-mediated IL-17 expression, in neutrophils, and include, e.g., corneal fungal infection; risk of corneal fungal infection; corneal ulcer; corneal ulcer resulting from fungal keratitis; corneal ulcer resulting from fungal infection; corneal fungal infection and related inflammation; keratitis; fungal keratitis; corneal inflammation; corneal disease; ocular disease; fungal-mediated corneal infection, corneal ulcer, corneal inflammation, or ocular ulcer, inflammation or infection; bacterial-mediated corneal infection, corneal ulcer, corneal inflammation, or ocular ulcer, inflammation or infection; microbial disease; bacterial infections; fungal infections; *Aspergillus* keratitis; *Fusarium* keratitis; cutaneous T cell lymphoma; lung inflammation; acute kidney ischemia-reperfusion injury; anthrax, including, cutaneous anthrax, inhalation anthrax, gastrointestinal

anthrax and injection anthrax; aspergillosis, including, pulmonary aspergillosis, chronic pulmonary aspergillosis (CPA), chronic aspergillosis, chronic cavitary pulmonary aspergillosis (CCPA), allergic bronchopulmonary aspergillosis (ABPA), allergic *Aspergillus* sinusitis, aspergilloma, invasive aspergillosis, chronic necrotizing aspergillosis and cutaneous (skin) aspergillosis; and histoplasmosis, including, systemic histoplasmosis. In particular embodiments, the fungus or fungal infection mediating the disease or disorder described above includes one or more of *Aspergillus*, *Fusarium*, *Alternaria*, *Candida*, *Curvularia* or *Histoplasma*.

[0081] Diseases and disorders mediated by IL-17 expression, and which are treatable using the compounds disclosed herein also include e.g., chronic graft-versus-host disease, acute graft-versus-host disease, celiac sprue, emphysema, lung fibrosis, giant cell arteritis, arteriosclerosis, hepatitis, chronic active hepatitis, alcoholic hepatitis, alcoholic liver fibrosis, alcoholic cirrhosis, viral hepatitis, hepatitis B viral liver disorder, autoimmune hepatitis, epidermal hyperplasia, cartilage inflammation, bone degradation, juvenile arthritis, juvenile rheumatoid arthritis, pauciarticular juvenile rheumatoid arthritis, polyarticular juvenile rheumatoid arthritis, systemic onset juvenile rheumatoid arthritis, spondyloarthritis, juvenile ankylosing spondylitis, juvenile enteropathic arthritis, juvenile reactive arthritis, juvenile Reiter's syndrome, seronegative enthesopathy and arthropathy (SEA) syndrome, juvenile dermatomyositis, juvenile psoriatic arthritis, juvenile scleroderma, juvenile systemic lupus erythematosus, juvenile vasculitis, pauciarticular rheumatoid arthritis, polyarticular rheumatoid arthritis, systemic onset rheumatoid arthritis, enteropathic arthritis, reactive arthritis, Reiter's syndrome, vasculitis, myositis, polymyositis, autoimmune myositis, osteoarthritis, polyarteritis nodosa, arteritis, polymyalgia rheumatica, sclerosis, primary biliary sclerosis, sclerosing cholangitis, enthesitis, enthesopathy, dermatitis, atopic eczema, eczema, atherosclerosis, Still's disease, Addison's disease, Raynaud's phenomenon, psoriatic epidermal hyperplasia, plaque psoriasis, guttate psoriasis, inverse psoriasis, pustular psoriasis, erythrodermic psoriasis, giant cell arteritis, noninfectious uveitis, Vogt-Koyanagi-Harada syndrome, mucosal leishmaniasis, Kawasaki disease, Hashimoto's thyroiditis, immune thrombocytopenic purpura (also known as immune thrombocytopenia, idiopathic immune thrombocytopenia, idiopathic thrombocytopenic thrombotic purpura, primary immune thrombocytopenia, idiopathic thrombocytopenic purpura, primary immune thrombocytopenic purpura, or autoimmune thrombocytopenic purpura), kidney inflammation, interstitial kidney inflammation, kidney disease, chronic kidney disease, renal failure, acute renal failure, end stage kidney disease, acute kidney injury, cisplatin induced acute renal failure, sepsis induced acute renal failure, nephritis, nephrotoxic nephritis, glomerulonephritis, acute glomerulonephritis, antineutrophil cytoplasmic autoantibody (ANCA) associated vasculitis, microscopic polyangiitis, granulomatosis with polyangiitis, Wegener's granulomatosis, amyotrophic lateral sclerosis, lupus nephritis, allergic eczema, transplant rejection, non-radiographic spondyloarthropathy, ophthalmic disorders, organ allograft rejection, fibroid lung, renal insufficiency, diabetic complications, diabetic nephropathy, diabetic retinopathy, diabetic retinitis, diabetic microangiopathy, insulinitis, tuberculosis, invasive staphylococci, invasive *Staphylococcus aureus* infection, inflammation after cataract surgery, allergic conjunctivitis, urticaria, chronic urticaria, allergic asthma, neutrophilic asthma, periodontal diseases, periodontitis, gingivitis, gum disease, diastolic cardiomyopathies, cardiac

infarction, myocarditis, chronic heart failure, angiostenosis, restenosis, reperfusion disorders, acute pancreatitis, chronic pancreatitis, asthma, bronchial asthma, acute respiratory distress syndrome, adult respiratory distress syndrome, inflammatory bone disease, inflammatory pulmonary disease, ischemic attack, transient ischemic attack, systemic inflammatory response syndrome, glaucoma, orbital cellulitis, sudden orbital inflammation, postoperative inflammation, posttraumatic inflammation, allergic inflammation, intestinal inflammation, mucosal inflammation, prostate inflammation, prostatitis, chronic pelvic pain syndrome, testicular inflammation, chronic testicular inflammation, orchitis, orchitis mediated infertility, liver disorder, liver injury, hepatotoxicity, pneumonia, meningitis, cystitis, pharyngolaryngitis, gastric mucosal injury, chronic pneumonia, pulmonary infarction, silicosis, pulmonary sarcoidosis, autoimmune anemia, autoimmune hemolytic anemia, Goodpasture's syndrome, sinusitis, chronic hypertrophic rhinitis, chronic inflammatory demyelinating polyneuropathy, mixed connective tissue disease, cognitive impairment, cognitive impairment in Alzheimer's disease, Parkinson's disease, spinal muscular atrophy, spinal cerebellar atrophy, progressive supranuclear palsy, Fisher syndrome, central nervous system lupus, encephalomyelitis, acute disseminated encephalomyelitis, multiple system atrophy, Huntington's disease, cerebrovascular dementia, diffuse Lewy body disease, cerebrovascular disorder, cerebral infarction, transient ischemic attack, intracerebral hemorrhage, vascular disease of the spinal cord, spinal cord infarction, Lambert-Eaton syndrome, muscular dystrophy, metabolic myopathy, inflammatory myopathy, inclusion body myositis, encephalitis, pemphigus, pemphigus vulgaris, profundus lupus erythematosus, chronic thyroiditis, autoimmune gastritis, sepsis, burn injury, pain, neuropathy, chronic pain, optic neuropathy, traumatic optic neuropathy, ischemic brain injury, deep venous thrombosis, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, abnormal immunoresponse, radiodermatitis, osteoporosis, parasitic infection, clonorchiasis, *Cryptosporidium* infection, *Streptococcus pneumoniae* carriage, chronic pneumococcal carriage, and an immune disorder associated with or arising from activity of pathogenic lymphocytes.

[0082] In one embodiment, a human patient is treated with a compound of the present invention and a pharmaceutically acceptable carrier, adjuvant, or vehicle, wherein said compound is present in an amount to treat or ameliorate one or more of the diseases and conditions recited above. In an alternative embodiment, the diseases and conditions treated or ameliorated by a compound of the present invention include, e.g., asthma, atopic dermatitis, acne, Crohn's disease, regional enteritis, ulcerative colitis, Sjögren's syndrome, uveitis, Behçet's disease, dermatomyositis, multiple sclerosis, ankylosing spondylitis, systemic lupus erythematosus (SLE), scleroderma, psoriasis, psoriatic arthritis (PsA), steroid resistant asthma and rheumatoid arthritis in the patient.

[0083] The present disclosure further relates to a combination therapy for treating or ameliorating a disease or a disorder described herein. In some embodiments, the combination therapy comprises administering at least one compound of the present invention in combination with one or more agents for treating or ameliorating inflammatory, metabolic and autoimmune diseases or disorders mediated by ROR γ . In some embodiments, the combination therapy comprises administering at least one compound of the present invention in

combination with one or more agents for the treatment of diseases including asthma, chronic obstructive pulmonary disease (COPD), bronchitis, allergic rhinitis, atopic dermatitis, contact dermatitis, acne, cystic fibrosis, allograft rejection, multiple sclerosis, scleroderma, arthritis, rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, systemic lupus erythematosus (SLE), psoriasis, Hashimoto's disease, pancreatitis, autoimmune diabetes, type I diabetes, autoimmune ocular disease, ulcerative colitis, Crohn's disease, regional enteritis, inflammatory bowel disease (IBD), inflammatory bowel syndrome (IBS), Sjögren's syndrome, optic neuritis, obesity, hepatosteatorrhea, adipose tissue-associated inflammation, insulin resistance, type II diabetes, neuromyelitis optica, myasthenia gravis, age related macular degeneration, dry eye, uveitis, Guillain-Barré syndrome, psoriasis, psoriatic arthritis (PsA), steroid resistant asthma, Graves' disease, scleritis, major depression, seasonal affective disorder, PTSD, bipolar disorder, autism, epilepsy, Alzheimer's, CNS disorders associated with altered sleep and/or circadian rhythms, endometriosis, obstructive sleep apnea syndrome (OSAS), Behçet's disease, dermatomyositis, polymyositis, graft versus host disease, primary biliary cirrhosis, liver fibrosis, non-alcoholic fatty liver disease (NAFLD), sarcoidosis, primary sclerosing cholangitis, autoimmune thyroid disease, autoimmune polyendocrine syndrome type I, autoimmune polyendocrine syndrome type II, celiac disease, neuromyelitis, juvenile idiopathic arthritis, systemic sclerosis, myocardial infarction, pulmonary hypertension, osteoarthritis, cutaneous leishmaniasis, sinonasal polyposis, and cancer, including but not limited to, lung cancer, gastric cancer, breast cancer and colon cancer.

[0084] The compounds disclosed herein may also be used in combination with immunotherapies for the treatment of a disease or disorder disclosed herein.

[0085] Combination therapy includes, e.g., co-administration of a compound disclosed herein and one or more other agents, sequential administration of a compound of the described and one or more other agents, administration of a composition containing a compound herein and one or more other agents, or simultaneous administration of separate compositions containing a compound described herein and one or more other agents.

[0086] The present disclosure further provides a method of treating a subject, such as a human, suffering from one of the abovementioned disorders or diseases.

[0087] In one aspect, ROR γ modulators disclosed in WO 2014/179564, WO 2015/116904, WO 2016/061160 and U.S. patent application publication Nos. U.S. 2016/0122318 and U.S. 2016/0122345 can also be used in the methods disclosed herein to treat or ameliorate, in a subject, one or more of the diseases and/or disorders and/or conditions recited herein. In one embodiment, a subject is treated with one or more ROR γ modulators disclosed in WO 2014/179564, WO 2015/116904, WO 2016/061160 and U.S. patent application publication Nos. U.S. 2016/0122318 and U.S. 2016/0122345 and a pharmaceutically acceptable carrier, adjuvant, or vehicle, wherein said ROR γ modulator is present in an amount to treat or ameliorate a disease or disorder selected from corneal fungal infection; risk of corneal fungal infection; corneal ulcer; corneal ulcer resulting from fungal keratitis; corneal ulcer resulting from fungal infection; corneal fungal infection and related inflammation; keratitis; fungal keratitis;

corneal inflammation; corneal disease; ocular disease; fungal-mediated corneal infection, corneal ulcer, corneal inflammation, or ocular ulcer, inflammation or infection; bacterial-mediated corneal infection, corneal ulcer, corneal inflammation, or ocular ulcer, inflammation or infection; microbial disease; bacterial infections; fungal infections; *Aspergillus* keratitis; *Fusarium* keratitis; cutaneous T cell lymphoma; lung inflammation; acute kidney ischemia-reperfusion injury; anthrax, including, cutaneous anthrax, inhalation anthrax, gastrointestinal anthrax and injection anthrax; aspergillosis, including, pulmonary aspergillosis, chronic pulmonary aspergillosis (CPA), chronic aspergillosis, chronic cavitary pulmonary aspergillosis (CCPA), allergic bronchopulmonary aspergillosis (ABPA), allergic *Aspergillus* sinusitis, aspergilloma, invasive aspergillosis, chronic necrotizing aspergillosis and cutaneous (skin) aspergillosis; histoplasmosis, including, systemic histoplasmosis; and prostate cancer.

[0088] In an alternative, a subject is treated with one or more ROR γ modulators disclosed in WO 2014/179564, WO 2015/116904, WO 2016/061160 and U.S. patent application publication Nos. U.S. 2016/0122318 and U.S. 2016/0122345 and a pharmaceutically acceptable carrier, adjuvant, or vehicle, wherein said ROR γ modulator is present in an amount to treat or ameliorate a disease or disorder selected from chronic graft-versus-host disease, acute graft-versus-host disease, celiac sprue, emphysema, lung fibrosis, giant cell arteritis, arteriosclerosis, hepatitis, chronic active hepatitis, alcoholic hepatitis, alcoholic liver fibrosis, alcoholic cirrhosis, viral hepatitis, hepatitis B viral liver disorder, autoimmune hepatitis, epidermal hyperplasia, cartilage inflammation, bone degradation, juvenile arthritis, juvenile rheumatoid arthritis, pauciarticular juvenile rheumatoid arthritis, polyarticular juvenile rheumatoid arthritis, systemic onset juvenile rheumatoid arthritis, spondyloarthritis, juvenile ankylosing spondylitis, juvenile enteropathic arthritis, juvenile reactive arthritis, juvenile Reiter's syndrome, seronegative enthesopathy and arthropathy (SEA) syndrome, juvenile dermatomyositis, juvenile psoriatic arthritis, juvenile scleroderma, juvenile systemic lupus erythematosus, juvenile vasculitis, pauciarticular rheumatoid arthritis, polyarticular rheumatoid arthritis, systemic onset rheumatoid arthritis, enteropathic arthritis, reactive arthritis, Reiter's syndrome, vasculitis, myositis, polymyositis, autoimmune myositis, osteoarthritis, polyarteritis nodosa, arteritis, polymyalgia rheumatica, sclerosis, primary biliary sclerosis, sclerosing cholangitis, enthesitis, enthesopathy, dermatitis, atopic eczema, eczema, atherosclerosis, Still's disease, Addison's disease, Raynaud's phenomenon, psoriatic epidermal hyperplasia, plaque psoriasis, guttate psoriasis, inverse psoriasis, pustular psoriasis, erythrodermic psoriasis, giant cell arteritis, noninfectious uveitis, Vogt-Koyanagi-Harada syndrome, mucosal leishmaniasis, Kawasaki disease, Hashimoto's thyroiditis, immune thrombocytopenic purpura (also known as immune thrombocytopenia, idiopathic immune thrombocytopenia, idiopathic thrombocytopenic thrombotic purpura, primary immune thrombocytopenia, idiopathic thrombocytopenic purpura, primary immune thrombocytopenic purpura, or autoimmune thrombocytopenic purpura), kidney inflammation, interstitial kidney inflammation, kidney disease, chronic kidney disease, renal failure, acute renal failure, end stage kidney disease, acute kidney injury, cisplatin induced acute renal failure, sepsis induced acute renal failure, nephritis, nephrotoxic nephritis, glomerulonephritis, acute glomerulonephritis, antineutrophil cytoplasmic autoantibody (ANCA) associated vasculitis, microscopic polyangiitis, granulomatosis with polyangiitis, Wegener's granulomatosis, amyotrophic lateral sclerosis, lupus nephritis, allergic eczema, transplant

rejection, non-radiographic spondyloarthropathy, ophthalmic disorders, organ allograft rejection, fibroid lung, renal insufficiency, diabetic complications, diabetic nephropathy, diabetic retinopathy, diabetic retinitis, diabetic microangiopathy, insulinitis, tuberculosis, invasive staphylococci, invasive *Staphylococcus aureus* infection, inflammation after cataract surgery, allergic conjunctivitis, urticaria, chronic urticaria, allergic asthma, neutrophilic asthma, periodontal diseases, periodontitis, gingivitis, gum disease, diastolic cardiomyopathies, cardiac infarction, myocarditis, chronic heart failure, angiostenosis, restenosis, reperfusion disorders, acute pancreatitis, chronic pancreatitis, asthma, bronchial asthma, acute respiratory distress syndrome, adult respiratory distress syndrome, inflammatory bone disease, inflammatory pulmonary disease, ischemic attack, transient ischemic attack, systemic inflammatory response syndrome, glaucoma, orbital cellulitis, sudden orbital inflammation, postoperative inflammation, posttraumatic inflammation, allergic inflammation, intestinal inflammation, mucosal inflammation, prostate inflammation, prostatitis, chronic pelvic pain syndrome, testicular inflammation, chronic testicular inflammation, orchitis, orchitis mediated infertility, liver disorder, liver injury, hepatotoxicity, pneumonia, meningitis, cystitis, pharyngolaryngitis, gastric mucosal injury, chronic pneumonia, pulmonary infarction, silicosis, pulmonary sarcoidosis, autoimmune anemia, autoimmune hemolytic anemia, Goodpasture's syndrome, sinusitis, chronic hypertrophic rhinitis, chronic inflammatory demyelinating polyneuropathy, mixed connective tissue disease, cognitive impairment, cognitive impairment in Alzheimer's disease, Parkinson's disease, spinal muscular atrophy, spinal cerebellar atrophy, progressive supranuclear palsy, Fisher syndrome, central nervous system lupus, encephalomyelitis, acute disseminated encephalomyelitis, multiple system atrophy, Huntington's disease, cerebrovascular dementia, diffuse Lewy body disease, cerebrovascular disorder, cerebral infarction, transient ischemic attack, intracerebral hemorrhage, vascular disease of the spinal cord, spinal cord infarction, Lambert-Eaton syndrome, muscular dystrophy, metabolic myopathy, inflammatory myopathy, inclusion body myositis, encephalitis, pemphigus, pemphigus vulgaris, profundus lupus erythematosus, chronic thyroiditis, autoimmune gastritis, sepsis, burn injury, pain, neuropathy, chronic pain, optic neuropathy, traumatic optic neuropathy, ischemic brain injury, deep venous thrombosis, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, abnormal immunoresponse, radiodermatitis, osteoporosis, parasitic infection, clonorchiasis, *Cryptosporidium* infection, *Streptococcus pneumoniae* carriage, chronic pneumococcal carriage, and an immune disorder associated with or arising from activity of pathogenic lymphocytes.

[0089] In some embodiments, the one or more ROR γ modulator disclosed in WO 2014/179564, WO 2015/116904, WO 2016/061160 and U.S. patent application publication Nos. U.S. 2016/0122318 and U.S. 2016/0122345 is administered in combination with one or more additional agent for treating the disease or disorder.

[0090] The present disclosure further relates to the use of provided compounds for the production of pharmaceutical compositions which are employed for the treatment and/or prophylaxis and/or amelioration of the diseases and disorders mentioned herein.

[0091] Compounds or compositions disclosed herein may be administered using any amount

and any route of administration effective for treating or lessening the severity of one or more of the diseases and conditions described herein. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like. Provided compounds are preferably formulated in unit dosage form for ease of administration and uniformity of dosage. The expression "unit dosage form" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present disclosure will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts.

[0092] Pharmaceutically acceptable compositions of this disclosure can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, provided compounds may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg and preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[0093] The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof, biopsied material obtained from a mammal or extracts thereof, and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

[0094] The amount of both, a provided compound and additional therapeutic agent (in those compositions which comprise an additional therapeutic agent as described above) that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

[0095] In those compositions which comprise an additional therapeutic agent, that additional therapeutic agent and the provided compound may act synergistically. Therefore, the amount of additional therapeutic agent in such compositions will be less than that required in a monotherapy utilizing only that therapeutic agent.

[0096] The amount of additional therapeutic agent present in the compositions of this disclosure will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent.

EXEMPLIFICATION

[0097] As depicted in the Examples below, in certain exemplary embodiments, compounds are prepared according to the following general procedures. It will be appreciated that, although the general methods depict the synthesis of certain compounds herein, the following general methods, and other methods known to one of ordinary skill in the art, can be applied to all compounds and subclasses and species of each of these compounds, as described herein.

GENERAL DESCRIPTION OF SYNTHESIS

[0098] The compounds disclosed herein can be readily prepared according to the following reaction schemes and examples, or modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures. Many of the reactions can also be carried out under microwave (MW) conditions or using conventional heating or utilizing other technologies such as solid phase reagents/scavengers or flow chemistry. In these reactions, it is also possible to make use of variants which are themselves known to those of ordinary skill in the art, but are not mentioned in greater detail. Furthermore, other methods for preparing compounds disclosed herein will be readily apparent to a person of ordinary skill in the art in light of the following reaction schemes and examples. In cases where synthetic intermediates and final products contain potentially reactive functional groups, for example amino, hydroxy, thiol and carboxylic acid groups, that may interfere with the desired reaction, it may be advantageous to employ protected forms of the intermediate. Methods for the selection, introduction and subsequent removal of protecting groups are well known to those skilled in the art. In the discussion below variables have the meanings indicated above unless otherwise indicated. The abbreviations used in these experimental details are listed below and additional ones should be known to a person skilled in the art of synthesis. In addition, one can refer to the following references for suitable methods of synthesis as described in March, *Advanced Organic Chemistry*, 3rd edition, John Wiley & Sons, 1985, Greene and Wuts, *Protective Groups in Organic Synthesis*, 2nd edition, John Wiley & Sons, 1991, and Richard Larock, *Comprehensive Organic Transformations*, 4th edition, VCH publishers Inc., 1989.

[0099] Generally, reagents in the reaction schemes are used in equimolar amounts; however, in certain cases it may be desirable to use an excess of one reagent to drive a reaction to completion. This is especially the case when the excess reagent can be readily removed by evaporation or extraction. Bases employed to neutralize HCl in reaction mixtures are generally used in slight to substantial excess (1.05 - 5 equivalents).

[0100] Where NMR data are presented, spectra were obtained on a Varian 400 (400 MHz) or 300 (300 MHz) and are reported as ppm downfield from tetramethylsilane with number of proton, multiplicities and coupling constants indicated parenthetically along with reference to deuterated solvent.

[0101] The invention is illustrated by way of the following examples, in which the following abbreviations may be employed.

Abbreviation	Meaning
ACN, MeCN, CH ₃ CN	acetonitrile
AIBN	azobisisobutyronitrile
aq	aqueous
Boc	<i>tert</i> -butoxycarbonyl or <i>t</i> -butoxycarbonyl
brine	saturated aqueous NaCl
c-Bu	cyclobutyl
Cbz	benzyloxy carbonyl
CeCl ₃	ceric chloride
Cs ₂ CO ₃	cesium carbonate
CuI	cuprous iodide
c-Pr	cyclopropyl
DCM or CH ₂ Cl ₂	methylene chloride
DIEA	diisopropyl ethyl amine
DMF	dimethyl formamide
DMS/Me ₂ S	dimethyl sulfide
DMSO	dimethyl sulfoxide
EDCI	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
EtI	ethyl iodide
Et	ethyl
Et ₂ O	ethyl ether
Et ₃ SiH	triethylsilane
Et ₃ N	triethylamine
EtOAc , EA, AcOEt	ethyl acetate
EtOH	ethanol
FeCl ₃	ferric chloride
h, hr	hour(s)
HATU	O-(7-azabenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium-hexafluorophosphate
HBTU	O-benzotriazole-1-yl- <i>N,N,N',N'</i> -tetramethyluronium-hexafluorophosphate
HCl	hydrochloric acid

Abbreviation	Meaning
H ₂ O	water
H ₂ O ₂	hydrogen peroxide
HPLC	high performance liquid chromatography
i-BuOCOCi	iso-butoxycarbonyl chloride
ICI	iodochloride
K ₂ CO ₃	potassium carbonate
K ₃ PO ₄	tripotassium phosphate
LC-MS	liquid chromatography-mass spectrometry
LDA	lithium diisopropylamide
LiCl	lithium chloride
LiOH	lithium hydroxide
MCPBA, <i>m</i> -CPBA	meta-chloroperoxybenzoic acid
MeOH	methanol
MeI	methyl iodide
Me	methyl
mg	milligram
MgSO ₄	magnesium sulfate (anhydrous)
min	minute(s)
mL	milliliters
mmol	millimoles
mp, m.p.	melting point
MS	mass spectrometry
MW, uwave	microwave
NaBH ₄	sodium borohydride
NaBH ₃ CN	sodium cyanoborohydride
NaH	sodium hydride
NaHCO ₃	sodium bicarbonate
NaOH	sodium hydroxide
NaOMe	sodium methoxide
Na ₂ S ₂ O ₃	sodium thiosulfate
Na ₂ S ₂ O ₅	sodium dithionate
Na ₂ SO ₄	sodium sulfate
NH ₄ OH	ammonium hydroxide

Abbreviation	Meaning
$(\text{NH}_4)_2\text{CO}_3$	ammonium carbonate
NH_4Cl	ammonium chloride
Na_2CO_3	sodium carbonate
NaHCO_3	sodium bicarbonate
NaH	sodium hydride
NBS	<i>N</i> -bromosuccinimide
<i>n</i> -BuLi	<i>n</i> -butyllithium
NMM	<i>N</i> -methyl-morpholine
NMP	<i>N</i> -methyl-pyrrolidin-2-one
OTf	trifluoromethanesulfonate
OTs	tosylate
PdCl_2dppf	[1,1-bis(diphenylphosphino)ferrocene] dichloropalladium(ii)
$\text{Pd}_2(\text{dba})_3$	tris(dibenzylideneacetone)dipalladium(0)
PE	petroleum ether
rt	room temperature
sat.	saturated
SFC	supercritical fluid chromatography
<i>t</i> -BuOK	potassium tert butoxide
<i>t</i> -BuLi	tert butyl lithium
<i>t</i> -BuOOH	tert butyl peroxide
TBAF	tetrabutylammonium fluoride
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TLC	thin layer chromatography
$\text{Ti}(\text{OEt})_4$	titanium tetra ethoxide
Zn	zinc
$\text{Zn}(\text{CN})_2$	zinc cyanide

LCMS Methods

Method 1

[0102]

HPLC System:	Waters ACQUITY	
Column:	Waters ACQUITY CSH™ C18 1.7 uM	
Guard column:	Waters Assy. Frit, 0.2uM, 2.1 mm.	
Column tem:	40 °C	
Mobile Phase:	A: TFA : Water (1 : 1000, v:v)	
	B: TFA : ACN (1: 1000, v:v)	

Gradient Program:

[0103]

Time (min)	B%
0	5
1.9	95
2.20	95
2.21	5
Flow Rate:	0.65 mL/min

Mass Spectrometer Parameters

[0104]

Mass Spectrometer	Waters SQD	
Ionization	Positive Electrospray Ionization (ESI)	
Mode	Scan (100-1400 m/z in every 0.2 second)	
ES Capillary Voltage:	3.5 kv	
ES Cone Voltage:	25 v	
Source Temperature	120 °C	
Disolvation Temperature:	500 °C	
Desolvation Gas Flow:	Nitrogen	Setting 650 (L/hr)
Cone Gas Flow:	Nitrogen	Setting 50 (L/hr)

Method 2

[0105]


5-95AB 1.5MIN		
Column	MERCK,RP-18e 25-2mm	
Mobile Phase	A:water(4L)+TFA(1.5mL)	
	B:acetonitrile(4L)+TFA(0.75mL)	
	TIME(min)	B%
	0	5
	0.7	95
	1.1	95
	1.11	5
	1.5	5
Flow Rate	1.5mL/min	
wavelength	UV 220nm	
Oven Temp	50°C	
MS ionization	ESI	

Method 3


[0106]

10-80CD_3MIN		
Column	Xbrige Shield RP-18,5um,2.1 *50mm	
Mobile Phase	A:water(1L)+NH3H2O(0.5mL)	
	B: acetonitrile	
	TIME(min)	B%
	0	10
	2	80
	2.48	80
	2.49	10
	3	10
Flow Rate	1.0mL/min	
wavelength	UV 220nm&254nm	
Oven Temp	30°C	
MS ionization	ESI	


[0110] In a third process, a compound of Formula I, wherein $X = \text{NHCO}$, is prepared from an amine of Formula 115 and a carboxylic acid of Formula 120, using peptide bond formation reagents such as EDC, HOBt or HATU. Alternatively, the carboxylic acid 120 may be converted to its acid chloride prior to reaction.




[0111] In a fourth process, an amine of Formula 115 is prepared from a compound of Formula 110 using NaN_3 in the presence of CuI or benzophenone imine in the presence of a palladium catalyst.



[0112] In a fifth process, a compound of Formula I is prepared from another compound of Formula I. For example a compound of Formula I, wherein L^2 is CH_2 or CHOH , is oxidized to a compound of Formula I, wherein L^2 is C=O , using, for example MnO_2 .




[0113] In an alternative, a compound of Formula I, wherein $\text{L}^2 = \text{CO}$, is reduced to a compound of Formula I, wherein $\text{L}^2 = \text{CHOH}$, using, for example NaBH_4 .




I (L = CO)

I (L = CHOH)


[0114] In an alternative, a compound of Formula I, wherein R^7 (or R^8) is CH_2OH is derivatized to provide a compound of Formula I, wherein R^7 (or R^8) is $CH_2OC(=O)NH_2$ by the action of $Cl_3CC(=O)NCO$, followed by K_2CO_3 in MeOH.



[0115] In another alternative, a compound of Formula I, wherein R^6 (or R^5) is $CONH_2$, is treated with trifluoroacetic anhydride in the presence of a base such as Et_3N or pyridine, to provide a compound of Formula I wherein R^6 (or R^5) is CN .




[0116] In a sixth process, a compound of Formula I, wherein A is N, is prepared from a compound of Formula 125 by treatment with acid, for example heating in HOAc at 100 °C.




[0117] In a seventh process, a compound of Formula 125 is prepared by S_NAr reaction, or Buchwald-Hartwig coupling, of an *ortho* halo nitro compound of Formula 7, wherein Hal = F, Cl, Br or I, with an amine of Formula 135 to give a compound of Formula 140. Reduction of 140, using for example zinc dust in the presence of NH_4Cl , water and THF, gives a diamine of Formula 145. Acylation of 145 with acids of Formula 150, mediated by peptide coupling


reagents such as HATU, gives compounds of Formula **125**. Alternatively, the acid chloride of **150** can be employed in the final step.




[0118] In an eighth process, a carboxylic acid of Formula **100** is prepared as follows. A compound of Formula **160** is prepared by $\text{S}_{\text{N}}\text{Ar}$ reaction, or Buchwald-Hartwig coupling, of an *ortho* halo nitro compound of Formula **155**, wherein Hal is F, Cl, Br or I and R^{101} is lower alkyl, with an amine of Formula **135** to give an *ortho* amino nitro compound of Formula **160**. Reduction of **160**, using for example zinc dust in the presence of NH_4Cl or hydrogen in the presence of palladium on carbon, gives a diamine of Formula **165**. Acylation of **165** with acids of Formula **150**, mediated by peptide coupling reagents such as HATU, gives amide compounds of Formula **170**. Alternatively, the acid chloride of **150** can be employed in this step. Treatment of **170** with acids, for example HOAc at 100°C or TFA at room temperature affords compounds of Formula **175**. Finally the ester moiety in **175** is hydrolyzed with MOH , wherein M = an alkali metal, to afford carboxylic acids of Formula **100**.




[0119] In a ninth process, an amine of Formula **115**, wherein A is N, is prepared by the following sequence. A dihalonitro compound of Formula **180**, wherein Hal¹ is fluoro or chloro and Hal² is bromo or iodo, is reacted with an amine of Formula **135** under S_NAr reaction conditions, for example i-Pr₂NEt in MeCN, to give a compound of Formula **185**. The nitro group in **185** is selectively reduced using for example SnCl₂ in EtOH or DMF, to afford a diamine of Formula **190**. Acylation of **190** with acids of Formula **150**, mediated by peptide coupling reagents such as HATU, gives amide compounds of Formula **195**. Alternatively, the acid chloride of **150** can be employed in this step. Treatment of **195** with acids, for example HOAc at 100 °C or TFA at room temperature, affords compounds of Formula **200**. Finally, Hal² in **200** is converted to an amino group using, for example, NaN₃ in the presence of CuI or benzophenone imine in the presence of a palladium source and a suitable ligand to give a compound of Formula **115**.




[0120] In a tenth process, a carboxylic acid of Formula **100**, wherein A is CH and R¹⁰¹ is lower alkyl, is prepared as follows. A halo benzene or pyridine of Formula **205**, wherein Hal² is bromine or iodine, is coupled to a terminal alkyne of Formula **210** under Sonogashira conditions e.g. PdCl₂(PPh₃)₂, CuI, Et₃N, to give a compound of Formula **215**. Treatment of **215** with a non nucleophilic base such as NaH effects ring closure to afford an indole or azaindole of Formula **220**. Finally, the ester moiety in **220** is hydrolyzed with MOH, wherein M = an alkali metal, to afford a carboxylic acid of Formula **100**.






[0121] In an eleventh process, a carboxylic acid of Formula **100**, wherein A and Y are both CH, is prepared as follows. A protected indole of Formula **225**, wherein PG is a suitable protecting group such as p-toluenesulfonyl and Hal² is bromine or iodine, is treated with a strong base such as LDA and an aldehyde of Formula **230** to give a secondary alcohol of Formula **235**. Reduction of the alcohol moiety of **235** with, for example Et₃SiH, provides a compound of Formula **240**. Carbonylation of **240** with carbon monoxide in the presence of an alcohol of Formula **245**, wherein R¹⁰¹ is lower alkyl, a palladium catalyst such as PdCl₂(dppf) or Pd(OAc)₂/dcpp, affords an ester of Formula **250**. Removal of the protecting group from **250** gives a compound of Formula **255** which is reacted with alkylating agent **260**, wherein R¹⁰⁰ is iodine, bromine, or trifluoromethanesulfonyloxy to yield a compound of Formula **265**. Finally the ester moiety in **265** is hydrolyzed with MOH, wherein M = an alkali metal, to afford a carboxylic acid of Formula **100**, wherein A and Y are both CH.



[0122] In a twelfth process, an amine of Formula **115** in which both A and Y are CH is prepared as follows. A suitably protected indole of Formula **240** is deprotected to give a compound of Formula **270**. Reaction of **270** with an alkylating agent of Formula **260**, wherein R¹⁰⁰ is iodine, bromine, or trifluoromethanesulfonyloxy, in the presence of a base affords a compound of Formula **275**. Finally, Hal² in **275** is converted to an amino group using, for example, NaN₃ in the presence of CuI or benzophenone imine in the presence of a palladium

source and a suitable ligand to give a compound of Formula **115**.




PREPARATION OF INTERMEDIATES

Preparation 1

4-(ethylsulfonyl)phenyl)methanamine

[0123]



Step 1

[0124] To a mixture of 4-methylbenzenethiol (100 g, 0.8 mol) in acetone (1 L) was added iodoethane (190 g, 1.2 mol) and potassium carbonate (220 g, 1.6 mol). The mixture was stirred at 60 °C overnight. The mixture was filtered and the filtrate was concentrated under reduced pressure to afford crude ethyl(*p*-tolyl)sulfane (120 g, 99%) as a yellow solid, which was used for the next step without further purification.

Step 2

[0125] To a solution of crude ethyl(*p*-tolyl)sulfane (35 g, 0.23 mol) in CH₂Cl₂ (1.5 L) was

added *m*-chloroperoxybenzoic acid (101 g, 0.59 mol) at 0 °C. The mixture was stirred at rt overnight. The mixture was filtered. The filtrate was added to saturated aqueous Na₂SO₃ (500 mL) slowly and then stirred for 0.5 h. After partitioning, the organic layer was washed with saturated aqueous NaHCO₃ (500 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude 1-(ethylsulfonyl)-4-methylbenzene (42.3 g, 100%) as a pale yellow solid. This material was used for the next step without further purification.

Step 3

[0126] To a solution of 1-(ethylsulfonyl)-4-methylbenzene (5 g, 25.7 mmol) in CCl₄ (30 mL) was added *N*-bromosuccinimide (5.54 g, 30.8 mmol) and azobisisobutyronitrile (0.46 g, 2.57 mmol). The mixture was stirred at 80 °C overnight. The mixture was filtered and the filtrate was concentrated under reduced pressure. The residue was added to water (50 mL) and extracted with EtOAc (3 x 30 mL). The combined organic layers were washed with water (2 x 40 mL) then brine (50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude 1-(bromomethyl)-4-(ethylsulfonyl)benzene (6.62 g, 98%) as a yellow solid. This material was used for the next step without further purification.


Step 4


[0127] To a solution of 1-(bromomethyl)-4-(ethylsulfonyl)benzene (6.62 g, 25.2 mmol) in MeOH (30 mL) was added a 28% aqueous ammonium hydroxide solution (30 mL). The mixture was stirred at rt overnight. The mixture was then concentrated under reduced pressure. The residue was purified by basic preparative HPLC method B separation to afford (4-(ethylsulfonyl)phenyl) methanamine (1.5 g, 30%) as a yellow solid. LC-MS t_R = 1.747 min in 0-30CD_3 min chromatography (Durashell C18, 2.1*30 mm, 3 μm), MS (ESI) m/z 200.0 [M + H]⁺ and 399.0 [2M + H]⁺. ¹H NMR (CDCl₃, 400 MHz): δ 7.85 (d, J = 8.0 Hz, 2H), 7.53 (d, J = 8.0 Hz, 2H), 3.98 (s, 2H), 3.10 (q, J = 7.6 Hz, 2H), 1.26 (t, J = 7.6 Hz, 3H).

Preparation 2

(*R*)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol (3:2 mixture of enantiomers, *R*:*S*)

[0128]





[0129] To a suspension of NaH (12.9 g, 0.322 mol, 60% in mineral oil) in dry THF (500 mL) was added a solution of ethane-1,2-diol (20.0 g, 0.322 mol) in dry THF (100 mL) dropwise. The mixture was stirred for 1 h at rt, then TBSCl (48.59 g, 0.322 mol) was added and the mixture was stirred for another 1 h at rt. The mixture was quenched with an aqueous K_2CO_3 solution (100 mL, 10%) and extracted with MTBE (3 x 300 mL). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography on silica gel (eluting with a gradient of 1% to 33% ethyl acetate in petroleum ether) to afford 2-((*tert*-butyldimethylsilyl)oxy)ethanol (55.0 g, 96%) as a colorless oil.

[0130] To a solution of $(COCl)_2$ (13.7 mL, 162.2 mmol) in anhydrous CH_2Cl_2 (400 mL) was added DMSO (25.1 mL, 353.9 mmol) at $-78\text{ }^\circ\text{C}$ under N_2 . After being stirred for 30 min, a solution of 2-((*tert*-butyldimethylsilyl)oxy)ethanol (26.0 g, 147.5 mmol) in anhydrous CH_2Cl_2 (100 mL) was added dropwise. The mixture was stirred at $-78\text{ }^\circ\text{C}$ for 30 min and then Et_3N (102.74 mL, 737.0 mmol) was added dropwise. After being stirred at $-78\text{ }^\circ\text{C}$ for 30 min, the reaction was allowed to warm to rt and stirred for 1 h. The reaction mixture was acidified with 2 N aqueous HCl solution to pH=4 and then extracted with CH_2Cl_2 (3 x 400 mL). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure to afford 2-((*tert*-butyldimethylsilyl)oxy)acetaldehyde (34.0 g, 100%) as a colorless oil. This material was used for the next step directly without further purification. To a solution of 2-((*tert*-butyldimethylsilyl)oxy)acetaldehyde (34.0 g, 147.5 mmol) and (*R*)-2-methylpropane-2-sulfonamide (19.6 g, 162.2 mmol) in DCM (400 mL) was added $CuSO_4$ (47.2 g, 295.0 mmol) at rt. The mixture was stirred at $25\text{ }^\circ\text{C}$ for 48 h. The mixture was filtered and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (eluting with a gradient of 1% to 10% ethyl acetate in petroleum ether) to afford (*R,E*)-*N*-(2-((*tert*-butyldimethylsilyl)oxy)ethylidene)-2-methylpropane-2-sulfonamide (30.0 g, 73%) as a yellow oil. $^1H\text{ NMR}$ ($CDCl_3$ 400 MHz): δ 7.96 (t, $J = 3.2\text{ Hz}$, 1H), 4.45 (d, $J = 3.2\text{ Hz}$, 2H), 1.11 (s, 9H), 0.82 (s, 9H), 0.00 (s, 6H).

[0131] To a solution of (4-bromophenyl)(ethyl)sulfane (1.88 g, 8.65 mmol) in anhydrous THF (15 mL) was added *n*-BuLi (14.4 mL, 36.0 mmol, 2.5 M in THF) at $-78\text{ }^\circ\text{C}$ under N_2 . The

reaction mixture was stirred at -78 °C for 0.5 h, then a solution of (*R,E*)-*N*-(2-((*tert*-butyldimethylsilyl)oxy)ethylidene)-2-methylpropane-2-sulfinamide (2.0 g, 7.2 mmol) in THF (1 mL) was added dropwise to the mixture at -78 °C. The reaction mixture was stirred at -78 °C for 0.5 h and was allowed to warm to rt. After 0.5 h at rt, the reaction mixture was quenched with water (10 mL) and extracted with ethyl acetate (3 x 20 mL). The combined organic layers were washed with H₂O (10 mL) then brine (10 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure.

[0132] The crude residue was purified by column chromatography on silica gel (eluting with a gradient of 5% to 10% ethyl acetate in petroleum ether) to afford a 3:2 mixture of (*R*)-*N*-((*R*)-2-((*tert*-butyldimethylsilyl)oxy)-1-(4-(ethylthio)phenyl)ethyl)-2-methylpropane-2-sulfinamide and (*R*)-*N*-((*S*)-2-((*tert*-butyldimethylsilyl)oxy)-1-(4-(ethylthio)phenyl)ethyl)-2-methylpropane-2-sulfinamide (2.86 g, 47.9 %) as a yellow oil. This 3:2 mixture of diastereomers was used for the subsequent reactions. ¹H NMR (CDCl₃ 400 MHz): δ 7.24-7.16 (m, 4H), 4.47-4.40 (m, 1H), 4.21 (s, 1H), 3.75-3.67 (m, 1H), 3.60-3.47 (m, 1H), 2.90 (q, *J* = 7.2 Hz, 2H), 1.27 (t, *J* = 7.2 Hz, 3H), 1.18 (s, 9H), 0.85 (s, 9H), 0.02 (s, 3H), 0.00 (s, 3H).

[0133] To a solution of (*R*)-*N*-((*R*)-2-((*tert*-butyldimethylsilyl)oxy)-1-(4-(ethylthio)phenyl)ethyl)-2-methylpropane-2-sulfinamide (2.86 g, 6.86 mmol), as a 3:2 mixture of diastereomers from previous step, in dioxane (10 mL) was added HCl in dioxane (20 mL, 4 M) at rt. The mixture was stirred for 3 h at rt and was then concentrated under reduced pressure to afford crude (*R*)-2-amino-2-(4-(ethylthio)phenyl)ethan-1-ol HCl salt (3.0 g, 100%) as a white solid. This crude material was a 3:2 mixture of enantiomers (*R*:*S*) and used for the next step without further purification.


[0134] To a solution of (*R*)-2-amino-2-(4-(ethylthio)phenyl)ethan-1-ol HCl salt (3.0 g, 6.86 mmol), as a 3:2 mixture of enantiomers / *R*:*S*, in H₂O (30 mL) was added Oxone® monopersulfate (8.4 g, 13.72 mmol) at rt. The mixture was stirred for 1.5 h at rt and was lyophilized directly. After lyophilization, the crude product was purified by flash column chromatography on silica gel (eluting with MeOH) to give crude (*R*)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol HCl salt (3:2 mixture of enantiomers, *R*:*S*) (1.25 g, 80%) as a yellow solid. ¹H NMR (CDCl₃ 400 MHz): δ 7.99 (d, *J* = 8.4 Hz, 2H), 7.73 (d, *J* = 8.0 Hz, 2H), 4.55-4.50 (m, 1H), 3.99-3.91 (m, 1H), 3.85-3.80 (m, 1H), 3.23 (q, *J* = 7.2 Hz, 2H), 1.21 (t, *J* = 7.2 Hz, 3H).

Preparation 3

(5-(ethylsulfonyl)pyridin-2-yl)methanamine

[0135]

EtSO₃Na, L-proline,




[0136] To a flame dried flask equipped with a stir bar was added *tert*-butyl ((5-bromopyridin-2-yl)methyl)carbamate (2.92 g, 10.2 mmol), ethane sulfinic acid sodium salt (2.36 g, 20.3 mmol), *L*-proline (234 mg, 2.03 mmol), copper (I) iodide (194 mg, 1.02 mmol) and sodium hydroxide (81.3 mg, 2.03 mmol). The flask was purged with N_2 , then DMSO (35 mL) was added. The reaction mixture was heated to 110°C and stirred for 15 h. The flask was then cooled to rt and the mixture was partitioned between EtOAc (150 mL) and saturated aqueous ammonium chloride (150 mL). The organic phase was separated, washed with brine (50 mL), dried over anhydrous MgSO_4 , filtered and concentrated under reduced pressure. The residue was purified by silica gel chromatography (eluting with 35% EtOAc in hexanes, gradient to 60%) to afford *tert*-butyl((5-bromopyridin-2-yl)methyl)carbamate (1.81 g, 59%). **LC-MS** $t_R = 0.74$ min in 1 min chromatography, **MS** (ESI) m/z 301.4 $[\text{M} + \text{H}]^+$. **^1H NMR** (CDCl_3 , 400 MHz): δ 9.02 (dd, $J = 0.8$ Hz, 2.0 Hz, 1H), 8.15 (dd, $J = 2.4$ Hz, 8.4 Hz, 1H), 7.49 (dd, $J = 0.8$ Hz, 8.4 Hz, 1H), 5.49 (broad s, 1H), 4.55 (d, $J = 7.0$ Hz, 2H), 3.15 (q, $J = 7.2$ Hz, 2H), 1.47 (s, 9H), 1.31 (t, $J = 7.2$ Hz, 3H).


[0137] To a solution of *tert*-butyl ((5-bromopyridin-2-yl)methyl)carbamate (1.81 g, 6.03 mmol) in MeOH (40 mL) at 0°C was added acetyl chloride (4.30 mL, 60.3 mmol) dropwise over 5 min. The solution was allowed to warm to rt and was stirred for 3 h. The mixture was concentrated under reduced pressure to yield 1.64 g (5-(ethylsulfonyl)pyridin-2-yl)methanamine bis-hydrochloride salt ($\sim 100\%$). This material was used directly for the next step without purification. **LC-MS** $t_R = 0.25$ min in 1 min chromatography, **MS** (ESI) m/z 201.2 $[\text{M} + \text{H}]^+$. **^1H NMR** (CD_3OD , 400 MHz): δ 9.09 (d, $J = 1.2$ Hz, 1H), 8.35 (dd, $J = 2.4$ Hz, 8.4 Hz, 1H), 7.49 (d, $J = 8.4$ Hz, 1H), 4.45 (s, 2H), 3.31 (q, $J = 7.2$ Hz, 2H), 1.26 (t, $J = 7.2$ Hz, 3H).

Preparation 4

2-(6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)acetic acid

[0138]






PREPARATION OF COMPOUNDS OF FORMULA I

[0139] Compounds of Formula I were prepared according to the general procedures outlined below.

Example 1

1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxamide (Compound IN-1)

[0140]



Step 1

[0141] To a solution of 5-bromo-1H-indole (5.0 g, 25.5 mmol) in anhydrous DMF (100 mL) was added NaH (1.2 g, 30.6 mmol, 60 % in mineral oil) under N₂. The reaction mixture was stirred at 0 °C for 1 h. 4-methylbenzene-1-sulfonyl chloride (5.8 g, 30.6 mmol) in anhydrous DMF (25 mL) was added dropwise at 0 °C and the reaction mixture was stirred at room temperature for 4 h. TLC (petroleum ether / EtOAc = 5/1) showed the reaction was completed. Water (50 mL) was added at 0 °C. Most solid was precipitated out. Then the solid was filtered and washed with water. The solid was collected and dried under reduced pressure to afford 5-bromo-1-tosyl-1H-indole (8.2 g, 92% yield) as a white solid. ¹H NMR (CDCl₃ 400 MHz): δ 7.86 (d, *J* = 8.4 Hz, 1H), 7.74 (d, *J* = 8.4 Hz, 2H), 7.66 (d, *J* = 1.6 Hz, 1H), 7.56 (d, *J* = 7.6 Hz, 1H), 7.38 (d, *J* = 8.4 Hz, 1H), 7.23 (d, *J* = 8.0 Hz, 2H), 6.59 (d, *J* = 3.2 Hz, 1H), 2.35 (s, 3H).

Step 2

[0142] To a solution of 5-bromo-1-tosyl-1H-indole (0.5 g, 1.43 mmol) in anhydrous toluene (5 mL) was added LDA (0.855 mL, 1.71 mmol, 2.0 M in THF) at -78 °C under N₂. After addition, the reaction mixture was stirred at -78 °C for 30 min. 4-(trifluoromethyl)benzaldehyde (298 mg, 1.71 mmol) in THF (3 mL) was added dropwise at -78 °C under N₂. The reaction mixture was stirred at room temperature for 2 h. TLC (petroleum ether / EtOAc = 5/1) showed the reaction was completed. The mixture was quenched with water (10 mL) and extracted with EtOAc (3 X 15 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether : ethyl acetate = 7 : 3 to afford (5-bromo-1-tosyl-1H-indol-2-yl)(4-(trifluoromethyl)phenyl)methanol (0.36 g, 48%) as a yellow solid. ¹H NMR (CDCl₃ 400 MHz): δ 7.97 (d, *J* = 8.8 Hz, 1H), 7.62 (d, *J* = 8.0 Hz, 2H), 7.59 (d, *J* = 8.0 Hz, 2H), 7.54 (d, *J* = 4.0 Hz, 1H), 7.51 (d, *J* = 8.0 Hz, 2H), 7.41 (d, *J* = 8.0 Hz, 1H), 7.21 (d, *J* = 8.4 Hz, 2H), 6.42 (d, *J* = 4.8 Hz, 1H), 6.13 (s, 1H), 3.64 (d, *J* = 5.2 Hz, 1H), 2.37 (s, 3H).

Step 3

[0143] To a solution of (5-bromo-1-tosyl-1H-indol-2-yl)(4-(trifluoromethyl)phenyl)methanol (260 mg, 0.5 mmol) in CH₂Cl₂ (10 mL) was added Et₃SiH (1.4 mL) and TFA (1.4 mL) at 0 °C under N₂. The reaction mixture was stirred at room temperature for 2 h. TLC (petroleum ether / EtOAc = 5/1) showed the reaction was completed. The mixture was quenched with water (10 mL) and extracted with DCM (3 X 10 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether: ethyl acetate = 96: 4 to afford 5-bromo-1-tosyl-2-(4-(trifluoromethyl)benzyl)-1H-indole (0.23 g, 91%) as a white solid. **LC-MS Method 2** *t_R* = 1.018 min in 5-95AB_1.5min chromatography (Merck RP-18e 25-

2mm), MS (ESI) m/z 507.8 $[M+H]^+$. ^1H NMR ($CDCl_3$ 400 MHz): δ 8.05 (d, J = 8.8 Hz, 1H), 7.55 (d, J = 4.0 Hz, 1H), 7.51 (d, J = 8.0 Hz, 3H), 7.43 (d, J = 8.0 Hz, 2H), 7.39 (dd, J = 4.0, 8.0 Hz, 1H), 7.28 (s, 1H), 7.12 (d, J = 8.0 Hz, 2H), 6.17 (s, 1H), 4.42 (s, 2H), 2.34 (s, 3H).

Step 4

[0144] To a solution of 5-bromo-1-tosyl-2-(4-(trifluoromethyl)benzyl)-1H-indole (300 mg, 0.59 mmol) in dry MeOH (20 mL) was added DMSO (1 mL) and Et_3N (1 mL). Then $Pd(dppf)Cl_2$ (20 mg, 0.024 mmol) was added under N_2 . The mixture was stirred at 75 °C under CO (50 psi) for 20 h. The reaction mixture was filtered and the filter cake was washed with EtOAc (20 mL). The combined solvents were removed under reduced pressure. The mixture was partitioned between ethyl acetate (30 mL) and water (30 mL). The organic layer was washed with brine, dried over anhydrous Na_2SO_4 , filtered and concentrated to afford crude product. The crude product was purified by column chromatography on silica gel eluting with petroleum ether : ethyl acetate = 93 : 7 to afford methyl 1-tosyl-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylate (142 mg, 49%) as a white solid. **LC-MS Method 2** t_R = 1.11 min in 5-95AB_1.5min chromatography (Merck RP-18e 25-2mm), MS (ESI) m/z 488.0 $[M+H]^+$.

Step 5

[0145] To a solution of methyl 1-tosyl-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylate (90 mg, 0.19 mmol) in MeOH (3 mL) was added KOH (207 mg, 3.69 mmol) and water (1.0 mL). The reaction mixture was stirred at 50 °C for 20 h. The solvents were removed. Water (5 mL) was added. The mixture was acidified with 1 *N* HCl solution to pH = 6 and most solid was precipitated out. The solid was filtered and then dried under reduced pressure to afford 2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylic acid (59 mg, 100%) as a pale yellow solid. **LC-MS Method 2** t_R = 0.754 min in 5-95AB_1.5min chromatography (Merck RP-18e 25-2mm), MS (ESI) m/z 319.9 $[M+H]^+$.

Step 6

[0146] To a solution of 2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylic acid (59 mg, 0.185 mmol) in EtOH (5 mL) was added $SOCl_2$ (1 mL). Then the mixture was stirred at 60 °C for 3.5 h. Then the reaction solution was concentrated under reduced pressure. The residue was purified by preparative TLC (petroleum ether : ethyl acetate = 5 : 1) to afford ethyl 2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylate (40 mg, 63%) as a red solid. **LC-MS Method 2** t_R = 0.996 min in 5-95AB_1.5min chromatography (Merck RP-18e 25-2mm), MS (ESI) m/z 347.9 $[M+H]^+$.

Step 7

[0147] To a solution of ethyl 2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylate (30 mg, 0.086 mmol) in dry THF (2 mL) was added KOH (14 mg, 0.258 mmol) and EtI (27 mg, 0.172 mmol). Then the mixture was stirred at 50 °C for 1 h. TLC (petroleum ether / EtOAc = 5/1) showed the reaction was completed. The mixture was quenched with water (5 mL) and extracted with EtOAc (3 X 5 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC (petroleum ether : ethyl acetate = 5 : 1) to afford ethyl 1-ethyl-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylate (17 mg, 53%) as a red solid. LC-MS Method 2 t_R = 0.938 min in 5-95AB_1.5min chromatography (Merck RP-18e 25-2mm), MS (ESI) m/z 376.0 [M+H]⁺.

Step 8

[0148] To a solution of ethyl 1-ethyl-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylate (18 mg, 0.048 mmol) in EtOH (2 mL) was added H₂O (0.5 mL) and NaOH (38 mg, 0.96 mmol). The reaction mixture was stirred at room temperature for 20 h. TLC (petroleum ether / EtOAc = 5/1) showed the reaction was completed. The solvent was removed and water (5 mL) was added. The mixture was acidified with 1 N HCl solution to pH = 5 - 6 and extracted with EtOAc (3X10 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated to afford crude 1-ethyl-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylic acid (16 mg, 96% yield) as a white solid, which was used for next step without further purification.

[0149] Step 9

[0150] To a mixture of 1-ethyl-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylic acid (16 mg, 0.046 mmol), (4-(ethylsulfonyl)phenyl)methanamine hydrochloride (16 mg, 0.069 mmol) and HATU (9 mg, 0.069 mmol) in dry CH₂Cl₂ (2 mL) was added Et₃N (14 mg, 0.138 mmol). Then the mixture was stirred at room temperature for 1 h. Water (8 mL) was added. The mixture was extracted with CH₂Cl₂ (3 X 10 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by basic preparative HPLC separation, then lyophilized directly to afford 1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxamide (IN-1, 17.60 mg, 72%) as a white solid. LC-MS Method 2 t_R = 0.822 min in 5-95AB_1.5min chromatography (MK RP-18e 25-2 mm), MS (ESI) m/z 529.0 [M+H]⁺. ¹H NMR (CDCl₃ 400 MHz): δ 8.07 (s, 1H), 7.86 (d, J = 8.0 Hz, 2H), 7.70 (dd, J = 4.0, 8.0 Hz, 1H), 7.58 (d, J = 8.0 Hz, 2H), 7.56 (d, J = 8.0 Hz, 2H), 7.33 (d, J = 8.0 Hz, 2H), 7.32 (d, J = 8.0 Hz, 1H), 6.64 (t, J = 6.0 Hz, 1H), 6.33 (s, 1H), 4.78 (d, J = 6.0 Hz, 2H), 4.20 (s, 2H), 4.07 (q, J = 7.2 Hz, 2H), 3.10

(q, $J = 7.6$ Hz, 2H), 1.27 (t, $J = 7.6$ Hz, 3H), 1.21 (t, $J = 7.6$ Hz, 3H).

[0151] Basic preparative HPLC method

Mobile phase A: water with 0.05% $\text{NH}_3 \cdot \text{H}_2\text{O}$

Mobile phase B: CH_3CN

Flow rate: 25 mL/min.







Detection: UV 220 nm / 254 nm

Column: Gemini 150*25mm*5 μm

Column temperature: 30 °C

Time in min	%A	%B
0.00	48	52
10.00	18	82
10.20	0	100
12.00	0	100

[0152] The following compounds are prepared by similar procedures:

	
Compound IN-2	Compound IN-3
	
Compound IN-4 Isomer 1 ^{a,b}	Compound IN-4 Isomer 2 ^{a,c}
	


Compound IN-5***	Compound IN-6
Compound IN-7	Compound IN-8
Compound IN-9 ^d	
^a The isomers were separated by SFC on a ChiralPak AS-H column; ^b Longer t_R isomer ^c Shorter t_R isomer; ^d Step 5 was carried out under a strict nitrogen atmosphere. *** Reference example	


Example 2 (reference example)

1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-pyrrolo[2,3-b]pyridine-5-carboxamide

(Compound AI-1)

[0153]





Step 1

[0154] To a solution of **1** (650 mg, 3.31 mmol) in THF (5 mL) was added a 1M solution of LiAlH₄ in THF (3.97 mL, 3.97 mmol) dropwise at room temperature. The reaction was refluxed for 3 hours (~80 °C oil bath). After cooling to room temp, Et₂O (5 mL) was added and the reaction was filtered through celite. The filtrate was carefully quenched at 0 °C using 1 N NaOH until the bubbling subsided and the mixture stirred at room temp for 10 minutes after quenching. Et₂O and H₂O were added for the workup. The Et₂O layer was separated and dried using MgSO₄. Carefully evaporation of the Et₂O gave product **2** as an oil. This material was used directly for the next step without purification.

Step 2

[0155] To a solution of crude **2** (from above), Et₃N (0.92 mL, 6.62 mmol) and DMAP (40 mg, 0.33 mmol) in THF (5 mL) was added TsCl (948 mg, 4.48 mmol) at room temp. The reaction stirred for 15 hours. A sat. NaHCO₃ solution (5 mL) and DCM (10 mL) was added for the workup. The DCM layer was separated, washed with brine, dried using Na₂SO₄ and evaporated. The crude material was purified by ISCO flash chromatography (eluting with 10% EtOAc in hexanes) to afford 250 mg of **3** as a crystalline solid (22%, 2 steps).

LC-MS t_R = 1.809 min in 2 min chromatography (UV); no ionization peak

¹H NMR (CDCl₃, 400 MHz): δ 7.93 (d, *J* = 8.4 Hz, 1H), 7.78 (d, *J* = 8.4 Hz, 1H), 7.41 (d, *J* = 8.4 Hz, 1H), 7.35 (d, *J* = 8.4 Hz, 1H), 3.84 (d, *J* = 6.4 Hz, 2H), 2.46 (s, 3H), 1.98-1.82 (m, 5H), 1.69-1.63 (m, 1H), 1.33-1.23 (m, 2H), 1.03-0.93 (m, 2H).

Step 3

[0156] To a suspension of lithium acetylide ethylenediamine complex (71 mg, 0.77 mmol) in DMSO (2 mL) was added a solution of **3** (200 mg, 0.60 mmol) in DMSO (1 mL) at room temp. The reaction stirred for 2 hours at room temp. Water (10 mL) and Et₂O (15 mL) were added

for the workup. The Et₂O layer was separated, washed with brine and dried using MgSO₄. Carefully evaporation of the Et₂O gave volatile product **4** as a colorless oil (34 mg, 30%). This crude material was used for the next step without further purification. **LC-MS** t_R = No UV or ionization peak observed on LCMS ¹H NMR (CDCl₃, 400 MHz): δ 3.48 (s, 1H), 2.61 (d, J = 2.8 Hz, 2H), 2.13-2.10 (m, 2H), 1.98-1.96 (m, 4H), 1.42-1.32 (m, 2H), 1.11-1.05 (m, 2H).

Step 4

[0157] To a solution of **5** (300 mg, 1.19 mmol) in THF (5 mL) was added cyclopropylamine (0.18 mL, 2.5 mmol) dropwise at room temp. The reaction stirred for 24 hours at room temp. Water and EtOAc were added for the workup. The EtOAc layer was separated, washed with brine, dried using Na₂SO₄ and evaporated. The crude material was purified by ISCO flash chromatography (eluting with 30% EtOAc in hexanes) to give compound **6** (158 mg, 49%). **LC-MS** t_R = 1.286 min in 2 min chromatography; MS (ESI) m/z 271.19 and 273.19 [M+H]⁺.

Step 5

[0158] A suspension of **6** (50 mg, 0.18 mmol), **4** (34 mg, 0.18 mmol), PdCl₂(PPh₃)₂ (6 mg, 0.01 mmol), CuI (4 mg, 0.02 mmol) and Et₃N (0.05 mL, 0.36 mmol) in THF (2 mL) was degassed with N₂ for 5 min at room temp. The reaction vessel was sealed and heated to 85 °C for 15 hours. The reaction mixture was filtered through celite and evaporated. The crude material was purified by ISCO flash chromatography (eluting with 30% EtOAc in hexanes) to afford compound **7** (23 mg, 34%). **LC-MS** t_R = 1.735 min in 2 min chromatography; MS (ESI) m/z 381.31 [M+H]⁺. ¹H NMR (CDCl₃, 400 MHz): δ 8.83 (s, 1H), 8.05 (s, 1H), 5.78 (s, 1H), 3.90 (s, 3H), 2.95-3.00 (m, 1H), 2.46 (d, J = 5.2 Hz, 2H), 2.07-2.04 (m, 4H), 1.65-1.63 (m, 2H), 1.47-1.37 (m, 2H), 1.22-1.13 (m, 2H), 0.97-0.95 (m, 2H), 0.62-0.63 (m, 2H).

Step 6


[0159] To a solution of **7** (20 mg, 0.05 mmol) in t-BuOH (1 mL) was added KOtBu (30 mg, 0.26 mmol) at room temp. The reaction was heated to 90 °C for 12 hours. 1 N NaOH (2 mL) and iPrOAc (3 mL) were added for the workup. The iPrOAc layer was separated and extracted with 1 N NaOH (2 mL). The NaOH layers were combined and carefully acidified at 0 °C using conc. HCl to adjust the pH to ~4. The product (**8**) was then extracted into EtOAc twice. The EtOAc layers were combined, dried using Na₂SO₄ and evaporated to give nearly pure crude product **8** (16 mg, 84%). **LC-MS** t_R = 1.477 min in 2 min chromatography; MS (ESI) m/z 367.32 [M+H]⁺. ¹H NMR (CDCl₃, 400 MHz): δ 8.80 (s, 1H), 8.25 (s, 1H), 6.06 (s, 1H), 2.97-2.94 (1H,

m), 2.63 (d, $J = 6.8$ Hz, 2H), 1.90-1.57 (m, 6H), 1.12-0.70 (m, 8H).

Step 7

[0160] A solution of **8** (16 mg, 0.04 mmol), 4-(ethylsulfonyl)benzylamine (13 mg, 0.07 mmol), EDC (13 mg, 0.07), $i\text{Pr}_2\text{NEt}$ (9 mg, 0.07 mmol) and HOBt (10 mg, 0.07 mmol) in DMF (1 mL) was stirred at room temp for 24 hours. H_2O and EtOAc were added for the workup. The EtOAc layer was separated, dried using Na_2SO_4 and evaporated to give crude product. Purification by Gilson HPLC afforded 4.38 mg of compound **AI-1** as the TFA salt. **LC-MS** $t_R = 1.652$ min in 2 min chromatography; **MS** (ESI) m/z 548.55 $[\text{M}+\text{H}]^+$. **^1H NMR** (CD_3OD , 400 MHz): δ 9.12 (t, $J = 6.4$ Hz, 1H), 8.66 (d, $J = 2.4$ Hz, 1H), 8.30 (d, $J = 1.6$ Hz, 1H), 7.83 (d, $J = 8.8$ Hz, 2H), 7.58 (d, $J = 8.8$ Hz, 2H), 6.32 (s, 1H), 4.66 (d, $J = 4.0$ Hz, 2H), 3.20-3.18 (m, 1H), 3.13 (q, $J = 7.6$ Hz, 2H), 2.84 (d, $J = 6.8$ Hz, 2H), 2.09-2.02 (m, 1H), 1.91-1.88 (m, 4H), 1.82-1.77 (m, 1H), 1.35-1.03 (m, 8H), 1.14 (t, $J = 7.6$ Hz, 3H). **^{19}F NMR** (CD_3OD , 400 MHz): δ - 75.39 (d)


[0161] The following compounds are prepared using analogous procedures:





Example 3

1-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1R,4R)-4-(trifluoromethyl)cyclohexyl)oxy)-1H-benzo[d]imidazole-5-carboxamide (Compound BE-1)

[0162]





Step 1

[0163] Compound 1 is known compound in the literature. Xu, Y.; Lin, H.; Ruan, X.; Yang, S.; Hao, G.; Yang, W.; Yang, G. Eur. J. Med. Chem. 2015, 92, 427-438.

[0164] A solution of 1 (500 mg, 2.16 mmol) in POCl₃ (3 mL) was stirred at 120 °C for 2 hours. Upon cooling, the POCl₃ was removed by rotovap to give the crude product. This crude product was taken up in 5 mL of DCM and washed with sat. NaHCO₃ (3 mL). The DCM layer was then dried using Na₂SO₄ and evaporated. The crude material was purified using ISCO flash chromatography (eluting with 50% EtOAc in hexanes) to afford 307 mg of 2 as a white solid (57%). LC-MS t_R = 1.308 min in 2 min chromatography; MS (ESI) m/z 251.10 [M+H]⁺.

¹H NMR (CDCl₃, 400 MHz): δ 8.34 (s, 1H), 8.06 (d, *J* = 8.8 Hz, 1H), 7.57 (d, *J* = 8.8 Hz, 1H), 3.98 (s, 3H), 3.32-3.26 (m, 1H), 1.38-1.33 (m, 2H), 1.22-1.18 (m, 2H).

Step 2

[0165] To a solution of 3 (134 mg, 0.80 mmol) in DMF (2 mL) was added 60% NaH in mineral oil (30 mg, 0.76 mmol) at room temperature. The resulting orange colored reaction stirred at room temperature for 30 min before adding a solution of 2 (100 mg, 0.40 mmol) in DMF (1 mL) dropwise at room temperature. The reaction stirred for 30 min at which point a sat. NH₄Cl solution (1 mL) was carefully added at 0 °C to quench the reaction. The workup was done using EtOAc, washing the EtOAc layer once with brine. The EtOAc layer was separated, dried using Na₂SO₄ and evaporated to give the crude product containing a mixture of 4 as a cis:trans mixture (~1:1) in addition to compound 5 as a cis:trans mixture (~1:1). This crude mixture of product was used directly for the next step without further purification. LC-MS t_R = Compound 4 / 1.706 min in 2 min chromatography; MS (ESI) m/z 383.25 [M+H]⁺.


Step 3

[0166] A solution of the crude mixture of compounds **4** and **5** (crude material used directly from previous step) was stirred in a solution of 2N NaOH (1 mL), MeOH (1 mL), and THF (2 mL) at room temperature for 24 hours. iPrOAc (5 mL) was added followed by 1 N NaOH (3 mL). The iPrOAc layer was separated and extracted again with 1 N NaOH. The aqueous layers were combined and acidified to pH ~4 at 0 °C using conc. HCl. The product (**6**) was then extracted into EtOAc (2 times, 5 mL each). The EtOAc layer was then dried using Na₂SO₄ and evaporated to afford crude product **6** (~1:1 cis:trans mixture of isomers), which was used directly for the next step without further purification. LC-MS *t_R* = 1.536 min in 2 min chromatography; MS (ESI) *m/z* 369.22 [M+H]⁺.

Step 4

[0167] A solution of **6** (23 mg, 0.06 mmol), (R)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol (21 mg, 0.06 mmol), EDC (13 mg, 0.06 mmol), iPr₂NEt (15 mg, 0.12 mmol) and HOBt (10 mg, 0.07 mmol) in DMF (1 mL) was stirred for 24 hours at room temperature. H₂O and EtOAc were added for the workup. The EtOAc layer was separated, dried using Na₂SO₄ and evaporated to give crude product **7**. Purification by Gilson HPLC afforded 3.21 mg of the pure trans isomer of **7** as the TFA salt, which was less polar on the HPLC compared to the cis isomer of **7**. LC-MS *t_R* = trans isomer / 1.455 min in 2 min chromatography; MS (ESI) *m/z* 580.51 [M+H]⁺. ¹H NMR (CD₃OD, 400 MHz): δ 8.67 (d, *J* = 7.2 Hz, 1H), 7.87 (s, 1H), 7.79 (d, *J* = 8.8 Hz, 2H), 7.69 (d, *J* = 8.4 Hz, 1H), 7.60 (d, *J* = 8.0 Hz, 2H), 7.43 (d, *J* = 8.4 Hz, 1H), 5.22-5.17 (m, 1H), 4.88-4.80 (m, 1H), 3.82 (d, *J* = 6.0 Hz, 2H), 3.09 (q, *J* = 7.6 Hz, 2H), 3.07-3.02 (m, 1H), 2.32-2.29 (m, 2H), 2.22-2.15 (m, 1H), 2.02-1.99 (m, 2H), 1.63-1.42 (m, 4H), 1.10 (t, *J* = 7.6 Hz, 3H), 1.08-1.06 (m, 2H), 0.94-0.90 (m, 2H). ¹⁹F NMR (CD₃OD, 400 MHz): δ -74.93 (d)

[0168] The following compounds are prepared using analogous procedures:




Cpd	Cy ²	L ²	R ²	R ⁷
BE-2	<i>trans</i> -4-CF ₃ -cyclohexyl	O	Et	CH ₂ OH
BE-3	<i>trans</i> -4-CF ₃ -cyclohexyl	O	c-Pr	H
BE-4.1 a	<i>trans</i> -4-CF ₃ -cyclohexyl	O	i-Bu	H

Cpd	Cy ²	L ²	R ²	R ⁷
BE-4.2a	<i>cis</i> -4-CF ₃ -cyclohexyl	O	i-Bu	H
BE-5	4-CF ₃ -phenyl	O	Et	H
BE-6	4-CF ₃ -phenyl	O	c-Pr	H
BE-7	4-CF ₃ -phenyl	O	c-Pr	CH ₂ OH
BE-8	4-CF ₃ -phenyl	O	Et	CH ₂ OH
BE-9	4-CF ₃ O-phenyl	O	Et	CH ₂ OH
BE-10	4-CF ₃ -phenyl	O	Et	CH ₂ NMe ₂
BE-11	4-CF ₃ -phenyl	O	Et	CH ₂ NH ₂
BE-12***	4,4-difluorocyclohexyl	CH ₂ O ^b	Et	CH ₂ OH
BE-13***	3,3-difluorocyclobutyl	CH ₂ O ^b	Et	CH ₂ OH
BE-14***	4-CF ₃ -phenyl	CH ₂ O ^b	Et	CH ₂ OH
BE-15***	4-CF ₃ -phenyl	CH ₂ O ^b	c-Pr	CH ₂ OH
BE-16***	4,4-difluorocyclohexyl	CH ₂ O ^b	c-Pr	CH ₂ OH
BE-17***	cyclopropyl	CH ₂ O ^b	Et	CH ₂ OH

^a Isomers were separated by chromatography. ^b Cy² is attached to CH₂ and O is attached to the 2-position of the benzimidazole.
 *** Reference example

[0169] The following compound is prepared by using (R)-2-amino-2-(5-(ethylsulfonyl)pyridin-2-yl)ethan-1-ol in Step 4.




Compound BE-70

Example 4

1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-((*trans*-4-

(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide (Compound BE-18.1) and 1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-((*cis*-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide (Compound BE-18.2)

[0170]



Step 1

[0171] To a mixture of 4-fluoro-3-nitrobenzoic acid (4.3 g, 23.2 mmol) and (4-(ethylsulfonyl)phenyl)methanamine (2.3 g, 11.6 mmol) in anhydrous CH₂Cl₂ (120 mL) was added EDCI (4.4 g, 23.2 mmol), HOBT (3.1 g, 23.2 mmol) and Et₃N (3.5 g, 34.8 mmol). The mixture was stirred at room temperature for 3 h under N₂ atmosphere. The mixture was quenched with H₂O (100 mL). The aqueous layer was extracted with CH₂Cl₂ (3 X 100 mL). The combined organic layers were washed with brine (300 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether/ethyl acetate 1/2 to afford N-(4-(ethylsulfonyl)benzyl)-4-fluoro-3-nitrobenzamide (4.0 g, 94%) as a yellow solid.

Step 2

[0172] To a mixture of N-(4-(ethylsulfonyl)benzyl)-4-fluoro-3-nitrobenzamide (2.0 g, 5.46 mmol) in EtOH (50 mL) was added cyclopropanamine (623 mg, 10.93 mmol) and Et₃N (2.76 g, 27.3 mmol). The mixture was stirred at 70 °C under N₂ atmosphere for 3 h. The mixture was

concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with CH₂Cl₂/MeOH 19/1 to afford 4-(cyclopropylamino)-N-(4-(ethylsulfonyl)benzyl)-3-nitrobenzamide (2.13 g, 97%) as a yellow solid.

Step 3

[0173] To a solution of 4-(cyclopropylamino)-N-(4-(ethylsulfonyl)benzyl)-3-nitrobenzamide (2.13 g, 5.3 mmol) in MeOH (50 mL) was added Pd/C (1.2 g, dry, 10% w/w). The mixture was stirred at room temperature under H₂(30 psi) for 2 h. The mixture was filtered. The filtrate was concentrated under reduced pressure to afford crude 3-amino-4-(cyclopropylamino)-N-(4-(ethylsulfonyl)benzyl)benzamide (1.88 g, 95%) as a pale yellow solid, which was used for the next step without further purification. **LC-MS** t_R = 0.691 min in 5-95 AB_1.5 min chromatography (Welch Merck RP-18e, 25-2 mm), MS (ESI) m/z 374.1 [M+H]⁺.

Step 4

[0174] To a mixture of 3-amino-4-(cyclopropylamino)-N-(4-(ethylsulfonyl)benzyl)benzamide (213 mg, 0.57 mmol) and 2-(4-(trifluoromethyl)cyclohexyl)acetic acid (80 mg, 0.38 mmol) in anhydrous CH₂Cl₂ (4 mL) was added HATU (217 mg, 0.57 mmol) and Et₃N (192 mg, 1.9 mmol). The mixture was stirred at room temperature for 3 h under N₂ atmosphere. The mixture was quenched with H₂O (5 mL) and extracted with CH₂Cl₂ (3 X 5 mL). The combined organic layers were washed with brine (15 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether/ethyl acetate = 1/3 to afford 4-(cyclopropylamino)-N-(4-(ethylsulfonyl)benzyl)-3-(2-(4-(trifluoromethyl)cyclohexyl)acetamido)benzamide (260 mg, 100%) as a pale yellow oil. **LC-MS** t_R = 0.888 min in 5-95 AB_1.5 min chromatography (Welch Merck RP-18e, 25-2 mm), MS (ESI) m/z 566.2 [M+H]⁺.

Step 5

[0175] To a solution of 4-(cyclopropylamino)-N-(4-(ethylsulfonyl)benzyl)-3-(2-(4-(trifluoromethyl)cyclohexyl)acetamido)benzamide (130 mg, 0.23 mmol) in DCE (5 mL) was added TFA (0.5 mL). The mixture was stirred at 85 °C for 2 h under N₂ atmosphere. H₂O (5 mL) was added and the mixture was extracted with CH₂Cl₂ (3 X 5 mL). The combined organic layers were washed with *sat. aq.* NaHCO₃ solution (15 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC (petroleum ether / acetone = 1/1), SFC separation (AD-H), basic preparative HPLC separation and then dry-freezing to afford 1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-((trans-

4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide (**BE-18.1**, HCl salt, 21.8 mg, 31%) and 1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-((cis-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide (**BE-18.2**, HCl salt, 7.5 mg, 11%) as white solids.

[0176] Before SFC separation: Isomer SFC t_R = 4.144 and 5.392 min in 10 min chromatography (Column: AD-H; Method Name: AD-H_4_40_2_35ML.M, ee = 53%).

SFC separation condition:

[0177] Instrument: Berger MultiGram™ SFC, Mettler Toledo Co, Ltd

Column: AD 250mm*30mm, 10um

Mobile phase: A: Supercritical CO₂, B: iPrOH(0.05%NH₃H₂O), A:B =60:40 at 80ml/min Column

Temp: 38 °C

Nozzle Pressure: 100 Bar

Nozzle Temp: 60 °C

Evaporator Temp: 20 °C

Trimmer Temp: 25 °C

Wavelength: 220 nm

[0178] BE-18.1 LC-MS Method 2 t_R = 0.803 min in 5-95AB_1.5 min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 548.2 [M+H]⁺. ¹H NMR (CDCl₃ 400 MHz): δ 8.11 (s, 1H), 7.86 (d, J = 8.4 Hz, 2H), 7.84 (d, J = 8.0 Hz, 1H), 7.60-7.53 (m, 3H), 6.74 (t, J = 5.6 Hz, 1H), 4.78 (d, J = 5.6 Hz, 2H), 3.30-3.20 (m, 1H), 3.11 (q, J = 7.6 Hz, 2H), 3.06 (d, J = 7.6 Hz, 2H), 2.56-2.45 (m, 1H), 2.25-2.08 (m, 1H), 1.85-1.70 (m, 8H), 1.36-1.30 (m, 2H), 1.28 (t, J = 7.6 Hz, 3H), 1.13-1.05 (m, 2H). **Isomer SFC** t_R = 4.133 min in 10 min chromatography (Column: AD-H; Method Name: AD-H_4_40_2_35ML.M, ee = 100%).

Basic preparative HPLC Method:

[0179]

Mobile phase A: water with 0.05% NH₃ H₂O solution

Mobile phase B: CH₃CN

Flow rate: 30 mL/min.

Detection: UV 220 nm

Column: Agela DuraShell C18 150*25*5um

Column temperature: 40 °C

Time in min	%A	%B
0.00	54	46
8.00	24	76
8.20	0	100
10.00	0	100

[0180] **BE-18.2 LC-MS** Method 2 t_R = 0.807 min in 5-95AB_1.5 min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 548.1 $[M+H]^+$. ^1H NMR ($CDCl_3$ 400 MHz): δ 8.11 (s, 1H), 7.87 (d, J = 8.4 Hz, 2H), 7.84 (d, J = 8.4 Hz, 1H), 7.60-7.54 (m, 3H), 6.74 (t, J = 6.0 Hz, 1H), 4.78 (d, J = 5.6 Hz, 2H), 3.30-3.20 (m, 1H), 3.11 (q, J = 7.6 Hz, 2H), 2.95 (d, J = 7.6 Hz, 2H), 2.20-1.90 (m, 6H), 1.43-1.31 (m, 4H), 1.28 (t, J = 7.6 Hz, 3H), 1.23-1.03 (m, 4H). **Isomer SFC** t_R = 5.360 min in 10 min chromatography (Column: AD-H; Method Name: AD-H_4_40_2.35ML.M, ee = 99%).

Basic preparative HPLC Method:

[0181]

Mobile phase A: water with 0.05% NH_3 H_2O solution

Mobile phase B: CH_3CN

Flow rate: 30 mL/min.


Detection: UV 220 nm

Column: Agela DuraShell C18 150*25*5um

Column temperature: 40 °C


Time in min	%A	%B
0.00	57	43
8.00	27	73
8.20	0	100
10.00	0	100

[0182] The following compounds are prepared using analogous procedures:



Cpd No.	*	R ²	R ⁷	R ⁸	A ³	R ^A
BE-19.1 ^a	<i>trans</i>	i-Bu	H	H	CH	Et
BE-19.2 ^a	<i>cis</i>	i-Bu	H	H	CH	Et
BE-20	<i>trans</i>	Et	H	H	CH	Et
BE-21	<i>trans</i>	c-Pr	CH ₂ OH	H	CH	Et
BE-22	<i>trans</i>	c-Pr	H	CH ₂ OH	CH	Et
BE-23	<i>trans</i>	Et	CH ₂ OH	H	CH	Et
BE-24	<i>trans</i>	Et	H	CH ₂ OH	CH	Et
BE-25	<i>trans</i>	c-Pr	CH ₂ CH ₂ OH	H	CH	Et
BE-26.1a	<i>trans</i>	i-Bu	H	H	N	Et
BE-26.2a	<i>cis</i>	i-Bu	H	H	N	Et
BE-27.1a	<i>trans</i>	c-Pr	H	H	N	Et
BE-27.2a	<i>cis</i>	c-Pr	H	H	N	Et
BE-28	<i>trans</i>	c-Pr	CH ₂ OH	H	N	Et
BE-29	<i>trans</i>	c-Pr	H	CH ₂ OH	N	Et
BE-30	<i>trans</i>	c-Pr	H	H	CH	Me
BE-31 ^{***}	<i>trans</i>	c-Pr	H	H	N	NHMe
BE-32	<i>trans</i>	c-Pr	CH ₂ OH	H	CH	Me
BE-33 ^{***}	<i>trans</i>	c-Pr	H	H	CH	NHMe
^a Isomer pairs were separated by chromatography ^{***} Reference example						


[0183] The following compounds are prepared using analogous procedures:



Cpd No.	R ²	R ⁷	R ⁸	A ³
BE-34	Me	H	H	CH
BE-35	Et	H	H	CH
BE-36	c-Pr	H	H	CH
BE-37	CF ₃ CH ₂	H	H	CH


Cpd No.	R ²	R ⁷	R ⁸	A ³
BE-38	c-Bu	H	H	CH
BE-39***	t-BuCH ₂	H	H	CH
BE-40	i-Pr	H	H	N
BE-41	i-Bu	H	H	CH
BE-42	t-Bu	H	H	N
BE-43	i-Bu	H	H	N
BE-44	i-Bu	H	Me	CH
BE-45	i-Bu	Me	H	CH
*** Reference example				













[0184] The following compounds are prepared using analogous procedures:








Cpd No.	Cy2	R ²	R ⁷
BE-46.1 ^a	<i>trans</i> -4-methoxycyclohexyl	c-Pr	H
BE-46.2 ^a	<i>cis</i> -4-methoxycyclohexyl	c-Pr	H
BE-47	4-(trifluoromethoxy)phenyl	c-Pr	CH ₂ OH
BE-48	4-(difluoromethoxy)phenyl	c-Pr	H
BE-49	1-(<i>t</i> -butoxycarbonyl)piperidin-4-yl	c-Pr	H
BE-50	1-(2,2,2-trifluoroethyl)piperidin-4-yl	c-Pr	H
BE-51	6-(trifluoromethyl)pyridin-3-yl	c-Pr	H
BE-52	5-chloropyridin-2-yl	c-Pr	H
BE-53	4-(trifluoromethyl)piperidin-1-yl	i-Bu	H
BE-54	4-(trifluoromethyl)piperidin-1-yl	c-Pr	CH ₂ OH
^a Isomers were separated by chromatography			



[0185] The following compounds are prepared using analogous procedures:



			
Compound Nos BE-55.1 and -55.2*** *isomers separated by chromatography, absolute configuration unknown			
			
			
Compounds BE-56.1, -56.2, -56.3 and -56.4 *isomers were separated by chromatography, absolute configuration unknown			
			
			
Compounds BE-57.1, -57.2, -57.3 and -57.4 * isomers were separated by chromatography, absolute configuration unknown			
			



		
Compound BE-58***	Compound BE-59***	Compound BE-60***
		
Compound BE-61 ***	Compound BE-62***	
		
Compounds BE-63.1 and -63.2*** * isomers were separated by chromatography, absolute configuration unknown		



[0186] The following compounds are prepared using analogous procedures:


	
Compound Nos BE-71.1 and -71.2*** *isomers separated by chromatography, <i>cis/trans</i> configuration not assigned	
*** Reference example	


Example 5

[0187] The following compounds were prepared from other compounds of Formula I:

	
---	---

	
Compound BE-64***	Compound BE-65 ***
By oxidation of BE63.1 and -63.2 with MnO ₂	By oxidation of BE-36 with MnO ₂
	
Compound BE-66	Compound BE-67
By reaction of BE-21 with Cl ₃ CC(=O)NCO, followed by treatment with K ₂ CO ₃ in MeOH	By reaction of BE-21 with Cl ₃ CC(=O)NCO, followed by treatment with K ₂ CO ₃ in MeOH

Example 6 (reference example)**2-(4-(ethylsulfonyl)phenyl)-N-(1-isobutyl-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazol-5-yl)acetamide (Compound BE-68)****[0188]****[0189]** The following compounds is prepared using analogous procedures:




Compound BE-69

Example 7 (reference example)

**3-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-((*cis*-4-(trifluoromethyl)cyclohexyl)methyl)-3H-imidazo[4,5-*b*]pyridine-6-carboxamide
(Compound AB-1.2)**

[0190]



Step 1

[0191] To a solution of methyl 6-chloro-5-nitronicotinate (500 mg, 2.32 mmol) in CH₃CN (15 mL) was added cyclopropanamine (265 mg, 4.63 mmol) and Et₃N (585 mg, 5.78 mmol). The mixture was stirred at room temperature for 16 h. After concentrated under reduced pressure and added with water (5 mL), the mixture was extracted with ethyl acetate (3 X 5 mL). The combined organic layers were washed with brine (10 mL), dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure to afford methyl 6-(cyclopropylamino)-5-nitronicotinate (550 mg, 100%) as a yellow solid.

Step 2

[0192] To a solution of methyl 6-(cyclopropylamino)-5-nitronicotinate (550 mg, 2.32 mmol) in MeOH/ THF / H₂O (40 mL, v/v = 1/1/1) was added Zn (1.5 g, 23.2 mmol) and NH₄Cl (1.25 g, 23.2 mmol). The resulting mixture was stirred at 25 °C for 4 h. The mixture was filtered and the filtrate was concentrated under reduced pressure. After added with water (15 mL), the mixture was extracted with ethyl acetate (3 X 15 mL). The combined organic layers were washed with brine (30 mL), dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure and purified by chromatography column on silica gel eluting with petroleum ether: ethyl acetate = 1:1 to give methyl 5-amino-6-(cyclopropylamino)nicotinate (480 mg, 100%) as a red oil.

Step 3

[0193] To a solution of methyl 5-amino-6-(cyclopropylamino)nicotinate (207 mg, 1.0 mmol) and 2-(4-(trifluoromethyl)cyclohexyl)acetic acid (210 mg, 1.0 mmol) in anhydrous DMF (8 mL) was added EDCI (287 mg, 1.5 mmol), HOBt (203 mg, 1.5 mmol) and Et₃N (303 mg, 3.0 mmol). The resulting mixture was stirred at room temperature for 16 h. After added with water (10 mL), the mixture was extracted with ethyl acetate (3 X 10 mL). The combined organic layers were washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure and purified by chromatography column on silica gel eluting with petroleum ether: ethyl acetate = 1:1 to give methyl 6-(cyclopropylamino)-5-(2-(4-(trifluoromethyl)cyclohexyl)acetamido)nicotinate (260 mg, 65%) as a red oil. **LC-MS** *t_R* = 0.795 min in 5-95 AB_1.5 min chromatography (Welch Xtimate MK RP-18e 25-2mm), MS (ESI) *m/z* 400.0 [M+H]⁺.

Step 4

[0194] Methyl 6-(cyclopropylamino)-5-(2-(4-(trifluoromethyl)cyclohexyl)acetamido)nicotinate (150 mg, 0.38 mmol) was dissolved in AcOH (10 mL). The resulting mixture was stirred at 120 °C for 1 h. The mixture was concentrated under reduced pressure. After added with water (10 mL) and adjusted to pH = 7 by saturated NaHCO₃ solution (5 mL), the mixture was extracted with ethyl acetate (3 X 10 mL). The combined organic layers were washed with brine (20 mL), dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure and purified by preparative TLC with petroleum ether: ethyl acetate = 1:2 to give the low polarity spot methyl 3-cyclopropyl-2-((*trans*-4-(trifluoromethyl)cyclohexyl)methyl)-3H-imidazo[4,5-*b*]pyridine-6-carboxylate (50 mg, 35%) and the high polarity spot methyl 3-cyclopropyl-2-((*cis*-4-(trifluoromethyl)cyclohexyl)methyl)-3H-imidazo[4,5-*b*]pyridine-6-carboxylate (16 mg, 11%) as colorless oil. **LC-MS** t_R = 0.897 min in 5-95 AB_1.5 min chromatography (Welch Xtimate MK RP-18e 25-2mm), MS (ESI) m/z 382.1 [M+H]⁺.

Step 5

[0195] To a solution of methyl 3-cyclopropyl-2-((*cis*-4-(trifluoromethyl)cyclohexyl)methyl)-3H-imidazo[4,5-*b*]pyridine-6-carboxylate (16 mg, 0.042 mmol) in anhydrous THF / MeOH (2 mL, v/v = 1/1) was added NaOH (0.7 mL, 1*N* in H₂O). The resulting mixture was stirred at room temperature for 2 h. The mixture was concentrated under reduced pressure. After adjusted to pH = 4 with 4*N* HCl solution, the mixture was extracted with ethyl acetate (3 X 2 mL). The combined organic layers were washed with brine (5 mL), dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure to give 3-cyclopropyl-2-((*cis*-4-(trifluoromethyl)cyclohexyl)methyl)-3H-imidazo[4,5-*b*]pyridine-6-carboxylic acid (15 mg, 100%) as a white solid. **LC-MS** t_R = 0.712 min in 5-95 AB_1.5 min chromatography (Welch Xtimate MK RP-18e 25-2mm), MS (ESI) m/z 367.9 [M+H]⁺.

Step 6

[0196] To a solution of 3-cyclopropyl-2-((*cis*-4-(trifluoromethyl)cyclohexyl)methyl)-3H-imidazo[4,5-*b*]pyridine-6-carboxylic acid (15 mg, 0.042 mmol) and (4-(ethylsulfonyl)phenyl)methanamine (25 mg, 0.126 mmol) in anhydrous DMF (2 mL) was added EDCI (40 mg, 0.21 mmol), HOBt (29 mg, 0.21 mmol) and Et₃N (42 mg, 0.42 mmol). The resulting mixture was stirred at room temperature for 16 h. After added with water (3 mL), the mixture was extracted with CH₂Cl₂ (3 X 5 mL). The combined organic layers were washed with brine (3 X 10 mL), dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure and purified by preparative TLC with ethyl acetate, then HCl preparative HPLC separation to give 3-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-((*cis*-4-(trifluoromethyl)cyclohexyl)methyl)-3H-imidazo[4,5-*b*]pyridine-6-carboxamide (**AB-1.2**, HCl salt, 9.3 mg, 38%) as a pale yellow solid. **AB-1.2 (HCl salt)** **LC-MS** t_R = 0.852 min in 5-95 AB_1.5

min chromatography (Welch Xtimate MK RP-18e 25-2mm), MS (ESI) m/z 549.1 $[M+H]^+$. ^1H NMR (CD_3OD 400 MHz): δ 9.13 (d, J = 2.0 Hz, 1H), 8.60 (d, J = 2.0 Hz, 1H), 7.89 (d, J = 8.4 Hz, 2H), 7.67 (d, J = 8.4 Hz, 2H), 4.75 (s, 2H), 3.63-3.55 (m, 1H), 3.28 (d, J = 7.2 Hz, 2H), 3.20 (q, J = 7.2 Hz, 2H), 2.24-2.06 (m, 2H), 2.05-1.93 (m, 4H), 1.50-1.25 (m, 9H), 1.21 (t, J = 7.6 Hz, 3H). **Isomer SFC** t_R = 4.82 min in 16 min chromatography (Column: AD-H; Method Name: AD-H_3UM_4_40_2.5ML.met, ee = 96.27%).

HCl preparative HPLC Method:

[0197]

Mobile phase A: water with 0.1% HCl solution

Mobile phase B: CH_3CN

Flow rate: 25 mL/min.



Detection: UV 220 nm / 254 nm

Column: Gemini 150*25mm*5um

Column temperature: 30 °C


Time in min	%A	%B
0.00	40	60
10.00	96	4
10.20	0	100
12.00	0	100

[0198] The following compounds are prepared using analogous procedures:

	
Compound AB-2	Compound AB-3

Example 8

(R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((4-

(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole-5-carboxamide (Compound IN-10)**[0199]****Step 1. 5-bromo-1-ethyl-1H-indole-2-carbaldehyde**

[0200] A mixture of 5-bromo-1H-indole-2-carbaldehyde (0.6420 g, 2.86 mmol), Cs_2CO_3 (4.060 g, 12.5 mmol), and EtI (1.0 mL, 12.5 mmol) in DMF (10 mL) was stirred at room temperature for 2 d. The reaction mixture was then quenched with saturated NH_4Cl and extracted twice with ethyl acetate. The combined organic layers were dried over Na_2SO_4 and evaporated. The residue was purified by flash chromatography on silica gel eluted with 0 $\rightarrow$ 40% ethyl acetate/hexanes to give 0.6775 g (94%) of 5-bromo-1-ethyl-1H-indole-2-carbaldehyde. LC-MS Method 1 t_R = 1.71 min, m/z 252, 254 (MH^+); ^1H NMR (CD_3OD , 400 MHz) δ 9.86 (s, 1H), 7.89 (s, 1H), 7.48 (m, 2H), 7.33 (s, 1H), 4.60 (q, J = 7.16 Hz, 2H), 1.33 (t, J = 7.22 Hz, 3H).

Step 2. 5-bromo-1-ethyl-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole

[0201] A mixture of 5-bromo-1-ethyl-1H-indole-2-carbaldehyde (0.4022 g, 1.60 mmol), 4-(trifluoromethyl)piperidine (0.5120 g, 3.34 mmol), and $\text{NaBH}(\text{OAc})_3$ (0.7330 g, 3.46 mmol) in 1,2-dichloroethane (15 mL) was stirred at room temperature for 23 h. The reaction mixture was then quenched with saturated NaHCO_3 and extracted three times with dichloromethane. The combined organic layers were dried over Na_2SO_4 and evaporated. The residue was purified by flash chromatography on silica gel eluted with 0 $\rightarrow$ 4% MeOH/DCM to afford 0.5769 g (93%) of 5-bromo-1-ethyl-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole. LC-MS

Method 1 t_R = 1.12 min, m/z 389, 391 (MH^+), 236, 238 ($M^+ - 152$); ^1H NMR (CD_3OD , 400 MHz) δ 7.76 (s, 1H), 7.44 (d, J = 8.78 Hz, 1H), 7.35 (d, J = 8.78 Hz, 1H), 6.77 (s, 1H), 4.61 (s, 2H), 4.36 (q, J = 7.13 Hz, 2H), 3.71-3.68 (m, 2H), 3.20-3.14 (m, 2H), 2.63-2.59 (m, 1H), 2.19-2.15 (m, 2H), 1.94-1.77 (m, 2H), 1.31 (t, J = 7.18 Hz, 3H); ^{19}F NMR (CD_3OD , 376 MHz) δ -75.56.

Step 3. 1-ethyl-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole-5-carboxylic acid


[0202] A mixture of 5-bromo-1-ethyl-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole (0.3196 g, 0.821 mmol), Molybdenumhexacarbonyl (0.3835 g, 1.45 mmol), Herrmann's palladacycle (0.1550 g, 0.165 mmol), Tri-*tert*-butylphosphonium tetrafluoroborate (0.1595 g, 0.550 mmol), water (0.4 mL), and DBU (0.4 mL) in dioxane (4 mL) was heated in microwave at 150 °C for 20 min. The reaction mixture was purified by reverse-phase HPLC (Phenomenex® Luna 5 μ C₁₈ (2) 250 × 21.20 mm column, 10% CH₃CN/H₂O, 0.1% CF₃COOH over 1.5 min, 10% → 90% CH₃CN/H₂O, 0.1% CF₃COOH over 8 min, and then 90% CH₃CN/H₂O, 0.1% CF₃COOH over 4 min, flow rate 25 mL/min) to yield 0.2676 g (70%) of 1-ethyl-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole-5-carboxylic acid as a TFA salt. LC-MS Method 1 t_R = 0.78 min, m/z 355 (MH^+), 202 ($M^+ - 152$); ^1H NMR (CD_3OD , 400 MHz) δ 8.38-8.37 (m, 1H), 7.96 (dd, J = 8.79, 0.88 Hz, 1H), 7.57 (d, J = 8.79 Hz, 1H), 6.93 (s, 1H), 4.63 (s, 2H), 4.41 (q, J = 7.13 Hz, 2H), 3.73-3.70 (m, 2H), 3.20-3.13 (m, 2H), 2.66-2.58 (m, 1H), 2.21-2.18 (m, 2H), 1.85-1.80 (m, 2H), 1.35 (t, J = 7.18 Hz, 3H); ^{19}F NMR (CD_3OD , 376 MHz) δ -75.58.

Step 4. (R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole-5-carboxamide

[0203] A mixture of 1-ethyl-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole-5-carboxylic acid (0.0495 g, 0.106 mmol), (R)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol (0.0516 g, 0.194 mmol), HATU (0.2154 g, 0.566 mmol), and Et₃N (0.3 mL) in DMF (1.5 mL) was stirred at room temperature for 2 h. The reaction mixture was purified by reverse-phase HPLC (Phenomenex® Luna 5 μ C₁₈ (2) 250 × 21.20 mm column, 10% CH₃CN/H₂O, 0.1% CF₃COOH over 1.5 min, 10% → 90% CH₃CN/H₂O, 0.1% CF₃COOH over 8 min, and then 90% CH₃CN/H₂O, 0.1% CF₃COOH over 4 min, flow rate 25 mL/min) and the fractions were then lyophilized to afford (R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole-5-carboxamide TFA salt as a white fluffy solid. LC-MS Method 1 t_R = 0.82 min, m/z 566 (MH^+); ^1H NMR (CD_3OD , 400 MHz) δ 8.72 (d, J = 7.03 Hz, 1H), 8.25 (m, 1H), 7.89 (d, J = 8.20 Hz, 2H), 7.85 (dd, J = 8.79, 0.87 Hz, 1H), 7.70 (d, J = 8.20 Hz, 2H), 7.60 (d, J = 8.79 Hz, 1H), 6.94 (s, 1H), 5.33-5.28 (m, 1H), 4.64 (s, 2H), 4.42 (q, J = 7.13 Hz, 2H), 3.93 (d, J = 6.15 Hz, 2H), 3.74-3.71 (m, 2H), 3.22-3.16 (m, 4H), 2.64-2.59 (m, 1H), 2.22-1.19 (m, 2H), 1.83-1.79 (m, 2H), 1.35 (t, J = 7.18 Hz, 3H), 1.21 (t, J = 7.46 Hz,


3H); ^{19}F NMR (CD_3OD , 376 MHz) δ -75.58.

[0204] The following compound is prepared by a similar procedure using (R)-2-amino-2-(5-(ethylsulfonyl)pyridin-2-yl)ethan-1-ol in Step 4.




Compound IN-11

[0205] The following compounds are prepared from 5-bromo-1-cyclopropyl-1H-indole-2-carbaldehyde using procedures similar to those in Steps 2-4.




Compound IN-12



Compound IN-13

[0206] The starting material for Step 2 for the preparation of these compounds was prepared as shown in below in Alternate Step 1.



Alternate Step 1: 5-Bromo-1-cyclopropyl-1H-indole-2-carbaldehyde


[0207] A mixture of 5-bromo-1H-indole-2-carbaldehyde (0.3482 g, 1.55 mmol), cyclopropylboronic acid (0.2948 g, 3.43 mmol), Cu(OAc)_2 (0.2930 g, 1.61 mmol), 2,2'-dipyridyl (0.2490 g, 1.59 mmol), and Na_2CO_3 (0.5135 g, 4.84 mmol) in dichloroethane (15 mL) was vigorously stirred at 60 $^\circ\text{C}$ for 3 h under air. The reaction mixture was cooled to room

temperature, and then quenched with saturated NH_4Cl and extracted twice with dichloromethane. The combined organic layers were dried over Na_2SO_4 and evaporated. The residue was purified by flash chromatography on silica gel eluted with 0 $\rightarrow$ 80% ethyl acetate/hexanes to afford 0.3690 g (90%) of 5-bromo-1-cyclopropyl-1H-indole-2-carbaldehyde. LC-MS Method 1 t_R = 1.70 min, m/z 264, 266 (MH^+).

Example 9

1-ethyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((trans)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide and 1-ethyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((cis)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide (Compounds IN-14.1 and -14.2))

[0208]



Step 1

[0209] To a solution of 5-chloro-2-methyl-1H-indole (1.5 g, 9.1 mmol) in anhydrous THF (20 mL) was added $n\text{-BuLi}$ (11 mL, 27.5 mmol, 2.5 M in hexane) dropwise at -70°C under N_2 . The

mixture was stirred at -70 °C under N₂ for 10 min, followed by adding t-BuOK (18 mL, 18 mmol, 1 M in THF) dropwise. The mixture was allowed to warm to -30 °C and stirred for 10 min, then cooled to -70 °C. A solution of 4-(trifluoromethyl)cyclohexanone (1.7 g, 10.2 mmol) in anhydrous THF (10 mL) was added dropwise. The mixture was stirred at -70 °C under N₂ for 2 h. TLC (petroleum ether: ethyl acetate = 5:1) showed 5-chloro-2-methyl-1H-indole was still remained and two new spot was found. The mixture was quenched with sat. NH₄Cl solution (10 mL), added H₂O (20 mL) and extracted with ethyl acetate (3 X 30 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether: ethyl acetate = 10:1-5:1 to afford 1-((5-chloro-1H-indol-2-yl)methyl)-4-(trifluoromethyl)cyclohexanol (1.2 g, 39%) as a pale yellow solid.

Step 2

[0210] The solution of 1-((5-chloro-1H-indol-2-yl)methyl)-4-(trifluoromethyl)cyclohexanol (1.2 g, 3.6 mmol) in TFA (3 mL) was heated to 100 °C and stirred for 2 h. TLC (petroleum ether: ethyl acetate = 5:1) showed the reaction was completed. The mixture was adjusted to pH = 7-8 by sat. NaHCO₃ solution and extracted with ethyl acetate (3 X 30 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether: ethyl acetate = 20:1 to afford 5-chloro-2-((4-(trifluoromethyl)cyclohexylidene)methyl)-1H-indole (600 mg, 55%) as a yellow solid. **LC-MS** *t_R* = 5.010 min in 10-80AB 7.0 min chromatography (Welch Xtimate 3um,C18, 2.1*30mm), MS (ESI) *m/z* 313.9 [M+H]⁺.

Step 3

[0211] To a solution of 5-chloro-2-((4-(trifluoromethyl)cyclohexylidene)methyl)-1H-indole (1 g, 3.2 mmol) in anhydrous DMF (15 mL) and anhydrous MeOH (20 mL) was added Pd(OAc)₂ (150 mg, 0.64 mmol), dcpp-2BF₄ (390 mg, 0.64 mmol) and K₂CO₃ (1.3 g, 9.4 mmol) under N₂, followed bubbled with CO (15 psi) for 10 min and stirred at 100 °C for 16 h under CO (50 psi). LCMS showed the desired product was found. The mixture was filtered, added H₂O (30 mL) and extracted with EtOAc (3 X 30 mL). The combined organic layers were washed with H₂O (3 X 50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether: ethyl acetate = 10:1 to afford methyl 2-((4-(trifluoromethyl)cyclohexylidene)methyl)-1H-indole-5-carboxylate (300 mg, 28%) as a yellow solid. **LC-MS** *t_R* = 4.475 min in 10-80 AB_7.0 min chromatography (Welch Xtimate 3um,C18, 2.1*30mm), MS (ESI) *m/z* 337.9 [M+H]⁺.

Step 4

[0212] To a solution of methyl 2-((4-(trifluoromethyl)cyclohexylidene)methyl)-1H-indole-5-carboxylate (300 mg, 0.89 mmol) in ethyl acetate (5 mL) was added Pd/C (50 mg, 10%wt) under H₂. The mixture was stirred at room temperature for 2 h under H₂ (15 psi). LCMS showed the reaction was completed. The mixture was filtered and concentrated under reduced pressure to afford crude methyl 2-((4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxylate (280 mg, 93%) as a yellow solid, which was used for next step without further purification. **LC-MS** Method 2 t_R = 0.885 min in 5-95 AB_1.5 min chromatography (Welch RP-18e, 25-2mm), MS (ESI) m/z 339.9 [M+H]⁺.

Step 5

[0213] To a solution of methyl 2-((4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxylate (280 mg, 0.83 mmol) in anhydrous THF (5 mL) and DMF (5 mL) was added Cs₂CO₃ (850 mg, 2.61 mmol) and EtI (480 mg, 2.61 mmol) under N₂. The mixture was stirred at 60 °C for 5 h under N₂. TLC (petroleum ether: ethyl acetate = 5:1) showed the reaction was completed. The mixture was added with EtOAc (10 mL) and washed with H₂O (3 X 10 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether: ethyl acetate = 10:1 to afford methyl 1-ethyl-2-((4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxylate (230 mg, 76%) as a yellow oil. **LC-MS** Method 2 t_R = 0.912 min in 5-95AB_1.5 min chromatography (Welch RP-18e, 25-2mm), MS (ESI) m/z 368.0 [M+H]⁺.

Step 6

[0214] To a solution of methyl 1-ethyl-2-((4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxylate (230 mg, 0.63 mmol) in MeOH (5 mL) was added NaOH solution (1.5 mL, 3.0 mmol, 2 M in H₂O). The mixture was stirred at 60 °C for 20 h. TLC (petroleum ether: ethyl acetate = 5:1) showed the reaction was completed. The mixture was adjusted to pH = 5-6 by HCl solution (2 M in H₂O) and extracted with EtOAc (3 X 5 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford crude 1-ethyl-2-((4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxylic acid (200 mg, 90%) as a yellow solid, which was used for next step without further purification.

Step 7

[0215] To a solution of 1-ethyl-2-((4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxylic acid (60 mg, 0.17 mmol) and (R)-2-amino-2-(5-(ethylsulfonyl)pyridin-2-yl) ethanol (90 mg, 0.34 mmol) in DMF (3 mL) was added Et₃N (52 mg, 0.51 mmol) and HATU (195 mg, 0.51 mmol). The mixture was stirred at 12 °C for 16 h. LCMS showed the reaction was completed. The mixture was added EtOAc (10 mL) and washed with H₂O (3 X 10 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC with ethyl acetate to afford a mixture of - ethyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((trans)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide and 1-ethyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((cis)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide (100 mg, 104%) as a yellow solid. The mixture was separated by SFC followed by basic preparative HPLC to give Compound IN-14.1 (6.90 mg, 7%, t_R = 4.276 min) and Compound IN-14.2 (21.40 mg, 21%, t_R = 3.977 min) as white solids.

Before SFC separation :

[0216] Isomer SFC t_R = 1.869 and 2.034 min in 3 min chromatography (Column: AD-H; Method Name: AD-H_3UM_4_5_40_4ML_3MIN.M, ee = 41.26%).

SFC separation condition:

[0217] Instrument: Thar 80

Column: AD(250mm*30mm,10um)

Mobile phase: A: Supercritical CO₂, B: MeOH (0.05% NH₃H₂O), A: B = 45: 45 at 80 mL/min

Column Temp: 38 °C

Nozzle Pressure: 100 Bar

Nozzle Temp: 60 °C

Evaporator Temp: 20 °C

Trimmer Temp: 25 °C

Wavelength: 220 nm

[0218] Compound IN-14.1 (6.90 mg, 7%) as a white solid

LC-MS t_R = 0.795 min in 5-95 AB_1.5 min chromatography (Welch MK RP-18e 25-2mm), MS

(ESI) m/z 566.1 [M+H]⁺ ¹H NMR (CDCl₃ 400 MHz): δ 9.06 (s, 1H), 8.20 (d, J = 8.4 Hz, 1H), 8.07 (s, 1H), 7.74 (d, J = 8.0 Hz, 1H), 7.70 (d, J = 7.6 Hz, 2H), 7.32 (d, J = 8.8 Hz, 1H), 6.32 (s, 1H), 5.49 (q, J = 7.2 Hz, 2H), 4.25 (d, J = 7.2 Hz, 1H), 4.16 (q, J = 7.6 Hz, 2H), 3.96 (q, J = 7.2 Hz, 1H), 3.72 (brs, 1H), 3.17 (q, J = 7.6 Hz, 2H), 2.66 (d, J = 6.8 Hz, 2H), 2.10-1.95 (m, 4H), 1.68-1.61 (m, 1H), 1.33 (t, J = 7.2 Hz, 3H), 1.32-1.29 (m, 3H), 1.29 (t, J = 7.2 Hz, 3H), 1.09-0.99 (m, 2H). **Isomer SFC t_R = 4.276 min** in 10 min chromatography (Column: AS-H; Method

Name: AS-H_S_3_5_40_3ML_8MIN_15CM.M, ee = 98.37%).

Basic preparative HPLC method

[0219]

Mobile phase A: water with 0.05% NH₃-H₂O solution

Mobile phase B: CH₃CN

Flow rate: 25 mL/min.

Detection: UV 220 nm / 254 nm

Column: Gemini 150*25 5u

Column temperature: 40 °C

Time in min	%A	%B
0.00	50	50
10.00	20	80
10.20	0	100
14.00	0	100

[0220] Compound IN-14.2 (21.40 mg, 21%) as a white solid

LC-MS t_R = 0.785 min in 5-95 AB_1.5 min chromatography (Welch MK RP-18e 25-2mm), MS (ESI) m/z 566.0 [M+H]⁺ ¹H NMR (CDCl₃ 400 MHz): δ 9.06 (s, 1H), 8.19 (d, J = 7.2 Hz, 1H), 8.08 (s, 1H), 7.73 (d, J = 8.4 Hz, 2H), 7.68 (d, J = 7.2 Hz, 1H), 7.32 (d, J = 8.4 Hz, 1H), 6.34 (s, 1H), 5.49 (q, J = 5.6 Hz, 1H), 4.23 (d, J = 7.2 Hz, 1H), 4.16 (q, J = 7.6 Hz, 2H), 3.96 (q, J = 7.2 Hz, 1H), 3.77 (q, J = 7.2 Hz, 1H), 3.17 (q, J = 7.6 Hz, 2H), 2.78 (d, J = 7.6 Hz, 2H), 2.14-1.95 (m, 2H), 1.74-1.64 (m, 8H), 1.37 (t, J = 7.2 Hz, 3H), 1.32 (t, J = 7.2 Hz, 3H). Isomer SFC t_R = 3.977 min in 10 min chromatography (Column: AS-H; Method Name: ASH_S_3_5_40_3ML_8MIN_15CM.M, ee = 100.00%)

Basic preparative HPLC method

[0221]

Mobile phase A: water with 0.05% NH₃-H₂O solution

Mobile phase B: CH₃CN

Flow rate: 25 mL/min.







Detection: UV 220 nm / 254 nm

Column: Phenomenex Gemini 150*25mm*10um

Column temperature: 40 °C

Time in min	%A	%B
0.00	50	50
10.00	20	80
10.20	0	100
14.00	0	100


[0222] The following compounds are prepared by substituting the appropriate amines in Step 7.

	
Compounds IN-15.1 and IN-15.2 *isomers separated by chromatography, <i>cis/trans</i> configuration not assigned	
	
Compounds IN-16.1 and IN-16.2 *isomers separated by chromatography, <i>cis/trans</i> configuration not assigned	
	
Compounds IN-17.1 and IN-17.2 *isomers separated by chromatography, <i>cis/trans</i> configuration not assigned	

Example 10

1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxamide (Compound IN-18)

[0223]



Step 1

[0224] To a mixture of ethyl 2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylate from Example 1 Step 6 (50 mg, 0.14 mmol), cyclopropylboronic acid (24 mg, 0.28 mmol), DMAP (50 mg, 0.42 mmol) and Cu(OAc)₂ (25 mg, 0.14 mmol) in dry toluene (2 mL) was added NaHMDS (0.42 mL, 0.42 mmol, 1N in THF) under N₂. Then the mixture was stirred at 95 °C for 2 h under O₂ (15 psi). TLC (petroleum ether/ethyl acetate = 5/1) showed the reaction was completed. The mixture was filtered and washed with ethyl acetate (15 mL). The filtrate was concentrated under reduced pressure. The residue was purified by pre-TLC (petroleum ether/ethyl acetate = 5 /1) to afford ethyl 1-cyclopropyl-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylate (18 mg, 33%) as a colorless oil. **LC-MS Method 2** t_R = 0.963 min in 5-95AB_1.5min chromatography (Merck RP-18e 25-2mm), MS (ESI) m/z 388.0 [M+H]⁺.

Step 2

[0225] To a solution of ethyl 1-cyclopropyl-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylate (20 mg, 0.05 mmol) in EtOH (3 mL) was added *aq.* NaOH solution (1 mL, 1.0 mmol, 1N). The reaction mixture was stirred at 26 °C for 20 h. TLC (petroleum ether/ethyl acetate = 5 /1) showed the reaction was not completed. Then the mixture was stirred at 50 °C for another 20 h. LCMS showed the reaction was completed. The solvent was removed and

water (10 mL) was added. The mixture was acidified with 1 N HCl solution to pH = 5 and extracted with ethyl acetate (3 X 10 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated to afford crude 1-cyclopropyl-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylic acid (17 mg, 94%) as a white solid, which was used for the next step without further purification.**LC-MS Method 2** *t_R* = 0.821 min in 5-95AB_1.5min chromatography (Merck RP-18e 25-2mm), MS (ESI) *m/z* 360.0 [M+H]⁺.

Step 3

[0226] To a mixture of 1-cyclopropyl-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxylic acid (17 mg, 0.047 mmol), (4-(ethylsulfonyl)phenyl)methanamine (17 mg, 0.07 mmol) and HATU (36 mg, 0.094 mmol) in dry CH₂Cl₂ (2 mL) was added Et₃N (14 mg, 0.14 mmol). Then the mixture was stirred at 26 °C for 2 h. LCMS showed the starting material was consumed completely. Water (5 mL) was added. The mixture was extracted with CH₂Cl₂ (3 X 10 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by neutral preparative HPLC separation, then basic preparative HPLC separation and then lyophilized directly to afford 1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxamide (Compound IN-18, 10.5 mg, 42%) as a white solid.

LC-MS Method 2 *t_R* = 0.829 min in 5-95AB_1.5min chromatography (MK RP-18e 25-2 mm), MS (ESI) *m/z* 541.0 [M+H]⁺. **¹H NMR** (CDCl₃ 400 MHz): δ 8.02 (s, 1H), 7.88 (d, *J* = 8.4 Hz, 2H), 7.67 (d, *J* = 8.4 Hz, 1H), 7.57 (d, *J* = 8.0 Hz, 5H), 7.34 (d, *J* = 8.0 Hz, 2H), 6.61 (t, *J* = 6.4 Hz, 1H), 6.28 (s, 1H), 4.79 (d, *J* = 6.4 Hz, 2H), 4.31 (s, 2H), 3.11 (q, *J* = 7.2 Hz, 2H), 3.00-2.90 (m, 1H), 1.28 (t, *J* = 7.2 Hz, 3H), 1.15-1.12 (m, 2H), 1.07-1.02 (m, 2H).

Neutral preparative HPLC method

[0227]

Mobile phase A: water with 10mM NH₄HCO₃

Mobile phase B: CH₃CN

Flow rate: 22 mL/min.

Detection: UV 220 nm / 254 nm

Column: Phenomenex Synergi C18 150*25*10um

Column temperature: 30 °C

Time in min	%A	%B
0.00	62	42
10.00	32	72

Time in min	%A	%B
10.20	0	100
13.00	0	100

Basic preparative HPLC method

[0228]

Mobile phase A: water with 0.05% ammonia hydroxide

Mobile phase B: CH₃CN

Flow rate: 25 mL/min.


Detection: UV 220 nm / 254 nm

Column: DuraShell 150*25mm*5um

Column temperature: 30 °C

Time in min	%A	%B
0.00	47	53
10.00	17	83
10.20	0	100
13.00	0	100

[0229] The compound below is prepared by a similar procedure using (R)-2-amino-2-(5-(ethylsulfonyl)pyridin-2-yl)ethan-1-ol in Step 3.




Compound IN-19

Example 11

(R)-2-(4-carbamoylbenzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide (Compound IN-20))

[0230]



Step 1

[0231] To a solution of 5-bromo-1-tosyl-1H-indole (1 g, 2.866 mmol) in anhydrous toluene (10 mL) was added dropwise LDA (1.92 mL, 3.84 mmol, 2 M in THF/heptane/ethylbenzene) at -78 °C under N₂. The mixture was stirred at -78 °C for 30 min. A solution of 4-formylbenzonitrile (1.126 g, 8.598 mmol) in anhydrous THF (5 mL) was added dropwise to the mixture at -78 °C under N₂. The reaction mixture was stirred at -78 °C-15 °C for 16 h under N₂. TLC (petroleum ether / ethyl acetate = 5/1) showed that most of 5-bromo-1-tosyl-1H-indole was consumed. The mixture was quenched with sat. NH₄Cl solution (50 mL) and extracted with ethyl acetate (3 X 50 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by re-crystallization from petroleum ether / ethyl acetate (2/1, 50 mL) to afford 4-((5-bromo-1-tosyl-1H-indol-2-yl)(hydroxy)methyl)benzonitrile (1.20 g, 87%) as a white solid. LC-MS Method 2 *t_R* = 0.873 min in 5-95AB_1.5min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) *m/z* 462.9 [M-NH₃+H]⁺.

Step 2

[0232] To a solution of 4-((5-bromo-1-tosyl-1H-indol-2-yl)(hydroxy)methyl)benzonitrile (1.2 g, 2.5 mmol) in CH₂Cl₂ (50 mL) was added Et₃SiH (6 mL) and TFA (6 mL). The mixture was stirred at room temperature for 4 h. LCMS showed that the reaction was completed. The mixture was quenched with water (80 mL) and extracted with CH₂Cl₂ (3 X 80 mL). The

combined organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether / ethyl acetate = 1/0-4/1 to afford 4-((5-bromo-1-tosyl-1H-indol-2-yl)methyl)benzonitrile (1.06 g, 91%) as a white solid. **LC-MS** Method 2 t_R = 0.945 min in 5-95AB_1.5min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 464.9 $[\text{M}+\text{H}]^+$.

Step 3

[0233] To a solution of 4-((5-bromo-1-tosyl-1H-indol-2-yl)methyl)benzonitrile (1.4 g, 3 mmol) in MeOH (50 mL) and DMSO (10 mL) was added $\text{Pd}(\text{dppf})\text{Cl}_2$ (661 mg, 0.905 mmol) and Et_3N (606 mg, 6 mmol). The mixture was stirred at 70 °C for 16 h under CO (50 psi). TLC (petroleum ether / ethyl acetate = 2/1) showed that most of 4-((5-bromo-1-tosyl-1H-indol-2-yl)methyl)benzonitrile was consumed. The mixture was concentrated under reduced pressure. The residue was diluted with ethyl acetate (200 mL) and filtered. The filtrate was washed with brine (3 X 200 mL), dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether / ethyl acetate = 1/0-2/1 to afford methyl 2-(4-cyanobenzyl)-1-tosyl-1H-indole-5-carboxylate (1.03 g, 77%) as a yellow solid. **LC-MS** Method 2 t_R = 0.889 min in 5-95AB_1.5min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 444.9 $[\text{M}+\text{H}]^+$.

Step 4

[0234] To a solution of methyl 2-(4-cyanobenzyl)-1-tosyl-1H-indole-5-carboxylate (1.03 g, 2.32 mmol) in MeOH (50 mL) and THF (10 mL) was added KOH (2.6 g, 4.64 mmol) and water (10 mL). The reaction mixture was stirred at 70 °C for 48 h under N_2 . LCMS showed the reaction was completed. The mixture was concentrated under reduced pressure. The residue was adjusted to pH = 5-6 with 1 N HCl solution. A lot of solids was precipitate out and filtered. The filter cake was washed with petroleum ether / ethyl acetate (3/1, 30 mL) and dried under reduced pressure to afford crude 2-(4-carbamoylbenzyl)-1H-indole-5-carboxylic acid (0.8 g, >100%, contained some salt) as a yellow solid, which was used for the next step directly without further purification. **LC-MS** Method 2 t_R = 0.595 min in 5-95AB_1.5min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 294.9 $[\text{M}+\text{H}]^+$.

Step 5

[0235] To a solution of 2-(4-carbamoylbenzyl)-1H-indole-5-carboxylic acid (400 mg, 1.36 mmol) in EtOH (10 mL) was added conc. H_2SO_4 (1.5 mL). Then the mixture was stirred at 80 °C for 3 h. LCMS showed that the starting material was consumed completely. The mixture

was basified with aq. NaHCO_3 solution to pH = 8. The aqueous layer was extracted with ethyl acetate (3 X 20 mL). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether / ethyl acetate = 5/1-0/1 to afford ethyl 2-(4-carbamoylbenzyl)-1H-indole-5-carboxylate (110 mg, 25%) as a yellow solid. **LC-MS** Method 2 t_R = 0.724 min in 5-95AB_1.5min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 323.1 $[\text{M}+\text{H}]^+$.

Step 6

[0236] To a solution of ethyl 2-(4-carbamoylbenzyl)-1H-indole-5-carboxylate (110 mg, 0.342 mmol) in anhydrous THF (5 mL) was added KOH (58 mg, 1.026 mmol) and EtI (107 mg, 0.684 mmol). Then the mixture was stirred at 40 °C for 4 h. LCMS showed the reaction was completed. The mixture was adjusted to pH = 5-6 with 1 N HCl solution and diluted with water (10 mL). The aqueous layer was extracted with ethyl acetate (3 X 15 mL). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel eluting with petroleum ether / ethyl acetate = 5/1-0/1 to afford ethyl 2-(4-carbamoylbenzyl)-1-ethyl-1H-indole-5-carboxylate (95 mg, 79%) as a yellow solid. **LC-MS** Method 2 t_R = 0.783 min in 5-95AB_1.5min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 350.9 $[\text{M}+\text{H}]^+$.

Step 7

[0237] To a solution of ethyl 2-(4-carbamoylbenzyl)-1-ethyl-1H-indole-5-carboxylate (95 mg, 0.271 mmol) in EtOH (5 mL) and H_2O (1 mL) was added KOH (304 mg, 5.43 mmol). The mixture was stirred at 40 °C for 6 h under N_2 . LCMS showed that the reaction was completed. The mixture was concentrated under reduced pressure. The residue was diluted with water (10 mL) and adjusted to pH = 5-6 with 1 N HCl solution. The aqueous layer was extracted with ethyl acetate (3 X 20 mL). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated to afford crude 2-(4-carbamoylbenzyl)-1-ethyl-1H-indole-5-carboxylic acid (240 mg, >100%, contained some salt) as a yellow solid, which was used for next step directly without further purification. **LC-MS** Method 2 t_R = 0.674 min in 5-95AB_1.5min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 322.9 $[\text{M}+\text{H}]^+$.

Step 8

[0238] To a solution of crude 2-(4-carbamoylbenzyl)-1-ethyl-1H-indole-5-carboxylic acid (87 mg, 0.271 mmol) in CH_2Cl_2 (5 mL) was added (R)-2-amino-2-(5-(ethylsulfonyl)pyridin-2-

yl)ethanol HCl salt (108 mg, 0.4065 mmol), EDCI (104 mg, 0.542 mmol), HOBt (73 mg, 0.542 mmol) and Et₃N (55 mg, 0.542 mmol). The mixture was stirred at room temperature for 16 h under N₂. LCMS showed that the reaction was completed. The mixture was quenched with water (15 mL) and extracted with CH₂Cl₂ (3 X 15 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC with CH₂Cl₂ / MeOH = 10/1 to afford crude (R)-2-(4-carbamoylbenzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide (40 mg, 27%) as a yellow solid. The crude product (14 mg, 0.026 mmol) was purified by basic preparative HPLC separation and dry-freezing directly to give (R)-2-(4-carbamoylbenzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide (Compound IN-20, 4.0 mg, 29%) as a white solid. **LC-MS** Method 2 **t_R** = 0.675 min in 5-95AB_1.5min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 535.1 [M+H]⁺. **¹H NMR** (CDCl₃ 400 MHz): δ 9.05 (d, *J* = 2.4 Hz, 1H), 8.20 (dd, *J* = 8.4 Hz, 2.0 Hz, 1H), 8.10 (s, 1H), 7.80-7.67 (m, 5H), 7.35-7.28 (m, 3H), 6.35 (s, 1H), 6.15-5.95 (brs, 1H), 5.65-5.53 (brs, 1H), 5.49 (q, *J* = 4.0 Hz, 1H), 4.24 (dd, *J* = 4.0, 11.2 Hz, 1H), 4.20 (s, 2H), 4.11-4.00 (m, 3H), 3.17 (q, *J* = 7.2 Hz, 2H), 1.34 (t, *J* = 7.2 Hz, 3H), 1.19 (t, *J* = 7.2 Hz, 3H). **Isomer SFC** **t_R** = 2.504 min in 5 min chromatography (Column: AD-H; Method Name: AD-H_3UM_4_40_4ML_5MIN, ee = 99%).

Basic preparative HPLC method

[0239]

Mobile phase A: water with 0.05% NH₄OH

Mobile phase B: CH₃CN

Flow rate: 25 mL/min.

Detection: UV 220 nm / 254 nm

Column: Phenomenex Gemini 150*25mm*10um


Column temperature: 30 °C

Time in min	%A	%B
0.00	80	20
10.00	50	50
10.20	0	100
13.00	0	100

Example 12

(R)-2-(4-cyanobenzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide (Compound IN-21)

[0240]



[0241] To a solution of (R)-2-(4-carbamoylbenzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide (40 mg, 0.0748 mmol) in CH_2Cl_2 (5 mL) was added TFAA (31 mg, 0.1495 mmol) and Et_3N (23 mg, 0.2244 mmol). The mixture was stirred at room temperature for 1.5 h. LCMS showed that the desired MS was observed. The mixture was quenched with water (10 mL) and extracted with CH_2Cl_2 (3 X 10 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC with CH_2Cl_2 / acetone = 2/1 to give crude product (15 mg, 39%) as a pale yellow solid. The crude product (10 mg) was further purified by basic preparative HPLC separation and dry-freezing directly to afford (R)-2-(4-cyanobenzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide (Compound IN-21, 2.00 mg) as a white solid. **LC-MS** Method 2 t_R = 0.743 min in 5-95AB_1.5min chromatography (Welch Merck RP-18e 25-2mm), MS (ESI) m/z 517.0 $[\text{M}+\text{H}]^+$. **^1H NMR** (CDCl_3 400 MHz): δ 8.98 (s, 1H), 8.13 (d, J = 8.4 Hz, 1H), 8.03 (s, 1H), 7.71-7.60 (m, 3H), 7.55 (d, J = 8.0 Hz, 2H), 7.30-7.18 (m, 3H), 6.27 (s, 1H), 5.46-5.39 (m, 1H), 4.24-4.15 (m, 1H), 4.13 (s, 2H), 4.05-3.93 (m, 3H), 3.63-3.53 (m, 1H), 3.09 (q, J = 7.6 Hz, 2H), 1.27 (t, J = 7.2 Hz, 3H), 1.12 (t, J = 7.2 Hz, 3H). **Isomer SFC** t_R = 1.063 min in 5 min chromatography (Column: AD-H; Method Name: AD-H_3UM_4_40_4ML_5MIN.M, ee = 98%).

Basic preparative HPLC method

[0242]

Mobile phase A: water with 0.05% NH_4OH

Mobile phase B: CH_3CN

Flow rate: 25 mL/min.

Detection: UV 220 nm / 254 nm

Column: Phenomenex Gemini 150*25mm*10um

Step 5

[0244] A stirred mixture of methyl 1-tosyl-2-((5-(trifluoromethyl)pyridin-2-yl)methyl)-1H-indole-5-carboxylate (135 mg, 0.28 mmol), LiOH.H₂O (116 mg, 2.8 mmol) and 2:1:1 MeOH/THF/H₂O (4 mL) was heated at 35 °C for 5 d open to the atmosphere. The mixture was diluted with EtOAc (90 mL), washed with 2.5% aq HCl (10 mL) and brine (10 mL), and dried over Na₂SO₄. Removal of the solvent left 2-(4-(trifluoromethyl)benzoyl)-1H-indole-5-carboxylic acid (91 mg, 99%). LC-MS Method 1 *t_R* = 1.43 min, *m/z* = 335.

Step 6

[0245] To a stirred mixture of 2-(4-(trifluoromethyl)benzoyl)-1H-indole-5-carboxylic acid (91 mg, 0.27 mmol), Cs₂CO₃ (178 mg, 0.54 mmol) and dry DMF (1.5 mL) was added iodoethane (44 µL, 0.54 mmol). The mixture was stirred at rt for 4 h, diluted with EtOAc (90 mL), washed with 2.5% aq HCl (10 mL) and brine (10 mL), and dried over Na₂SO₄. Removal of the solvent left a yellow solid (117 mg). Chromatography on a 12 g silica cartridge, eluted with a 0-60% EtOAc in hexanes gradient, afforded ethyl 1-ethyl-2-(5-(trifluoromethyl)picolinoyl)-1H-indole-5-carboxylate (29 mg, 27%) as an oil which was used directly. LC-MS Method 1 *t_R* = 1.94 min, *m/z* = 391.

Step 7

[0246] Ethyl 1-ethyl-2-(5-(trifluoromethyl)picolinoyl)-1H-indole-5-carboxylate (29 mg, 74 µmol) is treated under conditions analogous to those in Step 5 to provide 1-ethyl-2-(5-(trifluoromethyl)picolinoyl)-1H-indole-5-carboxylic acid (20 mg, 74%) which was used directly in the next step. LC-MS Method 1 *t_R* = 1.61 min, *m/z* = 363.

Step 8


[0247] To a stirred solution of 1-ethyl-2-(5-(trifluoromethyl)picolinoyl)-1H-indole-5-carboxylic acid (20 mg, 55 µmol), (R)-2-amino-2-(4-(ethylsulfonyl)phenyl)ethan-1-ol hydrochloride (22 mg, 83 µmol), *i*-Pr₂NEt (52 µL, 0.29 mmol) and CH₂Cl₂ (1 mL) was added solid HATU (42 mg, 0.11 mmol). The mixture was stirred for 2 h and concentrated. The residue was purified by prep HPLC to afford (R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)picolinoyl)-1H-indole-5-carboxamide (Compound IN-22, 23 mg, 72%) as a yellow solid. LC-MS Method 1 *t_R* = 1.46 min, *m/z* = 574. ¹H NMR (CD₃OD, 400 MHz) δ 1.20 (t,

3H), 1.43 (t, 3H), 3.18 (q, 2H), 3.92 (d, 2H), 4.73 (q, 2H), 5.31 (t, 1H), 7.63 (d, 1H), 7.66-7.74 (m, 3H), 7.87 (d, 2H), 7.93 (d, 1H), 8.31-8.39 (m, 2H), 9.04 (s, 1H).

Example 14 (reference example)

1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(hydroxy(5-(trifluoromethyl)pyridin-2-yl)methyl)-1H-indole-5-carboxamide (Compound IN-23)

[0248]



[0249] To a stirred solution of (R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)picolinoyl)-1H-indole-5-carboxamide (19 mg, 33 μ mol) in MeOH (1 mL) was added powdered NaBH₄ (~5 mg). The mixture was stirred for 1 h, and diluted with water (0.5 mL) and 5% aq HCl (0.5 mL). The solution was stirred for 15 min and purified directly by prep HPLC to afford the title compound (12 mg, 63%) as a solid. LC-MS Method 1 t_R = 1.26 min, m/z = 576. ¹H NMR (CD₃OD, 400 MHz) δ 1.20 (t, 3H), 1.28 (t, 3H), 2.17 (q, 2H), 3.91 (d, 2H), 4.38 (m, 2H), 5.17 (t, 1H), 6.16 (s, 1H), 6.19 (s, 1H), 7.45 (s, 2H), 7.65-7.78 (m, 3H), 7.89 (d, 2H), 7.97 (d, 1H), 8.09 (s, 1H), 8.13 (d, 1H), 8.84 (s, 1H).

LC-MS Data

Cpd No	Cpd Name	LC-MS Method	t_R (min)	Mass observed
IN-1	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxamide	2	0.82	529
IN-2	1-ethyl-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxamide	2	0.788	530.1
IN-3	(R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxamide	2	0.787	559.0
	(R)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-			

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
IN-4.1 and IN-4.2	yl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxamide	2	0.926	560.1
	(S)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxamide	2	0.774	560.1
IN-5 ***	(R)-2-((2,2-difluorobenzo[d][1,3]dioxol-5-yl)methyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide	2	0.891	572.1
IN-6	(R)-2-(4-(difluoromethoxy)benzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide	2	0.746	558.1
IN-7	(R)-2-(3-(difluoromethoxy)benzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide	4	1.130	558.0
IN-8	(R)-2-(4-chlorobenzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide	2	0.851	526.1
IN-9	(R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((5-(trifluoromethyl)pyridin-2-yl)methyl)-1H-indole-5-carboxamide	1	1.42	560
IN-10	(R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole-5-carboxamide	1	0.82	566
IN-11	(R)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole-5-carboxamide	1	0.79	567
IN-12	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole-5-carboxamide	1	0.91	548
IN-13	(R)-1-cyclopropyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-indole-5-carboxamide	1	0.84	578
	1-ethyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-			

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
IN-14.1 and IN-14.2	yl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide	2	0.785	566.0
	1-ethyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((1s,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide	2	0.795	566.1
IN-15.1 and IN-15.2	1-ethyl-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide	2	0.808	536.1
	1-ethyl-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-2-(((1s,4s)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide	2	0.800	536.1
IN-16.1 and IN-16.2	1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide	2	0.801	565.0
	1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1s,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide	2	0.798	565.1
IN-17.1 and IN-17.2	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide	2	0.838	535.0
	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1s,4s)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-indole-5-carboxamide	2	0.837	535.1
IN-19	(R)-1-cyclopropyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)benzyl)-1H-indole-5-carboxamide	2	0.783	572.2
IN-20	(R)-2-(4-carbamoylbenzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide	2	0.675	535.1
IN-21	(R)-2-(4-cyanobenzyl)-1-ethyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-1H-indole-5-carboxamide	2	0.743	517.0
IN-22 ***	(R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(5-(trifluoromethyl)picolinoyl)-1H-indole-5-	1	1.46	574

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
	carboxamide			
IN-23 ***	1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(hydroxy(5-(trifluoromethyl)pyridin-2-yl)methyl)-1H-indole-5-carboxamide	1	1.27	576
AI-1***	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-pyrrolo[2,3-b]pyridine-5-carboxamide	1	1.65	548
AI-2***	1-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-pyrrolo[2,3-b]pyridine-5-carboxamide	1	1.54	578
AI-3***	N-(4-(ethylsulfonyl)benzyl)-1-(2-methylallyl)-2-(4-(trifluoromethyl)benzyl)-1H-pyrrolo[2,3-b]pyridine-5-carboxamide	1	1.70	556
BE-1	1-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)oxy)-1H-benzo[d]imidazole-5-carboxamide	1	1.46	580.51 (M + 1)
BE-2	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)oxy)-1H-benzo[d]imidazole-5-carboxamide	1	1.51	538.47 (M + 1)
BE-3	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)oxy)-1H-benzo[d]imidazole-5-carboxamide	1	1.54	550.40 (M + 1)
BE-4.1 and BE-4.2	N-(4-(ethylsulfonyl)benzyl)-1-isobutyl-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)oxy)-1H-benzo[d]imidazole-5-carboxamide	1	1.68	566.38 (M + 1)
	N-(4-(ethylsulfonyl)benzyl)-1-isobutyl-2-(((1s,4s)-4-(trifluoromethyl)cyclohexyl)oxy)-1H-benzo[d]imidazole-5-carboxamide	1	1.64	566.34 (M + 1)
BE-5	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)phenoxy)-1H-benzo[d]imidazole-5-carboxamide	1	1.54	532.40 (M + 1)
BE-6	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)phenoxy)-1H-benzo[d]imidazole-5-carboxamide	1	1.57	544.46 (M + 1)

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
BE-7	(R)-1-cyclopropyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)phenoxy)-1H-benzo[d]imidazole-5-carboxamide	1	1.47	574.45 (M + 1)
BE-8	(R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)phenoxy)-1H-benzo[d]imidazole-5-carboxamide	1	1.44	562.53 (M + 1)
BE-9	(R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(4-(trifluoromethoxy)phenoxy)-1H-benzo[d]imidazole-5-carboxamide	1	1.49	578.47 (M + 1)
BE-10	(R)-N-(2-(dimethylamino)-1-(4-(ethylsulfonyl)phenyl)ethyl)-1-ethyl-2-(4-(trifluoromethyl)phenoxy)-1H-benzo[d]imidazole-5-carboxamide	2	0.82	589
BE-11	(R)-N-(2-amino-1-(4-(ethylsulfonyl)phenyl)ethyl)-1-ethyl-2-(4-(trifluoromethyl)phenoxy)-1H-benzo[d]imidazole-5-carboxamide	2	0.80	561
BE-12 ***	(R)-2-((4,4-difluorocyclohexyl)methoxy)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.83	550
BE-13 ***	(R)-2-((3,3-difluorocyclobutyl)methoxy)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.68	522, 544
BE-14 ***	(R)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((4-(trifluoromethyl)benzyl)oxy)-1H-benzo[d]imidazole-5-carboxamide	2	0.88	576
BE-15 ***	(R)-1-cyclopropyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((4-(trifluoromethyl)benzyl)oxy)-1H-benzo[d]imidazole-5-carboxamide	2	0.76	588
BE-16 ***	(R)-1-cyclopropyl-2-((4,4-difluorocyclohexyl)methoxy)-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.71	562
BE-17 ***	(R)-2-(cyclopropylmethoxy)-1-ethyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-1H-benzo[d]imidazole-5-carboxamide	3	1.70	472

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
BE-18.1 and BE-18.2	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.81	548
	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.80	548
BE-19.1 and BE-19.2	N-(4-(ethylsulfonyl)benzyl)-1-isobutyl-2-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.72	564
	N-(4-(ethylsulfonyl)benzyl)-1-isobutyl-2-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.72	564
BE-20	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.79	536
BE-21	1-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.68	578
BE-22	1-cyclopropyl-N-((S)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.67	578
BE-23	1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.66	566, 588
BE-24	1-ethyl-N-((S)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.65	566, 588
BE-25	1-cyclopropyl-N-((S)-1-(4-(ethylsulfonyl)phenyl)-3-hydroxypropyl)-2-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	4	0.94	592
	N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-1-			

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
BE-26.1 and BE-26.2	isobutyl-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.69	565
	N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-1-isobutyl-2-(((1s,4s)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.69	565
BE-27.1 and BE-27.2	1-cyclopropyl-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.65	549
	1-cyclopropyl-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-2-(((1s,4s)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.65	549
BE-28	1-cyclopropyl-N-((R)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.65	579
BE-29	1-cyclopropyl-N-((S)-1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(((1r,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.65	579
BE-30	1-cyclopropyl-N-(4-(methylsulfonyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.65	534
BE-31 ***	1-cyclopropyl-N-((5-(N-methylsulfamoyl)pyridin-2-yl)methyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	4	0.92	550
BE-32	1-cyclopropyl-N-((R)-2-hydroxy-1-(4-(methylsulfonyl)phenyl)ethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.66	564
BE-33 ***	1-cyclopropyl-N-(4-(N-methylsulfamoyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.69	549
BE-34	N-(4-(ethylsulfonyl)benzyl)-1-methyl-2-(4-(trifluoromethyl)benzyl)-1H-	2	0.67	516

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
	benzo[d]imidazole-5-carboxamide			
BE-35	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.70	530
BE-36	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.69	542
BE-37	N-(4-(ethylsulfonyl)benzyl)-1-(2,2,2-trifluoroethyl)-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.79	584
BE-38	1-cyclobutyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.70	556
BE-39 ***	N-(4-(ethylsulfonyl)benzyl)-1-neopentyl-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.75	572
BE-40	N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-1-isopropyl-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.76	545
BE-41	N-(4-(ethylsulfonyl)benzyl)-1-isobutyl-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.70	558
BE-42	1-(tert-butyl)-N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.78	559
BE-43	N-((5-(ethylsulfonyl)pyridin-2-yl)methyl)-1-isobutyl-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.82	559
BE-44	rel-(R)-N-(1-(4-(ethylsulfonyl)phenyl)ethyl)-1-isobutyl-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.73	572
BE-45	rel-(R)-N-(1-(4-(ethylsulfonyl)phenyl)ethyl)-1-isobutyl-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.74	572
BE-46.1 and BE-46.2	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4r)-4-methoxycyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	3	1.72	510
	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1s,4s)-4-methoxycyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.60	510
	(R)-1-cyclopropyl-N-(1-(4-			

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
BE-47	(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(4-(trifluoromethoxy)benzyl)-1H-benzo[d]imidazole-5-carboxamide	1	1.24	588.49 (M+1)
BE-48	1-cyclopropyl-2-(4-(difluoromethoxy)benzyl)-N-(4-(ethylsulfonyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.70	540, 562
BE-49	tert-butyl 4-((1-cyclopropyl-5-((4-(ethylsulfonyl)benzyl)carbamoyl)-1H-benzo[d]imidazol-2-yl)methyl)piperidine-1-carboxylate	1	1.09	581.59 (M + 1)
BE-50	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-((1-(2,2,2-trifluoroethyl)piperidin-4-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide	1	0.87	563.53 (M + 1)
BE-51	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-((6-(trifluoromethyl)pyridin-3-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide	1	1.18	543.47 (M + 1)
BE-52	2-((5-chloropyridin-2-yl)methyl)-1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.66	509, 511
BE-53	N-(4-(ethylsulfonyl)benzyl)-1-isobutyl-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide	1	1.04	565.38 (M + 1)
BE-54	(R)-1-cyclopropyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide	1	0.82	579.54 (M + 1)
BE-55.1 and BE-55.2 ***	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(1-(4-(trifluoromethyl)phenyl)ethyl)-1H-benzo[d]imidazole-5-carboxamide isomer 1	2	0.73	556
	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(1-(4-(trifluoromethyl)phenyl)ethyl)-1H-benzo[d]imidazole-5-carboxamide isomer 2	2	0.73	556
	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-((6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide isomer 1	2	0.64	538, 560
	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-((6-			

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
BE-56.1, BE-56.2, BE-56.3 and BE-56.4	(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide isomer 2	2	0.64	538, 560
	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-((6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide isomer 3	2	0.64	538, 560
	1-ethyl-N-(4-(ethylsulfonyl)benzyl)-2-((6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide isomer 4	2	0.75	538, 560
BE-57.1, BE-57.2, BE-57.3 and BE-57.4	1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide isomer 1	2	0.73	568
	1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide isomer 2	2	0.73	568
	1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide isomer 3	2	0.64	568, 590
	1-ethyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-((6-(trifluoromethyl)tetrahydro-2H-pyran-3-yl)methyl)-1H-benzo[d]imidazole-5-carboxamide isomer 4	2	0.63	568, 590
BE-58 ***	7-bromo-N-(4-(ethylsulfonyl)benzyl)-1-methyl-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide	1	1.57	600, 602
BE-59 ***	7-bromo-N-(4-(ethylsulfonyl)benzyl)-1-methyl-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	1	1.65	594, 596
BE-60 ***	N-(4-(ethylsulfonyl)benzyl)-7-methoxy-1-methyl-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazole-5-carboxamide	1	1.47	546
BE-61 ***	(R)-1-cyclopropyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)benzoyl)-1H-	1	1.50	586.54 (M + 1)

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
	benzo[d]imidazole-5-carboxamide			
BE-62 ***	(R)-1-cyclopropyl-N-(1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)piperidine-1-carbonyl)-1H-benzo [d]imidazole-5-carboxamide	2	0.66	593
BE-63.1 and BE-63.2 ***	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(hydroxy(trans-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide isomer 1	2	0.67	564, 586
	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(hydroxy(trans-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamide isomer 2	2	0.67	564, 586
BE-64 ***	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)cyclohexane-1-carbonyl)-1H-benzo[d]imidazole-5-carboxamide	2	0.80	562,584
BE-65 ***	1-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzoyl)-1H-benzo[d]imidazole-5-carboxamide	1	1.59	556.50 (M + 1)
BE-66	(R)-2-(1-cyclopropyl-2-(((1R,4R)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamido)-2-(4-(ethylsulfonyl)phenyl)ethyl carbamate	2	0.66	621
BE-67	(S)-2-(1-cyclopropyl-2-(((1R,4S)-4-(trifluoromethyl)cyclohexyl)methyl)-1H-benzo[d]imidazole-5-carboxamido)-2-(4-(ethylsulfonyl)phenyl)ethyl carbamate	2	0.66	621
BE-68 ***	2-(4-(ethylsulfonyl)phenyl)-N-(1-isobutyl-2-(4-(trifluoromethyl)benzyl)-1H-benzo[d]imidazol-5-yl)acetamide	1	1.22	558.30 (M + 1)
BE-69 ***	2-(4-(ethylsulfonyl)phenyl)-N-(1-isobutyl-2-(((4-(trifluoromethyl)piperidin-1-yl)methyl)-1H-benzo[d]imidazol-5-yl)acetamide	1	1.11	565.36 (M + 1)
BE-70	(R)-1-cyclopropyl-N-(1-(5-(ethylsulfonyl)pyridin-2-yl)-2-hydroxyethyl)-2-(4-(trifluoromethyl)phenoxy)-1H-benzo [d]imidazole-5 -carboxamide	2	0.75	575
	1-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1S,4S)-4-	2	0.81	594

Cpd No	Cpd Name	LC-MS Method	t _R (min)	Mass observed
BE-71.1 and BE-71.2 ***	(trifluoromethyl)cyclohexyloxy)methyl)-1H-benzo [d]imidazole-5 -carboxamide			
	1-cyclopropyl-N-((R)-1-(4-(ethylsulfonyl)phenyl)-2-hydroxyethyl)-2-(((1r,4R)-4-(trifluoromethyl)cyclohexyl)oxy)methyl)1H-benzo[d]imidazole-5-carboxamide	2	0.8	594
AB-1.1 ***	3-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1r,4r)-4-(trifluoromethyl)cyclohexyl)methyl)-3H-imidazo[4,5-b]pyridine-6-carboxamide	2	0.85	549
AB-1.2 ***	3-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(((1s,4s)-4-(trifluoromethyl)cyclohexyl)methyl)-3H-imidazo[4,5-b]pyridine-6-carboxamide	2	0.85	549
AB-2 ***	3-cyclopropyl-N-(4-(methylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzyl)-3H-imidazo[4,5-b]pyridine-6-carboxamide	1	1.35	529
AB-3 ***	3-cyclopropyl-N-(4-(ethylsulfonyl)benzyl)-2-(4-(trifluoromethyl)benzyl)-3H-imidazo[4,5-b]pyridine-6-carboxamide	1	1.42	543
*** Reference example				

BIOLOGICAL ASSAYS

Radio-Ligand ROR γ Binding Assay (Assay 1)

[0250] Compounds disclosed herein were tested for ability to bind to ROR γ in a cell-free competition assay with commercially available radio-ligand (RL), 25-hydroxy [26,27-³H]-cholesterol (PerkinElmer, Cat. # NET674250UC), for a ligand binding site on a recombinant ROR γ Ligand Binding Domain (LBD) protein expressed as a 6xHis-Glutathione-S-Transferase (GST) fusion. The assay was performed in 96-well SPA plates (PerkinElmer, Cat. # 1450-401) in 50 mM HEPES buffer, pH 7.4, containing 150 mM NaCl, 5 mM MgCl₂, 10% (v/v) glycerol, 2 mM CHAPS, 0.5 mM β -octylglucopyranoside and 5 mM DTT. Tested compounds were dissolved in DMSO, and semi-log (3.162x) serial dilutions of the compounds were prepared in the same solvent. Two μ L of the DMSO solutions were mixed with 28 μ L of 8.6 nM 25-hydroxy [26,27-³H]- cholesterol and 50 μ L of 24 nM ROR γ LBD. The plate was shaken at 700 rpm for 20 min and incubated for 10 min at rt, after which 40 μ L of poly-Lys YSi SPA beads

(PerkinElmer, Cat. # RPNQ0010) were added to achieve 50 µg of the beads per well. The plate was incubated on an orbital shaker for 20 min and then for 10 min without agitation at rt. SPA signal for tritium beta radiation was registered on PerkinElmer Microbeta plate reader. Percent inhibition values were calculated based on the high signal obtained with DMSO control and the low signal observed with 10 µM standard RORγ inverse agonist T0901317 (SigmaAldrich, Cat. # T2320). The percent inhibition vs. concentration data were fit into a four-parameter model, and IC₅₀ values were calculated from the fit as the concentrations corresponding to the inflection points on the dose-response curves. Inhibitory constants (K_i) were calculated using the following equation, where [RL] is the concentration in the assay and K_D is a dissociation constant of 25-hydroxy [26,27-³H]- cholesterol:

$$K_i = \frac{IC_{50}}{\left(1 + \frac{[RL]}{K_D}\right)}$$

RORγt 5xRORE Assay in Jurkat Cells (Assay 2)

[0251] Compounds disclosed herein were tested for RORγ inverse agonist activity in a cell-based, transcriptional activity assay. Secreted Nanoluc® luciferase was used as a reporter for transcriptional activity of the full-length RORγt in Jurkat cells (ATCC, Cat. # TIB-152). A reporter plasmid was constructed by inserting 5 repeats of the ROR Response Element (RORE) AAAGTAGGTCA (SEQ ID NO:1) into a commercially available promoterless plasmid pNL1.3[secNluc] (Promega, Cat. # N1021) using KpnI and HindIII restriction sites. The expression plasmid for RORγt was purchased (GeneCopoeia, Cat. # EX-T6988-M02). Jurkat cells (30 million cells) were transfected with 11 µg of EX-T6988-MO2 and 26 µg of the reporter plasmid in OptiMEM® media using Lipofectamine® LTX and Plus™ reagents (Life Technologies, Cat. # 15338-100). After 5-6 hrs of incubation at 37 °C/5% CO₂, the cells were collected, resuspended in phenol-red free RPMI media containing 10% (v/v) delipidated FBS (Hyclone, Cat. # SH30855.03) and dispensed into 96-well clear bottom tissue culture plates (CoStar, Cat. # 3603), at 80,000 cells per well. Tested compounds were added to the cells in the same media (final concentration of DMSO was 0.1% (v/v)), and the plates were incubated at 37 °C/5% CO₂ for 16-18 hrs. Luciferase activity in the conditioned supernatants was determined with NanoGlo® assay reagents (Promega, Cat.# N1130). Percent inhibition values were calculated based on the fully inhibited and non-inhibited (DMSO) controls, and the values were regressed against concentrations of the tested compounds to derive IC₅₀ values using a four-parameter non-linear fitting model.

[0252] The results of assays 1 and 2 are shown in **Table 1**.

Table 1

Cpd No	K _i (nM) (Assay 1)	IC ₅₀ (nM) (Assay 2)
IN-1	+++	+++
IN-2	+++	+++

Cpd No	Ki (nM) (Assay 1)	IC50 (nM) (Assay 2)
IN-3	+++	+++
IN-4.1	+++	+++
IN-4.2	+++	++
IN-5 ***	+++	+++
IN-6	+++	++
IN-7	+++	++
IN-8	+++	++
IN-9	+++	+++
IN-10	+++	+++
IN-11	+++	+++
IN-12	+++	+++
IN-13	+++	+++
IN-14.1	+++	+++
IN-14.2	+++	++
IN-15.1	+++	+++
IN-15.2	+++	+
IN-16.1	+++	+++
IN-16.2	+++	+
IN-17.1	+++	+++
IN-17.2	+++	+
IN-18	+++	+++
IN-19	+++	+++
IN-20	++	nt
IN-21	+++	++
IN-22 ***	+++	+++
IN-23 ***	++	nt
AI-1 ***	+++	+++
AI-2 ***	+++	+++
AI-3 ***	+++	++
BE-1	+++	+++
BE-2	+++	+++
BE-3	+++	++
BE-4.1	+++	++
BE-4.2	+++	+
BE-5	+++	+++

Cpd No	Ki (nM) (Assay 1)	IC50 (nM) (Assay 2)
BE-6	+++	+++
BE-7	+++	+++
BE-8	+++	+++
BE-9	+++	+++
BE-10	++	nt
BE-11	+++	+++
BE-12 ***	+++	+
BE-13 ***	+++	+
BE-14 ***	+++	+++
BE-15 ***	+++	+++
BE-16 ***	+++	+++
BE-17 ***	+++	+
BE-18.1	+++	+++
BE-18.2	+++	+
BE-19.1	+++	+++
BE-19.2	+++	+
BE-20	+++	+++
BE-21	+++	+++
BE-22	+++	++
BE-23	+++	+++
BE-24	++	nt
BE-25	+++	+++
BE-26.1	+++	+
BE-26.2	+++	+
BE-27.1	+++	+++
BE-27.2	+++	+
BE-28	+++	+++
BE-29	+	nt
BE-30	+++	+++
BE-31 ***	+++	+
BE-32	+++	+++
BE-33 ***	+++	+++
BE-34	++	++
BE-35	+++	+++

Cpd No	Ki (nM) (Assay 1)	IC50 (nM) (Assay 2)
BE-36	+++	+++
BE-37	+++	+++
BE-38	+++	++
BE-39 ***	+++	++
BE-40	++	nt
BE-41	+++	++
BE-42	++	nt
BE-43	+++	++
BE-44	++	nt
BE-45	+++	++
BE-46.1	+++	++
BE-46.2	+++	+
BE-47	+++	+++
BE-48	+++	+++
BE-49	+++	++
BE-50	++	++
BE-51	++	nt
BE-52	+++	+
BE-53	+++	+++
BE-54	+++	+++
BE-55.1 ***	++	+
BE-55.2 ***	+++	+++
BE-56.1	++	nt
BE-56.2	++	nt
BE-56.3	++	nt
BE-56.4	++	nt
BE-57.1	++	++
BE-57.2	+++	++
BE-57.3	++	nt
BE-57.4	++	nt
BE-58 ***	++	nt
BE-59 ***	++	nt
BE-60 ***	++	nt
BE-61 ***	+++	+++
BE-62 ***	++	nt

Cpd No	Ki (nM) (Assay 1)	IC50 (nM) (Assay 2)
BE-63.1 ***	+	nt
BE-63.2 ***	++	nt
BE-64 ***	+++	++
BE-65 ***	+++	+++
BE-66	+++	+++
BE-67	+++	+++
BE-68 ***	+++	++
BE-69 ***	++	++
BE-70	+++	+++
BE-71.1 ***	+++	+++
BE-71.2 ***	+++	+++
AB-1.1 ***	+++	+++
AB-1.2 ***	+++	+
AB-2 ***	++	nt
AB-3 ***	+++	+++
nt = not tested; + means > 1000 nM; ++ means 100 nM - 1000 nM; +++ means < 100 nM. *** Reference example		

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2014179564A \[0076\] \[0077\] \[0087\] \[0087\] \[0088\] \[0089\]](#)
- [WO2015116904A \[0076\] \[0077\] \[0087\] \[0087\] \[0088\] \[0089\]](#)
- [WO2016061160A \[0076\] \[0077\] \[0087\] \[0087\] \[0088\] \[0089\]](#)
- [US933468 \[0076\] \[0077\]](#)
- [US14933524B \[0076\] \[0077\]](#)
- [US20160122318A \[0087\] \[0087\] \[0088\] \[0089\]](#)

- US20160122345A [0087] [0087] [0088] [0089]


Non-patent literature cited in the description

- JETTENJOO Adv. Dev. Biol. 2006, 2006, vol. 16, 313-355 [0002]
- JETTEN Nucl. Recept. Signal., 2009, vol. 7, e003- [0002]
- MADDUR et al. Am. J. Pathol., 2012, vol. 181, 8-18 [0003]
- THIBOUTOT et al. J. Invest. Dermatol., 2014, vol. 134, 2307-10 [0003]
- AGAK et al. J. Invest. Dermatol., 2014, vol. 134, 2366-73 [0003]
- HIRATA et al. Endocrinol., 2010, vol. 151, 5468-5476 [0003]
- HIRATA et al. Fertil Steril., vol. 96, 1113-7 [0003]
- BEDOYA et al. Clin. Dev. Immunol., 2013, 986789- [0003]
- MIOSECKOLLS Nature Rev., 2012, vol. 11, 763-776 [0003]
- KORN et al. Annu. Rev. Immunol., 2009, vol. 27, 485-517 [0003]
- SOLT et al. ACS Chem. Biol., 2012, vol. 7, 1515-1519 [0003]
- IVANOV et al. Cell, 2006, vol. 126, 1121-1133 [0003]
- TAYLOR et al. J. Immunol., 2014, vol. 192, 3319-3327 [0004]
- TAYLOR et al. Nat. Immunol., 2014, vol. 15, 143-151 [0004]
- JETTEN op. cit., 2009, [0005]
- YE et al. Mediators Inflamm., 2012, 815308- [0006]
- TOSOLINI et al. Cancer Res., 2011, vol. 71, 1263-1271 [0007]
- SU et al. Immunol. Res., 2014, vol. 58, 118-124 [0007]
- CARMI et al. J. Immunol., 2011, vol. 186, 3462-3471 [0007]
- CHEN et al. Histopathology, 2013, vol. 63, 225-233 [0007]
- Nature Medicine, 2016, [0007]
- MEISSBURGER et al. EMBO Mol. Med., 2011, vol. 3, 637-651 [0008]
- CHANG et al. J. Exp. Pharmacol., 2012, vol. 4, 141-148 [0009]
- JETTEN et al. Frontiers Endocrinol., 2013, vol. 4, 1-8 [0009]
- HUHLITTMAN Eur. J. Immunol., 2012, vol. 42, 2232-2237 [0009]
- MARTINEZ et al. Ann. N.Y. Acad. Sci., 2008, vol. 1143, 188-211 [0009]
- PANTELYUSHIN et al. J. Clin. Invest., 2012, vol. 122, 2252-2256 [0009]
- JETTENUEDA Cell Death Differen., 2002, vol. 9, 1167-1171 [0009]
- SOLT et al. Curr. Opin. Lipidol., 2010, vol. 21, 204-211 [0009]
- MARCH Advanced Organic Chemistry John Wiley & Sons 19850000 [0098]
- GREENEWUTS Protective Groups in Organic Synthesis John Wiley & Sons 19910000 [0098]
- RICHARD LAROCK Comprehensive Organic Transformations VCH publishers Inc. 19890000 [0098]
- XU, Y.LIN, H.RUAN, X.YANG, S.HAO, G.YANG, W.YANG, G. Eur. J. Med. Chem., 2015, vol. 92, 427-438 [0163]

MODULATORER AF ROR-GAMMA


Patentkrav

1. En forbindelse med formlen IV eller V:



(IV);

eller



(V);

eller et farmakologisk acceptabelt salt deraf, hvori

A er CH eller N;

Y er CH

Cy¹ er fenyl eller pyridyl, hver erstattet med 1 til 3 grupper valgt uafhængigt fra R⁵, hvori mindst

én R⁵ er -SO₂-(C₁-C₃)alkyl

Cy² er valgt fra fenyl, pyridinyl, piperidinyl, cyclohexyl, tetrahydropyranyl, cyclopropyl og cyclobutyl,

hver eventuelt erstattet med 1 til 3 grupper valgt uafhængigt fra R⁶

R² er (C₁-C₄)alkyl, (C₂-C₄)alkenyl, (C₁-C₄)haloalkyl eller monocyklisk cycloalkyl;

R⁵ og R⁶ er hver valgt uafhængigt fra halogen, -CN, -ORᶜ, -NRᵈRᵉ, -S(O)ₖRᶜ, -NRᶜS(O)₂Rᶜ,

-S(O)₂NRᵈRᵉ, -C(=O)ORᶜ, -OC(=O)ORᶜ, -OC(=O)Rᶜ, -OC(=S)ORᶜ, -C(=S)ORᶜ, -OC(=S)Rᶜ, -C(=O)NRᵈRᵉ,

-NRᶜC(=O)Rᶜ, -C(=S)NRᵈRᵉ, -NRᶜC(=S)Rᶜ, -NRᶜC(=O)ORᶜ, -OC(=O)NRᵈRᵉ, -

NRᶜ(C=S)ORᶜ, -OC(=S)NRᵈRᵉ,

-NRᶜC(=O)NRᵈRᵉ, -NRᶜ(C=S)NRᵈRᵉ, -C(=S)Rᶜ, -C(=O)Rᶜ, (C₁-C₆)alkyl, cycloalkyl, -(CH₂)₁-

4-cycloalkyl, heterocyclyl, -(CH₂)₁-4-heterocyclyl, aryl, -NHC(=O)-heterocyclyl, -NHC(=O)-

cycloalkyl, -(CH₂)₁-4-aryl, heteroaryl

og -(CH₂)₁-4-heteroaryl,

hvori den portion af alkyl, cycloalkyl, heterocyclyl, aryl eller heteroaryl, som findes i hver af

nævnte (C₁-C₆)alkyl, cycloalkyl, -(CH₂)₁-4-cycloalkyl, heterocyclyl, -(CH₂)₁-4-heterocyclyl,

aryl, -(CH₂)₁-4-aryl, heteroaryl og

-(CH₂)₁-4-heteroaryl substituent for R⁶ er yderligere eventuelt erstattet med halogen, ORᶜ, -NO₂, -CN,

-NR^cC(=O)R^c, -NR^dR^e, -S(O)_kR^c, -C(=O)OR^c, -C(=O)NR^dR^e, -C(=O)R^c, (C₁-C₃)alkyl, halo(C₁-C₃)alkyl,

(C₁-C₃)alkoxy(C₁-C₃)alkyl, (C₁-C₃)alkoxy eller halo(C₁-C₃)alkoxy;

hver R^c er valgt uafhængigt fra hydrogen og (C₁-C₆)alkyl eventuelt erstattet med 1 til 3

halogen;

hver R^d og R^e er valgt uafhængigt fra hydrogen og (C₁-C₆)alkyl;

k er 0, 1 eller 2;

enhver heterocyclyl- eller heteroaryl-del af Cy¹ eller Cy² er yderligere eventuelt erstattet med =O; og

R⁷ er hydrogen, (C₁-C₂)alkyl, hydroxy(C₁-C₃)alkyl, -(C₁-C₃)alkylNH₂, -(C₁-C₃)alkyl-N-di(C₁-C₃)alkyl eller


-(C₁-C₃)alkoxy-C(O)-NH₂,

såfremt, at:


når A er N, Y is CH, R² er metyl, Cy¹ er fenyl og forbindelsen har formel IV, så er Cy² ikke fenyl

erstattet med CN.

2. Forbindelsen i krav 1, hvori forbindelsen har formlen VI eller VII:




eller




eller et farmakologisk acceptabelt salt deraf.

3. Forbindelsen i krav 1 eller 2, hvori R⁷ er hydroxy(C₁-C₃)alkyl.

4. Forbindelsen i ethvert af krav 1 til 3, hvori forbindelsen har formlen VIII eller IX:



eller



eller et farmakologisk acceptabelt salt deraf.

5. Forbindelsen i ethvert af krav 1 til 4, hvori R^2 er (C_1-C_4) alkyl eller cyclopropyl.

6. Forbindelsen i ethvert af kravene 1 til 5, hvori Cy^1 er pyridinyl erstattet med 1 til 3 grupper valgt uafhængigt fra R^5 , hvori mindst én R^5 er $-SO_2-(C_1-C_3)$ alkyl.

7. Forbindelsen i ethvert af krav 1 til 6, hvori Cy^2 er valgt fra fenyl og cyclohexyl, hver eventuelt erstattet med 1 til 3 grupper valgt uafhængigt fra R^6 .

8. Forbindelsen i ethvert af kravene 1 til 5 og 7, hvori Cy^1 er fenyl erstattet med 1 til 3 grupper valgt uafhængigt fra R^5 , hvori mindst én R^5 er $-SO_2-(C_1-C_3)$ alkyl; og Cy^2 er cyclohexyl erstattet med 1 til 3 grupper valgt uafhængigt fra R^6 .

9. Forbindelsen i ethvert af kravene 1-8, hvori

R^5 er valgt fra halogen, $-CN$, $-OR^c$, $-NR^dR^e$, $-NR^cS(O)_2R^c$, $-S(O)_2NR^dR^e$, $-C(=O)OR^c$, $-C(=O)NR^dR^e$,

$-NR^cC(=O)R^c$, $-NR^cC(=O)OR^c$, $-OC(=S)NR^dR^e$, $-C(=O)R^c$, $-SO_2-(C_1-C_3)$ alkyl og (C_1-C_4) alkyl eventuelt erstattet med halogen; og

R^6 er valgt fra halogen, $-CN$, $-OR^c$, $-NR^dR^e$, $-NR^cS(O)_2R^c$, $-S(O)_2NR^dR^e$, $-C(=O)OR^c$, $-OC(=O)OR^c$,

$-OC(=O)R^c$, $-C(=O)NR^dR^e$, $-NR^cC(=O)R^c$, $-C(=S)NR^dR^e$, $-NR^cC(=S)R^c$, $-NR^cC(=O)OR^c$, $-OC(=O)NR^dR^e$,

$-NR^c(C=S)OR^c$, $-OC(=S)NR^dR^e$, $-NR^cC(=O)NR^dR^e$, $-NR^c(C=S)NR^dR^e$, $-C(=S)R^c$, $-C(=O)R^c$, $-SO_2-(C_1-C_3)$ alkyl,

and (C_1-C_4) alkyl eventuelt erstattet med halogen.


10. Forbindelsen i ethvert af kravene 1-9, hvori

R^5 er $-SO_2-(C_1-C_3)$ alkyl;

R^6 er valgt fra halo, $-OR^c$, og (C_1-C_4) alkyl eventuelt erstattet med halogen; og R^c er (C_1-C_4) alkyl eventuelt erstattet med halogen.


11. Forbindelsen i ethvert af kravene 1 til 10, hvori R^6 er CF_3 .

12. Forbindelsen i krav 1, hvori forbindelsen har formelen:




eller et farmakologisk acceptabelt salt deraf.

13. Forbindelsen i krav 1, hvori forbindelsen har formelen:



eller et farmakologisk acceptabelt salt deraf.

14. Forbindelsen i krav 1, hvori forbindelsen har formelen:



eller et farmakologisk acceptabelt salt deraf.

15. En farmakologisk sammensætning, der omfatter en forbindelse i ethvert af kravene 1 til 14 eller et farmakologisk acceptabelt salt deraf og en farmakologisk acceptabelt bærer.

16. En forbindelse i henhold til ethvert af kravene 1 til 14 eller et farmaceutisk acceptabelt salt deraf til brug i medicin.

17. En forbindelse i ethvert af kravene 1 til 14 eller et farmakologisk acceptabelt salt deraf, til brug i behandlingen af en sygdom eller lidelse, som er valgt fra astma, kronisk obstruktiv lungesygdom (COPD), bronkitis, allergisk rinitis, atopisk dermatitis, kontaktdermatitis, acne, cystisk fibrose, afvisning af allotransplantation, multipel sclerose, skleroderma, arthritis, reumatoid arthritis, juvenil reumatoid arthritis, osteoarthritis, ankyloserende spondylitis, systemisk lupus erythematosus (SLE), psoriasis, Hashimotos sygdom, pankreatitis, autoimmun diabetes, diabetes type I, autoimmun okulær sygdom, ulcerativ colitis, Crohns sygdom, regional enteritis, inflammatorisk tarmsygdom (IBD), inflammatorisk tarmsyndrom (IBS), Sjögrens syndrom, optisk neuritis, fedme, hepatosteatoze, adipøs vævsassocieret inflammation,

- insulinresistens, diabetes type II, neuromyelitis optica, myasthenia gravis, aldersrelateret makulær degenerering, tørt øje, uveitis, Guillain-Barré-syndrom, psoriasis, psoriatisk artrit (PsA), steroidresistent astma, Graves' sygdom, scleritis, større depression, sæsonafhængig stemningsforstyrrelse, PTSD, bipolar lidelse, autisme, epilepsi, Alzheimers, lidelser i
- 5 centralnervesystem associeret med forandret søvn og/eller cirkadiske rytmer, endometriose, obstruktiv søvnapnøsyndrom (OSAS), Behçets sygdom, dermatomyositis, polymyocitis, transplantat-mod-vært-sygdom, primær biliær cirrose, leverfibrose, ikke-alkoholisk fedtleversygdom (NAFLD), sarkoidose, primær scleroserende cholangitis, autoimmun skjoldbruskkirtelsygdom, autoimmun polyendokrint syndrom type I, autoimmun polyendokrint
- 10 syndrom type II, cøliaki, neuromyelitis, juvenil idiopatisk artrit, systemisk sclerose, myokardieinfarkt, pulmonær hypertension, osteoarthritis, kutan leishmaniasis, sinonasal polypose og kræft.
- 18.** Forbindelsen til brug i henhold til krav 17 eller et farmakologisk acceptabelt salt deraf,
- 15 hvori sygdommen eller lidelsen er valgt fra astma, atopisk dermatitis, acne, Crohns sygdom, regional enteritis, ulcerativ colitis, Sjögrens syndrom, uveitis, Behçets sygdom, dermatomyositis, multipel sclerose, ankyloserende spondylitis, systemisk lupus erythematosus (SLE), skleroderma, psoriasis, psoriatisk artrit (PsA), steroidresistent astma og reumatoid arthritis.