



(86) Date de dépôt PCT/PCT Filing Date: 2014/02/22
(87) Date publication PCT/PCT Publication Date: 2014/08/28
(45) Date de délivrance/Issue Date: 2024/02/13
(85) Entrée phase nationale/National Entry: 2015/08/21
(86) N° demande PCT/PCT Application No.: US 2014/017867
(87) N° publication PCT/PCT Publication No.: 2014/130909
(30) Priorité/Priority: 2013/02/22 (US61/768,047)

(51) Cl.Int./Int.Cl. A61K 31/7105 (2006.01),
C12N 15/54 (2006.01), C12N 15/63 (2006.01),
C12N 15/85 (2006.01), C12N 15/87 (2006.01),
C12N 9/12 (2006.01), C12P 19/34 (2006.01)

(72) Inventeurs/Inventors:
RAMUNAS, JOHN, US;
YAKUBOV, EDUARD, US;
BLAU, HELEN M., US;
COOKE, JOHN, US

(73) Propriétaire/Owner:
THE BOARD OF TRUSTEES OF THE LELAND
STANFORD JUNIOR UNIVERSITY, US

(74) Agent: SMART & BIGGAR LP

(54) Titre : COMPOSES, COMPOSITIONS, PROCEDES, ET KITS LIES A L'EXTENSION DE TELOMERES
(54) Title: COMPOUNDS, COMPOSITIONS, METHODS, AND KITS RELATING TO TELOMERE EXTENSION

(57) Abrégé/Abstract:

Compounds and compositions for the transient expression of exogenous telomerase activity in a cell are provided. The compounds and compositions, which relate to a ribonucleic acid coding for a telomerase reverse transcriptase, are useful in the extension of telomeres in cells needing such treatment. Such cells include, for example, cells that contain shortened telomeres and cells from subjects that may benefit from telomere extension, for example subjects that suffer from, or are at risk of suffering from, age-related or other illnesses. Also provided are methods of extending telomeres through the administration of the provided compounds and compositions to animal cells, either in vitro or in vivo, and kits including the compounds and compositions and instructions for use.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number

WO 2014/130909 A1

(43) International Publication Date
28 August 2014 (28.08.2014)

(51) International Patent Classification:
C12P 19/34 (2006.01)

(21) International Application Number:
PCT/US2014/017867

(22) International Filing Date:
22 February 2014 (22.02.2014)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/768,047 22 February 2013 (22.02.2013) US

(71) Applicant (for all designated States except US): THE BOARD OF TRUSTEES OF THE LELAND STANFORD JUNIOR UNIVERSITY [US/US]; Office of Technology Licensing, 1705 El Camino Real, Palo Alto, California 94306-1106 (US).

(72) Inventors; and

(71) Applicants (for US only): RAMUNAS, John [CA/US]; 425 Olive Street, Menlo Park, California 94025 (US). YAKUBOV, Eduard [IL/US]; 2600 Gramercy Street, #332, Houston, Texas 77030 (US). BLAU, Helen M. [US/US]; 580 Cotton Street, Menlo Park, California 94025 (US). COOKE, John [US/US]; Apt. 845, Arcadian Kirby, 7550 Kirby Drive, Houston, Texas 77030 (US).

(74) Agent: ROISE, David A.; VLP Law Group LLP, 555 Bryant Street, Suite 820, Palo Alto, California 94301 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2014/130909 A1

(54) Title: COMPOUNDS, COMPOSITIONS, METHODS, AND KITS RELATING TO TELOMERE EXTENSION

(57) Abstract: Compounds and compositions for the transient expression of exogenous telomerase activity in a cell are provided. The compounds and compositions, which relate to a ribonucleic acid coding for a telomerase reverse transcriptase, are useful in the extension of telomeres in cells needing such treatment. Such cells include, for example, cells that contain shortened telomeres and cells from subjects that may benefit from telomere extension, for example subjects that suffer from, or are at risk of suffering from, age-related or other illnesses. Also provided are methods of extending telomeres through the administration of the provided compounds and compositions to animal cells, either in vitro or in vivo, and kits including the compounds and compositions and instructions for use.

Compounds, Compositions, Methods, and Kits Relating to Telomere Extension

Cross-Reference To Related Application

[0001] This application claims the benefit of U.S. Provisional Application No. 61/768,047, filed on February 22, 2013.

5

Background of the Invention

[0002] Telomeres comprise repetitive DNA sequences at the ends of linear chromosomes that, when sufficiently long, allow each chromosome end to form a loop that protects the ends from acting as double-stranded or single-stranded DNA breaks. Artandi & DePinho (2010) *Carcinogenesis* 31:9–18. Telomeres shorten over time, due in part to oxidative damage and incomplete DNA replication, eventually leading to critically short telomeres unable to form the protective loop, exposure of the chromosome ends, chromosome-chromosome fusions, DNA damage responses, and cellular senescence, apoptosis, or malignancy. O'Sullivan and Karlseder (2010) *Nat. Rev. Mol. Cell Biol.* 11:171–181; Calado *et al.* (2012) *Leukemia* 26:700–707; Artandi and DePinho (2010) *Carcinogenesis* 31:9–18.

[0003] The enzyme complex telomerase extends telomeres and comprises two essential components: the telomerase reverse transcriptase (TERT), and an RNA component known as telomerase RNA component (TERC). Other components of the telomerase complex include the proteins TCAB1, Dyskerin, Gar1, Nhp2, Nop10, and RHAU. Brouilette *et al.* (2003) *Arteriosclerosis, Thrombosis, and*

Vascular Biology 23:842-846. TERT is a limiting component of the telomerase complex, and thus treatments that increase TERT can increase telomerase activity. Telomerase activity is typically measured using the telomeric repeat amplification protocol (TRAP) assay, which quantifies the ability of a cell lysate or other sample 5 to extend a synthetic telomere-like DNA sequence.

[0004] As would be expected due to the importance of telomere length maintenance in preventing cellular senescence and apoptosis and resulting cellular dysfunction, genetic mutations of TERT and TERC are linked to fatal inherited diseases of inadequate telomere maintenance, including forms of idiopathic 10 pulmonary fibrosis, dyskeratosis congenita, and aplastic anemia. The effects of premature cellular senescence and apoptosis due to short telomeres in these diseases are devastating in themselves, and may be compounded by increased risk of cancer. Artandi and DePinho (2010) *Carcinogenesis* 31:9-18; Alter *et al.* (2009) *Blood* 113:6549-6557. In addition to abundant correlative data linking 15 short telomeres to cancer (Wentzensen *et al.* (2011) *Cancer Epidemiol. Biomarkers Prev.* 20:1238-1250), aplastic anemia provides some of the first direct evidence that critically short telomeres and resulting chromosomal instability predispose cells to malignant transformation in humans (Calado *et al.* (2012) *Leukemia* 26:700-707). There is evidence that short telomeres make the difference between 20 fatal and non-fatal muscular dystrophy (Sacco *et al.* (2010) *Cell* 143:1059-1071), and that telomere extension averts endothelial cell senescence (Matsushita *et al.* (2001) *Circ. Res.* 89:793-798), which is associated with atherosclerosis, 25 hypertension, and heart disease (Pérez-Rivero *et al.* (2006) *Circulation* 114:309-317). In addition to being implicated in these and other diseases, short telomeres also limit cell amplification for cell therapies and bioengineering applications. Mohsin *et al.* (2012) *Journal of the American College of Cardiology* doi:10.1016/j.jacc.2012.04.0474.

[0005] A natural product-derived telomerase activator, TA-65®, has been sold commercially as a nutraceutical by T.A. Sciences, Inc. Harley *et al.* (2011) 30 *Rejuvenation Research* 14:45-56. This compound purportedly turns on the endogenous hTERT gene, thus activating expression of native telomerase. It is not

clear, however, how this treatment overcomes the normal regulation of the native telomerase activity.

[0006] Human cells with little or no telomerase activity have been transfected with vectors encoding human TERT (hTERT). See, e.g., Bodnar *et al.* (1998)

5 *Science* 279:349-352. The transfected cells were found to express telomerase, to display elongated telomeres, to divide vigorously, and to display reduced senescence compared to cells lacking telomerase, but the genomic modification resulting from this treatment increases the risk and limits the utility of the approach.

10 [0007] A limited capacity to replicate is one of the defining characteristics of most normal cells. An end-point of this limited replicative process is senescence, an arrested state in which the cell remains viable but no longer divides. Senescent cells are often characterized by an altered pattern of gene expression, altered morphology, and reduced or abrogated ability to perform their normal functions.

15 [0008] The shortening of telomeres plays a direct role in cellular senescence in animal tissues during aging. Furthermore, there is accumulating evidence implicating short telomeres in a variety of diseases, including those described above. The prospect of preventing disease by telomere extension motivates a need for safe and effective treatments to extend telomeres in animal cells *in vivo* and/or 20 *in vitro*. Further, there is a need to safely and rapidly extend telomeres in cells for use in cell therapy, cell and tissue engineering, and regenerative medicine.

25 [0009] At the same time, however, there is a danger in the constitutive activation of telomerase activity. Indeed for cell therapy applications, avoiding the risk of cell immortalization is of paramount importance. To this end, transient, rather than constitutive, telomerase activity may be advantageous for safety, especially if the elevated telomerase activity is not only brief but extends telomeres rapidly enough that the treatment does not need to be repeated continuously. Current methods of extending telomeres include viral delivery of TERT under an inducible promoter, delivery of TERT using vectors based on adenovirus and adeno-associated virus, 30 and small molecule activators of telomerase. These methods risk either insertional mutagenesis, continual elevation of telomerase activity, or both.

[0010] Thus, there is strong motivation to develop a therapy that safely extends telomeres to potentially prevent, delay, ameliorate, or treat these and other conditions and diseases, to do the same for the gradual decline in physical form and function and mental function that accompanies chronological aging, and to 5 enable cell therapies and regenerative medicine. Such a therapy would be of great use in the rejuvenation of all animals, including humans, pets, livestock, zoo animals, and animals of endangered species.

Summary of the Invention

[0011] The present invention addresses these and other problems by providing an 10 alternative that offers the benefits of a highly transient increase in telomerase activity, combined with rapid telomere extension so that the treatment does not need to be continuous or even frequent, and with no risk of genomic insertional mutagenesis. Specifically, the invention provides compositions, methods, and kits for the extension of telomeres by the transient translation of exogenous telomerase 15 activity in a cell.

[0012] In one aspect, the invention provides compounds for the extension of telomeres comprising:

20 a synthetic ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase; wherein telomeres are extended within a cell treated with the compound.

[0013] In some embodiments, the telomerase reverse transcriptase is a mammalian, avian, reptilian, or fish telomerase reverse transcriptase or a variant that retains telomerase catalytic activity, and in specific embodiments is a human telomerase reverse transcriptase.

25 [0014] In some embodiments, the ribonucleic acid comprises a 5' cap, a 5' untranslated region, a 3' untranslated region, and a poly-A tail. The 5' cap may be non-immunogenic and the 5' cap may have been treated with phosphatase.

[0015] In preferred embodiments, the poly-A tail increases stability of the 30 ribonucleic acid.

[0016] In other preferred embodiments, the 5' untranslated region or the 3' untranslated region comprise a sequence from a stable mRNA or an mRNA that is

efficiently translated, or they both comprise a sequence from a stable mRNA or an mRNA that is efficiently translated.

[0017] In still other preferred embodiments, the 5' cap, the 5' untranslated region, or the 3' untranslated region stabilizes the ribonucleic acid, increases the rate of 5 translation of the ribonucleic acid, or modulates the immunogenicity of the ribonucleic acid.

[0018] In highly preferred embodiments, the at least one modified nucleoside modulates immunogenicity of the ribonucleic acid.

[0019] In some embodiments, the ribonucleic acid is a purified synthetic 10 ribonucleic acid.

[0020] In preferred embodiments, the synthetic ribonucleic acid is purified to remove immunogenic components.

[0021] In certain specific embodiments, the ribonucleic acid codes for a human, cat, dog, mouse, horse, cow, sheep, pig, African elephant, chicken, rat, zebrafish, 15 Japanese medaka, or chimpanzee telomerase reverse transcriptase, or a polypeptide with at least 95% sequence identity to the telomerase reverse transcriptase.

[0022] In another aspect, the invention provides compositions comprising a compound of the invention and a telomerase RNA component, which, in some embodiments, is a mammalian, avian, reptilian, or fish telomerase RNA 20 component. In more specific embodiments, the telomerase RNA component is a human telomerase RNA component. In some embodiments, the compounds and compositions of the invention further comprise a delivery vehicle.

[0023] In some embodiments, the delivery vehicle is an exosome, a lipid nanoparticle, a polymeric nanoparticle, a natural or artificial lipoprotein particle, a cationic lipid, a protein, a protein-nucleic acid complex, a liposome, a virosome, or 25 a polymer. In specific embodiments, the delivery vehicle is a cationic lipid.

[0024] In preferred embodiments, the delivery vehicle is non-immunogenic. In other preferred embodiments, the delivery vehicle is partly immunogenic. In particular, under some circumstances, it may be desirable for the vehicle to retain 30 some immunogenicity.

[0025] According to another aspect of the invention, methods of extending telomeres are provided, comprising the step of administering any of the disclosed

compounds or compositions to an animal cell, wherein at least one telomere is extended within the cell.

[0026] In some method embodiments, the cell has at least one shortened telomere prior to the administering step.

5 [0027] In some embodiments, the cell is from or in a subject suffering from or at risk of an age-related illness, an age-related condition, or an age-related decline in function or appearance.

[0028] In some embodiments, the cell is from or in a subject suffering from or at risk of cancer, heart disease, stroke, diabetes, diabetic ulcers, Alzheimer's disease, 10 osteoporosis, a decline in physical ability or appearance, physical trauma or chronic physical stress, psychological trauma or chronic psychological stress, reduced immune function, immunosenescence, or macular degeneration.

[0029] In some embodiments, the cell is a somatic cell of endodermal, mesodermal, or ectodermal lineage, or a germ line or embryonic cell.

15 [0030] In some embodiments, the cell is an induced pluripotent stem cell or a cell used to produce an induced pluripotent stem cell.

[0031] In some embodiments, the cell is a transdifferentiated cell or a cell used to produce a transdifferentiated cell.

20 [0032] In some embodiments, the cell is an isolated cell, and the administering step lasts no longer than 48 hours. In other embodiments, the cell is an isolated cell, and the administering step lasts at least 2 hours.

[0033] In some embodiments, the cell is an isolated cell, and the administering step is performed no more than four times. In other embodiments, the cell is an isolated cell, and the administering step is performed at least two times.

25 [0034] In some embodiments, the cell is an isolated cell, and the method further comprises the step of measuring telomerase activity in the cell. In specific embodiments, the administering step increases telomerase activity in the cell, and in even more specific embodiments, the telomerase activity is transiently increased by at least 5%. In other specific embodiments, the half-life of increased telomerase 30 activity is no longer than 48 hours.

[0035] In some embodiments, the method further comprises the step of measuring telomere length in the cell. In specific embodiments, average telomere length is increased by at least 0.1 kb.

5 [0036] In some embodiments, the cell is an isolated cell, and the method further comprises the step of measuring population doubling capacity in the cell. In specific embodiments, the population doubling capacity increases, in some cases by at least one population doubling.

[0037] In preferred embodiments, the cell is from or in a mammalian subject, and in even more preferred embodiments, the cell is from or in a human subject.

10 [0038] In some embodiments, the cell is an isolated cell, and in other embodiments, the cell is not an isolated cell. In some embodiments, the administering step comprises electroporation. In some embodiments, the at least one telomere is transiently extended within the cell.

15 [0039] According to yet another aspect of the invention, kits for extending telomeres in an animal cell are provided, the kits comprising any of the above compounds or compositions and instructions for using the compound or composition to extend telomeres.

20 [0040] In some embodiments, the kits further comprise packaging materials. In some specific embodiments, the packaging materials are air-tight. In some specific embodiments, the packaging materials comprise a metal foil container.

[0041] In some embodiments, the kits further comprise a desiccant, a culture medium, or an RNase inhibitor.

25 [0042] In some embodiments, the composition is sterile. In some embodiments, the kit comprises a compound of the invention, instructions for using the composition to extend telomeres, and a telomerase RNA component, a delivery vehicle, or both a telomerase RNA component and a delivery vehicle.

30 [0043] In yet another aspect, the invention provides methods of treatment, comprising administering a compound or composition of the invention to an animal subject in need of, or that may benefit from, telomere extension. In some embodiments, the animal subject suffers from or is at risk of a disease or condition resulting from shortened telomeres.

[0043a] In one embodiment, the invention provides a method of extending telomeres, comprising the step of: administering a compound to an animal cell wherein said administering is in vitro, wherein the compound comprises a synthetic ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase, wherein the at least one modified nucleoside comprised in the synthetic ribonucleic acid confers reduced immunogenicity to the synthetic ribonucleic acid, wherein the telomerase reverse transcriptase is expressed transiently in the cell, and wherein at least one telomere is extended within the cell.

5

[0043b] In one embodiment, the invention provides use of a compound comprising a synthetic ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase for extending telomeres in an animal cell, wherein the at least one modified nucleoside comprised in the synthetic ribonucleic acid confers reduced immunogenicity to the synthetic ribonucleic acid, and wherein the use is for transient expression of telomerase reverse transcriptase in the cell.

10

[0043c] In one embodiment, the invention provides a kit for extending telomeres in an animal cell, comprising: a compound comprising a synthetic ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase, wherein the at least one modified nucleoside comprised in the synthetic ribonucleic acid confers reduced immunogenicity to the synthetic ribonucleic acid; and instructions for using the compound for the transient expression of telomerase reverse transcriptase to extend telomeres.

15

[0043d] In one embodiment, the invention provides use of a compound comprising a synthetic ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase for treating an animal subject in need of, or that may benefit from, telomere extension, wherein the at least one modified nucleoside comprised in the synthetic ribonucleic acid confers reduced immunogenicity to the synthetic ribonucleic acid, and wherein the use is for transient expression of telomerase reverse transcriptase in a cell of the subject.

20

25

Brief Description of the Drawings

[0044] Figure 1. Transfection of MRC-5 cells with a modified ribonucleic acid (modRNA) encoding TERT, but not catalytically-inactive TERT (CI TERT), transiently increases telomerase activity. (a) Schematic of the modRNA construct and approach used in these studies. The modRNA construct encodes human TERT and has 5' and 3' UTRs that confer stability, such as from β -globin mRNA. CI TERT has a single bp mutation. (b) Transfection of TERT or CI TERT modRNA results in increased exogenous TERT modRNA levels in treated cells, but not untreated controls, above endogenous TERT RNA levels. (c) Levels of protein recognized by anti-TERT antibody are significantly higher in MRC-5 cells 24 hours after transfection with modRNA encoding TERT or CI TERT ($P<0.03$ or 0.01, respectively) than in untreated or vehicle-only cells. (d) Treatment with modRNA encoding TERT, but not CI TERT, causes transient increase in telomerase activity in MRC-5 cells.

[0045] Figure 2. Use of exosomes as a delivery vehicle for ribonucleic acid therapeutics.

[0046] Figure 3. Graphical illustration of the use of multiple rapid and transient telomerase treatments in the extension of telomeres in a cell.

[0047] Figure 4. Treatment with TERT modRNA rapidly extends telomeres in MRC-5 cells. (a) Treatment and assay schedule used in these studies. Treatments lasted 5 hours. (b) Image of TRF Southern blot (inset) and its quantification in which the chemoluminescent signal was normalized to account for the number of probes per unit of telomere length, and the average intensity of each pixel row in each gel lane was then plotted relative to telomere length ($n=3$ biological replicates for each treatment). (c) Averages of telomere length distributions in Figure 4b, revealing that telomeres in cells treated with TERT modRNA are significantly ($P<0.014$) longer than in untreated or vehicle only-treated controls. (d) Proportion of telomeres longer than an arbitrary threshold (4.2 kb) is greater in cells treated with TERT modRNA than in untreated and vehicle only-treated controls. (e) Representative fluorescence micrograph of metaphase spread of MRC-5 fibroblasts used in quantitative *in situ* hybridization (Q-FISH) to compare telomere lengths, showing telomere probe (light, punctate) and DNA (smooth shading). (f)

Quantification of Q-FISH measurements of telomere length in treated and control cells (n=15 cells for each of two biological replicates for each treatment), showing rapid telomere extension in treated cells. (Note that TRF measures the length of both telomeric and subtelomeric DNA, whereas Q-FISH measures only telomeric DNA, thus explaining the differences between the Q-FISH and TRF results.) (g) (i) Standard curve relating cumulative population doublings of MRC-5 cells following receipt from the supplier to telomere length as measured using Q-FISH to quantify average total telomere probe fluorescence per cell, which is linearly proportional to telomere length. (g) (ii) Quantification of average telomere length per cell in MRC-5 cells treated three times with TERT modRNA at 48 hour intervals, as measured using Q-FISH.

[0048] Figure 5. Brief treatment with TERT modRNA increases replicative capacity of, but does not immortalize, MRC-5 cells in a dose-dependent manner. (a) Growth curves of cells treated with TERT modRNA once, twice, or three times, or three times followed by an additional treatment 8 weeks after the first treatment. Controls comprise either no treatment, or treatment with vehicle only or CI TERT four times. (n=3 for each treatment). (b) Growth curves of cells treated with TERT modRNA and TERC RNA in a 1:5 molar ratio (n=3). (c) Treatment once, twice, or three times with TERT modRNA confers additional replicative capacity to MRC-5 cells beyond that of untreated cells, in a dose-dependent manner. The incremental increase in proliferative capacity conferred by the second and third treatments, delivered at 48 hours after the first treatment, is not as great as the increase in proliferative capacity conferred by the first treatment; however, an additional fourth treatment several weeks after the first three treatments confers as much additional proliferative capacity as did the first treatment, indicating that the timing of the treatments is important for optimizing the amount of proliferative capacity conferred by the treatment. Treatment three times with TERT modRNA and TERC modRNA together in a 1:5 molar ratio confers greater proliferative capacity than treatment three times with TERT modRNA alone. Treatment with vehicle only or CI TERT modRNA confers no additional replicative capacity. (n=3).

[0049] Figure 6. High efficiency of transfection of MRC-5 cells treated with modRNA encoding nuclear GFP (nGFP). (a) Percentage of cells with fluorescence more than two SD above the mean of untreated cells (n=10000). (b) Mean fluorescence (n=10000). (c) Fluorescence micrograph of nGFP in transfected 5 MRC-5 cells, counterstained with DAPI.

[0050] Figure 7. Dose-response of telomerase activity.

[0051] Figure 8. TERT modRNA treatment delays cell swelling in MRC-5 cells. (a) MRC-5 cells in early passages (left) are several times smaller area than cells in later passages (right), as shown in typical micrographs of PD 2 and PD 53 10 untreated cells as seen on a hemocytometer. (b) Early (PD 2) and mid (PD 35) passage MRC-5 cells exhibit little swelling, defined as having a diameter greater than 25 microns as measured visually on a hemocytometer (where one small square is 50 microns across), whereas a significantly greater fraction of late passage (PD 53) cells are swollen. In contrast, PD 53 cells that had been treated 15 with TERT modRNA, but not those treated with CI TERT modRNA, at PD 40 exhibit little swelling.

[0052] Figure 9. Growth curve for human microvascular dermal endothelial cells (HMDECs).

[0053] Figure 10. Effect of TERT modRNA treatment on cell number in 20 HMDECs. CO: control treatment; hTERT-CI: catalytically-inactive hTERT; hTERT-WT: wild-type hTERT.

[0054] Figure 11. Effect of TERT modRNA treatment on growth of HMDECs. UT: untreated; CO: control treatment; CI: catalytically-inactive hTERT; WT: wild-type hTERT.

25 [0055] Figure 12. Effect of TERT modRNA treatment on senescence in HMDECs. NT: untreated; hTERT-CI: catalytically-inactive hTERT; hTERT-WT: wild-type hTERT.

[0056] Figure 13. Increases in *TERT* protein and telomerase activity following 30 modified *TERT* mRNA delivery. (A) Schematic of modified mRNA comprising the coding sequence of the full length functional form of *TERT* or a catalytically-inactive (CI) form of *TERT*, flanked by untranslated regions (UTRs) of *HBB* and a 151 nt poly-A tail, synthesized using modified nucleotides pseudouridine and 5-

methylcytidine. **(B)** Transfection efficiency of myoblasts (n=2,000) treated with 0.8 μ g/ml modified mRNA encoding GFP measured by flow cytometry 24 h post-transfection exceeded 95% (additional plots in Figure 16A). **(C)** Total *TERT* protein levels were measured by quantitative Western blot (panel C, left; Figure 5 17). Quantification of *TERT* protein levels 24 hours after transfection with 1 μ g/ml of either *TERT* or CI *TERT* mRNA (n=3). Quantification of *TERT* protein in response to various doses of mRNA was measured at the single cell level by flow cytometry (n=10,000) (panel C, right). *P<0.05, **P<0.01 compared to untreated cells. Error bars represent s.e.m. **(D)** Detection of telomerase activity in 10 fibroblasts and myoblasts transfected with 1 μ g/ml modified *TERT* mRNA, as measured using the telomere repeat amplification protocol (TRAP). Arrow indicates internal controls for PCR efficiency.

[0057] Figure 14. Increases in telomere length and proliferative capacity following modified *TERT* mRNA (modRNA) delivery. **(A)** Mean telomere lengths in untreated fibroblasts decreased over time in culture as measured by MMqPCR 15 and by SpectraCell (correlation coefficient 0.97, P<0.001). Experiment was repeated twice with four technical replicates each. **(B)** Mean telomere lengths in fibroblasts transfected with 1 μ g/ml *TERT* modRNA, CI *TERT* mRNA, or vehicle only, once, twice, or three times in succession at 48 h intervals. Experiment was 20 repeated twice with four technical replicates each. **P<0.01, ***P<0.001 compared to vehicle only-treated cells. **(C)** Mean telomere lengths in myoblasts treated as in (B). Experiment was repeated twice with four technical replicates each. ***P<0.001 compared to vehicle only-treated cells. **(D)** Growth curves of fibroblasts treated as in (B), with green arrows indicating treatments times. 25 Growth curves were repeated twice with each population cultured in triplicate. Replicative capacity increased in a dose-dependent manner (right panel). *P<0.05, **P<0.01 compared to vehicle only-treated cells. **(E)** Proliferation capacity of myoblasts, treated as in (B) (green arrows). Growth curves were repeated twice, with each population cultured in triplicate. All data are presented as means \pm 30 s.e.m.

[0058] Figure 15. Transient reduction of senescence-associated markers following modified *TERT* mRNA delivery. **(A)** Quantification of β -gal-expressing

fibroblasts after modified *TERT* mRNA transfection three times in succession at 48 h intervals (green arrows). The control cells, comprising untreated, vehicle only, and CI *TERT* mRNA-treated populations, stopped expanding at PD 53, and the *TERT* modRNA-treated population stopped expanding at PD 80. Each experiment 5 was conducted twice with >50 cells per sample scored manually. Representative images show β -gal-stained *TERT* modRNA-treated fibroblasts at PD 53 (top) and PD 80 (bottom). Scale bar length is 200 microns. (B) Quantification of β -gal expression in myoblasts treated as in (A). Controls are as in (A). The control and *TERT* modRNA-treated populations stopped expanding at PD 8 and PD 11, 10 respectively. Each experiment was conducted twice, with >50 cells per sample scored manually. Representative images show myoblasts at PD 2 (top) and *TERT* modRNA-treated myoblasts at PD 11 (bottom). (C) Quantification of enlarged cells associated with replicative senescence in fibroblasts transfected three times with modified *TERT* mRNA. Population plateaus are as in (A). Controls are 15 vehicle only and CI *TERT* mRNA-treated. Each experiment was conducted twice, with >50 cells per sample scored manually. Representative images show untreated fibroblasts at PD 2 (top) and PD 53 (bottom). All data are presented as means \pm s.e.m. Scale bar length is 200 microns.

[0059] Figure 16. High efficiency transfection of modified mRNA into human 20 myoblasts. (A,B) Quantification of GFP fluorescent myoblasts (n=2000) transfected with 0.1–0.8 μ g/ml of modified mRNA encoding GFP as measured by flow cytometry 24 h after start of treatment. (C) Mean fluorescence of modified GFP mRNA-transfected myoblasts in response to increasing doses. (D) Quantification of exogenous *TERT* modRNA in fibroblasts 24 h after transfection 25 with 1 μ g/ml of *TERT* or CI *TERT* modRNA, as measured using RT-qPCR. Ratio of *TERT* to CI *TERT* was calculated using the Pfaffl method with RPL37A and GAPDH as reference genes (n=3). (E) Quantification of endogenous *TERT* modRNA in fibroblasts 24 h after transfection with 1 μ g/ml of *TERT* or CI *TERT* modRNA, as measured using qPCR, calculated as in (D). All data are presented as 30 means \pm s.e.m.

[0060] Figure 17. Expression of *TERT* after modified mRNA delivery. *TERT* protein expression in fibroblasts harvested 24 h after start of treatment with 1

$\mu\text{g}/\text{ml}$ *TERT* modRNA was measured by multiplexed infrared Western blot. The serial dilution of total protein was used to generate a standard curve to compare relative amounts of *TERT* protein to controls. The specificity of the *TERT* antibody used here has been extensively tested (Wu, 2006).

5 [0061] Figure 18. TRAP gel showing increased telomerase activity in mononuclear cells electroporated with *TERT* modRNA at a setting of 200 V, 25 uF, 1000 Ohm, in a 1 mm cuvette with 10 μl of cell suspension and *TERT* modRNA. The heat treated sample lane contains a primer dimer artifact of the method as indicated by the strong band at the fourth rung of the ladder.

10 [0062] Figure 19. Microscope images of mononuclear cells fluorescing after electroporation with modRNA encoding GFP.

[0063] Figure 20. Electroporation of GFP modRNA into human leukocytes (mononuclear cells) results in high transfection efficiency.

[0064] Figure 21. Image of gel from TRAP assay showing increasing telomerase activity in fibroblasts electroporated with increasing doses of *TERT* modRNA.

15 [0065] Figure 22. Fluorescent micrographs showing that CD8+ T-cells activated with CD3/CD28 and electroporated with GFP modRNA express GFP activity. Dark dots are CD3/CD28-coated beads.

[0066] Figure 23. Electroporation of human keratinocytes with *TERT* modRNA results in expression of telomerase activity.

20 [0067] Figure 24. Expression of a luciferase modRNA after *in vivo* delivery to a rodent.

Detailed Description of the Invention

25 [0068] Telomeres are DNA sequences at the ends of chromosomes that protect the ends of the chromosomes but that shorten over time. Critically short telomeres may cause cells to stop functioning correctly or to die. Critically short telomeres may also lead to chromosome fusions that may in turn lead to cancer. Even in the absence of a specific diagnosed disease, short telomeres are implicated in the gradual decline in function of the mind and body and in the appearance of aging.

30 [0069] At the same time, however, telomere shortening can play a protective role against cancer, for example in the situation where a cell acquires a mutation that

causes it to proliferate faster than normal cells. In that situation, telomere shortening prevents the cell from proliferating indefinitely and causing cancer. It is therefore not necessarily beneficial to extend telomeres continually.

[0070] In mammals, telomeres comprise tandem repeats of the sequence 5 TTAGGG, and in other animals such as birds, reptiles, and fish, the repeated sequence varies. In all of these types of animals, the telomeres are double stranded for many kilobases (kb). Average telomere lengths differ between species as shown in Table 1.

10 **Table 1: Average telomere length in adult fibroblasts of different species**

Species	Average telomere length (kb)
Cow	18
Sheep	18
Pig	15
Horse	14
Dog	15
Panda	25
Tiger	50
House mouse	40
Sonoran deer mouse	9
Norway rat	40
Naked mole rat	16
European white rabbit	50
Black-tailed jack rabbit	25
Spider monkey	7
Squirrel monkey	9
Rhesus monkey	16
Orangutang	10
Bonobo	10
Human	9
Indian elephant	15

African elephant	14
Cat	11

[0071] In humans, telomeres start out before birth with lengths of 15-20 kb, and at birth with lengths of 12-15 kb. Telomeres shorten rapidly during childhood, and then by about 0-100 bp per year on average in adulthood, a rate which varies 5 depending on the cell type, exposure to psychological or oxidative stress, and other factors.

[0072] Telomeres are part of the telomere complex, which protects the ends of chromosomes. The telomere complex also comprises a set of proteins collectively called Shelterin. Telomere complex proteins include POT1, TPP1, ATM, DAT, 10 TRF1, TRF2, Rap1, Rif1, TIN2, NBS, MRE17, and RAD50 and their homologs in different mammalian species. Podlevsky and Chen (2012) *Mutat. Res.* 730:3-11. In many species the telomere terminates in a single-stranded 3' overhang which inserts itself into the double stranded region, in association with telomere complex proteins, forming a loop within the telomere complex.

[0073] Telomeres shorten over time, due to oxidative damage and sister chromatid exchange, and also due to the end replication problem, in which the ends of chromosomes are not completely duplicated during mitosis. When telomeres become critically short, the telomere complex is no longer able to protect the chromosome ends, and the chromosome ends become "uncapped". Uncapping of 15 the chromosome ends may result in chromosome-chromosome fusions, which may in turn result in cancer. O'Sullivan and Karlseder (2010) *Nat. Rev. Mol. Cell Biol.* 11:171-181; Calado *et al.* (2012) *Leukemia* 26:700-707; Artandi and DePinho 20 (2010) *Carcinogenesis* 31:9-18. Uncapping can also result in the chromosome ends being recognized as damaged DNA, activating DNA damage responses and triggering cell apoptosis or senescence. Senescence is an arrested state in which 25 the cell remains viable but no longer divides, and senescent cells typically cease to perform their normal, pre-senescence, useful functions adequately or at all. Thus telomere shortening leads to tissue dysfunction, loss of physical ability and youthful appearance, loss of mental ability, and disease in part due to the 30 accumulation of senescent cells, and in part due to the loss of cells by apoptosis.

Indeed, aged people with short telomeres are approximately 200-750% more likely to develop myocardial infarction (200%) (von Zglinicki *et al.* (2000) *Laboratory Investigation; a Journal of Technical Methods and Pathology* 80:1739-1747), vascular dementia (200%) (Testa *et al.* (2011) *Diabetic Medicine* 28:1388-1394, 5 diabetes with complications (400%) (Blackburn *et al.* (2010) *Cancer Prevention Research* 3:394-402, cancer (Stern and Bryan (2008) *Cytogenetic and Genome Research* 122:243-254), stroke, Alzheimer's disease, infection (750%), idiopathic pulmonary fibrosis, and other disease. People with short telomeres in one tissue are likely to also have short telomeres in most of their other tissues, and thus short 10 telomeres correlate with increased risk for many diseases in one individual.

Takubo *et al.* (2010) *Geriatr Gerontol Int.* 10 Suppl 1:S197-206; doi: 10.1111/j.1447-0594.2010.00605.x. Short telomeres also limit cell replicative capacity which in turn limits cell therapies and regenerative therapies. Conversely, increasing telomere length in mice with short telomeres using virus-based genetic 15 engineering methods rejuvenates the mice by several parameters, including skin thickness and elasticity, liver function, and sense of smell. Jaskelioff (2011) *Nature* 469:102-107.

[0074] Since telomerase extends telomeres, a useful approach to extending telomeres is to increase the level of telomerase activity in cells. Many agents and 20 conditions have been reported to increase telomerase activity in cells, including the agents and conditions listed in Table 2.

Table 2: Examples of agents and conditions that increase telomerase activity

Type	Examples
Growth factors	EGF, IGF-1, FGF-2, VEGF (Liu <i>et al.</i> (2010) <i>Ageing Research Reviews</i> 9:245-256)
Genetic treatments	Viral delivery of DNA encoding TERT (Matsushita (2001) <i>Circ. Res.</i> 89:793-798; Hooijberg (2000) <i>J. Immunol.</i> 165:4239-4245); electroporation of plasmid encoding TERT (Bodnar <i>et al.</i> (1998) <i>Science</i> 279:349-352); transfection of mRNA encoding TERT (Sæbe-Larsen <i>et al.</i> (2002) <i>J. Immunol. Methods</i> 259:191-203)
Hormones	Estrogen (Imanishi <i>et al.</i> (2005) <i>Journal of Hypertension</i> 23:1699-1706), erythropoietin (Akiyama <i>et al.</i> (2011) <i>Leukemia Research</i> 35:416-418)
Physical treatments	UV radiation (Ueda <i>et al.</i> (1997) <i>Cancer Research</i> 57:370-374), hypoxia (Gladych <i>et al.</i> (2011) <i>Biochemistry and Cell Biology</i> 89:359-376)
Cytokines	IL-2, IL-4, IL-6, IL-7, IL-13, and IL-15 (Liu <i>et al.</i> (2010)

	<i>Ageing Research Reviews</i> 9:245-256)
Small molecules from plants	Resveratrol (Pearce <i>et al.</i> (2008) <i>Oncogene</i> 27:2365-2374), compounds extracted from <i>Astragalus membranaceus</i> including cycloastragenol (TAT2), TA-65, or TAT153 (Zvereva <i>et al.</i> (2010) <i>Biochemistry. Biokhimiia</i> 75:1563-1583; Harley <i>et al.</i> (2011) <i>Rejuvenation Research</i> 14:45-56)
Other	Inhibitors of Menin, SIP1 (Lin and Elledge (2003) <i>Cell</i> 113:881-889), pRB, p38 (Di Mitri <i>et al.</i> (2011) <i>Journal of Immunology</i> 187:2093-2100, p53, p73 (Beitzinger <i>et al.</i> (2006) <i>Oncogene</i> 25:813-826, MKRN1, CHIP, Hsp70 (Lee <i>et al.</i> (2010) <i>The Journal of Biological Chemistry</i> 285:42033-42045), androgens (Nicholls <i>et al.</i> (2011) <i>Protein & Cell</i> 2:726-738), and TGF-beta (Prade-Houdellier <i>et al.</i> (2007) <i>Leukemia</i> 21:2304-2310)

[0075] The treatment examples of Table 2 are not without undesired effects, however. For example, treatment with growth factors, hormones, or cytokines may cause side effects, may activate multiple signaling pathways, may cause unwanted cell replication, may trigger an immune response, and are generally non-specific. Genetic treatments using plasmids or viruses carry a risk of genomic modification by insertional mutagenesis and a risk of cancer. Transfection with unmodified RNA causes a strong immune response and has not been shown to extend telomeres. Physical treatments can damage genomic DNA. Treatment with small molecules from plants have been found to only extend telomeres in some subjects and cells, only extend telomeres very slowly, and require chronic delivery, therefore risking cancer.

[0076] The expression in cells of nucleic acid sequences encoding hTERT and TERC, and the use of these components themselves, have been proposed to be useful in the diagnosis, prognosis, and treatment of human diseases (see, *e.g.*, U.S. Patent Nos. 5,583,016 and 6,166,178), but telomere extension in a manner that is both rapid and transient, and thus potentially safe for the reasons described above, has not been demonstrated. Sæbe-Larssen *et al.* (2002) *J. Immunol. Methods* 259:191-203 reported the transfection of dendritic cells with mRNA encoding hTERT, and that such cells acquired telomerase activity, but the transfection used standard mRNA and resulted in a strong hTERT cytotoxic T lymphocyte (CTL) response rather than an extension of telomeres.

[0077] Furthermore, all existing small-molecule treatments are largely ineffective and slow (Harley *et al.* (2011) *Rejuvenation Research* 14:45-56),

primarily because they act through the catalytic component of telomerase, TERT, which is heavily regulated post-translationally, limiting existing treatments' effects to a small subset of cells, and excluding cells in interphase or G0 such as many stem and progenitor cells. Cifuentes-Rojas and Shippen (2012) *Mutat. Res.* 730:20-27; doi:10.1016/j.mrfmmm.2011.10.003; Cong *et al.* (2002) *Microbiology and Molecular Biology Reviews* 66:407-425. This regulation is mediated in part by interactions between components of the telomerase complex, the telomere complex, and other molecules. For example, TERT is phosphorylated or dephosphorylated at multiple sites by multiple kinases and phosphatases, and at some sites, phosphorylation results in increased telomerase activity (for example phosphorylation by Akt), while at others sites phosphorylation reduces telomerase activity (for example, phosphorylation by Src1 or cAbl). Also, TERT is ubiquitinated or deubiquitinated at specific sites. TERT also interacts with other proteins at specific sites on TERT, and these interactions can inactivate TERT (for example interactions with Pinx1 or cAbl), or transport TERT away from the chromosomes (for example, interactions with CRM1 and Pinx1), preventing or slowing telomere extension. Further, some proteins bind to telomeres or the telomere complex, blocking TERT (for example POT1), preventing telomere extension. Further, some proteins aid telomere extension indirectly, for example helicases and UPF1. Due to regulatory mechanisms, telomerase activity peaks during S phase of the cell cycle, and thus rapidly-dividing cells may tend to benefit more from treatments that increase telomerase activity. However, it is often desirable to keep cells in a slow-dividing or non-dividing state; for example, stem or progenitor cells are often slow-dividing, and thus may spend the majority of their time in interphase or G₀. Thus, existing treatments are slow and ineffective in most cell types generally, and in all cell types during interphase and G₀. Treatments that are slow are less safe, because they require treatment for a longer time. Since telomere-shortening provides a protective safety mechanism against run-away cell proliferation, such as in cancer, a treatment that extends telomeres rapidly is generally safer, because it may be delivered for short periods of time and infrequently, thus allowing the normal telomere-shortening safety mechanism to remain in effect for much of the time. Therefore a method capable of transiently

overcoming telomerase regulation to rapidly extend telomeres during a brief treatment is needed.

[0078] TERT regulates hundreds of genes including those listed in Table 3.

5 **Table 3: Examples of genes and pathways regulated by TERT**

Upregulated	
Type	Examples
Epigenetic state modulators	DNA 5-methylcytosine transferase I (Young <i>et al.</i> (2003) <i>The Journal of Biological Chemistry</i> 278:19904-19908)
Proto-oncogenes	Hepatocyte growth factor receptor (MET), AKT-2, CRK (Perrault <i>et al.</i> (2005) <i>Biochemical and Biophysical Research Communications</i> 335:925-936)
Differentiation, cell fate	Sox-13 (Perrault <i>et al.</i> (2005) <i>Biochemical and Biophysical Research Communications</i> 335:925-936), Wnt (Park <i>et al.</i> (2009) <i>Nature</i> 460:66-72)
Glycolysis	Phosphofructokinase (Bagheri <i>et al.</i> (2006) <i>Proc. Nat'l Acad. Sci. U.S.A.</i> 103:11306-11311), aldolase C (Bagheri <i>et al.</i> (2006) <i>Proc. Nat'l Acad. Sci. U.S.A.</i> 103:11306-11311)
Proliferation enhancers	Activating transcription factor-3, Xbox protein-1, FGF, EGFR (Smith <i>et al.</i> (2003) <i>Nature Cell Biology</i> 5:474-479), Insulin-like growth factor 2 (Perrault <i>et al.</i> (2005) <i>Biochemical and Biophysical Research Communications</i> 335:925-936), Wnt (Park <i>et al.</i> (2009) <i>Nature</i> 460:66-72), tp53bp1 (Perrault <i>et al.</i> (2005) <i>Biochemical and Biophysical Research Communications</i> 335:925-936), epiregulin (Lindvall <i>et al.</i> (2003) <i>Cancer Research</i> 63:1743-1747)
Metastasis-related genes	Mac-2 binding protein (Park <i>et al.</i> (2007) <i>International Journal of Cancer</i> 120:813-820)
Downregulated	
Type	Examples
Proliferation inhibitors	Interleukin 1 receptor antagonist, parathyroid hormone-related peptide, integrin-associated protein, TNF-related apoptosis-inducing ligand (Smith <i>et al.</i> (2003) <i>Nature Cell Biology</i> 5:474-479), IGF binding protein-5 (Perrault <i>et al.</i> (2005) <i>Biochemical and Biophysical Research Communications</i> 335:925-936), Melanoma inhibitory activity (Perrault <i>et al.</i> (2005) <i>Biochemical and Biophysical Research Communications</i> 335:925-936), p21, p53
Differentiation, cell fate	Transforming growth factor B2 (Perrault <i>et al.</i> (2005) <i>Biochemical and Biophysical Research Communications</i> 335:925-936)

[0079] In many cases, modulating the genes or pathways of Table 3 is undesirable because doing so can cause unwanted changes in cells. For example, TERT activates epigenetic regulators, which can change cell phenotype or interfere with efforts to reprogram or transdifferentiate cells for therapeutic purposes. TERT activates growth enhancers, but often proliferation is not desired, for example often stem cells with the most regenerative potential are those which

divide slowly. TERT modulates regulators of cell fate and differentiation, which can impair efforts to differentiate cells into specific cell types. TERT also activates proto-oncogenes, which could lead to cancer. Thus, it is desirable to minimize the amount of time during which TERT levels are artificially elevated, 5 including any treatment that extends telomeres using TERT. A treatment that extends telomeres by only transiently increasing telomerase activity levels is therefore needed.

[0080] In some cell types TERT has been shown to affect expression of other genes (Young *et al.* (2003) *J. Biol. Chem.* 278:19904-19908; Perrault *et al.* (2005) 10 *Biochem. Biophys. Res. Commun.* 335:925-936), and this may not be desirable in some cases. Thus, a treatment that minimizes the amount of time during which TERT levels are increased is needed.

Compounds

15 [0081] The present invention addresses these problems by providing in one aspect novel compounds for the transient expression of exogenous telomerase in a cell. The compounds comprise a synthetic ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase (TERT), wherein telomeres are extended within a cell treated with the compound.

20

Synthetic Ribonucleic Acids

[0082] The ribonucleic acids used in the transient expression of TERT according to various aspects of the instant invention comprise a ribonucleic acid coding for a TERT protein. The ribonucleic acids typically further comprise sequences that 25 affect the expression and/or stability of the ribonucleic acid in the cell. For example, as shown in Figure 1a, the ribonucleic acids may contain a 5' cap and untranslated region (UTR) to the 5' and/or 3' side of the coding sequence. The ribonucleic acids may further contain a 3' tail, such as a poly-A tail. The poly-A tail may, for example, increase the stability of the ribonucleic acid. In some 30 embodiments, the poly-A tail is at least 75 nucleotides, 100 nucleotides, 125 nucleotides, 150 nucleotides, or even longer.

[0083] In some embodiments, the 5' cap of the ribonucleic acid is a non-immunogenic cap. In some embodiments, the 5' cap may increase the translation of the ribonucleic acid. In some embodiments, the 5' cap may be treated with phosphatase to modulate the innate immunogenicity of the ribonucleic acid. In 5 some embodiments, the 5' cap is an anti-reverse cap analog ("ARCA"), such as a 3'-O-Me-m7G(5')ppp(5')G RNA cap structure analog.

[0084] As is well-known in the art, the above features, or others, may increase translation of the TERT protein encoded by the ribonucleic acid, may improve the stability of the ribonucleic acid itself, or may do both. In some embodiments, the 10 5' UTR and/or the 3' UTR are from a gene that has a very stable mRNA and/or an mRNA that is rapidly translated, for example, α -globin or β -globin, c-fos, or tobacco etch virus. In some embodiments, the 5' UTR and 3' UTR are from different genes, or are from different species than the species into which the 15 compositions are being delivered. The UTRs may also be assemblies of parts of UTRs from the mRNAs of different genes, where the parts are selected to achieve a certain combination of stability and efficiency of translation.

[0085] The ribonucleic acids of the invention are preferably nucleoside-modified 20 RNAs ("modRNA"). Most mature RNA molecules in eukaryotic cells contain nucleosides that are modified versions of the canonical unmodified RNA nucleosides, adenine, cytidine, guanosine, and uridine. Those modifications may prevent the RNA from being recognized as a foreign RNA. Karikó *et al.* (2005) 25 *Immunity* 23:165-175. Synthetic RNA molecules made using certain nucleosides are much less immunogenic than unmodified RNA. The immunogenicity can be reduced even further by purifying the synthetic modRNA, for example by using high performance liquid chromatography (HPLC). The modified nucleosides may be, for example, chosen from the nucleosides shown in Table 4. The nucleosides 30 are, in some embodiments, pseudouridine, 2-thiouridine, or 5-methylcytidine. Under some circumstances, it may be desirable for the modified RNA to retain some immunogenicity.

Table 4: Modified nucleosides found in eukaryotic RNA

symbol	common name
--------	-------------

m^1A	1-methyladenosine
m^6A	N^6 -methyladenosine
Am	2'- <i>O</i> -methyladenosine
i^6A	N^6 -isopentenyladenosine
io^6A	N^6 -(<i>cis</i> -hydroxyisopentenyl)adenosine
ms^2io^6A	2-methylthio- N^6 -(<i>cis</i> -hydroxyisopentenyl) adenosine
g^6A	N^6 -glyciny carbamoyl adenosine
t^6A	N^6 -threonyl carbamoyl adenosine
ms^2t^6A	2-methylthio- N^6 -threonyl carbamoyl adenosine
Ar(p)	2'- <i>O</i> -ribosyladenosine (phosphate)
m^6_2A	N^6,N^6 -dimethyladenosine
m^6Am	N^6 ,2'- <i>O</i> -dimethyladenosine
m^6_2Am	N^6,N^6 ,2'- <i>O</i> -trimethyladenosine
m^1Am	1,2'- <i>O</i> -dimethyladenosine
m^3C	3-methylcytidine
m^5C	5-methylcytidine
Cm	2'- <i>O</i> -methylcytidine
ac^4C	N^4 -acetylcytidine
f^5C	5-formylcytidine
m^4C	N^4 -methylcytidine
hm^5C	5-hydroxymethylcytidine
f^5Cm	5-formyl-2'- <i>O</i> -methylcytidine
m^1G	1-methylguanosine
m^2G	N^2 -methylguanosine
m^7G	7-methylguanosine
Gm	2'- <i>O</i> -methylguanosine
m^2_2G	N^2,N^2 -dimethylguanosine
Gr(p)	2'- <i>O</i> -ribosylguanosine (phosphate)
yW	wybutosine
o_2yW	peroxywybutosine
OHyW	hydroxywybutosine
OHyW*	undermodified hydroxywybutosine
imG	wyosine
$m^{2,7}G$	$N^2,7$ -dimethylguanosine
$m^{2,2,7}G$	$N^2,N^2,7$ -trimethylguanosine
I	inosine

m^1I	1-methylinosine
Im	2'- <i>O</i> -methylinosine
Q	queuosine
galQ	galactosyl-queuosine
manQ	mannosyl-queuosine
Ψ	pseudouridine
D	dihydrouridine
m^5U	5-methyluridine
Um	2'- <i>O</i> -methyluridine
m^5Um	5,2'- <i>O</i> -dimethyluridine
$m^1\Psi$	1-methylpseudouridine
Ψm	2'- <i>O</i> -methylpseudouridine
s^2U	2-thiouridine
ho^5U	5-hydroxyuridine
chm^5U	5-(carboxyhydroxymethyl)uridine
$mchm^5U$	5-(carboxyhydroxymethyl)uridine methyl ester
mcm^5U	5-methoxycarbonylmethyluridine
mcm^5Um	5-methoxycarbonylmethyl-2'- <i>O</i> -methyluridine
mcm^5s^2U	5-methoxycarbonylmethyl-2-thiouridine
ncm^5U	5-carbamoylmethyluridine
ncm^5Um	5-carbamoylmethyl-2'- <i>O</i> -methyluridine
$cmnm^5U$	5-carboxymethylaminomethyluridine
m^3U	3-methyluridine
$m^1acp^3\Psi$	1-methyl-3-(3-amino-3-carboxypropyl) pseudouridine
cm^5U	5-carboxymethyluridine
m^3Um	3,2'- <i>O</i> -dimethyluridine
m^5D	5-methyldihydrouridine
τm^5U	5-taurinomethyluridine
τm^5s^2U	5-taurinomethyl-2-thiouridine

[0086] Without intending to be bound by theory, the presence of the modified nucleosides enables modRNA to avoid activation of an immune response mediated by various receptors, including the Toll-like receptors and RIG-1. Non-
 5 immunogenic modRNA has been used as a therapeutic agent in mice via topical delivery. Kormann *et al.* (2011) *Nature Biotechnology* 29:154-157. The discovery

of nucleotide-modified mRNA facilitates the delivery of RNA-encoded therapeutic proteins, or mutants thereof, to cells, and the expression of those proteins in cells.

[0087] Accordingly, in some embodiments, the ribonucleic acids of the instant compositions comprise a pseudouridine, a 2-thiouridine, a 5-methylcytidine, or a 5 nucleoside from Table 4. In some embodiments, the ribonucleic acids comprise more than one of the above nucleosides or combination of the above nucleosides. In highly preferred embodiments, the ribonucleic acids comprise pseudouridine and 5-methylcytidine.

[0088] In some embodiments, an immune response to the modRNA may be 10 desired, and the RNA may be modified to induce an optimal level of innate immunity. In other embodiments, an immune response to the modRNA may not be desired, and the RNA may be modified in order to minimize such a reaction. The RNA can be modified for either situation.

[0089] The ribonucleic acids of the instant invention are preferably synthetic 15 ribonucleic acids. The term "synthetic", as used herein, means that the ribonucleic acids are in some embodiments prepared using the tools of molecular biology under the direction of a human, for example as described below. The synthetic ribonucleic acids may, for example, be prepared by *in vitro* synthesis using cellular extracts or purified enzymes and nucleic acid templates. The synthetic ribonucleic 20 acids may in some embodiments be prepared by chemical synthesis, either partially or completely. Alternatively, or in addition, the synthetic ribonucleic acids may in some embodiments be prepared by engineered expression in a cell, followed by disruption of the cell and at least partial purification of the ribonucleic acid. A synthetic ribonucleic acid is not, however, a naturally-occurring ribonucleic acid, 25 as it is expressed in an unmodified cell without extraction or purification.

[0090] The ribonucleic acids of the instant invention may be prepared using a 30 variety of standard techniques, as would be understood by one of ordinary skill in the art. In some embodiments, the ribonucleic acids may be prepared by *in vitro* synthesis, as described, for example, in U.S. Patent Application Publication Nos. 2009/0286852 and 2011/0143397. In some embodiments, the ribonucleic acids may be prepared by chemical synthesis. In some embodiments, the ribonucleic acids may be prepared by a combination of *in vitro* synthesis and chemical

synthesis. As described above, the term "synthetic" should be understood to include ribonucleic acids that are prepared either by chemical synthesis, by *in vitro* synthesis, by expression *in vivo* and at least partial purification, or by a combination of such, or other, chemical or molecular biological methods.

5 [0091] The ribonucleic acids of the instant invention may, in some embodiments, be purified. As noted above, purification may reduce immunogenicity of the ribonucleic acids and may be advantageous in some circumstances. See also U.S. Patent Application Publication No. 2011/0143397. In preferred embodiments, the ribonucleic acids are purified by HPLC or by affinity capture and elution.

10 [0092] The protein structure of TERT includes at least three distinct domains: a long extension at the amino-terminus (the N-terminal extension, NTE) that contains conserved domains and an unstructured linker region; a catalytic reverse-transcriptase domain in the middle of the primary sequence that includes seven conserved RT motifs; and a short extension at the carboxyl-terminus (the C-terminal extension, CTE). Autexier and Lue (2006) *Annu Rev Biochem.* 75:493-517. In some embodiments, the ribonucleic acid of the instant invention codes for a full-length TERT. In some embodiments, the ribonucleic acid codes for a catalytic reverse transcriptase domain of TERT. In some embodiments, the ribonucleic acid codes for a polypeptide having TERT activity.

15 [0093] The TERT encoded by the ribonucleic acids of the instant disclosure is preferably a mammalian, avian, reptilian, or fish TERT. More preferably, the TERT is a mammalian TERT, such as human TERT. Meyerson *et al.* (1997) *Cell* 90:785-795; Nakamura *et al.* (1997) *Science* 277:955-959; Wick *et al.* (1999) *Gene* 232:97-106. The amino acid sequence of two human TERT isoforms are available as NCBI Reference Sequences: NP_937983.2 and NP_001180305.1. Other non-limiting exemplary amino acid sequences usefully encoded by the ribonucleic acids of the instant compositions include TERT from cat (NCBI Reference Sequence: XP_003981636.1), dog (NCBI Reference Sequence: NP_001026800.1), mouse (NCBI Reference Sequence: NP_033380.1), cow (NCBI Reference Sequence: NP_001039707.1), sheep NCBI Reference Sequence: XP_004017220.1), pig (NCBI Reference Sequence: NP_001231229.1), African elephant (NCBI Reference Sequence: XP_003408191.1), chicken (NCBI

Reference Sequence: NP_001026178.1), rat (NCBI Reference Sequence: NP_445875.1), zebrafish (NCBI Reference Sequence: NP_001077335.1); Japanese medaka (NCBI Reference Sequence: NP_001098286.1); and chimpanzee (NCBI Reference Sequences: XP_003950543.1 and XP_003950544.1).

5 [0094] It should be understood that the ribonucleic acids of the instant invention may code for variants of any of the above-listed amino acid sequences, particularly variants that retain telomerase catalytic activity, including truncated variants. In some embodiments, the ribonucleic acids of the instant compositions code for one of the above-listed amino acid sequences or a sequence with at least 95% sequence 10 identity to that sequence. In some embodiments, the nucleic acids of the instant compositions code for one of the above-listed amino acid sequences or a sequence with at least 98%, 99%, 99.9%, or even higher sequence identity to that sequence.

15 [0095] It should also be understood that the instant ribonucleic acids may correspond to the native gene sequences coding for the above-listed TERT proteins or may correspond to variants that are made possible due to the redundancy of the genetic code, as would be understood by one of ordinary skill in the art. In some embodiments, the codon selection may be optimized to optimize protein expression using algorithms and methods known by those of ordinary skill in the art. Fath *et al.* (2011) *PLoS ONE* 6:3.

20

Compositions

25 [0096] In another aspect, the present invention provides compositions for the extension of telomeres in a cell, the compositions comprising a compound of the invention, as described above, and a further component. In some embodiments, the compositions further comprise a telomerase RNA component (TERC). (See also Table 6 below.) In some embodiments, the compositions further comprise a delivery vehicle.

Delivery Vehicles

30 [0097] As just noted, the compositions of the instant disclosure may further comprise a delivery vehicle for the ribonucleic acid. The delivery vehicle may, in some cases, facilitate targeting and uptake of the ribonucleic acid of the

composition to the target cell. In particular, the compositions of the instant disclosure may comprise any gene delivery vehicle known in the field, for example nanoparticles, liposomes, gene gun ballistic particles, viruses, cationic lipids, commercial products, such as Lipofectamine® RNAiMax, or other vehicles. In 5 some embodiments, the delivery vehicle is an exosome, a lipid nanoparticle, a polymeric nanoparticle, a natural or artificial lipoprotein particle, a cationic lipid, a protein, a protein-nucleic acid complex, a liposome, a virosome, or a polymer. In some preferred embodiments, the delivery vehicle is a cationic lipid formulation. Viral delivery is typically not preferred, however, as it can lead to insertional 10 mutagenesis.

[0098] In some preferred embodiments, the delivery vehicle is an exosome, a lipid nanoparticle, or a polymeric nanoparticle. In highly preferred embodiments, the delivery vehicle is an exosome. Exosomes are naturally-occurring lipid bilayer vesicles 40-100 nm in diameter. Exosomes contain a set of specific proteins, 15 including the membrane protein Lamp-1 and Lamp-2, which are particularly abundant. Lakhral and Wood (2011) *BioEssays: News and Reviews in Molecular, Cellular and Developmental Biology* 33:737-741. In 2007, exosomes were discovered to be natural carriers of RNA and protein, including over 1,300 types of mRNA and 121 types of non-coding microRNA. Exosomes can also transmit 20 mRNA between species: exposure of human cells to mouse exosomes carrying mouse mRNA results in translation in the human cells of the mouse mRNA.

[0099] As delivery vehicles for RNA, protein, or DNA, exosomes have a number of advantages over alternative vehicles. Specifically, exosomes can be generated from a patient's own cells, making them non-immunogenic – they are therefore not 25 attacked by antibodies, complement, coagulation factors, or opsonins. In addition, exosomes can be loaded with nucleic acids by electroporation, and they are naturally-occurring vehicles that carry mRNA and protein between human cells. Exosomes protect their RNA and protein cargo during transport, and the cargo is delivered directly into the cytosol. They can extravasate from the blood stream to 30 extravascular tissues, even crossing the blood-brain barrier, and they can be targeted. Furthermore, exosomes avoid being accumulated in untargeted organs, such as, for example, liver. Exosomes may therefore be used as cell-derived

"liposomes" to deliver therapeutic mRNA or other cargo in the treatment of disease. Mizrak *et al.* (2013) *Molecular Therapy* 21:101–108; doi:10.1038/mt.2012.161. A graphic illustration of an exosome delivering an mRNA to a cell is shown in Figure 2. See also van den Boorn *et al.* (2011) *Nature Biotechnology* 29:325-326.

5 [0100] Most cell types are believed to be capable of generating exosomes, and exosomes are found in most biological fluids including blood, saliva, urine, cerebrospinal fluid, breast milk, and amniotic fluid. Exosomes are produced by most cell types, in different abundance. Abundant exosomes, devoid of T-cell activators, can be derived from immature dendritic cells, which are present in 10 human blood. O'Doherty *et al.* (1994) *Immunology* 82:487-493. Exosomes may also be produced artificially, for example by combining recombinant exosomal proteins with lipids and phospholipids such as are found in exosomal membranes. Alternatively, exosomes may be constructed by *in vitro* self-assembly of liposomes 15 with a subset of exosomal surface proteins.

15 [0101] The drug delivery potential of exosomes was first demonstrated in 2011. Alvarez-Erviti *et al.* (2011) *Nature Biotechnology* 29:341-345. Specifically, exosomes were harvested from dendritic cells engineered to express a Lamp2B fusion protein fused to a 28 a.a. targeting ligand from rabies virus glycoprotein 20 (RVG), then electroporated siRNA into the exosomes and injected the exosomes into mice immunocompatible with the mice from which they obtained the dendritic cells. The exosomes were thus autologous, and did not generate an immune response, as measured by IL-6, IP-10, TNF- α , and IFN- α levels. Further, repeated doses over one month elicited similar responses, demonstrating that there was no 25 adaptive immune response either.

25 [0102] As described above, exosomes can be autologous and thus have low immunogenicity. Since modRNA also has low immunogenicity, the combination of modRNA as the ribonucleic acid and an exosome as the delivery vehicle in the compositions of the instant disclosure is particularly preferred. In these 30 embodiments, the disclosure thus provides a new way of delivering mRNA or modRNA to cells or tissues, using exosomes. Such delivery provides a useful method to temporarily increase the level of any protein in a cell *in vivo* using RNA

delivered in exosomes by intravenous or topical injection, and particularly in the delivery of an RNA encoding TERT. Accordingly, in preferred embodiments, the delivery vehicles of the instant compositions are non-immunogenic. Under some circumstances, however, it may be desirable for the vehicle to retain some 5 immunogenicity.

Additional Components

[0103] The compositions disclosed herein may further comprise additional components that either enhance the delivery of the composition to the target cell, 10 enhance the extension of telomeres within the cell, or both. For example, the compositions may further comprise one or more of the compounds and conditions of Table 2. As would be understood by one of ordinary skill in the art, combinations of active ingredients often display synergistic effects on a desired activity, such as, for example, the transient expression of exogenous telomerase 15 activity in a cell, and such combinations are understood to fall within the scope of the invention. Additional examples of proteins that may be included within the compositions of the instant disclosure are listed in Table 5. It should be understood that the compositions could either include the proteins themselves, or nucleic acid sequences, preferably RNAs or modRNAs, that encode these proteins, 20 or proteins with high sequence identity that retain the activity of the listed protein.

Table 5: Proteins usefully delivered in combination with TERT

Species	Protein	Activity	Advantages	Reference
Human	UPF1	Sustains telomere leading strand-replication	Increased rate or amount of telomere extension.	Chawla <i>et al.</i> (2011) <i>The EMBO Journal</i> 30:4047-4058
Human	HSP90	Prevents dephosphorylation of Akt kinase by PP2A. Akt needs to be phosphorylated to phosphorylate TERT. Also complexes with TERT and keeps TERT serine 823 phosphorylated, keeping TERT activated. Büchner <i>et al.</i> (2010) <i>Antioxidants & Redox Signaling</i> 13:551-558.	Increased TERT activity.	Haendeler <i>et al.</i> (2003) <i>FEBS Letters</i> 536:180-186; Büchner <i>et al.</i> (2010) <i>Antioxidants & Redox Signaling</i> 13:551-558
Human	Akt kinase (aka protein kinase B)	Complexes with TERT and HSP90, phosphorylates TERT at serine 823, increasing TERT	Increased TERT activity.	Wojtyla <i>et al.</i> (2011) <i>Molecular Biology Reports</i>

		activity. Büchner <i>et al.</i> (2010) <i>Antioxidants & Redox Signaling</i> 13:551-558		38:3339-3349; Büchner <i>et al.</i> (2010) <i>Antioxidants & Redox Signaling</i> 13:551-558
Human	Protein kinase C (PKC) (its various isoenzymes)	Phosphorylates TERT, allowing it to bind nuclear translocator.	Nuclear translocation.	Wojtyla <i>et al.</i> (2011) <i>Molecular Biology Reports</i> 38:3339-3349
Human	Shp-2	Inhibits phosphorylation of TERT Y707 by Src1, keeping TERT in nucleus. Transport of TERT to nucleus.	Nuclear translocation.	Wojtyla <i>et al.</i> (2011) <i>Molecular Biology Reports</i> 38:3339-3349
Human	TPP1	Recruits telomerase to the telomere.		Abreu <i>et al.</i> (2010) <i>Molecular and Cellular Biology</i> 30:2971-2982
Human	NFkB p65	Transport of TERT to nucleus.	Nuclear translocation.	Wojtyla <i>et al.</i> (2011) <i>Molecular Biology Reports</i> 38:3339-3349
Human	Rap1	regulator of telomere length.	Extension of telomeres.	O'Connor <i>et al.</i> (2004) <i>The Journal of Biological Chemistry</i> 279:28585-28591

[0104] Other examples of agents that may usefully be included within the compositions of the instant disclosure are listed in Table 6.

5

Table 6: Other agents usefully delivered in combination with TERT

Molecule	Activity	Advantages	Reference
Okadaic acid	Inhibits PP2A. PP2A dephosphorylates AKT and or TERT. AKT phosphorylates TERT, activating it.	Increased telomerase activity due to TERT phosphorylation.	Wojtyla <i>et al.</i> (2011) <i>Molecular Biology Reports</i> 38:3339-3349
TERRA or anti-sense TERRA (ARRET)	TERRA inhibits telomerase by binding to TERC, to which it is complementary.	Antisense TERRA, or ARRET, should increase telomerase activity by binding to TERRA, preventing it from binding to TERC.	Cifuentes-Rojas and Shippen (2012) <i>Mutat. Res.</i> 730:20-27; doi:10.1016/j.mrfmmm.2011.10.003
TERC RNA	RNA component of telomerase, essential for its	Increase telomerase activity.	

	function, may be second-most limiting factor after TERT in most cells.		
--	--	--	--

[0105] Since TERT is most active during certain phases of the cell cycle, the compositions of the instant disclosure may also optionally include one or more transient activators of cellular proliferation, in order to enhance the effectiveness of 5 the TERT treatment. Such agents may include, for example, an RNAi agent that transiently reduces the amounts of cell cycle inhibitors such as Rb or P19/Arf in the cell. Other transient activators of cellular proliferation may be usefully included in the instant compositions, as would be understood by one of ordinary skill in the art.

10

Methods of Extending Telomeres and Methods of Treatment

[0106] In another aspect, the instant disclosure provides methods of extending telomeres, comprising the step of administering any of the above-described compounds or compositions to a cell with shortened telomeres, wherein telomeres 15 are extended within the cell. The instant disclosure also provides methods of treatment, comprising the step of administering any of the above-described compounds or compositions to an animal subject in need of, or that may benefit from, telomere extension.

[0107] In preferred embodiments, the compounds or compositions are 20 administered to a cell, wherein the cell is an isolated cell or is part of a cell culture, an isolated tissue culture, an isolated organ, or the like (*i.e.*, the administration is *in vitro*).

[0108] In other preferred embodiments, the compounds or compositions are administered without isolating the cell or cells, the tissue, or the organ from the 25 subject (*i.e.*, the administration is *in vivo*). In some of these embodiments, the compound or composition is delivered to all, or almost all, cells in the subject's body. In some embodiments, the compound or composition is delivered to a specific cell or tissue in the subject's body.

[0109] In some embodiments, the subject is a mammal, bird, fish, or reptile. In 30 preferred embodiments, the subject is a mammal and more preferably a human. In

other preferred embodiments, the subject is a pet animal, a zoo animal, a livestock animal, or an endangered species animal. Examples of preferred subject species are listed in Table 7.

5 **Table 7: Subject animal species.**

Dog (<i>Canis lupus familiaris</i>)
Sheep (<i>Ovis aries</i>)
Domestic pig (<i>Sus scrofa domesticus</i>)
Domestic goat (<i>Capra aegagrus hircus</i>)
Cattle (<i>Bos primigenius taurus</i>)
Zebu (<i>Bos primigenius indicus</i>)
Cat (<i>Felis catus</i>)
Chicken (<i>Gallus gallus domesticus</i>)
Guinea pig (<i>Cavia porcellus</i>)
Donkey (<i>Equus africanus asinus</i>)
Domesticated duck (<i>Anas platyrhynchos domesticus</i>)
Water buffalo (<i>Bubalus bubalis</i>)
Horse (<i>Equus ferus caballus</i>)
Domesticated Silkmoth (<i>Bombyx mori</i>)
Domestic Pigeon (<i>Columba livia domestica</i>)
Domestic goose (<i>Anser anser domesticus</i>)
Llama (<i>Lama glama</i>)
Alpaca (<i>Vicugna pacos</i>)
Domesticated guineafowl (<i>Numida meleagris</i>)
Ferret (<i>Mustela putorius furo</i>)
Ringneck dove (<i>Streptopelia risoria</i>)
Bali cattle (<i>Bos javanicus domestica</i>)
Gayal (<i>Bos frontalis</i>)
Domesticated turkey (<i>Meleagris gallopavo</i>)
Goldfish (<i>Carassius auratus auratus</i>)
Domestic rabbit (<i>Oryctolagus cuniculus</i>)
Domestic Canary (<i>Serinus canaria domestica</i>)
Carabao (<i>Bubalus bubalis carabensis</i>)
Siamese fighting fish (<i>Betta splendens</i>)
Koi (<i>Cyprinus carpio haematopterus</i>)
Domesticated silver fox (<i>Vulpes vulpes</i>)
Domesticated hedgehog (<i>Atelerix albiventris</i>)
Society Finch (<i>Lonchura striata domestica</i>)
Yak (<i>Bos grunniens</i>)
Fancy rat and Lab rat
Domesticated Dromedary Camel (<i>Camelus dromedarius</i>)
Domesticated Bactrian Camel (<i>Camelus bactrianus</i>)
Guppy (<i>Poecilia reticulata</i> some strains)
Fancy mouse

[0110] For *in vitro* applications, the compounds or compositions may be administered using any suitable technique, as would be understood by those skilled in the fields of cell biology, cell culture, tissue culture, organ culture, or the like.

5 For *in vivo* applications, the compounds or compositions are usefully administered by injection, topical application, inhalation, or any other suitable administration technique, as would be understood by those of ordinary skill in the medical arts or the like.

[0111] As described above, cells usefully treated according to the methods of the 10 disclosure include cells, either in a subject (for *in vivo* administration) or from a subject (for *in vitro* administration), that may benefit from telomere extension.

Since short telomeres affect almost all cell types in most animals, telomere extension may benefit most animals. A telomere extension treatment that is transient and brief has the potential to be safe, as described above. Telomere 15 extension using a transient treatment as disclosed herein may therefore be of use in all or most individuals either as a preventive measure, for example to prevent or delay onset of the many diseases and conditions in which short telomeres are implicated, or as a treatment for those diseases and conditions. The treatment may benefit subjects at risk of age-related diseases or conditions, or who are already 20 suffering from such diseases, and may also benefit subjects who have experienced, are experiencing, or are at risk of experiencing physical trauma or chronic physical stress such as hard exercise or manual labor, or psychological trauma or chronic psychological stress, since all of these conditions cause telomere shortening; physical stress or trauma requires cell division in order to repair the resultant 25 damage, thus shortening telomeres, and these conditions may also cause oxidative stress, which also shortens telomeres. Such diseases and conditions include, for example, metabolic syndrome, diabetes, diabetic ulcers, heart disease, many forms of cancer, vascular dementia, Alzheimer's, stroke, age-related macular degeneration, immunosenescence, bone marrow failure, gastrointestinal ulcers, 30 cirrhosis, hernia, infection such as pneumonia secondary to impaired immune function, chronic infection, mild or severe cognitive impairment, impaired mobility, osteoporosis, osteoarthritis, rheumatoid arthritis, age-related anxiety,

balance disorders, tinnitus, Bell's palsy, cataracts, COPD, corneal abrasion, coronary artery disease, peripheral artery disease, conjunctivitis, chalazion, dehydration, depression, emphysema, various eye diseases, failure to thrive, flu, generalized anxiety disorder, glaucoma, hearing loss, loss of sense of taste, loss of 5 appetite, hip dislocation, memory loss, Parkinson's disease, spinal stenosis, urinary incontinence, vertebral fracture, and others.

[0112] Cells usefully treated according to the instant methods may also include 10 cells in or from a subject that suffers from an age-related illness or condition or that is at risk of suffering from such an illness or condition. Such age-related illnesses and conditions may include genetic diseases in which genes encoding components of the telomerase complex or telomere complex are mutated, thus leading to short telomeres. Such diseases include, for example, forms of idiopathic pulmonary fibrosis, dyskeratosis congenita, and aplastic anemia.

[0113] In the case of genetic diseases, the compositions may therefore further 15 include additional nucleic acids, such as the normal, functional versions of the coding sequences of the genes that are affected by such diseases, for example DKC1, TINF2, NOP10, NHP2, TERC, or other genes. Examples of such genes are listed in Table 8. See also Armanios (2009) *Ann. Rev. Genomics Hum. Genet.* 10: 45-61.

20

Table 8: Genes related to age-related diseases.

Gene	Diagnosis	Age of onset in years (typical)
<i>hTR</i> <i>hTERT</i>	Sporadic IPF 1-3% Familial IPF 8-15% Sporadic and familial aplastic anemia ~3-5% Autosomal dominant dyskeratosis congenital (DC)	Broad range of ages 5-77
<i>DKC1</i>	X-linked DC Hoyeraal-Hreiderasson	Less than 30 Less than 5
<i>TINF2</i>	Sporadic DC Autosomal dominant DC Hoyeraal-Hreiderasson	Less than 10 - Less than 5
<i>NOP10</i>	Autosomal Recessive DC	-
<i>NHP2</i>	Autosomal Recessive DC	-

[0114] In addition, the treatment may benefit subjects suffering from, or at risk of, other types of genetic diseases in which short telomeres may play a role, such as, for example, muscular dystrophy. In such diseases, the need for cell replication to address the problem caused by the genetic mutation shortens telomeres more rapidly than normal, resulting in more rapid telomere shortening than normal, which in turn exhausts the replicative capacity of cells, leading to tissue dysfunction, exacerbated or additional symptoms, disability, and often death. In addition, various types of cancer may be prevented or delayed by treatment with compounds of the invention, and indeed chromosome-chromosome fusions caused by critically short telomeres are believed to be a cause of cancer. Telomeres may also be selectively lengthened in healthy cells in an individual, while not lengthening telomeres in cancer cells, which may allow the instant compounds, compositions, and methods to be used, for example, to lengthen telomeres of the immune system to increase its ability to fight a cancer. Further, immune system cells may be harvested from an individual for treatment using the invention *ex vivo* followed by reintroduction into the individual.

[0115] In some embodiments, the cells treated according to the instant methods are from subjects where no disease state is yet manifested but where the subject is at risk for a condition or disease involving short telomeres, or where the cells contain shortened telomeres. In some embodiments, the age-related illness is simply old age. The instant treatments may also be used as a cosmetic aid, to prevent, delay, or ameliorate age-related deterioration in appearance of skin, hair, bone structure, posture, eye clarity, or other traits that decline with aging. For example, the treatments may be used to help maintain skin elasticity, thickness, smoothness, and appearance, since telomere extension improves these parameters. In cases of physical trauma such as a bone fracture or a tissue crush or cut injury or burn, the invention may be used to increase the lengths of telomeres in cells which participate in healing the trauma, to increase their replicative capacity. In cases of chronic physical stress, which causes telomere shortening, treatment with the invention may lengthen telomeres in affected cells increasing their replicative capacity and ability to repair tissue damage. Since telomere shortening accumulates over generations, for example in humans with haploinsufficiency of

telomerase components such as hTR or hTERT, Armanios (2009) *Annu. Rev. Genomics Hum. Genet.* 10:45-61, the treatments of the instant disclosure may be applied to germ line cells such as eggs, sperm, or their precursors, or to fertilized eggs or embryos, for example during *in vitro* fertilization procedures. The 5 treatments may also be useful for aiding other treatments of various diseases or conditions, for example, transdifferentiation of cells *in vivo*. The treatment methods may also be useful in advance of or during surgery or chemotherapy, or radiotherapy, to increase the ability of cells to replicate to repair damage resulting from these procedures.

10 [0116] The methods may also be useful for treating cells *in vitro* for various applications, including autologous or heterologous cell therapy, bioengineering, tissue engineering, growth of artificial organs, generation of induced pluripotent stem cells (iPSC), and cellular differentiation, dedifferentiation, or transdifferentiation. In these applications, cells may be required to divide many 15 times, which may lead to loss of telomere length, which may be counteracted by the invention before, during, or after the application.

[0117] In addition, various types of cancer may be considered age-related illnesses, particularly where the cancerous cells contain short telomeres. In some cases, the cells treated according to the instant methods are from subjects where no 20 disease state is yet manifested but where the cells contain shortened telomeres. In some embodiments, the age-related illness is simply the altered form and function typically associated with old chronological age in humans.

[0118] The cells usefully administered compounds or compositions of the instant disclosure according to the instant methods include cells from any tissue or cell 25 type that may suffer the effects of shortened telomeres or that may in any way benefit from lengthening of the cell's telomeres. Cells may include somatic cells or germ cells, as well as stem cells and other progenitor cells and/or undifferentiated cells. Cells may include tumor cells and non-tumor cells.

[0119] Examples of cells usefully administered compounds or compositions 30 according to the instant methods include cells that are derived primarily from endoderm, cells that are derived primarily from ectoderm, and cells that are derived primarily from mesoderm. Cells derived primarily from the endoderm

include, for example, exocrine secretory epithelial cells and hormone-secreting cells. Cells derived primarily from the ectoderm include, for example, cells of the integumentary system (*e.g.*, keratinizing epithelial cells and wet stratified barrier epithelial cells) and the nervous system (*e.g.*, sensory transducer cells, autonomic neuron cells, sense organ and peripheral neuron supporting cells, central nervous system neurons and glial cells, and lens cells). Cells derived primarily from the mesoderm include, for example, metabolism and storage cells, barrier-function cells (*e.g.*, cells of the lung, gut, exocrine glands, and urogenital tract), extracellular matrix cells, contractile cells, blood and immune system cells, germ cells, nurse cells, and interstitial cells. Accordingly, in some embodiments of the instant methods, the cell administered the composition is a somatic cell of endodermal, mesodermal, or ectodermal lineage. In some embodiments the cell is a germ line cell or an embryonic cell.

[0120] Specific examples of cells that may be administered a compound or composition according to the instant methods include, *e.g.*, salivary gland mucous cells, salivary gland serous cells, von Ebner's gland cells in tongue, mammary gland cells, lacrimal gland cells, ceruminous gland cells in ear, eccrine sweat gland dark cells, eccrine sweat gland clear cells, apocrine sweat gland cells, gland of Moll cells in eyelid, sebaceous gland cells, Bowman's gland cells in nose, Brunner's gland cells in duodenum, seminal vesicle cells, prostate gland cells, bulbourethral gland cells, Bartholin's gland cells, gland of Littre cells, uterus endometrium cells, isolated goblet cells of the respiratory and digestive tracts, stomach lining mucous cells, gastric gland zymogenic cells, gastric gland oxyntic cells, pancreatic acinar cells, paneth cells of the small intestine, type II pneumocytes of the lung, clara cells of the lung, anterior pituitary cells (*e.g.*, somatotropes, lactotropes, thyrotropes, gonadotropes, and corticotropes), intermediate pituitary cells (*e.g.*, those secreting melanocyte-stimulating hormone), magnocellular neurosecretory cells (*e.g.*, those secreting oxytocin or vasopressin), gut and respiratory tract cells, (*e.g.*, those secreting serotonin, endorphin, somatostatin, gastrin, secretin, cholecystokinin, insulin, glucagon, or bombesin), thyroid gland cells (*e.g.*, thyroid epithelial cells and parafollicular cells), parathyroid gland cells (*e.g.*, parathyroid chief cells and oxyphil cells), adrenal

gland cells (e.g., chromaffin cells and cells secreting steroid hormones such as mineralcorticoids and glucocorticoids), Leydig cells of testes, theca interna cells of the ovarian follicle, corpus luteum cells of the ruptured ovarian follicle, granulosa lutein cells, theca lutein cells, juxtaglomerular cells, macula densa cells of the 5 kidney, peripolar cells of the kidney, mesangial cells of the kidney, epidermal keratinocytes, epidermal basal cells, keratinocytes of fingernails and toenails, nail bed basal cells, medullary hair shaft cells, cortical hair shaft cells, cuticular hair shaft cells, cuticular hair root sheath cells, hair root sheath cells of Huxley's layer, hair root sheath cells of Henle's layer, external hair root sheath cells, hair matrix 10 cells, surface epithelial cells of the stratified squamous epithelium of the cornea, tongue, oral cavity, esophagus, anal canal, distal urethra, and vagina, basal cells of the epithelia of the cornea, tongue, oral cavity, esophagus, anal canal, distal urethra, and vagina, urinary epithelium cells (e.g., lining the urinary bladder and 15 urinary ducts), auditory inner hair cells of the organ of Corti, auditory outer hair cells of the organ of Corti, basal cells of the olfactory epithelium, cold-sensitive primary sensory neurons, heat-sensitive primary sensory neurons, Merkel cells of the epidermis, olfactory receptor neurons, pain-sensitive primary sensory neurons, photoreceptor cells of the retina in the eye (e.g., photoreceptor rod cells, photoreceptor blue-sensitive cone cells, photoreceptor green-sensitive cone cells, 20 and photoreceptor red-sensitive cone cells), proprioceptive primary sensory neurons, touch-sensitive primary sensory neurons, type I and II carotid body cells, type I and II hair cells of the vestibular apparatus of the ear, type I taste bud cells, cholinergic neural cells, adrenergic neural cells, peptidergic neural cells, inner and outer pillar cells of the organ of Corti, inner and outer phalangeal cells of the organ 25 of Corti, border cells of the organ of Corti, Hensen cells of the organ of Corti, vestibular apparatus supporting cells, taste bud supporting cells, olfactory epithelium supporting cells, Schwann cells, satellite glial cells, enteric glial cells, astrocytes, neuron cells, oligodendrocytes, spindle neurons, anterior lens epithelial cells, crystallin-containing lens fiber cells, hepatocytes, adipocytes (e.g., white fat 30 cells and brown fat cells), liver lipocytes, kidney parietal cells, kidney glomerulus podocytes, kidney proximal tubule brush border cells, loop of Henle thin segment cells, kidney distal tubule cells, kidney collecting duct cells, type I pneumocytes,

pancreatic duct cells, nonstriated duct cells (*e.g.*, principal cells and intercalated cells), duct cells (of seminal vesicle, prostate gland, etc.), intestinal brush border cells (with microvilli), exocrine gland striated duct cells, gall bladder epithelial cells, ductulus efferens nonciliated cells, epididymal principal cells, epididymal 5 basal cells, ameloblast epithelial cells, planum semilunatum epithelial cells of the vestibular apparatus of the ear, organ of Corti interdental epithelial cells, loose connective tissue fibroblasts, corneal fibroblasts (corneal keratocytes), tendon fibroblasts, bone marrow reticular tissue fibroblasts, other nonepithelial fibroblasts, pericytes, nucleus pulposus cells of the intervertebral disc,

10 cementoblasts/cementocytes, odontoblasts/odontocytes, hyaline cartilage chondrocytes, fibrocartilage chondrocytes, elastic cartilage chondrocytes, osteoblasts/osteocytes, osteoprogenitor cells, hyalocytes of the vitreous body of the eye, stellate cells of the perilymphatic space of the ear, hepatic stellate cells (Ito cells), pancreatic stelle cells, skeletal muscle cells (*e.g.*, red skeletal muscle cells 15 (slow) and white skeletal muscle cells (fast)), intermediate skeletal muscle cells, nuclear bag cells of the muscle spindle, nuclear chain cells of the muscle spindle, satellite cells, heart muscle cells (*e.g.*, ordinary heart muscle cells, nodal heart muscle cells, and Purkinje fiber cells, smooth muscle cells (various types), myoepithelial cells of the iris, myoepithelial cells of the exocrine glands,

20 erythrocytes, megakaryocytes, monocytes, connective tissue macrophages (various types), epidermal Langerhans cells, osteoclasts, dendritic cells, microglial cells, neutrophil granulocytes, eosinophil granulocytes, basophil granulocytes, hybridoma cells, mast cells, helper T cells, suppressor T cells, cytotoxic T cells, natural killer T cells, B cells, natural killer cells, reticulocytes, stem cells and 25 committed progenitors for the blood and immune system (various types), oogonia/oocytes, spermatids, spermatocytes, spermatogonium cells, spermatozoa, nurse cells, ovarian follicle cells, sertoli cells, thymus epithelial cells, and interstitial kidney cells.

[0121] In a preferred embodiment, the cells administered a compound or 30 composition according to the instant methods are stem or progenitor cells, since these cells give rise to other cells of the body. In another preferred embodiment, the cells treated are cells in which telomeres shorten more quickly than in other

cell types, for example endothelial cells, fibroblasts, keratinocytes, cells of the immune system including thyroid and parathyroid cells and leukocytes and their progenitors, cells of the intestines, liver, mucosal membrane cells, *e.g.* in the esophagus and colon, and cells of the gums and dental pulp.

5 [0122] Accordingly, in some embodiments, the cells are fibroblast cells, keratinocytes, endothelial cells, epithelial cells, or blood cells.

[0123] The administering step may be performed one or more times depending on the amount of telomere extension desired. In some embodiments of the instant methods, the cell is an isolated cell, and the administering step lasts no longer than 10 96 hours, no longer than 72 hours, no longer than 48 hours, no longer than 36 hours, no longer than 24 hours, no longer than 18 hours, no longer than 12 hours, no longer than 8 hours, no longer than 4 hours, or even shorter times. In some embodiments, the administering step lasts at least 2 hours, at least 4 hours, at least 8 hours, at least 12 hours, at least 18 hours, at least 24 hours, at least 36 hours, at 15 48 hours, or even longer times. In preferred embodiments, the administering step lasts no longer than 48 hours, no longer than 96 hours, or no longer than 1 week. In other preferred embodiments, the administering step lasts at least 2 hours. It should be understood that, in the case where administration is by transfection, the time for administration includes the time for the cell to recover 20 from the transfection method.

[0124] In some embodiments of the instant methods, the cell is an isolated cell, and the administering step is performed no more than 6 times, no more than 5 times, no more than 4 times, no more than 3 times, no more than 2 times, or even no more than 1 time. In some embodiments, the administering step is performed 25 not less than 2 times, not less than 3 times, not less than 4 times, not less than 5 times, not less than 6 times, or even more often.

[0125] In some embodiments, the administering step is performed once or a few times over a relatively brief period to re-extend telomeres, and then not performed for a prolonged period until telomeres need to be extended again. This cycle may 30 be repeated indefinitely. Such a treatment schedule allows telomeres to be periodically re-extended, with intervals in between administration steps during which telomeres shorten. Periodic treatment methods may be performed either by

in vivo administration or by *in vitro* administration, as desired. In some embodiments, the administering step in such a series is performed no more than 6 times, no more than 5 times, no more than 4 times, no more than 3 times, no more than 2 times, or even no more than 1 time. In some embodiments, the

5 administering step is performed not less than 2 times, not less than 3 times, not less than 4 times, not less than 5 times, not less than 6 times, or even more often. By varying the number of times the administering step is performed, and the dose of the compounds our compositions of the invention used, the amount of telomere extension achieved can be controlled.

10 [0126] In some embodiments, the methods of the instant disclosure further include the step of culturing the cell on a specific substrate, preferably an elastic substrate. Such substrates are known to prevent unwanted changes in the cell that would normally occur on other substrates due to the non-physiological elasticity of those substrates. See PCT International Publication No. WO2012/009682.

15 15 Elastic substrates may additionally promote cell survival.

[0127] Administration of the compounds or compositions of the instant disclosure results in the transient expression of a telomerase activity in the cell. The increased activity is readily measured by various assays, such as, for example, the Trapeze[®] RT telomerase detection kit (Millipore), which provides a sensitive, real-time *in vitro* assay using fluorimetric detection and quantitation of telomerase activity, although other measurement techniques are also possible. In some embodiments, the telomerase activity is increased by at least 5%, at least 10%, at least 20%, at least 30%, at least 50%, or even more. In preferred embodiments, the telomerase activity is increased by at least 5%.

20 25 [0128] As previously noted, one of the advantages of the instant techniques is that the expression of telomerase activity is transient in the treated cells. In particular, such transient expression is in contrast to previous techniques where a telomerase reverse transcriptase gene is inserted into the genomic sequence of the cell or otherwise permanently modifies the genetic make-up of the targeted cell and results in constitutive activity of the nucleic acid sequence.

[0129] Figure 3 graphically illustrates some of the advantages of the compounds, compositions, and methods disclosed herein. In particular, the speed of telomere extension made possible with these compounds, compositions, and methods enables telomere maintenance by very infrequent delivery of TERT modRNA.

5 The expressed telomerase activity rapidly extends telomeres in a brief period, before being turned over, thus allowing the protective anti-cancer mechanism of telomere-shortening to function most of the time. Between treatments, normal telomerase activity and telomere shortening is present, and therefore the anti-cancer safety mechanism of telomere shortening to prevent out-of-control

10 proliferation remains intact, while the risk of short telomere-related disease remains low. In contrast, the best existing small molecule treatment for extending telomeres requires chronic delivery, and thus presents a chronic cancer risk, and even then has a small, inconsistent effect on telomere length, with no detectable effect on telomere length at all in about half of patients.

15 [0130] Accordingly, in some embodiments of the instant methods, the expression of telomerase reverse transcriptase activity, *i.e.*, the half-life of telomerase activity, lasts no longer than 48 hours, no longer than 36 hours, no longer than 24 hours, no longer than 18 hours, no longer than 12 hours, no longer than 8 hours, no longer than 4 hours, or even shorter times. In some embodiments, the expression of

20 telomerase reverse transcriptase activity lasts at least 2 hours, at least 4 hours, at least 8 hours, at least 12 hours, at least 18 hours, at least 24 hours, at least 36 hours, at least 48 hours, or even longer times. In preferred embodiments, the expression of telomerase reverse transcriptase activity lasts no longer than 48 hours. In other preferred embodiments, the expression of telomerase reverse transcriptase activity

25 lasts at least 2 hours.

[0131] In some embodiments of the instant methods, the transient expression is independent of cell cycle.

[0132] As noted above, the transient expression of telomerase reverse transcriptase results in the extension of shortened telomeres in treated cells.

30 Telomere length can be readily measured using techniques such as terminal restriction fragment (TRF) length analysis, qPCR, MMqPCR, and Q-FISH, as would be understood by one of ordinary skill in the art. See, *e.g.*, Kimura *et al.*

(2010) *Nat Protoc.* 5:1596-607; doi: 10.1038/nprot.2010.124. In some embodiments, the instant methods increase telomere length in treated cells by at least 0.1 kb, at least 0.2 kb, at least 0.3 kb, at least 0.4 kb, at least 0.5 kb, at least 1 kb, at least 2 kb, at least 3 kb, at least 4 kb, at least 5 kb, or even more.

5 [0133] One of the advantages of the instant compounds, compositions, and methods, is the rapidity of extension of telomeres achieved by these techniques. The techniques allow treatments to be brief and thus safe because the normal protective telomere shortening mechanism remains intact for most of the time. Treatment with the compounds and compositions disclosed herein result in 10 delivery of tens or hundreds of copies of TERT modRNA per cell as measured by absolute RT-qPCR, which is substantially more than the average number of copies of endogenous TERT mRNA found even in cells with high telomerase activity. Typically such cells have less than one copy of TERT mRNA per cell (Yi *et al.* 15 (2001) *Nucl. Acids Res.* 29:4818-4825). Thus the treatments transiently introduce a large number of copies of modRNA encoding TERT to a cell resulting in rapid telomere extension. Without intending to be bound by theory, the large number of copies of modRNA encoding TERT may transiently overwhelm the inhibitory 20 regulatory mechanisms that normally prevent TERT, and other methods of telomere extension, from extending telomeres as rapidly as the compounds, compositions, and methods disclosed herein.

[0134] The transient expression of telomerase reverse transcriptase also results in an increased replicative capacity in treated cells. Increased replicative capacity is readily monitored in cells that are approaching replicative senescence by measuring additional population doublings in such cells. Senescent cells are not 25 stimulated to divide by passage in culture or treatment with serum. Senescent cells are further often characterized by the expression of pH-dependent β -galactosidase activity, expression of cell cycle inhibitors p53 and p19, and other altered patterns of gene expression, and an enlarged cell size. Absent treatment with the compounds and compositions of the instant disclosure, human lung fibroblast cells 30 typically double 50-60 times. With one set of one to three treatments lasting only a few days total, however, these cells achieve an additional 16-28 population doublings. If treated again several weeks later, additional proliferative capacity is

conferred again. This process of intermittent treatments to periodically re-extend telomeres may be applied additional times, with the interval between treatments depending on factors such as the rate of telomere shortening, the rate of cell divisions, and the amount of telomere extension provided by the treatment.

5 Likewise, human microvascular dermal endothelial cells from an aged individual, absent treatment with the instant compositions, may achieve only 1-2 population doublings, whereas treated cells may achieve 3, 4, or even more population doublings.

[0135] Accordingly, in some embodiments, the instant treatment methods
10 increase the number of population doublings by at least one, two, four, or even more population doublings. In some embodiments, the treatment methods increase the number of population doublings by at least 5, 10, 15, 20, or even more population doublings.

[0136] In some of the instant method embodiments, the compounds or
15 compositions of the invention are administered to the animal cell by electroporation. In specific embodiments, a compound of the invention is administered to the animal cell by electroporation in the absence of a delivery vehicle. In other specific embodiments a compound of the invention and a telomerase RNA component are administered to the animal cell by electroporation.

20

Kits

[0137] In another aspect, the instant disclosure provides ready-to-use kits for use in extending telomeres in a mammalian cell. The kits comprise any of the above-described compounds or compositions, together with instructions for their use. In
25 some embodiments, the kits further comprise packaging materials. In preferred embodiments, the packaging materials are air-tight. In these embodiments, the packaging materials may optionally be filled with an inert gas, such as, for example, nitrogen, argon, or the like. In some embodiments, the packaging materials comprise a metal foil container, such as, for example, a sealed aluminum
30 pouch or the like. Such packaging materials are well known by those of ordinary skill in the art.

[0138] In some embodiments, the kit may further comprise a desiccant, a culture medium, an RNase inhibitor, or other such components. In some embodiments, the kit may further comprise a combination of more than one of these additional components. In some kit embodiments, the composition of the kit is sterile.

5

Further Aspects

[0139] In yet another aspect, the invention provides novel compounds, compositions, kits, and methods according to the following numbered paragraphs:

1. A composition for the extension of telomeres comprising:

10 a ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase; and

a delivery vehicle for the ribonucleic acid;

wherein telomeres are extended within a cell treated with the composition.

2. The composition of paragraph 1, wherein the telomerase reverse

15 transcriptase is a mammalian, avian, reptilian, or fish telomerase reverse transcriptase or a variant that retains telomerase catalytic activity.

3. The composition of paragraph 2, wherein the telomerase reverse transcriptase is a human telomerase reverse transcriptase.

4. The composition of paragraph 1, wherein the ribonucleic acid comprises a 20 5' cap, a 5' untranslated region, a 3' untranslated region, and a poly-A tail.

5. The composition of paragraph 4, wherein the poly-A tail increases stability of the ribonucleic acid.

25 6. The composition of paragraph 4, wherein the 5' untranslated region or the 3' untranslated region comprise a sequence from a stable mRNA or an mRNA that is efficiently translated.

7. The composition of paragraph 4, wherein the 5' untranslated region and the 3' untranslated region both comprise a sequence from a stable mRNA or an mRNA that is efficiently translated.

30 8. The composition of paragraph 4, wherein the 5' cap, the 5' untranslated region, or the 3' untranslated region stabilizes the ribonucleic acid, increases the rate of translation of the ribonucleic acid, or reduces the immunogenicity of the ribonucleic acid.

9. The composition of paragraph 1, wherein the at least one modified nucleoside reduces immunogenicity of the ribonucleic acid.
10. The composition of paragraph 1, wherein the ribonucleic acid is a synthetic ribonucleic acid.
- 5 11. The composition of paragraph 10, wherein the synthetic ribonucleic acid is a purified synthetic ribonucleic acid.
12. The composition of paragraph 11, wherein the synthetic ribonucleic acid is purified to remove immunogenic components.
13. The composition of paragraph 1, wherein the ribonucleic acid codes for a
- 10 human, cat, dog, mouse, cow, sheep, pig, African elephant, chicken, rat, zebrafish, Japanese medaka, or chimpanzee telomerase reverse transcriptase, or a polypeptide with at least 95% sequence identity to the telomerase reverse transcriptase.
14. The composition of paragraph 1, wherein the composition further comprises a telomerase RNA component.
15. The composition of paragraph 14, wherein the telomerase RNA component is a mammalian, avian, reptilian, or fish telomerase RNA component.
16. The composition of paragraph 15, wherein the telomerase RNA component is a human telomerase RNA component.
17. The composition of paragraph 1, wherein the delivery vehicle is an
- 20 exosome, a lipid nanoparticle, a polymeric nanoparticle, a natural or artificial lipoprotein particle, a cationic lipid, a protein, a protein-nucleic acid complex, a liposome, a virosome, or a polymer.
18. The composition of paragraph 17, wherein the delivery vehicle is a cationic lipid.
- 25 19. The composition of paragraph 1, wherein the delivery vehicle is non-immunogenic.
20. A method of extending telomeres, comprising the step of: administering the composition of any one of paragraphs 1 to 19 to an animal cell, wherein at least one telomere is extended within the cell.
- 30 21. The method of paragraph 20, wherein the cell has at least one shortened telomere prior to the administering step.

22. The method of paragraph 20, wherein the cell is from a subject suffering from or at risk of an age-related illness, an age-related condition, or an age-related decline in function or appearance.
23. The method of paragraph 20, wherein the cell is from a subject suffering from or at risk of cancer, heart disease, stroke, diabetes, Alzheimer's disease, osteoporosis, a decline in physical ability or appearance, physical trauma or chronic physical stress, or psychological trauma or chronic psychological stress.
- 5 24. The method of paragraph 20, wherein the cell is a somatic cell of endodermal, mesodermal, or ectodermal lineage, or a germ line or embryonic cell.
- 10 25. The method of paragraph 20, wherein the cell is an induced pluripotent stem cell or a cell used to produce an induced pluripotent stem cell.
26. The method of paragraph 20, wherein the cell is a transdifferentiated cell or a cell used to produce a transdifferentiated cell.
- 15 27. The method of paragraph 20, wherein the cell is an isolated cell, and the administering step lasts no longer than 48 hours.
28. The method of paragraph 20, wherein the cell is an isolated cell, and the administering step is performed no more than four times.
29. The method of paragraph 20, wherein the cell is an isolated cell, and the method further comprises the step of measuring telomerase activity in the cell.
- 20 30. The method of paragraph 29, wherein the administering step increases telomerase activity in the cell.
31. The method of paragraph 30, wherein the telomerase activity is transiently increased by at least 5%.
32. The method of paragraph 30, wherein the half-life of increased telomerase 25 activity lasts no longer than 48 hours.
33. The method of paragraph 20, wherein the cell is an isolated cell, and the method further comprises the step of measuring average telomere length in the cell.
34. The method of paragraph 33, wherein average telomere length is increased by at least 0.1 kb.
- 30 35. The method of paragraph 20, wherein the cell is an isolated cell, and the method further comprises the step of measuring population doubling capacity in the cell.

36. The method of paragraph 35, wherein the population doubling capacity increases by at least 25%.
37. The method of paragraph 20, wherein the cell is from a mammalian subject.
38. The method of paragraph 37, wherein the cell is from a human subject.
- 5 39. The method of paragraph 20, wherein the cell is an isolated cell.
40. The method of paragraph 20, wherein the cell is not an isolated cell.
41. A kit for extending telomeres in an animal cell, comprising:
the composition of any one of paragraphs 1 to 19; and
instructions for using the composition to extend telomeres.
- 10 42. The kit of paragraph 41, further comprising packaging materials.
43. The kit of paragraph 42, wherein the packaging materials are air-tight.
44. The kit of paragraph 42, wherein the packaging materials comprise a metal foil container.
45. The kit of paragraph 41, further comprising a desiccant.
- 15 46. The kit of paragraph 41, further comprising a culture medium.
47. The kit of paragraph 41, further comprising an RNase inhibitor.
48. The kit of paragraph 41, wherein the composition is sterile.

[0140] It will be readily apparent to one of ordinary skill in the relevant arts that other suitable modifications and adaptations to the compounds, compositions, methods, and kits described herein may be made without departing from the scope of the invention or any embodiment thereof. Having now described the present invention in detail, the same will be more clearly understood by reference to the following Examples, which are included herewith for purposes of illustration only and are not intended to be limiting of the invention.

EXAMPLES

Example 1. Highly efficient telomere extension in human cells using modified mRNA encoding telomerase

30 [0141] Diseases of inadequate telomere length maintenance and the need for increased cell replicative capacity for cell therapies and bioengineering applications motivate development of safe methods for telomere extension.

Blackburn *et al.* (2010) *Cancer Prev Res (Phila)* 3:394–402; Calado *et al.* (2012) *Leukemia* 26:700–707; Alter *et al.* (2009) *Blood* 113:6549–6557; Mohsin *et al.* (2012) *Journal of the American College of Cardiology* doi:10.1016/j.jacc.2012.04.047. mRNA delivery to transiently increase the amount of protein encoded by the mRNA for therapeutic applications is facilitated by incorporation of modified nucleosides reduce immunogenicity and increase stability. Karikó *et al.* (2005) *Immunity* 23:165–175; Karikó *et al.* (2011) *Nucleic Acids Res.* 39:e142; doi: 10.1093/nar/gkr695.

5 [0142] To be therapeutically useful, a telomere extension treatment should ideally be non-immunogenic; specific; capable of being initiated before telomeres shorten to the critically short lengths that cause chromosomal instability and increased cancer risk (Wentzensen *et al.* (2011) *Cancer Epidemiol. Biomarkers Prev.* 20:1238–1250; Calado *et al.* (2012) *Leukemia* 26:700–707; Artandi and DePinho (2010) *Carcinogenesis* 31:9–18); transient and intermittent, to allow the normal anti-cancer telomere shortening mechanism to function almost continuously; effective even in slow-dividing cells such as some progenitor and stem cell populations; deliverable *in vitro* and *in vivo* using non-immunogenic vehicles; and, finite, enabling only enough additional cell divisions to potentially ameliorate or prevent diseases of inadequate telomere maintenance, or to enable 10 sufficient amplification of cells for cell therapies or bioengineering applications.

15 Existing treatments, including treatments using small molecules, do not meet all of these criteria. Harley *et al.* (2011) *Rejuvenation Res.* 14:45–56. Viral delivery of TERT, while possibly inducible, risks insertional mutagenesis and thus presents serious safety concerns. Treatments involving continuous telomerase 20 overexpression are potentially unsafe, because in a cell with an oncogenic mutation, either due to critically short telomeres and resulting chromosomal instability (O'Sullivan and Karlseder (2010) *Nat. Rev. Mol. Cell Biol.* 11:171–181) or another cause, constitutive telomerase expression, either due to a second mutation or a drug, may support malignancy by enabling unlimited proliferation.

25 Artandi and DePinho (2010) *Carcinogenesis* 31:9–18; Ding *et al.* (2012) *Cell* 148:896–907.

[0143] The criteria for safe telomere extension are met by an RNA-based approach, facilitated by the recent discovery that certain naturally-occurring nucleosides found in RNA increase the stability and reduce the immunogenicity of exogenous RNA when delivered to cells. Karikó *et al.* (2005) *Immunity* 23:165–5 175. Examples of such nucleosides are listed in Table 4. Without intending to be bound by theory, these modified versions of canonical nucleosides may allow Toll-like receptors of the innate immune system to distinguish endogenous modRNA from foreign RNA not containing these nucleotides. Delivery of sufficiently purified synthetic modRNA containing these non-canonical nucleosides to cells 10 results in increased stability of the modRNA and transient elevation of the protein encoded by the modRNA with reduced or abrogated immune response. Karikó *et al.* (2011) *Nucleic Acids Res.* 39:e142; doi: 10.1093/nar/gkr695. Thus, delivery of exogenous modRNA presents unprecedented opportunities for applications which require transient protein production. For example, biomimetic modRNA has been 15 transfected into fibroblasts, thus reprogramming them to pluripotent stem cells. Yakubov *et al.* (2010) *Biochem. Biophys. Res. Commun.* 394:189–193. Biomimetic modRNA has also been injected into mice to rescue a model of pulmonary surfactant protein deficiency and to elevate erythropoietin and hematocrit levels. Kormann *et al.* (2011) *Nat. Biotechnol.* 29:154–157. As 20 described above, the nucleosides used in the modRNAs used herein were chosen not only to increase the stability but also to reduce or abrogate immunogenicity and to increase translational efficiency of the RNA. Transfection of dendritic cells using an unmodified mRNA encoding hTERT, while resulting in an increase in telomerase activity within the cells also elicited a strong hTERT-specific cytotoxic 25 T lymphocyte response. Saeboe-Larsen *et al.* (2002) *Journal of Immunological Methods* 259: 191–203. Approaches using unmodified RNAs are therefore not likely to be effective in extending telomeres in cells.

[0144] Thus, delivery of modRNA encoding TERT to cells may be used to transiently increase levels of TERT protein and telomerase activity sufficiently to 30 extend telomeres and increase replicative capacity by a finite amount. The increased levels of telomerase activity occur rapidly enough to enable a telomere extension treatment that is brief and infrequent (see Figure 1a and Figure 3).

[0145] To demonstrate the approach, TERT modRNA was synthesized using ratios of canonical and non-canonical nucleosides that confer stability, efficient translation, and reduced or abrogated immunogenicity. Yakubov *et al.* (2010) *Biochem. Biophys. Res. Commun.* 394:189–193; Karikó *et al.* (2011) *Nucleic Acids Res.* 39:e142; doi: 10.1093/nar/gkr695. The synthetic modRNA contains the 5' and 3' UTRs of beta globin mRNA, which has a relatively long half-life. To further enhance stability the modRNA has a long, 151 nucleotide poly-A tail. To maximize the fidelity of the long TERT DNA template used in the *in vitro* transcription reaction to generate RNA, the DNA templates were generated using a 10 plasmid- rather than PCR-based approach.

[0146] To distinguish bona fide telomere extension from selection of cells with long telomeres from a heterogeneous starting population, a control modRNA was synthesized that encodes a catalytically inactive form of TERT (CI TERT) with the single residue mutation D712A, in which one of the triad of metal-coordinating 15 aspartates at the catalytic site of the reverse transcriptase domain is substituted with alanine, abrogating the catalytic activity of CI TERT but leaving it structurally intact to the extent of being able to bind template DNA, and as stable as TERT in reticulocyte lysate. Wyatt (2009) "Structure-Function Analysis of the Human Telomerase Reverse Transcriptase" University of Calgary, Ph.D. Thesis 20 (http://dspace.ucalgary.ca/bitstream/1880/47511/1/2009_Wyatt_PhD.pdf).

[0147] MRC-5 human fetal lung fibroblasts were chosen as test cells in this example because these and other human fibroblast strains have served for decades as workhorses in the telomere field and there is thus abundant data to inform experimental design and analysis. MRC-5 cells also have relatively low 25 endogenous telomerase activity, and exhibit telomere shortening and eventual senescence. Since oxidative stress increases the rate of telomere shortening in MRC-5 cells (von Zglinicki *et al.* (2000) *Free Radic. Biol. Med.* 28:64–74), and causes TERT to localize to the cytoplasm where it cannot extend telomeres (Ahmed *et al.* (2008) *J. Cell. Sci.* 121:1046–1053), the cells were cultured in 5% 30 rather than ambient oxygen. To further increase the likelihood of success the cells were also cultured in medium optimized for protein production in MRC-5 cells (Wu *et al.* (2005) *Cytotechnology* 49:95–107).

[0148] The efficiency of modRNA-based telomere extension is subject to several factors, including efficiency of transfection, translation, and folding into functional telomerase, and ability to at least transiently escape the extensive mechanisms of post-translational regulation that inhibit TERT and telomerase in many cell types including MRC-5 cells. While the transfection efficiencies with smaller modRNA species such as nGFP (0.8 kb) in MRC-5 cells are typically over 90% even with low concentrations of modRNA (Figure 6), the TERT open reading frame is relatively large (3399 bp), and the modRNA TERT construct includes UTRs and a poly-A tail, thus making the construct even larger (3751 bp). As shown in Figure 1b, however, both TERT and CI TERT were efficiently transfected into MRC-5 cells by cationic lipid (as measured by RT-PCR of mRNA harvested at end of treatment from MRC-5 cells treated with 1 ug/ml TERT modRNA for 5 hours). Further, delivery of TERT or CI TERT modRNA resulted in equivalent and significant (P<0.03 and <0.01, respectively) increases in the amount of protein recognized by anti-TERT antibody, with a size of approximately 122 kDa, close to the estimated size of TERT of 127 kDa (Figure 1c). Wick *et al.* (1999) *Gene* 232:97–106. Protein levels were measured by quantitative infrared fluorescence Western blot. TERT protein levels do not differ significantly between cells treated with TERT and cells treated with CI TERT (n=3).

[0149] To form functional telomerase, TERT must fold properly and form a complex with at least TERC. Telomerase is also heavily regulated post-translationally, including by cytoplasmic localization and inhibitory phosphorylation. Cifuentes-Rojas and Shippen (2012) *Mutat. Res.* 730:20-27; doi:10.1016/j.mrfmmm.2011.10.003. Indeed not all cells expressing TERT are able to maintain telomeres. Counter *et al.* (1998) *Proc. Natl. Acad. Sci. U.S.A.* 95:14723–14728. Delivery of TERT modRNA increased telomerase activity in a dose-dependent manner up to a maximum at a concentration of approximately 0.6 ug/ml (see Figure 7), and a concentration of 1 ug/ml was used for subsequent experiments. Telomerase activity levels increased rapidly but returned to baseline levels within approximately 48 hours, and CI TERT modRNA caused no change in telomerase activity (Figure 1d). Telomerase activity was measured by qPCR-based TRAPeze[®] RT assay (n=3).

[0150] Since post-translational regulation of TERT is mediated in part by cytoplasmic sequestration in a cell cycle-dependent manner, the subcellular localization of TERT in treated and untreated cells was also investigated.

Cifuentes-Rojas *et al.* (2011) *Mutat. Res.* doi:10.1016/j.mrfmmm.2011.10.003.

5 Consistent with the Western blot results, immunocytochemistry showed abundant, though possibly inhibited, protein recognized by TERT antibody in untreated MRC-5 cells, making it difficult to distinguish endogenous from exogenous TERT (data not shown).

[0151] Cells within approximately 10 population doublings (PD) of the onset of 10 replicative senescence were used to measure the effect of TERT modRNA treatment on telomere length and replicative capacity. The decision to use such cells was based on at least two factors. First, cells at this stage have relatively short telomeres, and since telomerase preferentially extends short telomeres, including in MRC-5 cells, treatment of cells at this stage should result in a larger, 15 more easily measured effect. Britt-Compton *et al.* (2009) *FEBS Lett.* 583:3076–3080. Second, by treating cells at this stage, the amount of time required to determine whether the treatment increased replicative capacity is reduced. Under the instant conditions, replicative senescence begins approximately 50 PD after receipt of MRC-5 cells from the supplier (see Methods for details), and thus it was 20 decided to treat cells at approximately PD 40. MRC-5 cells at this stage have average telomere lengths of approximately 5-7 kb, depending on culture history and PD on receipt from suppliers. Sitte *et al.* (1998) *Free Radic. Biol. Med.* 24:885–893; MacKenzie *et al.* (2000) *Exp. Cell Res.* 259:336–350.

[0152] Several considerations were taken into account in designing the treatment 25 schedule (Figure 4a). Rates of telomere extension obtained using viral transduction of TERT into MRC-5 cells vary but are on the order of about 0.2 kb per division. MacKenzie *et al.* (2000) *Exp. Cell Res.* 259:336–350. Rates of telomere shortening in MRC-5 cells vary with culture conditions and practices (Sitte *et al.* (1998) *Free Radic. Biol. Med.* 24:885–893; von Zglinicki *et al.* (2000) 30 *Free Radic. Biol. Med.* 28:64–74), but are approximately 0.1 kb per PD in ambient oxygen (Sitte *et al.* (1998) *Free Radic. Biol. Med.* 24:885–893). Cells were cultured in 5% oxygen, as MRC-5 cell telomeres shorten more slowly under less

oxidative conditions. von Zglinicki *et al.* (2000) *Free Radic. Biol. Med.* 28:64–74.

The PD time in PD 40 MRC-5 cells was found to be approximately 33 hours.

Given these data, it might be expected that multiple treatments would be required

in order to detect telomere extension within the resolution of the chosen methods

5 of telomere length measurement, telomere restriction fragment (TRF) analysis and

quantitative fluorescence *in situ* hybridization (Q-FISH). The interval between

treatments was chosen to be 48 hours since it was found that telomerase activity

levels returned to baseline levels by this time after a single treatment.

[0153] As shown in Figures 3(b)-(d), treatment of MRC-5 cells with hTERT

10 modRNA results in a significant increase in telomere length in these cells

compared to cells that are either untreated or treated with the delivery vehicle only.

Fluorescence micrographs of metaphase MRC-5 cells following treatment shows

the location of a telomere probe on the chromosomes in these cells (Figure 4(e)).

[0154] The effect of the treatment on telomere length was also measured by

15 quantitative fluorescence *in situ* hybridization (Q-FISH) (Figure 4(f)), because this

technique provides relatively high resolution (0.3 kb) and because the fluorescence

of Q-FISH telomere probes is directly proportional to telomere length. Lansdorp *et*

al. (1996) *Hum. Mol. Genet.* 5:685–691; Martens *et al.* (1998) *Nat. Genet.* 18:76–

80. Because replicative capacity is the functional parameter of most interest in

20 increasing as a result of extending telomeres, a standard curve relating population

doubling number of MRC-5 cells to telomere length as measured using Q-FISH

was constructed (Figure 4(g)(i)). After three treatments of PD 40 MRC-5 with

TERT mRNA at 48 hour intervals, average total telomere length per cell increased

by 56+-5% (n=15 cells for each of two biological replicates for each treatment;

25 error bars represent s.e.m.). Telomere lengths in treated PD 40 cells were similar

to those of untreated PD 3 cells (Figure 4(g)(ii)) (upper dashed line), and thus the

treatment extended telomeres by the amount by which telomere shorten in these

cells over 37 PD. In this number of PD under standard culture conditions, MRC-5

telomeres shorten by over >1 kb. Sitte *et al.* (1998) *Free Radic. Biol. Med.*

30 24:885–893. Consistent with this, the observed increase in telomere length of 56%

corresponds to an increase in average telomere length of approximately 0.6–2.5 kb,

based on the fact that PD 40 MRC-5 cells have average telomere lengths of

approximately 5-7 kb as measured using TRF (Sitte *et al.* (1998) *Free Radic. Biol. Med.* 24:885–893; MacKenzie *et al.* (2000) *Exp. Cell Res.* 259:336–350), which overestimates telomere length by approximately 2.5-4 kb, and thus actual average telomere lengths in PD 40 MRC-5 cells are in the range of 1-4.5 kb. Aubert *et al.* (2012) *Mutat. Res.* 730:59–67. Since the treatment lasted a total of 144 hours, and the population doubling time of the cells was approximately 33 h. during this time, the cells underwent approximately 4-5 PD. Thus, the rate of telomere extension was approximately 0.1-0.6 kb per PD, the upper range of which approaches the maximum rates ever reported following viral transduction of DNA encoding TERT, and the lower range of which is comparable to typical rates seen using viral delivery. As expected, the untreated cells have telomere lengths equivalent to PD 40 cells during generation of the standard curve (lower dashed line in Figure 4(g)).

[0155] Next, the effect of TERT mRNA treatment on cell replicative capacity was examined. As expected, untreated and vehicle only-treated cells senesced within the normal range of 50-60 PD (Figure 5a). In striking contrast, cells treated with TERT mRNA continued to proliferate beyond the PD at which the control populations reached replicative senescence in a dose-dependent fashion, with each additional treatment conferring significantly more extra PD.

[0156] Three treatments with TERT mRNA resulted in an increase in replicative capacity of 28+-1.5 PD. This result is consistent with an estimate that telomeres were extended by > 1 kb, as MRC-5 telomeres shorten by approximately 0.1 kb per PD (Sitte *et al.* (1998) *Free Radic. Biol. Med.* 24:885–893) in ambient (20%) oxygen, and the treated cells were cultured in 5% oxygen, in which telomeres shorten more slowly. The observed increase in replicative capacity of treated cells of 28 PD is equivalent to the loss of replicative capacity that occurs in normal fibroblasts over more than a decade of normal human aging. Takubo *et al.* (2010) *Geriatr Gerontol Int.* 10 Suppl 1:S197-206; doi: 10.1111/j.1447-0594.2010.00605.x.; Allsopp, R. C. *et al.* (1992) *Proc. Natl. Acad. Sci. U.S.A.* 89:10114–10118.

[0157] Importantly for therapeutic applications, all of the treated cells studied to date eventually senesced, and indeed senesced in fewer PD than untreated cells with equivalent telomere lengths, as might be expected since the treated cells have

undergone dozens of additional PD, compared to the untreated cells with similar-length telomeres, in which to accumulate non-telomeric DNA damage or other damage that might affect rates of telomere shortening or otherwise support induction of replicative senescence.

5 [0158] As MRC-5 cells approached and entered replicative senescence, they tended to swell to several times the diameter of early passages (Figure 8a). As with replicative senescence, this transition was delayed in cells treated with TERT modRNA, but not in cells receiving CI TERT modRNA or vehicle only (Figure 8b).

10 [0159] Consistent with the finding that CI TERT modRNA caused no increase in telomerase activity (Figure 1d), CI TERT modRNA treatment did not change telomere length distribution relative to untreated cells, and nor did CI TERT modRNA increase replicative capacity, despite causing an increase in TERT protein level equivalent to that caused by TERT modRNA (Figure 1c). Since the 15 only difference between TERT and CI TERT is a single-residue substitution which abrogates the ability of CI TERT to transfer nucleotides to telomeres, these results strongly support the hypothesis that the increase in telomere length and replicative capacity observed in cells treated with TERT modRNA is due to bona fide telomere extension in at least some of the treated cells, rather than selection of a 20 pre-existing subpopulation of cells with longer telomeres.

[0160] The results of this example show that delivery of TERT modRNA to human cells transiently elevates telomerase activity, rapidly extends telomeres, and increases replicative capacity by a finite, and thus potentially safe, amount. Purified biomimetic modRNA is also hypo-immunogenic. Karikó *et al.* (2011) 25 *Nucleic Acids Res.* 39:e142; doi: 10.1093/nar/gkr695. Thus modRNA delivery meets several important criteria for a therapeutic telomere extension treatment, and also has promise with regard to other criteria including specificity, targeted delivery, and ability to overcome post-translational regulation. Regarding specificity, TERT overexpression may also affect other genes such as Wnt (Park *et* 30 *al.* (2009) *Nature* 460:66–72), but such effects may be avoided by delivering modRNA encoding a TERT mutated at binding sites of factors that mediate any such non-specific effects. Regarding delivery, the recent discovery that in humans

modRNA is transported in exosomes between cells via blood and other body fluids enables exosome-based modRNA delivery for telomere extension and other applications. Lakhal and Wood (2011) *Bioessays* 33:737–741. Exosomes have been used successfully to deliver biomimetic modRNA to cells *in vitro* (data not shown). Regarding post-translational regulation, cell cycle-dependent and independent inhibitory post-translational regulation of TERT may be avoided by delivering modRNA encoding TERT mutated at one or more known sites that mediate those activities, such as the nuclear export sequence, or by co-delivering modRNA encoding other members of the telomerase or telomere complexes.

These approaches enable telomere extension even in slowly- or non-dividing cells, making TERT modRNA treatment appropriate for quiescent or slow-dividing stem and progenitor cell populations. modRNA-based telomere extension thus finds immediate use in increasing the replicative capacity of cells for research, and additionally for cell therapies, bioengineering applications, and *in vivo* treatments that address diseases and conditions of inadequate telomere maintenance.

[0161] In summary, modRNA encoding human telomerase reverse transcriptase (TERT) was delivered to MRC-5 human lung fibroblasts three times over 96 hours. Telomeres in treated cells were extended by >1 kb, an amount by which fibroblast telomeres shorten over more than a decade of aging in humans on average.

Takubo *et al.* (2010) *Geriatr Gerontol Int.* 10 Suppl 1:S197-206; doi: 10.1111/j.1447-0594.2010.00605.x. Telomerase activity returned to pre-treatment levels within 48 hours and the onset of replicative senescence in the treated cells was delayed by approximately 30 population doublings, in a dose-dependent manner. Thus delivery of telomerase RNA containing modified nucleosides to cells allows for the rapid and hypoimmunogenic or nonimmunogenic and RNA is a useful approach to telomere extension for diverse applications.

METHODS

[0162] modRNA template generation and synthesis. The wild type (WT) human TERT open reading frame (ORF) used to generate the DNA templates for modRNA synthesis is identical to the NCBI human TERT transcript variant 1 reference sequence NM_198253.2, and was generated by making the modification

G516D to the ORF of the pBABE-neo-hTERT plasmid (Addgene plasmid 1774). Residue 516 is in the QFP motif of the N-terminal extension of TERT, a motif associated with multimerization and RNA binding of TERT. The catalytically inactive TERT (CI TERT) mutant was generated from the WT TERT sequence by 5 introducing the mutation D712A. The WT and CI, TERT ORFs were inserted into the MCS of a starting plasmid containing the T7 promoter, the 5' UTR of human beta globin (hBB), the MCS, the 3' UTR of hBB, a 151 bp poly-A sequence, and a restriction site for linearization with class IIs enzyme following the poly-A sequence. The resulting intermediate plasmid was sequenced in at least 10 quadruplicate to ensure fidelity, linearized, and transcribed to capped RNA using the RNA polymerase from the MEGAscript T7 Kit (Ambion, Austin, Texas) and a custom nucleotide mix of canonical and non-canonical nucleotides (TriLink BioTechnologies) in which the final nucleotide concentrations per 40 ul IVT reaction were 7.5 mM for each of adenosine-5'-triphosphate (ATP), 5- 15 methylcytidine-5'-triphosphate (m5C), and pseudouridine-5'-triphosphate (Ψ), 1.5 mM for guanosine-5'-triphosphate (GTP), and 6 mM for the cap analog (ARCA, NEB), or a molar ratio of ATP:m5C:Ψ:GTP:ARCA of 1:1:1:0.2:0.8. To further decrease potential immunogenicity of the mRNA related to the 5'-3P-bearing fraction (~20% of total) the IVT products were treated with phosphatase (Antarctic 20 Phosphatase, NEB). The size and integrity of the modRNA products was verified using denaturing agarose gel electrophoresis.

[0163] Cells and cell culture. MRC-5 human fetal lung fibroblasts (ATCC CCL-171) were received from ATCC at passage 14, the current passage number of their distribution inventory. Since ATCC does not indicate the PD number, the PD 25 values cited herein refer to the number of PD after receipt of cells from ATCC, defined here as PD 0. To shorten their telomeres in preparation for telomere extension experiments, cells were cultured using ATCC guidelines in DMEM with 10% FBS, in ambient oxygen and 5% CO₂. Telomere extension treatment began at approximately PD 40 after receipt of cells from ATCC. At least 48 hours before 30 the start of modRNA treatment cells were transferred to 5% oxygen and DMEM medium containing 20% FBS and penicillin-streptomycin. Cells were treated at least 24 hours after plating and 24 hours before trypsinization.

[0164] modRNA transfection. Cells were transfected with 0.4 nM TERT modRNA and 2.0 nM TERC RNA unless otherwise specified, using Lipofectamine RNAiMax (Invitrogen), a cationic lipid, in Opti-MEM Reduced Serum Media (Invitrogen) for 4-5 hours, after which they were returned to normal medium.

5 [0165] Telomerase activity measurement. 24 hours after the start of transfection period, cells were washed with PBS, trypsinized, pelleted at 500 g for 5 min. and washed with PBS, repelleted, and then lysed following the instructions in the TRAPeze RT kit (Millipore). The reverse transcription and qPCR steps were carried out in a Roche LightCycler 480 II and an ABI 7900 HT. Telomerase 10 activity of samples was always compared to the reference standard provided with the kit, as we found that fold-increase was prone to be highly variable, probably due to the low telomerase activity in MRC-5 cells.

15 [0166] Immunocytochemistry. 24 hours after the start of transfection period, cells were washed with PBS, fixed in 2% paraformaldehyde for 20 minutes, washed three times with PBS, blocked in PBS containing 7.5% BSA and 0.1% Triton X-100 for 1 hour, washed three times, incubated overnight at 4oC on a rocker in anti-TERT antibody (ABCAM 32020 at 1:50 or Rockland 600-401-252S at 1:500) in PBS with 5% BSA, washed three times, incubated for one hour in secondary antibody, washed two times then two more times on a shaker for 3 20 minutes each time, then incubated in 0.1 ug/ml DAPI for 3 minutes, and washed four times in PBS.

25 [0167] Flow cytometry. Cells treated with 0.5-1.5 ug/ml of TERT modRNA and controls were trypsinized 22 h. after start of transfection, washed, fixed in 2% paraformaldehyde, permeabilized in 7.5% BSA with 0.1% Triton X-100 for 20 min., washed three times in PBS, blocked in 7.4% BSA for 1 h, incubated in anti-TERT antibody (ABCAM 32020 at 1:50) in PBS with 5% BSA, washed three times, incubated in secondary antibody (1:200) in PBS, then washed three times and resuspended in FACS buffer and analyzed on a Accuri C6 Flow Cytometer (Becton Dickinson). Mean fluorescence intensity was calculated as the average 30 fluorescence of all cells in treated and control samples. Data were analyzed using CFlow Plus software.

[0168] RT-PCR Total RNA was harvested using the RNeasy Mini kit (Qiagen) and converted to cDNA using the High-Capacity RNA-to-cDNA Kit (Invitrogen). cDNA was amplified using PCR with the following primers:

hTERT-F: 5'-GCCCTCAGACTTCAAGACCA (SEQ ID NO:1)

5 3'-hBB-R: 5'-AGGCAGAACCTCAGATGCTCA (SEQ ID NO:2)

GAPDH-F: 5'-GTGGACCTGACCTGCCGTCT (SEQ ID NO:3)

GAPDH-R: 5'-GGAGGAGTGGGTGTCGCTGT (SEQ ID NO:4)

[0169] Western blot. Protein was harvested by washing cells once with PBS and then lysing cells in RIPA buffer. Protein was run on NuPAGE Novex Tris-Acetate 10 Gels, transferred to PVDF membrane for 2 h. at 35 V, then hybridized to anti-alpha tubulin (Sigma at 1:10,000) and anti-TERT antibody (ABCAM 32020 at 1:1000 or Rockland 600-401-252S at 1:500) and anti-overnight at 4°C. Secondary detection was performed using infrared (680 nm and 800 nm) antibodies (LI-COR) and the Odyssey imager (LI-COR). Images were analyzed using ImageJ.

15 [0170] Q-FISH. Cells were incubated in 0.1 ug/ml colcemid for 4 hours before fixation and preparation of metaphase spreads following the procedure of Lansdorp *et al.*⁴¹ Slides were then stained using the Telomere PNA FISH Kit/FITC kit (Dako, Denmark), substituting AlexaFluor 555-labeled telomere probe (Bio-Synthesis, USA) for the kit probe. Each cell was imaged on a DeltaVision (Applied Precision, Inc., Washington) microscope using SoftWoRx software using either a 60X or 100X oil objective at 0.2 micron intervals over a range of 3 microns using a motorized stage, and custom software was used to identify the image in which each telomere spot was in focus and integrate its intensity in that in-focus image. Temporal variation in illumination intensity was compensated for 20 using the DeltaVision photosensor which recorded the illumination intensity of each image, and variation in spatial intensity and CCD dark current and read error was compensated for by acquiring flat field images and dark field images, respectively. At least 15 metaphase cells contributed to each replicate and each sample was measured in at least duplicate over at least two separate experiments.

25 [0171] Growth curves. Cells were harvested and counted on a hemocytometer as an automated counter was unable to accurately count the cells when the population became heterogeneous with respect to diameter as the population entered

replicative senescence. Images were acquired to allow measurement of cell diameters. PD were calculated as the log₂ of the ratio of cell numbers at end and start of each culture period between passages.

[0172] Statistics. Statistical analysis was performed using Microsoft Excel.
5 Error bars represent the mean \pm s.e.m. Telomere lengths were compared using T-tests with P < 0.05 (two-tailed) considered to be statistically significant.

Example 2. Effects of TERT modRNA treatment on human endothelial cells

[0173] The effects of TERT modRNA treatment is not limited to fibroblast cells.
10 As shown in Figure 9, human microvascular dermal endothelial cells ("HMDEC") display typical doubling curves, with early onset of senescence at approximately passage #12 and with cells completely senescent at passage #17. As shown in Figure 10, treatment of these cells at passage #18 with wild-type hTERT modRNA (right bar in each series) results in the reversal of senescence, whereas control treatment (left bar in each series), or treatment with mutant hTERT modRNA (middle bar in each series), had no effect on cellular senescence.
15

[0174] Figure 11 demonstrates the effects of various treatments on the growth of HMDECs. Specifically, cells treated three times with wild-type TERT modRNA (WT) between passage #10 and passage #11 continued to grow, whereas untreated 20 cells (UT), cells treated with carrier only (CO), and cells treated with a catalytically-inactive TERT (CI) stopped doubling.

[0175] Figure 12 shows the results of β -galactosidase (bGAL) staining of HMDECs at passage #14 following treatment with WT and CI modRNAs. This staining measures β -galactosidase activity at pH 6, which is a property of senescent 25 cells that is not found in presenescence, quiescent, or immortal cells. See Dimri *et al.* (1995) *Proc. Nat'l Acad. Sci. USA* 92:9363-7.

METHODS

[0176] The HMDECs Cumulative Population Doubling Curve was built using a 30 12-well plate format. Approximately 10^5 of HMDECs were at each time point replated in triplicates and cell counting was done on day 4 after the previous

replating. As shown in Figure 9, passage #12 was determined as the early onset point of HMDECs senescence.

[0177] As shown in Figure 10, approximately 10^5 HMDECs were transfected twice every other day with 0.75 mg of modRNAs encoding hTERT-CI, and -WT in 5 present of Carrier Only (CO, RNiMAX) control. Cell counting was done on Passages #19, 20, and 21 each time on day 5 after replating.

[0178] For the experiment shown in Figure 11, in the interval between passage #10 and #11 (and, as noted above, passage #12 corresponds to the early onset of HMDEC senescence), approximately 5×10^5 HMDECs per 75 cm³ flask were 10 cultivated on EBM-2 medium supported with EGM-2 MV supplement (both from Lonza, Walkersville, MD USA, catalogue nos. CC-3156 and CC-4176, respectively). Microvascular cells were transfected three times every other day with 5 μ g of modRNAs encoding hTERT-WT (WT) and hTERT-CI (CI) in the presence of untreated (UT) and carrier only (CO, RNAiMAX) control flasks. At 15 each time point, 5×10^5 cells were plated and, starting from passage #11, a cell count via hemocytometer was done on 5 days after replating. For each time point, each experimental condition was performed in triplicate.

[0179] For the experiments shown in Figure 12, approximately 10^5 of each NT, hTERT-CI and hTERT-WT HMDECs of passage #14 were treated as described in 20 Figure 10. The cells were then subjected to senescent cell analysis based on a histochemical stain for β -galactosidase activity at pH 6 using a Senescence Cells Histochemical Staining Kit (Sigma, catalogue no. CS0030). The left side of the slide shows representative images of non-treated (NT) cells and cells treated with hTERT-CI (CI) or hTERT-WT (WT) modRNAs. All images were obtained under 25 identical image-acquisition conditions. Each experimental condition was performed in triplicate, and the percent of cells positive for β -galactosidase activity (as represented in the chart) for each condition was calculated as an average of positive cells in three randomly captured areas.

Example 3. Further characterization of the effects of TERT modRNA treatment on human cells

[0180] In this example, the mRNA was transfected via a cationic lipid into primary human fibroblasts and myoblasts (Figure 13A), cells known to have limited proliferative capacity. Webster and Blau (1990) *Somat Cell Mol Genet.* 16:557–565; Hayflick and Moorhead (1961) *Exp Cell Res.* 25:585–621; Yakubov *et al.* (2010) *Biochem Biophys Res Commun.* 394:189–193. Transfection efficiency was determined using flow cytometric single cell quantitation of fluorescence following delivery of GFP mRNA, which showed that most cells (10) (>90%) were transfected even at relatively low concentrations of modified mRNA (0.1 µg/ml) (Figure 13B and Figure 16A-C). Treatment of cells with equal concentrations of exogenous *TERT* mRNA or mRNA encoding a catalytically inactive (CI) form of *TERT* resulted in internalization of similar amounts of mRNA (Figure 16D), as measured by RT-qPCR 24 h after the first treatment. CI *TERT* (15) has a substitution mutation at one of the triad of metal-coordinating aspartates at the catalytic site of the reverse transcriptase domain of *TERT*. As a result, CI *TERT* cannot add nucleotides to telomeres, yet remains structurally intact, able to bind template DNA, and exhibits stability comparable to wild type *TERT* in reticulocyte lysates. Wyatt (2009) *Structure-Function Analysis of the Human* (20) *Telomerase Reverse Transcriptase*. Neither *TERT* nor CI *TERT* mRNA treatment affected levels of endogenous *TERT* mRNA relative to untreated cells as measured by RT-qPCR (Figure 16E). Transfection with 1 µg/ml of either *TERT* or CI *TERT* mRNA resulted in equivalent 50% increases (P<0.05 and <0.01, respectively) in the amount of *TERT* protein in fibroblasts (Figure 13C, left panel). Wick *et al.* (25) (1999) *Gene.* 232:97–106; Ahmed *et al.* (2008) *J Cell Sci.* 121:1046–1053. The presence of endogenous *TERT* protein in cells with little endogenous telomerase activity is consistent with the relative abundance of inactive splice variants of *TERT* in many cell types and extensive post-translational inhibitory regulation of *TERT* activity as previously reported by others. Yi *et al.* (2001) *Nucleic Acids Res.* (30) 29:4818–4825; Cifuentes-Rojas and Shippen (2011) *Mutat Res.* [published online ahead of print: October 18, 2011]; doi:10.1016/j.mrfmmm.2011.10.003. Treatment with increasing amounts of *TERT* mRNA resulted in a dose-dependent increase of

TERT protein expression as measured in single cell assays by flow cytometry (Figure 13C, right panel).

[0181] *Telomerase activity is transiently increased.* To test whether modified *TERT* mRNA delivery resulted in the generation of functional *TERT* protein, 5 telomerase activity was quantified using a gel-based TRAP assay. Telomerase activity was detected in fibroblasts and myoblasts at all doses of *TERT* mRNA tested (0.25, 0.5, 1.0, and 2.0 μ g/ml), and was not detected in untreated cells or cells treated with either vehicle only or modified mRNA encoding CI *TERT*, even at the highest dose of 2.0 μ g/ml (Figure 13D). Although *TERT* requires *TERC* to 10 form a functional telomerase complex, delivery of *TERT* alone was sufficient to increase telomerase activity, consistent with previous findings that *TERT* is often limiting as *TERC* RNA copy number is high in many cell types lacking telomerase activity, including fibroblasts. Yi *et al.* (2001) *Nucleic Acids Res.* 29:4818–4825. A time course revealed that telomerase activity peaked at 24 hours and returned to 15 baseline levels within 48 hours after a single transfection. This time frame is consistent with previously reported half-lives of human *TERT* mRNA (2-4 h), human β -globin mRNA (17-18 h: our exogenous *TERT* mRNA is flanked by β -globin 5' and 3' UTRs), and telomerase activity in cells exposed to an inhibitor of protein synthesis (cell type dependent, but typically >24 h). Kabnick and 20 Housman (1988) *Mol Cell Biol.* 8:3244–3250; Holt *et al.* (1997) *Proc Natl Acad Sci U S A.* 94:10687–10692; Xu *et al.* (1999) *Br J Cancer.* 80:1156–1161.

[0182] *Lengthening of telomeres.* Telomere lengths in untreated fibroblasts declined over time (3 months) as expected (62) (Figure 14A) and was quantified using two different methods. The monochrome multiplex qPCR method 25 (MMqPCR) was used to assess length, and measurements were validated independently with a qPCR method performed by SpectraCell Laboratories, Inc. (correlation coefficient 0.97, P<0.001). Delivery of *TERT* mRNA three times in succession at 48-hour intervals to fibroblasts or myoblasts starting at population doubling (PD) 25 and 6, respectively, extended telomeres by 0.9 ± 0.1 kb ($22 \pm 3\%$), and 0.7 ± 0.1 kb ($12 \pm 2\%$), respectively (Figure 14B,C). Treatment with 30 vehicle only or CI *TERT* mRNA had no significant effect on telomere length

relative to untreated cells. The average rate of telomere extension in fibroblasts was 135 ± 15 bp/PD.

[0183] *Cell type-dependent increases in proliferative capacity.* To test the effect of modified *TERT* mRNA delivery and consequent telomere extension on cell 5 proliferative capacity, human fibroblasts were transfected either once, twice, or three times in succession. Treatments were delivered at 48-hour intervals. Untreated, vehicle only-treated, and CI *TERT* mRNA-treated fibroblasts exhibited an equivalent plateau in cell number after approximately 50-60 PD, whereas cells treated three times with *TERT* mRNA continued to proliferate for a finite 10 additional 28 ± 1.5 PD with an overall increase in cell number of 2.7×10^8 beyond untreated cells (Figure 14D, left panel). The effect was dose-dependent with each additional treatment conferring additional PD (Figure 14D, right panel). The incremental increase in proliferative capacity was greater with the first treatment than with the second or third treatments. Human myoblasts treated three times in 15 succession every 48 hours gained 3.4 ± 0.4 PD, equivalent to a 10-fold increase in cell number compared to untreated or vehicle treated controls (Figure 14E). Such differences in PD between myoblasts and fibroblasts are not unexpected, as prior studies found similar limited effects of *TERT* overexpression to a few PD and showed that this limitation was due to a p16-mediated growth arrest in human 20 myoblasts, in contrast to fibroblasts. Bodnar *et al.* (1998) *Science*. 279:349–352; Zhu *et al.* (2007) *Aging Cell*. 6:515–523. In both fibroblasts and myoblasts, vehicle only or CI *TERT* mRNA had no effect on proliferative capacity compared to untreated controls. These data show that delivery of modified *TERT* mRNA is an effective method for increasing PD in culture. Importantly, all of the treated 25 cells studied exhibited a significant increase in cell numbers, but eventually reached a plateau in their growth curves, demonstrating absence of immortalization.

[0184] *Transient reduction in markers of senescence.* As the fibroblast 30 populations stopped growing they exhibited markers of senescence including senescence-associated β -galactosidase (β -gal) staining and enlarged size (Figure 15A-C). Cristofalo and Kritchevsky (1969) *Med Exp Int J Exp Med*. 19:313–320; Dimri *et al.* (1995) *Proc Natl Acad Sci U S A*. 92:9363–9367; Cristofalo *et al.*

(2004) *Mech Ageing Dev.* 125:827–848; Lawless *et al.* (2010) *Exp Gerontol.* 45:772–778. These changes were transiently reduced in fibroblasts treated with *TERT* mRNA relative to untreated cells and cells receiving CI *TERT* mRNA or vehicle only. In accordance with findings by others, not all cells in the populations that had entered a growth plateau expressed β -galactosidase at detectable levels. Lawless *et al.* (2010) *Exp Gerontol.* 45:772–778; Binet *et al.* (2009) *Cancer Res.* 69:9183–9191. However, *TERT* mRNA-transfected fibroblasts and myoblasts expressed β -galactosidase to the same degree as the control cells of each type after the two populations reached a growth plateau. These data demonstrate that cells treated with *TERT* mRNA eventually and predictably cease division and express markers of senescence, and are therefore unlikely to be transformed.

[0185] This example demonstrates that transient delivery of *TERT* mRNA comprising modified nucleotides extends human telomeres and increases cell proliferative capacity without immortalizing the cells. The rate of telomere extension in fibroblasts observed here of 135 ± 15 bp/PD is comparable to rates reported using viral methods, from 94 to >150 bp/PD (22, 69). Modified *TERT* mRNA extended telomeres in fibroblasts in a few days by 0.9 ± 0.1 kb. Fibroblast telomere lengths have been reported to shorten over a human lifetime by approximately 1-2 kb on average. Allsopp *et al.* (1992) *Proc Natl Acad Sci U S A.* 89:10114–10118. Thus, modified *TERT* mRNA is efficacious, yet transient and non-integrating, overcoming major limitations of constitutively expressed viral *TERT* mRNA delivery.

[0186] Human cells of greatest interest are often limited in number, including stem cells for use in experimentation or regenerative medicine. This problem is currently being addressed by various methods including somatic nuclear transfer, viral methods for gene delivery, and the use of culture conditions that lessen the rate of telomere shortening. Le *et al.* (2013) *Cell Stem Cell.* [published online ahead of print: November 19, 2013]; doi:10.1016/j.stem.2013.11.005; Zimmermann and Martens (2008) *Cell Tissue Res.* 331:79–90; Mohsin *et al.* (2013) *Circ Res.* 113:1169–1179. The modified *TERT* mRNA treatment described here provides an advantageous complement or alternative to these methods that is brief, extends telomeres rapidly, and does not risk insertional mutagenesis. The

brevity of *TERT* mRNA treatment is particularly attractive in that it can avert the loss of stem cell phenotype that can occur over time in culture (Gilbert *et al.* (2010) *Science*. 329:1078–1081) and shorten the post-reprogramming stage of iPSC generation during which telomeres extend (Wang *et al.* (2012) *Cell Res.* 22:757–768). Such a method of extending telomeres has the potential to increase the utility of diverse cell types for modeling diseases, screening for ameliorative drugs, and use in cell therapies.

5 [0187] A spectrum of effects on proliferative capacity was observed for the cell types tested, in agreement with previous studies demonstrating different effects of 10 *TERT* overexpression on myoblast and fibroblast proliferative capacity. Bodnar *et al.* (1998) *Science*. 279:349–352; Zhu *et al.* (2007) *Aging Cell*. 6:515–523.

15 Moreover, the amount of telomere extension did not correlate with proliferative capacity. Thus, cell context determines the efficacy of *TERT* expression on proliferative capacity and an understanding of the factors mediating this effect is of 20 interest in overcoming this limitation. Factors that have been implicated in limiting myoblast proliferative capacity upon viral *TERT* overexpression include p16-mediated growth arrest, cell type and strain, and culture conditions. Zhu *et al.* (2007) *Aging Cell*. 6:515–523. More generally, the effect may be mediated by 15 non-telomeric DNA damage, age, and mitochondrial integrity. Sahin *et al.* (2011) *Nature*. 470:359–365; Mourkioti *et al.* (2013) *Nat Cell Biol*. 15:895–904; López- 25 Otín *et al.* (2013) *Cell*. 153:1194–1217. The absence of an increase in telomere length or cell proliferative capacity in CI *TERT* mRNA-transfected cells is consistent with the treatment acting through the catalytic site of *TERT* by which nucleotides are added directly to telomeres. *TERT* mRNA-treated cell populations increased in number exponentially for a period of time and then eventually ceased 25 expanding and exhibited markers of senescence to a similar degree as untreated populations, consistent with the absence of immortalization.

30 [0188] The transient non-integrating nature of modified mRNA and finite increase in proliferative capacity observed here render it safer than currently used viral or DNA vectors. Further, the method extends telomeres rapidly so that the treatment can be brief, after which the protective telomere shortening mechanism remains intact. This method can be used *ex vivo* to treat cell types that mediate

certain conditions and diseases, such as hematopoietic stem cells or progenitors in cases of immunosenescence or bone marrow failure. In addition, modified mRNA may be delivered to certain tissues *in vivo*. Kormann *et al.* (2011) *Nat Biotechnol.* 29:154–157. In summary, the rapid and safe method for rapid extension of 5 telomeres described here leads to delayed senescence and increased cell proliferative capacity without immortalizing human cells.

METHODS

[0189] *mRNA template generation and synthesis.* To generate modified mRNA 10 encoding GFP, *TERT*, and CI *TERT*, their respective open reading frames (ORFs) were inserted into the MCS of a starting plasmid containing the T7 promoter, the 5' UTR of human b-globin (*HBB*), the MCS, the 3' UTR of *HBB*, a 151 bp poly-A sequence, and a restriction site for linearization with a class II enzyme following the poly-A sequence. The resulting intermediate plasmids were sequenced, 15 linearized, and transcribed using the buffer and RNA polymerase from the MEGAscript T7 Kit (Ambion, Austin, Texas, USA), and a custom mix of canonical and non-canonical nucleotides (TriLink BioTechnologies, San Diego, CA, USA) in which the final nucleotide concentrations per 40 μ l IVT reaction were 7.5mM for each of adenosine-5'-triphosphate (ATP), 5-methylcytidine-5'-triphosphate (m5C), and pseudouridine-5'-triphosphate (Ψ), 1.5 mM for guanosine-5'-triphosphate (GTP), and 6mM for the cap analog (ARCA) (New England 20 Biolabs, Ipswich, MA, USA), or a molar ratio of ATP:m5C: Ψ :GTP:ARCA of 1:1:1:0.2:0.8. To further decrease potential immunogenicity of the mRNA related to the 5'-3P-bearing fraction, the IVT products were treated with Antarctic 25 Phosphatase (New England Biolabs). The size and integrity of the mRNA products were verified using denaturing agarose gel electrophoresis. The wild type human *TERT* ORF used to generate the DNA templates for mRNA synthesis is identical to the NCBI human *TERT* transcript variant 1 (reference sequence NM_198253.2). The ORF was generated from the pBABE-neo-hTERT plasmid 30 (Counter *et al.* (1998) *Proc Natl Acad Sci U S A.* 95:14723–14728) (plasmid 1774, Addgene, Cambridge, MA, USA). The pBABE-neo-hTERT plasmid had a non-silent mutation at residue 516 in the QFP motif of *TERT*, a motif associated with

multimerization and *TERT* interaction with *TERC* RNA, and thus to avoid the possibility of artifacts due to this mutation we made the sequence identical to the NCBI reference sequence by correcting the mutation with the change G516D. The CI *TERT* mutant was generated from the *TERT* sequence by introducing the 5 mutation D712A.

[0190] *Cell culture and treatment.* Human primary fetal lung MRC5 fibroblasts were obtained from ATCC (Manassas, VA, USA) at passage 14. ATCC does not indicate the PD number, thus, our PD values cited herein refer to the number of PD after receipt of cells from ATCC. MRC5 cells were cultured in DMEM with 20% 10 FBS and penicillin-streptomycin. Human 30 year-old primary skeletal muscle myoblasts (Lonza, Allendale, NJ, USA) were cultured in SkGM-2 media (Lonza) according to the vendor's instructions. Population doublings were calculated as the base 2 log of the ratio between cells harvested and cells plated at the previous passaging, and were considered to be zero if fewer cells were harvested than 15 plated. Cells were transfected with modified *TERT* mRNA using Lipofectamine RNAiMax (Life Technologies, Grand Island, NY, USA) prepared in OptiMEM Reduced Serum Media (Life Technologies, Grand Island, NY, USA) and added to the cells in a 1:5 v:v ratio with their normal media to achieve the final concentrations indicated herein.

20 [0191] *Telomerase activity measurement.* Twenty-four hours after the start of the transfection period, cells were harvested and lysed in CHAPS buffer. The TRAP assay was performed using a modified version of the TRAPeze kit (EMD Millipore, Billerica, MA, USA), in which the primers and polymerase were added after, rather than before, the step during which the artificial telomere substrate is 25 extended. The PCR program was 94°C 30s/59°C 30s/72°C 45s for 30 cycles, and the products were run on a 15% polyacrylamide gel in 0.5X TBE stained with SYBR Gold Nucleic Acid Gel Stain (Life Technologies, Grand Island, NY, USA). The time course of telomerase activity was performed using the TRAPeze RT kit (EMD Millipore, Billerica, MA, USA).

30 [0192] *Western blot.* Protein was harvested by washing cells once with PBS and then lysing cells in RIPA buffer (Cell Signaling Technology, Danvers MA, USA). Protein was run on NuPAGE Novex Tris-Acetate Gels (Life Technologies, Grand

Island, NY, USA), transferred to PVDF membrane for 2 h at 35V, then hybridized to anti- α tubulin (Sigma, St. Louis, MO, USA) at 1:10,000 and anti-TERT antibody (ABCAM, Cambridge, MA, USA, 32020 at 1:1000; or Rockland Immunochemicals, Gilbertsville, PA, USA, 600-401-252S at 1:500) and incubated 5 overnight at 4°C. Detection was performed using infrared (680 nm and 800 nm) antibodies (LI-COR, Lincoln, NE, USA) and the Odyssey imager (LI-COR). Total intensity of each band was quantified using ImageJ (NIH, Bethesda, MD, USA). The intensity of each TERT band was normalized by its corresponding α tubulin band.

10 [0193] *Flow cytometry.* Cells were harvested 24 h after transfection with the indicated doses (Figure 16A-C) of TERT mRNA and stained with anti-TERT antibody (Rockland Immunochemicals, Gilbertsville, PA, USA; 600-401-252S) at 1:500.

[0194] *Telomere length measurement by SpectraCell Laboratories, Inc.* 15 Genomic DNA was extracted using phenol chloroform and quantified using the Quant-iT™ PicoGreen® dsDNA Assay Kit (Life Technologies, Grand Island, NY, USA). Telomere length analysis was performed at SpectraCell Laboratories Inc. (Houston, TX, USA) using a CLIA approved, high throughput qPCR assay, essentially as described by Cawthon *et al.* Cawthon (2002) *Nucleic Acids Res.* 30(10):e47; Cawthon (2009) *Nucleic Acids Res.* 37(3):e21. The assay determines a relative telomere length by measuring the factor by which the sample differs from a reference DNA sample in its ratio of telomere repeat copy number to single gene (36B4) copy number. This ratio (T/S ratio) is thought to be proportional to the average telomere length. All samples were run in at least duplicate with at least one 20 negative control and two positive controls of two different known telomere lengths (high and low) and an average variance of up to 8% was seen. The results were reported as a telomere score equivalent to the average telomere length in kb.

25 [0195] *Telomere length measurement by MMqPCR.* Telomere length was measured using a modified version of the MMqPCR protocol developed by Cawthon (Cawthon (2009) *Nucleic Acids Res.* 37(3):e21) with the following 30 changes: Additional PCR pre-amplification cycles were added to make the telomere product amplify earlier, widening the gap between telomere and single-

copy gene signals; a mixture of two Taq polymerases was experimentally determined to result in better PCR reaction efficiencies than each on its own; reducing the SYBR Green concentration from 0.75X to 0.5X resulted in earlier signal. Genomic DNA was harvested from cells using the PureGene kit (Qiagen 5 Germantown, MD, USA) with RNase digestion, quantified using a NanoDrop 2000 (ThermoFisher Scientific, Waltham, MA, USA), and 10-40 ng was used per 15 μ l qPCR reaction performed in quadruplicate using a LightCycler 480 PCR System (Roche, Basel, Switzerland). A serial dilution of reference DNA spanning five points from 100 ng/ μ l to 1.23 ng/ μ l was included in each assay to generate a 10 standard curve required for sample DNA quantification. The final concentrations of reagents in each 15 μ l PCR reaction were: 20 mM Tris-HCl pH 8.4, 50 mM KCl, 3 mM MgCl₂, 0.2 mM each dNTP, 1 mM DTT, 1 M betaine (Affymetrix, Santa Clara, CA, USA), 0.5X SYBR Green I (Life Technologies, Grand Island, NY, USA), 0.1875U Platinum Taq (Life Technologies, Grand Island, NY, USA), 15 0.0625X Titanium Taq (Clontech), and 900 nM each primer (telg, telc, hbgu, and hbgd primer sequences specified in Cawthon (2009) *Nucleic Acids Res.* 37(3):e21. The thermal cycling program was: 2 minutes at 95°C; followed by 6 cycles of 15s at 95°C, 15s at 49°C; followed by 40 cycles of 15s at 95°C, 10s at 62°C, 15s at 72°C with signal acquisition, 15s at 84°C, and 10s at 88°C with signal acquisition. 20 The Roche LightCycler 480 software was used to generate standard curves and calculate the DNA concentrations of telomere and single-copy genes for each sample. T/S ratios were calculated for each sample replicate, and the result averaged to yield the sample T/S ratio which was calibrated using blinded replicate samples of reference cells sent to SpectraCell as described above. The 25 independently obtained relative values of T/S ratios measured using MMqPCR and by SpectraCell for the same samples were highly consistent (correlation coefficient = 0.97, P<0.001).
[0196] *Reverse transcription qPCR.* Primers were designed using Primer3 (Untergasser *et al.* (2012) *Nucleic Acids Res.* 40:e115) and are listed in Table 9 except where otherwise noted. Twenty-four hours after start of treatment, cells 30 were washed three times with PBS before harvesting in Buffer RLT (Qiagen, Germantown, MD, USA). RNA was converted to cDNA using High Capacity

RNA-to-cDNA Master Mix (Life Technologies, Grand Island, NY, USA). Endogenous *TERT* mRNA was amplified using a forward primer in the open reading frame of *TERT* and a reverse primer in the 3' UTR of endogenous *TERT* mRNA. Exogenous *TERT* mRNA was amplified using a forward primer in the 5 open reading frame of *TERT* mRNA and a reverse primer in the 3' UTR of *HBB* present in our exogenous *TERT* and CI *TERT* mRNA, but not in endogenous *TERT* mRNA. Relative levels were calculated using the Pfaffl method. Reference genes were *RPL37A* (using primers specified in Greber *et al.* (2011) *EMBO J.* 30:4874–4884) and *GAPDH*, neither of which exhibited a significant change in Ct value in 10 control or treated cells.

Table 9: Primer sequences.

Target	Forward primer (5'-3')	Reverse primer (5'-3')	Product length (bp)
Exogenous	hTERT (NM_198253.2)	3' UTR of HBB	162
TERT	GTCACCTACGTGCCACTCCT (SEQ ID NO:5)	AGCAAGAAAGCGAGCCAAT (SEQ ID NO:6)	
Endogenous	hTERT (NM_198253.2)	3' UTR of hTERT (NM_198253.2)	74
TERT	GCCCTCAGACTTCAAGACCA (SEQ ID NO:7)	GCTGCTGGTGTCTGCTCTC (SEQ ID NO:8)	
GAPDH	CAATGACCCCTTCATTGACC (SEQ ID NO:9)	TTGATTITGGAGGGATCTCG (SEQ ID NO:10)	159

[0197] *Senescence-associated β-galactosidase staining and cell size scoring.* β-gal staining was performed using the Senescence β-Galactosidase Staining Kit 15 (Cell Signaling Technology, Danvers MA, USA). At least 50 cells per population were scored in duplicate. Cell diameter was scored manually after trypsinization on a hemocytometer grid. Cristofalo and Kritchevsky (1969) *Med Exp Int J Exp Med.* 19:313–320.

20 [0198] *Statistics.* Student's T-tests and Pearson correlation coefficient calculations were performed using Microsoft Excel. Error bars represent the mean ± s.e.m.

Example 4. Delivery of TERT modRNA to cells by electroporation

[0199] The TERT modRNA compounds of the invention may also be delivered to cells by electroporation, as illustrated in Figures 18-20, using electroporation parameters, including the concentration of TERT modRNA, voltage wave form, 5 and electrode geometry appropriate for achieving optimal transfection efficiency and viability in a given cell type. Figure 21 shows the dependence of telomerase activity of the dose of TERT modRNA delivered by electroporation.

Example 5. Delivery of a modRNA to human blood cells

10 [0200] CD8+ T-cells were transfected with modRNA as follows. The buffy layer from whole human blood centrifuged on Lymphoprep (Axis-Shield) density gradient medium to obtain mononuclear cells which were washed twice and then depleted of non-CD8+ leukocytes using the Dynabeads Untouched Human CD8 T Cells Kit (Life Technologies). The CD8+ cells were stimulated using Dynabeads 15 Human T-Activator CD3.CD28 (Life Technologies) and cultured for 4 days using OpTimizer T-Cell media supplemented with 30,000 U/ml IL-2. The T-cells were then transfected with modRNA encoding nuclear GFP and TERT at a concentration of 50 μ g/ml in a volume of 20 μ l Nucleofector P3 solution containing Supplement 1. Cells were assayed for fluorescence and viability 24 20 hours after transfection. Results of the transfection are shown in Figure 22. The bright fluorescence demonstrates high copy number, with transfection efficiency over 90% with high viability.

25 **Example 6. Expression of telomerase in human keratinocytes using a telomerase modRNA**

[0201] As shown in Figure 23, electroporation of human keratinocytes with TERT modRNA results in the expression of telomerase activity in the cells. Human primary keratinocytes (Lonza) were suspended at a density of 3×10^7 30 cells/ml in OptiMEM media (Life Technologies) containing 50 μ g/ml of modRNA encoding either catalytically inactive (CI) TERT or wild type TERT. 10 μ l of the cell suspension was placed in a 1 mm gap cuvette and electroporated using a Gene

Pulser (BioRad) using 200 V, 1000 Ohms, and 25 microfarads. The cells were immediately returned to KGM-2 media (Lonza) and incubated for 24 hours before being harvested for measurement of telomerase activity using the gel-based TRAPeze assay (Millipore). Each 25 μ l TRAP reaction was performed using 1 μ g of total protein, with duplicate samples heated at 85°C for 10 minutes to inactivate telomerase. Untreated, electroporation only, and CI TERT samples served as negative controls, and 293T cells and TSR8 served as positive controls.

Example 6. Expression of a modRNA-encoded protein by delivery *in vivo*

10 [0202] modRNA delivered *in vivo* can result in the expression of a functional protein encoded by the modRNA. Kormann *et al.* (2012) *Nature Biotechnology* 29:154-157; Kariko *et al.* (2012) *Molecular Therapy* 20:948-93. This has been confirmed here by complexing 2 μ g of modRNA encoding luciferase with a cationic lipid vehicle (TransIT) and injecting intravenously in 50 μ l of OptiMEM (Life Technologies) in a rodent tail. The spleen was harvested and treated with luciferin and assayed for luciferase activity using an IVIS bioluminescence imager (Perkin-Elmer). As shown in Figure 24, luciferase activity was detected at the injection site and in the spleen (arrows).

15 20 [0203]

20 [0204] While specific examples have been provided, the above description is illustrative and not restrictive. Any one or more of the features of the previously described embodiments can be combined in any manner with one or more features of any other embodiments in the present invention. Furthermore, many variations 25 of the invention will become apparent to those skilled in the art upon review of the specification. The scope of the invention should, therefore, be determined by reference to the appended claims, along with their full scope of equivalents.

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 53896-6 Seq 09-NOV-15 v1.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

CLAIMS:

1. A method of extending telomeres, comprising the step of: administering a compound to an animal cell wherein said administering is in vitro, wherein the compound comprises a synthetic ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase, wherein the at least one modified nucleoside comprised in the synthetic ribonucleic acid confers reduced immunogenicity to the synthetic ribonucleic acid, wherein the telomerase reverse transcriptase is expressed transiently in the cell, and wherein at least one telomere is extended within the cell.
2. The method of claim 1, wherein the cell has at least one extensible telomere prior to the administering step.
3. The method of claim 1, wherein the cell is from a subject suffering from or at risk of an age-related illness, an age-related condition, or an age-related decline in function or appearance.
4. The method of claim 1, wherein the cell is a somatic cell of endodermal, mesodermal, or ectodermal lineage, or a germ line or embryonic cell.
5. The method of claim 1, wherein the cell is an induced pluripotent stem cell or a cell used to produce an induced pluripotent stem cell.
6. The method of claim 1, wherein the cell is a transdifferentiated cell or a cell used to produce a transdifferentiated cell.
7. The method of claim 1, wherein the cell is an isolated cell, and the administering step lasts no longer than 48 hours.
8. The method of claim 1, wherein the cell is an isolated cell, and the administering step lasts at least 2 hours.
9. The method of claim 1, wherein the cell is an isolated cell, and the administering step is performed no more than four times.

10. The method of claim 1, wherein the cell is an isolated cell, and the administering step is performed at least two times.

11. The method of claim 1, wherein the cell is an isolated cell, and the method further comprises the step of measuring telomerase activity in the cell.

12. The method of claim 1, wherein the administering step increases telomerase activity in the cell.

13. The method of claim 12, wherein the telomerase activity is transiently increased by at least 5%.

14. The method of claim 12, wherein the half-life of increased telomerase activity is no longer than 48 hours.

15. The method of claim 12, wherein the half-life of increased telomerase activity is at least 2 hours.

16. The method of claim 1, wherein the method further comprises the step of measuring telomere length in the cell.

17. The method of claim 1, wherein the administering step increases average telomere length in the cell.

18. The method of claim 17, wherein average telomere length in the cell is increased by at least 0.1 kb.

19. The method of claim 1, wherein the cell is an isolated cell, and the method further comprises the step of measuring population doubling capacity in the cell.

20. The method of claim 1, wherein the administering step increases population doubling capacity in the cell.

21. The method of claim 20, wherein the population doubling capacity increases by at least one population doubling.
22. The method of claim 1, wherein the cell is from a mammalian subject.
23. The method of claim 22, wherein the cell is from a human subject.
24. The method of claim 1, wherein the cell is an isolated cell.
25. The method of claim 1, wherein the cell is not an isolated cell.
26. The method of claim 1, wherein the administering step comprises electroporation.
27. The method of claim 1, wherein the at least one telomere is transiently extended within the cell.
28. Use of a compound comprising a synthetic ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase for extending telomeres in an animal cell, wherein the at least one modified nucleoside comprised in the synthetic ribonucleic acid confers reduced immunogenicity to the synthetic ribonucleic acid, and wherein the use is for transient expression of telomerase reverse transcriptase in the cell.
29. The use of claim 28, wherein the cell has at least one extensible telomere prior to the use of the compound.
30. The use of claim 28, wherein the cell is from a subject suffering from or at risk of an age-related illness, an age-related condition, or an age-related decline in function or appearance.
31. The use of claim 28, wherein the cell is a somatic cell of endodermal, mesodermal, or ectodermal lineage, or a germ line or embryonic cell.

32. The use of claim 28, wherein the cell is an induced pluripotent stem cell or a cell used to produce an induced pluripotent stem cell.

33. The use of claim 28, wherein the cell is a transdifferentiated cell or a cell used to produce a transdifferentiated cell.

34. The use of claim 28, wherein the cell is an isolated cell, and the compound is for administration over a period lasting no longer than 48 hours.

35. The use of claim 28, wherein the cell is an isolated cell, and the compound is for administration over a period lasting at least 2 hours.

36. The use of claim 28, wherein the cell is an isolated cell, and the compound is for administration no more than four times.

37. The use of claim 28, wherein the cell is an isolated cell, and the compound is for administration at least two times.

38. The use of claim 28, wherein the use is for increasing telomerase activity in the cell.

39. The use of claim 38, wherein the use is for transiently increasing telomerase activity by at least 5%.

40. The use of claim 38, wherein the use is for increasing a half-life of telomerase activity of no longer than 48 hours.

41. The use of claim 38, wherein the use is for increasing a half-life of telomerase activity of at least 2 hours.

42. The use of claim 28, wherein the use is for increasing average telomere length in the cell.

43. The use of claim 42, wherein the use is for increasing average telomere length in the cell by at least 0.1 kb.

44. The use of claim 28, wherein the use is for increasing population doubling capacity in the cell.

45. The use of claim 44, wherein the use is for increasing population doubling capacity by at least one population doubling.

46. The use of claim 28, wherein the cell is from a mammalian subject.

47. The use of claim 46, wherein the cell is from a human subject.

48. The use of claim 28, wherein the cell is an isolated cell.

49. The use of claim 28, wherein the cell is not an isolated cell.

50. The use of claim 28, wherein the use is for transient extension of at least one telomere within the cell.

51. A kit for extending telomeres in an animal cell, comprising:
a compound comprising a synthetic ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase, wherein the at least one modified nucleoside comprised in the synthetic ribonucleic acid confers reduced immunogenicity to the synthetic ribonucleic acid; and
instructions for using the compound for the transient expression of telomerase reverse transcriptase to extend telomeres.


52. The kit of claim 51, further comprising packaging materials.

53. The kit of claim 52, wherein the packaging materials are air-tight.

54. The kit of claim 52, wherein the packaging materials comprise a metal foil container.

55. The kit of claim 51, further comprising a desiccant.
56. The kit of claim 51, further comprising a culture medium.
57. The kit of claim 51, further comprising an RNase inhibitor.
58. The kit of claim 51, wherein the compound is sterile.
59. The kit of claim 51, further comprising:
a telomerase RNA component, a delivery vehicle, or both a telomerase RNA component and a delivery vehicle.
60. Use of a compound comprising a synthetic ribonucleic acid comprising at least one modified nucleoside and coding for a telomerase reverse transcriptase for treating an animal subject in need of, or that may benefit from, telomere extension, wherein the at least one modified nucleoside comprised in the synthetic ribonucleic acid confers reduced immunogenicity to the synthetic ribonucleic acid, and wherein the use is for transient expression of telomerase reverse transcriptase in a cell of the subject.
61. The use of claim 60, wherein the animal subject suffers from or is at risk of a disease resulting from shortened telomeres.
62. The use of claim 60, wherein the animal subject suffers from or is at risk of an age-related illness, an age-related condition, or an age-related decline in function or appearance.
63. The use of claim 60, wherein the use is for transient extension of telomeres within a cell of the subject animal.

Figure 1 (a)







Figure 1 (b)






Figure 1 (d)




Figure 2


Figure 3

Figure 4 (a)






Figure 4 (b)

Figure 4 (c)






Figure 4 (d)




Figure 4 (e)





Figure 4 (f)

Figure 4 (g)







Figure 5 (a)

Figure 5 (b)






Figure 5 (c)




Figure 6




Figure 7



**a**

Figure 8

**b**




Figure 9

Figure 10





Figure 11







Figure 12

Figure 13






Figure 15




Figure 16




Figure 17




Figure 18




Figure 19






Figure 20



Calculated Raw
values of Means
using X-Axis
channel(S) GFP

GFP - Panel 1

0	10.36
1	52.16

Figure 21




Figure 22




Figure 23




Figure 24

