



(51) International Patent Classification:

A61K 39/395 (2006.01) C07K 16/32 (2006.01)
A61K 45/06 (2006.01)

(21) International Application Number:

PCT/US2012/059683

(22) International Filing Date:

11 October 2012 (11.10.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/547,535	14 October 2011 (14.10.2011)	US
61/567,015	5 December 2011 (05.12.2011)	US
61/657,669	8 June 2012 (08.06.2012)	US
61/682,037	10 August 2012 (10.08.2012)	US
61/694,584	29 August 2012 (29.08.2012)	US

(71) Applicants: **GENENTECH, INC.** [US/US]; 1 DNA Way, South San Francisco, California 94080 (US). **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH). **ROCHE PRODUCTS LIMITED** [GB/GB]; 6 Falcon Way, Shire Park Hexagon Place, Welwyn Garden City Hertfordshire AL7 1TW (GB).

(72) Inventors; and

(71) Applicants : **ALAVATTAM, Sreedhara** [US/US]; c/o Genentech, Inc., 1 DNA Way, South San Francisco, California 94080 (US). **AMLER, Lukas C.** [DE/US]; c/o Genentech, Inc., 1 DNA Way, South San Francisco, California 94080 (US). **BENYUNES, Mark C.** [US/US]; c/o Genentech, Inc., 1 DNA Way, South San Francisco, California 94080 (US). **CLARK, Emma L.** [GB/GB]; c/o Roche Products Limited, 6 Falcon Way, Shire Park Hexagon Place, Welwyn Garden City Hertfordshire AL7 1TW (GB). **DE TOLEDO PELIZON, Christina H.** [BR/CA]; c/o F. Hoffmann-La Roche AG, Grenzacherstrasse 124, CH-4070 Basel (CH). **KWONG GLOVER, Zephania W.** [US/US]; c/o Genentech, Inc., 1 DNA Way, South San Francisco, California 94080 (US). **MITCHELL, Lada** [GB/CH]; c/o F. Hoffmann-La Roche AG, Grenzacherstrasse 124, CH-4070 Basel (CH). **RATNAYAKE, Jayantha** [GB/GB]; c/o Roche Products Limited, 6 Falcon Way, Shire Park Hexagon Place, Welwyn Garden City Hertfordshire AL7 1TW (GB). **ROSS, Graham A.** [GB/GB]; c/o Roche Products Limited, 6 Falcon Way, Shire Park Hexagon Place, Welwyn Garden City Hertfordshire AL7 1TW (GB). **WALKER, Ru-Amir** [US/US]; c/o Genentech, Inc., 1 DNA Way, South San Francisco, California 94080 (US).

(74) Agents: **LEE, Wendy M.** et al.; Genentech, Inc., 1 DNA Way, MS 49, South San Francisco, California 94080 (US).

[Continued on next page]

(54) Title: USES FOR AND ARTICLE OF MANUFACTURE INCLUDING HER2 DIMERIZATION INHIBITOR PERTUZUMAB




FIG. 8

(57) Abstract: The present application describes uses for and articles of manufacture including Pertuzumab, a first-in-class HER2 dimerization inhibitor. In particular, the application describes methods for extending progression free survival in a HER2-positive breast cancer patient population; combining two HER2 antibodies to treat HER2-positive cancer without increasing cardiac toxicity; treating early-stage HER2-positive breast cancer; treating HER2-positive cancer by co-administering a mixture of Pertuzumab and Trastuzumab from the same intravenous bag; treating HER2-positive metastatic gastric cancer; treating HER2-positive breast cancer with Pertuzumab, Trastuzumab and Vinorelbine; treating HER2-positive breast cancer with Pertuzumab, Trastuzumab and aromatase inhibitor; and treating low HER3 ovarian, primary peritoneal, or fallopian tube cancer. It also describes an article of manufacture comprising a vial with Pertuzumab therein and a package insert providing safety and/or efficacy data thereon; a method of making the article of manufacture; and a method of ensuring safe and effective use of Pertuzumab related thereto. In addition the application describes an intravenous (IV) bag containing a stable mixture of Pertuzumab and Trastuzumab suitable for administration to a cancer patient.



(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,

TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:

20 June 2013

USES FOR AND ARTICLE OF MANUFACTURE INCLUDING HER2 DIMERIZATION
INHIBITOR PERTUZUMAB

This non-provisional application filed under 37 CFR §1.53(b), claims the benefit under 35 USC §119(e) of U.S. Provisional Application Serial No. 61/547,535, filed on October 14, 2011, U.S. Provisional Application Serial No. 61/567,015, filed on December 5, 2011, U.S. Provisional Application Serial No. 61/657,669, filed on June 8, 2012, U.S. Provisional Application Serial No. 61/682,037, filed on August 10, 2012 and U.S. Provisional Application Serial No. 61/694,584, filed on August 29, 2012, which are incorporated by reference in entirety.

Field of the Invention

The present invention concerns uses for and articles of manufacture including Pertuzumab, a first-in-class HER2 dimerization inhibitor.

In particular, the invention concerns extending progression free survival in a HER2-positive breast cancer patient population; combining two HER2 antibodies to treat HER2-positive cancer without increasing cardiac toxicity; treating early-stage HER2-positive breast cancer; treating HER2-positive cancer by co-administering a mixture of Pertuzumab and Trastuzumab from the same intravenous bag; treating HER2-positive metastatic gastric cancer; treating HER2-positive breast cancer with Pertuzumab, Trastuzumab and Vinorelbine; treating HER2-positive breast cancer with Pertuzumab, Trastuzumab and aromatase inhibitor; and treating low HER3 ovarian, primary peritoneal, or fallopian tube cancer.

It also concerns an article of manufacture comprising a vial with Pertuzumab therein and a package insert providing safety and/or efficacy data thereon; a method of making the article of manufacture; and a method of ensuring safe and effective use of Pertuzumab related thereto.

In addition the invention concerns an intravenous (IV) bag containing a stable mixture of Pertuzumab and Trastuzumab suitable for administration to a cancer patient.

Background of the Invention

Members of the HER family of receptor tyrosine kinases are important mediators of cell growth, differentiation and survival. The receptor family includes four distinct members including epidermal growth factor receptor (EGFR, ErbB1, or HER1), HER2 (ErbB2 or p185^{neu}), HER3 (ErbB3) and HER4 (ErbB4 or tyro2). Members of the receptor family have been implicated in various types of human malignancy.

A recombinant humanized version of the murine anti-HER2 antibody 4D5 (huMAb4D5-8, rhuMAb HER2, Trastuzumab or HERCEPTIN[®]; U.S. Patent No. 5,821,337) is clinically active in

patients with HER2-overexpressing metastatic breast cancers that have received extensive prior anti-cancer therapy (Baselga *et al.*, *J. Clin. Oncol.* 14:737-744 (1996)).

Trastuzumab received marketing approval from the Food and Drug Administration September 25, 1998 for the treatment of patients with metastatic breast cancer whose tumors
5 overexpress the HER2 protein. At present, Trastuzumab is approved for use as a single agent or in combination with chemotherapy or hormone therapy in the metastatic setting, and as single agent or in combination with chemotherapy as adjuvant treatment for patients with early-stage HER2-positive breast cancer. Trastuzumab-based therapy is now the recommended treatment for patients with
10 HER2-positive early-stage breast cancer who do not have contraindications for its use (Herceptin[®] prescribing information; NCCN Guidelines, version 2.2011). Trastuzumab plus Docetaxel (or paclitaxel) is a registered standard of care in the first-line metastatic breast cancer (MBC) treatment setting (Slamon *et al.* *N Engl J Med.* 2001;344(11):783-792.; Marty *et al.* *J Clin Oncol.* 2005; 23(19):4265-4274).

While the administration of Trastuzumab has led to excellent results in the treatment of
15 breast cancer, recent data from a clinical trial of lapatinib appear to suggest that even with administration of Trastuzumab, HER2 plays an active role in tumor biology (Geyer *et al.*, *N Engl J Med* 2006; 355:2733-2743).

Patients treated with the HER2 antibody Trastuzumab are selected for therapy based on HER2 expression. See, for example, WO99/31140 (Paton *et al.*), US2003/0170234A1 (Hellmann, S.), and US2003/0147884 (Paton *et al.*); as well as WO01/89566, US2002/0064785, and
20 US2003/0134344 (Mass *et al.*). See, also, US Patent No. 6,573,043, US Patent No. 6,905,830, and US2003/0152987, Cohen *et al.*, concerning immunohistochemistry (IHC) and fluorescence *in situ* hybridization (FISH) for detecting HER2 overexpression and amplification. Thus, the optimal management of metastatic breast cancer now takes into account not only a patient's general
25 condition, medical history, and receptor status, but also the HER2 status.

Pertuzumab (also known as recombinant humanized monoclonal antibody 2C4 (rhuMab 2C4); Genentech, Inc, South San Francisco) represents the first in a new class of agents known as HER dimerization inhibitors (HDI) and functions to inhibit the ability of HER2 to form active heterodimers or homodimers with other HER receptors (such as EGFR/HER1, HER2, HER3 and
30 HER4). See, for example, Harari and Yarden *Oncogene* 19:6102-14 (2000); Yarden and Sliwkowski. *Nat Rev Mol Cell Biol* 2:127-37 (2001); Sliwkowski *Nat Struct Biol* 10:158-9 (2003); Cho *et al.* *Nature* 421:756-60 (2003); and Malik *et al.* *Pro Am Soc Cancer Res* 44:176-7 (2003).

Pertuzumab blockade of the formation of HER2-HER3 heterodimers in tumor cells has been demonstrated to inhibit critical cell signaling, which results in reduced tumor proliferation and
35 survival (Agus *et al.* *Cancer Cell* 2:127-37 (2002)).

Pertuzumab has undergone testing as a single agent in the clinic with a phase Ia trial in patients with advanced cancers and phase II trials in patients with ovarian cancer and breast cancer as

well as lung and prostate cancer. In a Phase I study, patients with incurable, locally advanced, recurrent or metastatic solid tumors that had progressed during or after standard therapy were treated with Pertuzumab given intravenously every 3 weeks. Pertuzumab was generally well tolerated. Tumor regression was achieved in 3 of 20 patients evaluable for response. Two patients had confirmed partial responses. Stable disease lasting for more than 2.5 months was observed in 6 of 21 patients (Agus *et al. Pro Am Soc Clin Oncol* 22:192 (2003)). At doses of 2.0-15 mg/kg, the pharmacokinetics of Pertuzumab was linear, and mean clearance ranged from 2.69 to 3.74 mL/day/kg and the mean terminal elimination half-life ranged from 15.3 to 27.6 days. Antibodies to Pertuzumab were not detected (Allison *et al. Pro Am Soc Clin Oncol* 22:197 (2003)).

US 2006/0034842 describes methods for treating ErbB-expressing cancer with anti-ErbB2 antibody combinations. US 2008/0102069 describes the use of Trastuzumab and Pertuzumab in the treatment of HER2-positive metastatic cancer, such as breast cancer. Baselga *et al., J Clin Oncol*, 2007 ASCO Annual Meeting Proceedings Part I, Vol. 25, No. 18S (June 20 Supplement), 2007:1004 report the treatment of patients with pre-treated HER2-positive breast cancer, which has progressed during treatment with Trastuzumab, with a combination of Trastuzumab and Pertuzumab. Portera *et al., J Clin Oncol*, 2007 ASCO Annual Meeting Proceedings Part I, Vol. 25, No. 18S (June 20 Supplement), 2007:1028 evaluated the efficacy and safety of Trastuzumab + Pertuzumab combination therapy in HER2-positive breast cancer patients, who had progressive disease on Trastuzumab-based therapy. The authors concluded that further evaluation of the efficacy of combination treatment was required to define the overall risk and benefit of this treatment regimen.

Pertuzumab has been evaluated in Phase II studies in combination with Trastuzumab in patients with HER2-positive metastatic breast cancer who have previously received Trastuzumab for metastatic disease. One study, conducted by the National Cancer Institute (NCI), enrolled 11 patients with previously treated HER2-positive metastatic breast cancer. Two out of the 11 patients exhibited a partial response (PR) (Baselga *et al., J Clin Oncol* 2007 ASCO Annual Meeting Proceedings; 25:18S (June 20 Supplement): 1004. The results of a Phase II neoadjuvant study evaluating the effect of a novel combination regimen of Pertuzumab and Trastuzumab plus chemotherapy (Docetaxel) in women with early-stage HER2-positive breast cancer, presented at the CTRC-AACR San Antonio Breast Cancer Symposium (SABCS), December 8-12, 2010, showed that the two HER2 antibodies plus Docetaxel given in the neoadjuvant setting prior to surgery significantly improved the rate of complete tumor disappearance (pathological complete response rate, pCR, of 45.8 percent) in the breast by more than half compared to Trastuzumab plus Docetaxel (pCR of 29.0 percent), $p=0.014$.

Patent Publications related to HER2 antibodies include: US Patent Nos. 5,677,171; 5,720,937; 5,720,954; 5,725,856; 5,770,195; 5,772,997; 6,165,464; 6,387,371; 6,399,063; 6,015,567; 6,333,169; 4,968,603; 5,821,337; 6,054,297; 6,407,213; 6,639,055; 6,719,971; 6,800,738; 8,075,890; 5,648,237; 7,018,809; 6,267,958; 6,685,940; 6,821,515; 7,060,268; 7,682,609; 7,371,376; 6,127,526;

6,333,398; 6,797,814; 6,339,142; 6,417,335; 6,489,447; 7,074,404; 7,531,645; 7,846,441; 7,892,549; 8,075,892; 6,573,043; 6,905,830; 7,129,051; 7,344,840; 7,468,252; 7,674,589; 7,919,254; 6,949,245; 7,485,302; 7,498,030; 7,501,122; 7,537,931; 7,618,631; 7,862,817; 7,041,292; 6,627,196; 7,371,379; 6,632,979; 7,097,840; 7,575,748; 6,984,494; 7,279,287; 7,811,773; 7,993,834; 8,076,066; 8,044,017; 5 7,435,797; 7,850,966; 7,485,704; 7,807,799; 8,142,784; 7,560,111; 7,879,325; 8,241,630; 7,449,184; 8,163,287; 7,700,299; 7,981,418; 8,247,397; and US 2010/0016556; US 2005/0244929; US 2001/0014326; US 2003/0202972; US 2006/0099201; US 2010/0158899; US 2011/0236383; US 2011/0033460; US 2008/0286280; US 2005/0063972; US 2006/0182739; US 2009/0220492; US 2003/0147884; US 2004/0037823; US 2005/0002928; US 2007/0292419; US 2008/0187533; US 10 2011/0250194; US 2012/0034213; US 2003/0152987; US 2005/0100944; US 2006/0183150; US 2008/0050748; US 2009/0155803; US 2010/0120053; US 2005/0244417; US 2007/0026001; US 2008/0160026; US 2008/0241146; US 2005/0208043; US 2005/0238640; US 2006/0034842; US 2006/0073143; US 2006/0193854; US 2006/0198843; US 2011/0129464; US 2007/0184055; US 2007/0269429; US 2008/0050373; US 2006/0083739; US 2009/0087432; US 2006/0210561; US 15 2002/0035736; US 2002/0001587; US 2008/0226659; US 2002/0090662; US 2006/0046270; US 2008/0108096; US 2007/0166753; US 2008/0112958; US 2009/0239236; US 2012/0034609; US 2012/0093838; US 2004/0082047; US 2012/0065381; US 2009/0187007; US 2011/0159014; US 2004/0106161; US 2011/0117096; US 2004/0258685; US 2009/0148402; US 2009/0099344; US 2006/0034840; US 2011/0064737; US 2005/0276812; US 2008/0171040; US 2009/0202536; US 20 2006/0013819; US 2012/0107391; US 2006/0018899; US 2009/0285837; US 2011/0117097; US 2006/0088523; US 2010/0015157; US 2006/0121044; US 2008/0317753; US 2006/0165702; US 2009/0081223; US 2006/0188509; US 2009/0155259; US 2011/0165157; US 2006/0204505; US 2006/0212956; US 2006/0275305; US 2012/0003217; US 2007/0009976; US 2007/0020261; US 2007/0037228; US 2010/0112603; US 2006/0067930; US 2007/0224203; US 2011/0064736; US 25 2008/0038271; US 2008/0050385; US 2010/0285010; US 2011/0223159; US 2008/0102069; US 2010/0008975; US 2011/0245103; US 2011/0246399; US 2011/0027190; US 2010/0298156; US 2011/0151454; US 2011/0223619; US 2012/0107302; US 2009/0098135; US 2009/0148435; US 2009/0202546; US 2009/0226455; US 2009/0317387; US 2011/0044977; US 2012/0121586.

Summary of the Invention

30 In a first aspect, the invention concerns a method for extending progression free survival in a HER2-positive breast cancer patient population by 6 months or more comprising administering Pertuzumab, Trastuzumab and chemotherapy (e.g. taxane, such as Docetaxel) to the patients in the population. Optionally the method results in an objective response rate of 80% or more in the patients in the population. The the breast cancer is optionally metastatic or locally recurrent, unresectable 35 breast cancer, or *de novo* Stage IV disease. In one embodiment, the patients in the population: have not received previous treatment or have relapsed after adjuvant therapy, have a left ventricular

ejection fraction (LVEF) of $\geq 50\%$ at baseline, and/or have an Eastern Cooperative Oncology Group performance status (ECOG PS) of 0 or 1. Optionally, the HER2-positive breast cancer is defined as immunohistochemistry (IHC) 3+ and/or fluorescence *in situ* hybridization (FISH) amplification ratio ≥ 2.0 . Optionally, the method reduces the risk of death by about 34% or more relative to a patient
5 treated with Trastuzumab and the chemotherapy.

In another aspect, the invention concerns a method of combining two HER2 antibodies to treat HER2-positive cancer without increasing cardiac toxicity in a HER2-positive cancer patient population, comprising administering Pertuzumab, Trastuzumab, and chemotherapy to the patients in the population. Optionally, cardiac toxicity in the patient population is monitored for incidence of
10 symptomatic left ventricular systolic dysfunction (LVSD) or congestive heart failure (CHF), or for decrease in left ventricular ejection fraction (LVEF). The HER2-positive cancer is optionally, breast cancer, for example metastatic or locally recurrent, unresectable breast cancer, or *de novo* Stage IV disease.

In another aspect, the invention concerns an article of manufacture comprising a vial with
15 Pertuzumab therein and a package insert, wherein the package insert provides the safety data in Table 3 or Table 4 and/or the efficacy data in Table 2, Table 5, Figure 8, or Figure 10.

The invention additionally concerns a method for making an article of manufacture comprising packaging together a vial with Pertuzumab therein and a package insert, wherein the package insert provides the safety data in Table 3 or Table 4 and/or the efficacy data in Table 2,
20 Table 5, Figure 8, or Figure 10.

In a related aspect, the invention concerns a method of ensuring safe and effective use of Pertuzumab comprising packaging together a vial with Pertuzumab therein and a package insert, wherein the package insert provides the safety data in Table 3 or Table 4 and/or the efficacy data in Table 2, Table 5, Figure 8, or Figure 10.

25 Optionally, the article of manufacture comprises a single-dose vial containing about 420mg of Pertuzumab.

Optionally, the package insert further comprises the warning box in Example 4.

Optionally, the package insert further provides the Overall Survival (OS) efficacy data in Example 9 or Table 14.

30 In another aspect, the invention concerns a method of treating early-stage HER2-positive breast cancer comprising administering Pertuzumab, Trastuzumab, and chemotherapy to a patient with the breast cancer, wherein the chemotherapy comprises anthracycline-based chemotherapy (for example, 5-FU, epirubicin, and cyclophosphamide (FEC)), or carboplatin-based chemotherapy (for example, Docetaxel and Carboplatin). Optionally, Pertuzumab is administered concurrently with the
35 anthracycline-based chemotherapy or the carboplatin-based chemotherapy. In one embodiment of this method, Pertuzumab administration does not increase cardiac toxicity relative to the treatment

without Pertuzumab. Such treatment of early-stage HER2-positive breast cancer optionally comprises neoadjuvant or adjuvant therapy.

The invention further concerns a method of treating HER2-positive cancer in a patient comprising co-administering a mixture of Pertuzumab and Trastuzumab from the same intravenous bag to the patient. Such method optionally further comprises administering chemotherapy to the patient.

In a related aspect, the invention provides an intravenous (IV) bag containing a stable mixture of Pertuzumab and Trastuzumab suitable for administration to a cancer patient. The mixture is optionally in saline solution; e.g. comprising about 0.9% NaCl or about 0.45% NaCl. The IV bag is optionally a 250mL 0.9% saline polyolefin or polyvinyl chloride infusion bag. In one embodiment, the IV bag which contains a mixture of about 420mg or of about 840mg of Pertuzumab and from about 200mg to about 1000mg of Trastuzumab. In one embodiment, the mixture is stable for up to 24 hours at 5°C or 30°C. Stability of the mixture can be evaluated by one or more assays selected from: color, appearance and clarity (CAC), concentration and turbidity analysis, particulate analysis, size exclusion chromatography (SEC), ion-exchange chromatography (IEC), capillary zone electrophoresis (CZE), image capillary isoelectric focusing (iCIEF), or potency assay.

The present invention provides a new treatment regimen for gastric cancer. In particular, the present invention concerns the treatment of HER2-positive gastric cancer in human subjects with a combination of Trastuzumab, Pertuzumab and at least one chemotherapy.

In one aspect, the invention concerns a method of treating HER2-positive gastric cancer in a human subject, comprising administering to the subject Pertuzumab, Trastuzumab, and a chemotherapy.

In one aspect, the invention concerns a method of treating gastric cancer in a human subject comprising administering Pertuzumab to the subject with gastric cancer, wherein Pertuzumab is administered at a dose of 840 mg in all treatment cycles.

In another aspect, the invention concerns a method of improving survival in a human subject with HER2-positive gastric cancer, comprising administering to the subject Pertuzumab, Trastuzumab, and a chemotherapy.

In yet another aspect, the invention concerns Pertuzumab for use in the treatment of HER2-positive gastric cancer in a human subject in combination with Trastuzumab and a chemotherapy.

In a further aspect, the invention concerns the use of Pertuzumab in the preparation of a medicament for the treatment of HER2-positive gastric cancer, wherein the treatment comprises administration of Pertuzumab in combination with Trastuzumab and a chemotherapy.

In a still further aspect, the invention concerns the use of Trastuzumab in the preparation of a medicament for the treatment of HER2-positive gastric cancer, wherein the treatment comprises administration of Trastuzumab in combination with Pertuzumab and a chemotherapy.

In another aspect, the invention concerns a kit comprising a container comprising Pertuzumab and instructions for administration of the Pertuzumab to treat HER2-positive gastric cancer in a subject in combination with Trastuzumab and a chemotherapy.

In yet another aspect, the invention concerns a kit comprising a container comprising
5 Trastuzumab and instructions for administration of the Trastuzumab to treat HER2-positive gastric cancer in a subject in combination with Pertuzumab and a chemotherapy.

In all aspects, the gastric cancer can, for example, be non-resectable locally advanced gastric cancer, or metastatic gastric cancer, or advanced, post-operatively recurrent gastric cancer, which may not be amenable to curative therapy by known methods. In all aspects, the gastric cancer
10 includes adenocarcinoma of the stomach or gastroesophageal junction. In all aspects, in a particular embodiment the patient did not receive prior anti-cancer treatment for metastatic gastric cancer. In all aspects, in a particular embodiment, the chemotherapy comprises administration of a platin and/or fluoropyrimidine. In certain embodiments, the platin is cisplatin. In other embodiments, the fluoropyrimidine comprises capecitabine and/or 5-fluorouracil (5-FU). In all aspects, the patient's
15 HER2-positive status may, for example, be IHC 3+ or IHC 2+/ISH+. In all aspects, in particular embodiments, the treatment improves survival, including overall survival (OS) and/or progression free survival (PFS) and/or response rate (RR). In all aspects, in particular embodiments, the patient has an ECOG PS of 0-1. In all aspects, treatment cycles are generally separated from each other by four weeks or less, or by three weeks or less, or by two weeks or less, or by one week or less.

In a particular aspect, the invention concerns a method of treating HER2-positive non-resectable or metastatic adenocarcinoma of the stomach or gastroesophageal junction in a human patient who did not receive prior chemotherapy for metastatic disease, except prior adjuvant or neoadjuvant therapy completed more than six months before the current treatment, comprising
20 administering Pertuzumab, Trastuzumab, cisplatin, and capecitabine and/or fluorouracil (5-FU) to the patient in an amount to improve progression free survival (PFS) and/or overall survival (OS),
25 wherein the patient has an ECOG PS of 0-1. In a particular embodiment, the patient did not receive prior treatment with a platin.

In another aspect, the invention concerns a method of improving progression free survival in a patient with HER2-positive non-resectable or metastatic adenocarcinoma of the stomach or
30 gastroesophageal junction comprising administering Pertuzumab to the patient in combination with Trastuzumab and chemotherapy.

In yet a further aspect, the invention concerns a method of treating HER2-positive breast cancer in a patient comprising administering Pertuzumab, Trastuzumab and vinorelbine to the patient. Optionally the Pertuzumab and Trastuzumab are co-administered to the patient from a single
35 intravenous bag. The breast cancer is optionally metastatic or locally advanced. In one embodiment, the patient has not previously received systemic non-hormonal anticancer therapy in the metastatic setting.

In another aspect, the invention concerns a method of treating HER2-positive breast cancer in a patient comprising administering Pertuzumab, Trastuzumab, and aromatase inhibitor (e.g. anastrozole or letrozole) to the patient. Optionally, the breast cancer is hormone receptor-positive advanced breast cancer, wherein the hormone receptor is estrogen receptor (ER) and/or progesterone receptor (PgR), for example. According to this embodiment of the invention, patient has not previously received systemic nonhormonal anticancer therapy in the metastatic setting. Moreover, the patient herein optionally receives induction chemotherapy (e.g. comprising taxane).

In an additional embodiment, the invention concerns a method of treating a cancer patient comprising administering to the patient an initial dose of 840mg of Pertuzumab followed every 3 weeks thereafter by a dose of 420mg of Pertuzumab, and further comprising re-administering an 840mg dose of Pertuzumab to the patient if the time between two sequential 420mg doses is 6 weeks or more. Optionally, the method further comprises administering 420mg of Pertuzumab every 3 weeks after the re-administered 840mg dose. In one embodiment, the cancer patient has HER2-positive breast cancer.

In a further aspect, the invention concerns a method for treating HER2-positive metastatic or locally recurrent breast cancer in a patient comprising administering Pertuzumab, Trastuzumab and taxoid (e.g. Docetaxel, Paclitaxel, or *nab*-paclitaxel) to the patient, wherein the patient has been previously treated with a Trastuzumab and/or lapatinib as adjuvant or neoadjuvant therapy.

In yet a further aspect, the invention concerns a method for treating low HER3 ovarian, primary peritoneal, or fallopian tube cancer in a patient comprising administering Pertuzumab and chemotherapy to the patient, wherein the chemotherapy comprises taxoid (e.g. paclitaxel) or topotecan.

In an additional aspect, the invention concerns a method for treating low HER3 ovarian, primary peritoneal, or fallopian tube cancer in a patient comprising administering Pertuzumab and chemotherapy to the patient, wherein the low HER3 cancer expresses HER3 mRNA at a concentration ratio equal or lower than about 2.81 as assessed by polymerase chain reaction (PCR). In one embodiment, the chemotherapy comprises gemcitabine, carboplatin, paclitaxel, docetaxel, topotecan, or pegylated liposomal doxorubicin (PLD). Optionally, the chemotherapy comprises paclitaxel or topotecan. In one embodiment, the cancer is epithelial ovarian cancer that is platinum-resistant or platinum-refractory.

Brief Description of the Drawings

Figure 1 provides a schematic of the HER2 protein structure, and amino acid sequences for Domains I-IV (SEQ ID Nos. 1-4, respectively) of the extracellular domain thereof.

Figures 2A and 2B depict alignments of the amino acid sequences of the variable light (V_L) (Fig. 2A) and variable heavy (V_H) (Fig. 2B) domains of murine monoclonal antibody 2C4 (SEQ ID Nos. 5 and 6, respectively); V_L and V_H domains of variant 574/Pertuzumab (SEQ ID Nos. 7 and 8, respectively), and human V_L and V_H consensus frameworks (hum κ 1, light kappa subgroup I; humIII, heavy subgroup III) (SEQ ID Nos. 9 and 10, respectively). Asterisks identify differences between

variable domains of Pertuzumab and murine monoclonal antibody 2C4 or between variable domains of Pertuzumab and the human framework. Complementarity Determining Regions (CDRs) are in brackets.

Figures 3A and 3B show the amino acid sequences of Pertuzumab light chain (Fig. 3A; SEQ ID NO. 11) and heavy chain (Fig. 3B; SEQ ID No. 12). CDRs are shown in bold. Calculated molecular mass of the light chain and heavy chain are 23,526.22 Da and 49,216.56 Da (cysteines in reduced form). The carbohydrate moiety is attached to Asn 299 of the heavy chain.

Figures 4A and 4B show the amino acid sequences of Trastuzumab light chain (Fig. 4A; SEQ ID NO. 13) and heavy chain (Fig. 4B; SEQ ID NO. 14), respectively. Boundaries of the variable light and variable heavy domains are indicated by arrows.

Figures 5A and 5B depict a variant Pertuzumab light chain sequence (Fig. 5A; SEQ ID NO. 15) and a variant Pertuzumab heavy chain sequence (Fig. 5B; SEQ ID NO. 16), respectively.

Figure 6 shows the study schema in Example 1. ECOG = Eastern Cooperative Oncology Group; PD = progressive disease. Notes: Trastuzumab, Pertuzumab, and Cisplatin are administered by IV infusion on Day 1 of each 3-week cycle. Capecitabine is administered orally twice daily, from the evening of Day 1 to the morning of Day 15 of each 3-week cycle. (a) HER2-positive tumor defined as either IHC 3+ or IHC 2+ in combination with ISH + (i.e., IHC 3+/ISH + or IHC 2+/ISH +); (b) Trastuzumab at a loading dose of 8 mg/kg for Cycle 1 and a dose of 6 mg/kg for subsequent cycles; (c) Pertuzumab on Day 1 of each cycle, at a loading dose of 840 mg for Cycle 1 and a dose of 420 mg for Cycles 2–6.

Figure 7 depicts enrollment, intent-to-treat and safety populations, and patient withdrawals in the study in Example 3.

Figure 8 is a Kaplan-Meier Curve of Progression-Free Survival (PFS) as assessed by an Independent Review Facility (IRF) for the study in Example 3.

Figure 9 depicts PFS by Patient Subgroup for the study in Example 3.

Figure 10 depicts overall survival for the study in Example 3.

Figure 11 is an overview of the dosing schedule in HER2-positive, neoadjuvant breast cancer, patients with low cardiac risk factors in Example 5. Additional radiotherapy, hormonal therapy and chemotherapy post surgery and during adjuvant Trastuzumab treatment were allowed if considered necessary by the investigator.

Figure 12 depicts mean change in LVEF (central readings) for the study in Example 5.

Figure 13 shows pathological complete response (pCR) for the study in Example 5.

Figure 14 depicts pathological complete response by hormone receptor status in the Example 5 study.

Figure 15 depicts Pertuzumab SEC profile of Pertuzumab/Trastuzumab mixture (840mg) at 30°C in 0.9% saline PO IV infusion bags (1) Time= 0; (2) Time= 24 hrs. Expanded view; full view (inset).

Figure 16 shows Trastuzumab SEC profile of Pertuzumab/Trastuzumab mixture (840mg) at 30°C in 0.9% saline PO IV infusion bags (1) Time= 0; (2) Time= 24 hrs. Expanded view; full view (inset).

Figure 17 shows Pertuzumab IEC profile of Pertuzumab/ Trastuzumab mixture at 30°C in 0.9% saline PO IV infusion bags (1) Time= 0; (2) Time= 24 hrs. Full view.

Figure 18 depicts Trastuzumab IEC profile of Pertuzumab/ Trastuzumab mixture at 30°C in 0.9% saline PO IV infusion bags (1) Time= 0; (2) Time= 24 hrs. Expanded view; Full view (inset).

Figure 19 depicts CE-SDS LIF non-reduced profile of Pertuzumab/ Trastuzumab mixture at 30°C in 0.9% saline PO IV infusion bags (1) Time= 0; (2) Time= 24 hrs. Expanded view.

Figure 20 shows CE-SDS LIF reduced profile of Pertuzumab/Trastuzumab mixture at 30°C in 0.9% saline PO IV infusion bags (1) Time= 0; (2) Time= 24 hrs. Expanded view.

Figure 21 is CZE of Pertuzumab/ Trastuzumab mixture at 30°C in 0.9% saline PO IV infusion bags (1) Time= 0; (2) Time= 24 hrs. Full view.

Figure 22 shows iCIEF of Pertuzumab/ Trastuzumab mixture at 30°C in 0.9% saline PO IV infusion bags (1) Time= 0; (2) Time= 24 hours. Full view.

Figure 23 shows potency dose response curves ($\mu\text{g/mL}$ versus RFU) of Pertuzumab/ Trastuzumab mixture, Pertuzumab alone, and Trastuzumab alone in 0.9% saline PO IV infusion bags (1) Time = 0; (2) Time = 24 hours.

Figure 24 depicts Pertuzumab SEC profile of Pertuzumab/Trastuzumab mixture (1560mg) in 0.9% saline IV infusion bags (1) PO 5°C T0 ; (2) PO 5°C T24 hrs; (3) PO 30°C T0; (4) PO 30°C T24 hrs; (5) PVC 5°C T0; (6) PVC 5°C T24 hrs; (7) PVC 30°C T0; (8) PVC 30°C T24 hrs. Expanded view; full view (inset).

Figure 25 shows Trastuzumab SEC profile of Pertuzumab/Trastuzumab mixture (1560mg) in 0.9% saline IV infusion bags (1) PO 5°C T0 ; (2) PO 5°C T24 hrs; (3) PO 30°C T0; (4) PO 30°C T24 hrs; (5) PVC 5°C T0; (6) PVC 5°C T24 hrs; (7) PVC 30°C T0; (8) PVC 30°C T24 hrs. Expanded view; full view (inset).

Figure 26 shows Pertuzumab IEC (Pertuzumab-fast) profile of Pertuzumab/Trastuzumab mixture (1560mg) in 0.9% saline IV infusion bags (1) PO 5°C T0 ; (2) PO 5°C T24 hrs; (3) PO 30°C T0; (4) PO 30°C T24 hrs; (5) PVC 5°C T0; (6) PVC 5°C T24 hrs; (7) PVC 30°C T0; (8) PVC 30°C T24 hrs. Full view.

Figure 27 shows Trastuzumab IEC profile of Pertuzumab/Trastuzumab mixture (1560mg) in 0.9% saline IV infusion bags (1) PO 5°C T0 ; (2) PO 5°C T24 hrs; (3) PO 30°C T0; (4) PO 30°C T24 hrs; (5) PVC 5°C T0; (6) PVC 5°C T24 hrs; (7) PVC 30°C T0; (8) PVC 30°C T24 hrs. Full view.

Figure 28 depicts study schema for Example 7.

Figure 29 shows study design for Example 8.

Figure 30 shows study design for Part 1 of Example 11.

Figure 31 shows study design for Part 2 of Example 11.

Figure 32 shows the samples taken and time points for the phase IIa gastric cancer (GC) study in Example 1.

5 Figure 33 shows the demographics of the patient population in the two arms of the GC study, treated with 420 mg (Arm A) or 840 mg (Arm B) of Pertuzumab.

Figure 34 shows the GC history of the patients in Arms A and B, respectively.

Figure 35 shows the GC patient disposition in Arms A and B, respectively.

Figure 36 shows the Overall Response Rate in Arms A and B, respectively, of the GC study.

10 Figure 37 shows the results of Pertuzumab Day 42 concentration assessment in gastric cancer (GC) versus metastatic breast cancer (MBC). Day 42 Ctrough is ~37% lower in GC (JOSHUA 840/420mg) vs. MBC (CLEO 840/420mg). JOSHUA 840/420mg and 840/840mg regimens both result in Day 42 Ctrough $\geq 20\mu\text{g/mL}$ in 90% of patients. JOSHUA 840/840mg regimen results in Day 42 Ctrough in GC comparable to that observed in MBC (CLEO 840/420mg)

15

Detailed Description of the Preferred Embodiments

Glossary of some abbreviations used herein: adverse drug reaction (ADR), adverse event (AE), alkaline phosphatase (ALP), absolute neutrophil count (ANC), area under the concentration-time curve (AUC), capillary zone electrophoresis (CZE), color, appearance and clarity (CAC),

20 CLinical Evaluation Of Pertuzumab And TRAstuzumab (CLEOPATRA), confidence interval (CI), chromogenic *in situ* hybridization (CISH), maximum concentration (C_{max}), complete response (CR), case report form (CRF), computed tomography (CT), common terminology criteria for adverse events (CTCAE), Docetaxel (D), dose limiting toxicity (DLT), ethics committee (EC), epirubicin, cisplatin, and 5-fluorouracil (ECF), echocardiogram (ECHO), epidermal growth factor receptor

25 (EGFR), European Union (EU), estrogen receptor (ER), 5-fluorouracil, methotrexate, and doxorubicin (FAMTX), fluorescence *in situ* hybridization (FISH), 5-fluorouracil (5-FU), hazard ratio (HR), human epidermal growth factor receptor (EGFR), gastric cancer (GC), good clinical practice (GCP), human epidermal growth factor receptor 2 (HER2), ion exchange chromatography (IEC), immunohistochemistry (IHC), independent review facility (IRF), institutional review board (IRB), *in*

30 *situ* hybridization (ISH), intravenous (IV), image capillary isoelectric focusing (iCIEF), left ventricular ejection fraction (LVEF), mitomycin C, cisplatin, and 5-fluorouracil (MCF), magnetic resonance image (MRI), metastatic breast cancer (MBC), multiple-gated acquisition (MUGA), not significant (NS), overall survival (OS), pathological complete response (pCR), polyolefin (PO), polyvinyl chloride (PVC), progressive disease (PD), progression free survival (PFS),

35 pharmacokinetic (PK), partial response (PR), progesterone receptor (PgR), response evaluation criteria in solid tumors (RECIST), serious adverse event (SAE), size exclusion chromatography (SEC), stable disease (SD), study management team (SMT), sterile water for injection (SWFI), time

to maximum plasma concentration (t_{\max}), upper limit of normal (ULN).

I. Definitions

5 The term “chemotherapy” as used herein refers to treatment comprising the administration of a chemotherapy, as defined hereinbelow.

“Survival” refers to the patient remaining alive, and includes overall survival as well as progression free survival.

10 “Overall survival” or “OS” refers to the patient remaining alive for a defined period of time, such as 1 year, 5 years, etc from the time of diagnosis or treatment. For the purposes of the clinical trial described in the example, overall survival (OS) is defined as the time from the date of randomization of patient population to the date of death from any cause.

15 “Progression free survival” or “PFS” refers to the patient remaining alive, without the cancer progressing or getting worse. For the purpose of the clinical trial described in the example, progression free survival (PFS) is defined as the time from randomization of study population to the first documented progressive disease, or unmanageable toxicity, or death from any cause, whichever occurs first. Disease progression can be documented by any clinically accepted methods, such as, for example, radiographical progressive disease, as determined by Response Evaluation Criteria in Solid Tumors (RECIST) (Therasse et al., *J Natl Ca Inst* 2000; 92(3):205-216), carcinomatous meningitis diagnosed by cytologic evaluation of cerebral spinal fluid, and/or medical photography to monitor chest wall recurrences of subcutaneous lesions.

20 By “extending survival” is meant increasing overall or progression free survival in a patient treated in accordance with the present invention relative to an untreated patient and/or relative to a patient treated with one or more approved anti-tumor agents, but not receiving treatment in accordance with the present invention. In a particular example, “extending survival” means
25 extending progression-free survival (PFS) and/or overall survival (OS) of cancer patients receiving the combination therapy of the present invention (e.g. treatment with a combination of Pertuzumab, Trastuzumab and a chemotherapy) relative to patients treated with Trastuzumab and the chemotherapy only. In another particular example, “extending survival” means extending progression-free survival (PFS) and/or overall survival (OS) of cancer patients receiving the
30 combination therapy of the present invention (e.g. treatment with a combination of Pertuzumab, Trastuzumab and a chemotherapy) relative to patients treated with Pertuzumab and the chemotherapy only.

An “objective response” refers to a measurable response, including complete response (CR) or partial response (PR).

35 By “complete response” or “CR” is intended the disappearance of all signs of cancer in response to treatment. This does not always mean the cancer has been cured.

“Partial response” or “PR” refers to a decrease in the size of one or more tumors or lesions,

or in the extent of cancer in the body, in response to treatment.

A “HER receptor” is a receptor protein tyrosine kinase which belongs to the HER receptor family and includes EGFR, HER2, HER3 and HER4 receptors. The HER receptor will generally comprise an extracellular domain, which may bind an HER ligand and/or dimerize with another HER receptor molecule; a lipophilic transmembrane domain; a conserved intracellular tyrosine kinase domain; and a carboxyl-terminal signaling domain harboring several tyrosine residues which can be phosphorylated. The HER receptor may be a “native sequence” HER receptor or an “amino acid sequence variant” thereof. Preferably the HER receptor is native sequence human HER receptor.

The expressions “ErbB2” and “HER2” are used interchangeably herein and refer to human HER2 protein described, for example, in Semba *et al.*, *PNAS (USA)* 82:6497-6501 (1985) and Yamamoto *et al.* *Nature* 319:230-234 (1986) (Genebank accession number X03363). The term “*erbB2*” refers to the gene encoding human ErbB2 and “*neu*” refers to the gene encoding rat p185^{neu}. Preferred HER2 is native sequence human HER2.

Herein, “HER2 extracellular domain” or “HER2 ECD” refers to a domain of HER2 that is outside of a cell, either anchored to a cell membrane, or in circulation, including fragments thereof. The amino acid sequence of HER2 is shown in Figure 1. In one embodiment, the extracellular domain of HER2 may comprise four domains: “Domain I” (amino acid residues from about 1-195; SEQ ID NO:1), “Domain II” (amino acid residues from about 196-319; SEQ ID NO:2), “Domain III” (amino acid residues from about 320-488; SEQ ID NO:3), and “Domain IV” (amino acid residues from about 489-630; SEQ ID NO:4) (residue numbering without signal peptide). See Garrett *et al.* *Mol. Cell.* 11: 495-505 (2003), Cho *et al.* *Nature* 421: 756-760 (2003), Franklin *et al.* *Cancer Cell* 5:317-328 (2004), and Plowman *et al.* *Proc. Natl. Acad. Sci.* 90:1746-1750 (1993), as well as Fig. 6 herein.

“HER3” or “ErbB3” herein refer to the receptor as disclosed, for example, in US Pat. Nos. 5,183,884 and 5,480,968 as well as Kraus *et al.* *PNAS (USA)* 86:9193-9197 (1989).

A “low HER3” cancer is one which expresses HER3 at a level less than the median level for HER3 expression in the cancer type. In one embodiment, the low HER3 cancer is epithelial ovarian, peritoneal, or fallopian tube cancer. HER3 DNA, protein, and/or mRNA level in the cancer can be evaluated to determine whether the cancer is a low HER3 cancer. See, for example, US Patent No. 7,981,418 for additional information about low HER3 cancer. Optionally, a HER3 mRNA expression assay is performed in order to determine that the cancer is a low HER3 cancer. In one embodiment, HER3 mRNA level in the cancer is evaluated, e.g. using polymerase chain reaction (PCR), such as quantitative reverse transcription PCR (qRT-PCR). Optionally, the cancer expresses HER3 at a concentration ratio equal or lower than about 2.81 as assessed qRT-PCR, e.g. using a COBAS z480® instrument.

A “HER dimer” herein is a noncovalently associated dimer comprising at least two HER receptors. Such complexes may form when a cell expressing two or more HER receptors is exposed

to an HER ligand and can be isolated by immunoprecipitation and analyzed by SDS-PAGE as described in Sliwkowski *et al.*, *J. Biol. Chem.*, 269(20):14661-14665 (1994), for example. Other proteins, such as a cytokine receptor subunit (*e.g.* gp130) may be associated with the dimer. Preferably, the HER dimer comprises HER2.

5 A "HER heterodimer" herein is a noncovalently associated heterodimer comprising at least two different HER receptors, such as EGFR-HER2, HER2-HER3 or HER2-HER4 heterodimers.

 A "HER antibody" is an antibody that binds to a HER receptor. Optionally, the HER antibody further interferes with HER activation or function. Preferably, the HER antibody binds to the HER2 receptor. HER2 antibodies of interest herein are Pertuzumab and Trastuzumab.

10 "HER activation" refers to activation, or phosphorylation, of any one or more HER receptors. Generally, HER activation results in signal transduction (*e.g.* that caused by an intracellular kinase domain of a HER receptor phosphorylating tyrosine residues in the HER receptor or a substrate polypeptide). HER activation may be mediated by HER ligand binding to a HER dimer comprising the HER receptor of interest. HER ligand binding to a HER dimer may activate a kinase domain of
15 one or more of the HER receptors in the dimer and thereby results in phosphorylation of tyrosine residues in one or more of the HER receptors and/or phosphorylation of tyrosine residues in additional substrate polypeptides(s), such as Akt or MAPK intracellular kinases.

 "Phosphorylation" refers to the addition of one or more phosphate group(s) to a protein, such as a HER receptor, or substrate thereof.

20 An antibody which "inhibits HER dimerization" is an antibody which inhibits, or interferes with, formation of a HER dimer. Preferably, such an antibody binds to HER2 at the heterodimeric binding site thereof. The most preferred dimerization inhibiting antibody herein is Pertuzumab or MAb 2C4. Other examples of antibodies which inhibit HER dimerization include antibodies which bind to EGFR and inhibit dimerization thereof with one or more other HER receptors (for example
25 EGFR monoclonal antibody 806, MAb 806, which binds to activated or "untethered" EGFR; see Johns *et al.*, *J. Biol. Chem.* 279(29):30375-30384 (2004)); antibodies which bind to HER3 and inhibit dimerization thereof with one or more other HER receptors; and antibodies which bind to HER4 and inhibit dimerization thereof with one or more other HER receptors.

 A "HER2 dimerization inhibitor" is an agent that inhibits formation of a dimer or
30 heterodimer comprising HER2.

 A "heterodimeric binding site" on HER2, refers to a region in the extracellular domain of HER2 that contacts, or interfaces with, a region in the extracellular domain of EGFR, HER3 or HER4 upon formation of a dimer therewith. The region is found in Domain II of HER2 (SEQ ID NO: 15). Franklin *et al. Cancer Cell* 5:317-328 (2004).

35 A HER2 antibody that "binds to a heterodimeric binding site" of HER2, binds to residues in Domain II (SEQ ID NO: 2) and optionally also binds to residues in other of the domains of the HER2 extracellular domain, such as domains I and III, SEQ ID NOs: 1 and 3), and can sterically hinder, at

least to some extent, formation of a HER2-EGFR, HER2-HER3, or HER2-HER4 heterodimer. Franklin *et al. Cancer Cell* 5:317-328 (2004) characterize the HER2-Pertuzumab crystal structure, deposited with the RCSB Protein Data Bank (ID Code IS78), illustrating an exemplary antibody that binds to the heterodimeric binding site of HER2.

5 An antibody that “binds to domain II” of HER2 binds to residues in domain II (SEQ ID NO: 2) and optionally residues in other domain(s) of HER2, such as domains I and III (SEQ ID NOs: 1 and 3, respectively). Preferably the antibody that binds to domain II binds to the junction between domains I, II and III of HER2.

 For the purposes herein, “Pertuzumab” and “rhuMAb 2C4”, which are used interchangeably, 10 refer to an antibody comprising the variable light and variable heavy amino acid sequences in SEQ ID NOs: 7 and 8, respectively. Where Pertuzumab is an intact antibody, it preferably comprises an IgG1 antibody; in one embodiment comprising the light chain amino acid sequence in SEQ ID NO: 11 or 15, and heavy chain amino acid sequence in SEQ ID NO: 12 or 16. The antibody is optionally produced by recombinant Chinese Hamster Ovary (CHO) cells. The terms “Pertuzumab” and 15 “rhuMAb 2C4” herein cover biosimilar versions of the drug with the United States Adopted Name (USAN) or International Nonproprietary Name (INN): Pertuzumab.

 For the purposes herein, “Trastuzumab” and rhuMAb4D5”, which are used interchangeably, refer to an antibody comprising the variable light and variable heavy amino acid sequences from within SEQ ID Nos: 13 and 14, respectively. Where Trastuzumab is an intact antibody, it preferably 20 comprises an IgG1 antibody; in one embodiment comprising the light chain amino acid sequence of SEQ ID NO: 13 and the heavy chain amino acid sequence of SEQ ID NO: 14. The antibody is optionally produced by Chinese Hamster Ovary (CHO) cells. The terms “Trastuzumab” and “rhuMAb4D5” herein cover biosimilar versions of the drug with the United States Adopted Name (USAN) or International Nonproprietary Name (INN): Trastuzumab.

25 The term “antibody” herein is used in the broadest sense and specifically covers monoclonal antibodies, polyclonal antibodies, multispecific antibodies (*e.g.* bispecific antibodies), and antibody fragments, so long as they exhibit the desired biological activity.

 “Humanized” forms of non-human (*e.g.*, rodent) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized 30 antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized 35 antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable

domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992). Humanized HER2 antibodies specifically include Trastuzumab (HERCEPTIN®) as described in Table 3 of U.S. Patent 5,821,337 expressly incorporated herein by reference and as defined herein; and humanized 2C4 antibodies such as Pertuzumab as described and defined herein.

An "intact antibody" herein is one which comprises two antigen binding regions, and an Fc region. Preferably, the intact antibody has a functional Fc region.

"Antibody fragments" comprise a portion of an intact antibody, preferably comprising the antigen binding region thereof. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragment(s).

"Native antibodies" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end. The constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains.

The term "hypervariable region" when used herein refers to the amino acid residues of an antibody which are responsible for antigen-binding. The hypervariable region generally comprises amino acid residues from a "complementarity determining region" or "CDR" (*e.g.* residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) and/or those residues from a "hypervariable loop" (*e.g.* residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). "Framework Region" or "FR" residues are those variable domain residues other than the hypervariable region residues as herein defined.

The term "Fc region" herein is used to define a C-terminal region of an immunoglobulin

heavy chain, including native sequence Fc regions and variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy chain Fc region is usually defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue.

Unless indicated otherwise, herein the numbering of the residues in an immunoglobulin heavy chain is that of the EU index as in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991), expressly incorporated herein by reference. The “EU index as in Kabat” refers to the residue numbering of the human IgG1 EU antibody.

A “functional Fc region” possesses an “effector function” of a native sequence Fc region. Exemplary “effector functions” include C1q binding; complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (*e.g.* B cell receptor; BCR), etc. Such effector functions generally require the Fc region to be combined with a binding domain (*e.g.* an antibody variable domain) and can be assessed using various assays as herein disclosed, for example.

A “native sequence Fc region” comprises an amino acid sequence identical to the amino acid sequence of an Fc region found in nature. Native sequence human Fc regions include a native sequence human IgG1 Fc region (non-A and A allotypes); native sequence human IgG2 Fc region; native sequence human IgG3 Fc region; and native sequence human IgG4 Fc region as well as naturally occurring variants thereof.

A “variant Fc region” comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one amino acid modification, preferably one or more amino acid substitution(s). Preferably, the variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, *e.g.* from about one to about ten amino acid substitutions, and preferably from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of the parent polypeptide. The variant Fc region herein will preferably possess at least about 80% homology with a native sequence Fc region and/or with an Fc region of a parent polypeptide, and most preferably at least about 90% homology therewith, more preferably at least about 95% homology therewith.

Depending on the amino acid sequence of the constant domain of their heavy chains, intact antibodies can be assigned to different “classes”. There are five major classes of intact antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into “subclasses” (isotypes),

e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the different classes of antibodies are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

5 A “naked antibody” is an antibody that is not conjugated to a heterologous molecule, such as a cytotoxic moiety or radiolabel.

An “affinity matured” antibody is one with one or more alterations in one or more hypervariable regions thereof which result an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). Preferred affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are produced by procedures known in the art. Marks *et al.* *Bio/Technology* 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by: Barbas *et al.* *Proc Nat. Acad. Sci, USA* 91:3809-3813 (1994); Schier *et al.* *Gene* 169:147-155 (1995); Yelton *et al.* *J. Immunol.* 155:1994-2004 (1995); Jackson *et al.*, *J. Immunol.* 154(7):3310-9 (1995); and Hawkins *et al.*, *J. Mol. Biol.* 226:889-896 (1992).

A “deamidated” antibody is one in which one or more asparagine residues thereof has been derivitized, e.g. to an aspartic acid, a succinimide, or an iso-aspartic acid.

The terms “cancer” and “cancerous” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth.

20 “Gastric cancer” specifically includes metastatic or locally advanced non-resectable gastric cancer, including, without limitation, histologically confirmed adenocarcinoma of the stomach or gastroesophageal junction with inoperable (non-resectable) locally advanced or metastatic disease, not amenable to curative therapy, and post-operatively recurrent advanced gastric cancer, such as adenocarcinoma of the stomach or gastroesophageal junction, when the intent of the surgery was to cure the disease.

An “advanced” cancer is one which has spread outside the site or organ of origin, either by local invasion or metastasis. Accordingly, the term “advanced” cancer includes both locally advanced and metastatic disease.

30 A “refractory” cancer is one which progresses even though an anti-tumor agent, such as a chemotherapy, is being administered to the cancer patient. An example of a refractory cancer is one which is platinum refractory.

A “recurrent” cancer is one which has regrown, either at the initial site or at a distant site, after a response to initial therapy, such as surgery.

35 A “locally recurrent” cancer is cancer that returns after treatment in the same place as a previously treated cancer.

A “non-resectable” or “unresectable” cancer is not able to be removed (resected) by surgery.

“Early-stage breast cancer” herein refers to breast cancer that has not spread beyond the

breast or the axillary lymph nodes. Such cancer is generally treated with neoadjuvant or adjuvant therapy.

“Neoadjuvant therapy” refers to systemic therapy given prior to surgery.

“Adjuvant therapy” refers to systemic therapy given after surgery.

5 “Metastatic” cancer refers to cancer which has spread from one part of the body (e.g. the breast) to another part of the body.

Herein, a “patient” or “subject” is a human patient. The patient may be a “cancer patient,” *i.e.* one who is suffering or at risk for suffering from one or more symptoms of cancer, in particular gastric or breast cancer.

10 A “patient population” refers to a group of cancer patients. Such populations can be used to demonstrate statistically significant efficacy and/or safety of a drug, such as Pertuzumab.

A “relapsed” patient is one who has signs or symptoms of cancer after remission. Optionally, the patient has relapsed after adjuvant or neoadjuvant therapy.

15 A cancer or biological sample which “displays HER expression, amplification, or activation” is one which, in a diagnostic test, expresses (including overexpresses) a HER receptor, has amplified HER gene, and/or otherwise demonstrates activation or phosphorylation of a HER receptor.

A cancer or biological sample which “displays HER activation” is one which, in a diagnostic test, demonstrates activation or phosphorylation of a HER receptor. Such activation can be determined directly (*e.g.* by measuring HER phosphorylation by ELISA) or indirectly (*e.g.* by gene
20 expression profiling or by detecting HER heterodimers, as described herein).

A cancer cell with “HER receptor overexpression or amplification” is one which has significantly higher levels of a HER receptor protein or gene compared to a noncancerous cell of the same tissue type. Such overexpression may be caused by gene amplification or by increased transcription or translation. HER receptor overexpression or amplification may be determined in a
25 diagnostic or prognostic assay by evaluating increased levels of the HER protein present on the surface of a cell (*e.g.* via an immunohistochemistry assay; IHC). Alternatively, or additionally, one may measure levels of HER-encoding nucleic acid in the cell, *e.g.* via *in situ* hybridization (ISH), including fluorescent *in situ* hybridization (FISH; see WO98/45479 published October, 1998) and chromogenic *in situ* hybridization (CISH; see, *e.g.* Tanner et al., *Am. J. Pathol.* 157(5): 1467–1472
30 (2000); Bella et al., *J. Clin. Oncol.* 26: (May 20 suppl; abstr 22147) (2008)), southern blotting, or polymerase chain reaction (PCR) techniques, such as quantitative real time PCR (qRT-PCR). One may also study HER receptor overexpression or amplification by measuring shed antigen (*e.g.*, HER extracellular domain) in a biological fluid such as serum (see, *e.g.*, U.S. Patent No. 4,933,294 issued June 12, 1990; WO91/05264 published April 18, 1991; U.S. Patent 5,401,638 issued March 28, 1995;
35 and Sias et al. *J. Immunol. Methods* 132: 73-80 (1990)). Aside from the above assays, various *in vivo* assays are available to the skilled practitioner. For example, one may expose cells within the body of the patient to an antibody which is optionally labeled with a detectable label, *e.g.* a radioactive

isotope, and binding of the antibody to cells in the patient can be evaluated, *e.g.* by external scanning for radioactivity or by analyzing a biopsy taken from a patient previously exposed to the antibody.

5 A “HER2-positive” cancer comprises cancer cells which have higher than normal levels of HER2. Examples of HER2-positive cancer include HER2-positive breast cancer and HER2-positive gastric cancer. Optionally, HER2-positive cancer has an immunohistochemistry (IHC) score of 2+ or 3+ and/or an *in situ* hybridization (ISH) amplification ratio ≥ 2.0 .

10 Herein, an “anti-tumor agent” refers to a drug used to treat cancer. Non-limiting examples of anti-tumor agents herein include chemotherapy agents, HER dimerization inhibitors, HER antibodies, antibodies directed against tumor associated antigens, anti-hormonal compounds, cytokines, EGFR-targeted drugs, anti-angiogenic agents, tyrosine kinase inhibitors, growth inhibitory agents and antibodies, cytotoxic agents, antibodies that induce apoptosis, COX inhibitors, farnesyl transferase inhibitors, antibodies that binds oncofetal protein CA 125, HER2 vaccines, Raf or ras inhibitors, liposomal doxorubicin, topotecan, taxene, dual tyrosine kinase inhibitors, TLK286, EMD-7200, Pertuzumab, Trastuzumab, erlotinib, and bevacizumab.

15 The “epitope 2C4” is the region in the extracellular domain of HER2 to which the antibody 2C4 binds. In order to screen for antibodies which bind essentially to the 2C4 epitope, a routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Preferably the antibody blocks 2C4's binding to HER2 by about 50% or more. Alternatively, epitope mapping can be performed to
20 assess whether the antibody binds essentially to the 2C4 epitope of HER2. Epitope 2C4 comprises residues from Domain II (SEQ ID NO: 2) in the extracellular domain of HER2. 2C4 and Pertuzumab binds to the extracellular domain of HER2 at the junction of domains I, II and III (SEQ ID NOs: 1, 2, and 3, respectively). Franklin *et al. Cancer Cell* 5:317-328 (2004).

25 The “epitope 4D5” is the region in the extracellular domain of HER2 to which the antibody 4D5 (ATCC CRL 10463) and Trastuzumab bind. This epitope is close to the transmembrane domain of HER2, and within Domain IV of HER2 (SEQ ID NO: 4). To screen for antibodies which bind essentially to the 4D5 epitope, a routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Alternatively, epitope mapping can be performed to assess whether the antibody binds
30 essentially to the 4D5 epitope of HER2 (*e.g.* any one or more residues in the region from about residue 529 to about residue 625, inclusive of the HER2 ECD, residue numbering including signal peptide).

“Treatment” refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with cancer as well as those in which cancer is to be
35 prevented. Hence, the patient to be treated herein may have been diagnosed as having cancer or may be predisposed or susceptible to cancer.

The term “effective amount” refers to an amount of a drug effective to treat cancer in the

patient. The effective amount of the drug may reduce the number of cancer cells; reduce the tumor size; inhibit (*i.e.*, slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (*i.e.*, slow to some extent and preferably stop) tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the cancer. To the extent the drug may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or cytotoxic. The effective amount may extend progression free survival (*e.g.* as measured by Response Evaluation Criteria for Solid Tumors, RECIST, or CA-125 changes), result in an objective response (including a partial response, PR, or complete response, CR), increase overall survival time, and/or improve one or more symptoms of cancer (*e.g.* as assessed by FOSI).

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (*e.g.* At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³² and radioactive isotopes of Lu), chemotherapeutic agents, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof.

A "chemotherapy" is use of a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents, used in chemotherapy, include alkylating agents such as thiotepa and CYTOXAN® cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; TLK 286 (TELCYTA™); acetogenins (especially bullatacin and bullatacinone); delta-9-tetrahydrocannabinol (dronabinol, MARINOL®); beta-lapachone; lapachol; colchicines; betulinic acid; a camptothecin (including the synthetic analogue topotecan (HYCAMTIN®), CPT-11 (irinotecan, CAMPTOSAR®), acetylcamptothecin, scoplectin, and 9-aminocamptothecin); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); podophyllotoxin; podophyllinic acid; teniposide; cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; bisphosphonates, such as clodronate; antibiotics such as the enediyne antibiotics (*e. g.*, calicheamicin, especially calicheamicin gammaII and calicheamicin omegaII (see, *e.g.*, Agnew, *Chem Intl. Ed. Engl.*, 33: 183-186 (1994)) and anthracyclines such as annamycin, AD 32, alcarubicin, daunorubicin, dexrazoxane, DX-52-1, epirubicin, GPX-100, idarubicin, KRN5500, menogaril, dynemicin, including dynemicin A, an esperamicin, neocarzinostatin chromophore and related chromoprotein enediyne antiobiotic

chromophores, aclacinomysins, actinomycin, anthramycin, azaserine, bleomycins, cactinomycin, carabacin, carminomycin, carzinophilin, chromomycinis, dactinomycin, detorubicin, 6-diazo-5-oxo-L-norleucine, ADRIAMYCIN® doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin, liposomal doxorubicin, and deoxydoxorubicin), esorubicin, 5 marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, and zorubicin; folic acid analogues such as denopterin, pteropterin, and trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, and thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, 10 dideoxyuridine, doxifluridine, enocitabine, and floxuridine; androgens such as calusterone, dromostanolone propionate, epitio stanol, mepitio stanone, and testolactone; anti-adrenals such as aminoglutethimide, mitotane, and trilostane; folic acid replenisher such as folinic acid (leucovorin); aceglatone; anti-folate anti-neoplastic agents such as ALIMTA®, LY231514 pemetrexed, dihydrofolate reductase inhibitors such as methotrexate, anti-metabolites such as 5-fluorouracil (5-FU) and its prodrugs such as UFT, S-1 and capecitabine, and thymidylate synthase inhibitors and 15 glycineamide ribonucleotide formyltransferase inhibitors such as raltitrexed (TOMUDEX^{RM}, TDX); inhibitors of dihydropyrimidine dehydrogenase such as eniluracil; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; 20 lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, OR); razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); 25 urethan; vindesine (ELDISINE®, FILDESIN®); dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxanes; chloranbucil; gemcitabine (GEMZAR®); 6-thioguanine; mercaptopurine; platinum; platinum analogs or platinum-based analogs such as cisplatin, oxaliplatin and carboplatin; vinblastine (VELBAN®); etoposide (VP-16); ifosfamide; mitoxantrone; vincristine (ONCOVIN®); vinca alkaloid; vinorelbine 30 (NAVELBINE®); novantrone; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; pharmaceutically acceptable salts, acids or derivatives of any of the above; as well as combinations of two or more of the above such as CHOP, an abbreviation for a combined therapy of cyclophosphamide, doxorubicin, vincristine, and prednisolone, and FOLFOX, an abbreviation for a 35 treatment regimen with oxaliplatin (ELOXATINTM) combined with 5-FU and leucovorin.

Also included in this definition are anti-hormonal agents that act to regulate or inhibit

hormone action on tumors such as anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX® tamoxifen), raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and FARESTON® toremifene; aromatase inhibitors; and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; as well as troxacitabine (a 1,3-dioxolane nucleoside cytosine analog); antisense oligonucleotides, particularly those that inhibit expression of genes in signaling pathways implicated in adherent cell proliferation, such as, for example, PKC-alpha, Raf, H-Ras, and epidermal growth factor receptor (EGF-R); vaccines such as gene therapy vaccines, for example, ALLOVECTIN® vaccine, LEUVECTIN® vaccine, and VAXID® vaccine; PROLEUKIN® rIL-2; LURTOTECAN® topoisomerase I inhibitor; ABARELIX® rmRH; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

A “taxane” is a chemotherapy which inhibits mitosis and interferes with microtubules. Examples of taxanes include Paclitaxel (TAXOL®; Bristol-Myers Squibb Oncology, Princeton, N.J.); cremophor-free, albumin-engineered nanoparticle formulation of paclitaxel or *nab*-paclitaxel (ABRAXANE™; American Pharmaceutical Partners, Schaumburg, Illinois); and Docetaxel (TAXOTERE®; Rhône-Poulenc Rorer, Antony, France).

An “anthracycline” is a type of antibiotic that comes from the fungus *Streptococcus peucetius*, examples include: Daunorubicin, Doxorubicin, and Epirubicin, etc.

“Anthracycline-based chemotherapy” refers to a chemotherapy regimen that consists of or include one or more anthracycline. Examples include 5-FU, epirubicin, and cyclophosphamide (FEC); 5-FU, doxorubicin, and cyclophosphamide (FAC); doxorubicin and cyclophosphamide (AC); epirubicin and cyclophosphamide (EC); etc.

For the purposes herein, “carboplatin-based chemotherapy” refers to a chemotherapy regimen that consists of or includes one or more Carboplatins. An example is TCH (Docetaxel/TAXOL®, Carboplatin, and Trastuzumab/HERCEPTIN®).

An “aromatase inhibitor” inhibits the enzyme aromatase, which regulates estrogen production in the adrenal glands. Examples of aromatase inhibitors include: 4(5)-imidazoles, aminoglutethimide, MEGASE® megestrol acetate, AROMASIN® exemestane, formestane, fadrozole, RIVISOR® vorozole, FEMARA® letrozole, and ARIMIDEX® anastrozole. In one embodiment, the aromatase inhibitor herein is letrozole or anastrozole.

An “antimetabolite chemotherapy” is use of an agent which is structurally similar to a metabolite, but can not be used by the body in a productive manner. Many antimetabolite chemotherapy interferes with the production of the nucleic acids, RNA and DNA. Examples of antimetabolite chemotherapeutic agents include gemcitabine (GEMZAR®), 5-fluorouracil (5-FU), capecitabine (XELODA™), 6-mercaptopurine, methotrexate, 6-thioguanine, pemetrexed, raltitrexed, arabinosylcytosine ARA-C cytarabine (CYTOSAR-U®), dacarbazine (DTIC-DOME®), azacytosine,

deoxycytosine, pyridimidine, fludarabine (FLUDARA®), cladribine, 2-deoxy-D-glucose etc.

By “chemotherapy-resistant” cancer is meant that the cancer patient has progressed while receiving a chemotherapy regimen (*i.e.* the patient is “chemotherapy refractory”), or the patient has progressed within 12 months (for instance, within 6 months) after completing a chemotherapy regimen.

The term “platin” is used herein to refer to platinum based chemotherapy, including, without limitation, cisplatin, carboplatin, and oxaliplatin.

The term “fluoropyrimidine” is used herein to refer to an antimetabolite chemotherapy, including, without limitation, capecitabine, floxuridine, and fluorouracil (5-FU).

A “fixed ” or “flat” dose of a therapeutic agent herein refers to a dose that is administered to a human patient without regard for the weight (WT) or body surface area (BSA) of the patient. The fixed or flat dose is therefore not provided as a mg/kg dose or a mg/m² dose, but rather as an absolute amount of the therapeutic agent.

A “loading” dose herein generally comprises an initial dose of a therapeutic agent administered to a patient, and is followed by one or more maintenance dose(s) thereof. Generally, a single loading dose is administered, but multiple loading doses are contemplated herein. Usually, the amount of loading dose(s) administered exceeds the amount of the maintenance dose(s) administered and/or the loading dose(s) are administered more frequently than the maintenance dose(s), so as to achieve the desired steady-state concentration of the therapeutic agent earlier than can be achieved with the maintenance dose(s).

A “maintenance” dose herein refers to one or more doses of a therapeutic agent administered to the patient over a treatment period. Usually, the maintenance doses are administered at spaced treatment intervals, such as approximately every week, approximately every 2 weeks, approximately every 3 weeks, or approximately every 4 weeks, preferably every 3 weeks.

“Infusion” or “infusing” refers to the introduction of a drug-containing solution into the body through a vein for therapeutic purposes. Generally, this is achieved via an intravenous (IV) bag.

An “intravenous bag” or “IV bag” is a bag that can hold a solution which can be administered via the vein of a patient. In one embodiment, the solution is a saline solution (e.g. about 0.9% or about 0.45% NaCl). Optionally, the IV bag is formed from polyolefin or polyvinyl chloride.

By “co-administering” is meant intravenously administering two (or more) drugs during the same administration, rather than sequential infusions of the two or more drugs. Generally, this will involve combining the two (or more) drugs into the same IV bag prior to co-administration thereof.

“Cardiac toxicity” refers to any toxic side effect resulting from administration of a drug or drug combination. Cardiac toxicity can be evaluated based on any one or more of: incidence of symptomatic left ventricular systolic dysfunction (LVSD) or congestive heart failure (CHF), or decrease in left ventricular ejection fraction (LVEF).

The phrase “without increasing cardiac toxicity” for a drug combination including

Pertuzumab refers to an incidence of cardiac toxicity that is equal or less than that observed in patients treated with drugs other than Pertuzumab in the drug combination (e.g. equal or less than that resulting from administration of Trastuzumab and the chemotherapy, e.g. Docetaxel).

5 A “vial” is a container suitable for holding a liquid or lyophilized preparation. In one embodiment, the vial is a single-use vial, e.g. a 20-cc single-use vial with a stopper.

A “package insert” is a leaflet that, by order of the Food and Drug Administration (FDA) or other Regulatory Authority, must be placed inside the package of every prescription drug. The leaflet generally includes the trademark for the drug, its generic name, and its mechanism of action; states its indications, contraindications, warnings, precautions, adverse effects, and dosage forms; and includes
10 instructions for the recommended dose, time, and route of administration.

The expression “safety data” concerns the data obtained in a controlled clinical trial showing the prevalence and severity of adverse events to guide the user regarding the safety of the drug, including guidance on how to monitor and prevent adverse reactions to the drug. Table 3 and Table 4 herein provide safety data for Pertuzumab. The safety data comprises any one or more (e.g. two,
15 three, four or more) of the most common adverse events (AEs) or adverse reactions (ADRs) in Tables 3 and 4. For example, the safety data comprises information about neutropenia, febrile neutropenia, diarrhea and/or cardiac toxicity as disclosed herein.

“Efficacy data” refers to the data obtained in controlled clinical trial showing that a drug effectively treats a disease, such as cancer. Efficacy data for Pertuzumab is provided in the examples
20 herein. As to HER2-positive metastatic or locally recurrent, unresectable breast cancer, efficacy data for Pertuzumab is found in Table 2, Table 5, Figure 8 and Figure 10 herein. The safety data comprises any one or more (e.g. two, three, four or more) of the primary endpoint (progression free survival, PFS, by IRF) and/or secondary endpoints (overall survival (OS); progression free survival (PFS) by investigator; objective response rate (ORR), including complete response (CR), partial
25 response (PR), stable disease (SD), and progressive disease (PD), and/or duration of response) in Table 2, Table 5, Figure 8 and Figure 10. For example, the efficacy data comprises information about progression free survival (PFS) and/or overall survival (OS) as disclosed herein.

By “stable mixture” when referring to a mixture of two or more drugs, such as Pertuzumab and Trastuzumab” means that each of the drugs in the mixture essentially retains its physical and
30 chemical stability in the mixture as evaluated by one or more analytical assays. Exemplary analytical assays for this purpose include: color, appearance and clarity (CAC), concentration and turbidity analysis, particulate analysis, size exclusion chromatography (SEC), ion-exchange chromatography (IEC), capillary zone electrophoresis (CZE), image capillary isoelectric focusing (iCIEF), and potency assay. In one embodiment, mixture has been shown to be stable for up to 24 hours at 5°C or
35 30°C.

A drug that is administered "concurrently" with one or more other drugs is administered during the same treatment cycle, on the same day of treatment as the one or more other drugs, and, optionally, at the same time as the one or more other drugs. For instance, for cancer therapies given every 3-weeks, the concurrently administered drugs are each administered on day-1 of a 3-week
5 cycle.

II. Antibody and Chemotherapy Compositions

The HER2 antigen to be used for production of antibodies may be, *e.g.*, a soluble form of the extracellular domain of a HER2 receptor or a portion thereof, containing the desired epitope. Alternatively, cells expressing HER2 at their cell surface (*e.g.* NIH-3T3 cells transformed to
10 overexpress HER2; or a carcinoma cell line such as SK-BR-3 cells, see Stancovski *et al. PNAS (USA)* 88:8691-8695 (1991)) can be used to generate antibodies. Other forms of HER2 receptor useful for generating antibodies will be apparent to those skilled in the art.

Various methods for making monoclonal antibodies herein are available in the art. For example, the monoclonal antibodies may be made using the hybridoma method first described by
15 Kohler *et al., Nature*, 256:495 (1975), by recombinant DNA methods (U.S. Patent No. 4,816,567).

The anti-HER2 antibodies used in accordance with the present invention, Trastuzumab and Pertuzumab, are commercially available.

(i) Humanized antibodies

Methods for humanizing non-human antibodies have been described in the art. Preferably, a
20 humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al., Nature*, 321:522-525 (1986); Riechmann *et al., Nature*, 332:323-327 (1988); Verhoeven *et al., Science*, 239:1534-1536 (1988)), by
25 substituting hypervariable region sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some hypervariable region residues and possibly some FR residues are
30 substituted by residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of
35 the rodent is then accepted as the human framework region (FR) for the humanized antibody (Sims *et*

al., *J. Immunol.*, 151:2296 (1993); Chothia *et al.*, *J. Mol. Biol.*, 196:901 (1987)). Another method uses a particular framework region derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta *et al.*, *J.*
5 *Immunol.*, 151:2623 (1993)).

It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and
10 humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence
15 the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

20 US Patent No. 6,949,245 describes production of exemplary humanized HER2 antibodies which bind HER2 and block ligand activation of a HER receptor.

Humanized HER2 antibodies specifically include Trastuzumab (HERCEPTIN®) as described in Table 3 of U.S. Patent 5,821,337 expressly incorporated herein by reference and as defined herein; and humanized 2C4 antibodies such as Pertuzumab as described and defined herein.

25 The humanized antibodies herein may, for example, comprise nonhuman hypervariable region residues incorporated into a human variable heavy domain and may further comprise a framework region (FR) substitution at a position selected from the group consisting of 69H, 71H and 73H utilizing the variable domain numbering system set forth in Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda,
30 MD (1991). In one embodiment, the humanized antibody comprises FR substitutions at two or all of positions 69H, 71H and 73H.

An exemplary humanized antibody of interest herein comprises variable heavy domain complementarity determining residues GFTFTDYTMX (SEQ ID NO: 17), where X is preferably D or S; DVNPNSGGSIYNQRFKG (SEQ ID NO:18); and/or NLGPSFYFDY (SEQ ID NO:19),
35 optionally comprising amino acid modifications of those CDR residues, *e.g.* where the modifications essentially maintain or improve affinity of the antibody. For example, an antibody variant for use in the methods of the present invention may have from about one to about seven or about five amino

acid substitutions in the above variable heavy CDR sequences. Such antibody variants may be prepared by affinity maturation, *e.g.*, as described below.

The humanized antibody may comprise variable light domain complementarity determining residues KASQDV SIGVA (SEQ ID NO:20); SASYX¹X²X³, where X¹ is preferably R or L, X² is preferably Y or E, and X³ is preferably T or S (SEQ ID NO:21); and/or QQYYIYPYT (SEQ ID NO:22), *e.g.* in addition to those variable heavy domain CDR residues in the preceding paragraph. Such humanized antibodies optionally comprise amino acid modifications of the above CDR residues, *e.g.* where the modifications essentially maintain or improve affinity of the antibody. For example, the antibody variant of interest may have from about one to about seven or about five amino acid substitutions in the above variable light CDR sequences. Such antibody variants may be prepared by affinity maturation, *e.g.*, as described below.

The present application also contemplates affinity matured antibodies which bind HER2. The parent antibody may be a human antibody or a humanized antibody, *e.g.*, one comprising the variable light and/or variable heavy sequences of SEQ ID Nos. 7 and 8, respectively (*i.e.* comprising the VL and/or VH of Pertuzumab). An affinity matured variant of Pertuzumab preferably binds to HER2 receptor with an affinity superior to that of murine 2C4 or Pertuzumab (*e.g.* from about two or about four fold, to about 100 fold or about 1000 fold improved affinity, *e.g.* as assessed using a HER2-extracellular domain (ECD) ELISA). Exemplary variable heavy CDR residues for substitution include H28, H30, H34, H35, H64, H96, H99, or combinations of two or more (*e.g.* two, three, four, five, six, or seven of these residues). Examples of variable light CDR residues for alteration include L28, L50, L53, L56, L91, L92, L93, L94, L96, L97 or combinations of two or more (*e.g.* two to three, four, five or up to about ten of these residues).

Humanization of murine 4D5 antibody to generate humanized variants thereof, including Trastuzumab, is described in U.S. Pat. Nos. 5,821,337, 6,054,297, 6,407,213, 6,639,055, 6,719,971, and 6,800,738, as well as Carter et al. PNAS (USA), 89:4285-4289 (1992). HuMAb4D5-8 (Trastuzumab) bound HER2 antigen 3-fold more tightly than the mouse 4D5 antibody, and had secondary immune function (ADCC) which allowed for directed cytotoxic activity of the humanized antibody in the presence of human effector cells. HuMAb4D5-8 comprised variable light (V_L) CDR residues incorporated in a V_L κ subgroup I consensus framework, and variable heavy (V_H) CDR residues incorporated into a V_H subgroup III consensus framework. The antibody further comprised framework region (FR) substitutions as positions: 71, 73, 78, and 93 of the V_H (Kabat numbering of FR residues; and a FR substitution at position 66 of the V_L (Kabat numbering of FR residues). Trastuzumab comprises non-A allotype human γ 1 Fc region.

Various forms of the humanized antibody or affinity matured antibody are contemplated. For example, the humanized antibody or affinity matured antibody may be an antibody fragment. Alternatively, the humanized antibody or affinity matured antibody may be an intact antibody, such

as an intact IgG1 antibody.

(ii) Pertuzumab compositions

In one embodiment of a HER2 antibody composition, the composition comprises a mixture of a main species Pertuzumab antibody and one or more variants thereof. The preferred embodiment herein of a Pertuzumab main species antibody is one comprising the variable light and variable heavy amino acid sequences in SEQ ID Nos. 7 and 8, and most preferably comprising a light chain amino acid sequence of SEQ ID No. 11, and a heavy chain amino acid sequence of SEQ ID No. 12 (including deamidated and/or oxidized variants of those sequences). In one embodiment, the composition comprises a mixture of the main species Pertuzumab antibody and an amino acid sequence variant thereof comprising an amino-terminal leader extension. Preferably, the amino-terminal leader extension is on a light chain of the antibody variant (*e.g.* on one or two light chains of the antibody variant). The main species HER2 antibody or the antibody variant may be an full length antibody or antibody fragment (*e.g.* Fab or F(ab')₂ fragments), but preferably both are full length antibodies. The antibody variant herein may comprise an amino-terminal leader extension on any one or more of the heavy or light chains thereof. Preferably, the amino-terminal leader extension is on one or two light chains of the antibody. The amino-terminal leader extension preferably comprises or consists of VHS-. Presence of the amino-terminal leader extension in the composition can be detected by various analytical techniques including, but not limited to, N-terminal sequence analysis, assay for charge heterogeneity (for instance, cation exchange chromatography or capillary zone electrophoresis), mass spectrometry, etc. The amount of the antibody variant in the composition generally ranges from an amount that constitutes the detection limit of any assay (preferably N-terminal sequence analysis) used to detect the variant to an amount less than the amount of the main species antibody. Generally, about 20% or less (*e.g.* from about 1% to about 15%, for instance from 5% to about 15%) of the antibody molecules in the composition comprise an amino-terminal leader extension. Such percentage amounts are preferably determined using quantitative N-terminal sequence analysis or cation exchange analysis (preferably using a high-resolution, weak cation-exchange column, such as a PROPAC WCX-10™ cation exchange column). Aside from the amino-terminal leader extension variant, further amino acid sequence alterations of the main species antibody and/or variant are contemplated, including but not limited to an antibody comprising a C-terminal lysine residue on one or both heavy chains thereof, a deamidated antibody variant, etc.

Moreover, the main species antibody or variant may further comprise glycosylation variations, non-limiting examples of which include antibody comprising a G1 or G2 oligosaccharide structure attached to the Fc region thereof, antibody comprising a carbohydrate moiety attached to a light chain thereof (*e.g.* one or two carbohydrate moieties, such as glucose or galactose, attached to one or two light chains of the antibody, for instance attached to one or more lysine residues), antibody comprising one or two non-glycosylated heavy chains, or antibody comprising a sialidated

oligosaccharide attached to one or two heavy chains thereof etc.

The composition may be recovered from a genetically engineered cell line, *e.g.* a Chinese Hamster Ovary (CHO) cell line expressing the HER2 antibody, or may be prepared by peptide synthesis.

- 5 For more information regarding exemplary Pertuzumab compositions, see US Patent Nos. 7,560,111 and 7,879,325 as well as US 2009/0202546A1.

(iii) *Trastuzumab compositions*

The Trastuzumab composition generally comprises a mixture of a main species antibody (comprising light and heavy chain sequences of SEQ ID NOS: 13 and 14, respectively), and variant forms thereof, in particular acidic variants (including deamidated variants). Preferably, the amount of such acidic variants in the composition is less than about 25%, or less than about 20%, or less than about 15%. See, U.S. Pat. No. 6,339,142. See, also, Harris et al., *J. Chromatography, B* 752:233-245 (2001) concerning forms of Trastuzumab resolvable by cation-exchange chromatography, including Peak A (Asn30 deamidated to Asp in both light chains); Peak B (Asn55 deamidated to isoAsp in one heavy chain); Peak 1 (Asn30 deamidated to Asp in one light chain); Peak 2 (Asn30 deamidated to Asp in one light chain, and Asp102 isomerized to isoAsp in one heavy chain); Peak 3 (main peak form, or main species antibody); Peak 4 (Asp102 isomerized to isoAsp in one heavy chain); and Peak C (Asp102 succinimide (Asu) in one heavy chain). Such variant forms and compositions are included in the invention herein.

20 (iv) *5-FU and Cisplatin*

There is no single, standard, globally accepted chemotherapeutic regimen for advanced gastric cancer, but 5-fluorouracil (5-FU) plus cisplatin is widely used for this indication. In Phase II studies in patients with no prior chemotherapy, 5-FU + cisplatin produced response rates of approximately 40% and median overall survival of 7-10.6 months (Lacave AJ, Baron FJ, Anton LM, et al. *Ann Oncol* 1991;2:751-754; Rougier P, Ducreux M, Mahjoubi M, et al. *Eur J Cancer* 1994;30A:1263-1269; Vanhoefer U, Wagner T, Lutz M, et al. *Eur J Cancer* 2001;37 Suppl 6: abstract S27.)

(v) *Capecitabine*

Capecitabine has been extensively tested in patients with advanced gastric cancer. Phase II efficacy results for capecitabine monotherapy show response rates of 19% and 26% and an overall survival of 8.1 and 10.0 months in studies by Koizumi et al 2003 (Koizumi W, Kurihara M, Sasai T, et al. *Cancer* 1993;72:658-62; Sakamoto J, Chin K, Kondo K, et al. *Anti-Cancer Drugs* 2006;17:2331-6. For capecitabine in combination with platinum, there are a number of studies showing response rates ranging from 28% to 65%, time to progression from 5.8 to 9 months, and overall survival from 10.1 to 12 months (Kang Y, Kang WK, Shin DB, et al. *J Clin Oncology* 2006;24 Suppl 18:abstract LBA4018; Park Y, Kim B, Ryoo B, et al. *Proc Am Soc Clin Oncol* 2006;24 Suppl 18: abstract 4079; Kim TW, Kang YK, Ahn JH, et al. *Ann Oncol* 2002;13:1893-8; Park YH, Kim BS, Ryoo BY, et al. *Br J Cancer* 2006;94:959-63).

III. Selecting Patients for Therapy

Detection of HER2 can be used to select patients for treatment in accordance with the present invention. Several FDA-approved commercial assays are available to identify HER2-positive cancer patients. These methods include HERCEPTEST[®] (Dako) and PATHWAY[®] HER2

5 (immunohistochemistry (IHC) assays) and PathVysion[®] and HER2 FISH pharmDx[™] (FISH assays). Users should refer to the package inserts of specific assay kits for information on the validation and performance of each assay.

For example, HER2 overexpression may be analyzed by IHC, *e.g.* using the HERCEPTEST[®] (Dako). Paraffin embedded tissue sections from a tumor biopsy may be subjected to the IHC assay
10 and accorded a HER2 protein staining intensity criteria as follows:

Score 0 no staining is observed or membrane staining is observed in less than 10% of tumor cells.

Score 1+ a faint/barely perceptible membrane staining is detected in more than 10% of the tumor cells. The cells are only stained in part of their membrane.

15 Score 2+ a weak to moderate complete membrane staining is observed in more than 10% of the tumor cells.

Score 3+ a moderate to strong complete membrane staining is observed in more than 10% of the tumor cells.

Those tumors with 0 or 1+ scores for HER2 overexpression assessment may be characterized
20 as HER2-negative, whereas those tumors with 2+ or 3+ scores may be characterized as HER2-positive.

Tumors overexpressing HER2 may be rated by immunohistochemical scores corresponding to the number of copies of HER2 molecules expressed per cell, and can be determined biochemically:

25 0 = 0-10,000 copies/cell,
1+ = at least about 200,000 copies/cell,
2+ = at least about 500,000 copies/cell,
3+ = at least about 2,000,000 copies/cell.

Overexpression of HER2 at the 3+ level, which leads to ligand-independent activation of the
30 tyrosine kinase (Hudziak *et al.*, *Proc. Natl. Acad. Sci. USA*, 84:7159-7163 (1987)), occurs in approximately 30% of breast cancers, and in these patients, relapse-free survival and overall survival are diminished (Slamon *et al.*, *Science*, 244:707-712 (1989); Slamon *et al.*, *Science*, 235:177-182 (1987)).

The presence of HER2 protein overexpression and gene amplification are highly correlated,
35 therefore, alternatively, or additionally, the use of *in situ* hybridization (ISH), *e.g.* fluorescent *in situ* hybridization (FISH), assays to detect gene amplification may also be employed for selection of

patients appropriate for treatment in accordance with the present invention. FISH assays such as the INFORM™ (sold by Ventana, Arizona) or PathVysion® (Vysis, Illinois) may be carried out on formalin-fixed, paraffin-embedded tumor tissue to determine the extent (if any) of HER2 amplification in the tumor.

5 Most commonly, HER2-positive status is confirmed using archival paraffin-embedded tumor tissue, using any of the foregoing methods.

Preferably, HER2-positive patients having a 2+ or 3+ IHC score or who are FISH or ISH positive are selected for treatment in accordance with the present invention.

See also US Patent No. 7,981,418 and Example 11 for alternative assays for screening
10 patients for therapy with Pertuzumab.

IV. Pharmaceutical Formulations

Therapeutic formulations of the HER2 antibodies used in accordance with the present invention are prepared for storage by mixing an antibody having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), generally in the form of lyophilized formulations or
15 aqueous solutions. Antibody crystals are also contemplated (see US Pat Appln 2002/0136719). Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as
20 octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine,
25 asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.* Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG). Lyophilized antibody formulations are described in WO 97/04801, expressly
30 incorporated herein by reference.

Lyophilized antibody formulations are described in U.S. Pat. Nos. 6,267,958, 6,685,940 and 6,821,515, expressly incorporated herein by reference. The preferred HERCEPTIN® (Trastuzumab) formulation is a sterile, white to pale yellow preservative-free lyophilized powder for intravenous
(IV) administration, comprising 440 mg Trastuzumab, 400 mg α,α -trehalose dehydrate, 9.9 mg L-
35 histidine-HCl, 6.4 mg L-histidine, and 1.8 mg polysorbate 20, USP. Reconstitution of 20 mL of bacteriostatic water for injection (BWFI), containing 1.1% benzyl alcohol as a preservative, yields a

multi-dose solution containing 21 mg/mL Trastuzumab, at pH of approximately 6.0. For further details, see the Trastuzumab prescribing information.

The preferred Pertuzumab formulation for therapeutic use comprises 30mg/mL Pertuzumab in 20mM histidine acetate, 120mM sucrose, 0.02% polysorbate 20, at pH 6.0. An alternate
5 Pertuzumab formulation comprises 25 mg/mL Pertuzumab, 10 mM histidine-HCl buffer, 240 mM sucrose, 0.02% polysorbate 20, pH 6.0.

The formulation of the placebo used in the clinical trials described in the Examples is equivalent to Pertuzumab, without the active agent.

The formulation herein may also contain more than one active compound as necessary for the
10 particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Various drugs which can be combined with the HER dimerization inhibitor are described in the Method Section below. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The formulations to be used for *in vivo* administration must be sterile. This is readily
15 accomplished by filtration through sterile filtration membranes.

V. Treatment Methods

In a first aspect of a treatment method herein, a method for extending progression free survival (PFS) in a HER2-positive breast cancer patient population by 6 months or more is provided which comprises administering Pertuzumab, Trastuzumab and chemotherapy (e.g. taxane such as
20 Docetaxel) to the patients in the population. Optionally, the patient population includes a suitable number of patients (e.g. 200 or more, 300 or more or 400 or more patients) so that a statistically significant extension of PFS in the population can be evaluated.

The phase III CLEOPATRA clinical data in Example 3 below show that median PFS assessed by investigators was 12.4 months with placebo plus Trastuzumab plus Docetaxel and 18.5
25 months with Pertuzumab plus Trastuzumab plus Docetaxel, thus the improvement in median PFS was 6 months or more (e.g. 6.1 months) relative to patients not receiving Pertuzumab (i.e. patients only receiving Trastuzumab and Docetaxel).

In an additional or alternative embodiment, a method of obtaining an objective response rate of 80% or more in a HER2-positive breast cancer patient population is provided which comprises
30 administering Pertuzumab, Trastuzumab and chemotherapy (e.g. taxane, such as Docetaxel) to the patients in the population.

In a related aspect, a method of combining two HER2 antibodies to treat HER2-positive cancer without increasing cardiac toxicity in a HER2-positive cancer patient population is provided which comprises administering Pertuzumab, Trastuzumab, and chemotherapy to the patients in the
35 population. Optionally, the patient population includes a suitable number of patients (e.g. 200 or more, 300 or more or 400 or more patients) so that a statistically significant assessment of lack of cardiac toxicity resulting from the combination can be made. The phase III CLEOPATRA clinical

data in Example 3 below show that combining Pertuzumab and Trastuzumab does not exacerbate cardiac toxicity. Cardiac toxicity can be monitored for incidence of symptomatic left ventricular systolic dysfunction (LVSD) or congestive heart failure (CHF), or decrease in left ventricular ejection fraction (LVEF), e.g. as disclosed in Example 3 below.

5 Optionally, the breast cancer is metastatic or locally recurrent, unresectable breast cancer, or *de novo* Stage IV disease, is defined as immunohistochemistry (IHC) 3+ and/or fluorescence *in situ* hybridization (FISH) amplification ratio ≥ 2.0 .

 Optionally, the patients in the population have not received previous treatment or have relapsed after adjuvant therapy, have a left ventricular ejection fraction (LVEF) of $\geq 50\%$ at baseline,
10 and/or have an Eastern Cooperative Oncology Group performance status (ECOG PS) of 0 or 1.

 In an alternative embodiment, the invention concerns a method of treating early-stage HER2-positive breast cancer comprising administering Pertuzumab, Trastuzumab, and chemotherapy to a patient with the breast cancer, wherein the chemotherapy comprises anthracycline-based chemotherapy, or carboplatin-based chemotherapy. This aspect of the invention is supported by the
15 clinical data in Example 5. In one embodiment, the chemotherapy comprises anthracycline-based chemotherapy, e.g. comprising 5-FU, epirubicin, and cyclophosphamide (FEC). In an alternative embodiment, the chemotherapy comprises carboplatin-based chemotherapy, e.g. comprising taxane (e.g. Docetaxel), Carboplatin in addition to HERCEPTIN®/Trastuzumab (e.g. TCH regimen). In one embodiment, Pertuzumab is administered concurrently with the anthracycline-based chemotherapy or
20 with the carboplatin-based chemotherapy, e.g. wherein the Pertuzumab, Trastuzumab and chemotherapy are administered in 3-week cycles with Pertuzumab, Trastuzumab and the chemotherapy being administered on day-1 of each cycle. The data in the examples herein demonstrates that Pertuzumab administration does not increase cardiac toxicity relative to the treatment without Pertuzumab (i.e. relative to Trastuzumab with anthracycline-based chemotherapy (e.g.
25 FEC) and no Pertuzumab; or relative to Trastuzumab with carboplatin-based chemotherapy and no Pertuzumab (i.e. TCH). The early-stage HER2-positive breast cancer therapy contemplated herein includes neoadjuvant and adjuvant therapy.

 The invention herein also concerns a method of treating HER2-positive cancer in a patient comprising co-administering a mixture of Pertuzumab and Trastuzumab from the same intravenous
30 bag to the patient. This embodiment is applicable to treatment of any HER2-positive cancer, including HER2-positive breast cancer, HER2-positive gastric cancer, HER2-positive metastatic or locally recurrent, unresectable breast cancer, or *de novo* Stage IV disease, early-stage HER2-positive breast cancer, etc. Optionally, this method further comprises administering chemotherapy to the patient.

35 In yet another embodiment, the treatment methods of the present invention comprise, consist essentially of, or consist of the administration of Pertuzumab, Trastuzumab and a chemotherapy, such as a platin (e.g. cisplatin) and/or a fluoropurimidine (e.g. capecitabine and/or 5-fluorouracil (5-FU))

to treat HER2-positive gastric cancer.

In particular, the treatment methods of the present invention comprise, consist essentially of, or consist of the administration of Pertuzumab, Trastuzumab, and a chemotherapy, such as a platin and/or a fluoropurimidine, e.g. cisplatin and/or capecitabine and/or 5-fluorouracil (5-FU), to a human
5 patient with metastatic gastric cancer, non-resectable locally advanced gastric cancer, or post-operatively recurrent gastric cancer. In certain embodiments, the gastric cancer is not amenable to curative therapy.

In an alternative embodiment, a method of treating HER2-positive breast cancer in a patient is provided comprising administering Pertuzumab, Trastuzumab and vinorelbine to the patient. The
10 breast cancer according to this embodiment is optionally metastatic or locally advanced. Optionally, the patient has not previously received systemic non-hormonal anticancer therapy in the metastatic setting.

In another aspect, the invention provides a method of treating HER2-positive breast cancer in a patient comprising administering Pertuzumab, Trastuzumab, and aromatase inhibitor (e.g.
15 anastrozole or letrozole) to the patient. According to this embodiment, the breast cancer is advanced breast cancer, including hormone receptor-positive breast cancer such as estrogen receptor (ER)-positive and/or progesterone receptor (PgR)-positive breast cancer. Optionally, the patient has not previously received systemic nonhormonal anticancer therapy in the metastatic setting. This treatment method optionally further comprises administering induction chemotherapy (e.g.
20 comprising taxane) to the patient.

Therapy in accordance with the present invention extends progression-free survival (PFS) and/or overall survival (OS) of the patient treated.

The antibodies and chemotherapeutic treatments are administered to a human patient in accord with known methods. Specific administration schedules and formulations are described in the
25 examples herein.

According to one embodiment, Pertuzumab is administered at a dose that produces a steady-state C_{min} of ≥ 20 $\mu\text{g/mL}$ in 90% of patients receiving Pertuzumab and Trastuzumab.

According to one particular embodiment of the invention, a Pertuzumab of approximately 840mg (loading dose) is administered, followed by one or more doses of approximately 420mg
30 (maintenance dose(s)) of the antibody. The maintenance doses are preferably administered about every 3 weeks, for a total of at least two doses, until clinical progressive disease, or unmanageable toxicity, preferably up to about 6, or 7, or 8, or 9, or 10, or 11, or 12, or 13, or 14, or 15, or 16, or 17 or more doses. Longer treatment periods, including more treatment cycles, are also contemplated.

According to another particular embodiment, Pertuzumab is administered at a dose of 840
35 mg for all treatment cycles.

Trastuzumab typically is administered as an intravenous loading dose of about 8 mg/kg, followed by the administration of 6 mg/kg doses in subsequent cycles. Trastuzumab is typically

administered every 3 weeks until clinical progressive disease or unmanageable toxicity, preferably up to about 17 or more doses.

In a particular embodiment, Trastuzumab is administered as an intravenous (IV) infusion on Day 1 of each treatment cycle until investigator-assessed disease progression or unmanageable toxicity, at a loading dose of 8 mg/kg for Cycle 1 and a dose of 6 mg/kg for subsequent cycles.

In another particular embodiment, Pertuzumab is administered as an IV infusion on Day 1 of each cycle, for a total of six cycles or until investigator assessed disease progression or unmanageable toxicity, whichever occurs first, either at a loading dose of 840 mg for Cycle 1 and a dose of 420 mg for the subsequent cycles, or a loading dose of 840 mg for Cycle 1 and a dose of 840 mg for the subsequent cycles.

For treating gastric cancer, Cisplatin 80 mg/m² is typically administered as an IV infusion on Day 1 of each cycle, for a total of at least six cycles.

For treating gastric cancer, Capecitabine 1000 mg/m² is typically administered orally twice daily, from the evening of Day 1 to the morning of Day 15 of each cycle, for a total of at least six cycles. Capecitabine administration may be prolonged at the discretion of the attending clinician after careful risk-benefit assessment for individual patients.

Dosages and schedules for chemotherapy used to treat HER2-positive breast cancer are disclosed in the examples below, but other dosages and schedules are known and contemplated according to the invention herein.

VI. Articles of Manufacture

One embodiment of an article of manufacture herein comprises an intravenous (IV) bag containing a stable mixture of Pertuzumab and Trastuzumab suitable for administration to a cancer patient. Optionally, the mixture is in saline solution; for example comprising about 0.9% NaCl or about 0.45% NaCl. An exemplary IV bag is a polyolefin or polyvinyl chloride infusion bag, e.g. a 250mL IV bag. According to one embodiment of the invention, the mixture includes about 420mg or about 840mg of Pertuzumab and from about 200mg to about 1000mg of Trastuzumab (e.g. from about 400mg to about 900mg of Trastuzumab).

Optionally, the mixture in the IV bag is stable for up to 24 hours at 5°C or 30°C. Stability of the mixture can be evaluated by one or more assays selected from the group consisting of: color, appearance and clarity (CAC), concentration and turbidity analysis, particulate analysis, size exclusion chromatography (SEC), ion-exchange chromatography (IEC), capillary zone electrophoresis (CZE), image capillary isoelectric focusing (iCIEF), and potency assay.

In an alternative embodiment, the invention provides an article of manufacture comprising a vial with Pertuzumab therein and a package insert, wherein the package insert provides the safety data in Table 3 or Table 4 and/or the efficacy data in Table 2, Table 5, Figure 8, or Figure 10. Optionally, the vial is a single-dose vial containing about 420mg of Pertuzumab. In one embodiment, the vial is provided inside a cardboard carton.

In a related aspect, the invention concerns a method of making an article of manufacture comprising packaging together a vial with Pertuzumab therein and a package insert, wherein the package insert provides the safety data in Table 3 or Table 4 and/or the efficacy data in Table 2, Table 5, Figure 8, or Figure 10.

5 In a further related aspect, the invention provides a method of ensuring safe and effective use of Pertuzumab comprising packaging together a vial with Pertuzumab therein and a package insert, wherein the package insert provides the safety data in Table 3 or Table 4 and/or the efficacy data in Table 2, Table 5, Figure 8, or Figure 10.

10 VII. Deposit of Biological Materials

The following hybridoma cell lines have been deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, USA (ATCC):

	Antibody Designation	ATCC No.	Deposit Date
	4D5	ATCC CRL 10463	May 24, 1990
15	2C4	ATCC HB-12697	April 8, 1999

Further details of the invention are illustrated by the following non-limiting Examples. The disclosures of all citations in the specification are expressly incorporated herein by reference.

20

EXAMPLE 1

Phase IIa Study Evaluating Pertuzumab in Combination with Trastuzumab and Chemotherapy in Patients with HER2-Positive Advanced Gastric Cancer

Despite a sharp worldwide decline in incidence and a reduction in mortality during the second half of the twentieth century, gastric cancer remains the world's second leading cause of cancer mortality, after lung cancer (Parkin, D. *Oncogene* 23:6329–40 (2004)). The incidence of gastric cancer varies widely according to geographic region (Kelley et al. *J Clin Epidemiol* 56:1–9 (2003); Plummer et al. *Epidemiology of gastric cancer*. In: Butlet et al., editors. *Mechanisms of carcinogenesis: contribution of molecular epidemiology*. Lyon: IARC Scientific Publications No 157, IARC (2004)). In Japan, Korea, China, and certain countries in Central and South America, the incidence is 20 to 95 cases per 100,000 men. In contrast, in the United States, India, and Thailand, the incidence is 4 to 8 cases per 100,000 men. The incidence in Western Europe ranges from 37 cases per 100,000 men in parts of Italy to 12 per 100,000 men in France. The incidence in women follows a similar geographic pattern but is about 50% lower than that in men. There are clear epidemiological differences between cancer localized to the gastric cardia (gastroesophageal junction) and that localized to the rest of the stomach. Cancer of the cardia accounts for 39% of gastric cancer cases in white men in the United States but only 4% of gastric cancers in men in Japan.

For reasons that are not clear, cancer of the gastric cardia and lower esophagus has increased rapidly in developed countries since the 1970s.

To date, the only potentially curative treatment for gastric cancer is surgery. Survival rates for gastric cancer improved significantly in Japan in recent years as a result of earlier detection and better surgical techniques (Inoue et al. *Postgrad Med J* 81:419–24 (2005)). However, in Western Europe and North America, gastric cancer is often diagnosed at a late stage when resection is no longer possible. Consequently, the overall 5-year survival in these populations does not exceed 25% (Ajani, J. *The Oncologist* 10 Suppl 3:49–58 (2005); Catalano et al. *Clin Rev Oncol/Hematol* 54: 209–41 (2005)).

Regardless of their geographic region, patients with unresectable disease due to locally advanced growth or metastatic spread have a poor prognosis, with overall 5-year survival within the range of 5%–15% (Cunningham et al., *Annals of Oncology* 16 Suppl 1:i22–3(2005)). For patients with unresectable disease at diagnosis and for patients with recurrent disease after surgery, the main therapeutic option is chemotherapy (National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology. Gastric cancer. Version 1. National Comprehensive Cancer Network, (2006)). Chemotherapy given with palliative intent has been shown to be superior to best supportive care in patients with advanced gastric cancer (Wagner et al. *J Clin Oncol* 24:2903–9 (2006)).

Study BO18255 (ToGA) was a randomized, open-label, multicenter, international, comparative Phase III trial designed to evaluate the efficacy and safety of Trastuzumab in combination with chemotherapy compared with chemotherapy alone as first-line therapy in patients with inoperable, locally advanced or recurrent and/or metastatic HER2-positive adenocarcinoma of the stomach or gastroesophageal junction. The primary objective of the study was to compare overall survival for patients treated with Trastuzumab combined with fluoropyrimidine (5-FU or capecitabine) plus cisplatin. The results from Study BO18255 demonstrated a significant clinical benefit when Trastuzumab was used in combination with chemotherapy in patients with gastric cancer. Overall survival, the primary endpoint, was significantly improved in the Trastuzumab plus chemotherapy arm compared with the chemotherapy alone arm ($p = 0.0045$, log-rank test; hazard ratio, 0.74). The median survival time was 13.8 months in the Trastuzumab plus chemotherapy arm and 11.1 months in the chemotherapy alone arm, and the risk of death was decreased by 26% for patients in the Trastuzumab plus chemotherapy arm. All other secondary endpoints demonstrated clinical significance with similar hazard and odds ratios. (See, e.g. Bang et al., *Lancet* 28; 376(9742):687–97 (2010)).

As a result of this study, Trastuzumab is now indicated, including the EU and United States, in combination with cisplatin plus capecitabine or 5-FU, for the treatment of patients with HER2-positive metastatic gastric or gastroesophageal junction adenocarcinoma who have not received prior treatment for metastatic disease.

At present, there is no single, standard, globally accepted chemotherapeutic regimen for

advanced gastric cancer. Despite the success of the ToGA trial, there is a great need for providing new and effective treatment options for this serious condition. In particular, there is a need for novel therapeutic approaches that aim to avoid treatment-related morbidity and/or to increase survival in gastric cancer patients. Accordingly, this example is a randomized, multicenter, open-label study evaluating two different doses of Pertuzumab in patients with HER2-positive adenocarcinoma of the stomach or gastroesophageal junction. Patients are randomized in a 1:1 ratio to two treatment arms. Patients in Arm A receive a Pertuzumab loading dose of 840 mg for Cycle 1 and a dose of 420 mg for Cycles 2–6, and patients in Arm B receive Pertuzumab 840 mg for all six cycles. Patients in both treatment arms receive Trastuzumab, cisplatin, and capecitabine. Study schema are in Figure 6. The length of the study is approximately 24 months (4 months for recruitment and 20 months of follow-up after last patient recruited). The end of study will be when progressive disease has occurred in all patients, or all patients have withdrawn or discontinued from the study, whichever is earlier.

Target population

The trial involves approximately 30 patients.

Patients must meet the following criteria for study entry:

- Histologically confirmed adenocarcinoma of the stomach or gastroesophageal junction with inoperable locally advanced or metastatic disease, not amenable to curative therapy.
- Patients with advanced disease who present with a recurrence post-operatively (when intent of surgery was cure) are also eligible for entry.
- Measurable disease, according to the Response Evaluation Criteria in Solid Tumors (RECIST), v1.1, assessed using imaging techniques (computed tomography (CT) or magnetic resonance imaging (MRI)), or non-measurable disease that can be followed.
- HER2-positive tumor defined as either IHC 3+ or IHC 2+ in combination with ISH +, as assessed by central laboratory on primary or metastatic tumor. ISH positivity is defined as a ratio of ≥ 2.0 for the number of HER2 gene copies to the number of signals for CEP17.
- Availability of formalin-fixed paraffin-embedded (FFPE) tissue with at least 5 mm of invasive tumor for central confirmation of HER2 eligibility is mandatory.
- Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 or 1.
- Baseline left ventricular ejection fraction (LVEF) $\geq 55\%$ (measured by echocardiogram (ECHO) or multiple-gated acquisition (MUGA) scan).
- Life expectancy of at least 3 months.
- Male or female.
- Age ≥ 18 years.
- Signed informed consent.

- For women of childbearing potential and male participants with partners of childbearing potential: agreement to use a highly effective non-hormonal form of contraception or two effective forms of non-hormonal contraception by the patient and/or partner.
- Contraception use must continue for the duration of study treatment and for at least 6 months after the last dose of study medication.

Patients who meet any of the following criteria are excluded from study entry:

- Previous chemotherapy for advanced or metastatic disease, except that prior adjuvant or neoadjuvant therapy is allowed if at least 6 months has elapsed between completion of adjuvant or neoadjuvant therapy and enrollment in the study.
- Adjuvant or neoadjuvant therapy with a platin is not allowed.
- Lack of physical integrity of the upper gastrointestinal tract or malabsorption syndrome (e.g., patients with partial or total gastrectomy can enter the study, but not those with a jejunostomy probe).
- Active (significant or uncontrolled) gastrointestinal bleeding.
- Residual relevant toxicity resulting from previous therapy (e.g., neurological toxicity of \geq Grade \geq 2 (NCI CTCAE)), with the exception of alopecia.
- Other malignancy within the last 5 years, except for carcinoma *in situ* of the cervix, or basal cell carcinoma.
- Any of the following abnormal laboratory tests immediately prior to randomization:
 Serum total bilirubin >1.5 times the upper limit of normal (ULN) or, for patients with known Gilberts syndrome, serum total bilirubin $> 2 \times$ ULN
 For patients with no liver and no bone metastases:
 AST or ALT $>2.5 \times$ ULN, and alkaline phosphatase (ALP) $>2.5 \times$ ULN
 In patients with liver metastases and no bone metastases: AST or ALT $>5 \times$ ULN, and ALP $>2.5 \times$ ULN
 In patients with liver metastases and bone metastases: AST or ALT $>5 \times$ ULN, and ALP $>10 \times$ ULN;
 In patients with bone metastases and no liver metastases: AST or ALT $>2.5 \times$ ULN, and ALP $>10 \times$ ULN
 Albumin <25 g/L
 Creatinine clearance <60 mL/min
 Total white blood cell (WBC) count $<2500/\mu\text{L}$ ($<2.5 \times 10^9/\text{L}$)
 Absolute neutrophil count (ANC) $<1500/\mu\text{L}$ ($<1.5 \times 10^9/\text{L}$)
 Platelets $<100,000/\mu\text{L}$ ($<100 \times 10^9/\text{L}$)
- Serious cardiac illness or medical conditions including but not confined to:
 history of documented heart failure or systolic dysfunction (LVEF $<50\%$);

high-risk uncontrolled arrhythmias, such as atrial tachycardia with a heart rate ≥ 100 /min at rest;

significant ventricular arrhythmia (ventricular tachycardia) or higher-grade AV block (second-degree AV block Type 2 (Mobitz II) or third-degree AV block);

5 angina pectoris requiring anti-anginal medication;

clinically significant valvular heart disease;

evidence of transmural infarction on ECG; poorly controlled hypertension (e.g., systolic blood pressure >180 mmHg or diastolic blood pressure >100 mmHg);

10 dyspnea at rest due to complications of advanced malignancy or other disease, or requirement for supportive oxygen therapy;

treatment with chronic or high-dose corticosteroid therapy;

inhaled steroids and short courses of oral steroids for anti-emesis or as an appetite stimulant are allowed;

15 clinically significant hearing abnormality; known dihydropyrimidine dehydrogenase deficiency;

history or clinical evidence of brain metastases; serious uncontrolled systemic intercurrent illness (e.g., infections or poorly controlled diabetes).

- Pregnant or lactating

20 Women of childbearing potential must have a negative serum pregnancy test within 7 days prior to randomization, irrespective of the method of contraception used.

- Radiotherapy within 4 weeks prior to start of study treatment, or within 2 weeks prior to start of study treatment if palliative radiotherapy is given to bone metastatic site peripherally and patient recovers from any acute toxicity.

- Major surgery within 4 weeks prior to start of study treatment, without complete recovery.

25 • Known active infection with HIV, hepatitis B virus, or hepatitis C virus.

- Known hypersensitivity to any of the study drugs.

- Inability to comply with follow-up testing or procedures, as determined by the investigator.

Investigational Medical Products: Dose, Route and Regimen

Treatment cycles are 3 weeks in length.

30 • Trastuzumab is administered as an intravenous (IV) infusion on Day 1 of each cycle until investigator-assessed disease progression or unmanageable toxicity, at a loading dose of 8 mg/kg for Cycle 1 and a dose of 6 mg/kg for subsequent cycles.

- Pertuzumab is administered as an IV infusion on Day 1 of each cycle, for a total of six cycles or until investigator assessed disease progression or unmanageable toxicity, whichever occurs first, as

35 follows for each arm:

Arm A: Patients receive Pertuzumab at a loading dose of 840 mg for Cycle 1 and a dose of 420 mg for Cycles 2–6.

Arm B: Patients receive Pertuzumab at 840 mg for Cycles 1–6.

Non-Investigational Medical Products

5 Treatment cycles are 3 weeks in length.

- Cisplatin 80 mg/m² is administered as an IV infusion on Day 1 of each cycle, for a total of six cycles.
- Capecitabine 1000 mg/m² is administered orally twice daily, from the evening of Day 1 to the morning of Day 15 of each cycle, for a total of six cycles. (Capecitabine may be prolonged at the discretion of the investigator after careful risk–benefit assessment for individual patients.)

10

-

Formulations

Formulation of Pertuzumab

Each lot of the recombinant antibodies produced for clinical purposes meets viral safety requirements and the United States Pharmacopeia and the European Pharmacopoeia requirements for sterility. Each lot meets the required specifications for identity, purity, and potency.

15

Pertuzumab is provided as a single-use formulation containing 30 mg/mL Pertuzumab formulated in 20-mM L-histidine-acetate (pH 6.0), 120-mM sucrose, and 0.02% polysorbate 20. Each 20-cc vial (14.0 mL solution per vial) contains approximately 420 mg of Pertuzumab.

Formulation of Trastuzumab

20

Investigational Trastuzumab is supplied as a freeze-dried preparation at a nominal content of 150 mg per vial in most countries (vial size varies by country).

Trastuzumab is formulated in histidine, trehalose, and polysorbate 20. Once reconstituted, each solution contains 21 mg/mL of active drug at a pH of approximately 6.0.

Assessments

25

Efficacy

Investigator-assessed tumor response will be used to summarize best overall response at the end of Cycles 3 and 6 for each treatment arm, defined as patients with a complete or partial response as determined by RECIST.

30

Safety

Safety will be assessed through summaries of adverse events, changes in laboratory test results, and changes in vital signs.

Pharmacokinetics/ Pharmacodynamics

- 5 Minimum (trough) serum concentration (C_{\min}) for Pertuzumab at Day 43 will be assessed. In addition, PK parameters such as CL, Vss, AUC, and half-life will be estimated. The evaluation of PK parameters from data collected up to Day 43 will enable modeling and simulation for an estimated dose that predicts a steady-state trough of ≥ 20 $\mu\text{g/mL}$ in 90% of patients.

Statistical Analyses

10 Pharmacokinetic Analyses

- Individual and mean serum Pertuzumab concentration–time data will be tabulated and plotted by dose level. The serum pharmacokinetics of Pertuzumab will be summarized by estimating total exposure (area under the curve (AUC)), maximum serum concentration (C_{\max}), minimum serum concentration (C_{\min}), time to steady-state C_{\max} and C_{\min} , total serum clearance, volume of distribution, and elimination half-life ($t_{1/2}$). Estimates for these parameters will be tabulated and summarized by descriptive statistics (mean, standard deviation, minimum, and maximum). Depending on the observed Pertuzumab serum concentration–time data, a population PK approach may be used to estimate the dose that will achieve the PK target concentrations.

- Observed C_{\max} and C_{\min} for Trastuzumab will be tabulated and summarized by descriptive statistics for each specified PK sampling timepoint. For all PK analyses, actual times of sample collection (rather than scheduled) will be used.

PK parameters (AUC, C_{\max} , $t_{1/2}$) of Pertuzumab will be calculated using non-compartmental methods, and the systemic clearance will be derived from the plasma concentrations via standard methods.

25 Analysis Populations

Intent-to-Treat Population

All randomized patients who receive at least one dose of study medication will be included in the intent-to-treat population (patients will be assigned to treatment groups as randomized for analysis purposes).

30 *Safety Population*

All patients who received at least one dose of study medication will be included in the safety-evaluable population (patients will be assigned to treatment groups as treated.)

Sample Size

- The purpose of this study is to assess C_{\min} for Pertuzumab at Day 43 in patients receiving two different Pertuzumab dose regimens. These data will then be analyzed using a population PK model

to identify a dose of Pertuzumab that will achieve a PK target steady-state trough concentration of $\geq 20 \mu\text{g/mL}$ in approximately 90% of the advanced gastric cancer patients. Analyses using the assumption that Pertuzumab behaves similarly to Trastuzumab in advanced gastric cancer suggest that with sample size of 15 patients per arm (total of 30 patients in this study), the dose to achieve the desired target concentration can be estimated with an acceptable degree of precision (coefficient of variation $< 15\%$).

Clinical Results

The clinical results of this phase IIa gastric cancer (GC) study are shown in Figures 32-37.

Figure 32 shows the samples taken and time points.

Figure 33 shows the demographics of the patient population in the two arms of the GC study, treated with 420 mg (Arm A) or 840 mg (Arm B) of Pertuzumab.

Figure 34 shows the GC history of the patients in Arms A and B, respectively.

Figure 35 shows the patient disposition in Arms A and B, respectively.

Figure 36 shows the Overall Response Rate in Arms A and B, respectively.

Safety Data

Diarrhea was the most common event occurring in 90% of subjects and was typically Grade 1 and 2 with onset in Cycle 1; no patient discontinued therapy because of diarrhea.

Grade ≥ 3 adverse events (AEs) ($>13\%$) included diarrhea, stomatitis, fatigue/asthenia, decreased appetite, hyponatremia, anemia, and neutopenia. With the exception of neutropenia and hyponatremia (higher in Arm A) and decreased appetite (higher in Arm B), incidence of these events was similar in the standard and high-dose Pertuzumab arms.

Asymptomatic change in ejection fraction (EF), neutropenic fever, rash, and drug hypersensitivity reaction were not associated with the higher dose of Pertuzumab.

Serious adverse events (SAEs) occurred in 60% of patients, and incidence was not associated with high-dose Pertuzumab.

Although more patients withdrew from treatment in Arm B, it is not clear that this was due to a higher Pertuzumab dose because events leading to treatment discontinuation were not uniform.

Pharmacokinetic (PK) Results

Figure 37 shows the results of Pertuzumab Day 42 concentration assessment in gastric cancer (GC) (JOSHUA) versus metastatic breast cancer (MBC) (CLEOPATRA).

Summary of the results

- Pertuzumab trough concentrations are lower in GC compared to MBC.
 - Between cycle concentrations (i.e. day 7, 14) are in-line with expected MBC concentrations as clearance is linear at these higher concentrations.
 - Trough levels with the 840/420 mg dose are about 37% lower compared to the

CLEOPATRA trial (Example 3), likely due to non-linear clearance at lower concentrations (incomplete receptor saturation).

- 840/840 mg dose in GC provides trough concentrations similar to 840/420 mg dose in MBC.
- Covariates have no impact on PK.

Conclusions

Based on Pertuzumab PK in GC, 840/840 mg dose will be used for the treatment of gastric cancer. This dose is expected to maintain trough levels above the target of $>20 \mu\text{g/mL}$ in 90% of patients, and provides similar trough levels as those observed in MBC.

EXAMPLE 2

Phase III Study Evaluating Pertuzumab in Combination with Trastuzumab and Chemotherapy in Patients with HER2-Positive Advanced Gastric Cancer

This is a Phase III, randomized, open-label, multicenter clinical study designed to assess the efficacy of Pertuzumab in combination with Trastuzumab and chemotherapy in patients with HER2-positive locally advanced or metastatic gastric cancer.

Patients in the treatment arm receive Trastuzumab, cisplatin, and capecitabine and/or 5-fluorouracil. In the other arm, patients are given either placebo or Pertuzumab.

Treatment regimens:

Pertuzumab:

840 mg dose for cycles 1-6.

Trastuzumab

8 mg/kg loading dose followed by 6 mg/kg q3w

Capecitabine

1000 mg/m^2 bid d1-14 q3w x 6

5-Fluorouracil

800 mg/m^2 day continuous iv infusion d1-5 q3w x 6

Cisplatin

800 mg/m^2 q3w x 6

Primary end point:

Overall survival (OS)

Secondary end points:

Progression-free survival (PFS), time to disease progression (TTP), objective response rate (ORR), Clinical Benefit Rate, Duration of Response, QoL, safety, pain intensity, analgesic consumption, weight change, pharmacokinetics.

Mail patient selection criteria

Inclusion criteria:

- Adenocarcinoma of stomach or GEJ
- Inoperable locally advanced and/or metastatic disease
- Measurable (RECIST), or non-measurable evaluable disease
- HER2-positive tumor: IHC 2+ or 3+ and/or ISH+
- Adequate organ function and ECOG performance status ≤ 2
- Written informed consent

Exclusion criteria

- Previous adjuvant chemotherapy within 6 months
- Chemotherapy for advanced disease
- Congestive heart failure or baseline LVEF $< 50\%$
- Creatinine clearance < 60 mL/min

It is expected that the treatment methods described herein, comprising the administration of Pertuzumab, Trastuzumab and chemotherapy(s), e.g. cisplatin and capecitabine, will meet the primary end point (OS). In particular, it is expected that the treatment methods herein will be therapeutically effective in the gastric cancer patients treated, for example, by extending survival, including overall survival (OS) and/or progression-free survival (PFS) and/or time to disease progression (TTP) and/or objective response rate (ORR) relative to treatment with Trastuzumab and chemotherapy only.

EXAMPLE 3

Results of a Phase III, Randomized, Double-Bind, Placebo-Controlled Registration Trial to Evaluate the Efficacy and Safety of Placebo + Trastuzumab + Docetaxel versus Pertuzumab + Trastuzumab + Docetaxel in Patients with Previously Untreated HER2-Positive Metastatic Breast Cancer (CLEOPATRA)

A protocol for evaluating Pertuzumab in HER2-positive metastatic breast cancer is found at <http://clinicaltrials.gov/ct2/show/NCT00567190> and in US 2009/0137387 as well as WO2009/154651.

This example concerns the clinical data obtained in the randomized, double-blind, placebo-controlled Phase III trial in patients with HER2-positive MBC, who had not received chemotherapy or biologic therapy for their metastatic disease. Patients were randomized 1:1 to receive placebo plus Trastuzumab plus Docetaxel or Pertuzumab plus Trastuzumab plus Docetaxel. The primary endpoint was progression-free survival (PFS), based on tumor assessments. PFS was defined as the time from randomization to the first documented radiographic progressive disease (PD) according to response evaluation criteria in solid tumors (RECIST) version 1.0 (Therasse et al. *J Natl Cancer Inst* 92:205-16 (2000)) or death from any cause, if within 18 weeks of the patient's last tumor assessment. Secondary endpoints included overall survival (OS), PFS by investigator assessment, objective response rate (ORR), and safety.

Patients: Eligible patients had centrally confirmed HER2-positive (defined as immunohistochemistry (IHC) 3+ and/or fluorescence *in situ* hybridization (FISH) amplification ratio ≥ 2.0) (Carlson et al. *J Natl Compr Canc Netw* 4 Suppl 3:S1-22 (2006)), locally recurrent, unresectable, or metastatic breast cancer, or *de novo* Stage IV disease. Patients were aged ≥ 18 years, had a left ventricular ejection fraction (LVEF) of $\geq 50\%$ at baseline (determined by echocardiogram or multiple gated acquisition), and an Eastern Cooperative Oncology Group performance status (ECOG PS) of 0 or 1. Patients may have received one hormonal treatment for MBC prior to randomization, or neoadjuvant or adjuvant systemic breast cancer therapy including Trastuzumab and/or taxanes, provided that they experienced a disease-free interval of ≥ 12 months between completion of neoadjuvant or adjuvant therapy and diagnosis of metastatic disease. Exclusion criteria included therapy for MBC (other than described above); central nervous system metastases; history of exposure to a cumulative dose of doxorubicin $>360 \text{ mg/m}^2$ or its equivalent; history of LVEF decline to $<50\%$ during or after prior Trastuzumab therapy; current uncontrolled hypertension; history of impaired cardiac function; impaired bone marrow, renal, or liver function; current known infection with HIV, HBV, or HCV; pregnancy; lactation; and refusal to use non-hormonal contraception.

Procedures: Patients received a loading dose of 8 mg/kg Trastuzumab, followed by a maintenance dose of 6 mg/kg every 3 weeks until investigator-assessed radiographic or clinical PD, or unmanageable toxicity. Docetaxel was administered every 3 weeks at a starting dose of 75 mg/m^2 , escalating to 100 mg/m^2 if tolerated. Per protocol, the investigator could reduce the dose by 25% to 55 mg/m^2 or 75 mg/m^2 (if the patient had been dose escalated) in order to manage tolerability. It was recommended that patients received at least 6 cycles of Docetaxel. Pertuzumab or placebo was given at a fixed loading dose of 840 mg, followed by 420 mg every 3 weeks until investigator-assessed radiographic or clinical PD, or unmanageable toxicity. In the case of chemotherapy discontinuation due to cumulative toxicity, antibody therapy was continued until PD, unacceptable toxicity, or withdrawal of consent. All drugs were administered intravenously.

Assessments: PFS was evaluated by standard RECIST-accepted methodology every 9 weeks by each center and by the IRF until IRF-assessed PD. Assessments of LVEF were performed at baseline, every 9 weeks during the treatment period, at treatment discontinuation, every 6 months in the first year after treatment discontinuation, then annually for up to 3 years in the follow-up period. Laboratory parameters and ECOG status were assessed at every cycle. Adverse events (AEs) were monitored continuously and graded according to NCI-CTCAE version 3.0. All cardiac events and serious adverse events (SAEs) that were ongoing at the time of treatment discontinuation were followed until resolution or stabilization up to 1 year after the final dose. Cardiac events and treatment-related SAEs with onset post-treatment

RESULTS

Study population: A total of 808 patients were enrolled and randomized to receive placebo plus Trastuzumab plus Docetaxel (n = 406) or Pertuzumab plus Trastuzumab plus Docetaxel (n = 402) (Figure 7). Baseline characteristics were similar between treatment arms (Table 1).

5 **Table 1: Baseline Characteristics of the Intent-to-Treat Population**

	Placebo + Trastuzumab + Docetaxel (n = 406)	Pertuzumab + Trastuzumab + Docetaxel (n = 402)
Sex, n (%)		
Female	404 (99.5)	402 (100.0)
Age, years		
Median	54.0	54.0
Range	27-89	22-82
Race, n (%)		
Asian	133 (32.8)	128 (31.8)
Black	20 (4.9)	10 (2.5)
White	235 (57.9)	245 (60.9)
Other*	18 (4.4)	19 (4.7)
Region, n (%)		
Asia	128 (31.5)	125 (31.1)
Europe	152 (37.4)	154 (38.3)
North America	68 (16.7)	67 (16.7)
South America	58 (14.3)	56 (13.9)
ECOG status, n (%)		
0	248 (61.1)	274 (68.2)
1	157 (38.7)	125 (31.1)
≥2	1 (0.2)	3 (0.7)
Prior treatment status, n (%)		
<i>De novo</i> MBC [†]	214 (52.7)	218 (54.2)
Prior adjuvant or neoadjuvant therapy	192 (47.3)	184 (45.8)
Prior Trastuzumab treatment, n (%)	41 (10.1)	47 (11.7)
Prior anthracycline treatment, n (%)	164 (40.4)	150 (37.3)
Prior taxane treatment, n (%)	94 (23.2)	91 (22.6)
Prior hormone treatment [†] , n (%)	107 (26.4)	114 (28.4)
Disease type at screening, n (%)		
Non-visceral	90 (22.2)	88 (21.9)
Visceral	316 (77.8)	314 (78.1)
Hormone receptor status, n (%)		
ER and/or PgR positive	199 (49.0)	189 (47.0)
ER and PgR negative	196 (48.3)	212 (52.7)
Unknown	11 (2.7)	1 (0.2)
HER2 status IHC, n (%)	405 (100)	401 (100)
0 and 1+	2 (0.5)	4 (1.0)
2+	32 (7.9)	47 (11.7)
3+	371 (91.6)	350 (87.3)
HER2 status FISH, n (%)	387 (100)	385 (100)
Positive	383 (99.0)	384 (99.7)
Negative	4 (1.0)	1 (0.3)

*Includes American Indian and Alaska Native

[†]No prior chemotherapy or biological therapy

‡In the neoadjuvant/adjuvant or metastatic setting

Progression-free survival: Treatment with Pertuzumab plus Trastuzumab plus Docetaxel significantly improved PFS-IRF, stratified by prior treatment status and region, compared with placebo plus Trastuzumab plus Docetaxel (HR = 0.62; 95% CI 0.51 to 0.75; $p < 0.0001$) (Figure 8). The median PFS-IVRF was prolonged by 6.1 months from 12.4 months with placebo plus Trastuzumab plus Docetaxel to 18.5 months with Pertuzumab plus Trastuzumab plus Docetaxel. The PFS benefit of Pertuzumab plus Trastuzumab plus Docetaxel treatment was observed across all predefined subgroups (Figure 9).

Assessment of PFS by investigators closely matched PFS-IRF. Median PFS assessed by investigators was 12.4 months with placebo plus Trastuzumab plus Docetaxel and 18.5 months with Pertuzumab plus Trastuzumab plus Docetaxel (HR = 0.65; 95% CI 0.54 to 0.78; $p < 0.0001$).

Key secondary efficacy endpoints: The interim analysis of OS took place when 43% of events ($n = 165$) that are planned for final OS analysis had occurred. More deaths occurred in the placebo plus Trastuzumab plus Docetaxel arm ($n = 96$; 23.6%) than in the Pertuzumab plus Trastuzumab plus Docetaxel arm ($n = 69$; 17.2%) (Figure 10). The HR (0.64; 95% CI 0.47 to 0.88; $p = 0.0053$) for OS did not meet the O'Brien-Fleming stopping boundary of the Lan-DeMets α -spending function for this interim analysis of survival ($HR \leq 0.603$, $p \leq 0.0012$), and therefore, was not statistically significant. However, the data showed a strong trend suggestive of a survival benefit in favor of Pertuzumab plus Trastuzumab plus Docetaxel. At the time of data cut-off, patients in both treatments arms had been followed for OS for a median of 19.3 months (Kaplan-Meier estimate). The ORR was 69.3% and 80.2% in the placebo plus Trastuzumab plus Docetaxel arm and Pertuzumab plus Trastuzumab plus Docetaxel arm, respectively. The difference in response rates between treatment arms was 10.8% (95% CI 4.2 to 17.5; $p = 0.0011$) (Table 2).

Table 2: Overall Response Rate

	Placebo + Trastuzumab + Docetaxel	Pertuzumab + Trastuzumab + Docetaxel
Patients with IRF-assessed measurable disease at baseline, n (%)	336 (100)	343 (100)
Objective response rate	233 (69.3)	275 (80.2)
Complete response rate	14 (4.2)	19 (5.5)
Partial response rate	219 (65.2)	256 (74.6)
Stable disease	70 (20.8)	50 (14.6)
Progressive disease	28 (8.3)	13 (3.8)
Unable to assess	2 (0.6)	2 (0.6)
No response assessment	3 (0.9)	3 (0.9)

IRF, independent review facility

Treatment exposure: The median number of cycles administered per patient was 15 and 18

with median time on treatment estimated to be 11.8 and 18.1 months for placebo plus Trastuzumab plus Docetaxel and for Pertuzumab plus Trastuzumab plus Docetaxel, respectively. Dose reductions were not permitted for placebo, Pertuzumab, or Trastuzumab. Patients received a median of eight cycles of Docetaxel in each arm. Based on the safety population, 61 (15.4%) patients in the placebo plus Trastuzumab plus Docetaxel arm received Docetaxel dose escalation to 100 mg/m² at any cycle compared with 48 (11.8%) patients in the Pertuzumab plus Trastuzumab plus Docetaxel arm. The median Docetaxel dose intensity was 24.8 mg/m²/week in the placebo plus Trastuzumab plus Docetaxel arm and 24.6 mg/m²/week in the Pertuzumab plus Trastuzumab plus Docetaxel arm. Reasons for permanent discontinuation of all study treatment are presented in Figure 7.

Tolerability and cardiac safety: The AE profile during the treatment period was generally balanced between treatment arms (Table 3). The incidence of the following AEs (all grades) was >5% higher with Pertuzumab plus Trastuzumab plus Docetaxel: diarrhea, rash, mucosal inflammation, febrile neutropenia, and dry skin.

Table 3: Adverse Events (All Grades) with ≥25% Incidence in Either Arm or ≥5% Difference Between Arms and Grade ≥3 Adverse Events with ≥2% Incidence in the Safety Population

	Placebo + Trastuzumab + Docetaxel (n = 397)	Pertuzumab + Trastuzumab + Docetaxel (n = 407)
Most common AEs (all grades), n (%)		
Diarrhea	184 (46.3)	272 (66.8)
Alopecia	240 (60.5)	248 (60.9)
Neutropenia	197 (49.6)	215 (52.8)
Nausea	165 (41.6)	172 (42.3)
Fatigue	146 (36.8)	153 (37.6)
Rash	96 (24.2)	137 (33.7)
Decreased appetite	105 (26.4)	119 (29.2)
Mucosal inflammation	79 (19.9)	113 (27.8)
Asthenia	120 (30.2)	106 (26.0)
Edema peripheral	119 (30.0)	94 (23.1)
Constipation	99 (24.9)	61 (15.0)
Febrile neutropenia	30 (7.6)	56 (13.8)
Dry skin	17 (4.3)	43 (10.6)
Grade ≥3 AEs with an incidence rate ≥2%, n (%)		
Neutropenia	182 (45.8)	199 (48.9)
Febrile neutropenia	30 (7.6)	56 (13.8)
Leukopenia	58 (14.6)	50 (12.3)
Diarrhea	20 (5.0)	32 (7.9)
Neuropathy peripheral	7 (1.8)	11 (2.7)
Anemia	14 (3.5)	10 (2.5)
Asthenia	6 (1.5)	10 (2.5)
Fatigue	13 (3.3)	9 (2.2)
Granulocytopenia	9 (2.3)	6 (1.5)
Left ventricular systolic dysfunction	11 (2.8)	5 (1.2)
Dyspnea	8 (2.0)	4 (1.0)

AE, adverse event

The incidence of the following grade ≥ 3 AEs was $>2\%$ higher with Pertuzumab plus Trastuzumab plus Docetaxel: neutropenia, febrile neutropenia, and diarrhea (Table 3). The incidence of grade ≥ 3 febrile neutropenia in patients from Asia was 12% in the placebo plus Trastuzumab plus Docetaxel arm and 26% in the Pertuzumab plus Trastuzumab plus Docetaxel arm; in all other geographical regions the incidence was $\leq 10\%$ in both arms.

LVSD (all grades) was reported more frequently in the placebo plus Trastuzumab plus Docetaxel arm compared with the Pertuzumab plus Trastuzumab plus Docetaxel arm (8.3% and 4.4%, respectively). Grade ≥ 3 LVSD was reported in 2.8% of patients receiving placebo plus Trastuzumab plus Docetaxel and in 1.2% of patients receiving Pertuzumab plus Trastuzumab plus Docetaxel. Among patients with a post-baseline LVEF assessment, LVEF declines of ≥ 10 percentage points from baseline to $<50\%$ at any stage during treatment were reported in 6.6% and 3.8% of patients in the placebo plus Trastuzumab plus Docetaxel arm and Pertuzumab plus Trastuzumab plus Docetaxel arm, respectively.

In the safety population, the majority of deaths in both treatment arms were attributed to PD (81 (20.4%) in the placebo arm, 57 (14.0%) in the Pertuzumab arm). Deaths due to causes other than PD were generally balanced and a similar number of patients died due to AEs (10 (2.5%) in the placebo arm, 8 (2.0%) in the Pertuzumab arm), with infections being the most common cause of death due to an AE.

DISCUSSION

These data show that the combination of the anti-HER2 monoclonal antibodies Pertuzumab and Trastuzumab with Docetaxel prolongs PFS in patients with HER2-positive MBC in the first-line setting. Treatment with Pertuzumab plus Trastuzumab plus Docetaxel exceeded expectations by resulting in a statistically significant reduction in PFS risk (HR = 0.62) and an improvement in median PFS of 6.1 months.

The combination was well tolerated and Pertuzumab did not increase rates of symptomatic or asymptomatic cardiac dysfunction. Before the data herein, it was expected that treatment with two HER2 antibodies would exacerbate cardiac toxicity. However, these data show this was not the case based on the tests herein for evaluating cardiac toxicity: incidence of symptomatic left ventricular systolic dysfunction (LVSD) including congestive heart failure (CHF), decrease in left ventricular ejection fraction (LVEF).

Pertuzumab-related AEs, including skin rash, mucosal inflammation, and dry skin, were mostly mild. There was an increased rate of grade ≥ 3 diarrhea and febrile neutropenia with Pertuzumab plus Trastuzumab plus Docetaxel treatment. The control arm in CLEOPATRA had a similar PFS to previous randomized studies that showed that the combination of Trastuzumab and Docetaxel in HER2-positive MBC had a median PFS of 11.7 months Marty et al. *J Clin Oncol*

23:4265-74 (2005).

Without being bound by any one theory, these data indicate that targeting HER2-positive tumors with two anti-HER2 monoclonal antibodies with complementary mechanisms of action results in a more comprehensive blockade of HER2 and highlight the clinical importance of preventing the ligand-dependent formation HER2 dimers to optimally silence HER2 signaling. This study has shown that combined HER2 blockade with Trastuzumab and Pertuzumab improves the outcome of patients with advanced HER2-positive disease in the first-line setting. These data are significant in that they support the first approved use of a HER2 dimerization inhibitor for therapy of HER2-positive cancer patients.

EXAMPLE 4
Article of Manufacture Including Pertuzumab

The phase III clinical data in Example 3 were used in the development of an article of manufacture comprising a vial (e.g. single-dose vial) with Pertuzumab therein and a package insert providing information about the safety and/or efficacy thereof, as well as a method of making an article of manufacture comprising packaging together Pertuzumab in a vial (e.g. single-dose vial) and a package insert with prescribing information regarding Pertuzumab on a package insert as herein below.

Pertuzumab is a sterile, clear to slightly opalescent, colorless to pale yellow liquid for IV infusion. Each single use vial contains 420 mg of Pertuzumab at a concentration of 30 mg/mL in 20 mM L-histidine acetate (pH 6.0), 120 mM sucrose and 0.02% polysorbate 20.

Pertuzumab is supplied in a single-dose vial containing preservative free liquid concentrate, at a concentration of 30 mg/mL ready for infusion. Each vial of Pertuzumab drug product contains a total of 420 mg Pertuzumab. Store vials in a refrigerator at 2°C to 8°C (36°F to 46°F) until time of use. Keep vial in the outer carton in order to protect from light.

FULL PRESCRIBING INFORMATION

WARNING: EMBRYO-FETAL TOXICITY

Exposure to PERTUZUMAB can result in embryo-fetal death and birth defects. Studies in animals have resulted in oligohydramnios, delayed renal development, and death. Advise patients of these risks and the need for effective contraception. (5.1, 8.1, 8.6)

1 INDICATIONS AND USAGE

Pertuzumab is indicated for use in combination with Trastuzumab and docetaxel for the treatment of patients with HER2-positive metastatic breast cancer who have not received prior anti-HER2 therapy or chemotherapy for metastatic disease.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Doses and Schedules

The initial dose of Pertuzumab is 840 mg administered as a 60-minute intravenous infusion, followed every 3 weeks thereafter by a dose of 420 mg administered as an intravenous infusion over 30 to 60 minutes. When administered with Pertuzumab, the recommended initial dose of Trastuzumab is 8 mg/kg administered as a 90-minute intravenous infusion, followed every 3 weeks thereafter by a dose of 6 mg/kg administered as an intravenous infusion over 30 to 90 minutes. When administered with Pertuzumab, the recommended initial dose of docetaxel is 75 mg/m² administered as an intravenous infusion. The dose may be escalated to 100 mg/m² administered every 3 weeks if the initial dose is well tolerated.

2.2 Dose Modification

For delayed or missed doses, if the time between two sequential infusions is less than 6 weeks, the 420 mg dose of Pertuzumab should be administered. Do not wait until the next planned dose. If the time between two sequential infusions is 6 weeks or more, the initial dose of 840 mg Pertuzumab should be re-administered as a 60-minute intravenous infusion followed every 3 weeks thereafter by a dose of 420 mg administered as an intravenous infusion over 30 to 60 minutes. The infusion rate of Pertuzumab may be slowed or interrupted if the patient develops an infusion-associated reaction. The infusion should be discontinued immediately if the patient experiences a serious hypersensitivity reaction [see Warnings and Precautions (5.2)].

20 Left Ventricular Ejection Fraction (LVEF):

Withhold Pertuzumab and Trastuzumab dosing for at least 3 weeks for either:

- a drop in LVEF to less than 40% or
- LVEF of 40% to 45% with a 10% or greater absolute decrease below pretreatment values [see Warnings and Precautions (5.2)]

25

Pertuzumab may be resumed if the LVEF has recovered to greater than 45% or to 40% to 45% associated with less than a 10% absolute decrease below pretreatment values.

30

If after a repeat assessment within approximately 3 weeks, the LVEF has not improved, or has declined further, discontinuation of Pertuzumab and Trastuzumab should be strongly considered, unless the benefits for the individual patient are deemed to outweigh the risks [see Warnings and Precautions (5.2)]. Pertuzumab should be withheld or discontinued if Trastuzumab treatment is withheld or discontinued. If docetaxel is discontinued, treatment with Pertuzumab

and Trastuzumab may continue. Dose reductions are not recommended for Pertuzumab. For docetaxel dose modifications, see docetaxel prescribing information.

2.3 Preparation for Administration

Administer as an intravenous infusion only. Do not administer as an intravenous push or bolus. Do not mix Pertuzumab with other drugs.

Preparation: Prepare the solution for infusion, using aseptic technique, as follows:

- Parenteral drug products should be inspected visually for particulates and discoloration prior to administration.
- Withdraw the appropriate volume of Pertuzumab solution from the vial(s).
- 10 • Dilute into a 250 mL 0.9% sodium chloride PVC or non-PVC polyolefin infusion bag.
- Mix diluted solution by gentle inversion. Do not shake.
- Administer immediately once prepared.
- If the diluted infusion solution is not used immediately, it can be stored at 2oC to 8oC for up to 24 hours.
- 15 • Dilute with 0.9% Sodium Chloride injection only. Do not use dextrose (5%) solution.

3 Dosage Forms and Strengths

Pertuzumab 420 mg/14 mL (30 mg/mL) in a single-use vial

4 Contraindications

None

20 5 Warnings and Precautions

5.1 Embryo-Fetal Toxicity

Pertuzumab can cause fetal harm when administered to a pregnant woman. Treatment of pregnant cynomolgus monkeys with Pertuzumab resulted in oligohydramnios, delayed fetal kidney development, and embryo-fetal death. If Pertuzumab is administered during pregnancy, or if the patient becomes pregnant while receiving this drug, the patient should be apprised of the potential hazard to a fetus [see *Use in Specific Populations* (8.1)]. Verify pregnancy status prior to the initiation of Pertuzumab. Advise patients of the risks of embryo-fetal death and birth defects and the need for contraception during and after treatment. Advise patients to contact their healthcare provider immediately if they suspect they may be pregnant. If Pertuzumab is administered during pregnancy or if a patient becomes pregnant while receiving Pertuzumab, immediately report exposure

to the Genentech Adverse Event Line at 1-888-835-2555. Encourage women who may be exposed during pregnancy to enroll in the MoTHER Pregnancy Registry by contacting 1-800-690-6720 [see *Patient Counseling Information (17)*]. Monitor patients who become pregnant during Pertuzumab therapy for oligohydramnios. If oligohydramnios occurs, perform fetal testing that is appropriate for gestational age and consistent with community standards of care. The efficacy of intravenous hydration in the management of oligohydramnios due to Pertuzumab exposure is not known.

5.2 Left Ventricular Dysfunction

Decreases in LVEF have been reported with drugs that block HER2 activity, including Pertuzumab. In the randomized trial, Pertuzumab in combination with Trastuzumab and docetaxel was not associated with increases in the incidence of symptomatic left ventricular systolic dysfunction (LVSD) or decreases in LVEF compared with placebo in combination with Trastuzumab and docetaxel [see *Clinical Studies (14.1)*]. Left ventricular dysfunction occurred in 4.4% of patients in the Pertuzumab-treated group and 8.3% of patients in the placebo-treated group. Symptomatic left ventricular systolic dysfunction (congestive heart failure) occurred in 1.0% of patients in the Pertuzumab-treated group and 1.8% of patients in the placebo-treated group [see *Adverse Reactions (6.1)*]. Patients who have received prior anthracyclines or prior radiotherapy to the chest area may be at higher risk of decreased LVEF. Pertuzumab has not been studied in patients with a pretreatment LVEF value of $\leq 50\%$, a prior history of CHF, decreases in LVEF to $< 50\%$ during prior Trastuzumab therapy, or conditions that could impair left ventricular function such as uncontrolled hypertension, recent myocardial infarction, serious cardiac arrhythmia requiring treatment or a cumulative prior anthracycline exposure to $> 360 \text{ mg/m}^2$ of doxorubicin or its equivalent. Assess LVEF prior to initiation of Pertuzumab and at regular intervals (e.g., every three months) during treatment to ensure that LVEF is within the institution's normal limits. If LVEF is $< 40\%$, or is 40% to 45% with a 10% or greater absolute decrease below the pretreatment value, withhold Pertuzumab and Trastuzumab and repeat LVEF assessment within approximately 3 weeks. Discontinue Pertuzumab and Trastuzumab if the LVEF has not improved or has declined further, unless the benefits for the individual patient outweigh the risks [see *Dosage and Administration (2.2)*].

5.3 Infusion-Associated Reactions, Hypersensitivity Reactions/Anaphylaxis

Pertuzumab has been associated with infusion and hypersensitivity reactions [see *Adverse Reactions (6.1)*]. An infusion reaction was defined in the randomized trial as any event described as hypersensitivity, anaphylactic reaction, acute infusion reaction or cytokine release syndrome occurring during an infusion or on the same day as the infusion. The initial dose of Pertuzumab was

given the day before Trastuzumab and docetaxel to allow for the examination of Pertuzumab-associated reactions. On the first day, when only Pertuzumab was administered, the overall frequency of infusion reactions was 13.0% in the Pertuzumab-treated group and 9.8% in the placebo-treated group. Less than 1% were grade 3 or 4. The most common infusion reactions ($\geq 1.0\%$) were pyrexia, chills, fatigue, headache, asthenia, hypersensitivity, and vomiting. During the second cycle when all drugs were administered on the same day, the most common infusion reactions in the Pertuzumab-treated group ($\geq 1.0\%$) were fatigue, dysgeusia, hypersensitivity, myalgia, and vomiting. In the randomized trial, the overall frequency of hypersensitivity/anaphylaxis reactions was 10.8% in the Pertuzumab-treated group and 9.1% in the placebo-treated group. The incidence of Grade 3 – 4 hypersensitivity/anaphylaxis reactions was 2% in the Pertuzumab-treated group and 2.5% in the placebo-treated group according to National Cancer Institute – Common Terminology Criteria for Adverse Events (NCI - CTCAE) (version 3). Overall, 4 patients in Pertuzumab-treated group and 2 patients in the placebo-treated group experienced anaphylaxis. Observe patients closely for 60 minutes after the first infusion and for 30 minutes after subsequent infusions of Pertuzumab. If a significant infusion-associated reaction occurs, slow or interrupt the infusion and administer appropriate medical therapies. Monitor patients carefully until complete resolution of signs and symptoms. Consider permanent discontinuation in patients with severe infusion reactions [*see Dosage and Administration (2.2)*].

5.4 HER2 Testing

Detection of HER2 protein overexpression is necessary for selection of patients appropriate for Pertuzumab therapy because these are the only patients studied and for whom benefit has been shown [*see Indications and Usage (1) and Clinical Studies (14)*]. In the randomized trial, patients with breast cancer were required to have evidence of HER2 overexpression defined as 3+ IHC by Dako HERCEPTEST[®] or FISH amplification ratio ≥ 2.0 by Dako HER2 FISH PHARMDX[™] test kit. Only limited data were available for patients whose breast cancer was positive by FISH, but did not demonstrate protein overexpression by IHC. Assessment of HER2 status should be performed by laboratories with demonstrated proficiency in the specific technology being utilized. Improper assay performance, including use of sub-optimally fixed tissue, failure to utilize specified reagents, deviation from specific assay instructions, and failure to include appropriate controls for assay validation, can lead to unreliable results.

6 Adverse Reactions

The following adverse reactions are discussed in greater detail in other sections of the label:

- Embryo-Fetal Toxicity [*see Warnings and Precautions (5.1)*]
- Left Ventricular Dysfunction [*see Warnings and Precautions (5.2)*]
- Infusion-Associated Reactions, Hypersensitivity Reactions/Anaphylaxis [*see Warnings and Precautions (5.3)*]

5 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice. In clinical trials, Pertuzumab has been evaluated in more than 1400 patients with various malignancies and treatment with

10 Pertuzumab was predominantly in combination with other anti-neoplastic agents.

The adverse reactions described in Table 4 were identified in 804 patients with HER2-positive metastatic breast cancer treated in the randomized trial. Patients were randomized to receive either Pertuzumab in combination with Trastuzumab and docetaxel or placebo in combination with Trastuzumab and docetaxel. The median duration of study treatment was 18.1 months for patients in

15 the Pertuzumab-treated group and 11.8 months for patients in the placebo-treated group. No dose adjustment was permitted for Pertuzumab or Trastuzumab. The rates of adverse events resulting in permanent discontinuation of all study therapy were 6.1% for patients in the Pertuzumab-treated group and 5.3% for patients in the placebo-treated group. Adverse events led to discontinuation of docetaxel alone in 23.6% of patients in the Pertuzumab-treated group and 23.2% of patients in the

20 placebo-treated group. Table 4 reports the adverse reactions that occurred in at least 10% of patients in the Pertuzumab-treated group. The most common adverse reactions (> 30%) seen with Pertuzumab in combination with Trastuzumab and docetaxel were diarrhea, alopecia, neutropenia, nausea, fatigue, rash, and peripheral neuropathy.

The most common NCI - CTCAE (version 3) Grade 3 – 4 adverse reactions (> 2%) were

25 neutropenia, febrile neutropenia, leukopenia, diarrhea, peripheral neuropathy, anemia, asthenia, and fatigue. An increased incidence of febrile neutropenia was observed for Asian patients in both treatment arms compared with patients of other races and from other geographic regions. Among Asian patients, the incidence of febrile neutropenia was higher in the Pertuzumab-treated group (26%) compared with the placebo-treated group (12%).

30

Table 4: Summary of Adverse Reactions Occurring in $\geq 10\%$ of Patients on the Pertuzumab Treatment Arm in the Randomized Trial

Body System/Adverse Reactions	Pertuzumab + Trastuzumab + docetaxel n=407		Placebo + Trastuzumab + docetaxel n=397	
	Frequency rate % All Grades %	Grades 3 – 4 %	Frequency rate % All Grades %	Grades 3 – 4 %
General disorders and administration site conditions				
Fatigue	37.6	2.2	36.8	3.3
Asthenia	26.0	2.5	30.2	1.5
Edema peripheral	23.1	0.5	30.0	0.8
Mucosal inflammation	27.8	1.5	19.9	1.0
Pyrexia	18.7	1.2	17.9	0.5
Skin and subcutaneous tissue disorders				
Alopecia	60.9	0.0	60.5	0.3
Rash	33.7	0.7	24.2	0.8
Nail disorder	22.9	1.2	22.9	0.3
Pruritus	14.0	0.0	10.1	0.0
Dry skin	10.6	0.0	4.3	0.0
Gastrointestinal disorders				
Diarrhea	66.8	7.9	46.3	5.0
Nausea	42.3	1.2	41.6	0.5
Vomiting	24.1	1.5	23.9	1.5
Constipation	15.0	0.0	24.9	1.0
Stomatitis	18.9	0.5	15.4	0.3
Blood and lymphatic system disorders				
Neutropenia	52.8	48.9	49.6	45.8
Anemia	23.1	2.5	18.9	3.5
Leukopenia	18.2	12.3	20.4	14.6
Febrile neutropenia*	13.8	13.0	7.6	7.3
Nervous system disorders				
Neuropathy peripheral	32.4	3.2	33.8	2.0
Headache	20.9	1.2	16.9	0.5
Dysgeusia	18.4	0.0	15.6	0.0
Dizziness	12.5	0.5	12.1	0.0
Musculoskeletal and connective tissue disorders				
Myalgia	22.9	1.0	23.9	0.8
Arthralgia	15.5	0.2	16.1	0.8
Infections and infestations				
Upper respiratory tract infection	16.7	0.7	13.4	0.0
Nasopharyngitis	11.8	0.0	12.8	0.3
Respiratory, thoracic and mediastinal disorders				
Dyspnea	14.0	1.0	15.6	2.0
Metabolism and nutrition disorders				
Decreased appetite	29.2	1.7	26.4	1.5
Eye disorders				

Lacrimation increased	14.0	0.0	13.9	0.0
Psychiatric disorders				
Insomnia	13.3	0.0	13.4	0.0

* In this table this denotes an adverse reaction that has been reported in association with a fatal outcome

The following clinically relevant adverse reactions were reported in < 10% of patients in the Pertuzumab-treated group:

- 5 **Skin and subcutaneous tissue disorders:** Paronychia (7.1% in the Pertuzumab-treated group vs. 3.5% in the placebo-treated group); **Respiratory, thoracic and mediastinal disorders:** Pleural effusion (5.2% in the Pertuzumab-treated group vs. 5.8% in the placebo-treated group); **Cardiac disorders:** Left ventricular dysfunction (4.4% in the Pertuzumab-treated group vs. 8.3% in the placebo-treated group) including symptomatic left ventricular systolic dysfunction (CHF) (1.0% in the Pertuzumab-treated group vs. 1.8% in the placebo-treated group); **Immune system disorders:** Hypersensitivity (10.1% in the Pertuzumab-treated group vs. 8.6% in placebo-treated group).

Adverse Reactions Reported in Patients Receiving Pertuzumab and Trastuzumab after Discontinuation of Docetaxel

- 15 In the randomized trial, adverse reactions were reported less frequently after discontinuation of docetaxel treatment. All adverse reactions in the Pertuzumab and Trastuzumab treatment group occurred in < 10% of patients with the exception of diarrhea (19.1%), upper respiratory tract infection (12.8%), rash (11.7%), headache (11.4%), and fatigue (11.1%).

6.2 Immunogenicity

- As with all therapeutic proteins, there is the potential for an immune response to Pertuzumab. Patients in the randomized trial were tested at multiple time-points for antibodies to Pertuzumab. Approximately 2.8% (11/386) of patients in the Pertuzumab-treated group and 6.2% (23/372) of patients in the placebo-treated group tested positive for anti-Pertuzumab antibodies. Of these 34 patients, none experienced anaphylactic/hypersensitivity reactions that were clearly related to the anti-therapeutic antibodies (ATA). The presence of Pertuzumab in patient serum at the levels expected at the time of ATA sampling can interfere with the ability of this assay to detect anti-Pertuzumab antibodies. In addition, the assay may be detecting antibodies to Trastuzumab. As a result, data may not accurately reflect the true incidence of anti-Pertuzumab antibody development. Immunogenicity data are highly dependent on the sensitivity and specificity of the test methods used. Additionally, the observed incidence of a positive result in a test method may be influenced by several factors, including sample handling, timing of sample collection, drug interference,

concomitant medication, and the underlying disease. For these reasons, comparison of the incidence of antibodies to Pertuzumab with the incidence of antibodies to other products may be misleading.

7 Drug Interactions

No drug-drug interactions were observed between Pertuzumab and Trastuzumab, or between
5 Pertuzumab and docetaxel.

8 Use in Specific Populations

8.1 Pregnancy

Pregnancy Category D

Risk Summary

10 There are no adequate and well-controlled studies of Pertuzumab in pregnant women. Based
on findings in animal studies, Pertuzumab can cause fetal harm when administered to a pregnant
woman. The effects of Pertuzumab are likely to be present during all trimesters of pregnancy.
Pertuzumab administered to pregnant cynomolgus monkeys resulted in oligohydramnios, delayed
fetal kidney development, and embryo-fetal deaths at clinically relevant exposures of 2.5 to 20-fold
15 greater than the recommended human dose, based on C_{max} . If Pertuzumab is administered during
pregnancy, or if a patient becomes pregnant while receiving Pertuzumab, the patient should be
apprised of the potential hazard to the fetus. If Pertuzumab is administered during pregnancy or if a
patient becomes pregnant while receiving Pertuzumab, immediately report exposure to the Genentech
Adverse Event Line at 1-888-835-2555. Encourage women who may be exposed during pregnancy
20 to enroll in the MoTHER Pregnancy Registry by contacting 1-800-690-6720 [*see Patient Counseling
Information (17)*].

Animal Data

Reproductive toxicology studies have been conducted in cynomolgus monkeys. Pregnant
monkeys were treated on Gestational Day (GD)19 with loading doses of 30 to 150 mg/kg
25 Pertuzumab, followed by bi-weekly doses of 10 to 100 mg/kg. These dose levels resulted in
clinically relevant exposures of 2.5 to 20-fold greater than the recommended human dose, based on
 C_{max} . Intravenous administration of Pertuzumab from GD19 through GD50 (period of
organogenesis) was embryotoxic, with dose-dependent increases in embryo-fetal death between
GD25 to GD70. The incidences of embryo-fetal loss were 33, 50, and 85% for dams treated with
30 bi-weekly Pertuzumab doses of 10, 30, and 100 mg/kg, respectively (2.5 to 20-fold greater than the
recommended human dose, based on C_{max}). At Cesarean section on GD100, oligohydramnios,

decreased relative lung and kidney weights and microscopic evidence of renal hypoplasia consistent with delayed renal development were identified in all Pertuzumab dose groups. Pertuzumab exposure was reported in offspring from all treated groups, at levels of 29% to 40% of maternal serum levels at GD100.

5 **8.3 Nursing Mothers**

It is not known whether Pertuzumab is excreted in human milk, but human IgG is excreted in human milk. Because many drugs are secreted in human milk and because of the potential for serious adverse reactions in nursing infants from Pertuzumab, a decision should be made whether to discontinue nursing, or discontinue drug, taking into account the elimination half-life of Pertuzumab and the importance of the drug to the mother [*See Warnings and Precautions (5.1), Clinical Pharmacology (12.3)*].

8.4 Pediatric Use

The safety and effectiveness of Pertuzumab have not been established in pediatric patients.

8.5 Geriatric Use

15 Of 402 patients who received Pertuzumab in the randomized trial, 60 patients (15%) were ≥ 65 years of age and 5 patients (1%) were ≥ 75 years of age. No overall differences in efficacy and safety of Pertuzumab were observed between these patients and younger patients. Based on a population pharmacokinetic analysis, no significant difference was observed in the pharmacokinetics of Pertuzumab between patients < 65 years ($n=306$) and patients ≥ 65 years ($n=175$).

20 **8.6 Females of Reproductive Potential**

Pertuzumab can cause embryo-fetal harm when administered during pregnancy. Counsel patients regarding pregnancy prevention and planning. Advise females of reproductive potential to use effective contraception while receiving Pertuzumab and for 6 months following the last dose of Pertuzumab. If Pertuzumab is administered during pregnancy or if a patient becomes pregnant while receiving Pertuzumab, immediately report exposure to the Genentech Adverse Event Line at 1-888-835-2555. Encourage women who may be exposed during pregnancy to enroll in the MoHER Pregnancy Registry by contacting 1-800-690-6720 [*see Patient Counseling Information (17)*].

8.7 Renal Impairment

Dose adjustments of Pertuzumab are not needed in patients with mild (creatinine clearance [CLcr] 60 to 90 mL/min) or moderate (CLcr 30 to 60 mL/min) renal impairment. No dose adjustment can be recommended for patients with severe renal impairment (CLcr less than 30 mL/min) because of the limited pharmacokinetic data available [see *Clinical Pharmacology* (12.3)].

8.8 Hepatic Impairment

No clinical studies have been conducted to evaluate the effect of hepatic impairment on the pharmacokinetics of Pertuzumab.

10 10 OVERDOSAGE

No drug overdoses have been reported with Pertuzumab to date.

11 Description

Pertuzumab is a recombinant humanized monoclonal antibody that targets the extracellular dimerization domain (Subdomain II) of the human epidermal growth factor receptor 2 protein (HER2). Pertuzumab is produced by recombinant DNA technology in a mammalian cell (Chinese Hamster Ovary) culture containing the antibiotic, gentamicin. Gentamicin is not detectable in the final product. Pertuzumab has an approximate molecular weight of 148 kDa. Pertuzumab is a sterile, clear to slightly opalescent, colorless to pale brown liquid for intravenous infusion. Each single use vial contains 420 mg of Pertuzumab at a concentration of 30 mg/mL in 20 mM L-histidine acetate (pH 6.0), 120 mM sucrose and 0.02% polysorbate 20.

12 Clinical Pharmacology

12.1 Mechanism of Action

Pertuzumab targets the extracellular dimerization domain (Subdomain II) of the human epidermal growth factor receptor 2 protein (HER2) and, thereby, blocks ligand-dependent heterodimerization of HER2 with other HER family members, including EGFR, HER3 and HER4. As a result, Pertuzumab inhibits ligand-initiated intracellular signaling through two major signal pathways, mitogen-activated protein (MAP) kinase and phosphoinositide 3-kinase (PI3K). Inhibition of these signaling pathways can result in cell growth arrest and apoptosis, respectively. In addition,

Pertuzumab mediates antibody-dependent cell-mediated cytotoxicity (ADCC). While Pertuzumab alone inhibited the proliferation of human tumor cells, the combination of Pertuzumab and Trastuzumab significantly augmented anti-tumor activity in HER2-overexpressing xenograft models.

12.2 Pharmacokinetics

5 Pertuzumab demonstrated linear pharmacokinetics at a dose range of 2 – 25 mg/kg. Based on a population PK analysis that included 481 patients, the median clearance (CL) of Pertuzumab was 0.24 L/day and the median half-life was 18 days. With an initial dose of 840 mg followed by a maintenance dose of 420 mg every three weeks thereafter, the steady-state concentration of Pertuzumab was reached after the first maintenance dose. The population PK analysis suggested no
10 PK differences based on age, gender, and ethnicity (Japanese vs. non-Japanese). Baseline serum albumin level and lean body weight as covariates only exerted a minor influence on PK parameters. Therefore, no dose adjustments based on body weight or baseline albumin level are needed. No drug-drug interactions were observed between Pertuzumab and Trastuzumab, or between Pertuzumab and docetaxel in a sub-study of 37 patients in the randomized trial. No dedicated renal impairment trial
15 for Pertuzumab has been conducted. Based on the results of the population pharmacokinetic analysis, Pertuzumab exposure in patients with mild (CLcr 60 to 90 mL/min, n=200) and moderate renal impairment (CLcr 30 to 60 mL/min, n=71) were similar to those in patients with normal renal function (CLcr greater than 90 mL/min, n=200). No relationship between CLcr and Pertuzumab exposure was observed over the range of observed CLcr (27 to 244 mL/min).

20 12.3 Cardiac Electrophysiology

The effect of Pertuzumab with an initial dose of 840 mg followed by a maintenance dose of 420 mg every three weeks on QTc interval was evaluated in a subgroup of 20 patients with HER2-positive breast cancer in the randomized trial. No large changes in the mean QT interval (i.e., greater than 20 ms) from placebo based on Fridericia correction method were detected in the trial. A small
25 increase in the mean QTc interval (i.e., less than 10 ms) cannot be excluded because of the limitations of the trial design.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have not been performed to evaluate the carcinogenic potential
30 of Pertuzumab. Studies have not been performed to evaluate the mutagenic potential of Pertuzumab. No specific fertility studies in animals have been performed to evaluate the effect of Pertuzumab. No

adverse effects on male and female reproductive organs were observed in repeat-dose toxicity studies of up to six months duration in cynomolgus monkeys.

14 Clinical Studies

14.1 Metastatic Breast Cancer

5 The randomized trial was a multicenter, double-blind, placebo-controlled trial of 808 patients with HER2-positive metastatic breast cancer. Breast tumor specimens were required to show HER2 overexpression defined as 3+ IHC or FISH amplification ratio ≥ 2.0 determined at a central laboratory. Patients were randomized 1:1 to receive placebo plus Trastuzumab and docetaxel or Pertuzumab plus Trastuzumab and docetaxel. Randomization was stratified by prior treatment (prior
10 or no prior adjuvant/neoadjuvant anti-HER2 therapy or chemotherapy) and geographic region (Europe, North America, South America, and Asia). Patients with prior adjuvant or neoadjuvant therapy were required to have a disease-free interval of greater than 12 months before trial enrollment. Pertuzumab was given intravenously at an initial dose of 840 mg, followed by 420 mg every 3 weeks thereafter. Trastuzumab was given intravenously at an initial dose of 8 mg/kg,
15 followed by 6 mg/kg every 3 weeks thereafter. Patients were treated with Pertuzumab and Trastuzumab until progression of disease, withdrawal of consent, or unacceptable toxicity. Docetaxel was given as an initial dose of 75 mg/m² by intravenous infusion every 3 weeks for at least 6 cycles. The docetaxel dose could be escalated to 100 mg/m² at the investigator's discretion if the initial dose was well tolerated.

20 At the time of the primary analysis, the mean number of cycles of study treatment administered was 16.2 in the placebo-treated group and 19.9 in the Pertuzumab-treated group.

 The primary endpoint of the randomized trial was progression-free survival (PFS) as assessed by an independent review facility (IRF). PFS was defined as the time from the date of randomization to the date of disease progression or death (from any cause) if the death occurred within 18 weeks of
25 the last tumor assessment. Additional endpoints included overall survival (OS), PFS (investigator-assessed), objective response rate (ORR) and duration of response.

 Patient demographic and baseline characteristics were balanced between the treatment arms. The median age was 54 (range 22 to 89 years), 59% were White, 32% were Asian, and 4% were Black. All were women with the exception of 2 patients. Seventeen percent of patients were
30 enrolled in North America, 14% in South America, 38% in Europe, and 31% in Asia. Tumor prognostic characteristics, including hormone receptor status (positive 48%, negative 50%), presence of visceral disease (78%) and non-visceral disease only (22%) were similar in the study arms. Approximately half of the patients received prior adjuvant or neoadjuvant anti-HER2 therapy or chemotherapy (placebo 47%, Pertuzumab 46%). Among patients with hormone receptor positive

tumors, 45% received prior adjuvant hormonal therapy and 11% received hormonal therapy for metastatic disease. Eleven percent of patients received prior adjuvant or neoadjuvant Trastuzumab.

The randomized trial demonstrated a statistically significant improvement in IRF-assessed PFS in the Pertuzumab-treated group compared with the placebo-treated group [hazard ratio (HR) = 0.62 (95% CI: 0.51, 0.75), $p < 0.0001$] and an increase in median PFS of 6.1 months (median PFS of 18.5 months in the Pertuzumab-treated group vs. 12.4 months in the placebo-treated group) (see Figure 8). The results for investigator-assessed PFS were comparable to those observed for IRF-assessed PFS. Consistent results were observed across several patient subgroups including age (< 65 or ≥ 65 years), race, geographic region, prior adjuvant/neoadjuvant anti-HER2 therapy or chemotherapy (yes or no), and prior adjuvant/neoadjuvant Trastuzumab (yes or no). In the subgroup of patients with hormone receptor-negative disease ($n=408$), the hazard ratio was 0.55 (95% CI: 0.42, 0.72). In the subgroup of patients with hormone receptor-positive disease ($n=388$), the hazard ratio was 0.72 (95% CI: 0.55, 0.95). In the subgroup of patients with disease limited to non-visceral metastasis ($n=178$), the hazard ratio was 0.96 (95% CI: 0.61, 1.52).

At the time of the PFS analysis, 165 patients had died. More deaths occurred in the placebo-treated group (23.6%) compared with the Pertuzumab-treated group (17.2%). At the interim OS analysis, the results were not mature and did not meet the pre-specified stopping boundary for statistical significance. See Table 5 and Figure 10.

Table 5: Summary of Efficacy from the Randomized Trial

Parameter	Pertuzumab + Trastuzumab + docetaxel n=402	Placebo + Trastuzumab + docetaxel n=406	HR (95% CI)	p-value
Progression-Free Survival (independent review)				
No. of patients with an event	191 (47.5%)	242 (59.6%)	0.62 (0.51, 0.75)	< 0.0001
Median months	18.5	12.4		
Overall Survival (interim analysis)				
No. of patients with an event	69 (17.2%)	96 (23.6%)	0.64 (0.47, 0.88)	0.0053*
Objective Response Rate (ORR)				
No. of patients analyzed				
Objective response (CR + PR)	343	336		
Complete response (CR)	275 (80.2%)	233 (69.3%)		
Partial Response (PR)	19 (5.5%)	14 (4.2%)		
	256 (74.6%)	219 (65.2%)		
Median Duration of Response (months)	20.2	12.5		

* The HR and p-value for the interim analysis of Overall Survival did not meet the pre-defined stopping boundary ($HR \leq 0.603$, $p \leq 0.0012$).

16 How Supplied/Storage and Handling**16.1 How Supplied**

Pertuzumab is supplied as a 420 mg/14 mL (30 mg/mL) single-use vial containing preservative-free solution. NDC 50242-145-01. Store vials in a refrigerator at 2°C to 8°C

5 (36°F to 46°F) until time of use. Keep vial in the outer carton in order to protect from light.

DO NOT FREEZE. DO NOT SHAKE.

17 Patient Counseling Information

Advise pregnant women and females of reproductive potential that Pertuzumab exposure can result in fetal harm, including embryo-fetal death or birth defects [*see Warnings and Precautions*

10 (5.1) and Use in Specific Populations (8.1)]

Advise females of reproductive potential to use effective contraception while receiving Pertuzumab and for 6 months following the last dose of Pertuzumab [*see Warnings and Precautions* (5.1) and Use in Special Populations (8.6)]

Advise nursing mothers treated with Pertuzumab to discontinue nursing or discontinue Pertuzumab, taking into account the importance of the drug to the mother [*see Use in Specific*

15 Populations (8.3)].

Encourage women who are exposed to Pertuzumab during pregnancy to enroll in the MoTHER Pregnancy Registry by contacting 1-800-690-6720 [*see Warnings and Precautions* (5.1) and Use in Specific Populations (8.1)]

20 Thus, the comprehensive phase III safety and efficacy data for Pertuzumab as in Example 3 provide for the article of manufacture in this example. This article of manufacture can be used in a method of ensuring safe and effective use of Pertuzumab to treat patients.

EXAMPLE 5**Early-Stage Breast Cancer Therapy with Pertuzumab**

25 Anthracyclines (generally used in combination with 5-FU and cyclophosphamide) have a central role in the management of breast cancer. Romond et al. *NEJM* 353(16): 1673-1684 (2005), and Poole et al. *NEJM* 355 (18): 1851-1852 (2006).

Taxanes are also integral in standard regimens for the treatment of breast cancer, used in combination with anthracyclines in a regimen known as TAC (Martin et al. *NEJM* 352 (22): 2302-

2313 (2005)) or in sequence with anthracyclines in a regimen known as AC->T (Romond et al., *supra*; Joensuu et al. *NEJM* 354 (8): 809-820 (2006)).

Carboplatin is both an active and well tolerated chemotherapy agent and there are studies in breast cancer which show clear efficacy in combination with a taxane and Trastuzumab in a regimen known as TCH (Slamon et al. *BCIRG 006*. SABS (2007); Robert et al. *J. Clin. Oncol.* 24: 2786-2792 (2006)). However, in metastatic breast cancer, there are negative data (Forbes et al. *BCIRG 007 Proc. Am. Soc. Clin. Oncol.* Abstract No. LBA516 (2006)).

A previous neoadjuvant study with Pertuzumab (NeoSphere) evaluated it in combination with Docetaxel and Trastuzumab (Gianni et al. *Cancer Research* 70 (24) (Suppl. 2) (December 2010)), but not in combination with anthracycline-based or carboplatin-based chemotherapy.

The following chemotherapy regimens were evaluated in this example:

FEC	Breast cancer therapy consisting of 5- <u>f</u> luorouracil, <u>e</u> pirubicin and <u>c</u> yclophosphamide.
FEC ->T	Sequential chemotherapy, consisting of courses of FEC chemotherapy followed by courses of Docetaxel.
TCH	Chemotherapy regimen for HER2-positive breast cancer combination comprising taxane (Docetaxel), Carboplatin, and Trastuzumab (HERCEPTIN®)

The treatment arms in this study were:

Arm A

Courses of 5-Fluorouracil, Epirubicin and Cyclophosphamide (FEC) followed by courses of Docetaxel (T) (FEC ->T) with Trastuzumab and Pertuzumab given from the start of the chemotherapy regimen (i.e. concurrently with the anthracycline)

5-Fluorouracil (500 mg/m²), epirubicin (100 mg/m²) followed by cyclophosphamide (600 mg/m²) for three cycles, followed by Docetaxel for three cycles with Trastuzumab (8 mg/kg on day 1 of the first treatment with epirubicin and 6 mg/kg every 3 weeks thereafter) and Pertuzumab (840 mg on day 1 of the treatment with FEC with 420 mg every 3 weeks thereafter). The starting dose for Docetaxel is 75 mg/m² for Cycle 4 (first Docetaxel cycle) then 100 mg/m² for Cycles 5-6, if no dose limiting toxicity occurs. All drugs will be administered by the IV route.

OR

Arm B

FEC ->T with Trastuzumab and Pertuzumab given from the start of the taxane treatment (i.e. following the anthracycline)

5-Fluorouracil (500 mg/m²), epirubicin (100 mg/m²) followed by cyclophosphamide (600 mg/m²) for three cycles, followed by Docetaxel for three cycles with Trastuzumab (8 mg/kg on day 1 of the first treatment with Docetaxel and 6 mg/kg every 3 weeks thereafter) and Pertuzumab (840 mg on day 1 on the first day of Docetaxel with 420 mg every 3 weeks thereafter). The starting dose for Docetaxel is 75 mg/m² for Cycle 4 (first Docetaxel cycle) then 100 mg/m² for Cycles 5-6, if no dose limiting toxicity occurs. All drugs will be administered by the IV route.

10 OR

Arm C

Taxane (Docetaxel), Carboplatin and Trastuzumab (TCH) with Pertuzumab, with both antibodies being given from the start of the chemotherapy.

Carboplatin (AUC6 using Calvert's Formula) followed by Docetaxel on day 1 with Trastuzumab (8 mg/kg on day 1 of the first treatment with Carboplatin and Docetaxel and 6 mg/kg every 3 weeks thereafter) and Pertuzumab (840 mg on day 1 with 420 mg every 3 weeks thereafter) for six cycles. The dose for Docetaxel is 75 mg/m² for all cycles. All drugs will be administered by the IV route.

All patients will receive Trastuzumab every three weeks for a total of one year from the start of treatment (from Cycle 1-17 for patients in Arms A and C and Cycle 4-20 for patients in Arm B) whether they receive additional chemotherapy or not.

Primary Objective

The primary objective was evaluated when all patients had received six cycles of neoadjuvant treatment, had their surgery and all necessary samples taken or were withdrawn from the study whichever is earlier.

Secondary Objectives

To make a preliminary assessment of the activity associated with each regimen as indicated by the complete pathological response rate.

To evaluate the safety profiles of each treatment regimen, including pre-operative (neoadjuvant) and post-operative (adjuvant) treatment.

To investigate the overall survival, the time to clinical response, time-to-response, disease free survival and progression free survival for each treatment arm.

To investigate the biomarkers that may be associated with primary and secondary efficacy endpoints in accordance with each treatment arm.

To investigate the rate of breast conservative surgery for all patients with T2-3 tumors for whom mastectomy was planned at diagnosis.

- 5 An overall assessment of the risk and benefit of each regimen will be made.

Overview of Study Design

10 This was a Phase II open-label, randomized, multi-center trial to evaluate the tolerability and activity associated with Trastuzumab and Pertuzumab when used in addition to anthracycline-based or carboplatin-based chemotherapy regimens as neoadjuvant therapy in patients with HER2-positive breast cancer which was early stage and >2cm in diameter or locally advanced or inflammatory (see Figure 11).

15 Six cycles of active chemotherapy were administered. However, if it was considered that patients required further therapy post surgery, it was suggested that those patients who had received FEC->T were given CMF (cyclophosphamide, methotrexate and 5-fluorouracil) and that those patients who had received TCH, but who are deemed to require further chemotherapy received FEC (5-fluorouracil, epirubicin and cyclophosphamide).

After the completion of surgery (and after the completion of post-operative chemotherapy if required), patients received radiotherapy as per local clinical standard and those patients whose tumors were estrogen-receptor positive received hormone manipulation as per local clinical standard.

20 In summary, all patients received at least 6 cycles of active chemotherapy and the two antibodies, Pertuzumab and Trastuzumab plus surgery and radiotherapy (as per local standard) plus any hormone manipulation indicated (as per local standard) and continued to receive Trastuzumab to one year in total.

25 Patients whose neoadjuvant study treatment was discontinued prior to surgery were managed as per local practice. Approximately 28 days after the last dose of study medication, patients were asked to perform a final safety assessment (called Final Visit).

Study Population

Overview

30 Female patients, aged 18 years or more, with early stage HER2-positive breast cancer whose primary tumors are > 2 cm with no metastases.

Inclusion Criteria

1. Female patients with locally advanced, inflammatory or early stage, unilateral and histologically confirmed invasive breast cancer. The initial breast cancer assessment should be performed by a physician with experience in surgery for breast cancer. Patients with inflammatory breast cancer must be able to have a core needle biopsy.
2. Primary tumor > 2cm in diameter.
3. HER2-positive breast cancer confirmed by a central laboratory. Tumors must be HER2 3+ by IHC or FISH/CISH + (FISH/CISH positivity mandatory for HER2 2+ tumors).
4. Availability of FFPE tissue (Buffered Formalin method of fixation will be accepted) for central confirmation of HER2 eligibility (FFPE tumor tissue will subsequently be used for assessing status of biomarkers).
5. Female patients, age ≥ 18 years.
6. Baseline LVEF $\geq 55\%$ (measured by echocardiography or MUGA).
7. Performance status ECOG ≤ 1 .
8. At least 4 weeks since major unrelated surgery, with full recovery.

Concomitant Medication and Treatment**Allowed Therapies**

Concomitant treatments are any prescription medications, over-the-counter preparations, herbal remedies or radiotherapy used by a patient in the interval beginning 7 days prior to the patient being recruited into the study and continuing through the study.

The following treatments are permitted during the study:

1. Acceptable methods of contraception must be used when the female patient or male partner is not surgical sterilized or does not meet the study definition of post-menopausal (≥ 12 months of amenorrhea).
2. H₁ and H₂ antagonist (e.g. diphenhydramine, cimetidine)
3. Analgesics (e.g. paracetamol/acetaminophen, meperidine, opioids)
4. Short term use of corticosteroids to treat or prevent allergic or infusion reactions
5. Antiemetics (approved prophylactic serotonin-antagonists, benzodiazepines, ondansetron etc)
6. Medication to treat diarrhea (e.g. loperamide)
7. Colony stimulating factors (e.g. G-CSF)
8. Estrogen receptor antagonist (e.g. tamoxifen) or aromatase inhibitors (e.g. anastrozole, exemestane) after completion of post-operative chemotherapy as per local practice.

Excluded Therapies

The following therapies are excluded during the treatment period of the study:

9. Anti-cancer therapies other than those administered in this study, including cytotoxic chemotherapy, radiotherapy, (except for adjuvant radiotherapy for breast cancer after completion of chemotherapy or additional adjuvant chemotherapy immediately post-surgery, if deemed necessary) immunotherapy, and biological anti-cancer therapy.
10. Any targeted therapy.
11. Treatment with steroids except for thyroid hormone replacement therapy and short term corticosteroid, in order to treat or prevent allergic or infusion reactions.
12. High doses of systemic corticosteroids. High dose is considered as > 20 mg of dexamethasone a day (or equivalent) for > 7 consecutive days.
13. Any investigational agent, except for those used for this study.
14. Initiation of herbal remedies. Herbal remedies initiated prior to study entry and continuing during the study are permitted and must be reported on the appropriate eCRF.
15. Any oral, injected or implanted hormonal methods of contraception.

RESULTS

- The baseline characteristics of the patients with HER2-positive early-stage breast cancer are provided in Table 6 below.

Table 6: Baseline Characteristics in the Safety Population

	FEC+H+P x3 → T+H+P x3 n = 72	FEC x3 → T+H+P x3 n = 75	TCH+P x6 n = 76
Median age, years (range)	49.0 (27-77)	49.0 (24-75)	50.0 (30-81)
ECOG PS 0, n (%) 1, n (%)	65 (91.5) 6 (8.5)	66 (88.0) 9 (12.0)	67 (88.2) 9 (11.8)
ER and/or PR-positive, n (%) ER and PR-negative, n (%)	39 (53.4) 34 (46.6)	35 (46.7) 40 (53.3)	40 (51.9) 37 (48.1)

Disease type, n (%)	53 (72.6)	54 (72.0)	49 (63.6)
Operable	15 (20.5)	17 (22.7)	24 (31.2)
Locally advanced	5 (6.8)	4 (5.3)	4 (5.2)
Inflammatory			
HER2 IHC 0 and 1+, n (%)	1 (1.4)	-	-
2+, n (%)	5 (6.8)	1 (1.3)	2 (2.6)
3+, n (%)	67 (91.8)	74 (98.7)	75 (97.4)
HER2 FISH-positive, n (%)	69 (94.5)	69 (92.0)	73 (94.8)
FISH-negative, n (%)	-	1 (1.3)	2 (2.6)
Unknown, n (%)	4 (5.5)	5 (6.7)	2 (2.6)

5 CBE, clinical breast examination; ECOG PS, Eastern Cooperative Oncology Group performance status; ER, estrogen receptor; FEC, 5-fluorouracil, epirubicin, cyclophosphamide; FISH, fluorescence *in situ* hybridisation; H, Trastuzumab; IHC, immunohistochemistry; P, Pertuzumab; PR, progesterone receptor; T, Docetaxel; TCH, Docetaxel/Carboplatin/Trastuzumab

Safety data are shown in Figure 12 and Tables 7 and 8 below.

Table 7: Cardiac Events Overall

	FEC+H+P x3 → T+H+P x3 n = 72	FEC x3 → T+H+P x3 n = 75	TCH+P x6 n = 76
Symptomatic LVSD (grade ≥3), n (%)	-	2 (2.7)	1 (1.3)
LVSD (all grades), n (%)	5 (6.9)	3 (4.0)	5 (6.6)
LVEF decline ≥10% points from baseline to <50%, n (%)	5 (6.9)	5 (6.7)	5 (6.6)

10 FEC, 5-fluorouracil, epirubicin, cyclophosphamide; H, Trastuzumab; LVEF, left ventricular ejection fraction; LVSD, left ventricular systolic dysfunction; P, Pertuzumab; T, Docetaxel; TCH, Docetaxel/Carboplatin/Trastuzumab

Table 8: Ten Most Common Adverse Events During Neoadjuvant Treatment Grade ≥ 3

Adverse event, n (%)	FEC+H+P x3 → T+H+P x3 n = 72	FEC x3 → T+H+P x3 n = 75	TCH+P x6 n = 76
Neutropenia	34 (47.2)	32 (42.7)	35 (46.1)
Febrile neutropenia	13 (18.1)	7 (9.3)	13 (17.1)
Leukopenia	14 (19.4)	9 (12.0)	9 (11.8)
Diarrhea	3 (4.2)	4 (5.3)	9 (11.8)
Anemia	1 (1.4)	2 (2.7)	13 (17.1)
Thrombocytopenia	-	-	9 (11.8)
Vomiting	-	2 (2.7)	4 (5.3)
Fatigue	-	-	3 (3.9)
Alanine aminotransferase inc.	-	-	3 (3.9)
Drug hypersensitivity	2 (2.8)	-	2 (2.6)

FEC, 5-fluorouracil, epirubicin, cyclophosphamide; H, Trastuzumab; P, Pertuzumab; T, Docetaxel; TCH, Docetaxel/Carboplatin/Trastuzumab

5

Efficacy data are provided in Figures 13 and 14 as well as Tables 9 and 10 below.

10

Table 9: Clinical Response Rate During Neoadjuvant Treatment

	FEC+H+P x3 → T+H+P x3 n = 73	FEC x3 → T+H+P x3 n = 75	TCH+P x6 n = 77
Objective response rate, n (%)	67 (91.8)	71 (94.7)	69 (89.6)
Complete response rate	37 (50.7)	21 (28.0)	31 (40.3)
Partial response rate	30 (41.1)	50 (66.7)	38 (49.4)
Stable disease, n (%)	3 (4.1)	1 (1.3)	5 (6.5)
Progressive disease, n (%)	-	1 (1.3)	-
No assessment, n (%)	3 (4.1)	2 (2.7)	3 (3.9)

FEC, 5-fluorouracil, epirubicin, cyclophosphamide; H, Trastuzumab; P, Pertuzumab; T, Docetaxel; TCH, Docetaxel/Carboplatin/Trastuzumab

5 Table 10: Breast Conserving Surgery in Patients for Whom Mastectomy was Planned

	FEC+H+P x3 → T+H+P x3 n = 46	FEC x3 → T+H+P x3 n = 36	TCH+P x6 n = 37
Achieved, n (%) (95% CI)	10 (21.7) (10.9-36.4)	6 (16.7) (6.4-32.8)	10 (27.0) (13.8-44.1)
Not achieved, n (%)	36 (78.3)	30 (83.3)	27 (73.0)

CI, confidence interval; FEC, 5-fluorouracil, epirubicin, cyclophosphamide; H, Trastuzumab; P, Pertuzumab; T, Docetaxel; TCH, Docetaxel/Carboplatin/Trastuzumab

CONCLUSIONS

- 10 • Results from this study indicate a low incidence of symptomatic and asymptomatic LVSD across all arms

- Concurrent administration of Pertuzumab plus Trastuzumab with epirubicin resulted in similar cardiac tolerability compared with sequential administration or the anthracycline-free regimen
- Neutropenia, febrile neutropenia, leukopenia and diarrhea were most frequently reported adverse events (grade ≥ 3) across all arms
- Regardless of chemotherapy chosen, the combination of Pertuzumab with Trastuzumab in the neoadjuvant setting resulted in high pathological complete response (pCR) rates (57 to 66%)
- TRYPHAENA supports the use of Pertuzumab and Trastuzumab plus anthracycline-based or carboplatin-based chemotherapy in the neoadjuvant and adjuvant settings of early-stage breast cancer.

EXAMPLE 6

Co-Administration of Pertuzumab and Trastuzumab

In the phase III clinical trials above Pertuzumab was administered by intravenous (IV) infusion in saline IV bags to patients with HER2-positive metastatic breast cancer followed by Trastuzumab and the chemotherapeutic agent Docetaxel also using saline IV infusions. The IV infusion process for Pertuzumab and Trastuzumab takes approximately 60 to 90 minutes each with a 30 to 60 minute patient observation period after each drug. Due to this treatment regimen per patient, a visit can take up to 7.5 hours total. As medical payments for both drugs and drug administration services have been under scrutiny in the recent past, there has been emphasis on business practices to shorten time and to increase medical resource utilization in clinical and hospital settings. Increased efficiency of patient care, compliance and treatment is expected by shortening the time patients spend in the clinic for each cycle of treatment.

As part of the phase III Pertuzumab clinical trials, Pertuzumab and Trastuzumab are administered through intravenous (IV) infusion to patients sequentially, i.e. one drug after the other. While Pertuzumab is given as a flat dose (420mg for maintenance, 840mg for loading), Trastuzumab is weight based (6mg/kg for maintenance doses). To increase convenience and minimize the in-clinic time for the patients, the feasibility of co-administering Pertuzumab with Trastuzumab in a single 250mL 0.9% saline polyolefin (PO) or polyvinyl chloride (PVC) IV infusion bag was assessed. The individual monoclonal antibodies have been demonstrated to be stable in infusion bags (PO and/or PVC) over 24 hours at 5°C and 30°C. In this study, the compatibility and stability of Pertuzumab (420mg and 840mg) mixed with either 420mg Trastuzumab (6mg/kg dose for a 70kg patient) or 720mg (6mg/kg for a 120kg patient) in IV bags for up to 24 hours at 5°C or 30°C was evaluated. The controls (i.e. Pertuzumab alone in an IV bag, Trastuzumab alone in an IV bag) and the monoclonal antibody (mAb) mixture samples were assessed using the existing Pertuzumab and Trastuzumab analytical methods, which include color, appearance and clarity (CAC), concentration and turbidity

by UV-spec scan, particulate analysis by HIAC-Royco, size exclusion chromatography (SEC), and ion-exchange chromatography (IEC). Additionally, capillary zone electrophoresis (CZE), image capillary isoelectric focusing (iCIEF), and potency (the Pertuzumab anti-proliferation assay only) was utilized to measure the admixtures containing 1:1 of Pertuzumab:Trastuzumab and their respective controls (420mg Pertuzumab only and 420mg Trastuzumab only) only as a representative case.

Results showed no observable differences by the above assays in the Pertuzumab/Trastuzumab mixtures between the time zero (T0) control and the sample stored up to 24 hours at either 5°C or 30°C. The physicochemical assays as listed above were able to detect both molecules as well as the minor variants in the drug mixture, though some overlaps of monoclonal antibody species were seen in the chromatograms. Furthermore, the drug mixture tested by the Pertuzumab specific inhibition of cell proliferation assay showed comparable potency before and after storage. The results from this study showed the Pertuzumab and Trastuzumab admixtures are physically and chemically stable in an IV infusion bag for up to 24 hours at 5°C or 30°C and can be used for clinical administration if necessary.

Dose I: 840mg Pertuzumab/Trastuzumab mixture (420mg Pertuzumab and 420mg Trastuzumab)

Sample Preparation: All procedures were performed aseptically under a laminar flow hood. PO IV infusion bag samples with three types of drug combinations were prepared for this study: 1) 420mg Pertuzumab/ 420mg Trastuzumab mixture, 2) 420mg Pertuzumab alone, and 3) 420mg Trastuzumab alone. The Pertuzumab and Trastuzumab alone samples served as controls.

Trastuzumab was reconstituted with 20mL of bacteriostatic water for injection (BWFI) and left on lab bench for approximately 15 minutes prior to use. To prepare the Pertuzumab/ Trastuzumab sample dose, 14mL of Pertuzumab (420mg) was diluted directly into the IV infusion bag that contained a nominal 250mL (\pm 25mL overage) 0.9% saline solution, without removing an equal amount of saline, followed by 20mL of the reconstituted Trastuzumab (420mg) using an 18 gauge needle at room temperature. The total concentration of the two proteins combined in the 250mL IV bag was expected to be approximately 3mg/mL. Similarly, the Pertuzumab (420mg) alone IV bag was prepared with 14mL of the 30mg/mL drug product directly diluted into an IV infusion bag. The final expected concentration was approximately 1mg/mL. The Trastuzumab (420mg) alone IV infusion bag was also prepared in the same manner except 20mL of the 21 mg/mL drug product was added into the bag. The final expected concentration was approximately 1mg/mL.

The PO IV bags were manually mixed thoroughly by a gentle back and forth rocking motion several times to ensure homogeneity. After mixing, 10mL of sample was removed with a syringe from each bag and stored in sterile 15cc falcon tubes to be used as the diluted sample control at time zero (T0). The IV bags were then stored covered in foil at 30°C for 24 hours (T24). Immediately after storage, the remainder of the sample was removed with a syringe from each bag and placed into

sterile 250 mL PETG containers. The T0 and T24 samples were held for up to 24 hours at 5°C or immediately analyzed by CAC, UV-spec scan (concentration and turbidity), SEC, IEC, CZE, iCIEF, HIAC-Royco, as well as potency. The product quality of the samples was tested by the Pertuzumab and Trastuzumab product specific SEC and IEC methods, while only the Pertuzumab specific potency method was performed. The other assays utilized were non-product specific methods. All assays were qualified for the intended testing in their respective molecules and used without further method optimization.

Dose II: 1560mg Pertuzumab/Trastuzumab mixture (840mg Pertuzumab and 720mg Trastuzumab)

Sample preparation: The upper range of the mAb co-administration dose was examined (1560mg total mixture: 840mg Pertuzumab and 720mg Trastuzumab) in PO and PVC IV infusion bag samples. In the event that an increase in protein aggregation is observed, the propensity of the formation of high molecular weight species (HMWS) would more likely occur at the upper dose of 1560mg total mAb rather than the mixture containing 840mg. To mitigate the risk during in-use conditions at the high dose range, both PO and PVC IV infusion bags were studied to ensure no interactions were seen.

Three types of drug combinations (mixture, 840mg Pertuzumab alone, and 720mg Trastuzumab alone) were prepared and handled similar to the dose I study. The Pertuzumab/Trastuzumab mixture contained 28mL of Pertuzumab (840mg) diluted directly into either PO or PVC IV infusion bags followed by 34mL of the reconstituted Trastuzumab (720mg) using an 18 gauge needle at room temperature. The total concentration of the two mAbs combined in a single 250mL IV bag was expected to be approximately 5mg/mL. For the controls, Pertuzumab and Trastuzumab alone IV infusion bag samples were prepared and handled similar to the dose I study, except 28mL of 30mg/mL Pertuzumab and 34mL of 21mg/mL Trastuzumab was directly diluted into each PO or PVC IV infusion bag. The final expected concentration was approximately 3mg/mL for the Pertuzumab (840mg) and Trastuzumab (720mg) alone samples. The bags were stored uncovered at either 5°C or 30°C for up to 24 hours. The T0 and T24 samples were analyzed immediately or held for up to 24 to 48 hours at 5°C by CAC, UV-spec scan (concentration and turbidity), SEC, IEC, and HIAC-Royco.

Details of the types of doses, IV infusion bags, dose & preparation, storage temperatures, and assays are summarized in Table 11.

Table 11: IV Bag Type, Dose, Preparation & Study Conditions

IV bag type (approx 250mL, 0.9%NaCl)	Total dose, concentration	Dilution (into approx 250mL IV bag)	Storage temperature (up to 24hrs)	Assays
Dose I (n=1)				
PO	840mg, approx 3mg/mL	Add 14mL P (~30mg/mL) + 20mL T (~21mg/mL)	30 °C	CAC, UV spec scan (concentration, turbidity), SEC, IEC, CZE, icIEF, HIAC- Royco, potency
	420mg ^b , approx 2mg/mL	Add 14mL P (~30mg/mL)		
	420mg ^b , approx 1mg/mL	Add 20mL T (~21mg/mL)		
Dose II (n=1)				
PO and PVC ^a	1560mg, approx 5mg/mL	Add 28mL P (~30mg/mL) + 34mL T (~21mg/mL)	5 °C 30 °C	CAC, UV spec scan (concentration & turbidity), SEC, IEC, HIAC-Royco
PO and PVC	840mg ^b , approx 3mg/mL	Add 28mL P (~30mg/mL)		
	720mg ^b , approx 3mg/mL	Add 34mL T (~21mg/mL)		

^a n=2^b control

P= Pertuzumab; T= Trastuzumab

Assays

All samples were held at 5°C or immediately analyzed. Typically, samples were analyzed within 24 to 48 hours of preparation and storage. The following assays were conducted to ascertain product quality and short term stability of Pertuzumab/ Trastuzumab mixture, Pertuzumab alone, and Trastuzumab alone samples diluted into saline IV infusion bags. Since several assays, i.e. SEC, IEC, CZE, icIEF, and potency, were not optimized for quantitative assessment of the mAb mixtures, only chromatographic or electropherographic overlays of these samples and their individual controls before and after storage at 5°C or 30°C are shown here. For consistency, no values, e.g. percent peak area, were calculated for all three sample types from the liquid chromatography and electrophoretic assays that were performed.

Color, Appearance, and Clarity (CAC)

The color, appearance, and clarity of the samples were determined by visual inspection under a white fluorescence light with black and white background at room temperature. A 3cc glass vial was filled with 1 mL of each sample for CAC testing. A negative control (purified water) with the corresponding sample volume was used for comparison.

UV- Vis Spectrophotometer Scan for Concentration Measurements

The concentration was determined by measurement of the UV-absorbance on an HP8453 spectrophotometer via volumetric sample preparation. The instrument was blanked with 0.9% saline.

Absorbance at A_{\max} (278nm or 279nm) and 320nm in a quartz cuvette with 1-cm path length were measured for each sample. The absorbance at 320nm is used to correct for background light scattering in solution. The concentration determination was calculated by using the absorptivity of 1.50 (mg/mL)⁻¹cm⁻¹ for both Pertuzumab and Trastuzumab molecules.

$$5 \quad \text{Protein Concentration (mg/mL)} = \frac{A_{\max} - A_{320}}{1.50} \times \text{Dilution Factor} \times \frac{1}{\text{cuvette pathlength}}$$

Size Exclusion Chromatography (SEC: Pertuzumab specific and Trastuzumab specific)

Each sample was injected into a TOSOHAAS® column G3000 SWXL, 7.8 X 300mm at ambient temperature on an AGILENT 1100® HPLC. The eluted peaks were monitored at 280nm. Chromatographic integrations were analyzed by the CHROMELEON® software. The autosampler
 10 temperature was held at 2-8°C throughout the run and mobile phases used were 0.2M potassium phosphate, 0.25mM potassium chloride, pH 6.2 and 100mM potassium phosphate, pH 6.8 for Pertuzumab-assay and Trastuzumab-assay, respectively. The recommended injection load as specified by the test procedure was 200 µg with an injection volume of 20 µL. The diluted 420mg sample was injected at a load less than the recommended amount due to the low concentration of the
 15 protein after dilution in the IV bags. The maximum injection volume of the HPLC sample loop was 100 µL, which limits the volume that is able to be injected at one time. As a result, the injection volumes were modified to 100 µL at 160 µg protein for the Pertuzumab alone and Trastuzumab alone samples (420 mg dose group) and 73 µL at 200 µg protein for the Pertuzumab/Trastuzumab mixture (840 mg dose group). Modification in the injection volumes have been utilized in previous IV bag
 20 studies and are necessary when handling low concentration samples.

Ion-exchange Chromatography (IEC)

The analysis of carboxypeptidase B (CpB)- digested Pertuzumab and Trastuzumab for charge heterogeneity was employed by IEC for each sample. For the Pertuzumab specific IEC, samples were either tested with regular IEC ("Pertuzumab-regular IEC") or a modified "fast" version
 25 of IEC ("Pertuzumab-IEC-fast") for high throughput, method for the purpose of these experiments. The IEC assays utilized the a DIONEX® WCX weak cation exchange column equilibrated with solvent A (20mM MES, 1mM Na₂EDTA pH 6.00) and solvent B (250mM sodium chloride in solvent A) monitored at 280nm for Pertuzumab-regular IEC and Pertuzumab-IEC-fast, whereas solvent A
 (10mM sodium phosphate, pH 7.5) and solvent B (100mM sodium chloride in Solvent A) monitored
 30 at 214nm was used for Trastuzumab on an AGILENT 1100® HPLC. The peaks were eluted at a flow rate of 0.8 mL/min with an increasing gradient of 18%-100% solvent B over 35 minutes and 90 minutes for Pertuzumab-regular-IEC and Pertuzumab-IEC- fast, respectively, and 15%-100% solvent B over 55 minutes for Trastuzumab-IEC. Column temperatures were maintained at either 34°C or

42°C and ambient for Pertuzumab-regular-IEC or Pertuzumab-fast-IEC and Trastuzumab-IEC, respectively, while the auto sampler temperature was held at 2-8°C throughout the run.

HIAC-ROYCO™ light obscuration for sub-visible particles

Particulate counts in the diluted drug product were carried out using the HIAC-ROYCO™ Liquid Particulate Counting System model 9703. Average cumulative numbers of particles at $\geq 10\mu\text{m}$ and $\geq 25\mu\text{m}$ per milliliter were tabulated in each sample using PHARMSPEC v2.0™. The test procedure was modified for a small-volume method, utilizing either four 1mL readings or four 0.4mL readings per a test session while discarding the first reading of each sample. The HIAC-ROYCO™ samples were degassed under vacuum for approximately 10-15 minutes each. The size below $10\mu\text{m}$ was not collected for this sample set.

UV- Vis Spectrophotometer Scan for Turbidity Measurements

The optical density of the samples from the IV bag (1mg/mL or 3mg/mL) was measured in a quartz cuvette with a 1-cm path length on a HP8453 spectrophotometer. The sample readings were blanked against purified water. The absorbance measurements were recorded at 340nm, 345nm, 350nm, 355nm, and 360nm and the turbidity was expressed as an average of these wavelengths.

Capillary Zone Electrophoresis, CZE

CZE was performed using a PROTEOMELAB PA800™ capillary electrophoresis system (Beckman Coulter) with neutral-coated capillary (50 μm x 50 cm). The buffer consisted of 40 mM ϵ -amino caproic acid/acetic acid, pH 4.5, 0.2% hydroxypropyl methyl cellulose (HPMC). Samples were diluted to 0.5 mg/mL in water and injected into the capillary at 1 psi for 10 seconds. Separation was performed using a voltage of 30 kV for 15 minutes, and the species were detected by UV at 214 nm.

CE-SDS-LIF, reduced and non-reduced

Each sample was derivatized with 5 carboxytetramethylrhodamine succinimidyl ester, a fluorescent dye. After removing the free dye through gel filtration (using NAP-5 columns), non-reduced samples were prepared by adding 40 mM iodoacetamide and heated at 70°C for 5 minutes. For the analysis of the reduced samples, the derivatized samples were mixed with SDS to a final concentration of 1% (v/v) and 10 mL of a solution containing 1 M DTT, and heated at 70°C for 20 minutes. The prepared samples were analyzed on a Beckman Coulter ProteomeLab PA800 system using a 50 mm I.D. 31.2 cm fused silica capillary maintained at 20°C throughout the analysis. Samples were introduced into the capillary by electrokinetic injection at 10 kV for 40 seconds. The separation was conducted at a constant voltage of 15 kV in the reversed polarity (negative to positive) mode using CE-SDS running buffer as the sieving medium. An argon ion laser operating at 488 nm was used for fluorescence excitation with the resulting emission signal monitored at 560 nm.

iCIEF

The distribution of charge variants of the Pertuzumab/Trastuzumab mixture, Pertuzumab alone, and Trastuzumab alone was assessed by iCIEF using an iCE280™ analyzer (Convergent Bioscience) with a fluorocarbon coated capillary cartridge (100 µm x 5 cm). The ampholyte solution consisted of a mixture of 0.35% methyl cellulose (MC), 0.47% Pharmalyte 3-10 carrier ampholytes, 2.66% Pharmalyte 8-10.5 carrier ampholytes, and 0.20% pI markers 7.05 and 9.77 in purified water. The anolyte was 80 mM phosphoric acid, and the catholyte was 100 mM sodium hydroxide, both in 0.10% methylcellulose. Samples were diluted in purified water and CpB was added to each diluted sample at an enzyme to substrate ratio of 1:100 followed by incubation at 37°C for 20 minutes. The CpB treated samples were mixed with the ampholyte solution and then focused by introducing a potential of 1500 V for one minute, followed by a potential of 3000 V for 10 minutes. An image of the focused Pertuzumab charge variants was obtained by passing 280 nm ultraviolet light through the capillary and into the lens of a charge coupled device digital camera. This image was then analyzed to determine the distribution of the various charge variants.

15 Anti-Proliferation Potency Assay

This test procedure is based on the ability of Pertuzumab to inhibit the proliferation of MDA MB 175 VII human breast carcinoma cells. Briefly, cells were seeded in 96-well tissue culture microtiter plates and incubated overnight at 37 °C under 5% CO₂ to allow cell attachment. The following day, the culture medium was removed and serial dilutions of each standard, controls, and sample(s) were added to the plates. The plates were then incubated for four days at 37 °C under 5% CO₂ and the relative number of viable cells was quantified indirectly using a redox dye, ALAMARBLUE® according to the manufacturer's protocol. Each sample was assayed in triplicate and the changes in color as measured by fluorescence were directly proportional to the number living cells in the culture. The absorbance of each well was then measured on a fluorescence 96-well plate reader. The results, expressed in relative fluorescence units (RFU), were plotted against the antibody concentration. No quantitative measurements were made, or possible, since there was no Pertuzumab/Trastuzumab mixture reference available. Therefore, the results are comparisons of the dose response curves only.

RESULTS AND DISCUSSION**30 Dose I: 840mg total Pertuzumab/ Trastuzumab mixture (420mg Pertuzumab and 420mg Trastuzumab)**

The product quality of the total 840mg Pertuzumab/Trastuzumab mixture (420mg Pertuzumab and 420mg Trastuzumab), Pertuzumab alone (420mg), and Trastuzumab alone (420mg) in IV infusion bags (n=1) before and after storage at 30°C for up to 24 hours was assessed by CAC, concentration measurements by UV-spec scan, turbidity, and HIAC Royco (Table 12). The

Pertuzumab and Trastuzumab alone IV infusion bags are considered controls that were also prepared to assess the ability of the assay to pick up the appropriate product attributes.

Table 12: Dose I 840mg: Stability data for Pertuzumab/Trastuzumab mixture, Pertuzumab, or Trastuzumab in 0.9% saline PO IV infusion bags (n=1)

5

Sample	IV bag type	Amount	Timepoint	Temp	CAC ^a	Conc.	Turbidity	Light Obscuration	
		mg	Hour(s)	°C	liquid	mg/mL	AU	total particles ≥ 10µm/mL	total particles ≥ 25µm/mL
pertuzumab/ trastuzumab mixture	PO	840	0	30	CL, CO	2.7	0.016	1	0
		840	24	30	CL, CO	2.7	0.016	6	0
pertuzumab	PO	420	0	30	CL, CO	1.4	0.012	3	0
		420	24	30	CL, CO	1.4	0.011	4	0
trastuzumab	PO	420	0	30	CL, CO	1.5	0.012	1	0
		420	24	30	CL, CO	1.5	0.011	6	0
saline only		-	-	-	-	-	-	2	1

^a Color, Appearance and Clarity: CL=clear ; SOPL= slightly opalescent, CO= colorless.

RT= room temperature

^a Color, appearance, and clarity: CL= clear; SOPL= slightly opalescent, CO= colorless

After storage, the Pertuzumab/Trastuzumab mixture, Pertuzumab alone, and Trastuzumab alone samples appeared as a clear and colorless liquid with no visible particles as observed by CAC. The concentration and turbidity measurements showed no measureable changes in any of the three sample types after 24 hours at 30°C. Particulate analysis by HIAC Royco detected no more than 6 particles greater than or equal to 10µm size and no particles greater than 25 µm size for Pertuzumab/Trastuzumab mixture, Pertuzumab alone, or Trastuzumab alone samples post storage. These results are comparable to the 0.9% saline only solution. The lack of visible precipitation or particulates indicates that the admixture and the controls are sufficiently stable upon dilution in the 0.9% saline IV infusion bags. The Pertuzumab/Trastuzumab mixture diluted in saline were run on SEC, both Pertuzumab and Trastuzumab specific methods, and showed comparable peak profiles between T0 and T24 (Figures 15 and 16). No increases were observed in the high molecular weight species (HMWS) and low molecular weight species (LMWS). Similarly, no changes were observed in the main peak in any sample. The main peak and the peak area of the HMWS and LMWS overlay and cannot be distinguished in the Pertuzumab/ Trastuzumab mixture due the size similarity between Pertuzumab and Trastuzumab (molecular weight approximately 150kD). Furthermore, comparison of T0 and T24 for both the Pertuzumab and Trastuzumab alone sample showed no observable changes in peak area or profile as detected by the two SEC methods listed above.

Two product specific methods for Pertuzumab or Trastuzumab IEC was utilized to analyze the Pertuzumab/Trastuzumab mixture (Figures 17 and 18). In the cation-exchange chromatography

assays, each molecule typically contain three distinct areas that are eluted based on relative charge, with the early eluting acidic variants, followed by the main peak, and lastly the late eluting basic variants. In the Pertuzumab and Trastuzumab alone chromatograms, the profile exhibiting the acidic variants, main peak, and the basic variants was observed and deemed comparable between the starting material and post storage at 30°C. These results are also consistent with prior studies conducted in saline IV infusion bags for the either Pertuzumab or Trastuzumab alone. For the Pertuzumab/ Trastuzumab mixture chromatogram, the Pertuzumab peaks elute first followed by the Trastuzumab peaks. Due to the nature of cation-exchange separation and the net charge difference between Pertuzumab (~pI 8.7) and Trastuzumab (~pI 8.9), two main peaks, or major charged species, are observed in the Pertuzumab/Trastuzumab mixture. In contrast, the SEC assay separates based on the hydrodynamic size of the molecule and show only one main peak due to the size similarity between Pertuzumab and Trastuzumab. The charged regions of each molecule appear to overlap with each other in the Pertuzumab/Trastuzumab mixture. Specifically, the Pertuzumab basic variants expected to elute at approximately 32 minutes and at 35 minutes appear to overlap with the main peak of Trastuzumab (Figures 17 and 18). Furthermore, the acidic variants of Trastuzumab expected to elute before the Trastuzumab main peak co-elute with the Pertuzumab basic variants and main peak. Despite the overlapping peak regions, the Pertuzumab/Trastuzumab mixture exhibited comparable chromatographic peak profiles before and after storage in IV saline bags for 24 hours at 30°C.

The Pertuzumab/Trastuzumab mixture, Pertuzumab alone, and Trastuzumab alone samples were also assayed on CE-SDS LIF under non-reduced conditions after storage for 24 hours at 30°C. The Pertuzumab/Trastuzumab mixture showed consistent peak profiles with no observable changes after storage compared to the starting material (Figures 19 and 20). A very slight baseline level variation attributed to noise is also observed and does not impact peak area. Similar to SEC, the non-reduced Pertuzumab/Trastuzumab mixture showed only one superimposed monomer constituting both the Pertuzumab and Trastuzumab main species. The Pertuzumab and Trastuzumab alone samples showed no changes at T0 compared to T24. However, individual molecular attributes, e.g. fragment peak level and species, between Pertuzumab/Trastuzumab mixture, Pertuzumab alone, and Trastuzumab alone was observed as expected.

Two major peaks known as the light chain (LC) and heavy chain (HC) are detected at 17 and 21.5 minutes, respectively, when Pertuzumab/ Trastuzumab mixture, Pertuzumab alone, and Trastuzumab alone was run on CE-SDS LIF reduced with DTT (Figure 20). No increase in fragmentation or concomitant decrease in the LC and HC was seen post storage at 30°C for the Pertuzumab/ Trastuzumab mixture. Furthermore, no detectable peak profile differences were noticed in the Pertuzumab and Trastuzumab alone samples post storage.

The charge separation assays CZE and iCIEF show comparable peak profiles for the Pertuzumab/Trastuzumab mixture after storage at 30°C (Figures 21 and 22). The Pertuzumab and Trastuzumab alone when compared to their respective T0 also showed consistent peak profiles with no changes after storage. Furthermore, the presence of various minor species was also observed, although no new peaks were detected upon dilution in the IV bag saline solution. As seen in the charge based IEC assay, two main peaks flanked by smaller overlapping peaks can be detected and was attributed to the difference in the molecular pI.

The potency results based on comparison of the dose response curve showed no impact on the potency of the Pertuzumab/Trastuzumab mixture stored at 30°C for 24 hours compared to its corresponding T0 dose response curve (Figure 23). The Trastuzumab alone showed little activity in the Pertuzumab potency assay. The Pertuzumab/Trastuzumab mixture dose response curve compared to the dose response curve of Pertuzumab or Trastuzumab alone showed that lower doses of the Pertuzumab/Trastuzumab mixture were needed to inhibit the growth of cells as compared to Pertuzumab alone, suggesting there may be an additive or synergistic effect on the inhibition of cell proliferation for the mixture.

Dose II: 1560mg total Pertuzumab/Trastuzumab mixture (840mg Pertuzumab and 720mg Trastuzumab)

In addition to the dose I study at 840mg total mAb, a higher dose of 1560mg mixture (840mg Pertuzumab and 720mg Trastuzumab), and their individual drug product controls (840mg Pertuzumab alone and 720mg Trastuzumab alone) was selected to investigate the impact of diluting these three mAb types in PO or PVC IV infusion bags at 5°C or 30°C for up to 24 hours. The product quality of these IV infusion bags before and after storage was assessed by CAC, UV-spec scan (concentration and turbidity), and HIAC- ROYCO™ are summarized in Table 13 and SEC and IEC are shown in Figures 24-27.

Table 13: Dose II 1560mg: Stability data for Pertuzumab/Trastuzumab mixture, Pertuzumab, or Trastuzumab in 0.9% saline PO IV infusion bags (n=1 for control; n=2 for mixture)

Sample	IV bag type	Amount	Temp	Timepoint	CAC ^a	Conc.	Turbidity	Light Obscuration	
		mg	°C	Hour(s)	liquid	mg/mL	AU	total particles ≥ 10um/mL	total particles ≥ 25um/mL
pertuzumab/ trastuzumab mixture	PO	1560	5	0	CL, CO	4.9	0.013	20	1
				24	CL, CO	4.7	0.016	15	1
		1560	30	0	CL, CO	5.0	0.014	8	0
				24	CL, CO	4.9	0.018	18	0
pertuzumab	PO	840	5	0	CL, CO	2.9	0.007	1	0
				24	CL, CO	2.9	0.005	6	0
		840	30	0	CL, CO	2.8	0.006	6	0
				24	CL, CO	2.8	0.004	9	0
trastuzumab	PO	720	5	0	CL, CO	2.6	0.004	4	0
				24	CL, CO	2.6	0.005	13	0
		720	30	0	CL, CO	2.5	0.007	19	0
				24	CL, CO	2.5	0.004	14	0
pertuzumab/ trastuzumab mixture	PVC	1560	5	0	CL, CO	4.9	0.016	18	0
				24	CL, CO	4.7	0.015	18	0
		1560	30	0	CL, CO	4.8	0.016	24	0
				24	CL, CO	4.8	0.012	17	0
pertuzumab	PVC	840	5	0	CL, CO	2.9	0.006	13	0
				24	CL, CO	2.7	0.004	10	0
		840	30	0	CL, CO	2.8	0.006	6	0
				24	CL, CO	2.8	0.006	11	0
trastuzumab	PVC	720	5	0	CL, CO	2.5	0.007	7	0
				24	CL, CO	2.5	0.004	9	0
		720	30	0	CL, CO	2.5	0.003	18	0
				24	CL, CO	2.5	0.005	19	0

^a Color, Appearance and Clarity: CL =clear ; SOPL= slightly opalescent, CO= colorless.

Two PO or PVC IV infusion bags each were prepared for the Pertuzumab/Trastuzumab mixture condition while only one IV infusion bag was prepared for the Pertuzumab and Trastuzumab alone samples.

Particulates from these bags were determined by visual observation, turbidity, and HIAC-Royco measurements. All samples appeared clear and colorless after storage at 5°C or 30°C for up to 24 hours. No visible particulate matter was observed and there was no significant change in the turbidity

post storage. For the Pertuzumab/ Trastuzumab mixture, Pertuzumab alone, and Trastuzumab alone, the HIAC-Royco showed comparable particle values before and after storage, with zero to 10 particles increase per milliliter at $\geq 10 \mu\text{m}$ and zero particle increase per milliliter at $\geq 25 \mu\text{m}$ for both PO and PVC IV infusion bags stored at either 5°C or 30°C. Similarly, the Pertuzumab and

5 Trastuzumab alone samples also exhibited no significant particle differences before and after storage in PO or PVC IV infusion bags. For all three sample types, the UV-spec scan showed no changes beyond normal assay variability in protein concentration, indicating the absence of protein adsorption or precipitation in the IV infusion bags between T0 and T24 hours at 5°C or 30°C storage.

10 The Pertuzumab/ Trastuzumab mixture, Pertuzumab alone, and Trastuzumab alone samples were analyzed using Pertuzumab or Trastuzumab specific SEC and IEC methods to assess their physical and chemical stability, respectively, as previously described. For the Pertuzumab/ Trastuzumab mixture, no changes in SEC were observed in the chromatographic profiles between the T0 and the T24 hour samples at 5°C or 30°C in either PO or PVC IV infusion bags (Figures 24 and 25), similar to the 840mg mixture dose I results. In addition, no increase or decrease in the high molecular weight species (HMWS), main peak, and low molecular weight species (LMWS) was observed, which
15 indicates a stable dosing solution at the upper ranges of protein content in 0.9% saline. Likewise, Pertuzumab alone and Trastuzumab alone samples also showed no changes after storage in the IV infusion bags.

IEC analysis, using both the Pertuzumab or Trastuzumab specific methods, of the Pertuzumab/
20 Trastuzumab mixture was used to assess chemical stability and showed comparable charge variant peak profiles with no observed changes relative to the initial time point after exposure to 5°C or 30°C in the PO or PVC IV infusion bags (Figures 26 and 27). Although a significant overlap of the charge variant species of the two mAbs were observed, these peaks species were not impacted from the increase in the mAb content of the IV infusion bag. Pertuzumab alone or Trastuzumab alone samples
25 in PO or PVC IV infusion bags showed no changes before and after exposure to 5°C or 30°C. These results are consistent with the 840mg dose I study.

CONCLUSION

All physicochemical assays indicate no significant changes in the mixtures (up to 840mg Pertuzumab and 720mg Trastuzumab for a 1560mg total dose) or in the individual Pertuzumab (up to
30 840mg) and Trastuzumab (up to 720mg) IV infusion bags (PO or PVC) for T0 to T24 hours at 5°C or 30°C. Furthermore, the potency of the mixture (up to 840mg) and the individual mAbs before and after storage were comparable. No differences were observed in the IV bags that contained the admixture of Pertuzumab and Trastuzumab when compared to the individual mAb components in IV bags over the course of this study. The current study also demonstrates that many of the assays used
35 to measure the individual mAbs were sufficient to qualitatively characterize the admixture.

EXAMPLE 7**Co-Administration of Pertuzumab and Trastuzumab, and
Combination Therapy with Vinorelbine**

This is a randomized, two-arm, open-label, multicenter Phase II trial to evaluate Pertuzumab in patients with HER2-positive advanced breast cancer (metastatic or locally advanced) who have not previously received systemic non-hormonal anticancer therapy in the metastatic setting. The study design is shown in Figure 28.

Patients are randomly assigned in a 2:1 ratio to one of two treatment arms:

- Pertuzumab given in combination with Trastuzumab and vinorelbine (**Arm A**)
- Trastuzumab and vinorelbine (control arm **Arm B**)

Arm A will consist of two cohorts as follows:

Cohort 1: (first 95 patients): Pertuzumab and Trastuzumab administered sequentially in separate infusion bags, followed by vinorelbine. Patients will receive Pertuzumab followed by Trastuzumab sequentially in separate infusion bags, followed by vinorelbine.

Pertuzumab (IV infusion)

Administered on Day 1 of the first treatment cycle as a loading dose of 840 mg, followed by 420 mg on Day 1 of each subsequent 3 weekly cycle.

Initial infusions of Pertuzumab will be administered over 90 (\pm 10) minutes and patients observed for at least 30 minutes from the end of infusion for infusion-related symptoms such as fever, chills etc. Interruption or slowing of the infusion may reduce such symptoms. If the infusion is well tolerated, subsequent infusions may be administered over 30 (\pm 10) minutes with patients observed for a further 30 minutes.

Trastuzumab (IV infusion)

Day 1 of the first treatment cycle as a loading dose of 8 mg/kg, followed by 6 mg/kg on Day 1 of each subsequent 3 weekly cycle; to be administered in line with product labeling.

Vinorelbine (IV infusion after Trastuzumab)

Day 1 and Day 8 of the first treatment cycle at a dose of 25 mg/m² followed by 30–35 mg/m² on Day 1 and Day 8 of each subsequent 3 weekly cycle; to be administered in line with product labeling.

Cohort 2: The second 95 patients will receive Pertuzumab and Trastuzumab administered together in a single infusion bag from Cycle 2 onwards, followed by vinorelbine.

Cycle 1 dosing

For the first cycle of treatment, Pertuzumab and Trastuzumab will be administered in separate infusion bags as described for Cohort 1.

Vinorelbine will be administered after Pertuzumab and Trastuzumab as described for Cohort 1.

Subsequent cycle dosing

- 5 If administration of all three drugs was well tolerated in Cycle 1, then on Day 1 of each subsequent 3 weekly treatment cycle, Pertuzumab 420 mg and Trastuzumab 6 mg/kg will be given together in a single infusion bag.

- 10 The first combined infusion of Pertuzumab and Trastuzumab should be administered over 90 (± 10) minutes with cardiac monitoring and close observation for infusion-associated reactions during the procedure, followed by a 60 minute observation period. If this first combined infusion is well tolerated, subsequent combined infusions can be administered over 60 (± 10) minutes followed by a 30 minute observation period with cardiac monitoring.

Vinorelbine will be administered after Pertuzumab and Trastuzumab as described for Cohort 1.

Control arm - Arm B

- 15 A total of 95 patients will be randomized to arm B.

Trastuzumab (IV infusion)

Day 1 of the first treatment cycle as a loading dose of 8 mg/kg, followed by 6 mg/kg on Day 1 of each subsequent 3 weekly cycle; to be administered in line with product labeling.

Vinorelbine (IV infusion after Trastuzumab)

- 20 Day 1 and Day 8 of the first treatment cycle at a dose of 25 mg/m² followed by 30–35 mg/m² on Day 1 and Day 8 of each subsequent 3 weekly cycle; to be administered in line with product labeling.

Efficacy Outcomes:

Primary

- 25 • To compare objective overall response rates (ORR) assessed by a blinded independent review committee (IRC) of Pertuzumab given in combination with Trastuzumab and vinorelbine versus Trastuzumab and vinorelbine

Secondary

- 30 • Within the Pertuzumab treatment group to compare the efficacy and safety of Pertuzumab and Trastuzumab administered together in a single infusion bag versus conventional sequential administration in separate infusion bags
- To compare Pertuzumab given in combination with Trastuzumab and vinorelbine versus Trastuzumab and vinorelbine with respect to:
- ORR assessed by the Investigator

- Time to response assessed by IRC and Investigator
- Duration of response assessed by IRC and Investigator
- Progression free survival (PFS)
- Time to progression (TTP)
- 5 ○ Overall survival (OS)
- Safety and tolerability
- Quality of life (EQ-5D and FACT-B questionnaires)

Inclusion Criteria

Patients must meet the following criteria to be eligible for this study according to the timing of the

10 Schedule of Assessments:

1. Female or male patients aged 18 years or older
2. Histologically or cytologically confirmed and documented adenocarcinoma of the breast with metastatic or locally advanced disease not amenable to curative resection
3. HER2-positive (defined as either immunohistochemistry (IHC) 3+ or in situ hybridization
15 (ISH) positive) as assessed by local laboratory on primary or metastatic tumor (ISH positivity is defined as a ratio of 2.0 or greater for the number of HER2 gene copies to the number of signals for CEP17, or for single probe tests, a HER2 gene count greater than 4).
4. At least one measurable lesion and/or non-measurable disease evaluable according to Response Evaluation Criteria In Solid Tumors (RECIST) version 1.1
- 20 5. ECOG performance status 0 or 1
6. Left ventricular ejection fraction (LVEF) of at least 50%
7. Negative pregnancy test in women of childbearing potential (premenopausal or less than 12 months of amenorrhea post-menopause, and who have not undergone surgical sterilization)
8. For women of childbearing potential who are sexually active, agreement to use a highly-
25 effective, non-hormonal form of contraception or two effective forms of non-hormonal contraception during and for at least 6 months post study treatment
9. Fertile males willing and able to use effective non-hormonal means of contraception (barrier method of contraception in conjunction with spermicidal jelly, or surgical sterilization) during and for at least 6 months post-study treatment
- 30 10. Life expectancy of at least 12 weeks

Exclusion Criteria

Patients who meet any of the following exclusion criteria will not be eligible for this study:

1. Previous systemic non-hormonal anticancer therapy in the metastatic or locally advanced breast cancer setting

2. Previous approved or investigative anti-HER2 agents in any breast cancer treatment setting, except Trastuzumab in the adjuvant or neoadjuvant setting
3. Disease progression while receiving Trastuzumab in the adjuvant or neoadjuvant setting
4. Disease-free interval from completion of adjuvant or neo-adjuvant systemic non-hormonal treatment to recurrent disease of less than 6 months
5. History of persistent grade 2 or higher (NCI-CTC, Version 4.0) hematological toxicity resulting from previous adjuvant or neoadjuvant therapy
6. Radiographic evidence of central nervous system (CNS) metastases as assessed by CT or MRI
7. Current peripheral neuropathy of grade 3 or greater (NCI-CTC, Version 4.0)
- 10 8. History of other malignancy within the last 5 years, except for carcinoma in situ of the cervix or basal cell carcinoma
9. Serious uncontrolled concomitant disease that would contraindicate the use of any of the investigational drugs used in this study or that would put the patient at high risk for treatment related complications
- 15 10. Inadequate organ function, evidenced by the following laboratory results:
 - Absolute neutrophil count $<1,500$ cells/mm³
 - Platelet count $<100,000$ cells/mm³
 - Hemoglobin <9 g/dL
 - Total bilirubin greater than upper limit of normal (ULN) (unless the patient has documented Gilbert's syndrome)
 - 20 • AST (SGOT) or ALT (SGPT) $>2.5 \times$ ULN
 - AST (SGOT) or ALT (SGPT) $>1.5 \times$ ULN with concurrent serum alkaline phosphatase $>2.5 \times$ ULN; Serum alkaline phosphatase may be $>2.5 \times$ ULN only if bone metastases are present and AST (SGOT) and ALT (SGPT) $<1.5 \times$ ULN
 - 25 • Serum creatinine >2.0 mg/dL or 177 μ mol/L
 - International normalized ratio (INR) and activated partial thromboplastin time or partial thromboplastin time (aPTT or PTT) $>1.5 \times$ ULN (unless on therapeutic coagulation)
11. Uncontrolled hypertension (systolic >150 mm Hg and/or diastolic >100 mm Hg) or clinically significant (i.e. active) cardiovascular disease: cerebrovascular accident (CVA)/stroke or myocardial infarction within 6 months prior to first study medication, unstable angina, congestive heart failure (CHF) of New York Heart Association (NYHA) grade II or higher, or serious cardiac arrhythmia requiring medication
- 30 12. Current known infection with HIV, HBV, or HCV
13. Dyspnea at rest due to complications of advanced malignancy, or other disease requiring continuous oxygen therapy
- 35

14. Major surgical procedure or significant traumatic injury within 28 days prior to randomization or anticipation of need for major surgery during the course of study treatment
 15. Receipt of intravenous (IV) antibiotics for infection within 14 days prior to randomization
 16. Current chronic daily treatment with corticosteroids (dose equivalent to or greater than
5 10 mg/day methylprednisolone), excluding inhaled steroids
 17. Known hypersensitivity to any of the study medications or to excipients of recombinant human or humanized antibodies
 18. History of receiving any investigational treatment within 28 days prior to randomization
 19. Concurrent participation in any clinical trial
- 10 It is anticipated that the treatment herein will demonstrate the safety and efficacy of co-administration of pertuzumab and Trastuzumab from the same intravenous (IV) bag to patients with HER2-positive cancer (exemplified by HER2-positive breast cancer), as well as the safety and efficacy of Pertuzumab in combination in vinorelbine according to any one or more of the primary or secondary efficacy outcomes above.

EXAMPLE 8

Pertuzumab Combined with Aromatase Inhibitors

- 15 This example is a randomized, two-arm, open-label, multicenter phase II study demonstrating the efficacy and safety of Pertuzumab given in combination with Trastuzumab plus an aromatase inhibitor in first line patients with HER2-positive and hormone receptor-positive advanced (metastatic or locally advanced) breast cancer. The study design is shown in Figure 29.

20 Primary Objectives

To compare progression-free survival (PFS) of Pertuzumab given in combination with Trastuzumab plus an aromatase inhibitor (AI) versus Trastuzumab plus an AI.

Secondary Objectives

- 25 To compare Pertuzumab given in combination with Trastuzumab plus an AI versus Trastuzumab plus an AI with respect to:
- Overall survival (OS)
 - Overall response rate (ORR)
 - Clinical benefit rate (CBR)
 - 30 - Duration of response
 - Time to response
 - Safety and tolerability
 - Quality of life (EQ-5D questionnaires)

Trial Design

Patients will be randomly assigned in a 1:1 ratio to one of two treatment arms:

- Pertuzumab in combination with Trastuzumab plus an AI (**Arm A**).
- Trastuzumab plus an AI (control arm **Arm B**).

5 At the investigator's discretion, patients may also receive induction chemotherapy (a taxane, either Docetaxel or paclitaxel), in combination with the assigned monoclonal antibody treatment arm up to the first 18 weeks of the treatment period. In patients receiving induction chemotherapy, treatment with the AI will start after the chemotherapy induction phase.

Stratification factors for analysis will be:

- 10 -
- Chosen to receive induction chemotherapy (Yes/No).
 - Time since adjuvant hormone therapy (<12 months, ≥12 months, or no prior hormone therapy).

Patients with HER2-positive and hormone receptor-positive (estrogen receptor (ER)-positive and/or progesterone receptor (PgR)-positive) advanced breast cancer (metastatic or locally advanced)
15 who have not previously received systemic nonhormonal anticancer therapy in the metastatic setting.

Inclusion Criteria

1. Age greater than or equal to 18 years.
2. Postmenopausal status >1 year (fulfilling one or more of National Comprehensive Cancer Network
20 (NCCN) guideline criteria, Version 2.2011).
3. Histologically or cytologically confirmed and documented adenocarcinoma of the breast with metastatic or locally advanced disease not amenable to curative resection.
4. HER2-positive (defined as either IHC 3+ or ISH positive) as assessed by local laboratory on primary or metastatic tumor (ISH positivity is defined as a ratio of 2.0 or greater for the number of
25 HER2 gene copies to the number of signals for CEP17, or for single probe tests, a HER2 gene count greater than 4).
5. Hormone receptor-positive defined as ER-positive and/or PgR-positive assessed locally as defined by institutional criteria.
6. At least one measurable lesion and/or non-measurable disease evaluable according to Response
30 Evaluation Criteria In Solid Tumors (RECIST) version 1.1.
7. ECOG performance status 0 or 1.
8. Left ventricular ejection fraction (LVEF) of at least 50%.
9. Life expectancy of at least 12 weeks.

Exclusion Criteria

- 35 1. Previous systemic non-hormonal anticancer therapy in the metastatic or locally advanced breast cancer setting.
2. Disease-free interval from completion of adjuvant or neo-adjuvant systemic non-hormonal

treatment to recurrence of within 6 months.

3. Previous approved or investigative anti-HER2 agents in any breast cancer treatment setting, except Trastuzumab and/or lapatinib in the neoadjuvant or adjuvant setting.
4. Disease progression while receiving Trastuzumab and/or lapatinib in the adjuvant setting.
- 5 5. History of persistent grade 2 or higher (NCI-CTC, Version 4.0) hematological toxicity resulting from previous adjuvant or neo-adjuvant therapy.
6. Radiographic evidence of central nervous system (CNS) metastases as assessed by CT or MRI.
7. Current peripheral neuropathy of grade 3 or higher (NCI-CTC, Version 4.0).
8. History of other malignancy within the last 5 years, except for carcinoma *in situ* of the cervix or
- 10 basal cell carcinoma.
9. Serious uncontrolled concomitant disease that would contraindicate the use of any of the investigational drugs used in this study or that would put the patient at high risk for treatment related complications.
10. Inadequate organ function, evidenced by the following laboratory results:
- 15 - Absolute neutrophil count $<1,500$ cells/mm³.
- Platelet count $<100,000$ cells/mm³.
- Hemoglobin <9 g/dL.
- Total bilirubin greater than the upper limit of normal (ULN) (unless the patient has documented Gilbert's syndrome).
- 20 - AST (SGOT) or ALT (SGPT) $>2.5 \times$ ULN.
- AST (SGOT) or ALT (SGPT) $>1.5 \times$ ULN with concurrent serum alkaline phosphatase $>2.5 \times$ ULN Serum alkaline phosphatase may be $>2.5 \times$ ULN only if bone metastases are present and AST (SGOT) and ALT (SGPT) $<1.5 \times$ ULN.
- Serum creatinine >2.0 mg/dL or $177 \mu\text{mol/L}$.
- 25 - International normalized ratio (INR) and activated partial thromboplastin time (aPTT) or partial thromboplastin time (PTT) $>1.5 \times$ ULN (unless on therapeutic coagulation).
11. Uncontrolled hypertension (systolic >150 mm Hg and/or diastolic >100 mm Hg) or clinically significant (i.e. active) cardiovascular disease: cerebrovascular accident (CVA)/stroke or myocardial infarction within 6 months prior to first study medication, unstable angina, congestive heart failure
- 30 (CHF) of New York Heart Association (NYHA) grade II or higher, or serious cardiac arrhythmia requiring medication.
12. Current known infection with HIV, HBV, or HCV.
13. Dyspnea at rest due to complications of advanced malignancy, or other disease requiring continuous oxygen therapy.
- 35 14. Major surgical procedure or significant traumatic injury within 28 days prior to randomization or anticipation of needed for major surgery during the course of study treatment.
15. Lack of physical integrity of the upper gastrointestinal tract, clinically significant malabsorption

syndrome, or inability to take oral medication.

16. Receipt of intravenous antibiotics for infection within 14 days prior to randomization.

17. Current chronic daily treatment with corticosteroids (dose of 10 mg/day methylprednisolone equivalent), excluding inhaled steroids.

5 18. Known hypersensitivity to any of the study medications or to excipients of recombinant human or humanized antibodies.

19. History of receiving any investigational treatment within 28 days prior to randomization.

20. Concurrent participation in any clinical trial.

10 **Arm A**

Pertuzumab (IV infusion)

Administered on Day 1 of the first treatment cycle as a loading dose of 840 mg, followed by 420 mg on Day 1 of each subsequent 3 weekly cycle.

15 Initial infusions of Pertuzumab will be administered over 90 (± 10) minutes and patients observed for at least 30 minutes from the end of infusion for infusion-related symptoms such as fever, chills etc. Interruption or slowing of the infusion may reduce such symptoms. If the infusion is well tolerated, subsequent infusions may be administered over 30 (± 10) minutes with patients observed for a further 30 minutes.

20 Trastuzumab (IV infusion administered after Pertuzumab) Day 1 of the first treatment cycle as a loading dose of 8 mg/kg, followed by 6 mg/kg on Day 1 of each subsequent 3 weekly cycle; to be administered in line with product labeling.

AI (oral)

Administered in line with product labeling (anastrozole: 1 mg once daily; letrozole: 2.5 mg once daily).

25 Induction chemotherapy

Patients receiving induction chemotherapy up to the first 18 weeks of the treatment period will receive a taxane (Docetaxel every 3 weeks or paclitaxel weekly), administered in line with the respective product labeling. Chemotherapy will be administered after the monoclonal antibody (Pertuzumab and/or Trastuzumab) infusions.

30 In patients receiving induction chemotherapy treatment with the AI will start after the chemotherapy induction phase.

Control arm - Arm B

Trastuzumab (IV infusion)

35 Day 1 of the first treatment cycle as a loading dose of 8 mg/kg, followed by 6 mg/kg on Day 1 of each subsequent 3 weekly cycle; to be administered in line with product labeling.

AI (oral)

Administered in line with product labeling (anastrozole: 1 mg once daily; letrozole: 2.5 mg once daily).

Induction chemotherapy

Same as for investigational arm.

5

Primary Efficacy Outcome

PFS (defined as the time from randomization until the first radiographically documented progression of disease or death from any cause, whichever occurs first).

Secondary Efficacy Outcome

- 10
- OS
 - ORR
 - CBR
 - Duration of response
 - Time to response

15

Safety

- Incidence and severity of adverse events (AEs) and serious adverse events (SAEs)
- Incidence of CHF
- LVEF over the course of the study
- 20 - Laboratory test abnormalities

It is anticipated that the the combination of Pertuzumab, Trastuzumab and AI will be safe and effective in the patient population and that the addition of Pertuzumab to Trastuzumab and an AI will extend progression-free survival (PFS) compared to Trastuzumab plus an AI without Pertuzumab.

EXAMPLE 9

Pertuzumab for Improving Overall Survival (OS) in Cancer Patients

25 **Background:** In the CLEOPATRA study in Example 3 above, 808 patients with HER2-positive first-line (1L) metastatic breast cancer (MBC) were randomized to treatment with Placebo+Trastuzumab+ Docetaxel (Pla+T+ D) or Pertuzumab+Trastuzumab+Docetaxel (P+T+D). The primary endpoint of independently reviewed progression-free survival was significantly improved with P+T+D vs Pla+T+D (hazard ratio (HR)=0.62; $P<0.0001$; medians, 18.5 vs 12.4 mths) (Example 3 above). This example includes a second interim overall survival (OS) analysis after

30 longer follow-up.

Methods: This interim overall survival (OS) analysis was performed applying the Lan-DeMets α -spending function with the O'Brien-Fleming (OBF) stopping boundary to maintain the overall Type I error at 5%. Based on the number of OS events observed, the OBF boundary for

35 statistical significance at this analysis was $P\leq 0.0138$. The log-rank test, stratified by prior treatment

status and geographic region, was used to compare OS between arms in the intention-to-treat population. The Kaplan-Meier approach was used to estimate the median OS in both arms; a stratified Cox proportional hazard model was used to estimate HR and 95% CIs. Subgroup analyses of OS were performed for the stratification factors and other key baseline characteristics.

- 5 **Results:** At the time of this analysis, median follow-up was 30 months and 267 deaths (69% of planned events for the final analysis) had occurred. The results showed a statistically significant improvement in OS in favor of P+T+D (HR=0.66; 95% confidence interval (CI), 0.52-0.84; $P=0.0008$). This HR represents a 34% reduction in the risk of death. The analysis achieved statistical significance and is therefore considered the confirmatory OS analysis. The median OS was 37.6 mths
10 in the Pla arm and has not yet been reached in the P arm. The treatment effect was generally consistent in predefined subgroups based on baseline variables and stratification factors, including: prior (neo)adjuvant therapy (HR=0.66; 95% CI, 0.46-0.94); no prior (neo)adjuvant therapy (HR=0.66; 95% CI, 0.47-0.93); prior (neo)adjuvant T (HR=0.68; 95% CI, 0.30-1.55); hormone receptor-negative disease (HR=0.57; 95% CI, 0.41-0.79); and hormone receptor-positive disease
15 (HR=0.73; 95% CI, 0.50-1.06). Kaplan-Meier estimates of OS rates show survival benefit with P+T+D at 1, 2, and 3 yrs.

Table 14: Overall Survival Benefit with Pertuzumab

	Pla+T+D	P+T+D	Δ
	Survival rates, %		
1 yr	89.0	94.4	5.4
2 yrs	69.4	80.7	11.3
3 yrs	50.4	65.8	15.4

- 20 The majority of pts received anti-cancer therapy after discontinuation of study treatment (64% Pla arm, 56% P arm). Subsequent therapy with HER2-directed agents (T, lapatinib, T emtansine) was balanced between arms. Causes of death remained unchanged from the first interim OS analysis, with the most common cause being progressive disease. Adverse events leading to death were rare and balanced between arms.

- 25 **Conclusions:** Treatment of patients with HER2-positive 1L MBC with P+T+D compared with Pla+T+D was associated with an improvement in OS, which was both statistically significant and clinically meaningful. These results show that combined HER2 blockade and chemotherapy using the P+T+D regimen can be considered a standard of care for patients with HER2-positive MBC in the 1L setting.

- 30 These data regarding OS can be included on the package insert with prescribing information regarding Pertuzumab in an article of manufacture as in Example 4 above, for example.

EXAMPLE 10**Pertuzumab and Trastuzumab with a Taxane as First-Line Therapy for Patients with HER2-Positive Advanced Breast Cancer (PERUSE)**

Background: Pertuzumab (P), a humanized monoclonal antibody, inhibits signaling downstream of HER2 by binding to the dimerization domain of the receptor and preventing heterodimerization with other HER family members. The epitope recognized by P is distinct from that bound by Trastuzumab (H) and so their complementary mechanisms of action result in a more comprehensive HER2 blockade. Data from the phase III trial CLEOPATRA showed significantly improved PFS in patients (pts) receiving P + H + docetaxel compared with H + docetaxel + placebo as first-line treatment for HER2-positive metastatic breast cancer (BC).

Trial design: This is a phase IIb, multicenter, open-label, single-arm study in pts with HER2-positive metastatic or locally recurrent BC who have not been treated with systemic nonhormonal anticancer therapy for metastatic cancer. Pts will receive, P: 840 mg initial dose, 420 mg q3w IV; H: 8 mg/kg initial dose, 6 mg/kg q3w IV; taxane: docetaxel, paclitaxel, or nab-paclitaxel according to local guidelines. Treatment will be administered until disease progression or unacceptable toxicity. A planned protocol amendment will allow hormone receptor-positive pts to receive endocrine therapy alongside P+H after completion of taxane therapy, in line with clinical practice.

Eligibility criteria: At baseline, pts must have an LVEF of $\geq 50\%$, an ECOG PS of 0, 1, or 2, a disease-free interval of ≥ 6 months, and must not have received prior anti-HER2 agents for the treatment of metastatic BC. Prior H and/or lapatinib in the (neo)adjuvant setting is permitted, providing there was no disease progression during treatment. Pts must not have experienced other malignancies within the last 5 yrs other than carcinoma *in situ* of the cervix or basal cell carcinoma. There must be no clinical or radiographic evidence of CNS metastases or clinically significant cardiovascular disease.

Specific aims: As H was not widely available in the (neo)adjuvant setting prior to CLEOPATRA recruitment, a relatively low proportion of pts in CLEOPATRA had previously received H. PERUSE will assess the safety and tolerability of P+H + choice of taxane as first-line therapy for pts with HER2-positive metastatic or locally advanced BC in a pt population likely to have experienced wider exposure to prior H therapy.

Statistical methods: The primary endpoints of the PERUSE study are safety and tolerability. Secondary endpoints include PFS, OS, ORR, CBR, duration of response, time to response and QoL. The final analysis will be performed when 1500 pts have been followed up for at least 12 months after the last pt receives last study treatment unless they have been lost to follow-up, withdrawn consent, or died, or if the study is prematurely terminated by the sponsor. Safety analyses are planned after enrollment of ~350, 700, and 1000 pts. Additionally, a data and safety monitoring board will review safety data after ~50 pts have been enrolled and then every 6 months.

It is anticipated that the Pertuzumab and Trastuzumab with a taxane will be effective as first-line therapy for patients with HER2-positive advanced breast cancer according to the protocol in this example.

EXAMPLE 11

Pertuzumab in Combination with Chemotherapy in Low HER3 Ovarian Cancer

Epithelial ovarian cancer, along with primary peritoneal, and fallopian tube carcinoma, is the fifth leading cause of cancer-related deaths in women in Europe (Bray et al. *Int. J. Cancer* 113:977-90 (2005)). Ovarian cancer is often not diagnosed until it has progressed to an advanced stage, at which point the standard treatment is surgical resection followed by chemotherapy. Although the addition of taxanes to platinum-based chemotherapy has resulted in approximately 80% of patients achieving complete response (CR), the disease recurs in most patients, and more than 50% of patients diagnosed with epithelial ovarian cancer eventually die from their disease (Du Bois et al. *Cancer* 115:1234-1244 (2009)). Following failure of platinum-based chemotherapy, there are few therapeutic options. Patients with platinum-sensitive disease (disease recurrence occurs more than 6 months after last cycle of platinum-based chemotherapy) are often retreated with platinum-based therapy and have a progression-free survival (PFS) of approximately 9-10; however, for patients with primary platinum-resistant disease, the prognosis is considerably worse. For these patients, re-treatment with platinum-based therapy or surgery is not reasonable, instead, patients with platinum-resistant are often treated with single-agent chemotherapy such as topotecan, pegylated liposomal doxorubicin (PLD), paclitaxel and gemcitabine.

Objective response rates for patients with platinum-resistant disease ranges between 10-20% while median progression free survival (PFS) ranges between 3.5-4 months. Platinum-resistant disease is not curable; the goals of treatment for these patients include palliation of symptoms, prolonged survival and improvements in quality of life (QoL). Overall, results from major clinical trials conducted over the last 20 years show that the median PFS for patients with advanced disease ranges between 16-23 months while median overall survival (OS) ranges between 31-65 months.

The majority of ovarian cancer cell lines and many ovarian cancer biopsy samples express all members of the HER family of receptors (Campiglio et al. *J. Cell Biochem* 73:522-32 (1999)). EGFR and HER2 have been studied the most extensively, and multiple agents targeting the receptor or associated intracellular tyrosine kinases have been tested.

In a recent study, quantitative HER2 protein analyses demonstrated that malignant ovarian tumors have significantly higher levels of HER2 compared with benign ovarian tumors and normal ovaries. Furthermore, a correlation between HER2 and HER3 protein levels has been seen (Steffensen et al. *Int J Oncol.* 33:195-204 (2008)). Studies in cell culture systems have shown that heregulin-activated HER3-HER2 heterodimers elicit the strongest proliferative and transformation responses of any possible receptor combination (Pinkas-Kramarski et al. *EMBO J.* 15:2452-67

(1996); Riese et al. *Mol Cell Biol* 15:5770-6 (1995). Erratum in: *Mol Cell Biol* 16:735 (1996)). The potency of these biologic responses is likely the result of the dual and efficient activation of the MAP kinase and PI3 kinase pathways. Furthermore, HER3 is the most potent activator of the PI3 kinase/AKT pathway (Olayioye et al. *EMBO J* 19:3159-67 (2000)). Studies in HER2-amplified breast cancer cell lines show that HER3 but not EGFR was critical for HER2 signaling, and that
5 HER3 inhibited growth in three-dimensional culture and induced rapid tumor regression of *in vivo* xenografts (Lee-Hoeflich et al. *Cancer Res* 68:5878-87 (2008)).

Additionally, HER3 expression has been implicated as a possible risk factor in ovarian cancer (Tanner et al. *J Clin Oncol* 24:4317-23 (2006)).

10 In a Phase II multicenter trial (TOC2689g) in patients with advanced ovarian cancer that recurred after treatment with or were refractory to platinum-based chemotherapy, patients who were enrolled in Cohort 1 (n=61) received a loading dose of 840 mg Pertuzumab, followed by 420 mg Pertuzumab on Day 1 of each 3-week cycle, and patients in Cohort 2 (n=62) received 1050 mg Pertuzumab on Day 1 of each 3-week cycle. Similar outcomes were observed in both cohorts in
15 terms of overall response rate and median PFS. Eight patients (4 in each cohort) had evidence of stable disease (SD) lasting at least 6 months. Median PFS and OS were 6.6 weeks and 52.7 weeks, respectively, for the overall population.

The results of this study led to two randomized Phase II trials in platinum-sensitive and platinum-resistant populations. In Study TOC3258g, the efficacy and safety of gemcitabine +
20 Pertuzumab versus gemcitabine + placebo were evaluated in patients with advanced ovarian, primary peritoneal, or fallopian tube cancer that was resistant to platinum-based chemotherapy (Amler et al. *J Clin Oncol* 26:5552 (2008)). The study allowed patients to cross over to receive Pertuzumab at the time of disease progression. There was a median PFS of 2.6 months in the gemcitabine + placebo arm and 2.9 months in the gemcitabine + Pertuzumab arm. Median OS was similar between the treatment
25 arms. Of the most common adverse events (AEs), those increased (by at least 6 patients) in the Pertuzumab-treated cohort included fatigue, nausea, diarrhea, back pain, dyspepsia, stomatitis, headache, epistaxis, rhinorrhea, rash, and Grade 3-4 neutropenia.

In Study BO17931, 149 patients with ovarian cancer who experienced a recurrence 6 months after a platinum-based therapy were randomized to receive a combination of paclitaxel and
30 carboplatin or gemcitabine with or without Pertuzumab. After 6 treatment cycles, chemotherapy was discontinued, and patients in the chemotherapy + Pertuzumab arm continued to receive Pertuzumab alone for up to 11 additional cycles (total of 17 cycles of Pertuzumab). There were no significant differences in the PFS or OS for the overall group. Median PFS was 34.1 weeks for the chemotherapy + Pertuzumab group versus 31.3 for the chemotherapy alone group; however, an
35 exploratory subset analyses of HER3 mRNA expression with a treatment-free interval of 6-12 months indicated a trend toward clinical benefit in patients who express high levels of HER3 mRNA (Kaye et al. *J Clin Oncol* 26:5520 (2008)).

Archival tissue samples from patients enrolled in both randomized Phase II studies were examined by quantitative reverse transcriptase polymerase chain reaction (qRT-PCR) for mRNA expression levels of the HER receptors EGFR, HER2, HER3, and two HER ligands: amphiregulin and betacellulin.

5 Only tumor HER3 mRNA expression was associated with a significant difference in PFS. For patients who achieved a clinical response, PRs were observed in 9 patients on the gemcitabine + Pertuzumab arm and 3 on the gemcitabine + placebo arm. Six of the gemcitabine + Pertuzumab patients with PRs had tumor HER3 mRNA levels lower than the median level. In contrast, no patients
10 in the gemcitabine + placebo arm whose tumor HER3 mRNA levels were lower than the median level of the study population experienced a PR. An additional 6 patients achieved PRs, and all of these patients had tumor HER3 mRNA levels at or above the median level of the study population. Of these patients, 3 received gemcitabine + Pertuzumab and 3 received gemcitabine + placebo, suggesting no effect of Pertuzumab in this population.

 Patients with low HER3 mRNA expression (lower than the median level of the study
15 population) demonstrated a PFS hazard ratio (HR) of 0.32 in contrast to 1.68 for patients with HER3 mRNA expression greater than or equal to the median level; i.e. the effect of adding Pertuzumab trended in the opposite direction. No significant benefit was detected in OS for patients with low HER3 mRNA expression; however, a trend toward greater OS was observed in patients receiving Pertuzumab. The OS for patients expressing high HER3 mRNA expression demonstrated an HR of
20 1.59.

 To assess the prognostic value, HER3 mRNA expression was correlated with PFS and OS for patients in the gemcitabine + placebo arm. Median PFS was 1.4 months for patients with low HER3 mRNA expression (n = 35), compared with 5.5 months for patients with high HER3 mRNA expression (n = 24). Similarly, median OS for patients with low HER3 mRNA expression was 8.4
25 months, compared with 18.2 months for patients with high HER3 mRNA expression.

 In Study BO17931, in patients with low HER3 mRNA expression (lower than the median level of this study population), no treatment effect was seen. However, in an exploratory analysis of patients with a treatment-free interval of 6-12 months, there was a trend toward benefit for the combination of chemotherapy with Pertuzumab in terms of PFS.

30

Overview of this Study

 This is a multicenter trial with two parts; a non-randomized safety run-in Part 1 and a randomized, double-blind Part 2.

Part 1 will be performed to assess safety and tolerability of Pertuzumab in a new
35 combination with two chemotherapeutic agents (topotecan or paclitaxel). Part 2 of the trial is a randomized, double-blind, placebo controlled, two-arm, multicenter, prospective trial of Pertuzumab in combination with chemotherapy (topotecan, paclitaxel, or gemcitabine). Patients will receive trial

medication until disease progression as per the Response Evaluation Criteria in Solid Tumors (RECIST) version 1.1, disease progression according to the Gynecologic Cancer Intergroup (GCIg) criteria of CA-125 assessable disease, unacceptable toxicity, withdrawal of consent, or death. PFS will be assessed in Part 1 of the trial, but due to small number of patients and PFS events per cohort, results will be descriptive only. The trial design for Part 1 of the study is provided in Figure 30.

In **Part 2** of the trial, patients will be randomized in a 1:1 ratio to receive either:

- **Arm A:** Pertuzumab in combination with chemotherapy (topotecan, paclitaxel, or gemcitabine), or
- **Arm B:** Pertuzumab-placebo plus chemotherapy (topotecan, paclitaxel, or gemcitabine).

The allocation of study medication will be double-blind with respect to whether the patient receives Pertuzumab or Pertuzumab-placebo. The chemotherapy agent allocated will be at the discretion of the investigator.

Stratification factors for Part 2 of the trial will be:

- Selected chemotherapy cohort (topotecan vs. paclitaxel vs. gemcitabine).
- Previous anti-angiogenic therapy (yes vs. no). If a patient has previously participated in a blinded trial with an anti-angiogenic agent, the patient will be enrolled in the same stratum with patients known to have previously received an anti-angiogenic agent.
- Treatment-free interval (TFI) since platinum therapy (strictly less than 3 months vs. 3 to 6 months inclusive, prior to first study treatment).

The trial design for Part 2 of the study is provided in Figure 31.

Primary Objectives of the Study:

Part 1: The primary objective for Part 1 of this study is to determine the safety and tolerability of Pertuzumab in combination with either topotecan or paclitaxel.

Part 2: The primary objective for Part 2 of this study is to determine if Pertuzumab plus chemotherapy is superior to placebo plus chemotherapy as measured by PFS.

Secondary Objectives of the Study:

Part 1: The secondary objective for Part 1 of this study is to evaluate descriptively the PFS of Pertuzumab in combination with either topotecan or paclitaxel.

Part 2: The secondary objectives for Part 2 of this study are to determine if Pertuzumab plus chemotherapy is superior to placebo plus chemotherapy with respect to:

- OS.
- Objective response rate.
- Biological progression-free interval (PFI_{BIO}).
- Safety and tolerability.
- QoL.

Efficacy Outcome Measures

The following efficacy outcome measures will be measured in **Part 1** of the trial:

- PFS, which is defined as the time from randomization into Part 1 of the trial, until disease progression per RECIST version 1.1 or according to GCIG criteria in CA-125 assessable disease or death from any cause, whichever occurs first.

The efficacy outcome measures for **Part 2** of the trial are as follows:

- PFS, which is defined as the time from randomization into Part 2 of the trial, until disease progression per RECIST version 1.1 or according to GCIG criteria in CA-125 assessable disease or death from any cause, whichever occurs first.
- OS, defined as the time from randomization into Part 2 of the trial until death from any cause.
- Objective response rate (ORR), which will be based on RECIST version 1.1 and assessed by the best (confirmed) overall response (BOR); defined as the best response recorded from the start of the treatment in Part 2 of the trial until disease progression/recurrence (taking as reference for PD, the smallest measurements recorded since the treatment started in Part 2 of the trial). Patients need to have two consecutive assessments of partial response (PR) or complete response (CR) to be a responder. PR or CR has to be confirmed by 2 consecutive tumor evaluations spaced at least 4 weeks apart. Only patients with measurable disease at baseline will be included in the analysis of objective response.
- Patients who have a response as per RECIST version 1.1 and using the 50% response criteria for CA-125 are defined as responders, whereas patients who only have response as defined per RECIST are defined as RECIST responders. Patients who do not have a response as per RECIST, but have a response defined using the 50% response criteria for CA-125 are defined as CA-125 responders.
- PFI_{BIO}, defined on the basis of a progressive serial elevation of serum CA-125 (assessed according to the CGIG criteria) as the time from the date of randomization into Part 2 of the trial to first documented increase in CA-125 levels to: two times the upper limit of normal (for patients with normal pretreatment CA-125 or elevated pretreatment CA-125 and initial normalization on-treatment), or two times the nadir value (for patients with elevated baseline CA-125 that did not normalize on-treatment).

Safety Outcome Measures

- In Part 1 of the study safety and tolerability will be assessed after all patients have received 3 cycles of treatment.

In addition, safety outcome measures for this study will be assessed in both Parts 1 and 2 of

the study, and are as follows:

- Incidence, nature, and severity of all AEs, serious adverse events (SAEs), AEs with NCI-CTCAE version 4.0 Grades ≥ 3 , and AEs that caused premature withdrawal from study medication.
- Premature withdrawal from the study and study treatment.
- 5 - Cardiac disorders / Incidence of congestive heart failure
- Laboratory test abnormalities.
- Left ventricular ejection fraction

Inclusion Criteria

- 10 1. Female patients aged 18 years or older.
2. Low HER3 mRNA expression levels (concentration ratio equal or lower than 2.81, as assessed by qRT-PCR on a COBAS z480® instrument).
3. Histologically or cytologically confirmed and documented epithelial ovarian cancer that is platinum-resistant or refractory (defined as progression within 6 months from completion of a
- 15 minimum of 4 platinum therapy cycles or progression during platinum therapy).
4. At least one measurable lesion and/or non-measurable disease according to RECIST version 1.1, or cancer antigen-125 (CA-125) assessable disease according to Gynecologic Center Intergroup (GCIG) criteria. The following histological types are eligible:
 - Adenocarcinoma not otherwise specified.
 - 20 - Clear cell adenocarcinoma.
 - Endometrioid adenocarcinoma.
 - Malignant Brenner's tumor.
 - Mixed epithelial carcinoma including malignant mixed Müllerian tumors.
 - Mucinous adenocarcinoma.
 - 25 - Serous adenocarcinoma.
 - Transitional cell carcinoma.
 - Undifferentiated carcinoma.
5. Eastern Cooperative Oncology Group (ECOG) performance status 0 to 2.
6. LVEF greater than or equal to 55%.

30

Pertuzumab Dosage and Administration

- Pertuzumab and Pertuzumab-placebo will be administered as an intravenous infusion on Day 1 of the first treatment cycle as a loading dose of 840, followed by 420 mg on Day 1 of each subsequent 3-weekly cycle. The initial infusion of Pertuzumab/Pertuzumab-placebo will be
- 35 administered over 60 minutes followed by a 60-minute observational period in a seated position if the infusion is well tolerated, subsequent infusions may be given over 30 minutes, followed by a 30-minute observational period, after which the chemotherapy agent will be administered. Pre-

medication should be implemented according to local practices and the chosen chemotherapy.

Topotecan Dosage and Administration

Topotecan should be administered 1.25 mg/m² as a 30-minute intravenous infusion daily on
5 Days 1–5 every 3 weeks, as per the directions in the summary of product characteristics.

Paclitaxel Dosage and Administration

Paclitaxel should be administered 80 mg/m² as a 1-hour i.v. infusion on Days 1, 8, 15 and
22. Pharmacists should follow the summary of product characteristics for information regarding the
10 preparation and administration of the 80 mg/m² dose.

Gemcitabine Dosage and Administration

Gemcitabine (Part 2 of the study only) should be administered 1000 mg/m² as a 30-minute
intravenous infusion on Days 1 and 8 every 3 weeks as per the directions described in the summary
15 of product characteristics.

HER3 mRNA Expression

Patients will be asked to specifically consent to the collection and testing of primary tumor
tissue samples to assess HER3 mRNA level, including mRNA and protein levels of other HER
20 family receptors e.g. HER2, before they provide consent to participate in the trial. Only patients who
have tumors expressing low levels of HER3 mRNA will be eligible to participate in the trial.

During the initial screening for HER3 mRNA levels, other receptors of the
HER family (e.g. EGFR, HER2, or HER4) will be assessed at the mRNA level and/or protein level in
parallel to the HER3 assessment, in order to obtain a more complete picture of the status of HER
25 family receptors by mRNA level.

The cut-off defined for study eligibility is defined as a concentration ratio of ≤ 2.81 as
assessed by qRT-PCR on a COBAS z480® instrument using the “COBAS® HER2 & HER3 (qRT-
PCR) mRNA expression assay” provided by Roche Molecular Diagnostics. The rationale for cut off
definition is based on a cut off modeling in previous studies as well as on a transformation function
30 that had to be introduced since the assay was switched to a new instrument; the COBAS z480®. It is
anticipated that 40-50% of screened patients will have HER3 mRNA levels below the cutoff of 2.81
and that 30% of patients expressing low levels of HER3 mRNA will be ineligible for enrollment
owing to other inclusion/exclusion criteria.

Submission of a formalin-fixed, paraffin-embedded tumor specimen of the primary tumor
35 from the original surgery will be required for all patients prior to screening; cytology specimens are
not acceptable replacements. Patients will be assessed for HER3 mRNA expression level, as well as
mRNA expression and protein expression levels of other HER family receptors by the use of a qRT-

PCR assay and IHC. Such assessment of HER receptor mRNA/protein expression will occur after obtaining the patient's informed consent at any time after the primary surgery and prior to screening.

It is anticipated that Pertuzumab in combination with topotecan or paclitaxel will be safe and effective in patients with epithelial ovarian, primary peritoneal, or fallopian tube cancer.

- 5 In addition, it is anticipated that Pertuzumab plus chemotherapy (topotecan, paclitaxel, or gemcitabine) will be superior to placebo plus chemotherapy in patients with epithelial ovarian, primary peritoneal, or fallopian tube cancer where efficacy is measured by PFS.

WHAT IS CLAIMED IS:

1. A method for extending progression free survival in a HER2-positive breast cancer patient population by 6 months or more comprising administering Pertuzumab, Trastuzumab and chemotherapy to the patients in the population.
2. The method of claim 1 which results in an objective response rate of 80% or more in the patients in the population.
3. The method of claim 1 or claim 2 wherein the chemotherapy comprises taxane.
4. The method of claim 3 wherein the taxane is Docetaxel.
5. The method of any one of claims 1 to 4 wherein the breast cancer is metastatic or locally recurrent, unresectable breast cancer, or *de novo* Stage IV disease.
6. The method of any one of claims 1 to 5 wherein the patients in the population have not received previous treatment or have relapsed after adjuvant therapy.
7. The method of any one of claims 1 to 6 wherein the HER2-positive breast cancer is defined as immunohistochemistry (IHC) 3+ and/or fluorescence *in situ* hybridization (FISH) amplification ratio ≥ 2.0 .
8. The method of any one of claims 1 to 7 wherein the patients in the population have a left ventricular ejection fraction (LVEF) of $\geq 50\%$ at baseline.
9. The method of any one of claims 1 to 8 wherein the patients in the population have an Eastern Cooperative Oncology Group performance status (ECOG PS) of 0 or 1.
10. A method of combining two HER2 antibodies to treat HER2-positive cancer without increasing cardiac toxicity in a HER2-positive cancer patient population, comprising administering Pertuzumab, Trastuzumab, and chemotherapy to the patients in the population.
11. The method of claim 10 wherein cardiac toxicity in the patient population is monitored for incidence of symptomatic left ventricular systolic dysfunction (LVSD) or congestive heart failure (CHF), or decrease in left ventricular ejection fraction (LVEF).
12. The method of claim 10 or claim 11 wherein the HER2-positive cancer is breast cancer.
13. The method of any one of claims 10 to 12 wherein the breast cancer is metastatic or locally recurrent, unresectable breast cancer, or *de novo* Stage IV disease.

14. An article of manufacture comprising a vial with Pertuzumab therein and a package insert, wherein the package insert provides the safety data in Table 3 or Table 4.
15. The article of manufacture of claim 14 wherein the package insert provides the efficacy data in Table 2, Table 5, Figure 8, or Figure 10.
16. The article of manufacture of claim 14 or claim 15 wherein the vial is a single-dose vial containing about 420mg of Pertuzumab.
17. A method for making an article of manufacture comprising packaging together a vial with Pertuzumab therein and a package insert, wherein the package insert provides the safety data in Table 3 or Table 4.
18. A method of ensuring safe and effective use of Pertuzumab comprising packaging together a vial with Pertuzumab therein and a package insert, wherein the package insert provides the safety data in Table 3 or Table 4 and the efficacy data in Table 2, Table 5, Figure 8, or Figure 10.
19. A method of treating early-stage HER2-positive breast cancer comprising administering Pertuzumab, Trastuzumab, and chemotherapy to a patient with the breast cancer, wherein the chemotherapy comprises anthracycline-based chemotherapy, or carboplatin-based chemotherapy.
20. The method of claim 19 wherein chemotherapy comprises anthracycline-based chemotherapy which comprises 5-FU, epirubicin, and cyclophosphamide (FEC).
21. The method of claim 20 wherein the Pertuzumab is administered concurrently with the anthracycline-based chemotherapy.
22. The method of claim 19 wherein the chemotherapy comprises carboplatin-based chemotherapy which comprises Docetaxel and Carboplatin.
23. The method of claim 22 wherein the Pertuzumab is administered concurrently with the carboplatin-based chemotherapy.
24. The method of any one of claims 19 to 23 wherein Pertuzumab administration does not increase cardiac toxicity relative to the treatment without Pertuzumab.
25. The method of any one of claims 19 to 24 which comprises neoadjuvant or adjuvant therapy.
26. A method of treating HER2-positive cancer in a patient comprising co-administering a mixture of Pertuzumab and Trastuzumab from the same intravenous bag to the patient.

27. The method of claim 26 which further comprises administering chemotherapy to the patient.
28. An intravenous (IV) bag containing a stable mixture of Pertuzumab and Trastuzumab suitable for administration to a cancer patient.
29. The IV bag of claim 28 wherein the mixture is in saline solution.
30. The IV bag of claim 29 wherein the saline solution comprises about 0.9% NaCl or about 0.45% NaCl.
31. The IV bag of any one of claims 28 to 30 which is a 250mL 0.9% saline polyolefin or polyvinyl chloride infusion bag.
32. The IV bag of any one of claims 28 to 31 which contains a mixture of about 420mg or about 840mg of Pertuzumab and from about 200mg to about 1000mg of Trastuzumab.
33. The IV bag of any one of claims 28 to 32 wherein the mixture is stable for up to 24 hours at 5°C or 30°C.
34. The IV bag of any one of claims 28 to 33 wherein stability has been evaluated by an assay selected from the group consisting of: color, appearance and clarity (CAC), concentration and turbidity analysis, particulate analysis, size exclusion chromatography (SEC), ion-exchange chromatography (IEC), capillary zone electrophoresis (CZE), image capillary isoelectric focusing (iCIEF), and potency assay.
35. A method of treating HER2-positive gastric cancer in a human subject comprising administering Pertuzumab, Trastuzumab, and chemotherapy to the subject with HER2-positive gastric cancer.
36. The method of claim 35 wherein the gastric cancer comprises non-resectable locally advanced gastric cancer, or metastatic gastric cancer, or advanced, post-operatively recurrent gastric cancer, which may not be amenable to curative therapy by known methods, or adenocarcinoma of the stomach or gastroesophageal junction.
37. The method of claim 35 or claim 36 wherein the patient has not received prior anti-cancer treatment for metastatic gastric cancer, has an Eastern Cooperative Oncology Group Performance Status Scale (ECOG PS) of 0-1, or has a HER2-positive status of IHC 3+ or IHC2+/ISH+.
38. The method of any one of claims 35 to 37 wherein the chemotherapy comprises platinum (cisplatin) and/or fluoropyrimidine (capecitabine or 5-fluorouracil (5-FU)).

39. The method of claim 38 wherein Pertuzumab, Trastuzumab, cisplatin, and capecitabine or 5-FU are administered.
40. The method of any one of claims 35 to 39 wherein Pertuzumab is administered at a dose of 840 mg in all treatment cycles.
41. The method of any one of claims 35 to 40 which improves overall survival (OS) relative to a patient treated with Trastuzumab and the chemotherapy only, or improves progression free survival (PFS) or response rate (RR) relative to treatment with Trastuzumab and chemotherapy only.
42. A method of treating gastric cancer in a human subject comprising administering Pertuzumab to the subject with gastric cancer, wherein Pertuzumab is administered at a dose of 840 mg in all treatment cycles.
43. The method of claim 42 wherein Pertuzumab is administered at the dose of 840 mg for six treatment cycles.
44. The method of claim 42 or claim 43 which maintains Pertuzumab trough levels above about 20 µg/mL in the subject.
45. The method of any one of claims 42 to 44 further comprising administering Trastuzumab and chemotherapy to the subject.
46. A method of treating HER2-positive non-resectable or metastatic adenocarcinoma of the stomach or gastroesophageal junction in a human patient who did not receive prior chemotherapy for metastatic disease, except prior adjuvant or neoadjuvant therapy completed more than six months before the current treatment, comprising administering Pertuzumab, Trastuzumab, cisplatin, and capecitabine or fluorouracil (5-FU) to the patient in an amount to improve progression free survival (PFS) and/or overall survival (OS), wherein the patient has an Eastern Cooperative Oncology Group Performance Status Scale (ECOG PS) of 0-1.
47. A method of improving progression free survival (PFS) in a human patient with HER2-positive non-resectable or metastatic adenocarcinoma of the stomach or gastroesophageal junction comprising administering Pertuzumab to the patient in combination with Trastuzumab and chemotherapy.
48. A method of treating HER2-positive breast cancer in a patient comprising administering Pertuzumab, Trastuzumab and vinorelbine to the patient.

49. The method of claim 48 wherein the Pertuzumab and Trastuzumab are co-administered to the patient from a single intravenous bag.
50. The method of claim 48 or claim 49 wherein the breast cancer is metastatic or locally advanced.
51. The method of any one of claims 48 to 50 wherein the patient has not previously received systemic non-hormonal anticancer therapy in the metastatic setting.
52. A method of treating HER2-positive breast cancer in a patient comprising administering Pertuzumab, Trastuzumab, and aromatase inhibitor to the patient.
53. The method of claim 52 wherein the aromatase inhibitor is anastrozole or letrozole.
54. The method of claim 52 or claim 53 wherein the breast cancer is hormone receptor-positive advanced breast cancer.
55. The method of claim 54 wherein the hormone receptor is estrogen receptor (ER) and/or progesterone receptor (PgR).
56. The method of any one of claims 52 to 55 wherein the patient has not previously received systemic nonhormonal anticancer therapy in the metastatic setting.
57. The method of any one of claims 52 to 56 wherein the patient receives induction chemotherapy.
58. The method of claim 57 wherein the induction chemotherapy comprises a taxane.
59. The method of claim 17 or claim 18 wherein the package insert further comprises the warning box in Example 4.
60. A method of treating a cancer patient comprising administering to the patient an initial dose of 840mg of Pertuzumab followed every 3 weeks thereafter by a dose of 420mg of Pertuzumab, and further comprising re-administering an 840mg dose of Pertuzumab to the patient if the time between two sequential 420mg doses is 6 weeks or more.
61. The method of claim 60 further comprising administering 420mg of Pertuzumab every 3 weeks after the re-administered 840mg dose.
62. The method of claim 60 or claim 61 wherein the cancer patient has HER2-positive breast cancer.

63. The method of any one of claims 1 to 13 which reduces the risk of death by about 34% or more relative to a patient treated with Trastuzumab and the chemotherapy.
64. The method of claim 17 or claim 18 wherein the package insert further provides the Overall Survival (OS) efficacy data in Example 9 or Table 14.
65. A method for treating HER2-positive metastatic or locally recurrent breast cancer in a patient comprising administering Pertuzumab, Trastuzumab and taxoid to the patient, wherein the patient has been previously treated with a Trastuzumab and/or lapatinib as adjuvant or neoadjuvant therapy.
66. The method of claim 65 wherein the taxoid is Docetaxel, Paclitaxel, or *nab*-paclitaxel.
67. The method of claim 65 wherein the taxoid is Paclitaxel or *nab*-paclitaxel.
68. The method of any one of claims 65 to 67 wherein the patient has been previously treated with Trastuzumab as neoadjuvant therapy.
69. A method for treating low HER3 ovarian, primary peritoneal, or fallopian tube cancer in a patient comprising administering Pertuzumab and chemotherapy to the patient, wherein the low HER3 cancer expresses HER3 mRNA at a concentration ratio equal or lower than about 2.81 as assessed by polymerase chain reaction (PCR).
70. The method of claim 69 wherein the chemotherapy comprises gemcitabine, carboplatin, paclitaxel, docetaxel, topotecan, or pegylated liposomal doxorubicin (PLD).
71. The method of claim 70 wherein the chemotherapy comprises paclitaxel or topotecan.
72. The method of any one of claims 69 to 71 wherein the cancer is epithelial ovarian cancer that is platinum-resistant or platinum-refractory.




FIG. 1

2 / 34

Variable Light

	10	20	30	40
2C4	DTVMTQSHKIMSTSVGDRVSITC	[KASQDVSIGVA]	WYQQR	P
	** **** *	*		*
574	DIQMTQSPSSLSASVGDRVTITC	[KASQDVSIGVA]	WYQQK	P
		* ** ***		
hum κI	DIQMTQSPSSLSASVGDRVTITC	[RASQISNYLA]	WYQQK	P

	50	60	0	80
2C4	GQSPKLLIY [SASYRYT]	GVPDRFTGSGSGTDFTLT	ISSVQA	
	** *	*	*	*
574	GKAPKLLIY [SASYRYT]	GVPSRFSGSGSGTDFTLT	ISLQP	
	* *****			
hum κI	GKAPKLLIY [AASSLES]	GVPSRFSGSGSGTDFTLT	ISLQP	

	90	100	
2C4	EDLAVYYC [QQYIYPYT]	FGGGTKLEIK (SEQ ID NO:5)	
	* *	* *	
574	EDFATYYC [QQYIYPYT]	FGQGTKVEIK (SEQ ID NO:7)	
	*** *		
hum κI	EDFATYYC [QQYNSLPWT]	FGQGTKVEIK (SEQ ID NO:9)	

FIG. 2A**Variable Heavy**

	10	20	30	40
2C4	EVQLQQSGPELVKPGTSVKISCKAS	[GFTFTDYM]	WVKQS	
	** ** * **** *		*	*
574	EVQLVESGGGLVQPGGSLRLS	CAAS [GFTFTDYM]	WVRQA	
		*** *		
hum III	EVQLVESGGGLVQPGGSLRLS	CAAS [GFTFSSYAMS]	WVRQA	

	50 a	60	70	80
2C4	HGKSLEWIG [DVNPNSGGSIYNQRFKG]	KASLTVD	RSSRIVYM	
	* * **	*** *	***** *	
574	PGKGLEWVA [DVNPNSGGSIYNQRFKG]	RFTLSVDR	SKNTLYL	
	***** *** ****	* * *		
hum III	PGKGLEWVA [VISGDGGSTYYADSVKG]	RFTISRDN	SKNTLYL	

	abc	90	100ab	110
2C4	ELRSLTFEDTAVYYCAR	[NLGPSFYFDY]	WGQGTTLTVSS (SEQ ID NO:6)	
	*** **		**	
574	QMNSLRAEDTAVYYCAR	[NLGPSFYFDY]	WGQGTTLTVSS (SEQ ID NO:8)	

hum III	QMNSLRAEDTAVYYCAR	[GRVGYSLYDY]	WGQGTTLTVSS (SEQ ID NO:10)	

FIG. 2B

3 / 34

Amino Acid Sequence for Pertuzumab Light Chain

1 10 20 30 40 50 60
 DIQMTQSPSSLSASVGDRVTITCKASQDVSIGVAWYQOKPGKAPKLLIYSASRYRTGVPS
 70 80 90 100 110 120
 RFSGSGSGTDFTLTISSLQPEDFATYYCQQYYIYPYTFGQGTKVEIKRTVAAPSVFIFPP
 130 140 150 160 170 180
 SDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLT
 190 200 210
 LSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 11)

FIG. 3A**Amino Acid Sequence for Pertuzumab Heavy Chain**

1 10 20 30 40 50 60
 EVQLVESGGGLVQPGGSLRLSCAASGFTFTDYMWDWVRQAPGKGLEWVADVNPNSGGSIY
 70 80 90 100 110 120
 NQRFKGRFTLSVDRSKNTLYLQMNSLRAEDTAVYYCARNLGPSFYFDYWGQGTLLTVSSA
 130 140 150 160 170 180
 STKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSG
 190 200 210 220 230 240
 LYSLSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGP
 250 260 270 280 290 300
 SVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS
 310 320 330 340 350 360
 TYRVSVELTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM
 370 380 390 400 410 420
 TKNQVSLTCLVKGFIYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQ
 430 440 448
 QGNVFSCSVMEALHNHYTQKSLSLSPG (SEQ ID NO: 12)

FIG. 3B

Trastuzumab Light Chain

1 DIQMTQSSSLASVGGDRVTITCRASQDVNTAVAWYQQKPGKAPK 45
46 LLISASFLYSGVPSRFSGSRSGTDFTLTISSLQPEDFATYYCQQ 90
91 HYTTPPTFGQGTKVEIK|RTVAAPSVFFIFPPSDEQLKSGTASVVC 135
136 LNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSYSSLSTLT 180
181 LSKADYEKHKVYACEVTHQGLSSSPVTKSFNRGEC (SEQ ID NO: 13) 210 214

4 / 34

FIG. 4A

5 / 34

Trastuzumab Heavy Chain

1 EVQLVESGGGLVQPGGSLRLSCAASGFNISKDTYIHWRQAPGKGL 45
46 EWVARIYPTNGYTRYADSVKGRFTISADTSKNTAYLQMNSLRAED 90
91 TAVYYCSRWG G D G F Y A M D Y W G Q G T L V T V S S A S T K G P S V F F L A P S S 135
136 K S T S G G T A A L G C L V K D Y F F E P V T V S W N S G A L T S G V H T F P A V L Q S S 180
181 G L Y S L S S V T V P S S S L G T Q T Y I C N V N H K P S N T K V D K K V E P K S C D K 225
226 T H T C P P C P A P E L L G G P S V F L F P P K P K D T L M I S R T P E V T C V V D V S 270
271 H E D P E V K F N W Y V D G V E V H N A K T K P R E E Q Y N S T Y R V S V L T V L H Q D 315
316 W L N G K E Y K C K V S N K A L P A P I E K T I S K A K G Q P R E P Q V Y T L P P S R E E 360
361 M T K N Q V S L T C L V K G F Y P S D I A V E W E S N G Q P E N N Y K T T P P V L D S D G 405
406 S F F L Y S K L T V D K S R W Q Q G N V F S C S V M H E A L H N H Y T Q K S L S L S P G 449

(SEQ ID NO: 14)

FIG. 4B

FIG. 5A

7/34



Pertuzumab Variant Heavy Chain

1 E V Q L V E S G G L V Q P G G S L R L S C A A S G F T F T D Y T M D W V R Q A P G K G L 45
46 E W V A D V N P N S G G S I Y N Q R F K G R F T L S V D R S K N T L Y L Q M N S L R A E D 90
91 T A V Y Y C A R N L G P S F Y F D Y W G Q G T L V T V S S A S T K G P S V F P L A P S S K 135
136 S T S G G T A A L G C L V K D Y F P E P V T V S W N S G A L T S G V H T F P A V L Q S S G 180
181 L Y S L S S V V T V P S S L G T Q T Y I C N V N H K P S N T K V D K K V E P K S C D K T 225
226 H T C P C P A P E L L G G P S V F L F P P K P K D T L M I S R T P E V T C V V D V S H 270
271 E D P E V K F N W Y V D G V E V H N A K T K P R E E Q Y N S T Y R V V S V L T V L H Q D W 315
316 L N G K E Y K C K V S N K A L P A P I E K T I S K A K G Q P R E P Q V Y T L P P S R E E M 360
361 T K N Q V S L T C L V K G F Y P S D I A V E W E S N G Q P E N N Y K T T P P V L D S D G S 405
406 F F L Y S K L T V D K S R W Q Q G N V F S C S V M H E A L H N H Y T Q K S L S L S P G K 449

(SEQ ID NO: 16)

FIG. 5B

8 / 34

Study Schema**FIG. 6****Schema HER2 Positive, Neoadjuvant Breast Cancer, Patients with Low Cardiac Risk Factors****FIG. 11**

Enrollment, Intent-to-Treat and Safety Populations, and Patient Withdrawals




FIG. 7




FIG. 8

11 / 34




FIG. 9







FIG. 10



14 / 34

**FIG. 13****FIG. 14**

Pertuzumab SEC Profile of Pertuzumab/Trastuzumab Mixture in Infusion Bags




FIG. 15

Trastuzumab SEC Profile of Pertuzumab/Trastuzumab Mixture in Infusion Bags




FIG. 16

Pertuzumab IEC Profile of Pertuzumab/Trastuzumab Mixture in Infusion Bags




FIG. 17

Trastuzumab IEC Profile of Pertuzumab/Trastuzumab Mixture in Infusion Bags




FIG. 18

CE-SDS LIF Non-reduced Profile of Pertuzumab/Trastuzumab Mixture in IV Infusion Bags




FIG. 19

CE-SDS LIF Reduced Profile of Pertuzumab/Trastuzumab Mixture in IV Infusion Bags




FIG. 20

CZE of Pertuzumab/Trastuzumab Mixture in IV Infusion Bags




FIG. 21

iCIEF of Pertuzumab/Trastuzumab Mixture in IV Infusion Bags





FIG. 22

23 / 34

**Potency Dose Response Curves of Pertuzumab/Trastuzumab Mixture,
Pertuzumab Alone, and Trastuzumab Alone in Infusion Bags**

**FIG. 23**

Pertuzumab SEC Profile of Pertuzumab/Trastuzumab Mixture in Infusion Bags




FIG. 24





FIG. 25

26 / 34

Pertuzumab IEC Profile of Pertuzumab/Trastuzumab Mixture in Infusion Bags

**FIG. 26**

Trastuzumab IEC Profile of Pertuzumab/Trastuzumab Mixture in Infusion Bags




FIG. 27




FIG. 29






FIG. 28

29 / 34

**FIG. 30****FIG. 31**

Samples Taken and Time Points

Cycle	Treatment Period									
	1			2			3			6
Day of cycle	1	8	15	1	8	15	1	1	1	1
Study Day	1	8	15	22	29	36	43	64	106	
	PRE ^a	EOI ^b		PRE ^a	EOI ^b		PRE ^a	EOI ^b	PRE ^a	EOI ^b
Trastuzumab	X	X					X	X	X	X
Pertuzumab	X	X	X ^c	X	X ^c	X ^c	X ^d	X	X	X
HER2 ECD	X									

EOI = end of infusion; PRE = pre-dose.

Note: Each blood sample will be 5 mL in volume. Separate PK samples will be collected for each drug (i.e., Trastuzumab & Pertuzumab)

- a. Sample can be collected up to 6 hours prior to administration of the specified drug (i.e., Trastuzumab or Pertuzumab).
- b. Sample can be collected up to 30 minutes after the end of infusion of the specified drug (i.e., Trastuzumab or Pertuzumab).
- c. Sample can be collected within ± 6 hours (relative to the start of infusion of the specified drug).
- d. Cycle 2 trough level = primary study endpoint - must be drawn on Study Day 43 (22 days after Cycle 2 dose) regardless of whether Cycle 3 dose is delayed or not given.

FIG. 32

31 / 34

Demographics

	Arm A 420mg (N=15)	Arm B 840mg (N=15)
Age (yr)		
Mean (SD)	60.6 (14.7)	57.3 (11.5)
Median	67.0	59.0
Min-Max	29-76	27-72
<65 (%)	6 (40.0)	10 (66.7)
≥65 (%)	9 (60.0)	5 (33.3)
Sex		
Male (%)	14 (93.3)	10 (66.7)
Female (%)	1 (6.7)	5 (33.3)
Race		
Asian (%)	9 (60.0)	8 (53.5)
White (%)	6 (40.0)	7 (46.7)
Weight (kg) at baseline		
Mean (SD)	67.55 (8.64)	67.69 (11.68)
Median	67.00	66.80
ECOG		
0	10 (66.7)	6 (40.0)
1	5 (33.3)	9 (60.0)

FIG. 33

32 / 34

Gastric Cancer History

	Pertuzumab 420mg (N=15)	Pertuzumab 840mg (N=15)
Extent of Disease on Entry		
n	15	15
Metastatic disease	13 (86.7%)	12 (80.0%)
Unresectable locally advanced disease	2 (13.3%)	3 (20.0%)
Primary Site		
n	15	15
Stomach	12 (80.0%)	13 (86.7%)
Gastroesophageal junction	3 (20.0%)	2 (13.3%)
Measurability		
n	15	15
Measurable disease	11 (73.3%)	13 (86.7%)
Non-measurable evaluable disease only	4 (26.7%)	2 (13.3%)
Histological Subtypes		
n	15	15
Intestinal	6 (40.0%)	5 (33.3%)
Diffuse	1 (6.7%)	3 (20.0%)
Not known	8 (53.3%)	7 (46.7%)

FIG. 34**Patient Disposition**

	Arm A Pertuzumab 840mg (N=15)	Arm B Pertuzumab 840mg (N=15)
Completed six cycles of pertuzumab (%)	5 (33.3)	5 (33.3)
Discontinued treatment (%)	1 (6.7)	3 (20.0)
Safety (%)	0	2 (13.3)
Adverse event (%)	0	1 (6.7)
Death (%)	0	1 (6.7)
Non-safety (%)	1 (6.7)	1 (6.7)
Physician decision (%)	0	1 (6.7)
Progression of disease (%)	1 (6.7)	0

FIG. 35

33 / 34


Overall Response Rate

	Arm A Pertuzumab 420mg (N=15)	Arm B Pertuzumab 840mg (N=15)
Total number of patients with tumor assessment at baseline	15	15
End of Cycle 3		
n	12	8
Partial response (%)	8 (66.7)	5 (62.5)
Stable disease (%)	2 (16.7)	2 (25.0)
Non-complete response/ non-progressive disease (%) ^a	1 (8.3)	1 (12.5)
Progressive disease (%)	1 (8.3)	0
End of Cycle 6		
n	4	4
Partial response (%)	3 (75.0)	3 (75.0)
Stable disease (%)	1 (25.0)	1 (25.0)

^a Non-complete response/non-progressive disease is stable disease in patients with non-measurable disease.

FIG. 36

34 / 34

FIG. 37**Pertuzumab - Day 42 C_{trough} (µg/mL)**

	JOSHUA 840/420mg	JOSHUA 840/840mg^a	CLEO 840/420mg
N	15	12	19
Mean (SD)	40.0 (17.3)	62.7 (29.1)	63.7 (46.8)
Median	36.7	65.9	56.0
Range	(15.8-78.7)	(17.1-135)	(19.3-245)
Geometric Mean	36.8	56.4	54.9
Estimated % of Patients ≥20 µg/mL ^b (95% CI)	91.6% (79.6-99.1)	98.6% (92.2-99.97)	94.1% ^c

^a Patient 8100 Day 42 concentration considered abnormally low (0.66 µg/mL) and was excluded from summary statistics

^b Estimated using Bootstrap analysis, assuming log-normal distribution

^c Based on observed concentrations

P4753R1WO.txt
Sequence Listing

<110> Genentech, Inc. et al.

<120> USES FOR AND ARTICLE OF MANUFACTURE INCLUDING HER2
DIMERIZATION INHIBITOR PERTUZUMAB

<130> P4753R1-WO

<141> 2012-10-11

<150> US 61/547,535

<151> 2011-10-14

<150> US 61/567,015

<151> 2011-12-05

<150> US 61/657,669

<151> 2012-06-08

<150> US 61/682,037

<151> 2012-08-10

<150> US 61/694,584

<151> 2012-08-29

<160> 22

<210> 1

<211> 195

<212> PRT

<213> Homo sapiens

<400> 1

Thr	Gln	Val	Cys	Thr	Gly	Thr	Asp	Met	Lys	Leu	Arg	Leu	Pro	Ala	1	5	10	15
Ser	Pro	Glu	Thr	His	Leu	Asp	Met	Leu	Arg	His	Leu	Tyr	Gln	Gly	20	25	30	
Cys	Gln	Val	Val	Gln	Gly	Asn	Leu	Glu	Leu	Thr	Tyr	Leu	Pro	Thr	35	40	45	
Asn	Ala	Ser	Leu	Ser	Phe	Leu	Gln	Asp	Ile	Gln	Glu	Val	Gln	Gly	50	55	60	
Tyr	Val	Leu	Ile	Ala	His	Asn	Gln	Val	Arg	Gln	Val	Pro	Leu	Gln	65	70	75	
Arg	Leu	Arg	Ile	Val	Arg	Gly	Thr	Gln	Leu	Phe	Glu	Asp	Asn	Tyr	80	85	90	
Ala	Leu	Ala	Val	Leu	Asp	Asn	Gly	Asp	Pro	Leu	Asn	Asn	Thr	Thr	95	100	105	
Pro	Val	Thr	Gly	Ala	Ser	Pro	Gly	Gly	Leu	Arg	Glu	Leu	Gln	Leu	110	115	120	
Arg	Ser	Leu	Thr	Glu	Ile	Leu	Lys	Gly	Gly	Val	Leu	Ile	Gln	Arg	125	130	135	
Asn	Pro	Gln	Leu	Cys	Tyr	Gln	Asp	Thr	Ile	Leu	Trp	Lys	Asp	Ile	140	145	150	
Phe	His	Lys	Asn	Asn	Gln	Leu	Ala	Leu	Thr	Leu	Ile	Asp	Thr	Asn	155	160	165	
Arg	Ser	Arg	Ala	Cys	His	Pro	Cys	Ser	Pro	Met	Cys	Lys	Gly	Ser				

170

175

180

Arg Cys Trp Gly Glu Ser Ser Glu Asp Cys Gln Ser Leu Thr Arg
185 190 195

<210> 2

<211> 124

<212> PRT

<213> Homo sapiens

<400> 2

Thr Val Cys Ala Gly Gly Cys Ala Arg Cys Lys Gly Pro Leu Pro
1 5 10 15

Thr Asp Cys Cys His Glu Gln Cys Ala Ala Gly Cys Thr Gly Pro
20 25 30

Lys His Ser Asp Cys Leu Ala Cys Leu His Phe Asn His Ser Gly
35 40 45

Ile Cys Glu Leu His Cys Pro Ala Leu Val Thr Tyr Asn Thr Asp
50 55 60

Thr Phe Glu Ser Met Pro Asn Pro Glu Gly Arg Tyr Thr Phe Gly
65 70 75

Ala Ser Cys Val Thr Ala Cys Pro Tyr Asn Tyr Leu Ser Thr Asp
80 85 90

Val Gly Ser Cys Thr Leu Val Cys Pro Leu His Asn Gln Glu Val
95 100 105

Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser Lys Pro
110 115 120

Cys Ala Arg Val

<210> 3

<211> 169

<212> PRT

<213> Homo sapiens

<400> 3

Cys Tyr Gly Leu Gly Met Glu His Leu Arg Glu Val Arg Ala Val
1 5 10 15

Thr Ser Ala Asn Ile Gln Glu Phe Ala Gly Cys Lys Lys Ile Phe
20 25 30

Gly Ser Leu Ala Phe Leu Pro Glu Ser Phe Asp Gly Asp Pro Ala
35 40 45

Ser Asn Thr Ala Pro Leu Gln Pro Glu Gln Leu Gln Val Phe Glu
50 55 60

Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp Pro
65 70 75

Asp Ser Leu Pro Asp Leu Ser Val Phe Gln Asn Leu Gln Val Ile
80 85 90

Arg Gly Arg Ile Leu His Asn Gly Ala Tyr Ser Leu Thr Leu Gln
95 100 105

Gly Leu Gly Ile Ser Trp Leu Gly Leu Arg Ser Leu Arg Glu Leu
110 115 120

P4753R1WO.txt

Gly Ser Gly Leu Ala Leu Ile His His Asn Thr His Leu Cys Phe
125 130 135
Val His Thr Val Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln
140 145 150
Ala Leu Leu His Thr Ala Asn Arg Pro Glu Asp Glu Cys Val Gly
155 160 165
Glu Gly Leu Ala

<210> 4
<211> 142
<212> PRT
<213> Homo sapiens

<400> 4
Cys His Gln Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro
1 5 10 15
Thr Gln Cys Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys
20 25 30
Val Glu Glu Cys Arg Val Leu Gln Gly Leu Pro Arg Glu Tyr Val
35 40 45
Asn Ala Arg His Cys Leu Pro Cys His Pro Glu Cys Gln Pro Gln
50 55 60
Asn Gly Ser Val Thr Cys Phe Gly Pro Glu Ala Asp Gln Cys Val
65 70 75
Ala Cys Ala His Tyr Lys Asp Pro Pro Phe Cys Val Ala Arg Cys
80 85 90
Pro Ser Gly Val Lys Pro Asp Leu Ser Tyr Met Pro Ile Trp Lys
95 100 105
Phe Pro Asp Glu Glu Gly Ala Cys Gln Pro Cys Pro Ile Asn Cys
110 115 120
Thr His Ser Cys Val Asp Leu Asp Asp Lys Gly Cys Pro Ala Glu
125 130 135
Gln Arg Ala Ser Pro Leu Thr
140

<210> 5
<211> 107
<212> PRT
<213> Mus musculus

<400> 5
Asp Thr Val Met Thr Gln Ser His Lys Ile Met Ser Thr Ser Val
1 5 10 15
Gly Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Ser
20 25 30
Ile Gly Val Ala Trp Tyr Gln Gln Arg Pro Gly Gln Ser Pro Lys
35 40 45
Leu Leu Ile Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp
50 55 60
Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile
65 70 75

Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln
80 85 90
Tyr Tyr Ile Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu
95 100 105
Ile Lys

<210> 6
<211> 119
<212> PRT
<213> Mus musculus

<400> 6
Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly
1 5 10 15
Thr Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Phe Thr Phe Thr
20 25 30
Asp Tyr Thr Met Asp Trp Val Lys Gln Ser His Gly Lys Ser Leu
35 40 45
Glu Trp Ile Gly Asp Val Asn Pro Asn Ser Gly Gly Ser Ile Tyr
50 55 60
Asn Gln Arg Phe Lys Gly Lys Ala Ser Leu Thr Val Asp Arg Ser
65 70 75
Ser Arg Ile Val Tyr Met Glu Leu Arg Ser Leu Thr Phe Glu Asp
80 85 90
Thr Ala Val Tyr Tyr Cys Ala Arg Asn Leu Gly Pro Ser Phe Tyr
95 100 105
Phe Asp Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser
110 115

<210> 7
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> Sequence is synthesized.

<400> 7
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
1 5 10 15
Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Ser
20 25 30
Ile Gly Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys
35 40 45
Leu Leu Ile Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Ser
50 55 60
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
65 70 75
Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
80 85 90
Tyr Tyr Ile Tyr Pro Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu
Page 4

Ile Lys

<210> 8
 <211> 119
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Sequence is synthesized.

<400> 8
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1 5 10 15
 Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Thr
 20 25 30
 Asp Tyr Thr Met Asp Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 35 40 45
 Glu Trp Val Ala Asp Val Asn Pro Asn Ser Gly Gly Ser Ile Tyr
 50 55 60
 Asn Gln Arg Phe Lys Gly Arg Phe Thr Leu Ser Val Asp Arg Ser
 65 70 75
 Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
 80 85 90
 Thr Ala Val Tyr Tyr Cys Ala Arg Asn Leu Gly Pro Ser Phe Tyr
 95 100 105
 Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 110 115

<210> 9
 <211> 107
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Sequence is synthesized.

<400> 9
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15
 Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser
 20 25 30
 Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys
 35 40 45
 Leu Leu Ile Tyr Ala Ala Ser Ser Leu Glu Ser Gly Val Pro Ser
 50 55 60
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
 65 70 75
 Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
 80 85 90
 Tyr Asn Ser Leu Pro Trp Thr Phe Gly Gln Gly Thr Lys Val Glu
 95 100 105

Ile Lys

<210> 10
 <211> 119
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Sequence is synthesized.

<400> 10
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1 5 10 15
 Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
 20 25 30
 Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 35 40 45
 Glu Trp Val Ala Val Ile Ser Gly Asp Gly Gly Ser Thr Tyr Tyr
 50 55 60
 Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser
 65 70 75
 Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
 80 85 90
 Thr Ala Val Tyr Tyr Cys Ala Arg Gly Arg Val Gly Tyr Ser Leu
 95 100 105
 Tyr Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 110 115

<210> 11
 <211> 214
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Sequence is synthesized.

<400> 11
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15
 Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Ser
 20 25 30
 Ile Gly Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys
 35 40 45
 Leu Leu Ile Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Ser
 50 55 60
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
 65 70 75
 Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
 80 85 90
 Tyr Tyr Ile Tyr Pro Tyr Thr Phe Gly Gln Gly Thr Lys Val Glu
 95 100 105
 Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro
 110 115 120

P4753R1WO.txt

Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu
125 130 135
Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val
140 145 150
Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu
155 160 165
Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr
170 175 180
Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu
185 190 195
Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn
200 205 210
Arg Gly Glu Cys

<210> 12
<211> 448
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized.

<400> 12
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
1 5 10 15
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Thr
20 25 30
Asp Tyr Thr Met Asp Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45
Glu Trp Val Ala Asp Val Asn Pro Asn Ser Gly Gly Ser Ile Tyr
50 55 60
Asn Gln Arg Phe Lys Gly Arg Phe Thr Leu Ser Val Asp Arg Ser
65 70 75
Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90
Thr Ala Val Tyr Tyr Cys Ala Arg Asn Leu Gly Pro Ser Phe Tyr
95 100 105
Phe Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala
110 115 120
Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
125 130 135
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp
140 145 150
Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu
155 160 165
Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly
170 175 180
Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu
Page 7

										185						190						195
Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	200	205	210					
Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	215	220	225					
His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	230	235	240					
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	245	250	255					
Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	260	265	270					
Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	275	280	285					
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	290	295	300					
Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	305	310	315					
Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	320	325	330					
Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	335	340	345					
Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	350	355	360					
Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	365	370	375					
Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	380	385	390					
Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	395	400	405					
Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	410	415	420					
Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	425	430	435					
Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly			440	445						

<210> 13
 <211> 214
 <212> PRT
 <213> Artificial sequence

 <220>
 <223> Sequence is synthesized.

 <400> 13
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val
 1 5 10 15
 Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Val Asn
 20 25 30

P4753R1WO.txt

Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys
35 40 45
Leu Leu Ile Tyr Ser Ala Ser Phe Leu Tyr Ser Gly Val Pro Ser
50 55 60
Arg Phe Ser Gly Ser Arg Ser Gly Thr Asp Phe Thr Leu Thr Ile
65 70 75
Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln
80 85 90
His Tyr Thr Thr Pro Pro Thr Phe Gly Gln Gly Thr Lys Val Glu
95 100 105
Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro
110 115 120
Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu
125 130 135
Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val
140 145 150
Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu
155 160 165
Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr
170 175 180
Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu
185 190 195
Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn
200 205 210
Arg Gly Glu Cys

<210> 14
<211> 449
<212> PRT
<213> Artificial sequence

<220>
<223> Sequence is synthesized.

<400> 14
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
1 5 10 15
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Asn Ile Lys
20 25 30
Asp Thr Tyr Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
35 40 45
Glu Trp Val Ala Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr
50 55 60
Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser
65 70 75
Lys Asn Thr Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
80 85 90
Thr Ala Val Tyr Tyr Cys Ser Arg Trp Gly Gly Asp Gly Phe Tyr
95 100 105

P4753R1WO.txt

Ala	Met	Asp	Tyr	Trp 110	Gly	Gln	Gly	Thr	Leu 115	Val	Thr	Val	Ser	Ser 120
Ala	Ser	Thr	Lys	Gly 125	Pro	Ser	Val	Phe	Pro 130	Leu	Ala	Pro	Ser	Ser 135
Lys	Ser	Thr	Ser	Gly 140	Gly	Thr	Ala	Ala	Leu 145	Gly	Cys	Leu	Val	Lys 150
Asp	Tyr	Phe	Pro	Glu 155	Pro	Val	Thr	Val	Ser 160	Trp	Asn	Ser	Gly	Ala 165
Leu	Thr	Ser	Gly	Val 170	His	Thr	Phe	Pro	Ala 175	Val	Leu	Gln	Ser	Ser 180
Gly	Leu	Tyr	Ser	Leu 185	Ser	Ser	Val	Val	Thr 190	Val	Pro	Ser	Ser	Ser 195
Leu	Gly	Thr	Gln	Thr 200	Tyr	Ile	Cys	Asn	Val 205	Asn	His	Lys	Pro	Ser 210
Asn	Thr	Lys	Val	Asp 215	Lys	Lys	Val	Glu	Pro 220	Lys	Ser	Cys	Asp	Lys 225
Thr	His	Thr	Cys	Pro 230	Pro	Cys	Pro	Ala	Pro 235	Glu	Leu	Leu	Gly	Gly 240
Pro	Ser	Val	Phe	Leu 245	Phe	Pro	Pro	Lys	Pro 250	Lys	Asp	Thr	Leu	Met 255
Ile	Ser	Arg	Thr	Pro 260	Glu	Val	Thr	Cys	Val 265	Val	Val	Asp	Val	Ser 270
His	Glu	Asp	Pro	Glu 275	Val	Lys	Phe	Asn	Trp 280	Tyr	Val	Asp	Gly	Val 285
Glu	Val	His	Asn	Ala 290	Lys	Thr	Lys	Pro	Arg 295	Glu	Glu	Gln	Tyr	Asn 300
Ser	Thr	Tyr	Arg	Val 305	Val	Ser	Val	Leu	Thr 310	Val	Leu	His	Gln	Asp 315
Trp	Leu	Asn	Gly	Lys 320	Glu	Tyr	Lys	Cys	Lys 325	Val	Ser	Asn	Lys	Ala 330
Leu	Pro	Ala	Pro	Ile 335	Glu	Lys	Thr	Ile	Ser 340	Lys	Ala	Lys	Gly	Gln 345
Pro	Arg	Glu	Pro	Gln 350	Val	Tyr	Thr	Leu	Pro 355	Pro	Ser	Arg	Glu	Glu 360
Met	Thr	Lys	Asn	Gln 365	Val	Ser	Leu	Thr	Cys 370	Leu	Val	Lys	Gly	Phe 375
Tyr	Pro	Ser	Asp	Ile 380	Ala	Val	Glu	Trp	Glu 385	Ser	Asn	Gly	Gln	Pro 390
Glu	Asn	Asn	Tyr	Lys 395	Thr	Thr	Pro	Pro	Val 400	Leu	Asp	Ser	Asp	Gly 405
Ser	Phe	Phe	Leu	Tyr 410	Ser	Lys	Leu	Thr	Val 415	Asp	Lys	Ser	Arg	Trp 420
Gln	Gln	Gly	Asn	Val 425	Phe	Ser	Cys	Ser	Val 430	Met	His	Glu	Ala	Leu 435
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	

440

445

<210> 15
 <211> 217
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Sequence is synthesized.

<400> 15
 Val His Ser Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 1 5 10 15
 Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln
 20 25 30
 Asp Val Ser Ile Gly Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys
 35 40 45
 Ala Pro Lys Leu Leu Ile Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly
 50 55 60
 Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 65 70 75
 Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Phe Ala Thr Tyr Tyr
 80 85 90
 Cys Gln Gln Tyr Tyr Ile Tyr Pro Tyr Thr Phe Gly Gln Gly Thr
 95 100 105
 Lys Val Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile
 110 115 120
 Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val
 125 130 135
 Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln
 140 145 150
 Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser
 155 160 165
 Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
 170 175 180
 Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 185 190 195
 Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys
 200 205 210
 Ser Phe Asn Arg Gly Glu Cys
 215

<210> 16
 <211> 449
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Sequence is synthesized.

<400> 16
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly
 1 5 10 15

Gly	Ser	Leu	Arg	Leu ₂₀	Ser	Cys	Ala	Ala	Ser ₂₅	Gly	Phe	Thr	Phe	Thr ₃₀
Asp	Tyr	Thr	Met	Asp ₃₅	Trp	Val	Arg	Gln	Ala ₄₀	Pro	Gly	Lys	Gly	Leu ₄₅
Glu	Trp	Val	Ala	Asp ₅₀	Val	Asn	Pro	Asn	Ser ₅₅	Gly	Gly	Ser	Ile	Tyr ₆₀
Asn	Gln	Arg	Phe	Lys ₆₅	Gly	Arg	Phe	Thr	Leu ₇₀	Ser	Val	Asp	Arg	Ser ₇₅
Lys	Asn	Thr	Leu	Tyr ₈₀	Leu	Gln	Met	Asn	Ser ₈₅	Leu	Arg	Ala	Glu	Asp ₉₀
Thr	Ala	Val	Tyr	Tyr ₉₅	Cys	Ala	Arg	Asn	Leu ₁₀₀	Gly	Pro	Ser	Phe	Tyr ₁₀₅
Phe	Asp	Tyr	Trp	Gly ₁₁₀	Gln	Gly	Thr	Leu	Val ₁₁₅	Thr	Val	Ser	Ser	Ala ₁₂₀
Ser	Thr	Lys	Gly	Pro ₁₂₅	Ser	Val	Phe	Pro	Leu ₁₃₀	Ala	Pro	Ser	Ser	Lys ₁₃₅
Ser	Thr	Ser	Gly	Gly ₁₄₀	Thr	Ala	Ala	Leu	Gly ₁₄₅	Cys	Leu	Val	Lys	Asp ₁₅₀
Tyr	Phe	Pro	Glu	Pro ₁₅₅	Val	Thr	Val	Ser	Trp ₁₆₀	Asn	Ser	Gly	Ala	Leu ₁₆₅
Thr	Ser	Gly	Val	His ₁₇₀	Thr	Phe	Pro	Ala	Val ₁₇₅	Leu	Gln	Ser	Ser	Gly ₁₈₀
Leu	Tyr	Ser	Leu	Ser ₁₈₅	Ser	Val	Val	Thr	Val ₁₉₀	Pro	Ser	Ser	Ser	Leu ₁₉₅
Gly	Thr	Gln	Thr	Tyr ₂₀₀	Ile	Cys	Asn	Val	Asn ₂₀₅	His	Lys	Pro	Ser	Asn ₂₁₀
Thr	Lys	Val	Asp	Lys ₂₁₅	Lys	Val	Glu	Pro	Lys ₂₂₀	Ser	Cys	Asp	Lys	Thr ₂₂₅
His	Thr	Cys	Pro	Pro ₂₃₀	Cys	Pro	Ala	Pro	Glu ₂₃₅	Leu	Leu	Gly	Gly	Pro ₂₄₀
Ser	Val	Phe	Leu	Phe ₂₄₅	Pro	Pro	Lys	Pro	Lys ₂₅₀	Asp	Thr	Leu	Met	Ile ₂₅₅
Ser	Arg	Thr	Pro	Glu ₂₆₀	Val	Thr	Cys	Val	Val ₂₆₅	Val	Asp	Val	Ser	His ₂₇₀
Glu	Asp	Pro	Glu	Val ₂₇₅	Lys	Phe	Asn	Trp	Tyr ₂₈₀	Val	Asp	Gly	Val	Glu ₂₈₅
Val	His	Asn	Ala	Lys ₂₉₀	Thr	Lys	Pro	Arg	Glu ₂₉₅	Glu	Gln	Tyr	Asn	Ser ₃₀₀
Thr	Tyr	Arg	Val	Val ₃₀₅	Ser	Val	Leu	Thr	Val ₃₁₀	Leu	His	Gln	Asp	Trp ₃₁₅
Leu	Asn	Gly	Lys	Glu ₃₂₀	Tyr	Lys	Cys	Lys	Val ₃₂₅	Ser	Asn	Lys	Ala	Leu ₃₃₀
Pro	Ala	Pro	Ile	Glu ₃₃₅	Lys	Thr	Ile	Ser	Lys ₃₄₀	Ala	Lys	Gly	Gln	Pro ₃₄₅
Arg	Glu	Pro	Gln	Val ₃₅₀	Tyr	Thr	Leu	Pro	Pro ₃₅₅	Ser	Arg	Glu	Glu	Met ₃₆₀

Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr
				365					370					375
Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu
				380					385					390
Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser
				395					400					405
Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln
				410					415					420
Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His
				425					430					435
Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys	
				440					445					

<210> 17
 <211> 10
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Sequence is synthesized.

<220>
 <221> Xaa
 <222> 10
 <223> Xaa is preferrably D or S

<400> 17
 Gly Phe Thr Phe Thr Asp Tyr Thr Met Xaa
 1 5 10

<210> 18
 <211> 17
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Sequence is synthesized.

<400> 18
 Asp Val Asn Pro Asn Ser Gly Gly Ser Ile Tyr Asn Gln Arg Phe
 1 5 10 15
 Lys Gly

<210> 19
 <211> 10
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Sequence is synthesized.

<400> 19
 Asn Leu Gly Pro Ser Phe Tyr Phe Asp Tyr
 1 5 10

<210> 20
 <211> 11
 <212> PRT
 <213> Artificial sequence

<220>

<223> Sequence is synthesized.

<400> 20

Lys Ala Ser Gln Asp Val Ser Ile Gly Val Ala
1 5 10

<210> 21

<211> 7

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized.

<220>

<221> Xaa

<222> 5

<223> Xaa is preferably R or L

<220>

<221> Xaa

<222> 6

<223> Xaa is preferably Y or E

<220>

<221> Xaa

<222> 7

<223> Xaa is preferably T or S

<400> 21

Ser Ala Ser Tyr Xaa Xaa Xaa
1 5

<210> 22

<211> 9

<212> PRT

<213> Artificial sequence

<220>

<223> Sequence is synthesized.

<400> 22

Gln Gln Tyr Tyr Ile Tyr Pro Tyr Thr
1 5