




AU9534245


(12) PATENT ABRIDGMENT (11) Document No. AU-B-34245/95
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 701064

- (54) Title
PHOTOSTABLE COSMETIC LIGHT SCREENING COMPOSITIONS
- International Patent Classification(s)
 (51)⁶ **A61K 007/44**
- (21) Application No. : **34245/95** (22) Application Date : **13.10.95**
- (30) Priority Data
- | | | |
|-----------------|-----------------|---------------------------------------|
| (31) Number | (32) Date | (33) Country |
| 94810601 | 14.10.94 | EP EUROPEAN PATENT OFFICE (EP) |
| 95102337 | 20.02.95 | EP EUROPEAN PATENT OFFICE (EP) |
- (43) Publication Date : **26.04.96**
- (44) Publication Date of Accepted Application : **21.01.99**
- (71) Applicant(s)
GIVAUDAN-ROURE (INTERNATIONAL) SA
- (72) Inventor(s)
PIERCLAUDIO BERNASCONI; HANS ULRICH GONZENBACH
- (74) Attorney or Agent
SPRUSON & FERGUSON , GPO Box 3898, SYDNEY NSW 2001
- (56) Prior Art Documents
US 5403944
US 4387089
- (57) Claim



1. Photostable, cosmetic light-screening composition, characterised in that it comprises, in a cosmetically acceptable vehicle containing at least one fatty phase, about 0,5 to about 5%, by weight, of a dibenzoylmethane type UV-A screening agent and at least about 0,1% to about 20%, by weight, of a polymer UV-B filter of the benzylidene malonate silicone type comprising an organosiloxane having at least one unit of the general formula



or



wherein




and

.. /2


V is



or



any other units present in the said siloxanes being those represented by the general formula



(III)

wherein R represents a C₁₋₈ alkyl or an aryl group, R¹ is a hydrogen atom or a C₁₋₅ alkyl group, R² is a hydrogen atom, a C₁₋₅ alkyl group or a group OR¹, R³ is a C₁₋₅ alkyl group, R'' represents a hydrogen atom, a monovalent C₁₋₈ hydrocarbon or halogenated hydrocarbon group, a has a value of 0, 1 or 2, b has a value of 0, 1, 2 or 3 and n has a value of from 1 to 6, provided that the group -U-O- or the group -V-O- and the two R² groups are linked to the aromatic ring (C₆...) at the para- and both meta-positions in relation to the group -CH=C[C(O)OR³]₂, the weight ratio of the silicone to the dibenzoylmethane derivative being not less than about 0.1, and not more than about 25.

AUSTRALIA
PATENTS ACT 1990

COMPLETE SPECIFICATION

FOR A STANDARD PATENT

ORIGINAL

Name and Address
of Applicant: Givaudan-Roure (International) SA
CH-1214 Vernier
Geneve
SWITZERLAND


Actual Inventor(s): Pierclaudio Bernasconi and Hans Ulrich Gonzenbach

Address for Service: Spruson & Ferguson, Patent Attorneys
Level 33 St Martins Tower, 31 Market Street
Sydney, New South Wales, 2000, Australia


Invention Title: Photostable Cosmetic Light Screening Compositions

The following statement is a full description of this invention, including the best method of performing it known to me/us:-

wherein R represents a C₁₋₈ alkyl or an aryl group, R¹ is a hydrogen atom or a C₁₋₅ alkyl group, R² is a hydrogen atom, a C₁₋₅ alkyl group or a group OR¹, R³ is a C₁₋₅ alkyl group, R'' represents a hydrogen atom, a monovalent C₁₋₈ hydrocarbon or halogenated hydrocarbon group, a has a value of 0, 1 or 2, b has a value of 0, 1, 2 or 3 and n has a value of from 1 to 6, provided that the




or the




10

groups and the two R² groups are linked to the aromatic ring (C₆...) at the para- and both meta-positions in relation to the group -CH=C[C(O)OR³]₂, and the two vertical strokes designate either the alkylene derivative




15 or the alkylidene derivative




in case of formula (I'),

or the saturated derivative with a linear chain




or the saturated derivative with a branched chain




in case of formula (II'),

the weight ratio of the silicone to the dibenzoylmethane derivative being not less than ca. 0.1, preferably not less than ca. 1.0, and not more than ca. 25, preferably not more than ca. 8 to 10.

An alternative designation for the subject matter of compounds I' and II' is thus




or




wherein

U is




or




and


V is




or



Formula I encompasses thus, as pointed out above, the alkylidene derivative Ia and the alkylene derivative Ib:







The preparation and the properties of these benzylidene malonate silicone types are described in WO 92/20690 of November 26, 1992.

The ratio of the alkylene to the alkylidene derivative is not critical at all, and is normally between ca. 5- ca. 50% : ca. 95-50% w/w.

Formula II encompasses the saturated linear (IIa) and saturated branched (IIb) derivatives:



The preparation and properties of these benzylidene malonate silicone types are described in EP 358 584 (USP 5053290) and FR 2 642 967-A1 (USP 5415854).

The suitable dimensions for the various parameters of the definition of the above captioned formula III are equally derivable from said WO publication. Thus, in the general formulae I and II the radicals can be defined as follows:


R may be for example methyl, ethyl, butyl or phenyl.

R" is hydrogen or a monovalent hydrocarbon or halogenated hydrocarbon group having up to 8 carbon atoms, for example alkyl, alkenyl, aryl, alkaryl, aralkyl and also halogen substituted alkyl, alkenyl, aryl, alkaryl and aralkyl are such groups. Particular examples include methyl, ethyl, vinyl, phenyl, tolyl, ethylphenyl, dimethylphenyl, benzyl, phenethyl and 3,3,3- trifluoropropyl, etc.

R³ denotes alkyl groups having up to 5 carbon atoms, as for example methyl, ethyl, propyl, isopropyl, butyl, secondary butyl, isobutyl, pentyl and neopentyl, etc.


Examples for the C₁₋₅ alkyl radicals of R¹ and R² are as given above. Suitable radicals OR¹ are methoxy, ethoxy, propoxy, isopropoxy, etc. From the above listing, it can also be deduced that the alkyl and the alkoxy radicals may in the present context represent straight-chain and branched radicals.

- 5 It is preferred that at least 80% of all R and R" groups are methyl groups, most preferably substantially all R and R" groups are methyl groups. It is also preferred that R¹ is either hydrogen, methyl or ethyl, most preferably hydrogen. Preferably each R² group is hydrogen or one R² group is a hydrogen, while the other one is an alkoxy group, preferably methoxy or
10 ethoxy. R³ is preferably methyl or ethyl, a is preferably 1 while b is preferably 2, making the organosilicon compound a substantially linear or cyclic diorgano-siloxane polymer. However, if the diorganosiloxane - see, e.g., Table 1, the second formula - is a substantially linear polymer at least two endblocking units must be present, thus requiring the presence of 2
15 units in which a has a value of 2, two units in which the value of b is 3 or one unit wherein a is 2 and one unit in which b is 3. n is preferably 1, 2 or 3. Suitable preferred polymers have therefore either the general formula





wherein R and R" are as defined above, X denotes a group Y or a group R"

and Y denotes an alkylene group of the formula




or the corresponding alkylidene derivative in case of formula (I),

or, in case of formula (II), one of the groups





In the above captioned formulae, r has a value of from 0 to 130, s has a value of from 0 to 20, whereby at least one X denotes Y in the case that s=0; t has a value of from 0 to 10, v has a value of from 1 to 10 and v+t has a value of
30 at least 3, and R¹, R² and R³ are as above.

In the substituent Y of the organosilicon compounds, the above captioned alkylene group




or the corresponding above captioned alkylidene derivative,

5 or the groups







may occupy the meta-position or the para-position of the aromatic ring (C₆...) in relation to the group $-CH=C[C(O)OR^3]_2$.

10 Preferably the para-position is thus occupied. The groups R² occupy the remaining two positions out of the para- and meta-positions in relation to the group $-CH=C[C(O)OR^3]_2$. Examples of preferred substituents Y thus include




or the corresponding alkylidene derivatives,
or the saturated linear structures, such as:





or the corresponding saturated branched derivatives,
namely those carrying the structure



- 5 The material organosilicon compounds have at least one unit falling within the general formulae (I) or (II), preferably at least 2. Suitable organosilicon compounds are polymeric materials which may be homopolymers consisting only of such units (I) or (II), or they may be copolymers containing both units (I) or (II) and units having the general
- 10 formula (III). The organosilicon compounds may vary from freely flowing liquids to highly viscous gum-like materials or resinous solids. Preferred, at least for cosmetic applications, are the liquid substantially linear organosiloxane homopolymers and copolymers, for example those having a viscosity (in c St) of from 100 to 20000mm²/s, more preferably 500 to
- 15 5000mm²/s as these are more easily mixed with other ingredients to make cosmetic compositions and as they will spread more easily onto the skin.

Organosilicon compounds which are especially preferred are those wherein the number of units (I) or (II) is limited to a maximum of 20% to 25% of the total number of siloxane units in the molecule. For maximum

20 efficiency in its U.V. absorbing property it is preferred that the number of

units (I) or (II) be limited to 10% to 12% or less of the total. The units of formula (I) or (II) may be distributed randomly in an organosiloxane polymer, they may be end-blocking units of the polymer or they may be located at the end of the polymer and pending in chain of the polymer at the same time. Units of the general formula (I) or (II) are conveniently situated at the end of the organosiloxane polymer forming one or more endblocking units of the polymer. In one class of the preferred organosilicon compounds which are substantially linear polyorganosiloxane polymers, both endblocking units have a structure represented by the general formula (I) or (II), while all other units are according to the general formula (III). The preferred organosilicon compounds have four units of the formula (I) or (II) and a larger number of units according to the general formula (III), e.g. 6 to 130, especially 8 to 80.

The organosilicon compounds are themselves effective in absorbing ultra violet radiation in the erythemic region (290 - 320nm) which makes them particularly suitable for use in cosmetic sunscreen preparations where absorption in the UV-B region is particularly desirable. Most preferred for this application are those that have a maximum absorbance at 300 - 320nm.

As far as the UV-A light screen agent of the novel combinations is concerned, the preferred compound is 4-tert. butyl-4'-methoxydibenzoylmethane, as disclosed e.g. in USP 4 387 089 or CH-Patent 642 536.

Other suitable compounds of this particular type are: 2-methyldibenzoylmethane, 4-methyl-dibenzoyl-methane, 4-isopropyldibenzoylmethane, 4-tert-butyldibenzoyl-methane, 2,4-dimethyldibenzoylmethane, 2,5-dimethyldibenzoylmethane, 4,4'-diisopropyldibenzoylmethane, 2-methyl-5-isopropyl-4'-methoxydibenzoylmethane, 2-methyl-5-tert-butyl-4'-methoxydibenzoylmethane, 2,4-dimethyl-4'-methoxydibenzoylmethane and 2,6-dimethyl-4-tert-butyl-4'-methoxydibenzoylmethane.

The function of the polymer filter containing formula I units is not only to provide the necessary so-called A + B total block in the final composition, in as far as the absorption of the UV-radiation is concerned, but also to photostabilize the involved UV-A screening agents, i.e. to guarantee a constant protection during prolonged exposure to the UV-light. This way, if a repeated application of the cosmetic formulation at various intervals is required, these intervals can be extended.

The present invention relates thus also to a process for stabilising dibenzoylmethane UV-A screening agents with respect to UV radiation of wavelengths between 290 and 400 nm, characterised in that 0,1 to 20 % by weight of the polymer filter is added to 0,5 to 5 % by weight of the dibenzoyl-
5 methane UV-A screening agent, the weight ratio of the silicone to the dibenzoylmethane derivative being not less than 0.1 and not more than 25.

The desired stabilization of the material UV-A filters is easily established by strictly parallel experiments with the respective UV-A filters and the novel combinations using an appropriately equipped Xenon lamp as
10 a solar simulator. Irradiated are standard preparations of the investigated products, e.g. solutions in cosmetic solvents, the resulting sunscreen being spread on quartz plates. The stabilizing effect is directly correlated to the difference in absorbance at $\lambda_{\max}$ before and after the irradiation. For an satisfactory effect, the structures of I and II are essential.

15 Even a slight variation of the structure in the molecules I and II leads to an unsatisfactory respective stabilization; under slight variation there is understood the "removal" of an ester group in I and II, thus resulting in cinnamates (namely the benzylidene acetates-CH=C-C(O)OR³), viz. EP 305059.

20 Both components of the present combination of the light-screening agents are lipophilic. The cosmetic formulations contain thus at least one fatty phase, and the formulations can consequently present themselves in the form of emulsions, lotions or gels.

Suitably the cosmetic screening composition takes the form of an oil, a
25 lotion, a gel, a solid stick, an emulsion, e.g. cream, milk or of a vesicular dispersion of ionic or nonionic amphiphilic lipids, an aerosol, a spray, a foam, a powder, a shampoo, a hair conditioner or lacquer or a make-up, etc.

The usual solvents known to the skilled practitioner can be used for the preparation of these forms, e.g. oils, waxes, alcohols, polyols, etc. The
30 preferred agents are fatty acids, esters, fatty alcohols, but also ethanol, isopropanol, propylene glycol, glycerine, etc.

The cosmetic formulations may contain further adjuvants, e.g. further solvents, thickeners, emollients, emulsifiers, humectants, tensides, preservatives, antifoams, fragrances, oils, waxes, lower polyols and
35 monohydric alcohols, propellants, silicones, colourings and pigments, etc.

Other UV-B-filters may also be incorporated. Examples are given in USP 4 387 089 mentioned above. Particularly in case of emulsions, such UV filters may, naturally, also be water-soluble derivatives. As still further suitable UV-B filters, microfine pigments, such as the usual micropigments
5 of metal oxides may be used.

In case of protection of the hairs, the suitable formulations are shampoos, conditioners, lotions, gels, emulsions, dispersions, lacquers, etc.

The preparation of all these formulations is well known to the skilled artisan in this field.

10 Examples of preferred polysiloxanes used for the present purpose are the ones of Table I below.

Further suitable polysiloxanes are those of Examples 3 to 6 of WO 92/20690, and also the product of Comparative Example 1 of WO 92/20690, the latter material representing a product encompassed by Formula II
15 above.


Table I

Illustrative Compounds and Formulae


The first column relates to the material Example of WO 92/20690 dealing with the preparation of products of this type.

4	s = ca. 4	r = ca. 60	R ² = H (most preferred, "Polysiloxane A")
4	s = ca. 15	r = ca. 127	R ² = H
4	s = ca. 15	r = ca. 116	R ² = H
4	s = ca. 11	r = ca. 51	R ² = H

4	s = ca. 4	r = ca. 60	R ² = OMe
4	s = ca. 15	r = ca. 127	R ² = OMe
4	s = ca. 15	r = ca. 116	R ² = OMe
4	s = ca. 11	r = ca. 51	R ² = OMe




3	r = ca. 7	R ² = H
3	r = ca. 7	R ² = OMe




The above compounds contain ca. 20% of the alkylene isomers; their preparation can be illustrated as follows:

- A. 39.84 g of [[4-(2-propynyloxy)phenyl]methylene]-diethyl ester were dissolved in 100 g of toluene and heated under nitrogen to about 70°C. 39.96 g of a hydrosiloxane having a degree of polymerisation of 8 and 25 mpc (mole %) SiH groups (9.7% SiH) were then added dropwise after a platinum complex was also added, giving 10^{-4} mole of Pt per mole of SiH of the hydrosiloxane. The mixture was heated to reflux and maintained until all SiH had disappeared of the infrared spectroscopic analysis. It was then allowed to cool to room temperature. The toluene was then evaporated to leave after washing 60.5 g of a brown, viscous polymer having the average structure
- 10 $R^4-[(CH_3)_2SiO]_8-R^4$, wherein R has the formula




Only 0.34% by weight of the total reaction product of unreacted [[4-(2-propynyloxy)phenyl]methylene]-diethyl ester was present in the end product.

- B. 13.28 g of [[4-(2-propynyloxy)phenyl]methylene]-diethyl ester were dissolved in 75 g of toluene and heated under nitrogen to about 70°C. 44 g of a hydrosiloxane having a degree of polymerisation of 65 and 6 mpc SiH groups (2.36% SiH) were then added dropwise after a platinum complex was also added, giving 10^{-4} mole of Pt per mole of SiH of the hydrosiloxane. The mixture was heated to reflux and maintained until all SiH had disappeared of the infrared spectroscopic analysis. It was then allowed to cool to room temperature. The toluene was then evaporated to leave after washing 52 g of a brown, viscous polymer having the average structure
- 20



wherein R^4 has the formula



25

("Polysiloxane A").

Examples

1. A sunscreen cream is prepared with the following ingredients:

A)	% (w/w)
5	Stearic acid (octadecanoic acid) 10.0
	Butylmethoxy dibenzoylmethane 2.0 (1-[4-(1,1-dimethylethyl)phenyl]-3-(4-methoxyphenyl)- 1,3-propanedione) (sold under the trade name PARSOL 1789 by Givaudan-Roure S.A.)
10	Glyceryl mono myristate 5.0 (tetradecanoic acid ester with 1,2,3-propanetriol)
	Cetyl alcohol ((hexadecanol) 2.0
	Coco-caprylate/caprate 10.0 (Mixture of esters of coconut alcohol and n-octanoic acid and n-decanoic acid) (sold under the trade name CETIOL LC by Henkel)
15	Polysiloxane A 10.0
	Dea Cetylphosphate 3.0 (Diethanolamine salt of hexadecyl phosphate) (sold under the trade name AMPHISOL by Givaudan-Roure S.A.)
20	EDTA Na ₂ (disodium ethylenediamine tetra acetic acid) 0.1
B	Deionized water 51.0
	Propyleneglycol (1,3-propanediol) 6.0
25	Mixture of parabens in phenoxyethanol 0.6 (Mixture of methyl, ethyl, propyl and butyl esters of 4-hydroxy benzoic acid) (sold under the trade name Phenonip® by Nipa Laboratoires Ltd.)
30	C Conventional fragrance 0.3

2. A sunscreen lotion is prepared with the following ingredients:

A		% (w/w)
	Glyceryl mono myristate	5.0
5	Cetyl alcohol	2.0
	Butylmethoxy dibenzoylmethane	1.0
	Isopropyl myristate (2-methylethyl tetradecanoate)	7.0
	Oleyl alcohol (octadecanol)	3.0
	Polysiloxane A	5.0
10	Dea Cetylphosphate	3.0
	EDTA Na ₂	0.1
B		
	Deionized water	67.0
	Propyleneglycol	6.0
15	Mixture of parabens in phenoxyethanol	0.6
C		
	Fragrance	0.3

3. A sunscreen lotion is prepared with the following ingredients:

A		% (w/w)
	Glyceryl mono stearate	4.0
5	Butylmethoxy dibenzoylmethane	1.0
	Cetyl alcohol	1.0
	Coco-caprylate/caprate	6.0
	Polysiloxane A	10.0
10	Potassium cetyl phosphate (potassium salt of hexadecyl phosphate) (sold under the trade name AMPHISOL K by Givaudan-Roure S.A.)	2.0
	EDTA Na ₂	0.1
15	B Deionized water	59.36
	Carbomer (1% dispersion in water, homo polymer of acrylic acid crosslinked with an allyl ether of sucrose) (sold under the trade name Carbopol 981® by B.F. Goodrich)	10.0
20	Propyleneglycol	5.0
	Potassium hydroxide, 10% solution	0.64
	Mixture of parabens in phenoxyethanol	0.6
25	C Fragrance	0.3


4. A water in silicone cream (a W/O formulation) is prepared with the following ingredients:

A		% (w/w)
5	Butylmethoxy dibenzoylmethane	1.0
	C _{12/15} alkyl benzoate (mixture of dodecyl & pentadecyl benzoate) (sold under the trade name FINSOLV TN by Finetex Corp.)	5.0
10	Silicone oil (copolyol of dimethicone and cyclomethicone) (sold under the trade name DC-3225 by Dow Corning)	10.0
	Silicone oil (mixture of cyclomethicone and dimethiconol) (sold under the trade name DC-1401 by Dow Corning)	10.0
	Polysiloxane A	10.0
15	EDTA Na ₂	0.1
B		
	Deionized water	56.0
	Sodium chloride	4.0
	Propyleneglycol	3.0
20	Mixture of parabens in phenoxyethanol	0.6
C		
	Fragrance	0.3


In the same way, the Polysiloxane A can be replaced in the above formulations by the Compound of Comparative Example 1 of WO 92/20690.

The claims defining the invention are as follows:

1. Photostable, cosmetic light-screening composition, characterised in that it comprises, in a cosmetically acceptable vehicle containing at least one fatty phase, about 0,5 to about 5%, by weight, of a dibenzoylmethane type UV-A screening agent and at least about 0,1% to about 20%, by weight, of a polymer UV-B filter of the benzylidene malonate silicone type comprising an organosiloxane having at least one unit of the general formula





or



10

wherein

U is




and


15 V is



or



any other units present in the said siloxanes being those represented by the general formula



20

wherein R represents a C₁₋₈ alkyl or an aryl group, R¹ is a hydrogen atom or a C₁₋₅ alkyl group, R² is a hydrogen atom, a C₁₋₅ alkyl group or a group OR¹, R³ is a C₁₋₅ alkyl group, R'' represents a hydrogen atom, a monovalent C₁₋₈ hydrocarbon or halogenated



hydrocarbon group, a has a value of 0, 1 or 2, b has a value of 0, 1, 2 or 3 and n has a value of from 1 to 6, provided that the group -U-O- or the group -V-O- and the two R² groups are linked to the aromatic ring (C₆...) at the para- and both meta-positions in relation to the group -CH=C[C(O)OR³]₂, the weight ratio of the silicone to the dibenzoylmethane derivative being not less than about 0.1, and not more than about 25.

2. A composition according to claim 1 wherein said dibenzoylmethane type UV-A screening agent is present in an amount of about 1 to about 4% by weight.

3. A composition according to claim 1 or claim 2 wherein said polymer UV-B filter of the benzylidene malonate silicon type is present in an amount of about 0.5 to about 15% by weight.

4. A composition according to any one of claims 1 to 3 wherein the weight ratio is not less than about 1.

5. A composition according to any one of claims 1 to 4 wherein the weight ratio is not more than about 8 to 10.

6. A composition according to any one of claims 1 to 5, wherein the ratio of the alkylene to the alkylidene derivative is about 5 - 50 : about 95-50 % w/w, or in case of the saturated derivatives, the linear derivatives is the predominant derivative.

7. A composition according to claim 6 wherein the ratio of the alkylene to the alkylidene derivative is about 20-30 : 80-70 % w/w.

8. A composition according to any one of claims 1 to 7, wherein at least 80% of all R and R" groups of (I) or (II) are methyl groups.

9. A composition according to any one of claims 1 to 8, wherein each R¹ of (I) is selected from hydrogen, methyl and ethyl.

10. A composition according to any one of the preceding claims, wherein at least one R² group of (I) or (II) is hydrogen, the other being hydrogen, methoxy or ethoxy.

11. A composition according to any one of the preceding claims, wherein R³ of (I) is methyl or ethyl.

12. A composition according to any one of the preceding claims, wherein each R¹ and R² group of (I) or (II) is hydrogen.

13. A composition according to any one of the preceding claims, wherein the polymer of I and II are linear or cyclic diorganosiloxane polymers.

14. A composition according to any one of the preceding claims, wherein the group U-O- or the group V-O- occupy the para-position on the aromatic ring in relation to the group -CH=C[C(O)OR³]₂, and both groups R² occupy the meta-positions on the aromatic ring in relation to the group -CH=C[C(O)OR³]₂.

15. A composition according to any one of the preceding claims, wherein the number of units of (I) or (II) are limited to a maximum of about 20% to about 25% of the total number of siloxane units in the molecule.



16. A composition according to any one of the preceding claims, wherein the number of units of (I) or (II) are limited to about 10% to about 12% or less of the total number of units.

17. A composition according to any one of the preceding claims, wherein units of the general formula (I) or (II) are situated at the end of the organosiloxane compound.

18. A composition according to any one of the preceding claims, wherein the polysiloxane of I or II has four end-blocking units of formula (I) or (II) and from 6 to 130 units of the general formula (III).

19. Composition according to any one of the preceding claims, characterised in that the dibenzoylmethane derivative is selected from the following compounds: 2-methyldibenzoylmethane, 4-methyl-dibenzoylmethane, 4-isopropyldibenzoylmethane, 4-tert-butyldibenzoyl-methane, 2,4-dimethyldibenzoylmethane, 2,5-dimethyldibenzoylmethane, 4,4'-diiso-propyldibenzoylmethane, 4-tert-butyl-4'-methoxydibenzoylmethane, 2-methyl-5-isoproyl-4'-methoxydibenzoylmethane, 2-methy-5-tert-butyl-4'-methoxydibenzoylmethane, 2,4-dimethyl-4'-methoxydibenzoylmethane and 2,6-dimethyl-4-tert-butyl-4'-methoxydibenzoylmethane.

20. Composition according to any one of the preceding claims, characterised in that the dibenzoylmethane derivative is 4-tert-butyl-4'-methoxydibenzoylmethane.

21. Composition according to any one of the preceding claims, characterised in that it takes the form of an oil, a lotion, a gel, a solid stick, an emulsion e.g. cream, milk or of a vesicular dispersion of ionic or nonionic amphiphilic lipids, an aerosol, a spray, a foam, a powder, a shampoo, a hair conditioner or lacquer or a make-up.

22. Composition according to any one of the preceding claims, characterised in that it contains, in addition, cosmetic adjuvants selected from solvents, thickeners, emollients, emulsifiers, humectants, tensides, preservatives, antifoams, fragrances, oils, waxes, lower polyols and monohydric alcohols, propellants, silicones, colourings and pigments.

23. Composition according to anyone of the preceding claims, characterised in that it contains, in addition, (a) further water-soluble or lipophilic UV screening agent(s).

24. Process for a cosmetic treatment of the human epidermis or the hairs in view of its protection against UV radiation of wavelengths between 290 and 400 nm, characterised in that it consists in applying to the skin or the hairs an effective quantity of a cosmetic screening composition as defined in any one of claims 1 to 23.

25. Process for stabilising dibenzoylmethane UV-A screening agents with respect to UV radiation of wavelengths between 290 and 400 nm, characterised in that 0.1 to 20% by weight of the polymer filter defined in claim 1 is added to 0.5 to 5% by weight of the dibenzoylmethane UV-A screening agent, the weight ratio of the silicone to the dibenzoylmethane derivative being not less than 0.1 and not more than 25.

26. Photostable, cosmetic light-screening composition, substantially as hereinbefore described with reference to any one of the Examples.



27. Process for stabilising dibenzoylmethane UV-A screening agents with respect to UV radiation of wavelengths between 290 and 400 nm, substantially as hereinbefore described with reference to any one of the Examples.

Dated 25 November, 1998
Givaudan-Roure (International) SA

5

Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON

W
B
A
A


8
0
1
A

AUSTRALIAN
PA Mh CE


Photostable Cosmetic Light Screening Compositions

Abstract

Photostable, cosmetic light screening compositions comprise, in a cosmetically acceptable vehicle containing at least one fatty phase, about 0.5 to about 5%, in particular about 1 to about 4% by weight, of a dibenzoylmethane type UV-A screening agent and at least about 0.1% to about 20%, in particular about 0.5 to about 15% by weight, of a polymer UV-B filter of the benzylidene malonate silicone type comprising an organosiloxane having at least one unit of the general formula




10 or



wherein U is $\text{-(CR}_2^1\text{)}_n$ or $\text{-(CR}_2^1\text{)}_n\text{-CH}(\text{CR}_2^1\text{)}_n$ and V is $\text{-CR}_2^1\text{-CHR}^1\text{-(CR}_2^1\text{)}_n$ or

$\text{-CR}_2^1\text{-CHR}_2^1$, any other units present in the said siloxanes being those represented by

the general formula



15

wherein R represents a C_{1-8} alkyl or an aryl group, R^1 is a hydrogen atom or a C_{1-5} alkyl group, R^2 is a hydrogen atom, a C_{1-5} alkyl group or a group OR^1 , R^3 is a C_{1-5} alkyl group, R^* represents a hydrogen atom, a monovalent C_{1-8} hydrocarbon or halogenated hydrocarbon group, a has a value of 0, 1 or 2, b has a value of 0, 1, 2 or 3 and n has a value of from 1 to 6, provided that the group -U-O- or the group -V-O- and the two R^2 groups are linked to the aromatic ring (C_6) at the para- and both meta-positions in relation to the group $\text{-CH=C[C(O)OR}^3\text{]}_2$, the weight ratio of the silicone to the dibenzoylmethane derivative being not less than ca. 0.1, preferably not less than ca. 1.0 and not more than ca. 26, preferably not more than ca. 8 to 10.